



US007205143B2

(12) **United States Patent**  
**Wonderling et al.**

(10) **Patent No.:** **US 7,205,143 B2**  
(45) **Date of Patent:** **Apr. 17, 2007**

(54) **FELINE IL-12 SINGLE CHAIN NUCLEIC ACID MOLECULES**

(75) Inventors: **Ramani S. Wonderling**, Waukegan, IL (US); **Karen L. Boroughs**, Fort Collins, CO (US)

(73) Assignee: **Heska Corporation**, Loveland, CO (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 237 days.

(21) Appl. No.: **10/821,670**

(22) Filed: **Apr. 9, 2004**

(65) **Prior Publication Data**

US 2004/0248261 A1 Dec. 9, 2004

**Related U.S. Application Data**

(62) Division of application No. 09/917,265, filed on Jul. 27, 2001, now Pat. No. 6,818,444.

(60) Provisional application No. 60/223,016, filed on Aug. 4, 2000.

(51) **Int. Cl.**

- C12N 15/00** (2006.01)
- C12N 15/19** (2006.01)
- C12N 15/12** (2006.01)
- C12N 15/62** (2006.01)
- C12N 15/86** (2006.01)
- C12N 15/85** (2006.01)

(52) **U.S. Cl.** ..... **435/252.3**; 435/325; 435/320.1; 536/23.1; 536/23.5

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

- 5,858,778 A 1/1999 Alnemri et al. .... 435/325
- 5,912,324 A 6/1999 Okamura et al. .... 530/351
- 5,914,253 A 6/1999 Okamura et al. .... 435/61.52
- 5,919,790 A 7/1999 Allen et al. .... 514/278
- 5,942,224 A 8/1999 Miwa et al. .... 424/85.7
- 5,945,310 A 8/1999 Young et al. .... 435/69.52
- 5,985,863 A 11/1999 Su et al. .... 438/22
- 6,013,268 A 1/2000 Reed ..... 424/269.1

**FOREIGN PATENT DOCUMENTS**

- WO WO 98/51327 11/1998
- WO WO 99/07851 2/1999
- WO WO 99/50425 10/1999
- WO WO 00/12555 3/2000

**OTHER PUBLICATIONS**

- Argyle, et al., Accession No. Y11133, Jul. 7, 1999.
- Argyle, et al., *Immunogenetics*, Jun. 1999, vol. 49, No. 6, pp. 541, abstract only.
- Belke-Louis, G.F., Accession No. U49085, Mar. 13, 1996.
- Belke-Louis, G.F., Accession No. U49100, Mar. 13, 1996.

Carr, et al., *Journal of Virology*, Oct. 1997, vol. 71, No. 10, pp. 7799-7803.

Cerretti, et al., Accession No. M87507, May 18, 1995.

Culhane, et al., Accession No. AJ222813, Aug. 11, 1998.

Johnson, et al., Accession No. AF031351, Nov. 3, 1998.

Keane, et al., Accession U14647, Jan. 27, 1996.

Lee, et al., *Human Gene Therapy*, Mar. 1, 1998, vol. 9, pp. 457-465.

Lieschke, et al., *Nature Biotechnology*, vol. 15, Jan. 1997, pp. 35-40.

Lode, et al., *Proc. Natl. Acad. Sci. USA*, vol. 95, Mar. 1998, pp. 2475-2480.

Molineaux, et al., Accession No. L28095, Jul. 27, 1994.

Mortola, et al., *J Vet Med Sci*, 1998, vol. 60, No. 11, pp. 1181-1185.

Nicolson, L., et al., Accession No. Y11131, Jul. 7, 1999.

Okamura, H., Accession No. D49949, Feb. 10, 1999.

Okano, F., Accession No. E15017, Jul. 28, 1999.

Okano, F., Accession No. E15018, Jul. 28, 1999.

Okano, et al., *Journal of Interferon and Cytokine Research*, 1997, vol. 17, pp. 713-718.

Okano, et al., *Journal of Interferon and Cytokine Research*, 1999, vol. 19, pp. 27-32.

Osaki, et al., *The Journal of Immunology*, 1998, vol. 160, pp. 1742-1749.

Penha-Goncalves, et al., Accession No. Y11132, Jul. 7, 1999.

Pirhonen, et al., *The Journal of Immunology*, 1999, vol. 162, pp. 7322-7329.

Rogerson, et al., *Ann N Y Acad Sci*, Oct. 31, 1996, vol. 795, pp. 354-356.

Sareneva et al., *The Journal of Immunology*, 1998, vol. 160, pp. 6032-6038.

Schijns, V.E.C.J., Accession No. Y07761, May 13, 1997.

Schijns, et al., Accession No. Y07762, May 13, 1997.

Taylor, et al., Accession No. AF135967, Jul. 12, 2000.

Ushio, S., Accession No. D49950, Feb. 10, 1999.

Wardlow, et al., Accession No. AF090119, Aug. 2, 1999.

Yoshimoto, et al., *The Journal of Immunology*, 1998, vol. 161, pp. 3400-3407.

Yoshimoto, et al., *Proc. Natl. Acad. Sci. USA*, Apr. 1997, vol. 94, pp. 3948-3953.

Foss, et al., 1999, *Scand. J. Immunol.*, vol. 50, pp. 596-604.

Peng, et al., 1999, *The Journal of Immunology*, vol. 163, pp. 250-258.

*Primary Examiner*—James Martinell

(57) **ABSTRACT**

The present invention relates to canine and feline proteins. In particular, the present invention discloses feline interleukin-18, feline caspase-1, feline interleukin-12 single chain and canine interleukin-12 single chain proteins. The present invention also includes feline interleukin-18, feline caspase-1, feline interleukin-12 single chain and canine interleukin-12 single chain nucleic acid molecules encoding such proteins, antibodies raised against such proteins and/or inhibitors of such proteins or nucleic acid molecules. The present invention also includes therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies and/or inhibitors, as well as their use to evaluate and regulate an immune response in an animal.

**19 Claims, No Drawings**

1

## FELINE IL-12 SINGLE CHAIN NUCLEIC ACID MOLECULES

### CROSS REFERENCE TO RELATED APPLICATIONS

This application is a divisional of co-pending U.S. application Ser. No. 09/917,265, filed Jul. 27, 2001; now U.S. Pat. No. 6,818,444 which claims priority to U.S. Provisional Application Ser. No. 60/223,016, filed Aug. 4, 2000; each entitled "CANINE AND FELINE PROTEINS, NUCLEIC ACID MOLECULES AND USES THEREOF".

### FIELD OF THE INVENTION

The present invention relates to canine and feline proteins. In particular, the present invention relates to feline interleukin-18 (IL-18), feline caspase-1 (casp-1), feline interleukin-12 (IL-12) single chain, and canine interleukin-12 (IL-12) single chain proteins and includes nucleic acid molecules encoding such proteins, antibodies raised against such proteins and/or inhibitors of such proteins or nucleic acid molecules. The present invention also includes therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies and/or inhibitors, as well as their use to evaluate and regulate an immune response in an animal.

### BACKGROUND OF THE INVENTION

Regulation of immune and inflammatory responses in animals is important in disease management. Immune responses can be regulated by modifying the activity of immunoregulatory molecules and immune cells. Examples of such immunoregulatory molecules include IL-18, caspase-1 and IL-12. These molecules have been found to play a role in the treatment of several disorders including allergy, cancer, and pathogenic infection.

Monocytes and macrophages represent the first line of defense against disease. Various diseases and infections activate transcriptional and posttranslational events in monocytes and macrophages, which lead to the production of cytokines such as IL-18 and IL-12. These cytokines in turn activate responses in T and B cells helping to eliminate pathogens and/or disease in a animal. Both IL-18 and IL-12 augment cellular immunity by stimulating T cells to produce interferon gamma (IFN- $\gamma$ ) which inhibits the production of IgE formation without compromising B cell proliferation. IL-18, formerly referred to as interferon gamma inducing factor (IGIF), stimulates T cells to produce IFN- $\gamma$  and has been isolated from humans, dogs, and mice. A cDNA encoding human IL-18 was isolated and used to express recombinant human IL-18 by Ushio et al., 1996, *J. Immunol.* 156, 4274-4279, GenBank accession number D49950. Feline IL-18 cDNA has a 85.8% homology to human IL-18 cDNA and feline IL-18 protein has a 81.7% homology to human IL-18 protein. A cDNA encoding canine IL-18 was isolated and used to express recombinant canine IL-18 by Okano et al., 1999, *J. Interferon Cytokine Res.* 19, 27-32, GenBank accession number Y11133. Feline IL-18 cDNA has a 90.7% homology to canine IL-18 cDNA and feline IL-18 protein has a 88.5% homology to canine IL-18 protein. A cDNA encoding murine IL-18 was isolated and used to express recombinant murine IL-18 by Okamura et al., 1995, *Nature* 378, 88-91, GenBank accession number D49949. Feline IL-18 cDNA has a 73.8% homology to murine IL-18 cDNA and feline IL-18 protein has a 70%

2

homology to murine IL-18 protein. A cDNA encoding rat IL-18 was isolated by Culhane, et al. *Mol. Psych.* 3, 362-366 (1998), GenBank accession number AJ222813. Feline IL-18 cDNA has a 73.4% homology with rat IL-18 cDNA, and feline IL-18 protein has a 70.7% homology with rat IL-18 protein. A cDNA encoding equine IL-18 was isolated by Nicolson, et al. (unpublished, direct submission to GenBank, accession number Y11131). Feline IL-18 cDNA has a 92% homology to equine IL-18 cDNA and feline IL-18 protein has a 89% homology to equine IL-18 protein. A cDNA encoding pig IL-18 was isolated by Penha-Goncalves, et al. (unpublished, direct submission to GenBank, accession number Y11132. Feline IL-18 cDNA has a 90.2% homology to pig IL-18 cDNA and feline IL-18 protein has a 85.9% homology to pig IL-18 protein. Expression of active IL-18 is controlled by caspase-1 (IL-1 $\beta$  converting enzyme). That is, IL-18 lacks a signal peptide so the precursor form of IL-18 (pro IL-18) is cleaved by caspase-1 resulting in a mature protein that is biologically active.

IL-12 is a heterodimer comprised of two subunits p40 and p35 which are covalently linked by a disulfide bond to form an active molecule. Simultaneous expression of the two subunits is necessary for the production of the biologically active heterodimer. Both human and murine p35 and p40 IL-12 single chain proteins (i.e., a single protein containing both p35 and p40 subunits) have been produced; see e.g., Lieschke et al., 1997, *Nature Biotechnology* 15, 35-40. Co-expression of the human and murine p35 and p40 cDNA subunits of IL-12 resulting in a biologically active IL-12 heterodimer was achieved by Gubler et al., 1991, *Proc. Natl. Acad. Sci. U.S.A.*, 88, 4143-4147 and Schoenhaut et al., 1992, *J. Immunol.*, 148, 3433-3440, respectively. cDNAs encoding canine IL-12 p35 and p40 subunits were isolated and co-transfected to express canine IL-12 by Okano et al., 1997, *J. Interferon Cytokine Res.* 17, 713-718. cDNAs encoding feline p35 and p40 subunits have been isolated and expressed; see, for example, Fehr et al., 1997, *DNA Seq.* 8, 77-82; Schijns et al., 1997, *Immunogenetics* 45, 462-463; Bush et al., 1994, *Molec. Immunol.* 31, 1373-1374. At the amino acid level, canine and feline IL-12 p40 subunit share 92.7 percent identity to each other; share 84.8 and 84.2 percent identity to human IL-12 p40, respectively; and share 67.4 and 68.9 percent identity to murine IL-12 p40, respectively. IL-12 shares some biological activities with IL-18 including IFN- $\gamma$  production in T cells. IL-18 and IL-12 in combination work synergistically to increase IFN- $\gamma$  production in T cells; as such these cytokines when utilized alone or in combination can be very effective in mediating IgE responses.

Caspase-1 may play a key role in the processing of IL-18 precursor in cells where IL-18 is produced. It may be that coexpression of caspase-1 along with IL-18 may be necessary for the proper processing of the IL-18 precursor and enhanced secretion of the processed IL-18 mature polypeptide. A cDNA encoding equine caspase-1 was isolated by Wardlow, et al. (unpublished; direct submission to GenBank, accession number AF090119). Feline caspase-1 cDNA has a 71% homology to equine caspase-1 cDNA and feline caspase-1 protein has a 48.8% homology to equine caspase-1 protein. A cDNA encoding equine caspase-1 was isolated by Cerretti, et al. (*Science* 256, p 97-100 (1992); GenBank accession number M87507). Feline caspase-1 cDNA has a 60% homology to human caspase-1 cDNA and feline caspase-1 protein has a 60% homology to human caspase-1 protein. A cDNA encoding rat caspase-1 (called interleukin-1 beta converting enzyme) was isolated by

Keane, et al. (Cytokine 7(2) 105–110 1995); GenBank accession number U14647). Feline caspase-1 cDNA has a 55.4% homology to rat caspase-1 cDNA and feline caspase-1 protein has a 40.2% homology to rat caspase-1 protein. A cDNA encoding murine caspase-1 was isolated by Molineaux, et al. (Proc Natl. Acad. Sci 90, 1809–1813, 1993); GenBank accession number L28095). Feline caspase-1 cDNA has a 55.7% homology to murine caspase-1 cDNA and feline caspase-1 protein has a 38.5% homology to murine caspase-1 protein. A cDNA encoding canine caspase-1 was isolated by Taylor, et al. (2000) *DNA Seq.* 10(6), pp 387–394; GenBank accession number AF135967). Feline caspase-1 cDNA has a 90% homology to canine caspase-1 cDNA.

To date, however, neither IL-18 nor caspase-1, nor the nucleic acid molecules encoding such proteins, have been isolated from cats. Neither have IL-12 single chain proteins been produced using feline or canine IL-12 subunits. As such there remains a need for compounds and methods to regulate an immune response in cats and dogs through manipulation of IL-18, caspase-1 and IL-12 single chain activities.

#### SUMMARY OF THE INVENTION

The present invention relates to canine and feline proteins, nucleic acid molecules encoding such proteins, antibodies raised against such proteins and/or inhibitors of such proteins or nucleic acid molecules. In a preferred embodiment, the present invention relates to feline interleukin-18 (IL-18), feline caspase-1 (casp-1), feline interleukin-12 (IL-12) single chain and canine interleukin-12 single chain proteins, nucleic acid molecules, antibodies and inhibitors. The present invention also includes therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies and/or inhibitors, as well as their use to evaluate and regulate an immune response in an animal.

One embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (a) an isolated nucleic acid molecule selected from the group consisting of (i) a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13; and (ii) a nucleic acid molecule comprising at least 70 contiguous nucleotides identical in sequence to at least 70 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, and SEQ ID NO:41; (b) an isolated nucleic acid molecule selected from the group consisting of (i) a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:25, and (ii) a nucleic acid molecule comprising at least 70 contiguous nucleotides identical in sequence to at least 70 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:25; (c) an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule comprising ((a)) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:26, SEQ ID NO:29, and a nucleic acid

sequence comprising at least 44 contiguous nucleotides identical in sequence to at least 44 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:26 and SEQ ID NO:29; ((b)) a nucleic acid linker of  $(XXX)_n$ , wherein  $n=0$  to 60; and ((c)) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:32, SEQ ID NO:35, and a nucleic acid molecule comprising at least 44 contiguous nucleotides identical in sequence to at least 44 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:32 and SEQ ID NO:35, such that said nucleic acid molecule of (i) encodes a feline IL-12 single chain protein; and (ii) a nucleic acid molecule comprising a nucleic acid sequence fully complementary to the coding strand of any of said nucleic acid molecules as set forth in (i); (d) an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule comprising ((a)) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:52 and SEQ ID NO:58, and a nucleic acid sequence comprising at least 47 contiguous nucleotides identical in sequence to at least 47 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:46 and SEQ ID NO:49; ((b)) a nucleic acid linker of  $(XXX)_n$ , wherein  $n=0$  to 60; and ((c)) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:49, and a nucleic acid molecule comprising at least 47 contiguous nucleotides identical in sequence to at least 47 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:46 and SEQ ID NO:49, such that said nucleic acid molecule of (i) encodes a canine IL-12 single chain protein; and (ii) a nucleic acid molecule comprising a nucleic acid sequence fully complementary to the coding strand of any of said nucleic acid molecules as set forth in (i); (e) an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule having a nucleic acid sequence that is at least 92 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13; and (ii) a nucleic acid molecule comprising a fragment of a nucleic acid molecule of (i) wherein said fragment is at least 80 nucleotides in length; (f) an isolated nucleic acid molecule selected from the group consisting of (i) a nucleic acid molecule having a nucleic acid sequence that is at least 85 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23 and SEQ ID NO:25, and (ii) a nucleic acid molecule comprising a fragment of a nucleic acid molecule of (i) wherein said fragment is at least 85 nucleotides in length; (g) an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule comprising ((a)) a nucleic acid molecule comprising a nucleic acid sequence that is at least 87 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:26 and SEQ ID NO:29, or a fragment thereof of at least 55 nucleotides in length; ((b)) a nucleic acid linker of  $(XXX)_n$ , wherein  $n=0$  to 60; and ((c)) nucleic acid molecule comprising a nucleic acid sequence that is at least 87 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:32 and SEQ ID NO:35, or a fragment thereof of at least 55 nucleotides in length, such that said nucleic acid molecule (i) encodes a feline IL-12 single chain protein; and (ii) a nucleic

acid molecule comprising a nucleic acid sequence fully complementary to the coding strand of a nucleic acid molecule as set forth in (i); and (h) an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule comprising ((a)) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:52 and SEQ ID NO:58, and a nucleic acid sequence comprising at least 55 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of NO:52 and SEQ ID NO:58; ((b)) a nucleic acid linker of (XXX)<sub>n</sub>, wherein n=0 to 60; and ((c)) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:49, and a nucleic acid molecule comprising at least 55 contiguous nucleotides identical in sequence to at least 55 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:32 and SEQ ID NO:35, such that said nucleic acid molecule of (i) encodes a canine IL-12 single chain protein; and (ii) a nucleic acid molecule comprising a nucleic acid sequence fully complementary to the coding strand of any of said nucleic acid molecules as set forth in (a), (b), (c), or (d). The present invention also includes recombinant molecules, recombinant viruses and recombinant cells comprising such IL-18, caspase-1, and IL-12 single chain nucleic acid molecules and methods to produce such nucleic acid molecules, recombinant molecules, recombinant viruses and recombinant cells.

Another embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (a) a nucleic acid molecule having a nucleic acid sequence encoding an IL-18 protein selected from the group consisting of: (i) a protein selected from the group consisting of ((a)) a protein having an amino acid sequence that is at least 92 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12, and ((b)) a protein comprising a fragment of a protein of ((a)), wherein said fragment is at least 30 amino acids in length; and (ii) a protein comprising at least 25 contiguous amino acids identical in sequence to at least 25 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12; (b) a nucleic acid molecule having a nucleic acid sequence encoding a caspase-1 protein selected from the group consisting of: (i) a protein selected from the group consisting of ((a)) a protein having an amino acid sequence that is at least 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and ((b)) a protein comprising a fragment of a protein of ((a)), wherein said fragment is at least 30 amino acids in length; and (ii) a protein comprising at least 25 contiguous amino acids identical in sequence to at least 25 contiguous amino acids selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24; (c) a nucleic acid molecule having a nucleic acid sequence encoding an IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to a IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p40 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino

acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30 and wherein said p35 subunit domain is selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p35 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36; (d) a nucleic acid molecule having a nucleic acid sequence encoding an IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to a IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p40 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59, and wherein said p35 subunit domain is selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p35 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50; (e) a nucleic acid molecule comprising a nucleic acid sequence fully complementary to the coding strand of any of said nucleic acid molecules as set forth in (a), (b), (c), or (d). The present invention also includes recombinant molecules, recombinant viruses and recombinant cells comprising such IL-18, caspase-1, and IL-12 single chain nucleic acid molecules and methods to produce such nucleic acid molecules, recombinant molecules, recombinant viruses and recombinant cells.

Another embodiment of the present invention is an isolated protein selected from the group consisting of: (a) an isolated IL-18 protein selected from the group consisting of: (i) an isolated protein of at least 25 amino acids in length, wherein said protein has an at least 25 contiguous amino acid region identical in sequence to a 25 contiguous amino acid region selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12; and (ii) an isolated protein having an amino acid sequence that is at least 92 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12 and a fragment thereof of at least 30 nucleotides; wherein said isolated protein has a function selected from the group consisting of (i) eliciting an immune response against an IL-18 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12, (ii) selectively binding to an antibody raised against an IL-18 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12,

and (iii) exhibiting IL-18 activity; (b) an isolated caspase-1 protein selected from the group consisting of: (i) an isolated protein of at least about 25 amino acids in length, wherein said protein has an at least 25 contiguous amino acid region identical in sequence to a 25 contiguous amino acid region selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24; and (ii) an isolated protein having an amino acid sequence that is at least 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24 and has a nucleic acid fragment thereof of at least 30 nucleotides; wherein said isolated protein has a function selected from the group consisting of (i) eliciting an immune response against a caspase-1 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, (ii) ii) selectively binding to an antibody raised against caspase-1 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and (iii) exhibiting caspase-1 activity; (c) an isolated IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to an IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p40 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30 and wherein said p35 subunit domain is selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p35 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36; wherein said isolated protein has a function selected from the group consisting of (i) eliciting an immune response against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:39 and SEQ ID NO:44, (ii) selectively binding to an antibody raised against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67, and (iii) exhibiting IL-12 activity; and (d) an isolated IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to an IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p40 subunit protein comprising at least 31 contiguous amino acids identical in

sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59; wherein said p35 subunit domain is selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p35 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50; and wherein said isolated protein has a function selected from the group consisting of (i) eliciting an immune response against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:62, and SEQ ID NO:67, (ii) selectively binding to an antibody raised against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67, and (iii) exhibiting IL-12 activity. The present invention also includes recombinant molecules, recombinant viruses and recombinant cells comprising such IL-18, caspase-1, and IL-12 single chain nucleic acid molecules and methods to produce such nucleic acid molecules, recombinant molecules, recombinant viruses and recombinant cells.

The present invention also includes an antibody that selectively binds to a protein of the present invention as well as methods to produce and use such proteins or antibodies. By selectively is meant an antibody that binds to a protein of the present invention, but does not bind a similar protein of another species.

One aspect of the present invention is a therapeutic composition that, when administered to an animal, regulates an immune response in the animal. Such a therapeutic composition includes at least one of the following protective compounds: an IL-18, caspase-1, or IL-12 single chain protein of the present invention, a mimotope of any of the proteins, a multimeric form of any of the proteins, an isolated nucleic acid molecule of the present invention, an antibody that selectively binds any of the proteins, and/or an inhibitor of a protein activity identified by its ability to inhibit the activity of any of the proteins. Also included is a method to regulate an immune response by administering such a therapeutic composition to an animal.

The present invention also includes a method to produce a protein of the present invention; such a method includes the step of culturing a recombinant cell capable of expressing a protein being encoded by a nucleic acid molecule of the present invention.

Another embodiment of the present invention is a method to identify a compound capable of regulating an immune response in an animal, a method selected from the group consisting of: (a) contacting an isolated feline IL-18 protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has T cell stimulating activity inducing T cells to make interferon gamma, and determining if the putative inhibitory compound inhibits that activity; (b) contacting an isolated feline caspase-1 protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the

protein cleaves precursor form of IL-18 resulting in biologically active IL-18, and determining if the putative inhibitory compound inhibits that activity; and (c) contacting an isolated IL-12 single chain protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has T cell proliferation stimulating activity, and determining if the putative inhibitory compound inhibits that activity.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides for isolated feline and canine proteins, nucleic acid molecules encoding such proteins, antibodies raised against such proteins and/or inhibitors of such proteins or nucleic acid molecules. Specifically, the present invention provides for isolated feline IL-18, feline caspase-1, and feline and canine IL-12 single chain proteins and nucleic acid molecules as well as antibodies raised against such proteins, and/or inhibitors of such proteins or nucleic acid molecules. Also included in the present invention is the use of these proteins, nucleic acid molecules, antibodies, inhibitors and/or other compounds derived therefrom as therapeutic compositions to regulate the immune response of an animal as well as in other applications, such as those disclosed below.

One embodiment of the present invention is an isolated protein that includes a feline IL-18 protein, a feline caspase-1 protein, a feline IL-12 single chain protein and/or a canine IL-12 single chain protein. As used herein, a feline and/or canine protein refers to a protein. As used herein, a protein of the present invention is a protein that is isolated from a felid or a canid or is derived therefrom and can be produced by methods known in the art, such as, for example, using recombinant DNA technology or by chemical synthesis. As such, a feline or canine protein of the present invention includes natural forms as well as any variants thereof, such as a feline or canine protein that has been altered in a manner known to those skilled in the art, such as those methods disclosed herein. As used herein, a feline or canine protein does not refer to a mouse or human protein.

Similarly, a feline or canine nucleic acid molecule of the present invention includes a feline IL-18 nucleic acid molecule, a feline caspase-1 nucleic acid molecule, a feline IL-12 single chain nucleic acid molecule and/or canine IL-12 single chain nucleic acid molecule. As used herein a feline or canine nucleic acid molecule of the present invention refers to a nucleic acid molecule that includes a nucleic acid molecule that encodes a protein of the present invention and/or a complement thereof. As such, a feline IL-18 nucleic acid molecule, a feline caspase-1 nucleic acid molecule, a feline IL-12 single chain nucleic acid molecule or a canine IL-12 single chain nucleic acid molecule of the present invention is a nucleic acid molecule that encodes a feline IL-18 protein, a feline caspase-1 protein, a feline IL-12 single chain protein or a canine IL-12 single chain protein, respectively, and/or that is a complement thereof. As used herein, a feline or canine nucleic acid molecule of the present invention is a nucleic acid molecule that is isolated from a felid or canid or is derived therefrom and can be produced using methods known in the art, such as, for example, recombinant DNA technology, or by chemical synthesis. As such, a feline or canine nucleic acid molecule of the present invention includes natural forms as well as any variants thereof, such as a feline or canine nucleic acid molecule that has been altered in a manner known to those skilled in the art, such as those methods disclosed herein. As

used herein, a feline or canine nucleic acid molecule does not refer to a mouse or human nucleic acid molecule.

According to the present invention, an isolated, or biologically pure, nucleic acid molecule or protein, is a nucleic acid molecule or protein that has been removed from its natural milieu. As such, "isolated" and/or "biologically pure" do not necessarily reflect the extent to which the nucleic acid molecule or protein has been purified. "Proteins" are defined as any compounds which comprise amino acids, including peptides, polypeptides and fusion proteins. It is to be noted that the term "a" or "an" entity refers to one or more of that entity; for example, a protein refers to one or more proteins or at least one protein. As such, the terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. Furthermore, an item "selected from the group consisting of" refers to one or more of the items in that group, including combinations thereof. The term "fragment" refers to any subset of the referent nucleic acid molecule. Furthermore, the term "linked in frame" refers to nucleic acid fragment joined to another nucleic acid fragment in a manner such that the molecule is able to be expressed when transformed into a host cell.

As used herein, a felid refers to any member of the felid family (i.e. the family Felidae), including, but not limited to, domestic cats, and wild cats such as tigers, lions, and lynx. Similarly, the term feline refers to "of the family Felidae".

As used herein, a canid refers to any member of the canid family (i.e. the family Canidae), including, but not limited to, domestic dogs, and wild canids such as wolves, foxes, and coyotes. Similarly, the term canine refers to "of the family Canidae".

Nucleic acid molecules of the present invention of known length isolated from *Felis catus* are denoted as follows: a feline IL-18 nucleic acid molecule is denoted as nFeIL-18<sub>x</sub>, wherein "x" refers to the number of nucleotides in that molecule; for example, nFeIL-18<sub>607</sub> refers to a feline IL-18 nucleic acid molecule of 607 nucleotides in length; and in a similar fashion, a feline Casp-1 nucleic acid molecule of length "x" is referred to as nFeCasp-1<sub>x</sub>, a feline IL-12 single chain nucleic acid molecule of length "x" is referred to as nFeIL-12<sub>x</sub>, a feline IL-12p35 subunit nucleic acid molecule of length "x" is referred to as nFeIL-12p35<sub>x</sub>, and a feline IL-12p40 subunit nucleic acid molecule of length "x" is referred to as nFeIL-12p40<sub>x</sub>. Similarly, *Felis catus* IL-18, Casp-1, IL-12 single chain, IL-12p35 subunit, and IL-12p40 subunit proteins of the present invention of known length are denoted PFeIL-18<sub>x</sub>, PFeCasp-1<sub>x</sub>, PFeIL-12<sub>x</sub>, PFeIL-12p35<sub>x</sub>, and PFeIL-12p40<sub>x</sub> respectively.

Nucleic acid molecules of the present invention of known length isolated from *Canis familiaris* are denoted as follows: a canine IL-12 single chain is denoted as nCaIL-12<sub>x</sub>, wherein "x" refers to the number of nucleotides in that molecule; for example, nCaIL-12<sub>1602</sub> refers to a canine IL-12 single chain nucleic acid molecule of 1602 nucleotides in length and in a similar fashion, a canine IL-12 single chain nucleic acid molecule of length "x" is referred to as nCaIL-12p35<sub>x</sub> and a canine IL-12p40 subunit nucleic acid molecule of length "x" is referred to as nCaIL-12p40<sub>x</sub>. Similarly, *Canis familiaris* IL-12 single chain proteins of the present invention of known length isolated from are denoted PCaIL-12<sub>x</sub>, PCaIL-12p35<sub>x</sub>, or PCaIL-12p40<sub>x</sub> respectively. The present invention includes nucleic acid molecules selected from the group consisting of nCaIL-12p35<sub>591</sub>, nCaIL-12p40<sub>2267</sub>, nCaIL-12p40<sub>1002</sub>, nCaIL-12p40<sub>987</sub>, nCaIL-12<sub>1599</sub>, and nCaIL-12<sub>1533</sub>.

The present invention includes nucleic acid molecules that include one or more of the following nucleic acid sequences: SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:43, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64 and/or SEQ ID NO:66, and/or a complements of these nucleic acid sequences, i.e. SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:10, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:40, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:60, SEQ ID NO:63, and/or SEQ ID NO:68, respectively. Complements are defined as two single strands of nucleic acid in which the nucleotide sequences are such that the strands will hybridize as a result of base pairing throughout their full length; i.e., these sequences are fully complementary. Such nucleic acid sequences are further described herein and can be easily determined by those skilled in the art. It should be noted that since nucleic acid sequencing technology is not entirely error-free, nucleic acid and protein sequences presented herein represent apparent nucleic acid and amino acid sequences of the isolated nucleic acid molecules and proteins, respectively, of the present invention.

As used herein, an isolated feline IL-18, feline caspase-1, feline IL-12p40 subunit, feline IL-12p35 subunit, feline IL-12 single chain, canine IL-12p40 subunit, canine IL-12p35, and/or canine IL-12 single chain protein of the present invention can be a full-length protein or any homolog of such a protein, including truncated forms of the protein. An isolated IL-18 protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response against, (or to) an IL-18 protein, whether the protein has IL-18 activity, such as T cell stimulating activity, or selectively binding to an antibody raised against an IL-18 protein. An isolated caspase-1 protein of the present invention may be identified in a straight-forward manner by the protein's ability to elicit an immune response against, (or to) a caspase-1 protein, whether the protein has caspase-1 activity, such as cleaving the precursor form of IL-18 resulting in a biologically active IL-18, or selectively binding to an antibody raised against a caspase-1 protein. A IL-12 single chain protein of the present invention may be identified in a straight-forward manner by the protein's ability to elicit an immune response against, (or to) an IL-12 protein, including the p35 or p40 subunits, whether the protein has IL-12 activity, such as T cell stimulating activity, or selectively binding to an antibody raised against an IL-12 protein, including the p35 or p40 subunits. Examples of protein homologs of the present invention includes proteins of the present invention in which amino acids have been deleted (e.g. a truncated version of the protein, such as a peptide), inserted, inverted, substituted and/or derivatized (e.g. by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation and/or addition of glycerophosphatidyl inositol) such that the protein homolog include at least one epitope capable of eliciting an immune response against the parent protein, where the term parent refers to the longer and/or full-length protein that the homolog is derived from. The ability of a protein to effect an immune response can be measured using techniques known to those skilled in the art. As used herein, the term "epitope" refers to the smallest portion of a protein capable of selec-

tively binding to the antigen binding site of an antibody. It is well accepted by those skilled in the art that the minimal size of a protein epitope capable of selectively binding to the antigen binding site of an antibody is about five or six to seven amino acids.

Proteins of the present invention include variants of a full-length protein of the present invention. Such variants include proteins that are less than full-length. As used herein, variants of the present invention refer to nucleic acid molecules that are naturally-occurring a defined below, and may result from alternative RNA splicing, alternative termination of an amino acid sequence or DNA recombination. Examples of variants include allelic variants as defined below. It is to be noted that a variant is an example of a homolog of the present invention.

Proteins of the present invention are encoded by nucleic acid molecules of the present invention. As used herein, an IL-18 nucleic acid molecule includes sequences related to a natural feline IL-18 gene. As used herein, a caspase-1 protein includes nucleic acid sequences related to a natural feline caspase-1 gene. As used herein, an IL-12 single chain nucleic acid molecule includes sequences related to a natural canine or feline IL-12 gene, IL-12p35 gene, and/or IL-12p40 gene. As used herein, a feline IL-18, a feline caspase-1, and or a feline or canine IL-12 single chain refers to the natural genomic elements that encode a feline IL-18, a feline caspase-1, and or a feline or canine IL-12 single chain, and includes all regions such as regulatory regions that control production of the protein encoded by the gene) such as, but not limited to, transcription, translation or post-translation control regions) as well as the coding region itself, and any introns or non-translated coding regions. As used herein, a gene that "includes" or "comprises" a sequence may include that sequence in one continuous array, or may include the sequence of fragmented exons. As used herein, the term "coding region" refers to a continuous linear array of nucleotides that translates into a protein. A full-length coding region is that region that is translated into a full-length, i.e., a complete protein as would be initially translated in its natural milieu, prior to any post-translational modifications.

In one embodiment of the present invention, isolated proteins are encoded by nucleic acid molecules that hybridize under stringent hybridization conditions to the non-coding strand of nucleic acid molecules encoding proteins. The minimal size of a protein of the present invention (4-6 amino acids) is a size sufficient to be encoded by a nucleic acid molecule capable of forming a stable hybrid, i.e., hybridizing under stringent hybridization conditions, with the complementary sequence of a nucleic acid molecule encoding the corresponding natural protein. The size of a nucleic acid molecule encoding such a protein is dependent on the nucleic acid composition and the percent homology between the nucleic acid molecule and the complementary nucleic acid sequence. It can easily be understood that the extent of homology required to form a stable hybrid under stringent conditions can vary depending on whether the homologous sequences are interspersed throughout a given nucleic acid molecule or are clustered, i.e. localized, in distinct regions on a given nucleic acid molecule.

The minimal size of a feline IL-18, feline caspase-1, and/or canine or feline IL-12 single chain protein homolog/portion/fragment of the present invention is a size sufficient to be encoded by a nucleic acid molecule capable of forming a stable hybrid (i.e. hybridize under stringent hybridization conditions) with the complementary sequence of a nucleic acid molecule encoding the corresponding natural protein. Stringent hybridization conditions are determined based on

defined physical properties of the gene to which the nucleic acid molecule is being hybridized, and can be defined mathematically. Stringent hybridization conditions are those experimental parameters that allow an individual skilled in the art to identify significant similarities between heterologous nucleic acid molecules. These conditions are well known to those skilled in the art. See, for example, Sambrook, et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press, and Meinkoth, et al., 1984, *Anal. Biochem.* 138, 267-284, each of which is incorporated herein by this reference. As explained in detail in the cited references, the determination of hybridization conditions involves the manipulation of a set of variables including the ionic strength ( $M$ , in moles/liter), the hybridization temperature ( $^{\circ}C$ ), the concentration of nucleic acid helix destabilizing agents, such as formamide, the average length of the shortest hybrid duplex ( $n$ ), and the percent G+C composition of the fragment to which an unknown nucleic acid molecule is being hybridized. For nucleic acid molecules of at least about 150 nucleotides, these variables are inserted into a standard mathematical formula to calculate the melting temperature, or  $T_m$ , of a given nucleic acid molecule. As defined in the formula below,  $T_m$  is the temperature at which two complementary nucleic acid molecule strands will disassociate, assuming 100% complementarity between the two strands:

$$T_m = 81.5^{\circ}C + 16.6 \log M + 0.41(\% G+C) - 500/n - 0.61(\% \text{ formamide}).$$

For nucleic acid molecules smaller than about 50 nucleotides, hybrid stability is defined by the dissociation temperature ( $T_d$ ), which is defined as the temperature at which 50% of the duplexes dissociate. For these smaller molecules, the stability at a standard ionic strength is defined by the following equation:

$$T_d = 4(G+C) + 2(A+T).$$

A temperature of  $5^{\circ}C$  below  $T_d$  is used to detect hybridization between perfectly matched molecules.

Also well known to those skilled in the art is how base pair mismatch, i.e. differences between two nucleic acid molecules being compared, including non-complementarity of bases at a given location, and gaps due to insertion or deletion of one or more bases at a given location on either of the nucleic acid molecules being compared, will affect  $T_m$  or  $T_d$  for nucleic acid molecules of different sizes. For example,  $T_m$  decreases about  $1^{\circ}C$  for each 1% of mismatched base pairs for hybrids greater than about 150 bp, and  $T_d$  decreases about  $5^{\circ}C$  for each mismatched base pair for hybrids below about 50 bp. Conditions for hybrids between about 50 and about 150 base pairs can be determined empirically and without undue experimentation using standard laboratory procedures well known to those skilled in the art. These simple procedures allow one skilled in the art to set the hybridization conditions, by altering, for example, the salt concentration, the formamide concentration or the temperature, so that only nucleic acid hybrids with greater than a specified % base pair mismatch will hybridize. Stringent hybridization conditions are commonly understood by those skilled in the art to be those experimental conditions that will allow about 30% or less base pair mismatch, i.e., at least about 85% identity. Because one skilled in the art can easily determine whether a given nucleic acid molecule to be tested is less than or greater than about 50 nucleotides, and can therefore choose the appropriate formula for determining hybridization conditions, he

or she can determine whether the nucleic acid molecule will hybridize with a given gene or specified nucleic acid molecule under stringent hybridization conditions and similarly whether the nucleic acid molecule will hybridize under conditions designed to allow a desired amount of base pair mismatch.

Hybridization reactions are often carried out by attaching the nucleic acid molecule to be hybridized to a solid support such as a membrane, and then hybridizing with a labeled nucleic acid molecule, typically referred to as a probe, suspended in a hybridization solution. Examples of common hybridization reaction techniques include, but are not limited to, the well-known Southern and northern blotting procedures. Typically, the actual hybridization reaction is done under non-stringent conditions, i.e., at a lower temperature and/or a higher salt concentration, and then high stringency is achieved by washing the membrane in a solution with a higher temperature and/or lower salt concentration in order to achieve the desired stringency.

In one embodiment, an IL-18 gene of the present invention includes the nucleic acid molecule SEQ ID NO:1, as well as the complement of SEQ ID NO:1. Nucleic acid sequence SEQ ID NO:1 represents the deduced sequence of the coding strand of a cDNA (complementary DNA) denoted herein as nucleic acid molecule nFeIL-18-N<sub>514</sub>, the production of which is disclosed in the Examples. SEQ ID NO:1 comprises an apparent partial coding region of nFeIL-18, coding for the N-terminal portion of feline IL-18 protein. The complement of SEQ ID NO:1 (represented herein by SEQ ID NO:3) refers to the nucleic acid sequence of the strand fully complementary to the strand having SEQ ID NO:1, which can easily be determined by those skilled in the art. Likewise, a nucleic acid sequence complement of any nucleic acid sequence of the present invention refers to the nucleic acid sequence complement of any nucleic acid sequence of the present invention that is fully complementary (i.e. can form a double helix with) the strand for which the sequence is cited. It should be noted that since nucleic acid sequencing technology is not entirely error-free, SEQ ID NO:1 (as well as other nucleic acid and protein sequences presented herein) represents an apparent nucleic acid sequence of the nucleic acid molecule encoding an immunoregulatory protein of the present invention.

Another IL-18 gene of the present invention includes the nucleic acid molecule SEQ ID NO:4, as well as the complement represented by SEQ ID NO:6. Nucleic acid sequence SEQ ID NO:4 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-18-C<sub>502</sub>, the production of which is disclosed in the examples. Nucleic acid nFeIL-18-C<sub>502</sub> represents an apparent partial coding region of FeIL-18, encoding a partial C-terminal region of the feline IL-18 protein. Another IL-18 gene of the present invention includes the nucleic acid molecule SEQ ID NO:7, as well as the complement represented by SEQ ID NO:10. Nucleic acid sequence SEQ ID NO:7 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-18<sub>607</sub>, the production of which is disclosed in the examples. Nucleic acid nFeIL-18<sub>607</sub> represents an apparent full-length coding region of the feline IL-18 protein. Another IL-18 gene of the present invention includes the nucleic acid molecule SEQ ID NO:9, as well as the complement represented by SEQ ID NO:41. Nucleic acid sequence SEQ ID NO:9 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFe IL-18<sub>576</sub>, the production of which is disclosed in the examples. Nucleic acid molecule nFe IL-18<sub>576</sub> rep-

resents the coding region for an apparent precursor protein to a mature feline IL-18 protein. Another IL-18 gene of the present invention includes the nucleic acid molecule SEQ ID NO:11, as well as the complement represented by SEQ ID NO:13. Nucleic acid sequence SEQ ID NO:11 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFe IL-18<sub>471</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nFe IL-18<sub>471</sub> represents the coding region for an apparent mature IL-18 protein. The putative cleavage site for the mature IL-18 protein is between amino acid positions 35 and 36 of SEQ ID NO:8, representing PFeIL-18<sub>192</sub>, which is the predicted amino acid sequence of the full-length IL-18 protein (i.e., containing signal, or leader, peptide). SEQ ID NO:12 represents the predicted amino acid sequence of the mature IL-18 protein (i.e., without the signal, or leader, sequence), also denoted as PFeIL-18<sub>157</sub>.

In another embodiment, a caspase-1 gene of the present invention includes the nucleic acid sequence SEQ ID NO:14, as well as the complement represented by SEQ ID NO:16. Nucleic acid sequence SEQ ID NO:14 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeCasp-1<sub>1233</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nFeCasp-1<sub>1233</sub> represents the coding region for an apparent full-length feline caspase-1 protein and includes a human primer sequence. Another caspase-1 protein of the present invention includes the nucleic acid sequence SEQ ID NO:17, as well as the complement represented by SEQ ID NO:19. Nucleic acid sequence SEQ ID NO:17 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeCasp-1-N<sub>526</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nFeCasp-1-N<sub>526</sub> represents the coding region for the apparent N-terminal region of the feline caspase-1 protein. Another caspase-1 protein of the present invention includes the nucleic acid molecule SEQ ID NO:20, as well as the complement represented by SEQ ID NO:22. Nucleic acid sequence SEQ ID NO:20 represents the deduced sequence of a coding strand of a cDNA denoted herein as nucleic acid molecule nFeCasp-1-C<sub>500</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nFeCasp-1-C<sub>500</sub> represents the coding region for the apparent C-terminal region of the feline caspase-1 protein. Another caspase-1 protein of the present invention includes the nucleic acid molecule SEQ ID NO:23, as well as the complement represented by SEQ ID NO:25. Nucleic acid sequence SEQ ID NO:23 represents the deduced sequence of a coding strand of a cDNA denoted herein as nucleic acid molecule nFeCasp-1<sub>1230</sub>, the production of which is disclosed in the examples. Nucleic acid molecule nFeCasp-1<sub>1230</sub> represents the coding region for the apparent full-length feline caspase-1 protein, denoted herein as PFeCasp-1<sub>410</sub>, represented herein as SEQ ID NO:24.

In another embodiment, feline IL-12 single chain proteins of the present invention contain both a mature IL-12 p35 subunit and a full-length IL-12 p40 subunit, joined by a linker. An IL-12 single chain gene of the present invention includes the nucleic acid sequence SEQ ID NO:38, as well as the complement represented by SEQ ID NO:40. Nucleic acid sequence SEQ ID NO:38 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12<sub>1599</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nFeIL-12<sub>1599</sub> represents the coding region encoding a single chain full-length feline IL-12 protein, which includes the coding region for a full-length (i.e. containing signal, or

leader, sequence) IL-12 p40 subunit, a linker of the present invention, and the coding region for a mature (i.e. not containing signal, or leader, sequence) IL-12 p35 subunit. SEQ ID NO:38 comprises a sequence that includes both the nucleic acid sequence SEQ ID NO:29 (nucleic acid sequence SEQ ID NO:29 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12p40<sub>987</sub>, which represents the coding region encoding the full-length feline IL-12 p40 subunit, whereas SEQ ID NO:31 represents the complement of SEQ ID NO:29) and SEQ ID NO:35 (nucleic acid sequence SEQ ID NO:35 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12p35<sub>591</sub>, which represents the coding region encoding the mature feline IL-12 p35 subunit, whereas SEQ ID NO:37 represents the complement of SEQ ID NO:35). Translation of SEQ ID NO:38 yields a predicted protein denoted herein as PFeIL-12<sub>533</sub>, also denoted as SEQ ID NO:39.

In another embodiment, feline IL-12 single chain proteins of the present invention contain both a mature L-12 p35 subunit and an mature IL-12 p40 subunit, joined by a linker. An IL-12 single chain gene of the present invention includes the nucleic acid sequence SEQ ID NO:43, as well as the complement represented by SEQ ID NO:45. Nucleic acid sequence SEQ ID NO:43 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12<sub>1533</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nFeIL-12<sub>1533</sub> represents the coding region encoding a single chain mature feline IL-12 protein. SEQ ID NO:33 comprises a sequence that includes both the nucleic acid sequence SEQ ID NO:26 (nucleic acid sequence SEQ ID NO:26 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12p40<sub>985</sub>, which represents the coding region encoding the mature feline IL-12 p40 subunit, whereas SEQ ID NO:28 represents the complement of SEQ ID NO:26) and SEQ ID NO:35 (nucleic acid sequence SEQ ID NO:35 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12p35<sub>591</sub>, which represents the coding region encoding the mature feline IL-12 p35 subunit, whereas SEQ ID NO:37 represents the complement of SEQ ID NO:35). Translation of SEQ ID NO:43 yields a predicted protein denoted herein as PFeIL-12<sub>511</sub>, also denoted as SEQ ID NO:44.

In another embodiment, canine IL-12 single chain proteins and nucleic acid molecules of the present invention contain both a mature IL-12 p35 subunit and a full-length IL-12 p40 subunit, joined by a linker. An IL-12 single chain gene of the present invention includes the nucleic acid sequence SEQ ID NO:61, as well as the complement represented by SEQ ID NO:63. Nucleic acid sequence SEQ ID NO:61 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaIL-12<sub>1599</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nCaIL-12<sub>1599</sub> represents the coding region encoding a single chain full-length canine IL-12 protein, which includes the coding region for a full-length (i.e. containing signal, or leader, sequence), IL-14 p40 subunit, a linker of the present invention, and the coding region for a mature, (i.e., not containing signal, or leader, sequence) IL-12 p35 subunit. SEQ ID NO:61 comprises a nucleic acid sequence that includes both the nucleic acid sequence SEQ ID NO:58 (nucleic acid SEQ ID NO:58 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaIL-12<sub>987</sub>,

which represents the coding region encoding the full-length canine IL-12 p40 subunit, whereas SEQ ID NO:60 represents the complement of SEQ ID NO:58) and SEQ ID NO:49 (nucleic acid sequence SEQ ID NO:49 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12p35<sub>591</sub>, which represents the coding region encoding the mature canine IL-12 subunit, whereas SEQ ID NO:51 represents the complement of SEQ ID NO:49. Translation of SEQ ID NO:61 yields a predicted protein denoted herein as PCaIL-12<sub>533</sub>, also denoted as SEQ ID NO:62.

In another embodiment, canine IL-12 single chain proteins and nucleic acid molecules of the present invention contain both a mature IL-12 p35 subunit and a mature IL-12 p40 subunit, joined by a linker. An IL-12 single chain gene of the present invention includes the nucleic acid sequence SEQ ID NO:66, as well as the complement represented by SEQ ID NO:68. Nucleic acid sequence SEQ ID NO:66 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaIL-12<sub>1533</sub>, the production of which is disclosed in the Examples. Nucleic acid molecule nCaIL-12<sub>1533</sub> represents the coding region encoding a single chain full-length canine IL-12 protein, which includes the coding region for a full-length (i.e. containing signal, or leader, sequence), IL-14 p40 subunit, a linker of the present invention, and the coding region for a mature, (i.e., not containing signal, or leader, sequence) IL-12 p35 subunit. SEQ ID NO:66 comprises a nucleic acid sequence that includes both the nucleic acid sequence SEQ ID NO:52 (nucleic acid SEQ ID NO:52 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaIL-12<sub>921</sub>, which represents the coding region encoding the mature canine IL-12 p40 subunit, whereas SEQ ID NO:54 represents the complement of SEQ ID NO:52) and SEQ ID NO:49 (nucleic acid sequence SEQ ID NO:49 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIL-12p35<sub>591</sub>, which represents the coding region encoding the mature canine IL-12 subunit, whereas SEQ ID NO:51 represents the complement of SEQ ID NO:49. Translation of SEQ ID NO:66 yields a predicted protein denoted herein as PCaIL-12<sub>511</sub>, also denoted as SEQ ID NO:67.

Nucleic acid molecules and proteins of the present invention having specific sequence identifiers are described in Table 1.

TABLE 1

Sequence identification numbers (SEQ ID NOs) and their corresponding nucleic acid molecule or proteins.	
SEQ ID NO.	description
1	nFeIL-18-N <sub>514</sub> coding strand
2	PFeIL-18-N <sub>133</sub>
3	nFeIL-18-N <sub>514</sub> complementary strand
4	nFeIL-18-C <sub>502</sub> coding strand
5	PFeIL-18-C <sub>154</sub>
6	nFeIL-18-C <sub>502</sub> complementary strand
7	nFeIL-18 <sub>607</sub> coding strand
8	PFeIL-18 <sub>192</sub>
9	nFeIL-18 <sub>576</sub> coding strand: coding sequence for full-length feline IL-18 protein
10	nFeIL-18 <sub>607</sub> complementary strand to SEQ ID NO: 7
11	nFeIL-18 <sub>471</sub> coding strand: coding sequence for mature feline IL-18 protein
12	PFeIL-18 <sub>157</sub>
13	nFeIL-18 <sub>471</sub> complementary strand
14	nFeCasp-1 <sub>1233</sub> coding strand

TABLE 1-continued

Sequence identification numbers (SEQ ID NOs) and their corresponding nucleic acid molecule or proteins.	
SEQ ID NO.	description
15	PFeCasp-1 <sub>410</sub>
16	nFeCasp-1 <sub>1233</sub> complementary strand
17	nFeCasp-1-N <sub>526</sub> coding strand
18	PFeCasp-1-N <sub>169</sub>
19	nFeCasp-1-N <sub>526</sub> complementary strand
20	nFeCasp-1-C <sub>500</sub> coding strand
21	PFeCasp-1-C <sub>120</sub>
22	nFeCasp-1-C <sub>500</sub> complementary strand
23	nFeCasp-1 <sub>1230</sub> coding strand: coding sequence for feline caspase-1 protein
24	PFeCasp-1 <sub>410</sub>
25	nFeCasp-1 <sub>1230</sub> complementary strand
26	nFeIL-12p40 <sub>921</sub> coding strand: coding sequence for feline mature IL-12p40 subunit
27	PFeIL-12p40 <sub>307</sub>
28	nFeIL-12p40 <sub>921</sub> complementary strand
29	nFeIL-12p40 <sub>987</sub> coding strand: coding sequence for feline full-length IL-12p40 subunit
30	PFeIL-12p40 <sub>329</sub>
31	nFeIL-12p40 <sub>987</sub> complementary strand
32	nFeIL-12p35 <sub>666</sub> coding strand: coding sequence for feline full-length IL-12p35 subunit
33	PFeIL-12p35 <sub>222</sub>
34	nFeIL-12p35 <sub>666</sub> complementary strand
35	nFeIL-12p35 <sub>591</sub> coding strand: coding sequence for feline mature IL-12p35 subunit
36	PFeIL-12p35-N <sub>187</sub>
37	nFeIL-12p35 <sub>591</sub> complementary strand
38	nFeIL-12 <sub>1599</sub> coding strand
39	PFeIL-12 <sub>533</sub>
40	nFeIL-12 <sub>1599</sub> complementary strand
41	nFeIL-18 <sub>576</sub> complementary strand to SEQ ID NO: 9
42	not used - inactive
43	nFeIL-12 <sub>1533</sub> coding strand
44	PFeIL-12 <sub>511</sub>
45	nFeIL-12 <sub>1533</sub> complementary strand
46	nCaIL-12p35 <sub>666</sub> coding strand: coding strand for canine full-length IL-12p35 subunit
47	PCaIL-12p35 <sub>222</sub>
48	nCaIL-12p35 <sub>666</sub> complementary strand
49	nCaIL-12p35 <sub>591</sub> coding strand
50	PCaIL-12p35 <sub>197</sub>
51	nCaIL-12p35 <sub>591</sub> complementary strand
52	nCaIL-12p40 <sub>921</sub> coding strand: coding sequence for mature form canine IL-12p40 subunit
53	PCaIL-12p40 <sub>307</sub>
54	nCaIL-12p40 <sub>921</sub> reverse complement
55	nFeIL-12p40-N <sub>985</sub> coding sequence
56	PCaIL-12p40-N <sub>328</sub>
57	nFeIL-12p40-N <sub>985</sub> complementary strand
58	nCaIL-12p40 <sub>987</sub> coding strand: coding sequence for full-length canine IL-12p40 subunit
59	PCaIL-12p40 <sub>329</sub>
60	nCaIL-12p40 <sub>987</sub> complementary strand
61	nCaIL-12 <sub>1599</sub> coding strand
62	PCaIL-12 <sub>533</sub>
63	nCaIL-12 <sub>1599</sub> complementary strand
64	not used - inactive
65	not used - inactive
66	nCaIL-12 <sub>1533</sub> coding strand
67	PCaIL-12 <sub>511</sub>
68	nCaIL-12 <sub>1533</sub> complementary strand
101	nFeIL-12p35-N <sub>561</sub> coding strand
102	PFeIL-12p35-N <sub>157</sub>
103	nFeIL-12p35-N <sub>561</sub> complementary strand
104	nCaIL-12p35 <sub>1455</sub> coding strand
105	PCaIL-12p35 <sub>222</sub>
106	nCaIL-12p35 <sub>1455</sub> complementary strand
107	nCaIL-12p40 <sub>2267</sub> coding strand
108	PCaIL-12p40 <sub>329</sub>
109	nCaIL-12p40 <sub>2267</sub> complementary strand

Particularly preferred nucleic acid molecules encoding feline IL-18 proteins are nFeIL-18-N<sub>514</sub>, nFeIL-18-C<sub>502</sub>,

nFeIL-18<sub>607</sub>, nFeIL-18<sub>576</sub>, and nFeIL-18<sub>471</sub>, the coding strands of which are represented by SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:9 and SEQ ID NO:11, respectively.

Particularly preferred nucleic acid molecules encoding feline caspase-1 proteins are nFeCasp-1<sub>1233</sub>, nFeCasp-1<sub>N<sub>526</sub></sub>, nFeCasp-1<sub>C<sub>500</sub></sub> and nFeCasp-1<sub>1230</sub>, the coding strands of which are represented by SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, and SEQ ID NO:23 respectively.

Particularly preferred nucleic acid molecules encoding canine and feline IL-12 p35 and p40 subunit proteins are nFeIL-12p40-N<sub>985</sub>, nFeIL-12p40<sub>987</sub>, nFeIL-12p40<sub>921</sub>, nFeIL-12p35<sub>666</sub>, nFeIL-12p35-N<sub>561</sub>, nFeIL-12p35<sub>591</sub>, nCaIL-12p35<sub>666</sub>, nCaIL-12p35<sub>1455</sub>, nCaIL-12p35<sub>591</sub>, nCaIL-12p40<sub>2267</sub>, nCaIL-12p40<sub>921</sub>, and nCaIL-12p40<sub>987</sub>. Coding strands of which are represented by SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:101, SEQ ID NO:104, and SEQ ID NO:107.

Additional preferred nucleic acid molecules encoding canine and feline IL-12 single chain proteins are nFeIL-12<sub>1533</sub>, nFeIL-12<sub>1599</sub>, nCaIL-12<sub>1533</sub>, nCaIL-12<sub>1599</sub>, the coding strands of which are represented by SEQ ID NO:43, SEQ ID NO:38, SEQ ID NO:61, and SEQ ID NO:66.

One embodiment of the present invention includes an isolated nucleic acid molecule that is selected from a group of nucleic acid molecules. One member of this group includes an isolated nucleic acid molecule that is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13; and a nucleic acid molecule comprising at least 70 contiguous nucleotides identical in sequence to at least 70 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13. Another member of this group of nucleic acid molecules includes an isolated nucleic acid molecule that is selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, and SEQ ID NO:25; and a nucleic acid molecule comprising at least 70 contiguous nucleotides identical in sequence to at least 70 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, and SEQ ID NO:25. Another member of this group of nucleic acid molecules includes an isolated nucleic acid molecule that is selected from the group consisting of a nucleic acid molecule comprising (a) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:26, SEQ ID NO:29, and a nucleic acid sequence comprising at least 44 contiguous nucleotides identical in sequence to at least 44 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:26 and SEQ ID NO:9; (b) a nucleic acid linker of (XXX)<sub>n</sub> wherein n=0 to 60; and (c) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:32, SEQ ID NO:35, and a nucleic acid molecule comprising at least 44 contiguous nucleotides identical in sequence to at least 44 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:32 and SEQ ID NO:35, such that a nucleic acid molecule of this particular group

encodes a feline IL-12 protein. Another member of this group of nucleic acid molecules includes an isolate nucleic acid molecule selected from the group consisting of a nucleic acid molecule comprising (a) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:52 and SEQ ID NO:58, and a nucleic acid sequence comprising at least 47 contiguous nucleotides identical in sequence to at least 47 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:46 and SEQ ID NO:49; (b) a nucleic acid linker of (XXX)<sub>n</sub> wherein n=0 to 60; and (c) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:46, SEQ ID NO:49, and a nucleic acid molecule comprising at least 47 contiguous nucleotides identical in sequence to at least 47 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:46 and SEQ ID NO:49, such that a nucleic acid molecule of this particular group encodes a canine IL-12 single chain protein.

The phrase, a nucleic acid molecule comprising at least "x" contiguous nucleotides identical in sequence to at least "x" contiguous nucleotides of a nucleic acid molecule selected from the group consisting of SEQ ID NO:"y", refers to an "x"-nucleotide in length nucleic acid molecule that is identical in sequence to an "x"-nucleotide portion of SEQ ID NO:"y", as well as to nucleic acid molecules that are longer in length than "x". The additional length may be in the form of nucleotides that extend from either the 5' or the 3' end(s) of the contiguous identical "x"-nucleotide portion. The 5' and/or 3' extensions can include one or more extensions that have no identity to an immunoregulatory molecule of the present invention, as well as extensions that show similarity or identity to cited nucleic acids sequences or portions thereof.

Preferred portions, or lengths, for feline IL-18, feline caspase-1, feline IL-12 single chain, and canine IL-12 single chain nucleic acid molecules of the present invention include nucleic acid molecules of at least 40 nucleotides in length, at least 43 nucleotides in length, at least 44 nucleotides in length, at least 47 nucleotides in length, at least 50 nucleotides in length, at least 55 nucleotides in length, at least 60 nucleotides in length, at least 65 nucleotides in length, at least 70 nucleotides in length, at least 75 nucleotides in length, at least 80 nucleotides in length, at least 85 nucleotides in length, at least 90 nucleotides in length, at least 95 nucleotides in length, at least 100 nucleotides in length, at least 120 nucleotides in length, at least 140 nucleotides in length, at least 160 nucleotides in length, at least 180 nucleotides in length, at least 200 nucleotides in length, at least 250 nucleotides in length, at least 300 nucleotides in length, at least 350 nucleotides in length, at least 400 nucleotides in length, at least 450 nucleotides in length, at least 500 nucleotides in length, at least 600 nucleotides in length, at least 700 nucleotides in length, at least 800 nucleotides in length, at least 900 nucleotides in length, and a full-length molecule. Particularly preferred portions, or lengths, of the nucleic acid molecules of the present invention include nucleic acids of at least 43 nucleotides, 44 nucleotides, 47 nucleotides, 70 nucleotides, and a full length molecule.

One embodiment of a protein and/or nucleic acid molecule of the present invention is a fusion nucleic acid and/or protein that includes either a feline IL-18, caspase-1, feline IL-12 single chain, and canine IL-12 single chain nucleic acid molecule and/or protein of the present invention domain, each attached to one or more fusion segments.

Suitable fusion segments for use with the present invention include, but are not limited to, segments that can: link two or more nucleic acids and/or proteins of the present invention, to form multimeric forms of a nucleic acid and/or protein of the present invention; enhance a nucleic acid molecule or protein's stability; enhance the biological activity of a nucleic acid molecule and/or protein of the present invention; and/or assist in purification a molecule of the present invention (e.g., by affinity chromatography). A suitable fusion segment can be a domain of any size that has the desired function (e.g., imparts increased stability, enhanced activity, and/or simplifies purification of a protein). Fusion segments can be joined to amino and/or carboxyl termini of the IL-18-containing domain, or the caspase-1 ligand-containing domain, or the IL-12 p40-containing domain, or the IL-12 p35-containing domain, or the IL-12 single chain-containing domain, of a protein and/or nucleic acid and can be susceptible to cleavage in order to enable straight-forward recovery of the protein and/or nucleic acid molecule. Fusion proteins are preferably produced by culturing a recombinant cell transformed with a fusion nucleic acid molecule that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of a feline IL-18, feline caspase-1, feline IL-12 p35 subunit, feline IL-12 p40 subunit, feline IL-12 single chain, canine IL-12 p35 subunit, canine IL-12 p40 subunit, and/or canine IL-12 single chain-containing domain. Preferred fusion segments include a metal binding domain (e.g., a poly-histidine segment); an immunoglobulin binding domain (e.g., Protein A; Protein G; T cell; B cell; Fc receptor or complement protein antibody-binding domains); a sugar binding domain (e.g., a maltose binding domain); and/or a "tag" domain (e.g., at least a portion of  $\beta$ -galactosidase, a strep tag peptide, a T7 tag peptide, a Flag<sup>TM</sup> peptide, or other domains that can be purified using compounds that bind to the domain, such as monoclonal antibodies). More preferred fusion segments include metal binding domains, such as a poly-histidine segment; a maltose binding domain; a strep tag peptide, such as that available from Biometra in Tampa, Fla.; and an S10 peptide.

The phrase, a nucleic acid linker, is a term known to those skilled in the art, and refers to a nucleic acid linker that can link, or attach, nucleic acid molecules, in such a manner that expression of the nucleic acid molecules produces one fusion protein as expression product. A linker can be any nucleotide sequence that directs expression of a single fusion polypeptide from a nucleotide molecule which includes two or more nucleic acid molecules of the present invention, wherein the fusion polypeptide has appropriate biological activity. Preferably, a nucleic acid linker of the present invention comprises nucleotides arranged in codons, (i.e., 3 nucleotides that, when transcribed, code for an amino acid residue), and the linker does not contain any stop codons in frame. A linker is represented herein as (XXX)<sub>n</sub>, where X is the designation of a variable nucleotide and n refers to the number of codons. The length of the nucleic acid linker may be of any length that permits expression of the fusion protein. More preferably, the length of the nucleic acid linker is from about 0 codons to about 60 codons, or from about 0 nucleotides to about 180 nucleotides. A particularly preferred linker includes SEQ ID NO:83. Appropriate biological activity includes the ability of such a fusion protein to elicit an immune response against a protein of the present invention, selectively binding an antibody raised against a protein of the present invention, and exhibiting the immunoregulatory activity of a protein of the present invention.

A single chain IL-12 protein of the present invention includes single chain IL-12 proteins comprising an IL-12 p35 subunit of the present invention at the N-terminus of the single chain protein and an IL-12 p40 subunit of the present invention at the C-terminus of the single chain protein, with the linker between the p35 subunit and the p40 subunit. Preferred single chain IL-12 proteins comprise an IL-12 p40 of the present invention at the N-terminus of the single chain protein and an IL-12 p40 subunit of the present invention at the C-terminus of the single chain protein, with the linker in between the subunits.

Another embodiment of the present invention includes an isolated nucleic acid molecule that is selected from the group consisting of: (i) a nucleic acid molecule having a nucleic acid sequence that is at least 92 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13; and (ii) a nucleic acid molecule comprising a fragment of a nucleic acid molecule of (i) wherein said fragment is at least 80 nucleotides in length. Preferred nucleic acid molecules include nucleic acid sequences that are at least 92%, at least 93%, at least 94%, more preferably at least 95% identical, and even more preferably at least 98% identical to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13. Preferred fragment lengths include fragments of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, and SEQ ID NO:13 which are at least 75 nucleotides in length, which are at least 80 nucleotides in length, which are at least 85 nucleotides in length, which are at least 90 nucleotides in length, which are at least 100 nucleotides in length, which are at least 120 nucleotides in length, which are at least 150 nucleotides in length, which are at least 200 nucleotides in length, which are at least 300 nucleotides in length, which are at least 400 nucleotides in length, which are at least 500 nucleotides in length, which are at least 600 nucleotides in length, and which preferably are full-length.

Another embodiment of the present invention includes an isolated nucleic acid molecule that is selected from the group consisting of: (i) a nucleic acid molecule having a nucleic acid sequence that is at least 85 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, and SEQ ID NO:25; and (ii) a nucleic acid molecule comprising a fragment of a nucleic acid molecule of (i) wherein said fragment is at least 85 nucleotides in length. Preferred nucleic acid molecules include nucleic acid sequences that are at least 85%, preferably at least 87%, more preferably at least 90%, even more preferably at least 95% identical to SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, and SEQ ID NO:25. Preferred fragment lengths include fragments of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, and SEQ ID NO:25 which are at least 70 nucleotides in length, which are at least 80 nucleotides in length, which are at least 85 nucleotides in length, which are at least 90 nucleotides in length, which are at least 100 nucleotides in length, which are at least 200 nucleotides in length, which are at least 300 nucleotides in length, which are at least 400 nucleotides in length, which

are at least 500 nucleotides in length, which are at least 600 nucleotides in length, or which preferably are full-length.

Another embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule comprising (a) a nucleic acid molecule comprising a nucleic acid sequence that is at least 87 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:26 and SEQ ID NO:29, or a fragment thereof of at least 55 nucleotides in length; (b) a nucleic acid linker of (XXX)<sub>n</sub>, wherein n=0 to 60, and (c) a nucleic acid molecule comprising a nucleic acid sequence that is at least 87 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:32 and SEQ ID NO:35, or a fragment thereof of at least 55 nucleotides in length, such that said nucleic acid molecule of (i) encodes a feline IL-12 single chain protein; and a nucleic acid molecule fully complementary to the coding strand of a nucleic acid molecule as set forth in (i). Preferred nucleic acid molecules include nucleic acid sequences that are at least 87%, at least 88%, at least 89%, more preferably at least 90%, even more preferably at least 95% identical to SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, and SEQ ID NO:35. Preferred fragment lengths include fragments of SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, and SEQ ID NO:35, which are at least 55 nucleotides in length, which are at least 60 nucleotides in length, which are at least about 65 nucleotides in length, which are at least 70 nucleotides in length, which are at least 80 nucleotides in length, which are at least 90 nucleotides in length, which are at least 100 nucleotides in length, which are at least 200 nucleotides in length, which are at least 300 nucleotides in length, which are at least 400 nucleotides in length, which are at least 500 nucleotides in length, which are at least 600 nucleotides in length, or which preferably are full-length.

Another embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (i) a nucleic acid molecule comprising (a) a nucleic acid molecule comprising a nucleic acid sequence that is at least 87 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:52 and SEQ ID NO:58, or a fragment thereof of at least 55 nucleotides in length; (b) a nucleic acid linker of (XXX)<sub>n</sub>, wherein n=0 to 60; and (c) a nucleic acid molecule comprising a nucleic acid sequence that is at least 87 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:46 and SEQ ID NO:49, or a fragment thereof of at least 55 nucleotides in length, such that said nucleic acid molecule of (i) encodes a canine IL-12 single chain protein; and a nucleic acid molecule fully complementary to the coding strand of a nucleic acid molecule as set forth in (i). Preferred nucleic acid molecules include nucleic acid sequences that are at least 87%, at least 88%, at least 89%, more preferably at least 90%, even more preferably at least 95% identical to SEQ ID NO:52, SEQ ID NO:58, SEQ ID NO:46, and SEQ ID NO:49. Preferred fragment lengths include fragments of SEQ ID NO:52, SEQ ID NO:58, SEQ ID NO:46, and SEQ ID NO:49, which are at least 55 nucleotides in length, which are at least 60 nucleotides in length, which are at least about 65 nucleotides in length, which are at least 70 nucleotides in length, which are at least 80 nucleotides in length, which are at least 90 nucleotides in length, which are at least 100 nucleotides in length, which are at least 200 nucleotides in length, which are at least 300 nucleotides in length, which are at least 400

nucleotides in length, which are at least 500 nucleotides in length, which are at least 600 nucleotides in length, or which preferably are full-length.

Preferred portions, or fragments, of a feline IL-18, feline caspase-1, canine or feline IL-12 single chain protein of the present invention include at least 15 amino acids, at least 20 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, at least 40 amino acids, at least 45 amino acids, at least 50 amino acids, at least 60 amino acids, at least 75 amino acids or at least 100 amino acids. An IL-18 or IL-12 single chain protein of the present invention can include at least a portion of an IL-18 or IL-12 single chain protein that is capable of binding to an IL-18 or IL-12 receptor, respectively. These receptors are known to those of skill in the art, and are described in Janeway et al., in *Immunobiology, the Immune System in Health and Disease*, Garland Publishing, Inc., NY, 1996 (which is incorporated herein by this reference in its entirety). The IL-18 or IL-12 receptor-binding portion of an IL-18 or IL-12 protein, respectively, can be determined by incubating the protein with an isolated IL-18 or IL-12 receptor, as appropriate, or a cell having an IL-18 or IL-12 receptor on its surface, as appropriate. IL-18 or IL-12 protein binding to purified IL-18 or IL-12 receptor, respectively, can be determined using methods known in the art including Biacore® screening, confocal immunofluorescent microscopy, immunoprecipitation, gel chromatography, determination of inhibition of binding of antibodies that bind specifically to the IL-18 or IL-12 binding domain of an IL-18 or IL-12 receptor, ELISA using an IL-18 or IL-12 receptor, respectively, labeled with a detectable tag such as an enzyme or chemiluminescent tag or yeast-2 hybrid technology. A caspase-1 protein of the present invention can include at least a portion of a caspase-1 protein that is capable cleaving pro-IL-18 to mature IL-18. The ability of the caspase-1 protein to cleave IL-18 can be determined by methods known in the art, including methods such as Biacore® screening, confocal immunofluorescent microscopy, immunoprecipitation, gel chromatography, determination of inhibition of cleavage upon binding of antibodies that bind specifically to either IL-18 or caspase-1, and enzymatic assays.

The present invention also includes mimetopes of feline IL-18, feline caspase-1, and canine and/or feline IL-12 single chain proteins of the present invention. As used herein, a mimetope of an immunoregulatory protein of the present invention refers to any compound that is able to mimic the activity of such a feline IL-18, feline caspase-1, and canine and/or feline IL-12 single chain protein, respectively, often because the mimetope has a structure that mimics the particular protein. Mimetopes can be, but are not limited to: peptides that have been modified to decrease their susceptibility to degradation such as all-D retro peptides; anti-idiotypic and/or catalytic antibodies, or fragments thereof; non-proteinaceous immunogenic portions of an isolated protein (e.g., carbohydrate structures); and/or synthetic or natural organic molecules, including nucleic acids. Such mimetopes can be designed using computer-generated structures of proteins of the present invention. Mimetopes can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides or other organic molecules, and screening such samples by affinity chromatography techniques using the corresponding binding partner.

Furthermore, it is known in the art that there are commercially available computer programs for determining the degree of similarity between two nucleic acid or protein sequences. These computer programs include various

known methods to determine the percentage identity and the number and length of gaps between hybrid nucleic acid molecules or proteins. Preferred methods to determine the percent identity among amino acid sequences and also among nucleic acid sequences include analysis using one or more of the commercially available computer programs designed to compare and analyze nucleic acid or amino acid sequences. These computer programs include, but are not limited to, the SeqLab® Wisconsin Package™ Version 10.0-UNIX sequence analysis software, available from Genetics Computer Group, Madison, Wis.; and DNAsis® sequence analysis software, version 2.0, available from Hitachi Software, San Bruno, Calif. Such software programs represent a collection of algorithms paired with a graphical user interface for using the algorithms. The DNAsis version 2.0 software and SeqLab Wisconsin Package Version 10.0-UNIX software, for example, employ a particular algorithm, the Needleman-Wunsch algorithm to perform pair-wise comparisons between two sequences to yield a percentage identity score, see Needleman, S. B. and Wunsch, C. D., 1970, *J. Mol. Biol.*, 48, 443, which is incorporated herein by reference in its entirety. Such algorithms, including the Needleman-Wunsch algorithm, are commonly used by those skilled in the nucleic acid and amino acid sequencing art to compare sequences. A preferred method to determine percent identity among amino acid sequences and also among nucleic acid sequences includes using the Needleman-Wunsch algorithm, available in the SeqLab Wisconsin Package Version 10.0-UNIX software (hereinafter "SeqLab"), using the Pairwise Comparison/Gap function with the nwsgapdna.cmp scoring matrix, the gap creation penalty and the gap extension penalties set at default values, and the gap shift limits set at maximum (hereinafter referred to as "SeqLab default parameters"). An additional preferred method to determine percent identity among amino acid sequences and also among nucleic acid sequences includes using the Higgins-Sharp algorithm, available in the DNAsis version 2.0 software (hereinafter "DNAsis"), with the gap penalty set at 5, the number of top diagonals set at 5, the fixed gap penalty set at 10, the k-tuple set at 2, the window size set at 5, and the floating gap penalty set at 10. A particularly preferred method to determine percent identity among amino acid sequences and also among nucleic acid sequences includes using the Needleman-Wunsch algorithm available in the DNAsis version 2.0 software, using the GCG default parameter function.

Another embodiment of the present invention includes a nucleic acid molecule that is selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:66, and SEQ ID NO:68, and a nucleic acid molecule comprising an allelic variant of a nucleic acid molecule comprising any of said nucleic acid sequences. An allelic variant of a feline and/or canine nucleic acid molecule of the present invention, including the particular SEQ ID NO's cited herein, is a gene that occurs at essentially the same locus (or loci) in the genome as the gene including the particular SEQ ID NO's cited herein, but which, due to natural variations caused by, for example, mutation or recombination, has a similar but not identical sequence. Also included in the term allelic variant are allelic variants of cDNAs derived from such genes. Because natural selection

typically selects against alterations that affect function, allelic variants usually encode proteins having similar activity to that of the protein encoded by the gene to which they are being compared. Allelic variants of genes or nucleic acid molecules can also comprise alterations in the 5' or 3' untranslated regions of the gene (e.g., in regulatory control regions), or can involve alternative splicing of a nascent transcript, thereby bringing alternative exons into juxtaposition. Allelic variants are well known to those skilled in the art and would be expected to be found within a given animal, since the respective genomes are diploid, and sexual reproduction will result in the reassortment of alleles. As such, a nucleic acid molecule of the present invention can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:9, SEQ ID NO:41, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:38, SEQ ID NO:40, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:61, SEQ ID NO:63, SEQ ID NO:66, and SEQ ID NO:68, and/or any other nucleic acid molecule cited herein.

In another embodiment of the present invention, a nucleic acid molecule of the invention is selected from the group consisting of (a) a nucleic acid molecule comprising a nucleic acid sequence that encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:62, and SEQ ID NO:67, and (b) a nucleic acid molecule comprising an allelic variant of a nucleic acid molecule encoding a protein having any of said amino acid sequences of (a).

Another embodiment of the present invention includes feline IL-18 nucleic acid molecules of the present invention, wherein said nucleic acid molecules encode a protein having a function selected from the group consisting of (i) eliciting an immune response against an IL-18 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12; (ii) selectively binding to an antibody raised against an IL-18 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12, and (iii) exhibiting IL-18 activity. Methods to elicit an immune response and to determine whether an antibody can selectively bind to a particular protein or antigen are known in the art, see, for example, Harlow, et al. (1988) *Antibodies, a Laboratory Manual*, Cold Spring Harbor Labs Press; Harlow, et al. is incorporated by reference herein in its entirety. Methods to determine whether an IL-18 protein has IL-18 activity are known in the art, and include determining whether IL-18 has the activity of stimulating T cells to produce interferon gamma (IFN- $\gamma$ ).

Another embodiment of the present invention includes caspase-1 nucleic acid molecules of the present invention that encode a protein having a function selected from the group consisting of (i) eliciting an immune response against a caspase-1 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, (ii) selectively binding to an antibody raised against caspase-1 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and (iii) exhibiting caspase-1

activity. Methods to elicit an immune response and to determine whether an antibody can selectively bind to a particular protein or antigen are known in the art, see, for example, Harlow, et al. (1988) *Antibodies, a Laboratory Manual*, Cold Spring Harbor Labs Press; Harlow, et al. is incorporated by reference herein in its entirety. Methods to determine whether a caspase-1 protein has caspase-1 activity are known in the art, and include, for example, determining if the caspase-1 protein has the ability to cleave the precursor form of IL-18 resulting in a biologically active IL-18.

Another embodiment of the present invention includes canine and feline IL-12 single chain nucleic acid molecules of the present invention, wherein a said nucleic acid molecule encodes a protein having a function selected from the group consisting of (i) eliciting an immune response against an IL-12 protein having an amino acid selected from the group consisting of SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:62, and SEQ ID NO:67, (ii) selectively binding to an antibody raised against an IL-12 protein having an amino acid sequence selected from the group consisting from the group of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67, and (iii) exhibiting IL-12 activity. Methods to elicit an immune response and to determine whether an antibody can selectively bind to a particular protein or antigen are known in the art, see, for example, Harlow, et al. (1988) *Antibodies, a Laboratory Manual*, Cold Spring Harbor Labs Press; Harlow, et al. is incorporated by reference herein in its entirety. Methods to determine whether an IL-12 protein has IL-12 activity are known in the art, and include determining if IL-12 has the activity of stimulating T cells to produce interferon gamma (IFN- $\gamma$ ).

A preferred nucleic acid molecule of the present invention includes a nucleic acid molecule selected from the group consisting of nFeIL-12p40-N<sub>985</sub>, nFeIL-12p40<sub>987</sub>, nFeIL-12p40<sub>921</sub>, nFeIL-12p35<sub>666</sub>, nFeIL-12p35<sub>561</sub>, nFeIL-12p35<sub>591</sub>, nCaIL-12p35<sub>666</sub>, nCaIL-12p35<sub>1455</sub>, nCaIL-12p35<sub>591</sub>, nCaIL-12p40<sub>2267</sub>, nCaIL-12p40<sub>921</sub>, nCaIL-12p40<sub>98</sub>, nFeIL-12<sub>1599</sub>, nFeIL-12<sub>1533</sub>, nCaIL-12<sub>1599</sub>, and nCaIL-12<sub>1533</sub>.

Another embodiment of the present invention includes an isolated nucleic acid molecule selected from the group consisting of: a nucleic acid molecule having a nucleic acid sequence encoding an IL-18 protein selected from the group consisting of: a protein selected from the group consisting of (a) a protein having an amino acid sequence that is at least 92 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12, and (b) a protein comprising a fragment of a protein of (a), wherein said fragment is at least 30 amino acids in length; and a protein comprising at least 25 contiguous amino acids identical in sequence to at least 25 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12. Preferred IL-18 proteins include proteins that are at least about 90 percent identical, preferably at least about 92 percent identical, preferably at least about 94 percent identical, preferably at least about 96 percent identical, and even more preferably at least about 98 percent identical to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12 or fragments thereof. Preferred fragments of IL-18 proteins include fragments of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12 that are at least about 20 amino acids in

length, at least about 30 amino acids in length, at least about 40 amino acids in length, at least about 50 amino acids in length, preferably at least about 75 amino acids in length, preferably at least about 100 amino acids in length, and more preferably are full-length. Preferred IL-18 proteins also include proteins that comprise at least 15 contiguous amino acids identical in sequence to at least 15 contiguous amino acids; at least 20 contiguous amino acids identical in sequence to at least 20 contiguous amino acids, preferably about 30 contiguous amino acids identical in sequence to at least 30 contiguous amino acids, preferably about 50 contiguous amino acids identical in sequence to at least 50 contiguous amino acids, preferably about 75 contiguous amino acids identical in sequence to at least 75 contiguous amino acids, preferably about 100 contiguous amino acids identical in sequence to at least 100 contiguous amino acids, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12.

Another embodiment of the present invention includes an isolated nucleic acid molecule selected from the group consisting of: a nucleic acid molecule having a nucleic acid sequence encoding caspase protein selected from the group consisting of: a protein selected from the group consisting of (a) a protein having an amino acid sequence that is at least 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and (b) a protein comprising a fragment of a protein of (a), wherein said fragment is at least 30 amino acids in length; and a protein comprising at least 25 contiguous amino acids identical in sequence to at least 25 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12. Preferred caspase-1 proteins include proteins that are at least about 85 percent identical, at least about 87 percent identical, preferably at least about 90 percent identical, preferably at least about 93 percent identical, more preferably at least about 95 percent identical, and even more preferably about 98 percent identical to SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24 or fragments thereof. Preferred fragments of caspase-1 proteins include fragments of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24 include fragments that are at least about 20 amino acids in length, at least about 30 amino acids in length, at least about 40 amino acids in length, at least about 50 amino acids in length, at least about 60 amino acids in length, preferably at least about 75 amino acids in length, preferably at least about 100 amino acids in length, and more preferably are full-length. Preferred caspase-1 proteins also include proteins that comprise at least 25 contiguous amino acids identical in sequence to at least 25 contiguous amino acids; at least 20 contiguous amino acids identical in sequence to at least 20 contiguous amino acids, preferably about 30 contiguous amino acids identical in sequence to at least 30 contiguous amino acids, preferably about 50 contiguous amino acids identical in sequence to at least 50 contiguous amino acids, preferably about 75 contiguous amino acids identical in sequence to at least 75 contiguous amino acids, preferably about 100 contiguous amino acids identical in sequence to at least 100 contiguous amino acids, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24.

Another embodiment of the present invention includes a nucleic acid molecule having a nucleic acid sequence encoding an IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to a IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of: (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p40 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30, and wherein said p35 domain is selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p35 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36. Preferred p40 subunit proteins and/or p35 subunit proteins include proteins that are at least about 84 percent identical, preferably at least about 87 percent identical, preferably at least about 90 percent identical, and even more preferably at least about 95 percent identical to SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, and SEQ ID NO:36 or fragments thereof. Preferred fragments of IL-12 single chain proteins include fragments of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, and SEQ ID NO:36 include fragments that are at least about 30 amino acids in length, at least about 40 amino acids in length, at least about 50 amino acids in length, at least about 60 amino acids in length, preferably at least about 75 amino acids in length, preferably at least about 100 amino acids in length, and more preferably are full-length. Preferred IL-12 single chain proteins also include proteins that comprise at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids, preferably about 30 contiguous amino acids identical in sequence to at least 30 contiguous amino acids, preferably about 50 contiguous amino acids identical in sequence to at least 50 contiguous amino acids, preferably about 75 contiguous amino acids identical in sequence to at least 75 contiguous amino acids, preferably about 100 contiguous amino acids identical in sequence to at least 100 contiguous amino acids, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, and SEQ ID NO:36.

Another embodiment of the present invention includes a nucleic acid molecule having a nucleic acid sequence encoding an IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to a IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of: (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p40 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino

acids of an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59, and wherein said p35 domain is selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p35 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50. Preferred p40 subunit proteins and/or p35 subunit proteins include proteins that are at least about 84 percent identical, preferably at least about 87 percent identical, preferably at least about 90 percent identical, and even more preferably at least about 95 percent identical to SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59 or fragments thereof. Preferred fragments of IL-12 single chain proteins include fragments of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, and SEQ ID NO:36 include fragments that are at least about 40 amino acids in length, at least about 50 amino acids in length, at least about 60 amino acids in length, at least about 70 amino acids in length, preferably at least about 80 amino acids in length, preferably at least about 100 amino acids in length, and more preferably are full-length. Preferred IL-12 single chain proteins also include proteins that comprise at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids, preferably about 35 contiguous amino acids identical in sequence to at least 35 contiguous amino acids, preferably about 50 contiguous amino acids identical in sequence to at least 50 contiguous amino acids, preferably about 75 contiguous amino acids identical in sequence to at least 75 contiguous amino acids, preferably about 100 contiguous amino acids identical in sequence to at least 100 contiguous amino acids, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59.

Another embodiment of the present invention includes a nucleic acid molecule comprising a nucleic acid sequence fully complementary to the coding strand of any of the nucleic acid molecules of the present invention. Another embodiment of the present invention includes a nucleic acid molecule that comprises a nucleic acid sequence that encodes a protein selected from the group consisting of an IL-18 protein, a caspase-1 protein, and an IL-12 single chain protein.

Another embodiment of the present invention includes a nucleic acid molecule that is selected from the group consisting of a nucleic acid molecule comprising a nucleic acid sequence encoding a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO: 8, SEQ ID NO:12, SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:62, and SEQ ID NO:67; and a nucleic acid molecule comprising an allelic variant of a nucleic acid molecule encoding a protein having any of said nucleic acid molecules set forth in this paragraph. In another embodiment, a nucleic acid molecule encoding an IL-12 single chain protein of the present invention further comprises a nucleic acid molecule encoding a linker.

The present invention also includes oligonucleotides, recombinant molecules, recombinant viruses and recombi-

nant cells comprising such nucleic acid molecules and methods to produce such nucleic acid molecules, oligonucleotides, recombinant molecules, recombinant viruses and recombinant cells.

Knowing the nucleic acid sequences of certain nucleic acid molecules of the present invention allows one skilled in the art to, for example, (a) make copies of those nucleic acid molecules, (b) obtain nucleic acid molecules including at least a portion of such nucleic acid molecules, e.g., nucleic acid molecules including full-length genes, full-length coding regions, regulatory control sequences, truncated coding regions, and (c) obtain other nucleic acid molecules. Such nucleic acid molecules can be obtained in a variety of ways including screening appropriate expression libraries with antibodies of the present invention; traditional cloning techniques using oligonucleotide probes of the present invention to screen appropriate libraries; and PCR amplification of appropriate libraries or DNA using oligonucleotide primers of the present invention. A preferred library to screen or from which to amplify nucleic acid molecules is a feline or canine mast library or a feline or canine peripheral blood mononuclear cell library. Techniques to clone and amplify genes are disclosed, for example, in Sambrook et al., *ibid*.

Oligonucleotides of the present invention can be RNA, DNA, or derivatives of either. The minimum size of such oligonucleotides is the size required for formation of a stable hybrid between an oligonucleotide and a complementary sequence on a larger nucleic acid molecule of the present invention, typically from about 12 to 15 to about 17 to 18 nucleotides depending on the GC/AT content. The present invention includes oligonucleotides that can be used as, for example, probes to identify nucleic acid molecules, primers to produce nucleic acid molecules, or therapeutic reagents to inhibit protein production or activity, e.g., as antisense-, triplex formation-, ribozyme- and/or RNA drug-based reagents. The present invention also includes the use of such oligonucleotides to protect animals from disease using one or more of such technologies. Appropriate oligonucleotide-containing therapeutic compositions can be administered to an animal using techniques known to those skilled in the art.

One embodiment of the present invention includes a recombinant vector, which includes at least one isolated nucleic acid molecule of the present invention, inserted into any vector capable of delivering the nucleic acid molecule into a host cell. Such a vector contains heterologous nucleic acid sequences, that is nucleic acid sequences that are not naturally found adjacent to nucleic acid molecules of the present invention and that preferably are derived from a species other than the species from which the nucleic acid molecule(s) are derived. The vector can be either RNA or DNA, either prokaryotic or eukaryotic, and typically is a virus or a plasmid. Recombinant vectors can be used in the cloning, sequencing, and/or otherwise manipulating of the nucleic acid molecules of the present invention.

One type of recombinant vector, referred to herein as a recombinant molecule, comprises a nucleic acid molecule of the present invention operatively linked to an expression vector. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the nucleic acid molecule is able to be expressed when transformed into a host cell. As used herein, an expression vector is a DNA or RNA vector that is capable of transforming a host cell and of effecting expression of a specified nucleic acid molecule. Preferably, the expression vector is also capable of replicating within the host cell. Expression vectors can be either prokaryotic or eukaryotic, and are typically viruses or plasmids. Expression vectors of

the present invention include any vectors that function, i.e., direct gene expression, in recombinant cells of the present invention, including in bacterial, fungal, insect, other animal, and plant cells. Preferred expression vectors of the present invention can direct gene expression in bacterial, yeast, insect and mammalian cells, and more preferably in the cell types disclosed herein.

In particular, expression vectors of the present invention contain regulatory sequences such as transcription control sequences, translation control sequences, origins of replication, and other regulatory sequences that are compatible with the recombinant cell and that control the expression of nucleic acid molecules of the present invention. In particular, recombinant molecules of the present invention include transcription control sequences. Transcription control sequences are sequences that control the initiation, elongation, and termination of transcription. Particularly important transcription control sequences are those that control transcription initiation, such as promoter, enhancer, operator and repressor sequences. Suitable transcription control sequences include any transcription control sequence that can function in at least one of the recombinant cells of the present invention. A variety of such transcription control sequences are known to those skilled in the art. Preferred transcription control sequences include those which function in bacterial, yeast, or insect and mammalian cells, such as, but not limited to, *tac*, *lac*, *trp*, *trc*, *oxy-pro*, *omp/lpp*, *rrnB*, bacteriophage lambda, such as lambda  $P_L$  and lambda  $p_R$  and fusions that include such promoters, bacteriophage T7, T7lac, bacteriophage T3, bacteriophage SP6, bacteriophage SP01, metallothionein, alpha-mating factor, *Pichia* alcohol oxidase, alphavirus subgenomic promoter, antibiotic resistance gene, baculovirus, *Heliothis zea* insect virus, vaccinia virus, herpesvirus, raccoon poxvirus, other poxvirus, adenovirus, cytomegalovirus, such as immediate early promoter, simian virus 40, retrovirus, actin, retroviral long terminal repeat, Rous sarcoma virus, heat shock, phosphate and nitrate transcription control sequences as well as other sequences capable of controlling gene expression in prokaryotic or eukaryotic cells. Additional suitable transcription control sequences include tissue-specific promoters and enhancers as well as lymphokine-inducible promoters, e.g., promoters inducible by interferons or interleukins.

A recombinant cell of the present invention includes any cell transformed with at least one of any nucleic acid molecule of the present invention. Suitable and preferred nucleic acid molecules as well as suitable and preferred recombinant molecules with which to transfer cells are disclosed herein. A recombinant cell is preferably produced by transforming a host cell with one or more recombinant molecules, each comprising one or more nucleic acid molecules of the present invention operatively linked to an expression vector containing one or more transcription control sequences, examples of which are disclosed herein.

Transformation of a nucleic acid molecule into a cell can be accomplished by any method by which a nucleic acid molecule can be inserted into the cell. Transformation techniques include, but are not limited to, transfection, electroporation, microinjection, lipofection, adsorption, and protoplast fusion. A recombinant cell may remain unicellular or may grow into a tissue, organ or a multicellular organism. It is to be noted that a cell line refers to any recombinant cell of the present invention that is not a transgenic animal. Transformed nucleic acid molecules of the present invention can remain extrachromosomal or can integrate into one or more sites within a chromosome of the transformed, i.e., recombinant, cell in such a manner that their ability to be

expressed is retained. Preferred nucleic acid molecules with which to transform a cell include nucleic acid molecules disclosed herein. Particularly preferred nucleic acid molecules with which to transform a cell include nFeIL-18-N<sub>514</sub>, nFeIL-18-C<sub>502</sub>, nFeIL-18<sub>607</sub>, nFeIL-18<sub>576</sub>, nFeIL-18<sub>471</sub>, nFeCasp-1<sub>1233</sub>, nFeCasp-1-N<sub>526</sub>, nFeCasp-1-C<sub>500</sub>, nFeCasp-1<sub>1230</sub>, nFeIL-12p40-N<sub>985</sub>, nFeIL-12p40<sub>987</sub>, nFeIL-12p40<sub>921</sub>, nFeIL-12p35<sub>666</sub>, nFeIL-12p35-N<sub>561</sub>, nFeIL-12p35<sub>591</sub>, nCaIL-12p35<sub>666</sub>, nCaIL-12p35<sub>1455</sub>, nCaIL-12p35<sub>591</sub>, nCaIL-12p40<sub>2267</sub>, nCaIL-12p40<sub>921</sub>, and nCaIL-12p40<sub>987</sub>, nCaIL-12<sub>1599</sub>, nCaIL-12<sub>533</sub>, nFeIL-12<sub>1599</sub>, and nFeIL-12<sub>1533</sub>.

Recombinant molecules of the present invention may also (a) contain secretory signals, i.e., signal segment nucleic acid sequences, to enable an expressed protein of the present invention to be secreted from the cell that produces the protein and/or (b) contain fusion sequences which lead to the expression of nucleic acid molecules of the present invention as fusion proteins. Examples of suitable signal segments include any signal segment capable of directing the secretion of a protein of the present invention. Preferred signal segments include, but are not limited to, tissue plasminogen activator (t-PA), interferon, interleukin, growth hormone, histocompatibility and viral envelope glycoprotein signal segments. Suitable fusion segments encoded by fusion segment nucleic acids are disclosed herein. In addition, a nucleic acid molecule of the present invention can be joined to a fusion segment that directs the encoded protein to the proteasome, such as a ubiquitin fusion segment. Eukaryotic recombinant molecules may also include intervening and/or untranslated sequences surrounding and/or within the nucleic acid sequences of nucleic acid molecules of the present invention.

Suitable host cells to transform include any cell that can be transformed with a nucleic acid molecule of the present invention. Host cells can be either untransformed cells or cells that are already transformed with at least one nucleic acid molecule, e.g., nucleic acid molecules encoding one or more proteins of the present invention and/or other proteins. Host cells of the present invention either can be endogenously, i.e., naturally, capable of producing proteins of the present invention or can be capable of producing such proteins after being transformed with at least one nucleic acid molecule of the present invention. Host cells of the present invention can be any cell capable of producing at least one protein of the present invention, and include bacterial, fungal, including yeast, insect, and other animal and plant cells. Preferred host cells include bacterial, mycobacterial, yeast, insect and mammalian cells. More preferred host cells include *Salmonella*, *Escherichia*, *Bacillus*, *Listeria*, *Saccharomyces*, *Pichia*, *Spodoptera*, *Mycobacteria*, *Trichoplusia*, BHK (baby hamster kidney) cells, MDCK cells (Madin-Darby canine kidney cell line), CRFK cells (Crandell feline kidney cell line), CV-1 cells (African monkey kidney cell line used, for example, to culture raccoon poxvirus), COS (e.g., COS-7) cells, and Vero cells. Particularly preferred host cells are *Escherichia coli*, including *E. coli* K-12 derivatives; *Salmonella typhi*; *Salmonella typhimurium*, including attenuated strains such as UK-1<sub>3987</sub> and SR-11<sub>4072</sub>; *Pichia*; *Spodoptera frugiperda*; *Trichoplusia ni*; BHK cells; MDCK cells; CRFK cells; CV-1 cells; COS cells; Vero cells; and non-tumorigenic mouse myoblast G8 cells (e.g., ATCC CRL 1246). Additional appropriate mammalian cell hosts include other kidney cell lines, other fibroblast cell lines, e.g., human, murine or chicken embryo

fibroblast cell lines, myeloma cell lines, Chinese hamster ovary cells, mouse NIH/3T3 cells, LMTK<sup>31</sup> cells and/or HeLa cells.

Recombinant cells of the present invention can also be co-transformed with one or more recombinant molecules including IL-18, caspase-1, IL-12 single chain nucleic acid molecules encoding one or more proteins of the present invention and one or more other nucleic acid molecules encoding other compounds.

Recombinant DNA technologies can be used to improve expression of transformed nucleic acid molecules by manipulating, for example, the number of copies of the nucleic acid molecules within a host cell, the efficiency with which those nucleic acid molecules are transcribed, the efficiency with which the resultant transcripts are translated, and the efficiency of post-translational modifications. Recombinant techniques useful for increasing the expression of nucleic acid molecules of the present invention include, but are not limited to, operatively linking nucleic acid molecules to high-copy number plasmids, integration of the nucleic acid molecules into one or more host cell chromosomes, addition of vector stability sequences to plasmids, substitutions or modifications of transcription control signals, e.g., promoters, operators, enhancers, substitutions or modifications of translational control signals, e.g., ribosome binding sites, Shine-Dalgarno sequences, modification of nucleic acid molecules of the present invention to correspond to the codon usage of the host cell, deletion of sequences that destabilize transcripts, and use of control signals that temporally separate recombinant cell growth from recombinant enzyme production during fermentation. The activity of an expressed recombinant protein of the present invention may be improved by fragmenting, modifying, or derivatizing nucleic acid molecules encoding such a protein.

Isolated feline IL-18, feline caspase-1, feline and/or canine IL-12 single chain proteins of the present invention can be produced in a variety of ways, including production and recovery of natural proteins, production and recovery of recombinant proteins, and chemical synthesis of the proteins. In one embodiment, an isolated protein of the present invention is produced by culturing a cell capable of expressing the protein under conditions effective to produce the protein, and recovering the protein. A preferred cell to culture is a recombinant cell of the present invention. Effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. An effective medium refers to any medium in which a cell is cultured to produce a protein of the present invention. Such a medium typically comprises an aqueous medium having assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present invention can be cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. Such culturing conditions are within the expertise of one of ordinary skill in the art. Examples of suitable conditions are included in the Examples section.

Depending on the vector and host system used for production, resultant proteins of the present invention may either remain within the recombinant cell; be secreted into the fermentation medium; be secreted into a space between two cellular membranes, such as the periplasmic space in *E. coli*; or be retained on the outer surface of a cell or viral membrane. The phrase recovering the protein, as well as

similar phrases, refers to collecting the whole fermentation medium containing the protein and need not imply additional steps of separation or purification. Proteins of the present invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization. Proteins of the present invention are preferably retrieved in substantially pure form. As used herein, substantially pure refers to a purity that allows for the effective use of the protein as a therapeutic composition or diagnostic. A therapeutic composition for animals, for example, should exhibit no substantial toxicity.

The present invention also includes isolated, i.e., removed from their natural milieu, antibodies that selectively bind to proteins of the present invention or a mimotope thereof, e.g., anti-feline IL-18, feline caspase-1, feline and canine IL-12 single antibodies. As used herein, the term selectively binds to a protein refers to the ability of antibodies of the present invention to preferentially bind to specified proteins and mimitopes thereof of the present invention. Binding can be measured using a variety of methods standard in the art including enzyme immunoassays, e.g., ELISA, immunoblot assays, etc.; see, for example, Sambrook et al., *ibid.*, and Harlow, et al., 1988, *Antibodies, a Laboratory Manual*, Cold Spring Harbor Labs Press; Harlow et al., *ibid.*, is incorporated herein by reference in its entirety. For example, an anti-feline IL-18 antibody of the present invention preferably selectively binds to a feline IL-18 protein in such a way as to inhibit the function of that protein.

The antibodies of the present invention bind to the proteins of the present invention, but not to similar proteins of other species. For instance, the antibodies that specifically bind feline IL-18 do not bind canine IL-18.

Isolated antibodies of the present invention can include antibodies in serum, or antibodies that have been purified to varying degrees. Antibodies of the present invention can be polyclonal or monoclonal, or can be functional equivalents such as antibody fragments and genetically-engineered antibodies, including single chain antibodies or chimeric antibodies that can bind to one or more epitopes.

A preferred method to produce antibodies of the present invention includes (a) administering to an animal an effective amount of a protein, peptide or mimotope of the present invention to produce the antibodies and (b) recovering the antibodies. In another method, antibodies of the present invention are produced recombinantly using techniques as heretofore disclosed to produce proteins of the present invention. Antibodies raised against defined proteins or mimitopes can be advantageous because such antibodies are not substantially contaminated with antibodies against other substances that might otherwise cause interference in a diagnostic assay or side effects if used in a therapeutic composition.

Antibodies of the present invention have a variety of potential uses that are within the scope of the present invention. For example, such antibodies can be used (a) to evaluate the immune status in felids and canids with diseases such as allergy, cancer and pathogen infections. Furthermore, antibodies of the present invention can be used to target cytotoxic agents to cells. Targeting can be accomplished by conjugating, i.e., stably joining, such antibodies to the cytotoxic agents using techniques known to those skilled in the art. Suitable cytotoxic agents are known to

those skilled in the art. Furthermore, antibodies of the present invention can be used to detect for example, feline IL-18, caspase-1, canine IL-12 single chain, and/or feline IL-12 single chain in a putative IL-18, caspase-1, canine IL-12 single chain, and/or feline IL-12 single chain containing biological sample, by contacting the putative IL-18, caspase-1, canine IL-12 single chain, and/or feline IL-12 single chain containing biological sample with the appropriate anti-IL-18, caspase-1, canine IL-12 single chain, and/or feline IL-12 single chain antibodies under conditions suitable for formation of an antigen-antibody complex, and then detecting said complex. Methods to detect said method are known to those skilled in the art and are contained herein.

The present invention includes proteins comprising SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:59, SEQ ID NO:62, and SEQ ID NO:67 as well as nucleic acid molecules encoding such proteins.

Preferred feline IL-18 proteins of the present invention include PFeIL-18-N<sub>133</sub>, PFeIL-18-C<sub>154</sub>, PFeIL-18<sub>192</sub>, and/or PFeIL-18<sub>157</sub>. In one embodiment, a preferred feline IL-18 protein of the present invention has an amino acid sequence that includes SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and/or SEQ ID NO:12 and is preferably encoded by a nucleic acid molecule having nucleic acid sequences SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:9 and/or SEQ ID NO:11. Such proteins are preferably encoded by a nucleic acid molecule comprising nFeIL-18-N<sub>514</sub>, nFeIL-18-C<sub>502</sub>, nFeIL-18<sub>607</sub>, nFeIL-18<sub>576</sub>, and/or nFeIL-18<sub>471</sub>.

Preferred feline caspase-1 proteins of the present invention include proteins encoded by a nucleic acid molecule comprising nFeCasp-1<sub>1233</sub>, nFeCasp-1-N<sub>526</sub>, nFeCasp-1-C<sub>500</sub>, and/or nFeCasp-1<sub>1230</sub>. Preferred feline caspase-1 proteins are PFeCasp-1<sub>410</sub>, PFeCasp-1-N<sub>169</sub>, and/or PFeCasp-1-C<sub>120</sub>. In one embodiment, a preferred feline caspase-1 protein of the present invention is encoded by SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, and/or SEQ ID NO:23, and, as such, has an amino acid sequence that includes SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21 and/or SEQ ID NO:24.

Preferred canine and feline IL-12 proteins of the present invention include proteins encoded by a nucleic acid molecule comprising nFeIL-12<sub>1599</sub>, nFeIL-12<sub>1533</sub>, nCaIL-12<sub>1599</sub>, and/or nCaIL-12<sub>1533</sub>. Preferred feline and canine IL-12 proteins are nFeIL-12p40-N<sub>985</sub>, nFeIL-12p40<sub>987</sub>, nFeIL-12p35<sub>666</sub>, nFeIL-12p35-N<sub>561</sub>, nFeIL-12p35<sub>591</sub>, nCaIL-12p35<sub>666</sub>, nCaIL-12p35<sub>1455</sub>, nCaIL-12p35<sub>591</sub>, nCaIL-12p40<sub>2267</sub>, nCaIL-12p40<sub>921</sub>, and nCaIL-12p40<sub>987</sub>. In one embodiment, a preferred canine and feline IL-12 single chain protein of the present invention is encoded by SEQ ID NO:38, SEQ ID NO:43, SEQ ID NO:61, and/or SEQ ID NO:66, and, as such, has an amino acid sequence that includes SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:62, and/or SEQ ID NO:67.

More preferred canine and feline IL-12 single chain proteins of the present invention include proteins encoded by a nucleic acid molecule comprising nFeIL-12p40-N<sub>985</sub>, nFeIL-12p40<sub>987</sub>, nFeIL-12p40<sub>921</sub>, nFeIL-12p35<sub>666</sub>, nFeIL-12p35-N<sub>561</sub>, nFeIL-12p35<sub>591</sub>, nCaIL-12p35<sub>666</sub>, nCaIL-12p35<sub>1455</sub>, nCaIL-12p35<sub>591</sub>, nCaIL-12p42<sub>2267</sub>, nCaIL-12p40<sub>921</sub>, nCaIL-12p40<sub>987</sub>, nCaIL-12<sub>1533</sub>, nCaIL-12<sub>1599</sub>, nFeIL-12<sub>1533</sub>, and nFeIL-12<sub>1599</sub>. Preferred feline and canine

IL-12 single chain proteins comprise PFeIL-12p40-N<sub>328</sub>, PFeIL-12p40<sub>329</sub>, PFeIL-12p40<sub>307</sub>, PFeIL-12p35<sub>222</sub>, PFeIL-12p35-N<sub>187</sub>, PFeIL-12p35<sub>197</sub>, PCaIL-12p35<sub>222</sub>, PCaIL-12p35<sub>197</sub>, PCaIL-12p40<sub>307</sub>, PCaIL-12p40<sub>329</sub>, PFeIL-12<sub>533</sub>, PFeIL-12<sub>511</sub>, PCaIL-12<sub>533</sub>, and PCaIL-12<sub>511</sub>. In one embodiment, a preferred canine and feline IL-12 single chain protein of the present invention is encoded by a nucleic acid comprising SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:101, SEQ ID NO:104, SEQ ID NO:107, SEQ ID NO:38, SEQ ID NO:43, SEQ ID NO:61, SEQ ID NO:66, and, as such, has an amino acid sequence that includes SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67.

As used herein, an isolated protein of the present invention can be a full-length protein or any homolog of such a protein. An isolated protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to bind to a receptor or a protein. Examples of protein homologs of the present invention include proteins of the present invention in which amino acids have been deleted (e.g., a truncated version of the protein, such as a peptide), inserted, inverted, substituted and/or derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation and/or addition of glycerophosphatidyl inositol) such that the protein homolog includes at least one epitope capable of eliciting an immune response against the parent protein, of binding to an antibody directed against the parent protein and/or of binding to the parent's receptor, where the term parent refers to the longer and/or full-length protein that the homolog is derived from. That is, when the homolog is administered to an animal as an immunogen, using techniques known to those skilled in the art, the animal will produce an immune response against at least one epitope of an immunoregulatory protein of the present invention, depending upon which protein is administered to an animal. The ability of a protein to effect an immune response can be measured using techniques known to those skilled in the art.

Homologs of proteins of the present invention can be the result of natural allelic variation, including natural mutation. Protein homologs of the present invention can also be produced using techniques known in the art including, but not limited to, direct modifications to the protein and/or modifications to the gene encoding the protein using, for example, classic or recombinant DNA techniques to effect random or targeted mutagenesis.

One embodiment of the present invention is an IL-18 protein selected from the group consisting of: (i) a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12; and (ii) a protein encoded by an allelic variant of a nucleic acid molecule encoding a protein selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:12. Another embodiment is a caspase-1 protein selected from the group consisting of: (i) a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24; and (ii) a protein encoded by an allelic variant of a nucleic acid molecule encoding a protein selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24. Yet another embodiment is a feline

IL-12 single chain protein selected from the group consisting of: (i) a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:38, and SEQ ID NO:44; and (ii) a protein encoded by an allelic variant of a nucleic acid molecule encoding a protein selected from the group consisting of SEQ ID NO:38, and SEQ ID NO:44; or a canine IL-12 single chain protein selected from the group consisting of: (i) a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:62 and SEQ ID NO:67; and (ii) a protein encoded by an allelic variant of a nucleic acid molecule encoding a protein selected from the group consisting of SEQ ID NO:62 and SEQ ID NO:67.

One embodiment of the present invention includes an isolated IL-18 protein selected from the group consisting of (i) an isolated protein of at least 25 amino acids in length, wherein said protein has an at least 25 contiguous amino acid region identical in sequence to a 25 contiguous amino acid region selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12; and (ii) an isolated protein having an amino acid sequence that is at least 92 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12, and a fragment thereof of at least 30 nucleotides. Preferred proteins have an at least 15 contiguous amino acid region identical with a 15 contiguous amino acid region, an at least 20 contiguous amino acid region identical with a 20 contiguous amino acid region, an at least 30 contiguous amino acid region identical with a 30 contiguous amino acid region, an at least 40 contiguous amino acid region identical with a 40 contiguous amino acid region, an at least 50 contiguous amino acid region contiguous with a 50 contiguous amino acid region, an at least 75 contiguous amino acid region contiguous with a 75 contiguous amino acid region, preferably an at least 100 contiguous amino acid region contiguous with a 100 contiguous amino acid region, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12. In another embodiment, preferred proteins have an amino acid sequence that is at least 90 percent identical, at least 92 percent identical, preferably at least 94 percent identical, preferably at least 96 percent identical, and even more preferably at least about 98 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12, and a fragment thereof of at least 20 amino acids, at least 30 amino acids, at least 50 amino acids, at least 75 amino acids, preferably at least 100 amino acids, and more preferably a full-length protein.

In a preferred embodiment, IL-18 proteins of the present invention has a function selected from the group consisting of: (i) eliciting an immune response against an IL-18 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12, (ii) selectively binding to an antibody raised against an IL-18 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, and SEQ ID NO:12, and (iii) exhibiting IL-18 activity.

One embodiment of the present invention includes an isolated caspase-1 protein selected from the group consisting of (i) an isolated protein of at least 25 amino acids in length, wherein said protein has an at least 25 contiguous amino acid region identical in sequence to a 25 contiguous amino acid region selected from the group consisting of SEQ ID

NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24; and (ii) an isolated protein having an amino acid sequence that is at least 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and a fragment thereof of at least 30 nucleotides. Preferred proteins have an at least 25 contiguous amino acid region identical with a 25 contiguous amino acid region, an at least 20 contiguous amino acid region identical with a 20 contiguous amino acid region, an at least 30 contiguous amino acid region identical with a 30 contiguous amino acid region, an at least 40 contiguous amino acid region identical with a 40 contiguous amino acid region, an at least 50 contiguous amino acid region contiguous with a 50 contiguous amino acid region, an at least 75 contiguous amino acid region contiguous with a 75 contiguous amino acid region, preferably an at least 100 contiguous amino acid region contiguous with a 100 contiguous amino acid region, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24. In another embodiment, preferred proteins have an amino acid sequence that is at least 85 percent identical, at least 88 percent identical, preferably at least 90 percent identical, and more preferably at least about 95 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and a fragment thereof of at least 30 amino acids, at least 50 amino acids, at least 75 amino acids, preferably at least 100 amino acids, and more preferably a full-length protein.

In a preferred embodiment, a caspase protein of the present invention has a function selected from the group consisting of: (i) eliciting an immune response against a caspase-1 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, (ii) selectively binding to an antibody raised against a caspase-1 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, and SEQ ID NO:24, and (iii) exhibiting caspase-1 activity.

One embodiment of the present invention includes an isolated IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to an IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p40 subunit protein comprising at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30. The p35 subunit is preferably selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:33 and SEQ ID NO:36, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 30 amino acids in length, and (iii) a p35 subunit protein comprising at least 23 contiguous amino acids identical in sequence to at least 23 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID

NO:33 and SEQ ID NO:36. Preferred amino acid sequences have an at least 23 contiguous amino acid region identical with a 23 contiguous amino acid region, an at least 30 contiguous amino acid region identical with a 30 contiguous amino acid region, an at least 40 contiguous amino acid region identical with a 40 contiguous amino acid region, an at least 50 contiguous amino acid region contiguous with a 50 contiguous amino acid region, an at least 75 contiguous amino acid region contiguous with a 75 contiguous amino acid region, preferably an at least 100 contiguous amino acid region contiguous with a 100 contiguous amino acid region, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:27 and SEQ ID NO:30. In another embodiment, preferred proteins have an amino acid sequence that is at least 84 percent identical, at least 86 percent identical, at least 88 percent identical, preferably at least 90 percent identical, and more preferably at least about 95 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, and a fragment thereof of at least 30 amino acids, at least 50 amino acids, at least 75 amino acids, preferably at least 100 amino acids, and more preferably a full-length protein.

In a preferred embodiment, an IL-12 single chain protein of the present invention has a function selected from the group consisting of: (i) eliciting an immune response against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67; (ii) selectively binding to an antibody raised against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67; and (iii) exhibiting IL-12 activity.

One embodiment of the present invention includes an isolated IL-12 single chain protein comprising an IL-12 p40 subunit domain linked to an IL-12 p35 subunit domain, wherein said p40 subunit domain is selected from the group consisting of (i) a p40 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59, (ii) a p40 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p40 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:53 and SEQ ID NO:59. The p35 subunit is preferably selected from the group consisting of (i) a p35 subunit protein having an amino acid sequence that is at least 84 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50, (ii) a p35 subunit protein comprising a fragment of a protein of (i), wherein said fragment is at least 40 amino acids in length, and (iii) a p35 subunit protein comprising at least 31 contiguous amino acids identical in sequence to at least 31 contiguous amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:47 and SEQ ID NO:50. Preferred amino acid sequences have an at least 23 contiguous amino acid region identical

with a 23 contiguous amino acid region, an at least 30 contiguous amino acid region identical with a 30 contiguous amino acid region, an at least 40 contiguous amino acid region identical with a 40 contiguous amino acid region, an at least 50 contiguous amino acid region contiguous with a 50 contiguous amino acid region, an at least 75 contiguous amino acid region contiguous with a 75 contiguous amino acid region, preferably an at least 100 contiguous amino acid region contiguous with a 100 contiguous amino acid region, and most preferably a full-length protein identical in sequence to a full-length protein of an amino acid sequence selected from the group consisting of SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59. In another embodiment, preferred proteins have an amino acid sequence that is at least 84 percent identical, at least 86 percent identical, at least 88 percent identical, preferably at least 90 percent identical, and more preferably at least about 95 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, and a fragment thereof of at least 30 amino acids, at least 50 amino acids, at least 75 amino acids, preferably at least 100 amino acids, and more preferably a full-length protein.

In a preferred embodiment, an IL-12 single chain protein of the present invention has a function selected from the group consisting of: (i) eliciting an immune response against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:62 and SEQ ID NO:67, (ii) selectively binding to an antibody raised against an IL-12 protein having an amino acid sequence selected from the group consisting of SEQ ID NO:27, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, and SEQ ID NO:59, SEQ ID NO:102, SEQ ID NO:105, SEQ ID NO:108, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:62, and/or SEQ ID NO:67, and (iii) exhibiting IL-12 activity.

One embodiment of the present invention is a therapeutic composition that, when administered to a animal in an effective manner, is capable of protecting that animal from a disease such as, for example, allergy, cancer or inflammation. Therapeutic compositions of the present invention include protective compounds that are capable of regulating feline IL-18, feline caspase-1, or feline or canine IL-12 protein amounts and/or activity. A protective compound of the present invention is capable of regulating feline IL-18, feline caspase-1, or feline or canine IL-12 activity and/or availability. Examples of protective compounds related to feline and canine proteins of the present invention include an isolated antibody that selectively binds to either feline IL-18, feline caspase-1, or feline or canine IL-12 or other inhibitors or activators of feline IL-18, feline caspase-1, or feline or canine IL-12 activity or amount. Other examples of protective compounds include an isolated nucleic acid molecule of the present invention; an isolated protein of the present invention; a mimetope of a protein of the present invention, a multimeric form of any of said proteins, or an inhibitor identified by its ability to inhibit the activity of any of said proteins; such an inhibitor can inhibit binding of the respective protein with its receptor, or inhibit the activity of the respective protein. Methods to perform such assays to measure binding and/or activity of protein of the present invention are known to those of skill in the art, and are described, for example, in Janeway et al., *ibid.* As such, these protective compounds may include antibodies, peptides, substrate analogs, and other large or small molecules which can be organic or inorganic. As used herein, a protective compound refers to a compound, that when

administered to an animal in an effective manner, is able to treat, ameliorate, and/or prevent a disease due to allergy, cancer or infection. Examples of proteins, nucleic acid molecules, antibodies and/or inhibitors of the present invention are disclosed herein.

The present invention also includes a therapeutic composition comprising at least one compound of the present invention in combination with at least one additional therapeutic compound. Examples of such compounds are disclosed herein.

The efficacy of a therapeutic composition of the present invention to protect an animal from a disease mediated by feline IL-18, feline caspase-1, or feline or canine IL-12 can be tested in a variety of ways including, but not limited to, detection of protective antibodies (using, for example, proteins or mimetopes of the present invention), detection of the amount of feline IL-18, feline caspase-1, or feline or canine IL-12, or detection of cellular immunity within the treated animal. Therapeutic compositions can be tested in animal models such as mice. Such techniques are known to those skilled in the art.

Therapeutic compounds of the present invention can be administered to any animal susceptible to such therapy, preferably to mammals, and more preferably to dogs, cats, humans, ferrets, horses, cattle, sheep and/or other pets, economic food animals, and/or zoo animals. Preferred animals include dogs and cats.

A therapeutic composition of the present invention is administered to an animal in an effective manner such that the composition is capable of regulating an immune response in that animal. Therapeutic compositions of the present invention can be administered to animals prior to the onset of a disease (i.e. as a preventative vaccine) and/or can be administered to animals after onset of a disease in order to treat the disease (i.e. as a therapeutic vaccine). Preferred diseases to prevent and/or treat include autoimmune diseases, allergic reactions, infectious diseases, tumor development, inflammatory diseases and/or graft rejection. In one embodiment, a therapeutic composition of the present invention is administered with an antigen to enhance an immune response against that antigen. Such administration can include, but is not limited to, oral, intravenous, intramuscular, intra ocular, mucosal, intranasal, subcutaneous, topical or transdermal application. In order to protect an animal from disease, a therapeutic composition of the present invention is administered to the animal in an effective manner such that the composition is capable of protecting that animal from a disease. Therapeutic compositions of the present invention can be administered to animals prior to disease in order to prevent disease and/or can be administered to animals after disease occurs. The exact dose, administration regimen, and administration route of therapeutic compositions of the present invention can be determined by one skilled in the art. A suitable single dose is a dose that is capable of regulating the immune response in an animal when administered one or more times over a suitable time period. For example, a preferred single dose of a protein, mimetope or antibody therapeutic composition is from about 1 microgram ( $\mu\text{g}$ ) to about 10 milligrams (mg) of the therapeutic composition per kilogram body weight of the animal. Booster vaccinations can be administered from about 2 weeks to several years after the original administration. Booster administrations preferably are administered when the immune response of the animal becomes insufficient to protect the animal from disease. A preferred administration schedule is one in which from about 10  $\mu\text{g}$  to about 1 mg of the therapeutic composition per kg body weight of

the animal is administered from about one to about two times over a time period of from about 2 weeks to about 12 months.

A therapeutic composition of the present invention can include at least one of the following: excipient, an adjuvant and a carrier. Therapeutic compositions of the present invention can be formulated in an excipient that the animal to be treated can tolerate. Examples of such excipients include water, saline, Ringer's solution, dextrose solution, Hank's solution, and other aqueous physiologically balanced salt solutions. Nonaqueous vehicles, such as fixed oils, sesame oil, ethyl oleate, or triglycerides may also be used. Other useful formulations include suspensions containing viscosity enhancing agents, such as sodium carboxymethylcellulose, sorbitol, or dextran. Excipients can also contain minor amounts of additives, such as substances that enhance isotonicity and chemical stability. Examples of buffers include phosphate buffer, bicarbonate buffer and Tris buffer, while examples of preservatives include thimerosal, or o-cresol, formalin and benzyl alcohol. Standard formulations can either be liquid injectables or solids which can be taken up in a suitable liquid as a suspension or solution for injection. Thus, in a non-liquid formulation, the excipient can comprise dextrose, human serum albumin, preservatives, etc., to which sterile water or saline can be added prior to administration.

Therapeutic compositions of the present invention can include an adjuvant. Adjuvants are agents that are capable of enhancing the immune response of an animal to a specific antigen. Suitable adjuvants include, but are not limited to, cytokines, chemokines, and compounds that induce the production of cytokines and chemokines (e.g., granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), colony stimulating factor (CSF), erythropoietin (EPO), interleukin 2 (IL-2), interleukin-3 (IL-3), interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 6 (IL-6), interleukin 7 (IL-7), interleukin 8 (IL-8), interleukin 10 (IL-10), interferon gamma, transforming growth factor beta, RANTES (regulated upon activation, normal T cell expressed and presumably secreted), macrophage inflammatory proteins (e.g., MIP-1 alpha and MIP-1 beta), and *Leishmania* elongation initiating factor (LEIF); bacterial components (e.g., endotoxins, in particular superantigens, exotoxins and cell wall components); aluminum-based salts; calcium-based salts; silica; polynucleotides; toxoids; serum proteins, viral coat proteins; block copolymer adjuvants (e.g., Hunter's Titermax™ adjuvant (Vaxcel™, Inc. Norcross, Ga.), Ribi adjuvants (Ribi ImmunoChem Research, Inc., Hamilton, Mont.); and saponins and their derivatives (e.g., Quil A (Superfos Biosector A/S, Denmark). Protein adjuvants of the present invention can be delivered in the form of the protein themselves or of nucleic acid molecules encoding such proteins using the methods described herein.

Therapeutic compositions of the present invention can include a carrier. Carriers include compounds that increase the half-life of a protective compound in the treated animal. Suitable carriers include, but are not limited to, polymeric controlled release vehicles, biodegradable implants, liposomes, other lipid or lipid containing formulations, including cationic lipids or lipid mixtures including cationic lipids, bacteria, viruses, other cells, oils, esters, and glycols.

A therapeutic composition can be a controlled release formulation that is capable of slowly releasing a protective compound of the present invention into an animal. As used herein, a controlled release formulation comprises a com-

position or protective compound of the present invention in a controlled release vehicle. Suitable controlled release vehicles include, but are not limited to, biocompatible polymers, other polymeric matrices, capsules, microcapsules, microparticles, bolus preparations, osmotic pumps, diffusion devices, liposomes, lipospheres, other lipids or lipid-containing formulations and transdermal delivery systems. Other controlled release formulations of the present invention include liquids that, upon administration to an animal, form a solid or a gel in situ. Preferred controlled release formulations are biodegradable, i.e., bioerodible.

A preferred controlled release formulation of the present invention is capable of releasing a composition of the present invention into the blood of the treated animal at a constant rate sufficient to attain therapeutic dose levels of the composition to regulate an immune response in an animal. The therapeutic composition is preferably released over a period of time ranging from about 1 to about 12 months. A controlled release formulation of the present invention is capable of effecting a treatment preferably for at least about 1 month, more preferably for at least about 3 months, even more preferably for at least about 6 months, even more preferably for at least about 9 months, and even more preferably for at least about 12 months.

According to one embodiment, a nucleic acid molecule of the present invention can be administered to an animal in a fashion to enable expression of that nucleic acid molecule into a therapeutic protein or therapeutic RNA (e.g. antisense RNA, ribozyme, triple helix forms or RNA drug) in the animal. Nucleic acid molecules can be delivered to an animal in a variety of methods including, but not limited to, (a) administering a naked (i.e. not packaged in a viral coat or cellular membrane) nucleic acid as a genetic vaccine (e.g. as naked DNA or RNA molecules, such is taught, for example, in Wolff et al., 1990, Science 247, p 1465-68) or (b) administering a nucleic acid molecule packaged as a recombinant virus vaccine or as a recombinant cell vaccine (i.e. the nucleic acid molecule is delivered by a viral or cellular vehicle).

One embodiment of a therapeutic composition of the present invention is a naked nucleic acid, a recombinant virus or a recombinant cell vaccine or therapy. Naked nucleic acid molecules of the present invention can be administered by a variety of methods. Suitable delivery methods include, for example, intramuscular injection, subcutaneous injection, intradermal injection, intradermal scarification, particle bombardment, oral application, topical application and nasal application, with intramuscular injection, intradermal injection, intradermal scarification and particle bombardment being preferred. A preferred single dose of a naked nucleic acid molecule ranges from about 1 nanogram (ng) to about 1 milligram (mg), depending on the route of administration and/or method of delivery, as can be determined by those skilled in the art. Examples of administration methods are disclosed, for example, in U.S. Pat. No. 5,204,253, by Bruner, et al., issued Apr. 20, 1993, PCT Publication No. WO 95/19799, published Jul. 27, 1995, by McCabe, and PCT Publication No. WO 95/05853, published Mar. 2, 1995, by Carson, et al. Naked nucleic acid molecules of the present invention can be contained in an aqueous excipient (e.g., phosphate buffered saline) and/or with a carrier (e.g., lipid-based vehicles), or it can be bound to microparticles (e.g., gold particles).

According to one embodiment, a nucleic acid molecule of the present invention can be administered to an animal in a fashion to enable expression of that nucleic acid molecule into a protective protein or protective RNA, e.g., antisense

RNA, ribozyme, triple helix form or RNA drug, in the animal. Nucleic acid molecules can be delivered to an animal in a variety of methods including, but not limited to, (a) administering a naked, i.e., not packaged in a viral coat or cellular membrane, nucleic acid as a genetic therapy or vaccine, e.g., as naked DNA or RNA molecules, such as is taught, for example in Wolff et al., 1990, *Science* 247, 1465–1468, or (b) administering a nucleic acid molecule packaged as a recombinant virus therapy or vaccine or as a recombinant cell therapy or vaccine, i.e., the nucleic acid molecule is delivered by a viral or cellular vehicle.

A genetic, i.e., naked nucleic acid, therapy vaccine of the present invention includes a nucleic acid molecule of the present invention and preferably includes a recombinant molecule of the present invention that preferably is replication or otherwise amplification, competent. A genetic therapy or vaccine of the present invention can comprise one or more nucleic acid molecules of the present invention operatively linked to a transcriptional control sequence in the form of, for example, a dicistronic recombinant molecule. Preferred viral vectors include those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, picornaviruses, and/or retroviruses, with those based on alphaviruses (such as Sindbis or Semliki forest virus) species-specific herpesviruses and/or poxviruses being particularly preferred. Any suitable transcription control sequence can be used, including those disclosed as suitable for protein production. Particularly preferred transcription control sequences include cytomegalovirus immediate early (preferably in conjunction with Intron-A), Rous sarcoma virus long terminal repeat, and tissue-specific transcription control sequences, as well as transcription control sequences endogenous to viral vectors if viral vectors are used. The incorporation of a “strong” polyadenylation signal is also preferred.

Genetic therapies and vaccines of the present invention can be administered in a variety of ways, with intramuscular, subcutaneous, intradermal, transdermal, intranasal, topical and oral routes of administration being preferred. A preferred single dose of a genetic therapy or vaccine ranges from about 1 nanogram (ng) to about 600 µg, depending on the route of administration and/or method of delivery, as can be determined by those skilled in the art. Suitable delivery methods include, for example, by injection, as drops, aerosolized and/or topically. Genetic therapies or vaccines of the present invention can be contained in an aqueous excipient, e.g., phosphate buffered saline, alone or in a carrier, e.g., lipid-based vehicles. One embodiment is a nucleic acid-lipid complex, preferably a nucleic acid-cationic lipid complex.

A recombinant virus therapy or vaccine of the present invention includes a recombinant molecule of the present invention that is packaged in a viral coat and that can be expressed in an animal after administration. Preferably, the recombinant molecule is packaging- or replication-deficient and/or encodes an attenuated virus. A number of recombinant viruses can be used, including, but not limited to, those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, picornaviruses, and retroviruses. Preferred recombinant virus vaccines are those based on alphaviruses, such as Sindbis virus, raccoon poxviruses, species-specific herpesviruses and species-specific poxviruses. An example of methods to produce and use recombinant virus therapies and vaccines is disclosed in U.S. Pat. No. 5,766,602, Xiong et al., issued Jun. 16, 1998; U.S. Pat. No. 5,753,235, Haanes et al., issued May 19, 1998; and U.S. Pat. No. 5,804,197, Haanes et al., issued Sep. 8, 1998, all of which are incorporated by reference herein in their entireties.

When administered to an animal, a recombinant virus therapy or vaccine of the present invention infects cells within the immunized animal and directs the production of a protective protein or RNA nucleic acid molecule that is capable of protecting the animal from a disease. For example, a recombinant virus vaccine comprising a feline IL-18 nucleic acid molecule of the present invention can be administered according to a protocol that results in the animal producing a sufficient immune response to protect itself from a disease mediated by IL-18. In another embodiment of the present invention a feline IL-18 nucleic acid molecule can be used as therapy to treat a disease. A recombinant virus vaccine comprising a feline IL-18 nucleic acid molecule can be administered to an animal with clinical signs of disease according to a protocol that results in reduction and/or termination of clinical signs of disease. A preferred single dose of a recombinant virus therapy or vaccine of the present invention is from about  $1 \times 10^4$  to about  $1 \times 10^8$  virus plaque forming units (pfu) per kilogram body weight of the animal. Administration protocols are similar to those described herein for protein-based vaccines, with subcutaneous, intramuscular, intranasal, topical and oral administration routes being preferred.

A recombinant cell therapy or vaccine of the present invention includes recombinant cells of the present invention that express at least one protein of the present invention. Preferred recombinant cells for this embodiment include *Salmonella*, *E. coli*, *Listeria*, *Mycobacterium*, *S. frugiperda*, yeast, (including *Saccharomyces cerevisiae* and *Pichia pastoris*), BHK, CV-1, myoblast G8, COS, e.g., COS-7, Vero, MDCK and CRFK recombinant cells. Recombinant cell therapy or vaccines of the present invention can be administered in a variety of ways but have the advantage that they can be administered orally, preferably at doses ranging from about  $10^8$  to about  $10^{12}$  cells per kilogram body weight. Administration protocols are similar to those described herein for protein-based vaccines. Recombinant cell vaccines can comprise whole cells, cells stripped of cell walls or cell lysates.

In one embodiment of the present invention, a method to regulate an immune response in an animal by administering the therapeutic compound to an animal preferably a canine or feline, wherein the composition comprises a component selected from the group consisting of an excipient, an adjuvant and a carrier.

Proteins of the present invention can be used to develop regulatory compounds including inhibitors and activators that, when administered to an animal in an effective manner, are capable of protecting that animal from disease mediated by IL-18, caspase-1 or IL-12. Preferred regulatory compounds derived from the present invention include inhibitors and activators. In accordance with the present invention, the ability of a regulatory compound, including an inhibitor or activator, of the present invention to protect a felid or canid from disease mediated by IL-18, caspase-1 or IL-12 refers to the ability of that compound to, for example, treat, ameliorate or prevent a disease mediated by IL-18, caspase-1 or IL-12 in that animal.

An IL-18, caspase-1 or IL-12 single chain inhibitor of the present invention is identified by its ability to bind to, modify, or otherwise interact with, an IL-18, caspase-1 or IL-12 single chain protein of the present invention, thereby inhibiting the activity of the protein. Suitable inhibitors of activity are compounds that inhibit the activity of the proteins of the present invention in at least one of a variety of ways: (1) by binding to or otherwise interacting with or otherwise modifying the protein binding, (2) by interacting

with other regions of the protein to inhibit activity, for example, by allosteric interaction, and (3) by binding to or otherwise interacting with or otherwise modifying a protein receptor binding site such that the protein is less likely to bind to the protein receptor binding site. Inhibitors of IL-18, caspase-1 and IL-12 single chain proteins are preferably relatively small compounds.

An embodiment of the present invention includes use of one of the following methods to identify a compound capable of regulating an immune response in an animal: (a) contacting an isolated feline IL-18 protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has T cell stimulating activity; and determining if the putative inhibitory compound inhibits the activity; (b) contacting an isolated feline caspase-1 protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein cleaves precursor IL-18 resulting in a biologically active mature IL-18; and determining if the putative inhibitory compound inhibits the activity; and (c) contacting an isolated IL-12 single chain protein with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has T cell proliferation stimulating activity; and determining if the putative inhibitory compound inhibits the activity.

A variety of methods are known to one skilled in the art to detect binding of an IL-18, caspase-1 or IL-12 protein to its binding partner (e.g., an antibody or receptor, as appropriate). Such methods can be used to detect IL-12, casp-1, or IL-18, or Abs or other binding partners thereof in a biological sample or to produce inhibitors of such interactions. Such methods include, but are not limited to an assay in which, for example, IL-18 and an IL-18 binding partner can interact and/or bind to each other, using, for example, the yeast two-hybrid system, see for example, Luban, et al. 1995, *Curr. Opin. Biotechnol.*, 6, 59–64; and identifying those proteins that specifically bind to the IL-18 protein binding domain. Additional methods to identify protein—protein interactions include Biacore® screening, confocal immunofluorescent microscopy, UV cross-linking, and immunoprecipitations. An example of a protein binding domain is an IL-18-binding domain, and a protein that would bind to an IL-18-binding domain would be IL-18. Additional teachings of general characteristics of reagents for use in the detection of binding between two moieties (e.g., between IL-18 and its receptor) as well as methods to produce and use such reagents are disclosed, for example, in U.S. Pat. No. 5,958,880, issued Sep. 28, 1999, by Frank et al.; and PCT International Publication No. WO 99/54349, published Oct. 28, 1999, by McCall et al.; each of these references is incorporated by reference herein in its entirety; furthermore, the disclosed reagents and methods are incorporated by reference herein in their entireties. It is to be noted that although the reagents and methods disclosed in each of the citations do not relate to the proteins, nucleic acid molecules, antibodies and inhibitors of the present invention per se, the disclosed reagents and methods are applicable by those skilled in the art to reagents, kits and detection methods of the present invention. Furthermore, proteins of the present invention can be used to develop regulatory compounds including inhibitors and activators that, when administered to a canid or felid in an effective manner are capable of protecting and treating that felid or canid from disease mediated by IL-18, caspase-1 or IL-12.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention. The following examples include a number

of recombinant DNA and protein chemistry techniques known to those skilled in the art; see, for example, Sambrook et al., *ibid.*

## EXAMPLE 1

Identification of the nucleic acid molecules of the feline IL-18 is unexpected because initial attempts to isolate feline IL-18 nucleic acid molecules using standard cDNA screening techniques were unsuccessful.

This example describes the isolation, sequencing and expression of nucleic acid molecules encoding feline IL-18 proteins of the present invention.

A. Feline IL-18 nucleic acid molecules were isolated as follows: A cDNA mitogen library was prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours as previously described in Example 2 of PCT Publication No. WO 99/61618, entitled “Canine and Feline Immunoregulatory Proteins, Nucleic Acid Molecules, and Uses Thereof,” inventors Gek-Kee Sim, Shumin Yang, Matthew Dreitz, and Ramani Wonderling, filed May 28, 1999, which is incorporated by reference herein in its entirety. An aliquot of this library was used as a template to isolate a feline IL-18 nucleic acid molecule by polymerase chain reaction (PCR). PCR amplification was performed using Amplitaq DNA polymerase™ (available from PE Applied Biosystems Inc, Foster City, Calif.). Two overlapping nucleic acid molecules encoding partial length feline IL-18 proteins were obtained by using IL-18 specific primers in combination with cDNA library vector specific primers. All primers came from Life Technologies, Gaithersburg, Md. The sequence of the vector forward primer (T3 primer) was 5' GCCAAGCTCG AAATTAACCC TCACTAAAGG 3' (SEQ ID NO:72), and that of the vector reverse primer (T7 primer) was 5' CGACGGCCAG TGAATTGTAA TAC-GACTC 3' (SEQ ID NO:73). The sequence of the IL-18-specific forward primer (IL-18 Forward 85) was 5'AGT-GATGAAG GCCTGGAATC AGATTACTTT G 3' (SEQ ID NO:74) and the sequence of the IL-18-specific reverse primer (IL-18 Reverse 435) was 5' ATGGCCTGGA ACACTTCTCT GAAAGAATAT GA 3' (SEQ ID NO:75). The first PCR amplification was done using T3 primer and IL-18 Reverse 435 primer and the second PCR amplification was done using IL-18 Forward 85 primer and T7 primer. The PCR profile for both reactions was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 30 seconds, then 59° C. for 30 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. The PCR products from both reactions were cloned into the TA-Cloning vector (available from Invitrogen, San Diego, Calif.) and the nucleic acid molecules were sequenced using an ABI Prism™ Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism™ dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). The PCR product from the first PCR amplification was sequenced and found to contain 514 nucleotides and was denoted herein as nFeIL-18<sub>514</sub> (5'-end partial clone) with a coding strand of SEQ ID NO:1, and a complementary strand of SEQ ID NO:3. The PCR product from the second PCR amplification was sequenced and found to contain 502 nucleotides and was denoted herein as nFeIL-18<sub>502</sub> (3'-end partial clone) with a coding strand of SEQ ID NO:4, and a complementary strand of SEQ ID NO:6. These two nucleic acid molecules shared more than 280 base pairs (bp) and together provided the sequence

for the complete feline IL-18 open reading frame. Translation of SEQ ID NO:1 suggests that nucleic acid molecule nFeIL-18-N<sub>514</sub> encodes an N-terminal portion of PFeIL-18-N protein, of about 133 amino acids, denoted herein as PFeIL-18-N<sub>133</sub>, the amino acid sequence of which is presented in SEQ ID NO:2, assuming an open reading frame having an initiation codon spanning from nucleotide 114 through nucleotide 116 of SEQ ID NO:1 and a stop codon spanning from nucleotide 510 through nucleotide 512 of SEQ ID NO:1. Translation of SEQ ID NO:4 suggests that nucleic acid molecule nFeIL-18-C<sub>502</sub> encodes an C-terminal portion of PFeIL-18-C protein, of about 154 amino acids, denoted herein as PFeIL-18-C<sub>154</sub>, the amino acid sequence of which is presented in SEQ ID NO:5, assuming an open reading frame having an initiation codon spanning from nucleotide 3 through nucleotide 5 of SEQ ID NO:4 and a stop codon spanning from nucleotide 462 through nucleotide 464 of SEQ ID NO:4.

Based on the sequence data obtained from these two nucleic acid molecules two new primers were made to isolate a cDNA encoding full-length feline IL-18. The IL-18 Full Forward primer sequence was 5' AACTATTGAG CACAGGGATA AAGATGACTG 3' (SEQ ID NO:76) and IL-18 Full Reverse primer sequence was 5' AATATCTAAT TCTTGTTTTG AACAGTGAAC ATT 3' (SEQ ID NO:77). The PCR amplification was performed using these two primers and Amplitaq DNA polymerase<sup>TM</sup> (available from PE Applied Biosystems Inc.) and an aliquot of the cDNA library prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours. The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 30 seconds, then 53° C. for 30 seconds, then 72° C. for 90 seconds; followed by a final extension at 72° C. for 7 minutes. The PCR product was cloned into the TA-Cloning vector (available from Invitrogen, San Diego, Calif.) and the nucleic acid molecule insert was sequenced using an ABI PRISM<sup>TM</sup> Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using PRISM<sup>TM</sup> dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). This PCR product the FeIL-18 containing region of which is denoted nFeIL-18<sub>607</sub> was found to encode a full-length FeIL-18 protein. The nucleotide sequence of the coding strand of nFeIL-18<sub>607</sub> is represented herein as SEQ ID NO:7, and its complement is denoted by SEQ ID NO:10. Translation of the open reading frame in SEQ ID NO:7, denoted herein as nFeIL-18<sub>576</sub>, the coding strand of which is denoted SEQ ID NO: 9, and the complementary strand denoted SEQ ID NO:41 suggests that feline IL-18 encodes a protein containing 192 amino acids, referred to herein as PFeIL-18<sub>192</sub>, with a SEQ ID NO:8. The nucleic acid sequence encoding feline IL-18 protein assumes an open reading frame in which the first codon spans from nucleotide 24 through 26 of SEQ ID NO:7, and the last codon spans from nucleotide 597 through nucleotide 599 of SEQ ID NO:7. The encoded protein has a predicted molecular weight of about 21.3 kiloDaltons (kDa) for the precursor protein. The IL-18 precursor protein does not contain a signal sequence; in order for IL-18 to be biologically active the precursor is cleaved by caspase-1. The putative caspase-1 cleavage site is between amino acid positions 35 and 36 of the feline IL-18 precursor protein. Nucleic acid molecule nFeIL-18<sub>471</sub>, which encodes the mature protein contains a coding strand with SEQ ID NO:11, and a complementary strand with SEQ ID NO:13. The amino acid sequence of the mature protein, denoted herein as PFeIL-

18<sub>157</sub> is SEQ ID NO:12 and the mature protein has a predicted molecular weight of about 17.4 kDa. Sequence analysis was performed using DNAsis<sup>TM</sup>, available from Hitachi Software, San Bruno, Calif. using the alignment settings of: gap penalty set at 5, k-tuple set at 3, number of top diagonals set at 5, window size set at 5, fixed gap penalty set at 10 and floating gap penalty set at 10.

B. In an attempt to express a mature feline IL-18 protein in a mammalian cell line, the region encoding only the mature IL-18 protein (SEQ ID NO:11) was isolated from the feline cDNA library described in Example 1A using the following primers: IL-18 MatNgo Forward primer which has the sequence 5' TATGCCGGCT ACTTTGGCAA GCT-TGAACAT AAACCTC 3' (SEQ ID NO:78) and IL-18 MatXho Reverse primer which has the sequence 5' GGC-CTCGAGC TAATCTTGT TTTGAACAGT GAACATT 3' (SEQ ID NO:79). The PCR amplification was performed using these two primers and Amplitaq DNA polymerase<sup>TM</sup> (available from PE Applied Biosystems Inc.) and an aliquot of the cDNA library prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours. The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 30 seconds, then 53° C. for 30 seconds, then 72° C. for 90 seconds; followed by a final extension at 72° C. for 7 minutes. The PCR products were digested with Ngo MI and Xho I restriction enzymes (available from New England Biolabs, Beverly, Mass.) and ligated downstream of nucleotides encoding a tissue plasminogen activator (tPA) signal sequence contained in the CMV-IntronA-tPA vector (available from Invitrogen). The construct was sequenced using an ABI Prism<sup>TM</sup> Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism<sup>TM</sup> dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). This construct encoded the mature feline IL-18 protein with the tPA signal sequence. When Chinese hamster ovary (CHO) cells (available from ATCC, Rockville, Md.) were transiently transfected with this construct, using techniques known to those skilled in the art and cell pellets and supernatants were harvested after 48 hrs. Western analysis was performed on the cell pellets and supernatant samples using a polyclonal antibody against human IL-18 (available from Biosource International, Camarillo, Calif.). A faint band of the expected size (about 17.4 kDa) was detected in the cell pellet and not in the supernatant, indicating that IL-18 is produced by this construct but it is not exported out of the cell at detectable levels. While not being bound by theory, it is believed that caspase-1 plays a key role in the processing of native IL-18 precursor in cells where IL-18 is produced, co-expression of full-length feline IL-18 along with the feline caspase-1 may be necessary for the proper processing of the IL-18 precursor and enhanced secretion of the processed IL-18 mature polypeptide.

#### EXAMPLE 2

This example describes the isolation and sequencing of nucleic acid molecules encoding feline caspase-1 proteins of the present invention.

Feline caspase-1 nucleic acid molecules were isolated as follows: A cDNA mitogen library was prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours as described in Example 1. An aliquot of this library was used as a template to isolate a feline caspase-1 by polymerase chain reaction (PCR). PCR amplification was

51

performed using Amplitaq DNA polymerase™ (available from PE Applied Biosystems Inc.). The forward and reverse primers were designed based on human caspase-1 sequences. The forward primer (Casp-1For) had a sequence of 5' ATGGCCGACA AGGTCCTGAA GGAGAAGA 3' (SEQ ID NO:80) and the reverse primer (Casp-1 Rev) had a sequence of 5' TTAATGTCCT GGAAGAGGT AGAAACATCT TGT 3' (SEQ ID NO:81). The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 45 seconds, then 53° C. for 45 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. The PCR product was cloned into the TA-Cloning vector (available from Invitrogen, San Diego, Calif.) and sequenced using an ABI Prism™ Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism™ dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). The PCR product was found to contain the complete full-length feline caspase-1 except for the primer region which was based on the human caspase-1 sequence. The nucleotide sequence of the coding strand of this PCR product is represented herein as nFeCasp-1<sub>1233</sub> with a SEQ ID NO: 14, and its complement is denoted by SEQ ID NO:16. Translation of SEQ ID NO:14 suggests that nucleic acid molecule nFeCasp-1<sub>1233</sub> encodes a full-length nFeCasp-1<sub>1233</sub> protein, of about 410 amino acids, denoted herein as PFeCasp-1<sub>410</sub>, the amino acid sequence of which is presented in SEQ ID NO:15, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 of SEQ ID NO:14 and a stop codon spanning from nucleotide 408 through nucleotide 410 of SEQ ID NO:14.

Additional primers were made based on the feline caspase-1 sequence of nFeCasp-1<sub>1233</sub> in order to obtain two nucleic acid molecules spanning the 5' and 3' end of the feline caspase-1 open reading frame. Two feline caspase-1 nucleic acid molecules were generated using feline caspase-1 specific primers in combination with cDNA library vector specific primers. The sequence of the vector forward primer (T3 primer) was 5' GCCAAGCTCG AAATTAACCC TACTAAAGG 3' (SEQ ID NO:72), and that of the vector reverse primer (T7 primer) was 5' CGACGGC-CAG TGAATTGTAA TACGACTC 3' (SEQ ID NO:73). The sequence of the feline caspase-1-specific forward primer (Casp 271 Forward) was 5' TCAAGCCCAC AATCTGGAAA TTCTCA 3'(SEQ ID NO:82) and the sequence of the feline caspase-1-specific reverse primer (Casp 895 Reverse) was 5' CTGGAGAGTC ACTGAT-CAAC AGTTCC 3'(SEQ ID NO:83). The first PCR amplification was done using T3 primer and Casp 895 Reverse primer and the second PCR amplification was done using Casp 271 Forward primer and T7 primer. The PCR profile for both reactions was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 45 seconds, then 52° C. for 45 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. The PCR products from both reactions that were greater than or equal to 1 kb were gel purified and cloned into the TA-Cloning vector (available from Invitrogen) and the nucleic acid molecules were sequenced using an ABI Prism™ Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism™ dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). The nucleic acid molecules obtained from these two PCR products represented two

52

nucleic acid molecules of feline caspase-1. The region of the first PCR amplification was sequenced and found to contain 527 nucleotides denoted herein as nFeCasp-1-N<sub>527</sub> (5'-end partial clone) with a coding strand of SEQ ID NO: 17, and a complementary strand of SEQ ID NO: 19. The region of the second PCR amplification was sequenced and found to contain 500 nucleotides denoted here as nFeCasp-1-C<sub>500</sub> (3'-end partial clone) with a coding strand of SEQ ID NO: 20, and a complementary strand of SEQ ID NO: 22. Translation of SEQ ID NO:17 suggests that nucleic acid molecule nFeCasp-1-N<sub>527</sub> encodes an N-terminal portion of PFeCasp-1-N protein, of about 169 amino acids, denoted herein as PFeCasp-1-N<sub>169</sub>, the amino acid sequence of which is presented in SEQ ID NO:18, assuming an open reading frame having an initiation codon spanning from nucleotide 18 through nucleotide 20 of SEQ ID NO:17 and a stop codon spanning from nucleotide 522 through nucleotide 524 of SEQ ID NO:17. Translation of SEQ ID NO:20 suggests that nucleic acid molecule nFeCasp-1-C<sub>500</sub> encodes an C-terminal portion of PFeCasp-1-C protein, of about 120 amino acids, denoted herein as PFeCasp-1-C<sub>120</sub>, the amino acid sequence of which is presented in SEQ ID NO:21, assuming an open reading frame having an initiation codon spanning from nucleotide 3 through nucleotide 5 of SEQ ID NO:20 and a stop codon spanning from nucleotide 360 through nucleotide 362 of SEQ ID NO:20

Based on the sequence data obtained from nucleic acid molecules nFeCasp-1-N<sub>527</sub> and nFeCasp-1-C<sub>500</sub>, two new primers were made to isolate a cDNA encoding full-length feline caspase-1. The feline caspase-1 full-length forward primer (CaspBamKozFor) sequence was 5' ACAAGGATCC ACCATGGCCG ACAAGGATCT GAAGGG 3' (SEQ ID NO:84) and feline caspase-1 full-length reverse primer (CaspXbaRev) sequence was 5' CGCCTCTAGA CCT-CAATTGC CAGGGAAGAG ATAGAAGTA 3' (SEQ ID NO:85). The PCR amplification was performed using these two primers and Amplitaq DNA polymerase™ (available from PE Applied Biosystems Inc.) and an aliquot of the cDNA mitogen library prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours. The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 45 seconds, then 52° C. for 45 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. The PCR product was cloned into the TA-Cloning vector (available from Invitrogen) and the nucleic acid molecule inserts were sequenced using an ABI Prism™ Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism™ dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). This PCR product the FeCaspase-1 containing region of which is denoted nFeCasp-1<sub>1230</sub> was found to encode the a full-length feline caspase-1 protein. The nucleotide sequence of the coding strand of nFeCasp-1<sub>1230</sub> is represented herein as SEQ ID NO 23, and its complement is denoted by SEQ ID NO:25. Translation of the open reading frame in SEQ ID NO:23, denoted here as nFeCasp-1<sub>1230</sub>, the coding strand of which is denoted SEQ ID NO:25 suggests that feline caspase-1 encodes a protein containing 410 amino acids, referred to herein as PFeCasp-1<sub>410</sub>, with a SEQ ID NO:24. The nucleic acid sequence encoding the protein assumes an open reading frame in which the first codon spans from nucleotide 1 through 3 of SEQ ID NO:23, and the last codon spans from nucleotide 1228 nucleotide 1230 of SEQ ID NO:23. The encoded protein has a predicted molecular weight of about 45.5 kDa. The feline caspase-1

protein is 9 amino acids longer than mouse and rat caspase-1 proteins, 6 amino acids longer than dog and human caspase-1 proteins, and 5 amino acids longer than horse caspase-1 proteins. Sequence analysis was performed using DNAsis™, available from Hitachi Software, San Bruno, Calif. using the alignment settings of: gap penalty set at 5, k-tuple set at 3, number of top diagonals set at 5, window size set at 5, fixed gap penalty set at 10 and floating gap penalty set at 10.

## EXAMPLE 3

This example describes the isolation and sequencing of nucleic acid molecules encoding feline IL-12 single chain proteins of the present invention.

A. A pBluescript-Linker plasmid was constructed as follows: Two complementary oligonucleotides, 60 nucleotides in length were synthesized. The oligonucleotides were allowed to hybridize to each other in solution producing a double stranded DNA fragment that would serve as a linker between the cDNAs encoding the p40 and p35 subunits of feline IL-12. The sequence of the sense linker was 5' CTGCAGTGGT GCGGGTGGCG GCGGATCTAG AAAGTGGCA ACCCCTACTC CATCCCCGGG 3' (SEQ ID NO:83) and the sequence of the antisense linker was 5' CCCGGGGATG GAGTAGGGGT TGGCAAGTTT CTA-GATCCGC CGCCACCGCC ACCACTGCAG 3' (SEQ ID NO:84). Equimolar amounts of sense linker and antisense linker were mixed and heated to 95° C. for 10 minutes in a heat block. The heat block containing the samples was removed from the heat source and allowed to cool to room temperature slowly, over a period of 4 hours. Then the hybridized oligonucleotides were digested with PstI and SmaI restriction enzymes (available from New England Biolabs, Beverly, Mass.) and ligated into pBluescript SK<sup>+</sup> vector (available from Stratagene, La Jolla, Calif.) digested with the same restriction enzymes to produce pBluescript-Linker plasmid. The presence of the linker in the ligated pBluescript-Linker plasmid was confirmed by sequencing conducted as described in Example 1. The pBluescript-Linker plasmid contained DNA coding for the following elements: (1) the last two C-terminal amino acid residues of the p40 subunit (i.e. C,S); (2) the seven amino acid residues of the linker (i.e. GGGGGGS) (SEQ ID NO:110); and (3) the first ten N-terminal amino acid residues of the mature p35 subunit mature protein (i.e. RNLPTPTPSP) (SEQ ID NO:111).

B. Feline IL-12 p40 nucleic acid molecule subunit was isolated as follows: A cDNA mitogen library was prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours as previously described in Example 1. An aliquot of this library was used as a template to isolate a feline IL-12 p40 nucleic acid molecule subunit by polymerase chain reaction (PCR). The PCR amplification was performed using Amplitaq DNA polymerase™ (available from PE Applied Biosystems Inc.). The sequence of the forward primer was 5' ATGCATCCTC AGCAGTTGGT CATCGCCT 3' (SEQ ID NO:85), and that of the reverse primer was 5' TGCAGGACAC GGATGCCAGTTGCT 3' (SEQ ID NO:86). The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 45 seconds, then 50° C. for 45 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. PCR products were cloned into the TA-Cloning vector (available from Invitrogen) and the nucleic acid molecule inserts were sequenced as described in Example 1. One of the sequenced PCR products

contained 985 nucleotides and was denoted herein as nFeIL-12 p40-N<sub>985</sub> with a coding strand of SEQ ID NO:55, and a complementary strand of SEQ ID NO:57. Translation of SEQ ID NO:55 suggests that nucleic acid molecule nFeIL-12p40-N<sub>985</sub> encodes an N-terminal portion of PFeIL-12p40-N protein, of about 328 amino acids, denoted herein as PFe IL-12p40-N<sub>328</sub>, the amino acid sequence of which is presented in SEQ ID NO:56, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 of SEQ ID NO:55 and a stop codon spanning from nucleotide 982 through nucleotide 984 of SEQ ID NO:55.

This nucleic acid molecule was used as a template for a subsequent PCR reaction to obtain a full-length nucleic acid molecule. The PCR amplification was performed using Amplitaq DNA polymerase™ (PE Applied Biosystems Inc, Foster City, Calif.). The sequence of the forward primer was 5' ACAGGTACCA TGCATCCTCA GCAGTTGGTC ATCGCCT 3' (SEQ ID NO:87), and that of the reverse primer was 5' CTAAGTGCAG GACACGGATG CCCAG 3' (SEQ ID NO:88). The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 35 cycles of the following: 94° C. for 30 seconds, then 50° C. for 30 seconds, then 72° C. for 90 seconds; followed by a final extension at 72° C. for 7 minutes. This PCR product, the Fe IL-12p40 single chain subunit containing region of which is denoted nFeIL-12 p40<sub>987</sub> was found to encode a full-length feline IL-12 p40 single chain subunit protein. The nucleotide sequence of the coding strand of nFeIL-12 p40<sub>987</sub> is represented herein as SEQ ID NO:29, and its complementary strand is denoted by SEQ ID NO:31. Translation of SEQ ID NO:29 suggests that nucleic acid molecule nFeIL-12p40<sub>987</sub> encodes a full-length PFeIL-12p40 protein of about 329 amino acids, denoted herein as PFe IL-12p40<sub>329</sub>, the amino acid sequence of which is presented in SEQ ID NO:30, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 of SEQ ID NO:29 and a stop codon spanning from nucleotide 985 through nucleotide 987 of SEQ ID NO:29. This PCR product was digested with Kpn I and Pst I restriction enzymes (available from New England Biolabs) and cloned into the pBluescript-Linker plasmid described in Example 3A. The resultant recombinant molecule is referred to as fep40-linker plasmid. There is a putative cleavage site on SEQ ID NO:30, yielding the coding region for a mature (i.e. lacking a signal or leader sequence) nFeIL-12p40<sub>921</sub>, denoted herein as SEQ ID NO:26, with the complement denoted SEQ ID NO:28. Translation of SEQ ID NO:26 yields a mature IL-12 p40 protein denoted PFeIL-12p40<sub>307</sub>, also denoted herein as SEQ ID NO:27.

C. A Feline IL-12 p35 nucleic acid molecule subunit was isolated as follows: A cDNA mitogen library was prepared from cat peripheral blood lymphocytes stimulated with ConA for 4 hours as previously described in Example 1. An aliquot of this library was used as a template to isolate feline IL-12 p35 subunit by polymerase chain reaction (PCR). The PCR amplification was performed using Amplitaq DNA polymerase™ (PE Applied Biosystems Inc, Foster City, Calif.). The sequence of the forward primer was 5' ATGT-GCCCGC CGCGTGGCC 3' (SEQ ID NO:89), and that of the reverse primer was 5' CTAGGAAGCA TTCAGATAGC TCATCAT 3' (SEQ ID NO:90). The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 43 cycles of the following: 94° C. for 45 seconds, then 50° C. for 45 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. PCR products were cloned into the TA-Cloning vector (available from

Invitrogen) and the nucleic acid molecules were sequenced as described in Example 1. One of the sequenced PCR products contained 666 nucleotides and was denoted herein as nFeIL-12-p35<sub>666</sub> with a coding strand of SEQ ID NO:32, and a complementary strand of SEQ ID NO:34. Translation of SEQ ID NO:32 suggests that nucleic acid molecule nFeIL-12p35<sub>666</sub> encodes a full-length PFeIL-12p35 protein of about 222 amino acids, denoted herein as PFeIL-12p35<sub>222</sub>, the amino acid sequence of which is presented in SEQ ID NO:33, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 of SEQ ID NO:32 and a stop codon spanning from nucleotide 664 through nucleotide 666 of SEQ ID NO:32. There is a putative cleavage site on SEQ ID NO:33, yielding the coding region for a mature (i.e. lacking a signal or leader sequence) nFeIL-12p35<sub>591</sub>, denoted herein as SEQ ID NO:35, with the complement denoted SEQ ID NO:37. Translation of SEQ ID NO:26 yields a mature IL-12 p35 protein denoted PFeIL-12p35<sub>197</sub>, also denoted herein as SEQ ID NO:36. SEQ ID NO:26 was digested with Sma I and Not I restriction enzymes (available from New England Biolabs) and cloned into the fep40-linker plasmid described in Example 3B digested with the same enzymes. The resultant recombinant molecule is referred to as fep40-linker-p35 mature plasmid.

D. The fep40-linker-p35 mature plasmid contained a nucleic acid molecule encoding a feline IL-12 single chain protein of the present invention inserted into the Kpn I and Not I sites of the pBluescript backbone. Nucleic acid molecule nFeIL-12<sub>599</sub> was sequenced as described in Example 1. The nucleotide sequence of the coding strand of nFeIL-12<sub>1599</sub> is represented herein as SEQ ID NO:38, and that of the complementary strand is SEQ ID NO:40. Translation of the open reading frame in SEQ ID NO:38, suggests that nFeIL-12<sub>1599</sub> encodes a protein containing 533 amino acids, referred to herein as pFeIL-12<sub>533</sub>, with an amino acid sequence denoted by SEQ ID NO:39. The nucleic acid sequence encoding the protein assumes an open reading frame in which the first codon spans from nucleotide 1 through 3 of SEQ ID NO:38 and the last codon spans from nucleotide 1597 nucleotide 1599 of SEQ ID NO:38. The encoded protein has a predicted molecular weight of about 59.2 kDa. The putative signal peptide cleavage site is between amino acid positions 22 and 23 of the p40 subunit. Nucleic acid molecule nFeIL-12<sub>1533</sub>, which encodes the mature protein contains a coding strand with SEQ ID NO:43, and a complementary strand with SEQ ID NO:45. The amino acid sequence of the mature protein, denoted herein as PFeIL-12<sub>511</sub> is SEQ ID NO:44 and the mature protein has a predicted molecular weight of about 56.8 kDa.

Chinese hamster ovary (CHO) cells (available from ATCC, Rockville, Md.) were transiently transfected with fep40-linker-p35 mature plasmid (containing SEQ ID NO:38) using techniques known to those skilled in the art, cell pellets and supernatants were harvested after 48 hrs. Western analysis was performed on the cell pellets and supernatant samples using a polyclonal antibody against human IL-12 (available from Biosource International, Camarillo, Calif.). A faint band of the expected size (about 59.2 kDa) was detected in the cell pellet and in the supernatant, indicating that IL-12 is produced by this construct at detectable levels.

#### EXAMPLE 4

This example describes the isolation and sequencing of nucleic acid molecules encoding canine IL-12 single chain proteins of the present invention.

A. A canine IL-12 p35 nucleic acid molecule subunit was isolated as follows: A cDNA mitogen library was prepared from canine peripheral blood lymphocytes (PBLs) stimulated with ConA for 4 hours as described in Example 1. Recombinant phage containing DNA encoding the p35 subunit were identified by nucleic acid hybridization using a P<sup>32</sup> radiolabeled probe. The p35 probe (nCaIL-12p35TA) was generated by PCR of total RNA, prepared from ConA-stimulated PBLs in the following manner. The sequence of the forward primer was 5' CCATCCTGGT CCTGCTAAG C 3' (SEQ ID NO:93) and the sequence of the reverse primer was 5' CCATCTGGTA CATCTTCAAG TC 3' (SEQ ID NO:94). PCR amplification was performed using Amplitaq DNA polymerase<sup>TM</sup> (available from PE Applied Biosystems Inc.) using the following profile: 95° C. for 2 minutes; then 30 cycles of 95° C. for 1 minute, 55° C. for 1 minute, and 72° C. for 1 minute; and a final extension at 72° C. for 10 minutes. The amplified DNA fragment was purified with Qiagen gel purification kit, available from Qiagen, La Jolla, Calif.) and PCR products were cloned into the TA cloning vector (available from Invitrogen Corporation, Carlsbad, Calif.), and the resulting clones were sequenced using an ABI Prism<sup>TM</sup> Model 377 Automatic DNA Sequencer (available from Perkin-Elmer Applied Biosystems Inc., Foster City, Calif.). DNA sequencing reactions were performed using Prism<sup>TM</sup> dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.). Phage DNA was permanently cross-linked to the nitrocellulose sheets using a Stratalinker<sup>®</sup> UV crosslinker (available from Stratagene). The plaque lifts were pre-hybridized in a solution of 6×SSC (20×SSC is 3.0 M NaCl and 0.3 M sodium citrate), 5× Denhardt's solution (50×SSC is 0.01 grams/milliliter Ficoll, type 400; 0.01 g/ml polyvinylpyrrolidone; and 0.01 g/ml bovine serum albumin, fraction V, all available from Sigma, St. Louis, Mo.), 0.5% sodium dodecyl sulfate (SDS), and 100 micrograms/ml denatured salmon sperm for 2 hours at 68° C. Denatured, radiolabeled probe was added to the pre-hybridization solution at a concentration of 1×10<sup>6</sup> cpm/ml and the hybridization continued for 18–24 hours at 68° C. Nonspecifically bound and unbound probe was removed by washing two times in 2×SSC with 0.1% SDS, 30 minutes each at 68° C. and one time in 1×SSC with 0.1% SDS, 60 minutes at 68° C. The hybridized plaque lifts were exposed to Kodak x-ray film for approximately 18 hours. Positive phage were plaque purified three times using the following hybridization protocol: phage plaques grown in solid top agar were lifted onto pure nitrocellulose sheets (available from Schleicher & Schuell, Keene, N.H.) then denatured and neutralized by soaking the sheets in 0.5 N NaOH/1.5 M NaCl, followed by 0.5 M Tris-HCl pH7.4/1.5 M NaCl. pBluescript plasmid, containing a cDNA encoding the full-length canine IL-12 p35 subunit, was excised from plaque purified phage using the ExAssist<sup>TM</sup> helper phage (available from Stratagene) following the manufacturers' instructions. The nucleotide sequence of that cDNA, denoted herein as nCa IL-12 p35<sub>1455</sub> was verified by sequencing as described in Example 1. The nucleic acid sequence of the coding strand of nCaIL-12p35<sub>1455</sub> represented as SEQ ID NO: 104, and its complementary strand is SEQ ID NO:106. Translation of SEQ ID NO:104 suggests that nucleic acid molecule nCaIL-12p35<sub>1455</sub> encodes an N-terminal portion of PCaIL-12p35 protein, of about 222 amino acids, denoted herein as PCa IL-12p35<sub>222</sub>, the amino acid sequence of which is presented in SEQ ID NO:105, assuming an open reading frame having an initiation codon spanning from nucleotide 232 through nucleotide

234 of SEQ ID NO:104 and a stop codon spanning from nucleotide 895 through nucleotide 897 of SEQ ID NO:104.

Nucleic acid molecule nCaIL-12p35<sub>1455</sub> was used as the template in PCR to obtain the coding region of the full-length form of canine IL-12 p35 subunit. The sequence of the forward primer was 5' AAAAAACCCG GGTATGTTCC AATGTTTCAA CCACTCCC 3' (SEQ ID NO:95) and the sequence of the reverse primer was 5' GCGGCCGCTC GAGTTAGGAA GAGTTC AAGT AGGACATCAT TCTATTGATG G 3' (SEQ ID NO:96). PCR was performed using Pfu DNA polymerase (available from Stratagene) as follows: 95° C. for 45 seconds; then 25 cycles of 95° C. for 45 seconds, 55° C. for 45 seconds, 72° C. for 1 minute; followed by a final extension at 72° C. for 10 minutes. The PCR product contains the nucleic acid sequence of canine IL-12 p35 subunit which encodes a full-length canine IL-12 p35 subunit protein. The nucleotide sequence of the coding strand of nCaIL-12p35<sub>666</sub> is represented herein as SEQ ID NO:46 and its complementary strand is denoted SEQ ID NO:48. Translation of SEQ ID NO:46 suggests that nucleic acid molecule nCaIL-12p35<sub>666</sub> encodes a mature PCaIL-12p35 single chain protein of about 222 amino acids, denoted herein as PCaIL-12p35<sub>222</sub>, the amino acid sequence of which is presented in SEQ ID NO:47, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 of SEQ ID NO:46 and a stop codon spanning from nucleotide 589 through nucleotide 591 of SEQ ID NO:46. The coding sequence for the mature polypeptide is encoded by SEQ ID NO:46, the coding region for a mature (i.e. lacking a signal or leader sequence) nCaIL-12p35<sub>591</sub>, denoted herein as SEQ ID NO:49, with the complement denoted SEQ ID NO:51. Translation of SEQ ID NO:49 yields a mature IL-12 p35 protein denoted PCaIL-12p35<sub>197</sub>, also denoted herein as SEQ ID NO:50. nCaIL-12p35<sub>591</sub> was digested with SmaI and XhoI restriction endonucleases (available from New England Biolabs) and ligated into the pBluescript-Linker plasmid described in Example 3A digested with the same enzymes. The resultant recombinant molecule is referred to as calinker-p35 mature plasmid.

B. Canine IL-12 p40 nucleic acid molecule subunit was isolated as follows: A cDNA mitogen library was prepared from canine peripheral blood lymphocytes stimulated with ConA for 4 hours as described in Example 1. Recombinant phage containing DNA encoding the p40 subunit were identified by nucleic acid hybridization using a P<sup>32</sup> radiolabeled probe. The p40 probe (nCaIL-12p40TA) was generated by PCR of total RNA, prepared from ConA stimulated PBLs in the following manner. The sequence of the forward primer was 5' CTTAAAGGAA CAGAAAGAAT CC 3' (SEQ ID NO:97) and the sequence of the reverse primer was 5' GGTATCCCA GCTGACCTC 3' (SEQ ID NO:98). PCR amplification was performed using Amplitaq DNA polymerase<sup>TM</sup> (available from PE Applied Biosystems Inc.) using the following profile: 95° C. for 2 minutes; then 30 cycles of 95° C. for 1 minute, 55° C. for 1 minute, and 72° C. for 1 minute; and a final extension at 72° C. for 10 minutes. The amplified DNA-fragment was purified with Qiagen gel purification kit, available from Qiagen, La Jolla, Calif.) and PCR products were cloned into the TA cloning vector (available from Invitrogen Corporation, Carlsbad, Calif.), and the resulting clones were sequenced using an ABI Prism<sup>TM</sup> Model 377 Automatic DNA Sequencer (available from Perkin-Elmer Applied Biosystems Inc., Foster City, Calif.). DNA sequencing reactions were performed using Prism<sup>TM</sup> dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems

Inc.). Phage DNA was permanently cross-linked to the nitrocellulose sheets using a Stratalinker<sup>®</sup> UV crosslinker (available from Stratagene). The plaque lifts were pre-hybridized in a solution of 6×SSC (20×SSC is 3.0 M NaCl and 0.3 M sodium citrate), 5× Denhardt's solution (50×SSC is 0.01 grams/milliliter Ficoll, type 400; 0.01 g/ml polyvinylpyrrolidone; and 0.01 g/ml bovine serum albumin, fraction V, all available from Sigma, St. Louis, Mo.), 0.5% sodium dodecyl sulfate (SDS), and 100 micrograms/ml denatured salmon sperm for 2 hours at 68° C. Denatured, radiolabeled probe was added to the pre-hybridization solution at a concentration of 1×10<sup>6</sup> cpm/ml and the hybridization continued for 18–24 hours at 68° C. Nonspecifically bound and unbound probe was removed by washing two times in 2×SSC with 0.1% SDS, 30 minutes each at 68° C. and one time in 1×SSC with 0.1% SDS, 60 minutes at 68° C. The hybridized plaque lifts were exposed to Kodak x-ray film for approximately 18 hours. Positive phage were plaque purified three times using the following hybridization protocol: phage plaques grown in solid top agar were lifted onto pure nitrocellulose sheets (available from Schleicher & Schuell, Keene, N.H.) then denatured and neutralized by soaking the sheets in 0.5 N NaOH/1.5 M NaCl, followed by 0.5 M Tris-HCl pH7.4/1.5 M NaCl. pBluescript plasmid, containing a cDNA encoding the full-length canine IL-12 p40 subunit, was excised from plaque purified phage using the ExAssist<sup>TM</sup> helper phage (available from Stratagene) following the manufacturers' instructions. The nucleotide sequence of that cDNA, denoted herein as nCaIL-12p40<sub>2267</sub> was verified by sequencing as described in Example 1. The nucleic acid sequence of the coding strand of nCaIL-12p40<sub>2267</sub> represented as SEQ ID NO:107, and its complementary strand is SEQ ID NO:109. Translation of SEQ ID NO:107 suggests that nucleic acid molecule nCaIL-12p40<sub>2267</sub> encodes an PCaIL-12p40 protein, of about 329 amino acids, denoted herein as PCaIL-12p40<sub>329</sub>, the amino acid sequence of which is presented in SEQ ID NO:108, assuming an open reading frame having an initiation codon spanning from nucleotide 154 through nucleotide 156 of SEQ ID NO:107 and a stop codon spanning from nucleotide 1138 through nucleotide 1140 of SEQ ID NO:107. Full length canine IL-12 p40 nucleic acid molecule was isolated as follows: A plasmid containing full-length canine IL-12 p40 nucleic acid molecule subunit (pCaIL-12p40) was used as a template to sub-clone canine IL-12 p40 subunit by polymerase chain reaction (PCR). PCR amplification was performed using Amplitaq DNA polymerase<sup>TM</sup> (available from PE Applied Biosystems Inc.). The sequence of the forward primer (Dog p40 KpnFor) was 5' CATAGGTACC ATGCACCCTC AGCAGTTGGT CATCTCC 3' (SEQ ID NO:99), and that of the reverse primer (Dog p40 NsiRev) was 5' ATCTAAATGC ATGACACAGA TGCCAGTC 3' (SEQ ID NO: 100). The PCR profile was as follows: one initial denaturation step at 94° C. for 5 minutes; then 35 cycles of the following: 94° C. for 30 seconds, then 55° C. for 30 seconds, then 72° C. for 2 minutes; followed by a final extension at 72° C. for 7 minutes. The PCR product contains the nucleic acid sequence of canine IL-12 p40 subunit along with its native signal sequence which encodes a canine full-length IL-12 p40 subunit protein. The nucleotide sequence of the coding strand of nCaIL-12 p40<sub>987</sub> is represented herein as SEQ ID NO:58 and its complementary strand is denoted SEQ ID NO:60. Translation of the open reading frame in SEQ ID NO:58, suggests that canine IL-12 p40 subunit encodes a protein containing 329 amino acids, referred to herein as PCaIL-12 p40<sub>329</sub> with an amino acid sequence denoted by SEQ ID NO:59. The resulting recom-

binant molecule is referred to as cap40-linker plasmid. The cap40-linker plasmid was digested with Kpn I and Pst I restriction enzymes (available from New England Biolabs) to remove the region encoding canine p40 mature protein. The PCR product containing the full-length canine p40 subunit (nCaIL-12 p40<sub>987</sub>) was digested with Kpn I and Nsi I restriction enzymes (available from New England Biolabs) and cloned into this digested plasmid. The coding sequence for the mature canine IL-12 p40 polypeptide is encoded by SEQ ID NO: 52, the coding region for a mature (i.e. lacking a signal or leader sequence) nCaIL-12p40<sub>921</sub>, with the complement denoted SEQ ID NO:53. Translation of SEQ ID NO:52 yields a mature IL-12 p40 protein denoted PCaIL-12p35<sub>307</sub>, also denoted herein as SEQ ID NO:53. nCaIL-12p40<sub>921</sub> was digested with SmaI and XhoI restriction endonucleases (available from New England Biolabs) and ligated into the pBluescript-Linker plasmid described in Example 3A digested with the same enzymes.

The resulting plasmid contains a nucleic acid molecule encoding a canine IL-12 single chain cloned at the Kpn I and Not I site into the pBluescript backbone. The complete canine IL-12 single chain insert was sequenced as described in Example 1. The nucleotide sequence of the coding strand of nCaIL-12-single chain<sub>1599</sub> is represented herein as SEQ ID NO:61, and its complement is denoted by SEQ ID NO:63. Translation of the open reading frame in SEQ ID NO:61, denoted here as nCaIL-12<sub>1599</sub> with a SEQ ID NO:64 suggests that canine IL-12-single chain encodes a protein

containing 533 amino acids, referred to herein as PCaIL-12<sub>533</sub>, with an amino acid sequence denoted by SEQ ID NO:62, assuming an open reading frame in which the first codon spans from nucleotide 1 through 3 of SEQ ID NO:61 and the last codon spans from nucleotide 1597 nucleotide 1599 of SEQ ID NO:61. The encoded protein has a predicted molecular weight of about 59.2 kDa for the precursor protein, and about 56.8 kDa for the mature protein. The putative signal peptide cleavage site is between amino acid positions 22 and 23 of the canine p40 subunit protein. Nucleic acid molecule nCaIL-12<sub>1533</sub>, which encodes the mature protein contains a coding strand with SEQ ID NO:66, and a complementary strand with SEQ ID NO:68. The amino acid sequence of the mature protein, denoted herein as pCaIL-12<sub>511</sub>, is SEQ ID NO:67 and the mature protein has a predicted molecular weight of about 56.8 kDa.

Chinese hamster ovary (CHO) cells (available from ATCC, Rockville, Md.) were transiently transfected with cap40-linker-p35 mature plasmid using techniques known to those skilled in the art and cell pellets and supernatants were harvested after 48 hrs. Western analysis was performed on the cell pellets and supernatant samples using a polyclonal antibody against human IL-12 (available from Biosource International, Camarillo, Calif.). A faint band of the expected size (about 59.2 kDa) was detected in the cell pellet and in the supernatant, indicating that IL-12 is produced by this construct at detectable levels.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 111

<210> SEQ ID NO 1

<211> LENGTH: 514

<212> TYPE: DNA

<213> ORGANISM: Felis catus

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (114)..(512)

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (470)..(470)

<223> OTHER INFORMATION: n = unknown at position 470

Xaa = unknown at position 119

<400> SEQUENCE: 1

```

gctaaaggcg ctctgtccac cttctgccat ctacacagct caggaaaaga aaggacctc      60
aaaccttcca gatcccttcc tctcttagga aactattgag cacagggata aag atg      116
                                     Met
                                     1
act gct ata cca gta gat gat tgc atc aac ttt gtg gga atg aaa ttt      164
Thr Ala Ile Pro Val Asp Asp Cys Ile Asn Phe Val Gly Met Lys Phe
      5              10              15
att gac aat aca ctt tac ttt gta gct gac agt gat gaa aac ctg gaa      212
Ile Asp Asn Thr Leu Tyr Phe Val Ala Asp Ser Asp Glu Asn Leu Glu
      20              25              30
aca gat tac ttt ggc aag ctt gaa cat aaa ctc tca atc tta cga aac      260
Thr Asp Tyr Phe Gly Lys Leu Glu His Lys Leu Ser Ile Leu Arg Asn
      35              40              45
ttg aac gac caa gtt ctc ttc att aac cag gga gat caa cct gtg ttt      308
Leu Asn Asp Gln Val Leu Phe Ile Asn Gln Gly Asp Gln Pro Val Phe
      50              55              60              65

```

-continued

```

gag gat atg cct gat tct gac tgt aca gat aat gca ccc cgg act gaa      356
Glu Asp Met Pro Asp Ser Asp Cys Thr Asp Asn Ala Pro Arg Thr Glu
              70              75              80

ttt atc ata tat atg tat aaa gat agc ctc act aga ggt ctg gca gta      404
Phe Ile Ile Tyr Met Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala Val
              85              90              95

acc atc tct gtg aat tat aag acc atg tct act ctc tcc tgt gag aac      452
Thr Ile Ser Val Asn Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu Asn
              100              105              110

aaa att att tcc ttt aan gga atg agt cct cct gag agt atc aat gat      500
Lys Ile Ile Ser Phe Xaa Gly Met Ser Pro Pro Glu Ser Ile Asn Asp
              115              120              125

gaa gga aat gac at                                                    514
Glu Gly Asn Asp
130

```

```

<210> SEQ ID NO 2
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (119)..(119)
<223> OTHER INFORMATION: The 'Xaa' at location 119 stands for Lys, or
Asn.

```

<400> SEQUENCE: 2

```

Met Thr Ala Ile Pro Val Asp Asp Cys Ile Asn Phe Val Gly Met Lys
 1              5              10              15

Phe Ile Asp Asn Thr Leu Tyr Phe Val Ala Asp Ser Asp Glu Asn Leu
 20              25              30

Glu Thr Asp Tyr Phe Gly Lys Leu Glu His Lys Leu Ser Ile Leu Arg
 35              40              45

Asn Leu Asn Asp Gln Val Leu Phe Ile Asn Gln Gly Asp Gln Pro Val
 50              55              60

Phe Glu Asp Met Pro Asp Ser Asp Cys Thr Asp Asn Ala Pro Arg Thr
 65              70              75              80

Glu Phe Ile Ile Tyr Met Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala
 85              90              95

Val Thr Ile Ser Val Asn Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu
 100             105             110

Asn Lys Ile Ile Ser Phe Xaa Gly Met Ser Pro Pro Glu Ser Ile Asn
 115             120             125

Asp Glu Gly Asn Asp
130

```

```

<210> SEQ ID NO 3
<211> LENGTH: 514
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (45)..(45)
<223> OTHER INFORMATION: n = unknown at position 45

```

<400> SEQUENCE: 3

```

atgtcatttc cttcatcatt gatactctca ggaggactca ttcnntaaa ggaataatt      60
ttgttctcac aggagagagt agacatggtc ttataattca cagagatggt tactgccaga      120
cctctagtga ggctatcttt atacatatat atgataaatt cagtccgggg tgcattatct      180
gtacagtcag aatcagggcat atcctcaaac acaggttgat ctccctggtt aatgaagaga      240

```

-continued

---

```

acttgctcgt tcaagtttcg taagattgag agtttatggt caagcttgcc aaagtaatct 300
gtttccaggt ttcatcact gtcagctaca aagtaaagtg tattgtcaat aaatttcatt 360
ccccaaaagt tgatgcaatc atctactggt atagcagtca tctttatccc tgtgctcaat 420
agtttcctaa gagaggaagg gatctggaag gtttgaggtc cctttctttt cctgagctgt 480
gtagatggca gaagtgggca ggagcgcctt tagc 514

```

```

<210> SEQ ID NO 4
<211> LENGTH: 502
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (3)..(464)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (126)..(126)
<223> OTHER INFORMATION: n = unknown at position 126
Xaa = unknown at position 42

```

```

<400> SEQUENCE: 4

```

```

gc aag ctt gaa cat aaa ctc tca atc tta cga aac ttg aac gac caa 47
  Lys Leu Glu His Lys Leu Ser Ile Leu Arg Asn Leu Asn Asp Gln
  1 5 10 15
gtt ctc ttc att aac cag gga gat caa cct gtg ttt gag gat atg cct 95
Val Leu Phe Ile Asn Gln Gly Asp Gln Pro Val Phe Glu Asp Met Pro
  20 25 30
gat tct gac tgt aca gat aat gca ccc cgg nct gaa ttt atc ata tat 143
Asp Ser Asp Cys Thr Asp Asn Ala Pro Arg Xaa Glu Phe Ile Ile Tyr
  35 40 45
atg tat aaa gat agc ctc act aga ggt ctg gca gta acc atc tct gtg 191
Met Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala Val Thr Ile Ser Val
  50 55 60
aat tat aag acc atg tct act ctc tcc tgt gag aac aaa att att tcc 239
Asn Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu Asn Lys Ile Ile Ser
  65 70 75
ttt aag gaa atg agt cct cct gag agt atc aat gat gaa gga aat gac 287
Phe Lys Glu Met Ser Pro Pro Glu Ser Ile Asn Asp Glu Gly Asn Asp
  80 85 90 95
atc ata ttc ttt cag aga agt gtt cca gga cat gat gat aag ata caa 335
Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asp Lys Ile Gln
  100 105 110
ttt gag tct tca ttg tac aag ggg tac ttt cta gct tgt gaa aaa gag 383
Phe Glu Ser Ser Leu Tyr Lys Gly Tyr Phe Leu Ala Cys Glu Lys Glu
  115 120 125
aaa gat ctt ttc aaa ctc att ttg aaa aaa aag gat gaa aat ggg gat 431
Lys Asp Leu Phe Lys Leu Ile Leu Lys Lys Lys Asp Glu Asn Gly Asp
  130 135 140
aag tcc ata atg ttc act gtt caa aac aag aat tagatattaa aattgcataa 484
Lys Ser Ile Met Phe Thr Val Gln Asn Lys Asn
  145 150
tttgaaaaaa aaaaaaaa 502

```

```

<210> SEQ ID NO 5
<211> LENGTH: 154
<212> TYPE: PRT
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (42)..(42)
<223> OTHER INFORMATION: The 'Xaa' at location 42 stands for Thr, Ala,
Pro, or Ser.

```

-continued

<400> SEQUENCE: 5

Lys Leu Glu His Lys Leu Ser Ile Leu Arg Asn Leu Asn Asp Gln Val  
 1 5 10 15  
 Leu Phe Ile Asn Gln Gly Asp Gln Pro Val Phe Glu Asp Met Pro Asp  
 20 25 30  
 Ser Asp Cys Thr Asp Asn Ala Pro Arg Xaa Glu Phe Ile Ile Tyr Met  
 35 40 45  
 Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala Val Thr Ile Ser Val Asn  
 50 55 60  
 Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu Asn Lys Ile Ile Ser Phe  
 65 70 75 80  
 Lys Glu Met Ser Pro Pro Glu Ser Ile Asn Asp Glu Gly Asn Asp Ile  
 85 90 95  
 Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asp Lys Ile Gln Phe  
 100 105 110  
 Glu Ser Ser Leu Tyr Lys Gly Tyr Phe Leu Ala Cys Glu Lys Glu Lys  
 115 120 125  
 Asp Leu Phe Lys Leu Ile Leu Lys Lys Lys Asp Glu Asn Gly Asp Lys  
 130 135 140  
 Ser Ile Met Phe Thr Val Gln Asn Lys Asn  
 145 150

<210> SEQ ID NO 6  
 <211> LENGTH: 502  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (377)..(377)  
 <223> OTHER INFORMATION: n = unknown at position 377

<400> SEQUENCE: 6

tttttttttt ttttcaaatt atgcaathtt aatatcctaat tcttgttttg aacagtgaac 60  
 attatggact tatccccatt ttcctccttt tttttcaaaa tgagtttgaa aagatctttc 120  
 tctttttcac aagctagaaa gtaccctctg tacaatgaag actcaaattg tatcttatca 180  
 tcatgtcctg gaacacttct ctgaaagaat atgatgtcat ttccttcatc attgatactc 240  
 tcaggaggac tcatttcctt aaaggaaata atttgttct cacaggagag agtagacatg 300  
 gtcttataat tcacagagat ggttactgcc agacctctag tgaggctatc tttatacata 360  
 tatatgataa attcagnccg ggggtgcatta tctgtacagt cagaatcagg catatcctca 420  
 aacacaggtt gatctccctg gttaatgaag agaacttggc cgttcaagtt tcgtaagatt 480  
 gagagtttat gttcaagctt gc 502

<210> SEQ ID NO 7  
 <211> LENGTH: 607  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (24)..(599)

<400> SEQUENCE: 7

aactattgag cacagggata aag atg act gct ata cca gta gat gat tgc atc 53  
 Met Thr Ala Ile Pro Val Asp Asp Cys Ile  
 1 5 10

-continued

---

```

aac ttt gtg gga atg aaa ttt att gac aat aca ctt tac ttt gta gct      101
Asn Phe Val Gly Met Lys Phe Ile Asp Asn Thr Leu Tyr Phe Val Ala
                      15                      20                      25

gac agt gat gaa aac ctg gaa aca gat tac ttt ggc aag ctt gaa cat      149
Asp Ser Asp Glu Asn Leu Glu Thr Asp Tyr Phe Gly Lys Leu Glu His
                      30                      35                      40

aaa ctc tca atc tta cga aac ttg aac gac caa gtt ctc ttc att aac      197
Lys Leu Ser Ile Leu Arg Asn Leu Asn Asp Gln Val Leu Phe Ile Asn
                      45                      50                      55

cag gga gat caa cct gtg ttt gag gat atg cct gat tct gac tgt aca      245
Gln Gly Asp Gln Pro Val Phe Glu Asp Met Pro Asp Ser Asp Cys Thr
                      60                      65                      70

gat aat gca ccc cgg act gaa ttt atc ata tat atg tat aaa gat agc      293
Asp Asn Ala Pro Arg Thr Glu Phe Ile Ile Tyr Met Tyr Lys Asp Ser
75                      80                      85                      90

ctc act aga ggt ctg gca gta acc atc tct gtg aat tat aag acc atg      341
Leu Thr Arg Gly Leu Ala Val Thr Ile Ser Val Asn Tyr Lys Thr Met
                      95                      100                      105

tct act ctc tcc tgt gag aac aaa att att tcc ttt aag gaa atg agt      389
Ser Thr Leu Ser Cys Glu Asn Lys Ile Ile Ser Phe Lys Glu Met Ser
110                      115                      120

cct cct gag agt atc aat gat gaa gga aat gac atc ata ttc ttt cag      437
Pro Pro Glu Ser Ile Asn Asp Glu Gly Asn Asp Ile Ile Phe Phe Gln
125                      130                      135

aga agt gtt cca gga cat gat gat aag ata caa ttt gag tct tca ttg      485
Arg Ser Val Pro Gly His Asp Asp Lys Ile Gln Phe Glu Ser Ser Leu
140                      145                      150

tac aag ggg tac ttt cta gct tgt gaa aaa gag aaa gat ctt ttc aaa      533
Tyr Lys Gly Tyr Phe Leu Ala Cys Glu Lys Glu Lys Asp Leu Phe Lys
155                      160                      165                      170

ctc att ttg aaa aaa aag gat gaa aat ggg gat aag tcc ata atg ttc      581
Leu Ile Leu Lys Lys Lys Asp Glu Asn Gly Asp Lys Ser Ile Met Phe
175                      180                      185

act gtt caa aac aag aat tagatatt      607
Thr Val Gln Asn Lys Asn
190

```

```

<210> SEQ ID NO 8
<211> LENGTH: 192
<212> TYPE: PRT
<213> ORGANISM: Felis catus

```

<400> SEQUENCE: 8

```

Met Thr Ala Ile Pro Val Asp Asp Cys Ile Asn Phe Val Gly Met Lys
1                      5                      10                      15

Phe Ile Asp Asn Thr Leu Tyr Phe Val Ala Asp Ser Asp Glu Asn Leu
20                      25                      30

Glu Thr Asp Tyr Phe Gly Lys Leu Glu His Lys Leu Ser Ile Leu Arg
35                      40                      45

Asn Leu Asn Asp Gln Val Leu Phe Ile Asn Gln Gly Asp Gln Pro Val
50                      55                      60

Phe Glu Asp Met Pro Asp Ser Asp Cys Thr Asp Asn Ala Pro Arg Thr
65                      70                      75                      80

Glu Phe Ile Ile Tyr Met Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala
85                      90                      95

Val Thr Ile Ser Val Asn Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu
100                      105                      110

Asn Lys Ile Ile Ser Phe Lys Glu Met Ser Pro Pro Glu Ser Ile Asn
115                      120                      125

```

-continued

---

Asp Glu Gly Asn Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His  
 130 135 140

Asp Asp Lys Ile Gln Phe Glu Ser Ser Leu Tyr Lys Gly Tyr Phe Leu  
 145 150 155 160

Ala Cys Glu Lys Glu Lys Asp Leu Phe Lys Leu Ile Leu Lys Lys Lys  
 165 170 175

Asp Glu Asn Gly Asp Lys Ser Ile Met Phe Thr Val Gln Asn Lys Asn  
 180 185 190

<210> SEQ ID NO 9  
 <211> LENGTH: 576  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 9

```
atgactgcta taccagtaga tgattgcatc aactttgtgg gaatgaaatt tattgacaat 60
acactttact ttgtagctga cagtgatgaa aacctgaaa cagattactt tggcaagctt 120
gaacataaac tctcaatctt acgaaacttg aacgaccaag ttctcttcat taaccaggga 180
gatcaacctg tgtttgagga tatgcctgat tctgactgta cagataatgc accccggact 240
gaatttatca tatatatgta taaagatagc ctactagag gtctggcagt aaccatctct 300
gtgaattata agaccatgtc tactctctcc tgtgagaaca aaattatttc cttaaggaa 360
atgagtcctc ctgagagtat caatgatgaa ggaaatgaca tcattattct tcagagaagt 420
gttccaggac atgatgataa gatacaattt gactcttcat tgtacaaggg gtactttcta 480
gcttgtgaaa aagagaaaga tcttttcaaa ctcatcttga aaaaaagga tgaaaatggg 540
gataagtcca taatgttcac tgttcaaac aagaat 576
```

<210> SEQ ID NO 10  
 <211> LENGTH: 607  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 10

```
aatatcfaat tcttgttttg aacagtgaac attatggact tatccccatt ttcaccttt 60
ttttcaaaa tgagtttgaa aagatcttct tctttttcac aagctagaaa gtacccttg 120
tacaatgaag actcaaatg tatcttatca tcatgtctct gaacacttct ctgaaagaat 180
atgatgtcat ttccttcac attgatactc tcaggaggac tcatttcctt aaaggaata 240
atcttgttct cacaggagag agtagacatg gtcttataat tcacagagat ggttactgcc 300
agacctctag tgaggctatc tttatacata tatatgataa attcagtcg ggtgcatta 360
tctgtacagt cagaatcagg catatctca aacacaggt gatctccctg gttaatgaag 420
agaacttggc cgttcaagtt tcgtaagatt gagagtttat gttcaagctt gccaaagtaa 480
tctgtttcca ggttttcac actgtcagct acaaagtaa gtgtattgtc aataaatttc 540
attcccaca agttgatgca atcatctact ggtatagcag tcacttttat cctgtgctc 600
aatagtt 607
```

<210> SEQ ID NO 11  
 <211> LENGTH: 471  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(471)

-continued

<400> SEQUENCE: 11

```

tac ttt ggc aag ctt gaa cat aaa ctc tca atc tta cga aac ttg aac      48
Tyr Phe Gly Lys Leu Glu His Lys Leu Ser Ile Leu Arg Asn Leu Asn
1           5           10           15

gac caa gtt ctc ttc att aac cag gga gat caa cct gtg ttt gag gat      96
Asp Gln Val Leu Phe Ile Asn Gln Gly Asp Gln Pro Val Phe Glu Asp
           20           25           30

atg cct gat tct gac tgt aca gat aat gca ccc cgg act gaa ttt atc      144
Met Pro Asp Ser Asp Cys Thr Asp Asn Ala Pro Arg Thr Glu Phe Ile
           35           40           45

ata tat atg tat aaa gat agc ctc act aga ggt ctg gca gta acc atc      192
Ile Tyr Met Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala Val Thr Ile
           50           55           60

tct gtg aat tat aag acc atg tct act ctc tcc tgt gag aac aaa att      240
Ser Val Asn Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu Asn Lys Ile
65           70           75           80

att tcc ttt aag gaa atg agt cct cct gag agt atc aat gat gaa gga      288
Ile Ser Phe Lys Glu Met Ser Pro Pro Glu Ser Ile Asn Asp Glu Gly
           85           90           95

aat gac atc ata ttc ttt cag aga agt gtt cca gga cat gat gat aag      336
Asn Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asp Lys
           100          105          110

ata caa ttt gag tct tca ttg tac aag ggg tac ttt cta gct tgt gaa      384
Ile Gln Phe Glu Ser Ser Leu Tyr Lys Gly Tyr Phe Leu Ala Cys Glu
           115          120          125

aaa gag aaa gat ctt ttc aaa ctc att ttg aaa aaa aag gat gaa aat      432
Lys Glu Lys Asp Leu Phe Lys Leu Ile Leu Lys Lys Lys Asp Glu Asn
           130          135          140

ggg gat aag tcc ata atg ttc act gtt caa aac aag aat      471
Gly Asp Lys Ser Ile Met Phe Thr Val Gln Asn Lys Asn
145           150           155

```

<210> SEQ ID NO 12  
 <211> LENGTH: 157  
 <212> TYPE: PRT  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 12

```

Tyr Phe Gly Lys Leu Glu His Lys Leu Ser Ile Leu Arg Asn Leu Asn
1           5           10           15

Asp Gln Val Leu Phe Ile Asn Gln Gly Asp Gln Pro Val Phe Glu Asp
           20           25           30

Met Pro Asp Ser Asp Cys Thr Asp Asn Ala Pro Arg Thr Glu Phe Ile
           35           40           45

Ile Tyr Met Tyr Lys Asp Ser Leu Thr Arg Gly Leu Ala Val Thr Ile
           50           55           60

Ser Val Asn Tyr Lys Thr Met Ser Thr Leu Ser Cys Glu Asn Lys Ile
65           70           75           80

Ile Ser Phe Lys Glu Met Ser Pro Pro Glu Ser Ile Asn Asp Glu Gly
           85           90           95

Asn Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asp Lys
           100          105          110

Ile Gln Phe Glu Ser Ser Leu Tyr Lys Gly Tyr Phe Leu Ala Cys Glu
           115          120          125

Lys Glu Lys Asp Leu Phe Lys Leu Ile Leu Lys Lys Lys Asp Glu Asn
           130          135          140

Gly Asp Lys Ser Ile Met Phe Thr Val Gln Asn Lys Asn
145           150           155

```

-continued

---

```

<210> SEQ ID NO 13
<211> LENGTH: 471
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 13

attcttggtt tgaacagtga acattatgga cttatcccca ttttcatcct tttttttcaa      60
aatgagtttg aaaagatcct tctctttttc acaagctaga aagtaccctt tgtacaatga      120
agactcaaat tgtatcttat catcatgtcc tggaacactt ctctgaaaga atatgatgtc      180
atttccttca tcattgatac tctcaggagg actcatttcc ttaaaggaaa taattttggt      240
ctcacaggag agagtagaca tggctttata attcacagag atggttactg ccagacctct      300
agtgaggcta tctttataca tatatatgat aaattcagtc cggggtgcat tatctgtaca      360
gtcagaatca ggcatacctt caaacacagg ttgatctccc tggttaatga agagaacttg      420
gtcgttcaag tttcgttaaga ttgagagttt atgttcaagc ttgccaaagt a              471

<210> SEQ ID NO 14
<211> LENGTH: 1233
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1230)

<400> SEQUENCE: 14

atg gcc gac aag gtc ctg aag gag aag agg aag cag ttc atc aac tca      48
Met Ala Asp Lys Val Leu Lys Glu Lys Arg Lys Gln Phe Ile Asn Ser
1 5 10 15

gtc ggc atg ggg acg gtc aac ggc ttg ctg gat gaa ctc ttt gag aaa      96
Val Gly Met Gly Thr Val Asn Gly Leu Leu Asp Glu Leu Phe Glu Lys
20 25 30

aac gtg ctg aac cag gag gag atg gag aga gta aaa tgt gaa aac gct      144
Asn Val Leu Asn Gln Glu Glu Met Glu Arg Val Lys Cys Glu Asn Ala
35 40 45

acc gtt atg gac aag gcc cga gct ctg atc gac agc gtc ctg cgg aaa      192
Thr Val Met Asp Lys Ala Arg Ala Leu Ile Asp Ser Val Leu Arg Lys
50 55 60

ggg cca cgg gcg tgc cag atc ttt atc tgt cac atc tgt gag gaa gac      240
Gly Pro Arg Ala Cys Gln Ile Phe Ile Cys His Ile Cys Glu Glu Asp
65 70 75 80

acc cac ctt gca gag acg ctg ggg ctc tcc tca agc cca caa tct gga      288
Thr His Leu Ala Glu Thr Leu Gly Leu Ser Ser Ser Pro Gln Ser Gly
85 90 95

aat tct cag aac acc acg gac tct gaa gta gcg ttt cct cct ctt cca      336
Asn Ser Gln Asn Thr Thr Asp Ser Glu Val Ala Phe Pro Pro Leu Pro
100 105 110

gcc agc gtg aat aac atg cct ggg ccg gct gag cca gaa gaa tct gta      384
Ala Ser Val Asn Asn Met Pro Gly Pro Ala Glu Pro Glu Glu Ser Val
115 120 125

gat gct ctc aag ctt tgt cct cgt gaa aac ttc gtg aaa ctg tgt aaa      432
Asp Ala Leu Lys Leu Cys Pro Arg Glu Asn Phe Val Lys Leu Cys Lys
130 135 140

cag agg gct gaa gag atc tac cca ata aag gag aga aag gat cgt act      480
Gln Arg Ala Glu Glu Ile Tyr Pro Ile Lys Glu Arg Lys Asp Arg Thr
145 150 155 160

cgt ctg gct ctc atc ata tgc aat acg acg ttc gat cat ctt tct ctc      528
Arg Leu Ala Leu Ile Ile Cys Asn Thr Thr Phe Asp His Leu Ser Leu
165 170 175

```

-continued

```

agg aag ggg gct gac ctt gac gtt gca ggg atg agg agg ctg ctt aca 576
Arg Lys Gly Ala Asp Leu Asp Val Ala Gly Met Arg Arg Leu Leu Thr
      180                      185                      190

gac ctt ggc tac agt gtg cac ata aaa gag gaa ctc act gct aag gac 624
Asp Leu Gly Tyr Ser Val His Ile Lys Glu Glu Leu Thr Ala Lys Asp
      195                      200                      205

atg gaa tca gag ctg agg gca ttt gct gcc cgt cca gag cac aag tcc 672
Met Glu Ser Glu Leu Arg Ala Phe Ala Ala Arg Pro Glu His Lys Ser
      210                      215                      220

tcg gac agc aca ttc ctg gtg ttc atg tct cat ggc atc ctg agt gga 720
Ser Asp Ser Thr Phe Leu Val Phe Met Ser His Gly Ile Leu Ser Gly
      225                      230                      235                      240

atc tgt ggg acg aag tac agc gct gaa gga gac cca gat gta ttg gct 768
Ile Cys Gly Thr Lys Tyr Ser Ala Glu Gly Asp Pro Asp Val Leu Ala
      245                      250                      255

tat gac acc atc ttc cag att ttc aac aac cgc aac tgc ctt agt cta 816
Tyr Asp Thr Ile Phe Gln Ile Phe Asn Asn Arg Asn Cys Leu Ser Leu
      260                      265                      270

aag gac aag ccc aag gtc atc atc gtc cag gcc tgc aga ggt gaa aat 864
Lys Asp Lys Pro Lys Val Ile Ile Val Gln Ala Cys Arg Gly Glu Asn
      275                      280                      285

ttg ggg gaa ctg ttg atc agt gac tct cca gcg gcc cca atg gac agc 912
Leu Gly Glu Leu Leu Ile Ser Asp Ser Pro Ala Ala Pro Met Asp Ser
      290                      295                      300

act tca cag atg ggt agc agc ctt tca cag gtg ggt gac aac cta gag 960
Thr Ser Gln Met Gly Ser Ser Leu Ser Gln Val Gly Asp Asn Leu Glu
      305                      310                      315                      320

gac gac gcc att tac aag gtc cac gtg gag aag gac ttc atc gct ttc 1008
Asp Asp Ala Ile Tyr Lys Val His Val Glu Lys Asp Phe Ile Ala Phe
      325                      330                      335

tgc tcc tcg acc cca cat cat gtg tct tgg aga gac gtg aac aag gga 1056
Cys Ser Ser Thr Pro His His Val Ser Trp Arg Asp Val Asn Lys Gly
      340                      345                      350

tct ctc ttc att aca caa ctc atc acg tgc ttc caa aag tat tcg tgg 1104
Ser Leu Phe Ile Thr Gln Leu Ile Thr Cys Phe Gln Lys Tyr Ser Trp
      355                      360                      365

tgc ttt cat ctg gag gaa gta ttt cgg aag gta caa cag tca ttt gaa 1152
Cys Phe His Leu Glu Glu Val Phe Arg Lys Val Gln Gln Ser Phe Glu
      370                      375                      380

aaa cca aat gtt aga gcc cag atg ccc acc att gaa cga cta tcc atg 1200
Lys Pro Asn Val Arg Ala Gln Met Pro Thr Ile Glu Arg Leu Ser Met
      385                      390                      395                      400

aca aga tgt ttc tac ctc ttc cca gga cat taa 1233
Thr Arg Cys Phe Tyr Leu Phe Pro Gly His
      405                      410

```

```

<210> SEQ ID NO 15
<211> LENGTH: 410
<212> TYPE: PRT
<213> ORGANISM: Felis catus

```

<400> SEQUENCE: 15

```

Met Ala Asp Lys Val Leu Lys Glu Lys Arg Lys Gln Phe Ile Asn Ser
1      5      10      15

Val Gly Met Gly Thr Val Asn Gly Leu Leu Asp Glu Leu Phe Glu Lys
      20      25      30

Asn Val Leu Asn Gln Glu Glu Met Glu Arg Val Lys Cys Glu Asn Ala
      35      40      45

Thr Val Met Asp Lys Ala Arg Ala Leu Ile Asp Ser Val Leu Arg Lys
      50      55      60

```

-continued

Gly	Pro	Arg	Ala	Cys	Gln	Ile	Phe	Ile	Cys	His	Ile	Cys	Glu	Glu	Asp	65	70	75	80
Thr	His	Leu	Ala	Glu	Thr	Leu	Gly	Leu	Ser	Ser	Ser	Pro	Gln	Ser	Gly	85	90	95	
Asn	Ser	Gln	Asn	Thr	Thr	Asp	Ser	Glu	Val	Ala	Phe	Pro	Pro	Leu	Pro	100	105	110	
Ala	Ser	Val	Asn	Asn	Met	Pro	Gly	Pro	Ala	Glu	Pro	Glu	Glu	Ser	Val	115	120	125	
Asp	Ala	Leu	Lys	Leu	Cys	Pro	Arg	Glu	Asn	Phe	Val	Lys	Leu	Cys	Lys	130	135	140	
Gln	Arg	Ala	Glu	Glu	Ile	Tyr	Pro	Ile	Lys	Glu	Arg	Lys	Asp	Arg	Thr	145	150	155	160
Arg	Leu	Ala	Leu	Ile	Ile	Cys	Asn	Thr	Thr	Phe	Asp	His	Leu	Ser	Leu	165	170	175	
Arg	Lys	Gly	Ala	Asp	Leu	Asp	Val	Ala	Gly	Met	Arg	Arg	Leu	Leu	Thr	180	185	190	
Asp	Leu	Gly	Tyr	Ser	Val	His	Ile	Lys	Glu	Glu	Leu	Thr	Ala	Lys	Asp	195	200	205	
Met	Glu	Ser	Glu	Leu	Arg	Ala	Phe	Ala	Ala	Arg	Pro	Glu	His	Lys	Ser	210	215	220	
Ser	Asp	Ser	Thr	Phe	Leu	Val	Phe	Met	Ser	His	Gly	Ile	Leu	Ser	Gly	225	230	235	240
Ile	Cys	Gly	Thr	Lys	Tyr	Ser	Ala	Glu	Gly	Asp	Pro	Asp	Val	Leu	Ala	245	250	255	
Tyr	Asp	Thr	Ile	Phe	Gln	Ile	Phe	Asn	Asn	Arg	Asn	Cys	Leu	Ser	Leu	260	265	270	
Lys	Asp	Lys	Pro	Lys	Val	Ile	Ile	Val	Gln	Ala	Cys	Arg	Gly	Glu	Asn	275	280	285	
Leu	Gly	Glu	Leu	Leu	Ile	Ser	Asp	Ser	Pro	Ala	Ala	Pro	Met	Asp	Ser	290	295	300	
Thr	Ser	Gln	Met	Gly	Ser	Ser	Leu	Ser	Gln	Val	Gly	Asp	Asn	Leu	Glu	305	310	315	320
Asp	Asp	Ala	Ile	Tyr	Lys	Val	His	Val	Glu	Lys	Asp	Phe	Ile	Ala	Phe	325	330	335	
Cys	Ser	Ser	Thr	Pro	His	His	Val	Ser	Trp	Arg	Asp	Val	Asn	Lys	Gly	340	345	350	
Ser	Leu	Phe	Ile	Thr	Gln	Leu	Ile	Thr	Cys	Phe	Gln	Lys	Tyr	Ser	Trp	355	360	365	
Cys	Phe	His	Leu	Glu	Glu	Val	Phe	Arg	Lys	Val	Gln	Gln	Ser	Phe	Glu	370	375	380	
Lys	Pro	Asn	Val	Arg	Ala	Gln	Met	Pro	Thr	Ile	Glu	Arg	Leu	Ser	Met	385	390	395	400
Thr	Arg	Cys	Phe	Tyr	Leu	Phe	Pro	Gly	His	405	410								

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 1233

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Felis catus

&lt;400&gt; SEQUENCE: 16

ttaatgtcct gggaagaggt agaaacatct tgtcatggat agtcggtcaa tgggtggcat	60
ctgggtccta acatttgggt tttcaaatga ctggtgtacc ttccgaaata ctctctccag	120
atgaaagcac cacgaatact tttggaagca cgtgatgagt tgtgtaatga agagagatcc	180

-continued

```

cttgttcacg tctctccaag acacatgatg tggggctcag gagcagaag cgatgaagtc 240
cttctccacg tggaccttgt aaatggcgtc gtcctctagg ttgtcaccca cctgtgaaag 300
gctgtaccac atctgtgaag tgctgtccat tggggccgct ggagagtcac tgatcaacag 360
ttccccaaa ttttcacctc tgcaggcctg gacgatgatg accttgggct tgtcctttag 420
actaaggcag ttgcggttgt tgaaaatctg gaagatggtg tcataagcca atacatctgg 480
gtctccttca gcgctgtact tegtcccaca gattccactc aggatgccat gagacatgaa 540
caccaggaat gtgctgtccg aggacttgtg ctctggacgg gcagcaaatg ccctcagctc 600
tgattccatg tccttagcag tgagttcctc ttttatgtgc aactgtagc caaggctctgt 660
aagcagcctc tcatccctg caacgtcaag gtcagcccc ttcttgagag aaagatgatc 720
gaacgtcgtg ttgcatatga tgagagccag acgagtacga tcctttctct cctttattgg 780
gtagatctct tcagccctct gtttacacag tttcacgaag tttcacgag gacaaagcct 840
gagagcatct acagattctt ctggctcagc cggcccaggc atgttattca cgctggctgg 900
aagaggagga aacgctactt cagagtccgt ggtgttctga gaatttcag attgtgggct 960
tgaggagagc cccagcgtct ctgcaaggtg ggtgtcttcc tcacagatgt gacagataaa 1020
gatctggcac gccctgggcc ctttccgcag gacgctgtcg atcagagctc gggccttgtc 1080
cataacggta gcgttttcac attttactct ctccatctcc tcctgttca gcacgttttt 1140
ctcaaagagt tcatccagca agccgttgac cgtcccatg ccgactgagt tgatgaactg 1200
cttctcttcc tccttcagga ccttgcgcg cat 1233

```

```

<210> SEQ ID NO 17
<211> LENGTH: 526
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (18)..(524)

```

<400> SEQUENCE: 17

```

ggcacgagca aaaagcc atg gcc gac aag gat ctg aag ggc aag agg aag 50
Met Ala Asp Lys Asp Leu Lys Gly Lys Arg Lys
1 5 10
cag ttc atc aac tca gtc ggc atg ggg acg gtc aac ggc ttg ctg gat 98
Gln Phe Ile Asn Ser Val Gly Met Gly Thr Val Asn Gly Leu Leu Asp
15 20 25
gaa ctc ttt gag aaa aac gtg ctg aac cag gag gag atg gag aga gta 146
Glu Leu Phe Glu Lys Asn Val Leu Asn Gln Glu Glu Met Glu Arg Val
30 35 40
aaa tgt gaa aac gct acc gtt atg gac aag gcc cga gct ctg atc gac 194
Lys Cys Glu Asn Ala Thr Val Met Asp Lys Ala Arg Ala Leu Ile Asp
45 50 55
agc gtc ctg cgg aaa ggg cca cgg gcg tgc cag atc ttt atc tgt cac 242
Ser Val Leu Arg Lys Gly Pro Arg Ala Cys Gln Ile Phe Ile Cys His
60 65 70 75
atc tgt gag gaa gac acc cac ctt gca gag acg ctg ggg ctc tcc tca 290
Ile Cys Glu Glu Asp Thr His Leu Ala Glu Thr Leu Gly Leu Ser Ser
80 85 90
agc cca caa tct gga aat tct cag aac acc acg gac tct gaa gta gcg 338
Ser Pro Gln Ser Gly Asn Ser Gln Asn Thr Thr Asp Ser Glu Val Ala
95 100 105
ttt cct cct ctt cca gcc agc gtg aat aac atg cct ggg ccg gct gag 386
Phe Pro Pro Leu Pro Ala Ser Val Asn Asn Met Pro Gly Pro Ala Glu
110 115 120

```

-continued

```

cca gaa gaa tct gta gat gct ctc aag ctt tgt cct cgt gaa aac ttc      434
Pro Glu Glu Ser Val Asp Ala Leu Lys Leu Cys Pro Arg Glu Asn Phe
    125                               130                               135

gtg aaa ctg tgt aaa cag agg gct gaa gag atc tac cca ata aag gag      482
Val Lys Leu Cys Lys Gln Arg Ala Glu Glu Ile Tyr Pro Ile Lys Glu
    140                               145                               150                               155

aga aag gat cgt act cgt ctg gct ctc atc ata tgc aat acg ac          526
Arg Lys Asp Arg Thr Arg Leu Ala Leu Ile Ile Cys Asn Thr
    160                               165
    
```

```

<210> SEQ ID NO 18
<211> LENGTH: 169
<212> TYPE: PRT
<213> ORGANISM: Felis catus
    
```

<400> SEQUENCE: 18

```

Met Ala Asp Lys Asp Leu Lys Gly Lys Arg Lys Gln Phe Ile Asn Ser
 1                               5                               10                               15

Val Gly Met Gly Thr Val Asn Gly Leu Leu Asp Glu Leu Phe Glu Lys
    20                               25                               30

Asn Val Leu Asn Gln Glu Glu Met Glu Arg Val Lys Cys Glu Asn Ala
    35                               40                               45

Thr Val Met Asp Lys Ala Arg Ala Leu Ile Asp Ser Val Leu Arg Lys
    50                               55                               60

Gly Pro Arg Ala Cys Gln Ile Phe Ile Cys His Ile Cys Glu Glu Asp
 65                               70                               75                               80

Thr His Leu Ala Glu Thr Leu Gly Leu Ser Ser Ser Pro Gln Ser Gly
    85                               90                               95

Asn Ser Gln Asn Thr Thr Asp Ser Glu Val Ala Phe Pro Pro Leu Pro
    100                              105                              110

Ala Ser Val Asn Asn Met Pro Gly Pro Ala Glu Pro Glu Glu Ser Val
    115                              120                              125

Asp Ala Leu Lys Leu Cys Pro Arg Glu Asn Phe Val Lys Leu Cys Lys
    130                              135                              140

Gln Arg Ala Glu Glu Ile Tyr Pro Ile Lys Glu Arg Lys Asp Arg Thr
 145                              150                              155                              160

Arg Leu Ala Leu Ile Ile Cys Asn Thr
    165
    
```

```

<210> SEQ ID NO 19
<211> LENGTH: 526
<212> TYPE: DNA
<213> ORGANISM: Felis catus
    
```

<400> SEQUENCE: 19

```

gtcgtattgc atatgatgag agccagacga gtacgatcct ttctctcctt tattgggtag      60
atctcttcag ccctctgttt acacagtttc acgaagtttt cagcaggaca aagcttgaga      120
gcatctacag attcttctgg ctcagccggc ccaggcatgt tattcacgct ggctggaaga      180
ggaggaaaacg ctacttcaga gtccgtggtg ttctgagaat ttccagattg tgggcttgag      240
gagagcccca gcgtctctgc aaggtgggtg tcttctctcac agatgtgaca gataaagatc      300
tggcacgccc gtggcccttt ccgcaggacg ctgtcgatca gagctcgggc cttgtccata      360
acggtagcgt tttcacattt tactctctcc atctctctct gggttcagcac gtttttctca      420
aagagttcat ccagcaagcc gttgaccgtc cccatgccga ctgagttgat gaactgcttc      480
ctcttgccct tcagatcctt gtcggccatg gctttttgct cgtgcc          526
    
```

-continued

---

```

<210> SEQ ID NO 20
<211> LENGTH: 500
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (3)..(362)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (473)..(473)
<223> OTHER INFORMATION: n = unknown at position 473

<400> SEQUENCE: 20

gg gaa ctg ttg atc agt gac tct cca gcg gcc cca atg gac agc act      47
  Glu Leu Leu Ile Ser Asp Ser Pro Ala Ala Pro Met Asp Ser Thr
    1             5             10             15

tca cag atg ggt agc agc ctt tca cag gtg ggt gac aac cta gag gac      95
  Ser Gln Met Gly Ser Ser Leu Ser Gln Val Gly Asp Asn Leu Glu Asp
    20             25             30

gac gcc att tac aag gtc cac gtg gag aag gac ttc atc gct ttc tgc     143
  Asp Ala Ile Tyr Lys Val His Val Glu Lys Asp Phe Ile Ala Phe Cys
    35             40             45

tcc tcg acc cca cat cat gtg tct tgg aga gac gtg aac aag gga tct     191
  Ser Ser Thr Pro His His Val Ser Trp Arg Asp Val Asn Lys Gly Ser
    50             55             60

ctc ttc att aca caa ctc atc acg tgc ttc caa aag tat tcg tgg tgc     239
  Leu Phe Ile Thr Gln Leu Ile Thr Cys Phe Gln Lys Tyr Ser Trp Cys
    65             70             75

ttt cat ctg gag gaa gta ttt cgg aag gta caa cag tca ttt gaa aaa     287
  Phe His Leu Glu Glu Val Phe Arg Lys Val Gln Gln Ser Phe Glu Lys
    80             85             90             95

cca aat gtt aga gcc cag atg ccc acc att gaa cga cta tcc atg aca     335
  Pro Asn Val Arg Ala Gln Met Pro Thr Ile Glu Arg Leu Ser Met Thr
    100            105            110

aga tac ttc tat ctc ttc cct ggc aat tgaaaatagc aatcatgggc         382
  Arg Tyr Phe Tyr Leu Phe Pro Gly Asn
    115            120

agtcagccc ttcttgacca acttgaaaa gtaccttagc tagcacaaca cactcattta   442

acgtttggtgta tctcaataaa aatgaaaaca nctaaaaaaaa aaaaaaaaaa aaaaaaaa   500

```

```

<210> SEQ ID NO 21
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Felis catus

<400> SEQUENCE: 21

Glu Leu Leu Ile Ser Asp Ser Pro Ala Ala Pro Met Asp Ser Thr Ser
 1             5             10             15

Gln Met Gly Ser Ser Leu Ser Gln Val Gly Asp Asn Leu Glu Asp Asp
 20             25             30

Ala Ile Tyr Lys Val His Val Glu Lys Asp Phe Ile Ala Phe Cys Ser
 35             40             45

Ser Thr Pro His His Val Ser Trp Arg Asp Val Asn Lys Gly Ser Leu
 50             55             60

Phe Ile Thr Gln Leu Ile Thr Cys Phe Gln Lys Tyr Ser Trp Cys Phe
 65             70             75             80

His Leu Glu Glu Val Phe Arg Lys Val Gln Gln Ser Phe Glu Lys Pro
 85             90             95

```

-continued

---

Asn Val Arg Ala Gln Met Pro Thr Ile Glu Arg Leu Ser Met Thr Arg  
 100 105 110

Tyr Phe Tyr Leu Phe Pro Gly Asn  
 115 120

<210> SEQ ID NO 22  
 <211> LENGTH: 500  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (28)..(28)  
 <223> OTHER INFORMATION: n = unknown at position 28

<400> SEQUENCE: 22

tttttttttt tttttttttt tttttagntg ttttcatttt tattgagata ccaaacgtta 60  
 aatgagtgtg ttgtgctagc taaggtaactt ttccaagttg gtcaagaagg gctggactgc 120  
 ccatgattgc ttttttcaat tgccagggaa gagatagaag tatcttgtca tggatagtcg 180  
 ttcaatgggtg ggcactctggg ctctaacatt tggtttttca aatgactggt gtaccttccg 240  
 aaatacttcc tccagatgaa agcaccacga atacttttgg aagcacgtga tgagttgtgt 300  
 aatgaagaga gatcccttgt tcacgtctct ccaagacaca tgatgtgggg tcgaggagca 360  
 gaaagcgtg aagtccttct ccacgtggac cttgtaaatg gcgtcgtcct ctaggttgtc 420  
 acccacctgt gaaaggctgc taccatctg tgaagtctg tccattgggg ccgctggaga 480  
 gtcactgatc aacagttccc 500

<210> SEQ ID NO 23  
 <211> LENGTH: 1230  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1230)

<400> SEQUENCE: 23

atg gcc gac aag gat ctg aag ggc aag agg aag cag ttc atc aac tca 48  
 Met Ala Asp Lys Asp Leu Lys Gly Lys Arg Lys Gln Phe Ile Asn Ser  
 1 5 10 15

gtc ggc atg ggg acg gtc aac ggc ttg ctg gat gaa ctc ttt gag aaa 96  
 Val Gly Met Gly Thr Val Asn Gly Leu Leu Asp Glu Leu Phe Glu Lys  
 20 25 30

aac gtg ctg aac cag gag gag atg gag aga gta aaa tgt gaa aac gct 144  
 Asn Val Leu Asn Gln Glu Glu Met Glu Arg Val Lys Cys Glu Asn Ala  
 35 40 45

acc gtt atg gac aag gcc cga gct ctg atc gac agc gtc ctg cgg aaa 192  
 Thr Val Met Asp Lys Ala Arg Ala Leu Ile Asp Ser Val Leu Arg Lys  
 50 55 60

ggg cca cgg gcg tgc cag atc ttt atc tgt cac atc tgt gag gaa gac 240  
 Gly Pro Arg Ala Cys Gln Ile Phe Ile Cys His Ile Cys Glu Glu Asp  
 65 70 75 80

acc cac ctt gca gag acg ctg ggg ctc tcc tca agc cca caa tct gga 288  
 Thr His Leu Ala Glu Thr Leu Gly Leu Ser Ser Ser Pro Gln Ser Gly  
 85 90 95

aat tct cag aac acc acg gac tct gaa gta gcg ttt cct cct ctt cca 336  
 Asn Ser Gln Asn Thr Thr Asp Ser Glu Val Ala Phe Pro Pro Leu Pro  
 100 105 110

gcc agc gtg aat aac atg cct ggg ccg gct gag cca gaa gaa tct gta 384  
 Ala Ser Val Asn Asn Met Pro Gly Pro Ala Glu Pro Glu Glu Ser Val  
 115 120 125

-continued

gat gct ctc aag ctt tgt cct cgt gaa aac ttc gtg aaa ctg tgt aaa	432
Asp Ala Leu Lys Leu Cys Pro Arg Glu Asn Phe Val Lys Leu Cys Lys	
130 135 140	
cag agg gct gaa gag atc tac cca ata aag gag aga aag gat cgt act	480
Gln Arg Ala Glu Glu Ile Tyr Pro Ile Lys Glu Arg Lys Asp Arg Thr	
145 150 155 160	
cgt ctg gct ctc atc ata tgc aat acg acg ttc gat cat ctt tct ctc	528
Arg Leu Ala Leu Ile Ile Cys Asn Thr Thr Phe Asp His Leu Ser Leu	
165 170 175	
agg aag ggg gct gac ctt gac gtt gca ggg atg agg agg ctg ctt aca	576
Arg Lys Gly Ala Asp Leu Asp Val Ala Gly Met Arg Arg Leu Leu Thr	
180 185 190	
gac ctt ggc tac agt gtg cac ata aaa gag gaa ctc act gct aag gac	624
Asp Leu Gly Tyr Ser Val His Ile Lys Glu Glu Leu Thr Ala Lys Asp	
195 200 205	
atg gaa tca gag ctg agg gca ttt gct gcc cgt cca gag cac aag tcc	672
Met Glu Ser Glu Leu Arg Ala Phe Ala Ala Arg Pro Glu His Lys Ser	
210 215 220	
tcg gac agc aca ttc ctg gtg ttc atg tct cat ggc atc ctg agt gga	720
Ser Asp Ser Thr Phe Leu Val Phe Met Ser His Gly Ile Leu Ser Gly	
225 230 235 240	
atc tgt ggg acg aag tac agc gct gaa gga gac cca gat gta ttg gct	768
Ile Cys Gly Thr Lys Tyr Ser Ala Glu Gly Asp Pro Asp Val Leu Ala	
245 250 255	
tat gac acc atc ttc cag att ttc aac aac cgc aac tgc ctt agt cta	816
Tyr Asp Thr Ile Phe Gln Ile Phe Asn Asn Arg Asn Cys Leu Ser Leu	
260 265 270	
aag gac aag ccc aag gtc atc atc gtc cag gcc tgc aga ggt gaa aat	864
Lys Asp Lys Pro Lys Val Ile Ile Val Gln Ala Cys Arg Gly Glu Asn	
275 280 285	
ttg ggg gaa ctg ttg atc agt gac tct cca gcg gcc cca atg gac agc	912
Leu Gly Glu Leu Leu Ile Ser Asp Ser Pro Ala Ala Pro Met Asp Ser	
290 295 300	
act tca cag atg ggt agc agc ctt tca cag gtg ggt gac aac cta gag	960
Thr Ser Gln Met Gly Ser Ser Leu Ser Gln Val Gly Asp Asn Leu Glu	
305 310 315 320	
gac gac gcc att tac aag gtc cac gtg gag aag gac ttc atc gct ttc	1008
Asp Asp Ala Ile Tyr Lys Val His Val Glu Lys Asp Phe Ile Ala Phe	
325 330 335	
tgc tcc tcg acc cca cat cat gtg tct tgg aga gac gtg aac aag gga	1056
Cys Ser Ser Thr Pro His His Val Ser Trp Arg Asp Val Asn Lys Gly	
340 345 350	
tct ctc ttc att aca caa ctc atc acg tgc ttc caa aag tat tcg tgg	1104
Ser Leu Phe Ile Thr Gln Leu Ile Thr Cys Phe Gln Lys Tyr Ser Trp	
355 360 365	
tgc ttt cat ctg gag gaa gta ttt cgg aag gta caa cag tca ttt gaa	1152
Cys Phe His Leu Glu Glu Val Phe Arg Lys Val Gln Gln Ser Phe Glu	
370 375 380	
aaa cca aat gtt aga gcc cag atg ccc acc att gaa cga cta tcc atg	1200
Lys Pro Asn Val Arg Ala Gln Met Pro Thr Ile Glu Arg Leu Ser Met	
385 390 395 400	
aca aga tac ttc tat ctc ttc cct ggc aat	1230
Thr Arg Tyr Phe Tyr Leu Phe Pro Gly Asn	
405 410	

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 410

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Felis catus

-continued

&lt;400&gt; SEQUENCE: 24

Met Ala Asp Lys Asp Leu Lys Gly Lys Arg Lys Gln Phe Ile Asn Ser  
 1 5 10 15  
 Val Gly Met Gly Thr Val Asn Gly Leu Leu Asp Glu Leu Phe Glu Lys  
 20 25 30  
 Asn Val Leu Asn Gln Glu Glu Met Glu Arg Val Lys Cys Glu Asn Ala  
 35 40 45  
 Thr Val Met Asp Lys Ala Arg Ala Leu Ile Asp Ser Val Leu Arg Lys  
 50 55 60  
 Gly Pro Arg Ala Cys Gln Ile Phe Ile Cys His Ile Cys Glu Glu Asp  
 65 70 75 80  
 Thr His Leu Ala Glu Thr Leu Gly Leu Ser Ser Ser Pro Gln Ser Gly  
 85 90 95  
 Asn Ser Gln Asn Thr Thr Asp Ser Glu Val Ala Phe Pro Pro Leu Pro  
 100 105 110  
 Ala Ser Val Asn Asn Met Pro Gly Pro Ala Glu Pro Glu Glu Ser Val  
 115 120 125  
 Asp Ala Leu Lys Leu Cys Pro Arg Glu Asn Phe Val Lys Leu Cys Lys  
 130 135 140  
 Gln Arg Ala Glu Glu Ile Tyr Pro Ile Lys Glu Arg Lys Asp Arg Thr  
 145 150 155 160  
 Arg Leu Ala Leu Ile Ile Cys Asn Thr Thr Phe Asp His Leu Ser Leu  
 165 170 175  
 Arg Lys Gly Ala Asp Leu Asp Val Ala Gly Met Arg Arg Leu Leu Thr  
 180 185 190  
 Asp Leu Gly Tyr Ser Val His Ile Lys Glu Glu Leu Thr Ala Lys Asp  
 195 200 205  
 Met Glu Ser Glu Leu Arg Ala Phe Ala Ala Arg Pro Glu His Lys Ser  
 210 215 220  
 Ser Asp Ser Thr Phe Leu Val Phe Met Ser His Gly Ile Leu Ser Gly  
 225 230 235 240  
 Ile Cys Gly Thr Lys Tyr Ser Ala Glu Gly Asp Pro Asp Val Leu Ala  
 245 250 255  
 Tyr Asp Thr Ile Phe Gln Ile Phe Asn Asn Arg Asn Cys Leu Ser Leu  
 260 265 270  
 Lys Asp Lys Pro Lys Val Ile Ile Val Gln Ala Cys Arg Gly Glu Asn  
 275 280 285  
 Leu Gly Glu Leu Leu Ile Ser Asp Ser Pro Ala Ala Pro Met Asp Ser  
 290 295 300  
 Thr Ser Gln Met Gly Ser Ser Leu Ser Gln Val Gly Asp Asn Leu Glu  
 305 310 315 320  
 Asp Asp Ala Ile Tyr Lys Val His Val Glu Lys Asp Phe Ile Ala Phe  
 325 330 335  
 Cys Ser Ser Thr Pro His His Val Ser Trp Arg Asp Val Asn Lys Gly  
 340 345 350  
 Ser Leu Phe Ile Thr Gln Leu Ile Thr Cys Phe Gln Lys Tyr Ser Trp  
 355 360 365  
 Cys Phe His Leu Glu Glu Val Phe Arg Lys Val Gln Gln Ser Phe Glu  
 370 375 380  
 Lys Pro Asn Val Arg Ala Gln Met Pro Thr Ile Glu Arg Leu Ser Met  
 385 390 395 400  
 Thr Arg Tyr Phe Tyr Leu Phe Pro Gly Asn  
 405 410

-continued

---

```

<210> SEQ ID NO 25
<211> LENGTH: 1230
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 25

attgccaggg aagagataga agtatcttgt catggatagt cgttcaatgg tgggcatctg      60
ggctctaaca ttgtgtttt caaatgactg ttgtacctc cgaataactt cctccagatg      120
aaagcaccac gaatactttt ggaagcacgt gatgagttgt gtaatgaaga gagatccctt      180
gttcacgtct ctccaagaca catgatgtgg ggctgaggag cagaaagcga tgaagtccct      240
ctccacgtgg acctgtgtaa tggcgtcgtc ctctaggttg tcaccacact gtgaaaggct      300
gctaccatc tgtgaagtgc tgtccattgg ggccgctgga gagtactga tcaacagttc      360
ccccaaattt tcacctctgc aggcctggac gatgatgacc ttgggcttgt cctttagact      420
aaggcagttg cgttcttga aaatctgaa gatggtgtca taagccaata catctgggtc      480
tccttcagcg ctgtacttcg tcccacagat tccactcagg atgccatgag acatgaacac      540
caggaatgtg ctgtccgagg acttctgctc tggacgggca gcaaatgccc tcagctctga      600
ttccatgtcc ttagcagtga gttcctctt tatgtgcaca ctgtagccaa ggtctgtaag      660
cagcctctc atccctgcaa cgtcaagtc agcccccttc ctgagagaaa gatgatcgaa      720
cgtcgtattg catatgatga gagccagacg agtacgatcc tttctctcct ttattgggta      780
gatctcttca gccctctgtt tacacagttt cacgaagttt tcacgaggac aaagcttgag      840
agcatctaca gattcttctg gctcagccgg cccaggcatg ttattcacgc tggctggaag      900
aggagaaac gctacttcag agtccgtggt gttctgagaa tttccagatt gtgggcttga      960
ggagagcccc agcgtctctg caaggtgggt gtcttctca cagatgtgac agataaagat     1020
ctggcacgcc cgtggccctt tccgcaggac gctgtcgatc agagctcggg ccttgtccat     1080
aacggtagcg ttttcacatt ttactctctc catctcctcc tggttcagca cgtttttctc     1140
aaagagttca tccagcaagc cgttgaccgt ccccatgccg actgagttga tgaactgctt     1200
cctcttgccc ttcagatcct tgcggccat                                     1230
    
```

```

<210> SEQ ID NO 26
<211> LENGTH: 921
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(921)

<400> SEQUENCE: 26

ata tgg gaa ctg gag aaa aac gtt tat gtt gta gag ttg gac tgg cac      48
Ile Trp Glu Leu Glu Lys Asn Val Tyr Val Val Glu Leu Asp Trp His
1          5          10          15

cct gat gcc ccc gga gaa atg gtg gtc ctc acc tgc aat act cct gaa      96
Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys Asn Thr Pro Glu
          20          25          30

gaa gat gac atc acc tgg acc tct gac cag agc agt gaa gtc cta ggc      144
Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln Ser Ser Glu Val Leu Gly
          35          40          45

tct ggt aaa act ctg acc atc caa gtc aaa gaa ttt gca gat gct ggc      192
Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Ala Asp Ala Gly
          50          55          60

cag tat acc tgt cat aaa gga ggc gag gtt ctg agc cat tcg ttc ctc      240
Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu Ser His Ser Phe Leu
65          70          75          80
    
```



-continued

---

Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu Ser His Ser Phe Leu  
 65 70 75 80

Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Arg  
 85 90 95

Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys  
 100 105 110

Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr  
 115 120 125

Asp Leu Lys Phe Thr Val Lys Ser Ser Arg Gly Ser Ser Asp Pro Gln  
 130 135 140

Glu Val Thr Cys Gly Ala Ala Thr Leu Ser Ala Glu Lys Val Arg Val  
 145 150 155 160

Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser  
 165 170 175

Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp  
 180 185 190

Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile  
 195 200 205

Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu Gln Leu Lys Pro  
 210 215 220

Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr  
 225 230 235 240

Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Gly Val Gln Val  
 245 250 255

Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Ser Val Asp Lys  
 260 265 270

Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln  
 275 280 285

Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asn Trp Ala Ser Val  
 290 295 300

Ser Cys Ser  
 305

<210> SEQ ID NO 28  
 <211> LENGTH: 921  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 28

actgcaggac acggatgccc agttgctcca ggatgagcta tagtagcggg ctctggcttg 60

cacgcggatc ttggcatcct tgtggcacac gaccttggct gaggtcttgt ccacggagag 120

tctgtctttc ttttctctgt tgttcttgcc ctggacctgt acgccaaatg ttaaggagaa 180

gtaggaatgt ggggtgctcc aggtgtcagg gtattcccag ctacttcca catgccgaga 240

atTTTTtaat ggcttcagtt gcaggttctt ggggtgggtcc ggTTTgatga tgtccctgat 300

gaagaagctg ctggtgtagt tttcgtactt gagcttgtga atagcgtcca ccaogacttc 360

aatgggtagg ctctcctcgg cagccgggca ggcactgccc tcctgacact ccaactgtgta 420

cttctataaa tcctctgtgt ccaactctgac cttctctgct gagagtgtcg ctgctccaca 480

agtcacctct tgggggtcag aggagocctct gotgcttttg acagtgaatt tcaaactcgg 540

actgattgcc gtcagccacc agcaggtgaa acgtccagaa taattctttg cctcacattt 600

tagaaagatc ttattttttg attctttctg ttcccttaag atatcagtg accaaaattcc 660

atcttccctt ttgtgtatca ggaggaacga atggctcaga acctcgcctc ctttatgaca 720

-continued

ggtatactgg ccagcatctg caaattcttt gacttggatg gtcagagttt taccagagcc 780  
 taggaacttca ctgctctggt cagaggtcca ggtgatgtca tcttcttcag gagtattgca 840  
 ggtgaggacc accattttctc cgggggcacg aggggtgccag tccaactcta caacataaac 900  
 gtttttctcc agttcccata t 921

<210> SEQ ID NO 29  
 <211> LENGTH: 987  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(987)

<400> SEQUENCE: 29

atg cat cct cag cag ttg gtc atc gcc tgg ttt tcc ctg gtt ttg ctg 48  
 Met His Pro Gln Gln Leu Val Ile Ala Trp Phe Ser Leu Val Leu Leu  
 1 5 10 15

gca cct ccc ctc atg gcc ata tgg gaa ctg gag aaa aac gtt tat gtt 96  
 Ala Pro Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asn Val Tyr Val  
 20 25 30

gta gag ttg gac tgg cac cct gat gcc ccc gga gaa atg gtg gtc ctc 144  
 Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu  
 35 40 45

acc tgc aat act cct gaa gaa gat gac atc acc tgg acc tct gac cag 192  
 Thr Cys Asn Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln  
 50 55 60

agc agt gaa gtc cta ggc tct ggt aaa act ctg acc atc caa gtc aaa 240  
 Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys  
 65 70 75 80

gaa ttt gca gat gct ggc cag tat acc tgt cat aaa gga ggc gag gtt 288  
 Glu Phe Ala Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val  
 85 90 95

ctg agc cat tcg ttc ctc ctg ata cac aaa aag gaa gat gga att tgg 336  
 Leu Ser His Ser Phe Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp  
 100 105 110

tcc act gat atc tta agg gaa cag aaa gaa tcc aaa aat aag atc ttt 384  
 Ser Thr Asp Ile Leu Arg Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe  
 115 120 125

cta aaa tgt gag gca aag aat tat tct gga cgt ttc acc tgc tgg tgg 432  
 Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp  
 130 135 140

ctg acg gca atc agt acc gat ttg aaa ttc act gtc aaa agc agc aga 480  
 Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Thr Val Lys Ser Ser Arg  
 145 150 155 160

ggc tcc tct gac ccc caa ggg gtg act tgt gga gca gcg aca ctc tca 528  
 Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser  
 165 170 175

gca gag aag gtc aga gtg gac aac agg gat tat aag aag tac aca gtg 576  
 Ala Glu Lys Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val  
 180 185 190

gag tgt cag gag ggc agt gcc tgc ccg gct gcc gag gag agc cta ccc 624  
 Glu Cys Gln Glu Gly Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro  
 195 200 205

att gaa gtc gtg gtg gac gct att cac aag ctc aag tac gaa aac tac 672  
 Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr  
 210 215 220

acc agc agc ttc ttc atc agg gac atc atc aaa ccg gac cca ccc aag 720  
 Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys  
 225 230 235 240

-continued

---

```

aac ctg caa ctg aag cca tta aaa aat tct cgg cat gtg gaa gtg agc      768
Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser
                245                250                255

tgg gaa tac cct gac acc tgg agc acc cca cat tcc tac ttc tcc tta      816
Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu
                260                265                270

aca ttt ggc gta cag gtc cag ggc aag aac aac aga gaa aag aaa gac      864
Thr Phe Gly Val Gln Val Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp
                275                280                285

aga ctc tcc gtg gac aag acc tca gcc aag gtc gtg tgc cac aag gat      912
Arg Leu Ser Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp
                290                295                300

gcc aag atc cgc gtg caa gcc aga gac cgc tac tat agc tca tcc tgg      960
Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp
305                310                315                320

agc aac tgg gca tcc gtg tcc tgc agt      987
Ser Asn Trp Ala Ser Val Ser Cys Ser
                325
    
```

```

<210> SEQ ID NO 30
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Felis catus
    
```

<400> SEQUENCE: 30

```

Met His Pro Gln Gln Leu Val Ile Ala Trp Phe Ser Leu Val Leu Leu
1                5                10                15

Ala Pro Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asn Val Tyr Val
                20                25                30

Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
                35                40                45

Thr Cys Asn Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln
50                55                60

Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
65                70                75                80

Glu Phe Ala Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
                85                90                95

Leu Ser His Ser Phe Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
                100                105                110

Ser Thr Asp Ile Leu Arg Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
                115                120                125

Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
130                135                140

Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Thr Val Lys Ser Ser Arg
145                150                155                160

Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser
                165                170                175

Ala Glu Lys Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val
                180                185                190

Glu Cys Gln Glu Gly Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro
195                200                205

Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr
210                215                220

Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys
225                230                235                240

Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser
                245                250                255
    
```

-continued

Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu  
 260 265 270  
 Thr Phe Gly Val Gln Val Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp  
 275 280 285  
 Arg Leu Ser Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp  
 290 295 300  
 Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp  
 305 310 315 320  
 Ser Asn Trp Ala Ser Val Ser Cys Ser  
 325

<210> SEQ ID NO 31  
 <211> LENGTH: 987  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 31

actgcaggac acggatgccc agttgctcca ggatgagcta tagtagcggg ctctggcctg 60  
 cacgcggatc ttggcatcct tgtggcacac gacctggct gaggtcttgt ccacggagag 120  
 tctgtctttc ttttctctgt tgttcttgcc ctggacctgt acgccaaatg ttaaggagaa 180  
 gtaggaatgt ggggtgctcc aggtgtcagg gtattcccag ctacttcca catgccgaga 240  
 attttttaat ggcttcagtt gcaggttcct ggggtgggtcc ggtttgatga tgtccctgat 300  
 gaagaagctg ctggtgtagt tttcgtactt gagcttgtga atagcgtcca ccacgacttc 360  
 aatgggtagg ctctcctcgg cagccgggca ggcactgccc tcctgacact cactgtgta 420  
 cttcttataa tcctctgtt cactctgac cttctctgct gagagtgtcg ctgctccaca 480  
 agtcaccctc tgggggtcag aggagcctct gctgcttttg acagtgaatt tcaaatcggg 540  
 actgattgcc gtcagccacc agcaggtgaa acgtccagaa taattctttg cctcacattt 600  
 tagaaagatc ttatttttgg attctttctg ttcccttaag atatcagtg accaaattcc 660  
 atcttccttt ttgtgtatca ggaggaacga atggctcaga acctgcctc ctttatgaca 720  
 ggtatactgg ccagcatctg caaattcttt gacttggatg gtcagagttt taccagagcc 780  
 taggacttca ctgctctggt cagaggtcca ggtgatgtca tcttcttcag gagtattgca 840  
 ggtgaggacc accatttctc cgggggcatc aggggtgccag tccaactcta caacataaac 900  
 gtttttctcc agttcccata tggccatgag gggaggtgcc agcaaaacca gggaaaacca 960  
 ggcgatgacc aactgctgag gatgcat 987

<210> SEQ ID NO 32  
 <211> LENGTH: 666  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(666)

<400> SEQUENCE: 32

atg tgc ccg ccg cgt ggc ctc ctc ctt gta acc atc ctg gtc ctg tta 48  
 Met Cys Pro Pro Arg Gly Leu Leu Leu Val Thr Ile Leu Val Leu Leu  
 1 5 10 15  
 aac cac ctg gac cac ctc agt ttg gcc agg aac ctc ccc aca ccc aca 96  
 Asn His Leu Asp His Leu Ser Leu Ala Arg Asn Leu Pro Thr Pro Thr  
 20 25 30  
 cca agc cca gga atg ttc cag tgc ctc aac cac tcc caa acc ctg ctg 144  
 Pro Ser Pro Gly Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu Leu  
 35 40 45



-continued

---

Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln  
 145 150 155 160

Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Val Asn  
 165 170 175

Ser Val Thr Val Pro Gln Asn Ser Ser Leu Glu Glu Pro Asp Phe Tyr  
 180 185 190

Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg  
 195 200 205

Ala Val Thr Ile Asn Arg Met Met Ser Tyr Leu Asn Ala Ser  
 210 215 220

<210> SEQ ID NO 34  
 <211> LENGTH: 666  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 34

ggaagcattc agatagctca tcattctatt gatggtcact gcacgaattc tgaaagcatg 60  
 aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggttcttcca gggaggagtt 120  
 ctgtggcaca gtcacactgt tgacattcag ggcctgtaac agctcatcaa tagctgtcag 180  
 catgttttga tccagaaaga tctgcctttt aggatccatt aacagctttg cattcatggc 240  
 cttgaaactcc acctggtaca tcttcaagtc ctcatagata ctgctaaggc acagggctcgt 300  
 cataaaaagag gtctttcttg aggccaggca actcccatta gttatcagag agatctctct 360  
 ggaagccagg caactctcat tcatgggtaa ttccagtggg aagcaggcct cactctgtct 420  
 ggttttatct tttgtgatat ctcatgatc aatctcttcg gaagtgcagg agtaaaattc 480  
 tagagtttgt ctggccttct gaagcgtggt gctgatggct cgcagcaggg tttgggagtg 540  
 gttgaggcac tggaacattc ctgggcttgg tgtgggtgtg gggagggtcc tggccaaact 600  
 gaggtggtcc aggtggttta acaggaccag gatggttaca aggaggaggc cagcgggcgg 660  
 gcacat 666

<210> SEQ ID NO 35  
 <211> LENGTH: 591  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(591)

<400> SEQUENCE: 35

agg aac ctc ccc aca ccc aca cca agc cca gga atg ttc cag tgc ctc 48  
 Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro Gly Met Phe Gln Cys Leu  
 1 5 10 15

aac cac tcc caa acc ctg ctg cga gcc atc agc aac acg ctt cag aag 96  
 Asn His Ser Gln Thr Leu Leu Arg Ala Ile Ser Asn Thr Leu Gln Lys  
 20 25 30

gcc aga caa act cta gaa ttt tac tcc tgc act tcc gaa gag att gat 144  
 Ala Arg Gln Thr Leu Glu Phe Tyr Ser Cys Thr Ser Glu Glu Ile Asp  
 35 40 45

cat gaa gat atc aca aaa gat aaa acc agc aca gtg gag gcc tgc tta 192  
 His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu  
 50 55 60

cca ctg gaa tta acc atg aat gag agt tgc ctg gct tcc aga gag atc 240  
 Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile  
 65 70 75 80

-continued

tct ctg ata act aat ggg agt tgc ctg gcc tcc aga aag acc tct ttt	288
Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe	
85 90 95	
atg acg acc ctg tgc ctt agc agt atc tat gag gac ttg aag atg tac	336
Met Thr Thr Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr	
100 105 110	
cag gtg gag ttc aag gcc atg aat gca aag ctg tta atg gat cct aaa	384
Gln Val Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys	
115 120 125	
agg cag atc ttt ctg gat caa aac atg ctg aca gct att gat gag ctg	432
Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu	
130 135 140	
tta cag gcc ctg aat gtc aac agt gtg act gtg cca cag aac tcc tcc	480
Leu Gln Ala Leu Asn Val Asn Ser Val Thr Val Pro Gln Asn Ser Ser	
145 150 155 160	
ttg gaa gaa ccg gat ttt tat aaa act aaa atc aag ctc tgc ata ctt	528
Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu	
165 170 175	
ctt cat gct ttc aga att cgt gca gtg acc atc aat aga atg atg agc	576
Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asn Arg Met Met Ser	
180 185 190	
tat ctg aat gct tcc	591
Tyr Leu Asn Ala Ser	
195	

<210> SEQ ID NO 36  
 <211> LENGTH: 197  
 <212> TYPE: PRT  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 36

Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro Gly Met Phe Gln Cys Leu	
1 5 10 15	
Asn His Ser Gln Thr Leu Leu Arg Ala Ile Ser Asn Thr Leu Gln Lys	
20 25 30	
Ala Arg Gln Thr Leu Glu Phe Tyr Ser Cys Thr Ser Glu Glu Ile Asp	
35 40 45	
His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu	
50 55 60	
Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile	
65 70 75 80	
Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe	
85 90 95	
Met Thr Thr Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr	
100 105 110	
Gln Val Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys	
115 120 125	
Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu	
130 135 140	
Leu Gln Ala Leu Asn Val Asn Ser Val Thr Val Pro Gln Asn Ser Ser	
145 150 155 160	
Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu	
165 170 175	
Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asn Arg Met Met Ser	
180 185 190	
Tyr Leu Asn Ala Ser	
195	

-continued

---

```

<210> SEQ ID NO 37
<211> LENGTH: 591
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 37
ggaagcattc agatagctca tcattctatt gatggtcact gcacgaattc tgaagcatg      60
aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggttcttcca aggaggagtt    120
ctgtggcaca gtcacactgt tgacattcag ggcctgtaac agctcatcaa tagctgtcag    180
catgttttga tccagaaaga tctgcctttt aggatccatt aacagctttg cattcatggc    240
cttgaactcc acctggtaca tcttcaagtc ctcatagata ctgctaaggc acagggtcgt    300
cataaaagag gtctttcttg aggccaggca actcccatta gttatcagag agatctctct    360
ggaagccagg caactctcat tcatggtaa ttccagtggg aagcaggcct ccactgtgct     420
ggttttatct tttgtgatat cttcatgatc aatctcttcg gaagtgcagg agtaaaattc    480
tagagtttgt ctggccttct gaagcgtggt gctgatggct cgcagcaggg tttgggagtg    540
gttgaggcac tgaacattc ctgggcttgg tgtgggtgtg gggaggttcc t              591

<210> SEQ ID NO 38
<211> LENGTH: 1599
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1599)

<400> SEQUENCE: 38
atg cat cct cag cag ttg gtc atc gcc tgg ctt tcc ctg gtt ttg ctg      48
Met His Pro Gln Gln Leu Val Ile Ala Trp Leu Ser Leu Val Leu Leu
1          5          10         15

gca cct ccc ctc atg gcc ata tgg gaa ctg gag aaa aac gtt tat gtt      96
Ala Pro Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asn Val Tyr Val
20         25         30

gta gag ttg gac tgg cac cct gat gcc ccc gga gaa atg gtg gtc ctc     144
Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
35         40         45

acc tgc aat act cct gaa gaa gat gac atc acc tgg acc tct gac cag     192
Thr Cys Asn Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln
50         55         60

agc agt gaa gtc cta ggc tct ggt aaa act ctg acc atc caa gtc aaa     240
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
65         70         75         80

gaa ttt gca gat gct ggc cag tat acc tgt cat aaa gga ggc gag gtt     288
Glu Phe Ala Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
85         90         95

ctg agc cat tcg ttc ctc ctg ata cac aaa aag gaa gat gga att tgg     336
Leu Ser His Ser Phe Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
100        105        110

tcc act gat atc tta agg gaa cag aaa gaa tcc aaa aat aag atc ttt     384
Ser Thr Asp Ile Leu Arg Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
115        120        125

cta aaa tgt gag gca aag aat tat tct gga cgt ttc acc tgc tgg tgg     432
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
130        135        140

ctg acg gca atc agt acc gat ttg aaa ttc act gtc aaa agc agc aga     480
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Thr Val Lys Ser Ser Arg
145        150        155        160

```

-continued

ggc tcc tet gac ccc caa gag gtg act tgt gga gca gcg aca ctc tca Gly Ser Ser Asp Pro Gln Glu Val Thr Cys Gly Ala Ala Thr Leu Ser 165 170 175	528
gca gag aag gtc aga gtg gac aac agg gat tat aag aag tac aca gtg Ala Glu Lys Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val 180 185 190	576
gag tgt cag gag ggc agt gcc tgc ccg gct gcc gag gag agc cta ccc Glu Cys Gln Glu Gly Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro 195 200 205	624
att gaa gtc gtg gtg gac gct att cac aag ctc aag tac gaa aac tac Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr 210 215 220	672
acc agc agc ttc ttc atc agg gac atc atc aaa ccg gac cca ccc aag Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys 225 230 235 240	720
aac ctg caa ctg aag cca tta aaa aat tct cgg cat gtg gaa gtg agc Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser 245 250 255	768
tgg gaa tac cct gac acc tgg agc acc cca cat tcc tac ttc tcc tta Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu 260 265 270	816
aca ttt ggc gta cag gtc cag ggc aag aac aac aga gaa aag aaa gac Thr Phe Gly Val Gln Val Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp 275 280 285	864
aga ctc tcc gtg gac aag acc tca gcc aag gtc gtg tgc cac aag gat Arg Leu Ser Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp 290 295 300	912
gcc aag atc cgc gtg caa gcc aga gac cgc tac tat agc tca tcc tgg Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp 305 310 315 320	960
agc aac tgg gca tcc gtg tcc tgc agt ggt ggc ggt ggc ggc gga tct Ser Asn Trp Ala Ser Val Ser Cys Ser Gly Gly Gly Gly Gly Gly Ser 325 330 335	1008
aga aac ttg cca acc cct act cca tcc ccg ggg atg ttc cag tgc ctc Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro Gly Met Phe Gln Cys Leu 340 345 350	1056
aac cac tcc caa acc ctg ctg cga gcc atc agc aac acg ctt cag aag Asn His Ser Gln Thr Leu Leu Arg Ala Ile Ser Asn Thr Leu Gln Lys 355 360 365	1104
gcc aga caa act cta gaa ttt tac tcc tgc act tcc gaa gag att gat Ala Arg Gln Thr Leu Glu Phe Tyr Ser Cys Thr Ser Glu Glu Ile Asp 370 375 380	1152
cat gaa gat atc aca aaa gat aaa acc agc aca gtg gag gcc tgc tta His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu 385 390 395 400	1200
cca ctg gaa tta acc atg aat gag agt tgc ctg gct tcc aga gag atc Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile 405 410 415	1248
tct ctg ata act aat ggg agt tgc ctg gcc tcc aga aag acc tct ttt Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe 420 425 430	1296
atg acg acc ctg tgc ctt agc agt atc tat gag gac ttg aag atg tac Met Thr Thr Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr 435 440 445	1344
cag gtg gag ttc aag gcc atg aat gca aag ctg tta atg gat cct aaa Gln Val Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys 450 455 460	1392
agg cag atc ttt ctg gat caa aac atg ctg aca gct att gat gag ctg Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu 465 470 475 480	1440

-continued

---

```

tta cag gcc ctg aat gtc aac agt gtg act gtg cca cag aac tcc tcc      1488
Leu Gln Ala Leu Asn Val Asn Ser Val Thr Val Pro Gln Asn Ser Ser
                485                          490                          495

ttg gaa gaa ccg gat ttt tat aaa act aaa atc aag ctc tgc ata ctt      1536
Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu
                500                          505                          510

ctt cat gct ttc aga att cgt gca gtg acc atc aat aga atg atg agc      1584
Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asn Arg Met Met Ser
                515                          520                          525

tat ctg aat gct tcc                                          1599
Tyr Leu Asn Ala Ser
    530

<210> SEQ ID NO 39
<211> LENGTH: 533
<212> TYPE: PRT
<213> ORGANISM: Felis catus

<400> SEQUENCE: 39
Met His Pro Gln Gln Leu Val Ile Ala Trp Leu Ser Leu Val Leu Leu
 1                5                10                15
Ala Pro Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asn Val Tyr Val
 20                25                30
Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
 35                40                45
Thr Cys Asn Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln
 50                55                60
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
 65                70                75                80
Glu Phe Ala Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
 85                90                95
Leu Ser His Ser Phe Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
 100               105               110
Ser Thr Asp Ile Leu Arg Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
 115               120               125
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
 130               135               140
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Thr Val Lys Ser Ser Arg
 145               150               155               160
Gly Ser Ser Asp Pro Gln Glu Val Thr Cys Gly Ala Ala Thr Leu Ser
 165               170               175
Ala Glu Lys Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val
 180               185               190
Glu Cys Gln Glu Gly Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro
 195               200               205
Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr
 210               215               220
Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys
 225               230               235               240
Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser
 245               250               255
Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu
 260               265               270
Thr Phe Gly Val Gln Val Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp
 275               280               285

```

-continued

---

Arg Leu Ser Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp  
 290 295 300

Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp  
 305 310 315 320

Ser Asn Trp Ala Ser Val Ser Cys Ser Gly Gly Gly Gly Gly Ser  
 325 330 335

Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro Gly Met Phe Gln Cys Leu  
 340 345 350

Asn His Ser Gln Thr Leu Leu Arg Ala Ile Ser Asn Thr Leu Gln Lys  
 355 360 365

Ala Arg Gln Thr Leu Glu Phe Tyr Ser Cys Thr Ser Glu Glu Ile Asp  
 370 375 380

His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu  
 385 390 395 400

Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile  
 405 410 415

Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe  
 420 425 430

Met Thr Thr Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr  
 435 440 445

Gln Val Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys  
 450 455 460

Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu  
 465 470 475 480

Leu Gln Ala Leu Asn Val Asn Ser Val Thr Val Pro Gln Asn Ser Ser  
 485 490 495

Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu  
 500 505 510

Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asn Arg Met Met Ser  
 515 520 525

Tyr Leu Asn Ala Ser  
 530

<210> SEQ ID NO 40  
 <211> LENGTH: 1599  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 40

ggaagcattc agatagctca tcattctatt gatggtcact gcacgaattc tgaagcatg 60

aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggttcttcca aggaggagt 120

ctgtggcaca gtcacactgt tgacattcag ggcctgtaac agctcatcaa tagctgtcag 180

catgttttga tccagaaaga tctgcctttt aggatccatt aacagctttg cattcatggc 240

cttgaactcc acctggtaca tcttcaagtc ctcatagata ctgctaaggc acagggctgt 300

cataaaaagag gtctttctgg aggccaggca actcccatta gttatcagag agatctctct 360

ggaagccagg caactctcat tcatgggtaa ttccagtggg aagcaggcct ccaactgtgct 420

ggttttatct tttgtgatat cttcatgatc aatctctctg gaagtgcagg agtaaaatc 480

tagagtttgt ctggccttct gaagcgtggt gotgatggct cgcagcaggg tttgggagtg 540

gttgaggcac tggaacatcc ccggggatgg agtaggggtt ggcaagtttc tagatccgcc 600

gccaccgccca ccaactgcagg acacggatgc ccagttgctc caggatgagc tatagtagcg 660

gtctctggct tgcacgggga tcttggcatc cttgtggcac acgaccttg ctgaggtctt 720

-continued

---

```

gtccacggag agtctgtctt tcttttctct gttgttcttg ccctggacct gtacgccaaa 780
tgtaaggag aagtaggaat gtgggggtgct ccagggtgtca gggatttccc agctcacttc 840
cacatgccga gaatttttta atggcttcag ttgcaggttc ttgggtgggt ccggtttgat 900
gatgtccctg atgaagaagc tgctgggtgta gttttcgtac ttgagcttgt gaatagcgtc 960
caccacgact tcaatgggta ggctctcctc ggcagccggg caggcactgc cctcctgaca 1020
ctccactgtg tacttcttat aatccctggt gtccactctg accttctctg ctgagagtgt 1080
cgctgctcca caagtcacct ctggggggtc agaggagcct ctgctgcttt tgacagtga 1140
tttcaaactg gtactgattg ccgtcagcca ccagcagggtg aaacgtccag aataattctt 1200
tgctcacat tttagaaga tcttattttt ggattctttc tgttccctta agatatcagt 1260
ggaccaaatt ccatcttctt ttttgtgat caggaggaac gaatggctca gaacctcgcc 1320
tcctttatga caggataact ggccagcatc tgcaaattct ttgacttggg tggtcagagt 1380
tttaccagag cctaggactt cactgctctg gtcagaggtc caggatgatgt catcttcttc 1440
aggagtattg caggtaggag ccaccatttc tccgggggca tcagggtgcc agtccaactc 1500
tacaacataa acgtttttct ccagttccca tatggccatg aggggaggtg ccagcaaac 1560
cagggaagc caggcagatga ccaactgctg aggatgcat 1599

```

```

<210> SEQ ID NO 41
<211> LENGTH: 576
<212> TYPE: DNA
<213> ORGANISM: Felis catus

```

```

<400> SEQUENCE: 41
aattctgttt ttgaacagtg aacattatgg acttatcccc attttcatcc ttttttttca 60
aaatgagttt gaaaagatct ttctcttttt cacaagctag aaagtacccc ttgtacaatg 120
aagactcaaa ttgtatctta tcatcatgtc ctggaacact tctctgaaag aatgatgatg 180
catttccttc atcattgata ctctcaggag gactcatttc cttaaaggaa ataattttgt 240
tctcacagga gagagtagac atggtcttat aattcacaga gatggttact gccagacctc 300
tagtgaggct atctttatac atatatatga taaattcagt ccgggggtgca ttatctgtac 360
agtcagaatc aggcataatc tcaaacacag gttgatctcc ctggttaatg aagagaactt 420
ggtcgttcaa gtttcgtaag attgagagtt tatgttcaag cttgccaaag taatctgttt 480
ccaggttttc atcactgtca gctacaaagt aaagtgtatt gtcaataaat ttcattccca 540
caaagttgat gcaatcatct actggtatag cagtca 576

```

```

<210> SEQ ID NO 42
<211> LENGTH: 0
<212> TYPE: DNA
<213> ORGANISM: Felis catus

```

```

<400> SEQUENCE: 42

```

```

000

```

```

<210> SEQ ID NO 43
<211> LENGTH: 1533
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1533)

```

-continued

&lt;400&gt; SEQUENCE: 43

ata tgg gaa ctg gag aaa aac gtt tat gtt gta gag ttg gac tgg cac	48
Ile Trp Glu Leu Glu Lys Asn Val Tyr Val Val Glu Leu Asp Trp His	
1 5 10 15	
cct gat gcc ccc gga gaa atg gtg gtc ctc acc tgc aat act cct gaa	96
Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys Asn Thr Pro Glu	
20 25 30	
gaa gat gac atc acc tgg acc tct gac cag agc agt gaa gtc cta ggc	144
Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln Ser Ser Glu Val Leu Gly	
35 40 45	
tct ggt aaa act ctg acc atc caa gtc aaa gaa ttt gca gat gct ggc	192
Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Ala Asp Ala Gly	
50 55 60	
cag tat acc tgt cat aaa gga ggc gag gtt ctg agc cat tcg ttc ctc	240
Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu Ser His Ser Phe Leu	
65 70 75 80	
ctg ata cac aaa aag gaa gat gga att tgg tcc act gat atc tta agg	288
Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Arg	
85 90 95	
gaa cag aaa gaa tcc aaa aat aag atc ttt cta aaa tgt gag gca aag	336
Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys	
100 105 110	
aat tat tct gga cgt ttc acc tgc tgg tgg ctg acg gca atc agt acc	384
Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr	
115 120 125	
gat ttg aaa ttc act gtc aaa agc agc aga ggc tcc tct gac ccc caa	432
Asp Leu Lys Phe Thr Val Lys Ser Ser Arg Gly Ser Ser Asp Pro Gln	
130 135 140	
gag gtg act tgt gga gca gcg aca ctc tca gca gag aag gtc aga gtg	480
Glu Val Thr Cys Gly Ala Ala Thr Leu Ser Ala Glu Lys Val Arg Val	
145 150 155 160	
gac aac agg gat tat aag aag tac aca gtg gag tgt cag gag ggc agt	528
Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser	
165 170 175	
gcc tgc ccg gct gcc gag gag agc cta ccc att gaa gtc gtg gtg gac	576
Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp	
180 185 190	
gct att cac aag ctc aag tac gaa aac tac acc agc agc ttc ttc atc	624
Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile	
195 200 205	
agg gac atc atc aaa ccg gac cca ccc aag aac ctg caa ctg aag cca	672
Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu Gln Leu Lys Pro	
210 215 220	
tta aaa aat tct cgg cat gtg gaa gtg agc tgg gaa tac cct gac acc	720
Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr	
225 230 235 240	
tgg agc acc cca cat tcc tac ttc tcc tta aca ttt ggc gta cag gtc	768
Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Gly Val Gln Val	
245 250 255	
cag ggc aag aac aac aga gaa aag aaa gac aga ctc tcc gtg gac aag	816
Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Ser Val Asp Lys	
260 265 270	
acc tca gcc aag gtc gtg tgc cac aag gat gcc aag atc cgc gtg caa	864
Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln	
275 280 285	
gcc aga gac cgc tac tat agc tca tcc tgg agc aac tgg gca tcc gtg	912
Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asn Trp Ala Ser Val	
290 295 300	

-continued

tcc tgc agt ggt ggc ggt ggc ggc gga tct aga aac ttg cca acc cct	960
Ser Cys Ser Gly Gly Gly Gly Gly Gly Ser Arg Asn Leu Pro Thr Pro	
305 310 315 320	
act cca tcc ccg ggg atg ttc cag tgc ctc aac cac tcc caa acc ctg	1008
Thr Pro Ser Pro Gly Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu	
325 330 335	
ctg cga gcc atc agc aac acg ctt cag aag gcc aga caa act cta gaa	1056
Leu Arg Ala Ile Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu	
340 345 350	
ttt tac tcc tgc act tcc gaa gag att gat cat gaa gat atc aca aaa	1104
Phe Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys	
355 360 365	
gat aaa acc agc aca gtg gag gcc tgc tta cca ctg gaa tta acc atg	1152
Asp Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met	
370 375 380	
aat gag agt tgc ctg gct tcc aga gag atc tct ctg ata act aat ggg	1200
Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly	
385 390 395 400	
agt tgc ctg gcc tcc aga aag acc tct ttt atg acg acc ctg tgc ctt	1248
Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe Met Thr Thr Leu Cys Leu	
405 410 415	
agc agt atc tat gag gac ttg aag atg tac cag gtg gag ttc aag gcc	1296
Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Val Glu Phe Lys Ala	
420 425 430	
atg aat gca aag ctg tta atg gat cct aaa agg cag atc ttt ctg gat	1344
Met Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp	
435 440 445	
caa aac atg ctg aca gct att gat gag ctg tta cag gcc ctg aat gtc	1392
Gln Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Val	
450 455 460	
aac agt gtg act gtg cca cag aac tcc tcc ttg gaa gaa ccg gat ttt	1440
Asn Ser Val Thr Val Pro Gln Asn Ser Ser Leu Glu Glu Pro Asp Phe	
465 470 475 480	
tat aaa act aaa atc aag ctc tgc ata ctt ctt cat gct ttc aga att	1488
Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile	
485 490 495	
cgt gca gtg acc atc aat aga atg atg agc tat ctg aat gct tcc	1533
Arg Ala Val Thr Ile Asn Arg Met Met Ser Tyr Leu Asn Ala Ser	
500 505 510	

<210> SEQ ID NO 44  
 <211> LENGTH: 511  
 <212> TYPE: PRT  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 44

Ile Trp Glu Leu Glu Lys Asn Val Tyr Val Val Glu Leu Asp Trp His	
1 5 10 15	
Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys Asn Thr Pro Glu	
20 25 30	
Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln Ser Ser Glu Val Leu Gly	
35 40 45	
Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Ala Asp Ala Gly	
50 55 60	
Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu Ser His Ser Phe Leu	
65 70 75 80	
Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Arg	
85 90 95	

-continued

---

Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys  
                   100  105  110

Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr  
                   115  120  125

Asp Leu Lys Phe Thr Val Lys Ser Ser Arg Gly Ser Ser Asp Pro Gln  
                   130  135  140

Glu Val Thr Cys Gly Ala Ala Thr Leu Ser Ala Glu Lys Val Arg Val  
                   145  150  155  160

Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser  
                                   165  170  175

Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp  
                                   180  185  190

Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile  
                                   195  200  205

Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu Gln Leu Lys Pro  
                   210  215  220

Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr  
                   225  230  235  240

Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Gly Val Gln Val  
                                   245  250  255

Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Ser Val Asp Lys  
                                   260  265  270

Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln  
                                   275  280  285

Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asn Trp Ala Ser Val  
                   290  295  300

Ser Cys Ser Gly Gly Gly Gly Gly Ser Arg Asn Leu Pro Thr Pro  
                   305  310  315  320

Thr Pro Ser Pro Gly Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu  
                                   325  330  335

Leu Arg Ala Ile Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu  
                                   340  345  350

Phe Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys  
                                   355  360  365

Asp Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met  
                   370  375  380

Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly  
                   385  390  395  400

Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe Met Thr Thr Leu Cys Leu  
                                   405  410  415

Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Val Glu Phe Lys Ala  
                                   420  425  430

Met Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp  
                                   435  440  445

Gln Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Val  
                   450  455  460

Asn Ser Val Thr Val Pro Gln Asn Ser Ser Leu Glu Glu Pro Asp Phe  
                   465  470  475  480

Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile  
                                   485  490  495

Arg Ala Val Thr Ile Asn Arg Met Met Ser Tyr Leu Asn Ala Ser  
                   500  505  510

-continued

---

```

<210> SEQ ID NO 45
<211> LENGTH: 1533
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 45
ggaagcattc agatagctca tcattctatt gatggtcact gcacgaattc tgaagcatg      60
aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggttcttcca aggaggagtt    120
ctgtggcaca gtcacactgt tgacattcag ggcctgtaac agctcatcaa tagctgtcag    180
catgttttga tccagaaaga tctgcctttt aggatccatt aacagctttg cattcatggc    240
cttgaactcc acctggtaca tcttcaagtc ctcatagata ctgctaaggc acaggggtcgt    300
cataaaagag gtctttcttg aggccaggca actcccatta gttatcagag agatctctct    360
ggaagccagg caactctcat tcatggttaa ttccagtggg aagcaggcct ccaactgtgct    420
ggtttttatct tttgtgatat cttcatgatc aatctctctg gaagtgcagg agtaaaattc    480
tagagtttgt ctggccttct gaagcgtggt gctgatggct cgcagcaggg tttgggagtg    540
gttgaggcac tgaacatcc cgggggatgg agtaggggtt ggcaagtttc tagatccgcc    600
gccaccgcca ccaactgcagg acacggatgc ccagttgctc caggatgagc tatagtagcg    660
gtctctggct tgcacgcgga tcttggcatc cttgtggcac acgacctgg ctgaggtcct    720
gtccacggag agtctgtcct tcttttctct gttgttcttg ccctggacct gtacgcaaaa    780
tgtaaggag aagtaggaat gtgggggtct ccagggtgca gggatttccc agctcacttc    840
cacatgccga gaatttttta atggcttcag ttgcaggttc ttgggtgggt cgggtttgat    900
gatgtccctg atgaagaagc tgctgggtga gttttcgtac ttgagcttgt gaatagcgtc    960
caccacgact tcaatgggta ggctctcctc ggcagccggg caggcactgc cctcctgaca   1020
ctccactgtg tacttcttat aatccctggt gtccactctg accttctctg ctgagagtgt   1080
cgctgtccca caagtacact cttgggggtc agaggagcct ctgctgcttt tgacagtгаа   1140
tttcaaatcg gtactgattg ccgtcagcca ccagcaggtg aaactgccag aataattctt   1200
tgcctcacat tttagaaga tcttattttt ggattctttc tgttccctta agatatcagt   1260
ggaccaaatt ccatcttctt ttttgtgtat caggaggaac gaatggctca gaacctcgcc   1320
tcctttatga caggtatact ggccagcatc tgcaaattct ttgacttggg tggtcagagt   1380
tttaccagag cctaggactt cactgctctg gtcagaggtc caggtgatgt catcttcttc   1440
aggagtattg caggtgagga ccaccatttc tccgggggca tcaggggtgcc agtccaactc   1500
tacaacataa acgtttttct ccagttccca tat                                  1533

```

```

<210> SEQ ID NO 46
<211> LENGTH: 666
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(666)

<400> SEQUENCE: 46

atg tgc ccg ccg cgc ggc ctc ctc ctt gtg acc atc ctg gtc ctg cta      48
Met Cys Pro Pro Arg Gly Leu Leu Val Thr Ile Leu Val Leu Leu
1          5          10          15

agc cac ctg gac cac ctt act tgg gcc agg agc ctc ccc aca gcc tca      96
Ser His Leu Asp His Leu Thr Trp Ala Arg Ser Leu Pro Thr Ala Ser
20          25          30

```

-continued

ccg agc cca gga ata ttc cag tgc ctc aac cac tcc caa aac ctg ctg	144
Pro Ser Pro Gly Ile Phe Gln Cys Leu Asn His Ser Gln Asn Leu Leu	
35 40 45	
aga gcc gtc agc aac acg ctt cag aag gcc aga caa act cta gaa tta	192
Arg Ala Val Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu Leu	
50 55 60	
tat tcc tgc act tcc gaa gag att gat cat gaa gat atc aca aag gat	240
Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp	
65 70 75 80	
aaa acc agc aca gtg gag gcc tgc tta cca ctg gaa tta acc atg aat	288
Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met Asn	
85 90 95	
gag agt tgc ctg gct tcc aga gag atc tct ttg ata act aac ggg agt	336
Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly Ser	
100 105 110	
tgc ctg gcc tct gga aag gcc tct ttt atg acg gtc ctg tgc ctt agc	384
Cys Leu Ala Ser Gly Lys Ala Ser Phe Met Thr Val Leu Cys Leu Ser	
115 120 125	
agc atc tat gag gac ttg aag atg tac cag atg gaa ttc aag gcc atg	432
Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Met Glu Phe Lys Ala Met	
130 135 140	
aac gca aag ctt tta atg gat ccc aag agg cag atc ttt ctg gat caa	480
Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln	
145 150 155 160	
aac atg ctg aca gct atc gat gag ctg tta cag gcc ctg aat ttc aac	528
Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Phe Asn	
165 170 175	
agt gtg act gtg cca cag aaa tcc tcc ctt gaa gag ccg gat ttt tat	576
Ser Val Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe Tyr	
180 185 190	
aaa act aaa atc aag ctc tgc ata ctt ctt cat gct ttc aga att cgt	624
Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg	
195 200 205	
gcg gtg acc atc gat aga atg atg agt tat ctg aat tct tcc	666
Ala Val Thr Ile Asp Arg Met Met Ser Tyr Leu Asn Ser Ser	
210 215 220	

&lt;210&gt; SEQ ID NO 47

&lt;211&gt; LENGTH: 222

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Canis familiaris

&lt;400&gt; SEQUENCE: 47

Met Cys Pro Pro Arg Gly Leu Leu Leu Val Thr Ile Leu Val Leu Leu	
1 5 10 15	
Ser His Leu Asp His Leu Thr Trp Ala Arg Ser Leu Pro Thr Ala Ser	
20 25 30	
Pro Ser Pro Gly Ile Phe Gln Cys Leu Asn His Ser Gln Asn Leu Leu	
35 40 45	
Arg Ala Val Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu Leu	
50 55 60	
Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp	
65 70 75 80	
Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met Asn	
85 90 95	
Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly Ser	
100 105 110	
Cys Leu Ala Ser Gly Lys Ala Ser Phe Met Thr Val Leu Cys Leu Ser	
115 120 125	

-continued

---

Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Met Glu Phe Lys Ala Met  
 130 135 140

Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln  
 145 150 155 160

Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Phe Asn  
 165 170 175

Ser Val Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe Tyr  
 180 185 190

Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg  
 195 200 205

Ala Val Thr Ile Asp Arg Met Met Ser Tyr Leu Asn Ser Ser  
 210 215 220

<210> SEQ ID NO 48  
 <211> LENGTH: 666  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 48

ggaagaattc agataactca tcattctatc gatggtcacc gcacgaattc tgaaagcatg 60  
 aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggctcttcaa gggaggattt 120  
 ctgtggcaca gtcacactgt tgaaattcag ggcctgtaac agctcatoga tagctgtcag 180  
 catgttttga tccagaaaga tctgcctctt gggatccatt aaaagctttg cgttcatggc 240  
 cttgaattcc atctggtaca tcttcaagtc ctcatagatg ctgctaaggc acaggaccgt 300  
 cataaaagag gcctttccag aggccaggca actcccgtta gttatcaaag agatctctct 360  
 ggaagccagg caactctcat tcatggtaa ttccagtggg aagcaggcct cactgtgct 420  
 ggttttatcc tttgtgatat cttcatgatc aatctcttcg gaagtgcagg aatataattc 480  
 tagagtttgt ctggccttct gaagcgtggt gctgacggct ctcagcagggt tttgggagtg 540  
 gttgaggcac tggaatattc ctgggctcgg tgaggctgtg gggaggctcc tggccaagt 600  
 aaggtggtcc aggtggctta gcaggaccag gatggtcaca aggaggaggc cgcgcggcgg 660  
 gcacat 666

<210> SEQ ID NO 49  
 <211> LENGTH: 591  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(591)

<400> SEQUENCE: 49

agg agc ctc ccc aca gcc tca ccg agc cca gga ata ttc cag tgc ctc 48  
 Arg Ser Leu Pro Thr Ala Ser Pro Ser Pro Gly Ile Phe Gln Cys Leu  
 1 5 10 15

aac cac tcc caa aac ctg ctg aga gcc gtc agc aac acg ctt cag aag 96  
 Asn His Ser Gln Asn Leu Leu Arg Ala Val Ser Asn Thr Leu Gln Lys  
 20 25 30

gcc aga caa act cta gaa tta tat tcc tgc act tcc gaa gag att gat 144  
 Ala Arg Gln Thr Leu Glu Leu Tyr Ser Cys Thr Ser Glu Glu Ile Asp  
 35 40 45

cat gaa gat atc aca aag gat aaa acc agc aca gtg gag gcc tgc tta 192  
 His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu  
 50 55 60

-continued

```

cca ctg gaa tta acc atg aat gag agt tgc ctg gct tcc aga gag atc      240
Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile
65              70              75              80

tct ttg ata act aac ggg agt tgc ctg gcc tct gga aag gcc tct ttt      288
Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Gly Lys Ala Ser Phe
85              90              95

atg acg gtc ctg tgc ctt agc agc atc tat gag gac ttg aag atg tac      336
Met Thr Val Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr
100             105             110

cag atg gaa ttc aag gcc atg aac gca aag ctt tta atg gat ccc aag      384
Gln Met Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys
115             120             125

agg cag atc ttt ctg gat caa aac atg ctg aca gct atc gat gag ctg      432
Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu
130             135             140

tta cag gcc ctg aat ttc aac agt gtg act gtg cca cag aaa tcc tcc      480
Leu Gln Ala Leu Asn Phe Asn Ser Val Thr Val Pro Gln Lys Ser Ser
145             150             155             160

ctt gaa gag ccg gat ttt tat aaa act aaa atc aag ctc tgc ata ctt      528
Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu
165             170             175

ctt cat gct ttc aga att cgt gcg gtg acc atc gat aga atg atg agt      576
Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asp Arg Met Met Ser
180             185             190

tat ctg aat tct tcc
Tyr Leu Asn Ser Ser      591
195
  
```

```

<210> SEQ ID NO 50
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris
  
```

<400> SEQUENCE: 50

```

Arg Ser Leu Pro Thr Ala Ser Pro Ser Pro Gly Ile Phe Gln Cys Leu
1              5              10             15

Asn His Ser Gln Asn Leu Leu Arg Ala Val Ser Asn Thr Leu Gln Lys
20             25             30

Ala Arg Gln Thr Leu Glu Leu Tyr Ser Cys Thr Ser Glu Glu Ile Asp
35             40             45

His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu
50             55             60

Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile
65             70             75             80

Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Gly Lys Ala Ser Phe
85             90             95

Met Thr Val Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr
100            105            110

Gln Met Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys
115            120            125

Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu
130            135            140

Leu Gln Ala Leu Asn Phe Asn Ser Val Thr Val Pro Gln Lys Ser Ser
145            150            155            160

Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu
165            170            175
  
```

-continued

---

Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asp Arg Met Met Ser  
 180 185 190

Tyr Leu Asn Ser Ser  
 195

<210> SEQ ID NO 51  
 <211> LENGTH: 591  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 51

ggaagaattc agataactca tcattctatc gatggtcacc gcacgaattc tgaagcatg 60  
 aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggctcttcaa gggaggattt 120  
 ctgtggcaca gtcacactgt tgaattcag ggcctgtaac agctcatcga tagctgtcag 180  
 catgttttga tccagaaaga tctgcctctt gggatccatt aaaagctttg cgttcatggc 240  
 cttgaattcc atctggtaca tcttcaagtc ctcatagatg ctgctaaggc acaggaccgt 300  
 cataaaaagag gcctttccag aggccaggca actcccgtta gttatcaaag agatctctct 360  
 ggaagccagg caactctcat tcatggttaa ttccagtggg aagcaggcct cactgtgct 420  
 ggttttatcc tttgtgatat cttcatgatc aatctcttcg gaagtgcagg aatataattc 480  
 tagagtttgt ctggccttct gaagcgtggt gctgacggct ctcagcaggg tttgggagtg 540  
 gttgaggcac tggaatattc ctgggctcgg tgaggctgtg gggaggctcc t 591

<210> SEQ ID NO 52  
 <211> LENGTH: 921  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(921)

<400> SEQUENCE: 52

ata tgg gaa ctg gag aaa gat gtt tat gtt gta gag ttg gac tgg cac 48  
 Ile Trp Glu Leu Glu Lys Asp Val Tyr Val Val Glu Leu Asp Trp His  
 1 5 10 15

cct gat gcc ccc gga gaa atg gtg gtc ctc acc tgc cat acc cct gaa 96  
 Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys His Thr Pro Glu  
 20 25 30

gaa gat gac atc act tgg acc tca gcg cag agc agt gaa gtc cta ggt 144  
 Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln Ser Ser Glu Val Leu Gly  
 35 40 45

tct ggt aaa act ctg acc atc caa gtc aaa gaa ttt gga gat gct ggc 192  
 Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Gly Asp Ala Gly  
 50 55 60

cag tat acc tgc cat aaa gga ggc aag gtt ctg agc cgc tca ctc ctg 240  
 Gln Tyr Thr Cys His Lys Gly Gly Lys Val Leu Ser Arg Ser Leu Leu  
 65 70 75 80

ttg att cac aaa aaa gaa gat gga att tgg tcc act gat atc tta aag 288  
 Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys  
 85 90 95

gaa cag aaa gaa tcc aaa aat aag atc ttt ctg aaa tgt gag gca aag 336  
 Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys  
 100 105 110

aat tat tct gga cgt ttc aca tgc tgg tgg ctg acg gca atc agt act 384  
 Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr  
 115 120 125

-continued

gat ttg aaa ttc agt gtc aaa agt agc aga ggc ttc tct gac ccc caa	432
Asp Leu Lys Phe Ser Val Lys Ser Ser Arg Gly Phe Ser Asp Pro Gln	
130 135 140	
ggg gtg aca tgt gga gca gtg aca ctt tca gca gag agg gtc aga gtg	480
Gly Val Thr Cys Gly Ala Val Thr Leu Ser Ala Glu Arg Val Arg Val	
145 150 155 160	
gac aac agg gat tat aag aag tac aca gtg gag tgt cag gag ggc agt	528
Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser	
165 170 175	
gcc tgc ccc tct gcc gag gag agc cta ccc atc gag gtc gtg gtg gat	576
Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp	
180 185 190	
gct att cac aag ctc aag tat gaa aac tac acc agc agc ttc ttc atc	624
Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile	
195 200 205	
aga gac atc atc aaa cca gac cca ccc aca aac ctg cag ctg aag cca	672
Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr Asn Leu Gln Leu Lys Pro	
210 215 220	
ttg aaa aat tct cgg cac gtg gag gtc agc tgg gaa tac ccc gac acc	720
Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr	
225 230 235 240	
tgg agc acc cca cat tcc tac ttc tcc ctg aca ttt tgc ata cag gcc	768
Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Ile Gln Ala	
245 250 255	
cag ggc aag aac aat aga gaa aag aaa gat aga ctc tgc gtg gac aag	816
Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Cys Val Asp Lys	
260 265 270	
acc tca gcc aag gtc gtg tgc cac aag gat gcc aag atc cgc gtg caa	864
Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln	
275 280 285	
gcc cga gac cgc tac tat agt tca tcc tgg agc gac tgg gca tct gtg	912
Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asp Trp Ala Ser Val	
290 295 300	
tcc tgc agt	921
Ser Cys Ser	
305	

<210> SEQ ID NO 53  
 <211> LENGTH: 307  
 <212> TYPE: PRT  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 53

Ile Trp Glu Leu Glu Lys Asp Val Tyr Val Val Glu Leu Asp Trp His	
1 5 10 15	
Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys His Thr Pro Glu	
20 25 30	
Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln Ser Ser Glu Val Leu Gly	
35 40 45	
Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Gly Asp Ala Gly	
50 55 60	
Gln Tyr Thr Cys His Lys Gly Gly Lys Val Leu Ser Arg Ser Leu Leu	
65 70 75 80	
Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys	
85 90 95	
Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys	
100 105 110	
Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr	
115 120 125	

-continued

Asp Leu Lys Phe Ser Val Lys Ser Ser Arg Gly Phe Ser Asp Pro Gln  
 130 135 140

Gly Val Thr Cys Gly Ala Val Thr Leu Ser Ala Glu Arg Val Arg Val  
 145 150 155 160

Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser  
 165 170 175

Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp  
 180 185 190

Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile  
 195 200 205

Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr Asn Leu Gln Leu Lys Pro  
 210 215 220

Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr  
 225 230 235 240

Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Ile Gln Ala  
 245 250 255

Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Cys Val Asp Lys  
 260 265 270

Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln  
 275 280 285

Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asp Trp Ala Ser Val  
 290 295 300

Ser Cys Ser  
 305

<210> SEQ ID NO 54  
 <211> LENGTH: 921  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris  
 <400> SEQUENCE: 54

actgcaggac acagatgccc agtcgctcca ggatgaacta tagtagcggg ctcgggcttg 60

cacgcggatc ttggcaccct tgtggcacac gacctggct gaggtcttgt ccacgcagag 120

tctatctttc ttttctctat tgttcttgcc ctgggcctgt atgcaaaatg tcagggagaa 180

gtaggaatgt ggggtgctcc aggtgtcggg gtattcccag ctgacctcca cgtgccgaga 240

atTTTTcaat ggcttcagct gcaggtttgt ggggtgggtct ggtttgatga tgtctctgat 300

gaagaagctg ctgggtgtagt tttcatactt gagcttgtga atagcatcca ccacgacctc 360

gatgggtagg ctctcctcgg cagaggggca ggcactgccc tcctgacact ccactgtgta 420

cttcttataa tcctctgtgt ccaactctgac cctctctgct gaaagtgtca ctgctccaca 480

tgtcaccctc tgggggtcag agaagcctct gctacttttg aactgaatt tcaaatcagt 540

actgattgcc gtcagccacc agcatgtgaa acgtccagaa taattctttg cctcacattt 600

cagaaagatc ttatttttgg attctttctg ttcctttaag atatcagtgg accaaattcc 660

atcttctttt ttgtgaatca acaggagtga gcggtcaga acctgcctc ctttatggca 720

ggtatactgg ccagcatctc caaattcttt gacttggatg gtcagagttt taccagaacc 780

taggacttca ctgctctcgc ctgagggtcca agtgatgtca tcttcttcag gggtagggca 840

ggtgaggacc accatttctc cgggggcatc aggggtccag tccaactcta caacataaac 900

atctttctcc agttcccata t 921

-continued

---

```

<210> SEQ ID NO 55
<211> LENGTH: 985
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(984)

<400> SEQUENCE: 55

atg cat cct cag cag ttg gtc atc gcc tgg ttt tcc ctg gtt ttg ctg      48
Met His Pro Gln Leu Val Ile Ala Trp Phe Ser Leu Val Leu Leu
1          5          10          15

gca cct ccc ctc atg gcc ata tgg gaa ctg gag aaa aac gtt tat gtt      96
Ala Pro Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asn Val Tyr Val
          20          25          30

gta gag ttg gac tgg cac cct gat gcc ccc gga gaa atg gtg gtc ctc     144
Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
          35          40          45

acc tgc aat act cct gaa gaa gat gac atc acc tgg acc tct gac cag     192
Thr Cys Asn Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln
          50          55          60

agc agt gaa gtc cta ggc tct ggt aaa act ctg acc atc caa gtc aaa     240
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
65          70          75          80

gaa ttt gca gat gct ggc cag tat acc tgt cat aaa gga ggc gag gtt     288
Glu Phe Ala Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
          85          90          95

ctg agc cat tcg ttc ctc ctg ata cac aaa aag gaa gat gga att tgg     336
Leu Ser His Ser Phe Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
          100         105         110

tcc act gat atc tta agg gaa cag aaa gaa tcc aaa aat aag atc ttt     384
Ser Thr Asp Ile Leu Arg Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
          115         120         125

cta aaa tgt gag gca aag aat tat tct gga cgt ttc acc tgc tgg tgg     432
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
          130         135         140

ctg acg gca atc agt acc gat ttg aaa ttc act gtc aaa agc agc aga     480
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Thr Val Lys Ser Ser Arg
          145         150         155         160

ggc tcc tet gac ccc caa ggg gtg act tgt gga gca gcg aca ctc tca     528
Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser
          165         170         175

gca gag aag gtc aga gtg gac aac agg gat tat aag aag tac aca gtg     576
Ala Glu Lys Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val
          180         185         190

gag tgt cag gag ggc agt gcc tgc ccg gct gcc gag gag agc cta ccc     624
Glu Cys Gln Glu Gly Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro
          195         200         205

att gaa gtc gtg gtg gac gct att cac aag ctc aag tac gaa aac tac     672
Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr
          210         215         220

acc agc agc ttc ttc atc agg gac atc atc aaa ccg gac cca ccc aag     720
Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys
          225         230         235         240

aac ctg caa ctg aag cca tta aaa aat tct cgg cat gtg gaa gtg agc     768
Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser
          245         250         255

tgg gaa tac cct gac acc tgg agc acc cca cat tcc tac ttc tcc tta     816
Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu
          260         265         270

```

-continued

---

```

aca ttt ggc gta cag gtc cag ggc aag aac aac aga gaa aag aaa gac      864
Thr Phe Gly Val Gln Val Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp
      275                      280                      285

aga ctc tcc gtg gac aag acc tca gcc aag gtc gtg tgc cac aag gat      912
Arg Leu Ser Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp
      290                      295                      300

gcc aag atc cgc gtg caa gcc aga gac cgc tac tat agc tca tcc tgg      960
Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp
      305                      310                      315                      320

agc aac tgg gca tcc gtg tcc tgc a      985
Ser Asn Trp Ala Ser Val Ser Cys
      325

<210> SEQ ID NO 56
<211> LENGTH: 328
<212> TYPE: PRT
<213> ORGANISM: Felis catus

<400> SEQUENCE: 56

Met His Pro Gln Gln Leu Val Ile Ala Trp Phe Ser Leu Val Leu Leu
 1          5          10          15

Ala Pro Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asn Val Tyr Val
 20          25          30

Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
 35          40          45

Thr Cys Asn Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln
 50          55          60

Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
 65          70          75          80

Glu Phe Ala Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
 85          90          95

Leu Ser His Ser Phe Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
 100         105         110

Ser Thr Asp Ile Leu Arg Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
 115         120         125

Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
 130         135         140

Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Thr Val Lys Ser Ser Arg
 145         150         155         160

Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser
 165         170         175

Ala Glu Lys Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val
 180         185         190

Glu Cys Gln Glu Gly Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro
 195         200         205

Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr
 210         215         220

Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys
 225         230         235         240

Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser
 245         250         255

Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu
 260         265         270

Thr Phe Gly Val Gln Val Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp
 275         280         285

```

-continued

Arg Leu Ser Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp  
 290 295 300  
 Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp  
 305 310 315 320  
 Ser Asn Trp Ala Ser Val Ser Cys  
 325

<210> SEQ ID NO 57  
 <211> LENGTH: 985  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 57

tgcaggacac ggatgccag ttgctccagg atgagctata gtgagcgtct ctggcttgca 60  
 cgcggatcctt ggcatccttg tggcacacga ccttggctga ggtcttgtcc acggagagtc 120  
 tgtctttctt ttctctgttg ttcttgcctt ggacctgtac gccaaatggt aaggagaagt 180  
 aggaatgtgg ggtgctccag gtgtcagggt attcccagct cacttccaca tgccgagaat 240  
 tttttaatgg cttcagttgc aggttcttgg gtgggtccgg tttgatgatg tcctgatga 300  
 agaagctgct ggtgtagttt tcgtacttga gcttgtgaat agcgtccacc acgacttcaa 360  
 tgggtaggct ctctcggca gccgggcagg cactgccctc ctgacactcc actgtgtact 420  
 tcttataatc cctggtgtcc actctgacct tctctgctga gagtgtcgtc gctccacaag 480  
 tcaccccttg ggggtcagag gagcctctgc tgcttttgac agtgaatttc aaatcggtac 540  
 tgattgccgt cagccaccag caggtgaaac gtccagaata attctttgcc tcacatttta 600  
 gaaagatcctt atttttggat tctttctggt cccttaagat atcagtggtc caaattccat 660  
 ctctcttttt gtgtatcagg aggaacgaat ggctcagaac ctgcctcctt ttatgacagg 720  
 tatactggcc agcatctgca aattctttga cttggatggt cagagtttta ccagagccta 780  
 ggacttcaact gctctggtca gaggtccagg tgatgtcatc ttcttcagga gtattgcagg 840  
 tgaggaccac catttctccg ggggcatcag ggtgccagtc caactctaca acataaacgt 900  
 ttttctccag ttcccatatg gccatgaggg gaggtgccag caaaaccagg gaaaaccagg 960  
 cgatgaccaa ctgctgagga tgcat 985

<210> SEQ ID NO 58  
 <211> LENGTH: 987  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(987)

<400> SEQUENCE: 58

atg cac cct cag cag ttg gtc atc tcc tgg ttt tcc ctc gtt ttg ctg 48  
 Met His Pro Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Leu Leu  
 1 5 10 15  
 gcg tct ccc ctc atg gcc ata tgg gaa ctg gag aaa gat gtt tat gtt 96  
 Ala Ser Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asp Val Tyr Val  
 20 25 30  
 gta gag ttg gac tgg cac cct gat gcc ccc gga gaa atg gtg gtc ctc 144  
 Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu  
 35 40 45  
 acc tgc cat acc cct gaa gaa gat gac atc act tgg acc tca gcg cag 192  
 Thr Cys His Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln  
 50 55 60

-continued

agc agt gaa gtc cta ggt tct ggt aaa act ctg acc atc caa gtc aaa	240
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys	
65 70 75 80	
gaa ttt gga gat gct ggc cag tat acc tgc cat aaa gga ggc aag gtt	288
Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Lys Val	
85 90 95	
ctg agc cgc tca ctc ctg ttg att cac aaa aaa gaa gat gga att tgg	336
Leu Ser Arg Ser Leu Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp	
100 105 110	
tcc act gat atc tta aag gaa cag aaa gaa tcc aaa aat aag atc ttt	384
Ser Thr Asp Ile Leu Lys Glu Lys Glu Ser Lys Asn Lys Ile Phe	
115 120 125	
ctg aaa tgt gag gca aag aat tat tct gga cgt ttc aca tgc tgg tgg	432
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp	
130 135 140	
ctg acg gca atc agt act gat ttg aaa ttc agt gtc aaa agt agc aga	480
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Ser Val Lys Ser Ser Arg	
145 150 155 160	
ggc ttc tct gac ccc caa ggg gtg aca tgt gga gca gtg aca ctt tca	528
Gly Phe Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Val Thr Leu Ser	
165 170 175	
gca gag agg gtc aga gtg gac aac agg gat tat aag aag tac aca gtg	576
Ala Glu Arg Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val	
180 185 190	
gag tgt cag gag ggc agt gcc tgc ccc tct gcc gag gag agc cta ccc	624
Glu Cys Gln Glu Gly Ser Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro	
195 200 205	
atc gag gtc gtg gtg gat gct att cac aag ctc aag tat gaa aac tac	672
Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr	
210 215 220	
acc agc agc ttc ttc atc aga gac atc atc aaa cca gac cca ccc aca	720
Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr	
225 230 235 240	
aac ctg cag ctg aag cca ttg aaa aat tct cgg cac gtg gag gtc agc	768
Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser	
245 250 255	
tgg gaa tac ccc gac acc tgg agc acc cca cat tcc tac ttc tcc ctg	816
Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu	
260 265 270	
aca ttt tgc ata cag gcc cag ggc aag aac aat aga gaa aag aaa gat	864
Thr Phe Cys Ile Gln Ala Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp	
275 280 285	
aga ctc tgc gtg gac aag acc tca gcc aag gtc gtg tgc cac aag gat	912
Arg Leu Cys Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp	
290 295 300	
gcc aag atc cgc gtg caa gcc cga gac cgc tac tat agt tca tcc tgg	960
Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp	
305 310 315 320	
agc gac tgg gca tct gtg tca tgc agt	987
Ser Asp Trp Ala Ser Val Ser Cys Ser	
325	

&lt;210&gt; SEQ ID NO 59

&lt;211&gt; LENGTH: 329

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Canis familiaris

-continued

<400> SEQUENCE: 59

Met His Pro Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Leu Leu  
 1 5 10 15  
 Ala Ser Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asp Val Tyr Val  
 20 25 30  
 Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu  
 35 40 45  
 Thr Cys His Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln  
 50 55 60  
 Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys  
 65 70 75 80  
 Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Lys Val  
 85 90 95  
 Leu Ser Arg Ser Leu Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp  
 100 105 110  
 Ser Thr Asp Ile Leu Lys Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe  
 115 120 125  
 Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp  
 130 135 140  
 Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Ser Val Lys Ser Ser Arg  
 145 150 155 160  
 Gly Phe Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Val Thr Leu Ser  
 165 170 175  
 Ala Glu Arg Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val  
 180 185 190  
 Glu Cys Gln Glu Gly Ser Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro  
 195 200 205  
 Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr  
 210 215 220  
 Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr  
 225 230 235 240  
 Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser  
 245 250 255  
 Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu  
 260 265 270  
 Thr Phe Cys Ile Gln Ala Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp  
 275 280 285  
 Arg Leu Cys Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp  
 290 295 300  
 Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp  
 305 310 315 320  
 Ser Asp Trp Ala Ser Val Ser Cys Ser  
 325

<210> SEQ ID NO 60  
 <211> LENGTH: 987  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 60

actgcatgac acagatgcc agtcgctcca ggtatgaacta tagtagcggg ctcgggcttg 60  
 cacgcgatc ttggcatcct tgtggcacac gaccttggt gaggtcttgt ccacgcagag 120  
 tctatctttc ttttctctat tgttcttgcc ctgggcctgt atgcaaatg tcagggagaa 180

-continued

```

gtagaaatgt ggggtgctcc aggtgtcggg gtattcccag ctgacctcca cgtgccgaga 240
atTTTTcaat ggcttcagct gcaggtttgt gggTgggtct ggTttgatga tgtctctgat 300
gaagaagctg ctggTgtagt tttcatactt gagcttTtga atagcatcca ccacgacctc 360
gatgggtagg ctctcctcgg cagaggggca ggcactgccc tcctgacact ccactgtgta 420
cttcttataa tcctctgttTt ccactctgac cctctctgct gaaagtTtca ctgctccaca 480
tgtcaccctc tgggggtcag agaagcctct gctacttttTg aactgaatt tcaaatcagt 540
actgattgcc gtcagccacc agcatgtgaa acgtccagaa taattctttTg cctcacattt 600
cagaaagatc ttatttttTg attctttctg ttcctttaag atatcagtTg accaaattcc 660
atcttctttt ttgtgaatca acaggagtga gcggctcaga acctTgcctc ctttatTgca 720
ggTactactg ccagcatctc caaattcttt gactTggatg gtcagagttt taccagaacc 780
taggacttca ctgctctcgg ctgaggtcca agtgatTtca tcttcttcag gggTatTgca 840
ggTgaggacc accatttctc cgggggcacTc agggTgccag tccaactcta caacataaac 900
atctttctcc agtTcccata Tggccatgag gggagacgcc agcaaaacga gggaaaacca 960
ggagatgacc aactgctgag ggtgcat 987

```

```

<210> SEQ ID NO 61
<211> LENGTH: 1599
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1599)

```

```

<400> SEQUENCE: 61
atg cac cct cag cag ttg gtc atc tcc tgg ttt tcc ctc gtt ttg ctg 48
Met His Pro Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Leu Leu
1 5 10 15
gcg tct ccc ctc atg gcc ata tgg gaa ctg gag aaa gat gtt tat gtt 96
Ala Ser Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asp Val Tyr Val
20 25 30
gta gag ttg gac tgg cac cct gat gcc ccc gga gaa atg gtg gtc ctc 144
Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
35 40 45
acc tgc cat acc cct gaa gaa gat gac atc act tgg acc tca gcg cag 192
Thr Cys His Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln
50 55 60
agc agt gaa gtc cta ggt tct ggt aaa act ctg acc atc caa gtc aaa 240
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
65 70 75 80
gaa ttt gga gat gct ggc cag tat acc tgc cat aaa gga ggc aag gtt 288
Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Lys Val
85 90 95
ctg agc cgc tca ctc ctg ttg att cac aaa aaa gaa gat gga att tgg 336
Leu Ser Arg Ser Leu Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
100 105 110
tcc act gat atc tta aag gaa cag aaa gaa tcc aaa aat aag atc ttt 384
Ser Thr Asp Ile Leu Lys Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
115 120 125
ctg aaa tgt gag gca aag aat tat tct gga cgt ttc aca tgc tgg tgg 432
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
130 135 140
ctg acg gca atc agt act gat ttg aaa ttc agt gtc aaa agt agc aga 480
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Ser Val Lys Ser Ser Arg
145 150 155 160

```

-continued

ggc ttc tet gac ccc caa ggg gtg aca tgt gga gca gtg aca ctt tca Gly Phe Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Val Thr Leu Ser 165 170 175	528
gca gag agg gtc aga gtg gac aac agg gat tat aag aag tac aca gtg Ala Glu Arg Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val 180 185 190	576
gag tgt cag gag ggc agt gcc tgc ccc tct gcc gag gag agc cta ccc Glu Cys Gln Glu Gly Ser Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro 195 200 205	624
atc gag gtc gtg gtg gat gct att cac aag ctc aag tat gaa aac tac Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr 210 215 220	672
acc agc agc ttc ttc atc aga gac atc atc aaa cca gac cca ccc aca Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr 225 230 235 240	720
aac ctg cag ctg aag cca ttg aaa aat tct cgg cac gtg gag gtc agc Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser 245 250 255	768
tgg gaa tac ccc gac acc tgg agc acc cca cat tcc tac ttc tcc ctg Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu 260 265 270	816
aca ttt tgc ata cag gcc cag ggc aag aac aat aga gaa aag aaa gat Thr Phe Cys Ile Gln Ala Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp 275 280 285	864
aga ctc tgc gtg gac aag acc tca gcc aag gtc gtg tgc cac aag gat Arg Leu Cys Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp 290 295 300	912
gcc aag atc cgc gtg caa gcc cga gac cgc tac tat agt tca tcc tgg Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp 305 310 315 320	960
agc gac tgg gca tct gtg tca tgc agt ggt ggc ggt ggc ggc gga tct Ser Asp Trp Ala Ser Val Ser Cys Ser Gly Gly Gly Gly Gly Ser 325 330 335	1008
aga aac ttg cca acc cct act cca tcc ccg ggt atg ttc caa tgt ttg Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro Gly Met Phe Gln Cys Leu 340 345 350	1056
aac cac tcc caa acc ttg ttg aga gcc gtc agc aac acg ctt cag aag Asn His Ser Gln Thr Leu Leu Arg Ala Val Ser Asn Thr Leu Gln Lys 355 360 365	1104
gcc aga caa act cta gaa tta tat tcc tgc act tcc gaa gag att gat Ala Arg Gln Thr Leu Glu Leu Tyr Ser Cys Thr Ser Glu Glu Ile Asp 370 375 380	1152
cat gaa gat atc aca aag gat aaa acc agc aca gtg gag gcc tgc tta His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu 385 390 395 400	1200
cca ctg gaa tta acc atg aat gag agt tgc ctg gct tcc aga gag atc Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile 405 410 415	1248
tct ttg ata act aac ggg agt tgc ctg gcc tct gga aag gcc tct ttt Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Gly Lys Ala Ser Phe 420 425 430	1296
atg acg gtc ctg tgc ctt agc agc atc tat gag gac ttg aag atg tac Met Thr Val Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr 435 440 445	1344
cag atg gaa ttc aag gcc atg aac gca aag ctt tta atg gat ccc aag Gln Met Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys 450 455 460	1392
agg cag atc ttt ctg gat caa aac atg ctg aca gct atc gat gag ctg Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu 465 470 475 480	1440

-continued

tta cag gcc ctg aat ttc aac agt gtg act gtg cca cag aaa tcc tcc	1488
Leu Gln Ala Leu Asn Phe Asn Ser Val Thr Val Pro Gln Lys Ser Ser	
485 490 495	
ctt gaa gag ccg gat ttt tat aaa act aaa atc aag ctc tgc ata ctt	1536
Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu	
500 505 510	
ctt cat gct ttc aga att cgt gcg gtg acc atc aat aga atg atg tcc	1584
Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asn Arg Met Met Ser	
515 520 525	
tac ttg aac tct tcc	1599
Tyr Leu Asn Ser Ser	
530	

<210> SEQ ID NO 62  
 <211> LENGTH: 533  
 <212> TYPE: PRT  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 62

Met His Pro Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Leu Leu	1 5 10 15
Ala Ser Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asp Val Tyr Val	20 25 30
Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu	35 40 45
Thr Cys His Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln	50 55 60
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys	65 70 75 80
Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Lys Val	85 90 95
Leu Ser Arg Ser Leu Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp	100 105 110
Ser Thr Asp Ile Leu Lys Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe	115 120 125
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp	130 135 140
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Ser Val Lys Ser Ser Arg	145 150 155 160
Gly Phe Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Val Thr Leu Ser	165 170 175
Ala Glu Arg Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val	180 185 190
Glu Cys Gln Glu Gly Ser Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro	195 200 205
Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr	210 215 220
Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr	225 230 235 240
Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser	245 250 255
Trp Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu	260 265 270
Thr Phe Cys Ile Gln Ala Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp	275 280 285

-continued

---

Arg Leu Cys Val Asp Lys Thr Ser Ala Lys Val Val Cys His Lys Asp  
 290 295 300

Ala Lys Ile Arg Val Gln Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp  
 305 310 315 320

Ser Asp Trp Ala Ser Val Ser Cys Ser Gly Gly Gly Gly Gly Ser  
 325 330 335

Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro Gly Met Phe Gln Cys Leu  
 340 345 350

Asn His Ser Gln Thr Leu Leu Arg Ala Val Ser Asn Thr Leu Gln Lys  
 355 360 365

Ala Arg Gln Thr Leu Glu Leu Tyr Ser Cys Thr Ser Glu Glu Ile Asp  
 370 375 380

His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu  
 385 390 395 400

Pro Leu Glu Leu Thr Met Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile  
 405 410 415

Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala Ser Gly Lys Ala Ser Phe  
 420 425 430

Met Thr Val Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr  
 435 440 445

Gln Met Glu Phe Lys Ala Met Asn Ala Lys Leu Leu Met Asp Pro Lys  
 450 455 460

Arg Gln Ile Phe Leu Asp Gln Asn Met Leu Thr Ala Ile Asp Glu Leu  
 465 470 475 480

Leu Gln Ala Leu Asn Phe Asn Ser Val Thr Val Pro Gln Lys Ser Ser  
 485 490 495

Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu  
 500 505 510

Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asn Arg Met Met Ser  
 515 520 525

Tyr Leu Asn Ser Ser  
 530

<210> SEQ ID NO 63  
 <211> LENGTH: 1599  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 63

```

ggaagagttc aagtaggaca tcattctatt gatggtcacc gcacgaattc tgaagcatg      60
aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggctcttcaa gggaggattt      120
ctgtggcaca gtcacactgt tgaattcag ggcctgtaac agctcatoga tagctgtcag      180
catgttttga tccagaaaga tctgcctctt gggatccatt aaaagctttg cgttcatggc      240
cttgaattcc atctgttaca tcttcaagtc ctcatagatg ctgctaaggc acaggaccgt      300
cataaaaagag gcctttccag aggccaggca actcccgtta gttatcaaag agatctctct      360
ggaagccagg caactctcat tcatgggtaa ttccagtggg aagcaggcct cactgtgct      420
ggttttatcc tttgtgatat cttcatgatc aatctctctg gaagtgcagg aatataattc      480
tagagtttgt ctggccttct gaagcgtggt gotgacggct ctcaacaagg tttgggagtg      540
gttcaaacat tggaacatac ccggggatgg agtaggggtt ggcaagtttc tagatccgcc      600
gccaccgcca cactgcatg acacagatgc ccagtcgctc caggatgaac tatagtagcg      660
gtctcgggct tgcacgggga tcttggcatc cttgtggcac acgaccttg ctgaggtctt      720
    
```

-continued

---

```

gtccacgcag agtctatctt tcttttctct attgttcttg ccctgggcct gtagcaaaa 780
tgtcaggag agtaggaat gtggggtgct ccagggtgctg gggattccc agctgacctc 840
cacgtgccga gaatTTTTca atggcttcag ctgcaggttt gtgggtgggt ctggtttgat 900
gatgtctctg atgaagaagc tgctgggtga gttttcatac ttgagcttgt gaatagcate 960
caccacgacc tcgatgggta ggctctcctc ggcagagggg caggcactgc cctcctgaca 1020
ctccactgtg tacttcttat aatccctggt gtccactctg accctctctg ctgaaagtgt 1080
cactgctcca catgtcacc cttgggggtc agagaagcct ctgctacttt tgacactgaa 1140
tttcaaatca gtactgattg ccgtcagcca ccagcatgtg aaactccag aataattctt 1200
tgctcacat ttcagaaaga tcttattttt ggattctttc tgttcttta agatatcagt 1260
ggaccaaatt ccatcttctt ttttgtgaat caacaggagt gagcggtca gaaccttgcc 1320
tcctttatgg caggataact gccagcatc tccaaattct ttgacttggg tggtcagagt 1380
tttaccagaa ctaggactt cactgctctg cgctgaggtc caagtgatgt catcttcttc 1440
aggggtatgg caggtgagga ccaccatttc tccgggggca tcagggtgcc agtccaactc 1500
tacaacataa acatctttct ccagttccca tatggccatg aggggagacg ccagcaaac 1560
gagggaaaac caggagatga ccaactgctg aggggtgat 1599

```

```

<210> SEQ ID NO 64
<211> LENGTH: 0
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```
<400> SEQUENCE: 64
```

```
000
```

```

<210> SEQ ID NO 65
<211> LENGTH: 0
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```
<400> SEQUENCE: 65
```

```
000
```

```

<210> SEQ ID NO 66
<211> LENGTH: 1533
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1533)

```

```
<400> SEQUENCE: 66
```

```

ata tgg gaa ctg gag aaa gat gtt tat gtt gta gag ttg gac tgg cac 48
Ile Trp Glu Leu Glu Lys Asp Val Tyr Val Val Glu Leu Asp Trp His
1 5 10 15
cct gat gcc ccc gga gaa atg gtg gtc ctc acc tgc cat acc cct gaa 96
Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys His Thr Pro Glu
20 25 30
gaa gat gac atc act tgg acc tca gcg cag agc agt gaa gtc cta ggt 144
Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln Ser Ser Glu Val Leu Gly
35 40 45
tct ggt aaa act ctg acc atc caa gtc aaa gaa ttt gga gat gct ggc 192
Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Gly Asp Ala Gly
50 55 60

```

-continued

cag tat acc tgc cat aaa gga ggc aag gtt ctg agc cgc tca ctc ctg Gln Tyr Thr Cys His Lys Gly Gly Lys Val Leu Ser Arg Ser Leu Leu 65 70 75 80	240
ttg att cac aaa aaa gaa gat gga att tgg tcc act gat atc tta aag Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys 85 90 95	288
gaa cag aaa gaa tcc aaa aat aag atc ttt ctg aaa tgt gag gca aag Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys 100 105 110	336
aat tat tct gga cgt ttc aca tgc tgg tgg ctg acg gca atc agt act Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr 115 120 125	384
gat ttg aaa ttc agt gtc aaa agt agc aga ggc ttc tct gac ccc caa Asp Leu Lys Phe Ser Val Lys Ser Ser Arg Gly Phe Ser Asp Pro Gln 130 135 140	432
ggg gtg aca tgt gga gca gtg aca ctt tca gca gag agg gtc aga gtg Gly Val Thr Cys Gly Ala Val Thr Leu Ser Ala Glu Arg Val Arg Val 145 150 155 160	480
gac aac agg gat tat aag aag tac aca gtg gag tgt cag gag ggc agt Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser 165 170 175	528
gcc tgc ccc tct gcc gag gag agc cta ccc atc gag gtc gtg gtg gat Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp 180 185 190	576
gct att cac aag ctc aag tat gaa aac tac acc agc agc ttc ttc atc Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile 195 200 205	624
aga gac atc atc aaa cca gac cca ccc aca aac ctg cag ctg aag cca Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr Asn Leu Gln Leu Lys Pro 210 215 220	672
ttg aaa aat tct cgg cac gtg gag gtc agc tgg gaa tac ccc gac acc Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr 225 230 235 240	720
tgg agc acc cca cat tcc tac ttc tcc ctg aca ttt tgc ata cag gcc Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Ile Gln Ala 245 250 255	768
cag gcc aag aac aat aga gaa aag aaa gat aga ctc tgc gtg gac aag Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Cys Val Asp Lys 260 265 270	816
acc tca gcc aag gtc gtg tgc cac aag gat gcc aag atc cgc gtg caa Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln 275 280 285	864
gcc cga gac cgc tac tat agt tca tcc tgg agc gac tgg gca tct gtg Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asp Trp Ala Ser Val 290 295 300	912
tca tgc agt ggt ggc ggt ggc ggc gga tct aga aac ttg cca acc cct Ser Cys Ser Gly Gly Gly Gly Gly Ser Arg Asn Leu Pro Thr Pro 305 310 315 320	960
act cca tcc ccg ggt atg ttc caa tgt ttg aac cac tcc caa acc ttg Thr Pro Ser Pro Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu 325 330 335	1008
ttg aga gcc gtc agc aac acg ctt cag aag gcc aga caa act cta gaa Leu Arg Ala Val Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu 340 345 350	1056
tta tat tcc tgc act tcc gaa gag att gat cat gaa gat atc aca aag Leu Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys 355 360 365	1104
gat aaa acc agc aca gtg gag gcc tgc tta cca ctg gaa tta acc atg Asp Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met 370 375 380	1152

-continued

aat gag agt tgc ctg gct tcc aga gag atc tct ttg ata act aac ggg	1200
Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly	
385 390 395 400	
agt tgc ctg gcc tct gga aag gcc tct ttt atg acg gtc ctg tgc ctt	1248
Ser Cys Leu Ala Ser Gly Lys Ala Ser Phe Met Thr Val Leu Cys Leu	
405 410 415	
agc agc atc tat gag gac ttg aag atg tac cag atg gaa ttc aag gcc	1296
Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Met Glu Phe Lys Ala	
420 425 430	
atg aac gca aag ctt tta atg gat ccc aag agg cag atc ttt ctg gat	1344
Met Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp	
435 440 445	
caa aac atg ctg aca gct atc gat gag ctg tta cag gcc ctg aat ttc	1392
Gln Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Phe	
450 455 460	
aac agt gtg act gtg cca cag aaa tcc tcc ctt gaa gag cag gat ttt	1440
Asn Ser Val Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe	
465 470 475 480	
tat aaa act aaa atc aag ctc tgc ata ctt ctt cat gct ttc aga att	1488
Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile	
485 490 495	
cgt gcg gtg acc atc aat aga atg atg tcc tac ttg aac tct tcc	1533
Arg Ala Val Thr Ile Asn Arg Met Met Ser Tyr Leu Asn Ser Ser	
500 505 510	

<210> SEQ ID NO 67  
 <211> LENGTH: 511  
 <212> TYPE: PRT  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 67

Ile Trp Glu Leu Glu Lys Asp Val Tyr Val Val Glu Leu Asp Trp His	
1 5 10 15	
Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys His Thr Pro Glu	
20 25 30	
Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln Ser Ser Glu Val Leu Gly	
35 40 45	
Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Gly Asp Ala Gly	
50 55 60	
Gln Tyr Thr Cys His Lys Gly Gly Lys Val Leu Ser Arg Ser Leu Leu	
65 70 75 80	
Leu Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys	
85 90 95	
Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys	
100 105 110	
Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr	
115 120 125	
Asp Leu Lys Phe Ser Val Lys Ser Ser Arg Gly Phe Ser Asp Pro Gln	
130 135 140	
Gly Val Thr Cys Gly Ala Val Thr Leu Ser Ala Glu Arg Val Arg Val	
145 150 155 160	
Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser	
165 170 175	
Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp	
180 185 190	
Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile	
195 200 205	

-continued

---

Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr Asn Leu Gln Leu Lys Pro  
 210 215 220

Leu Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr  
 225 230 235 240

Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Ile Gln Ala  
 245 250 255

Gln Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Cys Val Asp Lys  
 260 265 270

Thr Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln  
 275 280 285

Ala Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asp Trp Ala Ser Val  
 290 295 300

Ser Cys Ser Gly Gly Gly Gly Gly Ser Arg Asn Leu Pro Thr Pro  
 305 310 315 320

Thr Pro Ser Pro Gly Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu  
 325 330 335

Leu Arg Ala Val Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu  
 340 345 350

Leu Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys  
 355 360 365

Asp Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met  
 370 375 380

Asn Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly  
 385 390 395 400

Ser Cys Leu Ala Ser Gly Lys Ala Ser Phe Met Thr Val Leu Cys Leu  
 405 410 415

Ser Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Met Glu Phe Lys Ala  
 420 425 430

Met Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp  
 435 440 445

Gln Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Phe  
 450 455 460

Asn Ser Val Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe  
 465 470 475 480

Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile  
 485 490 495

Arg Ala Val Thr Ile Asn Arg Met Met Ser Tyr Leu Asn Ser Ser  
 500 505 510

<210> SEQ ID NO 68  
 <211> LENGTH: 1533  
 <212> TYPE: DNA  
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 68

ggaagagttc aagtaggaca tcattctatt gatggtcacc gcacgaattc tgaagcatg 60  
 aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggctcttcaa gggaggattt 120  
 ctgtggcaca gtcacactgt tgaaattcag ggcctgtaac agctcatcga tagctgtcag 180  
 catgttttga tccagaaaga tctgcctctt gggatccatt aaaagctttg cgttcatggc 240  
 cttgaattcc atctggtaca tcttcaagtc ctcatagatg ctgctaaggc acaggaccgt 300  
 cataaaagag gcctttccag aggccaggca actcccgtta gttatcaaag agatctctct 360  
 ggaagccagg caactctcat tcatggttaa ttccagtggg aagcaggcct ccaactgtgct 420

-continued

---

```

ggttttatcc tttgtgatat cttcatgac aatctcttcg gaagtgcagg aatataattc 480
tagagtttgt ctggccttct gaagcgtggt gctgacggct ctcaacaagg tttgggagtg 540
gttcaaacat tggaacatac cgggggatgg agtaggggtt ggcaagtttc tagatccgcc 600
gccaccgcc cactgcatg acacagatgc ccagtcgctc caggatgaac tatagtagcg 660
gtctcgggct tgcacgcgga tcttggcatc cttgtggcac acgacctgg ctgaggtcct 720
gtccacgcag agtctatctt tcttttctct attgttcttg ccctgggcct gtatgcaaaa 780
tgtcagggag aagtaggaat gtgggggtct ccagggtctg gggattccc agctgacctc 840
cacgtgccga gaatthttca atggcttcag ctgcaggttt gtgggtgggt ctggtttgat 900
gatgtctctg atgaagaagc tgctgggtga gttttcatac ttgagcttgt gaatagcatc 960
caccacgacc tcgatgggta ggctctcctc ggcagagggg caggcactgc cctcctgaca 1020
ctccactgtg tacttcttat aatccctggt gtccactctg accctctctg ctgaaagtgt 1080
cactgctcca catgtcacc cttgggggtc agagaagcct ctgctacttt tgacactgaa 1140
ttcaaatca gtactgattg ccgtcagcca ccagcatgtg aaactccag aataattctt 1200
tgctcaccat ttcagaaaga tcttattttt ggattctttc tgttccttta agatatcagt 1260
ggaccaaatt ccatcttctt ttttgtgaat caacaggagt gagcggctca gaacctgcc 1320
tcctttatgg caggataact ggccagcatc tccaaattct ttgacttga tggtcagagt 1380
ttaccagaa ctaggactt cactgctctg cgctgaggtc caagtgatgt catcttcttc 1440
aggggtatgg caggtgagga ccaccatttc tccgggggca tcagggtgcc agtccaactc 1500
tacaacataa acatctttct ccagttccca tat 1533

```

```

<210> SEQ ID NO 69
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

```

```
<400> SEQUENCE: 69
```

```
gccaagctcg aaattaacc tcactaaagg 30
```

```

<210> SEQ ID NO 70
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

```

```
<400> SEQUENCE: 70
```

```
cgacggccag tgaattgtaa tacgactc 28
```

```

<210> SEQ ID NO 71
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

```

```
<400> SEQUENCE: 71
```

```
agtgatgaag gcctggaatc agattacttt g 31
```

```

<210> SEQ ID NO 72
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence

```

-continued

---

```

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 72

atggcctgga acacttctct gaaagaatat ga                               32

<210> SEQ ID NO 73
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 73

aactattgag cacagggata aagatgactg                                   30

<210> SEQ ID NO 74
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 74

aatatctaatt tcttgttttg aacagtgaac att                             33

<210> SEQ ID NO 75
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 75

tatgccggct actttggcaa gcttgaacat aaactc                           36

<210> SEQ ID NO 76
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 76

ggcctcgagc taattcttgt tttgaacagt gaacatt                         37

<210> SEQ ID NO 77
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 77

atggccgaca aggtcctgaa ggagaaga                                    28

<210> SEQ ID NO 78
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 78

ttaatgtcct gggagaggt agaaacatct tgt                               33

```

-continued

---

<210> SEQ ID NO 79  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer  
  
<400> SEQUENCE: 79  
  
tcaagccccc aatctggaaa ttctca 26

<210> SEQ ID NO 80  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer  
  
<400> SEQUENCE: 80  
  
ctggagagtc actgatcaac agttcc 26

<210> SEQ ID NO 81  
<211> LENGTH: 36  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer  
  
<400> SEQUENCE: 81  
  
acaaggatcc accatggccg acaaggatct gaaggg 36

<210> SEQ ID NO 82  
<211> LENGTH: 39  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer  
  
<400> SEQUENCE: 82  
  
cgcctctaga cctcaattgc caggaagag atagaagta 39

<210> SEQ ID NO 83  
<211> LENGTH: 60  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer  
  
<400> SEQUENCE: 83  
  
ctgcagtgtt ggcggtggcg gcgatctag aaacttgcca acccctactc catccccggg 60

<210> SEQ ID NO 84  
<211> LENGTH: 60  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer  
  
<400> SEQUENCE: 84  
  
cccgggatg gagtaggggt tggcaagttt ctagatccgc cgccaccgcc accactgcag 60

<210> SEQ ID NO 85  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

-continued

---

<400> SEQUENCE: 85  
atgcacccctc agcagttggt catcgctt 28

<210> SEQ ID NO 86  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 86  
tgcaggacac ggatgcccag ttgct 25

<210> SEQ ID NO 87  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 87  
acaggtacca tgcacccctca gcagttggtc atcgctt 37

<210> SEQ ID NO 88  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 88  
ctaactgcag gacacggatg cccag 25

<210> SEQ ID NO 89  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 89  
atgtgcccg cgcgtggcc 19

<210> SEQ ID NO 90  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 90  
ctaggaagca ttcagatagc tcatcat 27

<210> SEQ ID NO 91  
<211> LENGTH: 39  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 91  
tatgaccgg ggatgttcca gtgcctcaac cactcccaa 39

-continued

---

```

<210> SEQ ID NO 92
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 92

atgactgcg g cgcctagga agcattcaga tagctcatca t          41

<210> SEQ ID NO 93
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 93

ccatcctggt cctgctaagc          20

<210> SEQ ID NO 94
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 94

ccatctggta catcttcaag tc          22

<210> SEQ ID NO 95
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 95

aaaaaacccg ggtatgttcc aatgtttcaa ccaactccc          38

<210> SEQ ID NO 96
<211> LENGTH: 51
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 96

gcggccgctc gagttaggaa gagttcaagt aggacatcat tctattgatg g          51

<210> SEQ ID NO 97
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 97

cttaaaggaa cagaaagaat cc          22

<210> SEQ ID NO 98
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Primer

```

-continued

---

<400> SEQUENCE: 98  
 ggtattccca gctgacctc 19

<210> SEQ ID NO 99  
 <211> LENGTH: 37  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 99  
 cataggtacc atgcaccctc agcagttggt catctcc 37

<210> SEQ ID NO 100  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Primer

<400> SEQUENCE: 100  
 atctaaatgc atgacacaga tgcccagtc 29

<210> SEQ ID NO 101  
 <211> LENGTH: 561  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(561)

<400> SEQUENCE: 101

ggg atg ttc cag tgc ctc aac cac tcc caa acc ctg ctg cga gcc atc 48  
 Gly Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu Leu Arg Ala Ile  
 1 5 10 15

agc aac acg ctt cag aag gcc aga caa act cta gaa ttt tac tcc tgc 96  
 Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu Phe Tyr Ser Cys  
 20 25 30

act tcc gaa gag att gat cat gaa gat atc aca aaa gat aaa acc agc 144  
 Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp Lys Thr Ser  
 35 40 45

aca gtg gag gcc tgc tta cca ctg gaa tta acc atg aat gag agt tgc 192  
 Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met Asn Glu Ser Cys  
 50 55 60

ctg gct tcc aga gag atc tct ctg ata act aat ggg agt tgc ctg gcc 240  
 Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala  
 65 70 75 80

tcc aga aag acc tct ttt atg acg acc ctg tgc ctt agc agt atc tat 288  
 Ser Arg Lys Thr Ser Phe Met Thr Thr Leu Cys Leu Ser Ser Ile Tyr  
 85 90 95

gag gac ttg aag atg tac cag gtg gag ttc aag gcc atg aat gca aag 336  
 Glu Asp Leu Lys Met Tyr Gln Val Glu Phe Lys Ala Met Asn Ala Lys  
 100 105 110

ctg tta atg gat cct aaa agg cag atc ttt ctg gat caa aac atg ctg 384  
 Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln Asn Met Leu  
 115 120 125

aca gct att gat gag ctg tta cag gcc ctg aat gtc aac agt gtg act 432  
 Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Val Asn Ser Val Thr  
 130 135 140

gtg cca cag aac tcc tcc ctg gaa gaa ccg gat ttt tat aaa act aaa 480  
 Val Pro Gln Asn Ser Ser Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys  
 145 150 155 160

-continued

atc aag ctg tgc ata ctt ctt cat gct ttc aga att cgt gca gtg acc 528  
 Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg Ala Val Thr  
 165 170 175

atc aat aga atg atg agc tat ctg aat gct tcc 561  
 Ile Asn Arg Met Met Ser Tyr Leu Asn Ala Ser  
 180 185

<210> SEQ ID NO 102  
 <211> LENGTH: 187  
 <212> TYPE: PRT  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 102

Gly Met Phe Gln Cys Leu Asn His Ser Gln Thr Leu Leu Arg Ala Ile  
 1 5 10 15  
 Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu Phe Tyr Ser Cys  
 20 25 30  
 Thr Ser Glu Ile Asp His Glu Asp Ile Thr Lys Asp Lys Thr Ser  
 35 40 45  
 Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met Asn Glu Ser Cys  
 50 55 60  
 Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly Ser Cys Leu Ala  
 65 70 75 80  
 Ser Arg Lys Thr Ser Phe Met Thr Thr Leu Cys Leu Ser Ser Ile Tyr  
 85 90 95  
 Glu Asp Leu Lys Met Tyr Gln Val Glu Phe Lys Ala Met Asn Ala Lys  
 100 105 110  
 Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln Asn Met Leu  
 115 120 125  
 Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Val Asn Ser Val Thr  
 130 135 140  
 Val Pro Gln Asn Ser Ser Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys  
 145 150 155 160  
 Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg Ala Val Thr  
 165 170 175  
 Ile Asn Arg Met Met Ser Tyr Leu Asn Ala Ser  
 180 185

<210> SEQ ID NO 103  
 <211> LENGTH: 561  
 <212> TYPE: DNA  
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 103

ggaagcattc agatagctca tcattctatt gatggtcact gcacgaattc tgaagcatg 60  
 aagaagtatg cagagcttga ttttagtttt ataaaaatcc ggttcttcca gggaggagtt 120  
 ctgtggcaca gtcacactgt tgacattcag ggcctgtaac agctcatcaa tagctgtcag 180  
 catgttttga tccagaaga tctgcctttt aggatccatt aacagctttg cattcatggc 240  
 cttgaactcc acctggtaca tcttcaagtc ctcatagata ctgctaaggc acagggtcgt 300  
 cataaaagag gtctttcttg aggccaggca actcccatta gttatcagag agatctctct 360  
 ggaagccagg caactctcat tcatggttaa ttccagtggg aagcaggcct ccaactgtgct 420  
 ggttttatct tttgtgatat cttcatgata aatctcttcg gaagtgcagg agtaaaattc 480  
 tagagtttgt ctggccttct gaagcgtggt gctgatggct cgcagcaggg tttgggagtg 540  
 gttgaggcac tggaacatcc c 561

-continued

```

<210> SEQ ID NO 104
<211> LENGTH: 1455
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (232)..(897)

<400> SEQUENCE: 104

ggcaccgagc aaaccccgcg gcccagctc cacgtgtcac cgagaagctg atgtagagag      60
agacagagag aaaaagcaag ccggacaccg ggtccccggg aaagtcctgg cgcgcctcgg      120
gcccaattata aaaatgtgac cccccgggtc ggcctcccac cgcgcgccctc ccctgcccgcg      180
tccgcagctc ggtgccagcg cccgcggggg tccacgcagc gcccgcccag c atg tgc      237
                                     Met Cys
                                     1

cgc cgc cgc ggc ctc ctc ctt gtg acc atc ctg gtc ctg cta agc cac      285
Pro Pro Arg Gly Leu Leu Leu Val Thr Ile Leu Val Leu Leu Ser His
      5                               10                               15

ctg gac cac ctt act tgg gcc agg agc ctc ccc aca gcc tca ccg agc      333
Leu Asp His Leu Thr Trp Ala Arg Ser Leu Pro Thr Ala Ser Pro Ser
      20                               25                               30

cca gga ata ttc cag tgc ctc aac cac tcc caa aac ctg ctg aga gcc      381
Pro Gly Ile Phe Gln Cys Leu Asn His Ser Gln Asn Leu Leu Arg Ala
      35                               40                               45                               50

gtc agc aac acg ctt cag aag gcc aga caa act cta gaa tta tat tcc      429
Val Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu Leu Tyr Ser
                               55                               60                               65

tgc act tcc gaa gag att gat cat gaa gat atc aca aag gat aaa acc      477
Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp Lys Thr
      70                               75                               80

agc aca gtg gag gcc tgc tta cca ctg gaa tta acc atg aat gag agt      525
Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met Asn Glu Ser
      85                               90                               95

tgc ctg gct tcc aga gag atc tct ttg ata act aac ggg agt tgc ctg      573
Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly Ser Cys Leu
      100                               105                               110

gcc tct gga aag gcc tct ttt atg acg gtc ctg tgc ctt agc agc atc      621
Ala Ser Gly Lys Ala Ser Phe Met Thr Val Leu Cys Leu Ser Ser Ile
      115                               120                               125                               130

tat gag gac ttg aag atg tac cag atg gaa ttc aag gcc atg aac gca      669
Tyr Glu Asp Leu Lys Met Tyr Gln Met Glu Phe Lys Ala Met Asn Ala
      135                               140                               145

aag ctt tta atg gat ccc aag agg cag atc ttt ctg gat caa aac atg      717
Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln Asn Met
      150                               155                               160

ctg aca gct atc gat gag ctg tta cag gcc ctg aat ttc aac agt gtg      765
Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Phe Asn Ser Val
      165                               170                               175

act gtg cca cag aaa tcc tcc ctt gaa gag ccg gat ttt tat aaa act      813
Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe Tyr Lys Thr
      180                               185                               190

aaa atc aag ctc tgc ata ctt ctt cat gct ttc aga att cgt gcg gtg      861
Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg Ala Val
      195                               200                               205                               210

acc atc gat aga atg atg agt tat ctg aat tct tcc taaaaagctg      907
Thr Ile Asp Arg Met Met Ser Tyr Leu Asn Ser Ser
      215                               220

aggctctctc cgactttaa gtcattccta taaaaatgtg aacccaaaag aatttttcat      967
aagatagggg ttaagaacca gggagggggg ggcttgacct ggtcctactt aagctagtac      1027

```

-continued

---

```

gataattctc atgcttgttt acattagttg ccactcaaat ttgaaaagat gtgactgtta 1087
tatcccacac gatgcctttg accaagtata tttoacattt actatggata agttaagtgt 1147
tcgtgagcaa attgctaaag aggaaaaatg tcctcaccga acatgttttt attttccctt 1207
taatagaaga gcaagacttt ataagctatt tctgtaccaa actgtttgta gaaacaaaca 1267
ctcaagcata atttatttaa aaatacttat ttatataatt ttgtgttcat gaaagcatgt 1327
gaattaatth atattttatt atgttatatt tattaagta tttattatca agtggatttg 1387
ggatatctta tgttctaaaa ataaaatgat tgagtagaaa aaaaaaaaaa aaaaaaaaaa 1447
aaaaaaaaa 1455
    
```

```

<210> SEQ ID NO 105
<211> LENGTH: 222
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris
    
```

<400> SEQUENCE: 105

```

Met Cys Pro Pro Arg Gly Leu Leu Leu Val Thr Ile Leu Val Leu Leu
 1             5             10             15
Ser His Leu Asp His Leu Thr Trp Ala Arg Ser Leu Pro Thr Ala Ser
          20             25             30
Pro Ser Pro Gly Ile Phe Gln Cys Leu Asn His Ser Gln Asn Leu Leu
          35             40             45
Arg Ala Val Ser Asn Thr Leu Gln Lys Ala Arg Gln Thr Leu Glu Leu
          50             55             60
Tyr Ser Cys Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp
65             70             75             80
Lys Thr Ser Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Met Asn
          85             90             95
Glu Ser Cys Leu Ala Ser Arg Glu Ile Ser Leu Ile Thr Asn Gly Ser
          100            105            110
Cys Leu Ala Ser Gly Lys Ala Ser Phe Met Thr Val Leu Cys Leu Ser
          115            120            125
Ser Ile Tyr Glu Asp Leu Lys Met Tyr Gln Met Glu Phe Lys Ala Met
          130            135            140
Asn Ala Lys Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln
145            150            155            160
Asn Met Leu Thr Ala Ile Asp Glu Leu Leu Gln Ala Leu Asn Phe Asn
          165            170            175
Ser Val Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe Tyr
          180            185            190
Lys Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg
          195            200            205
Ala Val Thr Ile Asp Arg Met Met Ser Tyr Leu Asn Ser Ser
          210            215            220
    
```

```

<210> SEQ ID NO 106
<211> LENGTH: 1455
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
    
```

<400> SEQUENCE: 106

```

tttttttttt tttttttttt tttttttttt tctactcaat cattttatth ttagaacata 60
agatatccca aatccacttg ataataaata ctttaataaa tataacataa ataaatataa 120
attaattcac atgcttttcat gaacacaaaa ttatataaat aagtattttt aaataaatta 180
    
```

-continued

```

tgcttgagtg tttgtttcta caaacagttt ggtacagaaa tagcttataa agtcttgctc 240
ttctattaaa gggaaaataa aaacatgttc ggtgaggaca tttttcctct ttagcaattt 300
gctcacgaac acttaactta tccatagtaa atgtgaaata tacttggtca aaggcatcgt 360
gtgggatata acagtcacat ctttcaaaa ttgagtggca actaatgtaa acaagcatga 420
gaattatcgt actagcttaa gtaggaccag gtcaagccac cccctccctg gttcttaacc 480
cctatcttat gaaaaattct tttgggttca catttttata ggaatgactt taaagtcgag 540
agagacctca gcttttttag aagaattcag ataactcadc attctatcga tggtcaccgc 600
acgaattctg aaagcatgaa gaagtatgca gagcttgatt ttagttttat aaaaatccgg 660
ctcttcaagg gaggatttct gtggcacagt cacactgttg aaattcaggg cctgtaacag 720
ctcatcgata gctgtcagca tgttttgatc cagaaagatc tgcctcttgg gatccattaa 780
aagctttgcg ttcattggcct tgaattccat ctggtacatc ttcaagtccat catagatgct 840
gctaaggcac aggaccgtca taaaagaggc ctttccagag gccaggcaac tcccgttagt 900
tatcaaaagag atctctctgg aagccaggca actctcattc atggttaatt ccagtggtaa 960
gcaggcctcc actgtgctgg ttttatcctt tgtgatatct tcatgatcaa tctcttcgga 1020
agtgcaggaa tataattcta gagtttgtct ggccttctga agcgtgttgc tgacggctct 1080
cagcagggtt tgggagtggt tgaggcactg gaatattcct gggctcggtg aggctgtggg 1140
gaggctcctg gcccagaata ggtggcccag gtggcttagc aggaccagga tggtcacaag 1200
gaggaggccg cgcggcgggc acatgctggg cgggcgctgc gtggaccctg gcgggcgctg 1260
gacgcggact gcggacgcgg caggggaggg cggcggtggg aggccgaccg ggggggtcac 1320
atttttataa ttggcccag gcgcgccagg actttcccgg gactccggtg tccggcttgc 1380
ttctctctc tgtctctctc tacatcagct tctcgggtgac acgtggagct gggcccgcgg 1440
ggtttgcctc gtgcc 1455

```

```

<210> SEQ ID NO 107
<211> LENGTH: 2267
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (154)..(1140)

```

<400> SEQUENCE: 107

```

ggcacgaggc aacttgagca agtcagtttc tagtttaagt ttccatcaa aagctcggta 60
ggagtagggt atataagctc cagtagcagt agcagcagca gcaggagact ccttttcaga 120
cccagggaac cttgcagcct ggccagaagc aag atg cat cct cag cag ttg gtc 174
Met His Pro Gln Gln Leu Val
1 5
atc tcc tgg ttt tcc ctc gtt ttg ctg gcg tct ccc ctc atg gcc ata 222
Ile Ser Trp Phe Ser Leu Val Leu Leu Ala Ser Pro Leu Met Ala Ile
10 15 20
tgg gaa ctg gag aaa gat gtt tat gtt gta gag ttg gac tgg cac cct 270
Trp Glu Leu Glu Lys Asp Val Tyr Val Val Glu Leu Asp Trp His Pro
25 30 35
gat gcc ccc gga gaa atg gtg gtc ctc acc tgc cat acc cct gaa gaa 318
Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys His Thr Pro Glu Glu
40 45 50 55
gat gac atc act tgg acc tca gcg cag agc agt gaa gtc cta ggt tct 366
Asp Asp Ile Thr Trp Thr Ser Ala Gln Ser Ser Glu Val Leu Gly Ser
60 65 70

```

-continued

ggt aaa act ctg acc atc caa gtc aaa gaa ttt gga gat gct ggc cag Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Gly Asp Ala Gly Gln 75 80 85	414
tat acc tgc cat aaa gga ggc aag gtt ctg agc cgc tca ctc ctg ttg Tyr Thr Cys His Lys Gly Gly Lys Val Leu Ser Arg Ser Leu Leu Leu 90 95 100	462
att cac aaa aaa gaa gat gga att tgg tcc act gat atc tta aag gaa Ile His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys Glu 105 110 115	510
cag aaa gaa tcc aaa aat aag atc ttt ctg aaa tgt gag gca aag aat Gln Lys Glu Ser Lys Asn Lys Ile Phe Leu Lys Cys Glu Ala Lys Asn 120 125 130 135	558
tat tct gga cgt ttc aca tgc tgg tgg ctg acg gca atc agt act gat Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Ala Ile Ser Thr Asp 140 145 150	606
ttg aaa ttc agt gtc aaa agt agc aga ggc ttc tct gac ccc caa ggg Leu Lys Phe Ser Val Lys Ser Ser Arg Gly Phe Ser Asp Pro Gln Gly 155 160 165	654
gtg aca tgt gga gca gtg aca ctt tca gca gag agg gtc aga gtg gac Val Thr Cys Gly Ala Val Thr Leu Ser Ala Glu Arg Val Arg Val Asp 170 175 180	702
aac agg gat tat aag aag tac aca gtg gag tgt cag gag ggc agt gcc Asn Arg Asp Tyr Lys Lys Tyr Thr Val Glu Cys Gln Glu Gly Ser Ala 185 190 195	750
tgc ccc tct gcc gag gag agc cta ccc atc gag gtc gtg gtg gat gct Cys Pro Ser Ala Glu Glu Ser Leu Pro Ile Glu Val Val Val Asp Ala 200 205 210 215	798
att cac aag ctc aag tat gaa aac tac acc agc agc ttc ttc atc aga Ile His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile Arg 220 225 230	846
gac atc atc aaa cca gac cca ccc aca aac ctg cag ctg aag cca ttg Asp Ile Ile Lys Pro Asp Pro Pro Thr Asn Leu Gln Leu Lys Pro Leu 235 240 245	894
aaa aat tct cgg cac gtg gag gtc agc tgg gaa tac ccc gac acc tgg Lys Asn Ser Arg His Val Glu Val Ser Trp Glu Tyr Pro Asp Thr Trp 250 255 260	942
agc acc cca cat tcc tac ttc tcc ctg aca ttt tgc ata cag gcc cag Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Ile Gln Ala Gln 265 270 275	990
ggc aag aac aat aga gaa aag aaa gat aga ctc tgc gtg gac aag acc Gly Lys Asn Asn Arg Glu Lys Lys Asp Arg Leu Cys Val Asp Lys Thr 280 285 290 295	1038
tca gcc aag gtc gtg tgc cac aag gat gcc aag atc cgc gtg caa gcc Ser Ala Lys Val Val Cys His Lys Asp Ala Lys Ile Arg Val Gln Ala 300 305 310	1086
cga gac cgc tac tat agt tca tcc tgg agc gac tgg gca tct gtg tcc Arg Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Asp Trp Ala Ser Val Ser 315 320 325	1134
tgc agt taggttccac ccccaggatg aatcttgag gaaagtga agatattatg Cys Ser	1190
caaaattttc taaggacaca ttgaagaggc tccaaaagtt attttctgcc taattttctt	1250
tttgtaaagg gtcattattg tgtcttcgca atatttttta catttaaag ccaaatgcc	1310
actgaaacaa tcagctactt tatttataga ttttcagcta gcaggctgcc actgacctta	1370
atgctattta aatatttaag taatttatgt atttattaat ttattgttat tgaacacttg	1430
tgtgccaaga tatattgat gtttcatacc ctcaggacct gatctgtaag gaataggccc	1490
tattatgcaa aatgtgaatt tatgtgttat ttatactgac aacttttcaa gcaagaatgt	1550

-continued

```

atcattttta tgacaaccag tgagcacaca atattatgat gccagcacca taatatattt 1610
gtgatggatg ggaacacaga ggtagttaaa tagagacatg gagacacgaa tccatttgag 1670
aagtttctg agacggagat gttagatcct gtatccataa agacttcctt gcggtggtgt 1730
tgataaagca attcagggcc acttgcatth ttaagcaagt ttagtthttg gatgcctgaa 1790
tttagaaaga cctgagacaa ataactcaaa ttgagattca gcttcagcca ccttgccagt 1850
ccccatcccc atctatctgt aagtcattgg agagtgaccc agggacactg taagtgtctg 1910
gaagtaaaaa ggtcttatga tccaagaggg agaaccaaca tggccaagca caaaaaattg 1970
tcagaatttc cagctgctcc ttaatagcca ggcaaaaaaa gcacatggat gcaaagaaaa 2030
tggccaagaa ttgcttactg gacagcgcaa gtgaacctga ctggtggatg tgaccagaaa 2090
gtgccaatcg ctgaggtgct acttttaagt aatgaatggt ctttctgtaa agtgatttca 2150
tttctttct gtttacttat ttgthtttgc attctgacaa tgcactaata aaaatataac 2210
tcttgthtgc aataataaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 2267

```

```

<210> SEQ ID NO 108
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

<400> SEQUENCE: 108

```

Met His Pro Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Leu Leu
1          5          10          15
Ala Ser Pro Leu Met Ala Ile Trp Glu Leu Glu Lys Asp Val Tyr Val
20          25          30
Val Glu Leu Asp Trp His Pro Asp Ala Pro Gly Glu Met Val Val Leu
35          40          45
Thr Cys His Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Ala Gln
50          55          60
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
65          70          75          80
Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Lys Val
85          90          95
Leu Ser Arg Ser Leu Leu Leu Ile His Lys Lys Glu Asp Gly Ile Trp
100         105         110
Ser Thr Asp Ile Leu Lys Glu Gln Lys Glu Ser Lys Asn Lys Ile Phe
115         120         125
Leu Lys Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
130         135         140
Leu Thr Ala Ile Ser Thr Asp Leu Lys Phe Ser Val Lys Ser Ser Arg
145         150         155         160
Gly Phe Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Val Thr Leu Ser
165         170         175
Ala Glu Arg Val Arg Val Asp Asn Arg Asp Tyr Lys Lys Tyr Thr Val
180         185         190
Glu Cys Gln Glu Gly Ser Ala Cys Pro Ser Ala Glu Glu Ser Leu Pro
195         200         205
Ile Glu Val Val Val Asp Ala Ile His Lys Leu Lys Tyr Glu Asn Tyr
210         215         220
Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Thr
225         230         235         240
Asn Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg His Val Glu Val Ser
245         250         255

```



-continued

```

agccaccagc atgtgaaacg tccagaataa ttctttgcct cacatttcag aaagatctta 1740
tttttgatt ctttctgttc ctttaagata tcagtggacc aaattccatc ttcttttttg 1800
tgaatcaaca ggagtgagcg gctcagaacc ttgcctcctt tatggcaggt atactggcca 1860
gcatctccaa attctttgac ttggatggtc agagttttac cagaacctag gacttcactg 1920
ctctgcgctg aggtccaagt gatgtcatct tcttcagggg tatggcaggt gaggaccacc 1980
atcttcccg gggcatcagg gtgccagtcc aactctacaa cataaacatc tttctccagt 2040
tcccataatg ccatgagggg agacgccagc aaaacgaggg aaaaccagga gatgaccaac 2100
tgctgaggat gcatcttgct tctggccagg ctgcaagggt ccctgggtct gaaacggagt 2160
ctctgctgc tgctgctact gctactggag cttatatacc ctactcctac cgagcttttg 2220
gatggaaact taaactagaa actgacttgt ccaagttgcc tcgtgcc 2267
    
```

```

<210> SEQ ID NO 110
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Linker
    
```

<400> SEQUENCE: 110

Gly Gly Gly Gly Gly Ser  
 1 5

```

<210> SEQ ID NO 111
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Felis catus
    
```

<400> SEQUENCE: 111

Arg Asn Leu Pro Thr Pro Thr Pro Ser Pro  
 1 5 10

What is claimed is: 40

1. An isolated nucleic acid molecule selected from the group consisting of:
  - (a) an isolated nucleic acid molecule comprising:
    - (i) a nucleic acid sequence encoding a feline p35 subunit protein; 45
    - (ii) a nucleic acid linker of (XXX)<sub>n</sub>, wherein n=0 to 60; and
    - (iii) a nucleic acid sequence encoding a feline p40 subunit protein; and 50
  - (b) an isolated nucleic acid molecule comprising a nucleic acid sequence fully complementary to the nucleic acid molecule set forth in (a).
2. The isolated nucleic acid molecule of claim 1, wherein said isolated nucleic acid molecule is selected from the group consisting of: 55
  - (a) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p35 subunit encoding nucleic acid sequence comprises at least 44 contiguous nucleotides identical in sequence to at least 44 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:32, SEQ ID NO:35 and SEQ ID NO:101; and 60
  - (b) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p40 subunit encoding 65

- nucleic acid sequence comprises at least 44 contiguous nucleotides identical in sequence to at least 44 contiguous nucleotides of a nucleic acid sequence selected from the group consisting of SEQ ID NO:26, SEQ ID NO:29 and SEQ ID NO:55.
3. The isolated nucleic acid molecule of claim 1, wherein said isolated nucleic acid molecule is selected from the group consisting of:
  - (a) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p35 subunit encoding nucleic acid sequence comprises a nucleotide sequence at least 90% identical to SEQ ID NO:32, SEQ ID NO:35 and SEQ ID NO:101;
  - (b) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p40 subunit encoding nucleic acid sequence comprises a nucleotide sequence at least 90% identical to SEQ ID NO:26, SEQ ID NO:29 and SEQ ID NO:55; and
  - (c) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said nucleic acid molecule comprises a nucleic acid sequence at least 90% identical to SEQ ID NO:38 or SEQ ID NO:43.

193

4. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule encodes a protein having a function selected from the group consisting of:

- (a) eliciting an immune response against an IL-12 protein having the amino acid sequence of SEQ ID NO: 39 or SEQ ID NO: 44;
- (b) selectively binding to an antibody raised against an IL-12 protein having the amino acid sequence of SEQ ID NO: 39 or SEQ ID NO: 44; and
- (c) exhibiting IL-12 activity.

5. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid linker comprises SEQ ID NO:83.

6. The isolated nucleic acid molecule of claim 1, wherein said isolated nucleic acid molecule is selected from the group consisting of:

- (a) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p35 subunit encoding nucleic acid sequence comprises SEQ ID NO:32, SEQ ID NO:35 or SEQ ID NO:101;
- (b) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p40 subunit encoding nucleic acid sequence comprises SEQ ID NO:26, SEQ ID NO:29 or SEQ ID NO:55; and
- (c) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said isolated nucleic acid molecule comprises SEQ ID NO:38 or SEQ ID NO:43.

7. The isolated nucleic acid molecule of claim 1, wherein said isolated nucleic acid molecule is selected from the group consisting of:

- (a) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p35 subunit encoding nucleic acid sequence consists of SEQ ID NO:32, SEQ ID NO:35 or SEQ ID NO:101;
- (b) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said p40 subunit encoding nucleic acid sequence consists of SEQ ID NO:26, SEQ ID NO:29 or SEQ ID NO:55; and
- (c) an isolated nucleic acid molecule comprising a feline p35 subunit encoding sequence and a feline p40 subunit encoding sequence, wherein said isolated nucleic acid molecule has a nucleic acid sequence consisting of SEQ ID NO:38 or SEQ ID NO:43.

8. An isolated nucleic acid molecule selected from the group consisting of:

- (a) an isolated nucleic acid molecule comprising:
  - (i) a first nucleic acid sequence encoding a protein comprising an at least 23 contiguous amino acid region identical in sequence to an at least 23 contiguous amino acid region from SEQ ID NO:33, SEQ ID NO:36 or SEQ ID NO:102;
  - (ii) a nucleic acid linker of (XXX)<sub>n</sub> wherein n=0 to 60; and
  - (iii) a second nucleic acid sequence encoding a protein comprising an at least 23 contiguous amino acid region identical in sequence to an at least 23 contiguous amino acid region from SEQ ID NO:27, SEQ ID NO:30 or SEQ ID NO:56; and

- (b) an isolated nucleic acid molecule comprising a nucleic acid sequence fully complementary to the nucleic acid molecule set forth in (a).

194

9. The isolated nucleic acid molecule of claim 8, wherein said isolated nucleic acid molecule is selected from the group consisting of:

- (a) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said first nucleic acid sequence encodes a protein comprising an amino acid sequence at least 90% identical to SEQ ID NO:33, SEQ ID NO:36 or SEQ ID NO:102;
- (b) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said second nucleic acid sequence encodes a protein comprising an amino acid sequence at least 90% identical to SEQ ID NO:27, SEQ ID NO:30 or SEQ ID NO:56; and
- (c) an isolated nucleic acid molecule encoding a protein comprising an amino acid sequence at least 90% identical to SEQ ID NO:39 or SEQ ID NO:44.

10. The isolated nucleic acid molecule of claim 8, wherein said nucleic acid molecule encodes a protein having a function selected from the group consisting of:

- (a) eliciting an immune response against an IL-12 protein having the amino acid sequence of SEQ ID NO:27, SEQ ID NO:30 or SEQ ID NO:56;
- (b) selectively binding to an antibody raised against an IL-12 protein having the amino acid sequence of SEQ ID NO:27, SEQ ID NO:30 or SEQ ID NO:56; and
- (c) exhibiting IL-12 activity.

11. The isolated nucleic acid molecule of claim 8, wherein said isolated nucleic acid molecule is selected from the group consisting of:

- (a) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said first nucleic acid sequence encodes an amino acid sequence comprising SEQ ID NO:33, SEQ ID NO:36 or SEQ ID NO: 102;
- (b) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said second nucleic acid sequence encodes an amino acid sequence comprising SEQ ID NO:27, SEQ ID NO:30 or SEQ ID NO: 56; and
- (c) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said isolated nucleic acid sequence encodes a protein comprising SEQ ID NO:39 or SEQ ID NO:44.

12. The isolated nucleic acid molecule of claim 8, wherein said isolated nucleic acid molecule is selected from the group consisting of:

- (a) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said first nucleic acid sequence encodes an amino acid sequence consisting of SEQ ID NO:33, SEQ ID NO:36 or SEQ ID NO: 102;
- (b) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said second nucleic acid sequence encodes an amino acid sequence consisting of SEQ ID NO:27, SEQ ID NO:30 or SEQ ID NO: 56; and
- (c) an isolated nucleic acid molecule comprising a first and second nucleic acid sequence, wherein said isolated nucleic acid sequence encodes a protein having the sequence of SEQ ID NO:39 or SEQ ID NO:44.

13. The isolated nucleic acid molecule of claim 8, wherein said nucleic acid linker comprises SEQ ID NO:83.

14. A recombinant virus comprising a nucleic acid molecule as set forth in claim 1.

**195**

15. A recombinant cell comprising a nucleic acid molecule as set forth in claim 1.

16. A kit comprising a nucleic acid molecule as set forth in claim 1.

17. A recombinant virus comprising a nucleic acid molecule as set forth in claim 8. 5

**196**

18. A recombinant cell comprising a nucleic acid molecule as set forth in claim 8.

19. A kit comprising a nucleic acid molecule as set forth in claim 8.

\* \* \* \* \*