

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/665, A61K 38/17, G01N 33/50		A1	(11) International Publication Number: WO 97/07212 (43) International Publication Date: 27 February 1997 (27.02.97)
(21) International Application Number: PCT/US96/13305		(74) Agent: POLLEY, Richard, J.; Klarquist, Sparkman, Campbell, Leigh & Winston, Suite 1600, One World Trade Center, 121 S.W. Salmon Street, Portland, OR 97204 (US).	
(22) International Filing Date: 12 August 1996 (12.08.96)			
(30) Priority Data: 08/514,541 11 August 1995 (11.08.95) US 08/553,058 3 November 1995 (03.11.95) US		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant: OREGON HEALTH SCIENCES UNIVERSITY [US/US]; 3181 S.W. Sam Jackson Park Road, Portland, OR 97201 (US).			
(72) Inventors: GRISEL, Judith, E.; 7419 S.W. 36th Street, Portland, OR 97219 (US). MOGL, Jeffrey, S.; 405 H N.E. 8th Street, Vancouver, WA 98665 (US). GRANDY, David, K.; 3738 S.W. Patton Road, Portland, OR 97221 (US). BUNZOW, James, R.; 9904 S.W. 56th Street, Portland, OR 97219 (US). CIVELLI, Olivier, Kirschgartenstrasse #36, CH-4147 Aesch (CH). REINSCHEID, Rainer, Klaus; Im Entenschwumm 1, D-79591 Eimeldingen (DE). NOTHACKER, Hans-Peter; In den Hofmatten 5b, D-79418 Schliengen (DE). MONSMA, Frederick, James; Hoehnstrasse 45, CH-4125 Riehen (CH).		<p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> </p>	

(54) Title: OPIOID ANTAGONISTS AND METHODS OF THEIR USE

(57) Abstract

The present invention relates to a novel mammalian anti-opioid receptor protein (OFQR), peptide ligands (such as OFQ) that bind to OFQR, and methods of using the OFQ peptide and analogues to reverse the physiologic effects of opiates such as morphine. The isolation, characterization and pharmacological use of the endogenous peptide ligand is described. A particular embodiment of the OFQ peptide is a heptadecapeptide having an FGGF aminoterminal motif. The peptide specifically binds to an OFQ receptor protein heterologously expressed in mammalian cells. The peptide does not bind with high affinity to μ , δ or κ receptors, but it antagonizes opioid mediated effects (such as analgesia and hypothermia) without increasing nociceptive sensitivity. Tyrosine substitution variants of the peptide ligand specifically bind to the opioid receptor and can be radioiodinated. Also provided are methods of making such peptide ligands and OFQR antagonists, and methods of using the ligands for diagnostic and therapeutic uses and for the identification of other naturally-occurring or synthetic opioid receptor ligands.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

OPIOID ANTAGONISTS AND METHODS OF THEIR USE

This invention was made at least in part with government support under National Institutes of Health grant RO1 MH48991. The government may have certain rights to this invention.

5

BACKGROUND OF THE INVENTION

Field of the Invention

This invention relates to receptors from mammalian species and ligands specific for such receptors, which are active in the antagonism of opioid action. Specifically, the invention relates to the isolation of an endogenous peptide ligand specific for a novel mammalian receptor, the 10 recognition of its anti-opioid properties, and use of the ligand for reversing physiologic effects of opiates such as morphine. The invention also relates to the construction of analogues, derivatives and peptide mimetics of this endogenous mammalian receptor ligand, and their use as opiate antagonists. Specifically provided is a mammalian hypothalamus-derived endogenous opioid receptor ligand, synthetic embodiments and analogues thereof, and methods of making and using 15 such ligands.

Background of the Invention

The use (and abuse) of opiates, such as opium and morphine, have been known since antiquity (*reviewed in* Brownstein, 1993, Proc. Natl. Acad. Sci. USA 90: 5391-5393). Since the nineteenth century, chemical characterization and synthesis of many morphine analogues have 20 been achieved in an effort to discover a compound having the analgesic effects of morphine but lacking its addictive potential. These efforts have heretofore been unsuccessful.

The biology behind the reasons for the analgesic and addictive properties of morphine and morphine-like compounds was first elucidated by the discovery of endogenous morphine-like compounds termed enkephalins (*see* DiChara & North, 1992, Trends in Pharmacol. Sci. 13: 185-25 193 *for review*). Accompanying this finding of an endogenous opioid was the biochemical evidence for a family of related but distinct opiate receptors, each of which displays a unique pharmacological profile of response to opiate agonists and antagonists (*see* McKnight & Rees, 1991, Neurotransmissions 7: 1-6 *for review*). To date, four distinct opiate receptors have been described by their pharmacological profiles and anatomical distribution: these comprise the μ , δ , 30 κ , and σ receptors (the σ receptor has been determined to be a non-opioid receptor displaying cross-reactivity with some opioid agonists).

Thus, mammalian opioid receptors are known in the art, and some of these proteins have been isolated biochemically and their corresponding genes have been recently cloned using genetic engineering means.

35 Kieffer *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89: 12048-12052 disclosed the isolation of a cDNA copy of the mouse δ -opioid receptor by expression cloning.

Evans *et al.*, 1992, Science 258: 1952-1955 disclosed the isolation of a cDNA copy of the mouse δ -opioid receptor by expression cloning.

Chen *et al.*, 1993, Molec. Pharmacol. 44: 8-12 disclosed the isolation of a cDNA copy of the rat μ -opioid receptor.

Yasuda *et al.*, 1993, Proc. Natl. Acad. Sci. USA 90: 6736-6740 disclosed the isolation of a cDNA copy of each of the mouse κ - and δ -opioid receptor.

5 Bzdega *et al.*, 1993, Proc. Natl. Acad. Sci. USA 90: 9305-9309 disclose the isolation and chromosomal location of the δ -opioid receptor in the mouse.

The present inventors have cloned, expressed and functionally characterized a novel mammalian receptor gene, disclosed in co-owned and co-pending U.S. Patent Application No. 08/149,093, filed November 8, 1993, which is hereby incorporated by reference in its entirety.

10 Specifically disclosed therein are nucleic acids encoding the novel mammalian receptor gene, recombinant expression constructs comprising this receptor gene, cells containing such constructs and expressing the novel receptor gene, and methods for making and using such nucleic acids, as well as constructs and cells for opioid detection and novel drug screening. The nucleic acid sequence of the gene and the deduced amino acid sequence of the cognate receptor protein were 15 also disclosed in this prior application.

The receptor gene was previously referred to in the co-pending case as an MSOR (methadone specific opioid receptor) gene because the receptor bound methadone with a high affinity. Such high affinity constitutes a specificity for methadone, although this term is not intended to imply that only methadone binds to the receptor. Indeed, as shown herein, an 20 endogenous ligand of the present invention binds to the same receptor with startlingly high affinity (producing a physiologic response when administered in a picomole or nanomole dose i.c.v.). Now that the endogenous ligand has been discovered, and disclosed in co-owned and co-pending U.S. Patent Application No. 08/514,451 filed August 11, 1995, the MSOR terminology is no longer needed. The ligand has been named orphanin FQ (OFQ) from the initial and terminal 25 amino acids of the disclosed sequence (F for phenylalanine and Q for glutamine). Hence the receptor will now be referred to as the orphanin FQ receptor (OFQR), which is an anti-opioid receptor (AOR).

A great advantage in efforts for developing novel psychotropic drugs which exert 30 their activity (analgesic and otherwise) *via* binding to mammalian opioid or anti-opioid receptors would be to identify the endogenous ligand(s) which bind to such receptors. Certain such opioid ligands have been isolated in the prior art, including the peptides comprising the endorphins and enkephalins (*see* Jaffe and Martin, 1990, "Opioid Analgesics and Antagonists", in Goodman and Gilman, eds., The Pharmacological Basis of Therapeutics, 7th ed. (Pergamon Press, Inc.: New York), Chapter 21, p. 491-531).

35 Although opiates are powerful therapeutic agents, their therapeutic index is sufficiently low to cause many problems in their clinical use. Such problems include stupor, respiratory compromise, or profound analgesia that may be undesired (and mask symptoms of an acute surgical emergency, such as appendicitis). Physical dependence on morphine or other opiates can

also develop, and may be difficult for a clinician to diagnose. It has been recognized that administration of an antagonist to reverse opiate effects is clinically useful in these circumstances, either for reducing physiologic effects of the opiate (undesired analgesia or apnea), or precipitating a withdrawal syndrome to demonstrate addiction.

5 In spite of the importance of opioid antagonists, progress has only slowly been made in developing additional members of this class of drugs. The best known such antagonist is naloxone, an organically synthesized molecule that resembles oxymorphone and is a competitive antagonist at μ , δ , κ and σ opioid receptors. Naltrexone appears to be another relatively pure antagonist, but with higher oral efficacy and a longer duration of action. These drugs are
10 believed to interact with and competitively bind to the opioid receptors to exert their antagonistic action. They reverse signs of opiate intoxication such as apnea and stupor, and they can also stimulate a withdrawal syndrome in drug dependent subjects characterized by anxiety, chills, abdominal cramping, tachycardia lacrimation and sweating. The search for additional opioid antagonists has continued, because of a clinical need for such drugs. Endogenous antagonists
15 have been particularly pursued because of their predicted potency, and the invaluable information they would provide about the body's physiologic pathways for developing tolerance and addiction. In spite of these advantages, such endogenous anti-opioids have eluded investigators for years.

SUMMARY OF THE INVENTION

20 The present invention takes advantage of the discovery that an anti-opioid biological system physiologically counteracts or antagonizes the actions mediated by the binding of opioids to opioid receptors. The OFQ anti-opioid receptor is the same receptor (MSOR) disclosed in the commonly assigned U.S. Patent Application Serial No. 08/149,093. The ligand of the present invention is the ligand disclosed in commonly assigned U.S. Patent Application Serial No. 08/514,451. The method of treatment of the present application was made possible by the present inventors' recognition that the receptor is an anti-opioid receptor that, when bound to a ligand of the present invention, antagonizes the physiologic effects of an opioid.

25 This invention provides small, readily-produced peptides that are ligands for a novel mammalian anti-opioid receptor protein having the amino acid sequence identified as SEQ ID Nos.: 5 and 6. Peptides of the invention are characterized as having the amino acid sequence identified as SEQ ID Nos.: 5 and 6 or a subsequence thereof, amino acid sequence variants of the sequence or subsequence, as well as analogues and derivatives thereof, that are ligands for the mammalian opioid receptor protein having the amino acid sequence identified as SEQ ID Nos.: 5 and 6, as well as analogues and derivatives thereof.

30 The method of the present invention antagonizes a physiological effect (such as analgesia or hypothermia) of an opioid (such as morphine or heroin) in an animal (such as a rodent or a human). The method includes administering to the animal a pharmaceutically effective amount of the peptide, which specifically binds to a receptor encoded by the DNA molecule set forth in

SEQ. ID. No. 3. The peptide is preferably an heptadecapeptide having an aminoterminal FGGF motif, and the peptide does not bind with high specificity to mu, delta, or kappa opioid receptors. It antagonizes physiologic opioid effects (such as analgesia or hypothermia) without increasing nociceptive sensitivity beyond the baseline sensitivity of the animal to painful stimuli in the 5 absence of the exogenous opioid or stress induced analgesia. Stress induced analgesia is a normal physiologic mechanism, believed to be mediated by endogenous opioids, for reducing pain perception in response to stressful environmental stimuli.

A pharmaceutically effective amount of the peptide is an amount sufficient to subjectively or objectively diminish the effects of the opiate. An example would be an amount sufficient to 10 reverse the degree of analgesia to painful stimuli induced by an opioid, or an amount sufficient to raise body temperature of an animal experiencing opiate induced hypothermia. In some of the described embodiments, this amount would be a picomolar or nanomolar icv dose. For example, a pharmaceutically effective dose in a mouse would be a 50 picomole to 10 nanomole intracranial ventricular (icv) dose, in 2 μ l vehicle. The present invention may be administered by many 15 routes, including intravenously (iv), subcutaneously (sc), or intrathecally (it). Given the marked anti-opioid potency of the peptide, relatively low doses of the peptide could even be given iv. Human doses given icv and intrathecally would be expected to be 1-10 nanomoles in a 2 microliter volume.

In yet other embodiments, the peptide may be an analogue, derivative or mimetic of the 20 peptides. An analogue is a peptide having the same anti-opioid activity as the disclosed peptides, but differing in primary amino acid sequence. The variations in the primary amino acid sequence could be made using the techniques described in this specification, and known in the art. A mimetic of the peptide is a molecule, either a peptide or other biopharmaceutical (such as an 25 organically synthesized compound) that specifically binds to a receptor encoded by SEQ. ID. No. 3 and antagonizes a physiological effect of an opioid in an animal.

In other embodiments of the invention, the method includes administering to the animal a second anti-opioid antagonist, wherein the second anti-opioid antagonist and anti-opioid peptide are administered in a sufficient amount in combination to antagonize physiological effects of the opioid. The first and second antagonists may be given either in series or administered together in 30 a mixture. For example, the anti-opioid peptide can be administered in a vehicle (such as isotonic saline solution) in admixture with naloxone (at a pharmaceutically effective dose of 1 mg/kg body weight of the animal). Alternatively, the anti-opioid peptide can be administered in combination with other anti-opioid peptides (such as analogues of the peptide of the present invention, or other endogenous or synthetic peptides that have an anti-opioid activity).

35 The peptides of the invention include linear and cyclized peptides, and synthetic analogues and variants thereof. Certain embodiments of such variants include substitution variants, wherein an amino acid residue at one or more positions in the peptide is replaced with a different amino acid residue (including atypical amino acid residues) from that found in the corresponding position

of the amino acid sequence of the parent peptide of the invention. Certain other embodiments of peptide variants of the invention include addition variants, wherein such variant peptides may include up to about a total of 10 additional amino acids, covalently linked to either the amino-terminal or carboxyl-terminal extent, or both, of the parent peptide of the present invention.

5 Such additional amino acids may also include atypical amino acids. Linear and cyclized embodiments of the amino acid substitution and addition variant peptides are also provided as peptides of the invention. In addition, peptides of the invention may be provided as fusion proteins with other functional targeting agents, such as immunoglobulin fragments. Derivatives of the peptides of the invention also include modifications of the amino- and carboxyl-termini and 10 amino acid side chain chemical groups such as amines, carboxylic acids, alkyl and phenyl groups.

In a first aspect, the invention provides peptides of the formula:

$(Xaa)_n\text{-Phe-Gly-Gly-Phe-(A}^1\text{)-(A}^2\text{)-(A}^3\text{)-(A}^4\text{)-(A}^5\text{)-(A}^6\text{)-(A}^7\text{)-(A}^8\text{)-(A}^9\text{)-(A}^{10}\text{)-(A}^{11}\text{)-(A}^{12}\text{)-Gln-(Xaa)}_m$

wherein A^1 is Thr, Leu or Met; A^2 is Gly, Arg or Thr; A^3 is Ala, Arg or Ser; A^4 is Arg, Ile, Glu or Gln; A^5 is Lys, Arg or Phe; A^6 is Ser, Pro or Lys; A^7 is Ala, Lys, Gln or Val; A^8 is Arg, 15 Leu, Thr or Val; A^9 is Lys, Pro or Thr; A^{10} is Tyr, Leu or Trp; A^{11} is Ala, Asp or Val; A^{12} is Asn, or Thr; (Xaa) is any amino acid; n and m are integers wherein $n + m$ is no more than 82, preferably no more than 20, or even no more than 10; and the amino acids are each individually in either the D or L stereochemical configuration and the peptide specifically binds to a mammalian opioid receptor having an amino acid sequence identified by SEQ ID No.: 3. In a 20 preferred embodiment, the peptide has the formula:

Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln (SEQ ID No.: 5).

In an another preferred embodiment, the peptide has the formula:

Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Tyr-Ala-Asn-Gln (SEQ ID No.: 6).

Naturally-occurring embodiments of the peptides, purified using well-established techniques from 25 the cells or tissues producing the peptide, and synthetic embodiments, are within the scope of the invention.

In other embodiments the peptides are of the general formula:

$(Xaa)_n\text{-Phe-Gly-Gly-Phe-(A}^1\text{)-(A}^2\text{)-(A}^3\text{)-(A}^4\text{)-(A}^5\text{)-(A}^6\text{)-(A}^7\text{)-(A}^8\text{)-(A}^9\text{)-(A}^{10}\text{)-(A}^{11}\text{)-(A}^{12}\text{)-Gln-(Xaa)}_m$

and having amino acid substituents as described above wherein such peptides are radiolabeled by 30 conjugation with or binding to a radioactive isotope. In a preferred embodiment, the peptide has the formula:

Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Tyr-Ala-Asn-Gln (SEQ ID No.: 6)

and the radiolabel is a radioisotope of iodine. In other preferred embodiments, the peptide is covalently linked to a radiolabel binding moiety and the radiolabel is a radioisotope of indium, gallium, technetium, rhenium or other useful radioisotope.

35 The invention also provides pharmaceutical compositions of the peptides of the invention.

In a first aspect, the peptides of the invention are provided in a pharmaceutical composition comprising an acceptable carrier or diluent, or a pharmaceutical dosage form such as a vial of

predetermined dose of the peptide, suspended in an injectable or inhalable vehicle such as sterile water or isotonic saline. Other dosage forms include tablets or capsules, particularly for synthetic mimetics more suitable for oral administration.

5 In a second aspect, such pharmaceutical compositions are provided in detectably-labeled embodiments, useful for diagnostic identification of sites of both normal and pathological peptide ligand receptor binding *in vivo* and *in vitro*. In such embodiments, the peptides of the invention are detectably labeled, for example, with a radioisotope such as I-123, I-125 or I-131, conjugated to the peptide *via, inter alia*, a tyrosine residue that is non-essential for receptor binding. Additional radioisotopes, such as In-111, Ga-67, Re-186, Re-188 and Tc-99m, can be conjugated 10 to such peptides using methods well-understood in the art. In such embodiments, the pharmaceutical composition is radiolabeled to an appropriate specific activity, and administered to an animal, preferably a human, at a diagnostically-effective and non-toxic dose. Methods and routes of administration may be selected by those with skill in the art based on well-established techniques and in a clinically-appropriate fashion. Alternatively, the radiolabeled ligands can be 15 used to identify specific brain regions where the ligand receptor is present.

DESCRIPTION OF THE DRAWINGS

Fig. 1 illustrates the nucleotide (SEQ ID No.: 3) and amino acid (SEQ ID No.: 4) sequences of the mammalian anti-opioid receptor of the invention.

20 Fig. 2 presents an amino acid sequence comparison between the novel mammalian OFQR protein of the invention (designated LC132) and the rat μ -opioid receptor, and the mouse δ - and κ -opioid receptor proteins.

Fig. 3 illustrates *in situ* hybridization of rat brain sections with a nucleic acid hybridization probe specific for the mammalian OFQR of the invention.

25 Fig. 4 presents affinity binding experiment results of 3 H-methadone binding to COS-7 cells (Panel A) and to COS-7 cells expressing the novel mammalian OFQR of the invention.

Fig. 5 shows the extent of inhibition of forskolin-stimulated cAMP accumulation in a CHO cell heterologously expressing OFQR of the invention by fractions of acetic acid-extracted peptides from porcine hypothalamus, wherein A_{280} represents absorbance at 280nm, shaded bars 30 represent results obtained with untransfected CHO cells and open bars represent results obtained with OFQR transfected CHO cells.

Fig. 6 shows the extent of inhibition of forskolin-stimulated cAMP accumulation in CHO cell heterologously expressing the OFQR of the invention by fractions of reverse-phase HPLC fractionated acetic acid-extracted peptides from porcine hypothalamus, wherein A_{214} represents 35 absorbance at 214nm, and shaded bars represents results obtained with OFQR transfected CHO cells.

Fig. 7 is a comparison of the amino acid sequence of orphanin FQ (OFQ) and opioid receptor binding peptides; identical amino acid residues shared between the peptides are shown in boldface.

Fig. 8 is a graph illustrating the extent of inhibition of forskolin-stimulated cAMP accumulation in OFQR transfected CHO cells by the peptides orphanin FQ (filled circles); Y14-orphanin FQ (filled squares); monoiodinated Y14-orphanin FQ (open squares) and Leu-Enkephalin (open circles). Orphanin FQ-treated untransfected CHO cells are shown as filled triangles.

Fig. 9 is a graph of (¹²⁵I)-labeled Y14-orphanin FQ binding to isolated membranes from OFQR transfected CHO cells. The inset is a Scatchard analysis of these results.

Fig. 10 shows the effects on horizontal activity (Panel A) and vertical activity (Panel B) in mice treated with varying concentrations of orphanin FQ peptide.

Fig. 11 illustrates the effects of orphanin FQ peptide on nociception in mice as determined using a hotplate assay.

Fig. 12 shows the effects of varying icv doses of orphanin FQ peptide on nociception in mice as determined using a tail-flick assay.

Fig. 13A is a bar graph, illustrating the anti-opioid effect of OFQ on stress induced analgesia, showing average tail withdrawal latency change in animals that have not received morphine analgesia, after administration of saline, naloxone, and OFQ.

Fig. 13B is a bar graph showing the results of a writhing test, demonstrating reversal of morphine analgesia by OFQ.

Fig. 14 is a graph that demonstrates reversal of morphine analgesia in animals by comparing tail withdrawal latency time in animals that have received varying doses of OFQ.

Fig. 15 is a graph of core body temperature vs. time, demonstrating attenuation of morphine hypothermia by OFQ.

Fig. 16 is a graph of withdrawal jumping precipitated by OFQ in morphine dependent mice, demonstrating precipitation of withdrawal signs in the mice.

Figs. 17A, 17B, and 17C is a series of graphs showing that OFQ does not bind with high affinity to mu, delta and kappa opioid receptors.

30

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is a method of antagonizing a physiologic effect of an opioid in an animal. As used in this specification, the terms "opiate" and "opioid" will be used as defined in Goodman and Gilman, The Pharmacological Basis of Therapeutics, 7th edition (hereinafter "Goodman and Gilman"), page 492. "Opiate" will be used to designate drugs derived from opium, such as morphine, codeine, and the many semi-synthetic congeners of morphine. "Opioid" will refer in a generic sense to all drugs or biological substances, natural and synthetic, with morphine-like actions. Hence the term "opioid" will include such synthetic morphine-like

compounds as propoxyphene, as well as the endogenous opioid peptides that include the enkephalins, endorphins, and dynorphins.

The term "anti-opioid" refers to a peptide or other chemical compound that antagonizes the physiological effects of an opioid. "Antagonizes" refers to a biochemical or pharmaceutical reversal or reduction of a physiologic effect. The peptides of the present invention specifically bind to an anti-opioid receptor, which is believed to exist separate from opioid receptors, and mediates physiologic antagonism of effects mediated by opioid receptors.

"High specificity" (selectivity) refers to binding with high affinity to a receptor, which in this specification shall be defined to bind with a K_i or IC_{50} of about 100 nM or less. In more particular embodiments wherein specific binding is particularly high, the K_i or IC_{50} is 10-100 nM, particularly 0.1 to 10 nM, and especially 0.1 nM or less. Methods of determining binding selectivity are well known, and for example are described in Goldstein, TIPS Reviews, pages 456-459, December 1987 (Vol. 8), which is incorporated by reference. The expression K_i is used as known in the art, and defined in J. Neurochem. 64:18-23 (1995), which is incorporated by reference. See particularly Table 1 of that incorporated reference.

As described in the incorporated reference, K_i values are calculated based on the IC_{50} value for the high affinity site using the equation of Cheng and Prusoff, Biochem. Pharmacol. 22:3099-3108 (1973): $K_i = IC_{50}/(1+[L]/K_D)$, where IC_{50} is the concentration of unlabeled ligand that inhibits 50% of the labeled ligand (L) binding, [L] is the total concentration of ligand provided in the assay, and the K_D is defined as the dissociation constant. IC_{50} may be determined by the method described in Example 12. The dissociation constant K_D is defined at Goodman and Gilman, page 41, which is incorporated by reference. Determination of the dissociation constant for an agonist receptor complex is well known in the art, and is, for example, determined by the methods described in Munson, Principles of Pharmacology (Chapman & Hall), 1995, Chapter 1, which is incorporated by reference. Determination of the affinity constant is particularly discussed at pages 17-18 thereof, and is a technique well known in the art. It may be determined from a conventional Scatchard analysis, wherein the slope of the Scatchard line is $-1/K_D$. Alternatively, K_D is determined from kinetic experiments in which $K_D = k_1/k_2$ where k_1 is the dissociation rate constant and k_2 is the association rate constant.

A physiologic effect of an opioid includes (without limitation) one or more of those biochemical changes and/or clinical signs and symptoms noted after administration of an opioid to an animal, and includes those effects enumerated at pages 497-504 of Goodman and Gilman. Such effects include one or more of inhibition of cellular cAMP accumulation, analgesia, drowsiness, changes in mood (including euphoria), pupillary responses, respiratory depression, nausea, peripheral vasodilation, hypothermia, and reduction of gastrointestinal motility. The invention also provides methods for designing OFQ receptor ligands and analogues, derivatives and mimetics thereof, using the amino acid sequence of the peptide ligands of the invention. Also

encompassed within the scope of this invention are such analogues, derivatives and mimetics produced by the methods of the invention.

In a second aspect, the invention provides nucleic acids encoding a novel mammalian opioid-specific receptor protein having an amino acid sequence identified as SEQ ID No.: 4, 5 recombinant eukaryotic expression constructs capable of expressing the novel mammalian opioid-specific receptor of the invention in cultures of transformed cells, as well as cultures of transformed eukaryotic cells that synthesize the receptor of the invention. The invention also provides homogeneous compositions of the receptor protein having an amino acid sequence identified as SEQ ID No.: 4, and antibodies against and epitopes of the receptor protein of the 10 invention. Methods for characterizing these receptor proteins and methods for using these proteins in the development of agents having pharmacological uses related to these receptors are also provided by the invention.

In this aspect of the invention is provided a nucleic acid having a nucleotide sequence encoding a mammalian OFQ receptor. In a preferred embodiment, the nucleic acid encodes 1452 15 nucleotides of the cDNA comprising 1101 nucleotides of coding sequence, 181 nucleotides of 5' untranslated sequence and 170 nucleotides of 3' untranslated sequence, depicted in Fig. 1 and identified as SEQ ID No.: 3. Encompassed in this aspect of the invention is the disclosed sequence and allelic variants of this sequence, either naturally occurring or the product of *in vitro* chemical or genetic modification.

20 The corresponding receptor protein, having the deduced amino acid sequence shown in Fig. 1 and identified as SEQ ID No.: 4, is also an aspect of the invention. A particular embodiment of this aspect of the invention is a homogeneous composition of the 47 kD mammalian OFQ receptor protein and derivatives thereof, said size being understood to be the size of the protein before any post-translational modifications. The amino acid sequence of the 47 25 kD receptor protein is depicted in Fig. 1 and identified as SEQ ID No.: 4.

This invention provides both nucleotide and amino acid sequence probes derived from the sequences herein provided, isolated from either cDNA or genomic DNA, as well as probes made synthetically with the sequence information derived therefrom. The invention specifically includes but is not limited to oligonucleotide, nick-translated, random primed, or *in vitro* 30 amplified probes made using cDNA or genomic clones embodying this aspect of the invention, and oligonucleotide and other synthetic probes synthesized chemically using the nucleotide sequence information of cDNA or genomic clone embodiments of the invention.

The present invention also includes synthetic peptides made using the nucleotide sequence information comprising the cDNA embodiments of the receptor protein embodiments of the 35 invention. The invention includes either naturally occurring or synthetic peptides which may be used as antigens for the production of receptor-specific antibodies, or useful as competitors of receptor molecules for agonist, antagonist or drug binding, or to be used for the production of inhibitors of the binding of agonists or antagonists or analogues thereof to such OFQ receptor

molecules. Particularly preferred embodiments of this aspect of the invention are such peptides that interact with the peptide ligand embodiment of the invention and enhance or inhibit peptide ligand binding to the receptor protein.

5 The present invention also provides antibodies against and epitopes of the mammalian opioid receptor molecules of the invention, including antisera and both polyclonal and monoclonal embodiments of such antibodies and hybridoma cell lines producing such monoclonal antibodies.

10 The present invention provides recombinant expression constructs comprising a nucleic acid encoding a mammalian OFQ receptor of the invention wherein the construct is capable of expressing the encoded OFQ in cultures of cells transformed with the construct. Preferred embodiments of such constructs comprise the OFQ receptor cDNA depicted in Fig. 1 (SEQ ID No.: 3), such constructs being capable of expressing the OFQ receptor encoded therein in cells transformed with the construct.

15 The invention also provides cultures of cells transformed with the recombinant expression constructs of the invention, each such culture being capable of and in fact expressing the mammalian OFQ receptor encoded in the transforming construct; and protein preparations of prokaryotic and eukaryotic cell membranes containing the OFQR protein of the invention, derived from cultures of prokaryotic or eukaryotic cells, respectively, transformed with the recombinant expression constructs of the invention.

20 The invention also provides methods for screening compounds for their ability to inhibit, facilitate or modulate the biochemical activity of the mammalian OFQ receptor molecules of the invention, for use in the *in vitro* screening of novel agonist and antagonist compounds. In preferred embodiments, cells transformed with a recombinant expression construct of the invention are contacted with such a compound, and the binding capacity of the compounds, as well as the effect of the compound on binding of other, known opioid agonists and antagonists, is 25 assayed. Additional preferred embodiments comprise quantitative analyses of such effects. The invention specifically provides a method for screening a compound for a capacity to bind to a mammalian OFQ receptor in cells expressing the receptor, and using a competitive assay between the compound to be tested and the peptide ligands of the invention. The method comprises the steps of:

30 (a) transforming a culture of eukaryotic or prokaryotic cells with a recombinant expression construct capable of expressing a mammalian OFQ receptor having an amino acid sequence identified as SEQ ID No.: 4, wherein the cells of the transformed cell culture express the receptor;

(b) assaying the transformed cell culture for binding of an amount of a detectably-labeled peptide according to Claim 1 in competition with varying amounts of the compound; and

(c) determining whether the compound competitively binds to the receptor by calculating the extent of inhibition of binding of the detectably-labeled peptide in the presence of the compound.

In additional embodiments of this method is included the additional step of:

5 (d) comparing the binding capacity of the compound with the binding capacities of additional compounds that are known to bind to mammalian opioid receptors, wherein said additional compounds comprise naturally-occurring and synthetic opioid receptor agonists and antagonists.

The present invention is also useful for the detection of analogues, agonists or antagonists, 10 known or unknown, of the mammalian OFQ receptor of the invention, either naturally occurring or embodied as a drug. In preferred embodiments, such analogues, agonists or antagonists may be detected in blood, saliva, semen, cerebrospinal fluid, plasma, lymph, or any other bodily fluid. In particularly preferred embodiments the peptide ligands of the invention are used in competitive binding assays to quantitatively evaluate novel ligand binding to the OFQ receptor.

15 Specific preferred embodiments of the present invention will become evident from the following more detailed description of certain preferred embodiments and the claims. The term "novel mammalian OFQ receptor" and "OFQR" as used herein refers to proteins comprising or consisting essentially of, and having substantially the same biological activity as, the protein encoded by the nucleic acid depicted in Fig. 1 (SEQ ID No.:3). This definition is intended to 20 encompass natural allelic variations in the disclosed OFQR sequence. Cloned nucleic acid provided by the present invention may encode OFQR protein of any species of origin, including, for example, mouse, rat, rabbit, cat, and human, but preferably the nucleic acid provided by the invention encodes OFQR receptors of mammalian, most preferably rat and human, origin.

The nucleic acids of the invention comprise nucleic acid hybridization probes comprising 25 DNA or RNA consisting essentially of the nucleotide sequence of the OFQ receptor, depicted in Fig. 1 (SEQ ID No.:3), or any portion thereof effective in nucleic acid hybridization. Mixtures of such nucleic acid hybridization probes are also within the scope of this embodiment of the invention. Nucleic acid probes as provided herein are useful for detecting OFQ receptor gene expression in cells and tissues using techniques well-known in the art, included but not limited to 30 Northern blot hybridization, *in situ* hybridization and Southern hybridization to reverse transcriptase-polymerase chain reaction product DNAs. The probes provided by the present invention, including oligonucleotide probes derived therefrom, are also useful for Southern hybridization of mammalian, preferably human, genomic DNA for screening for restriction fragment length polymorphisms (RFLPs) associated with certain genetic disorders.

35 The invention provides an isolated and purified, naturally-occurring, endogenous mammalian peptide ligand that specifically binds to the OFQ receptor of the invention. For the purposes of this invention it will be understood that the peptide is any biologically active molecule that specifically binds to an OFQ receptor of the invention. The peptides of the invention include

synthetic embodiments of the naturally-occurring peptide ligand isolated as described herein, as well as analogues, derivatives and variants of this peptide that specifically bind to the OFQR. Such analogues include substitution variants, wherein an amino acid is substituted conservatively with another amino acid that does not ablate the specific binding properties of the peptide ligand.

5 Specifically provided are substitution variants wherein the substituted amino acid is Leu¹⁴, wherein this residue is substituted with tyrosine.

Each peptide ligand of the invention is comprised of a sequence of amino acids. The term amino acid as used in this invention is intended to include all L- and D- amino acids, naturally occurring and otherwise.

10 Generally, those skilled in the art will recognize that peptides as described herein may be modified by a variety of chemical techniques to produce compounds having essentially the same activity as the unmodified peptide, and optionally having other desirable properties. For example, carboxylic acid groups of the peptide, whether carboxyl-terminal or sidechain, may be provided in the form of a salt of a pharmaceutically-acceptable cation or esterified to form a C₁-C₁₆ ester,

15 or converted to an amide of formula NR₁R₂ wherein R₁ and R₂ are each independently H or C₁-C₁₆ alkyl, or combined to form a heterocyclic ring, such as a 5- or 6- membered ring. Amino groups of the peptide, whether amino-terminal or sidechain, may be in the form of a pharmaceutically-acceptable acid addition salt, such as the HCl, HBr, acetic, benzoic, toluene sulfonic, maleic, tartaric and other organic salts, or may be modified to C₁-C₁₆ alkyl or dialkyl

20 amino or further converted to an amide.

Hydroxyl groups of the peptide sidechain may be converted to C₁-C₁₆ alkoxy or to a C₁-C₁₆ ester using well-recognized techniques. Phenyl and phenolic rings of the peptide sidechain may be substituted with one or more halogen atoms, such as fluorine, chlorine, bromine or iodine, or with C₁-C₁₆ alkyl, C₁-C₁₆ alkoxy, carboxylic acids and esters thereof, or amides of such carboxylic acids. Methylene groups of the peptide sidechains can be extended to homologous C₂-C₄ alkynes. Thiols can be protected with any one of a number of well-recognized protecting groups, such as acetamide groups. Those skilled in the art will also recognize methods for introducing cyclic structures into the peptides of this invention to select and provide conformational constraints to the structure that result in enhanced binding and/or stability. For example, a carboxyl-terminal or amino-terminal cysteine residue can be added to the peptide, so that when oxidized the peptide will contain a disulfide bond, thereby generating a cyclic peptide. Other peptide cyclizing methods include the formation of thioethers and carboxyl- and amino-terminal amides and esters.

30 It will be apparent to one skilled in the art that the anti-opioid activity of the peptides disclosed herein lies not in the precise amino acid sequence, but rather in the epitopes inherent in the amino acid sequences encoded by the DNA sequences. It will therefore also be apparent that it is possible to recreate the anti-opioid activity of one of these peptides by recreating the epitope, without necessarily recreating the exact amino acid sequence. This could be achieved either by

directly synthesizing the peptide (thereby circumventing the need to use the DNA sequences) or, alternatively, by designing a nucleic acid sequence that encodes for the epitope, but which differs, by reason of the redundancy of the genetic code, from the sequences disclosed herein. Similarly, the OFQR DNA sequence may also be varied, while still producing a functional OFQR protein.

5 Accordingly, the degeneracy of the genetic code further widens the scope of the present invention as it enables major variations in the nucleotide sequence of a DNA molecule while maintaining the amino acid sequence of the encoded protein. The genetic code and variations in nucleotide codons for particular amino acids is presented in Tables 1 and 2. Based upon the degeneracy of the genetic code, variant DNA molecules may be derived from the DNA sequences
10 disclosed herein using standard DNA mutagenesis techniques, or by synthesis of DNA sequences.

-14-

Table 1
The Genetic Code

5	First Position (5' end)	Second Position			Third Position (3' end)	
		T	C	A	G	
10						
15		Phe	Ser	Tyr	Cys	T
	T	Phe	Ser	Tyr	Cys	C
		Leu	Ser	Stop (och)	Stop	A
		Leu	Ser	Stop (amb)	Trp	G
20		Leu	Pro	His	Arg	T
	C	Leu	Pro	His	Arg	C
		Leu	Pro	Gln	Arg	A
		Leu	Pro	Gln	Arg	G
25		Ile	Thr	Asn	Ser	T
	A	Ile	Thr	Asn	Ser	C
		Ile	Thr	Lys	Arg	A
		Met	Thr	Lys	Arg	G
30		Val	Ala	Asp	Gly	T
	G	Val	Ala	Asp	Gly	C
		Val	Ala	Glu	Gly	A
		Val (Met)	Ala	Glu	Gly	G

"Stop (och)" stands for the ochre termination triplet, and "Stop (amb)" for the amber. ATG is the most common initiator codon; GTG usually codes for valine, but it can also code for methionine to initiate an mRNA chain.

Table 2
The Degeneracy of the Genetic Code

5	Number of Synonymous Codons	Amino Acid	Total Number of Codons
10	6	Leu, Ser, Arg	18
	4	Gly, Pro, Ala, Val, Thr	20
	3	Ile	3
	2	Phe, Tyr, Cys, His, Gln, Glu, Asn, Asp, Lys	18
15	1	Met, Trp	2
	Total number of codons for amino acids		61
	Number of codons for termination		<u>3</u>
	Total number of codons in genetic code		64

20

Additionally, standard mutagenesis techniques may be used to produce peptides which vary in amino acid sequence from the peptides encoded by the DNA molecules disclosed herein. However, such peptides will retain the essential characteristic of the peptides encoded by the DNA molecules disclosed herein, i.e. the ability to antagonize opioid action by binding to the OFQR. This characteristic can readily be determined by the assay technique described herein for cAMP accumulation. Screening for retention of anti-opioid effects is similarly achieved in a forthright manner using the methods of Examples 8-11 below. Variant peptides include those with variations in amino acid sequence, including minor deletions, additions and substitutions.

25

While the site for introducing an amino acid sequence variation is predetermined, the mutation *per se* need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed protein variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence as described above are well known.

30

In order to maintain a functional peptide, preferred peptide variants will differ by only a small number of amino acids from the peptides encoded by the native DNA sequences. Preferably, such variants will be amino acid substitutions of single residues. Substitutional variants are those in which at least one residue in the amino acid sequence has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Table 3 when it is desired to finely modulate the characteristics of the protein.

Table 3 shows amino acids which may be substituted for an original amino acid in a protein and

which are regarded as conservative substitutions. As noted, all such peptide variants are tested to confirm that they retain the ability to antagonize opiate action.

TABLE 3

5

Original Residue

Conservative Substitutions

10	Ala	ser
	Arg	lys
	Asn	gln, his
	Asp	glu
	Cys	ser
	Gln	asn
15	Glu	asp
	Gly	pro
	His	asn; gln
	Ile	leu, val
	Leu	ile; val
20	Lys	arg; gln; glu
	Met	leu; ile
	Phe	met; leu; tyr
	Ser	thr
	Thr	ser
25	Trp	tyr
	Tyr	trp; phe
	Val	ile; leu

30 Substantial changes in biological activity may be made by selecting substitutions that are less conservative than those in Table 2, i.e., selecting residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

35 Peptidomimetic and organomimetic embodiments are also hereby explicitly declared to be within the scope of the present invention, whereby the three-dimensional arrangement of the chemical constituents of such peptido- and organomimetics mimic the three-dimensional arrangement of the peptide backbone and component amino acid sidechains in the peptide, resulting in such peptido- and organomimetics of the peptides of this invention having substantial

40 biological activity. For computer modelling applications, a pharmacophore is an idealized, three-dimensional definition of the structural requirements for biological activity. Peptido- and organomimetics can be designed to fit each pharmacophore with current computer modelling software (computer aided drug design). The degree of overlap between the specific activities of pharmacophores remains to be determined. It will be understood that mimetics prepared using 45 such techniques that specifically bind to the OFQ receptor and developed using the peptide ligands of the invention will fall within the scope of the appended claims.

The peptides provided by the present invention can be chemically synthesized by any of a number of manual or automated methods of synthesis known in the art. Automated synthetic routines such as those available for use with automated peptide synthesizers are also intended to come within the scope of the present invention. Chemical derivatization, using the 5 methods disclosed in this specification or other methods well known in the art, of naturally- occurring peptides or peptides purified from mixtures of protein degradation products, degraded by enzymatic or chemical means, are also within the scope of this invention, as are peptides made by molecular or genetic engineering means. Preferably, solid phase peptide synthesis (SPPS) is carried out on a 0.25 millimole (mmole) scale using an Applied Biosystems Model 431A Peptide 10 Synthesizer and using 9-fluorenylmethyloxycarbonyl (Fmoc) amino-terminus protection, coupling with dicyclohexylcarbodiimide/hydroxybenzotriazole or 2-(1H-benzo-triazol-1-yl)-1,1,3,3- tetramethyluronium hexafluorophosphate/ hydroxybenzotriazole (HBTU/HOBT), and using *p*- hydroxymethylphenoxyethylpolystyrene (HMP) or Sasrin™ resin for carboxyl-terminus acids or Rink amide resin for carboxyl-terminus amides.

15 Fmoc-derivatized amino acids are prepared from the appropriate precursor amino acids by tritylation and triphenylmethanol in trifluoroacetic acid, followed by Fmoc derivitization as described by Atherton *et al.* (1989, Solid Phase Peptide Synthesis, IRL Press: Oxford).

20 Sasrin™ resin-bound peptides are cleaved using a solution of 1% TFA in dichloromethane to yield the protected peptide. Where appropriate, protected peptide precursors are cyclized between the amino- and carboxyl-termini by reaction of the amino-terminal free 25 amine and carboxyl-terminal free acid using diphenylphosphorylazide in nascent peptides wherein the amino acid sidechains are protected.

30 HMP or Rink amide resin-bound products are routinely cleaved and protected sidechain-containing cyclized peptides deprotected using a solution comprised of trifluoroacetic acid (TFA), optionally also comprising water, thioanisole, and ethanedithiol, in ratios of 100 : 5 : 5 : 2.5, for 0.5 - 3 hours at room temperature.

35 Crude peptides are purified by preparative high pressure liquid chromatography (HPLC) using a Waters Delta-Pak C18 column and gradient elution with 0.1% TFA in water modified with acetonitrile. After column elution, acetonitrile is evaporated from the eluted fractions, which are then lyophilized. The identity of each product so produced and purified is confirmed by fast atom bombardment mass spectroscopy (FABMS) or electrospray mass spectroscopy (ESMS).

The production of proteins such as the OFQ receptor proteins of the invention from cloned genes by genetic engineering means is well known in this art. The discussion which 40 follows is accordingly intended as an overview of this field, and is not intended to reflect the full state of the art.

45 DNA encoding an opioid receptor may be obtained, in view of the instant disclosure, by chemical synthesis, by screening reverse transcripts of mRNA from appropriate

cells or cell line cultures, by screening genomic libraries from appropriate cells, or by combinations of these procedures, as illustrated below. Screening of mRNA or genomic DNA may be carried out with oligonucleotide probes generated from the nucleic acid sequence information from the receptor (OFQR) disclosed herein. Probes may be labeled with any 5 detectable group and used in conventional hybridization assays, as described in greater detail in the Examples below. In the alternative, amino acid transporter-derived nucleic acid sequences may be obtained by use of the polymerase chain reaction (PCR) procedure, using PCR oligonucleotide primers corresponding to nucleic acid sequence information derived from an OFQR as provided herein. *See* U.S. Patent Nos. 4,683,195 to Mullis *et al.* and 4,683,202 to 10 Mullis.

OFQR protein may be synthesized in host cells transformed with a recombinant expression construct comprising a nucleic acid encoding the OFQ receptor cDNA. For the purposes of this invention, a recombinant expression construct is a replicable DNA construct in which a nucleic acid encoding an opioid receptor is operably linked to suitable control sequences 15 capable of effecting the expression of the opioid receptor in a suitable host. Generally, control sequences include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and sequences which control the termination of transcription and translation. *See*, Sambrook *et al.*, 1990, Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Press: New York).

20 Vectors useful for practicing the present invention include plasmids, viruses (including phage), retroviruses, and integratable DNA fragments (i.e., fragments integratable into the host genome by homologous recombination). A preferred vector is RcrVS (Invitrogen, San Diego, CA).

25 Cultures of cells derived from multicellular organisms are a desirable host for recombinant opioid receptor protein synthesis. Transformed host cells are cells which have been transformed or transfected with recombinant expression constructs made using recombinant DNA techniques and comprising nucleic acid encoding an amino acid transporter protein. Transformed host cells may express the OFQR protein; when expressed, the OFQ receptor of the invention will typically be located in the host cell membrane. *See*, Sambrook *et al.*, *ibid.* In principal, any 30 higher eukaryotic cell culture is useful, whether from vertebrate or invertebrate culture. However, mammalian cells are preferred, as illustrated in the Examples. Propagation of such cells in cell culture has become a routine procedure. *See* Tissue Culture, Academic Press, Kruse & Patterson, editors (1973). Examples of useful host cell lines are human 293 cells, VERO and HeLa cells, Chinese hamster ovary (CHO) cell lines, mouse Ltk cell lines and WI38, BHK, 35 COS-7, CF, and MDCK cell lines. CHO cells, COS-7 cells and Ltk cells are preferred.

The recombinant expression constructs of the present invention are useful to transform cells which do not ordinarily express an OFQ receptor to thereafter express the receptor. Such cells are useful as intermediates for making cell membrane preparations useful for

receptor binding activity assays, which are in turn useful for screening drugs or new peptides made in accordance with the present invention. The recombinant expression constructs of the present invention thus provide a method for obtaining reagents for screening potentially useful peptides and other drugs at advantageously lower cost than conventional animal screening 5 protocols. While not completely eliminating the need for ultimate *in vivo* activity and toxicology assays, the constructs and cultures of the invention provide an important first screening step for the vast number of potentially useful psychoactive drugs synthesized, discovered or extracted from natural sources each year.

The recombinant expression constructs of the present invention are useful to detect, 10 isolate, characterize and identify other novel endogenous receptors for anti-opioid agonists (such as OFQ) and antagonists found in plasma, serum, lymph, cerebrospinal fluid, seminal fluid, or other potential sources of such compounds.

The utility of the present invention for using the nucleic acids of the invention to produce cell membranes containing the receptor protein encoded thereby, and the demonstrated 15 utility of this aspect of the invention to permit the isolation, characterization and identification of a novel peptide, termed orphanin FQ (OFQ) herein, as an endogenous receptor-binding ligand, enables rational drug design of novel therapeutically-active drugs using currently-available techniques (see Walters, "Computer-Assisted Modeling of Drugs", in Klegerman & Groves, eds., 1993, Pharmaceutical Biotechnology, Interpharm Press: Buffalo Grove, IL, pp. 165-174).

20 The invention provides homogeneous compositions of a novel mammalian receptor protein produced by transformed eukaryotic cells as provided herein. Each such homogeneous composition is intended to be comprised of the OFQR protein that comprises at least 90% of the protein in such homogenous composition. The invention also provides membrane preparations from cells expressing the OFQR protein as the result of transformation with a recombinant 25 expression construct, as described herein.

Mammalian receptor proteins made from cloned genes in accordance with the present invention may be used for screening analogues of an anti-opioid system that operates in parallel with and in functional antagonism to the opioid system. The receptor proteins may also be useful to find agonists or antagonists of opioid binding, or for determining the amount of such 30 agonists or antagonists that are present in a solution of interest (e.g., blood plasma, cerebrospinal fluid or serum). For example, host cells may be transformed with a recombinant expression construct of the present invention, a mammalian OFQR expressed in those host cells, and the cells or membranes thereof used to screen compounds for their effect on anti-opioid receptor binding activity. By selection of host cells that do not ordinarily express an OFQR, pure 35 preparations of membranes containing the transporter can be obtained.

The invention also provides ligands for such OFQR proteins, including naturally-occurring ligands and synthetic embodiments thereof. Methods for identifying other, alternative ligands, and agonists and antagonists of OFQ peptide ligand binding are also provided by the

invention. Such methods include competitive binding assays, wherein the peptide ligands of the invention are detectably labeled and incubated under competitive binding conditions with varying amounts of any putative ligand compound to be tested. The extent of inhibition of labeled ligand binding is useful in characterizing the extent and affinity of binding of novel ligands to the 5 OFQR. Specificity of binding can also be determined using such assays. Preferably, the peptide ligand of the invention being used in such competition experiments is detectably labeled with a radioisotope, such as I-123, I-125 and I-131.

10 The invention also provides antibodies that are immunologically reactive to the opioid receptor protein or epitopes thereof provided by the invention. The antibodies provided by the invention may be raised using methods well known in the art.

15 The present invention also provides monoclonal antibodies that are immunologically reactive with an epitope derived from an OFQ receptor of the invention, or fragment thereof. Such antibodies are made using methods and techniques well known to those of skill in the art. Monoclonal antibodies provided by the present invention are produced by hybridoma cell lines, that are also provided by the invention and that are made by methods well known in the art.

20 The ligands of the invention are useful as diagnostic and therapeutic agents when constituted as pharmaceutical compositions with the appropriate carriers or diluents. In diagnostic embodiments, detectably-labeled peptide ligands are used in methods for diagnosing diseases or pathological conditions related to ligand binding of OFQ receptors *in vivo*. Similarly, therapeutic methods of treatment are encompassed by the invention and provided using pharmaceutical 25 compositions of such peptides administered *in vivo* in therapeutically-effective amounts.

30 Preparation of pharmaceutically acceptable compositions of the peptides of the present invention can be accomplished using methods well known to those with skill in the art. Any of the common carriers such as sterile saline solution, plasmas, etc., can be utilized with the peptides provided by the invention. Routes of administration include but are not limited to oral, intracranial ventricular (icv), intrathecal (it), intravenous (iv), parenteral, rectal, topical ophthalmic, subconjunctival, nasal, aural and transdermal. Peptides of the invention may be administered intravenously in any conventional medium for intravenous injection such as an aqueous saline medium, or in blood plasma medium. Such medium may also contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, buffers, preservatives and the like. Among the preferred media are normal saline and plasma.

35 Embodiments of the invention comprising medicaments can be prepared with conventional pharmaceutically acceptable carriers, adjuvants and counterions as would be known to those of skill in the art. The medicaments are preferably in the form of a unit dose in solid, semi-solid and liquid dosage forms such as tablets, pills, powders, liquid solutions or suspensions, and injectable and infusible solutions, for example a unit dose vial. Effective dosage ranges

included in the unit dose container vary from about 100 $\mu\text{g}/\text{kg}$ to about 10 mg/kg of body weight are contemplated.

The invention also provides methods for detecting the amount of an analyte in a solution wherein the analyte is an endogenous peptide ligand of the invention in a biological sample, such as blood, serum, plasma or cerebrospinal fluid. In such methods is provided a first mixture comprised of cells or membranes heterologously expressing the OFQ opioid receptor of the invention, a second mixture comprised of a standard amount of a detectably-labeled embodiment of the peptide ligands of the invention, and a third mixture comprised of a diagnostically-significant tissue sample or bodily fluid. From these mixtures is produced a specific binding reaction mixture by contacting the first, second and third mixtures and incubating the reaction mixture for a time sufficient to allow binding between the peptide ligand and the OFQ receptor. The extent of the binding reaction is determined by calculating the amount of the detectably-labeled peptide ligand of the invention bound to the OFQ receptor, and comparing the amount of binding of the peptide ligand to the OFQ receptor in the presence of the tissue sample or bodily fluid of the third provided mixture with the amount of binding found in the absence of the tissue sample or bodily fluid of the third provided mixture.

Throughout these examples, injections are sometimes given intrathecally or intraventricularly. The intrathecal injection method allows the application of small quantities of drug directly into the spinal cord, without surgical intervention. The mouse is briefly exposed to halothane, immediately prior to all injections.

Intrathecal injection is achieved with disposable 30 g, $\frac{1}{2}$ inch needles mated to a 10 μl luer tip syringe (Hamilton, Reno, NV) are used. Drug is administered in artificial cerebral spinal fluid and injection volumes are 2-5 μl . The mouse is placed on a flat surface, with its head in a towel cupped gently with the heel of the palm, and the thumb and forefinger lightly pushing down on the iliac crests with enough force to stretch the skin across the lower back. The needle is inserted into the tissue just medial to the L5 spinous process so that it enters the groove between the spinous and transverse processes. This site maximizes vertebral accessibility and minimizes the possibility of spinal damage (into the cauda equina). The needle is then moved carefully forward to the intervertebral space as the angle of the syringe is decreased from about 30° to 15°. The tip of the needle is inserted so that approximately 0.5 cm is within the vertebral column which is reliably indicated by a flick of the mouse's tail.

In a human, conventional lumbar puncture with a spinal needle can be used to access the epidural or intrathecal space.

For icv injections, a 26g $\frac{1}{2}$ inch needle is fitted with a piece of polyethylene tubing so that 3mm of uncovered needle enter the mouse's skull. (The sleeve ensures that the depth of the needle is appropriate for the depth of the lateral ventricle). Mice are briefly anesthetized with halothane. The needle is held at a 45° angle and inserted at Bregma, 1-2 mm lateral to midline. The needle readily punctures the skull at this suture, so no incision is required for accurate

placement. The needle tip is towards the tail, and the 3mm depth allows direct placement of the drug into a lateral ventricle.

The Examples which follow illustrate specific embodiments of the invention, and various uses thereof. They are set forth for explanatory purposes only, and are not to be taken as 5 limiting the invention.

EXAMPLE 1

Isolation of a Mammalian Opioid Receptor Probe by Random PCR Amplification of Rat Brain-derived cDNA Using 10 Degenerate Oligonucleotide Primers

In order to clone novel mammalian G-protein coupled receptors, cDNA prepared from RNA from different regions of mammalian (rat) brain was used as template for a polymerase chain reaction (PCR)-based random cloning experiment. PCR was performed using a pair of degenerate oligonucleotide primers derived from a mouse δ -opioid receptor (Kieffer *et al.*, 15 1992, Proc. Natl. Acad. Sci. USA 89:12048-12052; Evans *et al.*, 1992, Science 258:1952-1955). PCR products obtained in this experiment were characterized by nucleotide sequencing and used to isolate a full-length cDNA from a rat brain cDNA library.

The PCR amplification experiments were performed as described in co-owned and co-pending U.S. Serial No. 08/149,093, incorporated by reference, using the following primers:
20 Primer III (sense):

ATGAATTCAC(G/A/C/T)(A/G)T(G/C)ATGAG(C/T)GT(G/C)GAC(C/A)G(C/A)TA
(SEQ ID NO:1)

and

Primer VII (antisense):

25 TTGTCGAC(G/A)TA(G/A)AG(A/G)A(T/C)(G/A/C/T)GG(G/A)TT
(SEQ ID NO:2).

Amplified products of the PCR reaction were separated on a 1.0% agarose gel (*see* Sambrook *et al.*, *ibid.*), and fragments ranging in size from 400 basepairs (bps) to 750 bp were subcloned in the plasmid vector pBluescript (Stratagene, LaJolla, CA). A multiplicity of bacterial 30 colonies comprising each of the subcloned fragments were used to make bacterial colony lifts on nitrocellulose filters using conventional techniques (*see* Sambrook, *et al.*, *ibid.*). Such filters were hybridized with a [32 P]-dCTP-labeled radioactive nucleic acid probe comprising a full-length mouse δ -opioid receptor cDNA at a concentration of 1×10^6 cpm/mL under low stringency 35 hybridization conditions [35% formamide, 5X standard citrate saline (SSC; wherein 1X SSC is 0.15M NaCl/0.015M sodium citrate, pH 7.0), 5X Denhardt's solution (wherein 1X Denhardt's solution is 0.02g/mL each of bovine serum albumin, Ficoll and polyvinylpyrrolidone)] at 37°C overnight. After hybridization, the filters were washed in a solution of 2X SSC/0.1% sodium dodecyl sulfate (SDS) at 55°C and then exposed to X-ray film (XAR-5, Eastman-Kodak,

Rochester, NY) for 2 days at -70°C using tungsten-impregnated intensifying screens (DuPont-NEN, Wilmington, DE). Plasmid DNA from hybridizing clones was purified and the nucleotide sequence of the insert cDNA determined by the dideoxynucleotide chain termination method (Sanger *et al.*, 1977, Proc. Natl. Acad. Sci. USA 74:5463-5467) using Sequenase® (U.S. 5 Biochemical Corp., Cleveland, OH).

EXAMPLE 2

Isolation of a Novel Mammalian Opioid Receptor cDNA

One of the PCR products (termed LC132) was isolated and sequenced in this way 10 and found to have a high degree of homology to the mouse δ -opioid receptor sequence (Evans *et al.*, *ibid.* and Kieffer *et al.*, *ibid.*). A full-length cDNA clone corresponding to this PCR fragment was isolated from a cDNA library prepared in the cloning vector λ gt11 comprising 15 oligo(dT)-primed rat brain cDNA, as described in co-owned and co-pending U.S. Serial No. 07/149,093.

Nucleotide sequence analysis performed essentially as described (*see*, Sambrook, *ibid.*) revealed the sequence shown in Fig. 1 (SEQ ID No.:3). The putative protein product of 20 the gene is also shown in Fig. 1 (SEQ ID No.:4). The sequence was found to have an open reading frame comprising 1101 nucleotides encoding a protein 367 amino acids in length, and having a predicted molecular weight of 47 kilodaltons prior to post-translational modification.

The sequence immediately 5' to the proposed initiation codon was found to contain several 25 translation termination codons in-frame with the open reading frame, supporting the assignment of the translation start site. Predicted transmembrane domains [using the algorithm of Eisenberg *et al.* (1984, *J. Molec. Biol.* 179:125-142)] are boxed and identified by Roman numerals (I-VII), and three sites of possible N-linked glycosylation are identified in the amino-terminal portion of 30 the protein with solid triangles. Potential protein phosphorylation sites found in predicted cytoplasmic loops are marked with an asterisk. Further, a pair of cysteine residues conserved among known opioid receptors were found in the first and second predicted extracellular loops. On the basis of this analysis, this cloned nucleic acid was determined to be a novel mammalian receptor related to opioid receptors, such as the μ , δ and κ receptors.

The predicted amino acid sequences of this novel receptor, the rat μ -opioid receptor 35 (Chen *et al.*, *ibid.*), the mouse δ -opioid receptor (Evans *et al.*, *ibid.* and Kieffer *et al.*, *ibid.*) and the mouse κ -opioid receptor (Yasuda *et al.*, *ibid.*) are aligned in Fig. 2. Overbars indicate predicted transmembrane regions I through VII in the protein product of the genes. Amino acid residues that are found in common between all four mammalian opioid receptors are presented in boldface.

Overall, the novel mammalian receptor disclosed herein had 47% overall identity with the other mammalian opioid receptors, which similarity rose to 67% when only the predicted transmembrane domains were considered. Comparisons were made individually at each

transmembrane domain (TMI-TMVII), as an average over all transmembrane domains (TMavg) and as the average degree of amino acid sequence homology for each protein as a whole (avg/all). In total, 145 of the 367 residues are shared with the other mammalian opioid receptors, confirming the conclusion that the novel mammalian receptor disclosed herein is related to the opioid receptors. The inventors have now surprisingly discovered, however, that the receptor mediates an anti-opioid effect.

EXAMPLE 3

Construction of a Recombinant Expression Construct, DNA Transfection and Functional Expression of the Novel Mammalian Opioid Receptor

In order to biochemically characterize the novel mammalian opioid receptor described in Example 2, and to confirm that it encodes a novel opioid receptor, the cDNA was cloned into a mammalian expression construct, the resulting recombinant expression construct transfected into COS-7 cells (for transient expression assays) and mouse Ltk⁻ cells (for stable expression assays), and cell membranes (COS-7) or cell lines (Ltk⁻) were generated that expressed the receptor protein in cellular membranes at the cell surface. Such cells and membranes isolated from such cells were used for biochemical characterization experiments as described in U.S. Serial No. 149,093.

20 The entire coding region of the receptor cDNA insert was subcloned into the
RcRSV vector (Invitrogen, San Diego, CA) using conventional techniques (see Sambrook et al.,
ibid.). Such recombinant expression constructs were introduced into COS-7 cells using the
calcium-phosphate precipitation technique (Chen & Okayam, 1987, Molec. Cell Biol. 7:2745-
2752), the transfected cells allowed to express the receptor for between 24-96 hours, and then cell
25 membranes containing the receptor were isolated. The protein concentration was adjusted to a
15-80 μ g/sample for each of the binding studies described below.

These recombinant expression constructs were also introduced into Ltk⁻ cells using the calcium-phosphate precipitation technique, and stably-transfected clones were selected by growth in the mammalian neomycin analog G418 (Grand Island Biological Co., Long Island, NY), as the vector RcRSV contains a functional copy of a bacterial neomycin resistance gene. Stable cell lines were then selected for membrane binding studies based on mRNA expression levels of individual neomycin-resistant transfected clones determined by Northern analysis (see Sambrook et al., *ibid.*). Cell membranes were prepared and used as described above for COS-7 cell transfectants.

35 Specific binding assays using a variety of opioid receptor agonists and antagonists were performed on membranes from both transient and stable transfectants. Ligand binding experiments were performed essentially as described in Bunzow et al. (1988, *Nature* 336:783-787), and U.S. Serial No. 08/149,093). In binding experiments, increasing amounts of

membrane protein (from 15-80 μ g) were incubated with the radioactively-labeled opioid agonist or antagonist to be tested for 120 min at 22°C in a total volume of 1 ml. However, in these experiments no specific binding was found for the following compounds (their known receptor binding specificities are noted in parentheses): [3 H]-Tyr-dAla-Gly-Met-Phe-Gly-ol (DAMGO; μ -opioid receptor agonist), [3 H]-c[D-penicillamine², D-penicillamine⁵]enkephalin (DPDPE; δ agonist), [3 H]-U-69,593 (κ agonist), [3 H]-diprenorphine (μ agonist), [3 H]-bremacozine (κ agonist), [3 H]-dihydromorphine (μ agonist), [3 H]-ethylketocyclazocine (κ agonist) or [125 I]- β -endorphin. Although low levels of specific binding were seen using [3 H]-naloxone (μ antagonist), the significance of these results was compromised by the fact that untransfected COS-7 and Ltk⁻ cells also shown endogenous low levels of specific [3 H]-naloxone binding.

Surprisingly, however, specific binding was found using [3 H]-methadone. The results of Scatchard analysis of the methadone binding data are shown in Fig. 4.

For Scatchard analysis experiments, 0.25 ml aliquots of crude plasma membrane homogenate from transfected cell cultures was incubated in duplicate with increasing concentrations of [3 H]methadone (70.3 Ci/mmol; 10-3000 pM final concentration) under conditions described above. The estimated value for B_{max} was derived from these data using the LIGAND computer program. Panel A of Fig. 4 shows the results of radiolabeled methadone binding with untransfected COS-7 cells; similar results were found with Ltk⁻ cell membranes. These results demonstrate no or negligible amounts of endogenous methadone binding by these cell membranes. Panel B shows the results using COS-7 cells transfected with the RcRSV-LC132 expression construct. The levels of specific binding shown in this graph correspond to a dissociation constant (K_D) of about 10⁻¹⁰M for methadone and a B_{max} of about 400-450 femtomoles/ μ g protein for the novel mammalian opioid receptor expressed by these cells.

Thus, the novel mammalian opioid receptor disclosed herein has the heretofore unknown property of exhibiting specific binding to the opiate analog, methadone, while showing no specific binding to a variety of other known opioid receptor agonists and antagonists. These results support the conclusion that the receptor disclosed herein is a completely novel and heretofore unsuspected member of the opioid receptor family, termed herein therefore OFQ.

30

EXAMPLE 4

Brain Tissue Distribution of OFQR Expression

The distribution of mRNA corresponding to expression of the OFQ receptor gene in various regions of the rat brain was determined by *in situ* hybridization of rat brain slices, as described in U.S. Serial No. 08/149,093.

Results of these experiments are shown in Fig. 3. Panel A shows a section through the frontal cortex, preoptic area and caudate putamen; Panel B shows a section through the hypothalamus, thalamus and hippocampus; and Panel C shows a section through the pons and

cerebellum. These experiments localized high level OFQR expression in the hypothalamus (arcuate (Arc), posterior (PH), lateral (LH) and ventromedial (VMH) hypothalamic nuclei, Panel B), certain nuclei of the thalamus (paraventricular thalamic nuclei (PVP), Panel B), the medial habenula (MHb, Panel B), the CA regions of the hypothalamus, the dentate gyrus (DG, Panel B), 5 the locus coeruleus and certain cortical areas (medial preoptic area (MPA), Panel A and the cortex (CX), Panel B). Virtually no signal was seen in the caudate putamen (Cpu, Panel A) or cerebellum (Cb, Panel C). Strong hybridization was also detected in sections of the brainstem (Panel C) and the spinal cord (not shown).

These results demonstrate that the OFQ receptor disclosed herein is expressed in rat 10 brain in a variety of anatomically-distinct sites, suggesting an important role for this receptor in both higher brain function and central nervous system control of motor and sensory nerve signalling. The anti-opioid effects mediated by this receptor, however, were completely unpredictable.

15

EXAMPLE 5

Identification of an Endogenously-occurring Peptide Ligand for the OFQ Receptor

Due to the high level of sequence homology between the OFQ receptor and μ -, δ - and κ -opioid receptors found as disclosed in Example 2, it was appreciated that the novel 20 receptor disclosed herein might specifically recognize an endogenous peptide ligand, and respond to binding such a ligand using a second messenger signalling system similar to those found associated with the previously-described opioid receptors. Since the OFQR had been found to be expressed in cells in the central nervous system in Example 4, and since OFQR mRNA is highly expressed in hypothalamus, porcine hypothalamic homogenates were screened for receptor 25 binding activity. Inhibition of forskolin-stimulated, cyclic adenosine monophosphate (cAMP) accumulation in cells heterologously expressing OFQR was used as an assay, which assay had been used to characterize the interaction of opioid receptors and their ligands.

Acetic acid extracts of porcine hypothalamic tissues were prepared as follows. 4.5kg of freshly frozen porcine hypothalamic tissue were extracted in 9L of a solution of 0.5M 30 acetic acid/10mM ascorbic acid/1mM EDTA. The extract was centrifuged to remove insoluble debris and the supernatant absorbed batchwise onto a C₁₈silica matrix. Unbound material was removed by washing with water, and specifically-bound material was then eluted in a solution of 80% methanol. A total of 2L of methanolic eluate were then concentrated by rotary evaporation of a final volume of 44 mL, and material having a molecular weight less than 10 kilodaltons was 35 obtained by ultrafiltration using an Amicon Centriprep 10 column (Amicon, Beverly, MA). This material was applied in ten separate experiments to a cation exchange HPLC column (Protein Pak SP 8HR, 10x100mm, Waters) equilibrated with 10mM NH₄COO. The column was developed with a linear gradient of NH₄COO in 10% methanol at a flow rate of 1mL/min; a total of 80 1mL

fractions were collected. 10% (100 μ L) of each of 5 consecutive fractions were pooled and lyophilized, resulting in 16 pools. 5% of each pool was tested in duplicate for the capacity to inhibit forskolin-stimulated cAMP accumulation in cells heterologously expressing the OFQ opioid receptor.

5 Assays of ligand binding-associated inhibition of forskolin-stimulated cAMP accumulation in cells heterologously expressing the OFQ receptor were performed as follows. CHO cells, deficient in dihydrofolate reductase (*dhfr*) were transfected with OFQR-encoding cDNA cloned into the expression vector pRcRSV (Invitrogen), as described in Example 3 above, using calcium phosphate co-precipitation (Okayama and Chen, *ibid.*) Stably-transfected clones 10 were selected using G418, and screened for OFQ expression using a reverse transcriptase-PCR protocol (RT-PCR). One clone that tested positively for OFQR expression (LC-7) was used in ligand binding experiments.

15 For determination of cAMP levels, receptor-transfected CHO cells (i.e., LC-7) or untransfected CHO/*dhfr* cells, were plated in 24-well culture plates and grown to confluence. After removal of culture media, aliquots of HPLC fractions prepared as described above in a total 20 volume of 0.2mL DMEM containing 10mM HEPES buffer, pH 7.4, 1mM forskolin and 1mM Ro 20-1724 (Rolipram, RBI) were added per well and cells incubated at 37°C for 10 min. Cellular reactions were halted by the addition of 0.5mL ice-cold ethanol, and plates were stored for 12h or overnight at 80°C. Frozen plates were centrifuged and aliquots of the supernatant were removed 25 and dried onto 96-well plates for cAMP determinations. cAMP determinations were performed using a commercially-available kit (Biotrak SPA, Amersham) essentially according to the manufacturer's instructions.

25 The results of these experiments are shown in Fig. 5. Open bars represent results obtained with OFQ-expressing transfected CHO cells, and shaded bars represent results obtained with untransfected CHO cells. Pools 13 and 14 (corresponding to fractions 61-70) consistently showed adenyl cyclase inhibitory activity in transfected versus untransfected cells. The large increase in cAMP detected in fraction 2 is due to endogenous cAMP extracted from the tissue. Fractions 60-67 were found to contain most of the detected bioactivity, and were pooled for further purification by reverse-phase HPLC.

30 Reverse-phase HPLC was performed on pooled fractions 60-67 as follows. The complete bioactive material was loaded onto an octyl silica column (Superspher RP Select B, 2x125mm, Merck) and eluted at a flow rate of 0.12mL/min with a linear gradient of acetonitrile in 0.1% trifluoroacetic acid (TFA). Aliquots of fractions from this gradient were tested as described above, and the results of these assays shown in Fig. 6. Inhibition of forskolin-stimulated cAMP in OFQR-transfected CHO cells (shaded bars) was detected only in the major peak of eluted protein (measured as absorbance at 214nm (A₂₁₄)).

The isolated material having adenyl cyclase inhibitory activity was analyzed by mass spectrometry and sequenced by Edman degradation. The material was determined to be a peptide having the sequence:

Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln (SEQ ID No.: 5).

5 (Abbreviations for amino acids can be found in Zubay, *Biochemistry* 2d ed., 1988, MacMillan Publishing: New York, p. 33). Final yields of this peptide were about 200 picomoles from 4.5kg hypothalamic tissue (wet weight). A computer database search revealed that this peptide had not been reported, either as a unique entity or as part of a larger protein. This peptide is designated orphanin FQ (OFQ) herein.

10 The primary structure of this novel peptide was compared with the primary structure of a number of naturally-occurring opioid peptides, as shown in Fig. 7. The amino terminal tetrapeptide sequence motif YGGF of the known opioid receptor ligands is strikingly similar to the amino terminal sequence FGGF found in OFQ. Further, two clusters of basic amino acids found in the orphanin FQ peptide resemble the arrangement of positively-charged 15 residues in Dynorphin A and β -Endorphin. However, none of these peptides could be shown to bind to OFQR, suggesting a structural/functional divergence of the orphanin FQ peptide from these other opioid receptor ligands.

20 In order to verify that the isolated material had the observed properties of the pooled fraction from which it is isolated, a synthetic peptide having the deduced sequence was made and was found to be identical to the isolated peptide in its elution profile in reverse-phase HPLC assay and in molecular weight as determined by mass spectrometry. Moreover, the synthetic peptide was also found to inhibit adenyl cyclase production in forskolin-stimulated CHO cells transfected with and heterologously expressing OFQR, having an EC₅₀ of 1.58nM (\pm 0.666nM) and showing maximal inhibition (80%) at a concentration of about 100nM. This 25 high potency is comparable to that of other neuropeptides identified as the naturally-occurring ligands of other CNS-specific receptors. Thus, the isolated sequence, termed orphanin FQ herein, represents a novel endogenous peptide ligand for the OFQR.

30 To further characterize ligand binding of this novel ligand to OFQR, a series of peptide analogs were prepared that were tyrosine-substituted at different positions in the peptide sequence. Then cAMP inhibition assays were performed using both the tyrosine-substituted peptides and monoiodotyrosine substituted peptides. Iodinated peptide was synthesized using the chloramine T method of Hunter and Greenwood (1962, *Nature* 194: 495). From these studies it was determined that the peptide analog having tyrosine at position 14 (Y-14: SEQ ID No.: 6) and its monoiodo form (I-Y14) were OFQR agonists of almost equivalent potency to the orphanin FQ 35 peptide, having EC₅₀ values of 1nM and 3nM, respectively.

Fig. 8 shows a comparison of adenyl cyclase inhibitory activity of orphanin FQ (filled circles), Y14 orphanin FQ (filled squares), monoiodo Y14 orphanin FQ (open squares) and Leu-Enkephalin (open circles) in transfected CHO cells heterologously expressing OFQR;

untransfected CHO cells were treated with orphanin FQ peptide as a control (filled triangles). Data were normalized so that cAMP levels in forskolin-stimulated, untransfected CHO cells was equal to 100%. cAMP levels were determined with all incubations being done at least twice in triplicate. Fig. 8 shows the results of a representative experiment.

5 The Y14 peptide was used to quantitatively assay receptor binding affinity for the novel ligand. Membranes from OFQR transfected CHO cells (LC-7) were used at a concentration of 55 μ g membrane protein/assay. Membranes were incubated with increasing concentrations of (125 I)-Y14-orphanin FQ in a final volume of 0.2mL binding buffer (50 mM Hepes, pH 7.4, 10mM NaCl, 1mM MgCl₂, 2.5mM CaCl₂, 0.1% bovine serum albumin, 10 0.025% bacitracin) containing 1mg wheat germ agglutinin-coated SA beads (Amersham). Iodinated peptide was synthesized as above using the chloramine T method. The monoiodinated species was obtained as a single peak, having a specific activity of 2200 Ci/mmmole on the day of synthesis. Assays were performed in 96-well plates (OptiPlate, Caberra Packard), and the mixtures were incubated with shaking for 1h. Bound ligand-associated radioactivity were 15 determined by scintillation proximity (see, for example, Nelson, 1987, *Anal. Biochem.* 165: 287; Bosworth and Towers, 1989, *Nature* 341: 167) using a TopCount microplate scintillation counter (Canberra packard). Concentrations of free ligand were calculated by subtracting the amount of specifically-bound ligand from the total amount of radioligand added.

20 The results of these experiments are shown in Fig. 9. A Scatchard analysis is shown in the inset. The radiolabeled monoiodo-Y14-orphanin FQ peptide displayed saturable and displaceable binding to membranes of OFQR-expressing transfected cells, having a K_D (slope of Scatchard line = $-1/K_D$) of 0.1nM (80 picomole \pm .02 nanomole) and a B_{max} (Scatchard line abscissa intercept) of 800fmol/mg membrane protein. K_D was determined as in Munson, Principles of Pharmacology (Chapman & Hall), 1995, Chapter 1 (which is incorporated by 25 reference). These experiments demonstrate that the novel endogenous ligand peptide, orphanin FQ, and Y14-tyrosine substituted analogs thereof, bind saturably to OFQR with high affinity, further confirming the conclusion that this peptide is a naturally-occurring, endogenous ligand for this receptor. These experiments also indicate that the Y14-substituted peptide can be used as a radioligand to detect and quantify OFQ opioid receptor levels.

30

EXAMPLE 6

Functional Characterization of the Orphanin FQ Peptide

In order to study the physiological activity of the orphanin FQ peptide, the *in vivo* activity of the peptide was investigated on unrestricted animals, and specifically, on nociception in 35 such animals.

For these experiments, the peptide was administered intracerebroventricularly (icv) and intrathecally (it) in mice. Immediately after peptide administration, mice were placed in transparent boxes in groups of three, and behavioral signs recorded. Emphasis was placed on

- 30 -

signs indicative of depressant, stimulant and autonomic effects of the peptide (as described by Irwin, 1968, *Psychopharmacologia* 13: 222). In open-field observation experiments, the peptide at larger doses (e.g. above 1nmole/2 μ L/mouse) was noted to have a profound influence on locomotor activity, as well as a decrease in muscle tone, a loss of righting reflex, and ataxia. A 5 quantitative investigation showed a decrease of both horizontal and vertical locomotion following icv administration of the peptide at concentrations of 0.1-10nmole/2 μ L/mouse. These results are shown in figure 10. The effects of the peptide on horizontal and vertical locomotor activity was measured using Digiscan Animal Activity Monitors (Omnitech, Columbus, OH). The values for horizontal and vertical locomotor activity represent the total number of interruptions of the 10 horizontal and vertical sensors during the first 10 minutes following administration of the peptide. Data are shown as the mean \pm standard error of the mean.

On the other hand, the orphanin FQ peptide showed little analgesic effect in these mice, as recorded in a hot plate test, as shown in Fig. 11. Groups of 6-8 male MORO mice (weight=22g) were administered phosphate buffered saline (open circles) or varying doses of the 15 orphanin peptide. Reaction time represents the time taken for the mice to lick their paws. The hot plate was set to 58°C, and a cutoff time of 60 sec was used. No significant decrease in reaction time was detected, with the exception of mice administered the highest levels of peptide, 10nmole/mouse, icv (about .45 nmole/g) designated by an asterisk. This observation was most likely related to a decrease in locomotor activity and muscle tone. At all doses tested, all animals 20 exhibited normal toe- and tail-pinch reflexes, as shown in Fig. 12. No analgesic effect was observed when orphanin FQ was administered intrathecally at 2.5-10nmole/4 μ L/animal; however, at the highest dose, hind limb paralysis and decrease in locomotor activity was observed. No analgesic effect was observed in any mouse with this peptide at doses that did not produce 25 significant decreases in motor activity and muscle tone. These results indicate that, despite structural homology with analgesia-producing opioid receptor ligands as described in Example 5, the orphanin FQ peptide appears to be pharmacologically distinct and to lack appreciable analgesic activity (which is defined as no significant decrease in reaction time as measured in this Example 6).

EXAMPLE 7**Stress Induced Analgesia Reversed by Naloxone and OFQ**

In spite of the structural homology between OFQ and the opioid peptides, icv injected animals do not display unconfounded antinociception or analgesia, which would be 5 observable as a decreased sensitivity to pain. One report has been made that OFQ instead induces hyperalgesia, a supersensitivity to pain, in both the tail flick and hot-plate assays. Meunier et al., Nature 377,532-535 (1995). Contrary to the expectations of others, the present inventors have found that OFQ does not produce hyperalgesia, but instead potently and dose-dependently antagonizes opiate actions such as analgesia. Hence OFQ is observed to act as an 10 endogenous anti-opioid peptide that is capable of reversing effects of either endogenous or exogenous opioids in an animal, such as a mouse or human.

The anti-opioid effects of the OFQ peptide are demonstrated in this example, which shows the effects of OFQ in the tail-withdrawal (TW) test, an assay of response to acute thermal pain. The results are shown in Fig. 13A. Swiss-Webster mice of both sexes (20-35 g; Simonsen 15 Inc.; N=7-8 per group) were tested for nociceptive sensitivity to immersion of the distal half of the tail in 46°C water before and after (10, 20 and 30 min) subcutaneous (sc) injection with saline (Sal; 10ml/mg) and naloxone (NAL, 1mg/kg) and icv injection (under light halothane anesthesia) with Veh (2.5 microliter of artificial cerebrospinal fluid (CSF)) or 10 nmole OFQ. Separate groups of mice received only sc injections of saline or naloxone (labeled None to indicate no icv 20 injection in Fig. 13A).

The latency to respond to the heat stimulus by a vigorous flexion of the tail was measured to the nearest 0.1 s by an experienced observer blind to the drug condition of the animal. Mice were lightly restrained in a cloth/cardboard holder during testing. In the absence of a withdrawal reflex or other indication of distress, the tail was removed from the water after a 25 cut-off latency of 15 s. To improve accuracy, three separate TW latency determinations, separated by 20 s, were made and averaged at each test. In this and all subsequent experiments, no main effects nor interactions of sex were found to be significant, so data from both sexes were pooled. Because there were no changes in post-injection TW latencies in any group with repeated testing at 10, 20 and 30 min post-injection, data from these time points were pooled. An 30 ANOVA (analysis of variants) performed on averaged TW change data revealed significant main effects of sc injection and icv injection ($F_{2,44}=9.31$, $p<.05$, respectively). Planned contrasts demonstrate that the Sal/Veh group is significantly different from all others.

Although there was no evidence of hyperalgesia produced by OFQ, vehicle-treated animals displayed higher TW latencies (ie were less sensitive to pain) than OFQ-treated animals 35 ($p<.01$)(Fig. 13). The vehicle treated animals were displaying stress-induced-analgesia (SIA). SIA is mediated by endogenous opioids, hence the effect of both the opiate antagonist naloxone and OFQ on the SIA was investigated. Naloxone and OFQ both attenuated the 4 s post-injection increase in TW latency displayed by vehicle-treated animals ($p<.05$). The combined

administration of naloxone and OFQ had no additional effect. In no case did any mouse treated with OFQ or naloxone exhibit hyperalgesia; administration of these drugs instead merely returned pain sensitivity to baseline by reversing SIA. Hyperalgesia would have been detected by a shorter tail withdrawal latency than with vehicle alone, which was not observed with OFQ.

5

EXAMPLE 8

Reversal of morphine analgesia by OFQ in Writhing Test

To maximize the observation of hyperalgesia, the acetic acid AC assay was performed to determine the effect of OFQ on the abdominal constriction test. This test is an 10 extremely sensitive measure of tonic, chemically-induced pain stimulus. The results are given in Fig. 13B.

The AC assay was a modification of the technique originally described by Koster et al. Mice (N=9-15 per group) were acclimated to individual cylindrical Plexiglass observation chambers for > 30 minutes. They were then weighed and given an sc injection of saline (Sal; 10 15 ml/mg) or naloxone (NAL; 1 mg/kg and 10 mg/kg), or handled but given no injection (None). Immediately thereafter, mice were lightly anesthetized with halothane and given an icv injection of vehicle (VEH; 2.5 μ l of artificial CSF) or 2.5 nmole OFQ in 2.5 μ l vehicle (OFQ-2.5). A separate group of Sal-treated mice received no icv injection (None). Between 5 and 10 minutes 20 later, all mice were given an intraperitoneal (ip) injection of 0.9% acetic acid and placed back in their observation chambers. For the next 30 minutes, the number of abdominal constrictions (lengthwise stretches of the torso with a concomitant concave arching of the back) were counted and recorded. Four mice in separate chambers were observed simultaneously by a single experimenter blind to drug condition. ANOVA revealed a significant group differences.

The results are shown in Fig. 13B. The decreased number of constrictions in VEH 25 mice is believed to be due to SIA, which was reversed by NAL and OFQ-2.5. The 2.5 nmole icv dose of OFQ was pharmaceutically sufficient to inhibit the analgesic effect of morphine within 30 minutes after the OFQ was administered.

Example 9

Reversal of Morphine Analgesia by OFQ in Tail Withdrawal Latency

OFQ reverses the analgesic actions of morphine, which are mediated by the opioid SIA substrate. In this example, morphine is administered to mice immediately following 30 assessment of their baseline nociceptive sensitivity. It produced a profound analgesia that peaked at 30 minutes post-injection (Fig. 14) and lasted about 2 hours. But OFQ reversed this analgesia 35 in a dose-dependent manner (at doses of 2.5-25 nmole).

Swiss-Webster mice of both sexes (N=6-11 per group) were tested for baseline nociceptive sensitivity on the 49°C TW test (as described in Example 7), injected s.c. with morphine sulfate (MOR; 5 mg/kg), and then retested 20 min later to establish the existence of

morphine analgesia. Analgesic mice (those with TW latencies of at least double their baseline; all except 3) were then lightly anesthetized with halothane, injected i.c.v. with vehicle (Veh; 2.5 μ l artificial CSF) or OFQ (2.5, 5, 10 or 25 nmole in 2.5 μ l vehicle), and retested for TW latencies 10, 25, 40 and 100 min later by an experimenter blind to OFQ dose. The 25 nmol OFQ dose 5 produced marked atonia and flaccid paralysis in up to 50% of mice. These side-effects should not, however, be considered a confound of the present data because mice injected with this high OFQ dose actually reacted faster to the thermal stimulus.

Fig. 14 shows time-course data demonstrating morphine analgesia and its dose-dependent reversal by four doses of OFQ. Symbols represent mean (\pm SEM) TW latencies from 10 49°C water at each time point.

A linear regression analysis was used to calculate the dose of OFQ causing a 50% inhibition of the effect of 5 mg/kg morphine analgesia, and determined that this ED₅₀ was 7.5 nmol in a 2.5 μ l volume. The ED₅₀ is the dose required to produce a response of 50% of 15 maximal effect. A significant inhibition of analgesia would be a 50% inhibition of opiate analgesia. The ED₅₀ would be a pharmaceutically sufficient amount of OFQ to achieve significant inhibition.

- 34 -

EXAMPLE 10

Attenuation of morphine hypothermia by OFQ

This example demonstrates the ability of OFQ to functionally antagonize opioid effects other than analgesia. The non-nociceptive action of morphine studied here was the ability 5 to induce hypothermia in rodents at high doses. The DBA/2J mice strain was used to study attenuation of hypothermia because of its known robust hypothermic response.

Core body temperature of DBA/2J mice (15-30 g; The Jackson Laboratory; N = 7-8 per group) was assessed immediately before and 30 min after systemic administration of saline (Sal; 10ml/kg) or morphine (MOR; 20 mg/kg, i.p.). Immediately thereafter, mice were injected 10 i.c.v. with vehicle (Veh; 2.5 μ l artificial CSF) or 5 or 10 nmole OFQ in 2.5 μ l vehicle (OFQ-5 and OFQ-10), and retested 15, 30 and 60 min later.

As shown in Fig. 15, there was a 6°C decrease in body temperature produced by MOR, in addition to a transient hypothermia produced by halothane. The body temperature of saline-treated mice remained constant at about 38°C, except for a brief hypothermic episode from 15 exposure to halothane. OFQ itself did not affect body temperature. However, both doses of OFQ significantly reversed 20 mg/kg MOR hypothermia at 60-90 minutes post-MOR injection ($F_{4,33}=44.94, p < .001$). A reversal of hypothermia would be at least a 1°C increase in temperature under these conditions. The dose of OFQ given in this Example is pharmaceutically sufficient to achieve this reversal.

20 The high dose of morphine (20 mg/kg) used in this experiment produced Straub tail, which is a muscular rigidity of the tail accompanying exposure to opiates in virtually all mice. Exposure to doses of at least 5 nmole OFQ effectively eliminated Straub tail in the vast majority of animals within 1-2 minutes following icv injection. The 5 nmole icv dose in the 15-30g rat would be an example of a pharmaceutically sufficient amount to eliminate Straub tail in a 25 majority of animals within 1-2 minutes following icv injection. This finding further generalizes OFQ's functional antagonism of opiate mechanisms.

EXAMPLE 11

Precipitation of Opiate Withdrawal Syndrome

30 in Morphine Dependent Mice

Opiate tolerance, dependence and withdrawal can not be explained by simple alterations in opiate receptor function. Morphine tolerance and addiction is believed to involve action of (until-now) ill-defined anti-opioid activities. Specifically, morphine administration is believed to cause the release of anti-opioid peptides, which in turn may counteract the effects of 35 morphine and contribute to maintenance of physiologic homeostasis. An anti-opioid peptide should precipitate withdrawal symptoms by interfering with this homeostatic state. This Example illustrates that OFQ indeed causes an opiate withdrawal syndrome in morphine dependent mice, further supporting its role as an anti-opioid.

C57BL/6J mice of both sexes (15-25 g; The Jackson Laboratory; N=6-11 per group) were made dependent on morphine using a sustained-release preparation. The preparation was 35 mg/ml morphine sulfate in saline solution, vortexed with a 6:1 light mineral oil; mannide monooleate (Arlacel A, Sigma) mixture in a 1:7 ratio of morphine:oil solution. The 5 viscous white emulsion was administered s.c. at a dose of 200 mg/kg morphine (in an 8 ml/kg volume) using a 21-gauge needle. At the time of testing (5-6 h post-injection) morphine was still present based on Straub tail and 49°C TW latency elevations. Mice were administered s.c. saline (Sal; 10 ml/kg) or NAL (10 mg/kg) followed immediately by i.c.v. vehicle (Veh; 2.5 μ l artificial 10 CSF) or OFQ (0.1-10 nmole in 2.5 μ l vehicle). Since all doses of OFQ used appeared to be equally effective in the precipitation of withdrawal, these data were combined. Each animal was then placed on individual 17 x 27 cm platforms suspended 32 sm above a table top. The number of jumps was recorded for 20 min, as were occurrences of tremor/"wet-dog shakes" and the presence of ptosis. OFQ doses > 5 nmole produced atonia in some morphine-dependent mice that may have hindered jumping behavior.

15 Withdrawal jumping precipitated by OFQ is shown in Fig. 16. Bars represent mean (\pm SEM) number of withdrawal jumps in the 20-min testing period. Ratios above the error bars describe the number of animals exhibiting jumping behavior relative to the total number tested. The withdrawal syndrome produced by i.c.v. OFQ was unequivocal, albeit modest in comparison to that produced by systemic naloxone (NAL). Other withdrawal symptoms, 20 including wet-dog shakes and ptosis, were observed in mice treated with OFQ, but not in mice merely given injections of vehicle. Precipitation of withdrawal shall be defined as increasing total withdrawal jumps in the protocol of Example 11.

Example 12

25 Determination of Affinity of OFQ Receptor Agonists

The peptide of the present invention does not bind with high specificity to μ , κ or δ opioid receptors. This example will demonstrate that lack of receptor specificity, and demonstrate how specificity of ligand-receptor binding can be determined. [3 H]diprenorphine was used as a competitive antagonist to assay receptor binding affinity for OFQ. Membranes from 30 OFQR transfected CHO cells ((LC-7) were used at a concentration of 100 μ g membrane protein/assay.

Stable transfected-CHO cells were rinsed three times with 50 mM Tris-HCl buffer (pH 7.7), harvested and homogenized in the same buffer using a glass-Teflon homogenizer at 4°C. The homogenate was centrifuged at 42,000 g at 4°C for 15 minutes. The pellet was 35 resuspended in 50 mM Tris-HCl buffer and incubated at room temperature for one hour. Following pre-incubation, the homogenate was centrifuged as described above and the pellet was stored at 70°C until used.

The pellet was thawed, washed with 50 mM Tris-HCl buffer and the homogenate was centrifuged as above. The pellet was resuspended in the binding buffer and the protein concentration was adjusted to 100 μ g/ μ l as determined by a standard curve determined by Lowry assay using bovine serum albumin as the standard. In equilibrium competition experiments

5 [3H]diprenorphine was utilized as an antagonist radioligand to label μ , κ and δ receptors, respectively, in CHO cells expressing the μ , κ or δ receptors, and OFQ was used as a displacer. In addition, DAMGO, U50488H and DPDPE were also used as a displacer in the μ , κ and δ receptor expressed CHO cells in these assays.

10 Binding reactions were conducted in an incubation tube with 50 mM Tris-HCl buffer (pH 7.7) in the presence of 10 micromolar peptidase inhibitor bestatin at 25°C for 60 minutes in a total volume of 1 ml using a final [3H]diprenorphine concentration of 0.36 nM. All of the tubes contained the membrane preparation and [3H]diprenorphine. A ligand of interest was also added to each membrane/diprenorphine preparation. A subset of the tubes contained 10 micromolar naloxone as the ligand of interest, to permit determination of non-specific binding.

15 Another subset of the tubes contained OFQ in the presence of the [3H]diprenorphine to determine displacement of OFQ from each class of receptor by the diprenorphine displacer. Another subset of tubes contained DPDPE, while yet another subset contained U50488H as the ligand of interest. Each of the ligands of interest was present in a concentration from zero in order of magnitude up to 100 micromolar. The concentration of radioactive diprenorphine displacer was varied, in

20 different sets of tubes, across the range shown in Fig. 17.

At the end of the 60 minutes period of incubation, the incubation was terminated by vacuum filtration using a Brandel CellHarvester over polyethylenimine (0.5%)-soaked GF/B filters. The Cell Harvester has a manifold that simultaneously emptied out all of the reaction tubes in a controlled manner and deposits the contents on a filter disk that corresponds to each tube, washed 2 times with 2 ml each of ice-cold 50 mM Tris-HCl buffer, and the contents again deposited on the corresponding filter disk. Filter disks were placed in minivials, allowed to elute overnight in EcoLume scintillation fluid (ICN, Costa Mesa, CA), and then counted in a Beckman scintillation counter.

25 The resulting data were radioactive counts, which were proportional to percent total bound diprenorphine radioligand. The percent bound ligand of interest (e.g., OFQ) was inversely proportional to the amount of radioactivity detected. As shown in Fig. 17, the percent of total OFQ at the μ , κ and δ receptors was stable over a broad range of displacer concentrations. The DAMGO, DPDPE and U50488H ligands (known to bind to the μ , κ and δ receptors) were, however, competitively displaced.

30 35 For each set of ligand displacement studies, the results obtained with the scintillation counter were analyzed using an IBM computer system equipped with a GraphPad Inplot (from GraphPad, Inc., of San Diego, CA) software program. This program performs a

- 37 -

nonlinear regression fit of a logistic equation to the data, and produced the graphs shown in Fig. 17.

Using this data, the program further computes IC_{50} (the concentration of unlabeled ligand, such as OFQ) that inhibits 50% of the radioactively labeled ligand binding, such as [3H]diprenorphine. IC_{50} and K_i are related by the equation $K_i = IC_{50}/(1+[L]/K_D)$ where $[L]$ is the concentration of labeled ligand diprenorphine, and K_D is defined as the dissociation constant for [3H]diprenorphine. IC_{50} data are shown in the following Table 1:

TABLE 1

10 Affinity of opioid receptor agonists and OFQ at mu, delta and kappa receptors expressed in CHO cells

Compound	IC_{50}^* (nM)		
	μ -site	δ -site	κ -site
DAMGO	2.0±0.1		
DPDPE		0.59±0.01	
U50488H			1.6±0.2
OrphanFQ	2083±432	2246±339	754±89

20 * IC_{50} values were derived from competition experiments using an opioid antagonist radioligand [3H] diprenorphine (0.36 nM).

20

Data are mean SE values derived from three independent experiments. The ligand of interest has a low IC_{50} (≤ 100 nM) if it binds with high affinity to the receptor. All ligands of interest bound with high affinity to one of the opioid receptors, except for OFQ which had an IC_{50} greater 2000 at μ and δ , and 754 at the κ site. These values show that OFQ does not specifically bind to μ , κ and δ receptors. As used herein, specifically binding to a receptor shall mean having an affinity expressed by an IC_{50} of less than about 100nm. In preferred embodiments, the affinity is 0.1-10nm, preferably less than 1nm, and most preferably less than 0.1nm.

30

Example 13

The affinity of any of the peptides of the present invention can be determined by an affinity study as described in Example 12. Hence, an OFQ analogue with amino acid substitutions can be substituted for the OFQ of Example 12. An OFQR transfected CHO cell membrane would be substituted for the μ , κ and δ receptor expressing cell membranes of Example 12 to determine affinity of the peptide analogue to the OFQ receptor. This provides a

straightforward screening method for determining whether the peptide binds specifically (K_i or $IC_{50} \leq 100\text{nm}$) to the OFQR.

Example 14

5 **Protocol for Obtaining OFQ Anti-Opioid Antagonist and Using the Antagonist**

Antagonism of the OFQ/OFQR system will have substantial clinical utility.

Although the antagonist will not have analgesic actions itself, it is expected to be effective as an adjunct to morphine pharmacotherapy. By interfering with the body's homeostatic anti-opioid mechanism, the OFQR antagonist is expected to allow use of lower therapeutic doses of morphine to achieve greater analgesic effects. Lower doses of morphine have the advantage of producing fewer side-effects (such as respiratory depression or constipation), and will also help avoid development of tolerance or addiction that can be experienced following prolonged use of high doses of morphine. The OFQR antagonist is also expected to lessen the signs and symptoms of opiate withdrawal in addicts. Use of the antagonist in this manner has been made possible by the present inventors' recognition that the OFQ/OFQR interaction mediates an anti-opioid effect.

Although a method of using an anti-opioid antagonist has not been known to the art, oligonucleotides have previously been reported that antagonize the effect of OFQ. See Meunier et al., *Nature* 377:532-535 (1995), which is incorporated by reference. The oligonucleotides reported in that paper were antisense oligonucleotides to translated regions of the OFQR. Antisense oligonucleotide mAS[25,9] was the 17-mer 3'-GGAGAAAGGACGGGGTA-5' complementary to bases 9-25 of the translated region of the mouse ORL₁ mRNA. Control missense oligonucleotide hAS[25,9], was the 17-mer 3'-GGAGAAAGGGCGCGGCA-5' complementary to bases 9-25 of the translated region of the human ORL₁ mRNA. The oligonucleotides were dissolved in sterile physiological saline at the final concentration of 2 mg/ml. Male Swiss CD1 mice (20-25g; Charles River) were manually injected with 10 μl of solution directly into the lateral brain ventricle every day for 4 consecutive days.

Expression of the receptor was inhibited by the antisense oligonucleotides. Other oligonucleotides of similar length could be mapped to bases 9-25 (using the genetic code of Table 1) to provide similar antagonistic activity.

Other antagonists are used in the present method to treat an opiate withdrawal syndrome. A heptadecapeptide is synthesized with a sequence that is substantially homologous to the OFQ amino acid sequence, but with one or more amino acid substitutions made. The synthesized heptadecapeptide (or other oligopeptide) is then screened using the cAMP accumulation assay in OFQ transfected CHO cells. Peptides that do not inhibit cAMP accumulation (using the assay described in Example 5) are then selected as a partial agonist that acts as an antagonist. The selected peptides obtained by this straightforward assay are then administered to animals (as in Example 11) to demonstrate attenuation of opiate withdrawal, as

evidenced by reduction in withdrawal jumping compared to animals not receiving the OFQ antagonist.

Synthetic peptide combinatorial libraries may also be screened to find an agonist-antagonist peptide that does not inhibit cAMP accumulation in the disclosed assay. Screening is 5 performed using the techniques described in Houghten et al., Generation and use of synthetic peptide combinatorial libraries for basic research and drug discovery, *Nature* 354:84-86, incorporated by reference. The soluble, nonsupport bound peptide libraries appear useful in virtually all *in vitro* and *in vivo* assays, as reported by Ostresh, "Libraries from libraries: Chemical transformation of combinatorial libraries to extend the range and repertoire of chemical 10 diversity," *PNAS*. The successful use of these libraries has been reported for the development of receptor-active opioid peptides. See Dooley, "An All D-Amino Opioid Peptide with Central Analgesic Activity from a Combinatorial Library," *Science* 266:2019-2022, 1994, which is incorporated by reference.

Antagonists discovered by these techniques are tested for interference with opiate 15 withdrawal signs, as in Example 8.

It should be understood that the foregoing disclosure emphasizes certain specific embodiments of the invention and that all modifications or alternatives equivalent thereto are within the spirit and scope of the invention as set forth in the appended claims.

- 40 -

SEQUENCE LISTING

(1) GENERAL INFORMATION

(i) APPLICANTS: Grisel, Judith E.
5 Grandy, David K.
Mogil, Jeffrey S.
Bunzow, James R.
Civelli, Olivier
Reinscheid, Rainer Klaus
Nothacker, Hans-Peter
10 Monsma, Frederick James

(ii) TITLE OF INVENTION: Opioid Antagonists and Methods of Their Use

(iii) NUMBER OF SEQUENCES: 10

(iv) CORRESPONDENCE ADDRESS:
15 (A) ADDRESSEE: Klarquist Sparkman Campbell Leigh
& Whinston LLP

(B) STREET: 121 S.W. Salmon, Suite 1600

(C) CITY: Portland

(D) STATE: Oregon

20 (E) COUNTRY: USA

(F) ZIP: 97204

(v) COMPUTER READABLE FORM:
25 (A) MEDIUM TYPE: Floppy Disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version WP5.1 ASCII text

(vi) CURRENT APPLICATION DATA:
3.0 (A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

3.0 (vii) PRIOR APPLICATION DATA:
35 (A) APPLICATION NUMBER: 08/533,058
(B) FILING DATE: 3 November 1995
(C) APPLICATION NUMBER: 08/514,541
(D) FILING DATE: 11 August 1995

(viii) ATTORNEY/AGENT INFORMATION
40 (A) NAME: Richard J. Polley, Esq.
(B) REGISTRATION NUMBER: 28,107
(C) REFERENCE/DOCKET NUMBER: 899-45532/RJP

(ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (503) 226-7391
(B) TELEFAX: (503) 228-9446

- 41 -

(2) INFORMATION FOR SEQ ID NO: 1

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

5 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

(A) DESCRIPTION: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10 ATGAATTCAC NRTSATGAGY GTSGACHGHT A 31

(3) INFORMATION FOR SEQ ID NO: 2

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

15 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

(A) DESCRIPTION: DNA (genomic)

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TTGTCGACRT ARRAGRAYNG GRTT 24

(4) INFORMATION FOR SEQ ID NO: 3

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 1452 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

30 (A) DESCRIPTION: cDNA

(ix) FEATURE:

(A) NAME/KEY: 5'UTR

(B) LOCATION: 1..181

(ix) FEATURE:

35 (A) NAME/KEY: CDS

(B) LOCATION: 182..1282

(ix) FEATURE:

(A) NAME/KEY: 3'UTR

-42-

(B) LOCATION: 1283..1452

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCGAGGAGCC ATTCCCAGCC GCAGCAGACC CCAATCTAGA GTGAGAGTCA TTGCTCAGTC 60
 CACTGTGCTC CTGCCTGCC GCCTTCTGC TAAGCATTGG GGTCTATTT GCGCCAGCT120
 5 TCTGAAGAGG CTGTGTGTGC CGTTGGAGGA ACTGTACTGA GTGGCTTGC AGGGTGACAG180
 C ATG GAG TCC CTC TTT CCT GCT CCA TAC TGG GAG GTC TTG CAT GGC 226
 Met Glu Ser Leu Phe Pro Ala Pro Tyr Trp Glu Val Leu His Gly
 1 5 10 15
 AGC CAC TTT CAA GGG AAC CTG TCC CTC CTA AAT GAG ACC GTA CCC CAC 274
 10 Ser His Phe Gln Gly Asn Leu Ser Leu Leu Asn Glu Thr Val Pro His
 20 25 30
 CAC CTG CTC CTC AAT GCT AGT CAC AGC GCC TTC CTG CCC CTT GGA CTC 322
 His Leu Leu Leu Asn Ala Ser His Ser Ala Phe Leu Pro Leu Gly Leu
 35 40 45
 15 AAG GTC ACC ATC GTG GGG CTC ATC TTG GCT GTG TGC ATC GGG GGG CTC 370
 Lys Val Thr Ile Val Gly Leu Ile Leu Ala Val Cys Ile Gly Gly Leu
 50 55 60
 CTG GGG AAC TGC CTC GTC ATG TAT GTC ATC CTC AGG ACA CCC AAG ATG 418
 20 Leu Gly Asn Cys Leu Val Met Tyr Val Ile Leu Arg Thr Pro Lys Met
 65 70 75
 AAG ACA GCT ACC AAC ATT TAC ATA TTT AAT CTG GCA CTG GCT GAT ACC 466
 Lys Thr Ala Thr Asn Ile Tyr Ile Phe Asn Leu Ala Leu Ala Asp Thr
 80 85 90 95
 25 CTG GTC TTG CTA ACA CTG CCC TTC CAG GGC ACA GAC ATC CTA CTG GGC 514
 Leu Val Leu Leu Thr Leu Pro Phe Gln Gly Thr Asp Ile Leu Leu Gly
 100 105 110
 TTC TGG CCA TTT GGG AAA GCA CTC TGC AAG ACT GTC ATT GCT ATC GAC 562
 Phe Trp Pro Phe Gly Lys Ala Leu Cys Lys Thr Val Ile Ala Ile Asp
 115 120 125
 30 TAC TAC AAC ATG TTT ACC AGC ACT TTT ACT CTG ACC GCC ATG AGC GTA 610
 Tyr Tyr Asn Met Phe Thr Ser Thr Phe Thr Leu Thr Ala Met Ser Val
 130 135 140
 GAC CGC TAT GTG GCT ATC TGC CAC CCT ATC CGT GCC CTT GAT GTT CGG 658
 35 Asp Arg Tyr Val Ala Ile Cys His Pro Ile Arg Ala Leu Asp Val Arg
 145 150 155
 ACA TCC AGC AAA GCC CAG GCT GTT AAT GTG GCC ATA TGG GCC CTG GCT 706
 Thr Ser Ser Lys Ala Gln Ala Val Asn Val Ala Ile Trp Ala Leu Ala
 160 165 170 175
 TCA GTG GTT GGT GTT CCT GTT GCC ATC ATG GGT TCA GCA CAA GTG GAA 754
 40 Ser Val Val Gly Val Pro Val Ala Ile Met Gly Ser Ala Gln Val Glu
 180 185 190
 GAT GAA GAG ATC GAG TGC CTG GTG GAG ATC CCT GCC CCT CAG GAC TAT 802
 Asp Glu Glu Ile Glu Cys Leu Val Glu Ile Pro Ala Pro Gln Asp Tyr
 195 200 205
 45 TGG GGC CCT GTA TTC GCC ATC TGC ATC TTC CTT TTT TCC TTC ATC ATC 850
 Trp Gly Pro Val Phe Ala Ile Cys Ile Phe Leu Phe Ser Phe Ile Ile
 210 215 220
 CCT GTG CTG ATC ATC TCT GTC TGC TAC AGC CTC ATG ATT CGA CGA CTT 898
 50 Pro Val Leu Ile Ile Ser Val Cys Tyr Ser Leu Met Ile Arg Arg Leu
 225 230 235
 CGT GGT GTC CGT CTG CTT TCA GGC TCC CGG GAG AAG GAC CGA AAC CTG 946
 Arg Gly Val Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu
 240 245 250 255
 55 CGG CGT ATC ACT CGA CTG GTG CTG GTA GTG GTG GCT GTG TTT GTG GGC 994
 Arg Arg Ile Thr Arg Leu Val Leu Val Val Ala Val Phe Val Gly
 260 265 270

-43-

	TGC TGG ACG CCT GTG CAG GTG TTT GTC CTG GTT CAA GGA CTG GGT GTT	1042		
	Cys Trp Thr Pro Val Gln Val Phe Val Leu Val Gln Gly Leu Gly Val			
	275	280	285	
5	CAG CCA GGT AGT GAG ACT GCA GTT GCC ATC CTG CGC TTC TGC ACA GCC	1090		
	Gln Pro Gly Ser Glu Thr Ala Val Ala Ile Leu Arg Phe Cys Thr Ala			
	290	295	300	
	CTG GGC TAT GTC AAC AGT TGT CTC AAT CCC ATT CTC TAT GCT TTC CTG	1138		
	Leu Gly Tyr Val Asn Ser Cys Leu Asn Pro Ile Leu Tyr Ala Phe Leu			
	305	310	315	
10	GAT GAG AAC TTC AAG GCC TGC TTT AGA AAG TTC TGC TGT GCT TCA TCC	1186		
	Asp Glu Asn Phe Lys Ala Cys Phe Arg Lys Phe Cys Cys Ala Ser Ser			
	320	325	330	335
15	CTG CAC CGG GAG ATG CAG GTT TCT GAT CGT GTG CGG ACG ATT GCC AAG	1234		
	Leu His Arg Glu Met Gln Val Ser Asp Arg Val Arg Thr Ile Ala Lys			
	340	345	350	
	GAT GTT GGC CTT GGT TGC AAG ACT TCT GAG ACA GTA CCA CGG CCA GCA	1282		
	Asp Val Gly Leu Gly Cys Lys Thr Ser Glu Thr Val Pro Arg Pro Ala			
	355	360	365	
20	TGACTAGGCG TGGACCTGCC CATGGTGCCT GTCAGCCCAC AGAGCCCATC CTACACCCAA	342		
	CACGGAGCTC ACACAGGTCA CTGCTCTCTA GGTTGACCCCT GAACCTTGAG CATCTGGAG	402		
	CTTGAATGGC TTTCTTTTG GATCAGGATG CTCAGTCCTA GAGGAAGACC			
		1452		

(5) INFORMATION FOR SEQ ID NO: 4

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 367 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

(A) DESCRIPTION: protein

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Glu	Ser	Leu	Phe	Pro	Ala	Pro	Tyr	Trp	Glu	Val	Leu	His	Gly	Ser	
1										10					15	
His	Phe	Gln	Gly	Asn	Leu	Ser	Leu	Leu	Asn	Glu	Thr	Val	Pro	His	His	
														20	25	30
35	Leu	Leu	Leu	Asn	Ala	Ser	His	Ser	Ala	Phe	Leu	Pro	Leu	Gly	Leu	Lys
														35	40	45
	Val	Thr	Ile	Val	Gly	Leu	Ile	Leu	Ala	Val	Cys	Ile	Gly	Gly	Leu	Leu
														50	55	60
40	Gly	Asn	Cys	Leu	Val	Met	Tyr	Val	Ile	Leu	Arg	Thr	Pro	Lys	Met	Lys
	65					70					75				80	
	Thr	Ala	Thr	Asn	Ile	Tyr	Ile	Phe	Asn	Leu	Ala	Leu	Asp	Thr	Leu	
														85	90	95

-44-

Val Leu Leu Thr Leu Pro Phe Gln Gly Thr Asp Ile Leu Leu Gly Phe
 100 105 110
 Trp Pro Phe Gly Lys Ala Leu Cys Lys Thr Val Ile Ala Ile Asp Tyr
 115 120 125
 5 Tyr Asn Met Phe Thr Ser Thr Phe Thr Leu Thr Ala Met Ser Val Asp
 130 135 140
 Arg Tyr Val Ala Ile Cys His Pro Ile Arg Ala Leu Asp Val Arg Thr
 145 150 155 160
 10 Ser Ser Lys Ala Gln Ala Val Asn Val Ala Ile Trp Ala Leu Ala Ser
 165 170 175
 Val Val Gly Val Pro Val Ala Ile Met Gly Ser Ala Gln Val Glu Asp
 180 185 190
 Glu Glu Ile Glu Cys Leu Val Glu Ile Pro Ala Pro Gln Asp Tyr Trp
 195 200 205
 15 Gly Pro Val Phe Ala Ile Cys Ile Phe Leu Phe Ser Phe Ile Ile Pro
 210 215 220
 Val Leu Ile Ile Ser Val Cys Tyr Ser Leu Met Ile Arg Arg Leu Arg
 225 230 235 240
 20 Gly Val Arg Leu Leu Ser Gly Ser Arg Glu Lys Asp Arg Asn Leu Arg
 245 250 255
 Arg Ile Thr Arg Leu Val Leu Val Val Ala Val Phe Val Gly Cys
 260 265 270
 Trp Thr Pro Val Gln Val Phe Val Leu Val Gln Gly Leu Gly Val Gln
 275 280 285
 25 Pro Gly Ser Glu Thr Ala Val Ala Ile Leu Arg Phe Cys Thr Ala Leu
 290 295 300
 Gly Tyr Val Asn Ser Cys Leu Asn Pro Ile Leu Tyr Ala Phe Leu Asp
 305 310 315 320
 30 Glu Asn Phe Lys Ala Cys Phe Arg Lys Phe Cys Cys Ala Ser Ser Leu
 325 330 335
 His Arg Glu Met Gln Val Ser Asp Arg Val Arg Thr Ile Ala Lys Asp
 340 345 350
 Val Gly Leu Gly Cys Lys Thr Ser Glu Thr Val Pro Arg Pro Ala
 355 360 365
 35

(6) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 17 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

(A) DESCRIPTION: peptide

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Phe Gly Gly Phe Thr Gly Ala Arg Lys Ser Ala Arg Lys Leu Ala Asn
 1 5 10 15
 Gln

50 (7) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

-45-

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE:

5 (A) DESCRIPTION: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Phe Gly Gly Phe Thr Gly Ala Arg Lys Ser Ala Arg Lys Tyr Ala Asn
1 5 10 15
Gln

10

(8) INFORMATION FOR SEQ ID NO. 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids

- (B) TYPE: amino acid

15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO. 7

Tyr Gly Gly Phe Leu Arg Arg Ile Arg Pro Lys Leu Lys Trp Asp Asn Gln
1 5 10 15

20

(9) INFORMATION FOR SEQ ID NO. 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids

25 (B) TYPE: amino acid

- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO. 8

30

Tyr Gly Gly Phe Met Thr Ser Glu Lys Ser Gln Thr Pro Leu Val Thr
1 5 10 15

(10) INFORMATION FOR SEQ ID NO: 9

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids

- (B) TYPE: amino acid

- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO. 9

Tyr Gly Gly Phe Leu Arg Arg Gln Phe Lys Val Val Thr
1 5 10

-46-

(11) INFORMATION FOR SEQ ID NO: 10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO. 10

10 Tyr Gly Gly Phe Leu
1 5

What we claim is:

1. A method of antagonizing a physiological effect of an opioid in an animal, comprising administering to the animal a pharmaceutically effective amount of a peptide that has an activity of binding to a mammalian OFQ receptor with high specificity.
5
2. The method of claim 1 wherein the peptide has an FGGF motif.
3. The method of claim 1 wherein the peptide does not bind to mu, delta, or kappa opioid receptors with high specificity.
10
4. The method of claim 1 wherein the peptide antagonizes opioid analgesia without increasing nociceptive sensitivity.
5. The method of claim 1 wherein the mammalian OFQ receptor is encoded by the DNA molecule set forth in SEQ. ID. No. 3.
6. A method of antagonizing physiologic effects of an opioid in an animal, comprising the step of administering to the animal an amount of an anti-opioid peptide sufficient to reduce a physiological effect of the opioid, wherein the peptide:
15
(a) has an FGGF motif;
(b) binds with high specificity to a receptor encoded by SEQ. ID No. 3;
20
(c) does not bind to mu, delta or kappa opioid receptors with high specificity; and
(d) antagonizes opioid analgesia without increasing nociceptive sensitivity.
25
7. The method of claim 6, wherein the peptide antagonizes opiate induced hypothermia in animals.
8. The method of claim 6, wherein the peptide is administered to the animal to antagonize morphine induced analgesia.
9. The method of claim 6 wherein the FGGF motif is an amino-terminal motif.
30
10. The method of claim 6, wherein the peptide is a heptadecapeptide.
11. The method of claim 1, wherein the peptide is of the formula:
35 (Xaa)_n-Phe-Gly-Gly-Phe-(A¹)-(A²)-A³)-(A⁴)-(A⁵)-(A⁶)-(A⁷)-(A⁸)-(A⁹)-(A¹⁰)-(A¹¹)-(A¹²)-Gin-(Xaa)_m
wherein A¹ is Thr, Leu or Met;
A² is Gly, Arg or Thr;
A³ is Ala, Arg or Ser;
40 A⁴ is Arg, Ile, Glu or Gln;
A⁵ is Lys, Arg or Phe;
A⁶ is Ser, Pro or Lys;
A⁷ is Ala, Lys, Gln or Val;
A⁸ is Arg, Leu, Thr or Val;
45 A⁹ is Lys, Pro or Thr;
A¹⁰ is Tyr, Leu or Trp;

A¹¹ is Ala, Asp or Val;

A¹² is Asn, or Thr;

and

(Xaa) is any amino acid;

n and m are integers wherein n + m is no more than

5 82; and

wherein the amino acids are each individually in either the D or L stereochemical configuration.

10 12. The method of claim 11 wherein the peptide is selected from the group consisting of amino acid sequences selected from the group consisting of Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln and Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Tyr-Ala-Asn-Gln.

15 13. The method of claim 1, wherein the peptide is an analogue, derivative or mimetic of the peptide of claim 11.

15 14. The method of claim 6, further comprising administering to the animal a second anti-opioid antagonist, wherein the second anti-opioid antagonist and anti-opioid peptide are administered in a sufficient amount in combination to antagonize physiological effects of the opioid.

20 15. A method of antagonizing a physiologic effect of morphine on an animal, comprising the step of administering to the animal, in a sufficient amount to diminish the physiologic effect, a peptide having an amino acid sequence consisting of Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln.

25 16. A pharmaceutical dosage form of the peptide of claim 1.

17. The pharmaceutical dosage form of claim 16, comprising the peptide and a pharmaceutical carrier.

30 18. The pharmaceutical dosage form of claim 16, further comprising instructions for administering the pharmaceutical dosage form to an animal to diminish opioid intoxication.

35 19. A method of antagonizing opioid effects caused by exogenous administration of an opioid in an animal, comprising administering to the animal a pharmaceutically effective amount of an anti-opioid peptide sufficient to reduce analgesia induced by the exogenous opioid, wherein the anti-opioid peptide:

(a) is a heptadecapeptide having an aminoterminal FGGF motif;

40 (b) binds to a receptor encoded by SEQ. ID. No. 3 with an IC₅₀ of 100nm or less;

(c) does not bind to mu, delta or kappa opioid receptors with an IC₅₀ of 100nm or less;

(d) antagonizes opioid analgesia without increasing nociceptive sensitivity; and

-49-

(e) antagonizes morphine induced hypothermia in animals when given at a pharmaceutically effective dose to inhibit a morphine induced hypothermic response.

20. The method of claim 19 wherein the exogenous opioid is
5 morphine.

21. A peptide of formula:

$(Xaa)_n\text{-Phe-Gly-Gly-Phe-}(A^1)\text{-}(A^2)\text{-}(A^3)\text{-}(A^4)\text{-}(A^5)\text{-}(A^6)\text{-}(A^7)\text{-}(A^8)\text{-}(A^9)\text{-}(A^{10})\text{-}(A^{11})\text{-}(A^{12})\text{-Gln-}(Xaa)_m$

10 wherein A^1 is Thr, Leu or Met;

A^2 is Gly, Arg or Thr;

A^3 is Ala, Arg or Ser;

A^4 is Arg, Ile, Glu or Gln;

A^5 is Lys, Arg or Phe;

A^6 is Ser, Pro Lys;

15 A^7 is Ala, Lys, Gln or Val;

A^8 is Arg, Leu, Thr or Val;

A^9 is Lys, Pro or Thr;

A^{10} is Tyr, Leu or Trp;

20 A^{11} is Ala, Asp or Val;

A^{12} is Asn, or Thr;

and (Xaa) is any amino acid;

25 n and m are integers wherein $n + m$ is no more than 82; and wherein the amino acids are each individually in either the D or L stereochemical configuration and wherein the peptide specifically binds to a mammalian opioid receptor having an amino acid sequence identified by Seq ID No.:4.

22. A peptide according to claim 21 having the formula Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln, 30 wherein $n + m$ equals zero.

23. A peptide according to claim 21 having the formula Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Tyr-Ala-Asn-Gln, wherein $n + m$ equals zero.

24. The peptide of claim 23 that is radiolabeled with a 35 radioisotope of iodine.

25. A peptide according to claim 21 that is radiolabeled with a radioactive isotope.

26. A pharmaceutical composition comprising a peptide according to claims 21, 22, 23, 24 or 25 and a pharmaceutically-40 acceptable carrier or diluent.

27. A pharmaceutical composition of claim 26 that is a diagnostic agent.

28. A pharmaceutical composition of claim 26 that is a therapeutic agent.

45 29. A method of diagnosing a locomotor disease in an animal, comprising the step of administering a diagnostically

effective amount of the pharmaceutical composition of claim 27 to the animal.

30. A method of treating locomotor disease in an animal, comprising the step of administering a therapeutically effective 5 amount of the pharmaceutical composition of claim 28 to the animal.

31. A method of designing an MSOR opioid receptor ligant comprising the step of preparing an analogue, derivative or mimetic of the peptide of claim 21.

32. An analogue, derivative or mimetic of an MSOR opioid 10 receptor prepared by the method of claim 31.

33. A method of screening a compound for a capacity to bind to a mammalian MSOR opioid receptor in cells expressing the receptor, the method comprising the steps of:

15 (a) transforming a culture of eukaryotic or prokaryotic cells with a recombinant expression construct capable of expressing a mammalian MSOR opioid receptor having an amino acid sequence identified as SEQ ID No.:4, wherein the cells of the transformed cell culture express the opioid receptor;

20 (b) assaying the transformed cell culture for binding of an amount of a detectably-labeled peptide according to claim 21 in competition with varying amounts of the compound; and

25 (c) determining whether the compound competitively binds to the opioid receptor by calculating the extend of inhibition of binding of the detectably-labeled peptide in the presence of the compound.

34. A method of claim 33 comprising the additional step of:

30 (d) comparing the binding capacity of the compound with the binding capacities of additional compounds that are known to bind to mammalian opioid receptors, wherein said additional compounds comprise naturally-occurring and synthetic opioid receptor agonists and antagonists.

CCGAGGAGCCATTCCCAGCCGAGCAGACCCAACTAGAGTGAGAGTCATTGCTCAGTCCACTGTGCTCC 71
 TGCCTGCCGCCCTTCTGCTAAGCATGGGTCTATTTGCCGCCAGCTCTGAGAGGGCTGTGTGCCG 142
 TTGGAGGAACITGTACTGAGTGGCTTGCACGGTGACACCATGGAGTCCCTTCTGCTCCATACTGGAG 214
 M E S L F P A P Y W E
 CTCTTGATGGCAGCCACTTCAAGGGAACTCTCCCTCTAAATGAGACCGTACCCACCCACCTGCTCC 286
 V L Y G S H F Q G N L S L L N E T V P H H L L
 AATGCTAGTCACAGGCCCTCTGGACTCAAGGTACCATCTGAGGGCTACTTGGCTGTGTC 358
 N A S H S A F L P L G L K V T I V G L Y L A V C
 ATGGGGGGCTCTGGGAACCTGCTCGTCATGTATGTCATCTCAGGCACACCAAGATGAAGACAGCTACC 430
 I G G L L G N C L V M Y V I L R H T K M K T A T
 AACATTTACATATTAAATCTGGCACTGGCTGATACCTGGCTTGCTAACACTGCCCTCCAGGGCACACAC 502
 N I Y I F N L A L A D T L V L L T L P F Q G T D
 II ATCTACTGGCTCTGGCAATTGGGAATGCACTCTGCAAGACTGTCATGCTATCGACTACTAACATG 574
 I L L G F W P F G N A L C K T V I A D Y Y N M
 III TTTACCAAGCACTTTACTCTGACCCCATGAGCGTAGACCCCTATGTCATGCTATGCCACCCATCGCTGCC 646
 F T S T F T L T A M S V D R Y V A I C H P I R A
 CTTGATGTTGGACATCCAGCAAAGCCCAGCTGTTAATGTCATGCTATGGCCCTGGCTTCAGTGGTTGG 718
 L D P V R T S S K A Q A V N V A I W A L A S V V G
 * IV GTCCTGTTGCCATATGGGTCAGCACAAGTGGAAAGATGAAGAGATGAGTCAGTGGCTGGAGATCCCTGCC 790
 V P V A I M G S A Q V E D E E I E C L V E I P A
 CCTCAGGACTATTGGGCCCTGTATTGCACTGCACTTCCTTTCATCATCCCTGTGCTGATC 862
 P Q D Y W G P V F A I C I F L F S F I I P V L I
 V ATCTCTGCTCTGCTACAGCTCATGATTCGACCTGGACTCTGCTGCTCCGTGCTTCAGGCTCCGGAGAAG 934
 I S V C Y S L M I R R L R G V R L L S G S R E K
 GACCAAACCTGCCGGTATCACTGGACTGGTGCTGGTACTGGTGCTGCTGTTGGGGTGCTGGACGCC 1006
 D R N L R R I T R L V L V V A V F V G C W T P
 VI GTGCAGGTGTTGCTCTGGCTCAAGGACTGGGTGTTCAACCCAGGTAGTGAGACTGCACTGCCATCTGCC 1078
 V Q V F V L V Q G L G V Q P G S E T A V A I L R
 VII TCTCACAGCCCTGGCTATGCAACAGTGTCTCAATCCCATCTCTATGCTTCCGATGAGAACCTC 1150
 F C T A L G Y V N S C L N P I L Y A F L D E N F
 AAGGCCCTGCTTGAAGAAGTCTGCTGCTCATCCCTCCACGGGAGATGCAAGGTTCTGATGCTGCC 1222
 K A C F R K F C C A S S L H R E M Q V S D R V R
 *
 CGGATTGCCAAGGATGTTGCCCTGGTTGCAAGACTCTGAGACAGTACCAACGGCCACCATGACTAGGGCTG 1294
 S I A K D V G L G C K T S E T V P R P A
 GACCTGCCCATGGGCCCTGTCACCCACAGAGCCCATCTACACCCAAACACGGAGCTACACAGGTCACTGC 1366
 TCTCTAGGTGACCCCTGAACCTTGAGCATCTGGAGCCCTGAATGCTTTGGATCAGGATGCTCAGT 1438
 CCTAGAGGAAGACC

FIG. 1

Amino acid alignment

LC132

Rat μ -Opioid Receptor
 Mouse δ -Opioid Receptor
 Mouse κ -Opioid Receptor

MESLFPAPYWEVL
 MDSSTGPGNTSDCSDPLAQASCSPAPGSWLNL
 MELVPSARAELQSS
 MESPIQIFRGDPGPTCSPSACLLP

I

LC132 YGSHFQGNLSSLNETVPHLLLNAHSASFPLGLKVTIVGLYLAVCIGGLLNCL
 μ -OR HVDGNQSDPCGLNRTGLGGNDSLCPOTGSPSMVTAITIMALYSIVCVVGLFGNFL
 δ -OR PLVNLSDAFPSAFPSAGANASGSPGARSASSLALAITALYSAVCAVGLLGNVL
 κ -OR NSSSWFPNWAESDSNGSFGSEDQQLESAHISPAIPVIITAVYSVVFVVGVLGNSL

II

LC132 VMYVILRHTKMKTATNIIYIFNLALADTLVLLTLPFQGTDIILGFWPFGNALCKTV
 μ -OR VMYVIVRYTKMKTATNIIYIFNLALADALATSFLPFQSVNYLMGTWPFGTILCKIV
 δ -OR VMFGIVRYTKLKTATNIIYIFNLALADALATSFLPFQSAKYLMEWPFGELLCRAV
 κ -OR VMFVIIKYTKMKTATNIIYIFNLALADALVTTMPFQSAVYLMNSWPPGDVLCKIV

III

LC132 IAIDYYNMFTSTFTLAMSVDRYIAVCHPIRALDVRTSSKAQAVNVAIWALASVV
 μ -OR ISIDYYNMFTSIFTLCMTMSVDRYIAVCHPVKAALDFRTPRNAKIVNVCNWILSSAI
 δ -OR LSIDYYNMFTSIFTLTMMMSVDRYIAVCHPVKAALDFRTPAKAKIINICIWVLASGV
 κ -OR ISIDYYNMFTSIFTLTMMMSVDRYIAVCHPVKAALDFRTPLKAKIINICIWLLASSV

IV

LC132 GVPVAIMGSAQ VEDEBIECLVEIAP QDYWGPVFAICIFLFSIIIPVLIISV
 μ -OR GLPVFMATTK YRQGSIDCTLTFSHP TWYWENLLKICVFIFAFIMPILITV
 δ -OR GVPIMVMATQ PRDFAVVCMLQFPSP SWIWDITKICVFLFAFPVPIIITV
 κ -OR GISAIVLGGTKVREDVDVIECSLQFPDDEYSWWDFLHKICVFIFAFVIPVLIIV

V

LC132 CYSILMIRRLRGVRLLSGSREKDRNLLRRITRLVAVFVGCTPVQVFVLFVQGL
 μ -OR CYGLMILRLKSVRMLSGSKEKDRNLLRRITRMLVAVFIVCWTPIHIYVIKAL
 δ -OR CYGLMILRLRSVRLLSGSKEKDRSLLRITRMLVAVFIVCWTPIHIYVIKAL
 κ -OR CYTLMILRLKSVRLLSGSREKDRNLLRRITKLVAVFIVCWTPIHIYVIKAL

VI

LC132 GVQPGSETAVAIL RFCATLGYVNSCLNPILYAFDENFKACFRKFCASSLHRE
 μ -OR ITIPETTFQTVSW HFCIALGYTNSCLNPILYAFDENFKRCREFCIPTSSTIE
 δ -OR VDINRRDPLVVAALHLCIALGYANSCLNPILYAFDENFKRCFRQLCRTPCGRQE
 κ -OR GSTSHSTAALSSY YFCIALGYTNSCLNPILYAFDENFKRCFRDFCFPIKMRME


VII

LC132 MQVSDRVRSIAKDVGGLGCKTSETVPRPA 367
 μ -OR QQNSTRVRQNTREHPSTANTVDRTNHQLENLEAETAPLP 398
 δ -OR PGSLRPRQATTTRERVACTPSDGGGGAAA 372
 κ -OR RQSTNRVRNTVQDPASMRDVGGMNKPV 380

FIG. 2

3/12


FIG. 3



SUBSTITUTE SHEET (RULE 26)

FIG. 4A

3H-METHADONE BINDING TO COS-7 MEMBRANES

**FIG. 4B**

3H-METHADONE BINDING TO LC132 IN COS-7 MEMBRANES




FIG. 5





FIG. 6







FIG. 8




FIG. 9






FIG. 10A**FIG. 10B**

SUBSTITUTE SHEET (RULE 26)

FIG. 11





FIG. 12



10/12

FIG. 13A





FIG. 13B



11/12

FIG. 15





FIG. 16



12/12

FIG. 17A




FIG. 17B





FIG. 17C



INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/13305A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C07K14/665 A61K38/17 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	FEBS LETT, JUN 27 1994, 347 (2-3) P284-8, NETHERLANDS, XP002020226 BUNZOW JR ET AL: "Molecular cloning and tissue distribution of a putative member of the rat opioid receptor gene family that is not a mu, delta or kappa opioid receptor type." see the whole document --- WO,A,95 07983 (INDIANA UNIVERSITY FOUNDATION) 23 March 1995 see page 12, paragraph 2 - page 19, paragraph 2; figure SEQ.ID.16 ---	16-18, 31-34
Y	WO,A,95 07983 (INDIANA UNIVERSITY FOUNDATION) 23 March 1995 see page 12, paragraph 2 - page 19, paragraph 2; figure SEQ.ID.16 ---	16-18, 31-34

 Further documents are listed in the continuation of box C. Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

3 Date of the actual completion of the international search

4 December 1996

Date of mailing of the international search report

17.12.96

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/13305

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	INT J PEPT PROTEIN RES, NOV 1993, 42 (5) P411-9, DENMARK, XP002020227 KAWASAKI AM ET AL: "Syntheses, opioid binding affinities, and potencies of dynorphin A analogues substituted in positions, 1, 6, 7, 8 and 10." see the whole document ---	16-18, 31-34
A	J MED CHEM, JUL 1990, 33 (7) P1874-9, UNITED STATES, XP002020228 KAWASAKI AM ET AL: "Design and synthesis of highly potent and selective cyclic dynorphin A analogues." see the whole document ---	16-18, 31-34
A	25TH INTERNATIONAL NARCOTICS RESEARCH CONFERENCE: OPIOIDS AND THEIR RECEPTORS: CLONING AND BEYOND, NORTH FALMOUTH, MASSACHUSETTS, USA, JULY 16-21, 1994. REGULATORY PEPTIDES, 54 (1). 1994. 123-124., XP000610712 HRUBY V J ET AL: "Design of peptides and peptidomimetics for delta and kappa opioid receptor subtypes" see the whole document ---	16-18, 31-34
A	WO,A,91 00736 (ARIZONA TECH DEV) 24 January 1991 see claim 1 ---	16-18, 31-34
P,X	SCIENCE, NOV 3 1995, 270 (5237) P792-4, UNITED STATES, XP002020230 REINSCHEID RK ET AL: "Orphanin FQ: a neuropeptide that activates an opioidlike G protein-coupled receptor." see the whole document ---	21-28,32
P,X	BIOCHEM BIOPHYS RES COMMUN, DEC 14 1995, 217 (2) P539-45, UNITED STATES, XP002020239 SAITO Y ET AL: "N23K, a gene transiently up-regulated during neural differentiation, encodes a precursor protein for a newly identified neuropeptide nociceptin." see the whole document ---	21-28,32
P,X	NATURE, OCT 12 1995, 377 (6549) P532-5, ENGLAND, XP002020240 MEUNIER JC ET AL: "Isolation and structure of the endogenous agonist of opioid receptor-like ORL1 receptor [see comments]" see the whole document -----	21-28,32

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/13305

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: **1-15, 19-20, 29, 30**
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-15, 19-20, 29, 30 are directed to a method of treatment of, diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/13305

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9507983	23-03-95	NONE		
-----	-----	US-A-	5017689	21-05-91
WO-A-9100736	24-01-91	AT-T-	122566	15-06-95
		AU-B-	640158	19-08-93
		AU-A-	6077790	06-02-91
		CA-A-	2063712	07-01-91
		DE-D-	69019525	22-06-95
		DE-T-	69019525	18-01-96
		EP-A-	0486520	27-05-92
		JP-T-	4506525	12-11-92
-----	-----	-----	-----	-----