

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 February 2006 (09.02.2006)

PCT

(10) International Publication Number
WO 2006/014335 A2

(51) International Patent Classification:

C07K 16/24 (2006.01) G01N 33/574 (2006.01)
A61K 39/395 (2006.01) G01N 33/50 (2006.01)
C12Q 1/68 (2006.01) A61P 35/00 (2006.01)

(US). DANIEL, Dylan, L. [US/US]; 747 Page Street, San Francisco, CA 94117 (US). SHEK, Theresa [US/US]; 518 South Humboldt Street, San Mateo, CA 94402 (US).

(21) International Application Number:

PCT/US2005/023413

(74) Agents: NAIK, Paul et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(22) International Filing Date: 30 June 2005 (30.06.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/585,132 2 July 2004 (02.07.2004) US

(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ASHKENAZI, Avi J. [US/US]; 1400 Tarrytown Street, San Mateo, Ca 94402


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,

[Continued on next page]

(54) Title: COMPOSITIONS AND METHODS FOR TREATMENT OF NON-HODGKIN'S LYMPHOMA

(57) Abstract: The present invention is directed to compositions of matter useful for the treatment of non-hodgkin's lymphoma in mammals and to methods of using those compositions for the same.



WO 2006/014335 A2



SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished upon receipt of that report*

COMPOSITIONS AND METHODS FOR TREATMENT OF NON-HODGKIN'S LYMPHOMA

FIELD OF THE INVENTION

The present invention is directed to compositions of matter useful for the treatment of hematopoietic tumor in mammals and to methods of using those compositions of matter for the same.

BACKGROUND OF THE INVENTION

Malignant tumors (cancers) are the second leading cause of death in the United States, after heart disease (Boring et al., *CA Cancer J. Clin.* 43:7 (1993)). Cancer is characterized by the increase in the number of abnormal, or neoplastic, cells derived from a normal tissue which proliferate to form a tumor mass, the invasion of adjacent tissues by these neoplastic tumor cells, and the generation of malignant cells which eventually spread via the blood or lymphatic system to regional lymph nodes and to distant sites via a process called metastasis. In a cancerous state, a cell proliferates under conditions in which normal cells would not grow. Cancer manifests itself in a wide variety of forms, characterized by different degrees of invasiveness and aggressiveness.

Cancers which involve cells generated during hematopoiesis, a process by which cellular elements of blood, such as lymphocytes, leukocytes, platelets, erythrocytes and natural killer cells are generated are referred to as hematopoietic cancers. Lymphocytes which can be found in blood and lymphatic tissue and are critical for immune response are categorized into two main classes of lymphocytes: B lymphocytes (B cells) and T lymphocytes (T cells), which mediate humoral and cell mediated immunity, respectively.

B cells mature within the bone marrow and leave the marrow expressing an antigen-binding antibody on their cell surface. When a naive B cell first encounters the antigen for which its membrane-bound antibody is specific, the cell begins to divide rapidly and its progeny differentiate into memory B cells and effector cells called "plasma cells". Memory B cells have a longer life span and continue to express membrane-bound antibody with the same specificity as the original parent cell. Plasma cells do not produce membrane-bound antibody but instead produce the antibody in a form that can be secreted. Secreted antibodies are the major effector molecule of humoral immunity.

T cells mature within the thymus which provides an environment for the proliferation and differentiation of immature T cells. During T cell maturation, the T cells undergo the gene rearrangements that produce the T-cell receptor and the positive and negative selection which helps

determine the cell-surface phenotype of the mature T cell. Characteristic cell surface markers of mature T cells are the CD3:T-cell receptor complex and one of the coreceptors, CD4 or CD8.

In attempts to discover effective cellular targets for cancer therapy, researchers have sought to identify transmembrane or otherwise membrane-associated polypeptides that are specifically expressed on the surface of one or more particular type(s) of cancer cell as compared to on one or more normal non-cancerous cell(s). Often, such membrane-associated polypeptides are more abundantly expressed on the surface of the cancer cells as compared to on the surface of the non-cancerous cells. The identification of such tumor-associated cell surface antigen polypeptides has given rise to the ability to specifically target cancer cells for destruction via antibody-based therapies. In this regard, it is noted that antibody-based therapy has proved very effective in the treatment of certain cancers. For example, HERCEPTIN® and RITUXAN® (both from Genentech Inc., South San Francisco, California) are antibodies that have been used successfully to treat breast cancer and non-Hodgkin's lymphoma, respectively. More specifically, HERCEPTIN® is a recombinant DNA-derived humanized monoclonal antibody that selectively binds to the extracellular domain of the human epidermal growth factor receptor 2 (HER2) proto-oncogene. HER2 protein overexpression is observed in 25-30% of primary breast cancers. RITUXAN® is a genetically engineered chimeric murine/human monoclonal antibody directed against the CD20 antigen found on the surface of normal and malignant B lymphocytes. Both these antibodies are recombinantly produced in CHO cells.

In other attempts to discover effective cellular targets for cancer therapy, researchers have sought to identify (1) non-membrane-associated polypeptides that are specifically produced by one or more particular type(s) of cancer cell(s) as compared to by one or more particular type(s) of non-cancerous normal cell(s), (2) polypeptides that are produced by cancer cells at an expression level that is significantly higher than that of one or more normal non-cancerous cell(s), or (3) polypeptides whose expression is specifically limited to only a single (or very limited number of different) tissue type(s) in both the cancerous and non-cancerous state (e.g., normal prostate and prostate tumor tissue). Such polypeptides may remain intracellularly located or may be secreted by the cancer cell. Moreover, such polypeptides may be expressed not by the cancer cell itself, but rather by cells which produce and/or secrete polypeptides having a potentiating or growth-enhancing effect on cancer cells. Such secreted polypeptides are often proteins that provide cancer cells with a growth and/or survival advantage over normal cells and include such things as, for example, angiogenic factors, cellular adhesion factors, growth factors, and the like. Identification of antagonists of such non-membrane associated polypeptides would be expected to serve as effective therapeutic agents for the treatment of such cancers. Furthermore, identification of the expression pattern of such polypeptides would be useful for the diagnosis of particular cancers in mammals.

Despite the above-identified advances in mammalian cancer therapy, there is a great need for additional therapeutic agents capable of detecting the presence of tumor in a mammal and for effectively inhibiting neoplastic cell growth and/or survival, respectively. The present invention

discloses a polypeptide that is expressed at abnormally high levels in a great variety of non-Hodgkin's lymphomas, and provides methods and compositions for using this polypeptide as a diagnostic and/or therapeutic target in diagnosing and treating non-Hodgkin's lymphomas.

DISCLOSURE OF THE INVENTION

5 The present invention relates to the identification of a cellular polypeptide (and its encoding nucleic acid, variant or fragments thereof) which is specifically, differentially and/or highly expressed by non-Hodgkin's tumors compared to normal counterpart cell types. Interestingly and significantly, this cellular polypeptide is highly expressed across a wide spectrum of non-Hodgkin's lymphomas, including B and T-cell lymphomas. The invention is also in part based on substantial insight (as
10 described herein) into the differential regulation of the polypeptide with respect to specific subsets of non-Hodgkin's lymphoma types, which provides significant advantages in tailoring therapeutic approaches, if and where necessary, to selected subset(s) within the wide array of tumor types that collectively constitute the broad category of non-Hodgkin's lymphoma. This polypeptide is herein referred to as UNQ733 polypeptide and is expected to serve as an effective target for cancer therapy in
15 mammals.

 Accordingly, in one aspect, the invention provides a method for inhibiting the growth of a non-Hodgkin's lymphoma cell, wherein the method comprises contacting the cell with a UNQ733 antagonist which causes inhibition of the growth of the cell. In one embodiment, the UNQ733 antagonist binds to UNQ733 polypeptide. In one embodiment, the UNQ733 antagonist binds to
20 UNQ733 on a non-Hodgkin's lymphoma cell. In one embodiment, the UNQ733 antagonist causes death of a cell expressing and/or responsive to UNQ733. In one embodiment, the UNQ733 antagonist is an antibody, an oligopeptide, inorganic or organic small molecule. In one embodiment, binding of the antibody, oligopeptide, inorganic or organic small molecule to the UNQ733 polypeptide causes inhibition of the growth of a cell expressing and/or responsive to the polypeptide. In one embodiment,
25 the cell is a cancer cell and the UNQ733 antagonist causes death of the cell expressing and/or responsive to the UNQ733 polypeptide. In one embodiment, the cell is a cancer cell and binding of the antagonist to UNQ733 polypeptide causes death of the cell expressing and/or responsive to the polypeptide. In one embodiment, the UNQ733 antagonist is an antibody, oligopeptide, inorganic or organic small molecule and binding of the antibody, oligopeptide, inorganic or organic small molecule
30 to the UNQ733 polypeptide causes death of the cell expressing the polypeptide. In one embodiment, a UNQ733 antagonist inhibits activation of UNQ733, e.g. by modulating enzymatic processing of UNQ733 protein (e.g., by inhibiting a proprotein convertase that cleaves UNQ733). In one embodiment, an antibody is a monoclonal antibody, antibody fragment, chimeric antibody, humanized antibody, human antibody, multi-specific antibody or single-chain antibody. Antibodies, UNQ733
35 binding oligopeptides and UNQ733 binding inorganic or organic small molecules employed in the methods of the invention may optionally be conjugated to a growth inhibitory agent or cytotoxic agent such as a toxin, including, for example, a maytansinoid or calicheamicin, an antibiotic, a radioactive

isotope, a nucleolytic enzyme, or the like. In some embodiments of methods of the invention, a chemotherapeutic agent is also administered to the subject. The antibodies and UNQ733 binding oligopeptides employed in the methods of the invention may be produced in any suitable host cell, including for example CHO cells and bacterial cells.

5 In another aspect, the invention provides a method of therapeutically treating a mammal having non-Hodgkin's lymphoma, wherein the method comprises administering to the mammal a therapeutically effective amount of a UNQ733 antagonist, thereby resulting in the effective therapeutic treatment of the lymphoma. In one embodiment, the UNQ733 antagonist is an antibody, an oligopeptide, inorganic or organic small molecule. In one embodiment, the UNQ733 antagonist binds to UNQ733 polypeptide, thereby resulting in the effective therapeutic treatment of the tumor. In one
10 embodiment, a UNQ733 antagonist inhibits activation of UNQ733, for example by modulating enzymatic processing of UNQ733 protein (e.g., by inhibiting a proprotein convertase that cleaves UNQ733). In one embodiment, an antibody is a monoclonal antibody, antibody fragment, chimeric antibody, humanized antibody, human antibody, multi-specific antibody or single-chain antibody. Antibodies, UNQ733 binding oligopeptides and UNQ733 binding inorganic or organic small
15 molecules employed in the methods of the invention may optionally be conjugated to a growth inhibitory agent or cytotoxic agent such as a toxin, including, for example, a maytansinoid or calicheamicin, an antibiotic, a radioactive isotope, a nucleolytic enzyme, or the like. In some embodiments of methods of the invention, a chemotherapeutic agent is also administered to the subject. The antibodies and UNQ733 binding oligopeptides employed in the methods of the invention
20 may be produced in any suitable host cell, including for example CHO cells and bacterial cells.

In one aspect, the invention provides a method of inhibiting the growth of a non-Hodgkin's lymphoma comprising administering a UNQ733 antagonist to the lymphoma, whereby growth of the lymphoma is inhibited. In one aspect, the invention provides a method of inhibiting the growth of a
25 non-Hodgkin's lymphoma comprising administering a UNQ733 antagonist to a cell that expresses and/or is responsive to UNQ733, whereby growth of the lymphoma is inhibited. In one aspect, the invention provides a method of inhibiting growth of a non-Hodgkin's lymphoma comprising administering a UNQ733 antagonist to a cell present in and/or adjacent to the lymphoma, whereby growth of the lymphoma is inhibited. In one embodiment, said cell is not a non-Hodgkin's lymphoma cell (e.g., it is not a T or B cell); for example, said cell may be a stromal cell. In one embodiment, the
30 UNQ733 antagonist binds to UNQ733 polypeptide on a non-Hodgkin's lymphoma cell. In one embodiment, the UNQ733 antagonist causes death of a non-Hodgkin's lymphoma cell expressing and/or responsive to UNQ733. In one embodiment, the UNQ733 antagonist is an antibody, an oligopeptide, inorganic or organic small molecule. In one embodiment, binding of the antibody, oligopeptide, inorganic or organic small molecule to the UNQ733 polypeptide causes inhibition of the
35 growth of a non-Hodgkin's lymphoma cell expressing and/or responsive to the polypeptide. In one embodiment, the UNQ733 antagonist is an antibody, oligopeptide, inorganic or organic small molecule

and binding of the antibody, oligopeptide or organic small molecule to the UNQ733 polypeptide causes death of the non-Hodgkin's lymphoma cell expressing and/or responsive to the polypeptide. Optionally, the antibody is a monoclonal antibody, antibody fragment, chimeric antibody, humanized antibody, human antibody, multi-specific antibody or single-chain antibody. Antibodies, UNQ733 binding oligopeptides and UNQ733 binding inorganic or organic small molecules employed in the methods of the invention may optionally be conjugated to a growth inhibitory agent or cytotoxic agent such as a toxin, including, for example, a maytansinoid or calicheamicin, an antibiotic, a radioactive isotope, a nucleolytic enzyme, or the like. In one embodiment, a UNQ733 antagonist inhibits activation of UNQ733, for example by modulating enzymatic processing of UNQ733 protein (e.g., by inhibiting a proprotein convertase that cleaves UNQ733). In some embodiments of methods of the invention, a chemotherapeutic agent is also administered. The antibodies and UNQ733 binding oligopeptides employed in the methods of the invention may be produced in any suitable host cell, including for example CHO cells and bacterial cells.

In another aspect, the invention provides a method for treating or preventing a cell proliferative disorder associated with increased expression or activity of UNQ733, the method comprising administering to a subject in need of such treatment an effective amount of a UNQ733 antagonist. In one embodiment, the cell proliferative disorder is cancer and the UNQ733 antagonist is an anti-UNQ733 antibody, UNQ733 binding oligopeptide, UNQ733 binding inorganic or organic small molecule. In one embodiment, a UNQ733 antagonist inhibits activation of UNQ733, e.g. by modulating enzymatic processing of UNQ733 protein (e.g., by inhibiting a proprotein convertase that cleaves UNQ733). Effective treatment or prevention of the cell proliferative disorder may be a result of direct killing or growth inhibition of cells that express and/or are responsive to UNQ733 polypeptide or by antagonizing the cell growth potentiating activity of UNQ733 polypeptide. In one embodiment, the cell proliferative disorder is a non-Hodgkin's lymphoma. In one embodiment, the cell proliferative disorder is hyperplasia, which in one embodiment is diseased, non-malignant tonsillar tissue.

In one aspect the invention provides a method of diagnosing a non-Hodgkin's lymphoma of interest comprising determining the level of UNQ733 in a test sample of tissue cells comprising cells of origin for the lymphoma of interest and in a control sample of known cells of the same tissue origin, wherein a higher level of UNQ733 in the test sample, as compared to the control sample, is indicative of the presence of the lymphoma. In another aspect, the invention provides a method of determining whether an individual is at risk for non-Hodgkin's lymphoma comprising determining the level of UNQ733 in a test sample of tissue cells comprising cells of origin for the lymphoma of interest from said individual and in a control sample of known cells of the same tissue origin, wherein a higher level of UNQ733 in the test sample, as compared to the control sample, is an indication that the individual is at risk for the lymphoma. In one embodiment of methods of the invention, the level of UNQ733 is determined based on level of UNQ733 RNA in the test sample. In one embodiment of methods of the

invention, the level of UNQ733 is determined based on level of UNQ733 polypeptide in the test sample. In one embodiment of methods of the invention, the level of UNQ733 is determined based on the level of UNQ733 protein activity in the sample. In some embodiments, the method comprises exposing the sample to an antibody, oligopeptide, inorganic or organic small molecule that binds to the UNQ733 polypeptide and determining binding of the antibody, oligopeptide, inorganic or organic small molecule to the UNQ733 polypeptide in the sample. An antibody, UNQ733 binding oligopeptide, or UNQ733 binding inorganic or organic small molecule employed in the method may optionally be detectably labeled, attached to a solid support, or the like.

In one aspect, the invention provides a method for screening for biologically active agents for the treatment of non-Hodgkin's lymphoma comprising combining a candidate agent with a transgenic mammal having a genome comprising an integrated transgene encoding UNQ733 operably linked to a promoter, wherein said transgene results in said mammal acquiring non-Hodgkin's lymphoma and determining the effect of said agent on non-Hodgkin's lymphoma in said mammal. In yet another aspect, the invention provides a method for screening for biologically active agents for the treatment of non-Hodgkin's lymphoma comprising combining a candidate agent with a transgenic mammal cell culture, each cell of said culture comprising an integrated transgene encoding UNQ733 operably linked to a promoter, wherein said transgene results in said mammal acquiring non-Hodgkin's lymphoma, and determining the effect of said candidate agent on the transgenic mammal cell culture. In one embodiment of these methods, the candidate agent is an antibody or fragment thereof, an oligopeptide, or an inorganic or organic small molecule.

Yet another aspect of the present invention is directed to a method of binding an antibody, oligopeptide, inorganic or organic small molecule to a non-Hodgkin's lymphoma cell that expresses and/or is responsive to UNQ733, wherein the method comprises contacting said cell with said antibody, oligopeptide, inorganic or organic small molecule under conditions which are suitable for binding of the antibody, oligopeptide, or inorganic or organic small molecule to UNQ733 polypeptide and allowing binding therebetween. In one embodiment, binding of said antibody, oligopeptide, inorganic or organic small molecule to UNQ733 on the cell inhibits a UNQ733 biological function. In one embodiment, said antibody, oligopeptide, inorganic or organic small molecule does not compete with UNQ733 for binding to the cell. In one embodiment, said antibody, oligopeptide, inorganic or organic small molecule does not inhibit binding of UNQ733 to the cell. In one embodiment, said antibody, oligopeptide, inorganic or organic small molecule binds to UNQ733 bound on the cell and inhibits binding of free (unbound) UNQ733 to the cell. In one embodiment, the UNQ733 binding molecule is an antibody. In one embodiment, the antibody is designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody binds to

the same epitope on UNQ733 as the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody competes with the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034) for binding to UNQ733 (e.g., where UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell (*in vitro* or *in vivo*)).

In one aspect, the invention provides a method of targeting a therapeutic agent to a non-Hodgkin's lymphoma in a host, the method comprising administering to the host said therapeutic agent in a form that is linked to a molecule that binds UNQ733, whereby the agent is targeted to the lymphoma in the host. The molecule that binds UNQ733 can be any molecule capable of specifically binding to UNQ733 protein *in vivo*, for example an antibody, an oligopeptide, an inorganic or organic small molecule. In one embodiment, the molecule that binds UNQ733 is a molecule capable of specifically binding to UNQ733 located on a cell (either *in vitro* or *in vivo*), for example where UNQ733 is bound to the surface of a non-Hodgkin's lymphoma cell. In one embodiment, the molecule is an antibody. In one embodiment, the antibody is designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody binds to the same epitope on UNQ733 as the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody competes with the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034) for binding to UNQ733 (e.g., where UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell (*in vitro* or *in vivo*)). In one embodiment, the therapeutic agent is linked to the molecule that binds UNQ733 in the form of a multi-specific (e.g., bi-specific) antibody.

Other embodiments of the invention are directed to the use of (a) a UNQ733 polypeptide (processed or unprocessed by a proprotein convertase, singly or in combination), (b) a nucleic acid encoding a processed or unprocessed UNQ733 polypeptide or a vector or host cell comprising that nucleic acid, (c) an anti-UNQ733 polypeptide antibody, (d) a UNQ733 binding oligopeptide, (e) a UNQ733 binding inorganic or organic small molecule, or (f) an inhibitor of an *in vivo* enzyme (such as a proprotein convertase) that cleaves UNQ733 protein, in the preparation of a medicament useful for the therapeutic treatment of a non-Hodgkin's lymphoma.

In one aspect, the invention provides a UNQ733 binding molecule useful in a method of the invention as herein described. For example, a UNQ733 binding molecule can be a molecule (such as an antibody) that binds to UNQ733 and inhibits binding of UNQ733 to its receptor. In another example, a UNQ733 binding molecule can be a molecule (such as an antibody) that binds UNQ733 that is bound to a cell. In one embodiment, the binding molecule binds UNQ733 that is bound to a cell but does not inhibit UNQ733 binding to a cell. In yet another example, a UNQ733 binding molecule is a molecule (such as an antibody) that binds UNQ733 only when said UNQ733 is bound to a cell. In one embodiment, the binding molecule binds a first UNQ733 polypeptide only when said first polypeptide is bound to a cell but does not inhibit binding of a second UNQ733 polypeptide to a cell. In some embodiments, it is advantageous for a UNQ733 binding molecule of the invention to be conjugated to a therapeutic agent (e.g., a toxin, etc. as described in greater detail hereinbelow). In one aspect, the invention provides a composition comprising a UNQ733 antagonist, wherein the composition is suitable for administration to a subject having non-Hodgkin's lymphoma. In one embodiment, the composition comprises a therapeutically effective amount of the UNQ733 antagonist. In one embodiment, the composition further comprises a carrier which in some embodiments is a pharmaceutically acceptable carrier. In one embodiment, the UNQ733 antagonist is an antibody. In one embodiment, the antibody is designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody binds to the same epitope on UNQ733 as the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody competes with the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034).

for binding to UNQ733 (e.g., where UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell (*in vitro* or *in vivo*)).

In yet another aspect, the invention provides an article of manufacture comprising a container and a composition contained within the container, wherein the composition comprises a UNQ733 antagonist as described herein. The article may further optionally comprise a label affixed to the container, and/or a package insert included with the container, that refers to the use of the composition for therapeutic treatment of non-Hodgkin's lymphoma. In one embodiment, the invention provides a kit comprising a composition comprising a UNQ733 antagonist as described herein, optionally further comprising instruction for using the composition for treatment of non-Hodgkin's lymphoma.

In one aspect, the invention provides a composition comprising a UNQ733 polypeptide binding molecule, wherein the composition is suitable for detecting UNQ733 polypeptide in a test sample from an individual suspected of having non-Hodgkin's lymphoma. In one embodiment, the binding molecule is an antibody, oligopeptide, inorganic or organic small molecule. In one embodiment, the binding molecule is an antibody, wherein the antibody is capable of binding to UNQ733 in a detection assay such as western blot, immunoprecipitation and/or immunostaining. In one embodiment, the molecule is an antibody. In one embodiment, the antibody is designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody binds to the same epitope on UNQ733 as the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody competes with the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034) for binding to UNQ733 (e.g., where UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell (*in vitro* or *in vivo*)). In one embodiment, the composition further comprises a carrier.

In yet another aspect, the invention provides an article of manufacture comprising a container and a composition contained within the container, wherein the composition comprises a UNQ733 polypeptide binding molecule and is suitable for detecting UNQ733 polypeptide in a test sample from an individual suspected of having non-Hodgkin's lymphoma. The article may further optionally comprise a label affixed to the container, and/or a package insert included with the container, that

refers to the use of the composition for detection of non-Hodgkin's lymphoma. In one embodiment, the invention provides a kit comprising a composition comprising a UNQ733 polypeptide binding molecule, optionally further comprising instruction for using the composition for detecting non-Hodgkin's lymphoma. In one embodiment, the binding molecule is an antibody, wherein the antibody is capable of binding to UNQ733 in a detection assay such as western blot, immunoprecipitation and/or immunostaining. In one embodiment, the antibody is In one embodiment, the molecule is an antibody. In one embodiment, the antibody is designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody binds to the same epitope on UNQ733 as the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, the antibody competes with the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034) for binding to UNQ733 (e.g., where UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell (*in vitro* or *in vivo*)).

As described herein, an antibody of the invention can be in any suitable form for use in a method of the invention. For example, an antibody of the invention can be human, humanized or chimeric. In one embodiment, an antibody of the invention is the humanized or chimeric form of the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, an antibody of the invention is a humanized, chimeric or human antibody that binds to the same epitope on UNQ733 as the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, an antibody of the invention is a humanized, chimeric or human antibody that competes with the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC

deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) and/or 12H4.11.3 (ATCC deposit no. PTA-6034) for binding to UNQ733 (e.g., where UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell (*in vitro* or *in vivo*)). In one embodiment, an antibody of the invention comprises one, two, three, four, five or all of the hypervariable regions, hypervariable loops and/or complementarity determining region (CDR) sequences of the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034). In one embodiment, an antibody of the invention comprises one or both variable domains, or portion thereof, of the antibody designated 3E7.9.20 (ATCC deposit no. PTA-6026), 3F10.11.2 (ATCC deposit no. PTA-6027), 3H1.4.8 (ATCC deposit no. PTA-6028), 4A9.12.12 (ATCC deposit no. PTA-6029), 5A8.11.6 (ATCC deposit no. PTA-6030), 5F2.6.14 (ATCC deposit no. PTA-6031), 9D6.11.15 (ATCC deposit no. PTA-6032), 10G10.15.16 (ATCC deposit no. PTA-6033) or 12H4.11.3 (ATCC deposit no. PTA-6034).

In some embodiments, methods and compositions of the invention are directed to a non-Hodgkin's lymphoma that is a B-cell lymphoma (not-otherwise-specified - "NOS"), diffuse large B cell lymphoma, follicular lymphoma, small lymphocytic lymphoma, malignant lymphoma (NOS), malignant T cell lymphoma, anaplastic large cell lymphoma or mucosal associated lymphoid tissue lymphoma.

Yet further embodiments of the present invention will be evident to the skilled artisan upon a reading of the present specification.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 depicts GeneLogic microarray expression data of the UNQ733 mRNA.

FIGURE 2 depicts UNQ733 mRNA overexpression profile in non-Hodgkin's lymphomas assessed by in situ hybridization on tissue microarray.

FIGURE 3 depicts UNQ733 mRNA expression profile in normal human tissues as assessed by in situ hybridization.

FIGURES 4A-D depict UNQ733 mRNA expression in germinal centers and crypts.

FIGURE 5 depicts UNQ733 mRNA expression profile in neoplastic tissues.

FIGURES 6A-C depict cellular localization of UNQ733 mRNA in malignant cells of non-Hodgkin's lymphoma.

FIGURE 7 depicts Western blot analysis of UNQ733 protein.

FIGURE 8 depicts immunoprecipitation data for UNQ733 protein.

FIGURE 9 depicts immunohistochemistry data for UNQ733 protein. Expression is observed in a subset of cells in tonsil crypt (arrows) and in germinal centers (GC), most consistent with a follicular dendritic cell.

FIGURE 10 depicts flow cytometry data.

FIGURE 11 depicts the effect of a proprotein convertase on UNQ733. Lanes marked "v" refer to samples from vector mock transfectants. Lanes marked "733" refer to samples from UNQ733 transfectants.

FIGURE 12 depicts proportions of various UNQ733 protein species obtained when cells were contacted with or without furin inhibitor.

FIGURES 13A-B depict flow cytometry data for UNQ733 protein binding to lymphoma cells.

FIGURE 14 depicts flow cytometry data for detection of UNQ733 protein bound to non-Hodgkin's lymphoma cells.

FIGURES 15A-B depict binding of UNQ733 protein to C1R B lymphoblast cells in a dose-dependent manner (A) and competition binding data indicating specificity of binding of UNQ733 to these cells (B).

FIGURE 16 depicts UNQ733 binding on normal B cells that are unstimulated or stimulated with agonistic antibody to CD40 and recombinant IL-4.

FIGURE 17 shows a nucleotide sequence (SEQ ID NO:1) of a UNQ733 cDNA. Underlined codons in bold are the predicted start and stop codons, respectively.

FIGURE 18 shows the amino acid sequences (SEQ ID NOs:2-4) of native and variant UNQ733 polypeptides.

MODES FOR CARRYING OUT THE INVENTION

I. Definitions

The terms "UNQ733 polypeptide" and "UNQ733 protein" as used herein encompass native sequence polypeptides, polypeptide variants and fragments of a native sequence polypeptide and polypeptide variants (which are further defined herein). The UNQ733 polypeptide described herein may be that which is isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The terms "UNQ733 polypeptide" and "UNQ733 protein" also include variants of a UNQ733 polypeptide as disclosed herein.

A "native sequence UNQ733 polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding UNQ733 polypeptide derived from nature. In one embodiment, a native sequence UNQ733 polypeptide comprises the amino acid sequence of SEQ ID NO:2 (see Figure 18). In another embodiment, a native sequence UNQ733 polypeptide comprises an amino acid sequence lacking the signal peptide. In one embodiment, a native sequence UNQ733 polypeptide comprises the amino acid sequence of SEQ ID NO:3 (see Figure 18). In yet another embodiment, a native sequence UNQ733 polypeptide comprises an amino acid sequence resulting from enzymatic cleavage of the sequence of SEQ ID NO:2 with a proprotein convertase. In one embodiment, a native

sequence UNQ733 polypeptide comprises the amino acid sequence of SEQ ID NO:4 (see Figure 18). Such native sequence UNQ733 polypeptide can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence UNQ733 polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific UNQ733 polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. Native sequence UNQ733 is also reported in Marshall et al, J. Immunol. (2002), 169:2381-2389; PCT Pub. No. WO9931117; and PCT Pub. No. WO02/08288.

"UNQ733 polypeptide variant" means a UNQ733 polypeptide, preferably an active UNQ733 polypeptide, as defined herein having at least about 80% amino acid sequence identity with any of the native sequence UNQ733 polypeptide sequences as disclosed herein. Such UNQ733 polypeptide variants include, for instance, UNQ733 polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of a native amino acid sequence. Ordinarily, a UNQ733 polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% amino acid sequence identity, to a native sequence UNQ733 polypeptide sequence as disclosed herein. Ordinarily, UNQ733 variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600 amino acids in length, or more. Optionally, UNQ733 variant polypeptides will have no more than one conservative amino acid substitution as compared to a native UNQ733 polypeptide sequence, alternatively no more than 2, 3, 4, 5, 6, 7, 8, 9, or 10 conservative amino acid substitution as compared to the native UNQ733 polypeptide sequence.

"Percent (%) amino acid sequence identity" with respect to the UNQ733 polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific UNQ733 polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech,

Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1 below. The
5 ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or
10 comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment
15 program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid
20 sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "UNQ733", wherein "UNQ733" represents the amino acid sequence of a hypothetical UNQ733 polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "UNQ733" polypeptide of interest is being compared, and "X," "Y" and "Z" each represent different hypothetical amino acid residues. Unless specifically stated
25 otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

"UNQ733 variant polynucleotide" or "UNQ733 variant nucleic acid sequence" means a nucleic acid molecule which encodes a UNQ733 polypeptide, preferably an active UNQ733 polypeptide, as defined herein and which has at least about 80% nucleic acid sequence identity with a
30 nucleotide acid sequence encoding a native sequence UNQ733 polypeptide sequence as disclosed herein. Ordinarily, a UNQ733 variant polynucleotide will have at least about 80% nucleic acid sequence identity, alternatively at least about 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% nucleic acid sequence identity with a nucleic acid sequence encoding a native sequence UNQ733 polypeptide sequence as disclosed herein.
35 Variants do not encompass the native nucleotide sequence.

Ordinarily, UNQ733 variant polynucleotides are at least about 5 nucleotides in length, alternatively at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26,

27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, or 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length.

"Percent (%) nucleic acid sequence identity" with respect to a UNQ733 nucleic acid sequence (encoding a native sequence UNQ733 polypeptide, for example the nucleic acid sequence of SEQ ID NO:1 in Figure 17) disclosed herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the UNQ733 nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of

nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "UNQ733 DNA", wherein "UNQ733 DNA" represents a hypothetical UNQ733 nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "UNQ733 DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides. Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

In other embodiments, UNQ733 variant polynucleotides are nucleic acid molecules that encode a UNQ733 polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a UNQ733 polypeptide as disclosed herein. UNQ733 variant polypeptides may be those that are encoded by a UNQ733 variant polynucleotide.

"Isolated," means a molecule/compound (such as a polypeptide) that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with therapeutic uses for the molecule/compound (such as a polypeptide), and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In some embodiments, a polypeptide or oligopeptide will be purified (1) to a degree sufficient to obtain at least 10-15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by gel electrophoresis, for example SDS-PAGE under non-reducing or reducing conditions using, for example, Coomassie blue or silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the UNQ733 polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" polynucleotide is a polypeptide or oligopeptide-encoding nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide or oligopeptide-encoding nucleic acid. An isolated polypeptide or oligopeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide or oligopeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide or oligopeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide or oligopeptide-encoding nucleic acid molecule includes polypeptide or oligopeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide or oligopeptide where, for example, the nucleic acid molecule is in a chromosomal or extrachromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

5 Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

15 "Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

25 "Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) overnight hybridization in a solution that employs 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with a 10 minute wash at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) followed by a 10 minute high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

35 "Moderately stringent conditions" may be identified as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include

the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a UNQ733 polypeptide or anti-UNQ733 antibody fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues). Examples of tag polypeptides include detectable markers such as polyhistidine and human placenta alkaline phosphatase.

"Active" or "activity" for the purposes herein refers to form(s) of a UNQ733 polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring UNQ733 polypeptide, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring UNQ733 polypeptide other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring UNQ733 polypeptide and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring UNQ733 polypeptide.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native UNQ733 polypeptide disclosed herein. Suitable antagonist molecules specifically include antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native UNQ733 polypeptides, peptides, antisense oligonucleotides, small organic or inorganic small molecules, etc. Methods for identifying antagonists of a UNQ733 polypeptide may comprise contacting a UNQ733 polypeptide with a candidate antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the UNQ733 polypeptide.

"Treating" or "treatment" or "alleviation" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. A subject or mammal is successfully "treated" for a non-Hodgkin's lymphoma if, after receiving a therapeutic

amount of a UNQ733 antagonist according to the methods of the invention, the patient shows observable and/or measurable reduction in or absence of one or more of the following: reduction in the number of cancer cells or absence of the cancer cells; reduction in the tumor size; inhibition (i.e., slow to some extent and preferably stop) of cancer cell infiltration into peripheral organs including the spread of cancer into soft tissue and bone; inhibition (i.e., slow to some extent and preferably stop) of tumor metastasis; inhibition, to some extent, of tumor growth; and/or relief to some extent, one or more of the symptoms associated with the specific cancer; reduced morbidity and mortality, and improvement in quality of life issues. To the extent the UNQ733 antagonist may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. Reduction of these signs or symptoms may also be felt by the patient.

The above parameters for assessing successful treatment and improvement in the disease are readily measurable by routine procedures familiar to a physician. For cancer therapy, efficacy can be measured, for example, by assessing the time to disease progression (TTP) and/or determining the response rate (RR). Metastasis can be determined by staging tests and by bone scan and tests for calcium level and other enzymes to determine spread to the bone. CT scans can also be done to look for spread to the pelvis and lymph nodes in the area. Chest X-rays and measurement of liver enzyme levels by known methods are used to look for metastasis to the lungs and liver, respectively. Other routine methods for monitoring the disease include transrectal ultrasonography (TRUS) and transrectal needle biopsy (TRNB).

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Mammal" for purposes of the treatment of, alleviating the symptoms of a cancer refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or

dextrins; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN®, polyethylene glycol (PEG), and PLURONICS®.

By "solid phase" or "solid support" is meant a non-aqueous matrix to which an antibody, UNQ733 polypeptide binding oligopeptide or UNQ733 polypeptide binding organic or inorganic small molecule of the invention can adhere or attach. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a UNQ733 polypeptide, an antibody thereto or a UNQ733 polypeptide binding oligopeptide) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

A "small" molecule or "small" organic small molecule is defined herein to have a molecular weight below about 500 Daltons.

An "effective amount" of a UNQ733 antagonist as disclosed herein is an amount sufficient to carry out a specifically stated purpose. An "effective amount" may be determined empirically and in a routine manner, in relation to the stated purpose.

The term "therapeutically effective amount" refers to an amount of a UNQ733 antagonist effective to "treat" a disease or disorder in a subject or mammal. In the case of cancer, the therapeutically effective amount of the antagonist may reduce the number of cancer cells; reduce the tumor size; inhibit (i.e., slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (i.e., slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. See the definition herein of "treating". To the extent the antagonist may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic.

A "growth inhibitory amount" of a UNQ733 antagonist is an amount capable of inhibiting the growth of a cell, especially tumor, e.g., cancer cell, either *in vitro* or *in vivo*. A "growth inhibitory amount" of a UNQ733 antagonist for purposes of inhibiting neoplastic cell growth may be determined empirically and in a routine manner.

A "cytotoxic amount" of a UNQ733 antagonist is an amount capable of causing the destruction of a cell, especially tumor, e.g., cancer cell, either *in vitro* or *in vivo*. A "cytotoxic amount" of a UNQ733 antagonist for purposes of inhibiting neoplastic cell growth may be determined empirically and in a routine manner.

The term "antibody" is used in the broadest sense and specifically covers, for example, single anti-UNQ733 polypeptide monoclonal antibodies (including antagonist, binding and/or neutralizing antibodies), anti-UNQ733 polypeptide antibody compositions with polypeptopic specificity, polyclonal antibodies, single chain anti-UNQ733 polypeptide antibodies, and fragments of anti-UNQ733 polypeptide antibodies (see below) as long as they exhibit the desired biological or immunological activity. The term "immunoglobulin" (Ig) is used interchangeable with antibody herein.

An antibody useful in methods of the invention is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by gel electrophoresis such as SDS-PAGE under reducing or nonreducing conditions using, for example, Coomassie blue or silver stain.

The basic 4-chain antibody unit is a heterotetrameric glycoprotein composed of two identical light (L) chains and two identical heavy (H) chains (an IgM antibody consists of 5 of the basic heterotetramer unit along with an additional polypeptide called J chain, and therefore contain 10 antigen binding sites, while secreted IgA antibodies can polymerize to form polyvalent assemblages comprising 2-5 of the basic 4-chain units along with J chain). In the case of IgGs, the 4-chain unit is generally about 150,000 daltons. Each L chain is linked to a H chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. Each H and L chain also has regularly spaced intrachain disulfide bridges. Each H chain has at the N-terminus, a variable domain (V_H) followed by three constant domains (C_H) for each of the α and γ chains and four C_H domains for μ and ϵ isotypes. Each L chain has at the N-terminus, a variable domain (V_L) followed by a constant domain (C_L) at its other end. The V_L is aligned with the V_H and the C_L is aligned with the first constant domain of the heavy chain (C_{H1}). Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains. The pairing of a V_H and V_L together forms a single antigen-binding site. For the structure and properties of the different classes of antibodies, see, e.g., Basic and Clinical Immunology, 8th edition, Daniel P. Stites, Abba I. Terr and Tristram G. Parslow (eds.), Appleton & Lange, Norwalk, CT, 1994, page 71 and Chapter 6.

The L chain from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains (C_H), immunoglobulins can be assigned to different classes or isotypes. There are five classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, having heavy chains designated α , δ , ϵ , γ , and μ , respectively. The γ and α classes are

further divided into subclasses on the basis of relatively minor differences in C_H sequence and function, e.g., humans express the following subclasses: IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2.

The term "variable" refers to the fact that certain segments of the variable domains differ extensively in sequence among antibodies. The V domain mediates antigen binding and define specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the 110-amino acid span of the variable domains. Instead, the V regions consist of relatively invariant stretches called framework regions (FRs) of 15-30 amino acids separated by shorter regions of extreme variability called "hypervariable regions" that are each 9-12 amino acids long. The variable domains of native heavy and light chains each comprise four FRs, largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC).

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody which are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a "complementarity determining region" or "CDR" (e.g. around about residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the V_L, and around about 1-35 (H1), 50-65 (H2) and 95-102 (H3) in the V_H; Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the V_L, and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the V_H; Chothia and Lesk J. Mol. Biol. 196:901-917 (1987)).

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations which include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier "monoclonal" is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies useful in the present invention may be prepared by the hybridoma methodology first described by Kohler et al., Nature, 256:495 (1975), or may be made using recombinant DNA methods in bacterial, eukaryotic animal or plant cells (see, e.g., U.S. Patent

No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson et al., Nature, 352:624-628 (1991) and Marks et al., J. Mol. Biol., 222:581-597 (1991), for example.

The monoclonal antibodies herein include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see U.S. Patent No. 4,816,567; and Morrison et al., Proc. Natl. Acad. Sci. USA, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include "primatized" antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (*e.g.* Old World Monkey, Ape etc), and human constant region sequences.

An "intact" antibody is one which comprises an antigen-binding site as well as a C_L and at least heavy chain constant domains, C_{H1}, C_{H2} and C_{H3}. The constant domains may be native sequence constant domains (*e.g.* human native sequence constant domains) or amino acid sequence variant thereof. Preferably, the intact antibody has one or more effector functions.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (see U.S. Patent No. 5,641,870, Example 2; Zapata et al., Protein Eng. 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. The Fab fragment consists of an entire L chain along with the variable region domain of the H chain (V_H), and the first constant domain of one heavy chain (C_{H1}). Each Fab fragment is monovalent with respect to antigen binding, i.e., it has a single antigen-binding site. Pepsin treatment of an antibody yields a single large F(ab')₂ fragment which roughly corresponds to two disulfide linked Fab fragments having divalent antigen-binding activity and is still capable of cross-linking antigen. Fab' fragments differ from Fab fragments by having additional few residues at the carboxy terminus of the C_{H1} domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The Fc fragment comprises the carboxy-terminal portions of both H chains held together by disulfides. The effector functions of antibodies are determined by sequences in the Fc region, which region is also the part recognized by Fc receptors (FcR) found on certain types of cells.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (3 loops each from the H and L chain) that contribute the amino acid residues for antigen binding and confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

"Single-chain Fv" also abbreviated as "sFv" or "scFv" are antibody fragments that comprise the V_H and V_L antibody domains connected into a single polypeptide chain. Preferably, the sFv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994); Borrebaeck 1995, *infra*.

The term "diabodies" refers to small antibody fragments prepared by constructing sFv fragments (see preceding paragraph) with short linkers (about 5-10 residues) between the V_H and V_L domains such that inter-chain but not intra-chain pairing of the V domains is achieved, resulting in a bivalent fragment, i.e., fragment having two antigen-binding sites. Bispecific diabodies are heterodimers of two "crossover" sFv fragments in which the V_H and V_L domains of the two antibodies are present on different polypeptide chains. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

"Humanized" forms of non-human (*e.g.*, rodent) antibodies are chimeric antibodies that contain minimal sequence derived from the non-human antibody. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or non-human primate having the desired antibody specificity, affinity, and capability. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., Nature 321:522-525 (1986); Riechmann et al., Nature 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol. 2:593-596 (1992).

A "species-dependent antibody," e.g., a mammalian anti-human IgE antibody, is an antibody which has a stronger binding affinity for an antigen from a first mammalian species than it has for a homologue of that antigen from a second mammalian species. Normally, the species-dependent antibody "bind specifically" to a human antigen (i.e., has a binding affinity (K_d) value of no more than about 1×10^{-7} M, preferably no more than about 1×10^{-8} and most preferably no more than about 1×10^{-9} M) but has a binding affinity for a homologue of the antigen from a second non-human mammalian species which is at least about 50 fold, or at least about 500 fold, or at least about 1000 fold, weaker than its binding affinity for the human antigen. The species-dependent antibody can be of any of the various types of antibodies as defined above, but preferably is a humanized or human antibody.

A "UNQ733 polypeptide binding oligopeptide" is an oligopeptide that binds, preferably specifically, to a UNQ733 polypeptide as described herein. UNQ733 polypeptide binding oligopeptides may be chemically synthesized using known oligopeptide synthesis methodology or may be prepared and purified using recombinant technology. UNQ733 polypeptide binding oligopeptides are usually at least about 5 amino acids in length, alternatively at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 amino acids in length or more, wherein such oligopeptides are capable of binding, preferably specifically, to a UNQ733 polypeptide as described herein. UNQ733 polypeptide binding oligopeptides may be identified without undue experimentation using well known techniques. In this regard, it is noted that techniques for screening oligopeptide libraries for oligopeptides that are capable of specifically binding to a polypeptide target are well known in the art (see, e.g., U.S. Patent Nos. 5,556,762, 5,750,373, 4,708,871, 4,833,092, 5,223,409, 5,403,484, 5,571,689, 5,663,143; PCT Publication Nos. WO 84/03506 and WO84/03564; Geysen et al., Proc. Natl. Acad. Sci. U.S.A., 81:3998-4002 (1984); Geysen et al., Proc. Natl. Acad. Sci. U.S.A., 82:178-182 (1985); Geysen et al., in Synthetic Peptides as Antigens, 130-149 (1986); Geysen et al., J. Immunol. Meth., 102:259-274 (1987); Schoofs et al., J. Immunol., 140:611-616 (1988), Cwirla, S. E. et al. (1990) Proc. Natl. Acad. Sci. USA, 87:6378; Lowman, H.B. et al. (1991) Biochemistry, 30:10832; Clackson, T. et al. (1991) Nature, 352: 624; Marks, J. D. et al. (1991), J. Mol. Biol., 222:581; Kang, A.S. et al. (1991) Proc. Natl. Acad. Sci. USA, 88:8363, and Smith, G. P. (1991) Current Opin. Biotechnol., 2:668).

A "UNQ733 polypeptide binding organic or inorganic small molecule" is an organic or inorganic small molecule other than an oligopeptide or antibody as defined herein that binds, preferably specifically, to a UNQ733 polypeptide as described herein. A small molecule UNQ733 antagonist is preferably an organic small molecule. UNQ733 polypeptide binding organic may be identified and chemically synthesized using known methodology (see, e.g., PCT Publication Nos. WO00/00823 and WO00/39585). UNQ733 polypeptide binding organic small molecules are usually

less than about 2000 daltons in size, alternatively less than about 1500, 750, 500, 250 or 200 daltons in size, wherein such organic small molecules that are capable of binding, preferably specifically, to a UNQ733 polypeptide as described herein may be identified without undue experimentation using well known techniques. In this regard, it is noted that techniques for screening organic small molecule libraries for molecules that are capable of binding to a polypeptide target are well known in the art (see, e.g., PCT Publication Nos. WO00/00823 and WO00/39585).

An antibody, oligopeptide, or other organic or inorganic small molecule “which binds” an antigen of interest, e.g. a UNQ733 polypeptide, is one that binds the antigen with sufficient affinity such that the antibody, oligopeptide, or other organic or inorganic small molecule is useful as a therapeutic agent in targeting a cell or tissue expressing the antigen, and does not significantly cross-react with other proteins. In such embodiments, the extent of binding of the antibody, oligopeptide, or other organic or inorganic small molecule to a “non-target” protein will be less than about 10% of the binding of the antibody, oligopeptide, or other organic or inorganic small molecule to its particular target protein as determined by fluorescence activated cell sorting (FACS) analysis or radioimmunoprecipitation (RIA). With regard to the binding of an antibody, oligopeptide, or other organic or inorganic small molecule to a target molecule, the term “specific binding” or “specifically binds to” or is “specific for” a particular polypeptide or an epitope on a particular polypeptide target means binding that is measurably different from a non-specific interaction. Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule, which generally is a molecule of similar structure that does not have binding activity. For example, specific binding can be determined by competition with a control molecule that is similar to the target, for example, an excess of non-labeled target. In this case, specific binding is indicated if the binding of the labeled target to a probe is competitively inhibited by excess unlabeled target. The term “specific binding” or “specifically binds to” or is “specific for” a particular polypeptide or an epitope on a particular polypeptide target as used herein can be exhibited, for example, by a molecule having a K_d for the target of at least about 10^{-4} M, alternatively at least about 10^{-5} M, alternatively at least about 10^{-6} M, alternatively at least about 10^{-7} M, alternatively at least about 10^{-8} M, alternatively at least about 10^{-9} M, alternatively at least about 10^{-10} M, alternatively at least about 10^{-11} M, alternatively at least about 10^{-12} M, or greater. In one embodiment, the term “specific binding” refers to binding where a molecule binds to a particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

An antibody, oligopeptide, or other organic or inorganic small molecule that “inhibits the growth of tumor cells expressing UNQ733 polypeptide” or a “growth inhibitory” antibody, oligopeptide, or other organic or inorganic small molecule is one which results in measurable growth inhibition of non-Hodgkin’s lymphoma cells expressing or overexpressing the UNQ733 polypeptide. The UNQ733 polypeptide may be bound to the surface of a cell or may be in an extracellular environment. Preferred growth inhibitory anti-UNQ733 polypeptide antibodies, oligopeptides, or

organic or inorganic small molecules inhibit growth of UNQ733 polypeptide-expressing tumor cells by preferably greater than about 20%, preferably from about 20% to about 50%, preferably by greater than 50% (e.g., from about 50% to about 100%) as compared to the appropriate control, the control typically being tumor cells not treated with the antibody, oligopeptide, or other organic or inorganic small molecule being tested. In one embodiment, growth inhibition can be measured at an antibody concentration of about 0.1 to 30 $\mu\text{g/ml}$ or about 0.5 nM to 200 nM in cell culture, where the growth inhibition is determined 1-10 days after exposure of the tumor cells to the antibody. The antibody is growth inhibitory in vivo if administration of the anti-UNQ733 polypeptide antibody at about 1 $\mu\text{g/kg}$ to about 100 mg/kg body weight results in reduction in tumor size or tumor cell proliferation within about 5 days to 3 months from the first administration of the antibody, or within about 5 to 30 days.

A UNQ733 antagonist which "induces apoptosis" is one which induces programmed cell death as determined by binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies). The cell is usually one which overexpresses UNQ733 polypeptide. Preferably the cell is a tumor cell of a non-Hodgkin's lymphoma. Various methods are available for evaluating the cellular events associated with apoptosis. For example, phosphatidyl serine (PS) translocation can be measured by annexin binding; DNA fragmentation can be evaluated through DNA laddering; and nuclear/chromatin condensation along with DNA fragmentation can be evaluated by any increase in hypodiploid cells. Preferably, the UNQ733 antagonist which induces apoptosis is one which results in about 2 to 50 fold, preferably about 5 to 50 fold, preferably about 10 to 50 fold, induction of annexin binding relative to untreated cell in an annexin binding assay.

Antibody "effector functions" refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g., B cell receptor); and B cell activation.

"Antibody-dependent cell-mediated cytotoxicity" or "ADCC" refers to a form of cytotoxicity in which secreted Ig bound onto Fc receptors (FcRs) present on certain cytotoxic cells (e.g., Natural Killer (NK) cells, neutrophils, and macrophages) enable these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. The antibodies "arm" the cytotoxic cells and are absolutely required for such killing. The primary cells for mediating ADCC, NK cells, express Fc γ RIII only, whereas monocytes express Fc γ RI, Fc γ RII and Fc γ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an in vitro ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest

may be assessed in vivo, e.g., in a animal model such as that disclosed in Clynes et al. (USA) 95:652-656 (1998).

"Fc receptor" or "FcR" describes a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors. FcγRII receptors include FcγRIIA (an "activating receptor") and FcγRIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see review M. in Daëron, Annu. Rev. Immunol. 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, Annu. Rev. Immunol. 9:457-492 (1991); Capel et al., Immunomethods 4:25-34 (1994); and de Haas et al., J. Lab. Clin. Med. 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., J. Immunol. 117:587 (1976) and Kim et al., J. Immunol. 24:249 (1994)).

"Human effector cells" are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least FcγRIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source, e.g., from blood.

"Complement dependent cytotoxicity" or "CDC" refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system (C1q) to antibodies (of the appropriate subclass) which are bound to their cognate antigen. To assess complement activation, a CDC assay, e.g., as described in Gazzano-Santoro et al., J. Immunol. Methods 202:163 (1996), may be performed.

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated increase in cell number (generally referred to herein as cell growth), which can be due to abnormal increase in cell proliferation, abnormal decrease of cell death, or an imbalance of amounts of cell proliferation and cell death. Examples of cancer include, but are not limited to, hematopoietic cancers or blood-related cancers, such as lymphoma, leukemia, myeloma or lymphoid malignancies, but also cancers of the spleen and cancers of the lymph nodes.

The term "non-Hodgkin's lymphoma" or "NHL", as used herein, refers to a cancer of the lymphatic system other than Hodgkin's lymphomas. Hodgkin's lymphomas can generally be distinguished from non-Hodgkin's lymphomas by the presence of Reed-Sternberg cells in Hodgkin's lymphomas and the absence of said cells in non-Hodgkin's lymphomas. Examples of non-Hodgkin's

lymphomas encompassed by the term as used herein include any that would be identified as such by one skilled in the art (e.g., an oncologist or pathologist) in accordance with classification schemes known in the art, such as the Revised European-American Lymphoma (REAL) scheme as described in Color Atlas of Clinical Hematology, Third Edition; A. Victor Hoffbrand and John E. Pettit (eds.) (Harcourt Publishers Limited 2000) (see, in particular Fig. 11.57, 11.58 and/or 11.59). More specific examples include, but are not limited to, relapsed or refractory NHL, front line low grade NHL, Stage III/IV NHL, chemotherapy resistant NHL, precursor B lymphoblastic leukemia and/or lymphoma, small lymphocytic lymphoma, B cell chronic lymphocytic leukemia and/or prolymphocytic leukemia and/or small lymphocytic lymphoma, B-cell prolymphocytic lymphoma, immunocytoma and/or lymphoplasmacytic lymphoma, marginal zone B cell lymphoma, splenic marginal zone lymphoma, extranodal marginal zone - MALT lymphoma, nodal marginal zone lymphoma, hairy cell leukemia, plasmacytoma and/or plasma cell myeloma, low grade/follicular lymphoma, intermediate grade/follicular NHL, mantle cell lymphoma, follicle center lymphoma (follicular), intermediate grade diffuse NHL, diffuse large B-cell lymphoma, aggressive NHL (including aggressive front-line NHL and aggressive relapsed NHL), NHL relapsing after or refractory to autologous stem cell transplantation, primary mediastinal large B-cell lymphoma, primary effusion lymphoma, high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, Burkitt's lymphoma, precursor (peripheral) T-cell lymphoblastic leukemia and/or lymphoma, adult T-cell lymphoma and/or leukemia, T cell chronic lymphocytic leukemia and/or prolymphocytic leukemia, large granular lymphocytic leukemia, mycosis fungoides and/or Sezary syndrome, extranodal natural killer/T-cell (nasal type) lymphoma, enteropathy type T-cell lymphoma, hepatosplenic T-cell lymphoma, subcutaneous panniculitis like T-cell lymphoma, skin (cutaneous) lymphomas, anaplastic large cell lymphoma, angiocentric lymphoma, intestinal T cell lymphoma, peripheral T-cell (not otherwise specified) lymphoma and angioimmunoblastic T-cell lymphoma.

The terms "cell proliferative disorder" and "proliferative disorder" refer to disorders that are associated with some degree of abnormal cell proliferation. In one embodiment, the cell proliferative disorder is cancer.

"Tumor", as used herein, refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. The terms "cancer", "cancerous", "cell proliferative disorder", "proliferative disorder" and "tumor" are not mutually exclusive as referred to herein.

An antibody, oligopeptide or other organic small molecule which "induces cell death" is one which causes a viable cell to become nonviable. The cell is one which expresses a UNQ733 polypeptide and is of a cell type which specifically expresses or overexpresses a UNQ733 polypeptide. The cell may be cancerous or normal cells of the particular cell type. The UNQ733 polypeptide may be a transmembrane polypeptide expressed on the surface of a cancer cell or may be a polypeptide that is produced and secreted by a cancer cell. The cell may be a cancer cell, e.g., a B cell or T cell. Cell

death *in vitro* may be determined in the absence of complement and immune effector cells to distinguish cell death induced by antibody-dependent cell-mediated cytotoxicity (ADCC) or complement dependent cytotoxicity (CDC). Thus, the assay for cell death may be performed using heat inactivated serum (i.e., in the absence of complement) and in the absence of immune effector cells. To determine whether the antibody, oligopeptide or other organic small molecule is able to induce cell death, loss of membrane integrity as evaluated by uptake of propidium iodide (PI), trypan blue (see Moore et al. Cytotechnology 17:1-11 (1995)) or 7AAD can be assessed relative to untreated cells. Preferred cell death-inducing antibodies, oligopeptides or other organic small molecules are those which induce PI uptake in the PI uptake assay in BT474 cells.

A "UNQ733 polypeptide-expressing cell" is a cell which expresses an endogenous or transfected UNQ733 polypeptide, for example in a secreted form. UNQ733 polypeptide expression may be determined in a detection or prognostic assay by evaluating levels of the UNQ733 protein present in and/or on the surface of a cell, and/or secreted by the cell (e.g., via an immunohistochemistry assay using anti-UNQ733 polypeptide antibodies prepared against an isolated UNQ733 polypeptide which may be prepared using recombinant DNA technology from an isolated nucleic acid encoding the UNQ733 polypeptide; FACS analysis, etc.). Alternatively, or additionally, one may measure levels of UNQ733 polypeptide-encoding nucleic acid or mRNA in the cell, e.g., via fluorescent *in situ* hybridization using a nucleic acid based probe corresponding to a UNQ733 polypeptide-encoding nucleic acid or the complement thereof; (FISH; see WO98/45479 published October, 1998), Southern blotting, Northern blotting, or polymerase chain reaction (PCR) techniques, such as real time quantitative PCR (RT-PCR). One may also study UNQ733 polypeptide expression by measuring shed antigen in a biological fluid such as serum, e.g, using antibody-based assays (see also, e.g., U.S. Patent No. 4,933,294 issued June 12, 1990; WO91/05264 published April 18, 1991; U.S. Patent 5,401,638 issued March 28, 1995; and Sias et al., J. Immunol. Methods 132:73-80 (1990)). Aside from the above assays, various *in vivo* assays are available to the skilled practitioner. For example, one may expose cells within the body of the patient to an antibody which is optionally labeled with a detectable label, e.g., a radioactive isotope, and binding of the antibody to cells in the patient can be evaluated, e.g., by external scanning for radioactivity or by analyzing a biopsy taken from a patient previously exposed to the antibody.

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain

sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody, oligopeptide, or other organic or inorganic small molecule so as to generate a "labeled" antibody, oligopeptide, or other organic or inorganic small molecule. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g., At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu), chemotherapeutic agents e.g. methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents, enzymes and fragments thereof such as nucleolytic enzymes, antibiotics, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof, and the various antitumor or anticancer agents disclosed below. Other cytotoxic agents are described below. A tumoricidal agent causes destruction of tumor cells.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scoplectin, and 9-aminocamptothecin); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e. g., calicheamicin, especially calicheamicin gammaII and calicheamicin omegaII (see, e.g., Agnew, *Chem Intl. Ed. Engl.*, 33: 183-186 (1994)); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin

chromophore and related chromoprotein enediyne antiobiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitiostane, testolactone; anti-adrenals such as aminogluthethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); thiotepa; taxoids, *e.g.*, TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE™ Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Illinois), and TAXOTERE® doxetaxel (Rhône-Poulenc Rorer, Antony, France); chloranbucil; gemcitabine (GEMZAR®); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); oxaliplatin; leucovovin; vinorelbine (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine (XELODA®); pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovovin.

Also included in this definition are anti-hormonal agents that act to regulate, reduce, block, or inhibit the effects of hormones that can promote the growth of cancer, and are often in the form of systemic, or whole-body treatment. They may be hormones themselves. Examples include anti-

estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), EVISTA® raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® toremifene; anti-progesterones; estrogen receptor down-regulators (ERDs); agents that function to suppress or shut down the ovaries, for example, leutinizing hormone-releasing hormone (LHRH) agonists such as LUPRON® and ELIGARD® leuprolide acetate, goserelin acetate, buserelin acetate and triptorelin; other anti-androgens such as flutamide, nilutamide and bicalutamide; and aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN® exemestane, formestane, fadrozole, RIVISOR® vorozole, FEMARA® letrozole, and ARIMIDEX® anastrozole. In addition, such definition of chemotherapeutic agents includes bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), DIDROCAL® etidronate, NE-58095, ZOMETA® zoledronic acid/zoledronate, FOSAMAX® alendronate, AREDIA® pamidronate, SKELID® tiludronate, or ACTONEL® risedronate; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as THERATOPE® vaccine and gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; LURTOTECAN® topoisomerase 1 inhibitor; ABARELIX® rmRH; lapatinib ditosylate (an ErbB-2 and EGFR dual tyrosine kinase small-molecule inhibitor also known as GW572016); and pharmaceutically acceptable salts, acids or derivatives of any of the above.

A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a UNQ733 polypeptide-expressing non-Hodgkin's lymphoma cell, either *in vitro* or *in vivo*. Thus, the growth inhibitory agent may be one which significantly reduces the percentage of UNQ733 polypeptide-expressing cells in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxanes, and topoisomerase II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in The Molecular Basis of Cancer, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami et al. (WB Saunders: Philadelphia, 1995), especially p. 13. The taxanes (paclitaxel and docetaxel) are anticancer drugs both derived from the yew tree. Docetaxel (TAXOTERE®, Rhone-Poulenc Rorer), derived from the European yew, is a semisynthetic analogue of paclitaxel (TAXOL®, Bristol-Myers Squibb). Paclitaxel and docetaxel promote the assembly of

microtubules from tubulin dimers and stabilize microtubules by preventing depolymerization, which results in the inhibition of mitosis in cells.

"Doxorubicin" is an anthracycline antibiotic. The full chemical name of doxorubicin is (8S-cis)-10-[(3-amino-2,3,6-trideoxy- α -L-lyxo-hexapyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-5,12-naphthacenedione.

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor (VEGF); integrin; thrombopoietin (TPO); nerve growth factors such as NGF- β ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon - α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1a, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12; a tumor necrosis factor such as TNF- α or TNF- β ; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

Table 1

```

/*
 *
 * C-C increased from 12 to 15
5  * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M      -8      /* value of a match with a stop */

10 int      _day[26][26] = {
/*      A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */   { 2, 0,-2, 0, 0,-4, 1,-1,-1, 0,-1,-2,-1, 0,_M, 1, 0,-2, 1, 1, 0, 0,-6, 0,-3, 0},
/* B */   { 0, 3,-4, 3, 2,-5, 0, 1,-2, 0, 0,-3,-2, 2,_M,-1, 1, 0, 0, 0, 0,-2,-5, 0,-3, 1},
15 /* C */   {-2,-4,15,-5,-5,-4,-3,-3,-2, 0,-5,-6,-5,-4,_M,-3,-5,-4, 0,-2, 0,-2,-8, 0, 0,-5},
/* D */   { 0, 3,-5, 4, 3,-6, 1, 1,-2, 0, 0,-4,-3, 2,_M,-1, 2,-1, 0, 0, 0,-2,-7, 0,-4, 2},
/* E */   { 0, 2,-5, 3, 4,-5, 0, 1,-2, 0, 0,-3,-2, 1,_M,-1, 2,-1, 0, 0, 0,-2,-7, 0,-4, 3},
/* F */   {-4,-5,-4,-6,-5, 9,-5,-2, 1, 0,-5, 2, 0,-4,_M,-5,-5,-4,-3,-3, 0,-1, 0, 0, 7,-5},
/* G */   { 1, 0,-3, 1, 0,-5, 5,-2,-3, 0,-2,-4,-3, 0,_M,-1,-1,-3, 1, 0, 0,-1,-7, 0,-5, 0},
20 /* H */   {-1, 1,-3, 1, 1,-2,-2, 6,-2, 0, 0,-2,-2, 2,_M, 0, 3, 2,-1,-1, 0,-2,-3, 0, 0, 2},
/* I */   {-1,-2,-2,-2,-2, 1,-3,-2, 5, 0,-2, 2, 2,-2,_M,-2,-2,-2,-1, 0, 0, 4,-5, 0,-1,-2},
/* J */   { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,_M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */   {-1, 0,-5, 0, 0,-5,-2, 0,-2, 0, 5,-3, 0, 1,_M,-1, 1, 3, 0, 0, 0,-2,-3, 0,-4, 0},
/* L */   {-2,-3,-6,-4,-3, 2,-4,-2, 2, 0,-3, 6, 4,-3,_M,-3,-2,-3,-3,-1, 0, 2,-2, 0,-1,-2},
25 /* M */   {-1,-2,-5,-3,-2, 0,-3,-2, 2, 0, 0, 4, 6,-2,_M,-2,-1, 0,-2,-1, 0, 2,-4, 0,-2,-1},
/* N */   { 0, 2,-4, 2, 1,-4, 0, 2,-2, 0, 1,-3,-2, 2,_M,-1, 1, 0, 1, 0, 0,-2,-4, 0,-2, 1},
/* O */   {_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,
0,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M},
/* P */   { 1,-1,-3,-1,-1,-5,-1, 0,-2, 0,-1,-3,-2,-1,_M, 6, 0, 0, 1, 0, 0,-1,-6, 0,-5, 0},
30 /* Q */   { 0, 1,-5, 2, 2,-5,-1, 3,-2, 0, 1,-2,-1, 1,_M, 0, 4, 1,-1,-1, 0,-2,-5, 0,-4, 3},
/* R */   {-2, 0,-4,-1,-1,-4,-3, 2,-2, 0, 3,-3, 0, 0,_M, 0, 1, 6, 0,-1, 0,-2, 2, 0,-4, 0},
/* S */   { 1, 0, 0, 0, 0,-3, 1,-1,-1, 0, 0,-3,-2, 1,_M, 1,-1, 0, 2, 1, 0,-1,-2, 0,-3, 0},
/* T */   { 1, 0,-2, 0, 0,-3, 0,-1, 0, 0, 0,-1,-1, 0,_M, 0,-1,-1, 1, 3, 0, 0,-5, 0,-3, 0},
/* U */   { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,_M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
35 /* V */   { 0,-2,-2,-2,-2,-1,-1,-2, 4, 0,-2, 2, 2,-2,_M,-1,-2,-2,-1, 0, 0, 4,-6, 0,-2,-2},
/* W */   {-6,-5,-8,-7,-7, 0,-7,-3,-5, 0,-3,-2,-4,-4,_M,-6,-5, 2,-2,-5, 0,-6,17, 0, 0,-6},
/* X */   { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,_M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */   {-3,-3, 0,-4,-4, 7,-5, 0,-1, 0,-4,-1,-2,-2,_M,-5,-4,-4,-3,-3, 0,-2, 0, 0,10,-4},
40 /* Z */   { 0, 1,-5, 2, 3,-5, 0, 2,-2, 0, 0,-2,-1, 1,_M, 0, 3, 0, 0, 0, 0,-2,-6, 0,-4, 4}
};

```

45

50

55

Table 1 (cont')

```

/*
*/
#include <stdio.h>
5  #include <ctype.h>
    #define MAXJMP      16      /* max jumps in a diag */
    #define MAXGAP      24      /* don't continue to penalize gaps larger than this */
    #define JMPS        1024    /* max jmps in an path */
    #define MX          4       /* save if there's at least MX-1 bases since last jmp */
10  #define DMAT         3       /* value of matching bases */
    #define DMIS         0       /* penalty for mismatched bases */
    #define DINS0        8       /* penalty for a gap */
    #define DINS1        1       /* penalty per base */
    #define PINS0        8       /* penalty for a gap */
15  #define PINS1        4       /* penalty per residue */

struct jmp {
    short      n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
20 }; /* limits seq to 2^16 -1 */

struct diag {
    int      score; /* score at last jmp */
    long     offset; /* offset of prev block */
25  short    ijmp; /* current jmp index */
    struct jmp jp; /* list of jmps */
};

struct path {
30  int      spc; /* number of leading spaces */
    short    n[JMPS]; /* size of jmp (gap) */
    int      x[JMPS]; /* loc of jmp (last elem before gap) */
};

35  char      *ofile; /* output file name */
    char      *namex[2]; /* seq names: getseqs() */
    char      *prog; /* prog name for err msgs */
    char      *seqx[2]; /* seqs: getseqs() */
    int      dmax; /* best diag: nw() */
40  int      dmax0; /* final diag */
    int      dna; /* set if dna: main() */
    int      endgaps; /* set if penalizing end gaps */
    int      gapx, gapy; /* total gaps in seqs */
    int      len0, len1; /* seq lens */
45  int      ngapx, ngapy; /* total size of gaps */
    int      smax; /* max score: nw() */
    int      *xbm; /* bitmap for matching */
    long     offset; /* current offset in jmp file */
    struct diag *dx; /* holds diagonals */
50  struct path pp[2]; /* holds path for seqs */

char      *calloc(), *malloc(), *index(), *strcpy();
char      *getseq(), *g_calloc();

```

Table 1 (cont')

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
5  * where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
10 * Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
15 #include "nw.h"
#include "day.h"
static _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};
20 static _pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
25 };
main(ac, av)
main
{
    int    ac;
    char   *av[];
30
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;
40
    endgaps = 0;          /* 1 to penalize endgaps */
    ofile = "align.out";  /* output file */

    nw();                 /* fill in the matrix, get the possible jumps */
    readjumps();          /* get the actual jumps */
    print();              /* print stats, alignment */
50
    cleanup(0);           /* unlink any tmp files */
}

```

Table 1 (cont')

```

/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
5  * When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw()
10 {
    char      *px, *py;          /* seqs and ptrs */
    int       *ndely, *dely;     /* keep track of dely */
    int       ndelx, delx;       /* keep track of delx */
    int       *tmp;              /* for swapping row0, row1 */
15  int       mis;               /* score for each type */
    int       ins0, ins1;        /* insertion penalties */
    register  id;                /* diagonal index */
    register  ij;                /* jmp index */
    register  *col0, *col1;      /* score for curr, last row */
    register  xx, yy;            /* index into seqs */
20  dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));
    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
25  col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;
    smax = -10000;
    if (endgaps) {
30      for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
          col0[yy] = dely[yy] = col0[yy-1] - ins1;
          ndely[yy] = yy;
      }
      col0[0] = 0;      /* Waterman Bull Math Biol 84 */
35  }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
40  for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
45          if (xx == 1)
              col1[0] = delx = -(ins0+ins1);
          else
              col1[0] = delx = col0[0] - ins1;
          ndelx = xx;
50      }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
55      }
    }

```

Table 1 (cont')

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
    else
        mis += _day[*px-'A'][*py-'A'];
    /* update penalty for del in x seq;
    * favor new del over ongong del
    * ignore MAXGAP if weighting endgaps
    */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0+ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
    * favor new del over ongong del
    */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
    * mis over any del and delx over dely
    */
    id = xx - yy + len1 - 1;
    if (mis >= delx && mis >= dely[yy])
        col1[yy] = mis;

```

...nw

Table 1 (cont')

```

else if (delx >= dely[yy]) {
    col1[yy] = delx;
    ij = dx[id].ijmp;
5    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejumps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
20    col1[yy] = dely[yy];
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejumps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = -ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
35 if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        col1[yy] -= ins0+ins1*(len1-yy);
    if (col1[yy] > smax) {
        smax = col1[yy];
        dmax = id;
    }
}
45 }
if (endgaps && xx < len0)
    col1[yy-1] -= ins0+ins1*(len0-xx);
if (col1[yy-1] > smax) {
50     smax = col1[yy-1];
    dmax = id;
}

tmp = col0; col0 = col1; col1 = tmp;
}

55 (void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)col1);
}

```

Table 1 (cont')

```

/*
 *
 * print() -- only routine visible outside this module
 *
5  * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
10 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */
15 #include "nw.h"
#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */
extern _day[26][26];
20 int olen; /* set output line length */
FILE *fx; /* output file */

print()
{
25 int lx, ly, firstgap, lastgap; /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
30    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
35    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
40    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
45    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
50    lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

print

Table 1 (cont')

```

/*
 * trace back the best path, count matches
 */
5  static
   getmat(lx, ly, firstgap, lastgap)                                     getmat
       int    lx, ly;                                     /* "core" (minus endgaps) */
       int    firstgap, lastgap;                          /* leading trailing overlap */
   {
10      int      nm, i0, i1, siz0, siz1;
       char      outx[32];
       double    pct;
       register  n0, n1;
       register char *p0, *p1;
15      /* get total matches, score
       */
       i0 = i1 = siz0 = siz1 = 0;
       p0 = seqx[0] + pp[1].spc;
       p1 = seqx[1] + pp[0].spc;
20      n0 = pp[1].spc + 1;
       n1 = pp[0].spc + 1;
       nm = 0;
       while ( *p0 && *p1 ) {
           if (siz0) {
25                 p1++;
                     n1++;
                     siz0--;
           }
           else if (siz1) {
30                 p0++;
                     n0++;
                     siz1--;
           }
           else {
35                 if (xbm[*p0-'A']&xbm[*p1-'A'])
                     nm++;
                     if (n0++ == pp[0].x[i0])
                         siz0 = pp[0].n[i0++];
                     if (n1++ == pp[1].x[i1])
                         siz1 = pp[1].n[i1++];
40                 p0++;
                     p1++;
           }
       }
45      /* pct homology:
       * if penalizing endgaps, base is the shorter seq
       * else, knock off overhangs and take shorter core
       */
       if (endgaps)
50         lx = (len0 < len1)? len0 : len1;
       else
           lx = (lx < ly)? lx : ly;
       pct = 100.*(double)nm/(double)lx;
       fprintf(fx, "\n");
55      fprintf(fx, "<%d match%s in an overlap of %d: %.2f percent similarity\n",
           nm, (nm == 1)? "" : "es", lx, pct);

```

Table 1 (cont')

```

fprintf(fx, "<gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
5         ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);
    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
10         ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
15         "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
        smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
20         "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
        smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
25         "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
        firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
        lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "<endgaps not penalized\n");
}
static      nm;          /* matches in core -- for checking */
30 static      lmax;        /* lengths of stripped file names */
static      ij[2];        /* jmp index for a path */
static      nc[2];        /* number at start of current line */
static      ni[2];        /* current elem number -- for gapping */
static      siz[2];
35 static char *ps[2];      /* ptr to current element */
static char *po[2];      /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */
/*
40 * print alignment of described in struct path pp[]
*/
static
pr_align()
{
45     int      nn;          /* char count */
     int      more;
     register  i;

     for (i = 0, lmax = 0; i < 2; i++) {
50         nn = stripname(nameex[i]);
         if (nn > lmax)
             lmax = nn;
         nc[i] = 1;
         ni[i] = 1;
55         siz[i] = ij[i] = 0;
         ps[i] = seqx[i];

         po[i] = out[i];
     }

```

...getmat

pr_align

Table 1 (cont')

```

for (nn = nm = 0, more = 1; more; ) {
    for (i = more = 0; i < 2; i++) {
5         /*
           * do we have more of this sequence?
           */
           if (!*ps[i])
10                continue;
           more++;
           if (pp[i].spc) { /* leading space */
               *po[i]++ = ' ';
               pp[i].spc--;
           }
15         else if (siz[i]) { /* in a gap */
               *po[i]++ = '-';
               siz[i]--;
           }
           else { /* we're putting a seq element
20                */
               *po[i] = *ps[i];
               if (islower(*ps[i]))
                   *ps[i] = toupper(*ps[i]);
               po[i]++;
               ps[i]++;
25              /*
                 * are we at next gap for this seq?
                 */
               if (ni[i] == pp[i].x[ij[i]]) {
30                  /*
                     * we need to merge all gaps
                     * at this location
                     */
                     siz[i] = pp[i].n[ij[i]++];
                     while (ni[i] == pp[i].x[ij[i]])
35                          siz[i] += pp[i].n[ij[i]++];
                   }
                   ni[i]++;
               }
           }
40         if (++nn == olen || !more && nn) {
               dumpblock();
               for (i = 0; i < 2; i++)
                   po[i] = out[i];
45               nn = 0;
           }
       }
   }
   /*
50   * dump a block of lines, including numbers, stars: pr_align()
   */
   static
   dumpblock()
70       {
           register i;
           for (i = 0; i < 2; i++)
               *po[i]-- = '\0';
75       }
}

```

dumpblock

Table 1 (cont')

...dumpblock

```

5      (void) putc('\n', fx);
      for (i = 0; i < 2; i++) {
          if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
              if (i == 0)
                  nums(i);
              if (i == 0 && *out[1])
                  stars();
10             putline(i);
              if (i == 0 && *out[1])
                  fprintf(fx, star);
              if (i == 1)
                  nums(i);
15         }
    }
}
/*
 * put out a number line: dumpblock()
 */
20 static
nums(ix)
    int    ix;    /* index in out[] holding seq line */
{
25     char    nline[P_LINE];
    register i, j;
    register char *pn, *px, *py;
    for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
        *pn = ' ';
30     for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
35             if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
40             }
            else
                *pn = ' ';
            i++;
45         }
        *pn = '\0';
        nc[ix] = i;
        for (pn = nline; *pn; pn++)
            (void) putc(*pn, fx);
50     (void) putc('\n', fx);
}
/*
 * put out a line (name, [num], seq, [num]): dumpblock()
 */
55 static
putline(ix)
    int    ix;
    {
    putline

```

Table 1 (cont')

...putline

```

5      int          i;
      register char *px;

      for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
          (void) putc(*px, fx);
10     for (; i < lmax+P_SPC; i++)
          (void) putc(' ', fx);

      /* these count from 1:
      * ni[] is current element (from 1)
      * nc[] is number at start of current line
      */
15     for (px = out[ix]; *px; px++)
          (void) putc(*px&0x7F, fx);
      (void) putc('\n', fx);
20 }

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
25 static
stars()
{
    int          i;
    register char *p0, *p1, cx, *px;
30
    if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
        return;
    px = star;
35     for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
40         if (isalpha(*p0) && isalpha(*p1)) {
                if (xbm[*p0-'A']&xbm[*p1-'A']) {
                        cx = '*';
                        nm++;
                }
45             else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                        cx = '!';
                else
                        cx = ' ';
            }
50             else
                cx = ' ';
            *px++ = cx;
        }
        *px++ = '\n';
55     *px = '\0';
}

```

stars

Table 1 (cont')

```

/*
 * strip path or prefix from pn, return len: pr_align()
 */
5  static
   stripname(pn)                                stripname
   {
       char *pn; /* file name (may be path) */
       register char *px, *py;
10      py = 0;
       for (px = pn; *px; px++)
           if (*px == '/')
               py = px + 1;
15      if (py)
           (void) strcpy(pn, py);
       return(strlen(pn));
20  }

```

Table 1 (cont')

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
5  * g_calloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
10 #include <sys/file.h>

char *jname = "/tmp/homgXXXXXXX";          /* tmp file for jumps */
FILE *fj;
int cleanup();                             /* cleanup tmp file */
15 long lseek();
/*
 * remove any tmp file if we blow
 */
cleanup(i)                                cleanup
20     int i;
{
    if (fj)
        (void) unlink(jname);
    exit(i);
25 }
/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
30 */
char *
getseq(file, len)                        getseq
{
    char *file;      /* file name */
    int *len;        /* seq len */
35 {
    char line[1024], *pseq;
    register char *px, *py;
    int natgc, tlen;
    FILE *fp;
40     if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }
    tlen = natgc = 0;
45     while (fgets(line, 1024, fp)) {
        if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
50                 tlen++;
    }
    if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
55     }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

```

Table 1 (cont')

...getseq

```

py = pseq + 4;
*len = tlen;
5 rewind(fp);
while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
10        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
15            natgc++;
    }
}
*py++ = '\0';
*py = '\0';
20 (void) fclose(fp);
dna = natgc > (tlen/3);
return(pseq+4);
}
char *
25 g_malloc(msg, nx, sz)                                g_malloc
    char *msg; /* program, calling routine */
    int nx, sz; /* number and size of elements */
{
    char *px, *calloc();
30    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_malloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
35    }
    return(px);
}

/*
40 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
*/
readjmps()
{
    int fd = -1;
45    int siz, i0, i1;
    register i, j, xx;
    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
50            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
55        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;

```

Table 1 (cont')

...readjumps

```

    if (j < 0 && dx[dmax].offset && fj) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
    }

    else
        break;
}

if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}

if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;
        /* id = xx - yy + len1 - 1 */
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy++;
        ngapy -= siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}

else
    break;

}

/* reverse the order of jumps */
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}

for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}

if (fd >= 0)
    (void) close(fd);

if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}

```

Table 1 (cont')

```
/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)
{
    int ix;
    char *mktemp();
    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}
```

25

Table 2

UNQ733 polypeptide	XXXXXXXXXXXXXXXXXX (Length = 15 amino acids)
Comparison Protein	XXXXXXXXYYYYYYY (Length = 12 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the UNQ733 polypeptide) = 5 divided by 15 = 33.3%

Table 3

UNQ733 polypeptide	XXXXXXXXXXXX (Length = 10 amino acids)
Comparison Protein	XXXXXXXXYYYYYYZZYZ (Length = 15 amino acids)

45

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the UNQ733 polypeptide) = 5 divided by 10 = 50%

Table 4

UNQ733 DNA	NNNNNNNNNNNNNNNN (Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLLLL (Length = 16 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the UNQ733 polypeptide DNA sequence) = 6 divided by 14 = 42.9%

Table 5

UNQ733 DNA	NNNNNNNNNNNNNN (Length = 12 nucleotides)
Comparison DNA	NNNNLLLTVV (Length = 9 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the UNQ733 polypeptide DNA sequence) = 4 divided by 12 = 33.3%

II. Compositions and Methods of the Invention

A. Anti-UNQ733 Antibodies

In one embodiment, the present invention provides anti-UNQ733 polypeptide antibodies which may find use herein as therapeutic and/or diagnostic agents. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

1. Polyclonal Antibodies

Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen (especially when synthetic peptides are used) to a protein that is immunogenic in the species to be immunized. For example, the antigen can be conjugated to keyhole limpet hemocyanin

(KLH), serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor, using a bifunctional or derivatizing agent, e.g., maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

5 Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, e.g., 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later, the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later, the
10 animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

2. Monoclonal Antibodies

Monoclonal antibodies may be made using the hybridoma method first described by Kohler et
15 al., Nature, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as described above to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes
20 may be immunized *in vitro*. After immunization, lymphocytes are isolated and then fused with a myeloma cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium which medium preferably contains one or more substances that inhibit the growth or survival of the unfused,
25 parental myeloma cells (also referred to as fusion partner). For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the selective culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred fusion partner myeloma cells are those that fuse efficiently, support stable high-level
30 production of antibody by the selected antibody-producing cells, and are sensitive to a selective medium that selects against the unfused parental cells. Preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 and derivatives e.g., X63-Ag8-653 cells available from the American Type Culture Collection, Manassas, Virginia, USA.
35 Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); and Brodeur et

al., Monoclonal Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunosorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis described in Munson et al., Anal. Biochem., 107:220 (1980).

Once hybridoma cells that produce antibodies of the desired specificity, affinity, and/or activity are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, Monoclonal Antibodies: Principles and Practice, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal e.g., by i.p. injection of the cells into mice.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional antibody purification procedures such as, for example, affinity chromatography (e.g., using protein A or protein G-Sepharose) or ion-exchange chromatography, hydroxylapatite chromatography, gel electrophoresis, dialysis, etc.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce antibody protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra et al., Curr. Opinion in Immunol., 5:256-262 (1993) and Plückthun, Immunol. Revs. 130:151-188 (1992).

In a further embodiment, monoclonal antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty et al., Nature, 348:552-554 (1990). Clackson et al., Nature, 352:624-628 (1991) and Marks et al., J. Mol. Biol., 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks et al., Bio/Technology, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse et al., Nuc. Acids. Res. 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

The DNA that encodes the antibody may be modified to produce chimeric or fusion antibody polypeptides, for example, by substituting human heavy chain and light chain constant domain (C_H and C_L) sequences for the homologous murine sequences (U.S. Patent No. 4,816,567; and Morrison, et al., Proc. Natl Acad. Sci. USA, 81:6851 (1984)), or by fusing the immunoglobulin coding sequence with all or part of the coding sequence for a non-immunoglobulin polypeptide (heterologous polypeptide). The non-immunoglobulin polypeptide sequences can substitute for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

3. Human and Humanized Antibodies

The anti-UNQ733 antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeven et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies

in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity and HAMA response (human anti-
5 mouse antibody) when the antibody is intended for human therapeutic use. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable domain sequences. The human V domain sequence which is closest to that of the rodent is identified and the human framework region (FR) within it accepted for the humanized antibody (Sims et al., J. Immunol. 151:2296 (1993); Chothia et al., J. Mol. Biol.,
10 196:901 (1987)). Another method uses a particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter et al., Proc. Natl. Acad. Sci. USA, 89:4285 (1992); Presta et al., J. Immunol. 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high binding affinity for
15 the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-
20 dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for
25 the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

Various forms of a humanized anti-UNQ733 polypeptide antibody are contemplated. For example, the humanized antibody may be an antibody fragment, such as a Fab, which is optionally conjugated with one or more cytotoxic agent(s) in order to generate an immunoconjugate.

30 Alternatively, the humanized antibody may be an intact antibody, such as an intact IgG1 antibody.

As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining
35 region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array into such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g.,

Jakobovits et al., Proc. Natl. Acad. Sci. USA, 90:2551 (1993); Jakobovits et al., Nature, 362:255-258 (1993); Bruggemann et al., Year in Immuno. 7:33 (1993); U.S. Patent Nos. 5,545,806, 5,569,825, 5,591,669 (all of GenPharm); 5,545,807; and WO 97/17852.

Alternatively, phage display technology (McCafferty et al., Nature 348:552-553 [1990]) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B-cell. Phage display can be performed in a variety of formats, reviewed in, e.g., Johnson, Kevin S. and Chiswell, David J., Current Opinion in Structural Biology 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson et al., Nature, 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks et al., J. Mol. Biol. 222:581-597 (1991), or Griffith et al., EMBO J. 12:725-734 (1993). See, also, U.S. Patent Nos. 5,565,332 and 5,573,905.

As discussed above, human antibodies may also be generated by *in vitro* activated B cells (see U.S. Patents 5,567,610 and 5,229,275).

4. Antibody fragments

In certain circumstances there are advantages of using antibody fragments, rather than whole antibodies. The smaller size of the fragments allows for rapid clearance, and may lead to improved access to solid tumors.

Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al., Journal of Biochemical and Biophysical Methods 24:107-117 (1992); and Brennan et al., Science, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. Fab, Fv and ScFv antibody fragments can all be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of these fragments. Antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter et al., Bio/Technology 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Fab and F(ab')₂ fragment with increased *in vivo* half-life comprising a salvage receptor binding epitope residues are described in U.S. Patent No. 5,869,046.

Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; U.S. Patent No. 5,571,894; and U.S. Patent No. 5,587,458. Fv and sFv are the only species with intact combining sites that are devoid of constant regions; thus, they are suitable for reduced nonspecific binding during in vivo use. sFv fusion proteins may be constructed to yield fusion of an effector protein at either the amino or the carboxy terminus of an sFv. See Antibody Engineering, ed. Borrebaeck, supra. The antibody fragment may also be a "linear antibody", e.g., as described in U.S. Patent 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

5. Bispecific Antibodies

Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of a UNQ733 polypeptide as described herein. Other such antibodies may combine a UNQ733 polypeptide binding site with a binding site for another polypeptide. Alternatively, an anti-UNQ733 polypeptide antibody arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16), so as to focus and localize cellular defense mechanisms to the UNQ733 polypeptide-expressing and/or binding cell. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express and/or bind UNQ733 polypeptide. These antibodies possess a UNQ733 polypeptide binding arm and an arm which binds the cytotoxic agent (e.g., saporin, anti-interferon-α, vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g., F(ab')₂ bispecific antibodies).

WO 96/16673 describes a bispecific anti-ErbB2/anti-FcγRIII antibody and U.S. Patent No. 5,837,234 discloses a bispecific anti-ErbB2/anti-FcγRI antibody. A bispecific anti-ErbB2/Fcα antibody is shown in WO98/02463. U.S. Patent No. 5,821,337 teaches a bispecific anti-ErbB2/anti-CD3 antibody.

Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein et al., Nature 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker et al., EMBO J. 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. Preferably, the fusion is with an Ig heavy chain constant domain, comprising at least part

of the hinge, C_H2, and C_H3 regions. It is preferred to have the first heavy-chain constant region (C_H1) containing the site necessary for light chain bonding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host cell. This provides for greater flexibility in adjusting the mutual proportions of the three polypeptide fragments in
5 embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yield of the desired bispecific antibody. It is, however, possible to insert the coding sequences for two or all three polypeptide chains into a single expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios have no
10 significant affect on the yield of the desired chain combination.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from
15 unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology 121:210 (1986).

According to another approach described in U.S. Patent No. 5,731,168, the interface between a
20 pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second
25 antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies
30 have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

35 Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., Science 229:81 (1985) describe a procedure wherein intact antibodies are

proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent, sodium arsenite, to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets. Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a V_H connected to a V_L by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber et al., *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

6. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable

reagents for this purpose include iminothiolate and methyl-4-mercaptoputyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

7. Multivalent Antibodies

A multivalent antibody may be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The antibodies of the present invention can be multivalent antibodies (which are other than of the IgM class) with three or more antigen binding sites (e.g. tetravalent antibodies), which can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody. The multivalent antibody can comprise a dimerization domain and three or more antigen binding sites. The preferred dimerization domain comprises (or consists of) an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fc region. The preferred multivalent antibody herein comprises (or consists of) three to about eight, but preferably four, antigen binding sites. The multivalent antibody comprises at least one polypeptide chain (and preferably two polypeptide chains), wherein the polypeptide chain(s) comprise two or more variable domains. For instance, the polypeptide chain(s) may comprise VD1-(X1)_n-VD2-(X2)_n-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. For instance, the polypeptide chain(s) may comprise: VH-CH1-flexible linker-VH-CH1-Fc region chain; or VH-CH1-VH-CH1-Fc region chain. The multivalent antibody herein preferably further comprises at least two (and preferably four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a CL domain.

8. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, e.g., so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antibody. This may be achieved by introducing one or more amino acid substitutions in an Fc region of the antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp Med.* 176:1191-1195 (1992) and Shopes, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff et al., *Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design* 3:219-230 (1989). To increase the serum half life of the antibody, one may incorporate a salvage

receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Patent 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (e.g., IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule.

5 9. Immunoconjugates

The invention also pertains to immunoconjugates, or antibody-drug conjugates (ADC), comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, a drug, a growth inhibitory agent, a toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

10 The use of antibody-drug conjugates for the local delivery of cytotoxic or cytostatic agents, i.e. drugs to kill or inhibit tumor cells in the treatment of cancer (Syrigos and Epenetos (1999) Anticancer Research 19:605-614; Niculescu-Duvaz and Springer (1997) Adv. Drg Del. Rev. 26:151-172; U.S. patent 4,975,278) theoretically allows targeted delivery of the drug moiety to tumors, and intracellular accumulation therein, where systemic administration of these unconjugated drug agents may result in unacceptable levels of toxicity to normal cells as well as the tumor cells sought to be eliminated
15 (Baldwin et al., (1986) Lancet pp. (Mar. 15, 1986):603-05; Thorpe, (1985) "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review," in Monoclonal Antibodies '84: Biological And Clinical Applications, A. Pinchera et al. (ed.s), pp. 475-506). Maximal efficacy with minimal toxicity is sought thereby. Both polyclonal antibodies and monoclonal antibodies have been reported as useful
20 in these strategies (Rowland et al., (1986) Cancer Immunol. Immunother., 21:183-87). Drugs used in these methods include daunomycin, doxorubicin, methotrexate, and vindesine (Rowland et al., (1986) *supra*). Toxins used in antibody-toxin conjugates include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler et al (2000) Jour. of the Nat. Cancer Inst. 92(19):1573-1581; Mandler et al (2000) Bioorganic & Med. Chem. Letters
25 10:1025-1028; Mandler et al (2002) Bioconjugate Chem. 13:786-791), maytansinoids (EP 1391213; Liu et al., (1996) Proc. Natl. Acad. Sci. USA 93:8618-8623), and calicheamicin (Lode et al (1998) Cancer Res. 58:2928; Hinman et al (1993) Cancer Res. 53:3336-3342). The toxins may effect their cytotoxic and cytostatic effects by mechanisms including tubulin binding, DNA binding, or topoisomerase inhibition. Some cytotoxic drugs tend to be inactive or less active when conjugated to
30 large antibodies or protein receptor ligands.

 ZEVALIN® (ibritumomab tiuxetan, Biogen/Idec) is an antibody-radioisotope conjugate composed of a murine IgG1 kappa monoclonal antibody directed against the CD20 antigen found on the surface of normal and malignant B lymphocytes and ¹¹¹In or ⁹⁰Y radioisotope bound by a thiourea linker-chelator (Wiseman et al (2000) Eur. Jour. Nucl. Med. 27(7):766-77; Wiseman et al (2002) Blood 99(12):4336-42; Witzig et al (2002) J. Clin. Oncol. 20(10):2453-63; Witzig et al (2002) J. Clin. Oncol. 20(15):3262-69). Although ZEVALIN has activity against B-cell non-Hodgkin's Lymphoma (NHL), administration results in severe and prolonged cytopenias in most patients. MYLOTARG™

(gemtuzumab ozogamicin, Wyeth Pharmaceuticals), an antibody drug conjugate composed of a hu CD33 antibody linked to calicheamicin, was approved in 2000 for the treatment of acute myeloid leukemia by injection (Drugs of the Future (2000) 25(7):686; US Patent Nos. 4970198; 5079233; 5585089; 5606040; 5693762; 5739116; 5767285; 5773001). Cantuzumab mertansine (Immunogen, Inc.), an antibody drug conjugate composed of the huC242 antibody linked via the disulfide linker SPP to the maytansinoid drug moiety, DM1, is advancing into Phase II trials for the treatment of cancers that express CanAg, such as colon, pancreatic, gastric, and others. MLN-2704 (Millennium Pharm., BZL Biologics, Immunogen Inc.), an antibody drug conjugate composed of the anti-prostate specific membrane antigen (PSMA) monoclonal antibody linked to the maytansinoid drug moiety, DM1, is under development for the potential treatment of prostate tumors. The auristatin peptides, auristatin E (AE) and monomethylauristatin (MMAE), synthetic analogs of dolastatin, were conjugated to chimeric monoclonal antibodies cBR96 (specific to Lewis Y on carcinomas) and cAC10 (specific to CD30 on hematological malignancies) (Doronina et al (2003) Nature Biotechnology 21(7):778-784) and are under therapeutic development.

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re . Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, maytansinoids, a tricothecene, and CC1065, and the derivatives of these toxins that have toxin activity, are also contemplated herein.

Maytansine and maytansinoids

In one embodiment, an anti-UNQ733 polypeptide antibody (full length or fragments) of the invention is conjugated to one or more maytansinoid molecules.

Maytansinoids are mitototic inhibitors which act by inhibiting tubulin polymerization.

5 Maytansine was first isolated from the east African shrub *Maytenus serrata* (U.S. Patent No. 3,896,111). Subsequently, it was discovered that certain microbes also produce maytansinoids, such as maytansinol and C-3 maytansinol esters (U.S. Patent No. 4,151,042). Synthetic maytansinol and derivatives and analogues thereof are disclosed, for example, in U.S. Patent Nos. 4,137,230; 4,248,870; 4,256,746; 4,260,608; 4,265,814; 4,294,757; 4,307,016; 4,308,268; 4,308,269; 4,309,428; 10 4,313,946; 4,315,929; 4,317,821; 4,322,348; 4,331,598; 4,361,650; 4,364,866; 4,424,219; 4,450,254; 4,362,663; and 4,371,533, the disclosures of which are hereby expressly incorporated by reference.

Maytansinoid-antibody conjugates

In an attempt to improve their therapeutic index, maytansine and maytansinoids have been conjugated to antibodies specifically binding to tumor cell antigens. Immunoconjugates containing 15 maytansinoids and their therapeutic use are disclosed, for example, in U.S. Patent Nos. 5,208,020, 5,416,064 and European Patent EP 0 425 235 B1, the disclosures of which are hereby expressly incorporated by reference. Liu et al., Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996) described immunoconjugates comprising a maytansinoid designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic 20 towards cultured colon cancer cells, and showed antitumor activity in an *in vivo* tumor growth assay. Chari et al., Cancer Research 52:127-131 (1992) describe immunoconjugates in which a maytansinoid was conjugated via a disulfide linker to the murine antibody A7 binding to an antigen on human colon cancer cell lines, or to another murine monoclonal antibody TA.1 that binds the HER-2/*neu* oncogene. The cytotoxicity of the TA.1-maytansinoid conjugate was tested *in vitro* on the human breast cancer 25 cell line SK-BR-3, which expresses 3×10^5 HER-2 surface antigens per cell. The drug conjugate achieved a degree of cytotoxicity similar to the free maytansinoid drug, which could be increased by increasing the number of maytansinoid molecules per antibody molecule. The A7-maytansinoid conjugate showed low systemic cytotoxicity in mice.

Anti-UNQ733 polypeptide antibody-maytansinoid conjugates (immunoconjugates)

30 Anti-UNQ733 polypeptide antibody-maytansinoid conjugates are prepared by chemically linking an anti-UNQ733 polypeptide antibody to a maytansinoid molecule without significantly diminishing the biological activity of either the antibody or the maytansinoid molecule. An average of 3-4 maytansinoid molecules conjugated per antibody molecule has shown efficacy in enhancing cytotoxicity of target cells without negatively affecting the function or solubility of the antibody, 35 although even one molecule of toxin/antibody would be expected to enhance cytotoxicity over the use of naked antibody. Maytansinoids are well known in the art and can be synthesized by known techniques or isolated from natural sources. Suitable maytansinoids are disclosed, for example, in U.S.

Patent No. 5,208,020 and in the other patents and nonpatent publications referred to hereinabove. Preferred maytansinoids are maytansinol and maytansinol analogues modified in the aromatic ring or at other positions of the maytansinol molecule, such as various maytansinol esters.

There are many linking groups known in the art for making antibody-maytansinoid conjugates, including, for example, those disclosed in U.S. Patent No. 5,208,020 or EP Patent 0 425 235 B1, and Chari et al., Cancer Research 52:127-131 (1992). The linking groups include disulfide groups, thioether groups, acid labile groups, photolabile groups, peptidase labile groups, or esterase labile groups, as disclosed in the above-identified patents, disulfide and thioether groups being preferred.

Conjugates of the antibody and maytansinoid may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). Particularly preferred coupling agents include N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP) (Carlsson et al., Biochem. J. 173:723-737 [1978]) and N-succinimidyl-4-(2-pyridylthio)pentanoate (SPP) to provide for a disulfide linkage.

The linker may be attached to the maytansinoid molecule at various positions, depending on the type of the link. For example, an ester linkage may be formed by reaction with a hydroxyl group using conventional coupling techniques. The reaction may occur at the C-3 position having a hydroxyl group, the C-14 position modified with hydroxymethyl, the C-15 position modified with a hydroxyl group, and the C-20 position having a hydroxyl group. In a preferred embodiment, the linkage is formed at the C-3 position of maytansinol or a maytansinol analogue.

Calicheamicin

Another immunoconjugate of interest comprises an anti-UNQ733 polypeptide antibody conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. patents 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, 5,877,296 (all to American Cyanamid Company). Structural analogues of calicheamicin which may be used include, but are not limited to, γ_1^I , α_2^I , α_3^I , N-acetyl- γ_1^I , PSAG and θ_1^I (Hinman et al., Cancer Research 53:3336-3342 (1993), Lode et al., Cancer Research 58:2925-2928 (1998) and the aforementioned U.S. patents to American Cyanamid). Another anti-tumor drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily cross the

plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

Other cytotoxic agents

Other antitumor agents that can be conjugated to the anti-UNQ733 polypeptide antibodies of the invention include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. patents 5,053,394, 5,770,710, as well as esperamicins (U.S. patent 5,877,296).

Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published October 28, 1993.

The present invention further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

For selective destruction of the tumor, the antibody may comprise a highly radioactive atom. A variety of radioactive isotopes are available for the production of radioconjugated anti-UNQ733 antibodies. Examples include At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², Pb²¹² and radioactive isotopes of Lu. When the conjugate is used for detection, it may comprise a radioactive atom for scintigraphic studies, for example tc^{99m} or I¹²³, or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, mri), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

The radio- or other labels may be incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as tc^{99m} or I¹²³, Re¹⁸⁶, Re¹⁸⁸ and In¹¹¹ can be attached via a cysteine residue in the peptide. Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker et al (1978) Biochem. Biophys. Res. Commun. 80: 49-57 can be used to incorporate iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.


Conjugates of the antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-

difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science 238:1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026. The linker may be a “cleavable linker” facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker, 5
peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (Chari et al., Cancer Research 52:127-131 (1992); U.S. Patent No. 5,208,020) may be used.

The compounds of the invention expressly contemplate, but are not limited to, ADC prepared with cross-linker reagents: BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, 10
SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (e.g., from Pierce Biotechnology, Inc., Rockford, IL., U.S.A). See pages 467-498, 2003-2004 Applications Handbook and Catalog.

PREPARATION OF ANTIBODY DRUG CONJUGATES

15 In the antibody drug conjugates (ADC) of the invention, an antibody (Ab) is conjugated to one or more drug moieties (D), e.g. about 1 to about 20 drug moieties per antibody, through a linker (L). The ADC of Formula I may be prepared by several routes, employing organic chemistry reactions, conditions, and reagents known to those skilled in the art, including: (1) reaction of a nucleophilic group of an antibody with a bivalent linker reagent, to form Ab-L, via a covalent bond, followed by 20
reaction with a drug moiety D; and (2) reaction of a nucleophilic group of a drug moiety with a bivalent linker reagent, to form D-L, via a covalent bond, followed by reaction with the nucleophilic group of an antibody.



25 Nucleophilic groups on antibodies include, but are not limited to: (i) N-terminal amine groups, (ii) side chain amine groups, e.g. lysine, (iii) side chain thiol groups, e.g. cysteine, and (iv) sugar hydroxyl or amino groups where the antibody is glycosylated. Amine, thiol, and hydroxyl groups are nucleophilic and capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBT esters, haloformates, 30
and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups. Certain antibodies have reducible interchain disulfides, i.e. cysteine bridges. Antibodies may be made reactive for conjugation with linker reagents by treatment with a reducing agent such as DTT (dithiothreitol). Each cysteine bridge will thus form, theoretically, two reactive thiol nucleophiles. Additional nucleophilic groups can be introduced into antibodies through

the reaction of lysines with 2-iminothiolane (Traut's reagent) resulting in conversion of an amine into a thiol.

Antibody drug conjugates of the invention may also be produced by modification of the antibody to introduce electrophilic moieties, which can react with nucleophilic substituents on the linker reagent or drug. The sugars of glycosylated antibodies may be oxidized, e.g. with periodate
5 oxidizing reagents, to form aldehyde or ketone groups which may react with the amine group of linker reagents or drug moieties. The resulting imine Schiff base groups may form a stable linkage, or may be reduced, e.g. by borohydride reagents to form stable amine linkages. In one embodiment, reaction of the carbohydrate portion of a glycosylated antibody with either galactose oxidase or sodium meta-
10 periodate may yield carbonyl (aldehyde and ketone) groups in the protein that can react with appropriate groups on the drug (Hermanson, Bioconjugate Techniques). In another embodiment, proteins containing N-terminal serine or threonine residues can react with sodium meta-periodate, resulting in production of an aldehyde in place of the first amino acid (Geoghegan & Stroh, (1992) Bioconjugate Chem. 3:138-146; US 5362852). Such aldehyde can be reacted with a drug moiety or
15 linker nucleophile.

Likewise, nucleophilic groups on a drug moiety include, but are not limited to: amine, thiol, hydroxyl, hydrazide, oxime, hydrazine, thiosemicarbazone, hydrazine carboxylate, and arylhydrazide groups capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBt esters, haloformates, and acid
20 halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups.

Alternatively, a fusion protein comprising the anti-UNQ733 polypeptide antibody and cytotoxic agent may be made, e.g., by recombinant techniques or peptide synthesis. The length of DNA may comprise respective regions encoding the two portions of the conjugate either adjacent one
25 another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

In yet another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pre-targeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then
30 administration of a "ligand" (e.g., avidin) which is conjugated to a cytotoxic agent (e.g., a radionucleotide).

10. Immunoliposomes

The anti-UNQ733 polypeptide antibodies disclosed herein may also be formulated as
35 immunoliposomes. A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological

membranes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA 82:3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77:4030 (1980); U.S. Pat. Nos. 4,485,045 and 4,544,545; and WO97/38731 published October 23, 1997. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem. 257:286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon et al., J. National Cancer Inst. 81(19):1484 (1989).

B. UNQ733 polypeptide Binding Oligopeptides

UNQ733 polypeptide binding oligopeptides of the invention are oligopeptides that bind, preferably specifically, to a UNQ733 polypeptide as described herein. UNQ733 polypeptide binding oligopeptides may be chemically synthesized using known oligopeptide synthesis methodology or may be prepared and purified using recombinant technology. UNQ733 polypeptide binding oligopeptides are usually at least about 5 amino acids in length, alternatively at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 amino acids in length or more, wherein such oligopeptides that are capable of binding, preferably specifically, to a UNQ733 polypeptide as described herein. UNQ733 polypeptide binding oligopeptides may be identified without undue experimentation using well known techniques. In this regard, it is noted that techniques for screening oligopeptide libraries for oligopeptides that are capable of specifically binding to a polypeptide target are well known in the art (see, e.g., U.S. Patent Nos. 5,556,762, 5,750,373, 4,708,871, 4,833,092, 5,223,409, 5,403,484, 5,571,689, 5,663,143; PCT Publication Nos. WO 84/03506 and WO84/03564; Geysen et al., Proc. Natl. Acad. Sci. U.S.A., 81:3998-4002 (1984); Geysen et al., Proc. Natl. Acad. Sci. U.S.A., 82:178-182 (1985); Geysen et al., in Synthetic Peptides as Antigens, 130-149 (1986); Geysen et al., J. Immunol. Meth., 102:259-274 (1987); Schoofs et al., J. Immunol., 140:611-616 (1988), Cwirla, S. E. et al. (1990) Proc. Natl. Acad. Sci. USA, 87:6378; Lowman, H.B. et al. (1991) Biochemistry, 30:10832; Clackson, T. et al. (1991) Nature, 352: 624; Marks, J. D. et al. (1991), J. Mol. Biol., 222:581; Kang, A.S. et al. (1991) Proc. Natl. Acad. Sci. USA, 88:8363, and Smith, G. P. (1991) Current Opin. Biotechnol., 2:668).

In this regard, bacteriophage (phage) display is one well known technique which allows one to screen large oligopeptide libraries to identify member(s) of those libraries which are capable of specifically binding to a polypeptide target. Phage display is a technique by which variant

polypeptides are displayed as fusion proteins to the coat protein on the surface of bacteriophage particles (Scott, J.K. and Smith, G. P. (1990) Science, 249: 386). The utility of phage display lies in the fact that large libraries of selectively randomized protein variants (or randomly cloned cDNAs) can be rapidly and efficiently sorted for those sequences that bind to a target molecule with high affinity.

5 Display of peptide (Cwirla, S. E. et al. (1990) Proc. Natl. Acad. Sci. USA, 87:6378) or protein (Lowman, H.B. et al. (1991) Biochemistry, 30:10832; Clackson, T. et al. (1991) Nature, 352: 624; Marks, J. D. et al. (1991), J. Mol. Biol., 222:581; Kang, A.S. et al. (1991) Proc. Natl. Acad. Sci. USA, 88:8363) libraries on phage have been used for screening millions of polypeptides or oligopeptides for ones with specific binding properties (Smith, G. P. (1991) Current Opin. Biotechnol., 2:668). Sorting

10 phage libraries of random mutants requires a strategy for constructing and propagating a large number of variants, a procedure for affinity purification using the target receptor, and a means of evaluating the results of binding enrichments. U.S. Patent Nos. 5,223,409, 5,403,484, 5,571,689, and 5,663,143.

Although most phage display methods have used filamentous phage, lambdoid phage display systems (WO 95/34683; U.S. 5,627,024), T4 phage display systems (Ren et al., Gene, 215: 439 (1998);

15 Zhu et al., Cancer Research, 58(15): 3209-3214 (1998); Jiang et al., Infection & Immunity, 65(11): 4770-4777 (1997); Ren et al., Gene, 195(2):303-311 (1997); Ren, Protein Sci., 5: 1833 (1996); Efimov et al., Virus Genes, 10: 173 (1995)) and T7 phage display systems (Smith and Scott, Methods in Enzymology, 217: 228-257 (1993); U.S. 5,766,905) are also known.

Many other improvements and variations of the basic phage display concept have now been

20 developed. These improvements enhance the ability of display systems to screen peptide libraries for binding to selected target molecules and to display functional proteins with the potential of screening these proteins for desired properties. Combinatorial reaction devices for phage display reactions have been developed (WO 98/14277) and phage display libraries have been used to analyze and control bimolecular interactions (WO 98/20169; WO 98/20159) and properties of constrained helical peptides

25 (WO 98/20036). WO 97/35196 describes a method of isolating an affinity ligand in which a phage display library is contacted with one solution in which the ligand will bind to a target molecule and a second solution in which the affinity ligand will not bind to the target molecule, to selectively isolate binding ligands. WO 97/46251 describes a method of biopanning a random phage display library with an affinity purified antibody and then isolating binding phage, followed by a micropanning process

30 using microplate wells to isolate high affinity binding phage. The use of *Staphylococcus aureus* protein A as an affinity tag has also been reported (Li et al. (1998) Mol Biotech., 9:187). WO 97/47314 describes the use of substrate subtraction libraries to distinguish enzyme specificities using a combinatorial library which may be a phage display library. A method for selecting enzymes suitable for use in detergents using phage display is described in WO 97/09446. Additional methods of

35 selecting specific binding proteins are described in U.S. Patent Nos. 5,498,538, 5,432,018, and WO 98/15833.

Methods of generating peptide libraries and screening these libraries are also disclosed in U.S. Patent Nos. 5,723,286, 5,432,018, 5,580,717, 5,427,908, 5,498,530, 5,770,434, 5,734,018, 5,698,426, 5,763,192, and 5,723,323.

C. UNQ733 polypeptide Binding small molecules

UNQ733 polypeptide binding small molecules are preferably organic molecules other than oligopeptides or antibodies as defined herein that bind, preferably specifically, to a UNQ733 polypeptide as described herein. UNQ733 polypeptide binding organic small molecules may be identified and chemically synthesized using known methodology (see, e.g., PCT Publication Nos. WO00/00823 and WO00/39585). UNQ733 polypeptide binding organic small molecules are usually less than about 2000 daltons in size, alternatively less than about 1500, 750, 500, 250 or 200 daltons in size, wherein such organic small molecules that are capable of binding, preferably specifically, to a UNQ733 polypeptide as described herein may be identified without undue experimentation using well known techniques. In this regard, it is noted that techniques for screening organic small molecule libraries for molecules that are capable of binding to a polypeptide target are well known in the art (see, e.g., PCT Publication Nos. WO00/00823 and WO00/39585). UNQ733 polypeptide binding organic small molecules may be, for example, aldehydes, ketones, oximes, hydrazones, semicarbazones, carbazides, primary amines, secondary amines, tertiary amines, N-substituted hydrazines, hydrazides, alcohols, ethers, thiols, thioethers, disulfides, carboxylic acids, esters, amides, ureas, carbamates, carbonates, ketals, thioketals, acetals, thioacetals, aryl halides, aryl sulfonates, alkyl halides, alkyl sulfonates, aromatic compounds, heterocyclic compounds, anilines, alkenes, alkynes, diols, amino alcohols, oxazolidines, oxazolines, thiazolidines, thiazolines, enamines, sulfonamides, epoxides, aziridines, isocyanates, sulfonyl chlorides, diazo compounds, acid chlorides, or the like.

D. Screening for anti-UNQ733 polypeptide antibodies, UNQ733 polypeptide binding oligopeptides and UNQ733 polypeptide binding small molecules with the desired properties

Techniques for generating antibodies, oligopeptides and small molecules that bind to UNQ733 polypeptides have been described above. One may further select antibodies, oligopeptides or other small molecules with certain biological characteristics, as desired.

The growth inhibitory effects of an anti-UNQ733 antibody, oligopeptide or other small molecule of the invention may be assessed by methods known in the art, e.g., using cells which express a UNQ733 polypeptide either endogenously or following transfection with the UNQ733 polypeptide gene. For example, appropriate tumor cell lines and UNQ733 polypeptide-transfected cells may be treated with an anti-UNQ733 polypeptide monoclonal antibody, oligopeptide or other small molecule of the invention at various concentrations for a few days (e.g., 2-7) days and stained with crystal violet or MTT or analyzed by some other colorimetric assay. Another method of measuring proliferation would be by comparing ³H-thymidine uptake by the cells treated in the presence or absence an anti-UNQ733 polypeptide antibody, UNQ733 polypeptide binding oligopeptide or UNQ733 polypeptide binding small molecule of the invention. After treatment, the cells are harvested and the amount of

radioactivity incorporated into the DNA quantitated in a scintillation counter. Appropriate positive controls include treatment of a selected cell line with a growth inhibitory antibody known to inhibit growth of that cell line. Growth inhibition of tumor cells *in vivo* can be determined in various ways known in the art. The tumor cell may be one that overexpresses a UNQ733 polypeptide. The anti-UNQ733 polypeptide antibody, UNQ733 polypeptide binding oligopeptide or UNQ733 polypeptide binding organic small molecule will inhibit cell proliferation of a UNQ733 polypeptide-expressing tumor cell *in vitro* or *in vivo* by about 25-100% compared to the untreated tumor cell, more preferably, by about 30-100%, and even more preferably by about 50-100% or 70-100%, in one embodiment, at an antibody concentration of about 0.5 to 30 µg/ml. Growth inhibition can be measured at an antibody concentration of about 0.5 to 30 µg/ml or about 0.5 nM to 200 nM in cell culture, where the growth inhibition is determined 1-10 days after exposure of the tumor cells to the antibody. The antibody is growth inhibitory *in vivo* if administration of the anti-UNQ733 polypeptide antibody at about 1 µg/kg to about 100 mg/kg body weight results in reduction in tumor size or reduction of tumor cell proliferation within about 5 days to 3 months from the first administration of the antibody, preferably within about 5 to 30 days.

To select for an anti-UNQ733 polypeptide antibody, UNQ733 polypeptide binding oligopeptide or UNQ733 polypeptide binding organic small molecule which induces cell death, loss of membrane integrity as indicated by, e.g., propidium iodide (PI), trypan blue or 7AAD uptake may be assessed relative to control. A PI uptake assay can be performed in the absence of complement and immune effector cells. UNQ733 polypeptide-expressing tumor cells are incubated with medium alone or medium containing the appropriate anti-UNQ733 polypeptide antibody (e.g., at about 10µg/ml), UNQ733 polypeptide binding oligopeptide or UNQ733 polypeptide binding organic small molecule. The cells are incubated for a 3-day time period. Following each treatment, cells are washed and aliquoted into 35 mm strainer-capped 12 x 75 tubes (1ml per tube, 3 tubes per treatment group) for removal of cell clumps. Tubes then receive PI (10µg/ml). Samples may be analyzed using a FACSCAN® flow cytometer and FACSCONVERT® CellQuest software (Becton Dickinson). Those anti-UNQ733 polypeptide antibodies, UNQ733 polypeptide binding oligopeptides or UNQ733 polypeptide binding organic small molecules that induce statistically significant levels of cell death as determined by PI uptake may be selected as cell death-inducing anti-UNQ733 polypeptide antibodies, UNQ733 polypeptide binding oligopeptides or UNQ733 polypeptide binding organic small molecules.

To screen for antibodies, oligopeptides or other organic small molecules which bind to an epitope on a UNQ733 polypeptide bound by an antibody of interest, a routine cross-blocking assay such as that described in Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. This assay can be used to determine if a test antibody, oligopeptide or other organic small molecule binds the same site or epitope as a known anti-UNQ733 polypeptide antibody. Alternatively, or additionally, epitope mapping can be performed by methods known in the art. For example, the antibody sequence can be mutagenized such as by alanine

scanning, to identify contact residues. The mutant antibody is initially tested for binding with polyclonal antibody to ensure proper folding. In a different method, peptides corresponding to different regions of a UNQ733 polypeptide can be used in competition assays with the test antibodies or with a test antibody and an antibody with a characterized or known epitope.

5 E. Antibody Dependent Enzyme Mediated Prodrug Therapy (ADEPT)

The antibodies of the present invention may also be used in ADEPT by conjugating the antibody to a prodrug-activating enzyme which converts a prodrug (e.g., a peptidyl chemotherapeutic agent, see WO81/01145) to an active anti-cancer drug. See, for example, WO 88/07378 and U.S. Patent No. 4,975,278.

10 The enzyme component of the immunoconjugate useful for ADEPT includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form.

Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting
15 non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratin protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs;
20 β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as "abzymes", can be used to convert the prodrugs of the invention into free active drugs (see, e.g., Massey, Nature 328:457-458
25 (1987)). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes of this invention can be covalently bound to the anti-UNQ733 antibodies by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an
30 antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using recombinant DNA techniques well known in the art (see, e.g., Neuberger et al., Nature 312:604-608 (1984)).

F. Anti-UNQ733 polypeptide Antibody Variants

35 In addition to the anti-UNQ733 polypeptide antibodies described herein, it is contemplated that anti-UNQ733 polypeptide antibody variants can be prepared. Anti-UNQ733 polypeptide antibody variants can be prepared by introducing appropriate nucleotide changes into the encoding DNA, and/or

by synthesis of the desired antibody. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the anti-UNQ733 polypeptide antibody, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the anti-UNQ733 polypeptide antibodies described herein can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the antibody that results in a change in the amino acid sequence as compared with the native sequence antibody or polypeptide. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the anti-UNQ733 polypeptide antibody. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the anti-UNQ733 polypeptide antibody with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the parent sequence.

Anti-UNQ733 polypeptide antibody and UNQ733 polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native antibody or protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the anti-UNQ733 antibody or UNQ733 polypeptide.

Anti-UNQ733 antibody and UNQ733 polypeptide fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating antibody or polypeptide fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired antibody or polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, anti-UNQ733 antibody and UNQ733 polypeptide fragments share at least one biological and/or immunological activity with the native anti-UNQ733 polypeptide antibody or UNQ733 polypeptide disclosed herein.

In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity,

then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table 6

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp; Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

5 Substantial modifications in function or immunological identity of the anti-UNQ733 antibody or UNQ733 polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Amino acids may be grouped according to similarities in the properties of their side chains (in A. L. Lehninger, in Biochemistry, second ed., pp. 10 73-75, Worth Publishers, New York (1975)):

(1) non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M)

(2) uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q)

(3) acidic: Asp (D), Glu (E)

(4) basic: Lys (K), Arg (R), His(H)

Alternatively, naturally occurring residues may be divided into groups based on common side-chain properties:

(1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;

(2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

(3) acidic: Asp, Glu;

(4) basic: His, Lys, Arg;

(5) residues that influence chain orientation: Gly, Pro;

(6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the anti-UNQ733 polypeptide antibody or UNQ733 polypeptide variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244:1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

Any cysteine residue not involved in maintaining the proper conformation of the anti-UNQ733 polypeptide antibody or UNQ733 polypeptide also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the anti-UNQ733 antibody or UNQ733 polypeptide to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (e.g., a humanized or human antibody). Generally,

the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (e.g., 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e.g., binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and human UNQ733 polypeptide. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

Nucleic acid molecules encoding amino acid sequence variants of the anti-UNQ733 antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the anti-UNQ733 antibody.

G. Modifications of Anti-UNQ733 Antibodies and UNQ733 polypeptides

Covalent modifications of anti-UNQ733 polypeptide antibodies and UNQ733 polypeptides are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of an anti-UNQ733 antibody or UNQ733 polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the anti-UNQ733 polypeptide antibody or UNQ733 polypeptide. Derivatization with bifunctional agents is useful, for instance, for crosslinking anti-UNQ733 antibody or UNQ733 polypeptide to a water-insoluble support matrix or surface for use in the method for purifying anti-UNQ733 antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular

Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the anti-UNQ733 antibody or UNQ733 polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the antibody or polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence anti-UNQ733 antibody or UNQ733 polypeptide (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence anti-UNQ733 antibody or UNQ733 polypeptide. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Glycosylation of antibodies and other polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-aceylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the anti-UNQ733 antibody or UNQ733 polypeptide is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original anti-UNQ733 antibody or UNQ733 polypeptide (for O-linked glycosylation sites). The anti-UNQ733 antibody or UNQ733 polypeptide amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the anti-UNQ733 antibody or UNQ733 polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the anti-UNQ733 antibody or UNQ733 polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the anti-UNQ733 antibody or UNQ733 polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of

carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of anti-UNQ733 antibody or UNQ733 polypeptide comprises linking the antibody or polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337. The antibody or polypeptide also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980).

The anti-UNQ733 antibody or UNQ733 polypeptide of the present invention may also be modified in a way to form chimeric molecules comprising an anti-UNQ733 antibody or UNQ733 polypeptide fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the anti-UNQ733 antibody or UNQ733 polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl-terminus of the anti-UNQ733 antibody or UNQ733 polypeptide. The presence of such epitope-tagged forms of the anti-UNQ733 antibody or UNQ733 polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the anti-UNQ733 antibody or UNQ733 polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the anti-UNQ733 antibody or UNQ733 polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of

an anti-UNQ733 antibody or UNQ733 polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH₂ and CH₃, or the hinge, CH₁, CH₂ and CH₃ regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

H. Preparation of Anti-UNQ733 Antibodies and UNQ733 polypeptides

The description below relates primarily to production of anti-UNQ733 antibodies and UNQ733 polypeptides by culturing cells transformed or transfected with a vector containing anti-UNQ733 antibody- and UNQ733 polypeptide-encoding nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare anti-UNQ733 antibodies and UNQ733 polypeptides. For instance, the appropriate amino acid sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the anti-UNQ733 antibody or UNQ733 polypeptide may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the desired anti-UNQ733 antibody or UNQ733 polypeptide.

1. Isolation of DNA Encoding Anti-UNQ733 Antibody or UNQ733 Polypeptide

DNA encoding anti-UNQ733 antibody or UNQ733 polypeptide may be obtained from a cDNA library prepared from tissue believed to possess the anti-UNQ733 antibody or UNQ733 polypeptide mRNA and to express it at a detectable level. Accordingly, human anti-UNQ733 antibody or UNQ733 polypeptide DNA can be conveniently obtained from a cDNA library prepared from human tissue. The anti-UNQ733 antibody- or UNQ733 polypeptide-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding anti-UNQ733 antibody or UNQ733 polypeptide is to use PCR methodology [Sambrook et al., *supra*; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

Techniques for screening a cDNA library are well known in the art. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art,

and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for anti-UNQ733 antibody or UNQ733 polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., *supra*.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., *supra*, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see

Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA* ; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan'*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan'*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

Full length antibody, antibody fragments, and antibody fusion proteins can be produced in bacteria, in particular when glycosylation and Fc effector function are not needed, such as when the therapeutic antibody is conjugated to a cytotoxic agent (e.g., a toxin) and the immunoconjugate by itself shows effectiveness in tumor cell destruction. Full-length antibodies have greater half life in circulation. Production in *E. coli* is faster and more cost efficient. For expression of antibody fragments and polypeptides in bacteria, see, e.g., U.S. 5,648,237 (Carter et al.), U.S. 5,789,199 (Joly et al.), and U.S. 5,840,523 (Simmons et al.) which describes translation initiation regio (TIR) and signal sequences for optimizing expression and secretion, these patents incorporated herein by reference. After expression, the antibody is isolated from the *E. coli* cell paste in a soluble fraction and can be purified through, e.g., a protein A or G column depending on the isotype. Final purification can be carried out similar to the process for purifying antibody expressed e.g., in CHO cells.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for anti-UNQ733 antibody- or UNQ733 polypeptide-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include

Schizosaccharomyces pombe (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., Bio/Technology, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilarum* (ATCC 36,906; Van den Berg et al., Bio/Technology, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol., 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983]; Tilburn et al., Gene, 26:205-221 [1983]; Yelton et al., Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4:475-479 [1985]). Methylotropic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylotrophs, 269 (1982).

Suitable host cells for the expression of glycosylated anti-UNQ733 antibody or UNQ733 polypeptide are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells, such as cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442);

human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

Host cells are transformed with the above-described expression or cloning vectors for anti-UNQ733 antibody or UNQ733 polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding anti-UNQ733 antibody or UNQ733 polypeptide may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The UNQ733 polypeptide may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the anti-UNQ733 antibody- or UNQ733 polypeptide-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the anti-UNQ733 antibody- or UNQ733 polypeptide-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the anti-UNQ733 antibody- or UNQ733 polypeptide-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding anti-UNQ733 antibody or UNQ733 polypeptide.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytocrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

Anti-UNQ733 antibody or UNQ733 polypeptide transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma

virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the anti-UNQ733 antibody or UNQ733 polypeptide by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the anti-UNQ733 antibody or UNQ733 polypeptide coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-UNQ733 antibody or UNQ733 polypeptide.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of anti-UNQ733 antibody or UNQ733 polypeptide in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Culturing the Host Cells

The host cells used to produce the anti-UNQ733 antibody or UNQ733 polypeptide of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham et al., Meth. Enz. 58:44 (1979), Barnes et al., Anal. Biochem. 102:255 (1980), U.S. Pat. Nos. 4,767,704; 4,657,866; 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic compounds usually present at final

concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

5. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence UNQ733 polypeptide or against a synthetic peptide based on the DNA sequence provided herein or against exogenous sequence fused to UNQ733 polypeptide DNA and encoding a specific antibody epitope.

6. Purification of Anti-UNQ733 Antibody and UNQ733 polypeptide

Forms of anti-UNQ733 antibody and UNQ733 polypeptide may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of anti-UNQ733 antibody and UNQ733 polypeptide can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify anti-UNQ733 antibody and UNQ733 polypeptide from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the anti-UNQ733 antibody and UNQ733 polypeptide. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes,

Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular anti-UNQ733 antibody or UNQ733 polypeptide produced.

When using recombinant techniques, the antibody can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Carter et al., Bio/Technology 10:163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$ or $\gamma 4$ heavy chains (Lindmark et al., J. Immunol. Meth. 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss et al., EMBO J. 5:15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_H3 domain, the Bakerbond ABXTM resin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSETM chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (e.g., from about 0-0.25M salt).

I. Pharmaceutical Formulations

Therapeutic formulations of the anti-UNQ733 antibodies, UNQ733 polypeptide binding oligopeptides, UNQ733 polypeptide binding organic or inorganic small molecules and/or UNQ733 polypeptides used in accordance with the present invention are prepared for storage by mixing the antibody, polypeptide, oligopeptide or organic/inorganic small molecule having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as acetate, Tris, phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; tonicifiers such as trehalose and sodium chloride; sugars such as sucrose, mannitol, trehalose or sorbitol; surfactant such as polysorbate; salt-forming counter-ions such as sodium; metal complexes (e.g., Zn-protein complexes); and/or non-ionic surfactants such as TWEEN®, PLURONICS® or polyethylene glycol (PEG). The antibody preferably comprises the antibody at a concentration of between 5-200 mg/ml, preferably between 10-100 mg/ml.

The formulations herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, in addition to an anti-UNQ733 antibody, UNQ733 polypeptide binding oligopeptide, or UNQ733 polypeptide binding organic or inorganic small molecule, it may be desirable to include in the one formulation, an additional antibody, e.g., a second anti-UNQ733 antibody which binds a different epitope on the UNQ733 polypeptide, or an antibody to some other target such as a growth factor that affects the growth of the particular cancer. Alternatively, or additionally, the composition may further comprise a chemotherapeutic agent, cytotoxic agent, cytokine, growth inhibitory agent, anti-hormonal agent, and/or cardioprotectant. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and

nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. Ed. (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT® (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

J. Treatment with Anti-UNQ733 Antibodies, UNQ733 polypeptide Binding Oligopeptides and UNQ733 polypeptide Binding Organic/Inorganic Small Molecules

To determine UNQ733 polypeptide expression in the cancer, various detection assays are available. In one embodiment, UNQ733 polypeptide overexpression may be analyzed by immunohistochemistry (IHC). Paraffin embedded tissue sections from a tumor biopsy may be subjected to the IHC assay and accorded a UNQ733 polypeptide staining intensity criteria as follows:

Score 0 - no staining is observed or membrane staining is observed in less than 10% of tumor cells.

Score 1+ - a faint/barely perceptible membrane staining is detected in more than 10% of the tumor cells. The cells are only stained in part of their membrane.

Score 2+ - a weak to moderate complete membrane staining is observed in more than 10% of the tumor cells.

Score 3+ - a moderate to strong complete membrane staining is observed in more than 10% of the tumor cells.

Those tumors with 0 or 1+ scores for UNQ733 polypeptide expression may be characterized as not overexpressing UNQ733 polypeptide, whereas those tumors with 2+ or 3+ scores may be characterized as overexpressing UNQ733 polypeptide.

Alternatively, or additionally, FISH assays such as the INFORM® (sold by Ventana, Arizona) or PATHVISION® (Vysis, Illinois) may be carried out on formalin-fixed, paraffin-embedded tumor tissue to determine the extent (if any) of UNQ733 polypeptide overexpression in the tumor.

UNQ733 polypeptide overexpression or amplification may be evaluated using an *in vivo* detection assay, e.g., by administering a molecule (such as an antibody, oligopeptide or organic small molecule) which binds the molecule to be detected and is tagged with a detectable label (e.g., a radioactive isotope or a fluorescent label) and externally scanning the patient for localization of the label.

As described above, the anti-UNQ733 antibodies, oligopeptides and organic small molecules of the invention have various non-therapeutic applications. The anti-UNQ733 antibodies, oligopeptides and organic/inorganic small molecules of the present invention can be useful for staging of UNQ733 polypeptide-expressing cancers (e.g., in radioimaging). The antibodies, oligopeptides and organic small molecules are also useful for purification or immunoprecipitation of UNQ733 polypeptide from cells, for detection and quantitation of UNQ733 polypeptide *in vitro*, e.g., in an ELISA or a Western blot, to kill and eliminate UNQ733 polypeptide-expressing cells from a population of mixed cells as a step in the purification of other cells.

Currently, depending on the stage of the cancer, cancer treatment involves one or a combination of the following therapies: surgery to remove the cancerous tissue, radiation therapy, and chemotherapy. Anti-UNQ733 antibody, oligopeptide or organic small molecule therapy may be especially desirable in elderly patients who do not tolerate the toxicity and side effects of chemotherapy well and in metastatic disease where radiation therapy has limited usefulness. The tumor targeting anti-UNQ733 antibodies, oligopeptides and organic/inorganic small molecules of the invention are useful to alleviate UNQ733 polypeptide-expressing cancers upon initial diagnosis of the disease or during relapse. For therapeutic applications, the anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule can be used alone, or in combination therapy with, e.g., hormones, antiangiogens, or radiolabelled compounds, or with surgery, cryotherapy, and/or radiotherapy. Anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule treatment can be administered in conjunction with other forms of conventional therapy, either consecutively with, pre- or post-conventional therapy. Chemotherapeutic drugs such as TAXOTERE® (docetaxel), TAXOL® (paclitaxel), estramustine and mitoxantrone are used in treating cancer, in particular, in good risk patients. In the present method of the invention for treating or alleviating cancer, the cancer patient can be administered anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule in conjunction with treatment with the one or more of the preceding chemotherapeutic agents. In particular, combination therapy with paclitaxel and modified derivatives (see, e.g., EP0600517) is contemplated. The anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule will be administered with a therapeutically effective dose of the chemotherapeutic agent. In another embodiment, the anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule is administered in conjunction with chemotherapy to enhance the activity and efficacy of the chemotherapeutic agent, e.g., paclitaxel. The Physicians' Desk Reference (PDR) discloses dosages of these agents that have been used in treatment of various cancers. The dosing regimen and dosages of these aforementioned chemotherapeutic drugs that are therapeutically effective will depend on the particular cancer being treated, the extent of the disease and other factors familiar to the physician of skill in the art and can be determined by the physician.

In one particular embodiment, a conjugate comprising an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule conjugated with a cytotoxic agent is administered to the patient.

Preferably, the immunoconjugate bound to the UNQ733 protein is internalized by the cell, resulting in increased therapeutic efficacy of the immunoconjugate in killing the cancer cell to which it binds. In a preferred embodiment, the cytotoxic agent targets or interferes with the nucleic acid in the cancer cell. Examples of such cytotoxic agents are described above and include maytansinoids, calicheamicins, ribonucleases and DNA endonucleases.

The anti-UNQ733 antibodies, oligopeptides, organic/inorganic small molecules or toxin conjugates thereof are administered to a human patient, in accord with known methods, such as intravenous administration, e.g., as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous or subcutaneous administration of the antibody, oligopeptide or organic small molecule is preferred.

Other therapeutic regimens may be combined with the administration of the anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule. The combined administration includes co-administration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities. Preferably such combined therapy results in a synergistic therapeutic effect.

It may also be desirable to combine administration of the anti-UNQ733 antibody or antibodies, oligopeptides or organic/inorganic small molecules, with administration of an antibody directed against another tumor antigen associated with the particular cancer.

In another embodiment, the therapeutic treatment methods of the present invention involves the combined administration of an anti-UNQ733 antibody (or antibodies), oligopeptides or organic/inorganic small molecules and one or more chemotherapeutic agents or growth inhibitory agents, including co-administration of cocktails of different chemotherapeutic agents. Chemotherapeutic agents include estramustine phosphate, prednimustine, cisplatin, 5-fluorouracil, melphalan, cyclophosphamide, hydroxyurea and hydroxyureataxanes (such as paclitaxel and doxorubicin) and/or anthracycline antibiotics. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in Chemotherapy Service Ed., M.C. Perry, Williams & Wilkins, Baltimore, MD (1992).

The antibody, oligopeptide or organic/inorganic small molecule may be combined with an anti-hormonal compound; e.g., an anti-estrogen compound such as tamoxifen; an anti-progesterone such as onapristone (see, EP 616 812); or an anti-androgen such as flutamide, in dosages known for such molecules. Where the cancer to be treated is androgen independent cancer, the patient may previously have been subjected to anti-androgen therapy and, after the cancer becomes androgen independent, the anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule (and optionally other agents as described herein) may be administered to the patient.

Sometimes, it may be beneficial to also co-administer a cardioprotectant (to prevent or reduce myocardial dysfunction associated with the therapy) or one or more cytokines to the patient. In addition to the above therapeutic regimes, the patient may be subjected to surgical removal of cancer cells and/or radiation therapy, before, simultaneously with, or post antibody, oligopeptide or organic/inorganic small molecule therapy. Suitable dosages for any of the above co-administered agents are those presently used and may be lowered due to the combined action (synergy) of the agent and anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule.

For the prevention or treatment of disease, the dosage and mode of administration will be chosen by the physician according to known criteria. The appropriate dosage of antibody, oligopeptide or organic/inorganic small molecule will depend on the type of disease to be treated, as defined above, the severity and course of the disease, whether the antibody, oligopeptide or organic/inorganic small molecule is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, oligopeptide or organic/inorganic small molecule, and the discretion of the attending physician. The antibody, oligopeptide or organic/inorganic small molecule is suitably administered to the patient at one time or over a series of treatments. Preferably, the antibody, oligopeptide or organic/inorganic small molecule is administered by intravenous infusion or by subcutaneous injections. Depending on the type and severity of the disease, about 1 µg/kg to about 50 mg/kg body weight (e.g., about 0.1-15mg/kg/dose) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A dosing regimen can comprise administering an initial loading dose of about 4 mg/kg, followed by a weekly maintenance dose of about 2 mg/kg of the anti-UNQ733 antibody. However, other dosage regimens may be useful. A typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. The progress of this therapy can be readily monitored by conventional methods and assays and based on criteria known to the physician or other persons of skill in the art.

Aside from administration of the antibody protein to the patient, the present application contemplates administration of the antibody by gene therapy. Such administration of nucleic acid encoding the antibody is encompassed by the expression "administering a therapeutically effective amount of an antibody". See, for example, WO96/07321 published March 14, 1996 concerning the use of gene therapy to generate intracellular antibodies.

There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells; *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the site where the antibody is required. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells and the modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes

which are implanted into the patient (see, e.g., U.S. Patent Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. A commonly used vector for *ex vivo* delivery of the gene is a retroviral vector.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or adeno-associated virus) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are DOTMA, DOPE and DC-Chol, for example). For review of the currently known gene marking and gene therapy protocols see Anderson et al., Science 256:808-813 (1992). See also WO 93/25673 and the references cited therein.

The anti-UNQ733 antibodies of the invention can be in the different forms encompassed by the definition of "antibody" herein. Thus, the antibodies include full length or intact antibody, antibody fragments, native sequence antibody or amino acid variants, humanized, chimeric or fusion antibodies, immunoconjugates, and functional fragments thereof. In fusion antibodies an antibody sequence is fused to a heterologous polypeptide sequence. The antibodies can be modified in the Fc region to provide desired effector functions. As discussed in more detail in the sections herein, with the appropriate Fc regions, the naked antibody bound on the cell surface can induce cytotoxicity, e.g., via antibody-dependent cellular cytotoxicity (ADCC) or by recruiting complement in complement dependent cytotoxicity, or some other mechanism. Alternatively, where it is desirable to eliminate or reduce effector function, so as to minimize side effects or therapeutic complications, certain other Fc regions may be used.

In one embodiment, the antibody competes for binding or bind substantially to, the same epitope as the antibodies of the invention. Antibodies having the biological characteristics of the present anti-UNQ733 antibodies of the invention are also contemplated, specifically including the *in vivo* tumor targeting and any cell proliferation inhibition or cytotoxic characteristics.

Methods of producing the above antibodies are described in detail herein.

The present anti-UNQ733 antibodies, oligopeptides and organic/inorganic small molecules are useful for treating a UNQ733 polypeptide-expressing lymphoma, specifically non-Hodgkin's lymphoma, or alleviating one or more symptoms of the lymphoma in a mammal. Methods of the invention encompass usage of UNQ733 antagonists in the treatment and/or alleviation of symptoms of metastatic tumors associated with non-Hodgkin's lymphoma. The antibody, oligopeptide or organic/inorganic small molecule UNQ733 antagonist is able to bind to at least a portion of the cancer cells that express UNQ733 polypeptide in the mammal. In one embodiment, the antibody, oligopeptide or organic/inorganic small molecule is effective to destroy or kill UNQ733 polypeptide-expressing and/or -responsive tumor cells or inhibit the growth of such tumor cells, *in vitro* or *in vivo*,

upon binding to UNQ733 polypeptide. Such an antibody includes a naked anti-UNQ733 antibody (not conjugated to any agent). Naked antibodies that have cytotoxic or cell growth inhibition properties can be further harnessed with a cytotoxic agent to render them even more potent in tumor cell destruction. Cytotoxic properties can be conferred to an anti-UNQ733 antibody by, e.g., conjugating the antibody with a cytotoxic agent, to form an immunoconjugate as described herein. In some embodiments, the cytotoxic agent or a growth inhibitory agent is a small molecule. In some embodiments, toxins such as calicheamicin or a maytansinoid and analogs or derivatives thereof, are used.

The invention provides a composition comprising an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule of the invention, and a carrier. For the purposes of treating cancer, compositions can be administered to the patient in need of such treatment, wherein the composition can comprise one or more anti-UNQ733 antibodies present as an immunoconjugate or as the naked antibody. In a further embodiment, the compositions can comprise these antibodies, oligopeptides or organic/inorganic small molecules in combination with other therapeutic agents such as cytotoxic or growth inhibitory agents, including chemotherapeutic agents. The invention also provides formulations comprising an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule of the invention, and a carrier. In one embodiment, the formulation is a therapeutic formulation comprising a pharmaceutically acceptable carrier.

Another aspect of the invention is isolated nucleic acids encoding the anti-UNQ733 antibodies. Nucleic acids encoding both the H and L chains and especially the hypervariable region residues, chains which encode the native sequence antibody as well as variants, modifications and humanized versions of the antibody, are encompassed.

The invention also provides methods useful for treating non-Hodgkin's lymphoma or alleviating one or more symptoms of the lymphoma in a mammal, comprising administering a therapeutically effective amount of an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule to the mammal. The antibody, oligopeptide or organic/inorganic small molecule therapeutic compositions can be administered short term (acute) or chronic, or intermittent as directed by physician. Also provided are methods of inhibiting the growth of, and killing a UNQ733 polypeptide-expressing and/or -responsive cell.

The invention also provides kits and articles of manufacture comprising at least one anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule. Kits containing anti-UNQ733 antibodies, oligopeptides or organic/inorganic small molecules find use, e.g., for UNQ733 polypeptide cell killing assays, for purification or immunoprecipitation of UNQ733 polypeptide from cells. For example, for isolation and purification of UNQ733 polypeptide, the kit can contain an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule coupled to beads (e.g., sepharose beads). Kits can be provided which contain the antibodies, oligopeptides or organic/inorganic small molecules for detection and quantitation of UNQ733 polypeptide *in vitro*, e.g., in an ELISA or a Western blot.

Such antibody, oligopeptide or organic/inorganic small molecule useful for detection may be provided with a label such as a fluorescent or radiolabel.

K. Articles of Manufacture and Kits

Another embodiment of the invention is an article of manufacture containing materials useful for the treatment of a UNQ733 polypeptide expressing cancer, such as a non-Hodgkin's lymphoma. The article of manufacture comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for treating the cancer condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule of the invention. The label or package insert indicates that the composition is used for treating cancer. The label or package insert will further comprise instructions for administering the antibody, oligopeptide or organic/inorganic small molecule composition to the cancer patient. Additionally, the article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

Kits are also provided that are useful for various purposes, e.g., for UNQ733 polypeptide-expressing or cell killing assays, for purification or immunoprecipitation of UNQ733 polypeptide from cells. For isolation and purification of UNQ733 polypeptide, the kit can contain an anti-UNQ733 antibody, oligopeptide or organic/inorganic small molecule coupled to beads (e.g., sepharose beads). Kits can be provided which contain the antibodies, oligopeptides or organic/inorganic small molecules for detection and quantitation of UNQ733 polypeptide *in vitro*, e.g., in an ELISA or a Western blot. As with the article of manufacture, the kit comprises a container and a label or package insert on or associated with the container. The container holds a composition comprising at least one anti-UNQ733 polypeptide antibody, oligopeptide or organic/inorganic small molecule of the invention. Additional containers may be included that contain, e.g., diluents and buffers, control antibodies. The label or package insert may provide a description of the composition as well as instructions for the intended *in vitro* or detection use.

L. UNQ733 polypeptides and UNQ733 polypeptide-encoding nucleic acids - Specific forms and applications

Nucleotide sequences (or their complement) encoding UNQ733 polypeptides have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA probes. UNQ733 polypeptide-encoding nucleic acid will also be useful for the preparation of UNQ733 polypeptides by the

recombinant techniques described herein, wherein those UNQ733 polypeptides may find use, for example, in the preparation of anti-UNQ733 antibodies as described herein.

A full-length native sequence UNQ733 polypeptide gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate other cDNAs (for instance, those encoding naturally-occurring variants of UNQ733 polypeptide or UNQ733 polypeptide from other species) which have a desired sequence identity to a native UNQ733 polypeptide sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence UNQ733 polypeptide. By way of example, a screening method will comprise isolating the coding region of the UNQ733 polypeptide gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ³²P or ³⁵S, or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the UNQ733 polypeptide gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below. Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

Other useful fragments of the UNQ733 polypeptide-encoding nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target UNQ733 polypeptide mRNA (sense) or UNQ733 polypeptide DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of UNQ733 DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (Cancer Res. 48:2659, 1988) and van der Krol et al. (BioTechniques 6:958, 1988).

Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. Such methods are encompassed by the present invention. The antisense oligonucleotides thus may be used to block expression of a UNQ733 protein, wherein the UNQ733 protein may play a role in the induction of cancer in mammals. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable *in*

vivo (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

Preferred intragenic sites for antisense binding include the region incorporating the translation initiation/start codon (5'-AUG / 5'-ATG) or termination/stop codon (5'-UAA, 5'-UAG and 5'-UGA / 5'-TAA, 5'-TAG and 5'-TGA) of the open reading frame (ORF) of the gene. These regions refer to a portion of the mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation or termination codon. Other preferred regions for antisense binding include: introns; exons; intron-exon junctions; the open reading frame (ORF) or "coding region," which is the region between the translation initiation codon and the translation termination codon; the 5' cap of an mRNA which comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage and includes 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap; the 5' untranslated region (5'UTR), the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene; and the 3' untranslated region (3'UTR), the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene.

Specific examples of preferred antisense compounds useful for inhibiting expression of UNQ733 polypeptide include oligonucleotides containing modified backbones or non-natural internucleoside linkages. Oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides. Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotri-esters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and borano-phosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included. Representative United States patents that teach the preparation of phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496;

5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts. Representative United States patents that teach the preparation of such oligonucleosides include, but are not limited to, U.S. Pat. Nos.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, each of which is herein incorporated by reference.

In other preferred antisense oligonucleotides, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., Science, 1991, 254, 1497-1500.

Preferred antisense oligonucleotides incorporate phosphorothioate backbones and/or heteroatom backbones, and in particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a methylene (methylimino) or MMI backbone], -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein the native phosphodiester backbone is represented as -O-P-O-CH₂-] described in the above referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above referenced U.S. Pat. No. 5,602,240. Also preferred are antisense oligonucleotides having morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-alkyl, S-alkyl, or N-alkyl;

O-alkenyl, S-alkenyl, or N-alkenyl; O-alkynyl, S-alkynyl or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred antisense oligonucleotides comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂ CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₃)₂.

A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH₂-)_n group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂ NH₂), 2'-allyl (2'-CH₂-CH=CH₂), 2'-O-allyl (2'-O-CH₂-CH=CH₂) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-

5 methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ($-\text{C}\equiv\text{C}-\text{CH}_3$ or $-\text{CH}_2-\text{C}\equiv\text{CH}$) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine(1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one),
10 phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindeole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine,
15 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J. I., ed. John Wiley & Sons, 1990, and those disclosed by Englisch et al., Angewandte Chemie, International Edition, 1991, 30, 613. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention.
20 These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2.degree. C. (Sanghvi et al, Antisense Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar
25 modifications. Representative United States patents that teach the preparation of modified nucleobases include, but are not limited to: U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; 5,681,941 and 5,750,692, each of which is herein incorporated by reference.

30 Another modification of antisense oligonucleotides comprises chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides,
35 polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterol, lipids, cation lipids, phospholipids, cationic phospholipids, biotin, phenazine,

folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86, 6553-6556), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in U.S. patent application Ser. No. 09/334,130 (filed Jun. 15, 1999) and United States patents Nos.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the

case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Preferred chimeric antisense oligonucleotides incorporate at least one 2' modified sugar (preferably 2'-O-(CH₂)₂-O-CH₃) at the 3' terminal to confer nuclease resistance and a region with at least 4 contiguous 2'-H sugars to confer RNase H activity. Such compounds have also been referred to in the art as hybrids or gapmers. Preferred gapmers have a region of 2' modified sugars (preferably 2'-O-(CH₂)₂-O-CH₃) at the 3'-terminal and at the 5' terminal separated by at least one region having at least 4 contiguous 2'-H sugars and preferably incorporate phosphorothioate backbone linkages. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives. The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO_4 -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

Antisense or sense RNA or DNA molecules are generally at least about 5 nucleotides in length, alternatively at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, or 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length.

The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related UNQ733 polypeptide coding sequences.

Nucleotide sequences encoding a UNQ733 polypeptide can also be used to construct hybridization probes for mapping the gene which encodes that UNQ733 polypeptide and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as *in situ* hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

The UNQ733 polypeptide can be used in assays to identify other proteins or molecules involved in a binding interaction with the UNQ733 polypeptide. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors of the binding interaction. Screening assays can be designed to find lead compounds that mimic the biological activity of a native UNQ733 polypeptide or a receptor for UNQ733 polypeptide. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

Nucleic acids which encode UNQ733 polypeptide or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding UNQ733 polypeptide can be used to clone genomic DNA encoding UNQ733 polypeptide in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding UNQ733 polypeptide. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for UNQ733 polypeptide transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding UNQ733 polypeptide introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding UNQ733 polypeptide. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of UNQ733 polypeptide can be used to construct a UNQ733 gene "knock out" animal which has a defective or altered gene encoding UNQ733 polypeptide as a result of homologous recombination between the endogenous gene encoding UNQ733 polypeptide and altered genomic DNA encoding UNQ733 polypeptide introduced into an embryonic stem cell of the animal. For example, cDNA encoding UNQ733 polypeptide can be used to clone genomic DNA encoding UNQ733 polypeptide in accordance with established techniques. A portion of the genomic DNA encoding UNQ733 polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, Cell, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., Cell, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the UNQ733 polypeptide.

Nucleic acid encoding the UNQ733 polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve *in vivo* synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. "Gene therapy" includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes *in vivo*. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik *et al.*, Proc. Natl. Acad. Sci. USA 83:4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into

mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred *in vivo* gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., Trends in Biotechnology 11, 205-210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., J. Biol. Chem. 262, 4429-4432 (1987); and Wagner et al., Proc. Natl. Acad. Sci. USA 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., Science 256, 808-813 (1992).

The nucleic acid molecules encoding the UNQ733 polypeptides or fragments thereof described herein are useful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each UNQ733 nucleic acid molecule of the present invention can be used as a chromosome marker.

UNQ733 polypeptides and nucleic acid molecules of the invention may be used diagnostically for tissue typing, wherein UNQ733 polypeptides may be differentially expressed in one tissue as compared to another, preferably in a diseased tissue as compared to a normal tissue of the same tissue type. UNQ733 nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

This invention encompasses methods of screening compounds to identify those that prevent the effect of the UNQ733 polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the UNQ733 polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins, including e.g., inhibiting the expression of UNQ733 polypeptide from cells. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

All assays for antagonists are common in that they call for contacting the drug candidate with a UNQ733 polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the UNQ733 polypeptide or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the UNQ733 polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the UNQ733 polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

If the candidate compound interacts with but does not bind to a UNQ733 polypeptide, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, Nature (London), 340:245-246 (1989); Chien et al., Proc. Natl. Acad. Sci. USA, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, Proc. Natl. Acad. Sci. USA, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKERTM) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a UNQ733 polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular

component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

To assay for antagonists, the UNQ733 polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the UNQ733 polypeptide indicates that the compound is an antagonist to the UNQ733 polypeptide. Alternatively, antagonists may be detected by combining the UNQ733 polypeptide and a potential antagonist with membrane-bound UNQ733 polypeptide receptors or encoded receptors under appropriate conditions for a competitive inhibition assay. The UNQ733 polypeptide can be labeled, such as by radioactivity, such that the number of UNQ733 polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., Current Protocols in Immun., 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the UNQ733 polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the UNQ733 polypeptide. Transfected cells that are grown on glass slides are exposed to labeled UNQ733 polypeptide. The UNQ733 polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, labeled UNQ733 polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled UNQ733 polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with UNQ733 polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the UNQ733 polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the UNQ733 polypeptide.

Another potential UNQ733 antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature UNQ733 polypeptides herein, can be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res., 6:3073 (1979); Cooney et al., Science, 241: 456 (1988); Dervan et al., Science, 251:1360 (1991)), thereby preventing transcription and the production of the UNQ733 polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the UNQ733 polypeptide (antisense - Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, FL, 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of the UNQ733 polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the UNQ733 polypeptide, thereby blocking the normal biological activity of the UNQ733 polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, Current Biology, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is

designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, *supra*.

These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

Isolated UNQ733 polypeptide-encoding nucleic acid can be used for recombinantly producing UNQ733 polypeptide using techniques well known in the art and as described herein. In turn, the produced UNQ733 polypeptides can be employed for generating anti-UNQ733 antibodies using techniques well known in the art and as described herein.

Antibodies specifically binding a UNQ733 polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders, including cancer, in the form of pharmaceutical compositions.

If the UNQ733 polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco *et al.*, Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

MATERIALS & METHODS

5 *Cell Lines:* Chinese hamster ovary, DP12 (American Type Culture Collection (ATCC, Manassa, VA)) and PGSB (ATCC) mutants were used for transient expression and production of recombinant proteins. Jurkat T lymphoma, BJAB and Daudi B cell lymphomas, and Reh B cell leukemia were obtained from ATCC.

10 *RNA Expression Analysis:* For the analysis of UNQ733 mRNA expression in multiple human tumor and normal biopsy samples, Affymetrix data were obtained from Gene Logic, Inc. (Gaithersburg, MD). A total of 150 samples [6 normal samples; 62 cancer samples; and 82 non-cancer diseased samples] were analyzed. Gene Logic data were normalized using global scaling with a target intensity of 100. The Affymetrix data for UNQ733 in Fig.1 was generated from the U133 probe set ID 229152_at. For
15 in situ hybridization, a 368-bp ³³P-labeled antisense riboprobe was generated from a UNQ733 specific PCR product using a primer with oligonucleotide sequence 5'-CAG-GTG-ACC-AGG-TTT-ATC-GTG-3' (SEQ ID NO:5) and a sense control riboprobe with the primer 5'-AAT-ATT-CCA-GGG-CCA-GTC-ACT-3' (SEQ ID NO:6). Representative normal tissues and cancers including three lymphoma tissue microarrays (Cybrdi, Inc.; Frederick, MD) were also screened by in situ
20 hybridization.

Recombinant Proteins: UNQ733 cDNA was generated by PCR amplification from a human multi-tissue cDNA library. The PCR product was cloned into pSVID, a pRK based vector with a His₈-tag. PGSB cells (ATCC) were transfected using Fugene 6 (Roche; Indianapolis, IN) and conditioned media
25 collected after 7 days. His₈-tagged UNQ733 polypeptide was purified by Ni-chromatography followed by dialysis into a 10 mM citrate, pH 3.0 buffer. An N-terminal UNQ733 human placental alkaline phosphatase (UNQ733-AP) fusion protein was generated by inserting the entire UNQ733 coding sequence into a pRK vector containing the AP gene. CHO cells were transfected using Fugene 6 (Roche) and supernatant allowed to condition for 6 days. All experiments used 733-AP supernatant
30 and a control supernatant (Relt-AP) that was normalized for AP activity.

Monoclonal Anti-UNQ733 Antibodies: BALB/c mice were immunized 15 times with 1.0 µg of UNQ733-His8 resuspended in monophosphoryl lipid A/trehalose dicorynomycolate adjuvant (Corixa; Hamilton, MT) into each hind footpad at 3 to 4 days intervals. Three days after final boost, popliteal
35 lymph nodes were fused with myeloma cell line P3X63Ag8.653 (ATCC, Manassa, VA) using 50% polyethylene glycol and cultured in 96 well tissue culture plates. Culture supernatants were initially

screened for their ability to bind to UNQ733-His8. The selected hybridomas were then cloned by limiting dilution.

Antibody clones were deposited with the ATCC under the Budapest Treaty. American Type Culture Collection (ATCC) is located at 10801 University Boulevard, Manassas, Virginia 20110-2209.

Details of the deposits are as follows:

<u>Clone designation</u>	<u>Deposit Date</u>	<u>ATCC Deposit No.</u>
3E7.9.20	June 2, 2004	PTA-6026
3F10.11.2	June 2, 2004	PTA-6027
3H1.4.8	June 2, 2004	PTA-6028
4A9.12.12	June 2, 2004	PTA-6029
5A8.11.6	June 2, 2004	PTA-6030
5F2.6.14	June 2, 2004	PTA-6031
9D6.11.15	June 2, 2004	PTA-6032
10G10.15.16	June 2, 2004	PTA-6033
12H4.11.3	June 2, 2004	PTA-6034

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC § 122 and the Commissioner's rules pursuant thereto (including 37 CFR § 1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

Western blots and Immunoprecipitation: Purified His₈-tagged UNQ733 protein was subjected to 18% SDS-PAGE and immunoblotting (Invitrogen; Carlsbad, CA). Blots were incubated with 1 µg/ml anti-UNQ733 Mab overnight at 4°C. Rabbit anti-mouse Ig horseradish peroxidase (Amersham; Piscataway, NJ) was used as a secondary antibody, and the blots were developed using the SuperSignal

West Pico Luminol/Enhancer Solution (Pierce; Rockford, IL). Immunoprecipitation was done by adding 1 µg purified Mab to 50 ng of purified His₈-tagged UNQ733 in TBS [50 mM Tris, 150 mM NaCl], followed by protein A/G-agarose (Pierce), and incubated overnight at 4°C. The immunoprecipitates were recovered, washed with TBS, and subjected to 18% SDS-PAGE. Blots were incubated with India HisProbe (1:5000; Pierce) to detect His₈-tagged UNQ733, washed in TBS + 0.05% Tween-20 and developed as described above.

Immunohistochemistry: Frozen human tonsil sections (5µ) were fixed in -20°C acetone for 5 minutes followed by removal of OCT in TBST (Tris-buffered saline with Tween 20). Endogenous peroxidase, avidin and biotin were quenched using KPL blocking buffer (KPL, 37-00-84; Gaithersburg, MD) and Vector avidin/biotin blocking kit (Vector, SP-2001; Burlingame, CA) respectively, followed by TBST rinses. Slides were incubated for 30 minutes in 10% normal horse serum in 3% BSA/PBS for 30 minutes. Slides were then incubated in 10µg/ml anti-UNQ733 (clone 3H1) antibody for 60 minutes at room temperature. After washes in TBST, slides were incubated with 2.5µg/ml biotinylated horse anti-mouse secondary antibody (Vector, BA-2001) for 30 minutes at room temperature. Following TBST washes slides were incubated with Vectastain ABC Elite Reagents (Vector, PK-6100) for 30 minutes at room temperature, slides were then rinsed with TBST, incubated with Pierce Metal Enhanced DAB (Pierce, 1856090) for four minutes, and rinsed in water. Slides were then counterstained with Mayer's hematoxylin, dehydrated, mounted and coverslipped for bright field viewing.

Flow Cytometry: For analysis of UNQ733 expression in tonsil cells, fresh human tonsils tissue was minced and digested in 50 mg/ml of collagenase IV and 50 U/ml of DNase I (Sigma-Aldrich; St. Louis, MO, USA). The resulting cell suspension was subjected to negative selection of B cells with anti-CD19 microbeads on an autoMACS separator (Miltenyi Biotec; Auburn, CA). The CD19⁻ fraction was blocked with Fc blocking reagent (Miltenyi Biotec) followed by staining anti-CD14-FITC (clone M5E2; BD Pharmingen; San Diego, CA) and anti-CD21-PE (clone B-ly4; BD Pharmingen) in FACS wash [PBS + 0.5% bovine serum albumin] for 30 min at ambient temperature. Cells were washed in FACS wash and fixed with FACSlyse solution (BD Pharmingen) for 20 minutes at ambient temperature. Cells were washed with FACS wash and treated with FACS Perm 2 solution (BD Pharmingen). Cells were washed in FACS wash and incubated with 10 µg/ml biotinylated UNQ733-specific Mab 12H4 for 30 min at ambient temperature followed by washing and incubation with SA-APC (BD Pharmingen). Cells were analyzed on a BD FACSCaliber flow cytometer. FDC were gated as CD14⁺CD21⁺ cells and monocytes gated as CD14⁺CD21⁻ cells.

For analysis of His₈-tagged UNQ733 binding to lymphoma cell lines, cells were incubated with 50 µg/ml (4.5 µM) or 20 µg/ml (1.8 µM) His₈-tagged UNQ733 diluted in FACS wash at ambient temperature for 30 minutes. Cells were washed and bound His₈-tagged UNQ733 detected with 10 µg/ml anti-His₆(C-term)-FITC (Invitrogen). Alternately, His₈-tagged UNQ733 bound to cells was

detected with 10 µg/ml biotinylated UNQ733-specific Mab 12H4 followed by SA-PE (BD Pharmingen). For analysis of His₈-tagged UNQ733 binding to human peripheral blood B cells, peripheral blood from normal healthy human donors was obtained. Peripheral blood mononuclear cells were isolated using LSM lymphocyte separation medium (ICN; Costa Mesa, CA). B cells were purified using anti-CD19 microbeads and a MACS separator (Miltenyi Biotec). B cells were cultured in Iscove's modified Dulbecco's media + 10% fetal bovine serum in the presence or absence of 1 µg/ml anti-CD40 (clone 5C3; BD Pharmingen) and 10 ng/ml rIL-4 (Preprotech) for 48 hours before flow cytometric analysis as described above. B cells were gated by staining with anti-CD19-APC (clone HIB19; BD Pharmingen).

Protein Expression and Sequencing: PGSB cells were transiently transfected with His₈-tagged UNQ733 using Fugene 6 (Roche). The cultures were incubated in the presence or absence of 10 µM decanoyl-Arg-Val-Lys-Arg-chloromethylketone (Bachem; King of Prussia, PA), a proprotein convertase inhibitor, added to the cultures on day one. After 6 days in culture, supernatants were harvested, affinity precipitated with Ni-NTA (Qiagen; Valencia, CA). The Ni-NTA bound protein was subjected to 18% SDS-PAGE and transblotted to PVDF for N-terminal sequencing by Edman degradation on a Procise N-Terminal Sequencer (ABI; Foster City, CA).

Protein Binding Assay: BJAB cells were incubated with serial dilutions of UNQ733-AP supernatant or a single concentration of Relt-AP control supernatant. After incubation for 1 hour at 4°C, cells were washed three times with PBS, and transferred to a 96-well microplate. Cells were heated to 50°C for 1 hour to inactivate endogenous alkaline phosphatase activity. 1-Step PNPP substrate buffer (Pierce) was added, and the reaction was allowed to proceed for 30 minutes at ambient temperature in the dark. Cells were centrifuged and an aliquot of substrate buffer was removed for analysis of absorbance at 405nm on a SpectraMax 250 microplate reader (Molecular Devices; Sunnyvale, CA).

For competitive inhibition of UNQ733-AP binding, a fixed concentration (0.5X) of UNQ733-AP supernatant was used and simultaneously incubated with increasing concentrations of His₈-tagged UNQ733 or a vehicle control [10 mM citrate, pH 3.0 buffer].

RESULTS

UNQ733 is over-expressed in non-Hodgkin's lymphoma and hyperplastic tonsils

Oligonucleotide-based microarray expression data from over 4800 human biopsy samples were analyzed for genes that showed elevated mRNA expression relative to normal tissue. One such gene was designated UNQ733, and it was found to be over-expressed in tumors of lymphoid origin (Fig. 1). Within the lymphoid tumors analyzed, it was found significantly over-expressed in 29 of 62

(46.8%) non-Hodgkin's lymphomas. By this analysis over-expression of UNQ733 was also observed in 75 of 82 (91%) diseased, non-malignant tonsillar tissue designated hyperplastic by histopathology.

To further assess the prevalence of UNQ733 overexpression in non-Hodgkin's lymphoma, we obtained three lymphoma tissue microarrays (TMA) containing 173 non-Hodgkin's lymphoma tissue cores. *In situ* hybridization was carried out on the TMAs using a UNQ733-specific probe. In 66 of 173 (38%) individual tumors, UNQ733 mRNA expression was detected in the malignant cells. Overexpression of UNQ733 was observed in B cell lymphoma (NOS), diffuse large B cell lymphoma, follicular lymphoma, small lymphocytic lymphoma, malignant lymphoma (NOS), malignant T cell lymphoma, anaplastic large cell lymphoma and mucosal associated lymphoid tissue lymphoma.

In situ hybridization was carried out on normal and neoplastic human tissue. UNQ733 mRNA expression was observed in the germinal centers of all normal lymphoid tissues including the spleen, lymph nodes, tonsils, and Peyer's patches (Figure 3). In tonsils, UNQ733 expression was detected in the germinal centers (Figure 4A, 4B) and crypts (Figure 4C, 4D). Furthermore, UNQ733 expression was observed in the serous salivary glands (not shown). In neoplastic tissue, expression was observed in the malignant cells of 38% (3/8) of non-Hodgkin's lymphomas analyzed (Figures 5 & 6).

Monoclonal antibodies (MAb) against recombinant His₈-tagged UNQ733 were generated and characterized. A panel of 9 MAbs with reactivity to recombinant UNQ733 by ELISA was obtained and characterized in western blot assays, immunoprecipitation assays and immunohistochemistry assays (IHC) to assess binding to UNQ733 in cell-free and cell-associated environments. Two of the MAbs were found to detect recombinant UNQ733 in a western blot assay, 10G10 and 12H4 (Figure 7). These antibodies were able to detect recombinant His₈-tagged UNQ733 in the western blot. Most of the monoclonal antibodies were able to immunoprecipitate UNQ733 (Figure 8). The MAb 3H1 was tested for its ability to bind UNQ733 in frozen tonsillar tissue by IHC. The pattern of staining observed was identical to the localization of UNQ733 mRNA by *in situ* hybridization. Strong staining was observed in the crypts and weak staining in the germinal centers (Figure 9). The pattern of staining observed was consistent with a follicular dendritic cell (FDC).

To ascertain the identity of the cell type expressing UNQ733 in the tonsil, fresh tonsil tissue from surgical resections was obtained and used in flow cytometry to delineate the FDCs and monocytes by CD14 and CD21 expression. Biotinylated-12H4 MAb was used to detect intracellular UNQ733. Expression of UNQ733 protein was detected in the FDC population, but not in the monocytes (Figure 10), a finding consistent with the immunohistochemistry data.

UNQ733 is processed by a proprotein convertase

Proprotein convertases (PCs) are a family of proteolytic enzymes that cleave biologically inactive precursor proteins into their mature active forms. Members of this family of proteases recognize amino acid sequences with the motif (K/R)-X-X-(K/R)/. UNQ733 contains a putative PC cleavage site R-E-K-R/S³⁰. Two protein species are typically observed when recombinant UNQ733-

His₈ was analyzed by western blot (Figure 7). N-terminal sequencing of the proteins confirmed the presence of two species, the first beginning with the sequence ¹⁸FPVSQDQE, which was consistent with the predicted “full-length” protein after signal sequence cleavage. The second protein began with the sequence ³⁰SISDSDEL that would be the predicted product of PC cleavage of UNQ733. To confirm that the activity of a PC was responsible for the observed cleavage event, UNQ733-His₈ was expressed in the presence and absence of a PC inhibitor, decanoyl-RVKR-chloromethylketone (Figure 11). Multiple species were still observed, but the proportion of “full-length” UNQ733 was substantially increased in the PC inhibitor treated supernatants (Figure 12). This indicates that UNQ733 is processed by a PC.

UNQ733 binds to lymphoma cell lines and primary B cells

UNQ733 binding to lymphoma cell lines was assessed by flow cytometry. Binding of UNQ733-His₈ to both BJAB cells, a human B cell lymphoma (Figure 13A), and Jurkat cells, a human T cell lymphoma (Figure 13B) was detected by using a FITC conjugated anti-His₆ detecting antibody (Invitrogen). Antibody binding was observed for both cell lines, with the BJAB cells showing higher binding of the UNQ733-His₈ than the Jurkat cells. These studies were extended to test the panel of MAbs for the ability to detect binding of UNQ733 on a subset of human B cell lymphoma/leukemia cell lines. Several MAbs, including 3H1 and 10G10, were able to detect bound UNQ733-His₈ (not shown); 12H4 showed the best signal to noise ratio in detecting UNQ733-His₈ bound to BJAB, Reh and Daudi cells (Figure 14). To evaluate the specificity of the binding observed by the UNQ733-His₈ molecule, a UNQ733-human placenta alkaline phosphatase (UNQ733-AP) fusion protein was used to detect bound UNQ733, and UNQ733-His₈ used to function as a competitor. The UNQ733-AP protein in tissue culture supernatants was found to bind to the C1R B lymphoblast cell line in a dose-dependent manner while a control AP fusion protein, RELT-AP, showed no binding (Figure 15A). When tested in a competition assay, UNQ733-His₈ was observed to inhibit the binding of UNQ733-AP to C1R cells (Figure 15B). This indicated that the binding of UNQ733 to B lymphomas is specific.

UNQ733 is expressed in follicular dendritic cells in the germinal centers of secondary lymphoid organs. Because germinal centers are a site of B cell selection, class-switch recombination, and somatic hypermutation, the role of UNQ733 in non-neoplastic B cell biology was also ascertained. Peripheral blood mononuclear cells were obtained from normal healthy human donors, and the B cells were purified and analyzed for UNQ733 binding (Figure 16). It was observed that “resting” peripheral blood B cells from multiple human donors bound UNQ733. When those PB B cells were first stimulated with an agonistic MAb specific to CD40 and recombinant IL-4, the binding of UNQ733 was found to increase 2-5 fold depending on the donor (Figure 16). It appears that stimulation of the B cells with anti-CD40 results in increased surface expression of the receptor for UNQ733.

CLAIMSWHAT IS CLAIMED IS:

1. A method for inhibiting the growth of a non-Hodgkin's lymphoma comprising contacting the cell with a UNQ733 antagonist which causes inhibition of the growth of the cell.

2. A method of therapeutically treating a mammal having non-Hodgkin's lymphoma comprising administering to the mammal a therapeutically effective amount of a UNQ733 antagonist, thereby resulting in the effective therapeutic treatment of the lymphoma.

3. A method of inhibiting the growth of a non-Hodgkin's lymphoma comprising administering a UNQ733 antagonist to the lymphoma, whereby growth of the lymphoma is inhibited.

4. A method of inhibiting the growth of a non-Hodgkin's lymphoma comprising administering a UNQ733 antagonist to a cell that expresses and/or is responsive to UNQ733, whereby growth of the lymphoma is inhibited.

5. A method of inhibiting growth of a non-Hodgkin's lymphoma comprising administering a UNQ733 antagonist to a cell present in the lymphoma, whereby growth of the lymphoma is inhibited.

6. A method for treating or preventing a cell proliferative disorder associated with increased expression or activity of UNQ733, the method comprising administering to a subject in need of such treatment an effective amount of a UNQ733 antagonist.

7. The method of any of claims 1-6 wherein the UNQ733 antagonist binds to UNQ733 polypeptide.

8. The method of any of claims 1-6 wherein the UNQ733 antagonist binds to UNQ733 on a non-Hodgkin's lymphoma cell.

9. The method of any of claims 1-6 wherein the UNQ733 antagonist causes death of a cell expressing and/or responsive to UNQ733.

10. The method of any of claims 1-6 wherein the UNQ733 antagonist is an antibody, an oligopeptide, inorganic or organic small molecule.

11. The method of 10 wherein binding of the antibody, oligopeptide, inorganic or organic small molecule to the UNQ733 polypeptide causes inhibition of the growth of a cell expressing and/or responsive to the polypeptide.

12. The method of any of claims 1-6 wherein the cell is a cancer cell and the UNQ733 antagonist causes death of a cell expressing and/or responsive to the UNQ733 polypeptide.

13. The method of any of claims 1-6 wherein the cell is a cancer cell and binding of the antagonist to UNQ733 polypeptide causes death of a cell expressing and/or responsive to the polypeptide.

14. The method of any of claims 1-6 wherein the UNQ733 antagonist is an antibody, oligopeptide, inorganic or organic small molecule and binding of the antibody, oligopeptide, inorganic or organic small molecule to the UNQ733 polypeptide causes death of a cell expressing and/or responsive to the polypeptide.

15. The method of any of claims 1-6 wherein the UNQ733 antagonist inhibits activation of UNQ733.

16. The method of 15 wherein the antagonist inhibits activation of UNQ733 by modulating enzymatic processing of UNQ733 protein.

17. The method of 16 wherein enzymatic processing of the UNQ733 protein is modulated by inhibiting a proprotein convertase that cleaves UNQ733.

18. The method of any of claims 1-6 wherein the antagonist is a monoclonal antibody, antibody fragment, chimeric antibody, humanized antibody, human antibody, multi-specific antibody or single-chain antibody.

19. The method of any of claims 1-6 wherein the antagonist is conjugated to a growth inhibitory agent or cytotoxic agent.

20. The method of any of claims 1-6 wherein a chemotherapeutic agent is also administered to the subject.

21. The method of any of claims 1-6 wherein the antagonist comprises an antibody or binding oligopeptide produced in a CHO cell or bacterial cell.

22. The method of claim 6 wherein the cell proliferative disorder is cancer and the UNQ733 antagonist is an anti-UNQ733 antibody, UNQ733 binding oligopeptide, UNQ733 binding inorganic or organic small molecule.

23. The method of any of claims 1-6 wherein the antagonist causes direct killing or growth inhibition of cells that express and/or are responsive to UNQ733 polypeptide.

24. The method of any of claims 1-6 wherein the antagonist antagonizes the cell growth potentiating activity of UNQ733 polypeptide.

25. The method of any of claims 1-6 wherein the cell proliferative disorder is a non-Hodgkin's lymphoma.

26. The method of any of claims 1-6 wherein the cell proliferative disorder is hyperplasia.

27. The method of any of claims 1-6 wherein the hyperplasia is diseased, non-malignant tonsillar tissue.

28. A method of diagnosing a non-Hodgkin's lymphoma of interest comprising determining the level of UNQ733 in a test sample of tissue cells comprising cells of origin for the lymphoma of interest and in a control sample of known cells of the same tissue origin, wherein a higher level of UNQ733 in the test sample, as compared to the control sample, is indicative of the presence of the lymphoma.

29. A method of determining whether an individual is at risk for non-Hodgkin's lymphoma comprising determining the level of UNQ733 in a test sample of tissue cells comprising cells of origin for the lymphoma of interest from said individual and in a control sample of known cells of the same tissue origin, wherein a higher level of UNQ733 in the test sample, as compared to the control sample, is an indication that the individual is at risk for the lymphoma.

30. The method of claim 28 or 29 wherein the level of UNQ733 is determined based on level of UNQ733 RNA in the test sample.

31. The method of claim 28 or 29 wherein, the level of UNQ733 is determined based on level of UNQ733 polypeptide in the test sample.

32. The method of claim 28 or 29 wherein the level of UNQ733 is determined based on the level of UNQ733 protein activity in the test sample.

33. The method of claim 28 or 29 comprising exposing the sample to an antibody, oligopeptide, inorganic or organic small molecule that binds to the UNQ733 polypeptide and determining binding of the antibody, oligopeptide, inorganic or organic small molecule to the UNQ733 polypeptide in the sample.

34. The method of claim 33 wherein the antibody, UNQ733 binding oligopeptide, or UNQ733 binding inorganic or organic small molecule is detectably labeled and/or attached to a solid support.

35. A method for screening for biologically active agents for the treatment of non-Hodgkin's lymphoma comprising combining a candidate agent with a transgenic mammal having a genome comprising an integrated transgene encoding UNQ733 operably linked to a promoter, wherein said transgene results in said mammal acquiring non-Hodgkin's lymphoma and determining the effect of said agent on non-Hodgkin's lymphoma in said mammal.

36. A method for screening for biologically active agents for the treatment of non-Hodgkin's lymphoma comprising combining a candidate agent with a transgenic mammal cell culture, each cell of said culture comprising an integrated transgene encoding UNQ733 operably linked to a promoter, wherein said transgene results in said mammal acquiring non-Hodgkin's lymphoma, and determining the effect of said candidate agent on the transgenic mammal cell culture.

37. The method of claim 35 or 36 wherein the candidate agent is an antibody or fragment thereof, an oligopeptide, or an inorganic or organic small molecule.

38. A method of targeting a therapeutic agent to a non-Hodgkin's lymphoma in a host, the method comprising administering to the host said therapeutic agent in a form that is linked to a molecule that binds UNQ733, whereby the agent is targeted to the lymphoma in the host.

39. The method of claim 38 wherein the molecule that binds UNQ733 is an antibody, an oligopeptide, an inorganic or organic small molecule.

40. The method of claim 38 wherein the molecule that binds UNQ733 is capable of specifically binding to UNQ733 located on a cell *in vitro* or *in vivo*.

41. The method of claim 38 wherein the molecule that binds UNQ733 is an antibody.

42. The method of claim 41 wherein the antibody is produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034, or progeny thereof.

5 43. The method of claim 41 wherein the antibody binds to the same epitope on UNQ733 as the antibody having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.

10 44. The method of claim 41 wherein the antibody competes with the antibody having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034 for binding to UNQ733.

45. The method of claim 44 wherein UNQ733 is located in a cell free environment, is a secreted protein *in vivo* or *in vitro*, or is bound to a cell.

15 46. An article of manufacture comprising a container and a composition contained within the container, wherein the composition comprises a UNQ733 antagonist.

20 47. The article of manufacture of claim 46 comprising a label affixed to the container, and/or a package insert included with the container that refers to the use of the composition for therapeutic treatment of non-Hodgkin's lymphoma.

48. A kit comprising a composition comprising a UNQ733 antagonist, optionally further comprising instruction for using the composition for treatment of non-Hodgkin's lymphoma.

25 49. A composition comprising a UNQ733 polypeptide binding molecule, wherein the composition is suitable for detecting UNQ733 polypeptide in a test sample from an individual suspected of having non-Hodgkin's lymphoma.

30 50. The composition of claim 49 wherein the binding molecule is an antibody, oligopeptide, inorganic or organic small molecule.

51. The composition of claim 49 wherein the binding molecule is an antibody, wherein the antibody is capable of binding to UNQ733 in a detection assay.

35 52. The composition of claim 51 wherein the detection assay is western blot, immunoprecipitation or immunostaining.

53. The composition of claim 49 wherein the antibody is produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.

54. The composition of claim 49 wherein the antibody binds to the same epitope on UNQ733 as the antibody produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.

55. The composition of claim 49 wherein the antibody competes with the antibody produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034 for binding to UNQ733.

56. The composition of claim 49, further comprising a carrier.

57. An article of manufacture comprising a container and a composition contained within the container, wherein the composition comprises a UNQ733 polypeptide binding molecule and is suitable for detecting UNQ733 polypeptide in a test sample from an individual suspected of having non-Hodgkin's lymphoma.

58. The article of manufacture of claim 57 comprising a label affixed to the container, and/or a package insert included with the container, that refers to the use of the composition for detection of non-Hodgkin's lymphoma.

59. The article of manufacture of claim 57 wherein the binding molecule is an antibody, oligopeptide, inorganic or organic small molecule.

60. The article of manufacture of claim 59 wherein the binding molecule is an antibody, wherein the antibody is capable of binding to UNQ733 in a detection assay.

61. The article of manufacture of claim 60 wherein the detection assay is western blot, immunoprecipitation or immunostaining.

62. The article of manufacture of claim 59 wherein the antibody is produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.

63. The article of manufacture of claim 59 wherein the antibody binds to the same epitope on UNQ733 as the antibody produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.

5 64. The article of manufacture of claim 59 wherein the antibody competes with the antibody produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034 for binding to UNQ733.

10 65. A kit comprising a composition comprising a UNQ733 polypeptide binding molecule, optionally further comprising instruction for using the composition for detecting non-Hodgkin's lymphoma.

66. The kit of claim 65 wherein the binding molecule is an antibody.

15 67. The kit of claim 66 wherein the antibody is capable of binding to UNQ733 in a detection assay.

68. The kit of claim 67 wherein the detection assay is western blot, immunoprecipitation or immunostaining.

20 69. The kit of claim 66 wherein the antibody is produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.

25 70. The kit of claim 66 wherein the antibody binds to the same epitope on UNQ733 as the antibody produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034.


30 71. The kit of claim 66 wherein the antibody competes with the antibody produced by hybridoma having ATCC deposit no. PTA-6026, PTA-6027, PTA-6028, PTA-6029, PTA-6030, PTA-6031, PTA-6032, PTA-6033 or PTA-6034 for binding to UNQ733.

35 72. The method of any of claims 1-5 wherein the non-Hodgkin's lymphoma is a B-cell lymphoma, diffuse large B cell lymphoma, follicular lymphoma, small lymphocytic lymphoma, malignant lymphoma, malignant T cell lymphoma, anaplastic large cell lymphoma or mucosal associated lymphoid tissue lymphoma.

73. The composition of any of claims 49-56 wherein the non-Hodgkin's lymphoma is a B-cell lymphoma, diffuse large B cell lymphoma, follicular lymphoma, small lymphocytic lymphoma, malignant lymphoma, malignant T cell lymphoma, anaplastic large cell lymphoma or mucosal associated lymphoid tissue lymphoma.

1 / 18

FIG. 1



2 / 18

FIG. 2

Core Diagnosis	Negative	Positive	% Positive
B-cell lymphoma (NOS)	25	18	42
Diffuse large B cell lymphoma	45	29	39
Follicular lymphoma	1	4	80
Small lymphocytic lymphoma	3	3	50
Mantle cell lymphoma	1	0	0
Malignant lymphoma (NOS)	5	3	38
Burkitt's lymphoma	2	0	0
Malignant T cell lymphoma	15	6	29
Anaplastic large cell lymphoma	3	1	25
Mucosal associated lymphoid tissue lymphoma	7	2	22
Total	107	66	38

3 / 18

FIG. 3

Tissues	Expression	Comments
Aorta	-	
Breast	-	
Gall Bladder	-	
Prostate	-	
Lung	+	Germinal centers
Spleen	+	Germinal centers
Peyers Patches	+	Germinal centers
Lymph Node	+	Germinal centers
Salivary Gland	+	Serous cells
Tonsil	+	GCs and crypts

FIG. 4




FIG. 5

Tissues	No. Positive	Comments
Cartilage	0/4	
Adipose	0/4	
Liver	0/4	
Colon	0/4	
Salivary Gland	1/1 (100%)	Serous Adenoma
Breast (including TMA)	3/64 (5%)	Ductal epithelium
	2/64 (3%)	Lymphocyte infiltrate
NHL	3/8 (38%)	Malignant cells positive

FIG._6




FIG._7





FIG._8

8 / 18



9 / 18





FIG._9






FIG. 10
FDCs (CD14+CD21+)



Macrophages (CD14+CD21-)



7 yo male

61 yo male

FIG._11




FIG._12

% of each species






FIG._13



h733-his + anti-his

FIG. 14



h7333-his detected with 12H4-biotin & SA-PE

FIG._15




FIG._16




FIG._17

TAAAAACAGCTACAATAATTCCAGGGCCAGTCACCTTGCCCAATTTCTCATAAACAGCGTCAGAGAGAAAGAACTGACTGAAACGTTTGAGATGAAGA
AAGTTCTCCTCCTGATCACAGCCATCTTGGCAGTGGCTGTGTGGTTTCCCAAGTCTCTCAAGACCAGGAACGAGAAAAAAGAAAGTATCAGTGAC
AGCGATGAATTAGCTTCAGGGTTTTTTGTGTTCCTTACCCATATCCATTTGCCCCACTTCCACCAATTCCATTTCCAAGATTTCCATGGTT
TAGACGTAATTTTCCATATCCAATACCTGAATCTGCCCCCTACAACTCCCCTTCCCTAGCGAAAAAGTAAACAAAGAGGATAAGTCACGATAAAC
CTGGTCACCTGAAATTGAAATTGAGCCACTTCCCTTGAAGAAATCAAAATTCCTGTTAATAAAGAAAAACAAATGTAAATTGAAATAGCACACA
GCATTCTCTAGTCAATATCTTTAGTGATCTTTTAATAAACATGAAAGCAAGATTTTGGTTTTCTTAAATTCCACA (SEQ ID NO:1)

FIG._18

MKKVLLITAILAVGFPVSQDQEREKRSISDSDELASGFFVFPYPFRPLPPFPFRFP
WFRNFPPIPESAPTTPLPSEK (SEQ ID NO:2)

FPVSQDQEREKRSISDSDELASGFFVFPYPFRPLPPFPFRFPWFRNFPPIPESAPTTP
PLPSEK (SEQ ID NO:3)

SISDSDELASGFFVFPYPFRPLPPFPFRFPWFRNFPPIPESAPTTPLPSEK (SEQ
ID NO:4)