



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2252435 C 2004/03/23

(11)(21) **2 252 435**

(12) **BREVET CANADIEN  
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1997/04/16

(87) Date publication PCT/PCT Publication Date: 1997/10/23

(45) Date de délivrance/Issue Date: 2004/03/23

(85) Entrée phase nationale/National Entry: 1998/10/15

(86) N° demande PCT/PCT Application No.: US 1997/006485

(87) N° publication PCT/PCT Publication No.: 1997/038957

(30) Priorité/Priority: 1996/04/16 (60/015,521) US

(51) Cl.Int.<sup>6</sup>/Int.Cl.<sup>6</sup> C07C 31/125, C07C 233/46,  
C07C 303/24, C07C 51/23, C07C 29/16, C07C 305/04,  
C07C 303/02, C11D 1/14, C07C 53/126, C07C 309/12,  
C07C 2/06

(72) Inventeur/Inventor:  
CONNOR, DANIEL STEDMAN, US

(73) Propriétaire/Owner:  
THE PROCTER & GAMBLE COMPANY, US

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : FABRICATION D'UN TENSIOACTIF RAMIFIE

(54) Title: BRANCHED SURFACTANT MANUFACTURE

(57) **Abrégé/Abstract:**

Alpha-olefins are dimerised to form vinylidene olefins, which are then isomerised and reacted with CO/H<sub>2</sub> under OXO conditions or reacted directly with CO/H<sub>2</sub> under OXO conditions to give mid- to near-mid chain branched alcohols. These branched-chain alcohols are feedstocks for the production of alkyl sulphate, alkyl ethoxy sulphate, and alkyl carboxylate surfactants and other surfactants.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07C 2/06, 11/02, 29/16, 31/125, 303/24, 305/06, 51/16, 53/126, C11D 1/14</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/38957</b> <b>(43) International Publication Date:</b> 23 October 1997 (23.10.97)
<b>(21) International Application Number:</b> PCT/US97/06485 <b>(22) International Filing Date:</b> 16 April 1997 (16.04.97) <b>(30) Priority Data:</b> 60/015,521                      16 April 1996 (16.04.96)                      US <b>(71) Applicant (for all designated States except US):</b> THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> CONNOR, Daniel, Stedman [US/US]; 9217 Sagemeadow Drive, Cincinnati, OH 45251 (US). <b>(74) Agents:</b> REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> BRANCHED SURFACTANT MANUFACTURE  <b>(57) Abstract</b>  Alpha-olefins are dimerised to form vinylidene olefins, which are then isomerised and reacted with CO/H <sub>2</sub> under OXO conditions or reacted directly with CO/H <sub>2</sub> under OXO conditions to give mid- to near-mid chain branched alcohols. These branched-chain alcohols are feedstocks for the production of alkyl sulphate, alkyl ethoxy sulphate, and alkyl carboxylate surfactants and other surfactants.		



## BRANCHED SURFACTANT MANUFACTURE

FIELD OF THE INVENTION

5       The present invention relates to processes for manufacturing deterative surfactants, especially those containing branched-chain hydrophobic units.

BACKGROUND OF THE INVENTION

10       Conventional deterative surfactants comprise molecules having a water-solubilizing substituent (hydrophilic group) and an oleophilic substituent (hydrophobic group). Such surfactants typically comprise hydrophilic groups such as carboxylate, sulfate, sulfonate, amine oxide, polyoxyethylene, and the like, attached to an alkyl, alkenyl or alkaryl hydrophobe usually containing from about 10 to about 20 carbon atoms. Accordingly, the manufacturer of such surfactants must have access to a source of hydrophobe groups to which the desired hydrophile can be attached by chemical means. The earliest source of hydrophobe groups comprised the natural fats and oils, which were converted into soaps (i.e., carboxylate hydrophile) by saponification with base. Coconut oil and palm oil are still used to manufacture soap, as well as to manufacture the alkyl sulfate ("AS") class of surfactants. Other hydrophobes are available from petrochemicals, including alkylated benzene which is used to manufacture alkyl benzene sulfonate surfactants ("LAS").

15       The literature asserts that certain branched hydrophobes can be used to advantage in the manufacture of alkyl sulfate deterative surfactants; see, for example, U.S. 3,480,556 to deWitt, et al., November 25, 1969. However, it has been determined that the beta-branched surfactants described in the '556 patent are inferior with respect to certain solubility parameters, as evidenced by their Krafft temperatures. It has further been determined that surfactants having branching towards the center of carbon chain of the hydrophobe have much lower Krafft temperatures. See: "The Aqueous Phase Behavior of Surfactants", R.G. Laughlin, Academic Press, N.Y. (1994) p. 347. Accordingly, it has now been determined that such surfactants are preferred for use especially under cool or cold water washing conditions (e.g., 20°C-5°C).


25       One problem associated with the manufacture of deterative surfactants having hydrophobe groups with mid- or near-mid chain branching is the lack of a ready source of such hydrophobes. By the present invention, a process is described for manufacturing such branched hydrophobes and converting them into mid- or near-mid chain branched surfactants.

SUMMARY OF THE INVENTION

The present invention encompasses a process for preparing mid- to near mid-chain branched olefins (primarily, methyl branched at or near the mid-chain region). Such materials are then used as the basic feedstock which provides the hydrophobic portion of branched-chain deterative surfactants.

The process herein is illustrated by the following reaction sequence.

## 1) Alpha-Olefin Dimerization




wherein R and R' may be the same or different linear alkyl, and wherein R is C<sub>3</sub>-C<sub>7</sub>, preferably C<sub>5</sub> to C<sub>7</sub> linear alkyl, and R' is C<sub>3</sub>-C<sub>7</sub>, preferably C<sub>5</sub>-C<sub>7</sub> linear alkyl. For use in preparing surfactants in cleaning products such as laundry detergents, dishwashing liquids, and the like, R and R' are preferably the same or within one or two carbon atoms of each other in chain length. Some linear olefins may also result from the dimerization and these can optionally be removed using molecular sieves. Step 1 of the process herein is designed to provide branched olefins which preferably contain from about 12 to about 18 (avg.) total carbon atoms.

## 2) Alcohol Production

In Step 2 (Route A), the olefin mixture for Step 1 can be pre-randomized to enhance the ultimate formation of alcohols (i) and (ii) in subsequent Step 3. Alternatively (Step 2, Route B), this pre-randomization step can be deleted and the Oxo catalyst, itself, can randomize the final product among the three possible terminal positions.

Route A



or

Route B




The Oxo process to make alcohols is described in detail in Kirk-Othmer Encyclopedia of Chemical Technology, 4th Edition, Volume 1, pp. 903-8 (1991), Jacqueline I. Kroschwitz, Executive Editor, Wiley-Interscience, N.Y. The catalyst for this step is, for example, cobalt-carbon monoxide-organophosphine.

5        The alcohol mixture of Step (2) of the present process comprises branched-chain primary alcohol compounds of the following formulae for use in Step (3), below.



10        It is to be understood that when  $\text{CH}_2\text{OH}$  is substituted on R or R' it is primarily on their respective terminal carbons or to a lesser extent on their penultimate carbons. Desirably, minimal amounts of compounds of the formula (iii) are present in Step (3).

### 3) Surfactant Production



15        Advantageously, the present process results in no geminal branching, i.e., no "quaternary" carbon substitution. Steric hindrance will block inverse addition to vinylidene carbon with cobalt-carbon monoxide-organophosphine which otherwise would form a non-biodegradable quaternary carbon. Moreover, little (less than about 3%) vicinal branching occurs. Of course, some of the overall feedstock may  
20        remain unbranched. Typically, and preferably from the standpoint of cleaning performance and biodegradability, the present process provides hydrophobes with one near-central methyl in the case of isomers i) and ii).

All percentages, ratios and proportions herein are by weight, unless otherwise specified.

25

## DETAILED DESCRIPTION OF THE INVENTION

### 1) Olefin dimerization:

The present invention encompasses, in a process for preparing surfactant precursor hydrophobes from dimerization of two same or near same chain length alpha-olefins to form a detergent range vinylidene olefin. These alpha olefins C<sub>5</sub> to C<sub>10</sub>, preferably C<sub>7</sub> to C<sub>9</sub> are dimerized to give C<sub>10</sub> to C<sub>20</sub>, preferably C<sub>14</sub> to C<sub>18</sub> vinylidene olefins which upon Oxo reaction give C<sub>11</sub> to C<sub>21</sub>, preferably C<sub>15</sub> to C<sub>19</sub> alcohols. There are a number of processes for accomplishing said dimerization; see US 4,658,078, US 4,973,788; O.S. Vostrikova, A.G. Ibragimov, G.A. Tolstikov, L.M. Zelenova and U.M. Dzhemilev, Izv. Akad. Nauk SSSR, Ser. Khim. (1980), (10), 2330-2 [Chem. Abstr. 94:65032]; Jpn. Kokai Tokkyo Koho, 06228016 A2 [Chem. Abstr. 122:186930].

## 2) Alcohol Production

### Route A

part a) The carbon-carbon double bond of the vinylidene olefin is pre-isomerized using a method such as Shell uses to isomerize alpha-olefins in their SHOP process; see Kirk-Othmer Encyclopedia of Chemical Technology, 4th Edition, Volume 17, pp. 848-50 (1996), Jacqueline I. Kroschwitz, Executive Editor, Wiley-Interscience, N.Y. and Chemical Economics Handbook, pp. 681.5030K-L, Stanford Research Institute, Menlo Park, CA 94025, Oct. 1993.

part b) Oxo chemistry is used to convert the pre-isomerized vinylidene olefin (now largely internal olefin) to a primary alcohol mixture. For this an Oxo catalyst which isomerizes the double bond to alpha positions prior to carbonylation is desired as is the case using cobalt-carbonyl-phosphine catalysts in the one step process, see Kirk-Othmer Encyclopedia of Chemical Technology, 4th Edition, Volume 1, pp. 903-8 (1991). Route A, (that is including pre-isomerization) is undertaken to assure relatively high yields of alcohols i and ii versus alcohol iii. Note alcohol sulfates of i and ii are desired surfactants whereas that of iii may be deficient.

### Route B

This step utilizes the same Oxo catalyst on the vinylidene olefin directly without it pre-isomerization. This relies upon the catalyst to completely isomerize the carbon-carbon double bond of the vinylidene olefin prior to carbonylation. The object is to obtain as much i and ii relative to iii as is obtained in Route A.

## 4a

The present invention further seeks to encompass a process for preparing branched alkyl sulfate surfactants, which are produced by sulfating the alcohols prepared according to the previous process disclosed herein.

5 Other fatty alcohol-derived surfactants can also be made, e.g., alkyl ethoxyl sulfates (AES), alkyl polyglucosides (APG), etc. Note that surfactants other than alcohol sulfates or AES may be made by oxidizing said alcohol or its aldehyde intermediate into a carboxylate (i.e., a branched-chain soap). This soap can be an excellent surfactant and/or detergent builder in and of itself. This carboxylate can also be used as a feedstock and converted to branched acyl-taurates, -isethionates, -



sarcosinates, -N-methylglucamide or other acyl-derived surfactants using art-disclosed techniques.

#### INDUSTRIAL APPLICABILITY

5 Branched-chain surfactants of the type resulting from the present process can be used in all manner of cleaning compositions. Such compositions include, but are not limited to: granular, bar-form and liquid laundry detergents; liquid hand dishwashing compositions; liquid, gel and bar-form personal cleansing products; shampoos; dentifrices; hard surface cleaners, and the like. Such compositions can contain a variety of conventional deterative ingredients. The following listing of  
10 such ingredients is for the convenience of the formulator, and not by way of limitation of the types of ingredients which can be used with the branched-chain surfactants herein.

The branched-chain surfactants herein can be used in combination with detergency builders. Such builders include, for example, 1-10 micrometer zeolite A,  
15 polycarboxylate builders such as citrate, layered silicate builders such as "SKS-6" (Hoechst) and phosphate materials, especially sodium tripolyphosphate ("STPP"). Most laundry detergents typically comprise at least about 1% builder, more typically from about 5% to about 80% builder or mixtures of builders.

Enzymes, such as proteases, amylases, lipases, cellulases, peroxidases, and  
20 mixtures thereof, can be employed in detergent compositions containing the branched-chain surfactants. Typical detergent compositions comprise from about 0.001% to about 5% of commercial enzymes.

Detergent compositions can also contain polymeric soil release agents (SRA's). Such materials include, for example, anionic, cationic and non-charged  
25 monomer units, especially polyester materials. Preferred materials of this type include oligomeric terephthalate esters, sulfonated substantially linear ester oligomers comprising a backbone of terephthaloyl and oxyalkyleneoxy repeat units and phthaloyl-derived sulfonated terminal moieties. A variety of SRA's are described, for example, in U.S. 4,968,451; 4,711,730; 4,721,580; 4,702,857;  
30 4,877,896; 5,415,807; and in other literature references. Such soil release materials typically comprise from about 0.01% to about 10% of finished detergent compositions.

Detergent compositions may also optionally contain bleaching compositions comprising a bleaching agent and one or more bleach activators. If present,  
35 bleaching agents such as percarbonate or perborate (especially perborate monohydrate "PB1") typically are used at levels from about 1% to about 30% of finished detergent compositions. Bleach activators such as nonanoyloxy-benzene



sulfonate ("NOBS") and tetraacetyl ethylenediamine ("TAED"), and mixtures thereof, can be used to enhance the bleaching activity of materials such as perborate and percarbonate. If present, the amount of bleach activator will typically be from about 0.1% to about 60% of a bleaching composition comprising a bleaching agent-  
5 plus-bleach activator. Other bleaching agents such as the so-called "photoactivated" bleaches (see U.S. 4,033,718) can also be used. Sulfonated zinc phthalocyanine is an especially preferred photoactivated bleaching agent.

Detergent compositions can also contain clay soil removal/antiredeposition agents such as ethoxylated tetraethylene pentamine; see U.S. 4,597,898. Such  
10 materials typically comprise from about 0.01% to about 10% of fully-formulated laundry detergents.

Detergent compositions can also contain from about 0.1% to about 7% of polymeric dispersing agents, which are especially useful in the presence of zeolite and/or layered silicate builders. Such materials are known in the art (see U.S.  
15 3,308,067). Such materials include acrylate/malic-based copolymers, such as described in EP 193,360, as well as polyethylene glycol ("PEG").

Detergent compositions herein can also include various brighteners, dye transfer inhibiting agents (especially polymers of N-vinylpyrrolidone and N-vinylimidazole), suds suppressors (especially silicones), chelating agents such as  
20 nitrilotriacetate, ethylenediamine disuccinate, and the like. Such materials will typically comprise from about 0.5% to about 10%, by weight, of fully-formulated cleaning compositions.

Moreover, it is to be understood that the branched-chain surfactants prepared in the manner of the present invention may be used singly in cleaning compositions  
25 or in combination with other deterative surfactants. Typically, fully-formulated cleaning compositions will contain a mixture of surfactant types in order to obtain broad-scale cleaning performance over a variety of soils and stains and under a variety of usage conditions. One advantage of the branched-chain surfactants herein is their ability to be readily formulated in combination with other known surfactant  
30 types. Nonlimiting examples of additional surfactants which may be used herein typically at levels from about 1% to about 55%, by weight, include the unsaturated sulfates such as oleyl sulfate, the C<sub>10</sub>-C<sub>18</sub> alkyl alkoxy sulfates ("AE<sub>x</sub>S"; especially EO 1-7 ethoxy sulfates), C<sub>10</sub>-C<sub>18</sub> alkyl alkoxy carboxylates (especially the EO 1-5 ethoxycarboxylates), the C<sub>10</sub>-18 glycerol ethers, the C<sub>10</sub>-C<sub>18</sub> alkyl polyglycosides  
35 and their corresponding sulfated polyglycosides, and C<sub>12</sub>-C<sub>18</sub> alpha-sulfonated fatty acid esters. Nonionic surfactants such as the ethoxylated C<sub>10</sub>-C<sub>18</sub> alcohols and alkyl phenols, (e.g., C<sub>10</sub>-C<sub>18</sub> EO (1-10) can also be used. If desired, other

conventional surfactants such as the C<sub>12</sub>-C<sub>18</sub> betaines and sulfobetaines ("sultaines"), C<sub>10</sub>-C<sub>18</sub> amine oxides, and the like, can also be included in the overall compositions. The C<sub>10</sub>-C<sub>18</sub> N-alkyl polyhydroxy fatty acid amides can also be used. Typical examples include the C<sub>12</sub>-C<sub>18</sub> N-methylglucamides. See WO 92/06154. Other sugar-derived surfactants include the N-alkoxy polyhydroxy fatty acid amides, such as C<sub>10</sub>-C<sub>18</sub> N-(3-methoxypropyl) glucamide. The N-propyl through N-hexyl C<sub>12</sub>-C<sub>18</sub> glucamides can be used for low sudsing. C<sub>10</sub>-C<sub>20</sub> conventional soaps may also be used. If high sudsing is desired, the branched-chain C<sub>10</sub>-C<sub>16</sub> soaps may be used. C<sub>10</sub>C<sub>14</sub> alkyl benzene sulfonates (LAS), which are often used in laundry detergent compositions, can also be used with the branched surfactants herein.

The following Examples illustrate the use of branched-chain surfactants prepared according to the present invention in various cleaning compositions, but is not intended to be limiting thereof.

15

**EXAMPLE I**

Granular laundry detergents are prepared as follows.

		<b><u>A</u></b>	<b><u>B</u></b>	<b><u>C</u></b>
	<b>Blown Powder</b>			
	Zeolite A	30.0	22.0	6.0
20	Sodium sulfate	19.0	5.0	7.0
	Polyacrylate			
	LAS	13.0	11.0	21.0
	Branched AS*	9.0	8.0	8.0
	Silicate, Na	-	1.0	5.0
25	Soap	-	-	2.0
	Carbonate, Na	8.0	16.0	20.0
	<b>Spray On</b>			
	C <sub>14-15</sub> EO7	1.0	1.0	1.0
	<b>Dry additives</b>			
30	Protease	1.0	1.0	1.0
	Lipase	0.4	0.4	0.4
	Amylase	0.1	0.1	0.1
	Cellulase	0.1	0.1	0.1
	NOBS	-	6.1	4.5
35	PB1	1.0	5.0	6.0
	Sodium sulfate	-	6.0	-
	Moisture & Miscellaneous	-----Balance-----		



\*C<sub>12</sub>-C<sub>14</sub> methyl branched alkyl sulfate, prepared as disclosed above.

A bleach-containing nonaqueous liquid laundry detergent is prepared as follows.

### EXAMPLE II

5	<u>Component</u> <u>wt.)</u>	<u>Wt. %</u>	<u>Range (%)</u>
	<u>Liquid Phase</u>		
	Branched AS*	25.3	18-35
	C <sub>12-14</sub> , EO5 alcohol ethoxylate	13.6	10-20
10	Hexylene glycol	27.3	20-30
	Perfume	0.4	0-1.0
	<u>Solids</u>		
	Protease enzyme	0.4	0-1.0
	Na <sub>3</sub> Citrate, anhydrous	4.3	3-6
15	Sodium perborate (PB-1)	3.4	2-7
	Sodium nonanoyloxybenzene sulfonate (NOBS)	8.0	2-12
	Sodium carbonate	13.9	5-20
	Diethyl triamine pentaacetic acid (DTPA)	0.9	0-1.5
	Brightener	0.4	0-0.6
20	Suds Suppressor	0.1	0-0.3
	Minors	-----Balance-----	

\*C<sub>12</sub>-C<sub>16</sub> methyl branched alkyl sulfate, Na salt, prepared as disclosed above.

A hand dishwashing liquid is as follows.



EXAMPLE III

	<u>Ingredient</u>	<u>% (wt.)</u>	<u>Range (% wt.)</u>
	Branched AS*	13.0	5-15
5	Ammonium C <sub>12-13</sub> alkyl ethoxy sulfate	15.0	10-35
	Coconut amine oxide	2.6	2-5
	Betaine**/Tetronic 704®	0.87-0.10	0-2 (mix)
	Alcohol Ethoxylate C <sub>8</sub> E <sub>11</sub>	5.0	2-10
10	Ammonium xylene sulfonate	4.0	1-6
	Ethanol	4.0	0-7
	Ammonium citrate	0.06	0-1.0
	Magnesium chloride	3.3	0-4.0
	Calcium chloride	2.5	0-4.0
15	Ammonium sulfate	0.08	0-4.0
	Hydrogen peroxide	200 ppm	0-300 ppm
	Perfume	0.18	0-0.5
	Maxatase® protease	0.50	0-1.0
	Water and minors	-----Balance-----	
20	*C <sub>12</sub> C <sub>14</sub> methyl branched alkyl sulfate, triethanolammonium salt, prepared as disclosed above.		
	**Cocoalkyl betaine.		





## WHAT IS CLAIMED IS:

1. A process for preparing branched alcohols comprising the steps of:
  - (a) dimerizing alpha olefins of the formula  $RCH=CH_2$  and  $R'CH_2=CH_2$ , to form olefins of the formula:



wherein in the above formulas R and R' may be the same or different  $C_3$  to  $C_7$  linear alkyl substituents; followed by either

- (b) isomerizing the olefins from Step (a) and the subsequent reaction of said isomerized olefins with  $CO/H_2$  under Oxo conditions; or directly reacting the olefins from Step (a) with  $CO/H_2$  under Oxo conditions; and
- (c) recovering the desired branched alcohols of the formula:



and any aldehyde intermediates of the desired branched alcohols.

2. The process according to Claim 1 comprising the additional step of:
  - (d) sulfonating the desired branched alcohols to form branched alkyl sulfates.
  
3. The process according to Claim 1 comprising the additional steps of:
  - (d) ethoxylating the desired branched alcohols; and
  - (e) sulfating the product of step (d) to form branched alkyl ethoxy sulfates.
  
4. The process according to Claim 1 comprising the additional step of:
  - (d) oxidizing the desired branched alcohols and any aldehyde intermediates of the desired branched alcohols to form branched alkyl carboxylate surfactants.