

(43) International Publication Date  
27 June 2013 (27.06.2013)(51) International Patent Classification:  
*C07C 37/055* (2006.01)    *C07C 39/21* (2006.01)  
*C07C 37/62* (2006.01)(21) International Application Number:  
PCT/US201 1/066053(22) International Filing Date:  
20 December 2011 (20.12.2011)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **COLGATE-PALMOLIVE COMPANY** [US/US]; 300 Park Avenue, New York, New York 10022 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **NAIK, Ramesh** [IN/IN]; Plot No. 36B, Road No. 3&5, Jigani Industrial Area, Anekal Taluk, Bangalore, Karnataka 560105 (IN). **WALIKAR, Sanju** [IN/IN]; Plot No. 36B, Road No. 3&5, Jigani Industrial Area, Anekal Taluk, Bangalore, Karnataka 560105 (IN). **JEYARAMAN, Govindarajalu** [IN/IN]; Plot No. 36B, Road No. 3&5, Jigani Industrial Area, Anekal Taluk, Bangalore, Karnataka 560105 (IN). **RAMASWAMY RANGANATHAN, Koottungalmad-****hom** [IN/IN]; Plot No. 36B, Road No. 3&5, Jigani Industrial Area, Anekal Taluk, Bangalore, Karnataka 560105 (IN).(74) Agent: **HEBLE, Nikhil, A.**; COLGATE-PALMOLIVE COMPANY, Patent Department, 909 River Road, Piscataway, New Jersey 08855 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on nextpage]

(54) Title: PROCESSES FOR MAKING MAGNOOL ANALOGS



(57) Abstract: Described herein are high yield methods for making magnolol analogs which are 5,5'-dialkyl-biphenyl-2,2'-diols.

**WO 2013/095364 A1**       

---

**SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, Published:**  
GW, ML, MR, NE, SN, TD, TG).

— *with international search report (Art. 21(3))*

## PROCESSES FOR MAKING MAGNOLOL ANALOGS

### BACKGROUND

**[0001]** There is a need for safe, effective antibacterial and anti-inflammatory agents for use in oral care compositions. Magnolia extract is known to contain compounds having antibacterial and/or anti-inflammatory properties, and such compounds have been the focus of considerable interest for use in oral care compositions. The use of such compounds in oral care compositions is described, for example, in WO2001/085 116, WO 201 1/106492 and WO 201 1/106493, the contents of which application are incorporated herein by reference. Methods of synthesizing magnolol are disclosed, e.g., in WO 201 1/106003. Synthetic non-natural analogs of various components of magnolia extract are also known to have antibacterial activity, but the compounds are in some cases expensive to synthesize.

**[0002]** Magnolol analogs having lower alkyl in place of allyl are of particular interest, as are compounds wherein the alkyl or allyl side chains are ortho rather than para to the hydroxy groups (isomanolols). Tetrahydro-magnolol, (5,5'-dipropyl-biphenyl-2,2'-diol) is a broad spectrum antibacterial and anti-inflammatory agent with potential applications in oral care and personal care products. It has advantages over magnolol, e.g. in that it does not stain toothbrush bristles when used as an antibacterial/anti-inflammatory agent in a toothpaste. Existing synthetic methods for making magnolol derivatives, however, involve costly reagents and yields are low. There is a need for cheaper, higher yield synthetic procedures to make magnolol derivatives.

### SUMMARY

**[0003]** Previous synthetic approaches to making magnolol generally start with bromination of biphenyl-2,2'-diol, to get the 5,5'-dibromo- biphenyl-2,2'-diol, followed by O-protection, with methyl or other O-protecting group, reaction with allyl bromide to get the magnolol in protected form, deprotection to get magnolol, which can then be hydrogenated to obtain propyl magnolol. Analogous compounds are made analogously. We have found that on scale up, the step of O-protecting the 5,5'-dibromo- biphenyl-2,2'-diol is inefficient and slow. We have found that carrying out the O-protection step *before* the bromination step results in a more efficient reaction and higher yields.

[0004] The deprotection step is another expensive and yield limiting step in the existing processes. The methods reported for demethylation are often costly, and require sometimes very low temperatures (-78°C using **BBr3**) and sometimes reflux conditions. The reaction mixture is always difficult to separate and purify which results in low yield of magnolol. We have found that the use of an aluminium chloride / thiourea complex for deprotection does not require extreme temperatures or expensive reagents and results in high yields.

(0005] Finally, we avoid the need for a bromination step or a hydrogenation step or a difficult separation of alkyl analogs of magnolol from magnolol by introducing the alkyl group by Friedel-Crafts acylation of 2,2'-bianisole followed by Clemmensen reduction of the alkionyl derivative.

[0006] The invention thus provides a simple, relatively high yield synthesis for 5,5'-dialkyl-biphenyl-2,2'-diols, comprising

- (i) methylating biphenyl-2,2'-diol using dimethyl sulfate, to obtain 2,2'-dimethoxy-biphenyl;
- (ii) acylating the 2,2'-dimethoxy-biphenyl with alkionyl halide, for example propionyl chloride, to obtain the corresponding 2,2'-dimethoxy-5,5'-dialkionyl-biphenyl;
- (iii) reducing the 2,2'-dimethoxy-5,5'-dialkionyl-biphenyl to obtain 2,2'-dimethoxy-5,5'-dialkyl-biphenyl;
- (iv) demethylating the 2,2'-dimethoxy-5,5'-dipropyl-biphenyl by reaction with aluminium chloride and thiourea; and
- (v) recovering the 5,5'-dialkyl-biphenyl-2,2'-diol thus obtained;

wherein "alk" or "alkyl" refers to linear, branched or cyclic C<sub>2</sub>-io alkyl, for example selected from *t*-propyl, isopropyl,  $\alpha$ -butyl, and isobutyl, e.g., *n*-propyl.

[0007] Further areas of applicability of the present invention will become apparent from the detailed description provided hereinafter. It should be understood that the detailed description and specific examples, while indicating the preferred embodiment of the invention, are intended for purposes of illustration only and are not intended to limit the scope of the invention.

DETAILED DESCRIPTION

[0008] The invention thus provides a method (Method 1) for making 5,5'-dialkyl-biphenyl-2,2'-diols, for example 5,5'-dipropyl-biphenyl-2,2'-diol, comprising


- (i) methylating biphenyl-2,2'-diol using dimethyl sulfate, to obtain 2,2'-dimethoxy-biphenyl;
- (ii) acylating the 2,2'-dimethoxy-biphenyl with alkionyl halide to obtain 2,2'-dimethoxy-5,5'-dialkionyl-biphenyl;
- (iii) reducing the 2,2'-dimethoxy-5,5'-dialkionyl-biphenyl to obtain 2,2'-dimethoxy-5,5'-dialkyl-biphenyl;
- (iv) demethylating the 2,2'-dimethoxy-5,5'-dialkyl-biphenyl by reaction with aluminium chloride and thiourea; and
- (v) recovering the 5,5'-dialkyl-biphenyl-2,2'-diol thus obtained;

e.g., wherein "alk" or "alkyl" refers to linear, branched or cyclic  $C_{2-10}$  alkyl, for example selected from  $\alpha$ -propyl, isopropyl,  $\alpha$ -butyl, and isobutyl, e.g.,  $\alpha$ -propyl.

- 1.1. Method 1 wherein step (i) is carried out in aqueous media in presence of an inorganic base, e.g., sodium hydroxide or potassium hydroxide.
- 1.2. Any of the foregoing methods wherein step (ii) is carried out in an apolar aprotic solvent, e.g., ethylene dichloride, in the presence of an aluminum chloride catalyst.
- 1.3. Any of the foregoing methods wherein step (iii) is carried out in the presence of heat, a strong acid, e.g., HCl, and a zinc catalyst.
- 1.4. Any of the foregoing methods wherein step (iv) is carried out at temperatures between 30°C and 60°C.
- 1.5. Any of the foregoing methods wherein the product is 5,5'-dipropyl-biphenyl-2,2'-diol, comprising
  - (i) methylating biphenyl-2,2'-diol using dimethyl sulfate, to obtain 2,2'-dimethoxy-biphenyl;
  - (ii) acylating the 2,2'-dimethoxy-biphenyl with propionyl chloride to obtain 2,2'-dimethoxy-5,5'-dipropionyl-biphenyl;
  - (iii) reducing the 2,2'-dimethoxy-5,5'-dipropionyl-biphenyl to obtain 2,2'-dimethoxy-5,5'-dipropyl-biphenyl;

- (iv) demethylating the 2,2'-dimethoxy-5,5'-dipropyl-biphenyl by reaction with aluminium chloride and thiourea; and
- (v) recovering the 5,5'-dipropyl-biphenyl-2,2'-diol thus obtained.

[0009] The reaction scheme is as follows for 5,5'-dipropyl-biphenyl-2,2'-diol; other 5,5'-dialkyl-biphenyl-2,2'-diols are made analogously using the corresponding alkionyl halides in place of propionyl chloride:



[0010] As used throughout, ranges are used as shorthand for describing each and every value that is within the range. Any value within the range can be selected as the terminus of the range. In addition, all references cited herein are hereby incorporated by reference in their entireties. In the event of a conflict in a definition in the present disclosure and that of a cited reference, the present disclosure controls.

[0011] Unless otherwise specified, all percentages and amounts expressed herein and elsewhere in the specification should be understood to refer to percentages by weight. The amounts given are based on the active weight of the material.

### EXAMPLE

**Example I:** Synthesis of tetrahydro-magnolol (5,5'-dipropyl-biphenyl-2,2'-diol)

**[0012] Protection:** Biphenol is reacted in sodium hydroxide with sodium sulfate for 1-2 hours. Solid is separated, washed with water and dried at 60-65 °C

**[0013] Acylation:** Combine propionyl chloride with anhydrous aluminum chloride in 1,2-dichloroethane. Add bianisole at room temperature and allow to stir for 4-6 hours. Quench in diluted hydrochloric acid and separate organic layer. Remove solvent via distillation and crystallize from methanol.

**[0014] Reduction:** Reflux 3-5 hours the 4,4'-dipropionyl 2,2'-bianisole in hydrochloric acid with zinc powder. Remove methanol and add ethyl acetate with stirring. Remove zinc by filtration and solvent by distillation.

**[0015] De-methylation:** Add the product of the previous step to a mixture of aluminum chloride, thiourea and 1,2-dichloroethane slowly over 3 hours at 50°C. Maintain temperature and stirring for an additional 3-4 hours. Cool reaction and add to hydrochloric acid and separate phases. Add organic layer to charcoal, filter and distill solvent to recover title compound.

**[0016]** The overall yield is about 55%, with a purity of greater than 98%.

CLAIMS

1. A method for making 5,5'-dialkyl-biphenyl-2,2'-diols, comprising the following steps:
  - (i) methylating biphenyl-2,2'-diol using dimethyl sulfate, to obtain 2,2'-dimethoxy-biphenyl;
  - (ii) acylating the 2,2'-dimethoxy-biphenyl with alkionyl chloride to obtain 2,2'-dimethoxy-5,5'-dialkionyl-biphenyl;
  - (iii) reducing the 2,2'-dimethoxy-5,5'-dialkionyl-biphenyl to obtain 2,2'-dimethoxy-5,5'-dialkyl-biphenyl;
  - (iv) demethylating the 2,2'-dimethoxy-5,5'-dialkyl-biphenyl by reaction with aluminium chloride and thiourea; and
  - (v) recovering the 5,5'-dialkyl-biphenyl-2,2'-diol thus obtained.
2. The method of claim 1 wherein step (i) is carried out in aqueous media in presence of an inorganic base, e.g., sodium hydroxide or potassium hydroxide.
3. The method of any of the foregoing claims wherein step (ii) is carried out in an apolar aprotic solvent in the presence of aluminum chloride.
4. The method of any of the foregoing claims wherein step (iii) is carried out in the presence of heat, a strong acid, e.g., HC1, and a zinc catalyst.
5. The method of any of the foregoing claims wherein step (iv) is carried out at temperatures between 30°C and 60°C.
6. The method of any of the foregoing claims wherein "alk" or "alkyl" refers to linear, branched or cyclic  $C_2$ -<sup>10</sup> alkyl.
7. The method of any of the foregoing claims wherein the product is 5,5'-dipropyl-biphenyl-2,2'-diol, comprising the following steps:
  - (i) methylating biphenyl-2,2'-diol using dimethyl sulfate, to obtain 2,2'-dimethoxy-biphenyl;
  - (ii) acylating the 2,2'-dimethoxy-biphenyl with propionyl chloride to obtain 2,2'-dimethoxy-5,5'-dipropionyl-biphenyl;
  - (iii) reducing the 2,2'-dimethoxy-5,5'-dipropionyl-biphenyl to obtain 2,2'-dimethoxy-5,5'-dipropyl-biphenyl;
  - (iv) demethylating the 2,2'-dimethoxy-5,5'-dipropyl-biphenyl by reaction with aluminium chloride and thiourea; and

(v) recovering the 5,5'-dipropyl-biphenyl-2,2'-diol thus obtained.

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2011/066053

A. CLASSIFICATION OF SUBJECT MATTER  
INV. C07C37/055 C07C37/62 C07C39/21  
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2006/140880 AI (SUBRAMANYAM RAVI [US] ET AL) 29 June 2006 (2006-06-29) paragraphs [0019] , [0020] , [0046] -----	1-7



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
20 June 2012	28/06/2012

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Eberhard, Michael
--	---

**INTERNATIONAL SEARCH REPORT**

## Information on patent family members

International application No  
PCT/US2011/066053

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
us 2006140880	AI	29-06-2006	AR 052853 AI	04-04-2007
			AU 2005322192 AI	06-07-2006
			BR PI0519414 A2	20-01-2009
			CA 2591846 AI	06-07-2006
			EP 1830925 AI	12-09-2007
			EP 2266666 AI	29-12-2010
			RU 2349300 CI	20-03-2009
			US 2006140880 AI	29-06-2006
			US 2012052026 AI	01-03-2012
			Wo 2006071654 AI	06-07-2006