



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2438663 A1 2002/08/29

(21) **2 438 663**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2002/02/14
(87) Date publication PCT/PCT Publication Date: 2002/08/29
(85) Entrée phase nationale/National Entry: 2003/08/18
(86) N° demande PCT/PCT Application No.: JP 2002/001221
(87) N° publication PCT/PCT Publication No.: 2002/066499
(30) Priorité/Priority: 2001/02/19 (2001-41615) JP

(51) Cl.Int.⁷/Int.Cl.⁷ C07K 7/54, A61K 38/12, A61P 31/10,
C07K 7/06

(71) Demandeur/Applicant:
TAKARA BIO INC., JP

(72) Inventeurs/Inventors:
KUROME, TORU, JP;
AWAZU, NAOYUKI, JP;
TAKESAKO, KAZUTOH, JP;
KATO, IKUNOSHIN, JP

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PEPTIDE CYCLIQUE
(54) Title: CYCLIC PEPTIDE

(57) **Abrégé/Abstract:**

A novel cyclic peptide having an antifungal activity, its pharmacologically acceptable salts and antifungal medicinal compositions containing the same.



ABSTRACT

A novel cyclic peptide having an antifungal
activity, its pharmacologically acceptable salts and
5 antifungal medicinal compositions containing the same.

DESCRIPTION

CYCLIC PEPTIDE

5 Technical Field

The present invention relates to a novel cyclic peptide having an antifungal activity and a pharmacologically acceptable salt thereof as well as an antifungal pharmaceutical composition containing the same.

10

Background Art

Fungi infect man, animals, plants and the like to cause various diseases. Among these, fungi belonging to the genus Candida, the genus Cryptococcus, the genus Aspergillus and the like are known as the principal causative fungi of systemic mycoses.

15

Currently known antifungal agents that can be used for treatment or prevention of fungal infections are only a very few. The therapeutic drugs available for systemic mycoses in man and animals are amphotericin B, flucytosine, miconazole and fluconazole. However, therapeutic activities of these agents are unsatisfactory, either in efficacies, toxicities, antimicrobial spectra and the like.

20

Specifically, few agents are effective for fungi

25


belonging to the genus *Aspergillus* among fungi that cause systemic mycoses in man (e.g., *Candida albicans*, *Cryptococcus neoformans* and *Aspergillus fumigatus*). Therefore, a therapeutic agent that can cover the drawbacks associated with the conventional agents has been desired.

Objects of Invention

The main object of the present invention is to provide a novel cyclic peptide having an excellent antifungal activity among cyclic lipopeptides having D-allo-threonine or various D- α -amino acids having β -hydroxy that can be obtained by chemical synthesis.

Summary of Invention

In summary, the present invention relates to a cyclic peptide of general Formula (1) or a pharmacologically acceptable salt thereof:



wherein

m and n may be identical or different from each other, and are independently 0 or 1;

X is O or NH;

R¹ is hydrogen, C6-21 alkyl, substituted alkyl, alkenyl, substituted alkenyl, acylamino, aryl, substituted aryl, heterocyclic ring or substituted heterocyclic ring, wherein said substituted alkyl, substituted alkenyl, substituted aryl or substituted heterocyclic ring is substituted with at least one of hydroxyl group, C1-8 alkyl, C1-8 alkoxy, C2-8 alkenyl, C2-8 acyl, C2-8 acyloxy, C6-14 aryl, C6-14 aryloxy or C1-8 heterocyclic ring;

AA¹ to AA⁶ are independently an α - or β -amino acid;

AA⁷ is a D- α -amino acid having HO- in a side chain; and

AA⁸ is an α - or β -amino acid.

The present inventors have searched novel cyclic lipopeptides prepared by means of chemical synthesis for a compound having a strong activity on fungi that cause systemic mycoses in humans (e.g., *Candida albicans*, *Cryptococcus neoformans* and *Aspergillus fumigatus*). As a result, the present inventors have unexpectedly found that, among cyclic lipopeptides each consisting of eight α - or β -amino acid residues and one fatty acid residue having hydroxy group or amino group at the α - or β -position, a cyclic lipopeptide compound containing at least one D- α -amino acid residue having β -hydroxy, preferably D- α -amino

acid having erythro form β -hydroxy, or D-allo-threonine exhibits a strong activity on the above-mentioned fungi (including *Aspergillus fumigatus*) that cause systemic mycoses in humans for which few agents have been effective.
5 Thus, the present invention has been completed.

Brief Description of Drawings

Figure 1: a figure illustrating a general protocols of solid phase synthesis according to the present
10 invention.

Detailed Description of the Invention

The compound of Formula (1) is a cyclic lipopeptide consisting of one hydroxy acid or amino acid
15 residue having R^1 as a lipid portion as well as eight α - or β -amino acid residues, AA^1 to AA^8 , wherein m and n may be identical or different from each other, and are independently 0 or 1; and X is O or NH.

Said R^1 is hydrogen, C6-21 alkyl, substituted
20 alkyl, alkenyl, substituted alkenyl, acylamino, aryl, substituted aryl, heterocyclic ring or substituted heterocyclic ring.

The substituted alkyl, substituted alkenyl, substituted aryl or substituted heterocyclic ring is
25 substituted with at least one of hydroxyl group, C1-8 alkyl,

C1-8 alkoxy, C2-8 alkenyl, C2-8 acyl, C2-8 acyloxy, C6-14 aryl, C6-14 aryloxy or C1-8 heterocyclic ring.

Preferable alkyl for R^1 is exemplified by C6-21 linear or branched alkyl such as $\text{CH}_3(\text{CH}_2)_5-$, $\text{cycloC}_6\text{H}_{11}\text{CH}_2-$,
 5 $\text{CH}_3(\text{CH}_2)_6-$, $\text{CH}_3(\text{CH}_2)_7-$, $\text{CH}_3(\text{CH}_2)_8-$, $\text{CH}_3(\text{CH}_2)_9-$, $\text{CH}_3(\text{CH}_2)_{10}-$,
 $\text{CH}_3(\text{CH}_2)_{11}-$, $\text{CH}_3(\text{CH}_2)_{12}-$, $\text{CH}_3(\text{CH}_2)_{13}-$, $\text{CH}_3(\text{CH}_2)_{11}\text{CH}(\text{CH}_3)-$,
 $\text{CH}_3(\text{CH}_2)_{14}-$, $\text{CH}_3(\text{CH}_2)_{15}-$, $\text{CH}_3(\text{CH}_2)_{16}-$, $\text{CH}_3(\text{CH}_2)_{17}-$,
 $\text{CH}_3(\text{CH}_2)_{18}-$, $\text{CH}_3(\text{CH}_2)_{19}-$ or $\text{CH}_3(\text{CH}_2)_{20}-$.

$\text{CH}_3(\text{CH}_2)_{12}-$ and $\text{CH}_3(\text{CH}_2)_{11}\text{CH}(\text{CH}_3)-$ are preferable.
 10 Hydroxy acids having the former and the latter (wherein m is 0 and n is 1) are 3-hydroxy-hexadecanoic acid and 3-hydroxy-4-methyl-hexadecanoic acid, respectively.

Preferable alkenyl for R^1 is exemplified by C6-21 linear or branched alkenyl such as $\text{CH}_3(\text{CH}_2)_4\text{CH}=\text{CH}-(\text{CH}_2)_2-$
 15 or $\text{CH}_3(\text{CH}_2)_5\text{CH}=\text{CH}(\text{CH}_2)_4\text{CH}=\text{CH}(\text{CH}_2)_3-$.

Preferable acyl portion of acylamino for R^1 is exemplified by C6-21 linear or branched aliphatic acyl or aromatic acyl such as $\text{CH}_3(\text{CH}_2)_4\text{CO}-$, $\text{CH}_3(\text{CH}_3)\text{CH}(\text{CH}_2)_2\text{CO}-$,
 $\text{CH}_3(\text{CH}_2)_5\text{CO}-$, $\text{cycloC}_6\text{H}_{11}\text{CH}_2\text{CO}-$, $\text{CH}_3(\text{CH}_2)_6\text{CO}-$, $\text{CH}_3(\text{CH}_2)_7\text{CO}-$,
 20 $\text{CH}_3(\text{CH}_2)_8\text{CO}-$, $\text{CH}_3(\text{CH}_2)_9\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{10}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{11}\text{CO}-$,
 $\text{CH}_3(\text{CH}_2)_{12}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{13}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{14}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{15}\text{CO}-$,
 $\text{CH}_3(\text{CH}_2)_{16}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{17}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{18}\text{CO}-$, $\text{CH}_3(\text{CH}_2)_{19}\text{CO}-$,
 $\text{C}_6\text{H}_5\text{CO}-$, $\text{C}_6\text{H}_5-\text{C}_6\text{H}_4\text{CO}-$ or $\text{C}_{12}\text{H}_7\text{CO}-$.

Preferable aryl for R^1 is exemplified by phenyl,
 25 mesityl, tolyl, naphthyl or anthryl.

Preferable heterocyclic ring for R^1 is exemplified by a group such as furil, thienyl, pyrazolyl, imidazolyl, tetrazolyl, thiazolyl, thiadiazolyl or isooxazolyl.

5 One having hydroxyl group as a substituent is exemplified by $\text{CH}_3(\text{CH}_2)_{11}\text{C}(\text{OH})(\text{CH}_3)-$. Hydroxy acid having the same (wherein m is 0 and n is 1) is 3,4-dihydroxy-4-methyl-hexadecanoic acid.

10 Preferable alkyl, or alkyl portion of alkoxy, as a substituent is exemplified by C1-8 linear or branched alkyl such as CH_3- , CH_3CH_2- , $\text{CH}_3(\text{CH}_2)_2-$, $\text{CH}_3(\text{CH}_2)_3-$, $\text{CH}_3(\text{CH}_3)\text{CHCH}_2-$, $\text{CH}_3(\text{CH}_2)_4-$, $\text{CH}_3(\text{CH}_3)\text{CH}(\text{CH}_2)_2-$, $\text{CH}_3(\text{CH}_2)_5-$, $\text{cycloC}_6\text{H}_{11}\text{CH}_2-$, $\text{CH}_3(\text{CH}_2)_6-$ or $\text{CH}_3(\text{CH}_2)_7-$.

15 Preferable alkenyl as a substituent is exemplified by C2-8 linear or branched alkenyl such as $\text{CH}_2=\text{CHCH}_2-$, $\text{CH}_2=\text{CH}(\text{CH}_2)_3-$ or $\text{CH}_2=\text{CHCH}_2-$.

20 Preferable aryl, or aryl portion of aryloxy, as a substituent is exemplified by C6-14 aryl such as phenyl, mesityl, tolyl, naphtyl or anthryl. These aryl may have at least one group selected from C1-8 linear or branched alkyl, halogen (e.g., fluoro, chloro, bromo or iodo), hydroxy or above-mentioned aryl.

25 Preferable acyl, or acyl portion of acyloxy, as a substituent is exemplified by C2-8 linear or branched aliphatic acyl or aromatic acyl such as $\text{CH}_3\text{CO}-$, $\text{CH}_3\text{CH}_2\text{CO}-$,

$\text{CH}_3(\text{CH}_2)_2\text{CO}-$, $\text{CH}_3(\text{CH}_2)_3\text{CO}-$, $\text{CH}_3(\text{CH}_3)\text{CHCH}_2\text{CO}-$, $\text{CH}_3(\text{CH}_2)_4\text{CO}-$,
 $\text{CH}_3(\text{CH}_3)\text{CH}(\text{CH}_2)_2\text{CO}-$, $\text{CH}_3(\text{CH}_2)_5\text{CO}-$, $\text{cycloC}_6\text{H}_{11}\text{CH}_2\text{CO}-$,
 $\text{C}_6\text{H}_5\text{CO}-$, $\text{C}_6\text{H}_5-\text{C}_6\text{H}_4\text{CO}-$ or $\text{C}_{12}\text{H}_7\text{CO}-$.

Preferable heterocyclic ring as a substituent is
 5 exemplified by C1-8 heterocyclic ring such as a group
 selected from furil, thienyl, pyrazolyl, imidazolyl,
 tetrazolyl, thiazolyl, thiadiazolyl or isooxazolyl.

Said AA^1 is an α - or β -amino acid having C1-7
 linear or branched alkyl group in a side chain, preferably
 10 alanine (which has CH_3-).

Said AA^2 is an α - or β -amino acid having C1-7
 linear or branched alkyl group substituted with amino group,
 preferably ornithine (which has $\text{H}_2\text{N}-(\text{CH}_2)_3-$) or lysine
 (which has $\text{H}_2\text{N}-(\text{CH}_2)_4-$).

15 Said AA^3 is an amino acid having hydrogen or C1-7
 linear or branched alkyl group, or an amino acid having H-
 or C1-7 linear or branched alkyl group having $\text{HO}-$, $\text{HOOC}-$ or
 $\text{H}_2\text{NOC}-$, preferably glycine (which has H-), alanine (which
 has CH_3-), serine (which has HO-CH_2-), threonine (which has
 20 $\text{CH}_3\text{CH}(\text{OH})-$), glutamic acid (which has $\text{HOOC}-(\text{CH}_2)_2-$),
 aspartic acid (which has HOOC-CH_2-), glutamine (which has
 $\text{H}_2\text{NOC}-(\text{CH}_2)_2-$) or asparagine (which has $\text{H}_2\text{NOC-CH}_2-$).
 Alternatively, AA^3 is β -alanine.

Said AA^4 is an amino acid having C1-7 linear or
 25 branched alkyl group substituted with hydroxyl group,

preferably serine (which has $\text{HO-CH}_2\text{-}$) or threonine (which has $\text{CH}_3\text{CH(OH)-}$).

Said AA^5 is an amino acid having C1-7 linear or branched alkyl group substituted with hydroxyl group, preferably serine (which has $\text{HO-CH}_2\text{-}$) or threonine (which has $\text{CH}_3\text{CH(OH)-}$).

Said AA^6 is an amino acid having C1-7 linear or branched alkyl group having $\text{H}_2\text{NOC-}$, preferably glutamine (which has $\text{H}_2\text{NOC-(CH}_2\text{)}_2\text{-}$) or asparagine (which has $\text{H}_2\text{NOC-CH}_2\text{-}$).

Said AA^7 is a D-amino acid having C1-7 linear or branched alkyl group substituted with hydroxyl group, preferably D-allo-threonine (which has $\text{CH}_3\text{CH(OH)-}$) or D- β -hydroxyvaline (which has $\text{CH}_3(\text{CH}_3)\text{C(OH)-}$).

Said AA^8 is an α - or β -amino acid having hydrogen or C1-7 linear or branched alkyl group, or an amino acid having C1-7 linear or branched alkyl group substituted with hydroxyl group, preferably glycine (which has H-), alanine (which has $\text{CH}_3\text{-}$) or threonine (which has $\text{CH}_3\text{CH(OH)-}$).

Typical examples of the novel cyclic peptides represented by Formula (1') of the present invention include the compounds as shown in Table 1:

Formula (1'):




Table 1

	R ₁	X m,n	AA ³	AA ⁴	AA ⁵	AA ⁷	AA ⁸
1	CH ₃ (CH ₂) ₁₂ -	O 0,1	βAla	L-Ser	L-Ser	D-alloThr	Gly
2	CH ₃ (CH ₂) ₁₂ -	O 0,1	L-Ala	L-Ser	L-Ser	D-alloThr	Gly
3	CH ₃ (CH ₂) ₁₂ -	O 0,1	L-Asn	L-Ser	L-Ser	D-alloThr	Gly
4	CH ₃ (CH ₂) ₁₂ -	O 0,1	L-Asp	L-Ser	L-Ser	D-alloThr	Gly
5	CH ₃ (CH ₂) ₁₂ -	O 0,1	L-Glu	L-Ser	L-Ser	D-alloThr	Gly
6	CH ₃ (CH ₂) ₁₂ -	O 0,1	Gly	L-Ser	L-Ser	D-alloThr	Gly
7	CH ₃ (CH ₂) ₁₂ -	O 0,1	L-Ser	L-Ser	L-Ser	D-alloThr	Gly
8	CH ₃ (CH ₂) ₁₂ -	O 0,1	βAla	L-Ser	L-Ser	D-alloThr	D-alloThr
9	CH ₃ (CH ₂) ₁₂ -	O 0,1	D-alloThr	L-Ser	L-Ser	D-alloThr	Gly
10	CH ₃ (CH ₂) ₁₂ -	O 0,1	D-alloThr	L-Ser	L-Ser	D-alloThr	D-alloThr
11	CH ₃ (CH ₂) ₁₂ -	O 0,1	βAla	D-alloThr	L-Ser	D-alloThr	Gly
12	CH ₃ (CH ₂) ₁₂ -	O 0,1	βAla	L-Ser	D-alloThr	D-alloThr	Gly
13	CH ₃ (CH ₂) ₁₂ -	O 0,1	βAla	L-Ser	L-Ser	D-βHOVal	Gly
14	CH ₃ (CH ₂) ₁₂ -	O 0,1	βAla	L-Ser	L-Ser	D-βHOPhe	Gly
15	CH ₃ (CH ₂) ₁₀ CONH-	O 1,0	βAla	L-Ser	L-Ser	D-alloThr	Gly
16	CH ₃ (CH ₂) ₁₀ CO ₂ CH ₂ -	NH 1,0	βAla	L-Ser	L-Ser	D-alloThr	Gly

A D-α-amino acid residue having β-hydroxy can be
 10 introduced into the compound of the present invention
 according to a method in which an amino acid having β-
 hydroxy which has been introduced beforehand is used as a
 material, or a method in which β-hydroxy is introduced into
 a cyclic peptide which has been synthesized such that β-
 15 hydroxy can be introduced thereafter. In either case, a
 compound of interest can be prepared, for example, using an
 aldol condensation reaction between one selected from
 various aldehydes or ketones and a glycine residue or the

like.

If the introduction of β -hydroxy is carried out by preparing a D- α -amino acid having β -hydroxy, for example, Boc-Gly-OBzl or Fmoc-Gly-OBzl can be used as a material, and one selected from various aldehydes or ketones (acetaldehyde, acetone, etc.) can be used as an electrophilic agent to be reacted with the material. For example, a D- α -amino acid having β -hydroxy can be prepared according to a method in which Fmoc-Gly-OBzl is reacted with a strong base, lithium diisopropylamide (LDA), in an aprotic solvent (e.g., THF) in the presence of a metal salt (e.g., lithium chloride) to convert Fmoc-Gly-OBzl into a lithium enolate form; or a method in which Fmoc-Gly-OBzl which has been converted into a silyl enol ether form using trimethylsilyl chloride is stirred with an excess amount of an aldehyde or a ketone as an electrophilic agent for several hours at a low temperature. Alternatively, a cyclic peptide that has a residue such as glycine or β -alanine and in which functional groups of constituting amino acids side chain are protected can be prepared, and then glycine or β -alanine residue is converted into a desired β -hydroxy- α -amino acid residue by the reaction as described above. Preferably, the preparation is carried out by conversion of glycine or β -alanine residue into the lithium enolate form using a strong base such as LDA

followed by a reaction with an aldehyde or a ketone.

Aprotic solvents that can be used include dichloromethane and ethyl ether in addition to THF. Metal salts that can be used include magnesium chloride, cesium chloride, titanium chloride, aluminum chloride and a boride in addition to lithium chloride. By appropriately selecting the above to control the stereochemistry of asymmetric center, it is possible to obtain a mixture of diastereomers at an appropriate ratio generated due to an asymmetric carbon newly resulted in a reaction product, or the only desired one of the diastereomers with high yield. Strong bases that can be used for generating a metal enolate include methyllithium and $(n\text{-Bu})_2\text{BOSO}_2\text{CF}_3$ in addition to LDA. Furthermore, it is important to control the reaction temperature in order to obtain the desired stereochemically pure product. The reaction is usually carried out at a low temperature of -80°C to -20°C .


The desired compound can be obtained as a single optically active substance or a mixture of diastereomers by purifying the reaction mixture using reverse phase high performance liquid chromatography or the like. The desired compound can be obtained by removal, according to a known method or application thereof, of protecting groups used for the functional groups of constituting amino acids side chain and the functional groups on the aldehyde or the

ketone used as a material.

The compound of the present invention can be prepared according to a total synthesis utilizing a liquid phase synthetic method or a solid phase synthetic method.

5 In either of the synthetic methods, a method for cyclizing a linear peptide (e.g., one represented by general formula (2)) can be preferably utilized. A method in which two or three peptide fragments are synthesized and then condensed each other is desirable in order to prepare
10 such a linear peptide via a total synthesis utilizing a liquid phase synthetic method. Various linear peptides can be conveniently prepared also via a total synthesis utilizing a solid phase synthetic method.

15 Formula (2):



The total synthesis utilizing a solid phase
20 synthetic method is described with respect to Compound 1 represented by Formula (3) which is an example of a compound of general Formula (1) wherein $m = 0$; $n = 1$, R^1 is $\text{CH}_3(\text{CH}_2)_{12}\text{-}$; AA^1 is L-Ala; AA^2 is L-Orn; AA^3 is β -Ala; AA^4 is L-Ser; AA^5 is L-Ser; AA^6 is D-Asn; AA^7 is D-allo-Thr;
25 and AA^8 is Gly.

Formula (3) (Compound 1):

Cyclo[(±)-Hhd→L-Ala→L-Orn→β-Ala→L-Ser→L-Ser→D-Asn→D-allo-
5 Thr→Gly]

wherein (±)-Hhd represents (±)-hydroxyhexadecanoic acid;
and L-Orn represents L-ornithine.

A fragment Fmoc-Gly-Hhd-OH which has been
10 prepared beforehand and contains an ester bond is
immobilized according to a known method on a resin bead
commonly used for solid phase synthesis. A protected
linear peptide represented by Formula (4) is then
synthesized on a solid carrier such as a resin bead using a
15 conventional automated peptide synthesizer.

Formula (4):

H-L-Ala-L-Orn(Boc)-β-Ala-L-Ser(t-Bu)-L-Ser(t-Bu)-D-Asn-D-
20 allo-Thr-Gly-(±)-Hhd-OH

The protected linear peptide is cleaved from the
resin using an appropriate known cleavage reagent, and then
subjected to a cyclization reaction to obtain a protected
25 cyclic peptide represented by Formula (5). The Compound 1

of interest which is represented by Formula (3) is finally obtained by deprotection of protecting groups of the amino acid side chain contained in the protected cyclic peptide, using an appropriate known deprotection reagent.

5

Formula (5):

Cyclo[(±)-Hhd→L-Ala→L-Orn(Boc)→β-Ala→L-Ser(t-Bu)→L-Ser(t-Bu)→D-Asn→D-allo-Thr→Gly]

10

Any commercially available resins used for peptide synthesis or combinatorial synthesis can be used as a resin for the solid phase synthetic method for a synthesis of the linear peptide of general Formula (2). Preferably, 2-chlorotrityl-chloride resin (Cl-Trt(2-Cl)-Resin, 100-200 mesh, 1% divinylbenzene (DVB) crosslinkage) as described in Barlos et al. (Tetrahedron Letters, 30:3947, 1989) is used.

Formation of an ester bond between Gly⁹ and (±)-Hhd¹ is carried out, for example, according to a method in which a coupling agent such as PyBroP or HATU is used, preferably, a method in which dicyclohexylcarbodiimide (DCC) is used in the presence of a catalyst such as dimethylaminopyridine or 4-pyrrolidinopyridine.

25

A reaction for immobilization of a fragment such

as Fmoc-Gly-(±)-Hhd-OH, containing an ester bond between Gly⁹ and (±)-Hhd¹ synthesized as described above, onto 2-chlorotrityl-chloride resin (Cl-Trt(2-Cl)-Resin) is carried out in an organic solvent (e.g., dimethylformamide, tetrahydrofuran or dichloromethane) in the presence of a base (e.g., triethylamine, N-methylmorpholine or diisopropylethylamine) at a temperature of 0°C to 50°C. Preferably, the reaction is carried out in dichloromethane using diisopropylethylamine as a base. The choice of the solvent used for the reaction gives serious influence to the immobilization yield. Then, dichloromethane which is a solvent that results in a high swelling rate of a resin is preferably selected.

Elongation of a peptide chain on a resin can be carried out according to the Fmoc method or the Boc method, which may be carried out manually or using an automated synthesizer. Preferably, the elongation is carried out according to the Fmoc method using an automated synthesizer. An exemplary protocol of peptide synthesis according to the Fmoc method using an automated peptide synthesizer is shown in Figure 1.

The water-soluble carbodiimide (WSCD; EDC)-HOBt method, the PyBOP-HOBt method, the HATU method or the like can be used as a method for coupling of amino acids in the presence of a base such as diisopropylethylamine or N-

methyldmorpholine. Preferably, the coupling is carried out according to the PyBOP-HOBt method using N-methyldmorpholine as a base.

Although a reaction of selective deprotection of
5 N-Fmoc group proceeds using a wide range of bases, it is carried out using 10 to 100% piperidine, preferably a 30% piperidine/dimethylformamide solution.

After a synthesis according to the Fmoc method using an automated synthesizer is completed, a cleavage
10 reaction of the obtained peptide from a resin is usually carried out using a solution prepared by adding anisole, thioanisole, ethanedithiol or the like as a scavenger to trifluoroacetic acid, hydrochloric acid, trifluoromethanesulfonic acid or the like. Generally, the
15 protecting groups of amino acid side chain are also removed at the same time. If a peptide having protected amino acid is to be cleaved from a resin, the cleavage is carried out preferably using a mixture of acetic acid - trifluoroethanol - methylene chloride.

20 Both the amino group at the N terminus and the carboxyl group at the C terminus in the protected peptide cleaved from the resin are free groups. A cyclization between these free groups can be carried out by converting the peptide into a peptide having an activated carboxyl
25 group at the C terminus, and then subjecting it to a

reaction in the presence of a base. Generally, a temperature ranging from -30 to 60°C, preferably from -10 to 30°C is selected as a reaction temperature for the above-mentioned cyclization reaction although it varies depending on the selected activating group of carboxyl group. The reaction time for the cyclization is ranging from several minutes to several days, preferably from 30 minutes to four hours.

In the cyclization, for example, a method of conversion into N-hydroxysuccinimide (ONSu) or a method of activation with PyBroP, HATU or DCC-HOAt can be used for activating the carboxyl group at the C terminus. DIEA, triethylamine, pyridine or the like can be used as a base. Tetrahydrofuran, methylene chloride, dimethylformamide, dimethyl sulfoxide or the like can be used as a solvent for cyclization. Dimethyl sulfoxide is preferably used in a cyclization of the linear peptide of Formula (4).

If the carboxyl group at the C terminus is to be activated with HATU, the activated peptide can be obtained by reacting the peptide of Formula (4) (i.e., one having amino group in a free or salt form at the N terminus and free carboxyl group at the C terminus) with HATU in the presence of a base in a solvent (e.g., dimethyl sulfoxide) at a temperature ranging from -10 to 30°C. In this case, the reaction proceeds accompanying cyclization. The

cyclization yield can be further increased, for example, by precise control of the concentrations of the respective reaction components in the reaction mixture or the reaction temperature, or by appropriate selection of the solvent.

5 The compound of the present invention is prepared by cleaving a peptide having protected amino acid from a resin, carrying out a cyclization and then removing the protecting groups on the amino acid side chains using a mixture of an acid and a scavenger as described above.

10 Thus, the novel cyclic peptides whose minimum inhibitory concentrations (MICs, $\mu\text{g/ml}$) for *Candida albicans*, *Cryptococcus neoformans* and *Aspergillus fumigatus* are very low and having no toxicological effect can be obtained.

15 Description about salt form useful as medicine

 The cyclic peptide of the present invention can be used for medicine as it is or as a pharmacologically acceptable salt thereof. There is no specific limitation concerning the salt as long as it is pharmacologically
20 acceptable. Examples thereof include salts with mineral acids (hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, hydrofluoric acid, hydrobromic acid, etc.); organic acids (formic acid, acetic acid, tartaric acid, lactic acid, citric acid, fumaric acid, maleic acid,
25 succinic acid, methanesulfonic acid, ethanesulfonic acid,

benzenesulfonic acid, toluenesulfonic acid, naphthalenesulfonic acid, camphorsulfonic acid, etc.); and alkaline metals or alkaline earth metals (sodium, potassium, calcium, etc.).

5 If the cyclic peptide or a pharmacologically acceptable salt thereof of the present invention is to be administered as a medicine, it is administered as it is or as formulated into a pharmaceutical composition containing the compound, for example, at a concentration of 0.1 to
10 99.5%, preferably, 0.5 to 90% in a pharmaceutically acceptable, nontoxic and inert excipient to animals including humans.

 Examples of excipient include solid, semisolid or liquid diluents, fillers and other formulation auxiliaries.
15 One or more of them can be used.

 It is preferable to administer the said pharmaceutical composition in a unit dosage form. The said pharmaceutical composition can be administered orally, interstitially, topically (e.g., transdermally) or
20 transrectally. Of course, the pharmaceutical compositions should be administered in dosage forms suitable for the respective routes of administration.

 The dosage as an antifungal agent of the cyclic peptide or a pharmacologically acceptable salt thereof of
25 the present invention to be administered as a medicine

should be desirably adjusted according to the conditions of the patient (age, weight, etc.), the administration route, the nature and severity of the disease and the like. Usually, the amount of the active ingredient of the present invention to be administered to an adult human is 10 to 2000 mg/day. A dosage less than the above may be sufficient in some cases, while a dosage more than the above may be required in other cases. If a large amount is to be administered, the dosage may be preferably administered in portions several times a day.

The oral administration can be carried out using a solid, powder or liquid unit dosage form such as epipastics, powders, tablets, sugar coated tablets, capsules, drops, sublingual tablets or other dosage forms.

The interstitial administration can be carried out, for example, using a liquid unit dosage form (e.g., a solution or suspension) for subcutaneous, intramuscular or intravenous injection. Such dosage forms can be prepared by suspending or dissolving a predetermined amount of the cyclic peptide or a pharmacologically acceptable salt thereof of the present invention in an injectable nontoxic liquid carrier such as an aqueous or oleaginous medium, and then sterilizing the resulting suspension or solution.

The topical administration (e.g., transdermal administration) can be carried out using an external dosage

form such as liquid, cream, powder, paste, gel or ointment. Such a dosage form can be prepared by mixing a predetermined amount of the cyclic peptide or a pharmacologically acceptable salt thereof of the present invention with one or more of flavor, coloring agent, 5 filler, detergent, humectant, emollient, gelating agent, vehicle, preservative, stabilizer and the like which are acceptable for an external dosage form.

The transrectal administration can be carried out, 10 for example, using a suppository prepared by mixing a predetermined amount of the cyclic peptide or a pharmacologically acceptable salt thereof of the present invention with the solid having a low melting point such as higher esters (e.g., palmitic acid myristyl ester), 15 polyethyleneglycol, cacao fat or a mixture thereof.

The physicochemical and biological properties of typical compounds according to the present invention are as follows.

(1) Physicochemical properties

20 The physicochemical properties of the typical compounds 1-7 in the present invention as shown in Table 1 above are shown in Table 2 below.

Table 2

Compound no.	Molecular formula	FAB-MS
1	$C_{43}H_{76}N_{10}O_{14}$	957
2	$C_{43}H_{76}N_{10}O_{14}$	957
3	$C_{44}H_{77}N_{11}O_{15}$	1000
4	$C_{44}H_{76}N_{10}O_{16}$	1001
5	$C_{45}H_{78}N_{10}O_{16}$	1015
6	$C_{42}H_{74}N_{10}O_{14}$	943
7	$C_{43}H_{76}N_{10}O_{15}$	973

(2) Biological properties

The cyclic peptide of the present invention
5 exhibits high antifungal activities on *Candida albicans*,
Cryptococcus neoformans and *Aspergillus fumigatus* which are
fungi typically used for determining an antifungal activity.
For example, Compound 1 at a concentration of 12.5 µg/ml or
less can inhibit the growth of *Candida albicans* and
10 *Cryptococcus neoformans*.

No toxicity was observed when 100 mg/kg each of
the compounds was administered intraperitoneally to mice.

Examples

The following Examples illustrate the present
15 invention in more detail, but are not to be construed to
limit the scope thereof.

Example 1: Synthesis of cyclo(-L-Ala-L-Orn-β-Ala-

L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-(±)-Hhd-) (compound 1)

1) Phenacyl (±)-3-hydroxyhexadecanoate ((±)-Hhd-OPac) (Pac: phenacyl)

To a suspension of (±)-3-hydroxyhexadecanoic acid
5 (500 mg, 1.84 mmol) in ethyl acetate (3.6 ml) were added
phenacyl bromide (421 mg, 2.12 mmol) and triethylamine (294
mg, 2.12 mmol). The mixture was stirred at room
temperature overnight. The reaction mixture was washed
10 with water, saturated aqueous sodium hydrogencarbonate and
saturated aqueous sodium chloride, and then dried over
magnesium sulfate. Magnesium sulfate was filtered off and
the filtrate was concentrated under reduced pressure to
obtain 678 mg of the title compound as a pale yellow solid.

2) Fmoc-Gly-(±)-Hhd-OPac

15 To a suspension of (±)-Hhd-OPac prepared in 1)
above (300 mg, 0.77 mmol) in methylene chloride (2 ml) were
added Fmoc-Gly-OH (274 mg, 0.92 mmol),
dimethylaminopyridine (28.2 mg, 0.23 mmol) and
dicyclohexylcarbodiimide (190 mg, 0.92 mmol). The mixture
20 was stirred at room temperature overnight. Ethyl acetate
was added to the reaction mixture. The mixture was washed
with 10% citric acid, saturated aqueous sodium
hydrogencarbonate and saturated aqueous sodium chloride,
and then purified on silica gel medium pressure column
25 chromatography (40g, eluted with CHCl₃) to obtain 563 mg of

the title compound as a colorless solid. FAB-MS: m/z 670 [MH]⁺.

3) Fmoc-Gly-(±)-Hhd-OH

To a solution of Fmoc-Gly-(±)-Hhd-OPac prepared in 2) above (500 mg, 0.75 mmol) in 90% acetic acid/water (37.5 ml) were added zinc dust. The mixture was stirred at room temperature for 1.5 hours. The insoluble matter was then filtered off. Ethyl acetate was added to the filtrate. The mixture was washed with 10% citric acid, saturated aqueous sodium hydrogencarbonate and saturated aqueous sodium chloride to obtain 370 mg of the title compound.

4) Fmoc-Gly-(±)-Hhd-Trityl(2-Cl)-resin

To a solution of Fmoc-Gly-(±)-Hhd-OH prepared in 3) above (16.5 mg, 0.03 mmol) in methylene chloride (0.5 ml) were added 2-chlorotrityl-chloride-resin (50 mg, 100-200 mesh, Watanabe Kagaku Kogyo) and diisopropylethylamine (17.4 µl, 0.09 mmol). The mixture was allowed to stand at room temperature for 30 minutes. Methanol (40 µl) was added to the reaction mixture. The resin was then collected by filtration, washed with methylene chloride, dimethylformamide, isopropyl alcohol, methanol and diethyl ether, and dried under reduced pressure.

Calculation of amount of immobilized Fmoc-Gly-(±)-Hhd-OH: A 50% piperidine/methylene chloride solution (1 ml) was added to 10 mg of the resin prepared in 4) above.

The mixture was gently stirred at room temperature for 30 minutes. The reaction mixture was filtrated. The filtrate and the solvent used for washing the resin (methylene chloride) were combined and concentrated under reduced pressure. The residue was dissolved in methylene chloride (20 ml) and the solution was diluted 25 times with methylene chloride. The absorbances (A267 and A301) of the solution were measured to be 0.167 and 0.079, respectively. According to a conversion equation between the absorbance and the immobilized amounts, the amounts were calculated to be 0.48 mmol/g resin (using A267 data) and 0.51 mmol/g resin (using A301 data), respectively.

5) Fmoc-L-Ala-L-Orn(Boc)- β -Ala-L-Ser(t-Bu)-L-Ser(t-Bu)-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-Trtyl(2-Cl)-resin

Synthesis was carried out using 50 mg of the resin prepared in 4) above, Fmoc-D-allo-Thr-OH, Fmoc-D-Asn-OH, Fmoc-L-Ser(t-Bu)-OH, Fmoc-L-Ser(t-Bu)-OH, Fmoc- β -Ala-OH, Fmoc-L-Orn(Boc)-OH and Fmoc-L-Ala-OH (the amount of each Fmoc-amino acid used was 10 times compared with the amount of immobilized Fmoc-Gly-($\pm$)-Hhd-OH), and an automated peptide synthesizer PSSM-8 (Shimadzu) according to a conventional protocol of automated synthesis by the PyBOP-HOBt method. After confirming the completion of the synthetic protocol, the resin was washed with isopropyl alcohol and methanol, and dried under reduced pressure.

6) H-L-Ala-L-Orn(Boc)- β -Ala-L-Ser(t-Bu)-L-Ser(t-Bu)-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-OH

The resin obtained in 5) above was suspended in a mixture of acetic acid - trifluoroethanol - methylene chloride (1 : 1 : 8) (0.8 ml). The suspension was allowed to stand at room temperature for one hour. The resin was filtered off. The filtrate was concentrated under argon flow, and diethyl ether was then added thereto. The deposited crystals were collected by centrifugation (10,000 rpm, 5 minutes) to obtain 19.5 mg of the title compound. FAB-MS: m/z 1429[MH]⁺.

7) Cyclo(-L-Ala-L-Orn(Boc)- β -Ala-L-Ser(t-Bu)-L-Ser(t-Bu)-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-)

A solution of H-L-Ala-L-Orn(Boc)- β -Ala-L-Ser(t-Bu)-L-Ser(t-Bu)-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-OH obtained in 6) above (15 mg, 10.5 μ mol) in dimethyl sulfoxide (5.25 ml) was added to a solution of HATU (27.9 mg, 73.5 μ mol) and diisopropylethylamine (12.8 μ l, 73.5 μ mol) in dimethyl sulfoxide (5.25 ml), over five hours. The mixture was further stirred at room temperature for two hours, and lyophilized. FAB-MS: m/z 1411[MH]⁺, 1433[MNa]⁺.

8) Cyclo(-L-Ala-L-Orn- β -Ala-L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-)

A mixture of trifluoroacetic acid - anisole - ethanedithiol (94 : 5 : 1) (1 ml) was added to a residue

containing cyclo(-L-Ala-L-Orn(Boc)- β -Ala-L-Ser(t-Bu)-L-Ser(t-Bu)-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-) obtained in 7) above. The mixture was allowed to stand at room temperature for two hours. The reaction mixture was concentrated under argon flow, and ethyl ether was added thereto. The deposited crystals were collected by centrifugation (10,000 rpm, 5 minutes), re-dissolved in dimethyl sulfoxide and lyophilized to obtain the title compound. FAB-MS: m/z 957[MH]⁺.

Elementary analysis: theoretical (as C₄₃H₇₆N₁₀O₁₄·2H₂O), C: 52.00%, H: 8.12%, N: 14.13%; analytical, C: 51.96%, H: 8.15%, N: 14.10%.

Example 2: Synthesis of cyclo(-L-Ala-L-Orn-L-Ala-L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-) (Compound 2)

The title compound was obtained according to the procedure as described in Example 1, 5) to 8) except that Fmoc-L-Ala-OH was used in place of Fmoc- β -Ala-OH in 5). FAB-MS: m/z 957[MH]⁺.

Example 3: Synthesis of cyclo(-L-Ala-L-Orn-L-Asn-L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-) (Compound 3)

The title compound (4.8 mg) was obtained according to the procedure as described in Example 1, 5) to 8) except that Fmoc-L-Asn-OH was used in place of Fmoc- β -Ala-OH in 5). FAB-MS: m/z 1000[MH]⁺.

Example 4: Synthesis of cyclo(-L-Ala-L-Orn-L-Asp-

L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-(±)-Hhd-) (Compound 4)

The title compound was obtained according to the procedure as described in Example 1, 5) to 8) except that Fmoc-L-Asp-OH was used in place of Fmoc-β-Ala-OH in 5).

5 FAB-MS: m/z 1001[MH]⁺.

Example 5: Synthesis of cyclo(-L-Ala-L-Orn-L-Glu-L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-(±)-Hhd-) (Compound 5)

The title compound was obtained according to the procedure as described in Example 1, 5) to 8) except that

10 Fmoc-L-Glu-OH was used in place of Fmoc-β-Ala-OH in 5).

FAB-MS: m/z 1015[MH]⁺.

Example 6: Synthesis of cyclo(-L-Ala-L-Orn-Gly-L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-(±)-Hhd-) (Compound 6)

The title compound was obtained according to the procedure as described in Example 1, 5) to 8) except that

15 Fmoc-Gly-OH was used in place of Fmoc-β-Ala-OH in 5). FAB-MS: m/z 943[MH]⁺.

Example 7: Synthesis of cyclo(-L-Ala-L-Orn-L-Ser-L-Ser-L-Ser-D-Asn-D-allo-Thr-Gly-(±)-Hhd-) (Compound 7)

20 The title compound was obtained according to the procedure as described in Example 1, 5) to 8) except that Fmoc-L-Ser(t-Bu)-OH was used in place of Fmoc-β-Ala-OH in 5). FAB-MS: m/z 973[MH]⁺.

Example 8: Determination of antifungal activities

25 The antifungal activities of the compounds

synthesized in Examples 1 and 3 to 7 were measured by a liquid medium dilution method.

Candida albicans TIMM0136, Cryptococcus neoformans TIMM0354 or Aspergillus fumigatus TIMM1776 was inoculated into a medium containing one of the compounds diluted serially, and cultured at 30°C for two days. The concentration at which the growth of the fungus was almost completely inhibited was defined as a minimum inhibitory concentration (MIC, µg/ml). For the measurements, a medium containing 0.67% Yeast Nitrogen Base (Difco) and 1.0% glucose (YNBG medium) was used for the cultures of Candida albicans and Cryptococcus neoformans, and a medium containing 0.5% Brain Heart Infusion Bouillon (Nissui) (BHI medium) was used for the culture of Aspergillus fumigatus. The results are shown in Table 3.

As shown in Table 3, Compounds 1, 3, 4, 5, 6 and 7 exhibited high antifungal activities on Candida albicans (C.a.), Cryptococcus neoformans (C.n.) and Aspergillus fumigatus (A.f.).

Table 3

Compound	MIC ($\mu\text{g/ml}$)		
	C.a.	A.f.	C.n.
1	6.25	1.56	12.5
3	6.25	1.56	3.12
4	6.25	-	6.25
5	3.12	0.39	6.25
6	12.5	-	3.12
7	6.25	0.78	3.12

Example 9: Synthesis of analogs of Compound 1 having substitution of L-Thr or D-Thr for D-allo-Thr in AA⁷,
 5 cyclo(-L-Ala-L-Orn- β -Ala-L-Ser-L-Ser-D-Asn-AA⁷-Gly-($\pm$)-Hhd-), and comparison of antifungal activities thereof

1) Synthesis of the AA⁷:L-Thr-substituted analog

A compound having substitution of L-Thr in AA⁷ in the compound 1 (the AA⁷:L-Thr-substituted analog) was
 10 obtained according to the procedure as described in Example 1, 5) to 8) except that Fmoc-L-Thr(tBu)-OH was used in place of Fmoc-D-allo-Thr-OH in 5). FAB-MS: m/z 957[MH]⁺.

2) synthesis of the AA⁷:D-Thr-substituted analog

A compound having substitution of D-Thr in AA⁷ in the compound 1 (the AA⁷:D-Thr-substituted analog) was
 15 obtained according to the procedure as described in Example 1, 5) to 8) except that Fmoc-D-Thr(t-Bu)-OH was used in

place of Fmoc-D-allo-Thr-OH in 5). FAB-MS: m/z 957[MH]⁺.

3) Comparison of antifungal activities

The antifungal activities of the AA⁷:L-Thr-substituted analog and the AA⁷:D-Thr-substituted analog
5 against *Candida albicans*, *Cryptococcus neoformans* and *Aspergillus fumigatus* were measured as described in Example 8. As shown in Table 3, the compound having D-allo-Thr without substitution in AA⁷ (Compound 1) showed antifungal activities against *Candida albicans*, *Cryptococcus*
10 *neoformans* and *Aspergillus fumigatus* with MICs of 6.25 µg/ml, 12.5 µg/ml and 1.56 µg/ml, respectively. The AA⁷:L-Thr-substituted analog or the AA⁷:D-Thr-substituted analog did not show the activity against any of the fungi (MIC > 200 µg/ml).

15 Example 10: Preparation of a mixture of AA⁴-substituted analogs, cyclo(-L-Ala-L-Orn-β-Ala-AA⁴-L-Ser-D-Asn-D-allo-Thr-Gly-(±)-Hhd-).

A mixture of twenty title cyclic peptides each having one of L-Ala, L-Arg, L-Asn, L-Asp, L-Cys, L-Gln, L-
20 Glu, L-Gly, L-His, L-Ile, L-Leu, L-Lys, L-Met, L-Phe, L-Pro, L-Ser, L-Thr, L-Trp, L-Tyr and L-Val at AA⁴ of Compound 1 was prepared.

A synthetic protocol was carried out according to the procedure as described in Example 1, 5) to 8) except
25 that a mixture of 1/20 amounts (moles) each of Fmoc-L-Ala-

OH, Fmoc-L-Arg(Pmc)-OH, Fmoc-L-Asn(Trt)-OH, Fmoc-L-Asp(Ot-Bu)-OH, Fmoc-L-Cys(Trt)-OH, Fmoc-L-Gln(Trt)-OH, Fmoc-L-Glu(Ot-Bu)-OH, Fmoc-L-Gly-OH, Fmoc-L-His(Trt)-OH, Fmoc-L-Ile-OH, Fmoc-L-Leu-OH, Fmoc-L-Lys(Boc)-OH, Fmoc-L-Met-OH, 5 Fmoc-L-Phe-OH, Fmoc-L-Pro-OH, Fmoc-L-Ser(t-Bu)-OH, Fmoc-L-Thr(t-Bu)-OH, Fmoc-L-Trp-OH, Fmoc-L-Tyr(t-Bu)-OH and Fmoc-L-Val-OH than Fmoc-L-Ser(t-Bu)-OH was used in place of Fmoc-L-Ser(t-Bu)-OH used for the AA⁴ amino acid in Example 1, 5). From the FAB-MS spectrum of the finally obtained 10 residue, it was confirmed that the twenty title cyclic peptides were obtained.

Example 11: Preparation of a mixture of AA⁵ amino acid-substituted analogs, cyclo(-L-Ala-L-Orn- β -Ala-L-Ser-AA⁵-D-Asn-D-allo-Thr-Gly-($\pm$)-Hhd-)

15 A mixture of twenty title cyclic peptides each having one of L-Ala, L-Arg, L-Asn, L-Asp, L-Cys, L-Gln, L-Glu, L-Gly, L-His, L-Ile, L-Leu, L-Lys, L-Met, L-Phe, L-Pro, L-Ser, L-Thr, L-Trp, L-Tyr and L-Val at AA⁴ of Compound 1 was prepared.

20 A synthetic protocol was carried out according to the procedure as described in Example 1, 5) to 8) except that a mixture of 1/20 amounts (moles) each of Fmoc-L-Ala-OH, Fmoc-L-Arg(Pmc)-OH, Fmoc-L-Asn(Trt)-OH, Fmoc-L-Asp(Ot-Bu)-OH, Fmoc-L-Cys(Trt)-OH, Fmoc-L-Gln(Trt)-OH, Fmoc-L-Glu(Ot-Bu)-OH, Fmoc-L-Gly-OH, Fmoc-L-His(Trt)-OH, Fmoc-L- 25

Ile-OH, Fmoc-L-Leu-OH, Fmoc-L-Lys(Boc)-OH, Fmoc-L-Met-OH, Fmoc-L-Phe-OH, Fmoc-L-Pro-OH, Fmoc-L-Ser(t-Bu)-OH, Fmoc-L-Thr(t-Bu)-OH, Fmoc-L-Trp-OH, Fmoc-L-Tyr(t-Bu)-OH and Fmoc-L-Val-OH than Fmoc-L-Ser(t-Bu)-OH was used in place of Fmoc-L-Ser(t-Bu)-OH for introducing the AA⁵ amino acid in 5) . The FAB-MS spectrum of the finally obtained residue was analyzed to confirm that the twenty title cyclic peptides were contained.

Example 12: Preparation of a mixture of AA⁶ amino acid-substituted analogs, cyclo(-L-Ala-L-Orn- β -Ala-L-Ser-L-Ser-AA⁶-D-allo-Thr-Gly-($\pm$)-Hhd-)

A mixture of thirteen title cyclic peptides each having one of D-Ala, D-Asn, D-Asp, D-Glu, D-His, D-Ile, D-Leu, D-Phe, D-Ser, D-Thr, D-Trp, D-Tyr and D-Val at AA⁶ of Compound 1 was synthesized.

A synthesis protocol was carried out according to the procedure as described in Example 1, 5) to 8) except that a mixture of 1/13 amounts (moles) of Fmoc-D-Ala-OH, Fmoc-D-Asn(Trt)-OH, Fmoc-D-Asp(Ot-Bu)-OH, Fmoc-D-Glu(Ot-Bu)-OH, Fmoc-D-His(Trt)-OH, Fmoc-D-Ile-OH, Fmoc-D-Leu-OH, Fmoc-D-Phe-OH, Fmoc-D-Ser(t-Bu)-OH, Fmoc-D-Thr(t-Bu)-OH, Fmoc-D-Trp-OH, Fmoc-D-Tyr(t-Bu)-OH and Fmoc-D-Val-OH was used in place of Fmoc-D-Asn(Trt)-OH for introducing the AA⁶ amino acid in Example 1, 5) . The FAB-MS spectrum of the finally obtained residue was analyzed to confirm that the

thirteen title cyclic peptides were contained.


Example 13: Antifungal activities of mixtures of AA⁴, AA⁵ or AA⁶ amino acid-substituted analogs of Compound 1

The antifungal activities of the mixtures of AA⁴,
5 AA⁵ or AA⁶ amino acid-substituted analogs obtained in
Examples 10, 11 and 12 against *Candida albicans*,
Cryptococcus neoformans and *Aspergillus fumigatus* were
measured as described in Example 8. The mixture of AA⁴
amino acid-substituted analogs exhibited activities on
10 *Cryptococcus neoformans* and *Aspergillus fumigatus* with MICs
of 100 µg/ml and 50 µg/ml, respectively. The mixture of
AA⁵ amino acid-substituted analogs exhibited antifungal
activities on *Candida albicans*, *Cryptococcus neoformans* and
Aspergillus fumigatus with MICs of 25 µg/ml, 50 µg/ml and
15 25 µg/ml, respectively. The mixture of AA⁶ amino acid-
substituted analogs exhibited antifungal activities on
Candida albicans, *Cryptococcus neoformans* and *Aspergillus*
fumigatus with MICs of 50 µg/ml, 50 µg/ml and 25 µg/ml,
respectively.

CLAIMS

1. A cyclic peptide represented by general formula (I) or a pharmacologically acceptable salt thereof:

5



10 wherein

m and n may be identical or different from each other, and are independently 0 or 1;

X is O or NH;

15 R¹ is hydrogen, C6-21 alkyl, substituted alkyl, alkenyl, substituted alkenyl, acylamino, aryl, substituted aryl, heterocyclic ring or substituted heterocyclic ring, wherein said substituted alkyl, substituted alkenyl, substituted aryl or substituted heterocyclic ring is substituted with at least one of hydroxyl group, C1-8 alkyl, C1-8 alkoxy, C2-8 alkenyl, C2-8 acyl, C2-8 acyloxy, C6-14
20 aryl, C6-14 aryloxy or C1-8 heterocyclic ring;

AA¹ to AA⁶ are independently an α- or a β-amino acid;

25 AA⁷ is a D-α-amino acid having HO- in a side chain; and

AA⁸ is an α - or a β -amino acid.

2. The cyclic peptide of general formula (I) or a pharmacologically acceptable salt thereof according to claim 1,

5 wherein

AA¹ is a hydrophobic α - or β -amino acid;

AA² is an α - or β -amino acid having amino group in a side chain;

10 AA³ is β -alanine, glycine, or an α - or β -amino acid having HO-, HOOC- or H₂NOC- in a side chain;

AA⁴ is an α - or β -amino acid having HO- in a side chain;

AA⁵ is an α - or β -amino acid having HO- in a side chain;

15 AA⁶ is an α - or β -amino acid having H₂NOC- in a side chain;

AA⁷ is a D- α -amino acid having HO- in a side chain; and

20 AA⁸ is glycine or an α - or β -amino acid having HO- in a side chain.

3. The cyclic peptide of general formula (I) or a pharmacologically acceptable salt thereof according to claim 1,

wherein

25 AA¹ is alanine, valine or leucine;

AA² is lysine or ornithine;

AA³ is β -alanine, glycine, or an α - or β -amino acid having HO-, HOOC- or H₂NOC- in a side chain;

5 AA⁴ is an α -amino acid having HO- in a side chain;

AA⁵ is an α -amino acid having HO- in a side chain;

AA⁶ is asparagine or glutamine;

10 AA⁷ is a D- α -amino acid having HO- in a side chain; and

AA⁸ is glycine or an α -amino acid having HO- in a side chain.

4. An antifungal pharmaceutical composition which contains the cyclic peptide or a pharmacologically acceptable salt thereof defined by claim 1.

15

Fetherstonhaugh & Co.
Ottawa, Canada
Patent Agents

1 / 1

Figure 1

