

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2007316639 B2**

(54) Title
Crystalline modification of fipronil

(51) International Patent Classification(s)
A01N 47/02 (2006.01) **A01P 7/00** (2006.01)
A01N 25/00 (2006.01)

(21) Application No: **2007316639** (22) Date of Filing: **2007.11.05**

(87) WIPO No: **WO08/055882**

(30) Priority Data

(31) Number	(32) Date	(33) Country
06023437.4	2006.11.10	EP

(43) Publication Date: **2008.05.15**

(44) Accepted Journal Date: **2012.12.20**

(71) Applicant(s)
BASF SE

(72) Inventor(s)
Saxell, Heidi Emilia;Erk, Peter;Bhatt, Prashant M.;Desiraju, Gautam R.;Krohl, Thomas;Cox, Gerhard;Banerjee, Rahul;Taranta, Claude

(74) Agent / Attorney
Watermark Patent and Trade Marks Attorneys, Level 2 302 Burwood Road, HAWTHORN, VIC, 3122

(56) Related Art
CAIRA M, Topics in Current Chemistry, 1998, 198, pages 163-208
WO 2000/062616 A1

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 May 2008 (15.05.2008)

PCT

(10) International Publication Number
WO 2008/055882 A1

(51) International Patent Classification:
A01N 47/02 (2006.01) *A01P 7/00* (2006.01)
A01N 25/00 (2006.01)

[IN/US]; 3130 Sawtelle Blvd, APT 207, Los Angeles, Ca 90066 (US). **BHATT, Prashant M.** [IN/IN]; H-201, N.R.S. Hostel, University of Hyderabad, Hyderabad 500046 (IN).

(21) International Application Number:
PCT/EP2007/061896

(74) Common Representative: **BASF SE**; 67056 Ludwigshafen (DE).

(22) International Filing Date:
5 November 2007 (05.11.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
06023437.4 10 November 2006 (10.11.2006) EP

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **BASF SE** [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SAXELL, Heidi Emilia** [FI/DE]; Am Weidenschlag 5, 67071 Ludwigshafen (DE). **ERK, Peter** [DE/DE]; Carl-Spitzweg-Str. 5a, 67227 Frankenthal (DE). **TARANTA, Claude** [FR/DE]; Hahnenfussweg 8, 76297 Stutensee (DE). **KRÖHL, Thomas** [DE/DE]; Steinachstr. 30, 69198 Schriesheim (DE). **COX, Gerhard** [DE/DE]; Birkental 20, 67098 Bad Dürkheim (DE). **DESIRAJU, Gautam R.** [IN/IN]; B-6 Campus Quarters, Hyderabad 500 046 (IN). **BANERJEE, Rahul**

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(54) Title: CRYSTALLINE MODIFICATION OF FIPRONIL


(57) Abstract: The present invention relates to a crystalline modification of fipronil, to a process for the preparation of the same, to pesticidal and parasiticidal mixtures and compositions comprising said crystalline modification and to their use for combating pests and parasites.

WO 2008/055882 A1

CRYSTALLINE MODIFICATION OF FIPRONIL

The present invention relates to a novel crystalline modification of fipronil, to a process for the preparation of the same, to pesticidal and parasiticial mixtures and compositions comprising said crystalline modification and to their use for combating pests and parasites.

Fipronil (formula I) is an active compound for controlling certain insect and acarid pests, and parasites.



Various processes for the preparation of fipronil have been described, generally and in detail. Documents which give detailed preparation procedures are e.g. EP 295 117; EP 460 940; EP 484 165; EP 668 269; EP 967 206; EP 1 331 222; EP 1 374 061; US 5 631 381; CN 1374298; or J. of Heibei University of Science and Technology, Vol. 25 (2), Sum 69 (2004), Dok. Serial No. 1008-1542 (2004) 02-0018-03.

Characterization of the fipronil material obtained by the processes described in the prior art is usually done by ¹H-NMR analysis and/ or measurement of the melting point. The described melting points are in the range of from 187°C to 203°C, mostly in the range of from 195°C to 203°C. In the Pesticidal Manual, 13th Edition (2003), British Crop Protection Council, p.433, fipronil is described as a white solid with a melting point of 200 to 201°C, with technical fipronil having a melting point of 195.5 °C to 203°C. Observations of different crystalline forms of fipronil have not been described, let alone any characterization of a certain crystalline modification or a preparation procedure for obtaining a certain crystalline modification.

For the large-scale preparation and formulation of a market compound such as fipronil, it is of crucial importance to know whether different crystalline modifications (also frequently referred to as polymorphs) of a compound exist, how they can be obtained, and what their characteristic properties are. Crystalline modifications of one compound may have very different properties, for example with regard to solubility, rate of dissolution, suspension stability, stability during grinding, vapour pressure, optical and mechanical properties, hygroscopicity, crystal size, filtration properties, desiccation, density, melting point, degradation stability, stability against phase transformation into other crystalline modifications, colour, and even chemical reactivity.

For example, different crystalline modifications frequently manifest themselves in different forms of the crystals, such as needles or plates. This is of relevance for e.g a filtration step in the preparation procedure. Plates typically will clog the pores of a filter leading to loss of time and product and tedious and expensive cleaning work. Also, a crystalline modification being present as plates and a crystalline modification being present as needles can have significantly different bulk densities which has implications for storage and packaging. Another relevant aspect, especially in the production of pesticides, is whether the crystalline modification is present as a fine powder which can produce hazardous dusts, or as dust-free larger crystals.

Against this background, it has been an object of the present invention to find and characterize a novel crystalline modification of fipronil.

A preferred embodiment includes preparation procedures for the novel crystalline modification which reproducibly give the crystalline modification.

Another preferred embodiment includes preparation procedures which give the novel crystalline modification II in high yield.

Yet another preferred embodiment includes preparation procedures which give the novel crystalline modification essentially excluding other crystalline modification forms (i.e. in over 80% by weight).

Accordingly, a novel crystalline modification of fipronil, a process for its preparation, pesticidal and parasitocidal mixtures and compositions comprising it and its use for combating pests and parasites has been found. The novel crystalline modification of fipronil is defined as "novel crystalline modification II" throughout this application. It is present as long needles which provides for its easy filtration.

Also, most suprisingly, 3 other crystalline modifications of fipronil have been found, which are subject to co-pending patent applications. Especially

surprising was that the present crystalline modification II of fipronil does not melt but rather undergoes a phase transformation during heating into two thermodynamically more stable forms I and V and/or a mixture of them, and thus in a typical melting point measurement will give the melting points of these forms I and V or mixtures of them. Crystalline modification I has a very similar melting point as a second crystalline modification V (as described in co-pending patent applications), both melting points lying in the range of the melting points given in the prior art (i.e. 195 to 203°C) . Moreover, one further crystalline modification IV of fipronil, as described in a co-pending patent application, also undergoes a phase transformation. The solid forms of fipronil thus are part of a very complex crystallization scenario. It can be concluded that the melting points given in the literature in no way

can indicate which crystalline modification or crystalline modification mixtures were analyzed.

In T 605/02, the Technical Board of Appeal of the European Patent Authority ruled that, in the absence of a respective described preparation procedure, even the XRD pattern of a certain crystalline modification does not constitute prior art for lack of enablement. Thus, melting points given in documents published prior to the filing of this application cannot be regarded as prior art for the present invention as they do not enable the artisan to prepare the novel crystalline modification of fipronil.

10

Crystalline modification II of fipronil, in an X-ray powder diffractogram at 25°C, shows at least 4, in particular at least 5, especially 7 and preferably all of the following reflexes:

	(1) $d = 13.44 \pm 0.2 \text{ \AA}$	$2 \Theta = 6.6 \pm 0.2^\circ$
15	(2) $d = 7.84 \pm 0.1 \text{ \AA}$	$2 \Theta = 11.3 \pm 0.2^\circ$
	(3) $d = 5.50 \pm 0.07 \text{ \AA}$	$2 \Theta = 16.1 \pm 0.2^\circ$
	(4) $d = 5.14 \pm 0.05 \text{ \AA}$	$2 \Theta = 17.2 \pm 0.2^\circ$
	(5) $d = 4.95 \pm 0.05 \text{ \AA}$	$2 \Theta = 17.9 \pm 0.2^\circ$
	(6) $d = 3.95 \pm 0.05 \text{ \AA}$	$2 \Theta = 22.4 \pm 0.2^\circ$
20	(7) $d = 3.77 \pm 0.05 \text{ \AA}$	$2 \Theta = 23.5 \pm 0.2^\circ$
	(8) $d = 3.22 \pm 0.03 \text{ \AA}$	$2 \Theta = 27.6 \pm 0.2^\circ$
	(9) $d = 2.91 \pm 0.03 \text{ \AA}$	$2 \Theta = 30.8 \pm 0.2^\circ$

In a particularly preferred embodiment, the crystalline modification II exhibits a powder X-ray diffraction pattern substantially the same as the pattern shown in Figure 1.

Studies of single crystals of the crystalline modification II have shown that the crystal system is monoclinic and has the space group P 2(1)/c. The characteristic data of the crystal structure of the crystalline modification II are shown in Table 1:

30

Table 1: Crystallographic data and parameters of the crystalline modification II measured from

Parameter	Modification II
Class	Monoclinic
Space group	P 2(1)/c
a	8.606(1) Å
b	26.919(2) Å
c	16.086(1) Å
α	90°
β	102.066(1)°
γ	90°

Parameter	Modification II
Volume	3644.0(3) Å ³
Z	4
Temperature	-173.2°C
Density (calculated)	0.94 g/cm ³ ^a
R1, ωR2	0.081, 0.222

a,b,c = Length of the unit cell edges

α,β,γ = Angles of the unit cell

Z = Number of molecules in the unit cell

^aDensity calculated without the incorporated solvent

5

The single crystal structure further reveals, that modification II of fipronil has channel like tubes running through the crystal. These tubes are occupied by the corresponding crystallization solvent. This affects the powder diffraction patterns in a way that the X-ray reflections can be seen at slightly different 2Theta and d-spacing values (not more than ± 0.34° 2Theta) for many of the solvates. The peak intensities are independent of the incorporated solvent[B1]. Modification II can therefore be regarded as a solvate of the corresponding co-crystallized solvent.

As the solvents trapped in the crystal are removed, the crystalline modification II of fipronil undergoes an endothermic phase transformation. In a differential scanning calorimetry, DSC, measurement (heating rate 5 K / min) this is observed typically at around 130°C, with an onset at 100 to 110°C and completion at 138°C to 145°C. This phase transformation leads to crystalline modifications I and V and/or depending on the heating rate to mixtures of them. We give here as an example a differential scanning calorimetry (DSC) thermogram of the crystalline modification II of fipronil, recorded with 10 K/ min. It further contains two endotherms with maxima at 196°C and 203°C, representing the melting points of the crystalline modifications I (mp. 196°C) and V (mp. 203°C) of fipronil. It is shown in figure 2.

Solvents in the process for preparation of modification II are inert and consist mainly of benzene derivatives, such as benzene, which may be substituted by one or more groups selected from halogen, cyano, C₁-C₆-alkyl, C₁-C₆-alkoxy, halogenmethyl, and nitro, and other solvents, such as tetrahydrofuran (THF), 1,2-dichloroethane, and acetonitrile. Preferred benzene derivatives are fluorobenzene, benzonitrile, anisole, p-xylene, o-xylene, m-xylene, CF₃-benzene, n-butylbenzene, t-butylbenzene, s-butylbenzene, i-butylbenzene, chlorobenzene, 2-chlorotoluene, 4-chlorotoluene, 1,2-dichlorobenzene, 1,3-dichlorobenzene, 1,4-dichlorobenzene, 1,4-diisopropylbenzene, mesitylene, nitrobenzene, 4-nitrotoluene, n-propylbenzene, and toluene, and more preferably selected from mono-, di- or tri(C₁-C₆-alkyl) benzenes[B2], which may be halogenated.

The crystalline modification II of fipronil can be achieved preferably from the following solvents; tetrahydrofuran, 1,2-dichloroethane, acetonitrile, toluene, mono chloro benzene, 1,2-dichlorobenzene, ethyl benzene, mesitylene, nitrobenzene and CF₃-benzene.

The amount of co-crystallized solvent depends of the properties, such as size of the corresponding solvent. For toluene the maximum amount of solvent in the crystals is in the range of 5 w-%. The solvent amount can be analysed for example via thermogravimetry. A TGA-trace of crystalline fipronil modification II from toluene is shown in figure 3.

In addition to the various solvent adduct versions of modification II, it was further discovered, that from all of the above mentioned solvents also at least one other crystalline modification can be achieved via solvent crystallization. Several conditions lead also to mixtures of forms, but also polymorph or modification pure material can be prepared by applying delicate and controlled, solvent dependant methods. The applicant has described preparation methods of modifications I and V in co-pending patent applications[B3].

Also a further modification F-ST, a toluene solvate, is described in WO2007/069254. This solvate modification F-ST is different from the here described novel modification II which also contains co-crystallized toluene. This should be made clear by comparison of the powder diffraction patterns[B4] shown in Figure 1, and in WO2007/069254. It appears that F-ST is a stoichiometric solvate of fipronil and toluene in the ratio of about 2:1 respectively. In the case of toluene this means that there is about 8 w-% of toluene in the crystal lattice, which is about 3 w-% more than for crystalline modification II according to the current invention.

Further, structurally similar solvates to F-ST form can be prepared also with benzene, mono chloro benzene (MCB) as well as xylene (mixture).

Similar to F-ST, also crystalline modification II of fipronil can be used as starting material to prepare pure modification V via drying (tempering) above 100°C, preferably above 130°C. The drying time depends of the applied pressure. As modification II contains less toluene than F-ST, also less energy is needed in the drying (tempering) process.

In another embodiment, the present invention relates to the crystalline modification II having a fipronil content of at least 92 % by weight, particularly at least 96 % by weight and especially at least 98 % by weight.

This invention also relates to solid (compositions of) fipronil comprising the crystalline modification II as defined hereinabove and a form of fipronil being different from said crystalline modification II (herein also referred to as "fipronil form"), e.g. amorphous fipronil or fipronil of a crystalline modification different from crystalline modification II.

- 5 Preferably, the solid (compositions of) fipronil comprise the crystalline modification II in at least 85 % by weight, preferably in at least 90 % by weight, more preferably in at least 95 % by weight, and most preferably in at least 98 % by weight.

- 10 The crystalline modification II can be prepared using a process which comprises the following steps:

- step i) preparing a solution of a solid form of fipronil being different from the crystalline modification II in a solvent S preferably selected from tetrahydrofuran (THF), 1,2-dichloroethane (DCE), acetonitrile, toluene, monochlorobenzene (MCB),
15 1,2-dichlorobenzene (DCB), ethyl benzene, mesitylene, nitrobenzene and CF₃-benzene;
step ii) effecting crystallization of fipronil; and
step iii) isolating the resulting precipitate.

- 20 The preparation of modification II is favored by crystallization from concentrated solutions at elevated temperatures. Further, the resulting modification is dependent of especially the nucleation temperature followed by the formation of the first crystals. The beginning of the crystallization can be detected visually as clouding of the crystallization solution and/or with the aid of a turbidity probe. Usually the evaporation
25 or cooling crystallisation is effected not above 130°C, or close to the boiling point of the used solvent, and preferably not below 80°C.

- Thus crystallization of modification II can be done by crystallization of fipronil from the aforesaid solvents at constant temperature by evaporation and/or by adding at least a
30 polar solvent P to the above mentioned aromatic solvents. Such cooling crystallisation is advantageously started at a temperature close to the boiling point of the solvent, however, preferably not above 130°C or below 80°C. More precisely in the cases of THF, 1,2-dichloroethane and acetonitrile the crystallization temperature in an evaporation crystallization should be kept within a temperature window of 30 and 55°C, for toluene
35 within a temperature window of 90°C to 110°C, for MCB and 1,2-dichlorobenzene in between 70°C and 130°C, nitrobenzene and mesitylene in between 90°C and 130°C and for ethyl benzene in between 100°C and 130°C and for CF₃-benzene in between 90°C and 103°C.

- 40 Modification II can also be crystallised from mixtures of the above mentioned aromatic solvents and polar solvents P, advantageously in amounts up to 20 vol.-%, preferably 5 to 15 vol.-% of P. Such solvent P is preferably selected from the group of methanol,

ethanol, propan-1-ol, propan-2-ol (isopropanol), butan-1-ol (n-butanol), butan-2-ol, tert-butanol, 2-methyl-propan-1-ol (iso-butanol), 2-methyl-propan-2-ol, pentan-3-ol, 2-methyl butan-1-ol, 3-methyl butan-1-ol, 1,2-ethanediol, 1,3-propandiol, 1,2-propandiol, cyclohexanol, acetonitrile, propionitrile, acetone, butanone (methyl ethyl ketone),
5 pentan-2-one (methyl propyl ketone), pentan-3-one (diethylketone), 4-methyl-2-pentanone (isobutyl-methyl-ketone), 3-methyl-butan-2-one (iso-propyl-methyl-ketone), 3,3-dimethyl-2-butanone (tert-butyl-methyl-ketone), cyclohexanone, methylacetate, ethylacetate, isopropylacetate, N-butylacetate, isobutylacetate, diethylcarbonate, 2-butoxyethylacetate, dioxane, THF, diethylether, 2-methyl-THF, methyl-tert-butylether,
10 dimethylformamide, dimethylacetamide, dimethylsulfoxide (DMSO), nitromethane, and nitroethane.

Modification II can also be crystallized by effecting the crystallization by cooling concentrated solutions of the corresponding solvent. In a cooling crystallization the
15 crystallization begin needs to be within the aforesaid temperature ranges. Parameters effecting the nucleation temperature and beginning of crystallization are concentration, cooling rate and time of adding possible seeding crystals of the wanted modification to a supersaturated solution. The concentration depends naturally of the solubility of fipronil in the corresponding solvent. The cooling rate is typically in between 5 K/h and
20 20 K/h. The cooling rate may not exceed 1 K/minute.

A person skilled in the art is able to perform a cooling crystallization in a corresponding solvent so that these conditions are met.

25 If the crystallization is not affected in a way that the nucleation begins in the aforesaid narrow temperature ranges, other modifications or mixtures of modifications result.

Modification II can be obtained from a solution in THF, or DCE by evaporation at from 40°C to the boiling point of the solvent used. Evaporation from a THF solution at 20-
30 25°C yield a mixture of modifications I and II.

Alternatively modification II is obtainable from a solution in acetonitrile by cooling crystallization from 80 to 5°C (cf. example 4).

35 In a further embodiment of the invention modification II is obtainable from a solution in toluene by a cooling crystallization when nucleation and crystallization starts above 90°C (cf. example 5). In between 90 and 75°C mixtures of II and III (F-ST) result.

In a further embodiment of the invention modification II is obtainable from a solution in monochlorobenzene (MCB) also in a cooling crystallization (cf. example 6).
40

In a further embodiment of the invention modification II is obtainable from a solution in mesitylene (cf. example 7).

5 In a further embodiment of the invention modification II is obtainable from a solution in dichlorobenzene (DCB) in two concentrations.

In a further embodiment of the invention modification II is obtainable from a solution in nitrobenzene (cf. example 8).

10 In a further embodiment of the invention modification II is obtainable from a solution in ethyl benzene when crystallization took place above 100°C (cf. example 9). Preferably solutions with a fipronil content of 10 w.-% or more, more preferably of 15 w.-% or more are used.

15 In a further embodiment of the invention modification II is obtainable from a solution in CF₃-benzene at 90°C or more, preferably 95°C or above.

A detailed description of these steps is as follows:

20 Step i)

Suitable fipronil forms different from the crystalline modification II used in step i) are, for example, selected from amorphous fipronil or crystalline fipronil such as other triclinic or monoclinic forms, e.g. monoclinic fipronil of the space group C2/c, and also mixtures
25 of crystalline modifications of fipronil.

The fipronil form used as starting material in step i) preferably has a purity of at least 85 % by weight, in particular at least 90 % by weight and especially at least 95 % by weight. "Purity" means the absence of chemical compounds other than fipronil.

30

The solvent S used in step i) consists of either acetonitrile, DCB, DCE, ethyl benzene, MCB, mesitylene, nitrobenzene, THF, toluene, or CF₃-benzene, each pure or in combination with a polar solvent P.

35 In a preferred embodiment, the solvent S used in step i) consists of DCB, mesitylene, and nitrobenzene, each pure or in combination with a polar solvent P

In another preferred embodiment, the solvent S used in step i) consists of ethyl benzene, MCB, and toluene, each pure or in combination with a polar solvent P

40

In step i), the fipronil form different from the crystalline modification II will usually be incorporated into the solvent S as a solid with mixing at a concentration and temperature where the solvent S is capable of completely dissolving the fipronil form.

- 5 The amount of fipronil form dissolved in the solvent S depends, on the nature of the solvent S and on the dissolution temperature. The person skilled in the art will be able to determine suitable conditions by standard experiments.

Step ii)

10

In step ii) of the process of this invention, fipronil is then crystallized. Crystallization can be effected in a customary manner, for example by cooling the solution obtained in step i), by adding a solvent which reduces the solubility, or by concentrating the solution, or by a combination of the measures mentioned above.

15

In a preferred embodiment, step ii) is carried out in the presence of seed crystals of the crystalline modification II.

- 20 To achieve a conversion into the crystalline modification II which is as complete as possible, the crystallization is carried out over a period (duration of crystallization) of at least 1 h, in particular at least 3 h. Duration of crystallization is understood by the person skilled in the art as meaning the period of time between the beginning of the measure which initiates crystallization and the isolation of the fipronil by separating the crystalline material from the mother liquor.

25

In general, the crystallization is allowed to proceed to a point where at least 60%, preferably at least 70%, in particular at least 90% by weight, for example from 80 to 90% by weight, of the fipronil employed has crystallized out.

- 30 Concentration of the solution is effected by gradually removing the solvent S, such as by evaporation in vacuo, either at low temperature or at about 20°C to 25°C or at elevated temperature, and/or in the presence of a flow of an inert gas such as nitrogen or argon. The values of "low temperature" and "elevated temperature" depend, on the nature of the solvent S, however should not exceed 60°C.

35

In a preferred embodiment, the crystallization is effected by concentrating the solution.

Step iii)

- 40 In step iii) of the process of this invention, the crystalline modification II is isolated using customary techniques for separating solid components from liquids, for example by filtration, centrifugation or decanting. In general, the isolated precipitate will be washed,

- for example with the solvent S used for the crystallization. The washing can be carried out in one or more steps. The washing is typically carried out at temperatures lower than 30°C and in particular lower than 25°C, to keep the loss of the product of value as low as possible. The resulting crystalline fipronil or modification II can then be dried and
- 5 subjected to further processing.

The preparation process consisting of steps i) to step iii) can be repeated in order to achieve higher purities of fipronil.

- 10 The crystalline modification II is especially suitable for efficiently combating the following pests:

millipedes (Diplopoda) such as *Blaniulus* or *Narceus* spp;

- 15 insects (Insecta) such as:

- ants, bees, wasps, sawflies (Hymenoptera), e.g. *Atta capiguara*, *Atta cephalotes*, *Atta laevigata*, *Atta robusta*, *Atta sexdens*, *Atta texana*, *Crematogaster* spp., *Hoplocampa minuta*, *Hoplocampa testudinea*, *Monomorium pharaonis*, *Solenopsis geminata*,
- 20 *Solenopsis invicta*, *Solenopsis richteri*, *Solenopsis xyloni*, *Pheidole megacephala*, *Pogonomyrmex* species such as *Pogonomyrmex barbatus* and *Pogonomyrmex californicus*, *Dasymutilla occidentalis*, *Bombus* spp. *Vespula squamosa*, *Paravespula vulgaris*, *Paravespula pennsylvanica*, *Paravespula germanica*, *Dolichovespula maculata*, *Vespa crabro*, *Polistes rubiginosa*, *Camponotus floridanus*, and *Linepithema*
- 25 *humile*,

- beetles (Coleoptera), such as *Agrius sinuatus*, *Agriotes lineatus*, *Agriotes obscurus* and other *Agriotes* species, *Amphimallus solstitialis*, *Anisandrus dispar*, *Anthonomus grandis*, *Anthonomus pomorum*, *Aracanthus morei*, *Atomaria linearis*, *Blapstinus* species, *Blastophagus piniperda*, *Blitophaga undata*, *Bothynoderes punctiventris*,
- 30 *Bruchus rufimanus*, *Bruchus pisorum*, *Bruchus lentis*, *Byctiscus betulae*, *Cassida nebulosa*, *Cerotoma trifurcata*, *Ceuthorrhynchus assimilis*, *Ceuthorrhynchus napi*, *Chaetocnema tibialis*, *Conoderus vespertinus* and other *Conoderus* species, *Conorhynchus mendicus*, *Crioceris asparagi*, *Cylindrocopturus adspersus*, *Diabrotica*
- 35 *(longicornis) barberi*, *Diabrotica semi-punctata*, *Diabrotica speciosa*, *Diabrotica undecimpunctata*, *Diabrotica virgifera* and other *Diabrotica* species, *Eleodes* species, *Epilachna varivestis*, *Epitrix hirtipennis*, *Eutinobothrus brasiliensis*, *Hylobius abietis*, *Hypera brunneipennis*, *Hypera postica*, *Ips typographus*, *Lema bilineata*, *Lema melanopus*, *Leptinotarsa decemlineata*, *Limonius californicus* and other *Limonius*
- 40 species, *Lissorhoptus oryzophilus*, *Listronotus bonariensis*, *Melanotus communis* and other *Melanotus* species, *Meligethes aeneus*, *Melolontha hippocastani*, *Melolontha melolontha*, *Oulema oryzae*, *Ortiorrhynchus sulcatus*, *Oryzophagus oryzae*,

- Otiorrhynchus ovatus, Oulema oryzae, Phaedon cochleariae, Phyllotreta chrysocephala, Phyllophaga cuyabana and other Phyllophaga species, Phyllopertha horticola, Phyllotreta nemorum, Phyllotreta striolata, and other Phyllotreta species, Popillia japonica, Promecops carinicolis, Premnotypes voraz, Psylliodes species,
- 5 Sitona lineatus, Sitophilus granaria, Sternechus pinguis, Sternechus subsignatus, and Tanymechus palliatus and other Tanymechus species,
- Centipedes (Chilopoda), e.g. Scutigera coleoptrata,
- 10 Cockroaches (Blattaria - Blattodea), e.g. Blattella germanica, Blattella asahinae, Periplaneta americana, Periplaneta japonica, Periplaneta brunnea, Periplaneta fuliginosa, Periplaneta australasiae, and Blatta orientalis,
- Crickets, grasshoppers, locusts (Orthoptera), e.g. Acheta domestica, Gryllotalpa
- 15 gryllotalpa, Locusta migratoria, Melanoplus bivittatus, Melanoplus femurrubrum, Melanoplus mexicanus, Melanoplus sanguinipes, Melanoplus spretus, Nomadacris septemfasciata, Schistocerca americana, Schistocerca gregaria, Dociostaurus maroccanus, Tachycines asynamorus, Oedaleus senegalensis, Zonozelus variegatus, Hieroglyphus daganensis, Kraussaria angulifera, Calliptamus italicus, Chortoicetes
- 20 terminifera, and Locustana pardalina,
- fleas (Siphonaptera), e.g. Ctenocephalides felis, Ctenocephalides canis, Xenopsylla cheopis, Pulex irritans, Tunga penetrans, and Nosopsyllus fasciatus,
- 25 Flies, mosquitoes (Diptera), e.g. Aedes aegypti, Aedes albopictus, Aedes vexans, Agromyza oryzae, Anastrepha ludens, Anopheles maculipennis, Anopheles crucians, Anopheles albimanus, Anopheles gambiae, Anopheles freeborni, Anopheles leucosphyrus, Anopheles minimus, Anopheles quadrimaculatus, Calliphora vicina, Chrysomya bezziana, Chrysomya hominivorax, Chrysomya macellaria, Chrysops
- 30 discalis, Chrysops silacea, Chrysops atlanticus, Cochliomyia hominivorax, Contarinia sorghicola, Cordylobia anthropophaga, Culicoides furens, Culex pipiens, Culex nigripalpus, Culex quinquefasciatus, Culex tarsalis, Culiseta inornata, Culiseta melanura, Dacus cucurbitae, Dacus oleae, Dasineura brassicae, Delia antiqua, Delia coarctata, Delia platura, Delia radicum, Dermatobia hominis, Fannia canicularis,
- 35 Gasterophilus intestinalis, Geomyza Tripunctata, Glossina morsitans, Glossina palpalis, Glossina fuscipes, Glossina tachinoides, Haematobia irritans, Haplodiplosis equestris, Hippelates spp., Hypoderma lineata, Leptoconops torrens, Liriomyza sativae, Liriomyza trifolii, Lucilia caprina, Lucilia cuprina, Lucilia sericata, Lycoria pectoralis, Mansonia spp., Mayetiola destructor, Musca domestica, Muscina stabulans,
- 40 Oestrus ovis, Oestrus ovis, Opomyza florum, Oscinella frit, Pegomya hysocyami, Phlebotomus argentipes, Phorbia antiqua, Phorbia brassicae, Phorbia coarctata, Progonyx leyoscanii, Psila rosae, Psorophora columbiae, Psorophora discolor,

Prosimulium mixtum, Rhagoletis cerasi, Rhagoletis pomonella, Sarcophaga haemorrhoidalis, Sarcophaga sp., Simulium vittatum, Stomoxys calcitrans, Tabanus bovinus, Tabanus atratus, Tabanus lineola, Tabanus similis, Tetanops myopaeformis, Tipula olerace, and Tipula paludosa,

5

Heteropterans (Heteroptera), such as Acrosternum hilare, Blissus leucopterus, Cicadellidae such as Empoasca fabae, Chrysomelidae, Cyrtopeltis notatus, Delpahcidae, Dysdercus cingulatus, Dysdercus intermedius, Eurygaster integriceps, Euschistus impictiventris, Leptoglossus phyllopus, Lygus lineolaris, Lygus pratensis, 10 Nephrotettix species, Nezara viridula, Pentatomidae, Piesma quadrata, Solubea insularis and Thyanta perditor,

Aphids and other homopterans (Homoptera), e.g. Acyrthosiphon onobrychis, Adelges laricis, Aphidula nasturtii, Aphis fabae, Aphis forbesi, Aphis glycines, Aphis gossypii, 15 Aphis grossulariae, Aphis pomi, Aphis schneideri, Aphis spiraecola, Aphis sambuci, Acyrthosiphon pisum, Aulacorthum solani, Brachycaudus cardui, Brachycaudus helichrysi, Brachycaudus persicae, Brachycaudus prunicola, Brevicoryne brassicae, Capitophorus horni, Cerosipha gossypii, Chaetosiphon fragaefolii, Cryptomyzus ribis, Dreyfusia nordmannianae, Dreyfusia piceae, Dysaphis radicola, Dysaulacorthum 20 pseudosolani, Dysaphis plantaginea, Dysaphis pyri, Empoasca fabae, Hyalopterus pruni, Hyperomyzus lactucae, Macrosiphum avenae, Macrosiphum euphorbiae, Macrosiphon rosae, Megoura viciae, Melanaphis pyrarius, Metopolophium dirhodum, Myzodes (Myzus) persicae, Myzus ascalonicus, Myzus cerasi, Myzus varians, Nasonovia ribis-nigri, Nilaparvata lugens, Pemphigus bursarius, Pemphigus 25 populivenae, and other Pemphigus species, Perkinsiella saccharicida, Phorodon humuli, Psyllidae such as Psylla mali, Psylla piri and other Psylla species, Rhopalomyzus ascalonicus, Rhopalosiphum maidis, Rhopalosiphum padi, Rhopalosiphum insertum, Sappaphis mala, Sappaphis mali, Schizaphis graminum, Schizoneura lanuginosa, Sitobion avenae, Trialeurodes vaporariorum, Toxoptera 30 aurantiand, and Viteus vitifolii,

Lepidopterans (Lepidoptera), for example Agrotis ypsilon, Agrotis segetum and other Agrotis species, Alabama argillacea, Anticarsia gemmatilis, Argyresthia conjugella, Autographa gamma, Bupalus piniarius, Cacoecia murinana, Capua reticulana, 35 Cheimatobia brumata, Chilo suppressalis and other Chilo species, Choristoneura fumiferana, Choristoneura occidentalis, Cirphis unipuncta, Cnaphlocrocis medinalis, Cydia pomonella, Dendrolimus pini, Diaphania nitidalis, Diatraea grandiosella, Earias insulana, Elasmopalpus lignosellus, Eupoecilia ambiguella, Euxoa species, Evetria bouliana, Feltia subterranea, Galleria mellonella, Grapholitha funebrana, Grapholitha molesta, Heliothis armigera, Heliothis virescens, Heliothis zea, Hellula undalis, Hibernia defoliaria, Hyphantria cunea, Hyponomeuta malinellus, Keiferia lycopersicella, 40 Lambdina fiscellaria, Laphygma exigua, Lerodea eufala, Leucoptera coffeella,

- Leucoptera scitella, Lithocolletis blancardella, Lobesia botrana, Loxostege sticticalis, Lymantria dispar, Lymantria monacha, Lyonetia clerkella, Malacosoma neustria, Mamestra brassicae, Momphidae, Orgyia pseudotsugata, Ostrinia nubilalis, Panolis flammea, Pectinophora gossypiella, Peridroma saucia, Phalera bucephala,
- 5 Phthorimaea operculella, Phyllocnistis citrella, Pieris brassicae, Plathypena scabra, Plutella xylostella, Pseudoplusia includens, Rhyacionia frustrana, Scrobipalpula absoluta, Sesamia nonagrioides and other Sesamia species, Sitotroga cerealella, Sparganothis pilleriana, Spodoptera frugiperda, Spodoptera littoralis, Spodoptera litura, Thaumtopoea pityocampa, Tortrix viridana, Trichoplusia ni and Zeiraphera
- 10 canadensis,
- lice (Phthiraptera), e.g. Pediculus humanus capitis, Pediculus humanus corporis, Pthirus pubis, Haematopinus eurysternus, Haematopinus suis, Linognathus vituli, Bovicola bovis, Menopon gallinae, Menacanthus stramineus and Solenopotes
- 15 capillatus,
- orthopterans (Orthoptera), such as Acrididae, Acheta domestica, Forficula auricularia, Gryllotalpa gryllotalpa, Locusta migratoria, Melanoplus bivittatus, Melanoplus femur-rubrum, Melanoplus mexicanus, Melanoplus sanguinipes, Melanoplus spretus,
- 20 Nomadacris septemfasciata, Schistocerca americana, Schistocerca peregrina, Stauronotus maroccanus and Tachycines asynamorus,
- silverfish, firebrat (Thysanura), e.g. Lepisma saccharina and Thermobia domestica,
- 25 termites (Isoptera), such as Calotermes flavicollis, Coptotermes ssp., Dalbulus maidis, Heterotermes aureus, Leucotermes flavipes, Macrotermes gilvus, Reticulitermes ssp., Termes natalensis, Coptotermes formosanus,
- thrips (Thysanoptera), such as Frankliniella fusca, Frankliniella occidentalis,
- 30 Frankliniella tritici and other Frankliniella species, Scirtothrips citri, Thrips oryzae, Thrips palmi, Thrips simplex, and Thrips tabaci,
- ticks and parasitic mites (Parasitiformes): ticks (Ixodida), e.g. Ixodes scapularis, Ixodes holocyclus, Ixodes pacificus, Rhipicephalus sanguineus, Dermacentor andersoni,
- 35 Dermacentor variabilis, Amblyomma americanum, Amblyomma maculatum, Ornithodoros hermsi, Ornithodoros turicata and parasitic mites (Mesostigmata), e.g. Ornithonyssus bacoti and Dermanyssus gallinae,
- true bugs (Hemiptera), e.g. Cimex lectularius, Cimex hemipterus, Reduvius senilis,
- 40 Triatoma spp., Rhodnius prolixus, and Arilus critatus,

Arachnoidea, such as arachnids (Acarina), for example of the families Argasidae, Ixodidae and Sarcoptidae, such as *Amblyomma americanum*, *Amblyomma variegatum*, *Argas persicus*, *Boophilus annulatus*, *Boophilus decoloratus*, *Boophilus microplus*, *Dermacentor silvarum*, *Hyalomma truncatum*, *Ixodes ricinus*, *Ixodes rubicundus*,
 5 *Latrodectus mactans*, *Loxosceles reclusa*, *Ornithodoros moubata*, *Otobius megnini*, *Dermanyssus gallinae*, *Psoroptes ovis*, *Rhipicephalus appendiculatus*, *Rhipicephalus evertsi*, *Sarcoptes scabiei*, and Eriophyidae species such as *Aculus schlechtendali*, *Phyllocoptura oleivora* and Eriophyes sheldoni; Tarsonemidae species such as *Phytonemus pallidus* and *Polyphagotarsonemus latus*; Tenuipalpidae species such as
 10 *Brevipalpus phoenicis*; Tetranychidae species such as *Tetranychus cinnabarinus*, *Tetranychus kanzawai*, *Tetranychus pacificus*, *Tetranychus telarius* and *Tetranychus urticae*, *Panonychus ulmi*, *Panonychus citri*, and *Oligonychus pratensis*,

15 Earwigs (Dermaptera), e.g. *forficula auricularia*; and

Nematodes, including plant parasitic nematodes and nematodes living in the soil. Plant parasitic nematodes include, such as root knot nematodes, *Meloidogyne hapla*, *Meloidogyne incognita*, *Meloidogyne javanica*, and other *Meloidogyne* species; cyst-forming nematodes, *Globodera rostochiensis* and other *Globodera* species; *Heterodera*
 20 *avenae*, *Heterodera glycines*, *Heterodera schachtii*, *Heterodera trifolii*, and other *Heterodera* species; Seed gall nematodes, *Anguina* species; Stem and foliar nematodes, *Aphelenchoides* species; Sting nematodes, *Belonolaimus longicaudatus* and other *Belonolaimus* species; Pine nematodes, *Bursaphelenchus xylophilus* and other *Bursaphelenchus* species; Ring nematodes, *Criconema* species, *Criconemella*
 25 species, *Criconemoides* species, *Mesocriconema* species; Stem and bulb nematodes, *Ditylenchus destructor*, *Ditylenchus dipsaci* and other *Ditylenchus* species; Awl nematodes, *Dolichodorus* species; Spiral nematodes, *Helicotylenchus multicinctus* and other *Helicotylenchus* species; Sheath and sheathoid nematodes, *Hemicyclophora* species and *Hemicriconemoides* species; *Hirshmanniella* species; Lance nematodes,
 30 *Hoploaimus* species; false rootknot nematodes, *Nacobbus* species; Needle nematodes, *Longidorus elongatus* and other *Longidorus* species; Pin nematodes, *Paratylenchus* species; Lesion nematodes, *Pratylenchus neglectus*, *Pratylenchus penetrans*, *Pratylenchus curvatus*, *Pratylenchus goodeyi* and other *Pratylenchus* species; Burrowing nematodes, *Radopholus similis* and other *Radopholus* species;
 35 Reniform nematodes, *Rotylenchus robustus* and other *Rotylenchus* species; *Scutellonema* species; Stubby root nematodes, *Trichodorus primitivus* and other *Trichodorus* species, *Paratrachodorus* species; Stunt nematodes, *Tylenchorhynchus claytoni*, *Tylenchorhynchus dubius* and other *Tylenchorhynchus* species; Citrus nematodes, *Tylenchulus* species, Dagger nematodes, *Xiphinema* species and other
 40 plant parasitic nematode species.

Moreover, the crystalline modification II is especially useful for the control of crop pests, in particular of the Coleoptera, Lepidoptera and Acarina orders.

Moreover, the crystalline modification II is especially useful for the control of non-crop
5 pests (household, turf, ornamental). Non-crop pests are pests of the classes Chilopoda and Diplopoda and of the orders Isoptera, Diptera, Blattaria (Blattodea), Dermaptera, Hemiptera, Hymenoptera, Orthoptera, Siphonaptera, Thysanura, Phthiraptera, and Acarina.

10 For use according to the present invention, the crystalline modification II can be converted into the customary formulations, for example solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular intended purpose; in each case, it should ensure a fine and even distribution of the compound according to the invention.

15

The formulations are prepared in a known manner (see e.g. for review US 3,060,084, EP-A 707 445 (for liquid concentrates), Browning, "Agglomeration", Chemical Engineering, Dec. 4, 1967, 147-48, Perry's Chemical Engineer's Handbook, 4th Ed., McGraw-Hill, New York, 1963, pages 8-57 and et seq. WO 91/13546, US 4,172,714,
20 US 4,144,050, US 3,920,442, US 5,180,587, US 5,232,701, US 5,208,030, GB 2,095,558, US 3,299,566, Klingman, Weed Control as a Science, John Wiley and Sons, Inc., New York, 1961, Hance et al., Weed Control Handbook, 8th Ed., Blackwell Scientific Publications, Oxford, 1989 and Mollet, H., Grubemann, A., Formulation technology, Wiley VCH Verlag GmbH, Weinheim (Germany), 2001, 2. D. A. Knowles,
25 Chemistry and Technology of Agrochemical Formulations, Kluwer Academic Publishers, Dordrecht, 1998 (ISBN 0-7514-0443-8), for example by extending the active compound with auxiliaries suitable for the formulation of agrochemicals, such as solvents and/or carriers, if desired surfactants (e.g. adjuvants, emulsifiers, dispersing agents), preservatives, antifoaming agents, anti-freezing agents, for seed treatment
30 formulations also optionally colorants and/or binders and/or gelling agents.

Examples of suitable solvents are water, aromatic solvents (for example Solvesso products, xylene), paraffins (for example mineral oil fractions), alcohols (for example methanol, butanol, pentanol, benzyl alcohol), ketones (for example cyclohexanone,
35 gamma-butyrolactone), pyrrolidones (NMP, NOP), acetates (glycol diacetate), glycols, fatty acid dimethylamides, fatty acids and fatty acid esters. In principle, solvent mixtures may also be used.

Examples of suitable carriers are ground natural minerals (for example kaolins, clays,
40 talc, chalk) and ground synthetic minerals (for example highly disperse silica, silicates).

Suitable surfactants used are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid, phenolsulfonic acid, dibutyl naphthalenesulfonic acid, alkylarylsulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates, fatty acids and sulfated fatty alcohol glycol ethers, furthermore

5 condensates of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensates of naphthalene or of naphthalenesulfonic acid with phenol and formaldehyde, polyoxyethylene octylphenol ether, ethoxylated isooctylphenol, octylphenol, nonylphenol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ether, tristearylphenyl polyglycol ether, alkylaryl polyether alcohols, alcohol and fatty alcohol

10 ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignosulfite waste liquors and methylcellulose.

Substances which are suitable for the preparation of directly sprayable solutions,

15 emulsions, pastes or oil dispersions are mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, for example toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, methanol, ethanol, propanol, butanol, cyclohexanol, cyclohexanone, isophorone, highly polar

20 solvents, for example dimethyl sulfoxide, N-methylpyrrolidone or water.

Also anti-freezing agents such as glycerin, ethylene glycol, propylene glycol and bactericides can be added to the formulation.

25 Suitable antifoaming agents are for example antifoaming agents based on silicon or magnesium stearate.

Suitable preservatives are for example Dichlorophen und enzyalkoholhemiformal.

30 Seed Treatment formulations may additionally comprise binders and optionally colorants.

Binders can be added to improve the adhesion of the active materials on the seeds after treatment. Suitable binders are block copolymers EO/PO surfactants but also

35 polyvinylalcohols, polyvinylpyrrolidones, polyacrylates, polymethacrylates, polybutenes, polyisobutylenes, polystyrene, polyethyleneamines, polyethyleneamides, polyethyleneimines (Lupasol®, Polymin®), polyethers, polyurethans, polyvinylacetate, tylose and copolymers derived from these polymers.

40 Optionally, also colorants can be included in the formulation. Suitable colorants or dyes for seed treatment formulations are Rhodamin B, C.I. Pigment Red 112, C.I. Solvent

Red 1, pigment blue 15:4, pigment blue 15:3, pigment blue 15:2, pigment blue 15:1, pigment blue 80, pigment yellow 1, pigment yellow 13, pigment red 112, pigment red 48:2, pigment red 48:1, pigment red 57:1, pigment red 53:1, pigment orange 43, pigment orange 34, pigment orange 5, pigment green 36, pigment green 7, pigment
5 white 6, pigment brown 25, basic violet 10, basic violet 49, acid red 51, acid red 52, acid red 14, acid blue 9, acid yellow 23, basic red 10, basic red 108.

Examples of a gelling agent is carrageen (Satiage[®]).

10 Powders, materials for spreading and dustable products can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

Granules, for example coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active compounds to solid carriers.

15

Examples of solid carriers are mineral earths such as silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers, such as, for example, ammonium sulfate, ammonium phosphate, ammonium
20 nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.

In general, the formulations comprise from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active compound(s). In this case, the active compound(s) are
25 employed in a purity of from 90% to 100% by weight, preferably 95% to 100% by weight (according to NMR spectrum).

For seed treatment purposes, the respective formulations can be diluted 2-10 fold leading to concentrations in the ready to use preparations of 0.01 to 60% by weight
30 active compound by weight, preferably 0.1 to 40% by weight.

The crystalline modification II can be used as such, in the form of their formulations or the use forms prepared therefrom, for example in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes,
35 dustable products, materials for spreading, or granules, by means of spraying, atomizing, dusting, spreading or pouring. The use forms depend entirely on the intended purposes; they are intended to ensure in each case the finest possible distribution of the active compound(s) according to the invention.

40 Aqueous use forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances, as such or dissolved in an oil or solvent, can

be homogenized in water by means of a wetter, tackifier, dispersant or emulsifier. However, it is also possible to prepare concentrates composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

5

The active compound concentrations in the ready-to-use preparations can be varied within relatively wide ranges. In general, they are from 0.0001 to 10%, preferably from 0.01 to 1% per weight.

- 10 The active compound(s) may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply formulations comprising over 95% by weight of active compound, or even to apply the active compound without additives.

- 15 The following are examples of formulations: 1. Products for dilution with water for foliar applications. For seed treatment purposes, such products may be applied to the seed diluted or undiluted.

A) Water-soluble concentrates (SL, LS)

- 20 10 parts by weight of the active compound(s) are dissolved in 90 parts by weight of water or a water-soluble solvent. As an alternative, wetters or other auxiliaries are added. The active compound(s) dissolves upon dilution with water, whereby a formulation with 10 % (w/w) of active compound(s) is obtained.

B) Dispersible concentrates (DC)

- 25 20 parts by weight of the active compound(s) are dissolved in 70 parts by weight of cyclohexanone with addition of 10 parts by weight of a dispersant, for example polyvinylpyrrolidone. Dilution with water gives a dispersion, whereby a formulation with 20% (w/w) of active compound(s) is obtained.

- 30 C) Emulsifiable concentrates (EC)

15 parts by weight of the active compound(s) are dissolved in 80 parts by weight of xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5 parts by weight). Dilution with water gives an emulsion, whereby a formulation with 15% (w/w) of active compound(s) is obtained.

35

D) Emulsions (EW, EO, ES)

- 40 25 parts by weight of the active compound(s) are dissolved in 35 parts by weight of xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5 parts by weight). This mixture is introduced into 30 parts by weight of water by means of an emulsifier machine (e.g. Ultraturrax) and made into a homogeneous emulsion. Dilution with water gives an emulsion, whereby a formulation with 25% (w/w) of active compound(s) is obtained.

E) Suspensions (SC, OD, FS)

In an agitated ball mill, 20 parts by weight of the active compound(s) are comminuted with addition of 10 parts by weight of dispersants, wetters and 70 parts by weight of water or of an organic solvent to give a fine active compound(s) suspension. Dilution with water gives a stable suspension of the active compound(s), whereby a formulation with 20% (w/w) of active compound(s) is obtained.

F) Water-dispersible granules and water-soluble granules (WG, SG)

50 parts by weight of the active compound(s) are ground finely with addition of 50 parts by weight of dispersants and wetters and made as water-dispersible or water-soluble granules by means of technical appliances (for example extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active compound(s), whereby a formulation with 50% (w/w) of active compound(s) is obtained.

G) Water-dispersible powders and water-soluble powders (WP, SP, SS, WS)

75 parts by weight of the active compound(s) are ground in a rotor-stator mill with addition of 25 parts by weight of dispersants, wetters and silica gel. Dilution with water gives a stable dispersion or solution of the active compound(s), whereby a formulation with 75% (w/w) of active compound(s) is obtained.

H) Gel-Formulation (GF) (for seed treatment purposes only)

In an agitated ball mill, 20 parts by weight of the active compound(s) are comminuted with addition of 10 parts by weight of dispersants, 1 part by weight of a gelling agent/ wetters and 70 parts by weight of water or of an organic solvent to give a fine active compound(s) suspension. Dilution with water gives a stable suspension of the active compound(s), whereby a formulation with 20% (w/w) of active compound(s) is obtained.

2. Products to be applied undiluted for foliar applications. For seed treatment purposes, such products may be applied to the seed diluted.

I) Dustable powders (DP, DS)

5 parts by weight of the active compound(s) are ground finely and mixed intimately with 95 parts by weight of finely divided kaolin. This gives a dustable product having 5% (w/w) of active compound(s)

J) Granules (GR, FG, GG, MG)

0.5 part by weight of the active compound(s) is ground finely and associated with 95.5 parts by weight of carriers, whereby a formulation with 0.5% (w/w) of active

compound(s) is obtained. Current methods are extrusion, spray-drying or the fluidized bed. This gives granules to be applied undiluted for foliar use.

K) ULV solutions (UL)

- 5 10 parts by weight of the active compound(s) are dissolved in 90 parts by weight of an organic solvent, for example xylene. This gives a product having 10% (w/w) of active compound(s), which is applied undiluted for foliar use.

- 10 Conventional seed treatment formulations include for example flowable concentrates FS, solutions LS, powders for dry treatment DS, water dispersible powders for slurry treatment WS, water-soluble powders SS and emulsion ES and EC and gel formulation GF. These formulation can be applied to the seed diluted or undiluted. Application to the seeds is carried out before sowing, either directly on the seeds.

- 15 In a preferred embodiment a FS formulation is used for seed treatment. Typically, a FS formulation may comprise 1-800 g/l of active ingredient, 1-200 g/l surfactant, 0 to 200 g/l antifreezing agent, 0 to 400 g/l of binder, 0 to 200 g/l of a pigment and up to 1 liter of a solvent, preferably water.

- 20 The invention relates in particular to pesticidal or parasitocidal compositions in the form of an aqueous suspension concentrate (SC). Such suspension concentrates comprise the crystalline modification II in a finely divided particulate form, where the particles of the crystalline modification II are suspended in an aqueous medium. The size of the active compound particles, i.e. the size which is not exceeded by 90% by weight of the
25 active compound particles, is typically below 30 μm , in particular below 20 μm . Advantageously, at least 40% by weight and in particular at least 60% by weight of the particles in the SCs according to the invention have diameters below 2 μm .

- In addition to the active compound, suspension concentrates typically comprise
30 surfactants, and also, if appropriate, antifoam agents, thickeners, antifreeze agents, stabilizers (biocides), agents for adjusting the pH and anticaking agents.

- In such SCs, the amount of active compound, i.e. the total amount of the crystalline modification II and, if appropriate, further active compounds is usually in the range from
35 10 to 70% by weight, in particular in the range from 20 to 50% by weight, based on the total weight of the suspension concentrate.

- Preferred surfactants are anionic and nonionic surfactants. The amount of surfactants will generally be from 0.5 to 20% by weight, in particular from 1 to 15% by weight and
40 particularly preferably from 1 to 10% by weight, based on the total weight of the SCs according to the invention. Preferably, the surfactants comprise at least one anionic

surfactant and at least one nonionic surfactant, the ratio of anionic to nonionic surfactant typically being in the range from 10:1 to 1:10.

5 Examples of anionic surfactants include alkylaryl sulfonates, phenyl sulfonates, alkyl sulfates, alkyl sulfonates, alkyl ether sulfates, alkylaryl ether sulfates, alkyl polyglycol ether phosphates, polyaryl phenyl ether phosphates, alkyl sulfosuccinates, olefin sulfonates, paraffin sulfonates, petroleum sulfonates, taurides, sarcosides, fatty acids, alkyl naphthalenesulfonic acids, naphthalenesulfonic acids, lignosulfonic acids, condensates of sulfonated naphthalenes with formaldehyde or with formaldehyde and
10 phenol and, if appropriate, urea, and also condensates of phenolsulfonic acid, formaldehyde and urea, lignosulfite waste liquors and lignosulfonates, alkyl phosphates, alkylaryl phosphates, for example tristyryl phosphates, and also polycarboxylates, such as, for example, polyacrylates, maleic anhydride/olefin copolymers (for example Sokalan[®] CP9, BASF), including the alkali metal, alkaline
15 earth metal, ammonium and amine salts of the substances mentioned above. Preferred anionic surfactants are those which carry at least one sulfonate group, and in particular their alkali metal and their ammonium salts.

Examples of nonionic surfactants comprise alkylphenol alkoxyates, alcohol
20 alkoxyates, fatty amine alkoxyates, polyoxyethylene glycerol fatty acid esters, castor oil alkoxyates, fatty acid alkoxyates, fatty amide alkoxyates, fatty polydiethanolamides, lanolin ethoxyates, fatty acid polyglycol esters, isotridecyl alcohol, fatty amides, methylcellulose, fatty acid esters, alkyl polyglycosides, glycerol fatty acid esters, polyethylene glycol, polypropylene glycol, polyethylene
25 glycol/polypropylene glycol block copolymers, polyethylene glycol alkyl ethers, polypropylene glycol alkyl ethers, polyethylene glycol/polypropylene glycol ether block copolymers (polyethylene oxide/polypropylene oxide block copolymers) and mixtures thereof. Preferred nonionic surfactants are fatty alcohol ethoxyates, alkyl polyglycosides, glycerol fatty acid esters, castor oil alkoxyates, fatty acid alkoxyates,
30 fatty amide alkoxyates, lanolin ethoxyates, fatty acid polyglycol esters and ethylene oxide/propylene oxide block copolymers and mixtures thereof.

In particular, the SCs according to the invention comprise at least one surfactant which improves wetting of the plant parts by the aqueous application form (wetting agent) and
35 at least one surfactant which stabilizes the dispersion of the active compound particles in the SC (dispersant). The amount of wetting agent is typically in the range from 0.5 to 10% by weight, in particular from 0.5 to 5% by weight and especially from 0.5 to 3% by weight, based on the total weight of the SC. The amount of dispersant is typically from 0.5 to 10% by weight and in particular from 0.5 to 5% by weight, based on the total
40 weight of the SC.

Preferred wetting agents are of anionic or nonionic nature and selected, for example, from naphthalenesulfonic acids including their alkali metal, alkaline earth metal, ammonium and amine salts, furthermore fatty alcohol ethoxylates, alkyl polyglycosides, glycerol fatty acid esters, castor oil alkoxylates, fatty acid alkoxylates, fatty amide
5 alkoxylates, fatty polydiethanolamides, lanolin ethoxylates and fatty acid polyglycol esters.

Preferred dispersants are of anionic or nonionic nature and selected, for example, from polyethylene glycol/polypropylene glycol block copolymers, polyethylene glycol alkyl
10 ethers, polypropylene glycol alkyl ethers, polyethylene glycol/polypropylene glycol ether block copolymers, alkylaryl phosphates, for example tristyryl phosphates, lignosulfonic acids, condensates of sulfonated naphthalenes with formaldehyde or with formaldehyde and phenol and, if appropriate, urea, and also condensates of phenolsulfonic acid, formaldehyde and urea, lignosulfite waste liquors and
15 lignosulfonates, polycarboxylates, such as, for example, polyacrylates, maleic anhydride/olefin copolymers (for example Sokalan[®] CP9, BASF), including the alkali metal, alkaline earth metal, ammonium and amine salts of the substances mentioned above.

20 Viscosity-modifying additives (thickeners) suitable for the SCs according to the invention are in particular compounds which bestow upon the formulation pseudoplastic flow properties, i.e. high viscosity in the resting state and low viscosity in the agitated state. Suitable are, in principle, all compounds used for this purpose in suspension concentrates. Mention may be made, for example, of inorganic substances,
25 such as bentonites or attapulgites (for example Attaclay[®] from Engelhardt), and organic substances, such as polysaccharides and heteropolysaccharides, such as xanthan gum such as sold under the trademarks Kelzan[®] from Kelco, Rhodopol[®] 23 from Rhone Poulenc or Veegum[®] from R.T. Vanderbilt, and preference is given to using xanthan gum. Frequently, the amount of viscosity-modifying additives is from 0.1 to 5%
30 by weight, based on the total weight of the SC.

Antifoam agents suitable for the SCs according to the invention are, for example, silicone emulsions known for this purpose (Silikon[®] SRE, from Wacker, or Rhodorsil[®] from Rhodia), long-chain alcohols, fatty acids, defoamers of the type of aqueous wax
35 dispersions, solid defoamers (so-called Compounds), organofluorine compounds and mixtures thereof. The amount of antifoam agent is typically from 0.1 to 1% by weight, based on the total weight of the SC.

Bactericides may be added for stabilizing the suspension concentrates according to the
40 invention. Suitable bactericides are those based on isothiazolones, for example Proxel[®] from ICI or Acticide[®] RS from Thor Chemie or Kathon[®] MK from Rohm & Haas. The amount of bactericides is typically from 0.05 to 0.5% by weight, based on the total

weight of the SC.

Suitable antifreeze agents are liquid polyols, for example ethylene glycol, propylene glycol or glycerol. The amount of antifreeze agents is generally from 1 to 20% by weight, in particular from 5 to 10% by weight, based on the total weight of the suspension concentrate.

If appropriate, the SCs according to the invention may comprise buffers for regulating the pH. Examples of buffers are alkali metal salts of weak inorganic or organic acids, such as, for example, phosphoric acid, boric acid, acetic acid, propionic acid, citric acid, fumaric acid, tartaric acid, oxalic acid and succinic acid.

The invention relates in particular to pesticidal or parasitocidal compositions in the form of water-dispersible granules (WG) or a water dispersible powder (WP). Such formulations comprise the crystalline modification II in a finely divided particulate form, where the particles of the crystalline modification II are homogenized in a solid or powder form. The size of the active compound particles, i.e. the size which is not exceeded by 90% by weight of the active compound particles, is typically below 30 μm , in particular below 20 μm . Advantageously, at least 40% by weight and in particular at least 60% by weight of the particles in the WGs or WPs according to the invention have diameters below 5 μm .

In addition to the active compound, water-dispersible powders and water dispersible granules typically comprise surfactants, and also, if appropriate, antifoam agents, fillers, binders, and anticaking agents.

In such WGs and WPs, the amount of active compound, i.e. the total amount of the crystalline modification II and, if appropriate, further active compounds is usually in the range from 10 to 90% by weight, in particular in the range from 20 to 75% by weight, based on the total weight of the WG/WP.

Preferred surfactants are anionic and nonionic surfactants. The amount of surfactants will generally be from 0.5 to 20% by weight, in particular from 1 to 15% by weight and particularly preferably from 1 to 10% by weight, based on the total weight of the WGs or WPs according to the invention. Preferably, the surfactants comprise at least one anionic surfactant and at least one nonionic surfactant, the ratio of anionic to nonionic surfactant typically being in the range from 10:1 to 1:10.

Examples of anionic surfactants include alkylaryl sulfonates, phenyl sulfonates, alkyl sulfates, alkyl sulfonates, alkyl ether sulfates, alkylaryl ether sulfates, alkyl polyglycol ether phosphates, polyaryl phenyl ether phosphates, alkyl sulfosuccinates, olefin sulfonates, paraffin sulfonates, petroleum sulfonates, taurides, sarcosides, fatty acids,

alkylnaphthalenesulfonic acids, naphthalenesulfonic acids, lignosulfonic acids, condensates of sulfonated naphthalenes with formaldehyde or with formaldehyde and phenol and, if appropriate, urea, and also condensates of phenolsulfonic acid, formaldehyde and urea, lignosulfite waste liquors and lignosulfonates, alkyl
5 phosphates, alkylaryl phosphates, for example tristyryl phosphates, and also polycarboxylates, such as, for example, polyacrylates, maleic anhydride/olefin copolymers (for example Sokalan[®] CP9, BASF), including the alkali metal, alkaline earth metal, ammonium and amine salts of the substances mentioned above. Preferred anionic surfactants are those which carry at least one sulfonate group, and in particular
10 their alkali metal and their ammonium salts.

Examples of nonionic surfactants comprise alkylphenol alkoxylates, alcohol alkoxylates, fatty amine alkoxylates, polyoxyethylene glycerol fatty acid esters, castor oil alkoxylates, fatty acid alkoxylates, fatty amide alkoxylates, fatty
15 polydiethanolamides, lanolin ethoxylates, fatty acid polyglycol esters, isotridecyl alcohol, fatty amides, methylcellulose, fatty acid esters, alkyl polyglycosides, glycerol fatty acid esters, polyethylene glycol, polypropylene glycol, polyethylene glycol/polypropylene glycol block copolymers, polyethylene glycol alkyl ethers, polypropylene glycol alkyl ethers, polyethylene glycol/polypropylene glycol ether block
20 copolymers (polyethylene oxide/polypropylene oxide block copolymers) and mixtures thereof. Preferred nonionic surfactants are fatty alcohol ethoxylates, alkyl polyglycosides, glycerol fatty acid esters, castor oil alkoxylates, fatty acid alkoxylates, fatty amide alkoxylates, lanolin ethoxylates, fatty acid polyglycol esters and ethylene oxide/ propylene oxide block copolymers and mixtures thereof.

25 In particular, the WGs or WPs according to the invention comprise at least one surfactant which improves wetting of the formulation by the aqueous application form (wetting agent) and at least one surfactant which allows dispersion of the active compound particles in aqueous dilutions. The amount of wetting agent is typically in the
30 range from 0.5 to 10% by weight, in particular from 0.5 to 5% by weight and especially from 0.5 to 3% by weight, based on the total weight of the WG/WP. The amount of dispersant is typically from 0.5 to 10% by weight and in particular from 2.0 to 8% by weight, based on the total weight of the WG/WP.

35 Preferred wetting agents are of anionic or nonionic nature and selected, for example, from naphthalenesulfonic acids including their alkali metal, alkaline earth metal, ammonium and amine salts, furthermore fatty alcohol ethoxylates, alkyl polyglycosides, glycerol fatty acid esters, castor oil alkoxylates, fatty acid alkoxylates, fatty amide alkoxylates, fatty polydiethanolamides, lanolin ethoxylates and fatty acid polyglycol
40 esters.

Preferred dispersants are of anionic or nonionic nature and selected, for example, from polyethylene glycol/polypropylene glycol block copolymers, polyethylene glycol alkyl ethers, polypropylene glycol alkyl ethers, polyethylene glycol/polypropylene glycol ether block copolymers, alkylaryl phosphates, for example tristyryl phosphates, sodium
5 phosphates, sodium lauryl sulphate, modified cellulose gum, polyvinylpyrrolidinone, lignosulfonic acids, condensates of sulfonated naphthalenes with formaldehyde or with formaldehyde and phenol and, if appropriate, urea, and also condensates of phenol-sulfonic acid, formaldehyde and urea, lignosulfite waste liquors and lignosulfonates, polycarboxylates, such as, for example, polyacrylates, maleic anhydride/olefin
10 copolymers (for example Sokalan[®] CP9, BASF), including the alkali metal, alkaline earth metal, ammonium and amine salts of the substances mentioned above.

Antifoam agents suitable for the WGs or WPs according to the invention are, for example, tallow soap known for this purpose (Agnique Soap L, Foamaster Soap L),
15 long-chain alcohols, fatty acids, organofluorine compounds and mixtures thereof. The amount of antifoam agent is typically from 0.1 to 1% by weight, based on the total weight of the WG/WP.

Fillers, binders, or additional dispersing aids suitable for the WGs and WPs according to the invention typically make up the remainder of the formulation. These typically are for example kaolin or attapulgite clay, fumed or precipitated silica, diatomaceous earth, ammonium sulphate, or calcium silicate.

The crystalline modification II is effective through both contact and ingestion.
25

According to a preferred embodiment of the invention, the crystalline modification II is employed via soil application. Soil application is especially favorable for use against ants, termites, crickets, or cockroaches.

30 According to another preferred embodiment of the invention, for use against non-crop pests such as ants, termites, wasps, flies, mosquitoes, crickets, locusts, or cockroaches the crystalline modification II is prepared into a bait preparation.

The bait can be a liquid, a solid or a semisolid preparation (e.g. a gel). Solid baits can be formed into various shapes and forms suitable to the respective application e.g.
35 granules, blocks, sticks, disks. Liquid baits can be filled into various devices to ensure proper application, e.g. open containers, spray devices, droplet sources, or evaporation sources. Gels can be based on aqueous or oily matrices and can be formulated to particular necessities in terms of stickiness, moisture retention or aging characteristics.
40

The bait employed in the composition is a product which is sufficiently attractive to incite insects such as ants, termites, wasps, flies, mosquitoes, crickets etc. or cockroaches to eat it. This attractant may be chosen from feeding stimulants or para and/or sex pheromones. Suitable feeding stimulants are chosen, for example, from
5 animal and/or plant proteins (meat-, fish- or blood meal, insect parts, crickets powder, egg yolk), from fats and oils of animal and/or plant origin, or mono-, oligo- or polyorganosaccharides, especially from sucrose, lactose, fructose, dextrose, glucose, starch, pectin or even molasses or honey, or from salts such as ammonium sulfate, ammonium carbonate or ammonium acetate. Fresh or decaying parts of fruits, crops,
10 plants, animals, insects or specific parts thereof can also serve as a feeding stimulant. Pheromones are known to be more insect specific. Specific pheromones are described in the literature and are known to those skilled in the art.

Compositions of this invention may also contain other active ingredients, for example
15 other pesticides, insecticides, fungicides, herbicides, fertilizers such as ammonium nitrate, urea, potash, and superphosphate, phytotoxicants and plant growth regulators, safeners and nematicides. These additional ingredients may be used sequentially or in combination with the above-described compositions, if appropriate also added only immediately prior to use (tank mix). For example, the plant(s) may be sprayed with a
20 composition of this invention either before or after being treated with other active ingredients.

The following list of pesticidal or parasitocidal compounds which can be used together with the crystalline modification II according to the invention is intended to illustrate the
25 possible combinations, but not to impose any limitation:

A.1. Organo(thio)phosphates: acephate, azamethiphos, azinphos-methyl, chlorpyrifos, chlorpyrifos-methyl, chlorfenvinphos, diazinon, dichlorvos, dicrotophos, dimethoate, disulfoton, ethion, fenitrothion, fenthion, isoxathion, malathion, methamidophos,
30 methidathion, methyl-parathion, mevinphos, monocrotophos, oxydemeton-methyl, paraoxon, parathion, phenthoate, phosalone, phosmet, phosphamidon, phorate, phoxim, pirimiphos-methyl, profenofos, prothiofos, sulprophos, tetrachlorvinphos, terbufos, triazophos, trichlorfon;


35 A.2. Carbamates: alanycarb, aldicarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, fenoxycarb, furathiocarb, methiocarb, methomyl, oxamyl, pirimicarb, propoxur, thiodicarb, triazamate;

A.3. Pyrethroids: allethrin, bifenthrin, cyfluthrin, cyhalothrin, cyphenothrin,
40 cypermethrin, alpha-cypermethrin, beta-cypermethrin, zeta-cypermethrin, deltamethrin, esfenvalerate, etofenprox, fenpropathrin, fenvalerate, imiprothrin, lambda-cyhalothrin,


permethrin, prallethrin, pyrethrin I and II, resmethrin, silafluofen, tau-fluvalinate, tefluthrin, tetramethrin, tralomethrin, transfluthrin, profluthrin, dimefluthrin;

- A.4. Growth regulators: a) chitin synthesis inhibitors: benzoylureas: chlorfluazuron, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron; buprofezin, diofenolan, hexythiazox, etoxazole, clofentazine; b) ecdysone antagonists: halofenozide, methoxyfenozide, tebufenozide, azadirachtin; c) juvenoids: pyriproxyfen, methoprene, fenoxycarb; d) lipid biosynthesis inhibitors: spirotetramat, spiromesifen, spirotetramat;

A.5. Nicotinic receptor agonists/antagonists compounds: clothianidin, dinotefuran, imidacloprid, thiamethoxam, nitenpyram, acetamiprid, thiacloprid; the thiazol compound of formula Γ^1



- A.6. GABA antagonist compounds: acetoprole, endosulfan, ethiprole, fipronil, vanilprole, pyrafluprole, pyriprole, 5-Amino-1-(2,6-dichloro-4-trifluoromethyl-phenyl)-4-trifluoromethanesulfinyl-1H-pyrazole-3-carbothioic acid amide of formula Γ^2



- A.7. Macrocyclic lactone insecticides: abamectin, emamectin, milbemectin, lepimectin, spinosad;

A.8. METI I compounds: fenazaquin, pyridaben, tebufenpyrad, tolfeenpyrad, flufenerim;

A.9. METI II and III compounds: acequinocyl, fluacyprim, hydramethylnon;

A.10. Uncoupler compounds: chlorfenapyr;

A.11. Oxidative phosphorylation inhibitor compounds: cyhexatin, diafenthiuron, fenbutatin oxide, propargite;

A.12. Moulting disruptor compounds: cyromazine;


A.13. Mixed Function Oxidase inhibitor compounds: piperonyl butoxide;

A.14. Sodium channel blocker compounds: indoxacarb, metaflumizone,

A.15. Various: benclothiaz, bifenazate, cartap, flonicamid, pyridalyl, pymetrozine, sulfur, thiocyclam, flubendiamide, cyenopyrafen, flupyrazofos, cyflumetofen,

5 amidoflumet,

the anthranilamide compounds of formula Γ^3



wherein A¹ is CH₃, Cl, Br, I, X is C-H, C-Cl, C-F or N, Y' is F, Cl, or Br, Y'' is hydrogen, F, Cl, CF₃, B¹ is hydrogen, Cl, Br, I, CN, B² is Cl, Br, CF₃, OCH₂CF₃, OCF₂H, and R^B is hydrogen, CH₃ or CH(CH₃)₂, and the malononitrile compounds as described in JP 2002 284608, WO 02/89579, WO 02/90320, WO 02/90321, WO 04/06677, WO 04/20399, JP 2004 99597, WO 05/68423, WO 05/68432, or WO 05/63694, especially the malononitrile compounds CF₂HCF₂CF₂CF₂CH₂C(CN)₂CH₂CH₂CF₃ (2-(2,2,3,3,4,4,5,5-octafluoropentyl)-2-(3,3,3-trifluoropropyl)malononitrile),

10 CF₃(CH₂)₂C(CN)₂CH₂(CF₂)₅CF₂H (2-(2,2,3,3,4,4,5,5,6,6,7,7-Dodecafluoro-heptyl)-2-(3,3,3-trifluoro-propyl)-malononitrile), CF₃(CH₂)₂C(CN)₂(CH₂)₂C(CF₃)₂F (2-(3,4,4,4-Tetrafluoro-3-trifluoromethyl-butyl)-2-(3,3,3-trifluoro-propyl)-malononitrile),

15 CF₃(CH₂)₂C(CN)₂(CH₂)₂(CF₂)₃CF₃ (2-(3,3,4,4,5,5,6,6,6-Nonafluoro-hexyl)-2-(3,3,3-trifluoro-propyl)-malononitrile), CF₂H(CF₂)₃CH₂C(CN)₂CH₂(CF₂)₃CF₂H (2,2-Bis-(2,2,3,3,4,4,5,5-octafluoro-pentyl)-malononitrile), CF₃(CH₂)₂C(CN)₂CH₂(CF₂)₃CF₃ (2-(2,2,3,3,4,4,5,5,5-Nonafluoro-pentyl)-2-(3,3,3-trifluoro-propyl)-malononitrile),

20 CF₃(CF₂)₂CH₂C(CN)₂CH₂(CF₂)₃CF₂H (2-(2,2,3,3,4,4,4-Heptafluoro-butyl)-2-(2,2,3,3,4,4,5,5-octafluoro-pentyl)-malononitrile) and CF₃CF₂CH₂C(CN)₂CH₂(CF₂)₃CF₂H (2-(2,2,3,3,4,4,5,5-Octafluoro-pentyl)-2-(2,2,3,3,3-pentafluoro-propyl)-malononitrile).

25

The commercially available compounds of the group A may be found in The Pesticide Manual, 13th Edition, British Crop Protection Council (2003) among other publications. Thioamides of formula Γ^2 and their preparation have been described in WO 98/28279. Lepimectin is known from Agro Project, PJB Publications Ltd, November 2004.

30 Benclothiaz and its preparation have been described in EP-A1 454621. Methidathion and Paraoxon and their preparation have been described in Farm Chemicals Handbook, Volume 88, Meister Publishing Company, 2001. Acetoprole and its preparation have been described in WO 98/28277. Metaflumizone and its preparation have been described in EP-A1 462 456. Flupyrazofos has been described in Pesticide Science 54, 1988, p.237-243 and in US 4822779. Pyrafluprole and its preparation have been described in JP 2002193709 and in WO 01/00614. Pyriprole and its preparation

35

have been described in WO 98/45274 and in US 6335357. Amidoflumet and its preparation have been described in US 6221890 and in JP 21010907. Flufenimerim and its preparation have been described in WO 03/007717 and in WO 03/007718. Cyflumetofen and its preparation have been described in WO 04/080180.

- 5 Anthranilamides of formula I³ and their preparation have been described in WO 01/70671; WO 02/48137; WO 03/24222, WO 03/15518, WO 04/67528; WO 04/33468; and WO 05/118552. The malononitrile compounds CF₂HCF₂CF₂CF₂CH₂C(CN)₂CH₂CH₂CF₃ (2-(2,2,3,3,4,4,5,5-octafluoropentyl)-2-(3,3,3-trifluoropropyl)-malononitrile), CF₃(CH₂)₂C(CN)₂CH₂(CF₂)₅CF₂H (2-(2,2,3,3,4,4,5,5,6,6,7,7-Dodecafluoro-heptyl)-2-(3,3,3-trifluoro-propyl)-malononitrile),
- 10 CF₃(CH₂)₂C(CN)₂(CH₂)₂C(CF₃)₂F (2-(3,4,4,4-Tetrafluoro-3-trifluoromethyl-butyl)-2-(3,3,3-trifluoro-propyl)-malononitrile), CF₃(CH₂)₂C(CN)₂(CH₂)₂(CF₂)₃CF₃ (2-(3,3,4,4,5,5,6,6-Nonafluoro-hexyl)-2-(3,3,3-trifluoro-propyl)-malononitrile), CF₂H(CF₂)₃CH₂C(CN)₂CH₂(CF₂)₃CF₂H (2,2-Bis-(2,2,3,3,4,4,5,5-octafluoro-pentyl)-malononitrile), CF₃(CH₂)₂C(CN)₂CH₂(CF₂)₃CF₃ (2-(2,2,3,3,4,4,5,5,5-Nonafluoro-pentyl)-2-(3,3,3-trifluoro-propyl)-malononitrile), CF₃(CF₂)₂CH₂C(CN)₂CH₂(CF₂)₃CF₂H (2-(2,2,3,3,4,4,4-Heptafluoro-butyl)-2-(2,2,3,3,4,4,5,5-octafluoro-pentyl)-malononitrile) and CF₃CF₂CH₂C(CN)₂CH₂(CF₂)₃CF₂H (2-(2,2,3,3,4,4,5,5-Octafluoro-pentyl)-2-(2,2,3,3,3-pentafluoro-propyl)-malononitrile) have been described in WO 05/63694.

20

The following list of fungicidal compounds which can be used together with the crystalline modification II according to the invention is intended to illustrate the possible combinations, but not to impose any limitation:

- 25 Preferred are the binary mixtures containing modification II as compound I.

Preferred are the tertiary mixtures containing modification II as compound I, a compound IIA, and a compound IIB.

- 30 Preferred are the quaternary mixtures containing modification II as compound I, a compound IIA, and two compounds IIB1 and IIB2, resp.

- Especially preferred are binary mixtures containing modification II as compound I and a fungicidal compound IIA selected from the list comprising azoles: cyproconazole, di-
- 35 fenconazole, epoxiconazole, fenbuconazole, fluquinconazole, flutriafol, hexaconazole, ipconazole, metconazole, propiconazole, prothioconazole, tebuconazole, tetraconazole, triadimenol, triadimefon, triticonazole, cyazofamid, imazalil, prochloraz, triflumizol, benomyl, carbendazim, thiabendazole, ethaboxam, and hymexazole.

- 40 Especially preferred are binary mixtures containing modification II as compound I and a fungicidal compound IIA selected from the list comprising strobilurins: azoxystrobin, dimoxystrobin, enestroburin, fluoxastrobin, kresoxim-methyl, metominostrobin, picoxy-

strobil, pyraclostrobin, trifloxystrobin, methyl (2-chloro-5-[1-(3-methylbenzyloxy-imino)ethyl]benzyl)carbamate, methyl (2-chloro-5-[1-(6-methylpyridin-2-ylmethoxy-imino)ethyl]benzyl)carbamate, and methyl 2-(ortho-((2,5-dimethylphenoxy-methylene)phenyl)-3-methoxyacrylate;

5

Especially preferred are binary mixtures containing modification II as compound I and a fungicidal compound IIA selected from the list comprising carboxamides: boscalid,

carboxin, benalaxyl, fenhexamid, flutolanil, furametpyr, metalaxyl, mefenoxam (metalaxyl-M), ofurace, oxadixyl, oxycarboxin, penthiopyrad, thifluzamide, tiadinil, dimetho-

10 morph, fluopicolide (picobenzamid), diclocymet, N-(4'-bromobiphenyl-2-yl)-4-difluoromethyl-2-methylthiazole-5-carboxamide, N-(4'-trifluoromethylbiphenyl-2-yl)-4-di-

fluoromethyl-2-methylthiazole-5-carboxamide, N-(4'-chloro-3'-fluorobiphenyl-2-yl)-4-di-

fluoromethyl-2-methylthiazole-5-carboxamide, N-(3',4'-dichloro-4-fluorobiphenyl-2-yl)-3-

difluoromethyl-1-methylpyrazole-4-carboxamide, N-(3',4'-dichloro-5-fluorobiphenyl-2-

15 yl)-3-difluoromethyl-1-methylpyrazole-4-carboxamide; 3,4-dichloro-N-(2-cyanophenyl)isothiazol-5-carboxamide; N-(2',4'-difluorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-

1H-pyrazole-4-carboxamide; N-(2',4'-dichlorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-

1H-pyrazole-4-carboxamide; N-(2',4'-difluorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-

1H-pyrazole-4-carboxamide; N-(2',4'-dichloro-biphenyl-2-yl)-3-difluoromethyl-1-methyl-

20 1H-pyrazole-4-carboxamide; N-(2',5'-difluorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-

1H-pyrazole-4-carboxamide; N-(2',5'-dichlorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-

1H-pyrazole-4-carboxamide; N-(2',5'-difluorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-

1H-pyrazole-4-carboxamide; N-(2',5'-dichlorobiphenyl-2-yl)-3-di-fluoromethyl-1-methyl-

1H-pyrazole-4-carboxamide; N-(3',5'-difluoro-biphenyl-2-yl)-1-methyl-3-trifluoromethyl-

25 1H-pyrazole-4-carboxamide; N-(3',5'-dichlorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-

1H-pyrazole-4-carboxamide; N-(3',5'-difluorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-

1H-pyrazole-4-carboxamide; N-(3',5'-dichlorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-

1H-pyrazole-4-carboxamide; N-(3'-fluorobiphenyl-2-yl)-1-methyl-3-trifluoro-methyl-1H-

pyrazole-4-carboxamide, N-(3'-chlorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-1H-

30 pyrazole-4-carboxamide; N-(3'-fluorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-

pyrazole-4-carboxamide, N-(3'-chlorobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-

pyrazole-4-carboxamide; N-(2'-fluorobiphenyl-2-yl)-1-methyl-3-trifluoromethyl-1H-

pyrazole-4-carboxamide; N-(2'-chlorobiphenyl-2-yl)-1-methyl-3-trifluoro-methyl-1H-

pyrazole-4-carboxamide; N-(2'-fluorobiphenyl-2-yl)-3-difluoro-methyl-1-methyl-1H-

35 pyrazole-4-carboxamide; N-(2'-chlorbiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-

pyrazole-4-carboxamide; N-(2'-fluoro-4'-chloro-5'-methylbiphenyl-2-yl)-1-methyl-3-

trifluoromethyl-1H-pyrazole-4-carbox-amide; N-(3',4',5'-trifluorobiphenyl-2-yl)-1-methyl-

3-trifluoromethyl-1H-pyrazole-4-carboxamide; N-(3',4',5'-trifluorobiphenyl-2-yl)-1-

methyl-3-difluoromethyl-1H-pyrazole-4-carboxamide; N-(2',4',5'-trifluorobiphenyl-2-yl)-

40 1-methyl-3-difluoromethyl-1H-pyrazole-4-carboxamide; N-(3',4',5'-tri-fluorobiphenyl-2-

yl)-3-chlorofluoromethyl-1-methyl-1H-pyrazole-4-carbox-amide; N-[2-(1,1,2,3,3,3-hexa-

fluoropropoxy)phenyl]-1-methyl-3-trifluoro-methyl-1H-pyrazole-4-carboxamide; N-[2-

(1,1,2,3,3,3-hexafluoropropoxy)-phenyl]-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-[2-(2-chloro-1,1,2-trifluoroethoxy)phenyl]-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboxamide; N-[2-(2-chloro-1,1,2-trifluoroethoxy)phenyl]-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide; N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide; N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboxamide; N-(4'-(trifluoromethylthio)biphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide; N-(4'-(trifluoromethylthio)biphenyl-2-yl)-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboxamide; and 5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxylic acid [2-(1,2-dimethyl-propyl)-phenyl]-amide.

Especially preferred are binary mixtures containing modification II as compound I and a fungicidal compound IIA selected from the list comprising heterocyclic compounds: pyrimethanil, fenpiclonil, fludioxonil, aldimorph, dodemorph, fenpropimorph, tridemorph, iprodione, procymidone, famoxadone, fenamidone, octhilinone, probenazole, diclomezine, pyroquilon, proquinazid, tricyclazole, captan, captan, dazomet, fenoxanil, quin-oxyfen, 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine, 6-(3,4-dichloro-phenyl)-5-methyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-yl-amine, 6-(4-tert-butylphenyl)-5-methyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 5-methyl-6-(3,5,5-trimethyl-hexyl)-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 5-methyl-6-octyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 6-methyl-5-octyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 6-ethyl-5-octyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 5-ethyl-6-octyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 5-ethyl-6-(3,5,5-trimethyl-hexyl)-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 6-octyl-5-propyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 5-methoxymethyl-6-octyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, 6-octyl-5-trifluoromethyl-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine, and 5-trifluoromethyl-6-(3,5,5-trimethyl-hexyl)-[1,2,4]triazolo[1,5-a]pyrimidine-7-ylamine.

Especially preferred are binary mixtures containing modification II as compound I and a fungicidal compound IIA selected from the list comprising carbamates: mancozeb, maneb, metam, metiram, ferbam, propineb, thiram, zineb, ziram; diethofencarb, iprovalicarb, propamocarb, and methyl 3-(4-chlorophenyl)-3-(2-isopropoxycarbonylamino-3-methylbutylamino)propanoate.

Especially preferred are binary mixtures containing modification II as compound I and a fungicidal compound IIA selected from the list comprising: guazatine; streptomycin, validamycin A; binapacryl, dinocap, dinobuton; dithianon, isoprothiolane; fentin salts, such as fentin-acetate; edifenphos, iprobenfos, fosetyl, pyrazophos, chlorothalonil, dichlofluanid, flusulfamide, phthalide, quintozone, thiophanate-methyl, tolylfluanid; copper acetate, copper hydroxide, copper oxychloride, basic copper sulfate, sulfur; cyflufenamid, cymoxanil, dimethirimol, ethirimol, furalaxyl, metrafenone, and spiroxamine.

The active compounds IIA mentioned above, their preparation and their action against harmful fungi are generally known (cf.: <http://www.hclrss.demon.co.uk/index.html>); they are commercially available. The compounds named according to IUPAC, their
5 preparation and their fungicidal activity are likewise known from EP-A 12 01 648; EP-A 226 917; WO 98/46608; WO 99/24413; WO 2004/049804; WO 2003/066609; WO 2003/053145; WO 2003/14103; EP-A 10 35 122; EP-A 10 28 125; EP-A 71 792; EP-A 141 317; WO 2003/009687; WO 05/087771; WO 2005/087772; WO 2005/087773; WO 2006/087325; WO 2006/087325; WO 2006/092428; WO 2006/092428; WO
10 2006/087343; WO 2001/42223; WO 2005/34628; WO 2005/123689; WO 2005/123690; WO 2006/120219; PCT/EP2006/064991; WO 2007/017450, and EP Application No. 06123463.9

With respect to their intended use, the following tertiary and quaternary mixtures of
15 modification II as compound I are especially preferred:

Table1

Mixtures wherein compound IIA is trifloxystrobin, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

20

Table2

Mixtures wherein compound IIA is azoxystrobin, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

25 Table3

Mixtures wherein compound IIA is pyraclostrobin, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table4

30 Mixtures wherein compound IIA is boscalid, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table5

35 Mixtures wherein compound IIA is metalaxyl, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table6

Mixtures wherein compound IIA is metalaxyl-M, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

40

Table7

Mixtures wherein compound IIA is cyproconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

5 Table8

Mixtures wherein compound IIA is epoxiconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table9

- 10 Mixtures wherein compound IIA is fenbuconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table10

- 15 Mixtures wherein compound IIA is fluquinconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table11

- 20 Mixtures wherein compound IIA is flutriafol, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table12

Mixtures wherein compound IIA is ipconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

25 Table13

Mixtures wherein compound IIA is metconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table14

- 30 Mixtures wherein compound IIA is propiconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table15

- 35 Mixtures wherein compound IIA is prothioconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table16

Mixtures wherein compound IIA is tebuconazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table17

Mixtures wherein compound IIA is triadimenol, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

5 Table18

Mixtures wherein compound IIA is triticonazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table19

- 10 Mixtures wherein compound IIA is imazalil, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table20

- 15 Mixtures wherein compound IIA is prochloraz, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table21

- 20 Mixtures wherein compound IIA is carbendazim, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table22

Mixtures wherein compound IIA is thiabendazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

25 Table23

Mixtures wherein compound IIA is ethaboxam, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table24

- 30 Mixtures wherein compound IIA is hymexazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table25

- 35 Mixtures wherein compound IIA is pyrimethanil, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table26

Mixtures wherein compound IIA is fludioxonil, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table27

Mixtures wherein compound IIA is aldimorph, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

5 Table28

Mixtures wherein compound IIA is dodemorph, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table29

- 10 Mixtures wherein compound IIA is fenpropimorph, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table30

- 15 Mixtures wherein compound IIA is iprodione, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table31

- 20 Mixtures wherein compound IIA is captan, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table32

Mixtures wherein compound IIA is fenoxanil, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

25 Table33

Mixtures wherein compound IIA is probenazole, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table34

- 30 Mixtures wherein compound IIA is mancozeb, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table35

- 35 Mixtures wherein compound IIA is metiram, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table36

- 40 Mixtures wherein compound IIA is thiram, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table37

Mixtures wherein compound IIA is ziram, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

5 Table38

Mixtures wherein compound IIA is guazatin, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table39

- 10 Mixtures wherein compound IIA is thiophanate-methyl, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table40

- 15 Mixtures wherein compound IIA is chlorothalonil, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

Table41

Mixtures wherein compound IIA is metrafenone, and the combination of compounds IIB1 and IIB2 in each case corresponds to a row of Table Q.

20

Table Q

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-1	azoxystrobin	-	M-19	azoxystrobin	thiabendazole
M-2	azoxystrobin	boscalid	M-20	azoxystrobin	ethaboxam
M-3	azoxystrobin	metalaxyl	M-21	azoxystrobin	hymexazole
M-4	azoxystrobin	cyproconazole	M-22	azoxystrobin	pyrimethanil
M-5	azoxystrobin	epoxiconazole	M-23	azoxystrobin	fludioxonil
M-6	azoxystrobin	fenbuconazole	M-24	azoxystrobin	aldimorph
M-7	azoxystrobin	fluquinconazole	M-25	azoxystrobin	dodemorph
M-8	azoxystrobin	flutriafol	M-26	azoxystrobin	fenpropimorph
M-9	azoxystrobin	ipconazole	M-27	azoxystrobin	iprodione
M-10	azoxystrobin	metconazole	M-28	azoxystrobin	captan
M-11	azoxystrobin	propiconazole	M-29	azoxystrobin	fenoxanil
M-12	azoxystrobin	prothioconazole	M-30	azoxystrobin	probenazol
M-13	azoxystrobin	tebuconazole	M-31	azoxystrobin	mancozeb
M-14	azoxystrobin	triadimenol	M-32	azoxystrobin	metiram
M-15	azoxystrobin	triticonazole	M-33	azoxystrobin	thiram
M-16	azoxystrobin	imazalil	M-34	azoxystrobin	ziram
M-17	azoxystrobin	prochloraz	M-35	azoxystrobin	guazatin
M-18	azoxystrobin	carbendazim	M-36	azoxystrobin	thiophanate-

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
		methyl			
M-37	azoxystrobin	chlorothalonil	M-74	trifloxystrobin	thiophanate-methyl
M-38	azoxystrobin	metrafenone	M-75	trifloxystrobin	chlorothalonil
M-39	trifloxystrobin	-	M-76	trifloxystrobin	metrafenone
M-40	trifloxystrobin	boscalid	M-77	orysastrobin	-
M-41	trifloxystrobin	metalaxyl	M-78	orysastrobin	boscalid
M-42	trifloxystrobin	cyproconazole	M-79	orysastrobin	metalaxyl
M-43	trifloxystrobin	epoxiconazole	M-80	orysastrobin	cyproconazole
M-44	trifloxystrobin	fenbuconazole	M-81	orysastrobin	epoxiconazole
M-45	trifloxystrobin	fluquinconazole	M-82	orysastrobin	fenbuconazole
M-46	trifloxystrobin	flutriafol	M-83	orysastrobin	fluquinconazole
M-47	trifloxystrobin	ipconazole	M-84	orysastrobin	flutriafol
M-48	trifloxystrobin	metconazole	M-85	orysastrobin	ipconazole
M-49	trifloxystrobin	propiconazole	M-86	orysastrobin	metconazole
M-50	trifloxystrobin	prothioconazole	M-87	orysastrobin	propiconazole
M-51	trifloxystrobin	tebuconazole	M-88	orysastrobin	prothioconazole
M-52	trifloxystrobin	triadimenol	M-89	orysastrobin	tebuconazole
M-53	trifloxystrobin	triticonazole	M-90	orysastrobin	triadimenol
M-54	trifloxystrobin	imazalil	M-91	orysastrobin	triticonazole
M-55	trifloxystrobin	prochloraz	M-92	orysastrobin	imazalil
M-56	trifloxystrobin	carbendazim	M-93	orysastrobin	prochloraz
M-57	trifloxystrobin	thiabendazole	M-94	orysastrobin	carbendazim
M-58	trifloxystrobin	ethaboxam	M-95	orysastrobin	thiabendazole
M-59	trifloxystrobin	hymexazole	M-96	orysastrobin	ethaboxam
M-60	trifloxystrobin	pyrimethanil	M-97	orysastrobin	hymexazole
M-61	trifloxystrobin	fludioxonil	M-98	orysastrobin	pyrimethanil
M-62	trifloxystrobin	aldimorph	M-99	orysastrobin	fludioxonil
M-63	trifloxystrobin	dodemorph	M-100	orysastrobin	aldimorph
M-64	trifloxystrobin	fenpropimorph	M-101	orysastrobin	dodemorph
M-65	trifloxystrobin	iprodione	M-102	orysastrobin	fenpropimorph
M-66	trifloxystrobin	captan	M-103	orysastrobin	iprodione
M-67	trifloxystrobin	fenoxanil	M-104	orysastrobin	captan
M-68	trifloxystrobin	probenazol	M-105	orysastrobin	fenoxanil
M-69	trifloxystrobin	mancozeb	M-106	orysastrobin	probenazol
M-70	trifloxystrobin	metiram	M-107	orysastrobin	mancozeb
M-71	trifloxystrobin	thiram	M-108	orysastrobin	metiram
M-72	trifloxystrobin	ziram	M-109	orysastrobin	thiram
M-73	trifloxystrobin	guazatin	M-110	orysastrobin	ziram

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-111	orysastrobin	guazatin	M-148	pyraclostrobin	ziram
M-112	orysastrobin	thiophanate-methyl	M-149	pyraclostrobin	guazatin
M-113	orysastrobin	chlorothalonil	M-150	pyraclostrobin	thiophanate-methyl
M-114	orysastrobin	metrafenone	M-151	pyraclostrobin	chlorothalonil
M-115	pyraclostrobin	-	M-152	pyraclostrobin	metrafenone
M-116	pyraclostrobin	boscalid	M-153	boscalid	-
M-117	pyraclostrobin	metalaxyl	M-154	boscalid	metalaxyl
M-118	pyraclostrobin	cyproconazole	M-155	boscalid	cyproconazole
M-119	pyraclostrobin	epoxiconazole	M-156	boscalid	epoxiconazole
M-120	pyraclostrobin	fenbuconazole	M-157	boscalid	fenbuconazole
M-121	pyraclostrobin	fluquinconazole	M-158	boscalid	fluquinconazole
M-122	pyraclostrobin	flutriafol	M-159	boscalid	flutriafol
M-123	pyraclostrobin	ipconazole	M-160	boscalid	ipconazole
M-124	pyraclostrobin	metconazole	M-161	boscalid	metconazole
M-125	pyraclostrobin	propiconazole	M-162	boscalid	propiconazole
M-126	pyraclostrobin	prothioconazole	M-163	boscalid	prothioconazole
M-127	pyraclostrobin	tebuconazole	M-164	boscalid	tebuconazole
M-128	pyraclostrobin	triadimenol	M-165	boscalid	triadimenol
M-129	pyraclostrobin	triticonazole	M-166	boscalid	triticonazole
M-130	pyraclostrobin	imazalil	M-167	boscalid	imazalil
M-131	pyraclostrobin	prochloraz	M-168	boscalid	prochloraz
M-132	pyraclostrobin	carbendazim	M-169	boscalid	carbendazim
M-133	pyraclostrobin	thiabendazole	M-170	boscalid	thiabendazole
M-134	pyraclostrobin	ethaboxam	M-171	boscalid	ethaboxam
M-135	pyraclostrobin	hymexazole	M-172	boscalid	hymexazole
M-136	pyraclostrobin	pyrimethanil	M-173	boscalid	pyrimethanil
M-137	pyraclostrobin	fludioxonil	M-174	boscalid	fludioxonil
M-138	pyraclostrobin	aldimorph	M-175	boscalid	aldimorph
M-139	pyraclostrobin	dodemorph	M-176	boscalid	dodemorph
M-140	pyraclostrobin	fenpropimorph	M-177	boscalid	fenpropimorph
M-141	pyraclostrobin	iprodione	M-178	boscalid	iprodione
M-142	pyraclostrobin	captan	M-179	boscalid	captan
M-143	pyraclostrobin	fenoxanil	M-180	boscalid	fenoxanil
M-144	pyraclostrobin	probenazol	M-181	boscalid	probenazol
M-145	pyraclostrobin	mancozeb	M-182	boscalid	mancozeb
M-146	pyraclostrobin	metiram	M-183	boscalid	metiram
M-147	pyraclostrobin	thiram	M-184	boscalid	thiram

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-185	boscalid	ziram	M-222	metalaxyl	guazatin
M-186	boscalid	guazatin	M-223	metalaxyl	thiophanate-methyl
M-187	boscalid	thiophanate-methyl	M-224	metalaxyl	chlorothalonil
M-188	boscalid	chlorothalonil	M-225	metalaxyl	metrafenone
M-189	boscalid	metrafenone	M-226	cyproconazole	-
M-190	metalaxyl	-	M-227	cyproconazole	epoxiconazole
M-191	metalaxyl	cyproconazole	M-228	cyproconazole	fenbuconazole
M-192	metalaxyl	epoxiconazole	M-229	cyproconazole	fluquinconazole
M-193	metalaxyl	fenbuconazole	M-230	cyproconazole	flutriafol
M-194	metalaxyl	fluquinconazole	M-231	cyproconazole	ipconazole
M-195	metalaxyl	flutriafol	M-232	cyproconazole	metconazole
M-196	metalaxyl	ipconazole	M-233	cyproconazole	propiconazole
M-197	metalaxyl	metconazole	M-234	cyproconazole	prothioconazole
M-198	metalaxyl	propiconazole	M-235	cyproconazole	tebuconazole
M-199	metalaxyl	prothioconazole	M-236	cyproconazole	triadimenol
M-200	metalaxyl	tebuconazole	M-237	cyproconazole	triticonazole
M-201	metalaxyl	triadimenol	M-238	cyproconazole	imazalil
M-202	metalaxyl	triticonazole	M-239	cyproconazole	prochloraz
M-203	metalaxyl	imazalil	M-240	cyproconazole	carbendazim
M-204	metalaxyl	prochloraz	M-241	cyproconazole	thiabendazole
M-205	metalaxyl	carbendazim	M-242	cyproconazole	ethaboxam
M-206	metalaxyl	thiabendazole	M-243	cyproconazole	hymexazole
M-207	metalaxyl	ethaboxam	M-244	cyproconazole	pyrimethanil
M-208	metalaxyl	hymexazole	M-245	cyproconazole	fludioxonil
M-209	metalaxyl	pyrimethanil	M-246	cyproconazole	aldimorph
M-210	metalaxyl	fludioxonil	M-247	cyproconazole	dodemorph
M-211	metalaxyl	aldimorph	M-248	cyproconazole	fenpropimorph
M-212	metalaxyl	dodemorph	M-249	cyproconazole	iprodione
M-213	metalaxyl	fenpropimorph	M-250	cyproconazole	captan
M-214	metalaxyl	iprodione	M-251	cyproconazole	fenoxanil
M-215	metalaxyl	captan	M-252	cyproconazole	probenazol
M-216	metalaxyl	fenoxanil	M-253	cyproconazole	mancozeb
M-217	metalaxyl	probenazol	M-254	cyproconazole	metiram
M-218	metalaxyl	mancozeb	M-255	cyproconazole	thiram
M-219	metalaxyl	metiram	M-256	cyproconazole	ziram
M-220	metalaxyl	thiram	M-257	cyproconazole	guazatin
M-221	metalaxyl	ziram	M-258	cyproconazole	thiophanate-

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
		methyl	M-295	fenbuconazole	-
M-259	cyproconazole	chlorothalonil	M-296	fenbuconazole	fluquinconazole
M-260	cyproconazole	metrafenone	M-297	fenbuconazole	flutriafol
M-261	epoxiconazole	-	M-298	fenbuconazole	ipconazole
M-262	epoxiconazole	fenbuconazole	M-299	fenbuconazole	metconazole
M-263	epoxiconazole	fluquinconazole	M-300	fenbuconazole	propiconazole
M-264	epoxiconazole	flutriafol	M-301	fenbuconazole	prothioconazole
M-265	epoxiconazole	ipconazole	M-302	fenbuconazole	tebuconazole
M-266	epoxiconazole	metconazole	M-303	fenbuconazole	triadimenol
M-267	epoxiconazole	propiconazole	M-304	fenbuconazole	triticonazole
M-268	epoxiconazole	prothioconazole	M-305	fenbuconazole	imazalil
M-269	epoxiconazole	tebuconazole	M-306	fenbuconazole	prochloraz
M-270	epoxiconazole	triadimenol	M-307	fenbuconazole	carbendazim
M-271	epoxiconazole	triticonazole	M-308	fenbuconazole	thiabendazole
M-272	epoxiconazole	imazalil	M-309	fenbuconazole	ethaboxam
M-273	epoxiconazole	prochloraz	M-310	fenbuconazole	hymexazole
M-274	epoxiconazole	carbendazim	M-311	fenbuconazole	pyrimethanil
M-275	epoxiconazole	thiabendazole	M-312	fenbuconazole	fludioxonil
M-276	epoxiconazole	ethaboxam	M-313	fenbuconazole	aldimorph
M-277	epoxiconazole	hymexazole	M-314	fenbuconazole	dodemorph
M-278	epoxiconazole	pyrimethanil	M-315	fenbuconazole	fenpropimorph
M-279	epoxiconazole	fludioxonil	M-316	fenbuconazole	iprodione
M-280	epoxiconazole	aldimorph	M-317	fenbuconazole	captan
M-281	epoxiconazole	dodemorph	M-318	fenbuconazole	fenoxanil
M-282	epoxiconazole	fenpropimorph	M-319	fenbuconazole	probenazol
M-283	epoxiconazole	iprodione	M-320	fenbuconazole	mancozeb
M-284	epoxiconazole	captan	M-321	fenbuconazole	metiram
M-285	epoxiconazole	fenoxanil	M-322	fenbuconazole	thiram
M-286	epoxiconazole	probenazol	M-323	fenbuconazole	ziram
M-287	epoxiconazole	mancozeb	M-324	fenbuconazole	guazatin
M-288	epoxiconazole	metiram	M-325	fenbuconazole	thiophanate-methyl
M-289	epoxiconazole	thiram	M-326	fenbuconazole	chlorothalonil
M-290	epoxiconazole	ziram	M-327	fenbuconazole	metrafenone
M-291	epoxiconazole	guazatin	M-328	fluquinconazole	-
M-292	epoxiconazole	thiophanate-methyl	M-329	fluquinconazole	flutriafol
M-293	epoxiconazole	chlorothalonil	M-330	fluquinconazole	ipconazole
M-294	epoxiconazole	metrafenone	M-331	fluquinconazole	metconazole

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-332	fluquinconazole	propiconazole	M-369	flutriafol	prochloraz
M-333	fluquinconazole	prothioconazole	M-370	flutriafol	carbendazim
M-334	fluquinconazole	tebuconazole	M-371	flutriafol	thiabendazole
M-335	fluquinconazole	triadimenol	M-372	flutriafol	ethaboxam
M-336	fluquinconazole	triticonazole	M-373	flutriafol	hymexazole
M-337	fluquinconazole	imazalil	M-374	flutriafol	pyrimethanil
M-338	fluquinconazole	prochloraz	M-375	flutriafol	fludioxonil
M-339	fluquinconazole	carbendazim	M-376	flutriafol	aldimorph
M-340	fluquinconazole	thiabendazole	M-377	flutriafol	dodemorph
M-341	fluquinconazole	ethaboxam	M-378	flutriafol	fenpropimorph
M-342	fluquinconazole	hymexazole	M-379	flutriafol	iprodione
M-343	fluquinconazole	pyrimethanil	M-380	flutriafol	captan
M-344	fluquinconazole	fludioxonil	M-381	flutriafol	fenoxanil
M-345	fluquinconazole	aldimorph	M-382	flutriafol	probenazol
M-346	fluquinconazole	dodemorph	M-383	flutriafol	mancozeb
M-347	fluquinconazole	fenpropimorph	M-384	flutriafol	metiram
M-348	fluquinconazole	iprodione	M-385	flutriafol	thiram
M-349	fluquinconazole	captan	M-386	flutriafol	ziram
M-350	fluquinconazole	fenoxanil	M-387	flutriafol	guazatin
M-351	fluquinconazole	probenazol	M-388	flutriafol	thiophanate-methyl
M-352	fluquinconazole	mancozeb	M-389	flutriafol	chlorothalonil
M-353	fluquinconazole	metiram	M-390	flutriafol	metrafenone
M-354	fluquinconazole	thiram	M-391	ipconazole	-
M-355	fluquinconazole	ziram	M-392	ipconazole	metconazole
M-356	fluquinconazole	guazatin	M-393	ipconazole	propiconazole
M-357	fluquinconazole	thiophanate-methyl	M-394	ipconazole	prothioconazole
M-358	fluquinconazole	chlorothalonil	M-395	ipconazole	tebuconazole
M-359	fluquinconazole	metrafenone	M-396	ipconazole	triadimenol
M-360	flutriafol	-	M-397	ipconazole	triticonazole
M-361	flutriafol	ipconazole	M-398	ipconazole	imazalil
M-362	flutriafol	metconazole	M-399	ipconazole	prochloraz
M-363	flutriafol	propiconazole	M-400	ipconazole	carbendazim
M-364	flutriafol	prothioconazole	M-401	ipconazole	thiabendazole
M-365	flutriafol	tebuconazole	M-402	ipconazole	ethaboxam
M-366	flutriafol	triadimenol	M-403	ipconazole	hymexazole
M-367	flutriafol	triticonazole	M-404	ipconazole	pyrimethanil
M-368	flutriafol	imazalil	M-405	ipconazole	fludioxonil

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-406	ipconazole	aldimorph	M-443	metconazole	metiram
M-407	ipconazole	dodemorph	M-444	metconazole	thiram
M-408	ipconazole	fenpropimorph	M-445	metconazole	ziram
M-409	ipconazole	iprodione	M-446	metconazole	guazatin
M-410	ipconazole	captan	M-447	metconazole	thiophanate-methyl
M-411	ipconazole	fenoxanil	M-448	metconazole	chlorothalonil
M-412	ipconazole	probenazol	M-449	metconazole	metrafenone
M-413	ipconazole	mancozeb	M-450	propiconazole	-
M-414	ipconazole	metiram	M-451	propiconazole	prothioconazole
M-415	ipconazole	thiram	M-452	propiconazole	tebuconazole
M-416	ipconazole	ziram	M-453	propiconazole	triadimenol
M-417	ipconazole	guazatin	M-454	propiconazole	triticonazole
M-418	ipconazole	thiophanate-methyl	M-455	propiconazole	imazalil
M-419	ipconazole	chlorothalonil	M-456	propiconazole	prochloraz
M-420	ipconazole	metrafenone	M-457	propiconazole	carbendazim
M-421	metconazole	-	M-458	propiconazole	thiabendazole
M-422	metconazole	propiconazole	M-459	propiconazole	ethaboxam
M-423	metconazole	prothioconazole	M-460	propiconazole	hymexazole
M-424	metconazole	tebuconazole	M-461	propiconazole	pyrimethanil
M-425	metconazole	triadimenol	M-462	propiconazole	fludioxonil
M-426	metconazole	triticonazole	M-463	propiconazole	aldimorph
M-427	metconazole	imazalil	M-464	propiconazole	dodemorph
M-428	metconazole	prochloraz	M-465	propiconazole	fenpropimorph
M-429	metconazole	carbendazim	M-466	propiconazole	iprodione
M-430	metconazole	thiabendazole	M-467	propiconazole	captan
M-431	metconazole	ethaboxam	M-468	propiconazole	fenoxanil
M-432	metconazole	hymexazole	M-469	propiconazole	probenazol
M-433	metconazole	pyrimethanil	M-470	propiconazole	mancozeb
M-434	metconazole	fludioxonil	M-471	propiconazole	metiram
M-435	metconazole	aldimorph	M-472	propiconazole	thiram
M-436	metconazole	dodemorph	M-473	propiconazole	ziram
M-437	metconazole	fenpropimorph	M-474	propiconazole	guazatin
M-438	metconazole	iprodione	M-475	propiconazole	thiophanate-methyl
M-439	metconazole	captan	M-476	propiconazole	chlorothalonil
M-440	metconazole	fenoxanil	M-477	propiconazole	metrafenone
M-441	metconazole	probenazol	M-478	prothioconazole	-
M-442	metconazole	mancozeb			

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-479	prothioconazole	tebuconazole	M-516	tebuconazole	aldimorph
M-480	prothioconazole	triadimenol	M-517	tebuconazole	dodemorph
M-481	prothioconazole	triticonazole	M-518	tebuconazole	fenpropimorph
M-482	prothioconazole	imazalil	M-519	tebuconazole	iprodione
M-483	prothioconazole	prochloraz	M-520	tebuconazole	captan
M-484	prothioconazole	carbendazim	M-521	tebuconazole	fenoxanil
M-485	prothioconazole	thiabendazole	M-522	tebuconazole	probenazol
M-486	prothioconazole	ethaboxam	M-523	tebuconazole	mancozeb
M-487	prothioconazole	hymexazole	M-524	tebuconazole	metiram
M-488	prothioconazole	pyrimethanil	M-525	tebuconazole	thiram
M-489	prothioconazole	fludioxonil	M-526	tebuconazole	ziram
M-490	prothioconazole	aldimorph	M-527	tebuconazole	guazatin
M-491	prothioconazole	dodemorph	M-528	tebuconazole	thiophanate-methyl
M-492	prothioconazole	fenpropimorph	M-529	tebuconazole	chlorothalonil
M-493	prothioconazole	iprodione	M-530	tebuconazole	metrafenone
M-494	prothioconazole	captan	M-531	triadimenol	-
M-495	prothioconazole	fenoxanil	M-532	triadimenol	triticonazole
M-496	prothioconazole	probenazol	M-533	triadimenol	imazalil
M-497	prothioconazole	mancozeb	M-534	triadimenol	prochloraz
M-498	prothioconazole	metiram	M-535	triadimenol	carbendazim
M-499	prothioconazole	thiram	M-536	triadimenol	thiabendazole
M-500	prothioconazole	ziram	M-537	triadimenol	ethaboxam
M-501	prothioconazole	guazatin	M-538	triadimenol	hymexazole
M-502	prothioconazole	thiophanate-methyl	M-539	triadimenol	pyrimethanil
M-503	prothioconazole	chlorothalonil	M-540	triadimenol	fludioxonil
M-504	prothioconazole	metrafenone	M-541	triadimenol	aldimorph
M-505	tebuconazole	-	M-542	triadimenol	dodemorph
M-506	tebuconazole	triadimenol	M-543	triadimenol	fenpropimorph
M-507	tebuconazole	triticonazole	M-544	triadimenol	iprodione
M-508	tebuconazole	imazalil	M-545	triadimenol	captan
M-509	tebuconazole	prochloraz	M-546	triadimenol	fenoxanil
M-510	tebuconazole	carbendazim	M-547	triadimenol	probenazol
M-511	tebuconazole	thiabendazole	M-548	triadimenol	mancozeb
M-512	tebuconazole	ethaboxam	M-549	triadimenol	metiram
M-513	tebuconazole	hymexazole	M-550	triadimenol	thiram
M-514	tebuconazole	pyrimethanil	M-551	triadimenol	ziram
M-515	tebuconazole	fludioxonil	M-552	triadimenol	guazatin

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-553	triadimenol	thiophanate-methyl	M-589	imazalil	dodemorph
M-554	triadimenol	chlorothalonil	M-590	imazalil	fenpropimorph
M-555	triadimenol	metrafenone	M-591	imazalil	iprodione
M-556	triticonazole	-	M-592	imazalil	captan
M-557	triticonazole	imazalil	M-593	imazalil	fenoxanil
M-558	triticonazole	prochloraz	M-594	imazalil	probenazol
M-559	triticonazole	carbendazim	M-595	imazalil	mancozeb
M-560	triticonazole	thiabendazole	M-596	imazalil	metiram
M-561	triticonazole	ethaboxam	M-597	imazalil	thiram
M-562	triticonazole	hymexazole	M-598	imazalil	ziram
M-563	triticonazole	pyrimethanil	M-599	imazalil	guazatin
M-564	triticonazole	fludioxonil	M-600	imazalil	thiophanate-methyl
M-565	triticonazole	aldimorph	M-601	imazalil	chlorothalonil
M-566	triticonazole	dodemorph	M-602	imazalil	metrafenone
M-567	triticonazole	fenpropimorph	M-603	prochloraz	-
M-568	triticonazole	iprodione	M-604	prochloraz	carbendazim
M-569	triticonazole	captan	M-605	prochloraz	thiabendazole
M-570	triticonazole	fenoxanil	M-606	prochloraz	ethaboxam
M-571	triticonazole	probenazol	M-607	prochloraz	hymexazole
M-572	triticonazole	mancozeb	M-608	prochloraz	pyrimethanil
M-573	triticonazole	metiram	M-609	prochloraz	fludioxonil
M-574	triticonazole	thiram	M-610	prochloraz	aldimorph
M-575	triticonazole	ziram	M-611	prochloraz	dodemorph
M-576	triticonazole	guazatin	M-612	prochloraz	fenpropimorph
M-577	triticonazole	thiophanate-methyl	M-613	prochloraz	iprodione
M-578	triticonazole	chlorothalonil	M-614	prochloraz	captan
M-579	triticonazole	metrafenone	M-615	prochloraz	fenoxanil
M-580	imazalil	-	M-616	prochloraz	probenazol
M-581	imazalil	prochloraz	M-617	prochloraz	mancozeb
M-582	imazalil	carbendazim	M-618	prochloraz	metiram
M-583	imazalil	thiabendazole	M-619	prochloraz	thiram
M-584	imazalil	ethaboxam	M-620	prochloraz	ziram
M-585	imazalil	hymexazole	M-621	prochloraz	guazatin
M-586	imazalil	pyrimethanil	M-622	prochloraz	thiophanate-methyl
M-587	imazalil	fludioxonil	M-623	prochloraz	chlorothalonil
M-588	imazalil	aldimorph	M-624	prochloraz	metrafenone

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-625	carbendazim	-	M-662	thiabendazole	guazatin
M-626	carbendazim	thiabendazole	M-663	thiabendazole	thiophanate-methyl
M-627	carbendazim	ethaboxam	M-664	thiabendazole	chlorothalonil
M-628	carbendazim	hymexazole	M-665	thiabendazole	metrafenone
M-629	carbendazim	pyrimethanil	M-666	ethaboxam	-
M-630	carbendazim	fludioxonil	M-667	ethaboxam	hymexazole
M-631	carbendazim	aldimorph	M-668	ethaboxam	pyrimethanil
M-632	carbendazim	dodemorph	M-669	ethaboxam	fludioxonil
M-633	carbendazim	fenpropimorph	M-670	ethaboxam	aldimorph
M-634	carbendazim	iprodione	M-671	ethaboxam	dodemorph
M-635	carbendazim	captan	M-672	ethaboxam	fenpropimorph
M-636	carbendazim	fenoxanil	M-673	ethaboxam	iprodione
M-637	carbendazim	probenazol	M-674	ethaboxam	captan
M-638	carbendazim	mancozeb	M-675	ethaboxam	fenoxanil
M-639	carbendazim	metiram	M-676	ethaboxam	probenazol
M-640	carbendazim	thiram	M-677	ethaboxam	mancozeb
M-641	carbendazim	ziram	M-678	ethaboxam	metiram
M-642	carbendazim	guazatin	M-679	ethaboxam	thiram
M-643	carbendazim	thiophanate-methyl	M-680	ethaboxam	ziram
M-644	carbendazim	chlorothalonil	M-681	ethaboxam	guazatin
M-645	carbendazim	metrafenone	M-682	ethaboxam	thiophanate-methyl
M-646	thiabendazole	-	M-683	ethaboxam	chlorothalonil
M-647	thiabendazole	ethaboxam	M-684	ethaboxam	metrafenone
M-648	thiabendazole	hymexazole	M-685	hymexazole	-
M-649	thiabendazole	pyrimethanil	M-686	hymexazole	pyrimethanil
M-650	thiabendazole	fludioxonil	M-687	hymexazole	fludioxonil
M-651	thiabendazole	aldimorph	M-688	hymexazole	aldimorph
M-652	thiabendazole	dodemorph	M-689	hymexazole	dodemorph
M-653	thiabendazole	fenpropimorph	M-690	hymexazole	fenpropimorph
M-654	thiabendazole	iprodione	M-691	hymexazole	iprodione
M-655	thiabendazole	captan	M-692	hymexazole	captan
M-656	thiabendazole	fenoxanil	M-693	hymexazole	fenoxanil
M-657	thiabendazole	probenazol	M-694	hymexazole	probenazol
M-658	thiabendazole	mancozeb	M-695	hymexazole	mancozeb
M-659	thiabendazole	metiram	M-696	hymexazole	metiram
M-660	thiabendazole	thiram	M-697	hymexazole	thiram
M-661	thiabendazole	ziram			

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-698	hymexazole	ziram			methyl
M-699	hymexazole	guazatin	M-734	fludioxonil	chlorothalonil
M-700	hymexazole	thiophanate-methyl	M-735	fludioxonil	metrafenone
M-701	hymexazole	chlorothalonil	M-736	aldimorph	-
M-702	hymexazole	metrafenone	M-737	aldimorph	dodemorph
M-703	pyrimethanil	-	M-738	aldimorph	fenpropimorph
M-704	pyrimethanil	fludioxonil	M-739	aldimorph	iprodione
M-705	pyrimethanil	aldimorph	M-740	aldimorph	captan
M-706	pyrimethanil	dodemorph	M-741	aldimorph	fenoxanil
M-707	pyrimethanil	fenpropimorph	M-742	aldimorph	probenazol
M-708	pyrimethanil	iprodione	M-743	aldimorph	mancozeb
M-709	pyrimethanil	captan	M-744	aldimorph	metiram
M-710	pyrimethanil	fenoxanil	M-745	aldimorph	thiram
M-711	pyrimethanil	probenazol	M-746	aldimorph	ziram
M-712	pyrimethanil	mancozeb	M-747	aldimorph	guazatin
M-713	pyrimethanil	metiram	M-748	aldimorph	thiophanate-methyl
M-714	pyrimethanil	thiram	M-749	aldimorph	chlorothalonil
M-715	pyrimethanil	ziram	M-750	aldimorph	metrafenone
M-716	pyrimethanil	guazatin	M-751	dodemorph	-
M-717	pyrimethanil	thiophanate-methyl	M-752	dodemorph	fenpropimorph
M-718	pyrimethanil	chlorothalonil	M-753	dodemorph	iprodione
M-719	pyrimethanil	metrafenone	M-754	dodemorph	captan
M-720	fludioxonil	-	M-755	dodemorph	fenoxanil
M-721	fludioxonil	aldimorph	M-756	dodemorph	probenazol
M-722	fludioxonil	dodemorph	M-757	dodemorph	mancozeb
M-723	fludioxonil	fenpropimorph	M-758	dodemorph	metiram
M-724	fludioxonil	iprodione	M-759	dodemorph	thiram
M-725	fludioxonil	captan	M-760	dodemorph	ziram
M-726	fludioxonil	fenoxanil	M-761	dodemorph	guazatin
M-727	fludioxonil	probenazol	M-762	dodemorph	thiophanate-methyl
M-728	fludioxonil	mancozeb	M-763	dodemorph	chlorothalonil
M-729	fludioxonil	metiram	M-764	dodemorph	metrafenone
M-730	fludioxonil	thiram	M-765	fenpropimorph	-
M-731	fludioxonil	ziram	M-766	fenpropimorph	iprodione
M-732	fludioxonil	guazatin	M-767	fenpropimorph	captan
M-733	fludioxonil	thiophanate-	M-768	fenpropimorph	fenoxanil

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-769	fenpropimorph	probenazol	M-804	fenoxanil	metiram
M-770	fenpropimorph	mancozeb	M-805	fenoxanil	thiram
M-771	fenpropimorph	metiram	M-806	fenoxanil	ziram
M-772	fenpropimorph	thiram	M-807	fenoxanil	guazatin
M-773	fenpropimorph	ziram	M-808	fenoxanil	thiophanate-methyl
M-774	fenpropimorph	guazatin	M-809	fenoxanil	chlorothalonil
M-775	fenpropimorph	thiophanate-methyl	M-810	fenoxanil	metrafenone
M-776	fenpropimorph	chlorothalonil	M-811	probenazol	-
M-777	fenpropimorph	metrafenone	M-812	probenazol	mancozeb
M-778	iprodione	-	M-813	probenazol	metiram
M-779	iprodione	captan	M-814	probenazol	thiram
M-780	iprodione	fenoxanil	M-815	probenazol	ziram
M-781	iprodione	probenazol	M-816	probenazol	guazatin
M-782	iprodione	mancozeb	M-817	probenazol	thiophanate-methyl
M-783	iprodione	metiram	M-818	probenazol	chlorothalonil
M-784	iprodione	thiram	M-819	probenazol	metrafenone
M-785	iprodione	ziram	M-820	mancozeb	-
M-786	iprodione	guazatin	M-821	mancozeb	metiram
M-787	iprodione	thiophanate-methyl	M-822	mancozeb	thiram
M-788	iprodione	chlorothalonil	M-823	mancozeb	ziram
M-789	iprodione	metrafenone	M-824	mancozeb	guazatin
M-790	captan	-	M-825	mancozeb	thiophanate-methyl
M-791	captan	fenoxanil	M-826	mancozeb	chlorothalonil
M-792	captan	probenazol	M-827	mancozeb	metrafenone
M-793	captan	mancozeb	M-828	metiram	-
M-794	captan	metiram	M-829	metiram	thiram
M-795	captan	thiram	M-830	metiram	ziram
M-796	captan	ziram	M-831	metiram	guazatin
M-797	captan	guazatin	M-832	metiram	thiophanate-methyl
M-798	captan	thiophanate-methyl	M-833	metiram	chlorothalonil
M-799	captan	chlorothalonil	M-834	metiram	metrafenone
M-800	captan	metrafenone	M-835	thiram	-
M-801	fenoxanil	-	M-836	thiram	ziram
M-802	fenoxanil	probenazol	M-837	thiram	guazatin
M-803	fenoxanil	mancozeb			

Mixture No.	Compound IIB1	Compound IIB2	Mixture No.	Compound IIB1	Compound IIB2
M-838	thiram	thiophanate-methyl			methyl
M-839	thiram	chlorothalonil	M-848	guazatin	chlorothalonil
M-840	thiram	metrafenone	M-849	guazatin	metrafenone
M-841	ziram	-	M-850	thiophanate-methyl	-
M-842	ziram	guazatin	M-851	thiophanate-methyl	chlorothalonil
M-843	ziram	thiophanate-methyl	M-852	thiophanate-methyl	metrafenone
M-844	ziram	chlorothalonil	M-853	chlorothalonil	-
M-845	ziram	metrafenone	M-854	chlorothalonil	metrafenone
M-846	guazatin	-	M-855	metrafenone	-
M-847	guazatin	thiophanate-			

The crystalline modification II and the one or more compound(s) of groups A.1 – A.15 are usually applied in a weight ratio of from 500:1 to 1:100, preferably from 20:1 to 1:50, in particular from 5:1 to 1:20.

5

The afore-mentioned applies also to the ratios of combinations of modification II with fungicidal compounds IIA. Compounds IIB are usually combined with modification II in ratios from 100:1 to 1:100.

- 10 Depending on the desired effect, the application rates of the mixtures according to the invention are from 5 g/ha to 2000 g/ha, preferably from 50 to 1500 g/ha, in particular from 50 to 750 g/ha.

- 15 The crystalline modification II, the mixtures and the compositions according to the invention can be applied to any and all developmental stages, such as egg, larva, pupa, and adult. The pests may be controlled by contacting the target pest, its food supply, habitat, breeding ground or its locus with a pesticidally effective amount of the crystalline modification II, the mixtures or the compositions according to the invention.

- 20 "Locus" means a plant, seed, soil, area, material or environment in which a pest is growing or may grow.

- 25 In general, "pesticidally effective amount" means the amount of the crystalline modification II, the mixtures and the compositions according to the invention needed to achieve an observable effect on growth, including the effects of necrosis, death, retardation, prevention, and removal, destruction, or otherwise diminishing the occurrence and activity of the target organism. The pesticidally effective amount can

vary for the various mixtures / compositions used in the invention. A pesticidally effective amount of the mixtures / compositions will also vary according to the prevailing conditions such as desired pesticidal effect and duration, weather, target species, locus, mode of application, and the like.

5

The crystalline modification II, the mixtures and the compositions according to the invention can also be employed for protecting plants from attack or infestation by insects, acarids or nematodes comprising contacting a plant, or soil or water in which the plant is growing.

10

In the context of the present invention, the term plant refers to an entire plant, a part of the plant or the propagation material of the plant, that is, the seed or the seedling.

Plants which can be treated with the crystalline modification II, the mixtures and the compositions according to the invention include all genetically modified plants or transgenic plants, e.g. crops which tolerate the action of herbicides or fungicides or insecticides owing to breeding, including genetic engineering methods, or plants which have modified characteristics in comparison with existing plants, which can be generated for example by traditional breeding methods and/or the generation of mutants, or by recombinant procedures.

20

Some of the inventive mixtures and compositions have systemic action and can therefore be used for the protection of the plant shoot against foliar pests as well as for the treatment of the seed and roots against soil pests. The term seed treatment comprises all suitable seed treatment techniques known in the art, such as, but not limited to, seed dressing, seed coating, seed dusting, seed soaking, seed film coating, seed multilayer coating, seed encrusting, seed dripping, and seed pelleting.

25

The present invention also comprises seeds coated with or containing the crystalline modification II or the mixtures or the compositions according to the invention.

30

The term seed embraces seeds and plant propagules of all kinds including but not limited to true seeds, seed pieces, suckers, corms, bulbs, fruit, tubers, grains, cuttings, cut shoots and the like and means in a preferred embodiment true seeds.

35

Suitable seed is seed of cereals, root crops, oil crops, vegetables, spices, ornamentals, for example seed of durum and other wheat, barley, oats, rye, maize (fodder maize and sugar maize / sweet and field corn), soybeans, oil crops, crucifers, cotton, sunflowers, bananas, rice, oilseed rape, turnip rape, sugarbeet, fodder beet, eggplants, potatoes, grass, lawn, turf, fodder grass, tomatoes, leeks, pumpkin/squash, cabbage, iceberg lettuce, pepper, cucumbers, melons, Brassica species, melons, beans, peas, garlic,

40

onions, carrots, tuberous plants such as potatoes, sugar cane, tobacco, grapes, petunias, geranium/pelargoniums, pansies and impatiens.

5 In addition, the crystalline modification II, the mixtures and the compositions according to the invention may also be used for the treatment seeds from plants, which tolerate the action of herbicides or fungicides or insecticides or nematocides owing to breeding, mutation and/or genetic engineering methods.

10 For example, the crystalline modification II, the mixtures and the compositions according to the invention can be employed in transgenic crops which are resistant to herbicides from the group consisting of the sulfonylureas (EP-A-0257993, U.S. Pat. No. 5,013,659), imidazolinones (see for example US 6222100, WO0182685, WO0026390, WO9741218, WO9802526, WO9802527, WO 04/106529, WO 05/20673, WO 03/14357, WO 03/13225, WO 03/14356, WO 04/16073), glufosinate-type (see for
15 example EP-A-0242236, EP-A-242246) or glyphosate-type (see for example WO 92/00377) or in plants resistant towards herbicides selected from the group of cyclohexadienone/aryloxyphenoxypropionic acid herbicides (US 5,162,602 , US 5,290,696 , US 5,498,544 , US 5,428,001 , US 6,069,298 , US 6,268,550 , US 6,146,867 , US 6,222,099 , US 6,414,222) or in transgenic crop plants, for example
20 cotton, with the capability of producing *Bacillus thuringiensis* toxins (Bt toxins) which make the plants resistant to certain pests (EP-A-0142924, EP-A-0193259).

Furthermore, the crystalline modification II, the mixtures and the compositions according to the invention can be used also for the treatment of seeds from plants,
25 which have modified characteristics in comparison with existing plants consist, which can be generated, for example by traditional breeding methods and/or the generation of mutants, or by recombinant procedures). For example, a number of cases have been described of recombinant modifications of crop plants for the purpose of modifying the starch synthesized in the plants (e.g. WO 92/11376, WO 92/14827, WO
30 91/19806) or of transgenic crop plants having a modified fatty acid composition (WO 91/13972).

The seed treatment application of the crystalline modification II, the mixtures and the compositions according to the invention is carried out by spraying or dusting the seeds
35 before sowing of the plants and before emergence of the plants.

In the treatment of seeds the corresponding formulations are applied by treating the seeds with an effective amount of the crystalline modification II, the mixtures or the compositions according to the invention. Herein, the application rates of the crystalline
40 modification II are generally from 0.1 g to 10 kg per 100 kg of seed, preferably from 1 g

to 5 kg per 100 kg of seed, in particular from 1 g to 2.5 kg per 100 kg of seed. For specific crops such as lettuce and onions the rates can be higher.

5 The mixtures and the compositions according to the invention are effective through both contact (via soil, glass, wall, bed net, carpet, plant parts or animal parts), and ingestion (bait, or plant part) and through trophallaxis and transfer.

10 Preferred application methods are into water bodies, via soil, cracks and crevices, pastures, manure piles, sewers, into water, on floor, wall, or by perimeter spray application and bait.

15 According to another preferred embodiment of the invention, for use against non-crop pests such as ants, termites, wasps, flies, mosquitoes, crickets, locusts, or cockroaches the mixtures and the compositions according to the invention are prepared into a bait preparation.

20 The bait can be a liquid, a solid or a semisolid preparation (e.g. a gel). The bait employed in the mixtures/compositions is a product which is sufficiently attractive to incite insects such as ants, termites, wasps, flies, mosquitoes, crickets etc. or cockroaches to eat it. This attractant may be chosen from feeding stimulants or para and / or sex pheromones readily known in the art.

25 Methods to control infectious diseases transmitted by insects (e.g. malaria, dengue and yellow fever, lymphatic filariasis, and leishmaniasis) with the inventive mixtures and their respective compositions also comprise treating surfaces of huts and houses, air spraying and impregnation of curtains, tents, clothing items, bed nets, tsetse-fly trap or the like. Insecticidal compositions for application to fibers, fabric, knitgoods, nonwovens, netting material or foils and tarpaulins preferably comprise a composition including the inventive mixtures, optionally a repellent and at least one binder.

30 The crystalline modification II, the mixtures and the compositions according to the invention can be used for protecting wooden materials such as trees, board fences, sleepers, etc. and buildings such as houses, outhouses, factories, but also construction materials, furniture, leathers, fibers, vinyl articles, electric wires and cables etc. from
35 ants and/or termites, and for controlling ants and termites from doing harm to crops or human being (e.g. when the pests invade into houses and public facilities).

40 In the case of soil treatment or of application to the pests dwelling place or nest, the quantity of active ingredient ranges from 0.0001 to 500 g per 100 m², preferably from 0.001 to 20 g per 100 m².

Customary application rates in the protection of materials are, for example, from 0.01 g to 1000 g of active compound per m² treated material, desirably from 0.1 g to 50 g per m².

- 5 Insecticidal compositions for use in the impregnation of materials typically contain from 0.001 to 95 weight %, preferably from 0.1 to 45 weight %, and more preferably from 1 to 25 weight % of at least one repellent and / or insecticide.

- 10 For use in bait compositions, the typical content of active ingredient(s) is from 0.0001 weight % to 15 weight %, desirably from 0.001 weight % to 5% weight % of active compound. The composition used may also comprise other additives such as a solvent of the active material, a flavoring agent, a preserving agent, a dye or a bitter agent. Its attractiveness may also be enhanced by a special color, shape or texture.

- 15 For use in spray compositions, the content of the active ingredient(s) is from 0.001 to 80 weights %, preferably from 0.01 to 50 weight % and most preferably from 0.01 to 15 weight %.

- 20 For use in treating crop plants, the rate of application of the active ingredient(s) may be in the range of 0.1 g to 4000 g per hectare, desirably from 25 g to 600 g per hectare, more desirably from 50 g to 500 g per hectare.

- 25 It was also an object of the present invention to provide mixtures suitable for treating, controlling, preventing and protecting warm-blooded animals, including humans, and fish against infestation and infection by pests. Problems that may be encountered with pest control on or in animals and/or humans are similar to those described at the outset, namely the need for reduced dosage rates, and / or enhanced spectrum of activity and / or combination of knock-down activity with prolonged control and / or resistance management.

- 30 This invention also provides a method for treating, controlling, preventing and protecting warm-blooded animals, including humans, and fish against infestation and infection by pests of the orders Siphonaptera, Hymenoptera, Hemiptera, Orthoptera, Acarina, Phthiraptera, and Diptera, which comprises orally, topically or parenterally
35 administering or applying to said animals a pesticidally effective amount of the crystalline modification II, the mixtures and the compositions according to the invention.

- 40 The invention also provides a process for the preparation of a composition for treating, controlling, preventing or protecting a warm-blooded animal or a fish against infestation or infection by pests of the Siphonaptera, Hymenoptera, Hemiptera, Orthoptera, Acarina, Phthiraptera, and Diptera orders which comprises a pesticidally effective

amount of the crystalline modification II, the mixtures and the compositions according to the invention.

5 The above method is particularly useful for controlling and preventing infestations and infections in warm-blooded animals such as cattle, sheep, swine, camels, deer, horses, poultry, goats, dogs and cats as well as humans.

10 Infestations in warm-blooded animals and fish including, but not limited to, lice, biting lice, ticks, nasal bots, keds, biting flies, muscoid flies, flies, myiasitic fly larvae, chiggers, gnats, mosquitoes and fleas may be controlled, prevented or eliminated by the crystalline modification II, the mixtures and the compositions according to the invention.

15 For oral administration to warm-blooded animals, the crystalline modification II, the mixtures and the compositions according to the invention may be formulated as animal feeds, animal feed premixes, animal feed concentrates, pills, solutions, pastes, suspensions, drenches, gels, tablets, boluses and capsules. In addition, the crystalline modification II, the mixtures and the compositions according to the invention may be administered to the animals in their drinking water. For oral administration, the dosage
20 form chosen should provide the animal with 0.01 mg/kg to 100 mg/kg of animal body weight per day of the crystalline modification II, the mixtures and the compositions according to the invention.

25 Alternatively, the crystalline modification II, the mixtures and the compositions according to the invention may be administered to animals parenterally, for example, by intraruminal, intramuscular, intravenous or subcutaneous injection. The crystalline modification II, the mixtures and the compositions according to the invention may be dispersed or dissolved in a physiologically acceptable carrier for subcutaneous injection. Alternatively, the crystalline modification II, the mixtures and the compositions
30 according to the invention may be formulated into an implant for subcutaneous administration. In addition, the crystalline modification II, the mixtures and the compositions according to the invention may be transdermally administered to animals. For parenteral administration, the dosage form chosen should provide the animal with 0.01 mg/kg to 100 mg/kg of animal body weight per day of the crystalline modification
35 II, the mixtures and the compositions according to the invention.

The crystalline modification II, the mixtures and the compositions according to the invention may also be applied topically to the animals in the form of dips, dusts, powders, collars, medallions, sprays, spot-on and pour-on formulations. For topical
40 application, dips and sprays usually contain 0.5 ppm to 5000 ppm and preferably 1 ppm to 3000 ppm of the crystalline modification II. In addition, the crystalline

modification II may be formulated as ear tags for animals, particularly quadrupeds such as cattle and sheep.

5 The figure and examples below serve to illustrate the invention and are not to be understood as limiting it.

Figure 1: X-ray powder diffractogram of modification II

Figure 2: Differential scanning calorimetry thermogram of modification II

Figure 3: Thermogravimetric analysis of modification II

10 Figure 4: X-ray powder diffractograms of mixtures of modifications I and V

Preparation examples

Example 1: Characterization of a solid form of fipronil as starting material

15 All preparation procedures below were conducted with two samples of solid fipronil as starting materials which were obtained according to procedures as described in WO 2001/30760, with final crystallization of the product from a solvent mixture of MCB/ethanole (% by weight of ethanol at crystallization start: 13 %) at temperatures of 70°C to 35°C. This solid form in X-ray powder diffractogram studies proved to be
20 crystalline fipronil of a mixture of several crystalline modifications. This mixture has been characterized to consist of crystalline modification I and crystalline modification V, as for the first time identified and described in a co-pending patent application. A least squares refinement with the Topas program with simulated X-ray powder diffractogram patterns from single crystal data of form I and form V shows that in these two example
25 samples, the percentage of form I varies from 30% to 70 %. X-ray powder diffractograms of the two samples are shown in figure 4.

Irrespective of the sample of solid fipronil used as starting material, the crystallization examples given below yielded in modification II.

30

Example 2: Preparation of modification II by crystallization from THF

1,0 g of crystalline fipronil having a chemical purity of about 96 % by weight was dissolved in 25 ml of THF at 45 to 50°C_[B7] in a round bottomed flask. The solution was kept at this temperature while the solvent was slowly evaporated with a gentle flow of
35 inert N₂ gas. The solvent was left to evaporate for about 15 hours, after which the obtained crystalline material was filtered from some residual solvent on a paper filter. The material obtained has the X-ray powder diffractogram shown in Figure 1 with the reflexes listed in Table 2 below.

Example 3: Preparation of modification II by crystallization from 1,2-dichloroethane (DCE)

1,0 g of crystalline fipronil having a chemical purity of about 96 % by weight was dissolved in 25 ml of DCE at 45°C to 50°C in a round bottomed flask. Then the heating
5 was switched off and the oil bath was left to cool to 20°C to 25°C. The solvent was left to evaporate for about 15 hours. The material obtained has the X-ray powder diffractogram shown in Figure 1 with the reflexes listed in Table 2 below.

10 Example 4: Preparation of modification II by crystallization from acetonitrile
2,0 g of crystalline fipronil having a chemical purity of about 96 % by weight was dissolved in 4 ml of acetonitrile at 75°C to 80°C in a 10 ml test tube to give a clear solution. The solution was cooled down to 5°C in approximately 8h and left at 5°C over night. The supernatant was removed by using a pipet. The solid was dried on a filter
15 paper for a few hours and then analyzed to give within a 0.3° 2θ error marginal the X-ray powder diffractogram shown in Figure 1.

Example 5: Preparation of modification II by crystallization from toluene
0,7 g of fipronil having chemical purity of >98 % by weight was suspended in 10 ml of
20 toluene in a 30 ml glass reactor. Stirring was applied and the suspension was heated up to 110 °C where the solid was completely dissolved. The solution was then cooled down to 25 °C with a cooling rate of -1 K/min. The crystallization was observed to begin at 86 °C. The solid product was filtrated and dried on filtration paper over night and then analyzed to give within a 0.3° 2θ error marginal the X-ray powder diffractogram
25 shown in Figure 1. The content of fipronil in this experiment was 8.2 w-%.

Example 6: Preparation of modification II by crystallization from monochlorobenzene (MCB)
0,60 g of fipronil having chemical purity of >98 % by weight was suspended in 15 ml of
30 MCB. Stirring was applied and the suspension was heated up to 137 °C where the solid was completely dissolved. The solution was then cooled down to 25 °C with a cooling rate of -1 K/min. The crystallization was observed to begin at 76 °C. The solid product was filtrated and dried on filtration paper over night and then analyzed to give within a 0.3° 2θ error marginal the X-ray powder diffractogram shown in Figure 1.

35 Example 7: Preparation of modification II by crystallization from mesitylene
2g of fipronil having chemical purity of ~96 % by weight was dissolved at 15 ml of mesitylene. The solution was cooled down within 5 hours to 20-25°C and left to stand over night. The crystallized solid was filtered and dried in a vacuum oven (at 10 mbar)
40 for ~1.5 h at 40°C. The sample was analyzed to give within a 0.3° 2θ error marginal the X-ray powder diffractogram shown in Figure 1[B8].

Example 8: Preparation of modification II by crystallization from nitrobenzene
 2g of fipronil having chemical purity of about 96 % by weight was dissolved at 5 ml of nitrobenzene. The solution was cooled down within 5 hours to 20-25°C and left to stand over night. The crystallized solid was filtered and dried in a vacuum oven (at 10 mbar)
 5 for about 1.5 h at 40°C. The sample was analyzed to give within a 0.3° 2θ error marginal the X-ray powder diffractogram shown in Figure 1[B9].

Example 9: Preparation of modification II by crystallization from ethyl benzene
 1.5 g of fipronil having chemical purity of >98 % by weight was suspended in 10 ml of ethyl benzene. Stirring was applied and the suspension was heated up to 136°C giving
 10 a clear solution. The solution was cooled down to 25°C with a cooling rate of -1 K/min. The crystallization was detected to begin at 118°C. The solid product was filtrated and dried on filtration paper over night and then analyzed to give within a 0.3° 2θ error marginal the X-ray powder diffractogram shown in Figure 1.

Example 10: Preparation of modification II by crystallization from CF₃-benzene
 1.5 g of fipronil having chemical purity of >98 % by weight was suspended in 18.5 ml of CF₃-benzene. Stirring was applied and the suspension was heated up to 103 °C giving
 a clear solution. The solution was cooled down to 25°C with a cooling rate of -1 K/min.
 20 The crystallization was detected to begin at 97°C. The solid product was filtrated and dried on filtration paper over night at 20-25°C. The sample was analyzed to be within 0.3° 2θ error marginal the X-ray powder diffractogram shown in Figure 1.

Table 2: d-spacings and 2θ-angles of modification II

d (Å)	2θ
d = 13.44 ± 0.2 Å	6.6 ± 0.2°
d = 7.84 ± 0.1 Å	11.3 ± 0.2°
d = 5.50 ± 0.07 Å	16.1 ± 0.2°
d = 5.14 ± 0.05 Å	17.2 ± 0.2°
d = 4.95 ± 0.05 Å	17.9 ± 0.2°
d = 3.95 ± 0.05 Å	22.4 ± 0.2°
d = 3.77 ± 0.05 Å.	23.5 ± 0.2°
d = 3.22 ± 0.03 Å	27.6 ± 0.2°
d = 2.91 ± 0.03 Å	30.8 ± 0.2°

25

Analysis:

The X-ray powder diffractogram displayed in Figure 1 was recorded using a Siemens D-5000 diffractometer (manufacturer: Bruker AXS) in reflection geometry in the range
 30 from 2θ = 2°- 60° with increments of 0.02° using Cu-Kα radiation at 25°C. The 2θ values found were used to calculate the stated interplanar spacing d. In Figure 1, the

intensity of the peaks (y-axis: linear intensity in counts) is plotted versus the 2θ angle (x-axis in degrees 2θ).

The single crystal X-ray diffraction data was collected on a Bruker AXS CCD Detector using graphite $\text{Cu}_{K\alpha}$ radiation. The structure was solved by using direct methods, refined, and expanded by using Fourier techniques with the SHELX software package (G.M. Sheldrick, SHELX-97, Universität Göttingen, 1997). Absorption correction was performed with SADABS software.

Melting points indicated herein refer to values determined on a Mettler hot stage microscope and represent equilibrium melting points.

DSC was performed on a Mettler Toledo DSC 823 module in air atmosphere. Crystals taken from the mother liquor were blotted dry on filter paper and place in crimped but vented aluminum sample pans for the DCS experiment. The sample size in each case was 5 to 10 mg. The temperature range was typically 30°C to 230°C at a heating rate of $5^\circ\text{C}/\text{min}$.

TGA measurements were performed on a SEIKO Instrument in nitrogen atmosphere in platinum pans. The sample size in each case was $\sim 8\text{-}10$ mg. The temperature range was 30°C to 600°C at a heating rate $10^\circ\text{C}/\text{min}$.

Investigations of the effect of the temperature of nucleation and beginning of the crystallization to the modification of the crystalline end product carried out with Polyblock by HEL Ltd. The multi reactor crystallization system allows the monitoring of the crystallization process and change in turbidity with special reflectance turbidity probes by HEL. The heating/cooling mantle and thermostat "Julabo FP 50" as well as the turbidity probes were controlled with a PC.

Comprises when used in this specification is taken to specify the presence of stated features, integers, steps or components but does not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof.

58

Claims:

1. A crystalline modification II of fipronil which has an X-ray powder diffractogram showing, at 25°C, at least 5 of the following reflexes:

5

(1) $d = 13,44 \pm 0,2 \text{ \AA}$ (2) $d = 7,84 \pm 0,1 \text{ \AA}$ (3) $d = 5,50 \pm 0,07 \text{ \AA}$ (4) $d = 5,14 \pm 0,05 \text{ \AA}$

10

(5) $d = 4,95 \pm 0,05 \text{ \AA}$ (6) $d = 3,95 \pm 0,05 \text{ \AA}$ (7) $d = 3,77 \pm 0,05 \text{ \AA}$ (8) $d = 3,22 \pm 0,03 \text{ \AA}$ (9) $d = 2,91 \pm 0,03 \text{ \AA}$

15

2. The crystalline modification II according to claim 1 undergoing a phase transformation at 105°C to 145°C into two other crystalline modifications I and V of fipronil having melting points at 196 to 197°C and at 202 to 203°C, respectively.

20

3. The crystalline modification II according to claims 1 or 2 having a fipronil content of at least 98% by weight.

4. Solid fipronil comprising the crystalline modification II as defined in any of claims 1 to 3 and a form of fipronil being different from crystalline modification II.

25

5. Solid fipronil according to claim 4 comprising the crystalline modification II as defined in any of claims 1 to 3 in at least 85% by weight.

6. A process for preparing the crystalline modification II as defined in any of claims 1 to 3, comprising the steps of:

30

step i) preparing a solution of a solid form of fipronil being different from the crystalline modification II in a solvent S selected from tetrahydrofuran and 1,2-dichloroethane;

35

step ii) effecting crystallization of fipronil; and
step iii) isolating the resulting precipitate.

7. The process according to claim 6 or 7, wherein the solvent S is tetrahydrofuran.

8. The process according to claim 6 or 7, wherein the solvent S is 1,2-dichloroethane.

40

AMENDED SHEET

- 5 9. The process according to any of claims 6 to 8, wherein, in step ii), the crystallization of fipronil is effected by concentration of the solution obtained in step i).
- 10 10. The process according to any of claims 6 to 8, wherein, in step ii), the crystallization of fipronil is effected by adding a solvent which reduces the solubility.
- 15 11. The process according to any of claims 6 to 8, wherein step ii) is carried out in the presence of seed crystals of the crystalline modification II as defined in any of claims 1 to 3.
- 20 12. A synergistic pesticidal or parasitocidal mixture comprising, as active components, the crystalline modification II as defined in any of claims 1 to 3 and one or more pesticidal or parasitocidal compounds.
- 25 13. A pesticidal or parasitocidal composition, comprising the crystalline modification II as defined in any of claims 1 to 3 or the mixture as defined in claim 12 and pesticidally or parasitocidally acceptable carriers and/or auxiliaries.
- 30 14. The composition according to claim 13 in the form of an aqueous suspension concentrate.
- 35 15. The composition according to claim 13 in the form of water-dispersible granules.
- 40 16. The composition according to claim 13 in the form of a water-dispersible powder.
17. Use of the crystalline modification II as defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16 for controlling pests.
18. A method for controlling pests which comprises contacting the pests or their food supply, habitat, breeding grounds or their locus with a pesticidally effective amount of the crystalline modification II as defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16.
19. A method for protecting a plant from infestation and attack by pests which comprises applying to the foliage or stem of said plant a pesticidally effective

60

amount of the crystalline modification II as defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16.

- 5 20. A method as claimed in any of claims 22 or 23, wherein the crystalline modification II as defined in any of claims 1 to 3 or the solid fipronil as defined in claims 4 or 5 or the mixture as defined in claim 12 or the composition as defined in any of claims 13 to 16 are applied in an amount of from 5 g/ha to 2000 g/ha.
- 10 21. The use according to claim 17 or the method according to any of claims 18 to 20 wherein the pests are insects, arachnids or plant nematodes.
- 15 22. A method of the protection of seed comprising contacting the seeds with the crystalline modification II as defined in any of claims 1 to 3 or with the solid fipronil as defined in claims 4 or 5 or with the mixture as defined in claim 12 or with the composition as defined in any of claims 13 to 16 in pesticidally effective amounts.
- 20 23. A method as claimed in claim 23 wherein the crystalline modification II as defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16 is applied in an amount of from 0.1 g to 10 kg per 100 kg of seeds.
- 25 24. Seed comprising the crystalline modification II as defined in any of claims 1 to 3 or the solid fipronil as defined in claims 5 or 6 or the mixture as defined in claim 12 in an amount of from 0.1 g to 10 kg per 100 kg of seeds.
- 30 25. Use of the crystalline modification II as defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16 for combating parasites in and on animals.
- 35 26. A method for treating, controlling, preventing or protecting animals against infestation or infection by parasites which comprises orally, topically or parenterally administering or applying to the animals a parasitically effective amount of the crystalline modification II as defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16.
- 40 27. A process for the preparation of a composition for treating, controlling, preventing or protecting animals against infestation or infection by parasites which comprises a parasitically effective amount of the crystalline modification II as


AMENDED SHEET

61

defined in any of claims 1 to 3 or of the solid fipronil as defined in claims 4 or 5 or of the mixture as defined in claim 12 or of the composition as defined in any of claims 13 to 16.

AMENDED SHEET

Figure 1. Powder diffractogram of modification II






Figure 3. Differential Scanning Calorimetry Thermogram

