



(11) (21) (C) **2,199,133**
(86) 1995/08/24
(87) 1996/04/18
(45) 2000/07/18

(72) LAGNEMO, Hans, SE

(72) JIGSTAM, Monica, SE

(72) ODDSTIG, Cecilia, SE

(73) Eka Chemicals AB, SE

(51) Int.Cl.⁶ C11D 7/18, C11D 7/22, C11D 17/06, C11D 7/02

(30) 1994/10/07 (SE94/00944) SE

(30) 1994/10/07 (SE94/00943) SE

(30) 1994/11/03 (9403778-5) SE

(54) **AGENT DE BLANCHIMENT**

(54) **BLEACHING AGENT**

(57) L'invention concerne des particules, ou un de leurs sels, qui comprennent un composé peroxy, pouvant libérer du peroxyde d'hydrogène ou des acides peroxy dans des solutions aqueuses, et jusqu'à 15 % en poids d'un acide dicarboxylique comportant 5 à 10 atomes de carbone. L'invention concerne aussi l'utilisation de ces particules et une composition les comprenant.

(57) The invention relates to particles comprising a peroxy compound with capability of releasing hydrogen peroxide or peroxy acids in aqueous solutions and up to 15 % by weight of a dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof. The invention also relates to use of the particles and a composition comprising such particles.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :**C11D 3/39, C01B 15/10****A1**

(11) International Publication Number:

WO 96/11252

(43) International Publication Date:

18 April 1996 (18.04.96)

(21) International Application Number: **PCT/SE95/00959**(22) International Filing Date: **24 August 1995 (24.08.95)**

(30) Priority Data:

PCT/SE94/00943 7 October 1994 (07.10.94) WO

(34) Countries for which the regional or
international application was filed: SE et al.

PCT/SE94/00944 7 October 1994 (07.10.94) WO

(34) Countries for which the regional or
international application was filed: SE et al.

9403778-5 3 November 1994 (03.11.94) SE

(71) Applicant (for all designated States except US): **EKA NOBEL
AB [SE/SE]; S-445 80 Bohus (SE).**

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LAGNEMO, Hans
[SE/SE]; Kung Håkons gata 10, S-417 28 Göteborg (SE).
JIGSTAM, Monica [SE/SE]; Hästeviksvägen, Pl 785, S-423
41 Torslanda (SE). ODDSTIG, Cecilia [SE/SE]; Lotsgatan
10, S-414 58 Göteborg (SE).**(74) Agent: **JÖNSSON, Christer; Eka Nobel AB, Patent Dept., P.O.
Box 11556, S-100 61 Stockholm (SE).**(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE,
KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN,
MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT,
BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,
PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD,
SZ, UG).**2 1 9 9 1 3 3****Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*(54) Title: **BLEACHING AGENT**

(57) Abstract

The invention relates to particles comprising a peroxy compound with capability of releasing hydrogen peroxide or peroxy acids in aqueous solutions and up to 15 % by weight of a dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof. The invention also relates to use of the particles and a composition comprising such particles.

BLEACHING AGENT

The present invention relates to particles comprising a peroxy compound and a dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof. The invention also relates to
5 use of the particles and a composition containing such particles.

Powder detergent compositions often contain peroxy compounds as bleaching agents which release hydrogen peroxide or peroxy acids in aqueous solutions. However, many peroxy
10 compounds are not storage stable. Their decomposition is supposed to be catalysed by metal cations moving comparatively freely through the water normally being present in detergents, but is also facilitated by the alkaline pH (normally from about 8 to about 12) and by other components commonly present
15 in detergents, for example zeolites or bleaching activators such as TAED (tetraacetylene diamine), TAGU (tetraacetylglucoluril) or PAG (penta acetyl glucose).

In order to make environmental friendly detergents, it is desirable to use alkali metal carbonate peroxy hydrate,
20 commonly known as percarbonate, as a bleaching agent. However, the activity of percarbonate in a detergent decreases rapidly through decomposition if the detergent is stored at normal room temperature and humidity.

Many attempts to stabilize percarbonate have been done,
25 for example by mixing or coating with stabilizing substances such as sulfates, carbonates, borates, silicates or organic substances. Such stabilizing methods are described in the patent literature, for example in GB 1466799, GB 1538893, GB 1575792, EP 459625, US 3975280 and EP 573731.

30 US 4075116 describes preparation of percarbonate from sodium carbonate, hydrogen peroxide and different additives.

CH patent 659082 describes mixing bleaching agents with high amounts of inorganic carbonates and water soluble acids and thereby increasing the dissolution rate.

35 Japanese laid open patent application (Kokai) Sho 58-74508 describes a mixing percarbonate with silicate and phosphate, borate, citrate, tartrate or succinate of sodium and thereby increasing the dissolution rate.

US patent 5340496 describes stabilization of percarbon-

ate with certain combinations of inorganic salts and carboxylic acids, particularly long chained acids such as oleic acid.

The present invention seeks to provide particles comprising a peroxy compound, particularly alkali metal percarbonate, with improved storage stability, particularly in detergent compositions. The invention also seeks to provide particles comprising a peroxy compound only containing environmental friendly constituents.

In accordance with the invention there is provided particles comprising a peroxy compound with capability of releasing hydrogen peroxide or peroxy acids in aqueous solutions, the particles further comprising up to 15% by weight, preferably up to about 10% by weight, most preferably up to about 5% by weight of a preferably water soluble dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof.

The particles preferably comprise more than about 0.01% by weight, most preferably more than about 0.05% by weight of such a dicarboxylic acid or a salt thereof. The amount of the peroxy compound in the particles is suitably from about 10 to about 99% by weight, preferably more than 50% by weight and most preferably up to about 95% by weight. The dicarboxylic acid in the particles may be mixed with the peroxy compound and/or be included in a coating.

2a


It is assumed that the dicarboxylic acid acts as a chelating agent, but it has surprisingly been found that the dicarboxylic acids according to the invention gives superior storage stability compared to conventional chelating agents such as EDTA. Too high a concentration of dicarboxylic acids or salts thereof will however, decrease the thermal stability, particularly at high concentrations of the peroxy compounds.

It is preferred to use an alkali metal or an alkaline earth metal salt of the dicarboxylic acid. Alkali metals are preferably selected from sodium, potassium or mixtures thereof, and alkaline earth metals are preferably selected from calcium, magnesium or mixtures thereof. Sodium salts are particularly preferred. In the following description of the invention, the term dicarboxylic acid also refers to salts thereof.

Suitably the dicarboxylic acid comprises from 5 to 8

carbon atoms, preferably 5 to 7 carbon atoms. Preferably, the carboxylic acid groups are located at the ends of the carbon chain. Further, the dicarboxylic acid preferably does not comprise any hydroxyl groups and most preferably it does not
5 comprise any functional groups except the carboxylic acid groups.

The most preferred dicarboxylic acids satisfy the formula:



10 in which R is alkylene, which is preferred, or alkenylene, having from 3 to 8 carbon atoms, preferably from 3 to 5 carbon atoms. Preferably, R is a straight chain without branches.

Examples of useful dicarboxylic acids are glutaric acid adipic acid, pimelic acid, suberic acid, azelaic acid, sebacic
15 acid and mixtures thereof, of which glutaric acid, adipic acid and pimelic acid are particularly preferred. Glutaric acid is the most preferred acid.


It is to be understood that the particles may include two or more different dicarboxylic acids as described herein.
20 The particles may also include other environmentally acceptable stabilizers or inert substances.

The particles preferably contain silicate which further improves the stability and also enhances the mechanical strength. Silicate may be mixed with the peroxy compound
25 and/or be included in a coating. Preferably at least some of the silicate is mixed with the peroxy compound. The silicate is suitably an alkali metal silicate, preferably of sodium, potassium or mixtures thereof, most preferably of sodium. The molar ratio $\text{SiO}_2:\text{M}_2\text{O}$, M being an alkali metal, is preferably
30 from about 1 to about 3, most preferably from about 1 to about 2.5. The particles suitably contains from about 0.1 to about 20% by weight, preferably from about 0.2 to about 15% by weight, most preferably from about 0.5 to about 10% by weight of silicate expressed as sodium silicate.

35 It has been found that the stability is further improved if the particles contain a water soluble magnesium compound, preferably magnesium sulfate. The magnesium compound may be mixed with the peroxy compound and/or be included in a coating. Preferably at least some of the magnesium is mixed

with the peroxy compound, most preferably in combination with alkali metal silicate. The particles preferably contains from about 0.01 to about 5% by weight, most preferably from about 0.1 to about 3% by weight of magnesium compounds expressed as magnesium sulfate.

5 According to an embodiment of the invention, it has been found that a synergistic effect appears if the particles also comprise a chelating agent selected from hydroxy carboxylic acids satisfying the formula:



10 in which R' is CH₂OH or COOH, n is 2-6, and m is 0-n; or salts thereof.

It is preferred to use an alkali metal or an alkaline earth metal salt of the hydroxy carboxylic acid. Alkali metals are preferably selected from sodium, potassium or mixtures thereof, and alkaline earth metals are preferably selected from calcium, magnesium or mixtures thereof. Sodium salts are particularly preferred. In the following description of the invention, the term hydroxy carboxylic acid also refers to salts thereof.

20 The particles preferably contain from about 0.01 to about 15% by weight, most preferably from about 0.01 to about 10% by weight, particularly preferably from about 0.05 to about 5% by weight of hydroxy carboxylic acids of the above formula. The weight ratio dicarboxylic acid to hydroxy carboxylic acid is preferably from about 0.05:1 to about 4:1, most preferably from about 0.1:1 to about 1:1. The hydroxy carboxylic acid may be mixed with the peroxy compound and/or be included in a coating.

30 Preferably R' in the above formula is CH₂OH. It is also preferred that n is 4 or 5. Preferably, m is equal to n or n-2, the carbon chain thus being saturated or containing one double bond. A particularly preferred hydroxy carboxylic acid is glucoheptonic acid (R' is CH₂OH, n and m are 5) or alkali metals salts thereof. Other preferred salts of hydroxy carboxylic acids are alkali metal gluconate (R' is CH₂OH, n and m are 4) and alkali metal ascorbate (R' is CH₂OH, n is 4 and m is 2). Other useful hydroxy carboxylic acids include, for example, trihydroxy glutaric acid, tartaric acid or saccharic acid. It

is to be understood that the particles may include two or more different hydroxy carboxylic acids of the above formula.

Particles according to the invention have high stability even if uncoated, but the highest stability is normally achieved if the particles are coated. Such a coating may for example contain one or more dicarboxylic acid as described herein, hydroxy carboxylic acids as described herein, alkali metal silicate, water soluble magnesium compounds such as magnesium sulfate, alkali metal salts of carbonate, bicarbonate or sulfate, or environmental acceptable organic chelating agents, the different components being applied in one or several layers.

In one embodiment of the invention, a dicarboxylic acid as described herein is only present as mixed with the peroxy compound while a hydroxy carboxylic acid as described herein is only present in a coating, or vice versa.

In another embodiment both a dicarboxylic acid as described herein and a hydroxy carboxylic acid as described herein are mixed with the peroxy compound and/or are included in a coating.

Suitably, the particles have an average diameter from about 50 to about 3000 μm , preferably from about 100 to about 1600 μm . The preferred density is from about 600 to about 1500 g/l, particularly from about 800 to about 1100 g/l. It has been found that a high density as well as a high average particle size improves the storage stability.

The invention is particularly advantageous if the peroxy compound is alkali metal percarbonate, but also other peroxy compounds can be stabilized, for example alkali metal salts of perborates, peroxyulfates, peroxyphosphates or peroxysilicates, peroxy-carboxylic acids or peroxy-carboxylic acid releasing compounds such as di-acylated di-peroxy carboxylic acids (see WO 91/17143).

Particles according to the invention can be prepared by conventional methods. Ingredients to be mixed with the peroxy compound are preferably added in a granulation step, but may also be included directly when, for example, alkali metal percarbonate is produced from alkali metal carbonate and hydrogen peroxide. If alkali metal silicate is to be included,

it is preferably added in the form of an aqueous solution, and the dicarboxylic acid or a salt thereof and optionally a hydroxy carboxylic acid or a salt thereof is then preferably mixed therein before addition to the peroxy compound. Granulation may be performed by conventional methods well known among those skilled in the art, such as compacting, extruding, agglomeration in drum or disk, fluid-bed granulation, prilling or in different kinds of mixers. An optional coating step may be performed by spraying the particles with preferably aqueous solutions of the components to be applied, for example in a drum or a fluid bed.

The invention further concerns use of the described particles comprising a peroxy compound and a dicarboxylic acid as a bleaching agent, preferably in connection with washing of textiles or dishwashing. The washing water may be supplied with particles according to the invention being included in a detergent composition or being in the form of a separate bleaching powder. Preferably, washing water is supplied with particles in an amount so to obtain from about 0.001 to about 1 gram active oxygen per litre, which for example corresponds to from about 0.01 to about 6 grams sodium percarbonate.

Finally, the invention concerns a composition containing an inert filler, and/or one or several substances active at washing, which composition further contains particles comprising a peroxy compound and a dicarboxylic acid according to the invention, for example in an amount from about 1 up to about almost 100% by weight. The composition may comprise a mixture of particles containing different kinds of peroxy compounds. The substances active at washing may include detergent builders, surfactants, alkali generating substances, bleaching activators, enzymes, or any other substances commonly used in detergents. The detergent builder may for example be selected from phosphates such as tripolyphosphates, pyrophosphates or orthophosphates of alkali metals or ammonium, zeolites such as Zeolite A (e.g. Zeolite 4A), Zeolite B, Zeolite P, Zeolite X, or Zeolite HS, Zeolite MAP, silicates such crystalline layered disilicates (e.g. of the formula $\text{NaMSI}_{x+1} \cdot y\text{H}_2\text{O}$ wherein M is sodium or hydrogen, x is a number from 1.9 to 4 and y is a number from 0 to 20), amorphous disilicates (e.g. BritesilTM),

polycarboxylates, citrates, carbonates, bicarbonates, sesqui-carbonates, sulfates, borates or mixtures thereof. The surfac-tants are preferably selected from anionic surfactants, nonionic surfactants, soaps or mixtures thereof. Anionic
5 surfactants may for example be selected from linear alkyl-benzene sulfonates, secondary alkane sulfonates, alpha-sulfonated methyl fatty acid esters, alkylsulfates, alcohol-ethoxysulfates, alpha-olefine sulfonates, alkali metal sarcosinates or alkyl ester sulfonates. Nonionic surfactants
10 may for example be selected from alkoxyated compounds such as fatty alcohols, alkylphenols and alkylamines or from alkyl polyglucosides or polyhydroxy fatty acid amides. Soaps may for example be selected from sodium or potassium salts of tallow. Also cationic surfactants such as quaternary ammonium com-
15 pounds or imide azolinium salts, as well as amphoteric surfactants might be used. Alkali generating substances may for example be selected from carbonates, silicates, phosphates or mixtures thereof. The bleaching activators may for example be selected from N- or O-acylated compounds such as TAED,
20 TAGU, SNOBS (sodium nonoyl benzene sulfonate), PAG (penta acetyl glucose) or diacylated diperoxy carboxylic acids (see WO 91/17143). The filler may include any inert substance such as sodium sulfate. The composition may constitute a complete detergent or a bleaching powder to be added separately at
25 washing. The enzymes may for example be selected from amylases, neutral and alkaline proteases, lipases, esterases or cellulases.

A complete detergent for washing of textiles suitably contains from about 1 to about 40% by weight, preferably from
30 about 10 to about 30% by weight of particles according to the invention. Further, the detergent suitably contains a deter-gent builder, for example from about 5 to about 50% by weight, surfactants, for example from about 5 to about 35% by weight and an alkali generating substance, for example from about 5
35 to about 20% by weight. Preferably, the detergent contains from about 5 to about 20% by weight of anionic surfactants, from about 2 to about 15% by weight of nonionic surfactants, and from about 0.1 to about 5% by weight of soaps. The detergent may also contain bleaching activators, for example

from about 1 to about 10% by weight, enzymes, for example from about 0.5 to about 2.5% by weight, and fillers such as sodium sulfate, for example from about 5 to about 50% by weight. It is also possible to include chelating agents such as phosphonates, EDTA, NTA (nitrilo triacetic acid), iminodiacetic acid derivatives or EDDS (ethylenediamine-N,N-disuccinic acid, for example in an amount from about 0.1 to about 1% by weight. Additionally, the detergent may contain conventional components such as water glass, carboxy methyl cellulose, dispersants such as homo or copolymers of polycarboxylic acids, foam regulators, antioxidants, perfumes, colouring agents, optical brighteners and water (normally from about 3 to about 15% by weight). The detergent can be prepared with conventional methods such as dry mixing, agglomeration or spray drying. If the preparation involves spray drying, any heat sensible component such as particles comprising peroxy compounds, enzymes and perfumes should be added to the dried material.

A separate bleaching powder may comprise up to almost 100% by weight of particles containing peroxy compounds according to the invention, but preferably the content is from about 5 to about 90% by weight. The bleaching powder may comprise only one peroxy compound or a mixture of particles containing different kinds of peroxy compounds. It is advantageous if a hydrogen peroxide generating substance such as percarbonate, for example from about 10 to about 75% by weight, is used in combination with a bleaching activator such as TAED or TAGU, for example from about 2 to about 25% by weight. Also other bleaching activators such as a diacylated dipercarboxylic acid may be used, for example in an amount from about 2 to about 25% by weight. A bleaching powder may also contain detergent builders, for example from about 5 to about 90% by weight, surfactants, for example up to about 10% by weight, enzymes, for example up to about 2% by weight, or fillers, for example from about 5 to about 90% by weight. A preferred bleaching powder consists essentially of from about 30 to about 75% by weight of particles containing percarbonate, from about 10 to about 25% by weight of a bleaching activator, the balance preferably substantially being deter-

gent builders, fillers, surfactants, water or mixtures thereof.

A detergent for dishwashing may be in the form of a low alkaline detergent (pH of washing water intended to be about 10-11) suitably containing from about 2 to about 15% by weight of a bleaching agent comprising particles according to the invention, from about 5 to about 50% by weight of alkali metal disilicate, from 0 to about 40% by weight of alkali metal carbonate, from about 15 to about 50% by weight of builders such as sodium citrate and polycarboxylates or sodium tri polyphosphate (STPP), from about 0.5 to about 5% by weight of low foaming nonionic surfactants, from about 0.5 to about 5% by weight of enzymes and from about 1 to about 6% by weight of bleaching activators such as TAED. A dishwasher detergent may also be high alkaline (pH of washing water intended to be about 11-12) having a similar composition as the low alkaline detergent, the disilicate however being replaced with from about 20 to about 80% by weight of alkali metal metasilicate and the builder preferably substantially consisting of STPP.

The present invention makes it possible to prepare stable bleaching agents comprising peroxy compounds, particularly percarbonate, which may be used in detergents including zeolites such as zeolite 4A. The invention also makes it possible to decrease or exclude the use of for example EDTA or phosphonates which are less suitable from an environmental point of view.

The invention will now be described through the following examples which, however, not are intended to limit the scope of the invention. If not otherwise stated, all contents an percentages refer to content and percent by weight based on the entire particle.

EXAMPLE 1: Sodium percarbonate particles with a diameter of about 315-1250 μm and a density of about 900 g/l were prepared by granulating fine percarbonate particles in a rotating Eirich drum together with different additives supplied as aqueous solutions and drying the granules in a fluidized bed. In order to test the storage stability of the percarbonate, the prepared particles as well as reference particles of pure sodium percarbonate were included in a

standard detergent IEC-Z (Henkel) consisting of 9.7% linear alkyl benzenesulfonate ($C_{average} 11.5$), 5.2% ethoxylated C_{12-18} alcohol (EO₇), 3.6% sodium-soap, 32.5% zeolite 4A, 11.8% sodium carbonate, 5.2% sodium salt of copolymerisate of acrylic- and maleic acid (Sokolan CP₅), 3.4% sodium waterglass (ratio 3.3), 1.3% carboxy methyl cellulose, 0.3% EDTA, 0.3% optical brightener (stilbene type), 7.4% sodium sulfate, 12.2% water, 6.5% foam inhibitor (SIK), 0.5% enzyme prills protease (activity 300.00). Samples were prepared from 64 grams IEC-Z, 12 grams percarbonate particles and 4 grams TAED. A Quick Stability Index (QSI) was determined for each sample by measuring the amount of oxygen formed during 24 hours storage at 40°C and dividing the amount oxygen liberated by the relative content (%) of active oxygen in the original percarbonate. Thus, a low QSI-value means high stability. The results appear in the table below, in which the contents of the different constituents refer to % by weight based on the entire particle. The following abbreviations are used: SPC = sodium percarbonate; GA = sodium salt of glutaric acid; GH = sodium gluconate; AA = sodium salt of adipic acid; PA = sodium salt of pimelic acid; SDS = sodium disilicate; AO = active oxygen; QSI = quick stability index. Sokolan is a Trade-mark.

SPC	GA	AA	PA	AO	QSI
100%	-	-	-	14.5%	5.66
98%	2%	-	-	14.1%	2.64
98%	-	-2%	-	14.3	2.09
98%	-	-	2%	14.1%	2.38

EXAMPLE 2: Sodium percarbonate were prepared and tested as in Example 1. The results appear in the table below:

SPC	GA	GH	SDS	MgSO ₄	AO	QSI
97.2%	1.5%	-	1.2%	0.1%	14.0%	2.28
97.2%	-	1.5%	1.2%	0.1%	14.0%	2.59
97.2%	0.5%	1.0%	1.2%	0.1%	14.1%	1.63
97.2%	1.0%	0.5%	1.2%	0.1%	14.1%	2.14

EXAMPLE 3: Sodium percarbonate particles with a diameter of about 315-1250 μ m and a density of about 900 g/l were

prepared by granulation from fine percarbonate particles together with sodium salt of glutaric acid, sodium disilicate and magnesium sulfate supplied as aqueous solutions. The granulation was performed by extruding a mixture of the ingredients, crushing the string formed, rounding the particles obtained on a rotating disk and drying and sieving the rounded particles. For comparative purposes one sample was prepared without glutaric acid and one sample was prepared with EDTA instead of glutaric acid. The stability of the particles was tested in a standard IEC-Z detergent in the same way as in Example 1 and the results appear in the table below.

SPC	GA	EDTA	SDS	MgSO ₄	AO	QSI
99.2%	-	-	0.67%	0.12%	14.3%	3.4
99.0%	-	0.2%	0.67%	0.12%	14.4%	3.1
99.0%	0.2%	-	0.67%	0.12%	14.4%	2.6
98.8%	0.5%	-	0.63%	0.12%	14.2%	2.1

EXAMPLE 4: Sodium percarbonate particles with a diameter of 315-1250 μ m were coated with sodium salts of glutaric acid and/or glucoheptonic acid applied as aqueous solutions by spraying in a fluidized bed. For comparative purposes particles were also coated with benzoic acid and caproic acid. The stability of the particles was tested in a standard IEC-Z detergent in the same way as in Example 1 and the results appear in the table below.

SPC	GA	GH	AO	QSI
100%	-	-	14.5%	5.66
95%	5%	-	13.6%	0.65
95%	-	5%	13.9%	0.78
95%	4%	1%	13.6%	0.40
95%	3%	2%	13.4%	0.31
95%	2%	3%	13.3%	0.25
95%	1%	4%	13.6%	0.31
95%	5% benzoic acid		13.8%	2.31
95%	5% caproic acid		13.9%	2.27

EXAMPLE 5: Sodium percarbonate particles with a diameter

of 315-1250 μm were coated by spraying in a fluidized bed with a first aqueous solution of sodium silicate ($\text{SiO}_2\text{:Na}_2\text{O}$ ratio 1), sodium glucoheptonate and/or sodium salt of glutaric acid, and a second solution of magnesium sulfate and optionally sodium sulfate or sodium bicarbonate. In one experiment a third solution of sodium carbonate was applied. The stability of the particles was tested in a IEC-Z detergent in the same way as in Example 1 and the results appear in the table below:


	Exp. 1	Exp. 2	Exp. 3	Exp. 4
sodium percarbonate	85.1%	85.1%	85.1%	85.1%
glucoheptonic acid	-	0.75%	0.75%	0.75%
glutaric acid	1.5%	0.75%	0.75%	0.75%
sodium disilicate	2%	2%	2%	2%
MgSO_4	1.4%	1.4%	1.4%	1.4%
Na_2SO_4	10%	10%		-
NaHCO_3	-	-	10%	-
Na_2CO_3	-	-	-	10%
active oxygen	12.8%	12.9%	12.8%	12.8%
QSI	0.28	0.22	0	0.16

EXAMPLE 6: Sodium percarbonate particles prepared by granulation as in Example 3 contained 98.8% sodium percarbonate, 0.12% MgSO_4 , 0.63% sodium disilicate, 0.5% glutaric acid (as sodium salt), and had a content of active oxygen of 14.2%. The particles were coated as in Examples 5 with 1.4% MgSO_4 , 2% sodium disilicate, 10% Na_2CO_3 , and 1% glutaric acid (as sodium salt). The coated particles contained 12.6% of active oxygen. The stability of the particles was tested in a IEC-Z detergent in the same way as in Example 1 and it was found that QSI for the uncoated particles was 2.20 and QSI for the coated particles was 0.04.

CLAIMS

1. Particles comprising a peroxy compound with capability of releasing hydrogen peroxide or peroxy acids in aqueous solutions, wherein the particles comprise up to 15% by weight of a dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof.
2. Particles as claimed in claim 1, wherein said particles comprise up to 10% by weight of said dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof.
3. Particles as claimed in claim 1 or 2, wherein the particles comprise an alkali metal or alkaline earth metal salt of the dicarboxylic acid having from 5 to 10 carbon atoms.
4. Particles as claimed in claim 1, 2 or 3, wherein the dicarboxylic acid having from 5 to 10 carbon atoms or a salt thereof is mixed with the peroxy compound.
5. Particles as claimed in claim 4, wherein the peroxy compound and the dicarboxylic acid are granulated together.
6. Particles as claimed in claim 1, 2, 3, 4 or 5, wherein the dicarboxylic acid having 5 to 10 carbon atoms or a salt thereof is included in a coating.

7. Particles as claimed in any one of claims 1 to 6, wherein the dicarboxylic acid satisfies the formula:



in which R is alkylene or alkenylene having from 3 to 8 carbon atoms.

8. Particles as claimed in claim 7, wherein R is a straight chain without branched.

9. Particles as claimed in claim 8, wherein the dicarboxylic acid is selected from glutaric acid, adipic acid, pimelic acid, suberic acid, azelaic acid, sebacic acid or a mixture thereof.

10. Particles as claimed in claim 9, wherein the dicarboxylic acid is glutaric acid.

11. Particles as claimed in any one of claims 1 to 10, wherein the particles comprise silicate.

12. Particles as claimed in claim 11, wherein the particles comprise alkali metal silicate.

13. Particles as claimed in claim 11 or 12, wherein silicate is mixed with the peroxy compound.

14. Particles as claimed in claim 11, 12 or 13, wherein silicate is included in a coating.

15. Particles as claimed in any one of claims 1 to 14, wherein the particles comprise a water soluble magnesium compound.

16. Particles as claimed in any one of claims 1 to 15, wherein the particles comprise a hydroxy carboxylic acid satisfying the formula:



in which R' is CH₂OH or COOH, n is 2-6 and m is 0-n; or a salt thereof.

17. Particles as claimed in claim 16, wherein R' is CH₂OH, n is 4 or 5 and m is equal to n or n-2.

18. Particles as claimed in claim 17, wherein the hydroxy carboxylic acid is glucoheptonic acid.

19. Particles as claimed in any one of claims 1 to 15, wherein a hydroxy carboxylic acid or a salt as defined in claim 16, 17 or 18, is mixed with the peroxy compound.

20. Particles as claimed in any one of claims 16 to 19, wherein the hydroxy carboxylic acid or a salt as defined in claim 16, 17 or 18, is included in a coating.

21. Particles as claimed in any one of claims 1 to 20, wherein said dicarboxylic acid having from 5 to 10 carbon atoms and a hydroxy carboxylic acid as defined in claim 16, 17 or 18, are both mixed with the peroxy compound.

22. Particles as claimed in any one of claims 1 to 21, wherein a dicarboxylic acid having from 5 to 10 carbon atoms and a hydroxy carboxylic acid as defined in claim 16, 17 or 18, are both included in a coating.

23. Particles as claimed in any one of claims 1 to 22, wherein the peroxy compound is alkali metal percarbonate.

24. Particles as claimed in any one of claims 1 to 23, wherein the bulk density is from about 800 to about 1500 g/l.

25. Particles as claimed in claim 24, wherein the bulk density is from about 800 to about 1100 g/l.

26. Particles as claimed in any one of claims 1 to 25, wherein said particles comprise from 0.1 to 15%, by weight of said dicarboxylic acid or salt.

27. Particles as claimed in claim 26, comprising from 0.1 to 10%, by weight of said dicarboxylic acid or salt.

28. Particles as claimed in any one of claims 1 to 27, wherein said dicarboxylic acid is water soluble.

29. Use of particles according to any one of claims 1 to 28, as a bleaching agent in detergents.

30. A composition comprising an inert filler, at least one substance active at washing, which composition further contains particles according to any one of claims 1 to 28.

31. A detergent composition comprising particles according to any one of claims 1 to 28.

32. A detergent composition according to claim 31, comprising detergent ingredients.

33. A detergent composition as claimed in claim 31 or 32, further comprising zeolite as a builder.