



(12)

# Oversættelse af europæisk patent

Patent- og  
Varemærkestyrelsen

(51) Int.Cl.: **A 01 K 67/027 (2006.01)** **C 07 K 14/535 (2006.01)** **C 07 K 14/54 (2006.01)**

(45) Oversættelsen bekendtgjort den: **2019-10-07**

(80) Dato for Den Europæiske Patentmyndigheds  
bekendtgørelse om meddelelse af patentet: **2019-07-03**

(86) Europæisk ansøgning nr.: **17184813.8**

(86) Europæisk indleveringsdag: **2013-11-05**

(87) Den europæiske ansøgnings publiceringsdag: **2018-04-18**

(30) Prioritet: **2012-11-05 US 201261722437 P**

(62) Stamansøgningsnr: **13811637.1**

(84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(73) Patenthaver: **Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, NY 10591-6707, USA**  
**Yale University, Two Whitney Avenue, New Haven, CT 06510, USA**  
**Institute for Research in Biomedicine (IRB), Via Vincenzo Vela 6, 6501 Bellinzona, Schweiz**

(72) Opfinder: **Flavell, Richard, 283 Moose Hill Road, Guilford, CT 06437, USA**  
**STROWIG, Till, Leipziger Str. 232, 38124 Braunschweig, Tyskland**  
**MANZ, Markus G., Stettbachstrasse 9, 8702 Zollikon, Schweiz**  
**BORSOTTI, Chiara, 10 Kraft Avenue, Bronxville, NY 10708, USA**  
**Dhodapkar, Madhav, 105 McKinley Avenue, New Haven, CT 06515, USA**  
**Murphy, Andrew J., 10 Newton Court, Croton-on-hudson, NY 10520, USA**  
**Stevens, Sean, 3954 Camino Calma, San Diego, CA 92122, USA**  
**YANCOPOULOS, George D., 1519 Baptist Church Road, Yorktown Heights, NY 10598, USA**

(74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**

(54) Benævnelse: **Immundeficient genmodificeret gnaver og fremgangsmåder til anvendelse deraf**

(56) Fremdragne publikationer:  
**WO-A1-2013/063556**  
**WO-A2-2011/044050**  
**WO-A2-2012/040207**  
**WO-A2-2012/112544**  
**WO-A2-2014/039782**  
**R. W. J. GROEN ET AL: "Reconstructing the human hematopoietic niche in immunodeficient mice: opportunities for studying primary multiple myeloma", BLOOD, vol. 120, no. 3, 31 May 2012 (2012-05-31), pages e9-e16, XP055113167, ISSN: 0006-4971, DOI: 10.1182/blood-2012-03-414920**  
**PIERFRANCESCO TASSONE ET AL: "A clinically relevant SCID-hu in vivo model of human multiple myeloma", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 106, no. 2, 15 July 2005 (2005-07-15) , pages 713-716, XP002633148, ISSN: 0006-4971, DOI: 10.1182/BLOOD-2005-01-0373 [retrieved on 2005-04-07]**  
**T. WILLINGER ET AL: "Human IL-3/GM-CSF knock-in mice support human alveolar macrophage development**

and human immune responses in the lung", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 108, no. 6, 24 January 2011 (2011-01-24), pages 2390-2395, XP055058608, ISSN: 0027-8424, DOI: 10.1073/pnas.1019682108

A. RONGVAUX ET AL: "Human thrombopoietin knockin mice efficiently support human hematopoiesis in vivo", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 108, no. 6, 8 February 2011 (2011-02-08), pages 2378-2383, XP055112601, ISSN: 0027-8424, DOI: 10.1073/pnas.1019524108

TILL STROWIG ET AL: "Transgenic expression of human signal regulatory protein alpha in Rag2(-/-)gamma(-/-) mice improves engraftment of human hematopoietic cells in humanized mice + Supplementary Online Data", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, NATIONAL ACADEMY OF SCIENCES, US, vol. 108, no. 32 9 August 2011 (2011-08-09), pages 13218-13223, XP002681726, ISSN: 0027-8424, DOI: 10.1073/PNAS.1109769108 Retrieved from the Internet: URL:<http://www.pnas.org/content/108/32/13218> [retrieved on 2011-07-25]

C. RATHINAM ET AL: "Efficient differentiation and function of human macrophages in humanized CSF-1 mice", BLOOD, vol. 118, no. 11, 15 September 2011 (2011-09-15), pages 3119-3128, XP055112602, ISSN: 0006-4971, DOI: 10.1182/blood-2010-12-326926

KURUVILLA ET AL: "Dengue virus infection and immune response in humanized RAG2<-/->gammac<-/-> (RAG-hu) mice", VIROLOGY, ELSEVIER, AMSTERDAM, NL, vol. 369, no. 1, 3 November 2007 (2007-11-03), pages 143-152, XP022327268, ISSN: 0042-6822, DOI: 10.1016/J.VIROL.2007.06.005

OTOYA UEDA ET AL: "Novel genetically-humanized mouse model established to evaluate efficacy of therapeutic agents to human interleukin-6 receptor", SCIENTIFIC REPORTS, NATURE PUBLISHING GROUP, GB, vol. 3 1 January 2013 (2013-01-01), page 1196, XP002692003, ISSN: 2045-2322, DOI: 10.1038/SREP01196 Retrieved from the Internet: URL:<http://www.nature.com/srep/2013/130201/srep01196/full/srep01196.html> [retrieved on 2013-02-01]

LEONARD D. SHULTZ ET AL: "Humanized mice for immune system investigation: progress, promise and challenges", NATURE REVIEWS IMMUNOLOGY, vol. 12, no. 11, 1 November 2012 (2012-11-01), pages 786-798, XP055064740, ISSN: 1474-1733, DOI: 10.1038/nri3311

GARCIA SYLVIE ET AL: "Humanized mice: Current states and perspectives", IMMUNOLOGY LETTERS, ELSEVIER BV, NL, vol. 146, no. 1-2 30 August 2012 (2012-08-30), pages 1-7, XP002681730, ISSN: 0165-2478, DOI: 10.1016/J.IMLET.2012.03.009 Retrieved from the Internet: URL:<http://www.sciencedirect.com/science/article/pii/S0165247812001034> [retrieved on 2012-04-09]

Hua Yu ET AL: "A novel humanized mouse model with significant improvement of class-switched, antigen-specific antibody production", Blood, vol. 129, no. 8 11 January 2017 (2017-01-11), pages 959-969, XP055454421, DOI: 10.1182/blood-2016-04- Retrieved from the Internet: URL:<http://www.bloodjournal.org/content/129/8/959.full.pdf> [retrieved on 2018-02-26]

# DESCRIPTION

## CROSS REFERENCE TO RELATED APPLICATIONS

**[0001]** Pursuant to 35 U.S.C. § 119 (e), this application claims priority to the filing date of the United States Provisional Patent Application Serial No. 61/722,437 filed November 5, 2012.

## BACKGROUND OF THE INVENTION

**[0002]** The aim of biomedical research is to gain a better understanding of human physiology and to use this knowledge to prevent, treat or cure human diseases. Due to practical and ethical barriers to the experimentation on human subjects, many studies are conducted on small animal models, such as the mouse. Animal models of these human diseases are therefore needed.

**[0003]** For example, in the United States, around 20,000 patients are annually newly diagnosed with multiple myeloma (MM), a mostly incurable malignancy of antibody-secreting terminally differentiated B cells (Hideshima et al., 2007, *Nat Rev Cancer.* 7:585-98; Kuehl and Bergsagel, 2002, *Nat Rev Cancer.* 2:175-87). MM is characterized by the infiltration of malignant plasma cells in the bone marrow (BM) and clinical manifestations include bone disease, hypercalcemia, cytopenia, renal dysfunction, and peripheral neuropathy (Hideshima et al., 2007, *Nat Rev Cancer.* 7:585-98; Kuehl and Bergsagel, 2002, *Nat Rev Cancer.* 2:175-87). In most cases, MM is preceded by a premalignant condition called monoclonal gammopathy of undetermined significance (MGUS) that affects around 3% of persons older than 50 years (Landgren et al., 2009, *Blood* 113:5412-7). Complex heterogeneous genetic abnormalities characterize MM cells including changes in the karyotype as well as IgH translocations (Kuehl and Bergsagel, 2002, *Nat Rev Cancer.* 2:175-87; Zhan et al., 2006, *Blood* 108:2020-8). Plasma cell clones that are amplified in MGUS are thought to have genetic and phenotypic profiles similar to myelomatous plasma cells (Chng et al., 2005, *Blood* 106:2156-61; Fonseca et al., 2002, *Blood* 100:1417-24; Kaufmann et al., 2004, *Leukemia.* 18:1879-82). While mutations in cyclin D genes have been suggested to drive development of MM, the potential contributions of other factors have not been conclusively demonstrated (Bergsagel et al., 2005, *Blood* 106:296-303). Nonetheless, heritable genetic alterations are not the sole determinants of the behavior of MM cells. Instead, resistance towards drugs and aberrant biological responses towards cytokines are strongly influenced by interactions with the microenvironment offering an opportunity to develop novel therapeutics.

**[0004]** Like many other tumors, MM is characterized by heterogeneous cell populations strongly interacting with non-malignant stroma cells that create a supportive environment (De Raeve and Vanderkerken, 2005, *Histol Histopathol.* 20:1227-50; Dhodapkar, 2009, *Am J Hematol.* 84:395-6). The BM microenvironment for MM cells consists of a diverse extracellular

matrix (ECM) and of cellular components of both hematopoietic and non-hematopoietic origin. While the BM provides a protected environment for normal hematopoiesis, the interaction of MM cells with ECM proteins and accessory cells plays a crucial role in MM pathogenesis (De Raeve and Vanderkerken, 2005, *Histol Histopathol.* 20:1227-50; Dhodapkar, 2009, *Am J Hematol.* 84:395-6; Hideshima et al., 2007, *Nat Rev Cancer.* 7:585-98). Stroma cells, myeloid cells, osteoclasts, and osteoblasts produce growth factors such as interleukin 6 (IL-6), B-cell activating factor (BAFF), fibroblast growth factor, and stroma cell-derived factor 1a that activate signal pathways mediating migration, survival, and growth of MM cells. In particular, IL-6 produced by stroma cells, osteoclasts, and myeloid cells seems to be a crucial factor in the early stages and for pathogenesis of MM (De Raeve and Vanderkerken, 2005, *Histol Histopathol.* 20:1227-50). Similarly, upon interaction with MM cells, osteoclasts and dendritic cells produce BAFF and/or a proliferation-inducing ligand (APRIL) providing anti-apoptotic signals that also increase drug resistance (De Raeve and Vanderkerken, 2005, *Histol Histopathol.* 20:1227-50; Kukreja et al., 2006, *J Exp Med.* 203:1859-65).

**[0005]** The major events in cancer pathogenesis - uncontrolled proliferation, survival and spread of the malignant cells - depend on specific combinations of supportive cell types and soluble factors present in microenvironmental niches. Mouse models play an important role in characterizing key aspects of the driving forces of malignant transformation and disease in humans. However, they rarely represent the genetic complexity and clinicopathologic characteristics of human disease. While xenotransplantation of human tumors into immunocompromised mice has been extensively employed, reliable engraftment has typically been feasible only with highly aggressive tumors or cell lines.

**[0006]** The best models currently available to grow human tumor cells are severely immunodeficient mice that lack B cells, T cells, and NK cells. In the case of MM, engraftment of primary myeloma cells into these mice has been unsuccessful, but primary myeloma cells are able to engraft human fetal bone pieces upon co-transplantation into immunocompromised mice (Yaccoby et al., 1998, *Blood* 92:2908-13). In this model MM cells are found in the human bone, but are not detected in the mouse bone or in the periphery demonstrating high residual xenorejection and a need for the human BM microenvironment (Yaccoby et al., 1998, *Blood* 92:2908-13; Yaccoby and Epstein, 1999, *Blood* 94:3576-82). Proving its potential as in vivo model for MM, it was recently demonstrated that NOD/Scid/yc<sup>-/-</sup> mice allow the engraftment of several MM cell lines (Dewan et al., 2004, *Cancer Sci.* 95:564-8; Miyakawa et al., 2004, *Biochem Biophys Res Commun.* 313:258-62). However, even those mouse models with low xenorejection have constricted growth environments by virtue of a large number of factors that do not cross species barriers but are essential to support growth and survival of transformed cells (Manz, 2007). In vivo models that allow us to probe the complex pathogenic interplay between the tumor and its environment will be essential to design new drugs and therapies.

**[0007]** Therefore there is an unmet need to develop humanized non-human animals and methods to reliably grow and study human hematopoietic cells, including primary human hematopoietic tumor cells in mice. The present invention addresses these unmet needs in the art.

**SUMMARY OF THE INVENTION**

**[0008]** Genetically modified rodents are provided that may be used to model human hematopoietic cell development, function, or disease. The genetically modified rodents comprise a nucleic acid encoding human IL-6 operably linked to an *IL-6* promoter and express at least one of human M-CSF, human IL-3, human GM-CSF, human SIRPa or human TPO. The genetically modified rodent is immunodeficient. The rodent is engrafted with healthy or diseased human hematopoietic cells. Disclosed herein are methods for using the subject genetically modified rodents in modeling human hematopoietic cell development, function, and/or disease, as well as reagents and kits thereof that find use in making the subject genetically modified rodents and/or practicing the subject methods.

**[0009]** The genetically modified rodent does not express a functional native IL-6.

**[0010]** In some embodiments, the rodent is a mouse. In some such embodiments, the *IL-6* promoter to which the nucleic acid encoding human IL-6 is operably linked is the mouse *IL-6* promoter, and the human IL-6 gene is operably linked to the mouse *IL-6* promoter at the mouse *IL-6* locus.

**[0011]** The genetically modified rodent further comprises one or more additional nucleic acids selected from a nucleic acid encoding human SIRPa under the control of a *SIRPa* promoter; a nucleic acid encoding human M-CSF operably linked to an *M-CSF* promoter, wherein the rodent expresses human M-CSF; a nucleic acid encoding human IL-3 operably linked to an *IL-3* promoter, wherein the rodent expresses human IL-3; a nucleic acid encoding human GM-CSF operably linked to a *GM-CSF* promoter, wherein the rodent expresses human GM-CSF; and a nucleic acid encoding human TPO operably linked to a *TPO* promoter, wherein the rodent expresses human TPO. In some embodiments, the promoter is the human promoter for the gene. In other embodiments, the promoter is the rodent promoter for the gene. In some embodiments, the genetically modified rodent expresses the rodent's corresponding native protein. In other embodiments, the genetically modified rodent does not express the rodent's corresponding native protein.

**[0012]** The genetically modified rodent is immunodeficient for the rodent immune system. In some such embodiments, the immunodeficient genetically modified rodent does not express a recombination activating gene (RAG). In some such embodiments the immunodeficient genetically modified rodent does not express the IL2 receptor gamma chain (IL2rg, or "yc"). In some such embodiments the immunodeficient genetically modified rodent does not express either a RAG (e.g. RAG1, RAG2) or IL2rg.

**[0013]** The immunodeficient genetically modified rodent is engrafted with human hematopoietic cells to form a genetically modified and engrafted rodent. In one embodiment, the human hematopoietic cells are selected from human umbilical cord blood cells, human fetal

liver cells, and cells of a human hematopoietic cell line. In one embodiment, the human hematopoietic cells are CD34+ progenitor cells. In one embodiment, the human hematopoietic cells are cancer cells. In certain embodiments, the cancer cells are human multiple myeloma cells.

**[0014]** The genetically modified and engrafted rodent may give rise to a human cell selected from a CD34+ cell, a hematopoietic stem cell, a hematopoietic cell, a myeloid precursor cell, a myeloid cell, a dendritic cell, a monocyte, a granulocyte, a neutrophil, a mast cell, a thymocyte, a T cell, a B cell, a plasma cell, a platelet, and a combination thereof. The human cell may be present at 1 month, at 2 months, at 3 months, at 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, or 12 months after engraftment.

**[0015]** The genetically modified and engrafted rodent may give rise to a human hemato-lymphoid system that comprises human hematopoietic stem and progenitor cells, human myeloid progenitor cells, human myeloid cells, human dendritic cells, human monocytes, human granulocytes, human neutrophils, human mast cells, human thymocytes, human T cells, human B cells, human plasma cells, and human platelets. The human hemato-lymphoid system may be present at 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, or 12 months after engraftment.

**[0016]** Disclosed herein are, methods for generating a rodent engrafted with human hematopoietic cells. Disclosed are, methods for generating a rodent model of human immune cell development and function. Disclosed are, methods for generating a rodent model of human B cell development and function.

**[0017]** The methods comprise transplanting a population of human hematopoietic cells to a genetically modified rodent that is immunodeficient and expresses a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter. The animal does not express a functional native IL-6. In some embodiments, the IL-6 promoter is the rodent IL-6 promoter and the human IL-6 gene is operably linked to the rodent IL-6 promoter at the rodent IL-6 locus. In some embodiments, the rodent is a mouse. As such, in some embodiments, the IL-6 promoter to which the nucleic acid encoding human IL-6 is operably linked is the mouse *IL-6* promoter, and the human IL-6 gene is operably linked to the mouse *IL-6* promoter at the mouse *IL-6* locus.

**[0018]** In some embodiments, the transplanted population of hematopoietic cells comprises CD34+ cells. In some embodiments, the transplanted population of hematopoietic cells comprises cancer cells. In some embodiments, the transplanted population of cancer cells comprises multiple myeloma cells. In some embodiments, the transplanting comprises intrafemoral and/or intratibial injection.

**[0019]** The immunodeficient, genetically modified rodent expresses at least one additional human nucleic acid selected from the group consisting of a nucleic acid encoding human SIRPa operably linked to a SIRPa promoter; a nucleic acid encoding human M-CSF operably

linked to a M-CSF promoter; a nucleic acid encoding human IL-3 operably linked to an IL-3 promoter; a nucleic acid encoding human GM-CSF operably linked to a GM-CSF promoter; and a nucleic acid encoding human TPO operably linked to a TPO promoter.

**[0020]** In some aspects of the invention, engrafted, genetically modified rodent expressing a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter are provided, these engrafted, rodents having been prepared according to the methods described herein or as known in the art. In some embodiments, the engrafted, genetically modified rodent is an animal model of human B cell development and differentiation

**[0021]** Disclosed herein are methods that encompass the use of human hematopoietic cell-engrafted, genetically modified rodents of the subject disclosure. These methods include, for example, methods for the in vivo evaluation of the growth and differentiation of hematopoietic and immune cells, methods for the in vivo evaluation of human hematopoiesis, methods for the in vivo evaluation of cancer cells, methods for the in vivo assessment of an immune response, methods for the in vivo evaluation of vaccines and vaccination regimens, methods for the use in testing the effect of agents that modulate cancer cell growth or survival, methods for the in vivo evaluation of a treatment of cancer, and for use in testing the effect of agents that modulate hematopoietic and immune cell function. In certain embodiments, methods are provided for the in vivo production and collection of human antibodies. For example, in some disclosures, methods are provided for screening a candidate agent for the ability to treat a hematopoietic cancer. In some disclosures, the method comprises contacting a genetically modified rodent of the present disclosure that has been engrafted with human hematopoietic cancer cells with a candidate agent, and comparing the viability and/or proliferative rate of human hematopoietic cancer cells in the contacted engrafted, genetically modified rodent to the human hematopoietic cancer cells in a similarly engrafted, genetically modified rodent that is not contacted with candidate agent, wherein a decrease in the viability and/or rate of proliferation of the human hematopoietic cancer cells in the contacted engrafted, rodent indicates that the candidate agent will treat a hematopoietic cancer. These and other methods will be apparent to the ordinarily skilled artisan from the disclosure herein.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

**[0022]** The following detailed description of preferred embodiments of the invention will be better understood when read in conjunction with the appended drawings. It is emphasized that, according to common practice, the various features of the drawings are not to-scale. On the contrary, the dimensions of the various features are arbitrarily expanded or reduced for clarity. For the purpose of illustrating the invention, there are shown in the drawings embodiments which are presently preferred. It should be understood, however, that the invention is not limited to the precise arrangements and instrumentalities of the embodiments shown in the drawings. Included in the drawings are the following figures.

Figure 1 is a set of graphs depicting the results of experiments demonstrating the engraftment

of INA-6 cells in human IL-6 knock-in mice. Soluble IL-6R levels were measured in mice of the indicated genotypes transplanted with  $5 \times 10^6$  INA-6 cells intravenously. N indicates the number of transplanted mice per group.

Figure 2, comprising Figure 2A through Figure 2F, is a set of images depicting the histological analysis of femurs after intravenous engraftment of INA-6 cells.  $\text{Rag2}^{-/-}\text{Il2rg}^{\text{null}}\text{ Il6}^{\text{h/h}}\text{hSIRPa+}$  mice were sacrificed eight weeks after engraftment with  $5 \times 10^6$  INA-6 cells intravenously. Femurs were fixed in 10 % Formalin and decalcified. 10  $\mu\text{M}$  sections were stained with Toluidine blue or directly analyzed for GFP expression using a Leica Confocal microscope.

Figure 3 is a set of images and a graph depicting the analysis of lungs after intravenous engraftment of INA-6 cells.  $\text{Rag2}^{-/-}\text{Il2rg}^{\text{null}}\text{ Il6}^{\text{h/h}}\text{hSIRPa+}$  mice were sacrificed eight weeks after engraftment with  $5 \times 10^6$  intravenously injected INA-6 cells. Lung tissue was fixed in 10% Formalin and 10  $\mu\text{M}$  sections were directly analyzed for GFP expression using a Leica Confocal microscope. Images (left) depict a 10X (top) and 63X (bottom) magnification of sections. The graph (right) depicts human  $\text{Il6}$  gene expression measured in indicated tissues and normalized to murine  $\text{hprt}$  expression.

Figure 4 is a set of graphs depicting the results of experiments demonstrating the engraftment of INA-6 cells in human IL-6 knock-in mice. Soluble IL-6R levels were measured in mice of the indicated genotypes transplanted with  $5 \times 10^5$  INA-6 cells intrafemorally. Each line indicates an individual mouse.

Figure 5 is a set of images depicting the histological analysis of femurs after intrafemoral engraftment of INA-6 cells.  $\text{Rag2}^{-/-}\text{Il2rg}^{\text{null}}\text{ Il6}^{\text{h/h}}\text{hSIRPa+}$  mice were sacrificed four to six weeks after engraftment with  $5 \times 10^5$  intrafemorally injected INA-6 cells. Femurs were fixed in 10 % Formalin and decalcified. 10  $\mu\text{M}$  sections were stained with Toluidine blue.

Figure 6 is a set of images and a graph depicting the results of  $\mu\text{CT}$  analysis of murine femurs after INA-6 transplantation.  $\text{Rag2}^{-/-}\text{Il2rg}^{\text{null}}\text{ Il6}^{\text{h/h}}\text{hSIRPa+}$  and control mice were sacrificed four weeks after engraftment with  $5 \times 10^5$  intrafemorally injected INA-6 cells. Femurs were fixed in 70 % ethanol and analyzed using a murine  $\mu\text{CT}$ . Trabecular bone and tissue volumes were quantified to calculate the ratio between bone to tissue volume. \*:  $p < 0.01$  by student t-test.

Figure 7 is a graph depicting the results of  $\mu\text{CT}$  analysis of murine femurs after treatment with anti-myeloma drugs.  $\text{Rag2}^{-/-}\text{Il2rg}^{\text{null}}\text{ Il6}^{\text{h/h}}\text{hSIRPa+}$  were engrafted with  $5 \times 10^5$  intrafemorally injected INA-6 cells and treated biweekly with Velcade® or Zometa®, respectively. After four weeks, mice were sacrificed and femurs were fixed in 70 % ethanol for  $\mu\text{CT}$  analysis. Trabecular bone and tissue volumes were quantified to calculate the ratio between bone to tissue volume.

Figure 8 is a set of graphs depicting the results of FACS analysis of primary cell engraftment in  $\text{Rag2}^{-/-}\text{Il2rg}^{\text{null}}\text{hSIRPa+Tpo}^{\text{h/h}}\text{ Mcst}^{\text{h/h}}\text{ Il3/Gmcsf}^{\text{h/h}}\text{ Il6}^{\text{h/h}}$  mice. Mice were transplanted with

$1.5 \times 10^6$  intrafemorally injected CD3-depleted bone marrow cells and sacrificed twelve weeks later. Single-cell suspensions were generated from the injected femur, the collateral leg, and the spleen. Cells were stained for mCD45, hCD19, hCD38, and hCD138. FACS plots show events after gating on mCD45-negative cells. Numbers indicate frequency of CD38+CD138+ cells

Figure 9 depicts the results of experiments assessing the percentage of human hematopoietic (hCD45+) cells in blood in engrafted mice determined by flow cytometry. Horizontal bars indicate the respective mean frequencies.

Figure 10 depicts the results of experiments assessing the percentage of hCD45 of B (CD19+), T (CD3+) and myeloid (CD33+) cells in blood in engrafted mice determined by flow cytometry. Only mice with a hCD45 percentage higher than 2% are shown.

Figure 11, comprising Figures 11A and 11B, depicts the results of experiments assessing percentage (A) and number (B) of human CD45+ cells in BM, spleen and thymus of 20 week engrafted mice. Bars represent average  $\pm$  SEM of 4/5 mice for group.

Figure 12, comprising Figures 12A and 12B, depicts the results of FACS experiments assessing human cells. (A) Drawing showing the gating strategy used for separating different B cell populations by flow cytometry. (B) Percentage of different B cell subsets within the human CD45+CD19+ cells in 20 week old mice. Bars represent average  $\pm$  SEM of 4/5 mice for group.

Figure 13, comprising Figures 13A and 13B, depicts the results of FACS experiments assessing human cells. (A) Representative flow cytometric analysis of CD5+ B cells in human fetal liver (FL) and 20 week old mice. Numbers in the quadrants indicate percentages of cells. All plots are gated on human CD45+ cells. (B) Percentage of CD5 on human B cells in BM and spleen of 20 week engrafted mice. Bars represent average  $\pm$  SEM of 4/5 mice per group.

Figure 14 depicts the results of FACS experiments assessing human cells. (A) Representative flow cytometric analysis of CD27+ B cells in human fetal liver (FL) and 20 week old mice. Numbers in the quadrants indicate percentages of cells. All plots are gated on human CD45+ cells. (B) Percentage of CD27 on human B cells in BM and spleen of 20 week engrafted mice. Bars represent average  $\pm$  SEM of 4/5 mice per group.

Figure 15 depicts the results of experiments assessing the total human IgM and IgG levels in plasma samples of 12 (A) and 20 (B) week old mice. Horizontal bars indicate the respective geometric means. Mice with a PB human engraftment lower than 2% were excluded from the analysis.

## DETAILED DESCRIPTION

**[0023]** Genetically modified rodents are provided that may be used to model human

hematopoietic cell development, function, or disease. The genetically modified rodents comprise a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter and also express at least one of human M-CSF, human IL-3, human GM-CSF, human SIRPa or human TPO. The invention also relates to methods of generating and methods of using the genetically modified rodents described herein. In some embodiments, the genetically modified rodent is a mouse. In some embodiments, the genetically modified rodent described herein is engrafted with human hematopoietic cells, including either normal or neoplastic cells, or combinations thereof. In some embodiments, the genetically modified rodent described herein is engrafted with human multiple myeloma (MM) cells. In various embodiments, the human hematopoietic cell engrafted, genetically modified rodent of the invention are useful for the in vivo evaluation of the growth and differentiation of hematopoietic and immune cells, for the in vivo evaluation of human hematopoiesis, for the in vivo evaluation of cancer cells, for the in vivo assessment of an immune response, for the in vivo evaluation of vaccines and vaccination regimens, for the use in testing the effect of agents that modulate cancer cell growth or survival, for the in vivo evaluation of a treatment of cancer, for the in vivo production and collection of immune mediators, including human antibodies, and for use in testing the effect of agents that modulate hematopoietic and immune cell function.

**[0024]** These and other objects, advantages, and features of the invention will become apparent to those persons skilled in the art upon reading the details of the compositions and methods as more fully described below.

### **Definitions**

**[0025]** Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Such terms are found defined and used in context in various standard references illustratively including J. Sambrook and D. W. Russell, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press; 4th Ed., 2012; F. M. Ausubel, Ed., Short Protocols in Molecular Biology, Current Protocols; 5th Ed., 2002; B. Alberts et al., Molecular Biology of the Cell, 4th Ed., Garland, 2002; D. L. Nelson and M. M. Cox, Lehninger Principles of Biochemistry, 4th Ed., W.H. Freeman & Company, 2004; and Herdewijn, P. (Ed.), Oligonucleotide Synthesis: Methods and Applications, Methods in Molecular Biology, Humana Press, 2004. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described.

**[0026]** As used herein, each of the following terms has the meaning associated with it in this section.

**[0027]** The articles "a" and "an" are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

**[0028]** "About" as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of  $\pm 20\%$  or  $\pm 10\%$ , more preferably  $\pm 5\%$ , even more preferably  $\pm 1\%$ , and still more preferably  $\pm 0.1\%$  from the specified value, as such variations are appropriate to perform the disclosed methods.

**[0029]** The term "abnormal" when used in the context of organisms, tissues, cells or components thereof, includes those organisms, tissues, cells or components thereof that differ in at least one observable or detectable characteristic (e.g., age, treatment, time of day, etc.) from those organisms, tissues, cells or components thereof that display the "normal" (expected) respective characteristic. Characteristics which are normal or expected for one cell or tissue type, might be abnormal for a different cell or tissue type.

**[0030]** The term "antibody," as used herein, includes an immunoglobulin molecule which is able to specifically bind to a specific epitope on an antigen. Antibodies can be intact immunoglobulins derived from natural sources or from recombinant sources and can be immunoreactive portions of intact immunoglobulins. The antibodies in the present invention may exist in a variety of forms including, for example, polyclonal antibodies, monoclonal antibodies, intracellular antibodies ("intrabodies"), Fv, Fab and F(ab)2, as well as single chain antibodies (scFv), heavy chain antibodies, such as camelid antibodies, and humanized antibodies (Harlow et al., 1999, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY; Harlow et al., 1989, *Antibodies: A Laboratory Manual*, Cold Spring Harbor, New York; Houston et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; Bird et al., 1988. *Science* 242:423-426).

**[0031]** "Constitutive" expression includes a state in which a gene product is produced in a living cell under most or all physiological conditions of the cell.

**[0032]** A "coding region" of a gene includes the nucleotide residues of the coding strand of the gene and the nucleotides of the non-coding strand of the gene which are homologous with or complementary to, respectively, the coding region of an mRNA molecule which is produced by transcription of the gene. A "coding region" of a mRNA molecule also includes the nucleotide residues of the mRNA molecule which are matched with an anti-codon region of a transfer RNA molecule during translation of the mRNA molecule or which encode a stop codon. The coding region may thus include nucleotide residues comprising codons for amino acid residues which are not present in the mature protein encoded by the mRNA molecule (e.g., amino acid residues in a protein export signal sequence).

**[0033]** A "disease" includes a state of health of an animal wherein the animal cannot maintain homeostasis, and wherein if the disease is not ameliorated then the animal's health continues to deteriorate.

**[0034]** In contrast, a "disorder" in an animal includes a state of health in which the animal is able to maintain homeostasis, but in which the animal's state of health is less favorable than it would be in the absence of the disorder. Left untreated, a disorder does not necessarily cause

a further decrease in the animal's state of health.

**[0035]** A disease or disorder is "alleviated" if the severity of a symptom of the disease or disorder, the frequency with which such a symptom is experienced by a patient, or both, is reduced.

**[0036]** An "effective amount" or "therapeutically effective amount" of a compound includes that amount of compound which is sufficient to provide a beneficial effect to the subject to which the compound is administered. An "effective amount" of a delivery vehicle includes that amount sufficient to effectively bind or deliver a compound.

**[0037]** "Encoding" includes the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (i.e., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene encodes a protein if, for example, transcription and translation of mRNA corresponding to that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and the non-coding strand, used as the template for transcription of a gene or cDNA, can be referred to as encoding the protein or other product of that gene or cDNA.

**[0038]** As used herein "endogenous" includes any material from or produced inside an organism, cell, tissue or system.

**[0039]** As used herein, the term "exogenous" includes any material introduced from or produced outside an organism, cell, tissue or system.

**[0040]** The terms "expression construct" and "expression cassette" as used herein include a double-stranded recombinant DNA molecule containing a desired nucleic acid human coding sequence and containing one or more regulatory elements necessary or desirable for the expression of the operably linked coding sequence.

**[0041]** As used herein, the term "fragment," as applied to a nucleic acid or polypeptide, includes a subsequence of a larger nucleic acid or polypeptide. A "fragment" of a nucleic acid can be at least about 15 nucleotides in length; for example, at least about 50 nucleotides to about 100 nucleotides; at least about 100 to about 500 nucleotides, at least about 500 to about 1000 nucleotides, at least about 1000 nucleotides to about 1500 nucleotides; or about 1500 nucleotides to about 2500 nucleotides; or about 2500 nucleotides (and any integer value in between). A "fragment" of a polypeptide can be at least about 15 nucleotides in length; for example, at least about 50 amino acids to about 100 amino acids; at least about 100 to about 500 amino acids, at least about 500 to about 1000 amino acids, at least about 1000 amino acids to about 1500 amino acids; or about 1500 amino acids to about 2500 amino acids; or about 2500 amino acids (and any integer value in between).

**[0042]** As used herein, the terms "gene" and "recombinant gene" includes nucleic acid molecules comprising an open reading frame encoding a polypeptide. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of a given gene. Alternative alleles can be identified by sequencing the gene of interest in a number of different individuals. This can be readily carried out by using hybridization probes to identify the same genetic locus in a variety of individuals. Any and all such nucleotide variations and resulting amino acid polymorphisms or variations that are the result of natural allelic variation and that do not alter the functional activity are intended to be within the scope of the invention.

**[0043]** "Homologous" as used herein, includes the subunit sequence similarity between two polymeric molecules, e.g. between two nucleic acid molecules, e.g., two DNA molecules or two RNA molecules, or between two polypeptide molecules. When a subunit position in both of the two molecules is occupied by the same monomeric subunit, e.g., if a position in each of two DNA molecules is occupied by adenine, then they are homologous at that position. The homology between two sequences is a direct function of the number of matching or homologous positions, e.g. if half (e.g., five positions in a polymer ten subunits in length) of the positions in two compound sequences are homologous then the two sequences are 50% homologous, if 90% of the positions, e.g. 9 of 10, are matched or homologous, the two sequences share 90% homology. By way of example, the DNA sequences 5'-ATTGCC-3' and 5'-TATGGC-3' share 50% homology.

**[0044]** The terms "human hematopoietic stem and progenitor cells" and "human HSPC" as used herein, include human self-renewing multipotent hematopoietic stem cells and hematopoietic progenitor cells.

**[0045]** "Inducible" expression includes a state in which a gene product is produced in a living cell in response to the presence of a signal in the cell.

**[0046]** As used herein, an "instructional material" includes a publication, a recording, a diagram, or any other medium of expression which can be used to communicate the usefulness of a compound, composition, vector, or delivery system of the invention in the kit for effecting alleviation of the various diseases or disorders recited herein. Optionally, or alternately, the instructional material can describe one or more methods of alleviating the diseases or disorders in a cell or a tissue of a mammal. The instructional material of the kit of the invention can, for example, be affixed to a container which contains the identified compound, composition, vector, or delivery system of the invention or be shipped together with a container which contains the identified compound, composition, vector, or delivery system. Alternatively, the instructional material can be shipped separately from the container with the intention that the instructional material and the compound be used cooperatively by the recipient.

**[0047]** The term "nucleic acid" includes RNA or DNA molecules having more than one nucleotide in any form including single-stranded, double-stranded, oligonucleotide or

polynucleotide. The term "nucleotide sequence" includes the ordering of nucleotides in an oligonucleotide or polynucleotide in a single-stranded form of nucleic acid.

**[0048]** The term "operably linked" as used herein includes a polynucleotide in functional relationship with a second polynucleotide, e.g. a single-stranded or double-stranded nucleic acid moiety comprising the two polynucleotides arranged within the nucleic acid moiety in such a manner that at least one of the two polynucleotides is able to exert a physiological effect by which it is characterized, upon the other. By way of example, a promoter operably linked to the coding region of a gene is able to promote transcription of the coding region. Preferably, when the nucleic acid encoding the desired protein further comprises a promoter/regulatory sequence, the promoter/regulatory sequence is positioned at the 5' end of the desired protein coding sequence such that it drives expression of the desired protein in a cell. Together, the nucleic acid encoding the desired protein and its promoter/regulatory sequence comprise a "transgene."

**[0049]** The term "polynucleotide" as used herein includes a chain of nucleotides. Furthermore, nucleic acids are polymers of nucleotides. Thus, nucleic acids and polynucleotides as used herein are interchangeable. One skilled in the art has the general knowledge that nucleic acids are polynucleotides, which can be hydrolyzed into the monomeric "nucleotides." The monomeric nucleotides can be hydrolyzed into nucleosides. As used herein polynucleotides include, but are not limited to, all nucleic acid sequences which are obtained by any means available in the art, including, without limitation, recombinant means, i.e., the cloning of nucleic acid sequences from a recombinant library or a cell genome, using ordinary cloning technology and PCR, and the like, and by synthetic means.

**[0050]** As used herein, the terms "peptide," "polypeptide," and "protein" are used interchangeably, and include a compound comprised of amino acid residues covalently linked by peptide bonds. A protein or peptide must contain at least two amino acids, and no limitation is placed on the maximum number of amino acids that can comprise a protein's or peptide's sequence. Polypeptides include any peptide or protein comprising two or more amino acids joined to each other by peptide bonds. As used herein, the term includes both short chains, which also commonly are referred to in the art as peptides, oligopeptides and oligomers, for example, and to longer chains, of which there are many types. "Polypeptides" include, for example, biologically active fragments, substantially homologous polypeptides, oligopeptides, homodimers, heterodimers, variants of polypeptides, modified polypeptides, derivatives, analogs, fusion proteins, among others. The polypeptides include natural peptides, recombinant peptides, synthetic peptides, or a combination thereof. The term "peptide" typically refers to short polypeptides. The term "protein" typically refers to large polypeptides.

**[0051]** The term "progeny" as used herein includes a descendent or offspring and includes the differentiated or undifferentiated decedent cell derived from a parent cell. In one usage, the term progeny includes a descendent cell which is genetically identical to the parent. In another use, the term progeny includes a descendent cell which is genetically and phenotypically identical to the parent. In yet another usage, the term progeny includes a descendent cell that

has differentiated from the parent cell.

**[0052]** The term "promoter" as used herein includes a DNA sequence operably linked to a nucleic acid sequence to be transcribed such as a nucleic acid sequence encoding a desired molecule. A promoter is generally positioned upstream of a nucleic acid sequence to be transcribed and provides a site for specific binding by RNA polymerase and other transcription factors. In specific embodiments, a promoter is generally positioned upstream of the nucleic acid sequence transcribed to produce the desired molecule, and provides a site for specific binding by RNA polymerase and other transcription factors.

**[0053]** Ranges: throughout this disclosure, various aspects of the invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. This applies regardless of the breadth of the range.

**[0054]** A "recombinant polypeptide" includes one which is produced upon expression of a recombinant polynucleotide.

**[0055]** The term "regulatory element" as used herein includes a nucleotide sequence which controls some aspect of the expression of nucleic acid sequences. Exemplary regulatory elements illustratively include an enhancer, an internal ribosome entry site (IRES), an intron; an origin of replication, a polyadenylation signal (pA), a promoter, an enhancer, a transcription termination sequence, and an upstream regulatory domain, which contribute to the replication, transcription, post-transcriptional processing of a nucleic acid sequence. Those of ordinary skill in the art are capable of selecting and using these and other regulatory elements in an expression construct with no more than routine experimentation. Expression constructs can be generated recombinantly or synthetically using well-known methodology.

**[0056]** The term "specifically binds," as used herein with respect to an antibody, includes an antibody which recognizes a specific antigen, but does not substantially recognize or bind other molecules in a sample. For example, an antibody that specifically binds to an antigen from one species may also bind to that antigen from one or more species. But, such cross-species reactivity does not itself alter the classification of an antibody as specific. As another example, an antibody that specifically binds to an antigen may also bind to different allelic forms of the antigen. However, such cross reactivity does not itself alter the classification of an antibody as specific.

**[0057]** In some instances, the terms "specific binding" or "specifically binding", can be used in reference to the interaction of an antibody, a protein, or a peptide with a second chemical

species, to mean that the interaction is dependent upon the presence of a particular structure (e.g., an antigenic determinant or epitope) on the chemical species; for example, an antibody recognizes and binds to a specific protein structure rather than to proteins generally. If an antibody is specific for epitope "A", the presence of a molecule containing epitope A (or free, unlabeled A), in a reaction containing labeled "A" and the antibody, will reduce the amount of labeled A bound to the antibody.

**[0058]** The term "synthetic antibody" as used herein includes an antibody which is generated using recombinant DNA technology, such as, for example, an antibody expressed by a bacteriophage as described herein. The term should also be construed to mean an antibody which has been generated by the synthesis of a DNA molecule encoding the antibody and which DNA molecule expresses an antibody protein, or an amino acid sequence specifying the antibody, wherein the DNA or amino acid sequence has been obtained using synthetic DNA or amino acid sequence technology which is available and well known in the art.

**[0059]** "Variant" as the term is used herein, includes a nucleic acid sequence or a peptide sequence that differs in sequence from a reference nucleic acid sequence or peptide sequence respectively, but retains essential biological properties of the reference molecule. Changes in the sequence of a nucleic acid variant may not alter the amino acid sequence of a peptide encoded by the reference nucleic acid, or may result in amino acid substitutions, additions, deletions, fusions and truncations. Changes in the sequence of peptide variants are typically limited or conservative, so that the sequences of the reference peptide and the variant are closely similar overall and, in many regions, identical. A variant and reference peptide can differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A variant of a nucleic acid or peptide can be a naturally occurring such as an allelic variant, or can be a variant that is not known to occur naturally. Non-naturally occurring variants of nucleic acids and peptides may be made by mutagenesis techniques or by direct synthesis.

**[0060]** The term "genetically modified" includes an animal, the germ cells of which comprise an exogenous human nucleic acid or human nucleic acid sequence. By way of non-limiting examples a genetically modified animal can be a transgenic animal or a knock-in animal, so long as the animal comprises a human nucleic acid sequence.

**[0061]** As used herein, the term "transgenic animal" includes an animal comprising an exogenous human nucleic acid sequence integrated into the genome of the animal.

**[0062]** As used herein, by "knock-in" "knock in" or "knockin" includes a genetic modification that is targeted to a particular chromosomal locus of the non-human animal genome and inserts a nucleic acid of interest into that targeted locus. In some instances, the genetic modification replaces the genetic information encoded at the chromosomal locus in the non-human animal with a different DNA sequence.

#### **Genetically Modified Non-Human Animals**

**[0063]** In some aspects of the invention, a genetically modified rodent that expresses human IL-6 is provided. By human IL-6 (hIL6) it is meant the 184 amino acid protein the sequence for which is described at, e.g., Genbank Accession Nos. NM\_000600.3 and NP\_000591.1. Human IL-6 is a secreted protein that is produced by, for example, T cells, B cells, monocytes, macrophages, fibroblasts, keratinocytes, endothelial cells and myeloma cells. IL-6 acts through a cell surface heterodimeric receptor complex comprising a binding subunit (IL-6R) and a signal transducing subunit (gp130). gp130 is a common component of other receptors, such as the ones for IL-11, IL-27, LIF, whereas the IL-6R is predominantly restricted to hepatocytes, monocytes, activated B cells, resting T cells and myeloma cell lines. IL-6 plays a central role in hematopoiesis, in immune responses and in acute phase reactions, having been shown to be an important factor for the final maturation of B cells into antibody secreting cells (ASC), especially for the expansion of plasmablasts during the germinal center reaction in the T-dependent (TD) antibody response. IL-6 is required for T cell proliferation *in vitro* and for generation of cytotoxic T cells (CTL) *in vivo*, making them more responsive to IL-2.

**[0064]** The genetically modified rodent that expresses human IL-6 also expresses at least one additional human protein selected from human M-CSF, human IL-3, human GM-CSF, human TPO, and human SIRPa, or any combination thereof. In other words, the rodent that expresses human IL-6 may express one, two, three, four or all five of the human proteins selected from hM-CSF, hIL-3, hGM-CSF, hTPO, and hSIRPa. Genetically modified rodent that express hM-CSF, hIL-3, hGM-CSF, hTPO, and/or hSIRPa on which the subject rodent may be designed or from which the subject rodents may be generated are well known in the art, and are discussed in greater detail in, for example, US Application No. US 2013/0042330 and Rathinam et al. 2011, Blood 118:3119-28, disclosing knock-in mice that express human M-CSF; U.S. Patent No. 8,541,646 and Willinger et al. 2011, Proc Natl Acad Sci USA, 108:2390-2395, disclosing knock-in mice that express human IL-3 and human GM-CSF; U.S. Patent No. 8,541,646 and Rongvaux et al. 2011, Proc Natl Acad Sci USA, 108:2378-83, disclosing knock-in mice that express human TPO; and PCT Application No. WO 2012/040207 and Strowig et al. 2011, Proc Natl Acad Sci USA 108(32):13218-13223, disclosing transgenic mice that express human Sirpa.

**[0065]** In various embodiments, the nucleic acid encoding the human protein is operatively linked to one or more regulatory sequences in a manner which allows for transcription of the nucleic acid into mRNA and translation of the mRNA into the human protein. The term "regulatory sequence" is art-recognized and intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are known to those skilled in the art and are described in 1990, Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. In one embodiment, the human nucleic acid is expressed by the native regulatory elements of the human nucleic acid. In another embodiment, the human nucleic acid is expressed by the native regulatory elements of the corresponding nucleic acid of the rodent host animal.

**[0066]** Thus, in some embodiments, the nucleic acid encoding human IL-6 is operably linked to the rodent's IL-6 promoter. In other embodiments, the nucleic acid encoding human IL-6 is operably linked to the human IL-6 promoter. As another example, in some embodiments, the nucleic acid encoding human M-CSF is operably linked to the rodent's M-CSF promoter. In other embodiments, the nucleic acid encoding human M-CSF is operably linked to the human M-CSF promoter. As a third example, in some embodiments, the nucleic acid encoding human IL-3 is operably linked to the rodent's IL-3 promoter. In other embodiments, the nucleic acid encoding human IL-3 is operably linked to the human IL-3 promoter. As a fourth example, in some embodiments, the nucleic acid encoding human GM-CSF is operably linked to the rodent's GM-CSF promoter. In other embodiments, the nucleic acid encoding human GM-CSF is operably linked to the human GM-CSF promoter. As a fifth example, in some embodiments, the nucleic acid encoding human TPO is operably linked to the rodent's TPO promoter. In other embodiments, the nucleic acid encoding human TPO is operably linked to the human TPO promoter.

**[0067]** The skilled artisan will understand that the genetically modified rodents of the invention include genetically modified rodents that express at least one human nucleic acid from a promoter. Nonlimiting examples of ubiquitously expressed promoters useful in the invention include, but are not limited to, DNA pol II promoter, PGK promoter, ubiquitin promoter, albumin promoter, globin promoter, ovalbumin promoter, SV40 early promoter, the Rous sarcoma virus (RSV) promoter, retroviral LTR and lentiviral LTR, a beta-actin promoter, a ROSA26 promoter, a heat shock protein 70 (Hsp70) promoter, an EF-1 alpha gene encoding elongation factor 1 alpha (EF1) promoter, an eukaryotic initiation factor 4A (eIF-4A1) promoter, a chloramphenicol acetyltransferase (CAT) promoter and a CMV (cytomegalovirus) promoter. Promoter and enhancer expression systems useful in the invention also include inducible and/or tissue-specific expression systems. Non-limiting examples of tissue-specific promoters useful in the expression construct of the compositions and methods of the invention include a promoter of a gene expressed in the hematopoietic system, such as an IL-6 promoter, a M-CSF promoter, an IL-3 promoter, a GM-CSF promoter, a SIRPA promoter, a TPO promoter, an IFN- $\beta$  promoter, a Wiskott-Aldrich syndrome protein (WASP) promoter, a CD45 (also called leukocyte common antigen) promoter, a Flt-1 promoter, an endoglin (CD105) promoter and an ICAM-2 (Intracellular Adhesion Molecule 2) promoter. These and other promoters useful in the compositions and methods of the invention are known in the art as exemplified in Abboud et al. (2003, *J. Histochem & Cytochem.* 51:941-949), Schorpp et al. (1996, *NAR* 24:1787-1788), McBurney et al. (1994, *Devel. Dynamics*, 200:278-293) and Majumder et al. (1996, *Blood* 87:3203-3211). Further to comprising a promoter, one or more additional regulatory elements, such as an enhancer element or intron sequence, is included in various embodiments of the invention. Examples of enhancers useful in the compositions and methods of the invention include, but are not limited to, a cytomegalovirus (CMV) early enhancer element and an SV40 enhancer element. Examples of intron sequences useful in the compositions and methods of the invention include, but are not limited to, the beta globin intron or a generic intron. Other additional regulatory elements useful in some embodiments of the invention include, but are not limited to, a transcription termination sequence and an mRNA polyadenylation (pA) sequence.

**[0068]** The skilled artisan will also appreciate that in addition to the naturally occurring human nucleic acid and amino acid sequences, the terms human nucleic acid and human amino acid encompass variants of human nucleic acid and amino acid sequences as well. As used herein, the term "variant" defines either an isolated naturally occurring genetic mutant of a human or a recombinantly prepared variation of a human, each of which contain one or more mutations compared with the corresponding wild-type human. For example, such mutations can be one or more amino acid substitutions, additions, and/or deletions. The term "variant" also includes non-human orthologues. In some embodiments, a variant polypeptide of the present invention has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a wild-type human polypeptide.

**[0069]** The percent identity between two sequences is determined using techniques as those described elsewhere herein. Mutations can be introduced using standard molecular biology techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. One of skill in the art will recognize that one or more amino acid mutations can be introduced without altering the functional properties of human proteins.

**[0070]** Conservative amino acid substitutions can be made in human proteins to produce human protein variants. Conservative amino acid substitutions are art recognized substitutions of one amino acid for another amino acid having similar characteristics. For example, each amino acid may be described as having one or more of the following characteristics: electropositive, electronegative, aliphatic, aromatic, polar, hydrophobic and hydrophilic. A conservative substitution is a substitution of one amino acid having a specified structural or functional characteristic for another amino acid having the same characteristic. Acidic amino acids include aspartate, glutamate; basic amino acids include histidine, lysine, arginine; aliphatic amino acids include isoleucine, leucine and valine; aromatic amino acids include phenylalanine, glycine, tyrosine and tryptophan; polar amino acids include aspartate, glutamate, histidine, lysine, asparagine, glutamine, arginine, serine, threonine and tyrosine; and hydrophobic amino acids include alanine, cysteine, phenylalanine, glycine, isoleucine, leucine, methionine, proline, valine and tryptophan; and conservative substitutions include substitution among amino acids within each group. Amino acids may also be described in terms of relative size, alanine, cysteine, aspartate, glycine, asparagine, proline, threonine, serine, valine, all typically considered to be small.

**[0071]** Human variants can include synthetic amino acid analogs, amino acid derivatives and/or non-standard amino acids, illustratively including, without limitation, alpha-aminobutyric acid, citrulline, canavanine, cyanoalanine, diaminobutyric acid, diaminopimelic acid, dihydroxyphenylalanine, djenkolic acid, homoarginine, hydroxyproline, norleucine, norvaline, 3-phosphoserine, homoserine, 5-hydroxytryptophan, 1-methylhistidine, methylhistidine, and ornithine.

**[0072]** Human variants are encoded by nucleic acids having a high degree of identity with a nucleic acid encoding a wild-type human. The complement of a nucleic acid encoding a human

variant specifically hybridizes with a nucleic acid encoding a wild-type human under high stringency conditions. Nucleic acids encoding a human variant can be isolated or generated recombinantly or synthetically using well-known methodology.

**[0073]** In some embodiments, the genetically modified rodent that expresses a human nucleic acid sequence also expresses the corresponding non-human animal nucleic acid sequence. For example, and as described in greater detail below, in certain embodiments, the human nucleic acid sequence is randomly integrated into the genome of the rodent e.g. such that the rodent comprises the exogenous human nucleic acid sequence at a locus other than the rodent locus encoding the corresponding rodent protein. In other embodiments, the genetically modified rodent that expresses a human nucleic acid sequence does not express the corresponding rodent nucleic acid sequence. For example, and as described in greater detail below, in certain embodiments, the nucleic acid encoding the human protein is introduced into the rodent so as to replace genomic material encoding the corresponding rodent protein, rendering the rodent null for the corresponding rodent gene and deficient for the corresponding rodent protein. In other words, the rodent is a "knock-in" for the human gene.

**[0074]** Thus, in some embodiments, the genetically modified rodent that expresses human IL-6 also expresses rodent IL-6. In other embodiments, the genetically modified rodent that expresses human IL-6 does not express rodent IL-6. As a second example, in some embodiments, the genetically modified rodent that expresses human M-CSF also expresses rodent M-CSF. In other embodiments, the genetically modified rodent that expresses human M-CSF does not express rodent M-CSF. As a third example, in some embodiments, the genetically modified rodent that expresses human IL-3 also expresses rodent IL-3. In other embodiments, the genetically modified rodent that expresses human IL-3 does not express rodent IL-3. As a fourth example, in some embodiments, the genetically modified rodent that expresses human GM-CSF also expresses rodent GM-CSF. In other embodiments, the genetically modified rodent that expresses human GM-CSF does not express rodent GM-CSF. As a fifth example, in some embodiments, the genetically modified rodent that expresses human TPO also expresses rodent TPO. In other embodiments, the genetically modified rodent that expresses human TPO does not express rodent TPO.

**[0075]** The subject genetically modified rodent is immunodeficient. By "immunodeficient," it is meant that the rodent is deficient in one or more aspects of its native immune system, e.g. the rodent is deficient for one or more types of functioning host immune cells, e.g. deficient for non-human B cell number and/or function, non-human T cell number and/or function, non-human NK cell number and/or function, etc.

**[0076]** As one example, the immunodeficient rodent may have severe combined immune deficiency (SCID). SCID refers to a condition characterized by the absence of T cells and lack of B cell function. Examples of SCID include: X-linked SCID, which is characterized by gamma chain gene mutations or loss of the IL2RG gene and the lymphocyte phenotype T(-) B(+) NK(-); and autosomal recessive SCID characterized by Jak3 gene mutations and the lymphocyte phenotype T(-) B(+) NK(-), ADA gene mutations and the lymphocyte phenotype

T(-) B(-) NK(-), IL-7R alpha-chain mutations and the lymphocyte phenotype T(-) B(+) NK(+), CD3 delta or epsilon mutations and the lymphocyte phenotype T(-) B(+) NK(+), RAG1/RAG2 mutations and the lymphocyte phenotype T(-) B(-) NK(+), Artemis gene mutations and the lymphocyte phenotype T(-) B(-) NK(+), CD45 gene mutations and the lymphocyte phenotype T(-) B(+) NK(+), and *Prkdc*<sup>scid</sup> mutations (Bosma et al. (1989, *Immunogenetics* 29:54-56) and the lymphocyte phenotype T(-), B(-), lymphopenia, and hypoglobulinemia. As such, in some embodiments, the genetically modified immunodeficient rodent has one or more deficiencies selected from an IL2 receptor gamma chain deficiency, an ADA gene mutation, an IL7R mutation, a CD3 mutation, a RAG1 and/or RAG2 mutation, an Artemis mutation, a CD45 mutation, and a *Prkdc* mutation.

**[0077]** In certain embodiments, the subject rodent is a mouse or a rat.

**[0078]** In one embodiment, the rodent is selected from a mouse, a rat, and a hamster. In one embodiment, the rodent is selected from the superfamily Muroidea. In one embodiment, the genetically modified rodent is from a family selected from Calomyscidae (e.g., mouse-like hamsters), Cricetidae (e.g., hamster, New World rats and mice, voles), Muridae (true mice and rats, gerbils, spiny mice, crested rats), Nesomyidae (climbing mice, rock mice, with-tailed rats, Malagasy rats and mice), Platacanthomyidae (e.g., spiny dormice), and Spalacidae (e.g., mole rats, bamboo rats, and zokors). In a specific embodiment, the genetically modified rodent is selected from a true mouse or rat (family Muridae), a gerbil, a spiny mouse, and a crested rat. In one embodiment, the genetically modified mouse is from a member of the family Muridae.

**[0079]** In one embodiment, the subject genetically modified rodent is a rat. In one such embodiment, the rat is selected from a Wistar rat, an LEA strain, a Sprague Dawley strain, a Fischer strain, F344, F6, and Dark Agouti. In one another embodiment, the rat strain is a mix of two or more strains selected from the group consisting of Wistar, LEA, Sprague Dawley, Fischer, F344, F6, and Dark Agouti.

**[0080]** In another embodiment, the subject genetically modified rodent is a mouse, e.g. a mouse of a C57BL strain (e.g. C57BL/A, C57BL/An, C57BL/GrFa, C57BL/KaLwN, C57BL/6, C57BL/6J, C57BL/6ByJ, C57BL/6NJ, C57BL/10, C57BL/10ScSn, C57BL/10Cr, C57BL/Ola, etc.); a mouse of the 129 strain (e.g. 129P1, 129P2, 129P3, 129X1, 129S1 (e.g., 129S1/SV, 129S1/Svlm), 129S2, 129S4, 129S5, 129S9/SvEvH, 129S6 (129/SvEvTac), 129S7, 129S8, 129T1, 129T2); a mouse of the BALB strain; e.g., BALB/c; and the like. See, e.g., Festing et al. (1999) *Mammalian Genome* 10:836, see also, Auerbach et al (2000) Establishment and Chimera Analysis of 129/SvEv- and C57BL/6-Derived Mouse Embryonic Stem Cell Lines). In a specific embodiment, the genetically modified mouse is a mix of an aforementioned 129 strain and an aforementioned C57BL/6 strain. In another specific embodiment, the mouse is a mix of aforementioned 129 strains, or a mix of aforementioned BL/6 strains. In a specific embodiment, the 129 strain of the mix is a 129S6 (129/SvEvTac) strain. In yet another embodiment, the mouse is a mix of a BALB strain and another aforementioned strain.

**[0081]** Thus, for example, in some embodiments, the subject genetically modified rodent is an

immunodeficient mouse deficient in B cell number and/or function, and/or T cell number and/or function, and/or NK cell number and/or function (for example, due to an IL2 receptor gamma chain deficiency (i.e.,  $\gamma_c^{-/-}$ ) and/or a RAG deficiency), and having a genome that comprises a human nucleic acid, e.g. a nucleic acid encoding human IL-6, hM-CSF, hIL-3, hGM-CSF, hTPO, and/or hSIRPa, operably linked to its corresponding promoter, e.g. a *M-CSF*, *IL-3*, *GM-CSF*, *TPO* or *SIRPa* promoter, respectively, wherein the rodent expresses the encoded human protein(s).

**[0082]** In certain specific embodiments, the subject genetically modified rodent is an immunodeficient mouse comprising a nucleic acid encoding human IL-6 operably linked to an *IL-6* promoter at the mouse IL-6 locus, and a nucleic acid encoding human SIRPa operably linked to the human *SIRPa* promoter randomly integrated into the genome of the rodent (i.e., the mouse expresses mouse SIRPa), i.e. an immunodeficient hIL-6, hSirpa mouse, e.g. a *Rag2<sup>-/-</sup>IL2rg<sup>-/-</sup> IL-6<sup>h/+</sup> hSIRPa<sup>+</sup>* mouse or a *Rag2<sup>-/-</sup>IL2rg<sup>-/-</sup> IL-6<sup>h/h</sup> hSIRPa<sup>+</sup>* mouse. In some such embodiments, the mouse further comprises a nucleic acid encoding a human M-CSF operably linked to an *M-CSF* promoter, a nucleic acid encoding human IL-3 operably linked to an *IL-3* promoter, a nucleic acid encoding human GM-CSF operably linked to a *GM-CSF* promoter, and a nucleic acid encoding human TPO operably linked to a *TPO* promoter, i.e. an immunodeficient hIL-6, hSirpa, hM-CSF, hIL-3, hGM-CSF, hTPO mouse, e.g. a *Rag2<sup>-/-</sup>IL2rg<sup>-/-</sup> IL-6<sup>h/+</sup> M-CSF<sup>h/+</sup> IL-3<sup>h/+</sup> GM-CSF<sup>h/+</sup> TPO<sup>h/+</sup> hSIRPa<sup>+</sup>* mouse, a *Rag2<sup>-/-</sup>IL2rg<sup>-/-</sup> IL-6<sup>h/+</sup> M-CSF<sup>h/h</sup> IL-3<sup>h/h</sup> GM-CSF<sup>h/h</sup> TPO<sup>h/h</sup> hSIRPa<sup>+</sup>* mouse.

**[0083]** In certain specific embodiments, the subject genetically modified rodent is an immunodeficient mouse comprising a nucleic acid encoding human IL-6 operably linked to an *IL-6* promoter and deficient for mouse IL-6, a nucleic acid encoding human SIRPa operably linked to the human *SIRPa* promoter randomly integrated into the genome of the non-human animal (i.e., the mouse still expresses mouse SIRPa), a nucleic acid encoding human M-CSF operably linked to an *M-CSF* promoter and deficient for mouse M-CSF, a nucleic acid encoding human IL-3 operably linked to an *IL-3* promoter and deficient for mouse IL-3, a nucleic acid encoding human GM-CSF operably linked to a *GM-CSF* promoter and deficient for mouse GM-CSF, and a nucleic acid encoding human TPO operably linked to a *TPO* promoter and deficient for mouse TPO, i.e. a *Rag2<sup>-/-</sup>IL2rg<sup>-/-</sup> IL-6<sup>h/h</sup> M-CSF<sup>h/h</sup> IL-3<sup>h/h</sup> GM-CSF<sup>h/h</sup> TPO<sup>h/h</sup>, hSIRPa<sup>+</sup>* mouse.

#### **Methods of Making Genetically Modified Rodents**

**[0084]** The subject genetically modified rodents may be generated using any convenient method for the generation of genetically modified rodents, e.g. as known in the art or as described herein.

**[0085]** For example, a nucleic acid encoding the human protein of interest, e.g. IL-6, hM-CSF,

hIL-3, hGM-CSF, hTPO, or hSIRPa, may be incorporated into a recombinant vector in a form suitable for insertion into the genome of the host cell and expression of the human protein in a non-human host cell. In various embodiments, the recombinant vector includes the one or more regulatory sequences operatively linked to the nucleic acid encoding the human protein in a manner which allows for transcription of the nucleic acid into mRNA and translation of the mRNA into the human protein, as described above. It will be understood that the design of the vector may depend on such factors as the choice of the host cell to be transfected, the amount of human protein to be expressed, and/or how the encoding nucleic acid will integrate into the genome of the non-human host, e.g. as known in the art.

**[0086]** Any of various methods may then be used to introduce the human nucleic acid sequence into an animal cell to produce a genetically modified animal that expresses the human gene. Such techniques are well-known in the art and include, but are not limited to, pronuclear microinjection of oocytes, transformation of embryonic stem cells, homologous recombination and knock-in techniques. Methods for generating genetically modified animals that can be used include, but are not limited to, those described in Sundberg and Ichiki (2006, Genetically Engineered Mice Handbook, CRC Press), Hofker and van Deursen (2002, Genetically modified Mouse Methods and Protocols, Humana Press), Joyner (2000, Gene Targeting: A Practical Approach, Oxford University Press), Turksen (2002, Embryonic stem cells: Methods and Protocols in Methods Mol Biol., Humana Press), Meyer et al. (2010, Proc. Nat. Acad. Sci. USA 107:15022-15026), and Gibson (2004, A Primer Of Genome Science 2nd ed. Sunderland, Massachusetts: Sinauer), U.S. Pat. No. 6,586,251, Rathinam et al. (2011, Blood 118:3119-28), Willinger et al., (2011, Proc Natl Acad Sci USA, 108:2390-2395), Rongvaux et al., (2011, Proc Natl Acad Sci USA, 108:2378-83) and Valenzuela et al. (2003, Nat Biot 21:652-659).

**[0087]** For example, the subject genetically modified rodents can be created by introducing the nucleic acid encoding the human protein into an oocyte, e.g., by microinjection, and allowing the oocyte to develop in a female foster animal. In preferred embodiments, the construct comprising the human nucleic acid sequence is injected into fertilized oocytes. Fertilized oocytes can be collected from superovulated females the day after mating and injected with the expression construct. The injected oocytes are either cultured overnight or transferred directly into oviducts of 0.5-day p.c. pseudopregnant females. Methods for superovulation, harvesting of oocytes, expression construct injection and embryo transfer are known in the art and described in Manipulating the Mouse Embryo (2002, A Laboratory Manual, 3rd edition, Cold Spring Harbor Laboratory Press). Offspring can be evaluated for the presence of the introduced nucleic acid by DNA analysis (e.g., PCR, Southern blot, DNA sequencing, etc.) or by protein analysis (e.g., ELISA, Western blot, etc.). Such methods typically result in the random integration of the injected nucleic acid sequence -- in this instance, the construct comprising the nucleic acid encoding the human protein of interest - into the genome of the oocyte and hence the non-human animal, i.e. at a locus other than the locus in the host animal expressing the corresponding protein.

**[0088]** As another example, the construct comprising the nucleic acid encoding the human

protein may be transfected into stem cells (ES cells or iPS cells) using well-known methods, such as electroporation, calcium-phosphate precipitation, lipofection, etc. The cells can be evaluated for the presence of the introduced nucleic acid by DNA analysis (e.g., PCR, Southern blot, DNA sequencing, etc.) or by protein analysis (e.g., ELISA, Western blot, etc.). Cells determined to have incorporated the expression construct can then be introduced into preimplantation embryos. For a detailed description of methods known in the art useful for the compositions and methods of the invention, see Nagy et al., (2002, Manipulating the Mouse Embryo: A Laboratory Manual, 3rd edition, Cold Spring Harbor Laboratory Press), Nagy et al. (1990, Development 110:815-821), U.S. Pat. No. 7,576,259, U.S. Pat. No. 7,659,442, U.S. Pat. No. 7,294,754, and Kraus et al. (2010, Genesis 48:394-399). Such methods are typically used in the targeted integration of the transfected nucleic acid sequence -- in this instance, the construct comprising the nucleic acid encoding the rodent human protein of interest - into the genome of the stem cells and hence the rodent. Often, such methods result in the replacement of host genomic material, e.g. genomic material encoding the corresponding host protein, with the nucleic acid encoding the human protein of interest.

**[0089]** A genetically modified founder rodent can be used to breed additional animals carrying the genetic modification. Genetically modified rodents carrying a nucleic acid encoding the human protein(s) of the present disclosure can further be bred to other genetically modified rodents carrying other genetic modifications, or be bred to knockout animals, e.g., a knockout animal that does not express one or more of its genes.

**[0090]** In some embodiments, the genetically modified immunodeficient rodents comprise a genome that includes a nucleic acid encoding a human polypeptide operably linked to a promoter, wherein the rodent expresses the encoded human polypeptide. In various embodiments, the genetically modified immunodeficient rodents comprise a genome that comprises an expression cassette that includes a nucleic acid encoding at least one human polypeptide, wherein the nucleic acid is operably linked to a promoter and a polyadenylation signal and further contains an intron, and wherein the rodent expresses the encoded human polypeptide.

**[0091]** As discussed above, in some embodiments, the subject genetically modified rodent is an immunodeficient rodent. Genetically modified rodents that are immunodeficient and comprise one or more human cytokines, e.g. IL-6, M-CSF, IL-3, GM-CSF, TPO, and/or SIRPa, may likewise be generated using any convenient method for the generation of genetically modified rodents, e.g. as known in the art or as described herein, e.g. DNA injection of an expression construct into a preimplantation embryo or by use of stem cells, such as embryonic stem (ES) cells or induced pluripotent stem (iPS) cells, for example, comprising a mutant SCID gene allele that, when homozygous, will result in immunodeficiency, e.g. as described in greater detail above and in the working examples herein. Mice are then generated with the modified oocyte or ES cells using, e.g. methods described herein and known in the art, and mated to produce the immunodeficient mice comprising the desired genetic modification. As another example, genetically modified rodents can be generated in a non-immunodeficient background, and crossed to an animal comprising a mutant SCID gene allele that, when

homozygous, will result in immunodeficiency, and the progeny mated to create an immunodeficient animal expressing at least one human protein of interest.

**[0092]** Various embodiments of the invention provide genetically modified rodents that include a human nucleic acid in substantially all of their cells, as well as genetically modified rodents that include a human nucleic acid in some, but not all their cells. In some instances, e.g. targeted recombination, one copy of the human nucleic acid will be integrated into the genome of the genetically modified rodents. In other instances, e.g. random integration, multiple copies, adjacent or distant to one another, of the human nucleic acid may be integrated into the genome of the genetically modified rodents.

**[0093]** Thus, in some embodiments, the subject genetically modified rodent may be an immunodeficient rodent comprising a genome that includes a nucleic acid encoding a human polypeptide operably linked to the corresponding rodent promoter, wherein the rodent expresses the encoded human polypeptide. In other words, the subject genetically modified immunodeficient rodent comprises a genome that comprises an expression cassette that includes a nucleic acid encoding at least one human polypeptide, wherein the nucleic acid is operably linked to the corresponding rodent promoter and a polyadenylation signal, and wherein the rodent expresses the encoded human polypeptide.

### Utility

**[0094]** The genetically modified rodents provided in various embodiments of the present invention find many uses including, for example, for use as models of growth and differentiation of hematopoietic cells, for the *in vivo* evaluation of human hematopoiesis, for the *in vivo* evaluation of cancer cells, for *in vivo* study of an immune response, for *in vivo* evaluation of vaccines and vaccination regimens, for the use in testing the effect of agents that modulate cancer cell growth or survival, for the *in vivo* evaluation of a treatment of cancer, for *in vivo* production and collection of immune mediators, such as an antibody, and for use in testing the effect of agents that affect hematopoietic and immune cell function.

**[0095]** Towards this end, in some instances, the subject genetically modified rodent (a "host") is engrafted with at least one human hematopoietic cell. In some disclosures methods are provided for producing a rodent model for studies of the human hematopoietic system, comprising engrafting human hematopoietic cells into a subject genetically modified rodent (the "host"). In certain embodiments, methods are provided for engrafting human hematopoietic cells into the genetically modified rodent disclosed herein.

**[0096]** In some particular instances, the subject genetically modified rodent is engrafted with at least one human multiple myeloma cell. In some such disclosures, methods are provided for producing a rodent model for cancer studies, comprising engrafting human multiple myeloma cells into a subject genetically modified rodent. In some such embodiments, the invention is a method of engrafting human multiple myeloma cells into a subject genetically modified rodent.

The engrafted human multiple myeloma cells useful in the compositions and methods of the invention include any human multiple myeloma cell.

**[0097]** The human hematopoietic cells useful in the engraftment of the subject genetically modified rodents include any convenient human hematopoietic cell. Non-limiting examples of human hematopoietic cells useful in the invention include, but are not limited to, HSC, HSPC, leukemia initiating cells (LIC), and hematopoietic cells of any lineage at any stage of differentiation, including terminally differentiated hematopoietic cells of any lineage. In some instances, the human hematopoietic cell is a primary cell, where "primary cells", "primary cell lines", and "primary cultures" are used interchangeably herein to include acutely isolated cells, or cell cultures that have been derived from a subject and allowed to grow in vitro for a limited number of passages, i.e. splittings, of the culture. For example, primary cultures are cultures that may have been passaged 0 times, 1 time, 2 times, 4 times, 5 times, 10 times, or 15 times, but not enough times go through the crisis stage. In other embodiments, the human hematopoietic cell is from a cell line, that is, the cell is from a culture that is immortalized, e.g. it has been passaged more than about 15 times. In some instances, the hematopoietic cells that are engrafted comprise healthy cells. In other instances, the hematopoietic cells that are engrafted comprise diseased hematopoietic cells, e.g. cancerous hematopoietic cells, e.g. cancerous effector B cells, i.e. multiple myeloma cells. In some instances, the hematopoietic cells that are engrafted comprise both healthy and diseased cells, e.g. healthy B cells and cancerous effector B cells, healthy T cells and cancerous effector B cells, etc.

**[0098]** Hematopoietic cells, i.e. primary cells, cell lines generated therefrom, etc., can be derived from any tissue or location of a human donor, including, but not limited to, bone marrow, peripheral blood, liver, fetal liver, or umbilical cord blood. Such hematopoietic cells can be isolated from any human donor, including healthy donors, as well as donors with disease, such as cancer, including leukemia. Engraftment of hematopoietic cells in the subject genetically modified rodent is characterized by the presence of human hematopoietic cells in the engrafted rodent. In particular embodiments, engraftment of hematopoietic cells in the subject genetically modified rodent is characterized by the presence of differentiated human hematopoietic cells in the engrafted rodent in which hematopoietic cells are provided, as compared with appropriate control animals.

**[0099]** Isolation of human hematopoietic cells, administration of the human hematopoietic cells to a host animal and methods for assessing engraftment thereof are well-known in the art. Hematopoietic cells, including either normal and neoplastic cells, or combinations thereof, for administration to a host animal can be obtained from any tissue containing hematopoietic cells such as, but not limited to, umbilical cord blood, bone marrow, peripheral blood, cytokine or chemotherapy-mobilized peripheral blood and fetal liver. Exemplary methods of isolating human hematopoietic cells, of administering human hematopoietic cells to a host animal, and of assessing engraftment of the human hematopoietic cells in the host animal are described herein and in Pearson et al. (2008, Curr. Protoc. Immunol. 81:1-15), Ito et al. (2002, Blood 100:3175-3182), Traggiai et al. (2004, Science 304:104-107), Ishikawa et al. (2005, Blood 106:1565-1573), Shultz et al. (2005, J. Immunol. 174:6477-6489) and Holyoake et al. (1999,

Exp Hematol. 27:1418-27).

**[0100]** In some embodiments of the invention, the human hematopoietic cells, including either normal and neoplastic cells, or combinations thereof, are isolated from an original source material to obtain a population of cells enriched for a particular hematopoietic cell population (e.g., HSCs, HSPCs, LICs, CD34+, CD34-, lineage specific marker, cancer cell marker, etc.). The isolated hematopoietic cells may or may not be a pure population. In one embodiment, hematopoietic cells useful in the compositions and methods of the invention are depleted of cells having a particular marker. In another embodiment, hematopoietic cells useful in the compositions and methods of the invention are enriched by selection for a marker. In some embodiments, hematopoietic cells useful in the compositions and methods of the invention are a population of cells in which the selected cells constitute about 1-100% of the cells, although in certain embodiments, a population of cells in which the selected cells constitute fewer than 1% of total cells can also be used. In one embodiment, hematopoietic cells useful in the compositions and methods of the invention are depleted of cells having a particular marker, such as CD34. In another embodiment, hematopoietic cells useful in the compositions and methods of the invention are enriched by selection for a marker, such as CD34. In some embodiments, hematopoietic cells useful in the compositions and methods of the invention are a population of cells in which CD34+ cells constitute about 1-100% of the cells, although in certain embodiments, a population of cells in which CD34+ cells constitute fewer than 1 % of total cells can also be used. In certain embodiments, the hematopoietic cells useful in the compositions and methods of the invention are a T cell-depleted population of cells in which CD34+ cells make up about 1-3% of total cells, a lineage-depleted population of cells in which CD34+ cells make up about 50% of total cells, or a CD34+ positive selected population of cells in which CD34+ cells make up about 90% of total cells.

**[0101]** The number of hematopoietic cells administered is not considered limiting with regard to the generation of a human hematopoietic and/or immune system in a genetically modified non-human animal expressing at least one human gene. Thus, by way of non-limiting example, the number of hematopoietic cells administered can range from about  $1 \times 10^3$  to about  $1 \times 10^7$ , although in various embodiments, more or fewer can also be used. By way of another non-limiting example, the number of HSPCs administered can range from about  $3 \times 10^3$  to about  $1 \times 10^6$  CD34+ cells when the recipient is a mouse, although in various embodiments, more or fewer can also be used. For other species of recipient, the number of cells that need to be administered can be determined using only routine experimentation.

**[0102]** For example, in one embodiment, the genetically modified and treated mouse is engrafted with human hematopoietic cells or human hematopoietic stem cells (HSPCs) to form a genetically modified and engrafted mouse. In one embodiment, the hematopoietic cells are selected from human umbilical cord blood cells and human fetal liver cells. In one embodiment, engraftment is with about  $1-2 \times 10^5$  human CD34+ cells.

**[0103]** In some instances, administration of the hematopoietic cells (e.g., normal or neoplastic)

may be preceded by conditioning, e.g. either sub-lethal irradiation of the recipient animal with high frequency electromagnetic radiation, generally using gamma or X-ray radiation, or treatment with a radiomimetic drug such as busulfan or nitrogen mustard. Conditioning is believed to reduce numbers of host hematopoietic cells, create appropriate microenvironmental factors for engraftment of human hematopoietic cells, and/or create microenvironmental niches for engraftment of human hematopoietic cells. Standard methods for conditioning are known in the art, such as described herein and in J. Hayakawa et al, 2009, *Stem Cells*, 27(1):175-182. In one embodiment, the genetically modified mouse is treated so as to eliminate endogenous hematopoietic cells that may exist in the mouse. In one embodiment, the treatment comprises irradiating the genetically modified mouse. In a specific embodiment, newborn genetically modified mouse pups are irradiated sublethally. In a specific embodiment, newborn pups are irradiated 2 x 200 cGy with a four hour interval.

**[0104]** Hematopoietic cells (e.g., normal or neoplastic) can be administered into newborn or adult animals by administration via various routes, such as, but not limited to, intravenous, intrahepatic, intraperitoneal, intrafemoral and/or intratibial. Methods for engraftment of human hematopoietic cells, including either normal and neoplastic cells, or combinations thereof, in immunodeficient rodents are provided according to embodiments of the present invention which include providing human hematopoietic cells to the immunodeficient rodents, with or without irradiating the rodents prior to administration of the hematopoietic cells. Methods for engraftment of human hematopoietic cells in immunodeficient animals are provided according to embodiments of the present invention which include providing human hematopoietic cells, including either normal and neoplastic cells, or combinations thereof, to the genetically modified rodents of the invention, with or without, administering a radiomimetic drug, such as busulfan or nitrogen mustard, to the rodents prior to administration of the hematopoietic cells.

**[0105]** Engraftment of human hematopoietic cells, including either normal and neoplastic cells, or combinations thereof, in the genetically modified rodent of the invention can be assessed by any of various methods, such as, but not limited to, flow cytometric analysis of cells in the animals to which the human hematopoietic cells are administered at one or more time points following the administration of hematopoietic cells.

**[0106]** Generally, engraftment can be considered successful when the number (or percentage) of human hematopoietic cells, including either normal and neoplastic cells, or combinations thereof, present in the genetically modified rodent is greater than the number (or percentage) of human cells that were administered to the rodent at a point in time beyond the lifespan of the administered human hematopoietic cells. Detection of the progeny of the administered hematopoietic cells can be achieved by detection of human DNA in the recipient animal, for example, or by detection of intact human hematopoietic cells, such as by the detection of the human cell marker, such as human CD45, human CD34, or sIL-6R for example. Serial transfer of human hematopoietic cells from a first recipient into a secondary recipient, and engraftment of human hematopoietic cells in the second recipient, is a further optional test of engraftment in the primary recipient. Engraftment can be detected by flow cytometry as 0.05% or greater human CD45+ cells in the blood, spleen or bone marrow at 1-4 months after administration of

the human hematopoietic cells. A cytokine (e.g., GM-CSF) can be used to mobilize stem cells, for example, as described in Watanabe (1997, Bone Marrow Transplantation 19:1175-1181).

**[0107]** In one embodiment, the immunodeficient genetically modified and engrafted rodent gives rise to a human cell selected from a CD34+ cell, a hematopoietic stem cell, a hematopoietic cell, a myeloid precursor cell, a myeloid cell, a dendritic cell, a monocyte, a granulocyte, a neutrophil, a mast cell, a thymocyte, a T cell, a B cell, a platelet, and a combination thereof. In one embodiment, the human cell is present at 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, or 12 months after engraftment.

**[0108]** In one embodiment, the immunodeficient genetically modified and engrafted rodent gives rise to a human hemato-lymphoid system that comprises human hematopoietic stem and progenitor cells, human myeloid progenitor cells, human myeloid cells, human dendritic cells, human monocytes, human granulocytes, human neutrophils, human mast cells, human thymocytes, human T cells, human B cells, and human platelets. In one embodiment, the human hemato-lymphoid system is present at 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, or 12 months after engraftment.

**[0109]** In one embodiment, the immunodeficient genetically modified and engrafted rodent gives rise to a human hemato-lymphoid system that comprises cancerous human hematopoietic cells, for example neoplastic plasma (effector B) cells. In one embodiment, the cancerous human hematopoietic cells are present at 4 weeks, at 6 weeks, at 8 weeks, at 12 weeks, or at more than 12 weeks after engraftment. In certain embodiments, the cancerous human hematopoietic cells are present at 2 weeks, at 4 weeks, at 6 weeks, at 8 weeks, at 12 weeks, or at more than 12 weeks after engraftment.

**[0110]** Once engrafted with human hematopoietic cells, the subject genetically modified rodents find many uses in the art. For example, engrafted genetically modified rodents of the present disclosure are useful for studying the function of human hematopoietic cells in peripheral blood. As demonstrated in working example 2, genetically modified mice that are immunodeficient and comprise a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter at the IL-6 mouse locus (e.g., *Rag2*<sup>-/-</sup>*IL2rg*<sup>null</sup>*IL-6*<sup>h/h</sup> mice, *Rag2*<sup>-/-</sup>*IL2rg*<sup>null</sup>*IL-6*<sup>h/h</sup>*hSIRPa*<sup>+</sup> mice, and *Rag2*<sup>-/-</sup>*IL2rg*<sup>-/-</sup> *IL-6*<sup>h/h</sup> *M-CSF*<sup>h/h</sup> *IL-3*<sup>h/h</sup> *GM-CSF*<sup>h/h</sup> *TPO*<sup>h/h</sup>, *hSIRPa*<sup>+</sup>) support engraftment of human hematopoietic cells, e.g. CD34+ progenitor cells, into the peripheral blood and spleen better than immunodeficient mice that do not express human IL-6, i.e. *Rag2*<sup>-/-</sup>*IL2rg*<sup>null</sup> mice. Moreover, these genetically modified mice promote the differentiation of human hematopoietic cells more efficiently than immunodeficient mice that do not express human IL-6. For example, these genetically modified mice better promote the differentiation of CD5+ B cells and CD27+ B cells. CD5 is a protein found on a subset of IgM-secreting B cells called B-1 cells, and serves to mitigate activating signals from the B cell receptor so that the B-1 cells can only be activated by very strong stimuli (such as bacterial proteins) and not by normal tissue proteins. CD27 is a marker for memory B cells. Additionally, these genetically

modified mice support the development of better-functioning human hematopoietic cells than immunodeficient mice that do not express human IL-6. For example, B cells differentiate into IgG secreting plasma cells more rapidly in these genetically modified mice than in immunodeficient mice that do not express human IL-6. As such, engrafted genetically modified rodents of the present disclosure find use in studying hematopoietic cell development and function, and more particularly, B lymphocyte differentiation and function.

**[0111]** As another example, engrafted genetically modified rodents of the present disclosure are useful for studying hematopoietic cancers. As demonstrated in working example 1 below, genetically modified mice that are immunodeficient and comprise a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter at the mouse IL-6 locus, e.g. *Rag2*<sup>-/-</sup> *IL2rg*<sup>null</sup> *IL-6*<sup>h/h</sup> mice, *Rag2*<sup>-/-</sup> *IL2rg*<sup>null</sup> *IL-6*<sup>h/h</sup> *hSIRPa*<sup>+</sup> mice, and *Rag2*<sup>-/-</sup> *IL2rg*<sup>-/-</sup> *IL-6*<sup>h/h</sup> *M-CSF*<sup>h/h</sup> *IL-3*<sup>h/h</sup> *GM-CSF*<sup>h/h</sup> *TPO*<sup>h/h</sup>, *hSIRPa*<sup>+</sup>, engraft with primary human multiple myeloma cells and cells of human multiple myeloma cell lines, whereas immunodeficient mice that do not express human IL-6, i.e. *Rag2*<sup>-/-</sup> *IL2rg*<sup>null</sup> mice, do not. Expression of human SIRPa by the genetically modified host further improves the rate and extent of engraftment observed. Furthermore, engraftment of the multiple myeloma cells directly to bone of these immunodeficient, genetically modified mice disclosed herein reproduces the bone pathology typically associated with human multiple myeloma, e.g. bone destruction and resorption, e.g. as quantified by  $\mu$ CT scan.

**[0112]** As such, engrafted genetically modified rodents of the present disclosure find use in screening candidate agents to identify those that will treat hematopoietic cancers. The terms "treatment", "treating" and the like are used herein to generally include obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease. "Treatment" as used herein include any treatment of a disease in a mammal, and includes: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, i.e., arresting its development; or (c) relieving the disease, i.e., causing regression of the disease. Candidate agents of interest as therapeutics for hematopoietic cancers include those that may be administered before, during or after the onset of cancer. The treatment of ongoing disease, where the treatment stabilizes or reduces the undesirable clinical symptoms of the patient, is of particular interest. The terms "individual," "subject," "host," and "patient," are used interchangeably herein and include any mammalian subject for whom diagnosis, treatment, or therapy is desired, particularly humans.

**[0113]** As another example, engrafted genetically modified rodents of the present disclosure are useful for studying human pathogens, i.e. pathogens that infect humans; the response of the human immune system to infection by human pathogens; and the effectiveness of agents in protecting against and/or treating infection by human pathogens. The pathogen may be a virus, a fungus, a bacterium, etc. Non-limiting examples of viral pathogens include human or

porcine or avian influenza virus. Non-limiting examples of bacterial pathogens include mycobacterium, e.g. *Mycobacterium tuberculosis* (*M. tuberculosis*), and enterobacterium, e.g. *Salmonella typhi* (*S. typhi*). Examples of methods for infecting mice with *S. typhi* and for assessing infection may be found in, for example, US Published Application No. 2011/0200982. Examples of methods for infecting mice with *M. tuberculosis* and for assessing infection may be found in, for example, US Published Application No. 2011/0200982. Other examples of human pathogens that do not infect wild-type mice, or that infect wild-type mice but the infected mice do not model an immune response that a human mounts in response to the pathogen, will be well-known to the ordinarily skilled artisan. Such mouse models of pathogen infection are useful in research, e.g. to better understand the progression of human infection. Such mouse models of infection are also useful in drug discovery, e.g. to identify candidate agents that protect against or treat infection.

**[0114]** Engrafted genetically modified mice of the present disclosure also provide a useful system for screening candidate agents for desired activities *in vivo*, for example, to identify agents that are able to modulate (*i.e.*, promote or suppress) hematopoietic cell development and/or activity, e.g. the activity of B cells, T cells, NK cells, macrophages, neutrophils, eosinophils, basophils, etc., e.g. in a healthy or a diseased state, e.g. as cancerous cells, during pathogen infection, for example to identify novel therapeutics and/or develop a better understanding of the molecular basis of the development and function of the immune system; for agents that are toxic to hematopoietic cells, e.g. B cells, T cells, NK cells, macrophages, neutrophils, eosinophils, basophils, etc., and progenitors thereof; and for agents that prevent against, mitigate, or reverse the toxic effects of toxic agents on hematopoietic cells, e.g. B cells, T cells, NK cells, macrophages, neutrophils, eosinophils, basophils, etc., and progenitors thereof; etc. As yet another example, engrafted genetically modified rodents of the present disclosure provide a useful system for predicting the responsiveness of an individual to a disease therapy, e.g. by providing an *in vivo* platform for screening the responsiveness of an individual's immune system to an agent, e.g. a therapeutic agent, to predict the responsiveness of an individual to that agent.

**[0115]** In screening assays for biologically active agents, a human hematopoietic cell-engrafted genetically modified mouse of the present disclosure, e.g. an engrafted *Rag2*<sup>-/-</sup> *IL2rg*<sup>-/-</sup> *IL-6*<sup>h/h</sup> *hSIRPa*<sup>+</sup> mouse, an engrafted *Rag2*<sup>-/-</sup> *IL2rg*<sup>-/-</sup> *IL-6*<sup>h/h</sup> *M-CSF*<sup>h/h</sup> *IL-3*<sup>h/h</sup> *GM-CSF*<sup>h/h</sup> *TPO*<sup>h/h</sup>, *hSIRPa*<sup>+</sup> mouse, etc. is contacted with a candidate agent of interest and the effect of the candidate agent is assessed by monitoring one or more output parameters. These output parameters may be reflective of the viability of the cells, e.g. the total number of hematopoietic cells or the number of cells of a particular hematopoietic cell type, or of the apoptotic state of the cells, e.g. the amount of DNA fragmentation, the amount of cell blebbing, the amount of phosphatidylserine on the cell surface, and the like by methods that are well known in the art. Alternatively or additionally, the output parameters may be reflective of the differentiation capacity of the cells, e.g. the proportions of differentiated cells and differentiated cell types. Alternatively or additionally, the output parameters may be reflective of the function of the cells, e.g. the cytokines and chemokines produced by the cells, the antibodies (e.g. amount or type)

produced by the cells, the ability of the cells to home to and extravasate to a site of challenge, the ability of the cells to modulate, *i.e.* promote or suppress, the activity of other cells in vitro or in vivo, etc. Other output parameters may be reflective of the extent of damage induced by diseased hematopoietic cells, e.g. bone destruction and resorption induced by multiple myeloid cells. Yet other parameters may be reflective of the effect of the agent on infection, e.g. pathogen infection in the animal, e.g. the titer of pathogen in the mouse, the presence of granuloma in the mouse, etc., as relevant to the studies being performed.

**[0116]** Parameters are quantifiable components of cells, particularly components that can be accurately measured, desirably in a high throughput system. A parameter can be any cell component or cell product including cell surface determinant, receptor, protein or conformational or posttranslational modification thereof, lipid, carbohydrate, organic or inorganic molecule, nucleic acid, e.g. mRNA, DNA, etc. or a portion derived from such a cell component or combinations thereof. While most parameters will provide a quantitative readout, in some instances a semi-quantitative or qualitative result will be acceptable. Readouts may include a single determined value, or may include mean, median value or the variance, etc. Characteristically a range of parameter readout values will be obtained for each parameter from a multiplicity of the same assays. Variability is expected and a range of values for each of the set of test parameters will be obtained using standard statistical methods with a common statistical method used to provide single values.

**[0117]** Candidate agents of interest for screening include known and unknown compounds that encompass numerous chemical classes, primarily organic molecules, which may include organometallic molecules, inorganic molecules, genetic sequences, vaccines, antibiotics or other agents suspected of having antibiotic properties, peptides, polypeptides, antibodies, agents that have been approved pharmaceutical for use in a human, etc. An important aspect of the invention is to evaluate candidate drugs, including toxicity testing; and the like.

**[0118]** Candidate agents include organic molecules comprising functional groups necessary for structural interactions, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, frequently at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules, including peptides, polynucleotides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Included are pharmacologically active drugs, genetically active molecules, etc. Compounds of interest include chemotherapeutic agents, hormones or hormone antagonists, etc. Exemplary of pharmaceutical agents suitable for this invention are those described in, "The Pharmacological Basis of Therapeutics," Goodman and Gilman, McGraw-Hill, New York, N.Y., (1996), Ninth edition. Also included are toxins, and biological and chemical warfare agents, for example see Somani, S. M. (Ed.), "Chemical Warfare Agents," Academic Press, New York, 1992).

**[0119]** Candidate agents of interest for screening also include nucleic acids, for example,

nucleic acids that encode siRNA, shRNA, antisense molecules, or miRNA, or nucleic acids that encode polypeptides. Many vectors useful for transferring nucleic acids into target cells are available. The vectors may be maintained episomally, e.g. as plasmids, minicircle DNAs, virus-derived vectors such as cytomegalovirus, adenovirus, etc., or they may be integrated into the target cell genome, through homologous recombination or random integration, e.g. retrovirus derived vectors such as MMLV, HIV-1, ALV, etc. Vectors may be provided directly to the subject cells. In other words, the pluripotent cells are contacted with vectors comprising the nucleic acid of interest such that the vectors are taken up by the cells.

**[0120]** Methods for contacting cells, e.g. cells in culture or cells in a mouse, with nucleic acid vectors, such as electroporation, calcium chloride transfection, and lipofection, are well known in the art. Alternatively, the nucleic acid of interest may be provided to the cells via a virus. In other words, the cells are contacted with viral particles comprising the nucleic acid of interest. Retroviruses, for example, lentiviruses, are particularly suitable to the method of the invention. Commonly used retroviral vectors are "defective", i.e. unable to produce viral proteins required for productive infection. Rather, replication of the vector requires growth in a packaging cell line. To generate viral particles comprising nucleic acids of interest, the retroviral nucleic acids comprising the nucleic acid are packaged into viral capsids by a packaging cell line. Different packaging cell lines provide a different envelope protein to be incorporated into the capsid, this envelope protein determining the specificity of the viral particle for the cells. Envelope proteins are of at least three types, ecotropic, amphotropic and xenotropic. Retroviruses packaged with ecotropic envelope protein, e.g. MMLV, are capable of infecting most murine and rat cell types, and are generated by using ecotropic packaging cell lines such as BOSC23 (Pear et al. (1993) P.N.A.S. 90:8392-8396). Retroviruses bearing amphotropic envelope protein, e.g. 4070A (Danos et al. *supra*), are capable of infecting most mammalian cell types, including human, dog and mouse, and are generated by using amphotropic packaging cell lines such as PA12 (Miller et al. (1985) Mol. Cell. Biol. 5:431-437); PA317 (Miller et al. (1986) Mol. Cell. Biol. 6:2895-2902); GRIP (Danos et al. (1988) PNAS 85:6460-6464). Retroviruses packaged with xenotropic envelope protein, e.g. AKR env, are capable of infecting most mammalian cell types, except murine cells. The appropriate packaging cell line may be used to ensure that the cells of interest—in some instance, the engrafted cells, in some instance, the cells of the host, i.e. the genetically modified animal--are targeted by the packaged viral particles.

**[0121]** Vectors used for providing nucleic acid of interest to the subject cells will typically comprise suitable promoters for driving the expression, that is, transcriptional activation, of the nucleic acid of interest. This may include ubiquitously acting promoters, for example, the CMV- $\beta$ -actin promoter, or inducible promoters, such as promoters that are active in particular cell populations or that respond to the presence of drugs such as tetracycline. By transcriptional activation, it is intended that transcription will be increased above basal levels in the target cell by at least about 10 fold, by at least about 100 fold, more usually by at least about 1000 fold. In addition, vectors used for providing reprogramming factors to the subject cells may include genes that must later be removed, e.g. using a recombinase system such as Cre/Lox, or the cells that express them destroyed, e.g. by including genes that allow selective toxicity such as herpesvirus TK, bcl-xs, etc

**[0122]** Candidate agents of interest for screening also include polypeptides. Such polypeptides may optionally be fused to a polypeptide domain that increases solubility of the product. The domain may be linked to the polypeptide through a defined protease cleavage site, e.g. a TEV sequence, which is cleaved by TEV protease. The linker may also include one or more flexible sequences, e.g. from 1 to 10 glycine residues. In some embodiments, the cleavage of the fusion protein is performed in a buffer that maintains solubility of the product, e.g. in the presence of from 0.5 to 2 M urea, in the presence of polypeptides and/or polynucleotides that increase solubility, and the like. Domains of interest include endosomolytic domains, e.g. influenza HA domain; and other polypeptides that aid in production, e.g. IF2 domain, GST domain, GRPE domain, and the like. Additionally or alternatively, such polypeptides may be formulated for improved stability. For example, the peptides may be PEGylated, where the polyethyleneoxy group provides for enhanced lifetime in the blood stream. The polypeptide may be fused to another polypeptide to provide for added functionality, e.g. to increase the *in vivo* stability. Generally such fusion partners are a stable plasma protein, which may, for example, extend the *in vivo* plasma half-life of the polypeptide when present as a fusion, in particular wherein such a stable plasma protein is an immunoglobulin constant domain. In most cases where the stable plasma protein is normally found in a multimeric form, e.g., immunoglobulins or lipoproteins, in which the same or different polypeptide chains are normally disulfide and/or noncovalently bound to form an assembled multichain polypeptide, the fusions herein containing the polypeptide also will be produced and employed as a multimer having substantially the same structure as the stable plasma protein precursor. These multimers will be homogeneous with respect to the polypeptide agent they comprise, or they may contain more than one polypeptide agent.

**[0123]** The candidate polypeptide agent may be produced from eukaryotic cells, or may be produced by prokaryotic cells. It may be further processed by unfolding, e.g. heat denaturation, DTT reduction, etc. and may be further refolded, using methods known in the art. Modifications of interest that do not alter primary sequence include chemical derivatization of polypeptides, e.g., acylation, acetylation, carboxylation, amidation, etc. Also included are modifications of glycosylation, e.g. those made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing or in further processing steps; e.g. by exposing the polypeptide to enzymes which affect glycosylation, such as mammalian glycosylating or deglycosylating enzymes. Also embraced are sequences that have phosphorylated amino acid residues, e.g. phosphotyrosine, phosphoserine, or phosphothreonine. The polypeptides may have been modified using ordinary molecular biological techniques and synthetic chemistry so as to improve their resistance to proteolytic degradation or to optimize solubility properties or to render them more suitable as a therapeutic agent. Analogs of such polypeptides include those containing residues other than naturally occurring L-amino acids, e.g. D-amino acids or non-naturally occurring synthetic amino acids. D-amino acids may be substituted for some or all of the amino acid residues.

**[0124]** The candidate polypeptide agent may be prepared by *in vitro* synthesis, using conventional methods as known in the art. Various commercial synthetic apparatuses are

available, for example, automated synthesizers by Applied Biosystems, Inc., Beckman, etc. By using synthesizers, naturally occurring amino acids may be substituted with unnatural amino acids. The particular sequence and the manner of preparation will be determined by convenience, economics, purity required, and the like. Alternatively, the candidate polypeptide agent may be isolated and purified in accordance with conventional methods of recombinant synthesis. A lysate may be prepared of the expression host and the lysate purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, or other purification technique. For the most part, the compositions which are used will comprise at least 20% by weight of the desired product, more usually at least about 75% by weight, preferably at least about 95% by weight, and for therapeutic purposes, usually at least about 99.5% by weight, in relation to contaminants related to the method of preparation of the product and its purification. Usually, the percentages will be based upon total protein.

**[0125]** In some cases, the candidate polypeptide agents to be screened are antibodies. The term "antibody" or "antibody moiety" is intended to include any polypeptide chain-containing molecular structure with a specific shape that fits to and recognizes an epitope, where one or more non-covalent binding interactions stabilize the complex between the molecular structure and the epitope. The specific or selective fit of a given structure and its specific epitope is sometimes referred to as a "lock and key" fit. The archetypal antibody molecule is the immunoglobulin, and all types of immunoglobulins, IgG, IgM, IgA, IgE, IgD, etc., from all sources, e.g. human, rodent, rabbit, cow, sheep, pig, dog, other mammal, chicken, other avians, etc., are considered to be "antibodies." Antibodies utilized in the present invention may be either polyclonal antibodies or monoclonal antibodies. Antibodies are typically provided in the media in which the cells are cultured. Antibody production and screen is discussed in greater detail below.

**[0126]** Candidate agents may be obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds, including biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

**[0127]** Candidate agents are screened for biological activity by administering the agent to at least one and usually a plurality of samples, sometimes in conjunction with samples lacking the agent. The change in parameters in response to the agent is measured, and the result evaluated by comparison to reference cultures, e.g. in the presence and absence of the agent, obtained with other agents, etc. In instances in which a screen is being performed to identify candidate agents that will prevent, mitigate or reverse the effects of a toxic agent, the screen is typically performed in the presence of the toxic agent, where the toxic agent is added at the

time most appropriate to the results to be determined. For example, in cases in which the protective/preventative ability of the candidate agent is tested, the candidate agent may be added before the toxic agent, simultaneously with the candidate agent, or subsequent to treatment with the candidate agent. As another example, in cases in which the ability of the candidate agent to reverse the effects of a toxic agent is tested, the candidate agent may be added subsequent to treatment with the candidate agent. As mentioned above, in some instances, the "sample" is a genetically modified non-human animal that has been engrafted with cells, e.g. the candidate agent is provided to an immunodeficient animal, e.g. mouse, comprising a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter that has been engrafted with human hematopoietic cells. In some instances, the sample is the human hematopoietic cells to be engrafted, i.e. the candidate agent is provided to cells prior to engraftment into the immunodeficient genetically modified animal.

**[0128]** If the candidate agent is to be administered directly to the engrafted genetically modified animal, the agent may be administered by any of a number of well-known methods in the art for the administration of peptides, small molecules and nucleic acids to mice. For example, the agent may be administered orally, mucosally, topically, intradermally, or by injection, e.g. intraperitoneal, subcutaneous, intramuscular, intravenous, or intracranial injection, and the like. The agent may be administered in a buffer, or it may be incorporated into any of a variety of formulations, e.g. by combination with appropriate pharmaceutically acceptable vehicle. "Pharmaceutically acceptable vehicles" may be vehicles approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in mammals, such as humans. The term "vehicle" refers to a diluent, adjuvant, excipient, or carrier with which a compound of the invention is formulated for administration to a mammal. Such pharmaceutical vehicles can be lipids, e.g. liposomes, e.g. liposome dendrimers; liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like, saline; gum acacia, gelatin, starch paste, talc, keratin, colloidal silica, urea, and the like. In addition, auxiliary, stabilizing, thickening, lubricating and coloring agents may be used. Pharmaceutical compositions may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. The agent may be systemic after administration or may be localized by the use of regional administration, intramural administration, or use of an implant that acts to retain the active dose at the site of implantation. The active agent may be formulated for immediate activity or it may be formulated for sustained release. For some conditions, particularly central nervous system conditions, it may be necessary to formulate agents to cross the blood-brain barrier (BBB). One strategy for drug delivery through the blood-brain barrier (BBB) entails disruption of the BBB, either by osmotic means such as mannitol or leukotrienes, or biochemically by the use of vasoactive substances such as bradykinin. A BBB disrupting agent can be co-administered with the agent when the compositions are administered by intravascular injection. Other strategies to go through the BBB may entail the use of endogenous transport systems, including Caveolin-1 mediated transcytosis, carrier-mediated transporters such as glucose and amino acid carriers, receptor-mediated transcytosis for insulin or transferrin, and active efflux

transporters such as p-glycoprotein. Active transport moieties may also be conjugated to the therapeutic compounds for use in the invention to facilitate transport across the endothelial wall of the blood vessel. Alternatively, drug delivery of agents behind the BBB may be by local delivery, for example by intrathecal delivery, e.g. through an Ommaya reservoir (see e.g. US Patent Nos. 5,222,982 and 5385582); by bolus injection, e.g. by a syringe, e.g. intravitrally or intracranially; by continuous infusion, e.g. by cannulation, e.g. with convection (see e.g. US Application No. 20070254842); or by implanting a device upon which the agent has been reversably affixed (see e.g. US Application Nos. 20080081064 and 20090196903).

**[0129]** If the agent(s) are provided to cells prior to engraftment, the agents are conveniently added in solution, or readily soluble form, to the medium of cells in culture. The agents may be added in a flow-through system, as a stream, intermittent or continuous, or alternatively, adding a bolus of the compound, singly or incrementally, to an otherwise static solution. In a flow-through system, two fluids are used, where one is a physiologically neutral solution, and the other is the same solution with the test compound added. The first fluid is passed over the cells, followed by the second. In a single solution method, a bolus of the test compound is added to the volume of medium surrounding the cells. The overall concentrations of the components of the culture medium should not change significantly with the addition of the bolus, or between the two solutions in a flow through method.

**[0130]** A plurality of assays may be run in parallel with different agent concentrations to obtain a differential response to the various concentrations. As known in the art, determining the effective concentration of an agent typically uses a range of concentrations resulting from 1:10, or other log scale, dilutions. The concentrations may be further refined with a second series of dilutions, if necessary. Typically, one of these concentrations serves as a negative control, *i.e.* at zero concentration or below the level of detection of the agent or at or below the concentration of agent that does not give a detectable change in the phenotype.

**[0131]** An analysis of the response of cells in the engrafted genetically modified animal to the candidate agent may be performed at any time following treatment with the agent. For example, the cells may be analyzed 1, 2, or 3 days, sometimes 4, 5, or 6 days, sometimes 8, 9, or 10 days, sometimes 14 days, sometimes 21 days, sometimes 28 days, sometimes 1 month or more after contact with the candidate agent, *e.g.* 2 months, 4 months, 6 months or more. In some disclosures, the analysis comprises analysis at multiple time points. The selection of the time point(s) for analysis will be based upon the type of analysis to be performed, as will be readily understood by the ordinarily skilled artisan.

**[0132]** The analysis may comprise measuring any of the parameters described herein or known in the art for measuring cell viability, cell proliferation, cell identity, cell morphology, and cell function, particularly as they may pertain to cells of the immune cells. For example, flow cytometry may be used to determine the total number of hematopoietic cells or the number of cells of a particular hematopoietic cell type. Histochemistry or immunohistochemistry may be performed to determine the apoptotic state of the cells, *e.g.* terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) to measure DNA fragmentation, or

immunohistochemistry to detect Annexin V binding to phosphatidylserine on the cell surface. Flow cytometry may also be employed to assess the proportions of differentiated cells and differentiated cell types, e.g. to determine the ability of hematopoietic cells to differentiate in the presence of agent. ELISAs, Westerns, and Northern blots may be performed to determine the levels of cytokines, chemokines, immunoglobulins, etc. expressed in the engrafted genetically modified mice, e.g. to assess the function of the engrafted cells, to assess the survival of cancerous plasma cells, etc.  $\mu$ CT scans may be performed to determine the extent of damage induced by diseased hematopoietic cells, e.g. bone destruction and resorption induced by multiple myeloid cells. In vivo assays to test the function of immune cells, as well as assays relevant to particular diseases or disorders of interest such as diabetes, autoimmune disease, graft v. host disease, AMD, etc. may also be performed. See, e.g. Current Protocols in Immunology (Richard Coico, ed. John Wiley & Sons, Inc. 2012) and Immunology Methods Manual (I. Lefkovits ed., Academic Press 1997),.

**[0133]** So, for example, a method is disclosed for determining the effect of an agent on multiple myeloma, comprising administering the agent to a humanized IL-6 mouse, e.g. a *Rag2*<sup>-/-</sup>/*IL2rg*<sup>-/-</sup>/*IL-6*<sup>h/h</sup> mouse, that has been engrafted with human multiple myeloma cells; measuring a parameter of the viability and/or proliferative ability of the multiple myeloma cells over time in the presence of the agent; and comparing that measurement to the measurement from an engrafted humanized IL-6 mouse not exposed to the agent. The agent is determined to be anti-cancerous if it reduces the proliferation of and/or reduces the number of multiple myeloma cells in blood or a tissue of the mouse by at least 20%, 30%, 40% or more, in some instances 50%, 60%, 70% or more, e.g. 80%, 90% or 100%, i.e., to undetectable amounts, following a single administration or two or more administrations of the agent over a selected period of time. In a specific embodiment, the administration of the drug or combination of drugs is at least a week, 10 days, two week, three weeks, or four weeks after engraftment of the multiple myeloma cells.

**[0134]** Other examples of uses for the subject mice are provided elsewhere herein. Additional applications of the genetically modified and engrafted mice described in this disclosure will be apparent to those skilled in the art upon reading this disclosure.

#### Human Antibody Production

**[0135]** Also provided are compositions and methods useful for the production of human monoclonal antibodies from an engrafted immunodeficient rodent, as elsewhere described herein. In various embodiments, the methods comprise contacting an immunodeficient rodent with a human hematopoietic cell to generate an immune system-transplanted rodent (engrafted rodent), subsequently contacting the engrafted rodent with an antigen, collecting from the engrafted rodent a human cell producing a human antibody against the antigen, and isolating the antibody from the antibody producing cell.

**[0136]** In various embodiments, the invention comprises a method that includes establishing an antibody producing cell (e.g., a human B-cell) by a transformation method (e.g. EBV) or a cell fusion method (e.g. hybridoma). Preferably the antibody producing cell is capable of being maintained under suitable cell culture conditions for at least about 50 passages.

**[0137]** In various embodiments, the engrafted animal is a mouse or a rat.

**[0138]** In various embodiments of the invention, the human hematopoietic cell is CD34+ cell obtained from a human fetal liver, bone marrow, cord blood, peripheral blood, or spleen sample.

**[0139]** In various embodiments, the antigen at least one of: a peptide, a polypeptide, an MHC/peptide complex, DNA, a live virus, a dead virus or portion thereof, a live bacteria, a dead bacteria or portion thereof, or a cancer cell or portion thereof.

**[0140]** In some embodiments, the engrafted animal has been contacted with the antigen 1-5 months after the animal has been contacted with the human hematopoietic cell. In some embodiments, the engrafted animal is contacted only one time with the antigen, while in other embodiments, the engrafted animal is contacted two, three, four, five, six, seven, eight, or more times with the antigen.

**[0141]** In one embodiment, human antibody producing cell collected from the engrafted animal is a B cell. In various embodiments, the human antibody producing cell collected from the animal expresses on its surface at least one of: CD19, CD20, CD22, and CD27. The human antibody-producing cell of the invention can be recovered by removal of any suitable cellular components of the immune system from the animal. In various embodiments, the antibody-producing cell is removed from the engrafted animal by removal of at least one of the spleen, the lymph nodes, the peripheral blood, the bone marrow or portions thereof.

**[0142]** In various embodiments, the method of the invention employs a conventional hybridoma technology using a suitable fusion partner. In various embodiments, the fusion partner at least one cell selected from the group consisting of: MOPC21, P3X63AG8, SP2/0, NS-1, P3.X63AG8.653, F0, S194/5.XXO.BU-1, FOX-NY, SP2/0-Ag14, MEG-01, HEL, UT-7, M07e, MEG-A2, and DAMI, and cell lines derived from these cells.

**[0143]** Methods of isolating an antibody from the engrafted rodent of the invention are well known in the art. Isolation of the antibody from the antibody producing cell, the media in which the antibody producing cell is culture, and/or the ascites of the engrafted animal, can be performed according to the methods known in the art, such as, by way of example, chromatography and dialysis. In other various embodiments, the antibody can be isolated using one or more of immunoaffinity purification, ammonium sulphate precipitation, protein A/G purification, ion exchange chromatography and gel filtration. Such methods are described in Nau (1989, Optimization of monoclonal antibody purification, In: Techniques in Protein Chemistry, Hugli, T. (ed.), Academic Press, New York) and Coligan et al. (2005, Current

Protocols in Immunology, John Wiley & Sons, Inc.).

**[0144]** The antigen may be administered to the engrafted rodent by any suitable means known in art. In various embodiments, the antigen can be administered to the engrafted rodent by at least one of intrasplenically, intravenously, intraperitoneally, intradermally, intramuscularly, and subcutaneously. In some embodiments, the antigen is administered alone and in other embodiments, the antigen is administered in combination with appropriate immunomodulating agent or adjuvant. Examples of adjuvants useful in the methods of the invention include, but are not limited to, Complete Freund's Adjuvant (CFA), Incomplete Freund's Adjuvant (IFA), and Alum ( $\text{Al}_3(\text{OH})_4$ ).

### **Reagents and Kits**

**[0145]** Also disclosed are reagents and kits thereof for practicing one or more of the above-described methods. The subject reagents and kits thereof may vary greatly. The reagents or kits may comprise one or more reagents for use in the generation and/or maintenance of the subject genetically modified rodents. For example, the kit may comprise an immunodeficient mouse comprising a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter and a nucleic acid encoding human SIRPa operably linked to a SIRPa promoter; or a mouse comprising a nucleic acid encoding human IL-6 operably linked to an IL-6 promoter and further comprising a nucleic acid encoding human M-CSF operably linked to an M-CSF promoter; a nucleic acid encoding human IL-3 operably linked to an IL-3 promoter; a nucleic acid encoding human GM-CSF operably linked to a GM-CSF promoter; a nucleic acid encoding human TPO operably linked to a TPO promoter; and/or a nucleic acid encoding human SIRPa operably linked to a SIRPa promoter. The kit may comprise reagents for breeding such mice, e.g. primers for genotyping for the human IL-6 gene, for the human M-CSF gene, for the human IL-3 gene, for the human GM-CSF gene, for the human SIRPa gene, and/or for the human TPO gene, PCR buffer,  $\text{MgCl}_2$  solution, etc.

**[0146]** The reagents or kits may comprise one or more reagents for use in engrafting the subject genetically modified rodents, for example human hematopoietic cells, an enriched population of human hematopoietic progenitor cells, a hematopoietic cell line, a neoplastic hematopoietic cell line, etc. for transplantation into the subject genetically modified rodents, or reagents for preparing a population of hematopoietic cells, an enriched population of hematopoietic cells from a human, a hematopoietic cell line, a neoplastic hematopoietic cell line, etc. for transplantation into a subject genetically modified rodents.

**[0147]** The reagents or kits may include reagents for determining the viability and/or function of hematopoietic cells, e.g. in the presence/absence of a candidate agent, e.g. one or more antibodies that are specific for markers expressed by different types of hematopoietic cells, or reagents for detecting particular cytokines, chemokine, etc.. Other reagents may include culture media, culture supplements, matrix compositions, and the like.

**[0148]** In addition to the above components, the subject kits may further include instructions for practicing the subject methods. These instructions may be present in the subject kits in a variety of forms, one or more of which may be present in the kit. One form in which these instructions may be present is as printed information on a suitable medium or substrate, e.g., a piece or pieces of paper on which the information is printed, in the packaging of the kit, in a package insert, etc. Yet another means would be a computer readable medium, e.g., diskette, CD, etc., on which the information has been recorded. Yet another means that may be present is a website address which may be used via the internet to access the information at a removed site. Any convenient means may be present in the kits.

## **EXPERIMENTAL EXAMPLES**

**[0149]** The invention is further described in detail by reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified. Thus, the invention should in no way be construed as being limited to the following examples, but rather, should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

**[0150]** Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out the preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

**Example 1: Genetic humanization of cytokine genes enables engraftment of mice with human multiple myeloma cells.**

**[0151]** The data described herein demonstrate that the genetically modified non-human animals described herein represent a novel *in vivo* animal model for multiple myeloma.

**Materials and Methods**

**[0152] Mice.** Humanized IL-6 knock-in mice were generated by replacing 6.8 kb of the murine IL-6 gene locus with a 4.8-kb human IL-6 gene sequence containing exons 1 through 5 including 3' untranslated region of the human IL-6 gene.

**[0153]** Briefly, a targeting construct for replacing the mouse with the human IL-6 gene in a single targeting step was constructed using VELOCIGENE® genetic engineering technology (see, Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech*, 21(6):652-659). Mouse and human IL-6

DNA were obtained from bacterial artificial chromosome (BAC) RPCI-23 clone 368C3, and from BAC CTD clone 2369M23, respectively. Briefly, a NotI linearized targeting construct generated by gap repair cloning containing mouse IL-6 upstream and downstream homology arms flanking a 4.8 kb human IL-6 sequence extending from ATG in exon 1 to exon 5 with 16 nucleotides of 3' downstream sequence (genomic coordinates: NCBIh37.1: ch7:22,766,882 to 22,771,637) and a floxed neo selection cassette, was electroporated into Rag2<sup>+/-</sup> IL2rg<sup>Y/-</sup> ES cells. The parental ES cell line in which the RAG2 gene and IL2rg gene knockout was made was a commercially available V17 ES cell (BALB/cxL29 heterozygote). Correctly targeted ES cells may be electroporated with a transient Cre-expressing vector to remove the drug selection cassette.

**[0154]** Correctly targeted ES cell clones were identified by a loss-of-native-allele (LONA) assay (Valenzuela et al. 2003) in which the number of copies of the native, unmodified IL6 gene were determined by two TaqMan™ quantitative polymerase chain reactions (qPCRs) specific for sequences in the mouse IL6 gene that were targeted for deletion. The qPCR assays comprised the following primer-probe sets (written 5' to 3'): upstream forward primer, TTGCCGGTTT TCCCTTTCT C (SEQ ID NO:1); upstream reverse primer, AGGGAAGGCC GTGGTTGTC (SEQ ID NO:2); upstream probe, FAM-CCAGCATCAG TCCCAAGAAG GCAACT-BHQ (SEQ ID NO:3); downstream forward primer, TCAGAGTGTG GGCGAACAAA G (SEQ ID NO:4); downstream reverse primer, GTGGCAAAAG CAGCCTTAGC (SEQ ID NO:5); downstream probe, FAM-TCATTCCAGG CCCTTCTTAT TGCATCTG-BHQ (SEQ ID NO:6); in which FAM refers to the 5-carboxyfluorescein fluorescent probe and BHQ refers to the fluorescence quencher of the black hole quencher type (Biosearch Technologies). DNA purified from ES cell clones that have taken up the targeting vector and incorporated in their genomes was combined with TaqMan™ Gene Expression Master Mix (Life Technologies) according to the manufacturer's suggestions in a 384-well PCR plate (MicroAmp™ Optical 384-Well Reaction Plate, Life Technologies) and cycled in an Applied Biosystems Prism 7900HT, which collects fluorescence data during the course of the PCRs and determines a threshold cycle (C<sub>t</sub>), the fractional PCR cycle at which the accumulated fluorescence reaches a pre-set threshold. The upstream and downstream IL6-specific qPCRs and two qPCRs for non-targeted reference genes were run for each DNA sample. The differences in the C<sub>t</sub> values ( $\Delta C_t$ ) between each IL6-specific qPCR and each reference gene qPCR were calculated, and then the difference between each  $\Delta C_t$  and the median  $\Delta C_t$  for all samples assayed was calculated to obtain  $\Delta\Delta C_t$  values for each sample. The copy number of the IL-6 gene in each sample was calculated from the following formula: copy number = 2 • 2<sup>- $\Delta\Delta C_t$</sup> . A correctly targeted clone, having lost one of its native copies, will have an IL-6 gene copy number equal to one. Confirmation that the human IL-6 gene sequence replaced the deleted mouse IL-6 gene sequence in the humanized allele was confirmed by a TaqMan™ qPCR assay that comprises the following primer-probe sets (written 5' to 3'): the human forward primer, CCCCCACTCCACTGGAATTG (SEQ ID NO:7); the human reverse primer, GTTCAACCACAGGCCAGGAAAG (SEQ ID NO:8); and the human probe, FAM-AGCTACAACCTATTGGCATCCTGGCAA-BHQ (SEQ ID NO:9).

**[0155]** The upstream junction of the murine locus and the sequence containing the hIL-6 gene is designed to be within 5'-AATTAGAGAG TTGACTCCTA ATAAATATGA GACTGGGGAT

GTCTGTAGCT CATTCTGCTC TGGAGCCCCAC CAAGAACGAT AGTCAATTCC AGAAACCGCT ATGAACTCCT TCTCCACAAG TAAGTGCAGG AAATCCTTAG CCCTGGAACT GCCAGCGGCG GTCGAGCCCT GTGTGAGGGGA GGGGTGTGTG GCCCAGG (SEQ ID NO:10), wherein the final mouse nucleotide prior to the first nucleotide of the human gene is the "T" in CCGCT, and the first nucleotide of the human sequence is the first "A" in ATGAA. The downstream junction of the sequence containing the hIL-6 gene and the murine locus is designed to be within 5'-TTTTAAAGAA ATATTATAT TGTATTATA TAATGTATAA ATGGTTTTA TACCAATAAA TGGCATTAA AAAAATTCAG CAACTTGAG TGTGTCACGC TCCCAGGCTC GATAACTATA ACGGTCCTAA GGTAGCGACT CGAGATAACT T-3' (SEQ ID NO:11), wherein the final nucleotide of the human sequence is with the final "G" in TCACG and the first nucleotide of the mouse sequence is the first "C" in CTCCC; the downstream junction region also contained a loxP site at the 3' end (the beginning of which is shown) for removal of a floxed ubiquitin promoter-driven neo cassette. The junction of the neo cassette with the mouse IL-6 locus is designed to be within 5'-TATACGAAGT TATCCTAGGT TGGAGCTCCT AAGTTACATC CAAACATCCT CCCCCAAATC AATAATTAAG CACTTTTAT GACATGTAAA GTTAAATAAG AAGTGAAAGC TGCAGATGGT GAGTGAGA (SEQ ID NO:12), where the final "C" of AGCTC is the final nucleotide of the neo cassette; the first nucleotide of the mouse genome following the cassette is the initial "C" of CTAAG.

**[0156]** To generate a mouse comprising hIL-6 and lacking *Rag2* and *Il2rg*, correctly targeted ES cells are identified, and are introduced into preimplantation embryo using techniques known in the art.

**[0157]** Humanized IL-6 KI mice were then backcrossed to generate mice lacking *Rag2* and *Il2rg* and expressing hIL-6, and crossed to mice expressing a human SIRPa transgene (Strowig et al., 2011, Proc Natl Acad Sci USA, 108(32): 13218-13223) to generate mice deficient for *Rag2* and *Il2rg* as well as expressing both hIL-6 and hSIRPa (*Rag2*<sup>-/-</sup> *Il2rg*<sup>null</sup> *Il6*<sup>h/h</sup> *hSIRPa*<sup>+</sup>). In addition, *Rag2*<sup>-/-</sup>, *IL-2rg*<sup>Y/-</sup>, *hIL-6* KI mice were crossed with mice expressing human TPO (Rongvaux et al., 2011, Proc Natl Acad Sci USA, 108(6): 2378-2383), human IL-3 and human GM-CSF (Willinger et al, 2011, Proc Natl Acad Sci USA, 108(6): 2390-2395), and human M-CSF (Rathinam et al, 2011, Blood, 118(11): 3119-3128) as well as hSIRPa (Strowig et al., 2011, Proc Natl Acad Sci USA, 108(32): 13218-13223) to generate mice expressing a combination of these human proteins (*Rag2*<sup>-/-</sup> *Il2rg*<sup>null</sup> *hSIRPa*<sup>+</sup> *Tpo*<sup>h/h</sup> *Mcsf*<sup>h/h</sup> *Il3/Gmcsf*<sup>h/h</sup> *Il6*<sup>h/h</sup>).

**[0158]** Cell lines and primary cells. The multiple myeloma cell line INA-6 (Burger et al., 2001, Hematol J, 2(1): 42-53) was maintained in RPMI1640 medium supplemented with 20 % FCS, penicillin/streptomycin, L-glutamine, and 2.5 ng/ml of hIL-6 in a standard incubator at 37C and 5% CO<sub>2</sub>.

**[0159]** Primary cells from multiple myeloma patients were isolated from bone marrow aspirates after obtaining informed consent from patients. Mononuclear cells were purified by Ficoll density-gradient centrifugation and subsequently, different cell subsets were isolated by

Magnetic-activated cell sorting (MACS®). To obtain, T cell-depleted populations, CD3+ cells were depleted by negative selection on an AutoMACS system using anti-CD3 microbeads (Miltenyi Biotec). To obtain CD138+ cells, CD138+ cells were isolate by positive selection on an AutoMACS® system using anti-CD 138 microbeads (Miltenyi Biotec). Purity of cells after MACS® selection was analyzed by flow cytometry.

**[0160] Transplantation of cells**. For intrafemoral transplantation of INA6 cells, Rag2<sup>-/-</sup>Il2rg<sup>null</sup> Il6<sup>h/h</sup>hSIRPa<sup>+</sup> mice were irradiated twice with 200 rad from an X-ray source. Indicated amounts of cells were then transplanted into the femur of recipient mice. Briefly, mice were anaesthetized using Ketamine/Xylazine and a hole was drilled into the patellar surface of the femur using a 26 gauge needle. The cells were then slowly injected in a volume of 20  $\mu$ l using a 27 gauge needle. For transplantation of primary patient-derived cells, Rag2<sup>-/-</sup>Il2rg<sup>null</sup>hSIRPa<sup>+</sup> Tpo<sup>h/h</sup> Mcsf<sup>h/h</sup> Il3/Gmcsf<sup>h/h</sup> Il6<sup>h/h</sup> mice were irradiated twice with 150 rad from an X-ray source and transplantation was performed as described above.

**[0161] ELISA**. Commercial ELISA kits were used to measure the concentrations of human soluble IL-6R (R&D Systems), human Igk, and Ig $\lambda$  (Bethyl Laboratories). Detection of these proteins was performed according to manufacturer's instructions.

**[0162]  $\mu$ CT**. Femur morphometry was quantified using cone-beam microfocus x-ray computed tomography ( $\mu$ CT40; ScancoMedicalAG). Serial tomographic images were acquired, and 3D images were reconstructed and used to determine the parameters. Trabecular morphometry was characterized by measuring the bone volume fraction, trabecular thickness, trabecular number, and trabecular separation. Cortical measurements included average cortical thickness, cross-sectional area of cortical bone, subperiosteal cross-sectional area, and marrow area.

**[0163] Histology**. Femurs were stripped of soft tissue, fixed in 10% buffered formalin, dehydrated, and embedded in methyl methacrylate before being sectioned and stained with toluidine blue according to standard procedures.

## **Results**

**[0164] Engraftment of multiple myeloma cell line in mice with humanized IL-6 gene**. A human IL-6 dependent MM cell line (INA6-gfp) was utilized to evaluate if mice expressing human SIRPa, and IL-6 are suitable hosts for multiple myeloma (MM) cell lines. The INA6-gfp cell line shows high dependency on human microenvironment, i.e., human fetal bone chips, when transplanted in xenograft systems of scid-hu mice (Epstein et al., 2005, Methods Mol Med, 113: 183-190). Specifically, INA-6 cells are only able to engraft the human bone graft in scid-hu mice, suggesting dependence on a human bone marrow microenvironment, similar to primary MM cells (Tassone et al., 2005, Blood, 106(2): 713-716).

**[0165]** Hence, to directly test the potential of IL-6 humanization in enabling the growth of myeloma cells, INA-6 cells were transplanted intravenously into i) Rag2<sup>-1</sup>-Il2rg<sup>null</sup>, ii) Rag2<sup>-/-</sup>-Il2rg<sup>null</sup>-hSIRPa+, iii) Rag2<sup>-/-</sup>-Il2rg<sup>null</sup> Il6<sup>h/h</sup>, and iv) Rag2<sup>-/-</sup>-Il2rg<sup>null</sup> Il6<sup>h/h</sup>-hSIRPa+ mice. Engraftment was analyzed by measuring sIL-6R protein secreted by INA-6 cells in the blood. Engraftment was only detected in mice expressing human IL-6, (Figure 1) demonstrating that INA-6 cells were indeed able to engraft mice expressing human IL-6. Next, the location of INA-6 cells (modified to express GFP) in the engrafted mice was investigated by fluorescent microscopy. Few GFP+ cells was detected in the bone marrow (Figure 2), but an increased number of GFP+ cells was detected in the lung of engrafted mice (Figure 3).

**[0166]** Analysis of human Il6 gene expression revealed that the highest level of human Il6 gene expression was found in the lung (Figure 3), hence correlating with the presence of INA-6 cells in the lung. In summary, the data disclosed herein demonstrate the successful engraftment of INA-6 cells after intravenous injection (i.v.) into mice genetically modified to express human IL6 or human IL6 and SIRPa, suggesting that genetic humanization is able to overcome growth restrictions of human MM cells in mice.

**[0167]** The data in Figure 3 suggests that INA-6 cell do not home efficiently to the bone marrow and may grow instead at non-physiological sites. Therefore, it was next examined whether transplantation of the tumor cell line in its natural microenvironment is able to reproduce the pathology typically associated with human MM. To do so, intrabone injection of INA6 cells was tested. This strikingly resulted in bone destruction and resorption, which are aspects of the pathology seen in MM patients (Figure 4-6). Specifically, the loss of trabecular bone mass was observed by histology, which was quantified by  $\mu$ CT. Moreover, only limited metastases to peripheral sites, such as the lung, was observed, which leads to the conclusion that the model may be further explored to investigate novel drugs interfering with this pathology. To test this conclusion INA-6-engrafted mice were treated with Zometa or Velcade, two drugs commonly used to treat multiple myeloma patients. Strikingly, Zometa treatment of mice injected with INA-6 cells was able to reduce bone resorption compared to untreated mice as quantified by  $\mu$ CT (Figure 7).

**[0168]** The data disclosed herein demonstrate that humanization of the Il6 gene enables engraftment of multiple myeloma cell lines that typically require a human microenvironment. Engraftment recapitulates several pathological symptoms observed in patients including bone loss. Further, these symptoms can be treated using approved drugs highlighting the utility of this model to test new drugs.

**[0169]** Genetic humanization of cytokine genes enables engraftment of primary patient-derived multiple myeloma cells. Next, experiments were conducted to examine the transplantation of primary MM cells into genetically humanized mice. It has been previously demonstrated that humanization of multiple cytokines including thrombopoietin, IL-3, GM-CSF, and M-CSF as well as the macrophage inhibitory receptor SIRPa results in improved engraftment of human hematopoietic cells in immunodeficient mice (Strowig et al., 2011, Proc Natl Acad Sci USA, 108(32): 13218-13223; Rathinam et al, 2011, Blood, 118(11): 3119-3128; Rongvaux et al.,

2011, Proc Natl Acad Sci USA, 108(6): 2378-2383; Willinger et al, 2011, Proc Natl Acad Sci USA, 108(6): 2390-2395). Specifically, humanization of IL-3 and GM-CSF as well as M-CSF improved engraftment of myeloid cells that have been demonstrated to be important for specific aspects of MM pathology. Transgenic expression of hSIRPa improves human cell engraftment, but the SIRPa-CD47 axis has also been recently implicated in tumorigenesis. Previously generated humanized mice were combined with human IL-6 knock-in mice to generate *Rug2<sup>-/-</sup>**Il2rg<sup>null</sup>**hSIRPa<sup>+</sup>* *Tpo<sup>h/h</sup>* *Mcsf<sup>h/h</sup>* *Il3/Gmcsf<sup>h/h</sup>* *Il6<sup>h/h</sup>* mice. To evaluate the ability of this strain to support human cell engraftment, CD3-depleted bone marrow cells from MM patients was injected into the bone marrow of the mice. A high frequency of myeloma cells, as identified as CD138+CD38+CD19- cells, was detected in the injected bone, while only few cells were detected in the collateral bone (Figure 8).

**[0170]** These results suggest that the genetically humanized mice, described herein, support the engraftment of primary human MM cells *in vivo*. The present data demonstrates that the humanization of cytokines gene encoding IL-6, TPO, IL-3, GM-CSF, and/or M-CSF enable engraftment of primary multiple myeloma cells from patients that typically require a human microenvironment for successful transplantation.

**Example 2: Genetic humanization of the IL-6 gene enables engraftment of mice with human hematopoietic cells.**

### **Materials and Methods**

**[0171]** Mice. The humanized IL-6 KI mouse was generated as described above. The chimeric mice were first bred with BALB/c mice and then backcrossed in order to obtain offspring with hIL-6 in homozygosity. Mice with the same mixed BALB/c x 129 background were used as control.

**[0172]** Newborn pups (within first day of life) were sublethally irradiated by X-ray irradiation in a 4-hour interval with 2 x 150cGy. Four hours after the irradiation the mice were injected with 1-2x10<sup>5</sup> CD34<sup>+</sup> fetal liver (FL) cells in 20µl of PBS into the liver by using a 30-gauge needle (Hamilton Company, NV, USA). Mice were weaned at 3 weeks of age. The mice were maintained under specific pathogen-free conditions and all experiments were performed in compliance with Yale Institutional Animal Care and Use Committee protocols.

**[0173]** Analysis of human and mouse hematological cell populations. The mice were bled from retro-orbital plexus under isofluorane anesthesia at different times after transplantation. Plasma samples were collected and stored at -20°C for further Ig measurement. Red blood cells were lysed two times by using Ammonium-Chloride (ACK) lysing buffer and the remaining PBMC were resuspended in FACS buffer (PBS supplemented with 5%FBS and 5mM EDTA).

**[0174]** When the mice were killed, a single-cell suspension of cells was obtained from the bone marrow (BM), thymus and spleen. Samples were then stained with fluorochrome-labeled mAbs against mouse and human cell surface antigens according to the manufacturers' instructions. The following anti-human mAbs were used: CD45 (HI30), CD 19 (HIB19), CD3 (UCHT1), CD4 (RPA-T4), CD8 (HIT8a), CD20 (2H7), CD5 (UCHT2), CD27 (O323). CD24 (ML5), CD10 (HI10a), all from Biolegend, CA, USA; CD33 (WM53), CD38 (HIT2), IgM (G20-127), CD138 (M115) from BD Biosciences and CD34 (AC 136) from Miltenyi Biotec. The mouse cells were stained with an anti-murine CD45 Ab (30-F11, Biolegend). Samples were acquired on the LSRII (BD Biosciences) cytometer and analyzed on FlowJo (Treestar, OR, USA) software.

**[0175]** Measurement of human immunoglobulins. Total immunoglobulins (Igs) were measured in the plasma collected from the mice by ELISA. 96 well plates (Nunc, NY, USA) were coated at 4°C overnight with 20µg/ml purified goat anti-human IgM and IgG (Southern Biotechnology, AL, USA). After washing and blocking with PBS 1% bovine serum albumin (BSA, Sigma-Aldrich) appropriate dilutions of the samples were added for 2 hours at room temperature (RT). Plates were washed and incubated for 1 hour at RT with isotype specific secondary biotinylated antibodies (Southern Biotechnology) followed by Streptavidin-HRP (Pierce Protein Research Products, IL, USA). After a final wash the enzyme activity was determined using the TMB substrate solution followed by the stop solution (both from Pierce Protein Research Products). The absorbance was measured at 450nm. A sample of human serum from Bethyl (TX, USA) was used as reference.

**[0176]** Statistical analysis. All data were expressed as average  $\pm$  standard error of mean (SEM) with the exception of the Ab levels that were plotted as geometric mean. The non-parametric Mann-Whitney U test was used to determine statistical significance between two groups. Differences were considered significant when the p values were lower than 0.05.

## Results

**[0177]** Peripheral blood engraftment in human CD34 $^+$  cell transplanted RAG2 $^{-/-}$ γ $_{c/-}$  mice. The IL6h/h mice showed a higher peripheral blood (PB) engraftment throughout the time of the analysis when compared to the IL6m/m mice and their engraftment increased over time (Figure 9). There were no major differences in the composition of the human cells between the 2 groups of mice at any time point tested (Figure 10). In both groups B and T cell percentages were similar at week 8 and week 16-20 whereas at weeks 11-15 there was a higher percentage of B cells ( $69.23 \pm 3.97$  in IL6m/m and  $55.91 \pm 4.86$  in IL6h/h) than T cells ( $16.86 \pm 3.14$  in IL6m/m and  $30.26 \pm 6.23$  in IL6h/h). Myeloid CD33 $^+$  cells represented a minor component of the human cell and their percentage decreased over time.

**[0178]** Organ cell engraftment and composition of human CD34 $^+$  cell reconstituted RAG2 $^{-/-}$ γ $_{c/-}$  mice. Cells of human origin were found in the hemato-lymphoid organs, such as BM, spleen and thymus (Figure 11A). Interestingly the spleen of the IL6h/h mice displayed a greater

human engraftment than IL6m/m mice ( $61.75\% \pm 10.87$  versus  $23.56\% \pm 5.2$ ). These data were confirmed by the doubling in the absolute number of human cells ( $14.21 \times 10^6 \pm 1.38$  versus  $7.26 \times 10^6 \pm 0.66$ ) (Figure 11B).

**[0179]** B cell maturation in human CD34 $^{\pm}$  engrafted RAG2 $^{-/-}$ Yc $^{-/-}$  mice. The maturation stages of the human B cells were studied in the BM and spleen of the engrafted mice using the gating strategies illustrated in Figure 12A, based on the usage of a combination of CD24, CD38 and surface IgM antibodies. This strategy is particularly helpful in dissecting the presence of the transitional B cell subset. In BM the main compartment was made by the Pro-/Pre-B cells (Figure 12B) with no difference between the IL6m/m and the IL6h/h mice ( $76.04\% \pm 9.09$  and  $79.07\% \pm 3.43$  respectively). The spleen of both groups contained some mature B cells (~20%) but still a high percentage of immature/transitional cells ( $27.13\% \pm 8.99$  and  $36.45\% \pm 5.12$ ).

**[0180]** A significant percentage of B cells in the spleen were CD5 $^{+}$  (Figure 13B), a marker not commonly expressed on human BM and peripheral B cells but found in low percentage on the FL B cells (Figure 13A).

**[0181]** The IL6h/h mice showed a sharp increase in the percentage of the CD27 $^{+}$  B cells in the spleen when compared to the IL6m/m mice ( $24.73\% \pm 8.94$  versus  $9.46\% \pm 2.32$ ) but very few CD27 $^{+}$  cells were found in the BM (Figure 14A and 14B).

**[0182]** Antibody production in human CD34 $^{\pm}$  engrafted RAG2 $^{-/-}$ Yc $^{-/-}$  mice. As the B cells were found in the engrafted mice, the concentration of the human IgM and IgG in the plasma collected at 12 and 20 weeks after the human cell transplantation was then measured. Both IL6m/m and IL6h/h mice secreted human IgM and IgG (Figure 15).

**[0183]** In general there was a higher percentage of IL6h/h mice secreting antibodies compared to the IL6m/m ones with the former mice showing a smaller level of IgM level ( $14.69\mu\text{g/ml}$  versus  $33.66\mu\text{g/ml}$  at week 12 and  $18.25\mu\text{g/ml}$  versus  $1902\mu\text{g/ml}$  at week 20) but an increased level of IgG ( $243\mu\text{g/ml}$  versus  $496\mu\text{g/ml}$  at week 12 and  $553.6\mu\text{g/ml}$  versus  $2972\mu\text{g/ml}$  at week 20) (Figure 15A). Both IgM and IgG average level rose over time in IL6m/m mice (Figure 15B). Conversely the serum Igs remained steady in IL6h/h mice.

**[0184]** While this invention has been disclosed with reference to specific embodiments, it is apparent that other embodiments and variations of this invention may be devised by others skilled in the art without departing from the scope of the invention, as defined by the appended claims.

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

#### Patent documents cited in the description

- US61722437B [0001]
- US20130042330A [0064]
- US8541646B [0064] [0064]
- WO2012040207A [0064]
- US6586251B [0086]
- US7576259B [0088]
- US7659442B [0088]
- US7294754B [0088]
- US20110200982A [0113] [0113]
- US5222982A [0128]
- US5385582A [0128]
- US20070254842A [0128]
- US20080081064A [0128]
- US20090196903A [0128]

#### Non-patent literature cited in the description

- **HIDESHIMA et al.** *Nat Rev Cancer*, 2007, vol. 7, 585-98 [0003] [0003] [0004]
- **KUEHLBERGSAGEL** *Nat Rev Cancer*., 2002, vol. 2, 175-87 [0003]
- **KUEHLBERGSAGEL** *Nat Rev Cancer*, 2002, vol. 2, 175-87 [0003] [0003]
- **LANDGREN et al.** *Blood*, 2009, vol. 113, 5412-7 [0003]
- **ZHAN et al.** *Blood*, 2006, vol. 108, 2020-8 [0003]
- **CHNG et al.** *Blood*, 2005, vol. 106, 2156-61 [0003]
- **FONSECA et al.** *Blood*, 2002, vol. 100, 1417-24 [0003]
- **KAUFMANN et al.** *Leukemia*, 2004, vol. 18, 1879-82 [0003]
- **BERGSAGEL et al.** *Blood*, 2005, vol. 106, 296-303 [0003]
- **DE RAEVEVANDERKERKEN** *Histol Histopathol.*, 2005, vol. 20, 1227-50 [0004] [0004] [0004] [0004]
- **DHODAPKARA** *Am J Hematol.*, 2009, vol. 84, 395-6 [0004] [0004]
- **KUKREJA et al.** *J Exp Med.*, 2006, vol. 203, 1859-65 [0004]
- **YACCOBY et al.** *Blood*, 1998, vol. 92, 2908-13 [0006] [0006]

- **YACCOBYEPSTEIN**Blood, 1999, vol. 94, 3576-82 [0006]
- **DEWAN et al.**Cancer Sci., 2004, vol. 95, 564-8 [0006]
- **MIYAKAWA et al.**Biochem Biophys Res Commun., 2004, vol. 313, 258-62 [0006]
- **J. SAMBROOKD. W. RUSSELL**Molecular Cloning: A Laboratory ManualCold Spring Harbor Laboratory Press20120000 [0025]
- Short Protocols in Molecular Biology, Current Protocols20020000 [0025]
- **B. ALBERTS et al.**Molecular Biology of the Cell20020000 [0025]
- **D. L. NELSONM. M. COX**Lehninger Principles of BiochemistryW.H. Freeman & Company20040000 [0025]
- Oligonucleotide Synthesis: Methods and ApplicationsMethods in Molecular BiologyHumana Press20040000 [0025]
- **HARLOW et al.**Using Antibodies: A Laboratory ManualCold Spring Harbor Laboratory Press19990000 [0030]
- **HARLOW et al.**Antibodies: A Laboratory ManualCold Spring Harbor19890000 [0030]
- **HOUSTON et al.**Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 5879-5883 [0030]
- **BIRD et al.**Science, 1988, vol. 242, 423-426 [0030]
- **RATHINAM et al.**Blood, 2011, vol. 118, 3119-28 [0064] [0086]
- **WILLINGER et al.**Proc Natl Acad Sci USA, 2011, vol. 108, 2390-2395 [0064] [0086]
- **RONGVAUX et al.**Proc Natl Acad Sci USA, 2011, vol. 108, 2378-83 [0064] [0086]
- **STROWIG et al.**Proc Natl Acad Sci USA, 2011, vol. 108, 3213218-13223 [0064] [0157] [0157] [0169]
- Goeddel, Gene Expression Technology: Methods in EnzymologyAcademic Press19900000vol. 185, [0065]
- **ABBOUD et al.**J. Histochem & Cytochem., 2003, vol. 51, 941-949 [0067]
- **SCHORPP et al.**NAR, 1996, vol. 24, 1787-1788 [0067]
- **MCBURNEY et al.**Devel. Dynamics, 1994, vol. 200, 278-293 [0067]
- **MAJUMDER et al.**Blood, 1996, vol. 87, 3203-3211 [0067]
- **BOSMA et al.**Immunogenetics, 1989, vol. 29, 54-56 [0076]
- **FESTING et al.**Mammalian Genome, 1999, vol. 10, 836- [0080]
- **AUERBACH et al.**Establishment and Chimera Analysis of 129/SvEv- and C57BL/6-Derived Mouse Embryonic Stem Cell Lines, 2000, [0080]
- **SUNDBERGICHIKI**Genetically Engineered Mice HandbookCRC Press20060000 [0086]
- **HOKERVAN DEURSEN**Genetically modified Mouse Methods and ProtocolsHumana Press20020000 [0086]
- **JOYNER**Gene Targeting: A Practical ApproachOxford University Press20000000 [0086]
- **TURKSEN**Embryonic stem cells: Methods and Protocols in Methods Mol Biol.Humana Press20020000 [0086]
- **MEYER et al.**Proc. Nat. Acad. Sci. USA, 2010, vol. 107, 15022-15026 [0086]
- **GIBSON**Primer Of Genome ScienceSinauer20040000 [0086]
- **VALENZUELA et al.**Nat Biot, 2003, vol. 21, 652-659 [0086]
- Manipulating the Mouse EmbryoA Laboratory ManualCold Spring Harbor Laboratory Press20020000 [0087]
- **NAGY et al.**Manipulating the Mouse Embryo: A Laboratory ManualCold Spring Harbor Laboratory Press20020000 [0088]

- **NAGY et al.** Development, 1990, vol. 110, 815-821 [0088]
- **KRAUS et al.** Genesis, 2010, vol. 48, 394-399 [0088]
- **PEARSON et al.** Curr. Protoc. Immunol., 2008, vol. 81, 1-15 [0099]
- **ITO et al.** Blood, 2002, vol. 100, 3175-3182 [0099]
- **TRAGGIAI et al.** Science, 2004, vol. 304, 104-107 [0099]
- **ISHIKAWA et al.** Blood, 2005, vol. 106, 1565-1573 [0099]
- **SHULTZ et al.** J. Immunol., 2005, vol. 174, 6477-6489 [0099]
- **HOLYOAKE et al.** Exp Hematol., 1999, vol. 27, 1418-27 [0099]
- **J. HAYAKAWA et al.** Stem Cells, 2009, vol. 27, 1175-182 [0103]
- **WATANABE** Bone Marrow Transplantation, 1997, vol. 19, 1175-1181 [0106]
- **GOODMANGILMAN** The Pharmacological Basis of Therapeutics McGraw-Hill 19960000 [0118]
- Chemical Warfare Agents Academic Press 19920000 [0118]
- **PEAR et al.** P.N.A.S., 1993, vol. 90, 8392-8396 [0120]
- **MILLER et al.** Mol. Cell. Biol., 1985, vol. 5, 431-437 [0120]
- **MILLER et al.** Mol. Cell. Biol., 1986, vol. 6, 2895-2902 [0120]
- **DANOS et al.** PNAS, 1988, vol. 85, 6460-6464 [0120]
- Current Protocols in Immunology John Wiley & Sons, Inc. 20120000 [0132]
- Immunology Methods Manual Academic Press 19970000 [0132]
- Optimization of monoclonal antibody purification NAU Techniques in Protein Chemistry Academic Press 19890000 [0143]
- **COLIGAN et al.** Current Protocols in Immunology John Wiley & Sons, Inc. 20050000 [0143]
- **VALENZUELA et al.** High-throughput engineering of the mouse genome coupled with high-resolution expression analysis Nature Biotech, 2003, vol. 21, 6652-659 [0153]
- **RONGVAUX et al.** Proc Natl Acad Sci USA, 2011, vol. 108, 62378-2383 [0157] [0169]
- **WILLINGER et al.** Proc Natl Acad Sci USA, 2011, vol. 108, 62390-2395 [0157] [0169]
- **RATHINAM et al.** Blood, 2011, vol. 118, 113119-3128 [0157] [0169]
- **BURGER et al.** Hematol J, 2001, vol. 2, 142-53 [0158]
- **EPSTEIN et al.** Methods Mol Med, 2005, vol. 113, 183-190 [0164]
- **TASSONE et al.** Blood, 2005, vol. 106, 2713-716 [0164]

**Patentkrav**

**1.** Immundeficient, genmodificeret gnaver, der omfatter:

en nukleinsyre, der koder for humant IL-6, operativt forbundet med en IL-6-promoter, hvor gnaveren eksprimerer et humant IL-6-polypeptid og ikke eksprimerer et funktionelt nativt IL-6-polypeptid, og mindst en yderligere nukleinsyre, der er valgt fra gruppen bestående af:

i) en nukleinsyre, der koder for humant SIRPa, operativt forbundet med en SIRPa-promoter, hvor gnaveren eksprimerer et humant SIRPa-polypeptid;

ii) en nukleinsyre, der koder for humant M-CSF, operativt forbundet med en M-CSF-promoter, hvor gnaveren eksprimerer et humant M-CSF-polypeptid;

iii) en nukleinsyre, der koder for humant IL-3, operativt forbundet med en IL-3-promoter, hvor gnaveren eksprimerer et humant IL-3-polypeptid;

iv) en nukleinsyre, der koder for humant GM-CSF, operativt forbundet med en GM-CSF-promoter, hvor gnaveren eksprimerer et humant GM-CSF-polypeptid; og

v) en nukleinsyre, der koder for humant TPO, operativt forbundet med en TPO-promoter, hvor gnaveren eksprimerer et humant TPO-polypeptid, og hvor gnaveren omfatter en indpodning af humane hæmatopoietiske celler og producerer humane B-celler.

**2.** Genmodificeret gnaver ifølge krav 1, hvor IL-6-promoteren er gnaver-IL-6-promoteren, og nukleinsyren, der koder for humant IL-6, er operativt forbundet med gnaver-IL-6-promoteren ved gnaver-IL-6-locuset.

**3.** Genmodificeret gnaver ifølge krav 1 eller 2, hvor SIRPa-promoteren er gnaver-SIRPa-promoteren, M-CSF-promoteren er gnaver-M-CSF-promoteren, IL-3-promoteren er gnaver-IL-3-promoteren, GM-CSF-promoteren er gnaver-GM-CSF-promoteren, og TPO-promoteren er gnaver-TPO-promoteren.

**4.** Genmodificeret gnaver ifølge et af kravene 1-3, hvor gnaveren ikke eksprimerer et rekombinationsaktiverende gen (RAG), IL2-receptor-gamma-kæde (IL2rg) eller både et RAG og IL2rg.

5. Genmodificeret gnaver ifølge krav 4, hvor gnaveren ikke eksprimerer hverken et rekombinationsaktiviserende gen 2 (RAG2) eller et IL2rg.

5 6. Genmodificeret gnaver ifølge et af kravene 1-5, hvor de humane hämatopoietiske celler er CD34+-celler.

10 7. Genmodificeret gnaver ifølge et af kravene 1-6, hvor gnaveren er en mus.

10 8. Fremgangsmåde til fremstilling af humane antistoffer, hvilken fremgangsmåde omfatter:  
at bringe den immundeficiente, genmodificerede gnaver ifølge et af kravene 1-6 i kontakt med et antigen; og  
at indsamle en human B-celle, der producerer et humant antistof mod antigenet, fra gnaveren.

15 9. Fremgangsmåde ifølge krav 8, omfattende at isolere antistoffet fra den humane celle, der producerer det humane antistof.

20 10. Fremgangsmåde ifølge krav 8 eller 9, hvor indsamlingen omfatter at indsamle mindst en milt, lymfeknuder, perifert blod, knoglemarv eller dele deraf fra gnaveren.

25 11. Fremgangsmåde ifølge et af kravene 8-10, omfattende at producere en hybridomcellelinje fra den humane B-celle.

12. Fremgangsmåde ifølge krav 11, omfattende at isolere et humant antistof, der specifikt binder antigenet, fra hybridomcellelinjen.

30 13. Fremgangsmåde ifølge et af kravene 8-12, hvor antigenet er mindst et af et peptid, et MHC/peptid-kompleks, et DNA, en levende virus, en død virus eller en del deraf, en levende bakterie, en død bakterie eller en del deraf og en cancercelle eller en del deraf.

35 14. Fremgangsmåde ifølge et af kravene 8-13, hvor antigenet administreres på den podede, genmodificerede, immundeficiente mus ca. 1-5 måneder efter

indpodning af de humane hæmatopoietiske celler.

5      **15.** Fremgangsmåde ifølge et af kravene 8-14, hvor de humane hæmatopoietiske celler er CD34+-celler, der opnås fra en human føtal lever, knoglemarv, navlestrengsblod, perifert blod eller en miltprøve.

## DRAWINGS




Figure 1




Figure 2




Figure 3





Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12





Figure 13






Figure 15