



(86) Date de dépôt PCT/PCT Filing Date: 2011/12/14

(87) Date publication PCT/PCT Publication Date: 2012/06/21

(45) Date de délivrance/Issue Date: 2021/03/23

(85) Entrée phase nationale/National Entry: 2013/05/23

(86) N° demande PCT/PCT Application No.: GB 2011/052473

(87) N° publication PCT/PCT Publication No.: 2012/080727

(30) Priorités/Priorities: 2010/12/14 (GB1021161.3);
2011/06/01 (GB1109162.6)

(51) Cl.Int./Int.Cl. *A61K 31/437* (2006.01),
A61K 31/404 (2006.01), *A61P 25/28* (2006.01)

(72) Inventeurs/Inventors:

SHERIDAN, JOSEPH M., GB;
HEAL, JONATHAN R., GB;
HAMILTON, WILLIAM D.O., GB;
PIKE, IAN, GB


(73) Propriétaire/Owner:

ELECTROPHORETICS LIMITED, GB

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : 5-(1,3-BENZOXAZOL-2-YL)-4-(PYRIDINE-4-YL)PYRIMIDINE-2-AMINE ET SON UTILISATION COMME INHIBITEUR DE CASEINE KINASE 1 DELTA

(54) Title: 5-(1,3-BENZOXAZOL-2-YL)-4-(PYRIDIN-4-YL)PYRIMIDIN-2-AMINE AND ITS USE AS A CASEIN KINASE 1DELTA INHIBITOR



(57) Abrégé/Abstract:

The invention relates to pharmaceutical compositions comprising casein kinase 1 delta (CK1δ) and to the use of said inhibitors in the treatment of neurodegenerative disorders such as Alzheimer's disease.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2012/080727 A4

(43) International Publication Date
21 June 2012 (21.06.2012)

(51) International Patent Classification:

A61K 31/00 (2006.01) *A61K 31/506* (2006.01)
A61K 31/437 (2006.01) *A61P 25/28* (2006.01)
A61K 31/4439 (2006.01)

(21) International Application Number:

PCT/GB2011/052473

(22) International Filing Date:

14 December 2011 (14.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1021161.3 14 December 2010 (14.12.2010) GB
1109162.6 1 June 2011 (01.06.2011) GB

(71) Applicant (for all designated States except US): **ELECTROPHORETICS LIMITED** [GB/GB]; Coveham House, Downside Bridge Road, Cobham, Surrey KT11 3EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SHERIDAN, Joseph M.** [GB/GB]; c/o Electrophoretics Limited, Coveham House, Downside Bridge Road, Cobham, Surrey KT11 3EP (GB). **HEAL, Jonathan R.** [GB/GB]; c/o Electrophoretics Limited, Coveham House, Downside Bridge Road, Cobham, Surrey KT11 3EP (GB). **HAMILTON, William D.O.** [GB/GB]; c/o Electrophoretics Limited, Coveham House, Downside Bridge Road, Cobham, Surrey KT11 3EP (GB). **PIKE, Ian** [GB/GB]; c/o Electrophoretics Limited, Coveham House, Downside Bridge Road, Cobham, Surrey KT11 3EP (GB).

(74) Agents: **GIBSON, Mark** et al.; Sagittarius IP, Taylor House, 39 High Street, Marlow, Buckinghamshire SL7 1AF (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))

(88) Date of publication of the international search report:

23 August 2012

Date of publication of the amended claims: 26 October 2012

(54) Title: CASEIN KINASE 1DELTA (CK1DELTA) INHIBITORS



(57) Abstract: The invention relates to pharmaceutical compositions comprising casein kinase 1 delta (CK1δ) and to the use of said inhibitors in the treatment of neurodegenerative disorders such as Alzheimer's disease.

**5-(1,3-BENZOXAZOL-2-YL)-4-(PYRIDIN-4-YL)PYRIMIDIN-2-AMINE AND ITS USE AS A
CASEIN KINASE 1 DELTA INHIBITOR**

The invention relates to pharmaceutical compositions comprising casein kinase 1 delta (CK1δ) inhibitors and to the use of said inhibitors in the treatment of neurodegenerative disorders such as Alzheimer's disease.

Alzheimer's disease (AD, also known as senile dementia of the Alzheimer type (SDAT), primary degenerative dementia of the Alzheimer's type (PDDAT), or Alzheimer's) is the most common form of dementia. Most often, Alzheimer's disease is diagnosed in people over 65 years of age, although the less-prevalent early-onset Alzheimer's can occur much earlier. In 2006, there were 26.6 million sufferers worldwide. Alzheimer's is predicted to affect 1 in 85 people globally by 2050.

Alzheimer's disease is a neurodegenerative disease characterised by the presence of senile plaques and neurofibrillary tangles in the brain. The degree of dementia at death correlates better with neurofibrillary tangle numbers than with senile plaques counts. The presence of neurofibrillary tangles in neurons results in the death of those neurons, implying that prevention of tangle formation is an important therapeutic goal. The principal protein that forms the neurofibrillary tangle is the microtubule-associated protein, tau, which assembles into filaments that have the appearance of twisting about each other in pairs and are referred to as paired helical filaments (PHF). PHF are present in different locations in degenerating neurons in the Alzheimer brain and when many aggregate in the neuronal cell body, they produce the neurofibrillary tangle (Lee et al, Neurodegenerative Tauopathies, 2001, Annu Rev Neurosci. 24: 1121-59)

Intraneuronal deposits of tau in the form of typical neurofibrillary tangles of AD or other morphologically distinct tau aggregates in a number of other neurodegenerative diseases, is the basis for grouping these conditions as tauopathies. Thus, in addition to AD, the main examples of the tauopathies are frontotemporal dementia with Parkinsonism linked to chromosome 17 (FTDP-17), progressive supranuclear palsy (PSP), Pick's disease, corticobasal degeneration, and multisystem atrophy (MSA). The intracellular tau deposits (usually neuronal but can also be glial) are all filamentous and mostly in a hyperphosphorylated state compared to the level of phosphorylation of tau from control human brain. In the case of AD, this hyperphosphorylated tau is often referred to as PHF-tau because it is derived from the PHF.

Tau is a phosphoprotein, the function of phosphorylation remaining to be unequivocally


established. However, increased phosphorylation of tau on multiple serine and threonine residues reduces the ability of tau to promote microtubule assembly and to stabilise assembled microtubules, effects that have been demonstrated both *in vitro* and in cells. Many studies have shown that PHF-tau from AD brain is more heavily phosphorylated on serine and threonine than tau from control brain. This has been demonstrated partly by protein sequencing and partly by demonstrating that certain monoclonal antibodies only label either PHF-tau or non-phosphorylated tau and not PHF-tau; the epitopes for many of these antibodies have been mapped to particular phosphorylated residues present in PHF-tau and absent from control brain tau. The pathological tau from most other cases of other tauopathies seems to be similarly hyperphosphorylated to PHF-tau.

These findings strongly imply that similar abnormalities in regulating phosphorylation of tau are shared by all the tauopathies including AD.

A number of proline-directed and non-proline directed protein kinases have been suggested to have a role in the generation of PHF-tau in Alzheimer brain, including casein kinase 1. Mammalian casein kinase-1 exists as multiple isoforms CK1 α , CK1 β , CK1 γ 1, CK1 γ 2, CK1 γ 3, CK1 δ and CK1 ϵ . The role of CK1 δ as a potential tau kinase is of particular interest since it has been reported that CK1 δ protein is increased more than 30-fold in the hippocampus of Alzheimer brain compared to equivalent controls (Ghoshal, N. *et al* (1999) *Am. J. Pathol* 155, 1163-1172) while its mRNA content is increased 24-fold (Yasojima, K. *et al* (2000) *Brain Res* 865, 116-120) and CK1 has also been shown to be tightly associated with PHF (Kuret, J. *et al* (1997) *J. Neurochem* 69, 2506-2515). CK1 δ has also been reported to phosphorylate tau at two epitopes detecting using phospho-specific monoclonal antibodies to tau, and exogenous expression of CK1 δ in non-neuronal cells reduces binding of tau to microtubules (Li, G. *et al* (2004) *J. Biol. Chem.* 279, 15938-15945). Of note in the context of Alzheimer's disease is a report that CK1 activity is stimulated by amyloid beta-peptide (A β), a component of the senile neuritic plaques that, together with tangles, characterise Alzheimer brain (Chauhan, A. *et al* (1993) *Brain Res.* 629, 47-52). Additional evidence for possible involvement of CK1 in Alzheimer's disease comes from the reported influence of CK1 in the regulation of A β production in neurons (Flajolet, M. *et al* (2007) *PNAS USA* 104, 4159-4164). Further work has confirmed that at least 6 newly identified phosphorylation sites in PHF-tau (all on serine or threonine residues) can be generated by CK1 δ . The finding that a number of phosphorylation sites in PHF-tau for which CK1 is a strong candidate kinase, including three for which it is the only known kinase, implies that CK1 may make an important

contribution to the pathogenesis of Alzheimer's disease (Hanger *et al* (2007) J. Biol. Chem. 282, 23645-23654).

- There is therefore a need for CK1δ inhibitors which may be of potential therapeutic benefit in the treatment of neurodegenerative diseases, such as tauopathies including Alzheimer's disease, frontotemporal dementia with Parkinsonism linked to chromosome 17 (FTDP-17), progressive supranuclear palsy (PSP), Pick's disease, corticobasal degeneration, and multisystem atrophy (MSA).
- 10 According to a first aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (IB) or a pharmaceutically acceptable salt or solvate thereof:



(IB)

- 15 wherein

"Het B" represents a 5 membered heterocyclic ring system containing 1 to 3 heteroatoms selected from O, N or S, wherein said ring system is fused to one or more (e.g. 1-3) further rings to form a polycyclic ring system comprising up to 4 rings;


- Z represents a bond, $-C(R^{7b})(R^{8b})-$, $(CH_2)_2$, $-O-$, $-S-$, $-CH_2-O-$, $-(CH_2)_2-O-$, NR^{6b} , $-N(R^{6b})-$, $C(R^{7b})(R^{8b})-$, $-N(R^{6b})-(CH_2)_2-$, $-N(R^{6b})-(CH_2)_3-$, $-CH_2-N(R^{6b})-(CH_2)_2-$, $-N(R^{6b})-CO-$, $-CH_2-NH-$, $CO-(CH_2)_2-$, $-N(R^{6b})-CO-CH_2-$, $=N-$, $-N(R^{7b})-CH=$, $-C(H)(CN)-$, $-C(=N-NH-COC_{1-6} \text{ alkyl})-$, $-CH=C(R^{6b})-CO-$, $=CH-$, $-N=CH-$, $-N=C(Me)-$, $-C(R^{6b})=CH-$, $-NH-CO-C(=CH-\text{heteroaryl})-$, $-C(=C(R^{7b})(R^{8b}))-$, $-CH=CH-CO-N(R^{6b})-$, $-CH=C(R^{6b})-CO-NH-CH_2-$, $-CH=C(R^{6b})-NH-CO-$, $-CH=C(R^{6b})-CO-O-CH_2-$, $-CS-S-CH_2-$, $-NH-CS-NH-$, $-NH-CS-NH-CH_2-$, $-NH-CS-NH-(CH_2)_2-$, $-CH_2-N(CSNH_2)-CH_2-$, $-S-C(R^{5b})(R^{6b})-$, $-S-(CH_2)_2-O-$, SO_2 , $-NH-SO_2-$, $-CH_2-NH-SO_2-$, CO , $-CH_2-CO-$, $-(CH_2)_2-CO-$, $-O-CH_2-CO-$, $-(CH_2)_2-CO-$, COO , $-COO-C(R^{7b})CO-$, $-CH=C(R^{5b})-CONH-CH_2-$, $-CO-CH_2-N(R^{6b})-CO-$, $-CO-CH_2-C(R^{6b})-CH_2-CO-$, $-CO-CH_2-N(R^{6b})-CH_2-$, $-CO-NH-N=C(R^{7b})-$, $-S-CH_2-CO-$, $-S-CH_2-CO-N(R^{6b})-$, $-S-CH_2-CO-N(R^{6b})-CH_2-$, $-SO_2-N(R^{6b})-C(R^{7b})(R^{8b})-CONH-$, $-SO_2-N(R^{6b})-CH-(CH_2-\text{aryl})-CONH-CH_2-$, $-CH(-S-C_{1-6} \text{ alkyl})-C(Me)(OH)-$, $-CH_2-C(R^{6b})(OH)-$, $-C(OH)(CH(Me)(C_{3-8} \text{ cycloalkyl}))-(CH_2)-$, $-C(OH)(R^{6b})-CH_2-$, $-CH(Me)-NH-CO-CH_2-$, $-CO-N(R^{6b})-CH_2-$, $-C(H)(R^{6b})-CO-N(R^{5b})-CH_2-$, $-CO-N(R^{6b})-CH_2-$

- CH₂-, -CO-N(R^{6b})-CH₂-CH₂-CO-NH-CH₂-, -CO-NH-C(-CONH₂)=CH-, -CO-NH-CH(-CONH₂)-CH₂-, -CH₂-C(H)(Me)-CH₂-S-, -O-CH₂-CO-NH-, -CH₂-N(R^{6b})-CO-CH₂-O-, -N(R^{6b})-CO-CH₂-O-, -C(H)(-CH₂-aryl)-, -C(H)(-CH₂-heteroaryl)-, -C(NH-aryl)=N-N=CH-, -C(NH-aryl)=N-N=CH-, -NH-CO-CH₂-N(R^{6b})-, -NH-N=C(-aryl)-, -NH-N=C(-aryl)-CO-, -NH-C(=N-CO-C₁₋₆ alkyl)-NH-(CH₂)₂-, -C(-NH-aryl)=N-N=CH-, -NH-C(-NH-aryl)=N-CONH-, -C(=CH-aryl)-CONH-CH₂-, -CH=C(R^{6b})-CONH-, -CH(-CH₂-aryl)-NH-CO- or -CH(OH)-, wherein said aryl or heteroaryl groups of Z may be optionally substituted by one or more halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, NO₂ or hydroxyl groups;
- R^{5b} represents hydrogen, C₁₋₆ alkyl or cyano;
- 10 R^{6b} represents hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, cyano, COOH, -COOC₁₋₆ alkyl, C₃₋₈ cycloalkyl, -CH₂-C₃₋₈ cycloalkyl, aryl, heteroaryl, -C₁₋₆ alkylene-aryl, -CO-aryl, -O-CO-heteroaryl, -CO-heteroaryl or -C(R^{7b})(R^{8b})-heteroaryl, wherein said aryl groups of R^{6b} may be optionally substituted by one or more halogen or C₁₋₆ alkoxy groups;
- R^{7b} and R^{8b} independently represent hydrogen or C₁₋₆ alkyl;
- 15 R^{1b} represents aryl, C₃₋₈ cycloalkyl, monocyclic or bicyclic heterocyclyl or a monocyclic or bicyclic heteroaryl ring system, wherein R^{1b} may be substituted by one or more (e.g. 1, 2 or 3) R^{4b} groups;
- R^{4b} represents halogen, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₃₋₈ cycloalkyl, haloC₁₋₆ alkyl, hydroxyl, C₁₋₆ alkoxy, -O-C₁₋₆ alkenyl, haloC₁₋₆ alkoxy, -COOH, -CO-C₁₋₆ alkyl, -COO-C₁₋₆ alkyl, -CONH₂, -CH₂-CONH₂, -NH-C₁₋₆ alkyl, -NH-C₂₋₆ alkenyl, -NH-CO-C₁₋₆ alkyl, -CO-NH-C₁₋₆ alkyl, -O-CH₂-CO-NH-C₁₋₆ alkyl, -CH₂-CH₂-CO-NH-C₁₋₆ alkyl, -S-C₁₋₆ alkyl, -SO-C₁₋₆ alkyl, -SO₂-C₁₋₆ alkyl, -SO₂-NH₂, -SO₂-NH-C₁₋₆ alkyl, -S-CH₂-CO-C₂₋₆ alkenyl, -SO₂-OH, amino, cyano, NO₂, =O, -CO-NH-(CH₂)₂-OMe, -NH-C₃₋₈ cycloalkyl, -CH₂-CO-NH-C₃₋₈ cycloalkyl, -CO-heterocyclyl, -CO-heteroaryl, -COO-(CH₂)₂-heterocyclyl, -CH₂-aryl, -OCH₂-aryl, -OCH₂-heteroaryl, -CH₂-O-CO-aryl, -O-aryl, -NH-CO-aryl, -NH-CO-heteroaryl, -NH-CO-CH₂-aryl, -NH-aryl, aryl or heteroaryl groups, wherein said aryl, heterocyclyl or heteroaryl groups of R^{4b} may be optionally substituted by one or more halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, =S or hydroxyl groups and wherein said C₁₋₆ alkyl or C₂₋₆ alkenyl groups of R^{4b} may be optionally substituted by one or more hydroxyl, amino, cyano, C₁₋₆ alkoxy, CONH₂ or -COO-C₁₋₆ alkyl groups;
- 30 m represents an integer from 0 to 3;
- R^{2b} represents halogen, haloC₁₋₆ alkyl, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, hydroxyl, C₁₋₆ alkoxy, -S-C₁₋₆ alkyl, -CH₂-S-C₁₋₆ alkyl, -S-C₂₋₆ alkynyl, amino, cyano, NO₂, =O, =S, -SO₂-C₁₋₆ alkyl, -CONH₂, -CO-C₁₋₆ alkyl, -COO-C₁₋₆ alkyl, -NH-C₁₋₆ alkyl, -NH-CO-C₁₋₆ alkyl, -NH-CO-
- 35 CH=CH-CH₂-N(Me)₂, C₁₋₆ alkyl, -CO-NH-C₁₋₆ alkyl, -CO-NH-CH(Me)-COOH, -S-CH₂-CO-N(Et)₂, -NH-(CH₂)₂-OH, -NH-(CH₂)₃-OH, -NH-CH(Et)-CH₂-OH, -CO-NH-(CH₂)₃-OH, -

CH(CH₂OH)₂ or -S-CH₂-CO-NH-CO-NH-C₁₋₆ alkyl, wherein said C₁₋₆ alkyl groups of R^{2b} may be optionally substituted by one or more cyano or hydroxyl groups; with the proviso that the compound is other than compound number 54, 373, 458, 496, 585, 590, 594, 596-597, 601-602, 649, 703, 778, 877, 891, 910, 912, 926 and 962-963.

5

According to one particular aspect of the invention which may be mentioned there is provided a compound of formula (IB) or a pharmaceutically acceptable salt or solvate thereof:



(IB)

10

wherein

“Het B” represents a 5 membered heterocyclic ring system containing 1 to 3 heteroatoms selected from O, N or S, wherein said ring system is fused to one or more (e.g. 1-3) further rings to form a polycyclic ring system comprising up to 4 rings;

- 15 Z represents a bond, -C(R^{7b})(R^{8b})-, (CH₂)₂-, -O-, -S-, -CH₂-O-, -(CH₂)₂-O-, NR^{6b}-, -N(R^{6b})-C(R^{7b})(R^{8b})-, -N(R^{6b})-(CH₂)₂-, -N(R^{6b})-(CH₂)₃-, -CH₂-N(R^{6b})-(CH₂)₂-, -N(R^{6b})-CO-, -CH₂-NH-CO-(CH₂)₂-, -N(R^{6b})-CO-CH₂-, =N-, -C(H)(CN)-, -C(=N-NH-COC₁₋₆ alkyl)-, -CH=C(R^{6b})-CO-, =CH-, -N=CH-, -N=C(Me)-, -C(R^{6b})=CH-, -NH-CO-C(=CH-heteroaryl)-, -C=C(Me)₂-, -CH=CH-CO-N(R^{6b})-, -CH=C(R^{6b})-CO-NH-CH₂-, -CH=C(R^{6b})-NH-CO-, -CH=C(R^{6b})-CO-O-CH₂-, -CS-S-CH₂-, -NH-CS-NH-, -NH-CS-NH-CH₂-, -NH-CS-NH-(CH₂)₂-, -CH₂-N(CSNH₂)-CH₂-, -S-C(R^{5b})(R^{6b})-, -S-(CH₂)₂-O-, SO₂-, -NH-SO₂-, -CH₂-NH-SO₂-, CO-, -CH₂-CO-, -(CH₂)₂-CO-, -O-CH₂-CO-, -(CH₂)₂-CO-, COO-, -COO-C(R^{7b})CO-, -CH=C(R^{5b})-CONH-CH₂-, -CO-CH₂-N(R^{6b})-CO-, -CO-CH₂-C(R^{6b})-CH₂-CO-, -CO-CH₂-N(R^{6b})-CH₂-, -CO-NH-N=C(R^{7b})-, -S-CH₂-CO-, -S-CH₂-CO-N(R^{6b})-, -S-CH₂-CO-N(R^{6b})-CH₂-, -SO₂-N(R^{6b})-C(R^{7b})(R^{8b})-CONH-, -SO₂-N(R^{6b})-CH(-CH₂-aryl)-CONH-CH₂-, -CH(-S-C₁₋₆ alkyl)-C(Me)(OH)-, -CH₂-C(R^{6b})(OH)-, -C(OH)(CH(Me)(C₃₋₈ cycloalkyl))-CH₂-, -C(OH)(R^{6b})-CH₂-, -CH(Me)-NH-CO-CH₂-, -CO-N(R^{6b})-CH₂-, -C(H)(R^{6b})-CO-N(R^{5b})-CH₂-, -CO-N(R^{6b})-CH₂-CH₂-, -CO-N(R^{6b})-CH₂-CH₂-CO-NH-CH₂-, -CO-NH-C(-CONH₂)=CH-, -CO-NH-CH(-CONH₂)-CH₂-, -CH₂-C(H)(Me)-CH₂-S-, -O-CH₂-CO-NH-, -CH₂-N(R^{6b})-CO-CH₂-O-, -N(R^{6b})-CO-CH₂-O-, -C(H)(-CH₂-aryl)-, -C(H)(-CH₂-heteroaryl)-, -C(NH-aryl)=N-N=CH-, -C(NH-aryl)=N-N=CH-, -NH-CO-CH₂-N(R^{6b})-, -NH-N=C(-aryl)-, -NH-N=C(-aryl)-CO-, -NH-
- 20
- 25
- 30

C(=N-CO-C₁₋₆ alkyl)-NH-(CH₂)₂-, -C(-NH-aryl)=N-N=CH-, -NH-C(-NH-aryl)=N-CONH-, -C(=CH-aryl)-CONH-CH₂-, -CH=C(R^{6b})-CONH-, -CH(-CH₂-aryl)-NH-CO- or -CH(OH)-, wherein said aryl or heteroaryl groups of Z may be optionally substituted by one or more halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, NO₂ or hydroxyl groups;

5 R^{5b} represents hydrogen, C₁₋₆ alkyl or cyano;

R^{6b} represents hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, cyano, COOH, -COOC₁₋₆ alkyl, C₃₋₈ cycloalkyl, -CH₂-C₃₋₈ cycloalkyl, aryl, heteroaryl, -C₁₋₆ alkylene-aryl, -CO-aryl, -O-CO-heteroaryl, -CO-heteroaryl or -C(R^{7b})(R^{8b})-heteroaryl, wherein said aryl groups of R^{6b} may be optionally substituted by one or more halogen or C₁₋₆ alkoxy groups;

10 R^{7b} and R^{8b} independently represent hydrogen or C₁₋₆ alkyl;

R^{1b} represents aryl, C₃₋₈ cycloalkyl, monocyclic or bicyclic heterocyclyl or a monocyclic or bicyclic heteroaryl ring system, wherein R^{1b} may be substituted by one or more (e.g. 1, 2 or 3) R^{4b} groups;


15 R^{4b} represents halogen, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₃₋₈ cycloalkyl, haloC₁₋₆ alkyl, hydroxyl, C₁₋₆ alkoxy, -O-C₁₋₆ alkenyl, haloC₁₋₆ alkoxy, -COOH, -CO-C₁₋₆ alkyl, -COO-C₁₋₆ alkyl, -CONH₂, -CH₂-CONH₂, -NH-C₁₋₆ alkyl, -NH-C₂₋₆ alkenyl, -NH-CO-C₁₋₆ alkyl, -CO-NH-C₁₋₆ alkyl, -O-CH₂-CO-NH-C₁₋₆ alkyl, -CH₂-CH₂-CO-NH-C₁₋₆ alkyl, -S-C₁₋₆ alkyl, -SO-C₁₋₆ alkyl, -SO₂-C₁₋₆ alkyl, -SO₂-NH-C₁₋₆ alkyl, -S-CH₂-CO-C₂₋₆ alkenyl, -SO₂-OH, amino, cyano, NO₂, =O, -CO-NH-(CH₂)₂-OMe, -NH-C₃₋₈ cycloalkyl, -CH₂-CO-NH-C₃₋₈ cycloalkyl, -CO-heterocyclyl, -CO-heteroaryl, -COO-(CH₂)₂-heterocyclyl, -OCH₂-aryl, -OCH₂-heteroaryl, -CH₂-O-CO-aryl, -O-aryl, -NH-CO-aryl, -NH-CO-heteroaryl, -NH-CO-CH₂-aryl, -NH-aryl, aryl or heteroaryl groups, wherein said aryl, heterocyclyl or heteroaryl groups of R^{4b} may be optionally substituted by one or more halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, =S or hydroxyl groups and wherein said C₁₋₆ alkyl or C₂₋₆ alkenyl groups of R^{4b} may be optionally substituted by one or more hydroxyl, amino, cyano, C₁₋₆ alkoxy, CONH₂ or -COO-C₁₋₆ alkyl groups;

m represents an integer from 0 to 3;

20 R^{2b} represents halogen, haloC₁₋₆ alkyl, C₁₋₆ alkyl, hydroxyl, C₁₋₆ alkoxy, -S-C₁₋₆ alkyl, -CH₂-S-C₁₋₆ alkyl, -S-C₂₋₆ alkynyl, amino, cyano, NO₂, =O, =S, -SO₂-C₁₋₆ alkyl, -CONH₂, -CO-C₁₋₆ alkyl, -COO-C₁₋₆ alkyl, -NH-C₁₋₆ alkyl, -NH-CO-C₁₋₆ alkyl, -NH-CO-CH=CH-CH₂-N(Me)₂, C₁₋₆ alkyl, -CO-NH-C₁₋₆ alkyl, -CO-NH-CH(Me)-COOH, -S-CH₂-CO-N(Et)₂, -NH-(CH₂)₂-OH, -NH-(CH₂)₃-OH, -NH-CH(Et)-CH₂-OH, -CO-NH-(CH₂)₃-OH, -CH(CH₂OH)₂ or -S-CH₂-CO-NH-CO-NH-C₁₋₆ alkyl, wherein said C₁₋₆ alkyl groups of R^{2b} may be optionally substituted by one or more hydroxyl groups;

with the proviso that the compound is other than compound number 54, 373, 496 and 585; for use as a casein kinase 1 delta (CK1δ) inhibitor in the treatment of a neurodegenerative disorder, such as tauopathies.

- 5 According to one further particular aspect of the invention which may be mentioned there is provided a compound of formula (IB) or a pharmaceutically acceptable salt or solvate thereof:



(IB)

- 10 wherein

"Het B" represents a 5 membered heterocyclic ring system containing 1 to 3 heteroatoms selected from O, N or S, wherein said ring system is fused to one or more (e.g. 1-3)

further rings to form a polycyclic ring system comprising up to 4 rings;

Z represents a bond, $-C(R^{7b})(R^{8b})-$, $(CH_2)_2$, $-O-$, $-S-$, $-CH_2-O-$, $-(CH_2)_2-O-$, NR^{6b} , $-N(R^{6b})-$

- 15 $C(R^{7b})(R^{8b})-$, $-N(R^{6b})-(CH_2)_2-$, $-N(R^{6b})-(CH_2)_3-$, $-CH_2-N(R^{6b})-(CH_2)_2-$, $-N(R^{6b})-CO-$, $-CH_2-NH-$
 $CO-(CH_2)_2-$, $-N(R^{6b})-CO-CH_2-$, $=N-$, $-C(H)(CN)-$, $-C(=N-NH-COC_{1-6} \text{ alkyl})-$, $-CH=C(R^{6b})-$
 $CO-$, $=CH-$, $-N=CH-$, $-N=C(Me)-$, $-C(R^{6b})=CH-$, $-NH-CO-C(=CH-\text{heteroaryl})-$, $-C=C(Me)_2-$, $-$
 $CH=CH-CO-N(R^{6b})-$, $-CH=C(R^{6b})-NH-CO-$, $-CH=C(R^{6b})-CO-O-CH_2-$, $-CS-S-CH_2-$, $-NH-CS-$
 $NH-$, $-NH-CS-NH-CH_2-$, $-NH-CS-NH-(CH_2)_2-$, $-CH_2-N(CSNH_2)-CH_2-$, $-S-CH_2-$, $-S-(CH_2)_2-O-$
 SO_2 , $-NH-SO_2-$, $-CH_2-NH-SO_2-$, CO , $-CH_2-CO-$, $-(CH_2)_2-CO-$, $-O-CH_2-CO-$, $-(CH_2)_2-CO-$,
 COO , $-COO-C(R^{7b})CO-$, $-CH=C(R^{5b})-CONH-CH_2-$, $-CO-CH_2-N(R^{6b})-CO-$, $-CO-CH_2-C(R^{6b})-$
 CH_2-CO- , $-CO-CH_2-N(R^{6b})-CH_2-$, $-CO-NH-N=C(R^{7b})-$, $-S-CH_2-CO-$, $-S-CH_2-CO-N(R^{6b})-$, $-S-$
 $CH_2-CO-N(R^{6b})-CH_2-$, $-SO_2-N(R^{6b})-C(R^{7b})(R^{8b})-CONH-$, $-SO_2-N(R^{6b})-CH-(CH_2-\text{aryl})-$
 $CONH-CH_2-$, $-CH(-S-C_{1-6} \text{ alkyl})-C(Me)(OH)-$, $-CH_2-C(R^{6b})(OH)-$, $-C(OH)(CH(Me)(C_{3-8}$
 $25 \text{ cycloalkyl}))CH_2-$, $-C(OH)(R^{6b})-CH_2-$, $-CH(Me)-NH-CO-CH_2-$, $-CO-N(R^{6b})-CH_2-$, $-CO-$
 $N(R^{6b})-CH_2-CH_2-$, $-CO-N(R^{6b})-CH_2-CH_2-CO-NH-CH_2-$, $-CO-NH-C(-CONH_2)=CH-$, $-CO-NH-$
 $CH(-CONH_2)-CH_2-$, $-CH_2-C(H)(Me)-CH_2-S-$, $-O-CH_2-CO-NH-$, $-CH_2-N(R^{6b})-CO-CH_2-O-$, $-$
 $N(R^{6b})-CO-CH_2-O-$, $-C(H)(-CH_2-\text{aryl})-$, $-C(H)(-CH_2-\text{heteroaryl})-$, $-C(NH-\text{aryl})=N-N=CH-$, $-$
 $C(NH-\text{aryl})=N-N=CH-$, $-NH-N=C(-\text{aryl})-$, $-NH-N=C(\text{aryl})-CO-$, $-NH-C(=N-CO-C_{1-6} \text{ alkyl})-$
 $30 NH-(CH_2)_2-$, $-C(-NH-\text{aryl})=N-N=CH-$, $-NH-C(-NH-\text{aryl})=N-CONH-$, $-C(=CH-\text{aryl})-CONH-$
 CH_2- , $-CH=C(R^{6b})-CONH-$, $-CH(-CH_2-\text{aryl})-NH-CO-$ or $-CH(OH)-$, wherein said aryl or

heteroaryl groups of Z may be optionally substituted by one or more halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, NO₂ or hydroxyl groups;

R^{5b} represents hydrogen, C₁₋₆ alkyl or cyano;

R^{6b} represents hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, cyano, C₃₋₈ cycloalkyl, -CH₂-C₃₋₈

- 5 cycloalkyl, aryl, heteroaryl, -C₁₋₆ alkylene-aryl, -CO-aryl, -CO-heteroaryl or -C(R^{7b})(R^{8b})-heteroaryl, wherein said aryl groups of R^{6b} may be optionally substituted by one or more halogen or C₁₋₆ alkoxy groups;

R^{7b} and R^{8b} independently represent hydrogen or C₁₋₆ alkyl;

- R^{1b} represents aryl, C₃₋₈ cycloalkyl, monocyclic or bicyclic heterocyclyl or a monocyclic or
10 bicyclic heteroaryl ring system, wherein R^{1b} may be substituted by one or more (e.g. 1, 2 or 3) R^{4b} groups;

R^{4b} represents halogen, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₃₋₈ cycloalkyl, haloC₁₋₆ alkyl, hydroxyl, C₁₋₆ alkoxy, -O-C₁₋₆ alkenyl, haloC₁₋₆ alkoxy, -COOH, -CO-C₁₋₆ alkyl, -COO-C₁₋₆

- 15 NH-C₁₋₆ alkyl, -O-CH₂-CO-NH-C₁₋₆ alkyl, -CH₂-CH₂-CO-NH-C₁₋₆ alkyl, -S-C₁₋₆ alkyl, -SO-C₁₋₆ alkyl, -SO₂-C₁₋₆ alkyl, -SO₂-NH-C₁₋₆ alkyl, -S-CH₂-CO-C₂₋₆ alkenyl, -SO₂-OH, amino, cyano, NO₂, =O, -CO-NH-(CH₂)₂-OMe, -NH-C₃₋₈ cycloalkyl, -CO-heterocyclyl, -CO-heteroaryl, -COO-(CH₂)₂-heterocyclyl, -OCH₂-aryl, -OCH₂-heteroaryl, -CH₂-O-CO-aryl, -O-aryl, -NH-CO-heteroaryl, -NH-CO-CH₂-aryl, aryl or heteroaryl groups, wherein said aryl,
20 heterocyclyl or heteroaryl groups of R^{4b} may be optionally substituted by one or more halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, =S or hydroxyl groups and wherein said C₁₋₆ alkyl or C₂₋₆ alkenyl groups of R^{4b} may be optionally substituted by one or more hydroxyl, amino, cyano, C₁₋₆ alkoxy, CONH₂ or -COO-C₁₋₆ alkyl groups;

m represents an integer from 0 to 3;

- 25 R^{2b} represents halogen, haloC₁₋₆ alkyl, C₁₋₆ alkyl, hydroxyl, C₁₋₆ alkoxy, -S-C₁₋₆ alkyl, -CH₂-S-C₁₋₆ alkyl, -S-C₂₋₆ alkynyl, amino, cyano, NO₂, =O, =S, -SO₂-C₁₋₆ alkyl, -CONH₂, -CO-C₁₋₆ alkyl, -COO-C₁₋₆ alkyl, -NH-C₁₋₆ alkyl, -NH-CO-C₁₋₆ alkyl, -NH-CO-CH=CH-CH₂-N(Me)₂, C₁₋₆ alkyl, -CO-NH-C₁₋₆ alkyl, -CO-NH-CH(Me)-COOH, -S-CH₂-CO-N(Et)₂, -NH-(CH₂)₂-OH, -NH-(CH₂)₃-OH, -NH-CH(Et)-CH₂-OH, -CO-NH-(CH₂)₃-OH, -CH(CH₂OH)₂ or -S-CH₂-CO-
30 NH-CO-NH-C₁₋₆ alkyl, wherein said C₁₋₆ alkyl groups of R^{2b} may be optionally substituted by one or more hydroxyl groups;

for use as a casein kinase 1 delta (CK1δ) inhibitor in the treatment of a neurodegenerative disorder, such as tauopathies.

- 35 In one embodiment of the compound of formula (IB)

"Het B" represents a 5 membered heterocyclic ring system containing 1 to 3 heteroatoms selected from O, N or S, wherein said ring system is fused to a 6 membered ring to form a bicyclic heterocyclic ring system;

Z represents a bond, $-C(R^{7b})(R^{8b})-$, $-O-$, $-S-$, $-CH_2O-$, $-N(R^{6b})-C(R^{7b})(R^{8b})-$, $-N(R^{6b})-(CH_2)_2-$, $-N(R^{6b})-(CH_2)_3-$, $-N(R^{6b})-CO-$, $-N(R^{6b})-CO-CH_2-$, $-N(R^{7b})-CH=$, $=CH-$, $-N=CH-$, $-C(R^{6b})=CH-$, $-C(=C(R^{7b})(R^{8b}))-$, SO_2 , $-CH_2NH-SO_2-$, CO , $-O-CH_2-CO-$, $-SO_2-N(R^{6b})-C(R^{7b})(R^{8b})-CONH-$, $-SO_2-N(R^{6b})-CH(-CH_2-aryl)-CONH-CH_2-$, $-CH(-S-C_{1-6} alkyl)-C(Me)(OH)-$, $-C(H)(R^{6b})-CO-N(R^{5b})-CH_2-$, $-O-CH_2-CO-NH-$, $-N(R^{6b})-CO-CH_2-O-$, $-C(H)(-CH_2-aryl)-$, $-C(NH-aryl)=N-N=CH-$, $-NH-CO-CH_2-N(R^{6b})-$, $-NH-N=C(-aryl)-$, $-NH-C(=N-CO-C_{1-6} alkyl)-NH-(CH_2)_2-$, $-C(=CH-aryl)-CONH-CH_2-$ or $-CH(-CH_2-aryl)-NH-CO-$ wherein said aryl or heteroaryl groups of Z may be optionally substituted by one or more halogen, C_{1-6} alkyl, C_{1-6} alkoxy, NO_2 or hydroxyl groups;

R^{5b} represents hydrogen;

R^{6b} represents hydrogen, methyl, C_{1-6} alkoxy, $-COOH$, $-CO-aryl$, $-O-CO-heteroaryl$ or $-CO-heteroaryl$, wherein said aryl groups of R^{6b} may be optionally substituted by one or more halogen or C_{1-6} alkoxy groups;

R^{7b} and R^{8b} independently represent hydrogen or C_{1-6} alkyl;

R^{1b} represents a monocyclic aryl or heteroaryl ring system, wherein R^{1b} may be substituted by one or more (e.g. 1, 2 or 3) R^{4b} groups;

R^{4b} represents halogen, hydroxyl, $-O-C_{1-6} alkenyl$, $-COO-C_{1-6} alkyl$, $-NH-C_{1-6} alkyl$, $-SO_2NH_2$, amino, cyano, $=O$, $-CH_2CO-NH-C_{3-8} cycloalkyl$, $-CH_2-aryl$, $-OCH_2-heteroaryl$, $-O-aryl$, $-NH-CO-aryl$, $-NH-aryl$ or heteroaryl groups, wherein said aryl, heterocyclyl or heteroaryl groups of R^{4b} may be optionally substituted by one or more halogen, C_{1-6} alkyl, C_{1-6} alkoxy, $=S$ or hydroxyl groups and wherein said C_{1-6} alkyl or C_{2-6} alkenyl groups of R^{4b} may be optionally substituted by one or more hydroxyl, amino, cyano, C_{1-6} alkoxy, $CONH_2$ or $-COO-C_{1-6} alkyl$ groups;

m represents an integer from 0 to 2; and

R^{2b} represents halogen, halo C_{1-6} alkyl, C_{1-6} alkyl, C_{3-8} cycloalkyl, hydroxyl, C_{1-6} alkoxy, $-S-C_{1-6} alkyl$, amino, cyano, NO_2 , $=O$, $-CONH_2$, $-CO-C_{1-6} alkyl$, $-COO-C_{1-6} alkyl$, $C_{1-6} alkyl$, $-CO-NH-C_{1-6} alkyl$ or $-CO-NH-CH(Me)-COOH$, wherein said C_{1-6} alkyl groups of R^{2b} may be optionally substituted by one or more cyano or hydroxyl groups.

In one embodiment, Het B represents a 5 membered heterocyclic ring system containing 1 to 3 heteroatoms selected from O, N or S, wherein said ring system is fused to a 6 membered ring to form a bicyclic heterocyclic ring system. In a further embodiment, Het B represents benzoxazolyl, indolyl or indolizynyl.

In one embodiment, R^{5b} represents hydrogen.

In one embodiment, R^{6b} represents hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, -COOH, -CO-aryl, -
 5 O-CO-heteroaryl, -CO-heteroaryl or -C(R^{7b})(R^{8b})-heteroaryl, wherein said aryl groups of
 R^{6b} may be optionally substituted by one or more halogen or C₁₋₆ alkoxy groups.

In one embodiment, R^{1b} represents a monocyclic aryl or heteroaryl ring system, wherein
 R^{1b} may be substituted by one or more (e.g. 1, 2 or 3) R^{4b} groups. In a further
 10 embodiment, R^{1b} represents a monocyclic aryl group such as phenyl optionally substituted
 by one or more (e.g. 1) R^{4b} groups. In an alternative embodiment, R^{1b} represents a
 monocyclic heteroaryl group such as thienyl, pyrimidinyl or pyrazolinyl optionally
 substituted by one or more (e.g. 1 or 2) R^{4b} groups.

15 In one embodiment, R^{4b} represents halogen, hydroxyl, -O-C₁₋₆ alkenyl, -COO-C₁₋₆ alkyl, -
 NH-C₁₋₆ alkyl, -SO₂-NH₂, amino, cyano, =O, -CH₂-CO-NH-C₃₋₈ cycloalkyl, -CH₂-aryl, -
 OCH₂-heteroaryl, -O-aryl, -NH-CO-aryl, -NH-aryl or heteroaryl groups, wherein said aryl,
 heterocyclyl or heteroaryl groups of R^{4b} may be optionally substituted by one or more
 halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, =S or hydroxyl groups and wherein said C₁₋₆ alkyl or C₂₋₆
 20 alkenyl groups of R^{4b} may be optionally substituted by one or more hydroxyl, amino,
 cyano, C₁₋₆ alkoxy, CONH₂ or -COO-C₁₋₆ alkyl groups.

In a further embodiment, R^{4b} represents halogen (e.g. fluorine), amino or heteroaryl (e.g.
 pyridyl).

25

In one embodiment, Z represents a bond, -C(R^{7b})(R^{8b})-, -O-, -S-, -CH₂-O-, -N(R^{6b})-
 C(R^{7b})(R^{8b})-, -N(R^{6b})-(CH₂)₂-, -N(R^{6b})-(CH₂)₃-, -N(R^{6b})-CO-, -N(R^{6b})-CO-CH₂-, -N(R^{7b})-
 CH=, =CH-, -N=CH-, -C(R^{6b})=CH-, -C(=C(R^{7b})(R^{8b}))-, SO₂, -CH₂-NH-SO₂-, CO-, -O-CH₂-
 CO-, -SO₂-N(R^{6b})-C(R^{7b})(R^{8b})-CONH-, -SO₂-N(R^{6b})-CH(-CH₂-aryl)-CONH-CH₂-, -CH(-S-
 30 C₁₋₆ alkyl)-C(Me)(OH)-, -C(H)(R^{6b})-CO-N(R^{5b})-CH₂-, -O-CH₂-CO-NH-, -N(R^{6b})-CO-CH₂-O-,
 -C(H)(-CH₂-aryl)-, -C(NH-aryl)=N-N=CH-, -NH-CO-CH₂-N(R^{6b})-, -NH-N=C(-aryl)-, -NH-
 C(=N-CO-C₁₋₆ alkyl)-NH-(CH₂)₂-, -C(=CH-aryl)-CONH-CH₂- or -CH(-CH₂-aryl)-NH-CO-
 wherein said aryl or heteroaryl groups of Z may be optionally substituted by one or more
 halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, NO₂ or hydroxyl groups.

35

In a further embodiment, Z represents a bond or CO.

In one embodiment, m represents an integer from 0 to 2. In one embodiment, m represents 0. In an alternative embodiment, m represents 2.

- 5 In one embodiment, R^{2b} represents halogen, haloC₁₋₆ alkyl, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, hydroxyl, C₁₋₆ alkoxy, -S-C₁₋₆ alkyl, amino, cyano, NO₂, =O, -CONH₂, -CO-C₁₋₆ alkyl, -COO-C₁₋₆ alkyl, C₁₋₆ alkyl, -CO-NH-C₁₋₆ alkyl or -CO-NH-CH(Me)-COOH, wherein said C₁₋₆ alkyl groups of R^{2b} may be optionally substituted by one or more cyano or hydroxyl groups.

10

In a further embodiment, R^{2b} represents amino or -CONH₂.

- In one embodiment, the compound of formula (IB) is selected from any of compounds 2-3, 26-28, 30-33, 35, 47-48, 51, 57-60, 63-64, 78, 84, 113, 123, 127-129, 145, 155-157, 171-173, 204, 206-207, 210, 225, 227, 233, 235-236, 241-242, 244, 249, 269, 285, 288, 303, 307-312, 314-316, 320, 324-325, 333, 336, 351, 357-360, 374-375, 384-391, 396, 399-402, 404-405, 407-411, 414, 424-425, 427-428, 437, 448, 456-457, 482, 484-485, 489-491, 495, 497-498, 505, 507, 516, 519, 524, 526, 553, 559-560, 568, 570, 575, 609, 615-616, 618, 626-627, 638, 653, 669, 692-694, 705, 709, 712, 716, 719, 725, 734, 738, 740, 746, 749, 753-754, 756, 758-759, 767, 770, 777, 784-785, 790, 792, 796, 800-801, 804-805, 808, 819, 821, 827-828, 831, 833, 838, 844, 847, 857-858, 869, 872, 875, 933, 952, 955, 969, 987, 990 or 999 as described herein or a pharmaceutically acceptable salt or solvate thereof.

- 25 In a further embodiment, the compound of formula (IB) is selected from any of compounds 2-3, 26-28, 30, 32-33, 47-48, 51, 59-60, 84, 113, 123, 127, 129, 145, 155, 157, 172-173, 204, 206-207, 210, 225, 233, 235-236, 241, 244, 269, 285, 288, 307-311, 315-316, 320, 324-325, 333, 336, 351, 357-360, 374-375, 385-386, 388-391, 396, 399-402, 404-405, 407-410, 414, 424, 427-428, 437, 457, 482, 490, 495, 497-498, 505, 516, 519, 553, 559-560 or 568 as described herein or a pharmaceutically acceptable salt or solvate thereof.

- In a yet further embodiment, the compound of formula (IB) is selected from any of compounds 30, 314, 324-325, 391, 405, 626, 705, 753-754, 759, 770, 784, 808, 833 or 847 as described herein or a pharmaceutically acceptable salt or solvate thereof.

35

In a still yet further embodiment, the compound of formula (IB) is selected from any of compounds 324-325, 405, 754 or 847 as described herein or a pharmaceutically acceptable salt or solvate thereof.

- 5 In one embodiment, the compound of formula (IB) is selected from any of compounds 30, 288, 314, 324-325, 336, 374, 391, 405, 615-616, 626, 705, 740, 753-754, 756, 759, 770, 784, 808, 819, 833, 844, 847, 869, 872, 875, 933, 952, 955, 969, 987, 990 and 999 as described herein or a pharmaceutically acceptable salt or solvate thereof. The compounds of this embodiment were tested in the CK1δ inhibition assay as described
10 herein and exhibited inhibition of greater than 5%.

- In a further embodiment, the compound of formula (IB) is selected from any of compounds 324-325, 405, 754, 847, 952, 987, 990 and 999 as described herein or a pharmaceutically acceptable salt or solvate thereof. The compounds of this embodiment
15 were tested in the CK1δ inhibition assay as described herein and exhibited inhibition of greater than 50%.

- In a yet further embodiment, the compound of formula (IB) is selected from any one of compounds:
- 20 5-(1,3-benzoxazol-2-yl)-4-(pyridin-4-yl)pyrimidin-2-amine (Compound 324);
2-amino-3-[(thiophen-2-yl)carbonyl]indolizine-1-carboxamide (Compound 847);
2-[3-(pyridin-4-yl)-1H-pyrazol-4-yl]-1,3-benzoxazole (Compound 952);
2-amino-3-[(4-fluorophenyl)carbonyl]indolizine-1-carboxamide (Compound 987);
2-amino-3-benzoylindolizine-1-carboxamide (Compound 990); and
25 2-amino-1-[(4-fluorophenyl)carbonyl]-1H-indole-3-carboxamide (Compound 999);
or a pharmaceutically acceptable salt or solvate thereof.

- In a yet further embodiment, the compound of formula (IB) is selected from any of compounds 324, 952, 987, 990 and 999 as described herein or a pharmaceutically
30 acceptable salt or solvate thereof. The compounds of this embodiment were tested in the CK1δ inhibition assay as described herein and exhibited inhibition of greater than 90%.

- In a still yet further embodiment, the compound of formula (IB) is selected from any of compounds 324, 952, 987 and 999 as described herein or a pharmaceutically acceptable
35 salt or solvate thereof. The compounds of this embodiment were tested in a range of kinase inhibition assays and not only exhibited inhibition of greater than 90% in the CK1δ

inhibition assay as described herein, but also demonstrated significant and selective inhibition for CK1δ when compared with other kinases.

For example, compound number 324 (5-(1,3-benzoxazol-2-yl)-4-(pyridin-4-yl)pyrimidin-2-amine) demonstrated selectivity for CK1δ over ABL2/ARG, ALK4/ACVR1B, ALK5/TGFBR1, CDK5/p25, CK1a1, CK1g1, CK1g3, CLK2, c-SRC, EGFR, EPHA2, FGFR1, GSK3b, HGK/MAP4K4, JNK2, KDR/VEGFR2, LCK, MSK1/RPS6KA5, PDK1/PDPK1, PIM3, PKA, PKCa, PKCb2, RIPK2, ROCK1, TNIK and YES/YES1 each of which were inhibited at levels lower than 40%.

10

For example, compound number 952 (2-[3-(pyridin-4-yl)-1H-pyrazol-4-yl]-1,3-benzoxazole) demonstrated selectivity for CK1δ over ABL2/ARG, ALK4/ACVR1B, ALK5/TGFBR1, CDK5/p25, CK1g1, CK1g2, CK1g3, c-SRC, EGFR, EPHA2, FGFR1, KDR/VEGFR2, LCK, MSK1/RPS6KA5, PDK1/PDPK1, PIM3, PKA, PKCa, PKCb2, ROCK1 and YES/YES1 each of which were inhibited at levels lower than 40%.

15

For example, compound number 987 (2-amino-3-[(4-fluorophenyl)carbonyl]indolizine-1-carboxamide) demonstrated selectivity for CK1δ over ABL2/ARG, CDK5/p25, CK1g1, CK1g2, CK1g3, CLK2, c-SRC, FGFR1, GSK3b, HGK/MAP4K4, JNK2, KDR/VEGFR2, LCK, MSK1/RPS6KA5, PDK1/PDPK1, PIM3, PKCa, PKCb2, ROCK1 and TNIK each of which were inhibited at levels lower than 40%.

20

For example, compound number 999 (2-amino-1-[(4-fluorophenyl)carbonyl]-1H-indole-3-carboxamide) demonstrated selectivity for CK1δ over ABL2/ARG, CDK5/p25, CK1g1, CK1g2, CLK2, c-SRC, FGFR1, GSK3b, HGK/MAP4K4, KDR/VEGFR2, LCK, MSK1/RPS6KA5, PDK1/PDPK1, PIM3, PKCa, PKCb2 and ROCK1 each of which were inhibited at levels lower than 40%.

25

In a still yet further embodiment, the compound of formula (IB) is selected from any of compounds 324 and 987 as described herein or a pharmaceutically acceptable salt or solvate thereof. The compounds of this embodiment have been demonstrated to have a protective effect on cell viability as can be seen in the data presented herein and in particular within Figures 1 and 2. The compounds of this embodiment have also been demonstrated to inhibit phosphorylation of two different amino acid residues within Tau proteins (i.e. Ser 396 and Thr 391) as shown in Figures 4 and 5.

35

In a still yet further embodiment, the compound of formula (IB) is compound 324 as described herein or a pharmaceutically acceptable salt or solvate thereof. The compound of this embodiment has been demonstrated to have a protective effect on cell viability in a dose dependent manner as can be seen in the data presented herein and in particular within Figure 1. The compound of this embodiment has also been demonstrated to inhibit phosphorylation of two different amino acid residues within Tau proteins (i.e. Ser 396 and Thr 391) as shown in Figures 4A and 5.

In the present context, the term "pharmaceutically acceptable salt" is intended to indicate salts which are not harmful to the patient. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable metal salts and pharmaceutically acceptable alkaline addition salts. Acid addition salts include salts of inorganic acids as well as organic acids.

Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methanesulfonic, ethanesulfonic, tartaric, ascorbic, pantoic, bismethylene salicylic, ethanedithionyl, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci. 1977, **66**.

Examples of metal salts include

lithium, sodium, potassium, magnesium salts and the like. Examples of ammonium and alkylated ammonium salts include ammonium, methylammonium, dimethylammonium, trimethylammonium, ethylammonium, hydroxyethylammonium, diethylammonium, butylammonium, tetramethylammonium salts and the like.

Representative examples of alkaline salts include, for example, sodium, potassium, lithium, calcium, magnesium or ammonium or organic bases such as, for example, methylamine, ethylamine, propylamine, trimethylamine, diethylamine, triethylamine, N,N-dimethylethanolamine, tris(hydroxymethyl)aminomethane, ethanolamine, pyridine, piperidine, piperazine, picoline, dicyclohexylamine, morpholine, benzylamine, procaine, lysine, arginine, histidine, N-methylglucamine.

According to the invention, the compounds of formula (IB) can be in racemic forms, as well as in the form of pure enantiomers or non racemic (scalemic) mixture of enantiomers, including when the compounds of formula (IB) have more than one stereogenic centre. In case the compounds of formula (IB) have unsaturated carbon carbon double bonds, both the cis (Z) and trans (E) isomers and their mixtures belong to the invention.

References herein to "halogen" means a fluorine, chlorine, bromine or iodine atom.

References herein to "C₁₋₆ alkyl" means any linear, branched hydrocarbon groups having 1 to 6 carbon atoms, or cyclic hydrocarbon groups having 3 to 6 carbon atoms. Representative examples of such alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl, n-pentyl, isopentyl, neopentyl, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. References to "haloC₁₋₆alkyl" mean a C₁₋₆ alkyl group substituted by one or more halogen atoms as herein defined.

References herein to "C₁₋₆ alkylene" means a saturated divalent hydrocarbon chain having the specified number of member atoms. For example, C₁₋₆ alkylene refers to a bond or an alkylene group having from 1 to 6 member atoms. Alkylene groups may be straight or branched. Representative branched alkylene groups have one or two branches. Alkylene includes methylene, ethylene, propylene (n-propylene and isopropylene) and butylene (n-butylene, isobutylene, and t-butylene).

References herein to "C₂₋₆ alkenyl" means any linear, branched hydrocarbon groups of 2 to 6 carbon atoms, or cyclic hydrocarbon group having 3 to 6 carbon atoms having at least one double bond. Representative examples of such alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl.

References herein to "C₂₋₆ alkynyl" means any linear, or branched hydrocarbon groups of 2 to 6 carbon atoms, having at least one triple bond. Representative examples of such alkynyl groups include ethynyl, propargyl and butynyl.

References herein to 'C₁₋₆ alkoxy' means an -O-C₁₋₆ alkyl group wherein C₁₋₆ alkyl is as defined herein. Examples of such groups include methoxy, ethoxy, propoxy, butoxy, pentoxy or hexoxy and the like.

References herein to 'C₃₋₈ cycloalkyl' means a saturated monocyclic hydrocarbon ring of 3

to 8 carbon atoms. Examples of such groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl and the like.

References herein to 'aryl' means a C₆₋₁₂ monocyclic or bicyclic hydrocarbon ring wherein at least one ring is aromatic. Examples of such groups include phenyl, indyl or naphthyl and the like.

References herein to "heteroatom" means a nitrogen, sulphur, or oxygen atom.





References herein to "heterocyclyl" means a saturated or unsaturated non-aromatic ring containing from 1 to 4 heteroatoms as member atoms in the ring. Heterocyclyl groups containing more than one heteroatom may contain different heteroatoms. Heterocyclyl groups may be optionally substituted with one or more substituents as defined herein. Heterocyclyl groups are monocyclic ring systems or fused bicyclic or polycyclic ring systems or bicyclic structures known as heterocyclic "spiro" ring systems. In certain embodiments, heterocyclyl is saturated. In other embodiments, heterocyclyl is unsaturated and non-aromatic. Non-limiting examples of monocyclic heterocyclyl ring systems include pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, pyranlyl, tetrahydropyranlyl, dihydropyranlyl, tetrahydrothienyl, pyrazolidinyl, oxazolidinyl, thiazolidinyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, thiamorpholinyl, 1,3-dioxolanyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,3-oxathiolanyl, 1,3-oxathianyl, 1,3-dithianyl, and azetidiny.






References herein to "heteroaryl" means an aromatic ring containing from 1 to 4 heteroatoms as member atoms in the ring. Heteroaryl groups containing more than one heteroatom may contain different heteroatoms. Heteroaryl groups may be optionally substituted with one or more substituents as defined herein. Heteroaryl groups are monocyclic ring systems or are fused bicyclic or polycyclic ring systems. Monocyclic heteroaryl rings have 5 or 6 member atoms. Bicyclic heteroaryl rings have from 7 to 11 member atoms. Bicyclic heteroaryl rings include those rings wherein phenyl and a monocyclic heterocyclyl ring are attached forming a fused bicyclic ring system, and those rings wherein a monocyclic heteroaryl ring and a monocyclic cycloalkyl, cycloalkenyl, heterocyclyl, or heteroaryl ring are attached forming a fused bicyclic ring system. Non-limiting examples of heteroaryl includes pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, furanyl, furazanyl, thienyl, triazolyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, tetrazinyl, indolyl, isoindolyl, indoliziny, indazolyl, purinyl, quinolinyl, isoquinolinyl, quinoxaliny, quinazolinyl, pteridinyl, cinnolinyl, benzimidazolyl, benopyranly,






benzoxazolyl, benzofuranyl, isobenzofuranyl, benzothiazolyl, benzothienyl, furopyridinyl, and naphthyridinyl.






References herein to "heterocyclic ring system" mean either a heterocyclyl ring system or
5 a heteroaryl ring system as hereinbefore defined.






Representative compounds of formula (IB) include the compounds as set forth below:






Compound Number	Structure
2	
3	
26	
27	






28	
30	
31	
32	
33	






35	 <chem>c1ccc2c(c1)c(cnc2)n3ccccc3</chem>
47	 <chem>CCc1cccnc1N2C3=CC=CC=C3N=N2</chem>
48	 <chem>c1ccc(cc1)CC2C3=CC=CC=C3N=N2c4ccccc4</chem>
51	 <chem>Nc1ccc2c(c1)c(c[nH]2)CNc3ccccn3</chem>
54	 <chem>CN1C2=CC=C(C(=C2N1)S(=O)(=O)C)c3ccccc3N</chem>






57	
58	
59	
60	
63	






64	 <chem>Brc1ccc2nc3ccccc3nc2n1Cl</chem>
78	 <chem>N#Cc1c2ccccc2n1c3ccncc3Cl</chem>
84	 <chem>Clc1cc2nc3ccncc3n2c1c4ccncc4</chem>
113	 <chem>Clc1ccc(cc1)N2C(=O)Nc3cc(C(F)(F)F)cn32</chem>
123	 <chem>NC(=O)c1c2ccccc2n1c3cc(S(=O)(=O)c4ccccc4)cc(Cl)c3</chem>






127	
128	
129	
145	
155	






156	
157	
171	
172	
173	






204	
206	
207	
210	
225	






227	 <chem>Cc1c2nc3c(ncn3c2)nc4c1cnc4C5=CC(OC)=CC(OC)=C5</chem>
233	 <chem>COC1=CC=C(NC2=NC3=C(N2)N=CN3)C=C1</chem>
235	 <chem>COC1=CC=C(C=C1)CN(C2=CC=CC=C2)C3=C(C=C(C=C3C4=CC(=C(C=C4)C(=O)O)C5=CC(=C(C=C5)[N+]([O-])=O)N6=CC=CC=C6N7=CC=CC=C7N8=CC=CC=C8N9=CC=CC=C9N10=CC=CC=C10)N</chem>
236	 <chem>NC(=O)CC1=CC=CC=C1C2=CC=CC=C2C3=CC=CC=C3C4=CC(=C(C=C4)C(=O)O)C5=CC(=C(C=C5)[N+]([O-])=O)N6=CC=CC=C6N7=CC=CC=C7N8=CC=CC=C8N9=CC=CC=C9N10=CC=CC=C10</chem>
241	 <chem>Cc1c2nc3c(ncn3c2)nc4c1cnc4C5=CC(OC)=CC(OC)=C5</chem>






242	 <chem>Cc1nc2ccccc2n1C(=O)c3ccc4c(c3)OCO4</chem>
244	 <chem>Cc1nc2ccccc2n1C(=O)c3ccccc3c4c2c1OCCc5ccccc54</chem>
249	 <chem>O=C(c1ccc2c(c1)OCO2)c3ccc4c(c3)OCO4</chem>
269	 <chem>OCC(CO)N1C=NC2=C(N1C(=O)Nc3ccccc3)N=CN2</chem>
285	 <chem>COc1ccc(cc1)c2nc3ccccc3[nH]2C(=O)NCCCCO</chem>






288	 <chem>COC(=O)c1nc2c(c1)c3ccccc3[nH]2-c4ccc(OC)cc4</chem>
303	 <chem>CC1=CC=NC=C1C(=O)N(C1=CC=CC=C1C(=O)C2=CC=CC=C2)C(=O)C3=CC=C(NC4=CC=C5C(C4)OCCO5)C=C3</chem>
307	 <chem>CC1=NC2=C(C1)C(=CC=C2N3C(=O)NC3=CC=C4C(=C2)C=CC=N4)C5=CC=CC=N5</chem>
308	 <chem>Nc1ccc(cc1)N2C(=O)NC3=CC=CC=N32</chem>
309	 <chem>O=C1NC2=CC=CC=C2C1(Cc3ccncc3)C3=CC=CC=C3</chem>






310	 <chem>O=C1NC2=CC=CC=C2C1(Cc3ccccn3)O</chem>
311	 <chem>O=C1NC2=CC(Br)=CC=C2C1(Cc3ccccc3)O</chem>
312	 <chem>O=C1NC2=CC=CC=C2C1=C/C=C/c3ccc(Oc4ccccc4)cc3</chem>
314	 <chem>O=C1NC2=CC=CC=C2C1=N/C=C/c3c[nH]c4ccccc34</chem>
315	 <chem>O=C1NC2=CC=CC=C2C1=C/C=C/c3c[nH]c4ccccc34</chem>






316	 <chem>O=C1OC2=CC=C(C=C2C1/C=C/c3ccc(O)cc3)C=C4C=CC(=C4)O</chem>
320	 <chem>COc1cc(OC)c2cc3c(c1)c4ccccc4oc3n2C5=CC=CC=C5</chem>
324	 <chem>Nc1nc2c(ncn2C3=CC=CC=C3O4C=CC=CC=C4N3)cc5ccncc5</chem>
325	 <chem>COc1ccc(Nc2nc3c(ncn3c2C4=CC=CC=C4O5C=CC=CC=C54))cc1</chem>
333	 <chem>C1=CC=C2C(=C1)OC2OC3=CC=CC=C3N1N=CN=C1</chem>






336	 <chem>Nc1nc(N)nc(Cc2ccc3c(c2)OCO3)c1</chem>
351	 <chem>Oc1cccc(Nc2nc3ccccc3n2)c1</chem>
357	 <chem>CC(C)Sc1c2ccccc2n1N(c3ccc(OC)cc3)c4ccccc4</chem>
358	 <chem>N#CCSc1c2ccccc2n1N(c3ccc(OC)cc3)c4ccccc4</chem>
359	 <chem>OC(CCCNc1c2cc3c(c1)nnoc3cc2S4C5=CC=CC=C5O4)c6c([N+](=O)[O-])ccc7nnc67</chem>






360	
373	
374	
375	
384	






385	 <chem>CC1=CC=C(C=C1S2=NC=CC=C2S2)S(=O)(=O)NC(C(C)C)C(=O)NC3=CC=C(S)N3C</chem>
386	 <chem>CC1=CC=C(C=C1S2=NC=CC=C2S2)S(=O)(=O)NC(C(C)C)C(=O)NC3=CC=CC=S3</chem>
387	 <chem>c1ccc(cc1)CC(C(=O)NC2=CC=CC=N2)NC(=O)NS3=CC=C(C=C3)S4=NC=CC=C4S4</chem>
388	 <chem>c1ccc(cc1)CC(C(=O)NC2=CC=CC=C2)NC(=O)NS3=CC=C(C=C3)S4=NC=CC=C4S4</chem>
389	 <chem>c1ccc(cc1)CC(C(=O)NC2=CC=CC=C2)NC(=O)NS3=CC=C(C=C3)S4=NC=CC=C4S4</chem>






390	 <chem>CC1=CN=C(S1)C(=O)NCCNS(=O)(=O)c2ccccc2c3ncsc3</chem>
391	 <chem>NC(=O)COC1=NC2=CC=NC=C2S1NS(=O)(=O)c3ccc(NC(=O)COc4cnc5ccsc45)cc3</chem>
396	 <chem>CC(C)CC1=NC2=NC(=N1)SC2=N3C=CC=CC3</chem>
399	 <chem>COC1=CC(OC)=CC=C1CN(Cc2ccoc2)c3nc4c(ncn4C5CCCCC5)sc6ccccc36</chem>
400	 <chem>Cc1c2c(ncn2c3ccccc13)Nc4ccc5c(c4)OCO5</chem>






401	 <chem>Cc1c(C)c2nc3c(ncn3c2s1)Nc4ccc5c(c4)OCO5</chem>
402	 <chem>Cc1c2c(ncn2c1s1)Nc3ccc4c(c3)OCO4</chem>
404	 <chem>c1c2c(ncn2c1s1)Nc3ccc4c(c3)OCO4</chem>
405	 <chem>Oc1ccc(Nc2c3c(ncn3c2s1)C4CCCCC4)cc1</chem>
407	 <chem>COC(=O)c1ccc(Nc2c3c(ncn3c2s1)C4CCCCC4)cc1</chem>






408	
409	
410	
411	
414	






424	 <chem>CC(C)Cc1c2sc3ncnc3c2n1N4CCOCC4</chem>
425	 <chem>CC(C)(CS)C1=NC2=CC=CC=C2N1N3=CC=CC=C3S</chem>
427	 <chem>CC1=CC=C(C(=C1)N(C)C)C(=O)OCC2=NC3=CC=CC=C3S2</chem>
428	 <chem>O=C(OCC1=NC2=CC=CC=C2S1)N3CCc4ccccc4N3</chem>
437	 <chem>Nc1ncnc2c1nc3c2nc4ccccc4n3</chem>






448	 <chem>O=C1NC2=C(C(=N1)C3=CC=C(C=C3)Br)C4CCNCC42</chem>
456	 <chem>CC1=CC=C(C=C1)C2=NC(=CNC2=O)C3=CC=C(C=C3)C4CCNCC4</chem>
457	 <chem>CC1=CC=C(C=C1)C2=NC(=CNC2=O)C3=CC=C(C=C3)C4CCNCC4</chem>
458	 <chem>CC1=CC=C(C=C1)C2=NC(=CNC2=O)C3=CC=C(C=C3)C4CCNCC4</chem>
482	 <chem>ClC1=CC=C2C(=C1)C(=CN2)C3=CC=C(C=C3)C4CCNCC4</chem>






484	 <chem>Clc1ccccc1C=C(c2nc3ccccc3n2)C(=O)NCCc4ccc5c(c4)OCO5</chem>
485	 <chem>Fc1ccc(cc1)/C=C/C(=O)NCCc2ccccc2-c3nc4ccccc4[nH]3</chem>
489	 <chem>COc1cc(OC)c(OC)cc1-c2nn[nH]2</chem>
490	 <chem>Cc1nc2ccccc2n1C(=O)c3cc(O)ccc3</chem>
491	 <chem>c1ccc2c(c1)c(c[nH]2)CCNC(=O)c3ccc4c(c3)OCO4</chem>






495	 <chem>Cc1cnc2c(c1)sc(c2)C3=CN=CN3</chem>
496	 <chem>Cc1c2ncnc2sc1CNCC3=NC4=CC=CC=C4N3</chem>
497	 <chem>Cc1c2ncnc2sc1COCC(=O)Nc3ccc(C#N)cc3</chem>
498	 <chem>C1=CC=C2C(=C1)SC(=N2)C3=CC=CC=C3COCC(=O)Nc4ccccc4</chem>
505	 <chem>C1=CC=C2C(=C1)SC(=N2)C3=CC=CC=C3COCC(=O)Nc4ccc5c(c4)oc6ccccc6o5</chem>






507	 <chem>COc1ccc2c(c1)oc(CCN(C2)C(=O)Nc3ccsc3)c2</chem>
516	 <chem>COc1cc(OC)ccc1c2nc(C(=O)N(C)C(=O)[O-])c3cc4c(c2n3)c5ccccc5[nH]4</chem>
519	 <chem>CC1=C(C)OC2=NC=NC=C2N1Nc3ccncc3</chem>
524	 <chem>Cc1ccc(cc1OCc2nn[nH]2)n3c4ccccc4nn3</chem>
526	 <chem>BrC1=CC=C(C=C1)c2nc3c(ncn3C)C(F)C2</chem>






553	
559	
560	
568	
570	






575	 <chem>c1ccc2c(c1)oc3ccc(cc23)Oc4ccncc4</chem>
585	 <chem>Cc1cc(C)c2nc3c(c12)nn(C)C3CCCOc4ccc(OC)cc4</chem>
590	 <chem>O=C1NC(=O)C(=C2C(=C(N)C(=C2)Br)C3C(=O)NCCC3)C(=O)N1</chem>
594	 <chem>CC1=CC=C(C=C1)C2=CC3=C(N2)N(C(C)(C)C)N=C3N</chem>
596	 <chem>COc1cc(OC)c2cc3c(c1O2)C(=O)Nc4ccccc34</chem>

597	
601	
602	
609	
615	


616	 <chem>CCCCNC1=NC2=CC=CC=C2N3C1=NC=CC=C3C(=O)C4=CC=CC=C4</chem>
618	 <chem>COCCCN(C1=NC2=CC=CC=C2S1)C3=NC=CC=C3C4=CC=CC=C4N=C5C(C)=CC=C5N</chem>
626	 <chem>CC1=CC=CC=C1N2C=NC=C2C3=NC=CC=C3NC4=NC5=CC=CC=C5N4</chem>
627	 <chem>CC1=CC=C(C=C1N2C=NC=C2C3=NC=CC=C3)NC4=NC5=CC=CC=C5N4</chem>
638	 <chem>O=C1C(=CC=C(C=C1C2=NC3=CC=CC=C3N2)C4=NC=CC=C4N=C5C=CC=CC=C5N4)Cl</chem>

649	 <chem>Cc1nc2ncnc2n(c1)Sc1ccccc1NC(=O)c1ccsc1</chem>
653	 <chem>CCOC(=O)c1cc(s1)NC(=O)c2c3ccccc3n(c2)c4ccccc4</chem>
669	 <chem>Nc1nc2ccccc2n1C3Cc4ccccc4O3</chem>
692	 <chem>Nc1nc2ccccc2n1CCc3ccncc3c4ccc(F)cc4</chem>
693	 <chem>Nc1nc2ccccc2n1CNc3ccncc3</chem>


694	 <chem>Cc1nc2sc(cc12)c3ccccc3NC(=O)c4ccc(F)cc4</chem>
703	 <chem>Cc1nc2nc3ccncc3n2c1N(C)Cc4cc(OC)cc(OC)c4</chem>
705	 <chem>Cc1nc(C)c(Nc2nc3cc4ccccc4oc3cc2)c5ccccc5O</chem>
709	 <chem>COC(=O)c1cc2nc3ccncc3n2c1-c1ccnnc1C</chem>
712	 <chem>Cc1ccc2c(c1)OCO2Cc1ccc(NC(=O)C(OC(=O)c3ccsc3)c4cccnc4)cc1</chem>

716	 <chem>CN(C)C(=O)c1ccccc1c2nc(N)sc2</chem>
719	 <chem>C1CCCCC1NC(=O)CN(C(=O)c2cc3ccccc3o2)c4cc5ccccc5s4</chem>
725	 <chem>COC1=CC=CC=C1NCC2=CN(C)C=CN2</chem>
734	 <chem>CNC1=CC=CC=C1SC2=CN3C=CC=CC3O2</chem>
738	 <chem>O=C1C2C=CC=CC=C2N1C(=O)c3ccncc3C(=O)c4ccncc4</chem>


740




746




749













753













754













756	
758	
759	
767	
770	






777	
778	
784	
785	
790	






792	 <chem>Nc1nc2scn2c1S(=O)(=O)NCc3ccsc3</chem>
796	 <chem>CCc1cnc2c1nc(N)n2SCc3ccccn3</chem>
800	 <chem>CC(C)(S1C=NC2=C1N=CN2C(=O)O)C3=CC=CC=N3</chem>
801	 <chem>Nc1cc[nH]1NC2=CC=C3C(=C2)N=CN3</chem>
804	 <chem>CN1C(=O)OC2=CC=C(C=C2C(=O)c3cc(F)cc(F)c3)C4=CC=CC=C4</chem>






805	 <chem>CN1C(=O)C2=CC=C(C=C2N1C)C(=O)c3ccc(Cl)cc3</chem>
808	 <chem>CN1C=CC=C1C(=O)c2ccccc2C3=NC4=CC=CC=C4S3</chem>
819	 <chem>O=C(CC1=CC=C(C=C1)C2=NN3C=NC=N3S2)N4CCCCC4</chem>
821	 <chem>N#CC1=C(C(=O)N2C=CC=CC=N2)SC3=CC=CC=C13</chem>
827	 <chem>CC1=CC=C(C=C1N)C2=CC=CC=C2O2</chem>

828	 <chem>N#Cc1c2c(s1)CCNCC2C(=O)Nc3ccncc3</chem>
831	 <chem>O=C1C(=O)Nc2ccccc21C(=O)c3ccccc3</chem>
833	 <chem>C1=NC2=C(N1)N=CN=C2NCc3ccncc3</chem>
838	 <chem>O=C1C(=O)Nc2ccccc21C(=O)c3ccncc3</chem>
844	 <chem>Cc1nc2c(nc1O2)C(=O)Nc3ccncc3</chem>

847	 <chem>NC(=O)c1c(N)c(C(=O)Cc2ccsc2)n3ccccc13</chem>
857	 <chem>CC(=O)c1c(Cc2cc3c(c1)occc3O)nc4ccccc42</chem>
858	 <chem>CC(=O)c1c(Cc2ccncc2)nc3ccccc13</chem>
869	 <chem>C1=CC=C2C(=C1)S=C3N=CN=C3C2C1=CC=CC=C1NC1=CC=CC=N1</chem>
872	 <chem>Nc1nc2ccccc2n(Cc3ccccc3)c1</chem>

875	 <chem>Cc1ccncc1NCc2cnc3ccncc23</chem>
877	 <chem>CCOC(=O)C1=C(C2=CC=CC=C2S2)C3=CN=CN(C)S3</chem>
891	 <chem>CC1=CC=C2C(=C1)OC2N(S(=O)(=O)C)CC3=CC=NO3</chem>
910	 <chem>CC1=CC=C2C(=C1)OCC2N(S(=O)(=O)C)CC3=CC=NO3</chem>
912	 <chem>Cc1cc2nc3ccncc3c(=O)n2CCc4ccncc4</chem>

926	 <chem>CN1C=CC2=C1C=CC=C2NC(=O)CCN(C)S(=O)(=O)c3ccc(F)cc3</chem>
933	 <chem>CN1C(=O)N(C)c2ccccc2C1=O</chem>
952	 <chem>c1ccc2c(c1)oc(c2)c3c[nH]c3c4cccnc4</chem>
955	 <chem>Nc1nc2ccccc2n1Cc3ccccc3</chem>
962	 <chem>CN1C=NC=C1NC(=O)Oc2ccccc2C1=O</chem>

963	
969	
987	
990	
999	

According to a further aspect of the invention, there is provided a compound of formula (IB) for use as a casein kinase 1 delta (CK1δ) inhibitor in the treatment of a neurodegenerative disorder, such as tauopathies.

5 Compounds of formula 2-3, 26-28, 30-33, 35, 47-48, 51, 57-60, 63-64, 78, 84, 113, 123, 127-129, 145, 155-157, 171-173, 204, 206-207, 210, 225, 227, 233, 235-236, 241-242, 244, 249, 269, 285, 288, 303, 307-312, 314-316, 320, 324-325, 333, 336, 351, 357-360, 374-375, 384-391, 396, 399-402, 404-405, 407-411, 414, 424-425, 427-428, 437, 448, 456-457, 482, 484-485, 489-491, 495, 497-498, 505, 507, 516, 519, 524, 526, 553, 559-
10 560, 568, 570, 575, 609, 615-616, 618, 626-627, 638, 653, 669, 692-694, 705, 709, 712, 716, 719, 725, 734, 738, 740, 746, 749, 753-754, 756, 758-759, 767, 770, 777, 784-785, 790, 792, 796, 800-801, 804-805, 808, 819, 821, 827-828, 831, 833, 838, 844, 847, 857-858, 869, 872, 875, 933, 952, 955, 969, 987, 990 or 999 are either commercially available or may be prepared in accordance with known synthetic procedures.

15

According to a further aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (IB) for use in the treatment of a neurodegenerative disorder, such as tauopathies.

20 The pharmaceutical compositions of the invention may comprise, in addition to one of the above substances, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of administration, e. g. oral,
25 intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal routes.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant.

30 Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil.

Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

35

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction,

the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection.

5 Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

The compounds of formula (IB) are believed to be casein kinase 1 delta (CK1 δ) inhibitors. Certain compounds of formula (IB) have inhibitory activity of greater than 5%, in particular
10 greater than 10%, more particularly greater than 25%, yet more particularly greater than 50%, especially greater than 75%, such as greater than 90%. Such compounds may be useful in the treatment in neurodegenerative disorders such as tauopathies. Tauopathies are conditions which are characterised by neurofibrillary tangles or aggregates of the tau protein. Tauopathies are a recognised class of conditions known to those skilled in the art
15 and include Alzheimer's disease, frontotemporal dementia with Parkinsonism linked to chromosome 17 (FTDP-17), progressive supranuclear palsy (PSP), Pick's disease, corticobasal degeneration, multisystem atrophy (MSA), neurobasal degeneration with iron accumulation, type 1 (Hallervorden-Spatz), argyrophilic grain dementia, Down's syndrome, diffuse neurofibrillary tangles with calcification, dementia pugilistica,
20 Gerstmann-Straussler-Scheinker disease, myotonic dystrophy, Niemann-Pick disease type C, progressive subcortical gliosis, prion protein cerebral amyloid angiopathy, tangle only dementia, postencephalitic parkinsonism, subacute sclerosing panencephalitis, Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis/parkinsonism-dementia complex, non-Guamanian motor neuron disease with neurofibrillary tangles/dementia, and
25 Parkinson's disease. The intracellular tau deposits are usually neuronal or glial and are filamentous and generally in a hyperphosphorylated state as compared to the level of phosphorylation in tau from control human brain. In the case of AD, this hyperphosphorylated tau is often referred to a paired helical filament tau (PHF) tau because it is derived from the PHF. In one embodiment, the tauopathy comprises
30 Alzheimer's disease.

According to a further aspect of the invention, there is provided a method of treating a neurodegenerative disorder, such as tauopathies, which comprises administering a therapeutically effective amount of a compound of formula (IB).

35

Biological Data

1. CK1δ Inhibition Assay

The compounds of the invention may be tested for inhibition of casein kinase 1 delta (CK1δ) in accordance with the assay protocols described in US 2010/0152157, EP 1,636,375 or Hanger *et al* (2007) J. Biol. Chem. 282, 23645-23654. In particular, the assay was conducted in accordance with the following protocol :

Reaction Buffer:

Base Reaction buffer; 20 mM Hepes (pH 7.5), 10 mM MgCl₂, 1 mM EGTA, 0.02% Brij35, 0.02 mg/ml BSA, 0.1 mM Na₃VO₄, 2 mM DTT, 1% DMSO

It should be noted that required cofactors are added individually to each kinase reaction.

Reaction Procedure:

1. Prepare indicated substrate in freshly prepared Base Reaction Buffer as described above
2. Deliver any required cofactors to the substrate solution
3. Deliver indicated kinase into the substrate solution and gently mix
4. Deliver compounds in DMSO into the kinase reaction mixture
5. Deliver ³³P-ATP (specific activity 0.01 μCi/μl final) into the reaction mixture to initiate the reaction
6. Incubate kinase reaction for 120 min. at room temperature
7. Reactions are spotted onto P81 ion exchange paper (Whatman # 3698-915)
8. Wash filters extensively in 0.75% Phosphoric acid

Kinase information:

CK1d - Genbank Accession # NP_620693

Recombinant human full-length construct. GST-tagged, expressed in insect cells.

Final concentration in assay = 4 nM

Substrate: CK1tide

Substrate sequence: [KRRRAL[pS]VASLPGL]

Final substrate concentration in assay = 20 μM

It should be noted that no additional cofactors are added to the reaction mixture.

Compounds 30, 288, 314, 324-325, 336, 374, 391, 405, 615-616, 626, 705, 740, 753-754, 756, 759, 770, 784, 808, 819, 833, 844, 847, 869, 872, 875, 933, 952, 955, 969, 987, 990

and 999 were tested in the CK1δ inhibition assay and exhibited inhibition of greater than 5%.

In particular, compounds 324-325, 405, 754, 847, 952, 987, 990 and 999 exhibited
5 inhibition of greater than 50%.

Yet more particularly, compounds 324, 952, 987, 990 and 999 exhibited inhibition of greater than 90%.

10 **2. Measurement of Compound Effect on Ck1d-mediated Tau Phosphorylation**

The *in vivo* phosphorylation of Tau protein is complex with a number of putative protein kinases involved. It is widely accepted that the kinases GSK3b and CDK5 are significant players in the generation of PHF Tau, the pathogenic form found in neurofibrillary tangles in Alzheimer's disease. More recently, there has been growing evidence supporting the
15 role of other kinases, particularly CK1δ in Tau hyperphosphorylation *in vivo*. Hanger et al. 2007 (J. Biol. Chem. 282, 23645-23654) identified 37 phosphorylation sites in human PHF Tau and were able to recapitulate these *in vitro* using recombinant tau and various purified kinase preparations. These studies identified that certain sites were uniquely phosphorylated by CK1δ and that certain other sites required CK1δ and another kinase
20 with CK1δ providing upstream phosphorylation to render the targeted site available for the second kinase. Thus, to assess whether candidate compounds selective inhibit CK1δ activity either directly or through blocking its priming for other kinases a number of different screens have been developed. The general concept of these screens is provided in WO2005/001114.

25

To measure the effect of putative CK1δ inhibitors on the levels of CK1δ -mediated phosphorylation selected-reaction monitoring assays were performed that provide quantitative relative measurement of phosphate group occupancy at specific sites in transgenic human and endogenous murine forms of Tau.

30

The PhosphoTau SRM V2 assay measures total tau and relative phosphorylation levels at five of the most commonly studied sites on Tau and was obtained from Proteome Sciences plc (Cobham, England). None of the sites in the V2 assay is uniquely phosphorylated by CK1δ and there is a possibility that compound-induced inhibition of
35 phosphorylation measured by this method may be achieved through promiscuous inhibition of other kinases such as GSK3b and/or CDK5. To address this limitation,

Proteome Sciences has developed a V3 assay that measures total tau and two sites that are exclusively phosphorylated by CK1δ in addition to four others that have been shown to be phosphorylated *in vitro* by at least one other Tau kinase in addition to CK1δ. Table 1 lists the various sites covered and the candidate Tau kinases reported in Hanger et al. (2007).

Table 1: Tau phosphorylation sites covered by Tau Phosphorylation SRM V2 and V3 assays

Site number	Candidate Kinases
Assay V2	
Ser181	GSK3b
Ser199	CK2, GSK3b, PKA
Thr231	GSK3b, PKA
Ser262	CK1δ, GSK3b, PKA
Ser396	CK1δ, CK2, GSK3b
Assay V3	
Ser46	CK1δ, GSK3b
Thr50	CK1δ, GSK3b
Ser113*	CK1δ
Ser396	CK1δ, CK2, GSK3b
Ser404	CK1δ, CK2, GSK3b
Ser433*	CK1δ

Numbering based on human 2N4R tau.

* - CK1d unique site

SH-SY5Y-TMHT Cell Line

- 15 The SH-SY5Y-TMHT cell line (JSW Life Sciences, Graz, Austria) represents an *in vitro* model of tauopathy. The cell line is created by stably transfecting the human neuroblastoma derived SH-SY5Y cell line with a vector containing the full length human 2N4R Tau isoform which carries two common disease associated mutations (V337M/R406W).
- 20 In recent studies (Flunkert et al, Elevated Levels of Soluble Total and Hyperphosphorylated Tau Result in Early Behavioral Deficits and Distinct Changes in Brain Pathology in a New Tau Transgenic Mouse Model, 2013, Neurodegener Dis 11:194-205; Loeffler et al, Stable Mutated tau441 Transfected Sh-SY5Y Cells as Screening Tool of Alzheimer's Disease Drug Candidates, 2012, J. Mol. Neurosci. 47(1):192-203) both the SH-SY5Y-TMHT cell line and a transgenic mouse line carrying the same human transgene were shown to express high levels of human Tau which becomes hyperphosphorylated at multiple epitopes previously demonstrated to be phosphorylated

in various human tauopathies including Alzheimer's disease. Furthermore, in SH-SY5Y-TMHT cells exposed to different kinase inhibitors, including JNK-Inhibitor SP600125, and CK1 inhibitor IC261 levels of Tau phosphorylation at key pathogenic sites were reduced in patterns consistent with the known site-specificity of the targeted kinase. Thus, the SH-SY5Y-TMHT cell line is ideally suited to the screening of novel Tau kinase inhibitors.

Compound screening in SH-SY5Y-TMHT Cells

SH-SY5Y-TMHT cells are kept in culture medium (DMEM medium, 10% FCS, 1% NEAA, 1% L-Glutamine, 100 µg/ml Gentamycin, 300 µg/ml Geneticin G-418) for 2 days until 80-90% confluency. Cells are then differentiated in culture medium supplemented with 10 µM retinoic acid (RA) for 7 days changing medium every 2 to 3 days. Differentiated cells are seeded onto 6-well plates and 96-well plates at a cell density of 1.25×10^6 and 8×10^5 cells per well, respectively. On day 8 post-differentiation, the test compounds, reference compounds and vehicle control were added to the culture medium. After 6 h of compound exposure one plate of cells is subjected to a MTT assay to evaluate the effect of test and reference items on cell viability. Remaining wells are washed once with cold PBS and harvested in 300 µl RIPA-Buffer [50 mM Tris pH 7.4, 1% Nonident P40, 0.25% Na-deoxycholate, 150 mM NaCl, 1 mM EDTA, 1 µM NaF, 1 µM Na-ortho-vanadate, 80 mM Glycerophosphate, supplemented with freshly added protease (Calbiochem) and phosphatase (Sigma) inhibitor cocktail]. The cell suspension is transferred into a 1.5 ml tube and additionally lysed by sonication on ice. An aliquot of 20 µl is taken for the determination of the protein concentration (BCA assay). Subsequently, the lysates are snap frozen and stored at -80°C until shipment.

Two independent experiments in three (four) technical replicates are performed as depicted in Table 2.

Table 2

Experiment	Cells	Treatment	Concentration	Evaluation	
ExpA	SH-SY5Y-TMHT	Vehicle	-	MTT	BSA TauP V2 TauP V3
		Compound 324	10-5-1-0.5-0.1-0.05 µM		
		Compound 987	10-5-1-0.5-0.1-0.05 µM		
		PF670462	1-0.5-0.1 µM		

ExpB	SH-SY5Y-TMHT	Vehicle	-	MTT	BSA TauP V2 TauP V3
		Compound 324	10-5-1-0.5-0.1-0.05 μ M		
		Compound 987	10-5-1-0.5-0.1-0.05 μ M		
		PF670462	1-0.5-0.1 μ M		

Cell Viability Testing

To determine compound activity, it is necessary to control for potential cell toxicity of all molecules. Viability of cultures is determined by the MTT assay. This assay allows the measurement of the mitochondrial dehydrogenase activity which reduces yellow MTT to dark blue formazan crystals. Since this reaction is catalyzed in living cells only this assay is used for the determination of cell viability. MTT solution is added to each well in a final concentration of 0.5 mg/ml. After 2 hours, the MTT containing medium is aspirated. Cells are lysed in 3% SDS and the formazan crystals are dissolved in isopropanol/HCl. Optical density is measured with a plate-reader at wavelength 570 nm. Cell survival rate is expressed as optical density (OD). Values are calculated as percent of control values.

Quantitative determination of total protein content

Prior to assessment of specific Tau phosphorylation status the concentration of total protein in each cell lysate is determined using a standard BCA assay (Pierce Biotechnology, Rockford, USA). Briefly, 20 μ l of cell lysate was used in the assay according to the manufacturer's instructions.

Quantitative determination of total Tau & phosphorylated Tau

Mass spectrometric assays

Total cell lysates from TMHT cell lines treated with Compound 324, Compound 987, PF670462 and relevant vehicle control respectively are first subjected to 1-dimensional SDS-PAGE to purify the protein fraction. Stacking gels are loaded with approximately 100 μ g total protein based on BCA assay results. Gels are run until the total protein content forms a single discrete band in the stacking gel. Each protein band is then cut from the gel and digested with either trypsin or Asp-N and analysed using the PhosphoTau SRM assay V2 or V3 respectively. Each assay method quantifies the phosphorylation in pre-clinical material using a triple quadrupole mass spectrometer (TSQ Vantage, Thermo

Scientific, Hemel Hempstead, UK). Prior to SRM analysis phosphopeptides and pre-clinical samples were resolved by RP-chromatography (XBridge column, Waters, Manchester, UK) over a 9 minute gradient 0-30% ACN (buffer A; 0.1% FA, buffer B; ACN, 0.1% FA). Light and heavy versions for each peptide and phosphopeptide were
5 monitored by several SRM transitions, using optimised S Lens values and collision energy settings. The area under the SRM LC peak was used to quantitate the amount of analyte present in each cell lysate as a single point reference to the signal of the heavy peptide spike. An 11 point calibration curve of light phosphopeptides with each point in the curve spiked with 100 fmol heavy phosphopeptides was also produced to determine assay
10 characteristics (LOD, LOQ, precision and accuracy). For each specified tau population, the endogenous level of each tau phosphopeptide was quantified against its calibration curve (0.25-1000 fmol on column). Prior to LC-SRM analysis each tau population was spiked with 100fmol of the heavy phosphopeptide standards. All data was processed using Pinpoint software (Thermo Scientific) and results reported as pg phospho-
15 peptide/ μ g total protein.

Western Blotting

Lysates of treated cells were prepared in Laemmli buffer and 10 μ g loaded into each lane of a 10% Nu-PAGE gel (Invitrogen, UK) . Samples were run until the coomassie blue dye
20 front was within 1 cm of the bottom of the gel. The separated proteins were transferred onto nitrocellulose and blots developed using antibodies specific for total tau (Polyclonal Rabbit Anti-Human Tau, Dako, UK (cat # A0024)) and phospho-Threonine 231 (Tau (Phospho-Thr231) Antibody, Signalway Antibody, USA (cat # 11110)) respectively. In each case the bound antibody was detected using ECL Rabbit IgG, HRP-Linked (from
25 donkey) (GE Healthcare, UK (cat # NA934))

Results

Effect of test and reference compounds on the cell viability of SH-SY5Y-TMHT cells

30 Cell viability was determined in differentiated SH-SY5Y-TMHT cells by the MTT assay. Test and reference compounds were applied in a concentration range from 0.05 μ M to 10 μ M and from 0.1 μ M to 1 μ M, respectively. Upon 6 h of treatment, cell viability was evaluated. Figure 1 shows the effect of Compound 324 on the cell viability of SH-SY5Y-TMHT cells wherein the graph represents effect of Compound 324 on cell viability of SH-
35 SY5Y-TMHT cells in % of the vehicle control (VC, white bar). Statistical significance is indicated by * <0.05 , ** <0.01 , *** <0.001 as determined by One-Way ANOVA. Data are

shown from two independent experiments as group mean \pm SEM (n=8). It can be seen from Figure 1 that Compound 324 exhibited a protective effect on the cell viability of SH-SY5Y-TMHT cells in a dose dependent manner although the effect was only statistically significant at a concentration of 10 μ M. Figure 2 shows the effect of Compound 987 on the cell viability of SH-SY5Y-TMHT cells wherein the graph represents effect of Compound 987 on cell viability of SH-SY5Y-TMHT cells in % of the vehicle control (VC, white bar). Statistical significance is indicated by * <0.05 , ** <0.01 , *** <0.001 as determined by One-Way ANOVA. Data are shown from two independent experiments as group mean \pm SEM (n=8). It can be seen from Figure 2 that Compound 987 decreased the cell viability in the lower and higher concentration range. At a concentration of 1 and 0.5 μ M no cytotoxic effect was observed. Figure 3 shows the effect of PF670462 on the cell viability of SH-SY5Y-TMHT cells wherein the graph represents effect of PF670462 on cell viability of SH-SY5Y-TMHT cells in % of the vehicle control (VC, white bar). Statistical significance is indicated by * <0.05 , ** <0.01 , *** <0.001 as determined by One-Way ANOVA. Data are shown from two independent experiments as group mean \pm SEM (n=8). It can be seen from Figure 3 that the reference compound PF 670462 only displayed a significant protective effect on the cell viability of SH-SY5Y-TMHT cells at a concentration of 0.5 μ M.

Protein determination of SH-SY5Y-TMHT cells following treatment

Protein concentration of cell lysates of the treated SH-SY5Y-TMHT cells was determined using a standard BCA assay. Protein amount was determined from all samples in duplicates. The protein concentration of the samples was in the expected range according to the amount of cells seeded per 12-well plate ranging between 150 – 350 μ g/ml.

Determination of compound treatment effect on specific phosphorylation sites

Mass Spectrometric Assay

Testing of SH-SY5Y-TMHT cell lysates was performed using the PhosphoTau SRM assay V2 and V3. When the relative level of phosphorylation at each site is compared with the ratio in vehicle treated controls there was a distinct reduction in the level of phosphopeptide in cells treated with Compound 324 (data shown for 10 μ M) and Compound 987 (data shown for 10 μ M). An example showing reduction of phosphorylation on Serine 396 is shown in Figure 4. This Figure shows mass spectrometric determination of CK1d-selective compounds on phosphorylation of Serine 396 in SH-SY5Y-TMHT cells. Panel A shows cells treated with Vehicle Control (VC) or Compound 324 (T.I.1_10 μ M)

and Panel B shows cells treated with Vehicle Control (VC) or Compound 987 (T.I.2_10 μ M).

In cells exposed to the vehicle control approximately 83% of Tau is phosphorylated at S396. Treatment with 10 μ M Compound 324 reduced this to 38% whilst 10 μ M Compound 987 reduced pS396 levels to 24%. These results confirm the inhibition of pS396 by CK1d selective reagents.

Western Blot Assay

Levels of total Tau and Tau phosphorylated at Threonine 231 in SH-SY5Y-TMHT cell lysates treated with vehicle control, Compound 394 (10 μ M), Compound 987 (10 μ M) and PF670462 (5 μ M) were quantified by Western Blotting. Figure 5 shows the Western Blot measurement of pT231 (panel A) and total Tau (panel B) levels in SH-SY5Y-TMHT cells treated with selective CK1d inhibitors. As shown in Figure 5, all three compounds reduced the detectable level of pT231 in Tau protein whereas this epitope was strongly present in vehicle-treated cells. There was no significant difference in the detectable levels of total Tau between the preparations other than for the PF670462-treated lysate which appeared to contain marginally less total Tau than the others. These results confirm the inhibition of pT231 by CK1d selective reagents.

CLAIMS

1. A compound 5-(1,3-benzoxazol-2-yl)-4- (pyridin-4-yl)pyrimidin-2-amine or a pharmaceutically acceptable salt or solvate thereof.
- 5 2. A pharmaceutical composition comprising a compound 5-(1,3-benzoxazol-2-yl)-4- (pyridin-4-yl)pyrimidin-2-amine or a pharmaceutically acceptable salt or solvate thereof, and an excipient.
- 10 3. The compound of claim 1 or the pharmaceutical composition of claim 2 for use in therapy.
4. The compound of claim 1 or the pharmaceutical composition of claim 2 for use as a casein kinase delta (CK1δ) inhibitor in the treatment of a neurodegenerative disorder.
- 15 5. The compound or pharmaceutical composition of claim 4, wherein the neurodegenerative disorder is a tauopathy.
6. The compound or pharmaceutical composition of claim 5, wherein the tauopathy is selected from the group consisting of Alzheimer's disease, frontotemporal dementia with Parkinsonism linked to chromosome 17 (FTDP-17), progressive supranuclear palsy (PSP), Pick's disease, corticobasal degeneration, multisystem atrophy (MSA), neurobasal degeneration with iron accumulation, type 1 (Hallervorden-spatz), argyrophilic grain dementia, Down's syndrome, diffuse neurofibrillary tangles with calcification, dementia pugilistica, Gerstmann-Straussler-Scheinker disease, myotonic dystrophy, Niemann-Pick disease type C, progressive subcortical gliosis, prion protein cerebral amyloid angiopathy, tangle only dementia, postencephalitic parkinsonism, subacute sclerosing panencephalitis, Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis/parkinsonism-dementia complex, non-Guamanian motor neuron disease with neurofibrillary tangles/dementia, and
- 20 25 30 Parkinson's disease.
7. The compound or pharmaceutical composition of claim 5, wherein the tauopathy comprises Alzheimer's disease.

1/5




FIGURE 1

2/5




FIGURE 2

3/5




FIGURE 3

4/5




FIGURE 4

5/5




FIGURE 5

