



HU000032948T2

(19) **HU****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**(11) Lajstromszám: **E 032 948**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 12 760456**(22) A bejelentés napja: **2012. 03. 23.**

(96) Az európai bejelentés bejelentési száma:

EP 20120760456

(97) Az európai bejelentés közzétételi adatai:

EP 2688403 A1 **2012. 09. 27.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2688403 B1 **2017. 05. 31.**(51) Int. Cl.: **C07D405/14** (2006.01)**C07D311/96** (2006.01)**C07D405/04** (2006.01)**C07D409/14** (2006.01)**C07D409/12** (2006.01)**A61K 31/35** (2006.01)**C07D405/12** (2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 12/030327

(87) A nemzetközi közzétételi szám:

WO 12129495

(30) Elsőbbségi adatok:

201161466809 P **2011. 03. 23.** **US****201261596808 P** **2012. 02. 09.** **US**

(73) Jogosult(ak):

Trevena, Inc., King Of Prussia, PA 19406 (US)

(72) Feltaláló(k):

YAMASHITA, Dennis, Wayne Pennsylvania 19087 (US)**GOTCHEV, Dimitar, Hatboro Pennsylvania 19040 (US)****PITIS, Philip, North Wales Pennsylvania 19454 (US)****CHEN, Xiao-Tao, Furlong Pennsylvania 18925 (US)****LIU, Guodong, Jamison Pennsylvania 18929 (US)****YUAN, Catherine C.K., King of Prussia Pennsylvania****19406 (US)**

(74) Képviselő:

SBGK Szabadalmi Ügyvivői Iroda, Budapest

(54)

Opioid receptor ligandumok és eljárások azok alkalmazására és előállítására

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 688 403 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
31.05.2017 Bulletin 2017/22

(21) Application number: **12760456.9**

(22) Date of filing: **23.03.2012**

(51) Int Cl.:
C07D 405/14 ^(2006.01) **C07D 405/04** ^(2006.01)
C07D 405/12 ^(2006.01) **C07D 409/12** ^(2006.01)
C07D 409/14 ^(2006.01) **C07D 311/96** ^(2006.01)
A61K 31/35 ^(2006.01)

(86) International application number:
PCT/US2012/030327

(87) International publication number:
WO 2012/129495 (27.09.2012 Gazette 2012/39)

(54) **OPIOID RECEPTOR LIGANDS AND METHODS OF USING AND MAKING SAME**

OPIOIDREZEPTORLIGANDEN UND VERFAHREN ZU IHRER VERWENDUNG UND
HERSTELLUNG

LIGANDS DE RÉCEPTEURS OPÏOIDES, ET LEURS PROCÉDÉS D'UTILISATION ET DE
PRODUCTION

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

(30) Priority: **23.03.2011 US 201161466809 P**
09.02.2012 US 201261596808 P

(43) Date of publication of application:
29.01.2014 Bulletin 2014/05

(73) Proprietor: **Trevena, Inc.**
King Of Prussia, PA 19406 (US)

(72) Inventors:
• **YAMASHITA, Dennis**
Wayne
Pennsylvania 19087 (US)
• **GOTCHEV, Dimitar**
Hatboro
Pennsylvania 19040 (US)
• **PITIS, Philip**
North Wales
Pennsylvania 19454 (US)

- **CHEN, Xiao-Tao**
Furlong
Pennsylvania 18925 (US)
- **LIU, Guodong**
Jamison
Pennsylvania 18929 (US)
- **YUAN, Catherine C.K.**
King of Prussia
Pennsylvania 19406 (US)

(74) Representative: **Bassil, Nicholas Charles**
Kilburn & Strode LLP
20 Red Lion Street
London WC1R 4PJ (GB)

(56) References cited:
US-A1- 2006 217 372 US-A1- 2006 217 372
US-A1- 2009 247 561 US-A1- 2010 280 058
US-A1- 2010 280 058

- **POGOZHEVA ET AL.: 'Homology Modeling of
Opioid Receptor-Ligand Complexes Using
Experimental Constraints.' THE AAPS JOURNAL,**
[Online] vol. 7, no. 2, 2005, pages E434 - E448,
XP008142404 Retrieved from the Internet:
<URL:http://www.aapsj.org/artides/aapsj0702
/aapsj070243/aapsj070243.pdf>

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**FIELD**

5 **[0001]** This disclosure relates to a family of compounds acting as opioid receptor ligands. Such compounds may provide therapeutic benefit in the treatment of pain.

BACKGROUND

10 **[0002]** Opioid receptors (ORs) mediate the actions of morphine and morphine-like opioids, including most clinical analgesics. Three molecularly and pharmacologically distinct opioid receptor types have been described: δ , κ and μ . Furthermore, each type is believed to have sub-types. All three of these opioid receptor types appear to share the same functional mechanisms at a cellular level. For example, activation of the opioid receptors causes inhibition of adenylate cyclase, and recruits β -arrestin.


15 **[0003]** When therapeutic doses of morphine are given to patients with pain, the patients report that the pain is less intense, less discomforting, or entirely gone. In addition to experiencing relief of distress, some patients experience euphoria. However, when morphine in a selected pain-relieving dose is given to a pain-free individual, the experience is not always pleasant; nausea is common, and vomiting may also occur. Drowsiness, inability to concentrate, difficulty in mentation, apathy, lessened physical activity, reduced visual acuity, and lethargy may ensue.

20 **[0004]** There is a continuing need for new OR modulators to be used as analgesics. There is a further need for OR agonists as analgesics having reduced side effects. There is a further need for OR agonists as analgesics having reduced side effects for the treatment of pain, immune dysfunction, inflammation, esophageal reflux, neurological and psychiatric conditions, urological and reproductive conditions, medicaments for drug and alcohol abuse, agents for treating gastritis and diarrhea, cardiovascular agents and/or agents for the treatment of respiratory diseases and cough. US 2009/247561
 25 describes compounds that have affinity to the μ -opioid receptor and the ORL1-receptor, methods for their production, medications containing these compounds and the use of these compounds for the treatment of pain and other conditions.

SUMMARY

30 **[0005]** This disclosure describes opioid receptor (OR) ligands. It also describes methods of modulating opioid receptor activity using the compositions described herein. Certain compositions described herein act as opioid receptor agonists. Other compositions described herein act as opioid receptor antagonists.

[0006] Also described herein are compounds having the structure of Formula I:



I

55 **[0007]** In the structure above, variables A_1 , A_2 , A_3 , A_4 , A_5 , B_1 , B_2 , B_3 , B_4 , B_5 , and D_1 can be selected from the respective groups of chemical moieties later described. OR ligand derivatives and mimetics are also provided. Also provided are processes for preparing these compounds.

[0008] Also described herein are pharmaceutical compositions comprising one or more compounds as described herein a pharmaceutically acceptable carrier. Naturally, the compounds described herein can be employed in any form,


such as a solid or solution (e.g., aqueous solution) as is described further below. The compounds described herein, for example, can be obtained and employed in a lyophilized form alone or with suitable additives.

[0009] Also described herein are methods for treating pain and pain-related disorders. Such a method would comprise administering a therapeutically effective amount of one or more compounds described herein to a subject in need thereof.

Detailed Description

[0010] This disclosure describes a family of compounds, OR ligands, with a unique profile. The compounds described herein act as agonists or antagonists of opioid receptor (OR)-mediated signal transduction. The ligands of these receptors can be used to treat pathologies associated with ORs including pain and pain related disorders. Aspects of the invention for which protection is sought are as defined in the claims.

[0011] Compounds also comprise Formula I:



wherein: A₁ is null, CH₂, CHR₁, CR₁R₂, CH, CR₁, O, S, SO, SO₂, NH or NR₁; A₂ is null, CH₂, CHR₅, CR₅R₆, CH, CR₅, O, S, SO, SO₂, NH or NR₅; A₃ is null, CH₂, CHR₇, CR₇R₈, O, S, SO, SO₂, NH, NR₇, CH or CR₇; A₄ is null, CH₂, cycle of the formula C(CH₂)_n, where n = 2-5, CHR₉, CR₉R₁₀, O, S, SO, SO₂, NH, NR₉, CH or CR₉; and A₅ is null, CH₂, CHR₁₁, CR₁₁R₁₂, CH₂CH₂, CHR₁₁CH₂, CH₂CHR₁₁, CHR₁₁CHR₁₂, O, S, SO, SO₂, NH, NR₁₁, CH or CR₁₁.

[0012] No more than 2 out of 5 A_a (specifically A₁, A₂, A₃, A₄, A₅) can be null at the same time. The number of heteroatoms from A₁ to A₅ cannot exceed 2 at the same time, and O-O, S-O; S-S; S-N fragments in the ring structure are excluded from this composition.

[0013] The ring containing A₁, A₂, A₃, A₄, A₅ and the carbon connected to D₁ can be fused with another ring, such as benzene, pyridine, pyrimidine, furan, thiophene or pyridazine, but not limited to these examples, where the resulting bicyclic system is chemically stable and synthetically accessible. It is also understood that the above-mentioned fused rings could be multiply substituted with cyano, halogen, alkyl, branched alkyl, halogenated alkyl, hydroxyl, alkoxy, formyl, acetyl, amino, alkylamino, dialkylamino, mercaptanyl, alkylmercaptanyl, and other small substitution groups. The bonds between A₁ and A₂, A₂ and A₃, A₃ and A₄, A₄ and A₅ can independently be a single bond or a double bond. The bonds between A₁ and A₂, A₂ and A₃, A₃ and A₄, A₄ and A₅ cannot be a double bond at the same time.

[0014] A₂ and A₄ can be connected by a carbon bridge. Examples of such a bridge include -CH₂-, and -CH₂CH₂-.


[0015] B₁ is CH₂, CHR₁₃, CR₁₃R₁₄, O, S, SO, SO₂, NH, NR₁₃, CR₁₃ or CO. B₂ is CH₂, CHR₁₅, CR₁₅R₁₆, CR₁₅ or CO. B₃ is H, alkyl, branched alkyl, halogenated alkyl, aryl, arylalkyl, alkylcarbonyl, branched alkylcarbonyl, arylcarbonyl, alkoxy, alkoxy, or alkylsulfonyl. B₄ is null, C₁-C₆ alkyl, CH₂, CH₂CH₂, CHR₁₉, CR₁₉R₂₀ or CO. When B₄ is an alkyl one or more of the hydrogens can be replaced with a deuterium. B₅ is alkyl, branched alkyl, halogenated alkyl, carbocycle-substituted alkyl, aryl, carbocycle or arylalkyl.

[0016] Aryl, heteroaryl and carbocycle (non-aromatic)/heterocycle (non-aromatic with 1-3 heteroatoms, including O, N, S) are either unsubstituted, or substituted with small substitution groups. Small substitution groups can be cyano, halogen, alkyl, branched alkyl, halogenated alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino, mercaptanyl, alkylmercaptanyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, alkylcarbonyl, alkoxy, alkoxy, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, aryl, arylalkyl, carbocycle or carbocycle-alkyl. The small substitution groups may be selected from F, Cl, Br, CH₃, CH₂CH₃, CH₂F, CHF₂, CF₃, n-Pr, n-Bu, i-Bu, sec-Bu, i-Pr, t-Bu, CN, OH, OMe, OEt, O-iPr, OCF₃, NH₂, NHMe, NMe₂, methoxycarbonyl, methanesulfonyl, Ph, benzyl, MeSO₂, formyl, and acetyl.


[0017] Carbocycle may contain double bonds, but they should not be aromatic.

[0018] D₁ is an aryl or heteroaryl group or a carbocycle.


[0019] An aryl group is either a monocyclic aromatic group or a bicyclic aromatic group. A heteroaryl group may contain heteroatoms in the aromatic group. The following structures are some examples of representative aryl and heteroaryl groups, but the aryl and heteroaryl groups are not limited to those examples:




5




10




15




20



25




30




35

40





45 **[0020]** Carbocycle is either a monocyclic or a bicyclic non-aromatic ring system. The following structures are some examples of representative carbocycle, but the carbocycle is not limited to those examples:

50



55





wherein X_1 and X_2 in the carbocycle examples are independently O, S, N, NH or NR_{18} .

[0021] The aryl and heteroaryl groups can be independently mono or multiply substituted with cyano, halogen, alkyl, branched alkyl, halogenated alkyl, hydroxyl, alkoxy, amino, alkylamino, dialkylamino, mercaptanyl, alkylmercaptanyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, aryl, arylalkyl, carbocycle, carbocycle-alkyl, and/or other small substitution groups. In some instances the small substitution groups are selected from F, Cl, Br, CH_3 , CH_2CH_3 , CH_2F , CHF_2 , CF_3 , n-Pr, n-Bu, i-Bu, sec-Bu, i-Pr, t-Bu, CN, OH, OMe, OEt, O-iPr, OCF_3 , NH_2 , $NHMe$, NMe_2 , methoxycarbonyl, methanesulfonyl, Ph, benzyl, formyl, and acetyl.

[0022] D_1 is an aryl, heteroaryl or a carbocycle.

[0023] R_1 , R_2 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{18} , R_{19} , and R_{20} are independently: cyano, halogen, hydroxyl, alkoxy, alkyl, branched alkyl, halogenated alkyl, branched halogenated alkyl, aryl, arylalkyl, carbocycle, carbocycle-alkyl, alkylcarbonyl, branched alkylcarbonyl, halogenated alkylcarbonyl, branched halogenated alkylcarbonyl, arylcarbonyl or alkoxy carbonyl. R_1 , R_2 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{18} , R_{19} , and R_{20} may be independently F, Cl, Br, CH_3 , CH_2CH_3 , CH_2F , CHF_2 , CF_3 , n-Pr, n-Bu, i-Bu, sec-Bu, i-Pr, t-Bu, CN, OH, OMe, OEt, O-i-Pr, methoxycarbonyl, phenyl, benzyl, formyl or acetyl, whenever the resulting structure is stable.


[0024] R_1 and R_2 , R_5 and R_6 , R_7 and R_8 , R_9 and R_{10} , R_{11} and R_{12} , R_{13} and R_{14} , R_{15} and R_{16} , R_{19} and R_{20} , or R_{15} and R_{19} can form a monocycle.

[0025] Me is methyl; Et is ethyl; i-Pr is i-propyl; t-Bu is t-butyl; Ph is phenyl.

[0026] The following compounds can be excluded from the genus of compounds:

- 1) 2-[(2-[2-Ethyl-2-methyl-4-(4-methylphenyl)oxan-4-yl]ethyl)amino)methyl]phenol


5



10

2) 2-[(2-[2-Ethyl-4-(4-fluorophenyl)-2-methyloxan-4-yl]ethyl]amino)methyl]phenol


15



20

3) {2-[2,2-Dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(4-methoxyphenyl)methyl]amine


25



30

4) {2-[(4S*, 4R*)-2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(1R)-1-phenylethyl]amine


35



40

5) {2-[(4S*, 4R*)-2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(1S)-1-phenylethyl]amine


45




50

6) Benzyl({2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl})amine


55




7) 2-[(2-[2-Ethyl-4-(4-fluorophenyl)-2-methyloxan-4-yl]ethyl)amino)methyl]phenol




8) Benzyl[2-(2,2-dimethyl-4-phenyloxan-4-yl)ethyl]amine




9) {2-[2-Ethyl-4-(4-fluorophenyl)-2-methyloxan-4-yl]ethyl}[(4-methoxyphenyl)methyl]amine



10) [(3,4-Dimethoxyphenyl)methyl]{2-[4-(4-fluorophenyl)-2,2-dimethyloxan-4-yl]ethyl}amine




11) {2-[4-(4-Methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl}(1-phenylethyl)amine



12) [(4-Chlorophenyl)methyl]{2-[4-(4-methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl}amine


5



10

13) Benryl({2-[2-ethyl-4-(2-methoxyphenyl)-2-methyloxan-4-yl]ethyl})amine


15



20

14) [(3,4-dimethoxyphenyl)methyl]({2-[2-ethyl-4-(2-methoxyphenyl)-2-methyloxan-4-yl]ethyl})amine


25



30

15) 4-[(2-[4-(2-Methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl)amino)methyl]-N,N-dimethylaniline


35



40

16) Benzyl({2-[4-(4-fluorophenyl)-2,2-dimethyloxan-4-yl]ethyl})amine


45




50

17) {2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}(1-phenylethyl)amine


55




18) 2-(2,2-Dimethyl-4-phenyloxan-4-yl)ethyl [(4-methoxyphenyl)methyl]amine



19) 2-[4-(4-Fluorophenyl)-2,2-dimethyloxan-4-yl]ethyl [(4-methoxyphenyl)methyl]amine



20) [(3,4-Dimethoxyphenyl)methyl][2-(2,2-dimethyl-4-phenyloxan-4-yl)ethyl]amine




[0027] This disclosure also describes compounds having the structure of Formula II-1 and II-2:

5

10


15



II-1

20

25



II-2

30

wherein A_2 is CH_2 , CHR_5 , CR_5R_6 ; A_4 is CH_2 , CHR_9 , CR_9R_{10} or a cycle of the formula $C(CH_2)_n$, where $n = 2-5$.

[0028] Further R_5 , R_6 , R_9 , and R_{10} are independently CH_3 , CH_2CH_3 , CH_2F , CHF_2 , CF_3 , n -Pr, n -Bu, i -Bu, sec -Bu, i -Pr, t -Bu, or phenyl. Further, R_5 and R_6 , or R_9 and R_{10} can form a monocyclic carbocycle.

[0029] A_2 and A_4 can be connected by a carbon bridge. This bridge can be $-CH_2-$ or $-CH_2CH_2-$.

35

[0030] Further B_3 is selected from the following: H, alkyl, branched alkyl, aryl, arylalkyl, alkylcarbonyl, branched alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, and alkylsulfonyl. In some instances B_3 is C_1 - C_5 alkyl. In some instances B_3 is H.

[0031] Further B_4 is null, C_1 - C_6 alkyl, CH_2 , CH_2CH_2 , CHR_{19} , $CR_{19}R_{20}$ or CO. Further, R_{19} and R_{20} can form a monocycle of the formula $(CH_2)_n$, where $n = 2-4$. B_5 is alkyl, branched alkyl, carbocycle, carbocycle-substituted alkyl, aryl or arylalkyl.

[0032] Further D_1 is an aryl or heteroaryl. Examples of the aryl and heteroaryl groups are shown above.

40


[0033] Each aryl and heteroaryl group can be independently mono or multiply substituted with F, Cl, Br, CH_3 , CH_2CH_3 , CH_2F , CHF_2 , CF_3 , n -Pr, n -Bu, i -Bu, sec -Bu, i -Pr, t -Bu, CN, OH, OMe, OEt, O- i Pr, OCF_3 , NH_2 , $NHMe$, NMe_2 , methoxycarbonyl, Ph, benzyl, formyl, or acetyl. That is, each aryl and heteroaryl group may be multiply substituted with the same substituent (i.e., 2 chloro groups) or just be multiply substituted, albeit with different groups (e.g. an aryl or heteroaryl group with 1 chloro and 1 methyl group would be considered multiply substituted).

[0034] This disclosure also describes compounds having the structure of Formula III:

45

50

55



wherein A_2 is CH_2 , CHR_5 or CR_5R_6 ; A_4 is CH_2 , CHR_9 , CR_9R_{10} or a cycle of the formula $C(CH_2)_n$, where $n = 2-5$.

[0035] Further R_5 , R_6 , R_9 , and R_{10} are independently CH_3 , CH_2CH_3 , CH_2F , CHF_2 , CF_3 , n -Pr, n -Bu, i -Bu, sec -Bu, i -Pr, t -Bu, or phenyl. R_5 and R_6 , or R_9 and R_{10} can form a monocyclic carbocycle.

[0036] A_2 and A_4 can be connected by a carbon bridge. The bridge can be $-CH_2-$ or $-CH_2CH_2-$.



[0037] Further B_3 is selected from H, alkyl, branched alkyl, aryl, arylalkyl, alkylcarbonyl, branched alkylcarbonyl, arylcarbonyl, alkoxy carbonyl or alkylsulfonyl.


[0038] Further B_4 is null, C_1 - C_6 alkyl, CH_2 , CH_2CH_2 , CHR_{19} , $CR_{19}R_{20}$ or CO. Further, R_{19} and R_{20} can form a monocycle of the formula $(CH_2)_n$, where $n = 2-4$. B_5 is alkyl, branched alkyl, carbocycle, carbocycle-substituted alkyl, aryl or arylalkyl.

[0039] Further D_1 is an aryl or heteroaryl. Examples of the aryl and heteroaryl groups are shown above.

[0040] The aryl and heteroaryl groups can be mono or multiply substituted with F, Cl, Br, CH_3 , CH_2CH_3 , CH_2F , CHF_2 , CF_3 , n -Pr, n -Bu, i -Bu, sec -Bu, i -Pr, t -Bu, CN, OH, OMe, OEt, O- i Pr, OCF_3 , NH_2 , $NHMe$, NMe_2 , methoxycarbonyl, Ph, benzyl, formyl, or acetyl.

[0041] This disclosure also describes compounds having the structure of Formula IV-1, IV-2, or IV-3, V, or VI:





IV-3

wherein R_{21} and R_{22} are, independently, H or CH_3 ; A_4 is CH_2 , CR_9R_{10} or a cycle of the formula $\text{C}(\text{CH}_2)_n$, where $n = 2-5$.


[0042] Further R_9 and R_{10} are independently CH_3 or CH_2CH_3 .

[0043] Further B_3 is H, $\text{C}_1\text{-C}_6$ alkyl or branched alkyl.

[0044] Further B_4 is null, $\text{C}_1\text{-C}_6$ alkyl, CH_2 , CH_2CH_2 , or $-\text{CHCH}_3$.

[0045] B_5 is $-(\text{CH}_2)_n\text{CH}_3$, where $n = 2-3$, $-\text{C}(\text{CH}_3)_3$, cyclohexyl, cyclopentyl, aryl, heteroaryl or arylalkyl.


[0046] The aryl or heteroaryl group can be selected from the list below:







[0047] Each aryl or heteroaryl groups can be mono or multiply substituted with F, I, Cl, Br, CH_3 , CN, OH, OMe, OEt, OCF_3 , CF_3 , or methanesulfonyl.

[0048] Further, in some instances D_1 is a phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl which can be independently mono or multiply substituted with F, Cl, Br, OCF_3 , CF_3 , or CH_3 .


[0049] This disclosure also describes compounds having the structure of Formula V-1, V-2, V-3, VI-1, VI-2, or VI-3:



V-1,




or



VI-3

wherein D_1 is an aryl or heteroaryl; B_5 is an aryl, heteroaryl or carbocycle.

[0050] Each aryl or heteroaryl group is independently selected from the list below:




or

[0051] In some instances each aryl or heteroaryl group is independently mono or multiply substituted. Each aryl or heteroaryl group can be independently mono or multiply substituted with I, F, Cl, Br, CH_3 , CN, OH, OMe, OEt, OCF_3 , CF_3 , or methane sulfonyl. Further the carbocycle is cyclohexyl, cyclohexenyl or cyclopentyl.

[0052] D_1 is an optionally mono or multiply substituted aryl or heteroaryl. B_5 is an optionally mono or multiply substituted


aryl, heteroaryl or carbocycle. In some instances D_1 or B_5 is independently selected from the group consisting of:



and wherein the cabocycle is cyclohexyl, cyclohexenyl or cyclopentyl.

[0053] In some instances D₁ is optionally mono or multiply substituted phenyl, 2-pyridil, 3-pyridyl, or 4-pyridyl. In some instances, D₁ is optionally substituted with one or more of F, Cl, Br, I, OCF₃, CH₃, and CF₃. In some instances, D₁ is not substituted.

[0054] B₅ is optionally mono or multiply substituted





25




35

40



VII-2,


55



VII-3

are also described herein wherein D_1 is an optionally substituted heteroaryl or aryl, B_3 is H or alkyl, B_5 is an optionally substituted aryl or heteroaryl, and R_{26} and R_{27} are each hydrogen or an isotope thereof. In some instances, R_{26} and R_{27} are deuterium. In some instances, R_{26} or R_{27} are independently alkyl. In some instances, B_3 is C_1 - C_5 alkyl.


[0057] In some instances, the compound has a structure of Formula VIII or an enantiomer thereof



VIII,

wherein D_1 is an optionally substituted heteroaryl or aryl, B_3 is H or alkyl, B_5 is an optionally substituted aryl or heteroaryl, and R_{26} and R_{27} are each hydrogen or an isotope thereof. In some instances, R_{26} and R_{27} are deuterium. In some instances, R_{26} or R_{27} are independently alkyl. A_4 is as described herein. B_3 is C_1 - C_5 alkyl. The enantiomer is the R or S enantiomer at the carbon that is connected to D_1 .


[0058] In some embodiments, a compound has the structure of Formula IX or an enantiomer thereof



IX.

[0059] The enantiomer is the R or S enantiomer at the carbon that is connected to D_1 .


[0060] In some embodiments, a compound has the structure of Formula X or an enantiomer thereof




[0061] In some instances, the enantiomer is the R or S enantiomer at the carbon that is connected to D1.


[0062] In some of the structures described herein, D1 is an optionally substituted pyridyl group or phenyl group. In some instances, D1 is an optionally substituted 2-pyridyl, 3-pyridyl, or 4-pyridyl group or phenyl group. In some instances, D1 is optionally substituted with one or more of, H, OH, alkyl alcohol, halo, alkyl, amide, cyano, alkoxy, haloalkyl, or alkylsulfonyl. In some instances, D1 is optionally substituted with one or more of H, OH, Cl, Br, F, I, OMe, CN, CH₃, CF₃.

[0063] In some of the structures described herein, B₅ is an optionally substituted thiophene group. In some instances, B₅ is substituted with an alkoxy group. In some instances, B₅ is substituted with a C₁-C₅ alkoxy group. In some instances, B₅ is substituted with a methoxy group. In some instances, B₅ is




In some instances, B₅ is





wherein R_{23} , R_{24} , and R_{30} are each independently null, H, OH, cycle, aryl, branched or unbranched alkyl alcohol, halo, branched or unbranched alkyl, amide, cyano, alkoxy, haloalkyl, alkylsulfonyl, nitrite, alkylsulfanyl, and R_{25} is H or alkyl. In some instances, R_{23} and R_{24} together form a aryl or cycle that is attached to one or more of the atoms of B_5 . R_{23} , R_{24} , and R_{30} can also be further substituted. In some instances, R_{23} , R_{24} , and R_{30} are each independently H, NH_2 , OH, Cl, Br, F, I, OMe, CN, CH_3 , phenyl, C_3 - C_6 carbocycle, methanesulfonyl, CF_3 ,



wherein R₂₉ is H or an alkyl. In some instances, R₂₉ is a C₁-C₆ alkyl. In some instances, one of R₂₃, R₂₄, and R₃₀ is H. In some instances, at least one of R₂₃, R₂₄, and R₃₀ is H. In some instances, two of R₂₃, R₂₄, and R₃₀ are H.

[0064] The following compounds and others described herein have agonist activity for OR mediated signal transduction:

5 [(4-chlorophenyl)methyl]({2-[4-(4-methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl})amine[(3,4-dimethoxyphenyl)methyl]({2-[2,2-dimethyl-4-phenyloxan-4-yl]ethyl})amine2-[(2-[2-ethyl-2-methyl-4-(4-methylphenyl)oxan-4-yl]ethyl})amino)methyl]phenol[2-(2,2-dimethyl-4-phenyloxan-4-yl)ethyl]({2-fluorophenyl)methyl}amine4-[(2-[4-(2-methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl})amino)methyl]-N,N-dimethylaniline 2-[(2-[2-ethyl-4-(4-fluorophenyl)-2-methyl-oxan-4-yl]ethyl})amino)methyl]phenol[(3-methoxythiophen-2-1)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine.

[0065] Compounds, such as the ones described herein or selected from the compounds described in the Examples can be used in any of the methods described herein, including, but not limited to, treating pain.

[0066] Thus, also described herein are methods of generating agonist activity in OR mediated signal transduction through administration of one or more of the above recited compounds to a subject or subject in need thereof.

[0067] Various atoms in the compositions described herein can be isotopes that occur at lower frequency. Hydrogen can be replaced at any position in the compositions described herein with deuterium. Optionally, hydrogen can also be replaced with tritium. Carbon (¹²C) can be replaced at any position in the compositions described herein with ¹³C or ¹⁴C. Nitrogen (¹⁴N) can be replaced with ¹⁵N. Oxygen (¹⁶O) can be replaced at any position in the compositions described herein with ¹⁷O or ¹⁸O. Sulfur (³²S) can be replaced at any position in the compositions described herein with ³³S, ³⁴S or ³⁶S. Chlorine (³⁵Cl) can be replaced at any position in the compositions described herein with ³⁷Cl. Bromine (⁷⁹Br) can be replaced at any position in the compositions described herein with ⁸¹Br.

[0068] Selected compounds described herein are agonists and antagonists of Opioid Receptors (ORs). The ability of the compounds to stimulate OR mediated signaling may be measured using any assay known in the art to detect OR mediated signaling or OR activity, or the absence of such signaling/activity. "OR activity" refers to the ability of an OR to transduce a signal. Such activity can be measured, e.g., in a heterologous cell, by coupling an OR (or a chimeric OR) to a downstream effector such as adenylate cyclase.

[0069] A "natural ligand-induced activity" as used herein, refers to activation of the OR by a natural ligand of the OR. Activity can be assessed using any number of endpoints to measure OR activity.

[0070] Generally, assays for testing compounds that modulate OR-mediated signal transduction include the determination of any parameter that is indirectly or directly under the influence of a OR, e.g., a functional, physical, or chemical effect.

[0071] Samples or assays comprising ORs that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of inhibition. Control samples (untreated with inhibitors) are assigned a relative OR activity value of 100%. Inhibition of an OR is achieved when the OR activity value relative to the control is about 80%, 50%, or 25%. Activation of an OR is achieved when the OR activity value relative to the control (untreated with activators) is 110%, 150%, 200-500% (i.e., two to five fold higher relative to the control) or, 1000-3000% or higher.

[0072] The effects of the compounds upon the function of an OR can be measured by examining any of the parameters described above. Any suitable physiological change that affects OR activity can be used to assess the influence of a compound on the ORs and natural ligand-mediated OR activity. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as changes in intracellular second messengers such as cAMP.

[0073] Modulators of OR activity are tested using OR polypeptides as described above, either recombinant or naturally occurring. The protein can be isolated, expressed in a cell, expressed in a membrane derived from a cell, expressed in tissue or in an animal. For example, neuronal cells, cells of the immune system, transformed cells, or membranes can be used to test the GPCR polypeptides described above. Modulation is tested using one of the *in vitro* or *in vivo* assays described herein. Signal transduction can also be examined *in vitro* with soluble or solid state reactions, using a chimeric molecule such as an extracellular domain of a receptor covalently linked to a heterologous signal transduction domain, or a heterologous extracellular domain covalently linked to the transmembrane and or cytoplasmic domain of a receptor. Furthermore, ligand-binding domains of the protein of interest can be used *in vitro* in soluble or solid state reactions to assay for ligand binding.

[0074] Ligand binding to an OR, a domain, or chimeric protein can be tested in a number of formats. Binding can be performed in solution, in a bilayer membrane, attached to a solid phase, in a lipid monolayer, or in vesicles. Typically, in an assay described herein, the binding of the natural ligand to its receptor is measured in the presence of a candidate modulator. Alternatively, the binding of the candidate modulator may be measured in the presence of the natural ligand. Often, competitive assays that measure the ability of a compound to compete with binding of the natural ligand to the receptor are used. Binding can be tested by measuring, e.g., changes in spectroscopic characteristics (e.g., fluorescence,

absorbance, refractive index), hydrodynamic (e.g., shape) changes, or changes in chromatographic or solubility properties.

[0075] Modulators may also be identified using assays involving β -arrestin recruitment. β -arrestin serves as a regulatory protein that is distributed throughout the cytoplasm in unactivated cells. Ligand binding to an appropriate OR is associated with redistribution of β -arrestin from the cytoplasm to the cell surface, where it associates with the OR. Thus, receptor activation and the effect of candidate modulators on ligand-induced receptor activation, can be assessed by monitoring β -arrestin recruitment to the cell surface. This is frequently performed by transfecting a labeled β -arrestin fusion protein (e.g., β -arrestin-green fluorescent protein (GFP)) into cells and monitoring its distribution using confocal microscopy (see, e.g., Groarke et al., J. Biol. Chem. 274(33):23263-69 (1999)).

[0076] Another technology that can be used to evaluate OR-protein interactions in living cells involves bioluminescence resonance energy transfer (BRET). A detailed discussion regarding BRET can be found in Kroeger et al., J. Biol. Chem., 276(16):12736-43 (2001).

[0077] Other assays can involve determining the activity of receptors which, when activated by ligand binding, result in a change in the level of intracellular cyclic nucleotides, e.g., cAMP, by activating or inhibiting downstream effectors such as adenylate cyclase. Changes in intracellular cAMP can be measured using immunoassays. The method described in Offermanns & Simon, J. Biol. Chem. 270:15175-15180 (1995) may be used to determine the level of cAMP. Also, the method described in Felley-Bosco et al., Am. J. Resp. Cell and Mol. Biol. 11:159-164 (1994) may be used to determine the level of cGMP. Further, an assay kit for measuring cAMP is described in U.S. Pat. No. 4,115,538.

[0078] Transcription levels can be measured to assess the effects of a test compound on ligand-induced signal transduction. A host cell containing the protein of interest is contacted with a test compound in the presence of the natural ligand for a sufficient time to effect any interactions, and then the level of gene expression is measured. The amount of time to effect such interactions may be empirically determined, such as by running a time course and measuring the level of transcription as a function of time. The amount of transcription may be measured by using any method known to those of skill in the art to be suitable. For example, mRNA expression of the protein of interest may be detected using northern blots or their polypeptide products may be identified using immunoassays. Alternatively, transcription based assays using reporter genes may be used as described in U.S. Pat. No. 5,436,128. The reporter genes can be, e.g., chloramphenicol acetyltransferase, firefly luciferase, bacterial luciferase, β -galactosidase and alkaline phosphatase. Furthermore, the protein of interest can be used as an indirect reporter via attachment to a second reporter such as green fluorescent protein (see, e.g., Mistili & Spector, Nature Biotechnology 15:961-964 (1997)).

[0079] The amount of transcription is then compared to the amount of transcription in either the same cell in the absence of the test compound, or it may be compared with the amount of transcription in a substantially identical cell that lacks the protein of interest. A substantially identical cell may be derived from the same cells from which the recombinant cell was prepared but which had not been modified by introduction of heterologous DNA. Any difference in the amount of transcription indicates that the test compound has in some manner altered the activity of the protein of interest.

Pharmaceutical Compositions/ Formulations

[0080] Pharmaceutical compositions can be formulated by standard techniques using one or more physiologically acceptable carriers or excipients. The formulations may contain a buffer and/or a preservative. The compounds and their physiologically acceptable salts and solvates can be formulated for administration by any suitable route, including via inhalation, topically, nasally, orally, parenterally (e.g., intravenously, intraperitoneally, intravesically or intrathecally) or rectally in a vehicle comprising one or more pharmaceutically acceptable carriers, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard biological practice.

[0081] Pharmaceutical compositions can include effective amounts of one or more compound(s) described herein together with, for example, pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants and/or other carriers. Such compositions may include diluents of various buffer content (e.g., TRIS or other amines, carbonates, phosphates, amino acids, for example, glycineamide hydrochloride (especially in the physiological pH range), N-glycylglycine, sodium or potassium phosphate (dibasic, tribasic), etc. or TRIS-HCl or acetate), pH and ionic strength; additives such as detergents and solubilizing agents (e.g., surfactants such as Pluronics, Tween 20, Tween 80 (Polysorbate 80), Cremophor, polyols such as polyethylene glycol, propylene glycol, etc.), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimersol, benzyl alcohol, parabens, etc.) and bulking substances (e.g., sugars such as sucrose, lactose, mannitol, polymers such as polyvinylpyrrolidones or dextran, etc.); and/or incorporation of the material into particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Hyaluronic acid may also be used. Such compositions can be employed to influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of a compound described herein. See, e.g., Remington's Pharmaceutical Sciences, 18th Ed. (1990, Mack Publishing Co., Easton, Pa. 18042) pages 1435-1712. The compositions can, for example, be prepared in liquid form, or can be in dried powder, such as lyophilized form. Particular methods of

administering such compositions are described *infra*.

[0082] Where a buffer is to be included in the formulations described herein, the buffer can be selected from sodium acetate, sodium carbonate, citrate, glycylglycine, histidine, glycine, lysine, arginine, sodium dihydrogen phosphate, disodium hydrogen phosphate, sodium phosphate, and tris(hydroxymethyl)-aminomethane, or mixtures thereof. The buffer can also be glycylglycine, sodium dihydrogen phosphate, disodium hydrogen phosphate, and sodium phosphate or mixtures thereof.

[0083] Where a pharmaceutically acceptable preservative is to be included in a formulation of one of the compounds described herein, the preservative can be selected from phenol, m-cresol, methyl p-hydroxybenzoate, propyl p-hydroxybenzoate, 2-phenoxyethanol, butyl p-hydroxybenzoate, 2-phenylethanol, benzyl alcohol, chlorobutanol, and thiomerosal, or mixtures thereof. The preservative can also be phenol or m-cresol.

[0084] The preservative is present in a concentration from about 0.1 mg/ml to about 50 mg/ml, in a concentration from about 0.1 mg/ml to about 25 mg/ml, or in a concentration from about 0.1 mg/ml to about 10 mg/ml.

[0085] The use of a preservative in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0086] The formulation may further comprise a chelating agent where the chelating agent may be selected from salts of ethylenediaminetetraacetic acid (EDTA), citric acid, and aspartic acid, and mixtures thereof.

[0087] The chelating agent can be present in a concentration from 0.1 mg/ml to 5 mg/ml, from 0.1 mg/ml to 2 mg/ml or from 2 mg/ml to 5 mg/ml.

[0088] The use of a chelating agent in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0089] The formulation of the compounds described herein may further comprise a stabilizer selected from high molecular weight polymers and low molecular compounds where such stabilizers include, but are not limited to, polyethylene glycol (e.g. PEG 3350), polyvinylalcohol (PVA), polyvinylpyrrolidone, carboxymethylcellulose, different salts (e.g. sodium chloride), L-glycine, L-histidine, imidazole, arginine, lysine, isoleucine, aspartic acid, tryptophan, and threonine or any mixture thereof. The stabilizer can also be L-histidine, imidazole or arginine.

[0090] The high molecular weight polymer can be present in a concentration from 0.1 mg/ml to 50 mg/ml, from 0.1 mg/ml to 5 mg/ml, from 5 mg/ml to 10 mg/ml, from 10 mg/ml to 20 mg/ml, from 20 mg/ml to 30 mg/ml or from 30 mg/ml to 50 mg/ml.

[0091] The low molecular weight compound can be present in a concentration from 0.1 mg/ml to 50 mg/ml, from 0.1 mg/ml to 5 mg/ml, from 5 mg/ml to 10 mg/ml, from 10 mg/ml to 20 mg/ml, from 20 mg/ml to 30 mg/ml or from 30 mg/ml to 50 mg/ml.

[0092] The use of a stabilizer in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0093] The formulation of the compounds described herein may further include a surfactant. The surfactant may be selected from a detergent, ethoxylated castor oil, polyglycolized glycerides, acetylated monoglycerides, sorbitan fatty acid esters, poloxamers, such as 188 and 407, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene derivatives such as alkylated and alkoxyated derivatives (tweens, e.g. Tween-20, or Tween-80), monoglycerides or ethoxylated derivatives thereof, diglycerides or polyoxyethylene derivatives thereof, glycerol, cholic acid or derivatives thereof, lecithins, alcohols and phospholipids, glycerophospholipids (lecithins, cephalins, phosphatidyl serine), glyceroglycolipids (galactopyransoide), sphingophospholipids (sphingomyelin), and sphingoglycolipids (ceramides, gangliosides), DSS (docusate sodium, docusate calcium, docusate potassium, SDS (sodium dodecyl sulfate or sodium lauryl sulfate), dipalmitoyl phosphatidic acid, sodium caprylate, bile acids and salts thereof and glycine or taurine conjugates, ursodeoxycholic acid, sodium cholate, sodium deoxycholate, sodium taurocholate, sodium glycocholate, N-Hexadecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate, anionic (alkyl-aryl-sulphonates) monovalent surfactants, palmitoyl lysophosphatidyl-L-serine, lysophospholipids (e.g. 1-acyl-sn-glycero-3-phosphate esters of ethanolamine, choline, serine or threonine), alkyl, alkoxy (alkyl ester), alkoxy (alkyl ether)-derivatives of lysophosphatidyl and phosphatidylcholines, e.g. lauroyl and myristoyl derivatives of lysophosphatidylcholine, dipalmitoylphosphatidylcholine, and modifications of the polar head group, that is cholines, ethanolamines, phosphatidic acid, serines, threonines, glycerol, inositol, and the positively charged DODAC, DOTMA, DCP, BISHOP, lysophosphatidylserine and lysophosphatidylthreonine, zwitterionic surfactants (e.g. N-alkyl-N,N-dimethylammonio-1-propanesulfonates, 3-cholamido-1-propyldimethylammonio-1-propanesulfonate, dodecylphosphocholine, myristoyl lysophosphatidylcholine, hen egg lysolecithin), cationic surfactants (quarternary ammonium bases) (e.g. cetyl-trimethylammonium bromide, cetylpyridinium chloride), non-ionic surfactants, polyethyleneoxide/polypropyleneoxide block copolymers (Pluronic/Tetronics, Triton X-100, Dodecyl β -D-glucopyranoside) or polymeric surfactants (Tween-40, Tween-80, Brij-35), fusidic acid derivatives—(e.g. sodium tauro-dihydrofusidate etc.), long-chain fatty acids and salts thereof C6-C12 (e.g. oleic acid and caprylic acid), acylcarnitines and derivatives, N_{α} -acylated derivatives of lysine, arginine or histidine, or side-chain acylated derivatives of lysine or arginine, N_{α} -acylated derivatives of dipeptides comprising any combination of lysine, arginine or histidine and a neutral or acidic amino acid, N_{α} -acylated derivative of a tripeptide comprising any combination of a neutral amino acid and two charged amino acids, or the

surfactant may be selected from the group of imidazoline derivatives, or mixtures thereof.

[0094] The use of a surfactant in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0095] Pharmaceutically acceptable sweeteners can be part of the formulation of the compounds described herein. Pharmaceutically acceptable sweeteners include at least one intense sweetener such as saccharin, sodium or calcium saccharin, aspartame, acesulfame potassium, sodium cyclamate, alitame, a dihydrochalcone sweetener, monellin, stevioside or sucralose (4,1',6'-trichloro-4,1',6'-trideoxygalactosucrose), saccharin, sodium or calcium saccharin, and optionally a bulk sweetener such as sorbitol, mannitol, fructose, sucrose, maltose, isomalt, glucose, hydrogenated glucose syrup, xylitol, caramel, and honey.

[0096] Intense sweeteners are conveniently employed in low concentrations. For example, in the case of sodium saccharin, the concentration may range from 0.04% to 0.1% (w/v) based on the total volume of the final formulation, or is about 0.06% in the low-dosage formulations and about 0.08% in the high-dosage ones. The bulk sweetener can effectively be used in larger quantities ranging from about 10% to about 35%, or from about 10% to 15% (w/v).

[0097] The formulations of the compounds described herein may be prepared by conventional techniques, e.g. as described in Remington's Pharmaceutical Sciences, 1985 or in Remington: The Science and Practice of Pharmacy, 19th edition, 1995, where such conventional techniques of the pharmaceutical industry involve dissolving and mixing the ingredients as appropriate to give the desired end product.

[0098] The phrase "pharmaceutically acceptable" or "therapeutically acceptable" refers to molecular entities and compositions that are physiologically tolerable and preferably do not typically produce an allergic or similar untoward reaction, such as gastric upset, dizziness and the like, when administered to a human. As used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a State government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia (e.g., Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985)) for use in animals, and more particularly in humans.

[0099] Administration of the compounds described herein may be carried out using any method known in the art. For example, administration may be transdermal, parenteral, intravenous, intra-arterial, subcutaneous, intramuscular, intracranial, intraorbital, ophthalmic, intraventricular, intracapsular, intraspinal, intracisternal, intraperitoneal, intracerebroventricular, intrathecal, intranasal, aerosol, by suppositories, or oral administration. A pharmaceutical composition of the compounds described herein can be for administration for injection, or for oral, pulmonary, nasal, transdermal, ocular administration.

[0100] For oral administration, the pharmaceutical composition of the compounds described herein can be formulated in unit dosage forms such as capsules or tablets. The tablets or capsules may be prepared by conventional means with pharmaceutically acceptable excipients, including binding agents, for example, pregelatinised maize starch, polyvinylpyrrolidone, or hydroxypropyl methylcellulose; fillers, for example, lactose, microcrystalline cellulose, or calcium hydrogen phosphate; lubricants, for example, magnesium stearate, talc, or silica; disintegrants, for example, potato starch or sodium starch glycolate; or wetting agents, for example, sodium lauryl sulphate. Tablets can be coated by methods well known in the art. Liquid preparations for oral administration can take the form of, for example, solutions, syrups, or suspensions, or they can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives, for example, suspending agents, for example, sorbitol syrup, cellulose derivatives, or hydrogenated edible fats; emulsifying agents, for example, lecithin or acacia; non-aqueous vehicles, for example, almond oil, oily esters, ethyl alcohol, or fractionated vegetable oils; and preservatives, for example, methyl or propyl-p-hydroxybenzoates or sorbic acid. The preparations can also contain buffer salts, flavoring, coloring, and/or sweetening agents as appropriate. If desired, preparations for oral administration can be suitably formulated to give controlled release of the active compound.

[0101] For topical administration, the pharmaceutical composition of the compounds described herein can be formulated in a pharmaceutically acceptable vehicle containing 0.1 to 10 percent, or 0.5 to 5 percent, of the active compound(s). Such formulations can be in the form of a cream, lotion, sublingual tablet, aerosols and/or emulsions and can be included in a transdermal or buccal patch of the matrix or reservoir type as are conventional in the art for this purpose.

[0102] For parenteral administration, the compounds described herein are administered by either intravenous, subcutaneous, or intramuscular injection, in compositions with pharmaceutically acceptable vehicles or carriers. The compounds can be formulated for parenteral administration by injection, for example, by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, for example, in ampoules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain formulatory agents, for example, suspending, stabilizing, and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, for example, sterile pyrogen-free water, before use.

[0103] For administration by injection, the compound(s) can be used in solution in a sterile aqueous vehicle which may also contain other solutes such as buffers or preservatives as well as sufficient quantities of pharmaceutically acceptable salts or of glucose to make the solution isotonic. The pharmaceutical compositions of the compounds de-

scribed herein may be formulated with a pharmaceutically acceptable carrier to provide sterile solutions or suspensions for injectable administration. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspensions in liquid prior to injection or as emulsions. Suitable excipients are, for example, water, saline, dextrose, mannitol, lactose, lecithin, albumin, sodium glutamate, cysteine hydrochloride, or the like. In addition, if desired, the injectable pharmaceutical compositions may contain minor amounts of nontoxic auxiliary substances, such as wetting agents, pH buffering agents, and the like. If desired, absorption enhancing preparations (e.g., liposomes) may be utilized. Suitable pharmaceutical carriers are described in "Remington's pharmaceutical Sciences" by E. W. Martin.

[0104] For administration by inhalation, the compounds may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, for example, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base, for example, lactose or starch. For intranasal administration the compounds described herein may be used, for example, as a liquid spray, as a powder or in the form of drops.

[0105] The compounds can also be formulated in rectal compositions, for example, suppositories or retention enemas, for example, containing conventional suppository bases, for example, cocoa butter or other glycerides.

[0106] Furthermore, the compounds can be formulated as a depot preparation. Such long-acting formulations can be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0107] The compositions can, if desired, be presented in a pack or dispenser device that can contain one or more unit dosage forms containing the active ingredient. The pack can, for example, comprise metal or plastic foil, for example, a blister pack. The pack or dispenser device can be accompanied by instructions for administration.

[0108] The compounds described herein also include derivatives referred to as prodrugs, which can be prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved, either in routine manipulation or in vivo, to the parent compounds. Examples of prodrugs include compounds of the invention as described herein that contain one or more molecular moieties appended to a hydroxyl, amino, sulfhydryl, or carboxyl group of the compound, and that when administered to a patient, cleaves in vivo to form the free hydroxyl, amino, sulfhydryl, or carboxyl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of the invention. Preparation and use of prodrugs is discussed in T. Higuchi et al., "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

Dosages

[0109] The compounds described herein may be administered to a patient at therapeutically effective doses to prevent, treat, or control one or more diseases and disorders mediated, in whole or in part, by an OR-ligand interaction. Pharmaceutical compositions comprising one or more of compounds described herein may be administered to a patient in an amount sufficient to elicit an effective protective or therapeutic response in the patient. An amount adequate to accomplish this is defined as "therapeutically effective dose." The dose will be determined by the efficacy of the particular compound employed and the condition of the subject, as well as the body weight or surface area of the area to be treated. The size of the dose also will be determined by the existence, nature, and extent of any adverse effects that accompany the administration of a particular compound or vector in a particular subject.

[0110] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, for example, by determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and can be expressed as the ratio, LD₅₀/ED₅₀. Compounds that exhibit large therapeutic indices are used. While compounds that exhibit toxic side effects can be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue to minimize potential damage to normal cells and, thereby, reduce side effects.

[0111] The data obtained from cell culture assays and animal studies can be used to formulate a dosage range for use in humans. The dosage of such compounds lies within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration. For any compound described herein, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as

determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography (HPLC). In general, the dose equivalent of a modulator is from about 1 ng/kg to 10 mg/kg for a typical subject.

[0112] The amount and frequency of administration of the compounds described herein and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition. In general it is contemplated that an effective amount would be from 0.001 mg/kg to 10 mg/kg body weight, and in particular from 0.01 mg/kg to 1 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 0.01 to 500 mg, and in particular 0.1 mg to 200 mg of active ingredient per unit dosage form.

[0113] In some instances, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose. The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.01 mg to about 1000 mg, from about 0.01 mg to about 750 mg, from about 0.01 mg to about 500 mg, or from about 0.01 mg to about 250 mg, according to the particular application. The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total dosage may be divided and administered in portions during the day as required.

[0114] In some instances, one or more compounds described herein are administered with another compound. The administration may be sequentially or concurrently. The combination may be in the same dosage form or administered as separate doses. In some instances, the another compound is another analgesic or pain reliever. In some instances, the another compound is a non-opioid analgesic. Examples of useful non-opioid analgesics include, but are not limited to, non-steroidal anti-inflammatory agents, such as aspirin, ibuprofen, diclofenac, naproxen, benoxaprofen, flurbiprofen, fenoprofen, flubufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pramoprofen, muprofen, trioxaprofen, suprofen, aminoprofen, tiaprofenic acid, fluprofen, bucloxic acid, indomethacin, sulindac, tolmetin, zomepirac, tiopinac, zidometacin, acemetacin, fentiazac, clidanac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid, tolfenamic acid, diflusal, flufenisal, piroxicam, sudoxicam, isoxicam, and pharmaceutically acceptable salts thereof, and mixtures thereof. Other suitable non-opioid analgesics include the following, non-limiting, chemical classes of analgesic, antipyretic, nonsteroidal anti-inflammatory drugs: salicylic acid derivatives, including aspirin, sodium salicylate, choline magnesium trisalicylate, salsalate, diflunisal, salicylsalicylic acid, sulfasalazine, and olsalazine; para-aminophenol derivatives including acetaminophen and phenacetin; indole and indene acetic acids, including indomethacin, sulindac, and etodolac; heteroaryl acetic acids, including tolmetin, diclofenac, and ketorolac; anthranilic acids (fenamates), including mefenamic acid and meclofenamic acid; enolic acids, including oxicams (piroxicam, tenoxicam), and pyrazolidinediones (phenylbutazone, oxyphenanthartazone); and alkanones, including nabumetone. For a more detailed description of the NSAIDs, see Paul A. Insel, *Analgesic-Antipyretic and Anti-inflammatory Agents and Drugs Employed in the Treatment of Gout*, in Goodman & Gilman's *The Pharmacological Basis of Therapeutics* 617-57 (Perry B. Molinoff and Raymond W. Ruddon eds., 9.sup.th ed 1996); and Glen R. Hanson, *Analgesic, Antipyretic and Anti-Inflammatory Drugs* in Remington: *The Science and Practice of Pharmacy* Vol II 1196-1221 (A. R. Gennaro ed. 19.sup.th ed. 1995).

[0115] The compounds described herein can also be administered with Cox-II inhibitors. Examples of useful Cox-II inhibitors and 5-lipoxygenase inhibitors, as well as combinations thereof, are described in U.S. Pat. No. 6,136,839. Examples of Cox-II inhibitors include, but are not limited to, rofecoxib and celecoxib.

[0116] The compounds described herein can also be administered with antimigraine agents. Examples of useful antimigraine agents include, but are not limited to, alpropride, bromocriptine, dihydroergotamine, dolasetron, ergocornine, ergocorninine, ergocryptine, ergonovine, ergot, ergotamine, flumetoxone acetate, fonazine, ketanserin, lisuride, lomerizine, methylergonovine, methysergide, metoprolol, naratriptan, oxetorone, pizotyline, propranolol, risperidone, rizatriptan, sumatriptan, timolol, trazodone, zolmitriptan, and mixtures thereof.

[0117] The compounds described herein can also be administered with anti-constipation agents. Examples of anti-constipation agents include, but are not limited to, laxatives or stool softeners. Examples of anti-constipation agents include, but are not limited to, be docusate, poloxamer 188, psyllium, methylcellulose, carboxymethyl cellulose, polycarbophil, bisacodyl, castor oil, magnesium citrate, magnesium hydroxide, magnesium sulfate, dibasic sodium phosphate, monobasic sodium phosphate, sodium biphosphate or any combination thereof.

Medical Use

[0118] The compositions described herein may be useful for treating pain or pain associated disorders. The compositions described herein may be useful for treating immune dysfunction, inflammation, esophageal reflux, neurological

and psychiatric conditions, urological and reproductive conditions, medicaments for drug and alcohol abuse, agents for treating gastritis and diarrhea, cardiovascular agents and agents for the treatment of respiratory diseases and cough.

[0119] Methods of treating pain are described herein. One or more compound described herein are administered to a subject to treat the pain. In some instances, the pain can be post-operative pain. In some instances, the pain is caused by cancer. In some instances, the pain is neuropathic pain. In some instances, the pain is caused by trauma, such as but not limited to, blunt force trauma. In some instances, the pain is caused by inflammation.

[0120] The one or more compounds described herein can be administered by any suitable route, including, but not limited to, via inhalation, topically, nasally, orally, parenterally (e.g., intravenously, intraperitoneally, intravesically or intrathecally) or rectally in a vehicle comprising one or more pharmaceutically acceptable carriers, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard practice.

Definitions

[0121] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the compositions and compounds described herein, suitable methods and materials are described below. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only not intended to be limiting. Other features and advantages of the compositions and compounds described herein will be apparent from the following detailed description and claims.

[0122] The general chemical terms used throughout have their usual meanings. For example, the term alkyl refers to a branched or unbranched saturated hydrocarbon group. The term "*n*-alkyl" refers to an unbranched alkyl group. The term " C_x - C_y alkyl" refers to an alkyl group having from x to y carbon atoms, inclusively, in the branched or unbranched hydrocarbon group. By way of illustration, but without limitation, the term " C_1 - C_4 alkyl" refers to a straight chain or branched hydrocarbon moiety having from 1 to 4 carbon atoms, including methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl, and *tert*-butyl. The term " C_1 - C_4 *n*-alkyl" refers to straight chain hydrocarbon moieties having from 1 to 4 carbon atoms including methyl, ethyl, *n*-propyl, and *n*-butyl. C_x - C_y x can be from 1 to 10 and y is from 2 to 20. The term " C_3 - C_6 cycloalkyl" refers to cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. The term " C_3 - C_7 cycloalkyl" also includes cycloheptyl. Cycloalkylalkyl refers to cycloalkyl moieties linked through an alkyl linker chain, as for example, but without limitation, cyclopropylmethyl, cyclopropylethyl, cyclopropylpropyl, cyclopropylbutyl, cyclobutylmethyl, cyclobutylethyl, cyclobutylpropyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylpropyl, cyclohexylmethyl, cyclohexylethyl, and cyclohexylpropyl. Each alkyl, cycloalkyl, and cycloalkylalkyl group may be optionally substituted, such as, but not limited to, as specified herein. In some instances, the alkyl is a C_1 - C_3 , C_1 - C_4 , C_1 - C_6 , C_4 - C_6 , or C_1 - C_{10} alkyl.

[0123] The terms "alkoxy", "phenyloxy", "benzoyl" and "pyrimidinyl" refer to an alkyl group, phenyl group, benzyl group, or pyrimidinyl group, respectively, that is bonded through an oxygen atom. Each of these groups may be optionally substituted.

[0124] The terms "alkylthio", "phenylthio", and "benzylthio" refer to an alkyl group, phenyl group, or benzyl group, respectively, that is bonded through a sulfur atom. Each of these groups may be optionally substituted.

[0125] The term " C_1 - C_4 acyl" refers to a formyl group or a C_1 - C_3 alkyl group bonded through a carbonyl moiety. The term " C_1 - C_4 alkoxy carbonyl" refers to a C_1 - C_4 alkoxy group bonded through a carbonyl moiety.

[0126] The term "halo" refers to fluoro, chloro, bromo, or iodo. In some instances, the halo groups are fluoro, chloro, and bromo. In some instances, the halo groups are fluoro and chloro.


[0127] As used herein, "carbocycle" or "carbocyclic ring" is intended to mean, unless otherwise specified, any stable 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12-membered monocyclic, bicyclic or tricyclic ring, any of which can be saturated, unsaturated (including partially and fully unsaturated), or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl. As shown above, bridged rings are also included in the definition of carbocycle (e.g., [2.2.2]bicyclooctane). A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. In some instances, the bridges are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring can also be present on the bridge. Fused (e.g., naphthyl and tetrahydronaphthyl) and spiro rings are also included.

[0128] The term "heterocycle" is taken to mean a saturated or unsaturated 5- or 6-membered ring containing from 1 to 3 heteroatoms selected from nitrogen, oxygen and sulfur, said ring optionally being benzofused. Exemplary heterocycles include furanyl, thiophenyl (thienyl), pyrrolyl, pyrrolidinyl, pyridinyl, N-methylpyrrolyl, oxazolyl, isoxazolyl, pyrazolyl, imidazolyl, triazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, thiazolidinyl, N-acetylthiazolidinyl, pyrimidinyl, pyrazinyl, pyridazi-


nyl, and the like. Benzofused heterocyclic rings include isoquinolinyl, benzoxazolyl, benzodioxolyl, benzothiazolyl, quinolinyl, benzofuranyl, benzothiophenyl, indolyl, and the like, all of which may be optionally substituted, which also of course includes optionally substituted on the benzo ring when the heterocycle is benzofused.

[0129] The term "cycle" group is taken to mean a carbocyclic ring, a carbocycle or a heterocarbocycle.

[0130] As used herein, the phrase a "cycle of the formula" refers to a ring that can be formed with the variable referred to. For example, in the structure



wherein A can be a cycle of the formula $C(CH_2)_n$, where $n = 2-5$, it means that A is a carbon and forms a ring with itself with 2-5 CH_2 groups, which could also be represented structurally as



The variable "A" is not limited to carbon and can be another atom, such as, but not limited to, a heteroatom, but the context in which the variable is used will indicate the type of atom "A" could be. This is just a non-limiting example. Additionally, the ring that is formed with "A" can also be substituted. Exemplary substituents are described herein.

[0131] Heterocycles include, but are not limited to, pyridinyl, indolyl, furanyl, benzofuranyl, thiophenyl, benzodioxolyl, and thiazolidinyl, all of which may be optionally substituted.

[0132] As used herein, the term "aromatic heterocycle" or "heteroaryl" is intended to mean a stable 5, 6, 7, 8, 9, 10, 11, or 12-membered monocyclic or bicyclic aromatic ring which consists of carbon atoms and one or more heteroatoms, e.g., 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, independently selected from nitrogen, oxygen, and sulfur. In the case of bicyclic heterocyclic aromatic rings, only one of the two rings needs to be aromatic (e.g., 2,3-dihydroindole), though both can be (e.g., quinoline). The second ring can also be fused or bridged as defined above for heterocycles. The nitrogen atom can be substituted or unsubstituted (i.e., N or NR wherein R is H or another substituent, as defined). The nitrogen and sulfur heteroatoms can optionally be oxidized (i.e., $N \rightarrow O$ and $S(O)_p$, wherein $p = 1$ or 2). In certain compounds, the total number of S and O atoms in the aromatic heterocycle is not more than 1.

[0133] Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indolizyl, indolyl, 3H-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, methylenedioxyphenyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxindolyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranal, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinolizyl, quinoxalyl, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, tetrazolyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

[0134] Substituted alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, or alkylthio, means an alkyl, cycloalkyl, cycloalkylalkyl, alkoxy, or alkylthio group, respectively, substituted one or more times independently with a substituent selected from the group consisting of halo, hydroxy and C_1-C_3 alkoxy. By way of illustration, but without limitation, examples include trifluoromethyl, pentafluoroethyl, 5-fluoro-2-bromopentyl, 3-hydroxypropyloxy, 4-hydroxycyclohexyloxy, 2-bromoethylthio, 3-ethoxypropyloxy, 3-ethoxy-4-chlorocyclohexyl, and the like. Substitutions include substitution 1-5 times with halo, each independently selected, or substituted 1-3 times with halo and 1-2 times independently with a group selected from hydroxy and C_1-C_3 alkoxy, or substituted 1-3 times independently with a group selected from hydroxy and C_1-C_3 alkoxy, provided that no more than one hydroxy and/or alkoxy substituent may be attached through the same carbon.

[0135] The terms "substituted phenyl" and "substituted heterocycle" are taken to mean that the cyclic moiety in either case is substituted. They can be substituted independently with one or more substituents. They can be substituted independently with 1, 2, 3, 4, 5, 1-3, 1-4, or 1-5 substituents. The substitution can be, independently, halo, alkyl, such as, but not limited to, C₁-C₄ alkyl, alkoxy, such as but not limited to, C₁-C₄ alkoxy, and alkylthio, such as but not limited to, C₁-C₄ alkylthio, wherein each alkyl, alkoxy and alkylthio substituent can be further substituted independently with C₁-C₂ alkoxy or with one to five halo groups; or substituted with one substituent selected from the group consisting of phenyloxy, benzyloxy, phenylthio, benzylthio, and pyrimidinyl, wherein the phenyloxy, benzyloxy, phenylthio, benzylthio, and pyrimidinyl moiety can be further substituted with one to two substituents selected from the group consisting of halo, C₁-C₂ alkyl, and C₁-C₂ alkoxy; or substituted with one substituent selected from the group consisting of C₁-C₄ acyl and C₁-C₄ alkoxycarbonyl, and further substituted with zero to one substituent selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, and C₁-C₄ alkylthio. When a substituent is halo, in some instances, the halo groups are fluoro, chloro, and bromo. The halo can also be iodo.

[0136] DMF means N,N-dimethylformamide.

[0137] As used herein, the phrase "pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0138] By "pharmaceutical formulation" it is further meant that the carrier, solvent, excipients and salt must be compatible with the active ingredient of the formulation (e.g. a compound described herein). It is understood by those of ordinary skill in this art that the terms "pharmaceutical formulation" and "pharmaceutical composition" are generally interchangeable, and they are so used for the purposes of this disclosure.

[0139] As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids selected from 2-acetoxybenzoic, 2-hydroxyethane sulfonic, acetic, ascorbic, benzene sulfonic, benzoic, bicarbonic, carbonic, citric, edetic, ethane disulfonic, ethane sulfonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, glycolylarsanilic, hexylresorcinic, hydrabamic, hydrobromic, hydrochloric, hydroiodide, hydroxymaleic, hydroxynaphthoic, isethionic, lactic, lactobionic, lauryl sulfonic, maleic, malic, mandelic, methane sulfonic, napsylic, nitric, oxalic, pamoic, pantothenic, phenylacetic, phosphoric, polygalacturonic, propionic, salicylic, stearic, subacetic, succinic, sulfamic, sulfanilic, sulfuric, tannic, tartaric, and toluene sulfonic. The present disclosure includes pharmaceutically acceptable salts of any compound(s) described herein.

[0140] Pharmaceutically acceptable salts can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile, and the like. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing Company, Easton, PA, USA, p. 1445 (1990).

[0141] Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility, bioavailability, manufacturing, etc.) the compounds described herein can be delivered in prodrug form and can be administered in this form for the treatment of disease. "Prodrugs" are intended to include any covalently bonded carriers that release an active parent drug of described herein *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds described herein wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug is administered to a mammalian subject, it cleaves to form a free hydroxyl, free amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate, and benzoate derivatives of alcohol and amine functional groups in the compounds described herein.

[0142] "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

[0143] As used herein, "treating" or "treatment" includes any effect e.g., lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder, etc. "Treating" or "treatment" of a disease state means the treatment of a disease-state in a mammal, particularly in a human, and include: (a) inhibiting an existing disease-state, *i.e.*, arresting its development or its clinical symptoms; and/or (c) relieving the disease-state, *i.e.*, causing regression of the disease state.

[0144] As used herein, "preventing" means causing the clinical symptoms of the disease state not to develop *i.e.*,

inhibiting the onset of disease, in a subject that may be exposed to or predisposed to the disease state, but does not yet experience or display symptoms of the disease state.

[0145] As used herein, "mammal" refers to human and non-human patients.

[0146] As used herein, the term "therapeutically effective amount" refers to a compound, or a combination of compounds, described herein present in or on a recipient in an amount sufficient to elicit biological activity, e.g. pain relief. In some instances, the combination of compounds is a synergistic combination. Synergy, as described, for example, by Chou and Talalay, Adv. Enzyme Regul. vol. 22, pp. 27-55 (1984), occurs when the effect of the compounds when administered in combination is greater than the additive effect of the compounds when administered alone as a single agent. In general, a synergistic effect is most clearly demonstrated at sub-optimal concentrations of the compounds. Synergy can be in terms of lower cytotoxicity, increased decrease in pain, or some other beneficial effect of the combination compared with the individual components.

[0147] All percentages and ratios used herein, unless otherwise indicated, are by weight.

[0148] Throughout the description, where compositions are described as having, including, or comprising specific components, or where processes are described as having, including, or comprising specific process steps, it is contemplated that compositions described herein also consist essentially of, or consist of, the recited components, and that the processes described herein also consist essentially of, or consist of, the recited processing steps. Further, it should be understood that the order of steps or order for performing certain actions are immaterial so long as the process remains operable. Moreover, two or more steps or actions can be conducted simultaneously.

[0149] All enantiomers, diastereomers, and mixtures thereof, are included within the scope of compounds described herein. In some instances a composition comprising the *R* enantiomer is free or substantially free of the *S* enantiomer. In some instances, a composition comprising the *S* enantiomer is free or substantially free of the *R* enantiomer. In some instances, a composition comprises an enantiomeric excess of at least, or about, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% of either the *R* or the *S* enantiomer.


[0150] As used throughout this disclosure, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a composition" includes a plurality of such compositions, as well as a single composition, and a reference to "a therapeutic agent" is a reference to one or more therapeutic and/or pharmaceutical agents and equivalents thereof known to those skilled in the art, and so forth. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

[0151] The claimed compounds in this invention can be prepared from the procedures described in the schemes below.


Schemes

[0152] The following representative schemes illustrate how compounds described herein can be prepared. The specific solvents and reaction conditions referred to are also illustrative and are not intended to be limited. Compounds not described are either commercially available or are readily prepared by one skilled in the art using available starting materials.


Scheme 1: Synthesis of Spirocyclic Nitrile



Scheme 2: Converting the nitrile to the opioid receptor ligand (Approach 1)




[0153] The same scheme is applied to 1-7 and 1-8A.

Scheme 3: Converting the nitrile to the opioid receptor ligand (Approach 2)

n=1-2


R and R₁ are independent
 R and R₁ = phenyl, substituted phenyl, aryl,
 substituted aryl, pyridyl, substituted pyridyl,
 heteroaryl, substituted heteroaryl,
 carbocycle, heterocycle and etc.

[0154] The same scheme is applied to 1-7 and 1-8A.

Scheme 4: Synthesis of Non-Spirocyclic Nitrile


R = phenyl, substituted phenyl, aryl,
 substituted aryl, pyridyl, substituted pyridyl,
 heteroaryl, substituted heteroaryl,
 carbocycle, heterocycle and etc.

4-1 is selected from the group consisting of



[0155] Following a sequence outlined in Scheme 2 or 3, intermediate 4-4 can be converted to the opioid receptor ligands.


Scheme 5: Synthesis of Other Spirocyclic Derived Opioid Ligands



$n=1-2$

R and R₁ are independent
 R and R₁ = phenyl, substituted phenyl, aryl,
 substituted aryl, pyridyl, substituted pyridyl,
 heteroaryl, substituted heteroaryl,
 carbocycle, heterocycle and etc.

Scheme 5: Synthesis of Other Spirocyclic Derived Opioid Ligands




$n=1-2$


R and R₁ are independent
 R and R₁ = phenyl, substituted phenyl, aryl,
 substituted aryl, pyridyl, substituted pyridyl,
 heteroaryl, substituted heteroaryl,
 carbocycle, heterocycle and etc.

[0156] Other schemes can also be used. For example, the following schemes can be used alone or in combination with other schemes to prepare the compounds described herein.


Scheme 6: Allyltrimethylsilane Approach to Access the Quaternary Carbon Center




Scheme 7: N-linked Pyrrole Opioid Receptor Ligand




Scheme 8




Scheme 9




Scheme 10



[0157] In one embodiment, a process for preparing a compound having the structure of IV-




or a pharmaceutically acceptable salt thereof is provided, the process comprising contacting



with



under suitable conditions to form a compound having the structure of



IV-1.

In some instances, the process is performed at room temperature. In some instances, the process is performed in the presence of a borohydrate salt. In some instances, the process is performed in the presence of sodium borohydrate. Solvents can also be used to facilitate the preparation. The process can be modified to yield different alkyl groups, such as, but not limited to, the scheme shown in Scheme 10.

Examples

[0158] The following examples are illustrative, but not limiting, of the methods and compositions described herein.

Example 1:

Intermediate 1: methyl 2-cyano-2-(oxan-4-ylidene)acetate

[0159] A 50 ml round-bottom flask equipped with a Dean-Stark distillation setup and condenser was charged with tetrahydro-4H-pyran-4-one (4.61 ml, 50 mmol), methyl cyanoacetate (5.3 ml, 60 mmol), ammonium acetate (1 g, 13 mmol), acetic acid (0.57 ml, 10 mmol) and benzene (30 ml). The mixture was refluxed until no more water collected in the Dean-Stark (2 hours), cooled, benzene (30 ml) added and the organic layer washed with water (50 ml). The aqueous layer was extracted with CH_2Cl_2 (3x50 ml). The combined organic phase was washed with sat. NaHCO_3 (100 ml), brine (100 ml) dried (MgSO_4), filtered and concentrated. Purified by normal phase SiO_2 chromatography (10 to 60% EtOAc/hexanes) to afford methyl 2-cyano-2-(oxan-4-ylidene)acetate as a colorless oil (6.30g, 70%, m/z : 181.1 $[\text{M} + \text{H}]^+$ observed).

Intermediate 2: methyl 2-cyano-2-[4-(4-fluorophenyl)oxan-4-yl]acetate

[0160] A round bottom flask was equipped with a condenser, addition funnel and rubber septum with nitrogen inlet was charged with a solution of *p*-fluorophenylmagnesium bromide (2.0 M in diethyl ether, 1.99 ml, 3.97mmol) and CuI (63 mg, 0.331 mmol) in 10 ml dry diethyl ether (10 ml). Methyl 2-cyano-2-(oxan-4-ylidene)acetate (600 mg, 3.31 mmol) in diethyl ether (10 ml) was added drop-wise over 30 min while cooling the reaction flask in an ice bath. The mixture

was then stirred for 3h. The reaction mixture was poured into a 50 g ice/1 N HCl (25 ml) mixture. The product was extracted with Et₂O (3x50 ml), washed with brine (50 ml), dried (Na₂SO₄) and concentrated. Purified by normal phase SiO₂ chromatography (7% to 60% EtOAc/hexanes) to give methyl 2-cyano-2-[4-(4-fluorophenyl)oxan-4-yl]acetate as a white solid (730 mg, 80%, m/z: 277.1 [M + Na]⁺ observed).

Intermediate 3: 2-[4-(4-fluorophenyl)oxan-4-yl]acetonitrile

[0161] To a pre-dissolved solution of KOH (441 mg, 7.87 mmol) in ethylene glycol (20 ml) was added methyl 2-cyano-2-[4-(4-fluorophenyl)oxan-4-yl]acetate (1.09 g, 3.93 mmol). The mixture was heated to 120 °C for 3 h, and then cooled. H₂O was added (50 ml), the product extracted with Et₂O (3x50 ml), washed with H₂O (50 ml), dried over Na₂SO₄, filtered and concentrated. Purified by normal phase SiO₂ chromatography (5 to 40% EtOAc/hexanes) to give 2-[4-(4-fluorophenyl)oxan-4-yl]acetonitrile as a colorless oil (450 mg, 78%, m/z: 219.1 [M + H]⁺ observed).

Intermediate 4: 2-[4-(4-fluorophenyl)oxan-4-yl]ethan-1-amine

[0162] To a solution of 2-[4-(4-fluorophenyl)oxan-4-yl]acetonitrile (450 mg, 2.05 mmol) in anhydrous ether (15 ml) at 0 °C was added dropwise LAH (1.0 M in Et₂O, 4.1 ml, 4.11 mmol). After 2 h the reaction was quenched with 1 ml H₂O, 0.1 ml 15% NaOH and then 1 ml H₂O. The reaction mixture was extracted with Et₂O (3x20 ml), dried over Na₂SO₄ and concentrated to give 2-[4-(4-fluorophenyl)oxan-4-yl]ethan-1-amine as a yellow oil, which used without further purification (450 mg, 94%, m/z: 223.1 [M + H]⁺ observed).

Examples 2: benzyl({2-[4-(4-fluorophenyl)oxan-4-yl]ethyl})amine (Compound 8)

[0163] To a solution of 2-[4-(4-fluorophenyl)oxan-4-yl]ethan-1-amine (250 mg, 1.12 mmol) in anhydrous CH₂Cl₂ (5 ml) and Na₂SO₄ (159 mg, 1.12 mmol) at rt was added benzaldehyde (0.17 ml, 1.68 mmol). The reaction was stirred overnight. The reaction mixture was filtered and concentrated. The residue was dissolved in 5 ml MeOH at 0 °C and NaBH₄ added in one portion (51 mg, 1.34 mmol). The reaction was stirred at 0 °C for 1 h. The solution was then quenched with H₂O (10 ml), extracted with CH₂Cl₂ (3x20 ml), washed with brine (10 ml) and dried over Na₂SO₄. Purified by normal phase SiO₂ chromatography (0 to 10% MeOH/CH₂Cl₂) to give benzyl({2-[4-(4-fluorophenyl)oxan-4-yl]ethyl})amine as a colorless oil (200 mg, 60%, m/z: 314.2 [M + H]⁺ observed).

Intermediate 5: 2,2-dimethyl-4-(4-methylphenyl)oxan-4-ol

[0164] *n*-BuLi (26.3 ml, 1.6 M in hexane, 42 mmol) was added dropwise to a solution of 4-bromo-toluene (7.70 g, 45 mmol) in THF (100 ml) at -78 °C under N₂. The resulting mixture was stirred at -78 °C for 30 min and a solution of tetrahydro-2,2-dimethyl-4H-pyran-4-one (3.84 g, 30 mmol) in THF (20 ml) was added. The resulting mixture was stirred at -78 °C for another 20 min and quenched by adding MeOH (10 ml). The reaction was concentrated under vacuum and the resulting residue was diluted with EtOAc (500 ml) and washed with sat. NH₄Cl (250 ml), brine (250 ml), dried and concentrated to give 2,2-dimethyl-4-(4-methylphenyl)oxan-4-ol as a white solid, which was used without further purification (5.41 g, 82%).

[0165] ¹H NMR (400 MHz, CDCl₃) δ 7.36 - 7.26 (m, 2H), 7.11 (d, *J* = 8.0, 2H), 4.10 (td, *J* = 12.0, 2.2, 1H), 3.71 (ddd, *J* = 11.8, 5.0, 2.1, 1H), 2.28 (s, 3H), 2.11 (ddd, *J* = 13.7, 12.2, 5.0, 1H), 1.72 (dt, *J* = 14.2, 8.3, 2H), 1.58 (dq, *J* = 13.8, 2.2, 1H), 1.44 (s, 3H), 1.38 (s, 1H), 1.14 (s, 3H).

Intermediate 6: 2,2-dimethyl-4-(4-methylphenyl)-4-(prop-2-en-1-yl)oxane

[0166] Allyltrimethylsilane (4.34 ml, 27.2 mmol) was added to a solution of 2,2-dimethyl-4-(4-methylphenyl)oxan-4-ol (3.0 g, 13.6 mmol) in dry CH₂Cl₂ (100 ml) at 0 °C, followed by BF₃·OEt₂ (3.42 ml, 27.2 mmol). The resulting mix was stirred at 0 °C for 1h. The reaction was quenched with H₂O (10 ml) and diluted with CH₂Cl₂ (10 ml), and washed with sat. NaHCO₃ (20 ml), brine (20 ml), dried and concentrated. Purified by normal phase SiO₂ chromatography (5 to 40% EtOAc/hexanes) to give 2,2-dimethyl-4-(4-methylphenyl)-4-(prop-2-en-1-yl)oxane as a colorless oil, which was used crude (2.49 g, 75%).

Intermediate 7: 2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]acetaldehyde

[0167] O₃ gas was passed through a solution of 2,2-dimethyl-4-(4-methylphenyl)-4-(prop-2-en-1-yl)oxane (1.21 g, 5 mmol) in CH₂Cl₂ (50 ml) at -78 °C until the solution turned light blue (about 5 min). After additional 5 minutes, the reaction mix was purged with oxygen gas for 15 min before adding triphenylphosphine (2.62 g, 10 mmol). The reaction was

stirred at rt for 4h and concentrated. Purified by normal phase SiO₂ chromatography (10% to 60% EtOAc/hexanes) to give 2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]acetaldehyde as a colorless oil (641 mg, 52%).

[0168] ¹H NMR (400 MHz, CDCl₃) δ 9.42-9.27 (m, 1H), 7.26 (dd, *J* = 9.9, 8.0, 2H), 7.20 (t, *J* = 8.7, 2H), 3.94 - 3.75 (m, 2H), 2.69 (dd, *J* = 14.6, 2.5, 1H), 2.51 - 2.38 (m, 2H), 2.35 (s, 3H), 2.26 (dd, *J* = 13.9, 2.3, 1H), 1.84 (ddd, *J* = 14.3, 11.0, 4.6, 1H), 1.76 (d, *J* = 13.9, 1H), 1.23 (s, 3H), 0.73 (s, 3H).

Example 3: {2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(3-methylphenyl)methyl]amine (Compound 32)

[0169] A mixture of 2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]acetaldehyde (61.6 mg, 0.25 mmol), 3-methylbenzylamine (63 μl, 0.5 mmol) and acetic acid (50 μl, 8.6 mmol) in CH₂Cl₂ (3 ml) was stirred at rt for 1h before it adding sodium triacetoxyborohydride (106 mg, 0.50 mmol). The resulting mixture was stirred at rt for 18 h. The mix was concentrated and dissolved in MeOH and purified by HPLC to give {2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(3-methylphenyl)methyl]amine as a white solid (35 mg, 40%, *m/z*: 352.3 [M + H]⁺ observed).

Intermediate 8: methyl 2-cyano-2-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]acetate

[0170] A 100 ml round-bottom flask equipped with a Dean-Stark distillation setup and condenser was charged with 6-oxaspiro[4.5]decan-9-one (6 g, 39 mmol, which was prepared according to Hanschke, E. Chem. Ber. 1955, 88, 1053), methyl cyanoacetate (4.1 ml, 46.7 mmol), ammonium acetate (780 mg, 10.1 mmol), acetic acid (0.44 ml, 7.8 mmol) and benzene (40 ml). The mixture was refluxed until no more water collected in the Dean-Stark (2 hours), cooled, benzene (30 ml) added and the organic washed with water (50 ml). The aqueous layer was extracted with CH₂Cl₂ (3x50 ml). The combined organic phase was washed with sat. NaHCO₃ (100 ml), brine (100 ml) dried (MgSO₄), filtered and concentrated. Purified by normal phase SiO₂ chromatography (7% to 60% EtOAc/hexanes) to give methyl 2-cyano-2-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]acetate as a colorless oil (8.93g, 97.5%, *m/z* 235.1 [M + H]⁺ observed).

[0171] By the procedure for the preparation of intermediate 8 substituting 2,2-diethyloxan-4-one for 6-oxaspiro[4.5]decan-9-one, methyl 2-cyano-2-[(4Z)-2,2-diethyloxan-4-ylidene]acetate was prepared (*m/z* 237.1 [M + H]⁺ observed).

[0172] By the procedure for the preparation of intermediate 8 substituting 1-oxaspiro[5.5]undecan-4-one for 6-oxaspiro[4.5]decan-9-one, methyl 2-cyano-2-[(4Z)-1-oxaspiro[5.5]undecan-4-ylidene]acetate was prepared (*m/z* 249.1 [M + H]⁺ observed).

Intermediate 9: methyl 2-cyano-2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetate

[0173] A round bottom flask was equipped with a condenser, addition funnel and rubber septum with nitrogen inlet was charged with a solution of 4-fluoromagnesium bromide (2.0 M in diethyl ether, 7.5 ml, 12.5 mmol) and CuI (200 mg, 1.0 mmol) in 35 ml dry diethyl ether. Methyl 2-cyano-2-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]acetate (2.5g, 10.5 mmol) in diethyl ether (35 ml) was added drop-wise over 30 min while cooling the reaction flask in an ice bath. The mixture was then stirred at room temperature for 1h. The reaction mixture was poured into a 25 g ice/1 N HCl (20 ml) mixture. The product was extracted with Et₂O (3x50 ml), washed with brine (50 ml), dried (Na₂SO₄) and concentrated. Purified by normal phase SiO₂ chromatography (8% to 60% EtOAc/hexanes) to give methyl 2-cyano-2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetate as a colorless oil (3.24 g, 93%, *m/z* 331.2 [M + H]⁺ observed).

[0174] By the procedure described in the preparation of intermediate 9 substituting methyl 2-cyano-2-[(4Z)-2,2-diethyloxan-4-ylidene]acetate for methyl 2-cyano-2-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]acetate, methyl 2-cyano-2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]acetate was prepared (*m/z* 333.2 [M + H]⁺ observed).

[0175] By the procedure described in the preparation of intermediate 9 substituting methyl 2-cyano-2-[(4Z)-1-oxaspiro[5.5]undecan-4-ylidene]acetate for methyl 2-cyano-2-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]acetate, methyl 2-cyano-2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]acetate was prepared (*m/z* 345.2 [M + H]⁺ observed).

Intermediate 10: 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile

[0176] To a pre-dissolved solution of KOH (1.1g, 19.5 mmol) in ethylene glycol (50 ml) was added methyl 2-cyano-2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetate (3.24 g, 9.8 mmol). The mixture was heated to 120 °C for 3 h, then cooled. H₂O was added (50 ml), the product extracted with Et₂O (3x50 ml), washed with H₂O (50 ml), dried over Na₂SO₄, filtered and concentrated. (7% to 60% EtOAc/hexanes) to give methyl 2-cyano-2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetate (1.96 g, 73%, *m/z* 273.2 [M + H]⁺ observed).

[0177] 1.96 g of the enantiomers were separated by SFC on an AD-3 column using 15% MeOH (0.05% DEA) as a modifier to give 2-[(9S)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile as a colorless oil (faster eluting enantiomer, 635 mg, 24%, *m/z* 274.2 [M + H]⁺ observed) and 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile as a colorless oil (slower eluting enantiomer, 703 mg, 26%, *m/z* 273.2 [M + H]⁺ observed).

[0178] By the procedure described in the preparation of intermediate 10 substituting methyl 2-cyano-2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]acetate for methyl 2-cyano-2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetate, 2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]acetonitrile was prepared (m/z 275.2 $[M + H]^+$ observed).

[0179] By the procedure described in the preparation of intermediate 10 substituting methyl 2-cyano-2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]acetate for methyl 2-cyano-2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetate, 2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]acetonitrile was prepared (m/z 287.2 $[M + H]^+$ observed).

Intermediate 11: 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine

[0180] To a solution of 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (500 mg, 1.8 mmol) in anhydrous ether (30 ml) at 0 °C was added dropwise LAH (1.0 M in Et₂O, 3.7 ml, 3.7 mmol). The reaction was then warmed up to room temperature. After 2h the reaction was quenched with 1 ml H₂O, 0.2 ml 15% NaOH and then 1 ml H₂O. The reaction mixture was extracted with Et₂O (3x30 ml), dried over Na₂SO₄ and concentrated to give 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine as a yellow oil, which used without further purification (500 mg, 100%, m/z 277.2 $[M + H]^+$ observed).

[0181] By the procedure described in the preparation of intermediate 11 substituting 2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]acetonitrile for 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile, 2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethan-1-amine was prepared (m/z 279.2 $[M + H]^+$ observed).

[0182] By the procedure described in the preparation of intermediate 11 substituting 2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]acetonitrile for 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile, 2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethan-1-amine was prepared (m/z 291.2 $[M + H]^+$ observed).

Example 4: benzyl({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine (Compound 81)

[0183] To a solution of amine 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine (100 mg, 0.361 mmol) in anhydrous CH₂Cl₂ (6 ml) and Na₂SO₄ (256 mg, 1.80 mmol) at rt was added benzaldehyde (0.055 ml; 0.541 mmol). The reaction was stirred overnight. The reaction mixture was filtered and concentrated. The residue was dissolved in 6 ml MeOH at 0 °C and NaBH₄ added in one portion (16 mg, 0.433 mmol). The reaction was stirred at 0 °C for 1 h. The solution was then quenched with H₂O (20 ml), extracted with CH₂Cl₂ (3x30 ml), washed with brine (10 ml) and dried over Na₂SO₄. The mixture was purified by HPLC to give benzyl({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine as a white solid (121 mg, 92%, m/z 368.3 $[M + H]^+$ observed).

Intermediate 12: 2,2-diethyloxan-4-ol.

[0184] To a mixture of 3-butene-1-ol (19.8 ml; 233mmol) and 3-pentenone (12.3 ml; 116 mmol) was added 75% sulfuric acid (19.8; 334 mmol; prepared by diluting 79 ml of conc. sulfuric acid to 100 ml with distilled water) drop-wise at 0 °C. The reaction was allowed to warm to room temperature and stirred overnight. Water (70 ml) was added to the mixture then neutralized with NaOH (pellets) to pH 8 and extracted with diethyl ether (3x150 ml). The ether extract was washed with an aqueous sodium bisulfite solution (40 ml), dried over K₂CO₃ and the ether evaporated *in vacuo*. The residue was distilled under reduced pressure to give 2,2-diethyloxan-4-ol (4.89 g, 27%, B. Pt. 65-70 °C at 1mm Hg).

[0185] ¹H NMR (400 MHz, CDCl₃) δ 4.04 - 3.86 (m, 1H), 3.84 - 3.66 (m, 1H), 3.65 - 3.38 (m, 1H), 2.06 - 1.95 (m, 1H), 1.92 - 1.76 (m, 2H), 1.78 - 1.63 (m, 1H), 1.63 - 1.50 (m, 1H), 1.51 - 1.31 (m, 3H), 1.28 - 1.10 (m, 1H), 0.92 - 0.68 (m, 6H).

Intermediate 13: 2,2-diethyloxan-4-one

[0186] To a solution of crude 2,2-diethyloxan-4-ol (500mg, 3.2 mmol) in CH₂Cl₂ (10 ml) were added NMO (750 mg, 6.41 mmol) and 4A molecular sieves(2g). The solution was stirred for 30 mins and then TPAP (34 mg, 0.096 mmol) was added in one portion. The reaction was allowed to stir for 10 h. After checking the TLC, the alcohol was gone. It was filtered through a short pad of SiO₂. The filtrate was concentrated and purified by normal phase SiO₂ chromatography (0% to 50% EtOAc/hexanes) to give 2,2-diethyloxan-4-one (365 mg, 73%).

[0187] ¹H NMR (400 MHz, CDCl₃) δ 3.75 - 3.66 (m, 2H), 3.44 - 3.29 (m, 2H), 2.51 - 2.31 (m, 4H), 1.25-1.4 (m, 4H), 0.75 (m, 6H).

Intermediate 14: 2-(bromomagnesio)pyridine

[0188] Into a flask was placed isopropylmagnesium chloride 2.0M in THF (6 mL, 12 mmole), 2-bromopyridine (1.2 mL, 12 mmol) in anhydrous Et₂O (4 ml) added dropwise. The reaction mixture was stirred at rt. for 3h. The resulting mixture was used as is as 1M Grignard solution.

Example 5: dibenzyl({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine (Compound 225)

[0189] To a solution of 2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (30 mg, 0.13 mmol) in anhydrous CH₂Cl₂ (3 ml) and Na₂SO₄ (92.3 mg, 0.65 mmol) at rt was added 2.3 eq benzaldehyde (0.032 ml, 0.32 mmol); The reaction was stirred overnight. NaBH(OAc)₃ (6.6 mg, 0.31 mmol) added in one portion. The solution was then quenched with H₂O (10 ml), extracted with CH₂Cl₂ (3x20 ml), washed with brine (10 ml) and dried over Na₂SO₄. The solvent was evaporated *in vacuo* and the residue was purified by HPLC to obtain dibenzyl({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine (37.4 mg, 50%, m/z 458.3 [M + H]⁺ observed).

Example 6: {2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(3-methylphenyl)methyl]amine (Compound 122)

[0190] Following an analogous procedure described for Compound 81, Compound 122 was obtained from the corresponding intermediate after a chiral HPLC separation (The slower moving fraction on AD-3 column. The absolute configuration of Ex. 122 was determined by an X-ray crystallography).

Example 7: {2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[2-(pyridin-3-yl)ethyl]amine (Compound 75)

[0191] 1.0 M DIBAL solution in toluene (3.0 ml, 3 mmol) was added drop-wise to a solution of 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (350 mg, 1.4 mmol) in 7 mL toluene at -78 °C. The resulting mixture was stirred at -78 °C until completion (1.5 h). The reaction was then quenched with 5 eq of MeOH (0.28 mL) and 0.1 mL water, stir while warming, 175 mg Na₂SO₄ added, stir at room temp. 2h to give 310 mg (80%) of 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetaldehyde. LCMS m/z 250.6 (M + 1) observed.

[0192] To a solution of 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetaldehyde (50 mg, 0.19 mmole), 5 mL DCM and Na₂SO₄ (134 mg, 0.95 mmole) was added 2-(pyridin-3-yl)ethan-1-amine (31 mg, 0.25 mmole) and the reaction was stirred overnight. NaBH₄ (9.5 mg, 0.25 mmole) added, stir 10 minutes, 2 drops MeOH added, stir 1h, quenched with water, organics separated off and evaporated. The residue was passed through a Gilson reverse phase HPLC to give {2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[2-(pyridin-3-yl)ethyl]amine, 65.3 mg (71%). LCMS m/z 367.1 (M + 1) observed.

Example 8: 2-[(9R)-9-(2-{4H,5H,6H-thieno[2,3-c]pyrrol-5-yl}ethyl)-6-oxaspiro[4.5]decan-9-yl]pyridine (Compound 82)

[0193] To a stirred solution of 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine (0.030 g, 0.115 mmol; prepared by following a sequence described for Compound 81 in dried ACN (5.8 mL) was added 2,3-bis(bromomethyl)thiophene (31.1 mg, 0.115 mmol) followed by addition of K₂CO₃ (79.62 mg, 0.576 mmol). After 30 min, LCMS showed that the reaction was done and the major peak had the corresponding mass to the desired product. It was then subjected to HPLC purification. HPLC purification method: Luna acid medium column, 10-50% acetonitrile in H₂O over 15 min, followed by flashing with 100% acetonitrile, 0.1% TFA modifier was employed. The fractions containing the desired product were pooled, basified with 2N NaOH and extracted with DCM (3x20 mL). The combined organics were concentrated and purified with flash chromatography (10 g silica gel column, eluted by 0-10% MeOH in DCM, based upon TLC measurement: DCM/MeOH (10/1) R_f = 0.60) to afford 5 mg of 2-[(9R)-9-(2-{4H,5H,6H-thieno[2,3-c]pyrrol-5-yl}ethyl)-6-oxaspiro[4.5]decan-9-yl]pyridine as a colorless oil in 12% yield. LCMS m/z 369 (M + 1) observed.

Example 9: {2-[9-(1H-pyrazol-1-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine (Compound 26)

[0194] An oven-dried flask equipped with a Dean-Stork apparatus and condenser was cooled to rt under a stream of N₂ and was charged with 6-oxaspiro[4.5]decan-9-one (0.50 g, 3.24 mmol), (tert-butoxy)carbohydrazide (0.42 g, 3.24 mmol) and hexane (10 mL). The resulting solution was heated to reflux overnight.

[0195] It was cooled to rt and the solid collected by vacuum filtration. The solid was washed with hexane and air-dried to give (tert-butoxy)-N'-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]carbohydrazide (0.84 g, 96% yield). LCMS m/z 213 (M + 1-t-butyl) observed.

[0196] An oven-dried flask was charged with (tert-butoxy)-N'-[(9Z)-6-oxaspiro[4.5]decan-9-ylidene]carbohydrazide (0.42 g, 1.56 mmol) and THF. The solution was cooled to 0 °C and allylmagnesiumchloride (2.0 M, 1.60 mL) was added dropwise. The reaction was stirred at 0 °C for 1h and the warmed to rt overnight. LC-MS indicated the reaction didn't go to completion. Another 2 equivalent of allylmagnesiumchloride was added at rt. The solution was stirred for 1h before it was quenched with MeOH. The solution was diluted with DCM (60 mL) and H₂O (20 mL). A lot of precipitates were formed and the solid was filtered through a pad of celite. The organic was then separated and the aqueous layer was

extracted with 10 mL of EtOAc. The combined organic layers were concentrated and the residue was purified on 25 g Snap column (0-20% tOAc in Hex, 12 CV) to give (tert-butoxy)-N'-[9-(prop-2-en-1-yl)-6-oxaspiro[4.5]decan-9-yl]carbohydrazide (0.33 g, 68% yield). LCMS m/z 333 (M + Na) observed.

[0197] A solution of (tert-butoxy)-N'-[9-(prop-2-en-1-yl)-6-oxaspiro[4.5]decan-9-yl]carbohydrazide (0.33 g, 1.06 mmol) in 4 mL of EtOAc was added 4M HCl in dioxane at rt. The solution was stirred at rt until reaction completion, monitored by LC-MS (30 h). The solvent was then removed to give [9-(prop-2-en-1-yl)-6-oxaspiro[4.5]decan-9-yl]hydrazine (250 mg). LCMS m/z 211.1 (M + 1) observed.


[0198] A solution of [9-(prop-2-en-1-yl)-6-oxaspiro[4.5]decan-9-yl]hydrazine (250 mg, 1.0 mmol) in 4 mL of i-PrOH were added Et3N and 3-dimethylaminoacrolein. The solution was refluxed for 3h and then at 50 °C for 2d. The solvent removed and the residue was purified on 25 g Biotage snap column, eluted with 0-18% EtOAc in Hex (12CV) to give 1-[9-(prop-2-en-1-yl)-6-oxaspiro[4.5]decan-9-yl]-1H-pyrazole (80 mg, 31% yield). LCMS m/z 247.1 (M + 1) observed.

[0199] To a solution of 1-[9-(prop-2-en-1-yl)-6-oxaspiro[4.5]decan-9-yl]-1H-pyrazole (80 mg, 0.32 mmol) in DCM (5 mL) at -78 °C was bubbled with O₃ until the solution turned blue. The resulting solution was bubbled with N₂ for 5 min. To it was added PPh3 (168 mg, 0.64 mmol). And the solution was stirred for 4h at rt. After removal of the solvent, the residue was purified by flash column chromatography to give 2-[9-(1H-pyrazol-1-yl)-6-oxaspiro[4.5]decan-9-yl]acetaldehyde (15 mg, 23 % yield). LCMS m/z 249 (M + 1) observed.

[0200] To a mixture of 2-[9-(1H-pyrazol-1-yl)-6-oxaspiro[4.5]decan-9-yl]acetaldehyde (15 mg, 0.06 mmol) and thiophen-2-ylmethanamine (19 µL, 0.18 mmol) was stirred at rt for 1h before NaBH(OAc)₃ (25.4 mg, .12 mmol) was added. The solution stirred overnight. After removal of solvent, the residue was purified by HPLC to provide {2-[9-(1H-pyrazol-1-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine (17 mg, 61% yield) as a TFA salt. LCMS m/z 346 (M + 1) observed.

Example 10: Basic Procedure for making compounds of the formula:


[0201]



[0202] Following Scheme 8 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine, which can be prepared by following a sequence as described for Compound 81 (Compound 4) and a sequence similar to for Intermediate 11 reacts with an appropriately substituted heteroaromatic aldehyde or appropriately substituted aromatic aldehyde (1 equivalent) in the presence of an organic solvent (i.e. DCM, MeOH, EtOH) to form a corresponding imine, which is reduced by an appropriate reducing agent the compound. The (R)_n and the R_m refers to the optional substituents. Additionally, the phenyl groups can be replaced with other cycles or aryl groups as described herein.

Example 11: Basic procedures for making compounds of the formula:

[0203]



[0204] Following Scheme 9, 9-1, which can be prepared by following a sequence described for Compound 81 (Compound 4) and a sequence similar to for Intermediate 11, reacts with an appropriately substituted heteroaromatic aldehyde or appropriately substituted aromatic aldehyde (1 equivalent) in the presence of an organic solvent (i.e. DCM, MeOH, EtOH and etc) to form a corresponding imine, which is reduced by an appropriate reducing agent (i.e. NaBH₄) to give

the compound. The $(R)_n$ and the R_m refers to the optional substituents. Additionally, the phenyl groups can be replaced with other cycles or aryl groups as described herein.

Example 12: Opioid Receptor Ligands

[0205] The opioid receptor ligands and compounds listed in the following tables can be or were prepared according to the procedures described above from appropriate starting materials and appropriate reagents. Compounds that have been made lists NMR data and prophetic examples do not list NMR. data.

Table 1: Compounds with chemical name and characterization data			
Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
1	2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine	261.1	δ 8.58 (ddd, J = 4.8, 1.9, 0.9, 1H), 7.63 (m, 1H), 7.30 (m, 1H), 7.12 (ddd, J = 7.4, 4.8, 1.0, 1H), 3.76 (m, 2H), 2.55 (td, J = 11.6, 5.1, 1H), 2.46 (ddd, J = 13.7, 5.1, 2.7, 1H), 2.37 (dd, J = 13.7, 2.1, 1H), 2.14 (td, J = 11.6, 5.0, 1H), 1.92 (m, 2H), 1.70 (m, 4H), 1.46 (m, 4H), 1.13 (m, 1H), 0.71 (dt, J = 13.4, 8.8, 1H).
2	2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine	261.2	δ 8.58 (ddd, J = 4.8, 1.7, 0.7, 1H), 7.64 (td, J = 7.8, 1.9, 1H), 7.28 (m, 1H), 7.12 (ddd, J = 7.4, 4.8, 0.9, 1H), 3.76 (m, 2H), 2.55 (m, 1H), 2.46 (ddd, J = 13.7, 5.1, 2.7, 1H), 2.37 (m, 1H), 2.14 (m, 1H), 1.91 (m, 2H), 1.71 (m, 4H), 1.47 (m, 4H), 1.13 (m, 1H), 0.71 (m, 1H).
3	2-[9-(2-aminoethyl)-6-oxaspiro[4.5]decan-9-yl]pyridin-4-ol	277.1	δ 7.60 (d, J = 6.8, 1H), 7.60 (d, J = 6.8, 1H), 7.21 (s, 2H), 6.79 (d, J = 332.9, 3H), 6.54 (m, 5H), 6.37 (s, 1H), 6.29 (d, J = 6.5, 1H), 5.97 (m, 6H), 4.84 (d, J = 169.9, 3H), 3.69 (dt, J = 23.7, 11.7, 3H), 3.69 (dt, J = 23.7, 11.7, 3H), 3.40 (s, 2H), 3.40 (s, 2H), 2.64 (s, 1H), 2.64 (s, 1H), 2.32 (d, J = 12.0, 1H), 2.26 (dd, J = 46.7, 13.0, 2H), 2.20 (d, J = 13.9, 1H), 2.09 (d, J = 13.8, 1H), 2.09 (d, J = 13.8, 1H), 1.84 (t, J = 15.9, 2H), 1.60 (m, 15H), 1.55 (m, 11H), 0.89 (m, 2H), 0.89 (m, 1H).
4	6-[9-(2-aminoethyl)-6-oxaspiro[4.5]decan-9-yl]pyridin-3-ol	277.1	δ 8.05 (m, 1H), 7.01 (d, J = 8.6, 1H), 6.87 (dd, J = 8.6, 2.9, 1H), 5.37 (s, 2H), 3.66 (dd, J = 13.7, 7.2, 2H), 2.60 (ddd, J = 12.3, 10.1, 5.6, 1H), 2.22 (m, 3H), 1.88 (tt, J = 10.0, 7.9, 1H), 1.77 (d, J = 13.6, 1H), 1.58 (m, 4H), 1.37 (m, 5H), 1.08 (dd, J = 15.4, 4.9, 1H), 0.63 (dt, J = 13.7, 8.9, 1H).
5	6-[9-(2-aminoethyl)-6-oxaspiro[4.5]decan-9-yl]pyridin-2-ol	277.1	δ 7.38 (dd, J = 9.0, 7.1, 1H), 6.40 (d, J = 9.0, 1H), 6.09 (d, J = 7.1, 1H), 5.28 (s, 1H), 3.73 (s, 2H), 2.69 (m, 1H), 2.37 (m, 2H), 2.13 (m, 2H), 1.66 (m, 12H), 0.97 (dt, J = 12.4, 7.6, 1H).
6	2-[(9R)-9-(2-aminoethyl)-6-oxaspiro[4.5]decan-9-yl]-1-oxidopyridin-1-ium	277.1	δ 8.23 (m, 1H), 7.31 (dd, J = 5.8, 2.0, 2H), 7.23 (m, 1H), 4.03 (s, 2H), 3.81 (s, 2H), 3.27 (d, J = 13.9, 1H), 2.95 (td, J = 12.8, 5.1, 1H), 2.65 (td, J = 11.8, 5.1, 1H), 2.25 (s, 2H), 1.65 (ddd, J = 38.7, 17.6, 11.7, 9H), 1.24 (s, 1H), 0.84 (dt, J = 13.1, 8.8, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
7	benzyl({2-[1-(4-fluorophenyl)cyclohexyl]ethyl})amine	312.2	δ 9.72 (t, J = 1.5, 1H), 7.91 (m, 2H), 7.23 (m, 3H), 7.06 (m, 2H), 6.92 (m, 2H), 2.91 (t, J = 6.9, 2H), 2.44 (m, 4H), 2.09 (dd, J = 13.2, 5.3, 2H), 1.68 (m, 4H), 1.46 (m, 1H), 1.33 (dd, J = 15.4, 6.7, 4H).
8	benzyl({2-[4-(4-fluorophenyl)oxan-4-yl]ethyl})amine	314.2	δ 7.38 - 7.16 (m, 7H), 7.09 - 6.99 (m, 2H), 3.79 (ddd, J = 11.5, 5.7, 3.6, 2H), 3.64 (s, 2H), 3.56 (ddd, J = 11.6, 8.8, 2.8, 2H), 2.39 - 2.29 (m, 2H), 2.24 - 2.01 (m, 4H), 1.86 (ddd, J = 13.8, 7.9, 3.5, 2H).
9	[(2-methylphenyl)methyl]({2-[4-(4-methylphenyl)oxan-4-yl]ethyl})amine	324.2	δ 7.16 (m, 8H), 3.79 (ddd, J = 11.5, 5.2, 3.8, 2H), 3.58 (m, 4H), 2.41 (m, 2H), 2.36 (s, 3H), 2.27 (s, 3H), 2.16 (m, 2H), 1.86 (ddd, J = 12.3, 8.6, 4.6, 4H), 1.58 (s, 1H)
10	N-{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}aniline	324.3	δ 7.25 (dt, J = 5.9, 2.9, 3H), 7.11 (m, 2H), 6.98 (dd, J = 25.0, 8.2, 4H), 3.65 (dd, J = 8.9, 6.7, 2H), 2.95 (d, J = 4.6, 1H), 2.50 (d, J = 4.7, 1H), 2.23 (s, 3H), 2.10 (d, J = 13.9, 1H), 1.89 (m, 3H), 1.43 (m, 2H), 1.21 (m, 1H), 1.05 (s, 3H), 0.53 (s, 3H).
11	2-[(2-[4-(4-methylphenyl)oxan-4-yl]ethyl]amino)methyl] phenol	326.2	δ 7.15 (m, 5H), 6.88 (dd, J = 7.4, 1.3, 1H), 6.79 (dd, J = 8.1, 1.0, 1H), 6.73 (td, J = 7.4, 1.1, 1H), 3.76 (m, 4H), 3.54 (ddd, J = 11.7, 9.4, 2.5, 2H), 2.37 (m, 5H), 2.13 (m, 2H), 1.82 (m, 4H)
12	2-[(2-[4-(4-fluorophenyl)oxan-4-yl]ethyl]amino)methyl] phenol	330.2	δ 8.26 (s, 2H), 7.07 (m, 3H), 6.95 (dd, J = 14.3, 5.8, 2H), 6.86 (d, J = 6.3, 1H), 6.78 (d, J = 8.1, 1H), 6.71 (t, J = 7.4, 1H), 3.96 (d, J = 7.0, 2H), 3.80 (s, 2H), 3.64 (dt, J = 8.8, 3.9, 2H), 3.41 (t, J = 9.3, 2H), 2.45 (s, 2H), 1.95 (d, J = 14.7, 2H), 1.85 (m, 2H), 1.67 (m, 2H).
13	benzyl({2-[3-(pyridin-2-yl)-1-oxaspiro[4.4]nonan-3-yl]ethyl})amine	337.1	δ 9.76 (s, 2H), 8.59 (d, J = 4.7, 1H), 8.11 (t, J = 7.8, 1H), 7.69 (d, J = 8.1, 1H), 7.63 - 7.52 (m, 1H), 7.35 (s, 5H), 4.13 (d, J = 9.7, 1H), 4.03 (s, 2H), 3.91 (d, J = 9.7, 1H), 2.92 (d, J = 26.5, 2H), 2.53 (ddd, J = 14.6, 9.5, 5.4, 1H), 2.38 (dd, J = 19.0, 8.5, 2H), 2.28 (d, J = 13.6, 1H), 1.99 - 1.81 (m, 1H), 1.84 - 1.52 (m, 6H), 1.51 - 1.35 (m, 1H).
14	benzyl({2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl})amine	338.3	δ 8.54 (d, J = 226.0, 2H), 7.22 (q, J = 6.7, 3H), 7.03 (dd, J = 19.0, 8.3, 6H), 6.23 (d, J = 186.3, 2H), 3.69 (m, 4H), 2.66 (s, 1H), 2.25 (s, 4H), 2.10 (dd, J = 22.7, 13.3, 2H), 1.83 (m, 1H), 1.64 (m, 1H), 1.49 (m, 2H), 1.11 (s, 3H), 0.57 (s, 3H).
15	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}(pyridin-2-ylmethyl)amine	339.3	δ 8.48 (dd, J = 5.2, 0.9, 1H), 8.26 (s, 1H), 7.98 (dd, J = 7.8, 1.6, 1H), 7.59 (d, J = 7.9, 1H), 7.54 (m, 1H), 7.07 (s, 4H), 4.17 (q, J = 13.9, 2H), 3.73 (m, 2H), 2.88 (d, J = 4.8, 1H), 2.42 (d, J = 4.8, 1H), 2.21 (m, 4H), 2.10 (dd, J = 13.9, 2.1, 1H), 2.00 (d, J = 4.6, 1H), 1.78 (d, J = 4.6, 1H), 1.58 (m, 2H), 1.12 (s, 3H), 0.59 (s, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
16	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}(pyridin-3-ylmethyl)amine	339.3	δ 8.92 (s, 1H), 8.52 (s, 1H), 8.25 (d, J = 8.0, 1H), 7.67 (m, 1H), 7.08 (m, 4H), 5.92 (s, 4H), 4.09 (s, 2H), 3.71 (m, 2H), 2.85 (dd, J = 12.0, 7.9, 1H), 2.34 (m, 1H), 2.23 (m, 4H), 2.10 (d, J = 13.9, 1H), 1.94 (m, 1H), 1.74 (dd, J = 12.5, 4.3, 1H), 1.55 (m, 2H), 1.10 (s, 3H), 0.57 (s, 3H).
17	[(2-methoxyphenyl)methyl]({2-[4-(4-methylphenyl)oxan-4-yl]ethyl})amine	340.2	δ 7.21 (m, 1H), 7.13 (s, 4H), 7.07 (dd, J = 7.4, 1.7, 1H), 6.84 (ddd, J = 12.1, 9.3, 4.6, 2H), 3.78 (m, 5H), 3.63 (s, 2H), 3.54 (ddd, J = 11.6, 9.1, 2.7, 2H), 2.38 (d, J = 1.3, 1H), 2.32 (m, 5H), 2.10 (m, 2H), 1.84 (m, 4H)
18	(furan-3-ylmethyl)({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	341.1	δ 8.72 (d, J = 4.6, 1H), 8.23 (t, J = 7.3, 1H), 7.84 - 7.57 (m, 2H), 7.46 (s, 1H), 7.38 (t, J = 1.6, 1H), 7.28 (s, 1H), 3.89 (s, 2H), 3.82 (dt, J = 12.4, 4.2, 1H), 3.72 (dd, J = 16.1, 6.2, 1H), 2.96 (d, J = 4.4, 1H), 2.40 (ddd, J = 36.0, 24.7, 12.8, 4H), 2.20 (dd, J = 12.7, 4.8, 1H), 2.01 (d, J = 14.2, 1H), 1.95 - 1.77 (m, 2H), 1.69 (dd, J = 9.6, 4.4, 1H), 1.63 - 1.39 (m, 4H), 1.21 - 1.08 (m, 1H), 0.91-0.60 (m, 1H).
19	(1H-imidazol-2-ylmethyl)({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	341.1	δ 8.70 (d, J = 5.1, 1H), 8.40 (t, J = 7.9, 1H), 7.92 (d, J = 8.2, 1H), 7.87 - 7.74 (m, 1H), 7.31 (d, J = 18.0, 2H), 4.66 (d, J = 14.3, 1H), 4.49 (d, J = 14.3, 1H), 4.02 - 3.81 (m, 1H), 3.74 (d, J = 9.7, 1H), 3.10 (d, J = 4.9, 1H), 2.84 - 2.48 (m, 2H), 2.37 (t, J = 12.7, 3H), 2.17 -
			2.00 (m, 1H), 2.00 - 1.82 (m, 2H), 1.70 (s, 1H), 1.65 - 1.41 (m, 4H), 1.21 (s, 1H), 0.82 (d, J = 13.1, 1H).
20	(1,3-oxazol-4-ylmethyl)({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	342.1	δ 8.77 (dd, J = 5.5, 1.4, 1H), 8.26 (td, J = 8.0, 1.7, 1H), 7.90 (s, 1H), 7.82 (s, 1H), 7.79 - 7.60 (m, 2H), 4.08 (s, 2H), 3.86 (d, J = 12.9, 1H), 3.81 - 3.66 (m, 1H), 3.13 (d, J = 5.6, 1H), 2.76 - 2.60 (m, 1H), 2.48 (s, 1H), 2.42 - 2.25 (m, 3H), 2.16 - 2.01 (m, 1H), 1.89 (dd, J = 9.6, 4.0, 2H), 1.79 - 1.63 (m, 1H), 1.63 - 1.35 (m, 4H), 1.19 (s, 1H), 0.80 (d, J = 13.2, 1H).
21	{2-[3-(pyridin-2-yl)-1-oxaspiro[4.4]nonan-3-yl]ethyl}(thiophen-2-ylmethyl)amine	343	δ 10.01 (s, 3H), 8.62 (d, J = 4.5, 1H), 8.11 (td, J = 8.0, 1.4, 1H), 7.70 (d, J = 8.1, 1H), 7.63 - 7.46 (m, 1H), 7.33 (dd, J = 5.1, 1.0, 1H), 7.16 (d, J = 2.8, 1H), 7.00 (dd, J = 5.1, 3.6, 1H), 4.29 (s, 2H), 4.14 (d, J = 9.7, 1H), 3.92 (d, J = 9.7, 1H), 2.97 (qd, J = 18.1, 12.2, 2H), 2.53 (ddd, J = 14.4, 9.0, 5.7, 1H), 2.45 - 2.21 (m, 3H), 2.00 - 1.82 (m, 1H), 1.67 (tt, J = 22.6, 8.0, 6H), 1.44 (dd, J = 14.3, 10.0, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
22	{2-[3-(pyridin-2-yl)-1-oxaspiro[4.4]nonan-3-yl]ethyl}(thiophen-3-ylmethyl)amine	343	δ 11.15 (s, 2H), 9.70 (s, 2H), 8.64 (d, J = 4.4, 1H), 8.17 (td, J = 8.0, 1.5, 1H), 7.74 (d, J = 8.1, 1H), 7.63 (dd, J = 6.7, 5.7, 1H), 7.40 (dd, J = 2.8, 1.1, 1H), 7.33 (dd, J = 5.0, 3.0, 1H), 7.10 (dd, J = 5.0, 1.2, 1H), 4.23 - 4.07 (m, 3H), 3.94 (d, J = 9.8, 1H), 2.90 (d, J = 33.7, 2H), 2.67 - 2.50 (m, 1H), 2.50 - 2.24 (m, 3H), 1.91 (dd, J = 13.7, 4.9, 1H), 1.83 - 1.52 (m, 6H), 1.43 (td, J = 7.7, 3.9, 1H).
23	(cyclopentylmethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	343.3	δ 8.77 (d, J = 4.6, 2H), 8.26 (t, J = 7.6, 1H), 7.89 - 7.60 (m, 2H), 3.85 (dd, J = 8.5, 4.2, 1H), 3.73 (t, J = 10.1, 1H), 3.00 (s, 1H), 2.81 (s, 2H), 2.42 (dt, J = 23.0, 9.5, 4H), 2.25 (t, J = 10.8, 1H), 2.19 - 1.98 (m, 2H), 1.98 - 1.33 (m, 13H), 1.16 (s, 3H), 0.76 (dt, J = 13.1, 8.9, 1H).
24	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}(thiophen-2-ylmethyl)amine	344.2	δ 8.87 (d, J = 194.4, 2H), 3.91 (s, 3H), 3.69 (m, 2H), 2.66 (d, J = 7.9, 1H), 2.24 (m, 4H), 2.10 (ddd, J = 30.6, 14.0, 2.1, 2H), 1.84 (td, J = 12.5, 4.9, 1H), 1.65 (m, 1H), 1.49 (m, 2H), 1.11 (d, J = 6.1, 3H), 0.57 (s, 3H).
25	(2-[4-(4-fluorophenyl)oxan-4-yl]ethyl)[(2-methoxyphenyl)methyl]amine	344.2	δ 7.20 (ddd, J = 7.6, 4.8, 2.0, 3H), 7.03 (m, 3H), 6.84 (ddd, J = 11.7, 9.1, 4.5, 2H), 3.77 (m, 5H), 3.61 (s, 2H), 3.54 (ddd, J = 11.6, 8.8, 2.8, 2H), 2.27 (m, 2H), 2.08 (m, 2H), 1.84 (ddd, J = 10.5, 8.4, 3.0, 4H), 1.58 (s, 1H)
26	{2-[9-(1H-pyrazol-1-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	346	δ 9.87 (s, 1H), 9.00 (d, J = 145.4, 2H), 7.46 (dd, J = 12.7, 2.1, 2H), 7.28 (dd, J = 5.1, 1.1, 1H), 7.01 (d, J = 0.8, 1H), 6.93 (dd, J = 5.1, 3.5, 1H), 6.34 - 6.24 (m, 1H), 5.22 (s, 1H), 4.10 (q, J = 14.2, 2H), 3.68 (d, J = 2.7, 2H), 2.94 (s, 1H), 2.50 (s, 1H), 2.31 (s, 2H), 2.24 - 2.08 (m, 1H), 1.99 (dt, J = 14.7, 7.3, 1H), 1.93 -
			1.76 (m, 2H), 1.75 - 1.63 (m, 1H), 1.57 (ddd, J = 23.2, 14.0, 8.1, 1H), 1.51 (s, 4H), 1.17 - 1.04 (m, 1H), 0.69 (dt, J = 13.3, 8.7, 1H).
27	benzyl{2-[(9S)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	351.1	δ 8.67 (d, J = 4.7, 1H), 8.17 (t, J = 7.7, 1H), 7.64 (m, 2H), 7.35 (m, 5H), 6.51 (s, 4H), 4.72 (s, 1H), 3.94 (s, 2H), 3.75 (m, 2H), 2.95 (s, 1H), 2.49 (s, 1H), 2.33 (m, 3H), 2.19 (m, 1H), 1.98 (d, J = 14.1, 1H), 1.81 (dt, J = 13.4, 7.5, 2H), 1.68 (m, 1H), 1.49 (ddd, J = 20.8, 14.7, 7.2, 4H), 1.15 (s, 1H), 0.75 (m, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
28	benzyl({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	351.1	δ 8.61 (s, 1H), 8.18 (t, J = 7.7, 1H), 7.65 (m, 2H), 7.26 (m, 5H), 6.90 (d, J = 26.0, 4H), 3.88 (s, 2H), 3.72 (d, J = 12.7, 1H), 3.60 (t, J = 10.0, 1H), 2.90 (s, 1H), 2.39 (d, J = 34.6, 2H), 2.20 (t, J = 13.3, 3H), 1.92 (d, J = 14.8, 2H), 1.75 (m, 2H), 1.59 (d, J = 4.9, 1H), 1.41 (m, 4H), 1.08 (s, 1H), 0.68 (dt, J = 13.2, 9.0, 1H).
29	benzyl({2-[3-(pyridin-2-yl)-1-oxaspiro[4.5]decan-3-yl]ethyl})amine	351.1	δ 9.67 (s, 2H), 8.61 (s, 1H), 8.19 (t, J = 7.5, 1H), 7.80 (d, J = 8.1, 1H), 7.64 (s, 1H), 7.36 (s, 5H), 4.22 (d, J = 10.0, 1H), 4.05 (s, 2H), 3.98 (d, J = 10.0, 1H), 3.00 (s, 1H), 2.84 (s, 1H), 2.64 (s, 1H), 2.39 (d, J = 8.7, 1H), 2.18 (d, J = 13.6, 1H), 2.09 (d, J = 13.6, 1H), 1.75 - 1.52 (m, 4H), 1.33 (dd, J = 28.9, 16.2, 7H).
30	benzyl({2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	351.2	δ 8.49 (s, 1H), 8.03 (s, 1H), 7.53 (d, J = 8.0, 2H), 7.18 (m, 5H), 3.82 (s, 2H), 3.63 (s, 1H), 3.53 (dd, J = 23.8, 13.7, 1H), 2.84 (s, 1H), 2.38 (s, 1H), 2.27 (d, J = 7.4, 1H), 2.13 (d, J = 14.1, 3H), 1.84 (d, J = 14.2, 1H), 1.67 (m, 2H), 1.52 (d, J = 5.0, 1H), 1.32 (m, 4H), 1.01 (s, 1H), 0.61 (dt, J = 13.0, 8.9, 1H).
31	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(2-methylphenyl)methyl] amine	352.2	δ 7.09 (d, J = 8.3, 2H), 7.02 (ddd, J = 8.1, 6.1, 3.3, 6H), 3.69 (m, 2H), 3.47 (s, 2H), 2.41 (td, J = 10.8, 5.4, 1H), 2.25 (m, 4H), 2.11 (m, 5H), 1.75 (ddd, J = 13.2, 10.4, 5.2, 1H), 1.56 (m, 4H), 1.11 (s, 3H), 0.59 (s, 3H).
32	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(3-methylphenyl)methyl] amine	352.3	δ 9.13 (s, 1H), 8.69 (s, 1H), 7.04 (m, 6H), 6.86 (m, 2H), 3.65 (m, 6H), 2.59 (s, 1H), 2.12 (m, 9H), 1.83 (td, J = 12.4, 4.5, 1H), 1.64 (m, 1H), 1.48 (m, 2H), 1.10 (s, 3H), 0.57 (s, 3H).
33	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(4-methylphenyl)methyl] amine	352.3	δ 8.68 (d, J = 205.9, 2H), 7.02 (dd, J = 16.8, 9.0, 6H), 6.93 (d, J = 8.1, 2H), 3.67 (dd, J = 6.6, 2.7, 2H), 3.57 (s, 2H), 3.44 (s, 3H), 2.61 (s, 1H), 2.25 (d, J = 11.2, 3H), 2.17 (s, 3H), 2.08 (dd, J = 20.3, 14.0, 2H), 1.84 (m, 1H), 1.67 (d, J = 7.6, 1H), 1.48 (m, 2H), 1.09 (s, 3H), 0.56 (s, 3H).
34	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(1R)-1-phenylethyl]amine	352.3	δ 9.07 (dd, J = 228.2, 166.6, 2H), 7.24 (ddd, J = 9.3, 6.4, 3.4, 3H), 7.15 (m, 2H), 6.94 (m, 4H); 3.91 (s, 1H), 3.61 (dd, J = 7.0, 4.0, 2H), 2.42 (d, J = 33.9, 1H), 2.21 (d, J = 11.7, 6H), 2.00 (m, 2H), 1.82 (m, 1H), 1.62 (dd,
			J = 8.6, 4.1, 1H), 1.42 (m, 5H), 1.05 (s, 3H), 0.53 (d, J = 3.4, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
35	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(1S)-1-phenylethyl]amine	352.3	δ 8.92 (dd, J = 238.8, 174.0, 2H), 7.24 (m, 3H), 7.14 (td, J = 7.5, 2.2, 2H), 6.95 (m, 4H), 3.89 (d, J = 19.3, 1H), 3.62 (m, 2H), 2.96 (s, 2H), 2.42 (m, 1H), 2.21 (d, J = 11.6, 3H), 2.00 (m, 3H), 1.82 (m, 1H), 1.63 (m, 1H), 1.40 (m, 5H), 1.06 (s, 3H), 0.53 (d, J = 3.6, 3H).
36	benzyl({2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl})methylamine	352.3	δ 11.09 (s, 2H), 7.39 (m, 3H), 7.23 (m, 1H), 7.15 (m, 5H), 4.19 (dd, J = 25.7, 12.6, 1H), 3.91 (dd, J = 17.4, 8.4, 1H), 3.78 (m, 2H), 2.91 (d, J = 127.4, 1H), 2.56 (dd, J = 17.7, 7.2, 3H), 2.37 (d, J = 4.8, 3H), 2.24 (ddd, J = 22.0, 12.2, 2.2, 3H), 2.05 (m, 1H), 1.88 (td, J = 12.5, 4.7, 1H), 1.64 (m, 2H), 1.21 (s, 3H), 382.30.67 (d, J = 1.2, 3H).
37	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}(2-phenylethyl)amine	352.3	δ 9.06 (d, J = 128.6, 2H), 7.17 (m, 3H), 7.02 (m, 6H), 3.68 (dd, J = 11.8, 10.1, 2H), 2.77 (dt, J = 36.5, 30.4, 7H), 2.19 (m, 5H), 1.99 (m, 1H), 1.89 (td, J = 12.5, 4.6, 1H), 1.69 (m, 1H), 1.49 (m, 2H), 1.00 (s, 3H), 0.52 (s, 3H).
38	(pyrazin-2-ylmethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	353.1	δ 8.79 (dd, J = 5.6, 1.4, 1H), 8.68 - 8.54 (m, 2H), 8.51 (dd, J = 2.3, 1.6, 1H), 8.32 (td, J = 8.0, 1.6, 1H), 7.93 - 7.66 (m, 3H), 4.30 (s, 2H), 3.85 (dt, J = 12.3, 4.2, 1H), 3.72 (t, J = 9.9, 1H), 3.19 (td, J = 11.7, 5.2, 1H), 2.72 (td, J = 11.8, 4.0, 1H), 2.62 - 2.45 (m, 1H), 2.45 - 2.27 (m, 3H), 2.10 (d, J = 14.2, 1H), 2.00 - 1.79 (m, 2H), 1.69 (dt, J = 9.9, 6.6, 1H), 1.63 - 1.41 (m, 4H), 1.19 (dd, J = 12.6, 6.5, 1H), 0.78 (dt, J = 13.1, 8.9, 1H).
39	benzyl({2-[2,2-diethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl})amine	353.3	δ 8.71 (dd, J = 5.5, 1.4, 1H), 8.21 (td, J = 8.0, 1.7, 1H), 7.67 (m, 2H), 7.33 (m, 5H), 3.95 (s, 2H), 3.79 (m, 1H), 3.67 (d, J = 10.8, 1H), 3.01 (d, J = 5.2, 1H), 2.40 (m, 4H), 2.10 (s, 1H), 1.73 (t, J = 16.5, 2H), 1.55 (dd, J = 14.1, 7.5, 1H), 1.39 (dd, J = 14.1, 7.4, 1H), 0.81 (m, 5H), 0.56 (t, J = 7.3, 3H).
40	benzyl({2-[2,2,6,6-tetramethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl})amine	353.3	δ 8.74 - 8.62 (m, 1H), 8.24 (td, J = 8.1, 1.5, 1H), 7.87 (d, J = 8.2, 1H), 7.76-7.65 (m, 1H), 7.47 - 7.18 (m, 7H), 3.96 (s, 2H), 2.75 (s, 2H), 2.50 (d, J = 14.7, 2H), 2.43 - 2.28 (m, 2H), 1.89 (d, J = 14.8, 2H), 1.30 (s, 6H), 0.97 (s, 6H).
41	4-[(2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl)amino]methyl phenol	354.2	δ 8.58 (d, J = 187.3, 2H), 7.05 (q, J = 8.3, 4H), 6.91 (d, J = 8.3, 2H), 6.58 (d, J = 8.4, 2H), 3.67 (d, J = 10.4, 2H), 3.58 (s, 2H), 2.63 (d, J = 18.2, 1H), 2.26 (s, 4H), 2.07 (d, J = 14.3, 4H), 1.84 (t, J = 10.2, 1H), 1.49 (d, J = 13.9, 3H), 1.09 (s, 3H), 0.56 (s, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
42	2-[(2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl)amino)methyl] phenol	354.3	δ 8.15 (d, J = 107.7, 2H), 7.03 (dt, J = 26.2, 8.3, 5H), 6.82 (m, 2H), 6.64 (t, J = 7.4, 1H), 3.68 (m, 6H), 2.58 (s, 1H), 2.24 (d, J = 6.8, 4H), 2.05 (dd, J = 21.0, 14.9, 2H), 1.78 (d, J = 4.4, 1H), 1.58 (s, 1H), 1.44 (dd, J = 21.2, 9.8, 2H), 1.10 (d, J = 18.7, 3H), 0.57 (d, J = 24.3, 3H).
43	3-[(2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl)amino)methyl] phenol	354.3	δ 8.50 (d, J = 165.4, 2H), 7.02 (m, 5H), 6.68 (d, J = 7.5, 2H), 6.47 (d, J = 7.4, 1H), 3.67 (d, J = 9.4, 2H), 3.59 (s, 2H), 2.63 (s, 2H), 2.23 (s, 4H), 2.09 (dd, J = 27.3, 13.5, 2H), 1.84 (d, J = 7.8, 1H), 1.66 (d, J = 8.6, 1H), 1.52 (d, J = 13.9, 2H), 1.09 (s, 3H), 0.56 (s, 3H).
44	[(5-methylfuran-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	355.1	δ 8.73 (d, J = 4.2, 1H), 8.18 (td, J = 8.0, 1.5, 1H), 7.80 - 7.53 (m, 2H), 7.29 (s, 1H), 4.00 (d, J = 1.4, 2H), 3.83 (dt, J = 12.4, 4.3, 1H), 3.79 - 3.63 (m, 1H), 3.10 - 2.86 (m, 1H), 2.64 - 2.44 (m, 1H), 2.45 - 2.27 (m, 3H), 2.27 - 2.11 (m, 4H), 2.02 (d, J = 14.2, 1H), 1.95 - 1.77 (m, 2H), 1.68 (dd, J = 9.5, 4.1, 1H), 1.62 - 1.39 (m, 4H), 1.26 - 1.05 (m, 1H), 0.77 (dt, J = 13.3, 9.0, 1H).
45	[(5-methylfuran-2-yl)methyl]({2-[(9R)-9-(pyrazin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	356.1	δ 8.66 (s, 1H), 8.56 (s, 1H), 8.50 (s, 1H), 6.27 (d, J = 3.2, 1H), 6.05 - 5.83 (m, 1H), 3.94 (d, J = 1.9, 2H), 3.85 - 3.59 (m, 2H), 2.89 (d, J = 5.0, 1H), 2.49 (d, J = 5.1, 1H), 2.38 (t, J = 16.0, 2H), 2.24 (s, 4H), 2.02 (dd, J = 18.2, 6.8, 2H), 1.96 - 1.88 (m, 2H), 1.59 - 1.37 (m, 5H), 1.09 (s, 1H), 0.66 (d, J = 13.4, 1H).
46	benzyl({2-[(9R)-9-(thiophen-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	356.2	δ 9.56 (s, 1H), 9.11 (s, 1H), 7.31 (m, 3H), 7.23 (m, 2H), 7.19 (dd, J = 5.1, 1.0, 1H), 6.91 (dd, J = 5.1, 3.6, 1H), 6.74 (d, J = 3.5, 1H), 3.72 (m, 4H), 2.74 (m, 1H), 2.44 (m, 1H), 2.01 (d, J = 13.9, 2H), 1.95 (dd, J = 11.7, 5.0, 1H), 1.87 (m, 2H), 1.73 (s, 5H), 1.66 (m, 2H), 1.50 (m, 3H), 1.00 (dd, J = 13.6, 8.5, 1H).
47	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(2-fluorophenyl)methyl]amine	356.3	δ 7.15 (m, 1H), 7.06 (m, 5H), 6.94 (dt, J = 18.3, 8.1, 2H), 3.69 (t, J = 7.7, 2H), 3.60 (s, 2H), 2.44 (dd, J = 11.0, 5.2, 1H), 2.22 (d, J = 20.4, 4H), 2.11 (m, 2H), 1.77 (dd, J = 6.6, 4.0, 1H), 1.57 (qd, J = 10.9, 5.5, 3H), 1.11 (s, 3H), 0.59 (s, 3H).
48	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(3-fluorophenyl)methyl]amine	356.3	δ 8.79 (d, J = 198.9, 2H), 7.19 (m, 2H), 7.05 (d, J = 8.2, 2H), 7.00 (d, J = 8.4, 2H), 6.94 (td, J = 8.4, 2.2, 1H), 6.86 (d, J = 7.6, 1H), 6.79 (d, J = 8.9, 1H), 6.36 (s, 2H), 3.69 (m, 4H), 2.65 (s, 1H), 2.24 (s, 3H), 2.11 (ddd, J = 18.3, 15.7, 11.3, 3H), 1.81 (dt, J = 12.3, 6.2, 1H), 1.63 (m, 1H), 1.48 (m, 2H), 1.10 (s, 3H), 0.56 (s, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
49	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{(4-fluorophenyl)methyl}amine	356.3	δ 8.73 (d, J = 173.6, 2H), 7.03 (m, 6H), 6.88 (t, J = 8.5, 2H), 5.32 (s, 2H), 3.68 (m, 4H), 2.61 (s, 1H), 2.24 (s, 3H), 2.11 (m, 3H), 1.78 (dt, J = 12.3, 6.2, 1H), 1.61 (m, 1H), 1.47 (m, 2H), 1.10 (s, 3H), 0.56 (s, 3H).
50	benzyl(2-{9-cyclohexyl-6-oxaspiro[4.5]decan-9-yl}ethyl)amine	356.3	δ 9.09 (d, J = 38.9, 2H), 7.35 (m, 5H), 6.43 (s, 2H), 3.93 (s, 2H), 3.54 (m, 2H), 2.85 (s, 2H), 1.63 (m, 16H), 1.10 (m, 7H), 0.84 (q, J = 11.8, 2H).
51	{2-[3-(pyridin-2-yl)-1-oxaspiro[4.5]decan-3-yl]ethyl}(thiophen-2-ylmethyl)amine	357	δ 9.79 (s, 2H), 8.66 (s, 1H), 8.21 (t, J = 7.5, 1H), 7.80 (d, J = 8.1, 1H), 7.66 (s, 1H), 7.33 (d, J = 5.0, 1H), 7.16 (s, 1H), 7.06 - 6.98 (m, 1H), 4.29 (s, 2H), 4.23 (d, J = 9.9, 1H), 3.99 (d, J = 10.0, 1H), 3.00 (s, 1H), 2.87 (s, 1H), 2.63 (t, J = 9.5, 1H), 2.39 (d, J = 8.6, 1H), 2.20 (d, J = 13.5, 1H), 2.10 (d, J = 13.6, 1H), 1.77 - 1.49 (m, 4H), 1.47 - 1.19 (m, 6H).
52	{2-[3-(pyridin-2-yl)-1-oxaspiro[4.5]decan-3-yl]ethyl}(thiophen-3-ylmethyl)amine	357	δ 9.72 (s, 2H), 8.64 (s, 1H), 8.21 (t, J = 7.5, 1H), 7.80 (d, J = 8.1, 1H), 7.66 (t, J = 5.9, 1H), 7.44 - 7.31 (m, 2H), 7.09 (d, J = 4.8, 1H), 4.23 (d, J = 9.9, 1H), 4.10 (s, 2H), 3.99 (d, J = 10.0, 1H), 2.95 (s, 1H), 2.80 (s, 1H), 2.64 (s, 1H), 2.39 (d, J = 8.7, 1H), 2.21 (d, J = 13.7, 1H), 2.10 (d, J = 13.6, 1H), 1.77 - 1.50 (m, 4H), 1.49 - 1.22 (m, 6H).
53	{2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	357.1	δ 8.67 (d, J = 4.3, 1H), 8.14 (s, 1H), 7.66 (d, J = 8.2, 1H), 7.59 (s, 1H), 7.33 (dd, J = 5.1, 1.1, 1H), 7.12 (d, J = 2.7, 1H), 7.00 (dd, J = 5.1, 3.5, 1H), 4.22 (s, 2H), 3.80 (s, 1H), 3.72 (t, J = 9.8, 1H), 3.33 - 2.70 (m, 1H), 2.70 - 2.50 (m, 1H), 2.30 (d, J = 14.0, 3H), 2.19 (dd, J = 18.0, 7.1, 1H), 1.98 (d, J = 14.1, 1H), 1.83 (d, J = 4.6, 2H), 1.76 - 1.62 (m, 1H), 1.50 (dd, J = 20.1, 13.3, 5H), 1.16 (s, 1H), 0.75 (dt, J = 13.1, 9.1, 1H).
54	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	357.2	δ 8.67 (d, J = 4.3, 1H), 8.14 (s, 1H), 7.66 (d, J = 8.2, 1H), 7.59 (s, 1H), 7.33 (dd, J = 5.1, 1.1, 1H), 7.12 (d, J = 2.7, 1H), 7.00 (dd, J = 5.1, 3.5, 1H), 4.22 (s, 2H), 3.80 (s, 1H), 3.72 (t, J = 9.8, 1H), 3.33 - 2.70 (m, 1H), 2.70 - 2.50 (m, 1H), 2.30 (d, J = 14.0, 3H), 2.19 (dd, J = 18.0, 7.1, 1H), 1.98 (d, J = 14.1, 1H), 1.83 (d, J = 4.6, 2H), 1.76 - 1.62 (m, 1H), 1.50 (dd, J = 20.1, 13.3, 5H), 1.16 (s, 1H), 0.75 (dt, J = 13.1, 9.1, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
55	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-3-ylmethyl)amine	357.2	δ 8.73 (d, J = 5.0, 1H), 8.27 (t, J = 7.5, 2H), 7.88-7.62 (m, 2H), 7.48 - 7.23 (m, 1H), 7.04 (dd, J = 4.9, 1.0, 1H), 4.02 (s, 2H), 3.90 - 3.76 (m, 1H), 3.69 (t, J = 10.0, 1H), 2.95 (s, 1H), 2.62 - 2.12 (m, 4H), 2.13 - 1.95 (m, 1H), 1.95 - 1.76 (m, 2H), 1.68 (dt, J = 13.5, 7.9, 1H), 1.62 - 1.30 (m, 5H), 1.16 (dd, J = 13.2, 6.6, 1H), 0.76 (dt, J = 13.0, 8.9, 1H).
56	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(1,3-thiazol-2-ylmethyl)amine	358	δ 8.77 (d, J = 4.3, 1H), 8.27 (t, J = 7.3, 1H), 7.86-7.65 (m, 2H), 7.43 (d, J = 3.1, 1H), 7.28 (s, 1H), 4.56 - 4.39 (m, 2H), 3.79 (dddd, J = 21.9, 19.5, 10.8, 7.1, 2H), 3.19 (td, J = 11.5, 5.3, 1H), 2.81 - 2.63 (m, 1H), 2.62 - 2.43 (m, 1H), 2.43 - 2.26 (m, 3H), 1.14 - 1.99 (m, 1H), 2.00 - 1.79 (m, 2H), 1.79 - 1.63 (m, 1H), 1.63 - 1.38 (m, 4H), 1.20 (dd, J = 13.0, 6.5, 1H), 0.79 (dt, J = 13.0, 8.9, 1H).
57	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(1,3-thiazol-5-ylmethyl)amine	358	δ 8.76 (d, J = 4.7, 1H), 8.37 (td, J = 8.1, 1.4, 1H), 8.12 - 7.72 (m, 3H), 7.29 (s, 1H), 4.37 (s, 2H), 3.93 - 3.58 (m, 2H), 3.05 (td, J = 11.7, 5.1, 1H), 2.66-2.43 (m, 2H), 2.42-2.22 (m, 3H), 2.18-1.96 (m, 1H), 1.96-1.79 (m, 2H), 1.79 - 1.39 (m, 5H), 1.18 (dd, J = 12.1, 5.5, 1H), 0.77 (dt, J = 12.9, 8.9, 1H).
58	{2-[9-(pyrazin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	358	δ 8.63 (s, 1H), 8.55 (s, 1H), 8.49 (d, J = 2.3, 1H), 7.33 (dd, J = 5.1, 1.1, 1H), 7.08 (d, J = 2.6, 1H), 7.05 - 6.97 (m, 1H), 3.73 (d, J = 36.7, 2H), 3.17 - 2.73 (m, 1H), 2.54 - 2.43 (m, 1H), 2.35 (d, J = 13.0, 2H), 2.24 - 2.11 (m, 1H), 2.05-2.15 (m, 4H), 1.51 (s, 5H), 1.14 - 1.01 (m, 1H), 0.66 (s, 1H).
59	{2-[2,2-diethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl}(thiophen-3-ylmethyl)amine	359.2	δ 8.72 (dd, J = 5.5, 1.4, 1H), 8.21 (td, J = 8.0, 1.7, 1H), 7.68 (m, 2H), 7.35 (dd, J = 2.9, 1.2, 1H), 7.30 (m, 2H), 7.04 (dd, J = 5.0, 1.3, 1H), 4.02 (s, 2H), 3.80 (dd, J = 10.0, 6.3, 1H), 3.68 (d, J = 10.8, 1H), 3.00 (m, 1H), 2.42 (m, 4H), 2.08 (d, J = 4.4, 1H), 1.78 (s, 1H), 1.71 (d, J = 14.5, 1H), 1.56 (dd, J = 14.1, 7.5, 1H), 1.40 (dd, J = 14.1, 7.4, 1H), 0.81 (m, 5H), 0.57 (t, J = 7.3, 3H).
60	{2-[2,2-diethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl}(thiophen-2-ylmethyl)amine	359.2	δ 8.70 (dd, J = 5.4, 1.4, 1H), 8.15 (d, J = 1.6, 1H), 7.66 (d, J = 8.2, 1H), 7.60 (dd, J = 6.7, 5.5, 1H), 7.33 (dd, J = 5.1, 1.2, 1H), 7.11 (d, J = 2.6, 1H), 6.99 (dd, J = 5.1, 3.6, 1H), 3.73 (d, J = 44.0, 4H), 4.02 (s, 2H), 3.80 (dd, J = 10.0, 6.3, 1H), 3.68 (d, J = 10.8, 1H), 3.00 (m, 1H), 2.42 (m, 4H), 2.08 (d, J = 4.4, 1H), 1.78 (s, 1H), 1.71 (d, J = 14.5, 1H), 1.56 (dd, J = 14.1, 7.5, 1H), 1.40 (dd, J = 14.1, 7.4, 1H), 0.81 (m, 5H), 0.57 (t, J = 7.3, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
61	{2-[2,2,6,6-tetramethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl}(thiophen-2-ylmethyl)amine	359.2	δ 8.63 (dd, J = 5.6, 1.3, 1H), 8.18 (td, J = 8.1, 1.6, 1H), 7.81 (d, J = 8.2, 1H), 7.63 (dd, J = 6.8, 5.8, 1H), 7.36 - 7.16 (m, 2H), 7.05 - 6.96 (m, 2H), 6.88 (dd, J = 5.1, 3.6, 2H), 4.13 (s, 2H), 2.80 - 2.60 (m, 2H), 2.43 (d, J = 14.7, 2H), 2.33 - 2.17 (m, 2H), 1.81 (d, J = 14.8, 2H), 1.21 (d, J = 12.2, 6H), 0.89 (s, 6H).
62	{2-[2,2,6,6-tetramethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl}(thiophen-3-ylmethyl)amine	359.2	δ 8.62 (dd, J = 5.6, 1.4, 1H), 8.19 (td, J = 8.0, 1.7, 1H), 7.81 (d, J = 8.2, 2H), 7.67 - 7.60 (m, 1H), 7.27 (dd, J = 2.9, 1.2, 1H), 7.23 - 7.17 (m, 2H), 6.95 (dd, J = 5.0, 1.3, 1H), 3.95 (s, 2H), 2.62 (d, J = 8.1, 2H), 2.41 (d, J = 14.7, 2H), 2.34 - 2.08 (m, 2H), 1.82 (d, J = 14.8, 2H), 1.21 (d, J = 13.1, 6H), 0.89 (s, 6H).
63	{2-[9-(thiophen-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	362.2	δ 9.60 (s, 1H), 9.27 (s, 1H), 7.29 (dd, J = 5.1, 1.1, 2H), 7.21 (dd, J = 5.1, 1.0, 1H), 7.03 (d, J = 2.6, 1H), 6.94 (ddd, J = 9.9, 5.1, 3.6, 2H), 6.77 (dd, J = 3.6, 1.1, 1H), 4.03 (s, 2H), 3.74 (m, 2H), 2.80 (td, J = 11.9, 4.9, 1H), 2.50 (td, J = 11.8, 5.0, 1H), 1.96 (m, 4H), 1.71 (m, 4H), 1.48 (m, 6H), 1.00 (dt, J = 12.7, 8.1, 1H).
64	{2-[9-(thiophen-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-3-ylmethyl)amine	362.2	δ 9.46 (s, 1H), 9.23 (s, 1H), 7.27 (m, 2H), 7.21 (dd, J = 5.1, 1.0, 1H), 7.00 (dt, J = 7.5, 4.4, 1H), 6.93 (dd, J = 5.1, 3.5, 1H), 6.75 (dd, J = 3.6, 1.1, 1H), 3.85 (s, 2H), 3.74 (m, 2H), 2.73 (m, 1H), 2.43 (s, 1H), 2.12 (m, 1H), 2.03 (m, 2H), 1.96 (dd, J = 12.4, 7.6, 1H), 1.87 (m, 2H), 1.70 (m, 3H), 1.48 (m, 5H), 1.00 (dt, J = 12.8, 8.1, 1H).
65	(cyclopentylmethyl){2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}amine	362.3	δ 9.23 (m, 1H), 8.73 (m, 1H), 7.25 (dd, J = 8.9, 5.2, 2H), 7.07 (t, J = 8.6, 2H), 3.73 (d, J = 10.9, 2H), 2.69 (s, 2H), 2.10 (m, 4H), 1.78 (d, J = 18.1, 3H), 1.64 (m, 7H), 1.38 (s, 2H), 1.28 (s, 1H), 1.10 (d, J = 16.3, 3H), 0.84 (s, 4H), 0.53 (s, 3H).
66	(cyclopentylmethyl){2-[4-(4-fluorophenyl)-2,2,6,6-tetramethyloxan-4-yl]ethyl}amine	362.3	δ 8.64 (s, 2H), 7.22 (dd, J = 8.9, 5.1, 2H), 6.95 (t, J = 8.6, 2H), 3.25 (s, 2H), 2.61 (s, 2H), 2.43 (s, 2H), 2.24 (d, J = 14.3, 2H), 1.91 (m, 2H), 1.68 (m, 2H), 1.60 (d, J = 14.3, 2H), 1.49 (m, 4H), 1.18 (s, 6H), 1.03 (dd, J = 12.4, 7.3, 2H), 0.93 (s, 6H).
67	(2-{9-cyclohexyl-6-oxaspiro[4.5]decan-9-yl}ethyl)(thiophen-2-ylmethyl)amine	362.3	δ 9.21 (d, J = 25.7, 2H), 7.33 (dd, J = 5.1, 1.1, 2H), 7.14 (d, J = 2.7, 1H), 7.00 (dd, J = 5.1, 3.6, 1H), 4.19 (s, 2H), 3.56 (m, 2H), 2.92 (s, 2H), 1.65 (m, 17H), 1.12 (m, 7H), 0.87 (dd, J = 23.8, 11.9, 2H).
68	(2-{9-cyclohexyl-6-oxaspiro[4.5]decan-9-yl}ethyl)(thiophen-3-ylmethyl)amine	362.3	δ 9.07 (d, J = 31.8, 2H), 7.37 (ddd, J = 7.9, 3.9, 2.1, 2H), 7.10 (dd, J = 5.0, 1.3, 1H), 6.37 (s, 2H), 4.04 (s, 2H), 3.55 (m, 2H), 2.87 (s, 2H), 1.64 (m, 16H), 1.12 (m, 7H), 0.85 (q, J = 11.8, 2H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
69	2-{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}-2,3-dihydro-1H-isoindole	363.1	δ 8.77 (d, J = 4.0, 1H), 8.09 (td, J = 8.0, 1.7, 1H), 7.64 (d, J = 8.1, 1H), 7.55 (dd, J = 7.1, 5.8, 1H), 7.35 (dd, J = 5.6, 3.2, 2H), 7.24 (d, J = 3.6, 2H), 4.76 (m, 4H), 4.21 (brs, 1H), 3.77 (m, 2H), 3.30 (m, 1H), 2.80 (td, J = 12.3, 4.4, 1H), 2.49 (td, J = 12.9, 4.5, 1H), 2.38 (t, J = 15.1, 2H), 2.23 (td, J = 12.9, 4.2, 1H), 2.07 (d, J = 14.0, 1H), 1.87 (ddd, J = 24.1, 11.9, 7.1, 2H), 1.69 (m, 1H), 1.51 (dt, J = 24.2, 10.9, 4H), 1.15 (m, 1H), 0.78 (dt, J = 13.4, 9.0, 1H).
70	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}dipropylamine	364.4	δ 11.44 (s, 1H), 7.28 (m, 2H), 7.10 (m, 2H), 3.75 (m, 2H), 2.88 (m, 5H), 2.27 (m, 3H), 1.97 (td, J = 12.7, 3.9, 1H), 1.80 (td, J = 12.6, 4.9, 1H), 1.66 (m, 2H), 1.46 (m, 6H), 1.04 (m, 1H), 0.88 (m, 10H), 0.55 (m, 3H).
71	(2-phenylethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	365.1	δ 8.51 (dd, J = 5.3, 1.3, 1H), 8.04 (td, J = 7.9, 1.7, 1H), 7.56 (d, J = 8.1, 1H), 7.49 (dd, J = 7.1, 5.8, 1H), 7.25 - 7.12 (m, 6H), 7.10 - 7.03 (m, 2H), 3.88 - 3.47 (m, 3H), 3.01 (d, J = 7.5, 2H), 2.85 (t, J = 7.8, 2H), 2.44 (s, 1H), 2.38-2.17 (m, 3H), 2.17-1.99 (m, 1H), 1.92 (d, J = 14.1, 1H), 1.84 - 1.66 (m, 3H), 1.58 (d, J = 5.1, 1H), 1.40 (ddd, J = 15.2, 12.1, 8.9, 4H), 1.05 (d, J = 6.5, 1H), 0.65 (d, J = 13.4, 1H).
72	(2-phenylethyl){2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	365.3	δ 8.58 (d, J = 4.8, 1H), 8.07 (t, J = 7.9, 1H), 7.61 (s, 1H), 7.52 (dd, J = 12.0, 6.3, 1H), 7.27 (m, 3H), 7.20 (m, 2H), 4.04 (d, J = 3.2, 2H), 3.76 (ddd, J = 19.4, 12.6, 8.9, 2H), 3.05 (s, 1H), 2.53 (m, 2H), 2.29 (d, J = 43.6, 5H), 1.96 (d, J = 13.9, 1H), 1.80 (m, 2H), 1.68 (s, 1H), 1.50 (ddd, J = 20.5, 13.1, 7.0, 4H), 1.17 (s, 1H), 0.75 (m, 1H).
73	benzyl{2-[9-(6-methylpyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	365.7	δ 9.49 (s, 2H), 8.18 (t, J = 7.9, 1H), 7.55 (dd, J = 23.1, 7.8, 2H), 7.35 (s, 5H), 5.87 (s, 3H), 4.00 (s, 2H), 3.88 - 3.66 (m, 2H), 3.00 (s, 1H), 2.80 (s, 3H), 2.65 (d, J = 12.5, 1H), 2.53 (s, 1H), 2.31 (d, J = 14.3, 2H), 2.20 (d, J = 13.5, 1H), 2.11 - 2.00 (m, 1H), 1.97 - 1.80 (m, 2H), 1.70 (d, J = 5.3, 1H), 1.52 (ddd, J = 29.7, 17.1, 7.4, 4H), 1.28 (t, J = 7.1, 1H), 0.95 - 0.79 (m, 1H). (ddd, J = 29.7, 17.1, 7.4, 4H), 1.28 (t, J = 7.1, 1H), 0.92-0.77 (m, 1H).
74	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}(2-phenylpropan-2-yl)amine	366.3	¹ H NMR (400 MHz, 324.3, CDCl ₃) δ 8.50 (d, J = 223.4, 2H), 7.25 (s, 5H), 6.95 (d, 338.3 J = 8.1, 2H), 6.87 (d, J = 8.3, 2H), 5.69 (s, 3H), 3.62 (dd, J = 6.8, 2.5, 2H), 2.38 (dd, J = 15.7, 13.2, 1H), 2.22 (s, 3H), 1.98 (m, 2H), 1.80 (m, 2H), 1.63 (m, 1H), 1.56 (s, 3H), 1.51 (s, 3H), 1.47 (d, J = 14.1, 1H), 1.39 (dd, J = 10.5, 4.0, 1H), 1.06 (s, 3H), 0.53 (s, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
75	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{2-(pyridin-3-yl)ethyl}amine	367.1	δ 8.90 (s, 1H), 8.75 (d, J = 4.4, 1H), 8.61 (d, J = 5.2, 1H), 8.41-8.28 (m, 2H), 7.87 - 7.70 (m, 3H), 3.81 (s, 1H), 3.71 (s, 1H), 3.29 (t, J = 10.5, 3H), 2.97 (d, J = 7.3, 1H), 2.44 (s, 2H), 2.33 (t, J = 11.9, 2H), 2.21 (dt, J = 24.1, 11.9, 1H), 2.07 (d, J = 14.3, 1H), 1.88 (d, J = 10.3, 2H), 1.65 (dd, J = 16.4, 9.9, 1H), 1.60-1.44 (m, 5H), 1.19 (s, 1H), 0.81 (d, J = 13.1, 1H).
76	[(2-methylpyrimidin-5-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	367.1	δ 8.57 (s, 2H), 7.83 - 7.66 (m, 1H), 7.33 (s, 4H), 7.21 (dt, J = 10.8, 2.9, 1H), 3.93 (s, 1H), 3.69 (s, 2H), 2.65 (s, 1H), 2.40 - 2.20 (m, 3H), 2.09 (s, 2H), 1.87 (s, 2H), 1.76 - 1.50 (m, 3H), 1.42 (ddd, J = 33.3, 13.0, 3.9, 2H), 1.22 (td, J = 7.3, 1.9, 1H), 1.02 (s, 1H), 0.71 - 0.54 (m, 1H).
77	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{(2-methoxyphenyl)methyl}amine	368.3	δ 7.16 (m, 6H), 6.85 (dd, J = 18.0, 7.8, 2H), 3.80 (s, 3H), 3.61 (d, J = 1.9, 2H), 3.51 (s, 2H), 2.45 (d, J = 5.2, 1H), 2.35 (s, 4H), 2.15 (m, 2H), 1.81 (m, 1H), 1.66 (s, 4H), 1.20 (s, 3H), 0.69 (s, 3H).
78	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{(3-methoxyphenyl)methyl}amine	368.3	δ 9.28 (s, 1H), 8.80 (s, 1H), 7.10 (m, 1H), 7.01 (q, J = 8.4, 4H), 6.74 (dd, J = 8.2, 2.0, 1H), 6.65 (dd, J = 15.6, 4.8, 2H), 3.66 (m, 7H), 2.64 (s, 4H), 2.24 (s, 3H), 2.09 (m, 3H), 1.82 (m, 1H), 1.64 (m, 1H), 1.48 (ddd, J = 13.4, 9.8, 8.8, 2H), 1.10 (s, 3H), 0.57 (s, 3H).
79	benzyl{2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	368.3	δ 8.82 (d, J = 134.2, 2H), 7.31 (m, 3H), 7.16 (m, 4H), 7.00 (dd, J = 10.7, 6.5, 2H), 3.72 (m, 4H), 2.70 (s, 1H), 2.28 (s, 1H), 1.92 (m, 6H), 1.62 (m, 2H), 1.46 (m, 4H), 1.23 (m, 1H), 0.77 (dt, J = 13.6, 8.8, 1H).
80	benzyl{2-[(9S)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	368.3	δ 9.09 (s, 1H), 8.74 (s, 1H), 7.31 (m, 3H), 7.16 (m, 4H), 7.00 (t, J = 8.6, 2H), 3.73 (m, 4H), 2.67 (s, 1H), 2.26 (s, 1H), 2.02 (s, 2H), 1.94 (td, J = 12.6, 4.7, 1H), 1.85 (d, J = 13.9, 3H), 1.62 (s, 2H), 1.46 (dd, J = 7.8, 4.0, 4H), 1.24 (d, J = 12.7, 1H), 0.77 (dt, J = 13.6, 8.7, 1H).
81	benzyl{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	368.3	δ 7.24-7.17 (m, 2H), 7.16 - 7.09 (m, 3H), 7.01 (d, J = 7.8, 2H), 6.89 (d, J = 8.0, 2H), 3.68 (ddd, J = 11.8, 5.0, 1.3, 1H), 3.62 - 3.49 (m, 3H), 2.32 (t, J = 7.3, 2H), 2.25 (s, 3H), 2.22 - 2.13 (m, 1H), 1.93 (dtd, J = 15.7, 7.7, 3.8, 1H), 1.81 - 1.66 (m, 2H), 1.65 - 1.56 (m, 1H), 1.37 (d, J = 20.2, 1H), 1.20 - 1.05 (m, 2H), 1.01-1.02 (m, 2H), 0.86 (t, J = 12.7, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
82	2-[(9R)-9-(2-{4H,5H,6H-thieno[2,3-c]pyrrol-5-yl}ethyl)-6-oxaspiro[4.5]decan-9-yl]pyridine	369	δ 8.59 (ddd, J = 4.8, 1.9, 0.9, 1H), 7.64 (m, 1H), 7.32 (t, J = 5.9, 1H), 7.15 (d, J = 4.9, 1H), 7.12 (ddd, J = 7.5, 4.8, 1.0, 1H), 6.74 (d, J = 4.9, 1H), 3.80 (m, 4H), 3.68 (m, 2H), 2.63 (td, J = 11.6, 5.1, 1H), 2.49 (dd, J = 13.8, 2.2, 1H), 2.37 (dd, J = 13.7, 2.0, 1H), 2.16 (td, J = 11.6, 4.4, 1H), 2.05 (m, 1H), 1.79 (m, 3H), 1.62 (d, J = 7.8, 2H), 1.50 (m, 3H), 1.40 (m, 1H), 1.14 (ddd, J = 9.7, 7.6, 3.2, 1H), 0.72 (dt, J = 13.4, 8.9, 1H).
83	[(4,5-dimethylfuran-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	369.1	δ 10.28 (brs, 1H), 9.39 (brs, 1H), 8.70 (d, J = 4.6, 1H), 8.12 (t, J = 7.5, 1H), 7.65 (d, J = 8.1, 1H), 7.58 (m, 1H), 6.14 (s, 1H), 3.91 (q, J = 14.4, 2H), 3.75 (m, 2H), 2.95 (dd, J = 10.9, 5.9, 1H), 2.51 (t, J = 9.7, 1H), 2.33 (m, 3H), 2.10 (s, 3H), 1.99 (d, J = 14.1, 1H), 1.82 (m, 5H), 1.68 (m, 1H), 1.48 (m, 4H), 1.15 (m, 1H), 0.74 (dt, J = 13.2, 8.9, 1H).
84	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(pyridin-4-ylmethyl)amine	369.2	δ 8.53 (s, 2H), 7.78 (s, 3H), 7.29 - 7.05 (m, 6H), 6.96 (t, J = 8.4, 3H), 4.07 (s, 2H), 3.66 (d, J = 12.5, 2H), 2.83 (s, 1H), 2.37 (s, 1H), 2.11 (d, J = 13.7, 1H), 2.01 (d, J = 13.3, 2H), 1.83 (d, J = 14.0, 2H), 1.49 (t, J = 61.9, 9H), 1.17 (s, 2H), 0.70 (dt, J = 17.4, 8.9, 1H).
85	2-[(2-{4-(4-methoxyphenyl)-2,2-dimethyloxan-4-yl}ethyl)amino)methyl] phenol	370.3	δ 8.05 (d, J = 152.9, 2H), 7.08 (m, 1H), 7.01 (d, J = 8.9, 2H), 6.82 (m, 2H), 6.76 (d, J = 8.8, 2H), 6.69 (t, J = 7.3, 1H), 4.00 (s, 2H), 3.77 (s, 2H), 3.70 (s, 3H), 3.65 (dd, J = 6.9, 2.6, 2H), 2.61 (s, 1H), 2.23 (s, 1H), 2.04 (dd, J = 23.7, 13.9, 2H), 1.93 (s, 1H), 1.81 (td, J = 12.5, 4.9, 1H), 1.60 (td, J = 12.7, 4.8, 1H), 1.48 (d, J = 13.9, 2H), 1.08 (s, 3H), 0.55 (s, 3H).
86	benzyl({2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl})amine	370.3	δ 7.26 - 7.14 (m, 3H), 7.13 - 7.02 (m, 4H), 6.91 (t, J = 8.6, 2H), 3.69 - 3.47 (m, 4H), 2.51 (td, J = 12.2, 4.7, 1H), 2.14 - 1.94 (m, 3H), 1.83 (td, J = 12.7, 4.3, 1H), 1.64 (td, J = 12.6, 4.7, 1H), 1.56 - 1.35 (m, 3H), 1.27
			(tt, J = 27.2, 13.7, 1H), 0.95 (dq, J = 14.7, 7.4, 1H), 0.84 - 0.58 (m, 4H), 0.43 (t, J = 7.4, 3H).
87	benzyl({2-[4-(4-fluorophenyl)-2,2,6,6-tetramethyloxan-4-yl]ethyl})amine	370.3	δ 9.15 (s, 2H), 7.32 (m, 3H), 7.25 (m, 2H), 7.18 (dd, J = 7.3, 2.1, 2H), 6.99 (dd, J = 12.0, 5.3, 2H), 3.72 (s, 2H), 2.34 (dd, J = 53.2, 23.4, 2H), 1.91 (dd, J = 10.4, 6.5, 2H), 1.68 (d, J = 14.3, 2H), 1.27 (s, 6H), 1.02 (s, 6H).
88	[(2,3-dimethoxyphenyl)methyl]({2-[4-(4-methylphenyl)oxan-4-yl]ethyl})amine	370.3	δ 7.13 (s, 4H), 6.95 (m, 1H), 6.80 (dd, J = 8.2, 1.4, 1H), 6.72 (dd, J = 7.6, 1.4, 1H), 3.83 (s, 3H), 3.75 (m, 5H), 3.63 (s, 2H), 3.54 (ddd, J = 11.6, 9.1, 2.7, 2H), 2.31 (m, 5H), 2.10 (m, 3H), 1.82 (ddd, J = 13.3, 8.3, 3.7, 4H)

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
89	[(3-methylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	371.1	δ 9.68 (s, 1H), 8.75 (s, 1H), 8.16 (m, 1H), 7.74 (d, J = 27.0, 2H), 7.27 (d, J = 1.5, 1H), 6.85 (d, J = 5.1, 1H), 4.10 (m, 2H), 3.84 (d, J = 12.7, 1H), 3.66 (d, J = 10.3, 1H), 2.96 (m, 1H), 2.69 (m, 1H), 2.54 (m, 3H), 2.35 (m, 4H), 2.11 (d, J = 14.0, 1H), 1.87 (d, J = 10.3, 3H), 1.57 (m, 5H), 1.06 (m, 1H), 0.78 (d, J = 12.8, 1H).
90	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[2-(thiophen-2-yl)ethyl]amine	371.1	δ 8.80-8.66 (m, 1H), 8.45 - 8.25 (m, 1H), 7.84-7.63 (m, 2H), 7.16 (dd, J = 5.1, 1.1, 1H), 6.91 (dd, J = 5.1, 3.5, 1H), 6.83 (dd, J = 3.4, 0.9, 1H), 3.83 (tt, J = 13.7, 6.9, 1H), 3.69 (dd, J = 20.1, 10.1, 1H), 3.16 (s, 4H), 3.02 (s, 1H), 2.61 - 2.22 (m, 5H), 2.20 - 1.98 (m, 1H), 1.98 - 1.77 (m, 2H), 1.76 - 1.63 (m, 1H), 1.50 (tdd, J = 12.3, 10.9, 5.3, 4H), 1.17 (dd, J = 7.9, 5.2, 1H), 0.76 (dt, J = 13.0, 8.8, 1H).
91	[(2-methylthiophen-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	371.1	δ 8.68 (d, J = 5.4, 1H), 8.26 (s, 1H), 7.82 - 7.63 (m, 2H), 7.05 (t, J = 10.0, 1H), 6.94 (d, J = 5.3, 1H), 3.96 (s, 2H), 3.82 (s, 1H), 3.72 (s, 1H), 3.03 (s, 1H), 2.50 (d, J = 15.9, 2H), 2.39 (s, 3H), 2.30 (dd, J = 12.6, 7.5, 3H), 2.02 (d, J = 14.2, 1H), 1.92 - 1.79 (m, 2H), 1.70 (dt, J = 14.5, 10.2, 1H), 1.64 - 1.38 (m, 4H), 1.25 - 1.13 (m, 1H), 0.79 (d, J = 13.2, 1H).
92	[(5-methylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	371.2	δ 8.71 (d, J = 4.7, 1H), 8.14 (t, J = 7.6, 1H), 7.78-7.48 (m, 2H), 6.86 (d, J = 3.4, 1H), 6.78 - 6.53 (m, 1H), 4.09 (s, 2H), 3.76 (ddd, J = 40.6, 14.3, 7.2, 2H), 3.17 - 2.85 (m, 1H), 2.64 - 2.23 (m, 4H), 2.16 (dd, J = 16.4, 8.6, 1H), 1.99 (d, J = 14.2, 1H), 1.89 - 1.75 (m, 2H), 1.75 - 1.61 (m, 1H), 1.61 - 1.35 (m, 4H), 1.24-1.05 (m, 1H), 0.74 (dt, J = 13.2, 8.9, 1H).
93	{2-[9-(6-methylpyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-3-ylmethyl)amine	371.2	δ 9.47 (d, J = 86.3, 2H), 8.17 (t, J = 8.0, 1H), 7.58 (d, J = 8.0, 1H), 7.52 (d, J = 7.8, 1H), 7.39 (d, J = 1.9, 1H), 7.31 - 7.29 (m, 1H), 7.08 (dd, J = 5.0, 1.0, 1H), 6.43 (s, 3H), 4.11 - 3.95 (m, 2H), 3.91 - 3.67 (m, 2H), 2.97 (s, 1H), 2.81 (s, 3H), 2.61 (t, J = 12.6, 1H), 2.47 (t, J = 10.1, 1H), 2.43 - 2.15 (m, 3H), 2.15 - 1.99 (m, 1H), 1.87 (dd, J = 12.2, 6.8, 2H), 1.70 (dt, J = 12.7, 6.2, 1H), 1.63 - 1.40 (m, 4H), 1.28 - 1.20 (m, 1H), 0.84 (dt, J = 13.3, 9.0, 1H).
94	{2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}(1H-pyrrol-2-ylmethyl)amine	371.3	δ 7.09 (dd, J = 8.9, 5.1, 2H), 6.93 (dd, J = 11.7, 5.5, 2H), 6.71 (d, J = 2.3, 1H), 5.98 (s, 2H), 3.83 (s, 2H), 3.61 (m, 2H), 2.56 (m, 1H), 2.08 (t, J = 12.1, 3H), 1.68 (s, 3H), 1.48 (d, J = 14.6, 2H), 1.40 (d, J = 14.1, 2H), 1.29 (m, 3H), 1.05 (m, 3H), 0.58 (s, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
95	{2-[9-(6-methyl pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	371.3	δ 9.48 (s, 1H), 8.08 (t, J = 7.9, 1H), 7.48 (d, J = 8.0, 1H), 7.42 (d, J = 7.8, 1H), 7.22 (dd, J = 5.1, 0.8, 1H), 7.04 (d, J = 2.9, 1H), 6.88 (dd, J = 5.1, 3.5, 1H), 5.95 (s, 3H), 4.13 (s, 2H), 3.66 (ddd, J = 18.7, 12.8, 9.1, 2H), 2.91 (s, 1H), 2.71 (s, 3H), 2.60 - 2.40 (m, 2H), 2.18 (dd, J = 48.6, 14.1, 3H), 1.96 (d, J = 14.2, 1H), 1.88 - 1.68 (m, 2H), 1.71 - 1.54 (m, 1H), 1.56 - 1.31 (m, 4H), 1.20 - 1.05 (m, 1H), 0.83 - 0.63 (m, 1H).
96	[(4-methylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	371.3	δ 9.63 (s, 1H), 8.61 (d, J = 4.1, 1H), 8.08 (t, J = 7.8, 1H), 7.61 (d, J = 8.1, 1H), 7.53 (dd, J = 7.0, 5.6, 1H), 6.91 (s, 1H), 6.88 (s, 1H), 4.14 (m, 2H), 3.75 (dt, J = 19.0, 11.1, 2H), 3.02 (m, 1H), 2.61 (m, 1H), 2.40 (brs, 1H), 2.27 (m, 4H), 2.19 (d, J = 0.8, 3H), 1.95 (d, J = 14.0, 1H), 1.79 (m, 2H), 1.66 (dd, J = 12.1, 5.9, 1H), 1.47 (m, 4H), 1.16 (m, 1H), 0.74 (dt, J = 13.1, 8.9, 1H).
97	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(5-methylfuran-2-yl)methyl]amine	372	δ 7.12 (dd, J = 8.9, 5.2, 2H), 6.94 (t, J = 8.6, 2H), 6.10 (d, J = 3.1, 1H), 5.79 (dd, J = 3.1, 0.9, 1H), 3.77 (m, 2H), 3.72 - 3.49 (m, 2H), 2.63 (s, 1H), 2.19 (s, 1H), 2.13 - 2.08 (m, 3H), 2.06 (s, 1H), 1.98 (dd, J = 13.8, 1.3, 1H), 1.89 (td, J = 12.7, 4.5, 1H), 1.80 (dd, J = 13.1, 7.1, 2H), 1.71 (dd, J = 13.2, 6.0, 1H), 1.59 (ddd, J = 14.2, 9.4, 5.4, 2H), 1.50 - 1.28 (m, 4H), 1.25 - 1.09 (m, 1H), 0.71 (dt, J = 13.5, 8.8, 1H).
98	[(4-methyl-1,3-thiazol-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	372.1	δ 8.68 (dd, J = 5.3, 1.2, 1H), 8.25 (s, 1H), 8.09 (td, J = 8.0, 1.7, 1H), 7.63 (d, J = 8.1, 1H), 7.54 (dd, J = 7.1, 5.7, 1H), 6.94 (d, J = 0.9, 1H), 4.37 (m, 2H), 3.76 (m, 2H), 3.14 (td, J = 11.2, 5.9, 1H), 2.73 (td, J = 11.4, 4.7, 1H), 2.40 (m, 4H), 2.27 (m, 3H), 2.00 (m, 1H), 1.83 (ddd, J = 13.8, 9.3, 4.4, 2H), 1.66 (m, 1H), 1.49 (m, 4H), 1.19 (m, 1H), 0.78 (dt, J = 13.3, 9.0, 1H).
99	[(2-methyl-1,3-thiazol-5-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	372.1	δ 8.71 (d, J = 4.3, 1H), 8.33 (td, J = 8.0, 1.5, 1H), 7.77 (m, 2H), 7.69 (s, 1H), 5.53 (s, 1H), 4.28 (m, 2H), 3.78 (m, 2H), 3.04 (td, J = 11.4, 5.4, 1H), 2.73 (s, 3H), 2.56 (m, 2H), 2.30 (t, J = 15.3, 3H), 2.04 (m, 1H), 1.88 (ddd, J = 19.6, 11.5, 7.0, 2H), 1.68 (m, 1H), 1.49 (m, 4H), 1.18 (m, 1H), 0.77 (dt, J = 13.1, 9.0, 1H).
100	[(4-methyl-1,3-thiazol-5-yl)methyl]({2-[(9R)-9-	372.1	δ 13.17 (s, 1H), 9.91 (s, 1H), 8.88 (s, 1H), 8.69 (d, J = 4.9, 1H), 8.31 (t, J = 7.4, 1H), 7.75 (t, J = 7.9, 2H),

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
	(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine		4.25 (m, 2H), 3.77 (m, 2H), 3.04 (td, J = 11.5, 5.0, 1H), 2.57 (dt, J = 10.7, 7.9, 1H), 2.34 (m, 7H), 2.02 (m, 1H), 1.86 (ddd, J = 26.5, 13.3, 8.2, 2H), 1.66 (dt, J = 13.6, 8.4, 1H), 1.50 (m, 4H), 1.15 (dd, J = 13.2, 6.6, 1H), 0.73 (dt, J = 13.0, 8.9, 1H).
101	[(2-chlorophenyl)methyl]({ 2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}amine	372.2	δ 7.21 (m, 1H), 7.07 (m, 5H), 7.02 (d, J = 8.2, 2H), 3.69 (m, 2H), 3.58 (d, J = 1.0, 2H), 2.37 (td, J = 10.9, 5.3, 1H), 2.22 (m, 4H), 2.07 (ddd, J = 14.2, 9.9, 3.8, 2H), 1.74 (ddd, J = 13.2, 10.5, 5.1, 1H), 1.55 (m, 3H), 1.43 (s, 2H), 1.10 (s, 3H), 0.58 (s, 3H).
102	[(3-chlorophenyl)methyl]({ 2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}amine	372.2	δ 9.27 (d, J = 168.2, 2H), 7.19 (m, 2H), 7.11 (m, 2H), 7.04 (d, J = 8.2, 2H), 6.99 (d, J = 8.2, 3H), 3.67 (m, 2H), 3.58 (s, 2H), 2.57 (s, 1H), 2.33 (d, J = 12.1, 2H), 2.23 (s, 3H), 2.07 (m, 3H), 1.80 (td, J = 12.5, 4.6, 1H), 1.62 (m, 1H), 1.47 (m, 2H), 1.09 (s, 3H), 0.56 (s, 3H).
103	[(4-chlorophenyl)methyl]({ 2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}amine	372.2	δ 8.80 (d, J = 192.6, 2H), 7.19 (t, J = 4.2, 3H), 7.02 (m, 6H), 4.06 (s, 3H), 3.68 (dd, J = 12.4, 10.2, 4H), 2.62 (s, 1H), 2.24 (d, J = 13.6, 3H), 2.11 (ddd, J = 21.2, 15.6, 7.6, 3H), 1.80 (dt, J = 12.3, 6.3, 1H), 1.64 (m, 1H), 1.49 (m, 2H), 1.11 (s, 3H), 0.57 (s, 3H).
104	6-[9-{2-[(thiophen-2-ylmethyl)amino]ethyl}-6-oxaspiro[4.5]decan-9-yl]pyridin-3-ol	373	¹ H NMR (400 MHz, CD ₃ CN) δ 8.18 (t, J = 1.7, 1H), 8.11 (brs, 1H), 7.49 (dd, J = 5.1, 1.1, 1H), 7.34 (d, J = 1.7, 2H), 7.18 (d, J = 2.7, 1H), 7.06 (dd, J = 5.1, 3.6, 1H), 4.24 (s, 2H), 3.67 (m, 2H), 2.95 (m, 1H), 2.73 (brs, 1H), 2.51 (d, J = 4.3, 1H), 2.29 (t, J = 11.0, 2H), 2.08 (m, 2H), 1.84 (m, 2H), 1.72 (t, J = 8.5, 1H), 1.62 (dd, J = 14.4, 6.5, 2H), 1.48 (dt, J = 23.5, 7.0, 4H), 1.15 (m, 1H), 0.73 (dt, J = 12.7, 8.7, 1H).
105	6-[9-{2-[(thiophen-2-ylmethyl)amino]ethyl}-6-oxaspiro[4.5]decan-9-yl]pyridin-2-ol	373	δ 7.52 (d, J = 16.2, 1H), 7.29 (d, J = 1.1, 1H), 7.12 (d, J = 2.7, 1H), 6.97 (dd, J = 5.1, 3.6, 1H), 6.51 (d, J = 8.9, 1H), 6.27 (d, J = 7.2, 1H), 4.16 (s, 2H), 3.71 (s, 2H), 2.85 (dd, J = 13.9, 7.6, 1H), 2.68 (dd, J = 18.4, 9.5, 1H), 2.31 (m, 2H), 1.94 (d, J = 13.6, 2H), 1.59 (m, 10H), 0.90 (m, 1H).
106	[(5-methylthiophen-2-yl)methyl]({2-[2,2,6,6-tetramethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl}amine	373.2	δ 8.73 (dd, J = 5.5, 1.4, 2H), 8.24 (td, J = 8.0, 1.6, 1H), 7.87 (d, J = 8.2, 1H), 7.69 (dd, J = 7.0, 6.1, 1H), 6.83 (dd, J = 20.2, 3.4, 1H), 6.67 - 6.48 (m, 1H), 4.09 (s, 2H), 2.83 - 2.69 (m, 2H), 2.52 (dd, J = 19.1, 11.7, 3H), 2.41 (d, J = 0.5, 3H), 2.37 - 2.21 (m, 2H), 1.89 (d, J = 14.8, 2H), 1.31 (s, 6H), 0.98 (s, 6H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
107	2-(9-{2-[(thiophen-2-ylmethyl)amino]ethyl}-6-oxaspiro[4.5]decan-9-yl)pyridin-4-ol	373.2	δ 9.46 (m, 2H), 7.95 (d, J = 6.6, 1H), 7.25 (d, J = 5.1, 1H), 7.10 (s, 1H), 7.03 (t, J = 5.8, 2H), 6.90 (dd, J = 5.1, 3.6, 1H), 4.10 (s, 2H), 3.62 (m, 2H), 2.84 (s, 1H), 2.49 (s, 1H), 2.28 (s, 1H), 2.06 (dd, J = 44.3, 14.1, 3H), 1.66 (m, 4H), 1.35 (ddd, J = 72.6, 39.8, 18.9, 6H), 0.68 (s, 1H).
108	[(4-methylthiophen-2-yl)methyl]({2-[2,2,6,6-tetramethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl})amine	373.3	δ 8.75 (d, J = 4.6, 1H), 8.35 (td, J = 8.1, 1.3, 1H), 7.96 (d, J = 8.2, 1H), 7.86 - 7.74 (m, 1H), 6.95 - 6.80 (m, 2H), 4.14 (s, 2H), 2.87 - 2.68 (m, 2H), 2.52 (d, J = 14.8, 2H), 2.45 - 2.29 (m, 2H), 2.18 (d, J = 0.7, 3H), 1.93 (d, J = 14.9, 2H), 1.31 (s, 6H), 0.98 (s, 6H).
109	dibutyl({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	373.4	δ 8.78 (d, J = 4.6, 1H), 8.05 (t, J = 7.5, 1H), 7.62 (d, J = 8.0, 1H), 7.50 (m, 1H), 3.80 (m, 2H), 3.06 (t, J = 10.5, 1H), 2.90 (s, 4H), 2.42 (m, 4H), 2.02 (m, 2H), 1.83 (m, 2H), 1.68 (tt, J = 13.3, 6.8, 1H), 1.43 (m, 12H), 1.15 (dd, J = 13.2, 5.7, 1H), 0.91 (dt, J = 11.8, 7.1, 6H), 0.72 (dt, J = 13.3, 9.0, 1H).
110	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-3-ylmethyl)amine	374.2	δ 7.33 - 7.23 (m, 7H), 7.19 (dd, J = 8.9, 5.2, 2H), 7.04 (t, J = 8.6, 2H), 6.98 (dd, J = 5.0, 1.3, 1H), 3.84 (s, 2H), 3.79 - 3.69 (m, 2H), 2.67 (s, 1H), 2.19 - 1.74 (m, 22H), 1.66 (ddd, J = 14.0, 9.3, 4.6, 3H), 1.48 (ddd, J = 23.7, 15.2, 8.6, 4H), 1.28 (s, 1H), 0.99 - 0.64 (m, 1H).
111	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	374.2	δ 9.04 (d, J = 106.1, 2H), 7.21 (dd, J = 5.1, 1.1, 1H), 7.10 (m, 2H), 6.92 (m, 3H), 6.86 (dd, J = 5.1, 3.6, 1H), 3.93 (s, 2H), 3.64 (m, 3H), 2.63 (d, J = 7.9, 1H), 2.22 (t, J = 9.7, 1H), 2.05 (d, J = 14.1, 1H), 1.97 (d, J = 13.9, 1H), 1.88 (td, J = 12.7, 4.6, 1H), 1.75 (m, 3H), 1.57 (m, 2H), 1.38 (m, 3H), 1.17 (dd, J = 14.1, 6.1, 1H), 0.70 (dt, J = 13.6, 8.8, 1H).
112	(cyclopentylmethyl){2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}amine	374.3	δ 7.15 (dd, J = 8.9, 5.2, 2H), 6.96 (s, 2H), 3.64 (d, J = 13.0, 3H), 2.59 (s, 3H), 2.11 (m, 3H), 1.94 (dd, J = 10.4, 5.7, 2H), 1.68 (dd, J = 12.4, 4.8, 2H), 1.53 (m, 8H), 1.31 (d, J = 19.9, 4H), 1.03 (s, 7H), 0.65 (m, 1H).
113	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}(thiophen-3-ylmethyl)amine	376.2	δ 7.20 - 7.13 (m, 8H), 7.09 (dd, J = 8.9, 5.2, 2H), 6.93 (t, J = 8.6, 2H), 6.87 (dd, J = 4.9, 1.3, 1H), 3.70 (s, 2H), 3.61 (d, J = 2.3, 2H), 2.56 (s, 1H), 2.02 (d, J = 14.1, 3H), 1.75 (s, 11H), 1.44 (d, J = 14.2, 5H), 0.95 (dd, J = 14.5, 7.4, 1H), 0.73 (t, J = 7.5, 5H), 0.43 (t, J = 7.4, 4H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
114	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}(thiophen-2-ylmethyl)amine	376.2	δ 7.25 - 7.15 (m, 3H), 7.15 - 7.02 (m, 4H), 6.91 (t, J = 8.6, 2H), 3.82 - 3.36 (m, 4H), 2.51 (td, J = 12.2, 4.7, 1H), 2.12 - 1.94 (m, 3H), 1.83 (td, J = 12.7, 4.3, 1H), 1.64 (td, J = 12.6, 4.7, 1H), 1.55 - 1.35 (m, 3H), 1.28 (dq, J = 14.7, 7.4, 1H), 0.95 (dq, J = 14.7, 7.4, 1H), 0.80 - 0.64 (m, 4H), 0.43 (t, J = 7.4, 3H).
115	{2-[4-(4-fluorophenyl)-2,2,6,6-tetramethyloxan-4-yl]ethyl}(thiophen-3-ylmethyl)amine	376.2	δ 7.28 (m, 4H), 7.00 (ddd, J = 6.7, 6.3, 3.2, 3H), 3.82 (s, 3H), 2.46 (s, 1H), 2.28 (d, J = 14.3, 1H), 1.92 (m, 1H), 1.57 (m, 2H), 1.69 (d, J = 14.4, 2H), 1.28 (s, 6H), 1.02 (s, 6H).
116	{2-[4-(4-fluorophenyl)-2,2,6,6-tetramethyloxan-4-yl]ethyl}(thiophen-2-ylmethyl)amine	376.2	δ 7.29 (m, 3H), 7.01 (s, 4H), 3.98 (s, 2H), 2.50 (m, 2H), 2.30 (d, J = 14.2, 2H), 1.94 (m, 2H), 1.69 (d, J = 14.4, 2H), 1.28 (s, 6H), 1.03 (s, 6H).
117	benzyl({2-[9-(2-methoxyphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	380.3	δ 8.86 (d, J = 149.6, 2H), 7.25 - 7.19 (m, 3H), 7.18 - 7.12 (m, 1H), 7.09 (dd, J = 7.4, 2.0, 2H), 6.96 (dd, J = 7.8, 1.5, 1H), 6.85 - 6.75 (m, 2H), 3.74 - 3.63 (m, 7H), 2.55 (dd, J = 15.6, 7.9, 3H), 2.11 (d, J = 14.8, 2H), 1.75 - 1.46 (m, 5H), 1.46 - 1.32 (m, 3H), 1.32 - 1.22 (m, 1H), 1.17 (d, J = 4.1, 1H), 0.74 - 0.60 (m, 1H).
118	benzyl({2-[9-(6-methoxypyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	381.3	δ 9.43 (s, 1H), 9.20 (s, 1H), 7.52 (m, 2H), 7.30 (dd, J = 5.1, 1.8, 3H), 7.21 (m, 2H), 6.78 (d, J = 7.3, 1H), 6.57 (d, J = 8.1, 1H), 3.83 (s, 3H), 3.77 (s, 2H), 3.71 (dd, J = 7.8, 2.7, 2H), 2.77 (s, 1H), 2.32 (d, J = 13.6, 2H), 2.25 (d, J = 11.5, 1H), 2.06 (td, J = 11.9, 4.8, 1H), 1.76 (m, 3H), 1.59 (m, 3H), 1.47 (m, 3H), 1.38 (m, 1H), 1.15 (m, 1H), 0.70 (m, 1H).
119	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}[(2-methoxyphenyl)methyl]methylamine	382.3	δ 10.17 (m, 3H), 7.41 (tdd, J = 8.3, 4.8, 1.6, 1H), 7.13 (m, 5H), 6.93 (m, 2H), 4.20 (dd, J = 14.9, 5.8, 1H), 3.98 (ddd, J = 32.2, 12.9, 4.8, 1H), 3.80 (dd, J = 7.4, 2.6, 5H), 2.94 (d, J = 114.3, 1H), 2.35 (m, 9H), 2.05 (ddd, J = 17.1, 12.7, 6.5, 1H), 1.89 (dt, J = 12.8, 6.2, 1H), 1.67 (ddd, J = 22.2, 14.2, 5.0, 2H), 1.23 (d, J = 10.7, 3H), 0.69 (t, J = 9.5, 3H).
120	{2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(3-methylphenyl)methyl]amine	382.3	δ 8.90 (d, J = 138.8, 2H), 7.15 (tt, J = 13.7, 7.6, 4H), 6.97 (m, 4H), 3.70 (m, 4H), 2.67 (s, 1H), 2.27 (s, 4H), 2.00 (m, 3H), 1.82 (m, 3H), 1.63 (m, 2H), 1.46 (m, 4H), 1.24 (d, J = 9.6, 1H), 0.78 (dt, J = 13.6, 8.8, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
121	{2-[(9S)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[(3-methylphenyl)methyl] amine	382.3	δ 8.73 (d, J = 138.2, 2H), 7.16 (m, 4H), 7.00 (dd, J = 10.5, 6.7, 2H), 6.94 (m, 2H), 3.72 (m, 4H), 2.69 (m, 1H), 2.27 (s, 4H), 2.05 (m, 2H), 1.94 (td, J = 12.6, 4.7, 1H), 1.83 (m, 3H), 1.63 (ddd, J = 14.1, 9.6, 4.6, 2H), 1.47 (m, 4H), 1.23 (m, 1H), 0.78 (dt, J = 13.9, 8.9, 1H)
122	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[(3-methylphenyl)methyl] amine	382.3	δ 8.96 (d, J = 123.7, 2H), 7.15 (m, 4H), 6.98 (m, 4H), 3.71 (m, 4H), 2.66 (s, 1H), 2.25 (d, J = 14.0, 4H), 2.05 (m, 2H), 1.94 (td, J = 12.7, 4.6, 1H), 1.81 (m, 3H), 1.63 (ddd, J = 14.2, 7.7, 3.4, 2H), 1.47 (m, 4H), 1.23 (m, 1H), 0.77 (dt, J = 13.7, 8.9, 1H)
123	benzyl({2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl})amine	382.3	δ 7.23 (m, 3H), 7.10 (dd, J = 4.6, 2.6, 4H), 6.92 (s, 2H), 3.64 (s, 2H), 2.63 (m, 1H), 2.07 (t, J = 13.9, 3H), 1.74 (s, 2H), 1.48 (d, J = 8.3, 3H), 1.40 (d, J = 14.0, 2H), 1.29 (m, 3H), 1.06 (m, 4H), 0.57 (m, 1H).
124	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[(1R)-1-phenylethyl]amine	382.3	δ 7.47 - 7.32 (m, 3H), 7.31 - 7.22 (m, 2H), 7.11 (dd, J = 8.9, 5.2, 2H), 6.98 (t, J = 8.6, 2H), 6.28 (s, 2H), 4.03 (s, 1H), 3.79 - 3.58 (m, 2H), 2.51 (s, 1H), 2.19 (d, J =
			14.5, 1H), 2.07 - 1.90 (m, 3H), 1.89 - 1.71 (m, 3H), 1.72 - 1.32 (m, 9H), 1.32 - 1.10 (m, 1H), 0.78 (dt, J = 13.6, 8.8, 1H).
125	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[(1S)-1-phenylethyl]amine	382.3	δ 7.47 - 7.32 (m, 3H), 7.31 - 7.22 (m, 2H), 7.11 (dd, J = 8.9, 5.2, 2H), 6.98 (t, J = 8.6, 2H), 6.28 (s, 2H), 4.03 (s, 1H), 3.79 - 3.58 (m, 2H), 2.51 (s, 1H), 2.19 (d, J = 14.5, 1H), 2.07 - 1.90 (m, 3H), 1.89 - 1.71 (m, 3H), 1.72 - 1.32 (m, 9H), 1.32 - 1.10 (m, 1H), 0.78 (dt, J = 13.6, 8.8, 1H).
126	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{[(2-nitrophenyl)methyl]amine	383.3	δ 7.94 (dd, J = 8.1, 1.2, 1H), 7.53 (td, J = 7.6, 1.3, 1H), 7.40 (m, 2H), 7.15 (m, 4H), 3.80 (m, 4H), 2.48 (td, J = 10.9, 5.4, 1H), 2.32 (m, 4H), 2.18 (ddd, J = 12.7, 7.8, 3.7, 2H), 1.84 (ddd, J = 13.2, 10.4, 5.1, 1H), 1.63 (m, 4H), 1.21 (s, 3H), 0.69 (s, 3H).
127	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{[(3-nitrophenyl)methyl]amine	383.3	δ 9.09 (d, J = 219.1, 2H), 8.12 (dd, J = 8.2, 1.6, 1H), 8.01 (s, 1H), 7.45 (dt, J = 15.6, 7.7, 2H), 7.03 (q, J = 8.5, 4H), 3.87 (s, 2H), 3.69 (m, 2H), 3.42 (s, 1H), 3.22 (s, 2H), 2.73 (d, J = 4.5, 1H), 2.24 (d, J = 8.2, 4H), 2.12 (m, 2H), 1.85 (m, 1H), 1.69 (dd, J = 12.1, 4.5, 1H), 1.52 (m, 2H), 1.11 (s, 3H), 0.57 (s, 3H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
128	2-[(2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl] phenol	384.2	δ 8.36 (d, J = 129.4, 2H), 7.20 (dd, J = 11.0, 4.6, 1H), 7.14 (dd, J = 8.9, 5.1, 2H), 7.00 (t, J = 8.6, 2H), 6.92 (m, 2H), 6.79 (t, J = 7.1, 1H), 3.88 (s, 2H), 3.68 (m, 2H), 2.67 (m, 1H), 2.29 (m, 1H), 1.98 (m, 3H), 1.79 (m, 3H), 1.51 (m, 6H), 1.20 (s, 1H), 0.74 (dt, J = 13.8, 8.9, 1H)
129	{2-[4-(4-methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl}[(2-methoxyphenyl)methyl]amine	384.3	δ 8.47 (d, J = 196.5, 2H), 7.36 (td, J = 8.3, 1.7, 1H), 7.12 (dd, J = 9.5, 2.6, 2H), 7.08 (dd, J = 7.5, 1.6, 1H), 6.91 (td, J = 7.5, 0.8, 1H), 6.86 (d, J = 8.8, 3H), 5.77 (s, 2H), 3.91 (s, 2H), 3.82 (s, 3H), 3.79 (s, 3H), 3.77 (m, 2H), 2.76 (s, 1H), 2.33 (s, 1H), 2.16 (m, 2H), 1.96 (d, J = 4.6, 1H), 1.77 (d, J = 4.7, 1H), 1.59 (m, 2H), 1.19 (s, 3H), 0.66 (s, 3H).
130	[(5-ethylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	385.1	δ 8.73 (d, J = 4.6, 1H), 8.20 (t, J = 7.7, 2H), 7.80 - 7.55 (m, 2H), 6.88 (d, J = 3.4, 1H), 6.64 (d, J = 3.4, 1H), 4.11 (s, 2H), 3.81 (dd, J = 8.4, 4.3, 1H), 3.70 (t, J = 10.0, 1H), 3.00 (d, J = 4.6, 1H), 2.86 - 2.70 (m, 2H), 2.53 (t, J = 10.1, 1H), 2.45 - 2.25 (m, 3H), 2.18 (t, J = 10.0, 1H), 2.00 (d, J = 14.2, 1H), 1.93 - 1.75 (m, 2H), 1.68 (dd, J = 9.5, 4.4, 1H), 1.62 - 1.38 (m, 4H), 1.26 (t, J = 7.5, 3H), 1.20 - 1.07 (m, 1H), 0.75 (dt, J = 12.9, 8.8, 1H).
131	[(3,5-dimethylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	385.1	δ 9.45 (brs, 1H), 8.70 (d, J = 5.0, 1H), 8.26 (t, J = 7.7, 1H), 7.75 (d, J = 8.1, 1H), 7.70 (m, 1H), 6.46 (d, J = 0.8, 1H), 4.07 (s, 2H), 3.76 (ddd, J = 44.9, 13.9, 7.2, 2H), 3.05 (m, 1H), 2.58 (m, 1H), 2.43 (t, J = 10.6, 1H), 2.36 (d, J = 0.7, 3H), 2.24 (dd, J = 31.9, 17.7, 3H),
			2.03 (m, 4H), 1.85 (m, 2H), 1.66 (dd, J = 13.8, 8.8, 1H), 1.48 (m, 4H), 1.15 (d, J = 7.9, 1H), 0.75 (dt, J = 13.1, 8.9, 1H).
132	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}[(6-methylpyridin-3-yl)methyl]amine	385.3	δ 8.84 (s, 1H), 8.24 (d, J = 8.2, 1H), 7.53 (d, J = 8.2, 1H), 7.17 (m, 3H), 6.96 (t, J = 8.6, 2H), 4.08 (d, J = 13.9, 2H), 3.63 (d, J = 10.5, 2H), 2.84 (dd, J = 12.0, 8.2, 1H), 2.68 (s, 3H), 2.24 (m, 2H), 2.07 (d, J = 14.1, 1H), 1.96 (m, 1H), 1.74 (dd, J = 12.5, 8.6, 1H), 1.57 (m, 1H), 1.48 (d, J = 14.2, 1H), 1.41 (m, 1H), 1.28 (dd, J = 14.0, 7.4, 1H), 0.96 (dd, J = 14.5, 7.4, 1H), 0.73 (td, J = 7.3, 3.9, 4H), 0.44 (t, J = 7.4, 3H).
133	{2-[4-(4-fluorophenyl)-2,2,6,6-tetramethyloxan-4-yl]ethyl}[(6-methylpyridin-3-yl)methyl]amine	385.3	δ 8.85 (s, 1H), 8.24 (d, J = 8.2, 1H), 7.54 (d, J = 8.3, 2H), 7.24 (dd, J = 8.9, 5.1, 1H), 6.92 (m, 2H), 4.12 (s, 2H), 2.61 (m, 5H), 2.25 (d, J = 14.3, 2H), 1.91 (dd, J = 10.4, 6.2, 2H), 1.65 (d, J = 14.4, 2H), 1.19 (d, J = 8.9, 6H), 0.94 (s, 6H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
134	[(4,5-dimethylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	385.3	δ 9.46 (s, 1H), 8.62 (d, J = 4.2, 1H), 8.07 (t, J = 7.3, 1H), 7.60 (d, J = 8.1, 1H), 7.52 (m, 1H), 6.76 (s, 1H), 4.06 (q, J = 13.9, 2H), 3.75 (m, 2H), 3.01 (m, 1H), 2.57 (s, 1H), 2.29 (m, 7H), 2.19 (m, 1H), 2.04 (s, 3H), 1.95 (d, J = 14.0, 1H), 1.81 (m, 2H), 1.67 (d, J = 8.2, 1H), 1.47 (m, 4H), 1.15 (m, 1H), 0.74 (dt, J = 13.1, 8.8, 1H).
135	[(2,4-dimethyl-1,3-thiazol-5-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	386.1	δ 9.59 (s, 1H), 8.68 (dd, J = 5.6, 1.4, 1H), 8.35 (td, J = 8.0, 1.6, 1H), 7.80 (dd, J = 12.0, 7.0, 2H), 4.22 (m, 2H), 3.83 (dt, J = 12.5, 4.4, 1H), 3.72 (m, 1H), 3.05 (dt, J = 11.2, 5.6, 1H), 2.73 (s, 3H), 2.57 (m, 2H), 2.31 (m, 6H), 2.04 (m, 1H), 1.88 (ddd, J = 19.2, 11.4, 6.9, 2H), 1.68 (m, 1H), 1.52 (m, 4H), 1.19 (dd, J = 12.2, 5.9, 1H), 0.76 (dt, J = 13.1, 8.9, 1H).
136	{2-[9-(pyrazin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	386.1	δ 8.90 (s, 1H), 8.55 (s, 1H), 8.40 (s, 1H), 6.60 (s, 1H), 3.88 (d, J = 12.3, 2H), 3.79 - 3.66 (m, 1H), 3.58 (dd, J = 16.8, 6.5, 1H), 2.81 (s, 1H), 2.40 (s, 1H), 2.35 - 2.22 (m, 2H), 2.16 (s, 3H), 2.12 - 2.00 (m, 1H), 1.97 - 1.88 (m, 4H), 1.85 (t, J = 9.1, 1H), 1.75 - 1.49 (m, 3H), 1.49 - 1.27 (m, 4H), 0.98 (d, J = 11.4, 1H), 0.55 (dt, J = 13.3, 9.0, 1H).
137	[(4,5-dimethylfuran-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	386.1	δ 9.14 (s, 1H), 8.85 (s, 1H), 7.24 (ddd, J = 11.5, 6.2, 3.3, 2H), 7.05 (s, 2H), 6.06 (s, 1H), 3.89 - 3.66 (m, 4H), 2.72 (s, 1H), 2.29 (s, 1H), 2.22 - 2.13 (m, 1H), 2.11 (s, 4H), 1.85 (s, 7H), 1.76 - 1.62 (m, 2H), 1.60 - 1.36 (m, 4H), 1.33 - 1.24 (m, 1H), 0.82 (dt, J = 13.6, 8.8, 1H).
138	{2-[9-(2-methoxyphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	386.2	δ 8.90 (d, J = 150.1, 2H), 7.19 (dd, J = 3.7, 1.4, 1H), 7.18 - 7.14 (m, 1H), 6.99 (dd, J = 7.8, 1.5, 1H), 6.94-6.76 (m, 4H), 4.66 (s, 2H), 3.94 (s, 2H), 3.80 - 3.63
			(m, 5H), 2.73 - 2.45 (m, 3H), 2.30 - 2.08 (m, 2H), 1.76 - 1.48 (m, 5H), 1.39 (dt, J = 7.0, 6.3, 3H), 1.30 (d, J = 5.2, 1H), 1.18 (d, J = 4.1, 1H), 0.68 (dd, J = 8.7, 5.0, 1H).
139	{2-[9-(2-methoxyphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-3-ylmethyl)amine	386.2	δ 9.28 (d, J = 95.5, 2H), 7.18 - 7.12 (m, 3H), 6.97 (dd, J = 7.8, 1.5, 1H), 6.93 - 6.86 (m, 1H), 6.86 - 6.71 (m, 2H), 3.80 - 3.61 (m, 7H), 2.55 (dd, J = 19.5, 5.1, 3H), 2.12 (d, J = 12.8, 2H), 1.85 (s, 2H), 1.76 - 1.47 (m, 5H), 1.46 - 1.32 (m, 3H), 1.31-1.22 (m, 1H), 1.17 (d, J = 4.2, 1H), 0.74 - 0.60 (m, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
140	[(3-methoxythiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387	δ 11.70 (brs, 1H), 9.14 (d, J = 66.6, 2H), 8.72 (d, J = 4.3, 1H), 8.19 (td, J = 8.0, 1.4, 1H), 7.70 (d, J = 8.1, 1H), 7.63 (dd, J = 7.0, 5.8, 1H), 7.22 (d, J = 5.5, 1H), 6.78 (d, J = 5.6, 1H), 4.08 (m, 2H), 3.80 (m, 4H), 3.69 (dd, J = 11.2, 8.7, 1H), 2.99 (d, J = 4.8, 1H), 2.51 (t, J = 9.9, 1H), 2.35 (m, 3H), 2.18 (td, J = 13.5, 5.4, 1H), 1.99 (d, J = 14.2, 1H), 1.82 (m, 2H), 1.65 (m, 1H), 1.47 (m, 4H), 1.14 (m, 1H), 0.73 (dt, J = 13.2, 8.9, 1H).
141	[(3-methoxythiophen-2-yl)methyl]({2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387	δ 9.03 (d, J = 80.0, 2H), 8.75 (d, J = 5.3, 1H), 8.31 (t, J = 7.9, 1H), 7.76 (m, 2H), 7.26 (t, J = 4.0, 1H), 6.81 (d, J = 5.6, 1H), 4.12 (s, 2H), 3.82 (s, 4H), 3.69 (dd, J = 24.9, 14.9, 1H), 3.04 (s, 1H), 2.56 (s, 1H), 2.45 (dd, J = 17.7, 7.6, 1H), 2.29 (ddd, J = 17.8, 13.5, 5.8, 3H), 2.05 (d, J = 14.3, 1H), 1.87 (dt, J = 14.4, 6.7, 2H), 1.67 (ddd, J = 27.6, 16.0, 6.9, 1H), 1.52 (m, 4H), 1.20 (m, 1H), 0.78 (dt, J = 13.0, 8.9, 1H).
142	{2-[9-(6-methoxypyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-2-ylmethyl)amine	387.2	δ 9.37 (s, 1H), 9.11 (s, 0H), 7.55 (dd, J = 8.2, 7.5, 1H), 7.30 (dd, J = 5.1, 1.1, 1H), 7.03 (d, J = 2.6, 1H), 6.96 (dd, J = 5.1, 3.6, 1H), 6.81 (d, J = 7.3, 1H), 6.60 (d, J = 8.0, 1H), 4.07 (s, 2H), 3.86 (s, 3H), 3.73 (dd, J = 7.7, 2.7, 2H), 2.87 (m, 1H), 2.75 (brs, 1H), 2.47 (m, 1H), 2.32 (dd, J = 24.5, 13.6, 2H), 2.09 (m, 1H), 1.80 (m, 3H), 1.63 (dt, J = 15.1, 7.4, 2H), 1.49 (m, 3H), 1.39 (d, J = 4.5, 1H), 1.16 (m, 1H), 0.72 (dt, J = 13.4, 8.8, 1H).
143	{2-[9-(6-methoxypyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(thiophen-3-ylmethyl)amine	387.2	δ 9.40 (s, 1H), 9.21 (s, 1H), 7.53 (m, 1H), 7.28 (d, J = 3.0, 2H), 6.99 (dd, J = 4.8, 1.4, 1H), 6.80 (d, J = 7.4, 1H), 6.59 (d, J = 8.2, 1H), 3.86 (d, J = 6.4, 5H), 3.72 (dd, J = 7.7, 2.7, 2H), 2.78 (m, 1H), 2.30 (dd, J = 28.1, 12.5, 3H), 2.09 (m, 1H), 2.02 (brs, 1H), 1.79 (m, 3H), 1.61 (m, 2H), 1.47 (m, 4H), 1.16 (m, 1H), 0.71 (dt, J = 13.4, 8.7, 1H).
144	{2-[4-(4-chlorophenyl)-2,2-dimethyloxan-4-yl]ethyl}[(2-methoxyphenyl)methyl]	388.2	δ 8.48 (d, J = 152.7, 2H), 7.28 (td, J = 8.3, 1.7, 1H), 7.22 (dd, J = 6.6, 4.8, 2H), 7.06 (m, 2H), 6.97 (dd, J = 7.5, 1.6, 1H), 6.81 (ddd, J = 19.8, 13.2, 4.6, 2H), 6.03 (s, 1H), 3.82 (s, 2H), 3.66 (m, 5H), 2.64 (s, 1H), 2.15
	]amine		(s, 1H), 2.05 (ddd, J = 22.5, 14.1, 2.1, 2H), 1.85 (m, 1H), 1.72 (dd, J = 12.5, 4.7, 1H), 1.53 (m, 2H), 1.11 (s, 3H), 0.57 (s, 3H).
145	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(5-methylthiophen-2-yl)methyl]amine	388.2	δ 7.28 (s, 4H), 7.25 - 7.15 (m, 2H), 7.04 (t, J = 8.6, 2H), 6.77 (d, J = 3.5, 1H), 6.59 (dd, J = 3.4, 1.1, 1H), 3.91 (s, 2H), 3.85 - 3.64 (m, 2H), 2.73 (t, J = 9.7, 1H), 2.41 (d, J = 0.7, 3H), 2.37 - 1.75 (m, 18H), 1.67 (dd, J = 11.7, 7.1, 2H), 1.59 - 1.34 (m, 4H), 1.26 (s, 1H), 0.81 (dt, J = 14.0, 8.9, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
146	{2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}(thiophen-3-ylmethyl)amine	388.2	δ 7.18 (s, 1H), 7.15 (s, 1H), 7.10 (dd, J = 8.9, 5.2, 2H), 6.92 (dd, J = 10.8, 6.4, 2H), 6.87 (m, 1H), 3.67 (d, J = 35.8, 3H), 2.66 (m, 1H), 2.07 (s, 3H), 1.83 (m, 2H), 1.56 (s, 3H), 1.41 (d, J = 13.9, 2H), 1.33 (m, 3H), 1.02 (m, 4H), 0.58 (m, 1H).
147	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(3-methylthiophen-2-yl)methyl]amine	388.2	δ 9.01 (d, J = 137.9, 2H), 7.15 - 7.02 (m, 3H), 6.94 (t, J = 8.6, 2H), 6.77 - 6.63 (m, 1H), 4.82 (s, 1H), 3.83 (d, J = 19.1, 2H), 3.73 - 3.54 (m, 2H), 2.64 (s, 1H), 2.18 (d, J = 10.4, 1H), 2.12 - 1.64 (m, 9H), 1.65 - 1.50 (m, 2H), 1.50 - 1.27 (m, 4H), 1.27 - 1.08 (m, 1H), 0.69 (dt, J = 13.5, 8.8, 1H).
148	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(4-methylthiophen-2-yl)methyl]amine	388.2	δ 9.31 (d, J = 89.1, 2H), 7.15 - 7.05 (m, 2H), 6.93 (t, J = 8.6, 2H), 6.80 - 6.65 (m, 2H), 3.80 (s, 2H), 3.73 - 3.57 (m, 2H), 2.93 (s, 1H), 2.60 (s, 1H), 2.17 (s, 1H), 2.04 (dd, J = 16.0, 3.3, 4H), 1.91 (ddd, J = 17.5, 16.7, 9.1, 2H), 1.84 - 1.65 (m, 3H), 1.57 (ddd, J = 13.2, 9.0, 4.4, 2H), 1.50 - 1.25 (m, 4H), 1.17 (dd, J = 14.9, 5.0, 1H), 0.70 (dt, J = 13.6, 8.8, 1H).
149	{2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}(thiophen-2-ylmethyl)amine	388.3	δ 7.28 (s, 3H), 7.22 (dd, J = 8.6, 4.9, 2H), 7.01 (m, 2H), 4.01 (s, 2H), 3.74 (s, 1H), 2.26 (m, 1H), 1.73 (m, 11H), 1.52 (d, J = 14.1, 2H), 1.39 (m, 2H), 1.13 (s, 2H), 0.69 (m, 1H).
150	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(4-methyl-1,3-thiazol-2-yl)methyl]amine	389	δ 7.22 (dd, J = 8.9, 5.2, 2H), 7.02 (dd, J = 14.0, 5.4, 2H), 6.92 (d, J = 0.9, 1H), 4.25 (q, J = 14.7, 2H), 3.73 (m, 2H), 2.89 (td, J = 11.8, 4.8, 1H), 2.50 (td, J = 11.7, 5.0, 1H), 2.38 (d, J = 0.8, 3H), 2.15 (m, 1H), 2.08 (m, 2H), 1.98 (m, 1H), 1.91 (d, J = 13.9, 1H), 1.79 (d, J = 9.3, 1H), 1.69 (m, 2H), 1.48 (m, 5H), 1.25 (m, 1H), 0.81 (dt, J = 13.3, 8.7, 1H).
151	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}[(5-methylthiophen-2-yl)methyl]amine	390.2	δ 7.15 - 7.02 (m, 2H), 6.94 (t, J = 8.6, 2H), 6.67 (d, J = 3.5, 1H), 6.49 (s, 1H), 3.78 (s, 2H), 3.62 (dd, J = 10.4, 8.1, 3H), 2.61 (s, 1H), 2.30 (s, 4H), 2.08 (dd, J = 31.6, 14.0, 4H), 1.88 (d, J = 4.6, 1H), 1.79 - 1.34 (m, 19H), 1.29 (dd, J = 14.0, 7.4, 2H), 0.96 (dd, J = 14.5, 7.3, 1H), 0.74 (t, J = 7.5, 5H), 0.44 (t, J = 7.4, 4H).
152	[(5-chlorothiophen-2-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	391	δ 8.75 (d, J = 4.8, 1H), 8.24 (t, J = 7.7, 1H), 7.86 - 7.58 (m, 2H), 6.44 (d, J = 3.3, 1H), 6.28 (d, J = 3.3, 1H), 4.07 (s, 2H), 3.94 - 3.79 (m, 1H), 3.72 (t, J = 10.1, 1H), 3.01 (dd, J = 11.1, 6.0, 1H), 2.56 (t, J = 9.9, 1H), 2.49 - 2.11 (m, 4H), 2.05 (d, J = 14.1, 1H), 1.88 (ddd, J = 18.8, 11.0, 6.5, 2H), 1.78 - 1.31 (m, 5H), 1.31 - 1.07 (m, 1H), 0.77 (dt, J = 13.1, 8.9, 1H)

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
153	dibutyl({2-[4-(4-fluorophenyl)-2,2,6,6-tetramethyloxan-4-yl]ethyl})amine	392.4	δ 7.37 (m, 2H), 7.07 (m, 2H), 2.83 (dd, J = 16.3, 9.4, 4H), 2.68 (m, 2H), 2.38 (d, J = 14.3, 2H), 2.09 (s, 4H), 1.93 (m, 2H), 1.77 (d, J = 14.3, 2H), 1.33 (m, 10H), 1.05 (d, J = 8.6, 6H), 0.91 (t, J = 7.2, 6H).
154	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(2-phenylpropan-2-yl)amine	396.3	δ 7.37 (s, 5H), 7.28 (s, 0H), 7.17 - 6.99 (m, 3H), 6.93 (t, J = 8.6, 2H), 3.81 - 3.57 (m, 2H), 2.45 (d, J = 9.0, 1H), 2.04 - 1.72 (m, 7H), 1.66 (t, J = 10.7, 6H), 1.62 - 1.53 (m, 2H), 1.52 - 1.34 (m, 4H), 1.23 (s, 1H), 0.78 (d, J = 13.8, 1H).
155	{4H,5H,6H-cyclopenta[b]thiophen-2-ylmethyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	397.1	δ 9.57 (brs, 1H), 8.62 (d, J = 3.9, 1H), 8.02 (t, J = 7.1, 1H), 7.57 (d, J = 8.1, 1H), 7.48 (dd, J = 6.9, 5.5, 1H), 6.80 (s, 1H), 5.30 (brs, 1H), 4.06 (q, J = 14.1, 2H), 3.74 (m, 2H), 2.99 (m, 1H), 2.82 (t, J = 7.2, 2H), 2.65 (t, J = 7.2, 2H), 2.57 (m, 1H), 2.34 (ddd, J = 33.3, 21.0, 10.4, 5H), 2.16 (dd, J = 9.9, 5.6, 1H), 1.94 (d, J = 13.9, 1H), 1.78 (m, 2H), 1.66 (d, J = 8.0, 1H), 1.46 (ddd, J = 16.6, 12.7, 5.7, 4H), 1.14 (m, 1H), 0.72 (dt, J = 13.4, 9.0, 1H).
156	{2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}[(6-methylpyridin-3-yl)methyl]amine	397.3	δ 8.22 (d, J = 8.0, 1H), 7.49 (t, J = 16.4, 1H), 7.17 (m, 8H), 6.96 (t, J = 8.6, 2H), 4.09 (s, 2H), 3.66 (s, 4H), 2.84 (s, 1H), 2.68 (s, 3H), 2.29 (s, 1H), 2.20 (d, J = 13.2, 1H), 2.10 (d, J = 14.1, 1H), 1.93 (s, 1H), 1.73 (s, 1H), 1.59 (m, 1H), 1.45 (d, J = 14.0, 3H), 1.30 (m, 2H), 1.10 (m, 3H), 0.62 (d, J = 11.1, 1H).
157	[(2,3-dimethoxyphenyl)methyl]({2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl})amine	398.3	δ 7.05 (dd, J = 19.6, 8.3, 4H), 6.88 (m, 1H), 6.74 (dd, J = 8.2, 1.4, 1H), 6.62 (dd, J = 7.6, 1.4, 1H), 3.77 (s, 3H), 3.68 (m, 5H), 3.55 (d, J = 2.3, 2H), 2.37 (m, 1H), 2.22 (m, 4H), 2.06 (ddd, J = 13.8, 8.6, 4.1, 2H), 1.73 (dd, J = 6.6, 4.3, 1H), 1.56 (m, 4H), 1.10 (s, 3H), 0.58 (s, 3H).
158	[(2,4-dimethoxyphenyl)methyl]({2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl})amine	398.3	δ 8.09 (s, 1H), 7.68 (d, J = 33.5, 1H), 7.55 (s, 1H), 7.02 (q, J = 8.4, 4H), 6.86 (m, 1H), 6.32 (dd, J = 6.6, 2.2, 2H), 3.77 (d, J = 10.4, 2H), 3.69 (m, 8H), 2.67 (s, 1H), 2.24 (s, 4H), 2.10 (m, 2H), 1.87 (d, J = 4.5, 1H), 1.67 (d, J = 4.4, 1H), 1.51 (m, 2H), 1.10 (s, 3H), 0.57 (s, 3H).
159	{2-[9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(4-methoxyphenyl)methyl]amine	398.3	δ 9.06 (d, J = 131.9, 2H), 7.17 (m, 2H), 7.08 (d, J = 8.7, 2H), 7.00 (t, J = 8.6, 2H), 6.79 (d, J = 8.7, 2H), 3.69 (m, 7H), 2.62 (s, 1H), 2.20 (s, 1H), 1.99 (m, 3H), 1.81 (m, 3H), 1.62 (m, 2H), 1.46 (m, 4H), 1.24 (d, J = 9.5, 1H), 0.77 (dt, J = 13.4, 8.8, 1H)

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
160	[(5-propylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	399.1	δ 9.43 (s, 2H), 8.72 (d, J = 4.6, 1H), 8.21 (t, J = 7.3, 1H), 7.72 (d, J = 8.1, 2H), 6.88 (d, J = 3.5, 1H), 6.63 (d, J = 3.5, 1H), 4.11 (s, 2H), 3.87 - 3.65 (m, 2H), 3.00 (s, 1H), 2.71 (t, J = 7.5, 2H), 2.54 (s, 1H), 2.32 (s, 3H), 2.27 - 2.11 (m, 1H), 2.02 (s, 1H), 1.84 (dd, J = 16.6, 7.3, 2H), 1.64 (dd, J = 15.0, 7.4, 7H), 1.22 - 1.10 (m, 1H), 0.95 (t, J = 7.3, 3H), 0.83 - 0.72 (m, 1H).
161	1-{5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]thiophen-2-yl}ethan-1-ol	401.1	δ 9.38 (s, 2H), 8.76 (d, J = 4.6, 1H), 8.29 (t, J = 7.9, 1H), 7.84-7.69 (m, 2H), 6.92-6.74 (m, 4H), 5.02 (d, J = 6.4, 1H), 4.13 (s, 2H), 3.87 - 3.60 (m, 2H), 3.03 (s, 1H), 2.52 (s, 1H), 2.34 (t, J = 15.7, 3H), 2.20 (t, J = 12.6, 1H), 2.03 (dd, J = 14.2, 4.7, 1H), 1.96 - 1.78 (m, 2H), 1.81 - 1.65 (m, 1H), 1.65 - 1.43 (m, 7H), 1.15 (s, 1H), 0.77 (s, 1H).
162	6-[9-(2-[(4,5-dimethylthiophen-2-yl)methyl]amino)ethyl]-6-oxaspiro[4.5]decan-9-yl]pyridin-3-ol	401.1	¹ H NMR (400 MHz, CD ₃ CN) δ 8.18 (dd, J = 2.3, 1.2, 1H), 7.72 (s, 1H), 7.32 (d, J = 2.3, 2H), 6.82 (s, 1H), 4.10 (s, 2H), 3.67 (m, 2H), 2.95 (m, 1H), 2.50 (m, 1H), 2.32 (s, 3H), 2.27 (d, J = 13.9, 2H), 2.09 (m, 4H), 2.03 (m, 1H), 1.88 (m, 1H), 1.83 (t, J = 9.2, 2H), 1.71 (m, 1H), 1.63 (m, 2H), 1.48 (ddd, J = 16.6, 12.3, 7.6, 4H), 1.13 (dd, J = 11.7, 5.4, 1H), 0.72 (dt, J = 13.7, 9.0, 1H).
163	6-[9-(2-[(4,5-dimethylthiophen-2-yl)methyl]amino)ethyl]-6-oxaspiro[4.5]decan-9-yl]pyridin-2-ol	401.1	δ 7.49 (m, 1H), 6.76 (s, 1H), 6.51 (d, J = 8.9, 1H), 6.25 (d, J = 7.0, 1H), 3.99 (s, 2H), 3.71 (m, 2H), 2.83 (dd, J = 16.5, 11.3, 1H), 2.61 (dd, J = 17.0, 5.8, 1H), 2.27 (d, J = 21.1, 5H), 1.99 (m, 6H), 1.65 (m, 10H), 0.98 (dd, J = 18.1, 5.5, 1H).
164	2-[9-(2-[(4,5-dimethylthiophen-2-yl)methyl]amino)ethyl]-6-oxaspiro[4.5]decan-9-yl]pyridin-4-ol	401.2	δ 9.21 (d, J = 64.7, 2H), 8.00 (s, 1H), 7.07 (m, 2H), 6.67 (s, 1H), 3.95 (s, 2H), 3.62 (m, 2H), 2.84 (s, 1H), 2.44 (s, 1H), 2.27 (d, J = 12.2, 1H), 2.16 (s, 4H), 2.03 (d, J = 13.5, 2H), 1.94 (s, 3H), 1.83 (d, J = 13.9, 1H), 1.65 (m, 3H), 1.37 (m, 5H), 0.75 (s, 1H).
165	[(5-nitrothiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	402	δ 8.59 (d, J = 4.0, 1H), 8.15 (t, J = 7.0, 1H), 7.79 (d, J = 4.1, 1H), 7.66 (d, J = 8.2, 1H), 7.60 (m, 1H), 7.16 (d, J = 4.2, 1H), 4.23 (s, 2H), 3.78 (m, 2H), 3.04 (d, J = 6.0, 1H), 2.65 (m, 1H), 2.43 (d, J = 9.8, 1H), 2.29 (m, 3H), 1.98 (d, J = 14.1, 1H), 1.83 (d, J = 5.4, 2H), 1.67 (m, 1H), 1.48 (m, 4H), 1.16 (m, 1H), 0.75 (d, J = 13.2, 1H).
166	[(3,5-dimethylthiophen-2-yl)methyl]({2-[(9R)-9-	402.1	δ 7.19 (dd, J = 8.9, 5.1, 2H), 7.01 (dd, J = 13.7, 5.0, 2H), 6.43 (s, 1H), 3.87 (m, 2H), 3.72 (m, 2H), 3.02 (s, 1H), 2.72 (dd, J = 14.6, 8.9, 1H), 2.31 (dd, J = 31.1,

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
	(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl))amine		10.4, 4H), 2.15 (d, J = 13.8, 1H), 2.05 (d, J = 14.0, 1H), 1.98 (m, 4H), 1.87 (m, 2H), 1.77 (d, J = 9.7, 1H), 1.67 (ddd, J = 15.6, 10.3, 5.4, 2H), 1.46 (m, 4H), 1.25 (t, J = 7.1, 1H), 0.79 (dt, J = 13.7, 8.9, 1H).
167	[(5-ethylthiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl))amine	402.1	δ 7.21 (dd, J = 8.9, 5.2, 2H), 7.04 (t, J = 8.6, 2H), 6.79 (d, J = 3.5, 1H), 6.62 (d, J = 3.5, 1H), 3.92 (s, 2H), 3.80 - 3.67 (m, 3H), 2.82 - 2.67 (m, 2H), 2.32 (s, 1H), 2.16 (d, J = 14.3, 1H), 2.06 (s, 1H), 2.00 (td, J = 12.8, 4.9, 1H), 1.91 (d, J = 13.9, 2H), 1.84 - 1.75 (m, 1H), 1.69 (s, 2H), 1.50 (d, J = 3.7, 4H), 1.25 (t, J = 7.5, 4H), 0.81 (dt, J = 13.4, 8.7, 1H).
168	{2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}[(5-methylthiophen-2-yl)methyl]amine	402.3	δ 7.12 (m, 2H), 6.93 (s, 2H), 6.66 (d, J = 3.4, 1H), 6.49 (d, J = 2.5, 1H), 3.80 (s, 2H), 3.63 (s, 2H), 2.65 (m, 1H), 2.31 (s, 3H), 2.12 (m, 2H), 1.85 (m, 1H), 1.61 (s, 3H), 1.43 (d, J = 14.0, 2H), 1.33 (m, 3H), 1.03 (s, 4H), 0.59 (m, 1H).
169	[(4,5-dimethylthiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl))amine	402.3	δ 8.92 (d, J = 108.6, 2H), 7.15 - 7.05 (m, 2H), 6.93 (t, J = 8.6, 2H), 6.51 (s, 1H), 5.31 (s, 1H), 3.75 (s, 2H), 3.69 - 3.54 (m, 2H), 2.63 (s, 1H), 2.27 - 2.10 (m, 4H), 2.06 (d, J = 14.0, 1H), 1.98 (d, J = 13.9, 1H), 1.93-1.84 (m, 4H), 1.84 - 1.65 (m, 3H), 1.58 (ddd, J = 17.0, 8.4, 3.8, 2H), 1.51 - 1.27 (m, 4H), 1.17 (dd, J = 13.9, 6.2, 1H), 0.71 (dt, J = 13.6, 8.8, 1H).
170	{[5-(methylsulfanyl)thiophen-2-yl]methyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl))amine	403	δ 9.54 (s, 1H), 8.71 (d, J = 4.5, 1H), 8.26 (t, J = 7.2, 2H), 7.80 - 7.67 (m, 2H), 6.92 (dd, J = 21.5, 3.6, 2H), 4.15 (s, 2H), 3.76 (d, J = 40.3, 2H), 3.02 (td, J = 11.4, 5.3, 1H), 2.62 - 2.51 (m, 1H), 2.48 (s, 3H), 2.42 (s, 1H), 2.31 (t, J = 13.3, 3H), 2.03 (d, J = 14.2, 1H), 1.92 - 1.78 (m, 2H), 1.78 - 1.63 (m, 1H), 1.64 - 1.36 (m, 4H), 1.25 - 1.12 (m, 1H), 0.79 (s, 1H).
171	6-[9-(2-[(3-methoxythiophen-2-yl)methyl]amino)ethyl]-6-oxaspiro[4.5]decan-9-yl]pyridin-3-ol	403	¹ H NMR (400 MHz, CD ₃ CN) δ 8.15 (d, J = 1.5, 1H), 7.60 (s, 1H), 7.43 (d, J = 5.6, 1H), 7.31 (m, 2H), 6.97 (d, J = 5.6, 1H), 4.11 (s, 2H), 3.86 (s, 3H), 3.66 (dd, J = 7.8, 2.9, 2H), 2.97 (m, 1H), 2.51 (m, 1H), 2.28 (m, 2H), 2.02 (m, 1H), 1.87 (m, 2H), 1.80 (d, J = 13.5, 2H), 1.70 (d, J = 9.8, 1H), 1.61 (dd, J = 13.8, 7.1, 2H), 1.49 (m, 4H), 1.12 (m, 1H), 0.71 (d, J = 13.5, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
172	6-[9-(2-[[[3-methoxythiophen-2-yl)methyl]amino]ethyl)-6-oxaspiro[4.5]decan-9-yl]pyridin-2-ol	403	δ 9.47 (brs, 1H), 7.51 (dd, J = 9.0, 7.2, 1H), 7.23 (d, J = 5.6, 1H), 6.80 (d, J = 5.5, 1H), 6.52 (d, J = 8.9, 1H), 6.27 (d, J = 7.1, 1H), 4.10 (s, 2H), 3.82 (s, 3H), 3.73 (dd, J = 6.8, 3.4, 2H), 2.83 (dd, J = 11.9, 5.7, 1H), 2.60 (t, J = 10.0, 1H), 2.27 (t, J = 15.0, 2H), 2.00 (t, J = 12.3, 2H), 1.65 (m, 10H), 0.97 (d, J = 13.4, 1H).
173	2-[(9R)-9-(2-[[[3-methoxythiophen-2-yl)methyl]amino]ethyl)]	403.2	δ 9.84 (s, 1H), 8.76 (s, 1H), 8.32 (d, J = 5.3, 1H), 7.60 (t, J = 7.7, 1H), 7.52 (m, 1H), 7.41 (m, 1H), 7.25 (d, J = 5.5, 1H), 6.81 (d, J = 5.5, 1H), 4.16 (m, 2H), 3.82
	-6-oxaspiro[4.5]decan-9-yl]-1-oxidopyridin-1-ium		(m, 4H), 3.71 (m, 1H), 3.05 (d, J = 13.4, 2H), 2.85 (d, J = 9.1, 1H), 2.53 (s, 1H), 2.27 (d, J = 14.3, 1H), 2.14 (m, 1H), 1.99 (t, J = 11.3, 1H), 1.85 (m, 2H), 1.66 (ddd, J = 18.0, 10.0, 5.8, 1H), 1.51 (m, 4H), 1.22 (dd, J = 12.3, 6.0, 1H), 0.90 (dt, J = 13.0, 8.7, 1H).
174	2-[9-(2-[[[3-methoxythiophen-2-yl)methyl]amino]ethyl)-6-oxaspiro[4.5]decan-9-yl]pyridin-4-ol	403.2	δ 9.17 (d, J = 50.2, 2H), 8.00 (d, J = 6.5, 1H), 7.16 (d, J = 5.6, 3H), 6.72 (d, J = 5.6, 1H), 4.00 (s, 2H), 3.73 (s, 5H), 2.82 (s, 1H), 2.34 (d, J = 39.9, 2H), 2.11 (dd, J = 51.0, 13.1, 3H), 1.84 (d, J = 13.9, 1H), 1.43 (m, 9H), 0.75 (s, 1H).
175	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[[3-methoxythiophen-2-yl)methyl]amine	404	δ 7.24 (s, 3H), 7.03 (dd, J = 11.7, 5.6, 2H), 6.80 (d, J = 5.5, 1H), 4.00 (s, 2H), 3.81 (m, 5H), 2.78 (m, 1H), 2.39 (m, 1H), 2.17 (m, 1H), 2.06 (s, 2H), 1.86 (m, 2H), 1.66 (m, 3H), 1.51 (m, 3H), 1.26 (m, 2H), 0.80 (m, 1H).
176	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{[3-(trifluoromethyl)phenyl]methyl}amine	406.3	δ 9.43 (d, J = 141.7, 2H), 7.47 (d, J = 7.2, 1H), 7.39 (s, 1H), 7.31 (m, 2H), 6.99 (q, J = 8.3, 4H), 3.67 (m, 4H), 2.54 (d, J = 8.4, 1H), 2.20 (d, J = 7.1, 3H), 2.06 (m, 3H), 1.92 (s, 2H), 1.60 (td, J = 12.5, 4.7, 1H), 1.46 (m, 2H), 1.08 (s, 3H), 0.55 (s, 3H).
177	(1-benzothiophen-2-ylmethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	407.1	δ 8.51 (dd, J = 5.5, 1.3, 1H), 8.02 (d, J = 1.4, 1H), 7.73 - 7.52 (m, 3H), 7.43 (d, J = 1.0, 1H), 7.35 - 7.23 (m, 3H), 3.67 (s, 3H), 2.96 (td, J = 11.5, 5.7, 1H), 2.56-2.43 (m, 1H), 2.43 - 2.28 (m, 1H), 2.16 (d, J = 13.6, 3H), 1.89 (d, J = 14.2, 1H), 1.73 (ddd, J = 19.7, 11.9, 7.2, 2H), 1.55 (dt, J = 15.0, 5.7, 1H), 1.48 - 1.22 (m, 4H), 1.06 (s, 1H), 0.66 (dt, J = 13.2, 8.9, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
178	(1-benzothiophen-3-ylmethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	407.1	δ 11.71 (s, 2H), 9.34 (d, J = 85.8, 1H), 8.48 (d, J = 5.0, 1H), 8.10 (s, 1H), 7.71 (dd, J = 6.2, 2.8, 1H), 7.58 (ddd, J = 22.1, 9.6, 4.3, 3H), 7.47 (s, 1H), 7.36 - 7.24 (m, 2H), 4.12 (s, 2H), 3.64 (s, 2H), 2.93 (s, 1H), 2.51 - 2.23 (m, 2H), 2.13 (t, J = 14.3, 3H), 1.94 - 1.83 (m, 1H), 1.80 - 1.64 (m, 2H), 1.62 - 1.49 (m, 1H), 1.37 (dd, J = 39.4, 7.2, 4H), 1.06 (d, J = 13.0, 1H), 0.64 (dt, J = 13.1, 9.0, 1H).
179	[(5-chlorothiophen-2-yl)methyl]{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	408.2	δ 7.11 (dd, J = 8.9, 5.2, 2H), 6.94 (dd, J = 15.9, 7.2, 2H), 6.75 - 6.56 (m, 2H), 3.79 (s, 2H), 3.71 - 3.52 (m, 2H), 2.61 (s, 1H), 2.18 (s, 1H), 1.84 (dddd, J = 31.4, 25.9, 23.7, 13.1, 12H), 1.58 (td, J = 9.4, 4.6, 2H), 1.39 (ddd, J = 23.7, 14.8, 9.2, 5H), 1.17 (s, 2H), 0.69 (dd, J = 8.7, 5.1, 1H).
180	2-[(2-{2,2-dimethyl-4-[4-(trifluoromethyl)phenyl]oxan-4-yl}ethyl)amino]methyl} phenol	408.3	δ 8.34 (d, J = 45.4, 2H), 7.50 (d, J = 8.3, 2H), 7.24 (d, J = 8.2, 2H), 7.10 (s, 1H), 6.77 (m, 3H), 3.80 (s, 2H), 3.66 (d, J = 12.3, 2H), 3.31 (s, 3H), 2.63 (s, 1H), 2.09 (dd, J = 26.1, 13.9, 3H), 1.87 (t, J = 10.4, 1H), 1.71 (t, J = 10.4, 1H), 1.58 (d, J = 14.0, 2H), 1.10 (s, 3H), 0.53 (s, 3H).
181	[(5-chlorothiophen-2-yl)methyl]{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}amine	410.1	δ 7.12 (dd, J = 8.9, 5.2, 2H), 6.96 (t, J = 8.6, 2H), 6.69 (q, J = 3.8, 2H), 3.79 (s, 2H), 3.63 (dd, J = 12.2, 7.1, 2H), 2.63 (dd, J = 12.2, 7.5, 1H), 2.29 - 1.77 (m, 8H), 1.67 (td, J = 12.5, 4.7, 1H), 1.44 (dd, J = 24.5, 10.8, 3H), 1.31 (d, J = 7.5, 1H), 0.95 (s, 1H), 0.74 (t, J = 7.5, 4H), 0.44 (t, J = 7.4, 3H).
182	{[5-(2-methylpropyl)thiophen-2-yl]methyl}{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	413.1	δ 9.56 (brs, 1H), 8.66 (d, J = 4.7, 1H), 8.09 (t, J = 7.5, 1H), 7.62 (d, J = 8.1, 1H), 7.55 (m, 1H), 6.87 (d, J = 3.4, 1H), 6.59 (d, J = 3.4, 1H), 4.08 (m, 2H), 3.75 (m, 2H), 2.96 (d, J = 4.8, 1H), 2.57 (d, J = 7.0, 2H), 2.50 (t, J = 9.6, 1H), 2.31 (m, 3H), 2.14 (td, J = 13.5, 5.4, 1H), 1.96 (d, J = 14.1, 1H), 1.80 (m, 3H), 1.66 (m, 1H), 1.47 (m, 4H), 1.14 (d, J = 13.0, 1H), 0.89 (d, J = 6.6, 6H), 0.73 (dt, J = 13.6, 9.0, 1H).
183	[(5-butylthiophen-2-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	413.1	δ 10.87 (brs, 1H), 9.42 (brs, 1H), 8.70 (d, J = 4.8, 1H), 8.17 (t, J = 7.7, 1H), 7.68 (d, J = 8.1, 1H), 7.62 (m, 1H), 6.85 (d, J = 3.5, 1H), 6.60 (d, J = 3.4, 1H), 4.08 (s, 2H), 3.79 (m, 1H), 3.67 (t, J = 10.0, 1H), 2.97 (d, J = 4.3, 1H), 2.70 (t, J = 7.6, 2H), 2.50 (t, J = 9.9, 1H), 2.33 (m, 3H), 2.16 (td, J = 13.1, 5.0, 1H), 1.98 (t, J = 9.4, 1H), 1.80 (t, J = 9.6, 2H), 1.54 (m, 7H), 1.33 (dq, J = 14.5, 7.3, 2H), 1.14 (m, 1H), 0.90 (t, J = 7.3, 3H), 0.73 (dt, J = 13.0, 8.9, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
184	{4H,5H,6H-cyclopenta[b]thiophen-2-ylmethyl}{[2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl]amine	414	δ 7.20 (m, 2H), 7.01 (dd, J = 13.5, 4.7, 2H), 6.65 (s, 1H), 3.88 (s, 2H), 3.72 (m, 3H), 2.78 (t, J = 7.2, 3H), 2.61 (t, J = 7.2, 2H), 2.34 (dt, J = 14.5, 7.3, 3H), 2.15 (d, J = 14.1, 1H), 2.06 (d, J = 13.9, 1H), 1.99 (m, 1H), 1.89 (m, 2H), 1.78 (m, 1H), 1.67 (ddd, J = 18.6, 11.9, 7.0, 2H), 1.46 (m, 4H), 1.25 (m, 1H), 0.79 (dt, J = 13.4, 8.7, 1H).
185	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{2H,3H-thieno[3,4-b][1,4]dioxin-5-ylmethyl}amine	415	δ 9.37 (s, 1H), 8.65 (dd, J = 5.3, 1.3, 1H), 8.12 (td, J = 7.9, 1.6, 1H), 7.65 (d, J = 8.2, 1H), 7.56 (dd, J = 7.1, 5.7, 1H), 6.34 (s, 1H), 5.94 (s, 1H), 4.16 (dt, J = 8.2, 6.0, 4H), 4.05 (m, 2H), 3.77 (m, 2H), 3.06 (dd, J = 17.1, 11.1, 1H), 2.61 (t, J = 8.9, 1H), 2.29 (m, 4H), 1.99 (t, J = 8.8, 1H), 1.82 (ddd, J = 13.6, 9.4, 4.3, 2H), 1.67 (m, 1H), 1.48 (ddd, J = 14.5, 12.7, 6.9, 4H), 1.19 (m, 1H), 0.74 (dt, J = 13.3, 9.0, 1H).
186	{2-[2,2-dimethyl-4-(4-methylphenyl)oxan-4-yl]ethyl}{[2-methanesulfonylphenyl]methyl}amine	416.3	δ 8.66 (d, J = 167.7, 2H), 7.92 (m, 1H), 7.52 (m, 3H), 7.05 (s, 4H), 4.21 (s, 2H), 3.71 (m, 2H), 3.49 (s, 1H), 3.06 (s, 3H), 2.85 (s, 1H), 2.47 (d, J = 4.8, 1H), 2.20 (m, 4H), 2.09 (dd, J = 13.9, 2.1, 1H), 1.94 (d, J = 4.6, 1H), 1.72 (s, 1H), 1.55 (m, 2H), 1.10 (s, 3H), 0.57 (s, 3H).
187	[(4-bromofuran-2-yl)methyl]{[2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	419	δ 9.52 (s, 1H), 8.77 (s, 1H), 8.38 (s, 1H), 7.83 (d, J = 7.6, 2H), 7.40 (s, 1H), 6.51 (s, 1H), 4.09 (s, 2H), 3.78 (d, J = 48.0, 2H), 3.03 (s, 1H), 2.63 - 2.41 (m, 2H), 2.33 (dd, J = 28.3, 13.9, 3H), 2.09 (d, J = 14.2, 1H), 1.90 (s, 2H), 1.82 - 1.63 (m, 1H), 1.53 (ddd, J = 12.9, 10.9, 4.5, 4H), 1.19 (s, 1H), 0.79 (dt, J = 13.0, 8.9, 1H).
188	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(methylsulfonyl)thiophen-2-yl]methyl}amine	420	δ 7.22 (dd, J = 8.9, 5.2, 2H), 7.05 (t, J = 8.6, 2H), 6.85 (dd, J = 10.8, 3.7, 2H), 3.95 (s, 2H), 3.75 (d, J = 4.6, 2H), 2.75 (s, 1H), 2.47 (s, 3H), 2.32 (s, 1H), 2.17 (d, J = 14.4, 1H), 2.09 (d, J = 13.8, 1H), 1.99 (dt, J = 12.3, 6.4, 1H), 1.91 (d, J = 13.9, 2H), 1.80 (d, J = 10.5, 1H), 1.74 - 1.61 (m, 2H), 1.49 (dt, J = 18.7, 11.7, 4H), 1.26 (s, 1H), 0.89 - 0.75 (m, 1H).
189	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[6-(trifluoromethyl)pyridin-3-yl]methyl}amine	420.3	δ 8.83 - 8.56 (m, 2H), 8.35 (t, J = 7.6, 1H), 7.96 (dd, J = 19.3, 8.7, 1H), 7.87 - 7.75 (m, 2H), 7.71 (t, J = 9.2, 1H), 4.16 (s, 2H), 3.84 (dd, J = 8.5, 4.4, 1H), 3.71 (t, J = 10.0, 1H), 3.07 (dd, J = 11.7, 6.8, 1H), 2.55 (dt, J = 25.6, 11.9, 2H), 2.43 - 2.21 (m, 3H), 2.10 (d, J = 14.2, 1H), 1.90 (ddd, J = 26.1, 14.9, 6.7, 2H), 1.76 - 1.62 (m, 1H), 1.60 - 1.34 (m, 4H), 1.33 - 1.09 (m, 1H), 0.76 (dt, J = 12.8, 8.8, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
190	[(5-bromofuran-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	421	δ 8.75 (d, J = 4.8, 1H), 8.24 (t, J = 7.7, 1H), 7.86-7.58 (m, 2H), 6.44 (d, J = 3.3, 1H), 6.28 (d, J = 3.3, 1H), 4.07 (s, 2H), 3.94 - 3.79 (m, 1H), 3.72 (t, J = 10.1, 1H), 3.01 (dd, J = 11.1, 6.0, 1H), 2.56 (t, J = 9.9, 1H), 2.49 - 2.11 (m, 4H), 2.05 (d, J = 14.1, 1H), 1.88 (ddd, J = 18.8, 11.0, 6.5, 2H), 1.78 - 1.31 (m, 5H), 1.31-1.07 (m, 1H), 0.77 (dt, J = 13.1, 8.9, 1H)
191	(2-{2,2-dimethyl-4-[4-(trifluoromethyl)phenyl]oxan-4-yl}ethyl)[(2-methoxyphenyl)methyl]amine	422.3	δ 8.49 (d, J = 118.7, 2H), 7.50 (d, J = 8.3, 2H), 7.25 (dd, J = 10.3, 4.6, 3H), 6.96 (dd, J = 7.5, 1.5, 1H), 6.79 (ddd, J = 22.1, 14.4, 4.4, 2H), 6.08 (s, 1H), 3.84 (d, J = 9.2, 2H), 3.68 (m, 5H), 2.64 (s, 1H), 2.09 (m, 3H), 1.90 (m, 1H), 1.77 (dd, J = 12.7, 4.5, 1H), 1.58 (ddd, J = 14.0, 10.8, 10.1, 2H), 1.12 (s, 3H), 0.55 (s, 3H).
192	{2-[2,2,6,6-tetramethyl-4-(pyridin-2-yl)oxan-4-yl]ethyl}({[6-(trifluoromethyl)pyridin-3-yl]methyl})amine	422.3	δ 8.79 - 8.63 (m, 2H), 8.31 (t, J = 7.9, 1H), 8.05 - 7.90 (m, 2H), 7.87 - 7.61 (m, 2H), 4.16 (s, 2H), 2.82 (dd, J = 10.0, 6.6, 2H), 2.54 (d, J = 14.7, 2H), 2.46 - 2.30 (m, 2H), 1.95 (d, J = 14.8, 2H), 1.32 (s, 5H), 0.98 (s, 5H).
193	{[5-(furan-2-yl)thiophen-2-yl]methyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	423.1	δ 9.59 (s, 1H), 8.56 (d, J = 4.7, 1H), 8.05 (t, J = 7.4, 1H), 7.57 (d, J = 8.1, 1H), 7.46 (dd, J = 12.2, 6.3, 1H), 7.35 - 7.26 (m, 1H), 6.93 (dd, J = 19.9, 3.7, 2H), 6.46 - 6.30 (m, 2H), 4.08 (s, 2H), 3.78 - 3.54 (m, 2H), 3.00 - 2.81 (m, 1H), 2.46 (t, J = 9.7, 1H), 2.30 (t, J = 10.6, 1H), 2.13 (ddd, J = 17.3, 16.1, 9.3, 3H), 1.89 (d, J =
			14.2, 1H), 1.72 (ddd, J = 13.9, 9.5, 4.3, 2H), 1.54 (dd, J = 21.6, 14.5, 1H), 1.48 - 1.23 (m, 4H), 1.06 (d, J = 13.2, 1H), 0.65 (dt, J = 13.3, 8.9, 1H).
194	[(5-chlorothiophen-2-yl)methyl]({2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl})amine	423.2	δ 7.13 (dd, J = 8.9, 5.1, 2H), 6.95 (dd, J = 15.5, 6.8, 2H), 6.69 (q, J = 3.9, 2H), 3.81 (s, 2H), 3.62 (d, J = 13.8, 2H), 2.68 (m, 1H), 2.11 (dd, J = 22.2, 13.8, 3H), 1.84 (m, 1H), 1.54 (m, 4H), 1.30 (m, 4H), 1.05 (d, J = 11.4, 4H), 0.63 (m, 1H).
195	(1-benzothiophen-2-yl)methyl({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	424	δ 9.53 (d, J = 105.8, 2H), 7.77 - 7.66 (m, 2H), 7.36 - 7.32 (m, 2H), 7.15 (dd, J = 8.8, 5.2, 3H), 6.96 (t, J = 8.6, 2H), 3.96 (s, 2H), 3.75 - 3.63 (m, 2H), 2.75 (s, 1H), 2.33 (s, 1H), 2.19 - 2.16 (m, 0H), 2.15 - 1.71 (m, 6H), 1.71 - 1.29 (m, 6H), 1.22 (s, 1H), 0.77 (dt, J = 13.5, 9.0, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
196	(1-benzothiophen-3-ylmethyl)({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	424	δ 8.67 (d, J = 139.8, 2H), 7.76 - 7.61 (m, 1H), 7.56 - 7.39 (m, 1H), 7.34 - 7.25 (m, 3H), 6.98 (dd, J = 8.8, 5.1, 2H), 6.84 (t, J = 8.6, 2H), 3.89 (s, 2H), 3.71 - 3.54 (m, 2H), 2.66 (s, 1H), 2.21 (s, 1H), 2.07 - 1.89 (m, 2H), 1.89 - 1.60 (m, 4H), 1.60 - 1.45 (m, 2H), 1.44 - 1.24 (m, 4H), 1.19 - 1.07 (m, 1H), 0.67 (dt, J = 13.8, 8.9, 1H).
197	[(5-fluoro-1-benzothiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	425	δ 8.47 (s, 1H), 7.69 (s, 2H), 7.42 (d, J = 9.2, 1H), 7.30 (dd, J = 10.5, 9.5, 3H), 7.13 (s, 2H), 4.21 (d, J = 13.3, 2H), 3.73 (s, 2H), 3.16 - 2.91 (m, 1H), 2.82 - 2.52 (m, 1H), 2.27 (d, J = 14.8, 2H), 2.21 - 2.09 (m, 1H), 2.08 - 1.94 (m, 1H), 1.85 (d, J = 13.6, 1H), 1.65 (s, 4H), 1.43 (d, J = 38.4, 3H), 1.18 - 1.02 (m, 1H), 0.75 - 0.60 (m, 1H).
198	[(5-cyclopentylthiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	425.1	δ 12.19 - 12.13 (m, 0H), 8.69 (d, J = 4.7, 1H), 8.18 (s, 1H), 7.80 - 7.59 (m, 2H), 6.90 (d, J = 3.5, 1H), 6.67 (d, J = 2.9, 1H), 4.14 (s, 2H), 3.82 (d, J = 12.7, 2H), 3.73 (d, J = 9.7, 1H), 3.17 (t, J = 8.3, 1H), 3.02 (s, 1H), 2.58 (s, 1H), 2.31 (d, J = 14.1, 4H), 2.05 (dd, J = 33.0, 10.0, 3H), 1.90 - 1.74 (m, 4H), 1.68 (dt, J = 12.1, 9.1, 3H), 1.51 (ddd, J = 13.4, 10.8, 5.9, 6H), 1.18 (s, 1H), 0.79 (s, 1H).
199	[(4-phenylphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	427.3	δ 8.59 (d, J = 4.9, 1H), 8.18 (t, J = 7.4, 1H), 7.75 - 7.52 (m, 2H), 7.47 - 7.38 (m, 4H), 7.35 - 7.29 (m, 2H), 7.29 - 7.21 (m, 3H), 3.91 (s, 2H), 3.70 (dt, J = 12.3, 4.2, 1H), 3.57 (t, J = 9.7, 1H), 2.92 (s, 1H), 2.40 (dd, J = 26.0, 12.7, 2H), 2.30 - 2.04 (m, 3H), 2.04 - 1.84 (m, 1H), 1.76 (ddd, J = 27.2, 15.3, 6.8, 2H), 1.65 - 1.21 (m, 5H), 1.07 (dd, J = 14.4, 5.6, 1H), 0.67 (dt, J = 13.0, 9.0, 1H).
200	[(3-phenylphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	427.3	δ 8.44 (d, J = 4.1, 1H), 7.96 (t, J = 7.1, 1H), 7.53 - 7.43 (m, 5H), 7.42 - 7.26 (m, 3H), 7.19 (s, 3H), 3.94 (s, 1H), 3.82 - 3.45 (m, 2H), 2.73 (s, 2H), 2.44 (s, 1H), 2.31 (d, J = 10.6, 1H), 2.15 (d, J = 13.2, 3H), 1.86 (d, J = 14.1, 1H), 1.70 (t, J = 9.7, 2H), 1.56 (s, 1H), 1.50 - 1.22 (m, 5H), 1.06 (s, 1H), 0.66 (dd, J = 13.3, 9.0, 1H).
201	benzyl({2-[9-(4-bromophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	428.2	δ 9.51 (s, 1H), 9.15 (s, 1H), 7.42 (d, J = 8.6, 2H), 7.30 (m, 3H), 7.16 (dd, J = 7.3, 2.1, 2H), 7.06 (d, J = 8.7, 2H), 3.68 (m, 4H), 2.62 (m, 1H), 2.19 (m, 1H), 2.04 (dd, J = 22.4, 13.9, 2H), 1.93 (m, 1H), 1.85 (m, 3H), 1.60 (m, 2H), 1.45 (ddd, J = 21.1, 16.1, 8.8, 5H), 1.25 (m, 2H), 0.77 (dt, J = 13.2, 8.7, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
202	2-amino-4-chloro-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl] thiophene-3-carbonitrile	431	¹ H NMR (400 MHz, CD ₃ CN) δ 8.57 (dd, J = 4.9, 1.0, 1H), 7.83 (m, 1H), 7.48 (d, J = 8.1, 1H), 7.31 (ddd, J = 7.5, 4.9, 0.9, 1H), 6.13 (s, 2H), 4.06 (s, 2H), 3.69 (m, 2H), 2.96 (m, 1H), 2.42 (m, 2H), 2.08 (m, 2H), 1.92 (m, 1H), 1.87 (d, J = 13.5, 1H), 1.57 (m, 8H), 1.10 (m, 1H), 0.71 (m, 1H).
203	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{2H,3H-thieno[3,4-b][1,4]dioxin-5-ylmethyl}amine	432	δ 7.21 (dd, J = 9.0, 5.1, 2H), 7.03 (t, J = 8.6, 2H), 6.33 (s, 1H), 4.13 (s, 4H), 3.90 (s, 2H), 3.74 (m, 2H), 3.01 (brs, 1H), 2.77 (t, J = 13.7, 1H), 2.35 (m, 1H), 2.17 (d, J = 14.0, 1H), 2.02 (dt, J = 14.7, 9.5, 2H), 1.89 (m, 2H), 1.78 (d, J = 10.1, 1H), 1.68 (ddd, J = 16.9, 10.6, 5.8, 2H), 1.46 (ddd, J = 17.8, 10.0, 5.9, 4H), 1.25 (m, 1H), 0.79 (dt, J = 13.5, 8.8, 1H).
204	[(4-phenylthiophen-2-yl)methyl]{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)}amine	433.1	δ 9.71 (s, 4H), 8.53 (d, J = 5.0, 1H), 8.09 (t, J = 7.6, 1H), 7.62 (d, J = 8.1, 1H), 7.54 - 7.44 (m, 1H), 7.43 - 7.35 (m, 2H), 7.32 - 7.19 (m, 5H), 4.12 (s, 2H), 3.77 - 3.52 (m, 2H), 3.00 - 2.76 (m, 1H), 2.41 (dt, J = 25.0, 11.5, 2H), 2.18 (t, J = 17.1, 3H), 1.90 (d, J = 14.1, 1H), 1.73 (ddd, J = 19.6, 11.4, 6.9, 2H), 1.55 (dd, J = 10.0, 4.9, 1H), 1.48 - 1.26 (m, 4H), 1.04 (s, 1H), 0.72 - 0.56 (m, 1H).
205	[(5-phenylthiophen-2-yl)methyl]{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)}amine	433.1	δ 9.74 (brs, 1H), 7.62 (d, J = 8.1, 1H), 7.50 (m, 3H), 7.37 (m, 2H), 7.31 (m, 1H), 7.12 (d, J = 3.7, 1H), 7.04 (d, J = 3.7, 1H), 5.23 (brs, 1H), 4.19 (mz, 2H), 3.72 (m, 2H), 3.02 (d, J = 6.5, 1H), 2.59 (t, J = 9.1, 1H), 2.39 (t, J = 10.1, 1H), 2.22 (dd, J = 29.2, 10.0, 3H), 1.96 (d, J = 14.1, 1H), 1.80 (t, J = 11.0, 2H), 1.62 (dd, J = 14.1, 7.4, 1H), 1.44 (ddd, J = 16.8, 16.4, 7.5, 4H), 1.13 (m, 1H), 0.73 (dt, J = 12.7, 8.8, 1H).
206	[(5-methanesulfonylthiophen-2-yl)methyl]{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)}amine	435	δ 8.67 (d, J = 5.0, 1H), 8.32 (t, J = 8.0, 1H), 7.79 (d, J = 7.9, 2H), 7.59 (d, J = 3.8, 1H), 7.22 (d, J = 3.8, 1H), 4.31 (d, J = 6.2, 2H), 3.84 (s, 1H), 3.74 (s, 1H), 3.18 (s, 1H), 3.05 (s, 2H), 2.54 (t, J = 10.3, 2H), 2.31 (d, J = 13.3, 2H), 2.14 - 2.00 (m, 2H), 1.89 (d, J = 13.8, 3H), 1.81 - 1.64 (m, 1H), 1.64 - 1.37 (m, 3H), 1.28 (s, 2H), 0.81 (d, J = 13.2, 1H).
207	[(4-bromothiophen-3-yl)methyl]{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)}amine	435	δ 11.51 (s, 1H), 9.44 (s, 1H), 8.69 - 8.58 (m, 1H), 8.14 (td, J = 8.0, 1.6, 1H), 7.68 - 7.56 (m, 2H), 7.52 (d, J = 3.4, 1H), 7.21 (d, J = 3.3, 1H), 4.01 (s, 2H), 3.82 - 3.54 (m, 2H), 2.97 (td, J = 11.5, 5.7, 1H), 2.62 - 2.43 (m, 1H), 2.41 - 2.12 (m, 4H), 2.02 - 1.89 (m, 1H), 1.78 (ddd, J = 18.6, 11.9, 6.5, 2H), 1.60 (dt, J = 13.5, 7.7, 1H), 1.55 - 1.30 (m, 4H), 1.10 (d, J = 4.1, 0H), 0.67 (dt, J = 13.1, 8.9, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
208	[[4-bromothiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	435.1	δ 8.59 (d, J = 4.0, 1H), 8.13 (t, J = 7.1, 1H), 7.65 (d, J = 8.2, 1H), 7.59 (m, 1H), 7.23 (d, J = 1.4, 1H), 7.04 (d, J = 1.2, 1H), 4.19 (s, 2H), 3.75 (m, 2H), 3.01 (m, 1H), 2.84 (s, 1H), 2.60 (m, 1H), 2.40 (m, 1H), 2.25 (d, J = 13.0, 3H), 1.97 (d, J = 14.0, 1H), 1.83 (d, J = 9.4, 2H), 1.67 (m, 1H), 1.48 (dd, J = 24.0, 15.8, 4H), 1.17 (brs, 1H), 0.77 (m, 1H).
209	[[5-bromothiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	435.1	δ 8.61 (d, J = 4.3, 1H), 8.14 (t, J = 7.9, 1H), 7.65 (d, J = 8.1, 1H), 7.60 (m, 1H), 6.94 (d, J = 3.8, 1H), 6.89 (d, J = 3.8, 1H), 4.14 (s, 2H), 3.76 (m, 3H), 2.99 (m, 1H), 2.58 (m, 1H), 2.38 (d, J = 9.8, 1H), 2.26 (d, J = 13.9, 3H), 1.97 (d, J = 14.1, 1H), 1.82 (t, J = 9.7, 2H), 1.67 (s, 1H), 1.47 (m, 4H), 1.16 (s, 1H), 0.75 (dt, J = 13.4, 9.2, 1H).
210	[[2-bromothiophen-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	436	δ 8.66 (d, J = 5.3, 1H), 8.21 (d, J = 7.2, 1H), 7.85 - 7.58 (m, 2H), 7.33 (d, J = 5.7, 1H), 7.09 (d, J = 5.7, 1H), 4.02 - 3.63 (m, 3H), 3.10 - 2.97 (m, 2H), 2.61 (t, J = 9.1, 1H), 2.43 (d, J = 11.0, 1H), 2.30 (d, J = 13.6, 3H), 2.04 (s, 1H), 1.94 - 1.80 (m, 2H), 1.69 (s, 1H), 1.64 - 1.40 (m, 4H), 1.20 (s, 1H), 0.86 - 0.68 (m, 1H).
211	[[5-bromofuran-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	436	δ 9.07 (d, J = 116.7, 2H), 7.16 - 7.06 (m, 2H), 7.01 - 6.89 (m, 2H), 6.25 (d, J = 3.4, 1H), 6.17 (s, 1H), 3.83 (s, 2H), 3.76 - 3.58 (m, 2H), 2.66 (s, 1H), 2.23 (s, 1H), 2.09 (d, J = 14.0, 1H), 2.04 - 1.96 (m, 1H), 1.95 - 1.66 (m, 4H), 1.66 - 1.50 (m, 2H), 1.50 - 1.28 (m, 4H), 1.28 - 1.13 (m, 1H), 0.71 (dt, J = 13.6, 8.8, 1H).
212	(2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl){[6-(trifluoromethyl)pyridin-3-yl]methyl}amine	437.2	δ 8.63 (s, 1H), 7.83 (d, J = 8.3, 1H), 7.68 (d, J = 8.0, 1H), 7.29 (s, 1H), 7.21 (dd, J = 8.9, 5.1, 3H), 7.05 (s, 2H), 3.93 (s, 2H), 3.75 (dd, J = 11.3, 7.3, 2H), 2.84 - 2.58 (m, 1H), 2.44 - 2.04 (m, 10H), 2.02 - 1.75 (m, 5H), 1.74 - 1.56 (m, 3H), 1.59 - 1.33 (m, 5H), 1.33 - 1.19 (m, 1H), 0.78 (d, J = 13.6, 1H).
213	[[4-bromofuran-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	437.9	δ 7.38 (d, J = 0.6, 1H), 7.28 (s, 1H), 7.22 (d, J = 5.2, 2H), 7.08 (d, J = 8.5, 2H), 3.75 (dd, J = 11.7, 7.1, 2H), 2.73 (s, 1H), 2.30 (d, J = 4.5, 2H), 2.17 (d, J = 13.5, 1H), 2.10 (d, J = 13.9, 1H), 2.05 - 1.95 (m, 1H), 1.94 (s, 2H), 1.79 (d, J = 9.8, 1H), 1.74 - 1.62 (m, 2H), 1.49
			(dt, J = 16.4, 10.6, 4H), 1.28 (s, 2H), 0.80 (d, J = 13.7, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
214	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(thiophen-2-yl)thiophen-2-yl]methyl}amine	439	δ 8.52 (d, J = 5.3, 1H), 7.96 (t, J = 7.9, 1H), 7.50 (d, J = 8.1, 1H), 7.40 (dd, J = 16.2, 10.4, 1H), 7.24 - 7.10 (m, 1H), 7.03 (dd, J = 3.6, 1.0, 1H), 6.98 - 6.81 (m, 3H), 4.07 (s, 2H), 3.80 - 3.49 (m, 2H), 2.90 (d, J = 11.1, 2H), 2.16 (s, 5H), 1.87 (d, J = 14.0, 1H), 1.71 (dd, J = 11.5, 7.2, 2H), 1.54 (d, J = 6.1, 1H), 1.36 (ddd, J = 16.8, 12.9, 6.1, 5H), 1.05 (s, 1H), 0.82 - 0.54 (m, 1H).
215	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}{[6-(trifluoromethyl)pyridin-3-yl]methyl}amine	439.3	δ 8.62 (s, 1H), 7.84 (d, J = 8.2, 1H), 7.66 (d, J = 8.2, 1H), 7.20 (m, 1H), 7.04 (s, 2H), 3.90 (s, 2H), 3.71 (d, J = 12.1, 2H), 2.77 (m, 1H), 2.19 (m, 3H), 1.98 (m, 1H), 1.68 (m, 3H), 1.40 (d, J = 7.6, 2H), 1.04 (s, 1H), 0.83 (t, J = 7.5, 4H), 0.54 (d, J = 7.3, 3H).
216	[(5-chloro-1-benzothiophen-3-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	442	δ 8.61 (d, J = 4.9, 1H), 8.44 (s, 1H), 8.17 (s, 1H), 7.93 (d, J = 5.5, 1H), 7.69 (d, J = 8.1, 1H), 7.60 (s, 1H), 7.50 (d, J = 5.5, 1H), 7.28 (s, 1H), 3.82 (s, 3H), 3.17 (dd, J = 16.8, 10.9, 1H), 2.75 (t, J = 8.9, 1H), 2.47 (t, J = 9.7, 1H), 2.32 (d, J = 13.9, 3H), 2.10 - 1.98 (m, 1H), 1.87 (dd, J = 12.1, 7.1, 2H), 1.78 - 1.62 (m, 1H), 1.48 (dd, J = 23.5, 18.9, 5H), 1.18 (s, 1H), 0.77 (dt, J = 13.2, 9.0, 1H).
217	[(5-bromo-4-methylthiophen-2-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	449	δ 10.10 - 9.21 (m, 1H), 8.53 (d, J = 3.9, 1H), 7.90 (td, J = 7.9, 1.6, 1H), 7.45 (d, J = 8.1, 1H), 7.38 (dd, J = 7.0, 5.4, 1H), 6.69 (s, 1H), 4.02 - 3.86 (m, 2H), 3.74 - 3.55 (m, 2H), 2.85 (dd, J = 11.4, 5.9, 1H), 2.47 - 2.33 (m, 1H), 2.31 - 2.09 (m, 3H), 2.09 - 1.93 (m, 4H), 1.87 (d, J = 14.0, 1H), 1.69 (dt, J = 14.4, 6.1, 2H), 1.57 (d, J = 5.4, 1H), 1.38 (ddd, J = 26.7, 14.6, 8.4, 4H), 1.04 (s, 1H), 0.73 - 0.56 (m, 1H).
218	[(4-bromo-5-methylthiophen-2-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	449	δ 8.72 (d, J = 4.9, 1H), 8.26 (d, J = 7.7, 1H), 7.74 (dd, J = 16.7, 7.1, 2H), 6.92 (s, 1H), 4.21 (d, 1H), 3.90 - 3.78 (m, 2H), 3.74 (d, J = 9.6, 1H), 3.02 (s, 1H), 2.51 (dd, J = 52.4, 11.0, 2H), 2.39 - 2.16 (m, 6H), 2.05 (d, J = 13.8, 1H), 1.87 (d, J = 9.5, 2H), 1.69 (s, 1H), 1.63 - 1.41 (m, 4H), 1.24 (d, J = 30.9, 1H), 0.81 (s, 1H).
219	[(3-bromo-5-methylthiophen-2-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	449	δ 8.61 (d, J = 4.9, 1H), 8.02 (d, J = 7.9, 1H), 7.69 - 7.41 (m, 2H), 6.68 (d, J = 1.0, 1H), 4.23 (q, J = 14.2, 2H), 3.90 - 3.59 (m, 2H), 3.10 (s, 1H), 2.75 (m, 2H), 2.36 - 2.13 (m, 5H), 1.96 (d, J = 13.9, 1H), 1.82 (d, J = 9.9, 2H), 1.75 - 1.62 (m, 1H), 1.62 - 1.38 (m, 4H), 1.24 - 1.05 (m, 1H), 0.74 (d, J = 13.2, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
220	[(4-bromo-3-methylthiophen-2-yl)methyl]({2-[(9R)-9-yl]ethyl})amine	449	δ 8.61 (d, J = 5.2, 1H), 8.09 (t, J = 7.7, 1H), 7.71 - 7.49 (m, 2H), 7.30 (s, 1H), 4.21 (d, J = 4.3, 2H), 4.00 - 3.59 (m, 2H), 3.05 (s, 1H), 2.64 (s, 1H), 2.31 (d, J =
	(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine		14.5, 2H), 2.25 (d, J = 13.7, 2H), 2.18 (s, 3H), 1.96 (d, J = 13.9, 1H), 1.82 (dd, J = 12.0, 7.2, 2H), 1.68 (s, 1H), 1.61 - 1.40 (m, 4H), 1.17 (s, 1H), 0.92 - 0.64 (m, 1H).
221	{2-[4-(4-fluorophenyl)-1-oxaspiro[5.5]undecan-4-yl]ethyl}({[6-(trifluoromethyl)pyridin-3-yl]methyl})amine	451.2	δ 8.52 (s, 1H), 7.73 (d, J = 9.6, 1H), 7.57 (d, J = 8.0, 1H), 7.11 (dd, J = 9.0, 5.2, 2H), 6.94 (t, J = 8.4, 2H), 3.83 (s, 2H), 3.63 (d, J = 18.0, 2H), 2.69 (m, 1H), 2.12 (t, J = 13.9, 3H), 1.70 (m, 5H), 1.31 (d, J = 18.3, 4H), 1.03 (s, 4H), 0.57 (m, 1H).
222	[(4-bromothiophen-3-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	451.9	δ 9.26 (d, J = 136.7, 2H), 7.39 (dd, J = 22.6, 19.3, 2H), 7.10 (dd, J = 8.8, 5.2, 2H), 6.92 (dd, J = 10.6, 6.6, 2H), 3.82 (s, 2H), 3.71 - 3.53 (m, 2H), 2.64 (s, 1H), 2.20 (s, 1H), 2.05 (d, J = 14.1, 1H), 1.97 (d, J = 13.9, 1H), 1.89 (td, J = 12.6, 4.6, 1H), 1.83 - 1.64 (m, 3H), 1.57 (ddd, J = 14.0, 9.6, 4.7, 2H), 1.49 - 1.25 (m, 4H), 1.17 (d, J = 13.2, 1H), 0.69 (dt, J = 13.8, 8.8, 1H).
223	[(4-bromothiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	452.1	δ 9.38 (d, J = 89.0, 2H), 7.16 - 7.03 (m, 3H), 6.96 (t, J = 8.6, 2H), 6.84 (d, J = 1.3, 1H), 3.86 (s, 2H), 3.70 - 3.55 (m, 2H), 2.62 (dd, J = 12.1, 7.7, 1H), 2.18 (dd, J = 11.9, 7.8, 1H), 2.02 (dd, J = 32.5, 14.0, 2H), 1.91 - 1.63 (m, 4H), 1.64 - 1.50 (m, 2H), 1.49 - 1.25 (m, 4H), 1.16 (dd, J = 14.0, 6.1, 1H), 0.69 (dt, J = 13.5, 8.8, 1H).
224	[(5-bromothiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	452.1	δ 9.31 (d, J = 92.6, 2H), 7.15 - 7.04 (m, 2H), 6.95 (t, J = 8.6, 2H), 6.82 (d, J = 3.8, 1H), 6.67 (d, J = 3.8, 1H), 3.82 (s, 2H), 3.70 - 3.53 (m, 2H), 3.44 (s, 1H), 2.62 (dd, J = 12.0, 7.6, 1H), 2.18 (dd, J = 11.8, 7.9, 1H), 2.02 (dd, J = 31.6, 14.0, 2H), 1.93 - 1.64 (m, 4H), 1.57 (ddd, J = 12.1, 8.5, 3.8, 2H), 1.52 - 1.25 (m, 4H), 1.16 (dd, J = 14.9, 5.1, 1H), 0.69 (dt, J = 13.6, 8.8, 1H).
225	dibenzyl({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	458.3	δ 7.27 (m, 17H), 7.00 (dd, J = 8.9, 5.2, 2H), 6.86 (t, J = 8.6, 2H), 4.24 (s, 2H), 3.90 (m, 2H), 3.55 (d, J = 3.4, 2H), 2.59 (m, 1H), 2.22 (m, 12H), 1.86 (dd, J = 75.2, 14.8, 7H), 1.59 (dd, J = 44.5, 9.1, 2H), 1.38 (m, 6H), 1.18 (s, 1H), 1.11 (s, 1H), 0.68 (m, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
226	dibenzyl({2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl})amine	460.3	δ 7.27 (d, J = 34.4, 7H), 7.18 (s, 4H), 6.98 (dd, J = 8.9, 5.2, 2H), 6.84 (t, J = 8.6, 2H), 4.25 (s, 2H), 3.85 (d, J = 46.4, 2H), 3.53 (m, 2H), 2.57 (d, J = 4.7, 2H), 2.12 (d, J = 4.0, 2H), 1.97 (m, 3H), 1.73 (d, J = 4.8, 1H), 1.44 (s, 1H), 1.38 (dd, J = 13.8, 7.5, 3H), 1.23 (m, 1H), 0.90 (m, 1H), 0.70 (dt, J = 10.8, 7.4, 4H), 0.40 (t, J = 7.4, 3H).
227	[(4-bromo-3-methylthiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	465.9	δ 7.19 (dd, J = 8.9, 5.1, 2H), 7.04 (t, J = 8.6, 2H), 3.94 (d, J = 16.3, 2H), 3.72 (m, 2H), 2.72 (dd, J = 13.8, 6.5, 1H), 2.31 (m, 1H), 2.15 (d, J = 12.1, 2H), 2.07 (s, 3H), 1.90 (m, 5H), 1.65 (m, 2H), 1.47 (m, 4H), 1.25 (s, 1H), 0.78 (m, 1H).
228	[(4-bromo-5-methylthiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	465.9	δ 9.06 (d, J = 100.4, 2H), 7.15 - 7.04 (m, 2H), 6.95 (s, 2H), 6.68 (s, 1H), 3.80 (s, 2H), 3.73 - 3.57 (m, 2H), 2.64 (s, 1H), 2.21 (s, 4H), 2.07 (d, J = 14.1, 1H), 1.99 (d, J = 13.9, 1H), 1.94 - 1.64 (m, 4H), 1.64 - 1.51 (m, 2H), 1.51 - 1.26 (m, 4H), 1.17 (dd, J = 13.9, 6.3, 1H), 0.70 (dt, J = 13.7, 8.8, 1H).
229	[(3-bromo-5-methylthiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	466	δ 7.20 (m, 2H), 7.01 (dd, J = 11.1, 6.1, 2H), 6.61 (d, J = 1.1, 1H), 4.01 (s, 2H), 3.72 (m, 2H), 2.75 (m, 1H), 2.61 (brs, 1H), 2.41 (d, J = 0.9, 3H), 2.32 (m, 1H), 2.15 (d, J = 14.2, 1H), 2.07 (d, J = 13.9, 1H), 1.99 (m, 1H), 1.89 (m, 2H), 1.77 (m, 1H), 1.67 (ddd, J = 17.0, 10.6, 5.6, 2H), 1.46 (m, 4H), 1.25 (m, 1H), 0.78 (dt, J = 13.9, 8.9, 1H).
230	[(5-bromo-4-methylthiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	466.9	δ 7.20 (d, J = 5.2, 2H), 7.06 (d, J = 8.5, 2H), 6.85 (t, J = 3.6, 1H), 3.91 (s, 2H), 3.81 - 3.62 (m, 2H), 2.71 (s, 1H), 2.28 (s, 1H), 2.07 (s, 6H), 1.91 (d, J = 13.8, 2H), 1.79 (d, J = 10.3, 1H), 1.69 (ddd, J = 14.1, 9.4, 4.7, 2H), 1.59 - 1.37 (m, 4H), 1.28 (s, 1H), 0.80 (dd, J = 8.8, 4.9, 1H).
231	{2-[2,2-diethyl-4-(4-fluorophenyl)oxan-4-yl]ethyl}bis(thiophen-2-ylmethyl)amine	472.2	δ 7.31 (d, J = 4.9, 2H), 7.07 (dd, J = 8.9, 5.2, 2H), 6.99 (s, 2H), 6.95 (d, J = 4.5, 2H), 6.89 (t, J = 8.6, 2H), 4.24 (s, 2H), 3.58 (dt, J = 23.8, 6.6, 2H), 2.66 (m, 1H), 2.06 (d, J = 14.0, 4H), 1.82 (m, 2H), 1.51 (d, J = 14.3, 3H), 1.25 (m, 2H), 0.94 (dd, J = 14.6, 7.4, 1H), 0.74 (t, J = 7.5, 4H), 0.42 (t, J = 7.4, 3H).
232	[(4,5-dibromothiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	514.8	δ 8.61 (dd, J = 5.3, 1.3, 1H), 8.13 (td, J = 8.0, 1.7, 1H), 7.64 (d, J = 8.2, 1H), 7.60 (m, 1H), 6.94 (s, 1H), 4.40 (brs, 1H), 4.11 (s, 2H), 3.77 (ddd, J = 36.9, 13.7, 7.2, 2H), 2.98 (td, J = 11.3, 6.0, 1H), 2.54 (td, J = 11.2, 4.3, 1H), 2.38 (m, 1H), 2.22 (m, 3H), 1.98 (d, J = 14.0, 1H), 1.83 (dt, J = 18.5, 9.2, 2H), 1.68 (m, 1H), 1.48 (m, 4H), 1.21 (d, J = 37.1, 1H), 0.75 (dt, J = 13.1, 9.0, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
233	[(3,4-dibromothiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	514.8	δ 8.39 (d, J = 4.0, 1H), 7.67 (t, J = 7.0, 1H), 7.38 (s, 1H), 7.28 (d, J = 8.1, 1H), 7.18 (s, 1H), 4.22 (d, J = 18.2, 2H), 3.65 (dd, J = 11.2, 7.1, 2H), 3.09 - 2.85 (m, 1H), 2.60 (s, 1H), 2.22 (dd, J = 25.9, 13.8, 2H), 2.10 - 1.83 (m, 2H), 1.87 - 1.50 (m, 4H), 1.36 (dd, J = 18.7, 10.7, 3H), 1.02 (s, 2H), 0.68 - 0.50 (m, 1H).
234	[(4,5-dibromothiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	531.8	δ 7.20 (m, 2H), 7.05 (t, J = 8.6, 2H), 6.81 (s, 1H), 3.91 (s, 2H), 3.74 (m, 2H), 3.60 (brs, 1H), 2.74 (m, 1H), 2.31 (td, J = 12.1, 4.7, 1H), 2.15 (d, J = 14.1, 1H), 2.08 (d, J = 13.9, 1H), 1.88 (m, 4H), 1.67 (ddd, J = 15.1, 10.2, 5.0, 2H), 1.46 (ddd, J = 27.4, 14.5, 7.2, 4H), 1.24 (dd, J = 10.5, 5.6, 1H), 0.78 (dt, J = 13.5, 8.8, 1H).
235	[(3,4-dibromothiophen-2-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	531.8	δ 7.35 (s, 1H), 7.11 - 7.06 (m, 2H), 6.94 (dd, J = 14.3, 5.7, 2H), 4.05 (s, 2H), 3.75 - 3.56 (m, 2H), 2.70 (dd, J = 11.9, 7.4, 1H), 2.25 (dd, J = 11.7, 7.3, 1H), 2.08 (d, J = 14.7, 1H), 1.99 (d, J = 13.9, 1H), 1.95 - 1.65 (m, 4H), 1.65 - 1.50 (m, 2H), 1.50 - 1.29 (m, 4H), 1.16 (dd, J = 14.8, 7.3, 1H), 0.70 (dt, J = 13.6, 8.8, 1H).
236	[(2-fluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	369	δ 8.82 (s, 2H), 8.61 (dd, J = 4.8, 1.2, 1H), 7.85 (td, J = 7.8, 1.8, 1H), 7.51 (m, 3H), 7.30 (m, 3H), 5.22 (s, 2H), 4.12 (d, J = 5.3, 2H), 3.66 (m, 2H), 2.90 (d, J = 4.5, 1H), 2.39 (m, 3H), 2.08 (td, J = 12.8, 4.4, 1H), 1.54 (m, 7H), 1.02 (dd, J = 12.3, 5.8, 1H), 0.68 (dt, J = 13.3, 8.9, 1H).
237	[(2-bromophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	429	δ 8.93 (s, 2H), 8.60 (dd, J = 4.8, 1.2, 1H), 7.83 (td, J = 7.8, 1.9, 1H), 7.71 (dd, J = 8.0, 1.1, 1H), 7.50 (m, 3H), 7.33 (m, 2H), 4.18 (s, 2H), 3.65 (m, 2H), 2.94 (s, 1H), 2.43 (t, J = 12.2, 3H), 2.11 (td, J = 12.8, 4.4, 1H), 1.89 (m, 2H), 1.55 (m, 7H), 1.01 (m, 1H), 0.67 (dt, J = 13.3, 8.9, 1H).
238	[(2-chlorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	385	δ 8.75 (dd, J = 5.4, 1.2, 1H), 8.52 (s, 3H), 8.22 (td, J = 8.0, 1.7, 1H), 7.77 (d, J = 8.2, 1H), 7.67 (ddd, J = 7.5, 5.4, 0.9, 1H), 7.42 (m, 4H), 4.20 (d, J = 14.0, 2H), 3.72 (m, 2H), 3.05 (td, J = 12.0, 5.1, 1H), 2.53 (td, J = 12.0, 4.4, 1H), 2.36 (m, 3H), 2.17 (m, 1H), 2.01 (d, J = 14.2, 1H), 1.79 (ddd, J = 9.3, 6.7, 3.4, 2H), 1.52 (m, 5H), 1.17 (m, 1H), 0.78 (dt, J = 12.9, 8.8, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
239	[(2-methylphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	365.1	δ 8.81 (dd, J = 5.7, 1.3, 1H), 8.43 (td, J = 8.0, 1.7, 1H), 7.98 (s, 1H), 7.92 (d, J = 8.2, 1H), 7.86 (ddd, J = 7.6, 5.7, 1.0, 1H), 7.27 (m, 2H), 7.18 (m, 2H), 3.98 (s, 2H), 3.75 (m, 2H), 2.98 (d, J = 4.4, 1H), 2.44 (m, 2H), 2.36 (m, 5H), 2.25 (dd, J = 13.5, 5.4, 1H), 2.07 (d, J = 14.3, 1H), 1.84 (m, 2H), 1.55 (m, 5H), 1.23 (m, 1H), 0.83 (dt, J = 13.0, 8.8, 1H).
240	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[2-(trifluoromethyl)phenyl]methyl})amine	419.1	δ 8.71 (dd, J = 5.3, 1.2, 1H), 8.18 (td, J = 8.0, 1.7, 1H), 7.79 (d, J = 7.8, 1H), 7.75 (d, J = 8.2, 1H), 7.70 (m, 2H), 7.62 (ddd, J = 13.7, 6.8, 1.2, 2H), 6.71 (s, 3H), 4.23 (s, 2H), 3.75 (ddd, J = 17.6, 8.8, 3.7, 2H), 3.08 (m, 1H), 2.56 (m, 1H), 2.36 (m, 3H), 2.19 (m, 1H), 1.79 (dq, J = 7.2, 4.7, 2H), 1.53 (m, 5H), 1.19 (m, 1H), 0.79 (m, 1H).
241	2-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]phenol	367	δ 8.74 (m, 1H), 8.21 (td, J = 8.0, 1.8, 1H), 7.75 (d, J = 8.2, 1H), 7.66 (ddd, J = 7.6, 5.4, 1.0, 1H), 7.27 (m, 1H), 7.20 (dd, J = 7.6, 1.6, 1H), 6.90 (m, 4H), 4.05 (s, 2H), 3.72 (ddd, J = 12.4, 11.1, 5.4, 2H), 2.96 (d, J = 5.2, 1H), 2.35 (m, 4H), 2.13 (m, 1H), 1.78 (m, 2H), 1.51 (m, 5H), 1.15 (dd, J = 4.0, 2.0, 1H), 0.78 (m, 1H).
242	[(2-methoxyphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	381.1	δ 9.66 (s, 3H), 8.80 (dd, J = 5.5, 1.2, 1H), 8.31 (td, J = 8.0, 1.7, 1H), 7.77 (m, 3H), 7.42 (ddd, J = 15.9, 8.0, 1.6, 1H), 7.25 (dd, J = 7.5, 1.6, 1H), 7.04 (m, 1H), 6.96 (td, J = 7.5, 1.0, 1H), 4.04 (s, 2H), 3.85 (m, 4H), 3.73 (m, 2H), 2.97 (d, J = 4.9, 1H), 2.37 (m, 4H), 2.19 (dd, J = 13.2, 5.2, 1H), 2.04 (d, J = 14.1, 1H), 1.81 (ddd, J = 14.0, 9.5, 4.5, 2H), 1.81 (ddd, J = 14.0, 9.5, 4.5, 2H), 1.54 (m, 5H), 1.18 (m, 1H), 0.80 (m, 1H).
243	[(3-fluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	369	δ 8.77 (dd, J = 5.4, 1.5, 1H), 8.38 (s, 1H), 8.29 (td, J = 8.0, 1.7, 1H), 7.82 (d, J = 8.2, 1H), 7.74 (dd, J = 7.1, 6.1, 1H), 7.43 (ddd, J = 13.8, 7.5, 1.4, 1H), 7.19 (m, 3H), 6.90 (s, 3H), 4.03 (d, J = 2.0, 2H), 3.74 (m, 2H), 2.98 (dt, J = 11.4, 5.6, 1H), 2.42 (ddd, J = 29.2, 13.0, 3.8, 4H), 2.18 (m, 1H), 2.03 (d, J = 14.1, 1H), 1.81 (ddd, J = 13.9, 9.4, 4.5, 2H), 1.55 (m, 5H), 1.20 (ddd, J = 9.9, 6.9, 2.4, 1H), 0.80 (dt, J = 12.9, 8.8, 1H).
244	[(3-bromophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	431	δ 8.75 (dd, J = 5.4, 1.2, 1H), 8.41 (s, 1H), 8.25 (td, J = 8.0, 1.8, 1H), 7.79 (d, J = 8.2, 1H), 7.70 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.59 (m, 2H), 7.36 (ddd, J = 22.8, 10.9, 4.6, 2H), 6.76 (s, 3H), 4.01 (d, J = 2.3, 2H), 3.74 (ddd, J = 12.3, 11.0, 5.4, 2H), 2.97 (d, J = 5.0, 1H), 2.38 (m, 4H), 2.16 (m, 1H), 2.00 (m, 1H), 1.79 (ddd, J = 8.6, 7.8, 4.7, 2H), 1.53 (m, 5H), 1.20 (m, 1H), 0.80 (m, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
245	[(3-chlorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	385	δ 8.72 (dd, J = 5.4, 1.1, 1H), 8.57 (s, 1H), 8.19 (td, J = 8.0, 1.8, 1H), 7.74 (d, J = 8.2, 1H), 7.64 (ddd, J = 7.6, 5.4, 0.9, 1H), 7.37 (m, 5H), 3.99 (d, J = 2.3, 2H), 3.70 (m, 2H), 2.95 (m, 1H), 2.36 (m, 4H), 2.12 (td, J = 12.9, 5.1, 1H), 1.76 (ddd, J = 14.2, 9.3, 5.1, 2H), 1.50 (m, 5H), 0.77 (dt, J = 13.0, 8.9, 1H).
246	[(3-methylphenyl)methyl]	365	δ 8.81 (dd, J = 5.7, 1.3, 1H), 8.43 (td, J = 8.0, 1.7, 1H), 7.98 (s, 1H), 7.92 (d, J = 8.2, 1H), 7.86 (ddd, J = 7.6, 5.7, 1.0, 1H), 7.24 (dq, J = 19.8, 7.4, 4H), 3.98 (s, 2H), 3.75 (m, 2H), 2.98 (d, J = 4.4, 1H), 2.44 (m, 2H), 2.36 (m, 5H), 2.25 (dd, J = 13.5, 5.4, 1H), 2.07 (d, J = 14.3, 1H), 1.84 (m, 2H), 1.55 (m, 5H), 1.23 (m, 1H), 0.83 (dt, J = 13.0, 8.8, 1H).
247	methyl 3-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]benzoate	409.1	δ 8.78 (dd, J = 5.5, 1.3, 1H), 8.33 (td, J = 8.0, 1.7, 1H), 8.24 (s, 1H), 8.04 (m, 2H), 7.85 (d, J = 8.2, 1H), 7.78 (m, 1H), 7.63 (m, 2H), 7.55 (d, J = 7.7, 1H), 4.10 (d, J = 2.0, 2H), 3.90 (s, 3H), 3.75 (ddd, J = 12.2, 11.0, 5.4, 2H), 3.00 (dd, J = 11.5, 7.1, 1H), 2.40 (m, 4H), 2.21 (m, 1H), 2.04 (d, J = 14.2, 1H), 1.83 (ddd, J = 13.9, 9.2, 4.3, 2H), 1.51 (dddd, J = 17.6, 10.1, 8.1, 3.0, 5H), 1.21 (s, 1H), 0.82 (dd, J = 15.6, 6.6, 1H).
248	3-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]phenol	367	δ 8.77 (dd, J = 5.5, 1.2, 1H), 8.30 (td, J = 8.0, 1.7, 1H), 7.81 (s, 1H), 7.75 (ddd, J = 7.6, 5.5, 1.0, 1H), 7.23 (t, J = 8.1, 1H), 6.85 (dt, J = 3.2, 2.1, 3H), 6.65 (s, 3H), 3.96 (s, 2H), 3.73 (dd, J = 13.8, 7.3, 2H), 2.96 (s, 1H), 2.36 (m, 4H), 2.15 (ddd, J = 9.9, 8.5, 4.7, 1H), 2.03 (d, J = 14.2, 1H), 1.80 (dt, J = 11.2, 4.8, 2H), 1.52 (ddd, J = 21.7, 12.8, 7.4, 5H), 1.20 (m, 1H), 0.80 (d, J = 13.3, 1H).
249	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({3-(trifluoromethyl)phenyl)methyl}amine	419.1	δ 8.66 (m, 1H), 8.07 (td, J = 7.9, 1.8, 1H), 7.72 (m, 2H), 7.65 (m, 2H), 7.54 (m, 2H), 6.22 (s, 2H), 4.08 (d, J = 3.1, 2H), 3.71 (m, 2H), 2.97 (d, J = 5.0, 1H), 2.34 (dddd, J = 25.4, 19.6, 16.7, 4.4, 4H), 2.09 (m, 1H), 1.74 (m, 2H), 1.49 (m, 5H), 0.76 (m, 1H).
250	N-methyl-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]thiophene-2-carboxamide	414.1	δ 8.78 (dd, J = 5.6, 1.3, 1H), 8.36 (td, J = 8.0, 1.7, 1H), 7.86 (d, J = 8.2, 1H), 7.79 (ddd, J = 7.6, 5.6, 1.0, 1H), 7.38 (d, J = 3.8, 1H), 7.14 (d, J = 3.8, 1H), 7.06 (d, J = 3.9, 1H), 4.21 (s, 2H), 3.72 (m, 2H), 2.97 (td, J = 12.0, 5.1, 1H), 2.84 (d, J = 4.7, 3H), 2.34 (m, 4H), 2.16 (m, 1H), 2.03 (d, J = 14.2, 1H), 1.80 (dd, J = 12.2, 3.0, 2H), 1.50 (m, 5H), 1.20 (m, 1H), 0.82 (s, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
251	N-ethyl-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]thiophene-2-carboxamide	428.1	δ 8.78 (dd, J = 5.6, 1.2, 1H), 8.35 (td, J = 8.0, 1.7, 1H), 7.86 (d, J = 8.2, 1H), 7.79 (ddd, J = 7.6, 5.6, 0.9, 1H), 7.40 (d, J = 3.8, 1H), 7.14 (t, J = 10.7, 2H), 6.20 (s, 4H), 4.21 (d, J = 1.2, 2H), 3.72 (m, 2H), 3.34 (m, 2H), 2.97 (m, 1H), 2.39 (m, 4H), 2.18 (m, 1H), 2.03 (d, J = 14.2, 1H), 1.81 (m, 2H), 1.52 (m, 5H), 1.19 (m, 4H), 0.80 (m, 1H).
252	N-methyl-3-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]benzamide	408.1	δ 8.75 (dd, J = 5.4, 1.2, 1H), 8.24 (td, J = 8.0, 1.7, 1H), 7.89 (s, 1H), 7.77 (m, 2H), 7.69 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.49 (ddd, J = 18.3, 10.6, 4.6, 2H), 7.33 (s, 1H), 4.07 (s, 2H), 3.73 (m, 2H), 2.98 (m, 1H), 2.88 (d, J = 4.6, 3H), 2.47 (t, J = 10.7, 1H), 2.36 (dd, J = 12.8, 7.5, 3H), 2.16 (d, J = 4.8, 1H), 1.79 (m, 2H), 1.50 (ddd, J = 18.7, 13.3, 6.9, 5H), 1.19 (ddd, J = 8.3, 7.0, 1.8, 1H), 0.80 (d, J = 13.3, 1H).
253	N-ethyl-3-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]benzamide	422.1	δ 8.70 (dd, J = 5.3, 1.2, 1H), 8.43 (s, 1H), 8.14 (td, J = 7.9, 1.8, 1H), 7.89 (d, J = 1.4, 1H), 7.76 (dt, J = 7.3, 1.6, 1H), 7.71 (d, J = 8.2, 1H), 7.59 (ddd, J = 7.6, 5.3, 0.9, 1H), 7.46 (m, 2H), 7.37 (s, 1H), 6.55 (s, 3H), 4.05 (s, 2H), 3.70 (m, 2H), 3.36 (qd, J = 7.2, 5.7, 2H), 2.96 (d, J = 7.9, 1H), 2.45 (t, J = 10.2, 1H), 2.32 (dd, J = 21.2, 8.7, 3H), 2.11 (d, J = 5.2, 1H), 1.77 (m, 2H), 1.47 (m, 5H), 1.19 (m, 4H), 0.76 (d, J = 13.3, 1H).
254	[(4-methoxyphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	381.1	δ 9.09 (d, J = 86.1, 2H), 8.69 (d, J = 5.0, 1H), 8.29 (t, J = 7.7, 1H), 8.06 (s, 3H), 7.75 (m, 2H), 7.20 (d, J = 8.6, 2H), 6.81 (d, J = 8.6, 2H), 3.89 (s, 2H), 3.79 (m, 4H), 3.66 (m, 1H), 2.96 (s, 1H), 2.43 (dd, J = 23.4, 11.5, 2H), 2.27 (t, J = 16.0, 3H), 2.01 (d, J = 14.2, 1H), 1.83 (dd, J = 19.3, 9.5, 2H), 1.66 (m, 1H), 1.47 (m, 4H), 1.14 (d, J = 7.0, 1H), 0.75 (m, 1H).
255	4-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]phenol	367	δ 8.76 (dd, J = 5.5, 1.2, 1H), 8.26 (m, 1H), 7.80 (d, J = 8.2, 1H), 7.72 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.65 (s, 1H), 7.22 (m, 2H), 6.82 (m, 2H), 3.93 (s, 2H), 3.73 (dd, J = 10.7, 7.6, 2H), 2.94 (dd, J = 11.1, 5.9, 1H), 2.36 (m, 4H), 2.13 (m, 1H), 2.01 (m, 1H), 1.80 (d, J = 3.6, 2H), 1.51 (dd, J = 9.7, 5.6, 5H), 1.19 (m, 1H), 0.81 (s, 1H).
256	[(2,3-difluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387	δ 8.75 (dd, J = 5.4, 1.3, 1H), 8.26 (td, J = 8.0, 1.7, 1H), 7.80 (d, J = 8.2, 1H), 7.71 (ddd, J = 7.5, 5.5, 0.9, 1H), 7.34 (dtd, J = 10.0, 7.9, 1.9, 1H), 7.21 (m, 4H), 4.11 (s, 2H), 3.72 (m, 2H), 3.02 (td, J = 12.0, 5.1, 1H), 2.49 (td, J = 12.1, 4.3, 1H), 2.35 (m, 3H), 2.16 (m, 1H), 2.01 (d, J = 14.1, 1H), 1.79 (ddd, J = 11.2, 9.4, 4.1, 2H), 1.52 (m, 5H), 1.17 (m, 1H), 0.78 (dt, J = 12.9, 8.8, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
257	[(2,4-difluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387	δ 8.79 (dd, J = 5.5, 1.3, 1H), 8.38 (s, 1H), 8.32 (m, 1H), 7.84 (d, J = 8.2, 1H), 7.76 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.50 (dd, J = 14.8, 8.3, 1H), 7.03 (m, 2H), 4.08 (s, 2H), 3.74 (m, 2H), 3.02 (td, J = 12.0, 5.0, 1H), 2.42 (m, 4H), 2.18 (m, 1H), 2.03 (d, J = 14.2, 1H), 1.82 (ddd, J = 14.2, 9.6, 4.5, 2H), 1.56 (m, 5H), 1.19 (ddd, J = 7.0, 6.2, 2.8, 1H), 0.80 (dt, J = 12.9, 8.8, 1H).
258	[(2,5-difluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387	δ 8.74 (dd, J = 5.4, 1.2, 1H), 8.26 (td, J = 8.0, 1.7, 1H), 7.72 (m, 2H), 7.21 (dddd, J = 8.4, 7.0, 4.8, 1.8, 3H), 6.45 (s, 3H), 4.07 (s, 2H), 3.73 (ddd, J = 12.2, 11.1, 5.5, 2H), 3.02 (d, J = 5.2, 1H), 2.49 (d, J = 4.3, 1H), 2.36 (dt, J = 11.8, 4.5, 3H), 2.18 (dd, J = 12.3, 5.2, 1H), 2.00 (m, 1H), 1.79 (ddd, J = 13.9, 9.3, 4.4, 2H), 1.49 (m, 5H), 1.18 (m, 1H), 0.78 (d, J = 13.3, 1H).
259	[(2,6-difluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387.1	δ 8.79 (dd, J = 5.6, 1.3, 1H), 8.36 (m, 1H), 8.20 (s, 4H), 7.86 (d, J = 8.2, 1H), 7.79 (ddd, J = 7.6, 5.6, 1.0, 1H), 7.50 (tt, J = 8.5, 6.6, 1H), 7.04 (m, 2H), 4.13 (s, 2H), 3.72 (m, 2H), 3.05 (td, J = 12.0, 5.1, 1H), 2.52 (td, J = 12.1, 4.2, 1H), 2.37 (m, 3H), 2.19 (m, 1H), 2.04 (d, J = 14.2, 1H), 1.81 (m, 2H), 1.53 (m, 5H), 1.18 (m, 1H), 0.79 (dt, J = 12.8, 8.8, 1H).
260	[(3,4-difluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387.1	δ 8.84 (s, 1H), 8.79 (dd, J = 5.6, 1.3, 1H), 8.41 (td, J = 8.0, 1.6, 1H), 8.25 (s, 1H), 7.86 (ddd, J = 13.3, 7.6, 7.0, 2H), 7.30 (m, 3H), 3.99 (d, J = 1.7, 2H), 3.73 (m, 2H), 2.96 (dd, J = 12.1, 7.4, 1H), 2.38 (m, 4H), 2.21 (m, 1H), 2.05 (d, J = 14.2, 1H), 1.82 (ddd, J = 12.6, 9.0, 4.2, 2H), 1.53 (m, 5H), 1.21 (m, 1H), 0.81 (dt, J = 12.9, 8.8, 1H).
261	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	387	δ 8.77 (d, J = 5.4, 1H), 8.43 (s, 1H), 8.35 (d, J = 7.7, 1H), 7.83 (m, 2H), 7.03 (m, 3H), 4.01 (s, 2H), 3.74 (ddd, J = 27.7, 13.8, 7.4, 2H), 2.96 (m, 1H), 2.39 (m, 4H), 2.23 (dd, J = 13.3, 5.1, 1H), 2.02 (m, 1H), 1.81 (m, 2H), 1.52 (m, 5H), 1.21 (dd, J = 9.4, 5.3, 1H), 0.80 (dt, J = 12.9, 8.9, 1H).
262	[(2,3-dimethoxyphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	411.1	δ 8.75 (dd, J = 5.4, 1.2, 1H), 8.21 (td, J = 8.0, 1.8, 1H), 7.88 (s, 2H), 7.75 (d, J = 8.2, 1H), 7.66 (ddd, J = 7.6, 5.4, 0.9, 1H), 7.08 (dd, J = 9.0, 5.6, 2H), 6.87 (dd, J = 6.2, 3.0, 1H), 4.05 (s, 2H), 3.86 (d, J = 6.3, 6H), 3.73 (ddd, J = 12.5, 11.1, 5.4, 2H), 2.97 (s, 1H), 2.45 (s, 1H), 2.35 (m, 3H), 2.14 (m, 1H), 1.78 (ddd, J = 14.2, 6.0, 3.9, 2H), 1.49 (m, 5H), 1.16 (m, 1H), 0.77 (d, J = 13.3, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
263	[(3,4-dimethoxyphenyl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	411.1	δ 8.73 (dd, J = 5.5, 1.2, 1H), 8.24 (td, J = 8.0, 1.7, 1H), 8.00 (s, 1H), 7.78 (d, J = 8.2, 1H), 7.69 (ddd, J = 7.5, 5.5, 0.8, 1H), 6.96 (d, J = 1.0, 1H), 6.88 (d, J = 1.7, 2H), 6.55 (s, 3H), 3.93 (s, 2H), 3.78 (t, J = 7.5, 6H), 3.72 (m, 2H), 2.92 (s, 1H), 2.35 (m, 4H), 2.15 (m, 1H), 1.99 (d, J = 14.2, 1H), 1.79 (m, 2H), 1.49 (m, 5H), 1.18 (s, 1H), 0.79 (dd, J = 15.6, 6.6, 1H).
264	2-methoxy-4-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]phenol	397.1	δ 8.62 (dd, J = 5.0, 1.0, 1H), 7.94 (td, J = 7.9, 1.8, 1H), 7.57 (d, J = 8.1, 1H), 7.42 (m, 1H), 7.00 (s, 1H), 6.81 (d, J = 0.8, 2H), 3.93 (s, 2H), 3.84 (s, 3H), 3.70 (m, 3H), 2.93 (s, 1H), 2.36 (s, 3H), 2.17 (m, 1H), 1.90 (d, J = 13.7, 1H), 1.74 (m, 2H), 1.51 (s, 5H), 1.13 (m, 1H), 0.73 (dt, J = 13.2, 8.9, 1H).
265	[(5-fluoropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	370	δ 8.82 (s, 1H), 8.50 (dd, J = 34.5, 26.7, 3H), 7.93 (m, 2H), 7.74 (t, J = 9.7, 1H), 4.12 (d, J = 10.8, 2H), 3.76 (dd, J = 25.7, 11.8, 2H), 3.03 (d, J = 7.9, 1H), 2.39 (m, 5H), 2.09 (t, J = 13.0, 1H), 1.85 (d, J = 9.0, 2H), 1.60 (d, J = 44.8, 5H), 1.24 (s, 1H), 0.86 (d, J = 9.1, 1H).
266	[(5-bromopyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	430	δ 8.64 (m, 5H), 8.16 (s, 1H), 8.00 (d, J = 8.2, 1H), 7.95 (m, 1H), 4.10 (m, 2H), 3.76 (m, 2H), 3.02 (td, J = 12.4, 5.0, 1H), 2.49 (m, 2H), 2.29 (m, 3H), 2.12 (t, J = 10.2, 1H), 1.88 (ddd, J = 25.8, 12.8, 8.1, 2H), 1.57 (m, 5H), 1.26 (m, 1H), 0.86 (dt, J = 12.9, 8.9, 1H).
267	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	386	δ 8.80 (s, 1H), 8.63 (s, 1H), 8.49 (dd, J = 17.4, 10.6, 3H), 7.93 (m, 3H), 4.08 (s, 2H), 3.75 (dd, J = 29.6, 6.9, 2H), 2.99 (d, J = 11.6, 1H), 2.45 (m, 2H), 2.28 (m, 3H), 2.08 (d, J = 14.3, 1H), 1.85 (d, J = 7.5, 2H), 1.60 (m, 5H), 1.23 (s, 1H), 0.84 (d, J = 5.6, 1H).
268	[(5-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	382.1	δ 8.81 (d, J = 5.5, 1H), 8.48 (m, 3H), 7.95 (m, 3H), 4.22 (d, J = 13.4, 2H), 3.98 (s, 3H), 3.75 (ddd, J = 19.2, 12.7, 9.3, 2H), 3.04 (td, J = 11.6, 4.8, 1H), 2.42 (m, 7H), 2.09 (d, J = 14.3, 1H), 1.88 (m, 2H), 1.57 (m, 6H), 1.26 (d, J = 10.9, 1H), 0.85 (dt, J = 12.4, 8.7, 1H).
269	5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-3-carbonitrile	377.1	δ 8.96 (d, J = 15.0, 1H), 8.83 (t, J = 10.5, 2H), 8.53 (dt, J = 15.9, 8.0, 2H), 8.23 (d, J = 15.0, 1H), 7.97 (ddd, J = 13.4, 11.9, 7.5, 2H), 4.13 (m, 2H), 3.77 (m, 2H), 3.02 (m, 1H), 2.50 (ddd, J = 26.3, 14.4, 3.7, 2H), 2.31 (m, 3H), 2.13 (dd, J = 19.3, 11.3, 1H), 1.88 (ddd, J = 17.2, 11.0, 7.0, 2H), 1.58 (m, 5H), 1.27 (m, 1H), 0.85 (dt, J = 12.8, 8.7, 1H).

(continued)

Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
270	[(5-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	366	δ 8.71 (dd, J = 50.6, 19.3, 3H), 8.37 (m, 2H), 7.85 (m, 2H), 4.20 (d, J = 13.3, 2H), 3.74 (ddd, J = 11.9, 11.1, 5.6, 2H), 3.02 (m, 1H), 2.44 (m, 7H), 2.25 (dd, J = 12.5, 5.0, 1H), 1.84 (m, 2H), 1.57 (tdd, J = 24.6, 15.7, 8.5, 5H), 1.22 (d, J = 9.3, 1H), 0.83 (m, 1H).
271	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyridin-3-yl]methyl})amine	420.1	δ 8.96 (s, 1H), 8.81 (m, 2H), 8.45 (td, J = 8.1, 1.6, 2H), 8.21 (s, 1H), 7.90 (m, 2H), 4.16 (m, 2H), 3.77 (dtd, J = 12.7, 9.5, 5.3, 2H), 3.04 (td, J = 12.2, 5.1, 1H), 2.50 (m, 2H), 2.31 (ddd, J = 21.7, 14.1, 7.0, 3H), 2.12 (d, J = 12.6, 1H), 1.87 (ddd, J = 20.7, 12.7, 7.7, 2H), 1.58 (m, 5H), 1.26 (m, 1H), 0.85 (m, 1H).
272	{[6-chloro-5-(trifluoromethyl)pyridin-3-yl]methyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	454.1	δ 8.70 (m, 1H), 8.60 (d, J = 2.1, 1H), 8.28 (d, J = 2.2, 1H), 8.16 (td, J = 7.9, 1.8, 1H), 7.73 (d, J = 8.2, 1H), 7.62 (ddd, J = 7.6, 5.3, 1.0, 1H), 4.12 (m, 2H), 3.73 (m, 2H), 3.18 (brs, 1H), 2.99 (td, J = 12.0, 5.1, 2H), 2.49 (td, J = 12.0, 4.4, 1H), 2.35 (dd, J = 14.1, 1.9, 3H), 2.13 (ddd, J = 14.2, 12.1, 5.2, 1H), 1.79 (dd, J = 5.6, 3.7, 2H), 1.62 (dd, J = 7.8, 2.8, 1H), 1.51 (dd, J = 7.9, 4.1, 4H), 1.18 (m, 1H), 0.78 (dt, J = 13.2, 8.9, 1H).
273	{[2-fluoro-5-(trifluoromethyl)pyridin-3-yl]methyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	438.1	δ 8.73 (dd, J = 5.3, 1.2, 1H), 8.62 (s, 1H), 8.35 (dd, J = 8.5, 2.3, 1H), 8.22 (td, J = 8.0, 1.6, 1H), 7.78 (d, J = 8.2, 1H), 7.67 (dd, J = 6.9, 5.8, 1H), 4.25 (brs, 1H), 4.13 (m, 2H), 3.74 (ddd, J = 12.3, 11.0, 5.5, 2H), 3.05 (td, J = 11.9, 5.1, 1H), 2.54 (td, J = 12.0, 4.4, 1H), 2.35 (dt, J = 9.7, 5.3, 3H), 2.16 (ddd, J = 9.9, 8.8, 3.8, 1H), 2.01 (d, J = 14.1, 1H), 1.80 (m, 2H), 1.62 (m, 1H), 1.49 (m, 4H), 1.19 (m, 1H), 0.79 (dt, J = 13.1, 8.8, 1H).
274	{[6-fluoro-5-(trifluoromethyl)pyridin-3-yl]methyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	438.1	δ 8.58 (d, J = 4.0, 1H), 8.45 (s, 1H), 8.35 (d, J = 9.0, 1H), 7.85 (m, 1H), 7.51 (d, J = 8.1, 1H), 7.33 (dd, J = 7.4, 4.9, 1H), 4.12 (m, 2H), 3.70 (dd, J =
			8.8, 2.9, 2H), 2.98 (m, 2H), 2.47 (dd, J = 12.1, 7.4, 2H), 2.38 (t, J = 11.7, 2H), 2.18 (dd, J = 12.9, 4.8, 1H), 1.90 (d, J = 13.7, 1H), 1.70 (m, 2H), 1.60 (m, 1H), 1.50 (dt, J = 39.4, 20.7, 4H), 1.11 (m, 1H), 0.73 (dt, J = 13.5, 9.1, 1H).
275	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[3-(trifluoromethyl)pyridin-2-yl]methyl})amine	420.1	δ 9.29 (brs, 1H), 8.90 (s, 1H), 8.86 (d, J = 5.1, 1H), 8.80 (dd, J = 5.7, 1.2, 1H), 8.49 (td, J = 8.0, 1.7, 1H), 7.98 (d, J = 8.2, 1H), 7.91 (ddd, J = 7.6, 5.7, 1.0, 1H), 7.74 (d, J = 5.1, 1H), 4.27 (m, 2H), 3.81 (dt, J = 12.8, 4.6, 1H), 3.72 (m, 1H), 3.13 (td, J = 12.1, 5.1, 1H), 2.60 (td, J = 12.3, 4.1, 1H), 2.49 (m, 1H), 2.33 (m, 3H), 2.10 (d, J = 14.3, 1H), 1.85 (m, 2H), 1.65 (m, 1H), 1.52 (m, 4H), 1.25 (m, 1H), 0.84 (dt, J = 12.8, 8.8, 1H).

(continued)






Table 1: Compounds with chemical name and characterization data

Compound.	Name	MS (m/z) [M+H] ⁺	¹ H NMR
276	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	420.1	δ 8.81 (dd, J = 5.5, 1.2, 1H), 8.75 (d, J = 4.4, 1H), 8.32 (td, J = 8.0, 1.7, 1H), 8.16 (dd, J = 8.0, 0.7, 1H), 7.86 (d, J = 8.2, 1H), 7.77 (odd, J = 7.6, 5.5, 1.0, 1H), 7.59 (dd, J = 7.5, 5.0, 1H), 4.40 (m, 2H), 3.75 (m, 2H), 3.13 (td, J = 12.0, 5.3, 1H), 2.66 (td, J = 12.1, 4.5, 1H), 2.49 (ddd, J = 13.7, 11.9, 4.5, 1H), 2.41 (m, 1H), 2.32 (m, 2H), 2.07 (d, J = 14.0, 1H), 1.85 (ddd, J = 9.3, 7.7, 4.5, 2H), 1.64 (m, 1H), 1.51 (m, 4H), 1.22 (m, 1H), 0.82 (dt, J = 13.1, 8.9, 1H).
277	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-2-yl]methyl}amine	420.1	δ 8.80 (dd, J = 5.5, 1.3, 1H), 8.75 (d, J = 5.0, 1H), 8.31 (td, J = 8.0, 1.7, 1H), 7.84 (d, J = 8.2, 1H), 7.75 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.65 (M, 2H), 4.31 (m, 2H), 3.74 (m, 2H), 3.09 (td, J = 12.0, 5.2, 1H), 2.60 (td, J = 12.1, 4.4, 1H), 2.36 (m, 4H), 2.05 (d, J = 14.1, 1H), 1.82 (m, 2H), 1.64 (m, 1H), 1.50 (m, 4H), 1.20 (m, 1H), 0.81 (dt, J = 12.8, 8.8, 1H).
500	[(4-chlorophenyl)methyl]{2-[4-(4-methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl}amine		
501	[(3,4-dimethoxyphenyl)methyl][2-(2,2-dimethyl-4-phenyloxan-4-yl)ethyl]amine		
502	2-[(2-[2-ethyl-2-methyl-4-(4-methylphenyl)oxan-4-yl]ethyl)amino)methyl]phenol		
503	[2-(2,2-dimethyl-4-phenyloxan-4-yl)ethyl][(2-fluorophenyl)methyl]amine		
504	4-[(2-[4-(2-methoxyphenyl)-2,2-dimethyloxan-4-yl]ethyl)amino)methyl]-N,N-dimethylaniline		
505	2-[(2-[2-ethyl-4-(4-fluorophenyl)-2-methyloxan-4-yl]ethyl)amino)methyl]phenol		







Example 13: Opioid Receptor Ligands

[0206] The following compounds in Table 2 can also be prepared according to the procedures described above from appropriate starting materials and appropriate reagents and would be expected to also have similar properties and therapeutic effects as other compounds described herein. In addition to the specific structure shown the other isomers or enantiomers are included with the description herein. Compounds that have been made lists NMR data and prophetic examples do not list NMR data.






Table 2: Examples with chemical name and/or characterization data

Compound	Name	Structure and/or NMR Spectrum
506.	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(pyrimidin-5-ylmethyl)amine	MS: 353.2 ¹H NMR (400 MHz, CD₃CN) δ 9.16 (s, 1H), 8.78 (s, 2H), 8.70 (dd, J = 5.3, 1.1, 1H), 8.16 (td, J = 8.0, 1.8, 1H), 7.74 (d, J = 8.2, 1H), 7.62 (ddd, J = 7.6, 5.4, 0.9, 1H), 4.27 (brs, 1H), 4.04 (t, J = 7.7, 2H), 3.73 (m, 2H), 3.01 (td, J = 12.0, 5.1, 1H), 2.50 (td, J = 12.0, 4.4, 1H), 2.33 (m, 3H), 2.12 (ddd, J = 19.0, 11.7, 5.2, 1H), 1.99 (d, J = 10.1, 1H), 1.78 (m, 2H), 1.61 (m, 1H), 1.48 (m, 4H), 1.17 (m, 1H), 0.78 (dt, J = 13.1, 8.9, 1H). 
507.	[(2-methylpyrimidin-5-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
508.	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({2-(trifluoromethyl)pyrimidin-5-yl)methyl}amine	
509.	[(2-methoxypyrimidin-5-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 383.3 ¹H NMR (400 MHz, CD₃CN) δ 8.69 (dd, J = 5.2, 1.1, 1H), 8.54 (s, 2H), 8.10 (td, J = 7.9, 1.7, 1H), 7.69 (d, J = 8.1, 1H), 7.56 (dd, J = 6.7, 5.3, 1H), 3.98 (s, 5H), 3.71 (m, 3H), 3.50 (brs, 1H), 2.98 (td, J = 12.0, 5.0, 1H), 2.47 (td, J = 12.0, 4.3, 1H), 2.37 (m, 2H), 2.27 (m, 1H), 2.10 (m, 1H), 1.77 (m, 2H), 1.62 (m, 1H), 1.47 (dddd, J = 14.1, 12.4, 8.4, 4.9, 4H), 1.17 (m, 1H), 0.77 (dt, J = 13.1, 8.9, 1H). 
510.	(pyridazin-4-ylmethyl)({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	




(continued)

Compound	Name	Structure and/or NMR Spectrum
511.	[(6-methylpyridazin-4-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
512.	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[6-(trifluoromethyl)pyridazin-4-yl]methyl})amine	
513.	[(6-methoxypyridazin-4-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
514.	(pyrazin-2-ylmethyl)({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 353.3 ¹H NMR (400 MHz, CD₃CN) δ 8.74 (dd, J = 5.3, 1.1, 1H), 8.60 (d, J = 1.6, 2H), 8.55 (m, 1H), 8.16 (td, J = 7.9, 1.7, 1H), 7.73 (d, J = 8.2, 1H), 7.61 (ddd, J = 7.5, 5.3, 0.8, 1H), 7.13 (brs, 1H), 4.25 (m, 2H), 3.73 (m, 2H), 3.09 (td, J = 11.8, 5.4, 1H), 2.61 (td, J = 11.9, 4.6, 1H), 2.37 (m, 3H), 2.18 (ddd, J = 13.7, 11.6, 5.5, 1H), 1.99 (m, 1H), 1.77 (dd, J = 9.6, 4.4, 2H), 1.62 (m, 1H), 1.48 (m, 4H), 1.18 (m, 1H), 0.79 (dt, J = 13.1, 8.9, 1H). 
515.	[(6-methylpyrazin-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
516.	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[6-(trifluoromethyl)pyrazin-2-yl]methyl})amine	





(continued)

Compound	Name	Structure and/or NMR Spectrum
517.	[(6-methoxypyrazin-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
518.	[(5-methylpyrazin-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
519.	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyrazin-2-yl]methyl})amine	
520.	[(5-methoxypyrazin-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
521.	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(quinolin-3-ylmethyl)amine	MS: 402.3 ¹H NMR (400 MHz, CD₃CN) δ 9.90 (brs, 1H), 9.15 (d, J = 1.7, 1H), 8.89 (s, 1H), 8.77 (dd, J = 5.6, 1.3, 1H), 8.40 (td, J = 8.0, 1.6, 1H), 8.30 (d, J = 8.6, 1H), 8.16 (d, J = 8.2, 1H), 8.08 (ddd, J = 8.5, 7.0, 1.3, 1H), 7.90 (m, 2H), 7.81 (m, 1H), 4.36 (m, 2H), 3.74 (m, 2H), 3.06 (td, J = 12.0, 5.1, 1H), 2.57 (td, J = 12.2, 4.1, 1H), 2.45 (m, 1H), 2.29 (m, 3H), 2.08 (m, 1H), 1.98 (d, J = 2.5, 1H), 1.83 (m, 2H), 1.64 (ddd, J = 11.6, 8.7, 3.4, 1H), 1.50 (m, 4H), 1.23 (ddd, J = 10.4, 4.4, 2.4, 1H), 0.82 (dt, J = 12.9, 8.8, 1H). 








(continued)

Compound	Name	Structure and/or NMR Spectrum
522.	(1H-pyrazol-3-ylmethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 341.2 ¹H NMR (400 MHz, CD₃CN) δ 8.76 (dd, J = 5.5, 1.2, 1H), 8.28 (td, J = 8.0, 1.7, 1H), 7.80 (d, J = 8.2, 1H), 7.73 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.61 (d, J = 2.3, 1H), 6.32 (d, J = 2.3, 1H), 5.78 (brs, 1H), 4.09 (m, 2H), 3.72 (m, 2H), 2.98 (td, J = 12.0, 5.2, 1H), 2.47 (td, J = 12.1, 4.3, 1H), 2.36 (m, 3H), 2.16 (m, 1H), 2.02 (d, J = 14.2, 1H), 1.79 (m, 2H), 1.62 (m, 1H), 1.49 (m, 4H), 1.19 (m, 1H), 0.79 (dt, J = 12.9, 8.8, 1H). 
523.	[(1-methyl-1H-pyrazol-3-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 355.3 ¹H NMR (400 MHz, CD₃CN) δ 8.98 (brs, 1H), 8.73 (dd, J = 5.3, 1.1, 1H), 8.73 (dd, J = 5.3, 1.1, 1H), 8.16 (m, 2H), 7.72 (d, J = 8.2, 1H), 7.72 (d, J = 8.2, 1H), 7.62 (ddd, J = 7.5, 5.4, 0.8, 1H), 7.62 (ddd, J = 7.5, 5.4, 0.8, 1H), 7.47 (d, J = 2.2, 1H), 7.47 (d, J = 2.2, 1H), 6.25 (d, J = 2.2, 1H), 6.25 (d, J = 2.2, 1H), 4.02 (m, 2H), 3.80 (s, 3H), 3.72 (m, 2H), 2.98 (td, J = 11.8, 5.2, 1H), 2.48 (td, J = 11.9, 4.2, 1H), 2.33 (m, 3H), 2.12 (ddd, J = 13.5, 11.9, 5.4, 1H), 1.99 (m, 1H), 1.77 (m, 2H), 1.62 (m, 1H), 1.48 (m, 4H), 1.17 (m, 1H), 0.78 (dt, J = 13.1, 8.9, 1H). 
524.	[(5-methyl-1H-pyrazol-3-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 355.3 ¹H NMR (400 MHz, CD₃CN) δ 8.78 (dd, J = 5.5, 1.2, 1H), 8.34 (td, J = 8.0, 1.7, 1H), 7.79 (m, 2H), 6.07 (s, 1H), 5.95 (brs, 1H), 4.02 (m, 2H), 3.72 (m, 2H), 2.97 (td, J = 12.0, 5.1, 1H), 2.44 (ddd, J = 12.1, 10.0, 4.2, 1H), 2.34 (m, 3H), 2.26 (s, 3H), 2.18 (td, J = 13.1, 5.2, 1H), 2.03 (d, J = 14.2, 1H), 1.81 (ddd, J = 8.7, 7.4, 3.8, 2H), 1.63 (ddd, J = 14.6, 10.4, 4.6, 1H), 1.49 (m, 4H), 1.20 (m, 1H), 0.81 (dt, J = 12.9, 8.9, 1H). 





(continued)

Compound	Name	Structure and/or NMR Spectrum
525.	[[1,5-dimethyl-1H-pyrazol-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 369.3 ¹H NMR (400 MHz, CD₃CN) δ 12.13 (brs, 1H), 8.77 (dd, J = 5.4, 1.2, 1H), 8.28 (td, J = 8.0, 1.7, 1H), 8.00 (brs, 1H), 7.74 (m, 2H), 6.03 (s, 1H), 3.96 (m, 2H), 3.73 (m, 5H), 2.96 (td, J = 12.0, 5.2, 1H), 2.47 (td, J = 12.1, 4.2, 1H), 2.36 (m, 3H), 2.17 (m, 4H), 2.00 (m, 1H), 1.79 (m, 2H), 1.63 (ddd, J = 8.4, 7.6, 3.3, 1H), 1.50 (m, 4H), 1.19 (ddd, J = 10.1, 6.6, 1.8, 1H), 0.81 (dt, J = 12.9, 8.9, 1H). 
526.	(1H-pyrazol-4-ylmethyl)({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 341.2 ¹H NMR (400 MHz, CD₃CN) δ 8.73 (dd, J = 5.3, 1.1, 1H), 8.21 (td, J = 8.0, 1.7, 2H), 7.75 (d, J = 8.2, 1H), 7.66 (m, 3H), 7.56 (s, 1H), 3.96 (s, 2H), 3.73 (m, 2H), 2.91 (m, 1H), 2.32 (m, 4H), 2.08 (m, 1H), 1.99 (m, 1H), 1.78 (m, 2H), 1.62 (m, 1H), 1.49 (m, 4H), 1.19 (m, 1H), 0.78 (dt, J = 13.1, 8.8, 1H). 
527.	[(1-methyl-1H-pyrazol-4-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 355.2 ¹H NMR (400 MHz, CD₃CN) δ 8.78 (dd, J = 5.5, 1.3, 1H), 8.32 (td, J = 8.0, 1.7, 1H), 7.84 (d, J = 8.2, 1H), 7.77 (ddd, J = 7.6, 5.5, 0.9, 1H), 7.71 (brs, 1H), 7.55 (s, 1H), 7.43 (s, 1H), 3.91 (s, 2H), 3.81 (d, J = 11.7, 3H), 3.73 (m, 2H), 2.90 (dt, J = 11.7, 5.8, 1H), 2.35 (m, 4H), 2.14 (ddd, J = 10.8, 10.2, 5.2, 1H), 2.03 (d, J = 14.2, 1H), 1.80 (m, 2H), 1.62 (tdd, J = 8.7, 6.8, 2.7, 1H), 1.49 (m, 4H), 1.20 (m, 1H), 0.80 (dt, J = 12.9, 8.8, 1H). 
528.	[(5-methyl-1H-pyrazol-4-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	






(continued)

Compound	Name	Structure and/or NMR Spectrum
529.	[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
530.	[(5,6-difluoropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
531.	[(5-chloro-6-fluoropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
532.	[(5-bromo-6-fluoropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
533.	[(6-fluoro-5-iodopyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
534.	[(6-fluoro-5-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 384.3 ¹H NMR (400 MHz, CD₃CN) δ 8.73 (d, J = 4.4, 1H), 8.36 (s, 1H), 8.20 (d, J = 7.8, 1H), 8.01 (s, 1H), 7.77 (t, J = 7.1, 2H), 7.66 (m, 1H), 4.01 (s, 2H), 3.73 (m, 2H), 2.98 (dd, J = 11.6, 6.9, 1H), 2.36 (m, 5H), 2.26 (s, 3H), 2.16 (dd, J = 13.2, 5.1, 2H), 1.80 (m, 2H), 1.51 (m, 6H), 1.20 (dd, J = 8.7, 4.7, 1H), 0.79 (d, J = 13.3, 1H). 
535.	[(6-fluoro-5-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	

(continued)

Compound	Name	Structure and/or NMR Spectrum
536.	2-fluoro-5-[[{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amino)methyl]pyridine-3-carbonitrile	
537.	[(6-chloro-5-fluoropyridin-3-yl)methyl][{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 404.2 ¹H NMR (400 MHz, CD₃CN) δ 8.68 (dd, J = 5.2, 1.1, 1H), 8.24 (d, J = 1.9, 1H), 8.10 (td, J = 7.9, 1.8, 1H), 7.78 (dd, J = 9.0, 2.0, 1H), 7.69 (d, J = 8.2, 1H), 7.56 (ddd, J = 7.5, 5.3, 0.9, 1H), 4.75 (brs, 1H), 4.07 (m, 2H), 3.72 (m, 2H), 2.99 (td, J = 11.9, 5.2, 1H), 2.48 (td, J = 12.0, 4.5, 1H), 2.32 (m, 3H), 2.11 (m, 1H), 1.77 (m, 2H), 1.62 (m, 1H), 1.49 (m, 4H), 1.17 (m, 1H), 0.77 (dt, J = 13.1, 8.9, 1H). 
538.	[(5,6-dichloropyridin-3-yl)methyl][{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 422.2 ¹H NMR (400 MHz, CD₃CN) δ 8.51 (m, 1H), 8.33 (d, J = 2.1, 1H), 8.12 (d, J = 2.1, 1H), 7.76 (t, J = 7.9, 1H), 7.49 (d, J = 8.1, 1H), 7.23 (dd, J = 7.4, 4.9, 1H), 4.26 (d, J = 1.5, 2H), 3.57 (dd, J = 7.7, 3.0, 2H), 3.09 (td, J = 12.2, 4.6, 1H), 2.55 (td, J = 12.1, 4.6, 1H), 2.27 (dddd, J = 25.5, 17.3, 14.3, 3.4, 4H), 1.77 (m, 1H), 1.59 (m, 2H), 1.34 (m, 6H), 0.98 (dd, J = 11.4, 5.0, 1H), 0.60 (dt, J = 13.4, 9.0, 1H). 
539.	[(5-bromo-6-chloropyridin-3-yl)methyl][{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 466.1 ¹H NMR (400 MHz, CD₃CN) δ 8.64 (d, J = 5.1, 1H), 8.36 (d, J = 2.1, 1H), 8.24 (d, J = 2.1, 1H), 8.02 (m, 1H), 7.69 (d, J = 8.0, 1H), 7.47 (m, 1H), 3.59 (m, 2H), 3.17 (d, J = 4.7, 1H), 2.63 (d, J = 4.5, 1H), 2.34 (m, 4H), 2.12 (d, J = 4.8, 1H), 1.85 (d, J = 13.8, 1H), 1.66 (m, 2H), 1.35 (m, 6H), 1.02 (m, 1H), 0.66 (s, 








(continued)

Compound	Name	Structure and/or NMR Spectrum
540.	[(6-chloro-5-iodopyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
541.	[(6-chloro-5-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 400.2 ¹H NMR (400 MHz, CD₃CN) δ 8.70 (dd, J = 5.3, 1.2, 1H), 8.20 (t, J = 2.4, 1H), 8.17 (dd, J = 7.9, 1.7, 1H), 7.74 (t, J = 5.2, 2H), 7.63 (ddd, J = 7.5, 5.3, 0.9, 1H), 5.11 (s, 1H), 4.01 (m, 2H), 3.73 (m, 2H), 2.98 (td, J = 11.9, 5.1, 1H), 2.46 (td, J = 12.0, 4.2, 1H), 2.33 (m, 6H), 2.12 (ddd, J = 14.7, 10.5, 5.3, 1H), 1.99 (d, J = 6.9, 1H), 1.78 (m, 2H), 1.62 (m, 1H), 1.48 (m, 4H), 1.18 (m, 1H), 0.78 (dt, J = 13.1, 8.9, 1H). 
542.	[(6-chloro-5-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 416.2 ¹H NMR (400 MHz, CD₃CN) δ 8.73 (dd, J = 5.4, 1.2, 1H), 8.26 (td, J = 7.9, 1.6, 1H), 7.93 (d, J = 1.9, 1H), 7.80 (d, J = 8.2, 1H), 7.70 (dd, J = 7.1, 5.9, 1H), 7.52 (d, J = 1.9, 1H), 5.05 (brs, 1H), 4.04 (m, 2H), 3.90 (s, 3H), 3.74 (m, 2H), 2.98 (td, J = 12.0, 5.1, 1H), 2.40 (dddd, J = 19.5, 12.3, 9.6, 4.7, 3H), 2.16 (m, 1H), 1.99 (m, 1H), 1.80 (m, 2H), 1.63 (ddd, J = 14.5, 7.2, 3.0, 1H), 1.49 (m, 4H), 1.20 (m, 1H), 0.80 (dt, J = 12.9, 8.8, 1H). 
543.	2-chloro-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-3-carbonitrile	
544.	3-fluoro-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2-carbonitrile	

(continued)

Compound	Name	Structure and/or NMR Spectrum
545.	3-chloro-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2-carbonitrile	
546.	3-bromo-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2-carbonitrile	
547.	3-iodo-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2-carbonitrile	
548.	3-methyl-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2-carbonitrile	
549.	3-methoxy-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2-carbonitrile	
550.	5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-2,3-dicarbonitrile	
551.	5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]-3-(trifluoromethyl)pyridine-2-carbonitrile	
552.	{[5-fluoro-6-(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine}	





(continued)

Compound	Name	Structure and/or NMR Spectrum
553.	{[5-chloro-6-(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	
554.	{[5-bromo-6-(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	MS: 498.1 ¹H NMR (400 MHz, CD₃CN) δ 8.74 (dd, J = 5.4, 1.2, 1 H), 8.64 (d, J = 1.6, 1 H), 8.32 (d, J = 1.2, 1 H), 8.26 (td, J = 8.0, 1.6, 1H), 7.81 (d, J = 8.2, 1H), 7.71 (dd, J = 7.1, 6.0, 1H), 4.12 (m, 3H), 3.74 (m, 3H), 3.00 (td, J = 12.0, 5.1, 1H), 2.49 (td, J = 12.1, 4.2, 1H), 2.37 (ddd, J = 14.0, 11.9, 5.0, 3H), 2.16 (m, 1H), 2.02 (m, 1H), 1.81 (m, 2H), 1.63 (ddd, J = 14.4, 8.7, 4.7, 1H), 1.49 (m, 4H), 1.21 (m, 1H), 0.80 (dt, J = 13.0, 8.9, 1H). 
555.	{[5-iodo-6-(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	
556.	{[5-methyl-6-(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	
557.	{[5-methoxy-6-(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	
558.	5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]-2-(trifluoromethyl)pyridine-3-carbonitrile	
559.	{[5,6-bis(trifluoromethyl)pyridin-3-yl]methyl}{(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	





(continued)

Compound	Name	Structure and/or NMR Spectrum
560.	[(5-fluoro-6-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
561.	[(5-chloro-6-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
562.	[(5-bromo-6-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
563.	[(5-iodo-6-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
564.	[(5,6-dimethylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
565.	[(5-methoxy-6-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
566.	2-methyl-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-3-carbonitrile	
567.	{[6-methyl-5-(trifluoromethyl)pyridin-3-yl]methyl}({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	

(continued)

Compound	Name	Structure and/or NMR Spectrum
568.	[[5-fluoro-6-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 400.3 ¹H NMR (400 MHz, CD₃CN) δ 8.63 (dd, J = 5.3, 1.2, 1H), 8.25 (s, 1H), 8.12 (td, J = 8.0, 1.6, 1H), 7.83 (d, J = 1.9, 1H), 7.67 (d, J = 8.2, 1H), 7.57 (dd, J = 6.8, 5.7, 1H), 7.44 (dd, J = 11.1, 2.0, 1H), 3.88 (d, J = 6.7, 5H), 3.62 (m, 2H), 2.86 (dd, J = 11.5, 7.1, 1H), 2.26 (m, 4H), 2.05 (dd, J = 12.7, 5.0, 1H), 1.69 (ddd, J = 9.5, 8.0, 4.4, 2H), 1.69 (ddd, J = 9.5, 8.0, 4.4, 2H), 1.39 (m, 5H), 0.68 (d, J = 13.3, 1H). 
569.	[[5-chloro-6-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 416.2 ¹H NMR (400 MHz, CD₃CN) δ 8.65 (d, J = 5.4, 1H), 8.21 (s, 1H), 8.18 (d, J = 8.0, 1H), 7.96 (d, J = 2.1, 1H), 7.72 (m, 2H), 7.63 (t, J = 6.4, 1H), 3.87 (m, 5H), 3.62 (m, 2H), 2.85 (dd, J = 11.5, 7.2, 1H), 2.27 (m, 4H), 2.07 (d, J = 4.9, 1H), 1.91 (d, J = 14.1, 1H), 1.69 (m, 2H), 1.39 (m, 5H), 1.10 (m, 1H), 0.69 (d, J = 13.2, 1H). 
570.	[[5-bromo-6-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 460.2 ¹H NMR (400 MHz, CDCl₃) δ 8.79 (dd, J = 5.7, 1.3, 1H), 8.43 (td, J = 8.0, 1.7, 1H), 8.11 (d, J = 2.1, 1H), 7.98 (d, J = 2.1, 1H), 7.92 (d, J = 8.2, 1H), 7.86 (ddd, J = 7.6, 5.7, 1.0, 1H), 5.63 (brs, 1H), 3.97 (m, 5H), 3.75 (m, 2H), 2.96 (m, 1H), 2.42 (dq, J = 12.2, 4.1, 2H), 2.33 (d, J = 14.1, 2H), 2.21 (m, 1H), 2.06 (d, J = 14.2, 1H), 1.83 (m, 2H), 1.64 (ddd, J = 19.4, 10.1, 4.4, 1H), 1.50 (m, 4H), 1.23 (m, 1H), 0.82 (dt, J = 12.9, 8.9, 1H). 
571.	[[5-iodo-6-methoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	





(continued)

Compound	Name	Structure and/or NMR Spectrum
572.	[(6-methoxy-5-methylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 396.3 ¹H NMR (400 MHz, CDCl₃) δ 8.78 (dd, J = 5.6, 1.3, 1H), 8.38 (td, J = 8.0, 1.7, 1H), 7.96 (d, J = 2.2, 1H), 7.88 (d, J = 8.2, 1H), 7.81 (ddd, J = 7.6, 5.6, 1.0, 1H), 7.49 (d, J = 1.5, 1H), 5.36 (brs, 1H), 3.93 (m, 5H), 3.74 (m, 2H), 2.95 (dd, J = 11.4, 7.7, 1H), 2.39 (m, 4H), 2.21 (dd, J = 13.2, 5.4, 1H), 2.14 (m, 3H), 2.05 (d, J = 14.2, 1H), 1.82 (m, 2H), 1.63 (m, 1H), 1.50 (m, 4H), 1.21 (ddd, J = 10.5, 6.1, 2.5, 1H), 0.81 (dt, J = 12.9, 8.8, 1H). 
573.	[(5,6-dimethoxypyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
574.	2-methoxy-5-[(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amino)methyl]pyridine-3-carbonitrile	
575.	[(6-methoxy-5-(trifluoromethyl)pyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	





Example 14: Opioid Receptor Ligands

[0207] The following compounds in Table 3 can also be prepared according to the procedures described above from appropriate starting materials and appropriate reagents and would be expected to also have similar properties and therapeutic effects as other compounds described herein. In addition to the specific structure shown the other isomers or enantiomers are included with the description herein. Compounds that have been made lists NMR data and prophetic examples do not list NMR data.





Table 3: Opioid Receptor Ligands

Compound	Name	Structure
576	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-phenyl-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
577	{2-[(9R)-9-phenyl-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyridin-3-yl]methyl})amine	
578	{2-[(9R)-9-phenyl-6-oxaspiro[4.5]decan-9-yl]ethyl}({[4-(trifluoromethyl)pyridin-3-yl]methyl})amine	
579	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-phenyl-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	





(continued)

Compound	Name	Structure
580	[(3-methylphenyl)methyl]({2-[(9R)-9-phenyl-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
581	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
582	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyridin-3-yl]methyl})amin	
583	{2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[4-(trifluoromethyl)pyridin-3-yl]methyl})amine	





(continued)

Compound	Name	Structure
584	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-(4-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
585	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-[4-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
586	{2-[(9R)-9-[4-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-3-yl]methyl}amine	
587	{2-[(9R)-9-[4-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	





(continued)

Compound	Name	Structure
588	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-[4-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
589	[(3-methylphenyl)methyl]({2-[(9R)-9-[4-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
590	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-3-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
591	{2-[(9R)-9-(pyridin-3-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyridin-3-yl]methyl})amine	





(continued)

Compound	Name	Structure
592	{2-[(9R)-9-(pyridin-3-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	
593	[(3,5-difluorophenyl)methyl]{2-[(9R)-9-(pyridin-3-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	
594	[(3-methylphenyl)methyl]{2-[(9R)-9-(pyridin-3-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	
595	[(5-chloropyridin-3-yl)methyl]{2-[(9R)-9-(pyridin-4-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	





(continued)

Compound	Name	Structure
596	{2-[(9R)-9-(pyridin-4-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-3-yl]methyl}amine	
597	{2-[(9R)-9-(pyridin-4-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	
598	[(3,5-difluorophenyl)methyl]{2-[(9R)-9-(pyridin-4-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	
599	[(3-methylphenyl)methyl]{2-[(9R)-9-(pyridin-4-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	





(continued)

Compound	Name	Structure
600	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-(3-methylphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
601	{2-[(9R)-9-(3-methylphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-3-yl]methyl}amine	
602	{2-[(9R)-9-(3-methylphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	
603	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-(3-methylphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	





(continued)

Compound	Name	Structure
604	{2-[(9R)-9-(3-methylphenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}[(3-methylphenyl)methyl]amine	
605	[[5-chloropyridin-3-yl)methyl]{(2-[(9R)-9-[3-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl))amine	
606	{2-[(9R)-9-[3-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-3-yl]methyl}amine	
607	{2-[(9R)-9-[3-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	





(continued)

Compound	Name	Structure
608	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-[3-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
609	[(3-methylphenyl)methyl]({2-[(9R)-9-[3-(trifluoromethoxy)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
610	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-[4-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
611	{2-[(9R)-9-[4-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyridin-3-yl]methyl})amine	





(continued)

Compound	Name	Structure
612	{2-[(9R)-9-[4-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	
613	[(3,5-difluorophenyl)methyl]{2-[(9R)-9-[4-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	
614	[(3-methylphenyl)methyl]{2-[(9R)-9-[4-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	
615	[(5-chloropyridin-3-yl)methyl]{2-[(9R)-9-(3-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	




(continued)

Compound	Name	Structure
616	{2-[(9R)-9-(3-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-3-yl]methyl}amine	
617	{2-[(9R)-9-(3-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[4-(trifluoromethyl)pyridin-3-yl]methyl}amine	
618	[(3,5-difluorophenyl)methyl]{[2-[(9R)-9-(3-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	
619	{2-[(9R)-9-(3-fluorophenyl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{(3-methylphenyl)methyl}amine	



(continued)

Compound	Name	Structure
620	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-[3-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
621	{2-[(9R)-9-[3-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}({[5-(trifluoromethyl)pyridin-3-yl]methyl})amine	
622	{2-[(9R)-9-[3-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl}({[4-(trifluoromethyl)pyridin-3-yl]methyl})amine	
623	[(3,5-difluorophenyl)methyl]({2-[(9R)-9-[3-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	



(continued)

Compound	Name	Structure
624	[(3-methylphenyl)methyl]({2-[(9R)-9-[3-(trifluoromethyl)phenyl]-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
625	[(5-chloropyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 400.2 ¹H NMR (400 MHz, CD₃CN) δ 8.77 (dd, J = 5.6, 1.3, 1H), 8.67 (d, J = 2.0, 1H), 8.53 (s, 1H), 8.41 (td, J = 8.0, 1.6, 1H), 7.93 (m, 2H), 7.85 (m, 1H), 4.18 (s, 2H), 3.76 (ddd, J = 12.4, 11.3, 5.5, 2H), 3.09 (d, J = 5.1, 1H), 2.65 (s, 3H), 2.55 (m, 2H), 2.33 (m, 3H), 2.08 (d, J = 14.2, 1H), 1.84 (m, 2H), 1.53 (m, 5H), 1.21 (m, 1H), 0.77 (d, J = 13.2, 1H). 
626	methyl({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}){[5-(trifluoromethyl)pyridin-3-yl]methyl}amine	MS: 434.3 ¹H NMR (400 MHz, CD₃CN) δ 8.99 (s, 1H), 8.84 (s, 1H), 8.77 (m, 1H), 8.40 (td, J = 8.0, 1.6, 1H), 8.23 (s, 1H), 7.91 (d, J = 8.2, 1H), 7.84 (dd, J = 6.9, 6.3, 1H), 4.26 (s, 2H), 3.77 (m, 2H), 3.11 (d, J = 4.8, 1H), 2.65 (s, 3H), 2.57 (ddd, J = 17.4, 12.8, 8.9, 2H), 2.34 (dd, J = 19.0, 9.6, 3H), 2.09 (d, J = 14.2, 1H), 1.86 (m, 2H), 1.54 (m, 5H), 1.20 (dd, J = 9.5, 3.8, 1H), 0.77 (dd, J = 9.0, 4.1, 1H). 



(continued)

Compound	Name	Structure
627	[(5-chloropyridin-3-yl)methyl-(2H ₂)] {2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 388.2 ¹H NMR (400 MHz, CD₃CN) δ 9.51 (s, 1H), 8.48 (d, J = 1.9, 1H), 8.45 (d, J = 2.3, 1H), 8.42 (ddd, J = 4.8, 1.8, 0.8, 1H), 8.06 (m, 1H), 7.64 (td, J = 7.8, 1.8, 1H), 7.33 (d, J = 8.1, 1H), 7.12 (ddd, J = 7.4, 4.8, 0.7, 1H), 3.57 (m, 2H), 2.74 (td, J = 12.0, 4.7, 1H), 2.21 (m, 4H), 1.96 (dt, J = 12.4, 6.1, 1H), 1.76 (d, J = 13.8, 1H), 1.63 (dd, J = 9.9, 5.9, 1H), 1.40 (m, 6H), 0.95 (m, 1H), 0.59 (m, 1H). 
628	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-3-yl]methyl-(2H ₂)}amine	MS: 422.3 ¹H NMR (400 MHz, CD₃CN) δ 9.44 (s, 1H), 8.82 (d, J = 1.9, 1H), 8.78 (d, J = 1.2, 1H), 8.40 (ddd, J = 4.8, 1.8, 0.9, 1H), 8.31 (m, 1H), 7.61 (m, 1H), 7.31 (m, 1H), 7.09 (ddd, J = 7.4, 4.8, 1.0, 1H), 3.57 (m, 2H), 2.74 (m, 1H), 2.24 (m, 3H), 2.10 (m, 1H), 1.95 (dd, J = 12.5, 4.7, 1H), 1.75 (d, J = 13.6, 1H), 1.44 (m, 7H), 0.96 (s, 1H), 0.59 (m, 1H). 



(continued)

Compound	Name	Structure
629	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[6-(trifluoromethyl)pyridin-2-yl]methyl}amine	MS: 420.2 ¹H NMR (400 MHz, CD₃CN) δ 8.70 (dd, J = 5.2, 1.1, 1H), 8.53 (brs, 1H), 8.11 (td, J = 7.9, 1.7, 1H), 8.05 (t, J = 7.9, 1H), 7.80 (d, J = 7.8, 1H), 7.70 (d, J = 8.2, 1H), 7.57 (ddd, J = 8.3, 7.5, 4.4, 2H), 6.58 (brs, 1H), 4.29 (m, 2H), 3.73 (m, 2H), 3.09 (td, J = 11.8, 5.2, 1H), 2.60 (td, J = 11.9, 4.8, 1H), 2.36 (m, 3H), 2.16 (m, 1H), 1.99 (m, 1H), 1.77 (ddd, J = 14.0, 9.0, 5.1, 2H), 1.62 (m, 1H), 1.48 (m, 4H), 1.16 (ddd, J = 8.5, 7.0, 3.5, 1H), 0.78 (dt, J = 13.1, 8.9, 1H). 
630	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}{[5-(trifluoromethyl)pyridin-2-yl]methyl}amine	MS: 420.2 ¹H NMR (400 MHz, CD₃CN) δ 8.86 (d, J = 0.8, 1H), 8.77 (dd, J = 5.4, 1.2, 1H), 8.20 (m, 1H), 8.11 (dd, J = 8.3, 1.9, 1H), 7.77 (d, J = 8.2, 1H), 7.66 (ddd, J = 7.6, 5.4, 0.9, 1H), 7.53 (d, J = 8.3, 1H), 4.31 (m, 2H), 3.73 (m, 2H), 3.09 (td, J = 11.9, 5.4, 1H), 2.60 (td, J = 11.9, 4.6, 1H), 2.39 (m, 3H), 2.21 (ddd, J = 13.6, 11.8, 5.4, 1H), 2.02 (d, J = 14.0, 1H), 1.80 (ddd, J = 9.5, 8.3, 4.6, 2H), 1.63 (m, 1H), 1.49 (qdd, J = 13.9, 8.5, 3.5, 4H), 1.19 (m, 1H), 0.80 (dt, J = 13.1, 8.8, 1H). 



(continued)

Compound	Name	Structure
631	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(pyridin-2-ylmethyl)amine	MS: 352.3 ¹H NMR (400 MHz, CD₃CN) δ 10.49 (s, 1H), 8.81 (dd, J = 5.5, 1.2, 1H), 8.55 (dd, J = 3.7, 0.8, 1H), 8.30 (td, J = 8.0, 1.7, 1H), 7.91 (td, J = 7.8, 1.7, 1H), 7.83 (d, J = 8.2, 1H), 7.75 (ddd, J = 7.6, 5.5, 1.0, 1H), 7.45 (dd, J = 11.3, 6.5, 2H), 4.24 (m, 2H), 3.73 (m, 2H), 3.06 (td, J = 12.0, 5.2, 1H), 2.57 (td, J = 12.1, 4.4, 1H), 2.39 (m, 3H), 2.24 (m, 1H), 2.04 (d, J = 14.0, 1H), 1.82 (m, 2H), 1.63 (m, 1H), 1.50 (m, 4H), 1.19 (m, 1H), 0.81 (dt, J = 12.9, 8.8, 1H). 
632	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(pyridin-3-ylmethyl)amine	MS: 352.3 ¹H NMR (400 MHz, CD₃CN) δ 8.81 (s, 1H), 8.74 (m, 2H), 8.32 (d, J = 8.1, 1H), 8.26 (td, J = 8.0, 1.7, 1H), 7.80 (m, 2H), 7.70 (m, 1H), 4.18 (m, 2H), 3.73 (m, 2H), 3.02 (td, J = 12.0, 5.1, 1H), 2.51 (td, J = 12.1, 4.3, 1H), 2.36 (m, 3H), 2.15 (m, 1H), 2.01 (d, J = 14.1, 1H), 1.80 (ddd, J = 9.8, 8.2, 4.7, 2H), 1.62 (m, 1H), 1.48 (m, 4H), 1.19 (m, 1H), 0.80 (dt, J = 13.0, 8.8, 1H). 



(continued)

Compound	Name	Structure
633	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}(pyridin-4-ylmethyl)amine	MS: 352.3 ¹H NMR (400 MHz, CD₃CN) δ 8.73 (m, 3H), 8.20 (td, J = 8.0, 1.7, 1H), 7.82 (d, J = 6.5, 2H), 7.76 (d, J = 8.2, 1H), 7.65 (m, 1H), 4.22 (m, 2H), 3.73 (m, 2H), 3.03 (td, J = 12.0, 5.1, 1H), 2.53 (td, J = 12.1, 4.4, 1H), 2.37 (m, 3H), 2.16 (m, 1H), 2.00 (d, J = 14.2, 1H), 1.79 (m, 2H), 1.63 (ddd, J = 12.2, 8.8, 4.0, 1H), 1.49 (m, 4H), 1.19 (m, 1H), 0.80 (dt, J = 13.1, 8.9, 1H). 
634	(1H-imidazol-4-ylmethyl){2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 341.2 ¹H NMR (400 MHz, CD₃CN) δ 8.75 (dd, J = 5.4, 1.2, 1H), 8.54 (d, J = 1.0, 1H), 8.22 (td, J = 8.0, 1.6, 1H), 7.77 (d, J = 8.2, 1H), 7.67 (dd, J = 6.8, 5.6, 1H), 7.47 (s, 1H), 4.18 (s, 2H), 3.72 (m, 2H), 2.92 (td, J = 12.1, 5.0, 1H), 2.38 (m, 4H), 2.13 (m, 1H), 2.00 (m, 1H), 1.79 (m, 2H), 1.63 (m, 1H), 1.48 (m, 4H), 1.19 (m, 1H), 0.82 (dt, J = 13.1, 8.9, 1H). 



(continued)

Compound	Name	Structure
635	[(2-methylpyridin-4-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 366.3 ¹H NMR (400 MHz, CD₃CN) δ 8.71 (d, J = 1.9, 1H), 8.65 (dd, J = 5.1, 1.0, 1H), 8.18 (dd, J = 8.2, 2.1, 1H), 8.02 (td, J = 7.9, 1.8, 1H), 7.64 (d, J = 8.1, 2H), 7.49 (dd, J = 6.7, 5.2, 1H), 4.13 (m, 2H), 3.71 (m, 2H), 3.00 (td, J = 11.8, 5.1, 1H), 2.71 (s, 3H), 2.50 (td, J = 11.9, 4.6, 1H), 2.37 (m, 2H), 2.23 (m, 1H), 2.06 (dd, J = 12.0, 5.1, 1H), 1.76 (ddt, J = 14.1, 9.4, 3.8, 3H), 1.61 (dd, J = 16.6, 9.7, 1H), 1.49 (m, 4H), 1.16 (d, J = 11.3, 1H), 0.76 (dt, J = 13.1, 8.9, 1H). 
636	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({[2-(trifluoromethyl)pyridin-4-yl]methyl})amine	MS: 420.2 ¹H NMR (400 MHz, CD₃CN) δ 8.75 (d, J = 5.1, 2H), 8.30 (td, J = 8.1, 1.4, 1H), 7.84 (m, 2H), 7.74 (m, 1H), 7.63 (d, J = 4.7, 1H), 4.13 (m, 2H), 3.73 (m, 2H), 3.01 (td, J = 11.9, 5.0, 1H), 2.45 (m, 4H), 2.22 (td, J = 13.0, 5.0, 1H), 2.03 (d, J = 14.1, 1H), 1.81 (m, 2H), 1.63 (m, 1H), 1.49 (m, 4H), 1.21 (dd, J = 9.4, 5.1, 1H), 0.81 (dt, J = 12.8, 8.8, 1H). 





(continued)

Compound	Name	Structure
637	[(6-chloropyridin-3-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 386.2 ¹H NMR (400 MHz, CD₃CN) δ 8.69 (m, 1H), 8.38 (m, 1H), 8.12 (m, 1H), 7.81 (m, 1H), 7.70 (m, 1H), 7.58 (m, 1H), 7.45 (m, 1H), 4.43 (s, 1H), 4.06 (d, J = 13.9, 2H), 3.72 (m, 2H), 2.98 (td, J = 11.9, 5.1, 1H), 2.48 (td, J = 12.0, 4.4, 1H), 2.32 (m, 3H), 2.10 (m, 1H), 1.98 (d, J = 2.4, 1H), 1.77 (m, 2H), 1.61 (ddd, J = 15.0, 8.2, 4.0, 1H), 1.48 (m, 4H), 1.16 (ddd, J = 8.7, 7.1, 4.1, 1H), 0.77 (dt, J = 13.2, 8.9, 1H). 
638	[(1-methyl-1H-imidazol-2-yl)methyl]{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}amine	MS: 355.3 ¹H NMR (400 MHz, CD₃CN) δ 8.71 (ddd, J = 5.3, 1.7, 0.6, 1H), 8.14 (td, J = 8.0, 1.8, 1H), 7.73 (d, J = 8.2, 1H), 7.60 (ddd, J = 7.6, 5.3, 1.0, 1H), 7.43 (d, J = 1.9, 1H), 7.35 (d, J = 1.9, 1H), 4.33 (s, 2H), 3.80 (m, 3H), 3.72 (ddt, J = 15.3, 9.3, 3.1, 2H), 3.03 (td, J = 12.0, 4.9, 1H), 2.59 (td, J = 12.0, 4.6, 1H), 2.36 (m, 3H), 2.15 (m, 1H), 1.99 (m, 1H), 1.80 (m, 2H), 1.63 (ddd, J = 14.4, 9.9, 5.5, 1H), 1.49 (m, 4H), 1.19 (m, 1H), 0.83 (dt, J = 13.1, 8.9, 1H). 

(continued)

Compound	Name	Structure
639	(naphthalen-2-ylmethyl)((2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	MS: 401.3 ¹H NMR (400 MHz, CD₃CN) δ 8.57 (dd, J = 5.0, 1.0, 1H), 7.90 (m, 5H), 7.59 (m, 2H), 7.54 (d, J = 8.1, 1H), 7.48 (dd, J = 8.5, 1.7, 1H), 7.34 (m, 1H), 4.19 (s, 2H), 3.69 (dt, J = 8.9, 5.1, 3H), 3.48 (brs, 1H), 3.02 (s, 1H), 2.52 (s, 1H), 2.33 (m, 2H), 2.19 (m, 1H), 2.02 (m, 1H), 1.89 (t, J = 9.4, 1H), 1.70 (dq, J = 9.2, 5.1, 2H), 1.59 (m, 1H), 1.44 (m, 4H), 1.10 (m, 1H), 0.69 (dt, J = 13.1, 8.8, 1H). 
640	[(6-bromo-5-fluoropyridin-3-yl)methyl][(2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl)amine	MS: 448.2 ¹H NMR (400 MHz, CD₃CN) δ 8.68 (dd, J = 5.2, 1.2, 1H), 8.24 (d, J = 1.5, 1H), 8.12 (td, J = 7.9, 1.7, 1H), 7.71 (m, 2H), 7.58 (dd, J = 7.1, 5.7, 1H), 4.88 (s, 1H), 4.08 (d, J = 14.0, 2H), 3.72 (m, 2H), 2.98 (td, J = 11.9, 5.1, 1H), 2.48 (td, J = 12.0, 4.4, 1H), 2.32 (m, 3H), 2.11 (m, 1H), 1.98 (d, J = 2.5, 1H), 1.77 (m, 2H), 1.61 (m, 1H), 1.48 (m, 4H), 1.17 (m, 1H), 0.77 (dt, J = 13.1, 8.9, 1H). 






(continued)

Compound	Name	Structure
641	[(5-methanesulfonylpyridin-3-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	MS: 430.2 ¹H NMR (400 MHz, CD₃CN) δ 9.10 (d, J = 2.0, 1H), 8.87 (d, J = 1.8, 1H), 8.71 (dd, J = 5.3, 1.1, 1H), 8.37 (t, J = 2.0, 1H), 8.18 (td, J = 8.0, 1.8, 1H), 7.75 (d, J = 8.2, 1H), 7.63 (ddd, J = 7.6, 5.4, 0.9, 1H), 4.16 (m, 2H), 3.73 (m, 2H), 3.14 (s, 3H), 3.02 (td, J = 12.0, 5.2, 1H), 2.52 (m, 1H), 2.33 (m, 3H), 2.14 (m, 1H), 2.01 (m, 1H), 1.79 (m, 2H), 1.62 (m, 1H), 1.48 (m, 4H), 1.19 (m, 1H), 0.79 (dt, J = 13.1, 8.9, 1H). 
642	[2-(3-methylphenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
643	[2-(3-chlorophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
644	[2-(3-bromophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	






(continued)

Compound	Name	Structure
645	[2-(3-fluorophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
646	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({2-[3-(trifluoromethyl)phenyl]ethyl})amine	
647	[2-(3-methoxyphenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
648	[2-(4-methylphenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
649	[2-(4-chlorophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	

(continued)

Compound	Name	Structure
650	[2-(4-bromophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
651	[2-(4-fluorophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
652	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({2-[4-(trifluoromethyl)phenyl]ethyl})amine	
653	[2-(4-methoxyphenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
654	[2-(2-methylphenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	


(continued)

Compound	Name	Structure
655	[2-(2-chlorophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
656	[2-(2-bromophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
657	[2-(2-fluorophenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	
658	{2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl}({2-[2-(trifluoromethyl)phenyl]ethyl})amine	
659	[2-(2-methoxyphenyl)ethyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine	

Example 15: Synthesis of [(3-methoxythiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine (Compound 140).

Methyl 2-cyano-2-[6-oxaspiro[4.5]decan-9-ylidene]acetate (mixture of E and Z isomers)


[0208]



[0209] A mixture of 6-oxaspiro[4.5]decan-9-one (13.74 g, 89.1 mmol), methylcyanoacetate (9.4 ml, 106.9 mmol), ammonium acetate (1.79 g, 26.17 mmol) and acetic acid (1.02 ml, 17.8 mmol) in benzene (75 ml) was heated at reflux in a 250 ml round bottom flask equipped with a Dean-Stark and a reflux condenser. After 3h, TLC (25% EtOAc in hexane, PMA stain) showed the reaction was completed. After cooling, benzene (50 ml) was added and the layer was separated, the organic was washed by water (120 ml) and the aqueous layer was extracted by CH₂Cl₂ (3 x 120 ml). The combined organic was washed with sat'd NaHCO₃, brine, dried and concentrated and the residual was purified by flash chromatography (340 g silica gel column, eluted by EtOAc in hexane: 5% EtOAc, 2CV; 5-25%, 14CV; 25-40%, 8 CV) gave a mixture of E and Z isomers: methyl 2-cyano-2-[6-oxaspiro[4.5]decan-9-ylidene]acetate (18.37 g, 87.8 % yield, m/z 236.0 [M + H]⁺ observed) as a clear oil.

Methyl 2-cyano-2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetate


[0210]



A solution of 2-bromopyridine (14.4 ml, 150 mmol) in THF (75 ml) was added dropwise to a solution of isopropylmagnesium chloride (75 ml, 2M in THF) at 0°C under N₂, the mixture was then stirred at rt for 3h, copper iodide (2.59 g, 13.6 mmol) was added and allowed to stir at rt for another 30 min before a solution of a mixture of E and Z isomers of methyl 2-cyano-2-[6-oxaspiro[4.5]decan-9-ylidene]acetate (16 g, 150 mmol) in THF (60 ml) was added in 30 min. The mixture was then stirred at rt for 18h. The reaction mixture was poured into a 200 g ice/2 N HCl (100 ml) mixture. The product was extracted with Et₂O (3x300 ml), washed with brine (200 ml), dried (Na₂SO₄) and concentrated. The residual was purified by flash chromatography (100 g silica gel column, eluted by EtOAc in hexane: 3% 2CV; 3-25%, 12 CV; 25-40% 6CV gave methyl 2-cyano-2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetate (15.44 g, 72% yield, m/z 315.0 [M + H]⁺ observed) as an amber oil.

2-[9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile

[0211]




[0212] Ethylene glycol (300 ml) was added to methyl 2-cyano-2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ace-

tate (15.43 g, 49 mmol) followed by potassium hydroxide (5.5 g, 98 mmol), the resulting mix was heated to 120°C, after 3 h, the reaction mix was cooled and water (300 ml) was added, the product was extracted by Et₂O (3 x 400 ml), washed with water (200 ml), dried (Na₂SO₄) and concentrated, the residual was purified by flash chromatography (340 g silica gel column, eluted by EtOAc in hexane: 3% 2CV; 3-25%, 12 CV; 25-40% 6CV to give 2-[9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (10.37 g, 82% yield, m/z 257.0 [M + H]⁺ observed).

2-[(9R)-9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile


[0213]



The racemic 2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile was separated by chiral HPLC column under the following preparative-SFC conditions: Instrument: SFC-80 (Thar, Waters); Column: Chiralpak AD-H (Daicel); column temperature: 40 °C; Mobile phase: Methanol /CO₂=40/60; Flow: 70 g/min; Back pressure: 120 Bar; Cycle time of stack injection: 6.0min; Load per injection: 225 mg; Under these conditions, 2-[9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (4.0 g) was separated to provide the desired isomer, 2-[(9R)-9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (2.0 g, >99.5% enantiomeric excess) as a slow-moving fraction. The absolute (R) configuration of the desired isomer was later determined by an X-ray crystal structure analysis of Compound 140.

2-[(9R)-9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine

[0214]




[0215] LAH (1M in Et₂O, 20ml, 20 mmol) was added to a solution of 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]acetonitrile (2.56 g, 10 mmol) in Et₂O (100 ml, 0.1M) at 0°C under N₂. The resulting mix was stirred and allowed to warm to room temperature. After 2 h, LCMS showed the reaction had completed. The reaction was cooled at 0°C and quenched with water (1.12 ml), NaOH (10%, 2.24 ml) and another 3.36 ml of water. Solid was filtered and filter pad was washed with ether (3 x 20 ml). The combined organic was dried and concentrated to give 2-[(9R)-9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine (2.44 g, 94% yield, m/z 260.6 [M + H]⁺ observed) as a light amber oil.

[0216] Alternatively, 2-[(9R)-9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine was prepared by Raney-Nickel catalyzed hydrogenation.

[0217] An autoclave vessel was charged with 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl] acetonitrile and ammonia (7N solution in methanol). The resulting solution was stirred at ambient conditions for 15 minutes and treated with Raney 2800 Nickel, slurried in water. The vessel was pressurized to 30 psi with nitrogen and agitated briefly. The autoclave was vented and the nitrogen purge repeated additional two times. The vessel was pressurized to 30 psi with hydrogen and agitated briefly. The vessel was vented and purged with hydrogen two additional times. The vessel was pressurized to 85-90 psi with hydrogen and the mixture was warmed to 25-35 °C. The internal temperature was increased to 45-50 °C over 30-60 minutes. The reaction mixture was stirred at 45-50 °C for 3 days. The reaction was monitored by HPLC. Once reaction was deemed complete, it was cooled to ambient temperature and filtered through celite. The filter cake was washed with methanol (2 x). The combined filtrates were concentrated under reduced pressure at 40-45 °C. The resulting residue was co-evaporated with EtOH (3 x) and dried to a thick syrupy of 2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine.

[(3-Methoxythiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine

[0218]



[0219] Into a vial were added 2-[(9R)-9-(Pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethan-1-amine (500 mg, 1.92 mmole), 18 mL CH₂Cl₂ and sodium sulfate (1.3 g, 9.6 mmole). The 3-methoxythiophene-2-carboxaldehyde (354 mg, 2.4 mmole) was then added, and the mixture was stirred overnight. NaBH₄ (94 mg, 2.4 mmole) was added to the reaction mixture, stirred for 10 minutes, and then MeOH (6.0 mL) was added, stirred 1h, and finally quenched with water. The organics were separated off and evaporated. The crude residue was purified by a Gilson prep HPLC. The desired fractions collected and concentrated and lyophilized. After lyophilization, residue was partitioned between CH₂Cl₂ and 2N NaOH, and the organic layers were collected. After solvent was concentrated to half of the volume, 1.0 eq of 1N HCl in Et₂O was added, and majority of solvent evaporated under reduced pressure. The solid obtained was washed several times with Et₂O and dried to provide [(3-methoxythiophen-2-yl)methyl]({2-[(9R)-9-(pyridin-2-yl)-6-oxaspiro[4.5]decan-9-yl]ethyl})amine monohydrochloride (336 mg, 41% yield, m/z 387.0 [M + H]⁺ observed) as a white solid. The NMR for Compound 140 is described herein.

Example 16: Biological Example

Procedure for the Testing for Antinociception

[0220] The hot plate assay is adapted from the procedure originally described by O'Callaghan and Holtzman (JPET, 192, 497, 1975) and is commonly used to determine the potential analgesic efficacy of opioid agonists. The antinociceptive effect of the composition(s) described herein in the hot plate is expressed in %MPE (Maximum Possible Effect).

[0221] Rats (175-250g) or mice (20-30g) acclimated to the vivarium for at least 48 hr prior to behavioral testing. Test drugs were administered by the subcutaneous (SC) route. Animals were placed on the hot plate, which the temperature was set at 50-56°C, depending on the *in vitro* potency of the compound. A cutoff time of 30-60 seconds was used depending on the temperature of the hot plate so that the paws of the animal displaying analgesia, was not damaged by the heat stimulus. The cutoff time was considered a 100% response to the thermal insult. Prior to drug treatment, each animal was tested to determine the baseline response. Thirty minutes after drug administration, animals were re-tested. Dose response experiments were performed to evaluate the potency of the test compound when various doses were administered at the point when maximal analgesia is observed.

[0222] The %MPE was calculated according to the following formula:

$$\%MPE = [(Post\ drug\ latency - baseline\ latency) / (60\ or\ 30 - baseline\ latency)] \times 100$$

[0223] ED₅₀ values were calculated from the mean %MPE values for each group using log dose-response curves by least-squares regression analysis.

Table 4

COMPOUND	ED ₅₀ or %MPE
Morphine	3.8 mg/kg SC
Compound 81	100% at 10 mg/kg SC
Compound 122	1.1 mg/kg SC
Compound 28	1.2 mg/kg SC
Compound 145	5.9 mg/kg SC

[0224] Results are shown in **Table 4**. Naïve or control mice typically exhibit reaction times in the hot plate from 10-15 seconds. The ED₅₀ for morphine in the mouse hot plate was 3.8 mg/kg with full efficacy observed at a dose of 10 mg/kg SC. For comparison, Compound 122 and Compound 28 produced potent efficacy with an ED₅₀ of 1.1 and 1.2 mg/kg SC, respectively. These results demonstrate that Compound 122 and Compound 28 produced a more robust analgesic effect in the mouse hot plate assay compared to morphine.

Example 17: *In vivo* administration to humans (Prophetic Example)

[0225] One or more compounds will be administered in dosage range from 0.15 mg to 4 mg to human subject. The compound(s) will be administered as a continuous infusion over one hour. The dose may be escalated as deemed appropriate to obtain pain relief. Dose escalation will usually not exceed 5-fold as compared to the previous dose. Dosage amounts, however, may be repeated or decreased as deemed appropriate. The subjects will be tested for their ability to withstand or not appreciate pain as compared to a control (placebo) group.


[0226] The cold pain test has been shown to be a reproducible and sensitive measure of the effect of opiates and other centrally acting drugs (Van F and Rolan PE. The utility of the cold pain test to measure analgesia from intravenous morphine. *Br. J. Clin. Pharmacol.* 1996; 42: 663-664; ; Posner J. Pain Models in Healthy Volunteers. In: Nimmo WS, Tucker G, eds. *Clinical Measurement in Drug Evaluation*. 1991, Wolfe Publishing Limited, UK.; Wotherspoon HA, Kenny GNC, McArdle CS. Analgesic Efficacy of Controlled-Release DihydroCodeine. *Anaesthesia* 1991; 46: 915-917.; Lamb RJ, Mercer AJ, Posner J. The effect of lamotrigine (300 mg) and dipipanone (4 mg and 8 mg), alone and in combination, on the cold-pain test in healthy volunteers. *Br. J. Clin. Pharmacol.* 1994; 39: 539-588P.). In the test a subject's hand is immersed in cold water chilled to a range of 1 to 3 °C. The initial sensation of cold is replaced by a deep burning discomfort in the hand which is mediated by nociceptors in veins. The discomfort gradually builds to a plateau over approximately 90 seconds and then either persists or decreases slightly. The stimulus is easily controlled and the response is reproducible. The technique has been shown to be sensitive to different doses of analgesic drugs.

[0227] During the cold pain test, the subject will sit down and place his/her non-dominant hand into a stirred, thermostatically controlled water bath at about 2 °C. With the other hand the subject can adjust a visual analogue scale on a computer screen using the arrow keys on the keypad. The scale is labelled "no pain" at one end and "maximum pain" at the other end. The pointer will initially be at the "no pain" end and the subject will move the pointer across the line to rate their feelings continuously over the test period. At the end of 2 minutes the computer will automatically instruct the subject to remove his/her hand which can then be dried. The cold pain test has been used extensively in healthy volunteer studies and is non-invasive.

[0228] It is expected that the administration of the compound(s) will enable the human subject to feel no pain or less pain as compared to the control group.

Claims

1. A compound having a formula of Formula IV-1, or a pharmaceutically acceptable salt thereof,;



wherein:

R₂₁ and R₂₂ are independently H or CH₃;

A₄ is an optionally substituted cycle of the formula C(CH₂)_n, where n = 2-5, wherein the optional substitution is halo, cyano, C₁-C₆ alkyl, C₁-C₆ branched alkyl, halogenated C₁-C₆ alkyl, hydroxyl, C₁-C₆ alkyloxy, amino, or C₁-C₆ alkylamino;


B₃ is H or an optionally substituted C₁-C₆ alkyl, wherein the optional substitution is halo, hydroxy or C₁-C₃ alkoxy;

B₄ is C₁-C₆ alkylene;

D₁ is an optionally substituted monocyclic or bicyclic aryl or heteroaryl, wherein each ring is a 5, 6 or 7 membered ring and the optional substitution is halo, cyano, C₁-C₆ alkyl, C₁-C₆ branched alkyl, halogenated C₁-C₆ alkyl, hydroxyl, C₁-C₆ alkoxy, amino, or C₁-C₆ alkylamino; and


B₅ is an optionally substituted monocyclic or bicyclic aryl or heteroaryl, wherein each ring is a 5, 6, or 7 membered ring and the optional substitution is halo, cyano, C₁-C₆ alkyl, C₁-C₆ branched alkyl, halogenated C₁-C₆ alkyl, hydroxyl, C₁-C₆ alkoxy, amino, or C₁-C₆ alkylamino.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, having the structure of Formula VIII:




wherein R₂₆ and R₂₇ are independently H or an isotope thereof.

3. The compound of claim 2, or a pharmaceutically acceptable salt thereof, wherein the compound has a formula of Formula IX




4. The compound of claims 2 or 3, or a pharmaceutically acceptable salt thereof, wherein the compound has a formula of Formula VII-1




VII-1.

5. The compound of any of claims 2-4, or a pharmaceutically acceptable salt thereof, wherein the compound has a formula of Formula X




X.

6. The compound of any of claims 1-5, or a pharmaceutically acceptable salt thereof, wherein D_1 is an optionally substituted phenyl, wherein the optional substitution is halo, cyano, C_1 - C_6 alkyl, C_1 - C_6 branched alkyl, halogenated C_1 - C_6 alkyl, hydroxyl, C_1 - C_6 alkyloxy, amino, or C_1 - C_6 alkylamino or an optionally substituted pyridyl, wherein the optional substitution is halo, cyano, C_1 - C_6 alkyl, C_1 - C_6 branched alkyl, halogenated C_1 - C_6 alkyl, hydroxyl, C_1 - C_6 alkyloxy, amino, or C_1 - C_6 alkylamino.
7. The compound of any of claims 1-5, or a pharmaceutically acceptable salt thereof, wherein B_5 is an optionally substituted monocyclic or bicyclic heteroaryl, wherein each ring is a 5, 6, or 7 membered ring and the optional substitution is halo, cyano, C_1 - C_6 alkyl, C_1 - C_6 branched alkyl, halogenated C_1 - C_6 alkyl, hydroxyl, C_1 - C_6 alkyloxy, amino, or C_1 - C_6 alkylamino.
8. The compound of any of claims 1-5, or a pharmaceutically acceptable salt thereof, wherein B_5 is an optionally substituted aryl or heteroaryl selected from the group consisting of




5




10

15



20


25



and

30

35




wherein the optional substitution is halo, cyano, C₁-C₆ alkyl, C₁-C₆ branched alkyl, halogenated C₁-C₆ alkyl, hydroxyl, C₁-C₆ alkyloxy, amino, or C₁-C₆ alkylamino.

40


9. The compound of claims 1-5, or a pharmaceutically acceptable salt thereof, wherein B₅ is selected from the group consisting of


45

50



55






wherein

R₂₃, R₂₄, and R₃₀ are each independently null, H, OH, halo, branched or unbranched C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, amino, C₁-C₆ haloalkyl, or, R₂₃ and R₂₄ together form a aryl, heteroaryl or cycle that is attached to one or more of the atoms of B₅; and R₂₅ is H or an optionally substituted branched or unbranched alkyl, wherein the optional substitution is halo, hydroxy or C₁-C₃ alkoxy.

10. The compound of claim 9, or a pharmaceutically acceptable salt thereof, wherein R₂₃, R₂₄, and R₃₀ are each independently H, NH₂, OH, Cl, Br, F, I, OMe, CN, or CH₃.


11. The compound of claim 9, or a pharmaceutically acceptable salt thereof, wherein B₅ is




12. The compound of claim 9 or 11, or a pharmaceutically acceptable salt thereof, wherein R_{23} is C_1 - C_6 alkoxy.

13. The compound of any of claims 1-12, or a pharmaceutically acceptable salt thereof, wherein B_3 is H or C_1 - C_6 alkyl.


14. The compound of claim 1, or a pharmaceutically acceptable salt thereof, having the formula of:




15. The compound of claim 1, or a pharmaceutically acceptable salt thereof, having the formula of:




16. The compound of claim 1, or a pharmaceutically acceptable salt thereof, having the formula of:



17. The compound of claim 1, or a pharmaceutically acceptable salt thereof, having the formula of:




18. A pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt thereof, of any of claims 1-17 and a pharmaceutically acceptable carrier.
19. A compound or a pharmaceutically acceptable salt thereof, or composition of any of claims 1-18 for use in the treatment of pain.
20. A process of preparing a compound, or a pharmaceutically acceptable salt thereof, of claim 2, wherein R_{26} and R_{27} are H, the process comprising: contacting



with




under suitable conditions to form a compound having the structure of



Patentansprüche

1. Verbindung mit einer Formel der Formel IV-1, oder ein pharmazeutisch akzeptables Salz davon:



IV-1

worin:

R_{21} und R_{22} unabhängig H oder CH_3 sind;

A_4 ein optional substituierter Cyclus der Formel $C(CH_2)_n$ ist, worin $n = 2-5$, wobei die optionale Substitution Halogen, Cyano, C_1 - C_6 -Alkyl, verzweigtes C_{1-6} -Alkyl, halogeniertes C_{1-6} -Alkyl, Hydroxyl, C_{1-6} -Alkyloxy, Amino oder C_{1-6} -Alkylamino ist;


B_3 H oder ein optional substituiertes C_{1-6} -Alkyl ist, wobei die optionale Substitution Halogen, Hydroxy oder C_{1-3} -Alkoxy ist,

B_4 C_{1-6} -Alkyl ist;

D_1 ein optional substituiertes monocyclisches oder bicyclisches Aryl oder Heteroaryl ist, wobei jeder Ring ein 5-, 6- oder 7-gliedriger Ring ist und die optionale Substitution Halogen, Cyano, C_{1-6} -Alkyl, verzweigtes C_{1-6} -Alkyl, halogeniertes C_{1-6} -Alkyl, Hydroxyl, C_{1-6} -Alkyloxy, Amino oder C_{1-6} -Alkylamino ist; und

B_5 ein optional substituiertes monocyclisches oder bicyclisches Aryl oder Heteroaryl ist, wobei jeder Ring ein 5-, 6- oder 7-gliedriger Ring ist und die optionale Substitution Halogen, Cyano, C_{1-6} -Alkyl, verzweigtes C_{1-6} -Alkyl, halogeniertes C_{1-6} -Alkyl, Hydroxyl, C_{1-6} -Alkyloxy, Amino oder C_{1-6} -Alkylamino ist.


2. Die Verbindung nach Anspruch 1, oder ein pharmazeutisch akzeptables Salz davon, mit der Struktur der Formel VIII:




VIII,

worin R_{26} und R_{27} unabhängig H oder ein Isotop davon sind.


3. Die Verbindung nach Anspruch 2, oder ein pharmazeutisch akzeptables Salz davon, wobei die Verbindung eine Formel aufweist der Formel IX:



4. Die Verbindung nach Ansprüchen 2 oder 3, oder ein pharmazeutisch akzeptables Salz davon, wobei die Verbindung eine Formel aufweist der Formel VII-1:




5. Die Verbindung nach einem der Ansprüche 2-4, oder ein pharmazeutisch akzeptables Salz davon, wobei die Verbindung eine Formel aufweist der Formel X:




6. Die Verbindung nach einem der Ansprüche 1-5, oder ein pharmazeutisch akzeptables Salz davon, worin D₁ ein optional substituiertes Phenyl ist, wobei die optionale Substitution Halogen, Cyano, C₁-C₆-Alkyl, verzweigtes C₁₋₆-Alkyl, halogeniertes C₁₋₆-Alkyl, Hydroxyl, C₁₋₆-Alkyloxy, Amino oder C₁₋₆-Alkylamino ist, oder ein optional substituiertes Pyridyl ist, wobei die optionale Substitution Halogen, Cyano, C₁-C₆-Alkyl, verzweigtes C₁₋₆-Alkyl, halogeniertes C₁₋₆-Alkyl, Hydroxyl, C₁₋₆-Alkyloxy, Amino oder C₁₋₆-Alkylamino ist.
7. Die Verbindung nach einem der Ansprüche 1-5, oder ein pharmazeutisch akzeptables Salz davon, worin B₅ ein optional substituiertes monocyclisches oder bicyclisches Heteroaryl ist, wobei jeder Ring ein 5-, 6- oder 7-gliedriger Ring ist und die optionale Substitution Halogen, Cyano, C₁₋₆-Alkyl, verzweigtes C₁₋₆-Alkyl, halogeniertes C₁₋₆-Alkyl, Hydroxyl, C₁₋₆-Alkyloxy, Amino oder C₁₋₆-Alkylamino ist.
8. Die Verbindung nach einem der Ansprüche 1-5, oder ein pharmazeutisch akzeptables Salz davon, worin B₅ ein optional substituiertes Aryl oder Heteroaryl ist, das ausgewählt ist aus der Gruppe bestehend aus

5




10




15

20




25



30


35



40

und

45




50


wobei die optionale Substitution Halogen, Cyano, C₁-C₆-Alkyl, verzweigtes C₁₋₆-Alkyl, halogeniertes C₁₋₆-Alkyl, Hydroxyl, C₁₋₆-Alkyloxy, Amino oder C₁₋₆-Alkylamino ist.

55


9. Die Verbindung nach Ansprüchen 1-5, oder ein pharmazeutisch akzeptables Salz davon, worin B₅ ausgewählt ist aus der Gruppe bestehend aus



worin B₅ ausgewählt ist aus der Gruppe bestehend aus



und




worin

R_{23} , R_{24} und R_{30} jeweils unabhängig Null, H, OH, Halogen, verzweigtes oder unverzweigtes C_{1-6} -Alkyl, Cyano, C_{1-6} -Alkoxy, Amino, C_{1-6} -Haloalkyl sind, oder R_{23} und R_{24} zusammen ein Aryl, Heteroaryl oder einen Cyclus bilden, das/der an eines oder mehrere der Atome von B_5 angelagert ist; und R_{25} H oder ein optional substituiertes verzweigtes oder unverzweigtes Alkyl ist, wobei die optionale Substitution Halogen, Hydroxy oder C_{1-3} -Alkoxy ist.

10. Die Verbindung nach Anspruch 9, oder ein pharmazeutisch akzeptables Salz davon, worin R_{23} , R_{24} und R_{30} jeweils unabhängig H, NH_2 , OH, Cl, Br, F, I, OMe, CN oder CH_3 sind.


11. Die Verbindung nach Anspruch 9, oder ein pharmazeutisch akzeptables Salz davon, worin B_5 ist




12. Die Verbindung nach Anspruch 9 oder 11, oder ein pharmazeutisch akzeptables Salz davon, worin R_{23} C_{1-6} -Alkoxy ist.

13. Die Verbindung nach einem der Ansprüche 1-12, oder ein pharmazeutisch akzeptables Salz davon, worin B_3 H oder C_{1-6} -Alkyl ist.


14. Die Verbindung nach Anspruch 1, oder ein pharmazeutisch akzeptables Salz davon, mit der Formel:




15. Die Verbindung nach Anspruch 1, oder ein pharmazeutisch akzeptables Salz davon, mit der Formel:



16. Die Verbindung nach Anspruch 1, oder ein pharmazeutisch akzeptables Salz davon, mit der Formel:




17. Die Verbindung nach Anspruch 1, oder ein pharmazeutisch akzeptables Salz davon, mit der Formel:




18. Pharmazeutische Zusammensetzung, umfassend eine Verbindung, oder ein pharmazeutisch akzeptables Salz davon, nach einem der Ansprüche 1-17 und einen pharmazeutisch akzeptablen Träger.

19. Verbindung oder ein pharmazeutisch akzeptables Salz davon, oder Zusammensetzung nach einem der Ansprüche 1-18 zur Verwendung bei der Behandlung von Schmerzen.

20. Verfahren zur Herstellung einer Verbindung, oder eines pharmazeutisch akzeptablen Salzes davon, nach Anspruch 2, worin R_{26} und R_{27} H sind, welches Verfahren umfasst: Kontaktieren von




unter geeigneten Bedingungen, um eine Verbindung zu bilden mit der Struktur



Revendications

1. Composé ayant une formule de la formule IV-1, ou un sel pharmaceutiquement acceptable de celui-ci, :



IV-1

dans lequel :

R₂₁ et R₂₂ sont indépendamment H ou CH₃ ;

A₄ est un cycle éventuellement substitué de la formule C(CH₂)_n, où n = 2 à 5, la substitution éventuelle étant halo, cyan, alkyle en C₁-C₆, alkyle en C₁ à C₆ ramifié, alkyle en C₁ à C₆ halogéné, hydroxyle, alkyloxy en C₁ à C₆, amino, ou alkylamino en C₁ à C₆ ;


B₃ est H ou un alkyle en C₁ à C₆ éventuellement substitué, dans lequel la substitution éventuelle est halo, hydroxy ou alcoxy en C₁ à C₃ ;

B₄ est alkylène en C₁ à C₆ ;

D₁ est un aryle ou un hétéroaryle monocyclique ou bicyclique éventuellement substitué, chaque noyau étant un noyau à 5, 6 ou 7 éléments et la substitution éventuelle est halo, cyan, alkyle en C₁ à C₆, alkyle en C₁ à C₆ ramifié, alkyle en C₁ à C₆ halogéné, hydroxyle, alkyloxy en C₁ à C₆, amino ou alkylamino en C₁ à C₆ ; et

B₅ est un aryle ou un hétéroaryle monocyclique ou bicyclique éventuellement substitué, chaque noyau étant un noyau à 5, 6 ou 7 éléments et la substitution éventuelle est halo, cyan, alkyle en C₁ à C₆, alkyle en C₁ à C₆ ramifié, alkyle en C₁ à C₆ halogéné, hydroxyle, alkyloxy en C₁ à C₆, amino ou alkylamino en C₁ à C₆.


2. Composé selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, ayant la structure de la formule VIII :




VIII,

dans lequel R_{26} et R_{27} sont indépendamment H ou un isotope de celui-ci.


3. Composé selon la revendication 2, ou un sel pharmaceutiquement acceptable de celui-ci, le composé ayant une formule de la formule IX :



4. Composé selon la revendication 2 ou 3, ou un sel pharmaceutiquement acceptable de celui-ci, le composé ayant une formule de la formule VII-I :



5. Composé selon l'une quelconque des revendications 2 à 4, ou un sel pharmaceutiquement acceptable de celui-ci, le composé ayant une formule de la formule X




6. Composé selon l'une quelconque des revendications 1 à 5, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel D_1 est un phényle éventuellement substitué, dans lequel la substitution éventuelle est halo, cyan, alkyle en C_1 à C_6 , alkyle en C_1 à C_6 ramifié, alkyle en C_1 à C_6 halogéné, hydroxyle, alkyloxy en C_1 à C_6 , amino, ou alkylamino en C_1 à C_6 ou un pyridyle éventuellement substitué, la substitution éventuelle est halo, cyan, alkyle en C_1 à C_6 , alkyle en C_1 à C_6 ramifié, alkyle en C_1 à C_6 halogéné, hydroxyle, alkyloxy en C_1 à C_6 , amino, alkylamino


en C₁ à C₆.

7. Composé selon l'une quelconque des revendications 1 à 5, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel B₅ est un hétéroaryle monocyclique ou bicyclique éventuellement substitué, chaque noyau étant un noyau à 5, 6 ou 7 éléments et la substitution éventuelle est halo, cyan, alkyle en C₁ à C₆, alkyle en C₁ à C₆ ramifié, alkyle en C₁ à C₆ halogéné, hydroxyle, alkyloxy en C₁ à C₆, amino ou alkylamino en C₁ à C₆.

8. Composé selon l'une quelconque des revendications 1 à 5, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel B₅ est un aryle ou un hétéroaryle éventuellement substitué choisi dans le groupe constitué de




et de




la substitution éventuelle étant halo, cyano, alkyle en C₁ à C₆, alkyle en C₁ à C₆ ramifié, alkyle en C₁ à C₆ halogéné, hydroxyle, alkyloxy en C₁ à C₆, amino, ou alkylamino en C₁ à C₆.

9. Composé selon les revendications 1 à 5, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel B₅ est choisi dans le groupe constitué de



et de




dans lequel

R_{23} , R_{24} et R_{30} sont chacun indépendamment zéro, H, OH, halo, alkyle en C_1 à C_6 ramifié ou non ramifié, cyan, alcoxy en C_1 à C_6 , amino, haloalkyle en C_1 à C_6 , ou, R_{23} et R_{24} forment ensemble un aryle, un hétéroaryle ou un cycle qui est attaché à un ou plusieurs des atomes de B_5 ; et R_{25} est H ou un alkyle ramifié ou non ramifié éventuellement substitué, la substitution éventuelle étant halo, hydroxy ou alcoxy en C_1 à C_3 .

10. Composé selon la revendication 9, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel R_{23} , R_{24} et R_{30} sont chacun indépendamment H, NH_2 , OH, Cl, Br, F, I, OMe, CN, ou CH_3 .


11. Composé selon la revendication 9, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel B_5 est




12. Composé selon la revendication 9 ou 11, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel R_{23} est alcoxy en C_1 à C_6 .

13. Composé selon l'une quelconque des revendications 1 à 12, ou un sel pharmaceutiquement acceptable de celui-ci, dans lequel B_3 est H ou alkyle en C_1 à C_6 .


14. Composé selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, ayant la formule :




15. Composé selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, ayant la formule :



16. Composé selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, ayant la formule :




17. Composé selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, ayant la formule :




18. Composition pharmaceutique comprenant un composé, ou un sel pharmaceutiquement acceptable de celui-ci, selon l'une quelconque des revendications 1 à 17 et un support pharmaceutiquement acceptable.

19. Composé ou un sel pharmaceutiquement acceptable de celui-ci, ou une composition selon l'une quelconque des revendications 1 à 18 à utiliser dans le traitement de la douleur.

20. Processus de préparation d'un composé, ou d'un sel pharmaceutiquement acceptable de celui-ci, selon la revendication 2, dans lequel R_{26} et R_{27} sont H, le processus comprenant : la mise en contact de



dans des conditions appropriées pour former un composé ayant la structure de



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 2009247561 A [0004]
- US 4115538 A [0077]
- US 5436128 A [0078]
- US 6136839 A [0115]

Non-patent literature cited in the description

- **GROARKE et al.** *J. Biol. Chem.*, 1999, vol. 274 (33), 23263-69 [0075]
- **KROEGER et al.** *J. Biol. Chem.*, 2001, vol. 276 (16), 12736-43 [0076]
- **OFFERMANN ; SIMON.** *J. Biol. Chem.*, 1995, vol. 270, 15175-15180 [0077]
- **FELLEY-BOSCO et al.** *Am. J. Resp. Cell and Mol. Biol.*, 1994, vol. 11, 159-164 [0077]
- **MISTILI ; SPECTOR.** *Nature Biotechnology*, 1997, vol. 15, 961-964 [0078]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990, 1435-1712 [0081]
- Remington: The Science and Practice of Pharmacy. 1995 [0085] [0088] [0092] [0094] [0097]
- Remington's Pharmaceutical Sciences. 1985 [0097]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1985 [0098]
- **T. HIGUCHI et al.** Pro-drugs as Novel Delivery Systems. A.C.S. Symposium Series, vol. 14 [0108]
- Bioreversible Carriers in Drug Design. American Pharmaceutical Association and Pergamon Press, 1987 [0108]
- Analgesic-Antipyretic and Anti-inflammatory Agents and Drugs Employed in the Treatment of Gout. **PAUL A. INSEL.** Goodman & Gilman's The Pharmacological Basis of Therapeutics. 1996, 617-57 [0114]
- Analgesic, Antipyretic and Anti-Inflammatory Drugs. **GLEN R. HANSON.** Remington: The Science and Practice of Pharmacy. 1995, vol. II, 1196-1221 [0114]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1990, 1445 [0140]
- **CHOU ; TALALAY.** *Adv. Enzyme Regul.*, 1984, vol. 22, 27-55 [0146]
- **HANSCHKE, E.** *Chem. Ber.*, 1955, vol. 88, 1053 [0170]
- **VAN F ; ROLAN PE.** The utility of the cold pain test to measure analgesia from intravenous morphine. *Br. J. Clin. Pharmacol.*, 1996, vol. 42, 663-664 [0226]
- Pain Models in Healthy Volunteers. **POSNER J.** Clinical Measurement in Drug Evaluation. Wolfe Publishing Limited, 1991 [0226]
- **WOTHERSPOON HA ; KENNY GNC ; MCARDLE CS.** Analgesic Efficacy of Controlled-Release DihydroCodeine. *Anaesthesia*, 1991, vol. 46, 915-917 [0226]
- **LAMB RJ ; MERCER AJ ; POSNER J.** The effect of lamotrigine (300 mg) and dipipanone (4 mg and 8 mg), alone and in combination, on the cold-pain test in healthy volunteers. *Br. J. Clin. Pharmacol.*, 1994, vol. 39, 539-588P [0226]




716730/SZE

EP2688403

Opioid receptor ligandumok és eljárások azok alkalmazására és előállítására

Szabadalmi igénypontok

1. IV-1:




IV-1

képletű vegyület, vagy gyógyászatiilag elfogadható sója, ahol:

 R_{21} és R_{22} egymástól függetlenül H vagy CH_3 ; A_4 egy adott esetben szubsztituált $\text{C}(\text{CH}_2)_n$ képletű gyűrűs csoport, ahol $n = 2-5$, ahol az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6 szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport; B_3 H vagy egy adott esetben szubsztituált 1-6 szénatomos alkilcsoport, ahol az adott esetben jelenlévő szubsztituens halogénatom, hidroxil vagy 1-3 szénatomos alkoxycsoport; B_4 1-6 szénatomos alkilcsoport; D_1 egy adott esetben szubsztituált monociklusos vagy biciklusos aril vagy heteroaril, ahol mindegyik gyűrű is egy 5, 6 vagy 7-tagú gyűrű és az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6 szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport; és B_5 egy adott esetben szubsztituált monociklusos vagy biciklusos aril vagy heteroarilcsoport, ahol mindegyik gyűrű egy 5, 6, vagy 7-tagú gyűrű és az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6


szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport.

2. Az 1. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, mely VIII:




szerkezetű, ahol R_{26} és R_{27} egymástól függetlenül H vagy egy izotópja.

3. A 2. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol a vegyület IX:



képletű.


4. A 2. vagy 3. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol a vegyület VII-1:



VII-1.

képletű.

5. A 2-4. igénypontok bármelyike szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol a vegyület X:




X.

képletű.

6. Az 1-5. igénypontok bármelyike szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol D₁ egy adott esetben szubsztituált fenilcsoport, ahol az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6 szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport, vagy egy adott esetben szubsztituált piridilcsoport, ahol az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6 szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport.


7. Az 1-5. igénypontok bármelyike szerinti vegyület vagy gyógyászatilag elfogadható sója, ahol B_5 egy adott esetben szubsztituált monociklusos vagy biciklusos heteroarilcsoport, ahol mindegyik gyűrű egy 5, 6, vagy 7-tagú gyűrű és az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6 szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport.

8. Az 1-5. igénypontok bármelyike szerinti vegyület vagy gyógyászatilag elfogadható sója, ahol B_5 egy adott esetben szubsztituált,




képletű aril vagy heteroarilcsoport közül választott, és ahol az adott esetben jelenlévő szubsztituens halogénatom, ciano, 1-6 szénatomos alkil, 1-6 szénatomos elágazó alkil, halogénezett 1-6 szénatomos alkil, hidroxil, 1-6 szénatomos alkiloxi, amino, vagy 1-6 szénatomos alkilaminocsoport.

9. Az 1-5. igénypontok bármelyike szerinti vegyület vagy gyógyászatilag elfogadható sója, ahol B₅



és




csoportok közül választott, ahol

R_{23} , R_{24} , és R_{30} mindegyike egymástól függetlenül nulla, H, OH, halogénatom, elágazó vagy nem-elágazó 1-6 szénatomos alkil, ciano, 1-6 szénatomos alkoxi, amino, 1-6 szénatomos halogénalkilcsoport, vagy, R_{23} és R_{24} együtt egy aril, heteroaril vagy gyűrűs csoportot alkot, mely B_5 egy vagy több atomjához kapcsolódik; és R_{25} jelentése H vagy egy adott esetben szubsztituált elágazó vagy nem-elágazó alkilcsoport, ahol az adott esetben jelenlévő szubsztituens halogénatom, hidroxil vagy 1-3 szénatomos alkoxycsoport.

10. A 9. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol R_{23} , R_{24} , és R_{30} mindegyike egymástól függetlenül H, NH_2 , OH, Cl, Br, F, I, OMe, CN, vagy CH_3 csoport.

11. A 9. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol B_5




képletű csoport.

12. A 9. vagy 11. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol R_{23} 1-6 szénatomos alkoxycsoport.


13. Az 1-12. igénypontok bármelyike szerinti vegyület vagy gyógyászati lag elfogadható sója, ahol B_5 jelentése H vagy 1-6 szénatomos alkilcsoport.

14. Az 1. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, mely a:




képletű.

15. Az 1. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, mely a:




képletű.

16. Az 1. igénypont szerinti vegyület vagy gyógyászati lag elfogadható sója, mely a:



képletű.

17. Az 1. igénypont szerinti vegyület vagy gyógyászatilag elfogadható sója, mely a:




képletű.

18. Gyógyászati készítmény, mely egy 1-17. igénypontok bármelyike szerinti vegyületet és egy gyógyászatilag elfogadható hordozót tartalmaz.

19. Egy 1-18. igénypontok bármelyike szerinti vegyület, vagy gyógyászatilag elfogadható sója, vagy készítmény fájdalom kezelésében történő alkalmazásra.

20. Eljárás egy 2. igénypont szerinti vegyület vagy gyógyászatilag elfogadható sója előállítására, ahol R_{26} és R_{27} hidrogénatom, melynek során:


egy



képletű vegyületet egy



képletű vegyülettel hozunk érintkezésbe, egy



szerkezetű vegyület képződéséhez megfelelő körülmények között.