

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12Q 1/68, C12N 9/22, C07H 21/04</p>	<p>A1</p>	<p>(11) International Publication Number: WO 96/17086 (43) International Publication Date: 6 June 1996 (06.06.96)</p>
<p>(21) International Application Number: PCT/US95/15580 (22) International Filing Date: 1 December 1995 (01.12.95) (30) Priority Data: 08/349,023 2 December 1994 (02.12.94) US 08/472,194 7 June 1995 (07.06.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/472,194 (CIP) Filed on 7 June 1995 (07.06.95) (71) Applicant (for all designated States except US): THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10666 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): JOYCE, Gerald, F. [US/US]; 837 4th Street, Encinitas, CA 92024 (US). BREAKER, Ronald, R. [US/US]; 10116 Caminito Mulege, San Diego, CA 92126 (US).</p>	<p>(74) Agents: LOGAN, April, C. et al.; The Scripps Research Institute, 10666 North Torrey Pines Road, TPC-8, La Jolla, CA 92037 (US). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: ENZYMATIC DNA MOLECULES (57) Abstract The present invention discloses deoxyribonucleic acid enzymes - catalytic or enzymatic DNA molecules - capable of cleaving nucleic acid sequences or molecules, particularly RNA, in a site-specific manner, as well as compositions including same. Methods of making and using the disclosed enzymes and compositions are also disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

ENZYMATIC DNA MOLECULES

TECHNICAL FIELD

5 The present invention relates to nucleic acid enzymes or catalytic (enzymatic) DNA molecules that are capable of cleaving other nucleic acid molecules, particularly RNA. The present invention also relates to compositions containing the disclosed enzymatic DNA molecules and to methods of making and using such enzymes and compositions.

BACKGROUND

10 The need for catalysts that operate outside of their native context or which catalyze reactions that are not represented in nature has resulted in the development of "enzyme engineering" technology. The usual route taken in enzyme engineering has been a "rational design" approach, relying upon the understanding of natural enzymes to aid in the construction of new enzymes. Unfortunately, the state of proficiency in the areas of protein structure and chemistry is insufficient to make the generation of novel biological catalysts routine.

15 Recently, a different approach for developing novel catalysts has been applied. This method involves the construction of a heterogeneous pool of macromolecules and the application of an *in vitro* selection procedure to isolate molecules from the pool that catalyze the desired reaction. Selecting catalysts from a pool of macromolecules is not dependent on a comprehensive understanding of their structural and chemical properties. Accordingly, this process has been dubbed "irrational design" (Brenner and Lerner, PNAS USA 89: 5381-5383 (1992)).

25 Most efforts to date involving the rational design of enzymatic RNA molecules or ribozymes have not led to molecules with fundamentally new or improved catalytic function. However, the application of irrational design methods via a process we have described as "directed molecular evolution" or "*in vitro* evolution", which is patterned after Darwinian evolution of organisms in nature, has the potential to lead to the production of DNA molecules that have desirable functional characteristics.

30 This technique has been applied with varying degrees of success to RNA molecules in solution (see, e.g., Mills, et al., PNAS USA 58: 217 (1967); Green, et al., Nature 347: 406 (1990); Chowrira, et al., Nature 354: 320 (1991); Joyce, Gene 82: 83 (1989); Beaudry and Joyce, Science 257: 635-641 (1992); Robertson and Joyce, Nature 344: 467 (1990)), as well as to RNAs bound to a ligand that is attached to a solid support (Tuerk, et al., Science 249: 505 (1990); Ellington, et al., Nature 346: 818 (1990)). It has also been applied to peptides attached directly to a solid support (Lam, et al., Nature 354: 82 (1991)); and to peptide epitopes expressed within a viral coat

-2-

protein (Scott, et al., Science 249: 386 (1990); Devlin, et al., Science 249: 249 (1990); Cwirla, et al., PNAS USA 87: 6378 (1990)).

5 It has been more than a decade since the discovery of catalytic RNA (Kruger, et al., Cell 31: 147-157 (1982); Guerrier-Takada, et al., Cell 35: 849-857 (1983)). The list of known naturally-occurring ribozymes continues to grow (see Cech, in The RNA World, Gesteland & Atkins (eds.), pp. 239-269, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1993); Pyle, Science 261: 709-714 (1993); Symons, Curr. Opin. Struct. Biol. 4: 322-330 (1994)) and, in recent years, has been augmented by synthetic ribozymes obtained through *in vitro* evolution. (See, e.g., Joyce, Curr. Opin. Struct. Biol. 4: 331-336 (1994); Breaker & Joyce, Trends Biotech. 12: 268-275 (1994); Chapman & Szostak, Curr. Opin. Struct. Biol. 4: 618-622 (1994).)

10 It seems reasonable to assume that DNA can have catalytic activity as well, considering that it contains most of the same functional groups as RNA. However, with the exception of certain viral genomes and replication intermediates, nearly all of the DNA in biological organisms occurs as a complete duplex, precluding it from adopting a complex secondary and tertiary structure. Thus it is not surprising that DNA enzymes have not been found in nature.

15 Until the advent of the present invention, the design, synthesis and use of catalytic DNA molecules with nucleotide-cleaving capabilities has not been disclosed or demonstrated. Therefore, the discoveries and inventions disclosed herein are particularly significant, in that they highlight the potential of *in vitro* evolution as a means of designing increasingly more efficient catalytic molecules, including enzymatic DNA molecules that cleave other nucleic acids, particularly RNA.

BRIEF SUMMARY OF THE INVENTION

20 The present invention thus contemplates a synthetic or engineered (i.e., non-naturally-occurring) catalytic DNA molecule (or enzymatic DNA molecule) capable of cleaving a substrate nucleic acid (NA) sequence at a defined cleavage site. The invention also contemplates an enzymatic DNA molecule having an endonuclease activity.

30 In one preferred variation, the endonuclease activity is specific for a nucleotide sequence defining a cleavage site comprising single-stranded nucleic acid in a substrate nucleic acid sequence. In another preferred variation, the cleavage site is double-stranded nucleic acid. Similarly, substrate nucleic acid sequences may be single-stranded, double-stranded, partially single- or double-stranded, looped, or any combination thereof.

35 In another contemplated embodiment, the substrate nucleic acid sequence includes one or more nucleotide analogues. In one variation, the substrate nucleic acid

-3-

sequence is a portion of, or attached to, a larger molecule.

In various embodiments, the larger molecule is selected from the group consisting of RNA, modified RNA, DNA, modified DNA, nucleotide analogs, or composites thereof. In another example, the larger molecule comprises a composite of a
5 nucleic acid sequence and a non-nucleic acid sequence.

In another embodiment, the invention contemplates that a substrate nucleic acid sequence includes one or more nucleotide analogs. A further variation contemplates that the single stranded nucleic acid comprises RNA, DNA, modified RNA, modified DNA, one or more nucleotide analogs, or any composite thereof. In one embodiment of
10 the disclosed invention, the endonuclease activity comprises hydrolytic cleavage of a phosphoester bond at the cleavage site.

In various preferred embodiments, the catalytic DNA molecules of the present invention are single-stranded in whole or in part. These catalytic DNA molecules may preferably assume a variety of shapes consistent with their catalytic activity. Thus, in
15 one variation, a catalytic DNA molecule of the present invention includes one or more hairpin loop structures. In yet another variation, a catalytic DNA molecule may assume a shape similar to that of "hammerhead" ribozymes. In still other embodiments, a catalytic DNA molecule may assume a conformation similar to that of *Tetrahymena thermophila* ribozymes, e.g., those derived from group I introns.

Similarly, preferred catalytic DNA molecules of the present invention are able to
20 demonstrate site-specific endonuclease activity irrespective of the original orientation of the substrate molecule. Thus, in one preferred embodiment, an enzymatic DNA molecule of the present invention is able to cleave a substrate nucleic acid sequence that is separate from the enzymatic DNA molecule -- i.e., it is not linked to the
25 DNAzyme. In another preferred embodiment, an enzymatic DNA molecule is able to cleave an attached substrate nucleic acid sequence -- i.e., it is able to perform a reaction similar to self-cleavage.

The invention also contemplates enzymatic DNA molecules (catalytic DNA molecules, deoxyribozymes or DNAzymes) having endonuclease activity, whereby the
30 endonuclease activity requires the presence of a divalent cation. In various preferred, alternative embodiments, the divalent cation is selected from the group consisting of Pb^{2+} , Mg^{2+} , Mn^{2+} , Zn^{2+} , and Ca^{2+} . Another variation contemplates that the endonuclease activity requires the presence of a monovalent cation. In such alternative
35 embodiments, the monovalent cation is preferably selected from the group consisting of Na^+ and K^+ .

In various preferred embodiments of the invention, an enzymatic DNA molecule comprises a nucleotide sequence selected from the group consisting of SEQ ID NO 3,

-4-

SEQ ID NO 14; SEQ ID NO 15; SEQ ID NO 16; SEQ ID NO 17; SEQ ID NO 18; SEQ ID NO 19; SEQ ID NO 20; SEQ ID NO 21; and SEQ ID NO 22. In other preferred embodiments, a catalytic DNA molecule of the present invention comprises a nucleotide sequence selected from the group consisting of SEQ ID NO 23; SEQ ID NO 24; SEQ ID NO 25; SEQ ID NO 26; SEQ ID NO 27; SEQ ID NO 28; SEQ ID NO 29; SEQ ID NO 30; SEQ ID NO 31; SEQ ID NO 32; SEQ ID NO 33; SEQ ID NO 34; SEQ ID NO 35; SEQ ID NO 36; SEQ ID NO 37; SEQ ID NO 38; and SEQ ID NO 39.

Another preferred embodiment contemplates that a catalytic DNA molecule of the present invention comprises a nucleotide sequence selected from the group consisting of SEQ ID NO 50 and SEQ ID NO 51. In yet another preferred embodiment, a catalytic DNA molecule of the present invention comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS 52 through 101. As disclosed herein, catalytic DNA molecules having sequences substantially similar to those disclosed herein are also contemplated. Thus, a wide variety of substitutions, deletions, insertions, duplications and other mutations may be made to the within-described molecules in order to generate a variety of other useful enzymatic DNA molecules; as long as said molecules display site-specific cleavage activity as disclosed herein, they are within the boundaries of this disclosure.

In a further variation of the present invention, an enzymatic DNA molecule of the present invention preferably has a substrate binding affinity of about $1 \mu\text{M}$ or less. In another embodiment, an enzymatic DNA molecule of the present invention binds substrate with a K_D of less than about $0.1 \mu\text{M}$.

The present invention also discloses enzymatic DNA molecules having useful turnover rates. In one embodiment, the turnover rate is less than 5 hr^{-1} ; in a preferred embodiment, the rate is less than about 2 hr^{-1} ; in a more preferred embodiment, the rate is less than about 1 hr^{-1} ; in an even more preferred embodiment, the turnover rate is about 0.6 hr^{-1} or less.

In still another embodiment, an enzymatic DNA molecule of the present invention displays a useful turnover rate wherein the k_{obs} is less than 1 min^{-1} , preferably less than 0.1 min^{-1} ; more preferably, less than 0.01 min^{-1} ; and even more preferably, less than 0.005 min^{-1} . In one variation, the value of k_{obs} is approximately 0.002 min^{-1} or less.

The present invention also contemplates embodiments in which the catalytic rate of the disclosed DNA enzymes is fully optimized. Thus, in various preferred embodiments, the K_m for reactions enhanced by the presence of Mg^{2+} is approximately $0.5\text{-}20 \text{ mM}$, preferably about $1\text{-}10 \text{ mM}$, and more preferably about $2\text{-}5 \text{ mM}$.

The present invention also contemplates an embodiment whereby the nucleotide

sequence defining the cleavage site comprises at least one nucleotide. In various other preferred embodiments, a catalytic DNA molecule of the present invention is able to recognize and cleave a nucleotide sequence defining a cleavage site of two or more nucleotides.

5 In various preferred embodiments, an enzymatic DNA molecule of the present invention comprises a conserved core flanked by one or more substrate binding regions. In one embodiment, an enzymatic DNA molecule includes first and second substrate binding regions. In another embodiment, an enzymatic DNA molecule includes two or more substrate binding regions.

10 As noted previously, preferred catalytic DNA molecules of the present invention may also include a conserved core. In one preferred embodiment, the conserved core comprises one or more conserved regions. In other preferred variations, the one or more conserved regions include a nucleotide sequence selected from the group consisting of CG; CGA; AGCG; AGCCG; CAGCGAT; CTTGTTT; and CTTATTT (see, e.g., Fig. 3).

15 In one embodiment of the invention, an enzymatic DNA molecule of the present invention further comprises one or more variable or spacer nucleotides between the conserved regions in the conserved core. In another embodiment, an enzymatic DNA molecule of the present invention further comprises one or more variable or spacer nucleotides between the conserved core and the substrate binding region.

20 In one variation, the first substrate binding region preferably includes a nucleotide sequence selected from the group consisting of CATCTCT; GCTCT; TTGCTTTTT; TGTCTTCTC; TTGCTGCT; GCCATGCTTT (SEQ ID NO 40); CTCTATTTCT (SEQ ID NO 41); GTCGGCA; CATCTCTTC; and ACTTCT. In another preferred variation, the second substrate binding region includes a nucleotide sequence selected from the
25 group consisting of TATGTGACGCTA (SEQ ID NO 42); TATAGTCGTA (SEQ ID NO 43); ATAGCGTATTA (SEQ ID NO 44); ATAGTTACGTCAT (SEQ ID NO 45); AATAGTGAAGTGTT (SEQ ID NO 46); TATAGTGTA; ATAGTCGGT; ATAGGCCCGGT (SEQ ID NO 47); AATAGTGAGGCTTG (SEQ ID NO 48); and ATGNTG.

30 In various embodiments of the present invention, the substrate binding regions vary in length. Thus, for example, a substrate binding region may comprise a single nucleotide to dozens of nucleotides. However, it is understood that substrate binding regions of about 3-25 nucleotides in length, preferably about 3-15 nucleotides in length, and more preferably about 3-10 nucleotides in length are particularly preferred. In
35 various embodiments, the individual nucleotides in the substrate binding regions are able to form complementary base pairs with the nucleotides of the substrate molecules; in other embodiments, noncomplementary base pairs are formed. A mixture of complementary and noncomplementary base pairing is also contemplated as falling

-6-

within the scope of the disclosed embodiments of the invention.

In another preferred embodiment, a catalytic DNA molecule of the present invention may further comprise a third substrate binding region. In some preferred embodiments, the third region includes a nucleotide sequence selected from the group consisting of TGTT; TGTTA; and TGTTAG. Another preferred embodiment of the present invention discloses an enzymatic DNA molecule further comprising one or more variable or "spacer" regions between the substrate binding regions.

In another disclosed embodiment, the present invention contemplates a purified, synthetic enzymatic DNA molecule separated from other DNA molecules and oligonucleotides, the enzymatic DNA molecule having an endonuclease activity, wherein the endonuclease activity is specific for a nucleotide sequence defining a cleavage site comprising single- or double-stranded nucleic acid in a substrate nucleic acid sequence. In one variation, a synthetic (or engineered) enzymatic DNA molecule having an endonuclease activity is disclosed, wherein the endonuclease activity is specific for a nucleotide sequence defining a cleavage site consisting essentially of a single- or double-stranded region of a substrate nucleic acid sequence.

In yet another embodiment, the invention contemplates an enzymatic DNA molecule comprising a deoxyribonucleotide polymer having a catalytic activity for hydrolyzing a nucleic acid-containing substrate to produce substrate cleavage products. In one variation, the hydrolysis takes place in a site-specific manner. As noted previously, the polymer may be single-stranded, double-stranded, or some combination of both.

The invention further contemplates that the substrate comprises a nucleic acid sequence. In various embodiments, the nucleic acid sequence substrate comprises RNA, modified RNA, DNA, modified DNA, one or more nucleotide analogs, or composites of any of the foregoing. One embodiment contemplates that the substrate includes a single-stranded segment; still another embodiment contemplates that the substrate is double-stranded.

The present invention also contemplates an enzymatic DNA molecule comprising a deoxyribonucleotide polymer having a catalytic activity for hydrolyzing a nucleic acid-containing substrate to produce a cleavage product. In one variation, the enzymatic DNA molecule has an effective binding affinity for the substrate and lacks an effective binding affinity for the cleavage product.

In one preferred embodiment, the invention discloses a non-naturally-occurring enzymatic DNA molecule comprising a nucleotide sequence defining a conserved core flanked by recognition domains, variable regions, and spacer regions. Thus, in one preferred embodiment, the nucleotide sequence defines a first variable region contiguous

-7-

or adjacent to the 5'-terminus of the molecule, a first recognition domain located 3'-terminal to the first variable region, a first spacer region located 3'-terminal to the first recognition domain, a first conserved region located 3'-terminal to the first spacer region, a second spacer region located 3'-terminal to the first conserved region, a second conserved region located 3'-terminal to the second spacer region, a second recognition domain located 3'-terminal to the second conserved region, and a second variable region located 3'-terminal to the second recognition domain.

In another embodiment, the nucleotide sequence preferably defines a first variable region contiguous or adjacent to the 5'-terminus of the molecule, a first recognition domain located 3'-terminal to the first variable region, a first spacer region located 3'-terminal to the first recognition domain, a first conserved region located 3'-terminal to the first spacer region, a second spacer region located 3'-terminal to the first conserved region, a second conserved region located 3'-terminal to the second spacer region, a second recognition domain located 3'-terminal to the second conserved region, a second variable region located 3'-terminal to the second recognition domain, and a third recognition domain located 3'-terminal to the second variable region.

In one variation of the foregoing, the molecule includes a conserved core region flanked by two substrate binding domains; in another, the conserved core region comprises one or more conserved domains. In other preferred embodiments, the conserved core region further comprises one or more variable or spacer nucleotides. In yet another embodiment, an enzymatic DNA molecule of the present invention further comprises one or more spacer regions.

The present invention further contemplates a wide variety of compositions. For example, compositions including an enzymatic DNA molecule as described hereinabove are disclosed and contemplated herein. In one alternative embodiment, a composition according to the present invention comprises two or more populations of enzymatic DNA molecules as described above, wherein each population of enzymatic DNA molecules is capable of cleaving a different sequence in a substrate. In another variation, a composition comprises two or more populations of enzymatic DNA molecules as described hereinabove, wherein each population of enzymatic DNA molecules is capable of recognizing a different substrate. In various embodiments, it is also preferred that compositions include a monovalent or divalent cation.

The present invention further contemplates methods of generating, selecting, and isolating enzymatic DNA molecules of the present invention. In one variation, a method of selecting enzymatic DNA molecules that cleave a nucleic acid sequence (e.g., RNA) at a specific site comprises the following steps: (a) obtaining a population of putative enzymatic DNA molecules -- whether the sequences are naturally-occurring or

-8-

synthetic -- and preferably, they are single-stranded DNA molecules; (b) admixing nucleotide-containing substrate sequences with the aforementioned population of DNA molecules to form an admixture; (c) maintaining the admixture for a sufficient period of time and under predetermined reaction conditions to allow the putative enzymatic DNA molecules in the population to cause cleavage of the substrate sequences, thereby
5 producing substrate cleavage products; (d) separating the population of DNA molecules from the substrate sequences and substrate cleavage products; and (e) isolating DNA molecules that cleave substrate nucleic acid sequences (e.g., RNA) at a specific site from the population.

10 In a further variation of the foregoing method, the DNA molecules that cleave substrate nucleic acid sequences at a specific site are tagged with an immobilizing agent. In one example, the agent comprises biotin.

In yet another variation of the aforementioned method, one begins by selecting a sequence -- e.g., a predetermined "target" nucleotide sequence -- that one wishes to
15 cleave using an enzymatic DNA molecule engineered for that purpose. Thus, in one embodiment, the pre-selected (or predetermined) "target" sequence is used to generate a population of DNA molecules capable of cleaving substrate nucleic acid sequences at a specific site via attaching or "tagging" it to a deoxyribonucleic acid sequence containing one or more randomized sequences or segments. In one variation, the
20 randomized sequence is about 40 nucleotides in length; in another variation, the randomized sequence is about 50 nucleotides in length. Randomized sequences that are 1-40, 40-50, and 50-100 nucleotides in length are also contemplated by the present invention.

In one embodiment of the present invention, the nucleotide sequence used to
25 generate a population of enzymatic DNA molecules is selected from the group consisting of SEQ ID NO 4, 23, 50 AND 51. In another embodiment, the "target" or "substrate" nucleotide sequence comprises a sequence of one or more ribonucleotides -- see, e.g., the relevant portions of SEQ ID NOS 4 and 23, and SEQ ID NO 49. It is also contemplated by the present invention that a useful "target" or "substrate" nucleotide
30 sequence may comprise DNA, RNA, or a composite thereof.

The invention also contemplates methods as described above, wherein the
isolating step further comprises exposing the tagged DNA molecules to a solid surface having avidin linked thereto, whereby the tagged DNA molecules become attached to
the solid surface. As before, the substrate may be RNA, DNA, a composite of both, or
35 a molecule including nucleotide sequences.

The present invention also contemplates a method for specifically cleaving a substrate nucleic acid sequence at a particular cleavage site, comprising the steps of (a)

-9-

providing an enzymatic DNA molecule capable of cleaving a substrate nucleic acid sequence at a specific cleavage site; and (b) contacting the enzymatic DNA molecule with the substrate nucleic acid sequence to cause specific cleavage of the nucleic acid sequence at the cleavage site. In one variation, the enzymatic DNA molecule is a non-naturally-occurring (or synthetic) DNA molecule. In another variation, the enzymatic DNA molecule is single-stranded.

In still another variation of the foregoing method, the substrate comprises a nucleic acid. In various embodiments, the substrate nucleic acid comprises RNA, modified RNA, DNA, modified DNA, one or more nucleotide analogs, or composites of any of the foregoing. In yet another embodiment, the specific cleavage is caused by the endonuclease activity of the enzymatic DNA molecule. Alteration of reaction conditions -- e.g., the adjustment of pH, temperature, percent cation, percent enzyme, percent substrate, and percent product -- is also contemplated herein.

The present invention also contemplates a method of cleaving a phosphoester bond, comprising (a) admixing an catalytic DNA molecule capable of cleaving a substrate nucleic acid sequence at a defined cleavage site with a phosphoester bond-containing substrate, to form a reaction admixture; and (b) maintaining the admixture under predetermined reaction conditions to allow the enzymatic DNA molecule to cleave the phosphoester bond, thereby producing a population of substrate products. In one embodiment, the enzymatic DNA molecule is able to cleave the phosphoester bond in a site-specific manner. In another embodiment, the method further comprises the steps of (c) separating the products from the catalytic DNA molecule; and (d) adding additional substrate to the enzymatic DNA molecule to form a new reaction admixture.

The present invention also contemplates methods of engineering enzymatic DNA molecules that cleave phosphoester bonds. One exemplary method comprises the following steps: (a) obtaining a population of single-stranded DNA molecules; (b) introducing genetic variation into the population to produce a variant population; (c) selecting individuals from the variant population that meet predetermined selection criteria; (d) separating the selected individuals from the remainder of the variant population; and (e) amplifying the selected individuals.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a selective amplification scheme for isolation of DNAs that cleave a target RNA phosphoester. As shown, double-stranded DNA that contains a stretch of 50 random nucleotides (the molecule with "N₅₀" indicated above it) is amplified by PCR, employing a 5'-biotinylated DNA primer that is terminated at the 3' end by an adenosine ribonucleotide (rA). (The biotin label is indicated via the encircled letter "B".) This primer is extended by *Taq* polymerase to yield a DNA product that

-10-

contains a single embedded ribonucleotide. The resulting double-stranded DNA is immobilized on a streptavidin matrix and the unbiotinylated DNA strand is removed by washing with 0.2 N NaOH. After re-equilibrating the column with a buffered solution, the column is washed with the same solution with added 1 mM PbOAc. DNAs that undergo Pb²⁺-dependent self-cleavage are released from the column, collected in the eluant, and amplified by PCR. The PCR products are then used to initiate the next round of selective amplification.

Figure 2 illustrates self-cleavage activity of the starting pool of DNA (G0) and populations obtained after the first through fifth rounds of selection (G1 - G5), in the presence of lead cation (Pb²⁺). The symbol Pre represents 108-nucleotide precursor DNA (SEQ ID NO 4); Clv, 28-nucleotide 5'-cleavage product (SEQ ID NO 5); and M, primer 3a (SEQ ID NO 6), which corresponds in length to the 5'-cleavage product.

Figure 3 illustrates the sequence alignment of individual variants isolated from the population after five rounds of selection. The fixed substrate domain is shown at the top, with the target riboadenylate identified via an inverted triangle. Substrate nucleotides that are commonly involved in presumed base-pairing interactions are indicated by vertical bars. Sequences corresponding to the 50 initially-randomized nucleotides are aligned antiparallel to the substrate domain. All of the variants are 3'-terminated by the fixed sequence 5'-CGGTAAGCTTGGCAC-3' (not shown; SEQ ID NO 1). Nucleotides within the initially-randomized region that are presumed to form base pairs with the substrate domain are indicated on the right and left sides of the Figure; the putative base-pair-forming regions of the enzymatic DNA molecules are individually boxed in each sequence shown. Conserved regions are illustrated via the two large, centrally-located boxes.

Figures 4A and 4B illustrate DNA-catalyzed cleavage of an RNA phosphoester in an intermolecular reaction that proceeds with catalytic turnover. Figure 4A is a diagrammatic representation of the complex formed between the 19mer substrate (3'-TCACTATrAGGAAGAGATGG-5', SEQ ID NO 2) and 38mer DNA enzyme (5'-ACACATCTCTGAAGTAGCGCCCGTATAGTGACGCTA-3', SEQ ID NO 3). The substrate contains a single adenosine ribonucleotide ("rA", adjacent to the arrow), flanked by deoxyribonucleotides. The synthetic DNA enzyme is a 38-nucleotide portion of the most frequently occurring variant shown in Fig. 3. Highly-conserved nucleotides located within the putative catalytic domain are "boxed". As illustrated, one conserved sequence is "AGCG", while another is "CG" (reading in the 5'→3' direction).

Figure 4B shows an Eadie-Hofstee plot used to determine K_m (negative slope) and V_{max} (y-intercept) for DNA-catalyzed cleavage of [5'-³²P]-labeled substrate under conditions identical to those employed during *in vitro* selection. Initial rates of cleavage

-11-

were determined for reactions involving 5 nM DNA enzyme and either 0.125, 0.5, 1, 2, or 4 μ M substrate.

Figure 5 is a photographic representation showing a polyacrylamide gel demonstrating specific endoribonuclease activity of four families of selected catalytic DNAs. Selection of a Pb^{2+} -dependent family of molecules was repeated in a side-by-side fashion as a control (first group). In the second group, Zn^{2+} is used as the cation; in group three, the cation is Mn^{2+} ; and in the fourth group, the cation is Mg^{2+} . A fifth site on the gel consists of the cleavage product alone, as a marker.

As noted, there are three lanes within each of the aforementioned four groups. In each group of three lanes, the first lane shows the lack of activity of the selected population in the absence of the metal cation, the second lane shows the observed activity in the presence of the metal cation, and the third lane shows the lack of activity of the starting pool (G0).

Figures 6A and 6B provide two-dimensional illustrations of a "progenitor" catalytic DNA molecule and one of several catalytic DNA molecules obtained via the selective amplification methods disclosed herein, respectively. Figure 6A illustrates an exemplary molecule from the starting pool, showing the overall configuration of the molecules represented by SEQ ID NO 23. As illustrated, various complementary nucleotides flank the random (N_{40}) region. Figure 6B is a diagrammatic representation of one of the Mg^{2+} -dependent catalytic DNA molecules (or "DNAzymes") generated via the within-described procedures. The location of the ribonucleotide in the substrate nucleic acid is indicated via the arrow in both Figs. 6A and 6B.

Figure 7 illustrates some of the results of ten rounds of *in vitro* selective amplification carried out essentially as described in Example 5 hereinbelow. As shown, two sites and two families of catalysts emerged as displaying the most efficient cleavage of the target sequence. Cleavage conditions were essentially as indicated in Fig. 7, namely, 10mM Mg^{2+} , pH 7.5, and 37°C; data collected after the reaction ran for 2 hours is shown. Cleavage (%) is shown plotted against the number of generations (here, 0 through 10). The number/prevalence of catalytic DNA molecules capable of cleaving the target sequence at the indicated sites in the substrate is illustrated via the vertical bars, with cleavage at G!UAACUAGAGAU shown by the striped bars, and with cleavage at GUAACUA!GAGAU illustrated via the open (lightly-shaded) bars.

Figure 8 illustrates the nucleotide sequences, cleavage sites, and turnover rates of two catalytic DNA molecules of the present invention, clones 8-17 and 10-23. Reaction conditions were as shown, namely, 10mM Mg^{2+} , pH 7.5, and 37°C. The DNAzyme identified as clone 8-17 is illustrated on the left, with the site of cleavage of the RNA substrate indicated by the arrow. The substrate sequence (5' -

-12-

GGAAAAAGUAACUAGAGAUGGAAG - 3') -- which is separate from the DNAzyme (i.e., intermolecular cleavage is shown) -- is labeled as such. Similarly, the DNAzyme identified herein as 10-23 is shown on the right, with the site of cleavage of the RNA substrate indicated by the arrow. Again, the substrate sequence is indicated. For the 8-17 enzyme, the turnover rate was approximately 0.6 hr⁻¹; for the 10-23 enzyme, the turnover rate was approximately 1 hr⁻¹. Noncomplementary pairings are indicated with a closed circle (•), whereas complementary pairings are indicated with a vertical line (|).

Figure 9 further illustrates the nucleotide sequences, cleavage sites, and turnover rates of two catalytic DNA molecules of the present invention, clones 8-17 and 10-23. Reaction conditions were as shown, namely, 10mM Mg²⁺, pH 7.5, and 37°C. As in Fig. 8, the DNAzyme identified as clone 8-17 is illustrated on the left, with the site of cleavage of the RNA substrate indicated by the arrow. The substrate sequence (5' - GGAAAAAGUAACUAGAGAUGGAAG - 3') --which is separate from the DNAzyme (i.e., intermolecular cleavage is shown) -- is labeled as such. Similarly, the DNAzyme identified herein as 10-23 is shown on the right, with the site of cleavage of the RNA substrate indicated by the arrow. Again, the substrate sequence is indicated. For the 8-17 enzyme, k_{obs} was approximately 0.002 min⁻¹; for the 10-23 enzyme, the value of k_{obs} was approximately 0.01 min⁻¹. Noncomplementary pairings are indicated with a closed circle (•), whereas complementary pairings are indicated with a vertical line (|).

DETAILED DESCRIPTION

A. Definitions

As used herein, the term "deoxyribozyme" is used to describe a DNA-containing nucleic acid that is capable of functioning as an enzyme. In the present disclosure, the term "deoxyribozyme" includes endoribonucleases and endodeoxyribonucleases, although deoxyribozymes with endoribonuclease activity are particularly preferred. Other terms used interchangeably with deoxyribozyme herein are "enzymatic DNA molecule", "DNAzyme", or "catalytic DNA molecule", which terms should all be understood to include enzymatically active portions thereof, whether they are produced synthetically or derived from organisms or other sources.

The term "enzymatic DNA molecules" also includes DNA molecules that have complementarity in a substrate-binding region to a specified oligonucleotide target or substrate; such molecules also have an enzymatic activity which is active to specifically cleave the oligonucleotide substrate. Stated in another fashion, the enzymatic DNA molecule is capable of cleaving the oligonucleotide substrate intermolecularly. This complementarity functions to allow sufficient hybridization of the enzymatic DNA molecule to the substrate oligonucleotide to allow the intermolecular cleavage of the substrate to occur. While one-hundred percent (100%) complementarity is preferred,

-13-

complementarity in the range of 75-100% is also useful and contemplated by the present invention.

Enzymatic DNA molecules of the present invention may alternatively be described as having nuclease or ribonuclease activity. These terms may be used interchangeably herein.

The term "enzymatic nucleic acid" as used herein encompasses enzymatic RNA or DNA molecules, enzymatic RNA-DNA polymers, and enzymatically active portions or derivatives thereof, although enzymatic DNA molecules are a particularly preferred class of enzymatically active molecules according to the present invention.

The term "endodeoxyribonuclease", as used herein, is an enzyme capable of cleaving a substrate comprised predominantly of DNA. The term "endoribonuclease", as used herein, is an enzyme capable of cleaving a substrate comprised predominantly of RNA.

As used herein, the term "base pair" (bp) is generally used to describe a partnership of adenine (A) with thymine (T) or uracil (U), or of cytosine (C) with guanine (G), although it should be appreciated that less-common analogs of the bases A, T, C, and G (as well as U) may occasionally participate in base pairings. Nucleotides that normally pair up when DNA or RNA adopts a double stranded configuration may also be referred to herein as "complementary bases".

"Complementary nucleotide sequence" generally refers to a sequence of nucleotides in a single-stranded molecule or segment of DNA or RNA that is sufficiently complementary to that on another single oligonucleotide strand to specifically hybridize to it with consequent hydrogen bonding.

"Nucleotide" generally refers to a monomeric unit of DNA or RNA consisting of a sugar moiety (pentose), a phosphate group, and a nitrogenous heterocyclic base. The base is linked to the sugar moiety via the glycosidic carbon (1' carbon of the pentose) and that combination of base and sugar is a "nucleoside". When the nucleoside contains a phosphate group bonded to the 3' or 5' position of the pentose, it is referred to as a nucleotide. A sequence of operatively linked nucleotides is typically referred to herein as a "base sequence" or "nucleotide sequence", and their grammatical equivalents, and is represented herein by a formula whose left to right orientation is in the conventional direction of 5'-terminus to 3'-terminus, unless otherwise specified.

"Nucleotide analog" generally refers to a purine or pyrimidine nucleotide that differs structurally from A, T, G, C, or U, but is sufficiently similar to substitute for the normal nucleotide in a nucleic acid molecule. As used herein, the term "nucleotide analog" encompasses altered bases, different or unusual sugars (i.e. sugars other than the "usual" pentose), or a combination of the two. A listing of exemplary analogs

-14-

wherein the base has been altered is provided in section C hereinbelow.

"Oligonucleotide or polynucleotide" generally refers to a polymer of single- or double-stranded nucleotides. As used herein, "oligonucleotide" and its grammatical equivalents will include the full range of nucleic acids. An oligonucleotide will typically refer to a nucleic acid molecule comprised of a linear strand of ribonucleotides. The exact size will depend on many factors, which in turn depends on the ultimate conditions of use, as is well known in the art.

As used herein, the term "physiologic conditions" is meant to suggest reaction conditions emulating those found in mammalian organisms, particularly humans. While variables such as temperature, availability of cations, and pH ranges may vary as described in greater detail below, "physiologic conditions" generally comprise a temperature of about 35-40°C, with 37°C being particularly preferred, as well as a pH of about 7.0-8.0, with 7.5 being particularly preferred, and further comprise the availability of cations, preferably divalent and/or monovalent cations, with a concentration of about 2-15 mM Mg²⁺ and 0-1.0 M Na⁺ being particularly preferred. "Physiologic conditions", as used herein, may optionally include the presence of free nucleoside cofactor. As noted previously, preferred conditions are described in greater detail below.

B. Enzymatic DNA Molecules

In various embodiments, an enzymatic DNA molecule of the present invention may combine one or more modifications or mutations including additions, deletions, and substitutions. In alternative embodiments, such mutations or modifications may be generated using methods which produce random or specific mutations or modifications. These mutations may, for example, change the length of, or alter the nucleotide sequence of, a loop, a spacer region or the recognition sequence (or domain). One or more mutations within one catalytically active enzymatic DNA molecule may be combined with the mutation(s) within a second catalytically active enzymatic DNA molecule to produce a new enzymatic DNA molecule containing the mutations of both molecules.

In other preferred embodiments, an enzymatic DNA molecule of the present invention may have random mutations introduced into it using a variety of methods well known to those skilled in the art. For example, the methods described by Cadwell and Joyce (PCR Methods and Applications 2: 28-33 (1992)) are particularly preferred for use as disclosed herein, with some modifications, as described in the Examples that follow. (Also see Cadwell and Joyce, PCR Methods and Applications 3 (Suppl.): S136-S140 (1994).) According to this modified PCR method, random point mutations may be introduced into cloned genes.

-15-

The aforementioned methods have been used, for example, to mutagenize genes encoding ribozymes with a mutation rate of $0.66\% \pm 0.13\%$ (95% confidence interval) per position, as determined by sequence analysis, with no strong preferences observed with respect to the type of base substitution. This allows the introduction of random mutations at any position in the enzymatic DNA molecules of the present invention.

Another method useful in introducing defined or random mutations is disclosed in Joyce and Inoue, Nucleic Acids Research 17: 711-722 (1989). This latter method involves excision of a template (coding) strand of a double-stranded DNA, reconstruction of the template strand with inclusion of mutagenic oligonucleotides, and subsequent transcription of the partially-mismatched template. This allows the introduction of defined or random mutations at any position in the molecule by including polynucleotides containing known or random nucleotide sequences at selected positions.

Enzymatic DNA molecules of the present invention may be of varying lengths and folding patterns, as appropriate, depending on the type and function of the molecule. For example, enzymatic DNA molecules may be about 15 to about 400 or more nucleotides in length, although a length not exceeding about 250 nucleotides is preferred, to avoid limiting the therapeutic usefulness of molecules by making them too large or unwieldy. In various preferred embodiments, an enzymatic DNA molecule of the present invention is at least about 20 nucleotides in length and, while useful molecules may exceed 100 nucleotides in length, preferred molecules are generally not more than about 100 nucleotides in length.

In various therapeutic applications, enzymatic DNA molecules of the present invention comprise the enzymatically active portions of deoxyribozymes. In various embodiments, enzymatic DNA molecules of the present invention preferably comprise not more than about 200 nucleotides. In other embodiments, a deoxyribozyme of the present invention comprises not more than about 100 nucleotides. In still other preferred embodiments, deoxyribozymes of the present invention are about 20-75 nucleotides in length, more preferably about 20-65 nucleotides in length. Other preferred enzymatic DNA molecules are about 10-50 nucleotides in length.

In other applications, enzymatic DNA molecules may assume configurations similar to those of "hammerhead" ribozymes. Such enzymatic DNA molecules are preferably no more than about 75-100 nucleotides in length, with a length of about 20-50 nucleotides being particularly preferred.

In general, if one intends to synthesize molecules for use as disclosed herein, the larger the enzymatic nucleic acid molecule is, the more difficult it is to synthesize. Those of skill in the art will certainly appreciate these design constraints. Nevertheless, such larger molecules remain within the scope of the present invention.

-16-

It is also to be understood that an enzymatic DNA molecule of the present invention may comprise enzymatically active portions of a deoxyribozyme or may comprise a deoxyribozyme with one or more mutations, e.g., with one or more base-pair-forming sequences or spacers absent or modified, as long as such deletions, additions or modifications do not adversely impact the molecule's ability to perform as an enzyme.

The recognition domain of an enzymatic DNA molecule of the present invention typically comprises two nucleotide sequences flanking a catalytic domain, and typically contains a sequence of at least about 3 to about 30 bases, preferably about 6 to about 15 bases, which are capable of hybridizing to a complementary sequence of bases within the substrate nucleic acid giving the enzymatic DNA molecule its high sequence specificity. Modification or mutation of the recognition site via well-known methods allows one to alter the sequence specificity of an enzymatic nucleic acid molecule. (See, e.g., Joyce et al., Nucleic Acids Research 17: 711-712 (1989).)

Enzymatic nucleic acid molecules of the present invention also include those with altered recognition sites or domains. In various embodiments, these altered recognition domains confer unique sequence specificities on the enzymatic nucleic acid molecule including such recognition domains. The exact bases present in the recognition domain determine the base sequence at which cleavage will take place. Cleavage of the substrate nucleic acid occurs within the recognition domain. This cleavage leaves a 2', 3', or 2',3'-cyclic phosphate group on the substrate cleavage sequence and a 5' hydroxyl on the nucleotide that was originally immediately 3' of the substrate cleavage sequence in the original substrate. Cleavage can be redirected to a site of choice by changing the bases present in the recognition sequence (internal guide sequence). See Murphy et al., Proc. Natl. Acad. Sci. USA 86: 9218-9222 (1989).

Moreover, it may be useful to add a polyamine to facilitate recognition and binding between the enzymatic DNA molecule and its substrate. Examples of useful polyamines include spermidine, putrescine or spermine. A spermidine concentration of about 1 mM may be effective in particular embodiments, while concentrations ranging from about 0.1 mM to about 10 mM may also be useful.

In various alternative embodiments, an enzymatic DNA molecule of the present invention has an enhanced or optimized ability to cleave nucleic acid substrates, preferably RNA substrates. As those of skill in the art will appreciate, the rate of an enzyme-catalyzed reaction varies depending upon the substrate and enzyme concentrations and, in general, levels off at high substrate or enzyme concentrations. Taking such effects into account, the kinetics of an enzyme-catalyzed reaction may be described in the following terms, which define the reaction.

-17-

The enhanced or optimized ability of an enzymatic DNA molecule of the present invention to cleave an RNA substrate may be determined in a cleavage reaction with varying amounts of labeled RNA substrate in the presence of enzymatic DNA molecule. The ability to cleave the substrate is generally defined by the catalytic rate (k_{cat}) divided by the Michaelis constant (K_M). The symbol k_{cat} represents the maximal velocity of an enzyme reaction when the substrate approaches a saturation value. K_M represents the substrate concentration at which the reaction rate is one-half maximal.

For example, values for K_M and k_{cat} may be determined in this invention by experiments in which the substrate concentration [S] is in excess over enzymatic DNA molecule concentration [E]. Initial rates of reaction (v_o) over a range of substrate concentrations are estimated from the initial linear phase, generally the first 5% or less of the reaction. Data points are fit by a least squares method to a theoretical line given by the equation: $v = -K_M(v_o/[S]) + V_{max}$. Thus, k_{cat} and K_M are determined by the initial rate of reaction, v_o , and the substrate concentration [S].

In various alternative embodiments, an enzymatic DNA molecule of the present invention has an enhanced or optimized ability to cleave nucleic acid substrates, preferably RNA substrates. In preferred embodiments, the enhanced or optimized ability of an enzymatic DNA molecule to cleave RNA substrates shows about a 10- to 10⁹-fold improvement over the uncatalyzed rate. In more preferred embodiments, an enzymatic DNA molecule of the present invention is able to cleave RNA substrates at a rate that is about 10³- to 10⁷-fold improved over "progenitor" species. In even more preferred embodiments, the enhanced or optimized ability to cleave RNA substrates is expressed as a 10⁴- to 10⁶-fold improvement over the progenitor species. One skilled in the art will appreciate that the enhanced or optimized ability of an enzymatic DNA molecule to cleave nucleic acid substrates may vary depending upon the selection constraints applied during the *in vitro* evolution procedure of the invention.

Various preferred methods of modifying deoxyribozymes and other enzymatic DNA molecules and nucleases of the present invention are further described in Examples 1-3 hereinbelow.

C. Nucleotide Analogs

As noted above, the term "nucleotide analog" as used herein generally refers to a purine or pyrimidine nucleotide that differs structurally from A, T, G, C, or U, but is sufficiently similar to substitute for such "normal" nucleotides in a nucleic acid molecule. As used herein, the term "nucleotide analog" encompasses altered bases, different (or unusual) sugars, altered phosphate backbones, or any combination of these alterations. Examples of nucleotide analogs useful according to the present invention include those listed in the following Table, most of which are found in the approved listing of modified

bases at 37 CFR §1.822 (which is incorporated herein by reference).

Table 1
Nucleotide Analogs

	Abbreviation	Description
5	ac4c	4-acetylcytidine
	chm5u	5-(carboxyhydroxymethyl)uridine
	cm	2'-O-methylcytidine
10	cmnm5s2u	5-carboxymethylaminomethyl-2-thiouridine
	d	dihydrouridine
	fm	2'-O-methylpseudouridine
	galq	β , D-galactosylqueosine
	gm	2'-O-methylguanosine
15	l	inosine
	i6a	N6-isopentenyladenosine
	m1a	1-methyladenosine
	m1f	1-methylpseudouridine
	m1g	1-methylguanosine
20	m1l	1-methylinosine
	m22g	2,2-dimethylguanosine
	m2a	2-methyladenosine
	m2g	2-methylguanosine
	m3c	3-methylcytidine
25	m5c	5-methylcytidine
	m6a	N6-methyladenosine
	m7g	7-methylguanosine
	mam5u	5-methylaminomethyluridine
	mam5s2u	5-methoxyaminomethyl-2-thiouridine
30	manq	β , D-mannosylmethyluridine
	mcm5s2u	5-methoxycarbonylmethyluridine
	mo5u	5-methoxyuridine
	ms2i6a	2-methylthio-N6-isopentenyladenosine
	ms2t6a	N-((9- β -D-ribofuranosyl-2-methylthiopurine-6-yl)carbamoyl)threonine
35	mt6a	N-((9- β -D-ribofuranosylpurine-6-yl)N-methyl-carbamoyl)threonine

-19-

(Table 1, cont'd)

	Abbreviation	Description
5	mv	uridine-5-oxyacetic acid methylester
	o5u	uridine-5-oxyacetic acid (v)
	osyw	wybutoxosine
	p	pseudouridine
	q	queosine
10	s2c	2-thiocytidine
	s2t	5-methyl-2-thiouridine
	s2u	2-thiouridine
	s4u	4-thiouridine
	t	5-methyluridine
15	t6a	N-((9-β-D-ribofuranosylpurine-6-yl)carbamoyl)threoninetm 2'-O-methyl-5-methyluridine
	um	2'-O-methyluridine
	yw	wybutosine
	x	3-(3-amino-3-carboxypropyl)uridine, (acp3)u
20	araU	β, D-arabinosyl
	araT	β, D-arabinosyl

25 Other useful analogs include those described in published international application no. WO 92/20823 (the disclosures of which are incorporated herein by reference), or analogs made according to the methods disclosed therein. Analogs described in DeMesmaeker, et al., Angew. Chem. Int. Ed. Engl. **33**: 226-229 (1994); DeMesmaeker, et al., Synlett: 733-736 (Oct. 1993); Nielsen, et al., Science **254**: 1497-30 1500 (1991); and Idziak, et al., Tetrahedron Letters **34**: 5417-5420 (1993) are also useful according to the within-disclosed invention and said disclosures are incorporated by reference herein.

D. Methods of Engineering Enzymatic DNA Molecules

35 The present invention also contemplates methods of producing nucleic acid molecules having a predetermined activity. In one preferred embodiment, the nucleic acid molecule is an enzymatic DNA molecule. In another variation, the desired activity is a catalytic activity.

In one embodiment, the present invention contemplates methods of synthesizing

-20-

enzymatic DNA molecules that may then be "engineered" to catalyze a specific or predetermined reaction. Methods of preparing enzymatic DNA molecules are described herein; see, e.g., Examples 1-3 hereinbelow. In other embodiments, an enzymatic DNA molecule of the present invention may be engineered to bind small molecules or ligands, such as adenosine triphosphate (ATP). (See, e.g., Sassanfar, et al., Nature 364: 550-553 (1993).)

In another embodiment, the present invention contemplates that a population of enzymatic DNA molecules may be subjected to mutagenizing conditions to produce a diverse population of mutant enzymatic DNA molecules (which may alternatively be called "deoxyribozymes" or "DNAzymes"). Thereafter, enzymatic DNA molecules having desired characteristics are selected and/or separated from the population and are subsequently amplified.

Alternatively, mutations may be introduced in the enzymatic DNA molecule by altering the length of the recognition domains of the enzymatic DNA molecule. The recognition domains of the enzymatic DNA molecule associate with a complementary sequence of bases within a substrate nucleic acid sequence. Methods of altering the length of the recognition domains are known in the art and include PCR, for example; useful techniques are described further in the Examples below.

Alteration of the length of the recognition domains of an enzymatic DNA molecule may have a desirable effect on the binding specificity of the enzymatic DNA molecule. For example, an increase in the length of the recognition domains may increase binding specificity between the enzymatic DNA molecule and the complementary base sequences of an oligonucleotide in a substrate, or may enhance recognition of a particular sequence in a hybrid substrate. In addition, an increase in the length of the recognition domains may also increase the affinity with which it binds to substrate. In various embodiments, these altered recognition domains in the enzymatic DNA molecule confer increased binding specificity and affinity between the enzymatic DNA molecule and its substrate.

It has recently been noted that certain oligonucleotides are able to recognize and bind molecules other than oligonucleotides with complementary sequences. These oligonucleotides are often given the name "aptamers". For example, Ellington and Szostak describe RNA molecules that are able to bind a variety of organic dyes (Nature 346: 818-822 (1990)), while Bock, et al. describe ssDNA molecules that bind human thrombin (Nature 355: 564-566 (1992)). Similarly, Jellinek, et al. describe RNA ligands to basic fibroblast growth factor (PNAS USA 90: 11227-11231 (1993)). Thus, it is further contemplated herein that the catalytically active DNA enzymes of the present invention may be engineered according to the within-described methods to display a

-21-

variety of capabilities typically associated with aptamers.

One of skill in the art should thus appreciate that the enzymatic DNA molecules of this invention can be altered at any nucleotide sequence, such as the recognition domains, by various methods disclosed herein, including PCR and 3SR (self-sustained
5 sequence replication -- see Example 1 below). For example, additional nucleotides can be added to the 5' end of the enzymatic DNA molecule by including additional nucleotides in the primers.

Enzymatic DNA molecules of the present invention may also be prepared or engineered in a more non-random fashion via use of methods such as site-directed
10 mutagenesis. For example, site-directed mutagenesis may be carried out essentially as described in Morinaga, et al., Biotechnology 2: 636 (1984), modified as described herein, for application to deoxyribozymes. Useful methods of engineering enzymatic DNA molecules are further described in the Examples below.

In one disclosed embodiment, an enzymatic DNA molecule of the present
15 invention comprises a conserved core flanked by two substrate binding (or recognition) domains or sequences that interact with the substrate through base-pairing interactions. In various embodiments, the conserved core comprises one or more conserved domains or sequences. In another variation, an enzymatic DNA molecule further comprises a "spacer" region (or sequence) between the regions (or sequences) involved in base
20 pairing. In still another variation, the conserved core is "interrupted" at various intervals by one or more less-conserved variable or "spacer" nucleotides.

In various embodiments, the population of enzymatic DNA molecules is made up of at least 2 different types of deoxyribozyme molecules. For example, in one variation, the molecules have differing sequences. In another variation, the deoxyribozymes are
25 nucleic acid molecules having a nucleic acid sequence defining a recognition domain that is contiguous or adjacent to the 5'-terminus of the nucleotide sequence. In various alternative embodiments, enzymatic DNA molecules of the present invention may further comprise one or more spacer regions located 3'-terminal to the recognition domains, one or more loops located 3'-terminal to the recognition domains and/or spacer regions. In
30 other variations, a deoxyribozyme of the present invention may comprise one or more regions which are capable of hybridizing to other regions of the same molecule. Other characteristics of enzymatic DNA molecules produced according to the presently-disclosed methods are described elsewhere herein.

In other embodiments, mutagenizing conditions include conditions that introduce
35 either defined or random nucleotide substitutions within an enzymatic DNA molecule. Examples of typical mutagenizing conditions include conditions disclosed in other parts of this specification and the methods described by Joyce et al., Nucl. Acids Res. 17:

711-722 (1989); Joyce, Gene 82: 83-87(1989); and Beaudry and Joyce, Science 257: 635-41 (1992).

5 In still other embodiments, a diverse population of mutant enzymatic nucleic acid molecules of the present invention is one that contains at least 2 nucleic acid molecules that do not have the exact same nucleotide sequence. In other variations, from such a diverse population, an enzymatic DNA molecule or other enzymatic nucleic acid having a predetermined activity is then selected on the basis of its ability to perform the predetermined activity. In various embodiments, the predetermined activity comprises, without limitation, enhanced catalytic activity, decreased K_M , enhanced substrate
10 binding ability, altered substrate specificity, and the like.

Other parameters which may be considered aspects of enzyme performance include catalytic activity or capacity, substrate binding ability, enzyme turnover rate, enzyme sensitivity to feedback mechanisms, and the like. In certain aspects, substrate specificity may be considered an aspect of enzyme performance, particularly in
15 situations in which an enzyme is able to recognize and bind two or more competing substrates, each of which affects the enzyme's performance with respect to the other substrate(s).

Substrate specificity, as used herein, may refer to the specificity of an enzymatic nucleic acid molecule as described herein for a particular substrate, such as one
20 comprising ribonucleotides only, deoxyribonucleotides only, or a composite of both. Substrate molecules may also contain nucleotide analogs. In various embodiments, an enzymatic nucleic acid molecule of the present invention may preferentially bind to a particular region of a hybrid or non-hybrid substrate.

The term or parameter identified herein as "substrate specificity" may also
25 include sequence specificity; i.e., an enzymatic nucleic acid molecule of the present invention may "recognize" and bind to a nucleic acid substrate having a particular nucleic acid sequence. For example, if the substrate recognition domains of an enzymatic nucleic acid molecule of the present invention will only bind to substrate molecules having a series of one or two ribonucleotides (e.g., rA) in a row, then the
30 enzymatic nucleic acid molecule will tend not to recognize or bind nucleic acid substrate molecules lacking such a sequence.

With regard to the selection process, in various embodiments, selecting includes any means of physically separating the mutant enzymatic nucleic acids having a predetermined activity from the diverse population of mutant enzymatic nucleic acids.
35 Often, selecting comprises separation by size, by the presence of a catalytic activity, or by hybridizing the mutant nucleic acid to another nucleic acid, to a peptide, or some other molecule that is either in solution or attached to a solid matrix.

-23-

In various embodiments, the predetermined activity is such that the mutant enzymatic nucleic acid having the predetermined activity becomes labeled in some fashion by virtue of the activity. For example, the predetermined activity may be an enzymatic DNA molecule activity whereby the activity of the mutant enzymatic nucleic acid upon its substrate causes the mutant enzymatic nucleic acid to become covalently linked to it. The mutant enzymatic nucleic acid is then selected by virtue of the covalent linkage.

In other embodiments, selecting a mutant enzymatic nucleic acid having a predetermined activity includes amplification of the mutant enzymatic nucleic acid (see, e.g., Joyce, Gene 82: 83-87 (1989); Beaudry and Joyce, Science 257: 635-41 (1992)). Other methods of selecting an enzymatic nucleic acid molecule having a predetermined characteristic or activity are described in the Examples section.

E. Compositions

The invention also contemplates compositions containing one or more types or populations of enzymatic DNA molecules of the present invention; e.g., different types or populations may recognize and cleave different nucleotide sequences. Compositions may further include a ribonucleic acid-containing substrate. Compositions according to the present invention may further comprise lead ion, magnesium ion, or other divalent or monovalent cations, as discussed herein.

Preferably, the enzymatic DNA molecule is present at a concentration of about 0.05 μM to about 2 μM . Typically, the enzymatic DNA molecule is present at a concentration ratio of enzymatic DNA molecule to substrate of from about 1:5 to about 1:50. More preferably, the enzymatic DNA molecule is present in the composition at a concentration of about 0.1 μM to about 1 μM . Even more preferably, compositions contain the enzymatic DNA molecule at a concentration of about 0.1 μM to about 0.5 μM . Preferably, the substrate is present in the composition at a concentration of about 0.5 μM to about 1000 μM .

One skilled in the art will understand that there are many sources of nucleic acid-containing substrates including naturally-occurring and synthetic sources. Sources of suitable substrates include, without limitation, a variety of viral and retroviral agents, including HIV-1, HIV-2, HTLV-I, and HTLV-II.

Other suitable substrates include, without limitation, viral and retroviral agents including those comprising or produced by picornaviruses, hepadnaviridae (e.g., HBV, HCV), papillomaviruses (e.g., HPV), gammaherpesvirinae (e.g., EBV), lymphocryptoviruses, leukemia viruses (e.g., HTLV-I and -II), flaviviruses, togaviruses, herpesviruses (including alphaherpesviruses and betaherpesviruses), cytomegaloviruses (CMV), influenza viruses, and viruses and retroviruses contributing to immunodeficiency

-24-

diseases and syndromes (e.g., HIV-1 and -2). In addition, suitable substrates include viral and retroviral agents which infect non-human primates and other animals including, without limitation, the simian and feline immunodeficiency viruses and bovine leukemia viruses.

5 Magnesium ion, lead ion, or another suitable monovalent or divalent cation, as described previously, may also be present in the composition, at a concentration ranging from about 1-100 mM. More preferably, the preselected ion is present in the composition at a concentration of about 2 mM to about 50 mM, with a concentration of about 5 mM being particularly preferred. One skilled in the art will understand that the
10 ion concentration is only constrained by the limits of solubility of its source (e.g. magnesium) in aqueous solution and a desire to have the enzymatic DNA molecule present in the same composition in an active conformation.

 The invention also contemplates compositions containing an enzymatic DNA molecule of the present invention, hybrid deoxyribonucleotide-ribonucleotide molecules,
15 and magnesium or lead ion in concentrations as described hereinabove. As noted previously, other monovalent or divalent ions (e.g., Ca^{2+}) may be used in place of magnesium.

 Also contemplated by the present invention are compositions containing an enzymatic DNA molecule of the present invention, nucleic acid-containing substrate (e.g.
20 RNA), and a preselected ion at a concentration of greater than about 1 millimolar, wherein said substrate is greater in length than the recognition domains present on the enzymatic DNA molecule.

 In one variation, a composition comprises an enzymatic DNA molecule-substrate complex, wherein base pairing between an enzymatic DNA molecule and its substrate is
25 contiguous. In another embodiment, base pairing between an enzymatic DNA molecule and its substrate is interrupted by one or more noncomplementary pairs. In a variety of alternative embodiments, a composition of the present invention may further comprise a monovalent cation, a divalent cation, or both.

 In another variation, an enzymatic DNA molecule of the present invention is
30 capable of functioning efficiently in the presence or absence of a divalent cation. In one variation, a divalent cation is present and comprises Pb^{2+} , Mg^{2+} , Mn^{2+} , Zn^{2+} , or Ca^{2+} . Alternatively, an enzymatic DNA molecule of the present invention is capable of functioning efficiently in the presence or absence of monovalent cations. It is anticipated that monovalent or divalent cation concentrations similar to those described
35 herein for Pb^{2+} or Mg^{2+} will be useful as disclosed herein.

 Optionally, monovalent cations may also be present in addition to, or as "alternatives" for, divalent cations. For example, monovalent cations such as sodium

-25-

(Na⁺) or potassium (K⁺) may be present, either as dissociated ions or in the form of dissociable compounds such as NaCl or KCl.

In one embodiment, the concentration of monovalent cation present in the composition ranges from 0 - 1.0 M. In another embodiment, a monovalent cation is present in a concentration ranging from about 0-200 mM. In other embodiments, monovalent cations are present in a concentration ranging from about 1-100 mM. Alternatively, the concentration of monovalent cations ranges from about 2 mM - 50 mM. In still other embodiments, the concentration ranges from about 2 mM - 25 mM.

F. Methods of Using Enzymatic DNA Molecules

The methods of using enzymatic DNA molecules as disclosed herein are legion. As discussed previously, molecules capable of cleaving the bonds linking neighboring nucleic acids (e.g., phosphoester bonds) have numerous uses encompassing a wide variety of applications. For example, enzymatic DNA molecules having the within-disclosed capabilities, structures, and/or functions are useful in pharmaceutical and medical products (e.g., for wound debridement, clot dissolution, etc.), as well as in household items (e.g., detergents, dental hygiene products, meat tenderizers). Industrial utility of the within-disclosed compounds, compositions and methods is also contemplated and well within the scope of the present invention.

The present invention also describes useful methods for cleaving any single-stranded, looped, partially or fully double-stranded nucleic acid; the majority of these methods employ the novel enzymatically active nucleic acid molecules of the present invention. In various embodiments, the single-stranded nucleic acid segment or portion of the substrate (or the entire substrate itself) comprises DNA, modified DNA, RNA, modified RNA, or composites thereof. Preferably, the nucleic acid substrate need only be single-stranded at or near the substrate cleavage sequence so that an enzymatic nucleic acid molecule of the present invention can hybridize to the substrate cleavage sequence by virtue of the enzyme's recognition sequence.

A nucleic acid substrate that can be cleaved by a method of this invention may be chemically synthesized or enzymatically produced, or it may be isolated from various sources such as phages, viruses, prokaryotic cells, or eukaryotic cells, including animal cells, plant cells, yeast cells and bacterial cells. Chemically synthesized single- and double-stranded nucleic acids are commercially available from many sources including, without limitation, Research Genetics (Huntsville, AL).

RNA substrates may also be synthesized using an Applied Biosystems (Foster City, CA) oligonucleotide synthesizer according to the manufacturer's instructions. Single-stranded phage are also a source of nucleic acid substrates. (See, e.g., Messing et al., PNAS USA 74: 3642-3646 (1977), and Yanisch-Perron et al., Gene 33: 103-119

-26-

(1985).) Bacterial cells containing single-stranded phage would also be a ready source of suitable single-stranded nucleic acid substrates.

Single-stranded RNA cleavable by a method of the present invention could be provided by any of the RNA viruses such as the picornaviruses, togaviruses,
5 orthomyxoviruses, paramyxoviruses, rhabdoviruses, coronaviruses, arenaviruses or retroviruses. As noted previously, a wide variety of prokaryotic and eukaryotic cells may also be excellent sources of suitable nucleic acid substrates.

The methods of this invention may be used on single-stranded nucleic acids or single-stranded portions of looped or double-stranded nucleic acids that are present
10 inside a cell, including eukaryotic, procaryotic, plant, animal, yeast or bacterial cells. Under these conditions an enzymatic nucleic acid molecule (e.g., an enzymatic DNA molecule or deoxyribozyme) of the present invention could act as an anti-viral agent or a regulator of gene expression. Examples of such uses of enzymatic DNA molecules of the present invention are described further hereinbelow.

15 In the majority of methods of the present invention, cleavage of single-stranded nucleic acids occurs at the 3'-terminus of a predetermined base sequence. This predetermined base sequence or substrate cleavage sequence typically contains from 1 to about 10 nucleotides. In other preferred embodiments, an enzymatic DNA molecule of the present invention is able to recognize nucleotides either upstream, or upstream
20 *and* downstream of the cleavage site. In various embodiments, an enzymatic DNA molecule is able to recognize about 2-10 nucleotides upstream of the cleavage site; in other embodiments, an enzymatic DNA molecule is able to recognize about 2-10 nucleotides upstream and about 2-10 nucleotides downstream of the cleavage site. Other preferred embodiments contemplate an enzymatic DNA molecule that is capable
25 of recognizing a nucleotide sequence up to about 30 nucleotides in length, with a length up to about 20 nucleotides being even more preferred.

The within-disclosed methods allow cleavage at any nucleotide sequence by altering the nucleotide sequence of the recognition domains of the enzymatic DNA molecule. This allows cleavage of single-stranded nucleic acid in the absence of a
30 restriction endonuclease site at the selected position.

An enzymatic DNA molecule of the present invention may be separated from any portion of the single-stranded nucleic acid substrate that remains attached to the enzymatic DNA molecule by site-specific hydrolysis at the appropriate cleavage site. Separation of the enzymatic DNA molecule from the substrate (or "cleavage product")
35 allows the enzymatic DNA molecule to carry out another cleavage reaction.

Generally, the nucleic acid substrate is treated under appropriate nucleic acid cleaving conditions -- preferably, physiologic conditions -- with an effective amount of

-27-

an enzymatic DNA molecule of the present invention. If the nucleic acid substrate comprises DNA, cleaving conditions may include the presence of a divalent cation at a concentration of about 2-10mM.

5 An effective amount of an enzymatic DNA molecule is the amount required to cleave a predetermined base sequence present within the single-stranded nucleic acid. Preferably, the enzymatic DNA molecule is present at a molar ratio of DNA molecule to substrate cleavage sites of 1 to 20. This ratio may vary depending on the length of treating and efficiency of the particular enzymatic DNA molecule under the particular nucleic acid cleavage conditions employed.

10 Thus, in one preferred embodiment, treating typically involves admixing, in aqueous solution, the RNA-containing substrate and the enzyme to form a cleavage admixture, and then maintaining the admixture thus formed under RNA cleaving conditions for a time period sufficient for the enzymatic DNA molecule to cleave the RNA substrate at any of the predetermined nucleotide sequences present in the RNA. In
15 various embodiments, a source of ions is also provided -- i.e. monovalent or divalent cations, or both.

In one embodiment of the present invention, the amount of time necessary for the enzymatic DNA molecule to cleave the single-stranded nucleic acid has been predetermined. The amount of time is from about 1 minute to about 24 hours and will
20 vary depending upon the concentration of the reactants and the temperature of the reaction. Usually, this time period is from about 10 minutes to about 2 hours such that the enzymatic DNA molecule cleaves the single-stranded nucleic acid at any of the predetermined nucleotide sequences present.

25 The invention further contemplates that the nucleic acid cleaving conditions include the presence of a source of divalent cations (e.g., PbOAc) at a concentration of about 2-100 mM. Typically, the nucleic acid cleaving conditions include divalent cation at a concentration of about 2 mM to about 10 mM, with a concentration of about 5 mM being particularly preferred.

30 The optimal cationic concentration to include in the nucleic acid cleaving conditions can be easily determined by determining the amount of single-stranded nucleic acid cleaved at a given cation concentration. One skilled in the art will understand that the optimal concentration may vary depending on the particular enzymatic DNA molecule employed.

35 The present invention further contemplates that the nucleic acid cleaving conditions include a pH of about pH 6.0 to about pH 9.0. In one preferred embodiment, the pH ranges from about pH 6.5 to pH 8.0. In another preferred embodiment, the pH emulates physiological conditions, i.e., the pH is about 7.0-7.8, with a pH of about 7.5

-28-

being particularly preferred.

One skilled in the art will appreciate that the methods of the present invention will work over a wide pH range so long as the pH used for nucleic acid cleaving is such that the enzymatic DNA molecule is able to remain in an active conformation. An enzymatic DNA molecule in an active conformation is easily detected by its ability to cleave single-stranded nucleic acid at a predetermined nucleotide sequence.

In various embodiments, the nucleic acid cleaving conditions also include a variety of temperature ranges. As noted previously, temperature ranges consistent with physiological conditions are especially preferred, although temperature ranges consistent with industrial applications are also contemplated herein. In one embodiment, the temperature ranges from about 15°C to about 60°C. In another variation, the nucleic acid cleaving conditions include a temperature ranging from about 30°C to about 56°C. In yet another variation, nucleic acid cleavage conditions include a temperature from about 35°C to about 50°C. In a preferred embodiment, nucleic acid cleavage conditions comprise a temperature range of about 37°C to about 42°C. The temperature ranges consistent with nucleic acid cleaving conditions are constrained only by the desired cleavage rate and the stability of that particular enzymatic DNA molecule at that particular temperature.

In various methods, the present invention contemplates nucleic acid cleaving conditions including the presence of a polyamine. Polyamines useful for practicing the present invention include spermidine, putrescine, spermine and the like. In one variation, the polyamine is present at a concentration of about .01 mM to about 10 mM. In another variation, the polyamine is present at a concentration of about 1 mM to about 10 mM. Nucleic acid cleavage conditions may also include the presence of polyamine at a concentration of about 2 mM to about 5 mM. In various preferred embodiments, the polyamine is spermidine.

G. Vectors

The present invention also features expression vectors including a nucleic acid segment encoding an enzymatic DNA molecule of the present invention situated within the vector, preferably in a manner which allows expression of that enzymatic DNA molecule within a target cell (e.g., a plant or animal cell).

Thus, in general, a vector according to the present invention preferably includes a plasmid, cosmid, phagemid, virus, or phage vector. Preferably, suitable vectors comprise single-stranded DNA (ssDNA) -- e.g., circular phagemid ssDNA. It should also be appreciated that useful vectors according to the present invention need not be circular.

In one variation, nucleotide sequences flanking each of the additional enzymatic

-29-

DNA molecule-encoding sequences are preferably provided, which sequences may be recognized by the first enzymatic DNA molecule. The intervening or flanking sequences preferably comprise at least 1 nucleotide; more preferably, intervening or flanking sequences are about 2-20 nucleotides in length, with sequences of about 5-10 nucleotides in length being particularly preferred.

The addition of polynucleotide tails may also be useful to protect the 3' end of an enzymatic DNA molecule according to the present invention. These may be provided by attaching a polymeric sequence by employing the enzyme terminal transferase.

A vector according to the present invention includes two or more enzymatic DNA molecules. In one embodiment, a first enzymatic DNA molecule has intramolecular cleaving activity and is able to recognize and cleave nucleotide sequences to release other enzymatic DNA sequences; i.e., it is able to function to "release" other enzymatic DNA molecules from the vector. For example, a vector is preferably constructed so that when the first enzymatic DNA molecule is expressed, that first molecule is able to cleave nucleotide sequences flanking additional nucleotide sequences encoding a second enzymatic DNA molecule, a third enzymatic DNA molecule, and so forth. Presuming said first enzymatic DNA molecule (i.e., the "releasing" molecule) is able to cleave oligonucleotide sequences intramolecularly, the additional (e.g. second, third, and so on) enzymatic DNA molecules (i.e., the "released" molecules) need not possess characteristics identical to the "releasing" molecule. For example, in one embodiment, the "released" (i.e., the second, third, etc.) enzymatic DNA molecules are able to cleave specific RNA sequences, while the first ("releasing") enzymatic DNA molecule has nuclease activity allowing it to liberate the "released" molecules. In another embodiment, the "released" enzymatic DNA molecule has amide bond-cleaving activity, while the first ("releasing") enzymatic DNA molecule has nuclease activity.

Alternatively, the first enzymatic DNA molecule may be encoded on a separate vector from the second (and third, fourth, etc.) enzymatic DNA molecule(s) and may have intermolecular cleaving activity. As noted herein, the first enzymatic DNA molecule can be a self-cleaving enzymatic DNA molecule (e.g., a deoxyribozyme), and the second enzymatic DNA molecule may be any desired type of enzymatic DNA molecule. When a vector is caused to express DNA from these nucleic acid sequences, that DNA has the ability under appropriate conditions to cleave each of the flanking regions, thereby releasing one or more copies of the second enzymatic DNA molecule. If desired, several different second enzymatic DNA molecules can be placed in the same cell or carrier to produce different deoxyribozymes. It is also contemplated that any one or more vectors may comprise one or more ribozymes or deoxyribozymes in any combination of "releasing" and "released" enzymatic nucleic acid molecules, as long as

-30-

such a combination achieves the desired result: the release of enzymatic nucleic acid molecules that are capable of cleaving predetermined nucleic acid sequences.

5 Methods of isolating and purifying enzymatic DNA molecules of the present invention are also contemplated. In addition to the methods described herein, various purification methods (e.g. those using HPLC) and chromatographic isolation techniques are available in the art. See, e.g., the methods described in published international application no. WO 93/23569, the disclosures of which are incorporated herein by reference.

10 It should also be understood that various combinations of the embodiments described herein are included within the scope of the present invention. Other features and advantages of the present invention will be apparent from the descriptions hereinabove, from the Examples to follow, and from the claims.

EXAMPLES

15 The following examples illustrate, but do not limit, the present invention.

Example 1

In Vitro Evolution of Enzymatic DNA Molecules:

An Overview

20 *In vitro* selection and *in vitro* evolution techniques allow new catalysts to be isolated without *a priori* knowledge of their composition or structure. Such methods have been used to obtain RNA enzymes with novel catalytic properties. For example, ribozymes that undergo autolytic cleavage with lead cation have been derived from a randomized pool of tRNA^{Phe} molecules (Pan and Uhlenbeck, Biochemistry 31: 3887-3895 (1992)). Group I ribozyme variants have been isolated that can cleave DNA (Beaudry and Joyce, Science 257: 635-641 (1992)) or that have altered metal dependence (Lehman and Joyce, Nature 361: 182-185 (1993)). Starting with a pool of random RNA sequences, molecules have been obtained that catalyze a polymerase-like reaction (Bartel and Szostak, Science 261: 1411-1418 (1993)). In the present example, refinement of specific catalytic properties of an evolved enzyme via alteration of the selection constraints during an *in vitro* evolution procedure is described.

30 Darwinian evolution requires the repeated operation of three processes: (a) introduction of genetic variation; (b) selection of individuals on the basis of some fitness criterion; and (c) amplification of the selected individuals. Each of these processes can be realized *in vitro* (Joyce, Gene 82: 83 (1989)). A gene can be mutagenized by chemical modification, incorporation of randomized mutagenic oligodeoxynucleotides, or
35 inaccurate copying by a polymerase. (See, e.g., Cadwell and Joyce, in PCR Methods and Applications 2: 28-33 (1992); Cadwell and Joyce, PCR Methods and Applications 3

-31-

(Suppl.): S136-S140 (1994); Chu, et al., Virology 98: 168 (1979); Shortle, et al., Meth. Enzymol. 100: 457 (1983); Myers, et al., Science 229: 242 (1985); Matteucci, et al., Nucleic Acids Res. 11: 3113 (1983); Wells, et al., Gene 34: 315 (1985); McNeil, et al., Mol. Cell. Biol. 5: 3545 (1985); Hutchison, et al., PNAS USA 83: 710 (1986);
5 Derbyshire, et al., Gene 46: 145 (1986); Zakour, et al., Nature 295: 708 (1982);
Lehtovaara, et al., Protein Eng. 2: 63 (1988); Leung, et al., Technique 1: 11 (1989);
Zhou, et al., Nucl. Acids Res. 19: 6052 (1991).)

The gene product can be selected, for example, by its ability to bind a ligand or to carry out a chemical reaction. (See, e.g., Joyce, Id. (1989); Robertson and Joyce,
10 Nature 344: 467 (1990); Tuerk, et al., Science 249: 505 (1990).) The gene that corresponds to the selected gene product can be amplified by a reciprocal primer method, such as the polymerase chain reaction (PCR). (See, e.g., Saiki, et al., Science 230: 1350-54 (1985); Saiki, et al., Science 239: 487-491 (1988).)

Alternatively, nucleic acid amplification may be carried out using self-sustained
15 sequence replication (3SR). (See, e.g., Guatelli, et al., PNAS USA 87: 1874 (1990), the disclosures of which are incorporated by reference herein.) According to the 3SR method, target nucleic acid sequences may be amplified (replicated) exponentially *in vitro* under isothermal conditions by using three enzymatic activities essential to retroviral replication: (1) reverse transcriptase, (2) RNase H, and (3) a DNA-dependent
20 RNA polymerase. By mimicking the retroviral strategy of RNA replication by means of cDNA intermediates, this reaction accumulates cDNA and RNA copies of the original target.

In summary, if one is contemplating the evolution of a population of enzymatic DNA molecules, a continuous series of reverse transcription and transcription reactions
25 replicates an RNA target sequence by means of cDNA intermediates. The crucial elements of this design are (a) the oligonucleotide primers both specify the target and contain 5' extensions encoding the T7 RNA polymerase binding site, so that the resultant cDNAs are competent transcription templates; (b) cDNA synthesis can proceed to completion of both strands due to the degradation of template RNA in the
30 intermediate RNA-DNA hybrid by RNase H; and (c) the reaction products (cDNA and RNA) can function as templates for subsequent steps, enabling exponential replication.

If one is evolving enzymatic DNA molecules, various critical elements of this design are somewhat different, as disclosed in these Examples. For instance, (1) the oligonucleotide primers specify the target and are preferably "marked" or labeled in
35 some fashion -- e.g., via biotinylation -- so the resultant competent template strands are easily identified; and (2) the *in vitro* selection procedure used preferably depends upon the identification of the most favorable release mechanism.

-32-

A major obstacle to realizing Darwinian evolution *in vitro* is the need to integrate mutation and amplification, both of which are genotype-related, with selection, which is phenotype-related. In the case of nucleic acid enzymes, for which genotype and phenotype are embodied in the same molecule, the task is simplified.

5 A. Design of Enzymatic DNA Molecules

It is well known that single-stranded DNA can assume interesting tertiary structures. The structure of a "tDNA", for example, closely resembles that of the corresponding tRNA. (See Paquette, et al., Eur. J. Biochem. 189: 259-265 (1990).) Furthermore, it has been possible to replace as many as 31 of 35 ribonucleotides within a hammerhead ribozyme, while retaining at least some catalytic activity. (See Perreault, 10 et al., Nature 344: 565-567 (1990); Williams, et al., Proc. Natl. Acad. Sci. USA 89: 918-921 (1992); Yang, et al., Biochemistry 31: 5005-5009 (1992).)

In vitro selection techniques have been applied to large populations of random-sequence DNAs, leading to the recovery of specific DNA "aptamers" that bind a target ligand with high affinity (Bock, et al., Nature 355: 564-566 (1992); Ellington & Szostak, Nature 355: 850-852 (1992); Wyatt & Ecker, PNAS USA 91: 1356-1360 (1994)). Recently, two groups carried out the first NMR structural determination of an aptamer, a 15mer DNA that forms a G-quartet structure and binds the protein thrombin with high affinity (Wang, et al., Biochemistry 32: 1899-1904 (1993); Macaya, et al., 20 PNAS USA 90: 3745-3749 (1993)). These findings were corroborated by an X-ray crystallographic analysis (Padmanabhan, et al., J. Biol. Chem. 268: 17651-17654 (1993)).

The ability to bind a substrate molecule with high affinity and specificity is a prerequisite of a good enzyme. In addition, an enzyme must make use of 25 well-positioned functional groups, either within itself or a cofactor, to promote a particular chemical transformation. Furthermore, the enzyme must remain unchanged over the course of the reaction and be capable of operating with catalytic turnover. Some would add the requirement that it be an informational macromolecule, comprised of subunits whose specific ordering is responsible for catalytic activity. While these 30 criteria are open to debate on both semantic and chemical grounds, they serve to distinguish phenomena of chemical rate enhancement that range from simple solvent effects to biological enzymes operating at the limit of substrate diffusion (Albery & Knowles, Biochemistry 15: 5631-5640 (1976)).

As described in greater detail hereinbelow, we sought to develop a general 35 method for rapidly obtaining DNA catalysts and DNA enzymes, starting from random sequences. As an initial target, we chose a reaction that we felt was well within the capability of DNA: the hydrolytic cleavage of an RNA phosphodiester, assisted by a

-33-

divalent metal cofactor. This is the same reaction that is carried out by a variety of naturally-occurring RNA enzymes, including the hammerhead and hairpin motifs. (See, e.g., Forster A.C. & Symons R.H., Cell 49: 211-220 (1987); Uhlenbeck, Nature 328: 596-600 (1987); Hampel & Tritz, Biochemistry 28: 4929-4933 (1989)).

5 It has recently been shown that, beginning with a randomized library of tRNA molecules, one can obtain ribozymes that have Pb^{2+} -dependent, site-specific RNA phosphoesterase activity at neutral pH (Pan & Uhlenbeck, Biochemistry 31: 3887-3895 (1992); Pan & Uhlenbeck, Nature 358: 560-563 (1992)). This is analogous to the fortuitous self-cleavage reaction of yeast tRNA^{Phe} (Dirheimer & Werner, Biochimie 54: 10 127-144 (1972)), which depends on specific coordination of a Pb^{2+} ion at a defined site within the tRNA. (See Rubin & Sundaralingam, J. Biomol. Struct. Dyn. 1: 639-646 (1983); Brown, et al., Biochemistry 24: 4785-4801 (1985).)

As disclosed herein, our goals included the development of DNAs that could carry out Pb^{2+} -dependent cleavage of a particular RNA phosphoester, initially presented 15 within a short leader sequence attached to the 5' end of the DNA, and ultimately located within a separate molecule that could be cleaved in an intermolecular fashion with rapid catalytic turnover. These goals were successfully achieved, as described further below.

No assumptions were made as to how the DNA would interact with the target 20 phosphoester and surrounding nucleotides. Beginning with a pool of approximately 10^{14} random 50mer sequences, *in vitro* selection was allowed to run its course. After five rounds of selection carried out over four days, the population as a whole had attained the ability to cleave the target phosphoester in the presence of 1 mM Pb^{2+} at a rate of about 0.2 min^{-1} . This is an approximately 10^5 -fold increase compared to the 25 spontaneous rate of cleavage under the same reaction conditions.

Individuals were isolated from the population, sequenced, and assayed for catalytic activity. Based on this information, the reaction was converted to an intermolecular format and then simplified to allow site-specific cleavage of a 19mer substrate by a 38mer DNA enzyme, in a reaction that proceeds with a turnover rate of 1 30 min^{-1} at 23°C and pH 7.0 in the presence of 1 mM PbOAc.

B. In Vitro Selection Scheme

A starting pool of approximately 10^{14} single-stranded DNA molecules was generated, all of which contain a 5' biotin moiety, followed successively by a fixed domain that includes a single ribonucleotide, a potential catalytic domain comprised of 35 50 random deoxyribonucleotides, and a second fixed domain that lay at the 3' terminus (Fig. 1).

The pool was constructed by a nested PCR (polymerase chain reaction)

-34-

technique, beginning with synthetic DNA that contained 50 random nucleotides flanked by primer binding sites. The nested PCR primer was a 5'-biotinylated synthetic oligodeoxynucleotide with a 3'-terminal adenosine ribonucleotide.

5 Ribonucleotide-terminated oligonucleotides efficiently prime template-directed elongation in the context of the PCR (L.E. Orgel, personal communication), in this case giving rise to an extension product that contains a single embedded ribonucleotide.

Figure 1 illustrates a selective amplification scheme for isolation of DNAs that cleave a target RNA phosphoester. Double-stranded DNA containing a stretch of 50 random nucleotides is amplified via PCR, employing a 5'-biotinylated DNA primer (e.g.,
10 primer 3 -- 3a or 3b) terminated at the 3' end by an adenosine ribonucleotide (represented by the symbol "N" or "rA", wherein both N and rA represent an adenosine ribonucleotide). This primer is extended by *Taq* polymerase to yield a DNA product that contains a single embedded ribonucleotide. The resulting double-stranded DNA is immobilized on a streptavidin matrix and the unbiotinylated DNA strand is removed by
15 washing with 0.2 N NaOH. After re-equilibrating the column with a buffered solution, the column is washed with the same solution with added 1 mM PbOAc. DNAs that undergo Pb²⁺-dependent self-cleavage are released from the column, collected in the eluant, and amplified by PCR. The PCR products are then used to initiate the next round of selective amplification.

20 The PCR products were passed over a streptavidin affinity matrix, resulting in noncovalent attachment of the 5'-biotinylated strand of the duplex DNA. The nonbiotinylated strand was removed by brief washing with 0.2 N NaOH, and the bound strand was equilibrated in a buffer containing 0.5 M NaCl, 0.5 M KCl, 50 mM MgCl₂, and 50 mM HEPES (pH 7.0) at 23°C. Next, 1 mM PbOAc was provided in the same
25 buffer, allowing Pb²⁺-dependent cleavage to occur at the target phosphoester, thereby releasing a subset of the DNAs from the streptavidin matrix. In principle, an individual DNA might facilitate its own release by various means, such as disruption of the interaction between biotin and streptavidin or cleavage of one of the deoxyribonucleotide linkages. It was felt that cleavage of the ribonucleoside 3'-O-P
30 bond would be the most likely mechanism for release, based on the relative lability of this linkage, and that Pb²⁺-dependent hydrolytic cleavage would allow release to occur most rapidly. In principle, however, the *in vitro* selection procedure should identify the most favorable release mechanism as well as those individuals best able to carry out that mechanism.

35 DNA molecules released from the matrix upon addition of Pb²⁺ were collected in the eluant, concentrated by precipitation with ethanol, and subjected to nested PCR amplification. As in the construction of the starting pool of molecules, the first PCR

-35-

amplification utilized primers that flank the random region (primers 1 and 2) and the second utilized a 5'-biotinylated primer (primer 3b) that has a 3'-terminal riboadenylate, thereby reintroducing the target RNA phosphoester. The entire selective amplification procedure requires 3-4 hours to perform.

5 The molecules are purified in three ways during each round of this procedure: first, following PCR amplification, by extracting twice with phenol and once with chloroform / isoamyl alcohol, then precipitating with ethanol; second, following attachment of the DNA to streptavidin, by washing away all the nonbiotinylated molecules under strongly denaturing conditions; and third, following elution with Pb^{2+} ,
10 by precipitating with ethanol. There is no gel electrophoresis purification step, and thus no selection pressure constraining the molecules to a particular length.

C. Selection of Catalytic DNA

 We carried out five successive rounds of *in vitro* selection, progressively decreasing the reaction time following addition of Pb^{2+} in order to progressively increase
15 the stringency of selection. During rounds 1 through 3, the reaction time was 1 hour; during round 4, the reaction time was 20 minutes; and during round 5, it was 1 minute. The starting pool of single-stranded DNAs, together with the population of molecules obtained after each round of selection, was assayed for self-cleavage activity under conditions identical to those employed during *in vitro* selection (see Fig. 2).

20 For this assay, the molecules were prepared with a 5'- ^{32}P rather than a 5'-biotin moiety, allowing detection of both the starting material and the 5' cleavage product. Following a 5-minute incubation, there was no detectable activity in the initial pool (G0) or in the population obtained after the first and second rounds of selection. DNAs
25 obtained after the third round (G3) exhibited a modest level of activity; this activity increased steadily, reaching approximately 50% self-cleavage for the DNAs obtained after the fifth round of selection (G5). Cleavage was detected only at the target phosphoester, even after long incubation times. This activity was lost if Pb^{2+} was omitted from the reaction mixture.

 Figure 2 illustrates the self-cleavage activity of the starting pool of DNA (G0) and populations obtained after the first through fifth rounds of selection (G1 - G5).
30 Reaction mixtures contained 50 mM $MgCl_2$, 0.5 M NaCl, 0.5 M KCl, 50 mM HEPES (pH 7.0 at 23°C), and 3 nM [5'- ^{32}P]-labeled DNA, incubated at 23°C for 5 min either in the presence or in the absence of 1 mM $PbOAc$. The symbol **Pre** represents 108-nucleotide precursor DNA (SEQ ID NO 4); **Clv**, 28-nucleotide 5'-cleavage product (SEQ ID NO 5);
35 and **M**, primer 3a (SEQ ID NO 6), corresponding in length to the 5'-cleavage product.

 The 28-nucleotide 5' cleavage product (Clv) illustrated preferably has the sequence 5'-GGGACGAATTCTAATACGACTCACTATN-3', wherein "N" represents

-36-

adenosine ribonucleotide with an additional 2', 3'-cyclic phosphate on the 3' end (SEQ ID NO 5). In alternative embodiments, "N" represents adenosine ribonucleotide with an additional 2' or 3' phosphate on the 3' end of the molecule.

5 In Figure 2, the "G0" lane "Pre" band comprises a sampling of 108-nucleotide precursor DNAs that each include 50 random nucleotides. Therefore, any given "Pre" sampling will contain a wide variety of precursor DNAs, and each sampling will likely differ from previous and subsequent samplings. The "G1" through "G5" lanes contain "Pre" bands that are increasingly enriched for catalytic DNA molecules, but still contain a large number of different DNA sequences (i.e., differing in the 50 nucleotide
10 randomized domain). A sample of these different sequences from "G5 Pre" DNA is provided in Figure 3.

Shotgun cloning techniques were employed to isolate individuals from the G5 population; the complete nucleotide sequences of 20 of these subclones were then determined (see Fig. 3). (Also see, e.g., Cadwell and Joyce, in PCR Methods and Applications 2: 28-33 (1992); Cadwell and Joyce, PCR Methods and Applications 3 (Suppl.): S136-S140 (1994).) Of the 20 sequences, five were unique, two occurred twice, one occurred three times, and one occurred eight times. All of the individual variants share common sequence elements within the 50-nucleotide region that had
15 been randomized in the starting pool of DNA. They all contain two presumed template regions, one with complementarity to a stretch of nucleotides that lies just upstream from the cleavage site and the other with complementarity to nucleotides that lie at
20 least four nucleotides downstream. Between these two presumed template regions lies a variable domain of 1-11 nucleotides, followed by the fixed sequence 5'-AGCG-3', then a second variable domain of 3-8 nucleotides, and finally the fixed sequence 5'-CG-3' or
25 5'-CGA-3'. Nucleotides that lie outside of the two presumed template regions are highly variable in both sequence and length. In all of the sequenced subclones, the region corresponding to the 50 initially-randomized nucleotides remains a total of 50 nucleotides in length.

Figure 3 illustrates the sequence alignment of individual variants isolated from
30 the population after five rounds of selection. The fixed substrate domain (5'-GGGACGAATTCTAATACGACTCACTATrAGGAAGAGATGGCGAC-3', or 5'-GGGACGAATTCTAATACGACTCACTATNGGAAGAGATGGCGAC-3', where N represents adenosine ribonucleotide) (SEQ ID NO 13) is shown at the top, with the target riboadenylate identified with an inverted triangle. Substrate nucleotides that are
35 commonly involved in presumed base-pairing interactions are indicated by a vertical bar. Sequences corresponding to the 50 initially-randomized nucleotides are aligned antiparallel to the substrate domain. All of the variants are 3'-terminated by the fixed

-37-

sequence 5'-CGGTAAGCTTGGCAC-3' (SEQ ID NO 1) ("primer site"; not shown). Nucleotides within the initially-randomized region that are presumed to form base pairs with the substrate domain are indicated on the right and left sides of the Figure; the putative base-pair-forming (or substrate binding) regions of the enzymatic DNA molecules are individually boxed in each sequence shown. The highly-conserved nucleotides within the putative catalytic domain are illustrated in the two boxed columns.

While it is anticipated that additional data will be helpful in constructing a meaningful secondary structural model of the catalytic domain, we note that, like the hammerhead and hairpin ribozymes, the catalytic domain of our enzymatic DNA molecules appears to contain a conserved core flanked by two substrate binding regions (or recognition domains) that interact with the substrate through base-pairing interactions. Similar to the hammerhead and hairpin ribozymes, the catalytic DNAs also appear to require a short stretch of unpaired substrate nucleotides -- in this case 5'-GGA-3' -- between the two regions that are involved in base pairing.

It was also interesting to note that each of the nine distinct variants exhibited a different pattern of presumed complementarity with the substrate domain. In some cases, base pairing was contiguous, while in others it was interrupted by one or more noncomplementary pairs. The general tendency seems to be to form tighter interaction with the nucleotides that lie upstream from the cleavage site compared to those that lie downstream. Binding studies and site-directed mutagenesis analysis should enable us to gain further insights and to further substantiate this conjecture.

In order to gain further insight into the sequence requirements for catalytic function, the self-cleavage activity of six of the nine variants was tested and evaluated under the within-described selection conditions (see Fig. 3). Not surprisingly, the sequence that occurred in eight of the 20 subclones proved to be the most reactive, with a first-order rate constant of 1.4 min^{-1} . All of the studied variants were active in the self-cleavage assay and all gave rise to a single 5'-labeled product corresponding to cleavage at the target RNA phosphoester.

The dominant subclone was further analyzed under a variety of reaction conditions. Its self-cleavage activity was dependent on Pb^{2+} but was unaffected if Mg^{2+} was omitted from the reaction mixture. There was a requirement for a monovalent cation as well, which can be met by either Na^+ or K^+ . The reaction rate increased linearly with increasing concentration of monovalent cation over the range of 0 - 1.0 M ($r = 0.998$). Other variables that may affect the reaction, such as pH, temperature, and the presence of other divalent metals, are in the process of being evaluated further.

-38-

Example 2**Materials and Methods****A. Oligonucleotides and Oligonucleotide Analogs**

Synthetic DNAs and DNA analogs were purchased from Operon Technologies.

5 The 19-nucleotide substrate, 5'-pTCACTATrAGGAAGAGATGG-3' (or 5'-
pTCACTATNGGAAGAGATGG-3', wherein "N" represents adenosine ribonucleotide)
(SEQ ID NO 7), was prepared by reverse-transcriptase catalyzed extension of
5'-pTCACTATrA-3' (or 5'-pTCACTATN-3', wherein "N" represents adenosine
ribonucleotide) (SEQ ID NO 8), as previously described (Breaker, Banerji, & Joyce,
10 Biochemistry 33: 11980-11986 (1994)), using the template
5'-CCATCTTCTCCTATAGTGAGTCCGGCTGCA-3' (SEQ ID NO 9). Primer 3, 5'-
GGGACGAATTCTAATACGACTCACTATrA-3' (or 5'-
GGGACGAATTCTAATACGACTCACTATN-3', wherein "N" represents adenosine
ribonucleotide) (SEQ ID NO 6), was either 5'-labeled with [γ -³²P]ATP and T4
15 polynucleotide kinase (primer 3a) or 5'-thiophosphorylated with [γ -S]ATP and T4
polynucleotide kinase and subsequently biotinylated with *N*-iodoacetyl-*N'*-
biotinylhexylenediamine (primer 3b).

B. DNA Pool Preparation

The starting pool of DNA was prepared by PCR using the synthetic oligomer
20 5'-GTGCCAAGCTTACCG-N₅₀-GTCGCCATCTCTTCC-3' (SEQ ID NO 4), where N is an
equimolar mixture of G, A, T and C. A 2-ml PCR, containing 500 pmoles of the
randomized oligomer, 1,000 pmoles primer 1 (5'-GTGCCAAGCTTACCG-3', SEQ ID NO
10), 500 pmoles primer 2
(5'-CTGCAGAATTCTAATACGACTCACTATAGGAAGAGATGGCGAC-3', SEQ ID NO 11),
25 500 pmoles primer 3b, 10 μ Ci [α -³²P]dATP, and 0.2 U μ l⁻¹ *Taq* DNA polymerase, was
incubated in the presence of 50 mM KCl, 1.5 mM MgCl₂, 10 mM Tris-HCl (pH 8.3 at
23°C), 0.01% gelatin, and 0.2 mM of each dNTP for 1 min at 92°C, 1 min at 50°C, and
2 min at 72°C, then 5 cycles of 1 min at 92°C, 1 min at 50°C, and 1 min at 72°C. The
resulting mixture was extracted twice with phenol and once with chloroform / isoamyl
30 alcohol, and the DNA was isolated by precipitation with ethanol.

C. In Vitro Selection

The starting pool of DNA was resuspended in 500 μ L of buffer A (1 M NaCl and
50 mM HEPES (pH 7.0 at 23°C)) and was passed repeatedly over a streptavidin column
(AffiniTip Strep 20, Genosys, The Woodlands, TX). The column was washed with five
35 100- μ l volumes of buffer A, followed by five 100- μ l volumes of 0.2 N NaOH, then
equilibrated with five 100- μ l volumes of buffer B (0.5 M NaCl, 0.5 M KCl, 50 mM
MgCl₂, and 50 mM HEPES (pH 7.0 at 23°C)). The immobilized single-stranded DNA was

-39-

eluted over the course of 1 hr with three 20- μ l volumes of buffer B with added 1 mM PbOAc. The entire immobilization and elution process was conducted at 23°C. The eluant was collected in an equal volume of buffer C (50 mM HEPES (pH 7.0 at 23°C) and 80 mM EDTA) and the DNA was precipitated with ethanol.

5 The resulting DNA was amplified in a 100- μ L PCR containing 20 pmoles primer 1, 20 pmoles primer 2, 0.05 U μ l⁻¹ *Taq* polymerase, 50 mM KCl, 1.5 mM MgCl₂, 10 mM Tris-HCl (pH 8.3 at 23°C), 0.01% gelatin, and 0.2 mM of each dNTP for 30 cycles of 10 sec at 92°C, 30 sec at 50°C, and 30 sec at 72°C. The reaction products were extracted twice with phenol and once with chloroform / isoamyl alcohol, and the DNA
10 was recovered by precipitation with ethanol. Approximately 4 pmoles of the amplified DNA was added to a second, nested PCR containing 100 pmoles primer 1, 100 pmoles primer 3b, 20 μ Ci [α -³²P]dATP, and 0.1 U μ l⁻¹ *Taq* polymerase, in a total volume of 200 μ L that was amplified for 10 cycles of 1 min at 92°C, 1 min at 50°C, and 1 min at 72°C. The PCR products were once more extracted and precipitated, and the resulting
15 DNA was resuspended in 50 μ L buffer A, then used to begin the next round of selection.

The second and third rounds were carried out as above, except that the nested PCR at the end of the third round was performed in a 100- μ l volume. During the fourth round, the elution time following addition of Pb²⁺ was reduced to 20 min (two 20- μ L
20 elution volumes) and only half of the recovered DNA was used in the first PCR, which involved only 15 temperature cycles. During the fifth round, the elution time was reduced to 1 min (two 20- μ L elution volumes) and only one-fourth of the recovered DNA was used in the first PCR, which involved 15 temperature cycles. DNA obtained after the fifth round of selection was subcloned and sequenced, as described previously
25 (Tsang & Joyce, Biochemistry 33: 5966-5973 (1994)).

D. Kinetic Analysis of Catalytic DNAs

Populations of DNA and various subcloned individuals were prepared with a 5'-³²P label by asymmetric PCR in a 25- μ l reaction mixture containing 10 pmoles primer 3a, 0.5 pmoles input DNA, and 0.1 U μ l⁻¹ *Taq* polymerase, under conditions as described
30 above, for 10 cycles of 1 min at 92°C, 1 min at 50°C, and 1 min at 72°C. The resulting [5'-³²P]-labeled amplification products were purified by electrophoresis in a 10% polyacrylamide / 8 M gel.

Self-cleavage assays were carried out following preincubation of the DNA in buffer B for 10 min. Reactions were initiated by addition of PbOAc to 1 mM final
35 concentration and were terminated by addition of an equal volume of buffer C. Reaction products were separated by electrophoresis in a 10% polyacrylamide / 8 M gel. Kinetic assays under multiple-turnover conditions were carried out in buffer B that included 50

-40-

$\mu\text{g ml}^{-1}$ BSA to prevent adherence of material to the vessel walls. Substrate and enzyme molecules were preincubated separately for 5 min in reaction buffer that lacked Pb^{2+} , then combined, and the reaction was initiated by addition of PbOAc to a final concentration of 1 mM.

5

Example 3

Evolution of Deoxyribozymes That Cleave Intermolecularly

A. Conversion to an Intermolecular Format

10

Based on the variable pattern of presumed base-pairing interactions between the catalytic and substrate domains of the studied variants, it was felt that it would be reasonably straightforward to convert the DNA-catalyzed reaction to an intermolecular format. In doing so, we wished to simplify the two substrate-binding regions of the catalyst so that each would form an uninterrupted stretch of 7-8 base pairs with the substrate. In addition, we wished to provide a minimal substrate, limited to the two

15 base-pairing regions and the intervening sequence 5'-GGA-3' (Fig. 4A).

15

Figures 4A and 4B illustrate DNA-catalyzed cleavage of an RNA phosphoester in an intermolecular reaction that proceeds with catalytic turnover. Figure 4A is a diagrammatic representation of the complex formed between the 19mer substrate and 38mer DNA enzyme. The substrate contains a single adenosine ribonucleotide ("rA" or "N", adjacent to the arrow), flanked by deoxyribonucleotides. The synthetic DNA

20 enzyme is a 38-nucleotide portion of the most frequently occurring variant shown in Fig. 3. Highly-conserved nucleotides located within the putative catalytic domain are "boxed". As illustrated, one conserved sequence is "AGCG", while another is "CG" (reading in the 5'→3' direction).

20

25

Figure 4B shows an Eadie-Hofstee plot used to determine K_m (negative slope) and V_{max} (y-intercept) for DNA-catalyzed cleavage of [5'- ^{32}P]-labeled substrate under conditions identical to those employed during *in vitro* selection. Initial rates of cleavage were determined for reactions involving 5 nM DNA enzyme and either 0.125, 0.5, 1, 2, or 4 μM substrate.

30

35

In designing the catalytic domain, we relied heavily on the composition of the most reactive variant, truncating by two nucleotides at the 5' end and 11 nucleotides at the 3' end. The 15 nucleotides that lay between the two template regions were left unchanged and a single nucleotide was inserted into the 3' template region to form a continuous stretch of nucleotides capable of forming base pairs with the substrate. The substrate was simplified to the sequence 5'-ICACTATrA • GGAAGAGATGG-3' (or 5'-ICACTATN • GGAAGAGATGG-3', wherein "N" represents adenosine ribonucleotide) (SEQ ID NO 12), where the underlined nucleotides correspond to the two regions

-41-

involved in base pairing with the catalytic DNA molecule.

The simplified reaction system, employing a 38mer catalytic DNA molecule (catalyst) comprised entirely of deoxyribonucleotides and a 19mer substrate containing a single ribonucleotide embedded within an otherwise all-DNA sequence, allows efficient DNA-catalyzed phosphoester cleavage with rapid turnover. Over a 90-minute incubation in the presence of 0.01 μM catalyst and 1 μM substrate, 46% of the substrate is cleaved, corresponding to 46 turnovers of the catalyst. A preliminary kinetic analysis of this reaction was carried out, evaluated under multiple-turnover conditions. The DNA catalyst exhibits Michaelis-Menten kinetics, with values for k_{cat} and K_m of 1 min^{-1} and 2 μM , respectively (see Fig. 4B). The value for K_m is considerably greater than the expected dissociation constant between catalyst and substrate based on Watson-Crick interactions. The substrate was incubated under identical reaction conditions (but in the absence of the catalyst); a value for k_{uncat} of $4 \times 10^{-6} \text{ min}^{-1}$ was obtained. This is consistent with the reported value of $5 \times 10^{-3} \text{ min}^{-1}$ for hydrolysis of the more labile 1-nitrophenyl-1,2-propanediol in the presence of 0.5 mM Pb^{2+} at pH 7.0 and 37°C (Breslow & Huang, PNAS USA 88: 4080-4083 (1991)).

It is now presumed that the phosphoester cleavage reaction proceeds via a hydrolytic mechanism involving attack by the ribonucleoside 2'-hydroxyl on the vicinal phosphate, generating a 5' product with a terminal 2'(3')-cyclic phosphate and 3' product with a terminal 5'-hydroxyl. In support of this mechanism, the 3'-cleavage product is efficiently phosphorylated with T4 polynucleotide kinase and [γ - ^{32}P]ATP, consistent with the availability of a free 5'-hydroxyl (data not shown).

B. Discussion

After five rounds of *in vitro* selection, a population of single-stranded DNA molecules that catalyze efficient Pb^{2+} -dependent cleavage of a target RNA phosphoester was obtained. Based on the common features of representative individuals isolated from this population, a simplified version of both the catalytic and substrate domains was constructed, leading to a demonstration of rapid catalytic turnover in an intermolecular context. Thus the 38mer catalytic domain provides an example of a DNA enzyme, or what might be termed a "deoxyribozyme".

Referring to this molecule as an enzyme, based on the fact that it is an informational macromolecule capable of accelerating a chemical transformation in a reaction that proceeds with rapid turnover and obeys Michaelis-Menten kinetics, may not satisfy everyone's notion of what constitutes an enzyme. Some might insist that an enzyme, by definition, must be a polypeptide. If, however, one accepts the notion of an RNA enzyme, then it seems reasonable to adopt a similar view concerning DNA enzymes. Considering how quickly we were able to generate this molecule from a pool

-42-

of random-sequence DNAs, we expect that many other examples of synthetic DNA enzymes will appear in the near future.

The Pb^{2+} -dependent cleavage of an RNA phosphoester was chosen as an initial target for DNA catalysis because it is a straightforward reaction that simply requires the proper positioning of a coordinated Pb^{2+} -hydroxyl to facilitate deprotonation of the 2' hydroxyl that lies adjacent to the cleavage site. (See, e.g., Pan, et al., in The RNA World, Gesteland & Atkins (eds.), pp. 271-302, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1993).) Pb^{2+} is known to coordinate to the N7 position of purines, the O6 position of guanine, the O4 position of uracil, and the N3 position of cytosine (Brown, et al., Nature **303**: 543-546 (1993)). Thus, the differences in sugar composition and conformation of DNA compared to RNA seemed unlikely to prevent DNA from forming a well-defined Pb^{2+} -binding pocket.

A substrate that contains a single ribonucleotide within an otherwise all-DNA sequence was chosen because it provided a uniquely favored site for cleavage and insured that any resulting catalytic activity would be attributable solely to DNA. Substrate recognition appears to depend on two regions of base-pairing interactions between the catalyst and substrate. However, the unpaired substrate nucleotides, 5'-GGA-3', that lie between these two regions may play an important role in substrate recognition, metal coordination, or other aspects of catalytic function.

It is further anticipated that an all-RNA molecule, other RNA-DNA composites, and molecules containing one or more nucleotide analogs may be acceptable substrates. As disclosed herein, the within-described *in vitro* evolution procedures may successfully be used to generate enzymatic DNA molecules having the desired specificities; further analyses along these lines are presently underway.

In addition, studies to determine whether the presumed base-pairing interactions between enzyme and substrate are generalizable with respect to sequence are in progress, using the presently-described methods. The within-disclosed Pb^{2+} -dependent deoxyribozymes may also be considered model compounds for exploring the structural and enzymatic properties of DNA.

The methods employed in the present disclosure for the rapid development of DNA catalysts will have considerable generality, allowing us to utilize other cofactors to trigger the cleavage of a target linkage attached to a potential catalytic domain. In this regard, the development of Mg^{2+} -dependent DNA enzymes that specifically cleave target RNAs under physiological conditions is of interest, as is the development of DNA enzymes that function in the presence of other cations (see Example 4). Such molecules will provide an alternative to traditional antisense and ribozyme approaches for the specific inactivation of target mRNAs.

-43-

DNA thus joins RNA and protein on the list of biological macromolecules that are capable of exhibiting enzymatic activity. The full extent of DNA's catalytic abilities remains to be explored, but these explorations should proceed rapidly based on *in vitro* selection methods such as those employed in this study.

5 DNA enzymes offer several important advantages compared to other macromolecular catalysts. First, they are easy to prepare, in an era when most laboratories have access to an automated DNA synthesizer and the cost of DNA phosphoramidites has become quite modest. Second, they are very stable compounds, especially compared to RNA, thus facilitating their use in biophysical studies. Third, we expect that they can be adapted to therapeutic applications that at present make use of antisense DNAs that lack RNA-cleavage activity. *In vitro* selection could be carried out with DNA analogs, including compounds that are nuclease resistant such as phosphorothioate-containing DNA, so long as these analogs can be prepared in the form of a deoxynucleoside 5'-triphosphate and are accepted as a substrate by a DNA-dependent DNA polymerase. Finally, DNA enzymes offer a new window on our understanding of the macromolecular basis of catalytic function. It will be interesting, for example, to carry out comparative analyses of protein-, RNA-, and DNA-based enzymes that catalyze the same chemical transformation.

Example 4

20 Other Families of Catalytic DNAs

A starting pool of DNA was prepared by PCR essentially as described in Example 2.B. above, except that the starting pool of DNA comprised molecules containing 40 random nucleotides. Thus, the starting pool of DNA described herein was prepared by PCR using the synthetic oligomer 5' GGG ACG AAT TCT AAT ACG ACT CAC TAT rA GG AAG AGA TGG CGA CAT CTC N₄₀GT GAC GGT AAG CTT GGC AC 3' (SEQ ID NO 23), where N is an equimolar mixture of G, A, T and C, and where the DNA molecules were selected for the ability to cleave the phosphoester following the target rA. (See Figure 6A, also.)

30 Selective amplification was carried out in the presence of either Pb²⁺, Zn²⁺, Mn²⁺, or Mg²⁺, thereby generating at least four "families" of catalytic DNA molecules. As illustrated in Figure 5, catalytic DNA molecules demonstrating specific activity were generated in the presence of a variety of cations.

35 Figure 5 is a photographic representation showing a polyacrylamide gel demonstrating specific endoribonuclease activity of four families of selected catalytic DNAs. Selection of a Pb²⁺-dependent family of molecules was repeated in a side-by-side fashion as a control. In each group of three lanes, the first lane shows the lack of activity of the selected population in the absence of the metal cation, the second lane

-44-

shows the observed activity in the presence of the metal cation, and the third lane shows the lack of activity of the starting pool (G0). At present, the order of reactivity is observed to be $Pb^{2+} > Zn^{2+} > Mn^{2+} > Mg^{2+}$, mirroring the pK_a of the corresponding metal-hydroxide.

5 After either five (G5) or six (G6) rounds of selective amplification in the presence of the preselected divalent cation, the desired endonuclease activity was obtained. The following description of selective amplification in the presence of Mg^{2+} is intended to be exemplary.

10 Six rounds of *in vitro* selective amplification were carried out, following the method described in Example 2 hereinabove, except that the divalent metal used was 1 mM Mg^{2+} rather than 1 mM Pb^{2+} . (See also Breaker and Joyce, Chem. & Biol. 1: 223-229 (1994), incorporated by reference herein, which describes essentially the same procedure.)

15 Individual clones were isolated following the sixth round, and the nucleotide sequence of 24 of these clones was determined. All of the sequences began with: 5' GGG ACG AAT TCT AAT ACG ACT CAC TAT rA GG AAG AGA TGG CGA CA (SEQ ID NO 23 from position 1 to 44) and ended with: CGG TAA GCT TGG CAC 3' (SEQ ID NO 23 from position 93 to 107).

20 The segment in the middle, corresponding to TCTC N₄₀ GTGA (SEQ ID NO 23 from position 45 to 92) in the starting pool, varied as follows:

- (13) CCG CCC ACC TCT TTT ACG AGC CTG TAC GAA ATA GTG CTC TTG
TTA GTA T (SEQ ID NO 24)
- 25 (5) TCT CTT CAG CGA TGC ACG CTT GTT TTA ATG TTG CAC CCA TGT
IAG TGA (SEQ ID NO 25)
- (2) TCT CAT CAG CGA TTG AAC CAC TTG GTG GAC AGA CCC ATG TTA
GTG A (SEQ ID NO 26)
- (1) CCG CCC ACC TCT TTT ACG AGC CTG TAC GAA ATA GTG TTC TTG
TTA GTA T (SEQ ID NO 27)
- 30 (1) CCG CCC ACC TCT TTT ACG AGC CTG TAC GAA ATA GTG CTC TCG
TTA GTA T (SEQ ID NO 28)
- (1) TCT CAG ACT TAG TCC ATC ACA CTC TGT GCA TAT GCC TGC TTG
ATG TGA (SEQ ID NO 29)
- (1) -CT CTC ATC TGC TAG CAC GCT CGA ATA GTG TCA GTC GAT GTG A
35 (SEQ ID NO 30).

-45-

The initial number in parentheses indicates the number of clones having that particular sequence. Note that some mutations (highlighted in bold type) occurred at nucleotide positions other than those that were randomized initially.

5 The second sequence listed above (i.e., SEQ ID NO 25), which occurred in 5 of 24 clones, was chosen as a lead (i.e. principal) compound for further study. Its cleavage activity was measured in the presence of a 1 mM concentration of various divalent metals and 1 M NaCl at pH 7.0 and 23°C:

	metal	k_{obs} (min^{-1})
10	none	n.d.
	Mg²⁺	2.3×10^{-3}
	Mn ²⁺	6.8×10^{-3}
	Zn ²⁺	4.2×10^{-2}
	Pb ²⁺	1.1×10^{-2}

15

Thus, the lead compound is active in the presence of all four divalent metals, even though it was selected for activity in the presence of Mg²⁺. Conversely, DNA molecules that were selected for activity in the presence of Mn²⁺, Zn²⁺, or Pb²⁺ did not show any activity in the presence of Mg²⁺.

20

In addition, the population of DNAs obtained after six rounds of *in vitro* selection in the presence of Mg²⁺, when prepared as all-phosphorothioate-containing DNA analogs, showed Mg²⁺-dependent cleavage activity at an observed rate of $\sim 10^{-3} \text{ min}^{-1}$. The phosphorothioate-containing analogs were prepared enzymatically so as to have an *R_p* configuration at each stereocenter. Such compounds are relatively resistant to degradation by cellular nucleases compared to unmodified DNA.

25

The lead compound was re-randomized at 40 nucleotide positions (underlined), introducing mutations at a frequency of 15% (5% probability of each of the three possible base substitutions). The re-randomized population was subjected to seven additional rounds of *in vitro* selection. During the last four rounds, molecules that were reactive in the presence of 1 mM Pb²⁺ were removed from the population before the remainder were challenged to react in the presence of 1 mM Mg²⁺. Individual clones were isolated following the seventh round and the nucleotide sequence of 14 of these clones was determined. All of the sequences began with: 5' GGG ACG AAT TCT AAT ACG ACT CAC TAT **rA** GG AAG AGA TGG CGA CAT CTC (SEQ ID NO 23, from position 1 to 48), and ended with: GTG ACG GTA AGC TTG GCA C 3' (SEQ ID NO 23, from position 89 to 107).

30

35

The segment in the middle, corresponding to the 40 partially-randomized

-46-

positions (N_{40} , SEQ ID NO 23, from position 49 to 88), varied as follows:

- 5 (4) TAC AGC GAT TCA CCC TTG TTT AAG GGT TAC ACC CAT GTT A
(SEQ ID NO 31)
- (2) ATC AGC GAT TAA CGC TTG TTT CAA TGT TAC ACC CAT GTT A
(SEQ ID NO 32)
- (2) TTC AGC GAT TAA CGC TTA TTT TAG CGT TAC ACC CAT GTT A
(SEQ ID NO 33)
- 10 (1) ATC AGC GAT TCA CCC TTG TTT TAA GGT TGC ACC CAT GTT A
(SEQ ID NO 34)
- (1) ATC AGC GAT TCA CCC TTG TTT AAG CGT TAC ACC CAT GTT G
(SEQ ID NO 35)
- (1) ATC AGC GAT TCA CCC TTG TTT TAA GGT TAC ACC CAT GTT A
(SEQ ID NO 36)
- 15 (1) ATC AGC GAT TAA CGC TTA TTT TAG CGT TAC ACC CAT GTT A
(SEQ ID NO 37)
- (1) ATC AGC GAT TAA CGC TTG TTT TAG TGT TGC ACC CAT GTT A
(SEQ ID NO 38)
- (1) ATC AGC GAT TAA CGC TTA TTT TAG CAT TAC ACC CAT GTT A
20 (SEQ ID NO 39).

The number in parentheses indicates the number of clones having that particular sequence. Nucleotides shown in bold are those that differ compared to the lead compound.

25 Formal analysis of the cleavage activity of these clones is ongoing. The population as a whole exhibits Mg^{2+} -dependent cleavage activity at an observed rate of $\sim 10^{-2} \text{ min}^{-1}$, with a comparable level of activity in the presence of Pb^{2+} .

30 Figures 6A and 6B provide two-dimensional illustrations of a "progenitor" catalytic DNA molecule and one of several catalytic DNA molecules obtained via the selective amplification methods disclosed herein, respectively. Figure 6A illustrates an exemplary molecule from the starting pool, showing the overall configuration of the molecules represented by SEQ ID NO 23. As illustrated, various complementary nucleotides flank the random (N_{40}) region.

35 Figure 6B is a diagrammatic representation of one of the Mg^{2+} -dependent catalytic DNA molecules (or "DNAzymes") generated via the within-described procedures. The location of the ribonucleotide in the substrate nucleic acid is indicated via the arrow. (The illustrated molecule includes the sequence identified herein as SEQ

-47-

ID NO 25, as well as "beginning" and "ending" sequences of SEQ ID NO 23.)

Endonuclease activity is continuing to be enhanced in each of the
aforementioned "families" via *in vitro* evolution, as disclosed herein, so it is anticipated
that enzymatic DNA molecules of increasingly desirable specificities may be generated
successfully using the within-disclosed guidelines.

Example 5

Cleavage of Larger RNA Sequences

As an extension of the foregoing, we have developed DNA enzymes that cleave
an all-RNA substrate, rather than a single ribonucleotide embedded within an otherwise
all-DNA substrate as demonstrated above. (Also see R.R. Breaker & G.F. Joyce, Chem.
& Biol. 1: 223-229 (1994); R.R. Breaker & G.F. Joyce, Chem. & Biol. 2: 655-660
(1995)). As a target sequence, we chose a stretch of 12 highly-conserved nucleotides
within the U5 LTR region of HIV-1 RNA, having the sequence
5' GUAACUAGAGAU 3' (SEQ ID NO 49).

Following the methods described in the previous examples, we generated a pool
of 1014 DNA molecules that have the following composition:

5'- GGAAAA r(GUAACUAGAGAU) GGAAGAGATGGCGAC N₅₀
CGGTAAGCTTGGCAC -3' (SEQ ID NO 50),

where N is an equimolar mixture of the deoxyribonucleotides G, A, T, and C, and where
the sequence identified as "r(GUAACUAGAGAU)" is comprised of *ribonucleotides*.
(Optionally, one may alter the initial 5' nucleotide sequence, e.g., by adding an
additional dA residue to the sequence preceding the ribonucleotide portion at the 5' end,
thus causing the initial sequence to read "GGAAAAA" and causing SEQ ID NO 50 to be
99 residues in length. Clearly, this is but one example of the modifications that may be
made in order to engineer specific enzymatic DNA molecules, as disclosed in detail
herein.)

The enzymatic DNA molecules thus produced were selected for their ability to
cleave a phosphoester that lies within the embedded RNA target sequence. Ten rounds
of *in vitro* selective amplification were carried out, based on the enzymatic DNA
molecules' activity in the presence of 10 mM Mg²⁺ at pH 7.5 and 37°C. During the
selection process, there was competition for "preferred" cleavage sites as well as for the
"best" catalyst that cleaves at each such preferred site. Two sites and two families of
catalysts emerged as possessing the most efficient cleavage capabilities (see Fig. 7).

Figure 7 illustrates some of the results of ten rounds of *in vitro* selective
amplification carried out essentially as described herein. As shown, two sites and two
families of catalysts emerged as displaying the most efficient cleavage of the target
sequence. Cleavage conditions were essentially as indicated in Fig. 7, namely, 10mM

-48-

Mg²⁺, pH 7.5, and 37°; data collected after the reaction ran for 2 hours is shown. Cleavage (%) is shown plotted against the number of generations (here, 0 through 10). The number/prevalence of catalytic DNA molecules capable of cleaving the target sequence at the indicated sites in the substrate is illustrated via the vertical bars, with cleavage at G↓UAACUAGAGAU shown by the striped bars, and with cleavage at GUAACUA↓GAGAU illustrated via the open (lightly-shaded) bars. In Figure 7, as herein, the arrow (↓) indicates the site between two neighboring nucleotides at which cleavage occurs.

5
10
Various individuals from the population obtained after the 8th and 10th rounds of selective amplification were cloned. The nucleotide sequences of 29 individuals from the 8th round and 32 individuals from the 10th round were then determined (see Tables 2 and 3, respectively).

15
20
25
Under the heading "Nucleotide Sequence" in each of Tables 2 and 3 is shown the portion of each identified clone that corresponds to the 50 nucleotides that were randomized in the starting pool (i.e., N₅₀); thus, the entire nucleotide sequence of a given clone generally includes the nucleotide sequences preceding, following, and including the "N₅₀" segment, presuming the substrate sequence is attached and that self-cleavage has not occurred. For example, the entire sequence of a (non-self-cleaved) clone may generally comprise residue nos. 1-33 of SEQ ID NO 50, followed by the residues representing the randomized N₅₀ region, followed by residue nos. 84-98 of SEQ ID NO 50, or by residue nos. 1-34 of SEQ ID NO 51, followed by the residues representing the randomized N₅₀ region, followed by residue nos. 85-99 of SEQ ID NO 51. It is believed, however, that the N₅₀ (or N₄₀) region -- or a portion thereof -- of each clone is particularly important in determining the specificity and/or activity of a particular enzymatic DNA molecule. This is particularly evident in reactions in which the substrate and the DNAzyme are separate molecules (see, e.g., Figs. 8 and 9).

Clone numbers are designated as 8-x or 10-x for individuals obtained after the 8th or 10th rounds, respectively. SEQ ID NOS are also listed and correspond to the "N₅₀" region of each clone.

-49-

Table 2
Cloned Individuals from 8th Round of Amplification

Clone No.	SEQ ID NO	"N ₅₀ " Nucleotide Sequence (5'-3')
5	8-2 52	CCA ATA GTG CTA CTG TGT ATC TCA ATG CTG GAA ACA CGG GTT ATC TCC CG
	8-4 53	CCA AAA CAG TGG AGC ATT ATA TCT ACT CCA CAA AGA CCA CTT TTC TCC CG
10	8-5 ¹ 54	ATC CGT ACT AGC ATG CAG ACA GTC TGT CTG CTT TTT CAT TAC TCA CTC CC
	8-14 55	CAA TTC ATG ATG ACC AAC TCT GTC AAC ACG CGA ACT TTT AAC ACT GGC A
	8-17 ² 56	CTT CCA CCT TCC GAG CCG GAC GAA GTT ACT TTT TAT CAC ACT ACG TAT TG
15	8-3 57	GGC AAG AGA TGG CAT ATA TTC AGG TAA CTG TGG AGA TAC CCT GTC TGC CA
	8-6 58	CTA GAC CAT TCA CGT TTA CCA AGC TAT GGT AAG AAC TAG AAT CAC GCG TA
20	8-8 59	CGT ACA CGT GGA AAA GCT ATA AGT CAA GTT CTC ATC ATG TAC CTG ACC GC
	8-10 60	CAG TGA TAC ATG AGT GCA CCG CTA CGA CTA AGT CTG TAA CTT ATT CTA CC
	8-22 61	ACC GAA TTA AAC TAC CGA ATA GTG TGG TTT CTA TGC TTC TTC TTC CCT GA
25	8-11 62	CAG GTA GAT ATA ATG CGT CAC CGT GCT TAC ACT CGT TTT ATT AGT ATG TC
	8-21 63	CCC TAC AAC ACC ACT GGG CCC AAT TAG ATT AAC GCT ATT TTA TAA CTC G
30	8-12 64	CCA AAC GGT TAT AAG ACT GAA AAC TCA ATC AAT AGC CCA ATC CTC GCC C
	8-13 65	CAC ATG TAT ACC TAA GAA ATT GGT CCC GTA GAC GTC ACA GAC TTA CGC CA
	8-23 66	CAC AAC GAA AAC AAT CTT CCT TGG CAT ACT GGG GAG AAA GTC TGT TGT CC
35	8-40 67	CAC ACG AAC ATG TCC ATT AAA TGG CAT TCC GTT TTT CGT TCT ACA TAT GC

	8-24	68	CAG AAC GAG GGT CTT GTA AGA CTA CAC CTC CTC AGT GAC AAT AAT CCT G
	8-26	69	CAC TAC AGC CTG ATA TAT ATG AAG AAC AGG CAA CAA GCT TAT GCA CTG G
5	8-27	70	GGG TAC ATT TAT GAT TCT CTT ATA AAG AGA ATA TCG TAC TCT TTT CCC CA
	8-28	71	CCA AAG TAC ATT CCA ACC CCT TAT ACG TGA AAC TTC CAG TAG TTT CCT A
10	8-29	72	CTT GAA GAT CCT CAT AAG ACG ATT AAA CAA TCC ACT GGA TAT AAT CCG GA
	8-34	73	CGA ATA GTG TCC ATG ATT ACA CCA ATA ACT GCC TGC CTA TCA TGT TTA TG
	8-35	74	CCA AGA GAG TAT CGG ATA CAC TTG GAA CAT AGC TAA CTC GAA CTG TAC CA
15	8-36	75	CCA CTG ATA AAT AGG TAA CTG TCT CAT ATC TGC CAA TCA TAT GCC GTA
	8-37	76	CCC AAA TTA TAA ACA ATT TAA CAC AAG CAA AAG GAG GTT CAT TGC TCC GC
20	8-39	77	CAA TAA ACT GGT GCT AAA CCT AAT ACC TTG TAT CCA AGT TAT CCT CCC CC

¹ identical to 10-4, 10-40

25 ² identical to 8-20, 8-32, 8-38, 10-1, 10-34; 1 mutation to 10-11; 3 mutations to 10-29

Table 3

Cloned Individuals from 10th Round of Amplification

Clone No.	SEQ ID NO	"N ₅₀ " Nucleotide Sequence (5'-3')
30	10-3 ³ 78	CCG AAT GAC ATC CGT AGT GGA ACC TTG CTT TTG ACA CTA AGA AGC TAC AC
	10-10 79	CCA TAA CAA ATA CCA TAG TAA AGA TCT GCA TTA TAT TAT ATC GGT CCA CC
35	10-12 80	CAG AAC AAA GAT CAG TAG CTA AAC ATA TGG TAC AAA CAT ACC ATC TCG CA
	10-14 81	CCT TTA GTT AGG CTA GCT ACA ACG ATT TTT CCC TGC TTG GCA ACG ACA C

-51-

10-15 82 CTC CCT ACG TTA CAC CAG CGG TAC GAA TTT TCC ACG AGA GGT
AAT CCG CA

10-19 83 CGG CAC CTC TAG TTA GAC ACT CCG GAA TTT TTC CCC

10-39 84 CGG CAC CTC TAG TTA GAC ACT CCG GAA TTT TAG CCT ACC ATA
5 GTC CGG T

10-23 85 CCC TTT GGT TAG GCT AGC TAC AAC GAT TTT TCC CTG CTT GAA
TTG TA

10-27⁴ 86 CCC TTT GGT TAG GCT AGC TAC AAC GAT TTT TCC CTG CTT GAC
CTG TTA CGA

10 10-31 87 CCT TTA GTT AGG CTA GCT ACA ACG ATT TTT CCC TGC TTG GAA
CGA CAC

10-18 88 CAT GGC TTA ATC ATC CTC AAT AGA AGA CTA CAA GTC GAA TAT
GTC CCC CC

10-20 89 CAA CAG AGC GAG TAT CAC CCC CTG TCA ATA GTC GTA TGA AAC
15 ATT GGG CC

10-6 90 TAC CGA CAA GGG GAA TTA AAA GCT AGC TGG TTA TGC AAC CCT
TTT CGC A

10-7 91 CTC GAA ACA GTG ATA TTC TGA ACA AAC GGG TAC TAC GTG TTC
AGC CCC C

20 10-8 92 CCA ATA ACG TAA CCC GGT TAG ATA AGC ACT TAG CTA AGA TGT
TTA TCC TG

10-16 93 CAA TAC AAT CGG TAC GAA TCC AGA AAC ATA ACG TTG TTT CAG
AAT GGT CC

10-21 94 GCA ACA ACA AGA ACC AAG TTA CAT ACA CGT TCA TCT ATA CTG
25 AAC CCC CA

10-24 95 CCT TTG AGT TCC TAA ATG CCG CAC GGT AAG CTT GGC ACA CTT
TGA CTG TA

10-28 96 CAA AGA TCT CAC TTT GGA AAT GCG AAA TAT GTA TAT TCG CCC
TGT CTG C

30 10-33 97 CCA CGT AGA ATT ATC TGA TTT ATA ACA TAA CGC AGG ATA ACT
CTC GCC CA

10-35 98 CAC AAG AAA GTG TCG TCT CCA GAT ATT TGA GTA CAA GGA ACT
ACG CCC

10-36 99 CAT GAA GAA ATA GGA CAT TCT ACA GGC TGG ACC GTT ACT ATG
35 CCT GTA GG

10-37 100 CAT AGG ATA ATC ATG GCG ATG CTT ATG ACG TGT ACA TCT ATA
CCT T

-52-

10-38 101 CAG ATG ATC TTC CTT TAA AGA CTA CCC TTT AAA GAA ACA TAA
GGT ACC CC

- 5 ³ 1 mutation to 10-5
 ⁴ 1 mutation to 10-30

10 The self-cleavage activity of various clones was subsequently measured. Clones
8-5, 8-17, and 10-3 were found to cleave efficiently at the site 5' GUAACUAGAGAU
3', while clones 10-14, 10-19 and 10-27 were found to cleave efficiently at the site 5'
GUAACUAGAGAU 3'. When the RNA portion of the molecule was extended to the
sequence 5' GGAAAAGUAACUAGAGAUGGAAG 3' (residue nos. 1-24 of SEQ ID NO
15 51), clones 8-17, 10-14, and 10-27 retained full activity, while clones 8-5, 10-3, and
10-19 showed diminished activity. Subsequently, clone 10-23 was found to exhibit a
high level of activity in the self-cleavage reaction involving the extended RNA domain.

 It should also be noted, in the event one of skill in the relevant art does not
appreciate same, that the nucleotide sequences preceding and following the "N₅₀"
segments of the polynucleotide molecules engineered according to the teachings of the
20 present invention disclosure may be altered in a variety of ways in order to generate
enzymatic DNA molecules of particular specificities. For example, while residue nos. 1-
24 of SEQ ID NO 51 are described herein as RNA nucleotides, they may alternatively
comprise DNA, RNA, or composites thereof. (Thus, for example, SEQ ID NO 51 could
easily be altered so that nucleic acid residue nos. 1-7 would comprise DNA, residue nos.
25 8-19 would comprise RNA, residue nos. 20-99 would comprise DNA, and so on.)
Similarly, the nucleotides following the "N₅₀" region may comprise RNA, DNA, or
composites thereof. The length of the regions preceding and following the "N₅₀" (or
"N₄₀" -- see Example 4) region(s) may also be varied, as disclosed herein. Further,
sequences preceding and/or following N₅₀ or N₄₀ regions may be shortened, expanded,
30 or deleted in their entirety.

 Moreover, as noted above, we selected a specific region of HIV-1 RNA as the
target sequence in the methods described in this Example; such a sequence is not the
only sequence one may use as a target. Clearly, one of skill in the relevant art may
follow our teachings herein to engineer and design enzymatic DNA molecules with
35 specificity for other target sequences. As disclosed herein, such target sequences may
be constructed or inserted into larger sequences comprising DNA, RNA, or composites
thereof, as illustrated by SEQ ID NOS 50 and 51.

 The self-cleavage reaction was easily converted to an intermolecular cleavage

-53-

reaction by dividing the enzyme and substrate domains into separate molecules. Clones 8-17 and 10-23 were chosen as prototype molecules. Both were shown to act as DNA enzymes in the cleavage of a separate all-RNA substrate in a reaction that proceeds with multiple turnover (Fig. 8). The substrate binding arms were subsequently reduced to 7 base-pairs on each side of the unpaired nucleotide that demarcates the cleavage site (Fig. 9).

Figure 8 illustrates the nucleotide sequences, cleavage sites, and turnover rates of two catalytic DNA molecules of the present invention, clones 8-17 and 10-23. Reaction conditions were as shown, namely, 10mM Mg^{2+} , pH 7.5, and 37°C. The DNAzyme identified as clone 8-17 is illustrated on the left, with the site of cleavage of the RNA substrate indicated by the arrow. The substrate sequence (5' - GGAAAAAGUAACUAGAGAUGGAAG - 3') -- which is separate from the DNAzyme (i.e., intermolecular cleavage is shown) -- is labeled as such. Similarly, the DNAzyme identified herein as 10-23 is shown on the right, with the site of cleavage of the RNA substrate indicated by the arrow. Again, the substrate sequence is indicated. For the 8-17 enzyme, the turnover rate was approximately 0.6 hr^{-1} ; for the 10-23 enzyme, the turnover rate was approximately 1 hr^{-1} .

As illustrated in Fig. 8, the nucleotide sequence of the clone 8-17 catalytic DNA molecule capable of cleaving a separate substrate molecule was as follows:
5'-CTTCCACCTTCCGAGCCGGACGAAGTTACTTTTT-3' (residue nos. 1-34 of SEQ ID NO 56). In that same figure, the nucleotide sequence of the clone 10-23 catalytic DNA molecule capable of cleaving a separate substrate molecule was as follows:
5'-CTTTGGTTAGGCTAGCTACAACGATTTTTCC-3' (residue nos. 3-33 of SEQ ID NO 85).

Figure 9 further illustrates the nucleotide sequences, cleavage sites, and turnover rates of two catalytic DNA molecules of the present invention, clones 8-17 and 10-23. Reaction conditions were as shown, namely, 10mM Mg^{2+} , pH 7.5, and 37°C. As in Fig. 8, the DNAzyme identified as clone 8-17 is illustrated on the left, with the site of cleavage of the RNA substrate indicated by the arrow. The substrate sequence (5' - GGAAAAAGUAACUAGAGAUGGAAG - 3') --which is separate from the DNAzyme (i.e., intermolecular cleavage is shown) -- is labeled as such. Similarly, the DNAzyme identified herein as 10-23 is shown on the right, with the site of cleavage of the RNA substrate indicated by the arrow. Again, the substrate sequence is indicated. For the 8-17 enzyme, k_{obs} was approximately 0.002 min^{-1} ; for the 10-23 enzyme, the value of k_{obs} was approximately 0.01 min^{-1} .

As illustrated in Fig. 9, the nucleotide sequence of the clone 8-17 catalytic DNA molecule capable of cleaving a separate substrate molecule was as follows:

-54-

5'-CCACCTCCGAGCCGGACGAAGTTACT-3' (residue nos. 4-30 of SEQ ID NO 56). In that same figure, the nucleotide sequence of the clone 10-23 catalytic DNA molecule capable of cleaving a separate substrate molecule was as follows:

5'-CTAGTTAGGCTAGCTACAACGATTTTCC-3' (residue nos. 5-33 of SEQ ID NO 85, with "CTA" substituted for "TTG" at the 5' end).

5
10
The catalytic rate of the RNA-cleaving DNA enzymes has yet to be fully optimized. As disclosed above and as reported in previous studies, we have been able to improve the catalytic rate by partially randomizing the prototype molecule and carrying out additional rounds of selective amplification. We have found, however, that the K_m for Mg^{2+} is approximately 5 mM and 2 mM for the 8-17 and 10-23 DNA enzymes, respectively, measured at pH 7.5 and 37°C; this is certainly compatible with intracellular conditions.

15
The foregoing specification, including the specific embodiments and examples, is intended to be illustrative of the present invention and is not to be taken as limiting. Numerous other variations and modifications can be effected without departing from the true spirit and scope of the present invention.

-55-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (I) APPLICANT: The Scripps Research Institute
- (ii) TITLE OF INVENTION: ENZYMATIC DNA MOLECULES
- (iii) NUMBER OF SEQUENCES: 101
- (iv) CORRESPONDENCE ADDRESS:
- 10 (A) ADDRESSEE: The Scripps Research Institute
- (B) STREET: 10666 North Torrey Pines Road, TPC-8
- (C) CITY: La Jolla
- (D) STATE: California
- (E) COUNTRY: United States
- (F) ZIP: 92037
- 15 (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- 20 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER: PCT/US95/
- (B) FILING DATE: 01-DEC-1995
- (C) CLASSIFICATION:
- 25 (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/472,194
- (B) FILING DATE: 07-JUN-1995
- 30 (viii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/349,023
- (B) FILING DATE: 02-DEC-1994
- 35 (ix) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Logan, April C.
- (B) REGISTRATION NUMBER: 33,950
- (C) REFERENCE/DOCKET NUMBER: 463.2 PC

-56-

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (619) 554-2937

(B) TELEFAX: (619) 554-6312

5

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

10

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGGTAAGCTT GGCAC

15

20

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

25

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30

(ix) FEATURE:

(A) NAME/KEY: misc_difference

(B) LOCATION: replace(8, "")

(D) OTHER INFORMATION: /standard_name= "ADENOSINE
RIBONUCLEOTIDE"

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCACTATNAG GAAGAGATGG

20

5 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ACACATCTCT GAAGTAGCGC CGCCGTATAG TGACGCTA

38

20 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 80 base pairs

(B) TYPE: nucleic acid

25 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GTGCCAAGCT TACCGNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 60
NNNNNGTCGC CATCTCTTCC 80

35

-58-

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10

(ix) FEATURE:

15

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 28
- (D) OTHER INFORMATION: /standard_name= "2'3' CYCLIC
PHOSPHATE"

(ix) FEATURE:

20

- (A) NAME/KEY: misc_difference
- (B) LOCATION: replace(28, "")
- (D) OTHER INFORMATION: /standard_name= "ADENOSINE
RIBONUCLEOTIDE"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GGGACGAATT CTAATACGAC TCACTATN

28

(2) INFORMATION FOR SEQ ID NO:6:

30

(i) SEQUENCE CHARACTERISTICS:

35

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

-59-

(ix) FEATURE:

(A) NAME/KEY: misc_difference

(B) LOCATION: replace(28, "")

5 (D) OTHER INFORMATION: /standard_name= "ADENOSINE
RIBONUCLEOTIDE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

10 GGGACGAATT CTAATACGAC TCACTATN

28

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

(A) NAME/KEY: misc_difference

(B) LOCATION: replace(8, "")

25 (D) OTHER INFORMATION: /standard_name= "ADENOSINE
RIBONUCLEOTIDE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

30 TCACTATNGG AAGAGATGG

19

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 8 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

-60-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(ix) FEATURE:

(A) NAME/KEY: misc_difference

(B) LOCATION: replace(8, "")

(D) OTHER INFORMATION: /standard_name= "ADENOSINE
NUCLEOTIDE"

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCACTATN

8

15

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

20

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CCATCTCTTC CTATAGTGAG TCCGGCTGCA

30

(2) INFORMATION FOR SEQ ID NO:10:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

35

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

-61-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GTGCCAAGCT TACCG

15

5 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 43 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CTGCAGAATT CTAATACGAC TCACTATAGG AAGAGATGGC GAC

43

(2) INFORMATION FOR SEQ ID NO:12:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

30 (A) NAME/KEY: misc_difference

(B) LOCATION: replace(8, "")

(D) OTHER INFORMATION: /standard_name= "ADENOSINE
RIBONUCLEOTIDE"

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TCACTATNGG AAGAGATGG

19

-62-

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10

(ix) FEATURE:

- (A) NAME/KEY: misc_difference
(B) LOCATION: replace(28, "")
(D) OTHER INFORMATION: /standard_name= "ADENOSINE
15 RIBONUCLEOTIDE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGGACGAATT CTAATACGAC TCACTATNGG AAGAGATGGC GAC

43

20

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TCACACATCT CTGAAGTAGC GCCGCCGTAT GTGACGCTAG GGGTTCGCCT

50

35

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

-63-

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

10

GGGGGAACG CCGTAACAAG CTCTGAACTA GCGGTTGCGA TATAGTCGTA

50

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGGGACTCCG TAGCCCATTG CTTTTTGCAG CGTCAACGAA TAGCGTATTA

50

25

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CCACCATGTC TTCTCGAGCC GAACCGATAG TTACGTCATA CCTCCCGTAT

50

-64-

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GCCAGATTGC TGCTACCAGC GGTACGAAAT AGTGAAGTGT TCGTGACTAT 50

15

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ATAGGCCATG CTTTGGCTAG CGGCACCGTA TAGTGTACCT GCCCTTATCG 50

(2) INFORMATION FOR SEQ ID NO:20:

30

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

-65-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

TCTGCTCTCC TCTATTCTAG CAGTGCAGCG AAATATGTCG AATAGTCGGT 50

5 (2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TTGCCAGCA TAGTCGGCAG ACGTGGTGT AGCGACACGA TAGGCCCGT 50

(2) INFORMATION FOR SEQ ID NO:22:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

30

TTGCTAGCTC GGCTGAACTT CTGTAGCGCA ACCGAAATAG TGAGGCTTGA 50

(2) INFORMATION FOR SEQ ID NO:23:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 107 base pairs

(B) TYPE: nucleic acid

-66-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(ix) FEATURE:

(A) NAME/KEY: misc_difference

(B) LOCATION: replace(28, "")

(D) OTHER INFORMATION: /standard_name= "ADENOSINE

10

RIBONUCLEOTIDE"

/label= rA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

15

GGGACGAATT CTAATACGAC TCACTATNGG AAGAGATGGC GACATCTCNN NNNNNNNNNN 60

NNNNNNNNNN NNNNNNNNNN NNNNNNNNGT GACGGTAAGC TTGGCAC 107

(2) INFORMATION FOR SEQ ID NO:24:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 49 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

30

CCGCCACCT CTTTACGAG CCTGTACGAA ATAGTGCTCT TGTTAGTAT

49

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

35

(A) LENGTH: 48 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

-67-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TCTCTTCAGC GATGCACGCT TGTTTAAATG TTGCACCCAT GTTAGTGA

48

(2) INFORMATION FOR SEQ ID NO:26:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 46 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

15

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

20

TCTCATCAGC GATTGAACCA CTTGGTGGAC AGACCCATGT TAGTGA

46

(2) INFORMATION FOR SEQ ID NO:27:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 49 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

35

CCGCCCACCT CTTTACGAG CCTGTACGAA ATAGTGTCT TGTTAGTAT

49

(2) INFORMATION FOR SEQ ID NO:28:

-68-

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 49 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

10

CCGCCACCT CTTTACGAG CCTGTACGAA ATAGTGCTCT CGTTAGTAT

49

(2) INFORMATION FOR SEQ ID NO:29:

15

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

25

TTCAGACTT AGTCCATCAC ACTCTGTGCA TATGCCTGCT TGATGTGA

48

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30

35

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

-69-

CTCTCATCTG CTAGCACGCT CGAATAGTGT CAGTCGATGT GA

42

(2) INFORMATION FOR SEQ ID NO:31:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

15 TACAGCGATT CACCCTTGTT TAAGGGTTAC ACCCATGTTA

40

(2) INFORMATION FOR SEQ ID NO:32:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

30 ATCAGCGATT AACGCTTGTT TCAATGTTAC ACCCATGTTA

40

(2) INFORMATION FOR SEQ ID NO:33:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-70-

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

5 TTCAGCGATT AACGCTTATT TTAGCGTTAC ACCCATGTTA 40

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

20 ATCAGCGATT CACCCTTGTT TTAAGGTTGC ACCCATGTTA 40

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

ATCAGCGATT CACCCTTGTT TAAGCGTTAC ACCCATGTTG 40

35 (2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

-71-

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

10

ATCAGCGATT CACCCTTGTT TTAAGGTAC ACCCATGTTA

40

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

ATCAGCGATT AACGCTTATT TTAGCGTTAC ACCCATGTTA

40

25

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

-72-

ATCAGCGATT AACGCTTGTT TTAGTGTTGC ACCCATGTTA

40

(2) INFORMATION FOR SEQ ID NO:39:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

15 ATCAGCGATT AACGCTTATT TTAGCATTAC ACCCATGTTA

40

(2) INFORMATION FOR SEQ ID NO:40:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

GCCATGCTTT

10

30

(2) INFORMATION FOR SEQ ID NO:41:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-73-

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

5 CTCTATTTCT 10

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

TATGTGACGC TA 12

20

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

TATAGTCGTA 10

35 (2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

-74-

- (A) LENGTH: 11 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

10

ATAGCGTATT A

11

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 13 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

ATAGTTACGT CAT

13

25

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 14 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

-75-

AATAGTGAAG TGTT

14

(2) INFORMATION FOR SEQ ID NO:47:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

15 ATAGGCCCGG T

11

(2) INFORMATION FOR SEQ ID NO:48:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 14 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

AATAGTGAGG CTTG

14

30

(2) INFORMATION FOR SEQ ID NO:49:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-76-

- (ii) MOLECULE TYPE: RNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GUAACUAGAG AU

12

(2) INFORMATION FOR SEQ ID NO:50:

10

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 98 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

15

- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:
 - (A) NAME/KEY: misc_feature
 - (B) LOCATION: 7..18
 - (D) OTHER INFORMATION: /note= "Position 7-18 is RNA; the remainder of the sequence is DNA."

20

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GGAAAAGUAA CUAGAGAUGG AAGAGATGGC GACNNNNNNNN NNNNNNNNNNN NNNNNNNNNNN 60
 NNNNNNNNNNN NNNNNNNNNNN NNCCGGTAAG CTTGGCAC 98

30

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 99 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

35

-77-

- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:

- 5 (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..24
- (D) OTHER INFORMATION: /note= "Positions 1-24 is RNA; the remainder of the sequence is DNA."

- 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

```
GGAAAAAGUA ACUAGAGAUG GAAGAGATGG CGACNNNNNNN NNNNNNNNNN NNNNNNNNNN 60
NNNNNNNNNN NNNNNNNNNN NNNNCGGTAA GCTTGGCAC 99
```

- 15 (2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 50 base pairs
 - (B) TYPE: nucleic acid
 - 20 (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- 25 (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

```
CCAATAGTGC TACTGTGTAT CTCAATGCTG GAAACACGGG TTATCTCCCG 50
```

30

- (2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 50 base pairs
 - 35 (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

-78-

- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CCAAAACAGT GGAGCATTAT ATCTACTCCA CAAAGACCAC TTTTCTCCCG

50

(2) INFORMATION FOR SEQ ID NO:54:

10

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 50 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

15

- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

ATCCGTACTA GCATGCAGAC AGTCTGTCTG CTTTTTCATT ACTCACTCCC

50

25 (2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 49 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

30

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

CAATTCATGA TGACCAACTC TGTC AACACG CGAACTTTTA ACACTGGCA

49

-79-

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

5 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

15 CTTCCACCTT CCGAGCCGGA CGAAGTTACT TTTTATCACA CTACGTATTG 50

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

30 GGCAAGAGAT GGCATATATT CAGGTAAGT TGGAGATAACC CTGTCTGCCA 50

(2) INFORMATION FOR SEQ ID NO:58:

35 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

-80-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

10 CTAGACCATT CACGTTTACC AAGCTATGGT AAGAACTAGA ATCACGCGTA

50

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

25

CGTACACGTG GAAAAGCTAT AAGTCAAGTT CTCATCATGT ACCTGACCGC

50

(2) INFORMATION FOR SEQ ID NO:60:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

-81-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

5 CAGTGATACA TGAGTGCACC GCTACGACTA AGTCTGTAAC TTATTCTACC 50

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

20 ACCGAATTAA ACTACCGAAT AGTGTGGTTT CTATGCTTCT TCTTCCCTGA 50

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CAGGTAGATA TAATGCGTCA CCGTGCTTAC ACTCGTTTTA TTAGTATGTC 50

-82-

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

5

(A) LENGTH: 49 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

15

CCCTACAACA CCACTGGGCC CAATTAGATT AACGCTATTT TATAACTCG

49

(2) INFORMATION FOR SEQ ID NO:64:

(i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 49 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

30

CCAAACGGTT ATAAGACTGA AACTCAATC AATAGCCCAA TCCTCGCCC

49

(2) INFORMATION FOR SEQ ID NO:65:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

-83-

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)
- 5 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

10 CACATGTATA CCTAAGAAAT TGGTCCCGTA GACGTCACAG ACTTACGCCA 50

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
- 15 (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)
- 20 (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

25 CACAACGAAA ACAATCTTCC TTGGCATACT GGGGAGAAAAG TCTGTTGTCC 50

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
- 30 (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 35 (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO

-84-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

5 CACACGAACA TGCCATTAA ATGGCATTCC GTTTTTCGTT CTACATATGC 50

(2) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 49 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

20 CAGAACGAGG GTCTTGTAAG ACTACACCTC CTCAGTGACA ATAATCCTG 49

(2) INFORMATION FOR SEQ ID NO:69:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 49 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

CACTACAGCC TGATATATAT GAAGAACAGG CAACAAGCTT ATGCACTGG 49

-85-

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

15 GGGTACATTT ATGATTCTCT TATAAAGAGA ATATCGTACT CTTTTCCCCA 50

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 49 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

30 CCAAAGTACA TTCCAACCCC TTATACGTGA AACTTCCAGT AGTTTCCTA 49

(2) INFORMATION FOR SEQ ID NO:72:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid

-86-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

10 CTTGAAGATC CTCATAAGAC GATTAAACAA TCCACTGGAT ATAATCCGGA

50

(2) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

25

CGAATAGTGT CCATGATTAC ACCAATAACT GCCTGCCTAT CATGTTTATG

50

(2) INFORMATION FOR SEQ ID NO:74:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

-87-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

5 CCAAGAGAGT ATCGGATACA CTTGGAACAT AGCTAACTCG AACTGTACCA 50

(2) INFORMATION FOR SEQ ID NO:75:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 48 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

20 CCACTGATAA ATAGGTAACT GTCTCATATC TGCCAATCAT ATGCCGTA 48

(2) INFORMATION FOR SEQ ID NO:76:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

CCCAAATTAT AAACAATTTA ACACAAGCAA AAGGAGGTTT ATTGCTCCGC 50

-88-

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

15 CAATAAACTG GTGCTAAACC TAATACCTTG TATCCAAGTT ATCCTCCCCC 50

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

30 CCGAATGACA TCCGTAGTGG AACCTTGCTT TTGACTACTAA GAAGCTACAC 50

(2) INFORMATION FOR SEQ ID NO:79:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid

-89-

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (iii) HYPOTHETICAL: NO

- (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

10 CCATAACAAA TACCATAGTA AAGATCTGCA TTATATTATA TCGGTCCACC 50

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs

- (B) TYPE: nucleic acid

- (C) STRANDEDNESS: single

- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (iii) HYPOTHETICAL: NO

- (iv) ANTI-SENSE: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

25 CAGAACAAAG ATCAGTAGCT AACATATGG TACAAACATA CCATCTCGCA 50

(2) INFORMATION FOR SEQ ID NO:81:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 49 base pairs

- (B) TYPE: nucleic acid

- (C) STRANDEDNESS: single

- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (iii) HYPOTHETICAL: NO

-90-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

5 CCTTTAGTTA GGCTAGCTAC AACGATTTTT CCCTGCTTGG CAACGACAC 49

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 50 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

20 CTCCTACGT TACACCAGCG GTACGAATTT TCCACGAGAG GTAATCCGCA 50

(2) INFORMATION FOR SEQ ID NO:83:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

CGGCACCTCT AGTTAGACAC TCCGGAATTT TTCCC 36

-91-

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 49 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

15 CCGCACCTCT AGTTAGACAC TCCGGAATTT TAGCCTACCA TAGTCCGGT

49

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

30

CCCTTTGGTT AGGCTAGCTA CAACGATTTT TCCCTGCTTG AATTGTA

47

(2) INFORMATION FOR SEQ ID NO:86:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 base pairs
(B) TYPE: nucleic acid

-92-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

10 CCCTTTGGTT AGGCTAGCTA CAACGATTTT TCCCTGCTTG ACCTGTTACG A 51

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 48 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

25 CCTTTAGTTA GGCTAGCTAC AACGATTTTT CCCTGCTTGG AACGACAC 48

(2) INFORMATION FOR SEQ ID NO:88:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

-93-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

5 CATGGCTTAA TCATCCTCAA TAGAAGACTA CAAGTCGAAT ATGTCCCCC 50

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

20 CAACAGAGCG AGTATCACCC CCTGTCAATA GTCGTATGAA ACATTGGGCC 50

(2) INFORMATION FOR SEQ ID NO:90:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 49 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

TACCGACAAG GGAATTAAA AGCTAGCTGG TTATGCAACC CTTTTCGCA 49

-94-

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 49 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

15

CTCGAAACAG TGATATTCTG AACAAACGGG TACTACGTGT TCAGCCCCC

49

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

20

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

30

CCAATAACGT AACCCGGTTA GATAAGCACT TAGCTAAGAT GTTTATCCTG

50

(2) INFORMATION FOR SEQ ID NO:93:

35

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid

-95-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

10 CAATACAATC GGTACGAATC CAGAAACATA ACGTTGTTTC AGAATGGTCC

50

(2) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

25

GCAACAACAA GAACCAAGTT ACATACACGT TCATCTATAC TGAACCCCCA

50

(2) INFORMATION FOR SEQ ID NO:95:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

-96-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

5 CCTTTGAGTT CCTAAATGCC GCACGGTAAG CTTGGCACAC TTTGACTGTA 50

(2) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 49 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

20 CAAAGATCTC ACTTTGGAAA TCGGAAATAT GTATATTCGC CCTGTCTGC 49

(2) INFORMATION FOR SEQ ID NO:97:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 50 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

CCACGTAGAA TTATCTGATT TATAACATAA CGCAGGATAA CTCTCGCCCA 50

-97-

(2) INFORMATION FOR SEQ ID NO:98:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 48 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

15 CACAAGAAAG TGTCGTCTCC AGATATTTGA GTACAAGGAA CTACGCC

48

(2) INFORMATION FOR SEQ ID NO:99:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 50 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

30 CATGAAGAAA TAGGACATTC TACAGGCTGG ACCGTTACTA TGCCTGTAGG

50

(2) INFORMATION FOR SEQ ID NO:100:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 46 base pairs
(B) TYPE: nucleic acid

-98-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

10 CATAGGATAA TCATGGCGAT GCTTATGACG TGTACATCTA TACCTT

46

(2) INFORMATION FOR SEQ ID NO:101:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

25

CAGATGATCT TCCTTTAAAG ACTACCCTTT AAAGAAACAT AAGGTACCCC

50

-99-

We Claim:

1. A catalytic DNA molecule having site-specific endonuclease activity.
2. The catalytic DNA molecule of claim 1, wherein said endonuclease activity is specific for a nucleotide sequence defining a cleavage site comprising single-stranded nucleic acid in a substrate nucleic acid sequence.
3. The catalytic DNA molecule of claim 2, wherein said single stranded nucleic acid comprises RNA, DNA, modified RNA, modified DNA, nucleotide analogs, or composites thereof.
4. The catalytic DNA molecule of claim 2, wherein said substrate nucleic acid comprises RNA, DNA, modified RNA, modified DNA, nucleotide analogs, or composites thereof.
5. The catalytic DNA molecule of claim 2, wherein said endonuclease activity comprises hydrolytic cleavage of a phosphoester bond at said cleavage site.
6. The catalytic DNA molecule of claim 1, wherein said molecule is single-stranded.
7. The catalytic DNA molecule of claim 1, wherein said molecule includes one or more hairpin loop structures.
8. The catalytic DNA molecule of claim 1, wherein said substrate nucleic acid sequence is attached to said catalytic DNA molecule.
9. The catalytic DNA molecule of claim 1, wherein said substrate nucleic acid sequence is not attached to said catalytic DNA molecule.
10. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule comprises a nucleotide sequence selected from the group consisting of:
SEQ ID NO 3 and SEQ ID NOS 14 through 22.
11. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule comprises a nucleotide sequence selected from the group consisting of:
SEQ ID NOS 23 through 30.
12. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule comprises a nucleotide sequence selected from the group consisting of:
SEQ ID NOS 31 through 39.
13. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule comprises a nucleotide sequence selected from the group consisting of:
SEQ ID NOS 52 through 101.
14. The catalytic DNA molecule of claim 11, 12, or 13, wherein said endonuclease activity is enhanced by the presence of Mg^{2+} .
15. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule has a substrate binding affinity of about 1 μM or less.

-100-

16. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule binds substrate with a K_D of less than about $0.1 \mu\text{M}$.

17. The catalytic DNA molecule of claim 2, wherein said nucleotide sequence defining said cleavage site comprises at least one nucleotide.

5 18. The catalytic DNA molecule of claim 1, wherein said endonuclease activity is enhanced by the presence of a divalent cation.

19. The catalytic DNA molecule of claim 18, wherein said divalent cation is selected from the group consisting of Pb^{2+} , Mg^{2+} , Mn^{2+} , Zn^{2+} , and Ca^{2+} .

10 20. The catalytic DNA molecule of claim 1, wherein said endonuclease activity is enhanced by the presence of a monovalent cation.

21. The catalytic DNA molecule of claim 20, wherein said monovalent cation is selected from the group consisting of Na^+ and K^+ .

15 22. The catalytic DNA molecule of claim 1, wherein said catalytic DNA molecule comprises a conserved core flanked by first and second substrate binding regions.

23. The catalytic DNA molecule of claim 22, further comprising one or more spacer nucleotides between said conserved core and said substrate binding region.

24. The catalytic DNA molecule of claim 22, wherein said conserved core comprises one or more conserved regions.

20 25. The catalytic DNA molecule of claim 24, wherein said one or more conserved regions includes a nucleotide sequence selected from the group consisting of:

CG;

CGA;

AGCG;

25 AGCCG;

CAGCGAT;

CTTGTTT; and

CTTATTT.

30 26. The catalytic DNA molecule of claim 24, further comprising one or more variable or spacer nucleotides between said conserved regions in said conserved core.

27. The catalytic DNA molecule of claim 22, wherein said first substrate binding region includes a nucleotide sequence selected from the group consisting of:

CATCTCT;

GCTCT;

35 TTGCTTTTT;

TGTCTTCTC;

TTGCTGCT;

-101-

GCCATGCTTT;
CTCTATTTCT
GTCGGCA;
CATCTCTTC; and
5 ACTTCT.

28. The catalytic DNA molecule of claim 22, wherein said second substrate binding region includes a nucleotide sequence selected from the group consisting of:

TATGTGACGCTA;
TATAGTCGTA;
10 ATAGCGTATTA;
ATAGTTACGTCAT;
AATAGTGAAGTGTT;
TATAGTGTA;
ATAGTCGGT;
15 ATAGGCCCGGT;
AATAGTGAGGCTTG; and
ATGNTG.

29. The catalytic DNA molecule of claim 22, further comprising a third substrate binding region, wherein said third region includes a nucleotide sequence
20 selected from the group consisting of:

TGTT;
TGTTA; and
TGTTAG.

30. The catalytic DNA molecule of claim 29, further comprising one or more
25 spacer regions between said substrate binding regions.

31. A composition comprising two or more populations of catalytic DNA molecules according to claim 1, wherein each population of catalytic DNA molecules is capable of cleaving a different nucleotide sequence in a substrate.

32. A composition comprising two or more populations of catalytic DNA
30 molecules according to claim 1, wherein each population of catalytic DNA molecules is capable of recognizing a different substrate.

33. A method of selecting a catalytic DNA molecule that cleaves a substrate nucleic acid sequence at a specific site, comprising the following steps:

- a. obtaining a population of single-stranded DNA molecules;
- 35 b. admixing nucleotide-containing substrate molecules with said population of single-stranded DNA molecules to form an admixture;
- c. maintaining said admixture for a sufficient period of time and under

-102-

predetermined reaction conditions to allow single-stranded DNA molecules in said population to cause cleavage of said substrate sequences, thereby producing substrate cleavage products;

d. separating said population of single-stranded DNA molecules from said substrate sequences and substrate cleavage products; and

e. isolating single-stranded DNA molecules that cleave nucleotide-containing substrate at a specific site from said population.

34. The method of claim 33, wherein said substrate comprises RNA.

35. The method of claim 33, wherein said DNA molecules that cleave said substrate at a specific site are tagged with an immobilizing agent.

36. The method of claim 35, wherein said agent comprises biotin.

37. The method of claim 35, wherein said isolating step further comprises exposing said tagged DNA molecules to a solid surface having avidin linked thereto, whereby said tagged DNA molecules become attached to said solid surface.

38. A method of cleaving a phosphoester bond, comprising:

a. admixing a catalytic DNA molecule capable of cleaving a substrate nucleic acid sequence at a defined cleavage site with a phosphoester bond-containing substrate, to form a reaction admixture; and

b. maintaining said admixture under predetermined reaction conditions to allow said catalytic DNA molecule to cleave said phosphoester bond, thereby producing a population of substrate products.

39. The method of claim 38, further comprising the steps of

a. separating said products from said catalytic DNA molecule; and

b. adding additional substrate to said catalytic DNA molecule to form a new reaction admixture.

40. The method of claim 38, wherein said substrate comprises RNA.

41. The method of claim 38, wherein said predetermined reaction conditions include the presence of a monovalent cation, a divalent cation, or both.

42. A method of engineering catalytic DNA molecules that cleave phosphoester bonds, comprising the following steps:

a. obtaining a population of single-stranded DNA molecules;

b. introducing genetic variation into said population to produce a variant population;

c. selecting individuals from said variant population that meet predetermined selection criteria;

d. separating said selected individuals from the remainder of said variant population; and

-103-

e. amplifying said selected individuals.

43. A non-naturally-occurring catalytic DNA molecule comprising a nucleotide sequence defining a conserved core flanked by one or more recognition domains, variable regions, and spacer regions.

5 44. The catalytic DNA molecule of claim 43, wherein said nucleotide sequence defines a first variable region contiguous or adjacent to the 5'-terminus of the molecule, a first recognition domain located 3'-terminal to the first variable region, a first spacer region located 3'-terminal to the first recognition domain, a first conserved region located 3'-terminal to the first spacer region, a second spacer region located 3'-terminal to the first conserved region, a second conserved region located 3'-terminal to the second spacer region, a second recognition domain located 3'-terminal to the second conserved region, and a second variable region located 3'-terminal to the second recognition domain.

10 45. The catalytic DNA molecule of claim 43, wherein said nucleotide sequence defines a first variable region contiguous or adjacent to the 5'-terminus of the molecule, a first recognition domain located 3'-terminal to the first variable region, a first spacer region located 3'-terminal to the first recognition domain, a first conserved region located 3'-terminal to the first spacer region, a second spacer region located 3'-terminal to the first conserved region, a second conserved region located 3'-terminal to the second spacer region, a second recognition domain located 3'-terminal to the second conserved region, a second variable region located 3'-terminal to the second recognition domain, and a third recognition domain located 3'-terminal to the second variable region.




FIGURE 1

2 / 9




FIGURE 2

Target
ribodenylation

substrate: 3' CAGCCGATAGAGAAAGG ATATCACTCAGCATAAATCTTAAGCAGGG 5'
 ||||| |||||

SEQ ID NO # of clones

SEQ ID NO	# of clones	5' sequence	3' sequence
14	8	5' TCACA <u>CATCTCT</u> GAAAT	TAT - GTGA - - CGCTA GGGGTCCCT - 3'
15	3	5' GGGGGAAACCGGTAAACA <u>G-CTCT</u> GAACT	TATAGT - - - CGTA - 3'
16	2	5' CCGACTCCGTAGCCCA <u>TGCCTTTT</u> GC	- ATAG - - - - CGTATTA - 3'
17	2	5' CCACCA <u>TGCTTCTC</u> G	- ATAGTTA - - CGCAT ACCCCCGTAT - 3'
18	1	5' GCCAGA <u>TTCTG-CT</u> ACC	AATAGTGA - - GTGT CCGACIAT - 3'
19	1	5' ATAG <u>GCCATGCTT</u> GGCT	TATAGTG - - - - TA CCGCCCTTATCG - 3'
20	1	5' TCIGCTCTC <u>CTCTATTCT</u> AGCAGTCC	- ATAGT - - - CG - GT - 3'
21	1	5' TTGCCAGCATA <u>GTGGCA</u> GACGTGGTGT	- ATAG - G - CCCC - GT - 3'
22	1	5' TTCTAGCTCCGCCTGA <u>ACTTCT</u> GT	AATAGTGGAG - - GC - TTG A - 3'

3 / 9

Base-pair-forming regions (individually boxed sequences)

CONSERVED REGIONS

Base-pair-forming regions (individually boxed sequences)

FIGURE 3

4 / 9




FIGURE 4A




FIGURE 4B




FIGURE 5




FIGURE 6A

Mg²⁺-Dependent DNazyme






FIGURE 6B



10 mM Mg²⁺, pH 7.5, 37 °C, 2 h

FIGURE 7



turnover ≈ 0.6 hr⁻¹

turnover ≈ 1 hr⁻¹

10 mM Mg²⁺, pH 7.5, 37 °C

FIGURE 8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15580

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68; C12N 9/22; C07H 21/04
 US CL : 435/6, 91.31, 172.1; 536/23.1, 23.2, 24.5
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.31, 172.1; 536/23.1, 23.2, 24.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Dialog: CAS, BIOSIS, Derwent Biotechnology Abstracts, Medline; APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	Chemistry and Biology, Vol. 2, issued October 1995, Breaker et al., "A DNA enzyme with Mg ²⁺ -dependent RNA phosphoesterase activity", pages 655-660.	1-45

Further documents are listed in the continuation of Box C. See patent family annex.

<p>* Special categories of cited documents:</p> <p>*A* document defining the general state of the art which is not considered to be part of particular relevance</p> <p>*E* earlier document published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p>	<p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>* & * document member of the same patent family</p>
---	---

Date of the actual completion of the international search 25 MARCH 1996	Date of mailing of the international search report 09 APR 1996
--	--

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>John LeGUYADER</i> John LeGUYADER Telephone No. (703) 308-0196
---	---