

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 September 2011 (09.09.2011)

PCT

(10) International Publication Number
WO 2011/109322 A1

(51) International Patent Classification:

C11D 3/20 (2006.01) **C11D 3/00** (2006.01)
C11D 1/02 (2006.01)

(21) International Application Number:

PCT/US2011/026571

(22) International Filing Date:

1 March 2011 (01.03.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/310,538 4 March 2010 (04.03.2010) US

(71) Applicant (for all designated States except US): **THE PROCTER & GAMBLE COMPANY** [US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio 45202 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OH, Hiroshi** [US/US]; 8541 Meadow Bluff Court, Cincinnati, Ohio 45249 (US). **BATES, Tim** [US/US]; 1589 Forester Drive, Cincinnati, Ohio 45240 (US). **HECHT, Stacie, Ellen** [US/US]; 8238 Eagle Ridge Court, West Chester, Ohio 45069 (US).

(74) Common Representative: **THE PROCTER & GAMBLE COMPANY**; c/o Eileen L. Hughett, Global Patent

Services, 299 East Sixth Street, Sycamore Building, 4th Floor, Cincinnati, Ohio 45202 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))

(54) Title: DETERGENT COMPOSITION

(57) Abstract: Detergent compositions with improved sudsing properties comprising anionic surfactant and specific mono alcohol(s) in a specific ratio.



WO 2011/109322 A1

DETERGENT COMPOSITION

FIELD OF THE INVENTION

The present invention relates to detergent compositions with improved sudsing properties such as suds volume and suds mileage. The detergent compositions comprise anionic surfactant and specific mono alcohol(s) in a specific ratio.

BACKGROUND OF THE INVENTION

Detergent compositions must satisfy several criteria in order to be effective and fulfil the need of the consumer. The need for a composition to provide suds or not depends of its use. By way of example, in the case of automatic dish washing, it is desirable to have detergent composition that generates no or little suds to avoid the suds to overflow the dishwasher. On the contrary, in the case of detergent composition for washing fabrics by hand, the presence of suds in a cleaning operation may be desirable as it has long been used as a signal to the consumer that the detergent continues to be effective.

When the presence of suds is desirable, the quantity of suds generated, the mileage of the suds, and the aesthetics properties of the suds are not always optimised. There is as such a need to provide detergent composition having enhanced sudsing properties such as increased quantity and/or better mileage.

The inventors have found that, when used in the detergent compositions of the invention, a specific class of fatty alcohols was useful to fulfil at least partially one or more of the above mentioned needs.

This is particularly surprising as fatty alcohols are typically known for their anti-foaming properties and therefore are typically used in detergent compositions which application requires a low level of suds. For example, US3,629,122 is disclosing the use of fatty alcohols in low foaming compositions.

Unless otherwise specified, all percentage and ratio are in weight.

SUMMARY OF THE INVENTION

The present disclosure relates to a detergent composition comprising anionic surfactant and at least 0.1% by weight of mono alcohol(s) having a main chain comprising from 14 to 18

carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl, the weight ratio of said mono alcohol(s) to the total amount of anionic surfactant(s) comprised in the detergent composition being comprised between 0.01 and 0.07.

The invention also concerns the use of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl in a detergent composition to enhance the sudsing properties of said detergent composition.

DETAILED DESCRIPTION OF THE INVENTION

Mono-alcohol

The detergent composition of the invention comprises at least 0.1%, preferably from 0.15% to 3% or from 0.2% to 1%, typically from 0.25% to 0.8% or from 0.3% to 0.6% by weight of a mono alcohol or a mixture of mono alcohols having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl.

The mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl are preferably of formula:



with n being an integer comprised between 13 and 17, and with 2 or less, preferably 1 or less, typically none, of the CH₂ groups being substituted with an ethyl or methyl, preferably methyl, group.

The detergent composition of the invention may also comprise additional mono alcohol(s) having a main chain length of 13 carbon atoms or less or 19 carbon atoms or more and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl.

Preferably, the weight ratio of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl to the total amount of mono alcohol(s) having a main chain comprising from 9 to 20 carbon atoms and being linear or branched with 1 or 2

substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl comprised in the detergent composition is comprised between 0.7 and 1, in particular is comprised between 0.8 and 1 or between 0.9 and 1 or even 0.95 and 1.

Anionic surfactant

The detergent composition comprises an anionic surfactant or a plurality of anionic surfactants. The detergent composition may comprise from 0.5% to 50% by weight, preferably from 1% to 25% by weight, in particular from 3% to 20%, or from 5% to 15% by weight of anionic surfactant.

In particular, the anionic surfactant may comprise surfactants selected from alkyl ester sulfonate(s); linear, branched, and modified alkylbenzene sulfonate(s); C₁₀-C₁₈ alkyl alkoxy sulfates; C₁₀-C₂₀ primary, branched-chain and random alkyl sulfates; C₁₀-C₁₈ secondary (2,3) alkyl sulfates; C₁₀-C₁₈ alkyl alkoxy carboxylate(s); mid-chain branched alkyl sulfate(s); mid-chain branched alkyl alkoxy sulfate(s); alpha-olefin sulfonate(s); phosphate ester(s); and mixtures thereof.

Alkyl Ester Sulfonate Surfactant ("MES")

As used herein, "MES" refers to alkyl ester sulfonate surfactants, commonly used in methyl ester sulfonate form. MES surfactants useful herein include sulfonated fatty acid alkyl esters of the formula R-CH(SO₃⁻)-COOR', wherein R is, on the average, a C₆ to C₂₂ alkyl and R' is on the average a C₁ to C₈ alkyl.

The hydrophobic portion of these sulfonated alkyl esters have the sulfonate group at the α-position, i.e., the sulfonate group is positioned at the carbon atom adjacent to the carbonyl group. The alkyl portion of the hydrophobic portion, which corresponds to the R portion of the sulfonated fatty acid alkyl esters, is on the average a C₆ to C₂₂ alkyl. Preferably, the alkyl portion of this hydrophobic portion, R, has a straight-chain of an average length C₈ to C₁₆ hydrocarbon particularly when R' is methyl.

R', forming the ester portion of the sulfonated alkyl esters, is on the average a C₁ to C₈ alkyl. Preferably, R' is on the average a C₁ to C₆ alkyl, and most preferably a C₁ alkyl, i.e., methyl.

In one embodiment, the distribution is such that R is, on the average, a C₁₄ to C₁₆ alkyl (approximately, for example, a 95% C₁₄, 5% C₁₆ mixture) and R' is methyl. In another embodiment, the distribution is such that R is, on the average, a C₁₂ to C₁₆ alkyl (approximately, for example, a 3% C₁₂, 28% C₁₄, 69% C₁₆ mixture) and R' is methyl. In yet another embodiment, the distribution is such that R is, on the average, a C₁₀ to C₁₆ alkyl (approximately, for example, a 60% C₁₀, 35% C₁₂, 5% C₁₄ mixture) and R' is methyl. In yet another embodiment, the distribution is such that R is, on the average, a C₁₂ to C₁₄ alkyl (approximately, for example, a 65% C₁₂, 30% C₁₄ mixture). In yet a further embodiment, blends of the aforementioned distributions of R and R' may also be employed. In one embodiment, the methyl ester sulfonate has an average carbon length of about 16. In other embodiments, R' could be ethyl (C₂), n-propyl & i-propyl (C₃), n-butyl, i-butyl (C₄), n-pentyl (C₅) and n-hexyl (C₆).

Methods of making alkyl ester surfactants neutralized with an alkali metal or an alkaline earth metal have been well described and are known to those skilled in art the art. See, for example, U.S. Patent Nos.: 4,671,900; 4,816,188; 5,329,030; 5,382,677; 5,384,422; 5,475,134; 5,587,500; 6,780,830. MES as such is commercially available from Huish.


Linear, Branched, and Modified Alkylbenzene Sulfonate

Other suitable anionic surfactants useful herein include any of the conventional anionic surfactant types typically used in detergent products. These include alkyl benzene sulfonates as well as alkoxylated or non-alkoxylated alkyl sulfates.

Exemplary anionic surfactants are C₁₀₋₁₆ alkyl benzene sulfonates, preferably C₁₁₋₁₄ alkyl benzene sulfonates. In one embodiment, the alkyl group is linear and such linear alkyl benzene sulfonates are known as "LAS". Alkyl benzene sulfonates, and particularly LAS, are well known in the art. Such surfactants and their preparation are described for example in U.S. Patents 2,220,099 and 2,477,383. Preferred are the linear straight chain alkylbenzene sulfonates in which the average number of carbon atoms in the alkyl group is from about 11 to 14. Particularly, C₁₁-C₁₄, e.g., C₁₂, and LAS is a specific example of such surfactants.

Other exemplary alkylbenzene sulfonates include modified alkylbenzene sulfonate (MLAS) as discussed in WO 99/05243, WO 99/05242, WO 99/05244, WO 99/05082, WO 99/05084, WO 99/05241, WO 99/07656, WO 00/23549, and WO 00/23548.

MLAS may comprise a mixture, preferably consisting essentially of : (a) from about 15% to about 99%, preferably from about 15% to about 60%, more preferably from about 20% to about 40%, by weight of a mixture of branched alkylbenzene sulfonates having formula:



wherein L is an acyclic aliphatic moiety consisting of carbon and hydrogen, the L having two methyl termini and the L having no substituents other than A, R^1 and R^2 ; and wherein the mixture of branched alkylbenzene sulfonates contains two or more, preferably at least three, optionally more, of the branched alkylbenzene sulfonates differing in molecular weight of the anion of the formula (I), and wherein the mixture of branched alkylbenzene sulfonates has a sum of carbon atoms in R^1 , L and R^2 of from 9 to 15, preferably from 10 to 14; an average aliphatic carbon content, i.e., based on R^1 , L and R^2 and excluding A, of from about 10.0 to about 14.0, preferably from about 11.0 to about 13.0, more preferably from about 11.5 to about 12.5, carbon atoms; R^1 is C_1 - C_3 alkyl, preferably C_1 - C_2 alkyl, more preferably methyl; R^2 is selected from H and C_1 - C_3 alkyl, preferably H and C_1 - C_2 alkyl, more preferably H and methyl, more preferably H and methyl provided that in at least about 0.5, more preferably 0.7, more preferably 0.9 to 1.0 mole fraction of the branched alkylbenzene sulfonates, R^2 is H; A is a benzene moiety, typically A is the moiety- C_6H_4 -, with the SO_3^- moiety of Formula (I) in para-position to the L moiety, though in some proportion, usually no more than about 5%, preferably from 0 to 5% by weight, the SO_3^- moiety is ortho-to L; and (b) from about 1% to about 85%, preferably from about 40% to about 85%, more preferably from about 60% to about 80%, by weight of a mixture of nonbranched alkylbenzene sulfonates having formula:



wherein A is as defined hereinbefore and Y is an unsubstituted linear aliphatic moiety consisting of carbon and hydrogen having two methyl termini, and wherein the Y has a sum of carbon atoms of from 9 to 15, preferably from 10 to 14, and the Y has an average aliphatic carbon content of from about 10.0 to about 14.0, preferably from about 11.0 to about 13.0, more preferably 11.5 to 12.5 carbon atoms; and wherein the modified alkylbenzene sulfonate surfactant mixture is further characterized by a 2/3-phenyl index of from about 160 to about 275, preferably from about 170 to about 265, more preferably from about 180 to about 255; and also preferably wherein the modified alkylbenzene sulfonate surfactant mixture has a 2-methyl-2-phenyl index of less than about 0.3, preferably less than about 0.2, more preferably less than about 0.1, more preferably still, from 0 to 0.05.

C₁₀-C₁₈ Alkyl Alkoxy Sulfates

Another exemplary type of anionic surfactant includes ethoxylated alkyl sulfate surfactants. Such materials, also known as alkyl ether sulfates or alkyl polyethoxylate sulfates, are those which correspond to the formula:



wherein R' is a C₈-C₂₀ alkyl group and n is from about 1 to 20. In a specific embodiment, R' is C₁₀-C₁₈ alkyl and n is from about 0.1 to 15. In another embodiment, n is from about 1 to 15. In more specific embodiments, R' is a C₁₂-C₁₆, n is from about 1 to 6. In the disclosure herein, the designation "EOx" indicates that the alkoxy group is an ethoxy group, the integer "x" indicates the number of ethoxy groups in each chain.

The alkyl ether sulfates will generally be used in the form of mixtures comprising varying R' chain lengths and varying degrees of ethoxylation. Frequently, though the average n value may be more than zero, such mixtures will inevitably also contain some non-ethoxylated alkyl sulfate materials, i.e., individual surfactant molecules of the above ethoxylated alkyl sulfate formula wherein n=0 for that particular molecule.

C₁₀₋₂₀ Primary, Branched-Chain and Random Alkyl Sulfates

Alkyl sulfates may also be added separately to the compositions of this invention and used as or in any anionic surfactant component which may be present. Specific examples of alkyl sulfates surfactants are those produced by the sulfation of higher C₁₀-C₂₀ fatty alcohols. Conventional primary alkyl sulfate surfactants have the general formula:



wherein R is typically a linear C₁₀-C₂₀ alkyl group, which may be straight chain or branched chain. In specific embodiments, R is a C₁₀-C₁₅ alkyl, more specifically R is C₁₂-C₁₄.

C₁₀-C₁₈ Secondary (2,3) Alkyl Sulfates

Another anionic surfactant useful herein includes secondary (2,3) alkyl sulfates having formulae CH₃-(CH₂)_x-CH(OSO₃⁻)-CH₃ or CH₃-(CH₂)_y-CH(OSO₃⁻)-CH₂-CH₃.

Non-limiting examples of a preferred secondary alkyl sulfate include the one where x is at least about 7, preferably at least about 9, and y is an integer of at least 8, preferably at least about 9.

C₁₀-C₁₈ Alkyl Alkoxy Carboxylates

Another exemplary type of anionic surfactant includes ethoxylated alkyl carboxylate surfactants. Such materials, also known as alkyl ether carboxylates or alkyl polyethoxylate carboxylates, are those which correspond to the formula:




wherein R' is a C₈-C₂₀ alkyl group and n is an integer from about 1 to 20. In a specific embodiment, R' is C₁₀-C₁₈ alkyl and/or n is from about 1 to 15. In more specific embodiments, R' is a C₁₂-C₁₆ and/or n is from about 1 to 6.

The alkyl ether carboxylates will generally be used in the form of mixtures comprising varying R' chain lengths and varying degrees of ethoxylation. Frequently such mixtures will inevitably also contain some non-ethoxylated alkyl carboxylate materials, i.e., surfactants of the above alkyl ether carboxylate formula wherein n=0.

Mid-chain Branched Alkyl Sulfates

Exemplary anionic surfactants include mid-chain branched alkyl sulfates as discussed in U.S. Patents 6,020,303 and US 6,060,443.

Mid-chain branched alkyl sulfates may comprise at least about 0.5%, preferably at least about 5%, more preferably at least about 10%, most preferably at least about 20%, by weight of longer alkyl chain, mid-chain branched surfactant compounds of the formula:



wherein:

(a) A^b is a hydrophobic C₉ to C₂₂ (total carbons in the moiety), preferably from about C₁₂ to about C₁₈, mid-chain branched alkyl moiety having: (1) a longest linear carbon chain attached to the —X—B moiety in the range of from 8 to 21 carbon atoms; (2) one or more C₁₋₃ alkyl moieties branching from this longest linear carbon chain; (3) at least one of the branching alkyl moieties is attached directly to a carbon of the longest linear carbon chain at a position


within the range of position 2 carbon (counting from carbon #1 which is attached to the —X—B moiety) to position ω -2 carbon (the terminal carbon minus 2 carbons, i.e., the third carbon from the end of the longest linear carbon chain); and (4) the surfactant composition has an average total number of carbon atoms in the A^b —X moiety in the above formula within the range of greater than 14.5 to about 17.5 (preferably from about 15 to about 17);

(b) B is a hydrophilic moiety selected from sulfates; and

(c) X is selected from —CH₂— and —C(O)—.

Also preferred are mid-chain branched alkyl sulfates of the above formula wherein the A^b moiety does not have any quaternary substituted carbon atoms (i.e., 4 carbon atoms directly attached to one carbon atom).

Preferred mid-chain branched alkyl sulfates herein comprise longer alkyl chain, mid-chain branched surfactant compounds of the above formula wherein the A^b moiety is a branched primary alkyl moiety having the formula:



wherein the total number of carbon atoms in the branched primary alkyl moiety of this formula (including the R, R¹, and R² branching) is from 13 to 19; R, R¹, and R² are each independently selected from hydrogen and C₁-C₃ alkyl (preferably methyl), provided R, R¹, and R² are not all hydrogen and, when z is 0, at least R or R¹ is not hydrogen; w is from 0 to 13; x is from 0 to 13; y is from 0 to 13; z is from 0 to 13; and w+x+y+z is from 7 to 13.

Also preferred mid-chain branched alkyl sulfates comprise longer alkyl chain, mid-chain branched surfactant compounds of the above formula wherein the A^b moiety is a branched primary alkyl moiety having the formula selected from:




or mixtures thereof; wherein a, b, d, and e are integers, a+b is from 10 to 16, d+e is from 8 to 14 and wherein further when a+b=10, a is an integer from 2 to 9 and b is an integer from 1 to 8; when a+b=11, a is an integer from 2 to 10 and b is an integer from 1 to 9; when a+b=12, a is an integer from 2 to 11 and b is an integer from 1 to 10; when a+b=13, a is an integer from 2 to 12

and b is an integer from 1 to 11; when a+b=14, a is an integer from 2 to 13 and b is an integer from 1 to 12; when a+b=15, a is an integer from 2 to 14 and b is an integer from 1 to 13; when a+b=16, a is an integer from 2 to 15 and b is an integer from 1 to 14; when d+e=8, d is an integer from 2 to 7 and e is an integer from 1 to 6; when d+e=9, d is an integer from 2 to 8 and e is an integer from 1 to 7; when d+e=10, d is an integer from 2 to 9 and e is an integer from 1 to 8; when d+e=11, d is an integer from 2 to 10 and e is an integer from 1 to 9; when d+e=12, d is an integer from 2 to 11 and e is an integer from 1 to 10; when d+e=13, d is an integer from 2 to 12 and e is an integer from 1 to 11; when d+e=14, d is an integer from 2 to 13 and e is an integer from 1 to 12.

Mid-chain Branched Alkyl Alkoxy Sulfates

Still other exemplary anionic surfactants include mid-chain branched alkyl alkoxy sulfates as discussed in U.S. Patents 6,008,181 and US 6,020,303


Mid-chain branched alkyl alkoxy sulfates comprise from about 0.001% to about 100% of one or more (preferably a mixture of two or more) mid-chain branched primary alkyl alkoxy sulfates having the formula:



wherein the total number of carbon atoms in the branched primary alkyl moiety of this formula (including the R, R¹, and R² branching, but not including the carbon atoms in the EO/PO alkoxy moiety) is from 14 to 20, and wherein further for this surfactant mixture the average total number of carbon atoms in the branched primary alkyl moieties having the above formula is within the range of greater than 14.5 to about 17.5 (preferably from about 15 to about 17); R, R¹, and R² are each independently selected from hydrogen and C₁-C₃ alkyl (preferably methyl), provided R, R¹, and R² are not all hydrogen and, when z is 1, at least R or R¹ is not hydrogen; w is an integer from 0 to 13; x is an integer from 0 to 13; y is an integer from 0 to 13; z is an integer of at least 1; w+x+y+z is from 8 to 14; and EO/PO are alkoxy moieties including for example ethoxy, propoxy, butoxy, etc, preferably selected from ethoxy, propoxy, and mixed ethoxy/propoxy groups, most preferably ethoxy, wherein m is at least about 0.01, preferably within the range of from about 0.1 to about 30, more preferably from about 0.5 to about 10, and most preferably from about 1 to about 5. It is to be recognized that the (EO/PO)_m moiety may be either a distribution with average degree of alkoxylation corresponding to m, or it may be a

single specific chain with alkoxylation (e.g., ethoxylation and/or propoxylation) of exactly the number of units corresponding to m.


Preferably, the mid-chain branched alkyl alkoxy sulfates comprise a mixture of mid-chain branched primary alkyl alkoxyated sulfate surfactants, said mixture comprising at least about 5% by weight of two or more mid-chain branched primary alkyl alkoxyated sulfates having the formula:



or mixtures thereof; wherein a, b, d, and e are integers, a+b is from 10 to 16, d+e is from 8 to 14 and wherein further when a+b=10, a is an integer from 2 to 9 and b is an integer from 1 to 8; when a+b=11, a is an integer from 2 to 10 and b is an integer from 1 to 9; when a+b=12, a is an integer from 2 to 11 and b is an integer from 1 to 10; when a+b=13, a is an integer from 2 to 12 and b is an integer from 1 to 11; when a+b=14, a is an integer from 2 to 13 and b is an integer from 1 to 12; when a+b=15, a is an integer from 2 to 14 and b is an integer from 1 to 13; when a+b=16, a is an integer from 2 to 15 and b is an integer from 1 to 14; when d+e=8, d is an integer from 2 to 7 and e is an integer from 1 to 6; when d+e=9, d is an integer from 2 to 8 and e is an integer from 1 to 7; when d+e=10, d is an integer from 2 to 9 and e is an integer from 1 to 8; when d+e=11, d is an integer from 2 to 8 and e is an integer from 1 to 9; when d+e=12, d is an integer from 2 to 11 and e is an integer from 1 to 10; when d+e=13, d is an integer from 2 to 12 and e is an integer from 1 to 11; when d+e=14, d is an integer from 2 to 13 and e is an integer from 1 to 12; wherein for this surfactant mixture the average total number of carbon atoms in the branched primary alkyl moieties having the above formulas is within the range of greater than 14.5 to about 17.5; and wherein EO/PO are alkoxy moieties, preferably selected from ethoxy, propoxy, and mixed ethoxy/propoxy groups, wherein m is at least about 0.01, preferably within the range of from about 0.1 to about 30, more preferably from about 0.5 to about 10, and most preferably from about 1 to about 5.


The mid-chain branched alkyl alkoxy sulfates may comprise compounds of formula:





wherein: a is an integer from 2 to 11, b is an integer from 1 to 10, and a+b is 8 or 9; and EO/PO are alkoxy moieties, preferably selected from ethoxy, propoxy, and mixed ethoxy/propoxy groups, wherein m is at least about 0.01, preferably within the range of from about 0.1 to about 30, more preferably from about 0.5 to about 10, and most preferably from about 0.6 to about 5.


Also preferred herein are alkoxylated sulfate compounds of formula:



wherein: d and e are integers and d+e is 6 or 7; and wherein further when d+e=6, d is an integer from 2 to 5 and e is an integer from 1 to 4; when d+e=7, d is an integer from 2 to 6 and e is an integer from 1 to 5; and EO/PO are alkoxy moieties, preferably selected from ethoxy, propoxy, and mixed ethoxy/propoxy groups, wherein m is at least about 0.01, preferably within the range of from about 0.1 to about 30, more preferably from about 0.5 to about 10, and most preferably from about 0.6 to about 5.

Alpha-olefin Sulfonate


Other anionic surfactants useful in embodiments of the present disclosure include olefin sulfonates, which are compounds produced by the sulfonation of alpha-olefin by means of uncomplexed sulfur trioxide followed by neutralization of the acid reaction mixture under conditions such that sultones formed in the reaction are hydrolyzed to give corresponding hydroxyalkanesulfonates. The alpha-olefins from which the olefin sulfonates are derived are mono-olefins having from about 8 to about 24 carbon atoms, preferably from about 12 to about 16 carbon atoms. Preferably, they are straight chain olefins. Exemplary alpha-olefin sulfonates for use in the disclosure herein have the general formula:




where R is a linear or branched alkyl of about 8 to 20 carbon atoms. Examples of suitable alpha-olefins include 1-olefins such as 1-dodecene, 1-tetradecene, 1-hexadecene, 1-octadecene, 1-eicosene and 1-tetracosene.

Phosphate Esters

Other anionic surfactants useful in embodiments of the present disclosure include phosphate esters. Phosphate esters are any materials of the general formula:



wherein R and R' are C₆-C₂₀ alkyl or ethoxylated alkyl groups. Preferably R and R' are of the general formula:



wherein the alkyl substituent is C₁₀-C₁₆ and Y is between 0 and about 4. Most preferably the alkyl substituent of that formula is C₁₀-C₁₆ and Y is between about 2 and about 4. Such compounds may be prepared by known methods from phosphorus pentoxide, phosphoric acid, or phosphorus oxy halide and alcohols or ethoxylated alcohols.

It will be appreciated that the formula depicted represent mono- and di-esters, and commercial phosphate esters will generally comprise mixtures of the mono- and di-esters, together with some proportion of tri-ester.

Fatty acids

Preferably, the anionic surfactant comprises less than 3% or even less than 1% or 0.1% by weight of fatty acids. The detergent composition may comprise no fatty acids.

Fatty acids have the general formula:



wherein R is typically a C₉-C₂₁ alkyl group, which may be straight chain or branched chain. In specific embodiments, R is a C₉-C₁₇ alkyl, and more specifically R is C₁₁-C₁₅.

Exemplary fatty acids are selected from the group consisting of lauric acid, tridecylic acid, myristic acid, pentadecylic acid, palmitic acid, margaric acid, stearic acid, arachidic acid, phytanic acid, behenic acid, palmitoleic acid, oleic acid, elaidic acid, vaccenic acid, linoleic acid, cis-eleostearic acid, trans-eleosteric acid, linolenic acid, arachidonic acid and

combinations thereof. Fatty acids can be saturated or unsaturated. Unsaturated fatty acids typically having an iodine value from 15 to 25, preferably from 18 to 22 and a cis:trans isomer ratio from 1:1 to 200:1, preferably from 10:1 to 200:1.

Sources of fatty acid are coconut, soybean, tallow, palm, palm kernel, rapeseed, lard, sunflower, corn, safflower, canola, olive, peanut.

The weight ratio of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl to the total amount of anionic surfactant(s) comprised in the detergent composition is comprised between 0.01 and 0.07, preferably between 0.015 and 0.06, in particular between 0.02 and 0.05, for example above 0.03 and/or below 0.04.

Detergent composition

The detergent composition may be a laundry detergent composition or a dish washing detergent composition. Typically, the laundry detergent composition is formulated for use in an automatic washing machine or for hand-washing use, preferably for hand-washing use.

The detergent composition may be in solid form. The detergent composition is for example in particulate form, typically in free-flowing particulate form, although the composition may be in any liquid or solid form, in particular in the form of an agglomerate, granule, flake, extrudate, bar, tablet or any combination thereof. The composition can be made by methods such as dry-mixing, agglomerating, compaction, spray drying, pan-granulation, spheronization or any combination thereof. The solid composition typically has a bulk density of from 300 g/l to 1,500 g/l, typically from 500 g/l to 1,000 g/l.

The detergent composition may also be in the form of a liquid, gel, paste, dispersion, typically a colloidal dispersion or any combination thereof. Liquid compositions typically have a viscosity of from 500 mPa.s to 3,000 mPa.s, when measured at a shear rate of 20 s^{-1} at ambient conditions (20°C and 1 atmosphere), and typically have a density of from 800 g/l to 1300 g/l. If the composition is in the form of a dispersion, then it will typically have a volume average particle size of from 1 micrometer to 5,000 micrometers, typically from 1 micrometer to 50 micrometers. Typically, a Coulter Multisizer is used to measure the volume average particle size of a dispersion.

The detergent composition may be in unit dose form, including not only tablets, but also unit dose pouches wherein the composition is at least partially enclosed, typically completely enclosed, by a film such as a polyvinyl alcohol film.

The detergent composition may also be in the form of an insoluble substrate, for example a non-woven sheet, impregnated with detergent actives.

The detergent composition is preferably alkaline. A solution comprising 100 g of distilled water and 10 g of the detergent composition may have a pH above 8 or 9 or 10 at 50°C.

In addition to the anionic surfactant and the mono alcohol, the detergent composition may comprise one or more adjunct ingredient(s). The precise nature of these additional adjunct components, and levels of incorporation thereof, will depend on the physical form of the composition and the nature of the operation for which it is to be used. Suitable adjunct materials include, but are not limited to, builder, additional surfactants, flocculating aid, chelating agents, dye transfer inhibitors, enzymes, enzyme stabilizers, catalytic materials, bleach activators, hydrogen peroxide, sources of hydrogen peroxide, preformed peracids, polymeric dispersing agents, clay soil removal/anti-redeposition agents, brighteners, dyes, perfumes, structure elasticizing agents, fabric softeners, carriers, hydrotropes, processing aids, and/or pigments. In addition to the disclosure below, suitable examples of such other adjuncts and levels of use are found in U.S. Patent Nos. 5,576,282, 6,306,812 B1 and 6,326,348 B1 that are incorporated by reference. Such one or more adjuncts may be present as detailed below:

SURFACTANTS- the detergent composition may comprise one or more additional surfactant(s). The additional surfactant(s) may be selected from nonionic surfactants, cationic surfactants, ampholytic surfactants, zwitterionic surfactants, semi-polar nonionic surfactants and mixtures thereof.

The compositions of the invention may comprise non-ionic surfactant. Where present the non-ionic deterative surfactant(s) is generally present in amounts of from 0.5 to 20wt%, or from 2wt% to 4wt%.

The non-ionic deterative surfactant can be selected from the group consisting of: alkyl polyglucoside and/or an alkyl alkoxyated alcohol; C₁₂-C₁₈ alkyl ethoxylates, such as, NEODOL[®] non-ionic surfactants from Shell; C₆-C₁₂ alkyl phenol alkoxyates wherein the alkoxyate units are ethyleneoxy units, propyleneoxy units or a mixture thereof; C₁₂-C₁₈ alcohol and C₆-C₁₂ alkyl phenol condensates with ethylene oxide/propylene oxide block polymers such as Pluronic[®] from BASF; C₁₄-C₂₂ mid-chain branched alcohols, BA, as described in more detail

in US 6,150,322; C₁₄-C₂₂ mid-chain branched alkyl alkoxyates, BAEx, wherein x = from 1 to 35, as described in more detail in US 6,153,577, US 6,020,303 and US 6,093,856; alkylcelluloses as described in more detail in US 4,565,647, specifically alkylpolyglycosides as described in more detail in US 4,483,780 and US 4,483,779; polyhydroxy fatty acid amides as described in more detail in US 5,332,528, WO 92/06162, WO 93/19146, WO 93/19038, and WO 94/09099; ether capped poly(oxyalkylated) alcohol surfactants as described in more detail in US 6,482,994 and WO 01/42408; and mixtures thereof.

The composition may comprise a cationic deterative surfactant. When present, typically the composition comprises from 0.1wt% to 10 wt%, or from 1wt% to 2wt% cationic deterative surfactant.

Suitable cationic deterative surfactants are alkyl pyridinium compounds, alkyl quaternary ammonium compounds, alkyl quaternary phosphonium compounds, and alkyl ternary sulphonium compounds. The cationic deterative surfactant can be selected from the group consisting of: alkoxyate quaternary ammonium (AQA) surfactants as described in more detail in US 6,136,769; dimethyl hydroxyethyl quaternary ammonium surfactants as described in more detail in US 6,004,922; polyamine cationic surfactants as described in more detail in WO 98/35002, WO 98/35003, WO 98/35004, WO 98/35005, and WO 98/35006; cationic ester surfactants as described in more detail in US 4,228,042, US 4,239,660, US 4,260,529 and US 6,022,844; amino surfactants as described in more detail in US 6,221,825 and WO 00/47708, specifically amido propyldimethyl amine; and mixtures thereof.

Highly preferred cationic deterative surfactants are mono-C₈₋₁₀ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride, mono-C₁₀₋₁₂ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride and mono-C₁₀ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride. Cationic surfactants such as Praepagen HY (tradename Clariant) may be useful and may also be useful as a suds booster.

BUILDERS - The detergent composition may comprise one or more builders. When a builder is used, the subject composition will typically comprise from 1% to about 40%, typically from 2 to 25%, or even from about 5% to about 20%, or from 8 to 15% by weight of builder.

Builders include, but are not limited to, the alkali metal, ammonium and alkanolammonium salts of polyphosphates, alkali metal silicates, layered silicates, such as SKS-6 of Clariant®, alkaline earth and alkali metal carbonates, aluminosilicate builders, such as zeolite, and polycarboxylate compounds, ether hydroxypolycarboxylates, copolymers of maleic

anhydride with ethylene or vinyl methyl ether, 1, 3, 5-trihydroxy benzene-2, 4, 6-trisulphonic acid, and carboxymethyloxysuccinic acid, fatty acids, the various alkali metal, ammonium and substituted ammonium salts of polyacetic acids such as ethylenediamine tetraacetic acid and nitrilotriacetic acid, as well as polycarboxylates such as mellitic acid, succinic acid, citric acid, oxydisuccinic acid, polymaleic acid, benzene 1,3,5-tricarboxylic acid, carboxymethyloxysuccinic acid, and soluble salts thereof.

The detergent compositions of the present invention may comprise from 0 to 20%, in particular less than 15% or 10%, for example less than 5% of aluminosilicate builder(s).

The detergent composition of the present invention may comprise from 0 to 20%, in particular less than 15% or 10%, for example less than 5% of phosphate builder(s).

FLOCCULATING AID – The composition may further comprise a flocculating aid. Typically, the composition comprises at least 0.3% by weight of the composition of a flocculating aid. The composition may also be substantially free of flocculating aid. Typically, the flocculating aid is polymeric. Typically the flocculating aid is a polymer comprising monomer units selected from the group consisting of ethylene oxide, acrylamide, acrylic acid and mixtures thereof. Typically the flocculating aid is a polyethyleneoxide. Typically the flocculating aid has a molecular weight of at least 100,000 Da, in particular from 150,000 Da to 5,000,000 Da or even from 200,000 Da to 700,000 Da.

BLEACHING AGENT – The compositions of the present invention may comprise one or more bleaching agents. In general, when a bleaching agent is used, the compositions of the present invention may comprise from about 0.1% to about 50% or even from about 0.1% to about 25% bleaching agent by weight of the detergent composition. When present, suitable bleaching agents include bleaching catalysts, photobleaches for example Vitamin K3 and zinc or aluminium phthalocyanine sulfonate; bleach activators such as tetraacetyl ethylene diamine (TAED) and nonanoyloxybenzene sulphonate (NOBS); hydrogen peroxide; pre-formed peracids; sources of hydrogen peroxide such as inorganic perhydrate salts, including alkali metal salts such as sodium salts of perborate (usually mono- or tetra-hydrate), percarbonate, persulphate, perphosphate, persilicate salts and mixtures thereof, optionally coated, suitable coatings including inorganic salts such as alkali metal; and mixtures thereof.

The amounts of hydrogen peroxide source and peracid or bleach activator may be selected such that the molar ratio of available oxygen (from the peroxide source) to peracid is from 1:1 to 35:1, or even 2:1 to 10:1

FLUORESCENT WHITENING AGENT - The composition may contain components that may tint articles being cleaned, such as fluorescent whitening agent. When present, any fluorescent whitening agent suitable for use in a detergent composition may be used in the composition of the present invention. The most commonly used fluorescent whitening agents are those belonging to the classes of diaminostilbene-sulphonic acid derivatives, diarylpyrazoline derivatives and bisphenyl-distyryl derivatives.

Typical fluorescent whitening agents are Parawhite KX, supplied by Paramount Minerals and Chemicals, Mumbai, India; Tinopal[®] DMS and Tinopal[®] CBS available from Ciba-Geigy AG, Basel, Switzerland. Tinopal[®] DMS is the disodium salt of 4,4'-bis-(2-morpholino-4 anilino-s-triazin-6-ylamino) stilbene disulphonate. Tinopal[®] CBS is the disodium salt of 2,2'-bis-(phenyl-styryl) disulphonate.

FABRIC HUEING AGENTS- Fluorescent whitening agents emit at least some visible light. In contrast, fabric hueing agents alter the tint of a surface as they absorb at least a portion of the visible light spectrum. Suitable fabric hueing agents include dyes and dye-clay conjugates, and may also include pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Colour Index (C.I.) classifications of Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic Violet and Basic Red, or mixtures thereof.

POLYMERIC DISPERSING AGENTS – the compositions of the present invention can contain polymeric dispersing agents. These polymeric dispersing agents, if included, are typically at levels up to about 5%, typically from about 0.2% to about 2.5%, more typically from about 0.5% to about 1.5%. Suitable polymeric dispersing agents, include polymeric polycarboxylates, substituted (including quarternized and oxidized) polyamine polymers, and polyethylene glycols, such as: acrylic acid-based polymers having an average molecular of about 2,000 to about 10,000; acrylic/maleic-based copolymers having an average molecular weight of about 2,000 to about 100,000 and a ratio of acrylate to maleate segments of from about 30:1 to about 1:1; maleic/acrylic/vinyl alcohol terpolymers; polyethylene glycol (PEG) having a molecular weight of about 500 to about 100,000, typically from about 1,000 to about 50,000, more typically from about 1,500 to about 10,000; and water soluble or dispersible alkoxyated polyalkyleneamine materials.

POLYMERIC SOIL RELEASE AGENT - The compositions of the present invention can also contain polymeric soil release agent. polymeric soil release agent, or "SRA", have

hydrophilic segments to hydrophilize the surface of hydrophobic fibers such as polyester and nylon, and hydrophobic segments to deposit upon hydrophobic fibers and remain adhered thereto through completion of washing and rinsing cycles, thereby serving as an anchor for the hydrophilic segments. This can enable stains occurring subsequent to treatment with the SRA to be more easily cleaned in later washing procedures. Preferred SRA's include oligomeric terephthalate esters; sulfonated product of a substantially linear ester oligomer comprised of an oligomeric ester backbone of terephthaloyl and oxyalkyleneoxy repeat units and allyl-derived sulfonated terminal moieties covalently attached to the backbone; nonionic end-capped 1,2-propylene/polyoxyethylene terephthalate polyesters; an oligomer having empirical formula $(CAP)_2 (EG/PG)_5 (T)_5 (SIP)_1$ which comprises terephthaloyl (T), sulfoisophthaloyl (SIP), oxyethyleneoxy and oxy-1,2-propylene (EG/PG) units and which is typically terminated with end-caps (CAP), typically modified isethionates, as in an oligomer comprising one sulfoisophthaloyl unit, 5 terephthaloyl units, oxyethyleneoxy and oxy-1,2-propyleneoxy units in a defined ratio, typically about 0.5:1 to about 10:1, and two-end-cap units derived from sodium 2-(2-hydroxyethoxy)-ethanesulfonate; oligomeric esters comprising: (1) a backbone comprising (a) at least one unit selected from the group consisting of dihydroxy sulfonates, polyhydroxy sulfonates, a unit which is at least trifunctional whereby ester linkages are formed resulting in a branched oligomer backbone, and combinations thereof; (b) at least one unit which is a terephthaloyl moiety; and (c) at least one unsulfonated unit which is a 1,2-oxyalkyleneoxy moiety; and (2) one or more capping units selected from nonionic capping units, anionic capping units such as alkoxylated, typically ethoxylated, isethionates, alkoxylated propanesulfonates, alkoxylated propanedisulfonates, alkoxylated phenolsulfonates, sulfoaroyl derivatives and mixtures thereof. Preferred are esters of the empirical formula:



wherein CAP, EG/PG, PEG, T and SIP are as defined hereinabove, DEG represents di(oxyethylene)oxy units, SEG represents units derived from the sulfoethyl ether of glycerin and related moiety units, B represents branching units which are at least trifunctional whereby ester linkages are formed resulting in a branched oligomer backbone, a is from about 1 to about 12, b is from about 0.5 to about 25, c is from 0 to about 12, d is from 0 to about 10, b+c+d totals from about 0.5 to about 25, e is from about 1.5 to about 25, f is from 0 to about 12; e+f totals from about 1.5 to about 25, g is from about 0.05 to about 12; h is from about 0.01 to about 10, and a, b, c, d, e, f, g, and h represent the average number of moles of the corresponding units per mole

of the ester; and the ester has a molecular weight ranging from about 500 to about 5,000.; and; cellulosic derivatives such as the hydroxyether cellulosic polymers available as METHOCEL[®] from Dow; the C₁ -C₄ alkyl celluloses and C₄ hydroxyalkyl celluloses, see U.S. Pat. No. 4,000,093, issued Dec. 28, 1976 to Nicol et al., and the methyl cellulose ethers having an average degree of substitution (methyl) per anhydroglucose unit from about 1.6 to about 2.3 and a solution viscosity of from about 80 to about 120 centipoise measured at 20° C. as a 2% aqueous solution. Such materials are available as METOLOSE SM100[®] and METOLOSE SM200[®], which are the trade names of methyl cellulose ethers manufactured by Shinetsu Kagaku Kogyo KK.

ENZYME - The composition of the invention may further comprise an enzyme. When present in the detergent composition, the enzymes may be present at levels from about 0.00001% to about 2%, from about 0.0001% to about 1% or even from about 0.001% to about 0.5% or 0.02% enzyme protein by weight of the composition.

Examples of suitable enzymes include, but are not limited to, hemicellulases, peroxidases, proteases, cellulases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, mannanases, pectate lyases, keratinases, reductases, oxidases, phenoloxidases, lipoxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase, and amylases, or mixtures thereof.

ENZYME STABILIZERS - Enzymes for use in detergents can be stabilized by various techniques. The enzymes employed herein can be stabilized by the presence of water-soluble sources of calcium and/or magnesium ions in the finished compositions that provide such ions to the enzymes. In case of aqueous compositions comprising protease, a reversible protease inhibitor, such as a boron compound, can be added to further improve stability.

CATALYTIC METAL COMPLEXES – The compositions of the invention may comprise catalytic metal complexes. When present, one type of metal-containing bleach catalyst is a catalyst system comprising a transition metal cation of defined bleach catalytic activity, such as copper, iron, titanium, ruthenium, tungsten, molybdenum, or manganese cations, an auxiliary metal cation having little or no bleach catalytic activity, such as zinc or aluminum cations, and a sequester having defined stability constants for the catalytic and auxiliary metal cations, particularly ethylenediaminetetraacetic acid, ethylenediaminetetra(methylenephosphonic acid) and water-soluble salts thereof. Such catalysts are disclosed in U.S. 4,430,243.

If desired, the compositions herein can be catalyzed by means of a manganese compound. Such compounds and levels of use are well known in the art and include, for example, the manganese-based catalysts disclosed in U.S. 5,576,282.

Cobalt bleach catalysts useful herein are known, and are described, for example, in U.S. 5,597,936; U.S. 5,595,967. Such cobalt catalysts are readily prepared by known procedures, such as taught for example in U.S. 5,597,936, and U.S. 5,595,967.

Compositions herein may also suitably include a transition metal complex of ligands such as bispidones (WO 05/042532 A1) and/or macropolycyclic rigid ligands - abbreviated as "MRLs". As a practical matter, and not by way of limitation, the compositions and processes herein can be adjusted to provide on the order of at least one part per hundred million of the active MRL species in the aqueous washing medium, and will typically provide from about 0.005 ppm to about 25 ppm, from about 0.05 ppm to about 10 ppm, or even from about 0.1 ppm to about 5 ppm, of the MRL in the wash liquor.

Suitable transition-metals in the instant transition-metal bleach catalyst include, for example, manganese, iron and chromium. Suitable MRLs include 5,12-diethyl-1,5,8,12-tetraazabicyclo[6.6.2]hexadecane.

Suitable transition metal MRLs are readily prepared by known procedures, such as taught for example in WO 00/32601, and U.S. 6,225,464.

SOFTENING SYSTEM – the compositions of the invention may comprise a softening agent such as clay for softening through the wash. The composition may additionally comprise a charged polymeric fabric-softening boosting component.

COLORANT – the compositions of the invention may comprise a colorant, typically a dye or a pigment. Particularly, preferred dyes are those which are destroyed by oxidation during a laundry wash cycle. To ensure that the dye does not decompose during storage it is preferable for the dye to be stable at temperatures up to 40°C. The stability of the dye in the composition can be increased by ensuring that the water content of the composition is as low as possible. If possible, the dyes or pigments should not bind to or react with textile fibres. If the colorant does react with textile fibres, the colour imparted to the textiles should be destroyed by reaction with the oxidants present in laundry wash liquor. This is to avoid coloration of the textiles, especially over several washes. Particularly, preferred dyes include but are not limited to Basacid® Green 970 from BASF and Monastral blue from Albion.

SUDS SUPPRESSOR – Preferably, the detergent composition of the invention contains less than 3%, preferably up to 1%, and most preferably less than 0.1% or less than 0.01% or 0.001% or even 0.0001% of suds suppressor selected from the group consisting of trimethyl-, diethyl-, dipropyl-, dibutyl-, methylethyl-, phenylmethyl polysiloxane, and mixtures thereof. Preferably, the composition of the invention contain less than 3%, preferably up to 1%, and most preferably less than 0.1% or less than 0.01% or 0.001% or even 0.0001% of suds suppressor.

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as “40 mm” is intended to mean “about 40 mm”.

The following examples are given by way of illustration only and therefore should not be construed to limit the scope of the invention.

EXAMPLES

Example 1: Fabric care compositions

Ingredient	Composition A	Composition B	Composition C
	Concentration (weight percentages)		
LAS	10.	8	12
other anionic surfactants	3	4	2
non ionic surfactants	0.2	0.1	0.2
cationic surfactants	0.2		0.3
Mixture comprising C14-C18 mono alcohol ¹	0.45	0.25	0.7
sodium carbonate	15	13	11
sodium sulphate	55	57	59
silicate	3	4	1.8
zeolite	2	4	
polymer polyacrylate	2	1.5	2.5
polymer CMC	0.5	1	0.5
enzyme	0.5	0.5	0.5
bleach photobleach bleach activator	3.5	2	5
water, perfume, brightener, miscellaneous	qsp	qsp	qsp

¹C₁₄-C₁₈ mono alcohol is selected from:

- CO1695(95% of C₁₆OH cetyl alcohol) from P&G Chemicals
- CO1898 (98% of C₁₈OH stearyl alcohol) from P&G Chemicals
- CO1618(99% of C₁₆OH/C₁₈OH) from P&G Chemicals
- NeodolTM 67 (99.5% of C₁₄-C₁₈OH) from Shell Company
- NeodolTM 45 (99.5% of C₁₄-C₁₅OH) from Shell Company

- LialTM 145 (99.9% of C₁₄-C₁₈OH) from Sasol Company

The ratio C14-C18 mono alcohol/anionic surfactant in the previous composition depends on the percentage of C14-C18 mono alcohol in the mixture chosen and has been calculated below:

	Composition A	Composition B	Composition C
CO1695	0.033	0.020	0.048
Lial TM 145	0.035	0.021	0.050

What is claimed is:

1. Detergent composition comprising anionic surfactant and at least 0.1% by weight of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl, the weight ratio of said mono alcohol(s) to the total amount of anionic surfactant(s) comprised in the detergent composition being comprised between 0.01 and 0.07.

2. Detergent composition according to claim 1, wherein the weight ratio of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl to the total amount of anionic surfactant(s) comprised in the detergent composition is comprised between 0.02 and 0.05.

3. Detergent composition according to any one of the preceding claims comprising from 1% to 25% by weight of anionic surfactant, in particular from 5% to 15% by weight of anionic surfactant.

4. Detergent composition according to any one of the preceding claims comprising from 0.2% to 0.8%, in particular from 0.3% to 0.6%, by weight of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl.

5. Detergent composition according to any one of the preceding claims, wherein the weight ratio of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl to the total amount of mono alcohol(s) having a main chain comprising from 9 to 20 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl comprised in the detergent composition is comprised between 0.7 and 1, in particular is comprised between 0.9 and 1.

6. Detergent composition according to any one of the preceding claims, wherein the detergent composition is in solid form, in particular in powder form.

7. Detergent composition according to any one of the preceding claims, wherein the detergent composition is a fabric care composition.

8. Detergent composition according to any one of the preceding claims, comprising less than 3% by weight of fatty acid(s).

9. Detergent composition according to any one of the preceding claims, wherein a solution comprising 100 g of distilled water and 10 g of said detergent composition has a pH above 8 at 50°C.

AMENDED CLAIMS
received by the International Bureau on 26 Jul 2011 (26.07.2011)

- 5 1. Detergent composition comprising anionic surfactant and from 0.2% to 1% by weight of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl, the weight ratio of said mono alcohol(s) to the total amount of anionic surfactant(s) comprised in the detergent composition being comprised between 0.01 and 0.07.
- 10 2. Detergent composition according to claim 1, wherein the weight ratio of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl to the total amount of anionic surfactant(s) comprised in the detergent composition is comprised between
- 15 0.02 and 0.05.
3. Detergent composition according to any one of the preceding claims comprising from 1% to 25% by weight of anionic surfactant, in particular from 5% to 15% by weight of anionic surfactant.
- 20 4. Detergent composition according to any one of the preceding claims comprising from 0.2% to 0.8%, in particular from 0.3% to 0.6%, by weight of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl.
- 25 5. Detergent composition according to any one of the preceding claims, wherein the weight ratio of mono alcohol(s) having a main chain comprising from 14 to 18 carbon atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl to the total amount of mono alcohol(s) having a main chain comprising from 9 to 20 carbon
- 30 atoms and being linear or branched with 1 or 2 substituent(s), the 1 or 2 substituent(s) being methyl and/or ethyl comprised in the detergent composition is comprised between 0.7 and 1, in particular is comprised between 0.9 and 1.
6. Detergent composition according to any one of the preceding claims, wherein the
- 35 detergent composition is in solid form, in particular in powder form.

7. Detergent composition according to any one of the preceding claims, wherein the detergent composition is a fabric care composition.

5 8. Detergent composition according to any one of the preceding claims, comprising less than 3% by weight of fatty acid(s).

 9. Detergent composition according to any one of the preceding claims, wherein a solution comprising 100 g of distilled water and 10 g of said detergent composition has a pH
10 above 8 at 50°C.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/026571

A. CLASSIFICATION OF SUBJECT MATTER

INV. C11D3/20 C11D1/02 C11D3/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2007/232516 A1 (ANTHONY OLIVIER [FR] ET AL) 4 October 2007 (2007-10-04) claims example 2 page 1, paragraph 1 - page 2, paragraph 28 page 3, paragraph 52 - paragraph 65 -----	1-9
X	US 2 956 025 A (LEW HENRY Y) 11 October 1960 (1960-10-11) examples 8,11 claims column 1, line 15 - column 2, line 29 column 2, line 55 - line 59 column 3, line 21 - column 6, line 3 ----- -/--	1-9



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 May 2011

Date of mailing of the international search report

27/05/2011

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Neys, Patricia

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/026571

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 326 493 A (TAKAHATA MASAHIRO [JP] ET AL) 5 July 1994 (1994-07-05) examples 1,2 claims 1,4 column 1, line 6 - line 10 -----	1,2,5-8
X	EP 1 967 576 A1 (GOJO IND INC [US]) 10 September 2008 (2008-09-10) example 14 page 4, paragraph 30 - page 5, paragraph 34 -----	1-3,5,8
X	EP 0 388 810 A2 (KA0 CORP [JP]) 26 September 1990 (1990-09-26) example 4 -----	1,3-5,8
X	GB 965 248 A (SHELL INT RESEARCH) 29 July 1964 (1964-07-29) the whole document -----	1-8
A	Sasol North America Inc.: "ALFOL Alcohol C10-C20+ Blends", , December 2001 (2001-12), XP002637204, Retrieved from the Internet: URL: http://www.sasoltechdata.com/tds/ALC1020.pdf [retrieved on 2011-05-16] Alfol 1214; page 2 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2011/026571

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2007232516 A1	04-10-2007	EP 1805292 A1 FR 2875811 A1 WO 2006035154 A1 JP 2008514771 T	11-07-2007 31-03-2006 06-04-2006 08-05-2008
US 2956025 A	11-10-1960	NONE	
US 5326493 A	05-07-1994	DE 4240895 A1 GB 2262535 A JP 5156298 A	22-07-1993 23-06-1993 22-06-1993
EP 1967576 A1	10-09-2008	CA 2621986 A1	23-08-2008
EP 0388810 A2	26-09-1990	DE 69027468 D1 DE 69027468 T2 HK 29897 A JP 1964805 C JP 2247299 A JP 6099708 B PH 26509 A	25-07-1996 12-12-1996 21-03-1997 25-08-1995 03-10-1990 07-12-1994 07-08-1992
GB 965248 A	29-07-1964	BE 599155 A DE 1150171 B NL 247518 A	12-06-1963