

(19) **DANMARK**

(10) **DK/EP 2350279 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 15/113 (2010.01)** **A 61 K 31/711 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-02-15**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-01-27**
- (86) Europæisk ansøgning nr.: **09748890.2**
- (86) Europæisk indleveringsdag: **2009-10-22**
- (87) Den europæiske ansøgnings publiceringsdag: **2011-08-03**
- (86) International ansøgning nr.: **US2009061570**
- (87) Internationalt publikationsnr.: **WO2010048352**
- (30) Prioritet: **2008-10-22 US 196995 P** **2008-11-11 US 198931 P**
2009-02-15 WO PCT/IL2009/000179
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **Quark Pharmaceuticals, Inc., 6501 Dumbarton Circle, Fremont, California 94555, USA**
- (72) Opfinder: **SPIVAK, Igor, Rabbi Y. Ha-Nassi Street 8/16, 35425 Haifa, Israel**
SLAGER, Netanja, 176/4 Ben Yehuda Street, 53471 Tel Aviv, Israel
FEINSTEIN, Elena, 12/29 Hahagana Street, 76214 Rehovot, Israel
ALPERT, EVGENIA, Apartment 14, 25 Kobobi Street, 96757 Jerusalem, Israel
METT, Igor, 9/1 Paldi Street, 76248 Rehovot, Israel
BAR-ILAN, Amir, 12 Shoshanat Hacarmel Street, 34321 Haifa, Israel
KALINSKI, Hagar, 1 Haim Landau Street, 75701 Rishon-le-Zion, Israel
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **FREMGANGSMÅDER TIL BEHANDLING AF ØJENSYGDOMME**
- (56) Fremdragne publikationer:
WO-A2-2004/078940
WO-A2-2004/103389
WO-A2-2006/023544
WO-A2-2008/050329
WO-A2-2009/044392
US-A1- 2006 172 965
MUCKE H A M: "New ocular therapeutics: A view from the patenting perspective" IDRUGS, CURRENT DRUGS LTD, GB, vol. 10, no. 1, 1 January 2007 (2007-01-01), pages 37-41, XP009095914 ISSN: 1369-7056
SINGH M ET AL: "CELL-SPECIFIC CASPASE EXPRESSION BY DIFFERENT NEURONAL PHENOTYPES IN TRANSIENT RETINAL ESCHEMIA" JOURNAL OF NEUROCHEMISTRY, WILEY INTERSCIENCE, NEW YORK, NY, US, vol. 77, no. 2, 1 January 2001 (2001-01-01), pages 466-475, XP008047223 ISSN: 0022-3042
Z Ahmed ET AL: "Ocular neuroprotection by siRNA targeting caspase-2", Cell Death and Disease, vol. 2, no. 6,

Fortsættes ...

DESCRIPTION

RELATED APPLICATION

[0001] This application claims priority of U.S. Provisional Patent Applications Nos. 61/196995 filed Oct. 22, 2008 and 61/198931 filed Nov. 11, 2008 and of International Patent Application No. PCT/IL2009/000179 filed Feb. 15, 2009.

REFERENCE TO SEQUENCE LISTING

[0002] The sequences described in the specification (SEQ ID NOS:1-33,596) are being submitted with this application via the USPTO electronic filing system (EFS) in a text file titled, "188-PCT2.ST25.txt" created on October 20, 2009, file size 5952 KB, and are hereby incorporated by reference herein in their entirety.

FIELD OF THE INVENTION

[0003] The present invention relates to a non-invasive method of treating an eye disorder in a subject in need thereof comprising the step of topically administering to the surface of the eye of the subject a pharmaceutical composition comprising a therapeutic oligonucleotide directed to a target gene associated with loss of the retinal ganglion cell in the retina of the subject. Furthermore, the invention relates to non-invasive method of promoting retinal ganglion cell survival in subject suffering from an ocular disease, disorder or injury.

BACKGROUND OF THE INVENTION

[0004] Delivery of nucleic acid compounds to retinal tissue and in particular to retinal ganglion cells presents a big drug delivery challenge. Hitherto, eye drops have been considered to be useful primarily in the treatment of anterior segment disorders since it is has been shown that nucleic acids do not pass the cornea and insufficient drug concentrations reach the posterior ocular tissue (reviewed in del Amo and Urtti, 2008. Drug Discov Today 13(3/4):135-143 ; Fattal and Bochot, 2006. Adv Drug Del Rev 56:1203-1223).

[0005] A retinal ganglion cell (RGC) is a type of neuron located near the inner surface of the retina of the eye. Retinal ganglion cells receive visual information from photoreceptors and collectively transmit visual information from the retina to several regions in the brain. Further, there remains need for non-invasive method for inhibiting loss of retinal ganglion cells in a subject in need thereof. Various ocular diseases and disorders are characterized by death of retinal ganglion cells (RGCs). Accordingly, there remains a need for a non-invasive method for inhibiting loss of retinal ganglion cells (RGCs) in subjects that are suffering from an ocular disease, an ocular disorder or an ocular injury or are at risk of developing an ocular disease, an ocular disorder, or an ocular injury characterized and/or mediated by degeneration or death of retinal ganglion cells (RGCs).

SUMMARY OF THE INVENTION

[0006] The present invention discloses non-invasive methods of treating ocular diseases, disorders and injuries that are associated with degeneration or death of retinal ganglion cells (RGCs) and compositions useful in the methods. The methods as disclosed comprise topically administering to the surface of the eye of a subject a therapeutic oligonucleotide composition useful in promoting survival of retinal ganglion cells in a subject. Hitherto, oligonucleotides have been delivered to retinal ocular tissue by systemic delivery or intravitreal injection, methods associated with detrimental side effects and poor patient compliance, respectively. The present invention provides topical ophthalmic compositions comprising an oligonucleotide, and their use in non-invasive methods for down regulating expression of a target gene associated with loss of the retinal ganglion cell in the retina of a subject, for rescuing retinal ganglion cells from apoptosis and for treating eye disorders.

[0007] The present invention is directed to an oligoribonucleotide, which is a double stranded RNA compound having the structure:

5' 'B - GCCAGAAUGUGGAACUCCU 3' (sense strand; SEQ ID NO: 9015)

3' CGGUCUUACACCUUGAGGA 5' (antisense strand; SEQ ID NO: 9516)

wherein each A, C, U, and G is a ribonucleotide and each consecutive ribonucleotide is joined to the next ribonucleotide by a phosphodiester bond;

wherein the sense strand comprises, counting from the 5' terminus, an unmodified ribonucleotide at positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 and 19, a L-deoxycytidine at position 18, and an inverted abasic 5' cap; and

wherein the antisense strand comprises, counting from the 5' terminus, a 2'-O-Me sugar modified ribonucleotide at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and an unmodified ribonucleotide at positions 1, 3, 5, 7, 9, 10, 12, 14, 16 and 18.

[0008] The present invention is also directed to use in therapy of the subject oligoribonucleotide.

[0009] In one aspect the present invention provides a subject oligoribonucleotide for use in a method of non-invasive delivery to a retinal tissue in a subject suffering from an eye disorder, disease or injury comprising topically applying an ophthalmic composition comprising the oligonucleotide to the surface of the eye of the subject.

[0010] In another aspect the present invention provides a subject oligoribonucleotide for use in a method of non-invasive delivery of an oligonucleotide to a retinal ganglion cell in a subject suffering from an eye disorder comprising topically applying an ophthalmic composition comprising the oligonucleotide to the surface of the eye of the subject.

[0011] In yet another aspect, the present invention provides a subject oligoribonucleotide for use in a method of attenuating expression of a target gene associated with loss of a retinal ganglion cell in the retina in a subject suffering from an ocular disease, disorder or injury, which comprises topically (non-invasively) administering to the surface of the eye of the subject a pharmaceutical composition comprising at least one oligonucleotide directed to the target mRNA product of the target gene, in an amount and over a period of time effective to attenuate expression of the gene in the retina of the subject.

[0012] In a further aspect, the present invention provides a subject oligoribonucleotide for use in a method of treating a subject suffering from retinal ganglion cell loss or retinal ganglion cell damage and providing ocular neuroprotection to a subject suffering from or at risk of developing an eye disease, disorder or injury. The method comprises topically administering to the surface of the eye of the subject an ophthalmic pharmaceutical composition comprising at least one oligonucleotide directed to a target gene in the retina of the subject, in an amount and over a period of time effective to inhibit retinal ganglion cell loss or retinal ganglion cell damage in the subject.

[0013] In various embodiments the ophthalmic pharmaceutical composition is formulated as a cream, a foam, a paste, an ointment, an emulsion, a liquid solution including an eye drop, a gel, spray, a suspension, a microemulsion, microspheres, microcapsules, nanospheres, nanoparticles, lipid vesicles, liposomes, polymeric vesicles, a patch, or a contact lens. In some embodiments the pharmaceutical composition is formulated as an eye drop. In preferred embodiments the ophthalmic composition is to be administered to the eye, for example, by instillation of an eye drop or by administration of a mist.

[0014] The target ocular mRNA is selected from an mRNA polynucleotide set forth in any one of SEQ ID NOS:1-2.

[0015] The target gene is selected from a gene transcribed into an mRNA polynucleotide set forth in any one of SEQ ID NOS:1-2.

[0016] In some embodiments the at least one oligonucleotide is selected from chemically modified siRNA, unmodified siRNA, antisense, ribozyme, miRNA and shRNA. In preferred embodiments the at least one oligonucleotide is a chemically modified siRNA. The siRNA sense and antisense oligonucleotides are selected from sense and corresponding antisense oligonucleotide pairs set forth in SEQ ID NOS: 9015 and 9516.

[0017] In some embodiments the eye disorder, disease or injury is selected from glaucoma, dry eye, diabetic retinopathy (DR), diabetic macular edema (DME) or age related macular degeneration (AMD). In other embodiments the ocular disorder, disease or injury is optic neuritis, central retinal vein occlusion, branch retinal vein occlusion (BRVO). In further embodiments the eye disorder, disease or injury is retinitis pigmentosa (RP), ischemic optic neuropathy or optic nerve injury. In further embodiments ocular disorder, disease or injury is retinopathy of prematurity (ROP) retinal ganglion degeneration, macular degeneration, hereditary optic neuropathy, metabolic optic neuropathy, optic neuropathy due to a toxic agent or neuropathy caused by adverse drug reactions or vitamin deficiency. In yet another embodiment the disorder is vision loss associated with a tumor.

[0018] In various embodiments the at least one siRNA compound is delivered to the eye of a subject as a liquid solution, including an eye drop. In various embodiments the present invention provides a subject oligoribonucleotide for use in a method for attenuating expression of a target ocular gene associated with loss of a retinal ganglion cell in a subject suffering from an ocular disease, disorder or injury, comprising topically applying to the surface of the eye of the subject an ophthalmic pharmaceutical composition formulated as an eye drop.

[0019] In various embodiments expression of a target ocular mRNA is attenuated in an ocular cells in the retina of a subject suffering from an eye disease, disorder, or injury. In various embodiments the ocular cell includes but is not limited to an acinar cell of the lacrimal gland, a ductal cell of the lacrimal gland, a retinal ganglion cell (RGC), a retinal pigment epithelial (RPE) cell, a choroidal cell, a corneal cell, a cell of the ciliary process or a cell of the trabecular meshwork or a combination thereof.

[0020] The present description discloses a method for inhibiting loss of a retinal ganglion cell in a subject, comprising non-invasively administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA which down regulates expression of a target gene associated with loss of the retinal ganglion cell in the retina of the subject, thereby inhibiting loss of a retinal ganglion cell. In certain embodiments two or more target genes are down regulated by the method according to the invention. In certain embodiments attenuating expression of at least one target gene (ocular mRNA) associated with loss of the retinal ganglion cell in the retina of the subject confers upon the eye neuroprotective properties. The at least one ocular target gene is selected from SEQ ID NO:1-2.

[0021] In one embodiment the eye disorder is glaucoma. Thus the present invention provides a subject oligoribonucleotide for use in a method of inhibiting loss of a retinal ganglion cell in a subject suffering from glaucoma comprising topically (non-invasively) applying to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA which down regulates a target gene (target ocular mRNA) associated with loss of the retinal ganglion cell in the retina of the subject, thereby inhibiting loss of a retinal ganglion cell in the subject's eye. In preferred embodiments the siRNA is chemically modified. In some embodiments the chemically modified siRNA attenuates expression of gene (target ocular mRNA) in the retina of the subject's eye. In other embodiments the chemically modified siRNA attenuates expression of target ocular mRNA in the optic nerve of the subject's eye. In certain embodiments attenuating expression of at least one target gene (ocular mRNA) associated with loss of the retinal ganglion cell in the retina of the subject is effective to treat glaucoma. The at least one target gene is CASP2 (SEQ ID NO:1-2). In some embodiments the siRNA sense and antisense strands is the pair of oligonucleotide sequences set forth in SEQ ID NOS:9015 and 9516.

[0022] In another embodiment the eye disorder is dry-eye. Thus the present description discloses a method of inhibiting loss of a retinal ganglion cell in a subject suffering from dry-eye. The method comprises topically (non-invasively) administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA which down regulates a target gene (target ocular mRNA) associated with loss of the retinal ganglion cell in the retina of the subject, thereby inhibiting loss of a retinal ganglion cell in the subject's eye. In preferred embodiments the siRNA is chemically modified. In certain embodiments attenuating expression of at least one target ocular mRNA is effective to treat dry-eye disorder. In certain embodiments attenuating expression of at least one target gene associated with loss of retinal ganglion cells is effective to reduce the symptoms of dry-eye. In certain embodiments the at least one target gene (ocular target mRNA) is expressed in retinal ganglion cell in the eye of the subject. In certain embodiments the at least one target gene (ocular target mRNA) is expressed in the lacrimal gland in the eye of the subject. In certain embodiments the at least one target gene is selected from Table A2 set forth in any one of SEQ ID NO: 5 (p53), SEQ ID NO:8 - 10 (LRDD), SEQ ID NO:26 - 27 (SHC1) and SEQ ID NO:30 - 44 (FAS and FAS ligand). In some embodiments the target gene is selected from FAS, FAS ligand (FASL), p53, LRDD, PARP1, AIF (apoptosis inducing factor), NOS1, NOS2A, XIAP and SHC1-SHC. In certain preferred embodiments the target gene is set forth in any one of SEQ ID NO:36 - SEQ ID NO: 43 or SEQ ID NO:44. In some embodiments the siRNA sense and antisense strands are selected from pairs of oligonucleotide sequences set forth in SEQ ID NOS: 13,225-15,224.

[0023] In another embodiment the eye disorder is AMD, DR or DME. Thus, the present invention provides a subject oligoribonucleotide for use in a method of inhibiting loss of a retinal ganglion cell in a subject suffering from AMD, DR or DME. The method comprises topically (non-invasively) administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA which down regulates a target gene (target ocular mRNA) associated with loss of the retinal ganglion cell in the retina of the subject, thereby inhibiting loss of a retinal ganglion cell in the subject's eye. In preferred embodiments the siRNA is chemically modified. In certain embodiments attenuating expression of at least one target gene associated with loss of retinal ganglion cells is effective to treat AMD, DR or DME. In certain embodiments attenuating expression of at least one target gene associated with loss of retinal ganglion cells is effective to reduce the symptoms of AMD, DR or DME. In certain embodiments, the administration of at least one siRNA attenuates expression of at least one target gene (ocular mRNA) in a retinal ganglion cell in the subject's eye. In certain embodiments the administration of at least one siRNA attenuates expression of at least one target gene (target ocular mRNA) in the choroid in the subject's eye. The target

gene is set forth in any one of SEQ ID NOS:1-2.

[0024] In a further embodiment the eye disorder is Retinitis pigmentosa (RP). Thus, the present description discloses a method of inhibiting loss of a retinal ganglion cell in a subject suffering from RP. The method comprises topically (non-invasively) administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA which down regulates a target gene (target ocular mRNA) associated with loss of the retinal ganglion cell in the retina of the subject, thereby inhibiting loss of a retinal ganglion cell in the subject's eye. In preferred embodiments the siRNA is chemically modified. In certain embodiments attenuating expression of at least one target gene associated with loss of retinal ganglion cells is effective to RP. In certain embodiments attenuating expression of at least one target gene associated with loss of retinal ganglion cells is effective to reduce the symptoms of RP. In certain embodiments the at least one target ocular mRNA is product of a gene selected from a gene listed in Table A4 which is transcribed into mRNA set forth in any one of SEQ ID NOS: 3, 14, 26-35, 54-57. In some embodiments the target ocular mRNA is product of a gene selected from the group consisting of CASP1, CASP3, CASP12, RTP801, RTP801L, CAPNS1, PARP1, AIF, NOS1, NOS2, XIAP and SHC1-SHC. In certain preferred embodiments the siRNA targets mRNA set forth in SEQ ID NOS:56-57.

[0025] In another aspect the description discloses a method of rescuing a retinal ganglion cell from apoptosis in a subject, comprising non-invasively applying to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby rescuing retinal ganglion cell from apoptosis in the subject. In preferred embodiments the at least one siRNA is a chemically modified. In some embodiments the target gene is set forth in any one of SEQ ID NO:1- 58.

[0026] In another aspect, the description discloses a method for promoting survival of a retinal ganglion cell in a subject displaying signs or symptoms of an ocular neuropathy, comprising non-invasively administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby promoting survival of a retinal ganglion cell in the subject. In preferred embodiments the at least one siRNA is a chemically modified. In certain embodiments the signs or symptoms of an ocular neuropathy are mediated by apoptosis. In some embodiments the target gene is set forth in any one of SEQ ID NO:1- 58. In various embodiments the siRNA sense and antisense oligonucleotides are selected from sense and corresponding antisense oligonucleotides set forth in SEQ ID NOS:59-33,596.

[0027] In yet another aspect, the invention is directed to a subject oligoribonucleotide for use in a method for preventing, treating or alleviating the effects of an ocular disease associated with death of a retinal ganglion cell in a subject, comprising non-invasively applying to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby preventing, treating or alleviating the effects of an ocular disease associated with death of a retinal ganglion cell in the subject. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1-2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotide pair set forth in SEQ ID NOS:9015 and 9516.

[0028] Another aspect of the invention provides for a subject oligoribonucleotide for use in a method for treating or preventing retinal ganglion cell death in a subject, comprising non-invasively applying to the surface of the eye of the subject an ophthalmic pharmaceutical composition which comprises: (a) a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, and (b) a pharmaceutically acceptable excipient or carrier or mixture thereof, and thereby treating or preventing retinal ganglion cell death in the subject. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1-2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotides set forth in SEQ ID NOS:9015 and 9516.

[0029] In another aspect, the present invention is directed to a subject oligoribonucleotide for use in a method preventing retinal ganglion cell death mediated by elevated intraocular pressure (IOP) in the eye of a subject, comprising non-invasively administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby preventing retinal ganglion cell death in the subject. In preferred embodiments the at least one siRNA is a chemically modified. In a particular embodiment according to this method the subject is afflicted with glaucoma. The target gene is set forth in any one of SEQ ID NO:1-2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotides set forth in SEQ ID NOS:9015 and 9516.

[0030] The present invention further provides a subject oligoribonucleotide for use in a method of delaying, preventing or rescuing a retinal cell from death in a subject suffering from elevated IOP comprising non-invasively applying to the surface of the

eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene associated with death of the RGC in the retina of the subject, thereby delaying, preventing or rescuing the retinal cell from injury or death and wherein intraocular pressure (IOP) remains substantially elevated. In preferred embodiments the at least one siRNA is a chemically modified. In a particular embodiment according to this method the subject is afflicted with glaucoma. The target gene is set forth in any one of SEQ ID NO:1-2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotides set forth in SEQ ID NOS:9015 and 9516.

[0031] The present invention further provides a subject oligonucleotide for use in a method of treating a subject suffering from retinal ganglion cell loss or retinal ganglion cell damage, comprising non-invasively administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby treating the subject or reducing retinal ganglion cell death in the subject. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1-2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotides set forth in SEQ ID NOS:9015 and 9516.

[0032] The present invention further provides a subject oligoribonucleotide for use in a method for lowering retinal ganglion cell loss and providing ocular neuroprotection to a subject in need thereof, comprising non-invasively applying to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby lowering retinal ganglion cell loss and providing ocular neuroprotection to the subject. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1-2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotide pairs set forth in SEQ ID NOS:9015 and 9516.

[0033] The present invention further provides a subject oligoribonucleotide for use in a method for preventing visual field loss associated with loss of retinal ganglion cells in a subject, comprising non-invasively administering to the surface of the eye of the subject an ophthalmic composition comprising a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, thereby preventing visual field loss in the subject. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1- 58. In various embodiments the siRNA sense and antisense oligonucleotides are selected from sense and corresponding antisense oligonucleotide pair set forth in SEQ ID NOS: 9015 and 9516.

[0034] In another aspect, the present invention provides a subject oligoribonucleotide for use in an ophthalmic composition for non-invasive treatment of an ocular disease associated with loss of a retinal ganglion cell in a subject, comprising: (a) a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, and (b) a pharmaceutically acceptable excipient or carrier or mixture thereof. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1- 58. In various embodiments the siRNA sense and antisense oligonucleotides sense and corresponding antisense oligonucleotide pair set forth in SEQ ID NOS:9015 and 9516.

[0035] In yet another aspect, the present invention provides a topical ophthalmic pharmaceutical composition for non-invasive treatment of an ocular disease associated with pathological abnormalities/changes in the tissues of the visual system, comprising: (a) a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, wherein the target gene is set forth in any one of SEQ ID NO:1-2, and (b) a pharmaceutically acceptable excipient or carrier or mixture thereof. In preferred embodiments the at least one siRNA is a chemically modified. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotide pair set forth in SEQ ID NOS:9015 and 9516.

[0036] According to another aspect, the present invention is directed to a topical ophthalmic pharmaceutical composition for use in treating a subject afflicted with ocular disease associated with death of retinal ganglion cells, which comprises: (a) a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject, wherein the target gene is set forth in any one of SEQ ID NO:1- 2, and (b) a pharmaceutically acceptable excipient or carrier or mixture thereof. In preferred embodiments the at least one siRNA is a chemically modified. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotide pair set forth in SEQ ID NOS:9015 and 9516.

[0037] In various embodiments the ocular disease is selected from a group comprising glaucoma, dry eye, diabetic retinopathy (DR), diabetic macular edema (DME), age related macular degeneration (AMD), optic neuritis, central retinal vein occlusion, branch retinal vein occlusion, ischemic optic neuropathy, optic nerve injury, retinopathy of prematurity (ROP) or retinitis pigmentosa (RP), retinal ganglion degeneration, macular degeneration, hereditary optic neuropathy, metabolic optic neuropathy, neuropathy due to a toxic agent or that caused by adverse drug reactions or vitamin deficiency; and the composition is formulated as a cream, a foam, a paste, an ointment, an emulsion, a liquid solution, an eye drop, a gel, spray, a suspension, a

microemulsion, microspheres, microcapsules, nanospheres, nanoparticles, lipid vesicles, liposomes, polymeric vesicles, a patch, a biological insert. In a preferred embodiment the composition is formulated as an eye drop.

[0038] In another aspect the invention is directed to a packaged pharmaceutical preparation, comprising: (a) a pharmaceutical composition according to the invention in a container; and (b) instructions for using the composition to treat an ocular disease. In various embodiments the pharmaceutical composition according to present invention comprises a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject suffering from an ocular disease. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1- 2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotides set forth in SEQ ID NOS:9015 and 9516. In one particular embodiment according to this aspect of the invention, the pharmaceutical composition is for non-invasive treatment of an ocular disease associated with loss of a retinal ganglion cell in a subject. In another particular embodiment according to this aspect the invention, the pharmaceutical composition is for non-invasive treatment of an ocular disease associated with pathological abnormalities/changes in the tissues of the visual system. In yet another particular embodiment according to this aspect the invention, the pharmaceutical composition is for use in treating a subject afflicted with ocular disease associated with death of retinal ganglion cells.

[0039] According to another aspect, the invention is directed to use of a pharmaceutical composition according to the invention in the manufacture of a medicament for promoting retinal ganglion cell survival in a subject. In various embodiments the pharmaceutical composition according to present invention comprises a therapeutically effective amount of at least one siRNA to a target gene in the retina of the subject suffering from an ocular disease. In preferred embodiments the at least one siRNA is a chemically modified. The target gene is set forth in any one of SEQ ID NO:1- 2. In various embodiments the siRNA sense and antisense oligonucleotides are sense and corresponding antisense oligonucleotide pair set forth in SEQ ID NOS:9015 and 9516. In one particular embodiment according to this aspect the invention the pharmaceutical composition is for non-invasive treatment of an ocular disease associated with loss of a retinal ganglion cell in a subject. In another particular embodiment according to this aspect the invention the pharmaceutical composition is for non-invasive treatment of an ocular disease associated with pathological abnormalities/changes in the tissues of the visual system. In yet another particular embodiment according to this aspect the invention the pharmaceutical composition is for use in treating a subject afflicted with ocular disease associated with death of retinal ganglion cells.

[0040] In some embodiments the composition includes a viscosity enhancing agent. A viscosity enhancing agent is selected from for example a hydrophilic polymer including cellulose and cellulose derivatives methylcellulose and methylcellulose derivatives. Such agents include methylcellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, carboxymethyl cellulose, derivatives thereof, combinations thereof and their salts. In some embodiments the viscosity enhancing agent is methylcellulose. In certain embodiments the viscosity enhancing agent is provided at a concentration of about 0.01% to about 4%, in other embodiments the viscosity enhancing agent is provided at a concentration of about 0.1% to about 3%, or at a concentration of about 0.5% to about 2%.

[0041] In some embodiments the composition includes an agent providing osmotic balance, or a surfactant. Such agents include glycerol, ethylene glycol, poly(ethylene glycol), propylene glycol, sorbitol, mannitol, monosaccharide, disaccharides and oligosaccharides. The present invention further provides a subject oligoribonucleotide for use in a method of treating glaucoma in a subject in need of treatment that comprises topically (non-invasively) administering to the surface of the eye of the subject a therapeutically effective amount of at least one siRNA to a target gene in the eye of the subject, in an amount effective to treat glaucoma. In certain preferred embodiments the siRNA chemically modified siRNA. In certain embodiments the at least one siRNA inhibits expression of at least one gene expressed in the retina of the subject's eye. In certain embodiments inhibition of at least one gene confers upon the eye neuroprotective properties. The at least one gene is is CASP2 (SEQ ID NO:1-2). In some embodiments the siRNA sense and antisense strands are the sequences set forth in SEQ ID NOS:9015 and 9516.

[0042] The present description further discloses a method of treating dry-eye in a subject in need thereof, which comprises topically (non-invasively) applying to the surface of the eye of the subject a therapeutically effective amount of at least one siRNA which inhibits expression of at least one gene expressed in the eye of the subject in an amount effective to reduce the symptoms of dry eye. In certain preferred embodiments the siRNA is chemically modified siRNA. In certain embodiments the at least one gene is expressed in the lacrimal gland in the subject. In certain embodiments the at least gene is selected from Table A2 set forth in any one of SEQ ID NO: 5, SEQ ID NO:8-10, SEQ ID NO:26 - 27 and SEQ ID NO:30 - 44. In some embodiments the gene is selected from FAS, FAS ligand (FASL), p53, LRDD, PARP1, AIF (apoptosis inducing factor), NOS1, NOS2A, XIAP and SHC1-SHC. In certain preferred embodiments the gene is transcribed into mRNA set forth in any one of SEQ ID NO:36 - SEQ ID NO:44. In some embodiments the siRNA sense and antisense strands are selected from any one of sequences in Tables 16-17 set forth in SEQ ID NOS: 13,225-15,224.

[0043] The present invention further provides a subject oligoribonucleotide for use in a method of treating AMD, DR or DME in a subject in need thereof, which comprises topically (non-invasively) administering to the surface of the eye of the subject a therapeutically effective amount of at least one siRNA which inhibits expression of at least one gene expressed in the eye in the subject in an amount effective to reduce the symptoms of AMD, DR or DME. In certain preferred embodiments the siRNA is chemically modified siRNA. In certain embodiments the at least one siRNA inhibits expression of at least one gene expressed in the choroid in the subject's eye. The at least one target ocular mRNA set forth in any one of SEQ ID NOS:1-2.

[0044] In a further embodiment the eye disorder is Retinitis pigmentosa (RP). Thus, the present description discloses a method of attenuating expression of a target ocular RNA in the eye of a subject suffering RP. The method comprises topically (non-invasively) applying to the surface of the eye the subject a therapeutically effective amount of at least one siRNA to a target gene in the eye of the subject. In certain preferred embodiments the siRNA is chemically modified siRNA. In certain embodiments attenuating expression of at least one target gene (target ocular mRNA) is effective to treat RP. In certain embodiments attenuation of at least one target ocular mRNA is effective to reduce the symptoms of RP. In certain embodiments the at least one target ocular mRNA is product of a gene selected from a gene listed in Table A4 which is transcribed into mRNA set forth in any one of SEQ ID NOS: 3, 14, 26-35, 54-57. In some embodiments the target ocular mRNA is product of a gene selected from the group consisting of CASP1, CASP3, CASP12, RTP801, RTP801L, CAPNS1, PARP1, AIF, NOS1, NOS2, XIAP and SHC1-SHC. In certain preferred embodiments the siRNA targets mRNA set forth in SEQ ID NOS:56-57.

[0045] In various embodiments the at least one siRNA is chemically modified. In various embodiments the at least one siRNA comprises a sufficient number of consecutive nucleotides having a sequence of sufficient homology to a nucleic acid sequence present within the target mRNA to hybridize to the mRNA and attenuate expression of the mRNA in the eye of the subject.

[0046] In various embodiments the at least one siRNA comprises a sufficient number of consecutive nucleotides having a sequence of sufficient homology to a nucleic acid sequence present within the gene to hybridize to the gene and reduce or inhibit expression of the gene in the eye of the subject.

BRIEF DESCRIPTION OF THE FIGURES

[0047]

Figures 1A-1B (for illustrative purposes): Representative images of Cy3 labeled DDIT4 siRNA incorporated into murine retina following eye drop administration.

Figures 2A-2C (for illustrative purposes): Representative images of Cy3 labeled DDIT4 siRNA incorporated into ductal and acinar cells of the murine lacrimal gland following eye drop administration.

Figures 3A-3C (for illustrative purposes): Time course of accumulation of Cy3-siRNA in rat choroid after administration of eye drops, following 1 and 4 hours post administration.

Figure 4 (for illustrative purposes): Cy3-siRNA delivery to the trabecular meshwork and ciliary body one hour after eye drop administration to the eye.

Figures 5A-5B (for illustrative purposes): Confocal microscopy (magnification x60) of the retina 1 hour after eye drop administration of QM5 - siRNA targeted at the p53 gene. SiRNA in retina (retinal pigment epithelial cells, retinal ganglion cells) is shown by Cy3 fluorescence.

Figures 6A-6B (for illustrative purposes) are representative images of Cy3 labeled DDIT4 siRNA incorporated into mice retina. Figure 6B shows accumulation of DDIT4_1 Cy3-siRNA 1 hour post administration by eye drops. Choroid, outer nuclear layer, RPE and outer segment layer of photoreceptor cells show Cy3 staining.

Figure 7 (for illustrative purposes) shows accumulation of DDIT4_1 Dy-649/C6 -siRNA 1 hour post administration by eye drops in RGC cells by use of Dy-649/C6 staining.

Figure 8A-8C represent control siRNA FITC-CNL_1RD/CNL_1FD and scrambled3'cy3-CNL_1 delivery to retinal tissues by different staining methods.

Figures 9A-9B show delivery of different structures of Casp2 to the mouse retinal tissues.

Figures 10A -10B (for illustrative purposes) show TGASEII-FAM and HNOEL-FAM delivery.

Figure 11 (for illustrative purposes) shows retinal delivery of siRNA against p53 in PBS as positive control group. Figures 12A-12B show that in intact animals or when administering ED without siRNA no fluorescent signal is obtained in the retina.

Figures 13A-13D (for illustrative purposes) provide the results of an experiment showing knockdown of p53 in the retina. Figures 13A-13B show standard curves for measuring p53 protein levels. Figures 13C-13D show retinal p53 knockdown by eye drops.

DETAILED DESCRIPTION OF THE INVENTION

[0048] The present invention provides topical oligonucleotide compositions and non-invasive methods of use thereof for treating various eye diseases and disorders. In particular, the invention provides their use in methods for treatment of various eye diseases and disorders associated with gene expression in the eye of a subject suffering from the disease or disorder.

[0049] The present invention is based in part on the unexpected discovery that topical non-invasive administration of siRNA compositions targets certain ocular tissues and cell types and is active in those tissues and cells when delivered topically to the surface of the eye. The discovery is surprising in view of the known obstacles to siRNA delivery and provides non-invasive methods as a realistic alternative to intravitreal or systemic delivery.

[0050] For siRNA molecules to be effective in silencing mRNA of a target gene, the siRNA requires three levels of targeting: to the target tissue, to the target cell type and to the target subcellular compartment. The present invention now discloses non-invasive methods of treating eye diseases and disorders.

[0051] The present invention relates in general to compounds which down-regulate expression of genes expressed in ocular cells, particularly to novel small interfering RNAs (siRNAs), and to the use of these novel siRNAs in the treatment of a subject suffering from medical conditions associated with expression of those genes in eye tissues and cells.

[0052] Methods for attenuating expression of an ocular target mRNA and methods of treating disorders in the eye are discussed herein at length, and any of said molecules and/or compositions may be beneficially employed in the treatment of a subject suffering from any of said conditions.

[0053] The siRNAs of the present invention possess structures and modifications which may increase activity, increase stability, and or minimize toxicity; the novel modifications of the siRNAs of the present invention can be beneficially applied to double stranded RNA sequences useful in preventing or attenuating target gene expression, in particular the target genes discussed herein.

[0054] Details of a non-limited example of target genes per indication are presented in Tables A1- A4, hereinbelow.

Table A1: Target genes for treatment of glaucoma

Gene	Full name and Human Gene ID/SEQ ID NO. for mRNA polynucleotide
CASP2	caspase 2, apoptosis-related cysteine peptidase
	gi 39995058 ref NM_032982.2 (SEQ ID NO:1)
	gi 39995060 ref NM_032983.2 (SEQ ID NO:2)
RTP801	Homo sapiens DNA-damage-inducible transcript 4 (DDIT4), mRNA
	gi 56676369 ref NM_019058.2 (SEQ ID NO:3)
ASPP1	protein phosphatase 1, regulatory (inhibitor) subunit 13B (PPP1R13B)
	gi 121114286 ref NM_015316.2 (SEQ ID NO:4)
p53	tumor protein p53 gi:120407067,NM_000546.2(SEQIDNO:5)
TP53BP2	tumor protein p53 binding protein, 2
	gi 112799848 ref NM_001031685.2 (SEQ ID NO:6)
	gi 12799845 ref NM_005426.2 (SEQ ID NO:7)
LRDD	leucine-rich repeats and death domain containing
	variant 2 gi 61742781 ref NM_018494.3 (SEQ ID NO:8)

Gene	Full name and Human Gene ID/SEQ ID NO. for mRNA polynucleotide
	variant 1 gi 61742783 ref NM_145886.2 (SEQ ID NO:9)
	variant 3 gi 61742785 ref NM_145887.2 (SEQ ID NO:10)
CYBA	cytochrome b-245, alpha polypeptide
	gi 68509913 ref NM_000101.2 (SEQ ID NO:11)
BNIP3	BCL2/adenovirus E1B 19kDa interacting protein 3
	gi 7669480 ref NM_004052.2 (SEQ ID NO:12)
RAC1	ras-related C3 botulinum toxin substrate 1 (rho family, small GTP binding protein) (gi 38505164 ref NM_198829.1) (SEQ ID NO:13)
RTP801L	Homo sapiens DNA-damage-inducible transcript 4-like (DDIT4L), mRNA gi 34222182 ref NM_145244.2 (SEQ ID NO:14)
SPP1	secreted phosphoprotein 1
	variant 1 gi 91206461 ref NM_001040058.1 (SEQ ID NO:15)
	variant 2 gi 38146097 ref NM_000582.2 (SEQ ID NO:16)
	variant 3 gi 91598938 ref NM_001040060.1 (SEQ ID NO:17)
SOX9	Homo sapiens SRY (sex determining region Y)-box 9 (campomelic dysplasia, autosomal sex-reversal) (SOX9), mRNA
	gi 37704387 ref NM_000346.2 (SEQ ID NO:18)
ACHE	Homo sapiens acetylcholinesterase (Yt blood group) (ACHE), variant E4-E6, mRNA gi 88999567 ref NM_000665.3 (SEQ ID NO:19)
	variant E4-E5, mRNA gi 88999566 ref NM_015831.2 (SEQ ID NO:20)
ADRB1	Homo sapiens adrenergic, beta-1-, receptor (ADRB1), mRNA gi 110349783 ref NM_000684.2 (SEQ ID NO:21)
HTRA2	Htra serine peptidase 2
	v 1 gi:73747817, NM_013247 (SEQ ID NO:22)
	v 2 gi:73747818, NM_145074 (SEQ ID NO:23)
KEP1	Kelch-like ECH-associated protein 1
	v 1 gi:45269144 NM_203500 (SEQ ID NO:24)
	v 2 gi:45269143 NM_012289 (SEQ ID NO:25)
SHC1-SHC	Src homology 2 domain containing transforming prot. 1
	v 1 gi: 194239661 NM_183001 (SEQ ID NO:26)
	v 2 gi: 194239660 NM_003029 (SEQ ID NO:27)
CAPNS1	Homo sapiens calpain, small subunit 1 (CAPNS1),
	variant 1, mRNA gi 51599152 ref NM_001749.2 (SEQ ID NO:28)
	variant 2, mRNA gi 51599150 ref NM_001003962.1 (SEQ ID NO:29)
PARP1	Homo sapiens poly (ADP-ribose) polymerase 1 (PARP1), mRNA
	gi 156523967 ref NM_001618.3 (SEQ ID NO:30)
AIF (AIFM1)	Homo sapiens apoptosis-inducing factor, mitochondrion-associated, 1
	variant 4, mRNA gi 195927003 ref NM_001130846.1 (SEQ ID NO:31)
	variant 5, mRNA gi 195927005 ref NM_001130847.1 (SEQ ID NO:32)
NOS1	Homo sapiens nitric oxide synthase 1 (neuronal) (NOS1), mRNA
	gi 194239671 ref NM_000620.2 (SEQ ID NO:33)
NOS2A	Homo sapiens nitric oxide synthase 2, inducible (NOS2), mRNA
	gi 206597519 ref NM_000625.4 (SEQ ID NO:34)
XIAP	Homo sapiens X-linked inhibitor of apoptosis (XIAP), mRNA
	gi 32528298 ref NM_001167.21 (SEQ ID NO:35)

Table A2: Target genes for treatment of dry eye

Gene	Full name and Human Gene ID
FAS	CD95, TNF receptor superfamily, member 6
	v 1. gi 23510419 ref NM_000043.3 (SEQ ID NO:36)
	v 3. gi 23510422 ref NM_152872.1 (SEQ ID NO:37)
	v 2. gi 23510420 ref NM_152871.1 (SEQ ID NO:38)
	v 4. gi 23510424 ref NM_152873.1 (SEQ ID NO:39)
	v 8. gi 23510426 ref NM_152874.1 (SEQ ID NO:40)
	v 5. gi 23510428 ref NM_152875.1 (SEQ ID NO:41)
	v 7. gi 23510433 ref NM_152877.1 (SEQ ID NO:42)
	v 6. gi 23510430 ref NM_152876.1 (SEQ ID NO:43)
FAS ligand	TNF superfamily, member 6 (FASL)
	gi 4557328 ref NM_000639.1 (SEQ ID NO:44)
p53	tumor protein p53
	gi8400737, NM_000546.2 (SEQ ID NO:5)
LRDD	leucine-rich repeats and death domain containing
	gi 61742781 ref NM_018494.3 (SEQ ID NO:8)
	gi 61742783 ref NM_145886.2 (SEQ ID NO:9)
	gi 61742785 ref NM_145887.2 (SEQ ID NO:10)
PART1	Homo sapiens poly (ADP-ribose) polymerase 1 (PARP1), mRNA
	gi 156523967 ref NM_001618.3 (SEQ ID NO:30)
AIF (AIFM1)	Homo sapiens apoptosis-inducing factor, mitochondrion-associated, 1
	variant 4, mRNA gi 195927003 ref NM_001130846.1 (SEQ ID NO:31)
	variant 5, mRNA gi 195927005 ref NM_001130847.1 (SEQ ID NO:32)
NOS1	Homo sapiens nitric oxide synthase 1 (neuronal) (NOS1), mRNA
	gi 194239671 ref NM_000620.2 (SEQ ID NO:33)
NOS2A	Homo sapiens nitric oxide synthase 2, inducible (NOS2), mRNA
	gi 206597519 ref NM_000625.4 (SEQ ID NO:34)
XIAP	Homo sapiens X-linked inhibitor of apoptosis (XIAP), mRNA
	gi 32528298 ref NM_001167.2 (SEQ ID NO:35)
SHC1-SHC	Src homology 2 domain containing transforming prot. 1
	v 1 gi: 194239661 NM_183001 (SEQ ID NO:26)
	v 2 gi: 194239660 NM_003029 (SEQ ID NO:27)

Table A3: Target genes for treatment of DR, DME, AMD

Gene	Full name and Human Gene ID
	lectin galactoside-binding soluble 3
LGALS3	v1 gi:115430222 NM_002306 (SEQ ID NO:45)
	v2 gi:115430224 NR_003225 (SEQ ID NO:46)
SLC2A1	solute carrier family 2 (facilitated glucose transporter), member 1
	gi 166795298 ref NM_006516.1 (SEQ ID NO:49)
SLC2A2	solute carrier family 2 (facilitated glucose transporter), member 2
	gi 4557850 ref NM_000340.1 (SEQ ID NO:50)
SLC2A3	Homo sapiens solute carrier family 2 (facilitated glucose transporter), member 3 (GLUT3)
	gi 221136810 ref NM_006931.1 (SEQ ID NO:51)
SLC5A1	solute carrier family 5 (sodium/glucose cotransporter), member 1

Gene	Full name and Human Gene ID
	gi 208973247 ref NM_000343.2 (SEQ ID NO:52)
SORD	sorbitol dehydrogenase mRNA
	gi 156627570 ref NM_003104.3 (SEQ ID NO:53)
CTSD	Homo sapiens cathepsin D (CTSD)
	gi 23110949 ref NM_001909.3 (SEQ ID NO:47)
CASP2	caspase 2, apoptosis-related cysteine peptidase
	gi 39995058 ref NM_032982.2 (SEQ ID NO:1)
	gi 39995060 ref NM_032983.2 (SEQ ID NO:2)
RTP801	Homo sapiens DNA-damage-inducible transcript 4 (DDIT4), mRNA
	gi 56676369 ref NM_019058.2 (SEQ ID NO:3)
p53	tumor protein p53 gi8400737, NM_000546.2 (SEQ ID NO:5)
TP53BP2	tumor protein p53 binding protein, 2
	gi 112799848 ref NM_001031685.2 (SEQ ID NO:6)
	gi 112799845 ref NM_005426.2 (SEQ ID NO:7)
LRDD	leucine-rich repeats and death domain containing
	gi 61742781 ref NM_018494.3 (SEQ ID NO:8)
	gi 61742783 ref NM_145886.2 (SEQ ID NO:9)
	gi 61742785 ref NM_145887.2 (SEQ ID NO:10)
BNIP3	BCL2/adenovirus E1B 19kDa interacting protein 3
	gi 7669480 ref NM_004052.2 (SEQ ID NO:12)
RAC1	ras-related C3 botulinum toxin substrate 1 (rho family, small GTP binding protein) (gi 38505164 ref NM_198829.1) (SEQ ID NO:13)
AKR1B1	aldo-keto reductase family 1, member B1 (aldose reductase)
	gi 24497579 ref NM_001628.2 (SEQ ID NO:48)
KEAP1	Kelch-like ECH-associated protein 1
	v 1 gi:45269144 NM_203500 (SEQ ID NO:24)
	v 2 gi:45269143 NM_012289 (SEQ ID NO:25)
PARP1	Homo sapiens poly (ADP-ribose) polymerase 1 (PARP1), mRNA
	gi 156523967 ref NM_001618.3 (SEQ ID NO:30)
AIF	Homo sapiens apoptosis-inducing factor, mitochondrion-associated, 1
(AIFM1)	variant 4, mRNA gi 195927003 ref NM_001130846.1 (SEQ ID NO:31)
	variant 5, mRNA gi 195927005 ref NM_001130847.1 (SEQ ID NO:32)
NOS1	Homo sapiens nitric oxide synthase 1 (neuronal) (NOS1), mRNA
	gi 194239671 ref NM_000620.2 (SEQ ID NO:33)
NOS2A	Homo sapiens nitric oxide synthase 2, inducible (NOS2), mRNA
	gi 206597519 ref NM_000625.4 (SEQ ID NO:34)
XIAP	Homo sapiens X-linked inhibitor of apoptosis (XIAP), mRNA
	gi 32528298 ref NM_001167.2 (SEQ ID NO:35)
SHC1-	Src homology 2 domain containing transforming prot. 1
SHC	v 1 gi: 194239661 NM_183001 (SEQ ID NO:26)
	v 2 gi: 194239660 NM_003029 (SEQ ID NO:27)

Table A4: Examples of target genes for treatment of retinitis pigmentosa (RP)

Gene	Full name and Human Gene ID

Gene	Full name and Human Gene ID
CASP1	Homo sapiens caspase 1, apoptosis-related cysteine peptidase (interleukin 1, beta, convertase) (CASP1), mRNA
	variant epsilon, gi 73622117 ref NM_033295.2 (SEQ ID NO:54)
	variant alpha, gi 73622114 ref NM_033292.2 (SEQ ID NO:55)
CASP3	Homo sapiens caspase 3, apoptosis-related cysteine peptidase (CASP3), mRNA
	variant alpha, gi 73622121 ref NM_004346.3 (SEQ ID NO:56)
	variant beta, gi 73622122 ref NM_032991.2 (SEQ ID NO:57)
CASP12	Homo sapiens caspase 12 variant zeta (CASP12) mRNA, complete sequence; alternatively spliced gi 20069120 gb AF486846.1 (SEQ ID NO:58)
RTP801	Homo sapiens DNA-damage-inducible transcript 4 (DDIT4), mRNA
	gi 56676369 ref NM_019058.2 (SEQ ID NO:3)
RTP801L	Homo sapiens DNA-damage-inducible transcript 4-like (DDIT4L), mRNA
	gi 34222182 ref NM_145244.2 (SEQ ID NO:14)
CAPNS1	Homo sapiens calpain, small subunit 1 (CAPNS1),
	variant 1, mRNA gi 51599152 ref NM_001749.2 (SEQ ID NO:28)
	variant 2, mRNA gi 51599150 ref NM_001003962.1 (SEQ ID NO:29)
PARP1	Homo sapiens poly (ADP-ribose) polymerase 1 (PARP1), mRNA
	gi 156523967 ref NM_001618.3 (SEQ ID NO:30)
AIF (AIFM1)	Homo sapiens apoptosis-inducing factor, mitochondrion-associated, 1
	variant 4, mRNA gi 195927003 ref NM_001130846.1 (SEQ ID NO:31)
	variant 5, mRNA gi 195927005 ref NM_001130847.1 (SEQ ID NO:32)
NOS1	Homo sapiens nitric oxide synthase 1 (neuronal) (NOS1), mRNA
	gi 194239671 ref NM_000620.2 (SEQ ID NO:33)
NOS2A	Homo sapiens nitric oxide synthase 2, inducible (NOS2), mRNA
	gi 206597519 ref NM_000625.4 (SEQ ID NO:34)
XIAP	Homo sapiens X-linked inhibitor of apoptosis (XIAP), mRNA
	gi 32528298 ref NM_001167.2 (SEQ ID NO:35)
SHC1-SHC	Src homology 2 domain containing transforming prot. 1
	v 1 gi: 194239661 NM_183001 (SEQ ID NO:26)
	v 2 gi: 194239660 NM_003029 (SEQ ID NO:27)

[0055] "Variant" or "v" refer to transcript variant.

[0056] Tables A1-A4 provide the gi (GeneInfo identifier) and accession numbers for an example of polynucleotide sequences of human mRNA. The oligonucleotide inhibitors of the present invention are directed to CASP2. Inhibition of the genes in Tables A1, A2, A3 and A4 is useful in treating, *inter alia*, glaucoma, dry eye, diabetic retinopathy (DR), diabetic macular edema (DME), age related macular degeneration (AMD) and retinitis pigmentosa (RP), respectively.

Definitions

[0057] For convenience certain terms employed in the specification, examples and claims are described herein.

[0058] It is to be noted that, as used herein, the singular forms "a", "an" and "the" include plural forms unless the content clearly dictates otherwise.

[0059] Where aspects or embodiments of the invention are described in terms of Markush groups or other grouping of

alternatives, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the group.

[0060] An "inhibitor" is a compound which is capable of reducing the expression of a gene or the activity of the product of such gene to an extent sufficient to achieve a desired biological or physiological effect. The term "inhibitor" as used herein refers to one or more of an oligonucleotide inhibitor, including siRNA, antisense, shRNA, miRNA and ribozymes. Inhibition may also be referred to as attenuation of expression of mRNA, down-regulation, or for RNA interference (RNAi), silencing. The inhibitors disclosed herein are chemically modified siRNA compounds which incorporate modifications such as changes to the sugar moiety and/or the base moiety and/or the linkages between nucleotides in the oligonucleotide structure.

[0061] The term "inhibit" or "attenuate" as used herein refers to reducing the expression of a gene, a variant or product thereof or the activity of the product of such gene to an extent sufficient to achieve a desired biological or physiological effect. Inhibition may be complete or partial. For example "inhibition" of CASP2 gene means inhibition of the gene expression (transcription or translation) or polypeptide activity of one or more of the variants disclosed in any Table A1 or A3 or an SNP (single nucleotide polymorphism) or other variants thereof.

[0062] "Ocular tissue" refers to any tissue associated with structure of the eye and is intended to include, in a non-limiting manner, the sclera, the cornea, the choroid, the retina, the lacrimal gland, and the optic nerve.

[0063] "Ocular cell" refers to any cell associated with structures of the eye and lacrimal apparatus and is intended to include, in a non-limiting manner, retinal ganglion cell, a retinal pigment epithelial cell, a corneal cell, conjunctiva, anterior chamber, iris, ciliary process, retina, choroid and choroidal cells, trabecular meshwork, and the like. For example the trabecular meshwork includes the inner uveal meshwork, the corneoscleral meshwork and the juxtacanalicular tissue. The lacrimal gland includes the lacrimal gland per se, the inferior and superior lacrimal puncta, the inferior and superior lacrimal canal, the lacrimal sac and the like.

[0064] As used herein, the terms "polynucleotide" and "nucleic acid" may be used interchangeably and refer to nucleotide sequences comprising deoxyribonucleic acid (DNA), and ribonucleic acid (RNA). The terms should also be understood to include, as equivalents, analogs of either RNA or DNA made from nucleotide analogs. Throughout this application mRNA sequences are set forth as representing the corresponding genes. The terms "mRNA polynucleotide sequence" and mRNA are used interchangeably.

[0065] "Oligonucleotide", "oligoribonucleotide" or "oligomer" refers to a deoxyribonucleotide or ribonucleotide sequence from about 2 to about 50 nucleotides. Each DNA or RNA nucleotide of the sequence may be independently natural or synthetic, and or modified or unmodified. Modifications include changes to the sugar moiety, the base moiety and or the linkages between nucleotides in the oligonucleotide. The compounds of the present invention encompass molecules comprising deoxyribonucleotides, ribonucleotides, modified deoxyribonucleotides, modified ribonucleotides and combinations thereof.

[0066] The present invention provides methods and compositions for inhibiting expression of a target gene *in vivo*. In general, the method includes topically administering oligoribonucleotides, in particular small interfering RNAs (i.e., siRNAs) or a nucleic acid material that can produce siRNA in a cell, to target an mRNA of the genes set forth in Tables A1-A4; in an amount sufficient to down-regulate expression of a target gene by an RNA interference mechanism. In particular, the method can be used to inhibit expression of the gene for treatment of a subject suffering from an eye disorder or disease related to expression of that gene in ocular tissue or cell. In accordance with the present invention, the siRNA molecules or inhibitors of the target gene are used as drugs to treat various ocular pathologies.

[0067] "Nucleotide" is meant to encompass deoxyribonucleotides and ribonucleotides, which may be natural or synthetic, and or modified or unmodified. Modifications include changes and substitutions to the sugar moiety, the base moiety and/or the internucleotide linkages.

[0068] As used herein, the terms "non-pairing nucleotide analog" means a nucleotide analog which comprises a non-base pairing moiety including but not limited to: 6 des amino adenosine (Nebularine), 4-Me-indole, 3-nitropyrrole, 5-nitroindole, Ds, Pa, N3-Me ribo U, N3-Me riboT, N3-Me dC, N3-Me-dT, N1-Me-dG, N1-Me-dA, N3-ethyl-dC, N3-Me dC. In some embodiments the non-base pairing nucleotide analog is a ribonucleotide. In other embodiments it is a deoxyribonucleotide.

[0069] All analogs of, or modifications to, a nucleotide / oligonucleotide may be employed with the present invention, provided that said analog or modification does not substantially adversely affect the stability and function of the nucleotide / oligonucleotide. Acceptable modifications include modifications of the sugar moiety, modifications of the base moiety, modifications in the internucleotide linkages and combinations thereof.

[0070] What is sometimes referred to in the present invention as an "abasic nucleotide" or "abasic nucleotide analog" is more properly referred to as a pseudo-nucleotide or an unconventional moiety. A nucleotide is a monomeric unit of nucleic acid, consisting of a ribose or deoxyribose sugar, a phosphate, and a base (adenine, guanine, thymine, or cytosine in DNA; adenine, guanine, uracil, or cytosine in RNA). A modified nucleotide comprises a modification in one or more of the sugar, phosphate and or base. The abasic pseudo-nucleotide lacks a base, and thus is not strictly a nucleotide.

[0071] The term "capping moiety" as used herein includes abasic ribose moiety, abasic deoxyribose moiety, modifications abasic ribose and abasic deoxyribose moieties including 2' O alkyl modifications; inverted abasic ribose and abasic deoxyribose moieties and modifications thereof; C6-imino-Pi; a mirror nucleotide including L-DNA and L-RNA; 5'OMe nucleotide; and nucleotide analogs including 4',5'-methylene nucleotide; 1-(β -D-erythrofuransyl)nucleotide; 4'-thio nucleotide, carbocyclic nucleotide; 5'-amino-alkyl phosphate; 1,3-diamino-2-propyl phosphate, 3-aminopropyl phosphate; 6-aminohexyl phosphate; 12-aminododecyl phosphate; hydroxypropyl phosphate; 1,5-anhydrohexitol nucleotide; alpha-nucleotide; threo-pentofuransyl nucleotide; acyclic 3',4'-seco nucleotide; 3,4-dihydroxybutyl nucleotide; 3,5-dihydroxypentyl nucleotide, 5'-5'-inverted abasic moiety; 1,4-butanediol phosphate; 5'-amino; and bridging or non bridging methylphosphonate and 5'-mercapto moieties. A 2'-O-methyl sugar modified ribonucleotide is also referred to as 2'-OMe sugar modified or 2'-OMe modified ribonucleotide

[0072] Certain preferred capping moieties are abasic ribose or abasic deoxyribose moieties; inverted abasic ribose or abasic deoxyribose moieties; C6-amino-Pi; a mirror nucleotide including L-DNA and L-RNA.

[0073] The term "unconventional moiety" as used herein refers to abasic ribose moiety, an abasic deoxyribose moiety, a deoxyribonucleotide, a modified deoxyribonucleotide, a mirror nucleotide, a non-base pairing nucleotide analog and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate bond; bridged nucleic acids including LNA and ethylene bridged nucleic acids.

[0074] Abasic deoxyribose moiety includes for example abasic deoxyribose-3'-phosphate; 1,2-dideoxy-D-ribofuranose-3-phosphate; 1,4-anhydro-2-deoxy-D-ribitol-3-phosphate. Inverted abasic deoxyribose moiety includes inverted deoxyriboabasic; 3',5' inverted deoxyabasic 5'-phosphate.

[0075] In the context of the present invention, a "mirror" nucleotide (also referred to as a spieglemer), is a nucleotide analog with reverse chirality to the naturally occurring or commonly employed nucleotide, i.e., a mirror image of the naturally occurring or commonly employed nucleotide. The mirror nucleotide is a ribonucleotide (L-RNA) or a deoxyribonucleotide (L-DNA) and may further comprise at least one sugar or base modification and/or a backbone modification, such as a phosphorothioate or phosphonate moiety. US Patent No. 6,602,858 discloses nucleic acid catalysts comprising at least one L-nucleotide substitution. Mirror nucleotide includes for example L-DNA (L-deoxyriboadenosine-3'-phosphate (mirror dA); L-deoxyribocytidine-3'-phosphate (mirror dC); L-deoxyriboguanosine-3'-phosphate (mirror dG); L-deoxyribothymidine-3'-phosphate (mirror dT) and L-RNA (L-riboadenosine-3'-phosphate (mirror rA); L-ribocytidine-3'-phosphate (mirror rC); L-riboguanosine-3'-phosphate (mirror rG); L-ribouracil-3'-phosphate (mirror dU)

[0076] Modified deoxyribonucleotide includes, for example 5'OMe DNA (5-methyl-deoxyriboguanosine-3'-phosphate) which may be useful as a nucleotide in the 5' terminal position (position number 1); PACE (deoxyriboadenine 3' phosphonoacetate, deoxyribocytidine 3' phosphonoacetate, deoxyriboguanosine 3' phosphonoacetate, deoxyribothymidine 3' phosphonoacetate.

[0077] Bridged nucleic acids include LNA (2'-O,4'-C-methylene bridged Nucleic Acid adenosine 3' monophosphate, 2'-O,4'-C-methylene bridged Nucleic Acid 5-methyl-cytidine 3' monophosphate, 2'-O,4'-C-methylene bridged Nucleic Acid guanosine 3' monophosphate, 5-methyl-uridine (or thymidine) 3' monophosphate); and ENA (2'-O,4'-C-ethylene bridged Nucleic Acid adenosine 3' monophosphate, 2'-O,4'-C-ethylene bridged Nucleic Acid 5-methyl-cytidine 3' monophosphate, 2'-O,4'-C-ethylene bridged Nucleic Acid guanosine 3' monophosphate, 5-methyl-uridine (or thymidine) 3' monophosphate).

[0078] In some embodiments of the present disclosure a preferred unconventional moiety is an abasic ribose moiety, an abasic deoxyribose moiety, a deoxyribonucleotide, a mirror nucleotide, and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate bond.

[0079] According to one aspect the present invention provides inhibitory oligonucleotide compounds comprising unmodified and modified nucleotides. The compound comprises at least one modified nucleotide selected from the group consisting of a sugar modification, a base modification and an internucleotide linkage modification and may contain DNA, and modified nucleotides such as LNA (locked nucleic acid) including ENA (ethylene-bridged nucleic acid; PNA (peptide nucleic acid); arabinoside; PACE

(phosphonoacetate and derivatives thereof), mirror nucleotide, or nucleotides with a 6 carbon sugar.

[0080] The compound comprises a 2' modification on the sugar moiety of at least one ribonucleotide ("2' sugar modification").

[0081] Other stabilizing modifications are also possible (e.g. modified nucleotides added to a 3' or 5' terminus of an oligomer). In some embodiments the backbone of the oligonucleotides is modified and comprises phosphate-D-ribose entities but may also contain thiophosphate-D-ribose entities, triester, thioate, 2'-5' bridged backbone (also may be referred to as 5'-2'), PACE modified internucleotide linkage or any other type of modification.

[0082] Other modifications include additions to the 5' and/or 3' termini of the oligonucleotides. Such terminal modifications may be lipids, peptides, sugars or other molecules.

[0083] Possible modifications to the sugar residue are manifold and include 2'-O alkyl, locked nucleic acid (LNA), glycol nucleic acid (GNA), threose nucleic acid (TNA), arabinoside; alritol (ANA) and other 6-membered sugars including morpholinos, and cyclohexinyls.

[0084] LNA compounds are disclosed in International Patent Publication Nos. WO 00/47599, WO 99/14226, and WO 98/39352. Examples of siRNA compounds comprising LNA nucleotides are disclosed in Elmen et al., (NAR 2005. 33(1):439-447) and in International Patent Publication No. WO 2004/083430. Six-membered ring nucleotide analogs are disclosed in Allart, et al (Nucleosides & Nucleotides, 1998, 17:1523-1526; and Perez-Perez, et al., 1996, Bioorg. and Medicinal Chem Letters 6:1457-1460) Oligonucleotides comprising 6-membered ring nucleotide analogs including hexitol and alritol nucleotide monomers are disclosed in International patent application publication No. WO 2006/047842.

[0085] Backbone modifications, such as ethyl (resulting in a phospho-ethyl triester); propyl (resulting in a phospho-propyl triester); and butyl (resulting in a phospho-butyl triester) are also possible. Other backbone modifications include polymer backbones, cyclic backbones, acyclic backbones, thiophosphate-D-ribose backbones, amidates, and phosphonoacetate derivatives. Certain structures include siRNA compounds having one or a plurality of 2'-5' internucleotide linkages (bridges or backbone).

[0086] The present invention also relates to compounds which down-regulate expression of various genes, particularly to novel small interfering RNA (siRNA) compounds, and to the use of these novel siRNAs in the treatment of various ocular diseases and medical conditions of the eye.

[0087] For each gene there is a separate list of 19-mer oligomer sequences, which are prioritized based on their score in the proprietary algorithm as the best sequences for targeting the human gene expression. 21- or 23-mer siRNA sequences can be generated by 5' and/or 3' extension of the 19-mer sequences disclosed herein. Such extension is preferably complementary to the corresponding mRNA sequence. Certain 23-mer oligomers were devised by this method where the order of the prioritization is the order of the corresponding 19-mer. Table B provides human sense and corresponding antisense oligonucleotides useful in preparing siRNA. The abbreviations for cross species are: Ms: Mouse, Rb: Rabbit, Chmp: chimpanzee, Mnk: Monkey, Chn: chinchilla, GP: guinea-pig.

Ocular Diseases

[0088] Topical delivery of therapeutic oligonucleotide compounds is useful in the treatment of a broad spectrum of eye diseases and disorders. Certain of the compounds of the invention are useful in treating patients suffering from diseases and disorders in which neuroprotection of the optic nerve would be of benefit, for example in:

1. 1. open angle primary/secondary glaucoma
2. 2. multiple sclerosis (optic neuritis)
3. 3. central or branch retinal vein occlusion
4. 4. ischemic optic neuropathy (in status epilepticus, HIV-1 infection)
5. 5. optic nerve injury
6. 6. tumors extending into the suprasellar region (above the sella turcica)
7. 7. juxta chiasmal tumors (the visual loss associated with compression of the optic chiasm by pituitary tumors may be transient or permanent, possibly related to the extent of irreversible retrograde degeneration to the retinal ganglion cells.
8. 8. Retinoblastoma

Primary Open-Angle Glaucoma

[0089] The majority of the cases of glaucoma are the form known as primary-open-angle glaucoma POAG, also called chronic open-angle glaucoma). POAG results from a build up of aqueous humor fluid within the anterior chamber of the eye resulting in intraocular pressure (IOP). Elevated IOP, which can be measured by a "tonometry" test, results from fluid entering the eye and not enough fluid exiting the eye. Normally, fluid enters the eye by seeping out of the blood vessels in the ciliary body. This fluid eventually makes its way past the crystalline lens, through the pupil (the central opening in the iris), and into the irido-corneal angle, the anatomical angle formed where the iris and the cornea come together. Then the fluid passes through the trabecular meshwork in the angle and leaves the eye via the canal of Schlemm.

[0090] If excess fluid enters the eye, or if the trabecular meshwork "drain" gets clogged up (for instance, with debris or cells) so that not enough fluid is leaving the eye, the pressure builds up in what is known as "open angle glaucoma." Open angle glaucoma also can be caused when the posterior portion of the iris adheres to the anterior surface of the lens creating a "pupillary block", and preventing intraocular fluid from passing through the pupil into the anterior chamber.

[0091] If the angle between the iris and the cornea is too narrow or is even closed, then the fluid backs up, causing increased pressure in what is known as "closed angle glaucoma."

[0092] Untreated glaucoma eventually leads to optic atrophy and blindness.

Normal Tension Glaucoma

[0093] Intraocular eye pressure is normal (between 12 - 22 mmHg) in about 25 - 30% glaucoma cases in the US, a condition known as normal-tension glaucoma. (In Japan, the rates may be as high as 70%.) Other factors are present that cause optic nerve damage but do not affect IOP.

Closed-Angle Glaucoma

[0094] Closed-angle glaucoma (also called angle-closure glaucoma) is responsible for 15% of all glaucoma cases. It is less common than POAG in the U.S., but it constitutes about half of the world's glaucoma cases because of its higher prevalence among Asians. The iris is pushed against the lens, sometimes sticking to it, closing off the drainage angle. This can occur very suddenly, resulting in an immediate rise in pressure. It often occurs in genetically susceptible people when the pupil shrinks suddenly. Closed-angle glaucoma can also be chronic and gradual, a less common condition.

Congenital Glaucoma

[0095] Congenital glaucoma, in which the eye's drainage canals fail to develop correctly, is present from birth. It is very rare, occurring in about 1 in 10,000 newborns. This may be an inherited condition and often can be corrected with microsurgery.

[0096] In one aspect the present description discloses a method of attenuating expression of a target ocular mRNA in the eye of a subject suffering from glaucoma, comprising topically (non-invasively) administering to the surface of the eye of the subject an effective amount of at least one chemically modified siRNA and a pharmaceutically acceptable carrier. The at least one ocular target mRNA is a product of CASP2. In a currently preferred embodiment the siRNA is formulated for delivery as eye drops. The target ocular mRNA set forth in SEQ ID NOS: 1-2.

[0097] In another aspect the present description discloses a method of treating glaucoma in a subject in need of treatment, comprising topically (non-invasively) administering to the surface of the eye of the subject a therapeutically effective amount of at least one chemically modified siRNA which inhibits expression of a target gene in the eye of the subject. The gene is CASP2. In a currently preferred embodiment the siRNA is formulated for delivery as eye drops. The target ocular mRNA set forth in SEQ ID NOS:1-2.

Optic neuritis

[0098] Optic neuritis is an inflammation of the optic nerve that may affect the part of the nerve and disc within the eyeball (papillitis) or the portion behind the eyeball (retrobulbar optic neuritis). Optic neuritis may be caused by any of the following: demyelinating diseases such as multiple sclerosis or post infectious encephalomyelitis; systemic viral or bacterial infections including complications of inflammatory diseases (e.g., sinusitis, meningitis, tuberculosis, syphilis, chorioretinitis, orbital inflammation); nutritional and metabolic diseases (e.g., diabetes, pernicious anemia, hyperthyroidism); Leber's Hereditary Optic Neuropathy, a rare form of inherited optic neuropathy which mainly affects young men; toxins (tobacco, methanol, quinine, arsenic, salicylates, lead); and trauma.

Optic atrophy.

[0099] Optic atrophy is a hereditary or acquired loss of vision disorder that results from the degeneration of the optic nerve and optic tract nerve fibers. It may be acquired via occlusions of the central retinal vein or artery, arteriosclerotic changes, may be secondary to degenerative retinal disease, may be a result of pressure against the optic nerve, or may be related to metabolic diseases (e.g., diabetes), trauma, glaucoma, or toxicity (to alcohol, tobacco, or other poisons). Degeneration and atrophy of optic nerve fibers is irreversible, although intravenous steroid injections have been seen to slow down the process in some cases.

Papilledema

[0100] A swelling of the optic disc (papilla), most commonly due to an increase in intracranial pressure (tumor induced), malignant hypertension, or thrombosis of the central retinal vein. The condition usually is bilateral, the nerve head is very elevated and swollen, and pupil response typically is normal. Vision is not affected initially and there is no pain upon eye movement. Secondary optic nerve atrophy and permanent vision loss can occur if the primary cause of the papilledema is left untreated.

Ischemic optic neuropathy (ION)

[0101] A severely blinding disease resulting from loss of the arterial blood supply to the optic nerve (usually in one eye), as a result of occlusive disorders of the nutrient arteries. Optic neuropathy can be anterior, which causes a pale edema of the optic disc, or posterior, in which the optic disc is not swollen and the abnormality occurs between the eyeball and the optic chiasm. Ischemic anterior optic neuropathy usually causes a loss of vision that may be sudden or occur over several days. Ischemic posterior optic neuropathy is uncommon, and the diagnosis depends largely upon exclusion of other causes, chiefly stroke and brain tumor.

[0102] Other diseases and conditions include dry eye, diabetic retinopathy, diabetic macular edema and retinitis pigmentosa.

Dry eye

[0103] Dry eye, also known as keratoconjunctivitis sicca or keratitis sicca, is a common problem usually resulting from a decrease in the production of tear film that lubricates the eyes. Most patients with dry eye experience discomfort, and no vision loss; although in severe cases, the cornea may become damaged or infected.

[0104] Dry eye is a multifactorial disease of the tears and ocular surface that results in symptoms of discomfort, visual disturbance, and tear film instability with potential damage to the ocular surface. It is accompanied by increased osmolarity of the tear film and inflammation of the ocular surface.

[0105] The lacrimal gland is a multilobular tissue composed of acinar, ductal, and myoepithelial cells. The acinar cells account for 80% of the cells present in the lacrimal gland and are the site for synthesis, storage, and secretion of proteins. Several of these proteins have antibacterial (lysozyme, lactoferrin) or growth factor (epidermal growth factor, transforming growth factor α , keratocyte growth factor) properties that are crucial to the health of the ocular surface. The primary function of the ductal cells is

to modify the primary fluid secreted by the acinar cells and to secrete water and electrolytes. The myoepithelial cells contain multiple processes, which surround the basal area of the acinar and ductal cells, and are believed to contract and force fluid out of the ducts and onto the ocular surface.

Mechanisms of Lacrimal Gland Dysfunction

[0106] Apoptosis, hormonal imbalance, production of autoantibodies, alterations in signaling molecules, neural dysfunction, and increased levels of proinflammatory cytokines have been proposed as possible mediators of lacrimal gland insufficiency. One of the primary symptoms of Sjögrens Syndrome is dry eye. Apoptosis of the acinar and ductal epithelial cells of the lacrimal glands has been proposed as a possible mechanism responsible for the impairment of secretory function (Manganelli and Fietta, *Semin Arthritis Rheum.* 2003. 33(1):49-65). Without wishing to be bound by theory, apoptotic epithelial cell death may be due to activation of several apoptotic pathways involving Fas (Apo-1/CD95), FasL (FasL/CD95L), Bax, caspases, perforin, and granzyme B. Cytotoxic T cells through the release of proteases, such as perforin and granzyme B, or the interaction of FasL expressed by T cells with Fas on epithelial cells, can lead to apoptosis of the acinar cells.

[0107] The current treatment for dry eye is mainly local and symptomatic such as: tear supplementation with lubricants; tear retention with therapies such as punctal occlusion, moisture chamber spectacles or contact lenses; tear stimulation for example by secretagogues; biological tear substitutes; anti-inflammatory therapy (Cyclosporine, Corticosteroids, Tetracyclines); and dietary essential fatty acids.

Diabetic Retinopathy

[0108] Diabetic retinopathy is a complication of diabetes and a leading cause of blindness. It occurs when diabetes damages the tiny blood vessels inside the retina. Diabetic retinopathy has four stages:

- Mild Nonproliferative Retinopathy: microaneurysms in the retina's blood vessels.
- Moderate Nonproliferative Retinopathy. As the disease progresses, some blood vessels that nourish the retina are blocked.
- Severe Nonproliferative Retinopathy. Many more blood vessels are blocked, depriving several areas of the retina of a blood supply, which is overcome by the growth of new blood vessels.
- Proliferative Retinopathy. The new blood vessels grow along the retina and along the surface of the vitreous gel. When the vessels leak blood, severe vision loss and even blindness can result.

[0109] During pregnancy, diabetic retinopathy may be a problem for women with diabetes.

[0110] In one aspect the present description discloses a method of attenuating expression of a target ocular mRNA in the eye of a subject suffering from diabetic retinopathy, comprising topically (non-invasively) administering to the surface of the eye of the subject an effective amount of at least one chemically modified siRNA and a pharmaceutically acceptable carrier. The target ocular mRNA is product of a human gene having mRNA set forth in any one of SEQ ID NOS:1-2. In a currently preferred embodiment the siRNA is formulated for delivery as eye drops.

[0111] In another aspect the present description discloses a method of treating diabetic retinopathy in a subject in need of treatment, comprising topically (non-invasively) administering to the surface eye of the subject a therapeutically effective amount of at least one chemically modified siRNA which inhibits expression of a target gene in the eye of the subject. The target gene is a human gene having mRNA set forth in any one SEQ ID NOS:1-2. In a currently preferred embodiment the siRNA is formulated for delivery as eye drops.

[0112] Without wishing to be bound to theory, blood vessels damaged from diabetic retinopathy can cause vision loss in two ways: Fragile, abnormal blood vessels can develop and leak blood into the center of the eye, blurring vision. This is proliferative retinopathy and is the fourth and most advanced stage of the disease. Fluid can leak into the center of the macula, resulting in blurred vision. This condition is called macular edema. It can occur at any stage of diabetic retinopathy, although it is more likely to occur as the disease progresses and is known as diabetic macular edema (DME).

Age related Macular Degeneration (AMD)

[0113] The most common cause of decreased best-corrected, vision in individuals over 65 years of age in the United States is the retinal disorder known as age-related macular degeneration (AMD). The area of the eye affected by AMD is the macula, a small area in the center of the retina, composed primarily of photoreceptor cells. As AMD progresses, the disease is characterized by loss of sharp, central vision. So-called "dry" AMD accounts for about 85% - 90% of AMD patients and involves alterations in eye pigment distribution, loss of photoreceptors and diminished retinal function due to overall atrophy of cells. "Wet" AMD involves proliferation of abnormal choroidal vessels leading to clots or scars in the sub-retinal space. Thus, the onset of "wet" AMD occurs because of the formation of an abnormal choroidal neovascular network (choroidal neovascularization, CNV) beneath the neural retina. The newly formed blood vessels are excessively leaky. This leads to accumulation of subretinal fluid and blood leading to loss of visual acuity. Eventually, there is total loss of functional retina in the involved region, as a large disciform scar involving choroids and retina forms. While dry AMD patients may retain vision of decreased quality, wet AMD often results in blindness. (Hamdi & Kenney, *Frontiers in Bioscience*, e305-314, May 2003).

[0114] In one aspect the present description discloses a method of attenuating expression of a target ocular mRNA in the eye of a subject suffering from AMD or DME, comprising topically (non-invasively) administering to the surface of the eye of the subject an effective amount of at least one chemically modified siRNA and a pharmaceutically acceptable carrier. The target ocular mRNA is product of a human gene listed in Table A3, having mRNA set forth in any one of SEQ ID NOS:1-2. In a currently preferred embodiment the siRNA is formulated for delivery as eye drops.

[0115] In another aspect the present description discloses a method of treating AMD or DME in a subject in need of treatment, comprising topically (non-invasively) administering to the surface of the eye of the subject a therapeutically effective amount of at least one chemically modified siRNA which inhibits expression of a target gene in the eye of the subject. The target gene is a human gene having mRNA set forth in any one of SEQ ID NOS:1-2. In a currently preferred embodiment the siRNA is formulated for delivery as eye drops.

Additional Ocular conditions to be treated by compounds of the present invention

Viral and bacterial conditions

[0116] Viral and bacterial conditions relating to ocular tissues can be treated by the compounds of the present invention. Conjunctivitis and other eyelid diseases or conditions can be treated, in particular by administering according to the methods of the present invention oligonucleotides such as siRNAs which target genes which are essential for replication and/or survival of the organisms which cause such conditions.

Vision Loss Associated with Tumors

[0117] In another aspect the present invention provides method of treating vision loss associated with a tumor in a subject in need thereof which comprises topically and non-invasively administering to the surface of the eye of the subject a therapeutically effective amount of at least one chemically modified siRNA which inhibits expression of at least one gene associated with the tumor in the subject in an amount effective to treat the vision loss.

[0118] Tumors that cause vision loss, according to the present invention, include both malignant neoplasms (cancers) and benign tumors. Tumors include tumors of any ocular tissue or any type of ocular cell, including, but not limited to, Choroidal tumors, Conjunctival tumors, Eyelid tumors, Infiltrative Intraocular tumors, Iris tumors, Metastatic Ocular tumors, Optic Nerve tumors, Orbital tumors and Retinal tumors. More specifically, a non exhaustive list of tumors and cancers which the present invention aims to treat includes Choroidal Tumors such as Choroidal Hemangioma, Choroidal Melanoma, Choroidal Metastasis, Choroidal Nevus, Choroidal Osteoma, Ciliary Body Melanoma and Nevus of Ota; Conjunctival Tumors such as Conjunctival Kaposi's Sarcoma, Epibulbar Dermoid, Lymphoma of the Conjunctiva, Melanoma and PAM with Atypia, Pigmented Conjunctival Tumors, Pingueculum, Pterygium, Squamous Carcinoma and Intraepithelial Neoplasia of the Conjunctiva; Eyelid Tumors such as Basal Cell Carcinoma, Capillary Hemangioma, Hydrocystoma, Nevus at the Eyelid Margin, Seborrhic Keratosis, Malignant Melanoma of the Eyelid, Sebaceous Carcinoma of the Eyelid and Squamous Carcinoma of the Eyelid; Infiltrative Intraocular Tumors such as Chronic Lymphocytic Leukemia, Infiltrative Choroidopathy and Intraocular Lymphoma; Iris Tumors such as Anterior Uveal Metastasis, Iris Cysts, Iris Melanocytoma, Iris Melanoma and Pearl Cyst of the Iris; Metastatic Ocular Tumors such

as Metastatic Choroidal Melanoma; Optic Nerve Tumors such as Choroidal Melanoma Affecting the Optic Nerve, Circumpapillary Metastasis with Optic Neuropathy, Optic Nerve Melanocytoma and Optic Nerve Sheath Meningioma; Orbital Tumors such as Adenoid Cystic Carcinoma of the Lacrimal Gland, Cavernous Hemangioma, of the Orbit, Lymphangioma of the Orbit, Orbital Mucocele, Orbital Pseudotumor, Orbital Rhabdomyosarcoma, Periocular Hemangioma of Childhood and Sclerosing Orbital Pseudotumor; Retinal Tumors such as Retinal Pigment Epithelial (RPE) Hypertrophy, Retinal Pigment Epithelium (RPE) Tumors, Retinoblastoma and von Hippel Angioma.

Pharmaceutical Compositions

[0119] While it may be possible for the oligonucleotide compounds of the present invention to be administered as the raw chemical, it is preferable to administer them as a pharmaceutical composition. Accordingly the present invention provides a pharmaceutical composition comprising one or more of the compounds of the invention; and a pharmaceutically acceptable carrier. This composition may comprise a mixture of two or more different oligonucleotides / siRNAs.

[0120] The invention further provides a pharmaceutical composition comprising at least one compound of the invention covalently or non-covalently bound to one or more compounds of the description in an amount effective to inhibit one or more genes as disclosed above; and a pharmaceutically acceptable carrier. The compound may be processed intracellularly by endogenous cellular complexes to produce one or more oligoribonucleotides of the invention.

[0121] The invention further provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and one or more of the compounds of the invention in an amount effective to down-regulate expression in of a human gene in an eye of a subject suffering from an eye disease or disorder.

[0122] The present invention also provides for a process of preparing a pharmaceutical composition, which comprises:

providing one or more compounds of the invention ; and

admixing said compound with a pharmaceutically acceptable carrier.

[0123] The pharmaceutically acceptable carrier is preferably selected by one with skill in the art for ophthalmological administration.

[0124] In various embodiments the pharmaceutical composition of the invention comprises at least one siRNA compound of the invention, or salt thereof, up to 99% by weight, mixed with a physiologically acceptable ophthalmic carrier medium such as water, sodium chloride, buffer, saline (e.g. phosphate buffered saline (PBS)), mannitol, and the like, and combinations thereof, to form an aqueous, sterile ophthalmic suspension or solution.

[0125] The pharmaceutical composition further optionally comprises at least one ophthalmologically acceptable preservative, such as for example benzalkonium chloride. Further, the ophthalmic pharmaceutical composition may include an ophthalmologically acceptable surfactant to assist in dissolving the siRNA.

[0126] Ophthalmic pharmaceutical composition of the invention may be prepared by dissolving or admixing one or more of the interfering RNA with an ophthalmologically acceptable carrier, such as for example a physiologically acceptable isotonic aqueous buffer.

[0127] In a preferred embodiment, the siRNA compound used in the preparation of an ophthalmological composition is admixed with an ophthalmologically acceptable carrier in a pharmaceutically effective dose. In certain preferred embodiments the ophthalmologically acceptable carrier is PBS.

[0128] In some embodiments the pharmaceutical ophthalmic compositions of the invention further comprise additional pharmaceutically active agents or a combination of pharmaceutically active agent, such as non-steroidal anti-inflammatory drugs, corticosteroids, antibiotics, and the like.

[0129] Additionally, the description discloses a method of inhibiting the expression of a gene of the present invention by at least 20% as compared to a control comprising contacting an mRNA transcript of the gene of the present invention with one or more of

the compounds of the invention. In some embodiments an active siRNA compound inhibits gene expression at a level of at least 20%, 30%, 40%, 50%, 60% or 70% as compared to control. In certain preferred embodiments inhibition is at a level of at least 75%, 80% or 90% as compared to control.

Combination therapy

[0130] The compounds of the present invention can be administered alone or in combination with another therapeutic agent useful in treating an eye disorder or disease.

[0131] In some embodiments the pharmaceutical ophthalmic compositions of the invention further comprise additional pharmaceutically active agents or a combination of pharmaceutically active agent, such as oligonucleotide, e.g. siRNA, non-steroidal anti-inflammatory drug, corticosteroid, antibiotic, and the like.

[0132] In one embodiment, the co-administration of two or more therapeutic agents achieves a synergistic effect, i.e., a therapeutic effect that is greater than the sum of the therapeutic effects of the individual components of the combination. In another embodiment, the co-administration of two or more therapeutic agents achieves an additive effect.

[0133] The active ingredients that comprise a combination therapy may be administered together via a single dosage form or by separate administration of each active agent. In certain embodiments, the first and second therapeutic agents are administered in a single dosage form. The agents may be formulated into a single solution for topical administration. Alternatively, the first therapeutic agent and the second therapeutic agents may be administered as separate compositions. The first active agent may be administered at the same time as the second active agent or the first active agent may be administered intermittently with the second active agent. The length of time between administration of the first and second therapeutic agent may be adjusted to achieve the desired therapeutic effect. For example, the second therapeutic agent may be administered only a few minutes (e.g., 1, 2, 5, 10, 30, or 60 min) or several hours (e.g., 2, 4, 6, 10, 12, 24, or 36 hr) after administration of the first therapeutic agent. In certain embodiments, it may be advantageous to administer more than one dosage of one of the therapeutic agents between administrations of the second therapeutic agent. For example, the second therapeutic agent may be administered at 2 hours and then again at 10 hours following administration of the first therapeutic agent. Alternatively, it may be advantageous to administer more than one dosage of the first therapeutic agent between administrations of the second therapeutic agent. In certain embodiments it is preferred that the therapeutic effects of each active ingredient overlap for at least a portion of the duration of each therapeutic agent so that the overall therapeutic effect of the combination therapy is attributable in part to the combined or synergistic effects of the combination therapy.

Delivery

[0134] The siRNA molecules as disclosed herein are delivered to the target tissue of the eye by direct application of the molecules prepared with a carrier or a diluent.

[0135] The term "naked siRNA" refers to siRNA molecules that are free from any delivery vehicle or formulation that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. For example, siRNA in PBS or an acceptable ophthalmological formulation is "naked siRNA". In certain embodiments of the invention the siRNA is delivered as naked siRNA. For certain applications, a formulation that increases the residence time of the siRNA in the eye or nasal passage may be desired, for example, by addition of a polymer or viscosity enhancing agent.

[0136] Delivery systems aimed specifically at the enhanced and improved delivery of siRNA into mammalian cells have been developed, (see, for example, Shen et al FEBS Let. 2003, 539:111-114; Xia et al., Nat. Biotech. 2002, 20:1006-1010; Reich et al., Mol. Vision 2003, 9: 210-216; Sorensen et al., J. Mol. Biol. 2003, 327: 761-766; Lewis et al., Nat. Gen. 2002, 32: 107-108 and Simeoni et al., NAR 2003, 31,11: 2717-2724).

[0137] The pharmaceutically acceptable carriers, solvents, diluents, excipients, adjuvants and vehicles as well as implant carriers generally refer to inert, non-toxic solid or liquid fillers, diluents or encapsulating material not reacting with the active ingredients of the invention and they include liposomes and microspheres. Examples of delivery systems useful in the present invention include US Patent Nos. 5,225,182; 5,169,383; 5,167,616; 4,959,217; 4,925,678; 4,487,603; 4,486,194; 4,447,233; 4,447,224; 4,439,196; and 4,475,196. Many other such implants, delivery systems, and modules are well known to those skilled in the art.

[0138] A review of the considerations to be taken into account in the preparation of a pharmaceutical composition for topical ocular delivery or intranasal delivery, can be found in Bar-Ilan and Neumann, in Textbook of Ocular Pharmacology, Zimmerman et al eds., Lippencott-Raven 1997.

[0139] Topical routes of administration are preferably employed for providing the subject with an effective dosage of the therapeutic siRNA compounds. Dosage forms may include dispersions, suspensions, solutions, ointments and the like. In certain preferred embodiments the pharmaceutical composition of the invention comprising at least one siRNA is delivered as eye drops. In another embodiment the pharmaceutical composition of the invention comprising at least one siRNA is delivered as a spray or mist. For example US Patent No. 4,052,985 discloses ophthalmic spray applicators.

[0140] In certain embodiments of the present invention the siRNA reaches its target cell locally, for example direct contact or by diffusion through cells, tissue or intracellular fluid. The siRNAs or pharmaceutical compositions of the present invention are administered and dosed in accordance with good medical practice, taking into account the clinical condition of the individual patient, the disease to be treated, scheduling of administration, patient age, sex, body weight and other factors known to medical practitioners.

[0141] The "therapeutically effective dose" for purposes herein is thus determined by such considerations as are known in the art. The dose must be effective to achieve improvement including but not limited to an improved course of disease, more rapid recovery, improvement of symptoms, elimination of symptoms and other indicators as are selected as appropriate measures by those skilled in the art. The siRNA of the invention can be administered in a single dose or in multiple doses.

[0142] In general, the active dose of compound for humans is in the range of from 1ng/kg to about 20-100 mg/kg body weight per day, preferably about 0.01 mg/kg to about 2-10 mg/kg body weight per day, in a regimen of one dose per day or twice or three or more times per day for a single dose or multiple dose regimen. In certain embodiments the siRNA compounds are formulated for topical application to the eye as eye drops and comprise about 5 µg/µl to about 60 µg/µl by volume of the composition, about 6.6 µg/µl by volume of the composition, about 25 µg/µl by volume of the composition, about 33.3 µg/µl by volume of the composition, about 50 µg/µl by volume of the composition.

[0143] The pH of the formulation is about pH 5 to about pH 8, or about pH 5 to about pH 7, or from about pH 5 to about pH 6. In certain embodiments the pH is about Ph 5.9, about pH 6.15, about pH 6.25, about pH 6.3, about pH 6.5, about pH 7.25. The compounds of the present invention can be administered topically to the surface of the eye. It should be noted that the compound is preferably applied as the compound or as pharmaceutically acceptable salt active ingredient in combination with pharmaceutically acceptable carriers, solvents, diluents, excipients, adjuvants and or vehicles. As disclosed herein the preferred method of delivery is topical application of an ophthalmic composition to the eye.

[0144] Liquid forms may be prepared for drops or spray. The liquid compositions include aqueous solutions, with and without organic co-solvents, aqueous or oil suspensions, emulsions with edible oils, as well as similar pharmaceutical vehicles.

[0145] These compounds may be administered to humans and other animals for therapy by any suitable route of administration to the eye, as by, for example, a spray or drops, and topically, as by ointments, or drops.

Methods of Treatment

[0146] In one aspect, the present disclosure relates to a method for the treatment of a subject in need of treatment for an eye disease or disorder associated with expression of a gene listed in Tables A1-A4 in the eye of the subject, comprising topically and non-invasively administering to the subject an amount of a chemically modified siRNA which inhibits expression of at least one of the genes. In certain preferred embodiments more than one siRNA compound to one or more than one gene target is administered.

[0147] In preferred embodiments the subject being treated is a warm-blooded animal and, in particular, mammals including human.

[0148] The methods herein disclosed comprise topically and non-invasively administering to the eye of the subject one or more inhibitory compounds which down-regulate expression of the genes of Tables A1-A4; and in particular siRNA in a therapeutically effective dose so as to thereby treat the subject.

[0149] The term "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down, attenuate the related eye disorder as listed above. Those in need of treatment include those already experiencing the disease or condition, those prone to having the disease or condition, and those in which the disease or condition is to be prevented. The compounds of the invention may be administered before, during or subsequent to the onset of the eye disease or condition or symptoms associated therewith. In cases where treatment is for the purpose of prevention, then the present invention relates to a method for delaying the onset of or averting the development of the disease or disorder.

[0150] Ocular disorders include Acute Zonal Occult Outer Retinopathy, Adie Syndrome, Age Related Macular Degeneration, Amblyopia, Aniridia, Anisocoria, Anophthalmos, Aphakia, Blepharitis, Blepharoptosis, Blepharospasm, Blindness, Cataract, Chalazion, Chorioretinitis, Choroideremia, Coloboma, Conjunctival Diseases, Conjunctivitis, Corneal Diseases, Corneal Dystrophies, Corneal Edema, Corneal Ulcer, Diabetic Macular Edema, Diabetic Retinopathy, Diplopia, Distichiasis, Dry Eye Syndromes, Duane Retraction Syndrome, Ectropion, Endophthalmitis, Entropion, Esotropia, Exfoliation Syndrome, Exotropia, Eye Abnormalities, Eye Neoplasms, General Fibrosis Syndrome, Glaucomas, Gyrate Atrophy, Hemianopsia, Hermanski-Pudlak Syndrome, Hordeolum, Horner Syndrome, Hyperopia, Hyphema, Iritis, Kearns-Sayer Syndrome, Keratitis, Keratoconus, Lacrimal Apparatus Diseases, Lacrimal Duct Obstruction, Lens Diseases, Macular Degeneration, Nystagmus, Pathologic, Ocular Motility Disorders, Oculomotor Nerve Diseases, Ophthalmoplegia, Optic Atrophies, Hereditary, Optic Nerve Diseases, Optic Neuritis, Ischemic Optic Neuropathy, Orbital Cellulitis, Papilledema, Presbyopia, Pterygium, Pupil Disorders, Refractive Errors, Retinal Detachment, Retinal Diseases, Retinal Vein Occlusion, Retinal Blastoma, Retinitis Pigmentosa, Retinopathy of Prematurity, Retinoschisis, Scleritis, Scotoma, Strabismus, Sjögrens Syndrome, Thygeson's Superficial Punctate Keratitis, Trachoma, Uveitis.

Oligonucleotides

[0151] The oligonucleotides useful in the methods disclosed herein are double stranded oligonucleotides and siRNA compounds and include unmodified and chemically and/ or structurally modified compounds.

[0152] The selection and synthesis of siRNA corresponding to known genes has been widely reported; see for example Ui-Tei et al., J Biomed Biotechnol. 2006; 65052; Chalk et al., BBRC. 2004, 319(1):264-74; Sioud & Leirdal, Met. Mol Biol. 2004, 252:457-69; Levenkova et al., Bioinform. 2004, 20(3):430-2; Ui-Tei et al., NAR. 2004, 32(3):936-48. For examples of the use and production of modified siRNA see for example Braasch et al., Biochem. 2003, 42(26):7967-75; Chiu et al., RNA. 2003, 9(9):1034-48; PCT Publication Nos. WO 2004/015107 and WO 02/44321 and US Patent Nos. 5,898,031 and 6,107,094.

[0153] The present invention provides double-stranded oligonucleotides (e.g. siRNAs), which down-regulate the expression of a desired gene. A siRNA of the invention is a duplex oligoribonucleotide in which the sense strand is derived from the mRNA sequence of the desired gene, and the antisense strand is complementary to the sense strand. In general, some deviation from the target mRNA sequence is tolerated without compromising the siRNA activity (see e.g. Czauderna et al., NAR. 2003, 31(11):2705-2716). An siRNA of the invention inhibits gene expression on a post-transcriptional level with or without destroying the mRNA. Without being bound by theory, siRNA may target the mRNA for specific cleavage and degradation and/ or may inhibit translation from the targeted message.

[0154] In some aspects the siRNA is blunt ended, i.e. Z and Z' are absent, on one or both ends. More specifically, the siRNA may be blunt ended on the end defined by the 5'-terminus of the first strand and the 3'-terminus of the second strand, and/or the end defined by the 3'-terminus of the first strand and the 5'-terminus of the second strand.

[0155] In other aspects at least one of the two strands may have an overhang of at least one nucleotide at the 5'-terminus; the overhang may consist of at least one deoxyribonucleotide. At least one of the strands may also optionally have an overhang of at least one nucleotide at the 3'-terminus. The overhang may consist of from about 1 to about 5 nucleotides.

[0156] The length of RNA duplex is 19 ribonucleotides. Further, the length of each strand is 19, ribonucleotides.

[0157] The complementarity between said first strand and the target nucleic acid is perfect (100%). In some aspects the strands are substantially complementary, i.e. having one, two or up to three mismatches between said first strand and the target nucleic acid. Substantially complementary refers to complementarity of greater than about 84%, to another sequence. For example in a duplex region consisting of 19 base pairs one mismatch results in 94.7% complementarity, two mismatches results in about 89.5% complementarity and 3 mismatches results in about 84.2% complementarity, rendering the duplex region substantially complementary. Accordingly substantially identical refers to identity of greater than about 84%, to another sequence.

[0158] The first strand and the second strand may be linked by a loop structure, which may be comprised of a non-nucleic acid polymer such as, *inter alia*, polyethylene glycol. Alternatively, the loop structure may be comprised of a nucleic acid, including modified and non-modified ribonucleotides and modified and non-modified deoxyribonucleotides.

[0159] Further, the 5'-terminus of the first strand of the siRNA may be linked to the 3'-terminus of the second strand, or the 3'-terminus of the first strand may be linked to the 5'-terminus of the second strand, said linkage being via a nucleic acid linker typically having a length between 2-100 nucleobases, preferably about 2 to about 30 nucleobases.

[0160] In preferred aspects of the compounds of the invention having alternating ribonucleotides modified in at least one of the antisense and the sense strands of the compound, for 19-mer and 23-mer oligomers the ribonucleotides at the 5' and 3' termini of the antisense strand are modified in their sugar residues, and the ribonucleotides at the 5' and 3' termini of the sense strand are unmodified in their sugar residues. For 21-mer oligomers the ribonucleotides at the 5' and 3' termini of the sense strand are modified in their sugar residues, and the ribonucleotides at the 5' and 3' termini of the antisense strand are unmodified in their sugar residues, or may have an optional additional modification at the 3' terminus. As mentioned above, it is preferred that the middle nucleotide of the antisense strand is unmodified.

[0161] Additionally, the description discloses siRNA comprising a double stranded nucleic acid molecule wherein 1, 2, or 3 of the nucleotides in one strand or both strands are substituted thereby providing at least one base pair mismatch. The substituted nucleotides in each strand are preferably in the terminal region of one strand or both strands.

[0162] According to one preferred embodiment of the invention, the antisense and the sense strands of the oligonucleotide / siRNA are phosphorylated only at the 3'-terminus and not at the 5'-terminus. According to another preferred embodiment of the invention, the antisense and the sense strands are non-phosphorylated. According to yet another preferred embodiment of the invention, the 5' most ribonucleotide in the sense strand is modified to abolish any possibility of *in vivo* 5'-phosphorylation.

[0163] Any siRNA sequence disclosed herein can be prepared having any of the modifications / structures disclosed herein. The combination of sequence plus structure is novel and can be used in the treatment of the conditions disclosed herein.

siRNA Structures

[0164] The selection and synthesis of siRNA corresponding to known genes has been widely reported; (see for example Ui-Tei et al., J Biomed Biotech. 2006; 2006: 65052; Chalk et al., BBRC. 2004, 319(1): 264-74; Sioud & Leirdal, Met. Mol Biol.; 2004, 252:457-69; Levenkova et al., Bioinform. 2004, 20(3):430-2; Ui-Tei et al., NAR. 2004, 32(3):936-48).

[0165] For examples of the use of, and production of, modified siRNA see, for example, Braasch et al., Biochem. 2003, 42(26):7967-75; Chiu et al., RNA, 2003, 9(9):1034-48; PCT publications WO 2004/015107 (atugen AG) and WO 02/44321 (Tuschl et al). US Patent Nos. 5,898,031 and 6,107,094, teach chemically modified oligomers. US Patent Publication Nos. 2005/0080246 and 2005/0042647 relate to oligomeric compounds having an alternating motif and dsRNA compounds having chemically modified internucleoside linkages, respectively.

[0166] Other modifications have been disclosed. The inclusion of a 5'-phosphate moiety was shown to enhance activity of siRNAs in *Drosophila* embryos (Boutla, et al., Curr. Biol. 2001, 11:1776-1780) and is required for siRNA function in human HeLa cells (Schwarz et al., Mol. Cell, 2002, 10:537-48). Amarzguioui et al., (NAR, 2003, 31(2):589-95) showed that siRNA activity depended on the positioning of the 2'-O-methyl (2'-OMe) modifications. Holen et al (NAR. 2003, 31(9):2401-07) report that an siRNA having small numbers of 2'-OMe modified nucleosides gave good activity compared to wild type but that the activity decreased as the numbers of 2'-OMe modified nucleosides was increased. Chiu and Rana (RNA. 2003, 9:1034-48) teach that incorporation of 2'-OMe modified nucleosides in the sense or antisense strand (fully modified strands) severely reduced siRNA activity relative to unmodified siRNA. The placement of a 2'-OMe group at the 5'-terminus on the antisense strand was reported to severely limit activity whereas placement at the 3'-terminus of the antisense and at both termini of the sense strand was tolerated (Czauderna et al., NAR. 2003, 31(11):2705-16; WO 2004/015107). The molecules of the present invention offer an advantage in that they are non-toxic and may be formulated as pharmaceutical compositions for treatment of various diseases.

[0167] The nucleotides can be selected from naturally occurring or synthetic modified bases. Naturally occurring bases include adenine, guanine, cytosine, thymine and uracil. Modified bases of nucleotides include inosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl, 2-propyl and other alkyl adenines, 5-halo uracil, 5-halo cytosine, 6-aza cytosine and 6-aza thymine,

pseudo uracil, 4- thiouracil, 8-halo adenine, 8-aminoadenine, 8-thiol adenine, 8-thioalkyl adenines, 8-hydroxyl adenine and other 8-substituted adenines, 8-halo guanines, 8-amino guanine, 8-thiol guanine, 8-thioalkyl guanines, 8- hydroxyl guanine and other substituted guanines, other aza and deaza adenines, other aza and deaza guanines, 5-trifluoromethyl uracil and 5- trifluoro cytosine. Molecules comprising one or more abasic moiety (unconventional or pseudonucleotide) are encompassed by the present invention, as well as molecules comprising alternating RNA and DNA nucleotides.

[0168] In addition, analogues of polynucleotides can be prepared wherein the structure of one or more nucleotide is fundamentally altered and better suited as therapeutic or experimental reagents. An example of a nucleotide analog is a peptide nucleic acid (PNA) wherein the deoxyribose (or ribose) phosphate backbone in DNA (or RNA) is replaced with a polyamide backbone which is similar to that found in peptides. PNA analogs have been shown to be resistant to enzymatic degradation and to have extended lives *in vivo* and *in vitro*.

[0169] Possible modifications to the sugar residue are manifold and include 2'-O alkyl, locked nucleic acid (LNA), glycol nucleic acid (GNA), threose nucleic acid (TNA), arabinoside, altritol (ANA) and other, 6-membered sugars including morpholinos, and cyclohexinyls. Further, said molecules may additionally contain modifications on the sugar, such as 2' alkyl, 2' fluoro, 2'O allyl, 2'amine and 2'alkoxy. Additional sugar modifications are discussed herein.

[0170] LNA compounds are disclosed in International Patent Publication Nos. WO 00/47599, WO 99/14226, and WO 98/39352. Examples of siRNA compounds comprising LNA nucleotides are disclosed in Elmen et al., (NAR 2005. 33(1):439-447) and in PCT Patent Publication No. WO 2004/083430.

[0171] The compounds of the present invention can be synthesized using one or more inverted nucleotides, for example inverted thymidine or inverted adenine (for example see Takei, et al., 2002. JBC 277(26):23800-06).

[0172] Backbone modifications, such as ethyl (resulting in a phospho-ethyl triester); propyl (resulting in a phospho-propyl triester); and butyl (resulting in a phospho-butyl triester) are also possible. Other backbone modifications include polymer backbones, cyclic backbones, acyclic backbones, thiophosphate-D-ribose backbones, amidates, and phosphonoacetate derivatives. Certain structures include siRNA compounds having one or a plurality of 2'-5' internucleotide linkages (bridges or backbone).

[0173] Further, the inhibitory nucleic acid molecules of the present description may comprise one or more gaps and/or one or more nicks and/or one or more mismatches. Without wishing to be bound by theory, gaps, nicks and mismatches have the advantage of partially destabilizing the nucleic acid / siRNA, so that it may be more easily processed by endogenous cellular machinery such as DICER, DROSHA or RISC into its inhibitory components.

[0174] The molecules of the present invention may comprise siRNAs, synthetic siRNAs, shRNAs and synthetic shRNAs, in addition to other nucleic acid sequences or molecules which encode such molecules or other inhibitory nucleotide molecules.

[0175] The compounds of the present invention may further comprise an end modification. A biotin group may be attached to either the most 5' or the most 3' nucleotide of the first and/or second strand or to both ends. In a more preferred embodiment the biotin group is coupled to a polypeptide or a protein. It is also within the scope of the present invention that the polypeptide or protein is attached through any of the other aforementioned modifications.

[0176] The various end modifications as disclosed herein are preferably located at the ribose moiety of a nucleotide of the nucleic acid as disclosed herein. More particularly, the end modification may be attached to or replace any of the OH-groups of the ribose moiety, including but not limited to the 2'OH, 3'OH and 5'OH position, provided that the nucleotide thus modified is a terminal nucleotide. Inverted abasic or abasic are nucleotides, either deoxyribonucleotides or ribonucleotides which do not have a nucleobase moiety. This kind of compound is, inter alia, described in Sternberger, et al., (Antisense Nucleic Acid Drug Dev, 2002.12, 131-43).

[0177] In the context of the present invention, a gap in a nucleic acid refers to the absence of one or more internal nucleotides in one strand, while a nick in a nucleic acid refers to the absence of an internucleotide linkage between two adjacent nucleotides in one strand. Any of the molecules of the present invention may contain one or more gaps and/or one or more nicks. Further provided by the present invention is an siRNA encoded by any of the molecules disclosed herein, a vector encoding any of the molecules disclosed herein, and a pharmaceutical composition comprising any of the molecules disclosed herein or the vectors encoding them; and a pharmaceutically acceptable carrier.


[0178] Particular molecules to be administered according to the methods of the present invention are disclosed below under the

heading "structural motifs". For the sake of clarity, any of these molecules can be administered according to any of the methods of the present invention.

Structural motifs

[0179] As disclosed herein the siRNA compounds that are chemically and or structurally modified according to one of the following modifications set forth in Structures below or as tandem siRNA or RNAstar (see below) are useful in the methods of the present disclosure. Sense and antisense oligonucleotide pairs, set forth in SEQ ID NOS:59-33596, are useful in preparing corresponding siRNA compounds.

[0180] In one aspect the present description discloses a compound set forth as Structure (A):




wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of $(N)_x$ and $(N')_y$ is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein each of x and y is an integer between 18 and 40;

wherein each of Z and Z' may be present or absent, but if present is 1-5 consecutive nucleotides covalently attached at the 3' terminus of the strand in which it is present; and wherein the sequence of $(N)_x$ comprises an antisense sequence substantially complementary to about 18 to about 40 consecutive ribonucleotides in the mRNA of a gene expressed in the retina and associated with an ocular disease or disorder. In some embodiments the mRNA is set forth in any one of SEQ ID NOS:1-58.

[0181] In certain aspects the present description discloses a compound having structure B



wherein each of $(N)_x$ and $(N')_y$ is an oligomer in which each consecutive N or N' is an unmodified ribonucleotide or a modified ribonucleotide joined to the next N or N' by a covalent bond;

wherein each of x and y =19, 21 or 23 and $(N)_x$ and $(N')_y$ are fully complementary wherein alternating ribonucleotides in each of $(N)_x$ and $(N')_y$ comprise 2'-OMe sugar modified ribonucleotides;

wherein the sequence of $(N')_y$ is a sequence complementary to $(N)_x$; and wherein the sequence of $(N)_x$ comprises an antisense sequence substantially complementary to about 18 to about 40 consecutive ribonucleotides in the mRNA set forth in any one of SEQ ID NOS:1-58.

[0182] In some embodiments each of $(N)_x$ and $(N')_y$ is independently phosphorylated or non-phosphorylated at the 3' and 5' termini.


[0183] In certain aspects of the description, alternating ribonucleotides are modified in both the antisense and the sense strands of the compound.

[0184] In certain aspects wherein each of x and y =19 or 23, each N at the 5' and 3' termini of $(N)_x$ is modified; and each N' at the 5' and 3' termini of $(N')_y$ is unmodified.

[0185] In certain aspects wherein each of x and y =21, each N at the 5' and 3' termini of $(N)_x$ is unmodified; and each N' at the 5' and 3' termini of $(N')_y$ is modified.

[0186] In particular embodiments, when x and y =19, the siRNA consists of a 2'-OMe sugar modified ribonucleotides on the first, third, fifth, seventh, ninth, eleventh, thirteenth, fifteenth, seventeenth and nineteenth nucleotide of the antisense strand $(N)_x$, and 2'-OMe sugar modified ribonucleotides in the second, fourth, sixth, eighth, tenth, twelfth, fourteenth, sixteenth and eighteenth nucleotide of the sense strand $(N')_y$. In various embodiments these particular siRNA compounds are blunt ended at both termini.

[0187] In some aspects, the present description discloses a compound having Structure (C):



wherein each of N and N' is a nucleotide independently selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of (N)x and (N')y is an oligomer in which each consecutive nucleotide is joined to the next nucleotide by a covalent bond and each of x and y is an integer between 18 and 40;

wherein in (N)x the nucleotides are unmodified or (N)x comprises alternating 2' O Me sugar modified ribonucleotides and unmodified ribonucleotides; and wherein the ribonucleotide located at the middle position of (N)x being 2'OMe sugar modified or unmodified, preferably unmodified;

wherein (N')y comprises unmodified ribonucleotides further comprising one modified nucleotide at a terminal or penultimate position, wherein the modified nucleotide is selected from the group consisting of a mirror nucleotide, a bicyclic nucleotide, a 2'-sugar modified nucleotide, an alritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

wherein if more than one nucleotide is modified in (N')y, the modified nucleotides may be consecutive;

wherein each of Z and Z' may be present or absent, but if present is 1-5 deoxyribonucleotides covalently attached at the 3' terminus of any oligomer to which it is attached;

wherein the sequence of (N')y comprises a sequence substantially complementary to (N)x; and wherein the sequence of (N)x comprises an antisense sequence substantially complementary to about 18 to about 40 consecutive ribonucleotides in the mRNA set forth in any one of SEQ ID NOS:1-58.

[0188] In particular embodiments, x=y=19 and in (N)x each modified ribonucleotide is a 2'-OMe sugar modified and the ribonucleotide located at the middle of (N)x is unmodified. Accordingly, in a compound wherein x=19, (N)x comprises 2'-O-methyl sugar modified ribonucleotides at positions 1, 3, 5, 7, 9, 11, 13, 15, 17 and 19. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 5. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 2, 4, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 6. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 15. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 14. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 1, 2, 3, 7, 9, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 5. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 1, 2, 3, 5, 7, 9, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 6. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 1, 2, 3, 5, 7, 9, 11, 13, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 15. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 1, 2, 3, 5, 7, 9, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 14. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 2, 4, 6, 7, 9, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 5. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 1, 2, 4, 6, 7, 9, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 5. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 14, 16, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 15. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 1, 2, 3, 5, 7, 9, 11, 13, 14, 16, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudonucleotide for example in position 15. In other embodiments, (N)x comprises 2'-OMe modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 7. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 8. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 9. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 10. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 11. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 and may further comprise at least one abasic or inverted

abasic pseudo-nucleotide for example in position 12. In other embodiments, (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 15, 17 and 19 and may further comprise at least one abasic or inverted abasic pseudo-nucleotide for example in position 13.

[0189] In yet other embodiments (N)x comprises at least one nucleotide mismatch relative to the target gene. In certain preferred embodiments, (N)x comprises a single nucleotide mismatch on position 5, 6, or 14. In one embodiment of Structure (C), at least two nucleotides at either or both the 5' and 3' termini of (N'y are joined by a 2'-5' phosphodiester bond. In certain preferred embodiments $x=y=19$ or $x=y=23$; in (N)x the nucleotides alternate between 2'-OMe sugar modified ribonucleotides and unmodified ribonucleotides, and the ribonucleotide located at the middle of (N)x being unmodified; and three nucleotides at the 3' terminus of (N'y are joined by two 2'-5' phosphodiester bonds (set forth herein as Structure I). In other preferred embodiments, $x=y=19$ or $x=y=23$; in (N)x the nucleotides alternate between 2'-OMe sugar modified ribonucleotides and unmodified ribonucleotides, and the ribonucleotide located at the middle of (N)x being unmodified; and four consecutive nucleotides at the 5' terminus of (N'y are joined by three 2'-5' phosphodiester bonds. In a further embodiment, an additional nucleotide located in the middle position of (N)y is 2'-OMe sugar modified. In another preferred embodiment, in (N)x the nucleotides alternate between 2'-OMe sugar modified ribonucleotides and unmodified ribonucleotides, and in (N'y four consecutive nucleotides at the 5' terminus are joined by three 2'-5' phosphodiester bonds and the 5' terminal nucleotide or two or three consecutive nucleotides at the 5' terminus comprise 3'-O-methyl (3'-OMe)modifications.

[0190] In certain preferred embodiments of Structure C, $x=y=19$ and in (N'y, at least one position comprises an abasic or inverted abasic pseudo-nucleotide, preferably five positions comprises an abasic or inverted abasic pseudo-nucleotides. In various embodiments, the following positions comprise an abasic or inverted abasic: positions 1 and 16-19, positions 15-19, positions 1-2 and 17-19, positions 1-3 and 18-19, positions 1-4 and 19 and positions 1-5. (N'y may further comprise at least one LNA nucleotide.

[0191] In certain preferred embodiments of Structure C, $x=y=19$ and in (N'y the nucleotide in at least one position comprises a mirror nucleotide, a deoxyribonucleotide and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide bond;

[0192] In certain preferred embodiments of Structure C, $x=y=19$ and (N'y comprises a mirror nucleotide. In various embodiments the mirror nucleotide is an L-DNA nucleotide. In certain embodiments the L-DNA is L-deoxyribocytidine. In some embodiments (N'y comprises L-DNA at position 18. In other embodiments (N'y comprises L-DNA at positions 17 and 18. In certain embodiments (N'y comprises L-DNA substitutions at positions 2 and at one or both of positions 17 and 18. In certain embodiments (N'y further comprises a 5' terminal cap nucleotide such as 5'-O-methyl DNA or an abasic or inverted abasic pseudo-nucleotide as an overhang.

[0193] In yet other embodiments (N'y comprises at least one nucleotide mismatch relative to the target gene. In certain preferred embodiments, (N'y comprises a single nucleotide mismatch on position 6, 14, or 15 .

[0194] In yet other embodiments (N'y comprises a DNA at position 15 and L-DNA at one or both of positions 17 and 18. In that structure, position 2 may further comprise an L-DNA or an abasic pseudo-nucleotide.

[0195] Other embodiments of Structure C are envisaged wherein $x=y=21$ or wherein $x=y=23$; in these embodiments the modifications for (N'y discussed above instead of being on positions 15, 16, 17, 18 are on positions 17, 18, 19, 20 for 21 mer and on positions 19, 20, 21, 22 for 23 mer ; similarly the modifications at one or both of positions 17 and 18 are on one or both of positions 19 or 20 for a 21 mer and one or both of positions 21 and 22 for a 23 mer. All modifications in the 19 mer are similarly adjusted for the 21 and 23 mers.

[0196] According to various embodiments of Structure (C), in (N'y 2, 3, 4, 5, 6, 7, 8 , 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides at the 3' terminus are linked by 2'-5' internucleotide linkages In one preferred embodiment, four consecutive nucleotides at the 3' terminus of (N'y are joined by three 2'-5' phosphodiester bonds, wherein one or more of the 2'-5' nucleotides which form the 2'-5' phosphodiester bonds further comprises a 3'-O-methyl sugar modification. Preferably the 3' terminal nucleotide of (N'y comprises a 2'-O-methyl sugar modification. In certain preferred embodiments of Structure C, $x=y=19$ and in (N'y two or more consecutive nucleotides at positions 15, 16, 17, 18 and 19 comprise a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide bond. In various embodiments the nucleotide forming the 2'-5' internucleotide bond comprises a 3' deoxyribose nucleotide or a 3' methoxy nucleotide. In some embodiments the nucleotides at positions 17 and 18 in (N'y are joined by a 2'-5' internucleotide bond. In other embodiments the nucleotides at positions 16, 17, 18, 16-17, 17-18, or 16-18 in (N'y are joined by a 2'-5' internucleotide bond.

[0197] In certain embodiments (N)_y comprises an L-DNA at position 2 and 2'-5' internucleotide bonds at positions 16, 17, 18, 16-17, 17-18, or 16-18. In certain embodiments (N)_y comprises 2'-5' internucleotide bonds at positions 16, 17, 18, 16-17, 17-18, or 16-18 and a 5' terminal cap nucleotide.

[0198] According to various embodiments of Structure (C), in (N)_y 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides at either terminus or 2-8 modified nucleotides at each of the 5' and 3' termini are independently mirror nucleotides. In some embodiments the mirror nucleotide is an L-ribonucleotide. In other embodiments the mirror nucleotide is an L-deoxyribonucleotide. The mirror nucleotide may further be modified at the sugar or base moiety or in an internucleotide linkage.

[0199] In one preferred embodiment of Structure (C), the 3' terminal nucleotide or two or three consecutive nucleotides at the 3' terminus of (N)_y are L-deoxyribonucleotides.

[0200] In other embodiments of Structure (C), in (N)_y 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides at either terminus or 2-8 modified nucleotides at each of the 5' and 3' termini are independently 2' sugar modified nucleotides. In some embodiments the 2' sugar modification comprises the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments the 2' sugar modification comprises a methoxy moiety (2'-OMe).

[0201] In one series of preferred embodiments, three, four or five consecutive nucleotides at the 5' terminus of (N)_y comprise the 2'-OMe modification. In another preferred embodiment, three consecutive nucleotides at the 3' terminus of (N)_y comprise the 2'-O-methyl modification.

[0202] In some embodiments of Structure (C), in (N)_y 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides at either or 2-8 modified nucleotides at each of the 5' and 3' termini are independently bicyclic nucleotide. In various embodiments the bicyclic nucleotide is a locked nucleic acid (LNA). A 2'-O, 4'-C-ethylene-bridged nucleic acid (ENA) is a species of LNA (see below).

[0203] In various embodiments (N)_y comprises modified nucleotides at the 5' terminus or at both the 3' and 5' termini.

[0204] In some embodiments of Structure (C), at least two nucleotides at either or both the 5' and 3' termini of (N)_y are joined by P-ethoxy backbone modifications. In certain preferred embodiments $x=y=19$ or $x=y=23$; in (N)_x the nucleotides alternate between modified ribonucleotides and unmodified ribonucleotides, each modified ribonucleotide being modified so as to have a 2'-O-methyl on its sugar and the ribonucleotide located at the middle position of (N)_x being unmodified; and four consecutive nucleotides at the 3' terminus or at the 5' terminus of (N)_y are joined by three P-ethoxy backbone modifications. In another preferred embodiment, three consecutive nucleotides at the 3' terminus or at the 5' terminus of (N)_y are joined by two P-ethoxy backbone modifications.

[0205] In some embodiments of Structure (C), in (N)_y 2, 3, 4, 5, 6, 7 or 8, consecutive ribonucleotides at each of the 5' and 3' termini are independently mirror nucleotides, nucleotides joined by 2'-5' phosphodiester bond, 2' sugar modified nucleotides or bicyclic nucleotide. In one embodiment, the modification at the 5' and 3' termini of (N)_y is identical. In one preferred embodiment, four consecutive nucleotides at the 5' terminus of (N)_y are joined by three 2'-5' phosphodiester bonds and three consecutive nucleotides at the 3' terminus of (N)_y are joined by two 2'-5' phosphodiester bonds. In another embodiment, the modification at the 5' terminus of (N)_y is different from the modification at the 3' terminus of (N)_y. In one specific embodiment, the modified nucleotides at the 5' terminus of (N)_y are mirror nucleotides and the modified nucleotides at the 3' terminus of (N)_y are joined by 2'-5' phosphodiester bond. In another specific embodiment, three consecutive nucleotides at the 5' terminus of (N)_y are LNA nucleotides and three consecutive nucleotides at the 3' terminus of (N)_y are joined by two 2'-5' phosphodiester bonds. In (N)_x the nucleotides alternate between modified ribonucleotides and unmodified ribonucleotides, each modified ribonucleotide being modified so as to have a 2'-O-methyl on its sugar and the ribonucleotide located at the middle of (N)_x being unmodified, or the ribonucleotides in (N)_x being unmodified

[0206] In another embodiment of Structure (C), the present invention provides a compound wherein $x=y=19$ or $x=y=23$; in (N)_x the nucleotides alternate between modified ribonucleotides and unmodified ribonucleotides, each modified ribonucleotide being modified so as to have a 2'-O-methyl on its sugar and the ribonucleotide located at the middle of (N)_x being unmodified; three nucleotides at the 3' terminus of (N)_y are joined by two 2'-5' phosphodiester bonds and three nucleotides at the 5' terminus of (N)_y are LNA such as ENA.

[0207] In another embodiment of Structure (C), five consecutive nucleotides at the 5' terminus of (N)_y comprise the 2'-O-methyl sugar modification and two consecutive nucleotides at the 3' terminus of (N)_y are L-DNA.

[0208] In yet another aspect the present description discloses a compound wherein $x=y=19$ or $x=y=23$; (N)_x consists of unmodified ribonucleotides; three consecutive nucleotides at the 3' terminus of (N')_y are joined by two 2'-5' phosphodiester bonds and three consecutive nucleotides at the 5' terminus of (N')_y are LNA such as ENA.

[0209] According to other embodiments of Structure (C), in (N')_y the 5' or 3' terminal nucleotide, or 2, 3, 4, 5 or 6 consecutive nucleotides at either termini or 1-4 modified nucleotides at each of the 5' and 3' termini are independently phosphonocarboxylate or phosphinocarboxylate nucleotides (PACE nucleotides). In some embodiments the PACE nucleotides are deoxyribonucleotides. In some preferred embodiments in (N')_y, 1 or 2 consecutive nucleotides at each of the 5' and 3' termini are PACE nucleotides. Examples of PACE nucleotides and analogs are disclosed in US Patent Nos. 6,693,187 and 7,067,641 both incorporated by reference.

[0210] In additional aspects, the present description discloses a compound having Structure (D):

(D) 5'	(N) _x -Z 3'	antisense strand
3'	Z'-(N') _y 5'	sense strand

wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide or a modified deoxyribonucleotide;

wherein each of (N)_x and (N')_y is an oligomer in which each consecutive nucleotide is joined to the next nucleotide by a covalent bond and each of x and y is an integer between 18 and 40;

wherein (N)_x comprises unmodified ribonucleotides further comprising one modified nucleotide at the 3' terminal or penultimate position, wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an alritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

wherein (N')_y comprises unmodified ribonucleotides further comprising one modified nucleotide at the 5' terminal or penultimate position, wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an alritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

wherein in each of (N)_x and (N')_y modified and unmodified nucleotides are not alternating;

wherein each of Z and Z' may be present or absent, but if present is 1-5 deoxyribonucleotides covalently attached at the 3' terminus of any oligomer to which it is attached;

wherein the sequence of (N')_y is a sequence substantially complementary to (N)_x; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in the mRNA set forth in any one of SEQ ID NOS:1-58.

[0211] In one embodiment of Structure (D), $x=y=19$ or $x=y=23$; (N)_x comprises unmodified ribonucleotides in which two consecutive nucleotides linked by one 2'-5' internucleotide linkage at the 3' terminus; and (N')_y comprises unmodified ribonucleotides in which two consecutive nucleotides linked by one 2'-5' internucleotide linkage at the 5' terminus.

[0212] In some embodiments, $x=y=19$ or $x=y=23$; (N)_x comprises unmodified ribonucleotides in which three consecutive nucleotides at the 3' terminus are joined together by two 2'-5' phosphodiester bonds; and (N')_y comprises unmodified ribonucleotides in which four consecutive nucleotides at the 5' terminus are joined together by three 2'-5' phosphodiester bonds (set forth herein as Structure II).

[0213] According to various embodiments of Structure (D) 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 3' terminus of (N)_x and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N')_y are linked by 2'-5' internucleotide linkages.

[0214] According to one preferred embodiment of Structure (D), four consecutive nucleotides at the 5' terminus of (N')_y are joined by three 2'-5' phosphodiester bonds and three consecutive nucleotides at the 3' terminus of (N)_x are joined by two 2'-5' phosphodiester bonds. Three nucleotides at the 5' terminus of (N')_y and two nucleotides at the 3' terminus of (N)_x may also comprise 3'-O-methyl modifications.

[0215] According to various embodiments of Structure (D), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides starting at the ultimate or penultimate position of the 3' terminus of (N)_x and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N')_y are independently mirror nucleotides. In

some embodiments the mirror is an L-ribonucleotide. In other embodiments the mirror nucleotide is L-deoxyribonucleotide.

[0216] In other embodiments of Structure (D), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 3' terminus of (N)x and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N)y are independently 2' sugar modified nucleotides. In some embodiments the 2' sugar modification comprises the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments the 2' sugar modification comprises a methoxy moiety (2'-OMe).

[0217] In one preferred embodiment of Structure (D), five consecutive nucleotides at the 5' terminus of (N)y comprise the 2'-O-methyl modification and five consecutive nucleotides at the 3' terminus of (N)x comprise the 2'-O-methyl modification. In another preferred embodiment of Structure (D), ten consecutive nucleotides at the 5' terminus of (N)y comprise the 2'-O-methyl modification and five consecutive nucleotides at the 3' terminus of (N)x comprise the 2'-O-methyl modification. In another preferred embodiment of Structure (D), thirteen consecutive nucleotides at the 5' terminus of (N)y comprise the 2'-O-methyl modification and five consecutive nucleotides at the 3' terminus of (N)x comprise the 2'-O-methyl modification.

[0218] In some embodiments of Structure (D), in (N)y 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 3' terminus of (N)x and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N)y are independently a bicyclic nucleotide. In various embodiments the bicyclic nucleotide is a locked nucleic acid (LNA) such as a 2'-O, 4'-C-ethylene-bridged nucleic acid (ENA).

[0219] In various embodiments of Structure (D), (N)y comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;


[0220] In various embodiments of Structure (D), (N)x comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

[0221] In embodiments wherein each of the 3' and 5' termini of the same strand comprises a modified nucleotide, the modification at the 5' and 3' termini is identical. In another embodiment, the modification at the 5' terminus is different from the modification at the 3' terminus of the same strand. In one specific embodiment, the modified nucleotides at the 5' terminus are mirror nucleotides and the modified nucleotides at the 3' terminus of the same strand are joined by 2'-5' phosphodiester bond.

[0222] In one specific embodiment of Structure (D), five consecutive nucleotides at the 5' terminus of (N)y comprise the 2'-O-methyl modification and two consecutive nucleotides at the 3' terminus of (N)y are L-DNA. In addition, the compound may further comprise five consecutive 2'-O-methyl modified nucleotides at the 3' terminus of (N)x.

[0223] In various embodiments of Structure (D), the modified nucleotides in (N)x are different from the modified nucleotides in (N)y. For example, the modified nucleotides in (N)x are 2' sugar modified nucleotides and the modified nucleotides in (N)y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are mirror nucleotides and the modified nucleotides in (N)y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are nucleotides linked by 2'-5' internucleotide linkages and the modified nucleotides in (N)y are mirror nucleotides.

[0224] In additional aspects the present description discloses a compound having Structure (E):



wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide or a modified deoxyribonucleotide;

wherein each of (N)x and (N')y is an oligomer in which each consecutive nucleotide is joined to the next nucleotide by a covalent bond and each of x and y is an integer between 18 and 40;

wherein (N)x comprises unmodified ribonucleotides further comprising one modified nucleotide at the 5' terminal or penultimate position, wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

wherein (N')_y comprises unmodified ribonucleotides further comprising one modified nucleotide at the 3' terminal or penultimate position, wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;
 wherein in each of (N)_x and (N')_y modified and unmodified nucleotides are not alternating;
 wherein each of Z and Z' may be present or absent, but if present is 1-5 deoxyribonucleotides covalently attached at the 3' terminus of any oligomer to which it is attached;
 wherein the sequence of (N')_y is a sequence substantially complementary to (N)_x; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in the mRNA set forth in any one of SEQ ID NOS:1-58.

[0225] In certain preferred embodiments the ultimate nucleotide at the 5' terminus of (N)_x is unmodified.

[0226] According to various embodiments of Structure (E) 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N)_x, preferably starting at the 5' penultimate position, and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 3' terminus of (N')_y are linked by 2'-5' internucleotide linkages.

[0227] According to various embodiments of Structure (E), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides starting at the ultimate or penultimate position of the 5' terminus of (N)_x, preferably starting at the 5' penultimate position, and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides starting at the ultimate or penultimate position of the 3' terminus of (N')_y are independently mirror nucleotides. In some embodiments the mirror is an L-ribonucleotide. In other embodiments the mirror nucleotide is L-deoxyribonucleotide.

[0228] In other embodiments of Structure (E), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N)_x, preferably starting at the 5' penultimate position, and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 3' terminus of (N')_y are independently 2' sugar modified nucleotides. In some embodiments the 2' sugar modification comprises the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments the 2' sugar modification comprises a methoxy moiety (2'-OMe).

[0229] In some embodiments of Structure (E), in (N')_y 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 5' terminus of (N)_x, preferably starting at the 5' penultimate position, and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides starting at the ultimate or penultimate position of the 3' terminus of (N')_y are independently a bicyclic nucleotide. In various embodiments the bicyclic nucleotide is a locked nucleic acid (LNA) such as a 2'-O, 4'-C-ethylene-bridged nucleic acid (ENA).

[0230] In various embodiments of Structure (E), (N')_y comprises modified nucleotides selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, a nucleotide joined to an adjacent nucleotide by a P-alkoxy backbone modification or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage at the 3' terminus or at each of the 3' and 5' termini.


[0231] In various embodiments of Structure (E), (N)_x comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage at the 5' terminus or at each of the 3' and 5' termini.

[0232] In one embodiment where both 3' and 5' termini of the same strand comprise a modified nucleotide, the modification at the 5' and 3' termini is identical. In another embodiment, the modification at the 5' terminus is different from the modification at the 3' terminus of the same strand. In one specific embodiment, the modified nucleotides at the 5' terminus are mirror nucleotides and the modified nucleotides at the 3' terminus of the same strand are joined by 2'-5' phosphodiester bond.

[0233] In various embodiments of Structure (E), the modified nucleotides in (N)_x are different from the modified nucleotides in (N')_y. For example, the modified nucleotides in (N)_x are 2' sugar modified nucleotides and the modified nucleotides in (N')_y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)_x are mirror nucleotides and the modified nucleotides in (N')_y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)_x are nucleotides linked by 2'-5' internucleotide linkages and the modified nucleotides in (N')_y are mirror

nucleotides.

[0234] In additional aspects the present description discloses a compound having Structure (F):



wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide or a modified deoxyribonucleotide;

wherein each of (N)x and (N')y is an oligomer in which each consecutive nucleotide is joined to the next nucleotide by a covalent bond and each of x and y is an integer between 18 and 40;

wherein each of (N)x and (N')y comprise unmodified ribonucleotides in which each of (N)x and (N')y independently comprise one modified nucleotide at the 3' terminal or penultimate position wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, a nucleotide joined to an adjacent nucleotide by a P-alkoxy backbone modification or a nucleotide joined to an adjacent nucleotide by a 2'-5' phosphodiester bond;

wherein in each of (N)x and (N')y modified and unmodified nucleotides are not alternating;

wherein each of Z and Z' may be present or absent, but if present is 1-5 deoxyribonucleotides covalently attached at the 3' terminus of any oligomer to which it is attached;

wherein the sequence of (N')y is a sequence substantially complementary to (N)x; and wherein the sequence of (N)x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS: 1-58.

[0235] In some embodiments of Structure (F), $x=y=19$ or $x=y=23$; (N')y comprises unmodified ribonucleotides in which two consecutive nucleotides at the 3' terminus comprises two consecutive mirror deoxyribonucleotides; and (N)x comprises unmodified ribonucleotides in which one nucleotide at the 3' terminus comprises a mirror deoxyribonucleotide (set forth as Structure III).

[0236] According to various embodiments of Structure (F) 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 3' termini of (N)x and (N')y are linked by 2'-5' internucleotide linkages.

[0237] According to one preferred embodiment of Structure (F), three consecutive nucleotides at the 3' terminus of (N')y are joined by two 2'-5' phosphodiester bonds and three consecutive nucleotides at the 3' terminus of (N)x are joined by two 2'-5' phosphodiester bonds.

[0238] According to various embodiments of Structure (F), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides independently beginning at the ultimate or penultimate position of the 3' termini of (N)x and (N')y are independently mirror nucleotides. In some embodiments the mirror nucleotide is an L-ribonucleotide. In other embodiments the mirror nucleotide is an L-deoxyribonucleotide.

[0239] In other embodiments of Structure (F), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 3' termini of (N)x and (N')y are independently 2' sugar modified nucleotides. In some embodiments the 2' sugar modification comprises the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments the 2' sugar modification comprises a methoxy moiety (2'-OMe).

[0240] In some embodiments of Structure (F), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 3' termini of (N)x and (N')y are independently a bicyclic nucleotide. In various embodiments the bicyclic nucleotide is a locked nucleic acid (LNA) such as a 2'-O, 4'-C-ethylene-bridged nucleic acid (ENA).

[0241] In various embodiments of Structure (F), (N')y comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage at the 3' terminus or at both the 3' and 5' termini.

[0242] In various embodiments of Structure (F), (N)x comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage at the 3' terminus or at each of the 3' and 5' termini.

[0243] In one embodiment where each of 3' and 5' termini of the same strand comprise a modified nucleotide, the modification at the 5' and 3' termini is identical. In another embodiment, the modification at the 5' terminus is different from the modification at the 3' terminus of the same strand. In one specific embodiment, the modified nucleotides at the 5' terminus are mirror nucleotides and the modified nucleotides at the 3' terminus of the same strand are joined by 2'-5' phosphodiester bond.

[0244] In various embodiments of Structure (F), the modified nucleotides in (N)x are different from the modified nucleotides in (N')y. For example, the modified nucleotides in (N)x are 2' sugar modified nucleotides and the modified nucleotides in (N')y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are mirror nucleotides and the modified nucleotides in (N')y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are nucleotides linked by 2'-5' internucleotide linkages and the modified nucleotides in (N')y are mirror nucleotides.

[0245] In additional aspects the present description discloses a compound having Structure (

(G) 5'	(N)x -Z 3'	antisense strand
3'	Z'-(N')y 5'	sense strand

wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide or a modified deoxyribonucleotide;

wherein each of (N)x and (N')y is an oligomer in which each consecutive nucleotide is joined to the next nucleotide by a covalent bond and each of x and y is an integer between 18 and 40;

wherein each of (N)x and (N')y comprise unmodified ribonucleotides in which each of (N)x and (N')y independently comprise one modified nucleotide at the 5' terminal or penultimate position wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, a nucleotide joined to an adjacent nucleotide by a P-alkoxy backbone modification or a nucleotide joined to an adjacent nucleotide by a 2'-5' phosphodiester bond;

wherein for (N)x the modified nucleotide is preferably at penultimate position of the 5' terminal;

wherein in each of (N)x and (N')y modified and unmodified nucleotides are not alternating;

wherein each of Z and Z' may be present or absent, but if present is 1-5 deoxyribonucleotides covalently attached at the 3' terminus of any oligomer to which it is attached;

wherein the sequence of (N')y is a sequence substantially complementary to (N)x; and wherein the sequence of (N)x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in the mRNA set forth in any one of SEQ ID NOS:1-58.

[0246] In some embodiments of Structure (G), x=y=19.

[0247] According to various embodiments of Structure (G) 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 5' termini of (N)x and (N')y are linked by 2'-5' internucleotide linkages. For (N)x the modified nucleotides preferably starting at the penultimate position of the 5' terminal.

[0248] According to various embodiments of Structure (G), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides independently beginning at the ultimate or penultimate position of the 5' termini of (N)x and (N')y are independently mirror nucleotides. In some embodiments the mirror nucleotide is an L-ribonucleotide. In other embodiments the mirror nucleotide is an L-deoxyribonucleotide. For (N)x the modified nucleotides preferably starting at the penultimate position of the 5' terminal.

[0249] In other embodiments of Structure (G), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 5' termini of (N)x and (N')y are independently 2' sugar modified nucleotides. In some embodiments the 2' sugar modification comprises the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments the 2' sugar modification comprises a methoxy moiety (2'-OMe). In some preferred embodiments the consecutive modified nucleotides preferably begin at the penultimate position of the 5' terminus of (N)x.

[0250] In one preferred embodiment of Structure (G), five consecutive ribonucleotides at the 5' terminus of (N')y comprise a 2'-O-methyl modification and one ribonucleotide at the 5' penultimate position of (N')x comprises a 2'-O-methyl modification. In another preferred embodiment of Structure (G), five consecutive ribonucleotides at the 5' terminus of (N')y comprise a 2'-O-methyl modification and two consecutive ribonucleotides at the 5' terminal position of (N')x comprise a 2'-O-methyl modification.

[0251] In some embodiments of Structure (G), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 5' termini of (N)x and (N')y are bicyclic nucleotides. In

various embodiments the bicyclic nucleotide is a locked nucleic acid (LNA) such as a 2'-O, 4'-C-ethylene-bridged nucleic acid (ENA). In some preferred embodiments the consecutive modified nucleotides preferably begin at the penultimate position of the 5' terminus of (N)x.

[0252] In various embodiments of Structure (G), (N')y comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage at the 5' terminus or at each of the 3' and 5' termini.

[0253] In various embodiments of Structure (G), (N)x comprises a modified nucleotide selected from a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage at the 5' terminus or at each of the 3' and 5' termini.

[0254] In one embodiment where each of 3' and 5' termini of the same strand comprise a modified nucleotide, the modification at the 5' and 3' termini is identical. In another embodiment, the modification at the 5' terminus is different from the modification at the 3' terminus of the same strand. In one specific embodiment, the modified nucleotides at the 5' terminus are mirror nucleotides and the modified nucleotides at the 3' terminus of the same strand are joined by 2'-5' phosphodiester bond. In various embodiments of Structure (G), the modified nucleotides in (N)x are different from the modified nucleotides in (N')y. For example, the modified nucleotides in (N)x are 2' sugar modified nucleotides and the modified nucleotides in (N')y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are mirror nucleotides and the modified nucleotides in (N')y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are nucleotides linked by 2'-5' internucleotide linkages and the modified nucleotides in (N')y are mirror nucleotides.

[0255] In additional aspects, the present description discloses a compound having Structure (H):

(H) 5'	(N)x-Z 3'	antisense strand
3'	Z'-(N')y 5'	sense strand

wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide or a modified deoxyribonucleotide;

wherein each of (N)x and (N')y is an oligomer in which each consecutive nucleotide is joined to the next nucleotide by a covalent bond and each of x and y is an integer between 18 and 40;

wherein (N)x comprises unmodified ribonucleotides further comprising one modified nucleotide at the 3' terminal or penultimate position or the 5' terminal or penultimate position, wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

wherein (N')y comprises unmodified ribonucleotides further comprising one modified nucleotide at an internal position, wherein the modified nucleotide is selected from the group consisting of a bicyclic nucleotide, a 2' sugar modified nucleotide, a mirror nucleotide, an altritol nucleotide, or a nucleotide joined to an adjacent nucleotide by an internucleotide linkage selected from a 2'-5' phosphodiester bond, a P-alkoxy linkage or a PACE linkage;

wherein in each of (N)x and (N')y modified and unmodified nucleotides are not alternating;

wherein each of Z and Z' may be present or absent, but if present is 1-5 deoxyribonucleotides covalently attached at the 3' terminus of any oligomer to which it is attached;

wherein the sequence of (N')y is a sequence substantially complementary to (N)x; and wherein the sequence of (N)x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS:1-58.

[0256] In one embodiment of Structure (H), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides independently beginning at the ultimate or penultimate position of the 3' terminus or the 5' terminus or both termini of (N)x are independently 2' sugar modified nucleotides, bicyclic nucleotides, mirror nucleotides, altritol nucleotides or nucleotides joined to an adjacent nucleotide by a 2'-5' phosphodiester bond and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive internal ribonucleotides in (N')y are independently 2' sugar modified nucleotides, bicyclic nucleotides, mirror nucleotides, altritol nucleotides or nucleotides joined to an adjacent nucleotide by a 2'-5' phosphodiester bond. In some embodiments the 2' sugar modification comprises the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments the 2' sugar modified ribonucleotide comprises a methoxy moiety (2'-OMe).

[0257] In another embodiment of Structure (H), 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive ribonucleotides

independently beginning at the ultimate or penultimate position of the 3' terminus or the 5' terminus or 2-8 consecutive nucleotides at each of 5' and 3' termini of (N')y are independently 2' sugar modified nucleotides, bicyclic nucleotides, mirror nucleotides, altritol nucleotides or nucleotides joined to an adjacent nucleotide by a 2'-5' phosphodiester bond, and 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive internal ribonucleotides in (N)x are independently 2' sugar modified nucleotides, bicyclic nucleotides, mirror nucleotides, altritol nucleotides or nucleotides joined to an adjacent nucleotide by a 2'-5' phosphodiester bond.

[0258] In one embodiment wherein each of 3' and 5' termini of the same strand comprises a modified nucleotide, the modification at the 5' and 3' termini is identical. In another embodiment, the modification at the 5' terminus is different from the modification at the 3' terminus of the same strand. In one specific embodiment, the modified nucleotides at the 5' terminus are mirror nucleotides and the modified nucleotides at the 3' terminus of the same strand are joined by 2'-5' phosphodiester bond.

[0259] In various embodiments of Structure (H), the modified nucleotides in (N)x are different from the modified nucleotides in (N')y. For example, the modified nucleotides in (N)x are 2' sugar modified nucleotides and the modified nucleotides in (N')y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are mirror nucleotides and the modified nucleotides in (N')y are nucleotides linked by 2'-5' internucleotide linkages. In another example, the modified nucleotides in (N)x are nucleotides linked by 2'-5' internucleotide linkages and the modified nucleotides in (N')y are mirror nucleotides.

[0260] In one preferred embodiment of Structure (H), $x=y=19$; three consecutive ribonucleotides at the 9-11 nucleotide positions 9-11 of (N')y comprise 2'-O-methyl modification and five consecutive ribonucleotides at the 3' terminal position of (N)x comprise 2'-O-methyl modification.

[0261] The present invention provides a compound having Structure (I) set forth below:

(I) 5'	(N)x - Z 3'	(antisense strand)
3'	Z'-(N')y- z' 5'	(sense strand)

wherein each of N and N' is a ribonucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of (N)x and (N')y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' may be present or absent, but if present is independently 1-5 consecutive nucleotides covalently attached at the 3' terminus of the strand in which it is present;

wherein z' may be present or absent, but if present is a capping moiety covalently attached at the 5' terminus of (N')y;

wherein $x = y = 19$;

wherein (N)x comprises modified and unmodified ribonucleotides, each modified ribonucleotide having a 2'-O-methyl on its sugar, wherein N at the 3' terminus of (N)x is a modified ribonucleotide, (N)x comprises at least five alternating modified ribonucleotides beginning at the 3' end and at least nine modified ribonucleotides in total and each remaining N is an unmodified ribonucleotide;

wherein in (N')y at least one unconventional moiety is present, which unconventional moiety may be an abasic ribose moiety, an abasic deoxyribose moiety, a modified or unmodified deoxyribonucleotide, a mirror nucleotide, and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate bond; and

wherein the sequence of (N)x is substantially complementary to the sequence of (N')y; and wherein the sequence of (N)x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS:1-2.

[0262] The at least one unconventional moiety is present at positions 18 in (N')y.

[0263] The unconventional moiety is an L-DNA moiety.

[0264] In Structure (I) (N)x comprises nine alternating modified ribonucleotides further comprising a 2'O modified nucleotide at position 2. (N)x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17, 19.

[0265] In various embodiments z' is present and is selected from an abasic ribose moiety, a deoxyribose moiety; an inverted abasic ribose moiety, a deoxyribose moiety; C6-amino-Pi; a mirror nucleotide.

[0266] In another aspect the present description discloses a compound having Structure (J) set forth below:

(J) 5'	(N)x - Z 3' (antisense strand)
--------	--------------------------------

3' Z'-(N')_y-z" 5'(sense strand)

wherein each of N and N' is a ribonucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' may be present or absent, but if present is independently 1-5 consecutive nucleotides covalently attached at the 3' terminus of the strand in which it is present;

wherein z" may be present or absent but if present is a capping moiety covalently attached at the 5' terminus of (N')_y;

wherein x=18 to 27;

wherein y=18 to 27;

wherein (N)_x comprises modified or unmodified ribonucleotides, and optionally at least one unconventional moiety;

wherein in (N')_y at least one unconventional moiety is present, which unconventional moiety may be an abasic ribose moiety, an abasic deoxyribose moiety, a modified or unmodified deoxyribonucleotide, a mirror nucleotide, a non-base pairing nucleotide analog or a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate bond; and

[0267] wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS:1-58.

[0268] In some embodiments x=y=19. In other embodiments x=y=23. In some preferred embodiments (N)_x comprises modified and unmodified ribonucleotides, and at least one unconventional moiety.

[0269] In some embodiments in (N)_x the N at the 3' terminus is a modified ribonucleotide and (N)_x comprises at least 8 modified ribonucleotides. In other embodiments at least 5 of the at least 8 modified ribonucleotides are alternating beginning at the 3' end. In some embodiments (N)_x comprises an abasic moiety in one of positions 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15.

[0270] In some embodiments the at least one unconventional moiety in (N')_y is present at positions 15, 16, 17, or 18. In some embodiments the unconventional moiety is selected from a mirror nucleotide, an abasic ribose moiety and an abasic deoxyribose moiety. In some preferred embodiments the unconventional moiety is a mirror nucleotide, preferably an L-DNA moiety. In some embodiments an L-DNA moiety is present at position 17, position 18 or positions 17 and 18. In other embodiments the at least one unconventional moiety in (N')_y is an abasic ribose moiety or an abasic deoxyribose moiety.

[0271] In various embodiments of Structure (X) z" is present and is selected from an abasic ribose moiety, a deoxyribose moiety; an inverted abasic ribose moiety, a deoxyribose moiety; C6-amino-Pi; a mirror nucleotide.

[0272] In yet another aspect the present description discloses a compound having Structure (K) set forth below:

(K) 5' (N)_x-Z 3' (antisense strand)
3' Z'-(N')_y-z" 5' (sense strand)

wherein each of N and N' is a ribonucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' may be present or absent, but if present is independently 1-5 consecutive nucleotides covalently attached at the 3' terminus of the strand in which it is present;

wherein z" may be present or absent but if present is a capping moiety covalently attached at the 5' terminus of (N')_y;

wherein x=18 to 27;

wherein y=18 to 27;

wherein (N)_x comprises a combination of modified or unmodified ribonucleotides and unconventional moieties, any modified ribonucleotide having a 2'-O-methyl on its sugar;

wherein (N')_y comprises modified or unmodified ribonucleotides and optionally an unconventional moiety, any modified ribonucleotide having a 2'OMe on its sugar;

wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS:1-58; and wherein there are less than 15 consecutive nucleotides complementary to the mRNA.

[0273] In some embodiments x=y=19. In other embodiments x=y=23. In some preferred embodiments the at least one preferred

one unconventional moiety is present in (N)_x and is an abasic ribose moiety or an abasic deoxyribose moiety. In other embodiments the at least one unconventional moiety is present in (N)_x and is a non-base pairing nucleotide analog. In various embodiments (N')_y comprises unmodified ribonucleotides. In some embodiments (N)_x comprises at least five abasic ribose moieties or abasic deoxyribose moieties or a combination thereof. In certain embodiments (N)_x and/or (N')_y comprise modified ribonucleotides which do not base pair with corresponding modified or unmodified ribonucleotides in (N')_y and/or (N)_x.

[0274] In various aspects the present description discloses an siRNA set forth in Structure (L):

(L) 5' (N)_xZ 3' (antisense strand)
3' Z'-(N')_y 5' (sense strand)

wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' are absent;

wherein x=y=19;

wherein in (N')_y the nucleotide in at least one of positions 15, 16, 17, 18 and 19 comprises a nucleotide selected from an abasic pseudo-nucleotide, a mirror nucleotide, a deoxyribonucleotide and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide bond;

wherein (N)_x comprises alternating modified ribonucleotides and unmodified ribonucleotides each modified ribonucleotide being modified so as to have a 2'-O-methyl on its sugar and the ribonucleotide located at the middle position of (N)_x being modified or unmodified, preferably unmodified; and

wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS:1-58.

[0275] In some embodiments of Structure (L), in (N')_y the nucleotide in one or both of positions 17 and 18 comprises a modified nucleotide selected from an abasic pseudo-nucleotide, a mirror nucleotide and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide bond. In some embodiments the mirror nucleotide is selected from L-DNA and L-RNA. In various embodiments the mirror nucleotide is L-DNA.

[0276] In various embodiments (N')_y comprises a modified nucleotide at position 15 wherein the modified nucleotide is selected from a mirror nucleotide and a deoxyribonucleotide.


[0277] In certain embodiments (N')_y further comprises a modified nucleotide or pseudo nucleotide at position 2 wherein the pseudo nucleotide may be an abasic pseudo-nucleotide analog and the modified nucleotide is optionally a mirror nucleotide..

[0278] In various embodiments the antisense strand (N)_x comprises 2'-O-Me modified ribonucleotides at the odd numbered positions (5' to 3'; positions 1, 3, 5, 7, 9, 11, 13, 15, 17, 19). In some embodiments (N)_x further comprises 2'-O-Me modified ribonucleotides at one or both positions 2 and 18. In other embodiments (N)_x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17, 19.

[0279] Other embodiments of Structures (L), (I) and (J) are envisaged wherein x=y=21 or wherein x=y=23; in these embodiments the modifications for (N')_y discussed above instead of being in positions 17 and 18 are in positions 19 and 20 for 21-mer oligonucleotide and 21 and 22 for 23 mer oligonucleotide; similarly the modifications in positions 15, 16, 17, 18 or 19 are in positions 17, 18, 19, 20 or 21 for the 21-mer oligonucleotide and positions 19, 20, 21, 22, or 23 for the 23-mer oligonucleotide. The 2'-OMe modifications on the antisense strand are similarly adjusted. In some embodiments (N)_x comprises 2'-OMe sugar modified ribonucleotides at the odd numbered positions (5' to 3'; positions 1, 3, 5, 7, 9, 12, 14, 16, 18, 20 for the 21 mer oligonucleotide [nucleotide at position 11 unmodified] and 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23 for the 23 mer oligonucleotide [nucleotide at position 12 unmodified]. In other embodiments (N)_x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 10, 12, 14, 16, 18, 20 [nucleotide at position 11 unmodified for the 21 mer oligonucleotide and at positions 2, 4, 6, 8, 10, 13, 15, 17, 19, 21, 23 for the 23 mer oligonucleotide [nucleotide at position 12 unmodified].

[0280] In some embodiments (N')_y further comprises a 5' terminal cap nucleotide. In various embodiments the terminal cap moiety is selected from an abasic pseudo-nucleotide analog, an inverted abasic pseudo-nucleotide analog, an L-DNA nucleotide, and a C6-imine phosphate (C6 amino linker with phosphate at terminus).

[0281] In other aspects the present description discloses a compound having Structure (M) set forth below:



wherein each of N and N' is selected from a pseudo-nucleotide and a nucleotide;

wherein each nucleotide is selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' are absent;


wherein x = 18 to 27;

wherein y = 18 to 27;

wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS: 1-58;

wherein at least one of N is selected from an abasic pseudo nucleotide, a non-pairing nucleotide analog and a nucleotide mismatch to the mRNA of a target gene in a position of (N)_x such that (N)_x comprises less than 15 consecutive nucleotides complementary to the mRNA of a target gene.

[0282] In other aspects the present description discloses a double stranded compound having Structure (N) set forth below:



wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;


wherein Z and Z' are absent;

wherein each of x and y is an integer between 18 and 40;

wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS: 1-58;

wherein (N)_x, (N')_y or (N)_x and (N')_y comprise non base-pairing modified nucleotides such that (N)_x and (N')_y form less than 15 base pairs in the double stranded compound.

[0283] In other aspects the present description discloses a compound having Structure (O) set forth below:



wherein each of N is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of N' is a nucleotide analog selected from a six membered sugar nucleotide, seven membered sugar nucleotide, morpholino moiety, peptide nucleic acid and combinations thereof;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' are absent;

wherein each of x and y is an integer between 18 and 40;

wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS: 1-58.

[0284] In other aspects the present description discloses a compound having Structure (P) set forth below:



3' Z'-(N')_y 5' (sense strand)

wherein each of N and N' is a nucleotide selected from an unmodified ribonucleotide, a modified ribonucleotide, an unmodified deoxyribonucleotide and a modified deoxyribonucleotide;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein Z and Z' are absent;

wherein each of x and y is an integer between 18 and 40;

wherein one of N or N' in an internal position of (N)_x or (N')_y or one or more of N or N' at a terminal position of (N)_x or (N')_y comprises an abasic moiety or a 2' modified nucleotide;

wherein the sequence of (N)_x is substantially complementary to the sequence of (N')_y; and wherein the sequence of (N)_x comprises an antisense sequence having substantial complementarity to about 18 to about 40 consecutive ribonucleotides in mRNA set forth in any one of SEQ ID NOS:1-58.

[0285] In various embodiments (N')_y comprises a modified nucleotide at position 15 wherein the modified nucleotide is selected from a mirror nucleotide and a deoxyribonucleotide.

[0286] In certain embodiments (N')_y further comprises a modified nucleotide at position 2 wherein the modified nucleotide is selected from a mirror nucleotide and an abasic pseudo-nucleotide analog.

[0287] In various embodiments the antisense strand (N)_x comprises 2'-O-Me modified ribonucleotides at the odd numbered positions (5' to 3'; positions 1, 3, 5, 7, 9, 11, 13, 15, 17, 19). In some embodiments (N)_x further comprises 2'-O-Me modified ribonucleotides at one or both positions 2 and 18. In other embodiments (N)_x comprises 2'-OMe sugar modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17, 19.

[0288] The Structural motifs (A)-(P) described above are useful with any oligonucleotide pair (sense and antisense strands) to a mammalian or non-mammalian gene. In some embodiments the mammalian gene is a human gene preferably selected from the genes provided in Tables A1-A4, with mRNA set forth in SEQ ID NOS:1-58. In certain preferred embodiments the sense and antisense oligonucleotides of the siRNA are selected from any one of siRNA pairs set forth in SEQ ID NOS:59-33,596. Table A5 below shows certain preferred sense and antisense oligonucleotide pairs.

Table A5

TARGET GENE	SENSE (N') _y 5'-3'	ANTISENSE (N) _x 5'-3'
P53	GAGAAUUAUUUCACCCUUA	UGAAGGGUGAAAUAUUCUC
CASP2	GCCAGAAUGUGGAACUCCU	AGGAGUCCACAUUCUGGC
RTP801	GUGCCAACCUGAUGCAGCU	AGCAGCAUCAGGUUGGCAC
RTP801	UACUGUAGCAUGAAACAAA	UUUGUUUAUGCUACAGUA
RTP801	CAGUACUGUAGCAUGAAAC	GUUUCUAGCUACAGUACUG
TP53BP2	CACCCAGAGAACAUUUUAU	AAUAAAUGUUCUCUGGGUG
CYBA	UGGGGACAGAAGUACAUGA	UCAUGUACUUCUGUCCCCA
RAC1	GAGUCCUGCAUCAUUUGAA	UUCAAAUGAUGCAGGACUC
SPP1	GUGCCAUACCAGUUAACA	UGUUUAACUGGUAUGGCAC
SPP1	GCAAAAUGAAAGAGAACA	AUGUUCUCUUUCAUUUUGC
ASPP1	CGAACUCAGAGAAAUGUAA	UUACAUUUCUCUGAGUUCG
ASPP1	GGAGAAAAACGUACUGAAA	UUUCAGUACGUUUUUCUCC
SOX9	CCUUCAUGAAGAUGACCGA	UCGGUCAUCUUAUGAAGG

[0289] For all the above Structures (A)-(P), in various embodiments x = y and each of x and y is 19, 20, 21, 22 or 23. In preferred embodiments, x=y=19. In additional embodiments the compound comprises modified ribonucleotides in alternating positions wherein each N at the 5' and 3' termini of (N)_x are modified in their sugar residues and the middle ribonucleotide is not modified, e.g. ribonucleotide in position 10 in a 19-mer strand, position 11 in a 21 mer and position 12 in a 23-mer strand.

[0290] In some embodiments where x = y = 21 or x = y = 23 the position of modifications in the 19 mer are adjusted for the 21 and

23 mers with the proviso that the middle nucleotide of the antisense strand is preferably not modified.

[0291] For all the above Structures (A)-(P), in some embodiments, neither (N)x nor (N'y are phosphorylated at the 3' and 5' termini. In other embodiments either or both (N)x and (N'y are phosphorylated at the 3' termini. In yet another embodiment, either or both (N)x and (N'y are phosphorylated at the 3' termini using non-cleavable phosphate groups. In yet another embodiment, either or both (N)x and (N'y are phosphorylated at the terminal 2' termini position using cleavable or non-cleavable phosphate groups. These particular siRNA compounds are also blunt ended and are non-phosphorylated at the termini; however, comparative experiments have shown that siRNA compounds phosphorylated at one or both of the 3'-termini have similar activity *in vivo* compared to the non-phosphorylated compounds.

[0292] For all the above Structures (A)-(P), in some embodiments, the compound is blunt ended, for example wherein both Z and Z' are absent. In an alternative embodiment, the compound comprises at least one 3' overhang, wherein at least one of Z or Z' is present. Z and Z' independently comprises one or more covalently linked modified or non-modified nucleotides, for example inverted dT or dA; dT, LNA, mirror nucleotide and the like. In some embodiments each of Z and Z' are independently selected from dT and dTdT. siRNA in which Z and/or Z' is present have similar activity and stability as siRNA in which Z and Z' are absent.

[0293] In certain embodiments for all the above-mentioned Structures, the compound comprises one or more locked nucleic acids (LNA) also defined as bridged nucleic acids or bicyclic nucleotides. Preferred locked nucleic acids are 2'-O, 4'-C-ethylene nucleosides (ENA) or 2'-O, 4'-C-methylene nucleosides. Other examples of LNA and ENA nucleotides are disclosed in WO 98/39352, WO 00/47599 and WO 99/14226.

[0294] In certain embodiments for all the above-mentioned Structures, the compound comprises one or more altritol monomers (nucleotides), also defined as 1,5 anhydro-2-deoxy-D-altritol-hexitol (see for example, Allart, et al., 1998. Nucleosides & Nucleotides 17:1523-1526; Herdewijn et al., 1999. Nucleosides & Nucleotides 18:1371-1376; Fisher et al., 2007, NAR 35(4):1064-1074).

[0295] The present invention explicitly excludes compounds in which each of N and /or N' is a deoxyribonucleotide (D-A, D-C, D-G, D-T). In certain embodiments (N)x and (N'y may comprise independently 1, 2, 3, 4, 5, 6, 7, 8, or 9 deoxyribonucleotides. In certain embodiments the present description discloses a compound wherein each of N is an unmodified ribonucleotide and the 3' terminal nucleotide or 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides at the 3' terminus of (N'y are deoxyribonucleotides. In yet other embodiments each of N is an unmodified ribonucleotide and the 5' terminal nucleotide or 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides at the 5' terminus of (N'y are deoxyribonucleotides. In further embodiments the 5' terminal nucleotide or 2, 3, 4, 5, 6, 7, 8, or 9 consecutive nucleotides at the 5' terminus and 1, 2, 3, 4, 5, or 6 consecutive nucleotides at the 3' termini of (N)x are deoxyribonucleotides and each of N' is an unmodified ribonucleotide. In yet further embodiments (N)x comprises unmodified ribonucleotides and 1 or 2, 3 or 4 consecutive deoxyribonucleotides independently at each of the 5' and 3' termini and 1 or 2, 3, 4, 5 or 6 consecutive deoxyribonucleotides in internal positions; and each of N' is an unmodified ribonucleotide. In certain embodiments the 3' terminal nucleotide or 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides at the 3' terminus of (N'y and the terminal 5' nucleotide or 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 consecutive nucleotides at the 5' terminus of (N)x are deoxyribonucleotides. The present invention excludes compounds in which each of N and/or N' is a deoxyribonucleotide. In some embodiments the 5' terminal nucleotide of N or 2 or 3 consecutive of N and 1, 2, or 3 of N' is a deoxyribonucleotide. Certain examples of active DNA/RNA siRNA chimeras are disclosed in US patent publication 2005/0004064, and Ui-Tei, 2008 (NAR 36(7):2136-2151).

[0296] Unless otherwise indicated, in preferred embodiments of the structures discussed herein the covalent bond between each consecutive N and N' is a phosphodiester bond.

[0297] An additional novel molecule disclosed by the present description is an oligonucleotide comprising consecutive nucleotides wherein a first segment of such nucleotides encode a first inhibitory RNA molecule, a second segment of such nucleotides encode a second inhibitory RNA molecule, and a third segment of such nucleotides encode a third inhibitory RNA molecule. Each of the first, the second and the third segment may comprise one strand of a double stranded RNA and the first, second and third segments may be joined together by a linker. Further, the oligonucleotide may comprise three double stranded segments joined together by one or more linker.

[0298] Thus, one molecule disclosed by the present description is an oligonucleotide comprising consecutive nucleotides which encode three inhibitory RNA molecules; said oligonucleotide may possess a triple stranded structure, such that three double stranded arms are linked together by one or more linker, such as any of the linkers presented hereinabove. This molecule forms a "star"-like structure, and may also be referred to herein as RNAAstar. Such structures are disclosed in PCT patent publication WO 2007/091269, assigned to the assignee of the present invention.

[0299] A covalent bond refers to an internucleotide linkage linking one nucleotide monomer to an adjacent nucleotide monomer. A covalent bond includes for example, a phosphodiester bond, a phosphorothioate bond, a P-alkoxy bond, a P-carboxy bond and the like. The normal internucleoside linkage of RNA and DNA is a 3' to 5' phosphodiester linkage. In certain preferred embodiments a covalent bond is a phosphodiester bond. Covalent bond encompasses non-phosphorous-containing internucleoside linkages, such as those disclosed in WO 2004/041924 *inter alia*. Unless otherwise indicated, in preferred embodiments of the structures discussed herein the covalent bond between each consecutive N and N' is a phosphodiester bond.

[0300] For all of the structures above, in some embodiments the oligonucleotide sequence of (N)x is fully complementary to the oligonucleotide sequence of (N')y. In other embodiments (N)x and (N')y are substantially complementary. In certain embodiments (N)x is fully complementary to a target sequence. In other embodiments (N)x is substantially complementary to a target sequence.

[0301] In some embodiments, neither (N)x nor (N')y are phosphorylated at the 3' and 5' termini. In other embodiments either or both (N)x and (N')y are phosphorylated at the 3' termini (3' Pi). In yet another embodiment, either or both (N)x and (N')y are phosphorylated at the 3' termini with non-cleavable phosphate groups. In yet another embodiment, either or both (N)x and (N')y are phosphorylated at the terminal 2' termini position using cleavable or non-cleavable phosphate groups. Further, the inhibitory nucleic acid molecules of the present invention may comprise one or more gaps and/or one or more nicks and/or one or more mismatches. Without wishing to be bound by theory, gaps, nicks and mismatches have the advantage of partially destabilizing the nucleic acid / siRNA, so that it may be more easily processed by endogenous cellular machinery such as DICER, DROSHA or RISC into its inhibitory components.

[0302] In the context of the present invention, a gap in a nucleic acid refers to the absence of one or more internal nucleotides in one strand, while a nick in a nucleic acid refers to the absence of an internucleotide linkage between two adjacent nucleotides in one strand. Any of the molecules of the present invention may contain one or more gaps and/or one or more nicks.

[0303] The structures disclosed herein, when integrated into antisense and corresponding sense nucleic acid sequences to any target gene, provides siRNA compound useful in reducing expression of that target gene. The target gene is a mammalian or non-mammalian gene. The methods of the invention comprise topically and non-invasively administering to the eye of the subject one or more siRNA compounds which inhibit expression of a target gene in the eye of the subject.

siRNA Synthesis

[0304] The compounds of the present invention can be synthesized by any of the methods that are well-known in the art for synthesis of ribonucleic (or deoxyribonucleic) oligonucleotides. Such synthesis is, among others, described in Beaucage and Iyer, *Tetrahedron* 1992; 48:2223-2311; Beaucage and Iyer, *Tetrahedron* 1993; 49: 6123-6194 and Caruthers, et. al., *Methods Enzymol.* 1987; 154: 287-313; the synthesis of thioates is, among others, described in Eckstein, *Annu. Rev. Biochem.* 1985; 54: 367-402, the synthesis of RNA molecules is described in Sproat, in *Humana Press* 2005 edited by Herdewijn P.; Kap. 2: 17-31 and respective downstream processes are, among others, described in Pingoud et. al., in *IRL Press* 1989 edited by Oliver; Kap. 7: 183-208.

[0305] Other synthetic procedures are known in the art e.g. the procedures as described in Usman et al., *J. Am. Chem. Soc.*, 1987, 109:7845; Scaringe et al., *NAR*, 1990, 18:5433; Wincott et al., *NAR* 1995, 23:2677-2684; and Wincott et al., *Methods Mol. Bio.*, 1997, 74:59, and these procedures may make use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. The modified (e.g. 2'-O-methylated) nucleotides and unmodified nucleotides are incorporated as desired.

[0306] The oligonucleotides of the present invention can be synthesized separately and joined together post-synthetically, for example, by ligation (Moore et al., *Science* 1992, 256:9923; International Patent Publication No. WO 93/23569; Shabarova et al., *NAR* 1991, 19:4247; Bellon et al., *Nucleosides & Nucleotides*, 1997, 16:951; Bellon et al., *Bioconjugate Chem* 1997, 8:204), or by hybridization following synthesis and/or deprotection.

[0307] It is noted that a commercially available machine (available, *inter alia*, from Applied Biosystems) can be used; the oligonucleotides are prepared according to the sequences disclosed herein. Overlapping pairs of chemically synthesized fragments can be ligated using methods well known in the art (e.g., see US Patent No. 6,121,426). The strands are synthesized separately and then are annealed to each other in the tube. Then, the double-stranded siRNAs are separated from the single-stranded oligonucleotides that were not annealed (e.g. because of the excess of one of them) by HPLC. In relation to the siRNAs

or siRNA fragments of the present invention, two or more such sequences can be synthesized and linked together for use in the present invention.

[0308] The compounds of the invention can also be synthesized via tandem synthesis methodology, as described for example in US Patent Publication No. 2004/0019001 (McSwiggen), and in PCT Patent Publication No. WO 2007/091269 (assigned to the assignee of the present invention) wherein both siRNA strands are synthesized as a single contiguous oligonucleotide fragment or strand separated by a cleavable linker which is subsequently cleaved to provide separate siRNA fragments or strands that hybridize and permit purification of the siRNA duplex. The linker can be a polynucleotide linker or a non-nucleotide linker.

[0309] The present invention further provides for a pharmaceutical composition comprising two or more siRNA molecules for the treatment of any of the diseases and conditions mentioned herein, whereby said two molecules may be physically mixed together in the pharmaceutical composition in amounts which generate equal or otherwise beneficial activity, or may be covalently or non-covalently bound, or joined together by a nucleic acid linker of a length ranging from 2-100, preferably 2-50 or 2-30 nucleotides.

[0310] Thus, the siRNA molecules may be covalently or non-covalently bound or joined by a linker to form a tandem siRNA compound. Such tandem siRNA compounds comprising two siRNA sequences are typically about 38-150 nucleotides in length, more preferably 38 or 40-60 nucleotides in length, and longer accordingly if more than two siRNA sequences are included in the tandem molecule. A longer tandem compound comprised of two or more longer sequences which encode siRNA produced via internal cellular processing, e.g., long dsRNAs, is also envisaged, as is a tandem molecule encoding two or more shRNAs. Such tandem molecules are also considered to be a part of the present invention. A tandem compound comprising two or more siRNAs sequences of the invention is envisaged.

[0311] Additionally, the siRNA disclosed herein or any nucleic acid molecule comprising or encoding such siRNA can be linked or bound (covalently or non-covalently) to antibodies (including aptamer molecules) against cell surface internalizable molecules expressed on the target cells, in order to achieve enhanced targeting for treatment of the diseases disclosed herein. For example, anti-Fas antibody (preferably a neutralizing antibody) may be combined (covalently or non-covalently) with any of the siRNA compounds.

[0312] The compounds of the present invention can be delivered either directly or with viral or non-viral vectors. When delivered directly the sequences are generally rendered nuclease resistant. Alternatively the sequences can be incorporated into expression cassettes or constructs such that the sequence is expressed in the cell as discussed herein below. Generally the construct contains the proper regulatory sequence or promoter to allow the sequence to be expressed in the targeted cell. Vectors optionally used for delivery of the compounds of the present invention are commercially available, and may be modified for the purpose of delivery of the compounds of the present invention by methods known to one of skill in the art.

[0313] It is also envisaged that a long oligonucleotide (typically 25-500 nucleotides in length) comprising one or more stem and loop structures, where stem regions comprise the sequences of the oligonucleotides of the invention, may be delivered in a carrier, preferably a pharmaceutically acceptable carrier, and may be processed intracellularly by endogenous cellular complexes (e.g. by DROSHA and DICER as described above) to produce one or more smaller double stranded oligonucleotides (siRNAs) which are oligonucleotides of the invention. This oligonucleotide can be termed a tandem shRNA construct. It is envisaged that this long oligonucleotide is a single stranded oligonucleotide comprising one or more stem and loop structures, wherein each stem region comprises a sense and corresponding antisense siRNA sequence of the genes of the invention.

RNA interference

[0314] A number of PCT applications have recently been published that relate to the RNAi phenomenon. These include: PCT publication WO 00/44895; PCT publication WO 00/49035; PCT publication WO 00/63364; PCT publication WO 01/36641; PCT publication WO 01/36646; PCT publication WO 99/32619; PCT publication WO 00/44914; PCT publication WO 01/29058; and PCT publication WO 01/75164.

[0315] RNA interference (RNAi) is based on the ability of dsRNA species to enter a cytoplasmic protein complex, where it is then targeted to the complementary cellular RNA and specifically degrade it. The RNA interference response features an endonuclease complex containing an siRNA, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having a sequence complementary to the antisense strand of the siRNA duplex. Cleavage of the target RNA may take place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir et al., *Genes Dev.*, 2001, 15(2):188-200). In more detail, longer dsRNAs are digested into short (17-29 bp) dsRNA fragments (also

referred to as short inhibitory RNAs, "siRNAs") by type III RNases (DICER, DROSHA, etc.; Bernstein et al., Nature, 2001, 409(6818):363-6; Lee et al., Nature, 2003, 425(6956):415-9). The RISC protein complex recognizes these fragments and complementary mRNA. The whole process is culminated by endonuclease cleavage of target mRNA (McManus & Sharp, Nature Rev Genet, 2002, 3(10):737-47; Paddison & Hannon, Curr Opin Mol Ther. 2003, 5(3):217-24). (For additional information on these terms and proposed mechanisms, see for example Bernstein et al., RNA 2001, 7(11):1509-21; Nishikura, Cell 2001, 107(4):415-8 and PCT publication WO 01/36646).

[0316] Several groups have described the development of DNA-based vectors capable of generating siRNA within cells. The method generally involves transcription of short hairpin RNAs that are efficiently processed to form siRNAs within cells (Paddison et al. PNAS USA 2002, 99:1443-1448; Paddison et al. Genes & Dev 2002, 16:948-958; Sui et al. PNAS USA 2002, 8:5515-5520; and Brummelkamp et al. Science 2002, 296:550-553). These reports describe methods to generate siRNAs capable of specifically targeting numerous endogenously and exogenously expressed genes.

[0317] The invention has been described in an illustrative manner, and it is to be understood that the terminology used is intended to be in the nature of words of description rather than of limitation.

[0318] Obviously, many modifications and variations of the present invention are possible in light of the above teachings. It is, therefore, to be understood that within the scope of the appended claims, the invention can be practiced otherwise than as specifically described.

[0319] Throughout this application, various publications, including United States Patents, are referenced by author and year and patents by number. The disclosures of these publications and patents and patent applications describe the state of the art to which this invention pertains.

[0320] The present invention is illustrated in detail below with reference to examples, but is not to be construed as being limited thereto.

[0321] Citation of any document herein is not intended as an admission that such document is pertinent prior art, or considered material to the patentability of any claim of the present application. Any statement as to content or a date of any document is based on the information available to applicant at the time of filing and does not constitute an admission as to the correctness of such a statement.

EXAMPLES

[0322] Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the claimed invention in any way.

[0323] Standard molecular biology protocols known in the art not specifically described herein are generally followed essentially as in Sambrook et al., Molecular cloning: A laboratory manual, Cold Springs Harbor Laboratory, New-York (1989, 1992), and in Ausubel et al., Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, Maryland (1988), and as in Ausubel et al., Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, Maryland (1989) and as in Perbal, A Practical Guide to Molecular Cloning, John Wiley & Sons, New York (1988), and as in Watson et al., Recombinant DNA, Scientific American Books, New York and in Birren et al (eds) Genome Analysis: A Laboratory Manual Series, Vols. 1-4 Cold Spring Harbor Laboratory Press, New York (1998) and methodology as set forth in US Patent Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057. Polymerase chain reaction (PCR) was carried out as in standard PCR Protocols: A Guide To Methods And Applications, Academic Press, San Diego, CA (1990). In situ PCR in combination with Flow Cytometry (FACS) can be used for detection of cells containing specific DNA and mRNA sequences (Testoni et al., Blood 1996, 87:3822.) Methods of performing RT-PCR are well known in the art.

Cell culture

[0324] HeLa cells (American Type Culture Collection) were cultured as described in Czauderna, et al. (NAR, 2003. 31:670-82). Human keratinocytes were cultured at 37°C in Dulbecco's modified Eagle medium (DMEM) containing 10% FCS. The mouse cell line, B16V (American Type Culture Collection), was cultured at 37°C in Dulbecco's modified Eagle medium (DMEM) containing

10% FCS. Culture conditions were as described in (Methods Find Exp Clin Pharmacol. 1997, 19(4):231-9).

[0325] In each case, the cells were subject to the experiments as described herein at a density of about 50,000 cells per well and the double-stranded nucleic acid according to the present invention was added at a concentration of 20 nM, whereby the double-stranded nucleic acid was complexed using 1 µg/ml of a proprietary lipid as described below.

[0326] In the histochemical/microscopic figures, arrows were added to draw attention to staining of the tissue.

Animal Models

Model systems of glaucoma and of Retinal Ganglion Cells (RGC) death

ONC model in rats

[0327] Various animal models are useful for studying the effect of siRNA therapeutics in treating glaucoma. In the optic nerve crush model in rats the orbital optic nerve (ON) of anesthetized rats is exposed through a supraorbital approach, the meninges severed and all axons in the ON transected by crushing with forceps for 10 seconds, 2 mm from the lamina cribrosa. Testing active inhibitors of the invention (such as siRNA) for treating or preventing glaucoma is performed, for example, in the animal models described by Pease et al. (J. Glaucoma, 2006, 15(6):512-9. Manometric calibration and comparison of TonoLab and TonoPen tonometers in rats with experimental glaucoma and in normal mice).

[0328] Optic nerve crush (ONC) model in adult Wistar rats is also an accepted model for studying Retinal Ganglion Cells (RGC) death. The onset and kinetics of RGC death in this model are very reproducible; RGC apoptosis begins on day 4-5 after the ONC; massive RGC loss (about 50-60%) is observed on days 7-10 after the ONC; and 95% of the RGC loss is occurs by week 3-4 after the ONC. This model allows for establishment of the neuroprotective efficacy of test drugs *in vivo*.

[0329] In some non-limiting examples, siRNA compounds directed to genes shown in Tables A1-A4 are tested in this animal model which show that these siRNA compounds treat and/or prevent glaucoma and/or RGC death when topically and non-invasively delivered to the eye.

IOP model in rats

[0330] Intraocular pressure (IOP) is a measurement of the fluid pressure inside the eye. This fluid, called aqueous humor, is circulated and then drained out via specialized outflow pathways. If the drainage system does not function properly, as in prevalent forms of glaucoma, pressure inside the eye builds up. A model of ocular hypertension in Brown Norway rats developed by Dr. J. Morrison and collaborators at the Casey Eye Institute (Portland, Oregon) is used in this study. The Morrison model involves injection of hypertonic saline into an episcleral vein, leading to blockade of the aqueous humor outflow pathways. This procedure leads to gradual increase of eye pressure and progressive death of RGCs. Importantly, inner retinal atrophy, optic nerve degeneration, and optic nerve head remodeling observed in this model are similar to that seen in human glaucoma. Thus, the Morrison model is considered the best pre-clinical rodent model of glaucoma.

In vivo axotomy model in rats

[0331] In this model RGC apoptosis is induced by axotomy of the optic nerve (ON) in adult Sprague-Dawley rats. The onset and kinetics of RGC death in this model system are very reproducible and allow for the establishment of the neuroprotective efficacy of non-invasively administered siRNA compound *in vivo*. Using this method, the time course of RGC death follows a predictable course: cell death begins on day 5 and proceeds to the rapid loss of more than 90% of these neurons by 2 weeks.

Vehicle Formulations and Exemplary Eye Drop Formulations

[0332] The aqueous eye drop formulation optionally contain various additives incorporated ordinarily, such as buffering agents (e.g., phosphate buffers, borate buffers, citrate buffers, tartarate buffers, acetate buffers, amino acids, sodium acetate, sodium citrate and the like), isotonicities (e.g., saccharides such as sorbitol, glucose and mannitol, polyhydric alcohols such as glycerin, concentrated glycerin, polyethylene glycol and propylene glycol, salts such as sodium chloride), preservatives or antiseptics (e.g., benzalkonium chloride, benzethonium chloride, p-oxybenzoates such as methyl p-oxybenzoate or ethyl p-oxybenzoate, benzyl alcohol, phenethyl alcohol, sorbic acid or its salts, thimerosal, chlorobutanol and the like), solubilizing aids or stabilizing agents (e.g., cyclodextrins and their derivative, water-soluble polymers such as polyvinyl pyrrolidone) surfactants such as polysorbate 80 (Tween 80)), pH modifiers (e.g., hydrochloric acid, acetic acid, phosphoric acid, sodium hydroxide, potassium hydroxide, ammonium hydroxide and the like), chelating agents (e.g., sodium edetate, sodium citrate, condensed sodium phosphate) and the like.

[0333] The eye drop formulation in the form of an aqueous suspension may also contain suspending agents (e.g., polyvinyl pyrrolidone, glycerin monostearate) and dispersing agents (e.g., surfactants such as tyloxapol and polysorbate 80, ionic polymers such as sodium alginate), in addition to the additives listed above, thereby ensuring that the eye drop formulation is a further uniform microparticulate and satisfactorily dispersed aqueous suspension.

[0334] The ophthalmic ointment may comprise a known ointment base, such as purified lanolin, petrolatum, plastibase, liquid paraffin, polyethylene glycol and the like.

Exemplary eye drop formulation 1:

[0335] Formulation of siRNA compounds in PBS ("naked siRNA formulation") is typically prepared by dissolving dry siRNA in PBS. The formulation comprises at least one siRNA compound typically present in an amount ranging from about 5 µg/µl to about 60 µg/µl by volume of the composition.

[0336] In a non-limiting example formulation of an siRNA compound in PBS was prepared as follows: Under sterile conditions, 500 mg of dry siRNA were dissolved in 25 ml of sterile double distilled water (DDW), to achieve a clear 20 mg/ml solution. The solution was stored at -80°C until use. The 20 mg/ml stock solution in DDW was then brought to a working concentration of 100 µg/3µl, in PBS, as follows: 325 µl of 20 mg/ml stock of siRNA solution (6.5mg) were precipitates by 0.15M NaCl and EtOH, and dried under a tissue culture laminar (sterile conditions). 6.5 mg of dry siRNA were dissolved in 130µl PBS to form eye drop formulation 1.

Exemplary eye drop formulation 2:

[0337] In some embodiments at least one siRNA compound is formulated in tris(hydroxymethyl)aminomethane (TRIS) 1M, pH 8.0 (available e.g. from Sigma (catalog #T-1503)).

[0338] In a non-limiting example 121.1 g of TRIS base (Sigma # T1503) were dissolved in 700 ml of ddH₂O. Desired pH 8 was achieved with addition of concentrated HCl. DDW was added to have final 1L solution. Under sterile conditions, 42.426 mg of siRNA compound powder were dissolved in 2.1 ml of sterile double distilled water, to achieve a clear 20 mg/ml (1.5 mM) stock solution. The stock solution was stored at -80°C until use. Under sterile conditions, corresponding amounts of siRNA stock were lyophilized and re-suspended in corresponding amount of 1M TRIS pH 8.

Exemplary eye drop formulation 3:

[0339] The concentration of the viscosity-enhancing agents to be used is typically within the range of about 0.05 to about 5.0 % (w/v), and more desirably, from about 0.1 to about 3.0 % (w/v). One preferred ophthalmic composition comprises from about 0.01 to about 0.075 % (w/v) of an active ingredient; from about 0.15 to about 2.5 % (w/v) of polyoxyl 40 stearate; from about 0.1 to about 3.0 % (w/v) of a cellulose thickening agent selected from 25 hydroxypropyl methylcellulose or hydroxyethylcellulose; and from about 0.0001 % to about 0.01 % (w/v) of an anti-oxidant selected from butylated hydroxytoluene or sodium thiosulfate.

[0340] The combination of the surfactants such as polyoxyl 40 stearate or polyoxyethylene cetyl ether or polyoxyethylene octylphenyl ether with siRNA yields an eye-drop preparation causing less irritation to the eyes, providing better distribution in the eyes and having higher stability. The inclusion of an anti-oxidant and a cellulose-thickening agent further improves the distribution

of the active therapeutic agent (i.e. siRNA) in the eyes and the stability.

[0341] Other excipients may also be added, such as for example an isotonic agent, buffer, preservative, and or pH-controlling agent. Sterile purified water in appropriate amounts is present to obtain the desired eye-drop preparation.

[0342] The pH of the ophthalmic composition is within the range, which is normally used for ophthalmic preparations, as known in the art, but is desirably within the range of 5 to 8. The formulation further comprises at least one siRNA compound typically present in an amount ranging from about 6.6 µg/µl to about 50 µg/µl by volume of the composition.

Exemplary eye drop formulation 4:

[0343] In the topical delivery of drugs to the lens, drug retention on the eye surface is considered to be important. Without wishing to be bound to theory, increased retention on the ocular surface leads to increased ocular absorption of a drug through the cornea into the aqueous humor and subsequently the lens.

[0344] In some embodiments a topical suspension is preferred. Non-limiting examples of topical suspensions include:

1. (1) hydroxypropyl methylcellulose (HPMC, 0.5% w/v), (2) xanthan gum (0.5% w/v), (3) gellan gum (0.5% w/v), (4) carbopol (0.25% w/v), and (5) carbopol (0.25% w/v) - hydroxypropyl methylcellulose (HPMC) (0.25% w/v). Viscosity measurements are conducted with a viscometer.

[0345] The formulation further comprises at least one siRNA compound typically present in an amount ranging from about 5 µg/µl to about 60 µg/µl by volume of the composition.

Exemplary eye drop formulation 5:

[0346] In ophthalmic compositions, a chelating agent is optionally used to enhance preservative effectiveness. Suitable chelating agents are those known in the art, and, while not intending to be limiting, edetate (EDTA) salts like edetate disodium, edetate calcium disodium, edetate sodium, edetate trisodium, and edetate dipotassium are examples of useful chelating agents. It is understood that EDTA refers to a species having four carboxylic acid functional groups, and that these carboxylic acid groups may be protonated or deprotonated (i.e. in the salt form) depending upon the pH of the composition it is in.

[0347] Buffers are commonly used to adjust the pH to a desirable range for ophthalmic use. Generally, a pH of around 5-8 is desired, however, this may need to be adjusted due to considerations such as the stability or solubility of the therapeutically active agent or other excipients.

[0348] Another commonly used excipient in ophthalmic compositions is a viscosity-enhancing, or a thickening agent. Thickening agents are used for a variety of reasons, ranging from improving the form of the formulation for convenient administration to improving the contact with the eye to improve bioavailability. The viscosity-enhancing agent comprises a polymer containing hydrophilic groups such as monosaccharides, polysaccharides, ethylene oxide groups, hydroxyl groups, carboxylic acids or other charged functional groups. While not intending to limit some examples useful viscosity-enhancing agents are sodium carboxymethylcellulose, hydroxypropylmethylcellulose, povidone, polyvinyl alcohol, and polyethylene glycol..

[0349] The formulation further comprises at least one siRNA compound typically present in an amount ranging from about 5 µg/µl to about 60 µg/µl by volume of the composition.

Exemplary eye drop formulation 6 ("Formulation A"):

Formulation "A": Prepare 2 solutions, A and B.

[0350] Preparation of Solution A: 4% solution of methylcellulose in water. 0.4 g of methylcellulose 25 (ScienceLab.com, Cat#

SLM2050) dissolved in the final volume of 10ml of apyrogenic water for injections (WFI) in 50 mL sterile tube (Norbook, Cat# 7082-51). Put small sterile stirrer inside and close the cap. Keep in boiling water bath while stirring for at least 5 minutes until methylcellulose forms an opalescent solution.

[0351] Preparation of Solution B: 2% Glycerol; 0.02% (v/v) EDTA solution. Add 333.3 μ L of 60% glycerol (Sigma, Cat# G6279) & 2 μ L 0.5M EDTA, pH=8 (Sigma, Cat# E9884) (final - 100 μ M) to 9.665 ml WFI (Norbook, Cat# 7082-51). Final volume - 10 mL.

[0352] Prepare 2X (in regard to working concentration) solution of siRNA in Solution B. Note: if stock siRNA volume is substantially high, the final volume of Solution B is better kept lower than 10mL to allow volume adjustment later in 2X siRNA solution.

Preparation of working siRNA solution:

[0353] Cool solution A in hand ~ till 40-50°C (still liquid) and mix the desired volume with an equal volume of 2X solution of siRNA in Solution B.

[0354] Final siRNA formulation contains 2% methylcellulose, 1% glycerol and 0.01% (v/v) (50 μ M) EDTA in WFI. Final pH should be ~7.4 and osmolarity - similar to human tear film. Concentration of siRNA in final formulation is 33.3 mg/ml.

[0355] Must be prepared fresh once a week and aliquoted into the portions for daily usage. The aliquots are kept at 4°C. Before application, the dose is warmed for 20-30 min at room temperature.

[0356] The final concentration of the methylcellulose is about 0.1% to about 3% w/v about 0.1% to about 2% w/v, about 0.1% - 1.5% w/v methylcellulose, preferably about 2% w/v. The final concentration of glycerol is about 0.1% to about 5% v/v, about 0.5% to about 2%, preferably about 1% v/v. The final concentration of EDTA is about 0.001% to about 0.05%, about 0.005% to about 0.01%, preferably about 0.01% w/v.

Exemplary eye drop formulation 7:

[0357] In some embodiments siRNA is formulated in a commercially available lubricant eye drop formulation. Non-limiting example of commercially available lubricant eye drop formulation is Systane® available from Alcon Inc. Such commercially available lubricant eye drop formulations typically comprise polyethylene glycol and/ or propylene glycol, an antiseptic and / or antiviral agent such as boric acid, a gelling agent such as hydroxypropyl guar, potassium chloride and / or sodium chloride and / or magnesium chloride and/or calcium chloride and/or zinc chloride, preservatives such as Polyquad® and purified water. Under sterile conditions, 300 mg of siRNA powder is dissolved in 15 ml of sterile double distilled water, to achieve a clear 20 mg/ml solution. The solution is stored at (-)80°C until use. The 20 mg/ml stock solution in double distilled water is then brought to a working concentration of 100 μ g/3 μ L, in formulation, as follows:

To obtain a final formulated quantity of 1.7 mg siRNA: 85 μ L of 20 mg/ml stock of siRNA (1.7 mg), is precipitated by 0.15M NaCl and EtOH, dried under the tissue culture laminar (sterile conditions).

[0358] To achieve a siRNA solution of 33.3 mg/ml 1.7 mg dry siRNA is dissolved in 51 μ L Systane®.

siRNA Compounds Induce Knockdown (KD) of Target Genes

Example 1: *In Vitro* Testing of siRNA Compounds

[0359] About 1.5-2x10⁵ tested cells (HeLa cells and/or 293T cells for siRNA targeting human genes and NRK52 (normal rat kidney proximal tubule cells) cells and/or NMuMG cells (mouse mammary epithelial cell line) for siRNA targeting the rat/mouse gene) were seeded per well in 6 wells plate (70-80% confluent).

[0360] About 24 hours later, cells were transfected with siRNA compounds using the Lipofectamine™ 2000 reagent (Invitrogen) at final concentrations of 5 nM or 20 nM. The cells were incubated at 37°C in a CO₂ incubator for 72h.

[0361] As positive control for transfection phosphatase and tensin homolog (PTEN)-Cy3 labeled siRNA compounds were used. Various chemically modified blunt ended siRNA compounds having alternating modified and unmodified ribonucleotides (modified at the 2' position of the sugar residue in both the antisense and the sense strands, wherein the moiety at the 2' position is methoxy) and wherein the ribonucleotides at the 5' and 3' termini of the antisense strand are modified in their sugar residues, and the ribonucleotides at the 5' and 3' termini of the sense strand are unmodified in their sugar residues were tested. Another siRNA compound comprised a blunt ended structure having an antisense with an alternating pattern of methoxy moieties and a sense strand with three ribonucleotides linked by two 2'5' bridges at the 3' terminus; and another siRNA compound comprising antisense and sense strands having three ribonucleotides linked by 2'5' bridges at the 3' terminus was used. Some of the tested compounds comprised a blunt ended structure having an antisense with an alternating pattern of methoxy moieties and a sense strand with one or two L-nucleotides at the 3' terminal or 3' penultimate positions.

[0362] GFP siRNA compounds were used as negative control for siRNA activity. At 72h after transfection cells were harvested and RNA was extracted from cells. Transfection efficiency was tested by fluorescent microscopy. The percent of inhibition of gene expression using specific preferred siRNA structures was determined using qPCR analysis of a target gene in cells expressing the endogenous gene.

[0363] In general, the siRNAs having specific sequences that were selected for *in vitro* testing were specific for human and a second species such as non-human primate, rat or rabbit genes. Similar results are obtained using siRNAs having these RNA sequences and modified as described herein.

[0364] siRNAs targeted at genes related to apoptosis and neuroprotection were tested by a similar procedure and found active and able to induce knock down of their corresponding target genes.

[0365] siRNA Prepared for Non-Invasive Administration and Administered by Eye Drops is Delivered to the Target Retinal Tissue *in-vivo*

[0366] Example 2: Distribution of CY3-labeled siRNA in murine optic tissue (siRNA Formulated in PBS) as monitored by fluorescence microscopy and confocal microscopy The current study showed siRNA delivery *in-vivo* to the lacrimal gland and ocular structures via a topical ocular route.

Abbreviations: E.D.= Eye Drops; ISH= In-situ Hybridization

[0367] The objective of this study (for illustrative purposes) was to test the delivery of DDIT4-Cy3 labeled siRNA (siRNA compound targeting the RTP801 gene) to the lacrimal gland via non-invasive topical ocular route. The topical ocular route has been evaluated for the delivery of siRNA into the lacrimal gland, anterior chamber of the eye, the retina and the optic nerve.

Materials and Methods

[0368]

Table C1: Animals

Species/strain:	Mice/ ICR, males
Age:	6-10 weeks
Body Weight Range:	27-32 gr
Group Size:	2
Total number of animals:	26

[0369] DDIT4-Cy3 siRNA (siRNA against mouse RTP801) formulated as 20m/ml stock solution in PBS and stored at -80°C until use.

Experimental Procedure

[0370] The study included 6 experimental groups as described in Table C2 hereinbelow. Mice were treated with a single siRNA DDIT4-Cy3 as follows:

Group V (A and B): dose regime: 50µg/mouse/3µl/eye, administration route: Right Eye drops (E.D.);

Group Am-V-A, Am3-VIII, Am3-IX: dose regime: 50µg/mouse/3µl/eye, administration route: both eyes, Eye drops (E.D.);

Group VI-A: dose regime: 20µg/mouse/3µl/eye, administration route: Right eye, Eye drops (E.D.);

Group VII, Am-VII: non-treated control.

Table C2. Study Design

Group	SiRNA Type	Dose µg/mouse	Injected Volume (µl)	Route	Time point (hrs)	Group Size
V-A	DDIT4-Cy3	50.00 Right Eye	3.00	E.D.	1	2
Am-V-A	DDIT4-Cy3	50.00 per eye / Both Eyes	3.00	E.D.	1	2
V-B	DDIT4-Cy3	50.00 Right Eye	3.00	E.D.	4	2
VI-A	DDIT4-Cy3	20.00 Right Eye	3.00	E.D.	1	2
VII	None	none	None	none		5
Am-VII	None	none	None	none		1
Am3-VIII	DDIT4-Cy3	50.00 per eye / Both Eyes	3.00	E.D.	4	2
Am3-IX	DDIT4-Cy3	50.00 per eye / Both Eyes	3.00	E.D.	24	2

[0371] The siRNA doses were prepared under sterile conditions. siRNA aliquots were thawed for at least 30 min at room temperature prior to administration. Total amount per aliquot (per each mouse) included an additional 20% of the calculated volume. The designated siRNA dose was delivered in 3µl (Experimental groups V-VI) volume per eye.

[0372] Anesthesia: Mice were anesthetized with Ketamine/Xylazine mix as follows: 0.85 ml Ketamine + 0.15 ml Xylazine + 0.9 ml Saline, 0.1 ml mix solution/20 gr body weight (BW).

[0373] Eye drop (E.D.) delivery: A 3 µl sample volume was slowly dropped in the treated eye (corneal surface) using a blunt pipette tip; the animals were placed in a warm environment to prevent anesthesia-induced hypothermia, and were returned to cages after regaining consciousness.

[0374] Scheduled euthanasia: Following treatment, mice from all groups were euthanized according to the study design (Table C2, Time point termination).

[0375] Termination step was accomplished by cardiac puncture and blood collection; collected serum/plasma was stored (-20°C) for further siRNA blood detection analysis.

[0376] Tissue Collection: Left and right eyes including optic nerve, lacrimal glands (left and right) from all animals were excised, brain (front and middle parts of the brain, frontal oriented sections) from animals of groups, V-A one from group VII, Am3-VIII, Am3-IX were excised, embedded in optimal cutting temperature (OCT) compound and sectioned with a cryostat into 8 µm slices that were placed on microscope slides (Superfrost/Plus™). Sections were fixed in 95% EtOH for 7 minutes then counter stained with DAPI (1 µg/ml), incubated for 1 minute in absolute EtOH, incubated for 5 minutes in xylene, air dried and mounted.

[0377] Tissue Evaluation: Delivery of siRNA was evaluated using light microscopy and digital imaging. A tissue fragment was considered positive (i.e., successful Cy3 DDIT4 siRNA intracellular incorporation) only where histological (microscopic) examination showed a clear fluorescence signal within specific cells or structures such as acinar alone or in combination with ductal cells of the lacrimal gland. Background DAPI staining assisted in identification of tissue structure: in the case of anterior

eye chamber structures such as cornea, angle, ciliary body or posterior eye chamber such as retina (retinal pigmentum epithelium cells (RPE), retinal ganglion cells (RGC)). Delivery was considered positive if histological examination would show that all animals within one experimental group showed the same fluorescence signal within the investigated cell type/tissue. (i.e. time points or route administration).

Results & Discussion

[0378] All cryosections were analyzed under light microscopy (bright field, BF) and fluorescence confocal microscopy. The fluorescent signal was visualized in the following tissues:

Retina by E.D. delivery (retinal pigment epithelial cells, retinal ganglion cells) Figures 1A-1B: Representative images of Cy3 labeled DDIT4 siRNA incorporated into mice retina.

[0379] Figure 1A: Fluorescent microscopy (upper panel, magnification x40) of the retina 1 hour after E.D. administration. Right arrow identifies Cy3 stained retinal ganglion cells (RGC), left arrow identifies Cy3 stained retinal pigmentum epithelium cells (RPE). Confocal microscopy (lower panel, magnification x60) of the retinal ganglion cells layer. The lower set of 9 views show Cy3 labeled tissue (Left), bright filed (BF, center) and a merge of the two (M, right). Arrows point to the labeled RGCs.

[0380] Figures in 1B: Confocal microscopy of the retina 4 hours after E.D. administration (upper x40, lower x60). Cy3 staining of the RGC is prominent.

[0381] *Eye drop delivery of siRNA to the lacrimal glands (acinar cells, ductal cells)* Figures 2A-2D: Representative images of Cy3 labeled DDIT4 siRNA incorporated into ductal and acinar cells of the murine lacrimal gland.

[0382] Figure 2A: Fluorescent microscopy of the lacrimal gland 1 hour after E.D. administration (M=merged DAPI-Cy3 staining). Center arrow points to acinar cells, right arrow indicates ductal epithelial cells, the gray left arrow points to the lacrimal duct (magnification x60)

[0383] Figure 2B: Confocal microscopy of the lacrimal gland 1 hour after E.D administration (upper x40 and lower panel x60). Arrow in merged image M is pointing to lacrimal duct.

[0384] Figure 2C: Confocal microscopy of the lacrimal gland 1 hour E.D administration (x40). Arrows are pointing to acinar cells

[0385] Figure 2D: Confocal microscopy of the lacrimal gland 4 hours after E.D. (x60).

[0386] Figures 3A-3C show time course of accumulation of Cy3-siRNA in rat choroid after administration by eye drops, following 1 and 4 hours post administration. Choroid, outer nuclear layer, RPE and outer segment layer of photoreceptor cells show Cy3 staining.

[0387] Figure 4 shows Cy3-siRNA delivery to the trabecular meshwork and ciliary body one hour after administration by eye drops.

[0388] Different chemically modified siRNA (structural motives) were tested with mouse Cy3-labeled siRNA. Similar tissue distribution to that shown above was observed for all structures tested: Structure B having alternating natural ribonucleotides and 2'O-Me sugar modified ribonucleotides on both strands; Structure I having alternating natural ribonucleotides and 2'O-Me sugar modified ribonucleotides on the antisense strand and three ribonucleotides linked by 2'5' internucleotide linkages at the 3' end of the sense strand; and a structure having three ribonucleotides linked by 2'5' internucleotide linkages at the 3' end of each of the antisense and sense strands.

Example 3 (for illustrative purposes): Examination of Topical Ocular Route Delivery of Formulated siRNA in Rats (siRNA Formulated in PBS or in methyl cellulose 2% ("Formulation A"))

[0389] In the current study the ocular delivery of DDIT4-Cy3 siRNA formulated either in PBS or in Formulation A and applied by Eye Drops was studied in rats.

Table C3: Study Design

Group	siRNA	Dose µg/eye	Route of Adm. (Bilateral)	Adm. Volume (µl)	Formulation	Termination time	Group Size
I	DDIT4_1-Cy3	100.00	E.D.	3.00	Formulated	30 min.	1
Ia	DDIT4_1-Cy3	100.00	E.D.	3.00	PBS	30 min.	1
II	DDIT4_1-Cy3	100.00	E.D.	3.00	Formulated	1 hour	1
IIa	DDIT4_1-Cy3	100.00	E.D.	3.00	PBS	1 hour	1
III	DDIT4_1-Cy3	100.00	E.D.	3.00	Formulated	3 hours	1
IIIa	DDIT4_1-Cy3	100.00	E.D.	3.00	PBS	3 hours	1
IV	DDIT4_1-Cy3	100.00	E.D.	3.00	Formulated	6 hours	1
Iva	DDIT4_1-Cy3	100.00	E.D.	3.00	PBS	6 hours	1
V	DDIT4_1-Cy3	100.00	E.D.	3.00	Formulated	24 hours	1
Va	DDIT4_1-Cy3	100.00	E.D.	3.00	PBS	24 hours	1
VI	N/A	N/A	E.D.	3.00	vehicle	1 hours	1
VII	N/A Intact control	N/A	N/A	N/A	N/A	N/A	1

Experimental Design:**[0390]**

- 1) Group I, II, III, IV, V: dose regime: siRNA formulated in Formulation A; 100µg/rat/3µl/eye, administration route: Bilateral Eye drops (E.D.);
- 2) Group Ia, IIa, IIIa, IVa, Va: dose regime: siRNA formulated in PBS; 100µg/rat/3µl/eye, administration route: Bilateral Eye drops (E.D.);
- 3) Group VI: control group treated with the formulation vehicle of Formulation A (no siRNA); 100µg/rat/3µl/eye, administration route: Bilateral Eye drops (E.D.);
- 4) Group VII: intact control

[0391] The siRNA doses were prepared under sterile conditions. siRNA aliquots were thawed for at least 30 minutes at room temperature prior to administration. Total amount per aliquot (per each mouse) included an additional 20% of the calculated volume. The designated siRNA dose was delivered in 3µl volume per eye.

[0392] Anesthesia: Rats from groups I-VII were anesthetized with Equithesine (4ml/kg)

[0393] Termination step: At study termination, animals were deeply anesthetized by Equithesine (4ml/kg). Thereafter, rats were intracortically perfused with fresh 10% neutral buffered formalin using peristaltic pump standard regime.

[0394] Tissue Collection: Left and right eyes including optic nerve from all animals were enucleated, perforated and post fixed in 10% neutral buffered formalin for an additional 1 hour in room temperature with slow rotation, cryoprotected step-wise by sucrose gradient at 4°C overnight, following cryoprotection in optimal cutting temperature (OCT) compound at 4°C overnight with rotation, embedded in OCT compound and sectioned with a cryostat into 12 µm slices that were placed on microscope slides (Superfrost/Plus). Sections were counter stained with DAPI (1 µg/ml) and mounted.

[0395] Delivery of siRNA was evaluated using light microscopy and digital imaging. A tissue

[0396] fragment was considered positive (i.e., a successful Cy3 DDIT4_1 siRNA transfer in anterior eye chamber, retina and optic nerve delivery was considered positive) if histological examination would show that all animals within one experimental group showed the same fluorescent signal within the specific tissue/cell type.

[0397] Results: All cryosections were analyzed under light microscopy (bright field, BF) and fluorescence confocal microscopy. The fluorescent signal was visualized in retinal pigment epithelial cells and retinal ganglion cells. Notably, compounds formulated in PBS reached maximum distribution in ocular tissues 3 hours after administration, and were cleared from said tissues at the 6 hour time point. However, compounds formulated in Formulation A were maximally distributed in ocular tissues at 24 hours after administration.

Example 4: Ocular Distribution of siRNA (formulated in PBS) in the Cynomolgus Monkey (for illustrative purposes)

[0398]

Table C4: Experimental Design

Group	Animal number	Compound	Dose Volume	Termination after administration	Processing at termination
					Eyes
1	101	Cy3-DDIT4_1	500ug/20ul/eye	1 hour	4% PFA
2	251	DDIT4_1	500ug/20ul/eye	1 hour	Snap frozen in liquid nitrogen
3	351	Cy3-DDIT4_1	500ug/20ul/eye	5 hours	4% PFA
4	401	DDIT4_1	500ug/20ul/eye	5 hours	Snap frozen in liquid nitrogen
ED=Eye Drops, topically					

[0399] The aim of this study was to investigate the tissue distribution of siRNA formulated in PBS following single topical ocular administration to the surface of the eye of the Cynomolgus Macaca fascicularis monkey. All tissues were snap frozen in triplicate ~1 to ~2 g samples or as a whole, as appropriate.

RNA purification

[0400] Frozen samples were grinded to a fine powder under liquid nitrogen. A small amount of the powdered tissue was used for RNA extraction. The rest of the tissues were stored at - 80°C for further usage. PolyA RNA was extracted from each sample with MicroPoly(A)Purist mRNA purification Kit (#1919) and processed according to manufacturer's protocol for poly(A)RNA isolation from tissues or cells. The final yields and spectral characteristics of the RNA are summarized below in Table C5.

Table C5: polyA RNA from monkey

Tissue	260/280	260/230	Concentration µg/µl	Yield µg
Lymph Node	1.58	0.92	0.104	1.55
Spleen	1.74	1.27	0.175	2.62

[0401] One µg from the spleen and 0.7µg from the lymph node polyA RNA preparations were used for cDNA synthesis using random primer. Casp2 gene was successfully amplified from both cDNA preparations.

[0402] Results are summarized below in Table C6.

Table C6: Results

Treatment	Tissue analysed	Termination	Av	SD	Group size
ED(REDD14)	Neuronal Retina	1h	0.2	[+/-0]	1 (2 eyes)
ED(REDD14)	Neuronal Retina	5h	0.05	[+/- 0.03]	1 (2 eyes)
ED(REDD14)	RPE	1h	0.62	[+/-0.05]	1 (2 eyes)
ED(REDD14)	RPE	5h	0.1	[+/- 0.07]	1 (2 eyes)
ED(REDD14)	optic nerve	1h	8.9	[+/-2.6]	1 (2 eyes)
ED(REDD14):	optic nerve	5h	3	[+/-1.9]	1 (2 eyes)

[0403] Conclusion: SiRNA formulated in PBS following single topical ocular administration to the surface of the eye of the Cynomolgus Macaca fascicularis monkey reaches the neuronal retina and can be measured by the qPCR S&L method.

Example 5: Delivery of siRNA (formulated in TRIS) to target retinal tissue in mice

[0404] Examination of topical delivery of siRNA molecules labeled with different fluorophores.

[0405] Topical delivery to Mice of 20 ug of different fluorescently-labeled siRNAs formulated in tris(hydroxymethyl)aminomethane (TRIS) 1M, pH 8, 1 hr post-administration.

[0406] siRNAs tested:

Cy3-QM5; Cy3-DDIT4; Cy3.5-DDIT4; DDIT4_1 Dy-649/C6; FITC-CNL_1; 3'Cy3-CNL_1; Cy3-Casp2_4 L-DNA; Cy3-AS-Casp2_4; TGASEII-FAM; HNOEL-FAM. (QM5 is a rat/mouse siRNA that targets p53; CNL is control scrambled siRNA)

[0407] Exemplary chemically modified Casp2 siRNA compounds were as follows:

1. (comparative siRNA). Sense: GCCAGAAUGUGGAACUC2pC2pU, 17 and 18 are 2'-5'-bridge
Antisense: mAGmGAmGUmUCmCAmCAmUUmCUmGGmC-cy3
2. (comparative siRNA). Sense: GCCAGAAUGUGGAACUC;LdC;LdT 18 and 19 are L-DNA
Antisense: mAGmGAmGUmUCmCAmCAmUUmCUmGGmC-cy3
3. 3.Sense: GCCAGAAUGUGGAACUC;LdC;U18 is an L-DNA moiety
Antisense: mAGmGAmGUmUCmCAmCAmUUmCUmGGmC-cy3

[0408] LdC, LdT refer to L-DNA nucleotides, mA, mC, Mg, Mu refer to 2'-O-methoxy ribonucleotides.

[0409] Formulated (formulated compound): siRNA in 1M Tris pH 8.0 Of the following siRNA's: QM5-Cy3#1, QM5-Cy3#10, DDIT4_1-Cy3, DDIT4_1Cy3.5, Redd14(DDIT4_1) Dy-649/C6, FITC-CNL_1RD/CNL_1FD, scarmbled3'Cy3CNL 1, Cy3-AS-CASP2_4-Struc-L-DNA-s(2 residues at 3' of sense)-plus-alt AS, Cy3AS- CASP2_4-Struc2-5-s(2 residues at 3' of sense)-plus-alt AS, TGASEII-FAM, HNOEL-FAM.

Table C7: Formulations

siRNA	IDO	Final	siRNA Stock	Preparation
QM5-Cy3#1	116937	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8
QM5-Cy3#10	116938	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8
DDIT4_1-Cy3	117821	40µg	20mg/ml	2ul stock were lyophilized and resuspended in 7µl 1M Tris pH8
DDIT4_1-Cy3.5	112206	40µg	20mg/ml	2ul stock were lyophilized and resuspended in 7µl 1M Tris pH8
Redd14(DDIT4_1) Dy-649/C6	114029	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8
FITC-CNL_1RD/CNL_1 FD	110357	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8
scarmbled3'Cy3-CNL_1	110910	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8
Cy3-AS-CASP2_4-Struc-L-DNA-s(2 residues at 3' of sense)-plus-alt AS	122994	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8

siRNA	IDO	Final	siRNA Stock	Preparation
Cy3-AS-CASP2_4-Struc2-5-s(2 residues at 3' of sense)-plus-alt AS	122979	40µg	20mg/ml	2µl stock were lyophilized and resuspended in 7µl 1M Tris pH8
TGASEII-FAM	75755	11.62µg	0.75µg/µl	15.5µl stock were lyophilized and resuspended in 2µl 1M Tris pH8
HNOEL-FAM	75757	4.48µg	1.4µg/µl	3.2µl stock were lyophilized and resuspended in 0.784µl 1M Tris pH8
DDIT4_1-Cy3	117821	40µg	20mg/ml	2µl stock+0.7ul PBSx10+4.3µl DDW

[0410] Description of the test material: Under sterile conditions, corresponding amount of siRNA stock were lyophilized and re-suspended in corresponding amount of 1M Tris pH8 (see Table C7).

[0411] Quantity supplied: One vile for each tested siRNA (Synthesized by Biospring, AG)

[0412] Storage Conditions: frozen until use. Prior to use, samples were thawed and kept at room temperature for 30 minutes.

CONTROL ARTICLE(S)

[0413] Positive Control - DDIT4_1-Cy3 in PBS, Batch # : 117821

[0414] Description of the test material: Under sterile conditions, 42.426 mg of DDIT4_1-Cy3 powder (BioSpring) were dissolved in 2.1 ml of sterile double distilled water, to achieve a clear 20 mg/ml (1.5mM) solution. The solution was stored at -80°C until use.

[0415] Vehicle - 1M Tris pH 8.0 - Outsourcing from Sigma (catalog #T-1503). 121.1 g of TRIS base (Sigma # T1503) were dissolved in 700 ml of ddH₂O. Desire pH 8 was achieved with concentrated HCl. DDW was added to have final 1L solution.

[0416] Vehicle - PBSX10 - Outsourcing from Biological Industries (catalog #02-023-5A; (For 10x PBS) Batch # :619113)

TEST SYSTEM

[0417] Animals:

Species: Mice; Strain: RTP-801WT; CMF-608WT

Source: Harlan Laboratories, Jerusalem, Israel.

Age: 8-12 weeks; Body Weight Range: 17-28 gr

Sex: males; Group Size: 1; Total number of animals: 14

[0418] Animal Husbandry: Diet: Animals were provided an *ad libitum* commercial rodent diet and free access to drinking water. Environment: (i) Acclimatization of at least 5 days. (ii) All the animals were confined in a limited access facility with environmentally-controlled housing conditions throughout the entire study period, and maintained in accordance with approved standard operating procedures (SOPs).

EXPERIMENTAL DESIGN

[0419] General: The study included 14 experimental groups as described in Table C8:

Experimental groups 1-12 siRNA delivered by eye drops, group 13 the vehicle [TRIS 1M pH8 treated control delivered by eye drops and group 14 (non-treated control). Mice were treated with a single siRNA as follows:

- Groups 1-9 and 12: dose regime: 20 µg/eye/3µl/siRNA, group 10: dose regime 5.81 µg/eye/3µl/siRNA, group 11: dose regime 2.24 µg/eye/3µl/siRNA administration route: Eye Drop (E.D.);
- Group 13: vehicle (TRIS 1M pH8) 3µl/eye control
- Group 14: none treated control

Table C8. Study Design

Group	SiRNA type/bilateral	Dose µg/eye	Volume (µl)	Route	Time point (hrs)	Group Size
1	QM5-Cy3#1	20	3.00	E.D.	1	1
2	QM5-Cy3#10	20	3.00	E.D.	1	1
3	DDIT4_1-Cy3	20	3.00	E.D.	1	1
4	DDIT4_1-Cy3.5	20	3.00	E.D.	1	1
5	DDIT4_1 Dy-649/C6	20	3.00	E.D.	1	1
6	FITC-CNL_1RD/CNL_1FD	20	3.00	E.D.	1	1
7	scrambled3'cy3-CNL_1	20	3.00	E.D.	1	1
8	Cy3-AS-CASP2_4-Struc-L-DNA-s(2 residues at 3' of sense)-plus-alt AS	20	3.00	E.D.	1	1
9	CY3-AS-CASP2_4-Struc2-5-s(2 residues at 3' of sense)-plus-alt AS	20	3.00	E.D.	1	1
10	TGASEII-FAM	5.81	3.00	E.D.	1	1
11	HNOEL-FAM	2.24	3.00	E.D.	1	1
12	DDIT4_1-Cy3 in PBS vehicle (positive control)	20	3.00	E.D.	1	1
13	Vehicle (TRIS 1M pH8) (negative control)	N/A	3.00	E.D.	1	1
14	Non treated (negative control)	N/A	3.00	E.D.	1	1

[0420] The siRNA doses for delivery were prepared under sterile conditions and stored at -20°C. siRNA aliquots were thawed for at least 30 minutes at room temperature prior to administration. Total amount per aliquot (per each eye) included an additional 20% of the calculated volume. The designated siRNA dose was delivered in 3.5.µl (Experimental groups 1 to 9 and 12) volume per eye. The designated siRNA dose was delivered in 3.2.µl (Experimental groups 10 and 11) volume per eye.

[0421] Anesthesia: Mice were anesthetized with Ketamine/Xylazine mixture as follows (0.85ml Ketamine+0.15ml Xylazine + 0.9ml Saline, 0.1ml mix solution/20gr BW).

[0422] Eye drop delivery: A 3µl volume sample was slowly dropped in each eye (corneal surface) by blunt pipette tip; the animals were placed in a warm environment to prevent anesthesia-induced hypothermia, and were returned to its cage after it regaining consciousness.

[0423] Scheduled euthanasia: Mice from all groups were terminated according to the study design (Table C8, Time points termination).

[0424] Termination step was accomplished by cervical dislocation.

[0425] Tissue Collection: Left and right eyes including optic nerve, from all animals were enucleated, the retina was transversal dissected with the blade, the lens/vitreous were gently removed. The remaining eye cup was fixed with 4% PFA (in PBS pH 7.4) for 30 minutes, then infiltrated with 30% sucrose for 3 hours at 4°C. Washed 3X5 minutes with ice cold PBS pH7.4. It was then embedded in optimal cutting temperature (OCT) compound and sectioned longitudinal with a cryostat into 4 µm slices that were placed on microscope slides (Superfrost/Plus). Sections were counter stained with DAPI (1 µg/ml) and mounted.

[0426] A whole blood drop from all animals, was smeared on the slide with the cover slip, and covered. It was then glued with nail

polish

EVALUATION

[0427] Delivery of siRNA was evaluated using light microscopy and digital imaging. A tissue fragment was considered positive (i.e., a successful siRNA transfer\ intracellular incorporation occurred) only if histological (microscopic) examination showed clear fluorescence signal within specific cells or structures of the anterior or posterior chamber, and retina. Background DAPI staining was assisted in identification of tissue structure. Delivery was considered positive if histological examination was consistent within the group (bilateral identical cell type or structural staining), i.e. the histological examination would show the same fluorescent signal in all cell types/tissues (anterior eye chamber, retina and optic nerve) of the same experimental group.

Results

[0428] Figures 5A-5B: Confocal microscopy (magnification x60) of the retina 1 hour after eye drop administration of QM5 - siRNA targeted at the p53 gene. SiRNA in retina (retinal pigment epithelial cells, retinal ganglion cells) is shown by Cy3 fluorescence.

[0429] Figures 6A-6B are representative images of Cy3 labeled DDIT4 siRNA incorporated into mice retina.

[0430] Figure 6B shows accumulation of DDIT4_1 Cy3-siRNA 1 hour post administration by eye drops. Choroid, outer nuclear layer, RPE and outer segment layer of photoreceptor cells show Cy3 staining.

[0431] Figure 7 shows accumulation of DDIT4_1 Dy-649/C6 -siRNA 1 hour post administration by eye drops in RGC cells by use of Dy-649/C6 staining.

[0432] Figure 8A - 8C represent control siRNA FITC-CNL_1RD/CNL_1FD and scrambled3'cy3-CNL_1 delivery to retinal tissues by different staining methods. Figure 9A and 9B show delivery of different structures of Casp2 to the mouse retinal tissues.

[0433] Figures 10A and 10B show TGASEII-FAM and HNOEL-FAM delivery.

[0434] Figure 11 shows retinal delivery of siRNA against p53 in PBS as positive control group and

[0435] Figures 12A and 12B show that in intact animals or when administering ED without siRNA no fluorescent signal is obtained in the retina.

[0436] Conclusion: No sequence dependent differences, neither fluorophore nor gene dependent differences in the delivery to the retina were found. The study showed efficient delivery of Cy3 conjugated siRNA compounds directed to CASP2, RTP801, TIGASEII and p53 target genes, to ocular structures: RGC, RPE, photoreceptors, choroid.

Example 6: Quantification of Delivery of Casp2 siRNA (formulated in PBS or in MC 2% (Formulation "A")) to target retinal tissue in rats *in-vivo*

[0437] Examination of different eye Drop formulations for the delivery of siRNA to the retina. The objective of the study was the determination of CASP2_4_S510 siRNA quantity in rat retina at time points 1, 3, 6 & 24 hours following single topical administration in different eye drops (ED) formulations.

TEST ARTICLE

[0438] Substance (unformulated compound): CASP2_4_S510 (siRNA Against CASP2)

[0439] Supplied by Agilent (Manufacturer's catalog # QPI-1007, Batch #: Q02F08002N)

[0440] Description of the test material: CASP2_4_S510: S- inverted-Abasic 5'-cap, L-DNA 18/AS-AL. A 19-mer chemically

modified blunt-ended duplex having two separate strands, with a sense strand (SEN) comprising unmodified ribonucleotides (upper case letters), an L-deoxyribonucleotide at position 18 (bold, underlined) and inverted deoxybasic moiety (iB) present at the 5' terminus of the SEN strand; and with an antisense strand (AS) comprising unmodified ribonucleotides (upper case letters), and 2'OMe sugar modified ribonucleotides (lower case letters) at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 as shown in Formula I:

Formula I

SEN 5' iB-GCCAGAAUGUGGAACUCCU 3'

AS 3' cGgUcUuAcACcUuGaGgA 5'

Storage Conditions: -80°C

[0441] siRNA formulated in PBS : 33.3mg/ml CASP2_4_S510 in PBS (solution for eye drops)

[0442] Description of the test material: Under sterile conditions, 300 mg dry CASP2_4_S510 siRNA were dissolved in 15ml of sterile double distilled water, to achieve a clear 20 mg/ml solution. The solution was stored at -80°C until use. The 20mg/ml stock solution in DDW was then brought to a working concentration of 100µg/ 3µl, in PBS, as follows:

6.5mg CASP2_4_S510 siRNA: 325 µl of 20 mg/ml stock of CASP2_4 (6.5mg), were precipitated by 0.15M NaCl and EtOH, and dried under the tissue culture laminar (sterile conditions). siRNA solution 33.3mg/ml: 6.5mg dry siRNA were dissolved in 195µl PBsX1 Quantity prepared: 6.5mg / 195µl aliquoted into 4 tubes.

Storage Conditions: freshly prepared

[0443] siRNA formulated in Formulation A : 33.3mg/ml solution of CASP2_4_S510 siRNA in 2% (w/v) methylcellulose & 1% (v/v) sterile glycerol & 0.01% (w/v) EDTA solution in pyrogen free water (solution for eye drops)

[0444] Description of the test material: Under sterile conditions, 300 mg dry CASP2_4_S510 siRNA were dissolved in 15ml of sterile double distilled water, to achieve a clear 20 mg/ml solution. The solution was stored at -80°C until use. The 20mg/ml stock solution in DDW was brought to a working concentration of 100µg/3µl, in formulation, as follows:

6.5mg CASP2_4_S510 siRNA: 325 µl of 20 mg/ml stock of CASP2_4 (6.5mg), were precipitated by 0.15M NaCl and EtOH, and dried under the tissue culture laminar (sterile conditions).

[0445] siRNA solution 33.3mg/ml: 6.5mg dry siRNA were dissolved in 195µl Formulation A solution

[0446] Quantity prepared: 6.5mg / 195µl aliquoted into 4 tubes

Storage Conditions: freshly prepared

CONTROL ARTICLE(S)

[0447] Methyl cellulose formulation (no siRNA) - 2% methylcellulose & 1% v/v sterile glycerol & 0.01% w/v EDTA solution in pyrogen free water, was prepared as follows:

Solution A: 0.4 g of methylcellulose 25 (ScienceLab.com, Cat# SLM2050) dissolved in final volume of 10ml of hot boiled water (80-90°C), (Norbrook), cooled down to room temperature and added in a proportion of 1:1 (final concentration 2%) to solution B.

Solution B: 332 µl of 60% glycerol (Sigma, Cat# G6279) & 2 µl EDTA solution pH8 (prepared from Sigma, Cat# E9884) in 9.66 ml WFI (Norbrook).

[0448] 500 µL of Solution A mixed with 500 µL of Solution B to obtain: 2% methylcellulose & 1% v/v sterile Glycerol & 0.01% w/v EDTA solution in pyrogen free water (final pH was approximately 7.4 and osmolarity similar to human tear film).

[0449] PBS was supplied by Biological industries (Manufacturer catalog #02-023-5A (For 10x PBS); Batch # 619113)

TEST SYSTEM

[0450] Animals used: Species: Rats; Strain: Adult, Sprague-Dawley (SD)

Source: Harlan, Jerusalem Israel

Age: 8-10 weeks; Body Weight Range: 180-250gr

Sex: Male; Group Size: n=6/3; Total number of animals: 54

Animal Husbandry: Diet: Animals were provided an *ad libitum* commercial rodent diet and free access to drinking water.

Environment:

[0451]

1. (i) Acclimatization of at least 5 days.
2. (ii) All animals were confined in a limited access facility with environmentally-controlled housing conditions throughout the entire study period, and maintained in accordance with approved standard operating procedures (SOPs). Animals were provided *ad libitum* a commercial rodent diet (Harlan Teklad 2018S Global 18% Protein Rodent Diet) and filtered, chlorinated and acidified water.

[0452] Rats were kept in microisolator cages with filter top, 1-6 Rats / cage. The cages were maintained under controlled environmental- conditions of temperature (20-24 °C), relative humidity (30-70%), a 12- hr light / 12-hr dark cycle, monitored by the control computer, throughout the study period.

Experimental design

[0453] Study design: Retinal concentration of CASP2_4 siRNA following ED application of ocular formulation was determined by qPCR in study groups terminated at different time points after siRNA administration. Each experimental siRNA-treated group included 6 rats. ED was applied bilaterally as detailed in Study Design Table C9.

[0454] Termination time points post siRNA administration: Termination setup was performed according to the study design Table C9. Dosing and termination of experimental groups were performed on separate days.

Table C9: Study Design

Group Number	Group Size	Test Article	Dose of Bilateral ED [100 µg/eye/3ul]	Termination [hrs]	Method of Analysis
1	6	CASP2_4_S510	PBS formulated siRNA	1	qPCR
2	6	CASP2_4_S510	MC formulated	1	qPCR
3	6	CASP2_4_S510	PBS formulated siRNA	3	qPCR
4	6	CASP2_4_S510	MC formulated	3	qPCR
5	6	CASP2_4_S510	PBS formulated siRNA	6	qPCR
6	6	CASP2_4_S510	MC formulated	6	qPCR
7	6	CASP2_4_S510	PBS formulated siRNA	24	qPCR
8	6	CASP2_4_S510	MC formulated	24	qPCR
9	3	N/A	Vehicle MC	1	qPCR

Group Number	Group Size	Test Article	Dose of Bilateral ED [100 µg/eye/3ul]	Termination [hrs]	Method of Analysis
10	3	N/A	intact	N/A	qPCR

[0455] Anesthesia: In the course of the experiment, the animals were anesthetized with Equithesine (I.P. 4ml/kg) for ED application and/or before terminations.

[0456] Eye Drop Delivery: A 3µl sample volume of the test article or vehicle was applied to the corneal surface bilaterally to the anesthetized animals, by a blunt pipette tip (filter tips 10µl sterile (short)). The animals were placed in a warm environment to prevent anesthesia-induced hypothermia, and were returned to their cage after regaining consciousness (experimental Groups 7 and 8). Termination of animals from all study groups was according to study design. Tissues were collected according to study design.

[0457] Scheduled euthanasia: All animals were deeply anaesthetized and euthanized according to the study design (Table C9, Termination).

[0458] Perfusion setup: Rats were perfused transcardially with PBS (20-50 ml/min following 2-3 min).

[0459] Tissue Collection: After perfusion, both eyes (left and right) were enucleated and stored on ice. The eyes were dissected using a microscope, and optionally gross pathology was graded according to the sample grading scale. The cornea was dissected by a cut along the limbus, lens was gently removed, and the retina and vitreous were carefully separated from the sclera. Whole retinas and vitreous bodies (humor) were collected (Retina including: "neural retina" + Retinal Pigment Epithelium + Choroid) into two separate appropriate and properly marked test tubes. Dissected retinas were washed in a large volume of PBS (each retina in a separate tube with fresh PBS), extra liquid was removed with Kimwipe® and retinas were snap-frozen in liquid nitrogen. Retina and Vitreous body samples were subjected to RNA extraction.

EVALUATION

RNA Extraction:

[0460] Retina: RNA was extracted from each retina sample individually (left and right) by double extraction. The RNA was transferred for cDNA preparation and qPCR analyses.

[0461] Vitreous: Material for Casp2_4 siRNA detection from vitreous was obtained using the following protocol:

Each rat vitreous was around 10µl. 500µl of reagent A EZ-RNA II (Biological Industries Cat no. 20-410-100) were added to each vitreous. The sample was homogenized, and 10µg tRNA (1µl from 10 mg/ml stock) were added. The sample was stored for 5 minutes at room temperature. Subsequently : EZ-RNA II B: 400 µl and EZ-RNA II C: 90 µl were added and Mixed well. The sample was stored for 10 min at room temperature then centrifuged 12000g for 15 min at 4°C.

[0462] The upper phase was transferred to a fresh tube to which isopropanol: 500µl and Linear Acrylamide: 5µl were added. The sample was stored overnight at -20°C, centrifuged at 12000g for 20 min at 4°C, washed twice with 75% ethanol. The pellet was dissolved in 15 µl H₂O.

[0463] siRNA Quantification: The quantity of the siRNA in retinas and vitreous humor (siRNA quantification was performed per vitreous) was examined by qPCR siRNA quantification. qPCR was performed according to Quark's standard operating procedures. CASP2_4 siRNA and reference gene expression were tested. Results are summarized in Table C10.

Table C10: Results

Termination	Formulation	Eye	Group size	fmol siRNA/µg retinal RNA
1 hrs	Methyl Cellulose	Right	5	12.50
		Left	6	10.19

Termination	Formulation	Eye	Group size	fmol siRNA/ μ g retinal RNA
3 hrs	PBS	Right	6	8.08
		Left	6	4.77
	Methyl Cellulose	Right	5	1.76
		Left	5	2.47
6 hrs	PBS	Right	6	6.48
		Left	6	7.80
	Methyl Cellulose	Right	4	1.37
		Left	6	3.80
24 hrs	PBS	Right	6	3.61
		Left	6	1.04
	Methyl Cellulose	Right	6	0.11
		Left	6	0.21
	PBS	Right	6	0.03
		Left	6	0.04

[0464] Conclusion: The study showed efficient delivery of siRNA into retina 1 and 3 hours post administration. qPCR analysis showed that after application of 100 μ g siRNA in Formulation A, in average 11 fMol/ μ g of total RNA was obtained in the retina after 1 hour and 2 fMol/ μ g siRNA after 3 hours. When administering the same concentration in PBS, an amount of 6 fMol/ μ g of siRNA was obtained after 1 hour and 8 fMol/ μ g of tot RNA was obtained after 3 hours.

Example 7: Quantification of non-invasively administered siRNA compound (formulated in various formulations) in target retinal tissue in rats.

[0465] Examination of different eye drop formulations for non-invasive delivery of siRNA to the retina in vivo.

[0466] The objective of this study was the assessment of CASP2_4_S510 siRNA quantity in normal rat retina at 3 hours following single topical application of different siRNA formulations prepared for administration as eye drops (ED).

Test Article

[0467] Substance (unformulated compound) CASP2_4_S510 (siRNA Against CASP2) Supplied by Agilent (Manufacturer's catalog # QPI-1007; Batch #: Q02F08002N).

[0468] Description of the test material: CASP2_4_S510: S- inverted-Abasic 5'-cap, L-DNA 18/AS-AL. A 19-mer stabilized double strand RNA with inverted Abasic as 5'-cap, L-DNA at the 18 position of the sense strand and alternating 2'-OMe at positions 2, 4, 6, 8, 11, 13, 15, 17 & 19 on the antisense strand.

Storage Conditions: -80°C

[0469] siRNA formulated in PBS 33.3mg/ml CASP2_4_S510 in PBS (solution for eye drops) -

[0470] Description of the test material: Under sterile conditions, 300 mg dry CASP2_4_S510 siRNA were dissolved in 15ml of sterile double distilled water, to achieve a clear 20 mg/ml solution. The solution was stored at -80°C until use. The 20mg/ml stock solution in double distilled water was then brought to a working concentration of 100 μ g/ 3 μ l, in PBS, as follows:

1.7 mg CASP2_4_S510 siRNA: 85 μ l of 20 mg/ml stock of CASP2_4 (1.7 mg), were precipitated by 0.15M NaCl and EtOH, and dried under the tissue culture laminar (sterile conditions).

siRNA solution 33.3 mg/ml: 1.7 mg dry siRNA were dissolved in PBS x1 to achieve 51µl Quantity prepared: 1 tube of 1.7 mg in 51µl of PBS Storage Conditions: freshly prepared

siRNA formulated in commercial lubricant solution Systane® : 33.3mg/ml CASP2_4_S510 in Systane (solution for eye drops) - Group II. Purchased from Alcon Inc.; Batch #: Lot 165228F.

[0471] Description of the test material: Under sterile conditions, 300 mg of CASP2_4_S510 powder (Aligent, batch # Q02F08002N) were dissolved in 15 ml of sterile double distilled water, to achieve a clear 20 mg/ml solution. The solution was stored at -80°C until use. The 20 mg/ml stock solution in double distilled water was then brought to a working concentration of 100 µg/ 3µl, in formulation, as follows:

1.7 mg CASP2_4_S510 siRNA: 85 µl of 20 mg/ml stock of CASP2_4_S510 (1.7 mg), was precipitated by 0.15M NaCl and EtOH, and dried under the tissue culture laminar (sterile conditions).

[0472] siRNA solution 33.3 mg/ml: 1.7 mg dry siRNA was dissolved in 51µL commercial lubricant solution (i.e. Systane®).

[0473] Quantity prepared: 1 vial of 1.7 mg siRNA in 51µL Systane® solution.

[0474] SiRNA formulated in glycerol +EDTA solution and in methyl cellulose (MC) Solutions: 33.3 mg/ml solution of CASP2_4_S510 siRNA in Glycerol + EDTA formulation, and in 0.5%, 2% or 3% (w/v) methylcellulose & 1% (v/v) sterile glycerol & 0.01% (w/v) EDTA solution in pyrogen free water- Groups III - VI

[0475] Description of the test material: Under sterile conditions, 300 mg of CASP2_4_S510 powder (Aligent, batch # Q02F08002N) were dissolved in 15 ml of sterile double distilled water, to achieve a clear 20 mg/ml solution. The solution was stored at -80°C until use. The 20 mg/ml stock solution in double distilled water was then brought to a working concentration of 100µg/ 3µl, in formulation, as follows:

Seven vials of 1.7 mg CASP2_4_S510 siRNA: 85 µl of 20 mg/ml stock of CASP2_4_S510 (1.7 mg), was precipitated by 0.15M NaCl and EtOH, and dried under the tissue culture laminar (sterile conditions).

[0476] Group III: 0.5% MC formulation: 1:7 mixture of solution A and solution B [12.5µL of Solution A were mixed with 50 µL of Solution B and 37.5 µl WFI to obtain: 0.5% methylcellulose & 1% v/v sterile Glycerol & 0.01% w/v EDTA solution in pyrogen free water (final pH was approximately 7.4 and osmolarity similar to human tear film)]

[0477] siRNA formulated in 0.5% MC formulation 33.3 mg/ml: 1.7 mg dry siRNA were dissolved in 0.5% MC formulation that was prepared as described to achieve a 51µL formulated solution of siRNA in 0.5% MC formulation.

[0478] Group IV: 1% MC formulation: 1:3 mixture of solution A and solution B [25 µL of Solution A were mixed with 50 µL of Solution B and 25 µl of WFI to obtain: 1% methylcellulose & 1% v/v sterile Glycerol & 0.01% w/v EDTA solution in pyrogen free water (final pH was approximately 7.4 and osmolarity similar to human tear film)]

[0479] siRNA solution formulated in 1% MC formulation 33.3 mg/ml: 1.7 mg dry siRNA were dissolved in 1% MC formulation that was prepared as described to achieve a 51µL formulated solution of siRNA in 1% MC formulation.

[0480] Group V: 2% MC formulation: 1:1 mixture of solution A and solution B [50µL of Solution A were mixed with 50 µL of Solution B to obtain: 2% methylcellulose & 1% v/v sterile Glycerol & 0.01% w/v EDTA solution in pyrogen free water (final pH was approximately 7.4 and osmolarity similar to human tear film)]

[0481] siRNA solution formulated in 2% MC formulation 33.3 mg/ml: 1.7 mg dry siRNA were dissolved in 2% MC formulation that was prepared as described to achieve a 51µL formulated solution of siRNA in 2% MC formulation.

[0482] Group VI: 3% MC formulation: 1:1 mixture of solution A and solution C [50 µL of Solution A were mixed with 50 µL of Solution C to obtain: 3% methylcellulose & 1% v/v sterile Glycerol & 0.01% w/v EDTA solution in pyrogen free water (final pH was

approximately 7.4 and osmolarity similar to human tear film))

[0483] siRNA solution formulated in 3% MC formulation 33.3 mg/ml: 1.7 mg dry siRNA were dissolved in 3% MC formulation that was prepared as described to achieve a 51µL formulated solution of siRNA in 4% MC formulation.

[0484] Group VII: Glycerol + EDTA formulation: 50 µL WFI (Norbrook) were mixed with 50µL of Solution B to get: 1% v/v sterile Glycerol & 0.01% w/v EDTA solution in pyrogen free water.

[0485] siRNA solution formulated in Glycerol + EDTA formulation 33.3 mg/ml: 1.7 mg dry siRNA were dissolved in Glycerol + EDTA formulation that was prepared as described to achieve a 51µL formulated solution of siRNA in Glycerol + EDTA formulation.

[0486] Quantity prepared: 7 vials, as follows:

- 1 vial of 1.7 mg siRNA in 51 µL PBS formulation
- 1 vial of 1.7 mg siRNA in 51 µL Systane® formulation
- 1 vial of 1.7 mg siRNA in 51 µL 0.5% MC formulation
- 1 vial of 1.7 mg siRNA in 51µL 1% MC formulation
- 1 vial of 1.7 mg siRNA in 51µL 2% MC formulation
- 1 vial of 1.7 mg siRNA in 51µL 3% MC formulation
- 1 vial of 1.7 mg siRNA in 51µL Glycerol + EDTA formulation

Storage Conditions: freshly prepared

Control Article(s)

[0487]

1. 1. PBS - Supplied by Biological industries (Manufacturer catalog #02-023-5A (For 10x PBS); Batch # 619113).
2. 2. Systane® - commercially available eye drop solution ; Supplied by Alcon; Batch # : Lot 165228F; Quantity supplied: 15ml; Storage Conditions: RT; Expiration Date: 02/2011.

[0488] Formulation Solution A - methylcellulose solution in pyrogen free water Solution A) 0.4 g of methylcellulose 25 (ScienceLab.com, Cat# SLM2050) dissolved in final volume of 10ml of hot boiled water (80-90°C), (Norbrook), cooled down to room temperature.

[0489] Formulation Solution B - Sterile Glycerol & EDTA solution in pyrogen free water Solution B) 332 µl of 60% glycerol (Sigma, Cat# G6279) & 2 µl EDTA solution pH8 (prepared from Sigma, Cat# E9884) in 9.666 ml WFI (Norbrook).

[0490] Formulation Solution C - Concentrated methylcellulose solution in pyrogen free water Solution C) 0.6 g of methylcellulose 25 (ScienceLab.com, Cat# SLM2050) dissolved in final volume of 10ml of hot boiled water (80-90°C), (Norbrook), cooled down to room temperature.

Stability tests of siRNA compounds in the above Methyl Cellulose formulations:

[0491] The stability of Casp2_4 Q02F08002N siRNA was tested in Methyl cellulose formulations according to the following protocol:

siRNA was diluted in the different formulations containing Methyl cellulose, to a final concentration of 7µM siRNA. The 0%,0.5%,1%, 2% and 3% MC formulations were incubated at Room Temperature. In addition, the 3% MC formulation was also incubated at 37°C with and without nuclease inhibitor.

[0492] A 5-µL aliquot of each solution was transferred to 15 µL of 10x TBE-loading buffer after incubation at the following time points: 0,10', 0.5, 1, 1.5,3, 6h. The solution was then frozen in liquid nitrogen, and stored at -20°C.

[0493] 4 µL of each sample was loaded onto a non-denaturing 20% polyacrylamide gel and electrophoresis was performed at 80V for 2.5h.

[0494] For siRNA visualization the gel was stained with Ethidium bromide solution (1.0 µg/µL).

[0495] As a positive control for gel migration of a non-degraded siRNA, 5 µL of a 7 µM tested siRNA solution in PBS was transferred to 15 µL of 10x TBE-loading buffer and loaded onto the gel. Then, the sample was frozen in liquid nitrogen and stored at -20°C.

[0496] As a reference to the migration pattern of a degraded single strand (ss) siRNA, a non-relevant single strand siRNA was prepared.

[0497] Results: The Casp 2_4 siRNA compound was stable in all formulations.

The *in vivo* study:

Test System:

[0498] Animals used: Rats; Strain: Adult, Sprague -Dawley ; Modification: SD

[0499] Source: Harlan, Jerusalem Israel

[0500] Age: 8-10 weeks; Body Weight Range: 220-270 gr

[0501] Sex: Male; Group Size: n=6; Total number of animals: 54

[0502] Animal Husbandry: Diet: Animals were provided an *ad libitum* commercial rodent diet and free access to drinking water.

Environment:

[0503]

1. (i) Acclimatization of at least 5 days.
2. (ii) All the animals were confined in a limited access facility with environmentally-controlled housing conditions throughout the entire study period, and maintained in accordance with approved standard operating procedures (SOPs). Animals were provided *ad libitum* a commercial rodent diet (Harlan Teklad 2018S Global 18% Protein Rodent Diet) and filtered, chlorinated and acidified water.

[0504] Rats were kept in microisolator cages with filter top, 1-6 Rats / cage. The cages were maintained under controlled environmental conditions of temperature (20-24 °C), relative humidity (30-70%), a 12- hr light / 12-hr dark cycle, monitored by the control computer, throughout the study period.

Experimental design

[0505] Experimental setup included 9 experimental groups (6 rats per group). ED were applied bilaterally. Different formulations were used (Study Design Table C11). Termination time point (3hrs) post siRNA administration: Termination setup was performed according to the study design Table C11. Retinal concentration of CASP2_4 siRNA was determined by qPCR.

Table C11: Study Design

Group	Delivery Route	Formulation	siRNA Compound	Bilateral Application (dose/volume/eye)	Term. Time Point (hrs)	Group Size
I	ED	PBS	Casp2_4_S510	100 µg/3µl	3	6
II	ED	Systane	Casp2_4_S510	100 µg/3µl	3	6
III	ED	MC0.5%	Casp2_4_S510	100 µg/3µl	3	6
IV	ED	MC1%	Casp2_4_S510	100 µg/3µl	3	6
V	ED	MC2%	Casp2_4_S510	100 µg/3µl	3	6
VI	ED	MC3%	Casp2_4_S510	100 µg/3µl	3	6
VII	ED	Glycerol + EDTA *	Casp2_4_S510	100 µg/3µl	3	5
VIII	ED	PBS	-	-	3	6
IX	Intact	-	-	-	-	6
* Glycerol + EDTA formulation (MC free formulation) = 1% v/v sterile glycerol & 0.01% w/v EDTA solution in pyrogen free water						

[0506] Anesthesia: In the course of the experiment, All ED treated animals were anesthetized with Equithesine (I.P. 4ml/kg).

[0507] Eye Drop Delivery A 3 µl sample volume of the test article or vehicle were applied to the corneal surface bilaterally to the anesthetized animals, by a blunt pipette tip (filter tips 10µl sterile (short)). The animals were placed in a warm environment to prevent anesthesia-induced hypothermia. Termination of animals from all study groups was according to study design (time point 3 hrs). Tissues were collected according to study design.

[0508] Scheduled euthanasia: All animals were deeply anaesthetized and euthanized according to the study design (Table C11, Termination).

[0509] Perfusion setup: Rats were perfused transcardially with PBS (20-50 rpm/min following 2-3 min).

[0510] Tissue Collection: After perfusion, both eyes (left and right) were washed with PBS, were enucleated and stored on ice. The eyes were dissected using a binocular microscope, and optionally gross pathology was graded according to the sample grading scale. The cornea was dissected by a cut along the limbus, lens was gently removed, and the retina and vitreous were carefully separated from the sclera. Whole retinas and vitreous bodies (humor) were collected into two separate appropriate and properly marked test tubes. Dissected retinas were washed in a large volume of PBS (each retina in a separate tube with fresh PBS), extra liquid was removed with Kimwipes and retinas were snap-frozen in liquid nitrogen. Retina and Vitreous body samples were subjected to RNA extraction.

Evaluation:

[0511] RNA Extraction: Retina: RNA was extracted from each retina sample individually (left and right) by double extraction. The RNA was transferred for cDNA preparation and qPCR analyses.

[0512] Vitreous: Material for Casp2_4_S510 siRNA detection from vitreous was obtained using the following protocol: Each rat vitreous was around 10µl. 500µl of reagent A EZ-RNA II (Biological Industries Cat no. 20-410-100) were added to each vitreous. The sample was homogenized, and 10µg tRNA (1µl from 10 mg/ml stock) were added. The sample was stored for 5 minutes at room temperature. Subsequently EZ-RNA II B: 400 µl and EZ-RNA II C: 90 µl were added and mixed well. The sample thus obtained was stored for 10 min at room temperature then centrifuged 12000g for 15 min at 4°C.

[0513] The upper phase was transferred to a fresh tube to which isopropanol: 500µl and Linear Acrylamide: 5µl were added. The sample was stored overnight at -20°C, centrifuged at 12000g for 20 min at 4°C, washed twice with 75% ethanol. The pellet was dissolved in 15 µl DDW.

[0514] siRNA Quantification: The quantity of the siRNA in retinas and vitreous humor (siRNA quantification was performed per vitreous) was examined by qPCR siRNA according to Quark's standard operating procedures.

Results

[0515] The study showed that the following quantities of siRNA were detected in the rat retina three hours post application of siRNA non-invasively administered *in vivo* with the different formulations:

Table C12: CASP2 siRNA Normalized (miRNA) quantity in fmole/1 µg RNA

Group	Abbreviation	N	Mean	Std
IX	I-N/A	6	0.005	0.005
VIII	P-PBS	6	0.056	0.124
VII	C-MC(0)	6	6.303	3.921
XA	C-MC(1/2)	4	17.831	16.628
XIA	C-MC(1)	6	5.677	4.235
V	C-MC(2)	6	6.963	8.042
VI	C-MC(3)	5	7.653	6.674
II	C-Systane	5	5.109	2.492
I	C-PBS	6	4.228	6.758

[0516] MC(0)-MC(3) - formulations including increasing concentrations from 0 to 3% of Methyl Cellulose.

[0517] Conclusion: All groups that were treated by CASP2 siRNA show positive quantities of the siRNA (>4 fmole) in target retinal tissue. No significant difference (P-value=0.8425) were found between the formulation groups.

Non-Invasively Administered siRNA Induces in-vivo Knockdown (KD) of Target Gene Associated with Apoptosis in Target Ocular Tissue

Example 8 (for illustrative purposes): Determination of in-Vivo Knockdown Activity of siRNA Against p53 prepared for Non-invasive Administration by Eye Drops in Rat Retina (siRNA compound formulated in MC 2%(Formulation "A"))

Objective

[0518] The objective of the study was to determine knockdown activity of non-invasively delivered siRNA compound targeting p53 mRNA in the rat neural retina. The siRNA compound was formulated for non-invasive administration by eye drops. Knockdown activity was assessed by protein level determination using ELISA method.

Test article

[0519]

- a. Substance (unformulated siRNA compound): QM5 (siRNA Against Mouse/Rat p53)
- b. Formulated siRNA compound for non-invasive (eye drop) delivery (groups 2 & 5 in Table C13): 100µg/3µl solution of QM5 siRNA in 2% (w/v) methylcellulose & 1% (v/v) sterile glycerol & 0.01 % (w/v) EDTA solution in pyrogen free water.

3. c. Formulated siRNA compound for intravitreal injection (groups 1 & 4 in Table C13): 280 µg of QM5 siRNA in 140 µl of PBS
4. d. Formulated vehicle solution for non-invasive (eye drop) delivery (groups 3 & 6 in Table C13): 2% methylcellulose & 1% v/v sterile glycerol & 0.01% w/v EDTA solution in pyrogen free water.

Test System:

[0520] Adult Male, Sprague-Dawley (SD) Rats Harlan, Jerusalem Israel, 6-8 weeks old, 160-180 gr each.

Experimental Design

[0521] Study design: Unilateral axotomy (left eye, OS) was performed in each animal from groups 1-6.

[0522] Groups 2, 3, 5& 6: Test compound (100 µg test article in 3 µl of formulated vehicle - groups 2&5) or formulated vehicle alone (groups 3&6) were applied as eye drops every day starting on day 0 (immediately after axotomy). Experimental groups were terminated according to Table C13. Left and Right eye samples from group 7 served as an intact normal controls.

[0523] Groups 1 & 4: Three minutes following axotomy (at day 0), 20 µg of siRNA compound in 10 µl of PBS vehicle were applied by microinjection into the vitreous. The microinjection into the vitreous was performed perpendicular to the sclera, using an Insulin micro-injector (0.3 ml). Animals were terminated according to Table C13.

Table C13: Study Design:

Group	Group Size	Axotomy (unilat. OS)	Intravitreal injection (OS) (dose/volume/eye)	ED (OS) (dose/volume/eye)	Termination (days)
1	6	Yes	QM5 20µg/10 µl	N/A	1
2	6	Yes	N/A	QM5 100µg/3 µl	1
3	4	Yes	N/A	Formulated vehicle	1
4	6	Yes	QM5 20µg/10 µl	N/A	3
5	6	Yes	N/A	3x QM5 100µg/3 µl	3
6	4	Yes	N/A	3x Formulated vehicle	3
7	4	N/A	N/A	N/A	N/A

(see Tables C14-C17 hereinbelow and Figures 11A and 11B)

p53 signal according to Kidney W.C.E standard curve

Table C14: p53 signals values according to Kidney W.C.E standard curve

P53 Signal		retina	
Operation	Treatment	1 days	3 days
Axotomy	QM5 Intravitreal	1.00	1.84
Axotomy	QM5 Eye Drops	0.75	1.41
Axotomy	Vehicle Eye Drops	1.93	Missing
None	Non Treated	2.01	2.01

Table C15: % p53 signals ratio from non-treated animals

% signal ratio from non-treated		retina	
Axotomy	QM5 Intravitreal	50%	92%
Axotomy	QM5 Eye Drops	37%	70%
Axotomy	Vehicle Eye Drops	96%	0%

p53 signal according to GST-hp53

Table C16: p53 signal values according to GST-p53 standard curve

	P53 Signal	retina	
Operation	Treatment	1day	3days
Axotomy	QM5 Intravitreal	0.0159	0.0214
Axotomy	QM5 Eye Drops	0.0095	0.0149
Axotomy	Vehicle Eye Drops	0.0227	Missing
None	Non Treated	0.0233	0.0233

Table C17: % P53 signals ratio from non-treated animals

% Signal ratio from non-treated		retina	
Axotomy	QMS Intravitreal	8%	92%
Axotomy	QM5 Eye Drops	41%	64%
Axotomy	Vehicle Eye Drops	97%	Missing

[0524] Conclusion: These studies showed that non-invasively delivered siRNA compound targeting p53 mRNA in the rat neural retina induces knock down of the p53 protein.

siRNA Compounds Induce Ocular Neuronrotection *In-Vivo* in Animal Models of Ocular Neuronal Injury

Example 9 (for comparative purposes): Evaluation of ocular neuroprotective induction by siRNA compound targeting Caspase 2 in the ONC model after IVT injection

[0525] Study objectives: To evaluate ocular neuroprotective efficacy of siRNA targeting Caspase 2 in the ONC model following intravitreal injection(s)

Methods

Retrograde labelling of RGCs.

[0526] The Fluorogold tracer is transported retrogradely (from brain to eye) along RGC axons resulting in complete and specific labeling of all RGCs. For the purpose of this study, RGCs were labeled by application of the retrograde tracer FluoroGold (2%, Fluorochrome, Englewood, CO) in the superior colliculus. Briefly, a window was drilled in the scalp above the coordinates 6 mm rostral to the bregma and 1.2 mm lateral to the midline in both hemispheres. Using a Hamilton syringe, 3 µl of the FluoroGold was injected into superior colliculus 3.8 mm, 4 mm, and 4.2 mm below the bony surface at a rate 1 µl/min at each of the three depths. The needle was then slowly withdrawn and the skin was sutured. In adult rats, the time required to obtain full labeling of all RGCs following this procedure was ~ 1 week. For this reason, ONC was performed one week after retrograde labeling of RGCs.

[0527] Optic nerve crush. The orbital optic nerve (ON) of anaesthetized adults Wistar rats was exposed through a supraorbital approach, the meninges severed and all axons in the optic nerve (ON) transected by crushing with calibrated forceps for 10 seconds, 2 mm from the lamina cribrosa.

[0528] Intravitreal (IVT) injection. One or two 20 µg doses (in 10 µl of PBS) of test or control siRNAs or 10 µl of PBS vehicle were microinjected into the vitreous body 2 mm anterior to the nerve head, perpendicular to the sclera, using a glass micropipette. IVT administration is shown as a control and or for initial validation of a target gene.

[0529] Quantification of RGC survival. At termination, experimental animals were perfused transcardially with 4% paraformaldehyde. The eyes with the optic nerve were enucleated, the cornea was dissected with the blade and lens/vitreous were gently removed. Both retinas were dissected out, fixed for an additional 30 min and flat-mounted vitreal side up on a glass slide for examination of the ganglion cell layer. RGCs were examined under the fluorescence microscope with an UV filter (365/420 nm). The number of retrogradely fluorescent RGCs were determined by two different, independent and "blinded"

investigators counting them in 16 distinct areas (four areas per retinal quadrant at three different eccentricities of one-sixth, one-half, and five-sixths of the retinal radius). Microglia that may have incorporated FluoroGold after phagocytosis of dying RGCs was distinguished by their characteristic morphology and excluded from quantitative analyses.

Part 1

[0530] Test article: CASP2_4 siRNA - a double-stranded 19-mer oligonucleotide chemically modified by 2'O-methylation on both strands. Other compounds included siRNA chemically modified with L-DNA at the 3' terminus of the sense strand or 2'5' bridges at the 3' terminus. Targets the caspase2 gene in multiple species.

Control article: PBS

[0531] Design: Retinal ganglion cells (RGC) were selectively labeled first by application of the retrograde tracer FluoroGold to the superior colliculus. One week later, animals were subjected to optic nerve crush injury (ONC). The quantifications of surviving RGCs were carried out at 7 and 30 days after ONC by counting FluoroGold-labelled RGCs on flat-mounted retinas. Table C18 shows the groups treatments.

Table C18: Group Treatments

Groups	Right eye (n=4)	Left eye (n=4)	Termination
1 and 2	20 µg siCasp2_4 on day 0	20 µl PBS on day 0	Day 7
3 and 4	20 µg siCasp2_4 on days 0 and 10	20 µl PBS on days 0 and 10	Day 30

[0532] Results: The mean number of surviving RGCs in eyes treated with 20 µg Casp2_4 siRNA and subjected to optic nerve crush was 2040 ± 35 cells/mm² at 7 days and 298 ± 25 cells/mm² at 30 days post injury. These counts were significantly greater than the mean RGC counts in eyes treated with PBS and subjected to nerve crush, which was 941 ± 27 cells/mm² at 7 days and 41 ± 7 cells/mm² at 30 days. In the nonsurgical control eyes, the mean number of RGCs was 2253 ± 104 cells/mm², which is comparable to the average number of RGCs reported in literature. Table C19 provides the results of this study.

Table C19: Mean numbers of survived RGC at the different time points post injury

Time post ONC	7 days		30 days	
Treatment	PBS	Casp2_4	PBS	Casp2_4
Mean survival RGC (cells/mm ²)	941.347	2039.672	41.040	298.051
SD	27.038	34.766	7.219	25.401
% of RGC survival from total RGC	26.659	57.764	1.162	8.441
SD	0.766	0.985	0.204	0.719

[0533] All data are given as mean $\pm$ standard deviation. Values were compared using one-way analysis of variance (ANOVA) and considered as significantly different at $P < 0.02$.

[0534] Conclusions: The increased survival of the RGCs was not due to the neuroprotective effects of anesthesia (ketamine is an antagonist of the N-methyl-D-aspartate receptor, and xylazine is an $\alpha 2$ -adrenergic receptor agonist), since identically anesthetized animals in the control group had significantly lower RGC counts than did animals in the Casp2_4 siRNA treated group, but due to neuroprotective effect of silencing pro-apoptotic Caspase 2 activation and up regulation following ONC injury, by treatment with siRNA compound targeting the Caspase 2 gene.

Part 2

[0535] Test article: CASP2_4L siRNA - a double-stranded 19-mer oligonucleotide chemically modified by 2'O-methylation on the antisense strand and L-DNA on the sense strand. Targets the caspase 2 gene of multiple species.

Control articles:

[0536]

- PBS
- siRNA targeting GFP - a double-stranded 21-mer oligonucleotide stabilized by 2'O-methylation on both strands.
- CNL - siRNA with no match to any known mammalian transcript; a double-stranded 19-mer oligonucleotide chemically modified by 2'O-methylation on both strands.

[0537] Design. Retinal ganglion cells (RGC) were selectively labeled first by application of the retrograde tracer FluoroGold to the superior colliculus. One week later, animals were subjected to optic nerve crush injury (ONC). The quantifications of surviving RGCs were carried out at day 7 after ONC by counting FluoroGold-labelled RGCs on flat-mounted retinas. Test or control articles were injected at the time of ONC. Similar experiments were performed in order to test activity and efficacy of siRNA administered via eye drops (see next example). Table C20 shows groups treatments.

Table C20: Groups Treatments

Groups	Right eye (n=4)	Left eye (n=4)	Termination
1 and 2	20 µg siCasp2_4 on day 0	20 µl PBS on day 0	Day 7
3 and 4	20 µg siGFP on day 0	20 µl siCNL_1 on day 0	Day 7

[0538] Results: The mean number of surviving RGCs in eyes treated with 20 µg Casp2_4 siRNA and subjected to optic nerve crush was 2085 ± 40 cells/mm² at 7 days post injury. These counts were significantly greater than the mean RGC counts in eyes treated with either PBS or GFP siRNA or CNL_1 siRNA and subjected to nerve crush, which was 901 ± 50 cells/mm², 922 ± 38 cells/mm², 898 ± 42 cells/mm² respectively, at 7 days. In the nonsurgical control eyes, the mean number of RGCs was 2196 ± 110 cells/mm², which is comparable to the average number of RGCs reported in literature. Table C21 shows results. All data are given as mean $\pm$ standard deviation. Values were compared using one-way analysis of variance (ANOVA) and considered as significantly different at $P < 0.01$.

Table C21: Mean numbers of survival RGC 7 days post injury (n=4 retinas/group).

Treatment	PBS	siGFP	siCNL_1	siCasp2
Mean survival RGC (cells/mm ²)	901.3338	922.3666	898.4268	2084.815
SD	49.74134	38.04059	42.12429	40.03638
% of RGC survival from total RGC	25.52577	26.12142	25.44345	59.04197
SD	1.408675	1.07731	1.19296	1.13383

[0539] Conclusions: The increased survival of the RGCs was due to neuroprotective effect of silencing pro-apoptotic Caspase 2 activation and up-regulation following ONC injury, by treatment with siRNA targeting Caspase 2 gene. The Casp2_4 siRNA molecules having different structural modifications / motifs show similar neuroprotective effect on RGC survival range.

Example 10: Evaluation of ocular neuroprotective efficacy of siRNA compound targeting Caspase 2 in the IOP model

[0540] The objective of the current study was to establish the neuroprotective effect of a single injection of 20 µg of CASP2_4 compound or negative control GFP siRNAs in the pre-clinical rat glaucoma IOP model.

Study outline

[0541] In animals from all study groups, RGCs were retrograde labeled using the fluorescent tracer Dil (1,1'-dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine) applied to both superior colliculi. A week later, IOP was unilaterally increased in the left eyes of animals from groups 2 to 5. IOP was monitored every other day in the operated (left) and contralateral (right) eyes during 2 weeks. Two weeks following hypertonic saline injection, vehicle (PBS) or siRNA (20 µg in PBS, either siRNA targeting Casp2 or a negative control siRNA targeting the non-mammalian gene, Green Fluorescent Protein), was administered by intravitreal (IVT)

injection to the left (operated) eyes. The contralateral non-operated eyes received no IVT injection. In addition, two weeks after hypertonic saline injection, the animals from group 2 were sacrificed and DiI-labeled RGCs were counted in flat mounts of left retinas under fluorescence microscopy, in order to assess the extent of RGC loss at the time of siRNA injection. For all groups receiving IVT injection, IOP was measured once during the 3rd week of the experiment to confirm its increased state in the operated eyes. At 3 weeks of increased IOP, or at 1 week after the test and control article injections, the animals were sacrificed and DiI-labeled RGCs were counted in flat mounts of left retinas under fluorescence microscopy. Right retinas served as internal controls for labeling quality. A separate group of naive rats in which RGC's were retrogradely labeled at the experiment commence was used as an intact control to provide a reference for normal RGC densities. 4 rats from this group were sacrificed at 2 weeks of ocular hypertension (together with group 2) and the remaining 9 rats - at 3 weeks (together with groups 3 to 5). The experimental design is shown in Table C22 below.

Animals

[0542]

Species:	Rats	
Strain:	Brown NorwayRats	
Modification:	N/A	
Source:	Charles River(Canada)	
Age:	Adult ,retired breeders ,10-12 months of age	
Body Weight Range:	300-400g	
Sex:	Males	
Animal Husbandry:	Diet:	
	Environment:	All animal procedures were performed in accordance with the guidelines of the Canadian Council on Animal Care for the use of experimental animals (http://www.ccac.ca/).
	Environment:	(i) Acclimatization of at least 5 days.
		(ii) All the animals were confined in a limited-access facility with environmentally controlled housing conditions throughout the entire study period, and maintained in accordance with University of Montreal approved standard operating procedures (SOPs).

Materials and Equipment

[0543] Substance (unformulated compound) CASP2_4_S510 (CASP2_4; CASP2 siRNA; siRNA Against Caspase2 mRNA)

Supplied by Agilent

[0544] Description of the test material: A 19-mer chemically modified blunt-ended duplex having two separate strands, with a sense strand (SEN) comprising unmodified ribonucleotides (upper case letters), an L-deoxyribonucleotide at position 18 (bold, underlined) and inverted deoxyabasic moiety (iB) present at the 5' terminus of the SEN strand; and with an antisense strand (AS) comprising unmodified ribonucleotides (upper case letters), and 2'OMe sugar modified ribonucleotides (lower case letters) at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19 as shown in Formula 1.

Quantity supplied: 300 mg

Storage Conditions: -80°C

[0545] Control Substance (unformulated compound) GFP_5_S763 (siRNA Against GFP mRNA)

Outsourcing (manufacturer's name): Agilent

Manufacturer's catalog # N/A

Quantity supplied: 220.8150 mg

Storage Conditions: -80°C; Expiration Date: ND

Test /Control Article (formulated) CASP2_4 / GFP_5 20µg/5µL of PBS

[0546] 0.5 mg was diluted into 125µl to achieve a stock solution of 4µg/µl. These were subsequently aliquoted to 5µl per tube and stored at -80°C.

Vehicle - PBS.

[0547] Multicell Phosphate buffered saline, solution 1X, without calcium, without magnesium (CAT. No: 311-010-EL).

[0548] Fluorescence microscopy was performed using a Zeiss Axioskop 2 Plus microscope (Carl Zeiss Canada, Kirkland, QC), images were captured with a CCD camera (Retiga, Qimaging) and processed with Northern Eclipse image analysis software (Empix Imaging, Mississauga, ON). Microphotographs were taken at 25X magnification.

Experimental Procedure

[0549] Surgery was performed in adult male Brown Norway rats, retired breeders, between 10-12 months of age (300-400g), under general anesthesia by intraperitoneal injection of 1 mL/Kg standard rat cocktail consisting of Ketamine (100 mg/mL), xylazine (20 mg/mL) and acepromazine (10 mg/ mL).

[0550] Ocular hypertension surgery: Unilateral and chronic elevation of IOP was induced by using the Morrison model, involving an injection of hypertonic saline into a episcleral vein, leading to a blockade of the aqueous humor outflow pathways. This procedure lead to gradual increase of eye pressure and progressive death of RGS. All the animals in this study received only a single saline vein injection. The eye selected for the procedure was adapted with a plastic ring applied to the ocular equator to confine the saline injection to the limbal plexus. A micro needle (30-50 µm in diameter) was used to inject 50 µL of sterile 1.85 M NaCl solution through one episcleral vein. The plastic ring temporarily blocked off other episcleral veins forcing the saline solution into the Schlemm's canal to create isolated scarring. Animals were kept in a room with constant low fluorescent light (40-100 lux) to stabilize circadian IOP variation (8).

[0551] Measurement of intraocular pressure (IOP): IOP from glaucomatous and normal (contralateral) eyes was measured in awake animals using a calibrated tonometer (TonoPen XL, Medotronic Solan, Jacksonville, FL). IOP was measured every other day for the first two weeks after ocular hypertension surgery; and at least once between week 2 and week 3 after ocular hypertension surgery. Eyes become fragile after intraocular injections in conditions of high IOP, for this reason the number of IOP measurements was limited as this involves putting additional pressure on the cornea to get a reliable reading. The mean IOP (mmHg ± S.E.M) per eye was considered as the average of all IOP readings since the onset of pressure elevation. The maximum IOP measurements in each individual eye, glaucomatous or normal contralateral eye was defined as the peak IOP and this value was used to estimate the mean peak IOP for each group

[0552] Calculation of intraocular pressure (IOP): The mean IOP (mmHg ± S.E.M) per eye was considered as the average of all IOP readings since the onset of pressure elevation. The maximum IOP measurements in each individual eye, glaucomatous or normal contralateral eye was defined as the peak IOP and this value was used to estimate the mean peak IOP for each group. The positive integral IOP was calculated as the area under the IOP curve in the glaucomatous eye minus that of the fellow normal eye from ocular hypertension surgery to euthanasia. Integral IOP represents the total, cumulative IOP exposure throughout the entire experiment.

Study Design:

[0553] One week prior to induction of glaucoma, RGCs were retrogradely labeled using the fluorescent tracer Dil applied to both superior colliculi.

[0554] One week later, unilateral elevation of IOP was induced by injection of a hypertonic saline solution into an episcleral vein. This procedure was referred to as ocular hypertension surgery.

[0555] In a first experiment, prior to efficacy studies, the status of RGC loss in "no injection" groups at 2 and 3 weeks after ocular hypertension surgery (4 rats/group, total = 8 rats) was assessed.

[0556] Exactly 2 weeks after ocular hypertension surgery, a single intravitreal injection of each siRNA was performed.

[0557] Animals were euthanized and retinas prepared for analysis of RGC survival at exactly 3 weeks after ocular hypertension surgery.

[0558] The density of surviving RGCs was quantified in 12 standard retinal areas.

Table C22: Study Design

Group	n	Increased IOP	IVT Injection (µL/eye) 2 weeks post IOP induction	Dose Volume	Termination (Weeks Post IOP)
1 (Intact)	13 (4+9)	No	No	N/A	N/A
2	4	Yes	No	N/A	2
3	6	Yes	PBS Vehicle	5 µL	3
4	6	Yes	Negative Control siRNA, 20 µg	5 µL	3
5	6	Yes	CASP2 siRNA, 20 µg	5 µL	3
*NA - non-applicable, Group size: n=5					

[0559] Quantification of surviving RGC soma: Quantification of RGC bodies was performed in duplicate and in a masked fashion. For RGC density counts, rats were deeply anesthetized and then perfused transcardially with 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer and both eyes were immediately enucleated. Retinas were dissected and flat-mounted on a glass slide with the ganglion cell layer side up. Under fluorescence microscopy, Dil-labeled neurons were counted in 12 standard retinal areas as described.

RESULTS & DISCUSSION

[0560]

Table C23: Percent RGC from intact inference table

Treat	N	Mean	Std	CV	Dunnett P-value
PBS	6	62.37	5.86	9.40	
CASP2_4	6	81.71	14.28	17.48	0.05
2wIOP	4	71.25	6.75	9.47	0.19
GFPsiRNA	6	65.75	4.97	7.57	0.68
intact	13	100.00	2.84	2.84	0.00

Comparisons to PBS treated group

[0561] Preservation of RGC in CASP2_4 siRNA-treated eyes with increased IOP was significantly better than in the PBS treated group (1.31 fold higher from the PBS result; p-value = 0.05). In GFP siRNA treated group no significant difference was found (p-value=0.68) with respect to PBS treated group.

[0562] No significant difference between IOP (2 weeks) to IOP (3 weeks) treated by PBS (p-value=0.19).

Comparisons to IOP (2 weeks) group

[0563] The intact group has significantly higher RGC density per mm² than IOP (2w) group (p-value=0.00). No significant differences in RGC density per mm² were found between IOP (2w) group and other IOP groups that were sacrificed after 3 weeks (p-value range from 0.16 to 0.44).

Comparisons to Intact group

[0564] The intact group has significantly higher RGC density per mm² than IOP groups that were treated by PBS, GFP siRNA and IOP(2w) without treatment (p-value<0.01). No significant differences in RGC density per mm² were found between IOP group treated by CASP2_4 siRNA compared to intact group (p-value=0.06).

[0565] These data demonstrate that a siRNA targeting Caspase2 provides protection of RGCs from damage induced by increased intraocular pressure and its administration at 2 weeks of increased IOP leads to cessation of further RGC loss.

[0566] IOP measurements were carried out as described above.

[0567] IOP: The Dunnett approach was used for comparison of the results to PBS treated group. Additional parameters were calculated and analyzed such as: AUC (area under the curve), Mean IOP and Max(IOP) along the duration of measurements.

[0568] Most animals show significant increase in IOP at 1-2 weeks after ocular hypertension surgery.

[0569] No significant differences in IOP levels were found among operated eyes in treated groups (0.7746). Likewise, no differences in IOP levels were found among contra lateral eyes in treated groups and intact eyes.

[0570] The results show that Casp2 siRNA induced neuroprotection even as IOP remained elevated in all treatment groups.

[0571] Results: No significant differences between groups were found in all three calculated IOP parameters

Example 11: Evaluation of ocular neuroprotective efficacy of siRNA compound targeting Caspase 2 in the axotomy model

[0572] The purpose of the present efficacy studies was to use a model of RGC apoptosis induced by axotomy of the optic nerve (ON) in adult Sprague-Dawley rats. The onset and kinetics of RGC death in this model system are very reproducible and allow for the establishment of the neuroprotective efficacy of Casp2_4 siRNA *in vivo*. Using this method, the time course of RGC death follows a predictable course: cell death begins on day 5 and proceeds to the rapid loss of more than 90% of these neurons by 2 weeks.

Methods

[0573] Retrograde labeling of RGCs: For the purpose of this study, RGCs were labeled by application of the retrograde tracer FluoroGold (2%, Fluorochrome, Englewood, CO) in the superior colliculus. Briefly, both superior colliculi were exposed and a small piece of gelfoam soaked in FluoroGold was applied to their surface. In adult rats, the time required to obtain full labeling of all RGCs following this procedure is ~ 1 week. For this reason, optic nerve axotomy and intraocular injection of siRNA molecules were performed one week after retrograde labeling of RGCs.

[0574] Optic nerve axotomy: The entire population of RGCs were axotomized by transecting the optic nerve close to the eye (0.5 to 1 mm). Retinal fundus examination was routinely performed after each axotomy to check the integrity of the retinal circulation after surgery. Animals showing signs of compromised blood supply were excluded from the study.

[0575] For intravitreal injection 10 µg in 5 µl of PBS each of the reagents, either Casp2_4 siRNA or GFP siRNA were microinjected into the vitreous body 2 mm anterior to the nerve head, perpendicular to the sclera, using a glass micropipette at the time of surgery, day 0, and then repeated at day 7.

[0576] Quantification of RGC survival: Experimental and control animals were perfused transcardially with 4% paraformaldehyde

at 14 days after optic nerve axotomy. The left retinas (treated) and the right retinas (untreated controls) were dissected out, fixed for an additional 30 min and flat-mounted vitreal side up on a glass slide for examination of the ganglion cell layer. RGCs backfilled with FluoroGold were counted in 12 standard retinal areas. Microglia that may have incorporated FluoroGold after phagocytosis of dying RGCs were distinguished by their characteristic morphology and excluded from our quantitative analyses.

Experimental design

[0577] Test article: CASP2_4L siRNA: a double-stranded 19-mer oligonucleotide chemically modified by 2'O-methylation of sugar residues on the antisense strand and L-DNA on the sense strand. Targets caspase2 gene in multiple species.

Control articles

PBS

[0578] siRNA targeting GFP - a double-stranded 21-mer oligonucleotide chemically modified by 2'O-methylation on both strands.

[0579] CNL - siRNA with no match to any known mammalian transcript; a double-stranded 19-mer oligonucleotide chemically modified by 2'O-methylation on both strands. Table C24 shows groups treatments.

Table C24: Groups Treatments

SiRNA	SiRNA Dose	Time of administration (post axotomy)	Time of analysis (post axotomy)	No. of animals per group	Sample preparation
SiGFP, axotomy (OS)	10 ug X 2	Time 0	2 weeks	8	Flat mount
		1 week			
SiCasp2_4, Axotomy (OS)	10 ug X 2	Time 0	2 weeks	8	Flat mount
		1 week			

[0580] Results: The mean number of surviving RGCs in eyes treated with two intravitreal injections of 10 µg Casp2_4 siRNA and subjected to axotomy was 533 ± 24 cells/mm² at 14 days post injury. These counts were significantly greater than the mean RGC counts in eyes treated with GFP siRNA at the similar regiment and subjected to axotomy, which was 130 ± 7 cells/mm² at 14 days. In the nonsurgical control eyes, the mean number of RGCs was 2138 ± 91 cells/mm², which is comparable to the average number of RGCs reported in literature.

Table C25: Mean numbers of survival RGC 14 days post injury (n=6 retinas/group)

Treatment	GFP	Casp2_4
Mean survival RGC (cells/mm ²)	130	533
SE	7	24
% of RGC survival from total RGC	6	25
SE	0.32	1.12

[0581] Data analyses and statistics were performed using the GraphPad Instat software by one-way analysis of variance (ANOVA). All data are given as mean $\pm$ SE, significantly different at $P < 0.02$.

[0582] Conclusions: The increased survival of the RGCs was due to neuroprotective effect by silencing pro-apoptotic Caspase 2 activation and up regulation following axotomy by treatment with siRNA targeting Caspase 2 gene.

[0583] Discussion: In the present studies, the Casp2_4 siRNA was neuroprotective for at least 30 days in an optic nerve crush model and for 14 days in axotomy model of RGC loss. Optic nerve crush and axotomy experiments provide a realistic model of acute optic neuropathies.

Non-Invasive Administration of siRNA Compounds Induces Ocular Neuroprotection *In-Vivo* in Animal Models of

Ocular Neuronal Injury**Example 12: Evaluation of ocular neuroprotective efficacy of non-invasively administered siRNA compound targeting Caspase 2 in the ONC model**

[0584] Test article: CASP2_4 siRNA (compound of Formula I)

Control articles:

[0585]

- Methyl Cellulose
- CNL - siRNA with no match to any known mammalian transcript; a double-stranded 19-mer oligonucleotide stabilized by 2'O-methylation on both strands.

[0586] Design: Retinal ganglion cells (RGC) were selectively labeled first by application of the retrograde tracer FluoroGold to the superior colliculus. One week later, animals were subjected to optic nerve crush injury (ONC). Eye drops were applied every other day during one week (3 times over all). 100µg/3µl CNL_1, or Casp2_4 siRNA or 3µl of MC vehicle were applied, the first dose was applied 10 minutes after ONC. The quantifications of surviving RGCs were carried out at day 7 after ONC by counting FluoroGold-labelled RGCs on flat-mounted retinas.

[0587] Results: The mean number of surviving RGCs in eyes treated with 20 µg Casp2_4 siRNA and subjected to optic nerve crush was 445 ± 17 cells/mm² at 7 days post injury. These counts were significantly greater than the mean RGC counts in eyes treated with either PBS or CNL_1 siRNA and subjected to nerve crush, which were 337 ± 11 cells/mm², 341.6 ± 13 cells/mm², respectively, at 7 days. Table C26 shows results. All data are given as mean $\pm$ standard deviation. Values were compared using one-way analysis of variance (ANOVA) and considered as significantly different at $P < 0.01$.

Table C26: Mean numbers of survival RGCs 7 days post injury.

	CNL_1	Casp2_4	MC vehicle
Average	341.625	445.4375	337.5
SD	13.03265	17.222331	11.3389342
cells/mm ²	934.6785	1218.7073	923.392613
Sd	35.65705	47.119921	31.0230758
RGC/retina	26470.1	34513.789	26150.4788
SD	1009.808	1334.4362	878.573505
% total	26.4701	34.513789	26.1504788
SD	1.009808	1.3344362	0.87857351
Average	341.625	445.4375	337.5
SD	13.03265	17.222331	11.3389342

[0588] Conclusions: The increased survival of the RGCs was due to neuroprotective effect of silencing pro-apoptotic Caspase 2 following ONC injury by treatment with eye drops containing siRNA targeting Caspase 2 gene in a methyl cellulose formulation.

Example 13: Evaluation of ocular neuroprotective efficacy of non-invasively administered siRNA compound targeting Caspase 2 in the IOP model

Experimental setup:

[0589] Experimental animals: All animal procedures are performed in accordance with the guidelines of the Canadian Council on Animal Care for the use of experimental animals (<http://www.ccac.ca/>). Surgeries are performed in adult male Brown Norway rats, retired breeders, between 10-12 months of age (300-400 g), under general anesthesia by intraperitoneal injection of 1 ml/kg standard rat cocktail consisting of ketamine (100 mg/ml), xylazine (20 mg/ml) and acepromazine (10 mg/ml).

[0590] Retrograde labeling of RGCs: For neuronal survival experiments, RGCs are retrogradely labeled with 3% Dil (1,1'-diiododecyl-3,3',3',3'-tetramethyl-indocarbocyanine perchlorate; Molecular Probes, Junction City, OR), a fluorescent carbocyanine marker that persists for several months without fading or leakage and does not interfere with the function of labeled cells. For retrograde labeling, both superior colliculi, the main targets of RGCs in the brain, are exposed and a small piece of gelfoam (Pharmacia and Upjohn Inc., Mississauga, ON) soaked in Dil will be applied to their surface. Seven days after Dil application, the time required for labeling the entire RGC population, animals are subjected to ocular hypertension surgery as described below.

[0591] Ocular hypertension surgery: Unilateral and chronic elevation of IOP are induced using a method that involves injection of a hypertonic saline solution into an episcleral vein. All the animals involved in this study receive only a single saline vein injection. The eye selected for the procedure is adapted with a plastic ring applied to the ocular equator to confine the saline injection to the limbal plexus. A microneedle (30-50 μ m in diameter) is used to inject 50 μ l of sterile 1.85 M NaCl solution through one episcleral vein. The plastic ring temporarily blocks off other episcleral veins forcing the saline solution into the Schlemm's canal to create isolated scarring. Animals are kept in a room with constant low fluorescent light (40-100 lux) to stabilize circadian IOP variation.

[0592] Measurement of intraocular pressure (IOP): IOP from glaucomatous and normal (contralateral) eyes is measured in awake animals using a calibrated tonometer (TonoPen XL, Medtronic Solan, Jacksonville, FL). IOP is measured every other day for the first two weeks after ocular hypertension surgery; and at least once between week 2 and week 3 after ocular hypertension surgery. Eyes become fragile after intraocular injections in conditions of high IOP (see below), so it is advisable to limit the number of IOP measurements as this involves putting additional pressure on the cornea to get a reliable reading. The mean IOP (mm Hg $\pm$ S.E.M.) per eye is considered as the average of all IOP readings since the onset of pressure elevation. The maximum IOP measured in each individual eye, glaucomatous or normal contralateral eye is defined as the peak IOP and this value is used to estimate the mean peak IOP for each group. The positive integral IOP is calculated as the area under the IOP curve in the glaucomatous eye minus that of the fellow normal eye from ocular hypertension surgery to euthanasia. Integral IOP represents the total, cumulative IOP exposure throughout the entire experiment.

[0593] Intraocular injection of siRNA molecules: Intraocular injections are performed by infusing each siRNA compound at a concentration of 20 μ g into the vitreous chamber of the left eye (total injection volume: 5 μ l). The right eye serves as contralateral control. In addition, non-operated eyes from naive rats are used as intact controls. Intravitreal injections are performed using a 10- μ l Hamilton syringe adapted with a ~32-gauge glass needle. The needle tip is inserted into the superior hemisphere of the eye, at the level of the pars plana, at a 45° angle through the sclera into the vitreous body. This route of administration avoids retinal detachment or injury to eye structures, including the lens and the iris, which can release factors that induce RGC regeneration and survival. The injection is performed over a period of 2 min and the needle is kept in place for an additional 2 min, after which it is gently removed. Surgical glue (Indermill, Tyco Health Care, Mansfield, MA, USA) will be used to seal the site of injection.

Topical Eye Drop instillation of siRNA molecules:

[0594] Daily during week 3 after IOP induction a 3 μ l sample volume of 100 μ g of siRNA are applied to the corneal surface bilaterally to the anesthetized animals, by a blunt pipette tip (filter tips 10 μ l sterile (short)). The animals are placed in a warm environment to prevent anesthesia-induced hypothermia

[0595] Quantification of surviving RGC soma: Quantification of RGC bodies is performed in duplicate and in a masked fashion. For RGC density counts, rats are deeply anesthetized and then perfused transcardially with 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer and both eyes are immediately enucleated. Retinas are dissected and flat-mounted on a glass slide with the ganglion cell layer side up. Under fluorescence microscopy, DiI-labeled neurons is counted in 12 standard retinal areas as described.

[0596] IOP (intra ocular pressure) is induced and followed for 2 weeks. Then, CASP2_4 or control siRNA are administered by either Eye Drop (siRNA in a 2% methyl cellulose formulation) or IVT (siRNA in PBS) as detailed in the study design in Table C27.

Table C27: Study Design

Group	Treatment	Intravitreal injection (5 uL)		Eye drops (3 uL/eye)		Termination
		Dose	Frequency	Dose	Frequency	
1	CASP2_4	20 ug	Once a week starting at 2 weeks after IOP induction (total 3 times during weeks 3, 4, 5)	NA*	NA	5 weeks after IOP induction
2	SiGFP	20 ug	Once a week starting at 2 weeks after IOP induction (total 3 times during weeks 3,4, 5)	NA*	NA	5 weeks after IOP induction
3	CASP2_4	20 ug	Once after 2 weeks of IOP induction at the beginning of week 3	100 ug	Daily during weeks 4 and 5 after IOP induction	5 weeks after IOP induction
4	siGFP	20 ug	Once after 2 weeks of IOP induction at the beginning of week 3	100 ug	Daily during weeks 4 and 5 after IOP induction	5 weeks after IOP induction
5	CASP2_4			100 ug	Daily during week 3 after IOP induction	3 weeks after IOP induction
6	siGFP			100 ug	Daily during week3 after IOP induction	3 weeks after IOP induction
*NA - non-applicable, Group size: n=5						

[0597] The results show that non-invasive delivery of siRNA formulated in a 2% methyl cellulose formulation provides neuroprotection and increases neuronal survival.

Example 14: Rat Optic Nerve Crush (ONC) Model: Comparison of intravitreal siRNA delivery to topical eye drop delivery

[0598] For optic nerve transection the orbital optic nerve (ON) of anesthetized rats is exposed through a supraorbital approach, the meninges severed and all axons in the ON transected by crushing with forceps for 10 seconds, 2 mm from the lamina cribrosa.

[0599] The siRNA compounds are delivered alone or in combination in 5uL volume (20ug/uL) as eye drops. Immediately after optic nerve crush (ONC), 20 ug/10ul test siRNA or 10ul PBS is administered by IVT to one or both eyes of adult Wistar rats. After that, siRNA is applied to the eye by ED every other day and the levels of siRNA taken up into the dissected and snap frozen whole retinas at 5 hours and 1 day, and later at 2 days, 4 days, 7 days, 14 days and 21 days post injection is determined. Similar experiments are performed in order to test activity and efficacy of siRNA administered via eye drops.

[0600] Table C28 below shows an experimental procedure to determine efficacy of test siRNAs (siTEST1; siTEST2) alone or in combination, in the ONC model. Additional dosing parameters, concentrations, termination schedules, formulations and the like are contemplated.

Table C28: Study Design

Group (n=8)	Right eye (n=8)	Left eye (n=8)	termination
1 and 2	20 µg siTEST1 on days 0 and 7 (intravitreal)	20 µg siGFP on days 0 and 7 (intravitreal)	Day 21
3 and 4	20 µg siTEST1 on days 10 and 20 (intravitreal)	20 µg siGFP on days 10 and 20 (intravitreal)	Day 30
5 and 6	PBS (eye drops) every other day starting from day 0, 3 times/week)	PBS on days 0 and 7 (intravitreal)	Day 21

Group (n=8)	Right eye (n=8)	Left eye (n=8)	termination
7 and 8	50µg siTEST1 (eye drops) every other day starting from day 0, 3 times/week)	50µg siGFP (eye drops) every other day starting from day 0, 3 times/week)	Day 10
9 and 10	50µg siTEST1 (eye drops) every other day starting from day 0, 3 times/week)	50µg siGFP (eye drops) every other day starting from day 0, 3 times/week)	Day 21
11 and 12	50µg siTEST2 (eye drops) every other day starting from day 0, 3 times/week)	50µg siGFP (eye drops) every other day starting from day 0, 3 times/week)	Day 21
13 and 14	20 µg siTEST1+ 20 µg siTEST2 (eye drops) every other day starting from day 0, 3 times/week)	40 µg siGFP on days 0, 7 and 20 (eye drops) every other day starting from day 0, 3 times/week)	Day 21
15 and 16	20 µg siTEST1+ 20 µg siTEST2 on days 0, and 10 (intravitreal) n=4	40 µg siGFP on days 0 and 10 (intravitreal) n=4	Day 21

[0601] Results: According to the results that are obtained in this study non-invasive delivery of siRNA compound designed for down regulation of a target gene provides neuroprotection and increases neuronal survival in the retina.

[0602] Table B discloses oligonucleotide pairs of sense and antisense nucleic acids useful in synthesizing unmodified or chemically modified siRNA compounds. The table discloses the position of the sense strand along the mRNA for at least one variant.

Table B CASP2 - caspase 2, apoptosis-related cysteine peptidase

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
1	GAGUGAUGCCGGUAAAGAA	UUCUUUACCGGCAUCACUC	Cyn, Mnk	[1182-1200]
2	GGAAGGUGUCCAAAUUUAA	UUAAAUUUGGACACCUUCC	Mnk	[1965-1983]
3	GGGUCGACAUAGUAUGGAA	UCCAUAUCUAUGUCGACCC	Mnk	[4025-4043]
4	GGAGAGUGAUGCCGGUAAA	UUUACCGGCAUCACUCUCC	Cyn, Mnk	[1179-1197]
5	CAUCGAGGCUCUUGCUCAA	UUGAGCAAGAGCCUCGAUG	Mnk	[1305-1323]
6	GGAUCAUGUAAAUGCUCAA	UUGAGCAUUUACAUGAUCC	Mnk	[2505-2523]
7	AGGUGAACGCACUUAUCAA	UUGAUAAAGUGCGUUCACCU	Mnk	[1373-1391]
8	CCAGGAUUUGGUUGUUUAA	UAAAACAACCAAUCCUGG		[3229-3247]
9	CAAUCAACCUGCACUUCAU	AUGAAGUGCAGGUUGAUUG	Mnk	[2368-2386]
10	CACAGGAAAUUGCAAGAGAA	UUCUCUUGCAUUUCCUGUG	Cyn, Rat, Ms, GP, Chn, Mnk	[890-908]
11	GCCCAAGCCUACAGAACAA	UUGUUCUGUAGGCUUGGGGC	Cyn, Mnk	[1061-1079]
12	UGACCAGACUGCACAGGAA	UUCUGUGCAGUCUGGUCA	Cyn, Rat, GP, Dog, Mnk	[879-897]
13	CAAGGAUCGGGAAGGUUAA	AUAACCUUCCCGAUCCUUG	Mnk	[1389-1407]
14	CAACUUCCUGUUCUUUAA	UUAAAGAAGAGGAAGUUG	Mnk	[2663-2681]
15	GGAUUUGGUUGUUUAAAUA	UAUUUAAACAACCAAUCC		[3232-3250]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
16	CCACCAAUCUGUCUAGAA	UUCUAGACAGAUUUGGUGG	Mnk	[2474-2492]
17	CCUACAGAACAAACCAAAA	UUUUGGUUUUGUUCUGUAGG	Mnk	[1068-1086]
18	CAGGAAACUCUAAAAAAGA	UCUUUUUUAGAGUUUCCUG	Cyn, Mnk	[256-274]
19	CAUCAUCAGGAAACUCUAA	UUAGAGUUUCCUGAUGAUG	Mnk	[250-268]
20	ACAGGAAAUUGCAAGAGAAA	UUUCUCUUGCAUUUCCUGU	Cyn, Rat, Ms, GP, Chn, Mnk	[891-909]
21	AGGAAACUCUAAAAAAGAA	UUCUUUUUUAGAGUUUCCU	Cyn, Mnk	[257-275]
22	GAAGGUGUCCAAAUUUAAU	AUUAAAAUUGGACACCUUC		[1966-1984]
23	GCAUCCUCAUCAUCAGGAA	UUCCUGAUGAUGAGGAUGC	Mnk	[243-261]
24	GAGUUUGGAAGGUGUCCAA	UUGGACACCUUCCAAACUC	Mnk	[1959-1977]
25	UGCCCAUGGAAGUCUCAA	UUUGAGACUCCAUGGGCA	Mnk	[2179-2197]
26	GGACAUCAUCACCUUGGAA	UUCCAAGGUGAUGAUGUCC	Cyn, Dog, Mnk	[336-354]
27	CCUGACAAGUGAAGUUGUA	UACAACUUCACUUGUCAGG	Mnk	[1766-1784]
28	UGGUGUCCUUCUCCUCAA	UUGAGGAAGAAGGACACCA		[3491-3509]
29	CGUCCAUCUCUCUCCCUUA	UAAGGGAGAGAGAUGGACG	Mnk	[2332-2350]
30	CUAUGGUGUGGAUGGGAAA	UUUCCCAUCCACACCAUAG	Mnk	[996-1014]
31	CAAUAUACACGGAUUUAUA	UAAUAAUCCGUGUAUAUUG	Mnk	[3812-3830]
32	CGUUGAGCUGUGACUACGA	UCGUAGUCACAGCUCAACG	Rb, Dog, Mnk	[539-557]
33	UCAUGUAAAUGCUCAAAGA	UCUUUGAGCAUUUACAUGA		[2508-2526]
34	AGUGAUGCCGGUAAAAGAAA	UUUCUUUACCGGAUCACU	Cyn, Mnk	[1183-1201]
35	ACUCCUGAAUUUUUAUCAA	UUUGAUAAAAUUCAGGAGU	Mnk	[685-703]
36	GCACUUCACUGGAGAGAAA	UUUCUCUCCAGUGAAGUGC	Cyn, Rat, Ms, Dog, Mnk	[771-789]
37	CUACAAUCCUACCUUUUGA	UCAAAGGUAGGAWGUAG	Mnk	[3776-3794]
38	GACCAGACUGCACAGGAAA	UUUCCUGUGCAGUCUGGUC	Cyn, Rat, GP, Chn, Dog, Mnk	[880-898]
39	AAGCAAACAUGACUAGAGA	UCUCUAGUCAUGUUUGCUU		[2011-2029]
40	CCCUCAUUUAAAAUGUAAA	UUUACAUUUUAAAUGAGGG	Mnk	[3297-3315]
41	CUAUGACGUCCAUGUUCUA	UAGAACAUGGACGUCAUAG	Mnk	[858-876]
42	AGGACAUCAUCACCUUGGA	UCCAAGGUGAUGAUGUCCU	Cyn, Dog, Mnk	[335-353]
43	UCAUCAUCAGGAAACUCUA	UAGAGUUUCCUGAUGAUGA	Mnk	[249-267]
44	GCAGAAUUUUGCACAGUUA	UACUGUGCAAAAUUCUGC	Mnk	[912-930]
45	GACUGCACAGGAAAUAGAA	UUGCAUUUCCUGUGCAGUC	Cyn, Rat, GP, Chn, Mnk	[885-903]
46	CCAUCUCUCUCCCUUACUA	UAGUAAGGGAGAGAGAUGG	Mnk	[2335-2353]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
47	CUGCCUUUGGGUCGACAU	UAUGUCGACCCAAAGGCAG	Mnk	[4017-4035]
48	UGACAAGUGAAGUUGUAAA	UUUACAACUUCACUUGUCA	Mnk	[1768-1786]
49	CUGUCUGCCUUCAGGUGAA	UUCACCUGAAGGCAGACAG	Cyn, Mnk	[659-677]
50	GGAUGUGGAC CACAGUACU	AGUACUGUGGUCCACAUC	Cyn, Rb, Mnk	[813-831]
51	UAGUCAAGGUGCUCAAUAA	UUAUUGAGCACCUUGACUA	Mnk	[3959-3977]
52	CCUUUGGGUCGACAUAGUA	UACUAUGUCGACCCAAAGG	Mnk	[4020-4038]
53	GCAAGGAGAUGUCUGAAUA	UAUUCAGACAUCUCCUUGC	Mnk	[1433-1451]
54	CCAUGGAAGUCUCAAGAU	AUCUUUGAGACUCCAUGG	Mnk	[2182-2200]
55	GAAGAAGCAAACAUGACUA	UAGUCAUGUUUGCUUCUUC		[2007-2025]
56	CCAGCUCCUUUUCUGUGAA	UUCACAGAAAAGGAGCUGG	Mnk	[1679-1697]
57	GAAGAGCCUGACAAGUGAA	UUCACUUGUCAGGCUCUUC	Mnk	[1760-1778]
58	UCCUCAACUUGCUGCCUAA	UUAGGCAGCAAGUUGAGGA	Cyn, Mnk	[407-425]
59	CAUAGUGGGCCUUCAUUAA	UUAAUGAAGGCCACUAUG		[3714-3732]
60	CAUCAGGAAACUCUAAAAA	UUUUUAGAGUUUCCUGAUG	Cyn, GP, Mnk	[253-271]
61	CCGUGGAGAUGAGACUGAU	AUCAGUCUCAUCUCCACGG	Cyn, GP, Chn, Mnk	[1107-1125]
62	UCAUCAGGAAACUCUAAAA	UUUUUAGAGUUUCCUGAUGA	Mnk	[252-270]
63	AGUCAAGGUGCUCAAUAAA	UUUAUUGAGCACCUUGACU	Mnk	[3960-3978]
64	GGUGCAAGGAGAUGUCUGA	UCAGACAUCUCCUUGCACC	Mnk	[1430-1448]
65	UGAAGAGCCUGACAAGUGA	UCACUUGUCAGGCUCUUCA	Mnk	[1759-1777]
66	GCCUACAGAACAAACCAA	UUUGGUUUGUUCUGUAGGC	Mnk	[1067-1085]
67	AAUCCUACCUUUUGAUAAA	UUUAUCAAAAGGUAGGAUU	Mnk	[3780-3798]
68	GCCUUUGGGUCGACAUAGU	ACUAUGUCGACCCAAAGGC	Mnk	[4019-4037]
69	CAGGAAUGUUUCAGCUGCA	UGCAGCUGAAACAUUCCUG	Mnk	[1738-1756]
70	GAAGCCCUUUGC CUGUAGA	UCUACAGGCAAAGGGCUUC	Mnk	[1695-1713]
71	GGGAGAGGGCAUAUAAAUU	AAUUUAUAUGCCCUCUCCC		[1802-1820]
72	CCAUCAUCUCUGCCUUUGA	UCAAGGCAGAGAUGAUGG	Mnk	[1645-1663]
73	UGGAAGGUGUCCAAAUUUA	UAAAUUUGGACACCUUCCA	Mnk	[1964-1982]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
74	CUCCCUAGACAAUAAAGAU	AUCUUUAUUGUCUAGGGAG	Cyn, Mnk	[636-654]
75	AAACGAGGUUCCUGGUACA	UGUACCAGGAACCUCGUUU	Cyn, Mnk	[1288-1306]
76	CAUCCUCAUCAUCAGGAAA	UUUCCUGAUGAUGAGGAUG	Mnk	[244-262]
77	CCAACUUCCUGUUCUUUA	UAAAGAACAGGGAAGUUGG	Mnk	[2662-2680]
78	CUCUAGUCACCCUCUUCAA	UUGAAGAGGGUGACUAGAG	Cyn, Rat, Ms, Mnk	[830-848]
79	CAAGGUGCUCAAUAAAUAU	AUAUUUAUUGAGCACCUUG	Mnk	[3963-3981]
80	CUUGCUGCAGUGUCCAGAA	UUCUGGACACUGCAGCAAG	Mnk	[2035-2053]
81	CGACAGAUACUGUGGAACA	UGUUCCACAGUAUCUGUCG	Cyn, Mnk	[617-635]
82	GAACCCUUCUGCUUUCUUA	AUAGAAAGCAGAAGGGUUC	Mnk	[3275-3293]
83	GACCAUCUCCUAUCUUUUA	UAAAAGAUAGGAGAUGGUC	Mnk	[1902-1920]
84	UGUGGACCACAGUACUCUA	UAGAGUACUGUGGUCCACA	Cyn, Rb, Mnk	[816-834]
85	AACACUCCCUGACAAUAA	UUAUUGUCUAGGGAGUGUU	Cyn, Mnk	[632-650]
86	GAAACUGCUCCAGCUCCAA	UUGGAGCUGGAGCAGUUUC		[1011-1029]
87	AGUGAUUGCUWUAUUACA	UGUAAUAAAAGCAAUCACU	Mnk	[1864-1882]
88	CCUCCAGUCUGUUCUCACA	UGUGAGAACAGACUGGAGG		[3039-3057]
89	UGUUCCUUUUGACAAUUA	UAUAUUGUCAAAAGGAACA	Mnk	[3800-3818]
90	AAGCACUGAGGGAGACCAA	UUGGUCUCCUCAGUGCUU	Cyn, Mnk	[458-476]
91	CAGCUGUUGUUGAGCGAAU	AUUCGCUCAACAACAGCUG		[295-313]
92	CAAAUUJAAUGUAGACAUU	AAUGUCUACAUUAAAUUUG		[1975-1993]
93	UGUCCAAAUUUAAUGUAGA	UCUACAUUAAAUUUGGACA		[1971-1989]
94	GCUUCAGCAUGUACUCCCA	UGGGAGUACAUGCUGAAGC	Cyn, Mnk	[519-537]
95	CAAAAUGUUCUUCAUCCA	UGGAUGAAGAACAUUUUUG	Rat, Ms, Mnk	[1082-1100]
96	GAAGUAUUUGAGAGAGAGA	UCUCUCUCUCAAUACUUC	Mnk	[4041-4059]
97	GAAGCCUUGCACUCCUGAA	UUCAGGAGUGCAAGGCUUC	Cyn, Mnk	[675-693]
98	AAUGGUGAUUUUCUUGC UAA	UUAGCAAGAAAUACCAU	Mnk	[3370-3388]
99	GCAUUUUUAGCUCCUUGAA	UUCAAGGAGCUAAAAUUGC	Mnk	[3872-3890]
100	GGCUCUGAAGAAGCAAACA	UGUUUGCUUCUUCAGAGCC		[2001-2019]
101	UCCUGAAUUUUUAUCAAACA	UGUUUGAUAAAAUUCAGGA	Mnk	[687-705]
102	CCGACAUGCUGGUUAAGGU	ACCUUAACCAGCAUGUCGG	Rb, Mnk	[1358-1376]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
103	UGGAGAGAAAGAACUGGAA	UUCCAGUUCUUUCUCUCCA	Cyn, Mnk	[780-798]
104	CUAGGCCUGUAGUAGUCAA	UUGACUACUACAGGCCUAG	Mnk	[3947-3965]
105	GAGAUAAACUCCUUCACAU	AUGUGAAGGAAGUUAUCUC	Mnk	[3905-3923]
106	CUCCCUUCUUUACCUUCAU	AUGAAGGUAAAGAAGGGAG		[2436-2454]
107	CUACAGAACAAACCAAAAA	UUUUUGGUUUGUUCUGUAG	Rat, Mnk	[1069-1087]
108	CAGGAUUUGGUUGUUUAAA	UUUAAACAACCAAAUCCUG		[3230-3248]
109	CUCUGAGCUUGUUAUCUGA	UCAGUAAACAAAGCUCAGAG	Dog, Mnk	[2217-2235]
110	UAUCCCUUGCUGUUAUCUGAAA	UUUCAGAUAAAGCAGGGAUA	Mnk	[2639-2657]
111	GGGCCUUCAUUAAAUGUUU	AAACAUUUAUGAAGGCC		[3720-3738]
112	GCUUUACAGGAUCAUGUAA	UUACAUGAUCCUGUAAAGC		[2497-2515]
113	CUAGAGUGAGAGUUUGGAA	UUCCAAACUCUCACUCUAG	Mnk	[1950-1968]
114	GGCUGUAAAUGAGGGCAGA	UCUGCCCUCAUUUACAGCC	Mnk	[3651-3669]
115	CGACUUGAGUCUCCCUUUU	AAAAGGGAGACUCAAGUCG	Mnk	[555-573]
116	CCUUGAACCUCAUUAAAUG	CAUUUAAUGAGGUUCAAGG	Mnk	[3355-3373]
117	AACCAAAAAUGUUCUUCAU	AUGAAGAACAUUUUUGGUU	Rat, Mnk	[1079-1097]
118	CCAGUCUGUUCUCACAUAA	UUAUGUGAGAACAGACUGG		[3042-3060]
119	CUGACAAGUGAAGUUGUAA	UUACAACUUCACUUGUCAG	Mnk	[1767-1785]
120	CAUGCAUCCUCAUCAUCAG	CUGAUGAUGAGGAUGCAUG	Mnk	[240-258]
121	GAGGAUAUGUUGCUCACCA	UGGUGAGCAACAUUCCUC	Cyn, Mnk	[490-508]
122	GGUUGACCAACAAGAUGGA	UCCAUCUUGUUGGUCAACC	Mnk	[1131-1149]
123	CCUGCACUUCAUUAUCUAGA	UCUAGAUUAUGAAGUGCAGG	Mnk	[2375-2393]
124	CACUCCUGAAUUUUUAUCAA	UUGAUAAAAUUCAGGAGUG	Mnk	[684-702]
125	GAAACUGCAGAAUUUUGCA	UGCAAAAUUCUGCAGUUUC	Mnk	[906-924]
126	GGCUCUUGCUCUAGUGUUU	AAACACUUGAGCAAGAGCC	Mnk	[1311-1329]
127	UCUGAAGAAGCAAACAUGA	UCAUGUUUGCUUCUUCAGA		[2004-2022]
128	GCUCCUUGAAGACAUAUCA	UGAUUAUGUCUUAAGGAGC	Mnk	[3881-3899]
129	GUAUUAAACUCUCCUUUGA	UCAAGGAGAGUUUAAUAC	Mnk	[2562-2580]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
130	GAGCGAAUUGUUAGAACA	AUGUUCUAACAAUUCGCUC		[306-324]
131	CCAAAUAAAUGCUCUAU	AAUAUGAGCAUUUUAUUUGG	Mnk	[4104-4122]
132	GUGAUGCCGGUAAAGAAAA	UUUUCUUUACCGGCAUCAC	Cyn, Mnk	[1184-1202]
133	CUUCAUAUCUAGAUUUCUA	UAGAAAUCUAGAUUGAAG	Mnk	[2381-2399]
134	CCUUGCAUUUGUACCUAU	AUAAGGUACAAAUGCAAGG	Mnk	[2153-2171]
135	AAACCUUUUAUCCUGCUUA	UAAGCAGGGAUAAAGGUUU	Mnk	[2632-2650]
136	GUAAAUGCUCAAAGAUGUA	UACAUCUUUGAGCAUUUAC		[2512-2530]
137	ACAUGCUGGUUAAGGUGAA	UUCACCUUAACCAGCAUGU	Cyn, Rb, Mnk	[1361-1379]
138	CCGCCCUUUUUGUCCUAGA	UCUAGGACAAAAGGGCGG		[1936-1954]
139	CUUUUUCUCCUGAGAAA	UUUCUCAGGAAGGAAAAAG	Mnk	[2313-2331]
140	UCAUUAAAUGUUUGUUGAA	UUCAACAAACAUUUAAUGA		[3726-3744]
141	GACAUAGUAUGGAAGUAU	AAUACUCCAUACUAUGUC	Mnk	[4030-4048]
142	GGGCUAUGACGUCCAUGUU	AACAUGGACGUCAUAGCCC	Cyn, Mnk	[855-873]
143	GAUAAAUGUCCUUUUGA	UCAAAAAGGAACAUUUUAUC	Mnk	[3793-3811]
144	CCAGUGGAAGGACACUCU	AAGAGUGUCCUCCACUGG	Mnk	[2090-2108]
145	GUGUGAUAGAGCCUUUGAU	AUCAAAGGCUCUAUCACAC	Mnk	[1557-1575]
146	AUCAGGAAACUCUAAAAAA	UUUUUUAGAGUUUCCUGAU	Cyn, Mnk	[254-272]
147	AGCUGUGACUACGACUUGA	UCAAGUCGUAGUCACAGCU	Rb, Mnk	[544-562]
148	CAGAAUCCACCGGUGCAA	UUGCACCGGUGGAAUUCUG		[1418-1436]
149	CAAUGUGCACUUCACUGGA	UCCAGUGAAGUGCACAUUG	Cyn, Ms, Dog, Mnk	[765-783]
150	CAUGUAAAUGCUCAAAGAU	AUCUUUGAGCAUUUACAUG		[2509-2527]
151	CCCUUGAACCUCAUUAAAU	AUUUAAUGAGGUUCAAGGG	Mnk	[3354-3372]
152	UGGUGUUGAGCAAUGUGCA	UGCACAUUGCUCACACCA	Cyn, Mnk	[755-773]
153	CAAUCCUACCUUUUGAUAA	UUAUCAAAGGUAGGAUUG	Mnk	[3779-3797]
154	UGGAAAUGAGGGAGCUCAU	AUGAGCUCCCUCAUUUCCA	Ms, Rb, Mnk	[350-368]
155	UGCUUUACAGGAUCAUGUA	UACAUGAUCCUGUAAAGCA		[2496-2514]
156	GUUGUUGAGCGAAUUGUUA	UAACAAUUCGCUACAAC		[300-318]
157	GCAUUUGUACCUUAUUGAU	AUCAUAAGGUACAAAUGC	Mnk	[2157-2175]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
158	AAUCAACCUGCACUUCAUA	UAUGAAGUGCAGGUUGAUU	Mnk	[2369-2387]
159	AGGAUUUGGUUGUUUAAAU	AUUUAAACAACCAAUCCU		[3231-3249]
160	AAGAGAAACUGCAGAAUUU	AAAUUCUGCAGUUUCUCUU	Mnk	[902-920]
161	AACCUUAGCUACAAUCCUA	UAGGAUUGUAGCUAAGGUU	Mnk	[3768-3786]
162	UGUUUCAGCUGCAGUUGAA	UUCAACUGCAGCUGAAACA	Mnk	[1744-1762]
163	CAUUUAUUGCAGUUUAUUAU	AUAUAAACUGCAAUAAAUG	Mnk	[3592-3610]
164	GAUUGCUUUUAUUACAUAU	UAAUGUAAUAAAAGCAAUC	Mnk	[1867-1885]
165	CACCAAUCUGUCUAGAAU	AUUCUAGACAGAUUUGGUG	Mnk	[2475-2493]
166	ACUUAUCAAGGAUCGGGAA	UUCCCGAUCCUUGAUAAAGU	Cyn, Mnk	[1383-1401]
167	CCACAGGAGGUGUGAUAGA	UCUAUCACACCUCUGUGG	Mnk	[1548-1566]
168	UGCUCAAAGAUGUAAUGUA	UACAUUACAUCUUUGAGCA		[2517-2535]
169	CUUUGAUCUUCAGGAUGCA	UGCAUCCUGAAGAUCAAAG	Mnk	[1569-1587]
170	GCUACUUUUACUUGUUCAU	AUGAACAAGUAAAAGUAGC	Mnk	[3615-3633]
171	UGUCUAGAAUCCUGCUUUA	UAAAGCAGGAUUCUAGACA	Mnk	[2484-2502]
172	UCACCUUGGUUUUAUGAGAA	UUCUCAUAAACCAAGGUGA	Mnk	[3320-3338]
173	AACCUUUUAUCCUGCUUUAU	AUAAGCAGGGAUAAAGGUU	Mnk, Mnk	[2633-2651]
174	GGAGAGAAAGAACUGGAAU	AUUCAGUUCUUUCUCUCC	Mnk	[781-799]
175	GAGGUUCCUGGUACAUCGA	UCGAUGUACCAGGAACCUC	Mnk	[1292-1310]
176	GUCGACAGAUACUGUGGAA	UUCCACAGUAUCUGUCGAC	Cyn, Mnk	[615-633]
177	GCAGUUUAUAUAUAUGCUA	UAGCAUAUAUAUAAACUGC		[3600-3618]
178	CACUUAUCAAGGAUCGGGA	UCCCGAUCCUUGAUAAAGUG	Cyn, Mnk	[1382-1400]
179	AGCCUACAGAACAAACCAA	UUGGUUUUGUUCUGUAGGCU	Mnk	[1066-1084]
180	ACACUCCCUAGACAAUAAA	UUUAUUGUCUAGGGAGUGU	Cyn, Mnk	[633-651]
181	ACAUUGCCUUGCAUUUGUA	UACAAAUGCAAGGCAAUGU		[2147-2165]
182	CUUGCACUCCUGAAUUUUA	UAAAAUUCAGGAGUGCAAG	Cyn, Mnk	[680-698]
183	ACAUCAUCACCUUGGAAAU	AUUUCCAAGGUGAUGAUGU	Cyn, Dog, Mnk	[338-356]
184	CACGCGCUCAGACAUGAUA	UAUCAUGUCUGAGCGCGUG	Cyn, Mnk	[1224-1242]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058 :148-1506
185	CUGGAGAAGGACAUCAUCA	UGAUGAUGUCCUUCUCCAG	Cyn, Rb, Dog, Mnk	[328-346]
186	CCUUUUAUGAUCUUUGCCCA	UGGGCAAAGAUAUAAAGG	Mnk	[2166-2184]
187	GGAACACUCCCUAGACAAU	AUUGUCUAGGGAGUGUUC	Cyn, Mnk	[630-648]
188	CCCUGAGCAGUCUACUUCU	AGAAGUAGACUGCUCAGGG	Mnk	[3537-3555]
189	CAUGCGBAACACCAAACGA	UCGUUUGGUGUUCGCAUG	Mnk	[1275-1293]
190	UAGUAGUCAAGGUGCUGAA	UUGAGCACCUUGACUACUA	Mnk	[3956-3974]
191	CCUGCUUUCUCUUUCAGUA	UACUGAAAGAGAAAGCAGG		[2546-2564]
192	CAGUUGAAGAGCCUGACAA	UUGUCAGGCUCUUAACUG	Mnk	[1755-1773]
193	UGGAAAGAACCACGCAGGA	UCCUGCGUGGUUCUUUCCA	Mnk	[1146-1164]
194	UGUUGUUGAGCGAAUUGUU	AACAAUUCGCUCAACAACA		[299-317]
195	GUGCAAGGAGAUGUCUGAA	UUCAGACAUCUCCUUGCAC	Mnk	[1431-1449]
196	UUUACAGGAUCAUGUAAAU	AUUUACAUGAUCCUGUAAA		[2499-2517]
197	CAGGAUGCACGGUUUCUGU	ACAGAAACCGUGCAUCCUG	Mnk	[1579-1597]
198	CAUAUCUCUCCCUUCUUUA	UAAAGAAGGGAGAGUAUG		[2429-2447]
199	CCCAUGGUCCCUAGCAAAA	UUUUGCUAGGGACCAUGGG	Mnk	[3926-3944]
200	UCAAGGUGCUCAAUAAAU	UAUUUAUUGAGCACCUUGA	Mnk	[3962-3980]
201	AGAAGGACAUCACACCUU	AAGGUGAUGAUGUCCUUCU	Cyn, Dog, Mnk	[332-350]
202	CCCAUGGAAGUCUCAAAGA	UCUUUGAGACUCCAUGGG	Mnk	[2181-2199]
203	ACUGGAAUUUCGCUCUGGA	UCCAGAGCGAAAUUCCAGU	Cyn, Mnk	[792-810]
204	UCUGAGCUUUGUUACUGAA	UUCAGUAACAAAGCUCAGA	Dog, Mnk	[2218-2236]
205	GCAUUUAUUGCAGUUUAUA	UAUAAACUGCAAUAAUUGC	Mnk	[3591-3609]
206	GUUUGGAAGGUGUCCAAAU	AUUUGGACACCUUCCAAAC	Mnk	[1961-1979]
207	AAGAAAAGUUGCCGAAGAU	AUCUUCGGCAACUUUUCUU	Mnk	[1196-1214]
208	CAUUAGUUAAGAUGUCUGA	UCAGACAUCUUAACUAAUG	Mnk	[1881-1899]
209	GCAGAAACCUUGUUUGUUU	AAACAAACAAGGUUUCUGC	Mnk	[3665-3683]
210	AACUCUCCUUUGAUUAUUA	AUAAUAUCAAGGAGAGUU	Mnk	[2568-2586]
211	AAAUGUUCUUUUGACAAU	AUUGUCAAAAGGAACAUUU	Mnk	[3797-3815]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
212	GCAAAAUGCUAGGCCUGUA	UACAGGCCUAGCAUUUUGC	Mnk	[3939-3957]
213	GCACAGGAAAUGCAAGAGA	UCUCUUGCAUUUCCUGUGC	Rat, Ms, GP, Chn, Mnk	[889-907]
214	GCACAUAGUGGGCCUUCAU	AUGAAGGCCCCACUAUGUGC	AUGAAGGCCCCACUAUGUGC	[3711-3729]
215	AGCUUAUCUCCUGCUUCA	UGAAGCAGGGAGAUAAAGCU	Mnk	[2412-2430]
216	UAUUGUUUUAACCUAGAA	UUCUAGGUUGAAAACAAUA		[2615-2633]
217	GUGGAACACUCCCUAGACA	UGUCUAGGGAGUGUCCAC	Cyn, Dog, Mnk	[628-646]
218	CUCCCUGCUUCAUAUCUCU	AGAGAUUGAAGCAGGGAG	Mnk	[2419-2437]
219	CAUUUAAAAUGUAAACUCU	AGAGUUUACAUUUUAAAUG	Mnk	[3301-3319]
220	CGACAUAGUAUGGAAGUUA	AUACUCCAUACUAUGUCG	Mnk	[4029-4047]
221	AAUGUUUGUUGAAUAAAAG	CUUUUAUUAACAAACAUAU	Mnk	[3732-3750]
222	CCAAAUUUAAUGUAGACAU	AUGUCUACAUUAAUUUGG		[1974-1992]
223	UGAGAUAAACUCCUUCACA	UGUGAAGGAAGUUAUCUCA	Mnk	[3904-3922]
224	GAGCCUUUGAUCUUCAGGA	UCCUGAAGAUCAAAGGCUC	Mnk	[1565-1583]
225	CCUUUAUCCUGCUUAUCU	AGAUAAAGCAGGGAUAAAGG	Mnk	[2635-2653]
226	CGAAUUGUUAGAACAUCUU	AAGAUGUUCUAACAAUUCG		[309-327]
227	CUUUCUCUUUCAGUAUUA	UUAAUACUGAAAGAGAAAG		[2550-2568]
228	GAGAGAGAGAACCUUUCUA	UGGAAAGGUUCUCUCUCUC	Mnk	[4050-4068]
229	CUGAGCUUUGUACUGAAA	UUUCAGUAACAAAGCUCAG	Mnk	[2219-2237]
230	AUGUUUGUUGAAUAAAAGA	UCUUUUUAUUAACAAACAU	Mnk	[3733-3751]
231	AGUUAAGAUGUCUGAGAGA	UCUCUCAGACAUCUUAACU	Mnk	[1885-1903]
232	CUUCCUCAAGUCUUGACAA	UUGUCAAGACUUGAGGAAG		[3501-3519]
233	5CUCCUAUCUUUUUUUUAU	AUGGAAAUAAAAGAUAGGAG	Mnk	[1908-1926]
234	GGUGUGAUAGAGCCUUUGA	UCAAAGGCUCUAUCACACC	Mnk	[1556-1574]
235	GGCAGAAACCUUGUUUGU	ACAAACAAGGUUUCUGCCC	Mnk	[3663-3681]
236	GAGAUGUCUGAAUACUGCA	UGCAGUAUUCAGACAUCUC	Mnk	[1438-1456]
237	CCCACUUUCAUUCAAUCAA	UUGAAUUGAAUGAAAGUGGG	Mnk	[2356-2374]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
238	CCACUUUCAUUCAAUCAAC	GUUGAUUGAAUGAAAGUGG	Mnk	[2357-2375]
239	CAUCUCUCUCCCUUACUUAU	AUAGUAAGGGAGAGAGAUG	Mnk	[2336-2354]
240	UGAGCGAAUUGUUAGAACA	UGUUCUAACAAUUCGCUCA		[305-323]
241	UUAGCUCCUUGAAGACAUA	UAUGUCUUCAAGGAGCUAA	Mnk	[3878-3896]
242	ACAGUUACCUGCACACCGA	UCGGUGUGCAGGUAACUGU	Mnk	[924-942]
243	ACAGGAUCAUGUAAAUGCU	AGCAUUUACAUGAUCCUGU		[2502-2520]
244	CGUUCUUUUCAGUACUGCA	UGCAGUACUGAAGGGAACG	Mnk	[2064-2082]
245	AAAUGCUAGGCCUGUAGUA	UACUACAGGCCUAGCAUUU	Mnk	[3942-3960]
246	GUGUGGAUGGGAAACUGCU	AGCAGUUUCCCAUCCACAC	Mnk	[1001-1019]
247	CUGUCUUUUUCCUCCUGA	UCAGGAAGGAAAAAGACAG	Mnk	[2309-2327]
248	CCUAGACAAUAAAGAUGGU	ACCAUCUUUAUUGUCUAGG	Cyn, Mnk	[639-657]
249	UGGAAGUAAUUGAGAGAGA	UCUCUCUCAAUACUCCA	Mnk	[4039-4057]
250	CAGCCUUGGUUGGACCUAU	AUAGGUCCAACCAAGGCUG		[1716-1734]
251	GAACGCACUUAUCAAGGAU	AUCCUUGAUAAAGUGCGUUC	Mnk	[1377-1395]
252	GCGAAUUGUUAGAACAUCU	AGAUGUUCUAACAAUUCGC		[308-326]
253	AAGCCUACAGAACAAACCA	UGGUUUUGUUCUGUAGGCUU	Mnk	[1065-1083]
254	CGCCCUUUUUGUCCUAGAG	CUCUAGGACAAAAAGGGCG		[1937-1955]
255	GAGCCUGACAAGUGAAGUU	AACUUCACUUGUCAGGCUC	Mnk	[1763-1781]
256	GCCUGACAAGUGAAGUUGU	ACAACUUCACUUGUCAGGC	Mnk	[1765-1783]
257	GCUGUAAAUGAGGGCAGAA	UUCUGCCCUCAUUUACAGC	Mnk	[3652-3670]
258	UGUUGAGCGAAUUGUUAGA	UCUAACAAUUCGCUCAACA		[302-320]
259	AGCUGUUGUUGAGCGAAUU	AAUUCGCUCAACAACAGCU		[296-314]
260	CUGUGCGUCCCUUCAGUA	UACUGAAGGGAACGCACAG	Mnk	[2059-2077]
261	UAGCACUGGUGUUGAGCAA	UUGCUC AACACCAGUGCUA	Mnk	[749-767]
262	CGUCCAUGUUCUAUGUGAC	GUCACAUAGAACAUGGACG	Mnk	[864-882]
263	AGACCAUCUCCUUCUUUU	AAAAGAUAGGAGAUGGUCU	Mnk	[1901-1919]
264	GCUUUUAUUACAUAUAGUUA	UAACUAAUGUAAUAAAAGC	Mnk	[1871-1889]
265	AGCCUUUGAUCUUCAGGAU	AUCCUGAAGAUCAAAGGCU	Mnk	[1566-1584]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058 :148-1506
266	GGUCCCUAGCAAAAUGCUA	UAGCAUUUUGCUAGGGACC	Mnk	[3931-3949]
267	CAUUAAAUGGUGAUUUCUU	AAGAAAUCACCAUUUAAUG	Mnk	[3365-3383]
268	CAACCUUAGCUACAAUCCU	AGGAUUGUAGCUAAGGUUG	Mnk	[3767-3785]
269	GCAGUUGAAGAGCCUGACA	UGUCAGGCUCUUAACUGC	Mnk	[1754-1772]
270	CAGAAACCUUGUUUGUUUU	AAAACAAACAAGGUUUCUG	Mnk	[3666-3684]
271	CAUCUUCUGGAGAAGGACA	UGUCCUUCUCCAGAAGAUG	Mnk	[322-340]
272	CUUUGAGUGUGGGACUCCA	UGGAGUCCACACUCAAAAG	Mnk	[1658-1676]
273	UGCAUUUAUUGCAGUUUAU	AUAAACUGCAAUAAUGCA	Mnk	[3590-3608]
274	GAACCUCAUUAAAUGGUGA	UCACCAUUUAAUGAGGUUC	Mnk	[3359-3377]
275	CGUUGGUUGUUUCUCUGAG	CUCAGAGAAACAACCAACG		[2205-2223]
276	UGACAAUAUACACGGAUUA	UAAUCCGUGUAUAUUGUCA	Mnk	[3809-3827]
277	GCCUUGCAUUUGUACCUUA	UAAGGUACAAAUGCAAGGC	Mnk	[2152-2170]
278	UCUUCACCUUGGUUUAUGA	UCAUAAACCAAGGUGAAGA	Mnk	[3317-3335]
279	UCAGGAAACUCUAAAAAAG	CUUUUUUAGAGUUUCCUGA	Cyn, Mnk	[255-273]
280	UCUCCUAUCUUUUUUUCA	UGAAAUAAAAGAUAGGAGA	Mnk	[1907-1925]
281	GAUCGGGAAGGUUAUGCU	AGCAUAACCUUCCCGAUCC	Mnk	[1392-1410]
282	AGUCUGUUCUCACAUACA	UGUUAUGUGAGAACAGACU		[3044-3062]
283	AAGUAUUUGAGAGAGAGAA	UUCUCUCUCUCAAAUACUU	Mnk	[4042-4060]
284	AAAUGAGGGAGCUCAUCCA	UGGAUGAGCUCCCUCAUUU	Ms, Rb, Mnk	[353-371]
285	GUUCCUGCUUUCUCUUUCA	UGAAAGAGAAAGCAGGAAC		[2543-2561]
286	UGAAUAAAAGAGGGAAGAA	UUCUUCCUCUUUUUAUUA	Mnk	[3741-3759]
287	GAGCAAUGUGCACUUCACU	AGUGAAGUGCACAUUGCUC	Cyn, Ms , Dog, Mnk	[762-780]
288	CUUUGGGUCGACAUAGUAU	AUACUAUGUCGACCCAAAG	Mnk	[4021-4039]
289	GAGAAACGUCCAUCUCUCU	AGAGAGAUGGACGUUUCUC	Mnk	[2326-2344]
290	UAUCUAGAUUUCUAGAAAA	UUUUCUAGAAAUCUAGUA	Mnk	[2386-2404]
291	UGAGCUUUGUUACUGAAAU	AUUUCAGUAACAAAGCUCA	Mnk	[2220-2238]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058 :148-1506
292	AUCAUGUAAAUGCUCAAA	UUUGAGCAUUUACAUGAUC	Mnk	[2506-2524]
293	GCUCAAAGAUGUAAUGUAG	CUACAUUACAUCUUUGAGC		[2518-2536]
294	CCAUCUCCUAUCUUUUUAUU	AAUAAAAGAUAGGAGAUGG	Mnk	[1904-1922]
295	GGCCUGUAGUAGUCAAGGU	ACCUUGACUACUACAGGCC	Mnk	[3950-3968]
296	GACCACAGGAGGUGUGAUA	UAUCACACCUCUGUGGUC	Mnk	[1546-1564]
297	AGGUUUUCAGCUCUUUGA	UCAAAGAGCUGAAAAACCU	Mnk	[1031-1049]
298	GGUUAAGGUGAACGCACUU	AAGUGCGUUCACCUUAACC	Mnk	[1368-1386]
299	CUACCAGGCUGUAAAUGAG	CUCAUUUACAGCCUGGUAG	Mnk	[3645-3663]
300	GUUUCCCUAGACUCUGUAA	UUACAGAGUCUAGGGAAAC	Mnk	[2282-2300]
301	AGGAGAGUGAUGCCGGUAA	UUACCGGCAUCACUCUCCU	Cyn, Mnk	[1178-1196]
302	UCCCAUGGUCCCUAGCAAA	UUUGCUAGGGACCAUGGGA	Mnk	[3925-3943]
303	CUGUAAAUGAGGGCAGAAA	UUUCUGCCCUCAUUUACAG	Mnk	[3653-3671]
304	GAGAGAAAGAACUGGAAUU	AAUCCAGUUCUUUCUCUC	Cyn, Mnk	[782-800]
305	CCAUGGUCCCUAGCAAAAU	AUUUUGCUAGGGACCAUGG	Mnk	[3927-3945]
306	GUGGUGUCCUUCUCCUCA	UGAGGAAGAAGGACACCAC		[3490-3508]
307	GCUCCAAGAGGUUUUUCAG	CUGAAAAACCUCUUGGAGC		[1023-1041]
308	CAUGUUCUAUGUGACCAGA	UCUGGUCACAUAGAACAUG	Mnk	[868-886]
309	UUGUUUGGGUGGUUUGUGA	UCACAAACCACCCAAACAA		[3982-4000]
310	GAAAAGUUGCCGAAGAUGA	UCAUCUUCGGCAACUUUUC	Mnk	[1198-1216]
311	AUAGUAUGGAAGUAUUUGA	UCAAUACUCCAUACUAU	Mnk	[4033-4051]
312	CACCGUUGAGCUGUGACUA	UAGUCACAGCUAACGGUG	Rat, Ms, Rb, GP, Dog, Mnk	[536-554]
313	CUUUAGUGAUUGCUUUUAU	AUAAAAGCAAUCACUAAAG	Mnk	[1860-1878]
314	CUUUACAGGAUCAUGUAAA	UUUACAUGAUCCUGUAAAG		[2498-2516]
315	AAACCUUGUUUGUUUUUAUU	AAUAAAACAAACAAGGUUU	Mnk	[3669-3687]
316	GAAGCAAACAUGACUAGAG	CUCUAGUCAUGUUUGCUUC		[2010-2028]
317	UUGCACUCCUGAAUUUUUAU	AUAAAAUUCAGGAGUGCAA	Cyn, Mnk	[681-699]
318	GCCUGUCGACAGAUACUGU	ACAGUAUCUGUCGACAGGC	Cyn, Mnk	[611-629]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
319	GCAUGCAUCCUCAUCAUCA	UGAUGAUGAGGAUGCAUGC	Mnk	[239-257]
320	CCUACCAGGCUGUAAAUGA	UCAUUUACAGCCUGGUAGG	Mnk	[3644-3662]
321	ACUGCCCCAAGCCUACAGAA	UUCUGUAGGCUUGGGCAGU	Mnk	[1058-1076]
322	UGUGAAGCACUGAGGGAGA	UCUCCCUCAGUGCUUCACA	Mnk	[454-472]
323	AGUUUGGAAGGUGUCCAAA	UUUGGACACCUUCCAAACU	Mnk	[1960-1978]
324	ACUGAGGGAGACCAAGCAA	UUGCUUGGUCUCCCUCAGU	Cyn, Mnk	[462-480]
325	GACAUCAUCACCUUGGAAA	UUUCCAAGGUGAUGAUGUC	Cyn, Dog, Mnk	[337-355]
326	GGUCGACAUAGUAUGGAAG	CUUCCAUAUAUGUCGACC	Mnk	[4026-4044]
327	AACUCUUCACCUUGGUUUA	UAAACCAAGGUGAAGAGUU	Mnk	[3314-3332]
328	CUGAGAGACCAUCUCCUUA	AUAGGAGAUGGUCUCUCAG	Mnk	[1896-1914]
329	CAAUAAAGAUGGUCCUGUC	GACAGGACCAUCUUUAUUG	Mnk	[645-663]
330	UUUGGAAGGUGUCCAAAUU	AAUUUGGACACCUUCCAAA	Mnk	[1962-1980]
331	AGCCUUGGUUGGACCUAUU	AAUAGGUCCAACCAAGGCU		[1717-1735]
332	GAAUGUUUCAGCUGCAGUU	AACUGCAGCUGAAACAUUC	Mnk	[1741-1759]
333	GCUCUGAAGAAGCAAACAU	AUGUUUGCUUCUUCAGAGC		[2002-2020]
334	CUGUGACUACGACUUGAGU	ACUCAAGUCGUAGUCACAG	Rb, Mnk	[546-564]
335	CUUGCAUUUGUACCUUAUU	AAUAAGGUACAAAUGCAAG	Mnk	[2154-2172]
336	UUAUCUCCCUGCUUCAUUA	AUAUGAAGCAGGGAGAUAA	Mnk	[2415-2433]
337	GAUCUUUGCCCAUGGAAGU	ACUUCCAUGGGCAAAGAUC	Mnk	[2173-2191]
338	CCUUUUUGUCCUAGAGUGA	UCACUCUAGGACAAAAAGG	Mnk	[1940-1958]
339	GUGGGAUUCUCCCAGACUU	AAGUCUGGGAGAUUCCAC	Mnk	[1615-1633]
340	CUCCUUGAAGACAUUAUCAU	AUGAUAUGUCUUAAGGAG	Mnk	[3882-3900]
341	CAUGUACUCCCACCGUUGA	UCAACGGUGGGAGUACAUG	Cyn, Mnk	[526-544]
342	AUUAACUCUCCUUUGAUA	UAUCAAGGAGAGUUUAAU	Mnk	[2564-2582]
343	CAAGUGAAGUUGUAAACAC	GUGUUUACAACUUCACUUG	Mnk	[1771-1789]
344	GCUUUCUCUUUCAGUAUUA	UAAUACUGAAAGAGAAAGC		[2549-2567]
345	CCUAGAAACCUUUAUCCCU	AGGGAUAAAGGUUUCUAGG	Mnk	[2627-2645]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
346	GCCCAUGGAAGUCUCAAG	CUUUGAGACUCCAUGGGC	Mnk	[2180-2198]
347	GCACUUUAGUGAUUGCUIU	AAAGCAAUCACUAAAGUGC	Mnk	[1857-1875]
348	CAUUAAAUGUUUGUUGAAU	AUUCAACAAACAUUUAAUG		[3727-3745]
349	UGCCCAAGCCUACAGAACA	UGUUCUGUAGGCUUGGGCA	Mnk	[1060-1078]
350	AGGUGUGAUAGAGCCUIU	JAAAGGCUCUAUCACACCUC	Mnk	[1554-1572]
351	CAAGCUUUUGGGCUAUGAC	GUCAUAGCCCAAAGCUUG	Cyn, Mnk	[846-864]
352	CCUUCUGAGAAACGUCCA	UGGACGUUUCUCAGGAAGG	Mnk	[2319-2337]
353	CCUGAAUUUUAUCAAAACAC	GUGUUUGAUAAAAUUCAGG	Mnk	[688-706]
354	GAGACCAUCUCCUAUCUIU	AAAGAUAGGAGAUGGUCUC	Mnk	[1900-1918]
355	CUGCACAGGAAUUGCAAGA	UCUUGCAUUUCCUGUGCAG	Cyn, Rat, GP, Chn, Mnk	[887-905]
356	GUCCCUGAGCAGUCUACUU	AAGUAGACUGCUCAGGGAC	Mnk	[3535-3553]
357	GGAAGUAUUUGAGAGAGAG	CUCUCUCUCAAAUACUCC	Mnk	[4040-4058]
358	CACGGAUUUAUUUUGUAC	GUACAAAUAUAUCCGUG	Mnk	[3819-3837]
359	UCAAGGAUCGGGAAGGUUA	UAACCUUCCCGAUCCUUGA	Mnk	[1388-1406]
360	GGUGAUUUCUUGCUAAGCU	AGCUUAGCAAGAAUACACC	Mnk	[3373-3391]
361	GAGAGACCAUCUCCUAUCU	AGAUAGGAGAUGGUCUCUC	Mnk	[1898-1916]
362	AAAUGGUGAUUUCUUGCUA	UAGCAAGAAUACCAUUU	Mnk	[3369-3387]
363	CUCAGGUGGUGUCCUUCUU	AAGAAGGACACCACCUGAG		[3485-3503]
364	AGAAAAGCUUCCUAGCUUA	UAAGCUAGGAAGCUUUUCU	Mnk	[2399-2417]
365	CCAACAAGAUGGAAAGAAC	GUUCUUUCCAUCUUGUUGG	Mnk	[1137-1155]
366	GGUGUCCAAAUUUAAUGUA	UACAUUAAAUUUGGACACC		[1969-1987]
367	CAAGGAGAUGUCUGAAUAC	GUUUUCAGACAUCUCCUUG	Mnk	[1434-1452]
368	GGCAUUUUUAGCUCCUUGA	UCAAGGAGCUAAAAUGCC	Mnk	[3871-3889]
369	UGACGUCCAUGUUCUAUGU	ACAUAGAACAUGGACGUCA	Mnk	[861-879]
370	GUGAUUGCUIUUUAUUACAU	AUGUAAUAAAAGCAAUCAC	Mnk	[1865-1883]
371	CAAGAGAAACUGCAGAAUU	AAUUCUGCAGUUUCUCUUG	Mnk	[901-919]
372	CCAACAUUGCCUUGCAUUU	AAAUGCAAGGCAAUGUUGG		[2144-2162]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
373	UCGACAUAGUAUGGAAGUA	UACUUCCAUAACUAUGUCGA	Mnk	[4028-4046]
374	GCUGGUUAAGGUGAACGCA	UGCGUUCACCUUAACCAGC	Mnk	[1365-1383]
375	UAGUGGGCCUUCAUUAAAU	AUUUAAUGAAGGCCACUA		[3716-3734]
376	AAAGAAAAGUUGCCGAAGA	UCUUCGGCAACUUUUCUUU	Mnk	[1195-1213]
377	CUAGAAUCCUGCUUUACAG	CUGUAAAGCAGGAUUCUAG		[2487-2505]
378	AGAGUGAUGCCGGUAAAGA	UCUUUACCGGCAUCACUCU	Cyn, Mnk	[1181-1199]
379	AGAAGCAAACAUGACUAGA	UCUAGUCAUGUUUGCUUCU		[2009-2027]
380	UGCUACUUUUACUUGUUA	UGAACAAGUAAAAGUAGCA	Mnk	[3614-3632]
381	UACAUGUUUUGUUUUCAA	UUGAAAACAAUAACAUGUA		[2608-2626]
382	ACAUAGUAUGGAAGUAUUU	AAAUACUCCAUAACUAUGU	Mnk	[4031-4049]
383	AGUUGAAGAGCCUGACAAG	CUUGUCAGGCUCUUAACU	Mnk	[1756-1774]
384	AUGGGAAACUGCUCCAGCU	AGCUGGAGCAGUUUCCCAU	Mnk	[1007-1025]
385	UGGAACACUCCCUAGACAA	UUGUCUAGGGAGUGUCCA	Cyn, Dog, Mnk	[629-647]
386	AGAAUGUGGAACUCCUCAA	UUGAGGAGUCCACAUCU	Rat, Ms, GP, Chn, Mnk	[395-413]
387	UCUUUUUCCUCCUGAGAA	UUCUCAGGAAGGAAAAAGA	Mnk	[2312-2330]
388	CCUUGAAGACAUUCAUGU	ACAUGAUUAGUCUUAAGG	Mnk	[3884-3902]
389	AAACAUGACUAGAGACGCA	UGCGUCUCUAGUCAUGUUU	Mnk	[2015-2033]
390	CCUCUCAGCUUCUAAUUUU	AAAAUUAGAAGCUGAGAGG	Mnk	[3408-3426]
391	GACAAUUAACACGGAUUU	AUAAUCCGUGUAUAUUGUC	Mnk	[3810-3828]
392	AUAUCUAGAUUUCUAGAAA	UUUCUAGAAAUCUAGAUU	Mnk	[2385-2403]
393	CCUAGAGUGAGAGUUUGGA	UCCAAACUCUCACUCUAGG	Mnk	[1949-1967]
394	UUCUGAGCGGGCUUGUGAU	AUCACAAGCCCGCUCAGAA	Mnk	[1329-1347]
395	GACGCAGGAUAUUGGGAGU	ACUCCCAAUAUCCUGCGUC	Cyn, Mnk	[215-233]
396	GGUGCUCAAUAAAUUUUG	CAAAUAUUUAUUGAGCACC	Mnk	[3966-3984]
397	CUGUGGAACACUCCCUAGA	UCUAGGGAGUGUCCACAG	Cyn, Dog, Mnk	[626-644]
398	AGAGAAACUGCAGAAUUUU	AAAAUUCUGCAGUUUCUCU	Mnk	[903-921]
399	UGUUGAAUAAAAGAGGGAA	UUCCCUCUUUUUUAUCAACA	Mnk	[3738-3756]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
400	GAGAGAGAACCUUCCACU	AGUGGAAAGGUUCUCUCUC	Mnk	[4052-4070]
401	UGAACGCACUUAUCAAGGA	UCCUUGAUAAAGUGCGUUA	Mnk	[1376-1394]
402	ACGCGCUCAGACAUGAUU	AUAUCAUGUCUGAGCGCGU	Cyn, Mnk	[1225-1243]
403	CUUCACUGGAGAGAAAGAA	UUCUUUCUCUCCAGUGAAG	Cyn, Mnk	[774-792]
404	CCUUGUUUGUUUUAUUCAC	GUGAAUAAAACAAACAAGG	Mnk	[3672-3690]
405	AGAACAAACCAAAAUGUU	AACAUUUUUGGUUUGUUCU	Rat, Mnk	[1073-1091]
406	CGUGGAGAUGAGACUGAUC	GAUCAGUCUCAUCCACG	Cyn, GP, Chn, Mnk	[1108-1126]
407	CACUGGUGUUGAGCAAUGU	ACAUUGCUCACACCAGUG	Cyn, Mnk	[752-770]
408	UGUAAAUGCUCAAAGAUGU	ACAUCUUUGAGCAUUUACA		[2511-2529]
409	CAUCAUCUCUGCCUUUGAG	CUCAAAGGCAGAGAUGAUG	Mnk	[1696-1664]
410	CUACCUUUUGAUAAAAUGU	ACAUUUUAUCAAAGGUAG	Mnk	[3784-3802]
411	AAGGUGAACGCACUUAUCA	UGAUAAAGUGCGUUCACCUU	Mnk	[1372-1390]
412	GAACACUCCUAGACAAUA	UAUUGUCUAGGGAGUGUUC	Cyn, Mnk	[631-649]
413	ACCAGACUGCACAGGAAAU	AUUUCCUGUGCAGUCUGGU	Cyn, Rat, GP, Chn, Dog, Mnk	[881-899]
414	ACAAGUGAAGUUGUAAACA	UGUUUACAACUUCACUUGU	Mnk	[1770-1788]
415	CAUUCUUUCUCCUCCAGU	ACUGGAGGAAGAAAGAAUG	Mnk	[3028-3046]
416	CAGACAUGAUUGCGGCUA	UAGCCGCAUAUCAUGUCUG	Cyn, Mnk	[1232-1250]
417	CAACCUGCACUUCAUAUCU	AGAUAGAAGUGCAGGUUG	Mnk	[2372-2390]
418	CUUUGUCCUGCUUUCUCU	AGAGAAAGCAGGAACAAAG		[2539-2557]
419	CUUCUUUACCUUCAUUUCA	UGAAAUGAAGGUAAAGAAG		[2440-2458]
420	AACCCUUCUGCUUUCUAUU	AAUAGAAAGCAGAAGGGUU	Mnk	[3276-3294]
421	ACUUCACUGGAGAGAAAGA	UCUUUCUCUCCAGUGAAGU	Cyn, Rat, Ms, Dog, Mnk	[773-791]
422	GUCUCCUCUCAGCUUCUAA	UUAGAAGCUGAGAGGAGAC	Mnk	[3404-3422]
423	GGAAAGAACCACGCAGGAU	AUCCUGCGUGGUUCUUUCC	Mnk	[1147-1165]
424	ACUCCCUAGACAAUAAAGA	UCUUUAUUGUCUAGGGAGU	Cyn, Mnk	[635-653]
425	GUAAAGAAAAGUUGCCGAA	UUCGGCAACUUUUCUUUAC	Mnk	[1193-1211]
426	CUAUCUUUUUUUCAUUCA	UGAAUGAAAUAAAAGAUAG	Mnk	[1911-1929]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058:148-1506
427	AACCUGCACUUCAUAUCUA	UAGAUUGAAGUGCAGGUU	Mnk	[2373-2391]
428	AGCGGGCUUGUGAUUAUGCA	UGCAUAUCACAAGCCCGCU	Mnk	[1334-1352]
429	UGCAAGAGAAACUGCAGAA	UUCUGCAGUUUCUCUUGCA	Mnk	[899-917]
430	CUGGAGAGAAAGAACUGGA	UCCAGUUCUUUCUCUCCAG	Mnk	[779-797]
431	AAUAUACACGGAUUAUUAU	AUAAUAAUCCGUGUAUAUU	Mnk	[3813-3831]
432	UCUGAGAGACCAUCUCCUA	UAGGAGAUGGUCUCUCAGA	Mnk	[1895-1913]
433	UCUAUGGUGUGGAUGGGAA	UUCCCAUCCACACCAUAGA	Mnk	[995-1013]
434	CAGAAUGUGGAACUCCUCA	UGAGGAGUUCACAUUCUG	Rat, Ms, GP, Chn, Mnk	[394-412]
435	ACAGAAUCCACCGGUGCA	UGCACCGGUGGAAUUCUGU		[1417-1435]
436	CUUACUAUUCCACUUUCA	UGAAAGUGGGAAUAGUAAG	Mnk	[2347-2365]
437	GCAAGAGAAACUGCAGAAU	AUUCUGCAGUUUCUCUUGC	Mnk	[900-918]
438	GAAUCUCCAGACUUGUUU	AAACAAGUCUGGGAGAUUC	Mnk	[1619-1637]
439	UGGAGAAGGACAUCAUCAC	GUGAUGAUGUCCUUCUCCA	Rb, Dog, Mnk	[329-347]
440	GAGAACCUUCCACUCCCA	UGGGAGUGGAAAGGUUCUC	Mnk	[4056-4074]
441	ACUUCCCUGUUCUUUAAGA	UCUUAAGAAGCAGGGAAGU	Mnk	[2665-2683]
442	GUGUUUCCCUAGACUCUGU	ACAGAGUCUAGGGAAACAC	Mnk	[2280-2298]
443	AGGAAAUGCAAGAGAAACU	AGUUUCUCUUGCAUUUCCU	Rat, Ms, GP, Chn, Mnk	[893-911]
444	UUCAGUCCAGCUUUUGUA	UACAAAAGCUGGAACUGAA	Mnk	[1834-1852]
445	CGCGCUCAGACAUGAUUAUG	CAUAUCAUGUCUGAGCGCG	Cyn, Mnk	[1226-1244]
446	AUGUCUGAAUACUGCAGCA	UGCUGCAGUAUUCAGACAU	Mnk	[1441-1459]
447	UGGUUAAGGUGAACGCACU	AGUGCGUUCACCUUAACCA	Mnk	[1367-1385]
448	UGCAUUUGUACCUUAUUGA	UCAAUAAGGUACAAAUGCA	Mnk	[2156-2174]
449	CCCAAGCCUACAGAACAAA	UUUGUUCUGUAGGCUUGGG	Mnk	[1062-1080]
450	CUUUCUCCUCCAGUCUGU	ACAGACUGGAGGAAGAAAG	Mnk	[3032-3050]
451	AAGAUGUCUGAGAGACCAU	AUGGUCUCUCAGACAUCUU	Mnk	[1889-1907]
452	GCCUUGCACUCCUGAAUUU	AAAUUCAGGAGUGCAAGGC	Cyn, Mnk	[678-696]
453	CUUUAUCCUCCUGCUUAUCUG	CAGAUAAAGCAGGGAUAAAG	Mnk	[2636-2654]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058 :148-1506
454	UGUUAUUGUUUUAACCUA	UAGGUUGAAAACAAUAACA		[2612-2630]
455	AAAAGAGGGAAGAAGGCAA	UUGCCUUCUUCCCUCUUUU	Mnk	[3746-3764]
456	GAACCUUUCACUCCCACU	AGUGGGAGUGGAAAGGUUC	Mnk	[4058-4076]
457	GUCCAAAUUUAAUGUAGAC	GUCUACAUUAAAUUUGGAC		[1972-1990]
458	AGUUC CAGCUUUUGUAGAU	AUCUACAAAAGCUGGAACU	Mnk	[1837-1855]
459	GGAGAU GUCUGAAUACUGC	GCAGUAUUCAGACAUCUCC	Mnk	[1437-1455]
460	ACAGCUGUUGUUGAGCGAA	UUCGCUCAACAACAGCUGU		[294-312]
461	CAUCUCCUAUCUUUUUUUU	AAAUAAAAGAUAGGAGAUG	Mnk	[1905-1923]
462	CCCUGCUUAUCUGAAACUU	AAGUUUCAGAUAAAGCAGGG	Mnk	[2642-2660]
463	GGUGAACGCACUUAUCAAG	CUUGAUAAAGUGCGUUCACC	Mnk	[1374-1392]
464	AACUGCUCCAGCUCCAAGA	UCUUGGAGCUGGAGCAGUU		[1013-1031]
465	UGAGCAAUGUGCACUUCAC	GUGAAGUGCACAUUGCUCU	Cyn, Ms, GP,Dog,Mnk	[761-779]
466	CAUCAUGUACCAAGUGCUU	AAGCACUUGGUACAUGAUG	Mnk	[3691-3709]
467	UGAAGAAGCAAACAUGACU	AGUCAUGUUUGCUUCUUA		[2006-2024]
468	UACAGAACAACCAAAAAU	AUUUUUGGUUUUGUUCUGUA	Rat, Mnk	[1070-1088]
469	GUACAUCGAGGCUCUUGCU	AGCAAGAGCCUCGAUGUAC	Mnk	[1302-1320]
470	GAGCUUUGUUACUGAAAUG	CAUUUCAGUAACAAAGCUC	Mnk	[2221-2239]
471	ACCCUUCUGCUUUCUAUUA	UAAUAGAAAGCAGAAGGGU	Mnk	[3277-3295]
472	UUUAGUGAUUGCUUUUAUU	AAUAAAAGCAAUCACUAAA	Mnk	[1861-1879]
473	AACAGCUGUUGUUGAGCGA	UCGCUCAACAACAGCUGUU		[293-311]
474	GUUUCUCUGAGCUUUGUUA	UAACAAAGCUCAGAGAAAC	Mnk	[2213-2231]
475	CUCUAAAAAAGAACCGAGU	ACUCGGUUCUUUUUUAGAG	Cyn, Pig, Mnk	[263-281]
476	GAGAGUGAUGCCGGUAAAG	CUUUACCGGCAUCACUCUC	Cyn, Mnk	[1180-1198]
477	AGAGGUUUUUCAGCUCUUU	AAAGAGCUGAAAACCUCU	Mnk	[1029-1047]
478	AGGAUAUGUUGCUCACCAC	GUGGUGAGCAACAUAUCCU	Cyn, Mnk	[491-509]
479	CCUUUUCUGUGAAGCCCUU	AAGGGCUUCACAGAAAAGG	Mnk	[1685-1703]
480	UGAUGCCGGUAAAAGAAAAG	CUUUUCUUUACCGGCAUCA	Cyn, Mnk	[1185-1203]

No.	Sense siRNA	AntiSense siRNA	Other Sp	Human-39995058 :148-1506
481	CCUUCUCCUCAAGUCUUG	CAAGACUUGAGGAAGAAGG		[3497-3515]
482	CUGAGAAACGUCCAUCUCU	AGAGAUGGACGUUUCUCAG	Mnk	[2324-2342]
483	GAUUUGGUUGUUUAAAU	AUAUUUAAACAACCAAUC		[3233-3251]
484	UAUUAACUCUCCUUUGAU	AUCAAGGAGAGUUUAAUA	Mnk	[2563-2581]
485	UUCUGGAGAAGGACAUCAU	AUGAUGCCUUCUCCAGAA	Cyn, Mnk	[326-344]
486	AGACUGCACAGGAAUGCA	UGCAUUUCCUGUGCAGUCU	Cyn, Rat, GP, Chn, Mnk	[889-902]
487	GCUCAAUAAAUUUUGUUU	AAACAAAUUUUUAUUGAGC		[3969-3987]
488	AAACCAAAAUGUUCUUCU	UGAAGAACAUUUUUGGUUU	Rat, Mnk	[1078-1096]
489	CUCUGAAGAAGCAAACAUG	CAUGUUUGCUUCUUCAGAG		[2003-2021]
490	CUGUAGUAGUCAAGGUGCU	AGCACCUUGACUACUACAG	Mnk	[3953-3971]
491	UGCUUUUUAUACAUAUAGU	AACUAAUGUAAUAAAAGCA	Mnk	[1870-1888]
492	GUGCGUCCCUUCAGUACU	AGUACUGAAGGGAACGCAC	Mnk	[2061-2079]
493	CAGUCCAGCUUUUGUAGA	UCUACAAAAGCUGGAACUG	Mnk	[1836-1854]
494	GACAAGUGAAGUUGUAAAC	GUUUACAACUUCACUUGUC	Mnk	[1769-1787]
495	UUGCCCAUGGAAGUCUCAA	UUGAGACUCCAUGGGCAA	Mnk	[2178-2196]
496	UAGCUACAAUCCUACCUUU	AAAGGUAGGAUUGUAGCUA	Mnk	[3773-3791]
497	CCAUGUUCUAUGUGACCAG	CUGGUCACAUAGAACAUGG	Mnk	[867-885]
498	GUAGUCAAGGUGCUGAAUA	UAUUGAGCACCUUGACUAC	Mnk	[3958-3976]
499	CCACCUGGAGGAUAUGUUG	CAACAUAUCCUCCAGGUGG	Cyn, Mnk	[983-501]
500	UCUGAGCGGGCUUGUGAUA	UAUCACAAGCCCGCUCAGA	Cyn, Mnk	[1330-1348]
501	GCCAGAAUGUGGAACUCCU	AGGAGUCCACAUUCUGGC		

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US61196995A \[0001\]](#)
- [US61198931A \[0001\]](#)
- [IL 2009000179W \[0001\]](#)
- [US6602856B \[0075\]](#)
- [WO0047599A \[0084\] \[0170\] \[0293\]](#)
- [WO9914226A \[0084\] \[0170\] \[0293\]](#)
- [WO9839352A \[0084\] \[0170\] \[0293\]](#)
- [WO2004083430A \[0094\] \[0170\]](#)
- [WO2006047842A \[0084\]](#)
- [US5225182A \[0137\]](#)
- [US5169383A \[0137\]](#)
- [US5167616A \[0137\]](#)
- [US4959217A \[0137\]](#)
- [US4925678A \[0137\]](#)
- [US4487603A \[0137\]](#)
- [US4486194A \[0137\]](#)
- [US4447233A \[0137\]](#)
- [US4447224A \[0137\]](#)
- [US4439196A \[0137\]](#)
- [US4475196A \[0137\]](#)
- [US4052985A \[0139\]](#)
- [WO2004015107A \[0152\] \[0165\] \[0166\]](#)
- [WO0244321A \[0152\] \[0165\]](#)
- [US5898031A \[0152\] \[0165\]](#)
- [US6107094A \[0152\] \[0165\]](#)
- [US20050080246A \[0165\]](#)
- [US20050042647A \[0165\]](#)
- [US6693187B \[0209\]](#)
- [US7067641B \[0209\]](#)
- [US20050004064A \[0295\]](#)
- [WO2007091268A \[0298\] \[0308\]](#)
- [WO2004041924A \[0299\]](#)
- [WO9323569A \[0306\]](#)
- [US6121426A \[0307\]](#)
- [US20040019001A \[0308\]](#)
- [WO0044895A \[0314\]](#)
- [WO0049035A \[0314\]](#)
- [WO0063364A \[0314\]](#)
- [WO0136841A \[0314\]](#)
- [WO0136646A \[0314\] \[0315\]](#)
- [WO9932619A \[0314\]](#)
- [WO0044914A \[0314\]](#)
- [WO0129058A \[0314\]](#)
- [WO0175164A \[0314\]](#)
- [US4666828A \[0323\]](#)
- [US4683202A \[0323\]](#)
- [US4801531A \[0323\]](#)
- [US5192659A \[0323\]](#)
- [US5272057A \[0323\]](#)

Non-patent literature cited in the description

- **DEL AMOURTTI**Drug Discov Today, 2008, vol. 13, 3/4135-143 [0004]
- **FATTALBOCHOT**Adv Drug Del Rev, 2006, vol. 56, 1203-1223 [0004]

- ELMEN et al.NAR, 2005, vol. 33, 1439-447 [0084] [0170]
- ALLART et al.Nucleosides & Nucleotides, 1998, vol. 17, 1523-1526 [0084] [0294]
- PEREZ-PEREZ et al.Bioorg. and Medicinal Chem Letters, 1996, vol. 6, 1457-1460 [0084]
- MANGANELLI FiettaSemin Arthritis Rheum., 2003, vol. 33, 149-65 [0106]
- HAMDIKENNEYFrontiers in Bioscience, 2003, 305-314 [0113]
- SHEN et al.FEBS Let., 2003, vol. 539, 111-114 [0136]
- XIA et al.Nat. Biotech., 2002, vol. 20, 1006-1010 [0136]
- REICH et al.Mol. Vision, 2003, vol. 9, 210-216 [0136]
- SORENSEN et al.J. Mol. Biol., 2003, vol. 327, 761-766 [0136]
- LEWIS et al.Nat. Gen., 2002, vol. 32, 107-108 [0136]
- SIMEONI et al.NAR, 2003, vol. 31, 112717-2724 [0136]
- BAR-ILANNEUMANN et al.Textbook of Ocular PharmacologyLippencott-Raven19970000 [0138]
- UI-TEI et al.J Biomed Biotechnol., 2006, 65052- [0152]
- CHALK et al.BBRC., 2004, vol. 319, 1264-74 [0152] [0164]
- SIOUDLEIRDALMet. Mol Biol., 2004, vol. 252, 457-69 [0152] [0164]
- LEVENKOVA et al.Bioinform., 2004, vol. 20, 3430-2 [0152] [0164]
- UI-TEI et al.NAR., 2004, vol. 32, 3936-48 [0152] [0164]
- BRAASCH et al.Biochem., 2003, vol. 42, 267967-75 [0152] [0165]
- CHIU et al.RNA., 2003, vol. 9, 91034-48 [0152]
- CZAUDERNA et al.NAR., 2003, vol. 31, 112705-2716 [0153]
- UI-TEI et al.J Biomed Biotech. 2006, 2006, 65052- [0164]
- CHIU et al.RNA, 2003, vol. 9, 91034-48 [0165]
- BOUTLA et al.Curr. Biol., 2001, vol. 11, 1776-1780 [0166]
- SCHWARZ et al.Mol. Cell, 2002, vol. 10, 537-48 [0166]
- AMARZGUI OUI et al.NAR, 2003, vol. 31, 2589-95 [0166]
- HOLEN et al.NAR, 2003, vol. 31, 92401-07 [0166]
- CHIURANARNA, 2003, vol. 9, 1034-48 [0166]
- CZAUDERNA et al.NAR, 2003, vol. 31, 112705-16 [0166]
- TAKEI et al.JBC, 2002, vol. 277, 2623800-06 [0171]
- STERNBERGER et al.Antisense Nucleic Acid Drug Dev, 2002, vol. 12, 131-43 [0176]
- HERDEWIJN et al.Nucleosides & Nucleotides, 1999, vol. 18, 1371-1376 [0294]
- FISHER et al.NAR, 2007, vol. 35, 41064-1074 [0294]
- UI-TEINAR, 2008, vol. 36, 72136-2151 [0295]
- BEAUCAGELYERTetrahedron, 1992, vol. 48, 2223-2311 [0304]
- BEAUCAGEIYERTetrahedron, 1993, vol. 49, 6123-6194 [0304]
- CARUTHERSMETHODS Enzymol., 1987, vol. 154, 287-313 [0304]
- ECKSTEINAnnu. Rev. Biochem., 1985, vol. 54, 367-402 [0304]
- USMAN et al.J. Am. Chem. Soc., 1987, vol. 109, 7845- [0305]
- SCARINGE et al.NAR, 1990, vol. 18, 5433- [0305]
- WINCOTT et al.NAR, 1995, vol. 23, 2677-2684 [0305]
- WINCOTT et al.Methods Mol. Bio., 1997, vol. 74, 59- [0305]
- MOORE et al.Science, 1992, vol. 256, 9923- [0306]
- SHABAROVA et al.NAR, 1991, vol. 19, 4247- [0306]
- BELLON et al.Nucleosides & Nucleotides, 1997, vol. 16, 951- [0306]
- BELLON et al.Bioconjugate Chem, 1997, vol. 8, 204- [0306]
- ELBASHIR et al.Genes Dev., 2001, vol. 15, 2188-200 [0315]
- BERNSTEIN et al.Nature, 2001, vol. 409, 6818363-6 [0315]
- LEE et al.Nature, 2003, vol. 425, 6956415-9 [0315]
- MCMANUSSHARPNature Rev Genet, 2002, vol. 3, 10737-47 [0315]
- PADDISONHANNONCurr Opin Mol Ther., 2003, vol. 5, 3217-24 [0315]
- BERNSTEIN et al.RNA, 2001, vol. 7, 111509-21 [0315]
- NISHIKURACell, 2001, vol. 107, 4415-8 [0315]
- PADDISON et al.PNAS USA, 2002, vol. 99, 1443-1448 [0316]
- PADDISON et al.Genes & Dev, 2002, vol. 16, 948-958 [0316]
- SUI et al.PNAS USA, 2002, vol. 8, 5515-5520 [0316]
- BRUMMELKAMP et al.Science, 2002, vol. 296, 550-553 [0316]
- SAMBROOK et al.Molecular cloning: A laboratory manualCold Springs Harbor Laboratory19890000 [0323]
- AUSUBEL et al.Current Protocols in Molecular BiologyJohn Wiley and Sons19880000 [0323]

- **AUSUBEL et al.** Current Protocols in Molecular Biology John Wiley and Sons 1989 0000 [0323]
- **PERBALA** Practical Guide to Molecular Cloning John Wiley & Sons 1988 0000 [0323]
- **Recombinant DNA WATSON et al.** Scientific American Books, New York [0323]
- **Genome Analysis: A Laboratory Manual Series** Cold Spring Harbor Laboratory Press 1998 0000 vol. 1-4, [0323]
- **PCR Protocols: A Guide To Methods And Applications** Academic Press 1990 0000 [0323]
- **TESTONI et al.** Blood, 1996, vol. 87, 3822- [0323]
- **CZAUDERNA et al.** NAR, 2003, vol. 31, 670-82 [0324]
- **Methods Find Exp Clin Pharmacol.**, 1997, vol. 19, 4231-9 [0324]
- **PEASE et al.** J. Glaucoma, 2006, vol. 15, 6512-9 [0327]

Patentkrav

1. Dobbelstrenget RNA-forbindelse, der har strukturen:
5' iB - GCCAGAAUGUGGAACUCCU 3' (sensestreng; SEQ ID NO: 9015)
5 3' CGGUCUUACACCUUGAGGA 5' (antisensestreng; SEQ ID NO: 9516)
hvert hver A, C, U og G er et ribonukleotid og hvert konsekutivt ribonukleotid er forbundet med det næste ribonukleotid ved en phosphodiesterbinding;
hvor sensestrengen omfatter, tællende fra 5'-enden, et umodificeret ribonukleotid ved positionerne 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 og 19, et L-deoxycytidin ved
10 position 18, og en inverteret deoxyabasisk del (iB) 5'-cap; og
hvor antisensestrengen omfatter, tællende fra 5'-enden, et 2'-O-Me sukkermodificeret ribonukleotid ved positionerne 2, 4, 6, 8, 11, 13, 15, 17 og 19 og et umodificeret ribonukleotid ved positionerne 1, 3, 5, 7, 9, 10, 12, 14, 16 og 18; eller et farmaceutisk acceptabelt salt deraf.
- 15 2. Forbindelse ifølge krav 1, hvor den inverterede deoxyabasiske del (iB) som 5'-cap er en inverteret abasisk deoxyribosedel.
3. Forbindelse ifølge krav 1, hvor den inverterede deoxyabasiske del (iB) som 5'-cap er et 3',5' inverteret abasisk deoxyribose 5'-phosphat.
- 20 4. Forbindelse ifølge et hvilket som helst af kravene 1-3 til anvendelse i terapi.
5. Forbindelse ifølge et hvilket som helst af kravene 1-3 til anvendelse i terapi til behandling af en patient, der lider af en øjensygdom, en øjenlidelse, en øjenlæsion, et synsfelttab; eller til
25 tilvejebringelse af øjenneurobeskyttelse.
6. Forbindelse til anvendelse ifølge krav 5, hvor øjensygdommen, øjenlidelsen eller øjenlæsion er forbundet med patologiske anormaliteter/ændringer i vævet i synssystemet; og/eller hvor øjensygdommen, lidelsen eller læsionen omfatter neurodegeneration; og/eller hvor
30 øjensygdommen, lidelsen eller læsionen er forbundet med tab af retinale ganglieceller eller med læsion på retinale ganglieceller.
7. Forbindelse til anvendelse ifølge krav 6, hvor tabet af retinal gangliecelle eller læsionen på retinal gangliecelle er medieret af et forhøjet intraokulært tryk (IOP).
- 35 8. Forbindelse til anvendelse ifølge krav 5, hvor synsfelttabet er forbundet med retinal gangliecelledød.

9. Forbindelse til anvendelse ifølge krav 5, hvor øjensygdommen, øjenlidelsen eller øjenlæsionen er udvalgt fra gruppen bestående af okulær neuropati, forhøjet intraokulært tryk (IOP), glaukom, tørøje, Sjögrens Syndrom, diabetisk retinopati (DR), diabetisk makulært ødem (DME),

5 aldersrelateret makulær degeneration (AMD), opticusneuritis, retinal centralveneokklusion, retinal veneforgrningsokklusion, iskæmisk opticus neuropati, optisk nervelæsion, præmatur retinopati (ROP), retinitis pigmentosa (RP), retinal gangliedegeneration, makuladedegeneration, arvelig opticusneuropati, Lebers hereditære opticusneuropati, metabolisk opticusneuropati, neuropati på grund af et toksisk middel og neuropati forårsaget af en bivirkning eller en

10 vitaminmangel.
10. Sammensætning til anvendelse ifølge krav 9, hvor den iskæmiske opticusneuropati omfatter anterior iskæmisk opticusneuropati og/eller posterior iskæmisk opticusneuropati.
- 15 11. Sammensætning, der omfatter en forbindelse ifølge et hvilket som helst af kravene 1-3 eller et farmaceutisk acceptabelt salt deraf og en farmaceutisk acceptabel excipiens eller bærer eller blanding deraf.
- 20 12. Sammensætning, der omfatter en forbindelse ifølge et hvilket som helst af kravene 1-3 i en mængde effektiv til at nedregulere ekspresion af et CASP2-gen i en øjencelle udvalgt fra en gruppe bestående af en acinuscelle fra tårekirtlen, en duktal celle fra tårekirtlen, en retinal gangliecelle (RGC), en retinal pigmentepitel- (RPE) celle, en choroidacelle, en hornhindecelle, en celle fra den ciliære proces og en celle fra trabekelværket eller en kombination deraf.
- 25 13. Sammensætning ifølge krav 11, hvor forbindelsen er til stede ved en endelig koncentration på omkring 5 µg/µl til omkring 60 µg/µl i volumen af sammensætningen.
- 30 14. Sammensætning ifølge krav 11, hvor sammensætningen er formuleret som en creme, skum, en pasta, en salve, en emulsion, en flydende opløsning, øjendråber, en gel, spray, en suspension, en mikroemulsion, mikrosfærer, mikrokapsler, nanosfærer, nanopartikler, lipidvesikler, liposomer, polymeriske vesikler, et plaster eller en kontaktlinse.
15. Sammensætning ifølge krav 14, hvor sammensætningen er formuleret som øjendråber.
- 35 16. Sammensætning ifølge krav 14, hvor sammensætningen er formuleret som en flydende opløsning til intravitreal injektion.

DRAWINGS

Figure 1A




Figure 1B




Figure 1C




Figure 2A




Figure 2B




Figure 2C




Figure 2D




Figure 3A – 3C




Figure 4




Figure 5A




Figure 5B




Figure 6A




Figure 6B




Figure 7




Figure 8A




Figure 8B




Figure 8C




Figure 9A




Figure 9B




Figure 10A




Figure 10B




Figure 11





Figure 12A




Figure 12B




Figure 13A




Figure 13B




Figure 13C




Figure 13D

