



(86) Date de dépôt PCT/PCT Filing Date: 2011/07/27
(87) Date publication PCT/PCT Publication Date: 2012/07/02
(45) Date de délivrance/Issue Date: 2019/09/24
(85) Entrée phase nationale/National Entry: 2013/01/21
(86) N° demande PCT/PCT Application No.: US 2011/045524
(87) N° publication PCT/PCT Publication No.: 2012/015912
(30) Priorité/Priority: 2010/07/30 (US61/369,148)

(51) Cl.Int./Int.Cl. *C07K 1/18* (2006.01),
A61K 38/16 (2006.01), *A61K 47/68* (2017.01),
C07K 1/107 (2006.01), *C07K 14/21* (2006.01),
C07K 16/00 (2006.01), *C07K 16/28* (2006.01),
C07K 19/00 (2006.01)

(72) Inventeurs/Inventors:
LINKE, THOMAS, US;
WANG, WILLIAM K., US;
SHAH, AMBARISH, US;
SATHISH, HASIGE, US;
HUNTER, ALAN, US;
THOMPSON, CHRISTOPHER, US

(73) Propriétaire/Owner:

(54) Titre : PROCEDE DE PURIFICATION D'IMMUNOCONJUGUES OU DE POLYPEPTIDES ACTIFS
(54) Title: METHOD FOR PURIFYING ACTIVE POLYPEPTIDES OR IMMUNOCONJUGATES

(57) **Abrégé/Abstract:**

The present invention provides methods for isolating an active polypeptide or immunoconjugate by purification of a solution containing both the active polypeptide or immunoconjugate and an acidic variant thereof, such as a deamidated variant, using anion exchange chromatography.

(73) Propriétaires(suite)/Owners(continued):MEDIMMUNE, LLC, US

(74) Agent: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
2 February 2012 (02.02.2012)



(10) International Publication Number
WO 2012/015912 A1

(51) International Patent Classification:
B01D 15/04 (2006.01) *B01J 41/20* (2006.01)

(21) International Application Number:
PCT/US2011/045524

(22) International Filing Date:
27 July 2011 (27.07.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/369,148 30 July 2010 (30.07.2010) US

(71) Applicant (for all designated States except US): **MED-IMMUNE, LLC** [US/US]; One MedImmune Way, Gaithersburg, MD 20878 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LINKE, Thomas** [US/US]; 933 Isaac Chaney Court, Odenton, MD 21113 (US). **WANG, William, K.** [US/US]; 1404 Romeo Court, Mclean, VA 22102 (US). **SHAH, Ambarish** [IN/US]; 18306 Crestmount Road, Boyds, MD 20841 (US). **SATHISH, Hasige** [IN/US]; 12906 Barleycorn Terrace, Germantown, MD 20874 (US). **HUNTER, Alan** [US/US]; MedImmune, LLC, One MedImmune Way, Gaithersburg, MD 20878 (US). **THOMPSON, Christopher** [US/US]; MedImmune, LLC, One MedImmune Way, Gaithersburg, MD 20878 (US).

(74) Agents: **MUKHERJEE, Mita** et al.; MedImmune, LLC, One MedImmune Way, Gaithersburg, MD 20878 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: METHOD FOR PURIFYING ACTIVE POLYPEPTIDES OR IMMUNOCOJUGATES

(57) Abstract: The present invention provides methods for isolating an active polypeptide or immunoconjugate by purification of a solution containing both the active polypeptide or immunoconjugate and an acidic variant thereof, such as a deamidated variant, using anion exchange chromatography.

WO 2012/015912 A1

METHOD FOR PURIFYING ACTIVE POLYPEPTIDES OR IMMUNOCONJUGATES

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention provides methods for purifying an active polypeptide or immunoconjugate from a solution containing the polypeptide or immunoconjugate and an acidic variant thereof, wherein said acidic variant is a deamidated species of said polypeptide or immunoconjugate. The present invention also provides formulations containing such purified polypeptides or immunoconjugates.

Background Art

[0002] The large-scale, economic purification of proteins is a factor for the biopharmaceutical industry. Therapeutic proteins are typically produced using prokaryotic or eukaryotic cell lines that are engineered to express the protein of interest from a recombinant plasmid containing the gene encoding the protein. Separation of the desired protein from the mixture of components fed to the cells and cellular by-products to an adequate purity, *e.g.*, sufficient for use as a human therapeutic, poses a formidable challenge to biologics manufacturers for several reasons.

[0003] Manufacturers of protein-based pharmaceutical products must comply with strict regulatory standards, including extremely stringent purity requirements. To ensure safety, regulatory agencies, such as Food and Drug Administration (FDA), require that protein-based pharmaceutical products are substantially free from impurities, including both product related contaminants such as aggregates, fragments and variants of the recombinant protein and process related contaminants such as host cell proteins, media components, viruses, DNA and endotoxins. While various protein purification schemes are available and widely used in the biopharmaceutical industry, they typically include an affinity-purification step, such as Protein A purification in the case of antibodies, in order to reach a pharmaceutically acceptable degree of purity.

[0004] The development of a purification scheme applicable to a particular biomolecule or various biomolecules that is scalable, controllable, and that strategically employs the use of particular resins or a combination of resins will allow its integration into product

- 2 -

development at a very early stage in overall drug development. This approach to the design of a purification scheme can minimize costly changes to manufacturing processes which may otherwise be necessary later in drug development or, worse, after approval. As the process is scaled-up and approaches GMP production conditions, additional inherent complexities arise, including those associated with resin packing and buffer preparation. The manufacturing process, and its capacity, can be improved by simplifying the purification scheme, by eliminating process steps and maximizing throughput and productivity, while maintaining the integrity and purity of the molecule that is being purified. Therefore, it would be desirable and advantageous to start with a simple and efficient process that can produce a drug substance of high quality and safety.

[0005] One complexity associated with the purification of a drug product is the maintenance of potency throughout the purification process. Many factors can contribute to a reduction or inhibition of potency, including the modification of the drug product during the development process. Such modification can occur at various stages of the process, for example, when the protein is being expressed in the cell, or when a protein that has been isolated from a cell is subject to various conditions or buffers. The present invention provides a method for purifying an active polypeptide or immunoconjugate from a solution containing a modified variant of the polypeptide or immunoconjugate, where the presence of this modified variant results in an inhibition in potency of the final drug product.

BRIEF SUMMARY OF THE INVENTION

[0006] The present invention provides a method of purifying a polypeptide of interest from a solution containing the polypeptide and an acidic variant, such as a deamidated variant, of the polypeptide.

[0007] In particular, the present invention provides a method of purifying an active immunoconjugate, where the immunoconjugate is deamidated at one or more residues, and wherein the deamidation results in an inhibition of potency of said immunoconjugate, the method comprising: (a) contacting the immunoconjugate with an anion exchange AEX chromatography matrix; and (b) eluting the bound immunoconjugate from the AEX chromatography matrix with a high salt buffer, thereby separating the active immunoconjugate from the deamidated variant.

[0008] The invention also provides a method of producing a purified polypeptide from a solution comprising the polypeptide and an acidic variant of the polypeptide, where the acidic variant of the polypeptide results in an inhibition of potency of the polypeptide, the method comprising: (a) contacting the polypeptide with an anion exchange (AIEX) chromatography matrix; and (b) eluting the bound polypeptide from the AIEX chromatography matrix with a high salt buffer, thereby separating said polypeptide from the acidic variant and producing a purified polypeptide.

[0009] The invention further provides a method of producing a purified polypeptide or immunoconjugate from a solution comprising the polypeptide and an acidic variant of the polypeptide, the method comprising: (a) producing the polypeptide or immunoconjugate in a bacterial cell which expresses the polypeptide or immunoconjugate; (b) isolating inclusion bodies containing the polypeptide or immunoconjugate from the bacterial cells; (c) refolding the polypeptide or immunoconjugate isolated from the inclusion bodies; (d) contacting the composition containing the polypeptide or immunoconjugate with an AIEX chromatography matrix; and (e) eluting the bound polypeptide or immunoconjugate from the AIEX chromatography matrix with a high salt buffer, thereby purifying the polypeptide or immunoconjugate from the solution.

[0010] In certain embodiments, the acidic variant is a deamidated variant. In further embodiments, between about 75 to about 99% of the acidic or deamidated variant is removed during the purification process, in particular about 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99%.

[0011] The AIEX matrix of the invention contains quaternary amine or tertiary amine ion exchange groups, a quaternary amino (Q) group. In certain embodiments, the AIEX matrix is Q sepharose.TM

[0012] The polypeptide or immunoconjugate of the invention is eluted with a linear or step salt gradient. In certain embodiments, the linear salt gradient is from about 150mM NaCl in Tris/HCl, pH 8.0 to about 300 mM NaCl in Tris/HCl, pH 8.0, from about 175mM NaCl in Tris/HCl, pH 8.0 to about 275 mM NaCl in Tris/HCl, pH 8.0, or from about 192mM NaCl in Tris/HCl, pH 8.0 to about 245 mM NaCl in Tris/HCl, pH 8.0.

[0013] In one embodiment, the polypeptide or immunoconjugate of the invention comprises an antibody or antigen binding fragment thereof, where the antibody or antigen binding fragment comprises a Fab, a Fab', a F(ab')₂, a Fd, a single chain Fv or scFv, a

disulfide linked Fv, a V NAR domain, an IgNar, an intrabody, an IgGΔCH2, a minibody, a F(ab')₃, a tetrabody, a triabody, a diabody, a single-domain antibody, DVD-Ig, Fcab, mAb2, a (scFv)₂, or a scFv-Fc. In certain embodiments, the antibody or antigen binding fragment binds a cell surface receptor, such as the cell surface receptor is CD22. In further embodiments, the antibody or antigen binding fragment thereof comprises a VH and VL sequence, where the VH sequence is selected from the group consisting of SEQ ID NOs: 6-11 and the VL sequence is selected from the group consisting of SEQ ID NOs: 2, and 12-15.

[0014] In another embodiment, the polypeptide or immunoconjugate comprises a toxin, where the toxin is selected from the group consisting of: *Pseudomonas* exotoxin, ricin, abrin, diphtheria toxin and subunits thereof, as well as botulinum toxins A through F or variants, or derivatives thereof. In certain embodiments, the *Pseudomonas* exotoxin, or variant thereof has an amino acid sequence selected from the group consisting of SEQ ID NOs:16-22. In a particular embodiment, the immunoconjugate is the CAT-8015 immunotoxin comprising the VH-PE38 subunit of SEQ ID NO:1 and the VL subunit of SEQ ID NO:2.

[0015] The invention also provides a composition comprising a purified immunoconjugate having less than between about 25% and about 1% deamidated species, wherein said immunoconjugate is purified by any of the methods described above. The composition can have less than about 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, or 1% of the deamidated species present. In certain embodiments, the composition is a pharmaceutical composition comprising a purified polypeptide or immunoconjugate and a pharmaceutically acceptable carrier.

[0016] The invention also provides a formulation comprising 1 mg/mL CAT-8015 in 25 mM sodium phosphate, 4% sucrose, 8% glycine, 0.02% polysorbate 80 (PS80), pH 7.4. In further embodiments, the formulation is lyophilized.

[0017] The invention also provides a method of modifying the bioactivity of a polypeptide solution comprising a polypeptide and a deamidated variant, the method comprising separating the polypeptide from the deamidated variant by linear elution AIEX chromatography; and combining the purified polypeptide and deamidated variant in fixed quantities to obtain the desired bioactivity of the polypeptide solution.

[0017a] According to one aspect of the present invention, there is provided a method of producing a purified polypeptide from a solution comprising the polypeptide and a deamidated species of the polypeptide, wherein said deamidated species of the polypeptide results in an inhibition of potency of said polypeptide, the method comprising: (a) contacting the polypeptide with an anion exchange (AIEX) chromatography matrix; and (b) eluting the bound polypeptide from the AIEX chromatography matrix with a linear salt gradient that is from about 150 mM NaCl in Tris/HCl, pH 8.0 to about 300 mM NaCl in Tris/HCl, pH 8.0, thereby separating said polypeptide from the deamidated species of said polypeptide and producing a purified polypeptide; wherein the polypeptide is an immunoconjugate comprising an antibody or antigen binding fragment thereof that binds the cell surface receptor CD22 and comprising a *Pseudomonas* exotoxin, or variant thereof.

[0017b] According to another aspect of the present invention, there is provided a method of modifying the bioactivity of a polypeptide solution comprising a polypeptide and a deamidated species of the polypeptide, the method comprising separating the polypeptide from the deamidated species by linear elution AIEX chromatography; and combining the purified polypeptide and deamidated species in fixed quantities to obtain the desired bioactivity of the polypeptide solution.

BRIEF DESCRIPTIONS OF THE DRAWINGS

- [0018] Figure 1. A graph depicting an ion exchange chromatography (IEC) profile of a CAT-8015 reference standard. The pre-peak of CAT-8015 represents the majority of inactive deamidated, or iso-deamidated CAT-8015, while the main peak contains the majority of the active, intact CAT-8015 immunoconjugate.
- [0019] Figure 2. A graph depicting the correlation between the percent of relative potency of CAT-8015 and the percent of pre-peak in the sample.
- [0020] Figure 3. A graph depicting an elution profile of bench-scale purification of CAT-8015 by Q Sepharose HP Chromatography. CAT-8015 was purified using Q Sepharose HP. The majority of active, intact CAT-8015 resided in fractions D5, D7, and D9 (the main peak spanning from D3 to D12 as indicated above the peak).
- [0021] Figure 4. An SDS-PAGE analysis of QHP load and eluate pool samples (bench-scale purification). Lane 1 corresponds to the QHP load pool; Lane 2 corresponds to the QHP eluate pool; and Lane 3 corresponds to a CAT-8015 reference standard.
- [0022] Figure 5. Large-scale purification of CAT-8015 by Q Sepharose HP Chromatography. CAT-8015 was purified using Q Sepharose HP. As shown in the figure and Table 3, the majority of active, intact CAT-8015 resided in fractions 5, 6, and 7.
- [0023] Figure 6. An SDS-PAGE analysis of QHP load and eluate pool samples (large-scale purification). Lane 1 corresponds to the QHP load pool; and Lane 2 corresponds to the QHP eluate pool.
- [0024] Figure 7. A graph depicting percent Pre-Peak in HA Product as a Function of Solubilization pH as described in Example 6.

DETAILED DESCRIPTION OF THE INVENTION

- [0025] The invention provides methods for purifying an active polypeptide or immunoconjugate from a solution containing the polypeptide or immunoconjugate and an acidic variant thereof. In one embodiment, the acidic variant comprises a deamidated form of the polypeptide or immunoconjugate. In contrast to the expected elution behavior from an anion exchange column, the bulk of deamidated variants elute prior to intact

polypeptides under salt gradient elution conditions. Details of the methods are provided herein.

[0026] The terms "polypeptide," "peptide," "protein," and "protein fragment" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymers.

[0027] The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function similarly to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, gamma-carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, *e.g.*, an alpha carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs can have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions similarly to a naturally occurring amino acid. Negatively charged amino acids include aspartic acid (or aspartate) and glutamic acid (or glutamate). Positively charged amino acids include arginine, histidine, and lysine.

[0028] The "composition" to be purified herein comprises the polypeptide of interest and one or more impurities. The composition may be "partially purified" (*i.e.*, having been subjected to one or more purification steps, such as by non-affinity chromatography described herein) or may be obtained directly from a host cell or organism producing the polypeptide (*e.g.*, the composition may comprise harvested cell culture fluid).

[0029] The terms "polypeptide" or "polypeptide of interest" or "protein of interest" and "target protein" or "protein" are used interchangeably and refer to a protein or polypeptide such as an antibody or immunoconjugate (as defined herein) that is to be purified by a method of the invention from a mixture of proteins and, other materials such as an acidic variant of the polypeptide of interest.

- [0030] An "acidic variant" is a variant of a polypeptide or immunoconjugate which is more acidic (*e.g.*, as determined by cation exchange chromatography) than the polypeptide of interest. An example of an acidic variant is a deamidated variant.
- [0031] Deamidated proteins are that have had some or all of the free amide functional groups hydrolyzed to carboxylic acids, such as conversion of glutamines to glutamic acid. The rate of this reaction is dependent on the primary sequence, three-dimensional structure, pH, temperature, buffer type, ionic strength and other solution properties. Importantly, the deamidation reaction introduces a negative charge into the molecule. As described further below, the protein deamidation can have a negative impact on protein activity.
- [0032] As used herein, the terms "antibody" and "immunoglobulin" are used interchangeably in the broadest sense and include monoclonal antibodies (*e.g.*, full length or intact monoclonal antibodies), polyclonal antibodies, multivalent antibodies, multispecific antibodies (*e.g.*, bispecific antibodies so long as they exhibit the desired biological activity) and antibody fragments as described herein. The term "bispecific antibody" is intended to include any antibody that has two different binding specificities, *i.e.*, the antibody binds two different epitopes, which can be located on the same target antigen or, more commonly, on different target antigens.
- [0033] Native antibodies and immunoglobulins are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains (Clothia *et al.*, *J. Mol. Biol.* 186, 651-66, (1985); Novotny and Haber, *Proc. Natl. Acad. Sci. USA* 82, 4592-4596 (1985)). Five human immunoglobulin classes are defined on the basis of their heavy chain composition, and are named IgG, IgM, IgA,

- 8 -

IgE, and IgD. The IgG-class and IgA-class antibodies are further divided into subclasses, namely, IgG1, IgG2, IgG3, and IgG4, and IgA1 and IgA2. The heavy chains in IgG, IgA, and IgD antibodies have three constant region domains, that are designated CH1, CH2, and CH3, and the heavy chains in IgM and IgE antibodies have four constant region domains, CH1, CH2, CH3, and CH4. Thus, heavy chains have one variable region and three or four constant regions. Immunoglobulin structure and function are reviewed, for example, in Harlow *et al.*, Eds., Antibodies: A Laboratory Manual, Chapter 14, Cold Spring Harbor Laboratory, Cold Spring Harbor (1988).

[0034] The term "antibody fragment" refers to a portion of an intact antibody and refers to the antigenic determining variable regions of an intact antibody. Examples of antibody fragments include, but are not limited to Fab, Fab', F(ab')₂, Fv and single chain Fv fragments, linear antibodies, single chain antibodies, and multispecific antibodies formed from antibody fragments.

[0035] The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific and bind a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. That an antibody "selectively binds" or "specifically binds" means that the antibody reacts or associates more frequently, more rapidly, with greater duration, with greater affinity, or with some combination of the above to an epitope than with alternative substances, including unrelated proteins. "Selectively binds" or "specifically binds" means, for instance, that an antibody binds to a protein with a K_D of at least about 0.1 mM, but more usually at least about 1 μ M. "Selectively binds" or "specifically binds" means at times that an antibody binds to a protein at times with a K_D of at least about 0.1 μ M or better, and at other times at least about 0.01 μ M or better. Because of the sequence identity between homologous proteins in different species, specific binding can include an antibody that recognizes a tumor cell marker protein in more than one species.

[0036] The antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding

- 9 -

sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al*, *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984)).

[0037] "Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies can comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally will also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992). See also the following review articles and references cited therein: Vaswani and Hamilton, *Ann. Allergy, Asthma & Immunol.* 1:105-115 (1998); Harris, *Biochem. Soc. Transactions* 23: 1035-1038 (1995); Hurle and Gross, *Curr. Op. Biotech.* 5:428-433 (1994).

[0038] A "human antibody" is one that possesses an amino acid sequence that corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues.

[0039] The term "immunoconjugate" or "conjugate" or "immunotoxin" as used herein refers to a compound or a derivative thereof that is linked to a cell binding agent (e.g., an anti-CD22 antibody or fragment thereof) and is defined by a generic formula: C-L-A, wherein C = cytotoxin, L = linker, and A = cell binding agent (e.g., anti-CD22 antibody or antibody fragment). Immunoconjugates can also be defined by the generic formula in reverse order: A-L-C.

[0040] The term "cytotoxin" or "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g., At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu), chemotherapeutic agents e.g., methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents, enzymes and fragments thereof such as nucleolytic enzymes, antibiotics, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof, and the various antitumor or anticancer agents disclosed below. Examples of cytotoxic agents include, but are not limited to, abrin, ricin, *Pseudomonas* exotoxin (PE), diphtheria toxin (DT), botulinum toxin, or modified toxins thereof. For example, PE and DT are highly toxic compounds that typically bring about death through liver toxicity. PE and DT, however, can be modified into a form for use as an immunotoxin by removing the native targeting component of the toxin (e.g., domain Ia of PE or the B chain of DT) and replacing it with a different targeting moiety, such as an antibody.

[0041] In some embodiments, the toxin is *Pseudomonas* exotoxin. *Pseudomonas* exotoxin A (PE) is an extremely active monomeric protein (molecular weight 66 kD), secreted by *Pseudomonas aeruginosa*, which inhibits protein synthesis in eukaryotic cells through the inactivation of elongation factor 2 (EF-2) by catalyzing its ADP-ribosylation (catalyzing the transfer of the ADP ribosyl moiety of oxidized NAD onto EF-2).

[0042] The toxin contains three structural domains that act in concert to cause cytotoxicity. Domain Ia (amino acids 1-252) mediates cell binding. Domain II (amino acids 253-364) is responsible for translocation into the cytosol and domain III (amino acids 400-613) mediates ADP ribosylation of elongation factor 2, which inactivates the protein and causes cell death. The function of domain Ib (amino acids 365-399) remains

- 11 -

undefined, although a large part of it, amino acids 365-380, can be deleted without loss of cytotoxicity. See Siegall *et al.*, *J. Biol. Chem.* 264: 14256-14261 (1989).

[0043] The *Pseudomonas* exotoxins (PE) employed in the present invention include the native sequence, cytotoxic fragments of the native sequence, and conservatively modified variants of native PE and its cytotoxic fragments. Cytotoxic fragments of PE include those which are cytotoxic with or without subsequent proteolytic or other processing in the target cell (*e.g.*, as a protein or pre-protein). Cytotoxic fragments of PE include PE40, PE38, and PE35. PE40 is a truncated derivative of PE as previously described in the art. See, Pai *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:3358-62 (1991); Kondo *et al.*, *J. Biol. Chem.* 263:9470-9475 (1988). PE38 is a truncated PE composed of amino acids 253-364 and 381-613 of native PE. PE35 is a 35 kD carboxyl-terminal fragment of PE composed of a Met at position 280 followed by amino acids 281-364 and 381-613 of native PE. In one embodiment, the cytotoxic fragment PE38 is employed. PE38 is a pro-protein which can be activated to its cytotoxic form upon processing within a cell.

[0044] A "PE immunoconjugate" or "PE immunotoxin" is an immunoconjugate or immunotoxin comprising an antibody or antigen binding fragment thereof and a PE toxin or variant thereof.

[0045] By "purifying" a polypeptide or immunoconjugate from a composition comprising the polypeptide and one or more impurities is meant increasing the degree of purity of the polypeptide in the composition by removing (completely or partially) at least one impurity from the composition. According to the present invention, purification is performed without the use of an affinity chromatography step.

[0046] The term "chromatography" refers to the process by which a solute of interest in a mixture is separated from other solutes in a mixture as a result of differences in rates at which the individual solutes of the mixture migrate through a stationary medium under the influence of a moving phase, or in bind and elute processes.

[0047] The term "ion-exchange" and "ion-exchange chromatography" refers to the chromatographic process in which a solute of interest (such as a protein) in a mixture interacts with a charged compound linked (such as by covalent attachment) to a solid phase ion exchange material such that the solute of interest interacts non-specifically with the charged compound more or less than solute impurities or contaminants in the mixture. The contaminating solutes in the mixture elute from a column of the ion exchange

material faster or slower than the solute of interest or are bound to or excluded from the resin relative to the solute of interest. "Ion-exchange chromatography" specifically includes cation exchange, anion exchange, and mixed mode chromatography.

[0048] The phrase "ion exchange material" refers to a solid phase that is negatively charged (*i.e.*, a cation exchange resin) or positively charged (*i.e.*, an anion exchange resin). The charge may be provided by attaching one or more charged ligands to the solid phase, *e.g.*, by covalent linking. Alternatively, or in addition, the charge may be an inherent property of the solid phase (*e.g.*, as is the case for silica, which has an overall negative charge).

[0049] An "anion exchange resin" refers to a solid phase which is positively charged, thus having one or more positively charged ligands attached thereto. Any positively charged ligand attached to the solid phase suitable to form the anionic exchange resin can be used, such as quaternary amino groups. Commercially available anion exchange resins include DEAE cellulose, Poros PI 20, PI 50, HQ 10, HQ 20, HQ 50, D 50 from Applied Biosystems, Sartobind Q from Sartorius, MonoQ, MiniQ, Source 15Q and 30Q, Q, DEAE and ANX Sepharose Fast Flow, Q Sepharose High Performance, QAE SEPHADEX™ and FAST Q SEPHAROSE™ (GE Healthcare), WP PEI, WP DEAM, WP QUAT from J. T. Baker, Hydrocell DEAE and Hydrocell QA from Biochrom Labs Inc., UNOsphere Q, Macro-Prep DEAE and Macro-Prep High Q from Biorad, Ceramic HyperD Q, ceramic HyperD DEAE, Trisacryl M and LS DEAE, Spherox LS DEAE, QMA Spherosil LS, QMA Spherosil M and Mustang Q from Pall Technologies, DOWEX Fine Mesh Strong Base Type I and Type II Anion Resins and DOWEX MONOSPHER E 77, weak base anion from Dow Liquid Separations, Intercept Q membrane, Matrex Cellufine A200, A500, Q500, and Q800, from Millipore, Fractogel EMD TMAE, Fractogel EMD DEAE and Fractogel EMD DMAE from EMD, Amberlite weak strong anion exchangers type I and II, DOWEX weak and strong anion exchangers type I and II, Diaion weak and strong anion exchangers type I and II, Duolite from Sigma-Aldrich, TSK gel Q and DEAE 5PW and 5PW-HR, Toyopearl SuperQ-650S, 650M and 650C, QAE-550C and 650S, DEAE-650M and 650C from Tosoh, QA52, DE23, DE32, DE51, DE52, DE53, Express-Ion D and Express-Ion Q from Whatman.

[0050] By "solid phase" is meant a non-aqueous matrix to which one or more charged ligands can adhere. The solid phase may be a purification column, a discontinuous phase

of discrete particles, a membrane, or filter etc. Examples of materials for forming the solid phase include polysaccharides (such as agarose and cellulose); and other mechanically stable matrices such as silica (*e.g.*, controlled pore glass), poly(styrenedivinyl)benzene, polyacrylamide, ceramic particles and derivatives of any of the above.

[0051] The term "specific binding" as used herein, such as to describe interactions between a molecule of interest and a ligand bound to a solid phase matrix, refers to the generally reversible binding of a protein of interest to a ligand through the combined effects of spatial complementarity of protein and ligand structures at a binding site coupled with electrostatic forces, hydrogen bonding, hydrophobic forces, and/or van der Waals forces at the binding site. The greater the spatial complementarity and the stronger the other forces at the binding site, the greater will be the binding specificity of a protein for its respective ligand. Non-limiting examples of specific binding includes antibody-antigen binding, enzyme-substrate binding, enzyme-cofactor binding, metal ion chelation, DNA binding protein-DNA binding, regulatory protein-protein interactions, and the like.

[0052] The term "non-specific binding" as used herein, such as to describe interactions between a molecule of interest and a ligand or other compound bound to a solid phase matrix, refers to binding of a protein of interest to the ligand or compound on a solid phase matrix through electrostatic forces, hydrogen bonding, hydrophobic forces, and/or van der Waals forces at an interaction site, but lacking structural complementarity that enhances the effects of the non-structural forces. Examples of non-specific interactions include, but are not limited to, electrostatic, hydrophobic, and van der Waals forces as well as hydrogen bonding.

[0053] A "salt" is a compound formed by the interaction of an acid and a base. A salt useful for the invention include, but are not limited to acetate (*e.g.*, sodium acetate), citrate (*e.g.*, sodium citrate), chloride (*e.g.*, sodium chloride), sulphate (*e.g.*, sodium sulphate), or a potassium salt.

[0054] The term "detergent" refers to ionic and nonionic surfactants such as polysorbates (*e.g.*, polysorbates 20 or 80); poloxamers (*e.g.*, poloxamer 188); TritonTM; sodium dodecyl sulfate (SDS); sodium lauryl sulfate; sodium octyl glucoside; lauryl-, myristyl-, linoleyl-, or stearyl-sulfobetaine; lauryl-, myristyl-, linoleyl- or stearyl-sarcosine; linoleyl-, myristyl-, or cetyl-betaine; lauroamidopropyl-, cocamidopropyl-, linoleamidopropyl-,

myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-betaine (*e.g.*, lauroamidopropyl); myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-dimethylamine; sodium methyl cocoyl-, or disodium methyl oleoyl-taurate; and the MONAQUAT™ series (Mona Industries, Inc., Paterson, N.J.), Useful detergents is a polysorbate, such as polysorbate 20 (TWEEN 20®) or polysorbate 80 (TWEEN 80®).

[0055] A "buffer" used in the present invention is a solution that resists changes in pH by the addition of acid or base by the action of its acid-base conjugates components. Various buffers can be employed in a method of the present invention depending on the desired pH of the buffer and the particular step in the purification process [*see* Buffers. A Guide for the Preparation and Use of Buffers in Biological Systems, Gueffroy, D., ed. Calbiochem Corporation (1975)]. Non-limiting examples of buffer components that can be used to control the pH range desirable for a method of the invention include acetate, citrate, histidine, phosphate, ammonium buffers such as ammonium acetate, succinate, MES, CHAPS, MOPS, MOPSO, HEPES, Tris, and the like, as well as combinations of these TRIS-malic acid-NaOH, maleate, chloroacetate, formate, benzoate, propionate, pyridine, piperazine, ADA, PIPES, ACES, BES, TES, tricine, bicine, TAPS, ethanolamine, CHES, CAPS, methylamine, piperidine, O-boric acid, carbonic acid, lactic acid, butanedioic acid, diethylmalonic acid, glycylglycine, HEPPS, HEPPSO, imidazole, phenol, POPSO, succinate, TAPS, amine-based, benzylamine, trimethyl or dimethyl or ethyl or phenyl amine, ethylenediamine, or morpholine. Additional components (additives) can be present in a buffer as needed, *e.g.*, salts can be used to adjust buffer ionic strength, such as sodium chloride, sodium sulfate and potassium chloride; and other additives such as amino acids (such as glycine and histidine), chaotropes (such as urea), alcohols (such as ethanol, mannitol, glycerol, and benzyl alcohol), detergents (*see supra.*), and sugars (such as sucrose, mannitol, maltose, trehalose, glucose, and fructose). The buffer components and additives, and the concentrations used, can vary according to the type of chromatography practiced in the invention.

[0056] The "loading buffer" is that which is used to load the composition comprising the polypeptide molecule of interest and one or more impurities onto the ion exchange resin. The loading buffer has a conductivity and/or pH such that the polypeptide molecule of interest (and generally one or more impurities) is/are bound to the ion exchange resin or

- 15 -

such that the protein of interest flows through the column while the impurities bind to the resin.

[0057] The term "wash buffer" when used herein refers to a buffer used to wash or re-equilibrate the ion exchange resin, prior to eluting the polypeptide molecule of interest. Conveniently, the wash buffer and loading buffer may be the same, but this is not required.

[0058] The "elution buffer" is used to elute the polypeptide of interest from the solid phase. The conductivity and/or pH of the elution buffer is/are such that the polypeptide of interest is eluted from the ion exchange resin.

[0059] The "pI" or "isoelectric point" of a polypeptide refer to the pH at which the polypeptide's positive charge balances its negative charge. pI can be calculated from the net charge of the amino acid residues or sialic acid residues of attached carbohydrates of the polypeptide or can be determined by isoelectric focusing.

[0060] By "binding" a molecule to an ion exchange material is meant exposing the molecule to the ion exchange material under appropriate conditions (pH/conductivity) such that the molecule is reversibly immobilized in or on the ion exchange material by virtue of ionic interactions between the molecule and a charged group or charged groups of the ion exchange material.

[0061] By "washing" the ion exchange material is meant passing an appropriate buffer through or over the ion exchange material.

[0062] To "elute" a molecule (*e.g.*, polypeptide or impurity) from an ion exchange material is meant to remove the molecule therefrom by altering the ionic strength of the buffer surrounding the ion exchange material such that the buffer competes with the molecule for the charged sites on the ion exchange material.

[0063] As used in the present disclosure and claims, the singular forms "a," "an," and "the" include plural forms unless the context clearly dictates otherwise.

[0064] It is understood that wherever embodiments are described herein with the language "comprising," otherwise analogous embodiments described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0065] The term "and/or" as used in a phrase such as "A and/or B" herein is intended to include both "A and B," "A or B," "A," and "B." Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following

embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

Pseudomonas Exotoxin and Other Toxins

[0066] Toxins can be employed with antibodies of the present invention to yield immunotoxins. Exemplary toxins include ricin, abrin, diphtheria toxin and subunits thereof, as well as botulinum toxins A through F. These toxins are readily available from commercial sources (e.g., Sigma Chemical Company, St. Louis, Mo.). Diphtheria toxin is isolated from *Corynebacterium diphtheriae*. Ricin is the lectin RCA60 from *Ricinus communis* (Castor bean). The term also references toxic variants thereof. For example, see, U.S. Pat. Nos. 5,079,163 and 4,689,401. *Ricinus communis* agglutinin (RCA) occurs in two forms designated RCA₆₀ and RCA₁₂₀ according to their molecular weights of approximately 65 and 120 kD, respectively (Nicholson & Blaustein, *J. Biochem. Biophys. Acta* 266:543 (1972)). The A chain is responsible for inactivating protein synthesis and killing cells. The B chain binds ricin to cell-surface galactose residues and facilitates transport of the A chain into the cytosol (Olsnes, *et al.*, *Nature* 249:627-631 (1974) and U.S. Pat. No. 3,060,165).

[0067] Abrin includes toxic lectins from *Abrus precatorius*. The toxic principles, abrin a, b, c, and c, have a molecular weight of from about 63 to 67 kD and are composed of two disulfide-linked polypeptide chains A and B. The A chain inhibits protein synthesis; the B-chain (abrin-b) binds to D-galactose residues (see, Funatsu, *et al.*, *Agr. Biol. Chem.* 52:1095 (1988); and Olsnes, *Methods Enzymol.* 50:330-335 (1978)).

[0068] In preferred embodiments of the present invention, the toxin is *Pseudomonas* exotoxin (PE). The *Pseudomonas* exotoxin (or exotoxin A) is an exotoxin produced by *Pseudomonas aeruginosa*. The term "*Pseudomonas* exotoxin" as used herein refers to a full-length native (naturally occurring) PE or a PE that has been modified. Such modifications may include, but are not limited to, elimination of domain Ia, various amino acid deletions in domains Ib, II and III, single amino acid substitutions and the addition of one or more sequences at the carboxyl terminus such as KDEL (SEQ ID NO:3) and REDL (SEQ ID NO:4). See Siegall, *et al.*, *J. Biol. Chem.* 264:14256-14261 (1989). In a preferred embodiment, the cytotoxic fragment of PE retains at least 50%,

- 17 -

preferably 75%, more preferably at least 90%, and most preferably 95% of the cytotoxicity of native PE. In a most preferred embodiment, the cytotoxic fragment is more toxic than native PE.

[0069] Native *Pseudomonas* exotoxin A (PE) is an extremely active monomeric protein (molecular weight 66 kD), secreted by *Pseudomonas aeruginosa*, which inhibits protein synthesis in eukaryotic cells. The native PE sequence is provided in commonly assigned U.S. Pat. No. 5,602,095, incorporated herein by reference. The method of action is inactivation of the ADP-ribosylation of elongation factor 2 (EF-2). The exotoxin contains three structural domains that act in concert to cause cytotoxicity. Domain Ia (amino acids 1-252) mediates cell binding. Domain II (amino acids 253-364) is responsible for translocation into the cytosol and domain III (amino acids 400-613) mediates ADP ribosylation of elongation factor 2. The function of domain Ib (amino acids 365-399) remains undefined, although a large part of it, amino acids 365-380, can be deleted without loss of cytotoxicity. See Siegall, *et al.*, (1989), *supra*.

[0070] PE employed in the present invention include the native sequence, cytotoxic fragments of the native sequence, and conservatively modified variants of native PE and its cytotoxic fragments. PE variants useful in the invention are described in US 7,355,012, and WO 2007/016150 and WO 2009/032954. Cytotoxic fragments of PE include those which are cytotoxic with or without subsequent proteolytic or other processing in the target cell (*e.g.*, as a protein or pre-protein). Cytotoxic fragments of PE include PE40, PE38, and PE35.

[0071] In preferred embodiments, the PE has been modified to reduce or eliminate non-specific cell binding, frequently by deleting domain Ia as taught in U.S. Pat. No. 4,892,827, although this can also be achieved, for example, by mutating certain residues of domain Ia. U.S. Pat. No. 5,512,658, for instance, discloses that a mutated PE in which Domain Ia is present but in which the basic residues of domain Ia at positions 57, 246, 247, and 249 are replaced with acidic residues (glutamic acid, or "E")) exhibits greatly diminished non-specific cytotoxicity. This mutant form of PE is sometimes referred to as PE4E.

[0072] PE40 is a truncated derivative of PE as previously described in the art, with a deletion of domain Ia of the native PE molecule. See, Pai, *et al.*, *Proc. Nat'l Acad. Sci. USA* 88:3358-62 (1991); and Kondo, *et al.*, *J. Biol. Chem.* 263:9470-9475 (1988). PE35 is

a 35 kD carboxyl-terminal fragment of PE in which amino acid residues 1-279 have been deleted and the molecule commences with a Met at position 280 followed by amino acids 281-364 and 381-613 of native PE. PE35 and PE40 are disclosed, for example, in U.S. Pat. Nos. 5,602,095 and 4,892,827. PE4E is a form of PE where all of the domains of native PE are present, but where the basic residues of domain Ia at positions 57, 246, 247 and 249 are replaced with acidic residues (glutamine acid, or "E").

[0073] In some preferred embodiments, the cytotoxic fragment PE38 is employed. PE38 is a truncated PE pro-protein composed of amino acids 253-364 and 381-613 which is activated to its cytotoxic form upon processing within a cell (see *e.g.*, U.S. Pat. Nos. 5,608,039, 7,355,012, and Pastan et al., *Biochim. Biophys. Acta* 1333:C1-C6 (1997)).

[0074] As noted above, some or all of domain Ib may be deleted, and the remaining portions joined by a linker or directly by a peptide bond. Some of the amino portion of domain II may be deleted. And, the C-terminal end may contain the native sequence of residues 609-613 (REDLK) (SEQ ID NO:5), or may contain a variation found to maintain the ability of the construct to translocate into the cytosol, such as REDL (SEQ ID NO:4) or KDEL (SEQ ID NO:3), and repeats of these sequences. See, *e.g.*, U.S. Pat. Nos. 5,854,044; 5,821,238; and 5,602,095 and WO 99/51643. While in preferred embodiments, the PE is PE4E, PE40, or PE38, any form of PE in which non-specific cytotoxicity has been eliminated or reduced to levels in which significant toxicity to non-targeted cells does not occur can be used in the immunotoxins of the present invention so long as it remains capable of translocation and EF-2 ribosylation in a targeted cell.

Conservatively Modified Variants of PE

[0075] Conservatively modified variants of PE or cytotoxic fragments thereof have at least 80% sequence similarity, preferably at least 85% sequence similarity, more preferably at least 90% sequence similarity, and most preferably at least 95% sequence similarity at the amino acid level, with the PE of interest, such as PE38.

[0076] The term "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refer to those nucleic acid sequences which encode identical or

- 19 -

essentially identical amino acid sequences, or if the nucleic acid does not encode an amino acid sequence, to essentially identical nucleic acid sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given polypeptide. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

[0077] As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid.

[0078] *Pseudomonas* exotoxins employed in the invention can be assayed for the desired level of cytotoxicity by assays well known to those of skill in the art. Thus, cytotoxic fragments of PE and conservatively modified variants of such fragments can be readily assayed for cytotoxicity. A large number of candidate PE molecules can be assayed simultaneously for cytotoxicity by methods well known in the art. For example, subgroups of the candidate molecules can be assayed for cytotoxicity. Positively reacting subgroups of the candidate molecules can be continually subdivided and reassayed until the desired cytotoxic fragment(s) is identified. Such methods allow rapid screening of large numbers of cytotoxic fragments or conservative variants of PE.

Anti-CD22/PE immunoconjugates

[0079] In one embodiment, the polypeptide of interest comprises an antibody that specifically binds CD22. "CD22" refers to a lineage-restricted B cell antigen belonging to the Ig superfamily. It is expressed in 60-70% of B cell lymphomas and leukemias and

is not present on the cell surface in early stages of B cell development or on stem cells. See, e.g., Vaickus *et al.*, *Crit. Rev. Oncol/Hematol.* 11:267-297 (1991). In another embodiment, the polypeptide of interest is an antibody fragment that binds CD22 (e.g., Fab, or scFv).

[0080] As used herein, the term "anti-CD22" in reference to an antibody, refers to an antibody that specifically binds CD22 and includes reference to an antibody which is generated against CD22. In some embodiments, the CD22 is a primate CD22 such as human CD22. In one embodiment, the antibody is generated against human CD22 synthesized by a non-primate mammal after introduction into the animal of cDNA which encodes human CD22. In a further embodiment, the polypeptide of interest is a CD22 antibody immunoconjugate that comprises the PE38 exotoxin.

[0081] One example of a CD22/PE38 immunoconjugate is CAT-8015 described in International Patent Application Publication Nos. WO 98/41641 and WO2003/27135, US Patent Nos. 7,541,034, 7,355,012, and U.S. Publication No. 2007/0189962.

CAT-8015 is a recombinant immunotoxin protein composed of an antibody Fv fragment based on the murine anti-CD22 antibody RFB4 fused to a truncated form of the *Pseudomonas* exotoxin protein, PE38. The anti-CD22 Fv fragment consists of two domains, a V_L and a V_H, where the latter was modified to improve binding to the human CD22 target. The CAT-8015 protein is comprised of two independent polypeptides, the V_L chain (SEQ ID NO:2), and the V_H chain, fused at the C-terminus to the PE38 domain (V_H-PE38) (SEQ ID NO:1). Other V_L and V_H-PE38 sequences useful in this invention are described in US 7,541,034, 7,355,012, and 2007/0189962. Both domains were designed to each contain engineered cysteine residues that permit formation of an intermolecular disulfide bond. This feature increases the stability of the fusion protein.

[0082] The amino acid sequence of the V_H-P38 Subunit (SEQ ID NO:1) of CAT-8015 is the following:

```
MEVQLVESGGGLVKPGGSLKLSAASGFAFSIYDMSWVRQTPEKCLEWV
AYISSGGGTYYPDTVKGRFTISRDNKNTLYLQMSSLKSEDTAMYYCA
RHSGYGTHWGVLFAYWGQGTLVTVSAKASGGPEGGSLAALTAHQACHLP
LETFTTRHRQPRGWEQLEQCGYPVQRLVALYLAARLSWNQVDQVIRNALA
SPGSGGDLGEAIREQPEQARLALTAAESERFVRQGTGNDEAGAANGP
ADSGDALLERNYPTGAEFLGDGGDVSFSTRGTQNWTVERLLQAHQLEE
RGYVFVGYHGTFLEAAQSIVFGGVRARSQDLDAIWRGFYIAGDPALAYG
YAQDQEPDARGRI RNGALLRVYVPRSSLPGFYRTSLTLAAPEAAGEVER
```


- 21 -

LIGHPLPLRLDAITGPEEEGGRLETILGWPLAERTVVIPSAIPTDPRNV
GGDLDPSSIPDKEQAISALPDYASQPGKPPREDLK (SEQ ID NO:1)

[0083] The PE38 sequence is shown in bold, and the five amino acid linker between the V_H domain and the PE38 domain is shown underlined.

[0084] The amino acid sequence of the V_L Subunit (SEQ ID NO:2) of CAT-8015 is the following:

MDIQMTQTTSSLSASLGDRVTISCRASQDISNYLNWYQQKPDGTVKLLI
 YYTSILHSGVPSRFSGSGSGTDYSLTISNLEQEDFATYFCQQGNTLPWT
 FCGGTKLEIK (SEQ ID NO:2)

[0085] In further embodiments, the amino acid sequence of the V_H domain of the immunoconjugate is one of the following:

MEVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQTPEKCLEWV
 AYISSGGGTTYYPDTVKGRTISRDNANTLYLQMSSLKSEDTAMYYCA
 RHSGYGTHWGVLFAYWGQGLTVTVSA (SEQ ID NO:6)

MEVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQTPEKCLEWV
 AYISSGGGTTYYPDTVKGRTISRDNANTLYLQMSSLKSEDTAMYYCA
 RHSGYGYNWGVLFAYWGQGLTVTVSA (SEQ ID NO:7)

MEVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQTPEKCLEWV
 AYISSGGGTTYYPDTVKGRTISRDNANTLYLQMSSLKSEDTAMYYCA
 RHSGYGTTWGVLFAYWGQGLTVTVSA (SEQ ID NO:8)

MEVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQTPEKCLEWV
 AYISSGGGTTYYPDTVKGRTISRDNANTLYLQMSSLKSEDTAMYYCA
 RHSGYGSTYGVLFAYWGQGLTVTVSA (SEQ ID NO:9)

MEVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQTPEKCLEWV
 AYISSGGGTTYYPDTVKGRTISRDNANTLYLQMSSLKSEDTAMYYCA
 RHSGYGTHWGVLFAYWGQGLTVTVSA (SEQ ID NO:10)

- 22 -

MEVQLVESGGGLVKPGGSLKLSAASGFAFSIYDMSWVRQTPEKCLEWV
 AYISSGGGTTYYPDTVKGRTISRDNANTLYLQMSSLKSEDTAMYYCA
 RHSGYGSSYGVLFAIWGQGTLLTVSA (SEQ ID NO:11)

[0086] In additional embodiments, the amino acid sequence of the VL domain of the immunoconjugate is one of the following:

MDIQMTQTTSSLSASLGDRVTISCRASQDIARYLNWYQQKPDGTVKLLI
 YYTSILHSGVPSRFSGSGSGTDYSLTISNLEQEDFATYFCQQGNTLPWT
 FGCGTKLEIK (SEQ ID NO:12)

MDIQMTQTTSSLSASLGDRVTISCRASQDIHGYNWYQQKPDGTVKLLI
 YYTSILHSGVPSRFSGSGSGTDYSLTISNLEQEDFATYFCQQGNTLPWT
 FGCGTKLEIK (SEQ ID NO:13)

MDIQMTQTTSSLSASLGDRVTISCRASQDIGRYLNWYQQKPDGTVKLLI
 YYTSILHSGVPSRFSGSGSGTDYSLTISNLEQEDFATYFCQQGNTLPWT
 FGCGTKLEIK (SEQ ID NO:14)

MDIQMTQTTSSLSASLGDRVTISCRASQDIRGYLNWYQQKPDGTVKLLI
 YYTSILHSGVPSRFSGSGSGTDYSLTISNLEQEDFATYFCQQGNTLPWT
 FGCGTKLEIK (SEQ ID NO:15)

[0087] In certain other embodiments, the PE toxin of the immunoconjugate is a PE or variant thereof selected from the following:

[0088] Native PE

AEEAFDLWNECAKACVLDLKDGVRSRMSVDPADTNGQGVLYSMVL
 EGGNDALKLAIDNALSITSDGLTIRLEGGVEPNKPVRYSYTRQARGSW
 LNWLVPIGHEKPSNIKVFIHELNAGNQLSHMSPIYTIEMGDELLAKLAR
 DATFFVRAHESNEMQPTLAISHAGVSVVMAQTQPRREKRWSEWASGKVL
 CLLDPLDGVYNYLAQQRCNLDDTWEGKIYRVLAGNPAKHDLDIKPTVIS
 HRLHFPEGGSLAALTAHQACHLPLETFTRHRQPRGWEQLEQCGYPVQRL

- 23 -

VALYLAARLSWNQVDQVIRNALASPGSGGDLGEAIREQPEQARLALT
 AAESERFVRQGTGNDEAGAANGPADSGDALLERNYPTGAEFLGDGGDVS
 FSTRGTQNWTVRLLQHRQLEERGYVFGYHGTFLAAQSIVFGGVRA
 RSQDLDAIWRGFYIAGDPALAYGYAQDQEPDARGRIRNGALLRVYVPRS
 SLPGFYRTSLTLAAPEAAGEVERLIGHPLPLRLDAITGPEEEGGRLETI
 LGWPLAERTVVIPSAIPTDPRNVGGDLDPSSIPDKEQAISALPDYASQP
 GKPPREDLK (SEQ ID NO:16)

[0089]**PE40**

GGSLAALTAHQACHLPLETFTRHRQPRGWEQLEQCGYPVQRLVALYLAAR
 LSWNQVDQVIRNALASPGSGGDLGEAIREQPEQARLALTAAAESERFVR
 QGTGNDEAGANADVSLTCPVAAGECAGPADSGDALLERNYPTGAEFLG
 DGGDVSFSTRGTQNWTVRLLQHRQLEERGYVFGYHGTFLAAQSIVF
 GGVRRARSQDLDAIWRGFYIAGDPALAYGYAQDQEPDARGRIRNGALLRVY
 VPRSSLPGFYRTSLTLAAPEAAGEVERLIGHPLPLRLDAITGPEEEGGRL
 ETILGWPLAERTVVIPSAIPTDPRNVGGDLDPSSIPDKEQAISALPDYAS
 QPGKPPREDLK (SEQ ID NO:17)

[0090]**PE38**

GGSLAALTAHQACHLPLETFTRHRQPRGWEQLEQCGYPVQRLVALYLAAR
 LSWNQVDQVIRNALASPGSGGDLGEAIREQPEQARLALTAAAESERFVR
 QGTGNDEAGAANGPADSGDALLERNYPTGAEFLGDGGDVSFSTRGTQNW
 TVRLLQHRQLEERGYVFGYHGTFLAAQSIVFGGVRRARSQDLDAIWRG
 FYIAGDPALAYGYAQDQEPDARGRIRNGALLRVYVPRSSLPGFYRTSLTL
 AAPEAAGEVERLIGHPLPLRLDAITGPEEEGGRLETI
 LGWPLAERTVVIPSAIPTDPRNVGGDLDPSSIPDKEQAISALPDYASQPGKPPREDLK (SEQ
 ID NO:18)

[0091]**PE35**

MWEQLEQCGYPVQRLVALYLAARLSWNQVDQVIRNALASPGSGGDLGEAI
 REQPEQARLALTAAAESERFVRQGTGNDEAGAANGPADSGDALLERNYP
 TGAEFLGDGGDVSFSTRGTQNWTVRLLQHRQLEERGYVFGYHGTFL
 AAQSIVFGGVRRARSQDLDAIWRGFYIAGDPALAYGYAQDQEPDARGRIRN
 GALLRVYVPRSSLPGFYRTSLTLAAPEAAGEVERLIGHPLPLRLDAITGP

- 24 -

EEEGGRLETILGWPLAERTVVIPSAIPTDPRNVGGDLDPSSIPDKEQAIS
ALPDYASQPGKPPREDLK (SEQ ID NO:19)

[0092] PE-LR

RHRQPRGWEQLPTGAEFLGDGGDVSFSTRGTQNWTVRLLQAHRLQLEERG
YVFVGYHGTFLCAAQSIVFGGVRARSQDLDAIWRFYIAGDPALAYGYAQ
DQEPDARGRIRNGALLRVYVPRSSLPGFYRTSLTLAAPEAAGEVERLIGH
PLPLRLDAITGPEEEGGRLETILGWPLAERTVVIPSAIPTDPRNVGGDLDP
SSIPDKEQAISALPDYASQPGKPPREDLK (SEQ ID NO:20)

[0093] PE-LR-6X

RHRQPRGWEQLPTGAEFLGDGGDVSFSTRGTQNWTVRLLQAHRLQLEEGG
YVFVGYHGTFLCAAQSIVFGGVRARSQDLDAIWAGFYIAGDPALAYGYAQ
DQEPDAAGRIRNGALLRVYVPRSSLPGFYATSLTLAAPEAAGEVERLIGH
PLPLRLDAITGPEEAGGRLETILGWPLAERTVVIPSAIPTDPRNVGGDLDP
SSIPDSEQAISALPDYASQPGKPPREDLK (SEQ ID NO:21)

[0094] PE-38 (CAT-8015)

PEGGSLAALTAHQACHLPLETFTRHRQPRGWEQLEQCGYPVQRLVALYLA
ARLSWNQVDQVIRNALASPGSGDLGEAIREQPEQARLALTAAAESERF
VRQGTGNDEAGAANGPADSGDALLERNYPTGAEFLGDGGDVSFSTRGTQ
NWTVRLLQAHRLQLEERGYVVFVGYHGTFLCAAQSIVFGGVRARSQDLDAIW
RGFYIAGDPALAYGYAQDQEPDARGRIRNGALLRVYVPRSSLPGFYRTSL
TLAAPEAAGEVERLIGHPLPLRLDAITGPEEEGGRLETILGWPLAERTVV
IPSAIPTDPRNVGGDLDPSSIPDKEQAISALPDYASQPGKPPREDLK
(SEQ ID NO:22)

[0095] The PE toxin of the immunoconjugate is fused or conjugated to either the VH or VL domain directly or via a linker at either the N-terminus or the C-terminus of the VH or VL domain. An example of a linker is described above for CAT-8015 and corresponds to the amino acid sequence KASGG (SEQ ID NO: 23). Additional linkers can be readily generated by techniques known in the art.

- 25 -

Expression of a PE immunoconjugate

[0096] The PE immunoconjugate of the present invention is expressed in cells, such as bacterial cells, and then isolated from inclusion bodies. The PE immunoconjugate isolated from inclusion bodies is then further purified using downstream purification steps.

[0097] A variety of host-expression vector systems may be utilized to express the PE immunoconjugate of the present invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention *in situ*. These include but are not limited to microorganisms such as bacteria (*e.g.*, *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (*e.g.*, *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors containing antibody coding sequences; or mammalian cell systems (*e.g.*, COS, CHO, BLK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter).

[0098] DNA encoding each of the VL and VH-PE toxin (*e.g.*, VH-PE38) polypeptides is introduced into an expression vector by techniques well known in the art.

[0099] A "vector" refers to any vehicle for the cloning of and/or transfer of a nucleic acid into a host cell. A vector may be a replicon to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" refers to any genetic element (*e.g.*, plasmid, phage, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, *i.e.*, capable of replication under its

- 26 -

own control. The term "vector" includes vehicles for introducing the nucleic acid into a cell *in vitro*, *ex vivo* or *in vivo*. A large number of vectors known in the art may be used to manipulate nucleic acids, incorporate response elements and promoters, such as inducible promoters, into genes, etc. Possible vectors include, for example, plasmids such as pBR322 or pUC plasmid derivatives, or the Bluescript vector. For example, the insertion of the DNA fragments corresponding to response elements and promoters into a suitable vector can be accomplished by ligating the appropriate DNA fragments into a chosen vector that has complementary cohesive termini. Alternatively, the ends of the DNA molecules may be enzymatically modified or any site may be produced by ligating nucleotide sequences (linkers) into the DNA termini. Such vectors may be engineered to contain selectable marker genes that provide for the selection of cells. Such markers allow identification and/or selection of host cells that express the proteins encoded by the marker.

[00100] The term "expression vector" refers to a vector, plasmid or vehicle designed to enable the expression of an inserted nucleic acid sequence following transformation into the host. The cloned gene, *i.e.*, the inserted nucleic acid sequence, *e.g.*, a gene encoding an anti-CD22 VH, anti-CD22 VL, or anti-CD22 VH or VL fused to a PE toxin, is usually placed under the control of control elements such as a promoter, a minimal promoter, an enhancer, or the like. Initiation control regions or promoters, which are useful to drive expression of a nucleic acid in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving expression of these genes can be used in an expression vector, including but not limited to, viral promoters, bacterial promoters, animal promoters, mammalian promoters, synthetic promoters, constitutive promoters, tissue specific promoters, pathogenesis or disease related promoters, developmental specific promoters, inducible promoters, light regulated promoters; including, but are not limited to, the SV40 early (SV40) promoter region, the promoter contained in the 3' long terminal repeat (LTR) of Rous sarcoma virus (RSV), the E1A or major late promoter (MLP) of adenoviruses (Ad), the human cytomegalovirus (HCMV) immediate early promoter, the herpes simplex virus (HSV) thymidine kinase (TK) promoter, the baculovirus IE1 promoter, the elongation factor 1 alpha (EF1) promoter, the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) promoter, the phosphoglycerate kinase (PGK) promoter, the ubiquitin C (Ube) promoter, the albumin promoter, the

- 27 -

regulatory sequences of the mouse metallothionein-L promoter and transcriptional control regions, the ubiquitous promoters (HPRT, vimentin, β -actin, tubulin and the like), the promoters of the intermediate filaments (desmin, neurofilaments, keratin, GFAP, and the like), the promoters of therapeutic genes (of the MDR, CFTR or factor VIII type, and the like), pathogenesis or disease related-promoters. In addition, these expression sequences may be modified by addition of enhancer or regulatory sequences and the like.

[00101] The term "expression" refers to the biological production of a product encoded by a coding sequence. In most cases a DNA sequence, including the coding sequence, is transcribed to form a messenger-RNA (mRNA). The messenger-RNA is then translated to form a polypeptide product which has a relevant biological activity. Also, the process of expression may involve further processing steps to the RNA product of transcription, such as splicing to remove introns, and/or post-translational processing of a polypeptide product.

[00102] The VL and VH-PE38 polypeptides are expressed in cells, *e.g.*, bacterial cells, such as *E. coli*. The polypeptides are expressed, *e.g.*, in *E. coli* cells and isolated from inclusion bodies. In certain embodiments, the VL and VH-PE38 subunits are expressed in different cells. For example, the VL is expressed in one cell on a first vector and the VH-PE38 is expressed in a different cell on a second vector. Inclusion bodies from each cell line are recovered and solubilized. In certain embodiments, the inclusion bodies are solubilized at a pH in a range of about 9.0 to about 10.5. In further embodiments, the inclusion bodies are solubilized at a pH of 9.0, at a pH of 9.5, at a pH of 10.0 or a pH of 10.5. The solubilized VL and VH-PE38 inclusion bodies are combined to form an immunoconjugate comprising the VL and VH-PE38 subunits.

[00103] In other embodiments, the VL and VH-PE38 subunits are expressed in the same cell on different vectors, for example, the VL is expressed in one cell on a first vector, and the VH-PE38 is expressed in the same cell on a different vector. Inclusion bodies from the cell are recovered, solubilized and the VL and VH-PE38 subunits combined to form an immunoconjugate. In certain other embodiments, the VL and VH-PE38 subunits are expressed on the same vector in the same cell.

[00104] Downstream chromatography steps are utilized to further purify this immunoconjugate.

Chromatography Conditions

[00105] As appreciated in the art, load, wash, and elution conditions for use in the chromatography of the invention will depend on the specific chromatography media/ligands used. The process of the invention can, of course, be used in combination with other protein purification methodologies, such as salt precipitation, affinity chromatography, hydroxyapatite chromatography, reverse phase liquid chromatography, or any other commonly used protein purification technique. It is contemplated, however, that the process of the present invention will eliminate or significantly reduce the need for other purification steps.

[00106] Anionic exchange chromatography is also performed during chromatographic separation of the polypeptide of interest. As is well known in the art, anion exchangers may be based on various materials with respect to the matrix as well as to the attached charged groups. For example, the following matrices may be used, in which the materials mentioned may be more or less cross-linked: agarose based (such as SEPHAROSE Fast Flow[®] (such as Q-SEPHAROSE FF), and SEPHAROSE High Performance[®]); cellulose based (such as DEAE SEPHACEL[®]); silica based and synthetic polymer based, or resins such as SuperQ-650 (from TOSOH BIOSEP) and Macro High Q (from BIO-RAD). For the anion exchange resin, the charged groups which are covalently attached to the matrix may, *e.g.*, be diethylaminoethyl (DEAE), quaternary aminoethyl (QAE), and/or quaternary ammonium (Q). In certain embodiments, the resin is selected from the group including, but not limited to, Q Sepharose High Performance, Q Sepharose Fast Flow, DEAE Sepharose Fast Flow, Capto Q, Capto DEAE, Toyopearl SuperQ 650 (M), Toyopearl SuperQ 650 (S), Toyopearl DEAE 650 (M), Toyopearl DEAE 650 (S), TSKgel SuperQ-5PW (30), TSKgel SuperQ-5PW (20), TSKgel DEAE-5PW (30), TSKgel DEAE-5PW (20), EMD Chemicals: Fractogel EMD DEAE (S), Fractogel EMD DEAE (M), Fractogel EMD DMAE (S), Fractogel EMD DMAE (M), Fractogel EMD TMAE (S), Fractogel EMD TMAE (M), and Baker Bond XWP500 PolyQuat-35, SPE. In one embodiment of the present process, the anion exchange resin employed is Q-SEPHAROSE FF[®].

[00107] Although any of these resins may be used for small scale purification of antibodies, resins of certain size and lower cost are amenable to manufacturing scale

- 29 -

separation. If the size of the resin is too small, there is considerable back pressure generated in the system. In addition, the amount of polypeptide that can be purified is limited. If the resin is costly to make or purchase, it is not economically feasible/practical for use in large scale purification.

[00108] Thus, the resin used in the present invention must be of a certain size to provide efficient scale-up without being prohibitively expensive. "Manufacturing level purification" means purification of antibodies from a recombinant preparation on a scale that meets commercial scale production. The resin used in the predetermination step should be the same as that used in the final protocol for manufacturing level purification because one may not easily predict the variation in conditions necessary to separate the aggregates if the resin is changed. A particular resin that is useful in small scale or bench top purification may not be amenable to large scale purification. Such resins useful for the present invention include, *e.g.*, Q-SEPHAROSE HP. However, the skilled artisan would recognize other anion exchange resins useful for commercial scale production.

[00109] The volume of resin used when packing an anion exchange chromatography column is reflected by the dimensions of the column, *i.e.*, the diameter of the column and the height of the resin, and varies depending on, *e.g.*, the amount of antibody in the applied solution and the binding capacity of the resin used.

[00110] Before performing an anion exchange chromatography, the exchange resin may be equilibrated with a buffer. Any of a variety of buffers are suitable for the equilibration of exchange resin, *e.g.*, sodium acetate, sodium phosphate, TRIS(hydroxymethyl) amino-methane, TRIS, phosphate, bis-TRIS, and L-histidine. Persons skilled in the art will appreciate that numerous other buffers may be used for the equilibration as long as the pH and conductivity are about the same as for the applied antibody solution. When performing the "bind-washout" process, the equilibration buffers and the wash buffers are the same. When performing the "bind-elute" process, the elution buffers may be made of one or more buffer substances to control the pH. The salt used is, *e.g.*, a highly soluble salt, such as sodium chloride or potassium phosphate, but any salt may be used that maintains the functionality of the antibody and allows removal of the antibody monomer from the resin.

[00111] In performing the "bind-elute" process, the elution of the antibody monomers from the resin may be performed with a substantially non-denaturing buffer having a

- 30 -

pH and ionic strength sufficient to efficiently elute the monomeric antibody, thereby recovering an antibody-containing eluate, while leaving the aggregates bound to the resin. In this context, efficient elution means that at least 75%, or at least 80%, or at least 85%, or at least 90%, or at least 95% of the antibody loaded onto the resin is recovered. Only about 1.0%, preferably only 0.5%, most preferably less than 0.1% aggregates remain in the antibody preparation following ion exchange.

[00112] In one embodiment, the elution is carried out as a step gradient elution. In another embodiment, the elution is carried out in a linear gradient.

[00113] Surprisingly, deamidated variants of the immunoconjugate proteins eluted at higher salt concentration despite the apparent net increase of negative charge due to deamidation of an asparagine residue. Therefore, these reduced potency variants were separated from the more active proteins by the ion exchange chromatography described herein.

[00114] In certain embodiments of the invention, about 75% to about 99% of the acidic or deamidated variant present within the starting sample of the polypeptide or immunoconjugate is removed during the purification process. In other embodiments, at least 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% of the deamidated variant is removed. The composition comprising the active polypeptide or immunoconjugate thus has less than between about 25% and about 1% deamidated species, for example, less than 25%, less than 20%, less than 15%, less than 10%, less than 5%, less than 4%, less than 3%, less than 2% or less than 1%.

[00115] Deamidated variants of the invention include immunoconjugates comprising a PE toxin or variant thereof, wherein the deamidation occurs at one or more residues within the immunoconjugate, for example, at one or more residues within the PE toxin or variant thereof. In certain embodiments, deamidation occurs at 1, 2, 3, 4 or 5 residues within the immunoconjugate. In other embodiments, an immunoconjugate comprising a PE toxin or variant thereof is deamidated at 1, 2, 3, 4, or 5 residues within the PE toxin or variant thereof, for example at position 358 of SEQ ID NO:1, at position 495 of SEQ ID NO: 16, at position 243 of SEQ ID NO:17, at position 227 of SEQ ID NO:18, at position 200 of SEQ ID NO:19, at position 212 of SEQ ID NO: 20, at position 212 of SEQ ID NO: 21 or at position 229 of SEQ ID NO: 22.

- 31 -

[00116] In one embodiment, the salt concentration of the eluting buffer is sufficiently high to displace the antibody monomers from the resin without displacing the aggregates. However, it is contemplated that an increase in pH and a lower salt concentration can be used to elute the antibody monomers from the resin.

[00117] Any or all chromatographic steps of the present invention can be carried out by any mechanical means. Chromatography may be carried out, for example, in a column. The column may be run with or without pressure and from top to bottom or bottom to top. The direction of the flow of fluid in the column may be reversed during the chromatography process. Chromatography may also be carried out using a batch process in which the solid media is separated from the liquid used to load, wash, and elute the sample by any suitable means, including gravity, centrifugation, or filtration. Chromatography may also be carried out by contacting the sample with a filter that absorbs or retains some molecules in the sample more strongly than others. In the following description, the various embodiments of the present invention are described in the context of chromatography carried out in a column. It is understood, however, that use of a column is merely one of several chromatographic modalities that may be used, and the illustration of the present invention using a column does not limit the application of the present invention to column chromatography, as those skilled in the art may readily apply the teachings to other modalities as well, such as those using a batch process or filter.

[00118] A variety of different loading, washing and elution conditions can be used, as desired. In some embodiments, the initial loading conditions are adapted such that the protein (*e.g.*, antibody) eluted from the initial non-HT is applied directly to the HT column.

[00119] Elution can be achieved, for example, by changing the salt conditions in the liquid phase. For example, the salt and/or conductivity of the liquid phase is increased (linearly or step-wise) to a point at which the antibody elutes. Exemplary washing conditions include, *e.g.*, 10 mM phosphate, pH 6.7, with elution achieved by increasing the salt concentration (step-wise or in a linear fashion) (*e.g.*, to 10 mM phosphate, 1.5M NaCl, pH 6.7). All of the various embodiments or options described herein can be combined in any and all variations.

[00120] Before the sample is applied to the column, the column can be equilibrated in the buffer or salt that will be used to chromatograph the protein. As discussed below, chromatography (and loading of the protein to be purified) can occur in a variety of buffers or salts including sodium, potassium, ammonium, magnesium, calcium, chloride, fluoride, acetate, phosphate, and/or citrate salts and/or Tris buffer. Citrate buffers and salts are preferred by those skilled in the art for their ease of disposal. Such buffers or salts can have a pH of at least about 5.5. In some embodiments, equilibration may take place in a solution comprising a Tris or a sodium phosphate buffer. In some embodiments, equilibration takes place at a pH of at least about 5.5. Equilibration may take place at pHs between about 6.0 and about 8.6, preferably at pHs between about 6.5 and 7.5. Most preferably, the solution comprises a sodium phosphate buffer at a concentration of about 25 millimolar and at a pH of about 6.8.

[00121] The protein purification process of the present invention is applicable to removal of an acidic variant from any protein. Some proteins specifically contemplated for use with the invention include antibodies or fragments thereof. Other proteins include, but are not limited to recombinant fusion proteins comprising one or more constant antibody immunoglobulin domains, optionally an Fc portion of an antibody, and a protein of interest.

Formulations

[00122] Formulations of the purified polypeptides or immunoconjugates are prepared for storage and use by combining a purified polypeptide or immunoconjugate of the present invention with a pharmaceutically acceptable vehicle (*e.g.*, carrier, excipient) (Remington, The Science and Practice of Pharmacy 20th Edition Mack Publishing, 2000). Suitable pharmaceutically acceptable vehicles include, but are not limited to, nontoxic buffers such as phosphate, citrate, and other organic acids; salts such as sodium chloride; antioxidants including ascorbic acid and methionine; preservatives (*e.g.*, octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens, such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight polypeptides (*e.g.*, less than about 10 amino acid residues); proteins

such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; carbohydrates such as monosaccharides, disaccharides, glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and non-ionic surfactants such as TWEEN or polyethylene glycol (PEG).

[00123] The antibody and/or immunoconjugate compositions of this invention (*i.e.*, PE linked to an antibody), are particularly useful for parenteral administration, such as intravenous administration or administration into a body cavity or lumen of an organ. The compositions for administration will commonly comprise a solution of the antibody and/or immunoconjugate dissolved in a pharmaceutically acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers can be used, *e.g.*, buffered saline and the like. These solutions are sterile and generally free of undesirable matter. These compositions may be sterilized by conventional, well known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like. The concentration of fusion protein in these formulations can vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight and the like in accordance with the particular mode of administration selected and the patient's needs.

[00124] Thus, a typical pharmaceutical immunoconjugate composition for intravenous administration would be at a total treatment of about 0.3 to about 50 $\mu\text{g/kg}$ per day, in particular 20-50 $\mu\text{g/kg}$ per day with the dosage preferably administered continuously or allocated at three times per day. Actual methods for preparing administrable compositions will be known or apparent to those skilled in the art and are described in more detail in such publications as Remington's Pharmaceutical Science, 19th ed., Mack Publishing Company, Easton, Pa. (1995).

[00125] The composition including the present invention's immunoconjugate can be administered for therapeutic treatments. In therapeutic applications, compositions are administered to a patient suffering from a disease, in an amount sufficient to cure or at

- 34 -

least partially arrest the disease and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for this use will depend upon the severity of the disease and the general state of the patient's health.

[00126] Single or multiple administrations of the compositions may be administered depending on the dosage and frequency as required and tolerated by the patient. In any event, the composition should provide a sufficient quantity of the proteins of this invention to effectively treat the patient. The dosage can be administered three times a day every other day or continuously every other day for a cycle of, e.g., 21 days, but may be applied periodically until either a therapeutic result is achieved or until side effects warrant discontinuation of therapy. Generally, the dose should be sufficient to treat or ameliorate symptoms or signs of disease without producing unacceptable toxicity to the patient. An effective amount of the compound is that which provides either subjective relief of a symptom(s) or an objectively identifiable improvement as noted by the clinician or other qualified observer.

[00127] In one embodiment, the immunoconjugate is formulated as a pharmaceutical composition comprising at least one acceptable excipient. Pharmaceutically acceptable CAT-8015 immunoconjugate formulations include 0.5 mg/mL to 2.5 mg/mL CAT-8015, usually 1.0 mg/mL, 1.1 mg/mL, 1.2 mg/mL, 1.3 mg/mL, 1.4 mg/mL or 1.5 mg/mL in 25 mM sodium phosphate, 4% sucrose, 8% glycine, 0.02% polysorbate 80 (PS80), pH 7.4. In additional embodiments, the sodium phosphate can be in a range of 20 mM to 100 mM, 25 mM to 50 mM, or 25 mM to 35 mM; the sucrose can be at 2%, 3%, 4%, 5% or 6%; the glycine can be in the range of 5-10%, usually, 5%, 6%, or 7% ; the polysorbate 80 can be in a range from about 0.01% to about 1%, usually 0.01%, 0.02%, 0.03%, 0.04% or 0.05%; with a pH in the range of 6.5 to 8.0, usually at pH 7.2, 7.3, 7.4, 7.5 or 7.6. Other buffering agents known to one of ordinary skill in the art can also be utilized.

[00128] In certain embodiments of the invention, the formulation is lyophilized. The term "lyophilized" refers to any composition or formulation that is prepared in dry form by rapid freezing and dehydration, in the frozen state under high vacuum. "Lyophilizing" or "lyophilization" refers to a process of freezing and drying a solution. Lyophilized formulations or compositions are often made ready for use or reconstituted by addition of sterile distilled water. In certain embodiments, the lyophilized formulation of the invention is reconstituted into a vial.

[00129] For intravenous administration, a formulation of the invention, such as a liquid formulation or a formulation reconstituted from a lyophilized formulation is placed in a vial where the immunoconjugate in the formulation is present at concentrations as described above. This formulation is extracted from the vial and added to an intravenous (IV) bag solution, where the IV bag contains from about 30 mL to about 100 mL solution, usually 50 mL, 60 mL, 70 mL or 80 mL. A separate IV bag "protectant solution" can also be added to the total volume of the IV bag where the protectant solution contains polysorbate 80 in an amount such that the polysorbate 80 present in the final IV bag solution is in a range of 0.001% to about 3% polysorbate 80, usually in the range of about 0.01% to about 0.1%, and more usually at 0.01%, 0.02%, 0.03%, 0.04% or 0.05%. The protectant solution can be pre-formulated in a vial such that the polysorbate 80 is at a concentration of about 0.5% to about 5%, and can be 0.5%, 1.0%, 1.5%, 2.0%, 2.5%, 3.0%, 3.5%, 4.0%, 4.5% or 5.0%. The protectant solution prevents adsorption of the immunoconjugate or drug (*e.g.*, CAT-8015) to contact surfaces of the IV bag, thereby preventing or inhibiting the immunoconjugate or drug from sticking to the IV bag during administration and allowing the patient to receive the appropriate dosage of immunoconjugate or drug. The IV bag solution can be administered by infusion to the patient for various durations, usually 30 minutes to 1 hour, usually 30 minutes.

[00130] Among various uses of the immunoconjugates and formulations of the present invention are included a variety of disease conditions caused by specific human cells that may be eliminated by the toxic action of the protein. One application for the immunoconjugates of the invention is for the treatment of B cell malignancies or malignant B cells expressing CD22. Exemplary B cell malignancies include chronic B-lymphocytic cells (B-CLL), pediatric acute lymphocytic leukemia (pALL), follicular lymphoma (FL), diffuse large B cell lymphoma (DLBCL), Non Hodgkins lymphoma (NHL) and hairy cell leukemia (HCL).

[00131]

[00132] Although the present invention has been fully described in conjunction with several embodiments thereof with reference to the accompanying drawings, it is to be understood that various changes and modifications may be apparent to those skilled in the

- 36 -

art. Such changes and modifications are to be understood as included within the scope of the present invention as defined by the appended claims, unless they depart therefrom.

EXAMPLES

Example 1. Expression, recovery and inclusion body isolation of CAT-8015

[00133] Fermentation of separate cell lines containing CAT-8015 VL and CAT-8015 VH-PE38 expression vectors was performed. The fermentor was harvested by continuous centrifugation. The fermentor harvest was passed through a continuous centrifuge at 2 to 8°C at a rate of 0.5 to 0.8 L per minute and centrifuged at a speed of approximately 15,000 rpm. After centrifugation the cell paste was frozen at $\leq -70^{\circ}\text{C}$.

[00134] Following this treatment, the VH-PE38 and VL cell pastes were thawed for 12 to 24 hours at 2 to 8°C. The cells were lysed to release inclusion bodies containing the VL and VH-PE38 products. The resulting inclusion bodies were subsequently solubilized and the VH-PE38 and VL products obtained.

[00135] The product was concentrated to approximately 1 mg/mL (determined by Coomassie total protein assay) using a 30 kDa ultrafiltration hollow fiber cartridge. The retentate was then diafiltered with 5 to 6 volumes of 20 mM Tris, 100 mM urea pH 7.4 to achieve a conductivity of 2.5 to 3.0 mS/cm. This product was stored up to 72 hours at 2 to 8°C.

Example 2. Analytical-scale Purification of Active CAT-8015 by Anion Exchange Chromatography with High Performance Resins.

[00136] Expression of the VH-P38 subunit resulted in a formation of a deamidated variant of the subunit. The deamidation was found to occur in the PE38 portion of the immunoconjugate. Deamidation of the VH-P38 subunit resulted in decreased potency of the CAT-8015 protein. Surprisingly, the below described chromatographic conditions were successful in removing the deamidated variant, thus providing the ability to remove the inactive species during purification. Since the deamidation occurred in the PE38 portion of the fusion construct, the chromatographic conditions can be applied to removal of any deamidated variant of a PE conjugate.

51332-114

- 37 -

[00137] CAT-8015 was renatured from isolated inclusion bodies and subsequently purified by a 4-column process. Table 1 provides an overview of the renaturation and purification unit operations.

Table 1 - CAT-8015 production flowchart

Step	Unit Operation
1	Fermentation
2	Primary Recovery
3	Inclusion Body Isolation
4	Refold
5	Capture Step
6	Intermediate Purification Step I
7	Intermediate Purification Step II
8	Anion Exchange Chromatography
9	Formulation (Drug Substance)

[00138] Figure 1 shows the analytical ion exchange chromatography (IEC) profile of a sample of a CAT-8015 reference standard. As shown in the profile, a pre-peak emerges prior to the elution of the main peak. The individual fractions eluted from the IEC are assayed for CAT-8015 biological activity relative to a reference standard using an apoptosis bioassay that measures the ability of the test sample to induce dose dependent apoptotic death of the CD22 receptor-expressing Daudi cell line. Once bound to CD22 and internalized, CAT-8015 induces Daudi cells to undergo apoptosis via Caspase 3/7 activation that can be measured by Caspase-Glo™ 3/7 Assay System. The potency of the test sample is determined by dividing the 50% effective concentration (EC50) of the Reference Standard by the EC50 of the test sample. The results of the apoptosis bioassay

- 38 -

demonstrated that the relative potency of CAT-8015 is correlated with the percentage of pre-peak of CAT-8015 as diagrammed in Figure 1. Figure 2 shows the correlation between this relative potency and the percentage of pre-peak.

[00139] Pre-peak and main peak fractions from multiple IEC analysis were collected, pooled and subjected to peptide mapping and liquid chromatography-mass spectrometry/mass spectrometry (LC-MS/MS) analysis. Results were compared to those obtained from peptide mapping and LC-MS/MS experiments of purified CAT-8015. The analysis of purified CAT-8015 drug substance revealed that Asn-358 was partially deamidated to Asp-358 and iso-Asp-358 (Table 2). Asp-358 and iso-Asp-358 were found to be significantly enriched in the pre-peak fraction whereas the main peak fraction was enriched in intact CAT-8015 (Table 2). Taken together, the results demonstrate that deamidation at Asn-358 lead to a loss of relative potency in a cell based bioassay. The Asp-358 residue is present within the PE toxin portion of the immunoconjugate thus indicating that an immunoconjugate containing a PE toxin, or variant thereof, in which Asp-358 is present will likely be subject to deamidation and subject to a loss of potency or activity.

Table 2

Distribution of Amino Acid 358 in CAT-8015 Drug Substance, Pre-Peak and Main Peak Fractions based on Peptide Mapping and LC-MS/MS Analysis.

Amino Acid	Drug Substance (%)	Pre-Peak (%)	Main Peak (%)
N358	78.1	3.1	88.9
D358	11.9	44.0	2.2
iso-D358	10.0	52.9	8.9

D358 = deamidated Asn-358; iso-D358 = iso-deamidated Asn-358; N358 = Asn-358.

[00140] Separation of deamidated CAT-8015 from intact CAT-8015 was achieved by anion exchange chromatography with strong ion exchange groups such as Q (quaternary amino) coupled to small diameter resins such as Source 15 (particle diameter: 15µm; GE Healthcare) and Sepharose High Performance (particle diameter: 34µm; GE Healthcare). The application of small diameter chromatography resin in biomanufacturing processes is complicated by the generation of significant back pressures under typical operating conditions as defined by column geometry, flow rates and buffer composition. Based on these considerations and the requirement for a high resolution chromatography step Q

Sepharose High Performance was chosen for the separation of deamidated CAT-8015 from intact CAT-8015. Chromatography conditions were developed that achieved high resolution while maintaining operability at various manufacturing scales.

Example 3. Bench-scale purification of CAT-8015

[00141] The column was first pre-equilibrated with 5 column volumes (CVs) of Buffer C (Pre-Equilibration/Stripping Buffer: 10mM Tris/HCl, pH 8.0, 1.0M NaCl) and subsequently equilibrated with 5 CVs Buffer A (Equilibration Buffer: 10mM Tris/HCl, pH 8.0) at a linear flow rate of 100cm/hr. The chromatography resin was Q Sepharose High Performance (QHP, GE Healthcare) in a Millipore VantageTM column, 2.2cm x 19.5cm, and run on an AKTA Explorer. The intermediate purification product pool was prepared for loading onto the high performance anion exchange column by diafiltration with 10 volumes of Buffer A using a 10 kDa MWCO membrane. The diafiltered hydrophobic interaction product pool was loaded onto the QHP column at a linear flow rate of 100cm/hr, followed by a 2 CV re-equilibration step with Buffer A at the same flow rate. CAT-8015 was eluted with a 10 CV linear gradient from 35% Buffer B (Elution Buffer: 10mM Tris/HCl, pH 8.0, 500mM NaCl) to 55% Buffer B at a linear flow rate of 100cm/hr. Elution of product was monitored at 280nm. Fractions were collected and analyzed for % pre-peak by analytical ion exchange chromatography (IEC). Fractions containing less than 25% pre-peak were pooled. The QHP pool was analyzed for % pre-peak by analytical IEC on a strong anion exchange column. Relative potency was measured by an apoptosis bioassay as described above.

[00142] At pH 8.0, CAT-8015 strongly bound to the anion exchange resin with no protein detected in the flow-through fraction by absorbance at A280. After an initial wash step with 175mM NaCl in Tris/HCl, pH 8.0, CAT-8015 was eluted from the column with a linear salt gradient from 35% B (175mM NaCl in Tris/HCl, pH 8.0) to 55% B (275mM NaCl in Tris/HCl, pH 8.0). CAT-8015 eluted from the column between 39% B (192 mM NaCl in Tris/HCl, pH 8.0) and 49%B (245 mM NaCl in Tris/HCl, pH 8.0). Fig. 3 shows the QHP chromatography profile of CAT-8015.

[00143] Fractions were analyzed by analytical IEC. Table 3 shows the results for fractions eluted between 44.5% B (223mM NaCl) and 47.2% B (236mM NaCl).

Table 3
IEC Analysis of Collected QHP Fractions

Fraction No.	% Pre-Peak^a	% Main Peak
C12	41.0	59
D12	34.5	65.5
D11	27.0	73
D9	17.6	82.4
D7	15.9	84.1
D5	18.7	81.3
D3	21.9	78.1

^a Pre-peak contains >90% deamidated CAT-8015.

[00144] Table 3 demonstrates that anion exchange chromatography operated in a linear salt gradient elution mode is able to separate deamidated CAT-8015 from intact CAT 8015 in an effective manner.

[00145] Surprisingly, intact CAT-8015 was eluted at a higher salt concentration despite the apparent net increase of negative charge due to deamidation of an asparagine residue. This result is consistent with the chromatography profile observed by IEC. CAT-8015-containing samples were injected on an analytical anion exchange column (PL-SAX, Varian) equilibrated at pH 8.0 with a Tris/HCl buffer system and eluted by a combination of step and gradient elution steps (Fig.1).

[00146] Fractions were combined according to the pooling criteria of less than 25% pre-peak content. The QHP pool was analyzed for % pre-peak and relative potency by SDS-PAGE, analytical IEC and apoptosis bioassay. The SDS-PAGE analysis, as shown in Fig. 4, demonstrates that QHP load pool and eluate samples contained highly purified CAT-8015. However, the QHP load pool did not meet target specification for purity by IEC and bioactivity. Purity and potency measurements for the QHP load pool as compared to the QHP eluate pool, as presented in Table 4 below, demonstrate that the anion exchange chromatography step with QHP resulted in a significant increase in purity by IEC and relative potency of CAT-8015. The QHP load pool was generated from Intermediate Purification Step II.

Table 4**CAT-8015 Purity by IEC and Bioactivity**

Step	% Pre-Peak	% Main Peak	Relative Potency (%)
QHP Load Pool	53.8	46.2	52
QHP Eluate Pool	16.5	83.8	80

[00147] The QHP product pool was subsequently diafiltered into formulation buffer to generate CAT-8015 drug substance.

[00148] Thus, the manufacture of CAT-8015 drug substance requires the separation of deamidated CAT-8015 from active CAT-8015. The capability of anion exchange chromatography with high performance resins such as QHP to separate deamidated CAT-8015 from intact CAT-8015 and to increase bioactivity to target specifications is a prerequisite for the successful manufacture of CAT-8015 drug substance.

Example 4. Large-scale purification of CAT-8015

[00149] The column was first pre-equilibrated with 5 CVs Buffer C (Pre-Equilibration/Stripping Buffer: 10mM Tris/HCl, pH 8.0, 1.0M NaCl) at a linear flow rate of 66cm/hr and subsequently equilibrated with 5 CVs Buffer A at a linear flow rate of 76cm/hr. The chromatography resin was Q Sepharose High Performance (QHP, GE Healthcare), in a BP300, 30cm x 22cm column bed, run on a K Prime instrument. The intermediate purification product pool was prepared for loading onto the high performance anion exchange column by diafiltration with 10 volumes of Buffer A (Equilibration Buffer: 10mM Tris/HCl, pH 8.0) using a 10 kDa MWCO membrane. The diafiltered product pool was loaded onto the QHP column at a linear flow rate of 64 cm/hr, followed by a 2 CV re-equilibration step with Buffer A at 76cm/hr. CAT-8015 was eluted with a 10 CV linear gradient from 35% Buffer B (Elution Buffer: 10mM Tris/HCl, pH 8.0, 500mM NaCl) to 55% Buffer B at a linear flow rate of 76 cm/hr. Elution of product was monitored at 280nm. Fractions were collected and analyzed for % pre-peak by analytical ion exchange chromatography (IEC). Fractions containing less than 25% pre-peak were pooled. The QHP pool was analyzed for % pre-peak by analytical IEC on a strong anion exchange column. Relative potency was measured by an apoptosis bioassay.

- 42 -

[00150] Large scale anion exchange chromatography of CAT-8015 with QHP was performed as described above. QHP purification was carried out at reduced flow rates due to equipment constraints. CAT-8015 eluted from the column at conductivities between 22.3mS/cm and 26.4mS/cm. Figure 5 shows the QHP chromatography profile of CAT-8015 purified according to the method described above.

[00151] Fractions were analyzed by analytical IEC. Table 5 shows the results for fractions eluted at conductivities between 23.8 and 25.4mS/cm. Table 5 demonstrates that anion exchange chromatography operated in a linear salt gradient elution mode was able to separate deamidated CAT-8015 from intact CAT 8015. Separation of deamidated CAT-8015 from intact CAT-8015 took place within a conductivity range of less than 2mS/cm.

Table 5

IEC Analysis of Collected Fractions% Pre-peak purity

Fraction	% Pre-Peak	% Main Peak
1	55.1	44.9
2	39.0	61
3	30.1	69.9
4	25.1	74.9
5	18.7	81.3
6	14.7	85.3
7	16.4	83.6

[00152] Fractions 4-7 were combined according to the pooling criteria of less than 25% pre-peak content. The QHP pool was analyzed for % pre-peak and relative potency by SDS-PAGE and SEC, analytical IEC and apoptosis bioassay. The SDS-PAGE analysis, as shown in Fig. 6, demonstrates that QHP load pool and eluate samples contained highly purified CAT-8015. However, the QHP load pool did not meet target specification for purity by SEC, IEC and relative potency. Purity and potency measurements for the QHP load pool as compared to the QHP eluate pool, as presented in Tables 6 and 7 below, demonstrate that the anion exchange chromatography step with QHP resulted in a significant increase in purity by SEC, IEC and relative potency of CAT-8015. The QHP load pool was generated from Intermediate Purification Step II.

- 43 -

Table 6**CAT-8015 Purity by SEC**

Step	% Monomer	% Aggregate	% Other
QHP Load Pool	92.7	0.7	6.6
QHP Eluate Pool	99.0	1.0	0

Table 7**CAT-8015 Purity by IEC and Bioactivity**

Step	% Pre-Peak	% Main Peak	Relative Potency (%)
QHP Load Pool	50.3	49.7	51
QHP Eluate Pool	17	83	75

[00153] The QHP product pool was subsequently diafiltered into formulation buffer to generate CAT-8015 drug substance.

[00154] Examples 2-4 demonstrate the capacity of anion exchange chromatography with resins such as Q Sepharose High Performance to separate deamidated CAT-8015 from intact CAT-8015 and to increase its relative potency to meet target specifications (see Tables 4 and 6). Deamidated CAT-8015 differs from intact CAT-8015 by one additional negative charge. In contrast to the expected elution behavior from an anion exchange column, the bulk of deamidated CAT-8015 elutes prior to intact CAT-8015 under salt gradient elution conditions (see Tables 3 and 5). This unexpected elution pattern was observed at analytical scale, bench scale, and large scale anion exchange chromatography. This elution pattern was unexpected as the linear high salt elution buffer would be expected to result in a higher negative charge of the variant. Thus, Examples 2-4 demonstrate that using a linear elution buffer, a deamidated species can be removed from active immunoconjugates using anion-exchange chromatography. Separation of deamidated CAT-8015 from intact CAT-8015 took place within a particular range of conductivities, underscoring the need for high resolution anion exchange resins, careful control of elution conditions and in-process testing of collected fractions.

Example 5. Modifying the bioactivity of CAT-8015 formulations

[00155] The potency of CAT-8015 compositions was calibrated by mixing specific quantities of the deamidated pre-peak with the active main peak. To obtain compositions comprising a particular potency of CAT-8015, aliquots of CAT-8015 pre-peak product were combined with aliquots of CAT-8015 main peak product to achieve a composition with a particular potency of CAT-8015. By controlling the level of CAT-8015 potency in the composition, a CAT-8015 formulation was generated for administration of a particular volume of reconstituted CAT-8015 at a desired dose.

Example 6. Adjusting pH During Solubilization Results in Reduced Deamidation Species as Measured after Capture and Intermediate Purification Steps

[00156] While deamidated species can be removed from active immunoconjugates using the purification steps as described above, levels of deamidated species of CAT-8015 can also be effectively reduced by adjusting the pH at earlier steps in the purification process (*i.e.*, the refold step (Step 4 of Table 1 above). The refold procedure utilized to achieve a lower level of deamidated species of CAT-8015 includes the following substeps:

[00157] Refold Substep 1: Solubilization, Clarification and Concentration: VH-PE38 and VL inclusion bodies were thawed for 12-24 hours at room temperature (15-30°C). VH-PE38 and VL inclusion bodies were combined in a 1:1 molar ratio and adjusted to 15% (w/v) solids by adding 50 mM Tris, 20 mM EDTA, pH 7.4. The inclusion bodies were solubilized by adding 5 kg of inclusion body solubilization buffer (50 mM ethanolamine, 8 M urea, 0.5 M arginine, 2 mM EDTA, 10 mM DTE) for each kg of 15% (w/v) solids inclusion body suspension. The pH of the inclusion body solubilization buffer was varied between pH 9.0 and 10.5 in 0.5 pH unit increments. Solubilization was carried out for 2 hours at room temperature (15-30°C) with constant stirring. Solubilized inclusion bodies were clarified by depth filtration through a series of filters. The clarified filtrate was concentrated by tangential flow filtration to 5-6 g/L using a 5 kDa molecular weight cutoff (MWCO) ultra filtration membrane.

[00158] Refold Substep 2: Refold: The refolding of CAT-8015 was initiated by a 10-fold dilution of the clarified and concentrated inclusion body filtrate into pre-chilled (2-8°C) refolding buffer (50 mM ethanolamine, 1 M arginine, 2 mM EDTA, 0.91 mM oxidized glutathione, pH 9.5). The refold solution was maintained at 2-8°C for 48-72 hours with

- 45 -

continuous mixing. The refold was terminated by bringing the refold solution to room temperature (15-30°C) prior to concentration and diafiltration. The refold solution was concentrated by tangential flow filtration with a 10 kDa MWCO membrane, and diafiltered with 10 volumes of 20 mM potassium phosphate, pH 7.4. The concentrated and diafiltered refold solution was filtered through a 0.2 µm filter (TMAE load).

[00159] As part of the capture step (Step 5 of Table 1 above), the CAT-8015 preparation obtained from the refold procedure above was loaded onto a Fractogel TMAE column (EMD Biosciences or equivalent) equilibrated with 20 mM potassium phosphate, pH 7.4. After loading, the column was first washed with 20 mM potassium phosphate, pH 7.4, and then with 20 mM potassium phosphate, 0.1% (w/w) Triton X 100, pH 7.4, followed by a subsequent wash with 20 mM potassium phosphate, 100 mM sodium chloride, pH 7.4. The product was eluted from the column in reverse flow with 20 mM potassium phosphate, 200 mM sodium chloride pH 7.4. The column was stripped with 2 M sodium chloride, sanitized with 1 N sodium hydroxide and stored in 0.1 N sodium hydroxide at room temperature.

[00160] As part of the intermediate purification step 1, hydroxyapatite chromatography was performed. The hydroxyapatite chromatography step was operated as a flow-through chromatography step. The product obtained from the capture step above was loaded directly without any further adjustments onto a ceramic hydroxyapatite column (Bio-Rad Laboratories or equivalent) equilibrated with 400 mM potassium phosphate, 200 mM sodium chloride, pH 7.4, followed by 20 mM potassium phosphate, 200 mM sodium chloride, pH 7.4. Under the conditions of the chromatography, the product was collected in the flow-through fraction (HA product). The column was stripped with 400 mM potassium phosphate, 200 mM sodium chloride, pH 7.4, sanitized with 1 N sodium hydroxide and stored in 0.1 N sodium hydroxide at room temperature.

[00161] The percent pre-peak in the HA product from above was analyzed by high performance anion exchange chromatography. Table 8 and Figure 7 show the percent pre-peak in HA product as a function of solubilization pH. As shown in Table 8 and Figure 7, solubilizing the VH-PE38 and VL inclusion bodies at a lower pH leads to less deamidated CAT-8015 product. The capability of controlling CAT-8015 deamidation at an early step in the renaturation and purification process can increase overall process yield while maintaining the quality of the final purified drug substance.

- 46 -

Table 8**Percent Pre-Peak in HA Product as a Function of Solubilization pH**

Solubilization pH	9.0	9.5	10.0	10.5
Pre Peak (%)	9.8	14.5	22.1	31.8

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 51332-114 Seq 15-01-13 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> MEDIMMUNE, LLC

<120> METHOD FOR PURIFYING ACTIVE POLYPEPTIDES OR IMMUNOCONJUGATES

<130> MED0581.PCT

<140> PCT/US2011/045524

<141> 2011-07-27

<150> 61/369,148

<151> 2010-07-30

<160> 23

<170> PatentIn version 3.5

<210> 1

<211> 476

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 1

Met	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly
1				5					10					15	
Gly	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile
			20					25					30		
Tyr	Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Cys	Leu	Glu	Trp
		35					40					45			
Val	Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr
	50					55				60					
Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu
65					70				75					80	
Tyr	Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr
				85					90					95	

46b

Cys Ala Arg His Ser Gly Tyr Gly Thr His Trp Gly Val Leu Phe Ala
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Lys Ala Ser Gly
 115 120 125
 Gly Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys
 130 135 140
 His Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp
 145 150 155 160
 Glu Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu
 165 170 175
 Tyr Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg
 180 185 190
 Asn Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile
 195 200 205
 Arg Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala
 210 215 220
 Glu Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly
 225 230 235 240
 Ala Ala Asn Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn
 245 250 255
 Tyr Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe
 260 265 270
 Ser Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala
 275 280 285
 His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly
 290 295 300
 Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala
 305 310 315 320
 Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly
 325 330 335
 Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala
 340 345 350
 Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg
 355 360 365
 Ser Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro
 370 375 380
 Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu
 385 390 395 400
 Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu
 405 410 415
 Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser
 420 425 430
 Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser
 435 440 445
 Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala
 450 455 460
 Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 465 470 475

<210> 2

<211> 108

<212> PRT

<213> Artificial Sequence

46c

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 2

Met	Asp	Ile	Gln	Met	Thr	Gln	Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu
1				5					10					15	
Gly	Asp	Arg	Val	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Gln	Asp	Ile	Ser	Asn
			20					25					30		
Tyr	Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gly	Thr	Val	Lys	Leu	Leu
		35					40					45			
Ile	Tyr	Tyr	Thr	Ser	Ile	Leu	His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser
	50					55				60					
Gly	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu
65					70					75				80	
Gln	Glu	Asp	Phe	Ala	Thr	Tyr	Phe	Cys	Gln	Gln	Gly	Asn	Thr	Leu	Pro
			85					90						95	
Trp	Thr	Phe	Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys				
			100					105							

<210> 3

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 3

Lys	Asp	Glu	Leu
1			

<210> 4

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 4

Arg	Glu	Asp	Leu
1			

<210> 5

<211> 5

<212> PRT

<213> Artificial Sequence

46d

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 5

Arg	Glu	Asp	Leu	Lys
1				5

<210> 6

<211> 124

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 6

Met	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly
1				5					10					15	
Gly	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile
			20					25					30		
Tyr	Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Cys	Leu	Glu	Trp
		35					40					45			
Val	Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr
	50					55					60				
Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu
65					70				75					80	
Tyr	Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr
			85					90					95		
Cys	Ala	Arg	His	Ser	Gly	Tyr	Gly	Thr	His	Trp	Gly	Val	Leu	Phe	Ala
			100				105					110			
Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ala				
		115					120								

<210> 7

<211> 124

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 7

Met	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly
1				5					10					15	
Gly	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile
			20					25					30		
Tyr	Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Cys	Leu	Glu	Trp
		35					40					45			
Val	Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr
	50					55					60				

46e

Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu
 65 70 75 80
 Tyr Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr
 85 90 95
 Cys Ala Arg His Ser Gly Tyr Gly Tyr Asn Trp Gly Val Leu Phe Ala
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
 115 120

<210> 8
 <211> 124
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 polypeptide

<400> 8
 Met Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly
 1 5 10 15
 Gly Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ile
 20 25 30
 Tyr Asp Met Ser Trp Val Arg Gln Thr Pro Glu Lys Cys Leu Glu Trp
 35 40 45
 Val Ala Tyr Ile Ser Ser Gly Gly Gly Thr Thr Tyr Tyr Pro Asp Thr
 50 55 60
 Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu
 65 70 75 80
 Tyr Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr
 85 90 95
 Cys Ala Arg His Ser Gly Tyr Gly Thr Thr Trp Gly Val Leu Phe Ala
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
 115 120

<210> 9
 <211> 124
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 polypeptide

<400> 9
 Met Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly
 1 5 10 15
 Gly Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ile
 20 25 30
 Tyr Asp Met Ser Trp Val Arg Gln Thr Pro Glu Lys Cys Leu Glu Trp
 35 40 45
 Val Ala Tyr Ile Ser Ser Gly Gly Gly Thr Thr Tyr Tyr Pro Asp Thr
 50 55 60

Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu
65					70				75						80
Tyr	Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr
			85						90					95	
Cys	Ala	Arg	His	Ser	Gly	Tyr	Gly	Ser	Thr	Tyr	Gly	Val	Leu	Phe	Ala
			100					105					110		
Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ala				
		115					120								

<210> 10
 <211> 124
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic polypeptide

Met	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly
1			5					10					15		
Gly	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile
		20				25					30				
Tyr	Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Cys	Leu	Glu	Trp
	35					40					45				
Val	Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr
	50				55			60							
Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu
65					70				75						80
Tyr	Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr
			85					90						95	
Cys	Ala	Arg	His	Ser	Gly	Tyr	Gly	Thr	His	Trp	Gly	Val	Leu	Phe	Ala
			100				105						110		
Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ala				
		115					120								

<210> 11
 <211> 124
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic polypeptide

Met	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly
1			5					10					15		
Gly	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ile
		20				25					30				
Tyr	Asp	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Cys	Leu	Glu	Trp
	35					40					45				
Val	Ala	Tyr	Ile	Ser	Ser	Gly	Gly	Gly	Thr	Thr	Tyr	Tyr	Pro	Asp	Thr
	50				55			60							

46g

Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu
65					70				75						80
Tyr	Leu	Gln	Met	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr
			85						90					95	
Cys	Ala	Arg	His	Ser	Gly	Tyr	Gly	Ser	Ser	Tyr	Gly	Val	Leu	Phe	Ala
			100					105					110		
Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ala				
		115						120							

<210> 12
 <211> 108
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic polypeptide

Met	Asp	Ile	Gln	Met	Thr	Gln	Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu
1				5					10					15	
Gly	Asp	Arg	Val	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Gln	Asp	Ile	Ala	Arg
			20					25					30		
Tyr	Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gly	Thr	Val	Lys	Leu	Leu
		35					40					45			
Ile	Tyr	Tyr	Thr	Ser	Ile	Leu	His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser
	50					55					60				
Gly	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu
65					70				75						80
Gln	Glu	Asp	Phe	Ala	Thr	Tyr	Phe	Cys	Gln	Gln	Gly	Asn	Thr	Leu	Pro
			85					90						95	
Trp	Thr	Phe	Gly	Cys	Gly	Thr	Lys	Leu	Glu	Ile	Lys				
			100					105							

<210> 13
 <211> 108
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic polypeptide

Met	Asp	Ile	Gln	Met	Thr	Gln	Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu
1				5					10					15	
Gly	Asp	Arg	Val	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Gln	Asp	Ile	His	Gly
			20					25					30		
Tyr	Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gly	Thr	Val	Lys	Leu	Leu
		35					40					45			
Ile	Tyr	Tyr	Thr	Ser	Ile	Leu	His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser
	50					55					60				
Gly	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu
65					70				75						80

Gln Glu Asp Phe Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro
85 90 95
Trp Thr Phe Gly Cys Gly Thr Lys Leu Glu Ile Lys
100 105

```
<210> 14
<211> 108
<212> PRT
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

```

<400> 14
Met Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu
1          5          10          15
Gly Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Gly Arg
20          25          30
Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu
35          40          45
Ile Tyr Tyr Thr Ser Ile Leu His Ser Gly Val Pro Ser Arg Phe Ser
50          55          60
Gly Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu
65          70          75          80
Gln Glu Asp Phe Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro
85          90          95
Trp Thr Phe Gly Cys Gly Thr Lys Leu Glu Ile Lys
100          105

```

```
<210> 15
<211> 108
<212> PRT
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

```

<400> 15
Met Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu
1          5          10          15
Gly Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Arg Gly
20          25          30
Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu
35          40          45
Ile Tyr Tyr Thr Ser Ile Leu His Ser Gly Val Pro Ser Arg Phe Ser
50          55          60
Gly Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu
65          70          75          80
Gln Glu Asp Phe Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro
85          90          95
Trp Thr Phe Gly Cys Gly Thr Lys Leu Glu Ile Lys
100          105

```


<210> 16

<211> 597

<212> PRT

<213> Pseudomonas aeruginosa

<400> 16

Ala	Glu	Glu	Ala	Phe	Asp	Leu	Trp	Asn	Glu	Cys	Ala	Lys	Ala	Cys	Val
1				5					10					15	
Leu	Asp	Leu	Lys	Asp	Gly	Val	Arg	Ser	Ser	Arg	Met	Ser	Val	Asp	Pro
			20					25					30		
Ala	Ile	Ala	Asp	Thr	Asn	Gly	Gln	Gly	Val	Leu	His	Tyr	Ser	Met	Val
		35				40						45			
Leu	Glu	Gly	Gly	Asn	Asp	Ala	Leu	Lys	Leu	Ala	Ile	Asp	Asn	Ala	Leu
	50					55					60				
Ser	Ile	Thr	Ser	Asp	Gly	Leu	Thr	Ile	Arg	Leu	Glu	Gly	Gly	Val	Glu
65					70					75					80
Pro	Asn	Lys	Pro	Val	Arg	Tyr	Ser	Tyr	Thr	Arg	Gln	Ala	Arg	Gly	Ser
				85					90					95	
Trp	Ser	Leu	Asn	Trp	Leu	Val	Pro	Ile	Gly	His	Glu	Lys	Pro	Ser	Asn
			100					105						110	
Ile	Lys	Val	Phe	Ile	His	Glu	Leu	Asn	Ala	Gly	Asn	Gln	Leu	Ser	His
		115				120						125			
Met	Ser	Pro	Ile	Tyr	Thr	Ile	Glu	Met	Gly	Asp	Glu	Leu	Leu	Ala	Lys
	130					135					140				
Leu	Ala	Arg	Asp	Ala	Thr	Phe	Phe	Val	Arg	Ala	His	Glu	Ser	Asn	Glu
145					150					155					160
Met	Gln	Pro	Thr	Leu	Ala	Ile	Ser	His	Ala	Gly	Val	Ser	Val	Val	Met
				165					170					175	
Ala	Gln	Thr	Gln	Pro	Arg	Arg	Glu	Lys	Arg	Trp	Ser	Glu	Trp	Ala	Ser
			180					185					190		
Gly	Lys	Val	Leu	Cys	Leu	Leu	Asp	Pro	Leu	Asp	Gly	Val	Tyr	Asn	Tyr
	195						200					205			
Leu	Ala	Gln	Gln	Arg	Cys	Asn	Leu	Asp	Asp	Thr	Trp	Glu	Gly	Lys	Ile
	210					215					220				
Tyr	Arg	Val	Leu	Ala	Gly	Asn	Pro	Ala	Lys	His	Asp	Leu	Asp	Ile	Lys
225					230					235					240
Pro	Thr	Val	Ile	Ser	His	Arg	Leu	His	Phe	Pro	Glu	Gly	Gly	Ser	Leu
				245					250					255	
Ala	Ala	Leu	Thr	Ala	His	Gln	Ala	Cys	His	Leu	Pro	Leu	Glu	Thr	Phe
			260					265					270		
Thr	Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly
	275						280					285			
Tyr	Pro	Val	Gln	Arg	Leu	Val	Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser
	290					295					300				
Trp	Asn	Gln	Val	Asp	Gln	Val	Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly
305					310					315					320
Ser	Gly	Gly	Asp	Leu	Gly	Glu	Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala
			325						330					335	
Arg	Leu	Ala	Leu	Thr	Leu	Ala	Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg
			340					345					350		
Gln	Gly	Thr	Gly	Asn	Asp	Glu	Ala	Gly	Ala	Ala	Asn	Gly	Pro	Ala	Asp
	355					360						365			
Ser	Gly	Asp	Ala	Leu	Leu	Glu	Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe
	370					375					380				
Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn
385					390					395					400

46j

Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu Arg
 405 410 415
 Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala Gln
 420 425 430
 Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp Ala
 435 440 445
 Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr Gly
 450 455 460
 Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn Gly
 465 470 475 480
 Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe Tyr
 485 490 495
 Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val Glu
 500 505 510
 Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr Gly
 515 520 525
 Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu Gly Trp Pro Leu
 530 535 540
 Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro Thr Asp Pro Arg
 545 550 555 560
 Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro Asp Lys Glu Gln
 565 570 575
 Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro Gly Lys Pro Pro
 580 585 590
 Arg Glu Asp Leu Lys
 595

<210> 17

<211> 361

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 polypeptide

<400> 17

Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro
 1 5 10 15
 Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu
 20 25 30
 Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala
 35 40 45
 Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu
 50 55 60
 Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln
 65 70 75 80
 Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu
 85 90 95
 Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Asn
 100 105 110
 Ala Asp Val Val Ser Leu Thr Cys Pro Val Ala Ala Gly Glu Cys Ala
 115 120 125
 Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr
 130 135 140

46k

Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser Thr Arg
 145 150 155 160
 Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln
 165 170 175
 Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu
 180 185 190
 Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln
 195 200 205
 Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala
 210 215 220
 Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg
 225 230 235 240
 Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu
 245 250 255
 Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu Ala Ala
 260 265 270
 Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp
 275 280 285
 Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr Ile Leu
 290 295 300
 Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala Ile Pro
 305 310 315 320
 Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser Ile Pro
 325 330 335
 Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser Gln Pro
 340 345 350
 Gly Lys Pro Pro Arg Glu Asp Leu Lys
 355 360

<210> 18

<211> 345

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 polypeptide

<400> 18

Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His Leu Pro
 1 5 10 15
 Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu
 20 25 30
 Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr Leu Ala
 35 40 45
 Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn Ala Leu
 50 55 60
 Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg Glu Gln
 65 70 75 80
 Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu Ser Glu
 85 90 95
 Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala Ala Asn
 100 105 110
 Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr Pro Thr
 115 120 125

Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg
130						135					140				
Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln
145					150					155					160
Leu	Glu	Glu	Arg	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu
				165					170					175	
Glu	Ala	Ala	Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln
			180					185				190			
Asp	Leu	Asp	Ala	Ile	Trp	Arg	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala
	195						200				205				
Leu	Ala	Tyr	Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Arg	Gly	Arg
210					215					220					
Ile	Arg	Asn	Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu
225					230					235					240
Pro	Gly	Phe	Tyr	Arg	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala
				245					250					255	
Gly	Glu	Val	Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp
			260					265					270		
Ala	Ile	Thr	Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu
		275					280					285			
Gly	Trp	Pro	Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro
290					295					300					
Thr	Asp	Pro	Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro
305					310				315						320
Asp	Lys	Glu	Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro
				325					330					335	
Gly	Lys	Pro	Pro	Arg	Glu	Asp	Leu	Lys							
		340					345								

<210> 19

<211> 318

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 19

Met	Trp	Glu	Gln	Leu	Glu	Gln	Cys	Gly	Tyr	Pro	Val	Gln	Arg	Leu	Val
1				5				10						15	
Ala	Leu	Tyr	Leu	Ala	Ala	Arg	Leu	Ser	Trp	Asn	Gln	Val	Asp	Gln	Val
			20					25					30		
Ile	Arg	Asn	Ala	Leu	Ala	Ser	Pro	Gly	Ser	Gly	Gly	Asp	Leu	Gly	Glu
		35					40					45			
Ala	Ile	Arg	Glu	Gln	Pro	Glu	Gln	Ala	Arg	Leu	Ala	Leu	Thr	Leu	Ala
		50				55					60				
Ala	Ala	Glu	Ser	Glu	Arg	Phe	Val	Arg	Gln	Gly	Thr	Gly	Asn	Asp	Glu
65					70				75					80	
Ala	Gly	Ala	Ala	Asn	Gly	Pro	Ala	Asp	Ser	Gly	Asp	Ala	Leu	Leu	Glu
				85				90					95		
Arg	Asn	Tyr	Pro	Thr	Gly	Ala	Glu	Phe	Leu	Gly	Asp	Gly	Gly	Asp	Val
			100				105					110			
Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln	Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu
		115					120					125			

46m

Gln Ala His Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr
 130 135 140
 His Gly Thr Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val
 145 150 155 160
 Arg Ala Arg Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile
 165 170 175
 Ala Gly Asp Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro
 180 185 190
 Asp Ala Arg Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val
 195 200 205
 Pro Arg Ser Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala
 210 215 220
 Ala Pro Glu Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu
 225 230 235 240
 Pro Leu Arg Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg
 245 250 255
 Leu Glu Thr Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile
 260 265 270
 Pro Ser Ala Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp
 275 280 285
 Pro Ser Ser Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp
 290 295 300
 Tyr Ala Ser Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 305 310 315

<210> 20

<211> 230

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 polypeptide

<400> 20

Arg His Arg Gln Pro Arg Gly Trp Glu Gln Leu Pro Thr Gly Ala Glu
 1 5 10 15
 Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser Thr Arg Gly Thr Gln
 20 25 30
 Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His Arg Gln Leu Glu Glu
 35 40 45
 Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr Phe Leu Glu Ala Ala
 50 55 60
 Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg Ser Gln Asp Leu Asp
 65 70 75 80
 Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp Pro Ala Leu Ala Tyr
 85 90 95
 Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg Gly Arg Ile Arg Asn
 100 105 110
 Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser Ser Leu Pro Gly Phe
 115 120 125
 Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu Ala Ala Gly Glu Val
 130 135 140
 Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg Leu Asp Ala Ile Thr
 145 150 155 160

46n

Gly	Pro	Glu	Glu	Glu	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro
				165					170					175	
Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro
			180					185					190		
Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Lys	Glu
		195					200					205			
Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro
	210					215					220				
Pro	Arg	Glu	Asp	Leu	Lys										
225					230										

<210> 21
 <211> 230
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic polypeptide

<400> 21

Arg	His	Arg	Gln	Pro	Arg	Gly	Trp	Glu	Gln	Leu	Pro	Thr	Gly	Ala	Glu
1				5				10						15	
Phe	Leu	Gly	Asp	Gly	Gly	Asp	Val	Ser	Phe	Ser	Thr	Arg	Gly	Thr	Gln
			20					25					30		
Asn	Trp	Thr	Val	Glu	Arg	Leu	Leu	Gln	Ala	His	Arg	Gln	Leu	Glu	Glu
		35				40					45				
Gly	Gly	Tyr	Val	Phe	Val	Gly	Tyr	His	Gly	Thr	Phe	Leu	Glu	Ala	Ala
	50					55					60				
Gln	Ser	Ile	Val	Phe	Gly	Gly	Val	Arg	Ala	Arg	Ser	Gln	Asp	Leu	Asp
65				70				75						80	
Ala	Ile	Trp	Ala	Gly	Phe	Tyr	Ile	Ala	Gly	Asp	Pro	Ala	Leu	Ala	Tyr
			85					90					95		
Gly	Tyr	Ala	Gln	Asp	Gln	Glu	Pro	Asp	Ala	Ala	Gly	Arg	Ile	Arg	Asn
			100					105					110		
Gly	Ala	Leu	Leu	Arg	Val	Tyr	Val	Pro	Arg	Ser	Ser	Leu	Pro	Gly	Phe
	115					120						125			
Tyr	Ala	Thr	Ser	Leu	Thr	Leu	Ala	Ala	Pro	Glu	Ala	Ala	Gly	Glu	Val
	130					135						140			
Glu	Arg	Leu	Ile	Gly	His	Pro	Leu	Pro	Leu	Arg	Leu	Asp	Ala	Ile	Thr
145				150						155					160
Gly	Pro	Glu	Glu	Ala	Gly	Gly	Arg	Leu	Glu	Thr	Ile	Leu	Gly	Trp	Pro
			165					170						175	
Leu	Ala	Glu	Arg	Thr	Val	Val	Ile	Pro	Ser	Ala	Ile	Pro	Thr	Asp	Pro
			180					185					190		
Arg	Asn	Val	Gly	Gly	Asp	Leu	Asp	Pro	Ser	Ser	Ile	Pro	Asp	Ser	Glu
		195					200					205			
Gln	Ala	Ile	Ser	Ala	Leu	Pro	Asp	Tyr	Ala	Ser	Gln	Pro	Gly	Lys	Pro
	210					215					220				
Pro	Arg	Glu	Asp	Leu	Lys										
225					230										

<210> 22
 <211> 347

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 22

```

Pro Glu Gly Gly Ser Leu Ala Ala Leu Thr Ala His Gln Ala Cys His
1          5          10          15
Leu Pro Leu Glu Thr Phe Thr Arg His Arg Gln Pro Arg Gly Trp Glu
          20          25          30
Gln Leu Glu Gln Cys Gly Tyr Pro Val Gln Arg Leu Val Ala Leu Tyr
          35          40          45
Leu Ala Ala Arg Leu Ser Trp Asn Gln Val Asp Gln Val Ile Arg Asn
          50          55          60
Ala Leu Ala Ser Pro Gly Ser Gly Gly Asp Leu Gly Glu Ala Ile Arg
65          70          75          80
Glu Gln Pro Glu Gln Ala Arg Leu Ala Leu Thr Leu Ala Ala Ala Glu
          85          90          95
Ser Glu Arg Phe Val Arg Gln Gly Thr Gly Asn Asp Glu Ala Gly Ala
          100          105          110
Ala Asn Gly Pro Ala Asp Ser Gly Asp Ala Leu Leu Glu Arg Asn Tyr
          115          120          125
Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser
          130          135          140
Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
145          150          155          160
Arg Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
          165          170          175
Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
          180          185          190
Ser Gln Asp Leu Asp Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
          195          200          205
Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
          210          215          220
Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Arg Ser
225          230          235          240
Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu
          245          250          255
Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
          260          265          270
Leu Asp Ala Ile Thr Gly Pro Glu Glu Gly Gly Arg Leu Glu Thr
          275          280          285
Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
          290          295          300
Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
305          310          315          320
Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
          325          330          335
Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
          340          345

```

<210> 23

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 23

Lys Ala Ser Gly Gly

1

5

CLAIMS:

1. A method of producing a purified polypeptide from a solution comprising the polypeptide and a deamidated species of the polypeptide, wherein said deamidated species of the polypeptide results in an inhibition of potency of said polypeptide, the method comprising: (a) contacting the solution with an anion exchange (AIEX) chromatography matrix whereby the polypeptide binds to the AIEX chromatography matrix; and (b) eluting the bound polypeptide from the AIEX chromatography matrix with a linear salt gradient that is from about 150 mM NaCl in Tris/HCl, pH 8.0 to about 300 mM NaCl in Tris/HCl, pH 8.0, thereby separating said polypeptide from the deamidated species of said polypeptide and producing a purified polypeptide; wherein the polypeptide is an immunoconjugate comprising: (1) an antibody or antigen binding fragment thereof that binds the cell surface receptor CD22; and (2) a *Pseudomonas* exotoxin, or variant thereof.
2. The method of claim 1, wherein the AIEX matrix contains quaternary amine and tertiary amine ion exchange groups.
3. The method of claim 2, wherein the AIEX matrix contains a quaternary amino (Q) group.
4. The method of claim 3, wherein the AIEX matrix is Q sepharose™.
5. The method of any one of claims 1-4, wherein the linear salt gradient is from about 175 mM NaCl in Tris/HCl, pH 8.0 to about 275 mM NaCl in Tris/HCl, pH 8.0.
6. The method of any one of claims 1-4, wherein the linear salt gradient is from about 192 mM NaCl in Tris/HCl, pH 8.0 to about 245 mM NaCl in Tris/HCl, pH 8.0.
7. The method of any one of claims 1-6, wherein between about 75% to about 99% of the deamidated species is separated from said polypeptide.
8. The method of claim 7, wherein about 80% of the deamidated species is separated from said polypeptide.

9. The method of claim 7, wherein about 85% of the deamidated species is separated from said polypeptide.
10. The method of claim 7, wherein about 90% of the deamidated species is separated from said polypeptide.
11. The method of claim 7, wherein about 95% of the deamidated species is separated from said polypeptide.
12. The method of claim 7, wherein about 96%, 97%, 98%, or 99% of the deamidated species is separated from said polypeptide.
13. The method of any one of claims 1-12, wherein the antibody or antigen binding fragment comprises a Fab, a Fab', a F(ab')₂, a Fd, a single chain Fv or scFv, a disulfide linked Fv, a V-NAR domain, an IgNar, an intrabody, an IgGΔCH2, a minibody, a F(ab')₃, a tetrabody, a triabody, a diabody, a single-domain antibody, DVD-Ig, Fcab, mAb², a (scFv)₂, or a scFv-Fc.
14. The method of any one of claims 1-13, wherein said *Pseudomonas* exotoxin, or variant thereof has an amino acid sequence selected from the group consisting of SEQ ID NOs: 16-22.
15. The method of claim 14, wherein said *Pseudomonas* exotoxin, or variant thereof has the amino acid sequence of SEQ ID NO: 22.
16. The method of any one of claims 1-15, wherein said antibody or antigen binding fragment thereof comprises a VH and a VL sequence.
17. The method of claim 16, wherein said VH sequence is selected from the group consisting of SEQ ID NOs: 6-11.
18. The method of claim 16, wherein said VL sequence is selected from the group consisting of SEQ ID NOs: 2, and 12-15.

19. The method of any one of claims 1-18, wherein the immunoconjugate is the CAT-8015 immunotoxin comprising a V_H-PE38 subunit of SEQ ID NO: 1 and a V_L subunit of SEQ ID NO: 2.

20. A method of modifying the bioactivity of a polypeptide solution comprising a polypeptide and a deamidated species of the polypeptide, the method comprising separating the polypeptide from the deamidated species by linear elution AIEC chromatography; and combining the purified polypeptide and deamidated species in fixed quantities to obtain the desired bioactivity of the polypeptide solution.

21. The method of any one of claims 1-19, wherein said deamidated species of the polypeptide are present within the *Pseudomonas* exotoxin or variant thereof.

22. The method of claim 19, wherein said deamidated species of the polypeptide is deamidated at position 358 of SEQ ID NO: 1.

FIG. 1
Ion exchange chromatography profile of CAT-8015 reference standard






FIG. 2
Correlation between % Relative Potency and % Pre-Peak







4/6

FIG. 4

SDS-PAGE Analysis of QHP
Load and Eluate Pool Samples





6/6

FIG. 6

SDS-PAGE Analysis of QHP Load and Eluate Pool Samples




FIG. 7

Percent Pre-Peak in HA Product as a Function of Solubilization pH

