

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 January 2007 (18.01.2007)

PCT

(10) International Publication Number
WO 2007/008527 A2(51) International Patent Classification:
C07K 14/195 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2006/026183

(22) International Filing Date: 5 July 2006 (05.07.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/697,642 8 July 2005 (08.07.2005) US
60/801,835 19 May 2006 (19.05.2006) US

(71) Applicant (for all designated States except US): CHILDREN'S HOSPITAL INC. [US/US]; 700 Children's Drive, Columbus, OH 43205 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BAKALETZ, Lauren, O. [US/US]; 4825 Canterbury Court, Hilliard, OH 43026 (US). MUNSON, Robert, S., Jr. [US/US]; 4825 Canterbury Court, Hilliard, OH 43026 (US).

(74) Agent: SINTICH, Sharon, M.; MARSHALL, GERSTEIN & BORUN, 233 S. Wacker Drive, Suite 6300, Sears Tower, Chicago, IL 60606-6357 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2007/008527 A2

(54) Title: CHIMERIC VACCINE FOR HAEMOPHILUS INFLUENZAE-INDUCED DISEASE

(57) Abstract: The invention described herein relates to a chimeric protein comprising the NTHi twitching pilus major subunit protein (PilA) presenting a portion of the NTHi OMP P5 protein. The invention provides for vaccine compositions comprising the recombinant chimeric protein and methods of eliciting an immune response using the recombinant chimeric proteins of the invention.

CHIMERIC VACCINE FOR *HAEMOPHILUS INFLUENZAE*-INDUCED DISEASE

[0001] This application claims priority to U.S. Provisional Application No. 60/697,642 filed July 8, 2005 and U.S. Provisional Application No. 60/801,835 filed 5 May 19, 2006, all of which are incorporated by reference herein in its entirety.

Field of Invention

[0002] The invention described herein relates to a chimeric protein comprising the NTHi twitching pilus major subunit protein (PilA) presenting a portion of the NTHi OMP P5 protein. The invention provides for vaccine compositions comprising the 10 chimeric protein and methods of eliciting an immune response using the chimeric proteins of the invention.

Background

[0003] The clinical term for middle ear infections is otitis media (OM). According to Klein, *Vaccine*, 19 (Suppl. 1): S2-S8, 2000, OM is the most common reason for an 15 ill child to obtain healthcare and for a child in the United States to receive antibiotics or undergo general anesthesia. Statistics indicate that 24.5 million physician office visits were made for OM in 1990, representing a greater than 200% increase over those reported in the 1980s. While rarely associated with mortality, the morbidity 20 associated with OM is significant. Hearing loss is a common problem associated with this disease, often affecting a child's behavior, education and development of language skills (Baldwin, *Am. J. Otol.*, 14: 601-604, 1993; Hunter *et al.*, *Ann. Otol. Rhinol. Laryngol. Suppl.*, 163: 59-61, 1994; Teele *et al.*, *J. Infect. Dis.*, 162: 685-694, 1990). The socioeconomic impact of OM is also great, with direct and indirect costs of diagnosing and managing OM exceeding \$5 billion annually in the U.S. alone 25 (Kaplan *et al.*, *Pediatr. Infect. Dis. J.*, 16: S9-11, 1997).

[0004] OM is thought to result from infectious, environmental and host genetics factors. Bacteria such as *Haemophilus influenzae*, *Streptococcus pneumoniae* and *Moraxella catarrhalis* are the most common infectious organisms in OM. Acute OM 30 is a disease characterized by rapid onset and short duration of signs and symptoms of inflammation in the middle ear, while chronic OM refers to a condition that is defined by the relatively asymptomatic presence of fluid (or effusion) in the middle ear. However, in chronic OM, despite the absence of certain signs of acute infection (*i.e.*,

ear pain or fever), these abnormal middle ear fluids can persist for periods exceeding three months. Treatment of acute OM by antibiotic therapy is common, but multiple antibiotic-resistant bacteria have emerged in all three Genera of bacteria responsible for OM. Surgical management of chronic OM involves the insertion of

5 tympanostomy tubes through the tympanic membrane of the ear while a child is under general anesthesia. While this procedure is commonplace (prevalence rates are ~ 1 million tubes inserted per year in the U.S. Bright *et al.*, *Am. J. Public Health*, 83(7): 1026-8, 1993) and is highly effective in terms of relieving painful symptoms by draining the middle ear of accumulated fluids, it is invasive and carries incumbent 10 risks (Berman *et al.*, *Pediatrics*, 93(3):353-63, 1994; Bright *et al.*, *supra*; Cimons, *ASM News*, 60: 527-528; Paap, *Ann. Pharmacother.*, 30(11): 1291-7, 1996). There is thus a need for additional approaches to the management and, preferably, the prevention of OM.

15 [0005] OM vaccine development is most advanced for *S. pneumoniae*, the primary causative agent of acute OM (AOM), as evidenced by the recent approval and release of a seven-valent capsular-conjugate vaccine, PREVNAR® (Eskola and Kilpi, *Pediatr. Infect. Dis. J.* 16: S72-78, 2000). While PREVNAR® has been highly efficacious for invasive pneumococcal disease, coverage for OM has been disappointing (6-8%) with reports of an increased number of OM cases due to 20 serotypes not included in the vaccine (Black *et al.*, *Pediatr. Infect. Dis J.*, 19: 187-195, 2000; Eskola *et al.*, *Pediatr. Infect. Dis J.*, 19: S72-78, 2000; Eskola *et al.*, *N. Engl. J. Med.*, 344: 403-409, 2001; Snow *et al.*, *Otol. Neurotol.*, 23: 1-2, 2002).

25 [0006] *H. influenzae* is a gram-negative bacterium that, as noted above, plays a role in OM. Clinical isolates of *H. influenzae* are classified either as serotypes "a" through "f" or as non-typeable depending on the presence or absence, respectively, of type-specific polysaccharide capsules on the bacteria. A vaccine for *H. influenzae* type b has been developed. Like PREVNAR®, the type b *H. influenzae* vaccines target the polysaccharide capsule of this organism and thus the vaccine is comprised of capsule polysaccharide that has been conjugated to a protein carrier. Neither PREVNAR® or 30 the type b *H. influenzae* vaccine have any efficacy for NTHI-induced respiratory tract diseases, including OM. Less progress has been made for a vaccine for non-typeable *H. influenzae* (NTHi) which causes approximately 20% of acute OM in children and predominates in chronic OM with effusion (Coleman *et al.*, *Inf and Immunity*, 59(5),

1716-1722, 1991; Klein, *Pediatr. Infect. Dis. J.*, 16, S5-8, 1997; Spinola *et al.*, *J. Infect. Dis.*, 154, 100-109, 1986). NTHi can also cause pneumonia, sinusitis, septicemia, endocarditis, epiglottitis, septic arthritis, meningitis, postpartum and neonatal infections, postpartum and neonatal sepsis, acute and chronic salpingitis, 5 pericarditis, cellulitis, osteomyelitis, endocarditis, cholecystitis, intraabdominal infections, urinary tract infection, mastoiditis, aortic graft infection, conjunctivitis, Brazilian purpuric fever, occult bacteremia and exacerbation of underlying lung diseases such as chronic bronchitis, bronchiectasis and cystic fibrosis. A prototype NTHi isolate is the low passage isolate 86-028NP which was recovered from a child 10 with chronic OM. This strain has been well characterized *in vitro* (Bakaletz *et al.*, *Infect. Immun.*, 53: 331-5, 1988; Holmes *et al.*, *Microb. Pathog.*, 23: 157-66, 1997) as well as in chinchilla OM models (Bakaletz *et al.*, *Vaccine*, 15: 955-61, 1997; Suzuki *et al.*, *Infect. Immun.*, 62: 1710-8, 1994; DeMaria *et al.*, *Infect. Immun.*, 64: 5187-92, 1996). The NTHi strain 86-026NP was deposited with the American Type 15 Culture Collection, 10801 University Blvd., Manassas, VA 20110, on October 16, 2001 and assigned accession no. PTA-4764. A contig set from the genome of stain 86-028NP can be found at Columbus Children's Research Institute Center for Microbial Pathogenesis web site.

[0007] Adherence and colonization are acknowledged first steps in the pathogenesis 20 of *H. influenzae*-induced diseases. As such, *H. influenzae* express multiple adhesins including hemagglutinating pili, fimbriae and non-fimbrial adhesins (Gilsdorf *et al.*, *Pediatr Res* 39, 343-348, 1996; Gilsdorf., *Infect. Immun.*, 65, 2997-3002, 1997; and St. Geme III, *Cell. Microbiol.*, 4, 191-200, 2002). Notably, none of the adhesins described have previously been associated with a motility function. Moreover, *H. influenzae* do not express flagella which are also associated with motility. Twitching 25 motility is a flagella-independent form of bacterial translocation over moist surfaces and occurs by extension, tethering, and then retraction of polar structures known as type IV pili (Bardy ., *Microbiology*, 149, 295-304, 2003; Tonjum and Koomey, *Gene*, 192, 155-163, 1997; Wolfgang *et al.*, *EMBO J.*, 19, 6408-6418, ; Mattick, *Annu. Rev. Microbiol.*, 56, 289-314, 2002). Type IV pili are typically 5-7 nm in diameter, several 30 micrometers in length and comprised of a single protein subunit assembled into a helical conformation with ~5 subunits per turn (Bardy *et al.*, *Microbiology*, 149, 295-304, 2003; Wall and Kaiser, *Mol. Microbiol.*, 32, 1-10, 1999). Type IV pilin subunits

are usually 145-160 amino acids in length and may be glycosylated or phosphorylated. There are two classes of pilin subunits, type IVa and type IVb, which are distinguished from one another by the average length of the leader peptide and the mature subunit, which N-methylated amino acid occupies the N-terminal

5 position of the mature protein, and the average length of the D-region (for disulfide region). Most of the respiratory pathogens express class IVa pilins, whereas the enteropathogens more typically express class IVb pilins. Type IVa pili are distinguished by the presence of a highly conserved, hydrophobic N-terminal methylated phenylalanine.

10 [0008] Type IV pili serve as a means of rapid translocation over and colonization of new surfaces. Thus type IV pilus expression is important to both adherence and biofilm formation by many bacteria (Mattick, *Annu. Rev. Microbiol.*, 56, 289-314 2002; O'Toole and Kolter, *Mol. Microbiol.*, 30, 295-304, 1998; Klausen *et al.*, *Mol. Microbiol.*, 50, 61-68, 2003; Jesaitis *et al.*, *J. Immunol.*, 171, 4329-4339, 2003), as
15 well as virulence of *Neisseria* species, *Moraxella bovis*, *Vibrio cholerae*, enteropathogenic *Escherichia coli* and *Pseudomonas aeruginosa*, among others (O'Toole and Kolter, *supra*; Klausen *et al.*, *supra*; Klausen *et al.*, *Mol. Microbiol.*, 48, 1511-1524, 2003; Strom and Lory, *Annu. Rev. Microbiol.*, 47, 565-596, 1993). A biofilm is a complex organization of bacteria that are anchored to a surface via a
20 bacterially extruded matrix, comprised of exopolysaccharide or other substances. The matrix envelopes the bacteria and protects it from the human immune system. Ehrlich *et al.*, *JAMA*, 287(13), 1710-1715 (2002) describes biofilm formation by *H. influenzae*. It has been postulated that blocking the interaction between type IV pili and the human body can avoid or stop the bacterial infection (Meyer *et al.*, U.S.
25 Patent No. 6,268,171 issued July 31, 2001).

[0009] Type IV pilus expression is a complex and highly regulated bacterial function. In *P. aeruginosa*, the biogenesis and function of type IV pili is controlled by over forty genes (Strom and Lory, *supra*). To date, only a subset of the vast number of related type IV pilus genes (Tonjum and Koomey, *supra*; Darzins and
30 Russell, *Gene*, 192, 109-115, 1997) have been found in several members of the HAP (*Haemophilus*, *Actinobacillus* and *Pasteurella*) family (Stevenson *et al.*, *Vet. Microbiol.*, 92, 121-134, 2003; Doughty *et al.*, *Vet. Microbiol.*, 72, 79-90, 2000; Dougherty and Smith, *Microbiology*, 145, 401-409 1999), but neither expression of

type IV pili nor twitching motility has ever been described for any *H. influenzae* isolate. In fact, *H. influenzae* is classically described as a bacterium that does not express these structures (Friedrich *et al.* *Appl. Environ. Microbiol.*, 69, 3695-3700, 2003; Fussenegger *et al.*, *Gene*, 192, 125-134, 1997), despite the presence of a 5 cryptic gene cluster within the strain Rd genome (Fleischmann *et al.*, *Science*, 269, 496-512, 1995). Strain Rd is a non-encapsulated derivative of an *H. influenzae* serotype d organism (Zwahlen *et al.*, *Infect. Immun.*, 42, 708-715, 1983; Bendler and Goodgal, *J. Microbiol.*, 70, 411-422, 1972; Risberg *et al.*, *Eur. J. Biochem.*, 261, 171-180, 1999). Although strain Rd has some virulence properties, serotype d strains are 10 generally considered to be commensals; they do not frequently cause disease (Daines *et al.*, *J. Med. Microbiol.*, 52, 277-282, 2003). It is therefore important to make the distinction between disease-causing strains of *H. influenzae* and strain Rd.

[0010] Fimbriae, which are surface appendages found on non-typable *Haemophilus influenzae*, are produced by 100% of the bacteria recovered from the middle ears and 15 nasopharyngeal region of children with chronic otitis media. A vaccine comprised of fimbriae, a filamentous protein derived from the fimbriae of non-typable *Haemophilus influenzae* was previously developed and is useful in studying, preventing, or reducing the severity of otitis media. However, existing methodologies to isolate fimbriae protein from the bacterial outer membrane are tedious and time-consuming. 20 Similarly, purification of fimbriae expressed by the fimbriae gene in other host vector, is also tedious due to the homology between the fimbriae protein and the outer membrane proteins of the host vector.

[0011] The synthetic chimeric vaccine candidate, denoted as LB1 and described in U.S. Patent No. 5,843,464, has shown tremendous efficacy in multiple pre-clinical 25 vaccine trials in two rodent hosts. This synthetic peptide comprises a B-cell epitope of P5-fimbriae collinearly synthesized with a T-cell promiscuous epitope derived from a fusion protein of the measles vaccine. Whereas LB1 peptide has been shown to be efficacious in pre-clinical trials, there is concern about the ability to test and market a vaccine that contains a T-cell promiscuous epitope for intended use in very young 30 children. Therefore, there is a need to develop vaccine candidate that elicit a specific and controlled immune response to *H. influenzae*.

Summary of the Invention

[0012] The present invention relates to chimeric proteins comprising a portion of the Type IV pilus major subunit protein (PilA) of nontypeable *H. influenzae* (NTHi) and a portion of NTHi OMP P5 protein (also called P5-fimbrin, fimbrin or OMP P5-

5 homologous adhesin). In particular, the invention provides for chimeric proteins comprising PilA modified to present the B-cell epitope of the LB1 peptide. The invention also provides vaccine compositions comprising one or more chimeric proteins of the invention and methods of eliciting an immune response using the chimeric proteins of the invention.

10 [0013] The LB1 peptide is a 40 amino acid synthetic chimeric P5-fimbrin derived peptide (SEQ ID NO: 53) that induces an immunogenic response to NTHi and is advantageous because it does not require tedious purification techniques. The LB1 peptide comprises an N-terminal 19 amino acid peptide that is a B-cell epitope (SEQ ID NO: 4). The B-cell epitope was derived from the predicted surface-exposed loop 3
15 of an outer membrane protein (fimbrin) of NTHi denoted as OMP P5 (also called P5-fimbrin or OMP P5-homologous adhesin). The LB1 peptide further comprises a short 5-mer linker peptide and a 16-residue T cell promiscuous epitope. The T cell epitope was derived from a fusion protein of the measles virus. The T cell promiscuous epitope induces a very strong T cell response in individuals exposed to this epitope.

20 [0014] The present invention contemplates inserting a portion or fragment of the LB1 peptide into a safer and selective carrier protein that does not reduce the effectiveness of inducing a B-cell response. Preferably, the portion of the LB1 peptide is inserted into a carrier that itself also confers protection against NTHi-induced diseases. One such carrier that may induce protection against NTHi induced
25 diseases is the protein that comprises the NTHi Type IV pilus (twitching pilus) protein, also known as PilA protein (SEQ ID NO: 2). The PilA protein is encoded by the *pilA* gene (SEQ ID NO: 1).

[0015] The present invention provides for chimeric proteins comprising a portion of the LB1 peptide in order to present the peptide to induce an immunogenic
30 response. The invention contemplates presenting a portion of the LB1 peptide that is 12 to 35 amino acids, more preferably presenting a portion of the LB1 peptide that is 15 to 30 amino acids, and most preferably presenting a portion of the LB1 peptide that

is 18 to 19 amino acids and is a subunit of the fimbrin protein. A preferred portion of the LB1 peptide is the N-terminal amino acid sequence RSDYKFYEDANGTRDHKKKG (SEQ ID NO: 4).

[0016] In another embodiment, the invention provides for chimeric protein wherein 5 the PilA protein is modified to present a 24 amino acid peptide. The 24 amino acid peptide may comprise the B-cell epitope of the LB1 peptide modified as set out in the amino acid sequence of SEQ ID NO: 5 (LVRSDYKFYEDANGTRDHKKGRHT) in which a leucine and valine are added to the N terminus of the B-cell epitope of LB1 and an arginine, histidine and threonine are at the C terminus of the B-cell epitope of 10 LB1. These modifications to the B-cell epitope are contemplated to assist in protein folding and/or antigen presentation. The invention further contemplates any modifications to the B-cell epitope of LB1 that will assist in protein folding and/or antigen presentation.

[0017] The amino acid sequence of the surface exposed loop 3 of NTHi OMP P5 15 can vary between NTHi strains. The invention contemplates chimeric proteins comprising a portion of the PilA protein modified to present the B cell epitope of any variant amino acid sequence of loop 3 of the NTHi OMP P5. In particular, the invention provides for chimeric proteins wherein the PilA protein is modified to present one of the following variant NTHi OMP P5 amino acids sequences: 20 RSDYKLYNKNSSNSTLKNLGE (SEQ ID NO: 6), RSDYKLYNKNSSTLKDLGE (SEQ ID NO: 7) and RSDYKFYDNKRID (SEQ ID NO: 8). The variant peptides also may be presented with a leucine and valine added to the N terminus and an arginine, histidine and threonine added to the C terminus or any other modification to assist in protein folding and/or antigen presentation.

[0018] The chimeric proteins of the invention comprise the modified PilA amino 25 acids wherein the native PilA amino acids have been substituted with a portion of the LB1 peptide. In addition, the chimeric proteins of the invention comprise a modified PilA amino acid sequence wherein a portion of the LB1 peptide is inserted within and in addition to the native PilA amino acids. The chimeric proteins of the invention 30 have the ability to induce the formation of antibodies directed against two proteins and therefore are more effective and more specific vaccine candidates.

[0019] In one embodiment, the chimeric proteins comprise the mature amino acid sequence (residues 13-149) of the NTHi PilA protein (SEQ ID NO: 2) wherein a portion of the LB1 peptide is inserted between the cysteine residues at positions 62 and 72 of SEQ ID NO: 2 and may substitute the native amino acids, such as the 5 chimeric protein having the amino acid sequence of SEQ ID NO: 54. This chimeric protein comprises residues 40-149 of SEQ ID NO: 2 and has the B-cell epitope of LB1 (SEQ ID NO: 5) inserted between residues 62 and 72 of SEQ ID NO: 2. In another embodiment, the portion of the LB1 peptide is inserted between the cysteine residues at positions 131 and 144 of SEQ ID NO: 2 and may substitute the native 10 amino acids such as the protein having the amino acid sequence of SEQ ID NO: 55. This chimeric protein comprises residues 40-149 of SEQ ID NO: 2 and has the B-cell epitope of LB1 (SEQ ID NO: 5) inserted between residues 131 and 144 of SEQ ID NO: 2.

[0020] In another embodiment, the chimeric proteins comprise the mature amino 15 acid sequence (residues 13-149) of the NTHi PilA protein (SEQ ID NO: 2) wherein the portion of the LB1 peptide is inserted at the C-terminus of the PilA protein. For example, the chimeric protein of SEQ ID NO: 56 comprises residues 40-149 of SEQ ID NO: 2 and the B-cell epitope of LB1 (SEQ ID NO: 5) is inserted following residue 149 of SEQ ID NO: 2.

[0021] In another embodiment, the chimeric proteins comprise the mature amino 20 acid sequence (residues 13-149) of NTHi PilA protein (SEQ ID NO: 2) wherein the portion of the LB1 peptide is inserted at the N-terminus of the PilA protein. For example, the chimeric protein of SEQ ID NO: 57 comprises residues 40-149 of SEQ ID NO: 2 and the B-cell epitope of LB1 (SEQ ID NO: 5) is inserted before residue 40 25 of SEQ ID NO: 2.

[0022] In a further embodiment, the invention provides for chimeric proteins comprising a portion of the NTHi PilA protein and one or more of the LB1 peptides described herein. The chimeric proteins of the invention include those which present the same LB1 peptide more than once within a portion of the NTHi PilA protein and 30 those which present two or more different LB1 peptides within a portion of the NTHi PilA protein.

[0023] The invention further provides for chimeric proteins comprising a portion of the NTHi PilA protein and any antigenic protein that will elicit an immune response.

The NTHi Type IV Pilus (PilA) Polynucleotides and Polypeptides of the Invention

[0024] The chimeric proteins of the invention may comprise the full length or a portion of the major subunit of the NTHi Type IV Pilus which is encoded by the gene *pilA*. The PilA protein of the NTHi isolate 86-028NP is encoded by the nucleic acid sequence set out as SEQ ID NO: 2, which is described in U.S. patent application No. 11/019,005, incorporated by reference herein in its entirety. Also provided are polynucleotides encoding PilA polypeptides from NTHi clinical isolates 1728MEE, 10 1729MEE, 3224A, 10548MEE, 1060MEE, 1885MEE, 1714MEE, 1236MEE, 1128MEE and 214NP. The amino acid sequences of these PilA polypeptides are set out in SEQ ID NOS: 34, 36, 38, 40, 42, 44, 46, 48, 50 and 52 respectively. The possibility of alternative codon usage is specifically contemplated in polynucleotides encoding the polypeptides. In one embodiment, the polypeptides are respectively 15 encoded by the nucleotide sequences set out in SEQ ID NOS: 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51.

[0025] The invention provides for polynucleotides that hybridize under stringent conditions to (a) the complement of the nucleotide sequences set out in SEQ ID NOS: 1, 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51; (b) a polynucleotide which is an allelic 20 variant of any polynucleotides recited above; (c) a polynucleotide which encodes a species homolog of any of the proteins recited above; or (d) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptides of the present invention. PilA polynucleotides from other non-typeable *H. influenzae* strains and from *H. influenzae* strains a, b, c, e and f are specifically contemplated. 25 These polynucleotides can be identified and isolated by techniques standard in the art such as hybridization and polymerase chain reaction using part or all of the polynucleotides of SEQ ID NOS: 1, 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51 as probes or primers, respectively.

[0026] The polynucleotides of the invention also include nucleotide sequences that 30 are substantially equivalent to the polynucleotides recited above. Polynucleotides according to the invention can have, e.g., at least 65%, at least 70%, at least 75%, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89%, more typically at

least 90%, 91%, 92%, 93%, or 94% and even more typically at least 95%, 96%, 97%, 98% or 99% sequence identity to the NTHi polynucleotides recited above.

[0027] Included within the scope of the nucleic acid sequences of the invention are nucleic acid sequence fragments that hybridize under stringent conditions to the NTHi

5 nucleotide sequences of SEQ ID NOS: 1, 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51, or complements thereof, which fragment is greater than about 5 nucleotides, preferably 7 nucleotides, more preferably greater than 9 nucleotides and most preferably greater than 17 nucleotides. Fragments of, *e.g.*, 15, 17, or 20 nucleotides or more that are selective for (*i.e.*, specifically hybridize to any one of the PilA polynucleotides of the 10 invention) are contemplated. These nucleic acid sequence fragments capable of specifically hybridizing to an NTHi PilA polynucleotide of the invention can be used as probes to detect NTHi PilA polynucleotides of the invention and/or can differentiate NTHi PilA polynucleotides of the invention from other bacterial genes, and are preferably based on unique nucleotide sequences.

15 [0028] The term “stringent” is used herein to refer to conditions that are commonly understood in the art as stringent. Hybridization stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of stringent conditions for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium chloride,

20 0.0015M sodium citrate, and 50% formamide at 42°C. See Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, (Cold Spring Harbor, N.Y. 1989).

[0029] More stringent conditions (such as higher temperature, lower ionic strength, 25 higher formamide, or other denaturing agent) may also be used, however, the rate of hybridization will be affected. In instances wherein hybridization of

deoxyoligonucleotides is concerned, additional exemplary stringent hybridization conditions include washing in 6X SSC 0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos).

30 [0030] Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium

pyrophosphate, 0.1% sodium dodecylsulfate, NaDODSO₄, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or other non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the 5 stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4, however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. *See Anderson et al., Nucleic Acid Hybridisation: A Practical Approach*, Ch. 4, IRL Press Limited (Oxford, England). Hybridization conditions can be adjusted by one skilled in the art in order to 10 accommodate these variables and allow DNAs of different sequence relatedness to form hybrids.

[0031] As noted above, polynucleotides contemplated by the present invention are not limited to the specific PilA polynucleotides of SEQ ID NOS: 1, 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51, but also include, for example, allelic and species variations 15 thereof. Allelic and species variations can be routinely determined by comparing the sequence provided in SEQ ID NOS: 1, 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51, preferably the open reading frames therein, a representative fragment thereof, or a nucleotide sequence at least 90% identical, preferably 95% identical, to the open reading frames within SEQ ID NOS: 1, 33, 35, 37, 39, 41, 43, 45, 47, 49 and 51 with a 20 sequence from another isolate of the same species or another species. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux *et al.*, *Nucl. Acid. Res.*, 12: 387, 1984; Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Altschul 25 *et al.*, *J. Mol. Biol.*, 215: 403-410, 1990). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (*BLAST Manual*, Altschul *et al.* NCB/NLM/NIH Bethesda, MD 20894; Altschul *et al.*, *supra*). The well known Smith-Waterman algorithm may also be used to determine identity.

30 [0032] Polynucleotides of the invention may be isolated from natural sources or may be synthesized by standard chemical techniques, *e.g.*, the phosphotriester method described in Matteucci *et al.*, *J Am Chem Soc.*, 103: 3185 (1981).

[0033] The invention provides for chimeric proteins comprising a portion of NTHi PilA protein. In one embodiment the polypeptides comprise the NTHi 86-028NP amino acid sequences respectively set out in SEQ ID NO: 2. Polypeptides of the invention also include PilA polypeptides set out in SEQ ID NOS: 34, 36, 38, 40, 42, 5 44, 46, 48, 50 and 52. In additional embodiments, the PilA polypeptides of the invention are those of other non-typeable *H. influenzae* strains and from *H. influenzae* strains a, b, c, e and f.

[0034] Polypeptides of the invention specifically include peptide fragments (*i.e.*, peptides) or fragments of the PilA polypeptide that retain one or more biological or 10 immunogenic properties of a full length polypeptide of the invention. In one embodiment, PilA peptide fragments provided by the invention are designated TfpQ2, TfpQ3, TfpQ4 and OLP3 and respectively comprise amino acids 35 through 68 of SEQ ID NO: 2, amino acids 69 through 102 of SEQ ID NO: 2, amino acids 103 through 137 of SEQ ID NO: 2, and amino acids 21 through 35 of SEQ ID NO: 2. 15 Another PilA peptide fragment provided by the invention comprises amino acids 40 through 149 of SEQ ID NO: 2.

[0035] The invention also provides for chimeric proteins comprising a portion of a PilA polypeptide with one or more conservative amino acid substitutions that do not affect the biological and/or immunogenic activity of the PilA polypeptide. 20 Alternatively, the PilA polypeptides of the invention are contemplated to have conservative amino acids substitutions which may or may not alter biological activity. The term “conservative amino acid substitution” refers to a substitution of a native amino acid residue with a nonnative residue, including naturally occurring and nonnaturally occurring amino acids, such that there is little or no effect on the polarity 25 or charge of the amino acid residue at that position. For example, a conservative substitution results from the replacement of a non-polar residue in a polypeptide with any other non-polar residue. Further, any native residue in the polypeptide may also be substituted with alanine, according to the methods of “alanine scanning mutagenesis”. Naturally occurring amino acids are characterized based on their side 30 chains as follows: basic: arginine, lysine, histidine; acidic: glutamic acid, aspartic acid; uncharged polar: glutamine, asparagine, serine, threonine, tyrosine; and non-polar: phenylalanine, tryptophan, cysteine, glycine, alanine, valine, proline,

methionine, leucine, norleucine, isoleucine General rules for amino acid substitutions are set forth in Table 1 below.

Table 1
Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asn
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe,	Leu
Leu	Norleucine, Ile, Val, Met,	Leu
Lys	Arg, 1,4 Diaminobutyric	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Arg
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala,	Leu

5

[0036] The invention also provides for chimeric proteins comprising a portion of a variants of the NTHi PilA polypeptides of the present invention (e.g., a polypeptide exhibiting at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, 86%, 87%, 88%, 89%, at least about 90%, 91%, 92%, 93%, 10 94%, typically at least about 95%, 96%, 97%, more typically at least about 98%, or most typically at least about 99% amino acid identity to a polypeptide of SEQ ID NOS: 2, 34, 36, 38, 40, 42, 44, 46, 48, 50 and 52) that retain biological and/or immunogenic activity.

[0037] The invention contemplates that PilA polynucleotides of the invention may be inserted in a vector for amplification or expression. For expression, the polynucleotides are operatively linked to appropriate expression control sequences such as promoter and polyadenylation signal sequences. Further provided are host 5 cells comprising polynucleotides of the invention. Exemplary prokaryotic host cells include bacteria such as *E. coli*, *Bacillus*, *Streptomyces*, *Pseudomonas*, *Salmonella* and *Serratia*. Methods of producing polypeptides of the invention by growing the host cells and isolating polypeptide from the host cells or growth medium are specifically contemplated. Alternatively, polypeptides of the invention can be 10 prepared by chemical synthesis using standard means. Particularly convenient are solid phase techniques (see, e.g., Erikson *et al.*, *The Proteins* (1976) v. 2, Academic Press, New York, p. 255). Automated solid phase synthesizers are commercially available. In addition, modifications in the sequence are easily made by substitution, addition or omission of appropriate residues. For example, a cysteine residue may be 15 added at the carboxy terminus to provide a sulphydryl group for convenient linkage to a carrier protein, or spacer elements, such as an additional glycine residue, may be incorporated into the sequence between the linking amino acid at the C-terminus and the remainder of the peptide.

[0038] The term "isolated" refers to a substance removed from, and essentially free 20 of, the other components of the environment in which it naturally exists. For example, a polypeptide is separated from other cellular proteins or a DNA is separated from other DNA flanking it in a genome in which it naturally occurs.

[0039] Recombinant PilA protein (rPilA) may be generated to serve as a more readily renewable product. To do this, the published protocol of Keizer *et al.* (*J. 25 Biol. Chem.*, 276: 24186-14193, 2001), who studied a pilin which also had four Cys residues as it will be critical that rPilA similarly be properly folded so as to possess functional qualities of the native pilin subunit, is utilized. Briefly, a truncated pilin is engineered wherein the first 28 residues are removed from the N-terminus to prevent aggregation, and this truncated pilin will be further engineered to be transported to the 30 periplasm by means of the incorporation of an OmpA leader sequence in the construct. Using this strategy Keizer *et al.* generated a recombinant soluble monomeric *P. aeruginosa* pilin protein that was able to bind to its receptor (asialo GM1) in *in vitro* assays and decrease morbidity and mortality in mice when the

peptide was delivered 15 minutes prior to heterologous challenge. This soluble, monomeric, truncated form of NTHi PilA will be useful in the studies described herein.

[0040] The invention also provides for synthetic chimeric proteins. The chimeric 5 proteins may be synthesized, purified and sequenced using standard techniques. For example, the chimeric proteins may be assembled semi-manually by stepwise Fmoc-tert-butyl solid-phase synthesis and purified by HPLC. The composition and amino acid sequence of recombinant and synthetic chimeric proteins may be confirmed by amino acid analysis and/or mass spectral analysis.

10 Antibodies

[0041] The invention provides antibodies which bind to antigenic epitopes of the chimeric proteins of the invention. The antibodies may be polyclonal antibodies, monoclonal antibodies, antibody fragments which retain their ability to bind their unique epitope (e.g., Fv, Fab and F(ab)2 fragments), single chain antibodies and 15 human or humanized antibodies. Antibodies may be generated by techniques standard in the art using chimeric protein(s) of the invention or host cells expressing chimeric protein(s) of the invention as antigens.

[0042] The present invention provides for antibodies specific for the chimeric proteins of the present invention and fragments thereof, which exhibit the ability to 20 kill both *H. influenzae* bacteria and to protect humans from infection. The present invention also provides for antibodies specific for the chimeric proteins of the invention which reduce the virulence, inhibit adherence, inhibit biofilm formation, inhibit twitching motility, inhibit cell division, and/or inhibit penetration into the epithelium of *H. influenzae* bacteria and/or enhance phagocytosis of the *H. influenzae* 25 bacteria.

[0043] *In vitro* complement mediated bactericidal assay systems (Musher *et al.*, *Infect. Immun.* 39: 297-304, 1983; Anderson *et al.*, *J. Clin. Invest.* 51: 31-38, 1972) may be used to measure the bactericidal activity of anti-chimeric proteins antibodies.

[0044] It is also possible to confer short-term protection to a host by passive 30 immunotherapy via the administration of pre-formed antibody against a chimeric protein of the invention. Thus, antibodies of the invention may be used in passive immunotherapy. Human immunoglobulin is preferred in human medicine because a

heterologous immunoglobulin may provoke an immune response to its foreign immunogenic components. Such passive immunization could be used on an emergency basis for immediate protection of unimmunized individuals subject to special risks.

5 [0045] In another embodiment, antibodies of the invention may be used in the production of anti-idiotypic antibody, which in turn can be used as an antigen to stimulate an immune response against the chimeric protein epitopes or *H. influenzae* epitopes.

Methods for Eliciting an Immune Response and Compositions Therefor

10 [0046] The invention contemplates methods of eliciting in an individual an immune response to *H. influenzae* in an individual. In certain embodiments, the methods elicit an immune response to the chimeric proteins of the invention. These methods elicit one or more immune responses, including but not limited to, immune responses which inhibit bacterial replication, immune responses which block *H. influenzae* adherence to cells, immune responses which prevent *H. influenzae* twitching, immune responses that kill *H. influenzae* bacteria and immune responses which prevent biofilm formation. In one embodiment, the methods comprise a step of administering an immunogenic dose of a composition comprising one or more chimeric proteins of the invention. In another embodiment, the methods comprise administering an immunogenic dose of a composition comprising a cell expressing one or more chimeric proteins of the invention. In yet another embodiment, the methods comprise administering an immunogenic dose of a composition comprising one or more polynucleotides encoding one or more chimeric proteins of the invention. The polynucleotide may be a naked polynucleotide not associated with any other nucleic acid or may be in a vector such as a plasmid or viral vector (e.g., adeno-associated virus vector or adenovirus vector). The methods may be used in combination in a single individual. The methods may be used prior or subsequent to *H. influenzae* infection of an individual. The methods and compositions of the invention may be used to treat or prevent any pathological condition involving *H. influenzae* (typeable and nontypeable strains) such as OM, pneumonia, sinusitis, septicemia, endocarditis, epiglottitis, septic arthritis, meningitis, postpartum and neonatal infections, postpartum and neonatal sepsis, acute and chronic salpingitis, epiglottitis, pericarditis, cellulitis, osteomyelitis, endocarditis, cholecystitis, intraabdominal infections, urinary

tract infection, mastoiditis, aortic graft infection, conjunctivitis, Brazilian purpuric fever, occult bacteremia, chronic obstructive pulmonary disease and exacerbation of underlying lung diseases such as chronic bronchitis, bronchiectasis and cystic fibrosis.

[0047] In one embodiment of methods of the invention, a composition of the invention is administered as a priming dose followed by one or more booster doses. Co-administration of proteins or polypeptides that beneficially enhance the immune response such as cytokines (e.g., IL-2, IL-12, GM-CSF), cytokine-inducing molecules (e.g. Leaf) or co-stimulatory molecules is also contemplated.

[0048] An "immunogenic dose" of a composition of the invention is one that generates, after administration, a detectable humoral (antibody) and/or cellular (T cell) immune response in comparison to the immune response detectable before administration or in comparison to a standard immune response before administration. The invention contemplates that the immune response resulting from the methods may be protective and/or therapeutic. In a preferred embodiment, the antibody and/or

T cell immune response protects the individual from *H. influenzae* infection, particularly infection of the middle ear and/or the nasopharynx or lower airway. In this use, the precise dose depends on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc., but generally ranges from about 1.0 μ g to about 5000 μ g per 70 kilogram patient, more commonly from about 10 to about 500 μ g per 70 kg of body weight.

[0049] Humoral immune response may be measured by many well known methods, such as Single Radial Immunodiffusion Assay (SRID), Enzyme Immunoassay (EIA) and Hemagglutination Inhibition Assay (HAI). In particular, SRID utilizes a layer of a gel, such as agarose, containing the immunogen being tested. A well is cut in the gel and the serum being tested is placed in the well. Diffusion of the antibody out into the gel leads to the formation of a precipitation ring whose area is proportional to the concentration of the antibody in the serum being tested. EIA, also known as ELISA (Enzyme Linked Immunoassay), is used to determine total antibodies in the sample.

The immunogen is adsorbed to the surface of a microtiter plate. The test serum is exposed to the plate followed by an enzyme linked immunoglobulin, such as IgG. The enzyme activity adherent to the plate is quantified by any convenient means such as spectrophotometry and is proportional to the concentration of antibody directed against the immunogen present in the test sample. HAI utilizes the capability of an

immunogen such as viral proteins to agglutinate chicken red blood cells (or the like). The assay detects neutralizing antibodies, *i.e.*, those antibodies able to inhibit hemagglutination. Dilutions of the test serum are incubated with a standard concentration of immunogen, followed by the addition of the red blood cells. The 5 presence of neutralizing antibodies will inhibit the agglutination of the red blood cells by the immunogen. Tests to measure cellular immune response include determination of delayed-type hypersensitivity or measuring the proliferative response of lymphocytes to target immunogen.

10 [0050] The invention correspondingly provides compositions suitable for eliciting an immune response to chimeric proteins of the invention. As noted above, the compositions comprise one or more chimeric proteins, cells expressing one or more chimeric proteins, or one or more polynucleotides encoding one or more chimeric proteins. The compositions may also comprise other ingredients such as carriers and adjuvants.

15 [0051] In compositions of the invention, a chimeric protein may be fused to another protein when produced by recombinant methods. In one embodiment, the other protein may not, by itself, elicit antibodies, but it stabilizes the first protein and forms a fusion protein retaining immunogenic activity. In another embodiment, the fusion protein comprises another protein that is immunogenic, such as Glutathione-S- 20 transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the fusion protein and facilitate production and purification thereof. The other protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The other protein may be fused to either the amino or carboxy terminus of the chimeric proteins of the invention.

25 [0052] In other compositions of the invention, chimeric proteins may be otherwise linked to carrier substances. Any method of creating such linkages known in the art may be used. Linkages can be formed with hetero-bifunctional agents that generate a disulfide link at one functional group end and a peptide link at the other, such as a disulfide amide forming agent, *e.g.*, N-succidimidyl-3-(2-pyridyldithio) propionate (SPDP) (See, *e.g.*, Jansen *et al.*, *Immun. Rev.* 62:185, 1982) and bifunctional coupling agents that form a thioether rather than a disulfide linkage such as reactive esters of 6-maleimidocaproic acid, 2-bromoacetic acid, 2-iodoacetic acid, 4-(N-maleimido-methyl) cyclohexane-1-carboxylic acid and the like, and coupling agent which

activate carboxyl groups by combining them with succinimide or 1-hydroxy-2-nitro-4-sulfonic acid, for sodium salt such as succinimidyl 4-(N-maleimido-methyl) cyclohexane-1-carboxylate (SMCC).

[0053] The chimeric proteins may be formulated as neutral or salt forms.

5 Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the peptide) and which are formed with inorganic acids such as, *e.g.*, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, *e.g.*, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, and procaine.

10

[0054] Compositions of the invention may further comprise adjuvants. Known adjuvants include, for example, emulsions such as Freund's Adjuvants and other oil emulsions, *Bordetella pertussis*, MF59, purified saponin from *Quillaja saponaria* (QS21), aluminum salts such as hydroxide, phosphate and alum, calcium phosphate, (and other metal salts), gels such as aluminum hydroxide salts, mycobacterial products including muramyl dipeptides, solid materials, particles such as liposomes and virosomes. Examples of natural and bacterial products known to be used as adjuvants include monophosphoryl lipid A (MPL), RC-529 (synthetic MPL-like acylated monosaccharide), OM-174 which is a lipid A derivative from *E. coli*, holotoxins such as cholera toxin (CT) or one of its derivatives, pertussis toxin (PT) and heat-labile toxin (LT) of *E. coli* or one of its derivatives, and CpG oligonucleotides. Adjuvant activity can be affected by a number of factors, such as carrier effect, depot formation, altered lymphocyte recirculation, stimulation of T-lymphocytes, direct stimulation of B-lymphocytes and stimulation of macrophages.

15

20

25

[0055] Compositions of the invention are typically formulated as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients, which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, *e.g.*, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or

30

adjuvants, which enhance the effectiveness of the vaccine. The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly.

[0056] Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkalene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of 10 mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25-70%.

[0057] Compositions may also be administered through transdermal routes utilizing

15 jet injectors, microneedles, electroporation, sonoporation, microencapsulation, polymers or liposomes, transmucosal routes and intranasal routes using nebulizers, aerosols and nasal sprays. Microencapsulation using natural or synthetic polymers such as starch, alginate and chitosan, D-poly L-lactate (PLA), D-poly DL-lactic-coglycolic microspheres, polycaprolactones, polyorthoesters, polyanhydrides and

20 polyphosphazenes polyphosphazanes are useful for both transdermal and transmucosal administration. Polymeric complexes comprising synthetic poly-ornithate, poly-lysine and poly-arginine or amphipathic peptides are useful for transdermal delivery systems. In addition, due to their amphipathic nature, liposomes are contemplated for transdermal, transmucosal and intranasal vaccine delivery

25 systems. Common lipids used for vaccine delivery include N-(1)2,3-(dioleyl-dihydroxypropyl)-N,N,N, - trimethylammonium-methyl sulfate (DOTAP), dioleyloxy-propyl - trimethylammonium chloride DOTMA, dimystyloxypropyl-3-dimethyl-hydroxyethyl ammonium (DMRIE), dimethyldioctadecyl ammonium bromide (DDAB) and 9N(N',N-dimethylaminoethane) carbamoyl) cholesterol (DC-

30 Chol). The combination of helper lipids and liposomes will enhance up-take of the liposomes through the skin. These helper lipids include dioleoyl phosphatidylethanolamine (DOPE), dilauroylphosphatidylethanolamine (DLPE), dimyristoyl phosphatidylethanolamine (DMPE),

dipalmitoylphosphatidylethanolamine (DPPE). In addition, triterpenoid glycosides or saponins derived from the Chilean soap tree bark (*Quillaja saponaria*) and chitosan (deacetylated chitan) have been contemplated as useful adjuvants for intranasal and transmucosal vaccine delivery.

5 [0058] Formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

Methods of Inhibiting *H. influenzae*

[0059] Alternatively, the invention includes methods of inhibiting *H. influenzae* type IV pili function in an individual. The methods comprise administering to the individual, for example, one or more antibodies of the invention and/or one or more chimeric proteins of the invention; in an amount that inhibits function of the pili. *In vitro* assays may be used to demonstrate the ability to inhibit pili function.

10 Embodiments of these methods include, for example, methods using inhibitors of adherence mediated via type IV pili, inhibitors that disrupt existing biofilms mediated by type IV pili, and inhibitors of twitching.

[0060] Inhibition is contemplated for any pathological condition involving *H. influenzae*, for example, OM, pneumonia, sinusitis, septicemia, endocarditis, epiglottitis, septic arthritis, meningitis, postpartum and neonatal infections, 20 postpartum and neonatal sepsis, acute and chronic salpingitis, epiglottis, pericarditis, cellulitis, osteomyelitis, endocarditis, cholecystitis, intraabdominal infections, urinary tract infection, mastoiditis, aortic graft infection, conjunctivitis, Brazilian purpuric fever, occult bacteremia, chronic obstructive pulmonary disease and exacerbation of underlying lung diseases such as chronic bronchitis, bronchiectasis and cystic fibrosis.

25 [0061] Compositions comprising inhibitors of *H. influenzae* type IV pili function are provided. The compositions may consist of one of the foregoing active ingredients alone, may comprise combinations of the foregoing active ingredients or may comprise additional active ingredients used to treat bacterial infections. As discussed above, the compositions may comprise one or more additional ingredients 30 such as pharmaceutically effective carriers. Also as discussed above, dosage and frequency of the administration of the compositions are determined by standard techniques and depend, for example, on the weight and age of the individual, the route

of administration, and the severity of symptoms. Administration of the pharmaceutical compositions may be by routes standard in the art, for example, parenteral, intravenous, oral, buccal, nasal, pulmonary, rectal, intranasal, or vaginal.

Animal Model

5 [0062] Methods of the invention may be demonstrated in a chinchilla model widely accepted as an experimental model for OM. In particular, a chinchilla model of NTHi-induced OM has been well characterized (Bakaletz *et al.*, *J. Infect. Dis.*, 168: 865-872, 1993; Bakaletz and Holmes, *Clin. Diagn. Lab. Immunol.*, 4: 223-225, 1997; Suzuki and Bakaletz, *Infect. Immun.*, 62: 1710-1718, 1994; Mason *et al.*, *Infect. Immun.*, 71:3454-3462, 2003), and has been used to determine the protective efficacy of several NTHi outer membrane proteins, combinations of outer membrane proteins, chimeric synthetic peptide vaccine components, and adjuvant formulations against OM (Bakaletz *et al.*, *Vaccine*, 15: 955-961, 1997; Bakaletz *et al.*, *Infect. Immun.*, 67: 2746-2762, 1999; Kennedy *et al.*, *Infect. Immun.*, 68: 2756-2765, 2000; Kyd *et al.*, *Infect. Immun.*, 66:2272-2278, 2003; Novotny and Bakaletz, *J. Immunol.*, 171, 1978-1983, 2003).

10 [0063] In the model, adenovirus predisposes chinchillas to *H. influenzae*-induced OM media, which allowed for the establishment of relevant cell, tissue and organ culture systems for the biological assessment of NTHi (Bakaletz *et al.*, *J. Infect. Dis.*, 168: 865-72, 1993; Suzuki *et al.*, *Infect. Immunity* 62: 1710-8, 1994). Adenovirus infection alone has been used to assess the transudation of induced serum antibodies into the tympanum (Bakaletz *et al.*, *Clin. Diagnostic Lab Immunol.*, 4(2): 223-5, 1997) and has been used as a co-pathogen with NTHi, to determine the protective efficacy of several active and passive immunization regimens targeting various NTHi outer membrane proteins, combinations of OMPs, chimeric synthetic peptide vaccine components, and adjuvant formulations as vaccines against otitis media (Bakaletz *et al.*, *Infect Immunity*, 67(6): 2746-62, 1999; Kennedy *et al.*, *Infect. Immun.*, 68(5): 2756-65, 2000; Novotny *et al.*, *Infect Immunity* 68(4): 2119-28, 2000; Poolman *et al.*, *Vaccine* 19 (Suppl. 1): S108-15, 2000).

Brief Description of Drawing

[0064] Figure 1 provides the timeline of the immunization regimen, viral inoculation, bacterial challenge, and OM disease assessment period for the efficacy experiments described in Example 5.

5

Detailed Description of the Invention

[0065] The following examples illustrate the invention wherein Example 1 describes recombinant production of chimeric proteins of the invention, Example 2 describes assays to test the immunogenicity of the chimeric proteins of the invention, Example 3 describes assays for evaluating passive immunization, Example 4 10 describes assays for evaluating active immunization and Example 5 describes the evaluation of a chimeric protein of the invention.

Example 1

Synthesis of Chimeric Proteins

[0066] The chimeric proteins of the invention were produced using standard 15 recombinant methods. Initially, a gene-synthesis company, (Blue Heron Biotechnology Inc.) was contracted to make the initial plasmid based on the chimeric protein amino acid sequences described herein that were optimized for *E. coli* preferred codon usage. Briefly, the native NTHi pilin protein sequence was modified by truncating the N-terminus (residues 1-39 of SEQ ID NO: 2) and adding a HIS-tag 20 sequence and a thrombin cleavage site as set out in SEQ ID NO: 3. The HIS-tag was preceded by a sequence (MGSS) to assist in expression. The thrombin cleavage site allowed for release of the HIS-tag. These plasmids were then cloned into the *E. coli* expression vector pET-15b vector (Novagen). The plasmid were then transformed 25 into *E. coli* strain “Origami(DE3)” (available from Novagen) as the host for expression of soluble His-tagged chimeric proteins. Another *E.coli* host cell expression stain that may be used is Origami B(DE3) (Novagen).

[0067] The His-tagged variants of the chimeric proteins will be recovered by nickel 30 column chromatography, then used for initial studies to determine if they are reactive with antisera directed against any of the following: native OMP P5-fimbrin, LB1 (full length 40 amino acid peptide), LB1(1) (a synthetic peptide representing just the 19 amino acid B-cell epitope of LB1), recombinant PilA protein or native PilA protein. Once the His-tag is removed by thrombin site cleavage, the recombinant

chimeric proteins will be used as immunogens to determine their immunogenicity and protective capability.

[0068] Exemplary chimeric proteins of the invention have the sequences as set out in Table 2 below. The chimeric proteins having the amino acid sequences of SEQ ID

5 NOS: 10, 12 and 14 have been expressed by *E. coli* as described above.

Table 2

SEQ ID NO:	Chimeric Protein Amino Acid Sequence
9	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CRSDYKFYEDANGTRDHKKGCTGGKNGIAADITTAKGYVKSV TTSNGAITVKGDGTANMEYILQATGNAATGVTWTTCKGTD SLFPANFCGSVTQ
10	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CLVRSDYKFYEDANGTRDHKKGHTCTGGKNGIAADITTAKGY VKSVTTSNGAITVKGDGTANMEYILQATGNAATGVTWTTCK GTDASLFPANFCGSVTQ
11	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSNGAITVKG DGTLANMEYILQATGNAATGVTWTTCRSDYKFYEDANGTRD HKKGCGSVTQ
12	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSNGAITVKG DGTLANMEYILQATGNAATGVTWTTCLVRSDYKFYEDANGTR DHKKGRHTCGSVTQ
13	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSNGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQ RSDYKFYEDANGTRDHKKG
14	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSNGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQLVRSDYKFYEDANGTRDHKKGRHT
15	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CRSDYKLYNKNSSSNSTLKNLGETGGKNGIAADITTAKGYVK SVTTSNGAITVKGDGTANMEYILQATGNAATGVTWTTCKGT DASLFPANFCGSVTQ

SEQ ID NO:	Chimeric Protein Amino Acid Sequence
16	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CLVRSDYKLYNKNSSSNLKNLGERHTCTGGKNGIAADITTA KGYVKSVTTSGAITVKGDGTANMEYILQATGNAATGVTWT TCKGTDASLFPANFCGSVTQ
17	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCRSDYKLYNKNSSSN LKNLGERHTCGSVTQ
18	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCLVRSDYKLYNKNSSSN STLKNLGERHTCGSVTQ
19	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQRSDYKLYNKNSSSNSTLKNLGE
20	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQLVRSDYKLYNKNSSSNSTLKNLGERHT
21	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CRSDYKLYNKNSSSLKNLGECTGGKNGIAADITTAKGYVKSVT TSNGAITVKGDGTANMEYILQATGNAATGVTWTTCKGTDAS LFPANFCGSVTQ
22	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CLVRSDYKLYNKNSSSLKNLGERHTCTGGKNGIAADITTAKG YVKSVTTSGAITVKGDGTANMEYILQATGNAATGVTWTT KGTDASLFPANFCGSVTQ
23	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCRSDYKLYNKNSSTLKN LGECSVTQ
24	MGSSHHHHHSSGLVPRGSHMTKKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCLVRSDYKLYNKNSSTL KNLGERHTCGSVTQ

SEQ ID NO:	Chimeric Protein Amino Acid Sequence
25	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQRSDYKLYNKNSSTLKNLGE
26	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQLVRSDYKLYNKNSSTLKNLGERHT
27	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CRSDYKFYDNKRIDCTGGKNGIAADITTAKGYVKSVTTSGAI TVKGDGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANF CGSVTQ
28	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CLVRSDYKFYDNKRIDRHTCTGGKNGIAADITTAKGYVKSVTT SNGAITVKGDGTLANMEYILQATGNAATGVTWTTCKGTDASL FPANFCGSVTQ
29	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCRSDYKFYDNKRIDCGS VTQ
30	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCLVRSDYKFYDNKRIDR HTCGSVTQ
31	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQRSDYKFYDNKRID
32	MGSSHHHHSSGLVPRGSHMTKAAVSELLQASAPYKADVEL CVTSTNETTNCTGGKNGIAADITTAKGYVKSVTTSGAITVKG DGTLANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGSV TQLVRSDYKFYDNKRIDRHT

[0069] Additional exemplary chimeric proteins of the invention have the amino acid sequences as set out in Table 3 below. These chimeric proteins have been expressed by *E. coli* and purified using a HIS-tag, as described above. The chimeric proteins set out in Table 3 have the His tag sequence removed for use as an

immunogen. The chimeric protein having the amino acid sequence of SEQ ID NO: 56 was used in the studies described in Example 5.

Table 3

SEQ ID NO:	Chimeric Protein Amino Acid Sequence
54	GSHMTKKA AVSELLQASAPYKADVELCLVRS DYKFYEDANGT RDHKKGRHTCTGGKNGIAADITTA KGYVKS VTT SNGAITVKG DGT LANMEYILQATGNAATGVTWTTCKGTDASLFPANFCGS VTQ
55	GSHMTKKA AVSELLQASAPYKADVELCVYSTNETTNCTGGKN GIAADITTA KGYVKS VTT SNGAITVKG DGT LANMEYILQATGN AATGVTWTTCLVRS DYKFYEDANGTRDHKKGRHTCGSVTQ
56	GSHMTKKA AVSELLQASAPYKADVELCVYSTNETTNCTGGKN GIAADITTA KGYVKS VTT SNGAITVKG DGT LANMEYILQATGN AATGVTWTTCKGTDASLFPANFCGSVTQLVRS DYKFYEDAN GTRDHKKGRHT
57	GSHMLVRS DYKFYEDANGTRDHKKGRHTGPSLKLT KKA AVSE LLQASAPYKADVELCVYSTNETTNCTGGKNGIAADITTA KGYV KSVTT SNGAITVKG DGT LANMEYILQATGNAATGVTWTTCK GTDASLFPANFCGSVTQ

5

Example 2

Immunogenicity of Chimeric Proteins

[0070] Rabbits or chinchillas are immunized with the chimeric proteins. Rabbits receive an initial immunizing dose of 500 µg of a chimeric protein in complete Freund's adjuvant. The rabbits receive a second dose of 400 µg of the chimeric protein

10 21 days later. The rabbits receive a third dose of chimeric protein in complete Freund's adjuvant 42 days after the initial immunizing dose with 400 µg of the same peptide in either IFA or PBS (one rabbit per diluent). Sera are obtained 3 weeks after each dose. Chinchillas receive an initial immunizing dose of 10 µg of the chimeric protein in the adjuvant monophosphoryl lipid A (MPL). One month (~ 30 days) later,

15 chinchillas receive a second identical dose. The third and final dose is delivered ~ 30 days after the second dose. Sera are obtained ~10-14 days after each dose. The sera from all animals are assessed for titer and specificity against the LB1 peptide (40-mer), LB1(1), PilA protein and the chimeric proteins, by ELISA, Western blot and biosensor. Antisera are also tested against whole bacteria via flow cytometry (FACS) analysis.

Example 3Evaluating Passive Immunization

[0071] The protection conferred by an animal's immune response directed against the chimeric proteins of the invention is determined in a chinchilla model of experimental otitis media. Chinchillas are passively immunized with 5 ml/kg hyperimmune chinchilla or human serum directed against a chimeric protein of the invention. Control chinchillas receive normal chinchilla serum or normal human serum. Next the chinchillas receive first a viral co-pathogen intranasally, then a week later, an intranasal challenge with the NTHi bacteria. The chinchillas are examined and rated daily or every 2 days for up to 35 days after bacterial challenge. Immunized chinchillas receiving immune chinchilla or human serum display reduced tympanic membrane pathology and reduced or absent signs of infection of the middle ear space as determined by both video otoscopy and tympanometry. In this assay, the presence of middle ear fluids in chinchillas receiving chinchilla or human anti-chimeric protein serum is reduced when compared to controls.

Example 4Evaluating Active Immunization

[0072] Cohorts of 5-10 chinchillas each are actively immunized with either a saline control preparation, an adjuvant-only preparation, or one of the chimeric proteins of the invention that has been admixed with an appropriate adjuvant. The immunogens are assessed for endotoxin content prior to their use as an immunogen via a chromogenic Amoebocyte Lysate assay which is commercially available from Whittaker Bioproducts under the designation QCL-1000. The chinchillas are then subcutaneously injected with 10 µg immunogen in the adjuvant MPL (or another appropriate adjuvant). Then 30 days later they receive 10 µg of the same immunogen in MPL. Thirty days following the second immunization, these animals receive the final immunizing dose. Approximately 10-14 days after the final immunizing dose is delivered, chinchillas are challenged both transbulularly and intranasally with a strain of NTHi. The chinchillas are assessed over a 35-day period for: tympanic membrane pathology by video otoscopic examination and tympanometry; semiquantitation of NTHi recovered via epitympanic tap of the inferior bulla and passive lavage of the nasopharynx; and light microscopic examination of fixed middle ear mucosal

epithelium and tympanic membrane for histopathology. For example, chinchillas immunized with the chimeric proteins of the invention will have reduced tympanic membrane pathology, will be free of middle ears effusions or they will contain effusions that are culture-negative, there will be reduced or no biofilm present in the 5 tympanum and there will be minimal thickening of the middle ear mucosa, minimal osteoneogenesis and reduced presence of both red blood cells and inflammatory cells in the subepithelial space.

Example 5

Evaluation of Chimeric Proteins

10 [0073] The protective efficacy of the chimeric protein having the amino acid sequence of SEQ ID NO: 56 (referred to as “chim-V3” herein) was evaluated using the chinchilla passive-transfer, superinfection model of OM. This chimeric peptide comprised the B-cell epitope of the LB1 peptide (SEQ ID NO: 5) expressed after the C-terminal glutamine residue of recombinant PilA (residues 40-149 of SEQ ID NO: 2). To generate polyclonal antiserum for use in passive transfer efficacy studies, the chim-V3 protein was delivered to a cohort of adult chinchillas with the adjuvant, monophosphoryl lipid A (MPL) plus trehalose dimycolate (Corixa). A timeline depicting the immunization regimen is set out in Fig. 1. To generate immune serum pools, alert prone chinchillas were immunized subcutaneously 3 times with 30 µg of 15 chim-V3 plus 10 µg of MPL or 10 µg MPL alone every 21 days. At day 56, a terminal bleed of the inoculated animals was collected and serum was pooled for transfer to naïve juvenile animals. To study efficacy, a separate cohort of juvenile chinchillas was first challenged with adenovirus on day -7. Six days later (day -1), the pooled anti-chim-V3 immune serum was passively transferred to these 20 adenovirus-compromised animals. The following day (day 0), animals that received anti-chim-V3 serum by passive transfer were challenged with the bacterium, nontypeable *Haemophilus influenzae*. These animals were then monitored for incidence and severity of disease over a 26-day time-course (relative to bacterial challenge) by daily video otoscopy and tympanometry as well as Xenogen *in vivo* 25 imaging every other day.

[0074] The titer of anti-chim-V3 antibody was measured in the immune serum collected from the inoculated animals using an ELISA. This analysis demonstrated that the collected antiserum contained antibodies specific for the chim-V3 protein.

The presence of anti-chim-V3 antibodies in the collected antiserum was also confirmed using Western blot analysis.

[0075] FACS analysis was used to measure the ability of serum immunoglobulins from immunized animals to recognize surface exposed native structures expressed by NTHi 86-028 NP. NTHi bacteria were incubated with chim-V3 antiserum, washed, then incubated with naïve or immune FITC-Protein A, washed and analyzed by FACS analysis. Inoculation with the chim-V3 protein induced a significant increase in antibodies that were capable of recognizing the NTHi surface proteins or chim-V3 protein. The data obtained were dependent on both antibody titer and avidity as well as relative expression of both the type IV pilus and the OMP P5-homologous adhesion by NTHi when grown *in vitro*.

[0076] The luminescent reporter NTHi 86-028 NP pKMLN-1 was used to detect NTHi infection in the animals inoculated with chim-V3 protein using Xenogen *in vivo* real-time imaging. Growth curves of the luminescent strain NTHi 86-028 NP

pKMLN-1 and the parental strain NTHi 86-028 demonstrated that growth of the luminescent NTHi strain was comparable to the parental strain. Luminescent imaging of NTHi residing in the nasopharynx of the inoculated animals was readily accomplished however, due to the microaerophilic nature of the diseased middle ear, luminescence of NTHi present in the middle ear could not be monitored over the entire disease course because the luminescence is dependent on the availability of oxygen. Animals were monitored every other day for the presence of luminescent bacteria, and if bacteria were detected, this was recorded as a luminescent event. Luminescent infection was detected at least six days after challenge in the inoculated animals. The total number of luminescent events in the chim-V3 inoculated animals was less than the total number of luminescent events in the control animals (inoculated with MPL only).

[0077] Throughout the course of the study, daily video otoscopy and tympanometry was used to determine the percent of chinchilla middle ears with OM. Inoculation with chim-V3 caused 53% reduction in the number of animals with middle ears having OM as compared to control animals (inoculated with MPL only).

[0078] All of these studies demonstrate that the chim-V3 protein was immunogenic and anti-chim-V3 antibodies were protective in the chinchilla passive transfer-superinfection model of OM.

We claim:

1. A chimeric protein comprising a portion of PilA and a portion of the LB1 peptide

5

2. The chimeric protein of claim 1 wherein the portion of the LB1 peptide comprises the B-cell epitope of the LB1 peptide.

10 3. The chimeric protein of claim 2 wherein the portion of the LB1 peptide comprises the amino acid sequence of SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 or SEQ ID NO: 8.

15 4. The chimeric protein of claim 1 comprising the amino acid sequence of SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56 or SEQ ID NO: 57.

20

5. A polynucleotide encoding the chimeric protein of any one of claims 1 to 4.

25

6. A vector comprising a polynucleotide of claim 5.

7. A composition comprising a chimeric protein of any one of claims 1 to 4 and a pharmaceutically acceptable carrier.

30

8. An antibody that specifically binds to a chimeric protein of any one of claims 1 to 4.

9. A composition comprising an antibody of claim 8 and a pharmaceutically acceptable carrier.

10. A method for eliciting an immune response to NTHi bacteria comprising administering an immunogenic dose of one or more chimeric proteins of any one of claims 1 to 4 to a patient at risk of NTHi bacterial infection.


5 11. The method of claim 10 wherein the NTHi infection is in the middle ear.

12. A method of treating or preventing NTHi bacterial infection comprising administering an antibody of claim 8 to a patient in need thereof.

10 13. The method of claim 12 wherein the NTHi infection is in the middle ear.

1/1

Figure 1



SEQUENCE LISTING

<110> Bakalatz, et al.

<120> CHIMERIC VACCINE FOR HAEMOPHILUS INFLUENZAE-INDUCED DISEASE

<130> 28335/42096A

<150> US 60/801,835

<151> 2006-05-19

<150> US 60/697,642

<151> 2005-07-08

<160> 56

<170> PatentIn version 3.3

<210> 1

<211> 4345

<212> DNA

<213> H. influenzae

<220>

<221> CDS

<222> (403)..(849)

<400> 1

gcgatcatta aaattgacat attgcgtaat tcgcccattt cggtcgatca aacaatgtgc	60
tgaaaacacgc atttgataaa ttctgcaaa ataaggatga atcttaggat ctaattttcc	120
ttgaaaaaaaaa tcatccacat atccgcccgc aaattgttct ggcggcagac taatataatg	180
aataaccaat aaggaaatata cctgtggatt tgggcgttta tcgaagttag gtgactgaat	240
ttgccgacaa tccaatatac cttgttcaat atcttttagt ttttgcatac ttttttcctt	300
tttttgcgt caggatcgca gaaaaagtgc ggtcaatttt acaaacaaat ttttcctttt	360
cacaatgtcg tcgctaacaa aggcttaata aaaggaaaat ga atg aaa cta aca	414
Met Lys Leu Thr	
1	

aca cag caa acc ttg aaa aaa ggg ttt aca tta ata gag cta atg att	462
Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile Glu Leu Met Ile	
5 10 15 20	

gtg att gca att att gct att tta gcc act atc gca att ccc tct tat	510
Val Ile Ala Ile Ala Ile Leu Ala Thr Ile Ala Ile Pro Ser Tyr	
25 30 35	

caa aat tat act aaa aaa gca gcg gta tct gaa tta ctg caa gcg tca	558
Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala Ser	
40 45 50	

gcg cct tat aag gct gat gtg gaa tta tgt gta tat agc aca aat gaa	606
Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr Ser Thr Asn Glu	
55 60 65	

aca aca aac tgt acg ggt gga aaa aat ggt att gca gca gat ata acc	654
Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr	

70	75	80	
aca gca aaa ggc tat gta aaa tca gtg aca aca agc aac ggt gca ata Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile 85	90	95	702 100
aca gta aaa ggg gat ggc aca ttg gca aat atg gaa tat att ttg caa Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln 105	110		750 115
gct aca ggt aat gct gca aca ggt gta act tgg aca aca act tgc aaa Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys 120	125	130	798
gga acg gat gcc tct tta ttt cca gca aat ttt tgc gga agt gtc aca Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr 135	140	145	846
caa tgacgagcta tgctttactt catactcagc gtgttaaccgc tcaaaaatggc Gln			899
gagatcttta cgatctcgcc agatttatgg gaacgcaatc agcagcaaca atccttgctc ttgcggattt ttgctttgcc acttaaagaa gaaaataatc gtctttggct aggggttgat tctctctcca atcttcagc ttgtgaaacc attgcgttta taacaggaaa acctgtcgaa ccaattttgt tagaaagcag ccaactcaaa gaactgttac aacaacttac tccgcaccaa atgcaagtgg aagagcaagt taaattctat caacatcaag aaaccattt tgaacaagaa gatgatgaac ctgttatccg cttacttaat cagattttt aatctgcctt aaaaaaaaaat gcctctgata ttcatttaga aaccttggct gatcagtttc aagtgcgggt tagaattgat ggtgtttac aaccacaacc cttataataatc aaaatattcg ccaatcgtat tatttcacgc ttaaaattac tggctaaatt agatatttagt gaaaatcgac ttccacaaga tggacgattt caatttaaaa ccacttttc cgatattctt gatttcgccc tttcaacctt accaaccat tggggcgaaa aaatcgtgtt gcgagcgcaaa caaaataaaac ctgtagaact tagctttgct gaactgggta tgaccgaaaa tcagcaacaa gcatttcaac gctcacttag ccagccacaa ggattaattt tagtaaccgg ccccacagga agtggggaaaa gtatctcgct ttacaccgca cttcagtgcc taaatacgcc tgataaacat attatgaccg ctgaagatcc cattgaaatt gaacttgatg gtattattca aagccaaattt aatccgcaga ttggattaga ttttagccgt ctattcgttgc cttttttacg tcaagatccc gacatcatta tgcttaggtga aattcgagat gaagaaagtgc caaggattgc actacgtgcc gctcaaacgg gacatttgggt gctttcaact ttacataccca atgatgcaat atctgccatt tctcgcttac aacaactcgg tattcaacaa catgaaatttgc aaaaacagttt actactcgtc attgcacagc gtcttgcgtt aaaaatctgt ccaaagtgcgttgc gtggaaattt aataaattct tttgtgattgcc atcaagggttac tggagggcga atcggcgtgttgc atcaatttctt acattggcaaa cagaatgggttac tcaaaacggaa ttttgagaat 2159			

ttacgagaga	gtggtttggaa	aaaagtttagc	caaggcataa	cagatgagaa	agaaaattgaa	2219
cgtgtgttag	gtaaaaaactc	atgactaaaa	aactcttttta	ttatcaaggt	agtaacgcatt	2279
taaatcagaa	acaaaaaggc	tcaattatttgc	cggatacgaa	acaacaagcg	cactttcagt	2339
taataagccg	cgggcttact	cacatcaaatt	tacaacaaaa	ctggcaattt	ggggcaaaac	2399
ccaaaaattc	agaaatcagt	gaattactca	atcaatttagc	gacattgcta	cagtcgtaa	2459
ttccgttaaa	aaacagccta	caaattttgc	aacaaaatttgc	tactcaaattt	atgctcaaca	2519
aatggcttga	acgactgttt	caatccatttgc	aatctggctt	agcattctca	caagccatttgc	2579
aacaacaagg	aaaatatctc	acacaacaag	aaattcaact	gattcaagtg	ggagaaatgaa	2639
cagaaaaact	tgccgttagtt	tgtaaaaaaaa	tagccacgca	ccgttagtcaa	tctttggctt	2699
tacaacgcaa	attacagaaa	attatgttat	atccctcaat	ggtattggga	atttctctat	2759
tattgacact	cgcattactg	ctttttatcg	cgcctcaattt	tgctgaaaatg	tacagtggca	2819
ataatgcgga	gttaccaaca	ataaccgcaa	tattgctctc	aatatctaatt	ttccttaagc	2879
aaaatattgg	cattttgcta	tttttcgttt	ttagtttttt	tctattttat	tatttctatc	2939
taaaacgcca	gacttgggtt	catcaaaaga	aaaatcaact	tatttctatc	acgccttattt	2999
ttggcacaat	tcaaaagctt	tcacgttttag	tgaacttttag	tcaaagtttta	caaatttatgt	3059
tgcaggccgg	cgtaccgctt	aatcaggcac	tagacagttt	tcttcctcgc	acacaaactt	3119
ggcaaaccaa	gaaaacgctt	gtaaacgata	tggattttaga	taaagaagtg	cggtaattt	3179
tgcaatgggt	ttctcaaggc	tatgcgtttt	ctaatacggt	aagtagcgat	cttttcccga	3239
tggaaggcaca	acaaatgcta	caaattggcg	aacaaagcgg	aaaactcgct	ttgatgctag	3299
agcatatcgc	agataattac	caagaaaaac	ttaatcatca	aattgactta	ctctcacaaa	3359
tgctagaacc	attaatgatg	gtaatcatcg	gcagtcgtat	tggatttattt	atgatggaa	3419
tgtatttacc	tatctttaat	atgggatcag	ttattcaatg	atttacttca	caatgttttt	3479
attaggcggc	atcttaggaa	tcgcattgtg	gttctaccta	tctggtttttta	ttacgcattt	3539
gcagcaagag	atttatgcga	cttacgttga	attatttcca	caaaacagtt	ctccatttca	3599
accgcacttt	gcctctattt	aacaaaaaaa	gtgcggcat	attttgaggt	atttttttag	3659
tattggggtt	ggatttatata	ttttacaaat	tgccttcaaa	gattctattt	ttactgtatg	3719
gatccggactc	acacttatta	ttctttggac	aatcagttat	cttgattggc	actatcaact	3779
tatttctacg	acaccctgtt	tatggttact	tactctcggt	ttatttggcg	cagacaataa	3839
cttttcatttgc	ctaacgttat	ctgaaagcat	aaaaagtgcg	gctagttttt	ttattgtttt	3899
ctacgcaatc	tattggatttgc	caaaatgtta	ttatagaaaa	gaaggcttttgc	gacggggaga	3959
ttattggcttgc	gcaatggcat	taggaagttt	tattcattta	gaaaccttac	cgcacttttt	4019
attatttagcc	tcagtgccttgc	gaatatgttt	ttcgcttattt	cataaaaaga	aaaaagaattt	4079

tataacctttt	gccccttta	tgaacttac	ggctatcatt	atttatctcg	tcaaataatta	4139
cggatattaa	aaaggggaaa	acataatatt	tttcccttgt	tcttcataga	agtgcggttg	4199
tttttaccaa	cgtttcatca	cttcaaaaaa	ctttcgttg	gttttcgcca	tcatcagctt	4259
atcaatcaag	aattccattt	catccacttc	atccatttgg	ttaagaatct	tacgaagaat	4319
ccacattttt	tgtaattcgt	ccgctg				4345

<210> 2

<211> 149

<212> PRT

<213> H. influenzae

<400> 2

Met	Lys	Leu	Thr	Thr	Gln	Gln	Thr	Leu	Lys	Lys	Gly	Phe	Thr	Leu	Ile
1					5				10					15	

Glu	Leu	Met	Ile	Val	Ile	Ala	Ile	Ile	Ala	Ile	Leu	Ala	Thr	Ile	Ala
					20			25					30		

Ile	Pro	Ser	Tyr	Gln	Asn	Tyr	Thr	Lys	Lys	Ala	Ala	Val	Ser	Glu	Leu
							35		40			45			

Leu	Gln	Ala	Ser	Ala	Pro	Tyr	Lys	Ala	Asp	Val	Glu	Leu	Cys	Val	Tyr
					50			55			60				

Ser	Thr	Asn	Glu	Thr	Thr	Asn	Cys	Thr	Gly	Gly	Lys	Asn	Gly	Ile	Ala
					65			70		75			80		

Ala	Asp	Ile	Thr	Thr	Ala	Lys	Gly	Tyr	Val	Lys	Ser	Val	Thr	Thr	Ser
					85				90		95				

Asn	Gly	Ala	Ile	Thr	Val	Lys	Gly	Asp	Gly	Thr	Leu	Ala	Asn	Met	Glu
					100			105			110				

Tyr	Ile	Leu	Gln	Ala	Thr	Gly	Asn	Ala	Ala	Thr	Gly	Val	Thr	Trp	Thr
					115			120			125				

Thr	Thr	Cys	Lys	Gly	Thr	Asp	Ala	Ser	Leu	Phe	Pro	Ala	Asn	Phe	Cys
					130			135			140				

Gly	Ser	Val	Thr	Gln											
				145											

<210> 3

<211> 131

<212> PRT

<213> H. influenzae

<400> 3

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1 5 10 15

Arg Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln
20 25 30

Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr Ser Thr
35 40 45

Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp
50 55 60

Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly
65 70 75 80

Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile
85 90 95

Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Thr
100 105 110

Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser
115 120 125

Val Thr Gln
130

<210> 4

<211> 19

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic Peptide

<400> 4

Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp His
1 5 10 15

Lys Lys Gly

<210> 5

<211> 24

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic Peptide

<400> 5

Leu Val Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg
1 5 10 15

Asp His Lys Lys Gly Arg His Thr
20

<210> 6

<211> 22

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Peptide

<400> 6

Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Ser Asn Ser Thr
1 5 10 15

Leu Lys Asn Leu Gly Glu
20

<210> 7

<211> 19

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic Peptide

<400> 7

Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Thr Leu Lys Asp
1 5 10 15

Leu Gly Glu

<210> 8

<211> 13

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic Peptide

<400> 8

Arg Ser Asp Tyr Lys Phe Tyr Asp Asn Lys Arg Ile Asp
1 5 10

<210> 9

<211> 139

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric protein

<400> 9

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Arg Ser Asp Tyr Lys
35 40 45

Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp His Lys Lys Gly Cys Thr
50 55 60

Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr Thr Ala Lys Gly Tyr
65 70 75 80

Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile Thr Val Lys Gly Asp
85 90 95

Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln Ala Thr Gly Asn Ala
100 105 110

Ala Thr Gly Val Thr Trp Thr Cys Lys Gly Thr Asp Ala Ser Leu
115 120 125

Phe Pro Ala Asn Phe Cys Gly Ser Val Thr Gln
130 135

<210> 10

<211> 143

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric Protein

<400> 10

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Leu Val Arg Ser Asp
35 40 45

Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp His Lys Lys Gly
50 55 60

His Thr Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr Thr
65 70 75 80

Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile Thr
85 90 95

Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln Ala
100 105 110

Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys Gly Thr
115 120 125

Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr Gln
130 135 140

<210> 11

<211> 136

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric Protein

<400> 11

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Arg
100 105 110

Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp His Lys
115 120 125

Lys Gly Cys Gly Ser Val Thr Gln
130 135

<210> 12
<211> 141
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 12

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Cys Leu
100 105 110

Val Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp
115 120 125

His Lys Lys Gly Arg His Thr Cys Gly Ser Val Thr Gln
130 135 140

<210> 13
<211> 148
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 13

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
 100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
 115 120 125

Gln Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp
 130 135 140

His Lys Lys Gly
 145

<210> 14
 <211> 153
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 14

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Lys Asn Gly Ile Ala Ala Asp Ile

50

55

60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
 100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
 115 120 125

Gln Leu Val Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr
 130 135 140

Arg Asp His Lys Lys Gly Arg His Thr
 145 150

<210> 15
 <211> 142
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 15

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Arg Ser Asp Tyr Lys
 35 40 45

Leu Tyr Asn Lys Asn Ser Ser Asn Ser Thr Leu Lys Asn Leu Gly
 50 55 60

Glu Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr Thr Ala
 65 70 75 80

Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile Thr Val
 85 90 95

Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln Ala Thr
 100 105 110

Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys Gly Thr Asp
 115 120 125

Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr Gln
 130 135 140

<210> 16
 <211> 147
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 16

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Leu Val Arg Ser Asp
 35 40 45

Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Asn Ser Thr Leu Lys Asn
 50 55 60

Leu Gly Glu Arg His Thr Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala
 65 70 75 80

Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn
 85 90 95

Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr
 100 105 110

Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr
 115 120 125

Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser
 130 135 140

Val Thr Gln
 145

<210> 17
 <211> 139
 <212> PRT
 <213> Artificial sequence

<220>

<223> Chimeric Protein

<400> 17

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Arg
 100 105 110

Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Ser Asn Ser Thr Leu
 115 120 125

Lys Asn Leu Gly Glu Cys Gly Ser Val Thr Gln
 130 135

<210> 18

<211> 144

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric Protein

<400> 18

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Leu
 100 105 110

Val Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Ser Asn Ser
 115 120 125

Thr Leu Lys Asn Leu Gly Glu Arg His Thr Cys Gly Ser Val Thr Gln
 130 135 140

<210> 19
 <211> 151
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 19

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
 100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr

115

120

125

Gln Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Ser Asn Ser
130 135 140

Thr Leu Lys Asn Leu Gly Glu
145 150

<210> 20
<211> 156
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 20

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
115 120 125

Gln Leu Val Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Ser
130 135 140

Asn Ser Thr Leu Lys Asn Leu Gly Glu Arg His Thr
145 150 155

<210> 21
<211> 139

<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 21

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Arg Ser Asp Tyr Lys
35 40 45

Leu Tyr Asn Lys Asn Ser Ser Leu Lys Asn Leu Gly Glu Cys Thr
50 55 60

Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr Thr Ala Lys Gly Tyr
65 70 75 80

Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile Thr Val Lys Gly Asp
85 90 95

Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln Ala Thr Gly Asn Ala
100 105 110

Ala Thr Gly Val Thr Trp Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu
115 120 125

Phe Pro Ala Asn Phe Cys Gly Ser Val Thr Gln
130 135

<210> 22
<211> 145
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 22

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Leu Val Arg Ser Asp

35

40

45

Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Ser Thr Leu Lys Asn Leu Gly
 50 55 60

Glu Arg His Thr Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 65 70 75 80

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 85 90 95

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 100 105 110

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
 115 120 125

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
 130 135 140

Gln
 145

<210> 23
 <211> 136
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 23

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Arg
 100 105 110

Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Thr Leu Lys Asn Leu
 115 120 125

Gly Glu Cys Gly Ser Val Thr Gln
 130 135

<210> 24
 <211> 141
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 24

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Leu
 100 105 110

Val Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Thr Leu Lys
 115 120 125

Asn Leu Gly Glu Arg His Thr Cys Gly Ser Val Thr Gln
 130 135 140

<210> 25
 <211> 148
 <212> PRT
 <213> Artificial sequence

<220>

<223> Chimeric Protein

<400> 25

Met	Gly	Ser	Ser	His	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro	Arg
1				5					10					15	

Gly	Ser	His	Met	Thr	Lys	Lys	Ala	Ala	Val	Ser	Glu	Leu	Leu	Gln	Ala
			20				25						30		

Ser	Ala	Pro	Tyr	Lys	Ala	Asp	Val	Glu	Leu	Cys	Val	Thr	Ser	Thr	Asn
					35		40					45			

Glu	Thr	Thr	Asn	Cys	Thr	Gly	Gly	Lys	Asn	Gly	Ile	Ala	Ala	Asp	Ile
					50		55			60					

Thr	Thr	Ala	Lys	Gly	Tyr	Val	Lys	Ser	Val	Thr	Thr	Ser	Asn	Gly	Ala
					65		70			75			80		

Ile	Thr	Val	Lys	Gly	Asp	Gly	Thr	Leu	Ala	Asn	Met	Glu	Tyr	Ile	Leu
					85			90				95			

Gln	Ala	Thr	Gly	Asn	Ala	Ala	Thr	Gly	Val	Thr	Trp	Thr	Thr	Cys	Lys
					100			105				110			

Gly	Thr	Asp	Ala	Ser	Leu	Phe	Pro	Ala	Asn	Phe	Cys	Gly	Ser	Val	Thr
						115		120			125				

Gln	Arg	Ser	Asp	Tyr	Lys	Leu	Tyr	Asn	Lys	Asn	Ser	Ser	Thr	Leu	Lys
						130		135			140				

Asn	Leu	Gly	Glu
		145	

<210> 26

<211> 153

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric Protein

<400> 26

Met	Gly	Ser	Ser	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro	Arg
1				5				10					15	

Gly	Ser	His	Met	Thr	Lys	Lys	Ala	Ala	Val	Ser	Glu	Leu	Leu	Gln	Ala
					20			25					30		

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
115 120 125

Gln Leu Val Arg Ser Asp Tyr Lys Leu Tyr Asn Lys Asn Ser Ser Thr
130 135 140

Leu Lys Asn Leu Gly Glu Arg His Thr
145 150

<210> 27
<211> 133
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 27

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Arg Ser Asp Tyr Lys
35 40 45

Phe Tyr Asp Asn Lys Arg Ile Asp Cys Thr Gly Gly Lys Asn Gly Ile
50 55 60

Ala Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr
65 70 75 80

Ser Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met

85

90

95

Glu Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp
 100 105 110

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 115 120 125

Gly Ser Val Thr Gln
 130

<210> 28
 <211> 138
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 28

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Leu Val Arg Ser Asp
 35 40 45

Tyr Lys Phe Tyr Asp Asn Lys Arg Ile Asp Arg His Thr Cys Thr Gly
 50 55 60

Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val
 65 70 75 80

Lys Ser Val Thr Thr Ser Asn Gly Ala Ile Thr Val Lys Gly Asp Gly
 85 90 95

Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala
 100 105 110

Thr Gly Val Thr Trp Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe
 115 120 125

Pro Ala Asn Phe Cys Gly Ser Val Thr Gln
 130 135

<210> 29
 <211> 130

<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 29

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Arg
100 105 110

Ser Asp Tyr Lys Phe Tyr Asp Asn Lys Arg Ile Asp Cys Gly Ser Val
115 120 125

Thr Gln
130

<210> 30
<211> 135
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 30

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn

35

40

45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Leu
 100 105 110

Val Arg Ser Asp Tyr Lys Phe Tyr Asp Asn Lys Arg Ile Asp Arg His
 115 120 125

Thr Cys Gly Ser Val Thr Gln
 130 135

<210> 31
 <211> 142
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Chimeric Protein

<400> 31

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
 1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
 35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
 100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
115 120 125

Gln Arg Ser Asp Tyr Lys Phe Tyr Asp Asn Lys Arg Ile Asp
130 135 140

<210> 32
<211> 147
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric Protein

<400> 32

Met Gly Ser Ser His His His His Ser Ser Gly Leu Val Pro Arg
1 5 10 15

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
20 25 30

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Thr Ser Thr Asn
35 40 45

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
50 55 60

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
65 70 75 80

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
85 90 95

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
115 120 125

Gln Leu Val Arg Ser Asp Tyr Lys Phe Tyr Asp Asn Lys Arg Ile Asp
130 135 140

Arg His Thr
145

<210> 33
<211> 447
<212> DNA
<213> H. influenzae

<220>
 <221> CDS
 <222> (1)...(447)

<400> 33
 atg aaa tta aca aca cag caa acc ttg aaa aaa ggg ttt aca tta ata 48
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
 Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

att ccc tct tat caa aat tat act aaa aaa gca gcg gta tct gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

ctg caa gcg tca gcg cct tat aag gct gat gtg gaa tta tgt gta tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

agc aca aat gaa aca aca aac tgt acg ggt gga aaa aat ggt att gca 240
 Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

gca gat ata acc aca gca aaa ggc tat gta aaa tca gtg aca aca agc 288
 Ala Asp Ile Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
 85 90 95

aac ggt gca ata aca gta aaa ggg gat ggc aca ttg gca aat atg gaa 336
 Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
 100 105 110

tat att ttg caa gct aca ggt aat gct gca aca ggt gta act tgg aca 384
 Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

aca act tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc 432
 Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

gga agt gtc aca caa 447
 Gly Ser Val Thr Gln
 145

<210> 34
 <211> 149
 <212> PRT
 <213> H. influenzae

<400> 34
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu

35

40

45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
 85 90 95

Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
 100 105 110

Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

Gly Ser Val Thr Gln
 145

<210> 35
 <211> 447
 <212> DNA
 <213> H. influenzae

<220>
 <221> CDS
 <222> (1)..(447)

<400> 35
 atg aaa tta aca aca cag caa acc ttg aaa aaa ggg ttt aca tta ata 48
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
 Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

att ccc tct tat caa aat tat act aaa aaa gca gcg gta tct gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

ctg caa gcg tca gcg cct tat aag gct gat gtg gaa tta tgt gta tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

agc aca aat gaa aca aca aac tgt acg ggt gga aaa aat ggt att gca 240
 Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

gca gat ata acc aca gca aaa ggc tat gta aaa tca gtg aca aca agc 288

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser		
85	90	95
aac ggt gca ata aca gta aaa ggg gat ggc aca ttg gca aat atg gaa		336
Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu		
100	105	110
tat att ttg caa gct aca ggt aat gct gca aca ggt gta act tgg aca		384
Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr		
115	120	125
aca act tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc		432
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys		
130	135	140
gga agt gtc aca caa		447
Gly Ser Val Thr Gln		
145		
<210> 36		
<211> 149		
<212> PRT		
<213> H. influenzae		
<400> 36		
Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile		
1	5	10
15		
Glu Leu Met Ile Val Ile Ala Ile Ala Ile Leu Ala Thr Ile Ala		
20	25	30
Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu		
35	40	45
Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr		
50	55	60
Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala		
65	70	75
80		
Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser		
85	90	95
Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu		
100	105	110
Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr		
115	120	125
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys		
130	135	140

Gly Ser Val Thr Gln
145

<210> 37
<211> 447
<212> DNA
<213> H. influenzae

<220>
<221> CDS
<222> (1)..(447)

<400> 37
atg aaa tta aca aca cag caa acc ttg aaa aaa ggg ttt aca tta ata 48
Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
1 5 10 15
gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
20 25 30
att ccc tct tat caa aat tat act aaa aaa gca gcg gta tct gaa tta 144
Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
35 40 45
ctg caa gcg tca gcg cct tat aag gct gat gtg gaa tta tgt gta tat 192
Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
50 55 60
agc aca aat gaa aca aca aac tgt acg ggt gga aaa aat ggt att gca 240
Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
65 70 75 80
gca gat ata acc aca gca aaa ggc tat gta aaa tca gtg aca aca agc 288
Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
85 90 95
aac ggt gca ata aca gta aaa ggg gat ggc aca ttg gca aat atg gaa 336
Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
100 105 110
tat att ttg caa gct aca ggt aat gct gca aca ggt gta act tgg aca 384
Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
115 120 125
aca act tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc 432
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
130 135 140
gga agt gtc aca caa 447
Gly Ser Val Thr Gln
145

<210> 38
<211> 149
<212> PRT
<213> H. influenzae

<400> 38

Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
 85 90 95

Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
 100 105 110

Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

Gly Ser Val Thr Gln
 145

<210> 39
 <211> 447
 <212> DNA
 <213> H. influenzae

<220>
 <221> CDS
 <222> (1)..(447)

<400> 39
 atg aaa tta aca aca cag caa acc ttg aaa aaa ggg ttt aca tta ata 48
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15
 gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
 Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30
 att ccc tct tat caa aat tat act aaa aaa gca gcg gta tct gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

ctg caa gcg tca gcg cct tat aag gct gat gtg gaa tta tgt gta tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

agc aca aat gaa aca aca aac tgt acg ggt gga aaa aat ggt att gca 240
 Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

gca gat ata acc aca gca aaa ggc tat gta aaa tca gtg aca aca agc 288
 Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
 85 90 95

aac ggt gca ata aca gta aaa ggg gat ggc aca ttg gca aat atg gaa 336
 Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
 100 105 110

tat att ttg caa gct aca ggt aat gct gca aca ggt gta act tgg aca 384
 Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

aca act tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc 432
 Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

gga agt gtc aca caa 447
 Gly Ser Val Thr Gln
 145

<210> 40

<211> 149

<212> PRT

<213> H. influenzae

<400> 40

Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
 85 90 95

Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
 100 105 110

Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

Thr	Thr	Cys	Lys	Gly	Thr	Asp	Ala	Ser	Leu	Phe	Pro	Ala	Asn	Phe	Cys
130					135						140				

Gly Ser Val Thr Gln
145

<210> 41
<211> 447
<212> DNA
<213> *H. influenzae*

<220>
<221> CDS
<222> (1) .. (447)

<400> 41
atg aaa tta aca aca cag caa acc ttg aaa aaa ggt ttt aca tta atc 48
Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
1 5 10 15

```

gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
20          25          30

```

```

att cct tct tat caa aat tat acc aaa aaa gct tcg gta tcc gaa tta      144
Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ser Val Ser Glu Leu
            35          40          45

```

```

ctg caa gca tct gca cct tat aag gct gat gtg gaa tta tgt gta tat 192
Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
      50          55          60

```

```

agc aca aat gaa aca aca aac tgt acg ggt gga aaa aat ggt att gca 240
Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
65 70 75 80

```

```

gct gat att aca aca gca aaa ggc tat gta gcc tca gtg aaa act caa      288
Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
          85          90          95

```

```

tcg ggt ggc att aca gta aaa ggg aat ggc aca ttg gca aat atg gaa 336
Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu
          100          105          110

```

tat att ttg caa gct aaa ggt aat gct aca gca ggt gta act tgg aca 384
 Tyr Ile Leu Gln Ala Lys Gly Asn Ala Thr Ala Gly Val Thr Trp Thr
 115 120 125

```

aca acc tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc 432
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
130 135 140

```

aga agt gtc aca aaa 447
Arg Ser Val Thr Lys
145

<210> 42
 <211> 149
 <212> PRT
 <213> H. influenzae

 <400> 42

Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ser Val Ser Glu Leu
 35 40 45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

Ser Thr Asn Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala
 65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
 85 90 95

Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu
 100 105 110

Tyr Ile Leu Gln Ala Lys Gly Asn Ala Thr Ala Gly Val Thr Trp Thr
 115 120 125

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

Arg Ser Val Thr Lys
 145

<210> 43
 <211> 447
 <212> DNA
 <213> H. influenzae

<220>
 <221> CDS
 <222> (1)..(447)

<400> 43
 atg aaa tta aca aca cag caa acc acc ttg aaa aaa ggg ttt aca tta ata 48
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

gag cta atg att gtg att gca att att gct att tta gcc acc atc gca 96

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala			
20	25	30	
att cct tct tat aaa aat tat act aaa aaa gca gcg gta tct gaa tta			144
Ile Pro Ser Tyr Lys Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu			
35	40	45	
ctg caa gct tct gcg cct tat aag gct gat gtg gaa tta tgt gta tat			192
Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr			
50	55	60	
agc aca aat gaa ata aca aat tgt atg ggt gga aaa aat ggt att gca			240
Ser Thr Asn Glu Ile Thr Asn Cys Met Gly Gly Lys Asn Gly Ile Ala			
65	70	75	80
gct gat att aca aca gca aaa ggc tat gta gcc tca gtg aaa act caa			288
Ala Asp Ile Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln			
85	90	95	
tcg ggt ggc att acc gta aaa ggg gat ggc aca ttg gca aat atg gaa			336
Ser Gly Gly Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu			
100	105	110	
tat att ttg caa gct aca ggt aat gct gca gca ggt gta act tgg aca			384
Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Gly Val Thr Trp Thr			
115	120	125	
aca acc tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc			432
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys			
130	135	140	
gga agt atc aca caa			447
Gly Ser Ile Thr Gln			
145			

<210> 44
 <211> 149
 <212> PRT
 <213> H. influenzae

<400> 44

Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile

1

5

10

15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala

20

25

30

Ile Pro Ser Tyr Lys Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu

35

40

45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr

50

55

60

Ser Thr Asn Glu Ile Thr Asn Cys Met Gly Gly Lys Asn Gly Ile Ala

65

70

75

80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
 85 90 95

Ser Gly Gly Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
100 . 105 110

Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Ala Gly Val Thr Trp Thr
 115 120 125

Gly Ser Ile Thr Gln
145

<210> 45
<211> 447
<212> DNA
<213> *H. influenzae*

<220>
<221> CDS
<222> (1) .. (447)

<400> 45
atg aaa tta aca aca ctg caa acc ttg aaa aaa ggg ttt aca tta atc
Met Lys Leu Thr Thr Leu Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
1 5 10 15

```

gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
          20          25          30

```

att cct tct tat caa aat tat acc aaa aaa gct gcg gta tcc gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

ctg caa gct tct gcg cct tat aag gct gat gtg gaa tta tgc gtt tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Ile Cys Val Tyr
 50 55 60

```

agc aca ggc aaa cct tct act tgc tca gga gga agc aat gga att gca 240
Ser Thr Gly Lys Pro Ser Thr Cys Ser Gly Gly Ser Asn Gly Ile Ala
65           70           75           80

```

```

gct gat att acg aca gca aaa ggc tat gta gcc tca gtg aaa act caa      288
Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
85          90          95

```

tca ggt ggt att aca gta aaa ggg aat ggc aca ttg gca aat atg gaa	336
Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu	
100 105 110	

tat att ttg caa gct aca ggt aat gct gca aca ggt gta act tgg aca 384
 Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 . 125

```

aca act tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc      432
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
130          135          140

```

gga agt gtc aca caa 447
Gly Ser Val Thr Gln
145

<210> 46
<211> 149
<212> PRT
<213> H. influenzae

<400> 46

Met Lys Leu Thr Thr Leu Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
20 25 . 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
50 55 60

Ser Thr Gly Lys Pro Ser Thr Cys Ser Gly Gly Ser Asn Gly Ile Ala
65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
 85 90 95

Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu
100 105 110

Tyr Ile Leu Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
130 135 140

Gly Ser Val Thr Gln
145

<210> 47
<211> 447
<212> DNA
<213> *H. influenzae*

5220>

<221> CDS
 <222> (1)..(447)

<400> 47
 atg aaa tta aca aca ctg caa acc ttg aaa aaa ggg ttt aca tta atc 48
 Met Lys Leu Thr Thr Leu Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15
 gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
 Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30
 att cct tct tat caa aat tat acc aaa aaa gct gcg gta tcc gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45
 ctg caa gct tct gcg cct tat aag gct gat gtg gaa tta tgc gtt tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60
 agc aca ggc aaa ctt tct act tgc tca gga gga agc aat gga att gca 240
 Ser Thr Gly Lys Leu Ser Thr Cys Ser Gly Gly Ser Asn Gly Ile Ala
 65 70 75 80
 gct gat att aca aca gca aaa ggc tat gta gcc tca gtg aaa act caa 288
 Ala Asp Ile Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
 85 90 95
 tca ggt ggt att aca gta aaa ggg aat ggc aca ttg gca aat atg gaa 336
 Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu
 100 105 110
 tat att ttg caa gct aaa ggt aat gct aca gca ggt gta act tgg aca 384
 Tyr Ile Leu Gln Ala Lys Gly Asn Ala Thr Ala Gly Val Thr Trp Thr
 115 120 125
 aca acc tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc 432
 Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140
 gga agt gtc aca aaa 447
 Gly Ser Val Thr Lys
 145

<210> 48
 <211> 149
 <212> PRT
 <213> H. influenzae
 <400> 48

Met Lys Leu Thr Thr Leu Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

Ser Thr Gly Lys Leu Ser Thr Cys Ser Gly Gly Ser Asn Gly Ile Ala
 65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
 85 90 95

Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu
 100 105 110

Tyr Ile Leu Gln Ala Lys Gly Asn Ala Thr Ala Gly Val Thr Trp Thr
 115 120 125

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

Gly Ser Val Thr Lys
 145

<210> 49
 <211> 447
 <212> DNA
 <213> H. influenzae

<220>
 <221> CDS
 <222> (1)...(447)

<400> 49
 atg aaa tta aca aca cag caa acc ttg aaa aaa ggt ttt aca tta atc 48
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
 Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

att cct tct tat caa aat tat acc aaa aaa gct tcg gta tcc gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ser Val Ser Glu Leu
 35 40 45

ctg caa gct tcc gca cct tat aag tca gat gtg gaa tta tgc gtt tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ser Asp Val Glu Leu Cys Val Tyr
 50 55 60

agc aca ggc aaa cct tct act tgc tca gga gga agc aat gga att gca 240
 Ser Thr Gly Lys Pro Ser Thr Cys Ser Gly Gly Ser Asn Gly Ile Ala
 65 70 75 80

gct gat att aca aca gca aaa ggc tat gta gcc tca gtg aaa act caa 288
 Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln
 85 90 95

tca ggt ggt att aca gta aaa ggg aat ggc aca ttg gca aat atg gaa	336
Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu	
100 105 110	
tat att ttg caa gct aaa ggt aat gct aca gca ggt gta act tgg aca	384
Tyr Ile Leu Gln Ala Lys Gly Asn Ala Thr Ala Gly Val Thr Trp Thr	
115 120 125	
aca acc tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc	432
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys	
130 135 140	
aga agt gtc aca aaa	447
Arg Ser Val Thr Lys	
145	
<210> 50	
<211> 149	
<212> PRT	
<213> H. influenzae	
<400> 50	
Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile	
1 5 10 15	
Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala	
20 25 30	
Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ser Val Ser Glu Leu	
35 40 45	
Leu Gln Ala Ser Ala Pro Tyr Lys Ser Asp Val Glu Leu Cys Val Tyr	
50 55 60	
Ser Thr Gly Lys Pro Ser Thr Cys Ser Gly Gly Ser Asn Gly Ile Ala	
65 70 75 80	
Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Ala Ser Val Lys Thr Gln	
85 90 95	
Ser Gly Gly Ile Thr Val Lys Gly Asn Gly Thr Leu Ala Asn Met Glu	
100 105 110	
Tyr Ile Leu Gln Ala Lys Gly Asn Ala Thr Ala Gly Val Thr Trp Thr	
115 120 125	
Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys	
130 135 140	
Arg Ser Val Thr Lys	
145	

<210> 51
<211> 447
<212> DNA
<213> *H. influenzae*

<220>
<221> CDS
<222> (1) .. (447)

<400> 51
 atg aaa tta aca aca cag caa acc ttg aaa aaa ggg ttt aca tta ata 48
 Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
 1 5 10 15

 gag cta atg att gtg att gca att att gct att tta gcc act atc gca 96
 Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
 20 25 30

 att ccc tct tat caa aat tat act aaa aaa gcg gcg gta tct gaa tta 144
 Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
 35 40 45

 ctg caa gcg tca gcg cct tat aag gct gat gtg gaa tta tgt gta tat 192
 Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
 50 55 60

 agt aca ggt aaa cct tcc agt tgc tcg gga gga agc aat gga att gcg 240
 Ser Thr Gly Lys Pro Ser Ser Cys Ser Gly Gly Ser Asn Gly Ile Ala
 65 70 75 80

 gct gat att aca aca gca aaa ggc tat gta aaa tca gtg aca aca agc 288
 Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
 85 90 95

 aac ggt gca ata aca gta aaa ggg gat ggc aca ttg gca aat atg gaa 336
 Asn. Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
 100 105 110

 tat att ttg caa gcc agt ggt aat gct gca aca ggt gta act tgg aca 384
 Tyr Ile Leu Gln Ala Ser Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
 115 120 125

 aca act tgc aaa gga acg gat gcc tct tta ttt cca gca aat ttt tgc 432
 Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
 130 135 140

 gga agt gtc aca caa 447
 Gly Ser Val Thr Gln
 145

<210> 52
<211> 149
<212> PRT
<213> H. influenzae

<400> 52

Met Lys Leu Thr Thr Gln Gln Thr Leu Lys Lys Gly Phe Thr Leu Ile
1 5 10 15

Glu Leu Met Ile Val Ile Ala Ile Ile Ala Ile Leu Ala Thr Ile Ala
20 25 30

Ile Pro Ser Tyr Gln Asn Tyr Thr Lys Lys Ala Ala Val Ser Glu Leu
35 40 45

Leu Gln Ala Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr
50 55 60

Ser Thr Gly Lys Pro Ser Ser Cys Ser Gly Gly Ser Asn Gly Ile Ala
65 70 75 80

Ala Asp Ile Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser
85 90 95

Asn Gly Ala Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu
100 105 110

Tyr Ile Leu Gln Ala Ser Gly Asn Ala Ala Thr Gly Val Thr Trp Thr
115 120 125

Thr Thr Cys Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys
130 135 140

Gly Ser Val Thr Gln
145

<210> 53
<211> 40
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic Peptide

<400> 53

Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp His
1 5 10 15

Lys Lys Gly Pro Ser Leu Lys Leu Leu Ser Leu Ile Lys Gly Val Ile
20 25 30

Val His Arg Leu Glu Gly Val Glu
35 40

<210> 54
<211> 129
<212> PRT
<213> Artificial sequence

<220>

<223> Chimeric protein

<400> 54

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
1 5 10 15

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Leu Val Arg Ser Asp
20 25 30

Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg Asp His Lys Lys Gly
35 40 45

Arg His Thr Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr
50 55 60

Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile
65 70 75 80

Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln
85 90 95

Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys
100 105 110

Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr
115 120 125

Gln

<210> 55

<211> 126

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric protein

<400> 55

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
1 5 10 15

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr Ser Thr Asn
20 25 30

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
35 40 45

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala

50

55

60

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 65 70 75 80

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys
 85 90 95

Leu Val Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly Thr Arg
 100 105 110

Asp His Lys Lys Gly Arg His Thr Cys Gly Ser Val Thr Gln
 115 120 125

<210> 56

<211> 138

<212> PRT

<213> Artificial sequence

<220>

<223> Chimeric protein

<400> 56

Gly Ser His Met Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala
 1 5 10 15

Ser Ala Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr Ser Thr Asn
 20 25 30

Glu Thr Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile
 35 40 45

Thr Thr Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala
 50 55 60

Ile Thr Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu
 65 70 75 80

Gln Ala Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys
 85 90 95

Lys Gly Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val
 100 105 110

Thr Gln Leu Val Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala Asn Gly
 115 120 125

Thr Arg Asp His Lys Lys Gly Arg His Thr
 130 135

<210> 57
<211> 144
<212> PRT
<213> Artificial sequence

<220>
<223> Chimeric protein

<400> 57

Gly Ser His Met Leu Val Arg Ser Asp Tyr Lys Phe Tyr Glu Asp Ala
1 5 10 15

Asn Gly Thr Arg Asp His Lys Lys Gly Arg His Thr Gly Pro Ser Leu
20 25 30

Lys Leu Thr Lys Lys Ala Ala Val Ser Glu Leu Leu Gln Ala Ser Ala
35 40 45

Pro Tyr Lys Ala Asp Val Glu Leu Cys Val Tyr Ser Thr Asn Glu Thr
50 55 60

Thr Asn Cys Thr Gly Gly Lys Asn Gly Ile Ala Ala Asp Ile Thr Thr
65 70 75 80

Ala Lys Gly Tyr Val Lys Ser Val Thr Thr Ser Asn Gly Ala Ile Thr
85 90 95

Val Lys Gly Asp Gly Thr Leu Ala Asn Met Glu Tyr Ile Leu Gln Ala
100 105 110

Thr Gly Asn Ala Ala Thr Gly Val Thr Trp Thr Thr Cys Lys Gly
115 120 125

Thr Asp Ala Ser Leu Phe Pro Ala Asn Phe Cys Gly Ser Val Thr Gln
130 135 140