



(86) **Date de dépôt PCT/PCT Filing Date:** 2019/08/27
(87) **Date publication PCT/PCT Publication Date:** 2020/03/05
(45) **Date de délivrance/Issue Date:** 2023/05/16
(85) **Entrée phase nationale/National Entry:** 2021/02/05
(86) **N° demande PCT/PCT Application No.:** KR 2019/010894
(87) **N° publication PCT/PCT Publication No.:** 2020/045941
(30) **Priorités/Priorities:** 2018/08/27 (KR10-2018-0100359);
2019/08/26 (KR10-2019-0104641)


(51) **Cl.Int./Int.Cl.** **C07D 417/12** (2006.01),
A61K 31/4427 (2006.01), **A61K 31/4523** (2006.01),
A61K 31/496 (2006.01), **A61K 31/506** (2006.01),
A61P 35/00 (2006.01), **A61P 37/00** (2006.01),
C07D 417/14 (2006.01), **C07D 487/04** (2006.01)

(72) **Inventeurs/Inventors:**
KIM, WOL YOUNG, KR;
LEE, YEON IM, KR;
YOON, YOUN JUNG, KR;
PARK, JOON SEOK, KR;
EOM, DEOK KI, KR;
...

(73) **Propriétaire/Owner:**

(54) **Titre : NOUVEAU DERIVE D'AMINE HETEROCYCLIQUE ET COMPOSITION PHARMACEUTIQUE LE COMPRENANT**
(54) **Title: NOVEL HETEROCYCLIC AMINE DERIVATIVE AND PHARMACEUTICAL COMPOSITION COMPRISING SAME**

[Chemical Formula 1]



(57) **Abrégé/Abstract:**

The present disclosure relates to a novel heterocyclic amine derivative represented by the following Chemical Formula 1 and a pharmaceutical composition comprising the same, and the compound according to the present disclosure can be usefully used for the prevention or treatment of autoimmune diseases or cancers. (see Chemical formula 1) wherein, in Chemical Formula 1, R_1 , R_2 , X_1 , X_2 , L , Y , A and B are the same as defined in the specification.



(72) **Inventeurs(suite)/Inventors(continued):** BANG, KEUK CHAN, KR; JUNG, JAEHYUN, KR


(73) **Propriétaires(suite)/Owners(continued):** DAEWOONG PHARMACEUTICAL CO., LTD., KR

(74) **Agent:** RICHES, MCKENZIE & HERBERT LLP

ABSTRACT

The present disclosure relates to a novel heterocyclic amine derivative represented by the following Chemical Formula 1 and a pharmaceutical composition comprising the same, and the compound according to the present disclosure can be usefully used for the prevention or treatment of autoimmune diseases or cancers.

[Chemical Formula 1]



wherein, in Chemical Formula 1, R₁, R₂, X₁, X₂, L, Y, A and B are the same as defined in the specification.

【DESCRIPTION】

TITLE OF INVENTION

NOVEL HETEROCYCLIC AMINE DERIVATIVE AND
PHARMACEUTICAL COMPOSITION COMPRISING SAME

TECHNICAL FIELD

The present disclosure relates to a novel heterocyclic amine useful as BTK (Bruton's Tyrosine Kinase) inhibitor and a pharmaceutical composition comprising the same.

BACKGROUND OF ART

ITK(Interleukin-2 Tyrosine Kinase) and BTK(Bruton's Tyrosine Kinase) are a type of tyrosine kinase, together with Tec (tyrosine kinase expressed in hepatocellular carcinoma), RLK(Resting Lymphocyte Kinase) and BMX(Bone-Marrow tyrosine kinase gene on chromosome X), which that does not have a TEC family receptor and acts on various immune responses.

ITK is expressed not only in T cells but also in NK cells and mast cells, and plays an important role in T-cell proliferation and production of important cytokines such as IL-2, IL-4, IL-5, IL-10, IL-13 and IL-17 (Schaeffer et al. Nat. Immune 2001,2, 1183; Fowell et al. Immunity, 1999, 11, 399). T cells are activated by TCR signaling, and the activated T cells produce inflammatory cytokine and activate B cells and macrophages, causing autoimmune diseases such as RA (Sahu N. et al. Curr Top Med Chem. 2009, 9, 690). Previously, it was known that T cells are activated into

Th1 cells to induce RA diseases, but recently, it has been reported that not only Th17/Treg but also Th1 cells act as a pathogenesis of RA (J Leipe J. et al. Arthritis Rheum. 2010, 62, 2876). In addition, the ITK has been previously developed as an immunotherapeutic drug target such as asthma, but no ITK has been developed as a therapeutic for RA (Lo H. Y Expert Opin Ther Pat. 2010, 20, 459). Recently, however, it has been reported to regulate the development of Th17 and Treg cells via ITK^{-/-}mice, and it has ample potential as a therapeutic target for RA (Gomez-Rodriguez J. et al. J. Exp. Med. 2014, 211, 529).

In a study using the ITK inhibitor PRN694, a study on the reduction of TNF- α , which is a representative inflammatory cytokine of RA diseases, have been reported, confirming the possibility of development as a therapeutic agent by regulating Th17 expression via ITK inhibition (Zhong Y. et al. THE JOURNAL OF BIOLOGICAL CHEMISTRY 2015, 290, 5960).

BTK acts as a regulator of early B-cell development as well as of mature B-cell activation, signaling and survival. The B-cell is signaled by a B cell receptor (BCR) that recognizes an antigen attached to the surface of an antigen-presenting cell and is activated into a mature antibody-producing cell. However, aberrant signaling via BCR leads to abnormal B-cell proliferation and the formation of pathologic autoantibodies, and thereby can induce cancer, autoimmune and/or inflammatory diseases. Thus, in the abnormal B-cell proliferation, signaling via BCR may be blocked when BTK is deficient. Thus, inhibition of BTK can block B-cell mediated disease processes, and the use of BTK inhibitors may be a useful approach for the treatment of B-cell mediated diseases.

Furthermore, BTK can be expressed by other cells that may be associated with disease besides B-cells. For example, BTK is important components for Fc-gamma signaling in bone marrow cells, and is expressed by mast cells. Specifically, BTK-deficient bone marrow-induced mast cells exhibit impaired antigen-induced

degranulation, and inhibition of BTK activity is known to be useful for treating pathological mast cell responses such as allergy and asthma (Iwaki et al. J. Biol Chem. 2005 280: 40261). In addition, it is known that monocytes from XLA patients, in which BTK activity is absent, decreases in TNF alpha production following stimulation and thus TNF alpha-mediated inflammation could be inhibited by BTK inhibitors (see, Horwood et al., J. Exp. Med. 197: 1603, 2003).

At present, there has been no case where it has been developed as a substance that dually inhibits BTK and ITK. However, as the BTK inhibitor, WO 2008/039218 discloses 4-aminopyrazolo[3,4-d]pyrimidinylpiperidine derivatives, and WO2015/061247 discloses hetero compounds such as pyridine, pyrimidine, pyrazine and pyridazine, and WO2014/055934 discloses pyrimidinyl phenyl acrylamide derivatives. As the ITK inhibitor, WO2005/066335 discloses aminobenzimidazoles, WO2005/056785 discloses pyridones, WO2002/050071 discloses aminothiazole derivatives, and recently, WO2014/036016 discloses benzimidazole derivatives.

In view of the above, as a result of studying novel compounds, the present inventors has found that a compound having a chemical structure different from BTK, ITK inhibitors reported so far has excellent BTK and ITK dual-activity inhibitory effect, thereby completing the present disclosure. The compounds belonging to the present disclosure mainly have BTK and ITK inhibitory activity on their own, but do not exclude a possibility of exhibiting a pharmacological action as an efficacious agent by a special body environment or by products of metabolic process, after absorption into the body.

DETAILED DESCRIPTION OF THE INVENTION


TECHNICAL PROBLEM

It is one object of the present disclosure to provide a novel heterocyclic amine derivative useful as a BTK inhibitor, and a pharmaceutical composition comprising the same.

TECHNICAL SOLUTION

In order to achieve the above objects, there is provided a compound represented by the following Chemical Formula 1, or a pharmaceutically acceptable salt thereof:

[Chemical Formula 1]



wherein, in Chemical Formula 1,

B is a 5-membered or 6-membered heterocycle containing 1 to 3 heteroatoms each independently selected from the group consisting of N, O and S, provided that the 5-membered or 6-membered heterocycle includes at least one N,

R₁ is hydrogen, halogen, C₁₋₄ alkyl, or C₁₋₄ haloalkyl,

X₁ and X₂ are each independently N, or CR',

wherein R' is hydrogen or halogen,


L is a bond, C₁₋₄ alkylene, or -O-,


R₂ is cyano; C₁₋₄ alkyl; C₆₋₁₀ aryl; pyridinyl; morpholino; piperazinyl; or piperidinyl,

wherein, the piperazinyl and the piperidinyl are each independently, unsubstituted, or substituted with C₁₋₄ alkyl, C₁₋₄ alkyl substituted with cyano, C₁₋₄ alkyl substituted with amino, C₁₋₄ alkyl substituted with C₁₋₄ alkoxy, or -CO-(C₁₋₄ alkyl),

Y is a bond, -O-, -NH-, or -N(C₁₋₄ alkyl)-,

A is C₁₋₄ alkyl,





wherein R_3 is C_{1-4} alkyl, C_{1-4} haloalkyl, C_{2-4} alkenyl, C_{2-4} haloalkenyl, C_{2-4} alkynyl, or C_{2-4} haloalkynyl.

- 5 Preferably, B is thiazole, pyrazole, pyridine, or pyrimidine ring, and R_1 is hydrogen, chloro, methyl, or trifluoromethyl.

Preferably,

both X_1 and X_2 are CH;

- 10 one of X_1 and X_2 is CF and the other is CH; or
one of X_1 and X_2 is N and the other is CH.

More preferably,

both X_1 and X_2 are CH, or

X_1 is CH and X_2 is CF; or

- 15 X_1 is CH and X_2 is N; or
 X_1 is N and X_2 is CH.

Preferably, L is a bond, methylene, or -O-.

- 20 Preferably, R_2 is cyano; methyl; phenyl; pyridinyl; morpholino; piperazinyl substituted with methyl; piperazinyl substituted with ethyl; piperazinyl substituted with 2-cyanoethyl; piperazinyl substituted with 3-aminopropyl; piperazinyl substituted with 2-methoxyethyl; piperazinyl substituted with -CO-(methyl); unsubstituted piperidinyl; or piperidinyl substituted with methyl.


25

Preferably, Y is a bond, -O-, -NH-, or -N(methyl)-.

Preferably, R₃ is -CH₂CH₂Cl, -CH=CH₂, -CH=CHCH₃, -CH=CHCl, -C≡CH, or -C≡CCH₃.

5 Preferably, the compound represented by Chemical Formula 1 is represented by the following Chemical Formula 1-1:

[Chemical Formula 1-1]



wherein, in Chemical Formula 1-1,

10 R₁ is hydrogen, C₁₋₄ alkyl, or C₁₋₄ haloalkyl,

X₁ and X₂ are each independently N, or CR',

wherein R' is hydrogen or halogen,


L is a bond, C₁₋₄ alkylene, or -O-,


15 R₂ is cyano; C₁₋₄ alkyl; C₆₋₁₀ aryl; pyridinyl; morpholino; piperazinyl; or piperidinyl,

wherein the piperazinyl and the piperidinyl are each independently, unsubstituted, or substituted with C₁₋₄ alkyl, C₁₋₄ alkyl substituted with cyano, C₁₋₄ alkyl substituted with amino, C₁₋₄ alkyl substituted with C₁₋₄ alkoxy, or -CO-(C₁₋₄ alkyl),

Y is a bond, -O-, -NH-, or -N(C₁₋₄ alkyl)-,

20 A is C₁₋₄ alkyl,






wherein, R_3 is C_{1-4} haloalkyl, C_{2-4} alkenyl, C_{2-4} haloalkenyl, or C_{2-4} alkynyl.

- 5 Preferably, in Chemical Formula 1-1,
 R_1 is C_{1-4} alkyl,
 X_1 and X_2 are each independently N or CH,
 L is a bond, C_{1-4} alkylene, or -O-,
 R_2 is cyano; C_{1-4} alkyl; C_{6-10} aryl; pyridinyl; morpholino; piperazinyl; or
 10 piperidinyl,

wherein, the piperazinyl and the piperidinyl are each independently, unsubstituted, or substituted with C_{1-4} alkyl, C_{1-4} alkyl substituted with cyano, C_{1-4} alkyl substituted with C_{1-4} alkoxy, or -CO-(C_{1-4} alkyl),


Y is a bond, -O-, -NH-, or -N(C_{1-4} alkyl)-,



wherein, R_3 is C_{2-4} alkenyl, or C_{2-4} alkynyl.

- Also preferably, the compound represented by Chemical Formula 1 is
 20 represented by the following Chemical Formula 1-2:

[Chemical Formula 1-2]



wherein, in Chemical Formula 1-2,


R₁ is hydrogen, or halogen,

X₁ and X₂ are each independently N, or CH,

L is C₁₋₄ alkylene,

R₂ is morpholino,


Y is -NH-,



wherein, R₃ is C₂₋₄ alkenyl.

Further, preferably, the compound represented by Chemical Formula 1 is represented by the following Chemical Formula 1-3:

[Chemical Formula 1-3]




wherein, in Chemical Formula 1-3,

X₁ and X₂ are each independently N or CH,

L is C₁₋₄ alkylene,

R₂ is morpholino,


Y is -NH-,



wherein, R₃ is C₂₋₄ alkenyl.

Further, preferably, the compound represented by Chemical Formula 1 is
5 represented by the following Chemical Formula 1-4:

[Chemical Formula 1-4]




wherein, in Chemical Formula 1-4,

X₁ and X₂ are each independently N or CH,

10 L is C₁₋₄ alkylene,

R₂ is morpholino,

Y is -NH-,



wherein, R₃ is C₂₋₄ alkenyl.

In addition, the compounds of the present disclosure may exist in the form of salts, especially pharmaceutically acceptable salts. As salts, salts commonly used in the art, such as acid addition salts formed by pharmaceutically acceptable free acids can be used without limitation. The term "pharmaceutically acceptable salt" as used herein refers to any organic or inorganic addition salt of the compound represented by Chemical Formula 1, whose concentration is relatively non-toxic and harmless to a patient and activates effectively and whose side effects do not

degrade the beneficial efficacy of the above compound.

As the free acid, an organic acid and an inorganic acid can be used. Examples of the inorganic acids include hydrochloric acid, phosphoric acid, sulfuric acid, nitric acid, tartaric acid and the like. Examples of the organic acids include methanesulfonic acid, p-toluenesulfonic acid, acetic acid, trifluoroacetic acid, maleic acid, succinic acid, oxalic acid, benzoic acid, tartaric acid, fumaric acid, mandelic acid, propionic acid, citric acid, lactic acid, glycollic acid, gluconic acid, galacturonic acid, glutamic acid, glutaric acid, glucuronic acid, aspartic acid, ascorbic acid, carbonic acid, vanillic acid, hydroiodic acid and the like, but are not limited thereto. Preferably, the salt may be a hydrochloride salt.

Further, a pharmaceutically acceptable metal salt can be obtained by a conventional method using a base. For example, the compound represented by Chemical Formula 1 is dissolved in an excessive amount of an alkali metal hydroxide or an alkaline earth metal hydroxide solution, the non-soluble salt is filtered, and then the filtrate is evaporated and dried to obtain a pharmaceutically acceptable metal salt. At this time, it is particularly preferable to prepare a sodium salt, a potassium salt or a calcium salt as the metal salt.

In addition, a pharmaceutically unacceptable salt or solvate of the compound of Chemical Formula 1 may be used as an intermediate when preparing the compound of Chemical Formula 1, or the pharmaceutically acceptable salt or the solvate thereof.

Further, the compound represented by Chemical Formula 1 according to the present disclosure includes not only pharmaceutically acceptable salts thereof, but also solvates such as hydrates that can be prepared therefrom, and includes all possible stereoisomers, without being limited thereto. The solvate and the stereoisomer of the compound represented by Chemical Formula 1 may be prepared

from the compound of Chemical Formula 1 using common methods known in the art.

Further, the compound represented by Chemical Formula 1 according to the present disclosure may be prepared either in a crystalline form or in a non-crystalline form, and when the compound of Chemical Formula 1 is prepared in a crystalline form, it may be optionally hydrated or solvated. In the present disclosure, the compound represented by Chemical Formula 1 may not only include a stoichiometric hydrate, but also include a compound containing various amounts of water. The solvate of the compound represented by Chemical Formula 1 according to the present disclosure includes both stoichiometric solvates and non-stoichiometric solvates.

Representative examples of the compound represented by Chemical Formula 1 or a pharmaceutically acceptable salt thereof are as follows:

- 1) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 2) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,
- 3) 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 4) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 5) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 6) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,
- 7) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,
- 8) 1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,

- 9) (E)-1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)but-2-en-1-one,
- 10) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one,
- 5 11) 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,
- 12) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,
- 13) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 10 14) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)but-2-yn-1-one,
- 15) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,
- 15 16) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,
- 17) 1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 18) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 20 19) 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 20) (S)-1-(3-((4-((4-acetyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 25 21) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 22) (S)-1-(3-((4-((4-(2-methoxyethyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 23) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 30

- 24) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,
- 25) (R)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 5 26) (S)-1-(3-(methyl(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 27) N-(1-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)acrylamide,
- 28) 1-(6-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)octahydro-1H-pyrrolo[2,3-c]pyridin-1-yl)prop-2-en-1-one,
- 10 29) 1-(6-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)octahydro-1H-pyrrolo[2,3-c]pyridin-1-yl)but-2-yn-1-one,
- 30) 1-(6-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)-2-azaspiro[3.3]heptan-2-yl)prop-2-en-1-one,
- 15 31) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(piperidin-1-ylmethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 32) (S)-1-(3-((4-((4-ethylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 33) (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 20 34) (S)-3-(4-((2-((1-acryloylpiperidin-3-yl)amino)-6-((5-methylthiazol-2-yl)amino)pyridin-4-yl)methyl)piperazin-1-yl)propanenitrile,
- 35) (S)-1-(3-((4-methyl-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 25 36) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 37) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-2-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 38) (S)-2-((1-acryloylpiperidin-3-yl)amino)-6-(5-methylthiazol-2-yl)amino)isonicotinonitrile,
- 30

39) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-phenylpyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

40) (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-yn-1-one,

5 41) (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,

42) (S)-1-(3-((4-((4-ethylpiperazin-1-yl)methyl)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

10 43) (S)-1-(3-((4-((4-methylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

44) 5-methyl-N-(6-methyl-4-(morpholinomethyl)pyridin-2-yl)thiazol-2-amine,

45) (S)-3-chloro-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)propan-1-one,

15 46) (S,E)-3-chloro-1-(3-(6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,

47) (S)-1-(3-(4-((4-(3-aminopropyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,

20 48) (S)-1-(3-(6-(1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,

49) (S)-1-(3-(4-(morpholinomethyl)-6-(5-(trifluoromethyl)thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,

50) (S)-1-(3-(6-(5-chloro-1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,

25 51) (S)-1-(3-(4-(morpholinomethyl)-6-(thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,


52) (S)-1-(3-(3-fluoro-6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,

30 53) (S)-1-(3-(4-(morpholinomethyl)-6-(pyridin-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one, and


54) (S)-1-(3-(4-(morpholinomethyl)-6-(pyrimidin-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one.

In addition, according to the present disclosure, when A is not C₁₋₄ alkyl, the compound represented by Chemical Formula 1 may be prepared, for example, through Reaction Formula 1 below.

[Reaction Scheme 1]



In Reaction Scheme 1, A' is



10 A'' is and the rest are as defined above.

Step 1-1 is a step of reacting the compound represented by Chemical

Formula 1-1 and the compound represented by Chemical Formula 1-2 to prepare the compound represented by Chemical Formula 1-3. When the reaction is an amine substitution reaction, the reaction is preferably carried out in the presence of a palladium catalyst and a base, and when the reaction is a solvolysis reaction of an alkyl chloride due to a secondary alcohol, the reaction is preferably carried out in the presence of a base.

Step 1-2 is a step of reacting the compound represented by Chemical Formula 1-3 and the compound represented by Chemical Formula 1-4 to prepare the compound represented by Chemical Formula 1-5. The reaction is an amine substitution reaction, which is preferably carried out in the presence of a palladium catalyst and a base.

Step 1-3 is a step of removing a protecting group (BOC; tert-butyloxycarbonyl protecting group) from the compound represented by Chemical Formula 1-5 to prepare the compound represented by Chemical Formula 1-6. The reaction is preferably carried out under acidic conditions capable of removing the protecting group.


Step 1-4 is a step of reacting the compound represented by Chemical Formula 1-6 with the compound represented by Chemical Formula 1-7 to prepare the compound represented by Chemical Formula 1. The reaction is an amidation reaction, which is preferably carried out in the presence of a base.

Further, in Reaction Scheme 1, a reaction for protecting with a protecting group and a reaction for removing the protecting group may be added depending on each substituent.

According to another embodiment of the present disclosure, in the compound represented by Chemical Formula 1, when L is methylene and A is not C₁₋₄ alkyl, the

compound represented by Chemical Formula 1 may be prepared, for example, through Reaction Scheme 2 below.

[Reaction Scheme 2]



5 In Reaction Scheme 2, the definition of each substituent is the same as defined above.

Step 2-1 is a step of reacting the compound represented by Chemical Formula 2-1 and the compound represented by Chemical Formula 1-2 to prepare the compound represented by Chemical Formula 2-2. When the reaction is an amine substitution reaction, the reaction is preferably carried out in the presence of a palladium catalyst and a base, and when the reaction is a solvolysis reaction of an alkyl chloride due to a secondary alcohol, the reaction is preferably carried out in the presence of a base.

15 Step 2-2 is a step of reacting the compound represented by Chemical Formula 2-2 and the compound represented by Chemical Formula 2-3 to prepare the compound represented by Chemical Formula 2-4. The reaction is an amidation reaction, which is preferably carried out in the presence of an amidating reagent.

20 Step 2-3 is a step of reacting the compound represented by Chemical

Formula 2-4 and the compound represented by Chemical Formula 1-4 to prepare the compound represented by Chemical Formula 2-5. The reaction is an amine substitution reaction, which is preferably carried out in the presence of a palladium catalyst and a base.

5

Step 2-4 is a step of removing the protecting group (BOC) from the compound represented by Chemical Formula 2-5 and reducing ketone to prepare the compound represented by Chemical Formula 2-6. The reaction is preferably carried out under acidic conditions.

10

Step 2-5 is a step of reacting the compound represented by Chemical Formula 2-6 with the compound represented by Chemical Formula 1-7 to prepare the compound represented by Chemical Formula 1. The reaction is an amidation reaction, which is preferably carried out in the presence of a base.

15

Further, in Reaction Scheme 2, a reaction for protecting with a protecting group and a reaction for removing the protecting group may be added depending on each substituent.

20

In addition, according to the present disclosure, in the compound represented by Chemical Formula 1, the compound in which A is C₁₋₄ alkyl can be prepared using the same method as in step 1-1 and step 1-2 of Reaction Scheme 1, except that in step 1-1 of Reaction Scheme 1, the compound represented by $H-\overset{Y}{\curvearrowright}A$ is used instead of the compound represented by Chemical Formula 1-2.

25

The production method of each step described above can be more embodied in Examples described hereinafter.

According to a further embodiment of the present disclosure, there is provided a pharmaceutical composition for preventing or treating autoimmune diseases or

cancer diseases, which is effective for ITK and BTK inhibitory actions, comprising the compound represented by Chemical Formula 1, or a pharmaceutically acceptable salt, hydrate, solvate or isomer thereof as an active ingredient.

5 In this case, the autoimmune diseases include rheumatoid arthritis, systemic lupus erythematosus, childhood diabetes, psoriasis, aphthous stomatitis, chronic thyroiditis, acquired aplastic anemia, primary cirrhosis, ulcerative colitis, Behcet's disease, Crohn's disease, Silicosis, asbestosis, Sjogren's syndrome, Guillain-Barre syndrome, dermatomyositis, polymyositis, multiple sclerosis, autoimmune hemolytic
10 anemia, autoimmune encephalomyelitis, myasthenia gravis, Graves thyroid hyperplasia, nodular polyarteritis, ankylosing spondylitis, fibrositis, temporal arteritis, Wilson's disease, or Fanconi syndrome, and

the cancer may be blood cancer, extranodal marginal zone B-cell lymphoma, glioblastoma, lymphoplasmacytic lymphoma, acute myelogenous leukemia,
15 macroglobulinemia, B cell lymphoma, chronic lymphocytic leukemia, follicular lymphoma, non-hodgkin lymphoma, diffuse large B cell lymphoma, hairy cell leukemia, mantle cell lymphoma, glioblastoma, bladder cancer, pancreatic cancer, ovarian cancer, colorectal cancer, renal cancer, gastric cancer, transitional cell carcinoma, carcinoid tumor, breast cancer, non-small cell lung cancer, or multiple
20 myeloma.

As used herein, the term "prevention" refers to any act to delay or inhibit occurrence, spread or recurrence of the above-mentioned diseases by administration of the composition of the present disclosure, and "treatment" refers to
25 any act to improve or change the symptoms of the above diseases for the better by administration of the composition of the present disclosure.

The pharmaceutical composition according to the present disclosure can be formulated in types for oral or parenteral administrations according to a standard
30 pharmaceutical practice. These formulations may contain additives such as

pharmaceutically acceptable carrier, adjuvant or diluent in addition to the active ingredient.

Suitable carriers include, for example, physiological saline, polyethylene glycol, ethanol, vegetable oil, and isopropyl myristate and the like. Diluents include, for example, lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine and the like. but are not limited thereto. Further, the compounds of the present disclosure can be dissolved in oils, propylene glycol or other solvents commonly used in the preparation of injection solutions. Furthermore, the compounds of the present disclosure can be formulated in ointments or creams for topical application.

A preferred dose of the compound of the present disclosure may be varied according to the condition and weight of a patient, the severity of a disease, the type of a drug, and the route and duration of administration, but it may be suitably selected by those skilled in the art. In order to achieve the desirable effects, however, the compound of the present disclosure may be administered daily at a dose of 0.0001 to 100 mg/kg (body weight), and preferably 0.001 to 100 mg/kg (body weight). The administration may be performed once a day or in divided doses each day through an oral or parenteral route.

Depending on the method of administration, the pharmaceutical composition may contain the compound of the present disclosure in an amount of 0.001 to 99% by weight, preferably 0.01 to 60% by weight.

The pharmaceutical composition according to the present disclosure may be administered to mammals such as a rat, a mouse, a domestic animal, a human, through various routes. The administration may be carried out through all possible methods, for example, oral, rectal, intravenous, intramuscular, subcutaneous, intra-endometrial, intracerebroventricular injection.

ADVANTAGEOUS EFFECTS

The compound represented by Chemical Formula 1 according to the present disclosure or a pharmaceutically acceptable salt, hydrate, solvate or isomer thereof can be usefully used for the prevention or treatment of autoimmune diseases or

5 cancers.


DETAILED DESCRIPTION OF THE EMBODIMENTS

Below, the present disclosure will be described in more detail by way of examples. However, these examples are provided for illustrative purposes only, and

10 should not be construed as limiting the scope of the present disclosure to these examples.

Example 1: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one

15 **Step 1-1: Preparation of (2,6-dichloropyridin-4-yl)(morpholino)methanone**



2,6-Dichloroisonicotinic acid (10.0 g, 1.0 eq) was dissolved in dimethylformamide (100.0 mL), and then 1,1-carbonyldiimidazole (1.0 g, 1.2 eq) was

20 added thereto. The mixture was stirred at room temperature (25 ~ 30°C) for 1 hour under nitrogen gas, and then morpholine (5.4 mL, 1.2 eq) was added and stirred at the same temperature for 2 hours to complete the reaction. Ethyl acetate (200.0 mL) and water (200.0 mL) were added thereto, followed by extraction, and the aqueous layer was re-extracted three times using ethyl acetate (200.0 mL). The

25 ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:5) to give the title compound (13.0 g,


yield: 93.0%).

Step 1-2: Preparation of 4-((2,6-dichloropyridin-4-yl)methyl)morpholine



5 The intermediate (10.0 g, 1.0 eq) obtained in step 1-1 was dissolved in dichloromethane (100.0 mL), and then cooled to 0 to 10°C under nitrogen gas. 1M borane-tetrahydrofuran (115.0 mL, 3.0 eq) was slowly added dropwise. The mixture was stirred at room temperature for 12 hours to complete the reaction. The reaction solution was cooled to 0 to 10°C, and then 6N-hydrochloric acid aqueous solution
10 (256.0 mL, 20.0 eq) was slowly added dropwise, and then stirred at the same temperature for 1 hour. After adjusting the pH to 9~12 using 10N-sodium hydroxide aqueous solution, extraction was performed twice with dichloromethane. The dichloromethane layer was separated, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by
15 column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (8.1 g, yield: 90.0%).

Step 1-3: Preparation of tert-butyl 3-((6-chloro-4-(morpholinomethyl)pyridin-2-yl)amino)piperidine-1-carboxylate




20

After 1,4-dioxane (10.0 mL) was added to the intermediate (1.0 g, 1.0 eq) obtained in step 1-2 and dissolved, tris(dibenzylideneacetone)dipalladium(0) (465.8


mg, 0.2 eq) and Xantphos (1.5 g, 0.4 eq) were added thereto. Tert-butyl 3-aminopiperidine-1-carboxylate (780.0 μl , 1.0 eq) was added, and then sodium carbonate (1.3 g, 3.0 eq) was added, and the mixture was refluxed for 12 hours to complete the reaction. After cooling to 30°C or less, water (20.0 mL) and ethyl acetate (20.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate : hexane = 1:1) to give the title compound (900.0 mg, yield: 54.1%).

Step 1-4: Preparation of tert-butyl 3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidine-1-carboxylate




The intermediate (730.0 mg, 1.0 eq) obtained in step 1-3 was dissolved in 1,4-dioxane (14.0 mL). Palladium acetate (40.0 mg, 0.1 eq), Xantphos (204.7 mg, 0.2 eq), 5-methylthio-2-amine (203.6 mg, 1.0 eq), and cesium carbonate (1.7 g, 3.0 eq) were sequentially added. The mixture was reacted in a microwave reactor at 150°C for 30 minutes. Ethyl acetate (10.0 mL) and water (10.0 mL) were added, and the resulting solid was filtered to give the title compound (424.9 mg, yield 65.4%). After cooling to 30°C or less, water (15.0 mL) and ethyl acetate (15.0 mL) were added thereto, followed by layer separation. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (EA 100%) to give the title compound (564.0 mg, yield: 65.0%).

Step 1-5: Preparation of N²-(5-methylthiazol-2-yl)-4-(morpholinomethyl)-

N⁶-(piperidin-3-yl)pyridin-2,6-diamine

The intermediate (500.0 mg, 1.0 eq) obtained in step 1-4 was dissolved in dichloromethane (10.0 mL), and then cooled to 0 to 10°C. Trifluoroacetic acid (1.6 mL, 20.0 eq) was slowly added dropwise, and then stirred for 1 hour. After adjusting the pH to 9-12 using 12N-sodium hydroxide aqueous solution, the separated dichloromethane layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. Ethyl acetate (10.0 mL) was added to the resulting residue to form crystals for 30 minutes. The crystals were filtered and dried to give the title compound (357.5 mg, yield: 90.0%).


Step 1-6: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one

After the intermediate (350.0 mg, 1.0 eq) obtained in step 1-5 was dissolved in tetrahydrofuran (7.0 mL), water (7.0 mL) was added and sodium bicarbonate (226.8 mg, 3.0 eq) was added, and then cooled to 0 to 10°C. Acryloyl chloride (73.1 μ l, 1.0 eq) was slowly added dropwise, and then stirred for 30 minutes to complete the reaction. The layers were separated with dichloromethane, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting

residue was purified by column chromatography (dichloromethane: methanol = 15:1) to give the title compound (318.0 mg, yield: 80.0%).

¹H NMR(500 MHz, DMSO): 10.5(s, 1H), 6.94(s, 1H), 6.86-6.80(q, 1H), 6.50-6.49(d, 1H), 6.10-6.07(d, 1H), 6.04(s, 1H), 5.94(s, 1H), 5.66-5.64(d, 1H), 4.38-4.36(m, 1H), 4.18-4.16(m, 1H), 4.08-4.06(m, 1H), 3.55(m, 4H), 3.21(s, 3H), 2.88-2.83(m, 2H), 2.32(m, 4H), 2.28(s, 3H), 2.03(m, 2H), 1.30(m, 2H)


Example 2: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one



The title compound (15.0 mg, yield: 23.0%) was obtained in the same manner as in Example 1, except that in steps 1-3 of Example 1, tert-butyl 3-aminopyrrolidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.58-10.57(m, 1H), 6.94(s, 1H), 6.83-6.75(m, 1H), 6.63-6.48(m, 1H), 6.14-6.10(m, 1H), 6.08(s, 1H), 5.99-5.98(m, 1H), 5.67-5.59(m, 1H), 4.65-4.50(m, 1H), 3.99-3.97(m, 0.5H), 3.70-3.66(m, 1.5H), 3.55(m, 4H), 3.48(m, 1H), 3.35(m, 1H), 3.22(s, 2H), 2.34-2.32(m, 4H), 2.26-2.24(d, 3H), 2.25(m, 1H), 1.95-1.85(m, 1H)


Example 3: Preparation of 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (8.0 mg, yield: 53.0%) was obtained in the same manner as in Example 1, except that in step 1-3 of Example 1, tert-butyl 4-aminopiperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

5 ¹H NMR(500 MHz, DMSO): 10.54(s, 1H), 6.94(s, 1H), 6.93-6.83(m, 1H), 6.51-6.49(d, 1H), 6.10(d, 1H), 6.07(s, 1H), 5.94(s, 1H), 5.66-5.64(d, 1H), 4.38-4.36(m, 1H), 4.16(m, 1H), 4.08-4.02(m, 1H), 3.56(m, 4H), 3.21(s, 2H), 2.85(m, 1H), 2.61(m, 1H), 2.34-2.33(m, 4H), 2.28(s, 3H), 2.0(m, 2H), 1.30-1.21(m, 2H)

10 **Example 4: Preparation of (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one**




15 The title compound (10.0 mg, yield: 53.0%) was obtained in the same manner as in Example 1, except that in steps 1-3 of Example 1, tert-butyl (R)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

20 ¹H NMR(500 MHz, DMSO): 10.55-10.50(m, 1H), 6.91-6.90(m, 1H), 6.90-6.78(m, 0.5H), 6.47-6.56(m, 1.5H), 6.06-5.96(m, 3H), 5.65-5.67(m, 0.5H), 5.42-5.40(m, 0.5H), 4.42-4.40(m, 0.5H), 4.10-4.0(m, 1H), 3.90-3.87(m, 1.5H), 3.56(m, 4H), 3.20(s, 2H), 3.14-3.10(m, 1H), 2.68-2.63(m, 0.5H), 2.32(m, 4H), 2.19(s, 3H), 1.90-

2.0(m, 1H), 1.80(m, 1H), 1.50-1.40(m, 2.5H)


Example 5: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (13.0 mg, yield: 63.0%) was obtained in the same manner as in Example 1, except that in steps 1-3 of Example 1, tert-butyl (S)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.57(m, 1H), 6.91-6.90(m, 1H), 6.80-6.85(m, 0.5H), 6.70-6.40(m, 1.5H), 6.10-5.96(m, 3H), 5.65-5.63(d, 0.5H), 5.42-5.40(d, 0.5H), 4.42-4.40(m, 0.5H), 4.10-4.0(m, 1H), 3.90-3.87(m, 1.5H), 3.56(m, 4H), 3.20(s, 2H), 3.14-3.10(m, 1H), 2.68-2.63(m, 0.5H), 2.32(m, 4H), 2.19(s, 3H), 1.90-2.0(m, 1H), 1.80(m, 1H), 1.50-1.40(m, 2.5H)

Example 6: Preparation of (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one




The title compound (10.0 mg, yield: 58.0%) was obtained in the same manner as in Example 1, except that in step 1-3 of Example 1, tert-butyl (R) 3-

aminopyrrolidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

1H NMR(500 MHz, DMSO): 10.7(m, 1H), 6.94(s, 1H), 6.83-6.75(m, 1H), 6.63-6.47(m, 1H), 6.14-6.10(m, 2H), 6.09-3.08(m, 1H), 5.67-5.59(m, 1H), 4.67-4.50(1H), 3.97-3.96(m, 0.5H), 3.70(m, 1.5H), 3.55-3.54(m, 4H), 3.40(m, 1H), 3.38(m, 1H), 3.22(s, 2H), 2.32(m, 4H), 2.26-2.24(d, 3H), 2.20(m, 1H), 1.95-1.80(m, 1H)

Example 7: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one




The title compound (15.0 mg, yield: 63.0%) was obtained in the same manner as in Example 1, except that in step 1-3 of Example 1, tert-butyl (S) 3-aminopyrrolidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

1H NMR(500 MHz, DMSO): 10.7(m, 1H), 6.94(s, 1H), 6.83-6.75(m, 1H), 6.63-6.47(m, 1H), 6.14-6.10(m, 2H), 6.09-3.08(m, 1H), 5.67-5.59(m, 1H), 4.67-4.50(1H), 3.97-3.96(m, 0.5H), 3.70(m, 1.5H), 3.55-3.54(m, 4H), 3.40(m, 1H), 3.38(m, 1H), 3.22(s, 2H), 2.32(m, 4H), 2.26-2.24(d, 3H), 2.20(m, 1H), 1.95-1.80(m, 1H)


Example 8: Preparation of 1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one

Step 8-1: Preparation of methyl 6-((1-(tert-butoxycarbonyl)pyrrolidin-3-yl)amino)-2-chloropyrimidine-4-carboxylate




After methyl 2,4-dichloropyrimidine-6-carboxylate (500 mg, 1.0 eq) was dissolved in tetrahydrofuran (10.0 mL), diisopropylethylamine (1.5 eq) and tert-butyl 3-aminopyrrolidine-1-carboxylate (1.5 eq) were added thereto and then stirred at 80°C for 1 hour. Upon completion of the reaction, the mixture was cooled to 30°C or less, water (100.0 mL) and dichloromethane (100.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (640.0 mg, yield: 74.0%).

Step 8-2: Preparation of tert-butyl 3-((2-chloro-6-(morpholin-4-carbonyl)pyrimidin-4-yl)amino)pyrrolidine-1-carboxylate




After methyl 6-((1-tert-butoxycarbonyl)pyrrolidin-3-yl)amino)-2-chloropyrimidine-4-carboxylate (640.0 mg, 1.0 eq) obtained in step 8-1 was dissolved in tetrahydrofuran (10.0 mL), 1,5,7-triazabicyclo[4,4,0]dec-5-ene (0.3 eq) and morpholine (1.2 eq) were added thereto, and stirred at 60°C for 3 hours. Upon completion of the reaction, water (200.0 mL) and dichloromethane (200.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over sodium hydroxide and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 9:1) to give the title compound (470.0 mg, yield: 63.7%).

Step 8-3: Preparation of tert-butyl 3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholine-4-carbonyl)pyrimidin-4-yl)amino)pyrrolidine-1-carboxylate



After tert-butyl 3-((2-chloro-6-(morpholin-4-carbonyl)pyrimidin-4-yl)amino)pyrrolidine-1-carboxylate (450.0 mg, 1.0 eq) obtained in step 8-2 was dissolved in 1,4-dioxane (10.0 mL), palladium acetate (0.1 eq), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.2 eq), cesium carbonate (3.0 eq), 2-amino-5-methylthiazole (1.2 eq) were added thereto, and reacted in a microwave reactor (160°C, 30 min). Upon completion of the reaction, water (200.0 mL) and ethyl acetate (200.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 9:1) to give the title compound (410.0 mg, yield: 76.6%).


Step 8-4: Preparation of N²-(5-methylthiazol-2-yl)-6-(morpholinomethyl)-N⁴-(pyrrolidin-3-yl)pyrimidine-2,4-diamine



After tert-butyl 3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholin-4-carbonyl)pyrimidin-4-yl)amino)pyrrolidine-1-carboxylate (250.0 mg, 1.0 eq) obtained

in step 8-3 was dissolved in tetrahydrofuran (10.0 mL), 0.9M borane-tetrahydrofuran solution (5.0 eq) was added thereto, and stirred at 50°C for 5 hours. The reaction solution was cooled to 0°C, and then 6N aqueous hydrochloric acid solution (5.0 eq) was added and then stirred at 50°C for 12 hours. The reaction solution was again cooled to 0°C, and then the pH was adjusted to 12 using 12N sodium hydroxide aqueous solution, and then extracted with dichloromethane (200.0 mL) and water (200.0 mL). The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give the title compound (40.0 mg, yield: 20.9%).

Step 8-5: Preparation of 1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one




After N^2 -(5-methylthiazol-2-yl)-6-(morpholinomethyl)- N^4 -(pyrrolidin-3-

yl)pyrimidine-2,4-diamine (50.0 mg, 1.0 eq) obtained in step 8-4 was dissolved in tetrahydrofuran (1.6 mL) and water (0.4 mL), sodium bicarbonate (3.0 eq) and acryloyl chloride (1.1 eq) were added thereto, and stirred at 0°C for 30 minutes. Upon completion of the reaction, water (50.0 mL) and ethyl acetate (50.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 3:1) to give the title compound (7.0 mg, yield: 12.2%).

^1H NMR(500 MHz, CDCl_3): 6.95(s, 1H), 6.40-6.46(m, 3H), 6.18(s, 1H), 5.69-5.74(m, 1H), 3.91-4.10(m, 1H), 3.76(s, 2H), 3.50-3.75(m, 8H), 2.46-2.56(m, 6H), 2.36(s, 3H)

Example 9: Preparation of (E)-1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)but-2-en-1-one




5 The title compound (5.0 mg, yield: 8.5%) was obtained in the same manner as in Example 8, except that in step 8-5 of Example 8, crotonyl chloride was used instead of acryloyl chloride.

¹H NMR(500 MHz, CDCl₃): 6.94-7.00(m, 3H), 6.18(s, 1H), 6.13-6.16(m, 2H), 4.3(s, 1H), 3.75(s, 2H), 3.61-3.73(m, 8H), 2.56(s, 4H), 2.29-2.36(m, 5H), 1.86-1.91(m, 3H)

Example 10: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one


Step 10-1: Preparation of tert-butyl 3-((6-chloro-4-(morpholinomethyl)pyridin-2-yl)oxy)pyrrolidine-1-carboxylate



After tert-butyl 3-hydroxypyrrolidine-1-carboxylate (2.0 g, 1.0 eq) was dissolved in dimethylformamide (10.0 ml), potassium t-butoxide (1.4 g, 1.5 eq) was added thereto and stirred for 30 minutes. The intermediate (2.0 g, 1.0 eq) obtained in step 1-2 of Example 1 was added, and then the mixture was stirred at 60 to 80°C for 4 hours. After cooling to 30°C or less, water (40.0 mL) and ethyl acetate (40.0 mL) were added, and then the layers were separated. The ethyl acetate layer was

dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:5) to give the title compound (1.8 g, yield: 55.3%).


5 **Step 10-2: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one**



10 The title compound (13.2 mg, yield 65.5%) was obtained in the same manner as in Example 1, except that the intermediate obtained in step 10-1 was used instead of the intermediate obtained in step 1-3 of Example 1.

1H NMR(500 MHz, DMSO): 10.18(s, 1H), 6.62(m, 1H), 6.53(s, 1H), 6.04(m, 1H), 5.65(s, 1H), 5.58(m, 1H), 5.20(s, 1H), 4.44(s, 2H), 3.99(m, 1H), 3.71(m, 2H), 3.57(m, 4H), 3.50-3.49(m, 2H), 2.42(m, 4H), 2.04(m, 2H)

15 **Example 11: Preparation of 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one**




20 The title compound (8.5 mg, yield: 65.0%) was obtained in the same manner as in Example 10, except that in step 10-1 of Example 10, tert-butyl 4-hydroxypiperidine-1-carboxylate was used instead of tert-butyl 3-hydroxypyrrolidine-

1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.9(s, 1H), 7.00(s, 1H), 6.86-6.80(m, 1H), 6.53(s, 1H), 6.20(s, 1H), 6.11-6.08(d, 1H), 5.68-5.65(d, 1H), 5.36-5.32(m, 1H), 4.03(m, 1.5H), 3.90(m, 1.5H), 3.50(m, 4H), 3.45(m, 2H), 3.40(s, 2H), 2.34(m, 4H),
 5 2.31(s, 3H), 2.08-2.06(m, 2H), 1.61-1.59(m, 2H)

Example 12: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one




10 The title compound (9.5 mg, yield: 63.0%) was obtained in the same manner as in Example 10, except that in step 10-1 of Example 10, tert-butyl 3-hydroxypiperidine-1-carboxylate was used instead of tert-butyl 3-hydroxypyrrolidine-1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.994-10.92(m, 1H), 7.00-6.99(m, 1H), 6.95-
 15 6.85(m, 0.5H), 6.53(s, 1H), 6.60-6.5(m, 0.5H), 6.15-6.13(m, 1H), 6.10-5.96(m, 1H), 5.74(d, 0.5H), 5.43-5.45(d, 0.5H), 5.25-5.15(m, 1H), 4.01-3.95(m, 0.5H), 3.90-3.75(m, 2H), 3.70(m, 0.5H), 3.55(m, 4H), 3.40(s, 2H), 2.34(m, 4H), 2.37-2.20(s, 3H), 2.09-2.04(m, 1.5H), 1.97-1.78(m, 2.5H), 1.50(m, 1H)


20 **Example 13: Preparation of (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one**

Step 13-1: Preparation of (R)-N²-(5-methylthiazol-2-yl)-4-(morpholinomethyl)-N⁶-(piperidin-3-yl)pyridine-2,6-diamine



The title compound (150.0 mg, yield: 75.0%) was obtained in the same manner as in steps 1-3, 1-4, and 1-5 of Example 1, except that in steps 1-3 of Example 1, tert-butyl (R)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

Step 13-2: Preparation of (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one




After 2-butyric acid (21.6 mg, 1.0 eq) was dissolved in dimethylamide (1.0 mL), 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxidehexafluorophosphate (97.3 mg, 1.0 eq) was added thereto, and then stirred for 30 minutes. The intermediate (100.0 mg, 1.0 eq) obtained in step 13-1 was added and triethylamine (53.5 μ L, 1.5 eq) was added, and then stirred for 1 hour. Water (1.0 mL) and ethyl acetate (1.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 10:1) to give the title compound (64.0 mg, yield: 55.0%).

¹H NMR(500 MHz, DMSO): 10.59-10.58(d, 1H), 6.95-6.94(m, 1H), 6.84-

6.79(m, 1H), 6.10(m, 1H), 5.99-5.97(m, 1H), 4.57-4.56(m, 1H), 3.85-3.65(m, 2H), 3.55(m, 4H), 3.45-3.35(m, 4H), 3.20(s, 2H), 3.30(m, 4H), 2.26(s, 3H), 2.25-2.15(m, 2H), 2.0(d, 3H)


5 **Example 14: Preparation of 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)but-2-yn-1-one**



The title compound (58.6 mg, yield: 50.0%) was obtained in the same manner as in Example 13, except that in step 13-1 of Example 13, tert-butyl 3-aminopyrrolidine-1-carboxylate was used instead of tert-butyl (R)-3-aminopiperidine-1-carboxylate.

1H NMR(500 MHz, CDCl₃): 7.03(s, 1H), 6.11(s, 1H), 5.96(s, 1H), 4.56-4.54(d, 1H), 4.42-4.40(d, 1H), 3.73-3.71(m, 4H), 3.33(s, 2H), 3.30(m, 1H), 2.93(m, 1H), 2.45(m, 4H), 2.38(s, 3H), 1.47-1.40(m, 1H)

15 **Example 15: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one**




The title compound (496.0 mg, yield: 50.0%) was obtained in the same manner as in Example 10, except that in step 10-1 of Example 10, tert-butyl (S) 3-

hydroxypiperidine-1-carboxylate was used instead of tert-butyl 3-hydroxypyrrolidine-1-carboxylate.

1H NMR(500 MHz, DMSO): 10.95(m, 1H), 6.99-6.98(m, 1H), 6.75-6.85(m, 0.5H), 6.50(s, 1H), 6.4-6.5(m, 0.5H), 5.74-5.65(d, 1H), 6.45-6.43(d, 1H), 5.24-5.15(m, 1H), 4.02-4.00(m, 0.5H), 3.82-3.81(m, 2H), 3.78(m, 0.5H), 3.55(m, 4H), 3.50(m, 0.5H), 3.15-3.14(d, 2H), 2.32(m, 4H), 2.27-2.24(d, 3H), 2.06-1.96(m, 1.5H), 1.78-1.72(m, 2.5H), 1.51(m, 1H)


Example 16: Preparation of (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one



The title compound (15.0 mg, yield: 55.0%) was obtained in the same manner as in Example 10, except that in step 10-1 of Example 10, tert-butyl (R) 3-hydroxypiperidine-1-carboxylate was used instead of tert-butyl 3-hydroxypyrrolidine-1-carboxylate.

1H NMR(500 MHz, DMSO): 10.95(m, 1H), 6.99-6.98(m, 1H), 6.75-6.85(m, 0.5H), 6.50(s, 1H), 6.4-6.5(m, 0.5H), 5.74-5.65(d, 1H), 6.45-6.43(d, 1H), 5.24-5.15(m, 1H), 4.09-4.00(m, 0.5H), 3.82-3.81(m, 2H), 3.78(m, 0.5H), 3.56(m, 4H), 3.50(m, 0.5H), 3.15-3.14(d, 2H), 2.32(m, 4H), 2.27-2.25(d, 3H), 2.04(m, 1.5H), 1.87-1.72(m, 2.5H), 1.51(m, 1H)


Example 17: Preparation of 1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (10.0 mg, yield: 17.5%) was obtained in the same manner as in Example 8, except that in step 8-1 of Example 8, 3-amino-1-tert-butoxy-carbonylpiperidine was used instead of tert-butyl 3-aminopyrrolidine-1-carboxylate.

¹H NMR(500 MHz, CDCl₃): 7.25(s, 1H), 6.56-6.65(m, 1H), 6.25-6.39(m, 2H), 5.53(s, 0.5H), 5.22(s, 0.5H), 4.26(s, 1H), 4.09(s, 1H), 3.77(s, 4H), 3.72(s, 0.5H), 3.27-3.37(m, 4.5H), 2.54(s, 4H), 2.40(s, 3H), 2.14(s, 1H), 1.85(s, 1H), 1.66-1.67(m, 2H)


Example 18: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one



The title compound (12.1 mg, yield: 68.0%) was obtained in the same manner as in Example 13, except that in step 13-1 of Example 13, tert-butyl (S)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl (R)-3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.18(s, 1H), 6.53(s, 1H), 5.87(s, 1H), 5.85(s, 1H), 4.44(s, 2H), 3.71-3.47(m, 8H), 2.78(m, 1H), 2.42(m, 4H), 2.30(s, 3H), 1.87-1.58(m, 4H), 1.80(s, 3H)

Example 19: Preparation of 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one




The title compound (8.5 mg, yield: 53.0%) was obtained in the same manner as in Example 13, except that in step 13-1 of Example 13, tert-butyl 4-aminopiperidine-1-carboxylate was used instead of tert-butyl (R)-3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.18(s, 1H), 6.53(s, 1H), 5.87(s, 1H), 5.80(s, 1H), 4.44(s, 2H), 3.59-3.49(m, 8H), 2.68(m, 1H), 2.42(m, 4H), 2.30(s, 3H), 1.97-1.72(m, 4H), 1.80(s, 3H)


Example 20: Preparation of (S)-1-(3-((4-((4-acetylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one

Step 20-1: Preparation of 1-((2,6-dichloropyridin-4-yl)methyl)piperazine



The title compound (11.0 mg, yield: 86.0%) was obtained in the same manner as in steps 1-1 and 1-2 of Example 1, except that in step 1-1 of Example 1, tert-butyl piperazine-1-carboxylate was used instead of morpholine.


Step 20-2: Preparation of 1-(4-((2,6-dichloropyridin-4-yl)methyl)piperazin-1-yl)ethan-1-one



The intermediate (1.0 g, 1.0 eq) obtained in step 20-1 was dissolved in tetrahydrofuran (10.0 mL), and then triethylamine (1.1 mL, 2.0 eq) was added thereto. Acetyl chloride (434.6 μ l, 1.5 eq) was added thereto and stirred for 6 hours. After
 5 concentration, ethyl acetate (10.0 mL) and water (10.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 10:1) to give the title compound (1.0 g, yield: 85.5%).

10

Step 20-3: Preparation of (S)-1-(3-((4-((4-acetylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




The title compound (50.0 mg, yield: 45.0%) was obtained in the same
 15 manner as in Example 1, except that the intermediate obtained in step 20-2 of Example 20 was used instead of the intermediate obtained in step 1-2 of Example 1, and tert-butyl (S)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate in steps 1-3.

¹H NMR(500 MHz, DMSO): 10.55(m, 1H), 6.91-6.90(m, 1H), 6.85-6.75(m,
 20 0.5H), 6.57-6.48(m, 1.5H), 6.06-6.04(m, 1.5H), 5.99-5.96(m, 1.5H), 5.65-5.63(m,

0.5H), 5.43-5.41(m, 0.5H), 4.43-4.40(m, 0.5H), 4.14-4.10(m, 1H), 3.98-3.88(1.5H), 3.40(m, 4H), 3.24(s, 2H), 3.15-3.11(m, 2H), 2.67(m, 0.5H), 2.33-2.27(m, 4H), 2.19(s, 3H), 1.96(d, 3H), 1.79(m, 1H), 1.54-1.44(m, 2.5H)


5 Example 21: Preparation of (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one

Step 21-1: Preparation of methyl (S)-6-((1-(tert-butoxycarbonyl)piperidin-3-yl)amino)-2-chloropyrimidine-4-carboxylate



10 After methyl 2,4-dichloropyrimidine-6-carboxylate (5.0 g, 1.0 eq) was dissolved in tetrahydrofuran (100.0 mL), diisopropylethylamine (1.2 eq) and tert-butyl (S)-3-aminopiperidine-1-carboxylate (1.2 eq) were added thereto, and stirred at 80°C for 1 hour. Upon completion of the reaction, the mixture was cooled to 30°C or less, water (500.0 mL) and dichloromethane (500.0 mL) were added thereto, followed by
15 extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:3) to give the title compound (1.7 g, yield: 19.3%).


20 Step 21-2: Preparation of tert-butyl (S)-3-((2-chloro-6-(morpholine-4-carbonyl)pyrimidin-4-yl)amino)piperidine-1-carboxylate



After methyl (S)-6-((1-(tert-butoxycarbonyl)piperidin-3-yl)amino)-2-chloropyrimidine-4-carboxylate (1.7 g, 1.0 eq) obtained in step 21-1 was dissolved in


tetrahydrofuran (20.0 mL), 1,5,7-triazabicyclo[4,4,0]dec-5-ene (0.3 eq) and morpholine (1.2 eq) were added thereto and stirred at room temperature for 3 hours. Upon completion of the reaction, water (200.0 mL) and dichloromethane (200.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 19:1) to give the title compound (970.0 mg, yield: 55.7%).

Step 21-3: Preparation of tert-butyl 3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholine-4-carbonyl)pyrimidin-4-yl)amino)piperidine-1-carboxylate




After tert-butyl (S)-3-((2-chloro-6-(morpholin-4-carbonyl)pyrimidin-4-yl)amino)piperidine-1-carboxylate (950.0 mg, 1.0 eq) obtained in step 21-2 was dissolved in 1,4-dioxane (10.0 mL), palladium acetate (0.1 eq), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.2 eq), cesium carbonate (3.0 eq) and 2-amino-5-methylthiazole (1.2 eq) were added thereto and reacted in a microwave reactor (160°C, 30 min). Upon completion of the reaction, water (100.0 mL) and ethyl acetate (100.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 19:1) to give the title compound (900.0 mg, yield: 80.0%).

Step 21-4: Preparation of (S)-N²-(5-methylthiazol-2-yl)-6-(morpholinomethyl)-N⁴-(piperidin-3-yl)pyrimidine-2,4-diamine



After tert-butyl 3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholine-4-carbonyl)pyrimidin-4-yl)amino)piperidine-1-carboxylate (500.0 mg, 1.0 eq) obtained in step 21-3 was dissolved in tetrahydrofuran (10.0 mL), 0.9M borane-tetrahydrofuran solution (3.0 eq) was added thereto and stirred at 50°C for 5 hours. The reaction solution was cooled to 0°C, and then 6N aqueous hydrochloric acid solution (5.0 eq) was added and then stirred at 50°C for 12 hours. The reaction solution was again cooled to 0°C, and then the pH was adjusted to 12 using 12N aqueous sodium hydroxide solution, and extracted with dichloromethane (200.0 mL) and water (200.0 mL). The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give the title compound (270.0 mg, yield: 69.8%).

Step 21-5: Preparation of (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




After (S)-N²-(5-methylthiazol-2-yl)-6-(morpholinomethyl)-N⁴-(piperidin-3-yl)pyrimidine-2,4-diamine (270.0 mg, 1.0 eq) obtained in steps 21-4 was dissolved in tetrahydrofuran (4.0 mL) and water (1.0 mL), sodium bicarbonate (3.0 eq) and acryloyl chloride (1.2 eq) were added thereto and stirred at 0°C for 30 minutes. Upon

completion of the reaction, water (100.0 mL) and dichloromethane (100.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 5:1) to give the title compound (45.0 mg, yield: 14.6%).

¹H NMR(500 MHz, CDCl₃): 7.17(s, 1H), 6.57-6.61(m, 1H), 6.25-6.42(m, 2H), 5.53(s, 1H), 4.25(s, 1H), 4.10(s, 1H), 3.77(s, 4H), 3.72(s, 0.5H), 3.30-3.37(m, 4.5H), 2.54(s, 4H), 2.40(s, 3H), 2.15(s, 1H), 1.85(s, 1H), 1.66-1.67(m, 2H)

Example 22: Preparation of (S)-1-(3-((4-((4-(2-methoxyethyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one


Step 22-1: Preparation of 1-((2,6-dichloropyridin-4-yl)methyl)-4-(2-methoxyethyl)piperazine



After the intermediate (1.0 g, 1.0 eq) obtained in step 20-1 of Example 20 was dissolved in tetrahydrofuran (10.0 mL), triethylamine (1.1 mL, 2.0 eq) was added thereto. 1-Bromo-2-methoxyethane (572.0 μ l, 1.5 eq) was added and stirred for 6 hours. After concentration, ethyl acetate (10.0 mL) and water (10.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 10:1) to give the title compound (711.1 mg, yield: 60.0%).

Step 22-2: Preparation of (S)-1-(3-((4-((4-(2-methoxyethyl)piperazin-1-


yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (15.0 mg, yield: 68.0%) was obtained in the same manner as in Example 1, except that the intermediate obtained in step 22-1 of Example 22 was used instead of the intermediate obtained in step 1-2 of Example 1, and tert-butyl (S)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate in step 1-3.

¹H NMR(500 MHz, CDCl₃): 10.50(m, 1H), 7.04-7.02(m, 1.5H), 6.60(m, 0.5H), 6.44-6.49(m, 1H), 6.33-3.18(m, 3H), 5.50-5.48(m, 0.5H), 5.80(m, 0.5H), 4.52-4.05(m, 2H), 4.0-3.70(m, 2H), 3.47(s, 2H), 3.36(s, 3H), 2.74-2.52(m, 13H), 2.35(s, 3H), 2.12-2.01(m, 1H), 1.83(m, 1H), 1.67(m, 2H)

Example 23: Preparation of (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one




The title compound (25.0 mg, yield 10.7%) was obtained in the same manner as in Example 21, except that in steps 21-5 of Example 21, but-2-ynoyl chloride was

used instead of acryloyl chloride.

¹H NMR(500 MHz, CDCl₃): 7.17(d, 1H), 6.34(d, 1H), 5.30(s, 1H), 4.32-4.36(m, 1H), 4.22(s, 1H), 4.01(s, 1H), 3.78(s, 4H), 3.41-3.45(m, 1H), 3.39(d, 2H), 3.36(s, 1H), 2.54(s, 4H), 2.45(s, 3H), 2.17(s, 1H), 2.04(s, 3H), 1.62-1.72(m, 3H)

5


Example 24: Preparation of (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one



The title compound (15.0 mg, yield: 10.1%) was obtained in the same manner as in Example 21, except that in step 21-1 of Example 21, tert-butyl (S)-3-aminopyrrolidine-1-carboxylate was used instead of tert-butyl (S)-3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, CDCl₃): 11.4(s, 1H), 7.21(s, 1H), 6.38-6.47(m, 2H), 5.66-5.72(m, 1H), 5.34(s, 1H), 4.75-4.80(m, 1H), 3.91(d, 0.5H), 3.68-3.77(m, 8H), 3.51(d, 0.5H), 3.38(d, 2H), 2.54(s, 4H), 2.43(d, 3H), 2.17-2.26(m, 1H), 1.96-2.03(m, 1H).

Example 25: Preparation of (R)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




The title compound (7.0 mg, yield: 6.2%) was obtained in the same manner

as in Example 21, except that in step 21-1 of Example 21, tert-butyl (R)-3-aminopiperidine-1-carboxylate was used instead of tert-butyl (S)-3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, CDCl₃): 7.10(s, 1H), 6.50-6.61(m, 2H), 6.28-6.31(m, 3H),
 4.40(s, 1H), 4.20(s, 1H), 4.01(s, 1H), 3.77(s, 4H), 3.30-3.36(m, 4H), 2.53(s, 2H),
 2.37(s, 3H), 1.84(s, 2H), 1.70(m, 4H)

Example 26: Preparation of (S)-1-(3-(methyl(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




The title compound (145.0 mg, yield: 60.0%) was obtained in the same manner as in Example 1, except that in steps 1-3 of Example 1, tert-butyl (S)-3-(methylamino)piperidine-1-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, DMSO): 10.62-10.60(m, 1H), 6.92(s, 1H), 6.91-6.81(m, 0.5H), 6.66(m, 0.5H), 6.22(s, 1H), 6.09-5.99(m, 2H), 5.66-5.64(m, 0.5H), 5.50-5.48(m, 0.5H), 4.9(m, 0.5H), 4.8(m, 0.5H), 4.49-4.40(m, 1H), 4.08-3.92(m, 1H), 3.56(m, 4H), 3.30(s, 2H), 3.25-3.22(m, 0.5H), 2.86(s, 3H), 2.99-2.96(m, 0.5H), 2.80-2.77(m, 0.5H), 2.59-2.54(m, 0.5H), 2.34(m, 4H), 2.18(s, 3H), 1.83-1.77(m, 3H), 1.50(m, 1H)

Example 27: Preparation of N-(1-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)acrylamide


Step 27-1: Preparation of tert-butyl (1-(6-chloro-4-

(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)carbamate



After 4-((2,6-dichloropyridin-4-yl)methyl)morpholine (1.0 g, 3.8 mmol) was dissolved in N,N-dimethylformamide (8 mL), tert-butyl piperidin-4-ylcarbamate (0.9 g, 3.8 mmol) and cesium carbonate (1.3 g, 3.8 mmol) were added thereto, and then the mixture was stirred at reflux at 80°C for 12 hours. Upon completion of the reaction, the reaction mixture was diluted with ethyl acetate and washed with brine. The organic layer was collected, dried over anhydrous sodium sulfate, filtered, concentrated under reduced pressure, and purified by column chromatography (hexane/ethyl acetate = 1/1) to give the title compound (580 mg, yield: 35%).


Step 27-2: Preparation of tert-butyl (1-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)carbamate



After tert-butyl (1-(6-chloro-4-(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)carbamate (0.5 g, 1.3 mmol) obtained in step 27-1 was dissolved in 1,4-dioxane (9 mL), 5-methylthiazol-2-amine (0.2 g, 1.4 mmol), palladium acetate (0.06 g, 0.3 mmol), Xantphos (0.3 g, 0.5 mmol) and cesium carbonate (1.2 g, 3.8 mmol) were sequentially added thereto and then reacted in a microwave reactor at 150°C for 1 hour. Upon completion of the reaction, the reaction mixture was filtered through celite™, concentrated under reduced pressure, and then purified by column

chromatography (dichloromethane/methanol = 9/1) to give the title compound (106 mg, yield: 17%).


Step 27-3: Preparation of N-(6-(4-aminopyridin-1-yl)-4-(morpholinomethyl)pyridin-2-yl)-5-methylthiazol-2-amine



After tert-butyl (1-(6-((5-methylthiazol-2-yl)amino)-4-

(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)carbamate (0.1 g, 0.2 mmol) obtained in step 27-2 was dissolved in dichloromethane (11 mL), trifluoroacetic acid (229 mg, 3.0 mmol) was added thereto, and then stirred at 20°C for 2 hours. Upon completion of the reaction, 1N sodium hydroxide solution was added to adjust the pH to 7, diluted with ethyl acetate, and washed with brine. The organic layers were collected, dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to give the title compound (195 mg, yield: 100%).

Step 27-4: Preparation of N-(1-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)acrylamide




N-(6-(4-aminopyridin-1-yl)-4-(morpholinomethyl)pyridin-2-yl)-5-methylthiazol-2-amine (0.1 g, 0.2 mmol) obtained in step 27-3 was dissolved in

tetrahydrofuran (6 mL) and water (2 mL), and then cooled to 0°C, and sodium bicarbonate (0.08 g, 1.0 mmol) was added thereto. Acryloyl chloride (0.02 mL, 0.3 mmol) was slowly added to the reaction solution, and then stirred at 0°C for 10 minutes. Upon completion of the reaction, the reaction mixture was diluted with ethyl acetate, washed with brine, concentrated under reduced pressure, and purified by column chromatography (dichloromethane/methanol=9/1) to give the title compound (33 mg, yield: 37%).

¹H NMR(500 MHz, CDCl₃): 7.02(s, 1H), 6.30-6.32(m, 1H), 6.15-6.20(m, 1H), 6.01-6.10(m, 1H), 5.60-5.62(m, 1H), 5.41-5.45(m, 1H), 4.35-4.38(m, 2H), 4.15-4.20(m, 1H), 3.78(s, 4H), 3.38(s, 2H), 3.07-3.12(m, 2H), 2.45(s, 4H), 2.37(s, 3H), 2.10-2.12(m, 2H), 1.50-1.60(m, 2H).


Example 28: Preparation of 1-(6-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)octahydro-1H-pyrrolo[2,3-c]pyridin-1-yl)prop-2-en-1-one



The title compound (7 mg, yield 19%) was obtained in the same manner as in Example 27, except that in step 27-1 of Example 27, tert-butyl octahydro-1H-pyrrolo[2,3-c]pyridin-1-carboxylate was used instead of tert-butyl piperidin-4-ylcarbamate.

¹H NMR(500 MHz, CDCl₃): 6.97(s, 1H), 6.40-6.47(m, 2H), 6.07-6.20(m, 2H), 5.67-5.71(m, 1H), 4.32-4.42(m, 2H), 4.05-4.15(m, 2H), 3.70(s, 4H), 3.58-3.62(m, 2H), 3.50(s, 2H), 3.32(s, 2H), 2.95-3.05(m, 2H), 2.50-2.52(m, 1H), 2.47(s, 4H), 2.37(s, 3H), 1.85-1.90(m, 1H).


Example 29: Preparation of 1-(6-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)octahydro-1H-pyrrolo[2,3-c]pyridin-1-yl)but-2-yn-1-one



The title compound (12 mg, yield: 15%) was obtained in the same manner as in Example 27, except that in step 27-1 of Example 27, tert-butyl octahydroxy-1H-pyrrolo[2,3-c]pyridin-1-carboxylate was used instead of tert-butyl piperidin-4-ylcarbamate, and in step 27-4, but-2-ynoyl chloride was used instead of acryloyl chloride.

¹H NMR(500 MHz, CDCl₃): 6.98(s, 1H), 6.06-6.10(m, 2H), 4.20-4.25(m, 2H), 3.67(s, 3H), 3.60-3.65(m, 2H), 3.41-3.50(m, 2H), 3.31-3.37(m, 2H), 2.95-3.05(m, 2H), 2.39-2.50(m, 4H), 2.38(s, 3H), 2.27-2.35(m, 2H), 1.87-1.95(m, 4H), 1.75-1.87(m, 2H).

Example 30: Preparation of 1-(6-((6-((5-methylthiazol-2-yl)amino)-2-azaspiro[3.3]heptan-2-yl)prop-2-en-1-one




The title compound (5 mg, yield 11%) was obtained in the same manner as in Example 1, except that in step 1-3 of Example 1, tert-butyl 6-amino-2-

azaspiro[3.3]heptane-2-carboxylate was used instead of tert-butyl 3-aminopiperidine-1-carboxylate.

¹H NMR(500 MHz, CDCl₃): 6.95(s, 1H), 6.40-6.46(m, 2H), 6.25-6.30(m, 1H), 6.18(m, 1H), 5.69-5.74(m, 1H), 3.82(s, 2H), 3.72-3.77(m, 2H), 3.65-3.70(m, 2H), 3.10-3.12(m, 1H), 2.39-2.50(m, 4H), 2.36(s, 3H), 1.87-1.95(m, 4H), 1.85-1.87(m, 2H), 1.60-1.62(m, 2H).


Example 31: Preparation of (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(piperidin-1-yl)methyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (1.0 mg, yield: 1.5%) was obtained in the same manner as in Example 21, except that in step 21-2 of Example 21, piperidine was used instead of morpholine.

¹H NMR(500 MHz, CDCl₃): 7.05(s, 1H), 6.50-6.63(m, 2H), 6.21-6.28(m, 2H), 5.70(s, 0.5H), 5.54(s, 0.5H), 4.50-4.70(m, 1H), 4.20-4.40(m, 4H), 3.31(s, 2H), 3.20-3.30(m, 4H), 2.36(s, 3H), 1.33-1.72(m, 10H)

Example 32: Preparation of (S)-1-(3-((4-((4-ethylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




The title compound (0.5 mg, yield: 1.1%) was obtained in the same manner as in Example 21, except that in step 21-2 of Example 21, 1-ethylpiperazine was used instead of morpholine.


5 ¹H NMR(500 MHz, CDCl₃): 7.06(s, 1H), 6.50-6.63(m, 1H), 6.19-6.28(m, 2H), 5.71(s, 0.5H), 5.54(s, 0.5H), 4.20-4.44(m, 1H), 3.51-3.60(m, 1H), 3.37(s, 2H), 3.27-3.32(m, 1H), 2.52-2.57(m, 4H), 2.41-2.46(m, 4H), 2.36(s, 3H), 1.66-1.84(m, 8H), 1.08(t, 3H)

10 **Example 33: Preparation of (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one**


Step 33-1: Preparation of tert-butyl 4-((2,6-dichloropyridin-4-yl)oxy)piperidine-1-carboxylate



15 After 1-(tert-butoxycarbonyl)-4-hydroxypiperidine (5.5 g, 1.0 eq) was dissolved in dimethylformamide (50.0 mL), 60% sodium hydride (3.0 eq) was added and reacted at 0°C for 10 minutes, then 2,4,6-trichloropyridine (1.0 eq) was added and reacted for 30 minutes. Upon completion of the reaction, water (500.0 mL) and ethyl acetate (500.0 mL) were added thereto, followed by extraction. The separated
20 organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:5) to give the title compound (5.1 g, yield: 53.7%).


Step 33-2: Preparation of 2,6-dichloro-4-(piperidin-4-yloxy)pyridine

To tert-butyl 4-((2,6-dichloropyridin-4-yl)oxy)piperidine-1-carboxylate (5.0 g, 1.0 eq) obtained in step 33-1, 4M hydrochloric acid-dioxane solution (50.0 mL) was added and reacted at room temperature for 30 minutes. Upon completion of the reaction, the reaction solution was cooled at 0°C, and then the pH was adjusted to 12 using 12N sodium hydroxide aqueous solution, and water (250 mL) and ethyl acetate (500.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give the title compound (2.4 g, yield: 68.2%).

Step 33-3: Preparation of 2,6-dichloro-4-((1-methylpiperidin-4-yl)oxy)pyridine

After 2,6-dichloro-4-(piperidin-4-yloxy)pyridine (2.4 g, 1.0 eq) obtained in step 33-2 was dissolved in methanol (50.0 mL) and dichloromethane (50.0 mL), formaldehyde solution (1.0 eq), acetic acid (0.1 eq), and sodium triacetoxyborohydride (2.0 eq) were added thereto and reacted at room temperature for 30 minutes. Upon completion of the reaction, water (500.0 mL) and dichloromethane (500.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give the title compound (5.1 g, yield: 89.8%).


Step 33-4: Preparation of tert-butyl (S)-3-((6-chloro-4-((1-methylpiperidin-4-yl)oxy)pyridin-2-yl)amino)piperidine-1-carboxylate



5 After 2,6-dichloro-4-((1-methylpiperidin-4-yl)oxy)pyridine (2.0 g, 1.0 eq) obtained in step 33-3 was dissolved in 1,4-dioxane (20.0 mL), palladium acetate (0.1 eq), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.2 eq), cesium carbonate (3.0 eq), and 2-amino-5-methylthiazole (1.2 eq) were added thereto and reacted in a microwave reactor (150°C, 30 min). Upon completion of the reaction, water (250.0 mL) and dichloromethane (250.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 1:1) to give the title compound (900.0 mg, yield: 27.7%).

15


Step 33-5: Preparation of tert-butyl (S)-3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidine-1-carboxylate



20 After tert-butyl (S)-3-((6-chloro-4-((1-methylpiperidin-4-yl)oxy)pyridin-2-yl)amino)piperidine-1-carboxylate (900.0 mg, 1.0 eq) obtained in step 33-4 was dissolved in 1,4-dioxane (20.0 ml), palladium acetate (0.1 eq), 4,5-


bis(diphenylphosphino)-9,9-dimethylxanthene (0.2 eq), cesium carbonate (3.0 eq), and 2-amino-5-methylthiazole (1.1 eq) were added thereto, and reacted in a microwave reactor (160°C, 2hr). Upon completion of the reaction, water (250.0 mL) and ethyl acetate (250.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 1:1) to give the title compound (300.0 mg, yield: 28.3%).

Step 33-6: Preparation of (S)-4-((1-methylpiperidin-4-yl)oxy)-N²-(5-methylthiazol-2-yl)-N⁶-(piperidin-3-yl)pyridin-2,6-diamine



To tert-butyl (S)-3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidine-1-carboxylate (300.0 mg, 1.0 eq) obtained in steps 33-5, 1.25M hydrochloric acid-methanol solution (5.0 mL) was added and reacted at 50°C for 12 hours. Upon completion of the reaction, the reaction solution was cooled at 0°C, and then the pH was adjusted to 8~9 using saturated sodium bicarbonate aqueous solution, and water (100.0 mL) and ethyl acetate (100.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give the title compound (150.0 mg, yield: 62.5%).

Step 33-7: Preparation of (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




After (S)-4-((1-methylpiperidin-4-yl)oxy)-N²-(5-methylthiazol-2-yl)-N⁶-(piperidin-3-yl)pyridin-2,6-diamine (130.0 mg, 1.0 eq) obtained in steps 33-6 was dissolved in tetrahydrofuran (2.4 mL) and water (0.6 mL), sodium bicarbonate (3.0 eq) and acryloyl chloride (1.2 eq) were added and reacted at room temperature for 1 hour. Upon completion of the reaction, water (100.0 mL) and dichloromethane (100.0 mL) were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: methanol = 1:1) to give the title compound (1.0 mg, yield: 0.7%).

¹H NMR(500 MHz, CDCl₃): 6.99(s, 1H), 6.60-6.65(m, 0.5H), 6.46-6.51(m, 0.5H), 6.32(d, 0.5H), 6.24(d, 0.5H), 5.71(s, 1H), 5.53-5.59(m, 1H), 5.40(d, 2H), 4.46(s, 0.5H), 4.31(s, 2H), 4.29(s, 0.5H), 3.93(d, 1H), 3.86(s, 0.5H), 3.71(s, 0.5H), 3.48(s, 1H), 3.37-3.39(m, 1H), 2.69(s, 2H), 2.34(s, 3H), 2.30(s, 5H), 2.11(s, 1H), 2.00(s, 2H), 1.81(s, 3H), 1.65(s, 2H)

Example 34: Preparation of (S)-3-(4-((2-((1-acryloylpiperidin-3-yl)amino)-6-((5-methylthiazol-2-yl)amino)pyridin-4-yl)methyl)piperazin-1-yl)propanenitrile


Step 34-1: Preparation of 3-(4-((2,6-dichloropyridin-4-yl)methyl)piperazin-1-yl)propanenitrile



The title compound (7.1 g, yield: 57.2%) was obtained in the same manner as in steps 1-1 and 1-2 of Example 1, except that in step 1-1 of Example 1, 3-(piperazin-1-yl)propanenitrile was used instead of morpholine.

5


Step 34-2: Preparation of t-butyl (S)-3-((4-((4-(2-cyanoethyl)piperazin-1-yl)methyl)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidine-1-carboxylate



The title compound (400 mg, yield: 76.0%) was obtained in the same manner as in steps 1-3 and 1-4 of Example 1, except that the intermediate obtained in step 34-1 was used instead of the intermediate obtained in step 1-3 of Example 1, and t-butyl (S)-3-aminopiperidine-1-carboxylate was used instead of t-butyl 3-aminopiperidine-1-carboxylate.


15

Step 34-3: Preparation of (S)-3-(4-((2-((5-methylthiazol-2-yl)amino)-6-(piperidin-3-ylamino)pyridin-4-yl)methyl)piperazine-1-yl)propanenitrile



After tert-butyl (S)-3-((4-((4-(2-cyanoethyl)piperazin-1-yl)methyl)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidine-1-carboxylate (100.0 mg, 1.0 eq) obtained in step 34-2 was dissolved in dichloromethane (10.0 mL), trifluoroacetic acid (141.5 μ L, 10.0 eq) was added thereto at room temperature, and the reactant was allowed to react at room temperature for 2 hours. The reaction mixture was neutralized with 2.0M aqueous sodium hydroxide solution and then extracted with dichloromethane. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography to give 65.0 mg (yield: 77.6%) of the title compound as a brown solid.

Step 34-4: Preparation of (S)-3-(4-((2-((1-acryloylpiperidin-3-yl)amino)-6-((5-methylthiazol-2-yl)amino)pyridin-4-yl)methyl)piperazin-1-yl)propanenitrile




After (S)-3-(4-((2-((5-methylthiazol-2-yl)amino)-6-(piperidin-3-ylamino)pyridin-4-yl)methyl)piperazine-1-yl)propanenitrile (65.0 mg, 1.0 eq) obtained in step 34-3 was dissolved in tetrahydrofuran (5.0 mL) and water (1.0 mL), sodium hydrogen

carbonate (24.8 mg, 2.0 eq) was added at room temperature, and reacted for 30 minutes. Acryloyl chloride (24.0 μ L, 2.0 eq) was added to the mixture at room temperature. The reactant was allowed to react at room temperature for 10 minutes, then methanol was added, and water and ethyl acetate were added thereto, followed by extraction. The separated organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography to give 25.0 mg (yield: 34.1%) of the title compound as a brown solid.

¹H NMR(500 MHz, MeOD): 6.89(d, 1H), 6.84-6.79(m, 1H), 6.55-6.50(m, 1H), 6.20-6.03(m, 2H), 5.47(d, 1H), 4.61(d, 1H), 4.38-4.30(m, 1H), 4.29-4.18(m, 1H), 4.09-3.97 (m, 2H), 3.34 (s, 3H), 2.83 (t, 1H), 2.61-4.24 (m, 14H), 2.19-2.11(m, 1H), 1.97-1.88(m, 1H), 1.71-1.55(m, 1H).


Example 35: Preparation of (S)-1-(3-((4-methyl-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




The title compound (15 mg, yield: 21%) was obtained in the same manner as in Example 1, except that in steps 1-3 of Example 1, 2,6-dichloro-4-methylpyridine was used instead of 4-((2,6-dichloropyridin-4-yl)methyl)morpholine.

¹H NMR(500 MHz, CDCl₃): 6.94-6.98(m, 1H), 6.45-6.50(m, 1H), 6.20-6.25(m, 1H), 5.97-6.00(m, 1H), 5.75-5.85(m, 1H), 5.47-5.51(m, 1H), 4.22-4.40(m, 2H), 3.78-3.96(m, 2H), 3.70-3.72(m, 1H), 3.27-3.40(m, 2H), 2.36(s, 3H), 2.21(s, 3H), 1.85-1.90(m, 1H), 1.75-1.80(m, 1H).

Example 36: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one


Step 36-1: Preparation of 2,6-dichloro-4-(pyridin-3-ylmethyl)pyridine

After (2,6-dichloropyridin-4-yl)boronic acid (0.5 g, 2.6 mmol) was dissolved in 1,4-dioxane (13 mL) and water (1.6 mL), 3-(bromomethyl)pyridine hydrogen bromide (0.7 g, 1.6 mmol), potassium carbonate (1.8 g, 13.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.1 g, 0.2 mmol) were sequentially added thereto, and then the mixture was stirred at reflux at 110°C for 2 hours. Upon completion of the reaction, the reaction mixture was concentrated under reduced pressure and purified by column chromatography (hexane/ethyl acetate = 1/1) to give the title compound (420 mg, yield: 67%).

Step 36-2: Preparation of tert-butyl (S)-3-((6-chloro-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidine-1-carboxylate


After 2,6-dichloro-4-(pyridin-3-ylmethyl)pyridine (1.7 g, 7.1 mmol) obtained in step 36-1 was dissolved in 1,4-dioxane (24 mL), tert-butyl (S)-3-aminopiperidine-1-carboxylate (1.6 g, 7.8 mmol), palladium acetate (0.2 g, 0.7 mmol), Xantphos (0.8 g, 1.4 mmol) and sodium carbonate (2.3 g, 21.3 mmol) were sequentially added thereto, and then stirred at reflux at 100°C for 12 hours. Upon completion of the reaction, the reaction mixture was filtered through celite, concentrated under reduced pressure, and purified by column chromatography (ethyl acetate 100%) to give the title compound (280 mg, yield: 15%).

Step 36-3: Preparation of tert-butyl 3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidine-1-carboxylate



After tert-butyl (S)-3-((6-chloro-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidine-1-carboxylate (0.4 g, 2.3 mmol) obtained in step 36-2 was dissolved in 1,4-dioxane (6 mL), 5-methylthiazol-2-amine (0.1 g, 2.6 mmol), palladium acetate (0.02 g, 0.2 mmol), Xantphos (0.1 g, 0.5 mmol) and cesium carbonate (0.9 g, 7.0 mmol) were sequentially added thereto and reacted in a microwave reactor at 150°C for 1 hour. Upon completion of the reaction, the reaction mixture was filtered through celite, concentrated under reduced pressure, and purified by column chromatography (dichloromethane/methanol = 9/1) to give the title compound (360 mg, yield: 84%).


Step 36-4: Preparation of N²-(5-methylthiazol-2-yl)-N⁶-(piperidin-3-yl)-4-(pyridin-3-ylmethyl)pyridin-2,6-diamine



After tert-butyl 3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidine-1-carboxylate (0.2 g, 0.3 mmol) obtained in step 36-3 was dissolved in dichloromethane (2 mL), trifluoroacetic acid (0.5 mL, 6.6

mmol) was added thereto, and then stirred at 20°C for 2 hours. Upon completion of the reaction, 1N sodium hydroxide solution was added to adjust the pH to 7, diluted with ethyl acetate, and washed with brine. The organic layer was collected, dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to give the title compound (180 mg, yield: 100%).


Step 36-5: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



N²-(5-methylthiazol-2-yl)-N⁶-(piperidin-3-yl)-4-(pyridin-3-ylmethyl)pyridin-2,6-diamine (0.2 g, 0.4 mmol) obtained in step 36-4 was dissolved in tetrahydrofuran (6 mL) and water (2 mL), and then cooled to 0°C, and sodium bicarbonate (0.1 g, 1.1 mmol) was added thereto. Acryloyl chloride (0.05 mL, 0.6 mmol) was slowly added to the reaction solution, and then stirred at 0°C for 10 minutes. Upon completion of the reaction, the reaction mixture was diluted with ethyl acetate, washed with brine, concentrated under reduced pressure, and purified by column chromatography (dichloromethane/methanol = 9/1) to give the title compound (19 mg, yield: 25%).

¹H NMR(500 MHz, CDCl₃): 8.49(m, 2H), 7.45-7.48(m, 1H), 7.20-7.25(m, 1H), 6.78(s, 1H), 6.40-6.59(m, 1H), 6.18-6.30(m, 1H), 5.50-5.85(m, 3H), 4.34-4.48(m, 1H), 4.15-4.30(m, 2H), 4.85-4.89(m, 1H), 3.83(s 2H), 3.31-3.70(m, 3H), 2.32(s, 3H), 2.10-2.15(m, 1H), 1.80-1.85(m, 1H).


Example 37: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-2-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (11 mg, yield: 22%) was obtained in the same manner as in Example 36, except that in step 36-1 of Example 36, 2-(bromomethyl)pyridine hydrogen bromide was used instead of 3-(bromomethyl)pyridine hydrogen bromide.

¹H NMR(500 MHz, CDCl₃): 8.55(m, 1H), 7.59-7.61(m, 1H), 7.27-7.35(m, 2H), 6.90-6.95(s, 1H), 6.45-6.55(m, 1H), 6.20-6.25(m, 1H), 5.95-6.00(m, 1H), 5.85-5.90(m, 1H), 5.42-5.48(m, 1H), 4.18-4.30(m, 2H), 3.92(s, 2H), 3.85-3.90(m, 1H), 3.35-3.40(m, 2H), 2.37(s, 3H), 2.01-2.10(m, 1H), 1.85-1.90(m, 1H).


Example 38: Preparation of (S)-2-((1-acryloylpiperidin-3-yl)amino)-6-(5-methylthiazol-2-yl)isonicotinonitrile



The title compound (11 mg, yield: 22%) was obtained in the same manner as in Example 1, except that in step 1-3 of Example 1, 2,6-dichloroisonicotinonitrile was used instead of 4-((2,6-dichloropyridin-4-yl)methyl)morpholine.

¹H NMR(500 MHz, CDCl₃): 6.95-7.02(m, 1H), 6.20-6.45(m, 3H), 5.50-5.75(m, 1H), 4.95-5.05(m, 1H), 4.17-4.28(m, 1H), 3.80-4.10(m, 1H), 3.50-3.78(m, 3H), 2.37(s, 3H), 2.10-2.15(m, 1H), 1.80-1.87(m, 1H).


Example 39: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-phenylpyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one



The title compound (10 mg, yield: 22%) was obtained in the same manner as in Example 1, except that in step 1-3 of Example 1, 2,6-dichloro-4-phenylpyridine was used instead of 4-((2,6-dichloropyridin-4-yl)methyl)morpholine.

¹H NMR(500 MHz, CDCl₃): 7.52-7.60(m, 2H), 7.35-7.50(m, 3H), 6.95-7.02(s, 1H), 6.42-6.50(m, 1H), 6.17-6.37(m, 3H), 5.42-5.50(m, 1H), 4.42-4.50(m, 1H), 4.28-4.33(m, 1H), 3.89-3.95(m, 1H), 3.33-3.50(m, 2H), 2.30(s, 3H), 2.05-2.15(m, 2H), 1.81-1.90(m, 2H).


Example 40: Preparation of (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-yn-1-one



The title compound (10.0 mg, yield: 8.9%) was obtained in the same manner as in Example 33, except that in step 33-7 of Example 33, propioloyl chloride was used instead of acryloyl chloride.


¹H NMR(500 MHz, CDCl₃): 7.40(s, 1H), 6.97(s, 1H), 5.74(d, 1H), 5.55(d, 1H), 4.31-4.40(m, 2H), 4.16-4.21(m, 1H), 4.24(d, 1H), 3.82-3.87(m, 1H), 2.67(s, 2H), 2.35(t, 3H), 2.30(s, 6H), 1.94-1.99(m, 3H), 1.83-1.89(m, 4H), 1.65-1.74(m, 3H)

Example 41: Preparation of (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-yn-1-one

((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one

The title compound (40.0 mg, yield: 34.4%) was obtained in the same manner as in Example 33, except that in step 33-7 of Example 33, but-2-ynoyl chloride was used instead of acryloyl chloride.

¹H NMR(500 MHz, CDCl₃): 10.4(s, 1H), 6.99(s, 1H), 5.73(s, 0.5H), 5.71(s, 0.5H), 5.55(d, 0.5H), 5.53(d, 0.5H), 4.35-4.49(m, 1H), 4.28-4.29(m, 1H), 4.24(d, 1H), 3.82-3.86(m, 1H), 3.33-3.47(m, 2H), 2.70(s, 2H), 2.38(d, 3H), 2.30(s, 3H), 2.21-2.26(m, 3H), 2.09-2.13(m, 1H), 2.00(s, 2H), 1.80-1.83(m, 2H), 1.78(s, 3H), 1.61-1.67(m, 2H)


Example 42: Preparation of (S)-1-(3-((4-((4-ethylpiperazin-1-yl)methyl)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one

The title compound (50.0 mg, yield: 25.0%) was obtained in the same manner as in Example 22, except that in step 22-1 of Example 22, iodoethane was used instead of 1-bromo-2-methoxyethane.

¹H NMR(500 MHz, MeOH): 6.89-6.88(m, 1H), 6.85-6.45(m, 1H), 6.20-6.02(m, 3H), 5.80(m, 0.5H), 5.45(m, 0.5H), 4.60(m, 0.5H), 4.4-4.2(m, 1H), 4.15-4.00(1.5H), 3.40(s, 2H), 2.82-2.80(m, 2H), 2.50(m, 7H), 2.40(m, 2H), 2.27(s, 3H), 2.20-2.10(m,

1H), 1.92-1.80(m, 1H), 1.66-1.59(m, 3H), 1.10-1.07(t, 3H)

Example 43: Preparation of (S)-1-(3-((4-((4-methylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one




The title compound (61.0 mg, yield: 21.0%) was obtained in the same manner as in Example 22, except that in step 22-1 of Example 22, iodomethane was used instead of 1-bromo-2-methoxyethane.

¹H NMR(500 MHz, DMSO): 10.53-10.49(m, 1H), 6.91(m, 1H), 6.80-6.70(m, 0.5H), 6.54-6.46(m, 1.5H), 6.07-5.95(m, 3H), 5.65-5.63(m, 0.5H), 5.42-5.40(m, 0.5H), 4.40-4.50(m, 0.5H), 4.20-4.00(m, 1H), 3.90-3.87(m, 1.5H), 3.26(s, 2H), 3.12-3.08(m, 1H), 2.66-2.61(m, 0.5H), 2.41-2.31(m, 4H), 2.19(s, 3H), 2.13(s, 3H), 2.06-1.97(m, 3H), 1.81(m, 1.5H) 1.81(m, 1H), 1.54-1.44(m, 2H)


Example 44: Preparation of 5-methyl-N-(6-methyl-4-(morpholinomethyl)pyridin-2-yl)thiazol-2-amine

Step 44-1: Preparation of 4-((2-chloro-6-methylpyridin-4-yl)methyl)morpholine



The title compound (112.0 mg, yield: 46%) was obtained in the same manner as in step 1-1, except that in step 1-1 of Example 1, 2-chloro-6-methylisonicotinic acid was used instead of 2,6-dichloroisonicotinic acid.

Step 44-2: Preparation of 5-methyl-N-(6-methyl-4-(morpholinomethyl)pyridin-2-yl)thiazol-2-amine




5 The title compound (55.0 mg, yield: 54%) was obtained in the same manner as in Step 1-2 of Example 1 by using the intermediate obtained in step 44-1.

¹H NMR(500 MHz, CDCl₃): 7.10(s, 1H), 6.72(s, 1H), 6.67(s, 1H), 3.74-3.73(m, 4H), 3.43(s, 2H), 2.51(s, 3H), 2.46(m, 4H), 2.41(s, 3H)

10 **Example 45: Preparation of (S)-3-chloro-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)propan-1-one**

Step 45-1: Preparation of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one-hydrochloride




15 10.0 g of (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one as the material of Example 5 was dissolved in 200.0 ml of ethyl acetate, 3 equivalents of 1N-hydrochloric acid dissolved in ethyl acetate was added thereto. The mixture was

20 stirred at room temperature for 1 hour, filtered, and then dried under reduced

pressure for 12 hours at room temperature to give (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidine-1-yl)prop-2-en-1-one hydrochloride (11.6 g. yield: 85%).


5 Step 45-2: Preparation of (S)-3-chloro-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)propan-1-one



The material obtained in step 45-1 was stored at -20°C for 7 months. The material produced during the storage process was separated by column using a mixed solvent of methylene chloride and methanol in a ratio of 15:1 to give the title compound (30.0 mg, yield: 5%).

¹H NMR(500 MHz, CDCl₃): 7.03-7.0(d, 1H), 6.17-6.12(d, 1H), 5.98(s, 1H), 4.38-4.30(m, 1H), 4.20-4.11(m, 2H), 3.88-3.86(m, 1H), 3.75-3.70(m, 4H), 3.68-3.56(m, 1H), 3.43-3.39(m, 1H), 3.34(s, 2H), 2.91-2.80(m, 1H), 2.80-2.71(m, 1H), 2.62-2.45(m, 1H), 2.37(m, 4H), 2.12(s, 3H), 2.12-2.11(m, 1H), 2.10-2.12(m, 1H), 1.90-1.80(m, 2H)


Example 46: Preparation of (S,E)-3-chloro-1-(3-(6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one



The material obtained in step 45-1 was stored at -20°C for 7 months. The material produced during the storage process was separated by column using a mixed solvent of methylene chloride and methanol in a ratio of 15:1 to give the title compound (3.0 mg, yield: 0.5%).

5 ¹H NMR(500 MHz, CDCl₃): 6.99(s, 1H), 6.70-6.60(m, 1H), 6.4-6.2(m, 2H), 5.8-5.5(m, 1H), 3.9-3.8(m, 2H), 3.8-3.7(m, 4H), 3.6-3.7(m, 1H), 3.49(s, 2H), 3.39-3.41(m, 2H), 2.52(m, 4H), 2.33(s, 3H), 1.85-1.82(m, 2H), 1.68-1.60(m, 2H)

10 **Example 47: Preparation of (S)-1-(3-(4-((4-(3-aminopropyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one**




After (S)-3-(4-((2-((1-acryloylpiperidin-3-yl)amino)-6-((5-methylthiazol-2-yl)amino)pyridin-4-yl)methyl)piperazin-1-yl)propanenitrile (35.0 mg, 1.0 eq) as the material of Example 34 was dissolved in methanol (5.0 mL), 10% palladium/carbon was added thereto at room temperature, and reacted for 5 minutes. Then, the reaction mixture was filtered with methanol using celite. The separated solution was concentrated under reduced pressure and purified by column chromatography to obtain the title compound (4.0 mg, yield: 12.1%) as a brown solid.

20 ¹H NMR(500 MHz, MeOD): 6.57(s, 1H), 6.53-6.50(m, 1.5H), 6.08-5.98(m, 2.5H), 5.67-5.61(m, 0.5H), 5.42-5.38(m, 0.5H), 4.48-4.29(m, 0.5H), 4.20-4.06(m, 1H), 3.98-3.90(m, 1.5H), 3.29(s, 2H), 3.17-3.11(m, 1H), 2.65-2.62(m, 2.5H), 2.40-2.36(m, 6H), 2.23(s, 3H), 2.06-1.99(m, 3H), 1.80-1.79(m, 2.5H), 1.77(m, 2H) 1.54-1.40(m, 2H)


Example 48: Preparation of (S)-1-(3-(6-(1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one

Step 48-1: Preparation of tert-butyl (S)-3-((6-((1H-pyrazol-3-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidine-1-carboxylate




After (S)-tert-butyl 3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-carboxylate (41.0 mg, 1.0 eq) was dissolved in 1,4-dioxane (2.0 mL), tris(dibenzylideneacetone)dipalladium(0) (9.2 mg, 0.1 eq) and ($\pm$)-2,2'-bis(diphenylphosphino)-1,1'-binaphthalene (12.5 mg, 0.2 eq) were added thereto. 1H-pyrazol-3-amine (8.3 mg, 1.0 eq) was added and then cesium carbonate (97.7 mg, 3.0 eq) was sequentially added. The mixture was reacted in a microwave reactor at 130°C for 30 minutes. After cooling to 30°C or less, water (10.0 mL) and ethyl acetate (10.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (50.2 mg, yield: 99.8%).

Step 48-2: Preparation of (S)-4-(morpholinomethyl)-N²-(piperidin-3-yl)-N⁶-(1H-pyrazol-3-yl)pyridin-2,6-diamine



After the intermediate (50.0 mg, 1.0 eq) obtained in step 48-1 was dissolved in ethyl acetate (10.0 mL), 6N-hydrochloric acid aqueous solution (0.4 mL, 20.0 eq) was slowly added dropwise, and then stirred for 2 hours. After adjusting the pH to 9~12 using 12N-sodium hydroxide aqueous solution, the separated dichloromethane layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 10:1) to give the title compound (34.2 mg, yield: 87.9%).

Step 48-3: Preparation of (S)-1-(3-(6-(1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one




After the intermediate (20.0 mg, 1.0 eq) obtained in step 48-2 was dissolved in tetrahydrofuran (2.0 mL), water (1.0 mL) was added and sodium bicarbonate (14.1 mg, 3.0 eq) was added, and then cooled to 0 to 10°C. Acryloyl chloride (5.6 μ L, 1.0 eq) was slowly added dropwise, and then stirred for 30 minutes to complete the reaction. The layers were separated with dichloromethane, and then dried over

anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 15:1) to give the title compound (5.7 mg, yield: 15.4%).

¹H NMR(500 MHz, MeOD): 7.76 (d, 1H), 6.54(d, 1H), 6.17-6.13(m, 1H), 6.06(s, 2H), 6.02-5.92(m, 1H), 5.50-5.25(m, 1H), 3.83-3.74(m, 4H), 3.64(s, 2H), 2.65-2.27(m, 4H), 2.12-2.07(m, 1H), 1.73(m, 1H), 1.65-1.48(m, 2H), 1.43-1.20(m, 5H)

Example 49: Preparation of (S)-1-(3-(4-(morpholinomethyl)-6-(5-(trifluoromethyl)thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one


Step 49-1: Preparation of tert-butyl (S)-3-((4-(morpholinomethyl)-6-((5-(trifluoromethyl)thiazol-2-yl)amino)pyridin-2-yl)amino)piperidine-1-carboxylate



After (S)-tert-butyl 3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-carboxylate (200.0 mg, 1.0 eq) was dissolved in 1,4-dioxane (2.0 mL), palladium acetate (11.9 mg, 0.1 eq) and Xantphos (56.7 mg, 0.2 eq) were added thereto. 5-(Trifluoromethyl)thiazol-2-amine (81.8 mg, 1.0 eq) and cesium carbonate (476.0 mg, 3.0 eq) was sequentially added. The reactant was allowed to react in a microwave reactor at 150°C for 1 hour. After cooling to 30°C or less, water (10.0 mL) and ethyl acetate (10.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The


resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (68.2 mg, yield: 25.8%).

Step 49-2: Preparation of (S)-4-(morpholinomethyl)-N²-(piperidin-3-yl)-N⁶-(5-(trifluoromethyl)thiazol-2-yl)pyridin-2,6-diamine



After the intermediate (50.0 mg, 1.0 eq) obtained in step 48-1 was dissolved in acetate (10.0 mL), 6N-hydrochloric acid aqueous solution (0.4 mL, 20.0 eq) was slowly added dropwise, and then stirred for 2 hours. After adjusting the pH to 9~12 using 12N-sodium hydroxide aqueous solution, the separated dichloromethane layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 10:1) to give the title compound (27.0 mg, yield: 44.3%).


Step 49-3: Preparation of (S)-1-(3-(4-(morpholinomethyl)-6-(5-(trifluoromethyl)thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one



After the intermediate (20.0 mg, 1.0 eq) obtained in step 49-2 was dissolved in tetrahydrofuran (2.0 mL), water (1.0 mL) was added and sodium bicarbonate (17.2 mg, 3.0 eq) was added, and then cooled to 0 to 10°C. Acryloyl chloride (4.8 μ l, 1.0 eq) was slowly added dropwise, and then stirred for 30 minutes to complete the reaction. The layers were separated, then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 15:1) to give the title compound (3.6 mg, yield: 16.0%).

¹H NMR(500 MHz, MeOD): 7.66(d, 1H), 6.83-6.72(m, 0.5H), 6.55-6.44(m, 0.5H), 6.23(d, 1H), 6.16(d, 1H), 6.05(d, 0.5H), 5.73(d, 0.5H), 5.48(d, 0.5H), 4.54(d, 0.5H), 4.43-4.14(m, 1H), 4.03-3.93(m, 1.5H), 3.75-3.62(m, 4H), 3.38(s, 2H), 3.27-3.18(m, 1H), 2.86(t, 0.5H), 2.53-2.38(m, 4H), 2.28-2.12(m, 1H), 1.96-1.83(m, 1H), 1.72-1.49(m, 2.5H), 1.38-1.23(m, 1.5H)


Example 50: Preparation of (S)-1-(3-(6-(5-chloro-1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one



The title compound (14.0 mg, yield: 7.7%) was obtained in the same manner as in Example 48, except that in step 48-1, 5-chloro-1H-pyrazol-3-amine was used instead of 1H-pyrazol-3-amine.

¹H NMR(500 MHz, MeOD): 6.16-6.14(m, 1H), 6.13-6.11(m, 1H), 6.06(s, 2H), 6.03-6.02(m, 1H), 5.52-5.50(m, 1H), 3.73-3.70(m, 4H), 3.62 (s, 2H), 2.60-2.15(m, 4H), 2.32-2.06(m, 1H), 1.72-1.65(m, 1H), 1.65-1.50(m, 2H), 1.44-1.25(m, 5H)

Example 51: Preparation of (S)-1-(3-(4-(morpholinomethyl)-6-(thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one




The title compound (3.2 mg, yield: 21.1%) was obtained in the same manner as in Example 49, except that in step 49-1, thiazol-2-amine was used instead of 5-(trifluoromethyl) thiazol-2-amine.

¹H NMR(500 MHz, MeOD): 6.68-6.53(d, 1H), 6.38-6.30(d, 2H), 6.17-6.05(m, 2H), 5.89-5.83(d, 1H), 5.47-5.43(m, 0.5H), 5.37-5.32(m, 0.5H), 3.94-3.87(m, 1H), 3.80-3.66(m, 2H), 3.63-3.48(m, 2H), 2.58-2.16(m, 4H), 2.08-1.73(m, 4H), 1.65-1.55(m, 2H), 1.41-1.33(m, 4H)


Example 52: Preparation of (S)-1-(3-(3-fluoro-6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one

Step 52-1: Preparation of (2,6-dichloro-3-fluoropyridin-4-yl)(morpholino)methanone



After 2,6-dichloro-3-fluoronicotinic acid (500.0 mg, 1.0 eq) was dissolved in tetrahydrofuran (15.0 mL), 1,1-carbonyldiimidazole (463.3 mg, 1.2 eq) was added thereto. The mixture was stirred at room temperature (25 ~ 30°C) for 1 hour under nitrogen gas, and then morpholine (0.2 mL, 1.2 eq) was added and stirred at the same temperature for 2 hours to complete the reaction. Ethyl acetate (50.0 mL) and water (50.0 mL) were added thereto, followed by extraction, and the aqueous layer was re-extracted three times using ethyl acetate (50.0 mL). The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:5) to give the title compound (581.0 mg, 87.5%).


Step 52-2: Preparation of 4-((2,6-dichloro-3-fluoropyridin-4-yl)methyl)morpholine



The intermediate (500.0 mg, 1.0 eq) obtained in step 52-1 was dissolved in dichloromethane (20.0 mL), and then stirred at room temperature (25 to 30°C). 0.9M borane-tetrahydrofuran (6.0 mL, 3.0 eq) was slowly added dropwise. The mixture was stirred at room temperature for 12 hours to complete the reaction. The reaction solution was cooled to 0 to 10°C, and then 6N-hydrochloric acid aqueous solution (39.0 mL, 20.0 eq) was slowly added dropwise and then stirred at the same temperature for 1 hour. After adjusting the pH to 9 to 12 using 6N-sodium


hydroxide aqueous solution, the mixture was extracted twice with dichloromethane. The dichloromethane layer was separated, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (430.2 mg, yield: 85.9%).

Step 52-3: Preparation of tert-butyl (S)-3-((6-chloro-3-fluoro-4-(morpholinomethyl)pyridin-2-yl)amino)piperidine-1-carboxylate




To the intermediate (100.0 mg, 1.0 eq) obtained in step 52-2, 1,4-dioxane (2.0 mL) was added and dissolved, and then palladium acetate (9.3 mg, 0.1 eq) and Xantphos (43.4 mg, 0.2 eq) were added thereto. Tert-butyl (S)-3-aminopiperidine-1-carboxylate (75.5 mg, 1.0 eq) was added, and then cesium carbonate (325.8 mg, 3.0 eq) was sequentially added. The mixture was reacted in a microwave reactor at 140°C for 30 minutes. After cooling to 30°C or less, water (10.0 mL) and ethyl acetate (10.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (144.0 mg, yield: 89.4%).

Step 52-4: Preparation of tert-butyl (S)-3-((3-fluoro-6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidine-1-carboxylate



The intermediate (100.0 mg, 1.0 eq) obtained in step 52-3 was dissolved in 1,4-dioxane (2.0 mL). Palladium acetate (5.1 mg, 0.1 eq), Xantphos (24.3 mg, 0.2 eq), 5-methylthiano-2-amine (24.0 mg, 1.0 eq), and cesium carbonate (205.8 g, 3.0 eq) were sequentially added thereto. The mixture was reacted in a microwave reactor at 150°C for 30 minutes. After cooling to 30°C or less, water (10.0 mL) and ethyl acetate (10.0 mL) were added, and then the layers were separated. The ethyl acetate layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (ethyl acetate: hexane = 1:1) to give the title compound (86.2 mg, yield: 81.8%).

Step 52-5: Preparation of (S)-3-fluoro-N⁶-(5-methylthiazol-2-yl)-4-(morpholinomethyl)-N²-(piperidin-3-yl)pyridin-2,6-diamine




After the intermediate (80.0 mg, 1.0 eq) obtained in step 52-4 was dissolved in ethyl acetate (10.0 mL), 6N-hydrochloric acid aqueous solution (0.6 mL, 20.0 eq) was slowly added dropwise thereto, and then stirred for 2 hours. After adjusting the pH to 9~12 using 12N-sodium hydroxide aqueous solution, the separated

dichloromethane layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. Ethyl acetate (10.0 mL) was added to the resulting residue and crystals were produced for 30 minutes. The crystals were filtered and then dried to give the title compound (60.5 mg, yield: 99.9%).

5

Step 52-6: Preparation of (S)-1-(3-(3-fluoro-6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one




After the intermediate (50.0 mg, 1.0 eq) obtained in step 52-5 was dissolved in tetrahydrofuran (4.0 mL), water (1.0 mL) was added and sodium bicarbonate (31.0 mg, 3.0 eq) was added, and then cooled to 0 to 10°C. Acryloyl chloride (9.9 μ L, 1.0 eq) was slowly added dropwise, and then stirred for 30 minutes to complete the reaction. The layers were separated with dichloromethane, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane: methanol = 15:1) to give the title compound (26.5 mg, yield: 46.8%).

¹H NMR(500 MHz, MeOD): 6.88-6.80(m, 1H), 6.65(m, 1H), 6.21-6.15(m, 1H), 5.68(d, 1H), 5.37-5.28(m, 1H), 3.72-3.65(m, 4H), 3.53-3.48(s, 3H), 2.76-2.69(m, 2H), 2.52-2.42(m, 4H), 2.28-2.14 (m, 3H), 2.12-1.98(m, 2H), 1.66-1.53(m, 4H)

20


Example 53: Preparation of (S)-1-(3-(4-(morpholinomethyl)-6-(pyridin-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one



The title compound (11.0 mg, yield: 31.0%) was obtained in the same manner as in Example 48, except that in step 48-1, pyridin-2-amine was used instead of 1H-pyrazol-3-amine.

¹H NMR(500 MHz, MeOD): 8.50-8.45(t, 2H), 7.51(d, 1H), 6.80-6.75(m, 1H), 6.70-6.62(m, 0.5H), 6.24-6.16(m, 1.5H), 6.14-6.08(d, 0.5H), 5.77-5.51(m, 0.5H), 4.03-3.92 (m, 1H), 3.91-3.77(m, 2H), 3.74-3.65(m, 4H), 3.42-3.37(m, 3H), 3.26-3.18(m, 0.5H), 2.81-2.74(m, 0.5H), 2.54-2.40(m, 4H), 2.12-1.98(m, 1H), 1.89-1.81(m, 1H)

Example 54: Preparation of (S)-1-(3-(4-(morpholinomethyl)-6-(pyrimidin-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one



The title compound (24.2mg, 24.3%) was obtained in the same manner as in Example 1, except that in step 48-1, pyrimidine-2-amine was used instead of 1H-pyrazol-3-amine.

¹H NMR(500 MHz, MeOD): 8.15-8.10(m, 1H), 7.80(d, 1H), 7.62-7.56(m, 1H), 6.86-6.80 (m, 1H), 6.58-6.45(m, 1.5H), 6.24-6.19(m, 0.5H), 6.09(m, 1H), 6.06-6.00(m,

0.5H), 5.79-5.73(m, 0.5H), 4.01-3.80 (m, 4H) 3.73-3.65(m, 4H), 3.36(s, 2H), 2.52-2.41(m, 4H), 2.14-2.304(m, 1H), 1.94-1.86 (m, 2H), 1.73-1.56(m, 2H)

Experimental Example: Inhibitory activity against BTK and ITK

5 Inhibitory activities against BTK and ITK were measured for the compounds prepared in the above Examples as follows.

10 The inhibitory activities against BTK were evaluated using 'ADP-Glo™ + BTK Kinase enzyme system' kit (Promega Corporation). In a white 96-well plate, 10 μl of BTK enzyme prepared so as to have a final concentration of 1 ng/ μl was mixed with 5 μl of compounds having a final concentration of 1 μM in the case of evaluating a single concentration of compound and a concentration of 1000, 300, 100, 30, 10, 3, 1, 0.3, 0.1 and 0.03 nM in the case of IC₅₀ evaluation, and then reacted at room temperature for 15 minutes. 5 μl of substrate and 5 μl of ATP prepared so as to have a final concentration of 10 μM were added to the plate on which reactions were completed, and then allowed to react at 30°C for 1 hour. All wells of the plate were treated with 25 μl of ADP-Glo™ reagent and allowed to react at 30°C for 40 minutes. After that, all wells were treated with 50 μl of kinase detection buffer, and then reacted at 30°C for 30 minutes under light shielding conditions. For the plate on which all reactions were completed, luminescence was measured and the results were calculated. Evaluation was carried out in duplicate, and negative control and positive control were calculated depending on whether or not the enzyme was added without treatment of the compound. The IC₅₀ was calculated based on the calculated values.

25 The inhibitory activity against ITK was evaluated using 'ADP-Glo™ + ITK Kinase enzyme system' kit (Promega Corporation). In a white 96-well plate, 10 μl of ITK enzyme prepared so as to have a final concentration of 0.4 ng/ μl was mixed with 5 μl of compounds having a final concentration of 1 μM in the case of evaluating a single concentration of compound and a concentration of 1000, 300,

100, 30, 10, 3, 1, 0.3, 0.1 and 0.03 nM in the case of IC₅₀ evaluation, and then reacted at room temperature for 15 minutes. To the plate on which reactions were completed, 5 μ l of substrate and 5 μ l of ATP prepared so as to have a final concentration of 25 μ l were added and then allowed to react at 30°C for 1 hour.

- 5 All wells of the plate were treated with 25 μ l of ADP-Glo™ reagent and then allowed to react at 30°C for 40 minutes. After that, all wells were treated with 50 μ l of kinase detection buffer, and then allowed to react at 30°C for 30 minutes under light shielding conditions. For the plate on which all reactions were completed, luminescence was measured and the results were calculated. Evaluation was carried
- 10 out in duplicate, and negative control and positive control were calculated depending on whether or not the enzyme was added without treatment of the compound. The IC₅₀ was calculated based on the calculated values. The results are shown in Table 1 below.

15 [Table 1]


Example No.	Inhibitory activity		Example No.	Inhibitory activity	
	ITK IC ₅₀ (nM)	BTK IC ₅₀ (nM)		ITK IC ₅₀ (nM)	BTK IC ₅₀ (nM)
1	4.1	4.7	28	31.7	8.8
2	2.6	3.1	29	50.4	18.5
3	9.1	10.3	30	47.0	67.1
4	1.6	0.4	31	15.6	8.8
5	>200	151.6	32	5.6	3
6	36.0	27.7	33	3.9	1.8
7	3.5	3.6	34	1.7	1.1
8	71.7	72.5	35	13.4	2.1
9	>1000	>1000	36	1.8	3.9
10	>40	2.4	37	1.9	3.2
11	7.8	8.5	38	2.6	1.6

12	2	2	39	4.8	7.5
13	>200	>200	40	10.8	10.3
14	49.1	3.3	41	128.0	9.5
15	47.3	2.3	42	2.2	2.5
16	>1000	1.1	43	2.1	1.6
17	1.9	1.5	44	~1000	>1000
18	50.5	5.2	45	91.6	30.8
19	200.0	875.3	46	31.8	5.2
20	1.1	1 1	47	9.0	6.1
21	1.9	1.3	48	~1000	~1000
22	1.0	1.2	49	3.0	2.0
23	200.0	36.6	50	~1000	~1000
24	2.3	2.2	51	~1000	~1000
25	~1000	~1000	52	~ 1000	~ 1000
26	17.7	2.2	53	42.8	7.7
27	43.8	85.2	54	~ 1000	~1000

We claim:

1. A compound represented by the following Chemical Formula 1, or a pharmaceutically acceptable salt thereof:

[Chemical Formula 1]



wherein, in Chemical Formula 1,

B is a 5-membered or 6-membered heterocycle containing 1 to 3 heteroatoms each independently selected from the group consisting of N, O and S, provided that the 5-membered or 6-membered heterocycle includes at least one N,

R₁ is hydrogen, halogen, C₁₋₄ alkyl, or C₁₋₄ haloalkyl,

X₁ and X₂ are each independently N, or CR',


wherein R' is hydrogen or halogen,


L is C₁₋₄ alkylene or -O-,

R₂ is cyano; C₆₋₁₀ aryl; pyridinyl; morpholino; piperazinyl; or piperidinyl,

wherein, the piperazinyl and the piperidinyl are each independently, unsubstituted, or substituted with C₁₋₄ alkyl, C₁₋₄ alkyl substituted with cyano, C₁₋₄ alkyl substituted with amino, C₁₋₄ alkyl substituted with C₁₋₄ alkoxy, or -CO-(C₁₋₄ alkyl),

Y is a bond, -O-, -NH-, or -N(C₁₋₄ alkyl)-,





wherein R_3 is C_{1-4} alkyl, C_{1-4} haloalkyl, C_{2-4} alkenyl, C_{2-4} haloalkenyl, C_{2-4} alkynyl, or C_{2-4} haloalkynyl.

2. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

B is thiazole, pyrazole, pyridine, or pyrimidine ring, and

R_1 is hydrogen, chloro, methyl, or trifluoromethyl.

3. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

both X_1 and X_2 are CH; or

one of X_1 and X_2 is CF and the other is CH; or

one of X_1 and X_2 is N and the other is CH.

4. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

L is methylene or -O-.

5. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

R_2 is cyano; phenyl; pyridinyl; morpholino; piperazinyl substituted with methyl; piperazinyl substituted with ethyl; piperazinyl substituted with 2-cyanoethyl; piperazinyl substituted with 3-aminopropyl; piperazinyl substituted with 2-methoxyethyl; piperazinyl substituted with -CO-(methyl); unsubstituted piperidinyl; or piperidinyl substituted with methyl.

6. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

Y is a bond, -O-, -NH-, or -N(methyl)-.


7. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

R_3 is $-\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}=\text{CH}_2$, $-\text{CH}=\text{CHCH}_3$, $-\text{CH}=\text{CHCl}$, $-\text{C}\equiv\text{CH}$, or $-\text{C}\equiv\text{CCH}_3$.

8. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

the compound represented by Chemical Formula 1 is represented by the following Chemical Formula 1-1:

[Chemical Formula 1-1]



wherein, in Chemical Formula 1-1,

R_1 is hydrogen, C_{1-4} alkyl, or C_{1-4} haloalkyl,

X_1 and X_2 are each independently N, or CR' ,


wherein R' is hydrogen or halogen,

L is C_{1-4} alkylene or $-\text{O}-$,

R_2 is cyano; C_{6-10} aryl; pyridinyl; morpholino; piperazinyl; or piperidinyl,

wherein the piperazinyl and piperidinyl are each independently, unsubstituted, or substituted with C_{1-4} alkyl, C_{1-4} alkyl substituted with cyano, C_{1-4} alkyl substituted with amino, C_{1-4} alkyl substituted with C_{1-4} alkoxy, or $-\text{CO}-(\text{C}_{1-4} \text{ alkyl})$,

Y is a bond, $-\text{O}-$, $-\text{NH}-$, or $-\text{N}(\text{C}_{1-4} \text{ alkyl})-$,



wherein, R₃ is C₁₋₄ haloalkyl, C₂₋₄ alkenyl, C₂₋₄ haloalkenyl, or C₂₋₄ alkynyl.

9. The compound or a pharmaceutically acceptable salt thereof according to claim 8, wherein

R₁ is C₁₋₄ alkyl,


X₁ and X₂ are each independently N or CH,

L is C₁₋₄ alkylene or -O-,

R₂ is cyano; C₆₋₁₀ aryl; pyridinyl; morpholino; piperazinyl; or piperidinyl,

wherein, the piperazinyl and piperidinyl are each independently, unsubstituted, or substituted with C₁₋₄ alkyl, C₁₋₄ alkyl substituted with cyano, C₁₋₄ alkyl substituted with C₁₋₄ alkoxy, or -CO-(C₁₋₄ alkyl),

Y is a bond, -O-, -NH-, or -N(C₁₋₄ alkyl)-,




wherein, R₃ is C₂₋₄ alkenyl, or C₂₋₄ alkynyl.

10. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

the compound represented by Chemical Formula 1 is represented by the following Chemical Formula 1-2:

[Chemical Formula 1-2]



wherein, in Chemical Formula 1-2,


R₁ is hydrogen, or halogen,

X₁ and X₂ are each independently N, or CH,

L is C₁₋₄ alkylene,

R₂ is morpholino,

Y is -NH-,




wherein, R₃ is C₂₋₄ alkenyl.

11. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

the compound represented by Chemical Formula 1 is represented by the following Chemical Formula 1-3:

[Chemical Formula 1-3]




wherein, in Chemical Formula 1-3,

X₁ and X₂ are each independently N or CH,

L is C₁₋₄ alkylene,

R₂ is morpholino,

Y is -NH-,




wherein, R₃ is C₂₋₄ alkenyl.

12. The compound or a pharmaceutically acceptable salt thereof according to claim 1, wherein

the compound represented by Chemical Formula 1 is represented by the following Chemical Formula 1-4:

[Chemical Formula 1-4]




wherein, in Chemical Formula 1-4,

X₁ and X₂ are each independently N or CH,

L is C₁₋₄ alkylene,

R₂ is morpholino,

Y is -NH-,



wherein, R₃ is C₂₋₄ alkenyl.

13. A compound or a pharmaceutically acceptable salt thereof,
wherein the compound is any one selected from the group consisting of the following:

1) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

2) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,

3) 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

4) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

5) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-

yl)amino)piperidin-1-yl)prop-2-en-1-one,

6) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,

7) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,

8) 1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,

9) (E)-1-(3-((2-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-4-yl)amino)pyrrolidin-1-yl)but-2-en-1-one,

10) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one,

11) 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,

12) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,

13) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,

14) 1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)pyrrolidin-1-yl)but-2-yn-1-one,

15) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,

16) (R)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)oxy)piperidin-1-yl)prop-2-en-1-one,

17) 1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

- 18) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 19) 1-(4-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 20) (S)-1-(3-((4-((4-acetylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 21) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 22) (S)-1-(3-((4-((4-(2-methoxyethyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 23) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,
- 24) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)pyrrolidin-1-yl)prop-2-en-1-one,
- 25) (R)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(morpholinomethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 26) (S)-1-(3-(methyl(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 27) N-(1-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)piperidin-4-yl)acrylamide,
- 28) 1-(6-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)octahydro-1H-pyrrolo[2,3-c]pyridin-1-yl)prop-2-en-1-one,
- 29) 1-(6-(6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)octahydro-1H-pyrrolo[2,3-c]pyridin-1-yl)but-2-yn-1-one,
- 30) 1-(6-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-

yl)amino)-2-azaspiro[3.3]heptan-2-yl)prop-2-en-1-one,

31) (S)-1-(3-((4-((5-methylthiazol-2-yl)amino)-6-(piperidin-1-ylmethyl)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

32) (S)-1-(3-((4-((4-ethylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyrimidin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

33) (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

34) (S)-3-(4-((2-((1-acryloylpiperidin-3-yl)amino)-6-((5-methylthiazol-2-yl)amino)pyridin-4-yl)methyl)piperazin-1-yl)propanenitrile,

35) (S)-1-(3-((4-methyl-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

36) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-3-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

37) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(pyridin-2-ylmethyl)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

38) (S)-2-((1-acryloylpiperidin-3-yl)amino)-6-(5-methylthiazol-2-yl)amino)isonicotinonitrile,

39) (S)-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-phenylpyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

40) (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-yn-1-one,

41) (S)-1-(3-((4-((1-methylpiperidin-4-yl)oxy)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)but-2-yn-1-one,

42) (S)-1-(3-((4-((4-ethylpiperazin-1-yl)methyl)-6-((5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,

- 43) (S)-1-(3-((4-((4-methylpiperazin-1-yl)methyl)-6-(5-methylthiazol-2-yl)amino)pyridin-2-yl)amino)piperidin-1-yl)prop-2-en-1-one,
- 44) 5-methyl-N-(6-methyl-4-(morpholinomethyl)pyridin-2-yl)thiazol-2-amine,
- 45) (S)-3-chloro-1-(3-((6-((5-methylthiazol-2-yl)amino)-4-(morpholinomethyl)pyridin-2-yl)amino)piperidin-1-yl)propan-1-one,
- 46) (S,E)-3-chloro-1-(3-(6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 47) (S)-1-(3-(4-((4-(3-aminopropyl)piperazin-1-yl)methyl)-6-(5-methylthiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 48) (S)-1-(3-(6-(1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 49) (S)-1-(3-(4-(morpholinomethyl)-6-(5-(trifluoromethyl)thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 50) (S)-1-(3-(6-(5-chloro-1H-pyrazol-3-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 51) (S)-1-(3-(4-(morpholinomethyl)-6-(thiazol-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 52) (S)-1-(3-(3-fluoro-6-(5-methylthiazol-2-ylamino)-4-(morpholinomethyl)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one,
- 53) (S)-1-(3-(4-(morpholinomethyl)-6-(pyridin-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one, and
- 54) (S)-1-(3-(4-(morpholinomethyl)-6-(pyrimidin-2-ylamino)pyridin-2-ylamino)piperidin-1-yl)prop-2-en-1-one.

14. A pharmaceutical composition for preventing or treating autoimmune diseases or cancers, comprising the compound as defined in any one of claims 1 to 13 or a pharmaceutically acceptable salt thereof and a pharmaceutical acceptable carrier, adjuvant or diluent.

[Chemical Formula 1]

