



- (51) **International Patent Classification:**
C07K 14/605 (2006.01) C07K 14/575 (2006.01)
- (21) **International Application Number:**
PCT/EP2014/077337
- (22) **International Filing Date:**
11 December 2014 (11.12.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
13306713.2 13 December 2013 (13.12.2013) EP
- (71) **Applicant:** SANOFI [FR/FR]; 54 rue La Boetie, F-75008 Paris (FR).
- (72) **Inventors:** **KADEREIT, Dieter;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **LORENZ, Katrin;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **EVERS, Andreas;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **HAACK, Torsten;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **WAGNER, Michael;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **HENKEL, Bernd;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **LORENZ, Martin;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE). **STENGELIN, Siegfried;** c/o Sanofi-Aventis Deutschland GmbH, 65926 Frankfurt am Main (DE).
- (74) **Agent:** WEISS, Wolfgang; Weickmann & Weickmann, Richard-Strauss-Str. 80, 81679 Munchen (DE).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

Published:

- *with international search report (Art. 21(3))*
— *with sequence listing part of description (Rule 5.2(a))*



WO 2015/086729 A1

(54) **Title:** DUAL GLP-1/GIP RECEPTOR AGONISTS

(57) **Abstract:** The present invention relates to exendin-4 derivatives and their medical use, for example in the treatment of disorders of the metabolic syndrome, including diabetes and obesity, as well as reduction of excess food intake.

Dual GLP-1/GIP Receptor Agonists

5 Description

FIELD OF THE INVENTION

The present invention relates to exendin-4 peptide analogues which activate the
 10 glucagon-like peptide 1 (GLP-1) and the glucose-dependent insulintropic
 polypeptide (GIP) receptor and their medical use, for example in the treatment of
 disorders of the metabolic syndrome, including diabetes and obesity, as well as
 reduction of excess food intake.

15 BACKGROUND OF THE INVENTION

Exendin-4 is a 39 amino acid peptide which is produced by the salivary glands of the
 Gila monster (*Heloderma suspectum*) (Eng J. et al., *J. Biol. Chem.*, 267:7402-
 05,1 992). Exendin-4 is an activator of the glucagon-like peptide-1 (GLP-1) receptor,
 20 whereas it shows only very low activation of the GIP receptor and does not activate
 the glucagon receptor (see Table 1).

Table 1: Potencies of exendin-4 at human GLP-1, GIP and Glucagon receptors
 (indicated in pM) at increasing concentrations and measuring the formed cAMP as
 25 described in Methods.

SEQ ID NO:	peptide	EC50 hGLP-1 R [pM]	EC50 hGIP R [pM]	EC50 hGlucagon R [pM]
1	exendin-4	0.4	12500.0	>10000000

Exendin-4 shares many of the glucoregulatory actions observed with GLP-1. Clinical
 and non-clinical studies have shown that exendin-4 has several beneficial
 antidiabetic properties including a glucose dependent enhancement in insulin
 30 synthesis and secretion, glucose dependent suppression of glucagon secretion,
 slowing down gastric emptying, reduction of food intake and body weight, and an
 increase in beta-cell mass and markers of beta cell function (Gentilella R et al.,

Diabetes Obes Metab., 11:544-56, 2009; Norris SL et al., Diabet Med., 26:837-46, 2009; Bunck MC et al., Diabetes Care., 34:2041-7, 2011).

5 These effects are beneficial not only for diabetics but also for patients suffering from obesity. Patients with obesity have a higher risk of getting diabetes, hypertension, hyperlipidemia, cardiovascular and musculoskeletal diseases.

10 Relative to GLP-1 and GIP, exendin-4 is more resistant to cleavage by dipeptidyl peptidase-4 (DPP4) resulting in a longer half-life and duration of action in vivo (Eng J., Diabetes, 45 (Suppl 2):152A (abstract 554), 1996; Deacon CF, Horm Metab Res, 36: 761-5, 2004).

15 Exendin-4 was also shown to be much more stable towards degradation by neutral endopeptidase (NEP), when compared to GLP-1, glucagon or oxyntomodulin (Druce MR et al., Endocrinology, 150(4), 1712-1721, 2009).

20 Nevertheless, exendin-4 is chemically labile due to methionine oxidation in position 14 (Hargrove DM et al., Regul. Pept., 141 : 113-9, 2007) as well as deamidation and isomerization of asparagine in position 28 (WO 2004/035623).

The amino acid sequence of exendin-4 is shown as SEQ ID NO: 1:

HGEGTFTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH₂

25 The amino acid sequence of GLP-1 (7-36)-amide is shown as SEQ ID NO: 2:

HAEGTFTSDVSSYLEGQAAKEFIAWLVKGR-NH₂

30 Liraglutide is a marketed chemically modified GLP-1 analogue in which, among other modifications, a fatty acid is linked to a lysine in position 20 leading to a prolonged duration of action (Drucker DJ et al, Nature Drug Disc. Rev. 9, 267-268, 2010; Buse, JB et al., Lancet, 374:39-47, 2009).

The amino acid sequence of Liraglutide is shown as SEQ ID NO: 3:

HAEGTFTSDVSSYLEGQAAK((S)-4-Carboxy-4-hexadecanoylamino-buteryl-
)EFIAWLVRGRG-OH

5 GIP (glucose-dependent insulintropic polypeptide) is a 42 amino acid peptide that is released from intestinal K-cells following food intake. GIP and GLP-1 are the two gut enteroendocrine cell-derived hormones accounting for the incretin effect, which accounts for over 70% of the insulin response to an oral glucose challenge (Baggio LL, Drucker DJ. Biology of incretins: GLP-1 and GIP. Gastroenterology 2007; 132: 2131-2157).

GIP's amino acid sequence is shown as SEQ ID NO: 4:

YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGKKNWVKHNITQ-OH

15

Glucagon is a 29-amino acid peptide which is released into the bloodstream when circulating glucose is low. Glucagon's amino acid sequence is shown in SEQ ID NO: 5:

20 HSQGTFTSDYSKYLDSRRAQDFVQWLMNT-OH

During hypoglycemia, when blood glucose levels drop below normal, glucagon signals the liver to break down glycogen and release glucose, causing an increase of blood glucose levels to reach a normal level. Hypoglycemia is a common side effect of insulin treated patients with hyperglycemia (elevated blood glucose levels) due to diabetes. Thus, glucagon's most predominant role in glucose regulation is to counteract insulin action and maintain blood glucose levels.

30 Hoist (Hoist, J. J. Physiol. Rev. 2007, 87, 1409) and Meier (Meier, J. J. Nat. Rev. Endocrinol. 2012, 8, 728) describe that GLP-1 receptor agonists, such as GLP-1, liraglutide and exendin-4, improve glycemic control in patients with T2DM by reducing fasting and postprandial glucose (FPG and PPG). Peptides which bind and activate the GLP-1 receptor are described in patent applications WO1998/008871, WO2008/081418 and WO2008/023050, the contents of which are herein

incorporated by reference.

It has been described that dual activation of the GLP-1 and GIP receptors, e.g. by combining the actions of GLP-1 and GIP in one preparation, leads to a therapeutic principle with significantly better reduction of blood glucose levels, increased insulin secretion and reduced body weight in mice with T2DM and obesity compared to the marketed GLP-1 agonist liraglutide (e.g. VA Gault et al., Clin Sci (Lond), 121, 107-117, 2011). Native GLP-1 and GIP were proven in humans following co-infusion to interact in an additive manner with a significantly increased insulinotropic effect compared to GLP-1 alone (MA Nauck et al., J. Clin. Endocrinol. Metab., 76, 912-917, 1993).

Designing hybrid molecules which combine agonism on the GLP-1 receptor and the GIP receptor offers the therapeutic potential to achieve significantly better reduction of blood glucose levels, increased insulin secretion and an even more pronounced significant effect on body weight reduction compared to the marketed GLP-1 agonist liraglutide (e.g. VA Gault et al., Clin Sci (Lond), 121, 107-117, 2011).

Compounds of this invention are exendin-4 derivatives, which show agonistic activity at the GLP-1 and the GIP receptor and which have - among others - preferably the following modifications: Tyr at position 1 and lie at position 12.

Surprisingly, it was found that the modification of the selective GLP-1 R agonist Exendin-4 by Tyr in position 1 and lie in position 12 results in a peptide with high dual activity at the GLP-1 and GIP receptors. This observation is surprising, since the same modification in other GLP-1 agonists, such as GLP-1 itself, does not result in high activity at the GIP receptor, as shown in Table 2.

Table 2: Potencies of exendin-4 and GLP-1 peptide analogues at GLP-1 and GIP receptors (indicated in pM) at increasing concentrations and measuring the formed cAMP as described in Methods.

SEQ ID NO:	peptide	EC50 hGIP R [pM]	EC50 hGLP-1 R [pM]

6	Tyr(1)Ile(1 2)-exendin-4	93.9	1.3
7	Tyr(1)Ile(1 2)-GLP1	3660.0	5.0

Peptides which bind and activate both the GIP and the GLP-1 receptor, and improve glycaemic control, suppress body weight gain and reduce food intake are described in patent applications WO 2011/11 9657 A 1, WO 201 2/138941 A 1, WO 201 0/011439
5 A2, WO 201 0/1 48089 A 1, WO 201 1/094337 A 1, WO 201 2/0881 16 A2, the contents of which are herein incorporated by reference. These applications disclose that mixed agonists of the GLP-1 receptor and the GIP receptor can be designed as analogues of the native GIP or glucagon sequences.

10 Compounds of this invention are exendin-4 peptide analogues comprising leucine in position 10 and glutamine in position 13. Krstenansky et al. (Biochemistry, 25, 3833-3839, 1986) show the importance of residues 10 to 13 of glucagon for its receptor interactions and activation of adenylate cyclase. In the exendin-4 peptide analogues of this invention, several of the underlying residues are different from said of
15 glucagon. In particular, residues Tyr10 and Tyr13, are replaced by leucine in position 10 and glutamine, a non-aromatic polar amino acid, in position 13. This replacement, especially in combination with isoleucine in position 23 and glutamate in position 24 leads to exendin-4 derivatives with potentially improved biophysical properties as solubility or aggregation behavior in solution. The non-conservative replacement of
20 an aromatic amino acid with a polar amino acid in position 13 of an exendin-4 analogue surprisingly leads to peptides with high activity on the GIP.

Some compounds of this invention are exendin-4 peptide analogues which contain alpha, alpha-dialkylated amino acids with a basic side-chain in position 20.

25 Surprisingly, modification of the exendin-4 sequence with one of these amino acids leads to compounds with an improved biophysical profile, as solubility (in particular at low pH, especially at pH 4.5) or aggregation behaviour in solution, when the unnatural amino acid is incorporated at position 20. The resulting exendin-4 analogues thereby maintain their high activity at the GLP-1 receptor and the GIP
30 receptor. The incorporation of these unnatural amino acids also increases enzymatic stability of the peptides, potentially resulting in improved pharmacokinetic properties.

BRIEF SUMMARY OF THE INVENTION


5 Provided herein are exendin-4 analogues which potently activate the GLP-1 and the GIP receptor.

The invention provides a peptidic compound having the formula (I):



10

wherein Z is a peptide moiety having the formula (II)



15

X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and Met(O₂),

X16 represents an amino acid residue selected from Lys and Glu,

20

X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys, (R)Me-Lys and (S)Me-Orn,

X40 is absent or represents Lys,

R¹ represents NH₂,

25

R² represents t OH or NH₂,

or a salt or solvate thereof.

30

The compounds of the invention are GLP-1 and GIP receptor agonists as determined by the observation that they are capable of stimulating intracellular cAMP formation.

In vitro potency determination in cellular assays of agonists is quantified by determining the concentrations that cause 50% activation of maximal response (EC50) as described in Methods.

According to another embodiment, the peptidic compounds of the invention exhibit a relative activity of at least 0.07% (i.e. $EC_{50} < 1000$ pM), preferably at least 0.7% (i.e. $EC_{50} < 100$ pM), more preferably at least 1.4% (i.e. $EC_{50} < 50$ pM), and even more preferably at least 7% (i.e. $EC_{50} < 10$ pM) compared to that of GLP-1 (7-36) at the GLP-1 receptor ($EC_{50} = 0.7$ pM).

According to another embodiment, the peptidic compounds of the invention exhibit a relative activity of at least 0.04% (i.e. $EC_{50} < 1000$ pM), preferably at least 0.4% (i.e. $EC_{50} < 100$ pM), more preferably 0.8% (i.e. $EC_{50} < 50$ pM), and even more preferably 2% (i.e. $EC_{50} < 20$ pM) compared to that of natural GIP at the GIP receptor ($EC_{50} = 0.4$ pM).

The term "activity" as used herein preferably refers to the capability of a compound to activate the human GLP-1 receptor, the human GIP receptor and optionally the human glucagon receptor. More preferably the term "activity" as used herein refers to the capability of a compound to stimulate intracellular cAMP formation. The term "relative activity" as used herein is understood to refer to the capability of a compound to activate a receptor in a certain ratio as compared to another receptor agonist or as compared to another receptor. The activation of the receptors by the agonists (e.g. by measuring the cAMP level) is determined as described herein, e.g. as described in the examples.

According to one embodiment, the compounds of the invention have an EC_{50} for hGLP-1 receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less.

According to one embodiment, the compounds of the invention have an EC_{50} for hGIP receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less,

more preferably of 30 pM or less, and more preferably of 20 pM or less.

According to another embodiment, the compounds of the invention have an EC_{50} for hGLP-1 receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less, and/or an EC_{50} for hGIP receptor of 500 pM or less, preferably of 200 pM or less; more preferably of 150 pM or less, more preferably of 100 pM or less, more preferably of 90 pM or less, more preferably of 80 pM or less, more preferably of 70 pM or less, more preferably of 60 pM or less, more preferably of 50 pM or less, more preferably of 40 pM or less, more preferably of 30 pM or less, and more preferably of 20 pM or less.

15

In still another embodiment, the EC_{50} for both receptors, i.e. for the hGLP-1 receptor and for the hGIP receptor, is 500 pM or less, more preferably 200 pM or less, more preferably 150 pM or less, more preferably 100 pM or less, more preferably 90 pM or less, more preferably 80 pM or less, more preferably 70 pM or less, more preferably 60 pM or less, more preferably 50 pM or less, more preferably 40 pM or less, more preferably 30 pM or less, more preferably 20 pM or less.

20

The EC_{50} for hGLP-1 receptor, hGIP receptor and hGlucagon receptor may be determined as described in the Methods herein and as used to generate the results described in Example 3.

25

The compounds of the invention have the ability to reduce the intestinal passage, to increase the gastric content and/or to reduce the food intake of a patient. These activities of the compounds of the invention can be assessed in animal models known to the skilled person and also described herein in the Methods. Preferred compounds of the invention may increase the gastric content of mice, preferably of female NMRI-mice, if administered as a single dose, preferably subcutaneous dose, of 0.02 mg/kg body weight by at least 25%, more preferably by at least 30%, more preferably by at least 40%, more preferably by at least 50%, more preferably by at

30

least 60%, more preferably by at least 70%, more preferably by at least 80%.

Preferably, this result is measured 1 h after administration of the respective compound and 30 mins after administration of a bolus, and/or reduces intestinal
5 passage of mice, preferably of female NMRI-mice, if administered as a single dose, preferably subcutaneous dose, of 0.02 mg/kg body weight at least by 45%; more preferably by at least 50%, more preferably by at least 55%, more preferably by at least 60%, and more preferably at least 65%; and/or reduces food intake of mice, preferably of female NMRI-mice, over a period of 22 h, if administered as a single
10 dose, preferably subcutaneous dose of 0.01 mg/kg body weight by at least 10%, more preferably 15%, and more preferably 20%.

The compounds of the invention have the ability to reduce blood glucose level, and/or to reduce HbA1c levels of a patient. These activities of the compounds of the
15 invention can be assessed in animal models known to the skilled person and also described herein in the Methods.

Preferred compounds of the invention may reduce blood glucose level of mice, preferably in female leptin-receptor deficient diabetic db/db mice over a period of 24
20 h, if administered as a single dose, preferably subcutaneous dose, of 0.01 mg/kg body weight by at least 4 mmol/L; more preferably by at least 6 mmol/L, more preferably by at least 8 mmol/L. If the dose is increased to 0.1 mg/kg body weight a more pronounced reduction of blood glucose levels can be observed in mice over a period of 24 h, if administered as a single dose, preferably subcutaneous dose.
25 Preferably the compounds of the invention lead to a reduction by at least 7 mmol/L; more preferably by at least 9 mmol/L, more preferably by at least 11 mmol/L. The compounds of the invention preferably reduce the increase of HbA1c levels of mice over a period of 4 weeks, if administered at a daily dose of 0.01 mg/kg to about the
30 ignition value.

The compounds of the invention also have the ability to reduce body weight of a patient. These activities of the compounds of the invention can be assessed in animal models known to the skilled person and also described herein in the Methods.

Surprisingly, it was found that peptidic compounds of the formula (I) showed very potent GLP-1 and GIP receptor activation.

Furthermore, oxidation (in vitro or in vivo) of methionine, present in the core structure of exendin-4, can be avoided for peptidic compounds of the formula (I) if the Met in position 14 is replaced, for example by small lipophilic amino acids.

Further, compounds of the invention preferably have a high solubility at acidic and/or physiological pH values, e.g., at pH 4.5 and/or at pH 7.4 at 25 °C, in another embodiment at least 0.5 mg/ml and in a particular embodiment at least 1.0 mg/ml.

Furthermore, according to one embodiment, compounds of the invention preferably have a high stability when stored in solution. Preferred assay conditions for determining the stability is storage for 7 days at 25 °C or 40 °C in solution at pH 4.5, pH 7.4 or pH 9. The remaining amount of peptide is determined by chromatographic analyses as described in Methods and Examples. Preferably, after 7 days at 25°C or 40°C in solution at pH 4.5, pH 7.4 or pH 9, the remaining peptide amount is at least 80%, more preferably at least 85%, even more preferably at least 90% and even more preferably at least 95%.

Preferably, the compounds of the present invention comprise a peptide moiety Z (formula II) which is a linear sequence of 39-40 amino carboxylic acids, particularly α -amino carboxylic acids linked by peptide, i.e. carboxamide, bonds.

A further embodiment relates to a group of compounds, wherein

R^2 is NH_2 .

A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and Met(O₂),

X16 represents an amino acid residue selected from Lys and Glu,

X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys, (R)Me-Lys and (S)Me-Orn, particularly Arg

X40 is absent.

5 A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and Met(O₂),

X16 represents an amino acid residue selected from Lys and Glu,

X20 represents Arg,

10 X40 is absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Leu, Nle, Met(O) and Met(O₂),

15 X16 represents Lys,

X20 represents Arg,

X40 is absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

20 X14 represents Leu,

X16 represents Lys,

X20 represents an amino acid residue selected from Aib, (S)Me-Lys, (R)Me-Lys and (S)Me-Orn,

X40 is absent.

25

A further embodiment relates to a group of compounds, wherein

X14 represents Leu,

X16 represents Lys,

X20 represents an amino acid residue selected from Aib, (S)Me-Lys, and (S)Me-Orn,

30

X40 is absent.

A further embodiment relates to a group of compounds, wherein

X14 represents Leu,

X 16 represents Lys,

X20 represents an amino acid residue selected from (S)Me-Lys, and (S)Me-Orn,

X40 is absent.

5 A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and
Met(O₂), particularly Leu, Nle, Met(O) and Met(O₂),

X 16 represents Lys,

X20 represents Arg,

10 X40 is absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Met and Leu,

X 16 represents Glu,

15 X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys,
(R)Me-Lys and (S)Me-Orn,

X40 is absent or represents Lys.

A further embodiment relates to a group of compounds, wherein

20 X 14 represents Leu,

X 16 represents an amino acid residue selected from Lys and Glu,

X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys,
(R)Me-Lys and (S)Me-Orn,

X40 is absent.

25

A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Leu and Nle

X 16 represents an amino acid residue selected from Lys and Glu,

X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys,
30 (R)Me-Lys and (S)Me-Orn,

X40 is absent.

A further embodiment relates to a group of compounds, wherein

X14 represents an amino acid residue selected from Met, Met(O) and Met(O₂),

X16 represents an amino acid residue selected from Lys and Glu,
X20 represents Arg,
X40 is absent.

- 5 A further embodiment relates to a group of compounds, wherein
X14 represents an amino acid residue selected from Nle, Met(O) and Met(O₂),
X16 represents Lys,
X20 represents Arg,
X40 is absent or represents Lys.

10

Specific examples of peptidic compounds of formula (I) are the compounds of SEQ
ID NO: 8 - 20 as well as salts and solvates thereof.

- 15 Specific examples of peptidic compounds of formula (I) are the compounds of SEQ
ID NO: 10 - 14 and 16 - 20 as well as salts and solvates thereof.

Specific examples of peptidic compounds of formula (I) are the compounds of SEQ
ID NO: 10, 16, 17 and 19 as well as salts and solvates thereof.

- 20 In certain embodiments, i.e. when the compound of formula (I) comprises genetically
encoded amino acid residues, the invention further provides a nucleic acid (which
may be DNA or RNA) encoding said compound, an expression vector comprising
such a nucleic acid, and a host cell containing such a nucleic acid or expression
vector.

25

In a further aspect, the present invention provides a composition comprising a
compound of the invention in admixture with a carrier. In preferred embodiments, the
composition is a pharmaceutically acceptable composition and the carrier is a
pharmaceutically acceptable carrier. The compound of the invention may be in the
30 form of a salt, e.g. a pharmaceutically acceptable salt or a solvate, e.g. a hydrate. In
still a further aspect, the present invention provides a composition for use in a
method of medical treatment, particularly in human medicine.

In certain embodiments, the nucleic acid or the expression vector may be used as

therapeutic agents, e.g. in gene therapy.

The compounds of formula (I) are suitable for therapeutic application without an additionally therapeutically effective agent. In other embodiments, however, the
5 compounds are used together with at least one additional therapeutically active agent, as described in "combination therapy".

The compounds of formula (I) are particularly suitable for the treatment or prevention of diseases or disorders caused by, associated with and/or accompanied by
10 disturbances in carbohydrate and/or lipid metabolism, e.g. for the treatment or prevention of hyperglycemia, type 2 diabetes, impaired glucose tolerance, type 1 diabetes, obesity and metabolic syndrome. Further, the compounds of the invention are particularly suitable for the treatment or prevention of degenerative diseases, particularly neurodegenerative diseases.

15 The compounds described find use, inter alia, in preventing weight gain or promoting weight loss. By "preventing" is meant inhibiting or reducing when compared to the absence of treatment, and is not necessarily meant to imply complete cessation of a disorder.

20 The compounds of the invention may cause a decrease in food intake and/or increase in energy expenditure, resulting in the observed effect on body weight.

25 Independently of their effect on body weight, the compounds of the invention may have a beneficial effect on circulating cholesterol levels, being capable of improving lipid levels, particularly LDL, as well as HDL levels (e.g. increasing HDL/LDL ratio).

30 Thus, the compounds of the invention can be used for direct or indirect therapy of any condition caused or characterised by excess body weight, such as the treatment and/or prevention of obesity, morbid obesity, obesity linked inflammation, obesity linked gallbladder disease, obesity induced sleep apnea. They may also be used for treatment and prevention of the metabolic syndrome, diabetes, hypertension, atherogenic dyslipidemia, atherosclerosis, arteriosclerosis, coronary heart disease, or stroke. Their effects in these conditions may be as a result of or associated with their

effect on body weight, or may be independent thereof.

Preferred medical uses include delaying or preventing disease progression in type 2 diabetes, treating metabolic syndrome, treating obesity or preventing overweight, for decreasing food intake, increase energy expenditure, reducing body weight, delaying the progression from impaired glucose tolerance (IGT) to type 2 diabetes; delaying the progression from type 2 diabetes to insulin-requiring diabetes; regulating appetite; inducing satiety; preventing weight regain after successful weight loss; treating a disease or state related to overweight or obesity; treating bulimia; treating binge eating; treating atherosclerosis, hypertension, type 2 diabetes, IGT, dyslipidemia, coronary heart disease, hepatic steatosis, treatment of beta-blocker poisoning, use for inhibition of the motility of the gastrointestinal tract, useful in connection with investigations of the gastrointestinal tract using techniques such as X-ray, CT- and NMR-scanning.

15

Further preferred medical uses include treatment or prevention of degenerative disorders, particularly neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, Huntington's disease, ataxia, e.g. spinocerebellar ataxia, Kennedy disease, myotonic dystrophy, Lewy body dementia, multi-systemic atrophy, amyotrophic lateral sclerosis, primary lateral sclerosis, spinal muscular atrophy, prion-associated diseases, e.g. Creutzfeldt-Jacob disease, multiple sclerosis, telangiectasia, Batten disease, corticobasal degeneration, subacute combined degeneration of spinal cord, Tabes dorsalis, Tay-Sachs disease, toxic encephalopathy, infantile Refsum disease, Refsum disease, neuroacanthocytosis, Niemann-Pick disease, Lyme disease, Machado-Joseph disease, Sandhoff disease, Shy-Drager syndrome, wobbly hedgehog syndrome, proteopathy, cerebral β -amyloid angiopathy, retinal ganglion cell degeneration in glaucoma, synucleinopathies, tauopathies, frontotemporal lobar degeneration (FTLD), dementia, cadasil syndrome, hereditary cerebral hemorrhage with amyloidosis, Alexander disease, seipinopathies, familial amyloidotic neuropathy, senile systemic amyloidosis, serpinopathies, AL (light chain) amyloidosis (primary systemic amyloidosis), AH (heavy chain) amyloidosis, AA (secondary) amyloidosis, aortic medial amyloidosis, ApoAI amyloidosis, ApoAII amyloidosis, ApoAIV amyloidosis, familial amyloidosis of the Finnish type (FAF), Lysozyme amyloidosis, Fibrinogen amyloidosis, Dialysis amyloidosis, Inclusion body

myositis/myopathy, Cataracts, Retinitis pigmentosa with rhodopsin mutations, medullary thyroid carcinoma, cardiac atrial amyloidosis, pituitary prolactinoma, Hereditary lattice corneal dystrophy, Cutaneous lichen amyloidosis, Mallory bodies, corneal lactoferrin amyloidosis, pulmonary alveolar proteinosis, odontogenic
 5 (Pindborg) tumor amyloid, cystic fibrosis, sickle cell disease or critical illness myopathy (CIM).

Further medical uses include treatment of hyperglycemia, type 2 diabetes, obesity, particularly type 2 diabetes.

10

Further medical uses include treatment of bone related disorders, such as osteoporosis or osteoarthritis, etc., where increased bone formation and decreased bone resorption might be beneficial.

15

DETAILED DESCRIPTION OF THE INVENTION


Definitions





20 The amino acid sequences of the present invention contain the conventional one letter and three letter codes for naturally occurring amino acids, as well as generally accepted three letter codes for other amino acids, such as Aib (α -aminoisobutyric acid), Orn (ornithin), Dab (2,4-diamino butyric acid), Dap (2,3-diamino propionic acid), Nle (norleucine), GABA (γ -aminobutyric acid) or Ahx (ϵ -aminohexanoic acid).

25

Furthermore, the following codes were used for the amino acids shown in Table 3:

Table 3:

Name	Structure	code
(S)- α -methyl-lysine		(S)MeLys

(R)- α -methyl-lysine		(R)MeLys
(S)- α -methyl-ornithin		(S)MeOrn
L-methionine sulfoxide		L-Met(O)
L-methionine sulfone		L-Met(O2)

The term „native exendin-4" refers to native exendin-4 having the sequence HEGTFTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH2 (SEQ ID NO: 1).

5

The invention provides peptidic compounds as defined above.

The peptidic compounds of the present invention comprise a linear backbone of amino carboxylic acids linked by peptide, i.e. carboxamide bonds. Preferably, the amino carboxylic acids are α -amino carboxylic acids and more preferably L- α -amino carboxylic acids, unless indicated otherwise. The peptidic compounds preferably comprise a backbone sequence of 39-40 amino carboxylic acids.

For the avoidance of doubt, in the definitions provided herein, it is generally intended that the sequence of the peptidic moiety (II) differs from native exendin-4 at least at one of those positions which are stated to allow variation. Amino acids within the peptidic moiety (II) can be considered to be numbered consecutively from 0 to 40 in the conventional N-terminal to C-terminal direction. Reference to a „position" within peptidic moiety (II) should be constructed accordingly, as should reference to positions within native exendin-4 and other molecules, e.g., in exendin-4, His is at

20

position 1, Gly at position 2, ..., Met at position 14, ... and Ser at position 39.

In a further aspect, the present invention provides a composition comprising a compound of the invention as described herein, or a salt or solvate thereof, in admixture with a carrier.

The invention also provides the use of a compound of the present invention for use as a medicament, particularly for the treatment of a condition as described below.

The invention also provides a composition wherein the composition is a pharmaceutically acceptable composition, and the carrier is a pharmaceutically acceptable carrier.

Peptide synthesis

15

The skilled person is aware of a variety of different methods to prepare peptides that are described in this invention. These methods include but are not limited to synthetic approaches and recombinant gene expression. Thus, one way of preparing these peptides is the synthesis in solution or on a solid support and subsequent isolation and purification. A different way of preparing the peptides is gene expression in a host cell in which a DNA sequence encoding the peptide has been introduced. Alternatively, the gene expression can be achieved without utilizing a cell system. The methods described above may also be combined in any way.

25 A preferred way to prepare the peptides of the present invention is solid phase synthesis on a suitable resin. Solid phase peptide synthesis is a well established methodology (see for example: Stewart and Young, Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, Ill., 1984; E. Atherton and R. C. Sheppard, Solid Phase Peptide Synthesis. A Practical Approach, Oxford-IRL Press, New York, 1989).

30 Solid phase synthesis is initiated by attaching an N-terminally protected amino acid with its carboxy terminus to an inert solid support carrying a cleavable linker. This solid support can be any polymer that allows coupling of the initial amino acid, e.g. a trityl resin, a chlorotriyl resin, a Wang resin or a Rink resin in which the linkage of the carboxy group (or carboxamide for Rink resin) to the resin is sensitive to acid (when

Fmoc strategy is used). The polymer support must be stable under the conditions used to deprotect the α -amino group during the peptide synthesis.

5 After the first amino acid has been coupled to the solid support, the α -amino protecting group of this amino acid is removed. The remaining protected amino acids are then coupled one after the other in the order represented by the peptide sequence using appropriate amide coupling reagents, for example BOP, HBTU, HATU or DIC (N,N'-diisopropylcarbodiimide) / HOBt (1-hydroxybenzotriazol), wherein BOP, HBTU and HATU are used with tertiary amine bases. Alternatively, the liberated
10 N-terminus can be functionalized with groups other than amino acids, for example carboxylic acids, etc.

Usually, reactive side-chain groups of the amino acids are protected with suitable blocking groups. These protecting groups are removed after the desired peptides
15 have been assembled. They are removed concomitantly with the cleavage of the desired product from the resin under the same conditions. Protecting groups and the procedures to introduce protecting groups can be found in Protective Groups in Organic Synthesis, 3d ed., Greene, T. W. and Wuts, P. G. M., Wiley & Sons (New York: 1999).

20 Finally the peptide is cleaved from the resin. This can be achieved by using King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The raw material can then be purified by chromatography, e.g. preparative RP-HPLC, if necessary.

25

Potency

30 As used herein, the term "potency" or "in vitro potency" is a measure for the ability of a compound to activate the receptors for GLP-1, GIP or glucagon in a cell-based assay. Numerically, it is expressed as the "EC50 value", which is the effective concentration of a compound that induces a half maximal increase of response (e.g. formation of intracellular cAMP) in a dose-response experiment.

Therapeutic uses

According to one aspect, the compounds of the invention are for use in medicine, particularly in human medicine.

5

The compounds of the invention are agonists for the receptors for GLP-1 and for GIP receptor ("dual agonists"). Such peptides that are GIP/GLP-1 co-agonists may provide therapeutic benefit to address a clinical need for targeting the metabolic syndrome by allowing simultaneous treatment of diabetes and obesity.

10

Metabolic syndrome is a combination of medical disorders that, when occurring together, increase the risk of developing type 2 diabetes, as well as atherosclerotic vascular disease, e.g. heart disease and stroke. Defining medical parameters for the metabolic syndrome include diabetes mellitus, impaired glucose tolerance, raised
15 fasting glucose, insulin resistance, urinary albumin secretion, central obesity, hypertension, elevated triglycerides, elevated LDL cholesterol and reduced HDL cholesterol.

Obesity is a medical condition in which excess body fat has accumulated to the
20 extent that it may have an adverse effect on health and life expectancy and due to its increasing prevalence in adults and children it has become one of the leading preventable causes of death in modern world. It increases the likelihood of various other diseases, including heart disease, type 2 diabetes, obstructive sleep apnea, certain types of cancer, as well as osteoarthritis, and it is most commonly caused by
25 a combination of excess food intake, reduced energy expenditure, as well as genetic susceptibility.

Diabetes mellitus, often simply called diabetes, is a group of metabolic diseases in which a person has high blood sugar levels, either because the body does not
30 produce enough insulin, or because cells do not respond to the insulin that is produced. The most common types of diabetes are: (1) type 1 diabetes, where the body fails to produce insulin; (2) type 2 diabetes, where the body fails to use insulin properly, combined with an increase in insulin deficiency over time, and (3) gestational diabetes, where women develop diabetes due to their pregnancy. All

forms of diabetes increase the risk of long-term complications, which typically develop after many years. Most of these long-term complications are based on damage to blood vessels and can be divided into the two categories "macrovascular" disease, arising from atherosclerosis of larger blood vessels and "microvascular" disease, arising from damage of small blood vessels. Examples for macrovascular disease conditions are ischemic heart disease, myocardial infarction, stroke and peripheral vascular disease. Examples for microvascular diseases are diabetic retinopathy, diabetic nephropathy, as well as diabetic neuropathy.

10 The receptors for GLP-1 and GIP as well as glucagon are members of the family of 7-transmembrane-spanning, heterotrimeric G-protein coupled receptors. They are structurally related to each other and share not only a significant level of sequence identity, but have also similar mechanisms of ligand recognition and intracellular signaling pathways.

15

Similarly, the peptides GLP-1, GIP and glucagon share regions of high sequence identity/similarity. GLP-1 and glucagon are produced from a common precursor, proglucagon, which is differentially processed in a tissue-specific manner to yield e.g. GLP-1 in intestinal endocrine cells and glucagon in alpha cells of pancreatic islets. GIP is derived from a larger proGIP prohormone precursor and is synthesized and released from K-cells located in the small intestine.

20

The peptidic incretin hormones GLP-1 and GIP are secreted by intestinal endocrine cells in response to food and account for up to 70% of meal-stimulated insulin secretion. Evidence suggests that GLP-1 secretion is reduced in subjects with

25

impaired glucose tolerance or type 2 diabetes, whereas responsiveness to GLP-1 is still preserved in these patients. Thus, targeting of the GLP-1 receptor with suitable agonists offers an attractive approach for treatment of metabolic disorders, including diabetes. The receptor for GLP-1 is distributed widely, being found mainly in

30

pancreatic islets, brain, heart, kidney and the gastrointestinal tract. In the pancreas, GLP-1 acts in a strictly glucose-dependent manner by increasing secretion of insulin from beta cells. This glucose-dependency shows that activation of GLP-1 receptors is unlikely to cause hypoglycemia. Also the receptor for GIP is broadly expressed in peripheral tissues including pancreatic islets, adipose tissue, stomach, small intestine, heart, bone, lung, kidney, testis, adrenal cortex, pituitary, endothelial cells,

- trachea, spleen, thymus, thyroid and brain. Consistent with its biological function as incretin hormone, the pancreatic beta cell express the highest levels of the receptor for GIP in humans. There is some clinical evidence that the GIP-receptor mediated signaling could be impaired in patients with T2DM but the impairment of GIP-action is shown to be reversible and could be restored with improvement of the diabetic status. Of note, the stimulation of insulin secretion by both incretin hormones, GIP and GLP-1, is strictly glucose-dependent ensuring a fail-safe mechanism associated with a low risk for hypoglycemia.
- 5
- 10 At the beta cell level, GLP-1 and GIP have been shown to promote glucose sensitivity, neogenesis, proliferation, transcription of proinsulin and hypertrophy, as well as anti-apoptosis. A peptide with dual agonistic activity for the GLP-1 and the GIP receptor could be anticipated to have additive or synergistic anti-diabetic benefit. Other relevant effects of GLP-1 beyond the pancreas include delayed gastric emptying, increased satiety, decreased food intake, reduction of body weight, as well as neuroprotective and cardioprotective effects. In patients with type 2 diabetes, such extrapancreatic effects could be particularly important considering the high rates of comorbidities like obesity and cardiovascular disease. Further GIP actions in peripheral tissues beyond the pancreas comprise increased bone formation and decreased bone resorption as well as neuroprotective effects which might be beneficial for the treatment of osteoporosis and cognitive defects like Alzheimer's disease.
- 15
- 20
- Oxyntomodulin is a peptide hormone consisting of glucagon with a C-terminal extension encompassing eight amino acids. Like GLP-1 and glucagon, it is preformed in proglucagon and cleaved and secreted in a tissue-specific manner by endocrinal cells of the small bowel. Oxyntomodulin is known to stimulate both the receptors for GLP-1 and glucagon and is therefore the prototype of a dual agonist.
- 25
- 30 As GLP-1 and GIP are known for their anti-diabetic effects, it is conceivable that a combination of the activities of the two hormones in one molecule can yield a powerful medication for treatment of the metabolic syndrome and in particular its components diabetes and obesity.

Accordingly, the compounds of the invention may be used for treatment of glucose intolerance, insulin resistance, pre-diabetes, increased fasting glucose, type 2 diabetes, hypertension, dyslipidemia, arteriosclerosis, coronary heart disease, peripheral artery disease, stroke or any combination of these individual disease components.

In addition, they may be used for control of appetite, feeding and calory intake, increase of energy expenditure, prevention of weight gain, promotion of weight loss, reduction of excess body weight and altogether treatment of obesity, including morbid obesity.

Further disease states and health conditions which could be treated with the compounds of the invention are obesity-linked inflammation, obesity-linked gallbladder disease and obesity-induced sleep apnea.

Although all these conditions could be associated directly or indirectly with obesity, the effects of the compounds of the invention may be mediated in whole or in part via an effect on body weight, or independent thereof.

Further diseases to be treated are osteoporosis and neurodegenerative diseases such as Alzheimer's disease or Parkinson's disease, or other degenerative diseases as described above.

Compared to GLP-1, glucagon and oxyntomodulin, exendin-4 has beneficial physicochemical properties, such as solubility and stability in solution and under physiological conditions (including enzymatic stability towards degradation by enzymes, such as DPP-4 or NEP), which results in a longer duration of action in vivo. Therefore, exendin-4 might serve as good starting scaffold to obtain exendin-4 analogs with dual pharmacologies, e.g., GLP-1/GIP agonism.

Nevertheless, exendin-4 has also been shown to be chemically labile due to methionine oxidation in position 14 as well as deamidation and isomerization of asparagine in position 28. Therefore, stability might be further improved by substitution of methionine at position 14 and the avoidance of sequences that are

known to be prone to degradation via aspartimide formation, especially Asp-Gly or Asn-Gly at positions 28 and 29.

5 Pharmaceutical compositions

The term "pharmaceutical composition" indicates a mixture containing ingredients that are compatible when mixed and which may be administered. A pharmaceutical composition may include one or more medicinal drugs. Additionally, the
10 pharmaceutical composition may include carriers, buffers, acidifying agents, alkalizing agents, solvents, adjuvants, tonicity adjusters, emollients, expanders, preservatives, physical and chemical stabilizers e.g. surfactants, antioxidants and other components, whether these are considered active or inactive ingredients. Guidance for the skilled in preparing pharmaceutical compositions may be found, for
15 example, in Remington: The Science and Practice of Pharmacy, (20th ed.) ed. A. R. Gennaro A. R., 2000, Lippencott Williams & Wilkins and in R.C.Rowe et al (Ed), Handbook of Pharmaceutical Excipients, PhP, May 2013 update.

The exendin-4 peptide derivatives of the present invention, or salts thereof, are
20 administered in conjunction with an acceptable pharmaceutical carrier, diluent, or excipient as part of a pharmaceutical composition. A "pharmaceutically acceptable carrier" is a carrier which is physiologically acceptable (e.g. physiologically acceptable pH) while retaining the therapeutic properties of the substance with which it is administered. Standard acceptable pharmaceutical carriers and their
25 formulations are known to one skilled in the art and described, for example, in Remington: The Science and Practice of Pharmacy, (20th ed.) ed. A. R. Gennaro A. R., 2000, Lippencott Williams & Wilkins and in R.C.Rowe et al (Ed), Handbook of Pharmaceutical excipients, PhP, May 2013 update. One exemplary pharmaceutically acceptable carrier is physiological saline solution.

30

In one embodiment carriers are selected from the group of buffers (e.g. citrate/citric acid), acidifying agents (e.g. hydrochloric acid), alkalizing agents (e.g. sodium hydroxide), preservatives (e.g. phenol), co-solvents (e.g. polyethylene glycol 400), tonicity adjusters (e.g. mannitol), stabilizers (e.g. surfactant, antioxidants, amino

acids).

Concentrations used are in a range that is physiologically acceptable.

- 5 Acceptable pharmaceutical carriers or diluents include those used in formulations suitable for oral, rectal, nasal or parenteral (including subcutaneous, intramuscular, intravenous, intradermal, and transdermal) administration. The compounds of the present invention will typically be administered parenterally.
- 10 The term "pharmaceutically acceptable salt" means salts of the compounds of the invention which are safe and effective for use in mammals. Pharmaceutically acceptable salts may include, but are not limited to, acid addition salts and basic salts. Examples of acid addition salts include chloride, sulfate, hydrogen sulfate, (hydrogen) phosphate, acetate, citrate, tosylate or mesylate salts. Examples of basic
- 15 salts include salts with inorganic cations, e.g. alkaline or alkaline earth metal salts such as sodium, potassium, magnesium or calcium salts and salts with organic cations such as amine salts. Further examples of pharmaceutically acceptable salts are described in Remington: The Science and Practice of Pharmacy, (20th ed.) ed. A. R. Gennaro A. R., 2000, Lippencott Williams & Wilkins or in Handbook of
- 20 Pharmaceutical Salts, Properties, Selection and Use, e.d. P. H. Stahl, C. G. Wermuth, 2002, jointly published by Verlag Helvetica Chimica Acta, Zurich, Switzerland, and Wiley-VCH, Weinheim, Germany.

- The term "solvate" means complexes of the compounds of the invention or salts
- 25 thereof with solvent molecules, e.g. organic solvent molecules and/or water.

In the pharmaceutical composition, the exendin-4 derivative can be in monomeric or oligomeric form.

- 30 The term "therapeutically effective amount" of a compound refers to a nontoxic but sufficient amount of the compound to provide the desired effect. The amount of a compound of the formula I necessary to achieve the desired biological effect depends on a number of factors, for example the specific compound chosen, the intended use, the mode of administration and the clinical condition of the patient. An

appropriate "effective" amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation. For example the "therapeutically effective amount" of a compound of the formula (I) is about 0.01 to 50 mg/dose, preferably 0.1 to 10 mg/dose.

5

Pharmaceutical compositions of the invention are those suitable for parenteral (for example subcutaneous, intramuscular, intradermal or intravenous), oral, rectal, topical and peroral (for example sublingual) administration, although the most suitable mode of administration depends in each individual case on the nature and severity of the condition to be treated and on the nature of the compound of formula I used in each case.

10

Suitable pharmaceutical compositions may be in the form of separate units, for example capsules, tablets and powders in vials or ampoules, each of which contains a defined amount of the compound; as powders or granules; as solution or suspension in an aqueous or nonaqueous liquid; or as an oil-in-water or water-in-oil emulsion. It may be provided in single or multiple dose injectable form, for example in the form of a pen. The compositions may, as already mentioned, be prepared by any suitable pharmaceutical method which includes a step in which the active ingredient and the carrier (which may consist of one or more additional ingredients) are brought into contact.

15

20

In certain embodiments the pharmaceutical composition may be provided together with a device for application, for example together with a syringe, an injection pen or an autoinjector. Such devices may be provided separate from a pharmaceutical composition or prefilled with the pharmaceutical composition.

25

Combination therapy

30

The compounds of the present invention, dual agonists for the GLP-1 and GIP receptors, can be widely combined with other pharmacologically active compounds, such as all drugs mentioned in the Rote Liste 2013, e.g. with all antidiabetics mentioned in the Rote Liste 2013, chapter 12, all weight-reducing agents or appetite

suppressants mentioned in the Rote Liste 2013, chapter 1, all lipid-lowering agents mentioned in the Rote Liste 2013, chapter 58, all antihypertensives and nephroprotectives, mentioned in the Rote Liste 2013, or all diuretics mentioned in the Rote Liste 2013, chapter 36.

5

The active ingredient combinations can be used especially for a synergistic improvement in action. They can be applied either by separate administration of the active ingredients to the patient or in the form of combination products in which a plurality of active ingredients are present in one pharmaceutical preparation. When the active ingredients are administered by separate administration of the active ingredients, this can be done simultaneously or successively.

10

15

Most of the active ingredients mentioned hereinafter are disclosed in the USP Dictionary of USAN and International Drug Names, US Pharmacopeia, Rockville 2011 .

20

Other active substances which are suitable for such combinations include in particular those which for example potentiate the therapeutic effect of one or more active substances with respect to one of the indications mentioned and/or which allow the dosage of one or more active substances to be reduced.

Therapeutic agents which are suitable for combinations include, for example, antidiabetic agents such as:

25

Insulin and Insulin derivatives, for example: Glargine / Lantus[®], 270 - 330U/mL of insulin glargine (EP 2387989 A), 300U/mL of insulin glargine (EP 2387989 A), Glulisin / Apidra[®], Detemir / Levemir[®], Lispro / Humalog[®] / Liprolog[®], Degludec / DegludecPlus, Aspart, basal insulin and analogues (e.g. LY-2605541, LY2963016, NN1436), PEGylated insulin Lispro, Humulin[®], Linjeta, SuliXen[®], NN1 045, Insulin plus Symlin, PE0139, fast-acting and short-acting insulins (e.g. Linjeta, PH20, NN1 218, HinsBet), (APC-002)hydrogel, oral, inhalable, transdermal and sublingual insulins (e.g. Exubera[®], Nasulin[®], Afrezza, Tregopil, TPM 02, Capsulin, Oral-lyn[®], Cobalamin[®] oral insulin, ORMD-0801, NN1 953, NN1 954, NN1956, VIAtab, Oshadi

30

oral insulin). Additionally included are also those insulin derivatives which are bonded to albumin or another protein by a bifunctional linker.

5 GLP-1 , GLP-1 analogues and GLP-1 receptor agonists, for example: Lixisenatide / AVE001 0 / ZP1 0 / Lyxumia, Exenatide / Exendin-4 / Byetta / Bydureon / ITCA 650 / AC-2993, Liraglutide / Victoza, Semaglutide, Taspoglutide, Syncria / Albiglutide, Dulaglutide, rExendin-4, CJC-11 34-PC, PB-1 023, TTP-054, Langlenatide / HM-11260C, CM-3, GLP-1 Eligen, ORMD-0901 , NN-9924, NN-9926, NN-9927, Nodexen, Viador-GLP-1 , CVX-096, ZYOG-1 , ZYD-1 , GSK-2374697, DA-3091 , MAR-701 ,
10 MAR709, ZP-2929, ZP-3022, TT-401 , BHM-034. MOD-6030, CAM-2036, DA-1 5864, ARI-2651 ,ARI-2255, Exenatide-XTEN and Glucagon-Xten.

DPP-4 inhibitors, for example: Alogliptin / Nesina, Trajenta / Linagliptin / BI-1 356 / Ondero / Trajenta / Tradjenta / Trayenta / Tradzenta, Saxagliptin / Onglyza, Sitagliptin
15 / Januvia / Xelevia / Tesave / Janumet / Velmetia, Galvus / Vildagliptin, Anagliptin, Gemigliptin, Teneligliptin, Melogliptin, Trelagliptin, DA-1 229, Omarigliptin / MK-31 02, KM-223, Evogliptin, ARI-2243, PBL-1427, Pinoxacin.

SGLT2 inhibitors, for example: Invokana / Canaglifozin, Forxiga / Dapagliflozin,
20 Remoglifozin, Sergliflozin, Empagliflozin, Ipragliflozin, Tofogliflozin, Luseogliflozin, LX-4211 , Ertuglifozin / PF-04971 729, RO-4998452, EGT-0001442, KGA-3235 / DSP-3235, LIK066, SBM-TFC-039,

Biguanides (e.g. Metformin, Buformin, Phenformin), Thiazolidinediones (e.g.
25 Pioglitazone, Rivoglitazone, Rosiglitazone, Troglitazone), dual PPAR agonists (e.g. Aleglitazar, Muraglitazar, Tesaglitazar), Sulfonylureas (e.g. Tolbutamide, Glibenclamide, Glimiperide/Amaryl, Glipizide), Meglitinides (e.g. Nateglinide, Repaglinide, Mitiglinide), Alpha-glucosidase inhibitors (e.g. Acarbose, Miglitol, Voglibose), Amylin and Amylin analogues (e.g. Pramlintide, Symlin).

30

GPR1 19 agonists (e.g. GSK-263A, PSN-821 , MBX-2982, APD-597, ZYG-1 9, DS-8500), GPR40 agonists (e.g. Fasiglifam / TAK-875, TUG-424, P-1 736, JTT-851 , GW9508).

Other suitable combination partners are: Cycloset, inhibitors of 11-beta-HSD (e.g. LY2523199, BMS770767, RG-4929, BMS816336, AZD-8329, HSD-016, BI-135585), activators of glucokinase (e.g. TTP-399, AMG-151, TAK-329, GKM-001), inhibitors of DGAT (e.g. LCQ-908), inhibitors of protein tyrosinephosphatase 1 (e.g.

5 Trodusquemine), inhibitors of glucose-6-phosphatase, inhibitors of fructose-1,6-bisphosphatase, inhibitors of glycogen phosphorylase, inhibitors of phosphoenolpyruvate carboxykinase, inhibitors of glycogen synthase kinase, inhibitors of pyruvate dehydrokinase, alpha2-antagonists, CCR-2 antagonists, SGLT-1 inhibitors (e.g. LX-2761).

10

One or more lipid lowering agents are also suitable as combination partners, such as for example: HMG-CoA-reductase inhibitors (e.g. Simvastatin, Atorvastatin), fibrates (e.g. Bezafibrate, Fenofibrate), nicotinic acid and the derivatives thereof (e.g. Niacin), PPAR-(alpha, gamma or alpha/gamma) agonists or modulators (e.g. Aleglitazar),
15 PPAR-delta agonists, ACAT inhibitors (e.g. Avasimibe), cholesterol absorption inhibitors (e.g. Ezetimibe), Bile acid-binding substances (e.g. Cholestyramine), ileal bile acid transport inhibitors, MTP inhibitors, or modulators of PCSK9.

20 HDL-raising compounds such as: CETP inhibitors (e.g. Torcetrapib, Anacetrapid, Dalcatrapid, Evacetrapid, JTT-302, DRL-17822, TA-8995) or ABC1 regulators.

Other suitable combination partners are one or more active substances for the treatment of obesity, such as for example: Sibutramine, Tesofensine, Orlistat, antagonists of the cannabinoid-1 receptor, MCH-1 receptor antagonists, MC4
25 receptor agonists, NPY5 or NPY2 antagonists (e.g. Velneperit), beta-3-agonists, leptin or leptin mimetics, agonists of the 5HT2c receptor (e.g. Lorcaserin), or the combinations of bupropione/naltrexone, bupropione/zonisamide, bupropione/phentermine or pramlintide/metreleptin.

30 Other suitable combination partners are:

Further gastrointestinal peptides such as Peptide YY 3-36 (PYY3-36) or analogues thereof, pancreatic polypeptide (PP) or analogues thereof.

Glucagon receptor agonists or antagonists, GIP receptor agonists or antagonists, ghrelin antagonists or inverse agonists, Xenin and analogues thereof.

Moreover, combinations with drugs for influencing high blood pressure, chronic heart
5 failure or atherosclerosis, such as e.g.: Angiotensin II receptor antagonists (e.g. telmisartan, candesartan, valsartan, losartan, eprosartan, irbesartan, olmesartan, tasosartan, azilsartan), ACE inhibitors, ECE inhibitors, diuretics, beta-blockers, calcium antagonists, centrally acting hypertensives, antagonists of the alpha-2-
adrenergic receptor, inhibitors of neutral endopeptidase, thrombocyte aggregation
10 inhibitors and others or combinations thereof are suitable.

In another aspect, this invention relates to the use of a compound according to the invention or a physiologically acceptable salt thereof combined with at least one of the active substances described above as a combination partner, for preparing a
15 medicament which is suitable for the treatment or prevention of diseases or conditions which can be affected by binding to the receptors for GLP-1 and glucagon and by modulating their activity. This is preferably a disease in the context of the metabolic syndrome, particularly one of the diseases or conditions listed above, most particularly diabetes or obesity or complications thereof.

20 The use of the compounds according to the invention, or a physiologically acceptable salt thereof, in combination with one or more active substances may take place simultaneously, separately or sequentially.

25 The use of the compound according to the invention, or a physiologically acceptable salt thereof, in combination with another active substance may take place simultaneously or at staggered times, but particularly within a short space of time. If they are administered simultaneously, the two active substances are given to the patient together; if they are used at staggered times, the two active substances are
30 given to the patient within a period of less than or equal to 12 hours, but particularly less than or equal to 6 hours.

Consequently, in another aspect, this invention relates to a medicament which comprises a compound according to the invention or a physiologically acceptable salt

of such a compound and at least one of the active substances described above as combination partners, optionally together with one or more inert carriers and/or diluents.

- 5 The compound according to the invention, or physiologically acceptable salt or solvate thereof, and the additional active substance to be combined therewith may both be present together in one formulation, for example a tablet or capsule, or separately in two identical or different formulations, for example as so-called kit-of-parts.

10

METHODS

- 15 Abbreviations employed are as follows:

AA	amino acid
cAMP	cyclic adenosine monophosphate
Boc	tert-butyloxycarbonyl
20 BOP	(benzotriazol-1-yl)oxytris(dimethylamino)phosphonium hexafluorophosphate
BSA	bovine serum albumin
tBu	tertiary butyl
Dde	1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-ethyl
25 ivDde	1-(4,4-dimethyl-2,6-dioxocyclohexylidene)3-methyl-butyl
DIC	N,N'-diisopropylcarbodiimide
DIPEA	N,N-diisopropylethylamine
DMEM	Dulbecco's modified Eagle's medium
DMF	dimethyl formamide
30 EDT	ethanedithiol
FBS	fetal bovine serum
Fmoc	fluorenylmethyloxycarbonyl
HATU	0-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
35 HBSS	Hanks' Balanced Salt Solution

	HBTU	2-(1 H-benzotriazol-1 -yl)-1 ,1 ,3,3-tetramethyl-uronium hexafluorophosphate
	HEPES	2-[4-(2-hydroxyethyl)piperazin-1 -yl]ethanesulfonic acid
	HOBt	1-hydroxybenzotriazole
5	HOSu	N-hydroxysuccinimide
	HPLC	High Performance Liquid Chromatography
	HTRF	Homogenous Time Resolved Fluorescence
	IBMX	3-isobutyl-1-methylxanthine
	LC/MS	Liquid Chromatography/Mass Spectrometry
10	Palm	palmitoyl
	PBS	phosphate buffered saline
	PEG	polyethylene glycole
	PK	pharmacokinetic
	RP-HPLC	reversed-phase high performance liquid chromatography
15	TFA	trifluoroacetic acid
	Trt	trityl
	UPLC	Ultra Performance Liquid Chromatography
	UV	ultraviolet

20

General synthesis of peptidic compounds

Materials:

25 Different Rink-Amide resins (4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-norleucylaminomethyl resin, Merck Biosciences; 4-[(2,4-Dimethoxyphenyl)(Fmoc-amino)methyl]phenoxy acetamido methyl resin, Agilent Technologies) were used for the synthesis of peptide amides with loadings in the range of 0.3-0.4 mmol/g.

30 Fmoc protected natural amino acids were purchased from Protein Technologies Inc., Senn Chemicals, Merck Biosciences, Novabiochem, Iris Biotech, Nagase or Bachem. The following standard amino acids were used throughout the syntheses: Fmoc-L-Ala-OH, Fmoc-L-Arg(Pbf)-OH, Fmoc-L-Asn(Trt)-OH, Fmoc-L-Asp(OtBu)-OH, Fmoc-L-Cys(Trt)-OH, Fmoc-L-Gln(Trt)-OH, Fmoc-L-Glu(OtBu)-OH, Fmoc-Gly-OH, Fmoc-L-

His(Trt)-OH, Fmoc-L-Ile-OH, Fmoc-L-Leu-OH, Fmoc-L-Lys(Boc)-OH, Fmoc-L-Met-OH, Fmoc-L-Phe-OH, Fmoc-L-Pro-OH, Fmoc-L-Ser(tBu)-OH, Fmoc-L-Thr(tBu)-OH, Fmoc-L-Trp(Boc)-OH, Fmoc-L-Tyr(tBu)-OH, Fmoc-L-Val-OH.

- 5 In addition, the following special amino acids were purchased from the same suppliers as above: Fmoc-L-Lys(ivDde)-OH, Fmoc-Aib-OH, Fmoc-D-Ser(tBu)-OH, Fmoc-D-Ala-OH, Boc-L-His(Boc)-OH (available as toluene solvate) and Boc-L-His(Trt)-OH, Fmoc-L-Nle-OH, Fmoc-L-Met(O)-OH, Fmoc-L-Met(O₂)-OH, Fmoc-(S)MeLys(Boc)-OH, Fmoc-(R)MeLys(Boc)-OH, Fmoc-(S)MeOrn(Boc)-OH and Boc-L-Tyr(tBu)-OH.

The solid phase peptide syntheses were performed for example on a Prelude Peptide Synthesizer (Protein Technologies Inc) or similar automated synthesizer using standard Fmoc chemistry and HBTU/DIPEA activation. DMF was used as the
15 solvent. Deprotection : 20% piperidine/DMF for 2 x 2.5 min. Washes: 7 x DMF. Coupling 2:5:1 0 200 mM AA/ 500 mM HBTU / 2M DIPEA in DMF 2 x for 20 min. Washes: 5 x DMF.

All the peptides that had been synthesized were cleaved from the resin with King's
20 cleavage cocktail consisting of 82.5% TFA, 5% phenol, 5% water, 5% thioanisole, 2.5% EDT. The crude peptides were then precipitated in diethyl or diisopropyl ether, centrifuged, and lyophilized. Peptides were analyzed by analytical HPLC and checked by ESI mass spectrometry. Crude peptides were purified by a conventional preparative HPLC purification procedure.

25

Analytical HPLC / UPLC

Method A: detection at 215 nm

- column: Aeris Peptide, 3.6 μm , XB-C1 8 (250 x 4.6 mm) at 60 °C
30 solvent: H₂O+0.1 %TFA : ACN+0.1 %TFA (flow 1.5 ml/min)
gradient: 90:10 (0 min) to 90:10 (3 min) to 10:90 (43 min) to 10:90 (48 min) to 90:10 (49 min) to 90:10 (50 min)

Method B: detection at 220 nm

column: Zorbax, 5 μm , C 18 (250 x 4.6 mm) at 25 °C
solvent: H₂O+0.1 %TFA : 90% ACN + 10% H₂O +0.1 %TFA (flow 1.0 ml/min)
gradient: 100:0 (0 min) to 98:2 (2 min) to 30:70 (15 min) to 5:95 (20 min) to 0:100 (25 min) to 0:100 (30 min) to 98:2 (32 min) to 98:2 (35 min)

5

Method C1: detection at 210 - 225 nm, optionally coupled to a mass analyser Waters LCT Premier, electrospray positive ion mode

column: Waters ACQUITY UPLC® BEH™ C 18 1.7 μm (150 x 2.1 mm) at 50 °C
solvent: H₂O+1 %FA : ACN+1 %FA (flow 0.5 ml/min)

10 gradient: 95:5 (0 min) to 95:5 (1.80 min) to 80:20 (1.85 min) to 80:20 (3 min) to 60:40 (23 min) to 25:75 (23.1 min) to 25:75 (25 min) to 95:5 (25.1 min) to 95:5 (30min)

Method C2: detection at 210 - 225 nm, optionally coupled to a mass analyser Waters LCT Premier, electrospray positive ion mode

15 column: Waters ACQUITY UPLC® BEH™ C 18 1.7 μm (150 x 2.1 mm) at 50 °C
solvent: H₂O+1 %FA : ACN+1 %FA (flow 0.6 ml/min)
gradient: 95:5 (0 min) to 95:5 (1 min) to 65:35 (2 min) to 65:35 (3 min) to 45:55 (23 min) to 25:75 (23.1 min) to 25:75 (25 min) to 95:5 (25.1 min) to 95:5 (30 min)

20

Method C3: detection at 210 - 225 nm, optionally coupled to a mass analyser Waters LCT Premier, electrospray positive ion mode

25 column: Waters ACQUITY UPLC® BEH™ C 18 1.7 μm (150 x 2.1 mm) at 50 °C
solvent: H₂O+1 %FA : ACN+1 %FA (flow 1 ml/min)
gradient: 95:5 (0 min) to 95:5 (1 min) to 65:35 (2 min) to 65:35 (3 min) to 45:55 (20 min) to 2:98 (20.1 min) to 2:98 (25 min) to 95:5 (25.1 min) to 95:5 (30 min)

30 **Method C4:**

detection at 210 - 225 nm, optionally coupled to a mass analyser Waters LCT Premier, electrospray positive ion mode

column: Waters ACQUITY UPLC® BEH™ C 18 1.7 μm (150 x 2.1 mm) at 50 °C
solvent: H₂O+1 %FA : ACN+1 %FA (flow 1 ml/min)

gradient: 95:5 (0 min) to 95:5 (1.80 min) to 80:20 (1.85 min) to 80:20 (3 min) to 60:40 (23 min) to 2:98 (23.1 min) to 2:98 (25 min) to 95:5 (25.1 min) to 95:5 (30 min)

5 **Method D:** detection at 214 nm

column: Waters X-Bridge C18 3.5 μm 2.1 x 150 mm

solvent: H₂O+0.5%TFA :ACN (flow 0.55 ml/min)

gradient: 90:10 (0 min) to 40:60 (5 min) to 1:99 (15 min)

10 **Method E:** detection at 210 - 225 nm, optionally coupled to a mass analyser Waters LCT Premier, electrospray positive ion mode

column: Waters ACQUITY UPLC[®] BEH[™] C18 1.7 μm (150 x 2.1 mm) at 50 °C

solvent: H₂O+1 %FA :ACN+1 %FA (flow 0.9 ml/min)

gradient: 95:5 (0 min) to 95:5 (2min) to 35:65 (3 min) to 65:35 (23.5 min) to 5:95 (24 min) to 95:5 (26min) to 95:5 (30min)

15

General Preparative HPLC Purification Procedure:

The crude peptides were purified either on an Akta Purifier System or on a Jasco
20 semiprep HPLC System. Preparative RP-C18-HPLC columns of different sizes and
with different flow rates were used depending on the amount of crude peptide to be
purified. Acetonitrile + 0.05 to 0.1 % TFA (B) and water + 0.05 to 0.1 % TFA (A) were
employed as eluents. Alternatively, a buffer system consisting of acetonitrile and
water with minor amounts of acetic acid was used. Product-containing fractions were
25 collected and lyophilized to obtain the purified product, typically as TFA or acetate
salt.

Solubility and Stability-Testing of exendin-4 derivatives

Prior to the testing of solubility and stability of a peptide batch, its content was
30 determined. Therefore, two parameters were investigated, its purity (HPLC-UV) and
the amount of salt load of the batch (ion chromatography).

For solubility testing, the target concentration was 1.0 mg/mL pure compound.

Therefore, solutions from solid samples were prepared in different buffer systems

with a concentration of 1.0 mg/mL compound based on the previously determined content. HPLC-UV was performed after 2 h of gentle agitation from the supernatant, which was obtained by 20 min of centrifugation at 4000 rpm.

- 5 The solubility was then determined by comparison with the UV peak areas obtained with a stock solution of the peptide at a concentration of 2 mg/mL in pure water or a variable amount of acetonitrile (optical control that all of the compound was dissolved). This analysis also served as starting point (t_0) for the stability testing.
- 10 For stability testing, an aliquot of the supernatant obtained for solubility was stored for 7 days at 25°C or 40 °C. After that time course, the sample was centrifuged for 20 min at 4000 rpm and the supernatant was analysed with HPLC-UV.
- For determination of the amount of the remaining peptide, the peak areas of the target compound at t_0 and t_7 were compared, resulting in "% remaining peptide",
- 15 following the equation

$$\% \text{ remaining peptide} = [(\text{peak area peptide } t_7) \times 100] / \text{peak area peptide } t_0.$$

- The amount of soluble degradation products was calculated from the comparison of
- 20 the sum of the peak areas from all observed impurities reduced by the sum of peak areas observed at t_0 (i.e. to determine the amount of newly formed peptide-related species). This value was given in percentual relation to the initial amount of peptide at t_0 , following the equation:

- 25 $\% \text{ soluble degradation products} = \{[(\text{peak area sum of impurities } t_7) - (\text{peak area sum of impurities } t_0)] \times 100\} / \text{peak area peptide } t_0$

- The potential difference from the sum of "% remaining peptide" and "% soluble degradation products" to 100% reflects the amount of peptide which did not remain
- 30 soluble upon stress conditions following the equation

$$\% \text{ precipitate} = 100 - ([\% \text{ remaining peptide}] + [\% \text{ soluble degradation products}])$$

This precipitate includes non-soluble degradation products, polymers and/or fibrils,

which have been removed from analysis by centrifugation.

The chemical stability is expressed as "% remaining peptide".

5 Anion Chromatography

Instrument: Dionex ICS-2000, pre/column: Ion Pac AG-1 8 2 x 50 mm (Dionex)/AS1 8 2 x 250 mm (Dionex), eluent: aqueous sodium hydroxide, flow: 0.38 mL/min, gradient: 0-6 min: 22 mM KOH, 6-12 min: 22-28 mM KOH, 12-15 min: 28-50 mM KOH, 15-20min: 22mM KOH, suppressor: ASRS 300 2 mm, detection: conductivity.

10

As HPLC/UPLC method, method D or E has been used.

In vitro cellular assays for GIP receptor, GLP-1 receptor and glucagon receptor efficacy

15

Agonism of compounds for the receptors was determined by functional assays measuring cAMP response of HEK-293 cell lines stably expressing human GIP, GLP-1 or glucagon receptor.

20

cAMP content of cells was determined using a kit from Cisbio Corp. (cat. no. 62AM4PEC) based on HTRF (Homogenous Time Resolved Fluorescence). For preparation, cells were split into T175 culture flasks and grown overnight to near confluency in medium (DMEM / 10% FBS). Medium was then removed and cells washed with PBS lacking calcium and magnesium, followed by proteinase treatment

25

with accutase (Sigma-Aldrich cat. no. A6964). Detached cells were washed and resuspended in assay buffer (1 x HBSS; 20 mM HEPES, 0.1 % BSA, 2 mM IBMX) and cellular density determined. They were then diluted to 400000 cells/ml and 25 μ l aliquots dispensed into the wells of 96-well plates. For measurement, 25 μ l of test compound in assay buffer was added to the wells, followed by incubation for 30

30

minutes at room temperature. After addition of HTRF reagents diluted in lysis buffer (kit components), the plates were incubated for 1 hr, followed by measurement of the fluorescence ratio at 665 / 620 nm. In vitro potency of agonists was quantified by determining the concentrations that caused 50% activation of maximal response (EC50).

Bioanalytical screening method for quantification of exendin-4 derivatives in mice

Mice were dosed 1 mg/kg subcutaneously (s.c). The mice were sacrificed and blood samples were collected after 0.25, 0.5, 1, 2, 4, 8, 16 and 24 hours post application. Plasma samples were analyzed after protein precipitation via liquid chromatography mass spectrometry (LC/MS). PK parameters and half-life were calculated using WinonLin Version 5.2.1 (non-compartment model).

10 Glucose lowering in female diabetic dbdb-mice

Female diabetic dbdb-mice (BKS.Cg- +Leprdb/+Leprdb/OlaHsd) 10 weeks of age at study start were used. Mice were habituated to feeding and housing conditions for at least 2 weeks. 7 days prior to study start, HbA1c were determined to allocate mice to groups, aiming to spread low, medium and high HbA1c-values and in consequence the group-means (n = 8), as equally as possible. On the day of study, food was removed, directly before sampling for baseline glucose assessment (t = 0 min). Immediately afterwards, compounds or vehicle (phosphor buffered saline, PBS) were administered subcutaneously, 100 pg/kg, 10 ml/kg. Afterwards, blood samples were drawn by tail tip incision at 15, 30, 60, 90, 120, 150, 180, 240, 360, 480 min and 24 h. Food was re-offered after the 480 min-sampling. Data were analysed by 2-W-ANOVA on repeated measurements, followed by Dunnett's test as post-hoc assessment, level of significance $p < 0.05$.

25

EXAMPLES

The invention is further illustrated by the following examples.

Example 1:

30 Synthesis of SEQ ID NO: 10

The solid phase synthesis was carried out on Rink-resin with a loading of 0.29 mmol/g, 75-150 μm from the company Agilent Technologies. The Fmoc-synthesis

strategy was applied with HBTU/DIPEA-activation. The peptide was cleaved from the resin with King's cocktail (D. S. King, C. G. Fields, G. B. Fields, Int. J. Peptide Protein Res. 36, 1990, 255-266). The crude product was purified via preparative HPLC on a Waters column (XBridge, BEH1 30, Prep C 18 5 μ M) using an acetonitrile/water gradient (both buffers with 0.1 % TFA). The purified peptide was analysed by LCMS (Method C4). Deconvolution of the mass signals found under the peak with retention time 9.55 min revealed the peptide mass 4 164.1 which is in line with the expected value of 4 163.7.

10 In an analogous way, the following peptides were synthesized and characterized, see table 4.

Table 4: list of synthesized peptides and comparison of calculated vs. found molecular weight.

SEQ ID NO	calc. Mass	found mass	Rt [min]	Method
8	4182.7	4181.7	16.32	B
9	4181.7	4181.1	16.15	B
10	4163.7	4164.1	9.55	C4
11	4163.7	4163.6	5.25	C3
12	4291.9	4291.6	4.34	C3
13	4197.8	4198.1 4197.9	21.10 21.17	A
14	4213.8	4213.9	20.95	A
15	4164.6	4164.6	not determined	
16	4093.6	4092.8	7.74	C3
17	4150.7	4150.6	5.91	C3
18	4150.7	4150.6	5.53	C3
19	4136.6	4135.8	6.43	C3
20	4163.7	4163.9	19.37	C4

15

Example 2: Chemical stability and solubility

Solubility and chemical stability of peptidic compounds were assessed as described in Methods. The results are given in Table 5.

Table 5: Chemical stability and solubility

SEQ ID NO	Stability (pH4.5) at 25 °C [%]	Stability (pH7.4) at 25 °C [%]	Solubility (pH4.5) [µg/ml]	Solubility (pH7.4) [µg/ml]
1	100	77	934	>1000
8	97	97	842	>1000
11	100	100	911	>1000
12	100	100	943	848
14	100	92	>1000	>1000

5

Example 3: In vitro data on GLP-1 , GIP and glucagon receptor

Potencies of peptidic compounds at the GLP-1 , GIP and glucagon receptors were determined by exposing cells expressing human glucagon receptor (hGLUC R), human GIP (hGIP R) and human GLP-1 receptor (hGLP-1 R) to the listed compounds at increasing concentrations and measuring the formed cAMP as described in Methods.

The results for Exendin-4 derivatives with activity at the human GIP (hGIP R), human GLP-1 receptor (hGLP-1 R) and human glucagon receptor (hGLUC R) are shown in Table 6. All compounds are full agonists of the GIP and GLP-1 receptors.

20

Table 6. EC50 values of exendin-4 peptide analogues at GLP-1 , GIP and Glucagon receptors (indicated in pM)

SEQ ID NO	EC50 hGIP R [pM]	EC50 hGLP-1 R [pM]	EC50 hGLUC R [pM]
8	9.8	1.2	>1000000

9	3.1	1.1	>1000000
10	6.8	1.2	>1000000
11	6.7	1.0	>1000000
12	8.7	1.3	>1000000
13	8.1	0.4	>1000000
14	4.1	0.6	>1000000
15	7.3	1.4	>1000000
16	1.5	1.0	>1000000
17	1.8	1.1	>1000000
18	36.1	2.0	>1000000
19	2.2	1.2	>1000000
20	3.9	1.0	>1000000

Example 4: Pharmacokinetic testing

- 5 Pharmacokinetic profiles were determined as described in Methods. Calculated $T_{1/2}$ and C_{max} values are shown in Table 7.

Table 7. Pharmacokinetic profiles of exendin-4 derivatives.

SEQ ID NO	$T_{1/2}$ [h]	C_{max} [ng/ml]
8	0.3	1470.0

Table 8 : sequences

SEQ ID NO	sequence
1	H-G-E-G-T-F-T-S-D-L-S-K-Q-M-E-E-E-A-V-R-L-F-I-E-W-L-K-N-G-G-P-S-S-G-A-P-P-P-S-NH ₂
2	H-A-E-G-T-F-T-S-D-V-S-S-Y-L-E-G-Q-A-A-K-E-F-I-A-W-L-V-K-G-R-NH ₂
3	H-A-E-G-T-F-T-S-D-V-S-S-Y-L-E-G-Q-A-A-K((S)-4-Carboxy-4-hexadecanoylamino-butyl)-E-F-I-A-W-L-V-R-G-R-G
4	Y-A-E-G-T-F-I-S-D-Y-S-I-A-M-D-K-I-H-Q-Q-D-F-V-N-W-L-L-A-Q-K-G-K-K-N-D-W-K-H-N-I-T-Q
5	H-S-Q-G-T-F-T-S-D-Y-S-K-Y-L-D-S-R-R-A-Q-D-F-V-Q-W-L-M-N-T
6	Y-G-E-G-T-F-T-S-D-L-S-I-Q-M-E-E-E-A-V-R-L-F-I-E-W-L-K-N-G-G-P-S-S-G-A-P-P-P-S-NH ₂
7	Y-A-E-G-T-F-T-S-D-V-S-I-Y-L-E-G-Q-A-A-K-E-F-I-A-W-L-V-K-G-R
8	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-M-E-E-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
9	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-M-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
10	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
11	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-Nle-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
12	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-Nle-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-K-NH ₂
13	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-M(O)-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
14	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-M(O ₂)-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
15	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-E-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
16	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-E-E-A-V-Aib-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
17	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-E-E-A-V-(S)MeLys-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂

18	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-E-E-A-V-(R)MeLys-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
19	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-E-E-A-V-(S)MeOrn-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂
20	Y-Aib-E-G-T-F-T-S-D-L-S-I-Q-L-E-K-E-A-V-R-L-F-I-E-W-L-K-A-G-G-P-S-S-G-A-P-P-P-S-NH ₂

5. A compound according to any one of claims 1 - 4, wherein the peptidic compound exhibits a relative activity of at least 0.07% compared to that of GLP-1 (7-36) at the GLP-1 receptor.

5

6. A compound of any one of claims 1 - 5, wherein
X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and Met(O₂),
X16 represents an amino acid residue selected from Lys and Glu,
10 X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys, (R)Me-Lys and (S)Me-Orn, particularly Arg
X40 is absent.

7. A compound of any one of claims 1 - 5, wherein
15 X14 represents Leu,
X16 represents Lys,
X20 represents an amino acid residue selected from Aib, (S)Me-Lys, (R)Me-Lys and (S)Me-Orn,
X40 is absent.

20

8. A compound of any one of claims 1 - 5, wherein
X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and Met(O₂),
X16 represents an amino acid residue selected from Lys and Glu,
25 X20 represents Arg,
X40 is absent or represents Lys.

9. A compound of any one of claims 1 - 5, wherein
X14 represents an amino acid residue selected from Met, Leu, Nle, Met(O) and
30 Met(O₂),
X16 represents Lys,
X20 represents Arg,
X40 is absent or represents Lys.

- 10.A compound of any one of claims 1 - 5, wherein
X14 represents an amino acid residue selected from Met and Leu,
X16 represents Glu,
X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys,
5 (R)Me-Lys and (S)Me-Orn,
X40 is absent or represents Lys.
11. A compound of any one of claims 1 - 5, wherein
X14 represents an amino acid residue selected from Leu and Nle
10 X16 represents an amino acid residue selected from Lys and Glu,
X20 represents an amino acid residue selected from Arg, Aib, (S)Me-Lys,
(R)Me-Lys and (S)Me-Orn,
X40 is absent.
- 15 12.A compound of any one of claims 1 - 5, wherein
X14 represents an amino acid residue selected from Met, Met(O) and Met(O₂),
X16 represents an amino acid residue selected from Lys and Glu,
X20 represents Arg,
X40 is absent.
20
- 13.The compound of any one of claims 1 - 12, selected from the compounds of SEQ
ID NO: 8-20 or a salt or solvate thereof.
- 14.The compound of any one of claims 1 - 13, selected from the compounds of SEQ
25 ID NO: 10 - 14 and 16 - 20 as well as salts and solvates thereof.
- 15.The compound of any one of claims 1 - 14 for use in medicine, particularly in
human medicine.
- 30 16.The compound for use according to claim 15 which is present as an active agent
in a pharmaceutical composition together with at least one pharmaceutically
acceptable carrier.
- 17.The compound for use according to claim 15 or 16 together with at least one

additional therapeutically active agent, wherein the additional therapeutically active agent is selected from the series of Insulin and Insulin derivatives, GLP-1, GLP-1 analogues and GLP-1 receptor agonists, polymer bound GLP-1 and GLP-1 analogues, dual GLP1/glucagon agonists, PYY3-36 or analogues thereof, 5 pancreatic polypeptide or analogues thereof, Glucagon receptor agonists, GIP receptor agonists or antagonists, ghrelin antagonists or inverse agonists, Xenin and analogues thereof, DDP-IV inhibitors, SGLT2 inhibitors, dual SGLT2 / SGLT1 inhibitors, Biguanides Thiazolidinediones, dual PPAR agonists, Sulfonylureas, Meglitinides, alpha-glucosidase inhibitors, Amylin and Amylin analogues, GPR119 10 agonists, GPR40 agonists, GPR120 agonists, GPR142 agonists, systemic or low-absorbable TGR5 agonists, Cycloset, inhibitors of 11-beta-HSD, activators of glucokinase, inhibitors of DGAT, inhibitors of protein tyrosinephosphatase 1, inhibitors of glucose-6-phosphatase, inhibitors of fructose-1,6-bisphosphatase, inhibitors of glycogen phosphorylase, inhibitors of phosphoenol pyruvate 15 carboxykinase, inhibitors of glycogen synthase kinase, inhibitors of pyruvate dehydrogenase kinase, alpha2-antagonists, CCR-2 antagonists, modulators of glucose transporter-4, Somatostatin receptor 3 agonists, HMG-CoA-reductase inhibitors, fibrates, nicotinic acid and the derivatives thereof, nicotinic acid receptor 1 agonists, PPAR-alpha, gamma or alpha/gamma) agonists or 20 modulators, PPAR-delta agonists, ACAT inhibitors, cholesterol absorption inhibitors, bile acid-binding substances, IBAT inhibitors, MTP inhibitors, modulators of PCSK9, LDL receptor up-regulators by liver selective thyroid hormone receptor β agonists, HDL-raising compounds, lipid metabolism modulators, PLA2 inhibitors, ApoA-I enhancers, thyroid hormone receptor 25 agonists, cholesterol synthesis inhibitors, omega-3 fatty acids and derivatives thereof, active substances for the treatment of obesity, such as Sibutramine, Tesofensine, Orlistat, CB-1 receptor antagonists, MCH-1 antagonists, MC4 receptor agonists and partial agonists, NPY5 or NPY2 antagonists, NPY4 agonists, beta-3-agonists, leptin or leptin mimetics, agonists of the 5HT_{2c} 30 receptor, or the combinations of bupropione/naltrexone (CONTRAVE), bupropione/zonisamide (EMPATIC), bupropione/phentermine or pramlintide/metreleptin, QNEXA (Phentermine+ topiramate), lipase inhibitors, angiogenesis inhibitors, H3 antagonists, AgRP inhibitors, triple monoamine uptake inhibitors (norepinephrine and acetylcholine), MetAP2 inhibitors, nasal formulation

of the calcium channel blocker diltiazem, antisense against production of fibroblast growth factor receptor 4, prohibitin targeting peptide-1, drugs for influencing high blood pressure, chronic heart failure or atherosclerosis, such as angiotensin II receptor antagonists, ACE inhibitors, ECE inhibitors, diuretics, beta-blockers, calcium antagonists, centrally acting hypertensives, antagonists of the alpha-2-adrenergic receptor, inhibitors of neutral endopeptidase, thrombocyte aggregation inhibitors.

- 5
- 10
18. The compound for use according to claim 15 or 16 together with at least one additional therapeutically active agent, wherein the additional therapeutically active agent particularly is a GLP-1 agonist and/or insulin or an insulin analogue and/or a gastrointestinal peptide.
- 15
19. The compound for use according to any one of claims 15 - 18 for the treatment or prevention of hyperglycemia, type 2 diabetes, impaired glucose tolerance, type 1 diabetes, obesity, metabolic syndrome and neurodegenerative disorders, particularly for delaying or preventing disease progression in type 2 diabetes, treating metabolic syndrome, treating obesity or preventing overweight, for decreasing food intake, increase energy expenditure, reducing body weight, delaying the progression from impaired glucose tolerance (IGT) to type 2 diabetes; delaying the progression from type 2 diabetes to insulin-requiring diabetes; regulating appetite; inducing satiety; preventing weight regain after successful weight loss; treating a disease or state related to overweight or obesity; treating bulimia; treating binge eating; treating atherosclerosis, 20
- 25
- hypertension, IGT, dyslipidemia, coronary heart disease, hepatic steatosis, treatment of beta-blocker poisoning, use for inhibition of the motility of the gastrointestinal tract, useful in connection with investigations of the gastro-intestinal tract using techniques such as X-ray, CT- and NMR-scanning.
- 30
20. The compound for use according to any one of claims 15 - 19 for the treatment or prevention of hyperglycemia, type 2 diabetes, obesity.
- 35
21. A pharmaceutical composition comprising at least one compound according to any one of claims 1 - 14 or a physiologically acceptable salt or sovate of any of them, for use as a pharmaceutical.

22. A method of treating hyperglycemia, type 2 diabetes or obesity in a patient, the method comprising administering to the patient an effective amount of at least one compound of formula I according to any one of claims 1 - 14 and an effective amount of at least one other compound useful for treating hyperglycemia, type 2 diabetes or obesity.
- 5
23. A method as claimed in claim 21 wherein the effective amount of at least one compound of formula I according to claims 1 - 14 and the additional active ingredient are administered to the patient simultaneously.
- 10
24. A method as claimed in claim 21 wherein the effective amount of at least one compound of formula I according to claims 1 - 14 and the additional active ingredient are administered to the patient sequentially.
- 15

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2014/077337

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K14/605 C07K14/575
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal , WPI Data, Sequence Search , BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	wo 2008/081418 AI (COVX TECHNOLOGI ES IRELAND LTD [I E] ; BRADSHAW CURT [US] ; SAKAMURI SUKUM) 10 July 2008 (2008-07-10) cited in the applicati on claim 72; sequence Cl aim 26 -----	1-24
A	US 2011/237503 AI (ALSINA-FERNANDEZ JORGE [US] ET AL) 29 September 2011 (2011-09-29) cl aim 1; tabl es 1-2 -----	1-24
A	US 2011/166062 AI (DIMARCHI RICHARD D [US] ET AL) 7 July 2011 (2011-07-07) examples 25, 33; tabl es 3, 4; sequence 99 ----- -/- .	1-24

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 24 March 2015	Date of mailing of the international search report 01/04/2015
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Paresce, Donata
--	--

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2014/077337

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	wo 2012/167744 AI (BEIJING HANMI PHARMACEUTICAL CO LTD [CN] ; MA TAO [CN] ; XU SHUSEN [CN] ;) 13 December 2012 (2012-12-13) page 6 - page 8; claim 1 -----	1-24
A	wo 2011/024110 A2 (RINAT NEUROSCIENCE CORP [US] ; CHOU JOYCE CHING TSU [US] ; LIN CHIA-YANG) 3 March 2011 (2011-03-03) sequence 36 -----	1-24
A	PAN CLARK Q ET AL: "Design of a long acting peptide functioning as both a glucagon-1 like peptide-1 receptor agonist and a glucagon receptor antagonist", JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, US, vol . 281 , no. 18, 1 May 2006 (2006-05-01) , pages 12506-12515 , XP002529523 , ISSN: 0021-9258, DOI : 10.1074/JBC.M600127200 [retrieved on 2006-02-27] table 1 -----	1-24
A	wo 2013/164484 AI (ZEALAND PHARMA AS [DK]) 7 November 2013 (2013-11-07) claim 1; table 2 -----	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2014/077337

Patent document cited in search report	Publication date	Patent family member(s)	Publication date			
wo 2008081418	AI	10-07-2008	AU 2008203641 AI 10-07 -2008			
			BR PI0806308 A2 06-09 -2011			
			CA 2674112 AI 10-07 -2008			
			EP 2118131 AI 18-11 -2009			
			JP 5009376 B2 22-08 -2012			
			JP 2010514835 A 06-05 -2010			
			JP 2012184232 A 27-09 -2012			
			KR 20090096498 A 10-09 -2009			
			KR 20120083510 A 25-07 -2012			
			NZ 577686 A 25-11 -2011			
			US 2009098130 AI 16-04 -2009			
			wo 2008081418 AI 10-07 -2008			
			US 2011237503	AI	29-09-2011	AR 080592 AI 18-04 -2012
						AU 2011232597 AI 30-08 -2012
CA 2794664 AI 29-09 -2011						
CL 2012002635 AI 21-12 -2012						
CN 102821779 A 12-12 -2012						
CO 6630128 A2 01-03 -2013						
CR 20120457 A 16-10 -2012						
DO P2012000244 A 15-02 -2013						
EA 201270723 AI 30-01 -2013						
EC SP12012181 A 30-11 -2012						
EP 2552471 AI 06-02 -2013						
GT 201200263 A 05-12 -2013						
JP 2013523647 A 17-06 -2013						
KR 20120133402 A 10-12 -2012						
MA 34077 BI 05-03 -2013						
PE 05232013 AI 25-04 -2013						
SG 184289 AI 29-11 -2012						
TW 201201829 A 16-01 -2012						
US 2011237503 AI 29-09 -2011						
wo 2011119657 AI 29-09 -2011						
US 2011166062	AI	07-07-2011	AR 072159 AI 11-08 -2010			
			AU 2009274425 AI 28-01 -2010			
			CA 2729296 AI 28-01 -2010			
			CN 102105159 A 22-06 -2011			
			CO 6280539 A2 20-05 -2011			
			CR 11849 A 18-07 -2011			
			EA 201001801 AI 30-08 -2011			
			EC SP10010684 A 31-01 -2011			
			EP 2300035 A2 30-03 -2011			
			IL 209666 A 26-02 -2015			
			JP 2011524420 A 01-09 -2011			
			KR 20110040760 A 20-04 -2011			
			MA 32397 BI 01-06 -2011			
			NZ 589848 A 30-11 -2012			
			NZ 603445 A 28-03 -2014			
			PE 00562010 AI 26-01 -2010			
			SG 192405 AI 30-08 -2013			
			TW 201010729 A 16-03 -2010			
			US 2011166062 AI 07-07 -2011			
			wo 2010011439 A2 28-01 -2010			
wo 2012167744	AI	13-12-2012	CN 103764673 A 30-04-2014			
			EP 2718317 AI 16-04-2014			
			JP 2014516994 A 17-07-2014			

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2014/077337

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		KR 20140043780 A	10-04-2014
		US 2014162945 AI	12-06-2014
		WO 2012167744 AI	13-12-2012

W0 2011024110 A2	03-03-2011	AU 2010288191 AI	01-03-2012
		CA 2769835 AI	03-03-2011
		CN 102596229 A	18-07-2012
		EP 2470198 A2	04-07-2012
		JP 2011057670 A	24-03-2011
		KR 20120050460 A	18-05-2012
		RU 2012106200 A	10-10-2013
		US 2012148586 AI	14-06-2012
		WO 2011024110 A2	03-03-2011

W0 2013164484 AI	07-11-2013	AU 2013255752 AI	18-12-2014
		CA 2872315 AI	07-11-2013
		EP 2844670 AI	11-03-2015
		KR 20150006052 A	15-01-2015
		WO 2013164484 AI	07-11-2013
