



(86) Date de dépôt PCT/PCT Filing Date: 2006/04/27

(87) Date publication PCT/PCT Publication Date: 2006/11/02

(45) Date de délivrance/Issue Date: 2013/05/28

(85) Entrée phase nationale/National Entry: 2007/10/24

(86) N° demande PCT/PCT Application No.: US 2006/016087

(87) N° publication PCT/PCT Publication No.: 2006/116628

(30) Priorité/Priority: 2005/04/28 (US60/675,759)

(51) Cl.Int./Int.Cl. *A61K 47/48* (2006.01)

(72) Inventeurs/Inventors:

BIENIARZ, CHRISTOPHER, US;
WONG, JENNIFER, US;
LEFEVER, MARK, US;
KOSMEDER, JEROME W., US;
ASHWORTH-SHARPE, JULIA, US;
KERNAG, CASEY A., US


(73) Propriétaire/Owner:

VENTANA MEDICAL SYSTEMS, INC., US

(74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(54) Titre : CONJUGATS D'ANTICORPS PAR LIEURS HETEROBIFONCTIONNELS

(54) Title: ANTIBODY CONJUGATES VIA HETEROBIFUNCTIONAL PEG LINKERS



(57) Abrégé/Abstract:

Antibody/signal-generating moiety conjugates are disclosed that include an antibody covalently linked to a signal-generating moiety through a heterobifunctional polyalkyleneglycol linker. The disclosed conjugates show exceptional signal-generation in immunohistochemical and in situ hybridization assays on tissue sections and cytology samples. In one embodiment, enzyme-metallographic detection of nucleic acid sequences with hapten-labeled probes can be accomplished using the disclosed conjugates as a primary antibody without amplification.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 November 2006 (02.11.2006)

PCT

(10) International Publication Number
WO 2006/116628 A2

(51) International Patent Classification:

A61K 47/48 (2006.01)

(21) International Application Number:

PCT/US2006/016087

(22) International Filing Date: 27 April 2006 (27.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/675,759 28 April 2005 (28.04.2005) US

(71) Applicant (for all designated States except US): VENTANA MEDICAL SYSTEMS, INC. [US/US]; 1910 Innovation Park Drive, Tucson, Arizona 85755 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BIENIARZ, Christopher [US/US]; 2263 E. Quiet Canyon Drive, Tucson, Arizona 85718 (US). WONG, Jennifer [US/US]; 925 Lionel Court, Sparks, Nevada 89434 (US). LEFEVER, Mark [US/US]; 1022 Sonoran Heights Place, Tucson, Arizona 85748 (US). KOSMEDER, Jerome, W. [US/US];

5530 N. Moccasin Trail, Tucson, Arizona 85750 (US). ASHWORTH-SHARPE, Julia [US/US]; 10321 N. Bald Head Place, Tucson, Arizona 85737 (US). KERNAG, Casey, A. [US/US]; 1765 S. JONES BOULEVARD, #p-10, Tucson, Arizona 85713 (US).


(74) Agent: REHMS, Aden, A.; VENTANA MEDICAL SYSTEMS, INC., 1910 Innovation Park Drive, Tucson, Arizona 85755 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]


(54) Title: ANTIBODY CONJUGATES



A



B



C



D

(57) Abstract:

Antibody/signal-generating moiety conjugates are disclosed that include an antibody covalently linked to a signal-generating moiety through a heterobifunctional polyalkyleneglycol linker. The disclosed conjugates show exceptional signal-generation in immunohistochemical and in situ hybridization assays on tissue sections and cytology samples. In one embodiment, enzyme-metallographic detection of nucleic acid sequences with hapten-labeled probes can be accomplished using the disclosed conjugates as a primary antibody without amplification.

WO 2006/116628 A2

WO 2006/116628 A2

GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ANTIBODY CONJUGATES VIA HETEROBIFUNCTIONAL PEG LINKERS

Background of the Invention*1. Field*

The present invention relates to reagents and methods for detecting a molecule of interest in a biological sample. More particularly, the present invention relates to antibody conjugates and methods for using such conjugates to detect a molecule of interest in a biological sample such as a tissue section.

2. Background

Covalent conjugates of antibodies and signal-generating moieties can be used in immunoassays for detecting specific target molecules in biological samples. The antibody portion of such conjugates specifically binds to a target in the sample and the signal-generating moiety is utilized to provide a detectable signal that indicates the presence/and or location of the target. One type of conjugate that has become widely used, especially for immunohistochemical analysis, is a conjugate of an antibody and an enzyme (antibody-enzyme conjugate). A detectable signal is generated by adding a substrate to the sample and the enzyme portion of the conjugate converts the substrate to, for example, a colored, fluorescent or luminescent product at the site where the antibody portion is bound to its target.

Antibody-enzyme conjugates are typically prepared using polyfunctional (typically bifunctional) coupling reagents that are characterized by having at least two reactive groups, one of which is reacted with a functional group on the antibody and the other of which is reacted with a functional group on the enzyme. However, coupling can lead to inactivation of either or both of the antibody and the enzyme due to steric effects or because the coupling reagents react with functional groups located on portions of the enzyme or antibody that are critical for their function or specificity.

An approach to minimizing loss of antibody specificity and enzyme activity is to use a coupling scheme that is specific to particular amino acid residues on either or both of the antibody and the enzyme that are not associated with their functions. This approach is exemplified by the method for Fc-specific conjugation as described in U.S. Patent No. 5,191,066.

In this method, sulfhydryl groups (thiol groups) are introduced specifically to a glycosylated region of the Fc portion of an antibody and used along with a linker molecule to covalently attach an enzyme to the antibody. Since the Fc portion is not involved with the specific binding properties of the antibody, such conjugates retain greater specificity, which increases the detectable signal for a particular target molecule of interest and lowers background due to non-specific binding.

Although site specific conjugation can be used to help minimize loss of antibody specificity and enzyme activity due to loss of critical functional groups, such methods do not address loss of antibody specificity and enzyme activity that arise from steric effects such as those steric effects due to aggregation of multiple conjugates and from interactions between the antibody and the enzyme(s) in a conjugate. Detrimental steric effects also can arise due to unintended cross-linking between multiple enzymes,

antibodies and/or conjugates, which occurs during preparation of a conjugate composition.

One approach to minimizing loss of antibody specificity and enzyme activity due to steric effects is to increase the length of the coupling reagent in order that the antibody and enzyme are separated by a greater distance. This approach is exemplified by the methods and conjugation reagents disclosed in U.S. Patent No. 5,053,520. In this method, heterobifunctional linkers having extended alkyl, cycloalkyl, alkyl-cycloalkyl and aromatic portions are used to couple an antibody to an enzyme(s). Although such linkers contain more atoms and should provide greater separation between an antibody and an enzyme(s), it is believed that the hydrophobic nature of such linkers increases detrimental aggregation of conjugates in aqueous solution due to hydrophobic effects. In addition, such linkers are flexible enough to permit detrimental intra-conjugate interactions between the antibody and the enzyme(s) as a conjugate collapses in on itself to minimize its size due to hydrophobic effects.

An attempt to minimize detrimental aggregation between conjugates is described in U.S. Patent No. 4,810,638, which describes the use of homo-bifunctional, bis-maleimidopolyalkylene glycol linkers to prepare antibody-enzyme conjugates. However, use of such homo-bifunctional linkers can lead to cross-linking of antibodies, enzymes and/or conjugates during preparation of the conjugates. Cross-linking increases the average size and counteracts to some extent the increased water solubility imparted by using the glycol linker. Furthermore, cross-linking leads to lower monodispersity in a conjugate composition, which can have detrimental effects on consistency of results, especially in tissue and cell samples where detection of a target with a conjugate may be limited by diffusion through cell membranes.

Some heterobifunctional polyethylene glycol linkers are known, but there are no known attempts to use them as coupling reagents for forming antibody-enzyme conjugates. Rather, as disclosed in Chen et al. (Chen et al., "The use of bifunctional polyethylene glycol derivatives for coupling of proteins to and cross-linking of collagen matrices," *J. Mater. Sci. Mater. Med.*, 13: 1029-1035, 2002), such agents have been utilized to prepare degradable matrices to which active proteins are linked for the purposes of tissue engineering.

From the standpoint of increasing the signal generated by a given antibody conjugate it is desirable to conjugate multiple enzymes to a single antibody. However, as the number of enzymes linked to a single antibody increases, the likelihood increases that conjugate function will be impaired for steric reasons due to crowding of multiple enzymes around the single antibody. One approach to minimizing crowding of enzymes is to employ a scaffold to provide separation between enzymes and between enzymes and antibodies or antibody fragments. U.S. Patent Nos. 6,252,053 and 6,613,564, for example, describe the use of polylysine or dextran scaffolds to increase separation between enzymes, while still effectively increasing the number of enzyme molecules per specific binding component [specifically F(ab')₂ fragments]. While the approach described in these patents does increase the average number of signal-generating moieties per specific-binding component, the use of a polymeric scaffold (typically of low mono-dispersity) increases background and decreases reproducibility. The high molecular weight (typically greater >1 MDa) of such constructs can hinder diffusion and tissue/cell penetrability is diminished, thereby reducing signal.

What is needed, therefore, is an antibody/signal-generating conjugate composition that overcomes at least the described limitations of prior approaches. In

particular, antibody conjugates of enzyme (and methods of making the same) that are smaller and yet retain the high signal generating capacity of larger scaffolded conjugates are desirable.


Summary of the Invention

Antibody conjugates with signal generating moieties are disclosed, as are methods for making and using the conjugates. The disclosed antibody conjugates exhibit superior performance for detection of molecules of interest in biological samples, especially for detection of such molecules in tissue sections and cytology samples. In particular, disclosed antibody-enzyme conjugates retain high amounts of antibody specificity and enzymatic activity, and thereby provide more intense staining with less background than conjugates currently used for detection of antigens in biological samples.

In one aspect, a conjugate is disclosed that includes an antibody covalently linked to a signal-generating moiety through a heterobifunctional polyalkyleneglycol linker such as a heterobifunctional polyethyleneglycol (PEG) linker. In one embodiment, a disclosed conjugate includes an antibody and a signal-generating moiety covalently linked by a heterobifunctional PEG linker that includes a combination of two different reactive groups selected from a carbonyl-reactive group, an amine-reactive group, a thiol-reactive group and a photo-reactive group. In particular embodiments, the PEG linker includes a combination of a thiol reactive group and an amine-reactive group or a combination of a carbonyl-reactive group and an thiol-reactive group. In more particular embodiments, the thiol reactive group includes a maleimide group, the amine


reactive group includes an active ester and the carbonyl-reactive group includes a hydrazine derivative.

In even more particular embodiments, the disclosed conjugate has the general formula:



wherein Ab is an antibody, SM is a signal-generating moiety (for example, an enzyme) and $n = 1$ to 50 (such as $n = 2$ to 30, $n = 2$ to 20 or $n = 4$ to 12) and $s = 1$ to 10 (such as $s = 2$ to 6 or $s = 3$ to 4).

In other even more particular embodiments, a disclosed conjugate has the formula:



wherein Ab is an antibody, SM is a signal-generating moiety (such as an enzyme), $m = 1$ to 50 (such as $m = 2$ to 30, $m = 2$ to 20 or $m = 4$ to 12) and $t = 1$ to 10 (such as $t = 2$ to 6 or $t = 3$ to 4). In some instances, the hydrazide group of the PEG linker is bonded to the

carbon of an aldehyde group formed in the glycosylated portion of the antibody by oxidation.

In another aspect, methods for making the disclosed conjugates are provided. In one embodiment a method of making an antibody conjugate includes forming a thiolated antibody from an antibody; reacting a signal-generating moiety having an amine group with a PEG maleimide/active ester bifunctional linker to form an activated signal-generating moiety; and reacting the thiolated antibody with the activated signal-generating moiety to form the conjugate of the antibody and the signal-generating moiety. The thiolated antibody can be formed by reduction of intrinsic cystine bridges of the antibody with a reductant or can be formed by reacting the antibody with a reagent that introduces a thiol to the antibody.

In another embodiment, a method for making a disclosed antibody conjugate includes reacting an antibody with an oxidant to form an aldehyde-bearing antibody; reacting the aldehyde-bearing antibody with a PEG maleimide/hydrazide bifunctional linker to form a thiol-reactive antibody; and reacting the thiol-reactive antibody with a thiolated signal-generating moiety to form the antibody-signal-generating moiety conjugate. In a particular embodiment, reacting the antibody with an oxidant to form the aldehyde-bearing antibody includes oxidizing (such as with periodate, bromine or iodine) a glycosylated region of the antibody to form the aldehyde-bearing antibody.

In another aspect, PEG maleimide/hydrazide bifunctional linkers are disclosed that can be used in the disclosed methods to provide disclosed conjugates. In yet another aspect, methods are disclosed for detecting molecules in biological samples using disclosed conjugates. These and additional aspects, embodiments and features of

the disclosure will become apparent from the detailed description and examples that follow.

Brief Description of the Drawings

FIG. 1 is series of images of tissue sections immunohistochemically stained for Ki67 with a disclosed conjugate, in comparison to a scaffolded conjugate, both before and after storage at 45 °C for 7 days.

FIG. 2 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of bcl-2.

FIG. 3 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of CD15.

FIG. 4 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of CD20.

FIG. 5 is a series of images comparing the staining intensity of a disclosed conjugate and two scaffolded conjugates for immunohistochemical staining of CD23.

FIG. 6 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of CD57.

FIG. 7 is a series of images comparing the staining intensity of a disclosed conjugate and two scaffolded conjugates for immunohistochemical staining of cerbB2.

FIG. 8 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of cyclin D1.

FIG. 9 is a series of images comparing the staining intensity of a disclosed conjugate and two scaffolded conjugates for immunohistochemical staining of EGFR.

FIG. 10 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of ER.

FIG. 11 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of p53.

FIG. 12 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of PR.

FIG. 13 is a pair of images comparing the staining intensity of a disclosed conjugate and a scaffolded conjugate for immunohistochemical staining of PSA.

FIG. 14 is diagram outlining a scheme for enzyme metallographic detection of binding of a hapten-labeled nucleic acid probe to a target nucleic acid sequence that utilizes a disclosed antibody-enzyme conjugate.

FIG. 15 is a series of images of tissue sections treated for enzyme metallographic ISH detection of a nucleic acid sequence using a disclosed conjugate and a scaffolded conjugate, before and after storage both at 37 °C for 7 days and at 45 °C for 7 days.

FIG. 16 is a pair of graphs comparing the stability of a disclosed conjugate and a scaffolded conjugate in an enzyme metallographic detection scheme.

FIG. 17 is size-exclusion chromatogram comparing the effect of variations of antibody reduction time on the MW profile of a disclosed conjugate.

FIG. 18 is a size-exclusion chromatogram comparing the effect of variations of linker size and type on the MW profile of disclosed conjugates.

FIG. 19 is a series of images comparing the staining intensity of several disclosed conjugates compared to a conjugate prepared with an extended-length non-PEG linker.

FIG. 20 is a size-exclusion chromatogram comparing the effect of variations of linker excess on the MW profile of a disclosed conjugate.

FIG. 21 is a size-exclusion chromatogram comparing the effect of variations of horseradish peroxidase concentrations on the MW profile of a disclosed conjugate.

FIG. 22 is a size-exclusion chromatogram comparing the effect of variations of the ratio of antibody to horseradish peroxidase on the MW profile of a disclosed conjugate.

Detailed Description of Several Illustrative Embodiments

Further aspects of the invention are illustrated by the following non-limiting examples, which proceed with respect to the abbreviations and terms defined below.

I. Abbreviations

2-ME – 2-mercaptoethanol

2-MEA – 2-mercaptoethylamine

Ab - antibody

ALP – alkaline phosphatase

BSA – bovine serum albumin

DTE – dithioerythritol (*cis*-2,3-dihydroxy-1,4-dithiolbutane)

DTT – dithiothreitol (*trans*-2,3-dihydroxy-1,4-dithiolbutane)

EGFR – epidermal growth factor receptor

ER – estrogen receptor

HRP – horseradish peroxidase

IHC - immunohistochemistry

ISH –*in situ* hybridization

MAL – maleimide

NHS – *N*-hydroxy-succinimide

PEG – polyethylene glycol

PR – progesterone receptor

SAMSA – *S*-Acetylmercaptosuccinic acid

SATA – *N*-succinimidyl *S*-acetylthioacetate

SATP – Succinimidyl acetyl-thiopropionate

SM –signal-generating moiety

SMPT – Succinimidylloxycarbonyl- α -methyl- α -(2-pyridyldithio)toluene

SPDP – *N*-Succinimidyl 3-(2-pyridyldithio)propionate

TCEP - tris(carboxyethyl)phosphine

II. Terms

The terms “a,” “an” and “the” include both singular and plural referents unless the context clearly indicates otherwise.


The term “antibody” collectively refers to immunoglobulins or immunoglobulin-like molecules (including IgA, IgD, IgE, IgG and IgM, combinations thereof, and similar molecules produced during an immune response in any vertebrate, for example, in mammals such as humans, goats, rats, rabbits and mice) and antibody fragments that specifically bind to a molecule of interest (or a group of highly similar molecules of interest) to the substantial exclusion of binding to other molecules (for example, antibodies and antibody fragments that have a binding constant for the molecule of interest that is at least 10^3 M^{-1} greater, 10^4 M^{-1} greater or 10^5 M^{-1} greater than a binding constant for other molecules in a biological sample). Antibody fragments include

proteolytic antibody fragments [such as F(ab')₂ fragments, Fab' fragments, Fab'-SH fragments and Fab fragments as are known in the art], recombinant antibody fragments (such as sFv fragments, dsFv fragments, bispecific sFv fragments, bispecific dsFv fragments, diabodies, and triabodies as are known in the art), and camelid antibodies (see, for example, U.S. Patent Nos. 6,015,695; 6,005,079; 5,874,541; 5,840,526; 5,800,988; and 5,759,808).

The phrase "molecule of interest" refers to a molecule for which the presence, location and/or concentration is to be determined. Examples of molecules of interest include proteins and nucleic acid sequences labeled with haptens.

III. Overview

In one aspect, an antibody/signal-generating moiety conjugate is disclosed that includes an antibody covalently linked to a signal-generating moiety through a heterobifunctional polyalkyleneglycol linker having the general structure shown below:



wherein A and B include different reactive groups, x is an integer from 2 to 10 (such as 2, 3 or 4), and y is an integer from 1 to 50, for example, from 2 to 30 such as from 3 to 20 or from 4 to 12. One or more hydrogen atoms can be substituted for additional functional groups such as hydroxyl groups, alkoxy groups (such as methoxy and ethoxy), halogen atoms (F, Cl, Br, I), sulfato groups and amino groups (including mono- and di-substituted amino groups such as dialkyl amino groups).

A and B of the linker can independently include a carbonyl-reactive group, an amine-reactive group, a thiol-reactive group or a photo-reactive group, but are not the same. Examples of carbonyl-reactive groups include aldehyde- and ketone-reactive

groups like hydrazine derivatives and amines. Examples of amine-reactive groups include active esters such as NHS or sulfo-NHS, isothiocyanates, isocyanates, acyl azides, sulfonyl chlorides, aldehydes, glyoxals, epoxides, oxiranes, carbonates, aryl halides, imidoesters, anhydrides and the like. Examples of thiol-reactive groups include non-polymerizable Michael acceptors, haloacetyl groups (such as iodoacetyl), alkyl halides, maleimides, aziridines, acryloyl groups, vinyl sulfones, benzoquinones, aromatic groups that can undergo nucleophilic substitution such as fluorobenzene groups (such as tetra and pentafluorobenzene groups), and disulfide groups such as pyridyl disulfide groups and thiols activated with Ellman's reagent. Examples of photo-reactive groups include aryl azide and halogenated aryl azides. Additional examples of each of these types of groups will be apparent to those skilled in the art. Further examples and information regarding reaction conditions and methods for exchanging one type of reactive group for another are provided in Hermanson, "Bioconjugate Techniques," Academic Press, San Diego, 1996.

In a particular embodiment, a thiol-reactive group is other than vinyl sulfone.

In some embodiments, a thiol-reactive group of the heterobifunctional linker is covalently linked to the antibody and an amine-reactive group of the heterobifunctional linker is covalently linked to the signal-generating moiety, or vice versa. For example, a thiol-reactive group of the heterobifunctional linker can be covalently linked to a cysteine residue (such as formed by reduction of a cystine bridge) of the antibody or a thiol-reactive group of the heterobifunctional linker can be covalently linked to a thiol group that is introduced to the antibody, and the amine-reactive group is covalently linked to the signal-generating moiety.

Alternatively, an aldehyde-reactive group of the heterobifunctional linker can be covalently linked to the antibody and an amine-reactive group of the heterobifunctional linker can be covalently linked to the signal-generating moiety, or vice versa. In a particular embodiment, an aldehyde-reactive group of the heterobifunctional linker can be covalently linked to an aldehyde formed on a glycosylated portion of an antibody, and the amine-reactive group is covalently linked to the signal-generating moiety.

In yet other embodiments, an aldehyde-reactive group of the heterobifunctional linker is covalently linked to the antibody and a thiol-reactive group of the heterobifunctional linker is covalently linked to the signal-generating moiety, or vice versa.


Examples of signal-generating moieties include enzymes (such as horseradish peroxidase, alkaline phosphatase, acid phosphatase, glucose oxidase, β -galactosidase, β -glucuronidase or β -lactamase), fluorescent molecules (such as fluoresceins, coumarins, BODIPY dyes, resorufins, and rhodamines; additional examples can be found in *The Handbook — A Guide to Fluorescent Probes and Labeling Technologies*, Invitrogen Corporation, Eugene, OR), detectable constructs (such as fluorescent constructs like quantum dots, which can be obtained, for example, from Invitrogen Corporation, Eugene, OR; see, for example, U.S. Patent Nos. 6,815,064, 6,682,596 and 6,649,138),

metal chelates (such as DOTA and DPTA chelates of radioactive or paramagnetic metal ions like Gd^{3+}) and liposomes (such as liposomes sequestering fluorescent molecules).

When the signal-generating moiety includes an enzyme, a chromagenic compound, fluorogenic compound, or luminogenic compound is used in combination with the enzyme to generate a detectable signal (A wide variety of such compounds are


available, for example, from Molecular Probes, Inc., Eugene OR). Particular examples of chromogenic compounds include di-aminobenzidine (DAB), 4-nitrophenylphosphate (pNPP), fast red, bromochloroindolyl phosphate (BCIP), nitro blue tetrazolium (NBT), BCIP/NBT, fast red, AP Orange, AP blue, tetramethylbenzidine (TMB), 2,2'-azino-di-[3-ethylbenzothiazoline sulphonate] (ABTS), o -dianisidine, 4-chloronaphthol (4-CN), nitrophenyl- β -D-galactopyranoside (ONPG), o-phenylenediamine (OPD), 5-bromo-4-chloro-3-indolyl- β -galactopyranoside (X-Gal), methylumbelliferyl- β -D-galactopyranoside (MU-Gal), p-nitorphenyl- α -D-galactopyranoside (PNP), 5-bromo-4-chloro-3-indolyl- β -D-glucuronide (X-Gluc), 3-amino-9-ethyl carbazol (AEC), fuchsin, iodonitrotetrazolium (INT), tetrazolium blue and tetrazolium violet.

In particular embodiments the heterobifunctional linker of the conjugate has the formula:




wherein A and B include different reactive groups as before, x and y are as before, and X and Y are spacer groups, for example, spacer groups having between 1 and 10 carbons such as between 1 and 6 carbons or between 1 and 4 carbons, and optionally containing one or more amide linkages, ether linkages, ester linkages and the like. Spacers X and Y can be the same or different, and can be straight-chained, branched or cyclic (for example, aliphatic or aromatic cyclic structures), and can be unsubstituted or substituted. Functional groups that can be substituents on a spacer include carbonyl groups, hydroxyl groups, halogen (F, Cl, Br and I) atoms, alkoxy groups (such as methoxy and ethoxy), nitro groups, and sulfato groups.

In other particular embodiments, the heterobifunctional linker is a heterobifunctional polyethylene glycol linker having the formula:




wherein $n = 1$ to 50, for example, $n = 2$ to 30 such as $n = 3$ to 20 or $n = 4$ to 12. In more particular embodiments, a carbonyl of a succinimide group of this linker is covalently linked to an amine group on the signal-generating moiety and a maleimide group of the linker is covalently linked to a thiol group of the antibody, or vice versa. In other more particular embodiments, an average of between about 1 and about 10 signal moieties are covalently linked to an antibody.

In some particular embodiments, the heterobifunctional linker has the formula:




wherein $m = 1$ to 50, for example, $m = 2$ to 30 such as $m = 3$ to 20 or 4 to 12. In some more particular embodiments, a hydrazide group of the linker is covalently linked to an aldehyde group of the antibody and a maleimide group of the linker is covalently linked to a thiol group of the signal-generating moiety, or vice versa. In even more particular embodiments, the aldehyde group of the antibody is an aldehyde group formed in an Fc portion of the antibody by oxidation of a glycosylated region of the Fc portion of the antibody. In still other more particular embodiments, an average of between about 1 and about 10 signal-generating moieties are covalently linked to the antibody, such signal-generating moieties including enzymes, quantum dots and liposomes.

In other particular embodiments, a heterobifunctional PEG-linked antibody-signal-generating moiety conjugate comprises a conjugate having the formula:



wherein Ab is an antibody, SM is a signal-generating moiety and $n = 1$ to 50 (such as $n = 2$ to 30, $n = 2$ to 20 or $n = 4$ to 12) and $s = 1$ to 10 (such as $s = 2$ to 6 or $s = 3$ to 4).

In still other embodiments, a heterobifunctional PEG-linked antibody-signal-generating moiety conjugate comprises a conjugate having the formula:



wherein Ab is an antibody, SM is a signal-generating moiety, $m = 1$ to 50 (such as $m = 2$ to 30, $m = 2$ to 20 or $m = 4$ to 12) and $t = 1$ to 10 (such as $t = 2$ to 6 or $t = 3$ to 4).


Although the antibody used in the disclosed conjugates can specifically bind any particular molecule or particular group of highly similar molecules, in particular embodiments, the antibody comprises an anti-hapten antibody (which can be used to detect a hapten-labeled probe sequence directed to a nucleic acid sequence of interest) or an antibody the specifically binds to a particular protein or form of a particular protein (such as a phosphorylated form of a protein) that may be present in a sample. Haptens

are small organic molecules that are specifically bound by antibodies, although by themselves they will not elicit an immune response in an animal and must first be linked to a larger carrier molecule such as a protein or a poly-nucleic acid to generate an immune response. Examples of haptens include di-nitrophenol, biotin, and digoxigenin. In still other particular embodiments, the antibody comprises an anti-antibody antibody that can be used as a secondary antibody in an immunoassay. For example, the antibody can comprise an anti-IgG antibody such as an anti-mouse IgG antibody, an anti-rabbit IgG antibody or an anti-goat IgG antibody.

The disclosed antibody conjugates can be utilized for detecting molecules of interest in any type of binding immunoassay, including immunohistochemical binding assays. In one embodiment, the disclosed conjugates are used as a labeled primary antibody in an immunoassay, for example, a primary antibody directed to a particular molecule or a hapten-labeled molecule. Or, where the molecule of interest is multi-epitopic a mixture of conjugates directed to the multiple epitopes can be used. In another embodiment, the disclosed conjugates are used as secondary antibodies in an immunoassay (for example, directed to a primary antibody that binds the molecule of interest; the molecule of interest can be bound by two primary antibodies in a sandwich-type assay when multi-epitopic). In yet another embodiment, mixtures of disclosed conjugates are used to provide further amplification of a signal due to a molecule of interest bound by a primary antibody (the molecule of interest can be bound by two primary antibodies in a sandwich-type assay). For example, a first conjugate in a mixture is directed to a primary antibody that binds a molecule of interest and a second conjugate is directed to the antibody portion of the first conjugate, thereby localizing more signal-generating moieties at the site of the molecule of interest. Other types of

assays in which the disclosed conjugates can be used are readily apparent to those skilled in the art.

In another aspect, a heterobifunctional linker is disclosed having the formula:



wherein $m = 1$ to 50, for example, $m = 2$ to 30 such as $m=3$ to 20 or $m = 4$ to 12.


In yet another aspect, a method is disclosed for preparing an antibody-signal-generating moiety conjugate, the method including forming a thiolated antibody from an antibody; reacting a signal-generating moiety having an amine group with a PEG maleimide/active ester bifunctional linker to form an activated signal-generating moiety; and reacting the thiolated antibody with the activated signal-generating moiety to form the antibody-signal-generating moiety conjugate. A thiolated antibody can be formed by reacting the antibody with a reducing agent to form the thiolated antibody, for example, by reacting the antibody with a reducing agent to form a thiolated antibody having an average number of thiols per antibody of between about 1 and about 10. The average number of thiols per antibody can be determined by titration. Examples of reducing agents include reducing agents selected from the group consisting of 2-mercaptoethanol, 2-mercaptoethylamine, DTT, DTE and TCEP, and combinations thereof. In a particular embodiment the reducing agent is selected from the group consisting of DTT and DTE, and combinations thereof, and used at a concentration of between about 1 mM and about 40 mM.

Alternatively, forming the thiolated antibody includes introducing a thiol group to the antibody. For example, the thiol group can be introduced to the antibody by

reaction with a reagent selected from the group consisting of 2-Iminothiolane, SATA, SATP, SPDP, *N*-Acetylhomocysteinethiolactone, SAMSA, and cystamine, and combinations thereof (see, for example, Hermanson, "Bioconjugate Techniques," Academic Press, San Diego, 1996).

more particular embodiment, introducing the thiol group to the antibody includes reacting the antibody with an oxidant (such as periodate, I_2 , Br_2 , or a combination thereof) to convert a sugar moiety of the antibody into an aldehyde group and then reacting the aldehyde group with cystamine.


In other particular embodiments, reacting the signal-generating moiety with a PEG maleimide/active ester bifunctional linker to form an activated signal-generating moiety includes reacting the signal-generating moiety with a PEG maleimide/active ester having the formula:



wherein $n = 1$ to 50 , for example, $n = 2$ to 30 such as $n=3$ to 20 or $n = 4$ to 12 . The signal-generating moiety can, for example, be an enzyme (such as horseradish peroxidase or alkaline phosphatase).

In a further aspect, a method is disclosed for preparing an antibody-signal-generating moiety conjugate that includes reacting an antibody with an oxidant to form an aldehyde-bearing antibody; reacting the aldehyde-bearing antibody with a PEG maleimide/hydrazide bifunctional linker to form a thiol-reactive antibody; and reacting the thiol-reactive antibody with a thiolated signal-generating moiety to form the antibody-signal-generating moiety conjugate. In a particular embodiment, reacting the

antibody with an oxidant to form the aldehyde-bearing antibody includes oxidizing (such as with periodate) a glycosylated region of the antibody to form the aldehyde-bearing antibody. In a more particular embodiment, reacting an antibody with an oxidant to form an aldehyde-bearing antibody includes introducing an average of between about 1 and about 10 aldehyde groups per antibody. In another more particular embodiment, the PEG maleimide/hydrazide bifunctional linker used in the method has the formula:



wherein $m = 1$ to 50, for example, $m = 2$ to 30 such as $m=3$ to 20 or $m = 4$ to 12.

A thiolated signal-generating moiety can be formed from a signal-generating moiety by reacting the signal-generating moiety (such as an enzyme) with a reducing agent (such as a reducing agent selected from the group consisting of 2-mercaptoethanol, 2-mercaptoethylamine, DTT, DTE and TCEP, and combinations thereof) to form the thiolated signal-generating moiety, or by introducing a thiol group (for example, by reacting a signal generating moiety with a reagent selected from the group consisting of 2-Iminothiolane, SATA, SATP, SPDP, *N*-Acetylhomocysteinethiolactone, SAMSA, and cystamine, and combinations thereof).

In a still further aspect, a method is disclosed for detecting a molecule of interest in a biological sample that includes contacting the biological sample with a heterobifunctional PEG-linked antibody-signal-generating moiety conjugate and detecting a signal generated by the antibody-signal-generating moiety conjugate. The biological sample can be any sample containing biomolecules (such as proteins, nucleic

acids, lipids, hormones etc.), but in particular embodiments, the biological sample includes a tissue section (such as obtained by biopsy) or a cytology sample (such as a Pap smear or blood smear). In a particular embodiment, the heterobifunctional PEG-linked antibody-signal-generating moiety conjugate includes an antibody covalently linked to an enzyme such as horseradish peroxidase or alkaline phosphatase. In other particular embodiments, the heterobifunctional PEG-linked antibody-signal-generating moiety conjugate includes an antibody covalently linked to a detectable construct or a liposome.

In a more particular method, the signal-generating moiety comprises an enzyme such as alkaline phosphatase and the method further comprises contacting the biological sample with a water-soluble metal ion and a redox-inactive substrate of the enzyme that is converted to a redox-active agent by the enzyme, which redox-active agent reduces the metal ion causing it to precipitate. (see, for example, co-pending U.S. Patent Application No. 11/015,646, filed December 20, 2004, PCT Publication No. 2005/003777 and U.S. Patent Application Publication No. 2004/0265922).


In another particular embodiment the signal-generating moiety comprises an oxido-reductase enzyme (such as horseradish peroxidase) and the method further comprise contacting the biological sample with a water soluble metal ion, an oxidizing agent and a reducing agent (see, for example, U.S. Patent No. 6,670,113).

IV. Examples

The following non-limiting examples are provided to further illustrate certain aspects of the invention.


A. Preparation of antibody-signal-generating moiety conjugates using maleimide PEG active esters.

In one embodiment, a disclosed antibody signal-generating moiety conjugate is prepared according to the processes described in schemes 1 to 3 below, wherein the heterobifunctional polyalkylene glycol linker is a polyethylene glycol linker having an amine-reactive group (active ester) and a thiol-reactive group (maleimide). As shown in Scheme 1, a signal-generating moiety (such as an enzyme or a quantum dot) that has one or more available amine groups is reacted with an excess of the linker to form an activated signal-generating moiety.




Scheme 1

Thiol groups are introduced to the antibody by treating the antibody with a reducing agent such as DTT as shown in Scheme 2. For a mild reducing agent such as DTE or DTT, a concentration of between about 1 mM and about 40 mM (for example, a concentration of between about 5 mM and about 30 mM or between about 15 mM and about 25 mM) is utilized to introduce a limited number of thiols (such as between about 2 and about 6) to the antibody while keeping the antibody intact (which can be determined by size-exclusion chromatography).

**Scheme 2**


The components produced according to Schemes 1 and 2 are then combined to give a conjugate as shown in Scheme 3.

**Scheme 3**


Although Schemes 1-3 illustrate an optimal process for maleimide PEG active esters, wherein the signal-generating moiety is first activated by reacting an amine group

with the active ester of the linker to form an activated signal-generating moiety, it is also possible to first activate the antibody by reacting either an amine or a thiol on the antibody with the linker and then react the activated antibody with the signal generating moiety [having either a thiol or an amine to react with the remaining reactive group on the linker as appropriate]. Furthermore, although 3 signal-generating moieties are shown in Scheme 3, it is possible to link multiple antibodies to a single signal-generating moiety or any number of signal-generating moieties to a single antibody.

In an alternative embodiment, an antibody is activated for conjugation and then conjugated to a signal-generating moiety as shown in Schemes 4 and 5 below. In Scheme 4, the antibody is activated instead of the signal generating moiety as was shown in Scheme 1. In the particular embodiment of scheme 4, a sugar moiety (such as located in a glycosylated region of the Fc portion of the antibody) is first oxidized to provide an aldehyde group, which is then reacted with an aldehyde-reactive group of the linker (such as a hydrazide group of the illustrated maleimide/hydrazide PEG linker).

**Scheme 4**

Then, as shown in Scheme 5, a thiol-reactive group of the linker portion of the activated antibody (such as a maleimide group as illustrated) is then reacted with a thiol group on the signal generating moiety. Again, the process can be reversed, wherein the linker is first reacted with an aldehyde group on the signal-generating moiety (formed, for example, by oxidation of a sugar moiety) to form an activated signal generating moiety, and then the activated signal generating moiety can be reacted with a thiol group on the antibody. Furthermore, although Schemes 4 and 5 show only a single linker joining a single antibody and a single signal-generating moiety, it is to be understood that it is also possible to link multiple signal generating moieties to a single antibody or to link several antibodies to a one signal-generating moiety.



Scheme 5

B. Preparation of antibody-horseradish peroxidase conjugates

Activation of HRP

HRP can, for example, be activated for conjugation by treatment with a 100-fold molar excess of a bifunctional PEG linker having a maleimide group and an active ester group (for example, the MAL-PEG₄-NHS, MAL-PEG₈-NHS or MAL-PEG₁₂-NHS linkers available from Quanta Biodesign, Powell, OH) at ambient temperature (23 – 25 °C) for 60 minutes. After purification across a Superdex 200 10/300 GL column, excess linker-free HRP, typically with five to seven maleimides, is obtained with a 100-fold molar excess. An exemplary procedure is outlined below for production of an HRP antibody conjugate using a MAL-PEG₄-NHS linker. The number of maleimide groups on an activated HRP can be determined by the method described in detail in Example D.

HRP-PEG₄-maleimide(1): To a 4 mL amber vial was added 78.8 mg (100 eq.) of MAL-dPEG₄TM NHS ester (Quanta Biodesign, Powell, OH, F.W. = 513.50), followed by 2.46 mL (61.5 mg, 1.53 µM) of HRP (Horseradish Peroxidase, Pierce, Rockford, IL Lot FJ925901) as a 25 mg / mL solution in 0.1 M sodium phosphate, pH 7.5. The vial was then placed on an autorotator in the dark at ambient temperature (23 – 25 °C), and the amide bond forming reaction was allowed to proceed for 1 hour. A 400 µl aliquot was then removed for purification, and the remainder of the solution was temporarily stored at 4 °C. Pure HRP-PEG₄-maleimide was then obtained by fractionating the sample on an Akta PurifierTM fitted with a Superdex 10/300 column (Amersham, Piscataway, NJ) eluted with 0.1 M sodium phosphate, pH 7.5 at 1.0 mL / min. The HRP containing fractions were pooled to give 2.0 ml of a 4.52 mg / mL solution of HRP-

PEG₄-maleimide (90 % recovery) as measured by UV/VIS spectrophotometry using an extinction coefficient at 280 nm of a 1% solution (pH 6.5) of 6.52.

Introduction of Thiols to Antibodies

To activate an antibody, for example, an anti-mouse IgG or anti-rabbit IgG antibody, for conjugation an antibody can be incubated with 25 mmol DTT at ambient temperature (23 – 25 °C) for 25 minutes. After purification across a PD-10 SE column, DTT-free antibody, typically with two to six free thiols, is obtained (Scheme2). The exemplary procedure outlined for preparing goat anti-mouse IgG thiol is generally applicable to other antibodies. The number of thiols per antibody can be determined by the thiol assay described in Example D.

Goat anti-Mouse IgG-thiol (2): To a 8 mL amber vial was added 4.11 mL of Goat-anti-Mouse IgG (Bethyl, Montgomery, TX) as a 3.01 mg / mL solution in 0.1 M sodium phosphate, 1.0 mM EDTA, pH 6.5. To this solution was then added 216 µL of a freshly prepared 500 mM solution of the reducing agent DTT (1,4-Dithiothreitol, Sigma-Aldrich, St. Louis, MO). The vial was placed in the dark on an autorotator and the disulfide reduction was allowed to proceed for 25 minutes. The reaction solution was split into four equal volumes (due to the limited capacity of a desalting column used), and excess DTT was removed by passing each of the fractions across a PD-10 desalting column eluted with 0.1 M sodium phosphate, 1.0 mM EDTA, pH 6.5. The antibody containing fractions were combined to give 8.0 mL of a 1.22 mg / mL solution of DTT free Goat-anti-Mouse IgG-SH (78 % recovery) as measured by UV/VIS

spectrophotometry using an extinction coefficient at 280 nm of a 1% solution at pH 6.5 of 14.

HRP-Antibody Conjugation

To a thiolated antibody (such as anti-mouse IgG-thiol or anti-rabbit IgG-thiol), is added a three fold molar excess of HRP-PEG₄-maleimide. The reaction is then incubated at ambient temperature (23 – 25 °C) for 16 hours. After purification across a Superdex 200 10/300 GL SE column a conjugate, typically with an average of 2 or 3 HRPs per antibody, is obtained. The number of HRPs per antibody is determined by measuring the ratio of absorbances at 280 nm / 403 nm of the conjugate, and performing the calculations outlined in section Example D. An exemplary procedure is outlined below.

HRP-PEG₄-Goat-anti-Mouse IgG(3): To an 8 mL amber vial was added 4.0 mL of the Goat-anti-Mouse IgG-thiol solution (2) (1 eq., 4.88 mg, 0.0326 μmol) and 864 μL of the HRP-PEG₄-maleimide solution (1) (3 eq., 3.91 mg, 0.0976 μmol). The vial was then placed on an autorotator in the dark at ambient temperature (23 – 25 °C), and the Michael addition was allowed to proceed for 16 hours. HRP-PEG₄-Goat-anti-Mouse IgG conjugate devoid of free antibody and free HRP was then obtained by fractionating the sample on an Akta Purifier fitted with a Superdex 10/300 column (Amersham, Piscataway, NH) eluted with 0.1 M sodium phosphate, pH 7.5, at 0.9 ml / minute. After pooling fractions, 9.73 mL of a 1.04 mg / mL solution of conjugate was obtained as determined by Pierces' Coomassie Plus protein assay described in Example C. The conjugate was then stored in a cold room at 4 °C until use.

C. *MW Characterization of Antibody/Enzyme Conjugates*

To illustrate the superior monodispersity of the disclosed conjugates the MW profiles of a total of twelve examples of the disclosed conjugates (specifically, eight HRP-anti-mouse IgG conjugates and four HRP-anti-rabbit IgG conjugates) were determined by size-exclusion chromatography on an Akta Purifier fitted with a Superdex 200 10/300 GL column (Amersham, Piscataway, NJ) eluted with 0.1 M sodium phosphate buffer pH 7.5, 0.5-1.0 mL / min. Molecular weight calibration standards included: Aldolase (158 kDa), Catalase (232 kDa), Ferritin (440 kDa), Thyroglobin (669 kDa), Ribonuclease A (13.7 kDa), Chymotrypsinogen (25 kDa), Ovalbumin (43 kDa), and Albumin (67 kDa). The conjugates examined had an average MW between about 230 and about 330 kDa with an overall range of MWs for a given conjugate of approximately 190-550 kDa. Reinjection of purified conjugates demonstrated that conjugates were free of non-conjugated HRP and antibody.

D. *Analytical Procedures for Determining Conjugate*

The following representative methods may be used to determine maleimide and thiol content as well as the number of HRP molecules per conjugate.

*Total Protein Microplate Procedure (Pierce)*Equipment and Materials:

BSA	Pierce (Rockford, IL)
Coomasie Plus TM Reagent	Pierce (Rockford, IL)
Microtiter plate	BIO-TEK Synergy HT
Plate reader	

Procedure:

1. Turn on plate reader and let warm up for at least 30 minutes at 595 nm.
2. Prepare a set of BSA standards (1.0, 0.5, 0.25, and 0.125 mg /mL) in deionized water.
3. In triplicate, pipette 15ml of the Blank, and each standard or unknown into the appropriate microplate wells.
4. Add 300 ml of the Coomassie PlusTM Reagent to each well and mix with the plate shaker for 30 seconds.
5. Remove the plate from the shaker. For the most consistent results, incubate plate for 10 minutes at room temperature.
6. Measure the absorbance at 595 nm with the plate reader.
7. Subtract the average 595 nm measurement for the Blank replicates from the 595 nm measurements of all other individual standard and unknown sample replicates (done automatically by plate reader).
8. Prepare a standard curve by plotting the average Blank-corrected 595 nm measurements for each BSA standard versus its concentration in $\mu\text{g} / \text{mL}$. Use the standard curve to determine the protein concentration of each unknown sample (done by plate reader).

*Determination of Ab-Thiol and HRP-PEG₄-Maleimide Content*Equipment and Materials:

Mercaptoethanol J.T. Baker, Phillipsburg, NJ

Ellman's Reagent Pierce, Rockford, IL

Sodium phosphate

EDTA

Materials Preparation:

- Reaction Buffer: 0.1 M sodium phosphate; 1 mM EDTA, pH 8.0.
- Mercaptoethanol (BME): M.W. = 78.3, d = 1.114g / ml.

Procedure:

1. Turn on the plate reader and let warm up for at least 30 minutes at 412 nm.
2. Prepare working stock: 7 μ l BME into 5 ml Reaction Buffer
3. In triplicate, prepare a set of BME standards as described below.

Standard	Volume of Reaction buffer		Final Conc.
Standard stock	900 μ l	100 μ l of working stock	2 mM
Standard 1	500 μ l	500 μ l of Standard stock	1 mM
Standard 2	500 μ l	500 μ l of Standard 1	0.5 mM
Standard 3	500 μ l	500 μ l of Standard 2	0.25 mM
Standard 4	500 μ l	500 μ l of Standard 3	0.125mM
Standard 5	500 μ l	500 μ l of Standard 4	0.0625 mM

Standard 6	500 μ l	500 μ l of Standard 5	0.03125 mM
Standard 7	500 μ l	500 μ l of Standard 6	0.015625 mM
Standard 8 (blank)	1000 μ l		0 mM

4. If assaying HRP-PEG₄-MAL, add 160 μ l of sample to 160 μ l of Standard 1 and incubate for 30 minutes. This mixture serves as the unknown for HRP-PEG₄-MAL samples. Add 100 μ l of this unknown to the appropriate well as described in Step 5.
5. Add 100 μ l of each standard or unknown to the appropriate wells of a microtiter plate (attach template).
6. Prepare Ellman's Reagent Solution.
Ellman's Reagent Solution: Dissolve 8 mg Ellman's in 2 ml Reaction Buffer.
7. Add 20 μ l of Ellman's Reagent Solution to each well containing standard or unknown.
8. Mix and incubate at room temperature for 15 minutes.
9. Measure absorbance at 412 nm using the Plate reader.
10. If only raw data available, plot the values obtained for the standards to generate a standard curve.

Analysis:

Experimental concentrations (mM thiol) are determined from the standard curve, where the standard curve gives an equation: $Y = mX + b$, wherein $Y = OD_{412nm}$, $X =$ mM thiol, $m =$ slope (obtained from standard curve equation), and $b =$ x axis intercept (obtain from standard curve equation).

For each sample, the protein concentration in mM is determined by dividing the protein concentration in mg/ml (obtained from total protein assay) by the FW of the sample and multiplying by 1000. Then, the number of thiols per antibody molecule is obtained by dividing the mM thiol experimental concentration obtained from above by the protein concentration in mM obtained from the previous step. The number of maleimides per horseradish peroxidase molecule is determined by first subtracting the experimental mM thiol concentration obtained above from 0.5 mM, and then multiplying this difference by 2 and dividing by the protein concentration in mM.

A typical range for thiolation of an antibody is between about 1 and about 10 thiols per antibody molecules, for example, between about 2 and about 6 such as between about 2 and about 4. A typical range for the number of maleimide groups incorporated per HRP molecule is between about 1 and about 10, for example, between about 3 and about 8 such as between about 5 and about 7.

*Determination of the Number of HRPs Per Antibody*Constants

- HRP Molecular Weight = 40,000 Da
- Antibody Molecular Weight = 150,000 Da
- HRP 280 nm Extinction Coefficient of a 1 percent solution (1mg/mL) = 6.52

- Antibody 280 nm Extinction Coefficient of a 1 percent solution (1mg/mL) = 14
- HRP Absorbance at 403 nm / Absorbance at 280 nm = 2.90 (This value is measured for each different lot of HRP)

Calculations

- 1) Determine the 280 nm absorbance contribution to the conjugate due to HRP by measuring the conjugate absorbance at 403 nm and applying the equation: $\text{HRP Absorbance at 403 nm} / 2.90 = \text{HRP Absorbance at 280 nm}$.
- 2) From the value obtained in 1, determine the amount of HRP in mg/ml by applying the equation: $\text{HRP Absorbance at 280 nm} / 6.52 = [\text{HRP}] \text{ in mg/ml}$.
- 3) Determine the number of mM HRP by dividing the protein concentration in mg/ml (obtained from 2) by the FW (40,000) and multiplying by 1000.
- 4) Determine the 280 nm absorbance contribution to the conjugate due to secondary antibody by measuring the conjugate absorbance at 280 nm and subtracting the contribution due to HRP determined in 1.
- 5) From the value obtained in 4, determine the amount of HRP in mg/ml by applying the equation: $\text{Antibody Absorbance at 280 nm} / 14 = [\text{Antibody}] \text{ in mg/ml}$
- 6) Determine the number of mM antibody by dividing the antibody concentration in mg/ml (obtained from 5) by the FW (150,000) and multiplying by 1000.
- 7) Calculate the number of HRPs per secondary antibody by dividing the mMoles of HRP (determined in 3) by the number of mMoles of secondary antibody (determined in 6)

Determination of the Extinction Coefficient at 280 nm of a One Percent Solution of HRP-Antibody Conjugate

The determination of the extinction coefficient at 280 nm of a one percent (1 mg/mL) solution of HRP- antibody conjugate is determined by ascertaining the conjugate protein concentrations, and then measuring the absorbance at 280 nm. Protein concentrations can be measured according to the Pierce Coomassie assay described above.

E. Stability of Conjugates in Immunohistochemical Analyses

The stability at 45 °C of a cocktail of goat anti-mouse and goat anti-rabbit HRP conjugates in IHC was determined in an Avidin diluent with B5 blocker (Ventana Medical Systems, Inc, Tucson, AZ) and the results are shown in FIG. 1 A-D. Fixed, paraffin-embedded human tonsil tissue sections were probed using CD20/L26 (mouse) primary antibodies, followed by DAB detection with the cocktail of HRP conjugates according to a standard automated protocol on a BenchMark[®] XT autostainer (Ventana Medical Systems, Inc, Tucson, AZ). All slides were done in were done in triplicate. FIG. 1A shows typical results on Day 0 of the test; FIG. 1B shows typical results on Day 1 of the test; FIG. 1C shows typical results on Day 3 of the test; and FIG. 1D shows typical results on Day 7 of the test. Even at the high temperature of 45 °C, the disclosed conjugates were not completely degraded (30-40% loss of staining intensity) by day 7, demonstrating that the disclosed conjugates are highly stable.

Similar studies over a longer period were performed for storage at 2-8 °C, at 27°C, and at 37°C (no data shown), and further demonstrated the superior stability of the disclosed conjugates. In summary, at 2-8 °C no change in staining intensity was

observed between Day 0 and Week 2. At 27 °C virtually no change in staining intensity was observed between Day 0 and Week 2 for CD20. At 37 °C a ~ 25% loss in staining intensity was observed over a one week period, and a 30-50% loss in staining intensity was observed after 2 weeks for both CD20 and PSA. At Week 2 there is a 30-50% loss in staining intensity for both CD20 and PSA..

F. IHC Performance Assessment of Conjugates as Secondary Antibodies to Different Primary Antibodies

Goat anti-mouse IgG conjugate made with MAL-PEG₄-NHS linker, goat anti-rabbit IgG conjugate also made with the same linker, or a mixture of rabbit anti-mouse IgG and the two conjugates (“amplification”) was used as a secondary antibody reagent for detection of binding to tissue antigens of the primary antibodies that are listed below (available from Ventana Medical Systems, Inc, Tucson, AZ). Appropriate archival tissue sections were treated with these conjugates and developed using standard protocols for HRP signal generation (by addition of DAB) on an automated stainer (BenchMark[®] XT, Ventana Medical Systems, Inc, Tucson, AZ). A typical automated protocol includes deparaffinization, several rinse steps, addition of a reaction buffer, addition of the primary antibody, addition of the secondary antibody, addition of DAB and hydrogen peroxide, and addition of a counterstain.

Comparable (adjacent) tissue sections were stained with the disclosed conjugates and with polylysine-scaffolded HRP/F(ab')₂ conjugates (hereinafter “scaffolded conjugates”) used as the secondary antibody reagent. The scaffolded conjugates were either a second generation scaffolded conjugate (smaller, more homogeneous as determined by size-exclusion chromatography) or a first generation (larger, more

inhomogeneous as determined by size-exclusion chromatography). See, U.S. Patent Nos. 6,613,564 and 6,252,053 for a more complete description of the scaffolded conjugates.

Antibodies

Anti-bcl-2(clone 100/D5)	Anti-CD57(clone NK-1)
Anti-CD15(clone MMA)	Anti-CD23(clone 1B12)
Anti-CD20(clone L26)	Anti-ER(clone 6F11)
Anti-PR(clone 16)	Anti- p53(clone D07)
Anti-EGFR(clone 31G7)	Anti-Cyclin-d1(clone P2D11F11)
Anti-c-erbB-2(clone CB11)	Anti-PSA

**note: all were mouse antibodies, with the exception of PSA, which is a rabbit antibody.*

FIG. 2 shows the staining results for bcl-2 detection for the disclosed conjugate (FIG. 2A) and the second generation scaffolded conjugate (FIG 2B). The results demonstrate that higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 3 shows the staining results for CD-15 detection using the disclosed conjugate (FIG. 3A) and the second generation scaffolded conjugate (FIG 3B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 4 shows the staining results for CD-20 detection using the disclosed conjugate (amplification utilized, FIG. 4A) and the second generation scaffolded conjugate (FIG 4B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 5 shows the staining results for CD-23 detection using the disclosed conjugate (FIG. 5A), the second generation scaffolded conjugate (FIG 5B), and the first generation scaffolded conjugate (FIG. 5C). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections than is seen for both scaffolded conjugates.

FIG. 6 shows the staining results for CD57 detection using the disclosed conjugate (FIG. 6A) and the second generation scaffolded conjugate (FIG 6B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 7 shows the staining results for cerb-B2/CB11 detection using the disclosed conjugate (FIG. 7A), the second generation scaffolded conjugate (FIG 7B), and the first generation scaffolded conjugate (FIG. 7C). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections than is seen for both scaffolded conjugates.

FIG. 8 shows the staining results for cyclin D1 detection using the disclosed conjugate (FIG. 8A) and the second generation scaffolded conjugate (FIG 8B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 9 shows the staining results for EGFR detection using the disclosed conjugate (FIG. 9A), the second generation scaffolded conjugate (FIG 9B), and the first generation scaffolded conjugate (FIG. 9C). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections than is seen for both scaffolded conjugates.

FIG. 10 shows the staining results for ER detection using the disclosed conjugate (FIG. 10A) and the second generation scaffolded conjugate (FIG 10B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 11 shows the staining results for p53 detection using the disclosed conjugate (FIG. 11A) and the second generation scaffolded conjugate (FIG 11B). The results demonstrate comparable staining is achieved between the disclosed conjugate and the scaffolded conjugate in comparable tissue sections.

FIG. 12 shows the staining results for PR detection using the disclosed conjugate (FIG. 12A) and the second generation scaffolded conjugate (FIG 12B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

FIG. 13 shows the staining results for PSA detection using the disclosed conjugate (FIG. 13A) and the second generation scaffolded conjugate (FIG 13B). The results demonstrate higher intensity staining is achieved with the disclosed conjugate in comparable tissue sections.

In conclusion, the results of tissue testing of the disclosed conjugate detection compositions demonstrated that the disclosed conjugates perform significantly better for tissue staining than scaffolded conjugates.

**G. *Stability of Conjugates at 37 °C and 45 °C for Enzyme
Metallographic Detection of Nucleic Acid Sequences.***

Experiments were performed to assess the stability over time of a goat anti-rabbit IgG antibody-HRP (PEG4) conjugate at 45 °C and at 37 °C. In this instance, stability of the conjugates was assessed in an assay involving enzyme metallographic detection (EnzMet, Nanoprobes Inc., Yaphank, NY) of nucleic acid sequences. As illustrated in FIG. 14, biotin-labeled probe DNA was detected with a combination of an anti-biotin rabbit conjugate and anti-rabbit IgG conjugate. The conjugate mixture was stored in Stabilzyme SelectTM (Surmodics, Eden Prairie, MN) as the diluent. The stability of the second generation scaffolded conjugate discussed in Example D above, was also examined over the same time period.

FIG 15A shows a tissue stained with the disclosed conjugate at day 0, which may be compared to the tissue stained with the scaffolded conjugate at day 0 in FIG. 15B. FIG. 15C shows a tissue stained with the disclosed conjugate at day 7 after storage at 37 °C for 7 days, which may be compared to the tissue stained with the scaffolded conjugate at day 7 after storage at 37 °C for 7 days in FIG. 15D. FIG. 15E shows a tissue stained with the disclosed conjugate at day 7 after storage at 45 °C for 7 days, which may be compared to the tissue stained with the scaffolded conjugate at day 7 after storage at 45 °C for 7 days in FIG. 15F. The tissue staining intensity shown in the figures demonstrates the superior stability of the disclosed conjugate at both temperatures over a period of 7 days, with the scaffolded conjugate showing complete loss of staining ability after 7 days at the higher temperature.

The relative stability over time of the disclosed conjugate and the scaffolded conjugate for detecting single copy and for detecting multiple copies of a target DNA

sequence is shown in graphic form in FIG. 16A (37 °C) and FIG. 16B (45 °C). The graphs illustrate how much less effective the scaffolded conjugate is for enzyme metallography of both single and multiple copy targets, how the scaffolded conjugate is completely ineffective for single copy detection while the disclosed conjugate was effective for single copy detection even after many days of storage at elevated temperature, and how the disclosed conjugate maintains its ability for multiple copy detection over time at both temperatures while the scaffolded conjugate quickly loses its ability to amplify the gene signal at both temperatures.

H. Effect of Reaction Conditions on Conjugate Composition

The reproducibility with which a well defined antibody-HRP conjugate could be made was investigated by looking at the effect of DTT reduction time of the antibody, length of the linker as well as type, stoichiometry of linker added, HRP concentration in the coupling reaction, and the molar ratio of HRP to antibody. Size exclusion chromatography on an AKTA Purifier LC fitted with a Superdex 10/300 200 GL column (Amersham, Piscataway, NJ) was used make initial comparisons. The mobile phase used was phosphate buffered saline, pH = 7.5 with a flow rate of 1 ml/min.

Variation of DTT Reduction Time

Following the synthetic protocol for the conjugate previously outlined in Example B, a series of reactions were set up in which the incubation time with DTT (25mM) was varied. The following time points were tested: 15 min, 25 min, 35 min, and 60 min. After performing the coupling reaction between the antibody and maleimide derivatized HRP, the size exclusion chromatograms illustrated in FIG. 17. It

was evident that by changing the time period of the DTT treatment that the composition of the conjugate was not significantly altered. The staining obtained with these conjugates on tissue (tonsil, Ki-67) showed no significant change in staining specificity or intensity, with a 15 min DTT treatment being only slightly better than the rest. However, with the other three time points giving identical staining on tissue, this study indicates that the time sensitive nature of the DTT reduction is not overly critical in the production of a reproducible, active conjugate according to the disclosed methods.

Variation of Linker Length / Type

Following the procedure of Example B, a series of reactions were set up altering the linker type and size. The following linkers were used: LC-SMCC (16 atom hydrophobic linker, Pierce, Rockford IL), MAL-dPEG₈-NHS ester (34 atom hydrophilic linker, Quanta Biodesign, Inc., Powell OH), MAL-dPEG₁₂-NHS ester (46 atom hydrophilic linker, Quanta Biodesign, Inc., Powell OH), as well as the recommended MAL-dPEG₄-NHS ester (22 atom hydrophilic linker, Quanta Biodesign, Inc., Powell OH). Each of these linkers was used in a hundred-fold excess, in a buffer (0.1 M sodium phosphate, pH = 7.5) for 1 hour. The LC-SMCC was dissolved in dimethylformamide (DMF) and added to the HRP, but not exceeding 10% total volume of DMF in buffer. After coupling to the DTT-treated antibody, size exclusion chromatograms (FIG. 18) were obtained upon purification. Each of the three PEG linkers, based on retention volume, performed comparatively well, the LC-SMCC linker, however, showed less conjugation to the HRP (larger peak at ~16 min) and an overall smaller conjugate.

Differences were evident in the immunohistochemical tissue staining intensity (Ki-67 primary antibody/conjugate secondary antibody, with amplification, on tonsil tissue) afforded by the different conjugates (FIG. 19), and the LC-SMCC conjugate gave the lightest amount of staining. Each of the staining runs was done with the conjugates at equivalent 280 nm absorbances ($A_{280} = 0.075$), and therefore make the data directly comparable. The three PEG derived conjugates performed surprisingly better than the LC-SMCC (FIG 19A), and there were differences in the staining intensity afforded by each of them. It is clear from the figures that the PEG₁₂ (FIG. 19D) had the darkest overall staining followed by the PEG₈ (FIG. 19C) and then PEG₄ (FIG. 19D). As will be discussed further below with respect to *in situ* hybridization assays, the intense staining obtained with conjugates prepared with longer linkers can surprisingly obviate the need for amplification steps during staining.

Variation of Linker Stoichiometry

The synthesis of the HRP-IgG conjugate was carried out following the conjugation procedure of Example B, but the molar excess of MAL-PEG₄-NHS ester linker over the HRP amount was varied from a five-fold excess to a five hundred-fold excess. Analysis of the conjugates (500x, 250x, 100x, 50x, 25x, 10x, and 5x), after reaction with the DTT reduced Ab, carried out via size exclusion chromatography as described immediately above in this example, indicated that the conjugates synthesized using a larger excess of linker had a smaller, narrower size distribution range (FIG. 20). However, there did not seem to be a large difference in the overall size distribution for the conjugates ranging from 5x to 100x. Tissue staining (tonsil, Ki-67, not shown) for

each of these conjugates was roughly equivalent, where only 5x was slightly darker than other amounts.

Variation of HRP Concentration in Linker Coupling Reaction

Following the synthetic method outlined previously in example B, the effect of HRP concentration during the initial derivatization step was investigated. Stock solutions of HRP at the following concentrations: 5 mg/ml, 10 mg/ml, 15 mg/ml, 20 mg/ml, and 50 mg/ml, alongside the original protocol (25 mg/ml) concentration were used in the reactions. After the coupling step with the DTT-reduced antibody there was no difference in the overall size exclusion chromatograms for the synthesized conjugates (FIG. 21). In assaying the activity of the synthesized conjugates on tissue (tonsil, Ki-67), it was noticed that the staining specificity and intensity were identical for the conjugates synthesized using 5, 10, 15, 20, and 25 mg/ml HRP concentrations. However, the staining intensity decreased when the starting HRP concentration was increased to 50 mg/ml. It is concluded that the starting HRP concentration should stay between 10-25 mg/ml for the production level scale-up.

Variation of HRP / Ab Molar Ratios

The HRP / IgG conjugates were synthesized using the protocol outlined in Example B, but the ratio of the DTT-reduced antibody to the maleimide derivatized HRP was varied. The following ratios (Antibody / HRP) were tested: 3:1, 1:3, 1:2, 1:4, 1:5, 1:10, 1:20, as well as the recommended 1:3. The profiles in the size exclusion chromatographs (FIG. 22) show that as the relative amount of HRP increases, so does the overall size of the conjugate, with the 1:20 (Ab:HRP) giving the largest conjugate

and the 3:1 (Ab:HRP) generating the smallest. Each of these conjugates performed well on tissue (tonsil, Ki-67), with the 3:1 (Ab:HRP) producing the lightest amount of staining. The 1:3 (Ab:HRP) is a midrange point with good staining and produces relatively high yields with respect to the HRP.

I. Preparation of a Rabbit Anti-Biotin-HRP-PEG12 Conjugate and Its Use for Enzyme Metallographic In Situ Hybridization

HRP-PEG₁₂-maleimide (4): To a 4 mL amber vial was added 18.4 mg (100 eq.) of MAL-dPEG₁₂TM NHS ester (Quanta Biodesign, Powell, OH, F.W. = 865.92), followed by 341 μ L (8.52 mg, 0.213 μ M) of HRP (Horseradish Peroxidase, Pierce, Rockford, IL) as a 25 mg / mL solution in 0.1 M sodium phosphate, pH 7.5. The vial was then placed on an autorotator in the dark at ambient temperature (23 – 25 °C), and the amide bond forming reaction was allowed to proceed for 1 hour. A 340 μ L aliquot was then removed for purification. (The capacity of the Akta Purifier injection loop utilized was 500 μ L). Pure HRP-PEG₁₂-maleimide was then obtained by fractionating the sample on an Akta Purifier fitted with a Superdex 10/300 column eluted with 0.1 M sodium phosphate, pH 7.5 at 1.0 mL / min. The HRP containing fractions (F15-17) were pooled to give 1.5 ml of a 4.75 mg / mL solution of HRP-PEG₁₂-maleimide (83.6 % recovery) as measured on an UV/VIS spectrophotometer using the extinction coefficient at 280 nm of a 1% solution at pH 7.5 of 6.52.

Rabbit anti-Biotin thiol (5): To a 4 mL amber vial was added 2.0 mL of Rabbit anti-Biotin (Bethyl, Montgomery TX) as a 1.0 mg/mL solution. To this solution was then added 105.2 μ L of a freshly prepared 500 mM solution of the reducing agent DTT (1,4-Dithiothreitol). The vial was placed in the dark on an autorotator and the disulfide

reduction was allowed to proceed for 25 minutes. The reaction solution was split into two equal volumes (due to the limited capacity of the desalting columns), and the excess DTT was removed by passing each of the for fractions across a PD-10 desalting column eluted with 0.1 M sodium phosphate, 1.0 mM EDTA, pH 6.5. The antibody containing fractions (F4-5) were combined to give 4.0 mL of a 0.436 mg / mL solution of DTT free Rabbit anti-Biotin-SH (87.5 % recovery) as measured on a Agilent 8453 UV/VIS spectrophotometer using an extinction coefficient at 280 nm of a 1% solution at pH 6.5 of 14.


HRP- Antibody Conjugation(6): To the rabbit anti-biotin-IgG-thiol (5), was added a three fold molar excess of HRP-PEG₁₂-maleimide (4). The reaction was then incubated at ambient temperature (23 – 25 °C) overnight. After purification across a Superdex 200 10/300 GL SE column, 875 mg of conjugate with an average M.W. of 359 kD was obtained.

The enzyme metallographic procedure outlined in Example G was repeated using the PEG₁₂ anti-biotin conjugate as the primary antibody (i.e. no amplification), and resulted in surprisingly intense staining even though no amplification was employed. These results demonstrate that the use of long heterobifunctional PEG linkers (PEG₈ or greater, such as PEG₁₂ or greater) to prepare the disclosed conjugates surprisingly obviates the need for amplification schemes for IHC and ISH applications on tissue sections.

J. Maleimide/Hydrazide PEG-linker Synthesis


Scheme 6 shows a general method for preparing maleimide/hydrazide heterobifunctional PEG linkers. Briefly, a maleimide/active ester PEG linker (such as

obtained from Quanta Biodesign) is reacted with a protected hydrazine derivative, and then reacted with acid to yield the maleimide/hydrazide PEG linker.



Scheme 6


A specific synthesis of a maleimide/hydrazide PEG₄ linker is outlined in Scheme 7 below. To the active ester **7** (116mg, 1.0 eq.) in 5 ml dry dioxane was added 30 mg (1.0 eq.) of the Boc protected hydrazine **8** in 5 ml of dry dioxane over 1 hour. The reaction was then stirred at ambient temperature under dry nitrogen for 16 hours. The reaction mixture was fractionated by HPLC utilizing a Waters Delta 600 HPLC fitted with a 2996 photo-diode array detector and a Phenomenex luna 10 μ , C18(2), 100A, 250 x 30 mm column. The column was eluted with 30-60% ACN / water over 30 min at a flow rate of 12 mL / min. The desired Boc protected-PEG₄-maleimide **9** eluted at 38 minutes giving 50 mg of a thick yellow oil after drying under high vacuum. The final deprotected hydrazide **10** was then obtained by stirring the residue with 6 ml of anhydrous 2 N HCL / dioxane under dry nitrogen for 45 minutes. Concentration via rotary evaporation then gave 55 mg of the hydrazide-PEG₄-maleimide HCL salt.



Scheme 7


THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. An antibody-signal-generating moiety conjugate comprising an antibody covalently linked to a signal-generating moiety through a heterobifunctional PEG linker, wherein the conjugate has the formula:



wherein Ab is the antibody, SM is the signal-generating moiety, $n = 1$ to 50 and $s = 1$ to 10, and wherein said signal-generating moiety is an enzyme, fluorescent molecule, detectable construct, metal chelate or a liposome.

2. The conjugate of claim 1, wherein the linker, prior to conjugation, has the formula:



wherein $n = 1$ to 50.


3. The conjugate of claim 1, wherein the signal-generating moiety is an enzyme.

4. The conjugate of claim 3, wherein the enzyme is horseradish peroxidase or alkaline phosphatase.

5. The conjugate of claim 1, wherein $s = 2$ to 6.

6. The conjugate of claim 1, wherein $n = 4$ to 12.

7. A method for preparing an antibody-signal-generating moiety conjugate, comprising:
 reacting an antibody with a reducing agent to form a thiolated antibody;
 reacting a signal-generating moiety having an amine group with a PEG maleimide/active ester
 bifunctional linker to form an activated signal-generating moiety, wherein the PEG
 maleimide/active ester bifunctional linker has the formula:



wherein $n = 1$ to 50; and

reacting the thiolated antibody with the activated signal-generating moiety to form the antibody-signal-generating moiety conjugate, wherein said signal-generating moiety is an enzyme, fluorescent molecule, detectable construct, metal chelate or a liposome.

8. The method of claim 7, wherein reacting the antibody with the reducing agent to form the thiolated antibody comprises forming an antibody with an average number of thiols per antibody of between 1 and 10.

9. The method of claim 7, wherein the reducing agent is selected from the group consisting of 2-mercaptoethanol, 2-mercaptoethylamine, 1,4-Dithiothreitol (DTT), dithioerythritol (DTE) and tris(carboxyethyl)phosphine (TCEP), and combinations thereof.


10. The method of claim 9, wherein reacting the antibody with the reducing agent comprises reacting the antibody with a reducing agent selected from the group consisting of DTT and DTE, and combinations thereof.

11. The method of claim 10, wherein reacting the antibody with the reducing agent comprises reacting the antibody with the reducing agent at a concentration of between about 1 mM and about 40 mM.

12. The method of claim 7, wherein the signal-generating moiety is an enzyme.

13. The method of claim 12, wherein the enzyme is horseradish peroxidase or alkaline phosphatase.

14. A method for detecting a molecule of interest in a biological sample, comprising: contacting the biological sample with an antibody-signal-generating moiety conjugate comprising an antibody covalently linked to a signal-generating moiety through a heterobifunctional PEG linker, wherein the conjugate has the formula:



wherein Ab is the antibody, SM is the signal-generating moiety, $n = 1$ to 50 and $s = 1$ to 10, and wherein said signal-generating moiety is an enzyme, fluorescent molecule, detectable construct, metal chelate or a liposome; and detecting a signal generated by the antibody-signal-generating moiety conjugate that indicates the presence of the molecule of interest in the sample.

15. The method of claim 14, wherein the biological sample comprises a tissue section or a cytology sample.

16. The method of claim 14, wherein the signal-generating moiety is an enzyme.

17. The method of claim 16, wherein the enzyme is horseradish peroxidase or alkaline phosphatase.

18. The method of claim 16, wherein the method further comprises contacting the biological sample with a water-soluble metal ion and a redox-inactive substrate of the enzyme that is converted to a redox-active agent by the enzyme, which redox-active agent reduces the metal ion causing it to precipitate.

19. The method of claim 18, wherein the enzyme is alkaline phosphatase.

20. The method of claim 16, wherein the signal-generating moiety comprises an oxido-reductase enzyme and the method further comprises contacting the biological sample with a water soluble metal ion, and oxidizing agent and a reducing agent.

21. The method of claim 20, wherein the oxido-reductase enzyme is horseradish peroxidase.

22. The method of claim 14, wherein the antibody comprises an anti-antibody antibody.

23. The method of claim 14, wherein the antibody comprises an anti-hapten antibody, and the method further comprises contacting the sample with a hapten-labeled antibody or a hapten-labeled nucleic acid sequence.

24. The method of claim 14, wherein the signal-generating moiety comprises an enzyme and the method further comprises contacting the biological sample with a chromogenic, fluorogenic and/or luminogenic compound.

25. The conjugate of claim 1, wherein $n = 2$ to 30.

26. The conjugate of claim 1, wherein the signal-generating moiety comprises a quantum dot.

27. The method of claim 7, wherein the signal-generating moiety comprises a quantum dot.

28. The method of claim 14, wherein the signal-generating moiety comprises a quantum dot.

1/22



FIG. 1A



FIG. 1B



FIG. 1C




FIG. 1D

2/22




FIG. 2A




FIG. 2B

3/22




FIG. 3A




FIG. 3B

4/22




FIG. 4A




FIG. 4B

5/22




FIG. 5A




FIG. 5B




FIG. 5C

6/22




FIG. 6A




FIG. 6B

7/22




FIG. 7A




FIG. 7B




FIG. 7C

8/22




FIG. 8A



FIG. 8B

9/22




FIG. 9A



FIG. 9B




FIG. 9C

10/22




FIG. 10A




FIG. 10B

11/22




FIG. 11A




FIG. 11B

12/22




FIG. 12A




FIG. 12B

13/22




FIG. 13A




FIG. 13B

Nanoprobes EnzMet




FIG. 14

15/22




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 15D




FIG. 15E






FIG. 15F

16/22**FIG. 16A****FIG. 16B**

17/22




FIG. 17

18/22




FIG. 18

19/22




FIG. 19A




FIG. 19B




FIG. 19C




FIG. 19D

20/22




FIG. 20

21/22




FIG. 21

22/22




FIG. 22

