



US 20150353940A1

(19) **United States**

(12) **Patent Application Publication**

McClain et al.

(10) **Pub. No.: US 2015/0353940 A1**

(43) **Pub. Date: Dec. 10, 2015**

(54) **VECTORS FOR USE IN AN INDUCIBLE COEXPRESSION SYSTEM**

(71) Applicant: **ABSCI, LLC**, Portland, OR (US)

(72) Inventors: **Sean McClain**, Portland, OR (US); **Mark Valasek**, San Diego, CA (US); **Philip Barish**, Portland, OR (US); **Jeremy Minshull**, Menlo Park, CA (US)

(73) Assignee: **ABSCI, LLC**, Portland, OR (US)

(21) Appl. No.: **14/740,475**

(22) Filed: **Jun. 16, 2015**

Related U.S. Application Data

(63) Continuation-in-part of application No. 14/419,653, filed on Feb. 4, 2015, filed as application No. PCT/US2013/053562 on Aug. 5, 2013, Continuation-in-part of application No. PCT/US2014/014968, filed on

Feb. 5, 2014, which is a continuation-in-part of application No. PCT/US2013/053562, filed on Aug. 5, 2013.

(60) Provisional application No. 61/747,246, filed on Dec. 29, 2012, provisional application No. 61/679,751, filed on Aug. 5, 2012, provisional application No. 61/747,246, filed on Dec. 29, 2012, provisional application No. 61/679,751, filed on Aug. 5, 2012.

Publication Classification

(51) **Int. Cl.**
C12N 15/70 (2006.01)
C07K 16/24 (2006.01)

(52) **U.S. Cl.**
CPC *C12N 15/70* (2013.01); *C07K 16/241* (2013.01); *C12N 2830/002* (2013.01)

(57)

ABSTRACT

The present invention provides expression vectors for use in an inducible coexpression system, capable of controlled induction of expression of each gene product.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7

VECTORS FOR USE IN AN INDUCIBLE COEXPRESSION SYSTEM

REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. application Ser. No. 14/419,653, which is a national stage entry under 35 U.S.C. §371 of International Application No. PCT/US2013/053562 filed 5 Aug. 2013, which claims the benefit of priority under 35 U.S.C. §119(e) of U.S. Provisional Application No. 61/679,751, filed 5 Aug. 2012, and of U.S. Provisional Application No. 61/747,246, filed 29 Dec. 2012, the entire disclosures of all of which are incorporated by reference herein.

[0002] This application is also a continuation-in-part of International Application No. PCT/US2014/014968 filed 5 Feb. 2014, which claims the priority of International Application No. PCT/US2013/053562 filed 5 Aug. 2013, which claims the benefit of priority under 35 U.S.C. §119(e) of U.S. Provisional Application No. 61/679,751, filed 5 Aug. 2012, and of U.S. Provisional Application No. 61/747,246, filed 29 Dec. 2012, the entire disclosures of all of which are incorporated by reference herein.

REFERENCE TO THE SEQUENCE LISTING

[0003] This application includes a sequence listing submitted electronically, which is incorporated by reference herein.

FIELD OF THE INVENTION

[0004] The present invention is in the general technical fields of molecular biology and biotechnological manufacturing. More particularly, the present invention is in the technical field of recombinant protein expression.

BACKGROUND OF THE INVENTION

[0005] Production of biotechnological substances is a complex process, even more so when the desired product is a combination of molecules encoded by different genes, such as a multimeric protein formed from two or more different polypeptides. Successful coexpression of multiple gene products requires overcoming a number of challenges, which are compounded by the simultaneous expression of more than one gene product. Problems that must be overcome include creating compatible expression vectors when more than one type of vector is used; obtaining the correct stoichiometric ratio of products; producing gene products that are folded correctly and in the proper conformation with respect to binding partners; purifying the desired products away from cells and unwanted proteins, such as proteins that are folded incorrectly and/or are in an incorrect conformation; and minimizing the formation of inclusion bodies, as one aspect of maximizing the yield of the desired product(s). Many different approaches have been taken to address these challenges, but there is still a need for better coexpression methods.

[0006] Several inducible bacterial protein expression systems, including plasmids containing the lac and ara promoters, have been devised to express individual proteins. These systems have limited utility in the coexpression of difficult-to-express proteins as they fail to induce protein homogeneously within the entire growth culture population in wild-type *E. coli* (Khlebnikov and Keasling, "Effect of lacY expression on homogeneity of induction from the P_{lac} and P_{trc} promoters by natural and synthetic inducers", *Biotechnol Prog* 2002 May-June; 18(3): 672-674). When expression of

the transport proteins for inducers is dependent on the presence of inducer, as is the case for wild-type *E. coli* lac and ara systems, the cellular concentration of the inducer must reach a threshold level to initiate the production of transport proteins, but once that threshold has been reached, an uncontrolled positive feedback loop can occur, with the result being a high level of inducer in the cell and correspondingly high levels of expression from inducible promoters: the "all-or-none" phenomenon. Increasing the concentration of the inducer in the growth medium increases the proportion of cells in the population that are in high-expression mode. Although this type of system results in concentration-dependent induction of protein expression at the population scale, it is suboptimal for expression and production of proteins that require tight control of expression, including those that are toxic, have poor solubility, or require specific concentrations for other reasons.

[0007] Some efforts have been made to address the "all-or-none" induction phenomenon in single-promoter expression systems, by eliminating inducer-dependent transport of the inducer. One example is having a null mutation in the lactose permease gene (lacYam) and using an alternate inducer of the lac promoter such as IPTG (isopropyl-thio- β -D-galactoside), which can get through the cell membrane to some degree in the absence of a transporter (Jensen et al., "The use of lac-type promoters in control analysis", *Eur J Biochem* 1993 Jan. 15; 211(1-2): 181-191). Another approach is the use of an arabinose-inducible promoter in a strain deficient in the arabinose transporter genes, but with a mutation in the lactose permease gene, lacY(A117C), that allows it to transport arabinose into the cell (Morgan-Kiss et al., "Long-term and homogeneous regulation of the *Escherichia coli* araBAD promoter by use of a lactose transporter of relaxed specificity", *Proc Natl Acad Sci USA* 2002 May 28; 99(11): 7373-7377).

[0008] The components of individual protein expression systems are often incompatible, precluding their use in coexpression systems, as they may be adversely affected by 'crosstalk' effects between different inducer-promoter systems, or require mutually exclusive genomic modifications, or be subject to general metabolic regulation. An attempt to address the 'crosstalk' problem between the lac and ara inducible promoter systems included directed evolution of the AraC transcriptional activator to improve its ability to induce the araBAD promoter in the presence of IPTG, an inducer of the lac promoter (Lee et al., "Directed evolution of AraC for improved compatibility of arabinose- and lactose-inducible promoters", *Appl Environ Microbiol* 2007 September; 73(18): 5711-5715; *Epab* 2007 Jul. 20). However, the compatibility between expression vectors based on ara and lac inducible promoters is still limited due to the requirement for mutually exclusive genomic modifications: a lacY point mutation (lacY(A117C)) for homogenous induction of the araBAD promoter by arabinose, and a null lacY gene for homogenous induction of the lac promoter by IPTG. General metabolic regulation—for example, carbon catabolite repression (CCR)—can also affect the compatibility of inducible promoters. CCR is characterized by the repression of genes needed for utilization of a carbon-containing compound when a more preferred compound is present, as seen in the preferential use of glucose before other sugars. In the case of the ara and prp inducible promoter systems, the presence of arabinose reduces the ability of propionate to induce expression from the prpBCDE promoter, an effect believed to involve CCR (Park et al., "The mechanism of sugar-mediated

catabolite repression of the propionate catabolic genes in *Escherichia coli*”, Gene 2012 Aug. 1; 504(1): 116-121, Epub 2012 May 3). There is clearly a need for an inducible coexpression system that overcomes these problems.

SUMMARY OF THE INVENTION

[0009] The present invention provides expression constructs for use in inducible coexpression systems capable of controlled induction of each gene product component. One embodiment of the invention is an expression construct comprising two or more inducible promoters, wherein at least one of said inducible promoters is responsive to an inducer that is different than the inducer of another of said inducible promoters, and wherein none of the inducible promoters is an inducible promoter selected from the group consisting of: a tetracycline-inducible promoter, a copper-inducible promoter, and a methionine-inducible promoter. In some embodiments, the above expression construct does not comprise a lactose-inducible promoter, and in certain embodiments, the above expression construct does not comprise a promoter inducible by phosphate depletion. In additional embodiments, the expression construct is extrachromosomal. Further embodiments of the invention provide expression constructs comprising two or more inducible promoters, wherein at least one of said inducible promoters is responsive to an inducer that is different than the inducer of another of said inducible promoters, and (A) wherein each inducible promoter is selected from the group consisting of an L-arabinose-inducible promoter, a propionate-inducible promoter, a rhamnose-inducible promoter, a xylose-inducible promoter, a lactose-inducible promoter, and a promoter inducible by phosphate depletion; or (B) wherein at least one inducible promoter is selected from the group consisting of the araBAD promoter, the prpBCDE promoter, the rhaSR promoter, the xlyA promoter, the lacZYA promoter, and the phoA promoter; or (C) wherein the expression construct comprises at least one propionate-inducible promoter, which in some embodiments is the prpBCDE promoter, and at least one inducible promoter selected from the group consisting of an L-arabinose-inducible promoter, a rhamnose-inducible promoter, a xylose-inducible promoter, a lactose-inducible promoter, and a promoter inducible by phosphate depletion; or (D) wherein the expression construct comprises at least one L-arabinose-inducible promoter, which in some embodiments is the araBAD promoter, and at least one inducible promoter selected from the group consisting of a propionate-inducible promoter, a rhamnose-inducible promoter, a xylose-inducible promoter, a lactose-inducible promoter, and a promoter inducible by phosphate depletion; or (E) wherein at least one inducible promoter is a propionate-inducible promoter and at least one other inducible promoter is an L-arabinose-inducible promoter; or (F) wherein the expression construct comprises a nucleotide sequence having at least 80% (or at least 90%, or at least 95%) sequence identity to at least 50 (or at least 75, or at least 100) contiguous bases of nucleotides 4937 through 5185 of SEQ ID NO:15; or (G) wherein the expression construct comprises a nucleotide sequence having at least 80% (or at least 90%, or at least 95%) sequence identity to at least 50 (or at least 75, or at least 100) contiguous bases of nucleotides 2818 through 3151 of SEQ ID NO:15; or (H) wherein the expression construct comprises a nucleotide sequence having at least 80% (or at least 90%, or at least 95%) sequence identity to at least 50 (or at least 75, or at least 100) contiguous bases of nucleotides 2818 through

3151 of SEQ ID NO:15, and at least 80% (or at least 90%, or at least 95%) sequence identity to at least 50 (or at least 75, or at least 100) contiguous bases of nucleotides 4937 through 5185 of SEQ ID NO:15; or (I) wherein the expression construct comprises a nucleotide sequence having at least 80% (or at least 90%, or at least 95%) sequence identity to at least 3000 (or at least 3250) contiguous bases of nucleotides 1833 through 5304 of SEQ ID NO:15; or (J) wherein the expression construct comprises a nucleotide sequence having at least 80% (or at least 90%, or at least 95%) sequence identity to at least 5000 contiguous bases of SEQ ID NO:15; or (K) wherein the expression construct comprises nucleotides 1833 through 5304 of SEQ ID NO:15; or (L) wherein the expression construct comprises SEQ ID NO:15; or (M) wherein the expression construct further comprises a polynucleotide sequence encoding a transcriptional regulator that binds to an inducible promoter, wherein in some embodiments, the transcriptional regulator is selected from the group consisting of: AraC, PrpR, RhaR, and XylR; or (N) wherein the expression construct further comprises at least one polynucleotide sequence encoding at least one gene product to be transcribed from an inducible promoter.

[0010] Also provided by the invention is an expression construct comprising a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence having at least 97% sequence identity to at least 225 (or at least 240) contiguous bases of nucleotides 4937 through 5185 of SEQ ID NO:15; (b) a nucleotide sequence having at least 80% (or at least 90%, or at least 95%) sequence identity to at least 300 (or at least 350) contiguous bases of nucleotides 4937 through 5304 of SEQ ID NO:15; (c) a nucleotide sequence having at least 87% (or at least 90%, or at least 95%) sequence identity to at least 350 (or at least 375, or at least 400, or at least 425) contiguous bases of nucleotides 2818 through 3259 of SEQ ID NO:15; (d) a nucleotide sequence having at least 90% (or at least 95%) sequence identity to at least 400 (or at least 450, or at least 500) contiguous bases of nucleotides 10 through 1822 of SEQ ID NO:2; (e) nucleotides 2818 through 3259 of SEQ ID NO:15; (f) nucleotides 4937 through 5185 of SEQ ID NO:15; (g) nucleotides 2818 through 3259 of SEQ ID NO:15 and nucleotides 4937 through 5185 of SEQ ID NO:15; (h) nucleotides 10 through 1822 of SEQ ID NO:2; and (i) a nucleotide sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:15. In certain embodiments of the invention, the above expression construct comprises an inducible promoter selected from the group consisting of an L-arabinose-inducible promoter, which in some embodiments is the araBAD promoter; a propionate-inducible promoter, which in some embodiments is the prpBCDE promoter; a rhamnose-inducible promoter, which in some embodiments is the rhaSR promoter; a xylose-inducible promoter, which in some embodiments is the xlyA promoter; a lactose-inducible promoter, which in some embodiments is the lacZYA promoter; and a promoter inducible by phosphate depletion, which in some embodiments is the phoA promoter; and in some embodiments, the above expression construct further comprises at least one polynucleotide sequence encoding at least one gene product to be transcribed from an inducible promoter.

[0011] In additional embodiments of the invention, an expression construct is produced by a method comprising a step of inserting a polynucleotide sequence into a plasmid

selected from the group consisting of: pBAD240, pPRO43, pPRO44, pPRO45, pPRO430, pPRO430CloDF, and pSOL.

[0012] The present invention further provides a host cell comprising at least one expression construct of the invention as described in the above paragraphs, comprising at least one inducible promoter. In some embodiments of the invention, this host cell is a prokaryotic cell, and in some instances, it is an *E. coli* cell, which in certain embodiments is an *E. coli* ASE(DGH) cell. In other embodiments of the invention, the host cell is a eukaryotic cell, and in some instances it is a yeast cell, and in some further instances it is a *Saccharomyces cerevisiae* cell. Also provided by the invention is a host cell further comprising one or more of the following: (a) a deletion of the araBAD genes; (b) an altered gene function of the araE and araFGH genes; (c) a lacY(A177C) gene; (d) a reduced gene function of the prpB and prpD genes; (e) a reduced gene function of the sbm/scpA-ygfD/argK-ygfGH/scpBC genes; (f) a reduced gene function of the gor and trxB genes; (g) a reduced gene function of the AscG gene; (h) a polynucleotide encoding a form of DsbC lacking a signal peptide; (i) a polynucleotide encoding Erv1p; and (j) a polynucleotide encoding a chaperone. In certain additional embodiments, (A) the host cell has an alteration of gene function of at least one gene encoding a transporter protein for an inducer of at least one inducible promoter, and as another example, wherein the gene encoding the transporter protein is selected from the group consisting of araE, araF, araG, araH, rhaT, xylF, xylG, and xylH, or particularly is araE, or wherein the alteration of gene function more particularly is expression of araE from a constitutive promoter; (B) the host cell has a reduced level of gene function of at least one gene encoding a protein that metabolizes an inducer of at least one inducible promoter, and as further examples, wherein the gene encoding a protein that metabolizes an inducer of at least one said inducible promoter is selected from the group consisting of araA, araB, araD, prpB, prpD, rhaA, rhaB, rhaD, xylA, and xylB; and/or (C) the host cell has a reduced level of gene function of at least one gene encoding a protein involved in biosynthesis of an inducer of at least one inducible promoter, which gene in further embodiments is selected from the group consisting of scpA/sbm, argK/ygfD, scpB/ygfG, scpC/ygfH, rmlA, rmlB, rmlC, and rmlD.

[0013] In other aspects of the invention, a host cell is provided comprising two or more types of expression constructs, wherein the expression construct of each type comprises an inducible promoter, wherein the expression construct of each type comprises an inducible promoter that is not an inducible promoter of the expression construct of each other type, or wherein the expression construct of each type comprises an origin of replication that is different from the origin of replication of the expression construct of each other type.

[0014] In additional embodiments of the invention, at least one expression construct comprised by a host cell further comprises a polynucleotide sequence encoding a transcriptional regulator that binds to an inducible promoter; in some embodiments, the polynucleotide sequence encoding a transcriptional regulator and the inducible promoter to which said transcriptional regulator binds are in the same expression construct; and in further instances, the transcriptional regulator is selected from the group consisting of: AraC, PrpR, RhaR, and XylR; or in particular is AraC, or is PrpR.

[0015] A host cell is also provided by the invention, comprising at least one expression construct as described in the above paragraphs, which comprises at least one inducible

promoter, wherein the at least one expression construct further comprises at least one polynucleotide sequence encoding at least one gene product to be transcribed from an inducible promoter. Other examples of the invention include a host cell comprising at least one expression construct comprising at least one inducible promoter and at least one polynucleotide sequence encoding a gene product to be transcribed from an inducible promoter, wherein in certain embodiments at least one gene product is a polypeptide, or is a polypeptide that lacks a signal sequence, or is a polypeptide that forms at least one and fewer than twenty disulfide bonds, or at least two and fewer than seventeen disulfide bonds, or at least eighteen and fewer than one hundred disulfide bonds, or at least three and fewer than ten disulfide bonds, or at least three and fewer than eight disulfide bonds, or is a polypeptide that forms a number of disulfide bonds selected from the group consisting of one, two, three, four, five, six, seven, eight, and nine, or is an immunoglobulin heavy or light chain or a fragment thereof, or is an infliximab heavy or light chain or a fragment thereof.

[0016] Methods of producing gene products are also provided by the invention, such as by growing a culture of a host cell of the invention as described above; and in some embodiments by adding an inducer of at least one inducible promoter to the culture; and in further embodiments by purifying a gene product from the culture. In particular aspects of the invention, the gene product produced by the above method is a polypeptide that forms at least one disulfide bond, or forms at least two and fewer than seventeen disulfide bonds, or forms at least two and fewer than ten disulfide bonds, or forms a number of disulfide bonds selected from the group consisting of one, two, three, four, five, six, seven, eight, and nine, or is an immunoglobulin heavy or light chain or a fragment thereof. The present invention also provides a gene product produced by the above method, wherein the gene product is a polypeptide that lacks a signal peptide, and that in certain embodiments forms at least one disulfide bond, or at least two and fewer than seventeen disulfide bonds, or at least eighteen and fewer than one hundred disulfide bonds, or at least three and fewer than ten disulfide bonds, or at least three and fewer than eight disulfide bonds, or forms a number of disulfide bonds selected from the group consisting of one, two, three, four, five, six, seven, eight, and nine, or is an immunoglobulin heavy or light chain or a fragment thereof. A gene product produced by this method is also provided by the invention, and in some embodiments is part of a multimeric product, which in certain embodiments is an antibody, or in more particular instances, is an aglycosylated antibody, a chimeric antibody, or a human antibody.

[0017] Also provided by the systems and methods of the invention are kits comprising an expression construct as described in the above paragraphs; kits comprising a host cell, the host cell comprising at least one expression construct of the invention as described in the above paragraphs; and kits comprising a gene product or a multimeric product produced by growing a host cell of the invention, which in some embodiments includes adding at least one inducer to the culture, where in some embodiments the multimeric product is an antibody, or in more particular instances, is an aglycosylated antibody, a chimeric antibody, or a human antibody.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] FIG. 1 is a schematic illustration of the inducible coexpression system, which includes a host cell (1) comprising two different inducible expression vectors (3) and (4),

which express different gene products upon application of inducers (5), forming a multimeric product (6).

[0019] FIG. 2 is a schematic illustration of a particular use of the inducible coexpression system, in which the *E. coli* host cell genome (2) encodes a cytoplasmic form of the disulfide isomerase DsbC which lacks a signal peptide; the expression vector pBAD24 (or another expression vector containing an L-arabinose-inducible promoter, such as pBAD240) (3) provides L-arabinose-inducible expression of an immunoglobulin heavy chain, and the expression vector pPRO33 (or another expression vector containing a propionate-inducible promoter, such as pPRO43, pPRO430, pPRO430(CloDF13), pPRO44, or pPRO45) (4) provides propionate-inducible expression of an immunoglobulin light chain; forming upon induction (5) the multimeric antibody product (6).

[0020] FIG. 3 is a schematic illustration of an inducible coexpression system, which includes a host cell (1) comprising an inducible expression vector comprising two different inducible promoters (3) and (4), which express different gene products upon application of inducers (5), forming a multimeric product (6).

[0021] FIG. 4 is a schematic illustration of a particular use of the inducible coexpression system, in which the *E. coli* host cell genome (2) includes genetic alterations such as a cytoplasmic form of the disulfide isomerase DsbC which lacks a signal peptide; a multiple-promoter expression vector such as pSOL provides (3) L-arabinose-inducible expression of an immunoglobulin heavy chain and (4) propionate-inducible expression of an immunoglobulin light chain; forming upon induction (5) the multimeric antibody product (6).

[0022] FIG. 5 is a schematic representation of a vector ("dual vector", also named "pSOL") for use in an inducible coexpression system.

[0023] FIG. 6 shows a time course of coexpression of fluorescent proteins in bacterial cells; in this graph the fluorescence from a red fluorescent protein (RedFP), expressed from a propionate-inducible prpBCDE promoter, is shown. 'pSOL': pSOL-YellowFP-RedFP in *E. coli* ASE(DGH) cells. 'pPRO, pBAD': pBAD24-YellowFP/pPRO33-RedFP in *E. coli* ASE(DGH) cells. The inducer concentrations for each averaged group of samples are shown in the legend, with the propionate concentration listed first (0 mM, 5 mM, or 20 mM), followed by the L-arabinose concentration (0 micromolar, 6.66 micromolar, or 66.61 micromolar).

[0024] FIG. 7 shows a time course of coexpression of fluorescent proteins in bacterial cells; in this graph the fluorescence from a yellow fluorescent protein (YellowFP), expressed from an L-arabinose-inducible araBAD promoter, is shown. 'pSOL': pSOL-YellowFP-RedFP in *E. coli* ASE (DGH) cells. 'pBAD, pPRO': pBAD24-YellowFP/pPRO33-RedFP in *E. coli* ASE(DGH) cells. The inducer concentrations for each averaged group of samples are shown in the legend, with the L-arabinose concentration listed first (0 micromolar, 6.66 micromolar, or 66.61 micromolar), followed by the propionate concentration (0 mM, 5 mM, or 20 mM).

DETAILED DESCRIPTION OF THE INVENTION

[0025] The problem of incompatible coexpression system components is addressed by development of coordinated bacterial coexpression systems which utilize compatible homogeneously inducible promoter systems located on separate expression constructs and, in some embodiments, activated by different inducers. The advantages of the present invention

include, without limitation: 1) improved compatibility of components within the inducible coexpression system; 2) inducible expression of gene products that together form multimers, or other combinations of gene products (coexpression of two or more gene products); 3) improved control of gene product coexpression by independently titratable induction; 4) improved expression of gene product complexes and other products that are difficult to express such as multimeric products and products forming disulfide bonds; 5) streamlined optimization of gene product coexpression.

Coexpressed Gene Products.

[0026] The inducible coexpression systems of the invention are designed to coexpress two or more different gene products that contribute to a desired product. The desired product can be a multimer, formed from coexpressed gene products, or coexpression can be used to produce a combination of the desired product plus an additional product or products that assist in expression of the desired product.

[0027] A 'multimeric product' refers to a set of gene products that coassemble to carry out the function of the multimeric product, and does not refer to transitory associations between gene products and other molecules, such as modifying enzymes (kinases, peptidases, and the like), chaperones, transporters, etc. In certain embodiments of the invention, the multimeric products are heteromultimers. In many embodiments, the coexpressed gene products will be polypeptides that are subunits of multimeric proteins. However, it is also possible to use the inducible coexpression systems of the invention to coexpress multiple different non-coding RNA molecules, or a combination of polypeptide and non-coding RNA gene products. Non-coding RNA molecules, also called non-protein-coding RNA (ncpRNA), non-messenger RNA (nmRNA), and functional RNA (fRNA), include many different types of RNA molecules such as microRNAs that are not messenger RNAs and thus are not templates for the formation of polypeptides through translation.

[0028] Many biologically important products are formed from combinations of different polypeptide chains. In addition to antibodies and antibody fragments, other multimeric products that can be produced by the inducible coexpression methods of the invention include G-coupled protein receptors and ligand-gated ion channels such as nicotinic acetylcholine receptors, GABA_A receptors, glycine receptors, serotonin receptors, glutamate receptors, and ATP-gated receptors such as P2X receptors. The botulinum neurotoxin (often referred to as BoTN, BTX, or as one of its commercially available forms, BOTOX® (onabotulinumtoxinA)) is formed from a heavy chain and a light chain, linked by a disulfide bond (Simpson et al., "The role of the interchain disulfide bond in governing the pharmacological actions of botulinum toxin", J Pharmacol Exp Ther 2004 March; 308(3): 857-864, Epub 2003 Nov. 14). Another example of a product formed from different polypeptide chains is insulin, which in eukaryotes is first translated as a single polypeptide chain, folded, and then cleaved ultimately into two polypeptide chains held together by disulfide bonds. Efficient production of botulinum neurotoxin or of mature insulin in a single host cell are examples of uses of the inducible coexpression methods of the invention.

[0029] The methods of the invention are designed to produce gene products that have been correctly folded and/or assembled into functional products, and that have a desired number of disulfide bonds in the desired locations within such functional products (which can be determined by methods

such as that of Example 11). The number of disulfide bonds for a gene product such as a polypeptide is the total number of intramolecular and intermolecular bonds formed by that gene product when it is present in a desired functional product. For example, a light chain of a human IgG antibody typically has three disulfide bonds (two intramolecular bonds and one intermolecular bond), and a heavy chain of a human IgG antibody typically has seven disulfide bonds (four intramolecular bonds and three intermolecular bonds). In some embodiments, desired gene products are coexpressed with other gene products, such as chaperones, that are beneficial to the production of the desired gene product. Chaperones are proteins that assist the non-covalent folding or unfolding, and/or the assembly or disassembly, of other gene products, but do not occur in the resulting monomeric or multimeric gene product structures when the structures are performing their normal biological functions (having completed the processes of folding and/or assembly). Chaperones can be expressed from an inducible promoter or a constitutive promoter within an expression construct, or can be expressed from the host cell chromosome; preferably, expression of chaperone protein(s) in the host cell is at a sufficiently high level to produce coexpressed gene products that are properly folded and/or assembled into the desired product. Examples of chaperones present in *E. coli* host cells are the folding factors DnaK/DnaJ/GrpE, DsbC/DsbG, GroEL/GroES, IbpA/IbpB, Skp, Tig (trigger factor), and FkpA, which have been used to prevent protein aggregation of cytoplasmic or periplasmic proteins. DnaK/DnaJ/GrpE, GroEL/GroES, and ClpB can function synergistically in assisting protein folding and therefore expression of these chaperones in combinations has been shown to be beneficial for protein expression (Makino et al., "Strain engineering for improved expression of recombinant proteins in bacteria", *Microb Cell Fact* 2011 May 14; 10: 32). When expressing eukaryotic proteins in prokaryotic host cells, a eukaryotic chaperone protein, such as protein disulfide isomerase (PDI) from the same or a related eukaryotic species, is coexpressed or inducibly coexpressed with the desired gene product in certain embodiments of the invention.

Inducible Promoters.

[0030] The following is a description of inducible promoters that can be used in expression constructs for coexpression of gene products, along with some of the genetic modifications that can be made to host cells that contain such expression constructs. Examples of these inducible promoters and related genes are, unless otherwise specified, from *Escherichia coli* (*E. coli*) strain MG1655 (American Type Culture Collection deposit ATCC 700926), which is a substrain of *E. coli* K-12 (American Type Culture Collection deposit ATCC 10798). Table 1 lists the genomic locations, in *E. coli* MG1655, of the nucleotide sequences for these examples of inducible promoters and related genes. Nucleotide and other genetic sequences, referenced by genomic location as in Table 1, are expressly incorporated by reference herein. Additional information about *E. coli* promoters, genes, and strains described herein can be found in many public sources, including the online EcoliWiki resource, located at ecoliwiki.net.

[0031] Arabinose Promoter.

[0032] (As used herein, 'arabinose' means L-arabinose.) Several *E. coli* operons involved in arabinose utilization are inducible by arabinose—araBAD, araC, araE, and araFGH—but the terms 'arabinose promoter' and 'ara promoter' are typically used to designate the araBAD promoter. Several

additional terms have been used to indicate the *E. coli* araBAD promoter, such as P_{ara} , P_{araB} , P_{araBAD} , and P_{BAD} . The use herein of 'ara promoter' or any of the alternative terms given above, means the *E. coli* araBAD promoter. As can be seen from the use of another term, 'araC-araBAD promoter', the araBAD promoter is considered to be part of a bidirectional promoter, with the araBAD promoter controlling expression of the araBAD operon in one direction, and the araC promoter, in close proximity to and on the opposite strand from the araBAD promoter, controlling expression of the araC coding sequence in the other direction. The AraC protein is both a positive and a negative transcriptional regulator of the araBAD promoter. In the absence of arabinose, the AraC protein represses transcription from P_{BAD} , but in the presence of arabinose, the AraC protein, which alters its conformation upon binding arabinose, becomes a positive regulatory element that allows transcription from P_{BAD} . The araBAD operon encodes proteins that metabolize L-arabinose by converting it, through the intermediates L-ribulose and L-ribulose-phosphate, to D-xylulose-5-phosphate. For the purpose of maximizing induction of expression from an arabinose-inducible promoter, it is useful to eliminate or reduce the function of AraA, which catalyzes the conversion of L-arabinose to L-ribulose, and optionally to eliminate or reduce the function of at least one of AraB and AraD, as well. Eliminating or reducing the ability of host cells to decrease the effective concentration of arabinose in the cell, by eliminating or reducing the cell's ability to convert arabinose to other sugars, allows more arabinose to be available for induction of the arabinose-inducible promoter. The genes encoding the transporters which move arabinose into the host cell are araE, which encodes the low-affinity L-arabinose proton symporter, and the araFGH operon, which encodes the subunits of an ABC superfamily high-affinity L-arabinose transporter. Other proteins which can transport L-arabinose into the cell are certain mutants of the LacY lactose permease: the LacY(A177C) and the LacY(A177V) proteins, having a cysteine or a valine amino acid instead of alanine at position 177, respectively (Morgan-Kiss et al., "Long-term and homogeneous regulation of the *Escherichia coli* araBAD promoter by use of a lactose transporter of relaxed specificity", *Proc Natl Acad Sci USA* 2002 May 28; 99(11): 7373-7377). In order to achieve homogenous induction of an arabinose-inducible promoter, it is useful to make transport of arabinose into the cell independent of regulation by arabinose. This can be accomplished by eliminating or reducing the activity of the AraFGH transporter proteins and altering the expression of araE so that it is only transcribed from a constitutive promoter. Constitutive expression of araE can be accomplished by eliminating or reducing the function of the native araE gene, and introducing into the cell an expression construct which includes a coding sequence for the AraE protein expressed from a constitutive promoter. Alternatively, in a cell lacking AraFGH function, the promoter controlling expression of the host cell's chromosomal araE gene can be changed from an arabinose-inducible promoter to a constitutive promoter. In similar manner, as additional alternatives for homogenous induction of an arabinose-inducible promoter, a host cell that lacks AraE function can have any functional AraFGH coding sequence present in the cell expressed from a constitutive promoter. As another alternative, it is possible to express both the araE gene and the araFGH operon from constitutive promoters, by replacing the native araE and araFGH promoters with constitutive promoters in the host

chromosome. It is also possible to eliminate or reduce the activity of both the AraE and the AraFGH arabinose transporters, and in that situation to use a mutation in the LacY lactose permease that allows this protein to transport arabinose. Since expression of the lacY gene is not normally regulated by arabinose, use of a LacY mutant such as LacY (A177C) or LacY(A177V), will not lead to the 'all or none' induction phenomenon when the arabinose-inducible promoter is induced by the presence of arabinose. Because the LacY(A177C) protein appears to be more effective in transporting arabinose into the cell, use of polynucleotides encoding the LacY(A177C) protein is preferred to the use of polynucleotides encoding the LacY(A177V) protein.

[0033] Propionate Promoter.

[0034] The 'propionate promoter' or 'prp promoter' is the promoter for the *E. coli* prpBCDE operon, and is also called P_{prpB} . Like the ara promoter, the prp promoter is part of a bidirectional promoter, controlling expression of the prpB-CDE operon in one direction, and with the prpR promoter controlling expression of the prpR coding sequence in the other direction. The PrpR protein is the transcriptional regulator of the prp promoter, and activates transcription from the prp promoter when the PrpR protein binds 2-methylcitrate ('2-MC'). Propionate (also called propanoate) is the ion, $\text{CH}_3\text{CH}_2\text{COO}^-$, of propionic acid (or 'propanoic acid'), and is the smallest of the 'fatty' acids having the general formula $\text{H}(\text{CH}_2)_n\text{COOH}$ that shares certain properties of this class of molecules: producing an oily layer when salted out of water and having a soapy potassium salt. Commercially available propionate is generally sold as a monovalent cation salt of propionic acid, such as sodium propionate ($\text{CH}_3\text{CH}_2\text{COONa}$), or as a divalent cation salt, such as calcium propionate ($\text{Ca}(\text{CH}_3\text{CH}_2\text{COO})_2$). Propionate is membrane-permeable and is metabolized to 2-MC by conversion of propionate to propionyl-CoA by PrpE (propionyl-CoA synthetase), and then conversion of propionyl-CoA to 2-MC by PrpC (2-methylcitrate synthase). The other proteins encoded by the prpBCDE operon, PrpD (2-methylcitrate dehydratase) and PrpB (2-methylisocitrate lyase), are involved in further catabolism of 2-MC into smaller products such as pyruvate and succinate. In order to maximize induction of a propionate-inducible promoter by propionate added to the cell growth medium, it is therefore desirable to have a host cell with PrpC and PrpE activity, to convert propionate into 2-MC, but also having eliminated or reduced PrpD activity, and optionally eliminated or reduced PrpB activity as well, to prevent 2-MC from being metabolized. Another operon encoding proteins involved in 2-MC biosynthesis is the scpA-argK-scpBC operon, also called the sbm-ygfDGH operon. These genes encode proteins required for the conversion of succinate to propionyl-CoA, which can then be converted to 2-MC by PrpC. Elimination or reduction of the function of these proteins would remove a parallel pathway for the production of the 2-MC inducer, and thus might reduce background levels of expression of a propionate-inducible promoter, and increase sensitivity of the propionate-inducible promoter to exogenously supplied propionate. It has been found that a deletion of sbm-ygfD-ygfG-ygfH-ygfI, introduced into *E. coli* BL21(DE3) to create strain JSB (Lee and Keasling, "A propionate-inducible expression system for enteric bacteria", Appl Environ Microbiol 2005 November; 71(11): 6856-6862), was helpful in reducing background expression in the absence of exogenously supplied inducer, but this deletion also reduced overall expression from the prp

promoter in strain JSB. It should be noted, however, that the deletion sbm-ygfD-ygfG-ygfH-ygfI also apparently affects ygfl, which encodes a putative LysR-family transcriptional regulator of unknown function. The genes sbm-ygfDGH are transcribed as one operon, and ygfl is transcribed from the opposite strand. The 3' ends of the ygfh and ygfl coding sequences overlap by a few base pairs, so a deletion that takes out all of the sbm-ygfDGH operon apparently takes out ygfl coding function as well. Eliminating or reducing the function of a subset of the sbm-ygfDGH gene products, such as YgfG (also called ScpB, methylmalonyl-CoA decarboxylase), or deleting the majority of the sbm-ygfDGH (or scpA-argK-scpBC) operon while leaving enough of the 3' end of the ygfh (or scpC) gene so that the expression of ygfl is not affected, could be sufficient to reduce background expression from a propionate-inducible promoter without reducing the maximal level of induced expression.

[0035] Rhamnose Promoter.

[0036] (As used herein, 'rhamnose' means L-rhamnose.) The 'rhamnose promoter' or 'rha promoter', or P_{rhaSR} , is the promoter for the *E. coli* rhaSR operon. Like the ara and prp promoters, the rha promoter is part of a bidirectional promoter, controlling expression of the rhaSR operon in one direction, and with the rhaBAD promoter controlling expression of the rhaBAD operon in the other direction. The rha promoter, however, has two transcriptional regulators involved in modulating expression: RhaR and RhaS. The RhaR protein activates expression of the rhaSR operon in the presence of rhamnose, while RhaS protein activates expression of the L-rhamnose catabolic and transport operons, rhaBAD and rhaT, respectively (Wickstrum et al., "The AraC/XylS family activator RhaS negatively autoregulates rhaSR expression by preventing cyclic AMP receptor protein activation", J Bacteriol 2010 January; 192(1): 225-232). Although the RhaS protein can also activate expression of the rhaSR operon, in effect RhaS negatively autoregulates this expression by interfering with the ability of the cyclic AMP receptor protein (CRP) to coactivate expression with RhaR to a much greater level. The rhaBAD operon encodes the rhamnose catabolic proteins RhaA (L-rhamnose isomerase), which converts L-rhamnose to L-rhamnulose; RhaB (rhamnulokinase), which phosphorylates L-rhamnulose to form L-rhamnulose-1-P; and RhaD (rhamnulose-1-phosphate aldolase), which converts L-rhamnulose-1-P to L-lactaldehyde and DHAP (dihydroxyacetone phosphate). To maximize the amount of rhamnose in the cell available for induction of expression from a rhamnose-inducible promoter, it is desirable to reduce the amount of rhamnose that is broken down by catalysis, by eliminating or reducing the function of RhaA, or optionally of RhaA and at least one of RhaB and RhaD. *E. coli* cells can also synthesize L-rhamnose from alpha-D-glucose-1-P through the activities of the proteins RmlA, RmlB, RmlC, and RmlD (also called RfbA, RfbB, RfbC, and RfbD, respectively) encoded by the rmlBDACX (or rfbBDACX) operon. To reduce background expression from a rhamnose-inducible promoter, and to enhance the sensitivity of induction of the rhamnose-inducible promoter by exogenously supplied rhamnose, it could be useful to eliminate or reduce the function of one or more of the RmlA, RmlB, RmlC, and RmlD proteins. L-rhamnose is transported into the cell by RhaT, the rhamnose permease or L-rhamnose: proton symporter. As noted above, the expression of RhaT is activated by the transcriptional regulator RhaS. To make expression of RhaT independent of induction by rhamnose

(which induces expression of RhaS), the host cell can be altered so that all functional RhaT coding sequences in the cell are expressed from constitutive promoters. Additionally, the coding sequences for RhaS can be deleted or inactivated, so that no functional RhaS is produced. By eliminating or reducing the function of RhaS in the cell, the level of expression from the rhaSR promoter is increased due to the absence of negative autoregulation by RhaS, and the level of expression of the rhamnose catalytic operon rhaBAD is decreased, further increasing the ability of rhamnose to induce expression from the rha promoter.

[0037] Xylose Promoter.

[0038] (As used herein, 'xylose' means D-xylose.) The xylose promoter, or 'xyl promoter', or P_{xylA} , means the promoter for the *E. coli* xylAB operon. The xylose promoter region is similar in organization to other inducible promoters in that the xylAB operon and the xylFGHR operon are both expressed from adjacent xylose-inducible promoters in opposite directions on the *E. coli* chromosome (Song and Park, "Organization and regulation of the D-xylose operons in *Escherichia coli* K-12: XylR acts as a transcriptional activator", *J Bacteriol.* 1997 November; 179(22): 7025-7032). The transcriptional regulator of both the P_{xylA} and P_{xylF} promoters is XylR, which activates expression of these promoters in the presence of xylose. The xylR gene is expressed either as part of the xylFGHR operon or from its own weak promoter, which is not inducible by xylose, located between the xylH and xylR protein-coding sequences. D-xylose is catabolized by XylA (D-xylose isomerase), which converts D-xylose to D-xylulose, which is then phosphorylated by XylB (xylulokinase) to form D-xylulose-5-P. To maximize the amount of xylose in the cell available for induction of expression from a xylose-inducible promoter, it is desirable to reduce the amount of xylose that is broken down by catalysis, by eliminating or reducing the function of at least XylA, or optionally of both XylA and XylB. The xylFGHR operon encodes XylF, XylG, and XylH, the subunits of an ABC superfamily high-affinity D-xylose transporter. The xylE gene, which encodes the *E. coli* low-affinity xylose-proton symporter, represents a separate operon, the expression of which is also inducible by xylose. To make expression of a xylose transporter independent of induction by xylose, the host cell can be altered so that all functional xylose transporters are expressed from constitutive promoters. For example, the xylFGHR operon could be altered so that the xylFGH coding sequences are deleted, leaving XylR as the only active protein expressed from the xylose-inducible P_{xylF} promoter, and with the xylE coding sequence expressed from a constitutive promoter rather than its native promoter. As another example, the xylR coding sequence is expressed from the P_{xylA} or the P_{xylF} promoter in an expression construct, while either the xylFGHR operon is deleted and xylE is constitutively expressed, or alternatively an xylFGH operon (lacking the xylR coding sequence since

that is present in an expression construct) is expressed from a constitutive promoter and the xylE coding sequence is deleted or altered so that it does not produce an active protein.

[0039] Lactose Promoter.

[0040] The term 'lactose promoter' refers to the lactose-inducible promoter for the lacZYA operon, a promoter which is also called lacZp1; this lactose promoter is located at ca. 365603-365568 (minus strand, with the RNA polymerase binding ('-35') site at ca. 365603-365598, the Pribnow box ('-10') at 365579-365573, and a transcription initiation site at 365567) in the genomic sequence of the *E. coli* K-12 substrain MG1655 (NCBI Reference Sequence NC_000913.2, 11 Jan. 2012). In some embodiments, inducible coexpression systems of the invention can comprise a lactose-inducible promoter such as the lacZYA promoter. In other embodiments, the inducible coexpression systems of the invention comprise one or more inducible promoters that are not lactose-inducible promoters.

[0041] Alkaline Phosphatase Promoter.

[0042] The terms 'alkaline phosphatase promoter' and 'phoA promoter' refer to the promoter for the phoApsiF operon, a promoter which is induced under conditions of phosphate starvation. The phoA promoter region is located at ca. 401647-401746 (plus strand, with the Pribnow box ('-10') at 401695-401701 (Kikuchi et al., "The nucleotide sequence of the promoter and the amino-terminal region of alkaline phosphatase structural gene (phoA) of *Escherichia coli*", *Nucleic Acids Res* 1981 Nov. 11; 9(21): 5671-5678)) in the genomic sequence of the *E. coli* K-12 substrain MG1655 (NCBI Reference Sequence NC_000913.3, 16 Dec. 2014). The transcriptional activator for the phoA promoter is PhoB, a transcriptional regulator that, along with the sensor protein PhoR, forms a two-component signal transduction system in *E. coli*. PhoB and PhoR are transcribed from the phoBR operon, located at ca. 417050-419300 (plus strand, with the PhoB coding sequence at 417,142-417,831 and the PhoR coding sequence at 417,889-419,184) in the genomic sequence of the *E. coli* K-12 substrain MG1655 (NCBI Reference Sequence NC_000913.3, 16 Dec. 2014). The phoA promoter differs from the inducible promoters described above in that it is induced by the lack of a substance—intracellular phosphate—rather than by the addition of an inducer. For this reason the phoA promoter is generally used to direct transcription of gene products that are to be produced at a stage when the host cells are depleted for phosphate, such as the later stages of fermentation. In some embodiments, inducible coexpression systems of the invention can comprise a phoA promoter. In other embodiments, the inducible coexpression systems of the invention comprise one or more inducible promoters that are not phoA promoters.

TABLE 1

Genomic Locations of <i>E. coli</i> Inducible Promoters and Related Genes [1]		
Promoter or Gene	Genomic Location:	Comments:
araBAD promoter	[2] (ca. 70165)-70074 (minus strand)	Smith and Schleif [3]: RNA pol [4] binding ('-35') 70110-70104, Pribnow box ('-10') 70092-70085
araBAD operon	70075-65855 (minus strand)	Smith and Schleif [3]: transcript start 70075, araB ATG 70048; NCBI: araB end of TAA 68348; araA ATG 68337, end of TAA 66835; araD ATG 66550, end of TAA 65855
araC promoter	[2] (ca. 70166)-70241 (plus strand)	Smith and Schleif [3]: RNA pol binding ('-35') 70210-7021, Pribnow box ('-10') 70230-70236

TABLE 1-continued

Genomic Locations of <i>E. coli</i> Inducible Promoters and Related Genes [1]		
Promoter or Gene	Genomic Location:	Comments:
araC gene	70242-71265 (plus strand)	Miyada [5]: transcript start 70242, araC ATG 70387; NCBI: end of TAA 71265
araE promoter	[2] (ca. 2980349)-2980231 (minus strand)	Stoner and Schleif [6]: CRP binding 2980349-2980312, RNA pol binding ('-35') 2980269-2980264, Pribnow box ('-10') 2980244-2980239
araE gene	2980230-2978786 (minus strand)	Stoner and Schleif [6]: transcript start 2980230, ATG 2980204; NCBI: end of TGA 2978786
araFGH promoter	[2] (ca. 1984423)-1984264 (minus strand)	Hendrickson [7]: AraC binding ca. 1984423-ca. 1984414 and 1984326-1984317, CRP binding 1984315-1984297, RNA pol binding ('-35') 1984294-1984289, Pribnow box ('-10') 1984275-1984270
araFGH operon	1984263-1980578 (minus strand)	Hendrickson [7]: transcript start 1984263; NCBI: araF ATG 1984152, end of TAA 1983163; araG ATG 1983093, end of TGA 1981579; araH ATG 1981564, end of TGA 1980578
lacY gene	362403-361150 (minus strand)	Expressed as part of the lacZYA operon. NCBI: ATG 362403, end of TAA 361150
prpBCDE promoter	[2] ca. 347790-ca. 347870 (plus strand)	Keasling [8]: RNA pol binding ('-24') 347844-347848, Pribnow box ('-12') 347855-347859
prpBCDE operon	(ca. 347871)-353816 (plus strand)	Keasling [8]: inferred transcript start ca. 347871, prpB ATG 347906; NCBI: prpB end of TAA 348796; prpC ATG 349236, end of TAA 350405; prpD ATG 350439, end of TAA 351890; prpE ATG 351930, end of TAG 353816
prpR promoter	[2] ca. 347789-ca. 347693 (minus strand)	Keasling [8]: CRP binding 347775-347753, RNA pol binding ('-35') 347728-347723, Pribnow box ('-10') 347707-347702
prpR gene	(ca. 347692)-346081 (minus strand)	Keasling [8]: inferred transcript start ca. 347692, prpR ATG 347667; NCBI: end of TGA 346081
scpA-argK-scpBC (or sbm-ygfDGH) operon	3058872-3064302 (plus strand)	NCBI: scpA ATG 3058872, end of TAA 3061016; argK ATG 3061009, end of TAA 3062004; scpB ATG 3062015, end of TAA 3062800; scpC ATG 3062824, end of TAA 3064302
rhaBAD promoter	[2] (ca. 4095605)-4095496 (minus strand)	Wickstrum [9]: CRP binding 4095595-4095580, RNA pol binding ('-35') 4095530-4095525, Pribnow box ('-10') 4095506-4095501
rhaBAD operon	4095495-4091471 (minus strand)	Wickstrum [9]: transcript start 4095495, rhaB ATG 4095471; NCBI: rhaB end of TGA 4094002; rhaA ATG 4094005, end of TAA 4092746; rhaD ATG 4092295, end of TAA 4091471
rhaSR promoter	[2] (ca. 4095606)-4095733 (plus strand)	Wickstrum [9]: CRP binding 4095615-4095630, RNA pol binding ('-35') 4095699-4095704, Pribnow box ('-10') 4095722-4095727
rhaSR operon	4095734-4097517 (plus strand)	Wickstrum [9]: transcript start 4095734, rhaS ATG 4095759; NCBI: rhaS end of TAA 4096595; rhaR ATG 4096669, end of TAA 4097517
rfbBDACX (or rmlBDACX) operon	2111085-2106361 (minus strand)	NCBI: rfb GTG 2111085, end of TAA 2110000; rfbD ATG 2110000, end of TAA 2109101; rfbA ATG 2109043, end of TAA 2108162; rfbC ATG 2108162, end of TGA 2107605; rfbX ATG 2107608, end of TGA 2106361
rhaT promoter	[2] (ca. 4098690)-4098590 (minus strand)	Via [10]: CRP binding 4098690-4098675, RNA pol binding ('-35') 4098621-4098616, Pribnow box ('-10') 4098601-4098596
rhaT gene	4098589-4097514 (minus strand)	Via [10]: transcript start 4098589, rhaT ATG 4098548; NCBI: rhaT end of TAA 4097514
xylAB promoter	[2] (ca. 3728960)-3728831 (minus strand)	Song and Park [11]: CRP binding 3728919-3728901, RNA pol binding ('-35') 3728865-3728860, Pribnow box ('-10') 3728841-3728836
xylAB operon	3728830-3725940 (minus strand)	Song and Park [11]: transcript start 3728830, xylA ATG 3728788; NCBI: xylA end of TAA 3727466; xylB ATG 3727394, end of TAA 3725940
xylFGHR promoter	[2] (ca. 3728961)-3729091 (plus strand)	Song and Park [11]: RNA pol binding ('-35') 3729058-3729063, Pribnow box ('-10') 3729080-3729085

TABLE 1-continued

Genomic Locations of <i>E. coli</i> Inducible Promoters and Related Genes [1]		
Promoter or Gene	Genomic Location:	Comments:
xylFGHR operon	3729092-3734180 (plus strand)	Song and Park [11]; transcript start 3729092, xylF ATG 3729154; NCBI: xylF end of TAA 3730146, xylG ATG 3730224, end of TGA 3731765; xylH ATG 3731743, end of TGA 3732924; xylR ATG 3733002, end of TAG 3734180
xylE promoter	[2] ca. 4240482- ca. 4240320 (minus strand)	Davis and Henderson [12]; possible Pribnow box ('-10') 4240354-4240349, possible Pribnow box ('-10') 4240334-4240329
xylE gene	(ca. 4240319)- 4238802 (minus strand)	Davis and Henderson [12]; inferred transcript start ca. 4240319, xylE ATG 4240277, end of TAA 4238802

Notes for Table 1:

[1] All genomic sequence locations refer to the genomic sequence of *E. coli* K-12 substrain MG1655, provided by the National Center for Biotechnology Information (NCBI) as NCBI Reference Sequence NC_000913.2, 11 JAN. 2012.

[2] The location of the 5' (or 'upstream') end of the promoter region is approximated; for 'bidirectional' promoters, a nucleotide sequence location that is approximately equidistant between the transcription start sites is selected as the designated 5' end for both of the individual promoters. In practice, the promoter portion of an expression construct can have somewhat less sequence at its 5' end than the promoter sequences as indicated in the table, or it can have a nucleotide sequence that includes additional sequence from the region 5' (or 'upstream') of the promoter sequences as indicated in the table, as long as it retains the ability to promote transcription of a downstream coding sequence in an inducible fashion.

[3] Smith and Schleif, "Nucleotide sequence of the L-arabinose regulatory region of *Escherichia coli* K12", J Biol Chem 1978 Oct. 10; 253(19): 6931-6933.

[4] 'RNA pol' indicates RNA polymerase throughout the table.

[5] Miyada et al., "DNA sequence of the araC regulatory gene from *Escherichia coli* B/r", Nucleic Acids Res 1980 Nov. 25; 8(22): 5267-5274.

[6] Stoner and Schleif, "E. coli araE regulatory region araE codes for the low affinity L-arabinose uptake protein", GenBank Database Accession X00272.1, revision date 6 JUL. 1989.

[7] Hendrickson et al., "Sequence elements in the *Escherichia coli* araFGH promoter", J Bacteriol 1992 November; 174(21): 6862-6871.

[8] U.S. Pat. No. 8,178,338 B2; May 15, 2012; Keasling, Jay; FIG. 9.

[9] Wickström et al., "The AraC/XylS family activator RhaS negatively autoregulates rhaSR expression by preventing cyclic AMP receptor protein activation", J Bacteriol 2010 January; 192(1): 225-232.

[10] Via et al., "Transcriptional regulation of the *Escherichia coli* rhaF" gene", Microbiology 1996 July; 142(Pt 7): 1833-1840.

[11] Song and Park, "Organization and regulation of the D-xylose operons in *Escherichia coli* K-12: XylR acts as a transcriptional activator", J Bacteriol. 1997 November; 179(22): 7025-7032.

[12] Davis and Henderson, "The cloning and DNA sequence of the gene xylE for xylose-proton symport in *Escherichia coli* K12", J Biol Chem 1987 Oct. 15; 262(29): 13928-13932.

Expression Constructs.

[0043] Expression constructs are polynucleotides designed for the expression of one or more gene products of interest, and thus are not naturally occurring molecules. Expression constructs can be integrated into a host cell chromosome, or maintained within the host cell as polynucleotide molecules replicating independently of the host cell chromosome, such as plasmids or artificial chromosomes. An example of an expression construct is a polynucleotide resulting from the insertion of one or more polynucleotide sequences into a host cell chromosome, where the inserted polynucleotide sequences alter the expression of chromosomal coding sequences. An expression vector is a plasmid expression construct specifically used for the expression of one or more gene products. One or more expression constructs can be integrated into a host cell chromosome or be maintained on an extrachromosomal polynucleotide such as a plasmid or artificial chromosome. The following are descriptions of particular types of polynucleotide sequences that can be used in expression constructs for the coexpression of gene products.

[0044] Origins of Replication.

[0045] Expression constructs must comprise an origin of replication, also called a replicon, in order to be maintained within the host cell as independently replicating polynucleotides. Different replicons that use the same mechanism for replication cannot be maintained together in a single host cell through repeated cell divisions. As a result, plasmids can be categorized into incompatibility groups depending on the origin of replication that they contain, as shown in Table 2.

TABLE 2

Origins of Replication and Representative Plasmids for Use in Expression Constructs [1]			
Incompatibility Group:	Origin of Replication:	Copy Number:	Representative Plasmids (ATCC Deposit No.):
colE1, pMB1	colE1 pMB1 Modified pMB1	15-20 15-20 500-700	colE1 (ATCC 27138) pBR322 (ATCC 31344) pUC9 (ATCC 37252)
IncFII, pT181 F, P1, p15A, pSC101, R6K, RK2 [2]	R1(ts) p15A pSC101	15-120 18-22 ~5	pMOB45 (ATCC 37106) pACYC177 (ATCC 37031); pACYC184 (ATCC 37033); pPRO33 (Addgene 17810) [3] pSC101 (ATCC 37032); pGBM1 (ATCC 87497)
CloDF13 [4]	RK2 CloDF13	4-7 [2] 20-40 [4]	RK2 (ATCC 37125) pCDFDuet™-1 (EMD Millipore Catalog No. 71340-3)
ColA [4]	ColA	20-40 [4]	pCOLADuet™-1 (EMD Millipore Catalog No. 71406-3)
RSF1030 [4]	RSF1030 (also called NTP1)	>100 [4]	pRSFDuet™-1 (EMD Millipore Catalog No. 71341-3)

Notes for Table 2:

[1] Adapted from www.bio.davidson.edu/courses/Molbio/Protocols/ORIs.html, and Sambrook and Russell, "Molecular Cloning: A laboratory manual", 3rd Ed., Cold Spring Harbor Laboratory Press, 2001.

[2] Kües and Stahl, "Replication of plasmids in gram-negative bacteria", Microbiol Rev 1989 December; 53(4): 491-516.

[3] The pPRO33 plasmid (U.S. Pat. No. 8,178,338 B2; May 15, 2012; Keasling, Jay) is available from Addgene (www.addgene.org) as Addgene plasmid 17810.

[4] openwetware.org/wiki/CH391L/S12/Origins_of_Replication; accessed 3 Aug. 2013.

Origins of replication can be selected for use in expression constructs on the basis of incompatibility group, copy number, and/or host range, among other criteria. As described above, if two or more different expression constructs are to be used in the same host cell for the coexpression of multiple gene products, it is best if the different expression constructs contain origins of replication from different incompatibility groups: a pMB1 replicon in one expression construct and a p15A replicon in another, for example. The average number of copies of an expression construct in the cell, relative to the number of host chromosome molecules, is determined by the origin of replication contained in that expression construct. Copy number can range from a few copies per cell to several hundred (Table 2). In one embodiment of the invention, different expression constructs are used which comprise inducible promoters that are activated by the same inducer, but which have different origins of replication. By selecting origins of replication that maintain each different expression construct at a certain approximate copy number in the cell, it is possible to adjust the levels of overall production of a gene product expressed from one expression construct, relative to another gene product expressed from a different expression construct. As an example, to coexpress subunits A and B of a multimeric protein, an expression construct is created which comprises the colE1 replicon, the ara promoter, and a coding sequence for subunit A expressed from the ara promoter: 'colE1-P_{ara}-A'. Another expression construct is created comprising the p15A replicon, the ara promoter, and a coding sequence for subunit B: 'p15A-P_{ara}-B'. These two expression constructs can be maintained together in the same host cells, and expression of both subunits A and B is induced by the addition of one inducer, arabinose, to the growth medium. If the expression level of subunit A needed to be significantly increased relative to the expression level of subunit B, in order to bring the stoichiometric ratio of the expressed amounts of the two subunits closer to a desired ratio, for example, a new expression construct for subunit A could be created, having a modified pMB1 replicon as is found in the origin of replication of the pUC9 plasmid ('pUC9ori'): pUC9ori-P_{ara}-A. Expressing subunit A from a high-copy-number expression construct such as pUC9ori-P_{ara}-A should increase the amount of subunit A produced relative to expression of subunit B from p15A-P_{ara}-B. In a similar fashion, use of an origin of replication that maintains expression constructs at a lower copy number, such as pSC101, could reduce the overall level of a gene product expressed from that construct. Selection of an origin of replication can also determine which host cells can maintain an expression construct comprising that replicon. For example, expression constructs comprising the colE1 origin of replication have a relatively narrow range of available hosts, species within the Enterobacteriaceae family, while expression constructs comprising the RK2 replicon can be maintained in *E. coli*, *Pseudomonas aeruginosa*, *Pseudomonas putida*, *Azotobacter vinelandii*, and *Alcaligenes eutrophus*, and if an expression construct comprises the RK2 replicon and some regulator genes from the RK2 plasmid, it can be maintained in host cells as diverse as *Sinorhizobium meliloti*, *Agrobacterium tumefaciens*, *Caulobacter crescentus*, *Acinetobacter calcoaceticus*, and *Rhodobacter sphaeroides* (Kües and Stahl, "Replication of plasmids in gram-negative bacteria", *Microbiol Rev* 1989 December; 53(4): 491-516).

[0046] Similar considerations can be employed to create expression constructs for inducible coexpression in eukary-

otic cells. For example, the 2-micron circle plasmid of *Saccharomyces cerevisiae* is compatible with plasmids from other yeast strains, such as pSR1 (ATCC Deposit Nos. 48233 and 66069; Araki et al., "Molecular and functional organization of yeast plasmid pSR1", *J Mol Biol* 1985 Mar. 20; 182 (2): 191-203) and pKD1 (ATCC Deposit No. 37519; Chen et al., "Sequence organization of the circular plasmid pKD1 from the yeast *Kluyveromyces drosophilae*", *Nucleic Acids Res* 1986 Jun. 11; 14(11): 4471-4481).

[0047] Selectable Markers.

[0048] Expression constructs usually comprise a selection gene, also termed a selectable marker, which encodes a protein necessary for the survival or growth of host cells in a selective culture medium. Host cells not containing the expression construct comprising the selection gene will not survive in the culture medium. Typical selection genes encode proteins that confer resistance to antibiotics or other toxins, or that complement auxotrophic deficiencies of the host cell. One example of a selection scheme utilizes a drug such as an antibiotic to arrest growth of a host cell. Those cells that contain an expression construct comprising the selectable marker produce a protein conferring drug resistance and survive the selection regimen. Some examples of antibiotics that are commonly used for the selection of selectable markers (and abbreviations indicating genes that provide antibiotic resistance phenotypes) are: ampicillin (Amp^R), chloramphenicol (Cm^R or Cm^K), kanamycin (Kan^R), spectinomycin (Spc^R), streptomycin (Str^R), and tetracycline (Tet^R). Many of the representative plasmids in Table 2 comprise selectable markers, such as pBR322 (Amp^R, Tet^R); pMOB45 (Cm^R, Tet^R); pACYC177 (Amp^R, Kan^R); and pGBM1 (Spc^R, Str^R). The native promoter region for a selection gene is usually included, along with the coding sequence for its gene product, as part of a selectable marker portion of an expression construct. Alternatively, the coding sequence for the selection gene can be expressed from a constitutive promoter.

[0049] Inducible Promoter.

[0050] As described herein, there are several different inducible promoters that can be included in expression constructs as part of the inducible coexpression systems of the invention. Preferred inducible promoters share at least 80% polynucleotide sequence identity (more preferably, at least 90% identity, and most preferably, at least 95% identity) to at least 30 (more preferably, at least 40, and most preferably, at least 50) contiguous bases of a promoter polynucleotide sequence as defined in Table 1 by reference to the *E. coli* K-12 substrain MG1655 genomic sequence, where percent polynucleotide sequence identity is determined using the methods of Example 11. Under 'standard' inducing conditions (see Example 5), preferred inducible promoters have at least 75% (more preferably, at least 100%, and most preferably, at least 110%) of the strength of the corresponding 'wild-type' inducible promoter of *E. coli* K-12 substrain MG1655, as determined using the quantitative PCR method of De Mey et al. (Example 6). Within the expression construct, an inducible promoter is placed 5' to (or 'upstream of') the coding sequence for the gene product that is to be inducibly expressed, so that the presence of the inducible promoter will direct transcription of the gene product coding sequence in a 5' to 3' direction relative to the coding strand of the polynucleotide encoding the gene product.

[0051] Ribosome Binding Site.

[0052] For polypeptide gene products, the nucleotide sequence of the region between the transcription initiation

site and the initiation codon of the coding sequence of the gene product that is to be inducibly expressed corresponds to the 5' untranslated region ('UTR') of the mRNA for the polypeptide gene product. Preferably, the region of the expression construct that corresponds to the 5' UTR comprises a polynucleotide sequence similar to the consensus ribosome binding site (RBS, also called the Shine-Dalgarno sequence) that is found in the species of the host cell. In prokaryotes (archaea and bacteria), the RBS consensus sequence is GGAGG or GGAGGU, and in bacteria such as *E. coli*, the RBS consensus sequence is AGGAGG or AGGAGGU. The RBS is typically separated from the initiation codon by 5 to 10 intervening nucleotides. In expression constructs, the RBS sequence is preferably at least 55% identical to the AGGAGGU consensus sequence, more preferably at least 70% identical, and most preferably at least 85% identical, and is separated from the initiation codon by 5 to 10 intervening nucleotides, more preferably by 6 to 9 intervening nucleotides, and most preferably by 6 or 7 intervening nucleotides. The ability of a given RBS to produce a desirable translation initiation rate can be calculated at the website salis.psu.edu/software/RBSLibraryCalculatorSearchMode, using the RBS Calculator; the same tool can be used to optimize a synthetic RBS for a translation rate across a 100,000+ fold range (Salis, "The ribosome binding site calculator", Methods Enzymol 2011; 498: 19-42).

[0053] Multiple Cloning Site.

[0054] A multiple cloning site (MCS), also called a polylinker, is a polynucleotide that contains multiple restriction sites in close proximity to or overlapping each other. The restriction sites in the MCS typically occur once within the MCS sequence, and preferably do not occur within the rest of the plasmid or other polynucleotide construct, allowing restriction enzymes to cut the plasmid or other polynucleotide construct only within the MCS. Examples of MCS sequences are those in the pBAD series of expression vectors, including pBAD18, pBAD18-Cm, pBAD18-Kan, pBAD24, pBAD28, pBAD30, and pBAD33 (Guzman et al., "Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter", J Bacteriol 1995 July; 177 (14): 4121-4130); or those in the pPRO series of expression vectors derived from the pBAD vectors, such as pPRO18, pPRO18-Cm, pPRO18-Kan, pPRO24, pPRO30, and pPRO33 (U.S. Pat. No. 8,178,338 B2; May 15, 2012; Keasling, Jay). A multiple cloning site can be used in the creation of an expression construct: by placing a multiple cloning site 3' to (or downstream of) a promoter sequence, the MCS can be used to insert the coding sequence for a gene product to be coexpressed into the construct, in the proper location relative to the promoter so that transcription of the coding sequence will occur. Depending on which restriction enzymes are used to cut within the MCS, there may be some part of the MCS sequence remaining within the expression construct after the coding sequence or other polynucleotide sequence is inserted into the expression construct. Any remaining MCS sequence can be upstream or, or downstream of, or on both sides of the inserted sequence. A ribosome binding site can be placed upstream of the MCS, preferably immediately adjacent to or separated from the MCS by only a few nucleotides, in which case the RBS would be upstream of any coding sequence inserted into the MCS. Another alternative is to include a ribosome binding site within the MCS, in which case the choice of restriction enzymes used to cut within the MCS will determine whether the RBS is retained, and in what relation

to, the inserted sequences. A further alternative is to include a RBS within the polynucleotide sequence that is to be inserted into the expression construct at the MCS, preferably in the proper relation to any coding sequences to stimulate initiation of translation from the transcribed messenger RNA.

[0055] Expression from Constitutive Promoters.

[0056] Expression constructs of the invention can also comprise coding sequences that are expressed from constitutive promoters. Unlike inducible promoters, constitutive promoters initiate continual gene product production under most growth conditions. One example of a constitutive promoter is that of the Tn3 bla gene, which encodes beta-lactamase and is responsible for the ampicillin-resistance (Amp^R) phenotype conferred on the host cell by many plasmids, including pBR322 (ATCC 31344), pACYC177 (ATCC 37031), and pBAD24 (ATCC 87399). Another constitutive promoter that can be used in expression constructs is the promoter for the *E. coli* lipoprotein gene, lpp, which is located at positions 1755731-1755406 (plus strand) in *E. coli* K-12 substrain MG1655 (Inouye and Inouye, "Up-promoter mutations in the lpp gene of *Escherichia coli*", Nucleic Acids Res 1985 May 10; 13(9): 3101-3110). A further example of a constitutive promoter that has been used for heterologous gene expression in *E. coli* is the trpLEDCBA promoter, located at positions 1321169-1321133 (minus strand) in *E. coli* K-12 substrain MG1655 (Windass et al., "The construction of a synthetic *Escherichia coli* trp promoter and its use in the expression of a synthetic interferon gene", Nucleic Acids Res 1982 Nov. 11; 10(21): 6639-6657). Constitutive promoters can be used in expression constructs for the expression of selectable markers, as described herein, and also for the constitutive expression of other gene products useful for the coexpression of the desired product. For example, transcriptional regulators of the inducible promoters, such as AraC, PrpR, RhaR, and XylR, if not expressed from a bidirectional inducible promoter, can alternatively be expressed from a constitutive promoter, on either the same expression construct as the inducible promoter they regulate, or a different expression construct. Similarly, gene products useful for the production or transport of the inducer, such as PrpEC, AraE, or Rha, or proteins that modify the reduction-oxidation environment of the cell, as a few examples, can be expressed from a constitutive promoter within an expression construct. Gene products useful for the production of coexpressed gene products, and the resulting desired product, also include chaperone proteins, cofactor transporters, etc.

[0057] Signal Peptides.

[0058] Polypeptide gene products coexpressed by the methods of the invention can contain signal peptides or lack them, depending on whether it is desirable for such gene products to be exported from the host cell cytoplasm into the periplasm, or to be retained in the cytoplasm, respectively. Signal peptides (also termed signal sequences, leader sequences, or leader peptides) are characterized structurally by a stretch of hydrophobic amino acids, approximately five to twenty amino acids long and often around ten to fifteen amino acids in length, that has a tendency to form a single alpha-helix. This hydrophobic stretch is often immediately preceded by a shorter stretch enriched in positively charged amino acids (particularly lysine). Signal peptides that are to be cleaved from the mature polypeptide typically end in a stretch of amino acids that is recognized and cleaved by signal peptidase. Signal peptides can be characterized functionally by the ability to direct transport of a polypeptide, either co-

translationally or post-translationally, through the plasma membrane of prokaryotes (or the inner membrane of gram negative bacteria like *E. coli*), or into the endoplasmic reticulum of eukaryotic cells. The degree to which a signal peptide enables a polypeptide to be transported into the periplasmic space of a host cell like *E. coli*, for example, can be determined by separating periplasmic proteins from proteins retained in the cytoplasm, using a method such as that provided in Example 12.

Host Cells.

[0059] The inducible coexpression systems of the invention are designed to express multiple gene products; in certain embodiments of the invention, the gene products are coexpressed in a host cell. Examples of host cells are provided that allow for the efficient and cost-effective inducible coexpression of components of multimeric products. Host cells can include, in addition to isolated cells in culture, cells that are part of a multicellular organism, or cells grown within a different organism or system of organisms. In addition, the expression constructs of the inducible coexpression systems of the invention can be used in cell-free systems, such as those based on wheat germ extracts or on bacterial cell extracts, such as a continuous-exchange cell-free (CECF) protein synthesis system using *E. coli* extracts and an incubation apparatus such as the RTS ProteoMaster (Roche Diagnostics GmbH; Mannheim, Germany) (Jun et al., "Continuous-exchange cell-free protein synthesis using PCR-generated DNA and an RNase E-deficient extract", *Biotechniques* 2008 March; 44(3): 387-391).

[0060] Prokaryotic Host Cells.

[0061] In some embodiments of the invention, expression constructs designed for coexpression of gene products are provided in host cells, preferably prokaryotic host cells. Prokaryotic host cells can include archaea (such as *Haloferax volcanii*, *Sulfolobus solfataricus*), Gram-positive bacteria (such as *Bacillus subtilis*, *Bacillus licheniformis*, *Brevibacillus choshinensis*, *Lactobacillus brevis*, *Lactobacillus buchneri*, *Lactococcus lactis*, and *Streptomyces lividans*), or Gram-negative bacteria, including Alphaproteobacteria (*Agrobacterium tumefaciens*, *Caulobacter crescentus*, *Rhodobacter sphaeroides*, and *Sinorhizobium meliloti*), Betaproteobacteria (*Alcaligenes eutrophus*), and Gammaproteobacteria (*Acinetobacter calcoaceticus*, *Azotobacter vinelandii*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Pseudomonas putida*). Preferred host cells include Gammaproteobacteria of the family Enterobacteriaceae, such as *Enterobacter*, *Erwinia*, *Escherichia* (including *E. coli*), *Klebsiella*, *Proteus*, *Salmonella* (including *Salmonella typhimurium*), *Serratia* (including *Serratia marcescans*), and *Shigella*.

[0062] Eukaryotic Host Cells.

[0063] Many additional types of host cells can be used for the inducible coexpression systems of the invention, including eukaryotic cells such as yeast (*Candida shehatae*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, other *Kluyveromyces* species, *Pichia pastoris*, *Saccharomyces cerevisiae*, *Saccharomyces pastorianus* also known as *Saccharomyces carlsbergensis*, *Schizosaccharomyces pombe*, *Dekkera/Brettanomyces* species, and *Yarrowia lipolytica*); other fungi (*Aspergillus nidulans*, *Aspergillus niger*, *Neurospora crassa*, *Penicillium*, *Tolypocladium*, *Trichoderma reesia*); insect cell lines (*Drosophila melanogaster* Schneider 2 cells and *Spodoptera frugiperda* Sf9 cells); and mammalian cell

lines including immortalized cell lines (Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human embryonic kidney (HEK, 293, or HEK-293) cells, and human hepatocellular carcinoma cells (Hep G2)). The above host cells are available from the American Type Culture Collection.

Alterations to Host Cell Gene Functions.

[0064] Certain alterations can be made to the gene functions of host cells comprising inducible expression constructs, to promote efficient and homogeneous induction of the host cell population by an inducer. Preferably, the combination of expression constructs, host cell genotype, and induction conditions results in at least 75% (more preferably at least 85%, and most preferably, at least 95%) of the cells in the culture expressing gene product from each induced promoter, as measured by the method of Khlebnikov et al. described in Example 6. For host cells other than *E. coli*, these alterations can involve the function of genes that are structurally similar to an *E. coli* gene, or genes that carry out a function within the host cell similar to that of the *E. coli* gene. Alterations to host cell gene functions include eliminating or reducing gene function by deleting the gene protein-coding sequence in its entirety, or deleting a large enough portion of the gene, inserting sequence into the gene, or otherwise altering the gene sequence so that a reduced level of functional gene product is made from that gene. Alterations to host cell gene functions also include increasing gene function by, for example, altering the native promoter to create a stronger promoter that directs a higher level of transcription of the gene, or introducing a missense mutation into the protein-coding sequence that results in a more highly active gene product. Alterations to host cell gene functions include altering gene function in any way, including for example, altering a native inducible promoter to create a promoter that is constitutively activated. In addition to alterations in gene functions for the transport and metabolism of inducers, as described herein with relation to inducible promoters, and an altered expression of chaperone proteins, it is also possible to alter the carbon catabolite repression (CCR) regulatory system and/or the reduction-oxidation environment of the host cell.

[0065] Carbon Catabolite Repression (CCR).

[0066] The presence of an active CCR regulatory system within a host can affect the ability of an inducer to activate transcription from an inducible promoter. For example, when a host cell such as *E. coli* is grown in a medium containing glucose, genes needed for the utilization of other carbon sources, such as the araBAD and prpBCDE operons, are expressed at a low level if at all, even if the arabinose or propionate inducer is also present in the growth medium. There is also a hierarchy of utilization of carbon sources other than glucose: as in the case of the ara and prp inducible promoter systems, where the presence of arabinose reduces the ability of propionate to induce expression from the prpBCDE promoter (Park et al., "The mechanism of sugar-mediated catabolite repression of the propionate catabolic genes in *Escherichia coli*", *Gene* 2012 Aug. 1; 504(1): 116-121; *Epib* 2012 May 3). The CCR mechanism of the cell therefore makes it more difficult to use two or more carbon-source inducers in an inducible coexpression system, as the presence of the inducer that is the preferred carbon source will inhibit induction by less-preferred carbon sources. The Park et al. authors attempted to relieve the repression of the prp pro-

moter by arabinose, by using either a mutant *crp* gene that produces an altered cAMP receptor protein that can function independently of cAMP, or a deletion of PTS (phosphotransferase system) genes involved in the regulation of CCR; both approaches were largely unsuccessful. However, the PTS-knockout strain used by the Park et al. authors is based on strain TP2811, which is a deletion of the *E. coli* *ptsHI-crr* operon (Hernández-Montalvo et al., "Characterization of sugar mixtures utilization by an *Escherichia coli* mutant devoid of the phosphotransferase system", Appl Microbiol Biotechnol 2001 October; 57(1-2): 186-191). Deletion of the entire *ptsHI-crr* operon has been found to affect total cAMP synthesis more significantly than a deletion of just the *crr* gene (Levy et al., "Cyclic AMP synthesis in *Escherichia coli* strains bearing known deletions in the pts phosphotransferase operon", Gene 1990 Jan. 31; 86(1): 27-33). A different approach is to eliminate or reduce the function of *ptsG* gene in the host cell, which encodes glucose-specific EII A (EII A^{glc}), a key element for CCR in *E. coli* (Kim et al., "Simultaneous consumption of pentose and hexose sugars: an optimal microbial phenotype for efficient fermentation of lignocellulosic biomass", Appl Microbiol Biotechnol 2010 November; 88(5): 1077-1085, Epub 2010 Sep. 14). Another alteration in the genome of a host cell such as *E. coli*, which leads to increased transcription of the *prp* promoter, is to eliminate or reduce the gene function of the *ascG* gene, which encodes AscG. AscG is the repressor of the beta-D-glucoside-utilization operon *ascFB* under normal growth conditions, and also represses transcription of the *prp* promoter; disruption of the *AscG* coding sequence has been shown to increase transcription from the *prp* promoter (Ishida et al., "Participation of regulator AscG of the beta-glucoside utilization operon in regulation of the propionate catabolism operon", J Bacteriol 2009 October; 191(19): 6136-6144; Epub 2009 Jul. 24). A further alternative is to increase expression of the transcriptional regulator of promoters inducible by the less-preferred carbon-source inducer, by placing it either under the control of a strong constitutive promoter, or under the control of the more-preferred carbon-source inducer. For example, to increase the induction of genes needed for the utilization of the less-preferred carbon source xylose in the presence of the more-preferred arabinose, the coding sequence for XylR is placed into the *E. coli* *araBAD* operon (Groff et al., "Supplementation of intracellular XylR leads to cot utilization of hemicellulose sugars", Appl Environ Microbiol 2012 April; 78(7): 2221-2229, Epub 2012 Jan. 27). Host cells comprising inducible coexpression constructs therefore preferably include an increased level of gene function for transcriptional regulators of promoters inducible by the less-preferred carbon-source inducer(s), and an eliminated or reduced gene function for genes involved in the CCR system, such as *crr* and/or *ptsG* and/or *ascG*.

[0067] Cellular Transport of Cofactors.

[0068] When using the inducible coexpression systems of the invention to produce enzymes that require cofactors for function, it is helpful to use a host cell capable of synthesizing the cofactor from available precursors, or taking it up from the environment. Common cofactors include ATP, coenzyme A, flavin adenine dinucleotide (FAD), NAD⁺/NADH, and heme.

[0069] Host Cell Reduction-Oxidation Environment.

[0070] Many multimeric gene products, such as antibodies, contain disulfide bonds. The cytoplasm of *E. coli* and many other cells is normally maintained in a reduced state by the thioredoxin and the glutaredoxin/glutathione enzyme sys-

tems. This precludes the formation of disulfide bonds in the cytoplasm, and proteins that need disulfide bonds are exported into the periplasm where disulfide bond formation and isomerization is catalyzed by the Dsb system, comprising DsbABCD and DsbG. Increased expression of the cysteine oxidase DsbA, the disulfide isomerase DsbC, or combinations of the Dsb proteins, which are all normally transported into the periplasm, has been utilized in the expression of heterologous proteins that require disulfide bonds (Makino et al., "Strain engineering for improved expression of recombinant proteins in bacteria", Microb Cell Fact 2011 May 14; 10: 32). It is also possible to express cytoplasmic forms of these Dsb proteins, such as a cytoplasmic version of DsbC ('cDsbC'), that lacks a signal peptide and therefore is not transported into the periplasm. Cytoplasmic Dsb proteins such as cDsbC are useful for making the cytoplasm of the host cell more oxidizing and thus more conducive to the formation of disulfide bonds in heterologous proteins produced in the cytoplasm. The host cell cytoplasm can also be made more oxidizing by altering the thioredoxin and the glutaredoxin/glutathione enzyme systems directly: mutant strains defective in glutathione reductase (gor) or glutathione synthetase (gshB), together with thioredoxin reductase (trxB), render the cytoplasm oxidizing. These strains are unable to reduce ribonucleotides and therefore cannot grow in the absence of exogenous reductant, such as dithiothreitol (DTT). Suppressor mutations (ahpC*) in the gene *ahpC*, which encodes the peroxiredoxin AhpC, convert it to a disulfide reductase that generates reduced glutathione, allowing the channeling of electrons onto the enzyme ribonucleotide reductase and enabling the cells defective in gor and trxB, or defective in gshB and trxB, to grow in the absence of DTT. A different class of mutated forms of AhpC can allow strains, defective in the activity of gamma-glutamylcysteine synthetase (gshA) and defective in trxB, to grow in the absence of DTT; these include AhpC V164G, AhpC S71F, AhpC E173/S71F, AhpC E171Ter, and AhpC dup162-169 (Faulkner et al., "Functional plasticity of a peroxidase allows evolution of diverse disulfide-reducing pathways", Proc Natl Acad Sci USA 2008 May 6; 105(18): 6735-6740, Epub 2008 May 2). In such strains with oxidizing cytoplasm, exposed protein cysteines become readily oxidized in a process that is catalyzed by thioredoxins, in a reversal of their physiological function, resulting in the formation of disulfide bonds.

[0071] Another alteration that can be made to host cells is to express the sulphydryl oxidase Erv1p from the inner membrane space of yeast mitochondria in the host cell cytoplasm, which has been shown to increase the production of a variety of complex, disulfide-bonded proteins of eukaryotic origin in the cytoplasm of *E. coli*, even in the absence of mutations in gor or trxB (Nguyen et al., "Pre-expression of a sulphydryl oxidase significantly increases the yields of eukaryotic disulfide bond containing proteins expressed in the cytoplasm of *E. coli*", Microb Cell Fact 2011 Jan. 7; 10: 1). Host cells comprising inducible coexpression constructs preferably also express cDsbC and/or Erv1p, are deficient in trxB gene function, are also deficient in the gene function of either gor, gshB, or gshA, and express an appropriate mutant form of AhpC so that the host cells can be grown in the absence of DTT.

[0072] Glycosylation of Polypeptide Gene Products.

[0073] Host cells can have alterations in their ability to glycosylate polypeptides. For example, eukaryotic host cells can have eliminated or reduced gene function in glycosyltransferase and/or oligo-saccharyltransferase genes, impair-

ing the normal eukaryotic glycosylation of polypeptides to form glycoproteins. Prokaryotic host cells such as *E. coli*, which do not normally glycosylate polypeptides, can be altered to express a set of eukaryotic and prokaryotic genes that provide a glycosylation function (DeLisa et al., "Glycosylated protein expression in prokaryotes", WO2009089154A2, 2009 Jul. 16).

[0074] Available Host Cell Strains with Altered Gene Functions.

[0075] To create preferred strains of host cells to be used in the inducible coexpression systems and methods of the invention, it is useful to start with a strain that already comprises desired genetic alterations (Table 3).

TABLE 3

Host Cell Strains		
Strain:	Genotype:	Source:
<i>E. coli</i> TOP10	F- mcrA Δ(mrr-hsdRMS-mcrBC) φ80lacZAM15 ΔlacX74 recA1 araD139 Δ(ara-leu)7697 galU galK rpsL (Str ^R) endA1 nupG λ-	Invitrogen Life Technologies Catalog nos. C4040-10, C4040-03, C4040-06, C4040-50, and C4040-52
<i>E. coli</i> Origami™ 2	Δ(ara-leu)7697 ΔlacX74 ΔphoA PvUII phoR araD139 ahpC galE galK rpsL F'[lac ⁺ lacI ^q pro] gor522::Tn10 trxB (Str ^R , Tet ^R) fhuA2 [lon] ompT ahpC gal	Merck (EMD Millipore Chemicals) Catalog No. 71344
<i>E. coli</i> SHuffle® Express	λatt::NEB3-rlc-DsbC (Spec, lacI) AtrB sulA11 R(mcr-73::miniTn10-Tet ^S)2 [dcm] R(zgb-210::Tn10-Tet ^S) endA1 Δgor Δ(mcrC-mrr)14::IS10	New England Biolabs Catalog No. C3028H

[0076] Methods of Altering Host Cell Gene Functions.

[0077] There are many methods known in the art for making alterations to host cell genes in order to eliminate, reduce, or change gene function. Methods of making targeted disruptions of genes in host cells such as *E. coli* and other prokaryotes have been described (Muyrers et al., "Rapid modification of bacterial artificial chromosomes by ET recombination", Nucleic Acids Res 1999 Mar. 15; 27(6): 1555-1557; Datsenko and Wanner, "One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products", Proc Natl Acad Sci USA 2000 Jun. 6; 97(12): 6640-6645), and kits for using similar Red/ET recombination methods are commercially available (for example, the Quick & Easy *E. coli* Gene Deletion Kit from Gene Bridges GmbH, Heidelberg, Germany). In one embodiment of the invention, the function of one or more genes of host cells is eliminated or reduced by identifying a nucleotide sequence within the coding sequence of the gene to be disrupted, such as one of the *E. coli* K-12 substrain MG1655 coding sequences incorporated herein by reference to the genomic location of the sequence, and more specifically by selecting two adjacent stretches of 50 nucleotides each within that coding sequence. The Quick & Easy *E. coli* Gene Deletion Kit is then used according to the manufacturer's instructions to insert a polynucleotide construct containing a selectable marker between the selected adjacent stretches of coding sequence, eliminating or reducing the normal function of the gene. Red/ET recombination methods can also be used to replace a promoter sequence with that of a different promoter, such as a constitutive promoter, or an artificial promoter that is predicted to promote a certain level of transcription (De Mey et al., "Promoter knock-in: a novel

rational method for the fine tuning of genes", BMC Biotechnol 2010 Mar. 24; 10: 26). The function of host cell genes can also be eliminated or reduced by RNA silencing methods (Man et al., "Artificial trans-encoded small non-coding RNAs specifically silence the selected gene expression in bacteria", Nucleic Acids Res 2011 April; 39(8): e50, Epub 2011 Feb. 3). Further, known mutations that alter host cell gene function can be introduced into host cells through traditional genetic methods.

Inducible Coexpression Systems of the Invention

[0078] Inducible coexpression systems of the invention involve host cells comprising two or more expression constructs, where the expression constructs comprise inducible promoters directing the expression of gene products, and the host cells have altered gene functions that allow for homogeneous inducible expression of the gene products. FIG. 1 shows a schematic representation of an inducible coexpression system of the invention, with the following components: (1) host cell, (2) host genome (including genetic alterations), (3) an expression vector 'X' comprising an inducible promoter directing expression of a gene product, (4) a different expression vector 'Y' comprising an inducible promoter directing expression of another gene product, (5) chemical inducers of expression, and (6) the multimeric coexpression product. FIG. 3 shows a schematic representation similar to that shown in FIG. 1, with inducible promoters ((3) and (4)) and the coding sequences for the products expressed by the inducible promoters present on the same expression vector.

[0079] FIG. 2 shows a schematic representation of a particular example of an inducible coexpression system of the invention, utilizing the araBAD promoter on a pBAD24 (or pBAD240) expression vector in combination with a propionate-inducible promoter (prpBCDE promoter) on a pPRO33 (U.S. Pat. No. 8,178,338 B2; May 15, 2012; Keasling, Jay) (or pPRO43, pPRO430, pPRO430(CloDF13), pPRO44, or pPRO45) expression vector in an *E. coli* host cell housing the appropriate genomic alterations which allow for homogenously inducible expression. In this manner, tight control and optimization of expression of each component of a multimeric product can be achieved for use in a number of coexpression applications. In this embodiment, the host cell (1) is the Gram-negative bacterium *Escherichia coli*, commonly used in the art for protein expression. The host genome (2) is the genome of the host cell organism with mutations or other alterations that facilitate homogenously inducible protein coexpression, including expression of a cytoplasmic form of the disulfide isomerase DsbC which lacks a signal peptide. In one embodiment, the genomic alterations include both an araBAD operon knockout mutation, and either expression of araE and araFGH from constitutive promoters, or a point mutation in the lacY gene (A117C) in an araEFGH-deficient background, to facilitate homogenous induction of plasmid-based ara promoters with exogenously applied L-arabinose, and also an inactivated propionate metabolism gene, prpD, to facilitate homogenous induction of plasmid-based propionate promoters with exogenously applied propionate, which is converted to 2-methylcitrate in vivo.

[0080] Other genomic alterations that are useful for the inducible coexpression system, and may be introduced into the host cell, include without limitation: targeted inactivation of the scpA-argK-scpBC operon, to reduce background expression from the prpBCDE promoter; expression of the transcriptional regulator (prpR) for the less-preferred carbon-

source (propionate) from an L-arabinose-inducible promoter such as the araBAD promoter, and/or an eliminated or reduced gene function for genes involved in the CCR system, such as *crr* and/or *ptsG*, to avoid suppression by the CCR system of induction by propionate in the presence of L-arabinose; reductions in the level of gene function for glutathione reductase (*gor*) or glutathione synthetase (*gshB*), together with thioredoxin reductase (*trxB*), and/or expression of yeast mitochondrial sulphydryl oxidase *Erv1p* in the host cell cytoplasm, to provide a less strongly reducing environment in the host cell cytoplasm and promote disulfide bond formation; increased levels of expression, such as from a strong constitutive promoter, of chaperone proteins such as *DnaK/DnaJ/GrpE*, *DsbC/DsbG*, *GroEL/GroES*, *IbpA/IbpB*, *Skp*, *Tig* (trigger factor), and/or *FkpA*; and other mutations to reduce endogenous protease activity (such as that of the *Lon* and *OmpT* proteases) and recombinase activities.

[0081] As shown in FIG. 2, two compatible expression vectors (3, 4) are maintained in the host cell to allow for simultaneous expression (coexpression) of two different gene products. In this embodiment, one expression vector ('L-arabinose-induced expression vector') contains an L-arabinose-induced promoter, and is similar or identical to pBAD or related plasmids in which an araBAD promoter drives expression of an inserted expression sequence cloned into the multiple cloning site (MCS). The L-arabinose-induced expression vector also contains a coding sequence for an antibiotic-resistance gene (such as the *Tn3 bla* gene, which encodes beta-lactamase and confers resistance to ampicillin, or a gene encoding aminoglycoside 3'-phosphotransferase and conferring resistance to kanamycin) to facilitate selection of host cells (bacterial colonies) which contain an intact expression vector. An origin of replication (ORI) is required for propagation of the plasmid within bacterial host cells. The L-arabinose induced expression plasmid also contains a polynucleotide sequence encoding *araC*, a transcriptional regulator that allows for L-arabinose induction of the araBAD promoter and through transcriptional repression reduces 'leaky' background expression in the non-induced state. The other expression vector ('propionate-induced expression vector') is similar or identical to pPRO or related plasmids, in which a propionate-induced promoter drives expression of an inserted expression sequence cloned into the multiple cloning site (MCS). The plasmid also contains a coding sequence for an antibiotic-resistance gene (such as the *cat* gene, encoding chloramphenicol acetyltransferase, which confers resistance to chloramphenicol) to facilitate selection of host cells which contain an intact expression vector. An origin of replication (ORI) is required for propagation of the plasmid within bacterial host cells. In addition, the propionate-induced expression vector contains a polynucleotide sequence encoding *prpR*, a transcriptional regulator that allows for propionate (2-methylcitrate) induction of the *prpBCDE* promoter and reduces 'leaky' background expression in the non-induced state. To facilitate separate titration of induction, plasmid compatibility, and copropagation of the expression vectors, it is useful for the expression vectors to contain promoters responsive to different inducers, compatible origins of replication, and different antibiotic-resistance markers. In one embodiment of the invention, pBAD24 (pMB1 or 'pBR322' ORI, *Amp*^R) or a related expression vector such as pBAD240 (pMB1 ORI, *Kan*^R) containing an L-arabinose-inducible araBAD promoter is combined in a host cell with a pPRO33, pPRO43, pPRO430, or related expression vector (p15A ORI,

Cm^R) containing a propionate-inducible *prpBCDE* promoter. Compatible expression vectors containing a propionate-inducible *prpBCDE* promoter such as pPRO430(CloDF13), pPRO44 (RSF1030 ORI), or pPRO45 (CloDF13) can also be used in combination with the araBAD-promoter-containing expression vectors (pMB1 ORI). The expression vectors are copropagated and maintained using growth medium supplemented with the appropriate antibiotics: ampicillin, chloramphenicol, and/or kanamycin. In one embodiment, one expression vector comprises a polynucleotide sequence encoding the heavy chain of a full-length antibody, and the other expression vector comprises a polynucleotide sequence encoding the light chain of a full-length antibody, each coding sequence cloned in-frame into the MCS of the respective expression vector. For production of certain gene products such as antibodies, coding sequence optimization for the host organism (including adjustment for codon bias and GC-content, among other considerations) will determine the coding sequences to be inserted into the expression constructs of the coexpression system.

[0082] Referring again to FIG. 2, coexpression of gene products is induced by inexpensive exogenously applied chemical metabolites, L-arabinose and propionate (5). The level of induction of expression of each gene product is independently titrated with its own chemical inducer, thereby facilitating optimization of protein coexpression. This is useful for expression of protein complexes and proteins that require a binding partner for stabilization, and may facilitate expression of otherwise difficult to express proteins, such as those with poor solubility or cellular toxicity. In this example, upon induction, antibody heavy and short chains are each separately expressed, then the proteins join and form inter-chain disulfide bridges (within the cytoplasm of the bacterial host) which allows the formation and stabilization of full-length antibody comprised of the heavy and light chains. Proteins can be directed to various compartments of the host organism. For example, in *E. coli* the protein can be expressed in the cytoplasm, cell membrane, periplasm, or secreted into the medium. After an appropriate incubation time, cells and media are collected, and the total protein extracted, which includes the coexpressed gene products (6). After extraction, the desired product can be purified using a number of methods well known in the art depending on the nature of the gene products produced in the coexpression system (for example liquid chromatography). In the example shown in FIG. 2, the multimeric product (full-length antibody) is extracted and purified using chromatographic methods. Purified intact antibody is visualized on a non-denaturing gel using standard techniques, including protein-binding dyes or immunohistochemistry. The full-length antibody product can then be used for a number of research, diagnostic, or other applications.

[0083] FIG. 4 shows a schematic representation similar to that shown in FIG. 2, with an arabinose-inducible promoter (3), a propionate-inducible promoter (4), and coding sequences for antibody heavy and light chains present on the same expression vector. It is also possible to express the antibody heavy chain from the propionate-inducible promoter and the antibody light chain from the arabinose-inducible promoter.

Products Made by the Methods of the Invention

[0084] There is broad versatility in utilizing the inducible coexpression systems of the present invention in numerous coexpression applications, and in the properties of the products.

[0085] Glycosylation.

[0086] Gene products coexpressed by the methods of the invention may be glycosylated or unglycosylated. In one embodiment of the invention, the coexpressed gene products are polypeptides. Glycosylated polypeptides are polypeptides that comprise a covalently attached glycosyl group, and include polypeptides comprising all the glycosyl groups normally attached to particular residues of that polypeptide (fully glycosylated polypeptides), partially glycosylated polypeptides, polypeptides with glycosylation at one or more residues where glycosylation does not normally occur (altered glycosylation), and polypeptides glycosylated with at least one glycosyl group that differs in structure from the glycosyl group normally attached to one or more specified residues (modified glycosylation). An example of modified glycosylation is the production of “defucosylated” or “fucose-deficient” polypeptides, polypeptides lacking fucosyl moieties in the glycosyl groups attached to them, by expression of polypeptides in host cells lacking the ability to fucosylate polypeptides. Unglycosylated polypeptides are polypeptides that do not comprise a covalently bound glycosyl group. An unglycosylated polypeptide can be the result of deglycosylation of a polypeptide, or of production of an aglycosylated polypeptide. Deglycosylated polypeptides can be obtained by enzymatically deglycosylating glycosylated polypeptides, whereas aglycosylated polypeptides can be produced by expressing polypeptides in host cells that do not have the capability to glycosylate polypeptides, such as prokaryotic cells or cells in which the function of at least one glycosylation enzyme has been eliminated or reduced. In a particular embodiment, the coexpressed polypeptides are aglycosylated, and in a more specific embodiment, the aglycosylated polypeptides are coexpressed in prokaryotic cells such as *E. coli*.

[0087] Other Modifications of Gene Products.

[0088] Gene products coexpressed by the methods of the invention may be covalently linked to other types of molecules. Examples of molecules that may be covalently linked to coexpressed gene products, without limiting the scope of the invention, include polypeptides (such as receptors, ligands, cytokines, growth factors, polypeptide hormones, DNA-binding domains, protein interaction domains such as PDZ domains, kinase domains, antibodies, and fragments of any such polypeptides); water-soluble polymers (such as polyethylene glycol (PEG), carboxymethylcellulose, dextran, polyvinyl alcohol, polyoxyethylated polyols (such as glycerol), polyethylene glycol propionaldehyde, and similar compounds, derivatives, or mixtures thereof); and cytotoxic agents (such as chemotherapeutic agents, growth-inhibitory agents, toxins (such as enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof), and radioactive isotopes).

[0089] In addition, gene products to be coexpressed by the methods of the invention can be designed to include molecular moieties that aid in the purification and/or detection of the gene products. Many such moieties are known in the art; as one example, a polypeptide gene product can be designed to include a polyhistidine ‘tag’ sequence—a run of six or more histidines, preferably six to ten histidine residues, and most preferably six histidines—at its N- or C-terminus. The presence of a polyhistidine sequence on the end of a polypeptide allows it to be bound by cobalt- or nickel-based affinity media, and separated from other polypeptides. The polyhistidine tag sequence can be removed by exopeptidases. As

another example, fluorescent protein sequences can be expressed as part of a polypeptide gene product, with the amino acid sequence for the fluorescent protein preferably added at the N- or C-terminal end of the amino acid sequence of the polypeptide gene product. The resulting fusion protein fluoresces when exposed to light of certain wavelengths, allowing the presence of the fusion protein to be detected visually. A well-known fluorescent protein is the green fluorescent protein of *Aequorea victoria*, and many other fluorescent proteins are commercially available, along with nucleotide sequences encoding them.

[0090] Antibodies.

[0091] In one embodiment of the invention, the coexpressed gene products are antibodies. The term ‘antibody’ is used in the broadest sense and specifically includes ‘native’ antibodies, fully-human antibodies, humanized antibodies, chimeric antibodies, multispecific antibodies (such as bispecific antibodies), monoclonal antibodies, polyclonal antibodies, antibody fragments, and other polypeptides derived from antibodies that are capable of binding antigen. Unless indicated otherwise herein, the numbering of the residues in an immunoglobulin heavy chain (‘EU numbering’) is that of the EU index (the residue numbering of the human IgG1 EU antibody) as in Kabat et al., *Sequences of Proteins of Immunological Interest*, Fifth Edition, 1991, National Institute of Health, Bethesda, Md.

[0092] ‘Native’ antibodies are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of inter-chain disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at its N-terminal end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at its N-terminal end (V_L) and a constant domain at its C-terminal end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. The term ‘variable’ refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for an antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called hypervariable regions (HVRs) both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, connected by three HVRs, and with the HVRs from the other chain, contribute to the formation of the antigen-binding site of antibodies.

[0093] The term ‘Fc region’ refers to a C-terminal region of an immunoglobulin heavy chain, and includes native Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy-chain Fc region can be defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. Alternatively, the Fc region can be defined to extend from the N-terminal residue (Ala231) of the conserved C_H2 immunoglobulin domain to the C-terminus, and may include multiple con-

served domains such as C_{H2} , C_{H3} , and C_{H4} . The C-terminal lysine (residue 447 according to the EU numbering system) of the native Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. The Fc region of an antibody is crucial for recruitment of immunological cells and antibody dependent cytotoxicity (ADCC). In particular, the nature of the ADCC response elicited by antibodies depends on the interaction of the Fc region with receptors (FcRs) located on the surface of many cell types. Humans contain at least five different classes of Fc receptors. The binding of an antibody to FcRs determines its ability to recruit other immunological cells and the type of cell recruited. Hence, the ability to engineer antibodies with altered Fc regions that can recruit only certain kinds of cells can be critically important for therapy (US Patent Application 20090136936 A1, May 28, 2009, Georgiou, George). Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the CH2 domain of the Fc region. In certain embodiments, antibodies produced by the methods of the invention are not glycosylated or are aglycosylated, for example, due to a substitution at residue 297 of the Fc region, or to expression in a host cell that does not have the capability to glycosylate polypeptides. Due to altered ADCC responses, unglycosylated antibodies may stimulate a lower level of inflammatory responses such as neuroinflammation. Also, since an antibody having an aglycosylated Fc region has very low binding affinity for Fc receptors, such antibodies would not bind to the large number of immune cells that bear these receptors. This is a significant advantage since it reduces non-specific binding, and also increases the half-life of the antibody in vivo, making this attribute very beneficial in therapeutics.

[0094] The terms 'full-length antibody', 'intact antibody', and 'whole antibody' are used interchangeably to refer to an antibody in its substantially intact 'native' form, not antibody fragments as defined below. The terms particularly refer to an antibody with heavy chains that each comprise a variable domain and an Fc region. 'Antibody fragments' comprise a portion of an intact antibody, preferably comprising the antigen-binding region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, Fc, Fd, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules such as scFv; and multispecific antibodies formed from antibody fragments.

[0095] A 'human antibody' is one that possesses an amino-acid sequence corresponding to that of an antibody produced by a human. A 'chimeric' antibody is one in which a portion of the heavy and/or light chain is identical to, or shares a certain degree of amino acid sequence identity with, corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical to, or shares a certain degree of amino acid sequence identity with, corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies. A 'humanized' antibody is a chimeric antibody that contains minimal amino acid

residues derived from non-human immunoglobulin molecules. In one embodiment, a humanized antibody is a human immunoglobulin (recipient antibody) in which HVR residues of the recipient antibody are replaced by residues from an immunoglobulin HVR of a non-human species (donor antibody) such as mouse, rat, rabbit, or nonhuman primate. In some instances, FR residues of the human recipient antibody are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. The term 'monoclonal antibody' refers to an antibody obtained from a population of substantially homogeneous antibodies, in that the individual antibodies comprising the population are identical except for possible mutations, such as naturally occurring mutations, that may be present in minor amounts. Thus, the modifier 'monoclonal' indicates the character of the antibody as not being a mixture of discrete antibodies. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against the same single determinant on an antigen. In addition to their specificity, monoclonal antibody preparations are advantageous in that they are typically uncontaminated by other immunoglobulins.

[0096] The 'binding affinity' of a molecule such as an antibody generally refers to the strength of the sum total of non-covalent interactions between a single binding site of a molecule and its binding partner (such as an antibody and the antigen it binds). Unless indicated otherwise, 'binding affinity' refers to intrinsic binding affinity that reflects a 1:1 interaction between members of a binding pair (such as antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (Kd). Low-affinity antibodies (higher Kd) generally bind antigen slowly and tend to dissociate readily, whereas high-affinity antibodies (lower Kd) generally bind antigen faster and tend to remain bound longer. A variety of ways to measure binding affinity are known in the art, any of which can be used for purposes of the present invention. Specific illustrative methods for measuring binding affinity are described in Example 8. Antibodies and antibody fragments produced by and/or used in methods of the invention preferably have binding affinities of less than 100 nM, more preferably have binding affinities of less than 10 nM, and most preferably have binding affinities of less than 2 nM, as measured by a surface-plasmon resonance assay as described in Example 8.

[0097] Antibodies (Secondary) that Recognize Aglycosylated Antibodies.

[0098] Production of antibodies in *E. coli*-based or other prokaryotic expression systems without glycosylation enzymes will generally yield aglycosylated antibodies, which can be used as primary antibodies. In addition to using the inducible coexpression systems of the invention to produce aglycosylated primary antibodies, the inducible coexpression systems of the invention can also be used to efficiently produce secondary antibodies that specifically recognize aglycosylated primary antibodies. One aspect of the present invention is a secondary antibody system capable of detecting an unglycosylated or aglycosylated primary antibody for research, analytic, diagnostic, or therapeutic purposes. As one example, a secondary antibody system is provided with the following components: epitope, primary antibody, secondary antibody, and detection system. The epitope is a portion of an

antigen (usually a protein) which is the antigenic determinant that produces an immunological response when introduced into a live animal or is otherwise recognizable by an antibody. In practice, the epitope of interest may be present within a mixture or a tissue. In one embodiment, the epitope is a protein expressed in carcinoma cells in human tissue. The primary antibody is an antibody fragment, a single full-length antibody (monoclonal), or a mixture of different full-length antibodies (polyclonal), which recognizes and binds to the epitope, and preferably binds specifically to the epitope. A full-length antibody in this example comprises two heavy polypeptide chains and two light polypeptide chains joined by disulfide bridges. Each of the chains comprises a constant region (Fc) and a variable region (Fv). There are two antigen binding sites in the full-length antibody. In one embodiment of the present invention, the primary antibody is a full-length aglycosylated antibody (such as that produced in an *E. coli*-based expression system) which recognizes and binds an epitope of interest. The secondary antibody is an antibody fragment, a single full-length antibody (monoclonal), or a mixture of different full-length antibodies (polyclonal), which recognizes and binds to the aglycosylated primary antibody, and preferably binds specifically to the aglycosylated primary antibody. In one embodiment of the present invention, the secondary antibody is a full-length antibody which recognizes and binds the aglycosylated Fc portion of a full-length primary antibody. In this case, the antibody binding sites are selected and/or engineered to specifically recognize the Fc portion of the aglycosylated primary antibody, with or without the C-terminal lysine residue. In other embodiments, the secondary antibody could be engineered to recognize additional regions (epitopes) of the aglycosylated primary antibody, or additional engineered epitopes including but not limited to polypeptide sequences covalently attached to the primary antibody. The secondary antibody can be directed at single or multiple sites (epitopes) present on full-length aglycosylated antibodies molecules (including various immunoglobulin classes such as IgG, IgA, etc.) or antibody fragments such as Fc or Fab. Therefore, some secondary antibodies generated in this way would have broad specificity for any aglycosylated full-length antibody. The primary and secondary antibodies of the present invention can also include those produced by traditional methods (polyclonal antibody production using immunized rabbits or monoclonal antibody production using mouse hybridomas) and recombinant DNA technology such as phage display methods for identifying antigen-binding polypeptides.

[0099] Detection systems generally comprise an agent that is linked to or which binds the secondary antibody, enabling detection, visualization, and/or quantification of the secondary antibody. Various detection systems are well known in the art including but not limited to fluorescent dyes, enzymes, radioactive isotopes, or heavy metals. These may or may not involve direct physical linkage of additional polypeptides to the secondary antibody. Applications of this secondary antibody system include but are not limited to immunohistochemistry, Western blotting, and enzyme-linked immunosorbent assay (ELISA). For example, in one embodiment for use in immunohistochemistry, the epitope of interest would be present on a thin section of tissue, then an aglycosylated primary antibody would be applied to the tissue and allowed to bind the epitope. The unbound primary antibody would be removed, and then a secondary antibody capable of specifically binding the aglycosylated primary antibody is applied to

the tissue and allowed to bind to the primary antibody. The unbound secondary antibody would be removed, and then detection system reagents applied. For example, if the secondary antibody were linked to an enzyme, then colorogenic enzymatic substrates would be applied to the tissue and allowed to react. Direct microscopic or fluoroscopic visualization of the reactive enzymatic substrates could then be performed. Other detection methods are well known in the art. The advantages of a system using secondary antibodies that recognize aglycosylated antibodies include, without limitation, the following: 1) increased specificity in immunohistochemistry because the secondary antibody is designed to bind the aglycosylated Fc portion of the primary antibody which is not otherwise present in eukaryotic tissues; 2) decreased background staining because of increased specificity for the primary antibody; 3) decreased cost of secondary antibody system production because the primary and/or secondary antibodies can be generated in prokaryotes such as *E. coli*; and 4) avoiding unnecessary utilization of mammals, including mice and rabbits, because the entire process of antibody development can be performed in prokaryotes such as *E. coli*.

[0100] Enzymes Used in Industrial Applications.

[0101] Many industrial processes utilize enzymes that can be produced by the methods of the invention. These processes include treatment of wastewater and other bioremediation and/or detoxification processes; bleaching of materials in the paper and textile industries; and degradation of biomass into material that can be fermented efficiently into biofuels. In many instances it would be desirable to produce enzymes for these applications in microbial host cells or preferably in bacterial host cells, but the active enzyme is difficult to express in large quantities due to problems with enzyme folding and/or a requirement for a cofactor. In certain embodiments of the invention, the inducible coexpression methods of the invention are used to produce enzymes with industrial applications, such as arabinose- and xylose-utilization enzymes (e.g. xylose isomerase (EC 5.3.1.5)) or lignin-degrading peroxidases (e.g. lignin peroxidase (EC 1.11.1.14), manganese peroxidase (EC 1.11.1.13), versatile peroxidase (EC 1.11.1.16), or laccase (EC 1.10.3.2)).

Example 1

Introduction of Genomic Alterations into Host Cells to Facilitate Coexpression

[0102] As described above, certain changes in host cell gene expression can improve the coexpression of the desired gene product(s). Certain host cells, *E. coli* ASE(DGH) cells, were derived from *E. coli* SHuffle® Express cells, and their genotype can be expressed as: *E. coli* SHuffle® Express ΔaraBAD Asbm-ygfLΔGH ΔaraEp::J23104. The *E. coli* ASE (DGH) cells were produced as follows: deletions and alterations were made in the *E. coli* SHuffle® Express host cell genome by Gene Bridges GmbH (Heidelberg, Germany) using a recombineering method, described as deletion by counterselection, that seamlessly removes genomic sequences. A deletion of the host cell araBAD operon was made to reduce arabinose catabolism by the host cell, so that more of the arabinose inducer will be available for induction of a coexpressed gene product from an expression construct comprising the araBAD promoter. This deletion removes 4269 basepairs of the araBAD operon, corresponding to position 70,135 through 65,867 (minus strand) of the *E. coli*

genome (positions within genomic nucleotide sequences are all given as in Table 1), so that most of the native araBAD promoter through all but a few codons of the AraD coding region are removed. The nucleotide sequence (minus strand) around the deletion junction (position 70,136/position 65,866) is: TTAT[TACG. Another deletion was made within the sbm-ygfDGH (also called scpA-argK-scpBC) operon, eliminating the function of genes involved in the biosynthesis of 2-methylcitrate, to increase sensitivity of the host cell's propionate-inducible promoter to exogenously supplied propionate. The sbm-ygfDGH deletion removes 5542 basepairs (position 3,058,754 through 3,064,295 of the *E. coli* genome), taking out the sbm-ygfDGH promoter and all of the operon except for the last codon of the ygfH coding sequence, while leaving the adjacent ygfI coding sequence and stop codon intact. The nucleotide sequence (plus strand) around the deletion junction (position 3,058,753/position 3,064,296) is: ACAA[GGGT. In addition to these deletions made in the *E. coli* SHuffle® Express host cell genome, Gene Bridges GmbH introduced a point mutation in the genomic rpsL gene coding sequence, which extends on the minus strand from position 3,472,574 through 3,472,200, changing the A at position 3,472,447 to a G, altering the codon for Lys43 to a codon for Arg, which results in a streptomycin-resistant phenotype when the mutant rpsL-Arg43 gene is expressed. Another alteration to the host cell genome, allowing for more tightly controlled inducible expression as described above, is to make the araE promoter constitutive rather than responsive to arabinose. Most of the native araE promoter, including CRP-cAMP and AraC binding sites, was removed by deleting 97 basepairs (position 2,980,335 through 2,980,239 (minus strand)) and replacing that sequence with the 35-basepair sequence of the constitutive J23104 promoter (SEQ ID NO:1; the nucleotide sequence of J23104 was obtained from the partsregistry.org website, parts.igem.org/Main_Page). The resulting junction site sequences within the altered araE promoter are: TGAA[TTGA . . . TAGCITTC. An *E. coli* host cell, such as an *E. coli* SHuffle® Express host cell, with any of these genomic alterations, or any combination of them, can be employed in the inducible coexpression of gene products.

Example 2

Expression Vectors Comprising an Inducible Promoter

[0103] A. Expression Vector pPRO43

[0104] The expression vector pPRO43 (SEQ ID NO:2) is used for expressing gene products of interest from the propionate-inducible prpBCDE promoter, and was constructed with reference to the nucleotide sequence of the pPRO33 expression vector. The nucleotide sequence of pPRO33 was compiled from the sequences of the pBAD18 vector (GenBank Accession No. X81838.1), the *E. coli* genomic sequence of the prpR-P_{prpB} region, and the pBAD33 vector, as described in Guzman et al., "Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter", J Bacteriol 1995 July; 177(14): 4121-4130, and in U.S. Pat. No. 8,178,338 B2; May 15, 2012; Keasling, Jay. The nucleotide sequence of pPRO33 was confirmed by sequencing and is provided in SEQ ID NO:3.

[0105] In pPRO43 the nucleotide sequence encoding the transcriptional activator prpR has been optimized for expression in *E. coli* by DNA2.0 (Menlo Park, Calif.) using methods such as those described in Welch et al., "Design parameters to

control synthetic gene expression in *Escherichia coli*", PLoS One 2009 Sep. 14; 4(9): e7002; doi: 10.1371/journal.pone.0007002. The optimized prpR sequences in pPRO43 include the RBS and other sequences upstream of the prpR coding sequence, which is nucleotides 1593 through 7 of SEQ ID NO:2, on the opposite strand from that shown. The pPRO43 vector also has only one HindIII restriction site, which is in the multiple cloning site (MCS), in contrast with pPRO33 which has two HindIII sites, one in the MCS and a second in the prpR coding sequence.

B. Expression Vectors pPRO430 and pPRO430(CloDF13)

[0106] Expression vector pPRO430 (SEQ ID NO:4) was synthesized by DNA2.0, based on the nucleotide sequence of pPRO43 (SEQ ID NO:2). The pPRO430 vector is similar to pPRO43 in that both contain the p15 origin of replication, a gene conferring resistance to chloramphenical (Cm^R), the prpR coding sequence optimized for expression in *E. coli* described above, and a cloning site downstream of the propionate-inducible prpBCDE promoter into which a coding sequence can be inserted. The pPRO430 expression vector differs from pPRO43 in that it has an optimized RBS sequence—AGGAGGAAACATA (nucleotides 3566-3579 of SEQ ID NO:4)—upstream of the cloning site. The pPRO430 expression vector has also been streamlined relative to pPRO43 by the removal of some nucleotide sequences and the use of shorter terminators, with the result that the pPRO430 nucleotide sequence (SEQ ID NO:4) has only 3698 bases, compared to the pPRO43 nucleotide sequence (SEQ ID NO:2) at 5883 bases. The pPRO430(CloDF13) expression vector (SEQ ID NO:5) is identical to pPRO430 except that the p15 origin of replication in pPRO430, flanked by BfuAI restriction sites, is replaced by the higher-copy-number CloDF13 origin of replication in pPRO430(CloDF13).

C. Expression Vector pBAD240

[0107] The expression vector pBAD240 (SEQ ID NO:6) was synthesized by DNA2.0, based on the nucleotide sequence of pBAD24 (GenBank Database Accession No. X81837.1 (25 Oct. 1995)). The expression vector pBAD240 differs from pBAD24 in having an optimized RBS sequence—AGGAGGTAAAAA (nucleotides 3125-3136 of SEQ ID NO:6)—upstream of the cloning site into which a coding sequence can be inserted. The pBAD240 nucleotide sequence was also streamlined relative to pBAD24 by the removal of some nucleotide sequences and the use of shorter terminators, with the result that the pBAD240 nucleotide sequence (SEQ ID NO:6) has only 3255 bases, compared to the pBAD24 nucleotide sequence at 4542 bases. The pBAD240 expression vector also includes a gene conferring resistance to kanamycin (Kan^R), rather than the ampicillin-resistance gene that is in pBAD24.

D. Expression Vectors pPRO44 and pPRO45

[0108] Expression vectors pPRO44 (SEQ ID NO:7) and pPRO45 (SEQ ID NO:8) were created based on the nucleotide sequence of pPRO43 (SEQ ID NO:2), with different origins of replication to be used in the pPRO44 (RSF1030) and pPRO45 (CloDF13) expression vectors. Primers were designed for pPRO43, the RSF1030 origin of replication, and the CloDF13 origin of replication, in order to add a SpeI restriction site upstream and an AatII restriction site downstream of each origin of replication sequence. The pPRO43 and RSF1030 nucleotide sequences were amplified using those primers (SEQ ID Nos 9 and 10, and SEQ ID Nos 11 and 12, respectively), the amplified products were digested with SpeI and AatII, and the desired restriction fragments were gel

purified and ligated together to create pPRO44 (SEQ ID NO:7). The resulting pPRO44 expression vector was sequenced to confirm that no mutations were introduced through PCR amplification. To create pPRO45, the CloDF13 nucleotide sequence was amplified using primers (SEQ ID Nos 13 and 14) to add SpeI and AatII sites, the CloDF13 amplification product and pPRO44 were digested with SpeI and AatII restriction enzymes and the desired fragments were gel purified, and then ligated together to form pPRO45 (SEQ ID NO:8). The CloDF13 portion of pPRO45 was sequenced to confirm that no mutations were introduced through PCR amplification.

Example 3

Use of Dual-Promoter Vector pSOL for Inducible Coexpression of Fluorescent Proteins in Bacterial Cells

[0109] A. Construction of the Dual-Promoter pSOL Expression Vector.

[0110] An expression vector comprising two different inducible promoters, referred to as 'dual vector' or 'pSOL', is shown schematically in FIG. 5. This vector was synthesized by DNA2.0 (Menlo Park, Calif.) and contains several polynucleotide sequences, or 'elements', optimized for expression in *E. coli* host cells. A description of the polynucleotide elements utilized in pSOL is provided below as Table 4; the nucleotide sequence of pSOL is provided as SEQ ID NO:15.

TABLE 4

Polynucleotide Elements of pSOL Dual-Promoter Expression Vector		
Name of element:	Position of element in SEQ ID NO: 15 (or its complement):	Description of element:
BfuAI	complement (29-38)	BfuAI restriction site
Ori_pBR	39-651	pBR322 (pMB1) origin of replication
ROP	complement (703-894)	Coding sequence: <i>E. coli</i> regulatory protein rop (RNA one modulator rom)
BfuAI	895-900	BfuAI restriction site
Kan-R	complement (901-1710)	Coding sequence: kanamycin resistance protein
P_kan	complement (1711-1832)	Promoter for expression of kanamycin resistance protein
term.	complement	Terminator apFAB389 (BIOFAB, Emeryville, California)
apFAB389	(1833-1923)	
BsmBI	complement (1924-1929)	BsmBI restriction site
araC	complement (1930-2808)	Coding sequence: AraC transcriptional regulator protein
MKS	complement (2809-2817)	Coding sequence: MKS (methionine-lysine-serine)
rbs	complement (2818-2840)	Ribosome binding site
araC	complement (2959-2987)	Promoter for expression of AraC
promoter		
araBAD	3084-3111	Promoter from araBAD operon for expression of inserted coding sequences
promoter		
NheI	3134-3139	NheI restriction site
optimized	3140-3151	Optimized ribosome binding site (as in pBAD240)
RBS		
BsaI	complement (3158-3163)	BsaI restriction site
BsaI	3164-3169	BsaI restriction site
XbaI	3172-3177	XbaI restriction site

TABLE 4-continued

Polynucleotide Elements of pSOL Dual-Promoter Expression Vector		
Name of element:	Position of element in SEQ ID NO: 15 (or its complement):	Description of element:
term.	3178-3259	Terminator apFAB391 (BIOFAB, Emeryville, California)
apFAB391		
term.	complement (3260-3349)	Terminator apFAB391 (BIOFAB, Emeryville, California)
apFAB381		
opt. prpR	complement (3350-4936)	Optimized coding sequence: PrpR transcriptional regulator protein
prpB	5114-5129	Promoter from prpBCDE operon for expression of inserted coding sequences
promoter		
HindIII	5166-5171	HindIII restriction site
rbs	5172-5185	Optimized ribosome binding site (as in pPRO430)
BsaI	complement (5186-5191)	BsaI restriction site
BsaI	5192-5197	BsaI restriction site
Xho	5199-5204	XhoI restriction site
3 frame	5205-5215	Stop codons in all three forward reading frames
stop		
term.	5216-5304	Terminator apFAB390 (BIOFAB, Emeryville, California)
apFAB390		

[0111] In particular, the pSOL expression vector comprises two different inducible promoters that can be used for the inducible coexpression of proteins of interest: an arabinose-inducible araBAD promoter and a propionate-inducible prpBCDE promoter. Variants of pSOL can also be used for the inducible coexpression of proteins of interest, such as an expression vector based on pSOL in which the positions of the araBAD promoter and the prpBCDE promoter are switched relative to the origin of replication. Further useful variants of pSOL expression vectors include those in which the coding sequence for the AraC transcriptional activator and/or the coding sequence for the PrpR transcriptional activator are not present in the expression vector, but are instead expressed from a separate polynucleotide such as a different extrachromosomal element, or the host genome. Additional variants of pSOL expression vectors are those comprising a third inducible promoter inserted, for example, between pSOL nucleotide positions 5304 and 29 of SEQ ID NO:15, so that the third inducible promoter would be downstream of the prpB promoter and its associated cloning sites and terminator, and oriented in the same direction as the prpB promoter. The third inducible promoter in such variants of pSOL expression vectors could for example be a rhamnose-inducible promoter such as the rhaSR promoter, a xylose-inducible promoter such as the xylAB promoter, or a promoter inducible by phosphate depletion such as the phoA promoter.

B. Inducible Coexpression of Fluorescent Proteins.

[0112] To compare levels of coexpression from the dual-promoter pSOL vector with coexpression from a combination of pBAD24 and pPRO33 expression vectors, the pBAD24 and pPRO33 vectors were used to express a yellow fluorescent protein (YellowFP) and a red fluorescent protein (RedFP), respectively. The L-arabinose-inducible araBAD promoter in pSOL was used to express YellowFP, and the propionate-inducible prpBCDE promoter was used to express RedFP. The YellowFP has maximal emission at 528 nm-530 nm with excitation at 515 nm (Nagai et al., "A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications", Nat Biotechnol 2002

January; 20(1): 87-90), and the RedFP has maximal emission at 610 nm with excitation at 587 nm (Shaner et al., "Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein", Nat Biotechnol 2004 December; 22(12): 1567-1572; Epub 2004 Nov. 21).

[0113] To construct the pBAD24-YellowFP and pPRO33-RedFP plasmids, the coding sequences for YellowFP and RedFP were optimized for expression in *E. coli* by DNA 2.0. The optimized coding sequences for YellowFP and RedFP were digested with both NheI and SalI, and the digested inserts ligated into NheI/SalI-cut pBAD24 and pPRO33, respectively. In the case of pSOL, DNA2.0 used the amplification-based Electra cloning system to insert the optimized coding sequences for YellowFP and for RedFP into the pSOL expression vector downstream of the optimized ribosome binding sites in the araBAD promoter and prpBCDE promoter regions, respectively. The pBAD24-YellowFP and pPRO33-RedFP expression constructs were co-transformed into *E. coli* ASE(DGH) cells (*E. coli* ASE(DGH) cells are described in Example 1), and pSOL-YellowFP-RedFP expression constructs were also separately transformed into *E. coli* ASE(DGH) cells.

[0114] Cultures of pBAD24-YellowFP/pPRO33-RedFP in ASE(DGH) ('pBAD, pPRO') and pSOL-YellowFP-RedFP in ASE(DGH) ('pSOL') were grown overnight at 37 degrees C. with shaking at 275 RPM in LB medium containing chloramphenicol plus ampicillin or containing kanamycin, respectively. At an OD600 of 0.7, the cells were diluted in 2×6 mL of LB medium plus antibiotics to an OD600 of 0.01 and grown at 30 degrees C. with shaking at 275 RPM until reaching an OD600 of 0.75. Cells were pelleted at 3800×g for 7 minutes at 30 degrees C., and were resuspended in M9 medium plus antibiotics with no additional carbon source to an OD600 of 0.7. The resuspended cells were plated in a multi-well plate at 200 microliters per well as follows:

Rows A, B: pBAD24-YellowFP/pPRO33-RedFP in ASE(DGH) colony 1

Rows C, D: pBAD24-YellowFP/pPRO33-RedFP in ASE(DGH) colony 2

Rows E, F: pSOL-YellowFP-RedFP in ASE(DGH) colony 1

Row G, H: pSOL-YellowFP-RedFP in ASE(DGH) colony 2

[0115] Cells were induced by the addition of varying concentrations of propionate (by hand) and L-arabinose (by digital dispenser), with propionate concentrations of 0 mM, 1 mM, 2.5 mM, 5 mM, and 20 mM and L-arabinose concentrations of 0 micromolar, 0.67 micromolar, 3.33 micromolar, 6.66 micromolar, and 66.61 micromolar.

[0116] The plates were incubated in a Biotek Synergy™ 4 microplate reader (BioTek Instruments Inc., Winooski, Vt.) at 30 degrees C. using the "Fast" shake speed, and fluorescence was monitored every 15 minutes for 8 hours. Fluorescence values were averaged for samples cultured in the same inducer concentrations, and the fluorescence plotted over time as shown in FIGS. 6 and 7. FIG. 6 shows the levels of fluorescence from RedFP over time, expressed from a propionate-inducible prpBCDE promoter. The inducer concentrations for each averaged group of samples are shown in the legend, with the propionate concentration listed first, followed by the L-arabinose concentration. FIG. 7 shows the levels of fluorescence from YellowFP over time, expressed from an L-arabinose-inducible araBAD promoter. The inducer concentrations for each averaged group of samples are shown in the legend, with the L-arabinose concentration

listed first, followed by the propionate concentration. For each combination of inducer concentrations, the expression of the fluorescent protein being measured, as indicated by the level of fluorescence observed, was higher from the inducible promoter present on the pSOL dual-promoter expression vector than from the corresponding inducible promoter present in the pBAD24-YellowFP/pPRO33-RedFP combination. This increased level of fluorescent protein expression from pSOL relative to pBAD24 and pPRO33 was observed not only for the combinations of inducer concentrations shown in FIGS. 6 and 7, but also for the combinations of inducer concentrations that were omitted from the graphs for the sake of clarity.

Example 4

Inducible Coexpression of Infliximab

[0117] Infliximab is a chimeric monoclonal antibody that binds to TNF-alpha, an inflammatory cytokine, and is used in the treatment of conditions that involve TNF-alpha such as autoimmune diseases (Crohn's disease, rheumatoid arthritis, psoriasis, etc.). Infliximab is formed from a heavy chain (amino acid sequence shown as SEQ ID NO:16) and a light chain (amino acid sequence shown as SEQ ID NO:17); each of these chains has a variable domain sequence derived from mouse anti-TNF-alpha antibodies, and a human constant domain. Codon optimization for expression in *E. coli* and synthesis of polynucleotides encoding SEQ ID NOs 16 and 17 was performed by DNA2.0 (Menlo Park, Calif.).

[0118] The DNA2.0 Electra cloning method (www.dna20.com/products/expression-vectors/electra-system) is used to create the infliximab expression constructs. The expression construct formed by inserting the optimized coding sequence for the infliximab heavy chain into the Electra cloning site of the pBAD240 expression vector is pBAD240-Infliximab_HC, which has the nucleotide sequence shown as SEQ ID NO:18. The expression construct formed by inserting the optimized coding sequence for the infliximab light chain into the Electra cloning site of the pPRO430 expression vector is pPRO430-Infliximab_LC, which has the nucleotide sequence shown as SEQ ID NO:19. Both the optimized infliximab heavy and light chain coding sequences were cloned into the pSOL expression vector (SEQ ID NO:15) in a similar way, with the heavy chain expressed from the araBAD promoter, and the light chain from the prpBCDE promoter. The resulting pSOL-Infliximab expression vector has the nucleotide sequence shown in SEQ ID NO:20. The pBAD240-Infliximab_HC and pPRO430-Infliximab_LC expression constructs are used to cotransform *E. coli* ASE (DGH) cells through heat shock at 42 degrees C., followed by growth at 37 degrees C. overnight, and the pSOL-Infliximab expression construct is similarly transformed into *E. coli* ASE(DGH) cells, creating ASE(DGH)(pBAD240-Infliximab_HC/pPRO430-Infliximab_LC) cells and ASE(DGH) (pSOL-Infliximab) cells.

[0119] These cells are grown generally as described in Example 3, including the addition of selective compounds such as kanamycin and/or chloramphenicol as needed, and for induction of antibody expression cells are resuspended in M9 medium with no additional carbon source at an OD600 of approximately 0.7 (0.6-1.0). The cells are then induced by addition of arabinose (initially at concentrations including 0.1%) and propionate (initially at concentrations including 50 mM). Adjustment of the concentrations of arabinose and propionate can be made as described in Example 5. After

induction, the host cells in which antibodies have been produced are disrupted by chemical lysis using enzymes such as lysozyme, or by mechanical disruption methods such as sonication or microfluidization using a Microfluidics model M-110Y microfluidizer (Microfluidics International Corp., Westwood, Mass.). Centrifugation at 20,000 \times g for 15 minutes at room temperature is used to separate out the insoluble fraction, and the supernatant containing soluble protein including the expressed antibodies is collected.

[0120] The infliximab antibodies are detected and quantified using a capillary electrophoresis Western blot, run on a WES system (ProteinSimple, San Jose, Calif.), according to the manufacturer's instructions. Soluble protein extracts are loaded into the capillary set, proteins are electrophoretically separated by size, and then the antibodies in the samples are detected with a blocking step (instead of the use of a primary antibody), and incubation with an HRP-conjugated goat anti-human secondary antibody that recognizes human antibody heavy and light chains. Antibody detection is accomplished by addition of the chemiluminescent substrate to the capillary and the direct capture of the light emitted during the enzyme-catalyzed reaction. Molecular weight estimates are calculated using a standard curve generated using six biotinylated proteins ranging from 12 k to 230 kDa for each run. Fluorescent standards are included in the sample loading buffer, giving each sample an internal standard that is used to align the sample with the molecular weight standard.

[0121] To determine the amount of protein present at a given molecular weight, known amounts of a protein standard are run in some of the capillaries. In this case, serial dilutions are prepared of commercially available infliximab having a known protein concentration, starting for example at 10 micrograms/mL and diluted down to 1.0 nanograms/mL. Approximately five WES system capillaries are used to run the serial dilution. For each infliximab protein band in both the experimental and the serial dilution capillaries, a curve is generated by the WES system software representing the protein band's chemiluminescence intensity, and the area under each curve is evaluated, with a standard curve of these areas plotted for the infliximab protein bands in the infliximab serial dilution capillaries. To determine the concentration of the experimental samples, the area under each curve representing the chemiluminescence intensity of an experimental infliximab sample can be compared to the standard curve generated for the samples of known infliximab concentration.

[0122] The infliximab antibodies can be further purified as described in Example 7, and additional characterization of the infliximab antibodies is described in Example 8 (measurement of antibody binding affinity) and Example 9 (characterizing the disulfide bonds present in coexpression products).

Example 5

Titration of Coexpression by Varying Inducer Concentration

[0123] To optimize production of a multimeric product using the inducible coexpression systems of the invention, it is possible to independently adjust or titrate the concentrations of the inducers. Host cells containing L-arabinose-inducible and propionate-inducible expression constructs are grown to the desired density (such as an OD₆₀₀ of approximately 0.5) in M9 minimal medium containing the appropriate antibiotics, then cells are aliquoted into small volumes of M9 minimal medium, optionally prepared with no carbon

source such as glycerol, and with the appropriate antibiotics and varying concentrations of each inducer. The concentration of L-arabinose necessary to induce expression is typically less than 2%. In a titration experiment, the tested concentrations of L-arabinose can range from 2% to 1.5%, 1%, 0.5%, 0.2%, 0.1%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.005%, 0.002%, and 0.001%. The concentrations of L-rhamnose or D-xylose necessary to induce expression of L-rhamnose-inducible or D-xylose-inducible promoters are tested similarly, with the tested concentrations ranging from 5% to 0.01%. For each concentration 'x' of L-arabinose (or L-rhamnose or D-xylose) that is tested, the concentration of a different inducer such as propionate, added to each of the tubes containing concentration 'x' of the first inducer, is varied in each series of samples, which in the case of propionate range from 1 M to 750 mM, 500 mM, 250 mM, 100 mM, 75 mM, 50 mM, 25 mM, 10 mM, 5 mM, and 1 mM. Alternatively, titration experiments can start at a 'standard' combination of inducer concentrations, which is 0.002% of any of L-arabinose, L-rhamnose, or D-xylose, and/or 50 mM propionate, and test new combinations of inducer concentrations that vary from that of the 'standard' combination. Similar titration experiments can be performed with any combination of inducers used in an inducible coexpression system of the invention, including but not limited to L-arabinose, propionate, L-rhamnose, and D-xylose. After growth in the presence of inducers for 6 hours, the cells are pelleted, the desired product is extracted from the cells, and the yield of product per mass value of cells is determined by a quantitative immunological assay such as ELISA, or by purification of the product and quantification by UV absorbance at 280 nm.

[0124] It is also possible to titrate inducer concentrations using a high-throughput assay, in which the proteins to be expressed are engineered to include a fluorescent protein moiety, such as that provided by the mKate2 red fluorescent protein (Evrogen, Moscow, Russia), or the enhanced green fluorescent proteins from *Aequorea victoria* and *Bacillus cereus*. Another approach to determining the amount and activity of gene products produced by different concentrations of inducers in a high-throughput titration experiment, is to use a sensor capable of measuring biomolecular binding interactions, such as a sensor that detects surface plasmon resonance, or a sensor that employs bio-layer interferometry (BLI) (for example, an Octet® QK system from forteBIO, Menlo Park, Calif.).

Example 6

Measurement of the Strength of Promoters and the Homogeneity of Inducible Expression

[0125] The strength of a promoter is measured as the amount of transcription of a gene product initiated at that promoter, relative to a suitable control. For constitutive promoters directing expression of a gene product in an expression construct, a suitable control could use the same expression construct, except that the 'wild-type' version of the promoter, or a promoter from a 'housekeeping' gene, is used in place of the promoter to be tested. For inducible promoters, expression of the gene product from the promoter can be compared under inducing and non-inducing conditions.

A. Measuring Promoter Strength Using Quantitative PCR to Determine Levels of RNA Transcribed from the Promoter

[0126] The method of De Mey et al. ("Promoter knock-in: a novel rational method for the fine tuning of genes", BMC

Biotechnol 2010 Mar. 24; 10: 26) is used to determine the relative strength of promoters in host cells that can be grown in culture. Host cells containing an expression construct with the promoter to be tested, and control host cells containing a control expression construct, are grown in culture in triplicate. One-ml samples are collected at $OD_{600}=1.0$ for mRNA and protein collection. Total RNA extraction is done using an RNeasy mini kit (QIAGEN, The Netherlands). The purity of RNA is verified on a FA-agarose gel as recommended by QIAGEN and the RNA concentration is determined by measuring the absorbance at 260 nm. Two micrograms of RNA is used to synthesize cDNA using a random primer and RevertAid H Minus M-MuLV reverse transcriptase (Fermentas, Glen Burnie, Md.). The strength of the promoter is determined by RT-qPCR carried out in an iCycler IQ® (Bio-Rad, Eke, Belgium) using forward and reverse primers designed to amplify the cDNA corresponding to the transcript produced from the promoter. (For this purpose, the De Mey et al. authors used the Fw-ppc-qPCR and Rv-ppc-qPCR primers, and the primers Fw-rpoB-qPCR and Rv-rpoB-qPCR from the control housekeeping gene rpoB.) SYBR GreenER qPCR supermix (Life Technologies, Grand Island, N.Y.) is used to perform a brief UDG (uracil DNA glycosylase) incubation (50° C. for 2 min) immediately followed by PCR amplification (95° C. for 8.5 min; 40 cycles of 95° C. for 15 s and 60° C. for 1 min) and melting curve analysis (95° C. for 1 min, 55° C. for 1 min and 80 cycles of 55° C.+0.5° C./cycles for 10 s) to identify the presence of primer dimers and analyze the specificity of the reaction. This UDG incubation step before PCR cycling destroys any contaminating dU-containing products from previous reactions. UDG is then inactivated by the high temperatures during normal PCR cycling, thereby allowing the amplification of genuine target sequences. Each sample is performed in triplicate. The relative expression ratios are calculated using the "Delta-delta ct method" of PE Applied Biosystems (PerkinElmer, Forster City, Calif.).

B. Measuring Inducible Promoter Strength and Homogeneity of Induction Using a Fluorescent Reporter Gene

[0127] These experiments are performed using the methods of Khlebnikov et al. ("Regulatable arabinose-inducible gene expression system with consistent control in all cells of a culture", J Bacteriol 2000 December; 182(24): 7029-7034). Experiments measuring the induction of an inducible promoter are performed in C medium supplemented with 3.4% glycerol as a carbon source (Helmstetter, "DNA synthesis during the division cycle of rapidly growing *Escherichia coli* B/r", J Mol Biol 1968 Feb. 14; 31(3): 507-518). *E. coli* strains containing expression constructs comprising at least one inducible promoter controlling expression of a fluorescent reporter gene are grown at 37° C. under antibiotic selection to an optical density at 600 nm (OD600) of 0.6 to 0.8. Cells are collected by centrifugation (15,000×g), washed in C medium without a carbon source, resuspended in fresh C medium containing antibiotics, glycerol, and/or inducer (for the induction of gene expression) to an OD600 of 0.1 to 0.2, and incubated for 6 h. Samples are taken routinely during the growth period for analysis. Culture fluorescence is measured on a Versafluor Fluorometer (Bio-Rad Inc., Hercules, Calif.) with 360/40-nm-wavelength excitation and 520/10-nm-wavelength emission filters. The strength of expression from an inducible promoter upon induction can be expressed as the ratio of the maximum population-averaged fluorescence (fluorescence/OD ratio) of the induced cells relative to that of

control (such as uninduced) cells. To determine the homogeneity of induction within the cell population, flow cytometry is performed on a Beckman-Coulter EPICS XL flow cytometer (Beckman Instruments Inc., Palo Alto, Calif.) equipped with an argon laser (emission at a wavelength of 488 nm and 15 mW) and a 525-nm-wavelength band pass filter. Prior to the analysis, sampled cells are washed with phosphate-buffered saline that had been filtered (filter pore size, 0.22 micrometers), diluted to an OD600 of 0.05, and placed on ice. For each sample, 30,000 events are collected at a rate between 500 and 1,000 events/s. The percentage of induced (fluorescent) cells in each sample can be calculated from the flow cytometry data.

Example 7

Purification of Antibodies

[0128] Antibodies produced by the inducible coexpression systems of the invention are purified by centrifuging samples of lysed host cells at 10,000×g for 10 minutes to remove any cells and debris. The supernatant is filtered through a 0.45 micrometer filter. A 1-ml Recombinant Protein G—Sepharose® 4B column (Life Technologies, Grand Island, N.Y.) is set up to achieve flow rates of 1 ml/min, and is used with the following buffers: binding buffer: 0.02 M sodium phosphate, pH 7.0; elution buffer: 0.1 M glycine-HCl, pH 2.7; and neutralization buffer: 1 M Tris-HCl, pH 9.0. The column is equilibrated with 5 column volumes (5 ml) of binding buffer, and then the sample is applied to the column. The column is washed with 5-10 column volumes of the binding buffer to remove impurities and unbound material, continuing until no protein is detected in the eluent (determined by UV absorbance at 280 nm). The column is then eluted with 5 column volumes of elution buffer, and the column is immediately re-equilibrated with 5-10 column volumes of binding buffer.

Example 8

Measurement of Antibody Binding Affinity

[0129] The antibody binding affinity, expressed as "Kd" or "Kd value", is measured by a radiolabeled antigen-binding assay (RIA) performed with the Fab version of an antibody of interest and its antigen as described by the following assay. Production of the Fab version of a full-length antibody is well known in the art. Solution-binding affinity of Fabs for antigen is measured by equilibrating Fab with a minimal concentration of (125 I)-labeled antigen in the presence of a titration series of unlabeled antigen, then capturing bound antigen with an anti-Fab antibody-coated plate (see, for example, Chen et al., "Selection and analysis of an optimized anti-VEGF antibody: crystal structure of an affinity-matured Fab in complex with antigen", J Mol Biol 1999 Nov. 5; 293(4): 865-881). To establish conditions for the assay, microtiter plates (DYNEX Technologies, Inc., Chantilly, Va.) are coated overnight with 5 micrograms/ml of a capturing anti-Fab antibody (Cappel Labs, West Chester, Pa.) in 50 mM sodium carbonate (pH 9.6), and subsequently blocked with 2% (w/v) bovine serum albumin in PBS for two to five hours at room temperature (approximately 23° C.). In a non-adsorbent plate (Nunc #269620; Thermo Scientific, Rochester, N.Y.), 100 pM or 26 pM [125 I]-antigen are mixed with serial dilutions of a Fab of interest (e.g., consistent with assessment of the anti-VEGF antibody, Fab-12, in Presta et al., "Humanization of an

anti-vascular endothelial growth factor monoclonal antibody for the therapy of solid tumors and other disorders", Cancer Res 1997 Oct. 15; 57(20): 4593-4599). The Fab of interest is then incubated overnight; however, the incubation may continue for a longer period (e.g., about 65 hours) to ensure that equilibrium is reached. Thereafter, the mixtures are transferred to the capture plate for incubation at room temperature (e.g., for one hour). The solution is then removed and the plate washed eight times with 0.1% TWEEN-20TM surfactant in PBS. When the plates have dried, 150 microliters/well of scintillant (MICROSCINT-20TM; PerkinElmer, Waltham, Mass.) is added, and the plates are counted on a TOP-COUNTTM gamma counter (PerkinElmer) for ten minutes. Concentrations of each Fab that give less than or equal to 20% of maximal binding are chosen for use in competitive-binding assays.

[0130] Alternatively, the Kd or Kd value is measured by using surface-plasmon resonance assays using a BIACORE[®]-2000 or a BIACORE[®]-3000 instrument (BIACore, Inc., Piscataway, N.J.) at 25° C. with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIACore Inc.) are activated with N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 micrograms/ml (~0.2 micromolar) before injection at a flow rate of 5 microliters/minute to achieve approximately 10 RU of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% TWEEN 20TM surfactant (PBST) at 25° C. at a flow rate of approximately 25 microliters/min. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIACore[®] Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (Kd) is calculated as the ratio k_{off}/k_{on} . If the on-rate exceeds 10⁶ M⁻¹ s⁻¹ the surface-plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence-emission intensity (excitation=295 nm; emission=340 nm, 16 nm band-pass) at 25° C. of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow-equipped spectrophotometer (Aviv Instruments) or a 8000-series SLMAMINCOTM spectrophotometer (ThermoSpectronic) with a stirred cuvette.

[0131] Another method for determining the equilibrium dissociation constant (Kd) for antibody-antigen binding uses the Octet Red system (ForteBio, Pall Corporation, Port Washington, N.Y.) (www.fortebio.com/octet-RED96.html). The initial measurement step determines the baseline, followed by loading the His-tagged antigen at a concentration of 25 nM onto Ni-NTA biosensors for 10 minutes in 1X KB+ buffer (0.01% BSA, 0.002% Tween-20 in PBS, pH7.4), followed by another baseline measurement step (1X KB+ buffer only for 2 minutes). The sensor is then dipped into a well containing antibody (the association step) for 10 minutes, followed by a 20-minute wash in 1X KB+ buffer to measure dissociation. The equilibrium dissociation constant (Kd) is calculated as the ratio of k_{off}/k_{on} , with the Octet software determining the k_{off} and k_{on} rates.

Example 9

Characterizing the Disulfide Bonds Present in Coexpression Products

[0132] The number and location of disulfide bonds in protein coexpression products can be determined by digestion of the protein with a protease, such as trypsin, under non-reducing conditions, and subjecting the resulting peptide fragments to mass spectrometry (MS) combining sequential electron transfer dissociation (ETD) and collision-induced dissociation (CID) MS steps (MS2, MS3) (Nili et al., "Defining the disulfide bonds of insulin-like growth factor-binding protein-5 by tandem mass spectrometry with electron transfer dissociation and collision-induced dissociation", J Biol Chem 2012 Jan. 6; 287(2): 1510-1519; Epub 2011 Nov. 22).

Digestion of Coexpressed Protein.

[0133] To prevent disulfide bond rearrangements, any free cysteine residues are first blocked by alkylation: the coexpressed protein is incubated protected from light with the alkylating agent iodoacetamide (5 mM) with shaking for 30 minutes at 20° C. in buffer with 4 M urea. Alternatively and preferably, NEM is used as the alkylating reagent, with trypsin proteolysis in combination with reduction/alkylation conducted under denaturing conditions (6M GuaHCl). Following alkylation, the coexpressed protein is separated by non-reducing SDS-PAGE using precast gels. Alternatively, the coexpressed protein is incubated in the gel after electrophoresis with iodoacetamide or NEM, or without as a control. Protein bands are stained, de-stained with double-deionized water, excised, and incubated twice in 500 microliters of 50 mM ammonium bicarbonate, 50% (v/v) acetonitrile while shaking for 30 minutes at 20° C. Protein samples are dehydrated in 100% acetonitrile for 2 minutes, dried by vacuum centrifugation, and rehydrated with 10 mg/ml of trypsin or chymotrypsin in buffer containing 50 mM ammonium bicarbonate and 5 mM calcium chloride for 15 minutes on ice. Excess buffer is removed and replaced with 50 microliters of the same buffer without enzyme, followed by incubation for 16 hours at 37° C. or 20° C., for trypsin and chymotrypsin, respectively, with shaking. Digestions are stopped by addition of 3 microliters of 88% formic acid, and after brief vortexing, the supernatant is removed and stored at -20° C. until analysis. Alternative protein fragmentation methods (LysC, Glu-C, or CNBr) are used if trypsinolysis provides insufficient sequence coverage (<75%). Using the reducing agent TCEP (tris(2-carboxyethyl)phosphine) under acidic conditions in the presence of NEM provides access to fragments with partly intact disulfide linkages. The disulfide-intact digest map is compared to the reduced (DTT or TCEP) digest map.

Localization of Disulfide Bonds by Mass Spectrometry.

[0134] Peptides are injected onto a 1 mm×8 mm trap column (Michrom BioResources, Inc., Auburn, Calif.) at 20 microliters/minute in a mobile phase containing 0.1% formic acid. The trap cartridge is then placed in-line with a 0.5 mm×250 mm column containing 5 mm Zorbax SB-C18 stationary phase (Agilent Technologies, Santa Clara, Calif.), and peptides separated by a 2-30% acetonitrile gradient over 90 minutes at 10 microliters/minute with a 1100 series capillary HPLC (Agilent Technologies); alternatively, a C18 column suitable for UPLC is used. Peptides are analyzed using a LTQ

Velos linear ion trap with an ETD source (Thermo Fisher Scientific Inc., Waltham, Mass.). Electrospray ionization is performed using a Captive Spray source (Michrom Bioresources, Inc.), or preferably, an uncoated, pulled fused silica emitter (New Objective Inc., Woburn, Massachusetts) at 3.0 kV. Alternatively, analysis of medium-sized proteolytic fragments is performed using a Thermo LTQ-FT MS (7 Tesla) instrument, or a Synapt G2-Si quadrupole traveling wave ion mobility time-of-flight (ToF) mass spectrometer (Waters Corp., Milford, Mass.). Preferably, peptides are analyzed using an Orbitrap Fusion™ Tribrid™ mass spectrometer (Thermo Fisher Scientific). Disulfide-linked peptides have charge states of +4 or greater following trypsinization due to the presence of two N-termini and two basic residues (arginine or lysine) at the carboxy termini. These disulfide-linked peptides are preferentially isolated by the Orbitrap Fusion™ instrument so that the disulfide bonds can be broken using ETD fragmentation. Survey MS scans are followed by seven data-dependant scans consisting of CID and ETD MS2 scans on the most intense ion in the survey scan, followed by five MS3 CID scans on the first- to fifth-most intense ions in the ETD MS2 scan. CID scans use normalized collision energy of 35, and ETD scans use a 100 ms activation time with supplemental activation enabled. Minimum signals to initiate MS2 CID and ETD scans are 10,000, minimum signals for initiation of MS3 CID scans are 1000, and isolation widths for all MS2 and MS3 scans are 3.0 m/z. The dynamic exclusion feature of the software is enabled with a repeat count of 1, exclusion list size of 100, and exclusion duration of 30 seconds. Inclusion lists to target specific cross-linked species for collection of ETD MS2 scans are used. Separate data files for MS2 and MS3 scans are created by Bioworks 3.3 (Thermo Fisher Scientific) using ZSA charge state analysis. Matching of MS2 and MS3 scans to peptide sequences is performed by Sequest (V27, Rev 12, Thermo Fisher Scientific). The analysis is performed without enzyme specificity, a parent ion mass tolerance of 2.5, fragment mass tolerance of 1.0, and a variable mass of +16 for oxidized methionine residues. Results are then analyzed using the program Scaffold (V3_00_08, Proteome Software, Portland, Oreg.) with minimum peptide and protein probabilities of 95 and 99% being used. Software tools for data interpretation also include Proteome Discoverer™ 2.0 with the Disulfinator node (Thermo Fisher Scientific). Peptides from MS3 results are sorted by scan number, and cysteine containing peptides are identified from groups of MS3 scans produced from the five most intense ions observed in ETD MS2 scans. The identities of cysteine peptides participating in disulfide-linked species are further confirmed by manual examination of the parent ion masses observed in the survey scan and the ETD MS2 scan.

Example 10

Isolation of Coexpression Products from Bacterial Cell Periplasm, from Spheroplasts, and from Whole Cells

[0135] The inducible coexpression system of the invention can be used to express gene products that accumulate in different compartments of the cell, such as the cytoplasm or periplasm. Host cells such as *E. coli* or *S. cerevisiae* have an outer cell membrane or cell wall, and can form spheroplasts when the outer membrane or wall is removed. Coexpressed proteins made in such hosts can be purified specifically from the periplasm, or from spheroplasts, or from whole cells,

using the following method (Schoenfeld, "Convenient, rapid enrichment of periplasmic and spheroplasmic protein fractions using the new PeriPreps™ Periplasting Kit", Epicentre Forum 1998 5(1): 5; see www.epibio.com/newsletter/f5_1/f5_1pp.asp). This method, using the PeriPreps™ Periplasting Kit (Epicentre® Biotechnologies, Madison Wis.; protocol available at www.epibio.com/pdftechlit/107p10612.pdf), is designed for *E. coli* and other gram negative bacteria, but the general approach can be modified for other host cells such as *S. cerevisiae*.

[0136] 1. The bacterial host cell culture is grown to late log phase only, as older cell cultures in stationary phase commonly demonstrate some resistance to lysozyme treatment. If the expression of recombinant protein is excessive, cells may prematurely lyse; therefore, cell cultures are not grown in rich medium or at higher growth temperatures that might induce excessive protein synthesis. Protein expression is then induced; the cells should be in log phase or early stationary phase.

[0137] 2. The cell culture is pelleted by centrifugation at a minimum of 1,000×g for 10 minutes at room temperature. Note: the cells must be fresh, not frozen. The wet weight of the cell pellet is determined in order to calculate the amount of reagents required for this protocol.

[0138] 3. The cells are thoroughly resuspended in a minimum of 2 ml of PeriPreps Periplasting Buffer (200 mM Tris-HCl pH 7.5, 20% sucrose, 1 mM EDTA, and 30 U/microliter Ready-Lyse Lysozyme) for each gram of cells, either by vortex mixing or by pipeting until the cell suspension is homogeneous. Note: excessive agitation may cause premature lysing of the spheroplasts resulting in contamination of the periplasmic fraction with cytoplasmic proteins.

[0139] 4. Incubate for five minutes at room temperature. Ready-Lyse Lysozyme is optimally active at room temperature. Lysis at lower temperatures (0° C.-4° C.) requires additional incubation time; at such temperatures incubation times are extended 2- to 4-fold.

[0140] 5. Add 3 ml of purified water at 4° C. for each gram of original cell pellet weight (Step 2) and mix by inversion.

[0141] 6. Incubate for 10 minutes on ice.

[0142] 7. The lysed cells are pelleted by centrifugation at a minimum of 4,000×g for 15 minutes at room temperature.

[0143] 8. The supernatant containing the periplasmic fraction is transferred to a clean tube.

[0144] 9. To degrade contaminating nucleic acids, OmniCleave Endonuclease is optionally added to PeriPreps Lysis Buffer. Inclusion of a nuclease will generally improve the yield of protein and the ease of handling of the lysates, but addition of a nuclease is undesirable in some cases: for example, the use of a nuclease should be avoided if residual nuclease activity or transient exposure to the magnesium cofactor will interfere with subsequent assays or uses of the purified protein. The addition of EDTA to the lysate to inactivate OmniCleave Endonuclease, likewise, may interfere with subsequent assay or use of the purified protein. If nuclease is to be added, 2 microliters of OmniCleave Endonuclease and 10 microliters of 1.0 M MgCl₂ are diluted up to 1 ml with PeriPreps Lysis Buffer (10 mM Tris-HCl pH 7.5, 50 mM KCl, 1 mM EDTA, and 0.1% deoxycholate) for each milliliter of Lysis Buffer needed in Step 10.

[0145] 10. The pellet is resuspended in 5 ml of PeriPreps Lysis Buffer for each gram of original cell pellet weight.

[0146] 11. The pellet is incubated at room temperature for 10 minutes (if included, OmniCleave Endonuclease activity

will cause a significant decrease in viscosity; the incubation is continued until the cellular suspension has the consistency of water).

[0147] 12. The cellular debris is pelleted by centrifugation at a minimum of 4,000 \times g for 15 minutes at 4° C.

[0148] 13. The supernatant containing the spheroplast fraction is transferred to a clean tube.

[0149] 14. If OmniCleave Endonuclease was added to the PeriPreps Lysis Buffer, 20 microliters of 500 mM EDTA is added for each milliliter of the resultant spheroplast fraction, to chelate the magnesium (the final concentration of EDTA in the lysate is 10 mM). Following hydrolysis of nucleic acids with OmniCleave Endonuclease, lysates may contain substantial amounts of mono- or oligonucleotides. The presence of these degradation products may affect further processing of the lysate: for example, nucleotides may decrease the binding capacity of anion exchange resins by interacting with the resin.

[0150] The above protocol can be used to prepare total cellular protein with the following modifications. The cells pelleted in Step 2 can be fresh or frozen; at Step 4, the cells are incubated for 15 minutes; Steps 5 through 8 are omitted; at Step 10, 3 ml of PeriPreps Lysis Buffer is added for each gram of original cell pellet weight.

[0151] After preparation of periplasmic, or spheroplasmic, or whole-cell protein samples, the samples can be analyzed by any of a number of protein characterization and/or quantification methods. In one example, the successful fractionation of periplasmic and spheroplasmic proteins is confirmed by analyzing an aliquot of both the periplasmic and spheroplasmic fractions by SDS-PAGE (two microliters of each fraction is generally sufficient for visualization by staining with Coomassie Brilliant Blue). The presence of unique proteins or the enrichment of specific proteins in a given fraction indicates successful fractionation. For example, if the host cell contains a high-copy number plasmid with the ampicillin resistance marker, then the presence of β -lactamase (31.5 kDa) mainly in the periplasmic fraction indicates successful fractionation. Other *E. coli* proteins found in the periplasmic space include alkaline phosphatase (50 kDa) and elongation factor Tu (43 kDa). The amount of protein found in a given fraction can be quantified using any of a number of methods (such as SDS-PAGE and densitometry analysis of stained or labeled protein bands, scintillation counting of radiolabeled proteins, enzyme-linked immunosorbent assay (ELISA), or scintillation proximity assay, among other methods.) Comparing the amounts of a protein found in the periplasmic fraction as compared to the spheroplasmic fraction indicates the degree to which the protein has been exported from the cytoplasm into the periplasm.

Example 11

Determination of Polynucleotide or Amino Acid Sequence Similarity

[0152] Percent polynucleotide sequence or amino acid sequence identity is defined as the number of aligned symbols, i.e. nucleotides or amino acids, that are identical in both aligned sequences, divided by the total number of symbols in the alignment of the two sequences, including gaps. The degree of similarity (percent identity) between two sequences may be determined by aligning the sequences using the global alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as implemented by the National Center for

Biotechnology Information (NCBI) in the Needleman-Wunsch Global Sequence Alignment Tool, available through the website blast.ncbi.nlm.nih.gov/Blast.cgi. In one embodiment, the Needleman and Wunsch alignment parameters are set to the default values (Match/Mismatch Scores of 2 and -3, respectively, and Gap Costs for Existence and Extension of 5 and 2, respectively). Other programs used by those skilled in the art of sequence comparison may also be used to align sequences, such as, for example, the basic local alignment search tool or BLAST® program (Altschul et al., “Basic local alignment search tool”, *J Mol Biol* 1990 Oct. 5; 215(3): 403-410), as implemented by NCBI, using the default parameter settings described at the blast.ncbi.nlm.nih.gov/Blast.cgi website. The BLAST algorithm has multiple optional parameters including two that may be used as follows: (A) inclusion of a filter to mask segments of the query sequence that have low compositional complexity or segments consisting of short-periodicity internal repeats, which is preferably not utilized or set to ‘off’, and (B) a statistical significance threshold for reporting matches against database sequences, called the ‘Expect’ or E-score (the expected probability of matches being found merely by chance; if the statistical significance ascribed to a match is greater than this E-score threshold, the match will not be reported). If this ‘Expect’ or E-score value is adjusted from the default value (10), preferred threshold values are 0.5, or in order of increasing preference, 0.25, 0.1, 0.05, 0.01, 0.001, 0.0001, 0.00001, and 0.000001.

[0153] In practicing the present invention, many conventional techniques in molecular biology, microbiology, and recombinant DNA technology are optionally used. Such conventional techniques relate to vectors, host cells, and recombinant methods. These techniques are well known and are explained in, for example, Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* volume 152 Academic Press, Mc, San Diego, Calif.; Sambrook et al., *Molecular Cloning—A Laboratory Manual* (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 2000; and *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2006). Other useful references, for example for cell isolation and culture and for subsequent nucleic acid or protein isolation, include Freshney (1994) *Culture of Animal Cells, a Manual of Basic Technique*, third edition, Wiley-Liss, New York and the references cited therein; Payne et al. (1992) *Plant Cell and Tissue Culture in Liquid Systems* John Wiley & Sons, Inc. New York, N.Y.; Gamborg and Phillips (Eds.) (1995) *Plant Cell, Tissue and Organ Culture; Fundamental Methods* Springer Lab Manual, Springer-Verlag (Berlin Heidelberg N.Y.); and Atlas and Parks (Eds.) *The Handbook of Microbiological Media* (1993) CRC Press, Boca Raton, Fla. Methods of making nucleic acids (for example, by in vitro amplification, purification from cells, or chemical synthesis), methods for manipulating nucleic acids (for example, by site-directed mutagenesis, restriction enzyme digestion, ligation, etc.), and various vectors, cell lines, and the like useful in manipulating and making nucleic acids are described in the above references. In addition, essentially any polynucleotide (including labeled or biotinylated polynucleotides) can be custom or standard ordered from any of a variety of commercial sources.

[0154] The present invention has been described in terms of particular embodiments found or proposed to comprise certain modes for the practice of the invention. It will be appre-

ciated by those of ordinary skill in the art that, in light of the present disclosure, numerous modifications and changes can be made in the particular embodiments exemplified without departing from the intended scope of the invention.

[0155] All cited references, including patent publications, are incorporated herein by reference in their entirety. Nucleotide and other genetic sequences, referred to by published genomic location or other description, are also expressly incorporated herein by reference.

-continued

SEQUENCES PRESENTED IN THE SEQUENCE LISTING

SEQ ID	NO:	Length:	Type:	Organism:	Description; 'Other Information'
1	35	DNA	Artificial Sequence	J23104 promoter	
2	5883	DNA	Artificial Sequence	Expression vector pPRO43	
3	5882	DNA	Artificial Sequence	pPRO33 vector	
4	3698	DNA	Artificial Sequence	Expression vector pPRO430	
5	3610	DNA	Artificial Sequence	Expression vector pPRO430(CloDF13)	
6	3255	DNA	Artificial Sequence	Expression vector pBAD240	
7	5802	DNA	Artificial Sequence	Expression vector pPRO44	
8	5791	DNA	Artificial Sequence	Expression vector pPRO45	

SEQ ID	NO:	Length:	Type:	Organism:	Description; 'Other Information'
9	25	DNA	Artificial Sequence	pPRO_REV_SpeI primer	
10	28	DNA	Artificial Sequence	pPRO_FOR_AatII primer	
11	46	DNA	Artificial Sequence	RSF1030_FOR_SpeI primer	
12	52	DNA	Artificial Sequence	RSF1030_REV_AatII primer	
13	46	DNA	Artificial Sequence	CloDF13_FOR_SpeI primer	
14	56	DNA	Artificial Sequence	CloDF13_REV_AatII primer	
15	5304	DNA	Artificial Sequence	Dual-promoter vector, pSOL	
16	451	PRT	Artificial Sequence	Infliximab chimeric (murine variable domain, human constant domain) heavy chain	
17	215	PRT	Artificial Sequence	Infliximab chimeric (murine variable domain, human constant domain) light chain	
18	4594	DNA	Artificial Sequence	pBAD240-Infliximab_HC expression construct	
19	4332	DNA	Artificial Sequence	pPRO430-Infliximab_LC expression construct	
20	7288	DNA	Artificial Sequence	pSOL-Infliximab expression construct	

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 20

<210> SEQ ID NO 1

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: J23104 promoter

<400> SEQUENCE: 1

ttgacagcta gtcagtcct aggtattgtg cttagc

35

<210> SEQ ID NO 2

<211> LENGTH: 5883

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Expression vector pPRO43

<400> SEQUENCE: 2

atcgatttag cttttcagac gacgccaaaa ggtcgatcg gaaataccca aatagttggc 60
 cgcagccgtc ttgtcaccat taaacttctc aagcgcttgc tgcgggtca gcaaacgcgg 120
 agccggcgatc tttgcgcctc cacgcgcacg ctccggcagc agcaactgca tgaattgcgg 180
 agtcagatcc ggggtcggtt caacggacag gaacagcgcc aggcgttcca tcataattacg 240
 cagctcgccg atgttacccg gccagtcata atgcagcagc accgttgcg tgcctgcag 300
 accctggcgc agtgccgcag agaacgggtc gctcagggtc gccagcgaga ctttcaggaa 360
 agactccgccc agcggtaaaa tgtcgccgac acgttacgc aacggcgaaa gctgcagacg 420

-continued

cagaatgctc	aggcggtaga	acaggtcgcg	acgaaaacgg	ccctgttgc	tatccttcc	480
cagattgcag	tgggtcgccc	taatcacgcg	cacgtctacc	ggaaccgggt	gatgaccacc	540
gacgcgggctc	acttctttct	cttccagcac	acgcagcaga	cgggtttgc	atggcagcgg	600
catctcaccc	atctcgcca	ggaacaaggt	gcccgcgtg	gcaatttcaa	acaaaccagc	660
acggccaccc	cgacggctac	ccgtgaatgc	gcccttccg	tagccaaaca	gctcagctc	720
caggaggctt	tccgcgattt	caccgcaatt	aactgcaca	aacggatgag	atttcttacc	780
ctggcgggca	tcgtgacggg	cgaataactc	acgatgggat	gcttgcgcag	ccagttcc	840
acccgtacca	gtctcgcc	cgatcagaac	agccgcgc	ctacgtgc	atacgagaat	900
ggtctggcga	acttgctca	tttgagggt	ttggccca	atatcaccca	ggacataacg	960
ggtacgcagc	gcattacgcg	tcgcacgtg	ggtgttgc	cgcaggctca	ttctggtcat	1020
gtccaggcgc	tcgcacgc	cctgacgcac	cgttgcgc	ctgtagataa	agatgc	1080
catgcccgt	tcttcggca	agtccgtgat	cagaccgc	ccaaccacag	cctcggtacc	1140
gttcgcgtt	agttcgttga	tctggccacg	tgcac	tccgtatgt	agctgcgttgc	1200
gtccaggcgc	agattaaagg	tctttgaaa	cgcgacc	gcaggatcg	tttctggta	1260
ggtgacaa	ccaatcgagg	aggtcagttt	gcctgc	gccagegc	gcaagacatc	1320
gttaaccgc	ggttaatca	ggatcacgg	cacggacaga	cggttca	ggtaggcacc	1380
attgctaccc	gctgcgataa	tggcgtaca	acgctcg	gccag	tttgc	1440
ggtaacgg	ttctcgaa	ccagctgat	cgagt	ttcgc	ccagg	1500
caggcta	atcgcc	actcgac	acgggt	gacg	ctaacgttcc	1560
tttatcatcg	ttcaa	acgcg	gtgggtgt	catgg	gat	1620
atattgggtt	taaactt	tataattt	tgc	aatggaaa	cattgc	1680
aacgtta	gaaacgcata	tttgcgg	attt	catgc	tttgc	1740
at	taat	tttatt	aggc	atgc	tttgc	1800
gcaaaataaca	agttgataac	aagctagcga	attcgac	ggtac	ccgg	1860
agtcgac	caggcatc	agttgg	tttgg	tgaga	tttgc	1920
gatacagatt	aaatcagaac	gcaga	aggc	tctg	atggc	1980
tagcgcgt	gtcccac	acccatgc	gaactc	gtaa	acgc	2040
tggtagt	gggtct	atgcg	agg	actgc	cagg	2100
aggctcag	gaaagactgg	gcctt	ttatctgt	tttgc	tttgc	2160
tgagt	aaatccgc	ggagg	tga	acgt	ccgg	2220
ggcgggc	acgcccgc	taaa	actgc	ggc	ataaa	2280
cggatggc	ttttgcgtt	ctaca	actc	ttttctaa	at	2340
atgtatcc	tcatgagaca	ataacc	ctg	taatgc	tttgc	2400
atgtatg	tca	acatt	ccgt	gtcg	tttgc	2460
cctgtttt	ctcac	ccaga	aacg	ctgg	tttgc	2520
gcagcaaa	actt	aaact	gcgc	ctct	tttgc	2580
ggatggaggc	ggataa	agg	tcg	ccgc	ttccg	2640
ttattgc	taa	atct	gg	gtgg	tcgt	2700

-continued

ggccagatgg taagccctcc	cgtatcgtag ttatctacac	gacggggagt caggcaacta	2760				
tggatgaacg aaatagacag	atcgctgaga taggtgcctc	actgattaag cattggtaac	2820				
tgtcagacca agtttactca	tatatacttt agattgattt	acgcgcctg tagcggcgca	2880				
ttaagcgcgg cgggtgtgg	ggttacgcgc	agcgtgaccg ctacacttgc	2940				
gcgcgcgc	ctttcgttt	cttccctcc tttctcgcca	cgttcgcgg ctttccccgt	3000			
caagctctaa atcggggct	ccctttaggg	ttccgattt gtgcttaag	gcacctcgac	3060			
cccaaaaaac ttgatttgg	tgatggttca	cgtagtggc catgcctcg	atagacggtt	3120			
tttcgcctt tgacgttg	gtccacgttc	ttaatagtg	gactcttgc	3180			
acaacactca accctatctc	gggctattct	tttgatttat aagggat	tttgcattcg	3240			
gcctatttgt	taaaaaatga	gctgatttaa	caaaaattta acgcgaattt	taacaaaata	3300		
ttaacgttta caatttaaa	ggatcttagt	gaagatcctt	tttgataatc	tcatgaccaa	3360		
aatcccttaa cgtgagttt	cgttccactg	agcgtcagac	cccgtagaaa	agatcaaagg	3420		
atcttcttga gatcctttt	tttgcgcgt	aatctgctgc	ttgcacaa	aaaaaccacc	3480		
gctaccagcg	gtggttgtt	tgccggatca	agagctacca	actcttttc	cgaaggtaac	3540	
tggttcagc	agagcgcaga	taccaaatac	tgtccttc	gtgtacgcgt	agtttaggc	3600	
ccacttcaag	aactctgt	caccgcctac	atacctcg	ctgctaattc	tgttaccagt	3660	
caggcattt	agaagcacac	ggtcacactg	cttccggtag	tcaataaacc	ggtaaaccag	3720	
caatagacat	aagcggctat	ttaacgaccc	tgccctgaac	cgacgaccgg	gtcgaattt	3780	
ctttcgaatt	tctgcatttc	atccgcttat	tatcacttat	tcaggcgt	caaccaggcg	3840	
ttaagggca	ccaataactg	cctaaaaaa	attacgc	cccgcac	tcatecg	3900	
actgttgtaa	tccattaagc	attctgcgc	catgaa	gaccaaa	gcacatgaa	3960	
cctgaatcgc	cagcggc	atcaccttg	cgccttgcgt	ataatattt	ccatgtga	4020	
aaacggggc	gaagaagtt	tccatattgg	ccacgtttaa	atcaaactg	gtgaaactca	4080	
cccaggatt	ggctgagacg	aaaaacat	tctcaataaa	ccctttaggg	aaataggc	4140	
gttttca	gtaacacg	acatcttgc	aatatgtg	tagaaactg	cggaaatcg	4200	
cgttgtattc	actccagac	gatgaaac	tttcagttt	ctcatggaa	acggtgtaac	4260	
aagggtgaac	actatccat	atcaccag	caccgtctt	cattgcata	cggaattcc	4320	
gatgagcatt	catcaggcgg	gcaagaatgt	gaataaaggc	cggataaaac	ttgtgttat	4380	
ttttcttac	ggtctttaa	aaggcgtaa	tatccagctg	aacggctgg	ttataggtac	4440	
attgagcaac	tgactgaaat	gcctcaaaat	gttcttacg	atgcacatgg	gatata	4500	
cggtgtata	tccagtgatt	ttttctcca	ttttagctc	cttagctc	gaaaatctcg	4560	
ataactcaa	aaatacgc	ggtagtgc	ttatccatt	atggtaa	ttggac	4620	
ttacgtgcc	atcaacgt	catttcg	aaaagtggc	ccagggttc	ccggat	4680	
caggcacacc	aggat	tattctg	agtgat	cttc	cgtcaca	4740	
cgcggaaatgc	gtcggtgat	gtgc	actgt	gttgc	gtgtat	4800	
ggtgctccag	tggcttctgt	ttctatc	tgccctc	gttc	tgacgggt	4860	
gtgcgtaa	gc	aaaagc	cgccg	acat	agcgg	4920	
actatgttg	cactgat	gag	ggtgt	cagt	gaa	aaaaaggc	4980

-continued

tgcaccgggt	cgtcagcaga	atatgtgata	caggatata	tccgcttct	cgctca	tga	5040
ctcgctacgc	tcggcggttc	gactgcggcg	agcgaaatg	gcttacgaac	ggggcg	gaga	5100
tttcctggaa	gatgccagga	agataactta	caggaaatg	agagggccgc	ggcaa	agccg	5160
ttttccata	ggtccgccc	ccctgaca	catcacgaaa	tctgacgctc	aaatca	gttgg	5220
tggcgaaacc	cgacaggact	ataaagata	caggcggttc	cccctggcg	ctccctcg	tg	5280
cgtctctgt	ttctgcatt	tcgggttacc	gggtcattc	cgtgttatg	gccgcgtt	g	5340
tctcattcca	cgcgtacac	tca	gttccgg	gttggcgtt	cgctca	agg	5400
gcacgaaccc	cccggttca	cgaccgcgt	cgccat	ggtaactatc	gtcttg	gatc	5460
caacccggaa	agacatgaa	aagcaccact	ggcagcagcc	actggtaatt	gat	tttagagg	5520
agtttagtctt	gaagtcatgc	gcgggttaag	gctaaactga	aaggacaat	tttgg	tgact	5580
gcgtctcc	aaggcgttta	cctcggttca	aagagtgg	agtcagaga	ac	cttgc	5640
aacggccctg	caaggcggtt	tttcgtttt	cagagcaaga	gattacgc	agacca	aaac	5700
gatctcaaga	agatcatctt	attaatcaga	taaaatattt	gctcatg	gc	cgaa	5760
gagcccgatc	ttccccatcg	gtgtatgtgg	cgatata	gcacgcaacc	gcac	ctgtgg	5820
cgccgggtgat	gcggccacg	atgcgtccgg	cgtagaggat	ctgctcatgt	ttgac	agctt	5880
atc							5883

<210> SEQ ID NO 3
 <211> LENGTH: 5882
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pPRO33 vector

<400> SEQUENCE: 3

atcgattcag	ctttcagcc	gcccacaa	cgtcgccgg	ctgatgccta	aataattcgc		60
cgctgctgtt	ttatcgccat	taaatttctc	cagtgcctgt	tgtgggtca	gtaaggtgg		120
agcgggagtt	ttegcgcact	cgcgcgc	ttccggca	gtcgttgc	taaactgcgg		180
cgttaaatcc	ggegtcggtt	ccacacttaa	aaacageg	cgccat	tcatattgc		240
cagttcacga	atattgcctg	gca	gtgcagc	acat	ttgcctgt	aa	300
cccttggcgt	aatgcagcag	aaaatgggg	ggagag	gc	cagagaca	ctttcaaaaa	360
gcttccgccc	agcggaaagaa	tatccgcac	ccgc	tcg	agtgg	ggca	420
caaaatactc	agccgataaa	acagatc	ac	cgaaaacgt	ccttgc	tatcttctc	480
cagattgcag	tgagtggc	taatgac	cc	gaaacag	gt	gcccc	540
gacgcgggtg	acctctttt	cttcc	ccgc	cg	cg	aa	600
catttcgcca	atctcatcca	gaaac	ac	ccgttgg	gca	at	660
gcgac	ctcc	cg	tcg	cg	cc	aa	720
cagcagcgat	tcggcaatcg	ccccc	cg	ccgttgc	aa	tttgc	780
ctgtcgcc	tcgtggcg	caaaatattc	ccgat	gatc	gc	cttgc	840
gcccgtcccc	gttccccct	caatca	ac	ccgc	act	tttgc	900
agtctgc	cttgc	tctgtgg	tg	accg	at	atcgc	960
agtacgcagg	gcgttgc	gg	tggc	atcg	tcg	taac	1020

-continued

atccagcgca	tcgctgaacg	cctggcgcac	ggtggcgccg	gaatagataa	aaattccggt	1080
cattccggct	tcttctgcca	aatcggtaat	cagccctgcg	ccgaccacccg	cttcggtgcc	1140
gttagctttt	agctcgtaa	tctgcccgcg	tgcgctttcc	tcggtaatgt	agctacgtt	1200
gtcgaggcgc	aaattaaagg	tttttgaaa	cgccaccagc	gctggaatgg	tttcctgata	1260
ggtagacaacg	ccgatagaag	aggtgagttt	tccggctttt	gccagtgct	gtaacacatc	1320
gtagccgctc	ggtttaatca	aaataactgg	cactgacagg	cggtttca	ggtagcgcc	1380
gttagagcca	gcccgcgtga	tggcgtcaca	gcgttcgtt	gccagtttct	tgcggtatgt	1440
ggtaactgt	ttttcaaagc	caagctgaat	aggggtaatg	tccgcccagg	gatcaaactc	1500
gaggctgata	tcgcgaaaca	gctcgaacag	gcccgttaca	gataccgtcc	agataaccgg	1560
tttgcgtca	ttaagccgtg	gtggatgtgc	catagcgcac	cgcaaaagtt	agaaaccgaa	1620
tattgggttt	agtcttgttt	cataattgtt	gcaatgaaac	gccccgtt	attgcctgaa	1680
acgttaactg	aaacgcata	ttgcggatta	gttcatgact	ttatctctaa	caaattgaaa	1740
ttaaacattt	aattttatta	aggcaattgt	ggcacacccc	ttgcgttgc	tttataacg	1800
caaataacaa	gttgataaca	agotagcgaa	tgcgagctcg	gtaccgggg	atcccttaga	1860
gtcgacctgc	aggcatgca	gcttggctgt	tttggcggt	gagagaagat	tttcagcctg	1920
atacagatta	aatcagaacg	cagaagegg	ctgataaaac	agaatttgcc	tggcggcagt	1980
agcgcgggtgg	tcccacctga	ccccatgcgg	aactcagaag	tggaaacggcc	tagcgcgtat	2040
ggtagtgtgg	ggtctccca	tgcgagagta	gggaactgcc	aggcatcaaa	taaaacgaaa	2100
ggctcagtcg	aaagactggg	ccttcgttt	tatctgtt	ttgtcggtga	acgctctcct	2160
gagtaggaca	aatccgcgg	gagcggattt	gaacgttgcg	aagcaacggc	ccggagggtg	2220
gcgggcagga	cgcggccat	aaactgcgg	gcatcaaatt	aagcagaagg	ccatctgac	2280
ggatggcctt	tttgcgtt	tacaaactct	tttgtttatt	tttctaaata	cattcaaata	2340
tgtatccgt	catgagacaa	taaccctgt	aatgcgttca	ataatattgt	aaaaggaaga	2400
gtatgagtat	tcaacattt	cgtgtcgccc	ttattccctt	ttttgeggca	ttttgccttc	2460
ctgttttgc	tcacccagaa	acgctgggt	aagtaaaaaga	tgctgaatgt	cagttgggt	2520
cagcaaacta	ttaactggcg	aactacttac	tctagcttcc	cgcaacaat	taatagact	2580
gatggaggcg	gataaagtt	caggaccact	tctgcgtcg	gcccggccg	ctggctgggt	2640
tattgctgt	aaatctggag	ccggtgagcg	tgggtctcgc	ggtatcattt	cagcactgg	2700
gccagatgg	aagccctccc	gtatcgtagt	tatctacacg	acggggagtc	aggcaactat	2760
ggatgaacga	aatagacaga	tcgctgagat	aggtgcctca	ctgattaacg	attggtaact	2820
gtcagaccaa	gtttactcat	atatacttta	gattgattt	cgcccccgt	agcggcgcat	2880
taagcgcggc	gggtgtgggt	gttacgcgca	gcgtgaccgc	tacacttgcc	agcgccttag	2940
cgcggcgtcc	tttcgttttc	ttcccttcct	ttctcgccac	gttcggccgc	tttccccgtc	3000
aagctctaaa	tcgggggtc	cctttaggt	tcggatttag	tgctttacgg	cacctogacc	3060
ccaaaaaact	tgatttgggt	gatggttcac	gtagtgccc	atcgccctga	tagacgttt	3120
tccgccttt	gacgttggag	tccacgttct	ttaatagtgg	actcttgc	caaactggaa	3180
caacactcaa	ccctatctcg	ggctattctt	ttgatttata	agggattt	ccgatttgg	3240
cctattggtt	aaaaaatgag	ctgatttaac	aaaaatttaa	cgcgaattt	aacaaaat	3300

-continued

-continued

cgctcctcca	agccagttac	ctcggttcaa	agagttggta	gctcagagaa	ccttcgaaaa	5640
accgcctgc	aaggcggtt	tttcgtttc	agagcaagag	attacgcgca	gaccaaaacg	5700
atctcaagaa	gatcatctta	ttaatcagat	aaaatatttgc	ctcatgagcc	cgaagtggcg	5760
agcccgatct	tccccatcgg	tgatgtcggc	gatataggcg	ccagcaaccg	cacctgtggc	5820
gccggtgatg	ccggccacga	tgcgccggc	gtagaggatc	tgctcatgtt	tgacagctta	5880
tc						5882
<210> SEQ ID NO 4						
<211> LENGTH: 3698						
<212> TYPE: DNA						
<213> ORGANISM: Artificial Sequence						
<220> FEATURE:						
<223> OTHER INFORMATION: Expression vector pPRO430						
<400> SEQUENCE: 4						
ccccagaggc	aggctctagcg	gagtgtatac	tggcttacta	tgttggact	gatgagggtg	60
tcagtgaagt	gcttcatgtg	gcaggagaaa	aaaggctgca	ccgggtgcgtc	agcagaatat	120
gtgatacagg	atataattcg	cttcctcgct	cactgactcg	ctacgctcg	tcgttgcact	180
gcggcgagcg	gaaatggctt	acgaacgggg	cgagatttc	ctggaagatg	ccaggaagat	240
acttaacagg	gaagtgagag	ggccgcggca	aagccgtttt	tccataggt	ccgccccct	300
gacaaggcata	acgaaatctg	acgctcaat	cagtgggtgc	gaaacccgac	aggactataa	360
agataccagg	cgtttcccc	tggcggtc	ctcgtgcgt	ctcctgttcc	tgccttcgg	420
tttaccggtg	tcattccgct	gttatggcc	cgtttgc	attccacg	tgacactcag	480
ttccggtag	gcagttcgct	ccaagctgga	ctgtatgcac	gaacccccc	ttcagtcg	540
ccgctgcgc	ttatccgta	actatcgct	tgagtccaa	ccggaaagac	atgaaaaagc	600
accactggca	gcagccactg	gtaattgatt	tagaggagg	agtcttgc	tcatgcggc	660
gttaaggcta	aactgaaagg	acaagtttt	gtgactgcgc	tcctccaa	cagtacactc	720
gttcaaaga	gttggtagct	cagagaacct	tgc	ccctgcaag	gcggttttt	780
cgtttcaga	gcaagagatt	acgcgcagac	caaaacgatc	tcaagaagat	cattttatta	840
aacctgcctc	taaaaactt	tattacgccc	cgccctgc	ctcatacg	tactgttgc	900
attcattaag	cattctgcgc	acatgaa	catcacagac	ggcatgatg	acctgaatcg	960
ccagcggcat	cagcaccttgc	tgc	tataatattt	gccatagtg	aaaacgggg	1020
cgaagaagtt	gtccatatttgc	gccacgttta	aatcaaaact	ggtgaaactc	acccaggat	1080
tggctgagac	aaaaacata	ttctcaata	accctttag	gaaataggcc	aggtttgc	1140
cgtaaacacgc	cacatcttgc	gaatata	gtgaaactg	ccggaaatcg	tcgtggatt	1200
cactccagag	cgtgaaaac	gttgcgtt	gctcatggaa	aacgggtgaa	caagggtgaa	1260
cactatccca	tatcaccagc	tcaccgtt	tcattgc	acggaaactc	ggatgagcat	1320
tcatacg	ggcagaatg	tgaataaagg	ccggataaaa	cttgcgtt	tttttttta	1380
cggctttaa	aaaggccgt	atatccagct	gaacggctg	gttata	cattgagca	1440
ctgactgaaa	tgcctcaaaa	tgttcttac	gatgc	ccatg	ggatata	1500
atccagtgat	tttttctcc	atcatattct	tccttttca	atattattg	agcatttac	1560
agggttatttgc	tctcatgagc	ggatacatat	ttgaatgtat	ttagaaaaat	aaacaatag	1620

-continued

gggtcagtgt	tacaaccaat	taaccaattc	tgatctccctg	ttagtgggg	ttaatgccc	1680		
gaacgaagaa	aggcccaccc	gtgaagggtga	gccagtga	tggttacatt	ttctcttgag	1740		
ggtttagctt	ttcagacgc	gcca	aaaaggt	cgtacgtgaa	atacc	aaat	agttggcc	1800
agccgtctt	tcaccattaa	acttctcaag	cgcttgc	gggg	tca	gca	aacgcgg	1860
cggcgtctt	cg	gc	tc	tc	ac	g	cc	1920
cagatccggg	gtcg	gg	ct	ca	aa	cg	ac	1980
ctcgccgatg	tt	ac	cc	gg	gg	cc	gg	2040
ctggcgcagt	gc	cc	gc	ag	ga	gg	ct	2100
ctccggcc	ac	gg	gt	gg	gg	cc	gg	2160
aatgctcagg	cg	gt	ta	ga	ac	ca	gg	2220
attgcagtgg	gt	cg	cg	ct	aa	tt	cc	2280
gcgggtcact	tc	tt	tt	ct	tt	cc	tt	2340
ctcacccgatc	tc	gt	cc	ca	gg	gg	gg	2400
gccacccgca	cg	gg	tt	ac	cc	gg	cc	2460
caggcttcc	gc	g	at	tc	tt	cc	ct	2520
gcgggcateg	tg	ac	gg	gg	ga	tt	cc	2580
cgtaccagtc	tc	gc	ct	tc	ga	ac	tc	2640
ctggcgaact	tg	tc	cc	at	tt	tt	cc	2700
acgagcgc	ta	ac	gc	gt	gg	gg	cc	2760
cagggcgtcg	ct	ga	ac	gc	ct	gg	cc	2820
gcccgc	tt	tc	gg	cc	at	tt	cc	2880
cgtttcagt	tc	gt	tt	at	tc	tt	gg	2940
caggcgcaga	tt	aa	gg	gt	tt	tt	cc	3000
gacaacgcca	at	cg	gg	gg	tt	tt	cc	3060
ac	cg	ct	gg	cc	tt	cc	tt	3120
gctacccgc	tt	aa	at	gg	tt	cc	tt	3180
aaacggcttc	tc	ga	at	cc	gg	tt	cc	3240
gcta	at	gt	cc	gg	tt	cc	tt	3300
atcatcg	tc	ac	gc	gg	tt	cc	tt	3360
ttgggtttaa	at	tt	gt	tt	tc	tt	cc	3420
gttaactgaa	ac	gc	at	tt	tc	tt	cc	3480
aaacat	tt	taa	tt	aa	tt	cc	tt	3540
aataaca	at	tt	tt	aa	tt	cc	tt	3600
actagttgat	ag	at	gt	tt	cc	tt	cc	3660
tttttttga	c	c	t	t	a	a	a	3698

<210> SEQ ID NO 5
 <211> LENGTH: 3610
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Expression vector pPRO430(CloDF13)

<400> SEQUENCE: 5

cccgaggc aggtgcgcgtc cggacacata caaagttacc cacagattcc gtggataagc
agggactaa catgtgaggc aaaacagcag ggccgcgcg gtggcttt tccataggct
ccggccctctt gccagagttc acataaacag acgttttcc ggtgcatacg tggggccgt
gaggctcaac catgaatctg acagtaacggg cggaaacccga caggacttaa agatccccac
cgttccggc gggctcgctcc ctcttgcgtc ctcttgcgtc gaccctgcgcg tttaccggat
acctgttccg ccttttccccc ttacggaaag tggcgttcc ttcatacgact cacacactgg
tatctcggtt cgggttaggt cggtcgctcc aagctgggtt gtaagcaaga actccccgtt
cagcccgact gctgcgcctt atccggtaac tggttacttg agtccaaaccc ggaaaagcac
ggtaaaacgc cactggcagc agccattggg aactgggagt tcgcagagga tttgtttagc
taaacacgcg gttgtcttgc aagtgtgcgc caaagttccg ctacactggg aggacagatt
tgggtgtgt gctctgcgaa agccagttac cacggtaag cagttccca actgacttac
ccttcgtatca aaccacctcc ccagggtgtt ttgcgttca caggcaaaa gattacgcgc
agaaaaaaag gatctcaaga agatccttgc atcacctgca tctaaaaact tttattacgc
cccgccctgc cactcatcgc agtactgttgc taattcatta agcatttcgc cgacatggaa
gccccatcacag acggcatgat gaacotgaat cgccagcggc atcagcacct tggcgttgc
cgtataataat ttgcccatacg tggaaacggg ggcaagaag ttgtccatat tggccacgtt
taaatcaaaa ctggtaaac tcacccaggg attggctgag acgaaaaaca tatttcata
aaacccttta gggaaaatagg ccagggtttc acggtaacac gccacatctt gcaatata
gtgttagaaac tgccggaaat cgtcggttca ttcactccag agcgatgaaa acgtttcagt
ttgtctatgg aaaacgggtt aacaagggtt aacactatcc catatcacca gctcaccgtc
tttcattgca atacggaaact ccggatgagc attcatcagg cgggcaagaa tgtgaataaa
ggccggataaa aacttgtgtt tatttttctt taeggtctt taaaaggccg taatccag
ctgaacgggtc tgggtatagg tacattgagc aactgactga aatgcctca aatgttctt
acgatggcat tgggatataat caacgggtt aatccagtg attttttctt ccatacatatt
cttcctttt caatattattt gaagcattta tcagggtt tggcgttcatca gccgatacat
atttgaatgt atttagaaaa ataaacaaat aggggtcagt gttacaacca attaaccaat
tctgtatcc tggtagtgcg ggttaatgcg cggaaacgaa aaaggcccac ccgtgaaggt
gagccagtga gttggttaca ttttcttgc aggggtttagc ttttcagacg acgccaat
gtcgtagtgcg aaataccca atagttggcc gcagccgtct tggcgttcatca aatgttctca
agcgcttgcg ggggggtcag caaacgcggc gcccgcgtct tggcgttcatca acgcccac
tccggcgcgc gcaactgcgtc gaaattgcggc gtcagatccg gggcgttcatca aacggacagg
aacagcgcgc ggcgttccat catattacgc agtgcgcggc tggtagtgcg tggcgttcatca
tgcagcgcgc cgggttgcgtc cgcctgcaga ccctggcgcga gtggccgcaga gacgggtgcg
ctcagggtgcg ccagcgcgc gttcaggaaa gactccgcga gggtaaaat tggcgttcatca
cgttcacgcgca acggcgggag ctgcagacgc agaatgtcga ggcggtagaa caggtgcgcg
cgaaaacggc cctgttgcgtc atcccttcc agattgcgtc gggcgttcatca aatcgcgcg
2100

-continued

acgtcttaccg	gaaccgggtt	atgaccacccg	acgcgggtca	cttctttctc	ttccagcaca	2220	
cgcagcagac	gggttgcaa	tggcagcggc	atctcaccga	tctcgccag	gaacaagggt	2280	
ccgcccgtggg	caatccaaa	caaaccagca	cggccaccgc	gacggctacc	cgtgaatgcg	2340	
ccctcttcgt	agccaaacag	ctcagcttcc	agcaggctt	ccgcgattgc	accgcaatta	2400	
actgccacaa	acggatgaga	tttcttacc	tggcgggcat	cgtgacgggc	gaaatactca	2460	
cgtggattg	cttgcgcagc	cagttctta	ccgttaccag	tctcgccctc	gatcagaaca	2520	
gccgcgctgc	tacgatgcata	cagcagaatg	gtctggcgaa	cttgcgtccat	ttgagggctt	2580	
tggcccaagca	tatcacccag	gacataacgg	gtacgcagcg	cattacgcgt	cgcacatgtgg	2640	
gtgttgtggc	gcaggctcat	tctggatcatg	ttcaggggct	cgctgaacgc	ctgacgcacc	2700	
gttgcgcgc	tgttagataaa	gatgcccgtc	atgcccgtt	cttgcggccaa	gtccgtgatc	2760	
agacccgcac	caaccacacg	ctcggtacc	ttcgcttca	gttcgttgat	ctggcoacgt	2820	
gcacatccct	cggtaatgt	gctgcgttgg	ttcaggcgca	gattaaagg	ctttgaaac	2880	
gcgaccageg	caggatcg	ttctcggt	gtgacaacgc	caatcgagg	ggtcagttt	2940	
cctgccttcg	ccagcgcctg	caagacatcg	taaccgctcg	gtttaatcg	gatcacccgc	3000	
acggacagac	gggatttcag	gtaggccac	ttgctacc	ctgcgataat	ggcgtaacaa	3060	
cgctcggtgg	ccagctttt	gcaaatgt	gtaacggctt	tctcgaaacc	cagctgaatc	3120	
ggagtgtatgt	tcgcagg	atcaaactcc	aggctaatgt	cgcgaaaca	ctcgacaga	3180	
cggtgtacgc	taacggtca	aataactgg	ttatcatcg	tcaaacgcgg	ttgggtgtgc	3240	
atgggtgata	cctctgtt	agaaaccgaa	tattgggtt	aaacttgg	cataattgtt	3300	
gcaatgaaac	gcgggtgaaac	attgcctgaa	acgttaact	aaacgcata	ttgcggatta	3360	
gttcatgact	ttatctctaa	caaattgaa	ttaaacat	tttttatta	aggcaattgt	3420	
ggcacacccc	ttgtttgtc	tttatcaac	caaataacaa	gttgataaca	aaagcttgg	3480	
aggaaaacat	agagaccggt	ctctctcgag	taactagtt	atagagatca	agcctaacc	3540	
aactaagacc	ccgcacca	aagg	ttccggg	gttttttt	gacctaaaa	acataaccg	3600
ggagcagaca						3610	

<210> SEQ ID NO 6						
<211> LENGTH: 3255						
<212> TYPE: DNA						
<213> ORGANISM: Artificial Sequence						
<220> FEATURE:						
<223> OTHER INFORMATION: Expression vector pBAD240						
<400> SEQUENCE: 6						
ggccttctt	cggttagaaac	cccgagagca	ggtatcaaag	gatcttctt	agatcc	60
tttctgcgcg	taatctgct	cttgcaaaca	aaaaaaccac	cgctacc	agg	120
ttgcccgtc	aagagctacc	aactctttt	ccgaggtaac	tggcttc	agagcgc	180
taccaaatac	tgttcttca	gtgttagcgt	agtttagcc	ccacttca	aaactctgt	240
cacccctac	atacctcgct	ctgctaattcc	tgttacc	ggctgtgc	agtggcgata	300
agtcgtgtct	tacgggtt	gactcaagac	gatagtacc	ggataaggcg	cagcggtcg	360
gctgaacggg	gggttgcgtc	acacagccca	gcttggagcg	aacgacccat	accgaact	420
gataacctaca	gcgtgagcta	tgagaaagcg	ccacgcttcc	cgaagg	gaga	480

-continued

-continued

cggcagcagg ggatcatttt	gcttcagc catactttc	atactccgc cattcagaga	2820
agaaaaccaat tgcacatatt	gcatcagaca ttgccgtctc	tgcgtctttt actggcttt	2880
ctcgctaacc aaaccggtaa	cccgcttat taaaaggatt	ctgtaacaaa gcgggaccaa	2940
agccatgaca aaaacgcgta	acaaaagtgt ctataatcac	ggcagaaaag tccacattga	3000
ttatggcac ggcgtcacac	tttgctatgc catagcattt	ttatccataa gattagcgga	3060
tcctaccta	cgcttttat cgcaactctc	tactgtttct ccataccgt tttttggc	3120
tagcaggagg taaaaaaaaat	gtgagaccgg ttcggctca	gataactagt tgatcggtca	3180
gtttcacctg atttacgtaa	aaaccgcgtt cggcgggtt	ttgctttgg aggggcagaa	3240
agatgaatga ctgtc			3255

<210> SEQ ID NO 7
 <211> LENGTH: 5802
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Expression vector pPRO44

<400> SEQUENCE: 7

atcgattttag	cttttcagac	gacgcacaaa	ggtcgtacgt	gaaataccca	aatagttggc	60
cgcagccgtc	ttgtcacat	taaacttctc	aagcgcttgc	tgccgggtca	gcaaacgcgg	120
agccggcgtc	tttgcgcctc	cacgcgcag	ctccggcagc	agcaactgca	tgaattgcgg	180
agtcaagatcc	ggggtcggct	caacggacag	gaacagegac	aggcgttcca	tcatattacg	240
cagtcgcgg	atgttacccg	gccagtcata	atgcagcagc	accgtttcgc	tcgcctgcag	300
accctggcgc	agtgcgcag	agaacgggtgc	gctcagggtc	gccagegaga	ctttcaggaa	360
agactccgc	agcggtaaaa	tgtcggegac	acgttcacgc	aacgggggaa	gctgcagacg	420
cagaatgctc	aggcggtaga	acaggtcgcg	acgaaaacgg	ccctgttgca	tatccttcc	480
cagattgcag	tgggtcgegc	taatcaegc	cacgtctacc	gaaacccgg	gtgaccacc	540
gacgcgggtc	acttcttct	cttccagcac	acgcagcaga	cgggttgca	atggcagcgg	600
catctcaccc	atctcgteca	ggaacaagg	gcccgggtgg	gcaatttca	acaaaccacg	660
acggccaccc	cgacggctac	ccgtgaatgc	gccttctcg	tagccaaaca	gctcagctc	720
cagcaggctt	tccgcgattt	caccgcaatt	aactgccaca	aacggatgag	atttcttacc	780
ctggcgggca	tcgtgacggg	cgaaataactc	acgatggatt	gcttgcgcag	ccagttcctt	840
acccgtacca	gtctcgctt	cgatcagaac	agccgcgcgt	ctacgtcat	acagcagaat	900
ggctggcga	acttgcctca	tttgagggt	ttggccca	atatcacca	ggacataacg	960
ggtacgcgc	gcattacgcg	tcgcategtg	ggtgttgcgg	cgcagggtca	ttctggtcat	1020
gtccaggcgc	tcgtgacacg	cctgacgcac	cgttgcgcgc	ctgtagataa	agatgcgcgt	1080
catgcccgt	tcttcggcca	agtccgtgat	cagaccgcac	ccaaccacag	cctcggtacc	1140
gttcgttcc	agttcggtga	tctggccacg	tgcacatcc	tcggtaatgt	agctggttg	1200
gtccaggcgc	agattaaagg	tctttgaaa	cgcgaccacg	gcagggtatcg	tttcgttgcgt	1260
ggtacacgc	ccaatcgagg	aggtcagttt	gcctgcctc	gccagcgcct	gcaagacatc	1320
gtaaccgc	ggcttaatca	ggatcaccgg	cacggacaga	cgggatttca	ggtaggcacc	1380
attgctaccc	gctgcgataa	tggcgtaaca	acgctcgatg	gccagtttt	tgcgaatgta	1440

-continued

ggtaacggct ttctcgaaac ccagctgaat cggagtgtat ttcggccaggat gatcaaactc	1500
caggctaatg tcgcggaaaca actcgaacag acgggtgacg ctaacggtcc aaataactgg	1560
tttatcatcg ttcaaacgcg gtgggtgtgc catggtaat acctcctgtt aagaaaccga	1620
atattgggtt taaaacttgtt tcataattgt tgcaatgaaa cgcggtgaaa cattgctga	1680
aacgttaact gaaacgcata tttgcggatt agttcatgac tttatctcta acaaattgaa	1740
attaaacatt taatttatt aaggcaattt tggcacaccc cttgcttgc ctttatcaac	1800
gcaaataaca agttgataac aagctagcga attcgcgatc ggtacccggg gatcctctag	1860
agtcgacactg caggcatgca agttggctg ttttggcgga tgagagaaga ttttcagcct	1920
gatacagatt aaatcagaac gcagaagcgg tctgataaaa cagaatttgc ctggggcag	1980
tagcgcggtg gtcccacactg accccatgcc gaactcagaa gtgaaacgcc gtagcgcgcga	2040
tggtagtgtg gggtctcccc atgcgagagt agggaactgc caggcatcaa ataaaacgaa	2100
aggctcagtc gaaagactgg gccttcgtt ttatctgtt tttgtcggtg aacgctctcc	2160
tgagtaggac aaatccgcgg ggagcggatt tgaacgttgc gaagcaacgg cccggagggt	2220
ggcgggcagg acgcccccca taaaactgcca ggcataat taagcagaag gccatctga	2280
cggatggcct ttttgcgtt ctacaaactc ttttgtttat ttttctaaat acatccaaat	2340
atgtatccgc tcatgagaca ataaccctga taaatgttc aataatattt aaaaaggaag	2400
agtatgagta ttcaacacatt cctgtgtcc cttatccct ttttgcggc atttgcctt	2460
cctgttttg ctcacccaga aacgctgggtt aaagtaaaag atgctgaaga tcagttgggt	2520
gcagcaact attaactggc gaactactta ctctagctc cggcaacaa ttaatagact	2580
ggatggaggc ggataaaagtt gcaggaccac ttctgcgcgc ggccttccg gctggctgg	2640
ttattgctga taaatctgga gcccgtgacg gtgggtctcg cggatcattt gcagcactgg	2700
ggccagatgg taagccctcc cgtatcgtag ttatctacac gacggggagt caggcaacta	2760
tggatgaacg aaatagacag atcgctgaga taggtgcctc actgattaag cattggtaac	2820
tgtcagacca agtttactca tatatacttt agattgattt acgcgcctg tagcggcgca	2880
ttaagcgcgg cgggtgtggt ggtaacgcgc acgtgtaccc ctacacttgc cagcgcctta	2940
gcgcgcgcctc ctttcgcctt cttccctcc tttctegcca cgttcgcggg ctttccccgt	3000
caagctctaa atcgggggct ccctttaggg ttccgattt gtgttttacg gcacctcgac	3060
ccccaaaaac ttgatttggg tgatggttca cgttagtgggc catgcctctg atagacggtt	3120
tttcgcctt tgacgttgaa gtccacgttc ttaatagtg gactttgtt ccaaactgga	3180
acaacactca accctatctc gggctattct tttgatttt aaggatttt gccgatttcg	3240
gcctattggc taaaaaatga gctgatttaa caaaaattta acgcgaattt taacaaaata	3300
ttaacgttta caatttaaaa ggatcttaggtt gaagatccctt tttgataatc tcatgaccaa	3360
aatcccttaa cgtgagttt cgttccactg acgttcacac cccgtagaaa agatcaaagg	3420
atcttcttgc gatcttttt ttgtcgctg aatctgttc ttgcaaaaca aaaaacacc	3480
gctaccacgcg gtggttgtt tgccggatca agagctacca actcttttc cgaaggtaac	3540
tggcttcagc agagcgcaga taccaaatac tgccttcata gtgtagccgt agttagggca	3600
ccacttcaag aactctgttag caccgcctac atacctcgct ctgctaattcc tgttaccagt	3660
caggcatttga agaaggcacac ggtcacactg cttccggtag tcaataaacc ggtaaaccag	3720

-continued

```
<210> SEQ_ID NO 8
<211> LENGTH: 5791
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
```

-continued

<223> OTHER INFORMATION: Expression vector pPRO45

<400> SEQUENCE: 8

atcgatttag	ctttcagac	gacgccaaaa	ggtcgtacgt	gaaataccca	aatagttggc	60
cgcagccgtc	ttgtcaccat	taaacttctc	aagcgcttgc	tgcggggtca	gcaaacgcgg	120
agccggcgtc	tttgcgtct	cacgcgcagc	ctccggcagc	agcaactgca	tgaattgcgg	180
agtcaagatcc	ggggtcggct	caacggacag	gaacagcgcc	aggcgttcca	tcatattacg	240
cagctcgccg	atgttacccg	gccagtcata	atgcagcagc	accgtttcgc	tgcctgcag	300
accctggcgc	agtgccgcag	agaacgggtgc	gctcagggtct	gccagcgaga	ctttcaggaa	360
agactccgcc	agcggtaaaa	tgtcggegac	acgttacgc	aacgggggaa	gctgcagacg	420
cagaatgctc	aggcggtaga	acaggtcgcg	acgaaaacgg	ccctgttgca	tatccttcc	480
cagattgcag	tgggtcgccc	taatcacgcg	cacgttacc	ggaaccgggtt	gatgaccacc	540
gacgccccgtc	acttctttct	cttccagcac	acgcagcaga	cgggtttgca	atggcagcgg	600
catctcaccg	atctcgctca	ggaacaagggt	gcccgggtgg	gcaatttcaa	acaaaccaggc	660
acggccaccc	cgacggctac	ccgtgaatgc	gccctttcg	tagccaaaca	gctcagcttc	720
cagcaggctt	tccgcgattt	caccgcaatt	aactgcaca	aacggatgag	atttcttacc	780
ctggcgggca	tcgtgacggg	cgaaataactc	acgatggatt	gcttgcgcag	ccagttcctt	840
accegtacca	gtctcgctt	cgatcagaac	agccgcgcgt	ctacgtcat	acagcagaat	900
ggtctggcga	acttgcctca	tttgagggtct	ttggcccagc	atatcaccca	ggacataacg	960
ggtacgcgc	gcattacgcg	tcgcategtg	ggtgttgcgg	cgcaggctca	ttctggtcat	1020
gtccagggeg	tcgctgaacg	cctgacgcac	cgttgcgcg	ctgtagataa	agatgccgt	1080
catgcccgt	tcttcggcca	agtccgtat	cagaccgc	ccaaccacag	cctcggtacc	1140
gttcgcgttc	agttcggtga	tctggccacg	tgcacatccc	tcggtaatgt	agctgcgtt	1200
gtccaggcgc	agattaaagg	tcttttggaa	cgcgaccagc	gcagggtatcg	tttcctggta	1260
ggtgacaaacg	ccaatcgagg	aggtcagttt	gcctgccttc	gccagegcct	gcaagacatc	1320
gtaaccgc	ggttaatca	ggatcaccgg	cacggacaga	cgggattca	ggtaggcacc	1380
attgctaccc	gctgcgataa	tggcgtaca	acgctcggt	gccagcttt	tgcgaatgt	1440
ggtaacggct	ttctcgaaac	ccagctgat	cggagtgtat	ttcgcggatgt	gatcaaactc	1500
caggctaatt	tcgcggaaaca	actcgaacag	acgggtgacg	ctaacggtcc	aaataactgg	1560
tttatcatcg	ttcaaacgcg	gtgggtgtc	catggtaat	acctcctgtt	aagaaaccga	1620
atattgggtt	taaacttgc	tgcataattgt	tgcaatggaa	cgcgggtaaa	cattgcctga	1680
aacgttaact	gaaacgcata	tttgcggatt	agttcatgac	tttatctcta	acaaattgaa	1740
attaaacatt	taattttatt	aaggcaattt	tggcacaccc	cttgctttgt	ctttatcaac	1800
gcaataaca	agttgataac	aagctagcga	attcgagctc	ggtacccggg	gatcctctag	1860
agtgcaccc	caggcatgca	agttgggtt	ttttggcgga	tgagagaaga	ttttcagcct	1920
gatacagatt	aaatcagaac	gcagaagcgg	tctgataaaa	cagaatttgc	ctggcggcag	1980
tagcgcgg	gtccccaccc	accccatgca	gaactcagaa	gtgaaacgcc	gtagcgcgc	2040
tggtagtgc	gggtctccc	atgcgagagt	agggactgc	caggcatca	ataaaacgaa	2100
aggctcagtc	gaaagactgg	gccttcgtt	ttatctgtt	tttgcgggt	aacgcctcc	2160

-continued

tgagtaggac	aaatccgcgg	ggagcggatt	tgaacgttgc	gaagcaacgg	cccgagggt	2220
ggcgggcagg	acgcccgcga	taaaactgcca	ggcatcaa	taagcagaag	gccatctga	2280
cggatggcct	tttgcgttt	ctacaaactc	ttttgtttat	ttttctaaat	acattcaa	2340
atgtatccgc	tcatgagaca	ataaccctga	taaatgcttc	aataatattg	aaaaaggaag	2400
agtatgagta	ttcaacattt	ccgtgtcgcc	cttattccct	tttttgcggc	atttgcctt	2460
cctgttttg	ctcacccaga	aacgctggtg	aaagtaaaag	atgctgaaga	tcagttgggt	2520
gcagcaact	attaactggc	gaactactta	ctctagcttc	ccggcaacaa	ttaatagact	2580
ggatggaggc	ggataaaagtt	gcaggaccac	ttctgcgc	ggcccttccg	gctggctgg	2640
ttattgctga	taaatctgga	gcgggtgagc	gtgggtctcg	cggtatcatt	gcagcactgg	2700
ggccagatgg	taagccctcc	cgtatcgtag	ttatctacac	gacggggagt	caggcaacta	2760
tggatgaacg	aaatagacag	atcgctgaga	taggtgcctc	actgattaag	cattggtaac	2820
tgtcagacca	agtttactca	tatatacttt	agattgattt	acgcgcctg	tagcggcgca	2880
ttaagcgcgg	cgggtgtgg	ggttacgogc	agcgtgaccc	ctacacttc	cagcgcctca	2940
gcgcgcgc	ctttcgctt	cttcccttc	tttctcgcca	cgttgcgcgg	cttccccgt	3000
caagctctaa	atcgggggct	cccttttaggg	ttccgattta	gtgctttacg	gcacctcgac	3060
ccccaaaaac	ttgattttggg	tgtatggtca	cgtagtgggc	catcgcctg	atagacggtt	3120
tttcgcctt	tgacgttgg	gtccacgtt	tttaatagtg	gactcttgg	ccaaactgg	3180
acaacactca	accatatc	gggttattct	tttgatttat	aaggatattt	gccgatttc	3240
gcctatttgt	taaaaaatga	gctgatttaa	caaaaat	acgcgaattt	taacaaaata	3300
ttaacgttta	caataaaaa	ggatcttagg	gaagatcctt	tttgataatc	tcatgaccaa	3360
aatcccttaa	cgtgatgttt	cgttccact	agcgtcagac	cccgtagaaa	agatcaaagg	3420
atcttcttga	gatcctttt	ttctgcgcgt	aatctgctc	ttgcaacaa	aaaaaccacc	3480
gctaccageg	gtgggttgg	tgccggatca	agagctacca	actcttttc	cgaaggtaac	3540
tggcttcagc	agagcgcaga	taccaaatac	tgtccttcta	gtgtagccgt	agttaggcca	3600
ccacttcaag	aactctgtag	caccgcctac	atacctcgct	ctgctaatcc	tgttaccagt	3660
caggcattt	agaagcacac	ggtcacactg	cttccggtag	tcaataaacc	ggtaaaccag	3720
caatagacat	aagcggctat	ttaacgacc	tgcctgaac	cgacgaccgg	gtcgaattt	3780
cttcgaatt	tctgccattc	atccgctt	tatcactt	tcaggcgt	caaccaggcg	3840
ttaaggcga	ccaataactg	ccttaaaaaa	attacgcccc	gccctgcac	tcatcgagt	3900
actgttgtaa	tccattha	attctgcga	catggaagcc	atcacaac	gcatgatgaa	3960
cctgaatcgc	cagcggc	atcacctt	cgcccttgcgt	ataatattt	cccatggtga	4020
aaacggggc	gaagaagtt	tccatattgg	ccacgtttaa	atcaaactg	gtgaaactca	4080
cccaggatt	ggctgagacg	aaaaacat	tctcaataaa	ccctttaggg	aaataggcca	4140
gttttcacc	gtaacacgoc	acatcttgc	aatatatgt	tagaaactgc	cgaaatcg	4200
cgtggattc	actccagac	gatgaaaac	tttcagttt	ctcatggaa	acgggtgtaac	4260
aagggtgaac	actatccat	atcaccagct	caccgtctt	cattgcata	cggaattcc	4320
gatgagcatt	catcaggcgg	gcaagaatgt	gaataaaggc	cgataaaac	ttgtgtttat	4380
tttcttac	ggtctttaa	aaggccgtaa	tatccagctg	aacggctgg	ttataggtac	4440

-continued

attgagcaac	tgactgaaat	gcctcaaaat	gttctttacg	atgccatgg	gataatatcaa	4500
cggtggtata	tccagtgatt	ttttctcca	tttagcttc	cttagctcct	gaaaatctcg	4560
ataactcaa	aaatacgc	ccc	ggtagtgc	ttat	ttcattatgtt	4620
ttacgtgccc	atcaacgtct	catttcg	ccc	aaaagtggc	ccagggcttc	4680
cagggacacc	aggat	ttt	tattctcg	ca	gtgatcttc	4740
cgcaaaagtgc	gtcgggtgat	gtc	gccaact	tactgattt	gtgtatgt	4800
ggtgctccag	tggcttctgt	ttctatc	g	tgccctcct	gttcagtc	4860
gtgcgttaac	gcaaaagcac	cgccggacat	caactagtg	gc	tgccgaca	4920
tacccacaga	ttccgtggat	aagcagg	gg	cta	acatgt	4980
gcccgtggcg	ttttccata	gg	gtccg	ccc	gttcacataa	5040
ttccggtgca	tctgtggag	ccgt	gagg	gt	ca	5100
ccgacaggac	ttaaagatcc	ccaccgtt	cc	gggggtc	ctccctt	5160
ttccgacc	ccgttacc	ggata	cc	tc	tc	5220
gctttctat	agtcacaca	ctgg	ttatc	gg	tcgtgt	5280
ggctgtaa	agaactccc	cg	ttcagcc	g	actgtcg	5340
ctttagtcca	acc	cg	gg	aa	ccat	5400
gagttcgcag	aggat	ttt	tt	taa	aca	5460
ccggctacac	tg	gaagg	ac	atc	tc	5520
taagcagt	cc	caact	gt	tc	aaacc	5580
tttacagg	gc	aa	agatt	ac	atc	5640
gtctat	t	cat	gag	cc	gca	5700
atata	agg	gc	gc	cc	atc	5760
tagaggat	tc	at	gt	gt	ttt	5791

<210> SEQ ID NO 9
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPRO_REV_SpeI primer
<400> SEQUENCE: 9
gactgactag ttgatgtccg gcggt 25

<210> SEQ ID NO 10
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPRO_FOR_AatII primer
<400> SEQUENCE: 10
tcagctgacg tctat

ttgttgc 28
t catgagcc
<210> SEQ ID NO 11
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

```

<223> OTHER INFORMATION: RSF1030_FOR_SpeI primer
<400> SEQUENCE: 11
gcaaaagcac cgccggacat caactagtct tccgcttcct cgctca 46

<210> SEQ ID NO 12
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RSF1030_REV_AatII primer

<400> SEQUENCE: 12
gccacttcgg gctcatgagc aaatagacgt caacggaata gctgttcgtt ga 52

<210> SEQ ID NO 13
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CloDF13_FOR_SpeI primer

<400> SEQUENCE: 13
gcaaaagcac cgccggacat caactagtgc gctgcccaca cataca 46

<210> SEQ ID NO 14
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CloDF13_REV_AatII primer

<400> SEQUENCE: 14
gccacttcgg gctcatgagc aaatagacgt cgatcaaagg atcttcttga gatcct 56

<210> SEQ ID NO 15
<211> LENGTH: 5304
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Dual-promoter vector, pSOL

<400> SEQUENCE: 15
ggcctttctt cggtagaagt ctcccccaag aggcaggatct caaaggatct tcttgagatc 60
ctttttttctt gcgcgtaatc tgctgcttc aaacaaaaaa accaccgcta ccagcggtgg 120
tttggggcc ggtcaagag ctaccaactc ttttccgag gtaactggct tcagcagagc 180
gcagatacca aatactgttc ttcttagtta gcccgtatca ggccaccact tcaagaactc 240
tgttagcaccg cctacataacc tgcgtctgtc aatcctgtta ccagtggctg ctgcccgtgg 300
cgataagtctg tgcgtttaccg ggttggactc aagacgatag ttaccggata aggcgcagcg 360
gtcgggctga acgggggggtt cgtgcacaca gcccagctt gaggcaacga cctacaccga 420
actgagatac ctacagcgatc agatgttgcg aagcgccacg cttcccgaaag ggagaaaggc 480
ggacaggtat ccggtaagcg gcagggtcgg aacaggagag cgcacgagg agcttccagg 540
ggaaacgccc tggtatctt atagtcgttgcg cgggtttcgc cacctctgac ttgagcatcg 600
atttttgtga tgctcgatcg gggggccggag cctatggaaa aacgcccacca acgcagaaag 660
gcccacccga aggtgagccg ggtgattaca tttggccctt catcagaggt tttcacccgtc 720

```

-continued

atcacccgaaa	cgcgcgaggc	agctgcggta	aagctcatca	gcgtggcgt	gaagcgattc	780
acagatgtct	gcctgttcat	ccgcgtccag	ctcggtttagt	ttctccagaa	gcgttaatgt	840
ctggcttctg	ataaaagcggg	ccatgttaag	ggcggttttt	tcctgtttgg	tcataacctgc	900
ttagaaaaac	tcatcgagca	tcaaataatgaaa	ttgcaattta	ttcatatcag	gattatcaat	960
accatatttt	tgaaaaagcc	gtttctgttaa	tgaaggagaa	aactcaccga	ggcagttcca	1020
taggatggca	agatcctggt	atcggtctgc	gattccgact	cgtccaacat	caatacaacc	1080
tattaaatttc	ccctcgtcaa	aaataaggtt	atcaagtgag	aaatcaccat	gagtgcacgac	1140
tgaatccggt	gagaatggca	aaagtttatg	catttcttgc	cagacttgg	caacaggcca	1200
gccattacgc	tcgtcatcaa	aatcactcg	atcaacccaa	ccgttattca	ttcgtgattg	1260
cgcctgagcg	aggcgaaata	cgcgatcgct	gttaaaagga	caattacaaa	caggaatcg	1320
gtcaaccgg	cgcaggaaca	ctgcccagcg	atcaacaata	tttccacctg	aatcaggata	1380
ttcttctaat	accttggaaacg	ctgttttcc	ggggatcgca	gtgggtgagta	accatgcattc	1440
atcaggagta	cggataaaat	gcttgatgg	cggaagtggc	ataaattccg	tcagecagtt	1500
tagtctgacc	atctcatctg	taacatcatt	ggcaacgcta	ccttgcatt	gtttcagaaa	1560
caactctggc	gcacatcggt	tcccatacaa	gcgatagatt	gtcgcacctg	attgeccgac	1620
attatcgca	gcccatttat	accatataaa	atcagcatcc	atgttggaa	ttaatcgccg	1680
cctcgacgtt	tcccggttga	tatggctcat	agctcctgaa	aatctcgata	actcaaaaaaa	1740
tacgccccgt	agtgtatctt	tttcattatg	gtgaaagttg	gaaccttcta	cgtgcgcattc	1800
aagaagacgg	tcaaaagcct	ccggtcggag	gccgggagag	tgttcacccg	caaacaacag	1860
ataaaaacaaa	aggccccagtc	ttccgactga	gcctttgtt	ttatggatg	tctggcagtt	1920
cccgagacgt	tatgacaact	tgacggctac	atcattca	ttttcttcac	aaccggcacg	1980
gaactcgctc	gggctggccc	cggtgcattt	tttaataacc	cgcgagaaat	agagttgatc	2040
gtcaaaacca	acattgcac	cgacgggtgg	gataggcattc	cggggtggc	tcaaaagcag	2100
cttcgcctgg	ctgatacg	tttcctcg	ccagcttaag	acgctaattcc	ctaactgctg	2160
gcggaaaaga	tgtgacagac	gegacggcga	caagcaaaca	tgctgtcg	cgctggcgat	2220
atcaaaattt	ctgtctgcca	ggtgatcgct	gatgtactga	caagcctcgc	gtacccgatt	2280
atccatcggt	ggatggagcg	actcgtaat	cgcttccatg	cgccgcagta	acaattgctc	2340
aagcagattt	atcgccagca	gctccgaaata	gcgccttcc	cattgcgg	cgtaatgtat	2400
ttgccccaaac	aggtcgctga	aatgcggctg	gtgcgcattca	tccggggc	agaaccccgt	2460
attggcaaat	attgacggcc	agttaaagcca	ttcatgc	taggcgcgc	gacgaaagta	2520
aacccactgg	tgataccatt	cgcgagccct	cgatgcacga	ccgtatgtat	aatctctcc	2580
tggcgccgg	agcggaaat	cacccggc	gcaaaacaat	tctcgccct	gattttccac	2640
cacccctgt	ccgcaatgg	tgagattgag	aatataacct	ttcattccca	gcggcggtc	2700
gataaaaaaa	tcgagataac	cggtggcctc	aatcgccgtt	aaacccgcca	ccagatgggc	2760
atcaaacgag	tatcccgca	gcaggggatc	attttgcgt	tcagccata	ttttcatact	2820
cccgccattc	agagaagaaa	ccaattgtcc	atattgcattc	agacattgcc	gtctctcg	2880
cttttactgg	cttttctcg	taaccaaacc	ggtaaccccg	cttattaaaa	gcattctgt	2940
acaaagcg	accaaagcca	tgacaaaaac	cgtaacaaa	agtgtctata	atcacggcag	3000

-continued

aaaagtccac	attgattatt	tgcacggcgt	cacacttgc	tatgccatag	cattttatc	3060
cataagatta	gcggatccta	cctgacgctt	tttatacgaa	ctctctactg	tttctccata	3120
cccgaaaa	tgggctagca	ggaggtaaaa	aaaatgtgag	accggctcg	gtctagatcg	3180
gtcagttca	cctgatttac	gtaaaaaccc	gttcggccg	gtttttgctt	ttggaggggc	3240
agaaagatga	atgactgtct	ctcctgttag	tgagggtaa	tgcccgaaac	gaagaaaggc	3300
ccacccgtga	aggtgagcca	gtgagtttgt	tacatttct	cttgagggtt	tagtttca	3360
gacgacgcca	aaaggctcgta	cgtgaaatac	ccaaatagtt	ggccgcagcc	gtcttgcac	3420
cattaaactt	ctcaagcgct	tgctgcgggg	tcagcaaacg	cgagccggc	gtcttgcgc	3480
tctcacgcgc	cagctccggc	agcagcaact	gcatgaattt	cgagtcaga	tccggggtcg	3540
gctcaacgga	caggaacagc	gccagggcgtt	ccatcatatt	acgcagctcg	cgatgttac	3600
ccggccagtc	ataatgcgc	agoaccgttt	cgctcgctcg	cagaccctgg	cgcagtgcgc	3660
cagagaacgg	tgcgctcagg	gctgccagcg	agactttcag	gaaagactcc	gccagggta	3720
aaatgtcgcc	gacacgttca	cgcaacggcg	ggagctgcag	acgcagaatg	ctcaggcggt	3780
agaacaggc	gcgacgaaaa	cggccctgtt	gcatatccctc	ttccagattt	cagtgggtcg	3840
cgctaatac	gcccacgtct	accggaaacc	gttgcatttgc	accgaecgg	gtcaacttctt	3900
tctttccag	cacacgcgc	agacgggttt	gcaatggcag	cgccatctca	ccgatctcg	3960
ccaggaacaa	ggtgcgcgc	tggcaattt	caaacaacc	agcacggcca	ccgcgcacggc	4020
tacccgtgaa	tgegcctct	tegtageccaa	acagctcagc	ttccagcagg	cttccgcga	4080
ttgcaccgca	attaactgcc	acaaacggat	gagatttctt	accctggcg	gcategtgac	4140
ggggeaaata	ctcacgatgg	attgcttgcg	cagccagttc	cttaccctgt	ccagtctcg	4200
cttgcgtcag	aacagccgc	ctgcttaegtg	catacagcag	aatggcttgg	cgaacttgc	4260
ccatttgggg	gctttggccc	agcatatcac	ccaggacata	acgggtacgc	agcgcattac	4320
gcgtcgatc	gtgggtgttgc	tggcgccagg	tcatttctgtt	catgtccagg	gcgtcgatcg	4380
acgcctga	caccgttgc	gctgttgc	taaagatgcc	cgtcatgccc	gttcttcgg	4440
ccaagtccgt	gatcagaccc	gcaccaacca	cagccctgg	accgttcgt	ttcagtctgt	4500
tgatctggcc	acgtgcac	tcctcgtaa	tgtatgcgt	ttggccagg	cgcagattaa	4560
aggcttttgc	aaacgcgacc	agcgcaggaa	tcgtttctgt	gtaggtaca	acgccaatcg	4620
aggaggtc	tttgcctgc	ttcgccagcg	cctgcaagac	atcgtaacc	ctcggtttaa	4680
tcaggatcac	ccgcacggac	agacgggatt	tcaaggtaggc	accattgcta	cccgctgcga	4740
taatggcg	acaacgcgtc	ttggccagct	ttttgcgtat	gtaggtaca	gttttcgt	4800
aacccagctg	aatcggagtg	atgttcgc	ggtgtatcaaa	ctccaggct	atgtcgccg	4860
acaactcgaa	cagacgggtg	acgcttaacgg	tccaaataac	tggtttatca	tcgttcaa	4920
gcgggtgg	tgccatggtg	aatacctct	gttaagaaac	cgaatattgg	gtttaactt	4980
gtttcataat	tgttgcata	aaacgcgttgc	aaacattgc	tgaaacgtt	actgaaacgc	5040
atatttgcgg	attagttcat	gactttatct	ctaacaatt	gaaatcaa	atttat	5100
attaaggc	ttgtggcaca	ccccttgctt	tgtctttatc	aacgcaata	acaagtgtat	5160
aacaaaagct	taggagggaa	acatagagac	cggctctct	cgagtaacta	gttgcata	5220
atcaaggc	aacgaactaa	gaccccgca	ccgaaaggc	cgggggtttt	ttttgac	5280

-continued

aaaaaacataa ccgaggagca gaca 5304

<210> SEQ ID NO 16
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Infliximab chimeric (murine variable domain, human constant domain) heavy chain

<400> SEQUENCE: 16

Met Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15

Gly Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Ile Phe Ser Asn
 20 25 30

His Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp
 35 40 45

Val Ala Glu Ile Arg Ser Lys Ser Ile Asn Ser Ala Thr His Tyr Ala
 50 55 60

Glu Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser
 65 70 75 80

Ala Val Tyr Leu Gln Met Thr Asp Leu Arg Thr Glu Asp Thr Gly Val
 85 90 95

Tyr Tyr Cys Ser Arg Asn Tyr Tyr Gly Ser Thr Tyr Asp Tyr Trp Gly
 100 105 110

Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
 115 120 125

Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala
 130 135 140

Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
 145 150 155 160

Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
 165 170 175

Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
 180 185 190

Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His
 195 200 205

Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
 210 215 220

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
 225 230 235 240

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 245 250 255

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
 260 265 270

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 275 280 285

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
 290 295 300

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 305 310 315 320

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
 325 330 335

-continued

Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 340 345 350
 Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser
 355 360 365
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 370 375 380
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 385 390 395 400
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
 405 410 415
 Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
 420 425 430
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 435 440 445
 Pro Gly Lys
 450

<210> SEQ ID NO 17
 <211> LENGTH: 215
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Infliximab chimeric (murine variable domain,
 human constant domain) light chain

<400> SEQUENCE: 17

Met Asp Ile Leu Leu Thr Gln Ser Pro Ala Ile Leu Ser Val Ser Pro
 1 5 10 15
 Gly Glu Arg Val Ser Phe Ser Cys Arg Ala Ser Gln Phe Val Gly Ser
 20 25 30
 Ser Ile His Trp Tyr Gln Gln Arg Thr Asn Gly Ser Pro Arg Leu Leu
 35 40 45
 Ile Lys Tyr Ala Ser Glu Ser Met Ser Gly Ile Pro Ser Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser Ile Asn Thr Val Glu
 65 70 75 80
 Ser Glu Asp Ile Ala Asp Tyr Tyr Cys Gln Gln Ser His Ser Trp Pro
 85 90 95
 Phe Thr Phe Gly Ser Gly Thr Asn Leu Glu Val Lys Arg Thr Val Ala
 100 105 110
 Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
 115 120 125
 Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
 130 135 140
 Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
 145 150 155 160
 Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
 165 170 175
 Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
 180 185 190
 Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
 195 200 205
 Ser Phe Asn Arg Gly Glu Cys
 210 215

-continued

```
<210> SEQ ID NO 18
<211> LENGTH: 4594
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pBAD240-Infliximab_HC expression construct

<400> SEQUENCE: 18
ggctttctt cggtagaaac cccgagagca ggtatcaaag gatcttttg agatctttt 60
tttctgcgcg taatctgctg cttgcaaaca aaaaaaccac cgctaccagc ggtggttgt 120
ttgccccgtc aagagctacc aactctttt ccgaggtaac tggcttcagc agagcgcaga 180
taccaaatac tggcttctta gtgttagccgt agtttaggcac ccacttcaag aactctgtag 240
caccgcctac atacctcgct ctgctaattcc tggtaaccgt ggctgctgcc agtggcgata 300
agtcgtgtct taccgggttg gactcaagac gatagttacc ggataaggcg cagcggtcg 360
gctgaacggg ggggtcgtgc acacagccca gcttggagcg aacgacccat accgaactga 420
gataacctaca gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcgac 480
ggtatccggta aagcggcagg gtcggaaacag gagagcgcac gagggagctt ccagggggaa 540
acgcctggta tctttatagt cctgtcggtt ttcgcccaccc ctgacttggag catcgatttt 600
tgtgatgctc gtcagggggg cggagcgtat ggaaaaacgc cagcaacgc gaaaggccca 660
cccgaaggtg agccaggtga ttacatttg gcccctcatca gaggtttca ccgtcatcac 720
cgaaacgcgc gaggcagctg cggtaaagct catcagcgtg gtcgtgaagc gattcacaga 780
tgtctgcctg ttcatcccgcg tccagctcggt tgagtttctc cagaagcgtt aatgtctggc 840
ttctgataaa gcgggcccattt ttaagggcggtt tttttccctt tttggtcata cctgcttaga 900
aaaactcatc gacatcaag tggaaactgca atttattcat atcaggatttca tcaataccat 960
atttttgaaa aagccgtttc tggatgaaag gagaaaaactc accgaggcgat ttccatagga 1020
tggcaagatc ctggatcggtcg tctgcgttcc cgactcgatcc aacatcaata caacctattt 1080
atttccctc gtcaaaaata aggttatcaa gtgagaaatc accatgagtg acgactgaat 1140
ccgggtgagaa tggcaaaaatc ttatgcattt ctttccagac ttgttcaaca ggccagccat 1200
tacgctcgatc atcaaaaatca ctgcacccaa ccaaaaccgtt attcattcgat gattgcgcct 1260
gagcggagacg aaatacgcga tcgcccgttta aaggacaatt acaaacagga atcgaatgca 1320
accggcgcag gaacactgcc agcgcataa caatatttc acctgaatca ggatattttt 1380
ctaatacctg gaatgctgtt ttccctggaa tcgcgtgtt gagtaaccat gcatcatcag 1440
gagtaacggat aaaatgcttg atggtcggaa gaggcataaa ttccgtcagc cagtttagcc 1500
tgaccatctc atctgttaaca tcattggcaaa cgctacccctt gccatgtttc agaaacaact 1560
ctggcgcatc gggcttccca tacaatcgat agattgtcgcc acctgtatgc ccgcacattat 1620
cgcgagccca ttatataccca tataaatcgat catccatgtt ggaatttaat cgccggctcg 1680
agcaagacgt ttcccggttga atatggctca tagctcctga aaatctcgat aactcaaaaa 1740
atacgcggcgg tagtgatctt atttcattat ggtgaaagtt ggaaccttcc acgtgcgcgt 1800
caagtcaaaa gcctccgggg gagagtgttc accgcacaaac aacagataaa acaaaggcc 1860
cagtcctcccg actgagccctt ttgttttattt tgatgtctgg cagttccgaa gacgttatga 1920
caacttgacg gctacatcat tcacttttcc ttcacaacccg gcacggaaact cgctcgggct 1980
```

-continued

ggccccggtg	catttttaa	atacccgca	gaaatagagt	tgatcgtaa	aaccaacatt	2040
gcgaccgacg	gtggcgatag	gcatccggg	ggtgctaaa	agcagctcg	cctggctgat	2100
acgttggcc	tcgcgcacg	ttaagacgct	aatccctaac	tgctggcga	aaagatgtga	2160
cagacgcac	ggcgacaacg	aaacatgctg	tgcgacgctg	gcgatatacaa	aattgtgtc	2220
tgccaggtga	tcgctgatgt	actgacaacg	ctcgcttacc	cgattatcca	tcgggtggatg	2280
gagcgactcg	ttaatcgctt	ccatcgccg	cagtaacaat	tgctcaagca	gatttatcg	2340
cagcagctcc	gaatagcgcc	cttcccttg	cccgccgtta	atgatttgcc	caaacaggtc	2400
gctgaaatgc	ggctggtgcg	cttcatccgg	gcgaaagaac	cccgatattgg	caaattattga	2460
cggccagtt	agccattcat	gccagtaggc	gcccggacga	aagtaaaccc	actgggtgata	2520
ccattcgcga	gcctccggat	gacgaccgta	gtgatgaatc	tctcctggcg	ggaacacgcaa	2580
aatatcaccc	ggtcggcaaa	caaatttcg	tccctgat	ttcaccaccc	cctgaccgcg	2640
aatggtgaga	ttgagaatat	aacctttcat	tcccagcggt	cggtcgataa	aaaaatcgag	2700
ataaccgtt	gcctcaatcg	gcgttaaacc	cgccaccaga	tgggcattaa	acgagtatcc	2760
cggcagcagg	ggatcatttt	gchgcttcagc	catactttc	atactccgc	cattcagaga	2820
agaaaccaat	tgtccatatt	gcatcagaca	ttgcccgtctc	tgcgctttt	actggctt	2880
ctcgctaacc	aaaccggtaa	cccccgttat	taaaagcatt	ctgtaacaaa	gcgggaccaa	2940
agccatgaca	aaaacgcgta	acaaaagtgt	ctataatcac	ggcagaaaaag	tccacattga	3000
ttatggcac	ggcgctcacac	tttgctatgc	catagcattt	ttatccataa	gattagcgga	3060
tcctacctga	cgtttttat	cgcaactctc	tactgtttct	ccatacccg	ttttttggc	3120
tagcaggagg	taaaaaaaat	ggaagttaaa	ttagaagaaa	gcccggcgg	tttggtgcaa	3180
cctggcggtt	cgtatgaaat	gagctgcgtc	gcaagcggtt	tcatttttc	caaccactgg	3240
atgaactggg	tgcgccagtc	tccggaaaag	ggtctggaaat	gggttgcgg	gatccgtac	3300
aagagcatca	atagcgcac	gcattatgcc	gagagcgta	aaggccgtt	caccatttct	3360
cgtgacgaca	gcaaaagcgc	tgtgtatctg	caaattgaccg	acttgcgtac	cgaggacacg	3420
ggcggttatt	actgtcccg	caactattac	ggttccacct	acgactactg	ggggccagggt	3480
acgaccctga	cggttagtc	ggcgagcacc	aagggtccga	gcgttttcc	gctggcaccg	3540
agcagcaaaa	gcaccagcg	ttgttccgc	gcactgggtt	gcctggtgaa	agattacttc	3600
ccggaaacccg	ttactgttag	ctggaaacage	ggcgccgtga	cctctggcg	gcacacgttc	3660
ccggcagttc	tgcaaaggcag	cgccctgtac	tccctgtcca	gcgtcgta	cgtgcccgg	3720
agcagcctgg	gtacgcaaac	ctatatttg	aacgtcaatc	acaagccgag	caacacaaa	3780
gtggacaaaa	aagtgcgaaacc	gaaaagtcg	gataaaaccc	atacttgcc	gcccgtcccc	3840
gccccggagc	ttctgggtgg	tccaaggcgtt	tttctgttcc	cgccgaagcc	gaaagacacc	3900
ctgatgatca	gcccaccc	tgaggtgacc	tgtgtggtag	ttgacgttcc	ccacgaagat	3960
ccagagggtca	agtttaactg	gtatgtggat	ggcggtcgaa	ttcacaatgc	aaagaccaag	4020
ccgcgtgaag	aacagtataa	ctctacgtac	cgtgtcgta	gcgttctgac	tgttctgcac	4080
caggattggc	tgaacggcaa	agagtacaag	tgcaaggta	gcaataaagc	gctgeccggt	4140
ccgatcgaga	aaaccatttc	taaggctaa	ggtcagccgc	gtgagccgca	agtttacacc	4200
ctgcccaccga	gccgtgatga	gctgacgaaa	aatcaagtat	ctctgacactg	tctggtcaaa	4260

-continued

ggtttttacc caagcgatat cgcggttgaa tgggagagca acggccagcc ggagaataat	4320
tacaagacga cgectccggt gctggatagc gatgggtcg tttcctgt acaagtttgc	4380
acgggttataaa aaagccgttgc aacacagggtt aacgtgttct cctgtccgtt catgtatgt	4440
gcgcgtgcaca accattataac tcagaaaagc ctcagccgtt ccccggttaa ataagtctag	4500
ataacttagtt gatcggttag tttcacctga tttacgtaaa aacccgttcc ggcgggtttt	4560
tgcttttggaa ggggcagaaa gatgaatgac tgtc	4594

<210> SEQ ID NO 19
 <211> LENGTH: 4332
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pPRO430-Infliximab_LC expression construct

<400> SEQUENCE: 19	
ccccagaggc aggtcttagcg gagtgtataac tggcttacta tggacttgc gatgagggtg	60
tcaagtgaatgt gcttcatgtg gcaggagaaa aaaggctgca ccggctgcgc agcagaataat	120
gtgatacagg atatattccg cttcctcgct cactgactcg ctacgctcg tcgttgcact	180
gcggcgagcg gaaaatggctt acgaacgggg cggagatttc ctggaagatg ccaggaagat	240
acttaacagg gaagttagag ggccgcggca aagccgtttt tccataggtt ccgcggccct	300
gacaagcatc acgaaatctg acgctcaaat cagtgggtgc gaaacccgac aggactataa	360
agataccagg cgtttcccccc tggcggttcc ctctgtcgct ctctgttcc tgccttcgg	420
tttaccgggtg tcattccgtt gttatggccg cgtttgcctc attccacgccc tgacactcg	480
ttccgggttag gcagttcgctt ccaagctgga ctgtatgcac gaaccccccgg ttcagtcga	540
ccgctgcgcc ttatccggta actatcgctc tgagtccaaac ccggaaaagac atgcaaaagc	600
accactggca gcagccactg gtaattgatt tagaggagtt agtcttgaag tcatgcggcc	660
gttaaggcttta aactgaaagg acaagttttg gtgactgcgc tcctccaagc cagttacctc	720
gtttcaaaaga gttggtagct cagagaacct tcgaaaaacc gcccgttgc gcggtttttt	780
cggtttcaga gcaagagatt acgcgcagac caaaacgatc tcaagaagat catcttattaa	840
aacctgcctc taaaaacttt tattacgccc cgcctgttca ctcatcgatc tactgttgc	900
attcattaaag cattctgcgc acatggaagc catcacagac ggcgtatgtt acctgtatcg	960
ccagcggcat cagcaccttg tcgccttgcg tataatattt gcccatagtt gaaacgggg	1020
cgaagaagtt gtccatattt gccacgttta aatcaaaactt ggtgaaactc acccagggtt	1080
tggctgagac gaaaaacata ttctcaataa accctttagg gaaataggcc aggttttgc	1140
cgttaacacgc cacatcttgc gaatataatgtt gttagaaactc cggaaatcg tcgttgcatt	1200
cactccagag cgatgaaaac gtttcaatgtt gtcgtatggaa aacgggttgc caagggttgc	1260
cactatccca tattaccaggc tcaccgtctt tcattgcattt acggaaatcc ggatgagcat	1320
tcattcaggcg ggcaagaatgt tgaataaaagg ccggataaaa cttgtgttta tttttttttt	1380
cggcttttaa aaaggccgtt atatccagctt gaaacgggtcg gttataggtt cattgagca	1440
ctgactgaaa tgcctcaaaa tggcttttcgat gatggccatgg gatataatca acgggttgc	1500
atccagtgtt ttttttctcc atcatattct tcctttttca atattatgtt agcattatc	1560
agggttatttgc tctcatgac ggatacatat ttgtatgtt ttagaaaaat aaacaaatag	1620

-continued

gggtcagtgt	tacaaccaat	taaccaattc	tgatctcctg	ttagtgaggg	ttaatgcccc	1680	
gaacgaagaa	aggcccaccc	gtgaagggtga	gccagtgagt	tggttacatt	ttctttag	1740	
ggtttagctt	ttagacgcac	gcca	aaaagggt	cgtacgtgaa	atacccaaat	agttggccgc	1800
agccgtctt	tcaccattaa	acttctcaag	cgcttgcgtc	gggg	tca	gca aac	1860
cggcgtctt	gcgc	tctcac	gcccagc	cggc	agc a	actgcatga	1920
cagatccggg	gtcggctaa	cggacaggaa	cagcgc	cagg	cgatca	tattacgcag	1980
ctcgccggatg	ttacccggcc	agt	cataatg	cagc	acgc	gatc	2040
ctggcgcagt	gccc	agaga	acgg	tgcgt	cc	agactt	2100
ctccgccc	ggtaaaatgt	cggc	gacac	ttc	acgc	aaatgt	2160
aatgctcagg	cggtagaaca	gg	tcgcgac	aaa	acggccc	tgttgcata	2220
attgcagtgg	gtcgcgctaa	tcac	gcgcac	gtc	tacc	ggat	2280
gcgggtcact	tcttctt	ccag	cacac	cag	cac	ggat	2340
ctcaccgatc	tcgtcc	cagg	aaatgt	gccc	gtgg	caatac	2400
gccaccgcga	cggc	taccc	tgaat	gcgc	cc	atcc	2460
caggcttcc	gcgatt	gcac	cgca	tgc	acaa	atgt	2520
gcgggcatcg	tgac	gggcga	aata	tcac	tgat	ggat	2580
cgtaccagtc	tcgc	catt	tcaga	acac	cg	gcata	2640
ctggcgaact	tgct	ccattt	gagg	gcgtt	gccc	agcata	2700
acgc	acgc	ttac	gcgt	catc	gttgc	atgt	2760
cagggcgtcg	ctga	acgc	gac	gcgt	tgcc	gcgt	2820
gccc	gctt	tcgg	ccaa	gtcg	atcg	atcg	2880
cg	ttc	act	ggat	gtcg	atcg	atcg	2940
caggcgcaga	ttaa	agg	ttc	ttgc	ggat	cggt	3000
gacaacgc	atcg	aggagg	tca	gtttgc	cgct	cgca	3060
accg	ctcg	gc	ttaa	tcac	ggac	acgg	3120
gctacccgt	gcgata	atgg	tttgc	tc	ccgt	ggat	3180
aacggctt	tcgaa	acccca	gtc	aaatcg	gttgc	atgt	3240
gtaatgtcg	cgg	acaact	cga	acagac	gttgc	atcg	3300
atcatcg	tttc	aaac	cggt	gttgc	atcg	atcg	3360
ttgggtttaa	actt	gtttca	taat	gttgc	aatg	aaaac	3420
gttaactgaa	acgc	atattt	at	gtactt	atct	ctaa	3480
aaacatttaa	ttt	ttaa	gtca	attgt	gttgc	atgt	3540
aataacaagt	tgata	acaaa	ag	tttag	gaaat	tttgc	3600
gcccagccat	tct	gtctgtt	tccc	gggcg	agcgt	gttgc	3660
agttcgttgg	ttc	atcgatc	cac	tggtt	acgcgt	atcg	3720
tgattaagta	cgc	gagc	aa	gtat	ccgcgt	atcg	3780
gcggcacgg	c	ttc	acgc	gttgc	atcg	atcg	3840
actgccagca	gag	ccatt	t	ttgcgtt	tttgc	atcg	3900

-continued

aacgcaccgt ggcagcgcgcg tccgtttca ttttcctcc gtccgatgag caactgaaat	3960
cgggcacggc cagcgtcgtg tgtctgtta acaactcta cccgcgtgag gcgaaggtgc	4020
agtggaaagt ggacaacgcgc ctgcaatccg gtaatagcca ggaaagcgctc accgaacaag	4080
atagcaagga cagcacctac agoctgagct ctactctgac cctgagcaag gctgattatg	4140
agaaacacaa ggtctatgca tgtgaggtga cccatcaggg tctgtccagc cccgtcacc	4200
aaagcttcaa tcgcggtag tgctactcg agtaactagt tgatagagat caagcctaa	4260
cgaactaaga ccccccgcacc gaaaggccg ggggtttttt ttgacctaa aacataacc	4320
gaggagcaga ca	4332

<210> SEQ ID NO 20
 <211> LENGTH: 7288
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pSOL-Infliximab expression construct

<400> SEQUENCE: 20	
agctcctgaa aatctcgata actcaaaaaa tacgccccgt agtgcattta ttccattatg	60
gtgaaagttt gaaaccttta cgtgccgatc aagaagacgg tcaaaagcct ccggtcggag	120
gcggggagag tgttcaccga caaacaacag ataaaacaaa aggcccagtc ttccgactga	180
gcctttgtt ttatgtatc tctggcagtt cccgagacgt tatgacaact tgacggctac	240
atcattcact ttttcttac aaccggcagc gaactcgctc gggctggccc cgggtcattt	300
tttaataacc cgcgagaaat agagttgatc gtcaaaacca acattgcac cgacgggtgc	360
gataggcatc cgggtggtagc tcaaaagcag ctgcgcctgg ctgatacggtt ggtcctcg	420
ccagcttaag acgctaatcc ctaactgtctc gcggaaaaga tgtgacagac gcgacggcga	480
caagcaaaaca tgctgtgca cgctggcgat atcaaaatg ctgtctgca ggtgatcgct	540
gtatgtactga caagcctcg gtacccgatt atccatcggtt ggtgagacgc actcgtaat	600
cgcttcatg cgcccgacta acaattgctc aagcagattt atgcgcagca gctccgaaata	660
gcgccttcc ctttgcggcg cgttaatgtat tttccaaac aggtcgctga aatgcggctg	720
gtgcgttca tccggggcga agaaccggat attggcaaat attgacggcc agttaagcca	780
tccatgccag taggcgcgcg gacgaaagta aaccactgg tgataccatt cgcgagcctc	840
cgatgacga ccgtatgtat gaatctctcc tggcgaaac agcaaaat caccggctg	900
gcaaaacaaat ttcgtccct gattttccac caccggctga cccgaaatgg tgagattgag	960
aatataaccc ttcattccca ggggtcggtc gataaaaaaa tcgagataac cggtggctc	1020
aatcggcgat aaacccgcac ccagatgggc attaaacggat tatccggca gcaggggatc	1080
attttgcgtc tcagccatac ttttctactc cccgccccatc agagaagaaa ccaattgtcc	1140
atattgcatac agacattgcc gtctctcgat cttttactgg ctcttcgc taaccaaacc	1200
ggtaaccccg ctttattaaaaa gcattctgtat acaaaggccc accaaagcca tgacaaaaac	1260
gcgtaaacaaa agtgtctata atcacggcag aaaagtccac attgatttatt tgacggcgt	1320
cacactttgc tatggccatag ctttttatac cataagatata gggatccta cctgacgctt	1380
tttattcgcaa ctctctactg tttctccata cccgtttttt tgggctagca ggaggtaaaa	1440
aaaatggaaag ttaaattaga agaaaggccc ggcgggttgg tgcaacccgg cgggtcgat	1500

-continued

aagttgagct	gcgtcgcaag	cggttcatt	tttccaacc	actggatgaa	ctgggtgcgc	1560
cagtctccgg	aaaagggtct	ggaatgggtt	gcccggatcc	gtaccaagag	catcaatagc	1620
gcgcacgcatt	atgcccggat	cgtcaaaaggc	cgcttcacca	tttctcgta	cgacagcaaa	1680
agegcgtgtt	atctgcaat	gaccgacttg	cgatccggagg	acacgggcgt	gtattactgc	1740
tcccgcaact	attacggttc	cacctacgac	tactggggcc	agggtacgac	cctgaccgtt	1800
agctcggcga	gcaccaaggg	tcggagcgct	tttccgctgg	caccgagcag	caaaagcacc	1860
agcgggtggta	ccgcccgcact	gggttgcctg	gtgaaagatt	acttcccgga	accggttact	1920
gtgagctgga	acagcggcgc	gtcgacccct	ggcggtcaca	cgttccggc	agttctgcaa	1980
agcagcggcc	tgtactccct	gtccagcgct	gtcaccgtgc	cgagcagcag	cctgggtacg	2040
caaacctata	tttgcgtacgt	caatcacaag	ccgagcaaca	ccaaagtgga	caaaaaagtc	2100
gaaccgaaaa	gctgcgataa	aacccataact	tgtccgcccgt	gcccggcccc	ggagcttctg	2160
ggtggtccaa	cggtttttct	gttcccggcc	aagccgaaag	acaccctgtat	gtcggccgc	2220
acccctgagg	tgacccgtgt	ggtagttgac	gttcccacg	aagatccaga	ggtcaagttt	2280
aactggatgt	tggatggcgt	cgaagttcac	aatgcaaaaga	ccaagccgcg	tgaagaacag	2340
tataactcta	cgtaccgtgt	cgtgagcggt	ctgactgttc	tgcaccaggaa	ttggctgaac	2400
ggcaaaagagt	acaagtgc当地	ggtagcaat	aaagcgctgc	cggtcccgat	cgagaaaaacc	2460
atttctaagg	ctaaagggtca	gcccgcgtgag	ccgcaagttt	acaccctgccc	accgagccgt	2520
gatgagctga	cgaaaaatca	agtatctctg	acctgtctgg	tcaaagggtt	ttacccaagc	2580
gatatacgccg	ttgaatggga	gagcaacggc	cagccggaga	ataattacaa	gacgacgcct	2640
ccgggtctgg	atagcgatgg	ttcggttttc	ctgtacagca	agttgacggt	tgataaaagc	2700
cgttggcaac	agggttaacgt	gttctccctgt	tccgtcatgc	atgaaggcgt	gcacaaccat	2760
tataactcaga	aaagcctcag	cctgtccccc	ggtaaataag	tctagataac	tagttgatcg	2820
gtcagtttca	cctgatttac	gtaaaaacc	gttccggccg	gtttttgtct	ttggaggggc	2880
agaaaagatga	atgactgtct	ctctgttag	tgagggttaa	tgcccgaaac	gaagaaaaggc	2940
ccacccgtga	aggtgagcca	gtgagttgtt	tacattttct	cttgagggtt	tagtttca	3000
gacgacgcca	aaaggtcgta	cgtgaaatac	ccaaatagtt	ggccgcagcc	gtcttgcac	3060
cattaaactt	ctcaagcgct	tgctgcgggg	tcaagcaaaacg	cggagccggc	gtcttgcgc	3120
tctcacgcgc	cagctccggc	agcagcaact	gcatgaattg	cggagtcaga	tccgggtcgt	3180
gctcaacggc	caggaacgc	gccagggcgtt	ccatcatatt	acgcagctcg	cggtatgttac	3240
ccggccagtc	ataatgcagc	agcaccgttt	cgctcgctg	cagaccctgg	cgcagtgcgc	3300
cagagaacgg	tgcgttcagg	gtgtccagcg	agactttcag	gaaagactcc	gccagcggtt	3360
aaatgtcgcc	gacacgttca	cgcaacggcg	ggagctcgag	acgcagaatg	ctcaggcggt	3420
agaacagggtc	gcccggat	cgccctgtt	gcatatccctc	ttccagattg	cgtgggtcg	3480
cgctaatac	gcccacgtct	accggaaacc	gttgcgttacc	accgacgcgg	gtcacttctt	3540
tctttccag	cacacgcagc	agacgggttt	gcaatggcag	cggcatctca	ccgatctgt	3600
ccaggaacaa	ggtgcgcgcgc	tggcaattt	caaacaacc	agcacggcca	ccgcgcacggc	3660
tacccgtgaa	tgcgtccctct	tcgttagccaa	acagctcagc	ttccagcagg	ctttccgcga	3720
ttgcaccgca	attaaactgcc	acaaacggat	gagatttctt	accctggccg	gcacgtgac	3780

-continued

gggctaaata ctcacgatgg attgcttgcg cagccagtgc cttaccgta ccagtctcg	3840
cttcgatcag aacagccgcg ctgctacgtg catacagcag aatggctgg cgaacttgc	3900
ccatttgggg gctttggccc agcatatcac ccaggacata acgggtacgc agcgattac	3960
gcgtcgcatc gtgggtgtt tggcgacggc tcattctgtt catgtccagg gcgtcgctga	4020
acgctgtacg caccgttgcg gcgctgtaga taaagatgcc cgtcatgccc gcttcttcgg	4080
ccaagttcgt gatcagaccc gcaccaacca cagcctcggt accgttcgt ttcagttcgt	4140
tgtatctggcc acgtgcacatc tcctcggtaa tgttagctgcg ttgggtccagg cgccagattaa	4200
aggcttttg aaacgcgacc agcgcaggaa tcgtttccctg gttaggtaca acgccaatcg	4260
aggaggtcag tttgcctgcg ttgcgcagcg cctgcaagac atcgttaaccg ctgggtttaa	4320
tcaggatcac cggcacggac agacgggatt tcaggttaggc accattgcta cccgctgcga	4380
taatggcgtc acaacgctcg ttggccagct ttttgcgaat gttagttaacg gcttttcg	4440
aacccagctg aatcggagtg atgttcgcca ggtatcaaa ctccaggcta atgtcgccg	4500
acaactcgaa cagacgggtg acgctaacgg tccaaataac tggtttatca tcgttcaa	4560
gcgggtgggtg tgccatgggt aataacctct gttaaagaaac cgaatattgg gttttaactt	4620
gtttcataat tggtgcata gAACCGGGTG aaacattgaa tgaaacgttta actgaaacgc	4680
atatttgcgg attagttcat gactttatct ctaacaaatt gaaattaaac atttaatttt	4740
attaaggcaa ttgtggcaca ccccttgcgtt tgcttttatac aacgcaaata acaagttgat	4800
aacaaaagct taggaggaaa acatatggat attttattga cccaaagccc agccattctg	4860
tctgtttccc cggggcagcg tgtttctttt agctgcgtg caagccgtt cgttgggtca	4920
tcgatccact ggtaccaaca gcgtacgaaac ggtagccgcg gtctgtgtat taagtacgc	4980
agcgaaagca tgagcggat cccgagccgc ttcaagcgtt gccggcagcgg cacggacttc	5040
acgctgagca tcaatacggt tgaaagcgag gacatcgccg actattactg ccagcagac	5100
cattttggc cggttacattt tggcagcgtt accaatctgg aagttaaacg caccgtggca	5160
gcgcgtccg ttttcatattt tcctccgtcc gatgagcaac tgaaatcggg cacggcagc	5220
gtcggtgttc tggtgaacaa cttctacccg cgtgaggcga aggtgcgtt gaaagtggac	5280
aacgcgtgc aatccggtaa tagccaggaa agcgtcaccc aacaagatag caaggacagc	5340
acctacagcc tgagctctac tctgaccctg agcaaggctg attatgagaa acacaaggc	5400
tatgcacgtg aggtgaccca tcagggtctg tccagcccg tcaacaaag cttcaatcgc	5460
ggtgagtgtc aactcgagta actagttgtt agagatcaag cttcaacgaa ctaagacccc	5520
cgcaccgaaa ggtccggggg ttttttttga cttaaaaac ataaccgagg agcagacagg	5580
cctttcttcg gtatgttctt tcccccagag gcaggatca aaggatcttc ttgatgtt	5640
tttttctgc gcgtaatctg ctgcttgcac aaaaaaaaaa caccgttacc agcggtgggt	5700
tgtttggccg atcaagagct accaactctt tttccgaggat aactggcttc agcagacgc	5760
agataacaaa tactgtttt ctatgttagc cgtatgttcc accaccatcc aagaactctg	5820
tagcaccgac tacatacctc gctctgttacg tccgttacc agtggctgtt gccagtgcc	5880
ataagtcgtg tcttaccggg ttggactcaa gacgatgtt accggataag gcgacgggt	5940
cgggctgaac ggggggttcg tgcacacagc ccagcttggc gcgaaacgacc tacaccga	6000
tgagataacct acagcgtgag ctatgagaaa gcgccacgct tcccgaaaggg agaaaggcgg	6060

-continued

```

acaggtatcc ggtaagcggc agggtcggaa caggagagcg cacgaggagg cttccagggg 6120
gaaacgcctg gtatcttat agtcctgtcg ggtttcgcca cctctgactt gagcatcgat 6180
ttttgtatg ctctgtcaggg gggcggagcc tatggaaaaa cgccagcaac gcagaaaggc 6240
ccacccgaag gtgagccagg tgattacatt tggccctca tcagaggttt tcaccgtcat 6300
caccgaaacg cgcgaggcag ctgcggtaaa gtcatcagc gtggtegtga agcgattcac 6360
agatgtctgc ctgttcatcc gcttccagct cgttgagttt ctccagaagc gttaatgtct 6420
ggcttctgtat aaagcgggccc atgttaaggg cggtttttc ctgtttggtc atacctgctt 6480
agaaaaactc atcgagcatc aaatgaaatt gcaattttt catatcagga ttatcaatac 6540
catatttttggaaaagccgt ttctgtatag aaggagaaaa ctcaccgagg cagttccata 6600
ggatggcaag atcctggat cggtctgcga ttccgactcg tccaacatca atacaaccta 6660
ttaattttccc ctctgtcaaaa ataaggatcaatgtaatgaa atcaccatga gtgacgactg 6720
aatccgggtga gaatggcaaa agtttatgcata tttctttcca gacttgtca acaggccagc 6780
cattacgctc gtcatcaaaa tcactcgcata caaccaaacc gttatttcatt cgtgattgcg 6840
cctgagcggcag gcgaaataacg cgatcgctgt taaaaggaca attacaaaca ggaatcgagt 6900
gcaaccggcgc caggaacact gccagcgcata caacaatatt ttcacctgaa tcaggatatt 6960
cttctaatac ctggAACGCT gttttccggc ggatcgcaat ggtgagtaac catgcataat 7020
caggagttacg gataaaatgc ttgatggtcg gaagtggcat aaattccgtc agccagttt 7080
gtctgaccat ctcatctgtat acatcattgg caacgcatacc tttgccatgt ttcagaaaca 7140
actctggcgc atcgggcttc ccatacaacg gatagattgt cgcacctgtat tgcccgacat 7200
tatcgcgagc ccatttatacc ccatataaaat cagcatccat gttggaaattt aatcgoggcc 7260
tcgacgtttc ccgttgaata tggctcat 7288

```

What is claimed is:

1. An expression construct comprising two or more inducible promoters, wherein at least one of said inducible promoters is responsive to an inducer that is different than the inducer of another of said inducible promoters, and wherein none of the inducible promoters is an inducible promoter selected from the group consisting of: a tetracycline-inducible promoter, a copper-inducible promoter, and a methionine-inducible promoter.

2. The expression construct of claim 1, wherein the expression construct is extrachromosomal.

3. The expression construct of claim 1, wherein each inducible promoter is selected from the group consisting of an L-arabinose-inducible promoter, a propionate-inducible promoter, a rhamnose-inducible promoter, a xylose-inducible promoter, a lactose-inducible promoter, and a promoter inducible by phosphate depletion.

4. The expression construct of claim 3, wherein at least one inducible promoter is selected from the group consisting of the araBAD promoter, the prpBCDE promoter, the rhaSR promoter, the xlyA promoter, the lacZYA promoter, and the phoA promoter.

5. The expression construct of claim 1, wherein the expression construct comprises at least one propionate-inducible promoter and at least one inducible promoter selected from the group consisting of an L-arabinose-inducible promoter, a

rhamnose-inducible promoter, a xylose-inducible promoter, a lactose-inducible promoter, and a promoter inducible by phosphate depletion.

6. The expression construct of claim 5, wherein the propionate-inducible promoter is the prpBCDE promoter.

7. The expression construct of claim 5 comprising a nucleotide sequence having at least 80% sequence identity to at least 50 contiguous bases of nucleotides 4937 through 5185 of SEQ ID NO:15.

8. The expression construct of claim 1, wherein at least one inducible promoter is a propionate-inducible promoter and at least one other inducible promoter is an L-arabinose-inducible promoter.

9. The expression construct of claim 8 comprising a nucleotide sequence having at least 80% sequence identity to at least 50 contiguous bases of nucleotides 2818 through 3151 of SEQ ID NO:15, and at least 80% sequence identity to at least 50 contiguous bases of nucleotides 4937 through 5185 of SEQ ID NO:15.

10. The expression construct of claim 9 comprising SEQ ID NO:15.

11. A host cell comprising the expression construct of claim 1.

12. The host cell of claim 11 wherein the host cell is a prokaryotic cell.

13. The host cell of claim 12 wherein the host cell is *E. coli*.

14. The host cell of claim **13** wherein the host cell further comprises one or more of the following: (a) a deletion of the araBAD genes; (b) an altered gene function of the araE and araFGH genes; (c) a lacY(A177C) gene; (d) a reduced gene function of the prpB and prpD genes; (e) a reduced gene function of the sbm/scpA-ygfD/argK-ygfGH/scpBC genes; (f) a reduced gene function of the gor and trxB genes; (g) a reduced gene function of the AscG gene; (h) a polynucleotide encoding a form of DsbC lacking a signal peptide; (i) a polynucleotide encoding Erv1p; and (j) a polynucleotide encoding a chaperone.

15. The host cell of claim **13** wherein the host cell is *E. coli* ASE(DGH).

16. A host cell comprising the expression construct of claim **1**, wherein the expression construct further comprises at least one polynucleotide sequence encoding at least one gene product to be transcribed from an inducible promoter.

17. A method of producing a gene product, the method comprising growing a culture of the host cell of claim **16** and adding an inducer of at least one inducible promoter to the culture.

18. The method of claim **17**, wherein the method further comprises purifying the gene product from the culture.

19. A gene product produced by the method of claim **17**.

20. The gene product of claim **19** wherein the gene product is selected from the group consisting of: (a) an immunoglobulin heavy chain; (b) an immunoglobulin light chain; and (c) a fragment of any of (a)-(b).

21. The gene product of claim **20**, wherein the gene product is part of a multimeric product.

22. The multimeric product of claim **21**, wherein the multimeric product is an antibody.

23. A kit comprising the host cell of claim **11**.

24. An expression construct comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence having at least 97% sequence identity to at least 225 contiguous bases of nucleotides 4937 through 5185 of SEQ ID NO:15;

(b) a nucleotide sequence having at least 80% sequence identity to at least 300 contiguous bases of nucleotides 4937 through 5304 of SEQ ID NO:15;

(c) a nucleotide sequence having at least 87% sequence identity to at least 350 contiguous bases of nucleotides 2818 through 3259 of SEQ ID NO:15; and

(d) a nucleotide sequence having at least 90% sequence identity to at least 450 contiguous bases of nucleotides 10 through 1822 of SEQ ID NO:2.

* * * * *