

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3524603 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 D 471/04 (2006.01)** **A 61 K 31/436 (2006.01)** **A 61 K 31/4365 (2006.01)**
A 61 K 31/437 (2006.01) **A 61 K 31/4375 (2006.01)** **A 61 K 31/445 (2006.01)**
A 61 K 31/497 (2006.01) **A 61 K 31/505 (2006.01)** **A 61 K 31/506 (2006.01)**
A 61 K 31/517 (2006.01) **A 61 K 31/519 (2006.01)** **A 61 K 31/535 (2006.01)**
A 61 K 31/5355 (2006.01) **A 61 K 31/537 (2006.01)** **A 61 P 35/00 (2006.01)**
C 07 D 491/044 (2006.01) **C 07 D 491/107 (2006.01)** **C 07 D 495/04 (2006.01)**
C 07 D 498/08 (2006.01) **C 07 D 513/04 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-07-04**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2022-06-08**
- (86) Europæisk ansøgning nr.: **17883905.6**
- (86) Europæisk indleveringsdag: **2017-12-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-08-14**
- (86) International ansøgning nr.: **CN20171116186**
- (87) Internationalt publikationsnr.: **WO2018113584**
- (30) Prioritet: **2016-12-19 CN 201611177188** **2017-06-15 CN 201710453749**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Abbisko Therapeutics Co., Ltd., No. 3, Lane 898, HaLei Road , China (Shanghai) Pilot Free Trade Zone , Pudong New Area, Shanghai 201203, Kina**
- (72) Opfinder: **YANG, Fei, Floor 4 Building 6 Lane 576 Libing Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203, Kina**
DENG, Haibing, Floor 4 Building 6 Lane 576 Libing Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203, Kina
YING, Haiyan, Floor 4 Building 6 Lane 576 Libing Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203, Kina
YU, Hongping, Floor 4 Building 6 Lane 576 Libing Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203, Kina
CHEN, Zhui, Floor 4 Building 6 Lane 576 Libing Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203, Kina
XU, Yaochang, Floor 4 Building 6 Lane 576 Libing Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203, Kina
- (74) Fuldmægtig i Danmark: **Dennemeyer & Associates S.A, P.O. Box 700425, DE-81304 Munich, Tyskland**
- (54) Benævnelse: **FGFR4-HÆMMER, FREMGANGSMÅDE TIL FREMSTILLING DERAFF OG FARMACEUTISK ANVENDELSE DERAFF**
- (56) Fremdragne publikationer:

Fortsættes ...

WO-A1-2014/144737

WO-A1-2015/061572

WO-A1-2015/108992

WO-A1-2016/115412

WO-A2-2014/011900

CN-A- 104 540 809

CN-A- 105 307 657

CN-A- 105 658 642

QIUMENG ZHANG ET AL: "Design, Synthesis and Anti-Proliferative Activities of 2,6-Substituted Thieno[3,2-d]pyrimidine Derivatives Containing Electrophilic Warheads", MOLECULES, vol. 22, no. 5, 12 May 2017 (2017-05-12), pages 1-16, XP055594292, DOI: 10.3390/molecules22050788

DESCRIPTION

Technical field

[0001] The present invention belongs to the field of medicament synthesis, and in particular relates to an FGFR4 inhibitor, preparation method and pharmaceutical use thereof.

Technical background

[0002] Fibroblast growth factor (FGF) is a family of 22 structurally related polypeptides with diverse biological activities that can regulate cell proliferation, differentiation and migration, and play a major role in the process of limb development, angiogenesis, tissue repair, tumor formation and the like. (Eswarakumar et al., 2005 Cytokine Growth Factor Rev16: 139-149; Ornitz and Itoh, 2001 Genome Bio12: Reviews 3005).

[0003] The receptors for FGF (FGFR) belong to a family of RPTK of receptor tyrosine kinases. Four FGFRs, FGFR1, FGFR2, FGFR3 and FGFR4, have been identified to date (Ullrich and Schlessinger, 1990 Cell 61:203). The interaction between receptors and the corresponding ligands FGF leads to receptor dimerization and autophosphorylation, thereby initiating multiple downstream signaling cascades including MAPK and AKT (Powers et al., 2000 Endocr Relat Cancer7: 165-197).

[0004] FGFR1-3 has been found to be overexpressed, mutated or translocated in a variety of tumors (including myeloma, breast cancer, stomach cancer, colon cancer, bladder cancer, pancreatic cancer, and hepatocellular carcinoma), and considered to be driver gene in cancer (Chesi et al., 2001 Blood 97:729-726; Gowardhan et al., 2005 Br J Cancer 92: 320-327; Jaakkola et al., 1993 Int J Cancer 54:378-282; Jang et al., 2001 Cancer Res 61: 3541-3543). Some FGFR inhibitors have also been developed in the clinical and preclinical development process. However, previous studies have shown that FGFR1 can regulate the level of phosphate, so pan-FGFR inhibitors may pose safety concerns.

[0005] Hepatocellular carcinoma (HCC) is one of the leading causes of cancer-related deaths in China and is one of the fastest growing cancers every year (Shariff et al., 2009 Expert Rev Gastroenterol Hepato 13: 353-367). Currently, the first-line treatment option is sorafenib, there are no approved second-line treatment, and there is still a need for targeted therapy with anti-tumor agents.

[0006] Overexpression of FGF19 is present in 5-10% of hepatocellular carcinoma patients, whereas FGFR4 is a dominant FGFR present in human hepatocytes, and its high expression in hepatocytes is found to be associated with the aggressiveness of hepatocellular tumors. Therefore, FGFR4 plays a very important role in liver cancer. In addition, the interaction of

FGF19 and FGFR4 is also considered to be related to the aggressiveness of other cancers (such as gastric cancer, prostate cancer, lung cancer, colorectal cancer, pancreatic cancer, and ovarian cancer) (Ye et al, 2011 Cancer 5304-5313; Xu et al, 2011 BMC Cancer 11:84; Fawdar et al, 2013 PNAS 110:12426-12431).

[0007] At present, some FGFR inhibitors have entered into the clinical research stage as anti-tumor drugs, but mostly inhibitors against FGFR1, 2 and 3, with weaker inhibition of FGFR4 activity. The inhibition of FGFR1-3 has on-target side effects such as hyperphosphatemia. Highly selective inhibitor of FGFR4 can effectively treat cancer caused by abnormal FGFR4 signaling, and can avoid the side effects caused by FGFR1-3 inhibition such as hyperphosphatemia. Highly selective small molecule inhibitors against FGFR4 have significant application prospects in the field of anti-tumor targeted therapy. Therefore, the development of a novel anti-tumor agent that can selectively target FGFR4 as a good drug candidate will meet the needs of domestic liver cancer and other anti-tumor target therapy, and have the advantages of better safety and higher selectivity.

[0008] WO 2014/011900 discloses inhibitors of the fibroblast growth factor receptor.

[0009] Qiumeng Zhang et al. discloses design, synthesis and anti-proliferative activities of 2,6-substituted thiophen[3,2-d] pyrimidinderivatives.

[0010] WO 2014/144373 discloses heteroaryl compounds and uses thereof.

[0011] WO 2016/115412 discloses dual-warhead covalent inhibitors of FGFR-4.


[0012] WO 2015/061572 discloses inhibitors of the fibroblast growth factor receptor.

[0013] WO 2015/108992 discloses heterocyclic compounds and the use as FGFR-4 receptor inhibitors.

Summary of the invention

[0014] An object of the present invention is to provide an FGFR4 inhibitor, preparation method and pharmaceutical use thereof.

[0015] The first aspect of the invention provides a compound of formula (IVa-1), a stereoisomer or a pharmaceutically acceptable salt thereof as defined in the claims:



[0016] In the most preferred embodiment, the compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof includes, but is not limited to, the compounds defined in claim 3.

[0017] The second aspect of the invention provides a process for preparing the above compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof, as defined in claim 4.

[0018] The third aspect of the present invention provides a pharmaceutical composition comprising a therapeutically effective amount of the above compound of formula (IVa-1), stereoisomer or a pharmaceutically acceptable salt thereof, and pharmaceutically acceptable carrier.

[0019] The fourth aspect (not according to the present invention) provides use of the above compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof or the aforementioned pharmaceutical composition for preparing a medicament as an FGFR4 inhibitor.

[0020] The fifth aspect of the present invention provides use of the above compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof or the aforementioned pharmaceutical composition for preparing a medicament for treating cancer.

[0021] Preferably, the cancer is prostate cancer, liver cancer, pancreatic cancer, esophageal cancer, gastric cancer, lung cancer, breast cancer, ovarian cancer, colon cancer, skin cancer, glioblastoma or rhabdomyosarcoma.

[0022] The sixth aspect (not according to the invention) provides the above compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof or the aforementioned pharmaceutical composition for use as an FGFR4 inhibitor.

[0023] The seventh aspect of the invention provides the above compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof or the aforementioned pharmaceutical composition for use in the treatment of cancer.

[0024] Preferably, the cancer is prostate cancer, liver cancer, pancreatic cancer, esophageal cancer, gastric cancer, lung cancer, breast cancer, ovarian cancer, colon cancer, skin cancer, glioblastoma or rhabdomyosarcoma.

[0025] The eighth aspect (not according to the invention) provides a method for inhibiting FGFR4, which comprises administering to a patient in need thereof a therapeutically effective amount of the above compound of formula (IVa-1), the stereoisomer or pharmaceutically acceptable salt thereof, or the aforementioned pharmaceutical composition.

[0026] The ninth aspect (not according to the invention) provides a method for treating cancer, which comprises administering to a patient in need thereof treatment a therapeutically effective amount of the above compound of formula (IVa-I), the stereoisomer or pharmaceutically acceptable salt thereof, or the aforementioned pharmaceutical composition.

[0027] Preferably, the cancer is prostate cancer, liver cancer, pancreatic cancer, esophageal cancer, gastric cancer, lung cancer, breast cancer, ovarian cancer, colon cancer, skin cancer, glioblastoma or rhabdomyosarcoma.

[0028] The series of compounds developed by the present invention have strong inhibitory effects on FGFR4 kinase activity and very high selectivity, and can be widely used for preparing a medicament for treating cancer, especially prostate cancer, liver cancer, pancreatic cancer, esophageal cancer, stomach cancer, lung cancer, breast cancer, ovarian cancer, colon cancer, skin cancer, glioblastoma or rhabdomyosarcoma. The compounds are expected to be developed into a new generation medicaments of FGFR4 inhibitor.

[0029] It is to be understood that within the scope of the present invention, which is defined by the claims, the above various technical features of the present invention and the technical features specifically described hereinafter (as in the examples) may be combined with each other to constitute a new or preferred technical solution. Due to space limitations, they will not be described one by one.

Detailed description of the invention

[0030] Based on a long-term and in-depth study, the inventors have developed for the first time an FGFR4 inhibitor with a structure of the formula (IVa-1), the series of compounds have very strong inhibitory effects on FGFR4 kinase activity and very high selectivity, and could be widely used for preparing a medicament for treating cancer, especially prostate cancer, liver cancer, pancreatic cancer, esophageal cancer, stomach cancer, lung cancer, breast cancer, ovarian cancer, colon cancer, skin cancer, glioblastoma or rhabdomyosarcoma. These compounds will be expected to be developed into a new generation medicaments of FGFR4 inhibitor. On such basis, the present invention has been completed.

[0031] Detailed description: Unless otherwise stated, the following terms used in the specification and claims have the following meanings.


[0032] "Alkyl" means a straight or branched saturated aliphatic hydrocarbon group, for example, "C₁₋₈ alkyl" means a straight or branched alkyl having 1 to 8 carbon atoms, including but is not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, sec-butyl, n-pentyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 1-ethyl-2-methylpropyl, 1,1,2-trimethylpropyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1,3-dimethylbutyl, 2-ethylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,3-dimethylbutyl, n-heptyl, 2-methylhexyl, 3-methylhexyl, 4-

methylhexyl, 5-methylhexyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,2-dimethylpentyl, 3,3-dimethylpentyl, 2-ethylpentyl, 3-ethylpentyl, n-octyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylhexyl, 3,3-dimethylhexyl, 4,4-dimethylhexyl, 2-ethylhexyl, 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-ethylpentyl or various branched isomers thereof and so on.


[0033] Alkyl can be optionally substituted or unsubstituted, and when substituted, according to the claims.

[0034] "Cycloalkyl" means a saturated or partially unsaturated monocyclic or polycyclic hydrocarbon substituent, for example, "C₃₋₈ cycloalkyl" refers to a cycloalkyl having 3-8 carbon atoms, which may be a monocyclic cycloalkyl and a polycyclic cycloalkyl, wherein, monocyclic cycloalkyl includes, but is not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, cycloheptatrienyl, cyclooctyl and the like; and polycyclic cycloalkyl includes spiro, fused, and bridged cycloalkyls.

[0035] "Spirocycloalkyl" refers to a polycyclic group that shares a carbon atom (called a spiro atom) between the monocyclic rings. These groups may contain one or more double bonds, but none of the rings have a fully conjugated π -electron system. The spirocycloalkyl may be a monospirocycloalkyl, a bispirocycloalkyl or a polyspirocycloalkyl according to the number of common spiro atoms between the rings, spirocycloalkyl includes, but is not limited to:




[0036] "Fused cycloalkyl" means an all-carbon polycyclic group in which each ring shares an adjacent pair of carbon atoms with other rings in the system, wherein one or more of the rings may contain one or more double bonds, but none of the rings have a fully conjugated π -electron system. Depending on the number of rings, it may be bicyclic, tricyclic, tetracyclic or polycyclic, fused cycloalkyl includes but is not limited to:



[0037] "Bridged cycloalkyl" refers to an all-carbon polycyclic group in which any two rings share two carbon atoms that are not directly bonded, which may contain one or more double bonds,

but none of the rings have a fully conjugated pi-electron system. Depending on the number of rings, it may be bicyclic, tricyclic, tetracyclic or polycyclic, bridged cycloalkyl includes but is not limited to:




[0038] The ring of the cycloalkyl may be fused to a ring of aryl, heteroaryl or heterocycloalkyl, wherein the ring attached to the parent structure is a cycloalkyl, includes, but is not limited to indanyl, tetrahydronaphthyl, benzocycloheptyl and the likes.

[0039] The cycloalkyl can be optionally substituted or unsubstituted, and when substituted, according to the claims.


[0040] "Heterocyclyl" means a saturated or partially unsaturated monocyclic or polycyclic cyclic hydrocarbon substituent wherein one or more of the ring atoms are heteroatoms selected from nitrogen, oxygen or S(O)_r (wherein r is an integer of 0, 1, 2), but excluding ring moiety of -O-O-, -O-S- or -S-S-, and the remaining ring atoms are carbon atoms. For example, "5-10 membered heterocyclyl" means a cyclic group containing 5 to 10 ring atoms, and "3-10 membered heterocyclyl" means a cyclic group containing 3 to 10 ring atoms.

[0041] Monocyclic heterocyclyl includes, but is not limited to pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, homopiperazinyl and the likes.


[0042] Polycyclic heterocyclyl includes a spiro, fused, and bridged heterocyclyl. "Spiroheterocyclyl" refers to a polycyclic heterocyclyl that shares a carbon atom (called a spiro atom) between the monocyclic rings, wherein one or more of the ring atoms are heteroatoms selected from nitrogen, oxygen or S(O)_r (wherein r is an integer of 0, 1, 2), and the remaining ring atoms are carbon atoms. These groups may contain one or more double bonds, but none of the rings have a fully conjugated π-electron system. The spiroheterocyclyl may be a monospiroheterocyclyl, a bispiroheterocyclyl or a polyspiroheterocyclyl according to the number of common spiro atoms between the rings, spiroheterocyclyl includes, but is not limited to:




[0043] "Fused heterocyclyl" means a polycyclic heterocyclyl in which each ring shares an adjacent pair of carbon atoms with other rings in the system, wherein one or more of the rings may contain one or more double bonds, but none of the rings have a fully conjugated π -electron system, wherein one or more of the ring atoms are heteroatoms selected from nitrogen, oxygen or $S(O)_r$ (wherein r is an integer of 0, 1, 2), and the remaining ring atoms are carbon atoms. Depending on the number of rings, it may be bicyclic, tricyclic, tetracyclic or polycyclic, fused heterocyclyl includes, but is not limited to:



[0044] "Bridged heterocyclyl" refers to a polycyclic heterocyclyl in which any two rings share two carbon atoms that are not directly bonded, which may contain one or more double bonds, but none of the rings have a fully conjugated π -electron system, wherein one or more of the ring atoms are heteroatoms selected from nitrogen, oxygen or $S(O)_r$ (wherein r is an integer of 0, 1, 2), and the remaining ring atoms are carbon atoms. Depending on the number of rings, it may be bicyclic, tricyclic, tetracyclic or polycyclic, bridged heterocyclyl includes, but is not limited to:




[0045] The ring of the heterocyclyl may be fused to a ring of aryl, heteroaryl or cycloalkyl wherein the ring attached to the parent structure is a heterocyclyl, includes, but is not limited to:






[0046] The heterocyclyl can be optionally substituted or unsubstituted, and when substituted, according to the claims.

[0047] "Aryl" means an all-carbon monocyclic or fused polycyclic (ie, a ring that shares a pair of adjacent carbon atoms) group, and a polycyclic group having a conjugated π -electron system (i.e., a ring with adjacent pairs of carbon atoms), for example, "C₅₋₁₀ aryl" means an all-carbon aryl having 5-10 carbons, and "5-10 membered aryl" means an all-carbon aryl having 5-10 carbons, including but not limited to phenyl and naphthyl. The aryl ring may be fused to a ring of heteroaryl, heterocyclyl or cycloalkyl, wherein the ring attached to the parent structure is an aryl ring, includes, but is not limited to:



[0048] The aryl group can be optionally substituted or unsubstituted, and when substituted, the substituent is preferably one or more of the following groups, and independently substituted, according to the claims.

[0049] "Heteroaryl" refers to a heteroaromatic system containing 1 to 4 heteroatoms including a hetero atom selected from nitrogen, oxygen or S(O)_r (wherein r is an integer of 0, 1, 2), for example, 5-7 membered heteroaryl refers to a heteroaromatic system containing 5 to 7 ring atoms, and 5-10 membered heteroaryl refers to a heteroaromatic system containing 5 to 10 ring atoms, including but not limited to furyl, thiophenyl, pyridyl, pyrrolyl, N-alkylpyrrolyl, pyrimidinyl, pyrazinyl, imidazolyl, tetrazolyl group or the like. The heteroaryl ring may be fused to a ring of aryl, heterocyclyl or cycloalkyl wherein the ring attached to the parent structure is a heteroaryl ring, includes, but is not limited to:



[0050] The heteroaryl can be optionally substituted or unsubstituted, and when substituted,

according to the claims.

[0051] "Alkenyl" refers to an alkyl group as defined above consisting of at least two carbon atoms and at least one carbon-carbon double bond, for example, C₂₋₈ alkenyl refers to a straight or branched alkenyl containing 2 to 8 carbons. Alkenyl includes, but is not limited to vinyl, 1-propenyl, 2-propenyl, 1-, 2- or 3-butenyl, and the likes.

[0052] The alkenyl group can be substituted or unsubstituted, and when substituted, according to the claims.

[0053] "Alkynyl" refers to an alkyl group as defined above consisting of at least two carbon atoms and at least one carbon-carbon triple bond, for example, C₂₋₈ alkynyl refers to a straight or branched alkynyl containing 2 to 8 carbons. Alkynyl includes, but is not limited to ethynyl, 1-propynyl, 2-propynyl, 1-, 2- or 3-butyne, and the likes.

[0054] The alkynyl can be substituted or unsubstituted, and when substituted, according to the claims.

[0055] "Alkoxy" means -O-(alkyl), wherein alkyl is as defined above, for example, "C₁₋₈ alkoxy" refers to an alkoxy containing 1 to 8 carbons. Alkoxy includes, but is not limited to methoxy, ethoxy, propoxy, butoxy, and the likes.

[0056] The alkoxy can be substituted or unsubstituted, and when substituted, according to the claims.

[0057] "Cycloalkyloxy" means -O-(unsubstituted cycloalkyl), wherein cycloalkyl is as defined above, for example, "C₃₋₈ cycloalkyloxy" refers to a cycloalkyloxy containing 3 to 8 carbon atoms. Cycloalkyloxy includes, but is not limited to, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and the likes.

[0058] The cycloalkyloxy can be substituted or unsubstituted, and when substituted, according to the claims.

[0059] "3-10 membered heterocyclyloxy" means -O-(unsubstituted 3-10 membered heterocyclyl), wherein 3-10 membered heterocyclyl is as defined above, 3-10 membered heterocyclyloxy can be substituted or unsubstituted, and when substituted, according to the claims.

[0060] "C₅₋₁₀ aryloxy" means -O-(unsubstituted C₅₋₁₀ aryl), wherein C₅₋₁₀ aryl is as defined above, C₅₋₁₀ aryloxy can be substituted or unsubstituted, and when substituted, according to the claims.

[0061] "5-10 membered heteroaryloxy" means -O-(unsubstituted 5-10 membered heteroaryl),

wherein 5-10 membered heteroaryl is as defined above, 5-10 membered heteroaryloxy can be substituted or unsubstituted, and when substituted, according to the claims.

[0062] "C₁₋₈ alkanoyl" refers to a monovalent group obtained by removing hydroxyl from C₁₋₈ alkyl acid, is also generally referred to as "C₀₋₇-C(O)-", for example, "C₁-C(O)-" refers to acetyl; "C₂-C(O)-" refers to propionyl; and "C₃-C(O)-" refers to butyryl or isobutyryl.

[0063] "-C₀₋₈-S(O)_rR₁₀" means that the sulfur atom in -S(O)_rR₁₀ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0064] "-C₀₋₈-O-R₁₁" means that the oxygen atom in -O-R₁₁ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0065] "-C₀₋₈-C(O)OR₁₁" means that the carbonyl group in -C(O)OR₁₁ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0066] "-C₀₋₈-C(O)R₁₂" means that the carbonyl group in -C(O)R₁₂ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0067] "-C₀₋₈-O-C(O)R₁₂" means that the oxygen atom in -O-C(O)R₁₂ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0068] "-C₀₋₈-NR₁₃R₁₄" means that the nitrogen atom in -NR₁₃R₁₄ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0069] "-C₀₋₈-C(O)NR₁₃R₁₄" means that the carbonyl in -C(O)NR₁₃R₁₄ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0070] "-C₀₋₈-N(R₁₃)-C(O)R₁₂" means that the nitrogen atom in -N(R₁₃)-C(O)R₁₂ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0071] "-C₀₋₈-N(R₁₃)-C(O)OR₁₁" means that the nitrogen atom in -N(R₁₃)-C(O)OR₁₁ is bonded to C₀₋₈ alkyl, wherein C₀ alkyl means a bond, and C₁₋₈ alkyl is as defined above.

[0072] "C₁₋₈ haloalkyl" means a alkyl group having 1 to 8 carbon atoms, wherein any hydrogen atom on which is optionally substituted with F, Cl, Br or I, and includes, but is not limited to difluoromethyl, dichloromethyl, dibromomethyl, trifluoromethyl, trichloromethyl, tribromomethyl, and the likes.

[0073] "C₁₋₈ haloalkoxy" means an alkoxy having 1 to 8 carbon atoms, wherein any hydrogen atom on which is optionally substituted with F, Cl, Br or I, and includes, but is not limited to

difluoromethoxy, dichloromethoxy, dibromomethoxy, trifluoromethoxy, trichloromethoxy, tribromomethoxy, and the likes.

[0074] "Halogen" means F, Cl, Br or I. "THF" refers to tetrahydrofuran. "EA/EtOAc" refers to ethyl acetate. "MeOH" means methanol. "EtOH" refers to ethanol. "PivOH" refers to trimethylacetic acid. "DMSO" refers to dimethyl sulfoxide. "DMF" means N,N-dimethylformamide. "DIPEA" refers to diisopropylethylamine. "CH₃CN" means acetonitrile. "PE" means petroleum ether. "DCM/CH₂Cl₂" means dichloromethane. "DCE" refers to dichloroethane. "DMA" refers to dimethylacetamide. "Et₃N" refers to triethylamine. "NH₄Cl" means ammonium chloride. "NMP" refers to N-methylpyrrolidone. "HOAc" refers to acetic acid. "TFA" refers to trifluoroacetic acid. "MeI" means methyl iodide. "KI" means potassium iodide. "MsCl" refers to methylsulfonyl chloride. "SO₂Cl₂" means sulfonyl chloride. "POCl₃" refers to phosphorus oxychloride. "MeONa" means sodium methoxide. "NaHCO₃" refers to sodium bicarbonate. "Na₂SO₄" means sodium sulfate. "K₂CO₃" means potassium carbonate. "NaN₃" refers to sodium azide. "NaH" refers to sodium hydride. "CuI" refers to cuprous iodide. "PPA" refers to polyphosphoric acid. "m-CPBA" refers to m-chloroperoxybenzoic acid. "MnO₂" means manganese dioxide. "LiAlH₄" means lithium aluminum hydride. "LiOH" refers to lithium hydroxide. "NaOAc" refers to sodium acetate. "NaNO₂" refers to sodium nitrite. "AgNO₃" means silver nitrate. "Boc₂O" refers to di-tert-butyl dicarbonate. "LiCl" means lithium chloride. "Zn(CN)₂" refers to zinc cyanide. "IBX" means 2-iodoxybenzoic acid. "Pd/C" means palladium carbon. "Pd(OAc)₂" means palladium acetate. "PPh₃" means triphenylphosphine. "Pd(PPh₃)₂Cl₂" means palladium bis(triphenylphosphine) dichloride. "Pd₂(dba)₃" means tris(dibenzylideneacetone) dipalladium. "Pd(dppf)Cl₂" means [1,1'-bis(diphenylphosphino)ferrocene] palladium dichloride. "Pd(PPh₃)₄" means tetrakis(triphenylphosphine) palladium. "brett-phos" refers to dicyclohexyl[3,6-dimethoxy-2',4',6'-triisopropyl [1,1'-biphenyl]-2-yl]phosphine.

[0075] "Optional" or "optionally" means that the event or environment subsequently described may, but need not, occur, including where the event or environment occurs or does not occur. For example, "heterocyclyl optionally substituted by alkyl" means that an alkyl group may be, but is not necessarily, present, and the description includes the case where the heterocyclyl is substituted with an alkyl and the case where the heterocyclyl is not substituted with an alkyl.

[0076] "Substituted" means that one or more hydrogen atoms in a group are each independently substituted with a corresponding number of substituents. It goes without saying that a substituent is only in its possible chemical position, and those skilled in the art will be able to determine (by experiment or theory) possible or impossible substitution without undue efforts. For example, it may be unstable that an amino group or a hydroxyl group having a free hydrogen is attached with a carbon atom having an unsaturated bond (such as an olefin).

[0077] "Pharmaceutical composition" means a mixture comprising one or more of the compounds described herein, or a physiologically/pharmaceutically acceptable salt or pro-drug

thereof, and other chemical components, for example physiological/pharmaceutically acceptable carriers and excipients. The purpose of the pharmaceutical composition is to promote the administration to an organism, which facilitates the absorption of the active ingredient thereby exerting biological activities.

The present invention will be further described in detail below in conjunction with the embodiments which is not intended to limit the present invention. The present invention is also not limited to the contents of the embodiments, but defined by the claims.

[0078] The structure of the compound of the present invention is determined by nuclear magnetic resonance (NMR) or/and liquid chromatography-mass spectrometry (LC-MS). The NMR chemical shift (δ) is given in parts per million (ppm). The NMR is measured by a Bruker AVANCE-400 nuclear magnetic apparatus, and the solvent is deuterated dimethyl sulfoxide (DMSO- d_6), deuterated methanol (CD₃OD) and deuterated chloroform (CDCl₃), and the internal standard is tetramethylsilane (TMS).

[0079] The measurement of LC-MS is performed by using an Agilent 6120 mass spectrometer. The measurement of HPLC is performed by using an Agilent 1200 DAD high pressure liquid chromatograph (Sunfire C18 150 × 4.6 mm column) and a Waters 2695-2996 high pressure liquid chromatograph (Gimini C18 150 × 4.6 mm column).

[0080] The thin layer chromatography silica gel plate is Yantai Yellow Sea HSGF254 or Qingdao GF254 silica gel plate. The specification of TLC is 0.15 mm - 0.20 mm, and the specification for thin layer chromatography separation and purification is 0.4 mm - 0.5 mm. 200-300 mesh silica gel (Yantai Huanghai silica gel) as a carrier is generally used in column chromatography.


[0081] The starting materials in the examples of the present invention are known and commercially available or can be synthesized according to methods known in the art.

[0082] Unless otherwise stated, all reactions of the present invention are carried out under continuous magnetic stirring under a dry nitrogen or argon atmosphere, the solvent is a dry solvent, and the unit of the reaction temperature is degrees Celsius.


Preparation of intermediates

Intermediate 1: Preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2H-pyran [3,2-c] pyridin-2-one

[0083]

**Step 1: preparation of ethyl 6-chloro-4-methoxynicotinate**


[0084]



[0085] Ethyl 4,6-dichloronicotinate (10.0 g, 45.4 mmol) was added to anhydrous THF (100 mL), and cooled to 0 °C with ice water, then MeONa (2.8 g, 51.8 mmol) was added. After the addition was completed, the mixture was stirred at room temperature overnight. After the reaction was completed, the mixture was concentrated to remove THF. The crude product was dissolved in ethyl acetate (100 mL), washed twice with water and dried over anhydrous Na₂SO₄, filtrated and concentrated to obtain ethyl 6-chloro-4-methoxynicotinate (8.2 g, yield: 84%). MS (ESI): *m/z* 216.3 [M+1]⁺.

Step 2: preparation of (6-chloro-4-methoxypyridin-3-yl)methanol


[0086]



[0087] Ethyl 6-chloro-4-methoxynicotinate (8.2 g, 38.1 mmol) was dissolved in anhydrous THF (200 mL), then LiAlH₄ (3.0 g, 81.1 mmol) was added under ice-water bath. After the addition was completed, the mixture was stirred at room temperature for 2 h. After the reaction was completed, 2N aqueous NaOH (25 mL) was added for extraction. The solid residue was removed by filtration, and the filtrate was concentrated to obtain compound (6-chloro-4-methoxypyridin-3-yl)methanol (6.0 g, yield: 91%). MS (ESI): *m/z* 174.2 [M+1]⁺.

Step 3: preparation of 6-chloro-4-methoxynicotinaldehyde


[0088]



[0089] (6-chloro-4-methoxypyridin-3-yl)methanol (6.0 g, 34.6 mmol) was dissolved in acetone (100 mL), then IBX (12.0 g, 42.9 mmol) was added. The mixture was heated to reflux for 18 h. After the reaction was completed, the mixture was filtered and concentrated to obtain compound 6-chloro-4-methoxynicotinaldehyde (4.2 g, yield: 71%). MS (ESI): m/z 172.2 $[M+1]^+$.

Step 4: preparation of 6-chloro-4-hydroxynicotinaldehyde

[0090]




[0091] 6-chloro-4-methoxynicotinaldehyde (4.2 g, 24.5 mmol) was dissolved in 1,4-dioxane (30 ml), then concentrated hydrochloric acid (10 mL) was added. The mixture was heated to 90 °C for 16 h. After the reaction was completed, the mixture was concentrated. The crude product was separated by silica gel column chromatography to obtain compound 6-chloro-4-hydroxynicotinaldehyde (1.5 g, yield: 39%).

[0092] ^1H NMR (400MHz, CDCl_3): δ ppm 11.37 (s, 1H), 9.99 (s, 1H), 8.57 (s, 1H), 6.99 (s, 1H).

Step 5: preparation of 7-chloro-3-(3,5-dimethoxyphenyl)-2H-pyrano[3,2-c]pyridin-2-one

[0093]




[0094] 6-chloro-4-hydroxynicotinaldehyde (0.60 g, 3.82 mmol) was added to acetic anhydride (10 mL), then 2-(3,5-dimethoxyphenyl)acetic acid (0.80 g, 4.08 mmol) and triethylamine (1.5 g,

10.7 mmol) were added, the mixture was heated to 110 °C for 40 min. After the reaction was completed, the mixture was cooled to room temperature and concentrated. The obtained solids were washed with petroleum ether/ethyl acetate (3:1) to obtain compound d 7-chloro-3-(3,5-dimethoxyphenyl)-2H-pyrano[3,2-c]pyridin-2-one (0.35 g, yield: 29%). MS (ESI): m/z 318.3 $[M+1]^+$.

Step 6: preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2H-pyrano[3,2-c]pyridin-2-one


[0095]



[0096] 7-chloro-3-(3,5-dimethoxyphenyl)-2H-pyrano[3,2-c]pyridin-2-one (0.35 g, 1.1 mmol) was dissolved in anhydrous acetonitrile (10 mL), the mixture was cooled to -30 °C, then SO_2Cl_2 (1.0 g, 7.4 mmol) was added dropwise, the mixture was stirred at this temperature for 1 h. A saturated aqueous solution of NaHCO_3 was added to quench the reaction, and then acetonitrile was removed by concentration. After filtration, the solids were washed with water and then petroleum ether/ethyl acetate (3:1) to obtain 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2H-pyrano[3,2-c]pyridin-2-one (0.21 g, yield: 49%). MS (ESI): m/z 386.3 $[M+1]^+$.

Intermediate 2: Preparation of 2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido [3,4-d]pyrimidine


[0097]



Step 1: preparation of 2-chloro-5-nitroisonicotinic acid

[0098]






[0099] Chromium trioxide (40.0 g, 40 mmol) was added to a solution of 2-chloro-4-methyl-5-nitropyridine (20.0 g, 11.6 mmol) in sulfuric acid (200 mL) at 0 °C. After the addition was completed, the mixture was stirred at 0 °C for 1 h, then slowly warmed to room temperature and stirred overnight, and then poured into ice water (1 L) and filtrated to obtain 2-chloro-5-nitroisonicotinic acid (18 g, yield: 77%). MS (ESI): m/z 201.1 $[M-1]^-$.

Step 2: preparation of methyl 2-chloro-5-nitroisonicotinate


[0100]



[0101] Oxalyl chloride (12.7 g, 100 mmol) was added to a suspension of 2-chloro-5-nitroisonicotinic acid (16 g, 80 mmol) in dichloromethane (150 mL) at 0 °C. After the addition was completed, the mixture was stirred at room temperature for 3 h. Methanol (100 mL) was added, the reaction mixture was stirred at room temperature for another 4 h and then concentrated. The crude product was dissolved in dichloromethane (200 mL), the mixture was washed with sodium bicarbonate solution (100 mL*2), dried over anhydrous sodium sulfate and concentrated to obtain methyl 2-chloro-5-nitroisonicotinate (17.2 g, yield: 98%).

Step 3: preparation of methyl 2-(3,5-dimethoxyphenyl)-5-nitroisonicotinate

[0102]




[0103] 3,5-dimethoxyphenylboronic acid (8.47 g, 46 mmol), tetrakis(triphenylphosphine)palladium (5 g, 4.6 mmol) and sodium carbonate (5 g, 16 mmol) were added to a solution of methyl 2-chloro-5-nitroisonicotinate (10.0 g, 46.0 mmol) in the mixture of dioxane (200 mL) and water (50 mL). After the addition was completed, the mixture

was stirred under N_2 at 110 °C until the reaction of the starting materials was completed. The reaction solution was concentrated and separated by column chromatography (eluent: CH_2Cl_2/PE 20:1) to obtain compound methyl 2-(3,5-dimethoxyphenyl)-5-nitroisonicotinate (6 g, yield: 41%). MS (ESI): m/z 318.9 $[M+1]^+$.

Step 4: preparation of methyl 5-amino-2-(3,5-dimethoxyphenyl)isonicotinate


[0104]



[0105] Palladium carbon (10%, 500 mg) was added to a solution of methyl 2-(3,5-dimethoxyphenyl)-5-nitroisonicotinate (6 g, 18.8 mmol) in methanol (100 mL). Then the mixture was stirred under a hydrogen atmosphere at room temperature for 4 h, filtrated and concentrated to obtain methyl 5-amino-2-(3,5-dimethoxyphenyl)nitroisonicotinate (5 g, yield: 92.3%). MS (ESI): m/z 289.3 $[M+1]^+$.

Step 5: preparation of 6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,4(1H,3H)-dione


[0106]



[0107] A mixture of methyl 5-amino-2-(3,5-dimethoxyphenyl)isonicotinate (5 g, 17.4 mmol) and urea (12 g, 200 mmol) was heated to 160 °C and stirred for 4 h. Then the mixture was poured into ice water (100 mL) and filtered to obtain 6-(3,5-dimethoxyphenyl)pyrido [3,4-d]pyrimidine-2,4(1H,3H)-dione (6 g, yield: 99%). MS (ESI): m/z 300.3 $[M+1]^+$.

Step 6: preparation of 2,4-dichloro-6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine


[0108]



[0109] N,N-diethylaniline (3 mL) was added to a suspension of 6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,4(1H,3H)-dione (5.0 g, 16.7 mmol) in phosphorus oxychloride (50 mL). The mixture was then stirred at 110 °C overnight. The solvent was evaporated, ice water (200 mL) was added, and pH was adjusted to 7 with aqueous sodium bicarbonate solution. The aqueous solution was extracted for three times with EtOAc (50 mL). The organic phase was dried over anhydrous sodium sulfate, filtered, concentrated and separated by column chromatography [eluent: (EA: PE = 1: 5)] to obtain compound 2,4-dichloro-6-(3,5 - dimethoxyphenyl)pyrido[3,4-d]pyrimidine (2.5 g, yield: 44.5%). MS (ESI): *m/z* 336.2 [M+ 1]⁺.

Step 7: preparation of 2-chloro-6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine


[0110]



[0111] A concentrated aqueous ammonia (2 g) was added to a solution of 2,4-dichloro-6-(3,5 - dimethoxyphenyl)pyrido[3,4-d]pyrimidine (2.5 g, 7.5 mmol) in methanol (100 mL). The mixture was then stirred at 25 °C for 4 h. The pH was adjusted to 7 with hydrochloric acid (1N), then methanol is removed under vacuum, 2-chloro-6-(3,5-dimethoxyphenyl)pyrido[3, 4-d]pyrimidine-4-amine (2 g, yield: 84%) was obtained after filtration. MS (ESI): *m/z* 317. 1[M+1]⁺.

Step 8: preparation of 2-chloro-6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine

[0112]






[0113] Tert-butyl nitrite (720 mg, 6.2 mmol) was added to a solution of 2-chloro-6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine (1 g, 3.1 mmol) in tetrahydrofuran (100 mL). The mixture was then heated to reflux and stirred for 48 h. The reaction mixture was concentrated and separated by column chromatography [eluent: (EA: PE = 1: 5 -1: 2)] to obtain compound 2-chloro-6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (400 mg, yield: 41%).

[0114] ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.7(s, 1H), 9.53(s, 1H), 8.74(s, 1H), 7.4(s, 2H), 6.64(s, 1H), 3.88(s, 6H);

[0115] MS (ESI): m/z 302.0 $[\text{M}+1]^+$.

Step 9: preparation of 2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine


[0116]



[0117] The compound was prepared referring to the synthesis method of step 6 of intermediate 1.

Intermediate 3 and Intermediate 4: Preparation of 2-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine and 2-chloro-6-(2-fluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine

[0118]



[0119] 2-chloro-6-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (120 mg, 0.4 mmol) was dissolved in anhydrous acetonitrile (20 mL), then the mixture was cooled to 0 °C with ice-water

bath. 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate) (select-fluor, 283 mg, 0.8 mmol) was added dropwise. After the addition was completed, the reaction solution was warmed to room temperature for 6 h. The reaction was completed monitored by TLC. A saturated aqueous NaHCO_3 solution was added to quench the reaction, and then majority of acetonitrile was removed. The mixture was extracted with ethyl acetate, concentrated, and separated by silica gel column chromatography (PE/EA=10/11, adjusted by adding 10% DCM) to obtain 2-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (20 mg, yield: 15%), MS m/z (ESI): 338 $[\text{M}+\text{H}]^+$. At the same time, 2-chloro-6-(2-fluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (70 mg, yield: 55%) was also obtained, MS m/z (ESI): 320 $[\text{M}+\text{H}]^+$.


[0120] Intermediates 5-10 were prepared referring to the the synthesis method of intermediate 2.

Intermediate No.	Compound structure	Compound name	MS: m/z $[\text{M}+1]^+$
5		2-chloro-6-(2-chloro-6-fluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine	355
6		2-chloro-6-(2-fluoro-6-isopropyl-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine	362
7		2-chloro-6-(2-cyclopropyl-6-fluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine	360
8		2-chloro-6-(2-chloro-6-cyclopropyl-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine	377
9		2-chloro-6-(6-fluoro-7-methoxy-2,3-dihydrobenzo[b][1,4]dioxin-5-yl)pyrido[3,4-d]pyrimidine	348
10		2-chloro-6-(6-chloro-7-methoxy-2,3-dihydrobenzo[b][1,4]dioxin-5-yl)pyrido[3,4-d]pyrimidine	365

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺


Intermediate 11: Preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1H-pyrano[4,3-c]pyridin-1-one

[0121]



Step 1: preparation of methyl 5-bromo-2-chloroisonicotinate

[0122]




[0123] 5-bromo-2-chloroisonicotinic acid (30.0 g, 12.6 mmol) was dissolved in methanol (300 mL), then SOCl₂ (18.0 g, 15 mmol) was added, the mixture was heated to 75 °C for 8h. The reaction was completed monitored by LCMS. The mixture was cooled to room temperature and concentrated by reduced pressure, and then EtOAc (300 mL) was added to the residue. The mixture was washed with a saturated aqueous solution of NaHCO₃, dried over anhydrous Na₂SO₄, filtered and concentrated to obtain methyl 5-bromo-2-chloroisonicotinate (32.0 g, yield: 99%). MS (ESI): *m/z* 251.9 [M+1]⁺.

Step 2: preparation of methyl 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinate

[0124]






[0125] Methyl 5-bromo-2-chloro isonicotinate (30 g, 120 mmol) was dissolved in 1,4-dioxane (300 mL), and 3,5-dimethoxyphenylacetylene (20.4 g, 120 mmol), CuI (2.28 g, 12 mmol), Pd(dppf)Cl₂ (4.2 g, 6 mmol) and Et₃N (12.0 g, 120 mmol) were added, and the mixture was heated to 60 °C under N₂ for 6 h. The reaction was complete, and the mixture was filtered and concentrated. The crude product was separated by silica gel column chromatography (DCM: PE = 20: 1) to obtain compound methyl 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinate (22 g, yield: 55%). MS (ESI): *m/z* 332.1 [M+1]⁺.

Step 3: preparation of 7-chloro-3-(3,5-dimethoxyphenyl)-1H-pyrano[4,3-c]pyridin-1-one


[0126]



[0127] Methyl 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinate (21.0 g, 63 mmol) was added to PPA (200 mL), and the mixture was heated to 80 °C for 8 h. The reaction was completed monitored by LCMS. The reaction mixture was poured into ice water (1000 mL), and then filtered to obtain solid compound 7-chloro-3-(3,5-dimethoxyphenyl)-1H-pyrano[4,3-c]pyridin-1-one (15.0 g, yield: 75%). MS (ESI): *m/z* 318.2 [M+1]⁺.

Step 4: preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1H-pyrano[4,3-c]pyridin-1-one


[0128]



[0129] The compound was prepared referring to the synthetic method of step 6 of Intermediate 1.


Intermediate 12: Preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one

[0130]



Step 1: preparation of 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-1(2H)-one

[0131]




[0132] 7-chloro-3-(3,5-dimethoxyphenyl)-1H-pyrano[4,3-c]pyridin-1-one (15 g, 47.3 mmol) was dissolved in DMF (200 mL), then concentrated aqueous ammonia (150 mL) was added. The reaction mixture was heated to 80 °C for 48 h. The reaction was completed monitored by LCMS, and then filtered to obtain solid compound 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-1(2H)-one (7.5 g, yield: 50.5%).



[0133] ^1H NMR (400MHz, DMSO- d_6): δ ppm 11.7(m, 1H), 8.97(s, 1H), 8.01(s, 1H), 7.14(s, 1H), 6.97(s, 2H), 6.61(s, 1H), 3.86(s, 6H).

[0134] MS (ESI): m/z 317.2 $[\text{M}+1]^+$.

Step 2: preparation of 7-chloro-3-(3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one

[0135]






[0136] 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-1(2H)-one (2g, 6.32 mmol) was dissolved in DMF (30 mL), and NaH (758 mg, 18.95 mmol) was added under ice water bath. The mixture was stirred at 0 °C for 15 minutes, then MeI (8.967 g, 63.151 mmol) was added dropwise, the mixture was stirred at room temperature for 1 h. After the reaction was completed, it was quenched with water. The mixture was extracted for three times with ethyl acetate, and the organic phases were combined and washed with water, saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography (Eluent: PE/EtOAc = 2:1) to obtain compound 7-chloro-3-(3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one (1.35g, yield: 65%).

[0137] MS m/z (ESI): 331.0 [M+H]⁺.

Step 3: preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one







[0138]



[0139] The compound was prepared referring to the synthetic method of step 6 of Intermediate 1. MS m/z (ESI): 399.2 [M+H]⁺.


[0140] Intermediate 13-19 were prepared referring to the synthesis method of Intermediate 12.


Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
13		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1(2 H)-one	385
14		7-chloro-2-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1(2H)-one	440

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
15		N-(tert-butyl)-2-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-oxo-2,6-naphthyridin-2(1H)-yl)acetamide	499
16		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(2-methoxyethyl)-2,6-naphthyridin-1(2H)-one	444
17		2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-8(7H)-one	387
18		2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)-7-methylpyrido[3,4-d]pyrimidin-8(7H)-one	401
19		2-chloro-7-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-8(7H)-one	441

Intermediate 20: Preparation of 6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4(3H)-one


[0141]





Step 1: preparation of 5-amino-2-chloroisonicotinamide


[0142]



[0143] 5-amino-2-chloroisonicotinic acid (4.0 g, 23 mmol) was added to SOCl_2 (50 mL), the mixture was heated to 80 °C for 4 h. After the reaction was completed, the mixture was cooled to room temperature and concentrated. The residue was dissolved in anhydrous THF (20 mL), and then the mixture was cooled under ice water bath, a concentrated aqueous ammonia (100 mL) was added. The reaction was completed, and the mixture was extracted for five times with CH_2Cl_2 (100 mL). The organic phases were combined, dried over anhydrous sodium sulfate, filtrated and concentrated to obtain compound 5-amino-2-chloroisonicotinamide (3.5 g, yield: 88%).

Step 2: preparation of 6-chloro-2-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4(3H) - one

[0144]




[0145] 5-amino-2-chloroisonicotinamide (3.5 g, 20.4 mmol) and 3,5-dimethoxybenzaldehyde (3.7 g, 22.3 mmol) were dissolved in EtOH (50 mL), then HOAc (10 mL) was added, the mixture was heated to 80 °C for 2 days. The reaction liquid was concentrated, the residue was dissolved in the mixture of CH_2Cl_2 (100 mL) and THF (100 mL). Then MnO_2 (17.0 g, 87.0 mmol) was added, and the mixture was stirred at room temperature for 4 days. The reaction was completed, the mixture was filtered and concentrated. The residue was added to EtOAc (100 mL), stirred and filtrated to obtain compound 6-chloro-2-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4(3H)-one (4.0 g, yield: 63%).

[0146] ^1H NMR (400MHz, DMSO- d_6): δ ppm 12.78 (s, 1H), 8.91 (s, 1H), 7.95 (s, 1H), 7.44 (s, 1H), 7.43 (s, 1H), 3.84 (s, 6H).

Step 3: preparation of 6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4(3H)-one


[0147]



[0148] 6-chloro-2-(3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4(3H)-one (1.0 g, 3.2 mmol) was dissolved in CH_3CN (50 mL), then the mixture was cooled to $-20\text{ }^\circ\text{C}$, SO_2Cl_2 (0.85 g, 6.3 mmol) was added, the mixture was reacted at this temperature for 3 h. After the reaction was completed, a saturated aqueous solution of NaHCO_3 was added to quench the reaction, and the mixture was filtrated to obtain compound 6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4(3H)-one (1.1 g, 92%). MS (ESI): m/z 386 $[\text{M}+\text{H}]^+$.

Intermediate 21: Preparation of 6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-3-methylpyrido[3,4-d]pyrimidine-4(3H)-one


[0149]



[0150] 6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4(3H)-one (0.65 g, 1.68 mmol) was dissolved in DMF (20 mL), then the mixture was cooled to $0\text{ }^\circ\text{C}$ under ice water bath, NaH (134 mg, 3.36 mmol) was added, the mixture was stirred at room temperature for 2 h. The reaction was completed, water (50 mL) was added to quench the reaction. The mixture was filtrated, washed with water and dried to obtain compound 6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-3-methylpyrido[3,4-d]pyrimidine-4(3H)-one (0.65 g, yield:96%). MS m/z (ESI): 400 $[\text{M}+\text{H}]^+$.


Intermediate 22: Preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(2,2, 2-trifluoroethyl)-1,6-naphthyridin-2(1H)-one

[0151]



Step 1: preparation of ethyl 6-chloro-4-((2,2,2-trifluoroethyl)amino)nicotinate


[0152]



[0153] Ethyl 4,6-dichloronicotinate (4.0 g, 18.18 mmol) and trifluoroethylamine (2.7 g, 27.27 mmol) were dissolved in DMSO (50 mL), then Et₃N (5.5 g, 54.55 mmol) was added, the mixture was heated to 120 °C for 12 h. The reaction was completed, the mixture was cooled to room temperature, water (200 mL) was added, and then the mixture was extracted for three times with EtOAc (50 mL). The organic phase was dried over anhydrous sodium sulfate, filtrated and concentrated. The crude product was separated by silica gel column chromatography (PE:EA=3:1) to obtain compound ethyl 6-chloro-4-((2,2,2-trifluoroethyl)amino)nicotinate (1.5 g, yield: 29.1%). MS (ESI): *m/z* 283.0 [M+1]⁺.

Step 2: preparation of (6-chloro-4-((2,2,2-trifluoroethyl)amino)pyridin-3-yl)methanol

[0154]




[0155] Ethyl 6-chloro-4-((2,2,2-trifluoroethyl)amino)nicotinate (1.5 g, 5.32 mmol) was dissolved in dried THF (30 mL), then the mixture was cooled under ice water bath. LiAlH₄ (0.39 g, 10.64

mmol) was slowly added, and the mixture was stirred at 0 °C for 2 h. The reaction was completed, $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ was added to quench the reaction. The mixture was filtrated, and the filtrate was concentrated to obtain compound (6-chloro-4-((2,2,2-trifluoroethyl)amino)pyridin-3-yl)methanol (1.1 g, yield: 86%).

Step 3: preparation of 6-chloro-4-((2,2,2-trifluoroethyl)amino)nicotinaldehyde


[0156]



[0157] (6-chloro-4-((2,2,2-trifluoroethyl)amino)pyridin-3-yl)methanol (1.1 g, 4.58 mmol) was dissolved in the mixture of CH_2Cl_2 and THF (30 mL/10 mL), then MnO_2 (4.78 g, 54.9 mmol) was added, and the mixture was stirred at room temperature for 12 h. The reaction was completed, filtrated, and the filtrate was concentrated to obtain crude product 6-chloro-4-((2,2,2-trifluoroethyl)amino)nicotinaldehyde (0.86 g, yield: 79%).

Step 4: preparation of 7-chloro-3-(3,5-dimethoxyphenyl)-1-(2,2,2-trifluoroethyl)-1,6-naphthyridin-2(1H)-one

[0158]




6-chloro-4-((2,2,2-trifluoroethyl)amino)nicotinaldehyde (0.86 g, 3.61 mmol) was dissolved in DMF (20 mL), then methyl 3,5-dimethoxyphenylacetate (760 mg, 3.61 mmol) and K_2CO_3 (1.5 g, 10.84 mmol) were added, the mixture was heated to 110 °C for 3 h. After the reaction was completed, the mixture was concentrated and separated by silica gel column chromatography (PE: EA = 10: 1) to obtain 7-chloro-3-(3,5-dimethoxyphenyl)-1-(2,2,2-trifluoroethyl)-1,6-naphthyridin-2(1H)-one (960 mg, yield: 67%). MS (ESI): m/z 399.0 $[\text{M} + 1]^+$.

Step 5: preparation of 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(2,2,2-trifluoroethyl)-1,6-naphthyridin-2(1H)-one

[0159]






[0160] The compound was prepared referring to the synthetic method of step 6 of Intermediate 1. Intermediates 23-25 were prepared according to the synthesis method of Intermediate 22.

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
23		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-1,6-naphthyridin-2(1H)-one	400
24		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(tetrahydrofuran-3-yl)-1,6-naphthyridin-2(1H)-one	456
25		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(2-morpholinoethyl)-1,6-naphthyridin-2(1H)-one	499


Intermediate 26: Preparation of 3-chloro-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine

[0161]



Step 1: preparation of 5-bromo-2-chloroisonicotinaldehyde

[0162]




[0163] Diisopropylamine (8.95 g, 88 mmol) was dissolved in dried THF (100 mL) under N₂, then the mixture was cooled to -78 °C. N-butyllithium (50 mL, 78 mmol) was added dropwise, and then the mixture was stirred at 0 °C for 10 min. The mixture was cooled to -78 °C, then 2-chloro-5-bromopyridine (10.0 g, 52 mmol) was added, the mixture was stirred at -78 °C for 1 h, then dried DMF (11.4 g, 0.16 mol) was added dropwise, the mixture was stirred at this temperature for another 1 h. A ammonium chloride solution was added to quench the reaction, and the mixture was diluted with EtOAc (300 mL), washed for three times with water (50 mL), washed with saturated brine (100 mL), dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography (Eluent: PE/EtOAc 9:1) to obtain compound 5-bromo-2-chloroisonicotinaldehyde (7.0 g, yield: 61%).

[0164] ¹H NMR (400 MHz, CDC13): δ 10.3(s, 1H), 8.69 (s, 1H), 7.73 (s, 1H).

[0165] MS m/z (ESI): 252.0, 254.0, 256.0 [M+MeOH+H]⁺.

Step 2: preparation of 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinaldehyde

[0166]




[0167] 5-bromo-2-chloroisonicotinaldehyde (2.5 g, 11.3 mmol), 1-ethynyl-3,5-dimethoxybenzene (1.93 g, 11.9 mmol), DIPEA (3.66 g, 28.4 mmol), CuI (108 mg, 0.6 mmol), and Pd(PPh₃)₂Cl₂ (398 mg, 0.6 mmol) were added to 1,4-dioxane (50 mL) under N₂, the mixture was heated to 50 °C for 1 h. After the reaction was completed, the mixture was diluted with EtOAc (300 mL), washed successively with water (50 mL × 3) and saturated brine (100 mL), concentrated and separated by column chromatography (Eluent: PE/EtOAc 9:1) to obtain compound 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinaldehyde (3.0 g, yield: 75%). MS m/z (ESI): 302.2, 304.2 [M+H]⁺.

Step 3: preparation of 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinaldehyde

xime


[0168]



[0169] 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinaldehyde (0.6 g, 2 mmol), NaOAc (245 mg, 3.0 mmol) and hydroxylamine hydrochloride (207 mg, 3.0 mmol) were dissolved in the mixture of ethanol and 1,2-dichloroethane (20 mL/11.2 mL) under N_2 , the mixture was heated to 50 °C for 50 min. After the reaction was completed, the mixture was diluted with EtOAc (200 mL), washed successively with water (50 mL $\times$ 2) and saturated brine (80 mL), dried over anhydrous sodium sulfate, filtrated and concentrated to obtain compound 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinaldehydeoxime (612 mg, yield: 100%).

Step 4: preparation of 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-2-oxide

[0170]




[0171] 2-chloro-5-((3,5-dimethoxyphenyl)ethynyl)isonicotinaldehydeoxime (612 mg, 1.9 mmol) and $AgNO_3$ (66 mg, 0.38 mmol) were added to chloroform (20 mL), the mixture was heated to 60 °C for 1 h. After the reaction was completed, the mixture was concentrated and separated by column chromatography (Eluent: CH_2Cl_2 /MeOH 35:1) to obtain compound 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-2-oxide (580 mg, yield: 95%).

[0172] MS m/z (ESI): 317.2, 319.2 $[M+H]^+$.

Step 5: preparation of 3-chloro-7-(3,5-dimethoxyphenyl)-2,6-naphthyridine

[0173]






[0174] 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-2-oxide (200 mg, 0.63 mmol) was dissolved in dichloromethane (10 mL) under ice water bath, then phosphorus trichloride (0.7 mL, 1.4 mmol) was added, the mixture was stirred at room temperature overnight. The reaction was completed, the mixture was washed successively with a saturated aqueous solution of NaHCO_3 (20 mL), DCM (80 mL), saturated brine (50 mL), dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography (Eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 50:1) to obtain compound 3-chloro-7-(3,5-dimethoxyphenyl)-2,6-naphthyridine (60 mg, yield: 32%). MS m/z (ESI): 301.2, 303.2 $[\text{M}+\text{H}]^+$.

Step 6: preparation of 3-chloro-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine

[0175]




[0176] The compound was prepared referring to the synthesis method of Intermediate 1. Intermediates 27-32 were prepared referring to the synthesis method of Intermediate 26.

Intermediate No.	Compound structure	Compound name	MS: m/z $[\text{M}+1]^+$
27		3-chloro-7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridine	337
28		3-chloro-7-(2-chloro-3,5-dimethoxyphenyl)-2,6-naphthyridine	336
29		3-chloro-7-(2-chloro-6-fluoro-3,5-dimethoxyphenyl)-2,6-naphthyridine	354

Intermediate No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
30		3-chloro-7-(2-chloro-3,5-dimethoxy-6-methylphenyl)-2,6-naphthyridine	350
31		3-chloro-7-(2-chloro-6-isopropyl-3,5-dimethoxyphenyl)-2,6-naphthyridine	378
32		3-chloro-7-(2-chloro-6-cyclopropyl-3,5-dimethoxyphenyl)-2,6-naphthyridine	376


Intermediate 33: Preparation of 7-chloro-N-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine

[0177]



Step 1: preparation of 7-chloro-1-((cyclopropylmethyl)amino)-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine 2-oxide

[0178]




[0179] 7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine 2-oxide (4.0 g, 12.6 mmol) was dissolved in toluene (80 mL), cyclopropylmethylamine (7.19 g, 0.1 mol) and CuI (241 mg, 1.26 mmol) were added successively, the mixture was heated to 50 °C overnight under an oxygen atmosphere. After the reaction was completed, the reaction liquid was filtrated and concentrated. The residue was separated by column chromatography (Eluent: dichloromethane/methanol 25:1) to obtain compound 7-chloro-1-((cyclopropylmethyl)amino)-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine 2-oxide (2.8g, 57%).

[0180] MS m/z (ESI): 386.4 [M+H]⁺.

Step 2: preparation of 7-chloro-N-(cyclopropylmethyl)-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine

[0181]




[0182] 7-chloro-1-((cyclopropylmethyl)amino)-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine 2-oxide (2.8 g, 5.1 mmol) was dissolved in dichloromethane (40 mL), PCl₃ (3 mL, 6.1 mmol) was added dropwise under ice water bath, the mixture was stirred at room temperature for 1 h. A saturated solution of sodium hydrogencarbonate (50 mL) was added, then the mixture was extracted with dichloromethane (100 mL × 3), the organic phases were combined, dried over anhydrous sodium sulfate, concentrated and separated by column chromatography (Eluent: dichloromethane/methanol 50:1) to obtain compound 7-chloro-N-(cyclopropylmethyl)-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine (570 mg, 21%).

[0183] MS m/z (ESI): 370.4 [M+H]⁺.

Step 3: preparation of 7-chloro-N-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine

[0184]





[0185] 7-chloro-N-(cyclopropylmethyl)-3-(3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine (550 mg, 0.27 mmol) was added to acetonitrile (20 mL), then sulfonyl chloride (73 mg, 0.54 mmol) was added dropwise at $-30\text{ }^\circ\text{C}$, the mixture was stirred at this temperature for 1 h. A saturated solution of sodium hydrogencarbonate (30 mL) was added, then the mixture was extracted with ethyl acetate (50 mL $\times$ 3), the organic phases were combined, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography (Eluent: petroleum ether/ethyl acetate 10:1) to obtain compound 7-chloro-N-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine (337 mg, 52%).








[0186] MS m/z (ESI+APCI): 438.2/440.2 $[\text{M}+\text{H}]^+$.

[0187] Intermediates 34-131 were prepared referring to the synthesis method of Intermediate 33.

Intermediate No.	Compound structure	Compound name	MS: m/z $[\text{M}+1]^+$
34		7-chloro-N-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	438
35		7-chloro-N-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-di(methoxy- d_3)phenyl)-2,6-naphthyridine-1-amine	444
36		7-chloro-N-(cyclopropylmethyl)-3-(2,6-difluoro-3,5-di(methoxy- d_3)phenyl)-2,6-naphthyridine-1-amine	412
37		7-chloro-N-(cyclopropylmethyl)-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	406

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
38		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-methoxyethyl)-2,6-naphthyridine-1-amine	442
39		2-((7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)amino)ethan-1-ol	428
40		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(isopropylthio)ethyl)-2,6-naphthyridine-1-amine	486
41		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(ethylsulfonyl)ethyl)-2,6-naphthyridine-1-amine	504
42		N ¹ -(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-N ² ,N ² -dimethylethane-1,2-diamine	455
43		N ¹ -(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-N ³ ,N ³ -dimethylpropane-1,3-diamine	469
44		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-methyl-2,6-naphthyridine-1-amine	398
45		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-methyl-2,6-naphthyridine-1-amine	412

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		xyphenyl)-N-ethyl-2,6-naphthyridine-1-amine	
46		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2,2,2-trifluoroethyl)-2,6-naphthyridine-1-amine	466
47		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N,N-dimethyl-2,6-naphthyridine-1-amine	412
48		7-chloro-N-(cyclopropylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-methyl-2,6-naphthyridine-1-amine	452
49		N-(2-((7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)amino)ethyl)methanesulfonamide	505
50		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((tetrahydrofuran-2-yl)methyl)-2,6-naphthyridine-1-amine	468
51		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((tetrahydrofuran-3-yl)methyl)-2,6-naphthyridine-1-amine	468








Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
52		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(oxetan-3-ylmethyl)-2,6-naphthyridine-1-amine	454
53		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((tetrahydro-2H-pyran-4-yl)methyl)-2,6-naphthyridine-1-amine	482
54		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(oxetan-3-yl)-2,6-naphthyridine-1-amine	440
55		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(tetrahydro-2H-pyran-4-yl)-2,6-naphthyridine-1-amine	468
56		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(tetrahydrofuran-3-yl)-2,6-naphthyridine-1-amine	454
57		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-3-yl)methyl)-2,6-naphthyridine-1-amine	481
58		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-2-yl)methyl)-2,6-naphthyridine-1-amine	481

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
59		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(1-methylpyrrolidin-3-yl)-2,6-naphthyridine-1-amine	467
60		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylazetidin-3-yl)methyl)-2,6-naphthyridine-1-amine	467
61		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(1-methylazetidin-3-yl)-2,6-naphthyridine-1-amine	453
62		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(1-methylpiperidin-4-yl)-2,6-naphthyridine-1-amine	481
63		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylpiperidin-4-yl)methyl)-2,6-naphthyridine-1-amine	495
64		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-difluorocyclobutyl)-2,6-naphthyridine-1-amine	474
65		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-difluorocyclopentyl)-2,6-naphthyridine-1-amine	488








Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
66		7-chloro-N-(cyclopentylmethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	466
67		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-phenethyl-2,6-naphthyridine-1-amine	488
68		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)-2,6-naphthyridine-1-amine	478
69		2-(4-(((7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)amino)methyl)-1H-pyrazol-1-yl)ethan-1-ol	508
70		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)-2,6-naphthyridine-1-amine	522
71		N-benzyl-7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	474
72		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(4-methylpiperazin-1-yl)ethyl)-2,6-naphthyridine-1-amine	510







Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
73		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-morpholinoethyl)-2,6-naphthyridine-1-amine	497
74		2-((7-chloro-3-(2-chloro-3-methoxyphenyl)-2,6-naphthyridin-1-yl)amino)ethan-1-ol	364
75		2-((7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)amino)ethan-1-ol	360
76		2-((7-chloro-3-(2-fluoro-3-methoxyphenyl)-2,6-naphthyridin-1-yl)amino)ethan-1-ol	348
77		N ¹ -(7-chloro-3-(2-chloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-N ² ,N ² -dimethylethane-1,2-diamine	421
78		N ¹ -(7-chloro-3-(3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-N ² ,N ² -dimethylethane-1,2-diamine	387
79		N ¹ -(7-chloro-3-(2-fluoro-3-methoxyphenyl)-2,6-naphthyridin-1-yl)-N ² ,N ² -dimethylethane-1,2-diamine	375
80		7-chloro-3-(2-fluoro-3,5-dimethoxyphenyl)-N-((tetrahydrofuran-2-yl)methyl)-2,6-naphthyridine-1-amine	418









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		-2,6-naphthyridine-1-amine	
81		7-chloro-3-(2-fluoro-3-methoxyphenyl)-N-((tetrahydrofuran-2-yl)methyl)-2,6-naphthyridine-1-amine	388
82		(S)-7-chloro-N-(1-cyclopropylethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	452
83		(S)-7-chloro-N-(1-cyclopropylethyl)-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	420
84		7-chloro-N-cyclopropyl-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	424
85		7-chloro-N-cyclopropyl-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridine-1-amine	392
86		1-(azetidin-1-yl)-7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine	424
87		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3-methoxyazetidin-1-yl)-2,6-naphthyridine	454

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
88		7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-(3-methoxyazetidin-1-yl)-2,6-naphthyridine	422
89		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-N,N-dimethylazetidin-3-amine	467
90		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3-(trifluoromethyl)azetidin-1-yl)-2,6-naphthyridine	492
91		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3,3-dimethylazetidin-1-yl)-2,6-naphthyridine	452
92		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3,3-difluoroazetidin-1-yl)-2,6-naphthyridine	460
93		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-3-methylazetidin-3-ol	454
94		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3-methoxy-3-methylazetidin-1-yl)-2,6-naphthyridine	468



Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
95		7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-(3-methoxy-3-methylazetidin-1-yl)-2,6-naphthyridine	436
96		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-3-methylazetidin-3-carbonitrile	463
97		6-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2-oxa-6-azaspiro[3.3]heptane	466
98		6-(7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2-oxa-6-azaspiro[3.3]heptane	434
99		6-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-1-oxa-6-azaspiro[3.3]heptane	466
100		6-(7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-1-oxa-6-azaspiro[3.3]heptane	434
101		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(2-azaspiro[3.3]hept-2-yl)-2,6-naphthyridine	464

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
102		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(2-azaspiro[3.4]oct-2-yl)-2,6-naphthyridine	478
103		2-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-6-oxa-2-azaspiro[3.4]octane	480
104		2-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-7-oxa-2-azaspiro[3.5]nonane	494
105		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(pyrrolidin-1-yl)-2,6-naphthyridine	438
106		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3-methoxypyrrolidin-1-yl)-2,6-naphthyridine	468
107		7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-(3-methoxypyrrolidin-1-yl)-2,6-naphthyridine	436
108		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)pyrrolidine-3-carbonitrile	463

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
109		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3,3-difluoropyrrolidin-1-yl)-2,6-naphthyridine	474
110		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3-methoxy-3-methylpyrrolidin-1-yl)-2,6-naphthyridine	482
111		7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-(3-methoxy-3-methylpyrrolidin-1-yl)-2,6-naphthyridine	450
112		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-3-methylpyrrolidine-3-carbonitrile	477
113		1-(3-azabicyclo[3.1.0]hex-3-yl)-7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine	450
114		7-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2-oxa-7-azaspiro[3.4]nonane	494
115		4-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)methyl	454


Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		orpholine	
116		4-(7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)morpholine	422
117		4-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2-methylmorpholine	468
118		4-(7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2-methylmorpholine	436
119		4-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2,6-dimethylmorpholine	482
120		4-(7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-2,6-dimethylmorpholine	450
121		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)piperidin-4-ol	468
122		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(4-methoxypiperidin-1-yl)-2,6-naphthyridine	482

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
123		7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-(4-methoxypiperidin-1-yl)-2,6-naphthyridine	450
124		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(3-methoxypiperidin-1-yl)-2,6-naphthyridine	482
125		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-4-methylpiperidin-4-ol	482
126		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(4-methoxy-4-methylpiperidin-1-yl)-2,6-naphthyridine	496
127		7-chloro-3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-(4-methoxy-4-methylpiperidin-1-yl)-2,6-naphthyridine	464
128		1-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)-4-methylpiperidine-4-carbonitrile	491
129		N-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)cyclopropylcarboxamide	452
130		N-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)tetrahydro-2H-pyran-2-one	482

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		rahydrofuran 2-carboxamide	
131		N-(7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)sulfonamide	462


Intermediate 132: Preparation of 6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methoxy-2-(methylthio)pyrido[3,4-d]pyrimidine

[0188]



Step 1: preparation of methyl 5-((3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylate

[0189]




[0190] Methyl 5-bromo-2-(methylthio)pyrimidine-4-carboxylate (2631 mg, 10.0 mmol) and 1-ethynyl-3,5-dimethoxybenzene (1622 mg, 10.0 mmol) were dissolved in dried THF (50 mL). The gas was exchanged with N₂, Et₃N (2.8 mL, 20.0 mmol), Pd(PPh₃)₂Cl₂ (702 mg, 1.0 mmol), PPh₃ (525 mg, 2.0 mmol) and CuI (190 mg, 1.0 mmol) were added successively under N₂, the mixture was heated to 90 °C and stirred overnight. After the reaction was completed, the

mixture was cooled to room temperature, added with a saturated aqueous solution of NaHCO_3 (100 mL), extracted twice with EtOAc (100 mL), the organic phase was washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated, and separated by silica gel column chromatography (PE:EA 4:1) to obtain compound methyl 5-((3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylate (2.5 g, yield: 73%). MS m/z (ESI): 345.2 $[\text{M}+\text{H}]^+$.

Step 2: preparation of 5-((3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylic acid


[0191]



[0192] Methyl 5-((3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylate (600 mg, 1.742 mmol) was dissolved in methanol (15 mL), and then an aqueous lithium hydroxide monohydrate (366 mg, 8.711 mmol) in water (5 mL) was added, and the mixture was stirred at room temperature overnight. The reaction was completed. The organic solvent was evaporated by reduced pressure, EtOAc (30 mL) was added to the residue, and the pH was adjusted with 1 N hydrochloric acid solution to 3~4. The organic phase was separated, washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, and concentrated to obtain compound 5-((3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylic acid (584 mg, quantitative yield). MS m/z (ESI): 331.2 $[\text{M}+\text{H}]^+$.

Step 3: preparation of 6-(3,5-dimethoxyphenyl)-2-(methylthio)-8H-pyrano[3,4-d]pyrimidine-8-one

[0193]




5-((3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylic acid (584 mg, 1.768 mmol) was suspended in acetone (25 mL), and AgNO_3 (180 mg, 1.059 mmol) was added to the suspension. The mixture was stirred at room temperature for 4h, and green solid was precipitated, filtered to obtain 6-(3,5-dimethoxyphenyl)-2-(methylthio)-8H-pyrano[3,4-d]pyrimidine-8-one.

d]pyrimidine-8-one (600 mg, the crude product) MS m/z (ESI): 331.2 [M+H]⁺.

Step 4: preparation of 6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8(7H)-one


[0194]



[0195] 6-(3,5-dimethoxyphenyl)-2-(methylthio)-8H-pyrano[3,4-d]pyrimidine-8-one (600 mg, the crude product) was added to glacial acetic acid (50 mL), then ammonium acetate (2.1 g, 27.243 mmol) was added, the mixture was heated to 115 °C for 16 h. The reaction was completed, the mixture was cooled to room temperature, poured slowly into a saturated aqueous sodium bicarbonate solution, and extracted with EtOAc (100 mL). The organic phase was separated, and washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography (Eluent: CH₂Cl₂/MeOH 0~6%) to obtain compound 6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8(7H)-one (298 mg, two-step yield: 50%). MS m/z (ESI): 330.2 [M+H]⁺.

Step 5: preparation of 6-(3,5-dimethoxyphenyl)-8-methoxy-2-(methylthio)pyrido[3,4-d]pyrimidine

[0196]




[0197] 6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8(7H)-one (100 mg, 0.304 mmol) and silver carbonate (109 mg, 0.395 mmol) were suspended in toluene (3 mL), then methyl iodide (431 mg, 3.036 mmol) was added. The mixture was heated to 100 °C in a sealed tube for 3 h. The reaction was completed, the mixture was cooled to room temperature, diluted with EtOA, and filtrated to remove salt. The filtrate was concentrated and separated by column chromatography (Eluent: PE/EtOAc 10 - 80%) to obtain compound 6-(3,5-dimethoxyphenyl)-8-methoxy-2-(methylthio)pyrido[3,4-d]pyrimidine (47 mg, yield: 45%).

MS m/z (ESI): 344.3 [M+H]⁺.

Step 6: preparation of 6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methoxy-2-(methylthio)pyrido [3,4-d] pyrimidine

[0198]




[0199] The compound was prepared referring to the synthesis method of Intermediate 1. Intermediates 133-134 were prepared by referring to the synthesis method of example 132.

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
133		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-ethoxy-2,6-naphthyridine	413
134		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-ethoxy-2-(methylthio)pyrido[3,4-d]pyrimidine	426


Intermediate 135: Preparation of 6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-methoxyethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine

[0200]



Step 1: preparation of 8-chloro-6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine

[0201]




[0202] 6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8(7H)-one (20.0 mg, 0.061 mmol) and N,N-diisopropylethylamine (78 mg, 0.610 mmol) were added to acetonitrile (2 mL), then POCl₃ (0.8 mL) was added, the mixture was heated to 90 °C and stirred overnight. The solvent was removed by reduced pressure. The residue was diluted with EtOAc (10 mL) and washed with a saturated sodium bicarbonate solution. The organic phase was separated, washed with saturated brine, dried over anhydrous sodium sulfate, filtered, concentrated and separated by column chromatography (Eluent: PE/EtOAc 0-40%) to obtain compound 8-chloro-6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (6 mg, yield: 28%).

[0203] MS m/z (ESI): 348.2 [M+H]⁺.

Step 2: preparation of 6-(3,5-dimethoxyphenyl)-N-(2-methoxyethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine

[0204]




[0205] 8-chloro-6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (6 mg, 0.017 mmol) and N,N-diisopropylethylamine (6.5 mg, 0.052 mmol) were dissolved in acetonitrile (1.5 mL), then 2-methoxyethyl-1-amine (4 mg, 0.052 mmol) was added, the mixture was heated to 90 °C and stirred overnight. The reaction was completed, the mixture was cooled to room temperature, diluted with EtOAc (5 mL), washed with saturated brine, dried over anhydrous sodium sulfate, filtered, concentrated and separated by column chromatography (Eluent: PE/EtOAc 0-50%) to obtain compound 6-(3,5-dimethoxyphenyl)-N-(2-methoxyethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine.

(methylthio)pyrido[3,4-d]pyrimidine-8-amine (4 mg, yield: 61%).

[0206] MS m/z (ESI): 387.4 $[M+H]^+$.

Step 3: preparation of 6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-methoxyethyl)-2-(methylthio)pyrido [3,4-d] pyrimidine-8-amine


[0207]



[0208] The compound was prepared referring to the synthesis method of Intermediate 1.


Intermediate 139: Preparation of 4-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)morpholine

[0209]



Step 1: preparation of 2,4-difluoro-3-iodo-1,5-dimethoxybenzene

[0210]




[0211] 2,6-difluoro-3,5-dimethoxyaniline (27.0 g, 143 mmol) was added to 6.0 M hydrochloric acid solution (240 mL), and NaNO_2 aqueous solution (10.35 g, 150 mmol, 30 mL water) was slowly added dropwise under ice water for cooling within 25 min. After the addition was

completed, the mixture was reacted for another 15 min to obtain an orange-red suspension, and then added to an aqueous KI solution (94.9 g, 570 mmol, 150 mL water), the mixture was heated to room temperature and stirred for 30 min to precipitate a solid. The mixture was filtrated and washed with water to obtain a crude product. MeOH (60 mL) was added to the crude product, then the mixture was stirred at room temperature for 30 min, filtrated, and dried to obtain 2,4-difluoro-3-iodo-1,5-dimethoxybenzene (29.3 g, yield: 68%).

Step 2: preparation of (2,6-difluoro-3,5-dimethoxyphenylacetylene)trimethylsilane


[0212]



[0213] 2,4-difluoro-3-iodo-1,5-dimethoxybenzene (25.8 g, 86.0 mmol), trimethylsilylacetylene (36.5 mL, 258 mmol), CuI (817 mg, 4.3 mmol) and triethylamine (35.8 mL, 258 mmol) were added to DMF (250 mL) under N₂, then Pd(PPh₃)₂Cl₂ (3.15 g, 4.3 mmol) was added, and the mixture was heated to 50 °C for 2 h. The reaction was completed, a saturated aqueous NH₄Cl solution was added to quench the reaction, the mixture was extracted for three times with dichloromethane, and the organic phases were combined, dried over Na₂SO₄, filtrated and concentrated to obtain a crude product (27.0 g) which was used directly in the next step.

Step 3: preparation of 3-ethynyl-2,4-difluoro-1,5-dimethoxybenzene


[0214]



[0215] (2,6-difluoro-3,5-dimethoxyphenylacetylene)trimethylsilane (27.0 g, a crude product) was added to the mixture of THF and MeOH (200/200 mL), then aqueous NaOH solution (8.6 mL, 8.6 mmol, 1.0 N) was added, and the mixture was stirred at room temperature for 15 min. The reaction was completed, a saturated aqueous NH₄Cl solution was added to quench the reaction, and the mixture was extracted for three times with dichloromethane, and the organic phases were combined, dried over anhydrous sodium sulfate, filtrated and concentrated. MeOH (50 mL) was added to the crude product, and then the mixture was stirred at room temperature for 30 min, filtered to obtain the target product (15.0 g, two-step yield: 88%).

Step 4: preparation of methyl 5-((2,6-difluoro-3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylate


[0216]



[0217] 3-ethynyl-2,4-difluoro-1,5-dimethoxybenzene (10.0 g, 50.5 mmol) and methyl 5-bromo-2-methylthio-pyrimidine-4-carboxylate (13.0 g, 49.5 mmol) were dissolved in DMF (100 mL), then CuI (479 mg, 2.52 mmol), Pd(PPh₃)₄ (2.91 g, 2.52 mmol) and Et₃N (35.0 mL, 252.5 mmol) were added, the mixture was heated to 100 °C for 1.5 h under N₂. The reaction was completed, the mixture was cooled to room temperature, a saturated aqueous NH₄Cl solution was added to quench the reaction, then the mixture was extracted for three times with dichloromethane, and the organic phases were combined, dried over anhydrous sodium sulfate, filtrated and concentrated to obtain a crude product, then the crude product was separated by silica gel column chromatography (PE : EA : DCM = 10 : 2 : 1) to obtain the target product (15.4 g, yield: 82%). ¹H NMR (400 MHz, CDCl₃) δ 8.82 (s, 1H), 6.69 (t, *J* = 8.0 Hz, 1H), 4.03 (s, 3H), 3.90 (s, 6H), 2.63 (s, 3H).

Step 5: preparation of 5-((2,6-difluoro-3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylic acid


[0218]



[0219] Methyl 5-((2,6-difluoro-3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylate (30.0 g, 78.9 mmol) was dissolved in THF (300 mL), then LiOH/H₂O (236.8 mL, 236.8 mmol, 1 M) was added, the mixture was stirred at room temperature for 2 h. The reaction was completed, the mixture was concentrated to remove THF, then acidified to pH 3 with diluted hydrochloric acid to precipitate a solid. The mixture was filtrated, washed with water, and dried to obtain the target product (28.5 g, yield: 99%).

Step 6: preparation of 6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)-8H-pyran o[3,4-d] pyrimidine-8-one


[0220]



[0221] 5-((2,6-difluoro-3,5-dimethoxyphenyl)ethynyl)-2-(methylthio)pyrimidine-4-carboxylic acid (2.5 g, 6.83 mmol) was dissolved in DCE (50 mL), then TFA (0.5 mL) was added, the mixture was heated to reflux overnight. The reaction was completed, the mixture was concentrated, MeOH (50 mL) was added, then the mixture was stirred at room temperature for 30 min. The mixture was filtrated, and the solid was washed with MeOH (10 mL) to obtain the target product (2.0 g, yield: 80%). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.23 (s, 1H), 7.22 (t, $J = 8.4$ Hz, 1H), 7.17 (s, 1H), 3.93 (s, 6H), 2.63 (s, 3H).

Step 7: preparation of 6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8(7H)-one


[0222]



[0223] The compound was prepared referring to the synthesis method of step 4 of Intermediate 132.

Step 8: preparation of 8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine


[0224]



[0225] 6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8(7H)-one (1 g, 2.74 mmol) was dissolved in DCE (80mL), the mixture was heated to 90 °C, then phenylphosphonic dichloride (3.0 mL, 21.92 mmol) was added, the mixture was heated and stirred for 16h, and then cooled. pH was adjusted to neutral under ice bath. The mixture was extracted with DCM, and then separated by silica gel column chromatography (MeO H/DCM 1/20) to obtain compound 8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (930 mg, yield:88%). MS m/z (ESI): 384 [M+H]⁺.

Step 9: preparation of N-(cyclopropylmethyl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido [3,4-d] pyrimidine-8-amine

[0226]









[0227] The compound was prepared referring to the synthesis method of step 2 of Intermediate 135.








[0228] Intermediates 136-330 were prepared referring to the synthesis method of Intermediate 135 or 139.

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
136		N-(cyclopropylmethyl)-6-(2, 6-dichloro-3,5 -dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	451
137		N-(cyclopropylmethyl)-6-(2, 6-dichloro-3,5-di(methoxy- <i>d</i> ₃)phenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	457
138		N-(cyclopropylmethyl)-6-(2,6-difluoro-3,5-di(methoxy- <i>d</i> ₃)phenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	425

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
140		2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)amino)ethanol-1-ol	441
141		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(isopropylthio)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	499
142		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(ethylsulfonyl)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	518
143		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-methyl-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	411
144		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-ethyl-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	425
145		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N,N-dimethyl-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	425
146		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(2,2,2-trifluoroethyl)pyrido[3,4-d]pyrimidine-8-amine	479
147		N-(cyclopropylmethyl)-6-(2,6-di	465








Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-amine	
148		N ¹ -(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-N ² ,N ² -dimethylethane-1,2-diamine	468
149		N ¹ -(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-N ³ ,N ³ -dimethylpropane-1,3-diamine	482
150		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)amino)ethyl)sulfonamide	518
151		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-((tetrahydrofuran-2-yl)methyl)pyrido[3,4-d]pyrimidin-8-amine	481
152		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-((tetrahydrofuran-3-yl)methyl)pyrido[3,4-d]pyrimidin-8-amine	481
153		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(oxetan-3-ylmethyl)pyrido[3,4-d]pyrimidin-8-amine	467









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
154		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-((tetrahydro-2H-pyran-4-yl)methyl)pyrido[3,4-d]pyrimidine-8-amine	495
155		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(oxetan-3-yl)pyrido[3,4-d]pyrimidine-8-amine	453
156		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidine-8-amine	481
157		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(tetrahydrofuran-3-yl)pyrido[3,4-d]pyrimidine-8-amine	467
158		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-3-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	494
159		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-2-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	494
160		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(1-methylpyrrolidin-3-yl)-2-(methylthio)pyrido[3,4-d]p	480









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		2-methyl-6-(2-methoxy-6-chlorophenyl)-4-methylthio-1,2,3,4-tetrahydropyrimidin-8-amine	
161		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylazetidin-3-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	480
162		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(1-methylazetidin-3-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	466
163		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(1-methylpiperidin-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	494
164		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methylpiperidin-4-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	508
165		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-difluorocyclobutyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	487
166		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-difluorocyclopentyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	501

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
167		N-(cyclopentylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	479
168		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-phenethylpyrido[3,4-d]pyrimidine-8-amine	501
169		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-phenylpyrido[3,4-d]pyrimidine-8-amine	473
170		N-benzyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	487
171		N-(3-aminobenzyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	502
172		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	491
173		2-(4-(((6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)amino)methyl)-1H-pyrazol-1-yl)ethan-1-ol	521

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
174		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	535
175		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-neopentylpyrido[3,4-d]pyrimidine-8-amine	467
176		N ¹ -(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-N ³ -isopropylpropane-1,3-diamine	496
177		N ¹ -(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-N ⁴ -isopropylbutane-1,4-diamine	510
178		N ¹ -(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-N ⁴ ,N ⁴ -dimethylbutane-1,4-diamine	496
179		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(4-(pyrrolidin-1-yl)butyl)pyrido[3,4-d]pyrimidine-8-amine	522
180		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(4-methylpiperazin-1-yl)ethyl)pyrido[3,4-d]pyrimidine-8-amine	523








Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		2-(methylthio)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-8-amine	
181		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(2-morpholinoethyl)pyrido[3,4-d]pyrimidine-8-amine	510
182		N-(2-(3-aminopyrrolidin-1-yl)ethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	509
183		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(3-(dimethylamino)pyrrolidin-1-yl)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	537
184		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-(2-(dimethylamino)ethoxy)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	512
185		(1-(4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)amino)butyl)pyrrolidin-2-yl)methanol	552
186		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(4-(3,3-difluoropyrrolidin-1-yl)butyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	558








Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
187		6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(4-(3-methoxypyrrolidin-1-yl)butyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	552
188		N-(cyclopropylmethyl)-6-(2-fluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	401
189		N-(cyclopropylmethyl)-6-(2-fluoro-3-methoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	371
190		6-(2-chloro-3-(methoxy-d ₃)phenyl)-N-(cyclopropylmethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	390
191		N-(cyclopropylmethyl)-6-(2-fluoro-3-(methoxy-d ₃)phenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	374
192		6-(2-chloro-3,5-di(methoxy-d ₃)phenyl)-N-(cyclopropylmethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	423
193		N-(cyclopropylmethyl)-6-(2-fluoro-3,5-di(methoxy-d ₃)phenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	407

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
194		6-(2-chloro-3-methoxyphenyl)-2-(methylthio)-N-((tetrahydrofuran-2-yl)methyl)pyrido[3,4-d]pyrimidine-8-amine	417
195		6-(2-chloro-3-methoxyphenyl)-N-((1-methylpyrrolidin-2-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	430
196		6-(2-chloro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-2-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	460
197		6-(2-fluoro-3-methoxyphenyl)-N-((1-methylpyrrolidin-2-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	414
198		6-(2-chloro-3-methoxyphenyl)-N-(2-(4-methylpiperazin-1-yl)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	459
199		6-(2-chloro-3,5-dimethoxyphenyl)-N-(2-(4-methylpiperazin-1-yl)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	489
200		6-(2-chloro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidine-8-amine	447







Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		imidine-8-amine	
201		6-(3,5-dimethoxyphenyl)-N-(2-(4-methylpiperazin-1-yl)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	455
202		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(2-methoxyethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	423
203		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(tetrahydro-2H-pyran-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	449
204		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(1-methylpiperidin-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	462
205		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(2-(4-methylpiperazin-1-yl)ethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	491
206		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-2-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	462

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
207		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(1-methylpyrrolidin-3-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	448
208		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-((1-methylpyrrolidin-3-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	462
209		6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	459
210		6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)-N-((tetrahydrofuran-2-yl)methyl)pyrido[3,4-d]pyrimidine-8-amine	449
211		6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)-N-(tetrahydrofuran-3-yl)pyrido[3,4-d]pyrimidine-8-amine	435
212		7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-isopropoxy-2,6-naphthyridine	427
213		7-chloro-1-(cyclopropylmethoxy)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridine	439

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
214		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-isopropoxy-2-(methylthio)pyrido[3,4-d]pyrimidine	440
215		8-(cyclopropylmethoxy)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	452
216		(R)-N-(1-cyclopropylethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	465
217		(S)-N-(1-cyclopropylethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	465
218		(S)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-dimethylbut-2-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	481
219		(R)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-dimethylbut-2-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	481
220		(R)-N-(1-cyclopropylethyl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	433

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
221		(S)-N-(1-cyclopropylethyl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	433
222		(S)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(3,3-dimethylbut-2-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	449
223		(R)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-N-(3,3-dimethylbut-2-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	449
224		N-cyclopropyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	437
225		N-cyclopropyl-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine	405
226		8-(azetidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	437
227		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	467

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
228		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)-2-(methylsulfanyl)pyrido[3,4-d]pyrimidine	435
229		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-N,N-dimethylazetidin-3-amine	480
230		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(3-(trifluoromethyl)azetidin-1-yl)pyrido[3,4-d]pyrimidine	505
231		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-dimethylazetidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	465
232		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-difluoroazetidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	473
233		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-3-methylazetidin-3-ol	467
234		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylazetidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	481








Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
235		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylazetidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	449
236		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-3-methylazetidin-3-carbonitrile	476
237		6-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-2-oxa-6-azaspiro[3.3]heptane	479
238		6-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-2-oxa-6-azaspiro[3.3]heptane	447
239		6-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-1-oxa-6-azaspiro[3.3]heptane	479
240		6-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-1-oxa-6-azaspiro[3.3]heptane	447
241		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(2-azaspiro[3.3]heptan-2-yl)pyrido[3,4-d]pyrimidine	477









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		piro[3.3]hept-2-ylpyrido[3,4-d]pyrimidine	
242		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(2-azaspiro[3.4]oct-2-yl)pyrido[3,4-d]pyrimidine	491
243		2-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-6-oxa-2-azaspiro[3.4]octane	493
244		2-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-7-oxa-2-azaspiro[3.5]nonane	507
245		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(pyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	451
246		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxypyrrolidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	481
247		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxypyrrolidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	449

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
248		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)pyrrolidine-3-carbonitrile	476
249		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-difluoropyrrolidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	487
250		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylpyrrolidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	495
251		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylpyrrolidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	463
252		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-3-methylpyrrolidine-3-carbonitrile	490
253		8-(3-azabicyclo[3.1.0]hex-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	463
254		7-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-2-oxa-7-azaspiro[3.4]nonane	507

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
255		4-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)morpholine	467
256		4-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)morpholine	435
257		4-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfanyl)pyrido[3,4-d]pyrimidin-8-yl)-2-methylmorpholine	481
258		4-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-2-methylmorpholine	449
259		4-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-2,6-dimethylmorpholine	495
260		4-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-2,6-dimethylmorpholine	463
261		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)piperidin-4-ol	481









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
262		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-methoxypiperidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	495
263		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(4-methoxypiperidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	463
264		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxypiperidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	495
265		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-4-methylpiperidin-4-ol	495
266		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-methoxy-4-methylpiperidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	509
267		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(4-methoxy-4-methylpiperidin-1-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	477
268		1-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-4-methylpiperidine-4-carbonitrile	504









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
269		N-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)cyclopropyl carboxamide	465
270		N-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)tetrahydrofuran-2-carboxamide	495
271		N-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)sulfonamide	475
272		6-chloro-N-(cyclopropylmethyl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	439
273		6-chloro-N-(cyclopropylmethyl)-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	407
274		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-methoxyethyl)pyrido[3,4-d]pyrimidine-4-amine	443
275		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(tetrahydrofuran-3-yl)pyrido[3,4-d]pyrimidine-4-amine	455
276		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((tetrahydrofuran-3-yl)methyl)pyrido[3,4-d]pyrimidine-4-amine	469

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(2-methoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	
277		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidine-4-amine	469
278		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-(3,3-difluorocyclopentyl)pyrido[3,4-d]pyrimidine-4-amine	489
279		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-neopentylpyrido[3,4-d]pyrimidine-4-amine	455
280		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)pyrido[3,4-d]pyrimidine-4-amine	479
281		(S)-6-chloro-N-(1-cyclopropylethyl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	453
282		(S)-6-chloro-N-(1-cyclopropylethyl)-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	421
283		6-chloro-N-cyclopropyl-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	425









Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
284		6-chloro-N-cyclopropyl-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-4-amine	393
285		4-(azetidin-1-yl)-6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine	425
286		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidine	455
287		6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidine	423
288		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-N,N-dimethylazetidin-3-amine	468
289		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-(trifluoromethyl)azetidin-1-yl)pyrido[3,4-d]pyrimidine	493
290		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-dimethylazetidin-1-yl)pyrido[3,4-d]pyrimidine	453
291		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidine	461

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		midine	
292		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-3-methylazetidin-3-ol	455
293		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidine	469
294		6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidine	437
295		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-3-methylazetidin-3-carbonitrile	464
296		6-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2-oxa-6-azaspiro[3.3]heptane	467
297		6-(6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2-oxa-6-azaspiro[3.3]heptane	435
298		6-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2-oxa-6-azaspiro[3.3]heptane	467

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		pyrimidin-4-yl)-1-oxa-6-azaspiro[3.3]heptane	
299		6-(6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-1-oxa-6-azaspiro[3.3]heptane	435
300		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-azaspiro[3.3]hept-2-yl)pyrido[3,4-d]pyrimidine	465
301		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-azaspiro[3.4]oct-2-yl)pyrido[3,4-d]pyrimidine	479
302		2-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-6-oxa-2-azaspiro[3.4]octane	481
303		2-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-7-oxa-2-azaspiro[3.5]nonane	495
304		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(pyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	439
305		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	469

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		midine	
306		6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	437
307		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)pyrrolidine-3-carbonitrile	464
308		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-difluoropyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	475
309		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	483
310		6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidine	451
311		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-3-methylpyrrolidine-3-carbonitrile	478
312		4-(3-azabicyclo[3.1.0]hex-3-yl)-6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine	451


Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
313		7-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2-oxa-7-azaspiro[4.4]nonane	495
314		4-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)morpholine	455
315		4-(6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)morpholine	423
316		4-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2-methylmorpholine	469
317		4-(6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2-methylmorpholine	437
318		4-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2,6-dimethylmorpholine	483
319		4-(6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-2,6-dimethylmorpholine	451
320		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)piperidin-4-ol	469

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
321		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidine	483
322		6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidine	451
323		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidine	483
324		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-4-methylpiperidin-4-ol	483
325		6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidine	497
326		6-chloro-2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidine	465
327		1-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)-4-methylpiperidine-4-carbonitrile	492
328		N-(6-chloro-2-(2,6-dichloro-3,5-	453

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)cyclopropylcarboxamide	
329		N-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)tetrahydrofuran-2-carboxamide	483
330		N-(6-chloro-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)sulfonamide	463


Intermediate 331: Preparation of 8-cyclopropyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine

[0229]



Step 1: preparation of 8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine

[0230]




[0231] 8-chloro-6-(3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (500 mg, 1.43

mmol) was added to dichloromethane (30 mL), then the mixture was cooled to -30 °C. SO₂Cl₂ (0.35 mL, 4.31 mmol) was dissolved in dichloromethane (30 mL), and slowly added dropwise to the above reaction liquid with stirring. After the addition, the mixture was stirred for another 0.5 h. The reaction was completed, then quenched with a saturated aqueous sodium bicarbonate solution (50 mL). The mixture was extracted, washed with water and then saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography to obtain 8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (620 mg, 88%). MS m/z (ESI): 416.2 [M+H]⁺.

Step 2: preparation of 8-cyclopropyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine









[0232]










[0233] 8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (83 mg, 0.20 mmol), cyclopropyl boronic acid (26 mg, 0.30 mmol), Pd(OAc)₂ (5 mg, 0.02 mmol), PCy₃ (11 mg, 0.04 mmol) and K₃PO₄ (127 mg, 0.60 mmol) were added to the mixture of toluene and water (6:1, 5 mL), the mixture was heated to 100 °C and stirred over night. The reaction was completed, then the mixture was cooled to room temperature, diluted with EtOAc, washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by column chromatography to obtain 8-cyclopropyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (60 mg, 71%). MS m/z (ESI): 422.2 [M+H]⁺.

[0234] Intermediates 332-364 were prepared referring to the synthesis method of 331.

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
332		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methyl-2-(methylthio)pyrido[3,4-d]pyrimidine	396
333		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-ethyl-2-(methylthio)pyrido[3,4-d]pyrimidine	410

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
334		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-isopropyl-2-(methylthio)pyrido[3,4-d]pyrimidine	424
335		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-neopentylpyrido[3,4-d]pyrimidine	452
336		8-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	436
337		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-((tetrahydrofuran-3-yl)methyl)pyrido[3,4-d]pyrimidine	466
338		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-((tetrahydrofuran-2-yl)methyl)pyrido[3,4-d]pyrimidine	466
339		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidine	466
340		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-methylpyrrolidin-3-yl)methyl-2-(methylthio)pyrido[3,4-d]pyrimidine	479

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		(meth ylthio)pyrido[3,4-d]pyrimidine	
341		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpyrrolidin-2-yl)methyl)-2-(meth ylthio)pyrido[3,4-d]pyrimidine	479
342		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-methylpiperidin-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	479
343		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-methoxyethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	440
344		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methoxymethyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	426
345		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)methyl)cyclopropylamine	451
346		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)methyl)sulfonamide	489
347		N-((6-(2,6-dichloro-3,5-	517


Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)methyl)propane-2-sulfonamide	
348		N-(2-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)ethyl)propane-2-amine	467
349		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(2,2,2-trifluoroethyl)pyrido[3,4-d]pyrimidine	464
350		8-benzyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	472
351		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-phenylpyrido[3,4-d]pyrimidine	458
352		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(pyridin-4-yl)pyrido[3,4-d]pyrimidine	459
353		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(pyridin-3-yl)pyrido[3,4-d]pyrimidine	459
354		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-imidazol-2-yl)pyrido[3,4-d]pyrimidine	462

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		pyrazol-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	
355		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-pyrazol-4-yl)methyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	476
356		8-(1-cyclopropyl-1H-pyrazol-4-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	488
357		8-(1-(cyclopropylmethyl)-1H-pyrazol-4-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine	502
358		6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	506
359		2-(4-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-1H-pyrazol-1-yl)ethan-1-ol	492
360		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-(1-(tetrahydrofuran-3-yl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidine	518

Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d]pyrimidine	
361		6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)-8-phenethylpyrido[3,4-d]pyrimidine	486
362		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-pyrazol-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	430
363		6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)-2-(methylthio)pyrido[3,4-d]pyrimidine	474
364		2-(4-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidin-8-yl)-1H-pyrazol-1-yl)ethan-1-ol	460


Intermediate 365: Preparation of (3S,4S)-4-azidotetrahydro-2H-pyran-3-amine hydrochloride

[0235]



Step 1: preparation of tert-butyl ((3S,4S)-4-hydroxyltetrahydro-2H-pyran-3-yl)carbamate

[0236]




[0237] (3R,4R)-3-aminotetrahydro-2H-pyran-4-ol (5.00 g, 42.7 mmol) was dissolved in MeOH (100 mL), and then Et₃N (4.75 g, 47.0 mmol) and BoC₂O (10.25 g, 47.0 mmol) were successively added, the mixture was stirred at room temperature overnight. The reaction was completed monitored by LCMS. The mixture was concentrated to remove methanol, and the crude product was washed with petroleum ether (100 mL), and then filtrated to obtain tert-butyl ((3S,4S)-4-hydroxyltetrahydro-2H-pyran-3-yl)carbamate (8.50 g, yield: 92%).

[0238] ¹H NMR (400MHz, CDCl₃): δ ppm 4.68(m, 1H), 4.03(m, 1H), 3.93(m, 1H), 3.63(m, 1H), 3.45(m, 2H), 3.19(m, 1H), 2.81(m, 1H), 2.04(m, 1H), 1.68(m, 1H), 1.61(s, 9H).

Step 2: preparation of (3R,4R)-3-((tert-butoxycarbonyl)amino)tetrahydro-2H-pyran-4-yl methanesulfonate

[0239]





[0240] Tert-butyl ((3S,4S)-4-hydroxyltetrahydro-2H-pyran-3-yl)carbamate (8.5 g, 39.2 mmol) was dissolved in CH₂Cl₂ (100 mL), the mixture was cooled under ice water bath, and then Et₃N (4.30 g, 43.0 mmol) and MsCl (4.98 g, 43.0 mmol) were successively added, the mixture was stirred at 0 °C for 6 h. The reaction was completed, the mixture was washed with water (20 mL), then the organic phase was dried over anhydrous sodium sulfate, filtrated, and concentrated to obtain (3R,4R)-3-((tert-butoxycarbonyl)amino)tetrahydro-2H-pyran-4-yl methanesulfonate (10.2 g, yield: 88%).

[0241] ¹H NMR (400MHz, CDCl₃): δ ppm 4.90 (m, 1H), 4.76(m, 1H), 4.03(m, 1H), 3.98(m, 1H), 3.67(m, 2H), 3.47(m, 1H), 3.14(s, 3H), 2.18(m, 1H), 1.92(m, 1H), 1.44 (s, 9H).

Step 3: preparation of tert-butyl ((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)carbamate

[0242]






[0243] (3R,4R)-3-((tert-butoxycarbonyl)amino)tetrahydro-2H-pyran-4-ylmethanesulfonate (10.2 g, 34.6 mmol) was dissolved in dried DMF (100 mL), NaOAc (9.38 g, 69.0 mmol) and NaN₃ (4.49 g, 69.0 mmol) were successively added, the mixture was heated at 90 °C overnight. The reaction was completed, then the reaction solution was poured into water (200 mL) and extracted twice with EtOAc (50 mL), the organic phases were combined, washed twice with aqueous LiCl solution (50 mL), dried over anhydrous sodium sulfate, filtrated and concentrated to obtain compound tert-butyl ((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)carbamate (6.8 g, yield: 81%).

[0244] ¹H NMR (400MHz, CDCl₃): δ ppm 4.85(m, 1H), 3.89(m, 2H), 3.77(m, 1H), 3.63(m, 2H), 3.51(m, 1H), 1.93(m,2H), 1.45(s, 9H).

Step 4: preparation of (3S,4S)-4-azidotetrahydro-2H-pyran-3-amine hydrochloride

[0245]




[0246] Tert-butyl ((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)carbamate (6.8 g, 28.0 mmol) was added to the mixture of HCl and EtOAc (100 mL), the mixture was stirred at room temperature for 8 h. The reaction was completed monitored by LCMS. The mixture was concentrated to obtain compound (3S,4S)-4-azidotetrahydro-2H-pyran-3-amine hydrochloride (4.0 g, yield: 80%).

[0247] ¹H NMR (400MHz, DMSO-d₆): δ ppm 8.47(m, 3H), 4.36(m, 1H), 3.80(m, 2H), 3.63(m, 3H), 2.0(m, 2H).

Intermediate 366: Preparation of (3S,4R)-4-azidotetrahydrofuran-3-amine hydrochloride


[0248]



[0249] Intermediate 366 was prepared referring to the synthesis method of Intermediate 365 from (3S,4R)-4-aminotetrahydrofuran-3-ol.


Intermediate 367: Preparation of tert-butyl ((3R,4S)-4-aminotetrahydrofuran-3-yl)carbamate

[0250]



Step 1: preparation of tert-butyl ((3S,4R)-4-hydroxyltetrahydrofuran-3-yl)carbamate

[0251]



[0252] (3R,4S)-4-aminotetrahydrofuran-3-ol (5.00 g, 48.487 mmol) was dissolved in methanol (100 mL), then triethylamine (12.24 g, 121.2 mmol) was added, the mixture was cooled under ice water bath, then Boc_2O (11.64 g, 53.34 mmol) was added. After the addition, the mixture was stirred at room temperature for 24 h. The reaction was completed, the organic phase was concentrated to remove the majority solvent. Water (100 mL) was added, and the mixture was stirred to precipitate a large amount of white solids. The mixture was filtrated and washed with water to obtain compound tert-butyl ((3S,4R)-4-hydroxyltetrahydrofuran-3-yl)carbamate (8.00 g, yield: 81%).

[0253] ^1H NMR (400 MHz, Chloroform- d) δ 4.72 (s, 1H), 4.29 (m, 1H), 4.09 (m, 1H), 4.08 - 4.02 (m, 1H), 3.95 (s, 1H), 3.69 (m, 1H), 3.61 (m, 1H), 1.45 (s, 9H).

Step 2: preparation of tert-butyl ((3R,4S)-4-(1,3-dicarbonylisoindolin-2-yl)tetrahydrofuran-3-yl)carbamate

[0254]



[0255] Tert-butyl ((3S,4R)-4-hydroxyltetrahydrofuran-3-yl)carbamate (200 mg, 0.984 mmol), triphenylphosphine (620 mg, 2.364 mmol) and isoindoline-1,3-dione (173 mg, 1.181 mmol) were dissolved in tetrahydrofuran (10 mL), the mixture was cooled under ice water bath, then diisopropyl azodicarboxylate (717 mg, 3.546 mmol) was slowly added dropwise. After addition, the mixture was stirred under ice water bath for 1 h. The reaction liquid was directly concentrated and separated by column chromatography (Eluent: PE/EA 15 - 30%) to obtain a crude product compound tert-butyl ((3R,4S)-4-(1,3-dicarbonylisoindolin-2-yl)tetrahydrofuran-3-yl)carbamate (950 mg).

[0256] ^1H NMR (400 MHz, Chloroform- d) δ 7.87 (m, 2H), 7.75 (m, 2H), 4.84 (d, J = 9.5 Hz, 1H), 4.51 (t, J = 8.0 Hz, 1H), 4.37 (t, J = 8.3 Hz, 1H), 4.21 - 4.15 (m, 1H), 4.09 (t, J = 8.8 Hz, 1H), 3.85 (m, J = 9.2, 6.4 Hz, 1H), 1.11 (s, 9H).

[0257] MS m/z (ESI): 559.5 $[\text{M}+\text{H}]^+$.

Step 3: preparation of tert-butyl ((3R,4S)-4-aminotetrahydrofuran-3-yl)carbamate









[0258]



[0259] Tert-butyl ((3R,4S)-4-(1,3-dicarbonylisoindolin-2-yl)tetrahydrofuran-3-yl)carbamate (950 mg, 21%, 0.600 mmol) was dissolved in ethanol (6 mL), then hydrazine hydrate (45 mg, 0.900 mmol) was added, the mixture was heated to 75 °C and stirred for 2 h. The reaction was completed, then the mixture was concentrated and separated by column chromatography (Eluent: DCM/MeOH 0 - 5%) to obtain compound tert-butyl ((3R,4S)-4-aminotetrahydrofuran-3-yl)carbamate (120 mg, two-step yield: 60%).

[0260] ^1H NMR (400 MHz, Chloroform- d) δ 5.24 (s, 1H), 4.11 (s, 1H), 4.05 (m, 1H), 4.00 (m, 1H), 3.63 - 3.52 (m, 2H), 3.49 (m, 1H), 1.46 (s, 9H).

[0261] Intermediates 368-375 were prepared referring to the synthesis method of Intermediate 367.


Intermediate No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
368		tert-butyl ((3S,4S)-3-amino-4-((2,2,2-trichloroethoxy)carbonyl)amino)pyrrolidine-1-carboxylate	377
369		2,2,2-trichloroethyl ((3S,4R)-4-amino-1-methylpyrrolidin-3-yl)carbamate	291
370		2,2,2-trichloroethyl ((3S,4R)-1-acetyl-4-aminopyrrolidin-3-yl)carbamate	319
371		2,2,2-trichloroethyl ((3S,4R)-4-amino-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)carbamate	348
372		2,2,2-trichloroethyl ((3S,4R)-4-amino-1-(methylaminoformyl)pyrrolidin-3-yl)carbamate	334
373		2,2,2-trichloroethyl ((3S,4R)-4-amino-1-(oxetan-3-yl)pyrrolidin-3-yl)carbamate	333
374		2,2,2-trichloroethyl ((3S,4R)-4-amino-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)carbamate	343
375		2,2,2-trichloroethyl ((3S,4R)-4-amino-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)carbamate	357

Intermediate No.	Compound structure	Compound name	MS: m/z $[M+1]^+$
			

Preparation of specific examples


Example 1 Preparation of N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide

[0262]



Step 1: preparation of 3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-7-((2-methyl-6-nitrophenyl)amino)-2,6-naphthyridin-1(2H)-one


[0263]



[0264] 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one (150 mg, 0.375 mmol) was dissolved in 1,4-dioxane (20 mL), then 2-methyl-6-nitroaniline (114 mg, 0.751 mmol), cesium carbonate (367 mg, 1.126 mmol), brett-phos (161 mg, 0.3 mmol) and $\text{Pd}_2(\text{dba})_3$ (172 mg, 0.188 mmol) were added. The gas was exchanged with N_2 , and the mixture was heated to 120 °C for 2 h. The reaction was completed, and the mixture was concentrated and separated by silica gel column chromatography (Eluent: PE/EtOAc = 2 : 1) to obtain compound 3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-7-((2-methyl-6-nitrophenyl)amino)-2,6-naphthyridin-1(2H)-one (40 mg, yield: 21%). MS m/z (ESI): 516 $[M+H]^+$.

Step 2: preparation of 7-((2-amino-6-methylphenyl)amino)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one

[0265]




[0266] 3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-7-((2-methyl-6-nitrophenyl)amino)-2,6-naphthyridin-1(2H)-one (40 mg, 0.078 mmol) was dissolved in 10 mL of methanol, then 10 mg of Pd-C was added. The mixture was stirred at room temperature under H₂ for 30 min. The reaction was completed, and the mixture was concentrated and separated by silica gel column chromatography (Eluent: CH₂Cl₂/MeOH = 10:1) to obtain compound 7-((2-amino-6-methylphenyl)amino)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one (9.5 mg, yield:25%).

[0267] MS m/z (ESI): 486.4 [M+H]⁺.

Step 3: preparation of N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide

[0268]



[0269] 7-((2-amino-6-methylphenyl)amino)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-methyl-2,6-naphthyridin-1(2H)-one (9.5 mg, 0.02 mmol) was dissolved in the mixture of THF and H₂O (4 mL/1 mL), then sodium bicarbonate (9 mg, 0.104 mmol) was added, the mixture was cooled with ice water bath, and the solution of acryloyl chloride in THF (2 mg/2 mL, 0.02 mmol) was added dropwise. The mixture was stirred at 0 °C for 10 min. After the reaction was completed, the mixture was extracted for three times with dichloromethane. The organic phases were combined, washed with a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, filtrated and concentrated. The crude product was separated by








silica gel column chromatography, to obtain compound N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide (1 mg, yield: 9%).

[0270] ^1H NMR (400 MHz, CDCl_3) δ 8.60 (s, 1H), 8.30 (d, J = 8.1 Hz, 1H), 7.31 (s, 1H), 7.10 - 7.05 (m, 2H), 6.68 (s, 1H), 6.52 - 6.48 (m, 1H), 6.44 (s, 1H), 6.39 (s, 1H), 6.33 (d, J = 1.4 Hz, 1H), 6.19 (d, J = 10.2 Hz, 1H), 5.68 (dd, J = 10.2, 1.4 Hz, 1H), 3.99 (s, 6H), 3.25 (s, 3H), 2.21 (s, 3H).

[0271] MS m/z (ESI): 541.4 $[\text{M}+\text{H}]^+$.

[0272] Examples 2-75 were prepared referring to the synthesis method of Example 1.








Example No.	Compound structure	Compound name	MS: m/z $[\text{M}+1]^+$
2		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	525
3		N-(2-((6-(cyclopropylmethyl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	579
4		N-(2-((6-(2-(tert-butylamino)-2-oxoethyl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	638
5		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-oxo-3,4-dihydropyrido[3,4-d]pyrimidin-6-yl)amino)-3-methylphenyl)acrylamide	526
6		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-3-methyl-4-oxo-3,4-dihydropyrido[3,4-d]pyrimidin-6-yl)amino)-3-methylphenyl)acrylamide	540








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		3,4-dihydropyrido[3,4-d]pyrimidin-6-yl)amino)-3-methylphenyl)acrylamide	
7		N-(2-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-oxo-1H-pyrano[4,3-c]pyridin-7-yl)amino)-3-methylphenyl)acrylamide	526
8		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-7-methyl-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	540
9		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-methoxy-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	539
10		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-methoxypyrido[3,4-d]pyrimidin-6-yl)amino)-3-methylphenyl)acrylamide	540
11		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-3-(2-methoxyethyl)-4-oxo-3,4-dihydropyrido[3,4-d]pyrimidin-6-yl)amino)-3-methylphenyl)acrylamide	584
13		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	509
14		N-(2-((6-(2,6-dichloro-3,5-	510








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		dimethoxyphenyl)pyrido[3, 4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	
15		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3, 4-d]pyrimidin-2-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide	608
16		N-(2-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-(4-ethylpiperazin-1-yl)-3-methylphenyl)acrylamide	589
17		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-(4-ethylpiperazin-1-yl)-3-methylphenyl)acrylamide	621
18		N-(2-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide	575
19		N-(2-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-((4-methylpiperazin-1-yl)methyl)phenyl)acrylamide	575
20		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-(4-ethylpiperazin-1-yl)phenyl)acrylamide	607

Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
21		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-(2-(dimethylamino)ethoxy)phenyl)acrylamide	582
22		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-morpholinophenyl)acrylamide	580
23		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-((4-methylpiperazin-1-yl)methyl)phenyl)acrylamide	607
24		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-5-(1-ethylpiperidin-4-yl)phenyl)acrylamide	606
25		N-(2-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	578
26		N-(2-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-bis(methoxy-d ₃)phenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	584
27		N-(2-((5-((cyclopropylmethyl)amino)-7-(2,6-difluoro-3,5-bis(methoxy-d ₃)phenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	552

Example No.	Compound structure	Compound name	MS: m/z $[M+1]^+$
		bis(methoxy- d_3)phenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	
28		N-(2-((5-((cyclopropylmethyl)amino)-7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	546
29		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-hydroxyethyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	568
30		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-isopropylthioethyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	626
31		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-ethylsulfonyl)ethyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	644
32		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(dimethylamino)ethyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	595
33		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((3-(dimethylamino)propyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	609

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
34		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(methylamino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	538
35		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(ethylamino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	552
36		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2,2,2-trifluoroethyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	606
37		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(dimethylamino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	552
38		N-(2-((5-((cyclopropylmethyl)(methyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	592
39		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(methylsulfonamido)ethyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	645
40		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydro-2H-pyran-2-yl)methyl)(methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	608




Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	
41		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	608
42		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((oxetan-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	594
43		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydro-2H-pyran-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	622
44		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	580
45		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydro-2H-pyran-4-yl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	608
46		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	594

Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
			
47		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methylpyrrolidin-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	621
48		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methylpyrrolidin-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	621
49		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((1-methylpyrrolidin-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	607
50		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methylazetidin-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	607
51		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((1-methylazetidin-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	593
52		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((1-methylpiperidin-4-yl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	621
53		N-(2-((7-(2,6-dichloro-3,5-	635

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		dimethoxyphenyl)-5-(((1-methylpiperidin-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	
54		N-(2-(((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((3,3-difluorocyclobutyl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	614
55		N-(2-(((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((3,3-difluorocyclopentyl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	628
56		N-(2-(((5-((cyclopentylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	606
57		N-(2-(((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(phenethylamino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	628
58		N-(2-(((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	618
59		N-(2-(((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-(2-hydroxyethyl)-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	648


Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	
60		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	662
61		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-ethoxy-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	553
62		N-(2-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-isopropoxy-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	567
63		N-(2-((5-(cyclopropylmethoxy)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-3-methylphenyl)acrylamide	579
64		N-(3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide	500
65		N-(3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	544

Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
66		N-(3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methoxy-pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide	530
67		N-(3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide	499
68		N-(3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-methoxyethyl)amino)-2,6-naphthyridin-3-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide	572
69		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)thieno[2,3-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	515
70		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)thieno[3,2-c]pyridin-6-yl)amino)-3-methylphenyl)acrylamide	515
71		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)thiazolo[5,4-d]pyrimidin-5-yl)amino)-3-methylphenyl)acrylamide	516
72		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)thiazolo[4,5-c]pyridin-6-yl)amino)-3-methylphenyl)acrylamide	515
73		N-(2-((2-(2,6-dichloro-3,5-	526

Example No.	Compound structure	Compound name	MS: m/z $[M+1]^+$
		dimethoxyphenyl)-4-oxo-4 H-pyrano[2,3-c]pyridin-6-yl)amino)-3-methylphenyl)acrylamide	
74		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-8 H-pyrano[3,2-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	527
75		N-(2-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-3-methyl-4-oxo-3,4-dihydropyrido[3,4-d]pyrimidin-6-yl)amino)-3-methyl-5-(4-methylpiperazin-1-yl)phenyl)acrylamide	638


Example 76 Preparation of N-(3-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide

[0273]



Step 1: preparation of N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine

[0274]




[0275] N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine (210 mg, 0.465 mmol) was dissolved in dichloromethane (6 mL), then *m*-chloroperoxybenzoic acid (200 mg, 1.163 mmol) was added, and the mixture was stirred at room temperature for 18 h. After the reaction was completed, a saturated sodium sulfite solution (5 mL) was added, and the mixture was stirred for 5 min and extracted with dichloromethane, the organic phase was washed successively with a saturated sodium bicarbonate solution and then saturated brine, dried over anhydrous sodium sulfate, filtered, concentrated and separated by column chromatography (Eluent: PE/EA = 2/1) to obtain compound N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine (186 mg, yield: 82.7%).

[0276] MS m/z (ESI): 483.4 $[M+H]^+$.


Step 2: preparation of N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N²-(1-methyl-4-nitro-1H-pyrazol-3-yl)pyrido[3,4-d]pyrimidine-2,8-diamine

[0277]




[0278] N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine (90 mg, 0.186 mmol) was dissolved in dried DMF (5 mL), and NaH (15 mg, 0.372 mmol) was added at 0 °C, and the mixture was stirred at room temperature for 10 min. Then 1-methyl-4-nitro-1H-pyrazol-3-amine (32 mg, 0.223 mmol) was added, and the mixture was stirred at room temperature for 3 h. The mixture was quenched with a saturated aqueous ammonium chloride, extracted with ethyl acetate and separated by silica gel column chromatography (PE/EA = 1/1) to obtain compound N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N²-(1-methyl-4-nitro-1H-pyrazol-3-yl)pyrido[3,4-d]pyrimidine-2,8-diamine (100 mg, yield: 98%). MS m/z (ESI): 545 $[M+H]^+$.

Step 3: preparation of N²-(4-amino-1-methyl-1H-pyrazol-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine

(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine**[0279]**

[0280] N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N²-(1-methyl-4-nitro-1H-pyrazol-3-yl)pyrido[3,4-d]pyrimidine-2,8-diamine (40 mg, 0.074 mmol) was dissolved in methanol (10 mL), and then triethylamine (3 drops) and Pd/C (20 mg, 10% content) were successively added, and the mixture was stirred at room temperature under H₂ for 45 min. The reaction was completed and the mixture was filtrated and separated by silica gel column chromatography (DCM/MeOH) to obtain compound N²-(4-amino-1-methyl-1H-pyrazol-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine (30 mg, yield:80%). MS m/z (ESI): 515 [M+H]⁺.

Step 4: preparation of N-(3-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide








[0281]








[0282] N²-(4-amino-1-methyl-1H-pyrazol-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine (30 mg, 0.058 mmol) was dissolved in the mixture of THF and H₂O (8 mL/2 mL), then NaHCO₃ (33 mg, 0.39 mmol) was added, and acetyl chloride (7 mg, 0.078 mmol, dissolved in 1 mL of THF) was added dropwise under ice bath, and the mixture was stirred for 10 minutes. A saturated aqueous sodium bicarbonate solution (20 mL) was added, and the mixture was extracted with ethyl acetate and separated by silica gel column chromatography (PE/EA = 1/1) to obtain compound N-(3-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide (10 mg, yield:22%). MS m/z (ESI): 569 [M+H]⁺.

[0283] ^1H NMR (400 MHz, CDCl_3) δ 9.03 (s, 1H), 8.17 (s, 1H), 6.77 (s, 1H), 6.63 (s, 1H), 6.38 (d, J = 16.6 Hz, 2H), 6.33 - 6.24 (m, 1H), 5.70 (d, J = 10.1 Hz, 1H), 3.95 (s, 6H), 3.87 (s, 3H), 3.53-3.36(m, 2H), 1.19-1.10 (m, 1H), 0.55-0.41(m, 2H), 0.42-0.29 (m, 2H).







[0284] Examples 77-117 were prepared referring to the synthesis method of Example 76.







Example No.	Compound structure	Compound name	MS: m/z $[\text{M}+1]^+$
77		N-(2-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	579
78		N-(2-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-bis(methoxy- d_3)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	585
79		N-(2-((8-((cyclopropylmethyl)amino)-6-(2,6-difluoro-3,5-bis(methoxy- d_3)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	553
80		N-(2-((8-((cyclopropylmethyl)amino)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	547
81		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-hydroxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	569
82		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	583







Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		ide	
83		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(ethylsulfonyl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	645
84		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(dimethylamino)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	596
85		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3-(dimethylamino)propyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	610
86		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	539
87		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	553
88		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	607

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
89		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(dimethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	553
90		N-(2-((8-((cyclopropylmethyl)(methyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	593
91		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(methylsulfonylamido)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	646
92		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	609
93		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	609
94		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((oxetan-3-ylmethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	595


Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
95		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydro-2H-pyran-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	623
96		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	581
97		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	609
98		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	595
99		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	622
100		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	622

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
101		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpyrrolidin-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	608
102		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylazetidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	608
103		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylazetidin-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	594
104		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpiperidin-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	622
105		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpiperidin-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	636
106		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,3-difluorocyclobutyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	615

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
107		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,3-difluorocyclopentyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	629
108		N-(2-((8-((cyclopentylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	607
109		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(phenethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	629
110		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	619
111		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-(2-hydroxyethyl)-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	649
112		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	663


Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
113		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methoxypyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	540
114		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-ethoxypyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	554
115		N-(2-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-isopropoxypyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	568
116		N-(2-((8-(cyclopropylmethoxy)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-3-methylphenyl)acrylamide	
117		N-(3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methyl-1H-pyrazol-4-yl)acrylamide	573

Example 118 Preparation of (+)-N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide



Step 1: preparation of (±)-(3S,4R)-N³-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydrofuran-3,4-diamine


[0286]



[0287] 2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (40.0 mg, 0.108 mmol) and trans-tetrahydrofuran-3,4-diamine dihydrochloride (28.3 mg, 0.162 mmol) were dissolved in acetonitrile (2 mL), then N,N-diisopropylethylamine (70 mg, 0.543 mmol) was added, the mixture was heated to reflux for 16 h. After being cooled, the reaction liquid was diluted with EtOAc (10 mL), washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by PTLC (Eluent: CH₂Cl₂/MeOH = 10:1) to obtain compound (±)-(3S,4R)-N³-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydrofuran-3,4-diamine (30 mg, yield: 64%). MS m/z (ESI): 436.1 [M+H]⁺.

Step 2: preparation of (±)-N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide

[0288]




[0289] (±)-(3S,4R)-N³-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydrofuran-3,4-diamine (30.0 mg, 0.069 mmol) was dissolved in the mixture THF and H₂O (1.2/0.3 mL), NaHCO₃ (23.0 mg, 0.276 mmol) was added, and the mixture was cooled

under ice water bath, and then acryloyl chloride (6.8 mg, 0.076 mmol) was added. After addition, the mixture was stirred at 0 °C for 10 min. The reaction liquid was diluted with EtOAc (5 mL), washed with saturated brine, dried over anhydrous sodium sulfate, filtrated and concentrated. The crude product was separated by PTLC (Eluent: CH₂Cl₂/MeOH= 10:1) to obtain compound N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide (17.4 mg, yield: 52%).

[0290] ¹H NMR (400 MHz, CDCl₃) δ 9.21 (s, 1H), 9.16 (s, 1H), 7.55 (s, 1H), 6.68 (s, 1H), 6.45 (d, J = 7.3 Hz, 1H), 6.27 (dd, J = 17.0, 1.4 Hz, 1H), 6.07 (dd, J = 16.9, 10.3 Hz, 1H), 5.63 (dd, J = 10.2, 1.4 Hz, 1H), 4.94 (dd, J = 11.6, 4.9 Hz, 1H), 4.90 - 4.79 (m, 1H), 4.34 - 4.19 (m, 2H), 3.98 (s, 6H), 3.90 - 3.79 (m, 2H). MS m/z (ESI): 490.1 [M+H]⁺.


Example 119 Preparation of N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide

[0291]



Step 1: preparation of N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2-amine

[0292]




[0293] 2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (100 mg, 0.27 mmol) was dissolved in NMP (3 mL), and Na₂CO₃ (143 mg, 1.349 mmol) and (3S,4S)-4-azidotetrahydro-2H-pyran-3-amine hydrochloride (72 mg, 0.405 mmol) were added, the mixture was heated to 120 °C for 2 h. The reaction was completed, and the mixture was cooled to room temperature, added with water, and extracted for three times with ethyl acetate. The organic phases were combined, dried over anhydrous sodium sulfate, filtrated, concentrated

and separated by column chromatography (Eluent: petroleum ether/ethyl acetate 2:1) to obtain compound N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2-amine (38 mg, yield: 29%). MS m/z (ESI): 478.4 [M+ H]⁺.

Step 2: preparation of (3S,4S)-N³-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydro-2H-pyran-3,4-diamine


[0294]



[0295] N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2-amine (38 mg, 0.08 mmol) was dissolved in MeOH (8 mL), then Pd/C (10 mg) was added, and the mixture was stirred under H₂ at room temperature for 30 min. The reaction was completed, and the mixture was concentrated and separated by column chromatography (Eluent: dichloromethane/methanol 10: 1) to obtain compound (3S,4S)-N³-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydro-2H-pyran-3,4-diamine (12 mg, yield: 33%). MS m/z (ESI): 451.2 [M+H]⁺.

Step 3: preparation of N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide

[0296]



[0297] (3S,4S)-N³-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydro-2H-pyran-3,4-diamine (12 mg, 0.027 mmol) was dissolved in the mixture of THF and H₂O (4 mL/1 mL), then NaHCO₃ (12 mg, 0.141 mmol) was added, the mixture was cooled under ice water bath, and a solution of acryloyl chloride (3 mg, 0.027 mmol) in THF (2 mL) was added dropwise, and the mixture was stirred at low temperature for 10 min. After the reaction was completed, the mixture was extracted three times with dichloromethane. The organic phases








were combined, washed with saturated brine, dried over anhydrous sodium sulfate, concentrated, and separated by column chromatography (Eluent: dichloromethane/petroleum ether (10:1)) to obtain compound N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide (6.3 mg, yield: 47%).

[0298] ^1H NMR (400 MHz, CDCl_3) δ 9.17 (s, 2H), 7.56 (s, 1H), 6.68 (s, 1H), 6.25 (dd, J = 16.9, 1.4 Hz, 1H), 6.01 (dd, J = 17.0, 10.3 Hz, 1H), 5.60 (dd, J = 10.3, 1.4 Hz, 1H), 4.55 (s, 1H), 4.35 (s, 1H), 4.11 - 4.00 (m, 2H), 3.98 (s, 6H), 3.78 (d, J = 12.1 Hz, 1H), 3.64 (dd, J = 13.4, 10.9 Hz, 1H), 2.24-2.20 (m, 2H), 2.02-1.99 (m, 2H). MS m/z (ESI): 505.4 $[\text{M}+\text{H}]^+$.








[0299] Examples 120-420 and 752 were prepared referring to the synthesis method of Example 118 or 119.

Example No.	Compound structure	Compound name	MS: m/z $[\text{M}+1]^+$
120		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	458
121		N-((3R,4S)-4-((6-(2-chloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	456
122		N-((3R,4S)-4-((6-(2-chloro-3,5-dimethoxy-6-methylphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	470
123		N-((3R,4S)-4-((6-(2-chloro-6-fluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	474
124		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	489

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
125		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-methoxyethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	562
126		N-((3R,4S)-4-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	558
127		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	505
128		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	519
129		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-methoxy-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	519
130		N-((3R,4S)-4-((6-(2-chloro-6-cyclopropyl-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	496
13		N-((3R,4S)-4-((6-(2-chloro-6-isopropyl-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	498

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
132		N-((3R,4S)-4-((7-(2-chloro-6-isopropyl -3,5-dimethoxyphenyl)-2,6-naphthyridin-3 -yl)amino)tetrahydrofuran-3 -yl)acry lamide	497
133		N-((3R,4S)-4-((5-((cyclopropylmethyl) amino)-7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	526
134		N-((3R,4S)-4-((5-((cyclopropylmethyl) amino)-7-(2,6-dichloro-3,5-bis(methoxy -d ₃)phenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3 -yl)acrylamide	564
135		N-((3R,4S)-4-((5-((cyclopropylmethyl) amino)-7-(2,6-difluoro-3,5-bis(methoxy -d ₃)phenyl)-2,6-naphthyridin-3-yl)amin o)tetrahydrofuran-3 -yl)acrylamide	532
136		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dim ethoxyphenyl)-5-((2-hydroxyethyl)ami no)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3 -yl)acrylamide	548
137		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dim ethoxyphenyl)-5-((2-(ethylsulfonyl)eth yl)amino)-2, 6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	624
138		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dim ethoxyphenyl)-5-((2-(dimethylamino)et hyl)amino)-2,6-naphthyridin-3-yl)amin o)tetrahydrofuran-3 -yl)acrylamide	575








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
139		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(ethylamino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	532
140		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2,2,2-trifluoroethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	586
141		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(dimethylamino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	532
142		N-((3R,4S)-4-((5-((cyclopropylmethyl)(methyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	572
143		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(methylsulfonamido)ethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	625
144		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
145		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		1-methylamino)-2,6-naphthyridin-3-yl) amino)tetrahydrofuran-3-yl)acrylamide	
146		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((oxetan-3-ylmethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
147		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydro-2H-pyran-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
148		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	560
149		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
150		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydro-2H-pyran-4-yl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
151		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methylpiperidin-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	615








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
152		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((3,3-difluorocyclopentyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	608
153		N-((3R,4S)-4-((5-((cyclopentylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	586
154		N-((3R,4S)-4-((5-(benzylamino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	594
155		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	598
156		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-(2-hydroxyethyl)-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	628
157		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	642

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
	3,5-dim		
158		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(4-methylpiperazin-1-yl)ethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	630
159		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-morpholinoethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	617
160		N-((3R,4S)-4-((7-(2-chloro-3-methoxyphenyl)-5-((2-hydroxyethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	484
161		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-((2-hydroxyethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	516
162		N-((3R,4S)-4-((7-(2-fluoro-3-methoxyphenyl)-5-((2-hydroxyethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	468
163		N-((3R,4S)-4-((7-(2-chloro-3,5-dimethoxyphenyl)-5-((2-(dimethylamino)ethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	541
164		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-((2-hydroxyethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	543

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		(dimethylamino)ethylamino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	
165		N-((3R,4S)-4-((5-((2-(dimethylamino)ethyl)amino)-7-(2-fluoro-3-methoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	495
167		N-((3R,4S)-4-((7-(2-fluoro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	538
168		N-((3R,4S)-4-((7-(2-fluoro-3-methoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	508
169		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-ethoxy-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	533
170		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-isopropoxy-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	547
171		N-((3R,4S)-4-((5-(cyclopropylmethoxy)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	559







Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
172		N-((3 S,4S)-3-((6-(2-chloro-6-fluoro-3,5 - dimethoxyphenyl)pyrido[3,4- d]pyrimi din-2- yl)amino)tetrahydro-2H-pyran -4- yl)acrylamide	488
173		N-((3 S,4S)-3-((6-(6-chloro-7- methoxy-2,3-dihydrobenzo[b] [1,4]dioxin-5-yl)py rido[3,4- d]pyrimidin-2-yl)amino)tetrah ydro-2H-pyran-4-yl)acrylamide	498
174		N-((3S,4S)-3-((6-(2,6-dichloro- 3,5-dim ethoxyphenyl)-8-oxo-7, 8-dihydropyrid o[3,4-d]pyrimidin- 2-yl)amino)tetrahydr o-2H-pyran- 4-yl)acrylamide	520
175		N-((3S,4S)-3-((6-(2,6-dichloro- 3,5-dim ethoxyphenyl)-7-methyl- 8-oxo-7,8-dih ydropyrido[3,4- d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4- yl)acrylamide	534
176		N-((3S,4S)-3-((7- (cyclopropylmethyl)-6-(2,6- dichloro-3,5-dimethoxyphenyl)-8- oxo-7,8-dihydropyrido[3,4- d]pyrimid in-2- yl)amino)tetrahydro-2H-pyran -4- y l)acrylamide	574
177		N-((3S,4S)-3-((7-(2,6-dichloro- 3,5-dim ethoxyphenyl)-2,6- naphthyridin-3-yl)a mino)tetrahydro-2H-pyran-4- yl)acryla mide	503
178		N-((3S,4S)-3-((7-(2,6-dichloro- 3,5-dim ethoxyphenyl)-5-((2- methoxyethyl)ami no)-2,6-	576

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
179		N-((3S,4S)-3-((5-((cyclopropylmethyl) amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	572
180		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-methoxy-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	533
181		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	519
182		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	533
183		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)thieno[2,3-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	509
184		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)thiazolo[5,4-d]pyrimidin-5-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	510
185		N-((3S,4S)-3-((2-(2,6-dichloro-	509

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		3,5-dimethoxyphenyl)thiazolo[4,5-c]pyridin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
186		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-oxo-4H-pyrano[2,3-c]pyridin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	520
187		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-8H-pyrano[3,2-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	521
188		N-((3S,4S)-3-((6-(2-chloro-6-cyclopropyl-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	510
189		N-((3S,4S)-3-((6-(2-fluoro-6-isopropyl-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	496
190		N-((3S,4S)-3-((6-(2-cyclopropyl-6-fluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	493
191		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	472

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
192		N-((3S,4S)-3-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	471
193		N-((3S,4S)-3-((7-(2-chloro-6-cyclopropyl-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	509
194		N-((3S,4S)-3-((7-(2-cyclopropyl-6-fluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	493
195		N-((3S,4S)-3-((7-(2-fluoro-6-isopropyl-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	495
196		N-((3S,4S)-3-((7-(6-chloro-7-methoxy-2,3-dihydrobenzo[b][1,4]dioxin-5-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	496
197		N-((3S,4S)-3-((7-(6-fluoro-7-methoxy-2,3-dihydrobenzo[b][1,4]dioxin-5-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	481
198		N-((3S,4S)-3-((5-((cyclopropylmethyl)amino)-7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	540








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
199		N-((3S,4S)-3-((5-((cyclopropylmethyl) amino)-7-(2,6-dichloro-3,5-bis(methoxy-d ₃)phenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	578
200		N-((3S,4S)-3-((5-((cyclopropylmethyl) amino)-7-(2,6-difluoro-3,5-bis(methoxy-d ₃)phenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	546
201		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-hydroxyethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	562
202		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(dimethylamino)ethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
203		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(methylamino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	532
204		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(ethylamino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	546
205		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2,2,2-trifluoroethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	600








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
206		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(dimethylamino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	546
207		N-((3S,4S)-3-((5-((cyclopropylmethyl)(methyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	586
208		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(methylsulfonamido)ethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	639
209		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	602
210		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-3-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	602
211		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((oxetan-3-ylmethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	588
212		N-((3S,4S)-3-((7-(2,6-dichloro-	616








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
219		N-((3 S,4S)-3-((5-(benzylamino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	608
220		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	612
221		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-(2-hydroxyethyl)-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	642
222		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	656
223		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-(4-methylpiperazin-1-yl)ethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	644
224		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-morpholinoethyl)amino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	631

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
225		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-ethoxy-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	547
226		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-isopropoxy-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	561
227		N-((3S,4S)-3-((5-(cyclopropylmethoxy)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	573
228		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)pyrrolidin-3-yl)acrylamide	489
229		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	488
230		N-((3S,4R)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	455
231		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)amino)pyrrolidin-3-yl)acrylamide	505








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
232		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	504
233		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	518
234		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-8H-pyrano[3,4-d]pyrimidin-2-yl)amino)pyrrolidin-3-yl)acrylamide	506
235		N-((3S,4R)-4-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-oxo-1H-pyrano[4,3-c]pyridin-7-yl)amino)pyrrolidin-3-yl)acrylamide	505
236		N-((3S,4R)-4-((3-(2,6-difluoro-3,5-dimethoxyphenyl)-1-oxo-1H-pyrano[4,3-c]pyridin-7-yl)amino)pyrrolidin-3-yl)acrylamide	472
237		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	502
238		N-((3S,4R)-1-acetyl-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	530
239		N-((3S,4R)-1-acetyl-4-((7-(2,6-	559









Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		dichloro -3,5 - dimethoxyphenyl)-5-(methylamino)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	
240		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	559
241		(3S,4R)-3-acrylamido-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-N-methylpyrrolidine-1-carboxamide	545
242		(3S,4R)-3-acrylamido-4-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-N-methylpyrrolidine-1-carboxamide	614
243		N-((3S,4R)-1-acetyl-4-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	599
244		N-((3S,4R)-4-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	613
245		N-((3S,4R)-4-((5-((cyclopropylmethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	628

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		(dimethylamino)ethylpyrrolidin-3-yl)acrylamide	
246		N-((3S,4R)-4-((5-((cyclopropylmethyl) amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	623
247		N-((3S,4R)-4-((5-((cyclopropylmethyl) amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	637
248		N-((3S,4R)-4-((5-((cyclopropylmethyl) amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	571
249		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	601
250		(3S,4R)-3-acrylamido-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)-N-methylpyrrolidine-1-carboxamide	644
251		N-((3S,4R)-1-acetyl-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	629








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		di-3-yl)acrylamide	
252		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl) amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	643
253		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl) amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	658
254		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl) amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	653
255		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(((tetrahydrofuran-2-yl)methyl)amino)-2,6-naphthyridin-3-yl) amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	667
256		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl) amino)-1-methylpyrrolidin-3-yl)acrylamide	573
257		(3S,4R)-3-acrylamido-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-	616

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		naphthyridin-3-yl)amino)-N-methylpyrrolidine-1-carboxamide	
258		N-((3S,4R)-1-acetyl-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	601
259		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	615
260		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	630
261		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	625
262		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(oxetan-3-ylamino)-2,6-naphthyridin-3-yl)amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	639
263		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-2,6-naphthyridin-3-yl)acrylamide	587








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	
264		(3 S,4R)-3-acrylamido-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-N-methylpyrrolidine-1-carboxamide	630
265		N-((3S,4R)-1-acetyl-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)pyrrolidin-3-yl)acrylamide	615
266		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	629
267		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	644
268		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	639
269		N-((3S,4R)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((tetrahydrofuran-3-yl)amino)-2,6-naphthyridin-3-yl)amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	653

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	
270		N-((3R,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)piperidin-4-yl)acrylamide	503
271		N-((3R,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)piperidin-4-yl)acrylamide	470
272		N-((3R,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)piperidin-4-yl)acrylamide	502
273		N-((3R,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)amino)piperidin-4-yl)acrylamide	519
274		N-((3R,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)amino)piperidin-4-yl)acrylamide	518
275		N-((3R,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-7-methyl-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)amino)piperidin-4-yl)acrylamide	533
276		N-((3R,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-8H-pyrano[3,4-d]pyrimidin-2-yl)amino)piperidin-4-yl)acrylamide	520








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
277		N-((3R,4S)-3-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-oxo-1H-pyrano[4,3-c]pyridin-7-yl)amino)piperidin-4-yl)acrylamide	519
278		N-((3R,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-oxo-8H-pyrano[3,4-d]pyrimidin-2-yl)amino)piperidin-4-yl)acrylamide	487
279		N-((3R,4S)-4-((5-(((S)-1-cyclopropylethyl)amino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	572
280		N-((3R,4S)-4-((5-(((S)-1-cyclopropylethyl)amino)-7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	540
281		N-((3R,4S)-4-((5-(cyclopropylamino)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	544
282		N-((3R,4S)-4-((5-(cyclopropylamino)-7-(2,6-difluoro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	512
283		N-((3R,4S)-4-((5-(azetidin-1-yl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	544

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
284		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxyazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
285		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(3-methoxyazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	542
286		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-(dimethylamino)azetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
287		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-(trifluoromethyl)azetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	612
288		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3,3-dimethylazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	572
289		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3,3-difluoroazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	580
290		N-((3R,4S)-4-((7-(2,6-dichloro-	574

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		3,5-dimethoxyphenyl)-5-(3-hydroxy-3-methylazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	
291		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxy-3-methylazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
292		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(3-methoxy-3-methylazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	556
293		N-((3R,4S)-4-((5-(3-cyano-3-methylazetidin-1-yl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	583
294		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	586
295		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	554
296		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(1-oxa-6-azaspiro[3.3]heptan-6-yl)-2,6-	586

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	
297		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-diethoxyphenyl)-5-(1-oxa-6-azaspiro[3.3]heptan-6-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	554
298		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-diethoxyphenyl)-5-(2-azaspiro[3.3]heptan-2-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	584
299		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-diethoxyphenyl)-5-(2-azaspiro[3.4]octan-2-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	598
300		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-diethoxyphenyl)-5-(6-oxa-2-azaspiro[3.4]octan-2-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	600
301		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-diethoxyphenyl)-5-(7-oxa-2-azaspiro[3.5]nonan-2-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	614
302		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-diethoxyphenyl)-5-(pyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	558








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
303		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxypyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
304		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(3-methoxypyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	556
305		N-((3R,4S)-4-((5-(3-cyanopyrrolidin-1-yl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	583
306		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3,3-difluoropyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	594
307		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxy-3-methylpyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
308		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(3-methoxy-3-methylpyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	570
309		N-((3R,4S)-4-((5-(3-cyano-3-	597

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		methylpyrrolidin-1-yl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	
310		N-((3R,4S)-4-((5-(3-azabicyclo[3.1.0]hexan-3-yl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	570
311		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(2-oxa-7-azaspiro[4.4]nonan-7-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	614
312		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-morpholino-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
313		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-morpholino-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	542
314		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(2-methylmorpholino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
315		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(2-methylmorpholino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	556

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
316		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(2,6-dimethylmorpholino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
317		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(2,6-dimethylmorpholino)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	570
318		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(4-hydroxypiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
319		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(4-methoxypiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
320		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(4-methoxypiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	570
321		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxypiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
322		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(4-hydroxy-4-methylpiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	602








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)-2,6-naphthyridin-3 -yl)amino)tetrahydrofuran-3-yl)acrylamide	
323		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(4-methoxy-4-methylpiperidin-1-yl)-2,6-naphthyridin-3 -yl)amino)tetrahydrofuran-3-yl)acrylamide	616
324		N-((3R,4S)-4-((7-(2,6-difluoro-3,5-dimethoxyphenyl)-5-(4-methoxy-4-methylpiperidin-1-yl)-2,6-naphthyridin-3 -yl)amino)tetrahydrofuran-3-yl)acrylamide	584
325		N-((3R,4S)-4-((5-(4-cyano-4-methylpiperidin-1-yl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3 -yl)acrylamide	611
326		N-((7-(((3S,4R)-4-acrylamido-2,6-naphthyridin-1-yl)amino)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)cyclopropanecarboxamide	572
327		N-((7-(((3S,4R)-4-acrylamido-2,6-naphthyridin-1-yl)amino)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-1-yl)tetrahydrofuran-2-carboxamide	602
328		N-((3R,4S)-4-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(methylsulfonamido)-2,6-naphthyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	582

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
329		N-((3 S,4S)-3-((5-(cyclopropylamino)-7 -(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2 H-pyran-4-yl)acrylamide	558
330		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxyazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	588
331		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3,3-difluoroazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	594
332		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxy-3-methylazetidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	602
333		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(2-oxa-6-azaspiro[3.3]heptan-6-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	600
334		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(6-oxa-2-azaspiro[3.4]octan-2-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	614
335		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(7-oxa-2-azaspiro[3.5]nonan-2-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	628

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		naphthyridin-3 -yl)amin o)tetrahydro-2H-pyran-4-yl)acrylamide	
336		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxypyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	602
337		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(3-methoxy-3-methylpyrrolidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	616
338		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(2,6-dimethylmorpholino)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	616
339		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(4-hydroxy-4-methylpiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	616
340		N-((3S,4S)-3-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-(4-methoxy-4-methylpiperidin-1-yl)-2,6-naphthyridin-3-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	630
341		N-((3R,4S)-4-((4-((cyclopropylmethyl)amino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	559

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
342		N-((3R,4S)-4-((4-((cyclopropylmethyl) amino)-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	527
343		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	563
344		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
345		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
346		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
347		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((3,3-difluorocyclopentyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	609
348		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(neopentylamino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	575

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d[pyrimidin-6-yl]amino)tetrahydrofuran-3-yl)acrylamide	
349		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	599
350		N-((3R,4S)-4-((4-(((S)-1-cyclopropylethyl)amino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	573
351		N-((3R,4S)-4-((4-(((S)-1-cyclopropylethyl)amino)-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	541
352		N-((3R,4S)-4-((4-(cyclopropylamino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	545
353		N-((3R,4S)-4-((4-(cyclopropylamino)-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	513
354		N-((3R,4S)-4-((4-(azetidin-1-yl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	545








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
355		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
356		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	543
357		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-(dimethylamino)azetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
358		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-(trifluoromethyl)azetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	613
359		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-dimethylazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	573
360		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	581
361		N-((3R,4S)-4-((2-(2,6-dichloro-	575

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	
362		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
363		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
364		N-((3R,4S)-4-((4-(3-cyano-3-methylazetidin-1-yl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	584
365		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
366		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	555
367		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(1-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	587

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-6-yl amino)tetrahydrofuran-3-yl)acrylamide	
368		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(1-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	555
369		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-azaspiro[3.3]heptan-2-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	585
370		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-azaspiro[3.4]octan-2-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	599
371		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(6-oxa-2-azaspiro[3.4]octan-2-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	601
372		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(7-oxa-2-azaspiro[3.5]nonan-2-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	615
373		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(pyrrolidin-1-yl)pyridin-3-yl)amino)tetrahydrofuran-3-yl)acrylamide	559








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d[pyrimidin-6-yl]amino)tetrahydrofuran-3-yl)acrylamide	
374		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
375		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
376		N-((3R,4S)-4-((4-(3-cyanopyrrolidin-1-yl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	584
377		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-difluoropyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	595
378		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
379		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	571

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
380		N-((3R,4S)-4-((4-(3-cyano-3-methylpyrrolidin-1-yl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	598
381		N-((3R,4S)-4-((4-(3-azabicyclo[3.1.0]hexan-3-yl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
382		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-oxa-7-azaspiro[4.4]nonan-7-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	615
383		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-morpholinopyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
384		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-morpholinopyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	543
385		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-methylmorpholino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
386		N-((3R,4S)-4-((2-(2,6-difluoro-	557






Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		3,5-dimethoxyphenyl)-4-(2-methylmorpholino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	
387		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2,6-dimethylmorpholino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
388		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(2,6-dimethylmorpholino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
389		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-hydroxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
390		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
391		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
392		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)	603

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		tetrahydrofuran-3-yl)acrylamide	
393		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-hydroxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
394		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	617
395		N-((3R,4S)-4-((2-(2,6-difluoro-3,5-dimethoxyphenyl)-4-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	585
396		N-((3R,4S)-4-((4-(4-cyano-4-methylpiperidin-1-yl)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	612
397		N-(6-(((3S,4R)-4-acrylamidotetrahydrofuran-3-yl)amino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)cyclopropanecarboxamide	573
398		N-(6-(((3S,4R)-4-acrylamidotetrahydrofuran-3-yl)amino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-4-yl)tetrahydrofuran-2-carboxamide	603

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
399		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(methylsulfonylamino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	583
400		N-((3S,4S)-3-((4-((cyclopropylmethyl)amino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	573
401		N-((3S,4S)-3-((4-((cyclopropylmethyl)amino)-2-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	541
402		N-((3S,4S)-3-((4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	577
403		N-((3S,4S)-3-((4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
404		N-((3S,4S)-3-((4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603
405		N-((3S,4S)-3-((4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603


Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
406		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((3,3-difluorocyclopentyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	623
407		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(neopentylamino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
408		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	613
409		N-((3S,4S)-3-((4-(cyclopropylamino)-2-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	559
410		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
411		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	595

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
412		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603
413		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	601
414		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(6-oxa-2-azaspiro[3.4]octan-2-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	615
415		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(7-oxa-2-azaspiro[3.5]nonan-2-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	629
416		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603
417		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	617

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
418		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(2,6-dimethylmorpholino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	617
419		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-hydroxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	617
420		N-((3S,4S)-3-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	631
752		N-((3R,4S)-4-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-(methylamino)pyrido[3,4-d]pyrimidin-6-yl)amino)tetrahydrofuran-3-yl)acrylamide	519


Example 421 Preparation of N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide

[0300]



Step 1: preparation of N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine

[0301]




[0302] N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine-8-amine (210 mg, 0.465 mmol) was dissolved in dichloromethane (6 mL), then *m*-chloroperoxybenzoic acid (200 mg, 1.163 mmol) was added, and the mixture was stirred at room temperature for 18 h. After the reaction was completed, a saturated sodium sulfite solution was added, and then the mixture was stirred for 5 min and extracted with dichloromethane, the organic phase was washed with a saturated sodium bicarbonate and the saturated brine, dried over anhydrous sodium sulfate, filtered, concentrated, and separated by column chromatography (Eluent: PE/EA = 2/1) to obtain compound N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine (186 mg, yield: 82.7%).

[0303] MS *m/z* (ESI): 483.4 [M+H]⁺.

Step 2: preparation of (±)-N²-((3*S*,4*R*)-4-aminotetrahydrofuran-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine

[0304]




N-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine-8-amine (190 mg, 0.393 mmol) and (3*R*,4*S*)-tetrahydrofuran-3,4-diamine dihydrochloride (206 mg, 1.179 mmol) were dissolved in acetonitrile (6 mL), then *N,N*-diisopropylethylamine (507 mg, 3.93 mmol) was added, and the mixture was heated to reflux, and stirred for 20 h. The reaction was completed, and the mixture was cooled to room temperature. The reaction liquid was diluted with EtOAc (20 mL), washed with saturated brine, dried over anhydrous sodium sulfate, filtered, concentrated, and separated by TLC (developing agent: CH₂Cl₂/MeOH 10/1) to obtain compound (±)-N²-((3*S*,4*R*)-4-aminotetrahydrofuran-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine.

uran-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine (120 mg, yield: 60%).

[0305] MS m/z (ESI): 505.4 [M+H]⁺.

Step 3: preparation of (±)-N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide

[0306]





[0307] At 0 °C, acryloyl chloride (22.6 mg, 0.249 mmol) was added to the solution of (±)-N²-((3S,4R)-4-aminotetrahydrofuran-3-yl)-N⁸-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2,8-diamine (120 mg, 0.237 mmol) and sodium bicarbonate (79.6 mg, 0.948 mmol) in the mixture of tetrahydrofuran (6.4 mL) and water (1.6 mL). After addition, the mixture was stirred at 0 °C for 5 min. The reaction liquid was diluted with EtOAc (10 mL), washed with saturated brine, dried over anhydrous sodium sulfate, filtered, concentrated and separated by TLC (Eluent: CH₂Cl₂/MeOH = 20/1) to obtain compound (±)-N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide (93 mg, yield: 70%).

[0308] ¹H NMR (400 MHz, CDCl₃) δ 8.89 (s, 1H), 6.80 (brs, 1H), 6.68 (s, 1H), 6.62 (s, 1H), 6.38 (brs, 1H), 6.21 (dd, J = 17.0, 1.5 Hz, 1H), 6.02 (dd, J = 17.0, 10.2 Hz, 1H), 5.56 (dd, J = 10.1, 1.5 Hz, 1H), 4.93 - 4.83 (m, 2H), 4.30 - 4.18 (m, 2H), 3.96 (s, 6H), 3.90 (dd, J = 9.7, 3.2 Hz, 1H), 3.79 (dd, J = 9.2, 5.2 Hz, 1H), 3.52 - 3.37 (m, 2H), 0.88 (t, J = 6.0 Hz, 1H), 0.57 - 0.48 (m, 2H), 0.36 - 0.26 (m, 2H). MS m/z (ESI): 559.5 [M+H]⁺.

Step 4: preparation of N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide

[0309]






[0310] (±)-N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide (93 mg, 0.166 mmol) was separated by chiral HPLC to obtain N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide (43.5 mg, ee value >98%, yield: 46.8%) and N-((3S,4R)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide (39 mg, ee value >98%, yield: 41.9%).


Example 433 Preparation of N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide

[0311]



Step 1: preparation of 8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine


[0312]



[0313] The compound was prepared referring to the synthesis method of step 1 of Example 421.

Step 2: preparation of tert-butyl ((3R,4S)-4-((8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)carbamate


[0314]



[0315] 8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine (409 mg, 0.912 mmol) and tert-butyl ((3R,4S)-4-aminotetrahydrofuran-3-yl)carbamate (553 mg, 2.736 mmol) were dissolved in acetonitrile (50 mL), then trifluoroacetic acid (31 mg, 0.274 mmol) was added. The reaction liquid was stirred at 95 °C for 4 h. After being cooled, the mixture was diluted with ethyl acetate (50 mL), washed with a saturated sodium bicarbonate solution and saturated brine, dried over anhydrous sodium sulfate, concentrated and separated by column chromatography (CH₂Cl₂/MeOH 0 - 4%) to obtain compound tert-butyl ((3R,4S)-4-((8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)carbamate (414 mg, yield: 79.6%). MS m/z (ESI): 570.4, 572.4 [M+H]⁺.

Step 3: preparation of tert-butyl ((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)carbamate


[0316]



[0317] A solution of tert-butyl ((3R,4S)-4-((8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)carbamate (60 mg, 0.105 mmol) and methylamine (1.5 mL, 33% ethanol solution) in N-methylpyrrolidone (1 mL) was heated to 110 °C and stirred for 18 h. The reaction liquid was diluted with ethyl acetate (5 mL), washed with saturated brine, dried over anhydrous sodium sulfate, concentrated, and separated by using a preparative TLC (PE/EA = 1:1) to obtain compound tert-butyl ((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)carbamate (32 mg, yield: 54%). MS m/z (ESI): 565.4, 567.4 [M+H]⁺.

Step 4: preparation of N²-((3S,4R)-4-aminotetrahydrofuran-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N⁸-methylpyrido[3,4-d]pyrimidine-2,8-diamine trifluoroacetate


[0318]



[0319] A solution of tert-butyl ((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)carbamate (32 mg, 0.057 mmol) in the solution of trifluoroacetic acid in dichloromethane (2 mL, 20%) was stirred at room temperature for 1 h, and then concentrated to obtain an oil product which was directly used in the next step. MS m/z (ESI): 465.4, 467.4 $[M+H]^+$.

Step 5: preparation of N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide

[0320]




[0321] At 0 °C, acryloyl chloride (5.7 mg, 0.063 mmol) was added to a solution of N²-((3S,4R)-4-aminotetrahydrofuran-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)-N⁸-methylpyrido[3,4-d]pyrimidine-2,8-diamine trifluoroacetate (the crude product) and NaHCO₃ (57 mg, 0.684 mmol) in the mixture of tetrahydrofuran (3.2 mL) and water (0.8 mL). After addition, the mixture was stirred at 0 °C for 5 min. The reaction liquid was diluted with EtOAc (mL), washed with a saturated sodium bicarbonate and saturated brine, dried over anhydrous sodium sulfate, concentrated and separated by using a preparative TLC (CH₂Cl₂/MeOH 20:1) to obtain compound N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide (17.6 mg, yield: 59.5%).

[0322] ¹H NMR (400 MHz, CDCl₃) δ 8.88 (s, 1H), 6.67 (s, 1H), 6.62 (s, 1H), 6.30 (brs, 1H), 6.19 (d, J = 16.9 Hz, 1H), 5.97 (dd, J = 17.0, 10.3 Hz, 1H), 5.88 (brs, 1H), 5.54 (d, J = 10.4 Hz, 1H), 4.95 - 4.83 (m, 2H), 4.29 - 4.18 (m, 2H), 3.95 (s, 6H), 3.89 (dd, J = 9.7, 3.0 Hz, 1H), 3.77 (dd, J = 9.3, 5.5 Hz, 1H), 3.14 (d, J = 4.4 Hz, 3H). MS m/z (ESI): 519.4, 521.4 $[M+H]^+$.


Example 775 N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide

[0323]



Step 1: preparation of 8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine


[0324]



[0325] 8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylthio)pyrido[3,4-d]pyrimidine (930 mg, 2.42 mmol) was dissolved in DCM (50 mL), and *m*-CPBA (1.23g, 6.05 mmol) was added, and the mixture was stirred at room temperature for 2 h. The reaction was completed, and sodium thiosulfate was added to quench the reaction. The mixture was extracted with DCM and separated by silica gel column chromatography to obtain compound 8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine (800 mg, yield:79%). MS *m/z* (ESI): 416 [M+H]⁺.

Step 2: preparation of N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2-amine

[0326]




[0327] 8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)-2-(methylsulfonyl)pyrido[3,4-d]pyrimidine (1g, 2.41 mmol) was dissolved in the mixture of tert-butyl alcohol (80 mL) and DCE (20 mL),

then DIPEA (1.55 g, 12.05 mmol) was added, the mixture was heated to 90 °C and stirred overnight. The mixture was extracted with DCM and separated by silica gel column chromatography to obtain compound N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2-amine (650 mg, yield:56%). MS m/z (ESI): 478 [M+H]⁺.

Step 3: preparation of N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidine-2-amine


[0328]



[0329] N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine-2-amine (325 mg, 0.68 mmol), 3-methoxyazetidine hydrochloride (252 mg, 2.04 mmol) and DIPEA (439 mg, 3.4 mmol) were dissolved in n-butanol (15 mL), the mixture was heated to 100 °C for 4 h, and then the mixture was concentrated, extracted with ethyl acetate and separated by silica gel column chromatography to obtain compound N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidine-2-amine (350 mg, yield: 97%). MS m/z (ESI): 529 [M+H]⁺.

Step 4: preparation of (3S,4S)-N³-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydro-2H-pyran-3,4-diamine

[0330]




[0331] N-((3S,4S)-4-azidotetrahydro-2H-pyran-3-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidine-2-amine (350 mg, 0.662 mmol) was dissolved in the mixture of THF (10 mL) and water (1 mL), and then triphenylphosphine (521 mg, 1.99 mmol) was added, the mixture was heated to 80 °C and stirred for 16 h. The mixture was cooled, directly dried over anhydrous sodium sulfate, concentrated, and separated by silica gel

column chromatography to obtain compound (3S,4S)-N³-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydro-2H-pyran-3,4-diamine (290 mg, yield:87%). MS m/z (ESI): 503 [M+H]⁺.

Step 5: preparation of N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide

[0332]



[0333] (3S,4S)-N³-(6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)tetrahydro-2H-pyran-3,4-diamine (290 mg, 0.58 mmol) was dissolved in the mixture of THF (20 mL) and water (5 mL), then NaHCO₃ (243 mg, 2.89 mmol) was added, and acryloyl chloride solution (63 mg, 0.69 mmol, dissolved in 1 mL THF) was added dropwise at room temperature, the mixture was stirred at room temperature for 10min. The reaction was completed determined by TLC, a saturated aqueous solution of NaHCO₃ was added to quench the reaction, the mixture was extracted with ethyl acetate, concentrated and separated by silica gel column chromatography to obtain compound N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide (204 mg, yield:63%). MS m/z (ESI): 557 [M+H]⁺.

[0334] ¹H NMR (400 MHz, CDCl₃) δ 8.92 (s, 1H), 6.96 (s, 1H), 6.69 (t, J = 8.0 Hz, 1H), 6.65 (s, 1H), 6.25 (dd, J = 17.0, 1.4 Hz, 1H), 6.08 (s, 1H), 6.02 (dd, J = 16.9, 10.3 Hz, 1H), 5.60 (dd, J = 10.3, 1.4 Hz, 1H), 4.73 (brs, 1H), 4.56 (brs, 1H), 4.42 (d, J = 8.5 Hz, 1H), 4.39 - 4.22 (m, 4H), 4.05 (dd, J = 12.0, 4.5 Hz, 1H), 3.99 (d, J = 11.7 Hz, 1H), 3.92 (s, 6H), 3.75 (dd, J = 11.9, 1.6 Hz, 1H), 3.65 - 3.57 (m, 1H), 3.33 (s, 3H), 2.09 - 2.02 (m, 1H), 1.91 - 1.81 (m, 1H).








[0335] Examples 422-714 and 753-806 were prepared referring to the synthesis method of Example 433, 421 or 775.








Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
422		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	563







Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
423		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	520
424		N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	527
425		N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-bis(methoxy- d_3)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	565
426		N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-difluoro-3,5-bis(methoxy- d_3)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	533
427		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-hydroxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	549
428		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(isopropylthio)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	607
429		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(ethylsulfonyl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	625








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
430		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(methylsulfonyl)amino)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	626
431		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((dimethylamino)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	576
432		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((dimethylamino)propyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	590
434		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	533
435		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2,2,2-trifluoroethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
436		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(dimethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	533
437		N-((3R,4S)-4-((8-((cyclopropylmethyl)(methyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	573








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		5-(dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-ylamino)tetrahydrofuran-3-yl)acrylamide	
438A		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((R)-tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
438B		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((S)-tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
439		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
440		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((oxetan-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
441		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydro-2H-pyran-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
442		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	561

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
443		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
444		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
445		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
446		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
447		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpyrrolidin-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
448		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylazetidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	588

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
449		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylazetidin-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
450		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpiperidin-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	602
451		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpiperidin-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	616
452		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,3-difluorocyclobutyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	595
453		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,3-difluorocyclopentyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	609
454		N-((3R,4S)-4-((8-((cyclopentylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
455		N-((3R,4S)-4-((8-(benzylamino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	595








Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
		yrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	
456		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(phenethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	609
457		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(phenylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	581
458		N-((3R,4S)-4-((8-((3-aminobenzyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	610
459		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	599
460		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-(2-hydroxyethyl)-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	629
461		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	643

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
		yl)methyl)amin o)pyrido[3,4-d]pyrimidin-2-yl)amin o)tetrahydrofuran-3 -yl)acrylamide	
462		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(neopentylamino)pyrido[3,4-d]pyrimidin-2-yl)amino) tetrahydrofuran-3-yl)acrylamide	575
463		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((3 -(isopropylam ino)propyl)amino)pyrido[3,4-d]pyri midin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	604
464		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((4-(isopropylam ino)butyl)amino)pyrido[3,4-d]pyrimi din-2-yl)amino)tetrahydrofuran-3 -yl) acrylamide	618
465		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((4-(dimethylami no)butyl)amino)pyrido[3,4-d]pyrimi din-2-yl)amino)tetrahydrofuran-3 -yl) acrylamide	604
466		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((4-(pyrrolidin-1 -yl)butyl)amino)pyrido[3,4-d]pyrimi din-2-yl)amino)tetrahydrofuran-3 -yl) acrylamide	630
467		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((2-(4-methylpip erazin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofu ran-3-yl)acrylamide	631








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
			
468		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-morpholinoethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	618
469		N-((3R,4S)-4-((8-((2-(3-aminopyrrolidin-1-yl)ethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	617
470		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(3-(dimethylamino)pyrrolidin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	645
471		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(2-(dimethylamino)ethoxy)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	620
472		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(2-(hydroxymethyl)pyrrolidin-1-yl)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	660
473		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(3,3-difluoropyrrolidin-1-yl)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	666

Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
474		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(3-methoxypropyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	660
475		N-((3R,4S)-4-((6-(2-chloro-3,5-dimethoxyphenyl)-8-((cyclopropylmethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	525
476		N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2-fluoro-3-methoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	479
477		N-((3R,4S)-4-((6-(2-chloro-3-(methoxy-d ₃)phenyl)-8-((cyclopropylmethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	498
478		N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2-fluoro-3-(methoxy-d ₃)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	482
479		N-((3R,4S)-4-((6-(2-chloro-3,5-bis(methoxy-d ₃)phenyl)-8-((cyclopropylmethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	531
480		N-((3R,4S)-4-((8-((cyclopropylmethyl)amino)-6-(2-fluoro-3,5-bis(methoxy-d ₃)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	515

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
481		N-((3R,4S)-4-(((6-(2-chloro-3-methoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	525
482		N-((3R,4S)-4-(((6-(2-chloro-3-methoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	538
483		N-((3R,4S)-4-(((6-(2-fluoro-3-methoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	522
484		N-((3R,4S)-4-(((6-(2-chloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	568
485		N-((3R,4S)-4-(((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	570
486		N-((3R,4S)-4-(((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
487		N-((3R,4S)-4-(((6-(2-chloro-3,5-dimethoxyphenyl)-8-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	555

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	
488		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((2-(4-methylpiperazin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	599
489		N-((3R,4S)-4-((6-(2-chloro-3-methoxyphenyl)-8-((2-(4-methylpiperazin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	567
490		N-((3R,4S)-4-((6-(2-chloro-3,5-dimethoxyphenyl)-8-((2-(4-methylpiperazin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	597
491		N-((3R,4S)-4-((6-(3,5-dimethoxyphenyl)-8-((2-(4-methylpiperazin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	563
492		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	531
493		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((1-methylpiperidin-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	570

Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
494		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	567
495		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
496		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-ethoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	534
497		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-isopropoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	548
498		N-((3R,4S)-4-((8-(cyclopropylmethoxy)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	560
499		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	577
500		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	545

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
		d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
501		N-((3S,4S)-3-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	573
502		N-((3S,4S)-3-((8-((cyclopropylmethyl)amino)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	541
503		N-((3S,4S)-3-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-bis(methoxy-d3)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	579
504		N-((3S,4S)-3-((8-((cyclopropylmethyl)amino)-6-(2,6-difluoro-3,5-bis(methoxy-d3)phenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	547
505		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-hydroxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	563
506		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((isopropylthio)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	621
507		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-hydroxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	639

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		(ethylsulfonyl)ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
508		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(dimethylamino)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	590
509		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3-(dimethylamino)propyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	604
510		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	549
511		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	547
512		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2,2,2-trifluoroethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	601
513		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2,2,2-trifluoroethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	547

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
514		N-((3S,4S)-3-((8-((cyclopropylmethyl)(methyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	587
515		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(methylsulfonylamido)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	640
516		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603
517		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	571
518		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603
519		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((oxetan-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
520		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydro-2H-pyran-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	617







Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
521		N-((3 S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(oxetan-3-ylami no)pyrido[3,4-d]pyrimidin-2-yl)ami no)tetrahydro-2H-pyran-4-yl)acryla mide	575
522		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((tetrahydrofuran -3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-y l)acrylamide	589
523		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-d imethoxyphenyl)-8-((tetrahydrofuran -3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-y l)acrylamide	557
524		N-((3 S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyri midin-2-yl)amino)tetrahydro-2H-pyr an-4-yl)acrylamide	603
525		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-d imethoxyphenyl)-8-((tetrahydro-2H-pyran-4-yl)amino)pyrido[3,4-d]pyri midin-2-yl)amino)tetrahydro-2H-pyr an-4-yl)acrylamide	571
526		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(((1-methylpyrro lidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2 H-pyran-4-yl)acrylamide	616








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
527		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	584
528		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	616
529		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	584
530		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	602
531		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((1-methylpyrrolidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	570
532		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylazetidin-3-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	602
533		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-	588

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d imethoxyphenyl)-8-((1-methylazetidin-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
534		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpiperidin-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	616
535		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((1-methylpiperidin-4-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	584
536		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methylpiperidin-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	630
537		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,3-difluorocyclobutyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	609
538		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,3-difluorocyclopentyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	623
539		N-((3S,4S)-3-((8-((cyclopentylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	601







Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
540		N-((3S,4S)-3-((8-(benzylamino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	609
541		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(phenethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	623
542		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(phenylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	595
543		N-((3S,4S)-3-((8-((3-aminobenzyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	624
544		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	613
545		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(((1-methyl-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	581







Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
546		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-(2-hydroxyethyl)-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	643
547		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-(2-methoxyethyl)-1H-pyrazol-4-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	657
548		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(neopentylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
549		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3-(isopropylamino)propyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	618
550		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(isopropylamino)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	632
551		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(dimethylamino)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	618
552		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(pyrrolidin-1-yl)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	644

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		1-yl)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
553		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(4-methylpiperazin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	645
554		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-morpholinoethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	632
555		N-((3S,4S)-3-((8-((2-(3-aminopyrrolidin-1-yl)ethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	631
556		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(3-(dimethylamino)pyrrolidin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	659
557		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-(2-(dimethylamino)ethoxy)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	634
558		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(2-(hydroxymethyl)pyrrolidin-1-yl)ethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	674







Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)butyl)amino) pyrido[3,4-d]pyrimidin-2-yl)amino) tetrahydro-2H-pyran-4-yl)acrylamide	
559		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(3,3-difluoropyrrolidin-1-yl)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	680
560		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((4-(3-methoxypyrrolidin-1-yl)butyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	674
561		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-methoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	534
562		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-ethoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	548
563		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-isopropoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	562
564		N-((3S,4S)-3-((8-(cyclopropylmethoxy)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	574

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
565		N-((3 S,4R)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)pyrido[3,4 - d]pyrimi din-2-yl)amino)-1- methylpyrrolidin-3-yl)acrylamide	503
566		N-((3S,4R)-1-acetyl-4-((6-(2,6-dichl oro-3,5- dimethoxyphenyl)pyrido[3,4 - d]pyrimidin-2-yl)amino)pyrrolidin-3- yl)acrylamide	531
567		N-((3S,4R)-1-acetyl-4-((6-(2,6-dichl oro-3,5-dimethoxyphenyl)-8- (methyl amino)pyrido[3,4- d]pyrimidin-2-yl)a mino)pyrrolidin- 3-yl)acrylamide	560
568		N-((3 S,4R)-4-((6-(2,6-dichloro-3,5- d imethoxyphenyl)pyrido[3,4- d]pyrimi din-2-yl)amino)-1-(2- (dimethylamin o)ethyl)pyrrolidin-3- yl)acrylamide	560
569		(3 S,4R)-3-acrylamido-4-((6-(2,6- dic hloro-3,5- dimethoxyphenyl)pyrido[3 ,4- d]pyrimidin-2-yl)amino)-N -methy lpyrrolidine-1-carboxamide	546
570		N-((3S,4R)-4-((8-((cyclopropylmeth yl)amino)-6-(2,6-dichloro-3, 5-dimet hoxyphenyl)pyrido[3,4-d]pyrimidin- 2-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	572
571		(3 S,4R)-3-acrylamido-4-((8- ((cyclop ropylmethyl)amino)-6-(2,6- dichloro-3,5-	615

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-N-methylpyrrolidine-1-carboxamide	
572		N-((3S,4R)-1-acetyl-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)pyrrolidin-3-yl)acrylamide	600
573		N-((3S,4R)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	614
574		N-((3S,4R)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	629
575		N-((3S,4R)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	624
576		N-((3S,4R)-4-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	638
577		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)amino)-1-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(2,6-dichloro-3,5-dimethoxyphenyl)pyrrolidin-3-yl)acrylamide	602

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	
578		(3 S,4R)-3-acrylamido-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-N-methylpyrrolidine-1-carboxamide	645
579		N-((3S,4R)-1-acetyl-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)pyrrolidin-3-yl)acrylamide	630
580		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	644
581		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	659
582		N-((3 S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	654

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
583		N-((3 S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((tetrahydrofuran-2-yl)methyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	668
584		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	574
585		(3 S,4R)-3-acrylamido-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-N-methylpyrrolidine-1-carboxamide	617
586		N-((3 S,4R)-1-acetyl-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)pyrrolidin-3-yl)acrylamide	602
587		N-((3 S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(oxetan-3-yl)pyrrolidin-3-yl)acrylamide	616
588		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(oxetan-3-ylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	631
589		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-	626

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d imethoxyphenyl)-8-(oxetan-3 -ylami no)pyrido[3,4-d]pyrimidin-2-yl)ami no)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	
590		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(oxetan-3 -ylami no)pyrido[3,4-d]pyrimidin-2-yl)ami no)-1-(1-methyl-1H-pyrazol-4-yl)py rrolidin-3 -yl)acrylamide	640
591		N-((3 S,4R)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((tetrahydrofuran -3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-methylpyrrolidin-3-yl)acrylamide	588
592		(3 S,4R)-3-acrylamido-4-((6-(2,6-dic hloro-3,5-dimethoxyphenyl)-8-((tetr ahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-N-methylp yrrolidine-1-carboxamide	631
593		N-((3S,4R)-1-acetyl-4-((6-(2,6-dichl oro-3, 5-dimethoxyphenyl)-8-((tetrah ydrofuran-3-yl)amino)pyrido[3,4-d] pyrimidin-2-yl)amino)pyrrolidin-3-y l)acrylamide	616
594		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((tetrahydrofuran -3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(oxetan-3-yl)pyrrolidi n-3-yl)acrylamide	630







Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
595		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(2-(dimethylamino)ethyl)pyrrolidin-3-yl)acrylamide	645
596		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	640
597		N-((3S,4R)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)-1-(1-methyl-1H-pyrazol-4-yl)pyrrolidin-3-yl)acrylamide	654
598		N-((3R,4S)-4-((8-(((R)-1-cyclopropylethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	573
599		N-((3R,4S)-4-((8-(((R)-1-cyclopropylethyl)amino)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	541
600		N-((3R,4S)-4-((8-(((S)-1-cyclopropylethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	573
601		N-((3R,4S)-4-((8-(((S)-1-cyclopropylethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	541

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
		lethyl)amino)-6-(2,6-difluoro-3,5-di methoxyphenyl)pyrido[3,4-d]pyrimi din-2-yl)amino)tetrahydrofuran-3 - yl) acrylamide	
602		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(((R)-3,3 - dimeth ylbutan-2- yl)amino)pyrido[3,4-d]pyr imidin-2- yl)amino)tetrahydrofuran-3 - yl)acrylamide	589
603		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-d imethoxyphenyl)-8-(((R)-3,3 - dimeth ylbutan-2- yl)amino)pyrido[3,4-d]pyr imidin-2- yl)amino)tetrahydrofuran-3 - yl)acrylamide	557
604		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(((S)-3,3 - dimeth ylbutan-2- yl)amino)pyrido[3,4-d]pyr imidin-2- yl)amino)tetrahydrofuran-3 - yl)acrylamide	589
605		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-d imethoxyphenyl)-8-(((S)-3,3 - dimeth ylbutan-2- yl)amino)pyrido[3,4-d]pyr imidin-2- yl)amino)tetrahydrofuran-3 - yl)acrylamide	557
606		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8- methylpyrido[3, 4-d]pyrimidin-2- yl)amino)tetrahydro furan-3- yl)acrylamide	504
607		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8- ethylpyrido[3,4-d]pyrimidin-2- yl)amino)tetrahydrofu ran-3- yl)acrylamide	518








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
608		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-isopropylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	532
609		N-((3R,4S)-4-((8-cyclopropyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	530
610		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-neopentylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	560
611		N-((3R,4S)-4-((8-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	544
612		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
613		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-2-yl)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	574
614		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	574

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acryla mide	
615		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((1-methylpyrrol idin-3-yl)methyl)pyrido[3,4-d]pyrim idin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
616		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-((1-methylpyrrol idin-2-yl)methyl)pyrido[3,4-d]pyrim idin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
617		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-di methoxyphenyl)-8-(1-methylpiperidin -4-yl)pyrido[3,4-d]pyrimidin-2-yl)ami no)tetrahydrofuran-3-yl)acrylamide	587
618		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(methoxymethyl)pyrido[3,4-d]pyrimidin-2-yl)amino) tetrahydrofuran-3-yl)acrylamide	534
619		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(2-methoxyethyl)pyrido[3,4-d]pyrimidin-2-yl)amino) tetrahydrofuran-3-yl)acrylamide	548
620		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(2-(isopropylami no)ethyl)pyrido[3,4-d]pyrimidin-2-y l)amino)tetrahydrofuran-3-yl)acryla mide	575








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
621		N-((3R,4S)-4-((8-((cyclopropylamino)methyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	559
622		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylsulfonylmethyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	597
623		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(((1-methylethyl)sulfonamido)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	625
624		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2,2,2-trifluoroethyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	572
625		N-((3R,4S)-4-((8-benzyl-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	580
626		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-phenylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	566
627		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(pyridin-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	567

Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
628		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(pyridin-3-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	567
629		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	570
630		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methyl-1H-pyrazol-4-yl)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	584
631		N-((3R,4S)-4-((8-(1-cyclopropyl-1H-pyrazol-4-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	596
632		N-((3R,4S)-4-((8-(1-(cyclopropylmethyl)-1H-pyrazol-4-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	610
633		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	614
634		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	600








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		hydroxyethyl-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	
635		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-(tetrahydrofuran-3-yl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	626
636		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-phenethylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	594
637		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	538
638		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	582
639		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-(2-hydroxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	568
640		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-ethylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	532

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)acrylamide	
641		N-((3S,4S)-3-((8-(cyclopropylmethyl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	558
642		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	588
643		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	588
644		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	584
645		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((1-methyl-1H-pyrazol-4-yl)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	598
646		N-((3S,4S)-3-((8-(1-cyclopropyl-1H-pyrazol-4-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	610

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
647		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	628
648		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-(isopropylamino)ethyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
649		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-ethylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	500
650		N-((3S,4S)-3-((8-(cyclopropylmethyl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	526
651		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((tetrahydrofuran-3-yl)methyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	556
652		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(tetrahydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	556
653		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-methyl-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	552

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)amino)tetrahydro-2H-pyran-4-yl) acrylamide	
654		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	596
655		N-((3R,4S)-4-((8-(cyclopropylamino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	545
656		N-((3R,4S)-4-((8-(cyclopropylamino)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	513
657		N-((3R,4S)-4-((8-(azetidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	545
658		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
659		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	543

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
660		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-(dimethylamino)azetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	588
661		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-(trifluoromethyl)azetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	613
662		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-dimethylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	573
663		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	581
664		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
665		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
666		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-	557

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		3-yl)acrylamide	
667		N-((3R,4S)-4-((8-(3-cyano-3-methylazetidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	584
668		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
669		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(2-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	555
670		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(1-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
671		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-oxa-6-azaspiro[3.3]heptan-6-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	555
672		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-azaspiro[3.3]heptan-2-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	585

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
673		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-azaspiro[3.4]octan-2-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	599
674		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(6-oxa-2-azaspiro[3.4]octan-2-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	601
675		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(7-oxa-2-azaspiro[3.5]nonan-2-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	615
676		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(pyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	559
677		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
678		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
679		N-((3R,4S)-4-((8-(3-cyanopyrrolidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	584








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		amino)tetrahydrofuran-3-yl)acrylamide	
680		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-difluoropyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	595
681		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
682		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
683		N-((3R,4S)-4-((8-(3-cyano-3-methylpyrrolidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	598
684		N-((3R,4S)-4-((8-(3-azabicyclo[3.1.0]hexan-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
685		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-oxa-7-azaspiro[4.4]nonan-7-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	615

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
686		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-morpholinopyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	575
687		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-morpholinopyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	543
688		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2-methylmorpholino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
689		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(2-methylmorpholino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
690		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2,6-dimethylmorpholino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
691		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(2,6-dimethylmorpholino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
692		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-hydroxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d]pyrimidin-2-yl amino)tetrahydrofuran-3-yl)acrylamide	
693		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
694		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	571
695		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
696		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-hydroxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	603
697		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	617
698		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	585








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
699		N-((3R,4S)-4-((8-(4-cyano-4-methylpiperidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	612
700		N-(2-(((3S,4R)-4-acrylamidotetrahydrofuran-3-yl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-8-yl)cyclopropanecarboxamide	573
701		N-(2-(((3S,4R)-4-acrylamidotetrahydrofuran-3-yl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-8-yl)tetrahydrofuran-2-carboxamide	603
702		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(methylsulfonylamido)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	583
703		N-((3S,4S)-3-((8-(cyclopropylamino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	559
704		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
705		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	595

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
706		N-((3 S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(3-methoxy-3-m ethylazetidin-1-yl)pyrido[3,4-d]pyri midin-2-yl)amino)tetrahydro-2H-pyr an-4-yl)acrylamide	603
707		N-((3 S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(2-oxa-6- azaspir o[3.3]heptan-6-yl)pyrido[3,4-d]pyri midin-2-yl)amino)tetrahydro-2H-pyr an-4-yl)acrylamide	601
708		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(6-oxa-2- azaspir o[3.4]octan-2-yl)pyrido[3,4-d]pyrimi din-2-yl)amino)tetrahydro-2H-pyran -4-yl)acrylamide	615
709		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(7-oxa-2- azaspir o[3.5]nonan-2-yl)pyrido[3,4-d]pyrim idin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	629
710		N-((3 S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(3-methoxypyrro lidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl) acrylamide	603
711		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-d imethoxyphenyl)-8-(3-methoxy-3-m ethylpyrrolidin-1-yl)pyrido[3,4-d]py rimidin-2-yl)amino)tetrahydro-	617

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		2H-pyran-4-yl)acrylamide	
712		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(2,6-dimethylmorpholino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	617
713		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-hydroxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	617
714		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(4-methoxy-4-methylpiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	631
753		N-((3R,4S)-4-((8-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	524
754		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-bis(methoxy- d_3)phenyl)-8-ethoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	540
755		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3,5-dimethoxybenzyl)oxy)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	656








Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
756		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3,6-dihydro-2H-pyran-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	572
758		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((3-hydroxy-2,2-bis(hydroxymethyl)propyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	623
759		N-((3R,4S)-4-((8-(3,3-bis(hydroxymethyl)azetidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	605
760		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
761		N-((3R,4S)-4-((6-(2-chloro-3,5-dimethoxyphenyl)-8-(2-methylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	536
762		N-((3R,4S)-4-((8-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587
763		N-((3R,4S)-4-((8-(2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	587

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	
764		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	487
765		N-((3R,4S)-4-((8-amino-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	505
766		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-bis(methoxy-d3)phenyl)-8-((methyl-d3)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	528
767		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	501
768		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	549
769		N-((3R,4S)-4-((8-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	555
770		N-((3R,4S)-4-((6-(2-fluoro-3-methoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	439




Example No.	Compound structure	Compound name	MS: <i>m/z</i> [M+1] ⁺
		furan-3-yl)acrylamide	
771		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((methyl- <i>d</i> ₃)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	522
772		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	557
773		N-((3R,4S)-4-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	543
774		N-((3R,4S)-4-((8-(azetidin-1-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	513
775		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxyazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	557
776		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(7-oxa-2-azaspiro[3.5]nonan-2-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	615

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
777		N-((3R,4S)-4-((6-(2,6-difluoro-3-methoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	457
778		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
779		N-((3R,4S)-4-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydrofuran-3-yl)acrylamide	589
780		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-morpholinopyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
781		N-((3S,4S)-3-((8-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	601
782		N-((3S,4S)-3-((8-(2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	601
783		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylpyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	603

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
784		N-((3S,4S)-3-((8-(azetidin-1-yl)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	559
785		N-((3S,4S)-3-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	589
786		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(methylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	501
787		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(ethylamino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	515
788		N-((3S,4S)-3-((8-(cyclopropylamino)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	527
789		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-morpholinopyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	557
790		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	571


Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		an-4-yl)acrylamide	
791		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3,3-difluoroazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	563
792		N-((3S,4S)-3-((8-(azetidin-1-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	527
793		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-hydroxy-3-methylazetidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	557
794		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(3-methoxypyrrolidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	571
795		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(4-methoxypiperidin-1-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	585
796		N-((3S,4S)-3-((8-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	569

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
797		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((2,2,2-trifluoroethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	569
798		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-((3,3-dimethylbutan-2-yl)amino)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	571
799		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-methoxypyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	502
800		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-methylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	486
801		N-((3S,4S)-3-((8-cyclopropyl-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	512
802		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(piperidin-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	555
803		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	538
804		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-(1-(2-hydroxyethyl)-1H-pyrazol-4-yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	582

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		yl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	
805		N-((3S,4S)-3-((6-(2,6-difluoro-3,5-dimethoxyphenyl)-8-phenylpyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	548
806		N-((3S,4S)-3-((8-chloro-6-(2,6-difluoro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)amino)tetrahydro-2H-pyran-4-yl)acrylamide	506


Example 715 Preparation of N-(5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide

[0336]



Step 1: preparation of 6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(1-(2-methoxyethyl)-4-nitro-1H-pyrazol-5-yl)pyrido[3,4-d]pyrimidine


[0337]



[0338] 1-(2-methoxyethyl)-4-nitro-1H-pyrazole (84 mg, 0.49 mmol) and 2-chloro-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidine (200 mg, 0.54 mmol) were dissolved into DMA (15 mL), then CuI (112 mg, 0.59 mmol), PivOH (15 mg, 0.15 mmol), K₂CO₃ (88 mg, 0.64 mmol) and Pd(PPh₃)₂Cl₂ (34 mg, 0.05 mmol) were added, the mixture was heated to 140 °C under N₂ and stirred for 1 h. The reaction was completed, and the mixture was concentrated to remove the solvent by reduced pressure and separated by fast silica gel column (PE/EA 2:1) to obtain compound 6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(1-(2-methoxyethyl)-4-nitro-1H-pyrazol-5-yl)pyrido[3,4-d]pyrimidine (103 mg, yield 42%). MS m/z (ESI): 505 [M+1]⁺.

Step 2: preparation of 5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-amine


[0339]



[0340] 6-(2,6-dichloro-3,5-dimethoxyphenyl)-2-(1-(2-methoxyethyl)-4-nitro-1H-pyrazol-5-yl)pyrido[3,4-d]pyrimidine (103 mg, 0.20 mmol) was dissolved in the mixture of EtOH and H₂O (10/3 mL), then Fe powder (114 mg, 2.04 mmol) and NH₄Cl (109 mg, 2.04 mmol) were added, the mixture was heated to reflux and stirred for 2 h. After the reaction was completed, the reaction liquid was extracted for three times with dichloromethane. The organic phases were combined, washed with water, then a saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, filtrated and concentrated. The crude product was separated by fast silica gel column chromatography (CH₂Cl₂/MeOH 10:1) to obtain compound 5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-amine (12 mg, yield 12%). MS m/z (ESI): 476 [M+1]⁺.

Step 3: preparation of N-(5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide

[0341]
















[0342] 5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-amine (12 mg, 0.025 mmol) was dissolved in the mixture of THF and H₂O (4:1, 4 mL), then NaHCO₃ (11 mg, 0.13 mmol) was added, the mixture was cooled under ice water bath, Acryloyl chloride (2.3 mg, 0.025 mmol) was added, and the mixture was stirred at 0 °C for 10 min. After the reaction was completed, the reaction liquid was concentrated by reduced pressure, separated by silica gel column chromatography (CH₂Cl₂/MeOH 10:1) to obtain compound N-(5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide (4.8 mg, yield 36%). MS m/z (ESI): 529 [M+1]⁺.

[0343] ¹H NMR (400 MHz, CDCl₃) δ 10.71 (s, 1H), 9.63 (s, 1H), 9.60 (s, 1H), 8.61 (s, 1H), 7.80 (s, 1H), 6.72 (s, 1H), 6.49 (d, *J* = 16.7 Hz, 1H), 6.40 (dd, *J* = 16.9, 9.7 Hz, 1H), 5.84 (d, *J* = 9.7 Hz, 1H), 5.21 (t, *J* = 6.0 Hz, 2H), 4.00 (s, 7H), 3.89 (t, *J* = 5.9 Hz, 2H), 3.35 (s, 3H).

[0344] Examples 716-729 were prepared by referring to the synthesis method of Example 715.


Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
716		N-(5-(7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	544
717		N-(5-(7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	558
718		N-(5-(7-(2-chloro-6-fluoro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	542
719		N-(5-(6-(2-(tert-butylamino)-2-oxoethyl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	657

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
		amide	
720		N-(5-(6-(cyclopropylmethyl)-7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	598
721		N-(5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	545
722		N-(5-(6-(2,6-dichloro-3,5-dimethoxyphenyl)-7-methyl-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	559
723		N-(5-(7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	528
724		N-(5-(7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	601
725		N-(5-(7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-methoxy-2,6-naphthyridin-3-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	558
726		N-(5-(3-(2,6-dichloro-3,5-	545

Example No.	Compound structure	Compound name	MS:m/z [M+1] ⁺
		dimethoxyphen yl)-1-oxo-1H-pyrano[4,3-c]pyridin-7-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	
727		N-(5-(6-(2,6-dichloro-3,5-dimethoxyphen yl)-8-methoxypyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1H-pyrazol-4-yl)acrylamide	560
728		N-(5-(6-(2,6-dichloro-3,5-dimethoxyphen yl)-8-((2-methoxyethyl)amino)pyrido[3,4 -d]pyrimidin-2-yl)-1-(2-methoxyethyl)-1 H-pyrazol-4-yl)acrylamide	602
729		N-(5-(8-((cyclopropylmethyl)amino)-6-(2 ,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)-1-(2-methoxyethyl) -1H-pyrazol-4-yl)acrylamide	598


Example 730 Preparation of N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-yl)methyl)acrylamide

[0345]



Step 1: preparation of 3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-carbonitrile


[0346]



[0347] 7-chloro-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-1,6-naphthyridin-2(1H)-one (250.0 mg, 0.626 mmol), Zn(CN)_2 (110 mg, 0.938 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (34.7 mg, 0.063 mmol) were added to the mixture of DMF and H_2O (5 mL, 100:1), then $\text{Pd}_2(\text{dba})_3$ (28.7 mg, 0.031 mmol) was added under N_2 , the mixture was heated to 120 °C and stirred for 20 h. The reaction was completed, and the mixture was cooled to room temperature, diluted with water and extracted with EtOAc. The organic phase was washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by silica gel column chromatography (Eluent: PE/EtOAc = 10-25%) to obtain compound 3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-carbonitrile (75 mg, yield: 28%). MS m/z (ESI): 390.0 $[\text{M}+\text{H}]^+$.

Step 2: preparation of 7-(aminomethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-1,6-naphthyridin-2(1H)-one


[0348]



[0349] 3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-2-carbonyl-1,2-dihydro-1,6-naphthyridin-7-carbonitrile (15 mg, 0.038 mmol) was dissolved in methanol (3 mL), and concentrated hydrochloric acid (15 mL) and 10% Pd/C (catalytic amount) were added. the mixture was stirred at room temperature for 2h under H_2 , the reaction was completed. The reaction liquid was filtrated and evaporated in vacuo to obtain the crude product 7-(aminomethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-1,6-naphthyridin-2(1H)-one, which can be directly used in the next reaction. MS m/z (ESI): 394.0 $[\text{M}+\text{H}]^+$.

Step 3: preparation of N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-yl)methyl)acrylamide

[0350]



[0351] At 0 °C, acryloyl chloride (3.8 mg, 0.042 mmol) was added to a solution of 7-(aminomethyl)-3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-1,6-naphthyridin-2(1H)-one (the crude product, 0.038 mmol) and NaHCO₃ (19.0 mg, 0.230 mmol) in the mixture of THF and H₂O (0.8 mL/0.2 mL), the mixture was stirred for 10 min At 0 °C. The reaction liquid was diluted with EtOAc (5 mL), washed with saturated brine, dried over anhydrous sodium sulfate, filtrated, concentrated and separated by a preparative TLC (Eluent: CH₂Cl₂/MeOH = 10:1) to obtain compound N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-methyl-2-oxo-1,2-dihydro-1,6-naphthyridin-7-yl)methyl)acrylamide (1.1 mg, two-step yield: 6%).









[0352] ¹H NMR (400 MHz, Chloroform-d) δ 8.75 (s, 1H), 7.68 (s, 1H), 7.40 (s, 1H), 6.65 (s, 1H), 6.38 - 6.32 (m, 1H), 6.27 - 6.21 (m, 1H), 5.73 - 5.69 (m, 1H), 5.30 (s, 1H), 4.81 (d, J = 5.6 Hz, 2H), 3.96 (s, 6H), 3.78 (s, 3H);



[0353] MS m/z (ESI): 448.0 [M+H]⁺.

[0354] Examples 731-751 were prepared referring to the synthesis method of Example 730.

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
731		N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-oxo-1-(2,2,2-trifluoroethyl)-1,2-dihydro-1,6-naphthyridin-7-yl)methyl)acrylamide	516
732		N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-2-oxo-1-(tetrahydrofuran-3-yl)-1,2-dihydro-1,6-naphthyridin-7-yl)methyl)acrylamide	504
733		N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-(2-methoxyethyl)-2-oxo-1,2-dihydro-1,6-naphthyridin-7-yl)methyl)acrylamide	492

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
734		N-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-2,6-naphthyridin-3-yl)methyl)acrylamide	418
735		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)methyl)acrylamide	419
736		N-((3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-oxo-1H-pyrano[4,3-c]pyridin-7-yl)methyl)acrylamide	435
737		N-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)methyl)acrylamide	434
738		N-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-6-methyl-5-oxo-5,6-dihydro-2,6-naphthyridin-3-yl)methyl)acrylamide	448
739		N-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-methoxy-2,6-naphthyridin-3-yl)methyl)acrylamide	448
740		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)methyl)acrylamide	435
741		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-7-methyl-8-oxo-7,8-dihydropyrido[3,4-d]pyrimidin-2-yl)methyl)acrylamide	449
742		N-((6-(2,6-dichloro-3,5-dimethoxy	449

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		phenyl)-8-methoxypyrido[3,4-d]pyrimidin-2-yl)methyl)acrylamide	
743		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)thieno[2,3-d]pyrimidin-2-yl)methyl)acrylamide	424
744		N-((2-(2,6-dichloro-3,5-dimethoxyphenyl)thiazolo[5,4-d]pyrimidin-5-yl)methyl)acrylamide	425
745		N-((2-(2,6-dichloro-3,5-dimethoxyphenyl)thiazolo[4,5-c]pyridin-6-yl)methyl)acrylamide	424
746		N-((2-(2,6-dichloro-3,5-dimethoxyphenyl)-4-oxo-4H-pyrano[2,3-c]pyridin-6-yl)methyl)acrylamide	435
747		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-8H-pyrano[3,2-d]pyrimidin-2-yl)methyl)acrylamide	436
748		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-oxo-5,8-dihydropyrido[3,2-d]pyrimidin-2-yl)methyl)acrylamide	435
749		N-((6-(2,6-dichloro-3,5-dimethoxyphenyl)-8-((2-methoxyethyl)amino)pyrido[3,4-d]pyrimidin-2-yl)methyl)acrylamide	492
750		N-((8-((cyclopropylmethyl)amino)-6-(2,6-dichloro-3,5-dimethoxyphenyl)pyrido[3,4-d]pyrimidin-2-yl)methyl)acrylamide	488

Example No.	Compound structure	Compound name	MS: m/z [M+1] ⁺
		nyl)pyrido[3,4-d]pyrimidin-2-yl)methylacrylamide	
751		N-((7-(2,6-dichloro-3,5-dimethoxyphenyl)-5-((2-methoxyethyl)amino)-2,6-naphthyridin-3-yl)methyl)acrylamide	491

Biological test and evaluation

I. *In vitro* biochemical kinase analysis of FGFR4

[0355] FGFR4 Caliper Assay was used in the present invention to determine the inhibitory activities of the compounds against FGFR4. The detailed experimental procedure was as follows.

1. The kinase reaction in the present invention was carried out in a 384-well plate, and 12.5 μ M of FGFR4, 65 μ M of ATP and 1 μ M of peptide (5 Fluo Ahx KKKKEEIYFFFG NH₂) were respectively added into the following reaction system.
2. A reaction system is a mixture solution of 50 mM HEPES, pH 7.5, 1 mM DTT, 0.02% Tween 20, 0.02% BSA, 0.6% DMSO, 10 mM beta glycerol phosphate and 10 μ M sodium orthovanadate and 16 mM MgCl₂.
3. The reaction system was incubated at 30 °C for 40 minutes.
4. The reaction was terminated by adding a stop solution (100 mM HEPES, pH 7.5, 5% DMSO, 0.1% Caliper coating reagent, 10 mM EDTA and 0.015% Brij35).
5. The culture plate with the terminated kinase reaction was transferred to the Caliper LC 3000 workstation to read the data, the phosphorylated and unphosphorylated peptides were separated by using the Caliper microfluid migration shift technique, and the analyte was transferred by allowing a constant buffer flow through the chip, the migration of the substrate peptide was monitored by the labeled fluorescent signal, and the kinase activity was calculated by using the amount of the phosphate-based peptide formed.
6. Finally, IC₅₀ values were determined by non-linear regression analysis of percent inhibition at different compound concentrations. The test results for the enzymatic activities of the compounds of the specific examples were shown in Table 1.

II. FGFR4 cell proliferation experiment

[0356] Cell Titer Glo (CTG) experiment was used in the present invention to evaluate the functional effects of the compounds on cell proliferation. Huh7 hepatocellular carcinoma cells (Catalog No. TChU182) from the Chinese Academy of Sciences cell bank were cultured in DMEM with high glucose (Gibco, cat. No. 1773536), 10% fetal bovine serum (Gibco, 10099-141) at 37 °C, in a 5% CO₂ incubator. Compound-mediated inhibition of cell proliferation/survival was assessed by quantification of cellular ATP levels using CTG reagent (Promega, #G7573). The specific experimental procedure was as follows:

1. The cells were seeded into a tissue culture medium-treated 96-well plate (Costar #3904) at 3500 cells/well/90 μ L of fresh medium;
2. 10 μ L of medium containing a compound concentration of 10 fold of its final test concentration was added;
3. The dose effect was evaluated by a 5-fold serial dilution of the test compound, starting from 10 μ M.
4. After cells incubation for 3 days at 37 °C under 5% CO₂, the effect of the inhibitor on cell proliferation was quantified after adding 50 μ L of CTG and testing with luminescence.
5. The concentration of the compound (EC₅₀) leading to half maximal growth inhibition and the concentration of compound (Absolute IC₅₀) leading to absolute half growth inhibition were determined using a four-parameter curve fit in Graphpad Prism in a plate reader (M1000, Tecan). The test results of cell activities for the compounds of specific examples were shown in Table 1.

Table 1 The results of enzymatic activity and cell activity tests

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
Example 1	13.9	1527	2446	¹ H NMR (400 MHz, CDCl ₃) δ 8.60 (s, 1H), 8.30 (d, J = 8.1 Hz, 1H), 7.31 (s, 1H), 7.10 - 7.05 (m, 2H), 6.68 (s, 1H), 6.52 - 6.48 (m, 1H), 6.44 (s, 1H), 6.39 (s, 1H), 6.33 (d, J = 1.4 Hz, 1H), 6.19 (d, J = 10.2 Hz, 1H), 5.68 (dd, J = 10.2, 1.4 Hz, 1H), 3.99 (s, 6H), 3.25 (s, 3H), 2.21 (s, 3H).
Example 5	26.1	>10000	>10000	¹ H NMR (400 MHz, CDCl ₃) δ 9.21 (s, 1H), 8.72 (s, 1H), 8.23 (s, 3H), 7.34 (t, J = 7.8

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				Hz, 1H), 7.12 (d, J = 7.7 Hz, 1H), 7.02 (s, 1H), 6.70 (s, 1H), 6.42 - 6.25 (m, 2H), 5.71 (d, J = 9.8 Hz, 1H), 3.97 (s, 6H), 2.25 (s, 3H).
Example 7	28.9	>10000	>10000	¹ H NMR (400 MHz, CDCl ₃) δ 8.53 (s, 1H), 8.13 (s, 1H), 7.92 (s, 1H), 7.32 (t, J = 8.0 Hz, 1H), 7.13 (d, J = 7.6 Hz, 1H), 6.94 (s, 1H), 6.75 (s, 1H), 6.67 (s, 1H), 6.54 (s, 1H), 6.39 (d, J = 16.7 Hz, 1H), 6.25 - 6.13 (m, 1H), 5.73 (d, J = 10.0 Hz, 1H), 3.96 (s, 6H), 2.22 (s, 3H).
Example 9	138	89	1049	¹ H NMR (400 MHz, CDCl ₃) δ 8.91 (s, 1H), 8.42 (s, 1H), 8.20 (s, 1H), 7.34 (t, J = 7.9 Hz, 1H), 7.20 (s, 1H), 7.12 (d, J = 7.6 Hz, 1H), 6.65 (s, 2H), 6.35 (d, J = 15.3 Hz, 2H), 6.16 (dd, J = 17.2, 9.9 Hz, 1H), 5.68 (d, J = 10.2 Hz, 1H), 3.97 (s, 9H), 2.23 (s, 3H).
Example 10	105.3	3515	4817	¹ H NMR (400 MHz, CDCl ₃) δ 9.16 (s, 1H), 8.38 (d, J = 8.2 Hz, 1H), 8.17 (s, 1H), 7.36 (t, J = 7.9 Hz, 1H), 7.13 (d, J = 7.7 Hz, 1H), 6.98 (s, 1H), 6.64 (s, 1H), 6.56 (s, 1H), 6.37 (d, J = 16.8 Hz, 1H), 6.20 (dd, J = 16.9, 10.2 Hz, 1H), 5.70 (d, J = 10.2 Hz, 1H), 4.07 (s, 3H), 3.96 (s, 6H), 2.25 (s, 3H).
Example 11	100.7	4717	4717	¹ H NMR (400 MHz, CDCl ₃) δ 8.76 (s, 1H), 8.27 (d, J = 8.2 Hz, 1H), 8.14 (s, 1H), 7.49 (s, 1H), 7.33 (t, J = 7.9 Hz, 1H), 7.11 (d, J = 7.6 Hz, 1H), 6.94 (s, 1H), 6.69 (s, 1H), 6.43 - 6.34 (m, 1H), 6.26 (dd, J = 16.9, 10.1 Hz, 1H), 5.72 (dd, J = 10.1, 1.4 Hz, 1H), 3.98 (s, 6H), 3.51 (t, J = 6.4 Hz, 2H), 3.22 (s, 3H), 2.24 (s, 3H), 1.27 (d, J = 7.0 Hz, 2H).
Example 12	43.8	3352	3352	¹ H NMR (400 MHz, CDCl ₃) δ 8.38 (s, 1H), 8.02 (s, 1H), 7.57 (s, 1H), 7.33 (t, J = 7.9 Hz, 1H), 7.14 (d, J = 7.6 Hz, 1H), 6.64 (s, 1H), 6.41 (d, J = 16.6 Hz, 1H), 6.26 (dd, J = 16.6, 9.8 Hz, 1H), 6.07 (s, 1H), 5.74 (d, J

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				= 9.8 Hz, 1H), 3.95 (s, 6H), 2.08 (s, 3H).
Example 13	8.0	181.5	513.1	¹ H NMR (400 MHz, Methanol- <i>d</i> ₄) δ 9.05 (s, 1H), 8.98 (s, 1H), 7.72 (s, 1H), 7.68-7.66 (d, J = 8.0 Hz, 1H), 7.32-7.24 (m, 2H), 6.91 (s, 1H), 6.47 (s, 1H), 6.35-6.23 (m, 2H), 5.66-5.63 (dd, J = 8 Hz, 1H), 3.97 (s, 6H), 2.30 (s, 3H).
Example 64	53.3	54.2	268.5	¹ H NMR (400 MHz, CDCl ₃) δ 10.72 (s, 1H), 9.31 - 9.24 (m, 2H), 8.24 (s, 1H), 7.64 (s, 1H), 6.70 (s, 1H), 6.42 - 6.33 (m, 2H), 5.75 (dd, J = 9.0, 2.6 Hz, 1H), 5.38 - 5.32 (m, 1H), 3.99 (s, 6H), 3.87 (s, 3H).
Example 75	246.6	>10000	>10000	¹ H NMR (400 MHz, CDCl ₃) δ 8.77 (s, 1H), 8.08 (s, 1H), 7.97 (s, 1H), 6.89 (s, 1H), 6.75 (s, 1H), 6.62 (s, 1H), 6.54 - 6.50 (m, 1H), 6.28 (d, J = 16.6 Hz, 1H), 6.13 - 6.03 (m, 1H), 5.70 - 5.57 (m, 1H), 3.91 (s, 6H), 3.49 - 3.38 (m, 4H), 3.25 (s, 3H), 2.54 - 2.39 (m, 4H), 2.08 (s, 3H), 1.31 (s, 3H).
Example 76	24.0	23.8	62.6	¹ H NMR (400 MHz, CDCl ₃) δ 9.03 (s, 1H), 8.17 (s, 1H), 6.77 (s, 1H), 6.63 (s, 1H), 6.38 (d, J = 16.6 Hz, 2H), 6.33 - 6.24 (m, 1H), 5.70 (d, J = 10.1 Hz, 1H), 3.95 (s, 6H), 3.87 (s, 3H), 3.53-3.36 (m, 2H), 1.19-1.10 (m, 1H), 0.55-0.41 (m, 2H), 0.42-0.29 (m, 2H).
Example 118	80.5	179.4	332.8	¹ H NMR (400 MHz, CDCl ₃) δ 9.21 (s, 1H), 9.16 (s, 1H), 7.55 (s, 1H), 6.68 (s, 1H), 6.45 (d, J = 7.3 Hz, 1H), 6.27 (dd, J = 17.0, 1.4 Hz, 1H), 6.07 (dd, J = 16.9, 10.3 Hz, 1H), 5.63 (dd, J = 10.2, 1.4 Hz, 1H), 4.94 (dd, J = 11.6, 4.9 Hz, 1H), 4.90 - 4.79 (m, 1H), 4.34 - 4.19 (m, 2H), 3.98 (s, 6H), 3.90 - 3.79 (m, 2H).
Example 119	166.0	78.1	457.1	¹ H NMR (400 MHz, CDCl ₃) δ 9.17 (s, 2H), 7.56 (s, 1H), 6.68 (s, 1H), 6.25 (dd, J = 16.9, 1.4 Hz, 1H), 6.01 (dd, J = 17.0, 10.3 Hz, 1H), 5.60 (dd, J = 10.3, 1.4 Hz, 1H), 4.55 (s, 1H), 4.35 (s, 1H), 4.11 - 4.00 (m,

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				2H), 3.98 (s, 6H), 3.78 (d, J = 12.1 Hz, 1H), 3.64 (dd, J = 13.4, 10.9 Hz, 1H), 2.24-2.20 (m, 2H), 2.02-1.99 (m, 2H).
Example 120	303.0	155.6	754.2	¹ H NMR (400 MHz, CDCl ₃) δ 9.20 (s, 1H), 9.14 (s, 1H), 7.71 (s, 1H), 6.73 (t, J = 8.0 Hz, 1H), 6.46 (d, J = 7.0 Hz, 1H), 6.26 (d, J = 16.9 Hz, 2H), 6.04 (dd, J = 17.0, 10.3 Hz, 1H), 5.62 (d, J = 10.2 Hz, 1H), 4.93 (d, J = 6.0 Hz, 1H), 4.86 (t, J = 6.2 Hz, 1H), 4.28 (ddd, J = 18.7, 9.4, 6.2 Hz, 2H), 3.93 (s, 6H), 3.85 (ddd, J = 15.4, 9.5, 4.9 Hz, 2H).
Example 121	182.6	171.9	401.4	¹ H NMR (400 MHz, CDCl ₃) δ 9.20 (d, J = 9.2 Hz, 2H), 7.94 (s, 1H), 6.82 (d, J = 2.8 Hz, 1H), 6.60 (d, J = 2.7 Hz, 1H), 6.48 - 6.43 (m, 1H), 6.24 (t, J = 1.9 Hz, 1H), 6.18 (dd, J = 17.3, 10.4 Hz, 1H), 6.12 - 6.04 (m, 1H), 5.63 (dt, J = 10.3, 1.6 Hz, 1H), 4.97 - 4.85 (m, 3H), 4.28 (ddd, J = 20.2, 9.5, 6.3 Hz, 3H), 3.98 (s, 3H), 3.94 (s, 3H).
Example 124	61.6	302.7	556	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.12 (s, 1H), 9.02 (s, 1H), 8.11-8.09 (d, J = 8.0Hz, 1H), 7.67 (s, 1H), 7.03 (s, 1H), 6.92 (s, 1H), 6.70-6.68 (dd, J = 8Hz, 1H), 6.23-6.16 (dd, J = 16Hz, 1H), 6.01-5.97 (dd, J=16Hz, 1H), 5.52-5.48 (dd, J = 16Hz, 1H), 4.75-4.68 (m, 1H), 4.59-4.52 (m, 1H), 4.15-4.11(m, 1H), 4.06-4.03 (m, 1H), 3.98 (s, 6H), 3.74-3.67 (m, 2H).
Example 126	22.0	31.5	53.7	¹ H NMR (400 MHz, CDCl ₃) δ 8.68(s, 1H), 6.82 (s, 1H), 6.76 (s, 1H), 6.62(s, 1H), 6.44 (s, 1H), 6.22-6.18 (dd, J = 16Hz, 1H), 6.00-5.94 (dd, J = 16Hz, 1H), 5.55-5.52(dd, J=8Hz, 1H), 4.92-4.85 (m, 1H), 4.56-4.50 (m, 1H), 4.30-4.21 (m, 2H), 3.98 (s, 6H), 3.87-3.84 (dd, J = 12Hz, 1H), 3.77-3.74 (dd, J = 12Hz, 1H), 4.42-4.39 (m, 2H), 1.23-1.19 (m, 1H), 0.59-0.53 (m, 2H), 0.33-0.29 (m, 2H).

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
Example 421	12.4	5.4	10.0	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.80 (brs, 1H), 6.68 (s, 1H), 6.62 (s, 1H), 6.38 (brs, 1H), 6.21 (dd, J = 17.0, 1.5 Hz, 1H), 6.02 (dd, J = 17.0, 10.2 Hz, 1H), 5.56 (dd, J = 10.1, 1.5 Hz, 1H), 4.93 - 4.83 (m, 2H), 4.30 - 4.18 (m, 2H), 3.96 (s, 6H), 3.90 (dd, J = 9.7, 3.2 Hz, 1H), 3.79 (dd, J = 9.2, 5.2 Hz, 1H), 3.52 - 3.37 (m, 2H), 0.88 (t, J = 6.0 Hz, 1H), 0.57 - 0.48 (m, 2H), 0.36 - 0.26 (m, 2H).
Example 422	22.5	47.0	238.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.74 (s, 1H), 6.67 (s, 2H), 6.22 (d, J = 15.4 Hz, 1H), 5.56 (d, J = 10.4 Hz, 1H), 4.89 (s, 2H), 4.29 - 4.17 (m, 2H), 3.98 (s, 6H), 3.96 - 3.80 (m, 4H), 3.76 - 3.65 (m, 2H), 3.35 (s, 3H).
Example 423	31.7	98.0	347.6	¹ H NMR (400 MHz, CDCl ₃) δ 9.13 (s, 1H), 7.17 (s, 1H), 6.75 (s, 1H), 6.66 (s, 1H), 6.26 (d, J = 16.9 Hz, 1H), 6.12 (dd, J = 17.0, 10.2 Hz, 1H), 5.61 (d, J = 10.2 Hz, 1H), 5.03 - 4.93 (m, 1H), 4.93 - 4.85 (m, 1H), 4.31 - 4.20 (m, 2H), 4.14 (s, 3H), 3.98 (s, 6H), 3.88 (dd, J = 9.8, 4.0 Hz, 1H), 3.82 (dd, J = 9.3, 6.5 Hz, 1H).
Example 424	11.0	18.5	37.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.85 (s, 1H), 6.82 (t, J = 8.0 Hz, 1H), 6.15 (dd, J = 17.1, 2.5 Hz, 1H), 6.12 - 6.02 (m, 1H), 5.53 (dd, J = 9.5, 2.5 Hz, 1H), 4.99 (td, J = 6.3, 4.8 Hz, 1H), 4.87 (q, J = 6.5 Hz, 1H), 4.27 (dd, J = 9.4, 6.3 Hz, 1H), 4.16 (dd, J = 9.0, 6.9 Hz, 1H), 3.92 (m, 7H), 3.80 (dd, J = 9.0, 6.5 Hz, 1H), 3.45 (dd, J = 13.7, 7.0 Hz, 1H), 3.37 (d, J = 6.8 Hz, 1H), 1.33 - 1.24 (m, 1H), 0.55 - 0.41 (m, 2H), 0.42 - 0.29 (m, 2H).
Example 425	7.7	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 8.93 (s, 1H), 6.68 (s, 1H), 6.61 (s, 1H), 6.22 (d, J = 16.6 Hz, 2H), 6.00 (dd, J = 16.9, 10.2 Hz, 1H), 5.89 (s, 1H), 5.57 (d, J = 10.2 Hz, 1H), 4.89 (s, 2H), 4.30 - 4.20 (m, 2H), 3.90 (d, J

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				= 9.8 Hz, 1H), 3.82 - 3.72 (m, 1H), 3.42 (s, 2H), 1.26 - 1.21 (m, 1H), 0.58 - 0.47 (m, 2H), 0.36 - 0.27 (m, 2H).
Example 429	22.4	63.2	184.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.90 (s, 1H), 7.06 (s, 1H), 6.78 (s, 1H), 6.64 (s, 1H), 6.18 (d, J = 17.0 Hz, 1H), 6.11 (d, J = 7.2 Hz, 1H), 5.98 (dd, J = 17.0, 10.2 Hz, 1H), 5.81 (s, 1H), 5.56 (d, J = 10.3 Hz, 1H), 4.94 - 4.83 (m, 2H), 4.27 (dd, J = 9.6, 5.8 Hz, 1H), 4.20 (dd, J = 9.4, 6.1 Hz, 1H), 4.10 (s, 2H), 3.97 (s, 6H), 3.87 (dd, J = 9.6, 3.9 Hz, 1H), 3.81 (dd, J = 9.5, 5.0 Hz, 1H), 3.49 - 3.38 (m, 2H), 3.01 (q, J = 7.5 Hz, 2H), 1.27 (t, J = 7.4 Hz, 3H).
Example 430	19.1	32.0	93.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.60 (s, 1H), 7.24 (s, 1H), 6.81 (brs, 1H), 6.63 (s, 1H), 6.57 (s, 1H), 6.26 (brs, 1H), 6.16 - 6.03 (m, 2H), 5.53 (d, J = 9.6 Hz, 1H), 5.08 (s, 1H), 4.88 (s, 1H), 4.26 - 4.12 (m, 2H), 3.96 (d, J = 6.0 Hz, 6H), 3.93 - 3.88 (m, 2H), 3.81 (d, J = 10.1 Hz, 2H), 3.60 (d, J = 11.3 Hz, 2H), 3.42 (s, 1H), 2.94 (s, 3H).
Example 431	519.1	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 12.21 (s, 1H), 8.90 (s, 1H), 7.72 (s, 1H), 7.52 (s, 1H), 6.78 (s, 1H), 6.67 (s, 1H), 6.22 - 6.12 (m, 2H), 5.53 (d, J = 9.9 Hz, 1H), 5.00 - 4.88 (m, 2H), 4.33 - 4.27 (m, 2H), 4.26 - 4.19 (m, 2H), 3.97 (s, 6H), 3.94 - 3.87 (m, 2H), 3.49 (s, 2H), 2.87 (s, 6H).
Example 432	105.8	395.8	934.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.66 (s, 1H), 6.99 (s, 1H), 6.91 (s, 1H), 6.63 (s, 1H), 6.59 (s, 1H), 6.33 (d, J = 6.4 Hz, 1H), 6.18 (d, J = 5.9 Hz, 2H), 5.59 - 5.49 (m, 1H), 4.99 (s, 2H), 4.30 - 4.23 (m, 2H), 3.97 (s, 6H), 3.92 - 3.87 (m, 1H), 3.77 - 3.71 (m, 1H), 3.25 (s, 2H), 2.77 (s, 6H), 2.25 (s, 2H), 1.70 (s, 2H).
Example 433	7.6	19.3	26.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.88 (s, 1H), 6.67 (s, 1H), 6.62 (s, 1H), 6.30 (brs, 1H), 6.19 (d, J = 16.9 Hz, 1H), 5.97 (dd, J =

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				17.0, 10.3 Hz, 1H), 5.88 (brs, 1H), 5.54 (d, J = 10.4 Hz, 1H), 4.95 - 4.83 (m, 2H), 4.29 - 4.18 (m, 2H), 3.95 (s, 6H), 3.89 (dd, J = 9.7, 3.0 Hz, 1H), 3.77 (dd, J = 9.3, 5.5 Hz, 1H), 3.14 (d, J = 4.4 Hz, 3H).
Example 434	8.4	9.6	18.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.87 (s, 1H), 6.67 (s, 1H), 6.62 (s, 1H), 6.51 (s, 1H), 6.27 - 6.15 (m, 2H), 5.97 (dd, J = 16.9, 10.3 Hz, 1H), 5.79 (s, 1H), 5.56 (dd, J = 10.3, 1.4 Hz, 1H), 4.97 - 4.82 (m, 2H), 4.29 - 4.18 (m, 2H), 3.96 (s, 6H), 3.89 (dd, J = 9.6, 3.2 Hz, 1H), 3.76 (dd, J = 9.2, 5.5 Hz, 1H), 3.65 - 3.50 (m, 2H), 1.32 (t, J = 7.2 Hz, 3H).
Example 435	67.0	80	148.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.96 (s, 1H), 6.85 (s, 1H), 6.79 (s, 1H), 6.64 (s, 1H), 6.19 (d, J = 16.9 Hz, 1H), 6.13 (s, 1H), 6.06 (s, 1H), 5.98 (dd, J = 17.0, 10.3 Hz, 1H), 5.56 (d, J = 10.2 Hz, 1H), 4.93 (dd, J = 11.7, 4.8 Hz, 2H), 4.51 - 4.37 (m, 1H), 4.30 - 4.17 (m, 3H), 3.97 (s, 6H), 3.90 (dd, J = 9.7, 2.8 Hz, 1H), 3.78 (dd, J = 9.4, 5.0 Hz, 1H).
Example 436	>10000	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 9.33 (s, 1H), 6.59 (s, 1H), 6.56 (s, 1H), 6.45 (s, 1H), 6.22 (s, 1H), 6.14 - 6.00 (m, 1H), 5.62 (d, J = 10.2 Hz, 1H), 4.86 (s, 2H), 4.24 (s, 2H), 3.98 - 3.94 (m, 2H), 3.83 (s, 6H), 3.38 (s, 6H).
Example 438A	3.5	9.9	17.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.84 (brs, 1H), 6.70 (s, 1H), 6.63 (s, 1H), 6.42 (brs, 1H), 6.22 (dd, J = 17.0, 1.5 Hz, 1H), 6.11 - 6.01 (m, 1H), 5.85 (s, 1H), 5.57 (dd, J = 10.1, 1.5 Hz, 1H), 4.93 - 4.81 (m, 2H), 4.30 - 4.16 (m, 3H), 3.96 (s, 6H), 3.92 - 3.82 (m, 3H), 3.82 - 3.74 (m, 2H), 3.58 - 3.45 (m, 1H), 1.95 - 1.85 (m, 2H), 1.70 - 1.64 (m, 2H).
Example 438B	9.1	15.8	35.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.74 (s, 1H), 6.69 (s, 1H), 6.62 (s, 1H),

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				6.25 (dd, J = 16.9, 1.4 Hz, 1H), 6.20 (s, 1H), 6.03 (dd, J = 16.9, 10.3 Hz, 1H), 5.81 (s, 1H), 5.61 (d, J = 10.4 Hz, 1H), 4.93 - 4.81 (m, 2H), 4.31 - 4.16 (m, 3H), 3.96 (s, 6H), 3.93 - 3.84 (m, 2H), 3.82 - 3.72 (m, 3H), 3.59 (dd, J = 13.3, 6.7 Hz, 1H), 1.91 - 1.85 (m, 2H), 1.70 - 1.63 (m, 2H).
Example 439	4.6	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 8.88 (s, 1H), 6.72 (brs, 1H), 6.69 (s, 1H), 6.63 (s, 1H), 6.35 (brs, 1H), 6.19 (d, J = 17.1 Hz, 1H), 6.00 (dd, J = 17.0, 10.2 Hz, 1H), 5.89 (s, 1H), 5.55 (dd, J = 10.3, 1.5 Hz, 1H), 4.96 - 4.82 (m, 2H), 4.31 - 4.17 (m, 2H), 3.96 (s, 6H), 3.93 - 3.82 (m, 2H), 3.81 - 3.71 (m, 2H), 3.66 (dd, J = 8.9, 5.3 Hz, 1H), 3.63 - 3.51 (m, 2H), 2.82 - 2.69 (m, 1H), 2.13 - 2.00 (m, 1H), 1.77 - 1.72 (m, 1H).
Example 443	8.2	15.9	70.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.73 (s, 1H), 6.63 (s, 1H), 6.24 - 6.16 (m, 1H), 6.13 (brs, 1H), 5.96 (dd, J = 17.2, 10.3 Hz, 1H), 5.79 (brs, 1H), 5.59 - 5.51 (m, 1H), 4.95 - 4.84 (m, 2H), 4.71 (s, 1H), 4.30 - 4.15 (m, 2H), 4.13 - 4.04 (m, 1H), 4.05 - 3.98 (m, 1H), 3.97 (s, 6H), 3.92 - 3.82 (m, 3H), 3.76 (dd, J = 9.3, 5.7 Hz, 1H), 2.45 - 2.31 (m, 1H), 2.18 - 2.05 (m, 1H).
Example 444	7.0	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.70 (s, 1H), 6.63 (s, 1H), 6.42 - 6.33 (m, 1H), 6.20 (dd, J = 17.0, 1.4 Hz, 1H), 6.17 - 6.13 (m, 1H), 5.97 (dd, J = 17.0, 10.3 Hz, 1H), 5.75 (d, J = 6.5 Hz, 1H), 5.57 (dd, J = 10.4, 1.4 Hz, 1H), 4.95 - 4.82 (m, 2H), 4.29 - 4.18 (m, 3H), 4.03 - 3.98 (m, 2H), 3.97 (s, 6H), 3.89 (dd, J = 9.6, 3.6 Hz, 1H), 3.76 (dd, J = 9.3, 5.8 Hz, 1H), 3.58 - 3.48 (m, 2H), 2.24 - 2.10 (m, 2H), 1.73 - 1.64 (m, 2H).
Example 452	42.2	20.3	53.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.88 (brs, 1H), 6.77 (s, 1H), 6.64 (s, 1H), 6.17 (d, J = 16.8 Hz, 1H), 5.95 (m, 2H),

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				5.55 (d, J = 10.2 Hz, 1H), 5.02 - 4.94 (m, 1H), 4.90 (s, 1H), 4.46 (s, 1H), 4.31 - 4.18 (m, 2H), 3.97 (s, 6H), 3.91 (d, J = 10.8 Hz, 1H), 3.73 (t, J = 8.1 Hz, 1H), 3.11 (s, 2H), 2.80 (s, 1H), 2.69 (s, 1H).
Example 454	66.9	40.8	89.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.88 (s, 1H), 6.67 (s, 1H), 6.63 (s, 1H), 6.21 (d, J = 17.0 Hz, 1H), 6.09 - 5.97 (m, 1H), 5.57 (d, J = 10.2 Hz, 1H), 4.89 - 4.81 (m, 2H), 4.27 - 4.20 (m, 2H), 3.96 (s, 6H), 3.92 - 3.90 (m, 1H), 3.81 - 3.77 (m, 1H), 3.59 - 3.48 (m, 2H), 2.30 (dd, J = 15.2, 7.9 Hz, 1H), 1.83 - 1.76 (m, 2H), 1.59 (brd, J = 31.7 Hz, 4H), 1.28 (brs, 3H).
Example 456	82.4	163	299	¹ H NMR (400 MHz, CDCl ₃) δ 8.87 (s, 1H), 7.47 - 7.11 (m, 6H), 6.70 (s, 1H), 6.63 (s, 1H), 6.22 (d, J = 17.1 Hz, 2H), 6.09 - 5.75 (m, 2H), 5.58 (d, J = 10.3 Hz, 1H), 4.89 - 4.60 (m, 2H), 4.22 - 4.15 (m, 2H), 3.96 (s, 6H), 3.83-3.75 (m, 3H), 3.03 (q, J = 6.8 Hz, 2H).
Example 459	17.3	23.8	58.0	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.02 (s, 1H), 7.58 (s, 1H), 7.38 (s, 1H), 6.98 (s, 1H), 6.71 (s, 1H), 6.14 (dd, J = 17.1, 10.2 Hz, 1H), 5.94 (dd, J = 17.2, 2.2 Hz, 1H), 5.47 (dd, J = 10.4, 2.0 Hz, 1H), 4.86 (s, 1H), 4.67 (q, J = 6.1 Hz, 1H), 4.39 (td, J = 14.6, 13.6, 5.3 Hz, 2H), 4.12 (t, J = 7.9 Hz, 1H), 4.02 (dd, J = 8.8, 6.5 Hz, 1H), 3.97 (s, 6H), 3.81-3.68 (m, 4H), 3.66 (dd, J = 8.8, 5.9 Hz, 1H), 3.33 (s, 3H).
Example 462	14.5	24.6	38.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.66 (s, 1H), 6.64 (s, 1H), 6.62 (s, 1H), 6.30 (s, 1H), 6.23 (dd, J = 16.9, 1.4 Hz, 1H), 6.01 (dd, J = 16.9, 10.3 Hz, 1H), 5.84 (s, 1H), 5.59 (dd, J = 10.3, 1.4 Hz, 1H), 4.93 - 4.77 (m, 2H), 4.29 - 4.19 (m, 2H), 3.96 (s, 6H), 3.91 (dd, J = 9.5, 4.0 Hz, 1H), 3.79 (dd, J = 9.3, 5.5 Hz, 1H), 3.51 (dd, J = 13.1, 6.5 Hz, 1H), 3.36 (dd, J = 13.2, 5.9 Hz, 1H), 0.99 (s, 9H).

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
Example 465	46.4	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 8.87 (s, 1H), 6.67 (s, 1H), 6.62 (s, 1H), 6.18 (d, J = 17.1 Hz, 1H), 6.06 (d, J = 10.1 Hz, 1H), 5.86 (s, 1H), 5.54 (d, J = 10.1 Hz, 1H), 4.96 - 4.83(m, 2H), 4.30 - 4.19 (m, 2H), 3.96 (s, 6H), 3.89 (dd, J = 9.6, 3.7 Hz, 1H), 3.81 (dd, J = 9.2, 5.6 Hz, 1H), 3.59 (s, 2H), 2.76 (s, 2H), 2.53 (s, 6H), 1.26 (s, 4H).
Example 475	62.0	131.9	677.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 7.03 (s, 1H), 6.83 (d, J = 2.8 Hz, 1H), 6.54 (d, J = 2.8 Hz, 1H), 6.30 (s, 1H), 6.22 (dd, J = 16.8, 1.4 Hz, 1H), 6.00 (dd, J = 16.9, 10.3 Hz, 1H), 5.82 (s, 1H), 5.61 - 5.53 (m, 1H), 4.88 (q, J = 6.5, 6.0 Hz, 2H), 4.33 - 4.18 (m, 2H), 3.92 (s, 3H), 3.89 (s, 1H), 3.84 (s, 3H), 3.81 - 3.73 (m, 1H), 3.58 - 3.40 (m, 2H), 1.26 (s, 1H), 0.66 - 0.46 (m, 2H), 0.42 - 0.24 (m, 2H).
Example 498	8797	2393	>10000	¹ H NMR (400 MHz, CDCl ₃) δ 9.18 (s, 1H), 7.71 (d, J = 4.2 Hz, 1H), 7.53 (d, J = 8.4 Hz, 1H), 6.93 (s, 1H), 6.67 (s, 1H), 6.17 (d, J = 16.9 Hz, 1H), 6.06 (d, J = 11.2 Hz, 1H), 5.55 (d, J = 10.1 Hz, 1H), 4.95 (s, 1H), 4.63 (s, 1H), 4.39 (s, 2H), 4.31 (t, J = 6.7 Hz, 1H), 4.24 (s, 2H), 3.98 (s, 6H), 3.92 (s, 1H), 0.88 (t, J = 7.0 Hz, 1H), 0.76 - 0.65 (m, 2H), 0.47 (s, 2H).
Example 501	15.7	17.8	49.9	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.68 (s, 1H), 6.64 (s, 1H), 6.17 (d, J = 16.7 Hz, 1H), 5.96 (d, J = 12.1 Hz, 1H), 5.55 (d, J = 10.5 Hz, 1H), 4.75-4.60 (m, 1H), 4.40-4.25 (m, 1H), 4.10-4.02 (m, 1H), 4.01-3.90 (m, 3H), 3.96 (s, 6H), 3.77 (d, J = 11.9 Hz, 1H), 3.62 (t, J = 12.2 Hz, 2H), 1.98 - 1.86 (m, 2H), 0.88 (s, 1H), 0.55 (s, 2H), 0.35 (s, 2H).
Example 510	29	75.2	115.9	¹ H NMR (400 MHz, CDCl ₃) δ 8.87 (s, 1H), 6.67 (s, 1H), 6.62 (s, 1H), 6.48 (s, 1H), 6.20 (dd, J = 16.9, 1.4 Hz, 2H), 6.05 (d, J = 8.5 Hz, 1H), 5.91 (dd, J = 16.9, 10.3 Hz,

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				1H), 5.54 (dd, J = 10.4, 1.4 Hz, 1H), 4.70 (d, J = 8.7 Hz, 1H), 4.38 - 4.27 (m, 1H), 4.12 - 4.05 (m, 1H), 3.99 (d, J = 11.2 Hz, 1H), 3.96 (s, 6H), 3.75 (dd, J = 12.0, 1.7 Hz, 1H), 3.65 - 3.57 (m, 1H), 3.09 (d, J = 5.0 Hz, 3H), 1.97 - 1.86 (m, 2H).
Example 606	23.4	25.7	41.1	¹ H NMR (400 MHz, CDCl ₃) δ 9.18 (s, 1H), 7.44 (s, 1H), 6.67 (s, 1H), 6.30 - 6.24 (m, 1H), 6.17 - 6.08 (m, 1H), 5.63 (d, J = 10.7 Hz, 1H), 4.93 (s, 2H), 4.35 - 4.30 (m, 1H), 4.25 (dd, J = 9.4, 6.1 Hz, 1H), 3.97 (s, 6H), 3.92 (dd, J = 9.6, 3.1 Hz, 1H), 3.87 - 3.82 (m, 1H), 2.98 (s, 3H).
Example 609	NT	36.6	56.1	¹ H NMR (400 MHz, CDCl ₃) δ 9.07 (s, 1H), 6.64 (s, 1H), 6.55 (s, 1H), 6.25 (d, J = 16.8 Hz, 1H), 6.05 (dd, J = 16.9, 10.3 Hz, 1H), 5.86 (brs, 1H), 5.62 (dd, J = 10.3, 1.4 Hz, 1H), 4.90-4.87 (m, 2H), 4.82-4.78 (m, 1H), 4.31 (td, J = 10.3, 6.2 Hz, 2H), 3.97 (s, 6H), 3.95 - 3.68 (m, 2H), 3.28-3.24 (m, 1H), 1.27 - 1.21 (m, 2H), 1.10 (dd, J = 8.3, 3.0 Hz, 2H).
Example 629	2.9	2.6	4.0	¹ H NMR (400 MHz, CDCl ₃) δ 9.10 (s, 1H), 8.53 (s, 1H), 8.49 (s, 1H), 7.37 (s, 1H), 6.68 (s, 1H), 6.43 (d, J = 6.2 Hz, 1H), 6.20 (d, J = 16.9 Hz, 1H), 6.09 (s, 1H), 5.98 (dd, J = 17.0, 10.2 Hz, 1H), 5.57 (d, J = 10.3 Hz, 1H), 4.97 - 4.90 (m, 2H), 4.33 - 4.23 (m, 2H), 3.98 (s, 9H), 3.94 (dd, J = 9.8, 3.2 Hz, 1H), 3.82 (dd, J = 9.5, 4.8 Hz, 1H).
Example 637	4.2	10.1	17.3	¹ H NMR (400 MHz, Methanol- <i>d</i> ₄) δ 9.03 (s, 1H), 8.56 (s, 1H), 8.40 (s, 1H), 7.49 (s, 1H), 6.74 (t, J = 8.0 Hz, 1H), 6.03 (d, J = 5.9 Hz, 2H), 5.46 (t, J = 5.9 Hz, 1H), 4.87 (s, 2H), 4.20 (t, J = 7.8 Hz, 1H), 4.12 (dd, J = 9.1, 6.6 Hz, 1H), 3.93 (s, 3H), 3.86 (s, 6H), 3.83 (d, J = 4.0 Hz, 1H), 3.74 (dd, J = 9.2, 5.4 Hz, 1H).
Example 655	8.0	28.0	68.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.88 (s, 1H), 6.74 (s, 1H), 6.70 (s, 1H), 6.62 (s, 1H),

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				6.19 (dd, J = 16.9, 1.4 Hz, 2H), 5.95 (dd, J = 16.9, 10.3 Hz, 1H), 5.79 (s, 1H), 5.55 (dd, J = 10.3, 1.4 Hz, 1H), 4.93 - 4.81 (m, 2H), 4.29 - 4.17 (m, 2H), 3.96 (s, 6H), 3.88 (dd, J = 9.7, 3.3 Hz, 1H), 3.75 (dd, J = 9.2, 5.9 Hz, 1H), 3.01 - 2.84 (m, 1H), 0.87 - 0.79 (m, 2H), 0.76 - 0.69 (m, 1H), 0.68 - 0.58 (m, 1H).
Example 657	6.5	9.6	13.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.87 (s, 1H), 6.69 (s, 1H), 6.62 (s, 1H), 6.26 (dd, J = 16.8, 1.3 Hz, 2H), 6.06 (dd, J = 17.0, 10.3 Hz, 1H), 5.70 - 5.60 (m, 2H), 4.80-4.70 (m, 3H), 4.60-4.45 (m, 3H), 4.29 - 4.15 (m, 2H), 3.95 (s, 6H), 3.87 - 3.73 (m, 2H), 2.42 (q, J = 7.6 Hz, 2H).
Example 658	4.8	11.4	15.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.73 (s, 1H), 6.63 (s, 1H), 6.24 (d, J = 16.8 Hz, 1H), 6.13 (d, J = 10.9 Hz, 1H), 5.61 (d, J = 10.1 Hz, 1H), 4.84 - 4.57 (m, 4H), 4.54 - 4.07 (m, 5H), 3.95 (s, 6H), 3.89 - 3.72 (m, 2H), 3.30 (s, 3H).
Example 659	6.6	18.0	24.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.90 (s, 1H), 6.93 (s, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.34 (brs, 1H), 6.24 (dd, J = 16.8, 1.4 Hz, 1H), 6.05 (dd, J = 16.9, 10.3 Hz, 1H), 5.74 (brs, 1H), 5.62 (dd, J = 10.2, 1.4 Hz, 1H), 4.88 - 4.61 (m, 4H), 4.45 - 4.37 (m, 3H), 4.28 - 4.15 (m, 2H), 3.91 (s, 6H), 3.84 (dd, J = 9.5, 3.6 Hz, 1H), 3.77 (dd, J = 9.5, 5.1 Hz, 1H), 3.33 (s, 3H).
Example 663	4.5	2.3	3.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.95 (s, 1H), 6.88 (s, 1H), 6.64 (s, 1H), 6.25 (d, J = 16.9 Hz, 2H), 6.06 (dd, J = 17.0, 10.3 Hz, 1H), 5.93 (s, 1H), 5.63 (d, J = 10.3 Hz, 1H), 4.88 - 4.77 (m, 6H), 4.22 (dd, J = 9.5, 6.2 Hz, 2H), 3.96 (s, 6H), 3.82 (ddd, J = 17.2, 9.6, 5.0 Hz, 2H).
Example 664	4.9	3.7	4.9	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.04 (s, 1H), 8.06 (d, J = 8.1 Hz, 1H), 7.23 (brs, 1H), 6.97 (s, 1H), 6.77 (s, 1H), 6.23 (dd, J

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				= 17.1, 10.2 Hz, 1H), 6.02 (dd, J = 17.1, 2.1 Hz, 1H), 5.57 (d, J = 10.7 Hz, 1H), 5.55 (s, 1H), 4.74 - 4.58 (m, 2H), 4.21 (brs, 4H), 4.12 - 4.01 (m, 2H), 3.95 (s, 6H), 3.78 - 3.66 (m, 2H), 1.46 (s, 3H).
Example 665	5.7	9.9	11.8	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.06 (s, 1H), 8.06 (d, J = 8.3 Hz, 1H), 7.28 (s, 1H), 6.97 (s, 1H), 6.81 (s, 1H), 6.21 (dd, J = 17.1, 10.2 Hz, 1H), 6.00 (dd, J = 17.1, 2.2 Hz, 1H), 5.54 (dd, J = 10.1, 2.2 Hz, 1H), 4.76 - 4.60 (m, 2H), 4.17 (br s, 4H), 4.09 (d, J = 7.4 Hz, 1H), 4.05 - 4.00 (m, 1H), 3.95 (s, 6H), 3.75 (dd, J = 8.7, 5.9 Hz, 1H), 3.67 (dd, J = 9.0, 5.2 Hz, 1H), 3.21 (s, 3H), 1.49 (s, 3H).
Example 666	4.5	3.2	4.6	¹ H NMR (400 MHz, CDCl ₃) δ 8.90 (s, 1H), 6.93 (s, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.37 (s, 1H), 6.24 (d, J = 16.9 Hz, 1H), 6.04 (dd, J = 17.0, 10.3 Hz, 1H), 5.78 (s, 1H), 5.61 (d, J = 10.3 Hz, 1H), 4.84 - 4.73 (m, 2H), 4.56 - 4.41 (m, 2H), 4.41 - 4.26 (m, 2H), 4.26 - 4.17 (m, 2H), 3.91 (s, 6H), 3.85 (dd, J = 9.6, 3.5 Hz, 1H), 3.76 (dd, J = 9.4, 5.1 Hz, 1H), 3.30 (s, 3H), 1.56 (s, 3H).
Example 668	14.7	16	22.6	¹ H NMR (400 MHz, CDCl ₃) δ 8.82 (s, 1H), 6.70 (s, 1H), 6.66 (s, 1H), 6.56 (s, 1H), 6.16 (d, J = 17.0 Hz, 1H), 6.01 (dd, J = 17.1, 10.2 Hz, 1H), 5.53 (d, J = 10.0 Hz, 1H), 4.80-4.55(m, 8H), 4.18-4.15 (m, 2H), 3.88 (s, 6H), 3.84 - 3.70 (m, 4H).
Example 675	11.0	6.2	8.7	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.71 (s, 1H), 6.63 (s, 1H), 6.38 (brs, 1H), 6.24 (d, J = 16.9 Hz, 1H), 6.05 (dd, J = 16.9, 10.2 Hz, 1H), 5.78 (brs, 1H), 5.61 (dd, J = 10.2, 1.3 Hz, 1H), 4.87 - 4.75 (m, 2H), 4.42 - 4.15 (m, 6H), 3.95 (s, 6H), 3.86 (dd, J = 9.5, 3.9 Hz, 1H), 3.78 (dd, J = 9.4, 5.4 Hz, 1H), 3.72 - 3.60 (m, 4H), 1.86 (t, J = 5.2 Hz, 4H).

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
Example 677	7.4	3.7	5.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.80 (s, 1H), 6.63 (s, 1H), 6.54 (s, 1H), 6.41 (brs, 1H), 6.21 (d, <i>J</i> = 17.0 Hz, 1H), 5.99 (dd, <i>J</i> = 16.6, 10.2 Hz, 1H), 5.71 (brs, 1H), 5.53 (dd, <i>J</i> = 10.3, 3.5 Hz, 1H), 4.85-4.70 (m, 2H), 4.28-3.96 (m, 7H), 3.88 (s, 6H), 3.90-3.82 (m, 1H), 3.80-3.72 (m, 1H), 3.29 (d, <i>J</i> = 2.8 Hz, 3H), 2.07-1.94 (m, 2H).
Example 678	6.5	13.3	16.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.94 (s, 1H), 6.56 (s, 1H), 6.50 (d, <i>J</i> = 7.4 Hz, 1H), 6.18 (d, <i>J</i> = 16.9 Hz, 1H), 5.98 (dd, <i>J</i> = 16.9, 10.3 Hz, 2H), 5.54 (d, <i>J</i> = 9.7 Hz, 1H), 4.82 - 4.69 (m, 2H), 4.74 - 4.64 (m, 2H), 4.19 - 4.12 (m, 2H), 3.89 (s, 6H), 3.87 - 3.76 (m, 7H), 3.76 - 3.67 (m, 1H), 2.20 - 2.10 (m, 1H), 2.08 - 1.95 (m, 1H).
Example 686	8.2	5.4	8.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.94 (s, 1H), 6.56 (s, 1H), 6.50 (d, <i>J</i> = 7.4 Hz, 1H), 6.18 (d, <i>J</i> = 16.9 Hz, 1H), 5.98 (dd, <i>J</i> = 16.9, 10.3 Hz, 1H), 5.54 (d, <i>J</i> = 10.2 Hz, 1H), 4.81 - 4.73 (m, 1H), 4.70 - 4.67 (m, 1H), 4.20 - 4.12 (m, 2H), 3.89 (s, 6H), 3.88 - 3.67 (m, 10H).
Example 687	6.3	6.9	10.0	¹ H NMR (400 MHz, CDCl ₃) δ 9.03 (s, 1H), 7.25 (s, 1H), 6.70 (t, <i>J</i> = 7.9 Hz, 1H), 6.47 (brs, 1H), 6.25 (d, <i>J</i> = 16.9 Hz, 1H), 6.06 (dd, <i>J</i> = 16.9, 10.2 Hz, 1H), 5.62 (d, <i>J</i> = 10.3 Hz, 1H), 4.88 - 4.81 (m, 1H), 4.78 (brs, 1H), 4.23 (dd, <i>J</i> = 9.5, 6.2 Hz, 2H), 3.93 (s, 6H), 3.92 - 3.81 (m, 9H), 3.81 - 3.77 (m, 1H).
Example 690	8.0	8.4	12.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.98 (s, 1H), 6.99 (s, 1H), 6.64 (s, 1H), 6.55 (d, <i>J</i> = 7.1 Hz, 1H), 6.26 (dd, <i>J</i> = 17.1, 1.3 Hz, 1H), 6.11 - 6.04 (m, 1H), 6.04 (s, 1H), 5.63 (d, <i>J</i> = 10.2 Hz, 1H), 4.86 - 4.73 (m, 2H), 4.68 (d, <i>J</i> = 12.8 Hz, 1H), 4.61 (d, <i>J</i> = 12.8 Hz, 1H), 4.29 - 4.20 (m, 2H), 3.97 (s, 6H), 3.94 - 3.85 (m, 3H), 3.82 (dd, <i>J</i> = 9.4, 5.0 Hz, 1H), 2.79 - 2.68 (m, 2H), 1.23 (dd, <i>J</i> = 6.3,

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				2.2 Hz, 6H).
Example 693	5.6	1.3	2.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.99 (s, 1H), 6.99 (s, 1H), 6.63 (s, 1H), 6.53 (s, 1H), 6.23 (d, J = 16.9 Hz, 1H), 6.06 (dd, J = 17.0, 10.2 Hz, 1H), 5.94 (s, 1H), 5.60 (d, J = 10.2 Hz, 1H), 4.87 - 4.77 (m, 2H), 4.39 (d, J = 8.3 Hz, 1H), 4.30 (d, J = 13.2 Hz, 1H), 4.27 - 4.20 (m, 2H), 3.96 (s, 6H), 3.88 (dd, J = 9.5, 3.5 Hz, 1H), 3.81 (dd, J = 14.4, 5.1 Hz, 1H), 3.51 - 3.39 (m, 3H), 3.38 (s, 3H), 2.15 - 2.01 (m, 2H), 1.79 - 1.74 (m, 2H).
Example 694	4.8	10.0	12.0	¹ H NMR (400 MHz, CDCl ₃) δ 9.01 (s, 1H), 6.70 (t, J = 7.9 Hz, 1H), 6.22 (d, J = 16.8 Hz, 1H), 6.07 (dd, J = 17.0, 10.2 Hz, 1H), 5.59 (d, J = 10.2 Hz, 1H), 4.86 - 4.80 (m, 2H), 4.38 - 4.21 (m, 4H), 3.93 (s, 6H), 3.91 - 3.82 (m, 2H), 3.52 - 3.43 (m, 3H), 3.39 (s, 3H), 2.21 - 2.11 (m, 2H).
Example 704	8.5	17.2	20.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.76 (s, 1H), 6.65 (brs, 1H), 6.62 (s, 1H), 6.25 (d, J = 16.9 Hz, 1H), 6.05 (dd, J = 16.8, 10.4 Hz, 1H), 5.61 (d, J = 10.3 Hz, 1H), 4.69 (brs, 1H), 4.55 (brs, 1H), 4.46 - 4.23 (m, 5H), 4.10 - 3.99 (m, 2H), 3.95 (s, 6H), 3.74 (d, J = 11.9 Hz, 1H), 3.61 (t, J = 11.8 Hz, 1H), 3.32 (s, 3H), 2.10 - 1.98 (m, 1H), 1.90 - 1.80 (m, 1H).
Example 715	62.7	1917.5	1917.5	¹ H NMR (400 MHz, CDCl ₃) δ 10.71 (s, 1H), 9.63 (s, 1H), 9.60 (s, 1H), 8.61 (s, 1H), 7.80 (s, 1H), 6.72 (s, 1H), 6.49 (d, J = 16.7 Hz, 1H), 6.40 (dd, J = 16.9, 9.7 Hz, 1H), 5.84 (d, J = 9.7 Hz, 1H), 5.21 (t, J = 6.0 Hz, 2H), 4.00 (s, 7H), 3.89 (t, J = 5.9 Hz, 2H), 3.35 (s, 3H).
Example 723	78.5	338.1	1211.5	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.75 (s, 1H), 9.71 (s, 1H), 9.68 (s, 1H), 8.37 (s, 1H), 8.21 (s, 1H), 8.05 (s, 1H), 7.15 (s, 1H), 6.51 (dd, J = 17.0, 10.2 Hz, 1H), 6.26 (dd, J = 17.1, 2.2 Hz, 1H), 5.75 (dd, J =

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				10.1, 2.1 Hz, 1H), 4.56 (t, <i>J</i> = 5.6 Hz, 2H), 4.06 (s, 6H), 3.70 (t, <i>J</i> = 5.6 Hz, 2H), 3.14 (s, 3H).
Example 730	115.0	164	194	¹ H NMR (400 MHz, CDCl ₃) δ 8.75 (s, 1H), 7.68 (s, 1H), 7.40 (s, 1H), 6.65 (s, 1H), 6.38 - 6.32 (m, 1H), 6.27 - 6.21 (m, 1H), 5.73 - 5.69 (m, 1H), 5.30 (s, 1H), 4.81 (d, <i>J</i> = 5.6 Hz, 2H), 3.96 (s, 6H), 3.78 (s, 3H).
Example 752	990.9	1712	2631	¹ H NMR (400 MHz, CDCl ₃) δ 8.82 (s, 1H), 6.68 (s, 1H), 6.66 (s, 1H), 6.59 (s, 1H), 6.13 (dd, <i>J</i> = 16.8, 1.4 Hz, 1H), 5.94 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 5.51 (dd, <i>J</i> = 10.3, 1.4 Hz, 1H), 5.30 (s, 1H), 4.96 - 4.82 (m, 1H), 4.42 (s, 1H), 4.26 (dd, <i>J</i> = 9.8, 6.0 Hz, 1H), 4.18 (dd, <i>J</i> = 9.3, 7.4 Hz, 1H), 3.92 (s, 6H), 3.88 - 3.81 (m, 1H), 3.74 (dd, <i>J</i> = 9.2, 6.9 Hz, 1H), 3.13 (d, <i>J</i> = 4.6 Hz, 3H).
Example 753	19	130	212	¹ H NMR (400 MHz, Methanol- <i>d</i> ₄) δ 9.22 (s, 1H), 7.65 (s, 1H), 6.94 (s, 1H), 6.19 (dd, <i>J</i> = 17.1, 9.9 Hz, 1H), 6.10 (dd, <i>J</i> = 17.1, 2.2 Hz, 1H), 5.56 (dd, <i>J</i> = 9.9, 2.1 Hz, 1H), 5.01 (s, 2H), 4.29 (s, 1H), 4.17 (dd, <i>J</i> = 9.3, 6.2 Hz, 1H), 3.98 (s, 6H), 3.93 - 3.85 (m, 1H), 3.85 - 3.79 (m, 1H).
Example 754	13.6	18.0	54.0	¹ H NMR (400 MHz, CDCl ₃) δ 9.06 (d, <i>J</i> = 6.5 Hz, 1H), 7.11 (s, 1H), 6.64 (s, 1H), 6.25 (dd, <i>J</i> = 16.9, 1.5 Hz, 1H), 6.05 (dd, <i>J</i> = 17.0, 10.4 Hz, 1H), 5.60 (dd, <i>J</i> = 10.3, 1.5 Hz, 1H), 4.91 (brs, 1H), 4.79 (t, <i>J</i> = 6.5, 1H), 4.59 (q, <i>J</i> = 7.0, 2H), 4.35-4.19 (m, 2H), 3.93-3.68 (m, 2H) 1.48 (t, <i>J</i> = 7.1Hz, 3H).
Example 755	43.0	NT	NT	¹ H NMR (400 MHz, CDCl ₃) δ 8.84 (s, 1H), 6.84 (s, 1H), 6.67 (s, 1H), 6.58 - 6.49 (m, 3H), 6.30 (d, <i>J</i> = 8.5 Hz, 2H), 6.10 (d, <i>J</i> = 17.0 Hz, 2H), 6.04 - 5.83 (m, 2H), 5.48 (d, <i>J</i> = 10.1 Hz, 1H), 4.72 (s, 3H), 4.63 (d, <i>J</i> = 14.8 Hz, 1H), 4.17 - 4.04 (m, 2H), 3.89 (s, 6H), 3.78 (d, <i>J</i> = 6.2 Hz, 1H), 3.71 (s, 2H), 3.68 (s, 7H).

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
Example 756	20.2	19	31	¹ H NMR (400 MHz, CDCl ₃) δ 9.10 (s, 1H), 7.43 (s, 1H), 7.13 (s, 1H), 6.66 (s, 1H), 6.36 (s, 1H), 6.26 (dd, J = 16.9, 1.4 Hz, 1H), 6.04 (dd, J = 16.9, 10.3 Hz, 1H), 5.94 (s, 1H), 5.63 (dd, J = 10.3, 1.4 Hz, 1H), 4.89 - 4.78 (m, 2H), 4.45 (d, J = 2.8 Hz, 2H), 4.31 - 4.21 (m, 2H), 3.97 (s, 6H), 3.95 - 3.92 (m, 2H), 3.90 - 3.87 (m, 1H), 3.81 (dd, J = 9.5, 4.8 Hz, 1H), 2.94 - 2.77 (m, 2H).
Example 758	5.9	5.8	9.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (s, 1H), 6.71 (s, 1H), 6.62 (s, 1H), 6.42 (s, 1H), 6.20 (d, J = 17.0 Hz, 1H), 6.03 - 5.91 (m, 2H), 5.56 (d, J = 10.3 Hz, 1H), 4.93 (d, J = 7.1 Hz, 1H), 4.81 (s, 1H), 4.67 (s, 1H), 4.41 (s, 1H), 4.25-4.05 (m, 3H), 4.25 - 4.12 (m, 3H), 3.97 - 3.80 (m, 10H), 3.68 (s, 3H).
Example 759	5.7	42.2	61.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.89 (d, J = 11.1 Hz, 1H), 6.69 (s, 1H), 6.61 (s, 1H), 6.35 - 6.09 (m, 2H), 5.94 (dd, J = 17.0, 10.3 Hz, 1H), 5.59 (d, J = 10.1 Hz, 2H), 5.00 (p, J = 7.5 Hz, 1H), 4.83 (s, 1H), 4.31 - 4.18 (m, 2H), 4.16 (d, J = 8.1 Hz, 1H), 4.01 - 3.85 (m, 12H), 3.67 (d, J = 14.5 Hz, 2H).
Example 760	2.9	5.5	6.7	¹ H NMR (400 MHz, CDCl ₃) δ 8.86 (d, J = 5.2 Hz, 1H), 6.68 (d, J = 10.6 Hz, 1H), 6.61 (s, 1H), 6.42 - 6.21 (m, 1H), 6.14 (t, J = 18.1 Hz, 1H), 6.00 - 5.78 (m, 1H), 5.74 (s, 1H), 5.49 (dd, J = 29.1, 10.3 Hz, 1H), 4.91 (t, J = 7.6 Hz, 1H), 4.87 - 4.78 (m, 1H), 4.52 - 4.40 (m, 1H), 4.24 - 4.14 (m, 2H), 4.08 (d, J = 8.7 Hz, 1H), 3.95 (s, 6H), 3.92 - 3.83 (m, 2H), 3.80 - 3.70 (m, 2H), 2.03 - 1.89 (m, 2H), 1.54 - 1.43 (m, 3H).
Example 761	14.3	20.7	51.8	¹ H NMR (400 MHz, CDCl ₃) δ 9.14 (s, 1H), 8.64 (s, 1H), 8.53 (s, 1H), 7.74 (s, 1H), 6.88 (d, J = 2.7 Hz, 1H), 6.61 (d, J = 2.7 Hz, 1H), 6.20 (d, J = 16.9 Hz, 1H), 6.01 (s, 1H), 5.56 (d, J = 10.3 Hz, 1H), 5.01 - 4.91

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				(m, 2H), 4.32 - 4.23 (m, 2H), 4.02 (s, 3H), 3.99 - 3.95 (m, 1H), 3.95 (s, 3H), 3.87 (s, 3H), 3.84 - 3.82 (m, 1H).
Example 762	5.2	3.2	4.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.94 (s, 1H), 6.83 (s, 1H), 6.63 (s, 1H), 6.30 (s, 1H), 6.22 (d, J = 17.1 Hz, 1H), 6.04 (dd, J = 16.9, 10.2 Hz, 1H), 5.81 (s, 1H), 5.60 (d, J = 10.2 Hz, 1H), 4.87 - 4.80 (m, 1H), 4.80 - 4.73 (m, 1H), 4.70 (d, J = 6.4 Hz, 2H), 4.60 (d, J = 13.8 Hz, 2H), 4.27 (d, J = 13.6 Hz, 1H), 4.23 - 4.15 (m, 2H), 4.10 (d, J = 13.7 Hz, 1H), 3.96 (s, 6H), 3.86 (dd, J = 9.6, 4.0 Hz, 1H), 3.75 (dd, J = 9.4, 5.8 Hz, 1H), 3.23 (q, J = 7.2 Hz, 1H), 2.08 (d, J = 8.6 Hz, 1H).
Example 763	7.9	7.8	11.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.91 (s, 1H), 6.79 (s, 1H), 6.63 (s, 1H), 6.25 (dd, J = 23.9, 17.1 Hz, 2H), 5.63 (dd, J = 19.8, 10.3 Hz, 2H), 4.76 (s, 2H), 4.66 (s, 1H), 4.22 (dd, J = 15.1, 7.2 Hz, 3H), 4.14 - 4.10 (m, 1H), 3.97 (d, J = 2.0 Hz, 8H), 3.85 (d, J = 10.5 Hz, 3H), 3.75 (s, 1H), 2.00 (d, J = 9.2 Hz, 1H), 1.93 (s, 1H).
Example 764	7.0	49.4	60.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.78 (s, 1H), 6.84 (brs, 2H), 6.76 (s, 1H), 6.58 (t, J = 7.9 Hz, 1H), 6.39 (brs, 1H), 6.08 (d, J = 16.9 Hz, 1H), 5.93 (dd, J = 17.0, 10.2 Hz, 1H), 5.39 (d, J = 10.2 Hz, 1H), 4.81 (p, J = 6.6, 5.6 Hz, 2H), 4.15 - 4.05 (m, 2H), 3.85 - 3.81 (m, 1H), 3.81 (s, 6H), 3.71 (dd, J = 9.2, 5.8 Hz, 1H), 3.08 (s, 3H).
Example 765	158	420.8	656.5	¹ H NMR (400 MHz, CDCl ₃) δ 8.93 (s, 1H), 6.80 (s, 1H), 6.64 (s, 1H), 6.23 (d, J = 16.9 Hz, 1H), 6.08 - 5.91 (m, 4H), 5.75 (s, 1H), 5.58 (d, J = 10.4 Hz, 1H), 4.89 (s, 2H), 4.24 (ddd, J = 23.7, 9.4, 5.9 Hz, 2H), 3.96 (s, 6H), 3.88 (dd, J = 9.8, 3.4 Hz, 1H), 3.76 (dd, J = 9.2, 5.5 Hz, 1H).
Example 766	8.3	19.8	27.9	¹ H NMR (400 MHz, CDCl ₃) δ 8.80 (s, 1H), 6.60 (s, 1H), 6.54 (s, 1H), 6.33 (brs, 1H),

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				6.12 (d, J = 16.9 Hz, 1H), 5.96 (dd, J = 17.0, 10.3 Hz, 1H), 5.48 (dd, J = 10.3, 1.4 Hz, 1H), 4.83-4.78 (m, 2H), 4.17-4.06 (m, 2H), 3.84-3.81 (m, 1H), 3.71-3.68 (m, 1H).
Example 767	7	23.9	45.1	¹ H NMR (400 MHz, CDC1 ₃) δ 8.87 (s, 1H), 6.87 (s, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.60 (brs, 1H), 6.22 (s, 1H), 6.17 (s, 1H), 5.96 (dd, J = 16.9, 10.3 Hz, 1H), 5.80 (brs, 1H), 5.56 (d, J = 10.3 Hz, 1H), 4.94 - 4.83 (m, 2H), 4.30 - 4.18 (m, 2H), 3.91 (s, 6H), 3.88 (s, 1H), 3.75 (dd, J = 9.3, 5.9 Hz, 1H), 3.65 (s, 2H), 1.36 (t, J = 7.2 Hz, 3H).
Example 768	4	9.9	12.5	¹ H NMR (400 MHz, CDC1 ₃) δ 8.95 (s, 1H), 7.09 (s, 1H), 6.69 (t, J = 7.9 Hz, 1H), 6.26 (dd, J = 17.0, 1.3 Hz, 1H), 6.14 (d, J = 7.4 Hz, 1H), 6.03 (dd, J = 17.0, 10.3 Hz, 1H), 5.75 (d, J = 7.1 Hz, 1H), 5.63 (dd, J = 10.3, 1.3 Hz, 1H), 4.89 - 4.83 (m, 1H), 4.82 - 4.72 (m, 5H), 4.23 (dd, J = 9.5, 6.2 Hz, 2H), 3.92 (s, 6H), 3.81 (ddd, J = 19.8, 9.5, 4.8 Hz, 2H).
Example 769	3.7	8.4	10.0	¹ H NMR (400 MHz, CDC1 ₃) δ 8.94 (s, 1H), 7.04 (s, 1H), 6.69 (t, J = 7.8 Hz, 1H), 6.34 (brs, 1H), 6.22 (d, J = 16.9 Hz, 1H), 6.04 (dd, J = 17.1, 10.1 Hz, 1H), 5.60 (d, J = 10.2 Hz, 1H), 4.87 - 4.70 (m, 4H), 4.63 (d, J = 13.5 Hz, 2H), 4.30 (d, J = 13.6 Hz, 1H), 4.23 - 4.10 (m, 3H), 3.92 (s, 6H), 3.90 - 3.85 (m, 1H), 3.79 - 3.73 (m, 1H), 3.29 - 3.24 (m, 1H), 2.08 - 2.01 (m, 2H).
Example 770	2413.9	>10000	>10000	¹ H NMR (400 MHz, CDC1 ₃) δ 8.90 (s, 1H), 7.78 (t, J = 7.5 Hz, 1H), 7.31 (s, 1H), 7.17 (t, J = 8.0 Hz, 1H), 6.96 (t, J = 8.0 Hz, 1H), 6.49 (m, 1H), 6.27 (brs, 1H), 6.18 (d, J = 16.9 Hz, 1H), 5.93 (dd, J = 17.0, 10.3 Hz, 1H), 5.84, (brs, 1H), 5.53 (d, J = 10.3 Hz, 1H), 4.91-4.88 (m, 2H), 4.31 - 4.13 (m, 2H), 3.94 (s, 3H), 3.94-3.88(m, 1H), 3.82 - 3.70 (m, 1H), 3.21 (d, J = 4.8 Hz, 3H).

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
Example 771	7.9	25.5	32.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.88 (s, 1H), 6.67 (s, 1H), 6.63 (s, 1H), 6.22, (brs, 1H), 6.20 (dd, <i>J</i> = 17.0, 1.4 Hz, 1H), 5.97 (dd, <i>J</i> = 17.0, 10.3 Hz, 1H), 5.79 (brs, 1H), 5.55 (dd, <i>J</i> = 10.3, 1.5 Hz, 1H), 4.94 - 4.86 (m, 2H), 4.26-4.18 (m, 2H), 3.96 (s, 6H), 3.88 (dd, <i>J</i> = 9.8, 3.2 Hz, 1H), 3.81 - 3.72 (m, 1H).
Example 772	2.9	6.8	8.1	¹ H NMR (400 MHz, CDCl ₃) δ 8.77 (s, 1H), 6.81 (s, 1H), 6.59 (s, 1H), 6.57 (brs, 1H), 6.14 (d, <i>J</i> = 16.6 Hz, 1H), 5.96 (m, 1H), 5.85 (brs, 1H), 5.53 (d, <i>J</i> = 10.4 Hz, 1H), 4.87 - 4.66 (m, 2H), 4.53 - 4.23 (m, 4H), 4.18-4.08 (m, 2H), 3.83 (s, 6H), 3.82-3.69 (m, 2H), 1.95-1.75 (m, 2H), 1.51 (s, 3H).
Example 773	1.8	2.0	2.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.84 (s, 1H), 6.87 (d, <i>J</i> = 5.2 Hz, 1H), 6.66 (t, <i>J</i> = 8.1 Hz, 1H), 6.45 (brs, 1H), 6.14 (dd, <i>J</i> = 16.9, 12.4 Hz, 1H), 5.96 (dd, <i>J</i> = 16.9, 10.3 Hz, 1H), 5.82 (brs, 1H), 5.50 (dd, <i>J</i> = 20.2, 10.3 Hz, 1H), 4.93-4.85 (m, 1H), 4.84-4.78 (m, 1H), 4.43-4.34 (m, 2H), 4.19-3.77 (m, 5H), 3.90 (s, 6H), 3.72 - 3.69 (m, 1H), 1.49 (d, <i>J</i> = 23.8 Hz, 3H).
Example 774	3.2	6.9	8.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.87 (s, 1H), 6.88 (s, 1H), 6.67 (t, <i>J</i> = 7.9 Hz, 1H), 6.52 (brs, 1H), 6.24 (dd, <i>J</i> = 17.0, 1.5 Hz, 1H), 6.08 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 5.96 (brs, 1H), 5.61 (dd, <i>J</i> = 10.2, 1.4 Hz, 1H), 4.82 - 4.72 (m, 2H), 4.57 (brs, 4H), 4.26 - 4.16 (m, 2H), 3.91 (s, 6H), 3.85 (dd, <i>J</i> = 9.4, 3.9 Hz, 1H), 3.78 (dd, <i>J</i> = 9.4, 5.0 Hz, 1H), 2.42 (p, <i>J</i> = 7.6 Hz, 2H).
Example 775	8.9	20.8	27.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.96 (s, 1H), 6.69 (t, <i>J</i> = 8.0 Hz, 1H), 6.65 (s, 1H), 6.25 (dd, <i>J</i> = 17.0, 1.4 Hz, 1H), 6.08 (s, 1H), 6.02 (dd, <i>J</i> = 16.9, 10.3 Hz, 1H), 5.60 (dd, <i>J</i> = 10.3, 1.4 Hz, 1H), 4.73 (brs, 1H), 4.56 (brs, 1H), 4.42 (d, <i>J</i> = 8.5 Hz, 1H), 4.39 - 4.22 (m, 4H), 4.05 (dd, <i>J</i> =

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				12.0, 4.5 Hz, 1H), 3.99 (d, J = 11.7 Hz, 1H), 3.92 (s, 6H), 3.75 (dd, J = 11.9, 1.6 Hz, 1H), 3.65 - 3.57 (m, 1H), 3.33 (s, 3H), 2.09 - 2.02 (m, 1H), 1.91 - 1.81 (m, 1H).
Example 776	4.9	7.9	9.4	¹ H NMR (400 MHz, CDC1 ₃) δ 8.89 (s, 1H), 6.92 (s, 1H), 6.67 (t, J = 7.9 Hz, 1H), 6.38 (brs, 1H), 6.23 (dd, J = 17.0, 1.4 Hz, 1H), 6.03 (dd, J = 16.9, 10.3 Hz, 1H), 5.80 (brs, 1H), 5.61 (dd, J = 10.3, 1.4 Hz, 1H), 4.86 - 4.77 (m, 2H), 4.27 - 4.19 (m, 6H), 3.91 (s, 6H), 3.89 - 3.86 (m, 1H), 3.76 (dd, J = 9.3, 5.7 Hz, 1H), 3.69 - 3.66 (m, 4H), 1.86 (t, J = 5.2 Hz, 4H).
Example 777	NT	650.8	660	¹ H NMR (400 MHz, CDC1 ₃) δ 8.80 (s, 1H), 6.86-6.81 (m, 3H), 6.45 (brs, 1H), 6.13 (d, J = 16.9 Hz, 1H), 6.06 (brs, 1H), 5.86 (dd, J = 16.9, 10.3 Hz, 1H), 5.67 (brs, 1H), 5.49 (d, J = 10.4 Hz, 1H), 4.98 - 4.68 (m, 2H), 4.28-4.07 (m, 2H), 3.84 (s, 3H), 3.84-3.71 (m, 1H), 3.67 (dd, J = 9.2, 6.0 Hz, 1H), 3.08 (d, J = 4.8 Hz, 3H).
Example 778	NT	9.5	11.2	¹ H NMR (400 MHz, CDC1 ₃) δ 8.87 (s, 1H), 6.70 (s, 1H), 6.61 (s, 1H), 6.29 (brs, 1H), 6.11 (d, J = 17.1 Hz, 1H), 5.81 (m, 2H), 5.45 (d, J = 10.2 Hz, 1H), 5.07-4.84 (m, 2H), 4.55-4.40 (m, 1H), 4.22-4.15 (m, 2H), 4.05-3.75 (m, 3H), 3.95 (s, 6H), 3.91-3.71 (m, 2H), 2.05-1.96 (m, 2H), 1.44 (s, 3H).
Example 779	NT	7.5	8.6	¹ H NMR (400 MHz, CDC1 ₃) δ 8.87 (s, 1H), 6.70 (s, 1H), 6.61 (s, 1H), 6.43 (brs, 1H), 6.11 (d, J = 17.1 Hz, 1H), 5.97 (dd, J = 16.9, 10.3 Hz, 1H), 5.79 (brs, 1H), 5.45 (d, J = 10.2 Hz, 1H), 5.01-4.75 (m, 2H), 4.55-4.40 (m, 1H), 4.25-4.00 (m, 4H), 3.95 (s, 6H), 3.94-4.80 (m, 2H), 3.79-3.72 (m, 1H), 2.03-1.93 (m, 2H), 1.53 (s, 3H).
Example 786	17.0	38.4	47.6	¹ H NMR (400 MHz, CDC1 ₃) δ 8.87 (s, 1H), 6.86 (s, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.49 (s, 1H), 6.20 (dd, J = 16.9, 1.3 Hz, 1H), 6.11 (s, 1H), 6.02 (d, J = 8.6 Hz, 1H), 5.88

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				(dd, J = 17.0, 10.3 Hz, 1H), 5.53 (dd, J = 10.4, 1.4 Hz, 1H), 4.73 (d, J = 8.5 Hz, 1H), 4.38 - 4.26 (m, 1H), 4.08 (d, J = 12.1 Hz, 1H), 3.99 (d, J = 12.1 Hz, 1H), 3.92 (s, 6H), 3.75 (dd, J = 12.0, 1.7 Hz, 1H), 3.61 (m, 1H), 3.13 (s, 3H), 1.91 (m, 2H).
Example 787	45.0	86.0	102.8	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.89 (s, 1H), 6.90 (t, J = 8.1 Hz, 1H), 6.81 (s, 1H), 6.13 (dd, J = 17.1, 1.9 Hz, 1H), 5.96 (dd, J = 17.0, 10.3 Hz, 1H), 5.47 (dd, J = 10.2, 1.9 Hz, 1H), 4.24 (dt, J = 12.4, 4.2 Hz, 1H), 4.07 (dd, J = 11.9, 4.4 Hz, 1H), 3.99 - 3.93 (m, 1H), 3.90 (s, 6H), 3.78 (dd, J = 12.0, 2.0 Hz, 1H), 3.67 - 3.52 (m, 2H), 3.47 - 3.33 (m, 1H), 2.06 (qd, J = 12.5, 4.8 Hz, 1H), 1.77 - 1.66 (m, 1H), 1.46 - 1.26 (m, 5H).
Example 789	7.8	12.6	15.9	¹ H NMR (400 MHz, CDCl ₃) δ 9.02 (s, 1H), 7.24 (s, 1H), 7.00 (brs, 1H), 6.70 (t, J = 8.0 Hz, 1H), 6.25 (dd, J = 16.9, 1.3 Hz, 1H), 6.02 (dd, J = 17.0, 10.3 Hz, 1H), 5.57 (d, J = 10.3 Hz, 1H), 4.50 - 4.30 (m, 2H), 4.19 - 3.76 (m, 11H), 3.97 (s, 6H), 3.63 (t, J = 10.2 Hz, 1H), 2.32 - 1.75 (m, 2H).
Example 790	8.5	11.4	13.6	¹ H NMR (400 MHz, CDCl ₃) δ 8.93 (s, 1H), 6.97 (s, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.24 (dd, J = 17.1, 1.4 Hz, 1H), 6.04 (brs, 1H), 6.01 (dd, J = 16.9, 10.3 Hz, 1H), 5.63 - 5.53 (m, 1H), 4.62 - 4.31 (m, 4H), 4.22 - 3.95 (m, 4H), 3.92 (s, 6H), 3.78 - 3.72 (m, 1H), 3.66 - 3.57 (m, 1H), 3.28 (s, 3H), 2.13 - 2.05 (m, 1H), 1.90 - 1.78 (m, 1H), 1.53 (s, 3H).
Example 791	3.7	6.4	8.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.97 (s, 1H), 7.10 (s, 1H), 6.70 (t, J = 7.9 Hz, 1H), 6.48 (d, J = 7.2 Hz, 1H), 6.27 (dd, J = 17.0, 1.4 Hz, 1H), 6.18 - 6.08 (m, 1H), 5.98 (dd, J = 16.9, 10.3 Hz, 1H), 5.59 (dd, J = 10.3, 1.3 Hz, 1H), 4.80 (q, J = 12.0 Hz, 2H), 4.68 (q, J = 11.9 Hz, 2H), 4.44 (d, J = 10.0 Hz, 1H), 4.41 - 4.34 (m, 1H), 4.06 (dd, J = 11.9, 4.6

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				Hz, 1H), 3.99 (d, J = 11.8 Hz, 1H), 3.92 (s, 6H), 3.76 (dd, J = 11.9, 1.6 Hz, 1H), 3.62 (td, J = 12.0, 2.5 Hz, 1H), 2.02 (s, 1H), 1.92 - 1.81 (m, 1H).
Example 792	3.1	4.8	6.2	¹ H NMR (400 MHz, CDCl ₃) δ 8.91 (s, 1H), 6.92 (s, 1H), 6.83 (d, J = 6.8 Hz, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.26 (dd, J = 16.9, 1.5 Hz, 1H), 6.07 (dd, J = 17.0, 10.2 Hz, 1H), 6.02 (brs, 1H), 5.61 (dd, J = 10.3, 1.5 Hz, 1H), 4.53 (s, 2H), 4.48 - 4.36 (m, 3H), 4.35 - 4.29 (m, 1H), 4.05 (dd, J = 11.9, 4.6 Hz, 1H), 3.99 (d, J = 11.7 Hz, 1H), 3.91 (s, 6H), 3.74 (dd, J = 11.8, 1.6 Hz, 1H), 3.61 (td, J = 12.0, 2.5 Hz, 1H), 2.41 - 2.32 (m, 2H), 2.12 - 2.04 (m, 1H), 1.91 - 1.80 (m, 1H).
Example 793	5.3	7.9	9.9	¹ H NMR (400 MHz, CDCl ₃) δ 8.92 (s, 1H), 6.97 (s, 1H), 6.67 (t, J = 7.9 Hz, 1H), 6.63 (brs, 1H), 6.23 (dd, J = 17.0, 1.3 Hz, 1H), 6.04 (d, J = 8.8 Hz, 1H), 5.99 (dd, J = 17.1, 10.4 Hz, 1H), 5.57 (dd, J = 10.2, 1.4 Hz, 1H), 4.51 - 4.39 (m, 3H), 4.38 - 4.31 (m, 1H), 4.30 - 4.20 (m, 2H), 4.07 - 3.96 (m, 2H), 3.91 (s, 6H), 3.75 (dd, J = 11.8, 1.6 Hz, 1H), 3.65 - 3.56 (m, 1H), 2.05 - 1.98 (m, 1H), 1.91 - 1.80 (m, 1H), 1.56 (s, 3H).
Example 794	4.8	5.5	6.9	¹ H NMR (400 MHz, CDCl ₃) δ 8.90 (d, J = 3.0 Hz, 1H), 6.93 (d, J = 4.0 Hz, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.36 (brs, 1H), 6.26 - 6.15 (m, 1H), 6.05-5.90 (m, 2H), 5.60-5.50 (m, 1H), 4.50-4.30 (m, 2H), 3.92 (s, 6H), 4.23-3.80 (m, 7H), 3.70-3.65 (m, 1H), 3.60-3.55 (m, 1H), 3.34 (d, J = 7.0 Hz, 3H), 2.02-1.90 (m, 2H), 1.87-1.78 (m, 2H).
Example 795	7.5	9.5	11.8	¹ H NMR (400 MHz, CDCl ₃) δ 9.00 (s, 1H), 7.20 (s, 1H), 6.69 (t, J = 7.9 Hz, 1H), 6.22 (dd, J = 17.0, 1.5 Hz, 1H), 6.02 (dd, J = 17.0, 10.3 Hz, 1H), 5.55 (d, J = 10.8 Hz, 1H), 4.54 - 4.15 (m, 4H), 4.13-4.00 (m, 2H), 3.93 (s, 6H), 3.77 (d, J = 12.0 Hz, 1H), 3.70 - 3.57 (m, 1H), 3.52-3.38 (m,

Example No.	Enzymatic activity	Cell activity		¹ H-NMR
	FGFR4 IC ₅₀ (nM)	HuH-7 EC ₅₀ (nM)	HuH-7 Absolute IC ₅₀ (nM)	
				3H), 3.37 (s, 3H), 2.07-1.85 (m, 2H), 1.65 - 1.46 (m, 4H).
Example 796	23.0	38.0	47.8	¹ H NMR (400 MHz, CDCl ₃) δ 8.95 (s, 1H), 7.04 (s, 1H), 6.68 (t, J = 7.9 Hz, 1H), 6.25 (s, 1H), 6.20 (dd, J = 17.0, 1.4 Hz, 1H), 6.07 (s, 1H), 5.93 (dd, J = 16.9, 10.3 Hz, 1H), 5.56 (dd, J = 10.2, 1.4 Hz, 1H), 4.71 (d, J = 6.4 Hz, 2H), 4.61 (d, J = 13.6 Hz, 1H), 4.55 (d, J = 13.6 Hz, 1H), 4.45 (d, J = 8.9 Hz, 1H), 4.41 - 4.34 (m, 1H), 4.17 (d, J = 13.7 Hz, 1H), 4.07 (d, J = 13.5 Hz, 2H), 3.99 (d, J = 11.9 Hz, 1H), 3.92 (s, 6H), 3.73 (d, J = 11.6 Hz, 1H), 3.67 - 3.64 (m, 1H), 3.59 (td, J = 11.9, 2.6 Hz, 1H), 3.22 (q, J = 7.1 Hz, 1H), 1.97 (d, J = 13.7 Hz, 1H), 1.92 - 1.81 (m, 1H).
BLU-9931	24.7	549	1362	
BLU-554	18.0	22	126	
Note	"NT", i.e., "Not Tested", means that the compound is not tested. BLU-9931 and BLU-554 are positive compounds.			

[0357] It can be seen from the enzymatic activity data of the compounds of specific examples that the compounds of the present invention have a strong inhibitory effect on FGFR4 kinase activity. It can be seen from the cell activity data of the compounds of specific examples that the compounds of the present invention have a strong inhibitory effect on the proliferation activity of HuH-7 cells highly expressing FGFR4.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not

form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2014011900A](#) [0008]
- [WO2014144373A](#) [0010]
- [WO2016115412A](#) [0011]
- [WO2015061572A](#) [0012]
- [WO2015108992A](#) [0013]


Non-patent literature cited in the description

- **ESWARAKUMAR et al.** Cytokine Growth Factor Rev, 2005, vol. 16, 139-149 [0002]
- **ORNITZITOH** Genome Bio12: Reviews, 2001, 3005- [0002]
- **ULLRICHSCHLESSINGER** Cell, 1990, vol. 61, 203- [0003]
- **POWERS et al.** Endocr Relat Cancer, 2000, vol. 7, 165-197 [0003]
- **CHESI et al.** Blood, 2001, vol. 97, 729-726 [0004]
- **GOWARDHAN et al.** Br J Cancer, 2005, vol. 92, 320-327 [0004]
- **JAAKKOLA et al.** Int J Cancer, 1993, vol. 54, 378-282 [0004]
- **JANG et al.** Cancer Res, 2001, vol. 61, 3541-3543 [0004]
- **SHARIFF et al.** Expert Rev Gastroenterol Hepato, 2009, vol. 13, 353-367 [0005]
- **YE et al.** Cancer, 2011, 5304-5313 [0006]
- **XU et al.** BMC Cancer, 2011, vol. 11, 84- [0006]
- **FAWDAR et al.** PNAS, 2013, vol. 110, 12426-12431 [0006]

FGFR4-HÆMMER, FREMGANGSMÅDE TIL FREMSTILLING DERAFT OG FARMACEUTISK ANVENDELSE DERAFT


Patentkrav

1. Forbindelse med formlen (IVa-1), stereoisomer eller farmaceutisk acceptabelt salt deraf:



(IVa-1)

hvor X_5 er -CH- eller N,
m er 1 eller 0,



R_2 , R_3 , R_4 og R_5 hver er uafhængigt valgt fra gruppen bestående af H, deuterium, Cl, F, hydroxyl, methyl, isopropyl, cyclopropyl, 3-oxacyclobutyl, trifluormethyl, trideuteromethyl og -O- R_{11} ,

eller R_2 og R_4 , R_3 og R_5 sammen med de direkte bundne carbonatomer danner en 5-8-leddet heterocyclyl, idet heteroatomet er N eller O,

R_8 er valgt fra gruppen bestående af H, deuterium, halogen, cyano, nitro, azido, C_{1-8} -alkyl, C_{2-8} -alkenyl, C_{2-8} -alkynyl, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, - C_{0-8} -S(O) $_r$ R_{10} , - C_{0-8} -P(O)(R_{10}) $_2$, - C_{0-8} -O- R_{11} , - C_{0-8} -C(O)OR $_{11}$, - C_{0-8} -C(O)R $_{12}$, - C_{0-8} -O-C(O)R $_{12}$, - C_{0-8} -NR $_{13}$ R $_{14}$, - C_{0-8} -C(O)NR $_{13}$ R $_{14}$, - C_{0-8} -N(R $_{13}$)-C(O)R $_{12}$ og - C_{0-8} -N(R $_{13}$)-C(O)OR $_{11}$,

ovenstående grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, cyano, nitro, azido, C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, 3-8-leddet heterocyclyl, C_{5-8} -aryl, 5-8-leddet heteroaryl, - C_{0-4} -S(O) $_r$ R_{10} , - C_{0-4} -O- R_{11} , - C_{0-4} -C(O)OR $_{11}$, - C_{0-4} -C(O)R $_{12}$, - C_{0-4} -O-C(O)R $_{12}$, - C_{0-4} -NR $_{13}$ R $_{14}$, - C_{0-4} -C(O)NR $_{13}$ R $_{14}$, - C_{0-4} -N(R $_{13}$)-C(O)R $_{12}$ og - C_{0-4} -N(R $_{13}$)-C(O)OR $_{11}$, idet ovenstående grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, cyano, nitro, azido, C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, 3-8-leddet heterocyclyl, C_{5-8} -aryl, 5-8-leddet heteroaryl, - C_{0-4} -S(O) $_r$ R_{10} , - C_{0-4} -O- R_{11} , - C_{0-4} -

$C(O)OR_{11}$, $-C_{0-4}-C(O)R_{12}$, $-C_{0-4}-O-C(O)R_{12}$, $-C_{0-4}-NR_{13}R_{14}$, $-C_{0-4}-C(O)NR_{13}R_{14}$, $-C_{0-4}-N(R_{13})-C(O)R_{12}$ og $-C_{0-4}-N(R_{13})-C(O)OR_{11}$,

X_1 er $-CH-$ eller N ,

R_{10} er valgt fra gruppen bestående af H , deuterium, C_{1-8} -alkyl, C_{2-8} -alkenyl, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{1-8} -halogenalkyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, amino, mono- C_{1-8} -alkylamino, di- C_{1-8} -alkylamino og C_{1-8} -alkanoylamino,

R_{11} er valgt fra gruppen bestående af H , deuterium, C_{1-8} -alkyl, C_{2-8} -alkenyl, C_{2-8} -alkynyl, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl og 5-10-leddet heteroaryl, idet ovennævnte grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, cyano, C_{1-8} -alkyl, C_{1-8} -alkoxy, C_{1-8} -alkylthio, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, C_{1-8} -alkylsulfonyl, C_{1-8} -alkylsulfonylamino, amino, mono- C_{1-8} -alkylamino, di- C_{1-8} -alkylamino, $=O$ eller hydroxyl,

R_{12} er valgt fra gruppen bestående af H , deuterium, C_{1-8} -alkyl, C_{2-8} -alkenyl, C_{2-8} -alkynyl, C_{1-8} -alkoxy, C_{3-8} -cycloalkyl, C_{3-8} -cycloalkyloxy, 3-10-leddet heterocyclyl, 3-10-leddet heterocycliloxy, C_{5-10} -aryl, 5-10-leddet heteroaryl, C_{5-10} -aryloxy og 5-10-leddet heteroaryloxy, idet ovennævnte grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, cyano, C_{1-8} -alkyl, C_{1-8} -alkoxy, C_{1-8} -alkylthio, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, C_{1-8} -alkylsulfonyl, C_{1-8} -alkylsulfonylamino, amino, mono- C_{1-8} -alkylamino, di- C_{1-8} -alkylamino, $=O$ eller hydroxyl,

R_{13} og R_{14} hver er uafhængigt valgt fra gruppen bestående af H , deuterium, C_{1-8} -alkyl, C_{2-8} -alkenyl, C_{2-8} -alkynyl, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, C_{1-8} -alkylsulfonyl og C_{1-8} -alkanoyl, eller R_{13} og R_{14} sammen med det direkte bundne nitrogenatom danner en 4-10-leddet heterocyclyl,

idet ovennævnte grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, C_{1-8} -alkyl, C_{1-8} -alkoxy, C_{1-8} -alkylthio, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, C_{1-8} -alkylsulfonyl, C_{1-8} -alkylsulfonylamino, amino, mono- C_{1-8} -alkylamino, di- C_{1-8} -alkylamino, $=O$ eller hydroxyl,

r er 0, 1 eller 2,

hvor cycloalkyl betyder en mættet eller delvist umættet monocyklisk eller polycyklisk hydrocarbonsubstituent.


2. Forbindelse med formlen (IVa-1), stereoisomer eller farmaceutisk acceptabelt


salt deraf ifølge krav 1, hvor


R_8 er valgt fra gruppen bestående af H, deuterium, halogen, cyano, C_{1-8} -alkyl, C_{2-8} -alkenyl, C_{2-8} -alkynyl, C_{3-8} -cycloalkyl, 3-10-leddet heterocyclyl, C_{5-10} -aryl, 5-10-leddet heteroaryl, $-C_{0-8}-O-R_{11}$, $-C_{0-8}-NR_{13}R_{14}$, $-C_{0-8}-N(R_{13})-C(O)R_{12}$ og $-C_{0-8}-N(R_{13})-C(O)OR_{11}$,


ovenstående grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, cyano, nitro, azido, C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, 3-8-leddet heterocyclyl, C_{5-8} -aryl, 5-8-leddet heteroaryl, $-C_{0-4}-S(O)_rR_{10}$, $-C_{0-4}-O-R_{11}$, $-C_{0-4}-C(O)OR_{11}$, $-C_{0-4}-C(O)R_{12}$, $-C_{0-4}-O-C(O)R_{12}$, $-C_{0-4}-NR_{13}R_{14}$, $-C_{0-4}-C(O)NR_{13}R_{14}$, $-C_{0-4}-N(R_{13})-C(O)R_{12}$ og $-C_{0-4}-N(R_{13})-C(O)OR_{11}$, idet ovenstående grupper eventuelt yderligere er substitueret med en eller flere substituent, der er valgt fra gruppen bestående af deuterium, halogen, cyano, nitro, azido, C_{1-4} -alkyl, C_{2-4} -alkenyl, C_{2-4} -alkynyl, C_{1-4} -haloalkyl, C_{3-6} -cycloalkyl, 3-8-leddet heterocyclyl, C_{5-8} -aryl, 5-8-leddet heteroaryl, $-C_{0-4}-S(O)_rR_{10}$, $-C_{0-4}-O-R_{11}$, $-C_{0-4}-C(O)OR_{11}$, $-C_{0-4}-C(O)R_{12}$, $-C_{0-4}-O-C(O)R_{12}$, $-C_{0-4}-NR_{13}R_{14}$, $-C_{0-4}-C(O)NR_{13}R_{14}$, $-C_{0-4}-N(R_{13})-C(O)R_{12}$ og $-C_{0-4}-N(R_{13})-C(O)OR_{11}$.


3. Forbindelse med formlen (IVa-1), stereoisomer eller farmaceutisk acceptabelt salt deraf ifølge krav 1 eller 2, hvor forbindelsen er valgt fra gruppen bestående af:



































































































































































32




eller

4. Fremgangsmåde til fremstilling af forbindelsen med formelen (IVa-1), stereoisomeren eller det farmaceutisk acceptable salt deraf ifølge et hvilket som helst af kravene 1-3, hvor fremgangsmåden omfatter følgende trin:




(IVa-I)


eller



eller



eller



idet der eventuelt yderligere udføres en omdannelsesreaktion mellem de forskellige substituent i overensstemmelse med den forskellige substituent,

hvor X er en fraspaltelig gruppe, der er valgt fra gruppen bestående af Cl, Br, methylsulfanyl, methylsulfonyl og methoxy, R er valgt fra gruppen bestående af nitro, cyano og azido, Pg er en aminobeskyttelsesgruppe, der er valgt fra gruppen bestående af tert-butyloxycarbonyl, benzyloxycarbonyl, 2-biphenyl-2-propoxycarbonyl og p-toluensulfonyl, X₁, X₅, m, R₂, R₃, R₄, R₅, R₈ og Z er som beskrevet i krav 1.

5. Farmaceutisk sammensætning, der omfatter en terapeutisk virksom mængde af forbindelsen med formlen (IVa-1), stereoisomeren eller det farmaceutisk acceptable salt deraf ifølge et hvilket som helst af kravene 1-3 og en farmaceutisk acceptabel bærer.

6. Forbindelse med formlen (IVa-1), stereoisomer eller farmaceutisk acceptabelt salt deraf ifølge et hvilket som helst af kravene 1-3 eller farmaceutisk sammensætning ifølge krav 5 til anvendelse i behandlingen af cancer.

7. Forbindelse til anvendelse ifølge krav 6, hvor canceren er prostatacancer, levercancer, pancreascancer, øsofageal cancer, gastrisk cancer, lungecancer, brystcancer, ovariecancer, koloncancer, hudcancer, glioblastom eller rhabdomyosarkom.