



(43) International Publication Date
5 June 2014 (05.06.2014)

(10) International Publication Number
WO 2014/085208 A1

(51) International Patent Classification:
A61K 31/445 (2006.01)

(21) International Application Number:
PCT/US2013/071366

(22) International Filing Date:
22 November 2013 (22.11.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/730,144 27 November 2012 (27.11.2012) US

(71) Applicant: **MERCK SHARP & DOHME CORP.**
[US/US]; 126 East Lincoln Avenue, Rahway, New Jersey
07065-0907 (US).

(72) Inventors; and

(71) Applicants (for US only): **KUDUK, Scott, D.** [US/US];
770 Sumneytown Pike, West Point, Pennsylvania 19486
(US). **SKUDLAREK, Jason, W.** [US/US]; 770 Sum-
neytown Pike, West Point, Pennsylvania 19486 (US).

(74) Common Representative: **MERCK SHARP & DOHME
CORP.**; 126 East Lincoln Avenue, Rahway, New Jersey
07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: 2-PYRIDYLAMINO-4-NITRILE-PIPERIDINYL OREXIN RECEPTOR ANTAGONISTS

(57) Abstract: The present invention is directed to 2-pyridylamino-4-nitrile-piperidinyl compounds which are antagonists of orexin receptors. The present invention is also directed to uses of the 2-pyridylamino-4-nitrile-piperidinyl compounds described herein in the treatment or prevention of neurological and psychiatric disorders and diseases in which orexin receptors are involved. The present invention is also directed to pharmaceutical compositions comprising these compounds. The present invention is also directed to uses of these pharmaceutical compositions in the prevention or treatment of such diseases in which orexin receptors are involved.



WO 2014/085208 A1

TITLE OF THE INVENTION

2-PYRIDYLAMINO-4-NITRILE-PIPERIDINYL OREXIN RECEPTOR ANTAGONISTS

BACKGROUND OF THE INVENTION


5 The orexins (hypocretins) comprise two neuropeptides produced in the hypothalamus: the orexin A (OX-A) (a 33 amino acid peptide) and the orexin B (OX-B) (a 28 amino acid peptide) (Sakurai T. et al., Cell, 1998, 92, 573-585). Orexins are found to stimulate food consumption in rats suggesting a physiological role for these peptides as mediators in the central feedback mechanism that regulates feeding behavior (Sakurai T. et al., Cell, 1998, 92, 10 573-585). Orexins regulate states of sleep and wakefulness opening potentially novel therapeutic approaches for narcoleptic or insomniac patients (Chemelli R.M. et al., Cell, 1999, 98, 437-451). Orexins have also been indicated as playing a role in arousal, reward, learning and memory (Harris, et al., Trends Neurosci., 2006, 29 (10), 571-577). Two orexin receptors have been cloned and characterized in mammals. They belong to the super family of G-protein coupled 15 receptors (Sakurai T. et al., Cell, 1998, 92, 573-585): the orexin-1 receptor (OX or OX1R) is selective for OX-A and the orexin-2 receptor (OX2 or OX2R) is capable to bind OX-A as well as OX-B. The physiological actions in which orexins are presumed to participate are thought to be expressed via one or both of OX1 receptor and OX2 receptor as the two subtypes of orexin receptors.

SUMMARY OF THE INVENTION

20 The present invention is directed to 2-pyridylamino-4-nitrile-piperidinyll compounds which are antagonists of orexin receptors. The present invention is also directed to uses of the 2-pyridylamino-4-nitrile-piperidinyll compounds described herein in the treatment or 25 prevention of neurological and psychiatric disorders and diseases in which orexin receptors are involved. The present invention is also directed to pharmaceutical compositions comprising these compounds. The present invention is also directed to uses of these pharmaceutical compositions in the prevention or treatment of such diseases in which orexin receptors are involved.

DETAILED DESCRIPTION OF THE INVENTION

30 The present invention is directed to compounds of the formula I:



I

wherein:

A is selected from the group consisting of phenyl, naphthyl and heteroaryl;

5

R^{1a}, R^{1b} and R^{1c} are independently selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- (3) hydroxyl,
- 10 (4) $-(C=O)_m-O_n-C_{1-6}alkyl$, where m is 0 or 1, n is 0 or 1 (wherein if m is 0 or n is 0, a bond is present) and where the alkyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (5) $-(C=O)_m-O_n-C_{3-6}cycloalkyl$, where the cycloalkyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- 15 (6) $-(C=O)_m-C_{2-4}alkenyl$, where the alkenyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (7) $-(C=O)_m-C_{2-4}alkynyl$, where the alkynyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (8) $-(C=O)_m-O_n-phenyl$ or $-(C=O)_m-O_n-naphthyl$, where the phenyl or naphthyl is
- 20 unsubstituted or substituted with one or more substituents selected from R⁴,
- (9) $-(C=O)_m-O_n-heterocycle$, where the heterocycle is unsubstituted or substituted with one or more substituents selected from R⁴,
- (10) $-(C=O)_m-NR^{10}R^{11}$, wherein R¹⁰ and R¹¹ are independently selected from the group consisting of:
- 25 (a) hydrogen,
- (b) C₁₋₆alkyl, which is unsubstituted or substituted with R⁴,
- (c) C₃₋₆alkenyl, which is unsubstituted or substituted with R⁴,
- (d) C₃₋₆alkynyl, which is unsubstituted or substituted with R⁴,

- (e) C₃₋₆cycloalkyl which is unsubstituted or substituted with R⁴,
(f) phenyl, which is unsubstituted or substituted with R⁴, and
(g) heterocycle, which is unsubstituted or substituted with R⁴,
(11) -S(O)₂-NR¹⁰R¹¹,
5 (12) -S(O)_q-R¹², where q is 0, 1 or 2 and where R¹² is selected from the definitions of
R¹⁰ and R¹¹,
(13) -CO₂H,
(14) -CN, and
(15) -NO₂;

10 R³ is selected from C₁₋₆alkyl and C₃₋₆cycloalkyl, which is unsubstituted or substituted with
one or more substituents selected from R⁴;

R⁴ is selected from the group consisting of:

- 15 (1) hydroxyl,
(2) halogen,
(3) C₁₋₆alkyl,
(4) -C₃₋₆cycloalkyl,
(5) -O-C₁₋₆alkyl,
20 (6) -O(C=O)-C₁₋₆alkyl,
(7) -NH₂,
(7) -NH-C₁₋₆alkyl,
(8) -NO₂,
(9) phenyl,
25 (10) heterocycle,
(11) -CO₂H, and
(12) -CN;

R⁵ is selected from the group consisting of:

- 30 (1) hydrogen,
(2) halogen,
(3) hydroxyl,
(4) C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or
phenyl,


- (5) C₃₋₆cycloalkyl, which is unsubstituted or substituted with C₁₋₆alkyl, halogen, hydroxyl or phenyl,
 (6) -O-C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl, and
 5 (7) -S-C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl;

R⁶ is selected from the group consisting of:

- (1) hydrogen,
 10 (2) C₁₋₆alkyl, which is unsubstituted or substituted with oxo, halogen, hydroxyl or phenyl,
 (3) C₂₋₆alkenyl, and
 (4) -(CO)-O-C₁₋₆alkyl;

or a pharmaceutically acceptable salt thereof.


- 15 An embodiment of the present invention includes compounds of the formula Ia:



Ia

wherein R^{1a}, R^{1b}, R^{1c}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

- 20 An embodiment of the present invention includes compounds of the formula Ia':




Ia'

wherein R^{1a}, R^{1b}, R^{1c}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

5

An embodiment of the present invention includes compounds of the formula Ia'':




Ia''

wherein R^{1a}, R^{1b}, R^{1c}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

10


An embodiment of the present invention includes compounds of the formula Ib:



Ib

wherein R^{1a}, R^{1b}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

An embodiment of the present invention includes compounds of the formula Ib':




Ib'

wherein R^{1a}, R^{1b}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

5


An embodiment of the present invention includes compounds of the formula Ib'':



Ib''

wherein R^{1a}, R^{1b}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.


An embodiment of the present invention includes compounds of the formula Ic:



Ic

wherein R^{1a}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.


An embodiment of the present invention includes compounds of the formula Ic':



Ic'

wherein R^{1a}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

An embodiment of the present invention includes compounds of the formula Ic'':



Ic''

wherein R^{1a}, R³ and R⁵ are defined herein; or a pharmaceutically acceptable salt thereof.

An embodiment of the present invention includes compounds wherein A is selected from phenyl, pyridyl, thiophenyl, thiazolyl, isothiazolyl, and pyrazolyl. An embodiment of the present invention includes compounds wherein A is phenyl. An embodiment of the present invention includes compounds wherein A is pyridyl. An embodiment of the present invention includes compounds wherein A is thiophenyl. An embodiment of the present invention includes compounds wherein A is thiazolyl. An embodiment of the present invention includes compounds wherein A is isothiazolyl. An embodiment of the present invention includes compounds wherein A is pyrazolyl.

An embodiment of the present invention includes compounds wherein R^{1a}, R^{1b} and R^{1c} are independently selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- (3) hydroxyl,

- (4) C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,
- (5) -O-C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,
- 5 (6) heteroaryl, wherein heteroaryl is selected from triazolyl, tetrazolyl, oxazolyl, pyrrolyl, imidazolyl, indolyl, pyridyl, and pyrimidinyl, which is unsubstituted or substituted with halogen, hydroxyl, C₁₋₆alkyl, -O-C₁₋₆alkyl or -NO₂,
- (7) phenyl, which is unsubstituted or substituted with halogen, hydroxyl, C₁₋₆alkyl, -O-C₁₋₆alkyl or -NO₂,
- 10 (8) -O-phenyl, which is unsubstituted or substituted with halogen, hydroxyl, C₁₋₆alkyl, -O-C₁₋₆alkyl or -NO₂,
- (9) -CN, and
- (10) -NH-C₁₋₆alkyl, or -N(C₁₋₆alkyl)(C₁₋₆alkyl), which is unsubstituted or substituted with halogen, hydroxyl, C₁₋₆alkyl, and -O-C₁₋₆alkyl.

15 An embodiment of the present invention includes compounds wherein R^{1a}, R^{1b} and R^{1c} are independently selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- (3) hydroxyl,
- 20 (4) C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,
- (5) -O-C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,
- (6) -CN, and
- 25 (7) heteroaryl, wherein heteroaryl is selected from triazolyl, tetrazolyl, oxazolyl, pyrrolyl, imidazolyl, indolyl, pyridyl, and pyrimidinyl, which is unsubstituted or substituted with halogen, hydroxyl, C₁₋₆alkyl, -O-C₁₋₆alkyl or -NO₂.

An embodiment of the present invention includes compounds wherein R^{1a}, R^{1b} and R^{1c} are independently selected from the group consisting of:

- 30 (1) hydrogen,
- (2) halogen,
- (3) C₁₋₆alkyl, which is unsubstituted or substituted with halogen,
- (4) -O-C₁₋₆alkyl, which is unsubstituted or substituted with halogen,
- (5) -CN, and

- (6) heteroaryl, wherein heteroaryl is selected from triazolyl, tetrazolyl, oxazolyl, pyrrolyl, imidazolyl, indolyl, pyridyl, and pyrimidinyl.

An embodiment of the present invention includes compounds wherein R^{1c} is hydrogen, and R^{1a} and R^{1b} are independently selected from the group consisting of:

- 5 (1) hydrogen,
(2) fluoro,
(3) chloro,
(4) bromo,
(5) methyl,
10 (6) ethyl,
(7) trifluoromethyl, and
(8) heteroaryl, wherein heteroaryl is selected from triazolyl, tetrazolyl, oxazolyl, pyrrolyl, imidazolyl, indolyl, pyridyl, and pyrimidinyl.

An embodiment of the present invention includes compounds wherein R^{1c} is hydrogen, R^{1b} is hydrogen, and R^{1a} is independently selected from the group consisting of:

- 15 (1) triazolyl,
(2) tetrazolyl,
(3) oxazolyl,
(4) pyrrolyl,
20 (5) imidazolyl,
(6) pyridyl, and
(7) pyrimidinyl.

An embodiment of the present invention includes compounds wherein R^{1c} is hydrogen, R^{1b} is hydrogen, and R^{1a} is independently selected from the group consisting of:

- 25 (1) triazolyl,
(2) tetrazolyl, and
(3) pyrimidinyl.

An embodiment of the present invention includes compounds wherein R³ is C₁-6alkyl. An embodiment of the present invention includes compounds wherein R³ is C₃-6cycloalkyl. An embodiment of the present invention includes compounds wherein R³ is methyl or ethyl. An embodiment of the present invention includes compounds wherein R³ is methyl. An embodiment of the present invention includes compounds wherein R³ is (R)-methyl.

An embodiment of the present invention includes compounds wherein R⁵ is selected from the group consisting of:

- 35 (1) hydrogen,

- (2) chloro,
- (3) C₁₋₆alkyl, which is unsubstituted or substituted with halogen,
- (4) C₃₋₆cycloalkyl, which is unsubstituted or substituted with C₁₋₆alkyl or halogen,
- (5) -O-C₁₋₆alkyl, which is unsubstituted or substituted with halogen, and
- 5 (6) -S-C₁₋₆alkyl, which is unsubstituted or substituted with halogen.

An embodiment of the present invention includes compounds wherein R⁵ is selected from the group consisting of: hydrogen, methyl, cyclopropyl, chloro, -O-CH₃ and -S-CH₃.

10 An embodiment of the present invention includes compounds wherein R⁵ is hydrogen. An embodiment of the present invention includes compounds wherein R⁵ is methyl. An embodiment of the present invention includes compounds wherein R⁵ is cyclopropyl. An embodiment of the present invention includes compounds wherein R⁵ is chloro. An embodiment of the present invention includes compounds wherein R⁵ is -O-CH₃. An embodiment of the present invention includes compounds wherein R⁵ is -S-CH₃.

15 An embodiment of the present invention includes compounds wherein R⁶ is selected from the group consisting of:

- (1) hydrogen,
- (2) C₁₋₆alkyl,
- (3) C₂₋₆alkenyl, and
- 20 (4) -(CO)-O-C₁₋₆alkyl.

An embodiment of the present invention includes compounds wherein R⁶ is selected from the group consisting of: hydrogen, methyl, ethyl, propyl, propenyl, and -(CO)-O-CH₃.

25 An embodiment of the present invention includes compounds wherein R⁶ is hydrogen. An embodiment of the present invention includes compounds wherein R⁶ is methyl. An embodiment of the present invention includes compounds wherein R⁶ is ethyl. An embodiment of the present invention includes compounds wherein R⁶ is propyl. An embodiment of the present invention includes compounds wherein R⁶ is propenyl. An embodiment of the present invention includes compounds wherein R⁶ is
30 -(CO)-O-CH₃.

Certain embodiments of the present invention include a compound which is selected from the group consisting of the subject compounds of the Examples herein or a pharmaceutically acceptable salt thereof.

35 The compounds of the present invention may contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers,

diastereomeric mixtures and individual diastereomers. Additional asymmetric centers may be present depending upon the nature of the various substituents on the molecule. Each such asymmetric center will independently produce two optical isomers and it is intended that all of the possible optical isomers and diastereomers in mixtures and as pure or partially purified compounds are included within the ambit of this invention. The present invention is meant to comprehend all such isomeric forms of these compounds. Formula I shows the structure of the class of compounds without specific stereochemistry.

The independent syntheses of these diastereomers or their chromatographic separations may be achieved as known in the art by appropriate modification of the methodology disclosed herein. Their absolute stereochemistry may be determined by the x-ray crystallography of crystalline products or crystalline intermediates which are derivatized, if necessary, with a reagent containing an asymmetric center of known absolute configuration. If desired, racemic mixtures of the compounds may be separated so that the individual enantiomers are isolated. The separation can be carried out by methods well known in the art, such as the coupling of a racemic mixture of compounds to an enantiomerically pure compound to form a diastereomeric mixture, followed by separation of the individual diastereomers by standard methods, such as fractional crystallization or chromatography. The coupling reaction is often the formation of salts using an enantiomerically pure acid or base. The diastereomeric derivatives may then be converted to the pure enantiomers by cleavage of the added chiral residue. The racemic mixture of the compounds can also be separated directly by chromatographic methods utilizing chiral stationary phases, which methods are well known in the art. Alternatively, any enantiomer of a compound may be obtained by stereoselective synthesis using optically pure starting materials or reagents of known configuration by methods well known in the art.

As appreciated by those of skill in the art, halogen or halo as used herein are intended to include fluoro, chloro, bromo and iodo. Similarly, C₁₋₆, as in C₁₋₆alkyl is defined to identify the group as having 1, 2, 3, 4, 5 or 6 carbons in a linear or branched arrangement, such that C₁₋₈alkyl specifically includes methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, and hexyl. A group which is designated as being independently substituted with substituents may be independently substituted with multiple numbers of such substituents.

The term "heterocycle" as used herein includes both unsaturated and saturated heterocyclic moieties, wherein the unsaturated heterocyclic moieties (i.e. "heteroaryl") include benzoimidazolyl, benzimidazolonyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzothiazolyl, benzotriazolyl, benzothiophenyl, benzoxazepin, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, dihydroindolyl, indolaziny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl,

oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrahydroquinoxalinyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, and N-oxides thereof, and wherein the saturated heterocyclic moieties include azetidiny, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyridin-2-onyl, pyrrolidinyl, morpholinyl, tetrahydrofuranyl, thiomorpholinyl, and tetrahydrothienyl, and N-oxides thereof.

The present invention also includes all pharmaceutically acceptable isotopic variations of a compound of the Formula I in which one or more atoms is replaced by atoms having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes suitable for inclusion in the compounds of the invention include isotopes of hydrogen such as ^2H and ^3H , carbon such as ^{11}C , ^{13}C and ^{14}C , nitrogen such as ^{13}N and ^{15}N , oxygen such as ^{15}O , ^{17}O and ^{18}O , phosphorus such as ^{32}P , sulfur such as ^{35}S , fluorine such as ^{18}F , iodine such as ^{123}I and ^{125}I , and chlorine such as ^{36}Cl . Certain isotopically-labelled compounds of Formula I, for example those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, i.e. ^3H , and carbon-14, i.e. ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection. Substitution with heavier isotopes such as deuterium, i.e. ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life or reduced dosage requirements, and hence may be preferred in some circumstances. Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labelled compounds of Formula I can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples using appropriate isotopically-labelled reagents in place of the non-labelled reagent previously employed.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particular embodiments include the ammonium, calcium, magnesium, potassium, and sodium salts. Salts in the solid form may exist in more than one crystal structure, and may also be in the form of hydrates. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion

exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylene-diamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particular embodiments include the citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, fumaric, and tartaric acids. It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

Exemplifying the invention is the use of the compounds disclosed in the Examples and herein. Specific compounds within the present invention include a compound which selected from the group consisting of the compounds disclosed in the following Examples and pharmaceutically acceptable salts thereof and individual enantiomers or diastereomers thereof.

The subject compounds are useful in a method of antagonizing orexin receptor activity in a patient such as a mammal in need of such inhibition comprising the administration of an effective amount of the compound. The present invention is directed to the use of the compounds disclosed herein as antagonists of orexin receptor activity. In addition to primates, especially humans, a variety of other mammals may be treated according to the method of the present invention. The present invention is directed to a compound of the present invention or a pharmaceutically acceptable salt thereof that could be useful in medicine. The present invention may further be directed to a use of a compound of the present invention or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for antagonizing orexin receptor activity or treating the disorders and diseases noted herein in humans and animals.

The subject treated in the present methods is generally a mammal, such as a human being, male or female. The term "therapeutically effective amount" means the amount of the subject compound that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other

clinician. It is recognized that one skilled in the art may affect the neurological and psychiatric disorders by treating a patient presently afflicted with the disorders or by prophylactically treating a patient afflicted with the disorders with an effective amount of the compound of the present invention. As used herein, the terms "treatment" and "treating" refer to all processes wherein there may be a slowing, interrupting, arresting, controlling, or stopping of the progression of the neurological and psychiatric disorders described herein, but does not necessarily indicate a total elimination of all disorder symptoms, as well as the prophylactic therapy of the mentioned conditions, particularly in a patient who is predisposed to such disease or disorder. The terms "administration of" and or "administering a" compound should be understood to mean providing a compound of the invention or a prodrug of a compound of the invention to the individual in need thereof.

The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. Such term in relation to pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of the present invention and a pharmaceutically acceptable carrier. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The utility of the compounds in accordance with the present invention as orexin receptor OX1R and/or OX2R antagonists may be readily determined without undue experimentation by methodology well known in the art, including the "FLIPR Ca^{2+} Flux Assay" (Okumura et al., Biochem. Biophys. Res. Comm. 280:976-981, 2001). In a typical experiment the OX1 and OX2 receptor antagonistic activity of the compounds of the present invention was determined in accordance with the following experimental method. For intracellular calcium measurements, Chinese hamster ovary (CHO) cells expressing the rat orexin-1 receptor or the human orexin-2 receptor, are grown in Iscove's modified DMEM containing 2 mM L-glutamine, 0.5 g/ml G418, 1% hypoxanthine-thymidine supplement, 100 U/ml penicillin, 100 $\mu\text{g/ml}$ streptomycin and 10 % heat-inactivated fetal calf serum (FCS). The cells are seeded at 20,000 cells / well into Becton-Dickinson black 384-well clear bottom sterile plates coated with poly-D-

lysine. All reagents were from GIBCO-Invitrogen Corp. The seeded plates are incubated overnight at 37°C and 5% CO₂. Ala-6,12 human orexin-A as the agonist is prepared as a 1 mM stock solution in 1% bovine serum albumin (BSA) and diluted in assay buffer (HBSS containing 20 mM HEPES, 0.1% BSA and 2.5mM probenecid, pH7.4) for use in the assay at a final concentration of 70pM. Test compounds are prepared as 10 mM stock solution in DMSO, then diluted in 384-well plates, first in DMSO, then assay buffer. On the day of the assay, cells are washed 3 times with 100 µl assay buffer and then incubated for 60 min (37° C, 5% CO₂) in 60 µl assay buffer containing 1 µM Fluo-4AM ester, 0.02 % pluronic acid, and 1 % BSA. The dye loading solution is then aspirated and cells are washed 3 times with 100 µl assay buffer. 30 µl of that same buffer is left in each well. Within the Fluorescent Imaging Plate Reader (FLIPR, Molecular Devices), test compounds are added to the plate in a volume of 25 µl, incubated for 5 min and finally 25 µl of agonist is added. Fluorescence is measured for each well at 1 second intervals for 5 minutes and the height of each fluorescence peak is compared to the height of the fluorescence peak induced by 70 pM Ala-6,12 orexin-A with buffer in place of antagonist. For each antagonist, IC₅₀ value (the concentration of compound needed to inhibit 50 % of the agonist response) is determined. Alternatively, compound potency can be assessed by a radioligand binding assay (described in Bergman et. al. Bioorg. Med. Chem. Lett. 2008, 18, 1425 - 1430) in which the inhibition constant (K_i) is determined in membranes prepared from CHO cells expressing either the OX1 or OX2 receptor. The intrinsic orexin receptor antagonist activity of a compound which may be used in the present invention may be determined by these assays.

All of the final compounds of the following examples had activity in antagonizing the human orexin-2 receptor in the aforementioned assays with an IC₅₀ of about 0.1 nM to 100 nM. All of the final compounds of the following examples had activity in the radioligand binding assay with a K_i of about 0.1 nM to 100 nM against the orexin-2 receptor. All of the final compounds of the following examples had activity in the FLIPR assay with an IC₅₀ of about 0.1 nM to 100 nM against the orexin-2 receptor. Additional data is provided in the following Examples. Such a result is indicative of the intrinsic activity of the compounds in use as antagonists of orexin-1 receptor and/or the orexin-2 receptor. In general, one of ordinary skill in the art would appreciate that a substance is considered to effectively antagonize the orexin receptor if it has an IC₅₀ of less than about 50 µM, preferably less than about 100 nM. With respect to other piperidine compounds such as those disclosed in WO 2010/048012, it would be desirable that the present compounds exhibit unexpected properties, such as increased selectivity at the orexin-2 receptor relative to the orexin-1 receptor. For example, relative to certain compounds of WO2010/048012 that do not possess a 2-pyridyloxy-3-substituted-4-nitrile ring,

the compounds of the examples possess greater selectivity for the orexin-2 receptor than for the orexin-1 receptor.

The orexin receptors have been implicated in a wide range of biological functions. This has suggested a potential role for these receptors in a variety of disease processes in humans or other species. The compounds of the present invention could therefore potentially have utility in treating, preventing, ameliorating, controlling or reducing the risk of a variety of neurological and psychiatric disorders associated with orexin receptors, including one or more of the following conditions or diseases: sleep disorders, sleep disturbances, including enhancing sleep quality, improving sleep quality, increasing sleep efficiency, augmenting sleep maintenance; increasing the value which is calculated from the time that a subject sleeps divided by the time that a subject is attempting to sleep; improving sleep initiation; decreasing sleep latency or onset (the time it takes to fall asleep); decreasing difficulties in falling asleep; increasing sleep continuity; decreasing the number of awakenings during sleep; decreasing intermittent wakings during sleep; decreasing nocturnal arousals; decreasing the time spent awake following the initial onset of sleep; increasing the total amount of sleep; reducing the fragmentation of sleep; altering the timing, frequency or duration of REM sleep bouts; altering the timing, frequency or duration of slow wave (i.e. stages 3 or 4) sleep bouts; increasing the amount and percentage of stage 2 sleep; promoting slow wave sleep; enhancing EEG-delta activity during sleep; decreasing nocturnal arousals, especially early morning awakenings; increasing daytime alertness; reducing daytime drowsiness; treating or reducing excessive daytime sleepiness; increasing satisfaction with the intensity of sleep; increasing sleep maintenance; idiopathic insomnia; sleep problems; insomnia, hypersomnia, idiopathic hypersomnia, repeatability hypersomnia, intrinsic hypersomnia, narcolepsy, interrupted sleep, sleep apnea, wakefulness, nocturnal myoclonus, REM sleep interruptions, jet-lag, shift workers' sleep disturbances, dyssomnias, night terror, insomnias associated with depression, emotional/mood disorders, Alzheimer's disease or cognitive impairment, as well as sleep walking and enuresis, and sleep disorders which accompany aging; Alzheimer's sundowning; conditions associated with circadian rhythmicity as well as mental and physical disorders associated with travel across time zones and with rotating shift-work schedules, conditions due to drugs which cause reductions in REM sleep as a side effect; fibromyalgia; syndromes which are manifested by non-restorative sleep and muscle pain or sleep apnea which is associated with respiratory disturbances during sleep; conditions which result from a diminished quality of sleep; increasing learning; augmenting memory; increasing retention of memory; eating disorders associated with excessive food intake and complications associated therewith, compulsive eating disorders, obesity (due to any cause, whether genetic or environmental), obesity-related

disorders overeating, anorexia, bulimia, cachexia, dysregulated appetite control, hypertension, diabetes, elevated plasma insulin concentrations and insulin resistance, dyslipidemias, hyperlipidemia, endometrial, breast, prostate and colon cancer, osteoarthritis, obstructive sleep apnea, cholelithiasis, gallstones, heart disease, lung disease, abnormal heart rhythms and arrhythmias, myocardial infarction, congestive heart failure, coronary heart disease, acute and congestive heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ischemic or haemorrhagic stroke; subarachnoid haemorrhage; ulcers; allergies; benign prostatic hypertrophy; chronic renal failure; renal disease; impaired glucose tolerance; sudden death, polycystic ovary disease, craniopharyngioma, the Prader-Willi Syndrome, Frohlich's syndrome, GH-deficient subjects, normal variant short stature, Turner's syndrome, and other pathological conditions showing reduced metabolic activity or a decrease in resting energy expenditure as a percentage of total fat-free mass, e.g, children with acute lymphoblastic leukemia, metabolic syndrome, also known as syndrome X, insulin resistance syndrome, reproductive hormone abnormalities, sexual and reproductive dysfunction, such as impaired fertility, infertility, hypogonadism in males and hirsutism in females, fetal defects associated with maternal obesity, gastrointestinal motility disorders, intestinal motility dyskinesias, obesity-related gastro-esophageal reflux, hypothalamic diseases, hypophysis diseases, respiratory disorders, such as obesity-hypoventilation syndrome (Pickwickian syndrome), breathlessness, cardiovascular disorders, inflammation, such as systemic inflammation of the vasculature, arteriosclerosis, hypercholesterolemia, hyperuricaemia, lower back pain, gallbladder disease, gout, kidney cancer, increased anesthetic risk, reducing the risk of secondary outcomes of obesity, such as reducing the risk of left ventricular hypertrophy; diseases or disorders where abnormal oscillatory activity occurs in the brain, including depression, migraine, neuropathic pain, Parkinson's disease, psychosis and schizophrenia, as well as diseases or disorders where there is abnormal coupling of activity, particularly through the thalamus; enhancing cognitive function, including cognitive dysfunctions that comprise deficits in all types of attention, learning and memory functions occurring transiently or chronically in the normal, healthy, young, adult or aging population, and also occurring transiently or chronically in psychiatric, neurologic, cardiovascular and immune disorders; enhancing memory; increasing memory retention; increasing immune response; increasing immune function; hot flashes; night sweats; extending life span; schizophrenia; muscle-related disorders that are controlled by the excitation/relaxation rhythms imposed by the neural system such as cardiac rhythm and other disorders of the cardiovascular system; conditions related to proliferation of cells such as vasodilation or vasoconstriction and blood pressure; cancer; cardiac arrhythmia; hypertension; congestive heart failure; conditions of the genital/urinary system;

disorders of sexual function and fertility; adequacy of renal function; responsivity to anesthetics; mood disorders, such as depression or more particularly depressive disorders, for example, single episodic or recurrent major depressive disorders and dysthymic disorders, or bipolar disorders, for example, bipolar I disorder, bipolar II disorder and cyclothymic disorder, mood disorders due to a general medical condition, and substance-induced mood disorders; affective neurosis; depressive neurosis; anxiety neurosis; anxiety disorders including acute stress disorder, agoraphobia, generalized anxiety disorder, obsessive-compulsive disorder, panic attack, panic disorder, post-traumatic stress disorder, separation anxiety disorder, social phobia, specific phobia, substance-induced anxiety disorder and anxiety due to a general medical condition; acute neurological and psychiatric disorders such as cerebral deficits subsequent to cardiac bypass surgery and grafting, stroke, ischemic stroke, cerebral ischemia, spinal cord trauma, head trauma, perinatal hypoxia, cardiac arrest, hypoglycemic neuronal damage; Huntington's Chorea; Huntington's disease and Tourette syndrome; Cushing's syndrome/disease; basophile adenoma; prolactinoma; hyperprolactinemia; hypophysis tumor/adenoma; hypothalamic diseases; inflammatory bowel disease; gastric dyskinesia; gastric ulcers; Froehlich's syndrome; adrenohypophysis disease; hypophysis disease; adrenohypophysis hypofunction; adrenohypophysis hyperfunction; hypothalamic hypogonadism; Kallman's syndrome (anosmia, hyposmia); functional or psychogenic amenorrhea; hypopituitarism; hypothalamic hypothyroidism; hypothalamic-adrenal dysfunction; idiopathic hyperprolactinemia; hypothalamic disorders of growth hormone deficiency; idiopathic growth deficiency; dwarfism; gigantism; acromegaly; amyotrophic lateral sclerosis; multiple sclerosis; ocular damage; retinopathy; cognitive disorders; idiopathic and drug-induced Parkinson's disease; muscular spasms and disorders associated with muscular spasticity including tremors, epilepsy, convulsions, seizure disorders, absence seizures, complex partial and generalized seizures; Lennox-Gastaut syndrome; cognitive disorders including dementia (associated with Alzheimer's disease, ischemia, trauma, vascular problems or stroke, HIV disease, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeldt-Jacob disease, perinatal hypoxia, other general medical conditions or substance abuse); delirium, amnesic disorders or age related cognitive decline; schizophrenia or psychosis including schizophrenia (paranoid, disorganized, catatonic or undifferentiated), schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition and substance-induced psychotic disorder; dissociative disorders including multiple personality syndromes and psychogenic amnesias; substance-related disorders, substance use, substance abuse, substance seeking, substance reinstatement, all types of psychological and physical addictions and addictive behaviors, reward-related behaviors (including substance-

induced delirium, persisting dementia, persisting amnestic disorder, psychotic disorder or anxiety disorder; tolerance, addictive feeding, addictive feeding behaviors, binge/purge feeding behaviors, dependence, withdrawal or relapse from substances including alcohol, amphetamines, cannabis, cocaine, hallucinogens, inhalants, morphine, nicotine, opioids, phencyclidine, 5 sedatives, hypnotics or anxiolytics); appetite, taste, eating or drinking disorders; movement disorders, including akinesias and akinetic-rigid syndromes (including Parkinson's disease, drug-induced parkinsonism, postencephalitic parkinsonism, progressive supranuclear palsy, multiple system atrophy, corticobasal degeneration, parkinsonism-ALS dementia complex and basal ganglia calcification), chronic fatigue syndrome, fatigue, including Parkinson's fatigue, 10 multiple sclerosis fatigue, fatigue caused by a sleep disorder or a circadian rhythm disorder, medication-induced parkinsonism (such as neuroleptic-induced parkinsonism, neuroleptic malignant syndrome, neuroleptic-induced acute dystonia, neuroleptic-induced acute akathisia, neuroleptic-induced tardive dyskinesia and medication-induced postural tremor), Gilles de la Tourette's syndrome, epilepsy, and dyskinesias [including tremor (such as rest tremor, essential 15 tremor, postural tremor and intention tremor), chorea (such as Sydenham's chorea, Huntington's disease, benign hereditary chorea, neuroacanthocytosis, symptomatic chorea, drug-induced chorea and hemiballism), myoclonus (including generalised myoclonus and focal myoclonus), tics (including simple tics, complex tics and symptomatic tics), restless leg syndrome and dystonia (including generalised dystonia such as idiopathic dystonia, drug-induced dystonia, 20 symptomatic dystonia and paroxymal dystonia, and focal dystonia such as blepharospasm, oromandibular dystonia, spasmodic dysphonia, spasmodic torticollis, axial dystonia, dystonic writer's cramp and hemiplegic dystonia); neurodegenerative disorders including nosological entities such as disinhibition-dementia-parkinsonism-amyotrophy complex; pallido-ponto-nigral degeneration; epilepsy; seizure disorders; attention deficit/hyperactivity disorder (ADHD); 25 conduct disorder; migraine (including migraine headache); headache; hyperalgesia; pain; enhanced or exaggerated sensitivity to pain such as hyperalgesia, causalgia, and allodynia; acute pain; burn pain; atypical facial pain; neuropathic pain; back pain; complex regional pain syndrome I and II; arthritic pain; sports injury pain; pain related to infection e.g. HIV, post-chemotherapy pain; post-stroke pain; post-operative pain; neuralgia; emesis, nausea, vomiting; 30 gastric dyskinesia; gastric ulcers; Kallman's syndrome (anosmia); asthma; cancer; conditions associated with visceral pain such as irritable bowel syndrome, and angina; eating disorders; urinary incontinence; substance tolerance, substance withdrawal (including, substances such as opiates, nicotine, tobacco products, alcohol, benzodiazepines, cocaine, sedatives, hypnotics, etc.); psychosis; schizophrenia; anxiety (including generalized anxiety disorder, panic disorder, 35 and obsessive compulsive disorder); mood disorders (including depression, mania, bipolar

disorders); trigeminal neuralgia; hearing loss; tinnitus; neuronal damage including ocular damage; retinopathy; macular degeneration of the eye; emesis; brain edema; pain, including acute and chronic pain states, severe pain, intractable pain, inflammatory pain, neuropathic pain, post-traumatic pain, bone and joint pain (osteoarthritis), repetitive motion pain, dental pain, cancer pain, myofascial pain (muscular injury, fibromyalgia), perioperative pain (general surgery, gynecological), chronic pain, neuropathic pain, post-traumatic pain, trigeminal neuralgia, migraine and migraine headache and other diseases related to general orexin system dysfunction.

Thus, in certain embodiments the present invention may provide methods for: enhancing the quality of sleep; augmenting sleep maintenance; increasing REM sleep; increasing stage 2 sleep; decreasing fragmentation of sleep patterns; treating insomnia and all types of sleep disorders; treating or controlling sleep disturbances associated with diseases such as neurological disorders including neuropathic pain and restless leg syndrome; treating or controlling addiction disorders; treating or controlling psychoactive substance use and abuse; enhancing cognition; increasing memory retention; treating or controlling obesity; treating or controlling diabetes and appetite, taste, eating, or drinking disorders; treating or controlling hypothalamic diseases; treating or controlling depression; treating, controlling, ameliorating or reducing the risk of epilepsy, including absence epilepsy; treating or controlling pain, including neuropathic pain; treating or controlling Parkinson's disease; treating or controlling psychosis; treating or controlling dysthymic, mood, psychotic and anxiety disorders; treating or controlling depression, including major depression and major depression disorder; treating or controlling bipolar disorder; or treating, controlling, ameliorating or reducing the risk of schizophrenia, in a mammalian patient in need thereof which comprises administering to the patient a therapeutically effective amount of a compound of the present invention.

The subject compounds could further be of potential use in a method for the prevention, treatment, control, amelioration, or reduction of risk of the diseases, disorders and conditions noted herein. The dosage of active ingredient in the compositions of this invention may be varied, however, it is necessary that the amount of the active ingredient be such that a suitable dosage form is obtained. The active ingredient may be administered to patients (animals and human) in need of such treatment in dosages that will provide optimal pharmaceutical efficacy. The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment. The dose will vary from patient to patient depending upon the nature and severity of disease, the patient's weight, special diets then being followed by a patient, concurrent medication, and other factors which those skilled in the art will recognize. Generally, dosage levels of between 0.0001 to 10 mg/kg. of body weight daily are

administered to the patient, e.g., humans and elderly humans, to obtain effective antagonism of orexin receptors. The dosage range will generally be about 0.5 mg to 1.0 g. per patient per day which may be administered in single or multiple doses. In one embodiment, the dosage range will be about 0.5 mg to 500 mg per patient per day; in another embodiment about 0.5 mg to 200 mg per patient per day; and in yet another embodiment about 5 mg to 50 mg per patient per day. Pharmaceutical compositions of the present invention may be provided in a solid dosage formulation such as comprising about 0.5 mg to 500 mg active ingredient, or comprising about 1 mg to 250 mg active ingredient. The pharmaceutical composition may be provided in a solid dosage formulation comprising about 1 mg, 5 mg, 10 mg, 25 mg, 30 mg, 50 mg, 80 mg, 100 mg, 200 mg or 250 mg active ingredient. For oral administration, the compositions may be provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, such as 1, 5, 10, 15, 20, 25, 50, 75, 100, 150, 200, 250, 300, 400, 500, 600, 750, 800, 900, and 1000 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, such as once or twice per day. The compounds may be administered before bedtime. For example, the compounds may be administered about 1 Hour prior to bedtime, about 30 minutes prior to bedtime or immediately before bedtime.

The compounds of the present invention may be used in combination with one or more other drugs in the treatment, prevention, control, amelioration, or reduction of risk of diseases or conditions for which compounds of the present invention or the other drugs may have utility, where the combination of the drugs together are safer or more effective than either drug alone. Such other drug(s) may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of the present invention. When a compound of the present invention is used contemporaneously with one or more other drugs, a pharmaceutical composition in unit dosage form containing such other drugs and the compound of the present invention is contemplated. However, the combination therapy may also include therapies in which the compound of the present invention and one or more other drugs are administered on different overlapping schedules. It is also contemplated that when used in combination with one or more other active ingredients, the compounds of the present invention and the other active ingredients may be used in lower doses than when each is used singly. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition to a compound of the present invention. The above combinations include combinations of a compound of the present invention not only with one other active compound, but also with two or more other active compounds.

Likewise, compounds of the present invention may be used in combination with other drugs that are used in the prevention, treatment, control, amelioration, or reduction of risk of the diseases or conditions for which compounds of the present invention are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of the present invention. When a compound of the present invention is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of the present invention is contemplated. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients, in addition to a compound of the present invention.

The weight ratio of the compound of the present invention to the second active ingredient may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the present invention is combined with another agent, the weight ratio of the compound of the present invention to the other agent will generally range from about 1000:1 to about 1:1000, such as about 200:1 to about 1:200. Combinations of a compound of the present invention and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used. In such combinations the compound of the present invention and other active agents may be administered separately or in conjunction. In addition, the administration of one element may be prior to, concurrent to, or subsequent to the administration of other agent(s).

The compounds of the present invention may be administered in combination with other compounds which are known in the art to be useful for enhancing sleep quality and preventing and treating sleep disorders and sleep disturbances, including e.g., sedatives, hypnotics, anxiolytics, antipsychotics, antianxiety agents, antihistamines, benzodiazepines, barbiturates, cyclopyrrolones, GABA agonists, 5HT-2 antagonists including 5HT-2A antagonists and 5HT-2A/2C antagonists, histamine antagonists including histamine H3 antagonists, histamine H3 inverse agonists, imidazopyridines, minor tranquilizers, melatonin agonists and antagonists, melatonergic agents, other orexin antagonists, orexin agonists, prokineticin agonists and antagonists, pyrazolopyrimidines, T-type calcium channel antagonists, triazolopyridines, and the like, such as: adinazolam, allobarbitol, alonimid, alprazolam, amitriptyline, amobarbital, amoxapine, armodafinil, APD-125, bentazepam, benzocetamine, brotizolam, bupropion, busprione, butabarbital, butalbital, capromorelin, capuride, carbocloral, chloral betaine, chloral hydrate, chlordiazepoxide, clomipramine, clonazepam, cloperidone, clorazepate, clorethate, clozapine, conazepam, cyprazepam, desipramine, dexclamol, diazepam, dichloralphenazone,

divalproex, diphenhydramine, doxepin, EMD-281014, eplivanserin, estazolam, eszopiclone, ethchlorinol, etomidate, fenobam, flunitrazepam, flurazepam, fluvoxamine, fluoxetine, fosazepam, gaboxadol, glutethimide, halazepam, hydroxyzine, ibutamoren, imipramine, indiplon, lithium, lorazepam, lormetazepam, LY-156735, maprotiline, MDL-100907, 5 mecloqualone, melatonin, mephobarbital, meprobamate, methaqualone, methypylon, midaflur, midazolam, modafinil, nefazodone, NGD-2-73, nisobamate, nitrazepam, nortriptyline, ornortriptyline, oxazepam, paraldehyde, paroxetine, pentobarbital, perlapine, perphenazine, phenelzine, phenobarbital, prazepam, promethazine, propofol, protriptyline, quazepam, ramelteon, reclazepam, roletamide, secobarbital, sertraline, suproclone, TAK-375, temazepam, 10 thioridazine, tiagabine, tracazolate, tranlycypromaine, trazodone, triazolam, trepipam, tricetamide, triclofos, trifluoperazine, trimetozine, trimipramine, uldazepam, venlafaxine, zaleplon, zolazepam, zopiclone, zolpidem, and salts thereof, and combinations thereof, and the like, or the compound of the present invention may be administered in conjunction with the use of physical methods such as with light therapy or electrical stimulation.

15 In another embodiment, the subject compound may be employed in combination with other compounds which are known in the art, either administered separately or in the same pharmaceutical compositions, including, but are not limited to: insulin sensitizers including (i) PPAR γ antagonists such as glitazones (e.g. ciglitazone; darglitazone; englitazone; isaglitazone (MCC-555); pioglitazone; rosiglitazone; troglitazone; tularik; BRL49653; CLX-0921; 5-BTSD), 20 GW-0207, LG-100641, and LY-300512, and the like); (iii) biguanides such as metformin and phenformin; (b) insulin or insulin mimetics, such as biota, LP-100, novarapid, insulin detemir, insulin lispro, insulin glargine, insulin zinc suspension (lente and ultralente); Lys-Pro insulin, GLP-1 (73-7) (insulintropin); and GLP-1 (7-36)-NH₂); (c) sulfonylureas, such as acetohexamide; chlorpropamide; diabinese; glibenclamide; glipizide; glyburide; glimepiride; 25 gliclazide; glipentide; gliquidone; glisohamide; tolazamide; and tolbutamide; (d) α -glucosidase inhibitors, such as acarbose, adiposine; camiglibose; emiglitate; miglitol; voglibose; pradimicin-Q; salbostatin; CKD-711; MDL-25,637; MDL-73,945; and MOR 14, and the like; (e) cholesterol lowering agents such as (i) HMG-CoA reductase inhibitors (atorvastatin, itavastatin, fluvastatin, lovastatin, pravastatin, rivastatin, rosuvastatin, simvastatin, and other statins), (ii) bile acid 30 absorbers/sequestrants, such as cholestyramine, colestipol, dialkylaminoalkyl derivatives of a cross-linked dextran; Colestid®; LoCholest®, and the like, (ii) nicotiny alcohol, nicotinic acid or a salt thereof, (iii) proliferator-activator receptor α agonists such as fenofibric acid derivatives (gemfibrozil, clofibrate, fenofibrate and benzafibrate), (iv) inhibitors of cholesterol absorption such as stanol esters, beta-sitosterol, sterol glycosides such as tiqueside; and azetidinones such as 35 ezetimibe, and the like, and (acyl CoA:cholesterol acyltransferase (ACAT)) inhibitors such as

avasimibe, and melinamide, (v) anti-oxidants, such as probucol, (vi) vitamin E, and (vii) thyromimetics; (f) PPAR α agonists such as bezafibrate, benzaifibrate, ciprofibrate, clofibrate, etofibrate, fenofibrate, and gemfibrozil; and other fibric acid derivatives, such as Atromid®, Lipid® and Tricor®, and the like, and PPAR α agonists as described in WO 97/36579; (g) PPAR δ agonists, such as those disclosed in WO97/28149; (h) PPAR α/δ agonists, such as muraglitazar, and the compounds disclosed in US 6,414,002; (i) anti-obesity agents, such as (1) growth hormone secretagogues, growth hormone secretagogue receptor agonists/antagonists, such as NN703, hexarelin, MK-0677, SM-130686, CP-424,391, L-692,429, and L-163,255, and such as those disclosed in U.S. Patent Nos. 5,536,716, and 6,358,951, U.S. Patent Application Nos. 2002/049196 and 2002/022637, and PCT Application Nos. WO 01/56592 and WO 02/32888; (2) protein tyrosine phosphatase-1B (PTP-1B) inhibitors; (3) cannabinoid receptor ligands, such as cannabinoid CB₁ receptor antagonists or inverse agonists, such as rimonabant, taranabant, AMT-251, and SR-14778 and SR 141716A (Sanofi Synthelabo), SLV-319 (Solvay), BAY 65-2520 (Bayer) and those disclosed in U.S. Patent Nos. 5,532,237, 4,973,587, 5,013,837, 5,081,122, 5,112,820, 5,292,736, 5,624,941, 6,028,084, PCT Application Nos. WO 96/33159, WO 98/33765, WO98/43636, WO98/43635, WO 01/09120, WO98/31227, WO98/41519, WO98/37061, WO00/10967, WO00/10968, WO97/29079, WO99/02499, WO 01/58869, WO 01/64632, WO 01/64633, WO 01/64634, W002/076949, WO 03/007887, WO 04/048317, and WO 05/000809; (4) anti-obesity serotonergic agents, such as fenfluramine, dexfenfluramine, phentermine, and sibutramine; (5) β 3-adrenoreceptor agonists, such as AD9677/TAK677 (Dainippon/Takeda), CL-316,243, SB 418790, BRL-37344, L-796568, BMS-196085, BRL-35135A, CGP12177A, BTA-243, Trecadrine, Zeneca D7114, SR 59119A; (6) pancreatic lipase inhibitors, such as orlistat (Xenical®), Triton WR1339, RHC80267, lipstatin, tetrahydrolipstatin, teasaponin, diethylumbelliferyl phosphate, and those disclosed in PCT Application No. WO 01/77094; (7) neuropeptide Y1 antagonists, such as BIBP3226, J-115814, BIBO 3304, LY-357897, CP-671906, GI-264879A, and those disclosed in U.S. Patent No. 6,001,836, and PCT Patent Publication Nos. WO 96/14307, WO 01/23387, WO 99/51600, WO 01/85690, WO 01/85098, WO 01/85173, and WO 01/89528; (8) neuropeptide Y5 antagonists, such as GW-569180A, GW-594884A, GW-587081X, GW-548118X, FR226928, FR 240662, FR252384, 1229U91, GI-264879A, CGP71683A, LY-377897, PD-160170, SR-120562A, SR-120819A and JCF-104, and those disclosed in U.S. Patent Nos. 6,057,335; 6,043,246; 6,140,354; 6,166,038; 6,180,653; 6,191,160; 6,313,298; 6,335,345; 6,337,332; 6,326,375; 6,329,395; 6,340,683; 6,388,077; 6,462,053; 6,649,624; and 6,723,847, European Patent Nos. EP-01010691, and EP-01044970; and PCT International Patent Publication Nos. WO 97/19682, WO 97/20820, WO 97/20821, WO 97/20822, WO 97/20823, WO 98/24768; WO 98/25907; WO 98/25908; WO

98/27063, WO 98/47505; WO 98/40356; WO 99/15516; WO 99/27965; WO 00/64880, WO 00/68197, WO 00/69849, WO 01/09120, WO 01/14376; WO 01/85714, WO 01/85730, WO 01/07409, WO 01/02379, WO 01/02379, WO 01/23388, WO 01/23389, WO 01/44201, WO 01/62737, WO 01/62738, WO 01/09120, WO 02/22592, WO 0248152, and WO 02/49648; WO 5 02/094825; WO 03/014083; WO 03/10191; WO 03/092889; WO 04/002986; and WO 04/031175; (9) melanin-concentrating hormone (MCH) receptor antagonists, such as those disclosed in WO 01/21577 and WO 01/21169; (10) melanin-concentrating hormone 1 receptor (MCH1R) antagonists, such as T-226296 (Takeda), and those disclosed in PCT Patent Application Nos. WO 01/82925, WO 01/87834, WO 02/051809, WO 02/06245, WO 02/076929, 10 WO 02/076947, WO 02/04433, WO 02/51809, WO 02/083134, WO 02/094799, WO 03/004027; (11) melanin-concentrating hormone 2 receptor (MCH2R) agonist/antagonists; (12) orexin receptor antagonists, such as SB-334867-A, and those disclosed in patent publications herein; (13) serotonin reuptake inhibitors such as fluoxetine, paroxetine, and sertraline; (14) melanocortin agonists, such as Melanotan II; (15) Mc4r (melanocortin 4 receptor) agonists, such 15 as CHIR86036 (Chiron), ME-10142, and ME-10145 (Melacure), CHIR86036 (Chiron); PT-141, and PT-14 (Palatin); (16) 5HT-2 agonists; (17) 5HT2C (serotonin receptor 2C) agonists, such as BVT933, DPCA37215, WAY161503, R-1065, and those disclosed in U.S. Patent No. 3,914,250, and PCT Application Nos. WO 02/36596, WO 02/48124, WO 02/10169, WO 01/66548, WO 02/44152, WO 02/51844, WO 02/40456, and WO 02/40457; (18) galanin antagonists; (19) CCK 20 agonists; (20) CCK-A (cholecystokinin-A) agonists, such as AR-R 15849, GI 181771, JMV-180, A-71378, A-71623 and SR14613, and those disclosed in U.S. Patent No. 5,739,106; (21) GLP-1 agonists; (22) corticotropin-releasing hormone agonists; (23) histamine receptor-3 (H3) modulators; (24) histamine receptor-3 (H3) antagonists/inverse agonists, such as hioperamide, 3-(1H-imidazol-4-yl)propyl N-(4-pentenyl)carbamate, clobenpropit, iodophenpropit, imoproxifan, 25 GT2394 (Gliatech), and O-[3-(1H-imidazol-4-yl)propanol]-carbamates; (25) β -hydroxy steroid dehydrogenase-1 inhibitors (β -HSD-1); (26) PDE (phosphodiesterase) inhibitors, such as theophylline, pentoxifylline, zaprinast, sildenafil, amrinone, milrinone, cilostamide, rolipram, and cilomilast; (27) phosphodiesterase-3B (PDE3B) inhibitors; (28) NE (norepinephrine) transport inhibitors, such as GW 320659, despiramine, talsupram, and nomifensine; (29) ghrelin 30 receptor antagonists, such as those disclosed in PCT Application Nos. WO 01/87335, and WO 02/08250; (30) leptin, including recombinant human leptin (PEG-OB, Hoffman La Roche) and recombinant methionyl human leptin (Amgen); (31) leptin derivatives; (32) BRS3 (bombesin receptor subtype 3) agonists such as [D-Phe6,beta-Ala11,Phe13,Nle14]Bn(6-14) and [D-Phe6,Phe13]Bn(6-13)propylamide, and those compounds disclosed in Pept. Sci. 2002 Aug; 8(8): 35 461-75); (33) CNTF (Ciliary neurotrophic factors), such as GI-181771 (Glaxo-SmithKline),

SR146131 (Sanofi Synthelabo), butabindide, PD170,292, and PD 149164 (Pfizer); (34) CNTF derivatives, such as axokine (Regeneron); (35) monoamine reuptake inhibitors, such as sibutramine; (36) UCP-1 (uncoupling protein-1), 2, or 3 activators, such as phytanic acid, 4-[(E)-2-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1-propenyl]benzoic acid (TTNPB),
 5 retinoic acid; (37) thyroid hormone β agonists, such as KB-2611 (KaroBioBMS); (38) FAS (fatty acid synthase) inhibitors, such as Cerulenin and C75; (39) DGAT1 (diacylglycerol acyltransferase 1) inhibitors; (40) DGAT2 (diacylglycerol acyltransferase 2) inhibitors; (41) ACC2 (acetyl-CoA carboxylase-2) inhibitors; (42) glucocorticoid antagonists; (43) acyl-estrogens, such as oleoyl-estrone, disclosed in del Mar-Grasa, M. et al., Obesity Research,
 10 9:202-9 (2001); (44) dipeptidyl peptidase IV (DP-IV) inhibitors, such as isoleucine thiazolidide, valine pyrrolidide, NVP-DPP728, LAF237, P93/01, TSL 225, TMC-2A/2B/2C, FE 999011, P9310/K364, VIP 0177, SDZ 274-444, sitagliptin; and the compounds disclosed in US 6,699,871, WO 03/004498; WO 03/004496; EP 1 258 476; WO 02/083128; WO 02/062764; WO 03/000250; WO 03/002530; WO 03/002531; WO 03/002553; WO 03/002593; WO 03/000180;
 15 and WO 03/000181; (46) dicarboxylate transporter inhibitors; (47) glucose transporter inhibitors; (48) phosphate transporter inhibitors; (49) Metformin (Glucophage®); (50) Topiramate (Topimax®); (50) peptide YY, PYY 3-36, peptide YY analogs, derivatives, and fragments such as BIM-43073D, BIM-43004C (Olitvak, D.A. et al., Dig. Dis. Sci. 44(3):643-48 (1999)); (51) Neuropeptide Y2 (NPY2) receptor agonists such NPY3-36, N acetyl [Leu(28,31)] NPY 24-36,
 20 TASP-V, and cyclo-(28/32)-Ac-[Lys28-Glu32]-(25-36)-pNPY; (52) Neuropeptide Y4 (NPY4) agonists such as pancreatic peptide (PP), and other Y4 agonists such as 1229U91; (54) cyclooxygenase-2 inhibitors such as etoricoxib, celecoxib, valdecoxib, parecoxib, lumiracoxib, BMS347070, tiracoxib or JTE522, ABT963, CS502 and GW406381; (55) Neuropeptide Y1 (NPY1) antagonists such as BIBP3226, J-115814, BIBO 3304, LY-357897, CP-671906, GI-
 25 264879A; (56) Opioid antagonists such as nalmefene (Revex ®), 3-methoxynaltrexone, naloxone, naltrexone; (57) 11 β HSD-1 (11-beta hydroxy steroid dehydrogenase type 1) inhibitors such as BVT 3498, BVT 2733, and those disclosed in WO 01/90091, WO 01/90090, WO 01/90092, US 6,730,690 and US 2004-0133011; (58) aminorex; (59) amphetamine; (60) amphetamine; (61) benzphetamine; (62) chlorphentermine; (63) clobenzorex; (64) cloforex; (65) clominorex; (66) clortermine; (67) cyclexedrine; (68) dextroamphetamine; (69) diphenmethoxidine, (70) N-ethylamphetamine; (71) fenbutrazate; (72) fenisorex; (73) fenproporex; (74) fludorex; (75) fluminorex; (76) furfurylmethylamphetamine; (77) levamfetamine; (78) levophacetoperane; (79) mefenorex; (80) metamfepramone; (81) methamphetamine; (82) norpseudoephedrine; (83) pentorex; (84) phendimetrazine; (85) phenmetrazine; (86) picilorex; (87) phytopharm 57; and (88) zonisamide., (89) neuromedin U

and analogs or derivatives thereof, (90) oxyntomodulin and analogs or derivatives thereof, and (91) Neurokinin-1 receptor antagonists (NK-1 antagonists) such as the compounds disclosed in: U.S. Patent Nos. 5,162,339, 5,232,929, 5,242,930, 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, and 5,637,699.

5 In another embodiment, the subject compound may be employed in combination with an anti-depressant or anti-anxiety agent, including norepinephrine reuptake inhibitors (including tertiary amine tricyclics and secondary amine tricyclics), selective serotonin reuptake inhibitors (SSRIs), monoamine oxidase inhibitors (MAOIs), reversible inhibitors of monoamine oxidase (RIMAs), serotonin and noradrenaline reuptake inhibitors (SNRIs), corticotropin releasing factor (CRF) antagonists, α -adrenoreceptor antagonists, neurokinin-1 receptor antagonists, atypical anti-depressants, benzodiazepines, 5-HT_{1A} agonists or antagonists, especially 5-HT_{1A} partial agonists, and corticotropin releasing factor (CRF) antagonists. Specific agents include: amitriptyline, clomipramine, doxepin, imipramine and trimipramine; amoxapine, desipramine, maprotiline, nortriptyline and protriptyline; citalopram, duloxetine, 10 fluoxetine, fluvoxamine, paroxetine and sertraline; isocarboxazid, phenelzine, tranlycypromine and selegiline; moclobemide; venlafaxine; aprepitant; bupropion, lithium, nefazodone, trazodone and viloxazine; alprazolam, chlordiazepoxide, clonazepam, chlorazepate, diazepam, halazepam, 15 lorazepam, oxazepam and prazepam; buspirone, flesinoxan, gepirone and ipsapirone, and pharmaceutically acceptable salts thereof.

20 In another embodiment, the subject compound may be employed in combination with anti-Alzheimer's agents; beta-secretase inhibitors; gamma-secretase inhibitors; growth hormone secretagogues; recombinant growth hormone; HMG-CoA reductase inhibitors; NSAID's including ibuprofen; vitamin E; anti-amyloid antibodies; CB-1 receptor antagonists or CB-1 receptor inverse agonists; antibiotics such as doxycycline and rifampin; N-methyl-D- 25 aspartate (NMDA) receptor antagonists, such as memantine; cholinesterase inhibitors such as galantamine, rivastigmine, donepezil, and tacrine; growth hormone secretagogues such as ibutamoren, ibutamoren mesylate, and capromorelin; histamine H₃ antagonists; AMPA agonists; PDE IV inhibitors; GABA_A inverse agonists; or neuronal nicotinic agonists.

30 In another embodiment, the subject compound may be employed in combination with sedatives, hypnotics, anxiolytics, antipsychotics, antianxiety agents, cyclopyrrolones, imidazopyridines, pyrazolopyrimidines, minor tranquilizers, melatonin agonists and antagonists, melatonergic agents, benzodiazepines, barbiturates, 5HT-2 antagonists, and the like, such as: adinazolam, allobarbitol, alonimid, alprazolam, amitriptyline, amobarbital, amoxapine, bentazepam, benzoctamine, brotizolam, bupropion, buspirone, butabarbital, butalbital, capuride, 35 carbocloral, chloral betaine, chloral hydrate, chlordiazepoxide, clomipramine, clonazepam,

cloperidone, clorazepate, clorethate, clozapine, cyprazepam, desipramine, dexclamol, diazepam, dichloralphenazone, divalproex, diphenhydramine, doxepin, estazolam, ethchlorvynol, etomidate, fenobam, flunitrazepam, flurazepam, fluvoxamine, fluoxetine, fosazepam, glutethimide, halazepam, hydroxyzine, imipramine, lithium, lorazepam, lormetazepam, maprotiline, mecloqualone, melatonin, mephobarbital, meprobamate, methaqualone, midafur, midazolam, nefazodone, nisobamate, nitrazepam, nortriptyline, oxazepam, paraldehyde, paroxetine, pentobarbital, perlapine, perphenazine, phenelzine, phenobarbital, prazepam, promethazine, propofol, protriptyline, quazepam, reclazepam, roletamide, secobarbital, sertraline, suproclone, temazepam, thioridazine, tracazolate, tranlycypromaine, trazodone, triazolam, trepipam, tricetamide, triclofos, trifluoperazine, trimetozine, trimipramine, uldazepam, venlafaxine, zaleplon, zolazepam, zolpidem, and salts thereof, and combinations thereof, and the like, or the subject compound may be administered in conjunction with the use of physical methods such as with light therapy or electrical stimulation.

In another embodiment, the subject compound may be employed in combination with levodopa (with or without a selective extracerebral decarboxylase inhibitor such as carbidopa or benserazide), anticholinergics such as biperiden (optionally as its hydrochloride or lactate salt) and trihexyphenidyl (benzhexol) hydrochloride, COMT inhibitors such as entacapone, MOA-B inhibitors, antioxidants, A2a adenosine receptor antagonists, cholinergic agonists, NMDA receptor antagonists, serotonin receptor antagonists and dopamine receptor agonists such as alantemol, bromocriptine, fenoldopam, lisuride, naxagolide, pergolide and pramipexole.

In another embodiment, the subject compound may be employed in combination with acetophenazine, alantemol, benzhexol, bromocriptine, biperiden, chlorpromazine, chlorprothixene, clozapine, diazepam, fenoldopam, fluphenazine, haloperidol, levodopa, levodopa with benserazide, levodopa with carbidopa, lisuride, loxapine, mesoridazine, molindolone, naxagolide, olanzapine, pergolide, perphenazine, pimozide, pramipexole, risperidone, sulpiride, tetrabenazine, trihexyphenidyl, thioridazine, thiothixene or trifluoperazine.

In another embodiment, the subject compound may be employed in combination with a compound from the phenothiazine, thioxanthene, heterocyclic dibenzazepine, butyrophenone, diphenylbutylpiperidine and indolone classes of neuroleptic agent. Suitable examples of phenothiazines include chlorpromazine, mesoridazine, thioridazine, acetophenazine, fluphenazine, perphenazine and trifluoperazine. Suitable examples of thioxanthenes include chlorprothixene and thiothixene. An example of a dibenzazepine is clozapine. An example of a butyrophenone is haloperidol. An example of a diphenylbutylpiperidine is pimozide. An

example of an indolone is molindolone. Other neuroleptic agents include loxapine, sulpiride and risperidone.

In another embodiment, the subject compound may be employed in combination with a nicotine agonist or a nicotine receptor partial agonist such as varenicline, opioid antagonists (e.g., naltrexone (including naltrexone depot), antabuse, and nalmefene), dopaminergic agents (e.g., apomorphine), ADD/ADHD agents (e.g., methylphenidate hydrochloride (e.g., Ritalin® and Concerta®), atomoxetine (e.g., Strattera®), a monoamine oxidase inhibitor (MAOI), amphetamines (e.g., Adderall®)) and anti-obesity agents, such as apo-B/MTP inhibitors, 11Beta-hydroxy steroid dehydrogenase-1 (11Beta-HSD type 1) inhibitors, peptide YY3-36 or analogs thereof, MCR-4 agonists, CCK-A agonists, monoamine reuptake inhibitors, sympathomimetic agents, β 3 adrenergic receptor agonists, dopamine receptor agonists, melanocyte-stimulating hormone receptor analogs, 5-HT2c receptor agonists, melanin concentrating hormone receptor antagonists, leptin, leptin analogs, leptin receptor agonists, galanin receptor antagonists, lipase inhibitors, bombesin receptor agonists, neuropeptide-Y receptor antagonists (e.g., NPY Y5 receptor antagonists), thyromimetic agents, dehydroepiandrosterone or analogs thereof, glucocorticoid receptor antagonists, other orexin receptor antagonists, glucagon-like peptide-1 receptor agonists, ciliary neurotrophic factors, human agouti-related protein antagonists, ghrelin receptor antagonists, histamine 3 receptor antagonists or inverse agonists, and neuromedin U receptor agonists, and pharmaceutically acceptable salts thereof.

In another embodiment, the subject compound may be employed in combination with an anorectic agent such as aminorex, amphetamine, benzphetamine, chlorphentermine, clobenzorex, cloforex, clominox, clortermine, cyclohexedrine, dexfenfluramine, dextroamphetamine, diethylpropion, diphenmethoxidine, N-ethylamphetamine, fenbutrazate, fenfluramine, fenisorex, fenproporex, fludorex, fluminorex, furfurylmethylamphetamine, levamfetamine, levophacetoperane, mazindol, mefenorex, metamfepramone, methamphetamine, norpseudoephedrine, pentorex, phendimetrazine, phenmetrazine, phentermine, phenylpropanolamine, picilorex and sibutramine; selective serotonin reuptake inhibitor (SSRI); halogenated amphetamine derivatives, including chlorphentermine, cloforex, clortermine, dexfenfluramine, fenfluramine, picilorex and sibutramine; and pharmaceutically acceptable salts thereof.

In another embodiment, the subject compound may be employed in combination with an opiate agonist, a lipooxygenase inhibitor, such as an inhibitor of 5-lipoxygenase, a cyclooxygenase inhibitor, such as a cyclooxygenase-2 inhibitor, an interleukin inhibitor, such as an interleukin-1 inhibitor, an NMDA antagonist, an inhibitor of nitric oxide or an inhibitor of the

synthesis of nitric oxide, a non-steroidal antiinflammatory agent, or a cytokine-suppressing antiinflammatory agent, for example with a compound such as acetaminophen, aspirin, codiene, fentanyl, ibuprofen, indomethacin, ketorolac, morphine, naproxen, phenacetin, piroxicam, a steroidal analgesic, sufentanyl, sunlindac, tenidap, and the like. Similarly, the subject compound
5 may be administered with a pain reliever; a potentiator such as caffeine, an H₂-antagonist, simethicone, aluminum or magnesium hydroxide; a decongestant such as phenylephrine, phenylpropanolamine, pseudophedrine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxy-ephedrine; an antiitussive such as codeine, hydrocodone, caramiphen, carbetapentane, or dextramethorphan; a diuretic; and a sedating or
10 non-sedating antihistamine.

The compounds of the present invention may be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implant), by inhalation spray, nasal, vaginal, rectal, sublingual, or topical routes of administration and may be formulated, alone or together, in suitable dosage unit
15 formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration. In addition to the treatment of warm-blooded animals such as mice, rats, horses, cattle, sheep, dogs, cats, monkeys, etc., the compounds of the invention may be effective for use in humans.

The pharmaceutical compositions for the administration of the compounds of this
20 invention may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both,
25 and then, if necessary, shaping the product into the desired formulation. In the pharmaceutical composition the active object compound is included in an amount sufficient to produce the desired effect upon the process or condition of diseases. As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the
30 specified ingredients in the specified amounts.

Pharmaceutical compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide
35 pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in


admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. Compositions for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil. Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Oily suspensions may be formulated by suspending the active ingredient in a suitable oil. Oil-in-water emulsions may also be employed. Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Pharmaceutical compositions of the present compounds may be in the form of a sterile injectable aqueous or oleagenous suspension. The compounds of the present invention may also be administered in the form of suppositories for rectal administration. For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of the present invention may be employed. The compounds of the present invention may also be formulated for administered by inhalation. The compounds of the present invention may also be administered by a transdermal patch by methods known in the art.


Several methods for preparing the compounds of this invention are illustrated in the following Schemes and Examples. Starting materials are made according to procedures known in the art (e.g. PCT Patent Publications WO2001/68609, WO2004/085403, WO2005/118548, WO2008/147518, WO2009/143033 and WO2010/048012) or as illustrated herein. The following abbreviations are used herein: Me: methyl; Et: ethyl; t-Bu: *tert*-butyl; Ar: aryl; Ph: phenyl; Bn: benzyl; Ac: acetyl; Boc: *tert*-butoxy carbonyl; BSA: bovine serum albumin; CbzCl: benzylchloroformate; CDI: carbonyl diimidazole; DCM: dichloromethane; DCE: dichloroethane; DEAD: diethylazodicarboxylate; DIPEA: N,N-diisopropylethylamine; DMF: N,N-dimethylformamide; DMSO: dimethylsulfoxide; CH₂Cl₂: dichloromethane; EDC: N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide; Et₃N: triethylamine; EtOAc: ethylacetate; EtOH: ethanol; HCl: hydrogen chloride; HOAt: 1-hydroxy-7-aza-benzotriazole; HOBt: hydroxybenzotriazole hydrate; HPLC: high performance liquid chromatography; Hunig's base:

N,N-diisopropylethylamine; MeOH: methanol; MgSO₄: magnesium sulfate; MTBE: methyl tert-butyl ether; NaHCO₃: sodium bicarbonate; NaOH: sodium hydroxide; NMM: N-methylmorpholine; PyClu: 1-(chloro-1-pyrrolidinylmethylene)-pyrrolidinium hexafluorophosphate; rt: room temperature; SOCl₂: thionyl chloride; T3P: 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphorinane-2,4,6-trioxide; THF: tetrahydrofuran; TFA: trifluoroacetic acid. The compounds of the present invention can be prepared in a variety of fashions.

In some cases the final product may be further modified, for example, by manipulation of substituents. These manipulations may include, but are not limited to, reduction, oxidation, alkylation, acylation, and hydrolysis reactions which are commonly known to those skilled in the art. In some cases the order of carrying out the foregoing reaction schemes may be varied to facilitate the reaction or to avoid unwanted reaction products. The following examples are provided so that the invention might be more fully understood. These examples are illustrative only and should not be construed as limiting the invention in any way.

EXAMPLE 1





2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-Triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile

Step 1: (2*R*,5*S*)-Benzyl 2-methyl-5-((methanesulfonyl)oxy)piperidine-1-carboxylate (2)

5 A solution of (2*R*,5*S*)-benzyl 5-hydroxy-2-methylpiperidine-1-carboxylate (15.0 g, 60.2 mmol), 4-dimethylamino pyridine (0.74 g, 6.0 mmol), and CH_2Cl_2 (401 mL) was treated with DIEA (15.8 mL, 90.0 mmol) and cooled at 0 °C. Methanesulfonyl chloride (5.9 mL, 75 mmol) in CH_2Cl_2 (67 mL) was added dropwise and stirring continued at 0 °C for 2 h. The reaction was diluted with water and 1 N HCl and the separated organics were washed with brine,
10 dried over MgSO_4 , filtered, concentrated and dried, providing the crude title compound which was used without further purification. LRMS m/z (M+H) 328.3 found, 328.1 required.

Step 2: (2*R*,5*R*)-Benzyl 5-azido-2-methylpiperidine-1-carboxylate (3)

15 A solution of crude (2*R*,5*S*)-benzyl 2-methyl-5-((methanesulfonyl)oxy)piperidine-1-carboxylate (19.7 g, 60.2 mmol) and sodium azide (6.65 g, 102 mmol) in DMF (86 mL) was heated to 85 °C overnight. The reaction was diluted with EtOAc, washed 2x with water, 1x with brine, dried over magnesium sulfate, filtered, and concentrated, providing the crude title compound which was used without further purification. LRMS m/z (M+H) 275.3 found, 275.2 required.

20

Step 3: (2*R*,5*R*)-Benzyl 5-amino-2-methylpiperidine-1-carboxylate (4)

25 To a solution of (2*R*,5*R*)-benzyl 5-azido-2-methylpiperidine-1-carboxylate (15.2 g, 55.4 mmol) in THF (111 mL) was added trimethylphosphine (1.0 M in toluene, 55.4 mL, 55.4 mmol) dropwise over 20 min. The reaction was quenched with water (30 mL) and stirred at RT overnight. The mixture was diluted with dichloromethane, washed 2x with water, 1x with brine, dried over magnesium sulfate, filtered, and concentrated, providing the crude title compound as an orange oil which was used without further purification. LRMS m/z (M+H) 249.6 found, 249.3 required.

Step 4: (2*R*,5*R*)-Benzyl-5-((*tert*-butoxycarbonyl)amino-2-methylpiperidine-1-carboxylate (5)

A solution of (2*R*,5*R*)-Benzyl 5-amino-2-methylpiperidine-1-carboxylate (3.08 g, 12.4 mmol) and 4-dimethylaminopyridine (0.015 g, 0.12 mmol) in dichloromethane (62 mL) was treated with a solution of di-*tert*-butyl dicarbonate (2.91 mL, 12.5 mmol) in dichloromethane (20 mL) dropwise and stirred at RT overnight. The reaction was concentrated *in vacuo*, providing the crude title compound which was used without further purification. LRMS *m/z* (M+H) 349.4 found, 349.3 required.

Step 5: *tert*-Butyl ((3*R*,6*R*)-6-methylpiperidin-3-yl)carbamate (6)

A solution of (2*R*,5*R*)-benzyl-5-((*tert*-butoxycarbonyl)amino-2-methylpiperidine-1-carboxylate (4.32 g, 12.4 mmol) and palladium (10 wt % on activated carbon, 0.300 g, 2.82 mmol) in degassed ethanol (100 mL) and methanol (5 mL) was stirred for 72 h under an atmosphere of hydrogen gas. Additional palladium (10 wt % on activated carbon, 0.10 g, 0.94 mmol) was added, and the reaction was recharged with hydrogen gas and allowed to stir at RT overnight. The degassed mixture was then filtered over celite, washing with EtOH. The filtrate was concentrated to give the titled compound as a crude, light yellow solid which was used without further purification.

Step 6: *tert*-Butyl((3*R*,6*R*)-1-(2-(2*H*-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)carbamate (7)

A solution of 2-(2*H*-1,2,3-triazol-2-yl)benzoic acid (4.24 g, 22.40 mmol), *tert*-butyl ((3*R*,6*R*)-6-methylpiperidin-3-yl)carbamate (3.20 g, 14.93 mmol), 1-hydroxy-7-azabenzotriazole (4.06 g, 29.9 mmol), EDC (5.72 g, 29.9 mmol), and DIEA (15.65 mL, 90 mmol) in DMF (100 mL) was heated at 50 °C overnight. The reaction was diluted with saturated, aqu. NaHCO₃ and extracted 2x with EtOAc. The organics were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel gradient chromatography (0-80% EtOAc in hexanes), providing the title compound. LRMS *m/z* (M+H) 386.4 found, 386.2 required.

Step 7: (2-(2*H*-1,2,3-Triazol-2-yl)phenyl)((2*R*,5*R*)-5-amino-2-methylpiperidin-1-yl)methanone (8)

A solution of *tert*-butyl((3*R*,6*R*)-1-(2-(2*H*-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)carbamate (3.6 g, 9.3 mmol) in EtOAc (78 mL) was saturated with HCl (g) and stirred for 2 h. The reaction was concentrated and the residue basified with 1 N NaOH. The mixture was extracted 3x with EtOAc, then the organics were washed with brine, dried over

MgSO₄, filtered, and concentrated. The residue was purified by silica gel gradient chromatography (0-20% MeOH in CH₂Cl₂), providing the title compound. LRMS m/z (M+H) 286.3 found, 286.3 required.


5 Step 8: 2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile

A solution of (2-(2*H*-1,2,3-triazol-2-yl)phenyl)((2*R*,5*R*)-5-amino-2-methylpiperidin-1-yl)methanone (400 mg, 1.40 mmol) 2-fluoroisonicotinonitrile (342 mg, 2.80 mmol), and cesium carbonate (776 mg, 2.383 mmol) in DMSO (4673 μ L) was heated at 50 °C
 10 for 48 h. The reaction was cooled, diluted with water and saturated aqueous sodium bicarbonate, and extracted 2x with EtOAc. The combined organic fractions were washed with brine, dried over magnesium sulfate, filtered, and concentrated. The residue was purified by silica gel gradient chromatography (0-50% EtOAc in hexanes), providing the title compound. LRMS m/z (M+H) 388.4 found, 388.2 required.

15


TABLE 1

The following compound was prepared using the foregoing methodology and general procedure described in Example 1, but substituting the appropriate electrophile for 2-fluoroisonicotinonitrile (Step 8), as described in the foregoing Reaction Schemes and Examples.
 20 The requisite starting materials were commercially available, described in the literature or readily synthesized by one skilled in the art of organic synthesis from commercially available reagents using conventional reactions without undue experimentation.

Example	Structure	Name	HRMS
2		3-methoxy-2-([(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1-([2-(2 <i>H</i> -1,2,3-triazol-2-yl)phenyl]carbonyl)piperidin-3-yl]amino)pyridine-4-carbonitrile	C ₂₂ H ₂₃ N ₇ O ₂ [M+H] Calc'd 418.1988, found 418.1970

25

EXAMPLE 3



2-(((3*R*,6*R*)-1-(2-(2*H*-Tetrazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile

5 Step 1: (2*R*,5*R*)-benzyl 5-((4-cyanopyridin-2-yl)amino)-2-methylpiperidine-1-carboxylate (**9**)

A solution of (2*R*,5*R*)-benzyl 5-amino-2-methylpiperidine-1-carboxylate (150 mg, 0.604 mmol), 2-fluoroisonicotinonitrile (148 mg, 1.208 mmol), and cesium carbonate (335 mg, 1.027 mmol) in DMSO (2 mL) was heated at 80 °C overnight. The reaction was cooled, diluted with water and saturated aqueous sodium bicarbonate, and extracted 2x with EtOAc. The combined organic fractions were washed with brine, dried over MgSO_4 , filtered, and concentrated. The residue was purified by silica gel gradient chromatography (0-25% EtOAc in hexanes), providing the title compound. LRMS m/z ($M+H$) 351.4 found, 351.2 required.


Step 2: 2-(((3*R*,6*R*)-6-methylpiperidin-3-yl)amino)isonicotinonitrile (**10**)

15 A solution of (2*R*,5*R*)-benzyl 5-((4-cyanopyridin-2-yl)amino)-2-methylpiperidine-1-carboxylate (89 mg, 0.254 mmol) in CH_2Cl_2 (5.0 mL) was treated, at RT, with dimethylsulfide (507 μL , 6.86 mmol), then boron trifluoride etherate (321 μL , 1.22 mmol). After stirring overnight, the reaction was poured into 10% aq. NH_4OH and extracted 2x with CH_2Cl_2 . The combined organic fractions were washed with brine, dried over MgSO_4 , filtered, and concentrated, providing the crude title compound. LRMS m/z ($M+H$) 217.2 found, 217.1 required.

Step 3: 2-(((3*R*,6*R*)-1-(2-(2*H*-tetrazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile

A solution of 2-(((3*R*,6*R*)-6-methylpiperidin-3-yl)amino)isonicotinonitrile (55 mg, 0.254 mmol), 2-(2*H*-tetrazol-2-yl)benzoic acid (58.0 mg, 0.305 mmol), 1-hydroxy-7-azabenzotriazole (69.2 mg, 0.509 mmol), and EDC (97 mg, 0.509 mmol) in DMF (2.5 mL) was treated with DIEA (266 μ L, 1.526 mmol) and heated at 50 °C overnight. The reaction was cooled, diluted with water and saturated aqueous NaHCO₃, and extracted 2x with EtOAc. The combined organic fractions were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel gradient chromatography (0-85% EtOAc in hexanes), providing the title compound. HRMS *m/z* (M+H) 389.1818 found, 389.1833 required.

EXAMPLE 4



2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-Triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)-3-methylisonicotinonitrile

Step 1: (2*R*,5*R*)-benzyl 5-((4-cyano-3-methylpyridin-2-yl)amino)-2-methylpiperidine-1-carboxylate (11)

Prepared as per Example 3, Step 1, substituting 2-fluoro-3-methylisonicotinonitrile as the electrophile. The residue was purified by silica gel gradient chromatography (0-70% EtOAc in hexanes), providing the title compound. LRMS m/z (M+H) 365.4 found, 365.2 required.

Step 2: 3-methyl-2-(((3*R*,6*R*)-6-methylpiperidin-3-yl)amino)isonicotinonitrile (12)


A solution of (2*R*,5*R*)-benzyl 5-((4-cyano-3-methylpyridin-2-yl)amino)-2-methylpiperidine-1-carboxylate (31 mg, 0.085 mmol) in EtOH (1.7 mL) was evacuated 4x and backfilled with nitrogen each time. Palladium on carbon (6.3 mg, 6.0 μ mol) was added, and the reaction was evacuated and backfilled 2x from a hydrogen balloon. After stirring overnight, additional palladium on carbon was added and the reaction was again purged with hydrogen, and stirring continued for several more hours. The reaction was evacuated and backfilled with nitrogen, then diluted with EtOH and filtered through Celite. The filtrate was concentrated and dried to provide the title compound. LRMS m/z (M+H) 231.3 found, 231.2 required.

Step 3: 2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-Triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)-3-methylisonicotinonitrile


Prepared as per Example 3, Step 3, substituting 2-(2*H*-1,2,3-triazol-2-yl)benzoic acid as the carboxylic acid. The residue was purified by silica gel gradient chromatography (0-65% EtOAc in hexanes), then further purified by reversed-phase chromatography (10-75% MeCN in H₂O, 0.1% TFA), providing the title compound. HRMS m/z (M+H) 402.2024 found, 402.2037 required.

TABLE 2

The following compound was prepared using the foregoing methodology and general procedure described in Example 4, substituting the appropriate carboxylic acid for 2-(2*H*-1,2,3-triazol-2-yl)benzoic acid (Step 3), as described in the foregoing Reaction Schemes and Examples. The requisite starting materials were commercially available, described in the literature or readily synthesized by one skilled in the art of organic synthesis from commercially available reagents using conventional reactions without undue experimentation.

Example	Structure	Name	HRMS
5		3-methyl-2-{{[(3R,6R)-6-methyl-1-{{2-(2H-tetrazol-2-yl)phenyl}-carbonyl}-piperidin-3-yl]amino}pyridine-4-carbonitrile	C ₂₁ H ₂₂ N ₈ O [M+H] Calc'd 403.1992, found 403.1979

EXAMPLE 6



2-((3R,6R)-1-(2-(2H-1,2,3-Triazol-2-yl)thiophene-3-carbonyl)-6-methylpiperidin-3-ylamino)-3-methoxyisonicotinonitrile

Step 1: (2R,5R)-benzyl 5-(4-cyano-3-methoxypyridin-2-ylamino)-2-methylpiperidine-1-carboxylate

A mixture of 5-amino-2-methyl-piperidine-1-carboxylic acid benzyl ester (800 mg, 3.22 mmol), 2-chloro-3-methoxy-isonicotinonitrile (815 mg, 4.83 mmol), Pd(dppf)Cl₂ (235 mg, 0.32 mmol) and NaOtBu (618 mg, 6.44 mmol) in anhydrous THF (100 mL) was heated at 80 °C with stirring under nitrogen for 16 hr. The mixture was cooled to room temperature and concentrated *in vacuo*. The residue was purified by column chromatography (0-12.5% ethyl

acetate in petroleum ether), providing the title compound as yellow oil. LRMS m/z (M+H) 381.2 found, 381.18 required.

Step 2: 2-((3*R*,6*R*)-6-methylpiperidin-3-ylamino)-3-methoxypyridine-4-carbonitrile

To a solution of (2*R*,5*R*)-benzyl 5-(4-cyano-3-methoxypyridin-2-ylamino)-2-methylpiperidine-1-carboxylate (350mg, 0.92 mmol) in MeOH (25 mL) was added Pd/C (10%, 100 mg) at room temperature. The resulting mixture was stirred under hydrogen for 1 h. The mixture was filtered and the filtrate was concentrated *in vacuo*, providing the title compound as yellow oil. LRMS m/z (M+H) 247.2 found, 247.15 required

Step 3: 2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-triazol-2-yl)thiophene-3-carbonyl)-6-methylpiperidin-3-yl)amino)-3-methoxyisonicotinonitrile


To a solution of 2-((3*R*,6*R*)-6-methylpiperidin-3-ylamino)-3-methoxypyridine-4-carbonitrile (45 mg, 0.18 mmol) in anhydrous dichloromethane (20 mL) was added triethylamine (28 mg, 0.28 mmol) at ~5 °C. After stirring for 15 min, a solution of 2-[1,2,3]triazol-2-yl-thiophene-3-carbonyl chloride (prepared from 48.8 mg of 2-[1,2,3]triazol-2-yl-thiophene-3-carboxylic acid) in anhydrous CH₂Cl₂ (5 mL) was added via syringe at ~5 °C. The resulting mixture was stirred at 25 °C for 16 h. The mixture was concentrated *in vacuo* and purified by prep TLC (50% ethyl acetate in petroleum ether), providing the title compound as pale white solid. LRMS m/z (M+H) 424.2 found, 424.15 required.

TABLE 3

The following compound was prepared using the foregoing methodology and general procedure described in Example 6, substituting the appropriate acid for 2-(2*H*-1,2,3-triazol-2-yl)thiophene-3-carboxylic acid (Step 3), as described in the foregoing Reaction Schemes and Examples. The requisite starting materials were commercially available, described in the literature or readily synthesized by one skilled in the art of organic synthesis from commercially available reagents using conventional reactions without undue experimentation.

Example	Structure	Name	LRMS
7		3-methoxy-2-(((3 <i>R</i> ,6 <i>R</i>)-6-methyl-1-(2-(pyrimidin-2-yl)thiophene-3-carbonyl)piperidin-3-ylamino)isonicotinonitrile	C ₂₂ H ₂₂ N ₆ O ₂ S [M+H] calc 435.15 obs 435.1

EXAMPLE 8



- 5 2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-Triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)(methylamino)isonicotinonitrile



A solution of 2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile (Example 1, 8.3 mg, 0.021 mmol) in DMF (1 mL) was treated with sodium hydride (0.5 mg, 0.02 mmol) at RT. After a few mins, iodomethane (1.3 μ L, 0.021 mmol) was added. After 30 mins, additional NaH and iodomethane were added to complete the conversion. The reaction was quenched with H₂O, then diluted with saturated aqueous NaHCO₃, and extracted 2x with EtOAc. The combined organic fractions were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by reversed-phase chromatography (10-80% MeCN in H₂O, 0.1% TFA), providing the title compound. HRMS *m/z* (M+H) 402.2048 found, 402.2037 required.

TABLE 4


The following compounds were prepared using the foregoing methodology and general procedure described in Example 8, substituting the appropriate alkyl halide for iodomethane, as described in the foregoing Reaction Schemes and Examples. The requisite starting materials were commercially available, described in the literature or readily synthesized

by one skilled in the art of organic synthesis from commercially available reagents using conventional reactions without undue experimentation.

Example	Structure	Name	LRMS/HRMS
9		2-{methyl[(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1-{[2-(2 <i>H</i> -tetrazol-2-yl)phenyl]-carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile	C ₂₁ H ₂₂ N ₈ O [M+H] Calc'd 403.1989, found 403.1986
10		3-methoxy-2-{methyl[(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1-{[2-(2 <i>H</i> -1,2,3-triazol-2-yl)thiophen-3-yl]carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile	C ₂₁ H ₂₄ N ₇ O ₂ S [M+H] Calc'd 438.2, found 438.2
11		3-methoxy-2-(methyl{(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1-[(2-pyrimidin-2-ylthiophen-3-yl)carbonyl]piperidin-3-yl}amino)pyridine-4-carbonitrile	C ₂₃ H ₂₄ N ₆ O ₂ S [M+H] calc 449.17 found 449.2
12		2-{ethyl[(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1-{[2-(2 <i>H</i> -1,2,3-triazol-2-yl)phenyl]carbonyl}piperidin-3-yl]amino}-pyridine-4-carbonitrile	C ₂₃ H ₂₅ N ₇ O [M+H] Calc'd 416.2200, found 416.2193

13		2-[[[(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1- {[2-(2 <i>H</i> -1,2,3-triazol-2- yl)phenyl]carbonyl}piper- idin-3-yl](prop-2-en-1- yl)amino}pyridine-4- carbonitrile	C ₂₄ H ₂₅ N ₇ O [M+H] Calc'd 428.2193, found 428.2193
14		2-[[[(3 <i>R</i> ,6 <i>R</i>)-6-methyl-1- {[2-(2 <i>H</i> -1,2,3-triazol-2- yl)phenyl]carbonyl}piper- idin-3-yl](propyl)- amino}pyridine-4- carbonitrile	C ₂₄ H ₂₇ N ₇ O [M+H] Calc'd 430.2350, found 430.2338

EXAMPLE 15



N-((3*R*,6*R*)-1-(2-(2*H*-1,2,3-Triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)-N-(4-cyanopyridin-2-yl)acetamide

A solution of 2-(((3*R*,6*R*)-1-(2-(2*H*-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile (35 mg, 0.090 mmol) in CH₂Cl₂ (900 μL) was treated with DIEA (79 μL, 0.45 mmol) and acetyl chloride (16 μL, 0.23 mmol) at RT. After stirring overnight, additional DIEA and acetyl chloride were added and the reaction was heated at 35 °C for 8 h.


The reaction was quenched by addition of saturated, aq. NH₄Cl, then diluted with saturated aqueous NaHCO₃, and extracted 2x with EtOAc. The combined organic fractions were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by silica gel gradient chromatography (0-100% EtOAc in hexanes), providing the title compound. HRMS m/z (M+H) 403.1979 found, 403.1986 required.

The following table shows representative data for the compounds of the Examples as orexin receptor antagonists as determined by the foregoing assays wherein the hOX2R Ki is a measure of the ability of the test compound to inhibit the action of the human orexin-2 receptor and the hOX1R Ki is a measure of the ability of the test compound to inhibit the action of the rat orexin-1 receptor. The following table shows representative data for the compounds of the Examples as orexin receptor antagonists as determined by the assays herein.

Example	hOX2R FLIPR IC ₅₀ (nM)	hOX1R FLIPR IC ₅₀ (nM)
1	29	1742
2	11	267
3	9	1922
4	7	745
5	7	1149
6	11	827
7	47	384
8	20	2048
9	32	2236
10	39	> 10,000
11	34	> 10,000
12	23	1990
13	23	2548
14	13	330
15	1781	> 10,000


With respect to other piperidine compounds such as those disclosed in WO 2010/048012, it would be desirable that the present compounds exhibit unexpected properties, such as increased selectivity at the orexin-2 receptor relative to the orexin-1 receptor. For example, relative to certain compounds of WO2010/048012 that do not possess a 2-pyridylthio-piperidine, the compounds of the examples possess greater selectivity for the orexin-2 receptor than for the orexin-1 receptor.

For example, the following compounds are disclosed in WO 2010/048012:




Example 1-3

hOX2R IC₅₀ = 32 nM
hOX1R IC₅₀ = 375 nM
Selectivity **12x**




Example 1-4

hOX2R IC₅₀ = 47 nM
hOX1R IC₅₀ = 180 nM
Selectivity **3.8x**



Example 1-13 (F-2)


hOX2R IC₅₀ = 15 nM
hOX1R IC₅₀ = 16 nM
Selectivity **1x**




Example 1-19

hOX2R IC₅₀ = 26 nM
hOX1R IC₅₀ = 48 nM
Selectivity **1.8x**


Representative compounds herein are the compounds of Example 1, 3 and 4.



Example 1



Example 3



Example 4

hOX2R IC₅₀ = 29 nM

hOX2R IC₅₀ = 9 nM

hOX2R IC₅₀ = 7 nM

hOX1R IC₅₀ = 1742 nM

hOX1R IC₅₀ = 1992 nM

hOX1R IC₅₀ = 745 nM

Selectivity **60x**

Selectivity **221x**


Selectivity **106x**

As indicated by the data herein, the compounds of the present examples provide greater functional selectivity for the orexin-2 receptor over the orexin-1 receptor. The distinction in potency between the orexin-2 receptor and the orexin-1 receptor in the whole cell FLIPR functional assay provides enhanced predictive value for determining in vivo efficacy. Increasing the functional selectivity for the orexin-2 receptor reduces the potential for dual receptor antagonism in vivo. Such greater functional selectivity may provide benefits over other orexin receptor antagonists that are known in the art.

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various adaptations, changes, modifications, substitutions, deletions, or additions of procedures and protocols may be made without departing from the spirit and scope of the invention.

WHAT IS CLAIMED IS:

1. A compound of the formula I:



I

wherein:

A is selected from the group consisting of phenyl, naphthyl and heteroaryl;

R^{1a}, R^{1b} and R^{1c} are independently selected from the group consisting of:

- (1) hydrogen,
- (2) halogen,
- (3) hydroxyl,
- (4) $-(C=O)_m-O_n-C_{1-6}alkyl$, where m is 0 or 1, n is 0 or 1 (wherein if m is 0 or n is 0, a bond is present) and where the alkyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (5) $-(C=O)_m-O_n-C_{3-6}cycloalkyl$, where the cycloalkyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (6) $-(C=O)_m-C_{2-4}alkenyl$, where the alkenyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (7) $-(C=O)_m-C_{2-4}alkynyl$, where the alkynyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (8) $-(C=O)_m-O_n-phenyl$ or $-(C=O)_m-O_n-naphthyl$, where the phenyl or naphthyl is unsubstituted or substituted with one or more substituents selected from R⁴,
- (9) $-(C=O)_m-O_n-heterocycle$, where the heterocycle is unsubstituted or substituted with one or more substituents selected from R⁴,
- (10) $-(C=O)_m-NR^{10}R^{11}$, wherein R¹⁰ and R¹¹ are independently selected from the group consisting of:
 - (a) hydrogen,
 - (b) C₁₋₆alkyl, which is unsubstituted or substituted with R⁴,

- 5 (c) C₃-6alkenyl, which is unsubstituted or substituted with R⁴,
 (d) C₃-6alkynyl, which is unsubstituted or substituted with R⁴,
 (e) C₃-6cycloalkyl which is unsubstituted or substituted with R⁴,
 (f) phenyl, which is unsubstituted or substituted with R⁴, and
 (g) heterocycle, which is unsubstituted or substituted with R⁴,
 (11) -S(O)₂-NR¹⁰R¹¹,
 (12) -S(O)_q-R¹², where q is 0, 1 or 2 and where R¹² is selected from the definitions of
 R¹⁰ and R¹¹,
 (13) -CO₂H,
 10 (14) -CN, and
 (15) -NO₂;

R³ is selected from C₁-6alkyl and C₃-6cycloalkyl, which is unsubstituted or substituted with one or more substituents selected from R⁴;

15

R⁴ is selected from the group consisting of:

- (1) hydroxyl,
 (2) halogen,
 (3) C₁-6alkyl,
 20 (4) -C₃-6cycloalkyl,
 (5) -O-C₁-6alkyl,
 (6) -O(C=O)-C₁-6alkyl,
 (7) -NH₂,
 (7) -NH-C₁-6alkyl,
 25 (8) -NO₂,
 (9) phenyl,
 (10) heterocycle,
 (11) -CO₂H, and
 (12) -CN;

30

R⁵ is selected from the group consisting of:

- (1) hydrogen,
 (2) halogen,
 (3) hydroxyl,


- (4) C₁-6alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,
- (5) C₃-6cycloalkyl, which is unsubstituted or substituted with C₁-6alkyl, halogen, hydroxyl or phenyl,
- 5 (6) -O-C₁-6alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl, and
- (7) -S-C₁-6alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl;

10 R⁶ is selected from the group consisting of:

- (1) hydrogen,
- (2) C₁-6alkyl, which is unsubstituted or substituted with oxo, halogen, hydroxyl or phenyl,
- (3) C₂-6alkenyl, and
- 15 (4) -(CO)-O-C₁-6alkyl;

or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1 of the formula Ia:




20

Ia

or a pharmaceutically acceptable salt thereof.

3. The compound of Claim 1 of the formula Ia':




Ia'

or a pharmaceutically acceptable salt thereof.

5

4. The compound of Claim 1 of the formula Ib:



Ib

or a pharmaceutically acceptable salt thereof.

10

5. The compound of Claim 1 or a pharmaceutically acceptable salt thereof wherein A is selected from phenyl, pyridyl, thiophenyl, thiazolyl, isothiazolyl, and pyrazolyl.

6. The compound of Claim 1 or a pharmaceutically acceptable salt thereof wherein R^{1a}, R^{1b} and R^{1c} are independently selected from the group consisting of:

15

- (1) hydrogen,
- (2) halogen,
- (3) hydroxyl,
- (4) C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,

20

- (5) -O-C₁₋₆alkyl, which is unsubstituted or substituted with halogen, hydroxyl or phenyl,
(6) -CN, and
(7) heteroaryl, wherein heteroaryl is selected from triazolyl, tetrazolyl, oxazolyl, pyrrolyl, imidazolyl, indolyl, pyridyl, and pyrimidinyl, which is unsubstituted or substituted with halogen, hydroxyl, C₁₋₆alkyl, -O-C₁₋₆alkyl or -NO₂.

7. The compound of Claim 6 or a pharmaceutically acceptable salt thereof wherein R^{1c} is hydrogen, and R^{1a} and R^{1b} are independently selected from the group consisting of:

- (1) hydrogen,
(2) fluoro,
(3) chloro,
(4) bromo,
(5) methyl,
(6) ethyl,
(7) trifluoromethyl, and
(8) heteroaryl, wherein heteroaryl is selected from triazolyl, tetrazolyl, oxazolyl, pyrrolyl, imidazolyl, indolyl, pyridyl, and pyrimidinyl.

8. The compound of Claim 1 or a pharmaceutically acceptable salt thereof wherein R³ is methyl.

9. The compound of Claim 1 or a pharmaceutically acceptable salt thereof wherein R⁵ is selected from the group consisting of: hydrogen, methyl, cyclopropyl, chloro, -O-CH₃ and -S-CH₃.

10. The compound of Claim 1 or a pharmaceutically acceptable salt thereof wherein R⁶ is selected from the group consisting of:

- (1) hydrogen,
(2) C₁₋₆alkyl,
(3) C₂₋₆alkenyl, and
(4) -(CO)-O-C₁₋₆alkyl.

11. A compound which is selected from the group consisting of:

2-(((3R,6R)-1-(2-(2H-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile;

3-methoxy-2-{[(3R,6R)-6-methyl-1-{[2-(2H-1,2,3-triazol-2-yl)phenyl]carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile;

2-(((3R,6R)-1-(2-(2H-tetrazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)isonicotinonitrile;

2-(((3R,6R)-1-(2-(2H-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)amino)-3-methylisonicotinonitrile;

3-methyl-2-{[(3R,6R)-6-methyl-1-{[2-(2H-tetrazol-2-yl)phenyl]carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile;

2-(((3R,6R)-1-(2-(2H-1,2,3-triazol-2-yl)thiophene-3-carbonyl)-6-methylpiperidin-3-yl)amino)-3-methoxyisonicotinonitrile;

3-methoxy-2-((3R,6R)-6-methyl-1-(2-(pyrimidin-2-yl)thiophene-3-carbonyl)piperidin-3-yl)amino)isonicotinonitrile;

2-(((3R,6R)-1-(2-(2H-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)(methyl)amino)isonicotinonitrile;

2-{methyl[(3R,6R)-6-methyl-1-{[2-(2H-tetrazol-2-yl)phenyl]carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile;

3-methoxy-2-{methyl[(3R,6R)-6-methyl-1-{[2-(2H-1,2,3-triazol-2-yl)thiophen-3-yl]carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile;

3-methoxy-2-(methyl{(3R,6R)-6-methyl-1-[(2-pyrimidin-2-yl)thiophen-3-yl]carbonyl}piperidin-3-yl)amino)pyridine-4-carbonitrile;

2-{ethyl[(3R,6R)-6-methyl-1-{[2-(2H-1,2,3-triazol-2-yl)phenyl]carbonyl}piperidin-3-yl]amino}pyridine-4-carbonitrile;

2-{[(3R,6R)-6-methyl-1-{[2-(2H-1,2,3-triazol-2-yl)phenyl]carbonyl}piperidin-3-yl](prop-2-en-1-yl)amino}pyridine-4-carbonitrile;

2-{[(3R,6R)-6-methyl-1-{[2-(2H-1,2,3-triazol-2-yl)phenyl]carbonyl}piperidin-3-yl](propyl)amino}pyridine-4-carbonitrile;

N-((3R,6R)-1-(2-(2H-1,2,3-triazol-2-yl)benzoyl)-6-methylpiperidin-3-yl)-N-(4-cyanopyridin-2-yl)acetamide;

methyl 4-cyano-2-{[(3R,6R)-6-methyl-1-{[2-(2H-1,2,3-triazol-2-yl)phenyl]carbonyl}piperidin-3-yl]amino}pyridine-3-carboxylate; and

methyl 4-cyano-2-((3R,6R)-6-methyl-1-[(2-pyrimidin-2-yl)phenyl]carbonyl)piperidin-3-yl)amino)pyridine-3-carboxylate;

or a pharmaceutically acceptable salt thereof.

12. A pharmaceutical composition which comprises an inert carrier and a compound of Claim 1 or a pharmaceutically acceptable salt thereof.

5 13. A compound of Claim 1 or a pharmaceutically acceptable salt thereof for use in medicine.

14. Use of a compound of Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a sleep disorder.
10

15. A method for enhancing the quality of sleep in a mammalian patient in need thereof which comprises administering to the patient a therapeutically effective amount of the compound of Claim 1 or a pharmaceutically acceptable salt thereof.

15 16. A method for treating insomnia in a mammalian patient in need thereof which comprises administering to the patient a therapeutically effective amount of the compound of Claim 1 or a pharmaceutically acceptable salt thereof.

17. A method for treating or controlling obesity in a mammalian patient in need thereof which comprises administering to the patient a therapeutically effective amount of the compound of Claim 1 or a pharmaceutically acceptable salt thereof.
20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 13/71366

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/445 (2014.01)

USPC - 514/318

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): A61K 31/445 (2014.01)

USPC: 514/318

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC: 514/256; 514/277; 546/194; 544/333Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
PatBase, Google Scholar, PubWEST, SureChem
orexin receptor, antagonist, aminopyridine, isonictinonitrile, pyridin-4-nitrile, pyridin-4-carbonitrile

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2011/0201632 A1 (BRESLIN et al.) 18 August 2011 (18.08.2011) para [0184], [0229], Table 1; Table 2	1-17
Y	US 2009/0093478 A1 (BRANCH et al.) 09 April 2009 (09.04.2009) para [0308], Table; para [0355], [0361]	1-17
A, P	US 2013/0102619 A1 (KUDUK et al.) 25 April 2013 (25.04.2013) Entire Document	1-17

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 February 2014 (28.02.2014)

Date of mailing of the international search report

27 MAR 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774