

(19) **DANMARK**

(10) **DK/EP 3984996 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 D 239/48 (2006.01)** **A 61 K 31/395 (2006.01)** **A 61 K 31/505 (2006.01)**
A 61 K 31/715 (2006.01) **A 61 P 1/14 (2006.01)** **A 61 P 1/16 (2006.01)**
C 07 D 401/04 (2006.01) **C 07 D 403/04 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-10-21**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-10-09**
- (86) Europæisk ansøgning nr.: **21206067.7**
- (86) Europæisk indleveringsdag: **2017-08-04**
- (87) Den europæiske ansøgnings publiceringsdag: **2022-04-20**
- (30) Prioritet: **2016-08-12 WO PCT/CN2016/094833**
- (62) Stamansøgningsnr: **17838647.0**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285, USA**
- (72) Opfinder: **COATES, David Andrew, , Indianapolis, 46206-6288, USA**
QIN, Luo Heng, , Indianapolis, 46206-6288, USA
WEI, Yi, , Indianapolis, 46206-6288, USA
ZHOU, Jingye, , Indianapolis, 46206-6288, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Aminopyridin-SSAO-inhibitorer**
- (56) Fremdragne publikationer:
WO-A1-2013/163675
WO-A2-2007/120528
PETRA DUNKEL ET AL: "Semicarbazide-sensitive amine oxidase/vascular adhesion protein-1: a patent survey", EXPERT OPINION ON THERAPEUTIC PATENTS, vol. 21, no. 9, 15 June 2011 (2011-06-15), pages 1453 - 1471, XP055375413, ISSN: 1354-3776, DOI: 10.1517/13543776.2011.594040

DESCRIPTION

Description

[0001] This invention relates to amino pyrimidine compounds, pharmaceutically acceptable salts of the compounds, and therapeutic uses of the compound and salts.

[0002] Semicarbazide-sensitive amino oxidase/vascular adhesion protein-1 (SSAO)/VAP-1) is a member of the semicarbazide-sensitive amino oxidase family. SSAO/VAP-1 has been alternatively referred to as VAP-1 or SSAO. SSAO/VAP-1 is an enzyme that exists both as a membrane-bound and a soluble isoform; it is predominantly expressed from the endothelial cell surface, vascular smooth muscle and adipose cells. SSAO/VAP-1 participates in many cellular processes including glucose disposition, inflammation responses, and leukocyte recruitment. High activity levels of this enzyme are associated with diabetes, atherosclerosis, strokes, chronic kidney disease, and Alzheimer's disease, among other disorders. Recently SSAO/VAP-1 has been implicated in the pathogenesis of liver diseases such as fatty liver disease. Weston, C.J. et al., J Neural. Transm. 2011, 118, 1055-1064. Fatty liver disease (FLD) encompasses a spectrum of disease states where fat accumulates in the liver in excessive amounts and is accompanied by inflammation. FLD can lead to non-alcoholic fatty liver disease (NAFLD), which is characterized by insulin resistance. Unchecked NAFLD progresses to a persistent inflammatory response or non-alcoholic steatohepatitis (NASH), progressive liver fibrosis, and eventually to cirrhosis. Currently there is a need to provide alternative treatment therapies for liver diseases such as NAFLD and/or NASH.


[0003] It is thought that a SSAO/VAP-1 inhibitor will reduce liver inflammation and fibrosis and thereby provide a treatment for liver diseases, in particular, a treatment for NAFLD and/or NASH. The present invention provides compounds that inhibit the SSAO/VAP-1 enzyme and which may address one or more of the needs.

[0004] Dunkel et al. (Expert Opinion on Therapeutic Patents, vol. 21, no. 9, pp. 1453-1471) provides a patent survey for semicarbazide-sensitive amine oxidase/vascular adhesion protein-1.

[0005] WO 2007/120528 describes compositions and methods of using compositions for treatment of inflammatory diseases and immune disorders, including allylamino compounds which are inhibitors of SSAO and/or VAP-1.


[0006] WO 2013/163675 describes the preparation and pharmaceutical use of substituted 3-haloallylamine derivatives as SSAO/VAP-1 inhibitors.

[0007] A first aspect of the invention is a compound of the formula:




or a pharmaceutically acceptable salt thereof.

[0008] A second aspect of the invention is a compound of the formula:




or a pharmaceutically acceptable salt thereof.

[0009] A third aspect of the invention is a compound of the formula:



or a pharmaceutically acceptable salt thereof.

[0010] A fourth aspect of the invention is a compound of the formula:



or a pharmaceutically acceptable salt thereof.


[0011] A fifth aspect of the invention is a pharmaceutical composition comprising a compound according to any one of the first to fourth aspects and a pharmaceutically acceptable carrier, diluent or excipient.

[0012] A sixth aspect of the invention is a compound according to any one of the first to fourth aspects, or a pharmaceutical composition according to the fifth aspect, for use in therapy.

[0013] A seventh aspect of the invention is a compound according to any one of the first to fourth aspects, or a pharmaceutical composition according to the fifth aspect, for use in a method of treating a patient in need of treatment for non-alcoholic steatohepatitis wherein the method comprises administering to the patient an effective amount of a compound according to any one of the first to fourth aspects or a pharmaceutical composition according to the fifth aspect.

[0014] An eighth aspect of the invention is a compound according to any one of the first to fourth aspects, or a pharmaceutical composition according to the fifth aspect, for use in the treatment of a non-alcoholic steatohepatitis.

[0015] A ninth aspect of the invention is a compound of the formula



[0016] A tenth aspect of the invention is a method of preparing the compound of the ninth aspect, comprising reacting *tert*-butyl N-[(E)-2-(bromomethyl)-3-fluoro-allyl]carbamate with 2-chloropyrimidin-5-ol.

[0017] In one form, the present invention provides compounds of one of the first to fourth aspects as a free base. In another form, the present invention provides compounds of one of the first to fourth aspects as acid addition salts, such as a mono or di HCl addition salt(s) or a sulfonate salt, preferably a 4-methylbenzenesulfonate (a tosylate salt).

[0018] In another form, the present invention provides a compound for use in a method of treating a patient in need of treatment for a liver disorder. The method comprises administering to the patient an effective amount of a pharmaceutical composition that comprises the compound, or a pharmaceutically acceptable salt thereof.

[0019] In another form, the present invention provides a compound for use in a method of treating a patient in need of treatment for a liver disorder. The method comprises administering to the patient an effective amount of the compound, or a pharmaceutically acceptable salt thereof. Examples of liver disorders include liver inflammation, fibrosis, and steatohepatitis. In certain embodiments, the method comprises treating a patient in need of treatment for a liver disorder where the liver disorder is selected from: liver fibrosis, alcohol induced fibrosis, alcoholic steatosis, non-alcoholic fatty liver disease (NAFLD), and non-alcoholic steatohepatitis (NASH). In a particularly preferred embodiment, the method comprises treating a patient in need of treatment for non-alcoholic steatohepatitis (NASH).

[0020] In another form, the present invention provides a compound, or a pharmaceutically acceptable salt thereof, for use in therapy. In preferred embodiments, the therapy is for a liver disorder. Preferably, the therapy is for a liver disorder selected from: liver fibrosis, alcohol induced fibrosis, alcoholic steatosis, non-alcoholic fatty liver disease, and non-alcoholic steatohepatitis. In one embodiment, the therapy is for the treatment of liver fibrosis. In another embodiment, the therapy is for non-alcoholic fatty liver disease. In still yet another embodiment, the therapy is for non-alcoholic steatohepatitis.

[0021] In another form, the present invention provides a compound, or a pharmaceutically

acceptable salt thereof, for use in the treatment of a liver disorder. In one embodiment, the liver disorder is selected from: liver fibrosis, alcohol induced fibrosis, alcoholic steatosis, non-alcoholic fatty liver disease, and non-alcoholic steatohepatitis. In another embodiment, the liver disorder is non-alcoholic fatty liver disease or non-alcoholic steatohepatitis. In a particularly preferred embodiment, the liver disorder is non-alcoholic steatohepatitis (NASH).

[0022] The term "pharmaceutically-acceptable salt" as used herein refers a salt of a compound of the invention considered to be acceptable for clinical and/or veterinary use. Examples of pharmaceutically acceptable salts and common methodology for preparing them can be found in "Handbook of Pharmaceutical Salts: Properties, Selection and Use" P. Stahl, et al., 2nd Revised Edition, Wiley-VCH, 2011 and S.M. Berge, et al., "Pharmaceutical Salts", Journal of Pharmaceutical Sciences, 1977, 66(1), 1-19.

[0023] The pharmaceutical compositions for the present invention may be prepared using pharmaceutically acceptable additives. The term "pharmaceutically acceptable additive(s)" as used herein for the pharmaceutical compositions, refers to one or more carriers, diluents, and excipients that are compatible with the other additives of the compositions or formulations and not deleterious to the patient. Examples of pharmaceutical compositions and processes for their preparation can be found in "Remington: The Science and Practice of Pharmacy", Loyd, V., et al. Eds., 22nd Ed., Mack Publishing Co., 2012. Non-limiting examples of pharmaceutically acceptable carriers, diluents, and excipients include the following: saline, water, starch, sugars, mannitol, and silica derivatives; binding agents such as carboxymethyl cellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl-pyrrolidone; kaolin and bentonite; polyethyl glycols.

[0024] As used herein, the term "effective amount" refers to an amount that is a dosage, which is effective in treating a disorder, such as a liver disease including liver inflammation, fibrosis, and steatohepatitis. The attending physician, as one skilled in the art, can readily determine an effective amount by the use of conventional techniques and by observing results obtained under analogous circumstances. In determining an effective amount or dose of a compound, a number of factors are considered, including, but not limited whether the compound or its salt, will be administered; the co-administration of other agents, if used; the species of mammal; its size, age, and general health; the degree of involvement or the severity of the disorder; the response of the individual patient; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; the use of other concomitant medication; and other relevant circumstances.

[0025] As used herein, the terms "treating", "to treat", or "treatment", includes slowing, reducing, or reversing the progression or severity of an existing symptom, disorder, condition, or disease, which can include treating liver disease, such as, liver inflammation, fibrosis, and steatohepatitis.

[0026] As used herein, the term "patient" refers to a mammal, fowl, or fish. Preferably, the patient is a human or companion mammal, such as, a dog or cat or other domesticated

mammal, such as, a cow, pig, horse, sheep, rabbit, and goat.

[0027] A treating physician, veterinarian, or other medical person will be able to determine an effective amount of the compound for treatment of a patient in need. Preferred pharmaceutical compositions can be formulated as a tablet or capsule for oral administration, a solution for oral administration, or an injectable solution. The tablet, capsule, or solution can include a compound of the present invention in an amount effective for treating a patient in need of treatment.

[0028] The abbreviations used herein are defined according to Daub G.H., et al., "The Use of Acronyms in Organic Chemistry" *Aldrichimica Acta*, 1984, 17(1), 6-23. Other abbreviations are defined as follows: "ACN" refers to acetonitrile; "AUC" refers to area under the curve; "Boc" represents *tert*-butoxycarbonyl; "DCM" refers to dichloromethane; "DIPEA" refers to N,N-diisopropylethylamine; "DMF" refers to dimethylformamide; "DMSO" refers to dimethylsulfoxide; "EDTA" refers to ethylenediaminetetraacetic acid; "EGTA" refers to ethylene glycol tetraacetic acid; "ES/MS" refers to electrospray mass spectroscopy; "EtOAc" refers to ethyl acetate; "HEPES" refers to 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; "HPLC" refers to high performance liquid chromatography; "hr or hrs" refers to hour or hours; "HRP" refers to horse radish peroxidase; "IC₅₀" refers to the concentration of an agent which produces 50 % of the maximal inhibitory response possible for that agent (relative IC₅₀), or the concentration of an agent which produces 50 % inhibition of the target activity compared to placebo control (absolute IC₅₀); "MAOa and MAOb" refers to monoamine oxidase a and b isoform, respectively; "MeOH" refers to methyl alcohol or methanol; "min" refers to minutes; "MS" refers to mass spectroscopy; "PE" refers to petroleum ether; "R_t" refers to retention time; "SSAO" refers to semicarbazide-sensitive amine oxidase; and "hSSAO" refers to human SSAO.

[0029] The compounds of the present invention, or salts thereof, may be prepared by a variety of procedures, some of which are illustrated in the Preparations, and the Examples below. The product(s) of each step in the procedures below can be recovered by conventional methods including extraction, evaporation, precipitation, chromatography, filtration, trituration, and crystallization. Unless noted to the contrary, the reagents and starting materials are readily available.

[0030] In the preparations described herein the hydroxyl and amino functionalities can be protected to facilitate the synthesis of the compounds described herein. Examples of protecting functionalities can be found in "Greene's Protective Groups in Organic Synthesis," Wuts, P.G.M., et al., Eds. 5th Ed., John Wiley and Sons, 2014. Other functional groups that can be readily converted to the hydroxyl group or the amino group can be used. Such functional groups, preparations, and transformations of these groups can be found in "Comprehensive Organic Transformations: A Guide to Functional Group Preparations" by Larock, R.C., Wiley VCH, 1999 and in "March's Advanced Organic Chemistry: Reactions, Mechanisms and Structure," Smith, M.B., Ed., 7th Ed., Wiley-Interscience, 2013.

Chemical Synthesis Section

[0031] The following Preparations, Examples and Reference Examples further illustrate the invention and represent typical synthesis of the compounds of the invention. Subject-matter denoted "Reference Example" does not fall within the scope of the claims.

Preparation 1

tert-Butyl (3S)-3-methoxypyrrolidine-1-carboxylate

[0032] Add iodomethane (0.398 g, 2.80 mmol) to a mixture of *tert*-butyl (3S)-3-hydroxypyrrolidine-1-carboxylate (0.500 g, 2.67 mmol) and sodium hydride (60 mass % in mineral oil) (0.160 g, 4.01 mmol) in DMF (5 mL). Stir the resulting mixture at room temperature for 2 hours. Quench the reaction with saturated aqueous NH₄Cl aq. (30 mL) and extract with EtOAc (3×30 mL). Discard the aqueous layer. Combine the organic extracts and wash with brine, dry over Na₂SO₄, filter, and evaporate the filtrate to dryness to give the title compound (475 mg, 0.475 g, 88.4%). The crude material can be used in the next step without further purification. ES/MS (m/z): 224.2 (M+Na).

Preparation 2


(3 S)-3-Methoxypyrrolidine


[0033] Stir a solution of *tert*-butyl (3S)-3-methoxypyrrolidine-1-carboxylate (475 mg, 2.36 mmol) and trifluoroacetic acid (1 mL, 13.23 mmol) in DCM (3 mL) at room temperature for 1 hour. Concentrate the reaction mixture under vacuum to give the title compound (240 mg, 2.35 mmol, 99.5%), which can be used in the next step without further purification.

Preparation 3

1-(5-Benzyloxypyrimidin-2-yl)piperidin-4-ol

[0034]






[0035] Stir a mixture of 5-benzyloxy-2-chloro-pyrimidine (2.30 g, 9.90 mmol), piperidin-4-ol (1.23 g, 11.9 mmol) and DIPEA (3.88 mL, 29.7 mmol) in DMF (30 mL, 388 mmol) at 100 °C under a N₂ atmosphere for 17 hrs. Dilute the mixture with water (200 mL) and extract with EtOAc (3×50 mL). Combine the organic extracts; wash with brine (3×50 mL), dry over Na₂SO₄, filter, and concentrate the filtrate. Subject the residue to silica gel flash column chromatography eluting with a mixture of 60% EtOAc and 40% PE to give the title compound (2.00 g, 6.31 mmol, 63.7%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 1.40-1.60 (m, 2H), 1.90-1.99 (m, 2H), 3.17-3.29 (m, 2H), 3.85-3.95 (m, 1H), 4.25-4.39 (m, 2H), 5.09 (s, 2H), 7.38-7.52 (m, 5H), 8.14 (s, 2H).

Preparation 4

2-(4-Hydroxy-1-piperidyl)pyrimidin-5-ol

[0036]




[0037] Add palladium (10 mass % on carbon, 600 mg, 0.564 mmol) to 1-(5-benzyloxypyrimidin-2-yl)piperidin-4-ol (2.00 g, 6.31 mmol) in MeOH (40 mL, 989 mmol). Stir the resulting mixture at 15 °C under a hydrogen atmosphere (103 kPa) for 2 hrs. Filter the resulting mixture through diatomaceous earth. Concentrate the filtrate to give the title compound (0.70 g, 3.59 mmol, 56.8%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 1.23-1.38 (m, 2H), 1.60-1.77 (m, 2H), 2.65-2.90 (m, 2H), 3.00-3.25 (m, 2H), 3.50-3.70 (m, 2H), 4.05-4.16 (m, 1H), 7.90 (s, 2H).

Preparation 5

tert-Butyl N-[(E)-2-[(2-chloropyrimidin-5-yl)oxymethyl]-3-fluoro-allyl]carbamate

[0038]






[0039] Add K_2CO_3 (5.71 g, 41.3 mmol) to a solution of 2-chloropyrimidin-5-ol (5.01 g, 38.3 mmol) and *tert*-butyl N-[(E)-2-(bromomethyl)-3-fluoro-allyl]carbamate (3.67 g, 13.7 mmol) in DMF (25 mL). Stir the resulting solution at room temperature for 12 hours. Quench the reaction by adding water (80 mL) and EtOAc (100 mL). Separate the organic and aqueous phases. Extract the aqueous phase with EtOAc (3 × 100 mL). Combine all the organic extracts. Dry the combined organic extracts over Na_2SO_4 , filter, and concentrate the filtrate under vacuum to provide a residue. Subject the residue to silica gel flash chromatography eluting with a mixture of 30% EtOAc in hexanes to give the title compound as a white solid (4.15 g, 13.1 mmol, 96 % yield). ES/MS (*m/z*): 340 (*M*+*Na*).

Preparation 6

***tert*-Butyl
N-[(Z)-3-fluoro-2-[[2-(4-hydroxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate**

N-[(Z)-3-fluoro-2-[[2-(4-hydroxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate

[0040]




[0041] Stir a mixture of 2-(4-hydroxy-1-piperidyl)pyrimidin-5-ol (400 mg, 2.05 mmol), *tert*-butyl N-[(Z)-2-(bromomethyl)-3-fluoro-allyl]carbamate (1.10 g, 4.02 mmol) and K_2CO_3 (0.858 g, 6.15 mmol) in DMF (10 mL) at 60 °C for 3 hrs. Dilute the resulting mixture with EtOAc (50 mL). Sequentially wash the mixture with water (100 mL) then brine (2×50 mL). Dry the organic phase over Na_2SO_4 , filter, and concentrate the filtrate to provide a residue. Subject the residue to silica gel flash column chromatography eluting with a gradient of 65-75% EtOAc in PE to give the title compound (558 mg, 1.46 mmol, 71.2%) as a yellow gum. 1H NMR (400 MHz, $CDCl_3$) δ 1.25 (s, 9H), 1.40-1.65 (m, 4H), 1.89-1.98 (m, 2H), 3.23-3.30 (m, 2H), 3.91 (s, 1H), 3.89-3.97 (m, 1H), 4.36-4.40 (m, 2H), 4.64 (s, 2H), 4.76 (br, 1H), 6.62 (d, *J* = 84.0 Hz, 1H) 8.10 (s, 2H).

Preparation 7

***tert*-Butyl
N-[(E)-3-F-2-[[2-(4-hydroxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate**

N-[(E)-3-F-2-[[2-(4-hydroxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate

[0042]




[0043] Combine *tert*-butyl N-[(E)-2-[(2-chloropyrimidin-5-yl)oxymethyl]-3-fluoro-allyl]carbamate (21.3 g, 67.0 mmol), piperidin-4-ol (25.0 g, 235 mmol), and DIPEA (41 mL, 235 mmol) in 1,4-dioxane (200 mL). Heat the resulting mixture to 105 °C under a N₂ atmosphere for 7 hours. Concentrate the mixture and partition it between water (100 mL) and EtOAc (150 mL), then separate the phases. Extract the aqueous phase with EtOAc (100 mL). Combine all the organic phases. Wash with brine (100 mL), dry over Na₂SO₄, filter, and concentrate the filtrate to provide a residue. Dissolve the residue in EtOAc (21 mL), and then drop-wise add heptane (126 mL). Stir the resulting mixture at room temperature for 20 hours. Filter the mixture to collect the solid, rinse the solid with EtOAc/heptane (5:1, 21 mL). Dry the solid under vacuum to give the title compound (23.5 g, 61.5 mmol, 91.7%). ES/MS (m/z): 383 (M+H).

Preparation 8

***tert*-Butyl
N-[(E)-3-fluoro-2-[[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate**

N-[(E)-3-fluoro-2-[[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate

[0044]




[0045] Divide *tert*-butyl N-[(E)-2-[(2-chloropyrimidin-5-yl)oxymethyl]-3-fluoro-allyl]carbamate (4.10 g, 12.9 mmol) into 2 equal parts (2.05 g + 2.05 g) and place each part into a separate microwave vial (20 mL). Into each vial, add 4-methoxypiperidine (4.31 g, 37.3 mmol), 1,4-dioxane (30 mL, 15 mL) and DIPEA (6 mL, 34.4 mmol, 3 mL). Flush the vials with N₂ gas, seal, and heat to 120 °C for 12 hours by microwave. Combine the two reaction mixtures and

concentrate under vacuum to provide a residue. Subject the residue silica gel flash chromatography eluting with a mixture of 30% EtOAc in hexanes to give the title compound as a yellow oil (4.24 g, 10.2 mmol, 79 %). ES/MS (m/z): 397 (M+H).

Preparation 9

***tert*-Butyl** N-[(E)-3-fluoro-2-[[2-[(3S)-3-methoxypyrrolidin-1-yl]pyrimidin-5-yl]oxymethyl]allyl]carbamate

[0046]




[0047] Add *tert*-butyl N-[(E)-2-[(2-chloropyrimidin-5-yl)oxymethyl]-3-fluoro-allyl]carbamate (400 mg, 1.26 mmol) to a mixture of (3S)-3-methoxypyrrolidine (240 mg, 2.37 mmol) and K_2CO_3 (0.696 g, 5.04 mmol) in 1,4-dioxane (5 mL). Stir the resulting mixture at 120 °C under microwave conditions for 12 hours. Concentrate the mixture under vacuum to provide the title compound as a crude material, which can be used in the next step without further purification (481 mg, 1.26 mmol, 99.9%). ES/MS (m/z): 383.2 (M+H).

Preparation 10

***tert*-Butyl** N-[(E)-3-fluoro-2-[[2-[(3S)-3-hydroxypyrrolidin-1-yl]pyrimidin-5-yl]oxymethyl]allyl]carbamate

[0048]




[0049] Dissolve *tert*-butyl N-[(E)-2-[(2-chloropyrimidin-5-yl)oxymethyl]-3-fluoro-allyl]carbamate

(594.3 mg, 1.87 mmol) and (3S)-pyrrolidin-3-ol (488.9 mg, 5.61 mmol) in 1,4-dioxane (15 mL) and DIPEA (3 mL, 17.2 mmol). Flush the solution with N₂ gas, seal the vessel, and heat the mixture to 120 °C for 12 hours by microwave. Concentrate the resulting mixture under vacuum to provide a residue. Subject the residue to silica gel flash chromatography eluting with a gradient of 80-90% EtOAc in hexanes to give the title compound as a yellow foam (608.8 mg, 1.57 mmol, 84 %). ES/MS (m/z): 369 (M+H).

Example 1

1-[5-[(Z)-2-(Aminomethyl)-3-fluoro-allyloxy]pyrimidin-2-yl]piperidin-4-ol hydrochloride

[0050]




[0051] Add *tert*-butyl N-[(Z)-3-fluoro-2-[[2-(4-hydroxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate (558 mg, 1.46 mmol) to HCl (4 mol/L) and MeOH (10 mL). Stir the resulting mixture for 1.5 hrs at 10 °C. Concentrate the mixture under reduced pressure to provide a residue. Subject the residue to prep-HPLC column: (Phenomenex Synergi C18 150 30 mm, 4 µm) eluting with a gradient of 10 to 15% 0.05% aqueous HCl in ACN; flow rate: 25 mL/min, Rt: 5.95 min to give the title compound with a Z:E ratio of greater than 20:1 (456 mg, 1.43 mmol, 98.0%) as a yellow solid. ES/MS (m/z): 283.1 (M+H), ¹H NMR (400 MHz, d₄-MeOD) δ 1.56-1.78 (m, 2H), 1.92-2.13 (m, 2H), 3.61-3.72 (m, 2H), 3.75 (s, 2H), 3.95-4.06 (m, 1H), 4.11-4.25 (m, 2H), 4.89-4.93 (m, 2H), 7.19 (d, J = 80.4 Hz, 1H), 8.47 (s, 2H).

Example 2

1-[5-[(E)-2-(Aminomethyl)-3-fluoro-allyloxy]pyrimidin-2-yl]piperidin-4-ol dihydrochloride

[0052]




[0053] Combine *tert*-butyl N-[(E)-3-fluoro-2-[[2-(4-hydroxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate (31 g, 81.07 mmol), MeOH (20 mL, 494 mmol), and HCl in MeOH (120 mL, 4 mol/L, 486.4 mmol). Stir the resulting mixture for 30 hours under a N₂ atmosphere at room temperature. Add EtOAc (250 mL) drop-wise and stir the mixture for 30 minutes. Filter the mixture to collect the solid, rinse the solid with EtOAc (20 mL), and then dry the solid under vacuum to give the title compound with an E:Z ratio of greater than 20:1 (26.4 g, 72.8 mmol, 89.8%). ES/MS (m/z): 283 (M+H), ¹H NMR (500 MHz, d₆-DMSO) δ 1.28-1.36 (m, 2H), 1.73-1.76 (m, 2H), 3.18-3.25 (m, 2H), 3.55-3.60 (m, 2H), 3.68-3.75 (m, 1H), 4.18 (dt, J = 13.5, 4.5 Hz, 2H), 4.65 (d, J = 3.0 Hz, 2H), 6.55-7.10 (br, 2H), 7.27 (d, J = 82.0 Hz, 1H), 8.27 (s, 2H), 8.32-8.45 (br, 3H), ¹⁹F NMR (500 MHz, d₆-DMSO) δ 122.2 (s).

Reference Example 3

(E)-3-Fluoro-2-[[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]prop-2-en-1-amine, dihydrochloride

[0054]




[0055] Dissolve *tert*-butyl N-[(E)-3-fluoro-2-[[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl]carbamate (4.2374 g, 10.69 mmol) in HCl in MeOH (100 mL, 50 mmol, 0.5 mol/L). Heat the resulting clear solution to 60 °C for 4 hours. Concentrate the mixture under vacuum to provide a residue (3.95 g). Suspend the residue in MeOH (7 mL) and reflux the mixture to provide a clear solution. Cool the solution to room temperature to give needle crystals, and then cool the mixture to -20 °C. Filter the mixture to collect the solid, and wash the solid with cool MeOH to give the title compound as light yellow crystals (2.73 g, 7.01 mmol, 66 %). The resulting yellow crystals can be further purified via recrystallization with the same procedure described above to give the title compound as colorless, crystalline material with an E:Z ratio of greater than 20:1 ES/MS (m/z): 297 (M+H), ¹H NMR (500 MHz, d₆-DMSO) δ 1.33-1.40 (m, 2H), 1.84-1.89 (m, 2H), 3.26 (dt, J = 13.5, 9.5 Hz, 2H), 3.27 (s, 3H), 3.40-3.45 (m, 1H), 3.56-3.62 (m, 2H), 4.11 (dt, J = 13.5, 5.0 Hz, 2H), 4.62 (d, J = 3.0 Hz, 2H), 5.26-5.94 (br, 1H), 7.27 (d, J = 82.0 Hz, 1H), 8.26 (s, 2H), 8.21-8.31 (br, 3H), ¹⁹F NMR (500 MHz, d₆-DMSO) δ 122.1 (s).

Reference Example 4

(E)-3-fluoro-2-[[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]prop-2-en-1-amine

[0056]




[0057] Dissolve N-[(E)-3-Fluoro-2-[[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]allyl] carbamate (167.0 mg, 0.42 mmol) in 0.95M HCl solution in 18 mL EtOAc/MeOH (10:1 v/v). Stir the mixture overnight. Concentrate the resulting suspension under vacuum and dissolve the residue in water. Subject the residue to pre-HPLC; LC column: XBridge® C18 30 × 150 mm 5 µm; eluting with a gradient of 14 to 24% 10 mM NH₄HCO₃ aqueous solution in ACN from 0-11 minutes; column temperature: room temperature; flow rate: 35 mL/min, Rt = 7.8 minutes stopping after 17 min. monitor by UV. Collect and concentrate the appropriate fractions to provide a residue as an oil. The residue is dissolved in water and lyophilized to give the title compound as a white solid with an E:Z ratio greater than 20:1 (97 mg, 0.31 mmol, 74%, 95% purity). ES/MS (m/z): 297 (M+H), ¹H NMR (500 MHz, d₆-DMSO) δ 1.33-1.41 (m, 2H), 1.52-1.69 (br, 1H), 1.83-1.89 (m, 2H), 3.23-3.29 (m, 4H), 3.27 (s, 3H), 3.30-3.35 (br, 1H), 3.38-3.44 (m, 1H), 4.11 (dt, J = 13.5, 4.5 Hz, 2H), 4.55 (d, J = 4.5 Hz, 2H), 6.93 (d, J = 85.0 Hz, 1H), 8.22 (s, 2H), ¹⁹F NMR (500 MHz, d₆-DMSO) δ 131.8 (s).

Reference Example 4a

(2E)-3-Fluoro-2-({[2-(4-methoxypiperidin-1-yl)pyrimidin-5-yl]oxy}methyl)prop-2-en-1-amine 4-methylbenzenesulfonate (1:1)

[0058]



[0059] Dissolve (E)-3 -fluoro-2- [[2-(4-methoxy-1-piperidyl)pyrimidin-5-yl]oxymethyl]prop-2-en-1-amine (2.316 g, 7.81 mmol) in methyl acetate (3 mL) and stir at 1000 rpm at room temperature to give a light yellow solution. Add 4-toluenesulfonic acid monohydrate (1.62 g, 8.43 mmol) in a solution of methyl acetate (4 mL). The mixture becomes cloudy and a thick yellow slurry quickly forms. Filter the solid by vacuum filtration through filter paper. Rinse the filter cake with methyl acetate (4 mL) to give a white filter cake. Dry the solid under a vacuum air stream for 10 minutes and then in a vacuum oven at room temperature overnight to give the title compound (3.00 g, 81.8%).

X-Ray Powder Diffraction of Reference Example 4a,

[0060] The X-ray powder diffraction (XRD) pattern of crystalline (2E)-3-fluoro-2-([2-(4-methoxypiperidin-1-yl)pyrimidin-5-yl]oxy)methylprop-2-en-1-amine 4-methylbenzenesulfonate is obtained on a Bruker D4 Endeavor X-ray powder diffractometer, equipped with a CuK α source ($\lambda = 1.54060 \text{ \AA}$) and a Vantec detector, operating at 35 kV and 50 mA. The sample is scanned between 4 and 40° in 2 θ , with a step size of 0.009° in 2 θ and a scan rate of 0.5 seconds/step, and with 0.6 mm divergence, 5.28 fixed anti-scatter, and 9.5 mm detector slits. The dry powder is packed on a quartz sample holder and a smooth surface is obtained using a glass slide. The crystal form diffraction patterns are collected at ambient temperature and relative humidity. It is well known in the crystallography art that, for any given crystal form, the relative intensities of the diffraction peaks may vary due to preferred orientation resulting from factors such as crystal morphology and habit. Where the effects of preferred orientation are present, peak intensities are altered, but the characteristic peak positions of the polymorph are unchanged. See, e.g., The United States Pharmacopeia #23, National Formulary #18, pages 1843-1844, 1995. Furthermore, it is also well known in the crystallography art that for any given crystal form the angular peak positions may vary slightly. For example, peak positions can shift due to a variation in the temperature or humidity at which a sample is analyzed, sample displacement, or the presence or absence of an internal standard. In the present case, a peak position variability of ± 0.2 in 2 θ will take into account these potential variations without hindering the unequivocal identification of the indicated crystal form. Confirmation of a crystal form may be made based on any unique combination of distinguishing peaks (in units of ° 2 θ), typically the more prominent peaks. Collect the crystal form diffraction patterns at ambient temperature and relative humidity and adjust based on NIST 675 standard peaks at 8.853 and 26.774 degrees 2-theta.

[0061] A prepared sample of (2E)-3-fluoro-2-([2-(4-methoxypiperidin-1-yl)pyrimidin-5-yl]oxy)methylprop-2-en-1-amine 4-methylbenzenesulfonate is characterized by an XRD pattern using CuK α radiation as having diffraction peaks (2-theta values) as described in Table 1 below, and in particular having peaks at 18.6 in combination with one or more of the peaks selected from 22.4, 19.1, and 21.0; with a tolerance for the diffraction angles of $\pm 0.2^\circ$ in 2 theta alternatively the salt can be characterized by an XRD pattern having one or more peaks

at 18.6, 19.1, 21.0, 21.9, and 22.4 $\pm$ 0.2° in 2 theta, or 17.6, 11.0, 16.8, 18.6, 19.1, 21.0, 21.9, 22.4 and 26.1 $\pm$ 0.2° in 2 theta.


Table 1

X-ray powder diffraction peaks of (2E)-3-fluoro-2-([2-(4-methoxypiperidin-1-yl)pyrimidin-5-yl]oxy)methyl)prop-2-en-1-amine 4-methylbenzenesulfonate		
Peak	Angle (°2-Theta) $\pm$ 0.2°	Relative Intensity (% of most intense peak)
1	7.4	17.6%
2	11.0	23.4%
3	12.7	5.2%
4	16.8	10.1%
5	18.6	100.0%
6	19.1	42.2%
7	21.0	41.9%
8	21.9	31.6%
9	22.4	77.5%
10	26.2	18.6%

Example 5

(E)-3-Fluoro-2-[[2-[(3S)-3-methoxypyrrolidin-1-yl]pyrimidin-5-yl]oxymethyl]prop-2-en-1-amine

[0062]




[0063] Stir a solution of *tert*-butyl N-[(E)-3-fluoro-2-[[2-[(3S)-3-methoxypyrrolidin-1-yl]pyrimidin-5-yl]oxymethyl]allyl]carbamate (481 mg, 1.26 mmol) and trifluoroacetic acid (1 mL, 13.23 mmol) in DCM (3 mL) at room temperature for 1 hour. Concentrate the mixture under vacuum. Subject the residue to prep-HPLC: LC Column: XBridge® C18 30 $\times$ 150 mm 5 μ m; eluting with 5% 10mM aqueous NH_4HCO_3 in ACN from 0-2 minutes, then a gradient of 6-11% 10 mM

aqueous NH_4HCO_3 in ACN from 2-12 minutes; stop at 18 minutes; column temp: room temperature; flow rate: 35 mL/minutes, R_t = 10.6 minutes; monitor by UV to give the title compound (226 mg, 61.7%) as a white solid with an E:Z ratio greater than 20:1. ES/MS (m/z): 283.1 ($M+H$), ^1H NMR (500 MHz, CDCl_3) δ 1.12-1.78 (br, 2H), 2.07-2.16 (m, 2H), 3.37 (s, 3H), 3.53-3.68 (m, 6H), 4.07 (m, 1H), 4.44 (s, 2H), 6.57 (d, J = 83.0 Hz, 1H), 8.12 (s, 2H).

Example 6

(3S)-1-[5-[(E)-2-(Aminomethyl)-3-fluoro-allyloxy]pyrimidin-2-yl]pyrrolidin-3-ol;
dihydrochloride

[0064]



[0065] Dissolve *tert*-butyl N-[(E)-3-fluoro-2-[[2-[(3S)-3-hydroxypyrrolidin-1-yl]pyrimidin-5-yl]oxymethyl] allyl] carbamate (605.9 mg, 1.65 mmol) into a mixture of HCl in MeOH (30 mL, 15 mmol, 0.5 mol/L) and HCl in water (5 mL, 60 mmol, 12 mol/L). Stir the resulting clear solution overnight. Concentrate the reaction mixture under vacuum to provide a residue. Subject the residue to pre-HPLC: LC Column: XBridge[®] C18 30 × 150 mm 5 μm ; H_2O 10 mM NH_4HCO_3 ; at room temperature; eluting with 2% ACN for 0-2 minutes, then a gradient of 2-10% ACN for 2-10 minutes, at a flow rate: 35 mL/minutes, R_t = 8.0 minutes (monitored via UV detection); stopping at 16 mins. Collect the appropriate fractions and concentrate to provide the free base of the title compound. Dissolve the free base compound in 0.5 M HCl in MeOH (15 mL). Concentrate the solution, add water, and then lyophilize to give the title product as a light yellow solid with an E:Z ratio greater than 20:1 (352.7 mg, 0.982 mmol, 59 %). ES/MS (m/z): 269 ($M+H$), ^1H NMR (500 MHz, d_6 -DMSO) δ 1.87-1.94 (m, 1H), 1.98-2.05 (m, 1H), 3.44 (d, J = 11.5 Hz 1H), 3.51-3.61 (m, 5H), 4.39-4.42 (m, 1H), 4.66 (d, J = 3.0 Hz, 2H), 5.33-5.90 (br, 2H), 7.28 (d, J = 82.0 Hz, 1H), 8.35 (s, 2H), 8.35-8.44 (br, 3H), ^{19}F NMR (500 MHz, d_6 -DMSO) δ 121.9 (s).

Alternate Preparation of Example 6

[0066] Dissolve *tert*-butyl N-[(E)-3-fluoro-2-[[2-[(3S)-3-hydroxypyrrolidin-1-yl]pyrimidin-5-

yl]oxymethyl] allyl] carbamate (1.1601 g, 3.15 mmol) in a solution of HCl in EtOAc (50 mL, 50 mmol, 1.0 mol/L) (pre-mixed with 0.5 mol/L HCl in MeOH, 5 mL). Stir the resulting solution overnight. Concentrate the white suspension under vacuum to provide a white powder. Dissolve the white powder in water and lyophilize the solution to provide the title compounds as a light yellow solid (860.7 mg, 2.42 mmol, 77%).

[0067] The material prepared as Example 6 is dissolved in water (5 mL) and combined with the material of Alternate Example 6 in water (5 mL). Lyophilize the mixture to give the title compound with an E:Z ratio greater than 20:1 (1.151 g 3.27 mmol). ES/MS m/z: 269 (M+H), ¹H NMR (500 MHz, d₆-DMSO) δ 1.87-1.94 (m, 1H), 1.98-2.05 (m, 1H), 3.44 (d, J = 11.5 Hz 1H), 3.51-3.61 (m, 5H), 4.39-4.42 (m, 1H), 4.66 (d, J = 3.0 Hz, 2H), 5.33-5.90 (br, 2H), 7.28 (d, J = 82.0 Hz, 1H), 8.35 (s, 2H), 8.35-8.44 (br, 3H), ¹⁹F NMR (500 MHz, d₆-DMSO) δ 121.9 (s).

Biological Assays

SSAO/VAP-1 *In vitro* Activity

[0068] Amine oxidase activity of recombinant SSAO, MAOa, and MAOb isoforms are measured using the MAO-Glo™ assay kit from Promega (V1402). Test compounds (with DMSO as the vehicle, 0.5% v/v for SSAO) and the enzyme are incubated for 10 mins at room temperature before the addition of the luminogenic substrate. The substrate concentration is 10 μM for human recombinant SSAO. The assays are conducted in a pH 7.4 buffer (50 mM HEPES, 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1.4 mM MgCl₂, 0.001% Tween-20) in a well-plate. Oxidation of the substrate is conducted for 2 hrs before the addition of detecting reagent according to the manufacture's protocol. The IC₅₀ value of the tested compounds is calculated by fitting the dose response curve using a 4-parameter non-linear regression routine. The IC₅₀ values for the compounds of Reference Example 3, and Examples 5 and 6 are listed in Table 2.

Table 2

Ex	hSSAO Inhib Rel IC ₅₀ (nM)
3*	12 ± 1, n=5
5	32 ± 7, n=4
6	52 ± 5, n=7
*Reference example	

[0069] Data are presented as mean±SEM (SEM=standard error of the mean) for the number of assays (n).

[0070] The compounds of the Examples exhibited an IC_{50} for hSSAO of less than 60 nM. The compounds of the Examples exhibited an IC_{50} hMAOa and hMAOb more than 50 μ M and 200 μ M, respectively, indicating that the compounds of the Examples are selective for hSSAO over either hMAOa or hMAOb.

SSAO Target Engagement

[0071] The SSAO activity in rat plasma and liver tissues are measured using the MAO-Glo™ assay kit from Promega (V1402). The residual SSAO activity in rats after compound treatment is estimated by measuring the total amine oxidase activity in plasma or liver lysates that are insensitive to the presence of the MAO inhibitors Clogyline and Pargyline. Rats are administered the compound of Example 2 at the doses of 15, 3, 0.6, 0.12, 0.025, and 0.005 mg/kg. The control group is administered with the same volume (2 mL/kg) of the dosing vehicle (hydroxyethyl cellulose 1% w/v, 0.25% Tween 80). Plasma and liver at 2 or 24 hours post compound treatment are harvested and stored at -78 °C until analysis. Tissue lysates are prepared by homogenization in a lysis buffer (20 mM HEPES, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100 and 1x Roche Complete protease inhibitor tablet). Tissue particles are removed by centrifugation at 12,000 rpm at 4 °C for 30 minutes. 40 μ L of plasma or liver lysates is incubated with Clogyline (10 μ M) and Pargyline (10 μ M) for 20 minutes at room temperature before the addition of the luminogenic substrate (50 μ M) for 60 minutes. The product generated is quantified according to the manufacture's procedure. The fraction of activity that is insensitive to the presence of the MAO inhibitors is used as the surrogate for the residual SSAO activity. The compound of Example 2 was evaluated in the protocol essentially as described above administered at various doses. The results are listed in Table 3.

Table 3

SSAO Target Engagement for Example 2			
Dose (mg/kg)	SSAO Activity (%)		
	Plasma		Liver
	2 hour	24 hour	24 hour
Vehicle	100 $\pm$ 7	100 $\pm$ 9	100 $\pm$ 11
0.005	31 $\pm$ 3	70 $\pm$ 7	75 $\pm$ 18
0.025	21 $\pm$ 1	55 $\pm$ 9	68 $\pm$ 14
0.12	7 $\pm$ 1	35 $\pm$ 4	45 $\pm$ 8
0.6	4.3 $\pm$ 0.8	16.3 $\pm$ 2.6	16.2 $\pm$ 2.9
3	3.6 $\pm$ 1.0	8.8 $\pm$ 1.3	7.1 $\pm$ 1.7
15	-0.6 $\pm$ 0.3	6.6 $\pm$ 1.8	9.6 $\pm$ 2.6
Data are presented as mean $\pm$ SEM, n=6			

[0072] The results indicate that the compound of Example 2 dose-dependently inhibits SSAO activity in both rat plasma and liver.

Mouse Model of NASH and Fibrosis Induced by 3H Diet

[0073] Male C57BL/6N mice are fed with D09100301 diet (Research Diets, 40% fat, 2% cholesterol, 24% fructose, (the, high fat, high cholesterol and high fructose, the "3H diet") for 150 days. Each mouse is then singly housed after 5 days for an acclimation period. Plasma alanine aminotransferase (ALT) and cytokeratin 18 (CK18) are measured. After one week of recovery, the mice are randomized into 5 groups based on their ALT values, CK18 values, and body weight. Animals of each group are administered either vehicle (0.5% methylcellulose (MC) + 0.25% Tween 80 in distilled water) or the compound of Example 6 (at a 0.06, 2, 6, and 20 mg/kg dose) once daily in a volume of 5 mL/kg for 11 weeks.

[0074] Blood is collected from mice treated with the compound of Example 6 for 76 days 2 hours after the last dose. Compound levels in the plasma are analyzed by mass spectroscopy. The results are listed below in Table 4. Treated mice exhibit a dose-dependent increase in plasma compound levels. All groups of mice treated with Example 6 exhibit a significant decrease in ALT, indicating decreased hepatic lesions in those animals. Animals treated with 6 and 20 mg/kg of the compound of Example 6 also exhibit decreased triglyceride levels in blood.

[0075] At the completion of the study, the animals are sacrificed and their livers excised. Two sections of the left and right lobes are fixed in neutral buffered 10% formalin. Liver tissue slides are stained with hematoxylin and eosin (H&E), Sirius red, and Masson's Trichrome to prepare slides for pathological analysis. All specimens are examined microscopically and scored as a modified Brunt score NASH Activity Score. Scores are based on the grading scheme and end-points as described in Brunt E. M, et al., "Histopathology of nonalcoholic fatty liver disease," World J. of Gastroenterol, 2010, 16(42), 5286-5296. Group means are then calculated for each individual end-point. The following endpoints are used to characterize the fast food model of NASH in mice as modified from NASH endpoints (See Brunt, E. M. "Histopathology of nonalcoholic fatty liver disease," Clin Liver Dis., 2009, 13, 533-544 and Brunt, E. M, et al., "Nonalcoholic steatohepatitis: A proposal for grading and staging the histological lesions", Am J Gastroenterology, 1999, 94(9), 2467-2474.

[0076] Histopathological analysis of the livers from the mice treated with the compound of Example 6 is provided in Table 4. The results indicate that there is a significant decrease in hepatic inflammation, macrovesicular vacuolation, and perisinusoidal fibrosis in the mice that are treated with 20 mg/kg of the compound of Example 6.

Table 4

Efficacy Results for the Compound of Example 6				
Dose (mg/kg)	Nos. of Animals	Plasma Compound Level (ng/mL)	ALT (IU/L) ¹	Plasma TG (mg/dL) ¹
Vehicle	12	-	713±43	69±4
0.6	13	66±5	530±45*	55±6
2	13	163±21	483±44**	70±11
6	11	560±29	495±25*	46±4*
20	11	2117±219	466±37***	45±5*

¹MIXED model is applied to compare fold changes from baseline adjusted by baseline between compound treatment groups and vehicle group, (See "Generalized, Linear, and Mixed Models," McCulloch, C.E., and Searle, S.R., Eds. John Wiley and Sons, 2000 and "Mixed-Effects Models in S and S-PLUS", Pinheiro J.C., and Bates, D. M., Eds. Springer, 2000.) Data are presented as mean±SEM. * $p<0.05$; ** $p<0.01$; *** $p<0.001$

Table 5

Liver Histopathological Analysis for the Compound of Example 6						
Dose (mg/kg)	Score (0-3) ¹					
	Hepatic inflammation		Macrovesicular vacuolation		Perisinusoidal fibrosis	
	Left	Right	Left	Right	Left	Right
Vehicle	2.8±0.1	2.7±0.1	2.8±0.1	2.8±0.1	2.5±0.2	2.3±0.1
0.6	2.6±0.1	2.5±0.1	2.5±0.1	2.6±0.1	2.4±0.2	1.9±0.2
2	2.5±0.1	2.6±0.1	2.5±0.1	2.7±0.1	2.5±0.2	1.8±0.2
6	2.5±0.2	2.6±0.2	2.9±0.1	2.7±0.1	2.6±0.2	2.1±0.3
20	2.1±0.1**	2.1±0.1*	2.4±0.2	2.3±0.1*	1.9±0.2	1.5±0.2*

¹ Nonparametric test is applied to compare scores between compound treatment groups and vehicle group. Scores for left and right laterals are compared separately.
Data are presented as mean±SEM.
* $p<0.05$; ** $p<0.01$; *** $p<0.001$

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [WO2007120528A](#) [0005]
- [WO2013163675A](#) [0006]

Non-patent literature cited in the description

- **WESTON, C.J. et al.**J Neural. Transm., 2011, vol. 118, 1055-1064 [0002]
- **DUNKEL et al.**Expert Opinion on Therapeutic Patents, vol. 21, 91453-1471 [0004]
- **P. STAHL et al.**Handbook of Pharmaceutical Salts: Properties, Selection and UseWiley-VCH20110000 [0022]
- **Pharmaceutical SaltsS.M. BERGE et al.**Journal of Pharmaceutical Sciences19770000vol. 66, 1-19 [0022]
- **Remington: The Science and Practice of PharmacyMack Publishing Co**20120000 [0023]
- **DAUB G.H. et al.**The Use of Acronyms in Organic ChemistryAldrichimica Acta, 1984, vol. 17, 16-23 [0028]
- **Greene's Protective Groups in Organic SynthesisJohn Wiley and Sons**20140000 [0030]
- **LAROCK. R.C.**Comprehensive Organic Transformations: A Guide to Functional Group PreparationsWiley VCH19990000 [0030]
- **March's Advanced Organic Chemistry: Reactions, Mechanisms and StructureWiley-Interscience**20130000 [0030]
- **The United States Pharmacopeia #23, National Formulary #18, 1995, 1843-1844** [0060]
- **BRUNT E. M et al.**Histopathology of nonalcoholic fatty liver diseaseWorld J. of Gastroenterol, 2010, vol. 16, 425286-5296 [0075]
- **BRUNT, E. M.**Histopathology of nonalcoholic fatty liver diseaseClin Liver Dis., 2009, vol. 13, 533-544 [0075]
- **BRUNT, E. M et al.,** "Nonalcoholic steatohepatitis: A proposal for grading and staging the histological lesionsAm J Gastroenterology, 1999, vol. 94, 92467-2474 [0075]
- **Generalized, Linear, and Mixed ModelsJohn Wiley and Sons**20000000 [0076]
- **Mixed-Effects Models in S and S-PLUSSpringer**20000000 [0076]


Patentkrav

1. Forbindelse med formelen:



5 eller et farmaceutisk acceptabelt salt deraf.


2. Forbindelse med formelen:



eller et farmaceutisk acceptabelt salt deraf.


10

3. Forbindelse med formelen:



eller et farmaceutisk acceptabelt salt deraf.

15 4. Forbindelse med formelen:



eller et farmaceutisk acceptabelt salt deraf.


5. Farmaceutisk sammensætning omfattende en forbindelse ifølge et hvilket som helst af kravene 1 til 4 og en farmaceutisk acceptabel bærer, fortyndingsmiddel eller excipiens.

5 **6.** Forbindelse ifølge et hvilket som helst af kravene 1 til 4, eller en farmaceutisk sammensætning ifølge krav 5, til anvendelse i terapi.

7. Forbindelse ifølge et hvilket som helst af kravene 1 til 4, eller en farmaceutisk sammensætning ifølge krav 5, til anvendelse i en fremgangsmåde til behandling af en patient med behov for behandling for ikke-alkoholisk steatohepatitis, hvor fremgangsmåden omfatter indgivelse til patienten af en effektiv mængde af en forbindelse ifølge et hvilket som helst af kravene 1 til 4 eller en farmaceutisk sammensætning ifølge krav 5.

15 **8.** Forbindelse ifølge et hvilket som helst af kravene 1 til 4, eller en farmaceutisk sammensætning ifølge krav 5, til anvendelse i behandlingen af en ikke-alkoholisk steatohepatitis.

9. Forbindelse med formlen



10. Fremgangsmåde til fremstilling af forbindelsen ifølge krav 9, omfattende omsætning af tert-butyl-N-[(E)-2-(brommethyl)-3-fluor-allyl]carbamate med 2-chlorpyrimidin-5-ol.

11. Fremgangsmåde ifølge krav 10, hvor omsætningen eventuelt udføres i nærvær af kaliumcarbonat.

12. Fremgangsmåde ifølge krav 10 eller 11, hvorved omsætningen anvender dimethylformamid som opløsningsmiddel.