

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 October 2003 (30.10.2003)

PCT

(10) International Publication Number
WO 03/088940 A1

(51) International Patent Classification⁷: **A61K 7/075**

(21) International Application Number: PCT/US03/12265

(22) International Filing Date: 22 April 2003 (22.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/374,345 22 April 2002 (22.04.2002) US

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, OH
45202 (US).

(72) Inventors: **WELLS, Robert, Lee**; 4245 Rose Hill Avenue, Cincinnati, OH 45229 (US). **JOHNSON, Eric, Scott**; 4575 Aspen Drive, Hamilton, OH 45011 (US). **ROYCE, Douglas, Allan**; 10424 US 50, Aurora, IN 47001 (US).

(74) Agents: **REED, David, T.** et al.; The Procter & Gamble Company, 6110 Center Hill Road, Cincinnati, OH 45224 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SHAMPOO CONTAINING A CATIONIC POLYMER AND ANTI-DANDRUFF PARTICLES

(57) Abstract: The compositions of the present invention relate to improved shampoo compositions having from about 5 to about 50 weight percent of a detergent surfactant, at least about 0.1 weight percent of anti-dandruff particles, at least about 0.05 weight percent of a cationic polysaccharide polymer having a molecular weight of from about 10,000 to about 10,000,000 and a charge density from about 1.4 meq/gm to about 7.0 meq/gm, and at least about 20.0 weight percent of an aqueous carrier.



WO 03/088940 A1

SHAMPOO CONTAINING A CATIONIC POLYMER AND ANTI-DANDRUFF PARTICLES

FIELD

The present invention relates to a hair cleansing shampoo containing anti-dandruff particles. More specifically, it relates to a shampoo containing a cationic polymer having a charge density of at least 1.4 meq/g and anti-dandruff particles.

BACKGROUND

Shampoo compositions for cleaning the hair which also contain anti-dandruff agents are well known. Among the preferred types of anti-dandruff agents are particulate, crystalline anti-dandruff agents, such as sulfur, selenium disulfide, and heavy metal salts of pyridinethione. These particulates relieve dandruff symptoms when they are deposited on the scalp in the course of using the shampoo. Therefore, it is highly desirable to have rinse-off shampoo compositions capable of depositing an effective level of anti-dandruff particles to the scalp.

Compositions intended to deposit solid particle benefit agents to hair or skin surfaces are known; however, the efficiency of deposition has heretofore been unacceptable, requiring either an excess of the solid particle agent in the composition to affect delivery or an imperceivable or unacceptable level of the benefit to be obtained. The efficient deposition and retention of solid particle benefit agents is particularly difficult from compositions intended for cleansing or washing of surfaces, such as shampoos or other personal cleansing products, which contain surfactants and other ingredients which are used to solubilize, suspend and remove particle and oily substances from the surfaces treated therewith. It remains, nonetheless, highly desirable to provide the benefits and convenience afforded through deposition of anti-dandruff particles via use of a simple, washing composition.

Cleansing compositions containing cationic polymers to improve deposition of certain conditioning oils, such as silicone oils, capable of imparting conditioning or slip properties to surfaces treated therewith are known. These conditioning oils, however, are limited in the range of physical, optical, and aesthetic benefits they provide. Additionally, it is known that the viscosity, particle size and other factors associated with the conditioning oils can significantly affect their ability to deposit from cleansing compositions. It is also known to include solid particles in compositions containing cationic polymers; however, these particles are often added to modify the appearance or stability of the composition itself and are not deposited along with the conditioning oils or cationic polymers to the surface treated therewith. When deposition of solid particle benefit agents from washing compositions is intended, the compositions available

heretofore have suffered from the drawbacks of inefficient deposition requiring use of excess amounts of the particle agent or ineffective benefit delivery. It has also been attempted to make specific modifications to solid particle benefit agents to improve their deposition efficiency or retention from rinse-off compositions; however this approach can negatively impact the inherent properties, availability, utility, and cost of the solid particle benefit agents to be used.

It remains, therefore, highly desirable to have a rinse-off composition, preferably a cleansing composition, capable of containing and effectively depositing and retaining anti-dandruff particles on the scalp. It has now been discovered that select cationic polymers, when used in the cleansing compositions of the present invention, can surprisingly enhance the deposition and retention of anti-dandruff particles on the surfaces treated therewith.

SUMMARY

The present invention is directed to a shampoo composition comprising:

- a) from about 5 to about 50 weight percent of a deterative surfactant,
- b) at least about 0.1 weight percent of anti-dandruff particles,
- c) at least about 0.05 weight percent of a cationic polysaccharide polymer having a molecular weight of from about 10,000 to about 10,000,000 and a charge density from about 1.4 meq/gm to about 7.0 meq/gm, and
- d) at least about 20.0 weight percent of an aqueous carrier.

The present invention is further directed to a method of using the shampoo composition.

These and other features, aspects, and advantages of the present invention will become evident to those skilled in the art from a reading of the present disclosure.

DETAILED DESCRIPTION

While the specification concludes with claims which particularly point out and distinctly claim the invention, it is believed the present invention will be better understood from the following description.

The shampoo compositions of the present invention include deterative surfactant, anti-dandruff particles, a cationic polymer and an aqueous carrier. Each of these essential components, as well as preferred or optional components, are described in detail hereinafter.

All percentages, parts and ratios are based upon the total weight of the compositions of the present invention, unless otherwise specified. All such weights as they pertain to listed ingredients are based on the active level and, therefore, do not include solvents or by-products that may be included in commercially available materials, unless otherwise specified.

All molecular weights as used herein are weight average molecular weights expressed as grams/mole, unless otherwise specified.

The term "charge density", as used herein, refers to the ratio of the number of positive charges on a monomeric unit of which a polymer is comprised to the molecular weight of said monomeric unit. The charge density multiplied by the polymer molecular weight determines the number of positively charged sites on a given polymer chain.

Herein, "comprising" means that other steps and other ingredients which do not affect the end result can be added. This term encompasses the terms "consisting of" and "consisting essentially of". The compositions and methods/processes of the present invention can comprise, consist of, and consist essentially of the essential elements and limitations of the invention described herein, as well as any of the additional or optional ingredients, components, steps, or limitations described herein.

The term "polymer" as used herein shall include materials whether made by polymerization of one type of monomer or made by two (*i.e.*, copolymers) or more types of monomers.

The term "solid particle" as used herein means a particle that is not a liquid or a gas.

The term "suitable for application to human hair" as used herein, means that the compositions or components thereof so described are suitable for use in contact with human hair and the scalp and skin without undue toxicity, incompatibility, instability, allergic response, and the like.

The term "water soluble" as used herein, means that the polymer is soluble in water in the present composition. In general, the polymer should be soluble at 25° C at a concentration of 0.1% by weight of the water solvent, preferably at 1%, more preferably at 5%, most preferably at 15%.

All cited references are incorporated herein by reference in their entireties. Citation of any reference is not an admission regarding any determination as to its availability as prior art to the claimed invention.

A. Detersive Surfactant

The shampoo composition of the present invention includes a detersive surfactant. The detersive surfactant component is included to provide cleaning performance to the composition. The detersive surfactant component in turn comprises anionic detersive surfactant, zwitterionic or amphoteric detersive surfactant, or a combination thereof. Such surfactants should be physically and chemically compatible with the essential components described herein, or should not otherwise unduly impair product stability, aesthetics or performance.

Suitable anionic detersive surfactant components for use in the shampoo composition herein include those which are known for use in hair care or other personal care cleansing compositions. The concentration of the anionic surfactant component in the shampoo

composition should be sufficient to provide the desired cleaning and lather performance, and generally range from about 5% to about 50%, preferably from about 8% to about 30%, more preferably from about 10% to about 25%, even more preferably from about 12% to about 22%, by weight of the composition.

Preferred anionic surfactants suitable for use in the shampoo compositions are the alkyl and alkyl ether sulfates. These materials have the respective formulae ROSO_3M and $\text{RO}(\text{C}_2\text{H}_4\text{O})_x\text{SO}_3\text{M}$, wherein R is alkyl or alkenyl of from about 8 to about 18 carbon atoms, x is an integer having a value of from 1 to 10, and M is a cation such as ammonium, alkanolamines, such as triethanolamine, monovalent metals, such as sodium and potassium, and polyvalent metal cations, such as magnesium, and calcium. Solubility of the surfactant will depend upon the particular anionic detergent surfactants and cations chosen.

Preferably, R has from about 8 to about 18 carbon atoms, more preferably from about 10 to about 16 carbon atoms, even more preferably from about 12 to about 14 carbon atoms, in both the alkyl and alkyl ether sulfates. The alkyl ether sulfates are typically made as condensation products of ethylene oxide and monohydric alcohols having from about 8 to about 24 carbon atoms. The alcohols can be synthetic or they can be derived from fats, e.g., coconut oil, palm kernel oil, tallow. Lauryl alcohol and straight chain alcohols derived from coconut oil or palm kernel oil are preferred. Such alcohols are reacted with between about 0 and about 10, preferably from about 2 to about 5, more preferably about 3, molar proportions of ethylene oxide, and the resulting mixture of molecular species having, for example, an average of 3 moles of ethylene oxide per mole of alcohol, is sulfated and neutralized.

Specific non limiting examples of alkyl ether sulfates which may be used in the shampoo compositions of the present invention include sodium and ammonium salts of coconut alkyl triethylene glycol ether sulfate, tallow alkyl triethylene glycol ether sulfate, and tallow alkyl hexa-oxyethylene sulfate. Highly preferred alkyl ether sulfates are those comprising a mixture of individual compounds, wherein the compounds in the mixture have an average alkyl chain length of from about 10 to about 16 carbon atoms and an average degree of ethoxylation of from about 1 to about 4 moles of ethylene oxide.

Other suitable anionic detergent surfactants are the water-soluble salts of organic, sulfuric acid reaction products conforming to the formula $[\text{R}^1\text{-SO}_3\text{-M}]$ where R^1 is a straight or branched chain, saturated, aliphatic hydrocarbon radical having from about 8 to about 24, preferably about 10 to about 18, carbon atoms; and M is a cation described hereinbefore. Non limiting examples of such detergent surfactants are the salts of an organic sulfuric acid reaction product of a hydrocarbon of the methane series, including iso-, neo-, and n-paraffins, having from about 8 to about 24 carbon atoms, preferably about 12 to about 18 carbon atoms and a


sulfonating agent, e.g., SO_3 , H_2SO_4 , obtained according to known sulfonation methods, including bleaching and hydrolysis. Preferred are alkali metal and ammonium sulfonated C_{10} to C_{18} n-paraffins.

Still other suitable anionic deterative surfactants are the reaction products of fatty acids esterified with isethionic acid and neutralized with sodium hydroxide where, for example, the fatty acids are derived from coconut oil or palm kernel oil; sodium or potassium salts of fatty acid amides of methyl tauride in which the fatty acids, for example, are derived from coconut oil or palm kernel oil. Other similar anionic surfactants are described in U.S. Pat. Nos. 2,486,921; 2,486,922; and 2,396,278, which descriptions are incorporated herein by reference.

Other anionic deterative surfactants suitable for use in the shampoo compositions are the succinates, examples of which include disodium N-octadecylsulfosuccinate; disodium lauryl sulfosuccinate; diammonium lauryl sulfosuccinate; tetrasodium N-(1,2-dicarboxyethyl)-N-octadecylsulfosuccinate; diamyl ester of sodium sulfosuccinic acid; dihexyl ester of sodium sulfosuccinic acid; and dioctyl esters of sodium sulfosuccinic acid.

Other suitable anionic deterative surfactants include olefin sulfonates having about 10 to about 24 carbon atoms. In this context, the term "olefin sulfonates" refers to compounds which can be produced by the sulfonation of alpha-olefins by means of uncomplexed sulfur trioxide, followed by neutralization of the acid reaction mixture in conditions such that any sulfones which have been formed in the reaction are hydrolyzed to give the corresponding hydroxy-alkanesulfonates. The sulfur trioxide can be liquid or gaseous, and is usually, but not necessarily, diluted by inert diluents, for example by liquid SO_2 , chlorinated hydrocarbons, etc., when used in the liquid form, or by air, nitrogen, gaseous SO_2 , etc., when used in the gaseous form. The alpha-olefins from which the olefin sulfonates are derived are mono-olefins having from about 10 to about 24 carbon atoms, preferably from about 12 to about 16 carbon atoms. Preferably, they are straight chain olefins. In addition to the true alkene sulfonates and a proportion of hydroxy-alkanesulfonates, the olefin sulfonates can contain minor amounts of other materials, such as alkene disulfonates depending upon the reaction conditions, proportion of reactants, the nature of the starting olefins and impurities in the olefin stock and side reactions during the sulfonation process. A non limiting example of such an alpha-olefin sulfonate mixture is described in U.S. Patent 3,332,880, which description is incorporated herein by reference.

Another class of anionic deterative surfactants suitable for use in the shampoo compositions are the beta-alkyloxy alkane sulfonates. These surfactants conform to the formula



where R¹ is a straight chain alkyl group having from about 6 to about 20 carbon atoms, R² is a lower alkyl group having from about 1 to about 3 carbon atoms, preferably 1 carbon atom, and M is a water-soluble cation as described hereinbefore.

Preferred anionic deterative surfactants for use in the shampoo compositions include ammonium lauryl sulfate, ammonium laureth sulfate, triethylamine lauryl sulfate, triethylamine laureth sulfate, triethanolamine lauryl sulfate, triethanolamine laureth sulfate, monoethanolamine lauryl sulfate, monoethanolamine laureth sulfate, diethanolamine lauryl sulfate, diethanolamine laureth sulfate, lauric monoglyceride sodium sulfate, sodium lauryl sulfate, sodium laureth sulfate, potassium lauryl sulfate, potassium laureth sulfate, sodium lauryl sarcosinate, sodium lauroyl sarcosinate, lauryl sarcosine, cocoyl sarcosine, ammonium cocoyl sulfate, ammonium lauroyl sulfate, sodium cocoyl sulfate, sodium lauroyl sulfate, potassium cocoyl sulfate, potassium lauryl sulfate, triethanolamine lauryl sulfate, triethanolamine lauryl sulfate, monoethanolamine cocoyl sulfate, monoethanolamine lauryl sulfate, sodium tridecyl benzene sulfonate, sodium dodecyl benzene sulfonate, and combinations thereof.

Suitable amphoteric or zwitterionic deterative surfactants for use in the shampoo composition herein include those which are known for use in hair care or other personal care cleansing. Concentration of such amphoteric deterative surfactants preferably ranges from about 0.5% to about 20%, preferably from about 1% to about 10%, by weight of the composition. Non limiting examples of suitable zwitterionic or amphoteric surfactants are described in U.S. Pat. Nos. 5,104,646 (Bolich Jr. et al.), 5,106,609 (Bolich Jr. et al.), which descriptions are incorporated herein by reference.

Amphoteric deterative surfactants suitable for use in the shampoo composition are well known in the art, and include those surfactants broadly described as derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight or branched chain and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic water solubilizing group such as carboxy, sulfonate, sulfate, phosphate, or phosphonate. Preferred amphoteric deterative surfactants for use in the present invention include cocoamphoacetate, cocoamphodiacetate, lauroamphoacetate, lauroamphodiacetate, and mixtures thereof.

Zwitterionic deterative surfactants suitable for use in the shampoo composition are well known in the art, and include those surfactants broadly described as derivatives of aliphatic

quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight or branched chain, and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic group such as carboxy, sulfonate, sulfate, phosphate or phosphonate. Zwitterionics such as betaines are preferred.

The shampoo compositions of the present invention may further comprise additional surfactants for use in combination with the anionic deterative surfactant component described hereinbefore. Suitable optional surfactants include nonionic surfactants. Any such surfactant known in the art for use in hair or personal care products may be used, provided that the optional additional surfactant is also chemically and physically compatible with the essential components of the shampoo composition, or does not otherwise unduly impair product performance, aesthetics or stability. The concentration of the optional additional surfactants in the shampoo composition may vary with the cleansing or lather performance desired, the optional surfactant selected, the desired product concentration, the presence of other components in the composition, and other factors well known in the art.

Non limiting examples of other anionic, zwitterionic, amphoteric or optional additional surfactants suitable for use in the shampoo compositions are described in McCutcheon's, Emulsifiers and Detergents, 1989 Annual, published by M. C. Publishing Co., and U.S. Pat. Nos. 3,929,678, 2,658,072; 2,438,091; 2,528,378, which descriptions are incorporated herein by reference.

B. Anti-Dandruff Particles

The composition of the present invention includes anti-dandruff particles. The particles of the present invention preferably have a particle size of less than 300 μm . Typically, the particles will have a particle size from about 0.01 μm to about 80 μm , still more preferably from about 0.1 μm to about 70 μm , and even more preferably from about 1 μm to about 60 μm in diameter. Suitable, non-limiting examples of anti-dandruff particulates include: pyridinethione salts, selenium sulfide, particulate sulfur, and mixtures thereof. Preferred are pyridinethione salts.

1. Pyridinethione salts

Pyridinethione anti-dandruff particulates, especially 1-hydroxy-2-pyridinethione salts, are highly preferred particulate anti-dandruff agents for use in shampoo compositions of the present invention. The concentration of pyridinethione anti-dandruff particulate typically ranges from about 0.1% to about 4%, by weight of the composition, preferably from about 0.1% to about 3%, most preferably from about 0.3% to about 2%. Preferred pyridinethione salts include those formed from heavy metals such as zinc, tin, cadmium, magnesium, aluminum and zirconium, preferably zinc, more preferably the zinc salt of 1-hydroxy-2-pyridinethione (known

as "zinc pyridinethione" or "ZPT"), most preferably 1-hydroxy-2-pyridinethione salts in platelet particle form, wherein the particles have an average size of up to about 20 μ , preferably up to about 5 μ , most preferably up to about 2.5 μ . Salts formed from other cations, such as sodium, may also be suitable. Pyridinethione anti-dandruff agents are described, for example, in U.S. Pat. No. 2,809,971; U.S. Pat. No. 3,236,733; U.S. Pat. No. 3,753,196; U.S. Pat. No. 3,761,418; U.S. Pat. No. 4,345,080; U.S. Pat. No. 4,323,683; U.S. Pat. No. 4,379,753; and U.S. Pat. No. 4,470,982, all of which are incorporated herein by reference. It is contemplated that when ZPT is used as the anti-dandruff particulate in the shampoo compositions herein, that the growth or re-growth of hair may be stimulated or regulated, or both, or that hair loss may be reduced or inhibited, or that hair may appear thicker or fuller.

2. Selenium sulfide

Selenium sulfide is a particulate anti-dandruff agent suitable for use in the shampoo compositions of the present invention, effective concentrations of which range from about 0.1% to about 4%, by weight of the composition, preferably from about 0.3% to about 2.5%, more preferably from about 0.5% to about 1.5%. Selenium sulfide is generally regarded as a compound having one mole of selenium and two moles of sulfur, although it may also be a cyclic structure that conforms to the general formula Se_xS_y , wherein $x + y = 8$. Average particle diameters for the selenium sulfide are typically less than 15 μ m, as measured by forward laser light scattering device (e.g. Malvern 3600 instrument), preferably less than 10 μ m. Selenium sulfide compounds are described, for example, in U.S. Pat. No. 2,694,668; U.S. Pat. No. 3,152,046; U.S. Pat. No. 4,089,945; and U.S. Pat. No. 4,885,107, all of which descriptions are incorporated herein by reference.

3. Sulfur

Sulfur may also be used as a particulate anti-dandruff agent in the shampoo compositions of the present invention. Effective concentrations of the particulate sulfur are typically from about 1% to about 4%, by weight of the composition, preferably from about 2% to about 4%.

Other Anti-Microbial Actives

In addition to the anti-microbial active selected from pyridinethione salts, selenium sulfide, particulate sulfur, and mixtures thereof, the present invention may further comprise one or more anti-fungal or anti-microbial actives. Suitable anti-microbial actives include coal tar, sulfur, whitfield's ointment, castellani's paint, aluminum chloride, gentian violet, octopirox

(piroctone olamine), ciclopirox olamine, undecylenic acid and its metal salts, potassium permanganate, selenium sulphide, sodium thiosulfate, propylene glycol, oil of bitter orange, urea preparations, griseofulvin, 8-Hydroxyquinoline ciloquinol, thiobendazole, thiocarbamates, haloprogin, polyenes, hydroxypyridone, morpholine, benzylamine, allylamines (such as terbinafine), tea tree oil, clove leaf oil, coriander, palmarosa, berberine, thyme red, cinnamon oil, cinnamic aldehyde, citronellic acid, hinokitol, ichthyol pale, Sensiva SC-50, Elestab HP-100, azelaic acid, lyticase, iodopropynyl butylcarbamate (IPBC), isothiazalinones such as octyl isothiazalinone and azoles, and combinations thereof. Preferred anti-microbials include itraconazole, ketoconazole, selenium sulphide and coal tar.

Azoles

Azole anti-microbials include imidazoles such as benzimidazole, benzothiazole, bifonazole, butaconazole nitrate, climbazole, clotrimazole, croconazole, eberconazole, econazole, elubiol, fenticonazole, fluconazole, flutimazole, isoconazole, ketoconazole, lanconazole, metronidazole, miconazole, neticonazole, omoconazole, oxiconazole nitrate, sertaconazole, sulconazole nitrate, tioconazole, thiazole, and triazoles such as terconazole and itraconazole, and combinations thereof. When present in the composition, the azole anti-microbial active is included in an amount from about 0.01% to about 5%, preferably from about 0.1% to about 3%, and more preferably from about 0.3% to about 2%, by weight of the composition. Especially preferred herein is ketoconazole.

Keratolytic Agents

The present invention may further comprise one or more keratolytic agents such as Salicylic Acid.

Additional anti-microbial actives of the present invention may include extracts of melaleuca (tea tree) and charcoal. The present invention may also comprise combinations of anti-microbial actives. Such combinations may include octopirox and zinc pyrithione combinations, pine tar and sulfur combinations, salicylic acid and zinc pyrithione combinations, octopirox and climbazole combinations, and salicylic acid and octopirox combinations, and mixtures thereof.


C. Cationic Polymer

The composition of the present invention includes a cationic deposition polymer of sufficiently high cationic charge density to effectively enhance deposition of the anti-dandruff particle component described herein. Suitable cationic polymers will have cationic charge densities of at least about 1.4 meq/gm, preferably at least about 1.7 meq/gm, more preferably at least about 1.9 meq/gm, but also preferably less than about 7 meq/gm, more preferably less than about 5 meq/gm, at the pH of intended use of the shampoo composition, which pH will

generally range from about pH 3 to about pH 9, preferably between about pH 4 and about pH 8. The average molecular weight of such suitable cationic polymers will generally be between about 10,000 and 10 million, preferably between about 50,000 and about 5 million, more preferably between about 100,000 and about 3 million. The "cationic charge density" of a polymer, as that term is used herein, refers to the ratio of the number of positive charges on a monomeric unit of which the polymer is comprised to the molecular weight of said monomeric unit. The cationic charge density multiplied by the polymer molecular weight determines the number of positively charged sites on a given polymer chain.

The concentration of the cationic polymer in the shampoo composition ranges from about 0.05% to about 3%, preferably from about 0.075% to about 2.0%, more preferably from about 0.1% to about 1.0%, by weight of the shampoo composition. The weight ratio of cationic polymer to anti-dandruff particle (described hereinafter) in the shampoo compositions is preferably from about 2:1 to about 1:30, more preferably from about 1:1 to about 1:20, still more preferably from about 1:2 to about 1:10.

Cationic polymers useful in the present invention are polysaccharide polymers, such as cationic cellulose derivatives and cationic starch derivatives. Suitable cationic polysaccharide polymers include those which conform to the formula:



wherein A is an anhydroglucose residual group, such as a starch or cellulose anhydroglucose residual; R is an alkylene oxyalkylene, polyoxyalkylene, or hydroxyalkylene group, or combination thereof; R¹, R², and R³ independently are alkyl, aryl, alkylaryl, arylalkyl, alkoxyalkyl, or alkoxyaryl groups, each group containing up to about 18 carbon atoms, and the total number of carbon atoms for each cationic moiety (i.e., the sum of carbon atoms in R¹, R² and R³) preferably being about 20 or less; and X is an anionic counterion. Any anionic counterions can be used in association with the cationic polymers of the present invention so long as the polymers remain soluble in water, in the shampoo composition, or in a coacervate phase of the shampoo composition, and so long as the counterions are physically and chemically compatible with the essential components of the shampoo composition or do not otherwise unduly impair product performance, stability or aesthetics. Non limiting examples of such counterions include halides (e.g., chlorine, fluorine, bromine, iodine), sulfate and methylsulfate. The degree of cationic substitution in these polysaccharide polymers is typically from about 0.01-1 cationic groups per anhydroglucose unit.

Preferred cationic cellulose polymers salts of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide, referred to in the industry (CTFA) as Polyquaternium 10 and available from Amerchol Corp. (Edison, N.J., USA) as Polymer KG30M with a charge density of 1.9 and a molecular weight of ~ 1.25 million.

The cationic polymers herein are either soluble in the shampoo composition or are soluble in a complex coacervate phase in the shampoo composition formed by the cationic polymer and the anionic deterative surfactant component described hereinbefore. Complex coacervates of the cationic polymer can also be formed with other charged materials in the shampoo composition.

Coacervate formation is dependent upon a variety of criteria such as molecular weight, component concentration, and ratio of interacting ionic components, ionic strength (including modification of ionic strength, for example, by addition of salts), charge density of the cationic and anionic components, pH, and temperature. Coacervate systems and the effect of these parameters have been described, for example, by J. Caelles, et al., "Anionic and Cationic Compounds in Mixed Systems", *Cosmetics & Toiletries*, Vol. 106, April 1991, pp 49-54, C. J. van Oss, "Coacervation, Complex-Coacervation and Flocculation", *J. Dispersion Science and Technology*, Vol. 9 (5,6), 1988-89, pp 561-573, and D. J. Burgess, "Practical Analysis of Complex Coacervate Systems", *J. of Colloid and Interface Science*, Vol. 140, No. 1, November 1990, pp 227-238, which descriptions are incorporated herein by reference.

It is believed to be particularly advantageous for the cationic polymer to be present in the shampoo composition in a coacervate phase, or to form a coacervate phase upon application or rinsing of the shampoo to or from the hair. Complex coacervates are believed to more readily deposit on the hair. Thus, in general, it is preferred that the cationic polymer exist in the shampoo composition as a coacervate phase or form a coacervate phase upon dilution.

Techniques for analysis of formation of complex coacervates are known in the art. For example, microscopic analyses of the shampoo compositions, at any chosen stage of dilution, can be utilized to identify whether a coacervate phase has formed. Such coacervate phase will be identifiable as an additional emulsified phase in the composition. The use of dyes can aid in distinguishing the coacervate phase from other insoluble phases dispersed in the shampoo composition.

In the compositions of the present invention, it is believed that the tendency for high charge density cationic polymers to form relatively large coacervates of sizes ranging from about 20 microns to about 500 microns which are capable of effectively binding or flocculating with the particle and enhancing delivery to hair contributes to the superior deposition efficiency. Additionally, coacervates which have a cohesive character as evidenced by large, structured

flocs which retain a substantial amount of the particle component on dilution and resist deflocculation on exposure to shear enhance the deposition and retention of particles on hair.

D. Aqueous Carrier

The compositions of the present invention include an aqueous carrier. The level and species of the carrier are selected according to the compatibility with other components, and other desired characteristic of the product.

Carriers useful in the present invention include water and water solutions of lower alkyl alcohols. Lower alkyl alcohols useful herein are monohydric alcohols having 1 to 6 carbons, more preferably ethanol and isopropanol.

Preferably, the aqueous carrier is substantially water. Deionized water is preferably used. Water from natural sources containing mineral cations can also be used, depending on the desired characteristic of the product. Generally, the compositions of the present invention comprise from about 20% to about 99%, preferably from about 40% to about 98%, and more preferably from about 60% to about 98% aqueous carrier.

The pH of the present composition is preferably from about 4 to about 9, more preferably from about 4.5 to about 7.5. Buffers and other pH adjusting agents can be included to achieve the desirable pH.

E. Additional Components

The shampoo compositions of the present invention may further comprise one or more optional components known for use in hair care or personal care products, provided that the optional components are physically and chemically compatible with the essential components described herein, or do not otherwise unduly impair product stability, aesthetics or performance. Individual concentrations of such optional components may range from about 0.001% to about 10% by weight of the shampoo compositions.

Non-limiting examples of optional components for use in the shampoo composition include cationic polymers, conditioning agents (hydrocarbon oils, fatty esters, silicones), suspending agents, viscosity modifiers, dyes, nonvolatile solvents or diluents (water soluble and insoluble), pearlescent aids, foam boosters, additional surfactants or nonionic cosurfactants, pediculocides, pH adjusting agents, perfumes, preservatives, chelants, proteins, skin active agents, sunscreens, UV absorbers, and vitamins.

Conditioning agents

Conditioning agents include any material which is used to give a particular conditioning benefit to hair and/or skin. In hair treatment compositions, suitable conditioning agents are those which deliver one or more benefits relating to shine, softness, combability, antistatic properties, wet-handling, damage, manageability, body, and greasiness. The conditioning

agents useful in the shampoo compositions of the present invention typically comprise a water insoluble, water dispersible, non-volatile, liquid that forms emulsified, liquid particles or are solubilized by the surfactant micelles, in the anionic detergent surfactant component (described above). Suitable conditioning agents for use in the shampoo composition are those conditioning agents characterized generally as silicones (e.g. silicone oils, cationic silicones, silicone gums, high refractive silicones, and silicone resins), organic conditioning oils (e.g. hydrocarbon oils, polyolefins, and fatty esters) or combinations thereof, or those conditioning agents which otherwise form liquid, dispersed, particles in the aqueous surfactant matrix herein. Such conditioning agents should be physically and chemically compatible with the essential components of the composition, and should not otherwise unduly impair product stability, aesthetics or performance.

The concentration of the conditioning agent in the shampoo composition should be sufficient to provide the desired conditioning benefits, and as will be apparent to one of ordinary skill in the art. Such concentration can vary with the conditioning agent, the conditioning performance desired, the average size of the conditioning agent particles, the type and concentration of other components, and other like factors.

1. Silicones

The conditioning agent of the shampoo compositions of the present invention is preferably an insoluble silicone conditioning agent. The silicone conditioning agent particles may comprise volatile silicone, non-volatile silicone, or combinations thereof. Preferred are non-volatile silicone conditioning agents. If volatile silicones are present, it will typically be incidental to their use as a solvent or carrier for commercially available forms of non-volatile silicone materials ingredients, such as silicone gums and resins. The silicone conditioning agent particles may comprise a silicone fluid conditioning agent and may also comprise other ingredients, such as a silicone resin to improve silicone fluid deposition efficiency or enhance glossiness of the hair (especially when high refractive index (e.g. above about 1.46) silicone conditioning agents are used (e.g. highly phenylated silicones).

The concentration of the silicone conditioning agent typically ranges from about 0.01% to about 10%, by weight of the composition, preferably from about 0.1% to about 8%, more preferably from about 0.1% to about 5%, most preferably from about 0.2% to about 3%. Non-limiting examples of suitable silicone conditioning agents, and optional suspending agents for the silicone, are described in U.S. Reissue Pat. No. 34,584, U.S. Pat. No. 5,104,646, and U.S. Pat. No. 5,106,609, which descriptions are incorporated herein by reference. The silicone conditioning agents for use in the shampoo compositions of the present invention preferably have a viscosity, as measured at 25°C, from about 20 to about 2,000,000 centistokes ("cSk"),

more preferably from about 1,000 to about 1,800,000 csk, even more preferably from about 50,000 to about 1,500,000 csk, most preferably from about 100,000 to about 1,500,000 csk.


The dispersed silicone conditioning agent particles typically have a number average particle diameter ranging from about 0.01 μ m to about 50 μ m. For small particle application to hair, the number average particle diameters typically range from about 0.01 μ m to about 4 μ m, preferably from about 0.01 μ m to about 2 μ m, more preferably from about 0.01 μ m to about 0.5 μ m. For larger particle application to hair, the number average particle diameters typically range from about 4 μ m to about 50 μ m, preferably from about 6 μ m to about 30 μ m, more preferably from about 9 μ m to about 20 μ m, most preferably from about 12 μ m to about 18 μ m. Conditioning agents having an average particle size of less than about 5 μ m may deposit more efficiently on the hair. It is believed that small size particles of conditioning agent are contained within the coacervate that is formed between the anionic surfactant component (described above) and the cationic polymer component (described below), upon dilution of the shampoo.

Background material on silicones including sections discussing silicone fluids, gums, and resins, as well as manufacture of silicones, are found in *Encyclopedia of Polymer Science and Engineering*, vol. 15, 2d ed., pp 204-308, John Wiley & Sons, Inc. (1989), incorporated herein by reference.

a. Silicone oils

Silicone fluids include silicone oils, which are flowable silicone materials having a viscosity, as measured at 25°C, less than 1,000,000 csk, preferably from about 5 csk to about 1,000,000 csk, more preferably from about 10 csk to about 100,000 csk. Suitable silicone oils for use in the shampoo compositions of the present invention include polyalkyl siloxanes, polyaryl siloxanes, polyalkylaryl siloxanes, polyether siloxane copolymers, and mixtures thereof. Other insoluble, non-volatile silicone fluids having hair conditioning properties may also be used.

Silicone oils include polyalkyl or polyaryl siloxanes which conform to the following Formula (III):



wherein R is aliphatic, preferably alkyl or alkenyl, or aryl, R can be substituted or unsubstituted, and x is an integer from 1 to about 8,000. Suitable unsubstituted R groups for use in the shampoo compositions of the present invention include, but are not limited to: alkoxy, aryloxy, alkaryl, arylalkyl, arylalkenyl, alkamino, and ether-substituted, hydroxyl-substituted, and halogen-substituted aliphatic and aryl groups. Suitable R groups also include cationic amines and quaternary ammonium groups.


The aliphatic or aryl groups substituted on the siloxane chain may have any structure so long as the resulting silicones remain fluid at room temperature, are hydrophobic, are neither irritating, toxic nor otherwise harmful when applied to the hair, are compatible with the other components of the shampoo compositions, are chemically stable under normal use and storage conditions, are insoluble in the shampoo compositions herein, and are capable of being deposited on and conditioning the hair. The two R groups on the silicon atom of each monomeric silicone unit may represent the same or different groups. Preferably, the two R groups represent the same group.

Preferred alkyl and alkenyl substituents are C₁ to C₅ alkyls and alkenyls, more preferably from C₁ to C₄, most preferably from C₁ to C₂. The aliphatic portions of other alkyl-, alkenyl-, or alkynyl-containing groups (such as alkoxy, alkaryl, and alkamino) can be straight or branched chains, and are preferably from C₁ to C₅, more preferably from C₁ to C₄, even more preferably from C₁ to C₃, most preferably from C₁ to C₂. As discussed above, the R substituents can also contain amino functionalities (e.g. alkamino groups), which can be primary, secondary or tertiary amines or quaternary ammonium. These include mono-, di- and tri- alkylamino and alkoxyamino groups, wherein the aliphatic portion chain length is preferably as described above. The R substituents may also be substituted with other groups, such as halogens (e.g. chloride, fluoride, and bromide), halogenated aliphatic or aryl groups, hydroxy (e.g. hydroxy substituted aliphatic groups), and mixtures thereof. Suitable halogenated R groups could include, for example, tri-halogenated (preferably tri-fluoro) alkyl groups such as -R¹CF₃, wherein R¹ is a C₁ - C₃ alkyl. An example of such a polysiloxane includes, but is not limited to, polymethyl 3,3,3-trifluoropropylsiloxane.

Suitable R groups for use in the shampoo compositions of the present invention include, but are not limited to: methyl, ethyl, propyl, phenyl, methylphenyl and phenylmethyl. Specific non-limiting examples of preferred silicones include: polydimethyl siloxane, polydiethylsiloxane, and polymethylphenylsiloxane. Polydimethylsiloxane is especially preferred. Other suitable R groups include: methyl, methoxy, ethoxy, propoxy, and aryloxy. The three R groups on the end caps of the silicone may represent the same or different groups.

Non-volatile polyalkylsiloxane fluids that may be used include, for example, low molecular weight polydimethylsiloxanes. These siloxanes are available, for example, from the General Electric Company in their Viscasil R and SF 96 series, and from Dow Corning in their Dow Corning 200 series. Polyalkylaryl siloxane fluids that may be used, also include, for example, polymethylphenylsiloxanes. These siloxanes are available, for example, from the General Electric Company as SF 1075 methyl phenyl fluid or from Dow Corning as 556 Cosmetic Grade Fluid. Polyether siloxane copolymers that may be used include, for example, a polypropylene oxide modified polydimethylsiloxane (e.g., Dow Corning DC-1248) although ethylene oxide or mixtures of ethylene oxide and propylene oxide may also be used. The ethylene oxide and polypropylene oxide concentrations must be sufficiently low to prevent solubility in water and the composition described herein.


Alkylamino substituted silicones suitable for use in the shampoo compositions of the present invention include, but are not limited to, those which conform to the following general Formula (IV):




wherein x and y are integers. This polymer is also known as "amodimethicone."



b. Cationic silicones

Cationic silicone fluids suitable for use in the shampoo compositions of the present invention include, but are not limited to, those which conform to the general formula (V):




wherein G is hydrogen, phenyl, hydroxy, or C₁-C₈ alkyl, preferably methyl; a is 0 or an integer having a value from 1 to 3, preferably 0; b is 0 or 1, preferably 1; n is a number from 0 to 1,999, preferably from 49 to 149; m is an integer from 1 to 2,000, preferably from 1 to 10; the sum of n and m is a number from 1 to 2,000, preferably from 50 to 150; R₁ is a monovalent radical conforming to the general formula C_qH_{2q}L, wherein q is an integer having a value from 2 to 8 and L is selected from the following groups:






wherein R_2 is hydrogen, phenyl, benzyl, or a saturated hydrocarbon radical, preferably an alkyl radical from about C_1 to about C_{20} , and A^- is a halide ion.

An especially preferred cationic silicone corresponding to formula (V) is the polymer known as "trimethylsilylamodimethicone", which is shown below in formula (VI):



Other silicone cationic polymers which may be used in the shampoo compositions of the present invention are represented by the general formula (VII):



wherein R^3 is a monovalent hydrocarbon radical from C_1 to C_{18} , preferably an alkyl or alkenyl radical, such as methyl; R_4 is a hydrocarbon radical, preferably a C_1 to C_{18} alkylene radical or a C_{10} to C_{18} alkyleneoxy radical, more preferably a C_1 to C_8 alkyleneoxy radical; Q^- is a halide ion, preferably chloride; r is an average statistical value from 2 to 20, preferably from 2 to 8; s is an average statistical value from 20 to 200, preferably from 20 to 50. A preferred polymer of this class is known as UCARE SILICONE ALE 56™, available from Union Carbide.

c. Silicone gums


Other silicone fluids suitable for use in the shampoo compositions of the present invention are the insoluble silicone gums. These gums are polyorganosiloxane materials having a viscosity, as measured at 25°C, of greater than or equal to 1,000,000 csk. Silicone gums are described in U.S. Pat. No. 4,152,416; Noll and Walter, *Chemistry and Technology of Silicones*, New York: Academic Press (1968); and in General Electric Silicone Rubber Product Data

Sheets SE 30, SE 33, SE 54 and SE 76, all of which are incorporated herein by reference. The silicone gums will typically have a weight average molecular weight in excess of about 200,000, preferably from about 200,000 to about 1,000,000. Specific non-limiting examples of silicone gums for use in the shampoo compositions of the present invention include polydimethylsiloxane, (polydimethylsiloxane) (methylvinylsiloxane) copolymer, poly(dimethylsiloxane) (diphenyl siloxane)(methylvinylsiloxane) copolymer and mixtures thereof.

d. High refractive index silicones

Other non-volatile, insoluble silicone fluid conditioning agents that are suitable for use in the shampoo compositions of the present invention are those known as "high refractive index silicones," having a refractive index of at least about 1.46, preferably at least about 1.48, more preferably at least about 1.52, most preferably at least about 1.55. The refractive index of the polysiloxane fluid will generally be less than about 1.70, typically less than about 1.60. In this context, polysiloxane "fluid" includes oils as well as gums.

The high refractive index polysiloxane fluid includes those represented by general Formula (III) above, as well as cyclic polysiloxanes such as those represented by Formula (VIII) below:



wherein R is as defined above, and n is a number from about 3 to about 7, preferably from about 3 to about 5.

The high refractive index polysiloxane fluids contain an amount of aryl-containing R substituents sufficient to increase the refractive index to the desired level, which is described above. Additionally, R and n must be selected so that the material is non-volatile.

Aryl-containing substituents include those which contain alicyclic and heterocyclic five and six member aryl rings and those which contain fused five or six member rings. The aryl rings themselves can be substituted or unsubstituted. Substituents include aliphatic substituents, and may also include alkoxy substituents, acyl substituents, ketones, halogens (e.g., Cl and Br), amines, and the like. Examples of aryl-containing groups include, but are not limited to, substituted and unsubstituted arenes, such as phenyl, and phenyl derivatives, such as phenyls with C₁-C₅ alkyl or alkenyl substituents. Specific non-limiting examples include: allylphenyl, methyl phenyl and ethyl phenyl, vinyl phenyls (e.g. styrenyl), and phenyl alkynes (e.g. phenyl C₂-C₄

alkynes). Heterocyclic aryl groups include, but are not limited to, substituents derived from furan, imidazole, pyrrole, pyridine, and the like. Examples of fused aryl ring substituents include, but are not limited to, naphthalene, coumarin, and purine.

Generally, the high refractive index polysiloxane fluids will have a degree of aryl-containing substituents of at least about 15%, preferably at least about 20%, more preferably at least about 25%, even more preferably at least about 35%, most preferably at least about 50%. Typically, the degree of aryl substitution will be less than about 90%, more generally less than about 85%, preferably from about 55% to about 80%.

The high refractive index polysiloxane fluids are also characterized by relatively high surface tensions as a result of their aryl substitution. Generally, the polysiloxane fluids will have a surface tension of at least about 24 dynes/cm², typically at least about 27 dynes/cm². Surface tension, for purposes hereof, is measured by a de Nouy ring tensiometer according to Dow Corning Corporate Test Method CTM 0461 (23 November, 1971). Changes in surface tension can be measured according to the above test method or according to ASTM Method D 1331.

Preferred high refractive index polysiloxane fluids have a combination of phenyl or phenyl derivative substituents (most preferably phenyl), with alkyl substituents, preferably C₁-C₄ alkyl (most preferably methyl), hydroxy, or C₁-C₄ alkylamino (especially -R¹NHR²NH₂ wherein each R¹ and R² independently is a C₁-C₃ alkyl, alkenyl, and/or alkoxy). High refractive index polysiloxanes are available from Dow Corning, Huls America, and General Electric.

When high refractive index silicones are used in the shampoo compositions of the present invention, they are preferably used in solution with a spreading agent, such as a silicone resin or a surfactant, to reduce the surface tension by a sufficient amount to enhance spreading and thereby enhance the glossiness (subsequent to drying) of hair treated with the compositions. Generally, an amount of the spreading agent is used that is sufficient to reduce the surface tension of the high refractive index polysiloxane fluid by at least about 5%, preferably at least about 10%, more preferably at least about 15%, even more preferably at least about 20%, most preferably at least about 25%. Reductions in surface tension of the polysiloxane fluid/spreading agent mixture may improve shine of the hair.

Also, the spreading agent will preferably reduce the surface tension by at least about 2 dynes/cm², preferably at least about 3 dynes/cm², even more preferably at least about 4 dynes/cm², most preferably at least about 5 dynes/cm².

The surface tension of the mixture of the polysiloxane fluid and the spreading agent, at the proportions present in the final product, is preferably less than or equal to about 30 dynes/cm², more preferably less than or equal to about 28 dynes/cm², most preferably less than

or equal to about 25 dynes/cm². Typically, the surface tension will be in the range from about 15 dynes/cm² to about 30 dynes/cm², more typically from about 18 dynes/cm² to about 28 dynes/cm², and most generally from about 20 dynes/cm² to about 25 dynes/cm².

The weight ratio of the highly arylated polysiloxane fluid to the spreading agent will, in general, be from about 1000:1 to about 1:1, preferably from about 100:1 to about 2:1, more preferably from about 50:1 to about 2:1, most preferably from about 25:1 to about 2:1. When fluorinated surfactants are used, particularly high polysiloxane fluid to spreading agent ratios may be effective due to the efficiency of these surfactants. Thus, it is contemplated that ratios significantly above 1000:1 may be used.

Silicone fluids suitable for use in the shampoo compositions of the present invention are disclosed in U.S. Pat. No. 2,826,551, U.S. Pat. No. 3,964,500, U.S. Pat. No. 4,364,837, British Pat. No. 849,433, and *Silicon Compounds*, Petrarch Systems, Inc. (1984), all of which are incorporated herein by reference.

e. Silicone resins

Silicone resins may be included in the silicone conditioning agent of the shampoo compositions of the present invention. These resins are highly cross-linked polymeric siloxane systems. The cross-linking is introduced through the incorporation of trifunctional and tetrafunctional silanes with monofunctional or difunctional, or both, silanes during manufacture of the silicone resin. As is apparent to one of ordinary skill in the art, the degree of cross-linking that is required in order to result in a silicone resin will vary according to the specific silane units incorporated into the silicone resin. Generally, silicone materials which have a sufficient level of trifunctional and tetrafunctional siloxane monomer units (and hence, a sufficient level of cross-linking) such that they dry down to a rigid, or hard, film are considered to be silicone resins. The ratio of oxygen atoms to silicon atoms is indicative of the level of cross-linking in a particular silicone material. Silicone resins suitable for use in the shampoo compositions of the present invention generally have at least about 1.1 oxygen atoms per silicon atom. Preferably, the ratio of oxygen to silicon atoms is at least about 1.2:1.0. Silanes used in the manufacture of silicone resins include, but are not limited to: monomethyl-, dimethyl-, trimethyl-, monophenyl-, diphenyl-, methylphenyl-, monovinyl-, and methylvinyl-chlorosilanes, and tetra-chlorosilane, with the methyl-substituted silanes being most commonly utilized. Preferred resins are available from General Electric as GE SS4230 and GE SS4267. Commercially available silicone resins are generally supplied in a dissolved form in a low viscosity volatile or non-volatile silicone fluid. The silicone resins for use herein should be supplied and incorporated into the present compositions in such dissolved form, as will be readily apparent to one of ordinary skill in the art.

Silicone materials and silicone resins in particular, can conveniently be identified according to a shorthand nomenclature system known to those of ordinary skill in the art as "MDTQ" nomenclature. Under this system, the silicone is described according to presence of various siloxane monomer units which make up the silicone. Briefly, the symbol M denotes the monofunctional unit $(\text{CH}_3)_3\text{SiO}_{0.5}$; D denotes the difunctional unit $(\text{CH}_3)_2\text{SiO}$; T denotes the trifunctional unit $(\text{CH}_3)\text{SiO}_{1.5}$; and Q denotes the quadra- or tetra-functional unit SiO_2 . Primes of the unit symbols (e.g. M', D', T', and Q') denote substituents other than methyl, and must be specifically defined for each occurrence. Typical alternate substituents include, but are not limited to, groups such as vinyl, phenyls, amines, hydroxyls, and the like. The molar ratios of the various units, either in terms of subscripts to the symbols indicating the total number of each type of unit in the silicone (or an average thereof) or as specifically indicated ratios in combination with molecular weight complete the description of the silicone material under the MDTQ system. Higher relative molar amounts of T, Q, T' and/or Q' to D, D', M and/or M' in a silicone resin indicates higher levels of cross-linking. As discussed above, however, the overall level of cross-linking can also be indicated by the oxygen to silicon ratio.

Preferred silicone resins for use in the shampoo compositions of the present invention include, but are not limited to MQ, MT, MTQ, MDT and MDTQ resins. Methyl is a preferred silicone substituent. Especially preferred silicone resins are MQ resins, wherein the M:Q ratio is from about 0.5:1.0 to about 1.5:1.0 and the average molecular weight of the silicone resin is from about 1000 to about 10,000.

The weight ratio of the non-volatile silicone fluid, having refractive index below 1.46, to the silicone resin component, when used, is preferably from about 4:1 to about 400:1, more preferably from about 9:1 to about 200:1, most preferably from about 19:1 to about 100:1, particularly when the silicone fluid component is a polydimethylsiloxane fluid or a mixture of polydimethylsiloxane fluid and polydimethylsiloxane gum as described above. Insofar as the silicone resin forms a part of the same phase in the compositions hereof as the silicone fluid, i.e. the conditioning active, the sum of the fluid and resin should be included in determining the level of silicone conditioning agent in the composition.

2. Organic conditioning oils

The conditioning component of the shampoo compositions of the present invention may also comprise from about 0.05% to about 3%, by weight of the composition, preferably from about 0.08% to about 1.5%, more preferably from about 0.1% to about 1%, of at least one organic conditioning oil as the conditioning agent, either alone or in combination with other conditioning agents, such as the silicones (described above).

It is believed that these organic conditioning oils provide the shampoo composition with improved conditioning performance when used in combination with the essential components of the composition, and in particular when used in combination with cationic polymers (described below). The conditioning oils may add shine and luster to the hair. Additionally, they may enhance dry combing and dry hair feel. Most or all of these organic conditioning oils are believed to be solubilized in the surfactant micelles of the shampoo composition. It is also believed that this solubilization into the surfactant micelles contributes to the improved hair conditioning performance of the shampoo compositions herein.

The organic conditioning oils suitable for use as the conditioning agent herein are preferably low viscosity, water insoluble, liquids selected from the hydrocarbon oils, polyolefins, fatty esters, and mixtures thereof. The viscosity, as measured at 40°C, of such organic conditioning oils is preferably from about 1 centipoise to about 200 centipoise, more preferably from about 1 centipoise to about 100 centipoise, most preferably from about 2 centipoise to about 50 centipoise.

a. Hydrocarbon oils

Suitable organic conditioning oils for use as conditioning agents in the shampoo compositions of the present invention include, but are not limited to, hydrocarbon oils having at least about 10 carbon atoms, such as cyclic hydrocarbons, straight chain aliphatic hydrocarbons (saturated or unsaturated), and branched chain aliphatic hydrocarbons (saturated or unsaturated), including polymers and mixtures thereof. Straight chain hydrocarbon oils preferably are from about C₁₂ to about C₁₉. Branched chain hydrocarbon oils, including hydrocarbon polymers, typically will contain more than 19 carbon atoms.

Specific non-limiting examples of these hydrocarbon oils include paraffin oil, mineral oil, saturated and unsaturated dodecane, saturated and unsaturated tridecane, saturated and unsaturated tetradecane, saturated and unsaturated pentadecane, saturated and unsaturated hexadecane, polybutene, polydecene, and mixtures thereof. Branched-chain isomers of these compounds, as well as of higher chain length hydrocarbons, can also be used, examples of which include highly branched, saturated or unsaturated, alkanes such as the permethyl-substituted isomers, e.g., the permethyl-substituted isomers of hexadecane and eicosane, such as 2, 2, 4, 4, 6, 6, 8, 8-dimethyl-10-methylundecane and 2, 2, 4, 4, 6, 6-dimethyl-8-methylnonane, available from Permethyl Corporation. Hydrocarbon polymers such as polybutene and polydecene. A preferred hydrocarbon polymer is polybutene, such as the copolymer of isobutylene and butene. A commercially available material of this type is L-14 polybutene from Amoco Chemical Corporation.

b. Polyolefins

Organic conditioning oils for use in the shampoo compositions of the present invention can also include liquid polyolefins, more preferably liquid poly- α -olefins, most preferably hydrogenated liquid poly- α -olefins. Polyolefins for use herein are prepared by polymerization of C_4 to about C_{14} olefinic monomers, preferably from about C_6 to about C_{12} .

Non-limiting examples of olefinic monomers for use in preparing the polyolefin liquids herein include ethylene, propylene, 1-butene, 1-pentene, 1-hexene, 1-octene, 1-decene, 1-dodecene, 1-tetradecene, branched chain isomers such as 4-methyl-1-pentene, and mixtures thereof. Also suitable for preparing the polyolefin liquids are olefin-containing refinery feedstocks or effluents. Preferred hydrogenated α -olefin monomers include, but are not limited to: 1-hexene to 1-hexadecenes, 1-octene to 1-tetradecene, and mixtures thereof.

c. Fatty Esters

Other suitable organic conditioning oils for use as the conditioning agent in the shampoo compositions of the present invention include, but are not limited to, fatty esters having at least 10 carbon atoms. These fatty esters include esters with hydrocarbyl chains derived from fatty acids or alcohols (e.g. mono-esters, polyhydric alcohol esters, and di- and tri-carboxylic acid esters). The hydrocarbyl radicals of the fatty esters hereof may include or have covalently bonded thereto other compatible functionalities, such as amides and alkoxy moieties (e.g., ethoxy or ether linkages, etc.).

Suitable for use in the shampoo compositions of the present invention are alkyl and alkenyl esters of fatty acids having from about C_{10} to about C_{22} aliphatic chains, and alkyl and alkenyl fatty alcohol carboxylic acid esters having a C_{10} to about C_{22} alkyl and/or alkenyl alcohol-derived aliphatic chain, and mixtures thereof. Specific examples of preferred fatty esters include, but are not limited to: isopropyl isostearate, hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate, dihexyldecyl adipate, lauryl lactate, myristyl lactate, cetyl lactate, oleyl stearate, oleyl oleate, oleyl myristate, lauryl acetate, cetyl propionate, and oleyl adipate.

Other fatty esters suitable for use in the shampoo compositions of the present invention are mono-carboxylic acid esters of the general formula $R'COOR$, wherein R' and R are alkyl or alkenyl radicals, and the sum of carbon atoms in R' and R is at least 10, preferably at least 20. The mono-carboxylic acid ester need not necessarily contain at least one chain with at least 10 carbon atoms; rather the total number of aliphatic chain carbon atoms must be at least 10. Specific non-limiting examples of mono-carboxylic acid esters include: isopropyl myristate, glycol stearate, and isopropyl laurate.


Still other fatty esters suitable for use in the shampoo compositions of the present invention are di- and tri-alkyl and alkenyl esters of carboxylic acids, such as esters of C_4 to C_8

dicarboxylic acids (e.g. C₁ to C₂₂ esters, preferably C₁ to C₆, of succinic acid, glutaric acid, adipic acid, hexanoic acid, heptanoic acid, and octanoic acid). Specific non-limiting examples of di- and tri- alkyl and alkenyl esters of carboxylic acids include isocetyl stearyl stearate, diisopropyl adipate, and tristearyl citrate.


Other fatty esters suitable for use in the shampoo compositions of the present invention are those known as polyhydric alcohol esters. Such polyhydric alcohol esters include alkylene glycol esters, such as ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty acid esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters.

Still other fatty esters suitable for use in the shampoo compositions of the present invention are glycerides, including, but not limited to, mono-, di-, and tri-glycerides, preferably di- and tri-glycerides, most preferably triglycerides. For use in the shampoo compositions described herein, the glycerides are preferably the mono-, di-, and tri-esters of glycerol and long chain carboxylic acids, such as C₁₀ to C₂₂ carboxylic acids. A variety of these types of materials can be obtained from vegetable and animal fats and oils, such as castor oil, safflower oil, cottonseed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, lanolin and soybean oil. Synthetic oils include, but are not limited to, triolein and tristearin glyceryl dilaurate.

Other fatty esters suitable for use in the shampoo compositions of the present invention are water insoluble synthetic fatty esters. Some preferred synthetic esters conform to the general Formula (IX):



wherein R¹ is a C₇ to C₉ alkyl, alkenyl, hydroxyalkyl or hydroxyalkenyl group, preferably a saturated alkyl group, more preferably a saturated, linear, alkyl group; n is a positive integer having a value from 2 to 4, preferably 3; and Y is an alkyl, alkenyl, hydroxy or carboxy substituted alkyl or alkenyl, having from about 2 to about 20 carbon atoms, preferably from about 3 to about 14 carbon atoms. Other preferred synthetic esters conform to the general Formula (X):



wherein R² is a C₈ to C₁₀ alkyl, alkenyl, hydroxyalkyl or hydroxyalkenyl group; preferably a saturated alkyl group, more preferably a saturated, linear, alkyl group; n and Y are as defined above in Formula (X).

It is believed that the preferred synthetic esters provide improved wet hair feel when used in combination with the essential components of the shampoo compositions of the present invention, particularly when used in combination with the cationic polymer component (described below). These synthetic esters improve wet hair feel by reducing the slimy or excessively conditioned feel of wet hair that has been conditioned by a cationic polymer.

Specific non-limiting examples of suitable synthetic fatty esters for use in the shampoo compositions of the present invention include: P-43 (C₈-C₁₀ triester of trimethylolpropane), MCP-684 (tetraester of 3,3 diethanol-1,5 pentadiol), MCP 121 (C₈-C₁₀ diester of adipic acid), all of which are available from Mobil Chemical Company.

3. Other conditioning agents

Also suitable for use in the compositions herein are the conditioning agents described by the Procter & Gamble Company in U.S. Pat. Nos. 5,674,478, and 5,750,122, both of which are incorporated herein in their entirety by reference. Also suitable for use herein are those conditioning agents described in U.S. Pat. Nos. 4,529,586 (Clairol), 4,507,280 (Clairol), 4,663,158 (Clairol), 4,197,865 (L'Oreal), 4,217, 914 (L'Oreal), 4,381,919 (L'Oreal), and 4,422, 853 (L'Oreal), all of which descriptions are incorporated herein by reference.

Some other preferred silicone conditioning agents for use in the compositions of the present invention include: Abil[®] S 201 (dimethicone/sodium PG-propyldimethicone thiosulfate copolymer), available from Goldschmidt; DC Q2-8220 (trimethylsilyl amodimethicone) available from Dow Corning; DC 949 (amodimethicone, cetrimonium chloride, and Trideceth-12), available from Dow Corning; DC 749 (cyclomethicone and trimethylsiloxysilicate), available from Dow Corning; DC2502 (cetyl dimethicone), available from Dow Corning; BC97/004 and BC 99/088 (amino functionalized silicone microemulsions), available from Basildon Chemicals; GE SME253 and SM2115-D2_and SM2658 and SF1708 (amino functionalized silicone microemulsions), available from General Electric; siliconized meadowfoam seed oil, available from Croda; and those silicone conditioning agents described by GAF Corp. in U.S. Pat. No. 4,834,767 (quaternized amino lactam), by Biosil Technologies in U.S. Pat. No. 5,854,319 (reactive silicone emulsions containing amino acids), and by Dow

Corning in U.S. Pat. No. 4,898,585 (polysiloxanes), all of which descriptions are incorporated herein by reference.

Humectant

The compositions of the present invention may contain a humectant. The humectants herein are selected from the group consisting of polyhydric alcohols, water soluble alkoxyated nonionic polymers, and mixtures thereof. The humectants, when used herein, are preferably used at levels by weight of the composition of from about 0.1% to about 20%, more preferably from about 0.5% to about 5%.

Polyhydric alcohols useful herein include glycerin, sorbitol, propylene glycol, butylene glycol, hexylene glycol, ethoxylated glucose, 1, 2-hexane diol, hexanetriol, dipropylene glycol, erythritol, trehalose, diglycerin, xylitol, maltitol, maltose, glucose, fructose, sodium chondroitin sulfate, sodium hyaluronate, sodium adenosine phosphate, sodium lactate, pyrrolidone carbonate, glucosamine, cyclodextrin, and mixtures thereof.

Water soluble alkoxyated nonionic polymers useful herein include polyethylene glycols and polypropylene glycols having a molecular weight of up to about 1000 such as those with CTFA names PEG-200, PEG-400, PEG-600, PEG-1000, and mixtures thereof.

Commercially available humectants herein include: glycerin with tradenames STARTM and SUPEROLTM available from The Procter & Gamble Company, CRODEROL GA7000TM available from Croda Universal Ltd., PRECERINTM series available from Unichema, and a same tradename as the chemical name available from NOF; propylene glycol with tradename LEXOL PG-865/855TM available from Inolex, 1,2-PROPYLENE GLYCOL USP available from BASF; sorbitol with tradenames LIPONICTM series available from Lipo, SORBOTM, ALEXTM, A-625TM, and A-641TM available from ICI, and UNISWEET 70TM, UNISWEET CONCTTM available from UPI; dipropylene glycol with the same tradename available from BASF; diglycerin with tradename DIGLYCEROLTM available from Solvay GmbH; xylitol with the same tradename available from Kyowa and Eizai; maltitol with tradename MALBIT available from Hayashibara, sodium chondroitin sulfate with the same tradename available from Freeman and Bioiberica, and with tradename ATOMERGIC SODIUM CHONDROITIN SULFATE available from Atomergic Chemetals; sodium hyaluronate with tradenames ACTIMOIST available from Active Organics, AVIAN SODIUM HYALURONATE series available from Intergen, HYALURONIC ACID Na available from Ichimaru Pharcos; sodium adenosine phosphate with the same tradename available from Asahikasei, Kyowa, and Daiichi Seiyaku; sodium lactate with the same tradename available from Merck, Wako, and Showa Kako, cyclodextrin with tradenames CAVITRON available from American Maize, RHODOCAP series available from Rhone-

Poulenc, and DEXPEARL available from Tomen; and polyethylene glycols with the tradename CARBOWAX series available from Union Carbide.

Suspending Agent

The shampoo compositions of the present invention may further comprise a suspending agent at concentrations effective for suspending the particle, or other water-insoluble material, in dispersed form in the shampoo compositions or for modifying the viscosity of the composition. Such concentrations range from about 0.1% to about 10%, preferably from about 0.3% to about 5.0%, by weight of the shampoo compositions.

Suspending agents useful herein include anionic polymers and nonionic polymers. Useful herein are vinyl polymers such as cross linked acrylic acid polymers with the CTFA name Carbomer, cellulose derivatives and modified cellulose polymers such as methyl cellulose, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, nitro cellulose, sodium cellulose sulfate, sodium carboxymethyl cellulose, crystalline cellulose, cellulose powder, polyvinylpyrrolidone, polyvinyl alcohol, guar gum, hydroxypropyl guar gum, xanthan gum, arabia gum, tragacanth, galactan, carob gum, guar gum, karaya gum, carragheenin, pectin, agar, quince seed (*Cydonia oblonga* Mill), starch (rice, corn, potato, wheat), algae colloids (algae extract), microbiological polymers such as dextran, succinoglucan, pulleran, starch-based polymers such as carboxymethyl starch, methylhydroxypropyl starch, alginic acid-based polymers such as sodium alginate, alginic acid propylene glycol esters, acrylate polymers such as sodium polyacrylate, polyalkylene glycols having a molecular weight of more than about 1000, polyethylacrylate, polyacrylamide, polyethyleneimine, and inorganic water soluble material such as bentonite, aluminum magnesium silicate, laponite, hectonite, and anhydrous silicic acid.

Other Optional Components

The compositions of the present invention may contain also vitamins and amino acids such as: water soluble vitamins such as vitamin B1, B2, B6, B12, C, pantothenic acid, pantothenyl ethyl ether, panthenol, biotin, and their derivatives, water soluble amino acids such as asparagine, alanin, indole, glutamic acid and their salts, water insoluble vitamins such as vitamin A, D, E, and their derivatives, water insoluble amino acids such as tyrosine, tryptamine, and their salts.

The compositions of the present invention may also contain pigment materials such as inorganic, nitroso, monoazo, disazo, carotenoid, triphenyl methane, triaryl methane, xanthene, quinoline, oxazine, azine, anthraquinone, indigoid, thionindigoid, quinacridone, phthalocianine, botanical, natural colors, including: water soluble components such as those having C. I. Names: Acid Red 18, 26, 27, 33, 51, 52, 87, 88, 92, 94, 95, Acid Yellow 1, 3, 11, 23, 36, 40, 73,

Food Yellow 3, Food Green 3, Food blue 2, Food Red 1, 6, Acid Blue 5, 9, 74, Pigment Red 57-1, 53(Na), Basic Violet 10, Solvent Red 49, Acid orange 7, 20, 24, Acid Green 1, 3, 5, 25, Solvent Green 7, Acid Violet 9, 43; water insoluble components such as those having C. I. Names: Pigment Red 53(Ba), 49(Na), 49(Ca), 49(Ba), 49(Sr), 57, Solvent Red 23, 24, 43, 48, 72, 73, Solvent Orange 2, 7, Pigment Red 4, 24, 48, 63(Ca)3, 64, Vat Red 1, Vat blue 1, 6, Pigment Orange 1, 5, 13, Solvent Yellow 5, 6, 33, Pigment Yellow 1, 12, Solvent Green 3, Solvent Violet 13, Solvent Blue 63, Pigment Blue 15, titanium dioxides, chlorophyllin copper complex, ultramarines, aluminum powder, bentonite, calcium carbonate, barium sulfate, bismuthine, calcium sulfate, carbon black, bone black, chromic acid, cobalt blue, gold, ferric oxides, hydrated ferric oxide, ferric ferrocyanide, magnesium carbonate, manganous phosphate, silver, and zinc oxides.

The compositions of the present invention may also contain antimicrobial agents which are useful as cosmetic biocides and antidandruff agents including: water soluble components such as piroctone olamine, water insoluble components such as 3,4,4'- trichlorocarbanilide (trichlosan), triclocarban and zinc pyrithione.

The compositions of the present invention may also contain chelating agents such as: 2,2'-dipyridylamine; 1,10-phenanthroline {o-phenanthroline}; di-2-pyridyl ketone; 2,3-bis(2-pyridyl) pyrazine; 2,3-bis(2-pyridyl)-5,6-dihydropyrazine; 1,1'-carbonyldiimidazole; 2,4-bis(5,6-diphenyl-1,2,4-triazine-3-yl)pyridine; 2,4,6-tri(2-pyridyl)-1,3,5-triazine; 4,4'-dimethyl-2,2'-dipyridyl; 2,2'-biquinoline; di-2-pyridyl glyoxal {2,2'-pyridil}; 2-(2-pyridyl)benzimidazole; 2,2'-bipyrazine; 3-(2-pyridyl)5,6-diphenyl-1,2,4-triazine; 3-(4-phenyl-2-pyridyl)-5-phenyl-1,2,4-triazine; 3-(4-phenyl-2-pyridyl)-5,6-diphenyl-1,2,4-triazine; 2,3,5,6-tetrakis-(2'-pyridyl)-pyrazine; 2,6-pyridinedi-carboxylic acid; 2,4,5-trihydroxypyrimidine; phenyl 2-pyridyl ketoxime; 3-amino-5,6-dimethyl-1,2,4-triazine; 6-hydroxy-2-phenyl-3(2H)-pyridazinone; 2,4-pteridinediol {lumazine}; 2,2'-dipyridyl; and 2,3-dihydroxypyridine.

METHOD OF USE

The shampoo compositions of the present invention are used in a conventional manner for cleansing hair or skin and providing enhanced deposition of anti-dandruff particles and other benefits of the present invention. An effective amount of the composition for cleansing the hair or skin is applied to the hair or skin, that has preferably been wetted with water, and then rinsed off. Such effective amounts generally range from about 1g to about 50g, preferably from about 1g to about 20g. Application to the hair typically includes working the composition through the hair such that most or all of the hair is contacted with the composition.

This method for cleansing the hair and skin comprises the steps of:

a) wetting the hair and/or skin with water, b) applying an effective amount of the shampoo composition to the hair and/or skin, and c) rinsing the composition from the hair and/or skin using water. These steps can be repeated as many times as desired to achieve the desired cleansing and particle deposition benefits.

The following examples further describe and demonstrate the preferred embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration, and are not to be construed as limitations of the present invention since many variations thereof are possible without departing from its scope.

Examples

The shampoo compositions illustrated in the following Examples illustrate specific embodiments of the shampoo compositions of the present invention, but are not intended to be limiting thereof. Other modifications can be undertaken by the skilled artisan without departing from the spirit and scope of this invention. These exemplified embodiments of the shampoo composition of the present invention provide enhanced deposition efficiency benefits of the particles.

The shampoo compositions illustrated in the following Examples are prepared by conventional formulation and mixing methods, an example of which is set forth hereinbelow. All exemplified amounts are listed as weight percents and exclude minor materials such as diluents, preservatives, color solutions, imagery ingredients, botanicals, and so forth, unless otherwise specified.

The shampoo compositions of the present invention may be prepared using conventional formulation and mixing techniques. Where melting or dissolution of solid surfactants or wax components is required these can be added to a premix of the surfactants, or some portion of the surfactants, mixed and heated to melt the solid components, e.g., about 72° C. This mixture can then optionally be processed through a high shear mill and cooled, and then the remaining components are mixed in. The anti-dandruff particle component can be added either prior to processing through a high shear mill or preferably added as a predispersed suspension to this final mix , after cooling. The dimethicone may be prepared as a premix of desired particle size then added to the final mix. The compositions typically have a final viscosity of from about 2000 to about 20,000 cps. The viscosity of the composition can be adjusted by conventional techniques including addition of sodium chloride or ammonium xylenesulfonate as needed. The listed formulations, therefore, comprise the listed components and any minor materials associated with such components.

	Example 1	Example 2	Example 3	Example 4	Example 5	Example 6	Example 7
--	-----------	-----------	-----------	-----------	-----------	-----------	-----------

Ammonium Laureth Sulfate			12.00	12.00		12.00	12.00
Ammonium Lauryl Sulfate			2.00	2.00		2.00	2.00
Sodium Laureth Sulfate	12.00	12.00			12.00		
Sodium Lauryl Sulfate	2.00	2.00			2.00		
Cocamidopropyl Betaine	2.00	2.00		2.00	2.00	4.00	2.00
Sodium Lauroamphoacetate	2.00	2.00	4.00	2.00	2.00		2.00
Ethylene Glycol Distearate	1.50	1.50	1.50	1.50	1.50	1.50	1.50
CMEA	0.800	0.800	0.800	0.800	0.800	0.800	0.800
Cetyl Alcohol	0.900	0.900	0.600	0.600	0.900	0.600	0.600
Lauryl Alcohol	0.200	0.200			0.200		
Guar Hydroxypropyl Trimonium Chloride (1)					0.250		
Polyquaterium-10 (2)	0.500	0.500	0.500	0.500			0.500
Polyquaterium-10 (3)						0.500	
Dimethicone (4)		1.35				1.35	
Dimethicone (5)	1.00		1.00	1.00	1.00		1.00
Trimethylolpropane tricaprylate/tricaprate (6)							0.10
Hydrogenated Polydecene (7)			0.25			0.25	0.40
ZPT (8)	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Sodium Citrate			0.40	0.40		0.40	0.40
Citric Acid			0.39	0.39		0.39	0.39
Hydrochloric Acid	0.600	0.600	0.600	0.600	0.600	0.600	0.600
Sodium Xylenesulfonate	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Perfume	0.400	0.400	0.400	0.400	0.400	0.400	0.400
Sodium Benzoate	0.250	0.250	0.250	0.250	0.250	0.250	0.250
Kathon	0.0008	0.0008	0.0008	0.0008	0.0008	0.0008	0.0008
Benzyl Alcohol	0.0225	0.0225	0.0225	0.0225	0.0225	0.0225	0.0225
Water	q.s.	q.s.	q.s.	q.s.	q.s.	q.s.	q.s.

- (1) Jaguar C17 available from Rhodia
- (2) Polymer KG30M available from Amerchol/Dow Chemical
- (3) Polymer JR30M available from Amerchol/Dow Chemical
- (4) Viscasil 330M available from General Electric Silicones
- (5) DC1664 available from Dow Corning Silicones
- (6) Mobil P43, available from Mobil.
- (7) Puresyn 6, available from Mobil.
- (8) ZPT having an average particle size of about 2.5 μ m, available from Arch/Olin.


It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to one skilled in the art without departing from the scope of the present invention.

WHAT IS CLAIMED IS:

1. A shampoo composition comprising:

- a) from about 5 to about 50 weight percent of a deterative surfactant,
- b) at least about 0.1 weight percent of anti-dandruff particles,
- c) at least about 0.05 weight percent of a cationic polysaccharide polymer having a molecular weight of from about 10,000 to about 10,000,000 and a charge density from about 1.4 meq/gm to about 7.0 meq/gm, and
- d) at least about 20.0 weight percent of an aqueous carrier.

2. A shampoo composition according to Claim 1, wherein said cationic polysaccharide polymer conforms to the general formula:



- a) wherein A is an anhydroglucose residual group;
- b) wherein R is selected from the group consisting of alkylene oxyalkylene, polyoxyalkylene, hydroxyalkylene, and mixtures thereof;
- c) wherein R¹, R², and R³ are independently selected from the group consisting of alkyl, aryl, alkylaryl, arylalkyl, alkoxyalkyl, and alkoxyaryl; each group containing up to about 18 carbon atoms, and the sum of carbon atoms in R¹, R², and R³ is less than about 20; and
- d) wherein X is selected from the group consisting of chloride, fluoride, bromide, iodide, sulfate, methylsulfate, and mixtures thereof.

3. The composition of claim 1 wherein said cationic polysaccharide polymer has a charge density of from about 1.7 meq/gm to about 7 meq/gm.

4. The composition of claim 1 wherein said cationic polysaccharide polymer has a charge density of from about 1.9 meq/gm to about 5 meq/gm.

5. The composition of claim 1 wherein said anti-dandruff particles comprise a zinc salt of 1-hydroxy-2-pyridinethione.

6. The composition of claim 4 wherein said anti-dandruff particles comprise a zinc salt of 1-hydroxy-2-pyridinethione.
7. The composition of claim 1 wherein the weight ratio of said cationic polysaccharide polymer to said anti-dandruff particles is from about 1:1 to about 1:20.
8. A method of treating hair by administering a safe and effective amount of the composition according to claim 1.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 03/12265

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K7/075

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

PAJ, EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 22074 A (PROCTER & GAMBLE) 25 July 1996 (1996-07-25) examples IV,V,IX-XX page 3, line 16-20 page 15, line 26 -page 19, line 10 page 21, line 6 -page 22, line 5 ---	1-8
X	US 3 580 853 A (PARRAN JOHN J JR) 25 May 1971 (1971-05-25) column 1, line 58 -column 2, line 41 column 5, line 73 -column 6, line 55 column 7, line 50-67 column 8, line 49-75 --- -/--	1-8

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

13 August 2003

Date of mailing of the international search report

21/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hauss, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/12265

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 39683 A (OKUYAMA YOSHINARI ;HU FANG (JP); PROCTER & GAMBLE (US)) 12 August 1999 (1999-08-12) examples 1-3,7,9 claims 1-3,7-9 page 18, line 3-5 page 13, line 3,4 ---	1,3-8
X	WO 97 26854 A (PROCTER & GAMBLE) 31 July 1997 (1997-07-31) examples I-VI,VIII,X,XII,XIII,XVI claims page 14, paragraph 2 -page 15, paragraph 1 ---	1,3-8
X	WO 00 66081 A (PROCTER & GAMBLE) 9 November 2000 (2000-11-09) examples VII-XIV,XIX page 22, line 14 -page 23, line 8 claim 1 ---	1,3-8
P,X	WO 03 028683 A (PROCTER & GAMBLE) 10 April 2003 (2003-04-10) page 1, paragraph 1; claim 1 page 14, line 27 -page 15, line 15 page 17, paragraph 3 -page 20, paragraph 5 page 35, line 16 -page 36, paragraph 3 examples -----	1-8

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intel International Application No

PCT/US 03/12265

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9622074	A	25-07-1996	US 5624666 A	29-04-1997
			AU 713104 B2	25-11-1999
			AU 4696396 A	07-08-1996
			BR 9607157 A	11-11-1997
			CA 2210089 A1	25-07-1996
			CN 1169110 A	31-12-1997
			EP 0804141 A1	05-11-1997
			JP 10512568 T	02-12-1998
			WO 9622074 A1	25-07-1996
US 3580853	A	25-05-1971	AT 310905 B	25-10-1973
			BE 721384 A	25-03-1969
			CH 531041 A	30-11-1972
			DE 1792618 A1	25-11-1971
			FR 1588952 A	16-03-1970
			GB 1195158 A	17-06-1970
			IT 1046452 B	30-06-1980
			NL 6813826 A	31-03-1969
			SE 353738 B	12-02-1973
			US 3723325 A	27-03-1973
			US 3753916 A	21-08-1973
			US 3761417 A	25-09-1973
			US 3761418 A	25-09-1973
WO 9939683	A	12-08-1999	WO 9939683 A1	12-08-1999
WO 9726854	A	31-07-1997	AU 1691097 A	20-08-1997
			BR 9707216 A	20-07-1999
			CA 2242932 A1	31-07-1997
			CN 1213294 A	07-04-1999
			CZ 9802296 A3	16-12-1998
			EP 0920298 A1	09-06-1999
			HU 9900641 A2	28-07-1999
			JP 11503175 T	23-03-1999
			TR 9801453 T2	21-10-1998
			WO 9726854 A1	31-07-1997
			ZA 9700696 A	04-08-1997
WO 0066081	A	09-11-2000	AU 759514 B2	17-04-2003
			AU 4689100 A	17-11-2000
			CA 2370939 A1	09-11-2000
			CN 1353601 T	12-06-2002
			EP 1175202 A1	30-01-2002
			JP 2002543105 T	17-12-2002
			WO 0066081 A1	09-11-2000
			US 6451300 B1	17-09-2002
WO 03028683	A	10-04-2003	WO 03028683 A1	10-04-2003
			US 2003108501 A1	12-06-2003