

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 September 2003 (12.09.2003)

PCT

(10) International Publication Number
WO 03/074055 A1

(51) International Patent Classification⁷: **A61K 31/4425**,
31/4523, 31/4375, 45/06, A61P 11/06, 11/08

(21) International Application Number: PCT/EP03/01876

(22) International Filing Date: 25 February 2003 (25.02.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02004987.0 6 March 2002 (06.03.2002) EP

(71) Applicant (for all designated States except US): **ALTANA PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BEUME, Rolf** [DE/DE]; Bohlstrasse 13, 78465 Konstanz (DE). **BUND-SCHUH, Daniela** [DE/DE]; Rheingutstrasse 17, 78462 Konstanz (DE). **WEIMAR, Christian** [DE/DE]; Eichhornstr. 51, 78464 Konstanz (DE). **WOLLIN, Stefan-Lutz** [DE/DE]; Lehrenweg 15/4, 88709 Meersburg (DE).

(74) Common Representative: **ALTANA PHARMA AG**; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(81) Designated States (*national*): AE, AL, AU, BA, BR, CA, CN, CO, CU, DZ, EC, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, US, VN, YU, ZA, ZW.

(84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AL, AU, BA, BR, CA, CN, CO, CU, DZ, EC, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHARMACEUTICAL COMPOSITION OF A PDE4 OR A PDE3/4 INHIBITOR AND A HISTAMINE RECEPTOR ANTAGONIST

(57) Abstract: The invention relates to the combined administration of PDE4 or PDE3/4 inhibitors and histamine receptor antagonists for the treatment of respiratory diseases.



WO 03/074055 A1

PHARMACEUTICAL COMPOSITION OF A PDE4 OR A PDE3/4 INHIBITOR AND A HISTAMINE RECEPTOR ANTAGONIST

Field of application of the invention

The present invention relates to combinations of pharmaceutically active substances for use in the treatment of respiratory diseases.

The substances used in the combinations according to the invention are known active compounds from the PDE4 and PDE3/4 inhibitors class and active compounds from the class of the histamine receptor antagonists.

Known technical background

In the international application WO03/000289 compositions are described which comprise a PDE4 inhibitor and a H1-receptor antagonist and the use of these compositions for the manufacture of a medicament for the treatment of respiratory diseases.

Description of the invention

Asthma is a common inflammatory disease of the respiratory tract, accounting for 1 - 3% of all office visits, 500,000 hospital admissions per year and more pediatric hospital admissions than any other single illness in the US. Annually, more than 5000 children and adults die of asthma attacks in the United States (William E. S.; Goodman Gilman A.: The pharmacological Basis of Therapeutics, 9th Edition, pp. 152 & 659-682, Mc Graw Hill, New York 1996).

Asthma can no longer be viewed simply as a reversible airway obstruction. It should instead be considered primarily as an inflammatory illness that has bronchial hyperactivity and bronchospasm as its results. Allergen specific immunoglobulin E (IgE) is bound to the mast cells via Fc receptors. It is a fragment obtained by papain digestion of immunoglobulin molecules and contains most of the antigenic determinants. When an allergen comes into contact with IgE, the mast cells are activated and release a number of inflammatory mediators, which include granule contents like histamine, proteases, heparin, and tumor necrosis factor (TNF), a variety of lipid membrane derived molecules like prostaglandins, leukotrienes and platelet activating factor (PAF), and a number of cytokines like interleukin (IL)-1, 3, 4, 5, 6 and 8 and chemokines. An enormous variety of mediators are released which have more than one potent effect on airway inflammation.

- 2 -

As a result of vasodilation, increased vasopermeability and increased endothelial adhesiveness towards leukocytes further leads to an influx of inflammatory cells like lymphocytes, eosinophils and macrophages from blood circulation into the tissues. This in turn leads to the release of mediators which have further inflammatory effects (Rao A. R. et al. Recent Perspectives in the design of antiasthmatic agents, Pharmazie, 55, 7, 475-482, 2000).

Thus, it can be understood that it is unlikely that drugs affecting a single mediator can satisfactorily treat the disease alone. As asthma is one of the major diseases affecting mankind, there is a need to develop drugs which can affect a wide variety of mediators.

Therefore, it is the object of the present invention to make available respiratory tract therapeutics which fulfill the following conditions:

- Favorable simultaneous influence on several of the inflammatory mediators
- Marked bronchorelaxation and -dilatation
- Good oral availability
- Minor side effects
- Good suitability for long-term therapy
- Favorable influence on bronchial hyperreactivity

It has now been found that the combined use of a PDE4 or a PDE3/4 inhibitor and a histamine receptor antagonist outstandingly fulfills the above-mentioned conditions.

The invention thus relates to the combined use of a PDE4 or a PDE3/4 inhibitor and a histamine receptor antagonist in the treatment of respiratory diseases.

By the expression "PDE4 inhibitor" is meant a selective phosphodiesterase inhibitor, which inhibits preferentially the type 4 phosphodiesterase when compared to other known types of phosphodiesterase, e.g. type 1, 2, 3, 5 etc., whereby the compound has a lower IC_{50} (more potent) for the type 4 phosphodiesterase, such as where the IC_{50} for PDE4 inhibition is about factor 10 lower compared to IC_{50} for inhibition of other known type of phosphodiesterase, e.g. type 1, 2, 3, 5 etc.

























Analogously, the expression "PDE3/4 inhibitor" is defined. Methods to determine the activity and selectivity of a phosphodiesterase inhibitor are known to the person skilled in the art. In this connection it may be mentioned, for example, the methods described by Thompson et al. (Adv Cycl Nucl Res 10: 69-92, 1979), Giembycz et al. (Br J Pharmacol 118: 1945-1958, 1996) and the phosphodiesterase scintillation proximity assay of Amersham Pharmacia Biotech.


By the expression "histamine receptor antagonist" are meant H₁-receptor antagonists, particularly the so called H₁-receptor antagonists of the second generation (Mutschler, Arzneimittelwirkungen, 8. Edition, 2001, pages 456-461).

As possible PDE4 or PDE3/4 inhibitors within the meaning of the present invention may be mentioned, by way of example, those PDE4 or PDE3/4 inhibitors which are named expressis verbis as an example, or described or claimed generically in the following patent applications and patents:


DE 1545687, DE 2028869, DE 2123328, DE 2315801, DE 2402908, DE 2413935, DE 3900233, EP 0103497, EP 0139464, EP 0158380, EP 0163965, EP 0335386, EP 0389282, EP 0393500, EP 0428302, EP 0435811, EP 0449216, EP 0459505, EP 0470805, EP 0490823, EP 0506194, EP 0510562, EP 0511865, EP 0527117, EP 0553174, EP 0557016, EP 0626939, EP 0664289, EP 0671389, EP 0685474, EP 0685475, EP 0685479, EP 0731099, EP 0736532, EP 0738715, EP 0748805, EP 0763534, EP 0816357, EP 0819688, EP 0819689, EP 0832886, EP 0834508, EP 0848000, JP 92234389, JP 94329652, JP 95010875, JP 98072415, JP 98147585, US 5703098, US 5739144, WO 9117991, WO 9200968, WO 9212961, WO 9307146, WO 9315044, WO 9315045, WO 9318024, WO 9319068, WO 9319720, WO 9319747, WO 9319749, WO 9319751, WO 9325517, WO 9402465, WO 9412461, WO 9420455, WO 9422852, WO 9427947, WO 9500516, WO 9501338, WO 9501980, WO 9503794, WO 9504045, WO 9504046, WO 9505386, WO 9508534, WO 9509623, WO 9509624, WO 9509627, WO 9509836, WO 9514667, WO 9514680, WO 9514681, WO 9517392, WO 9517399, WO 9519362, WO 9520578, WO 9522520, WO 9524381, WO 9527692, WO 9535281, WO 9535283, WO 9535284, WO 9600218, WO 9601825, WO 9606843, WO 9603399, WO 9611690, WO 9611917, WO 9612720, WO 9631486, WO 9631487, WO 9635683, WO 9636595, WO 9636596, WO 9636611, WO 9636625, WO 9636626, WO 9636638, WO 9638150, WO 9639408, WO 9640636, WO 9703967, WO 9704779, WO 9705105, WO 9708143, WO 9709345, WO 9712895, WO 9718208, WO 9719078, WO 9720833, WO 9722585, WO 9722586, WO 9723457, WO 9723460, WO 9723461, WO 9724117, WO 9724355, WO 9725312, WO 9728131, WO 9730999, WO 9731000, WO 9732853, WO 9735854, WO 9736905, WO 9740032, WO 9743288, WO 9744036, WO 9744322, WO 9747604, WO 9748697, WO 9804534, WO 9805327, WO 9806692, WO 9806704, WO 9807715, WO 9808828, WO 9808830, WO 9808841, WO 9808844, WO 9809946, WO 9809961, WO 9811113, WO 9814448, WO 9818796, WO 9821207, WO 9821208, WO 9821209, WO 9822453, WO 9831674, WO 9840382, WO 9845268, WO 9855481, WO 9856756, WO 9905111, WO 9905112, WO 9505113, WO 9906404, WO 9918095, WO 9931071, WO 9931090, WO 9947505, WO 9957115, WO 9957118, WO 9964414, WO 0001695, WO 0012501, WO 0026208, WO 0042017, WO 0042018, WO 0042019, WO 0042020, WO 0042034, WO 0119818, WO 0130766, WO 0130777, WO 0151470, WO 02006270, WO 02066476, WO 02064584 and WO 02085885.


Exemplary PDE inhibitors are shown on the following pages with the aid of their formulae:








- 6 -

























































(Pumafentrin)



Roflumilast





In the above formulae there is given neither any stereochemical information nor are hydrogen atoms indicated [-O is accordingly -OH, >N is >NH or -N is NH₂]. Methyl groups, e.g. on the oxygen atoms, are indicated by lines].

Those PDE4 or PDE3/4 inhibitors are to be emphasized which are named expressis verbis as an example and/or claimed generically in the patent applications or patents EP 0163965, EP 0389282, EP 0393500, EP 0435811, EP 0482302, EP 0499216, EP 0506194, EP 0510562, EP 0528922, EP 0553174, EP 0731099, WO 9319749, WO 9500516, WO 9501338, WO 9600218, WO 9603399, WO 9611690, WO 9636625, WO 9636626, WO 9723457, WO 9728131, WO 9735854, WO 9740032, WO 9743288, WO 9809946, WO 9807715, WO 9808841, WO 9821207, WO 9821208, WO 9821209,

WO 9822453, WO 9831674, WO 9840382, WO 9855481, WO 9905111, WO 9905112, WO 9905113, WO 9931071, WO 9931090, WO 9947505, WO 9957115, WO 9957118, WO 9964414, WO 0001695, WO 0012501, WO 0042017, WO 0042018, WO 0042019, WO 0042020, WO 0042034, WO 0119818, WO 0130766, WO 0130777, WO 0151470 WO 02006270, WO 02066476, WO 02064584 and WO 02085885 and the compounds with the following research codes: CDC-998, D-4396, SCH-351591, IC-485, CC-1088 and KW-4490. Substances having good oral availability are preferred here.

Preferred PDE4 or PDE3/4 inhibitors are 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]-benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)-thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β-[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenyl-carbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST] and the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and KW-4490.

Further preferred PDE4 inhibitors are selected from the group consisting of
 (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
5-{4-[(4aS,8aR)-4-(3,4-Diethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-[1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl]-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester and
 2-{4-[(4aS, 8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide.

Particularly preferred PDE4 or PDE3/4 inhibitors are 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST] and (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE].

As possible histamine receptor antagonists within the meaning of the present invention may be mentioned, by way of example, (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)-phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-

yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-{2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl}-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLIN], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE].

Preferred histamine receptor antagonists are 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].

Particularly preferred histamine receptor antagonists are 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE] and ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE].

In the context of the present invention, unless otherwise stated, a pharmaceutically acceptable derivative of an active ingredient means a pharmaceutically acceptable salt or solvate (e. g. hydrate), a pharmaceutically acceptable solvate of such salt, a pharmaceutically acceptable N-oxide or a pharmaceutically acceptable salt or solvate of the latter.

Suitable pharmacologically tolerable salts here are on the one hand in particular water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)-benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 1-hydroxy-2-naphthoic acid, the acids being employed in salt preparation – depending on whether it is a mono- or polybasic acid and depending on which salt is desired – in an equimolar quantitative ratio or one differing therefrom. Furthermore, the active compounds mentioned can also be present as pure enantiomers or as enantiomer mixtures in any mixing ratio.

On the other hand, salts with bases are also suitable. Examples of salts with bases which may be mentioned are alkali metal (lithium, sodium, potassium) or calcium, aluminium, magnesium, titanium, ammonium, meglumine or guanidinium salts, where here too the bases are employed in salt preparation in an equimolar quantitative ratio or one differing therefrom.

Certain of the active ingredients used in the present invention are capable of existing in stereoisomeric forms. The invention encompasses all stereoisomers of the active ingredients and mixtures thereof including racemates. Tautomers and mixtures thereof of the active ingredients are also part of the present invention.

In accordance with the present invention, there is provided in a first aspect a pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from a histamine receptor antagonist and its pharmaceutically acceptable derivatives.

In a second aspect - which is an embodiment of the first aspect - there is provided a pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from 3-Cyclopropyl-methoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxy-benzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code:

ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN: AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN: CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research-Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088, KW-4490 and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]-pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE],

1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmaceutically acceptable derivatives.

In a third aspect - which is another embodiment of the first aspect - there is provided a pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from 3-Cyclopropyl-methoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β-[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBU-FELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATA-DINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEX-CHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBA-STINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXO-FENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]-propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-

piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLIN], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and their pharmaceutically acceptable derivatives.

In a forth aspect - which is still another embodiment of the first aspect - there is provided a Pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo[c][1,6]naphthyridine [INN: PUMAFENTRINE] and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmaceutically acceptable derivatives.

In the fifth aspect - which is a further embodiment of the first aspect - there is provided a pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-[1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl]-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl)-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl)-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRI-VASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chloro-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBA-

STINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLIN], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmaceutically acceptable derivatives.

In a sixth aspect - which is still a further embodiment of the first aspect - there is provided a pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-11⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-11⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-11⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-11⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-11⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-11⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)-piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmaceutically acceptable derivatives.

In a seventh aspect the invention provides a pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected from a histamine receptor antagonist and its pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.

In a eighth aspect - which is an embodiment of the seventh aspect - the invention provides a Pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyl-oxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyl-oxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β-[3-(Cyclopentyl-oxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFE-LONE], cis-[4-Cyano-4-(3-cyclopentyl-oxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088, KW-4490 and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]-ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-

benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.

In a ninth aspect - which is another embodiment of the seventh aspect - the invention provides a pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentylloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN: AROFYLLINE], 3-[3-(Cyclopentylloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN: CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentylloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research-Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentylloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)-phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-ben-

zo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-α-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-α-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-α-methyl-α-phenylbenzyl)-oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(α-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and their pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.

In a tenth aspect - which is still another embodiment of the seventh aspect - the invention provides a pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE] and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-α-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-α-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(α-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.

In the eleventh aspect - which is a further embodiment of the seventh aspect - the invention provides a pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

5-[4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl]-5-oxo-pentanoic acid,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,

(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester, 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)-phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)-oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPI-NASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]-ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPLENAMINE], and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.

In the twelfth aspect, which is a still further embodiment of the seventh aspect - the invention provides a pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-

benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-1H-benzo[5,6]cyclohepta[1,2-b]pyridin-1-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidiny]-N-methyl-amino-pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.

In a thirteenth aspect, the invention provides a kit comprising a preparation of a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from a histamine receptor antagonist and its pharmaceutically acceptable derivatives, and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

In a fourteenth aspect - which is an embodiment of the thirteenth aspect - the invention provides a kit comprising a preparation of a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarboxamido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN: CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research-Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088, KW-4490 and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN:

CETIRIZINE], (+)-2-{2-[(p-Chloro-alpha-methyl-alpha phenylbenzyl)oxy]ethyl}-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEX-CHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBA-STINE], [2-[4-bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXO-FENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]pro-pionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluoro-benzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTI-NE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-aze-pine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SU-DEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmaceutically acceptable derivatives, and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

In a fifteenth aspect - which is another embodiment of the thirteenth aspect - the invention provides a kit comprising a preparation of a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethyl-amino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetra-hydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxy-benzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cy-clopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyl-

oxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyl-oxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)-methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-di-benz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and their pharmaceutically acceptable derivatives, and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

In a sixteenth aspect - which is still another embodiment of the thirteenth aspect - the invention provides a kit comprising a preparation of a first active ingredient which is selected from 3-Cyclopropyl-methoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-

ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE] and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFE-NADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmaceutically acceptable derivatives, and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

In the seventeenth aspect – which is a further embodiment of the thirteenth aspect - the invention provides a kit comprising a preparation of a first active ingredient which is selected from (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

5-[4-[(4aS,8aR)-4-(3,4-Diethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl]-5-oxo-pentanoic acid,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,

(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-[1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl]-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)-phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazin-yl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]-ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenzo[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLIN], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]-pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-

1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmaceutically acceptable derivatives, and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

In the eighteenth aspect – which is a still further embodiment of the thirteenth aspect - the invention provides a kit comprising a preparation of a first active ingredient which is selected from

(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-[2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl]-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmaceutically acceptable derivatives, and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

It has been found that the administration of active ingredients according to the invention is advantageous because it results - in comparison to the administration of a single active ingredient from the PDE4 and PDE3/4 inhibitors or the histamine receptor antagonists class - in a reduced early allergic response and/or in a reduced late inflammatory airway response.

The pharmaceutical composition of the present invention may be prepared by mixing the first active ingredient with the second active ingredient. Therefore, in the nineteenth aspect of the present invention, there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmacologically acceptable derivatives, with a second active ingredient which is selected from a histamine receptor antagonist and its pharmacologically acceptable derivatives.

In a twentieth aspect - which is an embodiment of the nineteenth aspect - there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591],

3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β-[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research-Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088, KW-4490 and their pharmacologically acceptable derivatives, with a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-α-phenylbenzyl)-1-piperazinyl]-ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-α-methyl-α-phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]-cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]butyl]-α-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-α-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]-pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-α-methyl-α-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN:

SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmacologically acceptable derivatives.

In a twentyfirst aspect - which is another embodiment of the nineteenth aspect - there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzodiazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN: AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN: CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)-cyclohexane-1-carboxylic acid [INN: CILOMILAST], the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and their pharmacologically acceptable derivatives, with a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)-methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxe-

pino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and their pharmacologically acceptable derivatives.

In a twentysecond aspect - which is still another embodiment of the nineteenth aspect - there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active ingredient which is selected from 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyridin-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE] and their pharmacologically acceptable derivatives, with a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmacologically acceptable derivatives.

In a twentythird aspect - which is still another embodiment of the nineteenth aspect - there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active ingredient which is selected from
 (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-11⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-11⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,

(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-[1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl]-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmacologically acceptable derivatives, with a second active ingredient which is selected from (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRI-VASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-

b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFE-NADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXA-NOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE] and their pharmacologically acceptable derivatives.

In a twentyfourth aspect - which is still another embodiment of the nineteenth aspect - there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active ingredient which is selected from (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxy-phenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and their pharmacologically acceptable derivatives, with a second active ingredient which is selected from 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)-piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and their pharmacologically acceptable derivatives.

The first and second active ingredient may alternatively (other than in admixture as described above) be administered simultaneously, sequentially or separately to treat respiratory diseases. By sequential is meant that the first and second active ingredient are administered one immediately after the other. Separately means a time difference of administration of up to 24 h, preferably up to 12 h.

The present invention further provides the use of a pharmaceutical composition, or pharmaceutical product according to the invention in the manufacture of a medicament for the prophylaxis and/or treatment of a respiratory disease.

A further aspect of the present invention is a method for the treatment of a respiratory disease comprising administering to a patient in need thereof (a) an effective amount of a PDE4 inhibitor, a PDE3/4 inhibitor or a pharmaceutically acceptable derivative thereof and (b) an effective amount of a histamine receptor antagonist or a pharmaceutically acceptable derivative thereof.

Respiratory diseases which may be mentioned are in particular allergen- and inflammation- induced bronchial disorders (bronchitis, obstructive bronchitis, spastic bronchitis, allergic bronchitis, allergic asthma, bronchial asthma, COPD, allergic, seasonal and perennial rhinitis), which can be treated by the combination according to the invention also in the sense of a long-term therapy (if desired with appropriate adjustment of the dosage of the individual components to the needs at the time, for example needs subject to seasonally related variations).

The active ingredients may, and indeed will, as part of the pharmaceutical composition, the Pharmaceutical product or preparation, be used in admixture with one or more pharmaceutically acceptable auxiliaries and/or excipients.

Within the meaning of the present invention, "use" is preferably understood as meaning the oral administration of both active ingredients. The active ingredients also can be administered as a nasal spray, an aerosol, or as an inhaled powder. Further methods of administration, which may be mentioned are the parenteral (e. g. intravenous) and the transdermal administration of the active ingredients.

The invention encompasses on the one hand co-administering both drugs in one delivery form such as a fixed oral combination (putting both active ingredients in one tablet), as an inhaler (putting both active ingredients in the same inhaler) or as a free oral combination (putting both active ingredients in two separate tablets). On the other hand it encompasses also the administration of the drugs in two different delivery forms such as putting the PDE4 inhibitor into tablets and package them with an inhaler that contains the histamine receptor antagonist, or vice versa.

The person skilled in the art is familiar on the basis of his/her expert knowledge with which excipients or auxiliaries are suitable for the desired pharmaceutical composition, product or preparation. In addition to solvents, gel-forming agents, tablet excipients and other active compound carriers, it is possible to use, for example, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents,

preservatives, solubilizers, colorants or permeation promoters and complexing agents (e.g. cyclodextrins).

The pharmaceutical compositions or preparations according to the invention for oral administration are preferably in the form of tablets, coated tablets, capsules, emulsions, suspensions or solutions, the active ingredient content advantageously being between 0.1 and 95% and by appropriate choice of the excipients and the auxiliaries, it being possible to achieve a pharmaceutical administration form precisely tailored to the active ingredient(s) and/or to the desired onset of action (e.g. a sustained release form or an enteric form).

The active ingredients according to the invention are administered by inhalation preferably in the form of an aerosol; the aerosol particles of solid, liquid or mixed composition preferably having a diameter of 0.5 to 10 μm , advantageously of 2 to 6 μm .

Aerosol generation can be carried out, for example, by pressure-driven jet atomizers or ultrasonic atomizers, but advantageously by propellant-driven metered aerosols or propellant-free administration of micronized active compounds from inhalation capsules.

Depending on the inhaler system used, in addition to the active compounds the administration forms additionally contain the required excipients, such as, for example, propellants (e.g. Frigen in the case of metered aerosols), surface-active substances, emulsifiers, stabilizers, preservatives, flavorings, fillers (e.g. lactose in the case of powder inhalers) or, if appropriate, further active compounds.

For the purposes of inhalation, a large number of apparatuses are available with which aerosols of optimum particle size can be generated and administered, using an inhalation technique which is as right as possible for the patient. In addition to the use of adaptors (spacers, expanders) and pear-shaped containers (e.g. Nebulator®, Volumatic®), and automatic devices emitting a puffer spray (Autohaler®), for metered aerosols, in particular in the case of powder inhalers, a number of technical solutions are available (e.g. Diskhaler®, Rotadisk®, Turbohaler® or the inhaler described in European Patent Application EP 0 505 321), using which an optimal administration of active compound can be achieved.

Typical compositions for nasal delivery include those mentioned above for inhalation and further include non-pressurized compositions in the form of a solution or suspension in an inert vehicle such as water optionally in combination with conventional excipients such as buffers, anti-microbials, tonicity modifying agents and viscosity modifying agents which may be administered by nasal pump.

Typical transdermal formulations comprise a conventional aqueous or non aqueous vehicle, for example a cream, ointment, lotion or paste, or are in the form of a medicated plaster, patch or membrane.

For the above-mentioned therapeutic uses the dosages administered will, of course, vary with the first and second active ingredients employed, the mode of administration, the treatment desired and the disorder indicated.

However, in general, satisfactory results will be obtained when the total daily dosage of first active ingredient(s), the PDE4 respectively the PDE3/4 inhibitors, when taken oral is in the range from 1 - 2000 µg/kg of body weight. In the case of the particularly preferred PDE4 inhibitor ROFLUMILAST, the daily dosage is in a range from 1 - 20 µg/kg of body weight. The daily dosage for the particularly preferred PDE3/4 inhibitor PUMAFENTRINE is in a range from 300 - 1500 µg/kg of body weight.

The total daily dosage of the second active ingredient(s), the histamine receptor antagonists also can vary within a wide range (0.1 - 1500 µg/kg of body weight). In the case of the particularly preferred histamine receptor antagonist LORATADINE, the daily dosage when taken oral is in a range from 0.1 - 0.5 µg/kg of body weight.

Patent claims

1. A pharmaceutical composition comprising, in admixture, a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmaceutically acceptable derivatives, and a second active ingredient which is selected from a histamine receptor antagonist and its pharmaceutically acceptable derivatives.
2. A pharmaceutical composition according to claim 1, wherein the first and/or second active ingredient is in the form of a pharmaceutically acceptable salt, solvate, N-oxide or solvate of an salt or N-oxide.
3. A pharmaceutical composition according to claim 1 or 2, which is a fixed oral combination.
4. A pharmaceutical composition according to claim 1 or 2, which is a dry powder for use in a dry powder inhaler.
5. A pharmaceutical composition according to claim 1 or 2, which is an aqueous preparation for nasal administration.
6. A pharmaceutical composition according to claim 1 or 2, in which the PDE4 inhibitor, the PDE3/4 inhibitor or their pharmaceutically acceptable derivatives and the histamine receptor antagonist or its pharmaceutically acceptable derivative is combined with a propellant to form a composition which is delivered using a metered dose inhaler.
7. A pharmaceutical composition according to any of the claims 1 to 6, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarboxamido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide

[Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN: CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST] and the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and KW-4490, and the histamine receptor antagonist is selected from the group consisting of (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)-cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino-pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE].

8. A pharmaceutical composition according to any of the claims 1 to 6, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarboxamido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN: CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research -Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST] and the compounds with the research codes CDC-998, SH-636, D-4396, IC-485 and CC-1088, and the histamine receptor antagonist is selected from the group consisting of (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)-methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-

phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthene-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE].

9. A pharmaceutical composition according to any of the claims 1 to 6, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST] and (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], and the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].
10. A pharmaceutical composition according to claim 9, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof.
11. A pharmaceutical composition according to claim 9, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate, or N-oxide thereof, or solvate of an salt or N-oxide thereof.

12. A pharmaceutical composition according to any of the claims 1 to 6, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of
- (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1H⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1H⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1H⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1H⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- 5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
- (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- 4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
- 4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
- (cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and the histamine receptor antagonist is selected from the group consisting of (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)-piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidiny]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE].

13. A pharmaceutical composition according to any of the claims 1 to 6, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of
 (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
5-[4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl]-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanolamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-

[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].

14. A pharmaceutical composition according to claim 9 or 13, wherein the histamine receptor antagonist is selected from the group consisting of 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], a pharmaceutically acceptable salt or solvate thereof, or a solvate of an salt thereof.
15. Use of a pharmaceutical composition according to claim 1 in the manufacture of a medicament for the treatment of a respiratory diseases.
16. A process for the preparation of a pharmaceutical composition as defined in claim 1 which comprises mixing the first active ingredient with the second active ingredient.
17. A pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmaceutically acceptable derivatives, and a preparation of a second active ingredient which is selected from a histamine receptor antagonist and its pharmaceutically acceptable derivatives, for simultaneous, sequential or separate use in therapy.
18. A pharmaceutical product according to claim 17, wherein the first and/or second active ingredient is in the form of a pharmaceutically acceptable salt, solvate, N-oxide or solvate of an salt or N-oxide.
19. A pharmaceutical product according to claim 17 or 18, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241],

3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β -[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research - Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST] and the compounds with the research codes CDC-998, SH-636, D-4396, IC-485, CC-1088 and KW-4490, and the histamine receptor antagonist is selected from the group consisting of (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]-piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLINE], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxo-

xanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE] and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE].

20. A pharmaceutical product according to claim 17 or 18, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], 3,5-dichloro-4-[8-methoxy-2-(trifluoromethyl)quinolin-5-ylcarbox-amido]pyridine-1-oxide [Research Code: SCH-351591], 3-[3-(cyclopentyloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine [Research-Code: V-11294A], N-[9-methyl-4-oxo-1-phenyl-3,4,6,7-tetrahydropyrrolo[3,2,1-jk][1,4]benzo-diazepin-3(R)-yl]pyridine-4-carboxamide [Research-Code: CI-1018], 4-(3,4-dimethoxyphenyl)thiazole-2-carboxamide oxime [Research Code: ORG-20241], 3,7-dihydro-3-(4-chlorophenyl)-1-propyl-1H-purine-2,6-dione [INN AROFYLLINE], 3-[3-(Cyclopentyloxy)-4-methoxybenzylamino]-1H-pyrazole-4-methanol, N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-281], N-(3,5-dichloropyridin-4-yl)-2-[5-fluoro-1-(4-fluorobenzyl)-1H-indol-3-yl]-2-oxoacetamide [Research-Code: AWD-12-343], 8-Amino-1,3-bis(cyclopropylmethyl)xanthine [INN:CIPAMFYLLINE], Tetrahydro-5-[4-methoxy-3-[(1S,2S,4R)-2-norbornyloxy]phenyl]-2(1H)-pyrimidone [INN: ATIZORAM], β-[3-(Cyclopentyloxy)-4-methoxyphenyl]-1,3-dihydro-1,3-dioxo-2H-isoindole-2-propanamide [Research - Code: CDC-801], Methanesulfonic acid 2-(2,4-dichlorophenylcarbonyl)-3-ureidobenzo-furan-6-yl ester [INN: LIRIMILAST], (Z)-5-(3,5-di-tert-butyl-4-hydroxybenzylidene)-2-imidazothiazolidin-4-one [INN: DARBUFELONE], cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid [INN: CILOMILAST] and the compounds with the research codes CDC-998, SH-636, D-4396, IC-485 and CC-1088, and the histamine receptor antagonist is selected from the group consisting of (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidyliden)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chlor-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]-ethoxy]acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-

a)azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenzo[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVOCABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl]-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLIN], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-ylmethyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE] and N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELE-NAMINE].

21. A pharmaceutical product according to claim 17 or 18, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST] and (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], and the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]-pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].
22. A pharmaceutical product according to claim 21, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof.

23. A pharmaceutical product according to claim 21, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate, or N-oxide thereof, or solvate of an salt or N-oxide thereof.
24. A pharmaceutical product according to claim 17 or 18, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 5-{4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
2-[4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl]-2H-isopropyl-acetamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and the histamine receptor antagonist is selected from the group consisting of (E)-6-[(E)-3-(1-pyrrolidinyl)-1-p-tolylpropenyl]-2-pyridineacrylic acid [INN: ACRIVASTINE], 6,11-Dihydro-11-(1-methyl-4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin [INN: AZATADINE], 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (+)-(S)-4-[4-[1-(4-chlorophenyl)-1-(2-pyridyl)methoxy]piperidin-1-yl]-butanoic acid [INN: BEPOTASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], (+)-2-[2-[(p-Chloro-alpha-methyl-alpha phenylbenzyl)oxy]ethyl]-1-methylpyrrolidin [INN: CLEMASTINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], [3-(4-Chlorophenyl)-3-pyridin-2-yl-propyl]-dimethylamine [INN: DEXCHLORPHENIRAMINE], 4'-tert-butyl-4-[4-(diphenylmethoxy)piperidino]butyrophenone [INN: EBASTINE], [2-[4-[bis(p-fluorophenyl)methyl]-1-piperazinyl]ethoxy]-acetic acid [INN: EFLETIRIZINE], 1-(2-ethoxyethyl)-2-(hexahydro-4-methyl-1H-1,4-diazepin-1-yl)-benzimidazole [INN: EMEDASTINE], 3-amino-9,13b-dihydro-1H-dibenz[c,f]imidazo[1,5-a]azepine [INN: EPINASTINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], 3-[4-(8-fluoro-5,11-dihydrobenz[b]oxepino[4,3-b]pyridin-11-ylidene)-piperidin-1-yl]propionic acid [Research Code: HSR-609], (-)-(3S,4R)-1-[cis-4-cyano-4-(p-fluorophenyl)cyclohexyl]-3-methyl-4-phenylisonipecotic acid [INN: LEVO-CABASTINE], [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVO-CETIRIZINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE], 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: NORASTEMIZOLE], 3-(10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5-ylidene)-N-methyl-1-propanamine [INN: NORTRIPTYLIN], 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one [INN: PEMIROLAST], 8-chloro-11-[1-(5-methylpyridin-3-yl-methyl)piperidin-4-ylidene]-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridine [INN: RUPATADINE], 1-[2-[(p-chloro-alpha-methyl-alpha-phenylbenzyl)oxy]ethyl]hexahydro-1H-azepine [INN: SETASTINE], S-(7-carboxy-4-hexyl-9-oxoxanthen-2-yl)-S-methylsulfoximine [INN: SUDEXANOX], 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE], N-benzyl-N,N'-dimethyl-N-(2-pyridyl)-ethylenediamine [INN: TRIPELENAMINE], and 1-(4-fluorobenzyl)-2-(piperidin-4-ylamino)-1H-benzimidazole [INN: TECASTEMIZOLE].

25. A pharmaceutical product according to claim 17 or 18, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxyben-

zofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

5-{4-[(4aS,8aR)-4-(3,4-Diethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,

(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-2-[1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl]-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,

2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone

[INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidine-carboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].

26. A pharmaceutical product according to claim 21 or 25, the histamine receptor antagonist is selected from the group consisting of 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], a pharmaceutically acceptable salt or solvate thereof, or a solvate of an salt thereof.
27. Use of a pharmaceutical product according to claim 19 or 24 in the manufacture of a medicament for the treatment of respiratory diseases.
28. A kit comprising a preparation of a first active ingredient which is selected from a PDE4 inhibitor, a PDE3/4 inhibitor and their pharmaceutically acceptable derivatives, a preparation of a second active ingredient which is selected from a histamine receptor antagonist and its pharmaceutically acceptable derivatives, and instructions for simultaneous, sequential or separate administration to the patient in need thereof.
29. A kit according to claim 28, wherein the first and/or second active ingredient is in the form of a pharmaceutically acceptable salt, solvate, N-oxide or solvate of an salt or N-oxide.
30. A kit according to claim 28 or 29, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST] and (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], and the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)-methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-

fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].

31. A kit according to claim 30, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof.
32. A kit according to claim 30, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof.
33. A kit according to claim 28 or 29, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1l⁴-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (cis)-4-(3-Cyclopentylloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1l⁶-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one, (4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

5-{4-[(4aS,8aR)-4-(3,4-Diethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-5-oxo-pentanoic acid,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
(cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-[1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl]-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,

(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
 (4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
 1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,
 4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester,
 2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide and the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].

34. A kit according to claim 30 or 33, wherein the histamine receptor antagonist is selected from the group consisting of 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], a pharmaceutically acceptable salt or solvate thereof, or a solvate of an salt thereof.
35. A method for the treatment of a respiratory disease comprising administering to a patient in need thereof (a) an effective amount of a PDE4 inhibitor, a PDE3/4 inhibitor or a pharmaceutically acceptable derivative thereof and (b) an effective amount of a histamine receptor antagonist or a pharmaceutically acceptable derivative thereof.
36. A method according to claim 35, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST] and (-)-cis-9-ethoxy-8-methoxy-2-methyl-1,2,3,4,4a,10b-hexahydro-6-(4-diisopropylaminocarbonylphenyl)-benzo-[c][1,6]naphthyridine [INN: PUMAFENTRINE].

37. A method according to claim 35, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of
- (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3,4-Dimethoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(tetrahydrothiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3-Chloro-4-methoxyphenyl)-2-(1-oxo-hexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3,4-Diethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(2,3-Dihydro-2,2-dimethyl-7-methoxybenzofuran-4-yl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aR,8aS)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-(cis)-4-(3,4-Dimethoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (cis)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-2-(1,1-dioxohexahydro-1H-thiopyran-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(toluene-4-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-methanesulfonyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- (4aS,8aR)-2-(1-Acetyl-piperidin-4-yl)-4-(3,4-diethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- 5-[4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl]-5-oxo-pentanoic acid,
- (4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(1-pyridin-4-yl-methanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
- 4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,
- 4-[(4aS,8aR)-4-(3,4-Diethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid phenylamide,
- (cis)-4-[4-(7-Methoxy-2,2-dimethyl-2,3-dihydro-benzofuran-4-yl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidine-1-carboxylic acid tert-butylamide,

(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(5-dimethylamino-naphthalene-1-sulfonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-nitro-phenyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-4-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-2-{1-[2-(4-Amino-3,5-dichloro-phenyl)-2-oxo-ethyl]-piperidin-4-yl}-4-(3,4-dimethoxyphenyl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-[1-(1-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-yl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-naphthalen-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-thieno[2,3-d]pyrimidin-4-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyrimidin-2-yl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-oxo-2H-chromen-7-ylmethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
4-(3,4-Dimethoxyphenyl)-2-(1-isopropyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(2-morpholin-4-yl-2-oxo-ethyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-phenethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(morpholine-4-carbonyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-3-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-(1-pyridin-2-ylmethyl-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-[1-(2-morpholin-4-yl-ethanoyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
(4aS,8aR)-4-(3,4-Diethoxyphenyl)-2-(1-{2-[4-(2-dimethylamino-ethyl)-piperazin-1-yl]-ethanoyl}-piperidin-4-yl)-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-isopropyl-acetamide,
(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-2-[1-(4-1,2,3-thiadiazol-4-yl-benzyl)-piperidin-4-yl]-4a,5,8,8a-tetrahydro-2H-phthalazin-1-one,
1-(1-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-methanoyl)-4-ethyl-piperazine-2,3-dione,

4-(2-{4-[(4aS,8aR)-4-(3,4-Dimethoxy-phenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-ethanoylamino)-benzoic acid ethyl ester and
2-{4-[(4aS,8aR)-4-(3,4-Dimethoxyphenyl)-1-oxo-4a,5,8,8a-tetrahydro-1H-phthalazin-2-yl]-piperidin-1-yl}-2H-acetamide.

38. A method according to any of claims 35 to 37, wherein the histamine receptor antagonist is selected from the group consisting of 4-[(4-chlorophenyl)methyl]-2-(hexahydro-1-methyl-1H-azepin-4-yl)-1(2H)phthalazinone [INN: AZELASTINE], (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE], 8-chloro-6,11-dihydro-11-(4-piperidylidene)-5H-benzo[5,6]cyclohepta-[1,2-b]pyridine [INN: DESLORATADINE], (plus/minus)-p-[1-hydroxy-4-[4-(hydroxydiphenylmethyl)piperidino]-butyl]-alpha-methylhydratropic acid [INN: FEXOFENADINE], ethyl 4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidinecarboxylate [INN: LORATADINE], 2-[N-[1-(4-fluorobenzyl)-1H-benzimidazol-2-yl]-4-piperidinyl]-N-methyl-amino]pyrimidin-4(3H)-one [INN: MIZOLASTINE] and 1-(p-tert-butylphenyl)-4-[4'-(alpha-hydroxydiphenylmethyl)-1'-piperidyl]-butanol [INN: TERFENADINE].
39. A pharmaceutical composition according to any of the claims 1 to 7, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE] and its pharmaceutically acceptable derivatives.
40. A pharmaceutical product according to any of the claims 17 to 19, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE] and its pharmaceutically acceptable derivatives.
41. A pharmaceutical composition according to claim 9, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE] and its pharmaceutically acceptable derivatives.

42. A pharmaceutical product according to claim 21, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE] and its pharmaceutically acceptable derivatives.
43. Use of a pharmaceutical composition/product according to any of the claims 39 to 42 in the manufacture of a medicament for the treatment of respiratory diseases.
44. A kit according to claim 28 or 29, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE] and its pharmaceutically acceptable derivatives.
45. A kit according to claim 30, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE] and its pharmaceutically acceptable derivatives.
46. A method according to claim 35, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of [2-[4-[(R)-p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: LEVOCETIRIZINE] and its pharmaceutically acceptable derivatives.
47. A method according to claim 35, wherein the PDE4 inhibitor or PDE3/4 inhibitor is selected from the group consisting of 3-Cyclopropylmethoxy-4-difluoromethoxy-N-(3,5-dichloropyrid-4-yl)-benzamide [INN: ROFLUMILAST], a pharmaceutically acceptable salt, solvate or N-oxide thereof, or solvate of an salt or N-oxide thereof, and the histamine receptor antagonist is selected from the group consisting of (plus/minus)-[2-[4-(p-chloro-alpha-phenylbenzyl)-1-piperazinyl]ethoxy]-acetic acid [INN: CETIRIZINE] and its pharmaceutically acceptable derivatives.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/01876

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4425 A61K31/4523 A61K31/4375 A61K45/06 A61P11/06
A61P11/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 005 865 A (PANACEA BIOTEC LTD) 7 June 2000 (2000-06-07) paragraph '0003!	1-6, 15-18, 35
Y	paragraph '0023! - paragraph '0024! paragraph '0032! paragraph '0058! examples I, VII --- -/--	1-47

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

3 July 2003

Date of mailing of the international search report

24/07/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sindel, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/01876

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANDRI L ET AL: "Combined treatment of allergic rhinitis with terfenadine and nimesulide, a non-steroidal antiinflammatory drug" ALLERGIE ET IMMUNOLOGIE, NOUVELLES EDITIONS MEDICALES FRANCAISES, PARIS, FR, vol. 24, no. 8, October 1992 (1992-10), pages 313-314, 317-319, XP002102388 ISSN: 0397-9148 abstract page 314, column 1, last paragraph ---	1-3,17, 18,28, 29,35
X,P	WO 03 000289 A (NIALS ANTHONY TERENCE ;GLAXO GROUP LTD (GB); WARD PETER (GB); KNOW) 3 January 2003 (2003-01-03) abstract page 5, line 30 claims 1,2,5-8 example A ---	1-9,11, 14-21, 23,26, 27,35, 36, 38-43, 46,47
Y	US 5 962 464 A (HANDLEY DEAN A ET AL) 5 October 1999 (1999-10-05) column 1, line 54 - line 63 column 3, line 11 - line 25 ---	1-47
Y	HATZELMANN A ET AL: "ANTI-INFLAMMATORY AND IMMUNOMODULATORY POTENTIAL OF THE NOVEL PDE4 INHIBITOR ROFLUMILAST IN VITRO" JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, AMERICAN SOCIETY FOR PHARMACOLOGY AND, US, vol. 297, no. 1, 2001, pages 267-279, XP001024814 ISSN: 0022-3565 page 277, paragraph 2 page 277, paragraph 4 page 278, paragraph 2 ---	1-47
Y	WO 01 30777 A (BYK GULDEN LOMBERG CHEM FAB ;STERK JAN GEERT (NL)) 3 May 2001 (2001-05-03) examples 1-10 page 16 page 17, paragraph 6 page 18, paragraph 1 -----	1-47

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/01876

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1005865	A	07-06-2000	BR 9804993 A	06-06-2000
			CN 1253776 A	24-05-2000
			EP 1005865 A1	07-06-2000
WO 03000289	A	03-01-2003	WO 03000289 A1	03-01-2003
US 5962464	A	05-10-1999	US 5900421 A	04-05-1999
			US 2003109540 A1	12-06-2003
			US 6054463 A	25-04-2000
			US 2002040034 A1	04-04-2002
			AU 719907 B2	18-05-2000
			AU 6434898 A	26-08-1998
			BR 9807673 A	15-02-2000
			CN 1246793 T	08-03-2000
			EP 1005345 A1	07-06-2000
			JP 2002514202 T	14-05-2002
			NO 993847 A	27-09-1999
			WO 9834611 A1	13-08-1998
WO 0130777	A	03-05-2001	AU 1515101 A	08-05-2001
			BR 0014990 A	18-06-2002
			CA 2388119 A1	03-05-2001
			CN 1382137 T	27-11-2002
			CZ 20021457 A3	17-07-2002
			WO 0130777 A1	03-05-2001
			EP 1244654 A1	02-10-2002
			HU 0203487 A2	28-02-2003
			JP 2003512466 T	02-04-2003
			NO 20021959 A	29-05-2002
			SK -7232002 A3	10-09-2002
			TR 200201128 T2	21-08-2002
			US 6544993 B1	08-04-2003