



US00PP17071P2

(12) **United States Plant Patent**  
**Trees**

(10) **Patent No.:** **US PP17,071 P2**

(45) **Date of Patent:** **Aug. 29, 2006**

(54) **ANGELONIA PLANT NAMED**  
**'BALANGDARPI'**

(58) **Field of Classification Search** ..... Plt./263  
See application file for complete search history.

(50) Latin Name: *Angelonia angustifolia*  
Varietal Denomination: **Balangdarpi**

(56) **References Cited**  
PUBLICATIONS

(75) Inventor: **Scott C. Trees**, Shell Beach, CA (US)

European Plant Breeders' Rights application No. 2005/2330  
filed Nov. 16, 2005.

(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)

*Primary Examiner*—Kent Bell  
*Assistant Examiner*—Annette H Para  
(74) *Attorney, Agent, or Firm*—Audrey Charles

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

(21) Appl. No.: **11/223,630**

A new and distinct cultivar of *Angelonia* plant named  
'Balangdarpi' characterized by its dark pink-colored  
flowers, good basal branching character, and moderately  
vigorous growth habit.

(22) Filed: **Sep. 9, 2005**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./263**

**2 Drawing Sheets**

**1**

**2**

Latin name of genus and species of plant claimed: *Angelo-*  
*lonia angustifolia*.

Variety denomination: 'Balangdarpi'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of *Angelonia* plant botanically known as *Angelonia angus-*  
*tifolia* and hereinafter referred to by the cultivar name  
'Balangdarpi'.

The new cultivar was developed by the inventor in a  
controlled breeding program during May 2003 at Arroyo  
Grande, Calif. The objective of the breeding program was  
the development of *Angelonia* cultivars with freely  
branching, upright, and moderately vigorous growth habits,  
unique flower colors, and continuous flowering.

The female (seed) parent of the new cultivar was the  
proprietary *Angelonia angustifolia* breeding selection des-  
ignated 608-B, not patented, characterized by its dark pink-  
colored flowers, dark green-colored foliage, and tall upright  
growth habit. The male (pollen) parent of the new cultivar  
was the proprietary *Angelonia angustifolia* breeding selec-  
tion designated 608-A, not patented, characterized by its  
medium pink-colored flowers, dark green-colored foliage,  
and tall upright growth habit. The new cultivar was discov-  
ered and selected by the inventor as a single flowering plant  
within the progeny of the above stated cross-pollination  
during October 2003 in a controlled environment at Arroyo  
Grande, Calif.

Asexual reproduction of the new cultivar by terminal stem  
cuttings since October 2003 at West Chicago, Ill. has dem-  
onstrated that the new cultivar reproduces true to type with  
all the characteristics, as herein described, firmly fixed and  
retained through successive generations of such asexual  
propagation.

**SUMMARY OF THE INVENTION**

The following characteristics of the new cultivar have  
been repeatedly observed and can be used to distinguish

'Balangdarpi' as a new and distinct cultivar of *Angelonia*  
plant:

1. Dark pink-colored flowers;
2. Good basal branching character; and
3. Upright growth habit.

Plants of the new cultivar differ from plants of the female  
parent primarily in growth habit and from plants of the male  
parent primarily in growth habit and flower color.

Of the *Angelonia* cultivars known to the inventor, the  
most similar to 'Balangdarpi' is the *Angelonia* cultivar  
'Balangroki', not patented. However, in side-by-side  
comparisons, plants of the new cultivar differ from plants of  
'Balangroki' in the following characteristics:

1. Plants of the new cultivar have lighter colored flowers  
compared to plants of 'Balangroki'; and
2. Plants of the new cultivar have larger flowers, as  
measured by length and width, compared to plants of  
'Balangroki'.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs show, as nearly true as it  
is reasonably possible to make the same in color illustrations  
of this type, typical flower and foliage characteristics of the  
new cultivar. Colors in the photographs differ slightly from  
the color values cited in the detailed description, which  
accurately describe the colors of 'Balangdarpi'. The plants  
were grown in 10 cm pots for 11 weeks in a greenhouse at  
West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and  
flowering habit of 'Balangdarpi'.

FIG. 2 illustrates a close-up view of the inflorescence of  
'Balangdarpi'.

FIG. 3 illustrates a closer view of individual flowers of the  
inflorescence of 'Balangdarpi'.

FIG. 4 illustrates a close-up view of an individual flower  
of 'Balangdarpi'.

## DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where color terms of ordinary significance are used. The color values were determined on Apr. 13, 2005. The readings were taken between 1:00 p.m. and 3:00 p.m. under natural light conditions.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a double polycarbonate-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown in 10 cm pots for 11 weeks while utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 75°–85° F. (24°–29° C.) during the day and approximately 62°–70° F. (17°–21° C.) during the night. Greenhouse light levels were maintained at approximately 6,000 to 10,000 footcandles during the day.

Botanical classification: *Angelonia angustifolia*, cultivar Balangdarpi.

Parentage:

*Male (pollen) parent*.—Proprietary *Angelonia angustifolia* breeding selection 608-A, not patented.

*Female (seed) parent*.—Proprietary *Angelonia angustifolia* breeding selection designated 608-B, not patented.

Propagation:

*Type cutting*.—Terminal stem.

*Time to initiate roots*.—Approximately 7 to 9 days.

*Time to develop roots*.—Approximately 21 to 28 days.

*Root description*.—Fine and fibrous.

*Rooting habit*.—Freely branching.

Plant description:

*Crop time*.—Approximately 6 to 7 weeks from a rooted cutting.

*Habit of growth*.—Moderately vigorous with good branching.

*Form*.—Upright.

*Size*.—Height (from soil level to top of plant plane):

Approximately 26.4 cm. Diameter (area of spread):

Approximately 36.4 cm in diameter.

*Main branch*.—Quantity per plant: Approximately 5.

Shape: Square in cross section. Strength: Strong.

Length from soil level to base of raceme: Approximately 32.4 cm. Diameter: Approximately 3.1 mm.

Texture: Glabrous. Color: 145B. Internode length at

middle of branch: Approximately 2.1 cm.

*Foliage*.—Quantity of leaves per main branch:

Approximately 22. Type: Simple. Fragrance: None.

Arrangement: Opposite. Orientation to stem: Obtuse.

Shape: Elliptic. Margin: Serrate. Apex: Acuminate.

Base: Sessile. Length of leaf taken from middle of

branch: Approximately 7 cm. Width of leaf taken

from middle of branch: Approximately 1.3 cm. Texture

of upper and lower surfaces: Glabrous. Venation

pattern: Pinnate. Color of upper surface of young and

mature foliage: Darker than 137A with venation of

143B. Color of lower surface of young and mature foliage: 137C with venation of 145C.

Flowering description:

*Flowering habit*.—‘Balangdarpi’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year round in greenhouse environment.

*Time to first flower*.—Approximately 11 weeks from planting of rooted cutting.

*Lastingness of individual bloom*.—Approximately 7 to 10 days.

*Inflorescence type/description*.—Terminal racemes.

Length: 13.9 cm. Width: 4.8 cm. Number per plant:

Approximately 4 open racemes per plant. Number of

fully open flowers per raceme at any one time:

Approximately 13.

Flower description:

*Type/fragrance*.—Bi-labiate. No fragrance.

*Bud*.—Rate of opening: 3 to 4 days from first color to fully open. Shape: Globular. Length: Approximately 9.6 mm. Diameter: Approximately 7.3 mm. Color of upper surface: 48C with 51A at base. Color of lower surface: 48C with spots of N77A.

*Flower size/aspect*.—Length: Approximately 2.5 cm.

Width: Approximately 2.1 cm. Depth: Approx-

imately 6.2 mm. Aspect: Facing outward.

*Petals*.—Quantity: Five per flower, fused at base forming a throat, an upper lip having two petals and a lower lip having three petals, one central and one lateral on each side.

*Upper lip*.—Petal apex: Obtuse. Petal margin: Entire, undulate. Length of petals: Approximately 7.2 mm. Width of petals: Approximately 9.8 mm. Color of upper and lower surfaces: First open closest to 65A, fully mature 63C. Petal texture upper surface: Densely glandular-pubescent. Petal texture lower surface: Glabrous.

*Lower lip, lateral petals*.—Petal apex: Obtuse. Petal margin: Entire. Length of petals from throat: Approximately 8.2 mm. Width of petals: Approximately 9.4 mm. Color of upper surface: First open N155B with 65A at margin, fully mature 63C. Color of lower surface: First open N155A with 65A at margin, fully mature 63C. Petal texture upper and lower surfaces: Densely pubescent with very short glandular hairs.

*Lower lip, central petal*.—Petal apex: Obtuse. Petal margin: Entire, undulate. Length of petal from palate: Approximately 8 mm. Width of petal: Approximately 9.6 mm. Color of upper and lower surfaces: First open N155B with 65A at edges, fully mature 65A at margin only, 65D in center and at base. Petal texture upper and lower surfaces: Densely pubescent with very short glandular hairs. Gland color: 13B, translucent.

*Throat*.—Length: Approximately 5.7 mm. Width: Approximately 6.8 mm. Texture of inner and outer surfaces: Glabrous. Color of inner and outer surfaces: N81D with streaks of N81A. Palate color: N155D with spots at base of N81A. Palate texture: Glabrous, pubescent along margin. Teeth color: N155D with 145B at base.

*Pedicel*.—Strength: Good. Length: Approximately 7.7 mm. Diameter: Approximately 0.7 mm. Angle to stem: Acute. Texture: Glabrous. Color: 144B with overlay of 187A.

*Calyx*.—Shape: Five-pointed star, cupped. Width: Approximately 5.0 mm. Sepal number: 5. Sepal

## 5

shape: Lanceolate. Sepal margin: Entire. Sepal apex: Acuminate. Sepal length: Approximately 4.3 mm. Sepal width: Approximately 1.8 mm. Sepal texture of upper and lower surfaces: Glabrous. Sepal color of upper and lower surfaces: 143A.

*Reproductive organs.*—Androecium: Stamen quantity: Four per flower. Stamen length: Approximately 3.0 mm. Filament color: N155D. Anther shape: Bi-lobed. Anther length: Approximately 1.0 mm. Anther color: Closest to 24D. Amount of pollen: None observed. Gynoecium: Pistil quantity: One per flower. Pistil length: Approximately 3.0 mm. Stigma

## 6

length: Approximately 0.5 mm. Stigma color: 150D. Style length: Approximately 1.5 mm. Style color: 155B. Ovary diameter: Approximately 1.0 mm. Ovary texture: Puberulent. Ovary color: 144C.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Angelonia* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Angelonia* plant named 'Balangdarpi' substantially, as herein shown and described.

\* \* \* \* \*



FIG. 1




FIG. 2




FIG. 3




FIG. 4