



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 086 402 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:

06.04.2005 Bulletin 2005/14

(21) Application number: **00909163.8**

(22) Date of filing: **14.02.2000**

(51) Int Cl.7: **G03C 7/413**

(86) International application number:
PCT/EP2000/001186

(87) International publication number:
WO 2000/054106 (14.09.2000 Gazette 2000/37)

(54) **NEW COLOR PHOTOGRAPHIC DEVELOPER KIT**

NEUER SATZ FÜR FARBPHTOGRAPHISCHEN ENTWICKLER

NOUVEAU KIT POUR LA PREPARATION DE REVELATEUR PHOTOGRAPHIQUE CHROMOGENE

(84) Designated Contracting States:
DE FR GB

(30) Priority: **08.03.1999 FR 9902972**

(43) Date of publication of application:
28.03.2001 Bulletin 2001/13

(73) Proprietor: **EASTMAN KODAK COMPANY**
Rochester, New York 14650-2201 (US)

(72) Inventor: **THOMAS, Françoise, Marie,**
Kodak Industrie
71102 Chalon sur Saone Cedex (FR)

(74) Representative: **Weber, Etienne Nicolas et al**
Kodak Industrie,
Département Brevets,
CRT,
Zone Industrielle
71102 Chalon sur Saône Cedex (FR)

(56) References cited:
FR-A- 2 214 910 **GB-A- 1 468 015**
US-A- 3 721 563

Remarks:

The file contains technical information submitted
after the application was filed and not included in this
specification

EP 1 086 402 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**FIELD OF THE INVENTION**

5 **[0001]** This invention concerns a photographic developer kit for processing color photographic materials, and a method for preparing a developer from said kit.

BACKGROUND OF THE INVENTION

10 **[0002]** In the color photographic film processing, color developers are used that contain as their main constituent a color developer in an alkaline-medium. Such a color developer also contains other constituents such as permeabilizing agents, antifogging agents, preservatives, etc.

15 **[0003]** It is known that the different constituents of a color developer can be packaged separately. It is thus obtained kits for photographic processing, generally comprising several parts each containing one or more constituents of the developer. These different parts are mixed, and if necessary diluted by the end user, to obtain the ready-to-use color developer. Such packaging in kit form is necessary because the different constituents of the color developer become unstable once present together in the same solution. Further, photographic processing kits are designed to facilitate the preparation of the ready-to-use developer by a non-specialized end user.

20 **[0004]** For processing motion picture film, there exists a kit comprising two concentrated solutions, called concentrates, namely an alkaline concentrate and a concentrate containing the developer, and a solid part consisting of 3,5-dinitrobenzoic acid as a wet powder. This arrangement does not allow the automated manufacture of the kit, because the packaging of the powder is difficult to industrialize.


25 **[0005]** In addition, to obtain a homogeneous developer from these liquid concentrates and this solid part, it is necessary to mix the different components of the kit in a certain order, so that the powder dissolves completely.

SUMMARY OF THE INVENTION

30 **[0006]** One object of the present invention is to provide a color developer kit with a simplified packaging and improved stability, that is easier to industrialize. A further object of the invention is to provide the end user with a kit that allows a simple, rapid, and reproducible preparation of the ready-to-use color developer. A further object of the invention is to provide a kit that affords an efficient ready-to-use color developer.

[0007] These objects are achieved by the present invention, which concerns a photographic developer kit comprising two concentrates.

35 **[0008]** One of the concentrates, Concentrate (A), is a basic aqueous solution having a pH greater than or equal to 8 containing a compound of formula (I) :



(I)

45

where X is -COOH or -SO₃H, M is an alkali metal or ammonium, R is an alkyl group with preferably from 1 to 3 atoms of carbon, and n is 0, 1, 2 or 3.

50 **[0009]** The second concentrate, Concentrate (B), is an aqueous acidic solution containing a color developer of the paraphenylenediamine type.

[0010] The redox potential of compound (I) is preferably greater than -700 mV.

[0011] The redox potential of compound (I) is measured against a Ag/AgCl/KCl, 3M reference electrode.

55 **[0012]** The invention further concerns the use of this kit for the preparation of a ready-to-use color developer, and a method for the preparation of a ready-to-use color developer that comprises mixing the two concentrates (A) and (B) in any order.

[0013] In one embodiment, the position of the nitro group is meta or para to the X group, and n is 0, i.e., the benzene ring bears no alkyl group. In another embodiment, the nitro group is para or meta to the X group, R is an alkyl group having from 1 to 3 carbon atoms and n is 1, 2 or 3.

[0014] In the scope of the invention, the compound of formula (I) can be 3-nitrobenzoic acid (redox potential -650 mV), 4-nitrobenzoic acid (redox potential - 580 mV), 3-nitrobenzenesulfonic acid (redox potential -600 mV), and 4-nitrobenzenesulfonic acid. Because the compound (I) is dissolved in a basic solution, it occurs as a salt, for example a salt of sodium, potassium, lithium or ammonium.

[0015] In any cases the volume and concentration of the concentrate (A) will be adjusted to obtain a ready-to-use developer containing between 2×10^{-4} mole/l and 3×10^{-3} mole/l (0.350 g/l) of compound (I).

[0016] In a specific embodiment, the concentrate (A) contains the nitrobenzoic acid as its sodium salt at a concentration between 2.5×10^{-3} mole/l and 15×10^{-3} mole/l.

[0017] In a specific embodiment, the concentration of the sodium salt of the nitrobenzoic acid in the concentrate (A) is about 7.5×10^{-3} mole/l.

[0018] The kit of this invention has a particularly simple packaging because it comprises only two liquid concentrates. It allows a rapid and simple preparation of the ready-to-use color developer by the end user. Photographic materials processed from the kit of the invention exhibit good sensitometric results, in particular low fogging without reduced rapidity.

[0019] The concentrate (A) useful in this invention is a basic solution obtained from alkaline compounds such as sodium or potassium carbonate, borax, sodium or potassium hydroxide, or sodium metaborate, in aqueous solution. This concentrate (A) can contain chelating agents, water softeners such as aminopolycarboxylic acids, for example ethylenediaminetetra-acetic acid (EDTA), diethylenetriaminepenta-acetic acid (DTPA), isopropanoldiaminetetra-acetic acid (DPTA), aminopolyphosphonic acids, for example amino-N,N dimethylenephosphonic acids, hexametaphosphate, Dequest® (2000, 2006, 2010, etc.), and Versenex 80®.

[0020] The volume and the pH of the concentrate (A) will be adjusted to obtain a ready-to-use developer with a pH of at least 8, preferably between 10 and 12.

[0021] The developer used in the concentrate (B) is a p-phenylenediamine, for example 2-amino-5-diethylaminotoluene (known as CD2), 4-amino-N-ethyl-N-(β-methanesulfoamidoethyl)-m-toluidine (CD3), 4-amino-3-methyl-N-ethyl-N-(β-hydroxyethyl)aniline (CD4). CD2 is generally used in color developers for positive motion picture films, CD3 is generally used in color developers for negative motion picture films, and intermediate motion picture films.

[0022] The developer concentration and the volume of the concentrate (B) will be adjusted to obtain a ready-to-use developer containing a developer concentration of at least 7×10^{-3} mole/l, and preferably between 9×10^{-3} and 2×10^{-2} mole/l.

[0023] The concentrate (B) containing the color developer can contain other compounds such as for example antioxidants or surfactants. The antioxidants that can be used in the concentrate (B) are for example alkali metal sulfites, metabisulfites and bisulfites, sulfur compounds able to generate sulfite ions in aqueous solution, ascorbic acid and its derivatives, and hydroxylamine derivatives, etc.

[0024] In a specific embodiment, the concentrate (B) contains CD3 as color developer, and sulfite. The pH of this concentrate is kept acidic, and is less than or equal to 3.

[0025] These concentrates are mixed at the time of use either to prepare a color developer, or to prepare a replenishment or maintenance solution to maintain the efficiency of the developer during use. In the scope of the invention, the concentrates can be mixed in any order. To make the preparation of the ready-to-use developer even easier, the volumes of the concentrates (A) and (B) can be such that mixing the concentrates yields one liter of color developer without having to dilute the mixture.

[0026] The concentrates (A) and (B) can contain other compounds, for example antiseptics, heat stabilizers, developing activators such as thioether or oxothioether compounds, or benzylamine.

[0027] After mixing the concentrates, it may be necessary to adjust the pH to a value advantageously between 10.0 and 11.0, or to buffer the mixture to obtain a ready-to-use color developer.

[0028] In a specific embodiment, the kit of the present invention is designed for the preparation of a color developer for negative motion picture films such as Eastman Color Negative® marketed by Kodak. Conventionally, this process comprises a color development step in the presence of CD3, a bleaching step, and a fixing step. The bleaching step and the fixing step can be replaced by a single bleaching-fixing step. One or more washing baths can be inserted between these successive steps.

[0029] The present invention is illustrated by the following examples.

EXAMPLE 1: Preparation of concentrate A

[0030] In a vessel fitted with a magnetic stirrer containing 950 ml of distilled water, were added 4.0 g of NaBr, 125.8 g of Na_2CO_3 , 3.02 g of NaHCO_3 , and 19.3 g of DEQUEST 2006 chelating agent in 40% solution. To this stirred solution was added 7.5×10^{-3} moles of 3,5-dinitrobenzoic acid as indicated in Table I below. Nitrobenzoic acids marketed by Aldrich were used in dry powder form. This solution was stirred for 30 minutes to obtain complete dissolution of the nitrobenzoic acid in the solution. Demineralized water was added to this solution to obtain 1 liter of solution. The pH

of the solution was 10.9.

TABLE I

	Concentration in the concentrate A (g/l)
A - 3,5-dinitrobenzoic acid (control)	1.58
B - 3-nitrobenzoic acid	1.247
C - 4-nitrobenzoic acid	1.247
D - 3-nitrobenzene sulfonic acid	1.683

[0031] All these concentrations in g/l correspond to 7.5×10^{-3} mole/liter.

[0032] In this way the concentrate (A) was obtained containing the nitrobenzoic acid as its sodium salt.

EXAMPLE 2

[0033] A sample of the concentrate (A) prepared by the procedure of example 1 was kept in a plastic bottle at 50°C for times indicated in Table II below. Samples of the concentrate were taken at 9, 20, 30, 37 and 83 days, and the quantity of nitrobenzoic acid remaining (quantity of salt) was measured. The variation in the amount of nitrobenzoic acid between the freshly prepared concentrate and the concentrate at time t (expressed in %) is given in the table below. The quantity of nitrobenzoic acid was measured by HPLC (variability of measurement +/- 2%).

[0034] The following results were obtained:

TABLE II

number of days	Variation (%) of nitrobenzoic acid			
	AF-9	3NBA	4NBA	3NBSA
0	0	0	0	0
9	0	0	0	0
20	0	0	0	0
30	2.10	0	0	0
37	4.66	0	0	0
83	9.87	0	0	0

AF-9 : 3,5-dinitrobenzoic acid

3NBA : 3-nitrobenzoic acid

4NBA : 4-nitrobenzoic acid

3NBSA : 3-nitrobenzene sulfonic acid

[0035] No crystallization was observed during this time (crystallization test carried out at 5°C).

EXAMPLE 3

[0036] In this example, 1 liter of ready-to-use developer was prepared from the concentrate (A) prepared previously, and a concentrate (B) with the following composition:

Concentrate (B) (1 liter)	
demineralized water	912 ml
anhydrous sodium sulfite	52.9 g
CD3	116.3 g
pH at 25°C	3

[0037] To obtain 1 liter of ready-to-use developer (replenisher), 198.5 ml of concentrate (A) was mixed with 47.25 ml of the concentrate (B) of example 1.

[0038] An Eastman Color Intermediate 5274® negative color film was exposed through a step tablet with 21 density ranges, each of these ranges having an increment of 0.15 LogE with an exposure light of color temperature 2,850 K and a D1 illuminator (Tungsten) for 1/50 sec. The film was developed using an ECN-2® photographic process, which comprises the developer prepared in example 3, a bleaching bath, a fixing bath, and a final washing bath, the film and the process both being commercialized by Kodak.

[0039] By reading in the three colors blue, green and red on a densitometer, the following sensitometric results were obtained (freshly prepared baths).

	Red-sensitive layer	Green-sensitive layer	Blue-sensitive layer
Dmin (+/- 0.04)	0.180 (0.177)	0.560 (0.557)	0.926 (0.924)
Dmax (+/- 0.10)	1.467 (1.468)	2.050 (2.053)	2.327 (2.331)
Contrast (+ 0.04)	0.532 (0.534)	0.589 (0.590)	0.578 (0.573)
Speed (+ 4)	523.6 (523.5)	530.5 (530.1)	518.8 (519.9)

(): value obtained when the developer contained 3,5-dinitrobenzoic acid.

Dmin = minimal density corresponding to an unexposed part of the film (support + fog).

Dmax = maximal density of film.


Speed = $100(3 - \text{Log}E)$, E being the exposure at the density point Dmin + 1.

Contrast = slope of the straight line between the point of density Dmin + 0.20 and that of the exposure above 1.35 LogE.

Claims

1. A kit for a color photographic developer comprising only two concentrated solutions wherein

(i) one of the concentrated solutions is a basic solution having a pH greater than or equal to 8, and containing at least one compound of the formula:



(I)

wherein X is -COOM or -SO₃M with M selected from the group consisting of hydrogen, alkali metal counter ion and ammonium counter ions, R is an alkyl group having from 1 to 3 carbon atoms, and n is equal to 0, 1, 2 or 3, and


(ii) the second concentrated solution is an acidic aqueous solution having a pH less than or equal to 3 comprising a paraphenylenediamine color developing agent.

- 15
2. The kit according to claim 1 wherein the compounds of formula (I) have a redox potential greater than -700 mV measured against a Ag/AgCl/KCl, 3M reference electrode
- 20
3. The kit according to claim 1 wherein the compound of formula (I) is selected from the group consisting of 3-nitrobenzoic acid, 4-nitrobenzoic acid, 3-nitrobenzene sulfonic acid, and 4-nitrobenzene sulfonic acid.
- 25
4. The kit according to claim 1 wherein the compound (I) is present in the concentrated solution in a quantity between 2.5×10^{-3} mole/l and 15×10^{-3} mole/l.
5. The kit according to claim 1 wherein the color developing agent is present in the developer in a quantity between 9×10^{-3} and 2×10^{-2} mole/l.
- 30
6. Use of the kit as defined according to any of claims 1 to 5 for the preparation of a color developer for the development of color photographic materials.
7. Method of preparation of a color developer which comprises the steps of mixing the basic solution with the acidic solution as defined according to any of claims 1 to 5 in order to obtain the homogenous color developer.
- 35

Patentansprüche

- 40
1. Aus nur zwei konzentrierten Lösungen bestehender Satz für eine farbfotografische Entwicklerlösung, worin

(i) es sich bei der einen der konzentrierten Lösungen um eine basische Lösung mit einem pH-Wert gleich oder größer als 8 handelt, die mindestens eine Verbindung der Formel:



(I)

55 umfasst, in der X -COOM oder -SO₃M ist, M aus der aus Wasserstoff, Alkalimetall- und Ammonium-Gegenionen bestehenden Gruppe ausgewählt wird, R eine Alkyl-Gruppe mit 1 bis 3 Kohlenstoff-Atomen darstellt, und n gleich 0, 1, 2 oder 3 ist, und


EP 1 086 402 B1

(ii) die zweite konzentrierte Lösung eine saure wässrige Lösung mit einem pH-Wert kleiner als oder gleich 3 ist, die ein Paraphenylendiamin-Farbentwicklungsmittel umfasst.

- 5 2. Entwicklersatz nach Anspruch 1, worin die Verbindungen der Formel (I) ein gegen eine Ag/AgCl/KCl, 3M-Bezugselektrode gemessenes Redoxpotential aufweisen, das größer als -700 mV ist.
3. Entwicklersatz nach Anspruch 1, worin die Verbindung der Formel (I) aus der aus 3-Nitrobenzoesäure, 4-Nitrobenzoesäure, 3-Nitrobenzolsulfonsäure und 4-Nitrobenzolsulfonsäure bestehenden Gruppe ausgewählt wird.
- 10 4. Entwicklersatz nach Anspruch 1, worin die Verbindung (I) in der konzentrierten Lösung in Mengen zwischen $2,5 \times 10^{-3}$ mol/l und 15×10^{-3} mol/l vorliegt.
- 5 15 5. Entwicklersatz nach Anspruch 1, worin das Farbentwicklungsmittel in der Entwicklerlösung in Mengen zwischen 9×10^{-3} und 2×10^{-2} mol/l vorliegt.
6. Verwendung des Entwicklersatzes nach einem der Ansprüche 1 bis 5 für die Herstellung einer Farbentwicklerlösung zum Entwickeln von farbfotografischen Materialien.
- 20 7. Verfahren zur Herstellung einer Farbentwicklerlösung, mit dem Schritt: Vermischen der basischen Lösung mit der sauren Lösung nach einem der Ansprüche 1 bis 5 zur Gewinnung der homogenen Farbentwicklerlösung.

Revendications

- 25 1. Kit pour révélateur photographique chromogène comprenant uniquement deux solutions concentrées, où
(i) l'une des solutions concentrées est une solution basique ayant un pH égal ou supérieur à 8 et contenant au moins un composé représenté par la formule :



(I)

- 30 40 dans laquelle X est -COOM ou -SO₃M, M étant choisi parmi le groupe comprenant un atome d'hydrogène, un contre ion de métal alcalin et des contre ions ammonium, R est un groupe alkyle de 1 à 3 atomes de carbone et n est égal à 0, 1, 2 ou 3, et
45 (ii) la seconde solution concentrée est une solution aqueuse acide ayant un pH inférieur ou égal à 3 comprenant un développeur chromogène paraphénylènediamine.
2. Kit selon la revendication 1, dans lequel les composés de formule (I) ont un potentiel d'oxydo-réduction supérieur à -700 mV, mesuré contre une électrode de référence 3M Ag/AgCl/KCl.
- 50 3. Kit selon la revendication 1, dans lequel le composé de formule (I) est choisi parmi le groupe comprenant l'acide 3-nitrobenzoïque, l'acide 4-nitrobenzoïque, l'acide 3-nitrobenzène sulfonique et l'acide 4-nitrobenzène sulfonique.
4. Kit selon la revendication 1, dans lequel on utilise, dans la solution concentrée, une quantité de composé (I) comprise entre $2,5 \times 10^{-3}$ moles/l et 15×10^{-3} moles/l.
- 55 5. Kit selon la revendication 1, dans lequel on utilise, dans le révélateur, une quantité de développeur chromogène comprise entre 9×10^{-3} et 2×10^{-2} moles/l.

EP 1 086 402 B1

6. Utilisation du kit tel que défini selon l'une quelconque des revendications 1 à 5 pour la préparation d'un révélateur chromogène pour le développement de produits photographiques en couleurs.
7. Procédé de préparation d'un révélateur chromogène qui comprend le mélange de la solution basique et de la solution acide telles que définies selon l'une quelconque des revendications 1 à 5, afin d'obtenir un révélateur chromogène homogène.

5

10

15

20

25

30

35

40

45

50

55