



(43) International Publication Date
3 November 2016 (03.11.2016)

(10) International Publication Number
WO 2016/173996 A1

- (51) **International Patent Classification:**
A01P 3/00 (2006.01) *A01N 43/56* (2006.01)
- (21) **International Application Number:**
PCT/EP2016/059213
- (22) **International Filing Date:**
26 April 2016 (26.04.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
15165753.3 29 April 2015 (29.04.2015) EP
- (71) **Applicant:** **BAYER CROPSCIENCE AK-
TIENGESELLSCHAFT** [DE/DE]; Alfred-Nobel-Str. 50,
40789 Monheim am Rhein (DE).
- (72) **Inventors:** **HILLEBRAND, Stefan**; Lothringer Str. 22,
41462 Neuss (DE). **TAFFOREAU, Sylvain**; 50, Rue Phil-
ippe de Lassalle, 69004 Lyon (FR).
- (74) **Agent:** **BIP PATENTS**; c/o Bayer Intellectual Property
GmbH, Alfred-Nobel-Str. 10, 40789 Monheim am Rhein
(DE).
- (81) **Designated States** (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*

(54) **Title:** USE OF 2-{3-[2-(1-{[3,5-BIS(DIFLUOROMETHYL)-1H-PYRAZOL-1-YL]ACETYL} PIPERIDIN-4-YL)-1,3-
THIAZOL-4-YL]-4,5-DIHYDRO-1,2-OXAZOL-5-YL}-3-CHLOROPHENYL METHANESULFONATE

(57) **Abstract:** The present invention relates to the use of 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-
yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate (see Figure 1) and of compositions comprising
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-
chlorophenyl methanesulfonate and further fungicides or insecticides for controlling downy mildew diseases of monocotyledon
crops. Furthermore the treatment of seed is described to control downy mildew diseases of monocotyledon crops.



WO 2016/173996 A1

Use of 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate

The present invention relates to the use of 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate (see Figure 1) and of compositions comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and further fungicides or insecticides for controlling downy mildew diseases of monocotyledon crops. Furthermore the treatment of seed is described to control downy mildew diseases of monocotyledon crops.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate itself is known from WO WO2012/025557. 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate is a fungicide especially useful for the control of oomycetes such as *Phytophthora* species, for example *Phytophthora infestans* and *Plasmopara* species, for example *Plasmopara viticola*.

This invention now provides an advantageous use of compound of formula (I), 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate, and of compositions comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and further active ingredients against downy mildew in monocotyledon crops, such as *Peronosclerospora philippinensis* in corn, sorghum or oat, *Peronosclerospora* in sugarcane, *Sclerophthora rayssiae varzeae* in corn, *Sclerophthora macrocarpa* in corn, wheat, barley or rice, *Sclerospora graminicola* in millets, *Peronosclerospora maydis* in corn or teosinte, *Peronosclerospora sorghi* in sorghum.




Figure 1: 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control

Peronosclerospora philippinensis in corn, sorghum or oat.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control *Peronosclerospora* in sugarcane.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control *Sclerophthora rayssiae varzeae* in corn.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control *Sclerophthora macrocarpa* in corn, wheat, barley or rice.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control *Sclerospora graminicola* in millets.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control *Peronosclerospora maydis* in corn or teosinte.

2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticide are able to control *Peronosclerospora sorghi* in sorghum.

In conjunction with the present invention “controlling” denotes a significant reduction of the downy

mildew infestation in comparison to the untreated crop, more preferably the infestation is essentially diminished (50-79%), most preferably the infestation is totally suppressed (80-100%).

In an alternative embodiment of the present invention 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate
5 as such or formulations thereof can be applied in combination with further known fungicides, bactericides, acaricides, nematocides, herbicides, insecticides for broadening the activity spectrum and in order to prevent the development of resistance. In many instances, synergistic effects are obtained, i.e. the activity of the mixture exceeds the activity of the individual components.

According to the invention the expression "*composition*" stands for the various combinations of
10 compounds A) and B), for example in a single "ready-mix" form, in a combined spray mixture composed from separate formulations of the single active compounds, such as a "tank-mix", and in a combined use of the single active ingredients when applied in a sequential manner, i.e. one after the other with a reasonably short period, such as a few hours or days. Preferably the order of applying the compounds A) and B) is not essential for working the present invention.

15 Formulations

The present invention further relates to a composition for controlling unwanted microorganisms, comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate. These are preferably fungicidal compositions which comprise agriculturally suitable auxiliaries, solvents, carriers, surfactants or extenders.

20 According to the invention, a carrier is a natural or synthetic, organic or inorganic substance with which the active ingredients are mixed or combined for better applicability, in particular for application to plants or plant parts or seed. The carrier, which may be solid or liquid, is generally inert and should be suitable for use in agriculture.

Useful solid carriers include: for example ammonium salts and natural rock flours, such as kaolins, clays,
25 talc, chalk, quartz, attapulgit, montmorillonite or diatomaceous earth, and synthetic rock flours, such as finely divided silica, alumina and silicates; useful solid carriers for granules include: for example, crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, and also synthetic granules of inorganic and organic flours, and granules of organic material such as paper, sawdust, coconut shells, maize cobs and tobacco stalks; useful emulsifiers and/or foam-formers include: for example nonionic
30 and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates and also protein hydrolysates; suitable dispersants are nonionic and/or ionic substances, for example from the classes of the alcohol-POE and/or -POP ethers, acid and/or POP POE esters, alkylaryl and/or POP POE ethers, fat and/or POP POE adducts, POE- and/or POP-polyol derivatives, POE- and/or POP-sorbitan or -sugar adducts, alkyl

or aryl sulphates, alkyl- or arylsulphonates and alkyl or aryl phosphates or the corresponding PO-ether adducts. Additionally suitable are oligo- or polymers, for example those derived from vinylic monomers, from acrylic acid, from EO and/or PO alone or in combination with, for example, (poly)alcohols or (poly)amines. It is also possible to use lignin and its sulphonic acid derivatives, unmodified and modified
5 celluloses, aromatic and/or aliphatic sulphonic acids and also their adducts with formaldehyde.

The active ingredients can be converted to the customary formulations, such as solutions, emulsions, wettable powders, water- and oil-based suspensions, powders, dusts, pastes, soluble powders, soluble granules, granules for broadcasting, suspoemulsion concentrates, natural products impregnated with active ingredient, synthetic substances impregnated with active ingredient, fertilizers and also
10 microencapsulations in polymeric substances.

The active ingredients can be applied as such, in the form of their formulations or the use forms prepared therefrom, such as ready-to-use solutions, emulsions, water- or oil-based suspensions, powders, wettable powders, pastes, soluble powders, dusts, soluble granules, granules for broadcasting, suspoemulsion concentrates, natural products impregnated with active ingredient, synthetic substances impregnated with
15 active ingredient, fertilizers and also microencapsulations in polymeric substances. Application is accomplished in a customary manner, for example by watering, spraying, atomizing, broadcasting, dusting, foaming, spreading-on and the like. It is also possible to deploy the active ingredients by the ultra-low volume method or to inject the active ingredient preparation/the active ingredient itself into the soil. It is also possible to treat the seed of the plants.

20 The formulations mentioned can be prepared in a manner known per se, for example by mixing the active ingredients with at least one customary extender, solvent or diluent, emulsifier, dispersant and/or binder or fixing agent, wetting agent, a water repellent, if appropriate siccatives and UV stabilizers and if appropriate dyes and pigments, antifoams, preservatives, secondary thickeners, stickers, gibberellins and also other processing auxiliaries.

25 The present invention includes not only formulations which are already ready for use and can be deployed with a suitable apparatus to the plant or the seed, but also commercial concentrates which have to be diluted with water prior to use.

The 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate may be present as such or in their (commercial)
30 formulations and in the use forms prepared from these formulations as a mixture with other (known) active ingredients, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators, herbicides, fertilizers, safeners and/or semiochemicals.

The auxiliaries used may be those substances which are suitable for imparting particular properties to the

composition itself or and/or to preparations derived therefrom (for example spray liquors, seed dressings), such as certain technical properties and/or also particular biological properties. Typical auxiliaries include: extenders, solvents and carriers.

Suitable extenders are, for example, water, polar and nonpolar organic chemical liquids, for example from the classes of the aromatic and nonaromatic hydrocarbons (such as paraffins, alkylbenzenes, alkylnaphthalenes, chlorobenzenes), the alcohols and polyols (which may optionally also be substituted, etherified and/or esterified), the ketones (such as acetone, cyclohexanone), esters (including fats and oils) and (poly)ethers, the unsubstituted and substituted amines, amides, lactams (such as N-alkylpyrrolidones) and lactones, the sulphones and sulfoxides (such as dimethyl sulphoxide).

- 10 Liquefied gaseous extenders or carriers are understood to mean liquids which are gaseous at standard temperature and under standard pressure, for example aerosol propellants such as halohydrocarbons, or else butane, propane, nitrogen and carbon dioxide.

In the formulations it is possible to use tackifiers such as carboxymethylcellulose, natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids such as cephalins and lecithins and synthetic phospholipids. Further additives may be mineral and vegetable oils.

If the extender used is water, it is also possible to use, for example, organic solvents as auxiliary solvents. Useful liquid solvents are essentially: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example petroleum fractions, alcohols such as butanol or glycol and their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulphoxide, or else water.

Compositions comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate may additionally comprise further components, for example surfactants. Suitable surfactants are emulsifiers and/or foam formers, dispersants or wetting agents having ionic or nonionic properties, or mixtures of these surfactants. Examples thereof are salts of polyacrylic acid, salts of lignosulphonic acid, salts of phenolsulphonic acid or naphthalenesulphonic acid, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (preferably alkylphenols or arylphenols), salts of sulphosuccinic esters, taurine derivatives (preferably alkyl taurates), phosphoric esters of polyethoxylated alcohols or phenols, fatty esters of polyols, and derivatives of the compounds containing sulphates, sulphonates and phosphates, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates, protein hydrolysates, lignosulphite waste liquors and methylcellulose. The presence of a surfactant is

necessary if one of the active ingredients and/or one of the inert carriers is insoluble in water and when application is effected in water. The proportion of surfactants is between 5 and 40 per cent by weight of the inventive composition.

It is possible to use dyes such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyes such as alizarin dyes, azo dyes and metal phthalocyanine dyes, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Further additives may be perfumes, mineral or vegetable, optionally modified oils, waxes and nutrients (including trace nutrients), such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Additional components may be stabilizers, such as cold stabilizers, preservatives, antioxidants, light stabilizers, or other agents which improve chemical and/or physical stability.

If appropriate, other additional components may also be present, for example protective colloids, binders, adhesives, thickeners, thixotropic substances, penetrants, stabilizers, sequestering agents, complex formers. In general, the active ingredients can be combined with any solid or liquid additive commonly used for formulation purposes.

The formulations contain generally between 0.05 and 99% by weight, 0.01 and 98% by weight, preferably between 0.1 and 95% by weight, more preferably between 0.5 and 90% of active ingredient, most preferably between 10 and 70 per cent by weight.

The formulations described above can be used for controlling unwanted microorganisms, in which the compositions comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate are applied to the microorganisms and/or in their habitat.

Seed Treatment

The invention furthermore includes a method for treating seed.

A further aspect of the present invention relates in particular to seeds (dormant, primed, pregerminated or even with emerged roots and leaves) treated with 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate. The inventive seeds are used in methods for protection of seeds and emerged plants from the seeds from phytopathogenic harmful fungi. In these methods, seed treated with at least one inventive active ingredient is used.

The 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate are also suitable for the treatment of seeds and

young seedlings. A large part of the damage to crop plants caused by harmful organisms is triggered by the infection of the seeds before sowing or after germination of the plant. This phase is particularly critical since the roots and shoots of the growing plant are particularly sensitive, and even small damage may result in the death of the plant. Accordingly, there is great interest in protecting the seed and the germinating plant by using appropriate compositions.

It is also desirable to optimize the amount of the active ingredient used so as to provide the best possible protection for the seeds, the germinating plants and emerged seedlings from attack by phytopathogenic fungi, but without damaging the plants themselves by the active ingredient used. In particular, methods for the treatment of seed should also take into consideration the intrinsic phenotypes of transgenic plants in order to achieve optimum protection of the seed and the germinating plant with a minimum of crop protection compositions being employed.

The present invention therefore also relates to a method for protecting seeds, germinating plants and emerged seedlings against attack by animal pests and/or phytopathogenic harmful microorganisms by treating the seeds with an inventive composition. The invention also relates to the use of the compositions according to the invention for treating seeds for protecting the seeds, the germinating plants and emerged seedlings against animal pests and/or phytopathogenic microorganisms. The invention further relates to seeds which has been treated with an inventive composition for protection from animal pests and/or phytopathogenic microorganisms.

One of the advantages of the present invention is that the treatment of the seeds with these compositions not only protects the seed itself, but also the resulting plants after emergence, from animal pests and/or phytopathogenic harmful microorganisms. In this way, the immediate treatment of the crop at the time of sowing or shortly thereafter protect plants as well as seed treatment in prior to sowing. It is likewise considered to be advantageous that the inventive active ingredients or compositions can be used especially also for transgenic seed, in which case the plant which grows from this seed is capable of expressing a protein which acts against pests, herbicidal damage or abiotic stress. The treatment of such seeds with the inventive active ingredients or compositions, for example an insecticidal protein, can result in control of certain pests. Surprisingly, a further synergistic effect can be observed in this case, which additionally increases the effectiveness for protection against attack by pests., microorganisms, weeds or abiotic stress.

The 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate are suitable for protection of seed of any plant variety which is used in agriculture, in the greenhouse, in forests or in horticulture. More particularly, the seed is that of cereals (such as wheat, barley, rye, millet and oats), oilseed rape, maize, cotton, soybean, rice, potatoes, sunflower, beans, coffee, beet (e.g. sugar beet and fodder beet), peanut, vegetables (such as tomato, cucumber, onions and lettuce), lawns and ornamental plants. Of particular significance is the treatment of the seed of wheat, soybean, oilseed rape, maize and rice.

As also described below, the treatment of transgenic seed with the inventive active ingredients or compositions is of particular significance. This refers to the seed of plants containing at least one heterologous gene which allows the expression of a polypeptide or protein, e.g. having insecticidal properties. These heterologous genes in transgenic seeds may originate, for example, from microorganisms of the species *Bacillus*, *Rhizobium*, *Pseudomonas*, *Serratia*, *Trichoderma*, *Clavibacter*, *Glomus* or *Gliocladium*. These heterologous genes preferably originate from *Bacillus* sp., in which case the gene product is effective against the European corn borer and/or the Western corn rootworm. Particularly preferably, the heterologous genes originate from *Bacillus thuringiensis*.

In the context of the present invention, the inventive composition is applied to seeds either alone or in a suitable formulation. Preferably, the seed is treated in a state in which it is sufficiently stable for no damage to occur in the course of treatment. In general, seeds can be treated at any time between harvest and sometime after sowing. It is customary to use seed which has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh of the fruits. For example, it is possible to use seed which has been harvested, cleaned and dried down to a moisture content of less than 15% by weight. Alternatively, it is also possible to use seed which, after drying, for example, has been treated with water and then dried again, or seeds just after priming, or seeds stored in primed conditions or pre-germinated seeds, or seeds sown on nursery trays, tapes or paper.

When treating the seeds, it generally has to be ensured that the amount of the inventive composition applied to the seed and/or the amount of further additives is selected such that the germination of the seed is not impaired, or that the resulting plant is not damaged. This must be ensured particularly in the case of active ingredients which can exhibit phytotoxic effects at certain application rates.

The 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate can be applied directly, i.e. without containing any other components and without having been diluted. In general, it is preferable to apply the compositions to the seed in the form of a suitable formulation. Suitable formulations and methods for seed treatment are known to those skilled in the art. The 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate can be converted to the customary formulations relevant to on-seed applications, such as solutions, emulsions, suspensions, powders, foams, slurries or combined with other coating compositions for seed, such as film forming materials, pelleting materials, fine iron or other metal powders, granules, coating material for inactivated seeds, and also ULV formulations.

These formulations are prepared in a known manner, by mixing the active ingredients or active ingredient combinations with customary additives, for example customary extenders and solvents or diluents, dyes,

wetting agents, dispersants, emulsifiers, antifoams, preservatives, secondary thickeners, adhesives, gibberellins, and also water.

Useful dyes which may be present in the seed dressing formulations usable in accordance with the invention are all dyes which are customary for such purposes. It is possible to use either pigments, which
5 are sparingly soluble in water, or dyes, which are soluble in water. Examples include the dyes known by the names Rhodamine B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

Useful wetting agents which may be present in the seed dressing formulations usable in accordance with the invention are all substances which promote wetting and which are conventionally used for the formulation of active agrochemical ingredients. Usable with preference are alkylnaphthalenesulphonates,
10 such as diisopropyl- or diisobutyl naphthalenesulphonates.

Useful dispersants and/or emulsifiers which may be present in the seed dressing formulations usable in accordance with the invention are all nonionic, anionic and cationic dispersants conventionally used for the formulation of active agrochemical ingredients. Usable with preference are nonionic or anionic dispersants or mixtures of nonionic or anionic dispersants. Useful nonionic dispersants include especially ethylene
15 oxide/propylene oxide block polymers, alkylphenol polyglycol ethers and tristyrylphenol polyglycol ether, and the phosphated or sulphated derivatives thereof. Suitable anionic dispersants are especially lignosulphonates, polyacrylic acid salts and arylsulphonate/formaldehyde condensates.

Antifoams which may be present in the seed dressing formulations usable in accordance with the invention are all foam-inhibiting substances conventionally used for the formulation of active agrochemical
20 ingredients. Silicone antifoams and magnesium stearate can be used with preference.

Preservatives which may be present in the seed dressing formulations usable in accordance with the invention are all substances usable for such purposes in agrochemical compositions. Examples include dichlorophene and benzyl alcohol hemiformal.

Secondary thickeners which may be present in the seed dressing formulations usable in accordance with
25 the invention are all substances usable for such purposes in agrochemical compositions. Preferred examples include cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and finely divided silica.

Adhesives which may be present in the seed dressing formulations usable in accordance with the invention are all customary binders usable in seed dressing products. Preferred examples include
30 polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose.

The formulations for on-seed applications usable in accordance with the invention can be used to treat a wide variety of different kinds of seed either directly or after prior dilution with water. For instance, the

concentrates or the preparations obtainable therefrom by dilution with water can be used to dress the seed of cereals, such as wheat, barley, rye, oats, and triticale, and also seeds of maize, soybean, rice, oilseed rape, peas, beans, cotton, sunflowers, and beets, or else a wide variety of different vegetable seeds. The formulations usable in accordance with the invention, or the dilute preparations thereof, can also be used for seeds of transgenic plants. In this case, additional synergistic effects may also occur in interaction with the substances formed by expression.

For treatment of seeds with the formulations usable in accordance with the invention, or the preparations prepared therefrom by adding water, all mixing units usable customarily for on-seed applications are useful. Specifically, the procedure in on-seed applications is to place the seeds into a mixer, to add the particular desired amount of the formulations, either as such or after prior dilution with water, and to mix everything until all applied formulations are distributed homogeneously on the seeds. If appropriate, this is followed by a drying operation.

The application rate of the formulations usable in accordance with the invention can be varied within a relatively wide range. It is guided by the particular content of the active ingredients in the formulations and by the seeds. The application rates of each single active ingredient is generally between 0.001 and 15 g per kilogram of seed, preferably between 0.01 and 5 g per kilogram of seed.

GMO

As already mentioned above, it is possible to treat all plants and their parts in accordance with the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and also parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering methods, if appropriate in combination with conventional methods (Genetically Modified Organisms), and parts thereof are treated. The terms "parts" or "parts of plants" or "plant parts" have been explained above. More preferably, plants of the plant cultivars which are commercially available or are in use are treated in accordance with the invention. Plant cultivars are understood to mean plants which have new properties ("traits") and have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. They can be cultivars, varieties, bio- or genotypes.

The method of treatment according to the invention can be used in the treatment of genetically modified organisms (GMOs), e.g. plants or seeds. Genetically modified plants (or transgenic plants) are plants of which a heterologous gene has been stably integrated into genome. The expression "heterologous gene" essentially means a gene which is provided or assembled outside the plant and when introduced in the nuclear, chloroplastic or mitochondrial genome gives the transformed plant new or improved agronomic or other properties by expressing a protein or polypeptide of interest or by downregulating or silencing other gene(s) which are present in the plant (using for example, antisense technology, cosuppression technology,

RNA interference – RNAi – technology or microRNA – miRNA - technology). A heterologous gene that is located in the genome is also called a transgene. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

- Plants and plant cultivars which are preferably to be treated according to the invention include all plants
- 5 which have genetic material which impart particularly advantageous, useful traits to these plants (whether obtained by breeding and/or biotechnological means).

Plants and plant cultivars which are also preferably to be treated according to the invention are resistant against one or more biotic stresses, i.e. said plants show a better defense against animal and microbial pests, such as against nematodes, insects, mites, phytopathogenic fungi, bacteria, viruses and/or viroids.

- 10 Plants and plant cultivars which may also be treated according to the invention are those plants which are resistant to one or more abiotic stresses. Abiotic stress conditions may include, for example, drought, cold temperature exposure, heat exposure, osmotic stress, flooding, increased soil salinity, increased mineral exposure, ozone exposure, high light exposure, limited availability of nitrogen nutrients, limited availability of phosphorus nutrients, shade avoidance.
- 15 Plants and plant cultivars which may also be treated according to the invention, are those plants characterized by enhanced yield characteristics. Increased yield in said plants can be the result of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use, enhanced carbon assimilation, improved photosynthesis, increased germination efficiency and accelerated maturation. Yield can furthermore be affected by improved plant architecture (under stress and
- 20 non-stress conditions), including but not limited to, early flowering, flowering control for hybrid seed production, seedling vigor, plant size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as carbohydrate content and composition for example cotton or starch, protein content, oil content and
- 25 composition, nutritional value, reduction in anti-nutritional compounds, improved processability and better storage stability.

Plants that may be treated according to the invention are hybrid plants that already express the characteristic of heterosis or hybrid vigor which results in generally higher yield, vigor, health and resistance towards biotic and abiotic stresses).

- 30 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated according to the invention are herbicide-tolerant plants, i.e. plants made tolerant to one or more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a mutation imparting such herbicide tolerance.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are insect-resistant transgenic plants, i.e. plants made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such insect resistance.

- 5 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are tolerant to abiotic stresses. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance.

- Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention show altered quantity, quality and/or storage-stability of the
10 harvested product and/or altered properties of specific ingredients of the harvested product.

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as cotton plants, with altered fiber characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered fiber characteristics.

- 15 Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered oil profile characteristics.

- Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering)
20 which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered seed shattering characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered seed shattering characteristics and include plants such as oilseed rape plants with delayed or reduced seed shattering.

- Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering)
25 which may also be treated according to the invention are plants, such as Tobacco plants, with altered post-translational protein modification patterns.

Application Rates

- When using the 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate as fungicides, the application rates can
30 be varied within a relatively wide range, depending on the kind of application. The application rate of the inventive active ingredients is

- in the case of treatment of plant parts, for example leaves: from 0.1 to 10 000 g/ha, preferably from 10 to 1000 g/ha, more preferably from 50 to 300 g/ha (in the case of application by watering or dripping, it is even possible to reduce the application rate, especially when inert substrates such as rockwool or perlite are used);
- 5 • in the case of seed treatment: from 0.1 to 400 g per 100 kg of seed, preferably from 1 to 150 g per 100 kg of seed, more preferably from 2.5 to 50 g per 100 kg of seed, even more preferably from 5 to 25 g per 100 kg of seed;
- in the case of soil treatment: from 0.1 to 10 000 g/ha, preferably from 1 to 5000 g/ha.

The following examples are intended to illustrate the present invention.

10

15

20

25

30

35

Examples**Control of corn downy mildew (*Peronosclerospora maydis*)**

This example illustrates the efficacy of compound of formula (I) according to the invention against corn downy mildew (*Peronosclerospora maydis*).

- 5 A typical fungicide formulation containing 50 g of compound of formula (I) per liter was applied at several rates on corn seeds with conventional seed treatment method. Seeds were drilled in seedling trays; then young plants were transplanted to the field after 10 days. Plots received artificial inoculation with a suspension of spores of *Peronosclerospora maydis*. Trial design in the field consisted in randomized complete block with 20 m² plots and 3 replicates.
- 10 Assessments of disease incidence on plants were performed twice in order to evaluate the effect of fungicide treatments, when both crop and disease were well established. Results are presented in tables 1 and 2, below.

Table 1: Efficacy of compound of formula (I) against corn downy mildew (*Peronosclerospora maydis*) as seed treatment. Mean values of 3 replicates (a.i. = active ingredient)

composition	Rate g ai/ kg seeds	% disease incidence (% efficacy Abbott) 31 days after treatment	% disease incidence (% efficacy Abbott) 47 days after treatment
Untreated	-	51,7	66,7
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate EC50	0,1	26.7 (48.4)	33.3 (50.0)
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate EC50	0,5	2.3 (93.5)	16.7 (75.0)
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate EC50	2,5	0 (100)	1.7 (97.5)

15 Means followed by the same letter do not significantly differ ($P=0.05$) according to Duncan's Multiple range test

Table 2: Efficacy of compound of formula (I) against corn downy mildew (*Peronosclerospora maydis*) as seed treatment. Mean values of 3 replicates (a.i. = active ingredient)

composition	Rate g ai/ kg seeds	% disease incidence (% efficacy Abbott) 38 days after treatment	% disease incidence (% efficacy Abbott) 52 days after treatment
Untreated	-	15.8	19.2
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate EC50	0,1	6.7 (57.9)	7.5 (60.9)
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate EC50	0,5	10.8 (31.6)	8.3 (56.5)
2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate EC50	2,5	0 (100)	1.3 (93.5)

Means followed by the same letter do not significantly differ ($P=0.05$) according to Duncan's Multiple range test

Conclusion:

Efficacy results presented in tables 1 and 2, above, demonstrate the efficacy of 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate against corn downy mildew (*Peronosclerospora maydis*) when applied as seed treatment on corn at rates between 0.1 and 2.5 g a.i./kg seeds.

Claims:

1. Use of 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate for controlling downy mildew diseases of monocotyledon crops.
- 5 2. Use according to claim 1 for controlling *Peronosclerospora philippinensis* in corn sorghum or oat.
3. Use according to claim 1 for controlling *Peronosclerospora* in sugarcane.
4. Use according to claim 1 for controlling *Sclerophthora rayssiae varzeae* in corn.
5. Use according to claim 1 for controlling *Sclerophthora macrocarpa* in corn, wheat, barley and rice.
6. Use according to claim 1 for controlling *Sclerospora graminicola* in millets.
- 10 7. Use according to claim 1 for controlling *Peronosclerospora maydis* in corn or teosinte.
8. Use according to claim 1 for controlling *Peronosclerospora sorghi* in sorghum.
9. Use of combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticides for controlling downy mildew diseases of monocotyledon crops.
- 15 10. Use of combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticides for controlling *Peronosclerospora maydis* in corn or teosinte.
- 20 11. Use according to any of claims 1 to 10 for the treatment of seed.
12. Method of controlling downy mildew diseases of monocotyledon crops, characterized in that 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate is applied to the plant pathogen and/or their environment.
- 25 13. Method according to claim 12, characterized that the pathogen is selected from the group *Peronosclerospora philippinensis* in corn sorghum or oat, *Peronosclerospora* in sugarcane, *Sclerophthora rayssiae varzeae* in corn, *Sclerophthora macrocarpa* in corn, wheat, barley and rice, *Sclerospora graminicola* in millets, *Peronosclerospora maydis* in corn or teosinte, *Peronosclerospora sorghi* in sorghum.

14. Method of controlling downy mildew diseases of monocotyledon crops, characterized in that combinations comprising 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate and at least one further fungicide or insecticides is applied to the plant pathogen and/or their environment.
- 5 15. Method according to claim 14, characterized that the pathogen is *Peronosclerospora maydis* in corn or teosinte.
16. Method according to claim 12 or 14, wherein the treated crops are transgenic crops.
17. Method of controlling downy mildew diseases of monocotyledon crops, characterized in that plant seeds are treated with 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate according to
10 Claims 12 to 16.
18. Method according to Claim 17, wherein the plant seeds are derived from a transgenic plant.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2016/059213

A. CLASSIFICATION OF SUBJECT MATTER
INV. A01P3/00 A01N43/56
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2015/055707 A1 (BAYER CROPSOURCE AG [DE]) 23 April 2015 (2015-04-23) page 1, line 4 - line 9 page 5, line 32 - line 35 page 9, line 18 - line 21 page 15, line 36 - page 16, line 4 page 29, line 25 - page 30, line 4 page 31, line 23 - line 36 -----	1-18
X	WO 2015/055574 A1 (BAYER CROPSOURCE AG [DE]) 23 April 2015 (2015-04-23) page 1, line 2 - line 12 page 10, line 23 - page 13, line 15 page 14, line 1 - line 3 examples 1,2 ----- -/-	1-18



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 May 2016

Date of mailing of the international search report

27/05/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Zanobini, Alessandra

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2016/059213

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2014/206896 A1 (BAYER CROPS SCIENCE AG [DE]) 31 December 2014 (2014-12-31) page 1, line 24 - page 9, line 16 page 68 - page 72 page 58, line 22 - page 59, line 8 examples 1, 2 -----	1-18
A	WO 2013/127704 A1 (BAYER IP GMBH [DE]) 6 September 2013 (2013-09-06) page 2, line 7 - page 3, line 27 page 28, line 22 - page 35, line 33 page 42, line 16 - line 22 tables 3-6 -----	1-18
A	CN 104 336 040 A (SHAANXI MEIBANG PESTICIDES CO) 11 February 2015 (2015-02-11) the whole document -----	1-18
A	CN 104 336 048 A (SHAANXI MEIBANG PESTICIDES CO) 11 February 2015 (2015-02-11) the whole document -----	1-18
A	CN 104 336 023 A (SHAANXI MEIBANG PESTICIDES CO) 11 February 2015 (2015-02-11) the whole document -----	1-18
A	RAM P THAKUR ET AL: "Influence of dosage, storage time and temperature on efficacy of metalaxyl-treated seed for the control of pearl millet downy mildew", EUROPEAN JOURNAL OF PLANT PATHOLOGY, KLUWER ACADEMIC PUBLISHERS, DO, vol. 129, no. 2, 9 September 2010 (2010-09-09), pages 353-359, XP019872937, ISSN: 1573-8469, DOI: 10.1007/S10658-010-9679-9 abstract -----	1-18
A	GULYA T J: "EFFICACY OF SINGLE AND TWO-WAY FUNGICIDE SEED TREATMENTS FOR THE CONTROL OF METALAXYL-RESISTANT STRAINS OF PLASMOPARA HALSTEDII (SUNFLOWER DOWNY MILDEW)", BRIGHTON CROP PROTECTION CONFERENCE PESTS AND DISEASES, XX, XX, vol. 2, 1 January 2002 (2002-01-01), pages 575-580, XP009061404, ISSN: 0955-1506 abstract -----	1-18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2016/059213

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2015055707 A1	23-04-2015	CA 2927767 A1 WO 2015055707 A1	23-04-2015 23-04-2015
WO 2015055574 A1	23-04-2015	NONE	
WO 2014206896 A1	31-12-2014	CN 105473585 A EP 3013821 A1 TW 201542544 A WO 2014206896 A1	06-04-2016 04-05-2016 16-11-2015 31-12-2014
WO 2013127704 A1	06-09-2013	AR 090157 A1 CA 2865300 A1 CL 2014002250 A1 CN 104244716 A CO 7061086 A2 EP 2819518 A1 JP 2015511579 A KR 20140129230 A PE 17352014 A1 US 2015327549 A1 WO 2013127704 A1	22-10-2014 06-09-2013 23-01-2015 24-12-2014 19-09-2014 07-01-2015 20-04-2015 06-11-2014 12-11-2014 19-11-2015 06-09-2013
CN 104336040 A	11-02-2015	NONE	
CN 104336048 A	11-02-2015	NONE	
CN 104336023 A	11-02-2015	NONE	