





(86) Date de dépôt PCT/PCT Filing Date: 2014/06/20
(87) Date publication PCT/PCT Publication Date: 2015/01/08
(45) Date de délivrance/Issue Date: 2023/04/18
(85) Entrée phase nationale/National Entry: 2015/12/16
(86) N° demande PCT/PCT Application No.: US 2014/043423
(87) N° publication PCT/PCT Publication No.: 2015/002754
(30) Priorité/Priority: 2013/06/21 (US61/837,841)

(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
A61K 31/395 (2006.01), *A61K 31/4184* (2006.01),
A61K 31/423 (2006.01), *A61K 31/437* (2006.01),
A61K 31/444 (2006.01), *A61K 31/498* (2006.01),
A61K 31/4985 (2006.01), *A61K 31/517* (2006.01),
A61K 31/52 (2006.01), *C07D 403/04* (2006.01),
C07D 413/04 (2006.01), *C07D 413/14* (2006.01),
C07D 473/34 (2006.01)

(72) Inventeurs/Inventors:
DUFFY, BRYAN CORDELL, US;
LIU, SHUANG, US;
QUINN, JOHN FREDERICK, US;
...

(54) Titre : INHIBITEURS DE BROMODOMAINE BICYCLIQUES
(54) Title: BICYCLIC BROMODOMAIN INHIBITORS



(57) Abrégé/Abstract:

The invention provides novel substituted, bicyclic compounds which are useful as inhibitors of extra terminal domain BET protein function by binding to bromodomains, such as compounds of Formula IIb' or Formula IIId', or a stereoisomer, tautomer,

(72) **Inventeurs(suite)/Inventors(continued):** WANG, RUIFANG, US; JIANG, MAY XIAOWU, US;
MARTIN, GREGORY SCOTT, US; WAGNER, GREGORY STEVEN, US; YOUNG, PETER RONALD, US

(73) **Propriétaires(suite)/Owners(continued):**ZENITH EPIGENETICS LTD., CA

(74) **Agent:** TORYS LLP



(57) **Abrégé(suite)/Abstract(continued):**

pharmaceutically acceptable salt, or hydrate thereof. The pharmaceutical compositions comprising such compounds are for the prevention and treatment of diseases and conditions that are associated with bromodomain and extra terminal domain (BET) proteins.

(see formula IIb') (see formula II'd')

Abstract

The invention provides novel substituted, bicyclic compounds which are useful as inhibitors of extra terminal domain BET protein function by binding to bromodomains, such as compounds of Formula IIb' or Formula IIc', or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof. The pharmaceutical compositions comprising such compounds are for the prevention and treatment of diseases and conditions that are associated with bromodomain and extra terminal domain (BET) proteins.



DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 189

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 189

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

Bicyclic Bromodomain Inhibitors

[001] This application claims priority from U.S. Provisional Patent Application No. 61/837,841, filed June 21, 2013 .

[002] The invention provides novel compounds, pharmaceutical compositions containing such compounds, and their use in prevention and treatment of diseases and conditions associated with bromodomain and extra terminal domain (BET) proteins. Post-translational modifications (PTMs) of histones are involved in regulation of gene expression and chromatin organization in eukaryotic cells. Histone acetylation at specific lysine residues is a PTM that is regulated by histone acetylases (HATs) and deacetylases (HDACs). Peserico, A. and C. Simone, "Physical and functional HAT/HDAC interplay regulates protein acetylation balance," *J Biomed Biotechnol*, 2011:371832 (2011). Small molecule inhibitors of HDACs and HATs are being investigated as cancer therapy. Hoshino, I. and H. Matsubara, "Recent advances in histone deacetylase targeted cancer therapy" *Surg Today* 40(9):809-15 (2010); Vernarecci, S., F. Tosi, and P. Filetici, "Tuning acetylated chromatin with HAT inhibitors: a novel tool for therapy" *Epigenetics* 5(2):105-11 (2010); Bandyopadhyay, K., et al., "Spermidinyl-CoA-based HAT inhibitors block DNA repair and provide cancer-specific chemo- and radiosensitization," *Cell Cycle* 8(17):2779-88 (2009); Arif, M., et al., "Protein lysine acetylation in cellular function and its role in cancer manifestation," *Biochim Biophys Acta* 1799(10-12):702-16 (2010). Histone acetylation controls gene expression by recruiting protein complexes that bind directly to acetylated lysine via bromodomains. Sanchez, R. and M.M. Zhou, "The role of human bromodomains in chromatin biology and gene transcription," *Curr Opin Drug Discov Devel* 12(5):659-65 (2009). One such family, the bromodomain and extra terminal domain (BET) proteins, comprises Brd2, Brd3, Brd4, and BrdT, each of which contains two bromodomains in tandem that can independently bind to acetylated lysines, as reviewed in Wu, S.Y. and C.M. Chiang, "The double bromodomain-containing chromatin adaptor Brd4 and transcriptional regulation," *J Biol Chem* 282(18):13141-5 (2007).

[003] Interfering with BET protein interactions via bromodomain inhibition results in modulation of transcriptional programs that are often associated with diseases characterized by dysregulation of cell cycle control, inflammatory cytokine expression, viral transcription, hematopoietic differentiation, insulin transcription, and adipogenesis. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012). BET inhibitors are believed to be useful in the treatment of diseases or conditions related to systemic or tissue inflammation, inflammatory responses to infection or hypoxia, cellular activation and proliferation, lipid metabolism, fibrosis, and the prevention and treatment of viral infections. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat*

Rev Cancer 12(7):465-77 (2012); Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012).

[004] Autoimmune diseases, which are often chronic and debilitating, are a result of a dysregulated immune response, which leads the body to attack its own cells, tissues, and organs. Pro-inflammatory cytokines including IL-1 β , TNF- α , IL-6, MCP-1, and IL-17 are overexpressed in autoimmune disease. IL-17 expression defines the T cell subset known as Th17 cells, which are differentiated, in part, by IL-6, and drive many of the pathogenic consequences of autoimmune disease. Thus, the IL-6/Th17 axis represents an important, potentially druggable target in autoimmune disease therapy. Kimura, A. and T. Kishimoto, "IL-6: regulator of Treg/Th17 balance," *Eur J Immunol* 40(7):1830-5 (2010). BET inhibitors are expected to have anti-inflammatory and immunomodulatory properties. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012); Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012). BET inhibitors have been shown to have a broad spectrum of anti-inflammatory effects *in vitro* including the ability to decrease expression of pro-inflammatory cytokines such as IL-1 β , MCP-1, TNF- α , and IL-6 in activated immune cells. Mirguet, O., et al., "From ApoA1 upregulation to BET family bromodomain inhibition: discovery of I-BET151," *Bioorg Med Chem Lett* 22(8):2963-7 (2012); Nicodeme, E., et al., "Suppression of inflammation by a synthetic histone mimic," *Nature* 468(7327):1119-23 (2010); Seal, J., et al., "Identification of a novel series of BET family bromodomain inhibitors: binding mode and profile of I-BET151 (GSK1210151A)," *Bioorg Med Chem Lett* 22(8):2968-72 (2012). The mechanism for these anti-inflammatory effects may involve BET inhibitor disruption of Brd4 co-activation of NF- κ B-regulated pro-inflammatory cytokines and/or displacement of BET proteins from cytokine promoters, including IL-6. Nicodeme, E., et al., "Suppression of inflammation by a synthetic histone mimic," *Nature* 468(7327):1119-23 (2010); Zhang, G., et al., "Down-regulation of NF-kappaB Transcriptional Activity in HIV-associated Kidney Disease by BRD4 Inhibition," *J Biol Chem*, 287(34):8840-51 (2012); Zhou, M., et al., "Bromodomain protein Brd4 regulates human immunodeficiency virus transcription through phosphorylation of CDK9 at threonine 29," *J Virol* 83(2):1036-44 (2009). In addition, because Brd4 is involved in T-cell lineage differentiation, BET inhibitors may be useful in inflammatory disorders characterized by specific programs of T cell differentiation. Zhang, W.S., et al., "Bromodomain-Containing-Protein 4 (BRD4) Regulates RNA Polymerase II Serine 2 Phosphorylation in Human CD4+ T Cells," *J Biol Chem* (2012).

[005] The anti-inflammatory and immunomodulatory effects of BET inhibition have also been confirmed *in vivo*. A BET inhibitor prevented endotoxin- or bacterial sepsis-induced death and

cecal ligation puncture-induced death in mice, suggesting utility for BET inhibitors in sepsis and acute inflammatory disorders. Nicodeme, E., et al., "Suppression of inflammation by a synthetic histone mimic," *Nature* 468(7327):1119-23 (2010). A BET inhibitor has been shown to ameliorate inflammation and kidney injury in HIV-1 transgenic mice, an animal model for HIV-associated nephropathy, in part through inhibition of Brd4 interaction with NF- κ B. Zhang, G., et al., "Down-regulation of NF-kappaB Transcriptional Activity in HIV-associated Kidney Disease by BRD4 Inhibition," *J Biol Chem*, 287(34):8840-51 (2012). The utility of BET inhibition in autoimmune disease was demonstrated in a mouse model of multiple sclerosis, where BET inhibition resulted in abrogation of clinical signs of disease, in part, through inhibition of IL-6 and IL-17. R. Jahagirdar, S.M. et al., "An Orally Bioavailable Small Molecule RVX-297 Significantly Decreases Disease in a Mouse Model of Multiple Sclerosis," *World Congress of Inflammation*, Paris, France (2011). These results were supported in a similar mouse model where it was shown that treatment with a BET inhibitor inhibited T cell differentiation into pro-autoimmune Th1 and Th17 subsets *in vitro*, and further abrogated disease induction by pro-inflammatory Th1 cells. Bandukwala, H.S., et al., "Selective inhibition of CD4+ T-cell cytokine production and autoimmunity by BET protein and c-Myc inhibitors," *Proc Natl Acad Sci USA*, 109(36):14532-7 (2012).

[006] BET inhibitors may be useful in the treatment of a variety of chronic autoimmune inflammatory conditions. Thus, one aspect of the invention provides compounds, compositions, and methods for treating autoimmune and/or inflammatory diseases by administering one or more compounds of the invention or pharmaceutical compositions comprising one or more of those compounds. Examples of autoimmune and inflammatory diseases, disorders, and syndromes that may be treated using the compounds and methods of the invention include but are not limited to, inflammatory pelvic disease, urethritis, skin sunburn, sinusitis, pneumonitis, encephalitis, meningitis, myocarditis, nephritis (Zhang, G., et al., "Down-regulation of NF-kappaB Transcriptional Activity in HIV-associated Kidney Disease by BRD4 Inhibition," *J Biol Chem*, 287(34):8840-51 (2012)), osteomyelitis, myositis, hepatitis, gastritis, enteritis, dermatitis, gingivitis, appendicitis, pancreatitis, cholecystitis, agammaglobulinemia, psoriasis, allergy, Crohn's disease, irritable bowel syndrome, ulcerative colitis (Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012)), Sjogren's disease, tissue graft rejection, hyperacute rejection of transplanted organs, asthma, allergic rhinitis, chronic obstructive pulmonary disease (COPD), autoimmune polyglandular disease (also known as autoimmune polyglandular syndrome), autoimmune alopecia, pernicious anemia, glomerulonephritis, dermatomyositis, multiple sclerosis (Bandukwala, H.S., et al., "Selective inhibition of CD4+ T-cell cytokine production and autoimmunity by BET protein and c-Myc

inhibitors," *Proc Natl Acad Sci USA*, 109(36):14532-7 (2012)), scleroderma, vasculitis, autoimmune hemolytic and thrombocytopenic states, Goodpasture's syndrome, atherosclerosis, Addison's disease, Parkinson's disease, Alzheimer's disease, Type I diabetes (Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012)), septic shock (Zhang, G., et al., "Down-regulation of NF-kappaB Transcriptional Activity in HIV-associated Kidney Disease by BRD4 Inhibition," *J Biol Chem*, 287(34):8840-51 (2012)), systemic lupus erythematosus (SLE) (Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012)), rheumatoid arthritis (Denis, G.V., "Bromodomain coactivators in cancer, obesity, type 2 diabetes, and inflammation," *Discov Med* 10(55):489-99 (2010)), psoriatic arthritis, juvenile arthritis, osteoarthritis, chronic idiopathic thrombocytopenic purpura, Waldenstrom macroglobulinemia, myasthenia gravis, Hashimoto's thyroiditis, atopic dermatitis, degenerative joint disease, vitiligo, autoimmune hypopituitarism, Guillain-Barre syndrome, Behcet's disease, uveitis, dry eye disease, scleroderma, mycosis fungoides, and Graves' disease.

[007] BET inhibitors may be useful in the treatment of a wide variety of acute inflammatory conditions. Thus, one aspect of the invention provides compounds, compositions, and methods for treating inflammatory conditions including but not limited to, acute gout, giant cell arteritis, nephritis including lupus nephritis, vasculitis with organ involvement, such as glomerulonephritis, vasculitis, including giant cell arteritis, Wegener's granulomatosis, polyarteritis nodosa, Behcet's disease, Kawasaki disease, and Takayasu's arteritis.

[008] BET inhibitors may be useful in the prevention and treatment of diseases or conditions that involve inflammatory responses to infections with bacteria, viruses, fungi, parasites, and their toxins, such as, but not limited to sepsis, sepsis syndrome, septic shock (Nicodeme, E., et al., "Suppression of inflammation by a synthetic histone mimic," *Nature* 468(7327):1119-23 (2010)), systemic inflammatory response syndrome (SIRS), multi-organ dysfunction syndrome, toxic shock syndrome, acute lung injury, adult respiratory distress syndrome (ARDS), acute renal failure, fulminant hepatitis, burns, post-surgical syndromes, sarcoidosis, Herxheimer reactions, encephalitis, myelitis, meningitis, malaria, and SIRS associated with viral infections, such as influenza, herpes zoster, herpes simplex, and coronavirus. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012). Thus, one aspect of the invention provides compounds, compositions, and methods for treating these inflammatory responses to infections with bacteria, viruses, fungi, parasites, and their toxins described herein.

[009] Cancer is a group of diseases caused by dysregulated cell proliferation. Therapeutic approaches aim to decrease the numbers of cancer cells by inhibiting cell replication or by inducing

cancer cell differentiation or death, but there is still significant unmet medical need for more efficacious therapeutic agents. Cancer cells accumulate genetic and epigenetic changes that alter cell growth and metabolism, promoting cell proliferation and increasing resistance to programmed cell death, or apoptosis. Some of these changes include inactivation of tumor suppressor genes, activation of oncogenes, and modifications of the regulation of chromatin structure, including deregulation of histone PTMs. Watson, J.D., "Curing 'incurable' cancer," *Cancer Discov* 1(6):477-80 (2011); Morin, R.D., et al., "Frequent mutation of histone-modifying genes in non-Hodgkin lymphoma" *Nature* 476(7360):298-303 (2011).

[010] One aspect of the invention provides compounds, compositions, and methods for treating human cancer, including, but not limited to, cancers that result from aberrant translocation or overexpression of BET proteins (e.g., NUT midline carcinoma (NMC) (French, C.A., "NUT midline carcinoma," *Cancer Genet Cytogenet* 203(1):16-20 (2010) and B-cell lymphoma (Greenwald, R.J., et al., "E mu-BRD2 transgenic mice develop B-cell lymphoma and leukemia," *Blood* 103(4):1475-84 (2004)). NMC tumor cell growth is driven by a translocation of the Brd4 or Brd3 gene to the nutlin 1 gene. Filippakopoulos, P., et al., "Selective inhibition of BET bromodomains," *Nature* 468(7327):1067-73 (2010). BET inhibition has demonstrated potent antitumor activity in murine xenograft models of NMC, a rare but lethal form of cancer. The present disclosure also provides a method for treating human cancers, including, but not limited to, cancers dependent on a member of the myc family of oncoproteins including c-myc, MYCN, and L-myc. Vita, M. and M. Henriksson, "The Myc oncoprotein as a therapeutic target for human cancer," *Semin Cancer Biol* 16(4):318-30 (2006). These cancers include Burkitt's lymphoma, acute myelogenous leukemia, multiple myeloma, and aggressive human medulloblastoma. Vita, M. and M. Henriksson, "The Myc oncoprotein as a therapeutic target for human cancer," *Semin Cancer Biol* 16(4):318-30 (2006). Cancers in which c-myc is overexpressed may be particularly susceptible to BET protein inhibition; it has been shown that treatment of tumors that have activation of c-myc with a BET inhibitor resulted in tumor regression through inactivation of c-myc transcription. Dawson, M.A., et al., Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia. *Nature*, 2011. 478(7370): p. 529-33; Delmore, J.E., et al., "BET bromodomain inhibition as a therapeutic strategy to target c-Myc," *Cell* 146(6):904-17 (2010); Mertz, J.A., et al., "Targeting MYC dependence in cancer by inhibiting BET bromodomains," *Proc Natl Acad Sci USA* 108(40):16669-74 (2011); Ott, C.J., et al., "BET bromodomain inhibition targets both c-Myc and IL7R in high risk acute lymphoblastic leukemia," *Blood* 120(14):2843-52 (2012); Zuber, J., et al., "RNAi screen identifies Brd4 as a therapeutic target in acute myeloid leukaemia," *Nature* 478(7370):524-8 (2011).

[011] Embodiments of the invention include methods for treating human cancers that rely on BET proteins and pTEFb (Cdk9/CyclinT) to regulate oncogenes (Wang, S. and P.M. Fischer, "Cyclin-dependent kinase 9: a key transcriptional regulator and potential drug target in oncology, virology and cardiology," *Trends Pharmacol Sci* 29(6):302-13 (2008)), and cancers that can be treated by inducing apoptosis or senescence by inhibiting Bcl2, cyclin-dependent kinase 6 (CDK6) (Dawson, M.A., et al., "Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia," *Nature* 478(7370):529-33 (2011)), or human telomerase reverse transcriptase (hTERT). Delmore, J.E., et al., "BET bromodomain inhibition as a therapeutic strategy to target c-Myc," *Cell* 146(6):904-17 (2010); Ruden, M. and N. Puri, "Novel anticancer therapeutics targeting telomerase," *Cancer Treat Rev* (2012).

[012] BET inhibitors may be useful in the treatment of cancers including, but not limited to, adrenal cancer, acinic cell carcinoma, acoustic neuroma, acral lentiginous melanoma, acrospiroma, acute eosinophilic leukemia, acute erythroid leukemia, acute lymphoblastic leukemia, acute megakaryoblastic leukemia, acute monocytic leukemia, acute myeloid leukemia (Dawson, M.A., et al., "Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia," *Nature* 478(7370):529-33 (2011); Mertz, J.A., et al., "Targeting MYC dependence in cancer by inhibiting BET bromodomains," *Proc Natl Acad Sci USA* 108(40):16669-74 (2011); Zuber, J., et al., "RNAi screen identifies Brd4 as a therapeutic target in acute myeloid leukaemia," *Nature* 478(7370):524-8 (2011)), adenocarcinoma, adenoid cystic carcinoma, adenoma, adenomatoid odontogenic tumor, adenosquamous carcinoma, adipose tissue neoplasm, adrenocortical carcinoma, adult T-cell leukemia/lymphoma, aggressive NK-cell leukemia, AIDS-related lymphoma, alveolar rhabdomyosarcoma, alveolar soft part sarcoma, ameloblastic fibroma, anaplastic large cell lymphoma, anaplastic thyroid cancer, angioimmunoblastic T-cell lymphoma, angiomyolipoma, angiosarcoma, astrocytoma, atypical teratoid rhabdoid tumor, B-cell acute lymphoblastic leukemia (Ott, C.J., et al., "BET bromodomain inhibition targets both c-Myc and IL7R in highrisk acute lymphoblastic leukemia," *Blood* 120(14):2843-52 (2012)), B-cell chronic lymphocytic leukemia, B-cell prolymphocytic leukemia, B-cell lymphoma (Greenwald, R.J., et al., "E mu-BRD2 transgenic mice develop B-cell lymphoma and leukemia," *Blood* 103(4):1475-84 (2004)), basal cell carcinoma, biliary tract cancer, bladder cancer, blastoma, bone cancer, Brenner tumor, Brown tumor, Burkitt's lymphoma (Mertz, J.A., et al., "Targeting MYC dependence in cancer by inhibiting BET bromodomains," *Proc Natl Acad Sci USA* 108(40):16669-74 (2011)), breast cancer, brain cancer, carcinoma, carcinoma in situ, carcinosarcoma, cartilage tumor, cementoma, myeloid sarcoma, chondroma, chordoma, choriocarcinoma, choroid plexus papilloma, clear-cell sarcoma of the kidney, craniopharyngioma, cutaneous T-cell lymphoma, cervical cancer, colorectal

cancer, Degos disease, desmoplastic small round cell tumor, diffuse large B-cell lymphoma, dysembryoplastic neuroepithelial tumor, dysgerminoma, embryonal carcinoma, endocrine gland neoplasm, endodermal sinus tumor, enteropathy-associated T-cell lymphoma, esophageal cancer, fetus in fetu, fibroma, fibrosarcoma, follicular lymphoma, follicular thyroid cancer, ganglioneuroma, gastrointestinal cancer, germ cell tumor, gestational choriocarcinoma, giant cell fibroblastoma, giant cell tumor of the bone, glial tumor, glioblastoma multiforme, glioma, gliomatosis cerebri, glucagonoma, gonadoblastoma, granulosa cell tumor, gynandroblastoma, gallbladder cancer, gastric cancer, hairy cell leukemia, hemangioblastoma, head and neck cancer, hemangiopericytoma, hematological malignancy, hepatoblastoma, hepatosplenic T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, invasive lobular carcinoma, intestinal cancer, kidney cancer, laryngeal cancer, lentigo maligna, lethal midline carcinoma, leukemia, Leydig cell tumor, liposarcoma, lung cancer, lymphangioma, lymphangiosarcoma, lymphoepithelioma, lymphoma, acute lymphocytic leukemia, acute myelogenous leukemia (Mertz, J.A., et al., "Targeting MYC dependence in cancer by inhibiting BET bromodomains," *Proc Natl Acad Sci USA* 108(40):16669-74 (2011)), chronic lymphocytic leukemia, liver cancer, small cell lung cancer, non-small cell lung cancer, MALT lymphoma, malignant fibrous histiocytoma, malignant peripheral nerve sheath tumor, malignant triton tumor, mantle cell lymphoma, marginal zone B-cell lymphoma, mast cell leukemia, mediastinal germ cell tumor, medullary carcinoma of the breast, medullary thyroid cancer, medulloblastoma, melanoma (Miguel F. Segura, et al., "BRD4 is a novel therapeutic target in melanoma," *Cancer Research*. 72(8):Supplement 1 (2012)), meningioma, Merkel cell cancer, mesothelioma, metastatic urothelial carcinoma, mixed Mullerian tumor, mixed lineage leukemia (Dawson, M.A., et al., "Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia," *Nature* 478(7370):529-33 (2011)), mucinous tumor, multiple myeloma (Delmore, J.E., et al., "BET bromodomain inhibition as a therapeutic strategy to target c-Myc," *Cell* 146(6):904-17 (2010)), muscle tissue neoplasm, mycosis fungoides, myxoid liposarcoma, myxoma, myxosarcoma, nasopharyngeal carcinoma, neurinoma, neuroblastoma, neurofibroma, neuroma, nodular melanoma, NUT-midline carcinoma (Filippakopoulos, P., et al., "Selective inhibition of BET bromodomains," *Nature* 468(7327):1067-73 (2010)), ocular cancer, oligoastrocytoma, oligodendroglioma, oncocytoma, optic nerve sheath meningioma, optic nerve tumor, oral cancer, osteosarcoma, ovarian cancer, Pancoast tumor, papillary thyroid cancer, paraganglioma, pinealoblastoma, pineocytoma, pituicytoma, pituitary adenoma, pituitary tumor, plasmacytoma, polyembryoma, precursor T-lymphoblastic lymphoma, primary central nervous system lymphoma, primary effusion lymphoma, primary peritoneal cancer, prostate cancer, pancreatic

cancer, pharyngeal cancer, pseudomyxoma peritonei, renal cell carcinoma, renal medullary carcinoma, retinoblastoma, rhabdomyoma, rhabdomyosarcoma, Richter's transformation, rectal cancer, sarcoma, Schwannomatosis, seminoma, Sertoli cell tumor, sex cord-gonadal stromal tumor, signet ring cell carcinoma, skin cancer, small blue round cell tumors, small cell carcinoma, soft tissue sarcoma, somatostatinoma, soot wart, spinal tumor, splenic marginal zone lymphoma, squamous cell carcinoma, synovial sarcoma, Sezary's disease, small intestine cancer, squamous carcinoma, stomach cancer, testicular cancer, thecoma, thyroid cancer, transitional cell carcinoma, throat cancer, urachal cancer, urogenital cancer, urothelial carcinoma, uveal melanoma, uterine cancer, verrucous carcinoma, visual pathway glioma, vulvar cancer, vaginal cancer, Waldenstrom's macroglobulinemia, Warthin's tumor, and Wilms' tumor. Thus, one aspect of the inventions provides compounds, compositions, and methods for treating such cancers.

[013] BET inhibitors may be useful in the treatment of benign proliferative and fibrotic disorders, including benign soft tissue tumors, bone tumors, brain and spinal tumors, eyelid and orbital tumors, granuloma, lipoma, meningioma, multiple endocrine neoplasia, nasal polyps, pituitary tumors, prolactinoma, pseudotumor cerebri, seborrheic keratoses, stomach polyps, thyroid nodules, cystic neoplasms of the pancreas, hemangiomas, vocal cord nodules, polyps, and cysts, Castleman disease, chronic pilonidal disease, dermatofibroma, pilar cyst, pyogenic granuloma, juvenile polyposis syndrome, idiopathic pulmonary fibrosis, renal fibrosis, post-operative stricture, keloid formation, scleroderma, and cardiac fibrosis. See e.g., Tang, X et al., "Assessment of Brd4 Inhibition in Idiopathic Pulmonary Fibrosis Lung Fibroblasts and in Vivo Models of Lung Fibrosis," *Am J Pathology* in press (2013). Thus, one aspect of the invention provides compounds, compositions, and methods for treating such benign proliferative and fibrotic disorders.

[014] Cardiovascular disease (CVD) is the leading cause of mortality and morbidity in the United States. Roger, V.L., et al., "Heart disease and stroke statistics--2012 update: a report from the American Heart Association," *Circulation* 125(1):e2-e220 (2012). Atherosclerosis, an underlying cause of CVD, is a multifactorial disease characterized by dyslipidemia and inflammation. BET inhibitors are expected to be efficacious in atherosclerosis and associated conditions because of aforementioned anti-inflammatory effects as well as ability to increase transcription of ApoA-I, the major constituent of HDL. Mirguet, O., et al., "From ApoA1 upregulation to BET family bromodomain inhibition: discovery of I-BET151," *Bioorg Med Chem Lett* 22(8):2963-7 (2012); Chung, C.W., et al., "Discovery and characterization of small molecule inhibitors of the BET family bromodomains," *J Med Chem* 54(11):3827-38 (2011). Accordingly, one aspect of the invention

provides compounds, compositions, and methods for treating cardiovascular disease, including but not limited to atherosclerosis.

[015] Up-regulation of ApoA-I is considered to be a useful strategy in treatment of atherosclerosis and CVD. Degoma, E.M. and D.J. Rader, "Novel HDL-directed pharmacotherapeutic strategies," *Nat Rev Cardiol* 8(5):266-77 (2011). BET inhibitors have been shown to increase ApoA-I transcription and protein expression. Mirguet, O., et al., "From ApoA1 upregulation to BET family bromodomain inhibition: discovery of I-BET151," *Bioorg Med Chem Lett* 22(8):2963-7 (2012); Chung, C.W., et al., "Discovery and characterization of small molecule inhibitors of the BET family bromodomains," *J Med Chem* 54(11):3827-38 (2011). It has also been shown that BET inhibitors bind directly to BET proteins and inhibit their binding to acetylated histones at the ApoA-1 promoter, suggesting the presence of a BET protein repression complex on the ApoA-1 promoter, which can be functionally disrupted by BET inhibitors. It follows that, BET inhibitors may be useful in the treatment of disorders of lipid metabolism via the regulation of ApoA-I and HDL such as hypercholesterolemia, dyslipidemia, atherosclerosis (Degoma, E.M. and D.J. Rader, "Novel HDL-directed pharmacotherapeutic strategies," *Nat Rev Cardiol* 8(5):266-77 (2011)), and Alzheimer's disease and other neurological disorders. Elliott, D.A., et al., "Apolipoproteins in the brain: implications for neurological and psychiatric disorders," *Clin Lipidol* 51(4):555-573 (2010). Thus, one aspect of the invention provides compounds, compositions, and methods for treating cardiovascular disorders by upregulation of ApoA-1.

[016] BET inhibitors may be useful in the prevention and treatment of conditions associated with ischemia-reperfusion injury such as, but not limited to, myocardial infarction, stroke, acute coronary syndromes (Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012)), renal reperfusion injury, organ transplantation, coronary artery bypass grafting, cardio-pulmonary bypass procedures, hypertension, pulmonary, renal, hepatic, gastro-intestinal, or peripheral limb embolism. Accordingly, one aspect of the invention provides compounds, compositions, and methods for prevention and treatment of conditions described herein that are associated with ischemia-reperfusion injury.

[017] Obesity-associated inflammation is a hallmark of type II diabetes, insulin resistance, and other metabolic disorders. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012); Denis, G.V., "Bromodomain coactivators in cancer, obesity, type 2 diabetes, and inflammation," *Discov Med* 10(55):489-99 (2010). Consistent with the ability of BET inhibitors to inhibit inflammation, gene disruption of Brd2 in mice ablates inflammation and protects animals from obesity-induced insulin

resistance. Wang, F., et al., "Brd2 disruption in mice causes severe obesity without Type 2 diabetes," *Biochem J* 425(1):71-83 (2010). It has been shown that Brd2 interacts with PPAR γ and opposes its transcriptional function. Knockdown of Brd2 *in vitro* promotes transcription of PPAR γ -regulated networks, including those controlling adipogenesis. Denis, G.V., et al., "An emerging role for bromodomain-containing proteins in chromatin regulation and transcriptional control of adipogenesis," *FEBS Lett* 584(15):3260-8 (2010). In addition Brd2 is highly expressed in pancreatic β -cells and regulates proliferation and insulin transcription. Wang, F., et al., "Brd2 disruption in mice causes severe obesity without Type 2 diabetes," *Biochem J* 425(1):71-83 (2010). Taken together, the combined effects of BET inhibitors on inflammation and metabolism decrease insulin resistance and may be useful in the treatment of pre-diabetic and type II diabetic individuals as well as patients with other metabolic complications. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012). Accordingly, one aspect of the invention provides compounds, compositions, and methods for treatment and prevention of metabolic disorders, including but not limited to obesity-associated inflammation, type II diabetes, and insulin resistance.

[018] Host-encoded BET proteins have been shown to be important for transcriptional activation and repression of viral promoters. Brd4 interacts with the E2 protein of human papilloma virus (HPV) to enable E2 mediated transcription of E2-target genes. Gagnon, D., et al., "Proteasomal degradation of the papillomavirus E2 protein is inhibited by overexpression of bromodomain-containing protein 4," *J Virol* 83(9):4127-39 (2009). Similarly, Brd2, Brd3, and Brd4 all bind to latent nuclear antigen 1 (LANA1), encoded by Kaposi's sarcoma-associated herpes virus (KSHV), promoting LANA1-dependent proliferation of KSHV-infected cells. You, J., et al., "Kaposi's sarcoma-associated herpesvirus latency-associated nuclear antigen interacts with bromodomain protein Brd4 on host mitotic chromosomes," *J Virol* 80(18):8909-19 (2006). A BET inhibitor has been shown to inhibit the Brd4-mediated recruitment of the transcription elongation complex pTEFb to the Epstein-Barr virus (EBV) viral C promoter, suggesting therapeutic value for EBV-associated malignancies. Palermo, R.D., et al., "RNA polymerase II stalling promotes nucleosome occlusion and pTEFb recruitment to drive immortalization by Epstein-Barr virus," *PLoS Pathog* 7(10):e1002334 (2011). Also, a BET inhibitor reactivated HIV in models of latent T cell infection and latent monocyte infection, potentially allowing for viral eradication by complementary anti-retroviral therapy. Zhu, J., et al., "Reactivation of Latent HIV-1 by Inhibition of BRD4," *Cell Rep* (2012); Banerjee, C., et al., "BET bromodomain inhibition as a novel strategy for reactivation of HIV-1," *J Leukoc Biol* (2012); Bartholomeeusen, K., et al., "BET bromodomain inhibition activates transcription via a transient release of P-TEFb from 7SK

snRNP," *J Biol Chem* (2012); Li, Z., et al., "The BET bromodomain inhibitor JQ1 activates HIV latency through antagonizing Brd4 inhibition of Tat-transactivation," *Nucleic Acids Res* (2012).

[019] BET inhibitors may be useful in the prevention and treatment of episome-based DNA viruses including, but not limited to, human papillomavirus, herpes virus, Epstein-Barr virus, human immunodeficiency virus (Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012)), adenovirus, poxvirus, hepatitis B virus, and hepatitis C virus. Thus, the invention also provides compounds, compositions, and methods for treatment and prevention of episome-based DNA virus infections described herein.

[020] Some central nervous system (CNS) diseases are characterized by disorders in epigenetic processes. Brd2 haplo-insufficiency has been linked to neuronal deficits and epilepsy. Velisek, L., et al., "GABAergic neuron deficit as an idiopathic generalized epilepsy mechanism: the role of BRD2 haploinsufficiency in juvenile myoclonic epilepsy," *PLoS One* 6(8): e23656 (2011) SNPs in various bromodomain-containing proteins have also been linked to mental disorders including schizophrenia and bipolar disorders. Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012). In addition, the ability of BET inhibitors to increase ApoA-I transcription may make BET inhibitors useful in Alzheimer's disease therapy considering the suggested relationship between increased ApoA-I and Alzheimer's disease and other neurological disorders. Elliott, D.A., et al., "Apolipoproteins in the brain: implications for neurological and psychiatric disorders," *Clin Lipidol* 51(4):555-573 (2010). Accordingly, one aspect of the invention provides compounds, compositions, and methods for treating such CNS diseases and disorders.

[021] BRDT is the testis-specific member of the BET protein family which is essential for chromatin remodeling during spermatogenesis. Gaucher, J., et al., "Bromodomain-dependent stage-specific male genome programming by Brdt," *EMBO J* 31(19):3809-20 (2012); Shang, E., et al., "The first bromodomain of Brdt, a testis-specific member of the BET sub-family of double-bromodomain-containing proteins, is essential for male germ cell differentiation," *Development* 134(19):3507-15 (2007). Genetic depletion of BRDT or inhibition of BRDT interaction with acetylated histones by a BET inhibitor resulted in a contraceptive effect in mice, which was reversible when small molecule BET inhibitors were used. Matzuk, M.M., et al., "Small-Molecule Inhibition of BRDT for Male Contraception," *Cell* 150(4): 673-684 (2012); Berkovits, B.D., et al., "The testis-specific double bromodomain-containing protein BRDT forms a complex with multiple spliceosome components and is required for mRNA splicing and 3'-UTR truncation in round spermatids," *Nucleic Acids Res* 40(15):7162-75 (2012). These data suggest potential utility of BET inhibitors as a novel and

efficacious approach to male contraception. Thus, another aspect of the invention provides compounds, compositions, and methods for male contraception.

[022] Monocyte chemoattractant protein-1 (MCP-1, CCL2) plays an important role in cardiovascular disease. Niu, J. and P.E. Kolattukudy, "Role of MCP-1 in cardiovascular disease: molecular mechanisms and clinical implications," *Clin Sci (Lond)* 117(3):95-109 (2009). MCP-1, by its chemotactic activity, regulates recruitment of monocytes from the arterial lumen to the subendothelial space, where they develop into macrophage foam cells, and initiate the formation of fatty streaks which can develop into atherosclerotic plaque. Dawson, J., et al., "Targeting monocyte chemoattractant protein-1 signalling in disease," *Expert Opin Ther Targets* 7(1):35-48 (2003). The critical role of MCP-1 (and its cognate receptor CCR2) in the development of atherosclerosis has been examined in various transgenic and knockout mouse models on a hyperlipidemic background. Boring, L., et al., "Decreased lesion formation in CCR2^{-/-} mice reveals a role for chemokines in the initiation of atherosclerosis," *Nature* 394(6696):894-7 (1998); Gosling, J., et al., "MCP-1 deficiency reduces susceptibility to atherosclerosis in mice that overexpress human apolipoprotein B," *J Clin Invest* 103(6):773-8 (1999); Gu, L., et al., "Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low density lipoprotein receptor-deficient mice," *Mol Cell* 2(2):275-81 (1998); Aiello, R.J., et al., "Monocyte chemoattractant protein-1 accelerates atherosclerosis in apolipoprotein E-deficient mice," *Arterioscler Thromb Vasc Biol* 19(6):1518-25 (1999). These reports demonstrate that abrogation of MCP-1 signaling results in decreased macrophage infiltration to the arterial wall and decreased atherosclerotic lesion development.

[023] The association between MCP-1 and cardiovascular disease in humans is well-established. Niu, J. and P.E. Kolattukudy, "Role of MCP-1 in cardiovascular disease: molecular mechanisms and clinical implications," *Clin Sci (Lond)* 117(3):95-109 (2009). MCP-1 and its receptor are overexpressed by endothelial cells, smooth muscle cells, and infiltrating monocytes/macrophages in human atherosclerotic plaque. Nelken, N.A., et al., "Monocyte chemoattractant protein-1 in human atheromatous plaques," *J Clin Invest* 88(4):1121-7 (1991). Moreover, elevated circulating levels of MCP-1 are positively correlated with most cardiovascular risk factors, measures of coronary atherosclerosis burden, and the incidence of coronary heart disease (CHD). Deo, R., et al., "Association among plasma levels of monocyte chemoattractant protein-1, traditional cardiovascular risk factors, and subclinical atherosclerosis," *J Am Coll Cardiol* 44(9):1812-8 (2004). CHD patients with among the highest levels of MCP-1 are those with acute coronary syndrome (ACS). de Lemos, J.A., et al., "Association between plasma levels of monocyte chemoattractant protein-1 and long-term clinical outcomes in patients with acute coronary syndromes," *Circulation* 107(5):690-5 (2003). In addition to playing a role in the underlying

inflammation associated with CHD, MCP-1 has been shown to be involved in plaque rupture, ischemic/reperfusion injury, restenosis, and heart transplant rejection. Niu, J. and P.E. Kolattukudy, "Role of MCP-1 in cardiovascular disease: molecular mechanisms and clinical implications," *Clin Sci (Lond)* 117(3):95-109 (2009).


[024] MCP-1 also promotes tissue inflammation associated with autoimmune diseases including rheumatoid arthritis (RA) and multiple sclerosis (MS). MCP-1 plays a role in the infiltration of macrophages and lymphocytes into the joint in RA, and is overexpressed in the synovial fluid of RA patients. Koch, A.E., et al., "Enhanced production of monocyte chemoattractant protein-1 in rheumatoid arthritis," *J Clin Invest* 90(3):772-9 (1992). Blockade of MCP-1 and MCP-1 signaling in animal models of RA have also shown the importance of MCP-1 to macrophage accumulation and proinflammatory cytokine expression associated with RA. Brodmerkel, C.M., et al., "Discovery and pharmacological characterization of a novel rodent-active CCR2 antagonist, INCB3344," *J Immunol* 175(8):5370-8 (2005); Bruhl, H., et al., "Dual role of CCR2 during initiation and progression of collagen-induced arthritis: evidence for regulatory activity of CCR2+ T cells," *J Immunol* 172(2):890-8 (2004); Gong, J.H., et al., "An antagonist of monocyte chemoattractant protein 1 (MCP-1) inhibits arthritis in the MRL-lpr mouse model," *J Exp Med* 186(1):131-7 (1997); 65. Gong, J.H., et al., "Post-onset inhibition of murine arthritis using combined chemokine antagonist therapy," *Rheumatology (Oxford)* 43(1): 39-42 (2004).

[025] Overexpression of MCP-1, in the brain, cerebrospinal fluid (CSF), and blood, has also been associated with chronic and acute MS in humans. Mahad, D.J. and R.M. Ransohoff, "The role of MCP-1 (CCL2) and CCR2 in multiple sclerosis and experimental autoimmune encephalomyelitis (EAE)," *Semin Immunol* 15(1):23-32 (2003). MCP-1 is overexpressed by a variety of cell types in the brain during disease progression and contributes to the infiltration of macrophages and lymphocytes which mediate the tissue damage associated with MS. Genetic depletion of MCP-1 or CCR2 in the experimental autoimmune encephalomyelitis (EAE) mouse model, a model resembling human MS, results in resistance to disease, primarily because of decreased macrophage infiltration to the CNS. Fife, B.T., et al., "CC chemokine receptor 2 is critical for induction of experimental autoimmune encephalomyelitis," *J Exp Med* 192(6):899-905 (2000); Huang, D.R., et al., "Absence of monocyte chemoattractant protein 1 in mice leads to decreased local macrophage recruitment and antigen-specific T helper cell type 1 immune response in experimental autoimmune encephalomyelitis," *J Exp Med* 193(6):713-26 (2001).


[026] Preclinical data have suggested that small- and large-molecule inhibitors of MCP-1 and CCR2 have potential as therapeutic agents in inflammatory and autoimmune indications. Thus,

one aspect of the invention provides compounds, compositions, and methods for treating cardiovascular, inflammatory, and autoimmune conditions associated with MCP-1 and CCR2.

[027] Accordingly, the invention provides compounds that are useful for inhibition of BET protein function by binding to bromodomains, pharmaceutical compositions comprising one or more of those compounds, and use of these compounds or compositions in the treatment and prevention of diseases and conditions, including, but not limited to, cancer, autoimmune, and cardiovascular diseases. The compounds of the invention are defined by Formula Ia or Formula IIa:




Formula Ia



Formula IIa

or are stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, wherein:

A is selected from 5- or 6-membered monocyclic heterocycles fused to ring B;



B is a six-membered aromatic carbocycle or heterocycle;

Y is selected from N, C, and CH;

W_1 is selected from N and CR_1 ;

W_2 is selected from N and CR_2 ;

W_3 is selected from N and CR_3 ;

W₄ and **W₅** are independently selected from N, CH, and C or alternatively, **W₄** and **W₅** are both C (see, e.g., Formula Ia and Formula Ib below);

W_1 , W_2 , and W_3 may be the same or different from each other;

R_1 and R_2 are independently selected from hydrogen, deuterium, alkyl, -OH, -NH₂, -thioalkyl, alkoxy, ketone, ester, carboxylic acid, urea, carbamate, amino, amide, halogen, carbocycle, heterocycle, sulfone, sulfoxide, sulfide, sulfonamide, and -CN;



R_3 is selected from hydrogen, -NH₂, -CN, -N₃, halogen, and deuterium; or alternatively, R_3 is selected from -NO₂, -OMe, -OEt, -NHC(O)Me, NHSO₂Me, cycloamino, cycloamido, -OH, -SO₂Me, -SO₂Et, -CH₂NH₂, -C(O)NH₂, and -C(O)OMe;

X is selected from -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂O-, -CH₂CH₂NH-, -CH₂CH₂S-, -C(O)-, -C(O)CH₂-, -C(O)CH₂CH₂-, -CH₂C(O)-, -CH₂CH₂C(O)-, -C(O)NH-, -C(O)O-, -C(O)S-, -C(O)NHCH₂-, -C(O)OCH₂-, -C(O)SCH₂-, where one or more hydrogen may independently be replaced with deuterium, halogen, -CF₃, ketone, and where S may be oxidized to sulfoxide or sulfone; or alternatively, X may be selected from -NH-, -CH(OH)-, -CH(CH₃)-, and hydroxyl methyl, where one or more hydrogen may independently be replaced with deuterium, halogen, -CF₃, ketone, and where S may be oxidized to sulfoxide or sulfone;

R_4 is selected from 4-7 membered carbocycles and heterocycles; or alternatively, R_4 is a 3 membered carbocycle or heterocycle;

D_1 is selected from 5-membered monocyclic carbocycles and heterocycles; or alternatively, D_1 is a monocyclic heterocycle, where D_1 is attached to the B ring via a carbon atom that is part of a double bond;

with the proviso that if R_3 is hydrogen and A is a 5-membered ring, then D_1 cannot be




and with the proviso that if D_1 is

and R_2 and R_3 are hydrogen and R_1 is -OMe, then




the A-B bicyclic ring is not



and with the proviso that if D_1 is


and each of R_1 , R_2 , R_3 , are hydrogen, then the

A-B bicyclic ring is not




unless the B ring is substituted;


and with the proviso that if each of R_1 , R_2 , R_3 are hydrogen, then the A-B bicyclic ring is not




and with the proviso that if each of R_1 , R_2 , R_3 are hydrogen, then the A-B bicyclic ring is not



[028] In certain embodiments A is a five membered ring. In some embodiments Y is N or C. In some embodiments, R_1 and R_2 are independently selected from hydrogen, deuterium, alkyl, -OH, -NH₂, -thioalkyl, alkoxy, ketone, ester, carboxylic acid, urea, carbamate, amino, amide, halogen, sulfone, sulfoxide, sulfide, sulfonamide, and -CN. In some embodiments, the compound of Formula Ia is a compound of Formula Ib, i.e., wherein W_4 and W_5 of Formula I are both C.




Formula Ia




Formula Ib

In some embodiments, the compound of Formula Ia is a compound of Formula Ib, i.e., wherein W_4 and W_5 of Formula I are both C.



Formula IIa



Formula IIb

[029] In another aspect of the invention, a pharmaceutical composition comprising a compound of Formula Ia, Formula Ib, Formula IIa, and/or Formula IIb, or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof and one or more pharmaceutically acceptable carriers, diluents or excipients is provided.

[030] In yet another aspect of the invention there is provided a compound of Formula Ia, Formula Ib, Formula IIa, and/or Formula IIb, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, or a pharmaceutical composition comprising such compound, for use in therapy, in particular in the treatment of diseases or conditions for which a bromodomain inhibitor is indicated.

[031] In yet another aspect of the invention there is provided a compound of Formula Ia, Formula Ib, Formula IIa, and/or Formula IIb, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof in the manufacture of a medicament for the treatment of diseases or conditions for which a bromodomain inhibitor is indicated.

DEFINITIONS

[032] As used in the present specification, the following words, phrases and symbols are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise. The following abbreviations and terms have the indicated meanings throughout.

[033] As used herein, "cardiovascular disease" refers to diseases, disorders and conditions of the heart and circulatory system that are mediated by BET inhibition. Exemplary cardiovascular diseases, including cholesterol- or lipid-related disorders, include, but are not limited to, acute coronary syndrome, angina, arteriosclerosis, atherosclerosis, carotid atherosclerosis, cerebrovascular disease, cerebral infarction, congestive heart failure, congenital heart disease, coronary heart disease, coronary artery disease, coronary plaque stabilization, dyslipidemias, dyslipoproteinemias, endothelium dysfunctions, familial hypercholesterolemia, familial combined

hyperlipidemia, hypoalphalipoproteinemia, hypertriglyceridemia, hyperbetalipoproteinemia, hypercholesterolemia, hypertension, hyperlipidemia, intermittent claudication, ischemia, ischemia reperfusion injury, ischemic heart diseases, cardiac ischemia, metabolic syndrome, multi-infarct dementia, myocardial infarction, obesity, peripheral vascular disease, reperfusion injury, restenosis, renal artery atherosclerosis, rheumatic heart disease, stroke, thrombotic disorder, transitory ischemic attacks, and lipoprotein abnormalities associated with Alzheimer's disease, obesity, diabetes mellitus, syndrome X, impotence, multiple sclerosis, Parkinson's disease, and inflammatory diseases.

[034] As used herein, "inflammatory diseases" refers to diseases, disorders, and conditions that are mediated by BET inhibition. Exemplary inflammatory diseases, include, but are not limited to, arthritis, asthma, dermatitis, psoriasis, cystic fibrosis, post transplantation late and chronic solid organ rejection, multiple sclerosis, systemic lupus erythematosus, inflammatory bowel diseases, autoimmune diabetes, diabetic retinopathy, diabetic nephropathy, diabetic vasculopathy, ocular inflammation, uveitis, rhinitis, ischemia-reperfusion injury, post-angioplasty restenosis, chronic obstructive pulmonary disease (COPD), glomerulonephritis, Graves disease, gastrointestinal allergies, conjunctivitis, atherosclerosis, coronary artery disease, angina, and small artery disease.

[035] As used herein, "cancer" refers to diseases, disorders, and conditions that are mediated by BET inhibition. Exemplary cancers, include, but are not limited to, chronic lymphocytic leukemia and multiple myeloma, follicular lymphoma, diffuse large B cell lymphoma with germinal center phenotype, Burkitt's lymphoma, Hodgkin's lymphoma, follicular lymphomas and activated, anaplastic large cell lymphoma, neuroblastoma and primary neuroectodermal tumor, rhabdomyosarcoma, prostate cancer, breast cancer, NMC (NUT-midline carcinoma), acute myeloid leukemia (AML), acute B lymphoblastic leukemia (B-ALL), Burkitt's Lymphoma, B-cell lymphoma, melanoma, mixed lineage leukemia, multiple myeloma, pro-myelocytic leukemia (PML), non-Hodgkin's lymphoma, neuroblastoma, medulloblastoma, lung carcinoma (NSCLC, SCLC), and colon carcinoma.

[036] "Subject" refers to an animal, such as a mammal, that has been or will be the object of treatment, observation, or experiment. The methods described herein may be useful for both human therapy and veterinary applications. In one embodiment, the subject is a human.

[037] As used herein, "treatment" or "treating" refers to an amelioration of a disease or disorder, or at least one discernible symptom thereof. In another embodiment, "treatment" or "treating" refers to an amelioration of at least one measurable physical parameter, not necessarily discernible by the patient. In yet another embodiment, "treatment" or "treating" refers to inhibiting the progression of a disease or disorder, either physically, e.g., stabilization of a discernible

symptom, physiologically, e.g., stabilization of a physical parameter, or both. In yet another embodiment, "treatment" or "treating" refers to delaying the onset of a disease or disorder. For example, treating a cholesterol disorder may comprise decreasing blood cholesterol levels.

[038] As used herein, "prevention" or "preventing" refers to a reduction of the risk of acquiring a given disease or disorder.

[039] A dash ("-") that is not between two letters or symbols is used to indicate a point of attachment for a substituent. For example, -CONH₂ is attached through the carbon atom.

[040] By "optional" or "optionally" is meant that the subsequently described event or circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" encompasses both "aryl" and "substituted aryl" as defined below. It will be understood by those skilled in the art, with respect to any group containing one or more substituents, that such groups are not intended to introduce any substitution or substitution patterns that are sterically impractical, synthetically non-feasible and/or inherently unstable.

[041] As used herein, the term "hydrate" refers to a crystal form with either a stoichiometric or non-stoichiometric amount of water is incorporated into the crystal structure.

[042] The term "alkenyl" as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon double bond, such as a straight or branched group of 2-8 carbon atoms, referred to herein as (C₂-C₈)alkenyl. Exemplary alkenyl groups include, but are not limited to, vinyl, allyl, butenyl, pentenyl, hexenyl, butadienyl, pentadienyl, hexadienyl, 2-ethylhexenyl, 2-propyl-2-butenyl, and 4-(2-methyl-3-butene)-pentenyl.

[043] The term "alkoxy" as used herein refers to an alkyl group attached to an oxygen (-O-alkyl-). "Alkoxy" groups also include an alkenyl group attached to an oxygen ("alkenyloxy") or an alkynyl group attached to an oxygen ("alkynyloxy") groups. Exemplary alkoxy groups include, but are not limited to, groups with an alkyl, alkenyl or alkynyl group of 1-8 carbon atoms, referred to herein as (C₁-C₈)alkoxy. Exemplary alkoxy groups include, but are not limited to methoxy and ethoxy.

[044] The term "alkyl" as used herein refers to a saturated straight or branched hydrocarbon, such as a straight or branched group of 1-8 carbon atoms, referred to herein as (C₁-C₈)alkyl. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, and octyl.

[045] The term "alkynyl" as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon triple bond, such as a straight or branched group of 2-8 carbon atoms, referred to herein as (C₂-C₈)alkynyl. Exemplary alkynyl groups include, but are not limited to, ethynyl, propynyl, butynyl, pentynyl, hexynyl, methylpropynyl, 4-methyl-1-butynyl, 4-propyl-2-pentynyl, and 4-butyl-2-hexynyl.

[046] The term "amide" as used herein refers to the form -NR_aC(O)(R_b)- or -C(O)NR_bR_c, wherein R_a, R_b and R_c are each independently selected from alkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, haloalkyl, heteroaryl, heterocyclyl, and hydrogen. The amide can be attached to another group through the carbon, the nitrogen, R_b, or R_c. The amide also may be cyclic, for example R_b and R_c may be joined to form a 3- to 8-membered ring, such as 5- or 6-membered ring. The term "amide" encompasses groups such as sulfonamide, urea, ureido, carbamate, carbamic acid, and cyclic versions thereof. The term "amide" also encompasses an amide group attached to a carboxy group, e.g., -amide-COOH or salts such as -amide-COONa, an amino group attached to a carboxy group (e.g., -amino-COOH or salts such as -amino-COONa).

[047] The term "amine" or "amino" as used herein refers to the form -NR_dR_e or -N(R_d)R_e-, where R_d and R_e are independently selected from alkyl, alkenyl, alkynyl, aryl, arylalkyl, carbamate, cycloalkyl, haloalkyl, heteroaryl, heterocycle, and hydrogen. The amino can be attached to the parent molecular group through the nitrogen. The amino also may be cyclic, for example any two of R_d and R_e may be joined together or with the N to form a 3- to 12-membered ring (e.g., morpholino or piperidinyl). The term amino also includes the corresponding quaternary ammonium salt of any amino group. Exemplary amino groups include alkylamino groups, wherein at least one of R_d or R_e is an alkyl group. In some embodiments R_d and R_e each may be optionally substituted with hydroxyl, halogen, alkoxy, ester, or amino.

[048] The term "aryl" as used herein refers to a mono-, bi-, or other multi-carbocyclic, aromatic ring system. The aryl group can optionally be fused to one or more rings selected from aryls, cycloalkyls, and heterocyclyls. The aryl groups of this present disclosure can be substituted with groups selected from alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, ketone, nitro, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide, and thioketone. Exemplary aryl groups include, but are not limited to, phenyl, tolyl, anthracenyl, fluorenyl, indenyl, azulenyl, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl. Exemplary aryl groups also include, but are not limited to a monocyclic aromatic ring system, wherein the ring comprises 6 carbon atoms, referred to herein as "(C₆)aryl."

[049] The term "arylalkyl" as used herein refers to an alkyl group having at least one aryl substituent (e.g., -aryl-alkyl-). Exemplary arylalkyl groups include, but are not limited to, arylalkyls having a monocyclic aromatic ring system, wherein the ring comprises 6 carbon atoms, referred to herein as "(C₆)arylalkyl."

[050] The term "carbamate" as used herein refers to the form -R_gOC(O)N(R_h)-, -R_gOC(O)N(R_h)R_i-, or -OC(O)NR_hR_i, wherein R_g, R_h and R_i are each independently selected from alkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, haloalkyl, heteroaryl, heterocyclyl, and hydrogen. Exemplary carbamates include, but are not limited to, arylcarbamates or heteroaryl carbamates (e.g., wherein at least one of R_g, R_h and R_i are independently selected from aryl or heteroaryl, such as pyridine, pyridazine, pyrimidine, and pyrazine).

[051] The term "carbocycle" as used herein refers to an aryl or cycloalkyl group.

[052] The term "carboxy" as used herein refers to -COOH or its corresponding carboxylate salts (e.g., -COONa). The term carboxy also includes "carboxycarbonyl," e.g. a carboxy group attached to a carbonyl group, e.g., -C(O)-COOH or salts, such as -C(O)-COONa.

[053] The term "cyano" as used herein refers to -CN.

[054] The term "cycloalkoxy" as used herein refers to a cycloalkyl group attached to an oxygen.

[055] The term "cycloalkyl" as used herein refers to a saturated or unsaturated cyclic, bicyclic, or bridged bicyclic hydrocarbon group of 3-12 carbons, or 3-8 carbons, referred to herein as "(C₃-C₈)cycloalkyl," derived from a cycloalkane. Exemplary cycloalkyl groups include, but are not limited to, cyclohexanes, cyclohexenes, cyclopentanes, and cyclopentenes. Cycloalkyl groups may be substituted with alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, ketone, nitro, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide and thioketone. Cycloalkyl groups can be fused to other cycloalkyl saturated or unsaturated, aryl, or heterocyclyl groups.

[056] The term "dicarboxylic acid" as used herein refers to a group containing at least two carboxylic acid groups such as saturated and unsaturated hydrocarbon dicarboxylic acids and salts thereof. Exemplary dicarboxylic acids include alkyl dicarboxylic acids. Dicarboxylic acids may be substituted with alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydrogen, hydroxyl, ketone, nitro, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide and thioketone. Dicarboxylic acids include, but are not limited to succinic acid, glutaric acid, adipic acid, suberic acid, sebacic acid, azelaic acid, maleic acid, phthalic acid, aspartic acid, glutamic acid,

malonic acid, fumaric acid, (+)/(-)-malic acid, (+)/(-) tartaric acid, isophthalic acid, and terephthalic acid. Dicarboxylic acids further include carboxylic acid derivatives thereof, such as anhydrides, imides, hydrazides (for example, succinic anhydride and succinimide).

[057] The term “ester” refers to the structure $-C(O)O-$, $-C(O)O-R_j-$, $-R_kC(O)O-R_j-$, or $-R_kC(O)O-$, where O is not bound to hydrogen, and R_j and R_k can independently be selected from alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, cycloalkyl, ether, haloalkyl, heteroaryl, and heterocyclyl. R_k can be a hydrogen, but R_j cannot be hydrogen. The ester may be cyclic, for example the carbon atom and R_j , the oxygen atom and R_k , or R_j and R_k may be joined to form a 3- to 12-membered ring. Exemplary esters include, but are not limited to, alkyl esters wherein at least one of R_j or R_k is alkyl, such as $-O-C(O)-alkyl$, $-C(O)-O-alkyl-$, and $-alkyl-C(O)-O-alkyl-$. Exemplary esters also include aryl or heteroaryl esters, e.g. wherein at least one of R_j or R_k is a heteroaryl group such as pyridine, pyridazine, pyrimidine and pyrazine, such as a nicotinate ester. Exemplary esters also include reverse esters having the structure $-R_kC(O)O-$, where the oxygen is bound to the parent molecule. Exemplary reverse esters include succinate, D-argininate, L-argininate, L-lysinate and D-lysinate. Esters also include carboxylic acid anhydrides and acid halides.

[058] The terms “halo” or “halogen” as used herein refer to F, Cl, Br, or I.

[059] The term “haloalkyl” as used herein refers to an alkyl group substituted with one or more halogen atoms. “Haloalkyls” also encompass alkenyl or alkynyl groups substituted with one or more halogen atoms.

[060] The term “heteroaryl” as used herein refers to a mono-, bi-, or multi-cyclic, aromatic ring system containing one or more heteroatoms, for example 1-3 heteroatoms, such as nitrogen, oxygen, and sulfur. Heteroaryls can be substituted with one or more substituents including alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, ketone, nitro, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide and thioketone. Heteroaryls can also be fused to non-aromatic rings. Illustrative examples of heteroaryl groups include, but are not limited to, pyridinyl, pyridazinyl, pyrimidyl, pyrazyl, triazinyl, pyrrolyl, pyrazolyl, imidazolyl, (1,2,3)- and (1,2,4)-triazolyl, pyrazinyl, pyrimidyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, furyl, phenyl, isoxazolyl, and oxazolyl. Exemplary heteroaryl groups include, but are not limited to, a monocyclic aromatic ring, wherein the ring comprises 2-5 carbon atoms and 1-3 heteroatoms, referred to herein as “(C₂-C₅)heteroaryl.”

[061] The terms “heterocycle,” “heterocyclyl,” or “heterocyclic” as used herein refer to a saturated or unsaturated 3-, 4-, 5-, 6- or 7-membered ring containing one, two, or three heteroatoms independently selected from nitrogen, oxygen, and sulfur. Heterocycles can be

aromatic (heteroaryls) or non-aromatic. Heterocycles can be substituted with one or more substituents including alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, ketone, nitro, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide and thioketone. Heterocycles also include bicyclic, tricyclic, and tetracyclic groups in which any of the above heterocyclic rings is fused to one or two rings independently selected from aryls, cycloalkyls, and heterocycles. Exemplary heterocycles include acridinyl, benzimidazolyl, benzofuryl, benzothiazolyl, benzothieryl, benzoxazolyl, biotinyl, cinnolyl, dihydrofuryl, dihydroindolyl, dihydropyranyl, dihydrothienyl, dithiazolyl, furyl, homopiperidinyl, imidazolidinyl, imidazolyl, imidazolyl, indolyl, isoquinolyl, isothiazolidinyl, isothiazolyl, isoxazolidinyl, isoxazolyl, morpholinyl, oxadiazolyl, oxazolidinyl, oxazolyl, piperazinyl, piperidinyl, pyranyl, pyrazolidinyl, pyrazinyl, pyrazolyl, pyrazolinyl, pyridazinyl, pyridyl, pyrimidinyl, pyrimidyl, pyrrolidinyl, pyrrolidin-2-onyl, pyrrolinyl, pyrrolyl, quinolinyl, quinoxaloyl, tetrahydrofuryl, tetrahydroisoquinolyl, tetrahydropyranyl, tetrahydroquinolyl, tetrazolyl, thiadiazolyl, thiazolidinyl, thiazolyl, thienyl, thiomorpholinyl, thiopyranyl, and triazolyl.

[062] The terms "hydroxy" and "hydroxyl" as used herein refer to -OH.

[063] The term "hydroxyalkyl" as used herein refers to a hydroxy attached to an alkyl group.

[064] The term "hydroxyaryl" as used herein refers to a hydroxy attached to an aryl group.

[065] The term "ketone" as used herein refers to the structure -C(O)-R_n (such as acetyl, -C(O)CH₃) or -R_n-C(O)-R_o-. The ketone can be attached to another group through R_n or R_o. R_n or R_o can be alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl or aryl, or R_n or R_o can be joined to form a 3- to 12-membered ring.

[066] The term "monoester" as used herein refers to an analogue of a dicarboxylic acid wherein one of the carboxylic acids is functionalized as an ester and the other carboxylic acid is a free carboxylic acid or salt of a carboxylic acid. Examples of monoesters include, but are not limited to, to monoesters of succinic acid, glutaric acid, adipic acid, suberic acid, sebacic acid, azelaic acid, oxalic and maleic acid.

[067] The term "phenyl" as used herein refers to a 6-membered carbocyclic aromatic ring. The phenyl group can also be fused to a cyclohexane or cyclopentane ring. Phenyl can be substituted with one or more substituents including alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl,

heteroaryl, heterocyclyl, hydroxyl, ketone, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide and thioketone.

[068] The term "thioalkyl" as used herein refers to an alkyl group attached to a sulfur (-S-alkyl-).

[069] "Alkyl," "alkenyl," "alkynyl," "alkoxy," "amino" and "amide" groups can be optionally substituted with or interrupted by or branched with at least one group selected from alkoxy, aryloxy, alkyl, alkenyl, alkynyl, amide, amino, aryl, arylalkyl, carbamate, carbonyl, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, ketone, phosphate, sulfide, sulfinyl, sulfonyl, sulfonic acid, sulfonamide, thioketone, ureido and N. The substituents may be branched to form a substituted or unsubstituted heterocycle or cycloalkyl.

[070] As used herein, a suitable substitution on an optionally substituted substituent refers to a group that does not nullify the synthetic or pharmaceutical utility of the compounds of the present disclosure or the intermediates useful for preparing them. Examples of suitable substitutions include, but are not limited to: C₁₋₈ alkyl, alkenyl or alkynyl; C₁₋₆ aryl, C₂₋₅ heteroaryl; C₃₋₇ cycloalkyl; C₁₋₈ alkoxy; C₆ aryloxy; -CN; -OH; oxo; halo, carboxy; amino, such as -NH(C₁₋₈ alkyl), -N(C₁₋₈ alkyl)₂, -NH((C₆)aryl), or -N((C₆)aryl)₂; formyl; ketones, such as -CO(C₁₋₈ alkyl), -CO((C₆)aryl) esters, such as -CO₂(C₁₋₈ alkyl) and -CO₂(C₆ aryl). One of skill in art can readily choose a suitable substitution based on the stability and pharmacological and synthetic activity of the compound of the present disclosure.

[071] The term "pharmaceutically acceptable carrier" as used herein refers to any and all solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. The compositions may also contain other active compounds providing supplemental, additional, or enhanced therapeutic functions.

[072] The term "pharmaceutically acceptable composition" as used herein refers to a composition comprising at least one compound as disclosed herein formulated together with one or more pharmaceutically acceptable carriers.

[073] The term "pharmaceutically acceptable prodrugs" as used herein represents those prodrugs of the compounds of the present disclosure that are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, commensurate with a reasonable benefit / risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the present disclosure. A discussion is provided in Higuchi *et al.*, "Prodrugs as Novel Delivery Systems," *ACS Symposium Series*, Vol. 14, and in Roche, E.B., ed. *Bioreversible Carriers in Drug*

[074] The term "pharmaceutically acceptable salt(s)" refers to salts of acidic or basic groups that may be present in compounds used in the present compositions. Compounds included in the present compositions that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, including but not limited to sulfate, citrate, malate, acetate, oxalate, chloride, bromide, iodide, nitrate, sulfite, bisulfite, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. Compounds included in the present compositions that include an amino moiety may form pharmaceutically acceptable salts with various amino acids, in addition to the acids mentioned above. Compounds included in the present compositions, that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts and, particularly, calcium, magnesium, sodium, lithium, zinc, potassium, and iron salts.

[075] The compounds of the disclosure may contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as geometric isomers, enantiomers or diastereomers. The term "stereoisomers" when used herein consist of all geometric isomers, enantiomers or diastereomers. These compounds may be designated by the symbols "R" or "S," depending on the configuration of substituents around the stereogenic carbon atom. The present disclosure encompasses various stereoisomers of these compounds and mixtures thereof. Stereoisomers include enantiomers and diastereomers. Mixtures of enantiomers or diastereomers may be designated "(±)" in nomenclature, but the skilled artisan will recognize that a structure may denote a chiral center implicitly.

[076] Individual stereoisomers of compounds of the present disclosure can be prepared synthetically from commercially available starting materials that contain asymmetric or stereogenic centers, or by preparation of racemic mixtures followed by resolution methods well known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary,

(2) salt formation employing an optically active resolving agent, or (3) direct separation of the mixture of optical enantiomers on chiral chromatographic columns. Stereoisomeric mixtures can also be resolved into their component stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Stereoisomers can also be obtained from stereomerically-pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.


[077] Geometric isomers can also exist in the compounds of the present disclosure. The present disclosure encompasses the various geometric isomers and mixtures thereof resulting from the arrangement of substituents around a carbon-carbon double bond or arrangement of substituents around a carbocyclic ring. Substituents around a carbon-carbon double bond are designated as being in the "Z" or "E" configuration wherein the terms "Z" and "E" are used in accordance with IUPAC standards. Unless otherwise specified, structures depicting double bonds encompass both the *E* and *Z* isomers.

[078] Substituents around a carbon-carbon double bond alternatively can be referred to as "cis" or "trans," where "cis" represents substituents on the same side of the double bond and "trans" represents substituents on opposite sides of the double bond. The arrangements of substituents around a carbocyclic ring are designated as "cis" or "trans." The term "cis" represents substituents on the same side of the plane of the ring and the term "trans" represents substituents on opposite sides of the plane of the ring. Mixtures of compounds wherein the substituents are disposed on both the same and opposite sides of plane of the ring are designated "cis/trans."

[079] The compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be encompassed by the scope of the present disclosure, even though only one tautomeric structure is depicted.

EXEMPLARY EMBODIMENTS OF THE INVENTION


[080] The invention provides compounds and pharmaceutical composition comprising one or more of those compounds wherein the structure of the compound is defined by Formula Ia, Formula Ib, Formula IIa, and/or Formula IIb:



or is a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof,

wherein:

A is selected from optionally substituted 5- or 6 membered monocyclic heterocycles fused to ring B.



B is a six-membered aromatic carbocycle or heterocycle;

Y is selected from N and C:

W_1 is selected from N and CR_1 ;

W_2 is selected from N and CR_2 ;

W_3 is selected from N and CR_3 ;

W_4 and W_5 , if present, are independently selected from N, CH, and C;

W_1 , W_2 , and W_3 may be the same or different from each other;

X is selected from -NH-, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂O-, -CH₂CH₂NH-, -CH₂CH₂S-, -C(O)-, -C(O)CH₂-, -C(O)CH₂CH₂-, -CH₂C(O)-, -CH₂CH₂C(O)-, -C(O)NH-, -C(O)O-, -C(O)S-, -C(O)NHCH₂-, -C(O)OCH₂-, -C(O)SCH₂-, -CH(OH)-, and -CH(CH₃)- where one or more hydrogen may independently be replaced with deuterium, hydroxymethyl, halogen, -CF₃, ketone, and where S may be oxidized to sulfoxide or sulfone;



R₄ is selected from 3-7 membered carbocycles and heterocycles;

D₁ is selected from 5-membered monocyclic heterocycles, where D₁ is attached to the B ring via a carbon atom that is part of a doublebond within the D₁ ring.

R_1 and R_2 are independently selected from hydrogen, deuterium, alkyl, -OH, -NH₂, -thioalkyl, alkoxy, ketone, ester, carboxylic acid, urea, carbamate, amino, amide, halogen, sulfone, sulfoxide, sulfide, sulfonamide, and -CN;

R_3 is selected from hydrogen, $-NH_2$, $-CN$, $-N_3$, halogen, deuterium, $-NO_2$, $-OMe$, $-OEt$, $-NHC(O)Me$, $NHSO_2Me$, cycloamino, cycloamido, $-OH$, $-SO_2Me$, $-SO_2Et$, $-CH_2NH_2$, $-C(O)NH_2$, and $-C(O)OMe$;


with the proviso that if R_3 is hydrogen and **A** is a 5-membered ring, then D_1 cannot be



and with the proviso that if D_1 is

and R_2 and R_3 are hydrogen and R_1 is $-OMe$, then


the **A-B** bicyclic ring is not



and with the proviso that if D_1 is


and each of R_1 , R_2 , R_3 , are hydrogen, then the

A-B bicyclic ring is not




unless the **B** ring is substituted;

and with the proviso that if each of R_1 , R_2 , R_3 are hydrogen, then the **A-B** bicyclic ring is not




and with the proviso that if each of R_1 , R_2 , R_3 are hydrogen, then the **A-B** bicyclic ring is not



-Me, -CF₃, -Et, CH₃CH₂O-, CF₃CH₂-, -SMe, -SOMe, -SO₂Me, -CN,




[082] In some embodiments, compounds of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, are selected from




which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as NH(C₁-C₅), carbocycle (C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHHeterocycle(C₄-C₆), -NHcarbocycle(C₄-C₆)), heterocycle(C₄-C₆), carbocycle(C₄-C₆), halogen, -CN, -OH, -CF₃, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆); wherein X, R₄, and D₁ are as defined for any embodiment disclosed herein.

[083] In some embodiments, the compounds of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, are selected from




which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₆), -NHcarbocycle(C₄-C₆), heterocycle(C₄-C₆), carbocycle(C₄-C₆), halogen, -CN, -OH, -CF₃, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆); wherein X, R₄, and D₁ are as defined for any embodiment disclosed herein.

[084] In some embodiments, the compounds of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, are selected from




which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₆), -NHcarbocycle(C₄-C₆)), heterocycle(C₄-C₆), carbocycle(C₄-C₆), halogen, -CN, -OH, -CF₃, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆); wherein X, R₄, and D₁ are as defined for any embodiment disclosed herein.

[085] In some embodiments, compounds of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, are selected from




which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₆), -NHcarbocycle(C₄-C₆)), heterocycle(C₄-C₆), carbocycle(C₄-C₆), halogen, -CN, -OH, -CF₃, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆); wherein the definition of X, R₄, and D₁ are as defined for any embodiment disclosed herein.

[086] In some embodiments, compounds of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from



wherein Z is selected from hydrogen, deuterium, $-NH_2$, amino (such as $-NH(C_1-C_5)$, $-N(C_1-C_5)_2$, $-NHPh$, $-NHBn$, $-NHpyridyl$, $-NHheterocycle(C_4-C_6)$, $-NHcarbocycle(C_4-C_6)$), alkyl(C_1-C_6), thioalkyl(C_1-C_6), alkenyl(C_1-C_6), and alkoxy(C_1-C_6); carboxyl;




X is selected from $-CH_2-$ and $-CH(CH_3)-$; and

R_4 is a phenyl ring optionally substituted with groups independently selected with one or more groups independently selected from deuterium, alkyl(C_1-C_4), alkoxy(C_1-C_4), halogen, $-CF_3$, CN, and $-thioalkyl(C_1-C_4)$, wherein each alkyl, alkoxy, and thioalkyl may be optionally substituted with F, Cl, or Br.


[087] In certain embodiments, R_4 is a phenyl ring is optionally substituted with one or more alkyl(C_1-C_4) selected from methyl, ethyl, propyl, isopropyl, and butyl; alkoxy(C_1-C_4), selected from methoxy, ethoxy, and isopropoxy; halogen selected from F and Cl; and thioalkyl(C_1-C_4) selected from $-SMe$, $-SEt$, $-SPr$, and $-Sbu$.

[088] In some embodiments, the A-B bicyclic ring in the compound of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from




wherein Z is selected from hydrogen, deuterium, $-NH_2$, amino (such as $-NH(C_1-C_5)$, $-N(C_1-C_5)_2$, $-NHPh$, $-NHBn$, $-NHpyridyl$, $-NHheterocycle(C_4-C_6)$, $-NHcarbocycle(C_4-C_6)$), alkyl(C_1-C_6), thioalkyl(C_1-C_6), alkenyl(C_1-C_6), and alkoxy(C_1-C_6).

[089] In some embodiments, the A-B bicyclic ring in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from




[090] In some embodiments, the A-B bicyclic ring in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from, but not limited to




which may be optionally substituted with groups independently selected from hydrogen, deuterium, $-NH_2$, amino (such as $-NH(C_1-C_5)$, $-N(C_1-C_5)_2$, $-NHPh$, $-NHBn$, $-NHpyridyl$, $-NHheterocycle(C_4-C_7)$, $-NHcarbocycle(C_4-C_7)$), heterocycle(C_4-C_7), carbocycle(C_4-C_7), halogen, $-CN$, $-OH$, $-CF_3$, sulfone, sulfoxide, alkyl(C_1-C_6), thioalkyl(C_1-C_6), Alkenyl(C_1-C_6), alkoxy(C_1-C_6), ketone(C_1-C_6), ester, urea, carboxylic acid, carbamate, amide(C_1-C_6), oxo, and thio-oxo.

[091] In some embodiments of any of Formula Ia, Formula Ib, Formula IIa, and Formula IIb, or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, the A-B bicyclic ring, is selected from



which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₇), -NHcarbocycle(C₄-C₇)), heterocycle(C₄-C₇), carbocycle(C₄-C₇), halogen, -CN, -OH, -CF₃, sulfone, sulfoxide, alkyl(C₁-C₆), thioalkyl(C₁-C₆), Alkenyl(C₁-C₆), alkoxy(C₁-C₆), ketone(C₁-C₆), ester, urea, carboxylic acid, carbamate, amide(C₁-C₆), oxo, and thio-oxo.


[092] In some embodiments, the A-B bicyclic ring in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from



which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₇), -NHcarbocycle(C₄-C₇)), heterocycle(C₄-C₇), carbocycle(C₄-C₇), halogen, -CN, -OH, -CF₃, sulfone,

sulfoxide, alkyl(C₁-C₆), thioalkyl(C₁-C₆), Alkenyl(C₁-C₆), alkoxy(C₁-C₆), ketone(C₁-C₆), ester, urea, carboxylic acid, carbamate, amide(C₁-C₆), oxo, and thio-oxo.


[093] In some embodiments, the A-B bicyclic ring in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from



which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₇), -NHcarbocycle(C₄-C₇)), heterocycle(C₄-C₇), carbocycle(C₄-C₇), halogen, -CN, -OH, -CF₃, sulfone, sulfoxide, sulfonamide, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), alkoxy(C₁-C₆), ketone(C₁-C₆), ester, urea, carboxylic acid, carbamate, amide(C₁-C₆), oxo, and thio-oxo.

[094] In some embodiments, the A-B bicyclic ring in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically

acceptable salt, or hydrate thereof, is selected from




which may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₇), -NHcarbocycle(C₄-C₇)), heterocycle(C₄-C₇), carbocycle(C₄-C₇), halogen, -CN, -OH, -CF₃, sulfone, sulfoxide, sulfonamide, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), alkoxy(C₁-C₆), ketone(C₁-C₆), ester, urea, carboxylic acid, carbamate, amide(C₁-C₆), oxo, and thio-oxo.

[095] In some embodiments, the A ring in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 5-membered heterocycles fused to the B ring.

[096] In some embodiments, Y in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is nitrogen.

[097] In some embodiments, D_1 in the compound of any one of Formula I, Formula Ia, or Formula II or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from an 5-membered monocyclic heterocycle, such as, but not limited to:



which is optionally substituted with hydrogen, deuterium, alkyl(C_1-C_4) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C_1-C_4) (such as methoxy, ethoxy, isopropoxy), amino (such as $-NH_2$, $-NHMe$, $-NHEt$, $-NHPr$, $-NHBu$, $-NMe_2$, $NMeEt$, $-NEt_2$, $-NEtBu$, $-NHC(O)NHalkyl$), halogen (such as F, Cl), amide (such as $-NHC(O)Me$, $-NHC(O)Et$, $-C(O)NHMe$, $-C(O)NEt_2$, $-C(O)NiPr$), $-CF_3$, CN, $-N_3$, ketone (C_1-C_4) (such as acetyl, $-C(O)Et$, $-C(O)Pr$), $-S(O)Alkyl(C_1-C_4)$ (such as $-S(O)Me$, $-S(O)Et$), $-SO_2alkyl(C_1-C_4)$ (such as $-SO_2Me$, $-SO_2Et$, $-SO_2Pr$), $-thioalkyl(C_1-C_4)$ (such as $-SMe$, $-SEt$, $-SPr$, $-SBu$), $-COOH$, and/or ester (such as $-C(O)OMe$, $-C(O)OEt$, $-C(O)OBu$), each of which may be optionally substituted with hydrogen, F, Cl, Br, $-OH$, $-NH_2$, $-NHMe$, $-OMe$, $-SMe$, oxo, and/or thio-oxo.

[098] In some embodiments, D_1 in the compound of any one of Formula Ia, Ib, IIa, or IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is a monocyclic heterocycle optionally substituted with hydrogen, deuterium, alkyl(C_1-C_4) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C_1-C_4) (such as methoxy, ethoxy, isopropoxy), amino (such as $-NH_2$, $-NHMe$, $-NHEt$, $-NHPr$, $-NHBu$, $-NMe_2$, $NMeEt$, $-NEt_2$, $-NEtBu$, $-NHC(O)NHalkyl$), halogen (such as F, Cl), amide (such as $-NHC(O)Me$, $-NHC(O)Et$, $-C(O)NHMe$, $-C(O)NEt_2$, $-C(O)NiPr$), $-CF_3$, CN, $-N_3$, ketone (C_1-C_4) (such as acetyl, $-C(O)Et$, $-C(O)Pr$), $-S(O)Alkyl(C_1-C_4)$ (such as $-S(O)Me$, $-S(O)Et$), $-SO_2alkyl(C_1-C_4)$ (such as $-SO_2Me$, $-SO_2Et$, $-SO_2Pr$), $-thioalkyl(C_1-C_4)$ (such as $-SMe$, $-SEt$, $-SPr$, $-SBu$), $-COOH$, and/or ester (such as $-C(O)OMe$, $-C(O)OEt$, $-C(O)OBu$), each of which may be optionally substituted with hydrogen, F, Cl, Br, $-OH$, $-NH_2$, $-NHMe$, $-OMe$, $-SMe$, oxo, and/or thio-oxo.

[099] In some embodiments, D_1 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from a 5-membered monocyclic heterocycle containing one oxygen and one or two nitrogens, where the heterocycle is connected to the rest of the molecule via a carbon-carbon bond, and which is optionally substituted with hydrogen, deuterium, alkyl(C_1-C_4) (such as methyl,

ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)N₂Pr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), -COOH, and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0100] In some embodiments, D₁ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is an isoxazole optionally substituted with hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)N₂Pr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), -COOH, and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.


[0101] In some embodiments, D₁ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from an 5-membered monocyclic heterocycle, which is optionally substituted with hydrogen, deuterium, Alkyl(C₁-C₄), (such as methyl, ethyl, propyl), each of which may be optionally substituted with hydrogen, -OH, -F, and -NH₂.

[0102] In some embodiments, D₁ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from a 5-membered monocyclic heterocycle containing one oxygen and one or two nitrogens, where the heterocycle is connected to the rest of the molecule via a carbon-carbon bond, and which is optionally substituted with hydrogen, deuterium, Alkyl(C₁-C₄), (such as methyl, ethyl, propyl), each of which may be optionally substituted with hydrogen, -OH, -F, and -NH₂.

[0103] In some embodiments, D₁ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is an isoxazole or pyrazole optionally substituted with hydrogen, deuterium, Alkyl(C₁-C₄), (such as methyl, ethyl, propyl), each of which may be optionally substituted with hydrogen, -OH, -F, and -NH₂.

[0104] In some embodiments, D_1 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is

[0105] In some embodiments, D_1 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is In some embodiments, D_1 in the compound of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is



[0106] In some embodiments, W_1 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is CR_1 .

[0107] In some embodiments, W_2 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is CR_2 .

[0108] In some embodiments, at least one of W_1 and W_2 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is nitrogen.

[0109] In some embodiments, W_1 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is CH.

[0110] In some embodiments, W_2 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is CR_2 , where R_2 is selected from hydrogen, deuterium, -OH, -NH₂, methyl, halogen, and -CN.

[0111] In some embodiments, W_2 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is CH.

[0112] In some embodiments, W_4 and W_5 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof are carbon.

[0113] In some embodiments, at least one of W_4 and W_5 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is nitrogen.

[0114] In some embodiments, W_3 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is nitrogen.

[0115] In some embodiments, W_3 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is CR_3 , where R_3 is selected from hydrogen, $-NH_2$, and halogen.

[0116] In some embodiments, R_3 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof is selected from hydrogen and $-NH_2$.

[0117] In some embodiments, R_3 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is $-NH_2$.

[0118] In some embodiments, X in the compound of any one of Formula I, Formula Ia, or Formula II or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2O-$, $-CH_2CH_2NH-$, $-CH_2CH_2S-$, $-C(O)-$, $-C(O)NH-$, $-C(O)O-$, $-C(O)S-$, where one or more hydrogen may independently be replaced with deuterium, halogen, and where S may be oxidized to sulfoxide or sulfone.

[0119] In some embodiments, X in the compound of any one of Formula I, Formula Ia, or Formula II or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from $-CH_2-$ and $-C(O)-$.

[0120] In some embodiments, X is selected from $-CH_2-$, $-CH(CH_3)-$, $-CH(OH)-$, $-NH-$, $-CH_2CH_2-$, where one or more hydrogen may independently be replaced with deuterium or halogen.

[0121] In some embodiments, X is selected from $-CH_2-$, $CH(CH_3)-$, and $-NH-$, where one or more hydrogen may independently be replaced with deuterium or halogen.

[0122] In some embodiments, X is selected from $-CH_2-$, $-CH(CH_3)-$, where one or more hydrogen may independently be replaced with deuterium or halogen.

[0123] In some embodiments, X in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is $-CH_2-$.

[0124] In some embodiments, R_1 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate

thereof, is selected from hydrogen, deuterium, alkyl, -OH, -NH₂, -thioalkyl, alkoxy, ketone, ester, carboxylic acid, urea, carbamate, amino, amide, halogen, carbocycle, heterocycle, sulfone, sulfoxide, sulfide, sulfonamide, and and -CN.

[0125] In some embodiments, R₂ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from hydrogen, deuterium, alkyl, -OH, -NH₂, -thioalkyl, alkoxy, ketone, ester, carboxylic acid, urea, carbamate, amino, amide, halogen, carbocycle, heterocycle, sulfone, sulfoxide, sulfide, sulfonamide, and and -CN.

[0126] In some embodiments, R₁ and R₂ in the compound of any one of Formula I, Formula Ia, or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof are independently selected from hydrogen, deuterium, alkyl, -NH₂, -thioalkyl, alkoxy, amino, amide, halogen, carbocycle, heterocycle, and -CN.

[0127] In some embodiments, R₁ and R₂ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, are independently selected from hydrogen, deuterium, alkyl(C₁-C₆), -NH₂, -thioalkyl(C₁-C₆), alkoxy(C₁-C₆), amino, and amide.

[0128] In some embodiments, R₁ and R₂ are hydrogen.

[0129] In some embodiments, at least one of R₁, R₂, and R₃ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is not hydrogen.

[0130] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 5-6 membered carbocycles and heterocycles.


[0131] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 5-6 membered heterocycles.

[0132] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 5-6 membered heterocycles containing 1 or 2 nitrogens, such as unsubstituted and substituted pyrimidyl rings, which are optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂iPr, -NH₂Bu, -NHMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄)

(such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0133] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 6-membered heterocycles containing at least one nitrogen, such as unsubstituted and substituted pyridyl rings, which are optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂iPr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0134] In some embodiments, R₄ in the compound of any one of Formula I, Formula Ia, or Formula II or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from



[0135] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is an isoxazole or pyrazole optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂iPr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -

S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0136] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from a 5-membered heterocycle containing one or two nitrogens.

[0137] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 5-6 membered carbocycles, such as a phenyl ring optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0138] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is a phenyl ring optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0139] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from an aryl optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt,

-NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0140] In some embodiments, in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, -X-R₄ is selected from -CH₂Aryl.

[0141] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from a pyridyl optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NHET, -NHIPr, -NHBu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0142] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NHET, -NHIPr, -NHBu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)NiPr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.


[0143] In some embodiments, R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from 5-6 membered carbocycles.

[0144] In some embodiments, R_4 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is selected from a small cycloalkyl(C_3 - C_6) and phenyl ring optionally optionally substituted with one or more groups independently selected from deuterium, alkyl(C_1 - C_4) (such as methyl, ethyl, propyl, isopropyl, and butyl), alkoxy(C_1 - C_4) (such as methoxy, ethoxy, and isopropoxy), halogen (such as F and Cl), $-CF_3$, CN, and -thioalkyl(C_1 - C_4) (such as, e.g., -SMe, -SEt, -SPr, and -Sbu), wherein each alkyl, alkoxy, and thioalkyl may be optionally substituted with F, Cl, or Br.

[0145] In some embodiments, R_4 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is a phenyl ring optionally substituted with one or more groups independently selected from deuterium, alkyl(C_1 - C_4) (such as methyl, ethyl, propyl, isopropyl, and butyl), alkoxy(C_1 - C_4) (such as methoxy, ethoxy, and isopropoxy), halogen (such as F and Cl), $-CF_3$, CN, and -thioalkyl(C_1 - C_4) (such as, e.g., -SMe, -SEt, -SPr, and -Sbu), wherein each alkyl, alkoxy, and thioalkyl may be optionally substituted with F, Cl, or Br.


[0146] In some embodiments, R_4 in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is an aryl optionally substituted with one or more groups independently selected from deuterium, alkyl(C_1 - C_4) (such as methyl, ethyl, propyl, isopropyl, and butyl), alkoxy(C_1 - C_4) (such as methoxy, ethoxy, and isopropoxy), halogen (such as F and Cl), $-CF_3$, CN, and -thioalkyl(C_1 - C_4) (such as, e.g., -SMe, -SEt, -SPr, and -Sbu), wherein each alkyl, alkoxy, and thioalkyl may be optionally substituted with F, Cl, or Br.

[0147] In some embodiments, in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, the A-B bicyclic ring, is selected from



which may be optionally substituted with groups independently selected from hydrogen, deuterium, $-NH_2$, amino (such as $-NH(C_1-C_5)$, $-N(C_1-C_5)_2$, $-NHPh$, $-NHBn$, $-NHpyridyl$, $-NHheterocycle(C_4-C_7)$, $-NHcarbocycle(C_4-C_7)$), heterocycle(C_4-C_7), carbocycle(C_4-C_7), halogen, $-CN$, $-OH$, $-CF_3$, sulfone,


sulfoxide, alkyl(C₁-C₆), thioalkyl(C₁-C₆), Alkenyl(C₁-C₆), alkoxy(C₁-C₆), ketone(C₁-C₆), ester, urea, carboxylic acid, carbamate, amide(C₁-C₆), oxo, and thio-oxo;




X is selected from -CH₂- and -C(O)-;

R₄ is a phenyl ring optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)N₂Pr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SP₂, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0148] In some embodiments, in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate



may be optionally substituted with groups independently selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₇), -NHcarbocycle(C₄-C₇)), heterocycle(C₄-C₇), carbocycle(C₄-C₇), halogen, -CN, -OH, -CF₃, sulfone, sulfoxide, sulfonamide, alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), alkoxy(C₁-C₆), ketone(C₁-C₆), ester, urea, carboxylic acid, carbamate, amide(C₁-C₆), oxo, and thio-oxo.




X is selected from -CH₂-, -CH(CH₃)-, -CH(OH)-, and -NH-;

R₄ is a phenyl ring optionally substituted with groups independently selected from hydrogen, deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, isopropoxy), amino (such as -NH₂, -NHMe, -NH₂Et, -NH₂Pr, -NH₂Bu -NMe₂, NMeEt, -NEt₂, -NEtBu, -NHC(O)NHalkyl), halogen (such as F, Cl), amide (such as -NHC(O)Me, -NHC(O)Et, -C(O)NHMe, -C(O)NEt₂, -C(O)N₂Pr), -CF₃, CN, -N₃, ketone (C₁-C₄) (such as acetyl, -C(O)Et, -C(O)Pr), -S(O)Alkyl(C₁-C₄) (such as -S(O)Me, -S(O)Et), -SO₂alkyl(C₁-C₄) (such as -SO₂Me, -SO₂Et, -SO₂Pr), -


thioalkyl(C₁-C₄) (such as -SMe, -SEt, -SPr, -SBu), carboxyl (such as -COOH), and/or ester (such as -C(O)OMe, -C(O)OEt, -C(O)OBu), each of which may be optionally substituted with hydrogen, F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

[0149] In some embodiments, -X-R₄ in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, is -CH₂Aryl.

[0150] In some embodiments, in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, the A-B bicyclic ring is selected from



wherein Z is selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₆), -NHcarbocycle(C₄-C₆)), alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆), carboxyl;




D₁ is ; and


X is selected from -CH₂- and -CH(CH₃)-; and

R₄ is a phenyl ring optionally substituted with groups independently selected with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, CN, and -thioalkyl(C₁-C₄), wherein each alkyl, alkoxy, and thioalkyl may be optionally substituted with F, Cl, or Br.

[0151] In some embodiments, in the compound of any one of Formula Ia, Formula Ib, Formula IIa, and Formula IIb or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, the A-B bicyclic ring is selected from



wherein **Z** is selected from hydrogen, deuterium, -NH₂, amino (such as -NH(C₁-C₅), -N(C₁-C₅)₂, -NHPh, -NHBn, -NHpyridyl, -NHheterocycle(C₄-C₆), -NHcarbocycle(C₄-C₆)), alkyl(C₁-C₆), thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆); carboxyl;



X is selected from -CH₂- and -CH(CH₃)-; and

R₄ is a phenyl ring optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄) (such as methyl, ethyl, propyl, isopropyl, and butyl), alkoxy(C₁-C₄) (such as methoxy, ethoxy, and isopropoxy), halogen (such as F and Cl), -CF₃, CN, and -thioalkyl(C₁-C₄) (such as, e.g., -SMe, -SEt, -SPr, and -Sbu), wherein each alkyl, alkoxy, and thioalkyl may be optionally substituted with F, Cl, or Br.

[0152] In certain embodiments of the invention, the compound of Formula I, Formula Ia, or Formula II is selected from:

9-Benzyl-2-(3,5-dimethylisoxazol-4-yl)-9H-purin-6-amine;
 3-Benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one;
 1-Benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one;
 4-(3-Benzyl-3H-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole;
 4-(1-Benzyl-1H-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole;
 3-Benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[*d*]oxazol-2(3H)-one;
 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[*d*]imidazol-4-amine;
 1-Benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-benzo[*d*]imidazol-7-amine;
N,1-Dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[*d*]imidazol-4-amine;
 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one;
 1-Benzyl-7-(3,5-dimethylisoxazol-4-yl)quinoxalin-2(1H)-one; and
 1-Benzyl-7-(3,5-dimethylisoxazol-4-yl)-3,4-dihydroquinazolin-2(1H)-one.

[0153] In certain embodiments of the invention, the compound of Formula I, Formula Ia, or Formula II is selected from:

9-benzyl-2-(3,5-dimethylisoxazol-4-yl)-9H-purin-6-amine;
 3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one;
 1-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one;
 4-(3-benzyl-3H-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole;
 4-(1-benzyl-1H-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole;

3-benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[d]oxazol-2(3H)-one;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine;
 1-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-7-amine;
 N,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 1-benzyl-7-(3,5-dimethylisoxazol-4-yl)quinoxalin-2(1H)-one;
 1-benzyl-7-(3,5-dimethylisoxazol-4-yl)-3,4-dihydroquinazolin-2(1H)-one;
 4-(1-benzyl-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
 4-(1-(cyclopropylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one;
 4-amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-ethoxy-1H-benzo[d]imidazol-4-amine;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-4-nitro-1H-benzo[d]imidazol-2-amine;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N₂-ethyl-1H-benzo[d]imidazole-2,4-diamine;
 methyl 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carboxylate;
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carboxamide
 4-(aminomethyl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one
 5-(3,5-dimethylisoxazol-4-yl)-N-phenyl-1H-pyrrolo[3,2-b]pyridin-3-amine
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-b]pyridine 4-oxide
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-b]pyridin-5(4H)-one
 4-(3-benzyl-3H-imidazo[4,5-b]pyridin-5-yl)-3,5-dimethylisoxazole
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-benzo[d]imidazol-4-amine
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N-methyl-1H-benzo[d]imidazol-4-amine
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N,N-dimethyl-1H-benzo[d]imidazol-4-amine
 3,5-dimethyl-4-(1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 4-(1-benzyl-1H-imidazo[4,5-c]pyridin-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-c]pyridine 5-oxide
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-c]pyridin-4-amine
 4-(1-benzyl-3-bromo-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridine-3-carbaldehyde
 1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)ethanone

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-5-yl formate
 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benzamide
 4-(1-benzyl-3-nitro-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole
 3,5-dimethyl-4-(3-(4-(trifluoromethyl)benzyl)-3H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 3,5-dimethyl-4-(1-(4-(trifluoromethyl)benzyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 4-(3-(4-chlorobenzyl)-3H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(4-chlorobenzyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(3-(4-fluorobenzyl)-3H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 3,5-dimethyl-4-(3-(pyridin-2-ylmethyl)-3H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 3,5-dimethyl-4-(1-(pyridin-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 4-(1-(4-fluorobenzyl)-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(4-fluorobenzyl)-1H-pyrrolo[2,3-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(5-(4-fluorobenzyl)-5H-pyrrolo[2,3-b]pyrazin-3-yl)-3,5-dimethylisoxazole
 4-(1-(4-fluorobenzyl)-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-pyrrolo[2,3-b]pyridin-4-amine
 4-(1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-indazol-4-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-5(4H)-one
 3-((5-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)amino)benzonitrile
 4-(1-(4-fluorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-benzyl-2-ethoxy-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)-3,5-dimethylisoxazole
 4-(1-(2,4-dichlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(4-methoxybenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(cyclopropylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)acetamide
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)ethanesulfonamide
 4-(1-benzyl-4-methoxy-2-methyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 7-amino-3-benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[d]oxazol-2(3H)-one
 3,5-dimethyl-4-(2-methyl-1-(pyridin-3-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole

3,5-dimethyl-4-(2-methyl-1-(thiophen-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benzonitrile
 4-(1-benzyl-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)-N,N-dimethylmethanamine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[2,3-b]pyridin-4-amine
 3,5-dimethyl-4-(2-methyl-1-(pyridin-4-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine
 3,5-dimethyl-4-(2-methyl-1-((5-methylthiophen-2-yl)methyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 4-(1-((5-chlorothiophen-2-yl)methyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 5-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)thiophene-2-carbonitrile
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridine 4-oxide
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-5-yl acetate
 1-benzyl-6-(1,4-dimethyl-1H-pyrazol-5-yl)-2-methyl-4-nitro-1H-benzo[d]imidazole
 1-benzyl-6-(1,4-dimethyl-1H-pyrazol-5-yl)-2-methyl-1H-benzo[d]imidazol-4-amine
 4-(1-(4-chlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)phenol
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazole-4-carbonitrile
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carbonitrile
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-morpholino-1H-benzo[d]imidazol-4-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridine-3-carbonitrile
 4-(1-benzyl-3-chloro-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-amino-1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one
 1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one
 4-(1-benzyl-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(4-chlorobenzyl)-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-benzyl-2-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-benzo[d]imidazol-4-amine
 4-(1-(3,4-dichlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1H-benzo[d]imidazol-4-amine
 2-(azetidin-1-yl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine

3,5-dimethyl-4-(1-(thiophen-3-ylmethyl)-1H-pyrazolo[4,3-b]pyridin-6-yl)isoxazole
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)acetamide
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-amine
 1-(3,4-dichlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-indazol-4-amine
 6-(3,5-dimethylisoxazol-4-yl)-1-(4-methoxybenzyl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one
 4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(4-methoxybenzyl)-1H-benzo[d]imidazol-2(3H)-one
 1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 6-(3,5-dimethylisoxazol-4-yl)-1-(thiophen-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine
 3,5-dimethyl-4-(2-methyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-(tetrahydro-2H-pyran-4-yl)-1H-benzo[d]imidazole-
 2,4-diamine
 6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-4-
 yl)acetamide
 6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2-amine
 4-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine
 4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one
 4-(1-(cyclobutylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 4-(1-(cyclopentylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-(ethylamino)-1H-benzo[d]imidazol-4-
 yl)acetamide
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-ethoxy-1H-benzo[d]imidazol-4-yl)acetamide
 4-(1-benzyl-4-bromo-2-methyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-1H-benzo[d]imidazol-2(3H)-one
 4-(2-(azetidin-1-yl)-1-benzyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-((5-chlorothiophen-2-yl)methyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-
 2(3H)-one
 (S)-3,5-dimethyl-4-(2-methyl-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-6-yl)isoxazole
 (R)-3,5-dimethyl-4-(2-methyl-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-6-yl)isoxazole
 6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-2-amine

4-(1-benzyl-2-ethyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(4-hydroxybenzyl)-1H-benzo[d]imidazol-2(3H)-one
 N-(2-(azetidin-1-yl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-yl)acetamide
 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine
 1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine
 1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine
 6-(3,5-dimethylisoxazol-4-yl)-N2-ethyl-1-(1-phenylethyl)-1H-benzo[d]imidazole-2,4-diamine
 4-(1-benzyl-4-nitro-2-(pyrrolidin-1-yl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 4-(1-benzyl-2-(4-methylpiperazin-1-yl)-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-methoxyethyl)-4-nitro-1H-benzo[d]imidazol-2-amine
 4-(1-benzyl-2-cyclopropyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-(2-methoxyethyl)-1H-benzo[d]imidazole-2,4-diamine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-(pyrrolidin-1-yl)-1H-benzo[d]imidazol-4-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-(4-methylpiperazin-1-yl)-1H-benzo[d]imidazol-4-amine
 1-benzyl-N6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazole-4,6-diamine
 (S)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1H-benzo[d]imidazol-4-amine
 (R)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1H-benzo[d]imidazol-4-amine
 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine
 N,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-N-(pyridin-3-ylmethyl)-1H-benzo[d]imidazol-2-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-4-nitro-1H-benzo[d]imidazol-2-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-4-nitro-1H-benzo[d]imidazol-2(3H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-methyl-1H-benzo[d]imidazole-2,4-diamine
 N2,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2,4-diamine
 N,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine
 1-benzyl-2-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridine

N-(1-benzyl-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazol-4-amine
 4-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3,4-dihydroquinoxalin-2(1H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-(pyridin-3-ylmethyl)-1H-benzo[d]imidazole-2,4-diamine
 4-(1-benzyl-4-fluoro-2-methyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-4-nitro-1H-benzo[d]imidazol-2-amine
 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N2-ethyl-1H-benzo[d]imidazole-2,4-diamine
 4-amino-1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one
 4-amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-1H-benzo[d]imidazol-2(3H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-4-fluoro-1H-benzo[d]imidazol-2(3H)-one
 N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-4-yl)acetamide
 4-(1-benzyl-2-(4-methylpiperazin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-3,4-dihydroquinoxalin-2(1H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-methoxyethyl)-1H-imidazo[4,5-b]pyridin-2-amine
 4-(1-benzyl-2-methyl-4-(methylsulfonyl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-4-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine
 1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 (S)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-ol
 (R)-4-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-3,4-dihydroquinoxalin-2(1H)-one
 4-(1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine
 1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine
 4-amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2(3H)-thione
 (S)-4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one
 (R)-4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-7-methyl-1H-imidazo[4,5-b]pyridin-2(3H)-one
 4-(1-benzyl-2,7-dimethyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole

4-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)morpholine
 1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)azetidin-2-one
 1-benzyl-2-methyl-6-(1,3,5-trimethyl-1H-pyrazol-4-yl)-1H-benzo[d]imidazol-4-amine
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-3-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine
 4-(4-bromo-2-methyl-1-phenethyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 4-(4-bromo-2-methyl-1-(3-phenylpropyl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 4-(7-bromo-2-methyl-1-(3-phenylpropyl)-1H-benzo[d]imidazol-5-yl)-3,5-dimethylisoxazole
 4-(4-bromo-2-methyl-1-(2-phenoxyethyl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole
 4-(7-bromo-2-methyl-1-(2-phenoxyethyl)-1H-benzo[d]imidazol-5-yl)-3,5-dimethylisoxazole
 4-(1-(cyclohexylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(cyclopentylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 4-(1-(cyclobutylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine
 4-(1-benzyl-2-(pyrrolidin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole
 2-((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)ethanol
 1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)azetidin-3-ol
 1-benzyl-3-methyl-6-(1-methyl-1H-pyrazol-5-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one
 4-amino-1-benzyl-3-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-benzo[d]imidazol-2(3H)-one
 (4-bromo-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-1-yl)(phenyl)methanone
 1-benzyl-2-methyl-6-(5-methylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine
 1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one
 N-(1-benzyl-3-methyl-6-(1-methyl-1H-pyrazol-5-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-4-yl)acetamide
 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(4-methoxybenzyl)-1H-imidazo[4,5-b]pyridin-2-amine
 1-benzyl-2-methyl-6-(1-methyl-1H-1,2,3-triazol-5-yl)-1H-imidazo[4,5-b]pyridine
 4-((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)cyclohexanol
 4-(1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine

4-(2-(azetidin-1-yl)-1-(cyclopentylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole

4-(1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine

4-(2-(azetidin-1-yl)-1-(cyclobutylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole

N1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)-N2,N2-dimethylethane-1,2-diamine

4-(1-benzyl-2-(piperazin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole

1-benzyl-N-cyclopentyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-morpholinoethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

3-(((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)methyl)benzonitrile;

(R)-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

(S)-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

4-(1-benzyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridine-2-carboxamide;

1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

N1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)cyclohexane-1,4-diamine;

1-benzyl-N-(cyclohexylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(3-methoxypropyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(oxetan-3-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyrazin-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-(4-methylpiperazin-1-yl)ethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine;

1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-N-cyclohexyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(1-methylpiperidin-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

4-(1-benzyl-2-(pyridin-3-yloxy)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)-2-methylpropan-2-ol;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-(pyrrolidin-1-yl)ethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-(piperidin-1-yl)ethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

(R)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one;

4-(1-benzyl-7-methoxy-2-(trifluoromethyl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(thiazol-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2-carboximidamide;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2-carboxamide;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-((1-methylpiperidin-4-yl)methyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)azetidin-3-ol;

4-(1-benzyl-2-(pyridin-4-yloxy)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-3-yl)-1H-benzo[d]imidazol-2-amine; and

3-(1-benzyl-1H-benzo[d]imidazol-6-yl)-4-ethyl-1H-1,2,4-triazol-5(4H)-one;

or a stereoisomer, tautomer, salt, or hydrate thereof.

[0154] Another aspect of the invention provides a method for inhibition of BET protein function by binding to bromodomains, and their use in the treatment and prevention of diseases and conditions in a mammal (e.g., a human) comprising administering a therapeutically effective amount of a compound of Formula I, Formula Ia, and/or Formula II.

[0155] In one embodiment, because of potent effects of BET inhibitors *in vitro* on IL-6 and IL-17 transcription, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II may be used as therapeutics for inflammatory disorders in which IL-6 and/or IL-17 have been implicated in disease. The following autoimmune diseases are amenable to therapeutic use of BET inhibition by administration of a compound of Formula I, Formula Ia, and/or Formula II or stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate of Formula I, Formula Ia, and/or Formula II because of a prominent role of IL-6 and/or IL-17: Acute Disseminated Encephalomyelitis (Ishizu, T., et al., "CSF cytokine and chemokine profiles in acute disseminated encephalomyelitis," *J Neuroimmunol* 175(1-2): 52-8 (2006)), Agammaglobulinemia (Gonzalez-Serrano, M.E., et al., "Increased Pro-inflammatory Cytokine Production After Lipopolysaccharide Stimulation in Patients with X-linked Agammaglobulinemia," *J Clin Immunol* 32(5):967-74 (2012)), Allergic Disease (McKinley, L., et al., "TH17 cells mediate steroid-resistant airway inflammation and airway hyperresponsiveness in mice," *J Immunol* 181(6):4089-97 (2008)), Ankylosing spondylitis (Taylan, A., et al., "Evaluation of the T helper 17 axis in ankylosing spondylitis," *Rheumatol Int* 32(8):2511-5 (2012)), Anti-GBM/Anti-TBM nephritis (Ito, Y., et al., "Pathogenic significance of interleukin-6 in a patient with anti-glomerular basement membrane antibody-induced glomerulonephritis with multinucleated giant cells," *Am J Kidney Dis* 26(1):72-9 (1995)), Anti-phospholipid syndrome (Soltesz, P., et al., "Immunological features of primary anti-phospholipid syndrome in connection with endothelial dysfunction," *Rheumatology (Oxford)* 47(11):1628-34 (2008)), Autoimmune aplastic anemia (Gu, Y., et al., "Interleukin (IL)-17 promotes macrophages to produce IL-8, IL-6 and tumour necrosis factor- α in aplastic anaemia," *Br J Haematol* 142(1):109-14 (2008)), Autoimmune hepatitis (Zhao, L., et al., "Interleukin-17 contributes to the pathogenesis of autoimmune hepatitis through inducing hepatic interleukin-6 expression," *PLoS One* 6(4):e18909 (2011)), Autoimmune inner ear disease (Gloddek, B., et al., "Pharmacological influence on inner ear endothelial cells in relation to the pathogenesis of sensorineural hearing loss," *Adv Otorhinolaryngol* 59:75-83 (2002)), Autoimmune myocarditis (Yamashita, T., et al., "IL-6-mediated Th17 differentiation through ROR γ is essential for the initiation of experimental autoimmune myocarditis," *Cardiovasc Res* 91(4):640-8 (2011)), Autoimmune pancreatitis (Ni, J., et al., "Involvement of Interleukin-17A in Pancreatic Damage in Rat Experimental Acute Necrotizing Pancreatitis," *Inflammation* (2012)), Autoimmune retinopathy (Hohki, S., et al., "Blockade of interleukin-6 signaling suppresses experimental autoimmune uveoretinitis by the inhibition of inflammatory Th17 responses," *Exp Eye Res* 91(2):162-70 (2010)), Autoimmune thrombocytopenic purpura (Ma, D., et al., "Profile of Th17 cytokines (IL-17, TGF- β , IL-6) and Th1 cytokine (IFN- γ) in patients with immune thrombocytopenic purpura," *Ann Hematol* 87(11):899-904 (2008)), Behcet's Disease

(Yoshimura, T., et al., "Involvement of Th17 cells and the effect of anti-IL-6 therapy in autoimmune uveitis," *Rheumatology* (Oxford) 48(4):347-54 (2009)), Bullous pemphigoid (D'Auria, L., P. et al., "Cytokines and bullous pemphigoid," *Eur Cytokine Netw* 10(2):123-34 (1999)), Castleman's Disease (El-Osta, H.E. and R. Kurzrock, "Castleman's disease: from basic mechanisms to molecular therapeutics," *Oncologist* 16(4):497-511 (2011)), Celiac Disease (Lahdenpera, A.I., et al., "Up-regulation of small intestinal interleukin-17 immunity in untreated coeliac disease but not in potential coeliac disease or in type 1 diabetes," *Clin Exp Immunol* 167(2):226-34 (2012)), Churg-Strauss syndrome (Fujioka, A., et al., "The analysis of mRNA expression of cytokines from skin lesions in Churg-Strauss syndrome," *J Dermatol* 25(3):171-7 (1998)), Crohn's Disease (Holttä, V., et al., "IL-23/IL-17 immunity as a hallmark of Crohn's disease," *Inflamm Bowel Dis* 14(9):1175-84 (2008)), Cogan's syndrome (Shibuya, M., et al., "Successful treatment with tocilizumab in a case of Cogan's syndrome complicated with aortitis," *Mod Rheumatol* (2012)), Dry eye syndrome (De Paiva, C.S., et al., "IL-17 disrupts corneal barrier following desiccating stress," *Mucosal Immunol* 2(3):243-53 (2009)), Essential mixed cryoglobulinemia (Antonelli, A., et al., "Serum levels of proinflammatory cytokines interleukin-1beta, interleukin-6, and tumor necrosis factor alpha in mixed cryoglobulinemia," *Arthritis Rheum* 60(12):3841-7 (2009)), Dermatomyositis (Chevrel, G., et al., "Interleukin-17 increases the effects of IL-1 beta on muscle cells: arguments for the role of T cells in the pathogenesis of myositis," *J Neuroimmunol* 137(1-2):125-33 (2003)), Devic's Disease (Linhares, U.C., et al., "The Ex Vivo Production of IL-6 and IL-21 by CD4(+) T Cells is Directly Associated with Neurological Disability in Neuromyelitis Optica Patients," *J Clin Immunol* (2012)), Encephalitis (Kyburz, D. and M. Corr, "Th17 cells generated in the absence of TGF-beta induce experimental allergic encephalitis upon adoptive transfer," *Expert Rev Clin Immunol* 7(3):283-5 (2011)), Eosinophilic esophagitis (Dias, P.M. and G. Banerjee, "The Role of Th17/IL-17 on Eosinophilic Inflammation," *J Autoimmun* (2012)), Eosinophilic fasciitis (Dias, P.M. and G. Banerjee, "The Role of Th17/IL-17 on Eosinophilic Inflammation," *J Autoimmun* (2012)), Erythema nodosum (Kahawita, I.P. and D.N. Lockwood, "Towards understanding the pathology of erythema nodosum leprosum," *Trans R Soc Trop Med Hyg* 102(4):329-37 (2008)), Giant cell arteritis (Deng, J., et al., "Th17 and Th1 T-cell responses in giant cell arteritis," *Circulation* 121(7):906-15 (2010)), Glomerulonephritis (Ooi, J.D., et al., "Review: T helper 17 cells: their role in glomerulonephritis," *Nephrology* (Carlton) 15(5):513-21 (2010)), Goodpasture's syndrome (Ito, Y., et al., "Pathogenic significance of interleukin-6 in a patient with antglomerular basement membrane antibody-induced glomerulonephritis with multinucleated giant cells," *Am J Kidney Dis* 26(1):72-9 (1995)), Granulomatosis with Polyangiitis (Wegener's) (Nakahama, H., et al., "Distinct responses of interleukin-6 and other laboratory parameters to treatment in a patient with Wegener's granulomatosis," *Intern Med* 32(2):189-92 (1993)), Graves'

Disease (Kim, S.E., et al., "Increased serum interleukin-17 in Graves' ophthalmopathy," *Graefes Arch Clin Exp Ophthalmol* 250(10):1521-6 (2012)), Guillain-Barre syndrome (Lu, M.O. and J. Zhu, "The role of cytokines in Guillain-Barre syndrome," *J Neurol* 258(4):533-48 (2011)), Hashimoto's thyroiditis (Figuerola-Vega, N., et al., "Increased circulating pro-inflammatory cytokines and Th17 lymphocytes in Hashimoto's thyroiditis," *J Clin Endocrinol Metab* 95(2):953-62 (2009)), Hemolytic anemia (Xu, L., et al., "Critical role of Th17 cells in development of autoimmune hemolytic anemia," *Exp Hematol* (2012)), Henoch-Schonlein purpura (Jen, H.Y., et al., "Increased serum interleukin-17 and peripheral Th17 cells in children with acute Henoch-Schonlein purpura," *Pediatr Allergy Immunol* 22(8):862-8 (2011)), IgA nephropathy (Lin, F.J., et al., "Imbalance of regulatory T cells to Th17 cells in IgA nephropathy," *Scand J Clin Lab Invest* 72(3):221-9 (2012)), Inclusion body myositis (Baron, P., et al., "Production of IL-6 by human myoblasts stimulated with Abeta: relevance in the pathogenesis of IBM," *Neurology* 57(9):1561-5 (2001)), Type I diabetes (Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012)), Interstitial cystitis (Lamale, L.M., et al., "Interleukin-6, histamine, and methylhistamine as diagnostic markers for interstitial cystitis," *Urology* 68(4):702-6 (2006)), Kawasaki's Disease (Jia, S., et al., "The T helper type 17/regulatory T cell imbalance in patients with acute Kawasaki disease," *Clin Exp Immunol* 162(1):131-7 (2010)), Leukocytoclastic vasculitis (Min, C.K., et al., "Cutaneous leucoclastic vasculitis (LV) following bortezomib therapy in a myeloma patient; association with pro-inflammatory cytokines," *Eur J Haematol* 76(3):265-8 (2006)), Lichen planus (Rhodus, N.L., et al., "Proinflammatory cytokine levels in saliva before and after treatment of (erosive) oral lichen planus with dexamethasone," *Oral Dis* 12(2):112-6 (2006)), Lupus (SLE) (Mok, M.Y., et al., "The relation of interleukin 17 (IL-17) and IL-23 to Th1/Th2 cytokines and disease activity in systemic lupus erythematosus," *J Rheumatol* 37(10):2046-52 (2010)), Microscopic polyangitis (Muller Kobold, A.C., et al., "In vitro up-regulation of E-selectin and induction of interleukin-6 in endothelial cells by autoantibodies in Wegener's granulomatosis and microscopic polyangiitis," *Clin Exp Rheumatol* 17(4):433-40 (1999)), Multiple sclerosis (Jadidi-Niaragh, F. and Mirshafiey A., "Th17 cell, the new player of neuroinflammatory process in multiple sclerosis," *Scand J Immunol* 74(1):1-13 (2011)), Myasthenia gravis (Aricha, R., et al., "Blocking of IL-6 suppresses experimental autoimmune myasthenia gravis," *J Autoimmun* 36(2):135-41 (2011)), myositis (Chevrel, G., et al., "Interleukin-17 increases the effects of IL-1 beta on muscle cells: arguments for the role of T cells in the pathogenesis of myositis," *J Neuroimmunol* 137(1-2):125-33 (2003)), Optic neuritis (Icoz, S., et al., "Enhanced IL-6 production in aquaporin-4 antibody positive neuromyelitis optica patients," *Int J Neurosci* 120(1):71-5 (2010)), Pemphigus (Lopez-Robles, E., et al., "TNFalpha and IL-6 are mediators in the blistering process of pemphigus," *Int J Dermatol* 40(3):185-8 (2001)), POEMS syndrome

(Kallen, K.J., et al., "New developments in IL-6 dependent biology and therapy: where do we stand and what are the options?" *Expert Opin Investig Drugs* 8(9):1327-49 (1999)), Polyarteritis nodosa (Kawakami, T., et al., "Serum levels of interleukin-6 in patients with cutaneous polyarteritis nodosa," *Acta Derm Venereol* 92(3):322-3 (2012)), Primary biliary cirrhosis (Harada, K., et al., "Periductal interleukin-17 production in association with biliary innate immunity contributes to the pathogenesis of cholangiopathy in primary biliary cirrhosis," *Clin Exp Immunol* 157(2):261-70 (2009)), Psoriasis (Fujishima, S., et al., "Involvement of IL-17F via the induction of IL-6 in psoriasis," *Arch Dermatol Res* 302(7):499-505 (2010)), Psoriatic arthritis (Raychaudhuri, S.P., et al., IL-17 receptor and its functional significance in psoriatic arthritis," *Mol Cell Biochem* 359(1-2):419-29 (2012)), Pyoderma gangrenosum (Kawakami, T., et al., "Reduction of interleukin-6, interleukin-8, and anti-phosphatidylserine-prothrombin complex antibody by granulocyte and monocyte adsorption apheresis in a patient with pyoderma gangrenosum and ulcerative colitis," *Am J Gastroenterol* 104(9):2363-4 (2009)), Relapsing polychondritis (Kawai, M., et al., "Sustained response to tocilizumab, anti-interleukin-6 receptor antibody, in two patients with refractory relapsing polychondritis," *Rheumatology (Oxford)* 48(3):318-9 (2009)), Rheumatoid arthritis (Ash, Z. and P. Emery, "The role of tocilizumab in the management of rheumatoid arthritis," *Expert Opin Biol Ther*, 12(9):1277-89 (2012)), Sarcoidosis (Belli, F., et al., "Cytokines assay in peripheral blood and bronchoalveolar lavage in the diagnosis and staging of pulmonary granulomatous diseases," *Int J Immunopathol Pharmacol* 13(2):61-67 (2000)), Scleroderma (Radstake, T.R., et al., "The pronounced Th17 profile in systemic sclerosis (SSc) together with intracellular expression of TGFbeta and IFNgamma distinguishes SSc phenotypes," *PLoS One*, 4(6): e5903 (2009)), Sjogren's syndrome (Katsifis, G.E., et al., "Systemic and local interleukin-17 and linked cytokines associated with Sjogren's syndrome immunopathogenesis," *Am J Pathol* 175(3):1167-77 (2009)), Takayasu's arteritis (Sun, Y., et al., "MMP-9 and IL-6 are potential biomarkers for disease activity in Takayasu's arteritis," *Int J Cardiol* 156(2):236-8 (2012)), Transverse myelitis (Grabner, J.J., et al., "Interleukin-17 in transverse myelitis and multiple sclerosis," *J Neuroimmunol* 196(1-2):124-32 (2008)), Ulcerative colitis (Mudter, J. and M.F. Neurath, "IL-6 signaling in inflammatory bowel disease: pathophysiological role and clinical relevance," *Inflamm Bowel Dis* 13(8):1016-23 (2007)), Uveitis (Haruta, H., et al., "Blockade of interleukin-6 signaling suppresses not only th17 but also interphotoreceptor retinoid binding protein-specific Th1 by promoting regulatory T cells in experimental autoimmune uveoretinitis," *Invest Ophthalmol Vis Sci* 52(6):3264-71 (2011)), and Vitiligo (Bassiouny, D.A. and O. Shaker, "Role of interleukin-17 in the pathogenesis of vitiligo," *Clin Exp Dermatol* 36(3):292-7 115. (2011)). Thus, the invention includes compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof; pharmaceutical compositions

comprising one or more of those compounds; and methods of using those compounds or compositions for treating these diseases.

[0156] Acute and chronic (non-autoimmune) inflammatory diseases characterized by increased expression of pro-inflammatory cytokines, including IL-6, MCP-1, and IL-17, would also be amenable to therapeutic BET inhibition. These include, but are not limited to, sinusitis (Bradley, D.T. and S.E. Kountakis, "Role of interleukins and transforming growth factor-beta in chronic rhinosinusitis and nasal polyposis," *Laryngoscope* 115(4):684-6 (2005)), pneumonitis (Besnard, A.G., et al., "Inflammasome-IL-1-Th17 response in allergic lung inflammation" *J Mol Cell Biol* 4(1):3-10 (2012)), osteomyelitis (Yoshii, T., et al., "Local levels of interleukin-1beta, -4, -6 and tumor necrosis factor alpha in an experimental model of murine osteomyelitis due to staphylococcus aureus," *Cytokine* 19(2):59-65 2002), gastritis (Bayraktaroglu, T., et al., "Serum levels of tumor necrosis factor-alpha, interleukin-6 and interleukin-8 are not increased in dyspeptic patients with Helicobacter pylori-associated gastritis," *Mediators Inflamm* 13(1):25-8 (2004)), enteritis (Mitsuyama, K., et al., "STAT3 activation via interleukin 6 trans-signalling contributes to ileitis in SAMP1/Yit mice," *Gut* 55(9):1263-9. (2006)), gingivitis (Johnson, R.B., et al., "Interleukin-11 and IL-17 and the pathogenesis of periodontal disease," *J Periodontol* 75(1):37-43 (2004)), appendicitis (Latifi, S.Q., et al., "Persistent elevation of serum interleukin-6 in intraabdominal sepsis identifies those with prolonged length of stay," *J Pediatr Surg* 39(10):1548-52 (2004)), irritable bowel syndrome (Ortiz-Lucas, M., et al., "Irritable bowel syndrome immune hypothesis. Part two: the role of cytokines," *Rev Esp Enferm Dig* 102(12):711-7 (2010)), tissue graft rejection (Kappel, L.W., et al., "IL-17 contributes to CD4-mediated graft-versus-host disease," *Blood* 113(4):945-52 (2009)), chronic obstructive pulmonary disease (COPD) (Traves, S.L. and L.E. Donnelly, "Th17 cells in airway diseases," *Curr Mol Med* 8(5):416-26 (2008)), septic shock (toxic shock syndrome, SIRS, bacterial sepsis, etc) (Nicodeme, E., et al., "Suppression of inflammation by a synthetic histone mimic," *Nature* 468(7327):1119-23 (2010)), osteoarthritis (Chen, L., et al., "IL-17RA aptamer-mediated repression of IL-6 inhibits synovium inflammation in a murine model of osteoarthritis," *Osteoarthritis Cartilage* 19(6):711-8 (2011)), acute gout (Urano, W., et al., "The inflammatory process in the mechanism of decreased serum uric acid concentrations during acute gouty arthritis," *J Rheumatol* 29(9):1950-3 (2002)), acute lung injury (Traves, S.L. and L.E. Donnelly, "Th17 cells in airway diseases," *Curr Mol Med* 8(5):416-26 (2008)), acute renal failure (Simmons, E.M., et al., "Plasma cytokine levels predict mortality in patients with acute renal failure," *Kidney Int* 65(4):1357-65 (2004)), burns (Paquet, P. and G.E. Pierard, "Interleukin-6 and the skin," *Int Arch Allergy Immunol* 109(4):308-17 (1996)), Herxheimer reaction (Kaplanski, G., et al., "Jarisch-Herxheimer reaction complicating the treatment of chronic Q fever endocarditis: elevated TNFalpha and IL-6 serum levels," *J Infect* 37(1):83-4 (1998)), and SIRS

associated with viral infections (Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012)). Thus, the invention includes compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof; pharmaceutical compositions comprising one or more of those compounds; and methods of using those compounds or compositions for treating these diseases.

[0157] In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used for treating rheumatoid arthritis (RA) and multiple sclerosis (MS). Strong proprietary data exist for the utility of BET inhibitors in preclinical models of RA and MS. R. Jahagirdar, S.M. et al., "An Orally Bioavailable Small Molecule RVX-297 Significantly Decreases Disease in a Mouse Model of Multiple Sclerosis," *World Congress of Inflammation*, Paris, France (2011). Both RA and MS are characterized by a dysregulation of the IL-6 and IL-17 inflammatory pathways (Kimura, A. and T. Kishimoto, "IL-6: regulator of Treg/Th17 balance," *Eur J Immunol* 40(7):1830-5 (2010)) and thus would be especially sensitive to BET inhibition. In another embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II may be used for treating sepsis and associated afflictions. BET inhibition has been shown to inhibit development of sepsis, in part, by inhibiting IL-6 expression, in preclinical models in both published (Nicodeme, E., et al., "Suppression of inflammation by a synthetic histone mimic," *Nature* 468(7327):1119-23 (2010)) and proprietary data.

[0158] In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat cancer. Cancers that have an overexpression, translocation, amplification, or rearrangement c-myc or other myc family oncoproteins (MYCN, L-myc) are particularly sensitive to BET inhibition. Delmore, J.E., et al., "BET bromodomain inhibition as a therapeutic strategy to target c-Myc," *Cell* 146(6):904-17 (2010); Mertz, J.A., et al., "Targeting MYC dependence in cancer by inhibiting BET bromodomains," *Proc Natl Acad Sci USA* 108(40):16669-74 (2011). These cancers include, but are not limited to, B-acute lymphocytic leukemia, Burkitt's lymphoma, Diffuse large cell lymphoma, Multiple myeloma, Primary plasma cell leukemia, Atypical carcinoid lung cancer, Bladder cancer, Breast cancer, Cervix cancer, Colon cancer, Gastric cancer, Glioblastoma, Hepatocellular carcinoma, Large cell neuroendocrine carcinoma, Medulloblastoma, Melanoma, nodular, Melanoma, superficial spreading, Neuroblastoma, esophageal squamous cell carcinoma, Osteosarcoma, Ovarian cancer, Prostate cancer, Renal clear cell carcinoma, Retinoblastoma, Rhabdomyosarcoma, and Small cell lung

carcinoma. Vita, M. and M. Henriksson, "The Myc oncoprotein as a therapeutic target for human cancer," *Semin Cancer Biol* 16(4):318-30 (2006).

[0159] In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat cancers that result from an aberrant regulation (overexpression, translocation, etc) of BET proteins. These include, but are not limited to, NUT midline carcinoma (Brd3 or Brd4 translocation to nutlin 1 gene) (French, C.A., "NUT midline carcinoma," *Cancer Genet Cytogenet* 203(1):16-20 (2010)), B-cell lymphoma (Brd2 overexpression) (Greenwald, R.J., et al., "E mu-BRD2 transgenic mice develop B-cell lymphoma and leukemia," *Blood* 103(4):1475-84 (2004)), non-small cell lung cancer (BrdT overexpression) (Grunwald, C., et al., "Expression of multiple epigenetically regulated cancer/germline genes in nonsmall cell lung cancer," *Int J Cancer* 118(10):2522-8 (2006)), esophageal cancer and head and neck squamous cell carcinoma (BrdT overexpression) (Scanlan, M.J., et al., "Expression of cancer-testis antigens in lung cancer: definition of bromodomain testis-specific gene (BRDT) as a new CT gene, CT9," *Cancer Lett* 150(2):55-64 (2000)), and colon cancer (Brd4) (Rodriguez, R.M., et al., "Aberrant epigenetic regulation of bromodomain BRD4 in human colon cancer," *J Mol Med (Berl)* 90(5):587-95 (2012)).

[0160] In one embodiment, because BET inhibitors decrease Brd-dependent recruitment of pTEFb to genes involved in cell proliferation, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat cancers that rely on pTEFb (Cdk9/cyclin T) and BET proteins to regulate oncogenes. These cancers include, but are not limited to, chronic lymphocytic leukemia and multiple myeloma (Tong, W.G., et al., "Phase I and pharmacologic study of SNS-032, a potent and selective Cdk2, 7, and 9 inhibitor, in patients with advanced chronic lymphocytic leukemia and multiple myeloma," *J Clin Oncol* 28(18):3015-22 (2010)), follicular lymphoma, diffuse large B cell lymphoma with germinal center phenotype, Burkitt's lymphoma, Hodgkin's lymphoma, follicular lymphomas and activated, anaplastic large cell lymphoma (Bellan, C., et al., "CDK9/CYCLIN T1 expression during normal lymphoid differentiation and malignant transformation," *J Pathol* 203(4):946-52 (2004)), neuroblastoma and primary neuroectodermal tumor (De Falco, G., et al., "Cdk9 regulates neural differentiation and its expression correlates with the differentiation grade of neuroblastoma and PNET tumors," *Cancer Biol Ther* 4(3):277-81 (2005)), rhabdomyosarcoma (Simone, C. and A. Giordano, "Abrogation of signal-dependent activation of the cdk9/cyclin T2a complex in human RD rhabdomyosarcoma cells," *Cell Death Differ* 14(1):192-5 (2007)), prostate cancer (Lee, D.K., et al., "Androgen receptor interacts

with the positive elongation factor P-TEFb and enhances the efficiency of transcriptional elongation," *J Biol Chem* 276(13):9978-84 (2001)), and breast cancer (Bartholomeeusen, K., et al., "BET bromodomain inhibition activates transcription via a transient release of P-TEFb from 7SK snRNP," *J Biol Chem* (2012)).

[0161] In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat cancers in which BET-responsive genes, such as CDK6, Bcl2, TYRO3, MYB, and hTERT are up-regulated. Dawson, M.A., et al., "Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia," *Nature* 478(7370):529-33 (2011); Delmore, J.E., et al., "BET bromodomain inhibition as a therapeutic strategy to target c-Myc," *Cell* 146(6):904-17 (2010). These cancers include, but are not limited to, pancreatic cancer, breast cancer, colon cancer, glioblastoma, adenoid cystic carcinoma, T-cell prolymphocytic leukemia, malignant glioma, bladder cancer, medulloblastoma, thyroid cancer, melanoma, multiple myeloma, Barret's adenocarcinoma, hepatoma, prostate cancer, pro-myelocytic leukemia, chronic lymphocytic leukemia, mantle cell lymphoma, diffuse large B-cell lymphoma, small cell lung cancer, and renal carcinoma. Ruden, M. and N. Puri, "Novel anticancer therapeutics targeting telomerase," *Cancer Treat Rev* (2012); Kelly, P.N. and A. Strasser, "The role of Bcl-2 and its pro-survival relatives in tumorigenesis and cancer therapy" *Cell Death Differ* 18(9):1414-24 (2011); Uchida, T., et al., "Antitumor effect of bcl-2 antisense phosphorothioate oligodeoxynucleotides on human renal-cell carcinoma cells in vitro and in mice," *Mol Urol* 5(2):71-8 (2001).

[0162] Published and proprietary data have shown direct effects of BET inhibition on cell proliferation in various cancers. In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat cancers for which exist published and, for some, proprietary, *in vivo* and/or *in vitro* data showing a direct effect of BET inhibition on cell proliferation. These cancers include NMC (NUT-midline carcinoma), acute myeloid leukemia (AML), acute B lymphoblastic leukemia (B-ALL), Burkitt's Lymphoma, B-cell Lymphoma, Melanoma, mixed lineage leukemia, multiple myeloma, pro-myelocytic leukemia (PML), and non-Hodgkin's lymphoma. Filippakopoulos, P., et al., "Selective inhibition of BET bromodomains," *Nature* 468(7327):1067-73 (2010); Dawson, M.A., et al., "Inhibition of BET recruitment to chromatin as an effective treatment for MLL-fusion leukaemia," *Nature* 478(7370):529-33 (2011); Zuber, J., et al., "RNAi screen identifies Brd4 as a therapeutic target in acute myeloid leukaemia," *Nature* 478(7370):524-8 (2011); Miguel F. Segura, et al, "BRD4 is a novel

therapeutic target in melanoma," *Cancer Research*. 72(8):Supplement 1 (2012). The compounds of the invention have a demonstrated BET inhibition effect on cell proliferation *in vitro* for the following cancers: Neuroblastoma, Medulloblastoma, lung carcinoma (NSCLC, SCLC), and colon carcinoma.

[0163] In one embodiment, because of potential synergy or additive effects between BET inhibitors and other cancer therapy, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be combined with other therapies, chemotherapeutic agents, or anti-proliferative agents to treat human cancer and other proliferative disorders. The list of therapeutic agents which can be combined with BET inhibitors in cancer treatment includes, but is not limited to, ABT-737, Azacitidine (Vidaza)TM, AZD1152 (Barasertib), AZD2281 (Olaparib), AZD6244 (Selumetinib), BEZ235, Bleomycin Sulfate, Bortezomib (Velcade)TM, Busulfan (Myleran)TM, Camptothecin, Cisplatin, Cyclophosphamide (Clafen)TM, CYT387, Cytarabine (Ara-C), Dacarbazine, DAPT (GSI-IX), Decitabine, Dexamethasone, Doxorubicin (Adriamycin)TM, Etoposide, Everolimus (RAD001), Flavopiridol (Alvocidib), Ganetespib (STA-9090)TM, Gefitinib (Iressa)TM, Idarubicin, Ifosfamide (Mitoxana), IFNa2a (Roferon A)TM, Melphalan (Alkeran)TM, Methazolastone (temozolomide), Metformin, Mitoxantrone (Novantrone)TM, Paclitaxel, Phenformin, PKC412 (Midostaurin), PLX4032 (Vemurafenib), Pomalidomide (CC-4047), Prednisone (Deltasone)TM, Rapamycin, RevlimidTM (Lenalidomide), Ruxolitinib (INCB018424), Sorafenib (Nexavar)TM, SU11248 (Sunitinib), SU11274, Vinblastine, Vincristine (Oncovin)TM, Vinorelbine (Navelbine)TM, Vorinostat (SAHA), and WP1130 (Degrasyn).

[0164] In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat benign proliferative and fibrotic disorders, including, but not limited to, benign soft tissue tumors, bone tumors, brain and spinal tumors, eyelid and orbital tumors, granuloma, lipoma, meningioma, multiple endocrine neoplasia, nasal polyps, pituitary tumors, prolactinoma, pseudotumor cerebri, seborrheic keratoses, stomach polyps, thyroid nodules, cystic neoplasms of the pancreas, hemangiomas, vocal cord nodules, polyps, and cysts, Castleman disease, chronic pilonidal disease, dermatofibroma, pilar cyst, pyogenic granuloma, juvenile polyposis syndrome, idiopathic pulmonary fibrosis, renal fibrosis, post-operative stricture, keloid formation, scleroderma, and cardiac fibrosis. Tang, X et al., "Assessment of Brd4 Inhibition in Idiopathic Pulmonary Fibrosis Lung Fibroblasts and in Vivo Models of Lung Fibrosis," *Am J Pathology* in press (2013).

[0165] In one embodiment, because of their ability to up-regulate ApoA-1 transcription and protein expression (Mirguet, O., et al., "From ApoA1 upregulation to BET family bromodomain

inhibition: discovery of I-BET151," *Bioorg Med Chem Lett* 22(8):2963-7 (2012); Chung, C.W., et al., "Discovery and characterization of small molecule inhibitors of the BET family bromodomains," *J Med Chem* 54(11):3827-38 (2011)), BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat cardiovascular diseases that are generally associated with including dyslipidemia, atherosclerosis, hypercholesterolemia, and metabolic syndrome (Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012); Denis, G.V., "Bromodomain coactivators in cancer, obesity, type 2 diabetes, and inflammation," *Discov Med* 10(55):489-99 (2010)). In another embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II may be used to treat non-cardiovascular disease characterized by deficits in ApoA-1, including Alzheimer's disease. Elliott, D.A., et al., "Apolipoproteins in the brain: implications for neurological and psychiatric disorders," *Clin Lipidol* 51(4):555-573 (2010).

[0166] In one embodiment, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used in patients with insulin resistance and type II diabetes. Belkina, A.C. and G.V. Denis, "BET domain co-regulators in obesity, inflammation and cancer," *Nat Rev Cancer* 12(7):465-77 (2012); Denis, G.V., "Bromodomain coactivators in cancer, obesity, type 2 diabetes, and inflammation," *Discov Med* 10(55):489-99 (2010); Wang, F., et al., "Brd2 disruption in mice causes severe obesity without Type 2 diabetes," *Biochem J* 425(1):71-83 (2010); Denis, G.V., et al., "An emerging role for bromodomain-containing proteins in chromatin regulation and transcriptional control of adipogenesis," *FEBS Lett* 584(15):3260-8 (2010). The anti-inflammatory effects of BET inhibition would have additional value in decreasing inflammation associated with diabetes and metabolic disease. Alexandraki, K., et al., "Inflammatory process in type 2 diabetes: The role of cytokines," *Ann N Y Acad Sci* 1084:89-117 (2006).

[0167] In one embodiment, because of their ability to down-regulate viral promoters, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used as therapeutics for cancers that are associated with viruses including Epstein-Barr Virus (EBV), hepatitis virus (HBV, HCV), Kaposi's sarcoma associated virus (KSHV), human papilloma virus (HPV), Merkel cell polyomavirus, and human cytomegalovirus (CMV). Gagnon, D., et al., "Proteasomal degradation of the papillomavirus E2 protein is inhibited by overexpression of bromodomain-containing protein 4," *J Virol* 83(9):4127-39 (2009); You, J., et al.,

"Kaposi's sarcoma-associated herpesvirus latency-associated nuclear antigen interacts with bromodomain protein Brd4 on host mitotic chromosomes," *J Virol* 80(18):8909-19 (2006); Palermo, R.D., et al., "RNA polymerase II stalling promotes nucleosome occlusion and pTEFb recruitment to drive immortalization by Epstein-Barr virus," *PLoS Pathog* 7(10):e1002334 (2011); Poreba, E., et al., "Epigenetic mechanisms in virus-induced tumorigenesis," *Clin Epigenetics* 2(2):233-47. 2011. In another embodiment, because of their ability to reactivate HIV-1 in models of latent T cell infection and latent monocyte infection, BET inhibitors could be used in combination with anti-retroviral therapeutics for treating HIV. Zhu, J., et al., "Reactivation of Latent HIV-1 by Inhibition of BRD4," *Cell Rep* (2012); Banerjee, C., et al., "BET bromodomain inhibition as a novel strategy for reactivation of HIV-1," *J Leukoc Biol* (2012); Bartholomeeusen, K., et al., "BET bromodomain inhibition activates transcription via a transient release of P-TEFb from 7SK snRNP," *J Biol Chem* (2012); Li, Z., et al., "The BET bromodomain inhibitor JQ1 activates HIV latency through antagonizing Brd4 inhibition of Tat-transactivation," *Nucleic Acids Res* (2012.)

[0168] In one embodiment, because of the role of epigenetic processes and bromodomain-containing proteins in neurological disorders, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used to treat diseases including, but not limited to, Alzheimer's disease, Parkinson's disease, Huntington disease, bipolar disorder, schizophrenia, Rubinstein-Taybi syndrome, and epilepsy. Prinjha, R.K., J. Witherington, and K. Lee, "Place your BETs: the therapeutic potential of bromodomains," *Trends Pharmacol Sci* 33(3):146-53 (2012); Muller, S., et al., "Bromodomains as therapeutic targets," *Expert Rev Mol Med* 13:e29 (2011).

[0169] In one embodiment, because of the effect of BRDT depletion or inhibition on spermatid development, BET inhibitor compounds of Formula I, Formula Ia, and/or Formula II, stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof, or compositions comprising one or more of those compounds may be used as reversible, male contraceptive agents. Matzuk, M.M., et al., "Small-Molecule Inhibition of BRDT for Male Contraception," *Cell* 150(4): p. 673-684 (2012); Berkovits, B.D., et al., "The testis-specific double bromodomain-containing protein BRDT forms a complex with multiple spliceosome components and is required for mRNA splicing and 3'-UTR truncation in round spermatids," *Nucleic Acids Res* 40(15):7162-75 (2012).

Pharmaceutical Compositions

[0170] Pharmaceutical compositions of the present disclosure comprise at least one compound of Formulae I-II, or tautomer, stereoisomer, pharmaceutically acceptable salt or hydrate thereof formulated together with one or more pharmaceutically acceptable carriers. These

formulations include those suitable for oral, rectal, topical, buccal and parenteral (e.g., subcutaneous, intramuscular, intradermal, or intravenous) administration. The most suitable form of administration in any given case will depend on the degree and severity of the condition being treated and on the nature of the particular compound being used.

[0171] Formulations suitable for oral administration may be presented in discrete units, such as capsules, cachets, lozenges, or tablets, each containing a predetermined amount of a compound of the present disclosure as powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil emulsion. As indicated, such formulations may be prepared by any suitable method of pharmacy which includes the step of bringing into association at least one compound of the present disclosure as the active compound and a carrier or excipient (which may constitute one or more accessory ingredients). The carrier must be acceptable in the sense of being compatible with the other ingredients of the formulation and must not be deleterious to the recipient. The carrier may be a solid or a liquid, or both, and may be formulated with at least one compound described herein as the active compound in a unit-dose formulation, for example, a tablet, which may contain from about 0.05% to about 95% by weight of the at least one active compound. Other pharmacologically active substances may also be present including other compounds. The formulations of the present disclosure may be prepared by any of the well-known techniques of pharmacy consisting essentially of admixing the components.

[0172] For solid compositions, conventional nontoxic solid carriers include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talc, cellulose, glucose, sucrose, magnesium carbonate, and the like. Liquid pharmacologically administrable compositions can, for example, be prepared by, for example, dissolving or dispersing, at least one active compound of the present disclosure as described herein and optional pharmaceutical adjuvants in an excipient, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to thereby form a solution or suspension. In general, suitable formulations may be prepared by uniformly and intimately admixing the at least one active compound of the present disclosure with a liquid or finely divided solid carrier, or both, and then, if necessary, shaping the product. For example, a tablet may be prepared by compressing or molding a powder or granules of at least one compound of the present disclosure, which may be optionally combined with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, at least one compound of the present disclosure in a free-flowing form, such as a powder or granules, which may be optionally mixed with a binder, lubricant, inert diluent and/or surface active/dispersing agent(s). Molded tablets may be made by molding, in a

suitable machine, where the powdered form of at least one compound of the present disclosure is moistened with an inert liquid diluent.

[0173] Formulations suitable for buccal (sub-lingual) administration include lozenges comprising at least one compound of the present disclosure in a flavored base, usually sucrose and acacia or tragacanth, and pastilles comprising the at least one compound in an inert base such as gelatin and glycerin or sucrose and acacia.

[0174] Formulations of the present disclosure suitable for parenteral administration comprise sterile aqueous preparations of at least one compound of Formulae I-II or tautomers, stereoisomers, pharmaceutically acceptable salts, and hydrates thereof, which are approximately isotonic with the blood of the intended recipient. These preparations are administered intravenously, although administration may also be effected by means of subcutaneous, intramuscular, or intradermal injection. Such preparations may conveniently be prepared by admixing at least one compound described herein with water and rendering the resulting solution sterile and isotonic with the blood. Injectable compositions according to the present disclosure may contain from about 0.1 to about 5% w/w of the active compound.

[0175] Formulations suitable for rectal administration are presented as unit-dose suppositories. These may be prepared by admixing at least one compound as described herein with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

[0176] Formulations suitable for topical application to the skin may take the form of an ointment, cream, lotion, paste, gel, spray, aerosol, or oil. Carriers and excipients which may be used include Vaseline, lanoline, polyethylene glycols, alcohols, and combinations of two or more thereof. The active compound (i.e., at least one compound of Formulae I-IV or tautomers, stereoisomers, pharmaceutically acceptable salts, and hydrates thereof) is generally present at a concentration of from about 0.1% to about 15% w/w of the composition, for example, from about 0.5 to about 2%.

[0177] The amount of active compound administered may be dependent on the subject being treated, the subject's weight, the manner of administration and the judgment of the prescribing physician. For example, a dosing schedule may involve the daily or semi-daily administration of the encapsulated compound at a perceived dosage of about 1 μ g to about 1000 mg. In another embodiment, intermittent administration, such as on a monthly or yearly basis, of a dose of the encapsulated compound may be employed. Encapsulation facilitates access to the site of action and allows the administration of the active ingredients simultaneously, in theory producing a synergistic effect. In accordance with standard dosing regimens, physicians will readily determine optimum dosages and will be able to readily modify administration to achieve such dosages.

[0178] A therapeutically effective amount of a compound or composition disclosed herein can be measured by the therapeutic effectiveness of the compound. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being used. In one embodiment, the therapeutically effective amount of a disclosed compound is sufficient to establish a maximal plasma concentration. Preliminary doses as, for example, determined according to animal tests, and the scaling of dosages for human administration is performed according to art-accepted practices.

[0179] Toxicity and therapeutic efficacy can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compositions that exhibit large therapeutic indices are preferable.

[0180] Data obtained from the cell culture assays or animal studies can be used in formulating a range of dosage for use in humans. Therapeutically effective dosages achieved in one animal model may be converted for use in another animal, including humans, using conversion factors known in the art (see, e.g., Freireich et al., *Cancer Chemother. Reports* 50(4):219-244 (1966) and Table I for Equivalent Surface Area Dosage Factors).

Table I: Equivalent Surface Area Dosage Factors:

To: From:	Mouse (20 g)	Rat (150 g)	Monkey (3.5 kg)	Dog (8 kg)	Human (60 kg)
Mouse	1	1/2	1/4	1/6	1/12
Rat	2	1	1/2	1/4	1/7
Monkey	4	2	1	3/5	1/3
Dog	6	4	3/5	1	1/2
Human	12	7	3	2	1

[0181] The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. Generally, a therapeutically effective amount may vary with the subject's age, condition, and gender, as well as the severity of the medical condition in the subject. The dosage may be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment.

[0182] In one embodiment, a compound of Formulae I-II or a tautomer, stereoisomer, pharmaceutically acceptable salt or hydrate thereof, is administered in combination with another therapeutic agent. The other therapeutic agent can provide additive or synergistic value relative to the administration of a compound of the present disclosure alone. The therapeutic agent can be, for example, a statin; a PPAR agonist, e.g., a thiazolidinedione or fibrate; a niacin, a RVX, FXR or LXR agonist; a bile-acid reuptake inhibitor; a cholesterol absorption inhibitor; a cholesterol synthesis inhibitor; a cholesteryl ester transfer protein (CETP), an ion-exchange resin; an antioxidant; an inhibitor of AcylCoA cholesterol acyltransferase (ACAT inhibitor); a tyroprohostine; a sulfonylurea-based drug; a biguanide; an alpha-glucosidase inhibitor; an apolipoprotein E regulator; a HMG-CoA reductase inhibitor, a microsomal triglyceride transfer protein; an LDL-lowering drug; an HDL-raising drug; an HDL enhancer; a regulator of the apolipoprotein A-IV and/or apolipoprotein genes; or any cardiovascular drug.

[0183] In another embodiment, a compound of Formulae I and/or Formula II or a tautomer, stereoisomer, pharmaceutically acceptable salt or hydrate thereof, is administered in combination with one or more anti-inflammatory agents. Anti-inflammatory agents can include immunosuppressants, TNF inhibitors, corticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs), disease-modifying anti-rheumatic drugs (DMARDs), and the like. Exemplary anti-inflammatory agents include, for example, prednisone; methylprednisolone (Medrol®), triamcinolone, methotrexate (Rheumatrex®, Trexall®), hydroxychloroquine (Plaquenil®), sulfasalazine (Azulfidine®), leflunomide (Arava®), etanercept (Enbrel®), infliximab (Remicade®), adalimumab (Humira®), rituximab (Rituxan®), abatacept (Orencia®), interleukin—1, anakinra (Kineret™), ibuprofen, ketoprofen, fenoprofen, naproxen, aspirin, acetaminophen, indomethacin, sulindac, meloxicam, piroxicam, tenoxicam, lornoxicam, ketorolac, etodolac, mefenamic acid, meclofenamic acid, flufenamic acid, tolfenamic acid, diclofenac, oxaprozin, apazone, nimesulide, nabumetone, tenidap, etanercept, tolmetin, phenylbutazone, oxyphenbutazone, diflunisal, salsalate, olsalazine, or sulfasalazine.

EXAMPLES


[0184] General Methods. Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Proton nuclear magnetic resonance spectra were obtained on a Bruker AVANCE 300 spectrometer at 300 MHz or Bruker AVANCE 500 spectrometer at 500 MHz or a Bruker AVANCE 300 spectrometer at 300 MHz. Spectra are given in ppm (δ) and coupling constants, *J* values, are reported in hertz (Hz). Tetramethylsilane was used as an internal standard for ¹H nuclear magnetic resonance. Mass spectra analyses were performed on Waters Aquity UPLC Mass Spectrometer in ESI or APCI mode when appropriate, Agilent 6130A Mass Spectrometer in ESI,

APCI, or MultiMode mode when appropriate or Applied Biosystems API-150EX Spectrometer in ESI or APCI mode when appropriate. Silica gel chromatography was in general performed on a Teledyne Isco CombiFlash® Rf 200 system or a Teledyne Isco CombiFlash® Companion system.

[0185] Abbreviations: CDI: 1,1'-carbonyldiimidazole; DMAP: *N,N*-dimethylaminopropylamine; EDC: 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride; *m*-CPBA: 3-chloroperoxybenzoic acid; NBS: *N*-bromosuccinimide.

General Procedure A:


Preparation of 9-Benzyl-2-(3,5-dimethylisoxazol-4-yl)-9H-purin-6-amine (Example Compound 1)



[0186] Step 1: To a slurry of **1** (1.50 g, 8.84 mmol) in DMF (50 mL) was added potassium carbonate (3.64 g, 26.4 mmol) and benzyl chloride (1.01 mL, 8.84 mmol). The reaction was stirred at rt for 16 h. The reaction mixture was filtered, the filtrate was poured into water (100 mL) and stirred for 5 minutes. The solid was collected and dried to give **2** (1.60 g, 70%) as a yellow solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.26 (s, 1H), 7.80 (br s, 2H), 7.38–7.26 (m, 5H), 5.34 (s, 2H); ESI m/z 260 $[\text{M} + \text{H}]^+$.

[0187] Step 2: To a solution of **2** (260 mg, 1.0 mmol) in 1,4-dioxane (10 mL) and DMF (4 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (335 mg, 1.5 mmol), sodium carbonate (2.0 M in H_2O , 1.0 mL, 2.0 mmol) and tetrakis(triphenylphosphine)palladium(0) (116 mg, 0.1 mmol). The reaction mixture was purged with nitrogen and heated at 80 °C for 16 h. The mixture was diluted with methylene chloride (20 mL) and filtered. The filtrate was concentrated and purified by chromatography (silica gel, 0–5% methylene chloride/methanol) followed by trituration with EtOAc/hexanes to afford 9-benzyl-2-(3,5-dimethylisoxazol-4-yl)-9H-purin-6-amine (**Example Compound 1**) (110 mg, 34%) as a white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.29 (s, 1H), 7.36–7.28 (m, 7H), 5.38 (s, 2H), 2.73 (s, 3H), 2.51 (s, 3H); ESI m/z 321 $[\text{M} + \text{H}]^+$.

Preparation of 3-Benzyl-5-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one
(Example Compound 2)




[0188] Step 1: To a solution of **4** (500 mg, 2.66 mmol) in 1,4-dioxane (15 mL) was added CDI (517 mg, 3.19 mmol). The reaction was heated at 60 °C for 16 h. The solid was collected and dried to give **5** (340 mg, 60%) as a light purple solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.58 (br s, 1H), 11.02 (br s, 1H), 7.19 (d, *J* = 8.1 Hz, 1H), 7.13 (d, *J* = 8.1 Hz, 1H).

[0189] **Step 2:** To a solution of **5** (170 mg, 0.79 mmol) in 1,4-dioxane (12 mL) and DMF (6 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (352 mg, 1.58 mmol), sodium carbonate (2.0 M in H₂O, 1.19 mL, 2.37 mmol) and tetrakis(triphenylphosphine) palladium(0) (92 mg, 0.08 mmol). The reaction mixture was purged with nitrogen and heated at 80 °C for 16 h. The mixture was diluted with methylene chloride (20 mL) and filtered. The filtrate was concentrated and purified by chromatography (silica gel, 0–5% methylene chloride/methanol) to afford **6** (130 mg, 71%) as a white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.38 (br s, 1H), 10.90 (br s, 1H), 7.30 (d, *J* = 7.8 Hz, 1H), 7.07 (d, *J* = 8.1 Hz, 1H), 2.49 (s, 3H), 2.33 (s, 3H).

[0190] Step 3: To a solution of **6** (100 mg, 0.43 mmol) in DMF (10 mL) was added potassium carbonate (72 mg, 0.52 mmol) and di-*tert*-butyl dicarbonate (104 mg, 0.48 mmol). The reaction was stirred at rt for 16 h. To the reaction mixture was added potassium carbonate (72 mg, 0.52 mmol) and benzyl chloride (0.14 mL, 0.48 mmol). The reaction was stirred at rt for 16 h. The mixture was diluted with EtOAc (100 mL) and washed with brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–30% ethyl acetate/hexanes) afforded **6** (130 mg, 71%) as a colorless gum: ^1H NMR (300 MHz, DMSO- d_6) δ 7.97 (d, J = 8.1 Hz, 1H), 7.38–7.27 (m, 6H), 5.05 (s, 2H), 2.49 (s, 3H), 2.29 (s, 3H), 1.61 (s, 9H).

[0191] Step 4: A solution of **7** (130 mg, 0.31 mmol) in methylene chloride (4 mL) and TFA (2 mL) was stirred at rt for 2 h. The mixture was concentrated, the residue was dissolved in methylene chloride (100 mL), washed with saturated NaHCO₃ (50 mL × 2) and brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated to afford 3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (**Example Compound 2**) (81 mg, 81%) as an off-white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.31 (s, 1H), 7.40 (d, *J* = 7.8 Hz, 1H), 7.34–7.25 (m, 5 H), 7.15 (d, *J* = 7.8 Hz, 1H), 5.03 (s, 2H), 2.47 (s, 3H), 2.28 (s, 3H); ESI *m/z* 321 [M + H]⁺.

Preparation of 1-Benzyl-5-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (Example Compound 3)




[0192] Step 1: To a solution of **4** (500 mg, 2.66 mmol) and benzaldehyde (282 mg, 2.66 mmol) in methylene chloride (15 mL) was added acetic acid (319 mg, 5.32 mmol). The reaction was stirred at rt for 30 minutes, then NaBH(OAc)₃ (1.69 g, 7.98 mmol) was added. The reaction mixture was stirred at rt for 16 h. The mixture was diluted with methylene chloride (100 mL) and saturated aq. NaHCO₃ (50 mL) was added slowly. The organic layer was separated, dried over sodium sulfate, filtered and concentrated. The residue was triturated with methylene chloride/EtOAc to give **8** (401 mg, 54%) as a light brown solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.34–7.22 (m, 5H), 6.48 (d, *J* = 7.8 Hz, 1H), 6.40 (d, *J* = 7.8 Hz, 1H), 6.02 (br s, 2H), 5.54 (t, *J* = 5.7 Hz, 1H), 4.27 (d, *J* = 5.4 Hz, 2H).

[0193] Step 2: To a solution of **8** (400 mg, 1.44 mmol) in 1,4-dioxane (10 mL) was added CDI (514 mg, 3.17 mmol). The reaction was heated at 110 °C for 16 h. The reaction mixture was concentrated. Purification by chromatography (silica gel, 0–40% ethyl acetate/hexanes) afforded **9** (310 mg, 71%) as a white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.96 (s, 1H), 7.35–7.27 (m, 6H), 7.19 (d, *J* = 7.8 Hz, 1H), 5.02 (s, 2H).

[0194] Step 3: To a solution of **9** (310 mg, 1.02 mmol) in 1,4-dioxane (10 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (341 mg, 1.53 mmol), sodium carbonate (2.0 M in H₂O, 1.02 mL, 2.04 mmol) and tetrakis(triphenylphosphine)palladium(0) (59 mg, 0.05 mmol). The reaction mixture was purged with nitrogen and heated at 80 °C for 16 h. The mixture was diluted with methylene chloride (20 mL) and filtered. The filtrate was concentrated and the residue was purified by chromatography (silica gel, 0–80% EtOAc/hexanes) and trituration with EtOAc to afford 1-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (**Example Compound 3**) (202 mg, 62%) as a white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.76 (s, 1H), 7.44 (d, *J* = 7.8 Hz, 1H), 7.36–7.28 (m, 5H), 7.11 (d, *J* = 7.8 Hz, 1H), 5.05 (s, 2H), 2.49 (s, 3H), 2.32 (s, 3H); ESI *m/z* 321 [M + H]⁺.

General Procedure B


Preparation of 4-(3-Benzyl-3*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (**Example Compound 4**) and 4-(1-Benzyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (**Example Compound 5**).



[0195] Step 1: To a solution of **10** (400 mg, 2.0 mmol) in 1,4-dioxane (10 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (669 mg, 1.5 mmol), sodium carbonate (2.0 M in H₂O, 2.0 mL, 4.0 mmol) and tetrakis(triphenylphosphine)palladium(0) (116 mg, 0.1 mmol). The reaction mixture was purged with nitrogen and heated at 80 °C for 16 h. The mixture was concentrated and purified by chromatography (silica gel, 0–8% methylene chloride/methanol) followed by trituration with EtOAc/hexanes to afford **11** (228 mg, 53%) as a light yellow solid: ESI *m/z* 215 [M + H]⁺.

[0196] Step 2: To a solution of **11** (220 mg, 1.03 mmol) in CH₃CN (10 mL) was added potassium carbonate (426 mg, 3.09 mmol) and benzyl chloride (0.12 mL, 1.03 mmol). The reaction was stirred at rt for 16 h. The mixture was concentrated and purified by chromatography (silica gel, 0–10% methanol/methylene chloride) to afford 4-(3-benzyl-3*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (**Example Compound 4**) (34 mg, 11%) as an off-white solid: ¹H NMR (300 MHz, CDCl₃) δ 8.34 (d, *J* = 1.8 Hz, 1H), 8.14 (s, 1H), 7.99 (d, *J* = 1.8 Hz, 1H), 7.40–7.31 (m, 5H), 5.52 (s, 2H), 2.44 (s, 3H), 2.30 (s, 3H); ESI *m/z* 305 [M + H]⁺; 4-(1-benzyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (**Example Compound 5**) (39 mg, 12%) as an off-white solid: ¹H NMR (300 MHz, CDCl₃) δ 8.46 (d, *J* = 1.8 Hz, 1H), 8.29 (s, 1H), 7.40–7.37 (m, 3H), 7.34 (d, *J* = 2.1 Hz, 1H), 7.24–7.21 (m, 2H), 5.41 (s, 2H), 2.33 (s, 3H), 2.16 (s, 3H); ESI *m/z* 305 [M + H]⁺.

Preparation of 3-Benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[d]oxazol-2(3*H*)-one (Example Compound 6)



[0197] Step 1: To a solution of **13** (5.00 g, 22.9 mmol) in acetic acid (50 mL), ethanol (100 mL), and water (5 mL) was added iron powder (6.42 g, 115 mmol). The reaction was heated at 80 °C for 2 h under nitrogen. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–100% hexanes/ethyl acetate) to give **14** (3.27 g, 76%) as a brown solid: ¹H NMR (300 MHz, CDCl₃) δ 6.88 (d, *J* = 2.2 Hz, 1H), 6.77 (dd, *J* = 8.3, 2.3 Hz, 1H), 6.60 (d, *J* = 8.3 Hz, 1H), 6.00–5.20 (br s, 3H).

[0198] Step 2: To a solution of **14** (1.50 g, 7.98 mmol) in 1,4-dioxane (100 mL) was added 1,1'-carbonyldiimidazole (1.55 g, 9.58 mmol). The reaction was heated at 80 °C for 17 h under nitrogen. The mixture was cooled to room temperature and 2N aq. HCl (40 mL) was added. The solution was diluted with ethyl acetate (200 mL) and washed with brine (2 × 50 mL). The organic


layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/hexanes) afforded **15** (1.08 g, 63%) as an orange solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 11.81 (s, 1H), 7.27–7.25 (m, 3H).

[0199] Step 3: To a solution of **15** (150 mg, 0.701 mmol) in acetonitrile (10 mL) was added benzyl bromide (180 mg, 1.05 mmol) and potassium carbonate (193 mg, 1.40 mmol). The reaction was heated at 80 °C for 3 h. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–50% ethyl acetate/hexanes) to afford **16** (195 mg, 92%) as an off-white solid: ^1H NMR (500 MHz, CDCl_3) δ 7.41–7.30 (m, 5H), 7.22 (dd, J = 8.5, 1.7 Hz, 1H), 7.08 (d, J = 8.5 Hz, 1H), 6.97 (d, J = 1.6 Hz, 1H), 4.97 (s, 2H).

[0200] Step 4: To a solution of **16** (195 mg, 0.641 mmol) in 1,4-dioxane (10 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (172 mg, 0.769 mmol), potassium carbonate (177 mg, 1.28 mmol), and tetrakis(triphenylphosphine)palladium(0) (37 mg, 0.032 mmol). The reaction mixture was purged with nitrogen and heated at 100 °C for 4 h. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–30% ethyl acetate/hexanes). It was further purified by reverse phase HPLC on Polaris column eluting with 10–90% CH_3CN in H_2O to give 3-benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[d]oxazol-2(3H)-one (**Example Compound 6**) (115 mg, 56%) as an off-white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.47–7.42 (m, 3H), 7.40–7.34 (m, 2H), 7.34–7.28 (m, 1H), 7.23 (d, J = 1.6 Hz, 1H), 7.12 (dd, J = 8.2 Hz, 7.7 Hz, 1H), 5.07 (s, 2H), 2.33 (s, 3H), 2.15 (s, 3H); ESI m/z 321 $[\text{M} + \text{H}]^+$.


General Procedure C:

Preparation of 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-4-amine (Example Compound 7), 1-Benzyl-5-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-7-amine (Example Compound 8) and *N*,1-Dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-4-amine (Example Compound 9)



[0201] To a solution of 6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-4-amine **17** (290 mg, 1.27 mmol) in CH₃CN (15 mL) was added potassium carbonate (350 mg, 2.54 mmol) and benzyl chloride (200 mg, 1.59 mmol). The reaction mixture was stirred at 60 °C for 16 h. The mixture was diluted with methylene chloride (20 mL) and filtered through a layer of *Celite*. The filtrate was concentrated and purified by chromatography (silica gel, 0–10% CH₃OH/CH₂Cl₂) to afford 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-4-amine (**Example Compound 7**) (109 mg, 27%) as an off-white solid: ¹H NMR (300 MHz, CDCl₃) δ 7.95 (s, 1H), 7.37–7.34 (m, 3H), 7.23–7.20 (m, 2H), 6.46 (d, *J* = 1.2 Hz, 1H), 6.40 (d, *J* = 1.2 Hz, 1H), 5.34 (s, 2H), 2.31 (s, 3H), 2.16 (s, 3H); ESI MS *m/z* 319 [M + H]⁺; 1-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-7-amine (**Example Compound 8**) (19 mg, 4.7%) as an off-white solid: ¹H NMR (300 MHz, CDCl₃) δ 8.15 (s, 1H), 7.43–7.40 (m, 3H), 7.23 (d, *J* = 1.2 Hz, 1H), 7.20–7.17 (m, 2H), 6.39 (d, *J* = 1.2 Hz, 1H), 5.69 (s, 2H), 2.40 (s, 3H), 2.27 (s, 3H); ESI MS *m/z* 319 [M + H]⁺; *N*,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-4-amine (**Example Compound 9**) (40 mg, 8%) as an off-white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.27 (s, 1H), 7.40–7.18 (m, 10H), 6.62 (d, *J* = 1.2 Hz, 1H), 6.57 (t, *J* = 6.0 Hz, 1H), 5.97 (d, *J* = 1.2 Hz, 1H), 5.41 (s, 2H), 4.48 (d, *J* = 6.0 Hz, 2H), 2.12 (s, 3H), 1.94 (s, 3H); ESI MS *m/z* 409 [M + H]⁺.


Preparation of 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (Example Compound 10)



Example 10

[0202] 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (Example Compound 10) was prepared by following the method for the preparation of Example 3 affording the product (158 mg, 47%) as a white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 11.81 (s, 1H), 7.90 (d, $J = 2.1$ Hz, 1H), 7.44–7.25 (m, 6H), 5.05 (s, 2H), 2.34 (s, 3H), 2.16 (s, 3H); MM m/z 321 [$M + H$] $^+$.

Preparation of 1-Benzyl-7-(3,5-dimethylisoxazol-4-yl)quinoxalin-2(1*H*)-one (Example Compound 11)



[0203] Step 1: A solution of **18** (500 mg, 2.3 mmol), benzylamine (1.2 g, 11.4 mmol) and pyridine (5.0 mL) was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* and the product was purified by chromatography (silica gel, 0–10% ethyl acetate/ hexanes) to provide **19** (630 mg, 91%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 8.38 (s, 1H), 8.05 (d, $J = 9.1$ Hz, 1H), 7.40–7.32 (m, 5H), 7.01 (d, $J = 1.9$ Hz, 1H), 6.79 (dd, $J = 9.1, 1.9$ Hz, 1H), 4.51 (d, $J = 5.5$ Hz, 2H).


[0204] Step 2: A mixture of **19** (100 mg, 0.33 mmol), iron powder (127 mg, 2.28 mmol), ammonium chloride (27 mg, 0.5 mmol), water (0.5 mL) and ethanol (3 mL) was heated at reflux for 0.5 hour. The reaction mixture was cooled and filtered. The solvent was removed to provide **20** (90

mg, 100%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.40–7.29 (m, 5H), 6.81–6.77 (m, 2H), 6.61–6.58 (m, 1H), 4.27 (s, 2H), 3.41 (s, 1H); ESI m/z 278 $[\text{M} + \text{H}]^+$.

[0205] Step 3: To a mixture of **20** (100 mg, 0.36 mmol), triethylamine (48 mg, 0.47 mmol), CH_2Cl_2 (0.5 mL) and THF (1.0 mL) was added a solution of ethyl bromoacetate (78 mg, 0.47 mmol) in THF (1.0 mL) at room temperature. The reaction mixture was stirred for 18 hours and then heated to 75 °C for 1 hour. The reaction mixture was concentrated and the product purified by chromatography (silica gel, 0–30% ethyl acetate/ hexanes) to provide **21** (44 mg, 39%) as a tan solid: ^1H NMR (500 MHz, CDCl_3) δ 7.38–7.29 (m, 4H), 7.24–7.22 (m, 2H), 6.98–6.93 (m, 2H), 6.55 (d, J = 8.3 Hz, 1H), 5.13 (s, 2H), 4.05 (s, 2H); ESI m/z 318 $[\text{M} + \text{H}]^+$.

[0206] Step 4: A mixture of **21** (44 mg, 0.14 mmol), **3** (47 mg, 0.21 mmol), K_2CO_3 (39 mg, 0.28 mmol), tetrakis(triphenylphosphine)palladium(0) (8 mg, 0.01 mmol), 1,4-dioxane (3 mL) and water (0.5 mL) was heated at 90 °C for 16 hours. The reaction mixture was concentrated onto silica gel and the product purified by chromatography (silica gel, 0–50% ethyl acetate/ hexanes) to provide 1-benzyl-7-(3,5-dimethylisoxazol-4-yl)quinoxalin-2(1H)-one (**Example Compound 11**) (16 mg, 34%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 8.43 (s, 1H), 7.94 (d, J = 8.2 Hz, 1H), 7.35–7.32 (m, 2H), 7.29–7.27 (m, 1H), 7.21–7.18 (m, 3H), 7.04 (s, 1H), 5.51 (s, 1H), 2.16 (s, 3H), 2.02 (s, 3H); ESI m/z 332 $[\text{M} + \text{H}]^+$.

Preparation of 1-Benzyl-7-(3,5-dimethylisoxazol-4-yl)-3,4-dihydroquinazolin-2(1H)-one (**Example Compound 12**)



[0207] Step 1: To a solution of **22** (1.19 g, 5.53 mmol) and benzaldehyde (594 mg, 5.60 mmol) in CH_2Cl_2 (50 mL) and CH_3CN (50 mL) was added acetic acid (0.2 mL). The mixture was stirred at rt for 1 h. NaBH(OAc)_3 (3.52 g, 16.59 mmol) was added. The mixture was stirred at rt for 8 h. The reaction was quenched with saturated aq. NaHCO_3 (50 mL) and concentrated, the residue was suspended in EtOAc (300 mL), washed with brine (100 mL). The organic layer was separated, dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica

gel, 0–50% EtOAc/heptane) to afford **23** (201 mg, 12%) as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 8.75 (d, J = 5.7 Hz, 1H), 7.93 (br.s, 1H), 7.55 (d, J = 8.4 Hz, 1H), 7.38–7.31 (m, 6H), 6.76 (d, J = 1.8 Hz, 1H), 6.69 (dd, J = 8.4, 1.8 Hz, 1H), 4.39 (d, J = 6.0 Hz, 2H).


[0208] Step 2: To a solution of **23** (518 mg, 1.70 mmol) in THF (20 mL) was added $\text{BH}_3 \cdot \text{THF}$ (1.0 M in THF, 8.50 mL, 8.50 mmol). The mixture was heated to reflux for 16 h. MeOH (40 mL) was added slowly followed by 2 N HCl (40 mL). The mixture was heated to reflux for 3 h. NH_4OH (60 mL) was added, the mixture was extracted with EtOAc (200 mL $\times$ 3). The organic layer was separated, dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–10% MeOH/methylene chloride) to afford **24** (372 mg, 75%) as an colorless gum: ^1H NMR (300 MHz, DMSO- d_6) δ 7.32–7.21 (m, 5H), 6.98 (d, J = 7.8 Hz, 1H), 6.87 (t, J = 6.0 Hz, 1H), 6.65 (dd, J = 8.1, 2.1 Hz, 1H), 6.53 (d, J = 2.1 Hz, 1H), 4.33 (d, J = 5.7 Hz, 2H), 3.71 (s, 2H), 1.92 (br.s, 2H).

[0209] Step 3: Using the procedure used for Example Compound 3 step 2 starting with compound **24** (362 mg, 1.24 mmol) afforded **25** (325 mg, 85%) as a yellow solid: ^1H NMR (300 MHz, DMSO- d_6) δ 7.33–7.31 (m, 3H), 7.25–7.23 (m, 3H), 7.09 (d, J = 1.8 Hz, 2H), 6.86 (s, 1H), 5.05 (s, 2H), 4.35 (d, J = 1.5 Hz, 2H).

[0210] Step 4: Using the procedure used for Example Compound 3 step 3 starting with compound **25** (317 mg, 1.00 mmol) afforded Example Compound **12** (199 mg, 60%) as a white solid: ^1H NMR (500 MHz, DMSO- d_6) δ 7.34–7.21 (m, 7H), 6.90 (dd, J = 7.5, 1.0 Hz, 1H), 6.58 (d, J = 1.0 Hz, 1H), 5.09 (s, 2H), 4.43 (s, 2H), 2.06 (s, 3H), 1.89 (s, 3H); MM m/z 334 $[\text{M} + \text{H}]^+$.

General Procedure D:

Preparation of 4-(1-benzyl-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazol (Example Compound **13**)



[0211] Step 1: To a mixture of **26** (1.00 g, 5.32 mmol) and **3** (1.78 g, 7.98 mmol) in 1,4-dioxane (35 mL) and water (7.5 mL) was added potassium carbonate (1.47 g, 10.6 mmol) and


tetrakis(triphenylphosphine)palladium(0) (307 mg, 0.27 mmol). The reaction was stirred and heated at 90 °C for 16 h. The reaction mixture was diluted with methanol (20 mL) and silica gel (15 g) was added. The slurry was concentrated to dryness and the resulting powder was loaded onto silica gel and eluted with 0–90% ethyl acetate in hexanes. The clean product was concentrated to give **27** (939 mg, 70%) as a yellow-green solid: ^1H NMR (500 MHz, CDCl_3) δ 7.45 (t, J = 2.0 Hz, 1H), 6.78 (t, J = 2.0 Hz, 1H), 2.37 (s, 3H), 2.22 (s, 3H).

[0212] Step 2: To a solution of **27** (300 mg, 1.47 mmol) in 1,2-dichloroethane (15 mL) was added benzaldehyde (156 mg, 1.47 mmol) and glacial acetic acid (200 μL) at room temperature. After stirring for 17 h, CH_2Cl_2 (20 mL) then saturated aq. NaHCO_3 (20 mL, slowly) was added. The organic layer was separated and dried over Na_2SO_4 . The suspension was filtered and concentrated. The material was purified by chromatography (silica gel, 0–60% ethyl acetate in hexanes) to afford a yellow solid which was dissolved in methanol (10 mL), sodium borohydride (52 mg, 1.35 mmol) was added at room temperature. After stirring for 1 h, additional sodium borohydride (156 mg, 3.40 mmol) was added and the reaction stirred 1 h. A 2N aq. HCl solution was added to the mixture until pH 4 (2 mL) then a saturated NaHCO_3 solution was added to basify to pH 8 (2 mL). Water was added (10 mL) and the solution was extracted with ethyl acetate (3 $\times$ 100 mL). The ethyl acetate extracts were combined, dried over Na_2SO_4 , filtered and concentrated to afford **28** (401 mg, 93%) as a white solid: ^1H NMR (500 MHz, CDCl_3) δ 7.48 (s, 1H), 7.37–7.26 (m, 5H), 6.58 (s, 1H), 4.38 (s, 2H), 4.33 (br s, 2H), 3.77 (br s, 1H), 2.24 (s, 3H), 2.08 (s, 3H).

[0213] Step 3: To **28** (350 mg, 1.19 mmol) was added triethylorthoacetate (3.0 mL, 16.4 mmol) and sulfamic acid (1 mg). The mixture was heated to 100 °C for 1 h. The mixture was diluted with methanol (20 mL) and adsorbed onto silica gel (10 g). The material was purified by chromatography (silica gel, 0–60% ethyl acetate in hexanes then 0–5% methanol in CH_2Cl_2) to afford 4-(1-benzyl-2-methyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound **13**, 169 mg, 45%) as a white solid: ^1H NMR (500 MHz, CD_3OD) δ 8.32 (d, J = 1.0 Hz, 1H), 7.78 (d, J = 1.0 Hz, 1H), 7.36–7.29 (m, 3H), 7.20–7.17 (m, 2H), 5.56 (s, 2H), 2.69 (s, 3H), 2.36 (s, 3H), 2.18 (s, 3H); ESI m/z 319 $[\text{M} + \text{H}]^+$.

General Procedure E:

Preparation of 1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1*H*-benzo[d]imidazol-2(3*H*)-one (Example Compound **91) and 4-Amino-1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-2(3*H*)-one (Example Compound **90**)**



[0214] Step 1: To a solution of **29** (1.00 g, 4.61 mmol) in 1,4-dioxane (40 mL) and water (4 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (1.23 g, 5.53 mmol), potassium carbonate (1.27 g, 9.22 mmol), and tetrakis(triphenylphosphine)palladium(0) (266 mg, 0.231 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C overnight. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–30% ethyl acetate/hexanes) to give a yellow solid which was dissolved in acetic acid (15 mL), *N*-bromosuccinimide (753 mg, 4.23 mmol) was added at 0 °C. The reaction was warmed to room temperature and stirred overnight. The mixture was concentrated *in vacuo*. The residue was suspended in hot MeOH, cooled to room temperature and basified with 10% aq. NaHCO₃. The mixture was diluted with water and filtered. The filter cake was washed with water and dried *in vacuo* to afford **30** (1.10 g, 87%) as a yellow solid: ¹H NMR (500 MHz, CDCl₃) δ 8.04 (d, *J* = 2.1 Hz, 1H), 7.61 (d, *J* = 2.1 Hz, 1H), 6.69 (bs, 2H), 2.40 (s, 3H), 2.26 (s, 3H); ESI *m/z* 312 [M + H]⁺.


[0215] Step 2: To a solution of **30** (500 mg, 1.60 mmol) in toluene (50 mL) under nitrogen atmosphere was added 4-chlorobenzylamine (1.36 g, 9.62 mmol), cesium carbonate (1.04 g, 3.02 mmol), 2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl (114 mg, 0.240 mmol), and tris(dibenzylideneacetone)dipalladium(0) (146 mg, 0.160 mmol). The reaction mixture was heated at 90 °C overnight, cooled to room temperature, and purified by chromatography (silica gel, 0–50% ethyl acetate in hexanes) to afford **31** (290 mg, 49%) as a red solid: ESI *m/z* 373 [M + H]⁺.

[0216] Step 3: To a mixture of **31** (290 mg, 0.779 mmol) in 1,4-dioxane (10 mL) was added 1,1'-carbonyldiimidazole (630 mg, 3.89 mmol) and DMAP (a crystal). The reaction was heated in a sealed tube at 130 °C for 4 days. The mixture was concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give **Example Compound 91** (144 mg, 46%) as an orange solid: ¹H NMR (500 MHz, CD₃OD) δ 7.80 (d, *J* = 1.4 Hz, 1H), 7.40–7.35 (m, 4H), 7.24 (d, *J* = 1.4 Hz, 1H), 5.15 (s, 2H), 2.32 (s, 3H), 2.15 (s, 3H); ESI *m/z* 399 [M + H]⁺.

[0217] **Step 4:** To a solution of **Example Compound 91** (70 mg, 0.18 mmol) in tetrahydrofuran (10 mL) was added sodium dithionite (183 mg, 1.05 mmol) in water (10 mL). The reaction mixture was stirred at room temperature overnight and concentrated under vacuum. To the residue was added 2N HCl and heated to reflux, cooled to room temperature, and concentrated in vacuum. The residue was dissolved in MeOH and basified by conc. NH_4OH , concentrated, and purified by chromatography (silica gel, 0–100% hexanes/ethyl acetate). It was further purified by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O to give **Example Compound 90** (34 mg, 51%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.36–7.28 (m, 4H), 6.40 (d, $J = 1.4$ Hz, 1H), 6.25 (d, $J = 1.4$ Hz, 1H), 5.03 (s, 2H), 2.28 (s, 3H), 2.12 (s, 3H); ESI m/z 369 $[\text{M} + \text{H}]^+$.

General Procedure F:

Preparation of 4-(1-(cyclopropylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole (Example Compound 14) and 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine (Example Compound 75)



[0218] **Step 1:** A solution of **32** (488 mg, 2.10 mmol) and 2,4-pentanedione (421 mg, 4.21 mmol) in absolute ethanol (28 mL) and 5 N aq. HCl (7.8 mL) was heated to reflux for 3 h. The mixture was concentrated to dryness and ethyl acetate was added (200 mL). The solution was washed with saturated aq. NaHCO_3 (250 mL) and saturated aq. NaCl solution (250 mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 0–40% hexanes/ethyl acetate) to afford **33** (495 mg, 92%) as a orange solid: ^1H NMR (500 MHz, CDCl_3) δ 10.38 (br s, 1 H), 8.24 (d, $J = 2.0$ Hz, 1H), 8.12 (d, $J = 1.0$ Hz, 1H), 2.73 (s, 3H).

[0219] **Step 2:** To a mixture of **33** (200 mg, 0.78 mmol) and **3** (262 mg, 1.17 mmol) in 1,4-dioxane (6 mL) and water (1.5 mL) was added potassium carbonate (216 mg, 1.56 mmol) and tetrakis(triphenylphosphine)palladium(0) (45 mg, 0.04 mmol). The reaction was stirred and heated


at 90 °C for 17 h. The reaction mixture was diluted with methanol (20 mL) and silica gel (15 g) was added. The suspension was concentrated to dryness and the resulting powder was purified by chromatography (silica gel, 0–90% hexanes/ethyl acetate) to give **34** (187 mg, 88%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 8.00 (d, $J = 1.5$ Hz, 1H), 7.89 (s, 1H), 2.76 (s, 3H), 2.45 (s, 3H), 2.30 (s, 3H).

[0220] Step 3: To a solution of **34** (217 mg, 0.797 mmol), potassium carbonate (220 mg, 1.59 mmol), acetonitrile (5 mL) and DMF (1 mL) was added bromomethylcyclopropane (129 mg, 0.956 mmol) and the reaction was heated at 60 °C for 17 h. The material was cooled to room temperature and poured into a saturated aq. NaCl solution (30 mL). Ethyl acetate (100 mL) was added and the layers were separated. The ethyl acetate layer was washed with saturated aq. NaCl solution (2 $\times$ 20 mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 0–90% hexanes/ethyl acetate) to give **Example 14** (178 mg, 68%) as a yellow solid: ^1H NMR (500 MHz, CD_3OD) δ 8.03 (d, $J = 1.5$ Hz, 1H), 7.93 (d, $J = 1.5$ Hz, 1H), 4.27 (d, $J = 7.0$ Hz, 2H), 2.75 (s, 3H), 2.46 (s, 3H), 2.30 (s, 3H), 1.38–1.28 (m, 1H), 0.65–0.60 (m, 2H), 0.51–0.46 (m, 2H). ESI m/z 327 $[\text{M} + \text{H}]^+$

[0221] Step 4: To a solution of **Example Compound 14** (160 mg, 0.51 mmol) in THF (10 mL) was added a solution of sodium dithionite (446 mg, 2.56 mmol) in water (10 mL) dropwise over 5 min. The solution was stirred at room temperature for 16 h and the solvents were removed *in vacuo*. Methanol (20 mL) was added and the suspension stirred at room temperature for 3 h. The mixture was filtered and the filtrate was concentrated to dryness. A solution of 2N aq. HCl (10 mL) was added to the residue and was heated to reflux for 5 min. After concentration to dryness, methanol (20 mL) was added and the solution was adjusted to pH 8 using saturated aq. NaHCO_3 solution (10 mL). Silica gel was added (10 g) and the suspension was concentrated to dryness. The resulting powder was purified by chromatography (silica gel, 0–5% methanol/methylene chloride), the product was then purified by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O to give **Example Compound 75** (131 mg, 99%) as a white solid: ^1H NMR (500 MHz, CD_3OD) δ 6.70 (s, 1H), 6.44 (d, $J = 1.0$ Hz, 1H), 4.08 (d, $J = 6.5$ Hz, 2H), 2.61 (s, 3H), 2.40 (s, 3H), 2.25 (s, 3H), 1.30–1.19 (m, 1H), 0.62–0.53 (m, 2H), 0.45–0.40 (m, 2H). ESI m/z 297 $[\text{M} + \text{H}]^+$.

General Procedure G:

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one (Example Compound 15) and 4-amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one (Example Compound 16)



[0222] Step 1: To a solution of **32** (232 mg, 1.0 mmol) in 1,4-dioxane (5 mL) was added CDI (194 mg, 1.2 mmol). The reaction was heated at 60 °C for 16 h. The solid was collected and dried to give **35** (202 mg, 78%) as a brown yellow solid: ^1H NMR (300 MHz, DMSO- d_6) δ 11.83 (br s, 1H), 11.53 (br s, 1H), 7.86 (d, J = 1.8 Hz, 1H), 7.43 (d, J = 1.8 Hz, 1H).

[0223] Step 2: To a solution of **35** (200 mg, 0.78 mmol) in DMF (7 mL) was added potassium carbonate (118 mg, 0.85 mmol) and benzyl chloride (98 mg, 0.78 mmol). The reaction was stirred at rt for 16 h. The mixture was diluted with EtOAc (100 mL) and washed with brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–100% ethyl acetate/hexanes) afforded **36** (101 mg, 37%) as a yellow solid: ^1H NMR (300 MHz, DMSO- d_6) δ 12.15 (s, 1H), 7.90 (d, J = 0.9 Hz, 1H), 7.75 (d, J = 1.2 Hz, 1H), 7.36–7.28 (m, 5H), 5.10 (s, 2H).


[0224] Step 3: To a solution of **36** (100 mg, 0.29 mmol) in 1,4-dioxane (7 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (128 mg, 0.57 mmol), sodium carbonate (2.0 M in H₂O, 0.43 mL, 0.86 mmol) and tetrakis(triphenylphosphine)palladium(0) (34 mg, 0.03 mmol). The reaction mixture was purged with nitrogen and heated at 80 °C for 16 h. The mixture was diluted with methylene chloride (20 mL) and filtered. The filtrate was concentrated and purified by chromatography (silica gel, 10–50% ethyl acetate/hexanes) followed by trituration with ethyl acetate to afford **Example Compound 15** (70 mg, 66%) as a yellow solid: ^1H NMR (300 MHz, DMSO- d_6) δ 12.11 (s, 1H), 7.72 (d, J = 1.5 Hz, 1H), 7.50 (d, J = 1.5 Hz, 1H), 7.42–7.28 (m, 5H), 5.13 (s, 2H), 2.35 (s, 3H), 2.15 (s, 3H); ESI m/z 365 $[\text{M} + \text{H}]^+$.

[0225] Step 4: To a solution of **Example Compound 15** (52 mg, 0.14 mmol) in THF (5 mL) and water (4 mL) was added Na₂S₂O₄ (149 mg, 0.86 mmol). The mixture was stirred at rt for 4 h, 2N HCl (1 mL) was added, the mixture was heated to reflux for 15 minutes then cooled to rt. Na₂CO₃ was added slowly to adjust to pH 9. The mixture was extracted with CH₂Cl₂ (100 mL), the organic

layer was washed with brine (50 mL), filtered, concentrated and purified by chromatography (silica gel, 70–100% ethyl acetate/hexanes) to afford **Example Compound 16** (30 mg, 63%) as an off-white solid: ^1H NMR (500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 7.36–7.25 (m, 5H), 6.28 (s, 2H), 5.04 (s, 2H), 4.95 (s, 2H), 2.28 (s, 3H), 2.10 (s, 3H); ESI m/z 335 $[\text{M} + \text{H}]^+$.

General Procedure H:

Preparation of 4-(1-benzyl-4-bromo-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole (Example Compound 121)




[0226] Step 1: To a solution of **30** (1.09 g, 3.49 mmol) in tetrahydrofuran (30 mL) was added sodium dithionite (4.86 g, 28.0 mmol) in water (15 mL). The reaction mixture was stirred at room temperature overnight and concentrated under vacuum. The residue was dissolved in MeOH/water (1:1, 150 mL) and the solid was precipitated by removing some MeOH under vacuum. The solid was filtered, washed with water and dried under vacuum to afford **37** (440 mg, 34%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 6.85 (d, $J = 1.8$ Hz, 1H), 6.51 (d, $J = 1.8$ Hz, 1H), 4.00–3.60 (bs, 2H), 3.60–3.30 (bs, 2H), 2.36 (s, 3H), 2.23 (s, 3H); ESI m/z 282 $[\text{M} + \text{H}]^+$.

[0227] Step 2: To a solution of **37** (4.01 g, 14.2 mmol) in methanol (87 mL) was added triethyl orthoacetate (3.45 g, 21.3 mmol) and sulfamic acid (69 mg, 0.71 mmol). The reaction was stirred at room temperature for 5 h. The reaction mixture was diluted with water (50 mL), basified with NaHCO_3 and filtered. The solid was dried to afford **38** (4.2 g, 96%) as a brown solid: ^1H NMR (300 MHz, DMSO- d_6) δ 12.82 (br.s, 1H), 7.42 (d, $J = 1.5$ Hz, 1H), 7.31 (d, $J = 1.5$ Hz, 1H), 2.52 (s, 3H), 2.40 (s, 3H), 2.24 (s, 3H).

[0228] Step 3: The mixture of **38** (300 mg, 0.980 mmol), benzyl bromide (503 mg, 2.94 mmol), and potassium carbonate (676 mg, 4.90 mmol) in acetonitrile (50 mL) was heated in sealed tube at 75 °C overnight. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give **Example Compound**

121 (276 mg, 71%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.40–7.25 (m, 5H), 7.15 (d, J = 7.7 Hz, 2H), 5.51 (s, 2H), 2.64 (s, 3H), 2.32 (s, 3H), 2.15 (s, 3H); ESI m/z 396 $[\text{M} + \text{H}]^+$.


Preparation of 4-(1-benzyl-4-methoxy-2-methyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole (Example Compound 66)



[0229] A mixture of **Example 121** (80 mg, 0.20 mmol), NaOCH_3 (108 mg, 2.0 mmol) and CuI (57 mg, 0.30 mmol) in MeOH (1 mL) and DMF (3 mL) was purged with nitrogen and heated at 100°C for 6 h. The mixture was diluted with ethyl acetate (100 mL) and washed with brine (50 mL). The organic layer was dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 40–100% EtOAc /hexanes) to afford **Example Compound 66** (386 mg, 55%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.35–7.30 (m, 3H), 7.09–7.06 (m, 2H), 6.64 (d, J = 1.2 Hz, 1H), 6.53 (s, 1H), 5.32 (s, 2H), 4.03 (s, 3H), 2.66 (s, 3H), 2.33 (s, 3H), 2.19 (s, 3H); ESI m/z 348 $[\text{M} + \text{H}]^+$.

General procedure I:

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-4-nitro-1H-benzo[d]imidazol-2-amine (Example Compound 18) and 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N²-ethyl-1H-benzo[d]imidazole-2,4-diamine (Example Compound 19)




[0230] Step 1: A mixture of **Example Compound 15** (73 mg, 0.668 mmol) in POCl_3 (3 mL) was heated at 110°C for 16 h. The reaction mixture was concentrated, the residue was dissolved in CH_2Cl_2 (100 mL), washed with saturated NaHCO_3 (2×50 mL) and brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was dissolved in a solution of ethylamine in THF (2.0 M, 10 mL), and the mixture was heated at 70°C for 3 h. The reaction mixture was concentrated, the residue was purified by chromatography (silica gel, 20–60% EtOAc /hexanes) to afford **Example Compound 18** (113 mg, 43%) as an orange solid: ^1H NMR (300 MHz, CDCl_3) δ 7.84

(d, $J = 1.5$ Hz, 1H), 7.42–7.35 (m, 3H), 7.16–7.13 (m, 2H), 7.03 (d, $J = 1.5$ Hz, 1H), 5.15 (s, 2H), 4.29 (t, $J = 5.4$ Hz, 1H), 3.78–3.69 (m, 2H), 2.36 (s, 3H), 2.21 (s, 3H), 1.27 (t, $J = 7.5$ Hz, 3H); ESI m/z 392 $[M + H]^+$.

[0231] Step 2: To a solution of **Example Compound 18** (90 mg, 0.23 mmol) in THF (5 mL) and water (4 mL) was added $\text{Na}_2\text{S}_2\text{O}_4$ (240 mg, 1.38 mmol). The mixture was stirred at rt for 4 h, 2N HCl (1 mL) was added, the mixture was heated to reflux for 15 minutes then cooled to rt. Na_2CO_3 was added slowly to adjust to pH 9. The mixture was extracted with CH_2Cl_2 (100 mL), the organic layer was washed with brine (50 mL), dried over Na_2SO_4 , filtered, concentrated and purified by chromatography (silica gel, 0–10% methanol/ethyl acetate) to afford **Example Compound 19** (60 mg, 72%) as an off-white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.34–7.20 (m, 5H), 6.62 (t, $J = 5.4$ Hz, 1H), 6.30 (d, $J = 1.5$ Hz, 1H), 6.21 (d, $J = 1.5$ Hz, 1H), 5.19 (s, 2H), 4.83 (s, 2H), 3.47–3.38 (m, 2H), 2.28 (s, 3H), 2.11 (s, 3H), 1.22 (t, $J = 7.2$ Hz, 3H); ESI m/z 362 $[M + H]^+$.

General Procedure J:

Preparation of methyl 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carboxylate (**Example Compound 20**), 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carboxamide (**Example Compound 21**) and 4-(aminomethyl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one (**Example Compound 22**)



[0232] Step 1: To a solution of **39** (2.00 g, 8.70 mmol) in 1,4-dioxane (80 mL) and water (8 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (2.13 g, 9.57 mmol), potassium carbonate (2.40 g, 17.4 mmol) and tetrakis(triphenylphosphine)palladium(0) (502

mg, 0.435 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C overnight. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–50% ethyl acetate in hexanes) to afford **40** (1.43 g, 63%) as an off-white solid: ^1H NMR (500 MHz, CDCl_3) δ 7.74 (d, $J = 2.1$ Hz, 1H), 7.15 (dd, $J = 2.1, 8.4$ Hz, 1H), 6.73 (d, $J = 8.4$ Hz, 1H), 5.81 (s, 2H), 3.88 (s, 3H), 2.37 (s, 3H), 2.23 (s, 3H); ESI m/z 247 $[\text{M} + \text{H}]^+$.

[0233] Step 2: To a mixture of **40** (1.34 g, 5.45 mmol) in acetic acid (40 mL) was added *N*-bromosuccinimide (1.07 g, 5.99 mmol). The mixture was stirred at room temperature for 30 min and concentrated. The residue was dissolved in MeOH and neutralized to pH 7 with 10% sodium bicarbonate. The mixture was diluted with water, filtered. The filter cake was washed with water, and dried under vacuum to afford **41** (1.65 g, 93%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 7.74 (d, $J = 2.1$ Hz, 1H), 7.47 (d, $J = 2.1$ Hz, 1H), 6.43 (bs, 2H), 3.90 (s, 3H), 2.37 (s, 3H), 2.23 (s, 3H).

[0234] Step 3: To a solution of **41** (500 mg, 1.54 mmol) in toluene (40 mL) under nitrogen atmosphere was added benzylamine (823 mg, 7.69 mmol), cesium carbonate (1.00 g, 2.08 mmol), 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl (110 mg, 0.231 mmol), and tris(dibenzylideneacetone) dipalladium(0) (141 mg, 0.154 mmol). The reaction mixture was heated at 90 °C overnight, cooled to room temperature and purified by chromatography (silica gel, 0–20% ethyl acetate in hexanes) to afford **42** (310 mg, 57%) as a light brown solid: ^1H NMR (500 MHz, CDCl_3) δ 7.40–7.25 (m, 6H), 6.56 (d, $J = 1.8$ Hz, 1H), 5.68 (s, 2H), 4.36 (d, $J = 4.4$ Hz, 2H), 3.88 (s, 3H), 3.68 (s, 1H), 2.22 (s, 3H), 2.09 (s, 3H); ESI m/z 352 $[\text{M} + \text{H}]^+$.

[0235] Step 4: To a mixture of **42** (310 mg, 0.883 mmol) in 1,4-dioxane (10 mL) was added 1,1'-carbonyldiimidazole (244 mg, 2.12 mmol) and DMAP (one crystal). The reaction was heated in a sealed tube at 80 °C for 5 days. The mixture was concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give **Example Compound 20** (160 mg, 48%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.54 (d, $J = 1.5$ Hz, 1H), 7.37–7.24 (m, 5H), 7.07 (d, $J = 1.5$ Hz, 1H), 5.14 (s, 2H), 3.97 (s, 3H), 2.27 (s, 3H), 2.09 (s, 3H); HPLC >99%, $t_R = 15.0$ min; ESI m/z 378 $[\text{M} + \text{H}]^+$.


[0236] Step 5: To a mixture of **Example Compound 20** (50 mg, 0.13 mmol) in formamide (4 mL) was added potassium *tert*-butoxide (30 mg, 0.26 mmol). The mixture was heated in the microwave at 100 °C for 3 h, concentrated, and purified by chromatography (silica gel, 0–20% methanol in ethyl acetate) to afford **Example Compound 21** (13 mg, 26%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.41 (d, $J = 1.3$ Hz, 1H), 7.37–7.24 (m, 5H), 7.00 (d, $J = 1.4$ Hz, 1H), 5.13 (s, 2H), 2.28 (s, 3H), 2.11 (s, 3H); HPLC 98.3%, $t_R = 12.3$ min; ESI m/z 363 $[\text{M} + \text{H}]^+$.

[0237] Step 6: To a solution of **Example Compound 21** (40 mg, 0.11 mmol) in THF (10 mL) under nitrogen atmosphere was added sodium borohydride (38 mg, 0.99 mmol). The mixture was

heated to 65 °C and boron trifluoride diethyl etherate (0.2 mL) was added. The mixture was heated at 65 °C for 2 h. After cooling to room temperature, hydrochloride acid (2N, 5 mL) was added and the mixture stirred for 2 h. The mixture was basified with NaOH (2N, 5 mL), concentrated, and purified by chromatography (silica gel, 0–100% CMA in methylene chloride) (CMA = chloroform:methanol:concentrated ammonium hydroxide = 80:18:2). It was further purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH₃CN in H₂O to give **Example Compound 22** (16 mg, 42%) as an off-white solid: ¹H NMR (500 MHz, CD₃OD) δ 7.37–7.23 (m, 5H), 6.99 (d, *J* = 1.4 Hz, 1H), 6.77 (d, *J* = 1.4 Hz, 1H), 5.10 (s, 2H), 3.93 (s, 2H), 2.27 (s, 3H), 2.10 (s, 3H); ESI *m/z* 340 [M + H]⁺.


General Procedure K:

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine (Example Compound 55)



[0238] A mixture of **Example 121** (250 mg, 0.63 mmol), BocNH₂ (221 mg, 1.89 mmol), Xantphos (73 mg, 0.126 mmol), Pd₂(dba)₃ (58 mg, 0.063 mmol) and Cs₂CO₃ (720 mg, 2.21 mmol) in 1,4-dioxane (13 mL) was purged with nitrogen and heated at 100 °C for 18 h. The mixture was diluted with methylene chloride (200 mL) and filtered. The filtrate was concentrated and purified by chromatography (silica gel, 0–50% EtOAc/hexanes) to afford a light brown foam which was dissolved in CH₂Cl₂ (4 mL), TFA (2 mL) was added. The mixture was stirred at rt for 2 h, concentrated, the residue was dissolved in ethyl acetate (100 mL) and washed with saturated NaHCO₃ (50 mL × 2). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–10% MeOH/EtOAc) afforded **Example Compound 55** (146 mg, 88%) as an off-white solid: ¹H NMR (500 MHz, CDCl₃) δ 7.34–7.28 (m, 3H), 7.09–7.08 (m, 2H), 6.42 (d, *J* = 1.5 Hz, 1H), 6.36 (d, *J* = 1.5 Hz, 1H), 5.28 (s, 2H), 4.42 (br.s, 2H), 2.60 (s, 3H), 2.31 (s, 3H), 2.17 (s, 3H); ESI *m/z* 333 [M + H]⁺.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-pyrrolo[3,2-*b*]pyridine-3-carbonitrile (Example Compound 88) and 4-(1-benzyl-3-chloro-1*H*-pyrrolo[3,2-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 89)



[0239] Step 1: To a suspension of **43** (200 mg, 1.0 mmol) in CH_3CN (6 mL) was added ClSO_2NCO (360 mg, 2.5 mmol). The reaction mixture was stirred at 60 °C for 4 h. After the mixture was cooled to rt, DMF (1 mL) was added. The mixture was stirred at rt for 1 h. The mixture was diluted with 30% *i*-PrOH in CHCl_3 (50 mL) and washed with brine (20 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The crude was dissolved in CH_3CN (4 mL), potassium carbonate (280 mg, 2.0 mmol) and benzyl chloride (128 mg, 1.0 mmol) were added. The reaction was stirred at 70 °C for 16 h. The reaction mixture was filtered through a layer of celite,TM concentrated. The residue was purified by chromatography (silica gel, 0–50% ethyl acetate/hexanes) to afford **44** (16 mg, 5%) as a yellow oil and **45** (12 mg, 4%) as an off-white solid; **44**: ESI MS m/z 312 $[\text{M} + \text{H}]^+$; **45**: ESI MS m/z 321 $[\text{M} + \text{H}]^+$.


[0240] Step 2: Using the similar procedure used for General Procedure C step 1 on compound **44** (16 mg, 0.051 mmol) afforded Example Compound **88** (6 mg, 36%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 8.55 (s, 1H), 7.98 (s, 1H), 7.50 (s, 1H), 7.41–7.40 (m, 3H), 7.20–7.15 (m, 2H), 5.42 (s, 2H), 2.34 (s, 3H), 2.16 (s, 3H); ESI MS m/z 329 $[\text{M} + \text{H}]^+$.

[0241] Using the similar procedure used for General Procedure C step 1 on compound **45** (12 mg, 0.037 mmol) afforded Example Compound **89** (8 mg, 64%) as a yellow solid: ^1H NMR (300

MHz, CDCl₃) δ 8.49 (s, 1H), 7.55 (s, 1H), 7.50 (s, 1H), 7.38–7.36 (m, 3H), 7.18–7.16 (m, 2H), 5.36 (s, 2H), 2.34 (s, 3H), 2.16 (s, 3H); ESI MS m/z 338 [M + H]⁺.

General Procedure M:

Preparation of 5-(3,5-dimethylisoxazol-4-yl)-*N*-phenyl-1*H*-pyrrolo[3,2-*b*]pyridin-3-amine (Example Compound 23)



[0242] **Step 1:** To a solution of 46 (500 mg, 2.54 mmol) in 1,4-dioxane (10 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (792 mg, 3.56 mmol), sodium carbonate (538 mg in 2 mL H₂O, 5.08 mmol) and tetrakis(triphenylphosphine)palladium(0) (294 mg, 0.25 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 16 h. The mixture was filtered through a layer of *Celite*. The filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded 47 (700 mg, >100%) as a yellow oil: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.4 (s, 1H), 7.85 (dd, *J* = 8.1, 0.9 Hz, 1H), 7.68 (t, *J* = 3.0 Hz, 1H), 7.23 (d, *J* = 8.1 Hz, 1H), 6.58 (d, *J* = 2.1 Hz, 1H), 2.49 (s, 3H), 2.37 (s, 3H).

[0243] **Step 2:** To a solution of 47 (700 mg, 2.54 mmol) in DMF (8 mL) at 0 °C was added NBS (497 mg, 2.79 mmol). The reaction mixture was stirred at 0 °C for 2 h. The mixture was diluted with methylene chloride (50 mL) and washed with brine (20 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded 48 (660 mg, 89%) as a brown solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.8 (s, 1H), 7.92 (d, *J* = 6.0 Hz, 1H), 7.90 (s, 1H), 7.36 (d, *J* = 8.4 Hz, 1H), 2.49 (s, 3H), 2.37 (s, 3H); ESI m/z 292 [M + H]⁺.


[0244] **Step 3:** To a solution of 48 (250 mg, 0.86 mmol) in CH₂Cl₂ (5 mL) was added NEt₃ (130 mg, 1.28 mmol), DMAP (12 mg, 0.1 mmol) and di-*tert*-butyl dicarbonate (224 mg, 1.03 mmol). The reaction was stirred at rt for 16 h. The reaction mixture was concentrated. Purification by chromatography (silica gel, 0–30% ethyl acetate/hexanes) afforded 49 (210 mg, 70%) as an off-white

solid: ^1H NMR (300 MHz, CDCl_3) δ 8.43 (d, $J = 5.4$ Hz, 1H), 7.93 (s, 1H), 7.34 (d, $J = 5.1$ Hz, 1H), 2.64 (s, 3H), 2.50 (s, 3H), 1.69 (s, 9H).

[0245] Step 4: To a solution of **49** (100 mg, 0.26 mmol) in 1,4-dioxane (5 mL) under nitrogen atmosphere was added aniline (71 mg, 0.76 mmol), cesium carbonate (250 mg, 0.76 mmol), X-phos (24 mg, 0.05 mmol), and tris(dibenzylideneacetone)dipalladium(0) (23 mg, 0.03 mmol). The reaction mixture was heated at 90 °C for 16 h. The mixture was diluted with methylene chloride (10 mL) and filtered through a layer of *Celite*. The filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/hexanes) gave a red oil which was dissolved in methylene chloride (5 mL), TFA (2 mL) was added, the mixture was stirred at rt for 2 h. The mixture was concentrated, the residue was dissolved in methylene chloride (100 mL), washed with saturated NaHCO_3 (50 mL $\times$ 2) and brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded **Example Compound 23** (47 mg, 64%) as a yellow solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.1 (d, $J = 1.8$ Hz, 1H), 7.82 (d, $J = 8.4$ Hz, 1H), 7.61 (d, $J = 2.7$ Hz, 1H), 7.43 (s, 1H), 7.25 (d, $J = 8.4$ Hz, 1H), 7.09 (d, $J = 8.4$ Hz, 1H), 7.07 (d, $J = 7.2$ Hz, 1H), 6.85 (d, $J = 7.5$ Hz, 2H), 6.60 (t, $J = 7.2$ Hz, 1H), 2.48 (s, 3H), 2.29 (s, 3H); ESI MS m/z 305 $[\text{M} + \text{H}]^+$.

General Procedure N:

Preparation of 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-*b*]pyridine-4-oxide (**Example Compound 24**) and 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-*b*]pyridin-5(4H)-one (**Example Compound 25**)




[0246] Step 1: To a solution of **Example Compound 53** (85 mg, 0.25 mmol) in CH_2Cl_2 (3 mL) was added *m*-CPBA (160 mg, 0.5 mmol). The reaction mixture was stirred at rt for 7 h. The mixture was diluted with methylene chloride (50 mL) and washed with 10% $\text{Na}_2\text{S}_2\text{O}_3$ solution (10

mL), 2N NaOH solution (10 mL) and brine (10 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–70% ethyl acetate/dichloromethane) afforded **Example Compound 24** (60 mg, 67%) as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 8.21 (d, J = 0.9 Hz, 1H), 7.83 (d, J = 0.9 Hz, 1H), 7.40–7.35 (m, 2H), 7.20–7.14 (m, 2H), 5.59 (s, 2H), 2.69 (s, 3H), 2.45 (s, 3H), 2.27 (s, 3H); ESI MS m/z 353 $[\text{M} + \text{H}]^+$.

[0247] Step 2: A solution of **Example Compound 24** (32 mg, 0.091 mmol) in Ac_2O (3 mL) was heated at 130 °C for 2 h. The mixture was concentrated. The residue was diluted with 1:1 $\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (10 mL) and stirred at 80 °C for 10 h. The reaction mixture was concentrated. Purification by chromatography (silica gel, 0–5% methanol/dichloromethane) afforded **Example Compound 25** (20 mg, 63%) as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 12.0 (s, 1H), 8.07 (s, 1H), 7.36–7.31 (m, 2H), 7.19–7.13 (m, 2H), 5.45 (s, 2H), 2.30 (s, 6H), 2.14 (s, 3H); ESI MS m/z 353 $[\text{M} + \text{H}]^+$.

Preparation of 4-(3-benzyl-3H-imidazo[4,5-b]pyridin-5-yl)-3,5-dimethylisoxazole (Example Compound 26)



[0248] Step 1: To a solution of **50** (560 mg, 2.57 mmol) in CH_3CN (15 mL) was added K_2CO_3 (887 mg, 6.43 mmol) and benzyl chloride (484 mg, 2.83 mmol). The reaction was heated at 60 °C for 16 h. The mixture was diluted with ethyl acetate (100 mL), filtered and concentrated to give **51** (790 mg, 100%) as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 8.58 (br s, 1H), 8.24 (d, J = 8.4 Hz, 1H), 7.46–7.35 (m, 5H), 6.82 (d, J = 8.7 Hz, 1H), 4.82 (d, J = 5.7 Hz, 2H).


[0249] Step 2: To a solution of **51** (790 mg, 2.56 mmol) in 1,4-dioxane (25 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (1.14 g, 5.12 mmol), sodium carbonate (2.0 M in H_2O , 3.84 mL, 7.68 mmol) and tetrakis(triphenylphosphine)palladium(0) (300 mg, 0.26 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 8 h. The mixture was diluted with methylene chloride (200 mL) and filtered. The filtrate was concentrated

and purified by chromatography (silica gel, 0–20% EtOAc/hexanes) to afford **52** (500 mg, 60%) as a yellow oil: ^1H NMR (300 MHz, DMSO- d_6) δ 9.09 (t, J = 6.0 Hz, 1H), 8.51 (d, J = 8.4 Hz, 1H), 7.32–7.20 (m, 5H), 6.96 (d, J = 8.7 Hz, 1H), 4.85 (d, J = 6.3 Hz, 2H), 2.47 (s, 3H), 2.25 (s, 3H); ESI m/z 325 $[\text{M} + \text{H}]^+$.

[0250] Step 3: To a solution of **52** (500 mg, 1.54 mmol) in THF (15 mL) and water (12 mL) was added $\text{Na}_2\text{S}_2\text{O}_4$ (1.61 g, 9.24 mmol). The mixture was stirred at rt for 5 h; 2 N HCl (10 mL) was added, and the mixture was heated to reflux for 15 minutes then cooled to rt. Na_2CO_3 was added slowly to adjust to pH 9. The mixture was extracted with ethyl acetate (100 mL), the organic layer was washed with brine (50 mL), filtered and concentrated to give **53** (460 mg, 100%) as a brown oil: ^1H NMR (300 MHz, DMSO- d_6) δ 7.33–7.18 (m, 5H), 6.78 (d, J = 7.5 Hz, 1H), 6.52 (d, J = 7.5 Hz, 1H), 6.29 (t, J = 5.7 Hz, 1H), 4.94 (s, 2H), 4.60 (d, J = 5.7 Hz, 2H), 2.36 (s, 3H), 2.17 (s, 3H); ESI m/z 295 $[\text{M} + \text{H}]^+$.

[0251] Step 4: A solution of **53** (150 mg, 0.51 mmol), trimethylorthoformate (81 mg, 0.765 mmol) and sulfamic acid (3 mg) in MeOH (5 mL) was heated to reflux for 4 h. The mixture was concentrated, the residue was purified by chromatography (silica gel, 30–100% ethyl acetate/hexanes) to afford **Example Compound 26** (100 mg, 65%) as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 8.67 (s, 1H), 8.17 (d, J = 8.1 Hz, 1H), 7.44 (d, J = 8.1 Hz, 1H), 7.36–7.27 (m, 5H), 5.52 (s, 2H), 2.54 (s, 3H), 2.34 (s, 3H); ESI m/z 305 $[\text{M} + \text{H}]^+$.

Preparation of 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-benzo[d]imidazol-4-amine (Example Compound 27), 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N-methyl-1H-benzo[d]imidazol-4-amine (Example Compound 28) and 6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N,N-dimethyl-1H-benzo[d]imidazol-4-amine (Example Compound 29)




[0252] Example Compound 27 was made followed by the similar procedure described for **Example 7**: ^1H NMR (300 MHz, DMSO- d_6) δ 8.23 (s, 1H), 7.42 (dd, J = 8.0, 6.0 Hz, 2H), 7.17 (dd, J = 9.0, 9.0 Hz, 2H), 6.62 (s, 1H), 6.32 (s, 1H), 5.40 (s, 4H), 2.33 (s, 3H), 2.16 (s, 3H); ESI m/z 337 $[\text{M} + \text{H}]^+$.

[0253] To a solution of **Example Compound 27** (35 mg, 0.10 mmol) in methylene chloride (5 mL), was added a 37% solution of formaldehyde in water (8.5 μL) and acetic acid (1 drop). The solution was stirred for 45 min, sodium triacetoxyborohydride (66 mg, 0.31 mmol) was added and

the mixture stirred for 16 h. The mixture was diluted with methylene chloride (20 mL) and neutralized with saturated sodium bicarbonate (5 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated *in vacuo*. The residue was purified by chromatography (silica gel, 0–75% ethyl acetate/methylene chloride) to afford **Example Compound 28** as a white solid (8 mg, 22%) and **Example Compound 29** as a clear solid (7 mg, 18%). **Example Compound 28**: ^1H NMR (500 MHz, DMSO- d_6) δ 8.22 (s, 1H), 7.43 (dd, J = 8.8, 5.5 Hz, 2H), 7.16 (dd, J = 8.8, 5.5 Hz, 2H), 6.65 (d, J = 1.0 Hz, 1H), 6.09 (d, J = 1.0 Hz, 1H), 5.85 (q, J = 5.0 Hz, 1H), 5.41 (s, 2H), 2.83 (d, J = 5.5 Hz, 3H), 2.35 (s, 3H), 2.17 (s, 3H); ESI m/z 351 $[\text{M} + \text{H}]^+$; **Example 29**: ^1H NMR (500 MHz, DMSO- d_6) δ 8.28 (s, 1H), 7.41 (dd, J = 8.5, 5.5 Hz, 2H), 7.17 (dd, J = 9.0, 9.0 Hz, 2H), 6.85 (d, J = 1.0 Hz, 1H), 6.25 (d, J = 1.0 Hz, 1H), 5.43 (s, 2H), 3.18 (s, 6H), 2.35 (s, 3H), 2.18 (s, 3H); ESI m/z 365 $[\text{M} + \text{H}]^+$.

Preparation of 4-(1-benzyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 30)



[0254] Step 1: To a suspension of 3-amino-5-bromo-2-nitropyridine (**54**, 780 mg, 3.58 mmol) and potassium carbonate (2.28 g, 16.5 mmol) in dry acetonitrile (50 mL) was added 1-(bromoethyl)benzene (1.22 g, 6.60 mmol). The mixture was heated to 80 °C for 48 h then water (20 mL) and ethyl acetate (20 mL) were added. The layers were separated and the aqueous layer was extracted with ethyl acetate (2 × 20 mL). The combined ethyl acetate fractions were dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 0–40% ethyl acetate in hexanes) to afford **55** (219 mg, 19%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 8.14 (d, J = 5.0 Hz, 1H), 7.84 (d, J = 2.0 Hz, 1H), 7.40–7.29 (m, 6H), 4.64 (quint, J = 6.5 Hz, 1H), 1.67 (d, J = 7.0 Hz, 3H).


[0255] Step 2: To a mixture of **55** (261 mg, 0.81 mmol) and **3** (217 mg, 0.97 mmol) in 1,4-dioxane (7 mL) and water (1.5 mL) was added potassium carbonate (224 mg, 1.62 mmol) and

tetrakis(triphenylphosphine)palladium(0) (47 mg, 0.04 mmol). The reaction was stirred and heated at 90 °C for 17 h. The reaction mixture was diluted with methanol (20 mL) and silica gel (15 g) was added. The suspension was concentrated to dryness and the resulting powder was loaded onto silica gel and eluted with 0–50% ethyl acetate in hexanes. The clean product was concentrated to give **56** (226 mg, 82%) as a yellow solid: ¹H NMR (500 MHz, CDCl₃) δ 8.19 (d, *J* = 4.5 Hz, 1H), 7.77 (d, *J* = 2.0 Hz, 1H), 7.40–7.28 (m, 5H), 6.89 (d, *J* = 2.0 Hz, 1H), 4.66 (quint, *J* = 5.0 Hz, 1H), 2.10 (s, 3H), 1.94 (s, 3H), 1.71 (d, *J* = 7.0 Hz, 3H).

[0256] Step 3: To a solution of **56** (226 mg, 0.67 mmol) in THF (20 mL) was added a solution of sodium dithionite (698 mg, 4.01 mmol) in water (20 mL) dropwise over 5 min. The solution was stirred at room temperature for 16 h and the solvents were removed *in vacuo*. Methanol (20 mL) was added and the suspension stirred at room temperature for 3 h. The mixture was filtered and the filtrate was concentrated to dryness. A solution of 2N aq. HCl was added to the residue and was heated to reflux for 5 min. After concentration to dryness, methanol was added (10 mL) and the solution was adjusted to pH 8 using saturated aq. NaHCO₃ solution (20 mL). Silica gel was added (10 g) and the suspension was concentrated to dryness. The resulting powder was loaded onto silica gel and eluted with 0–70% ethyl acetate in hexanes. The clean product was concentrated to give **57** (96 mg, 47%) as a beige solid: ¹H NMR (500 MHz, CDCl₃) δ 7.42 (d, *J* = 2.0 Hz, 1H), 7.33–7.30 (m, 4H), 7.25–7.22 (m, 1H), 6.34 (d, *J* = 1.5 Hz, 1H), 4.44 (quint, *J* = 5.0 Hz, 1H), 4.36 (br s, 2H), 3.70 (br s, 1H), 2.07 (s, 3H), 1.89 (s, 3H), 1.58 (d, *J* = 6.5 Hz, 3H).

[0257] Step 4: A mixture of **57** (47 mg, 0.15 mmol), trimethylorthoformate (2 mL, 18.3 mmol) and sulfamic acid (1 mg) were heated in a sealed tube at 100 °C for 30 min. The mixture was cooled, concentrated and loaded onto silica gel and eluted with 0–20% ethyl acetate in hexanes. The resulting material was purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH₃CN in H₂O to afford (**Example Compound 30**) (19 mg, 39%) as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.76 (s, 1H), 8.36 (d, *J* = 2.0 Hz, 1H), 7.65 (d, *J* = 2.5 Hz, 1H), 7.40–7.30 (m, 5H), 4.44 (q, *J* = 7.0 Hz, 1H), 2.29 (s, 3H), 2.10 (s, 3H), 2.06 (d, *J* = 7.0 Hz, 3H). ESI *m/z* 319 [M + H]⁺.

Preparation of 4-(1-benzyl-1*H*-imidazo[4,5-*c*]pyridin-6-yl)-3,5-dimethylisoxazole (**Example Compound 31**), 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*c*]pyridine 5-oxide (**Example 32**) and 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*c*]pyridin-4-amine (**Example Compound 33**)



[0258] Step 1: To a solution of **58** (1.00 g, 5.76 mmol) in 1,4-dioxane (40 mL) and water (4 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (1.93 g, 8.64 mmol), potassium carbonate (1.59 g, 11.5 mmol), and tetrakis(triphenylphosphine)palladium(0) (333 mg, 0.288 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C overnight. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to afford **59** (1.42 g, >99%) as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 9.26 (s, 1H), 6.67 (s, 1H), 6.90–6.00 (bs, 2H), 2.61 (s, 3H), 2.44 (s, 3H); ESI m/z 235 $[\text{M} + \text{H}]^+$.

[0259] Step 2: A mixture of **59** (710 mg, 3.03 mmol), benzyl bromide (778 mg, 4.55 mmol), and potassium carbonate (836 mg, 6.06 mmol) in acetonitrile (30 mL) was heated in sealed tube at 90 °C overnight. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–30% ethyl acetate in hexanes) to afford **60** (303 mg, 30%) as a brown solid: ^1H NMR (500 MHz, CDCl_3) δ 9.26 (s, 1H), 8.68 (s, 1H), 7.50–7.10 (m, 5H), 6.50 (s, 1H), 4.65 (d, $J = 4.1$ Hz, 2H), 2.39 (s, 3H), 2.19 (s, 3H); ESI m/z 325 $[\text{M} + \text{H}]^+$.

[0260] Step 3: To a solution of **60** (300 mg, 0.926 mmol) in tetrahydrofuran (10 mL) was added sodium dithionite (967 mg, 5.56 mmol) in water (10 mL). The reaction mixture was stirred at room temperature overnight and concentrated under vacuum. The residue was suspended in MeOH

and the solid was filtered, washed with MeOH, and the filtrate concentrated under vacuum. To the residue was added 2N HCl and heated to just boiling, cooled to room temperature and concentrated under vacuum. The residue was dissolved in MeOH and basified with 10% NaHCO₃, concentrated and purified by chromatography (silica gel, 0–20% methanol in ethyl acetate) to afford **61** (150 mg, 55%) as a gray solid: ¹H NMR (500 MHz, CDCl₃) δ 7.99 (s, 1H), 7.40–7.28 (m, 5H), 6.39 (s, 1H), 4.64 (s, 1H), 4.43 (d, *J* = 5.4 Hz, 2H), 3.15 (s, 2H), 2.33 (s, 3H), 2.21 (s, 3H); ESI *m/z* 295 [M + H]⁺.

[0261] Step 4: To a solution of **61** (150 mg, 0.51 mmol) in ethanol (5 mL) was added trimethylorthoformate (81 mg, 0.77 mmol) and sulfamic acid (1 mg, 0.01 mmol). The reaction was heated in a sealed tube at 90 °C overnight. The mixture was concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give **Example Compound 31** (143 mg, 92%) as a yellow solid: ¹H NMR (500 MHz, CD₃OD) δ 9.00 (d, *J* = 1.0 Hz, 1H), 8.05 (s, 1H), 7.48 (d, *J* = 1.0 Hz, 1H), 7.40–7.30 (m, 5H), 5.58 (s, 2H), 2.40 (s, 3H), 2.25 (s, 3H); ESI *m/z* 305 [M + H]⁺.


[0262] Step 5: To a mixture of **Example Compound 31** (100 mg, 0.329 mmol) in dichloromethane (5 mL) was added 3-chloroperoxybenzoic acid (264 mg, 77% with water, 1.18 mmol). The mixture was stirred at room temperature overnight, concentrated and purified by chromatography (silica gel, 0–20% methanol in ethyl acetate) to afford **Example Compound 32** (127 mg, >99%) as an off-white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.92 (s, 1H), 8.61 (s, 1H), 7.67 (s, 1H), 7.45–7.25 (m, 5H), 6.57 (s, 2H), 2.28 (s, 3H), 2.17 (s, 3H); ESI *m/z* 321 [M + H]⁺.

[0263] Step 6: To a mixture of phosphorus oxybromide (268 mg, 0.938 mmol) in DMF (2 mL) was added **Example 32** (100 mg, 0.313 mmol) in DMF (6 mL). The mixture was stirred at room temperature for 10 min and heated at 100 °C for 1 h. After cooling to room temperature, water and MeOH were added. The mixture was neutralized to pH 7 by addition of 10% sodium bicarbonate and concentrated. The residue was purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to afford **62** (30 mg, 25%) as an off-white solid: ¹H NMR (500 MHz, CDCl₃) δ 8.09 (s, 1H), 7.43–7.35 (m, 3H), 7.23–7.19 (m, 2H), 7.03 (s, 1H), 5.38 (s, 2H), 2.47 (s, 3H), 2.31 (s, 3H); ESI *m/z* 383 [M + H]⁺.

[0264] Step 7: To a solution of **62** (30 mg, 0.078 mmol) in toluene (10 mL) under nitrogen atmosphere was added *tert*-butyl carbamate (27 mg, 0.23 mmol), cesium carbonate (51 mg, 0.16 mmol), 2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl (6 mg, 0.01 mmol) and tris(dibenzylideneacetone) dipalladium(0) (7 mg, 0.008 mmol). The reaction mixture was heated at 90 °C overnight, cooled to room temperature, and purified by chromatography (silica gel, 0–20% methanol in ethyl acetate). It was further purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH₃CN in H₂O to give **Example Compound 33** (10 mg, 40%) as an off-white solid: ¹H

NMR (500 MHz, CD₃OD) δ 8.21 (s, 1H), 7.42–7.25 (m, 5H), 6.70 (s, 1H), 5.46 (s, 2H), 2.39 (s, 3H), 2.24 (s, 3H); HPLC 96.9%, t_R = 10.1 min; ESI m/z 320 [M + H]⁺.


Preparation of 4-(1-benzyl-3-bromo-1*H*-pyrrolo[3,2-*b*]pyridin-6-yl)-3,5-dimethylisoxazole.
(Example Compound Compound 34)



[0265] Step 1: To a solution of **46** (1.0 g, 5.08 mmol) in 1,4-dioxane (50 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (1.47 g, 6.6 mmol), sodium carbonate (1.10 g in 8 mL H₂O, 10.2 mmol) and tetrakis(triphenylphosphine)palladium(0) (587 mg, 0.51 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 16 h. The mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded **63** (850 mg, 79%) as a yellow solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.4 (s, 1H), 8.30 (t, J = 2.1 Hz, 1H), 7.75 (dd, J = 1.8, 0.9 Hz, 1H), 7.70 (t, J = 3.0 Hz, 1H), 6.61–6.59 (m, 1H), 2.42 (s, 3H), 2.24 (s, 3H).

[0266] Step 2/3: To a solution of **63** (500 mg, 2.35 mmol) in DMF (10 mL) at 0 °C was added NBS (500 mg, 2.82 mmol). The reaction mixture was stirred at 0 °C for 2 h. The mixture was diluted with methylene chloride (50 mL) and washed with brine (20 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The crude **64** was carried forward. To a solution of **64** (300 mg, 1.03 mmol) in DMF (1 mL) and CH₃CN (10 mL) was added potassium carbonate (283 mg, 2.06 mmol) and benzyl chloride (130 mg, 1.03 mmol). The reaction was stirred at 70 °C for 16 h. The mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography silica gel, 0–50% ethyl acetate/dichloromethane) afforded **Example Compound 34** (200 mg, 51%) as an off-white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.33 (d, J = 1.5 Hz, 1H), 7.86 (s, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.34–7.24 (m, 5H), 5.48 (s, 2H), 2.35 (s, 3H), 2.17 (s, 3H); ESI MS m/z 382 [M + H]⁺.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridine-3-carbaldehyde (Example Compound 35)




Example 35

[0267] Step 1: To a mixture of **46** (300 mg, 1.5 mmol) and hexamethylenetetramine (0.32 g, 2.25 mmol) was added AcOH (2 mL). The reaction mixture was stirred at 120 °C for 6 h and was quenched with H₂O (5 mL). The precipitate was collected by filtration to afford **65** (190 mg, 56%) as a yellow solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.4 (s, 1H), 10.1 (s, 1H), 8.58 (d, *J* = 2.1 Hz, 1H), 8.47 (s, 1H), 8.18 (d, *J* = 2.1 Hz, 1H).

[0268] Step 2: To a solution of **65** (190 mg, 0.84 mmol) in 1,4-dioxane (5 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (245 mg, 1.09 mmol), sodium carbonate (178 mg in 1 mL H₂O, 1.68 mmol) and tetrakis(triphenylphosphine)palladium(0) (97 mg, 0.08 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 16 h. The mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded **66** (135 mg, 67%) as an off-white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.5 (s, 1H), 10.2 (s, 1H), 8.51 (d, *J* = 1.8 Hz, 1H), 8.49 (d, *J* = 3.0 Hz, 1H), 7.92 (d, *J* = 1.8 Hz, 1H), 2.44 (s, 3H), 2.26 (s, 3H); ESI MS *m/z* 242 [M + H]⁺.


[0269] Step 3: To a solution of **66** (92 mg, 0.38 mmol) in DMF (0.5 mL) and CH₃CN (5 mL) was added potassium carbonate (105 mg, 0.76 mmol) and benzyl chloride (58 mg, 0.46 mmol). The reaction was stirred at 70 °C for 16 h. The reaction mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded **Example Compound 35** (72 mg, 57%) as an off-white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.2 (s, 1H), 8.73 (s, 1H), 8.53 (d, *J* = 1.8 Hz, 1H), 8.11 (d, *J* = 1.8 Hz, 1H), 7.44–7.30 (m, 5H), 5.59 (s, 2H), 2.40 (s, 3H), 2.21 (s, 3H); ESI MS *m/z* 332 [M + H]⁺.

Preparation of 1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-pyrrolo[3,2-*b*]pyridin-3-yl)-*N,N*-dimethylmethanamine (Example Compound 72)



[0270] A solution of **Example Compound 35** (54 mg, 0.16 mmol), dimethylamine (0.25 mL, 2M in THF, 0.49 mmol) and NaBH(OAc)₃ (104 mg, 0.49 mmol) in CH₂Cl₂ (3 mL) was stirred at room temperature for 16 h. The reaction mixture was concentrated under reduced pressure. The crude reaction mixture was purified by chromatography (silica gel, 0–10% methanol/dichloromethane) to provide **Example Compound 72** (42 mg, 71%) as an off-white solid: ¹H NMR (300 MHz, CDCl₃) δ 8.34 (d, *J* = 1.8 Hz, 1H), 8.30 (s, 1H), 7.36–7.32 (m, 4H), 7.21–7.18 (m, 2H), 5.39 (s, 2H), 4.50 (s, 2H), 2.86 (s, 6H), 2.32 (s, 3H), 2.16 (s, 3H); ESI MS *m/z* 361 [M + H]⁺.

Preparation of 1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-pyrrolo[3,2-*b*]pyridin-3-yl)ethanone (Example Compound 36)




[0271] Step 1: To a suspension of AlCl₃ (313 mg, 2.35 mmol) in CH₂Cl₂ (20 mL) was added **63** (100 mg, 0.47 mmol) and AcCl (184 mg, 2.35 mmol). The reaction mixture was stirred at rt for 6 h. The reaction was quenched with methanol (10 mL) carefully and the pH adjusted to neutral with solid Na₂CO₃. The mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–10% methanol/dichloromethane) afforded **67** (82 mg, 68%) as an off-white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.8 (s, 1H), 8.67 (s, 1H), 8.57 (s, 1H), 8.21 (s, 1H), 2.71 (s, 3H), 2.45 (s, 3H), 2.26 (s, 3H); ESI MS *m/z* 256 [M + H]⁺.

[0272] Step 2: To a solution of **67** (62 mg, 0.24 mmol) in DMF (0.5 mL) and CH₃CN (5 mL) was added potassium carbonate (67 mg, 0.48 mmol) and benzyl chloride (37 mg, 0.29 mmol). The reaction was stirred at 70 °C for 16 h. The reaction mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl


acetate/dichloromethane) afforded **Example Compound 36** (30 mg, 36%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 8.59 (d, $J = 1.5$ Hz, 1H), 8.22 (s, 1H), 7.45 (d, $J = 1.8$ Hz, 1H), 7.40–7.36 (m, 3H), 7.21–7.18 (m, 2H), 5.40 (s, 2H), 2.89 (s, 3H), 2.34 (s, 3H), 2.17 (s, 3H); ESI MS m/z 346 $[\text{M} + \text{H}]^+$.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-5-yl formate (Example Compound 37)



[0273] **Step 1:** A solution of **Example Compound 56** (165 mg, 0.52 mmol) in DMF (2 mL) was added POCl_3 (159 mg, 1.03 mmol). The reaction mixture was heated at 100 °C for 2 h and concentrated. The residue was dissolved in CH_2Cl_2 (100 mL), washed with saturated NaHCO_3 (2 $\times$ 20 mL) and brine (20 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded **Example Compound 37** (81 mg, 45%) as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 9.90 (s, 1H), 7.62 (s, 1H), 7.43–7.41 (m, 3H), 7.28 (s, 1H), 7.22–7.18 (m, 3H), 5.31 (s, 2H), 2.22 (s, 3H), 2.10 (s, 3H); ESI MS m/z 348 $[\text{M} + \text{H}]^+$.


Preparation of 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benzamide (Example Compound 38)



[0274] To a solution of **Example Compound 70** (100 mg, 0.29 mmol) in ethanol (3 mL) was added 2N sodium hydroxide in water (1.46 mL, 2.9 mmol). The mixture was heated to 85 °C for 20 min, then cooled to room temperature, and neutralized with 2 mL of acetic acid. The mixture was basified (pH 8) with solid sodium carbonate, diluted in methylene chloride (100 mL), washed

with brine (20 mL), and dried over anhydrous sodium sulfate. After filtration, the filtrate was concentrated *in vacuo* and purified by chromatography (silica gel, 0–20% methanol/methylene chloride) to afford **Example Compound 38** as a white solid (71 mg, 68%): ^1H NMR (300 MHz, DMSO- d_6) δ 8.35 (d, J = 1.8 Hz, 1H), 7.99 (d, J = 2.1 Hz, 1H), 7.94 (br s, 1H), 7.83 (d, J = 8.4 Hz, 2H), 7.37 (br s, 1H), 7.27 (d, J = 8.4 Hz, 2H), 5.61 (s, 2H), 2.60 (s, 3H), 2.39 (s, 3H), 2.21 (s, 3H); ESI m/z 362 $[\text{M} + \text{H}]^+$.


Preparation of 4-(1-benzyl-3-nitro-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 39)



[0275] Step 1: To a solution of **63** (100 mg, 0.47 mmol) in H_2SO_4 (0.5 mL) at 0 °C was added HNO_3 (35 mg, 0.47 mmol). The reaction mixture was stirred at 0 °C for 1 h. The reaction mixture was diluted with H_2O (10 mL) and adjusted to neutral pH with 6N NaOH solution. The solution was extracted with CH_2Cl_2 (30 mL). The organic layer was dried, filtered and concentrated. Purification by chromatography (silica gel, 0–10% methanol/dichloromethane) afforded **68** (82 mg, 68%) as a yellow solid: ^1H NMR (300 MHz, DMSO- d_6) δ 12.9 (s, 1H), 8.85 (s, 1H), 8.58 (d, J = 2.1 Hz, 1H), 7.95 (d, J = 1.8 Hz, 1H), 2.45 (s, 3H), 2.26 (s, 3H); ESI MS m/z 259 $[\text{M} + \text{H}]^+$.

[0276] Step 2: To a solution of **68** (82 mg, 0.32 mmol) in DMF (0.5 mL) and CH_3CN (5 mL) was added potassium carbonate (88 mg, 0.64 mmol) and benzyl chloride (44 mg, 0.35 mmol). The reaction was stirred at 70 °C for 16 h. The reaction mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–50% ethyl acetate/dichloromethane) afforded **Example Compound 39** (68 mg, 61%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 8.74 (s, 1H), 8.47 (s, 1H), 7.56 (s, 1H), 7.45–7.42 (m, 3H), 7.27–7.26 (m, 2H), 5.47 (s, 2H), 2.35 (s, 3H), 2.17 (s, 3H); ESI MS m/z 349 $[\text{M} + \text{H}]^+$.

**Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-ethoxy-1*H*-benzo[*d*]imidazol-4-amine
(Example Compound 17)**



[0277] Step 1: A mixture of **37** (200 mg, 0.709 mmol) in tetraethoxymethane (340 mg, 1.77 mmol) was heated at 100°C for 4 h. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–50% ethyl acetate in hexanes) to afford **69** (177 mg, 74%) as a yellow solid: ^1H NMR (500 MHz, CD_3OD) δ 7.30–7.15 (m, 2H), 4.57 (q, $J = 7.0$ Hz, 2H), 2.39 (s, 3H), 2.23 (s, 3H), 1.47 (t, $J = 7.0$ Hz, 3H); ESI m/z 336 $[\text{M} + \text{H}]^+$.

[0278] Step 2: To a solution of **69** (250 mg, 0.74 mmol) in CH_3CN (8 mL) and DMF (2 mL) was added K_2CO_3 (155 mg, 0.82 mmol) and benzyl chloride (104 mg, 0.82 mmol). The reaction was heated at 60°C for 16 h. The mixture was diluted with ethyl acetate (100 mL), filtered and concentrated. The residue was purified by chromatography (silica gel, 0–30% EtOAc/hexanes) to afford **70** (200 mg, 63%) as an off-white solid and **71** (87 mg, 27%) as a colorless oil: **70**: ^1H NMR (300 MHz, CDCl_3) δ 7.34–7.29 (m, 3H), 7.21–7.18 (m, 3H), 6.77 (d, $J = 1.5$ Hz, 1H), 5.16 (s, 2H), 4.75 (q, $J = 7.5$ Hz, 2H), 2.29 (s, 3H), 2.14 (s, 3H), 1.50 (t, $J = 7.0$ Hz, 3H); **71**: ^1H NMR (300 MHz, CDCl_3) δ 7.37 (d, $J = 1.5$ Hz, 1H), 7.34–7.28 (m, 3H), 7.18 (d, $J = 7.5$ Hz, 2H), 7.12 (d, $J = 1.5$ Hz, 1H), 5.60 (s, 2H), 4.63 (q, $J = 7.0$ Hz, 2H), 2.41 (s, 3H), 2.28 (s, 3H), 1.45 (t, $J = 7.0$ Hz, 3H).

[0279] Step 3: A mixture of **70** (100 mg, 0.235 mmol), BocNH_2 (82 mg, 0.705 mmol), Xantphos (28 mg, 0.048 mmol), $\text{Pd}_2(\text{dba})_3$ (22 mg, 0.024 mmol) and Cs_2CO_3 (268 mg, 0.823 mmol) in 1,4-dioxane (8 mL) was purged with nitrogen and heated at 100°C for 18 h. The mixture was diluted with methylene chloride (200 mL) and filtered. The filtrate was concentrated and purified by chromatography (silica gel, 0–30% EtOAc/hexanes) to afford **72** (90 mg, 83%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.74 (br s, 1H), 7.41 (s, 1H), 7.32–7.29 (m, 3H), 7.22–7.19 (m, 2H), 6.51 (d, $J = 1.5$ Hz, 1H), 5.14 (s, 2H), 4.64 (q, $J = 7.2$ Hz, 2H), 2.32 (s, 3H), 2.17 (s, 3H), 1.49 (t, $J = 7.2$ Hz, 3H), 1.46 (s, 9H).

[0280] Step 4: A solution of **72** (90 mg, 0.195 mmol) in TFA (1 mL) and CH_2Cl_2 (2 mL) was stirred at rt for 1 h. The mixture was concentrated, the residue was dissolved in ethyl acetate (100 mL) and washed with saturated NaHCO_3 (50 mL $\times$ 2). The organic layer was dried over sodium


sulfate, filtered and concentrated. Purification by chromatography (silica gel, 40–100% EtOAc/hexanes) afforded **Example Compound 17** (51 mg, 72%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.35–7.20 (m, 5H), 6.33 (d, $J = 1.5$ Hz, 1H), 6.30 (d, $J = 1.5$ Hz, 1H), 5.13 (s, 2H), 4.68 (q, $J = 6.9$ Hz, 2H), 4.30 (br s, 2H), 2.30 (s, 3H), 2.16 (s, 3H), 1.49 (t, $J = 7.2$ Hz, 3H); ESI m/z 363 $[\text{M} + \text{H}]^+$.

Preparation of 4-(1-benzyl-2-ethoxy-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 59)



[0281] To a mixture of **28** (50 mg, 0.17 mmol) and tetraethyl orthocarbonate (131 mg, 0.68 mmol) was added sulfamic acid (3 mg, 0.034 mmol). The mixture was then heated to 100 °C for 8 h, then diluted with ethyl acetate (30 mL), washed with brine (15 mL), dried over anhydrous sodium sulfate, and concentrated *in vacuo*. The residue was purified by chromatography (silica gel, 0 – 10% methanol/methylene chloride) to afford **Example Compound 59** (24 mg, 41%) as an off-white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.75 (d, $J = 1.2$ Hz, 1H), 7.38–7.22 (m, 5H), 7.18 (d, $J = 1.5$ Hz, 1H), 4.99 (s, 2H), 4.34 (q, $J = 7.2$ Hz, 2H), 2.37 (s, 3H), 2.18 (s, 3H), 1.42 (t, $J = 7.2$ Hz, 3H); ESI m/z 349 $[\text{M} + \text{H}]^+$.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazole-4-carbonitrile (Example Compound 85)




[0282] Compound **73** was prepared by following the method for General Procedure J steps 1 to 3 starting with 2-amino-5-bromobenzonitrile. Using the procedure used for General Procedure D step 3 on compound **73** (30 mg, 0.09 mmol) afforded **Example Compound 85** (10 mg, 31%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.63 (d, $J = 1.5$ Hz, 1H), 7.60 (d, $J = 1.5$ Hz,

1H), 7.38–7.27 (m, 3H), 7.19–7.14 (m, 2H), 5.57 (s, 2H), 2.69 (s, 3H), 2.32 (s, 3H), 2.16 (s, 3H); ESI m/z 343 $[M + H]^+$.

General Procedure O:


Preparation of *N*-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-4-yl)acetamide (Example Compound 111)



[0283] A solution of **Example Compound 16** (34 mg, 0.10 mmol), acetic anhydride (12 mg, 0.12 mmol) and *i*-Pr₂NEt (26 mg, 0.20 mmol) in THF (3 mL) was stirred at rt for 16 h. The mixture was concentrated, and the residue was purified by chromatography (silica gel, 0–5% methanol/EtOAc) to afford **Example Compound 111** (28 mg, 74%) as a white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.78 (s, 1H), 9.85 (s, 1H), 7.60–7.46 (m, 5H), 7.28 (d, *J* = 1.2 Hz, 1H), 7.06 (d, *J* = 1.2 Hz, 1H), 5.22 (s, 2H), 2.51 (s, 3H), 2.33 (s, 3H), 2.27 (s, 3H); ESI m/z 377 $[M + H]^+$.

General Procedure P:

Preparation of 6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one (**Example Compound 110**) and 4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one (**Example Compound 115**)



[0284] **Step 1:** To a solution of **30** (1.00 g, 3.21 mmol) in toluene (70 mL) under nitrogen atmosphere was added benzyl amine (1.94 g, 16.0 mmol), potassium *tert*-butoxide (539 mg, 4.82 mmol), 2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl (229 mg, 0.482 mmol), and tris(dibenzylideneacetone) dipalladium(0) (293 mg, 0.321 mmol). The reaction mixture was heated at 90 °C overnight, cooled to room temperature, and purified by chromatography (silica gel, 0–50%


ethyl acetate in hexanes) to afford **74** (700 mg, 62%) as a red-brown solid: ^1H NMR (500 MHz, CDCl_3) δ 7.50 (d, $J = 1.8$ Hz, 1H), 7.70–7.22 (m, 5H), 6.41 (d, $J = 1.6$ Hz, 1H), 6.07 (s, 2H), 4.48 (q, $J = 3.5$ Hz, 1H), 3.65 (s, 1H), 2.05 (s, 3H), 1.90 (s, 3H), 1.62 (d, $J = 6.6$ Hz, 3H); ESI m/z 353 $[\text{M} + \text{H}]^+$.

[0285] Step 2: To a mixture of **74** (600 mg, 1.70 mmol) in 1,4-dioxane (40 mL) was added 1,1'-carbonyldiimidazole (2.76 mg, 17.0 mmol) and DMAP (a crystal). The reaction was heated in a sealed tube at 120 °C for 2 days. The mixture was concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give **Example Compound 110** (420 mg, 65%) as an orange solid: ^1H NMR (500 MHz, CD_3OD) δ 7.75 (d, $J = 1.3$ Hz, 1H), 7.44 (d, $J = 7.7$ Hz, 2H), 7.38 (t, $J = 7.7$ Hz, 2H), 7.31 (t, $J = 7.7$ Hz, 1H), 6.88 (d, $J = 1.3$ Hz, 1H), 5.88 (q, $J = 7.1$ Hz, 1H), 2.20 (s, 3H), 2.02 (s, 3H), 1.91 (d, $J = 7.2$ Hz, 3H); ESI m/z 377 $[\text{M} - \text{H}]^+$.

[0286] Step 3: To a solution of **Example Compound 110** (100 mg, 0.265 mmol) in tetrahydrofuran (10 mL) was added sodium dithionite (276 mg, 1.59 mmol) in water (10 mL). The reaction mixture was stirred at room temperature overnight and concentrated under vacuum. The residue was added 2N HCl and heated to just boiling, cooled to room temperature, and concentrated in vacuum. The residue was dissolved in MeOH and basified by conc. NH_4OH , concentrated, and purified by chromatography (silica gel, 0–100% hexanes/ethyl acetate). It was further purified by reverse phase HPLC on a Polaris C_{18} column eluted with 10–90% CH_3CN in H_2O to give **Example Compound 115** (49 mg, 53%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.42–7.32 (m, 4H), 7.26 (t, $J = 6.9$ Hz, 1H), 6.35 (s, 1H), 5.94 (s, 1H), 5.78 (q, $J = 7.2$ Hz, 1H), 2.17 (s, 3H), 2.00 (s, 3H), 1.86 (d, $J = 7.2$ Hz, 3H); ESI m/z 349 $[\text{M} + \text{H}]^+$.

General Procedure Q:

Preparation of 4-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine (**Example Compound 114**)




[0287] A mixture of **Example Compound 10** (90 mg, 0.28 mmol) and phosphorus (V) oxychloride (1 mL) was heated to 110 °C for 5 h, then cooled to room temperature. The mixture was concentrated, dissolved with methylene chloride (75 mL), and washed with saturated sodium bicarbonate solution (20 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was dissolved in a 2.0 M solution of morpholine in tetrahydrofuran (5.6 mL, 11.2 mmol) and the mixture was heated to 75 °C for 3 h. The reaction mixture was

concentrated, and the residue was purified by chromatography (silica gel, 0–5% methanol/methylene chloride), and then triturated with ethyl acetate/hexanes to afford **Example Compound 114** (62 mg, 57%) as a white solid: ^1H NMR (500 MHz, CDCl_3) δ 8.24 (d, J = 2.0 Hz, 1H), 7.41–7.34 (m, 3H), 7.15 (d, J = 6.5 Hz, 2H), 7.06 (d, J = 1.0 Hz, 1H), 5.26 (s, 2H), 3.83 (t, J = 4.5 Hz, 4H), 3.50 (t, J = 4.5 Hz, 4H), 2.29 (s, 3H), 2.11 (s, 3H); ESI m/z 390 $[\text{M} + \text{H}]^+$.

General Procedure R:

Preparation of 1-(3,4-dichlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (Example Compound 101)




[0288] Compound 75 was prepared according to General Procedure D, steps 1-2.

[0289] To a solution of 75 (218 mg, 0.60 mmol) in 1,4-dioxane (5 mL) was added 1,1'-carbonyldiimidazole (117 mg, 0.72 mmol), and the mixture was heated to 100 °C for 16 h. The mixture was diluted with methylene chloride (70 mL), and washed with brine (20 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–10% methanol/methylene chloride) to afford **Example Compound 101** (155 mg, 66%) as a white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 11.83 (s, 1H), 7.92 (d, J = 1.5 Hz, 1H), 7.73 (d, J = 2.0 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 7.53 (d, J = 2.0 Hz, 1H), 7.35 (dd, J = 8.5, 2.0 Hz, 1H), 5.05 (s, 2H), 2.37 (s, 3H), 2.19 (s, 3H); ESI m/z 389 $[\text{M} + \text{H}]^+$.

General Procedure S:

Preparation of (S)-3,5-dimethyl-4-(2-methyl-4-nitro-1-(1-phenylethyl)-1*H*-benzo[d]imidazol-6-yl)isoxazole (Example Compound 125) and (S)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1*H*-benzo[d]imidazol-4-amine (Example Compound 143)




[0290] Compound **76** was prepared by following the method of General Procedure P step 1 starting with (S)-1-phenylethanamine.

[0291] **Step 1:** Using the procedure used in General Procedure F step 1 starting with compound **76** (140 mg, 0.40 mmol) afforded **Example Compound 125** (108 mg, 72%) as a yellow solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.87 (d, $J = 1.5$ Hz, 1H), 7.42-7.30 (m, 6H), 6.11 (q, $J = 7.2$ Hz, 1H), 2.74 (s, 3H), 2.23 (s, 3H), 2.04 (s, 3H), 1.94 (d, $J = 6.9$ Hz, 3H); ESI MS m/z 377 $[\text{M} + \text{H}]^+$.

[0292] **Step 2:** Using the procedure used in General Procedure P step 3 starting with compound **Example Compound 125** (80 mg, 0.21 mmol) afforded **Example Compound 143** (53 mg, 72%) as an off-white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.39-7.26 (m, 5H), 6.23 (d, $J = 1.5$ Hz, 1H), 6.14 (d, $J = 1.2$ Hz, 1H), 5.86 (q, $J = 7.2$ Hz, 1H), 5.26 (s, 2H), 2.58 (s, 3H), 2.20 (s, 3H), 2.02 (s, 3H), 1.86 (d, $J = 6.9$ Hz, 3H); ESI MS m/z 347 $[\text{M} + \text{H}]^+$.


General Procedure T:

Preparation of 4-(1-benzyl-2-(pyridin-3-yloxy)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole (**Example Compound 236**)



[0293] A mixture of **Example Compound 10** (100 mg, 0.31 mmol) and phosphorus (V) oxychloride (1 mL) was heated to 110 °C for 5 h, then cooled to room temperature. The mixture was concentrated, dissolved with methylene chloride (75 mL), and washed with saturated sodium bicarbonate solution (20 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was dissolved in *N,N*-dimethylformamide (2.5 mL), 3-hydroxypyridine (109 mg, 1.15 mmol) and potassium carbonate (175 mg, 1.27 mmol) were added. The mixture was heated to 100 °C for 16 h, then diluted with ethyl acetate (75 mL), washed with brine (2 × 25 mL), dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–10% methanol/methylene chloride) to afford **Example Compound 236** (58 mg, 47%) as a light brown solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.74 (d, $J = 2.7$ Hz, 1H), 8.57 (dd, $J = 4.5, 0.9$ Hz, 1H), 8.27 (d, $J = 1.8$ Hz, 1H), 8.02–7.98 (m, 2H), 7.59 (dd, $J = 8.4, 4.5$ Hz, 1H), 7.47 (d, $J = 6.9$ Hz, 2H), 7.42–7.30 (m, 3H), 5.53 (s, 2H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 398 $[\text{M} + \text{H}]^+$.


Preparation of 6-(3,5-dimethylisoxazol-4-yl)-*N*-ethyl-4-nitro-1-(1-phenylethyl)-1*H*-benzo[d]imidazol-2-amine (Example Compound 127) and 6-(3,5-dimethylisoxazol-4-yl)-*N*²-ethyl-1-(1-phenylethyl)-1*H*-benzo[d]imidazole-2,4-diamine (Example Compound 134)



[0294] **Step 1:** To Example Compound 110 (200 mg, 0.529 mmol) was added phosphorus(V) oxychloride (2 mL, 21.5 mmol) and *N,N*-dimethylformamide (one drop). The reaction was heated at 90 °C overnight. The mixture was concentrated, the residue was dissolved in tetrahydrofuran (5 mL), ethylamine (10 mL, 1M in tetrahydrofuran) was added. The reaction mixture was heated in a sealed tube at 70 °C for 2 days. The mixture was concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give Example Compound 127 (40 mg, 19%) as a yellow solid: ¹H NMR (500 MHz, CD₃OD) δ 7.70 (d, *J* = 1.5 Hz, 1H), 7.45–7.30 (m, 5H), 6.72 (d, *J* = 1.5 Hz, 1H), 5.86 (q, *J* = 7.0 Hz, 1H), 3.72 (q, *J* = 7.2 Hz, 2H), 2.17 (s, 3H), 1.98 (s, 3H), 1.90 (d, *J* = 7.0 Hz, 3H), 1.36 (t, *J* = 7.2 Hz, 3H); ESI *m/z* 406 [*M* – H]⁺.

[0295] **Step 2:** To a solution of Example Compound 127 (35 mg, 0.086 mmol) in tetrahydrofuran (10 mL) was added sodium dithionite (90 mg, 0.52 mmol) in water (10 mL). The reaction mixture was stirred at room temperature overnight and concentrated under vacuum. The residue was added 2N HCl and heated to just boiling, cooled to room temperature, and concentrated in vacuum. The residue was dissolved in MeOH and basified by conc. NH₄OH, concentrated, and purified by chromatography (silica gel, 0–100% hexanes/ethyl acetate). It was further purified by reverse phase HPLC on a Polaris C₁₈ column eluting with 10–90% CH₃CN in H₂O to give Example Compound 134 (15 mg, 47%) as an off-white solid: ¹H NMR (500 MHz, CD₃OD) δ 7.40–7.25 (m, 5H), 6.31 (d, *J* = 1.5 Hz, 1H), 5.92 (d, *J* = 1.5 Hz, 1H), 5.72 (q, *J* = 6.9 Hz, 1H), 3.53 (q, *J* = 7.2 Hz, 2H), 2.15 (s, 3H), 1.99 (s, 3H), 1.86 (d, *J* = 7.0 Hz, 3H), 1.33 (t, *J* = 7.2 Hz, 3H); ESI *m/z* 376 [*M* + H]⁺.


Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-4-nitro-1*H*-benzo[d]imidazol-2(3*H*)-one (Example Compound 150) and 4-amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-1*H*-benzo[d]imidazol-2(3*H*)-one (Example Compound 162)



[0296] **Step 1:** A mixture of **Example Compound 15** (73 mg, 0.20 mmol), CH_3I (85 mg, 0.60 mmol) and K_2CO_3 (110 mg, 0.8 mmol) in DMF (3 mL) was stirred at rt for 16 h. The reaction mixture was diluted with EtOAc (100 mL) and washed with brine (3×50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was triturated with EtOAc/hexanes to afford **Example Compound 150** (65 mg, 86%) as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 7.48 (d, $J = 1.5$ Hz, 1H), 7.35–7.30 (m, 5H), 6.84 (d, $J = 1.5$ Hz, 1H), 5.15 (s, 2H), 3.65 (s, 3H), 2.26 (s, 3H), 2.09 (s, 3H); ESI m/z 379 $[\text{M} + \text{H}]^+$.


[0297] **Step 2:** To a solution of **Example Compound 150** (57 mg, 0.15 mmol) in THF (5 mL) and water (4 mL) was added $\text{Na}_2\text{S}_2\text{O}_4$ (153 mg, 0.90 mmol). The mixture was stirred at rt for 4 h, 2N HCl (1 mL) was added, the mixture was heated to reflux for 15 minutes. After cooled to rt, Na_2CO_3 was added slowly to adjust to pH 9. The mixture was extracted with CH_2Cl_2 (100 mL), the organic layer was washed with brine (50 mL), filtered, concentrated and purified by chromatography (silica gel, 0–10% methanol/ethyl acetate) to afford **Example Compound 162** (60 mg, 72%) as a white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.36–7.24 (m, 5H), 6.40 (d, $J = 1.5$ Hz, 1H), 6.39 (d, $J = 1.8$ Hz, 1H), 5.08 (s, 2H), 4.99 (s, 2H), 3.62 (s, 3H), 2.29 (s, 3H), 2.12 (s, 3H); ESI m/z 349 $[\text{M} + \text{H}]^+$. HPLC >99%

Preparation of 4-(1-benzyl-2-methyl-4-(methylsulfonyl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole (Example Compound 168)



[0298] A mixture of **Example Compound 121** (100 mg, 0.25 mmol), sodium methanesulfonate (39 mg, 0.38 mmol), CuI (5 mg, 0.025 mmol), L-proline (6 mg, 0.05 mmol) and NaOH (2 mg, 0.05 mmol) in DMSO (3 mL) was heated at 150 °C in a microwave reactor for 2 h. The mixture was diluted with ethyl acetate (100 mL) and washed with brine (50 mL). The organic layer was dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 50–100% EtOAc/hexanes) to afford **Example Compound 168** (13 mg, 13%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 7.75 (d, $J = 1.5$ Hz, 1H), 7.37–7.33 (m, 3H), 7.24 (d, $J = 1.5$ Hz, 1H), 7.11–7.08 (m, 2H), 5.39 (s, 2H), 3.54 (s, 3H), 2.73 (s, 3H), 2.31 (s, 3H), 2.16 (s, 3H); ESI m/z 396 $[\text{M} + \text{H}]^+$. HPLC 92.3%.

Preparation of 4-(1-benzyl-2,7-dimethyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 181)




[0299] Step 1: To a solution of 77 (4.4 g, 16.5 mmol) in 1,4-dioxane (100 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (4.4 g, 19.8 mmol), Na_2CO_3 (2.0 M in H_2O , 25 mL, 50.0 mmol) and tetrakis(triphenylphosphine)palladium(0) (959 mg, 0.83 mmol). The reaction mixture was purged with nitrogen and heated at 80 °C for 16 h. The mixture was diluted with EtOAc (100 mL) and washed with brine (50 mL). The organic layer was dried over Na_2SO_4 and filtered. The filtrate was concentrated and then purified by chromatography (silica gel, 0–60% ethyl acetate/hexanes) to afford 78 (2.64 g, 57%) as an off-white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.71 (s, 1H), 6.32 (s, 2H), 2.22 (s, 3H), 2.08 (s, 3H), 2.02 (s, 3H).

[0300] Step 2: A mixture of 78 (1.3 g, 4.61 mmol), benzylamine (2.51 mL, 23.05 mmol), X-phos (658 mg, 1.38 mmol), $\text{Pd}_2(\text{dba})_3$ (632 mg, 0.69 mmol) and $t\text{-BuOK}$ (774 mg, 6.92 mmol) in toluene (50 mL) was purged with nitrogen for 10 minutes and then heated at 90 °C for 18 h. The mixture was diluted with methylene chloride (200 mL) and filtered. The filtrate was concentrated and purified by chromatography (silica gel, 0–100% EtOAc/hexanes) to afford 79 (125 mg, 9%) as a brown gum: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.38 (s, 1H), 7.31–7.22 (m, 5H), 5.68 (s, 2H), 4.28 (t, J = 7.5 Hz, 1H), 4.01 (d, J = 7.0 Hz, 2H), 2.14 (s, 3H), 1.93 (s, 3H), 1.74 (s, 3H).

[0301] Step 3: To a solution of 79 (80 mg, 0.26 mmol) in triethylorthoacetate (2 mL) was added AcOH (0.2 mL). The mixture was heated to 120 °C for 2 h. The mixture was concentrated, the residue was dissolved in EtOAc (100 mL) and washed with saturated NaHCO_3 (50 mL $\times$ 2). The organic layer was dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 0–10% MeOH/ethyl acetate) to afford Example Compound 181 (39 mg,


45%) as an off-white solid: ^1H NMR (300 MHz, CDCl_3) δ 8.23 (s, 1H), 7.37–7.31 (m, 3H), 6.95–6.92 (m, 2H), 5.58 (s, 2H), 2.64 (s, 3H), 2.23 (s, 3H), 2.22 (s, 3H), 2.06 (s, 3H); ESI m/z 333 $[\text{M} + \text{H}]^+$.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-7-methyl-1H-imidazo[4,5-b]pyridin-2(3H)-one (Example Compound 180)



[0302] A mixture of **79** (31 mg, 0.10 mmol) and CDI (33 mg, 0.2 mmol) in dioxane (3 mL) was heated to 120 °C for 16 h. The mixture was concentrated, the residue was purified by chromatography (silica gel, 50–100% ethyl acetate/hexanes) to afford **Example Compound 180** (10 mg, 30%) as an off-white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.89 (s, 1H), 7.74 (s, 1H), 7.38–7.24 (m, 3H), 7.17–7.14 (m, 2H), 5.26 (s, 2H), 2.16 (s, 3H), 2.01 (s, 3H), 1.99 (s, 3H); ESI m/z 335 $[\text{M} + \text{H}]^+$.

Preparation of 3,5-dimethyl-4-(2-methyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole (Example Compound 108)




[0303] Step 1: To a suspension of **27** (660 mg, 3.23 mmol) in acetonitrile (33 mL) was added (1-bromoethyl)benzene (658 mg, 3.55 mmol) and potassium carbonate (893 mg, 6.46 mmol). The mixture was heated to 60 °C for 16 hours, then cooled, diluted with methylene chloride (120 mL) and washed with brine (40 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–10% methanol/methylene chloride) to afford **57** (256 mg, 26%) as white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.36 (d, J = 1.5 Hz, 2H), 7.30 (t, J = 7.5 Hz, 2H), 7.20–7.17 (m, 2H), 6.15 (d, J = 2.0 Hz, 1H), 5.82 (s, 2H), 5.40 (d, J = 5.5 Hz, 1H), 4.51–4.45 (m, 1H), 2.05 (s, 3H), 1.84 (s, 3H), 1.48 (d, J = 7.0 Hz, 3H).

[0304] Step 2: To a solution of **57** (41 mg, 0.13 mmol) in triethylorthoacetate (0.24 mL, 1.33 mmol) was added acetic acid (20 μL , 0.36 mmol). The mixture was heated to 100 °C for 1 h, then one drop of concentrated HCl was added. The mixture was heated to 100 °C for 10 min. The

mixture was basified with saturated sodium bicarbonate, diluted with methylene chloride (45 mL) and washed with brine (20 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–3% methanol/methylene chloride) followed by trituration with methylene chloride/hexanes to afford **Example Compound 108** (11 mg, 28%) as white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.27 (d, $J = 2.0$ Hz, 1H), 7.44 (d, $J = 2.0$ Hz, 1H), 7.40–7.36 (m, 4H), 7.33–7.30 (m, 1H), 6.01 (q, $J = 7.0$ Hz, 1H), 2.70 (s, 3H), 2.26 (s, 3H), 2.06 (s, 3H), 1.93 (d, $J = 7.0$ Hz, 3H); ESI m/z 333 $[\text{M} + \text{H}]^+$.

Preparation of 6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one (Example Compound 112) and 6-(3,5-dimethylisoxazol-4-yl)-*N*-ethyl-1-(1-phenylethyl)-1H-imidazo[4,5-*b*]pyridin-2-amine (Example Compound 113)




[0305] Step 1: To a suspension of **57** (250 mg, 0.81 mmol) in 1,4-dioxane (6 mL), was added 1,1'-carbonyldiimidazole (158 mg, 0.97 mmol). The mixture was purged with nitrogen for 5 min, and then heated to 100 °C for 16 h. The mixture was diluted with methylene chloride (100 mL), filtered and concentrated. The residue was purified by chromatography (silica gel, 0–5% methanol/methylene chloride) then triturated with methylene chloride/hexanes to afford **Example Compound 112** (258 mg, 95%) as off-white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 11.78 (s, 1H), 7.87 (d, $J = 2.0$ Hz, 1H), 7.44 (d, $J = 7.5$ Hz, 2H), 7.36 (t, $J = 7.5$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 1H), 7.09 (d, $J = 2.0$ Hz, 1H), 5.72 (q, $J = 7.0$ Hz, 1H), 2.26 (s, 3H), 2.06 (s, 3H), 1.84 (d, $J = 7.0$ Hz, 3H); ESI m/z 335 $[\text{M} + \text{H}]^+$.

[0306] Step 2: A mixture of **Example Compound 112** (100 mg, 0.30 mmol) and phosphorus (V) oxychloride (1 mL) was heated to 110 °C for 5 h, and cooled to room temperature. The reaction mixture was concentrated, diluted with methylene chloride (75 mL), and washed with saturated sodium bicarbonate solution (20 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was dissolved in a 2.0 M solution of ethylamine in tetrahydrofuran (6.0 mL, 12.0 mmol) and the mixture was heated to 75 °C for 7 h. The reaction mixture was concentrated, and the residue was purified by chromatography (silica gel, 0–5% methanol/methylene chloride), then triturated with ethyl acetate/hexanes to afford **Example Compound 113** (52 mg, 49%) as a white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.90 (d, $J = 2.0$ Hz,

1H), 7.40–7.28 (m, 6H), 6.81 (d, $J = 2.0$ Hz, 1H), 5.84 (q, $J = 7.0$ Hz, 1H), 3.54–3.48 (m, 2H), 2.20 (s, 3H), 1.99 (s, 3H), 1.83 (d, $J = 7.0$ Hz, 3H), 1.27 (t, $J = 7.0$ Hz, 3H); ESI m/z 362 $[M + H]^+$.

Preparation of 6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one (Enantiomer A) (Example Compound 218) and 6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one (Enantiomer B) (Example Compound 219)




[0307] Example Compound 112 (87 mg) was separated by SFC chiral HPLC (Chiralpak AS-H, 30 mm × 250 mm, mobile phase 30% EtOH in CO₂ (0.2% Et₂NH), 120 bar, flow rate 80 mL/min) to afford Example Compound 218 (Enantiomer A) (41 mg, 46%) and Example Compound 219 (Enantiomer B) (41 mg, 46%) as off-white solids.

[0308] Example Compound 218 (Enantiomer A): ¹H NMR (500 MHz, DMSO-*d*₆) δ 11.77 (s, 1H), 7.87 (d, $J = 2.0$ Hz, 1H), 7.44 (d, $J = 7.5$ Hz, 2H), 7.37 (t, $J = 7.5$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 1H), 7.09 (d, $J = 2.0$ Hz, 1H), 5.72 (q, $J = 7.5$ Hz, 1H), 2.26 (s, 3H), 2.06 (s, 3H), 1.84 (d, $J = 7.5$ Hz, 3H); ESI m/z 335 $[M + H]^+$; HPLC (Chiralcel OD, 4.6 mm × 250 mm, 10% EtOH in heptane, 1 mL/min) >99%, $t_R = 9.4$ min.

[0309] Example Compound 219 (Enantiomer B): ¹H NMR (500 MHz, DMSO-*d*₆) δ 11.78 (s, 1H), 7.87 (d, $J = 1.5$ Hz, 1H), 7.44 (d, $J = 7.5$ Hz, 2H), 7.36 (t, $J = 7.5$ Hz, 2H), 7.29 (t, $J = 7.5$ Hz, 1H), 7.08 (d, $J = 2.0$ Hz, 1H), 5.72 (q, $J = 7.5$ Hz, 1H), 2.26 (s, 3H), 2.06 (s, 3H), 1.84 (d, $J = 7.5$ Hz, 3H); ESI m/z 335 $[M + H]^+$; HPLC (Chiralcel OD, 4.6 mm × 250 mm, 10% EtOH in heptane, 1 mL/min) >99%, $t_R = 10.9$ min.

Preparation of 3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-1H-benzo[d]imidazol-2(3H)-one (Example Compound 122)




[0310] Step 1: To a solution of **20** (214 mg, 0.77 mmol) in 1,4-dioxane (5 mL) was added 1,1'-carbonyldiimidazole (150 mg, 0.93 mmol) and the mixture was heated to 100 °C for 15 h. The mixture was concentrated and purified by chromatography (silica gel, 0–20% ethyl acetate/hexanes) to afford **80** (142 mg, 61%) as a white solid; ¹H NMR (500 MHz, DMSO-*d*₆) δ 11.13 (s, 1H), 7.35–7.25 (m, 6H), 7.12 (dd, *J* = 8.5, 2.0 Hz, 1H), 6.94 (d, *J* = 8.0 Hz, 1H), 5.01 (s, 2H).

[0311] Step 2: To a solution of **80** (100 mg, 0.33 mmol) in 1,4-dioxane (5 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (110 mg, 0.49 mmol), potassium carbonate (91 mg, 0.66 mmol), and water (1 mL). The mixture was purged with nitrogen for 10 min, tetrakis(triphenylphosphine)palladium(0) (19 mg, 0.016 mmol) was added, and the mixture was heated to 90 °C for 16 h. The mixture was diluted with methylene chloride (100 mL), and washed with brine (30 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–5% methanol/methylene chloride) then triturated with ethyl acetate/hexanes to afford **81** (55 mg, 52%) as a white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.07 (s, 1H), 7.40–7.23 (m, 5H), 7.06 (d, *J* = 8.1 Hz, 1H), 7.02 (s, 1H), 6.95 (dd, *J* = 7.8, 1.5 Hz, 1H), 5.03 (s, 2H), 2.30 (s, 3H), 2.13 (s, 3H); ESI *m/z* 320 [*M* + *H*]⁺.

[0312] Step 3: To a solution of **81** (36 mg, 0.11 mmol) in acetonitrile (3 mL) was added potassium carbonate (109 mg, 0.79 mmol) and iodoethane (80 mg, 0.56 mmol), then the mixture was heated to 40 °C for 48 h. The mixture was diluted with methylene chloride (75 mL), and washed with brine (20 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–20% ethyl acetate/methylene chloride), then triturated with ethyl acetate/hexanes to afford **Example Compound 122** (14 mg, 37%) as a yellow-white solid: ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.37 (d, *J* = 7.5 Hz, 2H), 7.33 (t, *J* = 7.0 Hz, 2H), 7.29 (d, *J* = 8.0 Hz, 1H), 7.26 (t, *J* = 7.0 Hz, 1H), 7.09 (d, *J* = 1.5 Hz, 1H), 7.03 (dd, *J* = 8.0, 1.5 Hz, 1H), 5.08 (s, 2H), 3.94 (q, *J* = 7.0 Hz, 2H), 2.31 (s, 3H), 2.13 (s, 3H), 1.26 (t, *J* = 7.0 Hz, 3H); ESI *m/z* 348 [*M* + *H*]⁺.

Preparation of 1-benzyl-*N*⁶-(3,5-dimethylisoxazol-4-yl)-2-methyl-1*H*-benzo[*d*]imidazole-4,6-diamine (Example Compound 142)



[0313] Step 1: To a suspension of **33** (790 mg, 3.09 mmol) in acetonitrile (15 mL) was added benzyl chloride (703 mg, 5.55 mmol) and potassium carbonate (1.07 g, 7.71 mmol). The reaction mixture was heated to 60 °C for 16 h, then concentrated, and the residue was purified by chromatography (silica gel, 0–30% ethyl acetate/hexanes) to afford **82** (813 mg, 76%) as a yellow


solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.33 (d, $J = 1.8$ Hz, 1H), 8.12 (d, $J = 1.8$ Hz, 1H), 7.39–7.27 (m, 3H), 7.13 (d, $J = 6.6$ Hz, 2H), 5.62 (s, 2H), 2.60 (s, 3H).

[0314] **Step 2:** To a solution of **82** (150 mg, 0.43 mmol) in toluene (5 mL) was added **83** (73 mg, 0.65 mmol), cesium carbonate (282 mg, 0.87 mmol) and XPhos (41 mg, 0.087 mmol). The solution was purged with nitrogen for 5 min, then tris(dibenzylideneacetone)dipalladium(0) (40 mg, 0.043 mmol) was added and heated to 110 °C for 16 h. The mixture was filtered through celite and concentrated, the residue was purified by chromatography (silica gel, 0–7% methanol/methylene chloride) to afford **84** (80 mg, 49%) as a brown oil: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.59 (s, 1H), 7.34–7.28 (m, 4H), 7.06 (d, $J = 7.0$ Hz, 2H), 6.76 (d, $J = 2.5$ Hz, 1H), 5.44 (s, 2H), 2.54 (s, 3H), 2.13 (s, 3H), 1.91 (s, 3H).

[0315] **Step 3:** To a solution of **84** (78 mg, 0.21 mmol) in tetrahydrofuran (5 mL) was added a solution of sodium dithionite (215 mg, 1.24 mmol) in water (4 mL). The mixture was stirred at room temperature for 2 h, the 2N HCl (1 mL) was added, the mixture was heated to reflux for 15 min. The mixture was basified by sodium carbonate, and extracted with methylene chloride (50 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–10% methanol/methylene chloride) to afford **Example Compound 142** (38 mg, 53%) as a red-brown solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.31 (t, $J = 7.5$ Hz, 2H), 7.25 (t, $J = 7.5$ Hz, 1H), 7.04 (d, $J = 7.5$ Hz, 2H), 6.69 (s, 1H), 5.73 (d, $J = 2.0$ Hz, 1H), 5.60 (d, $J = 2.0$ Hz, 1H), 5.18 (s, 2H), 5.05 (s, 2H), 2.38 (s, 3H), 2.13 (s, 3H), 1.92 (s, 3H); ESI m/z 348 $[\text{M} + \text{H}]^+$.

General Procedure U:


Preparation of 1-benzyl-2-methyl-6-(5-methylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine (Example Compound 201)



[0316] To a solution of **82** (100 mg, 0.29 mmol) in 1,4-dioxane (5 mL) was added 5-methylisoxazole-4-boronic acid pinacol ester (91 mg, 0.43 mmol), sodium carbonate (80 mg, 0.58 mmol), water (1 mL), and tetrakis(triphenylphosphine)palladium(0) (17 mg, 0.01 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 5 h. The mixture was diluted with methylene chloride (70 mL), washed with brine (25 mL), dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–5% ethyl

acetate/methylene chloride) to a yellow solid which was dissolved in THF (4 mL), a solution of sodium dithionite (159 mg, 0.91 mmol) in water (2 mL) was added and the mixture was stirred at room temperature for 2 h. 2 N HCl (1 mL) was added to the mixture, and the mixture was heated to reflux for 15 min. The mixture was basified by saturated aqueous sodium bicarbonate solution, and extracted with methylene chloride (40 mL $\times$ 2). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–8% methanol/methylene chloride) and triturated with ethyl acetate/hexanes to afford **Example Compound 201** (12 mg, 25%) as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 8.69 (d, J = 0.6 Hz, 1H), 7.36–7.26 (m, 3H), 7.15 (d, J = 6.9 Hz, 2H), 6.78 (d, J = 1.5 Hz, 1H), 6.47 (d, J = 1.5 Hz, 1H), 5.40 (s, 2H), 5.33 (s, 2H), 2.50 (s, 3H), 2.47 (s, 3H); ESI m/z 319 $[\text{M} + \text{H}]^+$.

Preparation of *N*-(1-benzyl-2-methyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazol-4-amine (Example Compound 155)




[0317] Step 1: To a suspension of 2,3-diamino-5-bromopyridine **26** (1.5 g, 7.98 mmol) in methylene chloride (80 mL) was added benzaldehyde (931 mg, 8.78 mmol) and acetic acid (40 drops). The mixture was stirred at room temperature for 16 h, then washed with saturated sodium bicarbonate solution (40 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was dissolved in methanol (50 mL) and sodium borohydride (815 mg, 21.5 mmol) was slowly added. The mixture was stirred at room temperature for 1 h. The mixture was diluted with methylene chloride (100 mL), washed with saturated sodium bicarbonate (40 mL), dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–10% methanol/methylene chloride) to afford **85** (1.12 g, 51%) as an off-white solid: ^1H NMR (500 MHz, DMSO- d_6) δ 7.35–7.34 (m, 4H), 7.28–7.23 (m, 2H), 6.54 (d, J = 2.0 Hz, 1H), 5.78 (s, 2H), 5.73 (t, J = 5.5 Hz, 1H), 4.30 (d, J = 5.5 Hz, 2H).

[0318] Step 2: To a suspension of **85** (970 mg, 3.49 mmol) in triethylorthoacetate (5.66 g, 37.9 mmol) was added acetic acid (539 μL , 9.42 mmol). The mixture was heated to 100 °C for 40 min. The reaction mixture was basified with saturated sodium bicarbonate (8 mL), diluted with methylene chloride (50 mL), and washed with saturated sodium bicarbonate (30 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated. The residue was purified by chromatography (silica gel, 0–8% methanol/methylene chloride) to afford **86** (305 mg, 30%) as a

light brown solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.41 (d, $J = 2.0$ Hz, 1H), 8.29 (d, $J = 2.0$ Hz, 1H), 7.35 (t, $J = 7.0$ Hz, 2H), 7.30 (t, $J = 7.0$ Hz, 1H), 7.15 (d, $J = 7.0$ Hz, 2H), 5.52 (s, 2H), 2.55 (s, 3H).


[0319] Step 3: To a solution of **86** (80 mg, 0.26 mmol) in toluene (5 mL), was added **83** (44 mg, 0.40 mmol), cesium carbonate (173 mg, 0.53 mmol), and XPhos (25 mg, 0.053 mmol). The solution was purged with nitrogen for 5 min, then tris(dibenzylideneacetone)dipalladium(0) (24 mg, 0.026 mmol) was added. The mixture was heated to 110 °C for 16 h. The reaction mixture was diluted with methylene chloride (20 mL), filtered through celite, and concentrated. The residue was purified by chromatography (silica gel, 0–10% methanol/methylene chloride) then triturated with methylene chloride/hexanes to afford **Example Compound 155** (40 mg, 45%) as a light-brown solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.88 (d, $J = 2.5$ Hz, 1H), 7.34–7.30 (m, 3H), 7.27 (t, $J = 7.0$ Hz, 1H), 7.05 (d, $J = 7.0$ Hz, 2H), 6.71 (d, $J = 2.5$ Hz, 1H), 5.38 (s, 2H), 2.47 (s, 3H), 2.14 (s, 3H), 1.92 (s, 3H); ESI m/z 334 $[\text{M} + \text{H}]^+$.

Preparation of 1-benzyl-2-methyl-6-(1-methyl-1H-1,2,3-triazol-5-yl)-1H-imidazo[4,5-b]pyridine (Example Compound 206)




[0320] To a solution of **86** (100 mg, 0.33 mmol) in 1,4-dioxane (5 mL) was added 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-1,2,3-triazole (138 mg, 0.66 mmol), K_2CO_3 (137 mg, 0.99 mmol), water (1 mL), and tetrakis(triphenylphosphine)palladium(0) (19 mg, 0.02 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 16 h. The mixture was diluted with ethyl acetate (70 mL), washed with brine (25 mL), dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–8% methanol/methylene chloride) followed by trituration with methylene chloride/hexanes to afford **Example Compound 206** (14 mg, 14%) as a white solid; ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.54 (d, $J = 2.5$ Hz, 1H), 8.27 (d, $J = 2.0$ Hz, 1H), 7.96 (s, 1H), 7.35 (t, $J = 7.0$ Hz, 2H), 7.29 (t, $J = 7.0$ Hz, 1H), 7.21 (d, $J = 7.0$ Hz, 2H), 5.58 (s, 2H), 4.07 (s, 3H), 2.60 (s, 3H); ESI m/z 305 $[\text{M} + \text{H}]^+$.

Preparation of 1-benzyl-2-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridine (Example Compound 154)




[0321] 1-Benzyl-2-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridine (**Example Compound 154**) was prepared by following the similar method for the preparation of **Example Compound 206** as an off-white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.48 (d, $J = 2.0$ Hz, 1H), 8.14 (d, $J = 2.0$ Hz, 1H), 7.50 (d, $J = 2.0$ Hz, 1H), 7.35 (t, $J = 7.0$ Hz, 2H), 7.29 (t, $J = 7.0$ Hz, 1H), 7.21 (d, $J = 7.0$ Hz, 2H), 6.46 (d, $J = 2.0$ Hz, 1H), 5.57 (s, 2H), 3.83 (s, 3H), 2.60 (s, 3H); ESI m/z 304 $[\text{M} + \text{H}]^+$.

Preparation of 4-(1-benzyl-2-cyclopropyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 138**)**



[0322] To a solution of diamine **28** (100 mg, 0.340 mmol) in 1,4-dioxane (2 mL) was added cyclopropanecarboxaldehyde (29 mg, 0.408 mmol) and acetic acid (0.67 mL). The mixture was heated at 110 °C for 24 h. The mixture was then diluted with methylene chloride and washed with saturated sodium bicarbonate. The organic layer was then dried with sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–5% methanol/methylene chloride) to afford **Example Compound 138** (68 mg, 58%) as an off-white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.29 (d, $J = 2.1$ Hz, 1H), 7.95 (d, $J = 2.0$ Hz, 1H), 7.37–7.33 (m, 2H), 7.30–7.28 (m, 3H), 5.67 (s, 2H), 2.38 (s, 3H), 2.37–2.35 (m, 1H), 2.20 (s, 3H), 1.13–1.11 (m, 4H); ESI m/z 345 $[\text{M} + \text{H}]^+$. HPLC >99%.

Preparation of 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1*H*-benzo[d]imidazol-2(3*H*)-one (Example Compound 145), 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-*N*-ethyl-4-nitro-1*H*-benzo[d]imidazol-2-amine (Example Compound 159), 4-Amino-1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazol-2(3*H*)-one (Example Compound 161) and 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-*N*-ethyl-1*H*-benzo[d]imidazole-2,4-diamine (Example Compound 160)



[0323] Step 1: To a mixture of **32** (1.50 g, 6.46 mmol) and **3** (2.16 g, 9.70 mmol) in 1,4-dioxane (40 mL) and water (4 mL) was added potassium carbonate (1.79 g, 12.9 mmol) and tetrakis(triphenylphosphine)palladium(0) (373 mg, 0.32 mmol). The reaction was stirred and heated at 90°C for 17 h. The reaction mixture was diluted with methanol (20 mL) and silica gel (20 g) was added. The suspension was concentrated to dryness and the resulting powder was loaded onto silica gel and eluted with 0–50% ethyl acetate in hexanes. The clean product was concentrated to give **87** (585 mg, 36%) as a brown solid: ^1H NMR (500 MHz, CDCl_3) δ 7.62 (d, $J = 2.0$ Hz, 1H), 6.81 (d, $J = 2.0$ Hz, 1H), 6.01 (br s, 2H), 3.52 (br s, 2H), 2.39 (s, 3H), 2.25 (s, 3H).

[0324] Step 2: To a solution of **87** (250 mg, 1.01 mmol), a catalytic amount of DMAP and 1,4-dioxane (4 mL) in a pressure tube was added 1,1'-carbonyldiimidazole (327 mg, 2.01 mmol). The tube was sealed and heated to 80°C for 17 h. The reaction mixture was diluted with methanol (20 mL) and silica gel (10 g) was added. The suspension was concentrated to dryness and the resulting powder was loaded onto silica gel (40 g) and eluted with 0–70% ethyl acetate in hexanes. The clean

product was concentrated to give **88** (167 mg, 60%) as an orange solid: ^1H NMR (500 MHz, CDCl_3) δ 7.74 (d, $J = 1.5$ Hz, 1H), 7.17 (d, $J = 1.5$ Hz, 1H), 2.43 (s, 3H), 2.28 (s, 3H).

[0325] Step 3: To a solution of **88** (309 mg, 1.13 mmol), potassium carbonate (312 mg, 2.25 mmol), acetonitrile (5 mL) and DMF (2 mL) in a pressure tube was added (bromomethyl)cyclopropane (183 mg, 1.35 mmol) and the reaction was sealed and heated at 80 °C for 17 h. The material was cooled to room temperature and poured into a saturated aq. NaCl solution (30 mL). Ethyl acetate (100 mL) was added and the layers were separated. The ethyl acetate layer was washed with saturated aq. NaCl solution (2×100 mL), dried over Na_2SO_4 , filtered and the filtrate was concentrated. The resulting oil in CH_2Cl_2 (10 mL) was loaded onto silica gel (80 g) and eluted with 0–40% ethyl acetate in hexanes. The clean product was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example Compound 145** (88 mg, 35%) as a yellow solid: ^1H NMR (500 MHz, CD_3OD) δ 7.82 (d, $J = 1.5$ Hz, 1H), 7.52 (d, $J = 1.0$ Hz, 1H), 3.87 (d, $J = 7.0$ Hz, 2H), 2.45 (s, 3H), 2.29 (s, 3H), 1.30–1.18 (m, 1H), 0.60–0.52 (m, 2H), 0.47–0.43 (m, 2H). ESI m/z 329 $[\text{M} + \text{H}]^+$. HPLC >99%.


[0326] Step 4: A solution of **Example Compound 145** (171 mg, 0.521 mmol) in phosphorus(V) oxychloride (4 mL) was placed in a sealed tube and heated at 110 °C for 8 h. The solvent was removed *in vacuo* and a saturated aq. NaHCO_3 solution (5 mL) was added. The mixture was extracted with ethyl acetate (2×20 mL) and the combined extracts were dried over Na_2SO_4 , filtered and the filtrate was concentrated. THF (5 mL) and 2.0M ethylamine solution in THF were then added and the reaction was heated at 70 °C for 12 h. The reaction was concentrated to dryness and the residue diluted with CH_2Cl_2 (5 mL). The resulting solution was loaded onto silica gel (40 g) and eluted with 0–80% ethyl acetate in hexanes. The clean product was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example Compound 159** (105 mg, 57%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 7.78 (d, $J = 1.5$ Hz, 1H), 7.44 (d, $J = 1.5$ Hz, 1H), 4.03 (d, $J = 6.5$ Hz, 2H), 3.67 (q, $J = 7.0$ Hz, 2H), 2.44 (s, 3H), 2.29 (s, 3H), 1.33 (t, $J = 7.0$ Hz, 3H), 1.30–1.18 (m, 1H), 0.60–0.52 (m, 2H), 0.47–0.41 (m, 2H). ESI m/z 356 $[\text{M} + \text{H}]^+$. HPLC >99%.

[0327] Step 5: A solution of **Example Compound 145** (59 mg, 0.215 mmol) in THF (10 mL) was added a solution of sodium dithionite (225 mg, 1.29 mmol) in water (10 mL) dropwise over 5 min. The solution was stirred at room temperature for 16 h and the solvents were removed *in vacuo*. Methanol (20 mL) was added and the suspension stirred at room temperature for 3 h. The mixture was filtered and the filtrate was concentrated to dryness. A solution of 2N aq. HCl (10 mL) was added to the residue and heated to reflux for 5 min. After concentration to dryness, methanol

was added (10 mL) and the solution was adjusted to pH 8 using saturated aq. NaHCO_3 solution (15 mL). Silica gel was added (10 g) and the suspension was concentrated to dryness. The resulting powder was loaded onto silica gel and eluted with 0–4% methanol in methylene chloride. The clean product was then purified by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example Compound 161** (32 mg, 50%) as a white solid: ^1H NMR (500 MHz, CD_3OD) δ 6.49 (d, $J = 1.5$ Hz, 1H), 6.42 (d, $J = 1.5$ Hz, 1H), 3.75 (d, $J = 6.5$ Hz, 2H), 2.39 (s, 3H), 2.24 (s, 3H), 1.28–1.18 (m, 1H), 0.56–0.48 (m, 2H), 0.44–0.39 (m, 2H). ESI m/z 299 $[\text{M} + \text{H}]^+$. HPLC 97.4%.

[0328] Step 6: A solution of **Example Compound 159** (90 mg, 0.253 mmol) in THF (10 mL) was added a solution of sodium dithionite (265 mg, 1.52 mmol) in water (10 mL) dropwise over 5 min. The solution was stirred at room temperature for 16 h and the solvents were removed *in vacuo*. Methanol (20 mL) was added and the suspension stirred at room temperature for 3 h. The mixture was filtered and the filtrate was concentrated to dryness. A solution of 2N aq. HCl (10 mL) was added to the residue and heated to reflux for 5 min. After concentration to dryness, methanol was added (10 mL) and the solution was adjusted to pH 8 using saturated aq. NaHCO_3 solution (15 mL). Silica gel was added (10 g) and the suspension was concentrated to dryness. The resulting powder was loaded onto silica gel and eluted with 0–4% methanol in methylene chloride. The clean product was then purified by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example Compound 160** (61 mg, 74%) as a white solid: ^1H NMR (500 MHz, CD_3OD) δ 6.49 (d, $J = 1.5$ Hz, 1H), 6.37 (d, $J = 1.5$ Hz, 1H), 3.88 (d, $J = 6.5$ Hz, 2H), 3.48 (q, $J = 7.0$ Hz, 2H), 2.39 (s, 3H), 2.24 (s, 3H), 1.28–1.18 (m, 1H), 0.53–0.48 (m, 2H), 0.40–0.35 (m, 2H). ESI m/z 326 $[\text{M} + \text{H}]^+$. HPLC >99%.


Preparation of 4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(4-hydroxybenzyl)-1H-benzo[d]imidazol-2(3H)-one (Example Compound 129).



[0329] To a solution of **Example Compound 104** (54 mg, 0.15 mmol) in dichloromethane (5 mL) under nitrogen atmosphere was added boron tribromide (0.45 mL, 1M in dichloromethane, 0.45 mmol). The reaction mixture was stirred at room temperature overnight, treated with methanol, and concentrated in vacuum. The residue was dissolved in methanol, basified with

ammonium hydroxide, concentrated in vacuum, and purified by chromatography (silica gel, 0–20% methanol in ethyl acetate). It was further purified by reverse phase HPLC on a Polaris C₁₈ column eluting with 10–90% CH₃CN in H₂O to give **Example Compound 129** (31 mg, 59%) as an off-white solid: ¹H NMR (500 MHz, CD₃OD) δ 7.17 (d, *J* = 8.6 Hz, 2H), 6.72 (d, *J* = 8.6 Hz, 2H), 6.39 (d, *J* = 1.3 Hz, 1H), 6.26 (d, *J* = 1.3 Hz, 1H), 4.94 (s, 2H), 2.28 (s, 3H), 2.12 (s, 3H); HPLC >99%, *t_R* = 11.0 min; ESI *m/z* 351 [M + H]⁺.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1*H*-benzo[*d*]imidazol-4-ol (Example Compound 173)



[0330] Step 1: To a solution of **89** (5.00 g, 32.5 mmol) and triethylamine (9.04 mL, 65.0 mmol) in *N,N*-dimethylformamide (150 mL) was added *tert*-butylchlorodimethylsilane (5.86 g, 39.0 mmol) at room temperature. The reaction mixture was stirred at room temperature for 1h and diluted with ethyl acetate. The mixture was washed with water, brine, dried over sodium sulfate, and filtered. The filtrate was concentrated to afford **90** (8.59 g, 98%) as a brown oil: ¹H NMR (300 MHz, CDCl₃) δ 7.75 (dd, *J* = 1.3, 8.9 Hz, 1H), 6.89 (dd, *J* = 1.2, 7.6 Hz, 1H), 6.53 (dd, *J* = 8.8, 7.6 Hz, 1H), 6.45–6.15 (bs, 2H), 1.03 (s, 9H), 0.28 (s, 6H).

[0331] Step 2: To a solution of **90** (8.59 g, 32.1 mmol) in acetic acid (120 mL) was added *N*-bromosuccinimide (6.28 g, 35.3 mmol) at room temperature. The reaction mixture was stirred at room temperature for 30 min and then concentrated. The residue was dissolved in methanol and basified with 5% aqueous sodium bicarbonate. The precipitate formed was filtered, washed with water, and dried under vacuum to afford **91** (8.56 g, 76%) as an orange solid: ¹H NMR (500 MHz, CDCl₃) δ 7.91 (d, *J* = 2.1 Hz, 1H), 6.96 (d, *J* = 2.1 Hz, 1H), 6.50–6.12 (bs, 2H), 1.03 (s, 9H), 0.30 (s, 6H).

[0332] Step 3: To a solution of **91** (5.00 g, 14.4 mmol) in tetrahydrofuran (60 mL) was added platinum on carbon (1.00 g, 5% Pt on carbon). The reaction mixture was stirred under hydrogen atmosphere at room temperature overnight. The mixture was filtered, washed with MeOH, and the filtrate was concentrated to afford **92** (5.65 g, >99%) as a dark brown oil: ^1H NMR (500 MHz, CDCl_3) δ 6.51 (d, $J = 2.0$ Hz, 1H), 6.46 (d, $J = 2.0$ Hz, 1H), 3.50–2.50 (bs, 4H), 1.01 (s, 9H), 0.24 (s, 6H); ESI m/z 317 $[\text{M} + \text{H}]^+$.

[0333] Step 4: To a solution of **92** (2.00 g, 6.31 mmol) in ethanol (50 mL) was added triethylorthoacetate (3.07 g, 18.9 mmol) and sulfamic acid (1 mg, 0.01 mmol). The reaction was heated in a sealed tube at 80 °C overnight. The mixture was concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to afford **93** (2.07 g, 96%) as a light red solid: ^1H NMR (500 MHz, CDCl_3) δ 8.75 (s, 1H), 7.45 (s, 1H), 6.78 (s, 1H), 3.61 (s, 3H), 1.03 (s, 9H), 0.28 (s, 6H); ESI m/z 341 $[\text{M} + \text{H}]^+$.


[0334] Step 5: A mixture of **93** (200 mg, 0.587 mmol), benzyl bromide (150 mg, 0.880 mmol), and potassium bicarbonate (113 mg, 0.822 mmol) in acetonitrile (20 mL) was heated at 45 °C for 2 days. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–30% ethyl acetate in hexanes) to afford **94** (303 mg, 30%) as a brown solid: ^1H NMR (500 MHz, CDCl_3) δ 7.36–7.26 (m, 3H), 7.01 (d, $J = 8.2$ Hz, 2H), 6.97 (d, $J = 1.6$ Hz, 1H), 6.81 (d, $J = 1.6$ Hz, 1H), 5.22 (s, 2H), 2.50 (s, 3H), 1.05 (s, 9H), 0.30 (s, 6H); ESI m/z 431 $[\text{M} + \text{H}]^+$.

[0335] Step 6: To a solution of **94** (75 mg, 0.17 mmol) in 1,4-dioxane (10 mL) and water (1 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (58 mg, 0.26 mmol), potassium bicarbonate (70 mg, 0.70 mmol), and tetrakis(triphenylphosphine)palladium(0) (10 mg, 0.0087 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 2h. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to give **95** (53 mg, 70%) as an off-white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.33 (t, $J = 6.3$ Hz, 2H), 7.27 (t, $J = 5.1$ Hz, 1H), 7.14 (d, $J = 7.1$ Hz, 2H), 6.89 (d, $J = 1.3$ Hz, 1H), 6.58 (d, $J = 1.3$ Hz, 1H), 5.45 (s, 2H), 2.59 (s, 3H), 2.32 (s, 3H), 2.16 (s, 3H), 1.05 (s, 9H), 0.30 (s, 6H); HPLC >99%, $t_R = 16.4$ min; ESI m/z 448 $[\text{M} + \text{H}]^+$.

[0336] Step 7: A mixture of **95** (48 mg, 0.11 mmol) and potassium carbonate (30 mg, 0.22 mmol) in acetonitrile (10 mL) was heated in a sealed tube at 80 °C overnight. The reaction mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–20% methanol in ethyl acetate). It was further purified by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O to give **Example Compound 173** (32 mg, 87%) as an off-white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 9.84 (s, 1H), 7.33 (t, $J = 7.6$ Hz, 2H), 7.26 (t, $J = 7.3$ Hz, 1H), 7.18


(d, $J = 7.1$ Hz, 2H), 6.86 (d, $J = 1.3$ Hz, 1H), 6.47 (d, $J = 1.3$ Hz, 1H), 5.42 (s, 2H), 2.52 (s, 3H), 2.33 (s, 3H), 2.15 (s, 3H); ESI m/z 334 $[M + H]^+$.

Preparation of 4-Amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2(3H)-thione (Example Compound 177).



[0337] A mixture of **Example Compound 16** (34 mg, 0.10 mmol) and Lawesson's reagent (202 mg, 0.5 mmol) was heated to 180 °C in microwave reactor for 2 h. The mixture was concentrated, the residue was purified by chromatography (silica gel, 0–40% EtOAc/hexanes) followed by chromatography (C_{18} , 10–70% CH_3CN /water) to give **Example Compound 177** (13 mg, 37%) as an off-white solid: ^1H NMR (300 MHz, $DMSO-d_6$) δ 12.56 (s, 1H), 7.45–7.42 (m, 2H), 7.34–7.25 (m, 3H), 6.44 (d, $J = 1.2$ Hz, 1H), 6.39 (d, $J = 1.5$ Hz, 1H), 5.44 (s, 4H), 2.29 (s, 3H), 2.11 (s, 3H); ESI m/z 351 $[M + H]^+$. HPLC 98.6%

Preparation of 1-benzyl-3-methyl-6-(1-methyl-1H-pyrazol-5-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one (Example Compound 198) and 4-amino-1-benzyl-3-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-benzo[d]imidazol-2(3H)-one (Example Compound 199).




[0338] Compound 96 was prepared by following the similar method for the preparation of **Example Compound 15** using 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole.

[0339] Step 1: A mixture of **96** (70 mg, 0.20 mmol), CH_3I (85 mg, 0.60 mmol) and K_2CO_3 (110 mg, 0.8 mmol) in DMF (3 mL) was stirred at rt for 16 h. The reaction mixture was diluted with EtOAc (100 mL) and washed with brine (3 $\times$ 50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 20–70% ethyl acetate/hexanes) to afford **Example Compound 198** (50 mg, 68%) as a yellow solid: ^1H NMR (300

MHz, CDCl₃) δ 7.66 (d, J = 1.5 Hz, 1H), 7.50 (d, J = 1.8 Hz, 1H), 7.36–7.30 (m, 5H), 7.02 (d, J = 1.5 Hz, 1H), 6.27 (d, J = 1.2 Hz, 1H), 5.16 (s, 2H), 3.69 (s, 3H), 3.65 (s, 3H); ESI m/z 364 [M + H]⁺.


[0340] Step 2: To a solution of **Example Compound 198** (45 mg, 0.12 mmol) in THF (5 mL) and water (4 mL) was added Na₂S₂O₄ (129 mg, 0.74 mmol). The mixture was stirred at rt for 4 h, 2N HCl (1 mL) was added, the mixture was heated to reflux for 15 minutes then cooled to rt. Na₂CO₃ was added slowly to adjust to pH 9. The mixture was extracted with CH₂Cl₂ (100 mL), the organic layer was washed with brine (50 mL), filtered, concentrated and purified by chromatography (silica gel, 0–10% methanol/ethyl acetate) to afford **Example Compound 199** (37 mg, 90%) as a white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.39 (d, J = 1.8 Hz, 1H), 7.35–7.24 (m, 5H), 6.56 (d, J = 1.5 Hz, 1H), 6.54 (d, J = 1.5 Hz, 1H), 6.20 (d, J = 1.8 Hz, 1H), 5.15 (s, 2H), 5.01 (s, 2H), 3.72 (s, 3H), 3.63 (s, 3H); ESI m/z 334 [M + H]⁺.

Preparation of 4-(1-benzyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 220)




[0341] To a solution of **28** (100 mg, 0.34 mmol) and tetrahydro-2H-pyran-4-carboxylic acid (65 mg, 0.51 mmol) in CH₂Cl₂ was added EDC (131 mg, 0.68 mmol), *i*-Pr₂NEt (132 mg, 1.02 mmol) and DMAP (10 mg). The reaction mixture was stirred at rt for 16 h. The mixture was diluted with EtOAc (100 mL), washed with brine (50 mL) and saturated NaHCO₃ (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was dissolved in AcOH (2 mL) and heated to reflux for 5 h. The mixture was concentrated, the residue was dissolved in EtOAc (100 mL), washed with saturated NaHCO₃ (2 × 50 mL) and brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–10% MeOH/EtOAc) to give **Example Compound 220** (47 mg, 36%) as a light brown solid: ¹H NMR (300 MHz, CDCl₃) δ 8.41 (d, J = 1.8 Hz, 1H), 7.38–7.32 (m, 3H), 7.24 (d, J = 2.1 Hz, 1H), 7.08–7.05 (m, 2H), 5.42 (s, 2H), 4.12 (dd, J = 11.7, 1.8 Hz, 2H), 3.52 (td, J = 11.7, 1.8 Hz, 2H), 3.20–3.12 (m, 1H), 2.36–2.23 (m, 5H), 2.14 (s, 3H), 1.83–1.78 (m, 2H); ESI m/z 389 [M + H]⁺.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-*b*]pyridine-2-carboxamide (Example Compound 221)



[0342] A mixture of **28** (300 mg, 1.02 mmol) and methyl 2,2,2-trimethoxyacetate (1.5 mL) was heated to 120 °C for 16 h. The mixture was purified by chromatography (silica gel, 20–80% EtOAc/hexanes) to give a brown solid. The solid was dissolved in CH₃NH₂/THF (2 M) (3 mL) and heated 80 °C for 16 h. The mixture was concentrated, the residue was purified by chromatography (C₁₈, 10–70% CH₃CN/water) to give **Example Compound 221** (45 mg, 12%) as an off-white solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.31 (q, *J* = 4.5 Hz, 1H), 8.27 (d, *J* = 1.8 Hz, 1H), 7.54 (d, *J* = 1.8 Hz, 1H), 7.36–7.24 (m, 5H), 5.54 (s, 2H), 3.00 (d, *J* = 4.8 Hz, 3H), 2.21 (s, 3H), 2.00 (s, 3H); ESI *m/z* 362 [M + H]⁺.

Preparation of 1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one (Example Compound 171)




[0343] Step 1: To a solution of **85** (1.14 g, 4.09 mmol) in 1,4-dioxane (41 mL) was added 1,1'-carbonyldiimidazole (796 mg, 4.91 mmol). The reaction mixture was heated at 110 °C for 16 h. The reaction mixture was concentrated. Purification by chromatography (silica gel, 0–100% ethyl acetate/hexanes) afforded **97** (1.03 g, 83%) as a white solid: ¹H NMR (500 MHz, DMSO-*d*₆) δ 11.89 (s, 1H), 8.00 (d, *J* = 2.0 Hz, 1H), 7.68 (d, *J* = 2.0 Hz, 1H), 7.37–7.32 (m, 4H), 7.30–7.26 (m, 1H), 5.02 (s, 2H).

[0344] Step 2: To a solution of **97** (334 mg, 1.09 mmol) in 1,4-dioxane (11 mL) was added 1-methyl-1H-pyrazole-5-boronic acid pinacol ester (457 mg, 2.20 mmol), sodium carbonate (1.0 M in H₂O, 3.29 mL, 3.29 mmol) and tetrakis(triphenylphosphine)palladium(0) (127 mg, 0.1 mmol). The reaction mixture was purged with nitrogen and heated at 90 °C for 32 h. The mixture was diluted with methylene chloride (80 mL), washed with brine (40 mL), dried over sodium sulfate, filtered and concentrated. The residue was purified by chromatography (silica gel, 0–5% methanol/methylene chloride) followed by trituration with EtOAc to afford **Example Compound 171** (173 mg, 52%) as a


white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 11.87 (s, 1H), 8.04 (d, $J = 1.5$ Hz, 1H), 7.57 (d, $J = 1.5$ Hz, 1H), 7.46 (d, $J = 2.0$ Hz, 1H), 7.38 (d, $J = 7.5$ Hz, 2H), 7.34 (t, $J = 7.5$ Hz, 2H), 7.27 (t, $J = 7.0$ Hz, 1H), 6.37 (d, $J = 1.5$ Hz, 1H), 5.06 (s, 2H), 3.77 (s, 3H); ESI m/z 306 $[\text{M} + \text{H}]^+$.

Preparation of *N*-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-pyrrolo[3,2-*b*]pyridin-3-yl)acetamide (Example Compound 99)



[0345] A solution of **Example Compound 39** (100 mg, 0.29 mmol) in EtOH (3 mL) and AcOH (1 mL) was added iron powder (162 mg, 2.9 mmol). The reaction mixture was heated at 80 °C for 1 h. It was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–5% methanol/dichloromethane) afforded **Example Compound 99** (28 mg, 27%) as a red solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.2 (s, 1H), 8.32 (d, $J = 1.8$ Hz, 1H), 8.23 (s, 1H), 7.97 (d, $J = 1.8$ Hz, 1H), 7.32–7.25 (m, 5H), 5.45 (s, 2H), 2.40 (s, 3H), 2.22 (s, 3H), 2.12 (s, 3H); ESI MS m/z 361 $[\text{M} + \text{H}]^+$.


Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-pyrrolo[3,2-*b*]pyridin-3-amine (Example Compound 100)



[0346] To a solution of **Example Compound 39** (100 mg, 0.29 mmol) in EtOH (3 mL) and H_2SO_4 (0.5 mL) was added iron powder (162 mg, 2.9 mmol). The reaction mixture was heated at 80 °C for 1 h. It was diluted with EtOH (20 mL), adjusted to pH 7 by 6 N aq. NaOH. The mixture was filtered through a layer of *Celite* and the filtrate was concentrated. Purification by chromatography (silica gel, 0–5% methanol/dichloromethane) afforded **Example Compound 100** (12 mg, 13%) as a

red solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.18 (d, $J = 1.8$ Hz, 1H), 7.82 (d, $J = 1.8$ Hz, 1H), 7.33–7.21 (m, 5H), 7.06 (s, 1H), 5.30 (s, 2H), 4.26 (s, 2H), 2.37 (s, 3H), 2.21 (s, 3H); ESI MS m/z 319 $[\text{M} + \text{H}]^+$.

Preparation of 4-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3,4-dihydroquinoxalin-2(1H)-one (Example Compound 156)




[0347] Step 1: 4-Bromo-2-fluoro-1-nitrobenzene (1.00 g, 4.54 mmol), ethyl 2-(benzylamino)acetate (0.87 g, 4.5 mmol), and potassium carbonate (0.78 g, 5.7 mmol) in ethanol (15 mL) and water (11 mL) were heated at 85 °C for 10 h then stirred at rt for 8 h. The reaction mixture was diluted with water and brine then washed with methylene chloride. The resultant aqueous layer was filtered to afford **99** as an orange solid (1.28 g, 72%): ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.57 (d, $J = 8.6$ Hz, 1H), 7.37–7.21 (m, 6H), 6.97 (dd, $J = 8.7, 2.0$ Hz, 1H), 4.52 (s, 2H), 3.40 (s, 2H).

[0348] Step 2: To a solution of **99** (1.28 g, 3.51 mmol) in acetic acid (14 mL) at rt was added iron (470 mg, 8.4 mmol) and the resultant slurry was heated to 90 °C for 2.25 h. The mixture was cooled to rt and filtered through Celite, rinsing with methylene chloride. The filtrate was concentrated in vacuo and the resultant oil was partitioned between methylene chloride and saturated aqueous sodium bicarbonate. The aqueous layer was extracted with methylene chloride and the combined organic layers were dried with sodium sulfate, concentrated in vacuo, and purified by flash column chromatography (silica gel, 0–100% ethyl acetate/methylene chloride) to afford **100** as a white solid (430 mg, 39% yield): ^1H NMR (300 MHz, CDCl_3) δ 8.74 (br s, 1H), 7.39–7.26 (m, 5H), 6.89–6.85 (m, 2H), 6.62 (d, $J = 8.0$ Hz, 2H), 4.39 (s, 2H), 3.80 (s, 2H).


[0349] Step 3: Using the similar procedure used for Example Compound 7 step 1 on compound **100** afforded Example Compound 156 as a white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 10.58 (s, 1H), 7.38–7.34 (m, 4H), 7.30–7.23 (m, 1H), 6.87 (d, $J = 7.9$ Hz, 1H), 6.65 (d, $J = 7.9$ Hz, 1H), 6.51 (s, 1H), 4.46 (s, 2H), 3.86 (s, 2H), 2.15 (s, 3H), 1.97 (s, 3H); ESI m/z 334 $[\text{M} + \text{H}]^+$.

Preparation of 4-benzyl-6-(1-methyl-1*H*-pyrazol-5-yl)-3,4-dihydroquinoxalin-2(1*H*)-one (Example Compound 166)



[0350] Using the similar procedure used for **Example Compound 7** step 1 on compound **100** afforded **Example Compound 166** as a white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 10.62 (s, 1H), 7.37–7.33 (m, 5H), 7.29–7.25 (m, 1H), 6.90 (d, $J = 7.9$ Hz, 1H), 6.80 (dd, $J = 7.9, 1.8$ Hz, 1H), 6.70 (d, $J = 1.6$ Hz, 1H), 6.18 (d, $J = 1.8$ Hz, 1H), 4.49 (s, 2H), 3.83 (s, 2H), 3.58 (s, 3H); ESI m/z 319 $[\text{M} + \text{H}]^+$.

Preparation of (*R*)-4-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-3,4-dihydroquinoxalin-2(1*H*)-one (Example Compound 174)




[0351] **Step 1:** 4-Bromo-2-fluoro-1-nitrobenzene (0.50 g, 2.3 mmol), (*R*)-methyl 2-(benzylamino)propanoate (0.55 g, 2.3 mmol), and potassium carbonate (0.47 g, 3.4 mmol) in ethanol (8 mL) and water (6 mL) were heated at 85 °C for 10 h then stirred at rt for 8 h. The reaction mixture was diluted with water and filtered. The pH of the filtrate was adjusted to ~4 with 6*N* aqueous HCl and the resultant slurry was re-filtered to afford **101** as a sticky orange solid (not weighed; used directly in the next step).

[0352] **Step 2:** Using the similar procedure used for **Example Compound 156** step 2 on compound **101** afforded compound **102** as a white solid (430 mg, 39% yield): ^1H NMR (500 MHz,

DMSO- d_6) δ 10.57 (br s, 1H), 7.39–7.25 (m, 5H), 6.87–6.66 (m, 3H), 4.60 (d, J = 15.5 Hz, 1H), 4.29 (d, J = 15.2 Hz, 1H), 3.85 (q, J = 6.9 Hz, 1H), 1.08 (d, J = 6.7 Hz, 3H).

[0353] Step 3: Using the similar procedure used for **Example Compound 156** step 3 on compound **102** afforded **Example Compound 174** as an off-white solid: ^1H NMR (500 MHz, DMSO- d_6) δ 10.53 (s, 1H), 7.37–7.32 (m, 4H), 7.26–7.23 (m, 1H), 6.88 (d, J = 7.9 Hz, 1H), 6.66 (dd, J = 7.9, 1.7 Hz, 1H), 6.42 (d, J = 1.5 Hz, 1H), 4.54 (d, J = 15.6 Hz, 1H), 4.37 (d, J = 15.7 Hz, 1H), 3.98 (q, J = 6.7 Hz, 1H), 2.11 (s, 3H), 1.93 (s, 3H), 1.12 (d, J = 6.7 Hz, 3H); ESI m/z 348 $[\text{M} + \text{H}]^+$.

Preparation of 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one (Example Compound 118) and 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-*N*-ethyl-1H-imidazo[4,5-*b*]pyridin-2-amine (Example Compound 131)



[0354] Step 1: To a stirred solution of **26** (2.00 g, 10.6 mmol) in dry CH_2Cl_2 (50 mL) was added glacial acetic acid (0.61 mL, 10.8 mmol) and cyclopropanecarboxaldehyde (0.81 mL, 12.3 mmol). The solution was stirred at room temperature for 1 h and was cooled to 0 °C. Sodium borohydride (1.21 g, 31.8 mmol) was added carefully and the reaction was allowed to warm to room temperature. After stirring at ambient temperature for 15 h, saturated aq. NaHCO_3 (20 mL) was added to basify and then the mixture was extracted with CH_2Cl_2 (2×100 mL). The combined methylene chloride layers were dried over Na_2SO_4 , filtered and the filtrate was concentrated to a brown residue. The residue was diluted with CH_2Cl_2 (20 mL), the solution was loaded onto silica gel (120 g) and eluted with 0–70% ethyl acetate in hexanes to afford **103** (330 mg, 13%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 7.62 (d, J = 2.0 Hz, 1H), 6.83 (d, J = 1.5 Hz, 1H), 4.17 (br s, 2H), 3.39 (br s, 1H), 2.90 (d, J = 5.0 Hz, 1H), 2.89 (d, J = 5.0 Hz, 1H), 1.19–1.07 (m, 1H), 0.63–0.56 (m, 2H), 0.27–0.22 (m, 2H).


[0355] Step 2: To a mixture of **103** (300 mg, 1.24 mmol) and **3** (415 mg, 1.86 mmol) in 1,4-dioxane (10 mL) and water (2.5 mL) was added potassium carbonate (343 mg, 2.48 mmol) and tetrakis(triphenylphosphine)palladium(0) (76 mg, 0.062 mmol). The reaction was stirred and heated at 90 °C for 17 h. The mixture was diluted with methanol (20 mL) and silica gel (10 g) was added.

The suspension was concentrated to dryness and the resulting powder was loaded onto silica gel (80 g) and eluted with 0–80% ethyl acetate in hexanes. The clean product was concentrated to give **104** (312 mg, 97%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 7.48 (d, J = 1.5 Hz, 1H), 6.61 (d, J = 1.5 Hz, 1H), 4.27 (br s, 2H), 3.39 (br s, 1H), 2.92 (t, J = 6.0 Hz, 2H), 2.38 (s, 3H), 2.24 (s, 3H), 1.18–1.09 (m, 1H), 0.63–0.56 (m, 2H), 0.28–0.22 (m, 2H).

[0356] Step 3: To a solution of **104** (310 mg, 1.20 mmol), a catalytic amount of DMAP and 1,4-dioxane (4 mL) in a pressure tube was added 1,1'-carbonyldiimidazole (390 mg, 2.40 mmol). The tube was sealed and heated to 80 °C for 2 h. The reaction mixture was diluted with methanol (20 mL) and silica gel (10 g) was added. The suspension was concentrated to dryness and the resulting powder was loaded onto silica gel (40 g) and eluted with 0–80% ethyl acetate in hexanes. The clean product was concentrated to give 1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2(3*H*)-one (275 mg, 81%) as a yellow solid. A 50 mg sample was then purified by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example Compound 118** (37 mg) as a white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.90 (d, J = 1.5 Hz, 1H), 7.50 (d, J = 1.5 Hz, 1H), 3.81 (d, J = 7.0 Hz, 2H), 2.42 (s, 3H), 2.26 (s, 3H), 1.31–1.20 (m, 1H), 0.60–0.53 (m, 2H), 0.44–0.38 (m, 2H); ESI m/z 285 $[\text{M} + \text{H}]^+$.

[0357] Step 4: A solution of **Example Compound 118** (220 mg, 0.774 mmol) in phosphorus(V) oxychloride (3 mL) was placed in a sealed tube and heated at 110 °C for 6 h. The solvent was removed *in vacuo* and a saturated aq. NaHCO_3 solution (5 mL) was added. The mixture was extracted with ethyl acetate (2×20 mL) and the combined extracts were dried over Na_2SO_4 , filtered and the filtrate was concentrated. THF (5 mL) and 2.0 M ethylamine solution in THF (6 mL, 12.0 mmol) were then added and the reaction was heated at 70 °C for 17 h. The reaction was concentrated to dryness and the residue diluted with CH_2Cl_2 (5 mL). The resulting solution was loaded onto silica gel (40 g) and eluted with 0–80% ethyl acetate in hexanes. The clean product was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example Compound 131** (91 mg, 38%) as a white solid: ^1H NMR (500 MHz, CDCl_3) δ 7.93 (d, J = 2.0 Hz, 1H), 7.48 (d, J = 1.5 Hz, 1H), 3.98 (d, J = 6.5 Hz, 2H), 3.57 (q, J = 7.0 Hz, 2H), 2.42 (s, 3H), 2.26 (s, 3H), 1.30 (t, J = 7.0 Hz, 3H), 1.29–1.19 (m, 1H), 0.59–0.52 (m, 2H), 0.45–0.39 (m, 2H); ESI m/z 312 $[\text{M} + \text{H}]^+$.

Preparation of 4-(1-(cyclohexylmethyl)-2-methyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 191), 4-(1-(cyclopentylmethyl)-2-methyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 192) and 4-(1-(cyclobutylmethyl)-2-methyl-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 193)



[0358] Step 1: A mixture of 2,3-diamino-5-bromopyridine (10.0 g, 0.053 mol), cyclohexanecarboxaldehyde (6.08 g, 0.054 mol) and glacial acetic acid (3.05 mL) in dry CH₂Cl₂ (250 mL) was stirred for 1.5 h at room temperature. Sodium borohydride (6.06 g, 0.159 mol) was added portionwise over 20 min and the mixture was stirred for 17 h at room temperature. Saturated aq. NaHCO₃ was added until the mixture reached pH 8 (70 mL) and the aqueous layer extracted with CH₂Cl₂ (100 mL). The combined CH₂Cl₂ layers were combined, washed with water (500 mL), dried over Na₂SO₄, filtered and concentrated. The brown solid was taken up in methanol (100 mL) and silica gel (40 g) was added. The suspension was concentrated to dryness and the material was purified by chromatography (silica gel, 0–50% EtOAc/hexane then 0–10% EtOAc/CH₂Cl₂) to afford **105a** (1.30 g, 9%) as a brown-gray solid: ¹H NMR (500 MHz, CDCl₃) δ 7.60 (d, *J* = 2.0 Hz, 1H), 6.85 (d, *J* = 2.0 Hz, 1H), 4.11 (br s, 2H), 3.28 (br s, 1H), 2.88 (d, *J* = 5.0 Hz, 2H), 1.88–1.64 (m, 4H), 1.70–1.52 (m, 1H), 1.38–1.15 (m, 4H), 1.10–0.96 (m, 2H).

[0359] **105b** was prepared starting with cyclopentanecarbaldehyde (14% yield; brown-gray solid): ¹H NMR (500 MHz, CDCl₃) δ 7.60 (d, *J* = 2.0 Hz, 1H), 6.86 (d, *J* = 2.0 Hz, 1H), 4.14 (br s, 2H), 3.28 (br s, 1H), 2.99–2.93 (m, 2H), 2.23–2.11 (m, 1H), 1.88–1.71 (m, 2H), 1.70–1.53 (m, 4H), 1.32–1.23 (m, 2H).

[0360] **105c** was prepared starting with cyclobutanecarbaldehyde (15% yield; brown-gray solid): ¹H NMR (500 MHz, CDCl₃) δ 7.61 (d, *J* = 2.0 Hz, 1H), 6.86 (d, *J* = 2.0 Hz, 1H), 4.12 (br s, 2H), 3.14

(br s, 1H), 3.09–3.02 (m, 2H), 2.67–2.52 (m, 1H), 2.18–2.11 (m, 2H), 2.07–1.86 (m, 2H), 1.80–1.71 (m, 2H).

[0361] Step 2: To a mixture of **105a** (500 mg, 1.76 mmol), 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (589 mg, 2.64 mmol), potassium carbonate (487 mg, 3.52 mmol), water (4 mL) and 1,4-dioxane (16 mL) was added tetrakis(triphenylphosphine)palladium (0) and the mixture was heated to 90 °C for 17 h. The two phase mixture was diluted with methanol (20 mL) and silica gel was added. After concentrating to dryness the material was purified by chromatography (silica gel, 0–80% EtOAc/hexane) to afford **106a** (551 mg, 99%) as a brown solid: ¹H NMR (500 MHz, CDCl₃) δ 7.47 (d, *J* = 2.0 Hz, 1H), 6.62 (d, *J* = 2.0 Hz, 1H), 4.25 (br s, 2H), 3.34 (br s, 1H), 2.92 (t, *J* = 6.0 Hz, 2H), 2.38 (s, 3H), 2.25 (s, 3H), 1.88–1.67 (m, 4H), 1.67–1.56 (m, 1H), 1.33–1.19 (m, 4H), 1.10–0.96 (m, 2H).

[0362] 106b was prepared starting with **105b** (96% yield; brown-gray solid): ¹H NMR (500 MHz, CDCl₃) δ 7.47 (d, *J* = 1.5 Hz, 1H), 6.64 (d, *J* = 1.5 Hz, 1H), 4.25 (br s, 2H), 3.28 (br s, 1H), 2.99 (t, *J* = 6.0 Hz, 1H), 2.38 (s, 3H), 2.24 (s, 3H), 2.24–2.17 (m, 1H), 1.90–1.81 (m, 2H), 1.72–1.55 (m, 4H), 1.38–1.22 (m, 2H).

[0363] 106c was prepared starting with **105c** (95% yield; brown-gray solid): ¹H NMR (500 MHz, CDCl₃) δ 7.65 (d, *J* = 1.5 Hz, 1H), 6.64 (d, *J* = 2.0 Hz, 1H), 4.26 (br s, 2H), 3.18 (br s, 1H), 3.09 (t, *J* = 6.0 Hz, 1H), 2.67–2.58 (m, 1H), 2.20–2.12 (m, 2H), 2.02–1.86 (m, 2H), 1.82–1.72 (m, 2H).


[0364] Step 3: A solution of **106a** (100 mg, 0.33 mmol), triethylorthoacetate (5 mL) and glacial acetic acid (0.10 mL) was heated in a sealed tube for 24 hours at 80 °C. The mixture was evaporated to dryness and methanol (10 mL), saturated aq. NaHCO₃ (5 mL) and silica gel (10 g) were added. After concentrating to dryness the resulting powder was loaded onto silica gel and eluted with 0–5% methanol in methylene chloride. The clean product was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH₃CN in H₂O and the clean fractions were frozen and lyophilized to give **Example Compound 191** (56 mg, 52%) as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.30 (d, *J* = 1.5 Hz, 1H), 7.96 (d, *J* = 2.0 Hz, 1H), 4.14 (d, *J* = 7.5 Hz, 2H), 2.69 (s, 3H), 2.44 (s, 3H), 2.28 (s, 3H), 1.95–1.82 (m, 1H), 1.76–1.50 (m, 5H), 1.29–1.07 (m, 5H); ESI *m/z* 325 [M + H]⁺.

[0365] Starting with **106b**, **Example Compound 192** (31 mg, 29%) was prepared as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.30 (d, *J* = 2.0 Hz, 1H), 7.98 (d, *J* = 2.0 Hz, 1H), 4.26 (d, *J* = 8.0 Hz, 2H), 2.71 (s, 3H), 2.49–2.38 (m, 1H), 2.44 (s, 3H), 2.28 (s, 3H), 1.80–1.68 (m, 4H), 1.66–1.57 (m, 2H), 1.40–1.27 (m, 2H); ESI *m/z* 311 [M + H]⁺.

[0366] Starting with **106c**, **Example Compound 193** (33 mg, 30%) was prepared as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.30 (d, *J* = 1.5 Hz, 1H), 8.00 (d, *J* = 1.5 Hz, 1H), 4.33 (d, *J* = 7.0 Hz,

2H), 2.92–2.80 (m, 1H), 2.70 (s, 3H), 2.45 (s, 3H), 2.28 (s, 3H), 2.10–1.98 (m, 2H), 1.96–1.81 (m, 4H); ESI m/z 297 $[M + H]^+$.


Preparation of 1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one (Example Compound 202) and 1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2(3H)-one (Example Compound 203)



[0367] A solution of **106b** (1.30 g, 4.54 mmol), 1,1'-carbonyldiimidazole (1.47 g) and *N,N*-dimethylaminopyridine (5 mg) in 1,4-dioxane (16 mL) was heated at 80 °C for 2 h and cooled to room temperature. To the mixture was added silica gel (10 g) and methanol (20 mL) and the suspension was concentrated to a dry powder. This material was loaded onto silica gel (80 g) and eluted with 0–90% ethyl acetate in hexanes to give 1.08 g (76%) of **Example Compound 202** as a yellow solid. A 100 mg sample of the product was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH₃CN in H₂O and the clean fractions were frozen and lyophilized to give **Example Compound 202** as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 7.90 (d, *J* = 1.5 Hz, 1H), 7.47 (d, *J* = 2.0 Hz, 1H), 3.86 (d, *J* = 7.5 Hz, 2H), 2.52–2.38 (m, 1H), 2.41 (s, 3H), 2.25 (s, 3H), 1.78–1.68 (m, 4H), 1.60–1.52 (m, 2H), 1.41–1.30 (m, 2H); ESI m/z 313 $[M + H]^+$.

[0368] Starting with **106c**, **Example Compound 203** (76% yield, white solid) was synthesized in a similar procedure as **Example Compound 202**: ¹H NMR (500 MHz, CD₃OD) δ 7.89 (d, *J* = 1.5 Hz, 1H), 7.46 (d, *J* = 2.0 Hz, 1H), 3.94 (d, *J* = 7.0 Hz, 2H), 2.86–2.77 (m, 1H), 2.41 (s, 3H), 2.25 (s, 3H), 2.08–1.98 (m, 2H), 1.94–1.80 (m, 4H); ESI m/z 299 $[M + H]^+$.


Preparation of 4-(1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-*b*]pyridin-2-yl)morpholine (Example Compound 208) and 4-(2-(azetidin-1-yl)-1-(cyclopentylmethyl)-1H-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example Compound 209)



[0369] A solution of **Example Compound 202** (175 mg, 0.56 mmol) and phosphorus(V) oxychloride (4 mL) was heated to 110 °C for 17 h. The reaction was concentrated in vacuo and saturated aq. NaHCO₃ (5 mL) and ethyl acetate (20 mL) were added. The ethyl acetate layer was separated, dried over Na₂SO₄, filtered and the filtrate was concentrated to a dark yellow solid. The solid was dissolved in THF (5 mL) and morpholine (732 mg, 8.40 mmol) was added. The stirred solution was heated to 70 °C for 17 h. To the cooled mixture was added silica gel (5 g) and methanol (20 mL) and the suspension was concentrated to a dry powder. This material was loaded onto silica gel (40 g) and eluted with 0–3% methanol in methylene chloride to give 143 mg (67%) of product as an off-white solid. The product sample was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH₃CN in H₂O and the clean fractions were frozen and lyophilized to give **Example Compound 208** as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 8.17 (d, *J* = 1.5 Hz, 1H), 7.81 (d, *J* = 2.0 Hz, 1H), 4.14 (d, *J* = 7.5 Hz, 2H), 3.87 (t, *J* = 5.0 Hz, 4H), 3.41 (t, *J* = 5.0 Hz, 4H), 2.58–2.49 (m, 1H), 2.43 (s, 3H), 2.27 (s, 3H), 1.75–1.66 (m, 2H), 1.62–1.50 (m, 4H), 1.30–1.19 (m, 2H). ESI *m/z* 382 [M + H]⁺.

[0370] **Example Compound 209** was synthesized using a similar procedure as was used for **Example Compound 208**; **Example Compound 209** was collected as a white solid (166 mg, 84%): ¹H NMR (500 MHz, CD₃OD) δ 8.00 (d, *J* = 1.5 Hz, 1H), 7.59 (d, *J* = 1.5 Hz, 1H), 4.42–4.37 (m, 4H), 4.01 (d, *J* = 8.0 Hz, 2H), 2.57–2.44 (m, 2H), 2.50–2.41 (m, 1H), 2.41 (s, 3H), 2.25 (s, 3H), 1.76–1.51 (m, 6H), 1.32–1.22 (m, 2H). ESI *m/z* 352 [M + H]⁺.

Preparation of 4-(1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2-yl)morpholine (Example 210) and 4-(2-(azetidin-1-yl)-1-(cyclobutylmethyl)-1*H*-imidazo[4,5-*b*]pyridin-6-yl)-3,5-dimethylisoxazole (Example 211)




[0371] **Example 210** and **Example 211** were synthesized using a similar procedure as was used for **Example 208**.

[0372] **Example 210** collected as white solid (176 mg, 82% yield): ¹H NMR (500 MHz, CD₃OD) δ 8.16 (d, *J* = 1.5 Hz, 1H), 7.80 (d, *J* = 2.0 Hz, 1H), 4.24 (d, *J* = 7.0 Hz, 2H), 3.88 (t, *J* = 5.0 Hz, 4H), 3.41 (t, *J* = 5.0 Hz, 4H), 2.93–2.82 (m, 1H), 2.43 (s, 3H), 2.27 (s, 3H), 1.98–1.91 (m, 2H), 1.90–1.76 (m, 4H). ESI *m/z* 368 [M + H]⁺.

[0373] **Example 211** collected as white solid (180 mg, 91% yield): ¹H NMR (500 MHz, CD₃OD) δ 7.99 (d, *J* = 2.0 Hz, 1H), 7.61 (d, *J* = 2.0 Hz, 1H), 4.38 (m, 4H), 4.10 (d, *J* = 7.0 Hz, 2H), 2.88–


2.79 (m, 1H), 2.57–2.48 (m, 2H), 2.41 (s, 3H), 2.25 (s, 3H), 2.04–1.95 (m, 2H), 1.95–1.78 (m, 4H). ESI m/z 338 $[M + H]^+$.

Preparation of 1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-*N*-(tetrahydro-2*H*-pyran-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2-amine (Example 222)



[0374] A solution of **Example 202** (175 mg, 0.56 mmol) and phosphorus (V) oxychloride (4 mL) was heated to 110 °C for 17 h. The reaction was concentrated in vacuo and saturated aq. NaHCO_3 (5 mL) and ethyl acetate (20 mL) were added. The ethyl acetate layer was separated, dried over Na_2SO_4 , filtered and the filtrate was concentrated to a dark yellow solid. The solid was dissolved in propionitrile (5 mL) and 4-aminotetrahydropyran (283 mg, 28.0 mmol) was added. The stirred solution was heated to 180 °C in a microwave reactor for 6 h. To the cooled mixture was added silica gel (10 g) and methanol (20 mL) and the suspension was concentrated to a dry powder. This material was loaded onto silica gel (40 g) and eluted with 0–3% methanol in methylene chloride to give a yellow solid. The material was then purified by reverse phase HPLC on a Polaris column eluting with 10–90% CH_3CN in H_2O and the clean fractions were frozen and lyophilized to give **Example 222** (70 mg, 31%) as a white solid: ^1H NMR (500 MHz, CD_3OD) δ 7.94 (d, $J = 1.5$ Hz, 1H), 7.50 (d, $J = 2.0$ Hz, 1H), 4.17–4.05 (m, 1H), 4.05 (d, $J = 8.0$ Hz, 2H), 4.02–3.97 (m, 2H), 3.57 (t, $J = 11.75$ Hz, 2H), 2.44–2.36 (m, 1H), 2.41 (s, 3H), 2.25 (s, 3H), 2.08–2.00 (m, 2H), 1.78–1.64 (m, 6H), 1.62–1.54 (m, 2H), 1.38–1.25 (m, 2H). ESI m/z 396 $[M + H]^+$.


Preparation of 1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-*N*-(tetrahydro-2*H*-pyran-4-yl)-1*H*-imidazo[4,5-*b*]pyridin-2-amine (Example Compound 223)



[0375] **Example Compound 223** was synthesized using a similar procedure as was used for **Example Compound 222**. **Example Compound 223** collected as white solid (45 mg, 20% yield): ^1H NMR (500 MHz, CD_3OD) δ 7.93 (d, $J = 2.0$ Hz, 1H), 7.52 (d, $J = 2.0$ Hz, 1H), 4.17–4.05 (m, 1H), 4.10

(d, $J = 7.5$ Hz, 2H), 4.03–3.97 (m, 2H), 3.56 (t, $J = 11.75$ Hz, 2H), 2.86–2.78 (m, 1H), 2.41 (s, 3H), 2.25 (s, 3H), 2.08–1.92 (m, 8H), 1.75–1.64 (m, 2H). ESI m/z 382 $[M + H]^+$.

Preparation of 4-(1-benzyl-7-methoxy-2-(trifluoromethyl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole (Example Compound 241)




[0376] Step 1: To a solution of **107** (136 mg, 0.627 mmol) in THF (6 mL) was added di-tert-butyl dicarbonate (137 mg, 0.627 mmol) and the reaction was stirred at rt for 16 h. The reaction was then concentrated and the residue was purified by chromatography (silica gel, 0–25% ethyl acetate/hexanes) to afford an off-white solid which was dissolved in CH_2Cl_2 (3 mL), benzaldehyde in CH_2Cl_2 (2 mL) was added followed by AcOH (2 drops). The reaction was stirred at rt for 1 h and $\text{NaBH}(\text{OAc})_3$ (283 mg, 1.34 mmol) was added. The reaction was then stirred at rt for 16 h. The reaction was quenched with saturated NaHCO_3 and extracted with CH_2Cl_2 (2 $\times$ 50 mL). The combined organics were dried with Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 0–30% ethyl acetate/hexanes) to afford **108** (97 mg, 38%) as an off-white solid: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.43 (s, 1H), 7.32–7.26 (m, 4H), 7.23–7.00 (m, 1H), 6.95 (s, 2H), 4.87 (t, $J = 6.9$ Hz, 1H), 4.31 (d, $J = 6.9$ Hz, 2H), 3.64 (s, 3H), 1.42 (s, 9H).

[0377] Step 2: To a solution of **108** (135 mg, 0.332 mmol) in CH_2Cl_2 (5 mL) at 0 °C was added TFA (0.51 mL, 6.63 mmol) and the reaction was warmed to room temperature and stirred for 16 h. The reaction was then concentrated to afford **109** (114 mg, 90%): ESI m/z 385 $[M + H]^+$.

[0378] Step 3: Using the procedure used in General Procedure B step 1 starting with compound **109** (114 mg, 0.296 mmol) afforded **Example Compound 241** (45 mg, 38%) as an off white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.72 (d, $J = 8.4$ Hz, 1H), 7.36–7.26 (m, 4H), 7.03–7.00 (m, 2H), 5.81 (s, 2H), 3.13 (s, 3H), 2.27 (s, 3H), 2.09 (s, 3H); ESI m/z 402 $[M + H]^+$.

Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazole-2-carboximidamide (Example Compound 243) and 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1*H*-benzo[d]imidazole-2-carboxamide (Example Compound 244)




[0379] Step 1: To a solution of **20** (3.00 g, 10.8 mmol) in 1,4-dioxane (60 mL) and water (6 mL) was added 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (2.90 g, 13.0 mmol), tetrakis(triphenylphosphine)palladium(0) (624 mg, 0.54 mmol) and potassium carbonate (2.98 g, 21.6 mmol). The reaction mixture was purged with nitrogen and heated at 90°C for 18 h. The mixture was cooled to room temperature, concentrated and purified by chromatography (silica gel, 0–20% ethyl acetate in hexanes) to afford **110** (3.18 g, 99%) as a yellow solid: ^1H NMR (500 MHz, CDCl_3) δ 7.38 (d, $J = 8.3$ Hz, 2H), 7.34 (t, $J = 7.3$ Hz, 2H), 7.28 (t, $J = 7.1$ Hz, 1H), 6.78 (d, $J = 7.8$ Hz, 1H), 6.55 (dd, $J = 1.8, 7.7$ Hz, 1H), 6.43 (d, $J = 1.8$ Hz, 1H), 4.35 (s, 2H), 3.88 (s, 1H), 3.42 (s, 2H), 2.23 (s, 3H), 2.11 (s, 3H); ESI m/z 294 $[\text{M} + \text{H}]^+$.

[0380] Step 1: To a solution of **110** (100 mg, 0.34 mmol) in acetic acid (2 mL) was added methyl 2,2,2-trichloroacetimidate (66 mg, 0.38 mmol) at room temperature. The reaction mixture was stirred at room temperature for 1 h and then water was added. The precipitate formed was collected by filtration, the filter cake was washed with water, and dried under vacuum at 40°C to afford **111** (110 mg, 77%) as an off-white solid: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.93 (dd, $J = 0.4, 8.4$ Hz, 1H), 7.40–7.25 (m, 4H), 7.19–7.11 (m, 3H), 5.96 (s, 2H), 2.21 (s, 3H), 2.03 (s, 3H); ESI m/z 422 $[\text{M} + \text{H}]^+$.

[0381] Step 2: To a solution of **111** (100 mg, 0.238 mmol) in ethanol (1 mL) was added concentrated ammonium hydroxide (1 mL). The reaction mixture was heated at 120°C for 1 h. The mixture was cooled to room temperature and concentrated. The residue was purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes then to 20% methanol in ethyl acetate) followed by reverse phase HPLC on a Polaris C_{18} column eluting with 10–90% CH_3CN in H_2O to afford Example Compound **243** (21 mg, 25%) and Example Compound **244** (29 mg, 35%) as an off-white

solids. **Example Compound 243**: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.77 (d, J = 8.3 Hz, 1H), 7.49 (s, 1H), 7.36 (s, 1H), 7.33–7.19 (m, 6H), 6.58 (s, 2H), 6.27 (s, 2H), 2.32 (s, 3H), 2.15 (s, 3H); ESI m/z 346 $[\text{M} + \text{H}]^+$; **Example Compound 244**: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.38 (s, 1H), 7.92 (s, 1H), 7.82 (d, J = 8.5 Hz, 1H), 7.63 (d, J = 1.0 Hz, 1H), 7.33–7.28 (m, 5H), 7.27–7.22 (m, 1H), 6.02 (s, 2H), 2.35 (s, 3H), 2.18 (s, 3H); ESI m/z 347 $[\text{M} + \text{H}]^+$.


Preparation of 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-*N*-(pyridin-3-yl)-1*H*-benzo[d]imidazol-2-amine (Example Compound 248)



[0382] Step 1: A solution of **81** (500 mg, 1.57 mmol) and phosphorus(V) oxychloride (2 mL) was heated to 100 °C for 17 h. The reaction was concentrated in vacuo and saturated aq. NaHCO_3 (5 mL) and ethyl acetate (20 mL) were added. The ethyl acetate layer was separated, dried over Na_2SO_4 , filtered and concentrated. The residue was purified by chromatography (silica gel, 0–30% ethyl acetate in hexanes) to afford **112** (415 mg, 78%) as a light brown oil: ESI m/z 338 $[\text{M} + \text{H}]^+$.

[0383] Step 2: A mixture of **112** (20 mg, 0.06 mmol), pyridin-3-amine (28 mg, 0.30 mmol) and $p\text{-TsOH}\cdot\text{H}_2\text{O}$ (22 mg, 0.12 mmol) in NMP was heated at 190 °C in a microwave reactor for 2 h. The mixture was concentrated, and the residue was purified by chromatography (silica gel, 0–100% ethyl acetate in hexanes) to afford **Example Compound 248** as an light brown oil: ESI m/z 396 $[\text{M} + \text{H}]^+$.

Preparation of 3-(1-benzyl-1*H*-benzo[d]imidazol-6-yl)-4-ethyl-1*H*-1,2,4-triazol-5(4*H*)-one (Example Compound 249)





[0384] **Step 1:** A solution of **113** (1.20 g, 4.51 mmol) and hydrazine monohydrate (3.27 mL, 67.65 mmol) in EtOH (20 mL) was heated to reflux for 16 h. The mixture was cooled to rt, the precipitate was collected by filtration, the filter cake was dried to afford **114** (1.02 g, 85%) as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 9.74 (s, 1H), 8.54 (s, 1H), 8.07 (s, 1H), 7.73–7.67 (m, 2H), 7.38–7.26 (m, 5H), 5.54 (s, 2H), 4.47 (s, 2H).

[0385] **Step 2:** A suspension of **114** (500 mg, 1.88 mmol) and ethylisocyanate (160 mg, 2.26 mmol) in THF was stirred at rt for 5 h. The mixture was filtered, the filter cake was washed with ethyl acetate, and dried to afford **115** (610 mg, 96%) as a white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 10.09 (s, 1H), 8.57 (s, 1H), 8.14 (s, 1H), 7.81–7.79 (m, 2H), 7.72 (d, J = 8.4 Hz, 1H), 7.38–7.28 (m, 5H), 6.47 (t, J = 5.4 Hz, 1H), 5.55 (s, 2H), 3.09–3.00 (m, 2H), 1.00 (t, J = 7.2 Hz, 3H).

[0386] **Step 3:** A suspension of **115** (337 mg, 1.0 mmol) in 3 N NaOH (5 mL) was heated to reflux for 16 h. The mixture was adjusted to pH 8 by 2 N HCl, and then was extracted with CH_2Cl_2 (3 $\times$ 50 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The residue was triturated with EtOAc/ CH_2Cl_2 to afford **Example Compound 249** as an off-white solid: ^1H NMR (300 MHz, DMSO- d_6) δ 11.85 (s, 1H), 8.59 (s, 1H), 7.81–7.76 (m, 2H), 7.43 (dd, J = 8.1, 1.5 Hz, 1H), 7.35–7.28 (m, 5H), 5.58 (s, 2H), 3.63 (q, J = 7.2, Hz 2H), 0.98 (t, J = 7.2 Hz, 3H); ESI m/z 320 $[\text{M} + \text{H}]^+$.

Table 2: Example Compounds

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
1	9-benzyl-2-(3,5-dimethylisoxazol-4-yl)-9H-purin-6-amine		A	^1H NMR (300 MHz, DMSO- d_6) δ 8.29 (s, 1H), 7.36–7.28 (m, 7H), 5.38 (s, 2H), 2.73 (s, 3H), 2.51 (s, 3H); ESI m/z 321 $[\text{M} + \text{H}]^+$.	96.6
2	3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		No general procedure	^1H NMR (300 MHz, DMSO- d_6) δ 11.31 (s, 1H), 7.40 (d, J = 7.8 Hz, 1H), 7.34–7.25 (m, 5H), 7.15 (d, J = 7.8 Hz, 1H), 5.03 (s, 2H), 2.47 (s, 3H), 2.28 (s, 3H); ESI m/z 321 $[\text{M} + \text{H}]^+$.	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
3	1-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		No general procedure	^1H NMR (300 MHz, DMSO- d_6) δ 11.76 (s, 1H), 7.44 (d, J = 7.8 Hz, 1H), 7.36-7.28 (m, 5H), 7.11 (d, J = 7.8 Hz, 1H), 5.05 (s, 2H), 2.49 (s, 3H), 2.32 (s, 3H); ESI m/z 321 $[\text{M} + \text{H}]^+$.	>99
4	4-(3-benzyl-3H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.62 (s, 1H), 8.36 (br s, 1H), 7.65 (s, 1H), 7.45 (s, 5H), 5.96 (s, 2H), 2.34 (s, 3H), 2.17 (s, 3H); ESI m/z 305 $[\text{M} + \text{H}]^+$.	>99
5	4-(1-benzyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.62 (s, 1H), 8.36 (br s, 1H), 7.65 (s, 1H), 7.45 (s, 5H), 5.96 (s, 2H), 2.34 (s, 3H), 2.17 (s, 3H); ESI m/z 305 $[\text{M} + \text{H}]^+$.	>99
6	3-benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[d]oxazol-2(3H)-one		No general procedure	^1H NMR (500 MHz, DMSO- d_6) δ 7.47-7.42 (m, 3H), 7.40-7.34 (m, 2H), 7.34-7.28 (m, 1H), 7.23 (d, J = 1.6 Hz, 1H), 7.12 (dd, J = 8.2 Hz, 7.7 Hz, 1H), 5.07 (s, 2H), 2.33 (s, 3H), 2.15 (s, 3H); ESI m/z 321 $[\text{M} + \text{H}]^+$.	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
7	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine		C	^1H NMR (300 MHz, CDCl_3) δ 7.95 (s, 1H), 7.37-7.34 (m, 3H), 7.23-7.20 (m, 2H), 6.46 (d, J = 1.2 Hz, 1H), 6.40 (d, J = 1.2 Hz, 1H), 5.34 (s, 2H), 2.31 (s, 3H), 2.16 (s, 3H); ESI MS m/z 319 $[\text{M} + \text{H}]^+$	>99
8	1-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-7-amine		C	^1H NMR (300 MHz, CDCl_3) δ 8.15 (s, 1H), 7.43-7.40 (m, 3H), 7.23 (d, J = 1.2 Hz, 1H), 7.20-7.17 (m, 2H), 6.39 (d, J = 1.2 Hz, 1H), 5.69 (s, 2H), 2.40 (s, 3H), 2.27 (s, 3H); ESI MS m/z 319 $[\text{M} + \text{H}]^+$	95.2
9	N,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine		C	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.27 (s, 1H), 7.40-7.18 (m, 10H), 6.62 (d, J = 1.2 Hz, 1H), 6.57 (t, J = 6.0 Hz, 1H), 5.97 (d, J = 1.2 Hz, 1H), 5.41 (s, 2H), 4.48 (d, J = 6.0 Hz, 2H), 2.12 (s, 3H), 1.94 (s, 3H); ESI MS m/z 409 $[\text{M} + \text{H}]^+$	>99
10	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		No general procedure	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.81 (s, 1H), 7.90 (d, J = 2.1 Hz, 1H), 7.44-7.25 (m, 6H), 5.05 (s, 2H), 2.34 (s, 3H), 2.16 (s, 3H); MM m/z 321 $[\text{M} + \text{H}]^+$	>99





Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
11	1-benzyl-7-(3,5-dimethylisoxazol-4-yl)quinoxalin-2(1H)-one		No general procedure	¹ H NMR (300 MHz, CDCl ₃) δ 8.43 (s, 1H), 7.94 (d, J = 8.2 Hz, 1H), 7.35-7.32 (m, 2H), 7.29-7.27 (m, 1H), 7.21-7.18 (m, 3H), 7.04 (s, 1H), 5.51 (s, 2H), 2.16 (s, 3H), 2.02 (s, 3H); ESI m/z 332 [M + H] ⁺ .	>99
12	1-benzyl-7-(3,5-dimethylisoxazol-4-yl)-3,4-dihydroquinazolin-2(1H)-one		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.34-7.21 (m, 7H), 6.90 (dd, J = 7.5, 1.0 Hz, 1H), 6.58 (d, J = 1.0 Hz, 1H), 5.09 (s, 2H), 4.43 (s, 2H), 2.06 (s, 3H), 1.89 (s, 3H); MM m/z 334 [M + H] ⁺ .	>99
13	4-(1-benzyl-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	¹ H NMR (500 MHz, CD ₃ OD) δ 8.32 (d, J = 1.0 Hz, 1H), 7.78 (d, J = 1.0 Hz, 1H), 7.36-7.29 (m, 3H), 7.20-7.17 (m, 2H), 5.56 (s, 2H), 2.69 (s, 3H), 2.36 (s, 3H), 2.18 (s, 3H); ESI m/z 319 [M + H] ⁺ .	>99
14	4-(1-(cyclopropylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole		F	¹ H NMR (500 MHz, CD ₃ OD) δ 8.03 (d, J = 1.5 Hz, 1H), 7.93 (d, J = 1.5 Hz, 1H), 4.27 (d, J = 7.0 Hz, 2H), 2.75 (s, 3H), 2.46 (s, 3H), 2.30 (s, 3H), 1.38-1.28 (m, 1H), 0.65-0.60 (m, 2H), 0.51-0.46 (m, 2H). ESI m/z 327 [M + H] ⁺ .	97.3

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
15	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one		G	^1H NMR (300 MHz, DMSO- d_6) δ 12.11 (s, 1H), 7.72 (d, J = 1.5 Hz, 1H), 7.50 (d, J = 1.5 Hz, 1H), 7.42-7.28 (m, 5H), 5.13 (s, 2H), 2.35 (s, 3H), 2.15 (s, 3H); ESI m/z 365 $[\text{M} + \text{H}]^+$.	98.5
16	4-amino-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one		G	^1H NMR (500 MHz, DMSO- d_6) δ 10.44 (s, 1H), 7.36-7.25 (m, 5H), 6.28 (s, 2H), 5.04 (s, 2H), 4.95 (s, 2H), 2.28 (s, 3H), 2.10 (s, 3H); ESI m/z 335 $[\text{M} + \text{H}]^+$.	98.6
17	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-ethoxy-1H-benzo[d]imidazol-4-amine		No general procedure	^1H NMR (300 MHz, CDCl_3) δ 7.35-7.20 (m, 5H), 6.33 (d, J = 1.5 Hz, 1H), 6.30 (d, J = 1.5 Hz, 1H), 5.13 (s, 2H), 4.68 (q, J = 6.9 Hz, 2H), 4.30 (br.s, 2H), 2.30 (s, 3H), 2.16 (s, 3H), 1.49 (t, J = 7.2 Hz, 3H); ESI m/z 363 $[\text{M} + \text{H}]^+$.	99
18	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-4-nitro-1H-benzo[d]imidazol-2-amine		F	^1H NMR (300 MHz, CDCl_3) δ 7.84 (d, J = 1.5 Hz, 1H), 7.42-7.35 (m, 3H), 7.16-7.13 (m, 2H), 7.03 (d, J = 1.5 Hz, 1H), 5.15 (s, 2H), 4.29 (t, J = 5.4 Hz, 1H), 3.78-3.69 (m, 2H), 2.36 (s, 3H), 2.21 (s, 3H), 1.27 (t, J = 7.5 Hz, 3H); ESI m/z 392 $[\text{M} + \text{H}]^+$.	99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
19	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-ethyl-1H-benzo[d]imidazole-2,4-diamine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.20 (m, 5H), 6.62 (t, J = 5.4 Hz, 1H), 6.30 (d, J = 1.5 Hz, 1H), 6.21 (d, J = 1.5 Hz, 1H), 5.19 (s, 2H), 4.83 (s, 2H), 3.47-3.38 (m, 2H), 2.28 (s, 3H), 2.11 (s, 3H), 1.22 (t, J = 7.2 Hz, 3H); ESI m/z 362 [M + H] ⁺ .	96.8
20	methyl 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carboxylate		J	¹ H NMR (500 MHz, CD ₃ OD) δ 7.54 (d, J = 1.5 Hz, 1H), 7.37-7.24 (m, 5H), 7.07 (d, J = 1.5 Hz, 1H), 5.14 (s, 2H), 3.97 (s, 3H), 2.27 (s, 3H), 2.09 (s, 3H); ESI m/z 378 [M + H] ⁺ .	>99
21	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carboxamide		I	¹ H NMR (500 MHz, CD ₃ OD) δ 7.41 (d, J = 1.3 Hz, 1H), 7.37-7.24 (m, 5H), 7.00 (d, J = 1.4 Hz, 1H), 5.13 (s, 2H), 2.28 (s, 3H), 2.11 (s, 3H); ESI m/z 363 [M + H] ⁺ .	98.3
22	4-(aminomethyl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2(3H)-one		J	¹ H NMR (500 MHz, CD ₃ OD) δ 7.37-7.23 (m, 5H), 6.99 (d, J = 1.4 Hz, 1H), 6.77 (d, J = 1.4 Hz, 1H), 5.10 (s, 2H), 3.93 (s, 2H), 2.27 (s, 3H), 2.10 (s, 3H); ESI m/z 349 [M + H] ⁺ .	93.9

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
23	5-(3,5-dimethylisoxazol-4-yl)-N-phenyl-1H-pyrrolo[3,2-b]pyridin-3-amine		M	¹ H NMR (300 MHz, DMSO-d ₆) δ 11.1 (d, J = 1.8 Hz, 1H), 7.82 (d, J = 8.4 Hz, 1H), 7.61 (d, J = 2.7 Hz, 1H), 7.43 (s, 1H), 7.25 (d, J = 8.4 Hz, 1H), 7.09 (d, J = 8.4 Hz, 1H), 7.07 (d, J = 7.2 Hz, 1H), 6.85 (d, J = 7.5 Hz, 2H), 6.60 (t, J = 7.2 Hz, 1H), 2.48 (s, 3H), 2.29 (s, 3H); ESI MS m/z 305 [M + H] ⁺ .	>99
24	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-b]pyridine 4-oxide		N	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.21 (d, J = 0.9 Hz, 1H), 7.83 (d, J = 0.9 Hz, 1H), 7.40-7.35 (m, 2H), 7.20-7.14 (m, 2H), 5.59 (s, 2H), 2.69 (s, 3H), 2.45 (s, 3H), 2.27 (s, 3H); ESI MS m/z 353 [M + H] ⁺ .	>99
25	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-b]pyridin-5(4H)-one		N	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.0 (s, 1H), 8.07 (s, 1H), 7.36-7.31 (m, 2H), 7.19-7.13 (m, 2H), 5.45 (s, 2H), 2.30 (s, 6H), 2.14 (s, 3H); ESI MS m/z 353 [M + H] ⁺ .	96.2



Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
26	4-(3-benzyl-3H-imidazo[4,5-b]pyridin-5-yl)-3,5-dimethylisoxazole		No general procedure	^1H NMR (300 MHz, DMSO- d_6) δ 8.67 (s, 1H), 8.17 (d, J = 8.1 Hz, 1H), 7.44 (d, J = 8.1 Hz, 1H), 7.36-7.27 (m, 5H), 5.52 (s, 2H), 2.54 (s, 3H), 2.34 (s, 3H); ESI m/z 305 $[\text{M} + \text{H}]^+$.	98
27	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-benzo[d]imidazol-4-amine		C	^1H NMR (300 MHz, DMSO- d_6) δ 8.23 (s, 1H), 7.42 (dd, J = 8.0, 6.0 Hz, 2H), 7.17 (dd, J = 9.0, 9.0 Hz, 2H), 6.62 (s, 1H), 6.32 (s, 1H), 5.40 (s, 4H), 2.33 (s, 3H), 2.16 (s, 3H); ESI m/z 337 $[\text{M} + \text{H}]^+$.	>99
28	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N-methyl-1H-benzo[d]imidazol-4-amine		No general procedure	^1H NMR (500 MHz, DMSO- d_6) δ 8.22 (s, 1H), 7.43 (dd, J = 8.8, 5.5 Hz, 2H), 7.16 (dd, J = 8.8, 5.5 Hz, 2H), 6.65 (d, J = 1.0 Hz, 1H), 6.09 (d, J = 1.0 Hz, 1H), 5.85 (q, J = 5.0 Hz, 1H), 5.41 (s, 2H), 2.83 (d, J = 5.5 Hz, 3H), 2.35 (s, 3H), 2.17 (s, 3H); ESI m/z 351 $[\text{M} + \text{H}]^+$.	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
29	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N,N-dimethyl-1H-benzo[d]imidazol-4-amine		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.28 (s, 1H), 7.41 (dd, J = 8.5, 5.5 Hz, 2H), 7.17 (dd, J = 9.0, 9.0 Hz, 2H), 6.85 (d, J = 1.0 Hz, 1H), 6.25 (d, J = 1.0 Hz, 1H), 5.43 (s, 2H), 3.18 (s, 6H), 2.35 (s, 3H), 2.18 (s, 3H); ESI m/z 365 [M + H] ⁺ .	98.1
30	3,5-dimethyl-4-(1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 8.76 (s, 1H), 8.36 (d, J = 2.0 Hz, 1H), 7.65 (d, J = 2.5 Hz, 1H), 7.40-7.30 (m, 5H), 4.44 (q, J = 7.0 Hz, 1H), 2.29 (s, 3H), 2.10 (s, 3H), 2.06 (d, J = 7.0 Hz, 3H). ESI m/z 319 [M + H] ⁺ .	98.6
31	4-(1-benzyl-1H-imidazo[4,5-c]pyridin-6-yl)-3,5-dimethylisoxazole		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 9.00 (d, J = 1.0 Hz, 1H), 8.05 (s, 1H), 7.48 (d, J = 1.0 Hz, 1H), 7.40-7.30 (m, 5H), 5.58 (s, 2H), 2.40 (s, 3H), 2.25 (s, 3H); ESI m/z 305 [M + H] ⁺ .	98.6
32	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-c]pyridine 5-oxide		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 8.92 (s, 1H), 8.61 (s, 1H), 7.67 (s, 1H), 7.45-7.25 (m, 5H), 6.57 (s, 2H), 2.28 (s, 3H), 2.17 (s, 3H); ESI m/z 321 [M + H] ⁺ .	98.7





Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
33	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-c]pyridin-4-amine		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 8.21 (s, 1H), 7.42-7.25 (m, 5H), 6.70 (s, 1H), 5.46 (s, 2H), 2.39 (s, 3H), 2.24 (s, 3H); ESI m/z 320 [M + H] ⁺ .	96.9
34	4-(1-benzyl-3-bromo-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 8.33 (d, J = 1.5 Hz, 1H), 7.86 (s, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.34-7.24 (m, 5H), 5.48 (s, 2H), 2.35 (s, 3H), 2.17 (s, 3H); ESI MS m/z 382 [M + H] ⁺ .	>99
35	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridine-3-carbaldehyde		No general procedure	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.2 (s, 1H), 8.73 (s, 1H), 8.53 (d, J = 1.8 Hz, 1H), 8.11 (d, J = 1.8 Hz, 1H), 7.44-7.30 (m, 5H), 5.59 (s, 2H), 2.40 (s, 3H), 2.21 (s, 3H); ESI MS m/z 332 [M + H] ⁺ .	>99
36	1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)ethanone		No general procedure	¹ H NMR (300 MHz, CDCl ₃) δ 8.59 (d, J = 1.5 Hz, 1H), 8.22 (s, 1H), 7.45 (d, J = 1.8 Hz, 1H), 7.40-7.36 (m, 3H), 7.21-7.18 (m, 2H), 5.40 (s, 2H), 2.89 (s, 3H), 2.34 (s, 3H), 2.17 (s, 3H); ESI MS m/z 346 [M + H] ⁺ .	>99





Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
37	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-5-yl formate		No general procedure	¹ H NMR (300 MHz, CDCl ₃) δ 9.90 (s, 1H), 7.62 (s, 1H), 7.43-7.41 (m, 3H), 7.28 (s, 1H), 7.22-7.18 (m, 3H), 5.31 (s, 2H), 2.22 (s, 3H), 2.10 (s, 3H); ESI MS m/z 348 [M + H] ⁺ .	>99
38	4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benzamide		No general procedure	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.35 (d, J = 1.8 Hz, 1H), 7.99 (d, J = 2.1 Hz, 1H), 7.94 (br.s, 1H), 7.83 (d, J = 8.4 Hz, 2H), 7.37 (br.s, 1H), 7.27 (d, J = 8.4 Hz, 2H), 5.61 (s, 2H), 2.60 (s, 3H), 2.39 (s, 3H), 2.21 (s, 3H); ESI m/z 362 [M + H] ⁺ .	>99
39	4-(1-benzyl-3-nitro-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole		No general procedure	¹ H NMR (300 MHz, CDCl ₃) δ 8.74 (s, 1H), 8.47 (s, 1H), 7.56 (s, 1H), 7.45-7.42 (m, 3H), 7.27-7.26 (m, 2H), 5.47 (s, 2H), 2.35 (s, 3H), 2.17 (s, 3H); ESI MS m/z 349 [M + H] ⁺ .	>99
40	3,5-dimethyl-4-(3-(4-(trifluoromethyl)benzyl)-3H-imidazo[4,5-b]pyridin-6-yl)isoxazole		B	¹ H NMR (300 MHz, CDCl ₃) δ 8.33 (d, J = 2.1 Hz, 1H), 8.15 (s, 1H), 8.00 (d, J = 2.1 Hz, 1H), 7.64 (d, J = 8.1 Hz, 2H), 7.45 (d, J = 8.1 Hz, 2H), 5.58 (s, 2H), 2.44 (s, 3H), 2.30 (s, 3H); MM m/z 373 [M + H] ⁺	98.3

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
41	3,5-dimethyl-4-(1-(4-(trifluoromethyl)benzyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.49 (d, J = 2.1 Hz, 1H), 8.29 (s, 1H), 7.66 (d, J = 8.1 Hz, 2H), 7.34-7.30 (m, 3H), 5.50 (s, 2H), 2.33 (s, 3H), 2.16 (s, 3H); MM m/z 373 $[\text{M} + \text{H}]^+$	98.9
42	4-(3-(4-chlorobenzyl)-3H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.32 (d, J = 2.1 Hz, 1H), 8.11 (s, 1H), 7.98 (d, J = 2.11 Hz, 1H), 7.37-7.27 (m, 4H), 5.48 (s, 2H), 2.44 (s, 3H), 2.29 (s, 3H); MM m/z 339 $[\text{M} + \text{H}]^+$	>99
43	4-(1-(4-chlorobenzyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.47 (d, J = 2.1 Hz, 1H), 8.25 (s, 1H), 7.37 (d, J = 8.7 Hz, 2H), 7.32 (d, J = 2.1 Hz, 1H), 7.16 (d, J = 8.7 Hz, 2H), 5.39 (s, 2H), 2.35 (s, 3H), 2.18 (s, 3H); MM m/z 339 $[\text{M} + \text{H}]^+$	>99
44	4-(3-(4-fluorobenzyl)-3H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.33 (d, J = 2.1 Hz, 1H), 8.10 (s, 1H), 7.98 (d, J = 2.1 Hz, 1H), 7.38-7.33 (m, 2H), 7.09-7.03 (m, 2H), 5.48 (s, 2H), 2.44 (s, 3H), 2.30 (s, 3H); MM m/z 323 $[\text{M} + \text{H}]^+$	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
45	4-(1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		B	¹ H NMR (300 MHz, CDCl ₃) δ 8.47 (d, J = 2.1 Hz, 1H), 8.25 (s, 1H), 7.34 (d, J = 2.1 Hz, 1H), 7.24-7.19 (m, 2H), 7.09 (t, J = 8.7 Hz, 2H), 5.38 (s, 2H), 2.35 (s, 3H), 2.18 (s, 3H); MM m/z 323 [M + H] ⁺	98.4
46	3,5-dimethyl-4-(3-(pyridin-2-ylmethyl)-3H-imidazo[4,5-b]pyridin-6-yl)isoxazole		B	¹ H NMR (300 MHz, CDCl ₃) δ 8.62-8.59 (m, 1H), 8.33 (s, 1H), 8.31 (d, J = 2.1 Hz, 1H), 7.98 (d, J = 2.1 Hz, 1H), 7.71-7.65 (m, 1H), 7.33-7.23 (m, 2H), 5.63 (s, 2H), 2.43 (s, 3H), 2.29 (s, 3H); MM m/z 306 [M + H] ⁺	95.5




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
47	3,5-dimethyl-4-(1-(pyridin-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		B	^1H NMR (300 MHz, CDCl_3) δ 8.62-8.59 (m, 1H), 8.46 (d, J = 2.1 Hz, 1H), 8.34 (s, 1H), 7.72-7.66 (m, 1H), 7.59 (d, J = 2.1 Hz, 1H), 7.31-7.27 (m, 1H), 7.13 (d, J = 7.8 Hz, 1H), 5.51 (s, 2H), 2.38 (s, 3H), 2.22 (s, 3H); MM m/z 306 $[\text{M} + \text{H}]^+$	98.3
48	4-(1-(4-fluorobenzyl)-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole		A: using 6-bromo-1H-pyrrolo[3,2-b]pyridine as starting material	^1H NMR (300 MHz, CD_3OD) δ 8.26 (d, J = 1.8 Hz, 1H), 7.78 (dd, J = 0.9, 1.8 Hz, 1H), 7.75 (d, J = 3.3 Hz, 1H), 7.29-7.24 (m, 2H), 7.08-7.02 (m, 2H), 6.70 (dd, J = 0.6, 3.3 Hz, 1H), 5.47 (s, 2H), 2.36 (s, 3H), 2.19 (s, 3H); ESI MS m/z 322 $[\text{M} + \text{H}]^+$	97.6
49	4-(1-(4-fluorobenzyl)-1H-pyrrolo[2,3-b]pyridin-6-yl)-3,5-dimethylisoxazole		A: using 6-bromo-1H-pyrrolo[2,3-b]pyridine as starting material	^1H NMR (300 MHz, CD_3OD) δ 8.04 (d, J = 8.1 Hz, 1H), 7.46 (d, J = 3.6 Hz, 1H), 7.26-7.21 (m, 3H), 7.04-6.98 (m, 2H), 6.55 (d, J = 3.6 Hz, 1H), 5.50 (s, 2H), 2.53 (s, 3H), 2.37 (s, 3H); ESI MS m/z 322 $[\text{M} + \text{H}]^+$	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
50	4-(5-(4-fluorobenzyl)-5H-pyrrolo[2,3-b]pyrazin-3-yl)-3,5-dimethylisoxazole		A: using 3-bromo-5H-pyrrolo[2,3-b]pyrazine as starting material	^1H NMR (300 MHz, CD_3OD) δ 8.54 (s, 1H), 7.91 (d, J = 3.6 Hz, 1H), 7.35-7.30 (m, 2H), 7.08-7.02 (m, 2H), 6.72 (d, J = 3.6 Hz, 1H), 5.52 (s, 2H), 2.60 (s, 3H), 2.42 (s, 3H); ESI MS m/z 323 $[\text{M} + \text{H}]^+$.	>99
51	4-(1-(4-fluorobenzyl)-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole		A: using 6-bromo-1H-pyrazolo[4,3-b]pyridine as starting material	^1H NMR (300 MHz, CD_3OD) δ 8.50 (d, J = 1.8 Hz, 1H), 8.28 (d, J = 0.9 Hz, 1H), 8.05 (dd, J = 1.8, 1.2 Hz, 1H), 7.36-7.31 (m, 2H), 7.08-7.02 (m, 2H), 5.70 (s, 2H), 2.42 (s, 3H), 2.25 (s, 3H); ESI MS m/z 323 $[\text{M} + \text{H}]^+$.	98.5
52	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-pyrrolo[2,3-b]pyridin-4-amine		A: using 6-bromo-1H-pyrrolo[2,3-b]pyridin-4-amine as starting material	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.29-7.24 (m, 3H), 7.15-7.09 (m, 2H), 6.55 (d, J = 3.6 Hz, 1H), 6.35 (s, 1H), 6.33 (s, 2H), 5.33 (s, 2H), 2.49 (s, 3H), 2.32 (s, 3H); ESI MS m/z 337 $[\text{M} + \text{H}]^+$.	>99
53	4-(1-(4-fluorobenzyl)-3-methyl-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole		A: using 6-bromo-3-methyl-1H-pyrazolo[4,3-b]pyridine as starting material	^1H NMR (300 MHz, CD_3OD) δ 8.45 (d, J = 1.8 Hz, 1H), 7.98 (d, J = 1.8 Hz, 1H), 7.34-7.29 (m, 2H), 7.08-7.02 (m, 2H), 5.61 (s, 2H), 2.65 (s, 3H), 2.42 (s, 3H), 2.25 (s, 3H); ESI MS m/z 337 $[\text{M} + \text{H}]^+$.	96.7

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
54	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-indazol-4-amine		B: using 6-bromo-1H-indazol-4-amine as starting material	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.13 (d, J = 0.6 Hz, 1H), 7.32-7.23 (m, 5H), 6.70 (s, 1H), 6.11 (d, J = 1.2 Hz, 1H), 5.97 (s, 2H), 5.53 (s, 2H), 2.37 (s, 3H), 2.19 (s, 3H); ESI MS m/z 319 [M + H] ⁺ .	>99
55	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine		K	¹ H NMR (500 MHz, CDCl ₃) δ 7.34-7.28 (m, 3H), 7.09-7.08 (m, 2H), 6.42 (d, J = 1.5 Hz, 1H), 6.36 (d, J = 1.5 Hz, 1H), 5.28 (s, 2H), 4.42 (br.s, 2H), 2.60 (s, 3H), 2.31 (s, 3H), 2.17 (s, 3H); ESI m/z 333 [M + H] ⁺ .	99
56	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-5(4H)-one		N	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.15 (d, J = 1.5 Hz, 1H), 7.83 (d, J = 3.5 Hz, 1H), 7.64 (s, 1H), 7.34-7.32 (m, 5H), 6.75 (d, J = 2.5 Hz, 1H), 5.50 (s, 2H), 2.39 (s, 3H), 2.20 (s, 3H); ESI MS m/z 320 [M + H] ⁺ .	>99
57	3-((5-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)amino)benzo nitrile		M	¹ H NMR (300 MHz, DMSO-d ₆) δ 11.5 (s, 1H), 7.98 (s, 2H), 7.78 (s, 1H), 7.36-7.25 (m, 2H), 7.11-7.07 (m, 1H), 7.01-6.99 (m, 2H), 2.46 (s, 3H), 2.26 (s, 3H); ESI MS m/z 330 [M + H] ⁺ .	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
58	4-(1-(4-fluorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	^1H NMR (300 MHz, DMSO- d_6) δ 8.34 (d, J = 1.8 Hz, 1H), 7.99 (d, J = 2.1 Hz, 1H), 7.32-7.26 (m, 2H), 7.22-7.15 (m, 2H), 5.53 (s, 2H), 2.61 (s, 3H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 337 $[\text{M} + \text{H}]^+$.	98.9
58	4-(1-benzyl-2-ethoxy-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		No general procedure	^1H NMR (300 MHz, DMSO- d_6) δ 7.75 (d, J = 1.2 Hz, 1H), 7.38 - 7.22 (m, 5H), 7.18 (d, J = 1.5 Hz, 1H), 4.99 (s, 2H), 4.34 (q, J = 7.2 Hz, 2H), 2.37 (s, 3H), 2.18 (s, 3H), 1.42 (t, J = 7.2 Hz, 3H); ESI m/z 349 $[\text{M} + \text{H}]^+$.	>99
60	4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)-3,5-dimethylisoxazole		D	^1H NMR (300 MHz, DMSO- d_6) δ 8.35 (d, J = 1.8 Hz, 1H), 7.93 (d, J = 2.1 Hz, 1H), 5.37 (s, 2H), 2.56 (s, 3H), 2.41 (s, 3H), 2.33 (s, 3H), 2.23 (s, 3H), 1.91 (s, 3H); ESI m/z 338 $[\text{M} + \text{H}]^+$.	>99
61	4-(1-(2,4-dichlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	^1H NMR (300 MHz, DMSO- d_6) δ 8.36 (d, J = 2.1 Hz, 1H), 7.88 (d, J = 1.8 Hz, 1H), 7.74 (d, J = 2.1 Hz, 1H), 7.38 (dd, J = 8.4, 2.1 Hz, 1H), 6.77 (d, J = 8.4 Hz, 1H), 5.61 (s, 2H), 2.54 (s, 3H), 2.38 (s, 3H), 2.19 (s, 3H); ESI m/z 387 $[\text{M} + \text{H}]^+$.	>99





Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
62	4-(1-(4-methoxybenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	^1H NMR (300 MHz, DMSO- d_6) δ 8.33 (d, J = 1.8 Hz, 1H), 7.98 (d, J = 2.1 Hz, 1H), 7.21 (d, J = 8.7 Hz, 2H), 6.90 (d, J = 8.7 Hz, 2H), 5.46 (s, 2H), 3.71 (s, 3H), 2.61 (s, 3H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 349 $[\text{M} + \text{H}]^+$.	>99
63	4-(1-(cyclopropylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	^1H NMR (300 MHz, DMSO- d_6) δ 8.31 (d, J = 2.1 Hz, 1H), 8.05 (d, J = 2.1 Hz, 1H), 4.17 (d, J = 7.2 Hz, 2H), 2.65 (s, 3H), 2.44 (s, 3H), 2.26 (s, 3H), 1.31-1.18 (m, 1H), 0.54-0.48 (m, 2H), 0.46-0.41 (m, 2H); ESI m/z 283 $[\text{M} + \text{H}]^+$.	97.4
64	N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)acetamide		K	^1H NMR (300 MHz, CDCl_3) δ 8.59 (br.s, 1H), 8.20 (s, 1H), 7.38-7.31 (m, 3H), 7.09-7.06 (m, 2H), 6.76 (d, J = 1.2 Hz, 1H), 5.34 (s, 2H), 2.65 (s, 3H), 2.35 (s, 3H), 2.31 (s, 3H), 2.21 (s, 3H); ESI m/z 375 $[\text{M} + \text{H}]^+$.	97.4

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
65	N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-yl)ethanesulfonamide		K	^1H NMR (300 MHz, CDCl_3) δ 7.71 (br.s, 1H), 7.39-7.30 (m, 4H), 7.12-7.09 (m, 2H), 6.79 (d, J = 1.2 Hz, 1H), 5.33 (s, 2H), 3.21 (q, J = 7.5 Hz, 2H), 2.64 (s, 3H), 2.35 (s, 3H), 2.20 (s, 3H), 1.42 (t, J = 7.5 Hz, 3H); APCI m/z 425 $[\text{M} + \text{H}]^+$.	95.7
66	4-(1-benzyl-4-methoxy-2-methyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole		No general procedure	^1H NMR (300 MHz, CDCl_3) δ 7.35-7.30 (m, 3H), 7.09-7.06 (m, 2H), 6.64 (d, J = 1.2 Hz, 1H), 6.53 (s, 1H), 5.32 (s, 2H), 4.03 (s, 3H), 2.66 (s, 3H), 2.33 (s, 3H), 2.19 (s, 3H); ESI m/z 348 $[\text{M} + \text{H}]^+$.	93.7
67	7-amino-3-benzyl-5-(3,5-dimethylisoxazol-4-yl)benzo[d]oxazol-2(3H)-one		G	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.43-7.30 (m, 5H), 6.40 (d, J = 1.5 Hz, 1H), 6.39 (d, J = 1.5 Hz, 1H), 5.58 (s, 2H), 4.99 (s, 2H), 2.31 (s, 3H), 2.13 (s, 3H); ESI m/z 336 $[\text{M} + \text{H}]^+$.	97.6

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
68	3,5-dimethyl-4-(2-methyl-1-(pyridin-3-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		D	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.58 (d, J = 1.8 Hz, 1H), 8.51 (dd, J = 4.7, 1.8 Hz, 1H), 8.35 (d, J = 2.1 Hz, 1H), 8.03 (d, J = 2.1 Hz, 1H), 7.60 (dt, J = 8.1, 1.8 Hz, 1H), 7.37 (ddd, J = 7.8, 4.8, 0.6 Hz, 1H), 5.60 (s, 2H), 2.64 (s, 3H), 2.40 (s, 3H), 2.21 (s, 3H); ESI m/z 320 [M + H] ⁺	96.5
69	3,5-dimethyl-4-(2-methyl-1-(thiophen-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		D	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.34 (d, J = 2.1 Hz, 1H), 8.11 (d, J = 2.1 Hz, 1H), 7.48 (dd, J = 5.1, 1.2 Hz, 1H), 7.25 (dd, J = 3.1, 1.2 Hz, 1H), 7.00 (dd, J = 5.1, 3.3 Hz, 1H), 5.75 (s, 2H), 2.67 (s, 3H), 2.44 (s, 3H), 2.26 (s, 3H); ESI m/z 325 [M + H] ⁺	>99
70	4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benz onitrile		D	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.36 (d, J = 2.1 Hz, 1H), 7.98 (s, J = 2.1 Hz, 1H), 7.83 (d, J = 8.4 Hz, 2H), 7.36 (d, J = 8.4 Hz, 2H), 5.67 (s, 2H), 2.57 (s, 3H), 2.39 (s, 3H), 2.21 (s, 3H); ESI m/z 344 [M + H] ⁺	98.3





Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
71	4-(1-benzyl-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole		B: using 6-bromo-1H-pyrrolo[3,2-b]pyridine as starting material	^1H NMR (300 MHz, CDCl_3) δ 8.36 (d, $J = 1.8$ Hz, 1H), 7.54 (d, $J = 2.7$ Hz, 1H), 7.41 (s, 1H), 7.36-7.32 (m, 3H), 7.16-7.13 (m, 2H), 6.88 (s, 1H), 5.38 (s, 2H), 2.33 (s, 3H), 2.16 (s, 3H); ESI MS m/z 304 $[\text{M} + \text{H}]^+$.	>99
72	1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)-N,N-dimethylmethanamine		L	^1H NMR (300 MHz, CDCl_3) δ 8.34 (d, $J = 1.8$ Hz, 1H), 8.30 (s, 1H), 7.36-7.32 (m, 4H), 7.21-7.18 (m, 2H), 5.39 (s, 2H), 4.50 (s, 2H), 2.86 (s, 6H), 2.32 (s, 3H), 2.16 (s, 3H); ESI MS m/z 361 $[\text{M} + \text{H}]^+$.	98.3
73	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[2,3-b]pyridin-4-amine		B: using 6-bromo-1H-pyrrolo[2,3-b]pyridin-4-amine as starting material	^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.31-7.20 (m, 6H), 6.56 (d, $J = 3.6$ Hz, 1H), 6.35 (s, 1H), 6.32 (s, 2H), 5.35 (s, 2H), 2.49 (s, 3H), 2.32 (s, 3H); ESI MS m/z 319 $[\text{M} + \text{H}]^+$.	>99
74	3,5-dimethyl-4-(2-methyl-1-(pyridin-4-ylmethyl)-imidazo[4,5-b]pyridin-6-yl)isoxazole		D	^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.53 (dd, $J = 3.0, 1.5$ Hz, 2H), 8.36 (d, $J = 2.0$ Hz, 1H), 7.96 (d, $J = 2.0$ Hz, 1H), 7.12 (d, $J = 6.0$ Hz, 2H), 5.62 (s, 2H), 2.57 (s, 3H), 2.39 (s, 3H), 2.20 (s, 3H); ESI m/z 320 $[\text{M} + \text{H}]^+$.	98.9




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
75	1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazole-4-amine		F	¹ H NMR (500 MHz, CD ₃ OD) δ 6.70 (s, 1H), 6.44 (d, J = 1.0 Hz, 1H), 4.08 (d, J = 6.5 Hz, 2H), 2.61 (s, 3H), 2.40 (s, 3H), 2.25 (s, 3H), 1.30-1.19 (m, 1H), 0.62-0.53 (m, 2H), 0.45-0.40 (m, 2H). ESI m/z 297 [M + H] ⁺ .	>99
76	3,5-dimethyl-4-(2-methyl-1-((5-methylthiophen-2-yl)methyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		D	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.34 (d, J = 2.1 Hz, 1H), 8.09 (d, J = 2.1 Hz, 1H), 7.04 (d, J = 3.6 Hz, 1H), 6.66 (dd, J = 2.1, 1.2 Hz, 1H), 5.65 (s, 2H), 2.66 (s, 3H), 2.44 (s, 3H), 2.34 (d, J = 0.6 Hz, 3H), 2.27 (s, 3H); ESI m/z 339 [M + H] ⁺ .	98.1
77	4-(1-((5-chlorothiophen-2-yl)methyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.35 (d, J = 2.1 Hz, 1H), 8.12 (d, J = 2.1 Hz, 1H), 7.13 (d, J = 3.6 Hz, 1H), 7.02 (d, J = 3.6 Hz, 1H), 5.70 (s, 2H), 2.66 (s, 3H), 2.44 (s, 3H), 2.27 (s, 3H); ESI m/z 359 [M + H] ⁺ .	96.3

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
78	5-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)thiophene-2-carbonitrile		D	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.36 (d, J = 2.0 Hz, 1H), 8.11 (d, J = 2.0 Hz, 1H), 7.87 (d, J = 4.0 Hz, 1H), 7.31 (d, J = 4.0 Hz, 1H), 5.86 (s, 2H), 2.65 (s, 3H), 2.43 (s, 3H), 2.26 (s, 3H); ESI m/z 350 [M + H] ⁺ .	>99
79	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridine 4-oxide		N	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.28 (s, 1H), 8.05 (s, 1H), 7.83 (s, 1H), 7.49-7.45 (m, 2H), 7.13-7.07 (m, 2H), 6.00 (s, 2H), 2.48 (s, 3H), 2.32 (s, 3H); ESI MS m/z 339 [M + H] ⁺ .	>99
80	6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-5-yl acetate		N: using Example 59 as starting material	¹ H NMR (300 MHz, CDCl ₃) δ 8.34 (s, 1H), 8.07 (s, 1H), 7.43-7.38 (m, 2H), 7.12-7.06 (m, 2H), 5.46 (s, 2H), 2.31 (s, 3H), 2.19 (s, 3H), 2.16 (s, 3H); ESI MS m/z 381 [M + H] ⁺ .	>99
81	1-benzyl-6-(1,4-dimethyl-1H-pyrazol-5-yl)-2-methyl-4-nitro-1H-benzo[d]imidazole		F	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.04 (d, J = 1.5 Hz, 1H), 7.95 (d, J = 1.5 Hz, 1H), 7.37-7.29 (m, 4H), 7.23-7.21 (m, 2H), 5.6 (s, 2H), 3.69 (s, 3H), 2.68 (s, 3H), 1.93 (s, 3H); ESI m/z 362 [M + H] ⁺ .	99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
82	1-benzyl-6-(1,4-dimethyl-1H-pyrazol-5-yl)-2-methyl-1H-benzo[d]imidazole-4-amine		F	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.36-7.27 (m, 4H), 7.20-7.17 (m, 2H), 6.62 (d, J = 1.2 Hz, 1H), 6.30 (d, J = 1.2 Hz, 1H), 5.40 (s, 2H), 5.36 (s, 2H), 3.62 (s, 3H), 2.51 (s, 3H), 1.89 (s, 3H); ESI m/z 332 [M + H] ⁺ .	98.4
83	4-(1-(4-chlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.34 (d, J = 1.8 Hz, 1H), 7.98 (d, J = 1.8 Hz, 1H), 7.42 (d, J = 8.4 Hz, 2H), 7.24 (d, J = 8.4 Hz, 2H), 5.55 (s, 2H), 2.59 (s, 3H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 353 [M + H] ⁺ .	>99
84	4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)phenol		D	¹ H NMR (500 MHz, DMSO-d ₆) δ 9.45 (s, 1H), 8.31 (d, J = 2.0 Hz, 1H), 7.95 (d, J = 2.0 Hz, 1H), 7.09 (d, J = 8.5 Hz, 2H), 6.71 (d, J = 8.5 Hz, 2H), 5.39 (s, 2H), 2.61 (s, 3H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 335 [M + H] ⁺ .	>99
85	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazole-4-carbonitrile		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 7.63 (d, J = 1.5 Hz, 1H), 7.60 (d, J = 1.5 Hz, 1H), 7.38-7.27 (m, 3H), 7.19-7.14 (m, 2H), 5.57 (s, 2H), 2.69 (s, 3H), 2.32 (s, 3H), 2.16 (s, 3H); ESI m/z 343 [M + H] ⁺ .	>99





Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
86	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazole-4-carbonitrile		J: using 2-amino-5-bromobenzonitrile as starting material	^1H NMR (500 MHz, CD_3OD) δ 7.38-7.25 (m, 6H), 7.10 (d, J = 1.5 Hz, 1H), 5.13 (s, 2H), 2.27 (s, 3H), 2.09 (s, 3H); ESI m/z 345 $[\text{M} + \text{H}]^+$.	>99
87	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-morpholino-1H-benzo[d]imidazole-4-amine		I	^1H NMR (300 MHz, CDCl_3) δ 7.35-7.27 (m, 3H), 7.18-7.15 (m, 2H), 6.36 (s, 1H), 6.23 (d, J = 0.9 Hz, 1H), 5.22 (s, 2H), 4.29 (br.s, 2H), 3.83 (t, J = 4.5 Hz, 4H), 3.25 (br.s, 4H), 2.27 (s, 3H), 2.13 (s, 3H); ESI m/z 404 $[\text{M} + \text{H}]^+$.	>99
88	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridine-3-carbonitrile		No general procedure	^1H NMR (300 MHz, CDCl_3) δ 8.55 (s, 1H), 7.98 (s, 1H), 7.50 (s, 1H), 7.41-7.40 (m, 3H), 7.20-7.15 (m, 2H), 5.42 (s, 2H), 2.34 (s, 3H), 2.16 (s, 3H); ESI MS m/z 329 $[\text{M} + \text{H}]^+$.	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
89	4-(1-benzyl-3-chloro-1H-pyrrolo[3,2-b]pyridin-6-yl)-3,5-dimethylisoxazole		No general procedure	¹ H NMR (300 MHz, CDCl ₃) δ 8.49 (s, 1H), 7.55 (s, 1H), 7.50 (s, 1H), 7.38-7.36 (m, 3H), 7.18-7.16 (m, 2H), 5.36 (s, 2H), 2.34 (s, 3H), 2.16 (s, 3H); ESI MS m/z 338 [M + H] ⁺ .	>99
90	4-amino-1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one		E	¹ H NMR (500 MHz, CD ₃ OD) δ 7.36-7.28 (m, 4H), 6.40 (d, J = 1.4 Hz, 1H), 6.25 (d, J = 1.4 Hz, 1H), 5.03 (s, 2H), 2.28 (s, 3H), 2.12 (s, 3H); HPLC >99%, t _R = 13.4 min; ESI m/z 369 [M + H] ⁺ .	>99
91	1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one		E	¹ H NMR (500 MHz, CD ₃ OD) δ 7.80 (d, J = 1.4 Hz, 1H), 7.40-7.35 (m, 4H), 7.24 (d, J = 1.4 Hz, 1H), 5.15 (s, 2H), 2.32 (s, 3H), 2.15 (s, 3H); HPLC 98.7%, t _R = 16.5 min; ESI m/z 399 [M + H] ⁺ .	98.7
92	4-(1-benzyl-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole		A	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.55 (d, J = 1.8 Hz, 1H), 8.38 (d, J = 0.9 Hz, 1H), 8.27 (dd, J = 1.8 Hz, 1.0 Hz, 1H), 7.32-7.26 (m, 5H), 5.72 (s, 2H), 2.45 (s, 3H), 2.27 (s, 3H); ESI m/z 305 [M + H] ⁺ .	98.7




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
93	4-(1-(4-chlorobenzyl)-1H-pyrazolo[4,3-b]pyridin-6-yl)-3,5-dimethylisoxazole		A	¹ H NMR (500 MHz, CDCl ₃) δ 8.48 (d, J = 1.0 Hz, 1H), 8.34 (s, 1H), 7.41 (s, 1H), 7.33-7.30 (m, 2H), 7.19-7.16 (m, 2H), 5.60 (s, 2H), 2.38 (s, 3H), 2.22 (s, 3H); ESI m/z 374 [M + H] ⁺ .	98.8
94	1-benzyl-2-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-benzo[d]imidazol-4-amine		U	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.39 (d, J = 1.5 Hz, 1H), 7.33 (t, J = 7.0 Hz, 2H), 7.26 (t, J = 7.0 Hz, 1H), 7.16 (d, J = 7.0 Hz, 2H), 6.76 (d, J = 1.5 Hz, 1H), 6.44 (d, J = 1.5 Hz, 1H), 6.22 (d, J = 2.0 Hz, 1H), 5.41 (s, 2H), 5.36 (s, 2H), 3.76 (s, 3H), 3.31 (s, 3H); ESI m/z 318 [M + H] ⁺ .	>99
95	4-(1-(3,4-dichlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.35 (d, J = 2.0 Hz, 1H), 8.00 (d, J = 2.0 Hz, 1H), 7.61 - 7.59 (m, 2H), 7.13 (dd, J = 8.5, 2.0 Hz, 1H), 5.56 (s, 2H), 2.61 (s, 3H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 387 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
96	6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1H-benzo[d]imidazol-4-amine		K	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.39-7.28 (m, 5H), 6.24 (s, 1H), 6.15 (s, 1H), 5.86 (q, J = 6.9 Hz, 1H), 5.26 (s, 2H), 2.58 (s, 3H), 2.20 (s, 3H), 2.02 (s, 3H), 1.86 (d, J = 6.9 Hz, 3H); ESI m/z 347 [M + H] ⁺ .	>99
97	2-(azetidin-1-yl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.17 (m, 5H), 6.38 (d, J = 1.5 Hz, 1H), 6.27 (d, J = 1.5 Hz, 1H), 5.16 (s, 2H), 5.02 (s, 2H), 4.08 (t, J = 7.5 Hz, 4H), 2.34-2.24 (m, 5H), 2.12 (s, 3H); ESI m/z 374 [M + H] ⁺ .	98.8
98	3,5-dimethyl-4-(1-(thiophen-3-ylmethyl)-1H-pyrazolo[4,3-b]pyridin-6-yl)isoxazole		A	¹ H NMR (500 MHz, CDCl ₃) δ 8.48 (d, J = 1.7 Hz, 1H), 8.39 (s, 1H), 7.47 (s, 1H), 7.34-7.32 (m, 1H), 7.23-7.21 (m, 1H), 6.97 (dd, J = 5.0 Hz, 1.3 Hz, 1H), 5.67 (s, 2H), 2.39 (s, 3H), 2.23 (s, 3H); ESI m/z 311 [M + H] ⁺ .	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
99	N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-yl)acetamide		No general procedure	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.2 (s, 1H), 8.32 (d, J = 1.8 Hz, 1H), 8.23 (s, 1H), 7.97 (d, J = 1.8 Hz, 1H), 7.32-7.25 (m, 5H), 5.45 (s, 2H), 2.40 (s, 3H), 2.22 (s, 3H), 2.12 (s, 3H); ESI MS m/z 361 [M + H] ⁺ .	96.7
100	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-pyrrolo[3,2-b]pyridin-3-amine		No general procedure	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.18 (d, J = 1.8 Hz, 1H), 7.82 (d, J = 1.8 Hz, 1H), 7.33-7.21 (m, 5H), 7.06 (s, 1H), 5.30 (s, 2H), 4.26 (s, 2H), 2.37 (s, 3H), 2.21 (s, 3H); ESI MS m/z 319 [M + H] ⁺ .	84.2
101	1-(3,4-dichlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		R	¹ H NMR (500 MHz, DMSO-d ₆) δ 11.83 (s, 1H), 7.92 (d, J = 1.5 Hz, 1H), 7.73 (d, J = 2.0 Hz, 1H), 7.61 (d, J = 8.0 Hz, 1H), 7.53 (d, J = 2.0 Hz, 1H), 7.35 (dd, J = 8.5, 2.0 Hz, 1H), 5.05 (s, 2H), 2.37 (s, 3H), 2.19 (s, 3H); ESI m/z 389 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
102	1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-indazol-4-amine		C	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.14 (d, J = 0.8 Hz, 1H), 7.38-7.34 (m, 2H), 7.28-7.24 (m, 2H), 6.69 (s, 1H), 6.12 (d, J = 1.1 Hz, 1H), 5.94 (s, 2H), 5.53 (s, 2H), 2.37 (s, 3H), 2.20 (s, 3H); ESI m/z 353 [M + H] ⁺ .	>99
103	6-(3,5-dimethylisoxazol-4-yl)-1-(4-methoxybenzyl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one		E	¹ H NMR (500 MHz, CD ₃ OD) δ 7.78 (d, J = 1.5 Hz, 1H), 7.31 (d, J = 8.7 Hz, 2H), 7.23 (d, J = 1.5 Hz, 1H), 6.90 (d, J = 8.7 Hz, 2H), 5.09 (s, 2H), 3.75 (s, 3H), 2.32 (s, 3H), 2.14 (s, 3H); ESI m/z 395 [M + H] ⁺ .	>99
104	4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(4-methoxybenzyl)-1H-benzo[d]imidazol-2(3H)-one		E	¹ H NMR (500 MHz, CD ₃ OD) δ 7.26 (d, J = 8.6 Hz, 2H), 6.87 (d, J = 8.6 Hz, 2H), 6.39 (d, J = 1.4 Hz, 1H), 6.26 (d, J = 1.4 Hz, 1H), 4.97 (s, 2H), 3.74 (s, 3H), 2.28 (s, 3H), 2.12 (s, 3H); HPLC 93.0%, t _R = 12.2 min; ESI m/z 365 [M + H] ⁺ .	93.0
105	1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		R	¹ H NMR (500 MHz, DMSO-d ₆) δ 11.81 (s, 1H), 7.91 (d, J = 1.5 Hz, 1H), 7.45 (d, J = 2.0 Hz, 1H), 7.43-7.39 (m, 4H), 5.04 (s, 2H), 2.35 (s, 3H), 2.17 (s, 3H); ESI m/z 355 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
106	6-(3,5-dimethylisoxazol-4-yl)-1-(thiophen-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		R	¹ H NMR (500 MHz, DMSO-d ₆) δ 11.77 (s, 1H), 7.91 (d, J = 2.0 Hz, 1H), 7.57 (d, J = 2.0 Hz, 1H), 7.44 (dd, J = 5.0, 1.0 Hz, 1H), 7.26 (dd, J = 3.5, 1.0 Hz, 1H), 6.97 (dd, J = 5.0, 3.5 Hz, 1H), 5.24 (s, 2H), 2.39 (s, 3H), 2.21 (s, 3H); ESI m/z 327 [M + H] ⁺ .	98.6
107	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine		Q	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.95 (d, J = 1.5 Hz, 1H), 7.37-7.31 (m, 4H), 7.28-7.23 (m, 3H), 5.30 (s, 2H), 3.51-4.53 (m, 2H), 2.33 (s, 3H), 2.14 (s, 3H), 1.23 (t, J = 7.0 Hz, 3H); ESI m/z 348 [M + H] ⁺ .	>99
108	3,5-dimethyl-4-(2-methyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.27 (d, J = 2.0 Hz, 1H), 7.44 (d, J = 2.0 Hz, 1H), 7.40-7.36 (m, 4H), 7.33-7.30 (m, 1H), 6.01 (q, J = 7.0 Hz, 1H), 2.70 (s, 3H), 2.26 (s, 3H), 2.06 (s, 3H), 1.93 (d, J = 7.0 Hz, 3H); ESI m/z 333 [M + H] ⁺ .	97.7

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
109	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-(tetrahydro-2H-pyran-4-yl)-1H-benzo[d]imidazole-2,4-diamine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.21 (m, 5H), 6.48 (d, J = 7.8 Hz, 1H), 6.29 (d, J = 1.5 Hz, 1H), 6.21 (d, J = 1.5 Hz, 1H), 5.23 (s, 2H), 4.83 (s, 2H), 4.04-3.96 (m, 1H), 3.89 (dd, J = 11.4, 2.7 Hz, 2H), 3.42 (td, J = 11.4, 2.7 Hz, 2H), 2.28 (s, 3H), 2.11 (s, 3H); 1.98 (dd, J = 12.3, 2.7 Hz, 2H), 1.62-1.49 (m, 2H), ESI m/z 418 [M + H] ⁺ .	>99
110	6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazole-2(3H)-one		P	¹ H NMR (500 MHz, CD ₃ OD) δ 7.75 (d, J = 1.3 Hz, 1H), 7.44 (d, J = 7.7 Hz, 2H), 7.38 (t, J = 7.7 Hz, 2H), 7.31 (t, J = 7.7 Hz, 1H), 6.88 (d, J = 1.3 Hz, 1H), 5.88 (q, J = 7.1 Hz, 1H), 2.20 (s, 3H), 2.02 (s, 3H), 1.91 (d, J = 7.2 Hz, 3H); ESI m/z 377 [M - H] ⁺ .	>99
111	N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-4-yl)acetamide		O	¹ H NMR (300 MHz, DMSO-d ₆) δ 10.78 (s, 1H), 9.85 (s, 1H), 7.60-7.46 (m, 5H), 7.28 (d, J = 1.2 Hz, 1H), 7.06 (d, J = 1.2 Hz, 1H), 5.22 (s, 2H), 2.51 (s, 3H), 2.33 (s, 3H), 2.27 (s, 3H); ESI m/z 377 [M + H] ⁺ .	98.8




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
112	6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 11.78 (s, 1H), 7.87 (d, J = 2.0 Hz, 1H), 7.44 (d, J = 7.5 Hz, 2H), 7.36 (t, J = 7.5 Hz, 2H), 7.29 (t, J = 7.5 Hz, 1H), 7.09 (d, J = 2.0 Hz, 1H), 5.72 (q, J = 7.0 Hz, 1H), 2.26 (s, 3H), 2.06 (s, 3H), 1.84 (d, J = 7.0 Hz, 3H); ESI m/z 335 [M + H] ⁺ .	>99
113	6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2-amine		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.90 (d, J = 2.0 Hz, 1H), 7.40-7.28 (m, 6H), 6.81 (d, J = 2.0 Hz, 1H), 5.84 (q, J = 7.0 Hz, 1H), 3.54-3.48 (m, 2H), 2.20 (s, 3H), 1.99 (s, 3H), 1.83 (d, J = 7.0 Hz, 3H), 1.27 (t, J = 7.0 Hz, 3H); ESI m/z 362 [M + H] ⁺ .	>99
114	4-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine		Q	¹ H NMR (500 MHz, CDCl ₃) δ 8.24 (d, J = 2.0 Hz, 1H), 7.41-7.34 (m, 3H), 7.15 (d, J = 6.5 Hz, 2H), 7.06 (d, J = 1.0 Hz, 1H), 5.26 (s, 2H), 3.83 (t, J = 4.5 Hz, 4H), 3.50 (t, J = 4.5 Hz, 4H), 2.29 (s, 3H), 2.11 (s, 3H); ESI m/z 390 [M + H] ⁺ .	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
115	4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-benzo[d]imidazol-2(3H)-one		P	¹ H NMR (500 MHz, CD3OD) δ 7.42-7.32 (m, 4H), 7.26 (t, J = 6.9 Hz, 1H), 6.35 (s, 1H), 5.94 (s, 1H), 5.78 (q, J = 7.2 Hz, 1H), 2.17 (s, 3H), 2.00 (s, 3H), 1.86 (d, J = 7.2 Hz, 3H); ESI m/z 349 [M + H] ⁺ .	>99
116	4-(1-(cyclobutylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole		F	¹ H NMR (500 MHz, DMSO-d6) δ 8.09 (d, J = 1.5 Hz, 1H), 7.91 (d, J = 1.5 Hz, 1H), 4.37 (d, J = 7.0 Hz, 2H), 2.80-2.75 (m, 1H), 2.67 (s, 3H), 2.45 (s, 3H), 1.94 (s, 3H), 1.95-1.90 (m, 2H), 1.86-1.77 (m, 4H); ESI m/z 341 [M + H] ⁺ .	>99
117	4-(1-(cyclopentylmethyl)-2-methyl-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole		F	¹ H NMR (500 MHz, DMSO-d6) δ 8.06 (d, J = 1.5 Hz, 1H), 7.91 (d, J = 1.5 Hz, 1H), 4.29 (d, J = 7.5 Hz, 2H), 2.68 (s, 3H), 2.45 (s, 3H), 2.37 (m, 1H), 2.27 (s, 3H), 1.71-1.58 (m, 4H), 1.57-1.47 (m, 2H), 1.33-1.27 (m, 2H); ESI m/z 355 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
118	1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 7.90 (d, J = 1.5 Hz, 1H), 7.50 (d, J = 1.5 Hz, 1H), 3.81 (d, J = 7.0 Hz, 2H), 2.42 (s, 3H), 2.26 (s, 3H), 1.31-1.20 (m, 1H), 0.60-0.53 (m, 2H), 0.44-0.38 (m, 2H). ESI m/z 285 [M + H] ⁺ .	>99
119	N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-(ethylamino)-1H-benzo[d]imidazol-4-yl)acetamide		O	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.37 (s, 1H), 7.60 (s, 1H), 7.35-7.20 (m, 5H), 6.93 (t, J = 5.4 Hz, 1H), 6.80 (s, 1H), 5.29 (s, 2H), 3.57-3.48 (m, 2H), 2.31 (s, 3H), 2.15 (s, 3H), 2.13 (s, 3H), 1.23 (t, J = 7.2 Hz, 3H); ESI m/z 404 [M + H] ⁺ .	99.0
120	N-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-ethoxy-1H-benzo[d]imidazol-4-yl)acetamide		O	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.64 (s, 1H), 7.73 (s, 1H), 7.37-7.27 (m, 5H), 7.11 (s, 1H), 5.25 (s, 2H), 4.65 (q, J = 7.2 Hz, 2H), 2.35 (s, 3H), 2.18 (s, 3H), 2.16 (s, 3H), 1.43 (t, J = 7.2 Hz, 3H); ESI m/z 405 [M + H] ⁺ .	>99
121	4-(1-benzyl-4-bromo-2-methyl-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole		H	¹ H NMR (500 MHz, CD ₃ OD) δ 7.40-7.25 (m, 5H), 7.15 (d, J = 7.7 Hz, 2H), 5.51 (s, 2H), 2.64 (s, 3H), 2.32 (s, 3H), 2.15 (s, 3H); ESI m/z 396 [M + H] ⁺ .	>99




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
122	3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1-ethyl-1H-benzo[d]imidazol-2(3H)-one		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.37 (d, J = 7.5 Hz, 2H), 7.33 (t, J = 7.0 Hz, 2H), 7.29 (d, J = 8.0 Hz, 1H), 7.26 (t, J = 7.0 Hz, 1H), 7.09 (d, J = 1.5 Hz, 1H), 7.03 (dd, J = 8.0, 1.5 Hz, 1H), 5.08 (s, 2H), 3.94 (q, J = 7.0 Hz, 2H), 2.31 (s, 3H), 2.13 (s, 3H), 1.26 (t, J = 7.0 Hz, 3H); ESI m/z 348 [M + H] ⁺ .	94.6
123	4-(2-(azetidin-1-yl)-1-benzyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		Q	¹ H NMR (500 MHz, CDCl ₃) δ 8.07 (s, 1H), 7.43-7.37 (m, 3H), 7.13 (d, J = 6.5 Hz, 2H), 7.05 (s, 1H), 5.23 (s, 2H), 4.49 (t, J = 7.0 Hz, 4H), 2.54 (quin, J = 7.5 Hz, 2H), 2.30 (s, 3H), 2.10 (s, 3H); ESI m/z 360 [M + H] ⁺ .	>99
124	1-((5-chlorothiophen-2-yl)methyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one		R	¹ H NMR (500 MHz, DMSO-d ₆) δ 11.81 (s, 1H), 7.92 (d, J = 2.0 Hz, 1H), 7.63 (d, J = 1.5 Hz, 1H), 7.15 (d, J = 4.0 Hz, 1H), 6.99 (s, J = 4.0 Hz, 1H), 5.17 (s, 2H), 2.40 (s, 3H), 2.22 (s, 3H); ESI m/z 361 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
125	(S)-3,5-dimethyl-4-(2-methyl-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-6-yl)isoxazole		S	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.87 (d, J = 1.5 Hz, 1H), 7.42-7.30 (m, 6H), 6.11 (q, J = 7.2 Hz, 1H), 2.74 (s, 3H), 2.23 (s, 3H), 2.04 (s, 3H), 1.94 (d, J = 6.9 Hz, 3H); ESI MS m/z 377 [M + H] ⁺ .	>99
126	(R)-3,5-dimethyl-4-(2-methyl-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-6-yl)isoxazole		S	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.87 (d, J = 1.5 Hz, 1H), 7.42-7.30 (m, 6H), 6.11 (q, J = 7.2 Hz, 1H), 2.74 (s, 3H), 2.23 (s, 3H), 2.04 (s, 3H), 1.94 (d, J = 6.9 Hz, 3H); ESI MS m/z 377 [M + H] ⁺ .	98.3
127	6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-4-nitro-1-(1-phenylethyl)-1H-benzo[d]imidazol-2-amine		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 7.70 (d, J = 1.5 Hz, 1H), 7.45-7.30 (m, 5H), 6.72 (d, J = 1.5 Hz, 1H), 5.86 (q, J = 7.0 Hz, 1H), 3.72 (q, J = 7.2 Hz, 2H), 2.17 (s, 3H), 1.98 (s, 3H), 1.90 (d, J = 7.0 Hz, 3H), 1.36 (t, J = 7.2 Hz, 3H); ESI m/z 406 [M + H] ⁺ .	96.3




Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
128	4-(1-benzyl-2-ethyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		D	¹ H NMR (500 MHz, CDCl ₃) δ 8.42 (d, J = 1.7 Hz, 1H), 7.39-7.33 (m, 3H), 7.30 (d, J = 1.6 Hz, 1H), 7.10-7.09 (m, 2H), 5.41 (s, 2H), 3.08 (q, J = 7.5 Hz, 2H), 2.32 (s, 3H), 2.15 (s, 3H), 1.51 (t, J = 7.5 Hz, 3H); ESI m/z 333 [M + H] ⁺ .	>99
129	4-amino-6-(3,5-dimethylisoxazol-4-yl)-1-(4-hydroxybenzyl)-1H-benzo[d]imidazol-2(3H)-one		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 7.17 (d, J = 8.6 Hz, 2H), 6.72 (d, J = 8.6 Hz, 2H), 6.39 (d, J = 1.3 Hz, 1H), 6.26 (d, J = 1.3 Hz, 1H), 4.94 (s, 2H), 2.28 (s, 3H), 2.12 (s, 3H); ESI m/z 351 [M + H] ⁺ .	>99
130	N-(2-(azetidin-1-yl)-1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazol-4-yl)acetamide		O	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.69 (s, 1H), 7.36-7.16 (m, 6H), 6.92 (s, 1H), 5.26 (s, 2H), 4.18 (t, J = 7.5 Hz, 4H), 2.35-2.27 (m, 5H), 2.15 (s, 3H), 2.14 (s, 3H); ESI m/z 416 [M + H] ⁺ .	98.2

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
131	1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine		No general procedure	¹ H NMR (500 MHz, CDCl ₃) δ 7.93 (d, J = 2.0 Hz, 1H), 7.48 (d, J = 1.5 Hz, 1H), 3.98 (d, J = 6.5 Hz, 2H), 3.57 (q, J = 7.0 Hz, 2H), 2.42 (s, 3H), 2.26 (s, 3H), 1.30 (t, J = 7.0 Hz, 3H), 1.29-1.19 (m, 1H), 0.59-0.52 (m, 2H), 0.45-0.39 (m, 2H). ESI m/z 312 [M + H] ⁺ .	>99
132	1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine		F	¹ H NMR (500 MHz, CD ₃ OD) δ 6.70 (d, J = 1.5 Hz, 1H), 6.43 (d, J = 1.5 Hz, 1H), 4.18 (d, J = 7.0 Hz, 2H), 2.85-2.79 (m, 1H), 2.60 (s, 3H), 2.40 (s, 3H), 2.25 (s, 3H), 2.06-1.98 (m, 2H), 1.94-1.82 (m, 4H); ESI m/z 311 [M + H] ⁺ .	98.5
133	1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazol-4-amine		F	¹ H NMR (500 MHz, CD ₃ OD) δ 6.69 (d, J = 1.5 Hz, 1H), 6.44 (d, J = 1.5 Hz, 1H), 4.10 (d, J = 7.5 Hz, 2H), 2.61 (s, 3H), 2.50-2.40 (m, 1H), 2.40 (s, 3H), 2.25 (s, 3H), 1.80-1.65 (m, 4H), 1.64-1.55 (m, 2H), 1.42-1.28 (m, 2H); ESI m/z 325 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
134	6-(3,5-dimethylisoxazol-4-yl)-N2-ethyl-1-(1-phenylethyl)-1H-benzo[d]imidazole-2,4-diamine		No general procedure	¹ H NMR (500 MHz, CD3OD) δ 7.40-7.25 (m, 5H), 6.31 (d, J = 1.5 Hz, 1H), 5.92 (d, J = 1.5 Hz, 1H), 5.72 (q, J = 6.9 Hz, 1H), 3.53 (q, J = 7.2 Hz, 2H), 2.15 (s, 3H), 1.99 (s, 3H), 1.86 (d, J = 7.0 Hz, 3H), 1.33 (t, J = 7.2 Hz, 3H); ESI m/z 376 [M + H] ⁺ .	>99
135	4-(1-benzyl-4-nitro-2-(pyrrolidin-1-yl)-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole			¹ H NMR (300 MHz, DMSO-d ₆) δ 7.74 (d, J = 1.5 Hz, 1H), 7.55 (d, J = 1.5 Hz, 1H), 7.37-7.24 (m, 3H), 7.15-7.12 (m, 2H), 5.60 (s, 2H), 3.69 (t, J = 6.9 Hz, 4H), 2.34 (s, 3H), 2.16 (s, 3H), 1.92-1.88 (m, 4H); ESI m/z 418 [M + H] ⁺ .	96.8
136	4-(1-benzyl-2-(4-methylpiperazin-1-yl)-4-nitro-1H-benzo[d]imidazol-6-yl)-3,5-dimethylisoxazole			¹ H NMR (300 MHz, DMSO-d ₆) δ 7.82 (d, J = 1.5 Hz, 1H), 7.59 (d, J = 1.5 Hz, 1H), 7.37-7.28 (m, 3H), 7.22-7.19 (m, 2H), 5.45 (s, 2H), 3.40 (t, J = 4.8 Hz, 4H), 2.45 (t, J = 4.5 Hz, 4H), 2.33 (s, 3H), 2.21 (s, 3H), 2.13 (s, 3H); ESI m/z 447 [M + H] ⁺ .	98.5

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
137	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-methoxyethyl)-4-nitro-1H-benzo[d]imidazole-2-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.84 (t, J = 5.1 Hz, 1H), 7.67 (d, J = 1.5 Hz, 1H), 7.44 (d, J = 1.5 Hz, 1H), 7.36-7.25 (m, 5H), 5.41 (s, 2H), 3.73-3.67 (m, 2H), 3.61-3.57 (m, 2H), 3.27 (s, 3H), 2.33 (s, 3H), 2.15 (s, 3H); ESI m/z 422 [M + H] ⁺ .	97.4
138	4-(1-benzyl-2-cyclopropyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 8.29 (d, J = 2.1 Hz, 1H), 7.95 (d, J = 2.0 Hz, 1H), 7.37-7.33 (m, 2H), 7.30-7.28 (m, 3H), 5.67 (s, 2H), 2.38 (s, 3H), 2.37-2.35 (m, 1H), 2.20 (s, 3H), 1.13-1.11 (m, 4H); ESI m/z 345 [M + H] ⁺ .	>99
139	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N2-(2-methoxyethyl)-1H-benzo[d]imidazole-2,4-diamine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.33-7.20 (m, 5H), 6.76 (t, J = 5.1 Hz, 1H), 6.32 (d, J = 1.2 Hz, 1H), 6.21 (d, J = 1.5 Hz, 1H), 5.21 (s, 2H), 4.84 (s, 2H), 3.56 (s, 4H), 3.28 (s, 3H), 2.29 (s, 3H), 2.11 (s, 3H); ESI m/z 392 [M + H] ⁺ .	97.5

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
140	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-(pyrrolidin-1-yl)-1H-benzo[d]imidazole-4-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.24 (m, 3H), 7.18-7.15 (m, 2H), 6.35 (d, J = 1.5 Hz, 1H), 6.28 (d, J = 1.2 Hz, 1H), 5.42 (s, 2H), 4.98 (s, 2H), 3.47 (t, J = 6.9 Hz, 4H), 2.29 (s, 3H), 2.12 (s, 3H), 1.88-1.84 (m, 4H); ESI m/z 388 [M + H] ⁺ .	>99
141	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-2-(4-methylpiperazin-1-yl)-1H-benzo[d]imidazole-4-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.34-7.20 (m, 5H), 6.35 (d, J = 1.5 Hz, 1H), 6.29 (d, J = 1.2 Hz, 1H), 5.22 (s, 2H), 5.16 (s, 2H), 3.14 (t, J = 4.8 Hz, 4H), 2.50 (t, J = 4.5 Hz, 4H), 2.27 (s, 3H), 2.23 (s, 3H), 2.10 (s, 3H); ESI m/z 417 [M + H] ⁺ .	97.8
142	1-benzyl-N6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-benzo[d]imidazole-4,6-diamine		No general procedure	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.31 (t, J = 7.5 Hz, 2H), 7.25 (t, J = 7.5 Hz, 1H), 7.04 (d, J = 7.5 Hz, 2H), 6.69 (s, 1H), 5.73 (d, J = 2.0 Hz, 1H), 5.60 (d, J = 2.0 Hz, 1H), 5.18 (s, 2H), 5.05 (s, 2H), 2.38 (s, 3H), 2.13 (s, 3H), 1.92 (s, 3H); ESI m/z 348 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
143	(S)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1H-benzo[d]imidazol-4-amine		S	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.39-7.26 (m, 5H), 6.23 (d, J = 1.5 Hz, 1H), 6.14 (d, J = 1.2 Hz, 1H), 5.86 (q, J = 7.2 Hz, 1H), 5.26 (s, 2H), 2.58 (s, 3H), 2.20 (s, 3H), 2.02 (s, 3H), 1.86 (d, J = 6.9 Hz, 3H); ESI MS m/z 347 [M + H] ⁺ .	>99
144	(R)-6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1-(1-phenylethyl)-1H-benzo[d]imidazol-4-amine		S	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.39-7.26 (m, 5H), 6.23 (d, J = 1.5 Hz, 1H), 6.14 (d, J = 1.2 Hz, 1H), 5.86 (q, J = 7.2 Hz, 1H), 5.26 (s, 2H), 2.58 (s, 3H), 2.20 (s, 3H), 2.02 (s, 3H), 1.86 (d, J = 6.9 Hz, 3H); ESI MS m/z 347 [M + H] ⁺ .	>99
145	1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2(3H)-one		No general procedure	¹ H NMR (500 MHz, CD ₃ OD) δ 7.82 (d, J = 1.5 Hz, 1H), 7.52 (d, J = 1.0 Hz, 1H), 3.87 (d, J = 7.0 Hz, 2H), 2.45 (s, 3H), 2.29 (s, 3H), 1.30-1.18 (m, 1H), 0.60-0.52 (m, 2H), 0.47-0.43 (m, 2H). ESI m/z 329 [M + H] ⁺ .	>99

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
146	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine		Q	¹ H NMR (500 MHz, DMSO-d ₆) δ 7.96 (d, J = 2.0 Hz, 1H), 7.42 (d, J = 1.0 Hz, 1H), 7.40-7.36 (br s, 1H), 7.35-7.31 (m, 2H), 7.28-7.23 (m, 3H), 5.29 (s, 2H), 3.00 (d, J = 4.6 Hz, 3H), 2.34 (s, 3H), 2.15 (s, 3H); ESI m/z 334 [M + H] ⁺ .	>99
147	N,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-1H-benzo[d]imidazol-2-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.25 (t, J = 5.4 Hz, 1H), 7.69 (s, 1H), 7.50 (s, 1H), 7.39-7.22 (m, 10H), 5.44 (s, 2H), 4.77 (d, J = 5.7 Hz, 2H), 2.35 (s, 3H), 2.16 (s, 3H); ESI m/z 454 [M + H] ⁺ .	97.9
148	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-4-nitro-N-(pyridin-3-ylmethyl)-1H-benzo[d]imidazol-2-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.65 (d, J = 1.5 Hz, 1H), 8.47 (dd, J = 4.8, 1.5 Hz, 1H), 8.30 (t, J = 6.0 Hz, 1H), 7.81 (dt, J = 7.8, 1.8 Hz, 1H), 7.70 (d, J = 1.5 Hz, 1H), 7.51 (d, J = 1.5 Hz, 1H), 7.38-7.21 (m, 6H), 5.42 (s, 2H), 4.76 (d, J = 5.7 Hz, 2H), 2.34 (s, 3H), 2.16 (s, 3H); ESI m/z 455 [M + H] ⁺ .	98.5

Example Compound	Chemical Name	Structure	General procedure	Characterization	Purity HPLC (%)
149	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-4-nitro-1H-benzo[d]imidazole-2-amine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.68-7.66 (m, 2H), 7.45 (d, J = 1.5 Hz, 1H), 7.37-7.22 (m, 5H), 5.37 (s, 2H), 3.06 (d, J = 4.8 Hz, 3H), 2.34 (s, 3H), 2.16 (s, 3H); ESI m/z 378 [M + H] ⁺ .	>99
150	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-3-methyl-4-nitro-1H-benzo[d]imidazole-2(3H)-one		No general procedure	¹ H NMR (300 MHz, CDCl ₃) δ 7.48 (d, J = 1.5 Hz, 1H), 7.35-7.30 (m, 5H), 6.84 (d, J = 1.5 Hz, 1H), 5.15 (s, 2H), 3.65 (s, 3H), 2.26 (s, 3H), 2.09 (s, 3H); ESI m/z 379 [M + H] ⁺ .	>99
151	1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N ² -methyl-1H-benzo[d]imidazole-2,4-diamine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.33-7.20 (m, 5H), 6.63 (br.s, 1H), 6.32 (s, 1H), 6.23 (s, 1H), 5.17 (s, 2H), 4.86 (s, 2H), 2.94 (d, J = 4.5 Hz, 3H), 2.29 (s, 3H), 2.12 (s, 3H); ESI m/z 348 [M + H] ⁺ .	>99
152	N ² ,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-benzo[d]imidazole-2,4-diamine		I	¹ H NMR (300 MHz, DMSO-d ₆) δ 7.37-7.22 (m, 11H), 6.35 (s, 1H), 6.22 (s, 1H), 5.26 (s, 2H), 4.83 (s, 2H), 4.65 (d, J = 5.7 Hz, 2H), 2.29 (s, 3H), 2.12 (s, 3H); ESI m/z 424 [M + H] ⁺ .	>99

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 189

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 189



NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

What is claimed is:

1. A compound of Formula IIb' or Formula IIc':



or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein:


Rings **A** and **B** may be optionally substituted with groups independently selected from deuterium, -NH₂, amino, heterocycle(C₄-C₆), carbocycle(C₄-C₆), halogen, -CN, -OH, -CF₃, alkyl(C₁-C₆), -thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆);

X is selected from -NH-, -CH₂-, -CH₂CH₂-, -CH(CH₃)-, -CH₂CH₂CH₂-, -CH₂CH₂O-, -CH₂CH₂NH-, -CH₂CH₂S-, -C(O)-, -C(O)CH₂-, -C(O)CH₂CH₂-, -CH₂C(O)-, -CH₂CH₂C(O)-, -C(O)NH-, -C(O)O-, -C(O)S-, -C(O)NHCH₂-, -C(O)OCH₂-, and -C(O)SCH₂-, wherein one or more hydrogen(s) may independently be replaced with deuterium, hydroxy, methyl, halogen, -CF₃, or ketone, and where S may be oxidized to sulfoxide or sulfone;

R₄ is selected from cycloalkyl(C₃-C₆) and a phenyl ring, each optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with F, Cl, or Br; or alternatively


R₄ is a phenyl ring optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo; or alternatively

- R₄** is selected from 5- or 6-membered heterocycles containing 1 or 2 nitrogens, optionally substituted with groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo; and
- D₁** is selected from the following 5-membered monocyclic heterocycles:



which are optionally substituted with deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), -COOH, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

2. The compound according to claim 1, selected from Formula IIb”:



or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein:

Z is selected from hydrogen, deuterium, -NH₂, amino, alkyl(C₁-C₆), -thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆).


3. The compound according to claim 1 or 2, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **R₄** is selected from cycloalkyl(C₃-C₆) and a phenyl ring, each of which is optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN,

and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with F, Cl, or Br.


4. The compound according to claim 1 or 2, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **R₄** is a phenyl ring optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.
5. The compound according to claim 1 or 2, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **R₄** is selected from 5- or 6-membered heterocycles containing 1 or 2 nitrogens, optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone(C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.
6. The compound according to claim 5, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **R₄** is selected from pyrimidyl, pyridyl, isoxazole, and pyrazole, optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.
7. The compound according to any one of claims 1 to 6, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **D₁** is isoxazole optionally substituted with deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), -COOH,

and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo.

8. The compound according to any one of claims 1 to 6, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **D₁** is selected from




9. The compound according to any one of claims 1 to 8, or a stereoisomer, tautomer,




10. The compound according to any one of claims 1 to 9, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **X** is selected from -CH₂-, -CH(CH₃)-, -CH(OH)-, -NH-, and -CH₂CH₂-, where one or more hydrogen(s) may independently be replaced with deuterium or halogen.
11. The compound according to any one of claims 1 to 10, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **X** is selected from -CH₂-, -CH(CH₃)-, and -NH-, where one or more hydrogen(s) may independently be replaced with deuterium or halogen.
12. The compound according to any one of claims 1 to 11, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **X** is selected from -CH₂- and -CH(CH₃)-, where one or more hydrogen(s) may independently be replaced with deuterium or halogen.
13. The compound according to any one of claims 1 to 4 and 7 to 12, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **-X-R₄** is -CH₂-phenyl.

14. A compound according to claim 1, selected from Formula IIb”:



or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein:


- Z** is selected from hydrogen, deuterium, -NH₂, amino, alkyl(C₁-C₆), -thioalkyl(C₁-C₆), alkenyl(C₁-C₆), and alkoxy(C₁-C₆);



- X** is selected from -CH₂- and -CH(CH₃)-; and

- R₄** is a phenyl ring optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with F, Cl, and/or Br.


15. A compound of Formula IIb”:



or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein:


- X** is selected from -NH-, -CH₂-, -CH₂CH₂-, -CH(CH₃)-, -CH₂CH₂CH₂-, -CH₂CH₂O-, -CH₂CH₂NH-, -CH₂CH₂S-, -C(O)-, -C(O)CH₂-, -C(O)CH₂CH₂-, -CH₂C(O)-, -CH₂CH₂C(O)-, -C(O)NH-, -C(O)O-, -C(O)S-, -C(O)NHCH₂-, -C(O)OCH₂-, and -C(O)SCH₂-, wherein one or more hydrogen(s) may independently be replaced with deuterium, hydroxy, methyl, halogen, -CF₃, ketone, and where S may be oxidized to sulfoxide or sulfone;

- R₄** is selected from cycloalkyl(C₃-C₆) and a phenyl ring, each optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with F, Cl, or Br; or alternatively
- R₄** is a phenyl ring optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo; or alternatively
- R₄** is selected from 5- or 6-membered heterocycles containing 1 or 2 nitrogens, optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone(C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), carboxyl, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo; and
- D₁** is selected from the following 5-membered monocyclic heterocycles:




which are optionally substituted with deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), amino, halogen, amide, -CF₃, -CN, -N₃, ketone (C₁-C₄), -S(O)alkyl(C₁-C₄), -SO₂alkyl(C₁-C₄), -thioalkyl(C₁-C₄), -COOH, and/or ester, each of which may be optionally substituted with F, Cl, Br, -OH, -NH₂, -NHMe, -OMe, -SMe, oxo, and/or thio-oxo; and


- Z** is selected from:



16. The compound according to claim 15, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, wherein **Z** is selected from:



17. A compound of Formula IIIb':



or a pharmaceutically acceptable salt thereof,

wherein:


Ring **B** may be optionally substituted with one or more groups independently selected from deuterium and halogen;

X is selected from -CH₂- and -CH(CH₃)-, wherein one or more hydrogens may be independently replaced with deuterium or halogen;

Z is selected from -NH₂ and amino; and

R₄ is selected from 4-7 membered carbocycles and heterocycles optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with halogen.

18. The compound according to claim 17, or a pharmaceutically acceptable salt thereof, wherein **X** is -CH₂-, and one or more hydrogens may be independently replaced with deuterium or halogen.
19. The compound according to claim 17 or 18, or a pharmaceutically acceptable salt thereof, wherein **R₄** is selected from cyclopentyl optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with halogen.
20. The compound according to claim 17 or 18, or a pharmaceutically acceptable salt thereof, wherein **R₄** is selected from pyridyl optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with halogen.
21. The compound according to claim 17 or 18, or a pharmaceutically acceptable salt thereof, wherein **R₄** is selected from phenyl optionally substituted with one or more groups independently selected from deuterium, alkyl(C₁-C₄), alkoxy(C₁-C₄), halogen, -CF₃, -CN, and -thioalkyl(C₁-C₄), wherein each alkyl(C₁-C₄), alkoxy(C₁-C₄), and -thioalkyl(C₁-C₄) may be optionally substituted with halogen.
22. The compound according to any one of claims 17 to 21, or a pharmaceutically acceptable salt thereof, wherein **Z** is selected from H₂N-, MeNH-, EtNH-, PhCH₂NH-,



23. A compound selected from:

- 3-benzyl-5-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
- 4-(1-benzyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
- 4-(1-benzyl-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 4-(3-benzyl-3H-imidazo[4,5-b]pyridin-5-yl)-3,5-dimethylisoxazole;
- 3,5-dimethyl-4-(1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;
- 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benzamide;
- 3,5-dimethyl-4-(1-(4-(trifluoromethyl)benzyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;
- 4-(1-(4-chlorobenzyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 4-(1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 3,5-dimethyl-4-(1-(pyridin-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;
- 4-(1-(4-fluorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 4-(1-benzyl-2-ethoxy-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)-3,5-dimethylisoxazole;
- 4-(1-(2,4-dichlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 4-(1-(4-methoxybenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 4-(1-(cyclopropylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;
- 3,5-dimethyl-4-(2-methyl-1-(pyridin-3-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;
- 3,5-dimethyl-4-(2-methyl-1-(thiophen-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;
- 4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)benzonitrile;
- 3,5-dimethyl-4-(2-methyl-1-(pyridin-4-ylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;

3,5-dimethyl-4-(2-methyl-1-((5-methylthiophen-2-yl)methyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;

4-(1-((5-chlorothiophen-2-yl)methyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

5-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)thiophene-2-carbonitrile;

4-(1-(4-chlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

4-((6-(3,5-dimethylisoxazol-4-yl)-2-methyl-1H-imidazo[4,5-b]pyridin-1-yl)methyl)phenol;

4-(1-(3,4-dichlorobenzyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-(3,4-dichlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

6-(3,5-dimethylisoxazol-4-yl)-1-(thiophen-2-ylmethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine;

3,5-dimethyl-4-(2-methyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-6-yl)isoxazole;

6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

4-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine;

1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

4-(2-(azetidin-1-yl)-1-benzyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-((5-chlorothiophen-2-yl)methyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

4-(1-benzyl-2-ethyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-(cyclopropylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine;

4-(1-benzyl-2-cyclopropyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine;

N,1-dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-2-methyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridine;

4-(1-benzyl-2-(4-methylpiperazin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(2-methoxyethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-4-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

4-(1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine;

1-benzyl-6-(1-methyl-1H-pyrazol-5-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-7-methyl-1H-imidazo[4,5-b]pyridin-2(3H)-one;

4-(1-benzyl-2,7-dimethyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-3-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine;

4-(1-(cyclohexylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

4-(1-(cyclopentylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

4-(1-(cyclobutylmethyl)-2-methyl-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

4-(1-benzyl-2-(pyrrolidin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

2-((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)ethanol;

1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

1-benzyl-2-methyl-6-(1-methyl-1H-1,2,3-triazol-5-yl)-1H-imidazo[4,5-b]pyridine;

4-((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)cyclohexanol;

4-(1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine;

4-(2-(azetidin-1-yl)-1-(cyclopentylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

4-(1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)morpholine;

4-(2-(azetidin-1-yl)-1-(cyclobutylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

N1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)-N2,N2-dimethylethane-1,2-diamine;

4-(1-benzyl-2-(piperazin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-benzyl-N-cyclopentyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

(R)-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

(S)-6-(3,5-dimethylisoxazol-4-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

4-(1-benzyl-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole;

1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-(cyclobutylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

N1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)cyclohexane-1,4-diamine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(3-methoxypropyl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(oxetan-3-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

6-(3,5-dimethylisoxazol-4-yl)-1-(4-fluorobenzyl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine;

1-(4-chlorobenzyl)-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-N-cyclohexyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(1-methylpiperidin-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine;

1-((1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)amino)-2-methylpropan-2-ol; and

1-(1-benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-yl)azetidin-3-ol;

or stereoisomers, tautomers, pharmaceutically acceptable salts, or hydrates thereof.

24. 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-ethyl-1H-imidazo[4,5-b]pyridin-2-amine or a pharmaceutically acceptable salt thereof.

25. 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-methyl-1H-imidazo[4,5-b]pyridin-2-amine or a pharmaceutically acceptable salt thereof.

26. N,1-Dibenzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine or a pharmaceutically acceptable salt thereof.

27. 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-N-(pyridin-3-ylmethyl)-1H-imidazo[4,5-b]pyridin-2-amine or a pharmaceutically acceptable salt thereof.

28. 4-(1-Benzyl-2-(pyrrolidin-1-yl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole or a pharmaceutically acceptable salt thereof.

29. 4-(2-(Azetidin-1-yl)-1-(cyclopentylmethyl)-1H-imidazo[4,5-b]pyridin-6-yl)-3,5-dimethylisoxazole or a pharmaceutically acceptable salt thereof.

30. 1-Benzyl-6-(3,5-dimethylisoxazol-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine or a pharmaceutically acceptable salt thereof.

31. 1-(cyclopentylmethyl)-6-(3,5-dimethylisoxazol-4-yl)-N-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyridin-2-amine or a pharmaceutically acceptable salt thereof.

32. A pharmaceutical composition comprising the compound according to any one of claims 1 to 31 or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, and a pharmaceutically acceptable carrier.

33. Use of a compound according to any one of claims 1 to 31, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or hydrate thereof, or a pharmaceutical composition according to claim 32, for treating a disease or disorder selected from an autoimmune disease or disorder, an inflammatory disease or disorder, cancer, a benign proliferative

or fibrotic disorder, a cardiovascular disease or disorder, a metabolic disease or disorder, HIV, and a neurological disease or disorder.


34. The use according to claim 33, wherein the autoimmune disease or disorder is selected from Acute Disseminated Encephalomyelitis, Agammaglobulinemia, Allergic Disease, Ankylosing spondylitis, Anti-GBM/Anti-TBM nephritis, Anti-phospholipid syndrome, Autoimmune aplastic anemia, Autoimmune hepatitis, Autoimmune inner ear disease, Autoimmune myocarditis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune thrombocytopenic purpura, Behcet's Disease, Bullous pemphigoid, Castleman's Disease, Celiac Disease, Churg-Strauss syndrome, Crohn's Disease, Cogan's syndrome, Dry eye syndrome, Essential mixed cryoglobulinemia, Dermatomyositis, Devic's Disease, Encephalitis, Eosinophilic esophagitis, Eosinophilic fasciitis, Erythema nodosum, Giant cell arteritis, Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis (Wegener's), Graves' Disease, Guillain-Barre syndrome, Hashimoto's thyroiditis, Hemolytic anemia, Henoch-Schonlein purpura, IgA nephropathy, Inclusion body myositis, Type I diabetes, Interstitial cystitis, Kawasaki's Disease, Leukocytoclastic vasculitis, Lichen planus, Systemic lupus erythematosus (SLE), Microscopic polyangitis, Multiple sclerosis, Myasthenia gravis, myositis, Optic neuritis, Pemphigus, POEMS syndrome, Polyarteritis nodosa, Primary biliary cirrhosis, Psoriasis, Psoriatic arthritis, Pyoderma gangrenosum, Relapsing polychondritis, Rheumatoid arthritis, Sarcoidosis, Scleroderma, Sjogren's syndrome, Takayasu's arteritis, Transverse myelitis, Ulcerative colitis, Uveitis, and Vitiligo.
35. The use according to claim 33, wherein the inflammatory disease or disorder is selected from sinusitis, pneumonitis, osteomyelitis, gastritis, enteritis, gingivitis, appendicitis, irritable bowel syndrome, tissue graft rejection, chronic obstructive pulmonary disease (COPD), septic shock, toxic shock syndrome, SIRS, bacterial sepsis, osteoarthritis, acute gout, acute lung injury, acute renal failure, burns, Herxheimer reaction, and SIRS associated with viral infections.
36. The use according to claim 33, wherein the disease or disorder is cancer.

37. The use according to claim 33 or 36, wherein the cancer is selected from B-acute lymphocytic leukemia, Burkitt's lymphoma, diffuse large cell lymphoma, multiple myeloma, primary plasma cell leukemia, atypical carcinoid lung cancer, bladder cancer, breast cancer, cervix cancer, colon cancer, gastric cancer, glioblastoma, hepatocellular carcinoma, large cell neuroendocrine carcinoma, medulloblastoma, melanoma, neuroblastoma, esophageal squamous cell carcinoma, osteosarcoma, ovarian cancer, prostate cancer, renal clear cell carcinoma, retinoblastoma, rhabdomyosarcoma, small cell lung carcinoma, NUT midline carcinoma, non-small cell lung cancer, head and neck squamous cell carcinoma, chronic lymphocytic leukemia, follicular lymphoma, diffuse large B cell lymphoma with germinal center phenotype, Hodgkin's lymphoma, activated anaplastic large cell lymphoma, primary neuroectodermal tumor, pancreatic cancer, adenoid cystic carcinoma, T-cell prolymphocytic leukemia, malignant glioma, thyroid cancer, Barret's adenocarcinoma, pro-myelocytic leukemia, and mantle cell lymphoma.
38. The use according to claim 33 or 36, wherein the cancer exhibits overexpression, translocation, amplification, or rearrangement of a myc family oncoprotein.
39. The use according to claim 33 or 36, wherein the cancer results from aberrant regulation of BET proteins.
40. The use according to claim 33 or 36, wherein the cancer relies on pTEFb (Cdk9/cyclin T) and BET proteins to regulate oncogenes.
41. The use according to claim 33 or 36, wherein the cancer is associated with upregulation of BET responsive genes CDK6, Bcl2, TYRO3, MYB and/or hTERT.
42. The use according to claim 33 or 36, wherein the cancer is associated with a viral infection.
43. The use according to claim 38, wherein the cancer exhibiting overexpression, translocation, amplification, or rearrangement of a myc family oncoprotein is selected from B-acute lymphocytic leukemia, Burkitt's lymphoma, Diffuse large cell lymphoma, Multiple myeloma, Primary plasma cell leukemia, Atypical carcinoid lung cancer,


Bladder cancer, Breast cancer, Cervix cancer, Colon cancer, Gastric cancer, Glioblastoma, Hepatocellular carcinoma, Large cell neuroendocrine carcinoma, Medulloblastoma, Nodular melanoma, Superficial spreading melanoma, Neuroblastoma, Esophageal squamous cell carcinoma, Osteosarcoma, Ovarian cancer, Prostate cancer, Renal clear cell carcinoma, Retinoblastoma, Rhabdomyosarcoma, and Small cell lung carcinoma.

44. The use according to claim 39, wherein the cancer resulting from aberrant regulation of BET proteins is selected from NUT midline carcinoma, B-cell lymphoma, non-small cell lung cancer, esophageal cancer, head and neck squamous cell carcinoma, and colon cancer.
45. The use according to claim 40, wherein the cancer relying on pTEFb (Cdk9/cyclin T) and BET proteins to regulate oncogenes is selected from chronic lymphocytic leukemia, multiple myeloma, follicular lymphoma, diffuse large B cell lymphoma with germinal center phenotype, Burkitt's lymphoma, Hodgkin's lymphoma, anaplastic large cell lymphoma, neuroblastoma, primary neuroectodermal tumor, rhabdomyosarcoma, prostate cancer, and breast cancer.
46. The use according to claim 41, wherein the cancer associated with upregulation of BET responsive genes CDK6, Bcl2, TYRO3, MYB, and/or hTERT is selected from pancreatic cancer, breast cancer, colon cancer, glioblastoma, adenoid cystic carcinoma, T-cell prolymphocytic leukemia, malignant glioma, bladder cancer, medulloblastoma, thyroid cancer, melanoma, multiple myeloma, Barret's adenocarcinoma, hepatoma, prostate cancer, pro-myelocytic leukemia, chronic lymphocytic leukemia, mantle cell lymphoma, diffuse large B-cell lymphoma, small cell lung cancer, and renal carcinoma.
47. The use according to claim 42, wherein the viral infection is selected from Epstein-Barr Virus, hepatitis B virus, hepatitis C virus, Kaposi's sarcoma associated herpesvirus, human papilloma virus, Merkel cell polyomavirus, and human cytomegalovirus.
48. The use according to claim 36, wherein the compound is for administration in combination with another anticancer agent.

49. The use according to claim 48, wherein the anticancer agent is selected from ABT-737, Azacitidine (Vidaza[™]), AZD1152 (Barasertib), AZD2281 (Olaparib), AZD6244 (Selumetinib), BEZ235, Bleomycin Sulfate, Bortezomib (Velcade[™]), Busulfan (Myleran[™]), Camptothecin, Cisplatin, Cyclophosphamide (Clafen[™]), CYT387, Cytarabine (Ara-C), Dacarbazine, DAPT (GSI-IX), Decitabine, Dexamethasone, Doxorubicin (Adriamycin[™]), Etoposide, Everolimus (RAD001), Flavopiridol (Alvocidib), Ganetespib (STA-9090), Gefitinib (Iressa[™]), Idarubicin, Ifosfamide (Mitoxana), IFNa2a (Roferon A[™]), Melphalan (Alkeran[™]), Methazolastone (temozolomide), Metformin, Mitoxantrone (Novantrone[™]), Paclitaxel, Phenformin, PKC412 (Midostaurin), PLX4032 (Vemurafenib), Pomalidomide (CC-4047), Prednisone (Deltasone[™]), Rapamycin, Revlimid[™] (Lenalidomide), Ruxolitinib (INCB018424), Sorafenib (Nexavar[™]), SU11248 (Sunitinib), SU11274, Vinblastine, Vincristine (Oncovin[™]), Vinorelbine (Navelbine[™]), Vorinostat (SAHA), and WP1130 (Degrasyn).



IIb'



IIc'