



US 20150246969A1

(19) **United States**

(12) **Patent Application Publication**
Hartle et al.

(10) **Pub. No.: US 2015/0246969 A1**

(43) **Pub. Date: Sep. 3, 2015**

(54) **TREATMENT FOR RHEUMATOID
ARTHRITIS**

(30) **Foreign Application Priority Data**

Sep. 20, 2012 (EP) 12185235.4

(71) Applicant: **MORPHOSYS AG**,
Martinsried/Planegg (DE)

(72) Inventors: **Stefan Hartle**, Martinsried/Planegg
(DE); **Stephane Leclair**,
Martinsried/Planegg (DE); **Amgad**
Shebl, Martinsried/Planegg (DE); **Stefan**
Steidl, Martinsried/Planegg (DE)

(73) Assignee: **MORPHOSYS AG**,
Martinsried/Planegg (DE)

(21) Appl. No.: **14/429,996**

(22) PCT Filed: **Sep. 19, 2013**

(86) PCT No.: **PCT/EP2013/069501**

§ 371 (c)(1),

(2) Date: **Mar. 20, 2015**

Related U.S. Application Data

(60) Provisional application No. 61/703,871, filed on Sep.
21, 2012.

Publication Classification

(51) **Int. Cl.**

C07K 16/24 (2006.01)

A61K 31/519 (2006.01)

A61K 39/395 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 16/243** (2013.01); **A61K 39/3955**

(2013.01); **A61K 31/519** (2013.01); **C07K**

2317/76 (2013.01); **C07K 2317/21** (2013.01)

(57) **ABSTRACT**

The present invention provides anti-GM-CSF antibodies for use in the treatment of rheumatoid arthritis. Anti-GM-CSF antibodies, in particular MOR103, are administered to patients suffering from rheumatoid arthritis at dosages that are beneficial in a clinical setting.

Figure 1

MOR103:

Variable Heavy Chain Peptide (CDRs are bold and underlined):

QVQLVESGGGLVQPGGSLRLSCAAS**GFTFSSYWMN**WVRQAPGKGLEWVS**GIENKYAGG**
ATYYAASVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR**GFGTDF**WGQGLTVTVSS

Variable Heavy Chain DNA:

CAGGTGCAGCTGGTTCGAGTCTGGCGGCGGACTGGTGCAGCCTGGCGGCAGCCTGAG
ACTGAGCTGTGCCGCCAGCGGCTTCACCTTCAGCAGCTACTGGATGAACTGGGTGAGG
CAGGCCCCCTGGCAAGGGCCTGGAGTGGGTGTCCGGCATCGAGAACAAAGTATGCCGGC
GGAGCCACCTACTACGCCGCCAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACAAC
AGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGAGGGCCGAGGACACCGCCGTG
TACTACTGTGCCAGGGGCTTCGGCACCGATTTCTGGGGCCAGGGCACCCCTGGTGACA
GTCAGCTCA

Variable Light Chain Peptide (CDRs are bold and underlined):

DIELTQPPSVSVAPGQTARISC**SGDSIGKKYAY**WYQQKPGQAPVLVIY**KKRPS**GIPERFSG
SNSGNTATLTISGTQAEDEADYYC**SAWGDKGM**VFGGGTKLTVLGQ

Variable Heavy Chain DNA:

GACATCGAGCTGACCCAGCCCCCAGCGTGTCTGTGGCCCCTGGCCAGACCGCCCGG
ATCAGCTGCTCCGGCGACAGCATCGGCAAGAAGTACGCCTACTGGTATCAGCAGAAGC
CCGGCCAGGCCCCCGTGCTGGTGATCTACAAGAAGCGGCCAGCGGCATCCCCGAGC
GGTTCAGCGGCAGCAACAGCGGCAACACCGCCACCCTGACCATCAGCGGCACCCAGG
CCGAGGACGAGGCCGACTACTACTGCTCCGCCTGGGGCGACAAGGGCATGGTGTGTTG
GCGGCGGAACAAAGTTAACCGTGCTGGGGCAG

Figure 2




Figure 3





Figure 4

TREATMENT FOR RHEUMATOID ARTHRITIS

FIELD OF THE INVENTION

[0001] The present invention provides anti-GM-CSF antibodies for use in the treatment of rheumatoid arthritis, and methods for the treatment of rheumatoid arthritis using such antibodies. Anti-GM-CSF antibodies, in particular MOR103, are administered to patients suffering from rheumatoid arthritis at dosages that are beneficial in a clinical setting.

BACKGROUND OF THE INVENTION

[0002] Rheumatoid arthritis (RA) is a chronic systemic inflammatory disease that affects 0.5% to 1% of the adult population worldwide. RA primarily affects the joints and is characterized by chronic inflammation of the synovial tissue, which eventually leads to the destruction of cartilage, bone and ligaments and can cause joint deformity. RA has a peak incidence between 40 and 60 years of age and affects primarily women. The cause of RA is not known; however, certain histocompatibility antigens are associated with poorer outcomes. Nonsteroidal anti-inflammatory drugs (NSAIDs) provide only symptomatic relief. Disease-modifying antirheumatic drugs (DMARDs), the cornerstone of RA treatment throughout all stages of the disease, maintain or improve physical function and retard radiographic joint damage. Pro-inflammatory cytokines, such as tumor necrosis factor- α (TNF α), interleukin-1, interleukin-6 and granulocyte macrophage colony stimulating factor (GM-CSF), which lead to the activation and proliferation of immune cells, are found to be increased in the inflamed joint.

[0003] More recently, biological compounds, such as antibodies, that target tumor necrosis factor α (TNF α), B-cells, or T-cells have been used to treat RA, but still many patients fail to respond to these therapies. Colony-stimulating factors (CSFs) have been suggested for a potential point of intervention for inflammatory disorders, such as RA (reviewed e.g. in Nat Rev Immunol (2008) 8, 533-44) or Nat Rev Rheumatol (2009) 5, 554-9). One of such CSF is granulocyte-macrophage colony-stimulation factor (GM-CSF).

[0004] MOR103 is a fully human anti-GM-CSF antibody (Mol Immunol (2008) 46, 135-44; WO 2006/122797). MOR 103 is also in a clinical Phase Ib trial for multiple sclerosis. The present invention describes the development of a clinically efficacious treatment regimen comprising MOR103 for RA.

SUMMARY OF THE INVENTION

[0005] In one aspect, the present invention provides an anti-GM-CSF antibody for use in the treatment of a patient suffering from rheumatoid arthritis, wherein said antibody is administered to said patient in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0006] In another aspect, the present invention also provides a method to treat a patient suffering from rheumatoid arthritis, said method comprising administering to said patient an anti-GM-CSF antibody in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous admin-

istration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0007] In an embodiment, the anti-GM-CSF antibody is administered intravenously, optionally at a dosage of at least 1.0 mg/kg, or at a dose of about 1.0 mg/kg or about 1.5 mg/kg. In an embodiment, the anti-GM-CSF antibody is administered weekly, over at least four weeks.

[0008] In an embodiment, the anti-GM-CSF antibody is administered subcutaneously, optionally at a dose of at least 2.0 mg/kg, or at a dose of about 2.0 mg/kg, about 3.0 mg/kg or about 4.0 mg/kg. In an embodiment, the anti-GM-CSF antibody is administered biweekly, monthly or bimonthly. In another embodiment, the antibody is administered at a fixed dose of about 75 mg, of about 100 mg, of about 150 mg, of about 200 mg, of about 300 mg or of about 400 mg. Administration of fixed doses may be every week, every second week, every third week, every fourth week or every sixth week.

[0009] In an embodiment, the dosage of anti-GM-CSF antibody administered to said patient and frequency of said administration is sufficient to provide and maintain a serum concentration of said antibody at at least 2 μ g/ml in said patient over the duration of said treatment.

[0010] In another aspect, the present invention provides an anti-GM-CSF antibody, wherein said anti-GM-CSF antibody is an antibody comprising an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIENKYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGDKGM (SEQ ID NO.: 7) for use in the treatment of a patient suffering from rheumatoid arthritis, wherein said antibody is administered to said patient in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0011] In another aspect, the present invention provides an anti-GM-CSF antibody, wherein said anti-GM-CSF antibody is an antibody comprising an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIENKYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGDKGM (SEQ ID NO.: 7) for use in the treatment of a patient suffering from rheumatoid arthritis, wherein said antibody is administered intravenously at a dose of about 1.0 mg/kg or at a dose of about 1.5 mg/kg and wherein said antibody is administered weekly over at least four weeks.

[0012] In another aspect, the present invention provides an anti-GM-CSF antibody for use in the treatment of a patient suffering from rheumatoid arthritis, wherein said antibody is administered to said patient in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg or at least 1.5 mg/kg when administered weekly over at least four weeks, and wherein said anti-GM-CSF antibody is administered in combination with a DMARD, such as methotrexate.

[0013] In an embodiment, the administration of said antibody to achieve such a therapeutically effective amount comprises the administration of said antibody intravenously at a dose at least 0.6, at least 0.7, at least 0.8, at least 0.9 or at least 1.0 mg/kg. In other embodiments, the antibody of the present invention is administered intravenously at a dose of about 1.0 mg/kg or a dose of about 1.5 mg/kg. Administration may be monthly, biweekly (every two weeks) or weekly.

[0014] In another aspect, the present invention provides an anti-GM-CSF antibody for use in the treatment of a patient suffering from rheumatoid arthritis, wherein said antibody is administered to said patient subcutaneously in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg or at least 1.5 mg/kg when administered weekly over at least four weeks, and wherein said anti-GM-CSF antibody is administered in combination with a DMARD, such as methotrexate.

[0015] In an embodiment, the administration of said antibody to achieve such a therapeutically effective amount comprises the administration of said antibody subcutaneously at a dose of at least 1.0, at least 1.5, at least 2.0, at least 2.5, at least 3.0, at least 3.5 or at least 4.0 mg/kg. In other embodiments, the antibody of the present invention is administered subcutaneously at a dose of about 2.0 mg/kg, a dose of about 3.0 mg/kg or a dose of about 4.0 mg/kg. Administration may be monthly, biweekly (every two weeks) or weekly.

[0016] In an embodiment, the administration of said antibody to achieve such a therapeutically effective amount comprises the administration of said antibody subcutaneously at a fixed dose of about 40 mg, at a fixed dose of 75 mg, at a fixed dose of 100 mg, at a fixed dose of 140 mg, at a fixed dose of 150 mg, at a fixed dose of 180 mg, at a fixed dose of 200 mg, at a fixed dose of 280 mg, at a fixed dose of 300 mg or at a fixed dose of 400 mg. Administration of fixed doses may be every week, every second week, every third week, every fourth week or every sixth week.

[0017] In another aspect, the present invention provides a method of treating a patient suffering from rheumatoid arthritis, said method comprising administering to said patient an anti-GM-CSF antibody subcutaneously at

[0018] (i) a dose of at least 1.0 mg/kg, or

[0019] (ii) a fixed dose of between 40 mg and 400 mg.

The anti-GM-CSF antibody may be administered to said patient in a manner to achieve to a serum concentration of said antibody at at least 2 µg/ml in said patient over the duration of said treatment. The antibody may be administered to said patient in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0020] In another aspect, the present invention provides an anti-GM-CSF antibody for inhibiting progression of structural joint damage in a rheumatoid arthritis patient comprising administering to said patient said antibody in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] FIG. 1 shows the amino acid sequence and the DNA sequence of MOR04357.

[0022] FIG. 2 shows the mean changes of the DAS28 score after four weeks (left panel) and after eight weeks (right panel) of treatment compared. DAS28 score changes are compared to baseline levels, i.e. disease status prior to treatment.

[0023] FIG. 3 shows the average ACR20 score of all treatment arms after four weeks. An increase of the ACR20 scores corresponds to an improvement of the severity of disease.

[0024] FIG. 4 shows the average ACR20 score of all treatment arms after eight weeks. An increase of the ACR20 scores corresponds to an improvement of the severity of disease.

DESCRIPTION

[0025] The terms “GM-CSF” and “GMCSF” refer to the protein known as GM-CSF or Granulocyte-macrophage colony-stimulating factor, having the following synonyms: Colony-stimulating factor 2, CSF2, GMCSF, GM-CSF, Granulocyte-macrophage colony-stimulating factor, MGC131935, MGC138897, Molgramostin, Sargramostim. Human GM-CSF has the amino acid sequence of (UniProt P04141):

(SEQ ID NO.: 1)
MWLQSLLLLTGTVACISAPARSPSPSTQPEWHVNAIQEARLLNLSRD
TAAEMNETVEVISEMFDLQEPCLQTRLELYKQGRLSLTKLKGPLTM
MASHYKQHCPTPETSCATQIITFESFKENLKDFFLVIPFDCWEPVQE

[0026] “MOR103” is an anti-GM-CSF antibody whose amino acid sequence and DNA sequence is provided in FIG. 1. “MOR103” and “MOR04357” and “MOR4357” are used as synonyms to describe the antibody shown in FIG. 1. MOR04357 comprises an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIENKYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGDKGM (SEQ ID NO.: 7). MOR04357 comprises a variable heavy chain of the sequence QVQLVESGGGLVQPGGSLRLSCAASG-FTFSSYWMNWRQAPGKGLEWVSGIENKYAGGATYYAASVKGRTISRDNSKNT-LYLQMNSLAEDTAVYYCARGFGTDF-WGQGTLLVTVSS (SEQ ID NO.: 8) and a variable light chain of the sequence DIETQPPSVSVAPGQTARISCSGDSIGKKYAYWYQKPGQAPVLVIYKKRPSGIPERFSGNSGNTALTISGTQAEDEADYYCSAWGDKGMVFGGGTKLTVLGQ (SEQ ID NO.: 9).

[0027] In certain embodiments, the antibody used in the present invention is an antibody specific for GM-CSF. In other embodiments, the antibody used in the present invention is an antibody specific for a polypeptide encoding an amino acid sequence comprising SEQ ID NO.: 1.

[0028] As used herein, “specifically for” or “specifically binding to” refers to an antibody selectively or preferentially binding to GM-CSF. Preferably the binding affinity for antigen is of K_d value of 10⁻⁹ mol/l or lower (e.g. 10⁻¹⁰ mol/l), preferably with a K_d value of 10⁻¹⁰ mol/l or lower (e.g. 10⁻¹²).

mol/l). The binding affinity is determined with a standard binding assay, such as surface plasmon resonance technique (BIAcore®).

[0029] In certain embodiments, the antibody used in the present invention is MOR103. In other embodiments, the antibody used in the present invention is an antibody comprising an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIENKYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGDKGM (SEQ ID NO.: 7). In other embodiments, the antibody used in the present invention is an antibody comprising a variable heavy chain of the sequence QVQLVESGGGLVQPQGLRLSCAASGFTFSSYWMNWVRQAPGKGLEWVSGIENKYAGGATYYAASVKGRTISRDNKNTLYLQMNSLR AEDTAVYYCARGFGTDFWGQGTTLTVSS (SEQ ID NO.: 8) and a variable light chain of the sequence DIELTQPPSVSVAPGQTARISCSGDSIGKKYAYWYQQKPGQAPVLVIYKKRPSGIPERFSGNSGNTAITLTISGTQAEDADYYCSAWGDKGMVFGGGLTKLTVLGQ (SEQ ID NO.: 9). In other embodiments, the antibody used in the present invention is an antibody which cross-competes with an antibody comprising an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIENKYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGDKGM (SEQ ID NO.: 7). In other embodiments, the antibody used in the present invention is an antibody which binds to the same epitope like an antibody specific for GM-CSF comprising an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIENKYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGDKGM (SEQ ID NO.: 7).

[0030] The term “antibody” is used in the broadest sense and specifically covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies) formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

[0031] “Antibody fragments” herein comprise a portion of an intact antibody which retains the ability to bind antigen. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[0032] The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variants that may arise during production of the monoclonal antibody, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes),

each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are uncontaminated by other immunoglobulins.

[0033] The monoclonal antibodies herein specifically include “chimeric” antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity.

[0034] “Humanized” forms of non-human (e.g., murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable regions correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence, except for FR substitution(s) as noted above. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin.

[0035] A “human antibody” herein is one comprising an amino acid sequence structure that corresponds with the amino acid sequence structure of an antibody obtainable from a human B-cell, and includes antigen-binding fragments of human antibodies. Such antibodies can be identified or made by a variety of techniques, including, but not limited to: production by transgenic animals (e.g., mice) that are capable, upon immunization, of producing human antibodies in the absence of endogenous immunoglobulin; selection from phage display libraries expressing human antibodies or human antibody; generation via in vitro activated B; and isolation from human antibody producing hybridomas.

[0036] In certain embodiments, the antibody used in the present invention is a monoclonal antibody.

[0037] In other embodiments, the antibody used in the present invention is a chimeric, a humanized or a human antibody. In preferred embodiments, the antibody used in the present invention is a human antibody.

[0038] In certain embodiments, the antibody used in the present invention is administered in combination with an additional drug that treats RA.

[0039] The additional drug may be one or more medications, and include, for example, immunosuppressive agents, non-steroidal anti-inflammatory drugs (NSAIDs), disease modifying anti-rheumatic drugs (DMARDs) such as meth-

otrexate (MTX), anti-B-cell surface marker antibodies, such as anti-CD20 antibodies (e.g. rituximab), TNF-alpha-inhibitors, corticosteroids, and co-stimulatory modifiers, or any combination thereof. Optionally, the second or additional drug is selected from the group consisting of non-biological DMARDs, NSAIDs, and corticosteroids.

[0040] These additional drugs are generally used in the same dosages and with administration routes as used hereinbefore and hereinafter. If such additional drugs are used at all, preferably, they are used in lower amounts than if the first medicament were not present, especially in subsequent dosings beyond the initial dosing with the first medicament, so as to eliminate or reduce side effects caused thereby. The combined administration of an additional drug includes co-administration (concurrent administration), using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents (medicaments) simultaneously exert their biological activities.

[0041] The term "DMARD" refers to "Disease-Modifying Anti-Rheumatic Drugs" and includes among others hydroxychloroquine, sulfasalazine, methotrexate, leflunomide, azathioprine, D-penicillamine, gold salts (oral), gold salts (intramuscular), minocycline, cyclosporine including cyclosporine A and topical cyclosporine, and TNF-inhibitors, including salts, variants, and derivatives thereof. Exemplary DMARDs herein are non-biological, i.e. classic, DMARDs, including, azathioprine, chloroquine, hydroxychloroquine, leflunomide, methotrexate and sulfasalazine.

[0042] Methotrexate is an especially preferred DMARD of the present invention. Therefore, in certain embodiments, the antibody used in the present invention is administered in combination with a DMARD. In other embodiments, the antibody used in the present invention is administered in combination with methotrexate.

[0043] A "TNF-inhibitor" as used herein refers to an agent that inhibits, to some extent, a biological function of TNF-alpha, generally through binding to TNF-alpha and/or its receptor and neutralizing its activity. Examples of TNF inhibitors include etanercept (ENBREL®), infliximab (REMICADE®), adalimumab (HUMIRA®), certolizumab pegol (CIMZIA®), and golimumab (SIMPONI®).

[0044] "Treatment" of a patient or a subject refers to both therapeutic treatment and prophylactic or preventative measures. The terms "effective amount" or "therapeutically effective" refer to an amount of the antibody that is effective for treating rheumatoid arthritis. Such effective amount can result in any one or more of reducing the signs or symptoms of RA (e.g. achieving ACR20), reducing disease activity (e.g. Disease Activity Score, DAS20), slowing the progression of structural joint damage or improving physical function. In one embodiment, such clinical response is comparable to that achieved with intravenously administered anti-GM-CSF antibody.

[0045] The antibody of the present invention may be administered in different suitable forms. Potential forms of administration include systemic administration (subcutaneous, intravenous, intramuscular), oral administration, inhalation, transdermal administration, topical application (such as topical cream or ointment, etc.) or by other methods known in the art. The doses (in mg/kg) specified in the present invention refer to milligrams of antibody per kilogram of body weight of the patient. In vitro cell based assays showed that an anti-

GM-CSF antibody (MOR103) is capable of inhibiting several GM-CSF mediated responses. Evaluated responses include TF-1 cell proliferation, STAT5 phosphorylation, polymorphonuclear neutrophils (PMN) migration, PMN up-regulation of CD11b, monocyte up-regulation of MHC II, and eosinophil survival. Complete inhibitory effects were generally reached at concentrations of about 0.2 µg/ml anti-GM-CSF antibody. GM-CSF concentrations up to 1 ng/ml were applied in such studies. As a reference, GM-CSF levels in the synovial fluid of RA patients were reported to be <500 pg/ml. It is reasonable to consider that similar GM-CSF concentrations as used in these in vitro studies are present in affected tissues of RA patients

[0046] To effectively treat RA it may be important for an anti-GM-CSF antibody to penetrate the synovium. There is evidence to suggest that monoclonal antibodies can distribute into the synovium when dosed subcutaneously or intravenously. Based on a predicted penetration rate of 30%, continuous GM-CSF production and considering patient heterogeneity, the minimal or sub-optimal clinical effect level in RA patients is anticipated to be at a serum concentration of approximately 2 µg/ml antibody (thus, approximately 10-fold higher than the inhibitory concentration derived from in vitro studies).

[0047] A specific anti-GM-CSF antibody (MOR103) has been administered to patients with active rheumatoid arthritis who received 4 intravenous weekly doses of 0.3, 1, and 1.5 mg/kg. The anti-GM-CSF antibody showed significant clinical efficacy on DAS28, EULAR, ACR20, ACR50, ACR70 and tender joint counts following once a week dosing with 1 and 1.5 mg/kg as compared to placebo.

[0048] In certain embodiments, the antibody of the present invention is administered intravenously. In other embodiments, the antibody of the present invention is administered subcutaneously.

[0049] From other therapeutic antibodies it is known that a concentration that leads to a certain level of the antibody in the blood when administered intravenously corresponds to about 50-76% of the blood concentration achieved when the same antibody concentration is administered subcutaneously (Meibohm, B.: Pharmacokinetics and Pharmacodynamics of Biotech Drugs, Wiley-VCH, 2006). For MOR103 this ratio was determined to be 52%, i.e. a given concentration administered subcutaneously leads to a blood concentration which is equivalent to about 52% of the blood concentration when the same given concentration is administered intravenously. Therefore, the concentration of a subcutaneous formulation needs to be about twice as high to achieve the same drug blood level as compared to an intravenous formulation.

[0050] In certain embodiments the blood level to be achieved in a patient is equal or higher compared to the blood concentration achieved with intravenous administration of the antibody of the present invention at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0051] In alternative embodiments said blood concentration to be achieved is equal or higher compared to the blood concentration achieved with intravenous administration of the antibody of the present invention at a doses of at least 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 mg/kg when administered weekly over at least four weeks. In alternative embodiments the blood level to be achieved in a patient is equal or higher compared to the blood concentration achieved with intravenous administration of the antibody of the present invention at a dose of at least 1.0 mg/kg when administered weekly over at least two

weeks or at least three weeks. In alternative embodiments the blood level to be achieved in a patient is equal or higher compared to the blood concentration achieved with intravenous administration of the antibody of the present invention at a dose of at least 1.0 mg/kg when administered biweekly over at least two weeks or at least four weeks.

[0052] In certain embodiments, the antibody of the present invention is administered intravenously. In other embodiments, the antibody of the present invention is administered intravenously at a dose at least 0.6, at least 0.7, at least 0.8, at least 0.9 or at least 1.0 mg/kg. In other embodiments, the antibody of the present invention is administered intravenously at a dose of about 1.0 mg/kg or a dose of about 1.5 mg/kg.

[0053] In certain embodiments, the antibody of the present invention is administered subcutaneously. Various dosing regimen have been simulated using the subcutaneous delivery of MOR103 in order to produce plasma concentrations that are similar those obtained after 1 mg/kg iv, a dose that was efficacious in RA. The majority of simulations produce trough concentration values greater than 2 ug/mL, a value that is believed to be the minimum blood concentration that is required to produce efficacy in the context of an anti-GM-CSF antibody. These studies indicate that subcutaneous doses of 1, 2, 3 and 4 mg/kg can produce plasma concentration similar to 1 mg/kg, IV depending on the dosing frequency.

[0054] In other embodiments, the antibody of the present invention is administered subcutaneously at a dose at least 1.0, at least 1.5, at least 2.0, at least 2.5, at least 3.0, at least 3.5 or at least 4.0 mg/kg. In other embodiments, the antibody of the present invention is administered subcutaneously at a dose of about 2.0 mg/kg, a dose of about 3.0 mg/kg or a dose of about 4.0 mg/kg. In certain embodiments, the antibody of the present invention is subcutaneously administered biweekly, monthly or bimonthly.

[0055] In other embodiments, the antibody of the present invention is administered subcutaneously at a fixed dose. In such "fixed dose" treatment the antibody is administered at a certain, fixed, concentration, i.e. without taking into account a patient's body weight. In certain embodiments, the antibody of the present invention is administered at a fixed dose of between 40 mg and 400 mg, optionally at a fixed dose of 75 mg, at a fixed dose of 100 mg, at a fixed dose of 140 mg, at a fixed dose of 150 mg, at a fixed dose of 180 mg, at a fixed dose of 200 mg, at a fixed dose of 280 mg, at a fixed dose of 300 mg or at a fixed dose of 400 mg. Administration of fixed doses may be every week, every second week, every third week, every fourth week or every sixth week. Typically, the antibody will be administered weekly at a fixed dose.

[0056] In an embodiment, the antibody will be administered weekly, at a fixed subcutaneous dose of 40, 56, 70, 75, 100, 140, 150, 180, 200, 210, or 280 mg.

[0057] In an embodiment, the antibody will be administered biweekly, at a fixed subcutaneous dose of 70, 75, 100, 140, 150, 180, 200, 210, 280 or 300 mg.

[0058] In an embodiment, the antibody will be administered monthly, at a fixed subcutaneous dose of 100, 140, 150, 180, 200, 210, 280, 300, 320, 350, 360 or 400 mg.

[0059] In an embodiment, the antibody is administered in a dose sufficient to maintain trough concentration of antibody of at least 2 ug/mL. The trough concentration of antibody may be maintained at 2.0 ug/mL, 2.5 ug/mL, 3.0 ug/mL, 3.5 ug/mL, 4.0 ug/mL, 4.5 ug/mL or 5.0 ug/mL, during the course of therapy.

[0060] In alternative embodiments, the antibody will be administered weekly, at a fixed subcutaneous dose of 28 or 35 mg,

[0061] In certain embodiments, the present invention provides an anti-GM-CSF antibody for use in the treatment of a patient suffering from rheumatoid arthritis, wherein said antibody is administered to said patient in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0062] In certain embodiments, the present invention provides a method to treat a patient suffering from rheumatoid arthritis, said method comprising administering to said patient an anti-GM-CSF antibody in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0063] In certain embodiments, the present invention provides an anti-GM-CSF antibody for inhibiting progression of structural joint damage in a rheumatoid arthritis patient comprising administering to said patient said antibody in a manner to achieve a therapeutically effective antibody level in the blood of said patient equal or higher compared to the intravenous administration of said antibody at a dose of at least 1.0 mg/kg when administered weekly over at least four weeks.

[0064] The terms "drug" and "medicament" refer to an active drug to treat rheumatoid arthritis or joint damage or symptoms or side effects associated with RA. The term "pharmaceutical formulation" refers to a preparation which is in such form as to permit the biological activity of the active ingredient or ingredients, i.e. the antibody of the present invention, to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered. Such formulations are sterile.

[0065] The antibody herein is preferably recombinantly produced in a host cell transformed with nucleic acid sequences encoding its heavy and light chains (e.g. where the host cell has been transformed by one or more vectors with the nucleic acid therein). The preferred host cell is a mammalian cell, most preferably a PER.C6 cell.

[0066] Therapeutic formulations of the antibody of the present invention are prepared for storage by mixing the antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, histidine and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes {e.g. Zn-protein complexes}; and/or non-ionic

surfactants such as TWEEN™ (such as Tween-80), PLURONICS™ or polyethylene glycol (PEG).

[0067] In certain embodiments, the present invention provides a pharmaceutical composition comprising an antibody of the present invention and a pharmaceutically acceptable carrier and/or excipient for use in any of the methods provided in the present invention. In certain embodiments, the formulation for the antibody of the present invention consists of 30 mM histidine, pH 6.0, 200 mM sorbitol and 0.02% Tween-80. In other embodiments, the formulation for the antibody of the present invention consists of PBS, pH 6.2 (0.2 g/l KCl, 0.96 g/l KH_2PO_4 , 0.66 g/l $\text{Na}_2\text{HPO}_4 \times 7\text{H}_2\text{O}$, 8 g/l NaCl).

EXAMPLES

Example 1

Design and Concept of a Clinical Phase Ib/Phase IIa Trial

[0068] A multi-center, randomized, double-blinded, placebo-controlled study to evaluate the safety, preliminary clinical activity and immunogenicity of multiple doses of MOR103 administered intravenously to patients with active rheumatoid arthritis was conducted.

[0069] Primary outcome measures were the adverse event rate and the safety profile. Secondary outcome measures included DAS28 scores, ACR scores and EULAR28 response criteria.

[0070] The clinical trial comprised three treatment arms. In each treatment arm patient received either placebo or MOR103. The MOR103 doses were 0.3 mg/kg body weight for treatment arm 1, 1.0 mg/kg body weight for treatment arm 2 and 1.5 mg/kg body weight for treatment arm 3. MOR103 and placebo were administered intravenously, weekly with 4 doses in total.

[0071] Summary of the treatment arms:

Arms	Assigned Interventions
Experimental: Group 1: MOR103, experimental Biological: MOR103 0.3 mg/kg or placebo	Drug: MOR103 MOR103 0.3 mg/kg or placebo iv x 4 doses
Experimental: Group 2: MOR103, experimental Biological: MOR103 1.0 mg/kg or placebo	Drug: MOR103 MOR103 1.0 mg/kg or placebo iv x 4 doses
Experimental: Group 3: MOR103, experimental Biological: MOR103 1.5 mg/kg or placebo	Drug: MOR103 MOR103 1.5 mg/kg or placebo iv x 4 doses

[0072] Eligible for participation in the study were patients of 18 years and older and of either sex (male and female). Healthy volunteers were not accepted.

[0073] Inclusion criteria were as follows:

[0074] Rheumatoid arthritis (RA) per revised 1987 ACR criteria

[0075] Active RA: ≥ 3 swollen and 3 tender joints with at least 1 swollen joint in the hand, excluding the PIP joints

[0076] CRP > 5.0 mg/L (RF and anti-CCP seronegative); CRP > 2 mg/l (RF and/or anti-CCP seropositive)

[0077] DAS28 ≤ 5.1

[0078] Stable regimen of concomitant RA therapy (NSAIDs, steroids, non-biological DMARDs).

[0079] Negative PPD tuberculin skin test

Exclusion criteria were as follows:

[0080] Previous therapy with B or T cell depleting agents other than Rituximab (e.g. Campath). Prior treatment with Rituximab, TNF-inhibitors, other biologics (e.g. anti-IL-1 therapy) and systemic immunosuppressive agents is allowed with a washout period.

[0081] Any history of ongoing, significant or recurring infections

[0082] Any active inflammatory diseases other than RA

[0083] Treatment with a systemic investigational drug within 6 months prior to screening

[0084] Women of childbearing potential, unless receiving stable doses of methotrexate or leflunomide

[0085] Significant cardiac or pulmonary disease (including methotrexate-associated lung toxicity)

[0086] Hepatic or renal insufficiency

Example 2

Patient Recruitment and Patient Population

[0087] Clinical sites for patient recruitment, screening and treatment were located in Bulgaria, Germany, the Netherlands, Poland and the Ukraine.

[0088] 96 patients were included in the trial. 27 patients received placebo, 24 patients received MOR103 at a dose of 0.3 mg/kg, 22 patients received MOR103 at a dose of 1.0 mg/kg and 23 patients received MOR103 at a dose of 1.5 mg/kg. The average age and the average Body Mass Index (BMI) was about the same for all treatment groups. Key characteristics are summarized in the following Table:

Characteristic	MOR 103 Active Treatment Groups				
	Placebo N = 27	0.3 mg/kg N = 24	1.0 mg/kg N = 22	1.5 mg/kg N = 23	Total active N = 69
Age	53.8	57.4	49	53	53.3
BMI	26.3	26.3	26.1	25.7	26.0
Gender- Female	19 (70%)	21 (88%)	17 (77%)	18 (78%)	56 (81%)
White	27	24	22	23	69

[0089] 90% of all patients of the study were previously treated with DMARDs. The most commonly used DMARD was methotrexate (75% of all patients). The rate of previous treatment with DMARDs was comparable in all treatment arms.

[0090] Prior to administration of MOR103 or the placebo the disease activity of all patients was measured according to accepted guidelines by calculating the DAS28 score, a 28-joint Disease Activity Score (see e.g. Ann Rheum Dis (2009) 68, 954-60). DAS28 score is a validated and commonly used tool to quantify the disease status of RA patients. The average DAS28 score was comparable for all treatment arms.

Example 3

Safety Profile

[0091] Based on the available observed safety data, MOR103 showed a favorable safety profile among all doses tested. The key observations are as follows:

[0092] No deaths were observed during the conduct of the trial

[0093] No infusion related reactions were observed

[0094] Two serious adverse events (SAEs) were observed:
 [0095] One patient in the placebo group developed paronychia
 [0096] One patient in the 0.3 mg/kg treatment arm developed pleurisy
 [0097] More treatment-emergent adverse effects (TEAEs) were observed in the placebo group (25.9%) than in the active groups (14.5%)
 [0098] Most TEAEs were mild
 [0099] No severe TEAEs were observed in the active groups
 [0100] In summary, it can be concluded that treatment with MOR103 at all doses tested is safe. Two serious adverse events were observed, both none in the treatment arms that showed clinical efficacy (see below). Sub-cutaneous administration of MOR103 at a dose that leads to an antibody drug level in the blood of patients equivalent to the intravenous application of the present study is expected to show a similar safety profile.

Example 4

Efficacy—DAS28

[0101] 4 weeks and 8 weeks after the first administration of MOR103 (or placebo) the DAS28 scores of all patients was determined. A decrease in DAS28 scores correlates to diminished disease severity. Results are shown in FIG. 2 as the mean changes compared to baseline, i.e. disease status prior to treatment.
 [0102] The placebo group only shows marginal changes. Patients treated with MOR103 at 0.3 mg/kg showed a slight decrease in DAS28 scores, indicating slightly less severity of the disease. In contrast, patients treated with MOR103 at 1.0 mg/kg or with 1.5 mg/kg showed a significant decrease in DAS28 scores, indicating the high efficacy of MOR103 at these doses.

Example 5

Efficacy—ACR20

[0103] As another measure of efficacy the ACR20 criteria were used. ACR criteria measure improvement in tender or swollen joint counts and improvement in certain other parameters. The procedure to measure ACR scores is highly standardized. The present clinical trial applied the respective applicable guidelines. Results are depicted in FIGS. 3 and 4. A higher score corresponds to an improvement in the severity of the disease.
 [0104] In line with the results of the DAS28 scores (see Example 4), also the ACR scores show a strong clinical improvement of patients' condition upon treatment with either 1.0 mg/kg MOR103 or 1.5 mg/kg MOR103. The

improvement after 4 weeks is highly significant for the 1.0 mg/kg group ($p < 0.0001$). Taken together, the ACR20 scores confirm the surprising finding that the efficacy of MOR103 can already be shown with a comparably low number of patients in each treatment arm and a comparably short treatment period.

Example 6

Clinical Trial with Additional Doses of MOR103

[0105] The clinical trial set out herein above is repeated with additional doses of MOR103. MOR103 is administered to patients intravenously at a dose of 0.5 mg/kg (treatment arm 1) and 0.75 mg/kg (treatment arm 2). All other parameters are identical to Example 1.
 [0106] Both treatment arms show a favorable safety profile and demonstrate clinical efficacy as measured by DAS28 scores and ACR20 scores.

Example 7

Clinical Trial with a Sub-Cutaneous Formulation of MOR103

[0107] The clinical trial set out herein above is repeated with a sub-cutaneous formulation of MOR103. In order to achieve similar levels of MOR103 in the blood of patients as observed for intravenous treatment, the sub-cutaneous dose of MOR103 is increased.
 [0108] In different treatment arms MOR103 is administered to patients at 1.5 mg/kg, 2.0 mg/kg, 3.0 mg/kg and 4.0 mg/kg. The drug is administered sub-cutaneously, either biweekly, monthly or bimonthly. All other parameters are identical to Example 1.
 [0109] All treatment arms show a favorable safety profile and demonstrate clinical efficacy as measured by DAS28 scores and ACR20 scores.

Example 8

Clinical Trial with a Sub-Cutaneous Formulation of MOR103 at a Fixed Dose

[0110] Example 7 is repeated with a fixed dose of MOR103. In different treatment arms MOR103 is administered to patients at fixed dose of 75 mg, of 100 mg, of 150 mg, of 200 mg, of 300 mg and of 400 mg. The drug is administered sub-cutaneously every week, every second week, every fourth week or every sixth week. All other parameters are identical to the Examples described herein above.
 [0111] All treatment arms show a favorable safety profile and demonstrate clinical efficacy as measured by DAS28 scores and ACR20 scores.

SEQUENCE LISTING

```
<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1
<211> LENGTH: 144
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens
```

-continued

<400> SEQUENCE: 1

Met Trp Leu Gln Ser Leu Leu Leu Leu Gly Thr Val Ala Cys Ser Ile
1 5 10 15
Ser Ala Pro Ala Arg Ser Pro Ser Pro Ser Thr Gln Pro Trp Glu His
20 25 30
Val Asn Ala Ile Gln Glu Ala Arg Arg Leu Leu Asn Leu Ser Arg Asp
35 40 45
Thr Ala Ala Glu Met Asn Glu Thr Val Glu Val Ile Ser Glu Met Phe
50 55 60
Asp Leu Gln Glu Pro Thr Cys Leu Gln Thr Arg Leu Glu Leu Tyr Lys
65 70 75 80
Gln Gly Leu Arg Gly Ser Leu Thr Lys Leu Lys Gly Pro Leu Thr Met
85 90 95
Met Ala Ser His Tyr Lys Gln His Cys Pro Pro Thr Pro Glu Thr Ser
100 105 110
Cys Ala Thr Gln Ile Ile Thr Phe Glu Ser Phe Lys Glu Asn Leu Lys
115 120 125
Asp Phe Leu Leu Val Ile Pro Phe Asp Cys Trp Glu Pro Val Gln Glu
130 135 140

<210> SEQ ID NO 2

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 2

Gly Phe Thr Phe Ser Ser Tyr Trp Met Asn
1 5 10

<210> SEQ ID NO 3

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 3

Gly Ile Glu Asn Lys Tyr Ala Gly Gly Ala Thr Tyr Tyr Ala Ala Ser
1 5 10 15

Val Lys Gly

<210> SEQ ID NO 4

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 4

Gly Phe Gly Thr Asp Phe
1 5

<210> SEQ ID NO 5

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 5

Ser Gly Asp Ser Ile Gly Lys Lys Tyr Ala Tyr
1 5 10

<210> SEQ ID NO 6

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 6

Lys Lys Arg Pro Ser
1 5

<210> SEQ ID NO 7

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 7

Ser Ala Trp Gly Asp Lys Gly Met
1 5

<210> SEQ ID NO 8

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 8

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Trp Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Gly Ile Glu Asn Lys Tyr Ala Gly Gly Ala Thr Tyr Tyr Ala Ala
50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
65 70 75 80

Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85 90 95

Tyr Cys Ala Arg Gly Phe Gly Thr Asp Phe Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> SEQ ID NO 9

<211> LENGTH: 106

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: polypeptide

<400> SEQUENCE: 9

-continued

Asp	Ile	Glu	Leu	Thr	Gln	Pro	Pro	Ser	Val	Ser	Val	Ala	Pro	Gly	Gln
1				5					10					15	
Thr	Ala	Arg	Ile	Ser	Cys	Ser	Gly	Asp	Ser	Ile	Gly	Lys	Lys	Tyr	Ala
			20					25					30		
Tyr	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Val	Leu	Val	Ile	Tyr
		35					40					45			
Lys	Lys	Arg	Pro	Ser	Gly	Ile	Pro	Glu	Arg	Phe	Ser	Gly	Ser	Asn	Ser
		50				55				60					
Gly	Asn	Thr	Ala	Thr	Leu	Thr	Ile	Ser	Gly	Thr	Gln	Ala	Glu	Asp	Glu
65					70					75				80	
Ala	Asp	Tyr	Tyr	Cys	Ser	Ala	Trp	Gly	Asp	Lys	Gly	Met	Val	Phe	Gly
				85					90					95	
Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln						
			100					105							

1-20. (canceled)

21. A method of treating a patient suffering from rheumatoid arthritis, said method comprising administering to said patient an anti-GM-CSF antibody subcutaneously at a dose selected from the group consisting of:

- (i) a dose of at least 1.0 mg/kg; and
- (ii) a fixed dose of between 40 mg and 400 mg.

22. The method of claim **21**, wherein said antibody is administered at a dose of at least 2.0 mg/kg.

23. The method of claim **21**, wherein said antibody is administered at a fixed dose of 75 mg.

24. The method of claim **21**, wherein said antibody is administered in a manner selected from the group consisting of:

- (i) biweekly;
- (ii) monthly; and
- (iii) bimonthly.

25. The method of claim **21**, wherein said antibody is administered in a manner selected from the group consisting of:

- (i) every week;
- (ii) every second week;
- (iii) every third week;
- (iv) every fourth week; and
- (v) every sixth week.

26. The method of claim **25**, wherein said antibody is administered weekly.

27. The method of claim **21**, wherein said anti-GM-CSF antibody is administered in combination with a Disease-modifying antirheumatic drug (DMARD).

28. The method of claim **27**, where in the DMARD is methotrexate.

29. A method of treating a patient suffering from rheumatoid arthritis, said method comprising administering to said patient an anti-GM-CSF antibody subcutaneously at a dose selected from the group consisting of:

- (i) a dose of at least 1.0 mg/kg; and
 - (ii) a fixed dose of between 40 mg and 400 mg;
- wherein said anti-GM-CSF antibody is an antibody comprising an HCDR1 region of sequence GFTFSSYWMN (SEQ ID NO.: 2), an HCDR2 region of sequence GIEN-KYAGGATYYAASVKG (SEQ ID NO.: 3), an HCDR3 region of sequence GFGTDF (SEQ ID NO.: 4), an

LCDR1 region of sequence SGDSIGKKYAY (SEQ ID NO.: 5), an LCDR2 region of sequence KKRPS (SEQ ID NO.: 6), and an LCDR3 region of sequence SAWGD-KGM (SEQ ID NO.: 7).

30. The method of claim **21**, said anti-GM-CSF antibody further comprising a pharmaceutically acceptable carrier.

31. The method of claim **30**, wherein the pharmaceutically acceptable carrier is selected from the group consisting of: buffers; antioxidants; preservatives; alkyl parabens; low molecular weight (less than about 10 residues) polypeptides; proteins; hydrophilic polymers; amino acids; sugars including monosaccharides, disaccharides, and other carbohydrates; chelating agents; sugars; salt-forming counter-ions; metal complexes; and non-ionic surfactants.

32. The method of claim **31**, wherein the buffer is selected from the group consisting of: phosphate, citrate, histidine, and other organic acids.

33. The method of claim **31**, wherein the sugar is selected from the group consisting of: sucrose, mannitol, trehalose and sorbitol.

34. The method of claim **31**, wherein the non-ionic surfactant is selected from the group consisting of: TWEEN™ (such as Tween-80), PLURONIC™, and polyethylene glycol (PEG).

35. The method of claim **31**, wherein the buffer is histidine; the sugar is sorbitol; and the non-ionic surfactant is Tween-80.

36. The method of claim **31**, wherein the pharmaceutically acceptable carrier comprises 30 mM histidine, pH 6.0, 200 mM sorbitol, and 0.02% Tween-80.

37. A method of treating a patient suffering from rheumatoid arthritis, said method comprising administering to said patient a pharmaceutical composition comprising an anti-GM-CSF antibody; a buffer which is histidine; a sugar which is sorbitol; and a non-ionic surfactant which is Tween-80 subcutaneously at a dose selected from the group consisting of:

- (iii) a dose of at least 1.0 mg/kg; and
- (iv) a fixed dose of between 40 mg and 400 mg.

38. A pharmaceutical composition comprising:

- (i) an anti-GM-CSF antibody;
- (ii) a buffer selected from the group consisting of: phosphate, citrate, histidine and other organic acids;

(iii) a sugar selected from the group consisting of: sucrose, mannitol, trehalose and sorbitol; and

(iv) a non-ionic surfactant selected from the group consisting of: TWEENTM (such as Tween-80), PLURON-ICSTM, and polyethylene glycol (PEG).

39. The pharmaceutical composition according to claim **38**, wherein the buffer is histidine; the sugar is sorbitol; and the non-ionic surfactant is Tween-80.

40. A pharmaceutical composition comprising an anti-GM-CSF antibody and a pharmaceutically acceptable carrier, wherein the carrier consists of 30 mM histidine, pH 6.0, 200 mM sorbitol and 0.02% Tween-80.

* * * * *