

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 200054355 B2**
(10) Patent No. **770652**

(54) Title
An integrin heterodimer and an alpha subunit thereof

(51) ⁶ International Patent Classification(s)
C07K 014/705 C07K 016/28
A61K 038/17

(21) Application No: 200054355 (22) Application Date: 2000.05.31

(87) WIPO No: W000/75187

(30) Priority Data

(31) Number	(32) Date	(33) Country
9902056	1999.06.03	SE

(43) Publication Date : 2000.12.28
(43) Publication Journal Date : 2001.03.08
(44) Accepted Journal Date : 2004.02.26

(71) Applicant(s)
Cartela AB

(72) Inventor(s)
Donald Gullberg

(74) Agent/Attorney
Griffith Hack,GPO Box 4164,SYDNEY NSW 2001

(56) Related Art
GENINI ET AL., (1996) INT. J. CANCER 66:571-577
GULLBERG ET AL. (1995) DEVELOPMENT DYNAMICS 204:57-65

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number
WO 00/75187 A1

(51) International Patent Classification: C07K 14/705,
A61K 38/17, C07K 16/28

(21) International Application Number: PCT/SE00/01135

(22) International Filing Date: 31 May 2000 (31.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9902056-2 3 June 1999 (03.06.1999) SE

(71) Applicant (for all designated States except US): **ACTIVE**
BIOTECH AB (SE/SE); Box 724, S-220 07 Lund (SE).
BMC, B12 SE-221 84 Lund, (SE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): GULLBERG, Donald
[SE/SE]; Björkgatan 3F, S-753 28 Uppsala (SE).

(74) Agent: AWAPATENT AB; Box 5117, S-200 71 Malmö
(SE).

(81) Designated States (national): AE, AG, AL, AM, AT, AT
(utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH,

CN, CR, CU, CZ, CZ (utility model), DE, DE (utility
model), DK, DK (utility model), DM, DZ, EE, EE (utility
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (utility
model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 00/75187 A1

(54) Title: AN INTEGRIN HETERODIMER AND AN ALPHA SUBUNIT THEREOF

(57) Abstract: A recombinant or isolated integrin heterodimer comprising a novel subunit $\alpha 11$ in association with a subunit β is described. The integrin or the subunit $\alpha 11$ can be used as marker or target of all types of cells. The integrin or subunit $\alpha 11$ thereof can be used as marker or target in different physiological or therapeutic methods. They can also be used as active ingredients in pharmaceutical compositions and vaccines.

AN INTEGRIN HETERODIMER AND AN ALPHA SUBUNIT THEREOFFIELD OF THE INVENTION

The present invention relates to a recombinant or isolated integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , the subunit $\alpha 11$ thereof, homologues and
5 fragments of said integrin and of said subunit $\alpha 11$,
processes of producing the same, polynucleotides and
oligonucleotides encoding the same, vectors and cells
comprising the same, binding entities binding
specifically to binding sites of the same, and the use of
10 the same.

BACKGROUND OF THE INVENTION

Integrins are heterodimers composed of non-covalently associated α - and β -chains which connect cells to the extracellular matrix or to other cells (1). In addition
15 to acting as mechanical links between the cytoskeleton and extracellular ligands, integrins are signal transducing receptors which influence processes such as cell proliferation, cell migration and cell differentiation (2-4). Integrins can be grouped into subfamilies based on
20 shared β -chains, shared ligand binding properties, or shared structural features of the α -chains. Currently 17 α -chains and 8 β -chains have been identified (5). Of the subfamilies with shared β -chains, the $\beta 1$ subfamily has the most members. To date, 11 integrin α -chains associated with the $\beta 1$ -chain have been identified and characterized, $\alpha 1$ - $\alpha 10$ and αv (5).

Several integrins bind the sequence RGD in their respective ligands (1). Of those integrins identified so far, $\alpha 4$ -, $\alpha 5$ -, $\alpha 8$ -, αIIb - and αv -chains form heterodimers
30 that mediate RGD-dependent interactions. The ligands containing RGD are generally found in the interstitial type of extracellular matrix. Major non-RGD dependent ligands include various collagen and laminin isoforms. Although both collagens and laminins contain the RGD

sequence in their primary sequences, these RGD sequences are cryptic (6-9) and normally not accessible to cells in the native proteins, but they may be exposed during growth and reorganization events of the extracellular matrix.

Another subdivision of integrins can be made based on structural similarities of the α -chains. A number of integrins contain an extracellular I-domain (10,11) which is homologous to collagen binding A-domains present in von Willebrand factor (12). The I-domain constitutes an inserted domain of approximately 200 amino acids which is present in 8 known integrins ($\alpha 1$, $\alpha 2$, $\alpha 10$, αL , αM , αX , αD and αE) (5,10). Structural analysis of integrin I-domains crystallized in the presence of Mg^{2+} have revealed the presence of a characteristic "MIDAS" (metal ion dependent adhesion site) motif, shown to be critical for ligand binding (13). Integrin α -chains containing the I-domain are not cleaved into heavy and light chains, although the rat $\alpha 1$ chain possesses a proteolytic cleavage site near the membrane spanning region (14,15). For I-domain integrins the principal ligand binding sites are found within the I-domain (10). Known ligands for I-domains found within the $\beta 1$ integrin subfamily include laminins and collagens ($\alpha 1\beta 1$ and $\alpha 2\beta 1$ integrins) (16-19), and Echo-virus ($\alpha 2\beta 1$ integrin) (20).

Structure comparisons have suggested that integrins fold into a so-called 7-bladed β -propeller structure which forms one globular domain with the ligand binding region on the upper surface (21). The I-domain is inserted between blade 2 and 3 in this propeller and divalent cation binding sites are located on the lower surface in blades 4-7 (22,23). Studies of $\beta 2$ integrins have revealed that proper folding of the $\beta 2$ -chain is dependent on the presence of the αL -chain but that the I-domain folds independently of other structural elements in the α - and β -chains (24). In integrin α -chains, a less conserved stalk region separates the predicted β -pro-

pellor from the short transmembrane region. This stalk region is possibly involved in transducing conformational changes between the extracellular and intracellular regions, as well as mediating protein-protein interactions.

5 Although integrins take part in cell signalling events, the cytoplasmic tail is short and lacks enzymatic activity. The sequence GFFKR is conserved in a majority of integrin α -subunits cytoplasmic tails and has been shown to be important for calreticulin binding (25).

10 Cellular interactions with the extracellular matrix during muscle formation and in muscular dystrophy have received increased interest during the past years. In the early 1960's a mutant was described in *Drosophila* which was characterized by the detachment of muscles from
15 their attachment points at the time of the first embryonic muscle contraction, causing the embryos to assume a spheroid shape (26). The mapping of the molecular defect in the lethal mysospheroid mutant in 1988 to an integrin β -chain (27), was the first evidence for a role
20 of integrins in maintaining muscle integrity. More recently, refined analysis of *Drosophila* mutants have indicated distinct roles for integrins in muscle endpoint attachments and sarcomere structure (28). The *Drosophila* integrins are all cleaved α -chains and share many features with vertebrate integrins such as the ability to
25 cluster into focal contacts (29).

The finding that inactivation of the $\alpha 7$ integrin gene in mouse (30), as well as mutations in the human ITGA7 gene (31), both cause muscular dystrophy affecting
30 mainly muscle attachment points, indicates a striking conservation of integrin function during evolution. Of the 11 members of the $\beta 1$ subfamily, $\alpha 7$ exists as a major integrin α -chain (32,33) associated with the $\beta 1 D$ integrin chain in the adult skeletal muscle sarcolemma (34).
35 Intriguingly, mutations in the basement membrane protein laminin $\alpha 2$ -chain (35-37) cause a more severe disease than that observed for the laminin receptor integrin $\alpha 7 \beta 1$

(30). This indicates that other receptors for laminins exist in muscle.

A novel integrin has recently been identified on cultured human fetal muscle cells (38). The present invention is related to, inter alia, the cloning and characterization of this novel I-domain containing, $\beta 1$ -associated integrin chain, which is expressed in muscle tissues.

SUMMARY OF THE INVENTION

10 The full-length cDNA for this integrin subunit, $\alpha 11$, has now been isolated. The open reading frame of the cDNA encodes a precursor of 1188 amino acids. The predicted mature protein of 1166 amino acids contains 7 conserved FG-GAP repeats, an I-domain with a MIDAS motif, a short
15 transmembrane region and a unique cytoplasmic domain of 24 amino acids containing the sequence GFFRS. $\alpha 11$, like other I-domain integrins, lacks a dibasic cleavage site for generation of a heavy and a light chain, and contains three potential divalent cation binding sites in re-
20 peats 5-7. The presence of 22 inserted amino acids in the extracellular stalk portion (amino acids 804-826) distinguishes the $\alpha 11$ integrin sequence from other integrin α -chains. Amino acid sequence comparisons reveal the highest identity of 42% with $\alpha 10$ integrin chain. Immuno-
25 precipitation with antibodies to $\alpha 11$ integrin captures a 145 kD protein, distinctly larger than the 140 kD $\alpha 2$ integrin chain when analyzed by SDS-PAGE under non-reducing conditions. Fluorescence in situ hybridization maps the integrin $\alpha 11$ gene to chromosome 15q23, in the
30 vicinity of an identified locus for Bardet-Biedl syndrome. Based on Northern blotting integrin $\alpha 11$ mRNA levels are high in adult human uterus and in heart, and intermediate in skeletal muscle and some other tissues tested. During in vitro myogenic differentiation, $\alpha 11$
35 mRNA and protein are up-regulated. Studies of ligand binding properties show that $\alpha 11\beta 1$ binds collagen type

I Sepharose and cultured muscle cells localize $\alpha 11 \beta 1$ into focal contacts on collagen type I.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates in its different aspects to the following:

A recombinant or isolated integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID NO. 1, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein.

The invention also encompasses integrin homologues of said integrin, isolated from other species, such as bovine integrin heterodimer comprising a subunit $\alpha 11$ in association with a subunit β , preferably $\beta 1$, as well as homologues isolated from other types of human cells or from cells originating from other species.

The term "homologues" in the context of the present invention is meant to imply proteins of a common evolutionary origin, having identical or similar functions, specifically requiring evidence based on gene structure and not merely a similarity of protein structure.

The invention also encompasses a process of producing a recombinant integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID NO. 1, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which process comprises the steps of

a) isolating a polynucleotide comprising a nucleotide sequence coding for an integrin subunit $\alpha 11$, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein,

b) constructing an expression vector comprising the isolated polynucleotide,

c) transforming a host cell with said expression vector,

d) culturing said transformed host cell in a culture medium under conditions suitable for expression of integrin subunit $\alpha 11$, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, in said transformed host cell, and, optionally,

e) isolating the integrin subunit $\alpha 11$, or homologues or fragments thereof, from said transformed host cell or said culture medium. The transformation can be performed *in vitro*, *in situ* or *in vivo*.

In further aspects, the invention encompasses:

- A process of providing an integrin subunit $\alpha 11$, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, whereby said subunit is isolated from a cell in which it is naturally present.

- An isolated polynucleotide comprising a nucleotide coding for said integrin subunit $\alpha 11$, or for homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which polynucleotide comprises essentially the nucleotide sequence shown in SEQ ID NO. 1 or suitable parts thereof.

- An isolated polynucleotide or oligonucleotide which hybridises to a polynucleotide or oligonucleotide encoding said integrin subunit $\alpha 11$ or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, wherein said isolated polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit $\alpha 10$.

- A vector comprising a polynucleotide or oligonucleotide coding for said integrin subunit $\alpha 11$, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which polynucleotide or oligonucleotide comprises the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof.

- A vector comprising a polynucleotide or oligonucleotide which hybridises to a DNA or RNA encoding an integrin subunit $\alpha 11$ or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit $\alpha 10$.

- A cell containing the vector as defined above.

- A cell generated during the process as defined above, in which a polynucleotide or oligonucleotide coding for said integrin subunit $\alpha 11$, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which polynucleotide or oligonucleotide comprises essentially the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof, has been stably integrated in the cell genome.

- An isolated protein comprising a binding site of the amino acid sequence of the integrin subunit $\alpha 11$, or of homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, said binding sites having the capability of binding specifically to entities chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

- Binding entities having the capability of binding specifically to integrin subunit $\alpha 11$ comprising the amino acid sequence of SEQ ID NO. 1 or to homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, preferably chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

- A recombinant or isolated integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , in which the subunit $\alpha 11$ comprises essentially the amino acid sequence

shown in SEQ ID NO. 1, or homologues and fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein. Said subunit β is preferably $\beta 1$.

- 5 - A process of producing a recombinant integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , in which the subunit $\alpha 11$ comprises essentially the amino acid sequence shown in SEQ ID NO. 1, or homologues or fragments thereof wherein the fragments include at least one binding
10 site of the integrin subunit $\alpha 11$ protein, which process comprises the steps of

- a) isolating one polynucleotide comprising a nucleotide sequence coding for a subunit $\alpha 11$ of an integrin heterodimer and, optionally, another
15 polynucleotide comprising a nucleotide sequence coding for a subunit β of an integrin heterodimer, or polynucleotides or oligonucleotides coding for homologues or fragments thereof having similar biological activity,

- b) constructing an expression vector comprising said
20 isolated polynucleotide coding for said subunit $\alpha 11$ optionally in combination with an expression vector comprising said isolated nucleotide coding for said subunit β ,

- c) transforming a host cell with said expression
25 vector or vectors, which transformation may be performed *in vitro*, *in situ* or *in vivo*,

- d) culturing said transformed host cell in a culture medium under conditions suitable for expression of an integrin heterodimer comprising a subunit $\alpha 11$ and a
30 subunit β , or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, in said transformed host cell, and, optionally,

- e) isolating the integrin heterodimer comprising a
35 subunit $\alpha 11$ and a subunit β , or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, or the $\alpha 11$

subunit thereof from said transformed host cell or said culture medium.

- A process of providing an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or homologues or
5 fragments thereof having similar biological activity, wherein said integrin heterodimer is isolated from a cell in which it is naturally present.

- A cell containing

i) a first vector, said first vector comprising a
10 polynucleotide or oligonucleotide coding for a subunit $\alpha 11$ of an integrin heterodimer, or for homologues or parts thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which
polynucleotide or oligonucleotide comprises essentially
15 the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof, and

ii) a second vector, said second vector comprising a polynucleotide or oligonucleotide coding for a subunit β of an integrin heterodimer, or for homologues or fragments
20 thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein.

- Binding sites of an integrin heterodimer as defined above, or of homologues or fragments thereof, said binding sites having the capability of binding specifically to
25 entities chosen among the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

- Binding entities having the capability of binding
30 specifically to said integrin heterodimer, or to homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, or a subunit $\alpha 11$ thereof. Said subunit β is preferably $\beta 1$. The binding entities are preferably chosen
35 among the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.

- A fragment of the integrin subunit $\alpha 11$, which fragment is a peptide chosen from the group comprising peptides of the cytoplasmic domain, especially a peptide comprising essentially the amino acid sequence
5 KLGFFRSARRRREPGLDTPKVLE, of the I-domain, especially a peptide comprising essentially the amino acid sequence from about amino acid No. 159 to about amino acid No. 355 of SEQ ID No. 1, and the extracellular extension region, especially a peptide comprising essentially the amino acid
10 sequence from about amino acid No. 804 to about amino acid No. 826 of SEQ ID No. 1.

- A method of producing a fragment of the integrin subunit $\alpha 11$ as defined above, which method comprises a sequential addition of amino acids. This method comprises
15 adding and removing protective groups in a manner known by the man skilled in the art.

- A polynucleotide or oligonucleotide coding for a fragment of the integrin subunit $\alpha 11$ as defined above.

- Binding sites of a fragment as defined above, said
20 binding sites having the capability of binding specifically to entities chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.

- Binding entities having the capability of binding
25 specifically to a fragment as defined of the human integrin subunit $\alpha 11$ as defined above. Preferably, said binding entities are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, and fragments thereof.

- A process of using an integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID No. 1 or an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β , or a homologue or fragment of said integrin or subunit wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as a marker or target molecule of cells or tissues expressing said integrin subunit $\alpha 11$, which cells or tissues are of animal including human origin. Especially, said subunit β is $\beta 1$.

In embodiments of this process, said fragment is a peptide chosen from the above defined group.

In one embodiment of said process, the cells are chosen from the group comprising fibroblasts, muscle cells, chondrocytes, osteoblasts, mesenchymally derived cells and stem cells.

Especially, said process is used during pathological conditions involving said subunit $\alpha 11$. Said pathological conditions comprise in one embodiment damage of muscles, muscle dystrophy, fibrosis or wound healing. In another embodiment, said pathological conditions comprise damage of cartilage and/or bone, or cartilage and/or bone diseases. In a still further embodiment, said pathological conditions comprise trauma, rheumatoid arthritis, osteoarthritis or osteoporosis.

In a further embodiment, said process is a process for detecting the formation of cartilage during embryonic development, or for detecting physiological or therapeutic reparation of cartilage and/or muscle, or for selection and analysis, or for sorting, isolating or purification of chondrocytes and/or muscle cells, or for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes, respectively, or of muscle or muscle cells during transplantation of muscle or muscle cells, respectively, or for studies of differentiation of chondrocytes or muscle cells.

Said process may be and *in vitro*, an *in situ* or an *in vivo* process.

- A process of using binding entities having the capability of binding specifically to binding sites of an integrin subunit $\alpha 11$ as defined above, or of an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β , or
- 5 to homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as markers or target molecules of cells or tissues expressing said integrin subunit $\alpha 11$, which cells or tissues are of animal including human origin.
- 10 Especially, said subunit β is $\beta 1$.

In embodiments of this process, said fragment is as defined above.

- In one embodiment, said process is a process for detecting the presence of an integrin subunit $\alpha 11$
- 15 comprising the amino acid sequence shown in SEQ ID NO. 1, or of an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β , or of homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein.

- 20 Furthermore, embodiments of this process encompass similar embodiments as defined above in connection with the process of using the integrin subunit $\alpha 11$ as a marker or target molecule.

- A process for detecting the presence of an integrin
- 25 subunit $\alpha 11$, or of a homologue or fragment of said integrin subunit, as defined above, on cells, whereby a polynucleotide or oligonucleotide chosen from the group comprising essentially a polynucleotide or oligonucleotide as shown in SEQ ID NO.1 is used as a marker under
- 30 hybridisation conditions, wherein said polynucleotide or oligonucleotide fails to hybridise to a DNA or RNA encoding an integrin subunit $\alpha 10$. Said cells may be chosen from the group comprising muscle cells.

- In embodiments of this process, said fragment is as
- 35 defined above.

Furthermore, embodiments of this process encompass similar embodiments as defined above in connection with

the process of using the integrin subunit $\alpha 11$ as a marker or target molecule.

- A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody
5 which is capable of using an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or a homologue or fragment of said integrin or subunit $\alpha 11$ wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as a
10 target molecule.

- A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of stimulating cell surface expression or activation of an integrin heterodimer comprising a subunit
15 $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit $\alpha 11$ wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein. In one embodiment, said composition is for use in stimulating, inhibiting or
20 blocking the formation of muscles, cartilage, bone or blood vessels.

- A vaccine comprising as an active ingredient at least one member of the group comprising an integrin heterodimer, which heterodimer comprises a subunit $\alpha 11$ and
25 a subunit β , or the subunit $\alpha 11$ thereof, and homologues or fragments of said integrin or subunit $\alpha 11$ wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, and a polynucleotide and an oligonucleotide coding for said integrin subunit $\alpha 11$.

30 - A method of gene therapy, whereby a vector comprising a polynucleotide or oligonucleotide coding for a subunit $\alpha 11$ of an integrin heterodimer, or for homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit
35 $\alpha 11$ protein, which polynucleotide or oligonucleotide comprises essentially the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof, and optionally a second vector

comprising a polynucleotide or oligonucleotide coding for a subunit β of said integrin heterodimer, is administered to a subject suffering from pathological conditions involving said subunit $\alpha 11$.

- 5 - A method of using binding entities having the capability of binding specifically to binding sites of a integrin subunit $\alpha 11$ comprising substantially the amino acid sequence shown in SEQ ID NO. 1, or of an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β , or
10 to homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, for promoting adhesion of cells.

- A method of using an integrin heterodimer comprising an integrin subunit $\alpha 11$ and a subunit β , or the
15 subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit $\alpha 11$ wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as a target for anti-adhesive drugs or molecules in tissues where adhesion impairs the function of the
20 tissue.

- A method of in vitro detecting the presence of integrin binding entities, comprising interaction of an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or
25 fragments of said integrin or subunit wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, with a sample, thereby causing said integrin, subunit $\alpha 11$, or homologue or fragments thereof, to modulate the binding to its natural
30 ligand or other integrin binding proteins present in said sample.

- A method of in vitro studying consequences of the interaction of a human heterodimer integrin comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof,
35 or homologues or fragments of said integrin or subunit wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, with an integrin binding

entity and thereby initiate a cellular reaction. In one embodiment of this method, the consequences of said interactions are measured as alterations in cellular functions.

- 5 - A method of using a polynucleotide or oligonucleotide encoding an integrin subunit $\alpha 11$ or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein as a target molecule.

- 10 One embodiment of this method comprises hybridising a polynucleotide or oligonucleotide to the DNA or RNA encoding the integrin subunit $\alpha 11$ or homologue or fragment thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which
- 15 polynucleotide or oligonucleotide fails to hybridise to a polynucleotide or oligonucleotide encoding an integrin subunit $\alpha 10$.

- A method of using binding entities having the capability of binding specifically to an integrin subunit $\alpha 10$ comprising the amino acid sequence shown in SEQ ID NO. 1 or SEQ ID NO. 2, or an integrin heterodimer comprising said subunit $\alpha 10$ and a subunit β , or to homologues or fragments thereof having similar biological activity, for
- 20 promoting adhesion of chondrocytes and/or osteoblasts to surfaces of implants to stimulate osseointegration.

- 25 - A method of using an integrin heterodimer comprising an integrin subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 10$ thereof, or homologues or fragments of said integrin or subunit $\alpha 10$, as a target for anti-adhesive
- 30 drugs or molecules in tendon, ligament, skeletal muscle or other tissues where adhesion impairs the function of the tissue.

- A method of stimulating, inhibiting or blocking the formation of cartilage or bone, comprising administration
- 35 to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or

the subunit $\alpha 1$ thereof, or homologues or fragments of said integrin or subunit $\alpha 1$ wherein the fragments include at least one binding site of the integrin subunit $\alpha 1$ protein, as a target molecule.

5

EXPERIMENTAL PROCEDURES

Cell cultures

10 The human fetal myoblast/myotube cultures were derived from clone G6 originating from a thigh muscle of a 73-day old aborted fetus ((39); referred to as G6 hereafter). Cultures of G6 and 2.5 years postnatal human satellite cells XXVI, a gift from Dr. Helen Blau (Stanford University, CA), were grown as reported earlier (39). Human rhabdomyosarcoma cell lines RD (ATCC No. CCL-136) and A204 (ATCC No. CRL-7900) were grown in DMEM 15 (Swedish Agricultural University, Uppsala) supplemented with 10% fetal calf serum.

RNA isolation and cDNA synthesis

20 Total RNA from G6 and XXVI myoblasts, the same cells differentiated for 3 or 7 days, and RD and A204 cell lines, was isolated using the RNeasy Midi kit (Qiagen)



according to the manufacturer's instructions. Poly-A RNA was extracted from total RNA of G6 and XXVI cells using Dynabeads mRNA DIRECT kit (DYNAL A.S., Norway).

PCR based cloning and generation of human $\alpha 11$ probes

- 5 First strand cDNA was generated from 1 μ g of G6 mRNA using a reverse transcription PCR-kit (Perkin-Elmer). Advantage cDNA Polymerase Mix (Clontech) was used in PCR amplifications using two different pairs of primers:
- (1) 5' ACG GGA GAC GTG TAC AAG TG 3' (forward), 5'-AAA
- 10 GTG CTG AAC CTC CAC CC-3' (reverse) and (2) 5'-CAC CAT CCA CCA GGC TAT GC -3' (forward), 5'-TTA GCG TTC CGT TAT AAA CA -3' (reverse). The PCR conditions were: 94°C, 4 min. ("hot start"); 94°C, 30 s; 55°C, 30 s; and 72°C, 1 min., for 25 cycles. Two products, named PCR1 and PCR2,
- 15 were obtained (figure 1), subcloned into the plasmid vector TA (Invitrogen), and sequenced. A single product of 1,4 kb in size, named PCR 3 (figure 1), was amplified using primers 1 (forward) and 2 (reverse), and human heart Marathon-Ready cDNA (Clontech) as template. Anneal-
- 20 ing temperatures in the applied touch-down program were: 68°C, 1 min., 5 cycles; 65°C, 1 min., 5 cycles; 60°C, 1 min., 25 cycles. Other steps were as described above. After the final cycle the reactions were extended for additional 7 min. at 72°C followed by a hold step at 4°C.
- 25 To obtain a sequence covering the 5' end, Rapid Amplification of cDNA Ends (RACE) was employed according to the manufacturer's instructions (Marathon cDNA Amplification kit, Clontech) using cDNA prepared from G6 mRNA and the gene specific antisense primer: 5'-CTT GGA GAA CCT GAA
- 30 GTT GGA GTT GAC -3'. Amplification was carried out applying the "touch-down" program (see above). To identify relevant products, 10 μ l of each RACE product was resolved on 1% agarose gel and subjected to Southern blot analysis as described previously (40). PCR2 (see above)
- 35 was labeled with [α -³²P]dCTP (Amersham Pharmacia Biotech, Sweden) using the RedyPrimeII DNA labeling system (Amersham Pharmacia Biotech, UK), and used as a hybridi-

16

zation probe. One specific signal was detected. Corresponding cDNA was purified (Gel Extraction kit, Quagen), cloned into the TA vector and sequenced (see figure 1).

Screening of cDNA libraries

5 A λ ZAP custom made G6 cDNA library (Stratagene, USA) was screened with PCR2 (see above) as a probe. The screening procedure (carried out as described in (40)) resulted in two clones representing the 5' non-coding region and the beginning of the coding part of integrin α 11 (figure 10 1). To obtain an additional sequence, a human uterus 5'-stretch λ gt11 cDNA library (Clontech) was screened with a mixture of PCR1 and PCR2 as probes. The probes were labeled with [α - 32 P]dCTP using the Ready-To-Go DNA labeling beads (Amersham Pharmacia Biotech, Sweden). Three clones 15 (1.1-1.3 in figure 1) representing parts of α 11 cDNA, were obtained. Rescreening of the human uterus 5'-stretch λ gt11 cDNA library with the probe λ 290 (corresponding to 2183-2473 in Fig. 1) yielded three more clones (2.1-2.3, figure 1) covering the rest of α 11 cDNA. Positive clones were 20 plaque purified, the phage DNA isolated using the Lambda Midi kit (Qiagen) and then sub-cloned into the Bluescript SK or pUC19 plasmid vectors before sequencing.

Northern hybridization

25 A filter containing 6 μ g of the poly-A RNA from G6 and XXVI cells and 10 μ g of the total RNA from RD and A204 cell lines, and a Human Multiple Tissue Northern Blot containing poly-A RNA from adult human tissues (Clontech), were hybridized at 68°C in ExpressHyb solution (Clontech) with probes labeled as described above. 30 The probes used were PCR1, PCR2, cDNA clone 1.3 (figure 1), 3RA (1.8 kb cDNA specific for human integrin α 1 mRNA, a generous gift from E.E. Marcantonio (Columbia University, New York), a 1.1 kb cDNA clone recognizing human G3PHD mRNA and a 1.8 kb cDNA clone recognizing human β - 35 actin (both from Clontech).

cDNA sequencing and sequence analysis

All PCR fragments and cDNA clones were sequenced on both strands either manually (29) or using ABI 310 Genetic Analyzer automatic sequencer. Sequences were analyzed with the aid of MacVector™ 6.0, DNA Star, Faktura™NEW 1.2.0, and Sequence Navigator 1.0.1 software programs. A distance tree of all I-domain containing integrin α subunits was assembled using SEAVIEW and PHYLO-WIN softwares (41). Percent similarity between every two members in the I-domain integrin subfamily was calculated by a formula $I=(1-D)\times 100$, where "I" is identity and "D" is distance.

Antibodies

A polyclonal antiserum (α 11 cyt) was produced against the peptide CRREPGLDPTPKVLE from the integrin α 11 cytoplasmic domain. Peptide synthesis and conjugation to Keyhole limpet hemocyanin, immunization of rabbits and affinity purification was performed at Innovagen AB (Lund, Sweden). The monoclonal antibody Mab 13 against integrin β 1 was obtained from S.K. Akiyama (NIEHS, NIH). Monoclonal antibodies to integrin α 1 (clone FB12, sold as MAB 1973) and integrin α 2 (clone BHA2.1 sold as Mab 1998) were both obtained from Chemicon, Temecula, CA. The monoclonal antibody to vinculin (clone hVIN-1) was from Sigma (Saint Louis, MO, USA). Secondary fluorescent antibodies (CY3™-coupled goat-anti rabbit IgG and FITC-coupled goat anti-mouse IgG of multiple labeling grade) were from Jackson ImmunoResearch Laboratories, Inc. (West Grove, PA, USA).

Immunoprecipitation and SDS-PAGE

G6 and XXVI cells were labeled with [35 S] cysteine/methionine and subjected to immunoprecipitation and SDS-PAGE as reported previously (38). The two-step procedure used to dissociate integrin heterodimers was carried out as follows. After incubation of samples with β 1 antibody and capture with GammaBind G Sepharose (Amersham Pharmacia Biotech, Uppsala, Sweden), 100 μ l of 1% SDS was

18

added to the washed beads which were then boiled for 5 minutes. 10 mM Tris-HCl, pH 7.4, 0.15 M NaCl and 1% Triton X-100 was added to a final volume of 1ml and the lysate was incubated with GammaBind G Sepharose for 1 hour. The incubation with GammaBind G was performed in order to ensure that no reactive $\beta 1$ antibodies remained. After removal of GammaBind G Sepharose, $\alpha 11$ integrin antibody was added for additional 2 hours, followed by capture with protein A Sepharose (Amersham Pharmacia Biotech) and boiling in SDS-PAGE sample buffer.

Chromosomal localization

Chromosomal localization of the human integrin $\alpha 11$ was performed by using a combination of FISH (Fluorescent In Situ Hybridization) technique and DAPI (4',6-diamidino-2-phenylindole) banding essentially as described earlier (42). As a hybridization probe, the 1.4 kb RT-PCR product PCR3 was used.

Surface iodination and affinity chromatography

Cultured XXVI cells were surface iodinated as described (38). Labeled cells were solubilized in 1 ml of solubilization buffer (10 mM Tris-HCl pH 7.4, 15 mM NaCl, 1% Triton X-100, 1mM $MgCl_2$, 1 mM $CaCl_2$, 1mM $MnCl_2$), centrifuged at 14000 g for 20 min., and soluble membrane proteins were applied to a collagen type I Sepharose (bovine collagen type I from Vitrogen (Collagen Corp., Palo Alto) coupled to CNBr-activated Sepharose CL-4B at 3 mg/ml gel as described (14)), equilibrated in solubilization buffer. Following a one hour incubation the column was washed extensively with buffer A (10 mM Tris-HCl pH 7.4, 50 mM NaCl, 1 mM $MnCl_2$, 0.1% Triton X-100) and by 10 column volumes of buffer A without NaCl. Bound proteins were eluted with 20 mM EDTA, 10 mM Tris-HCl pH 7.4, 0.1% Triton X-100. Peak fractions were pooled and concentrated by immunoprecipitation with $\beta 1$ integrin and $\alpha 11$ integrin antibodies as described under Immunoprecipitation and SDS-PAGE. Eluted fractions and captured

19

proteins were analyzed on 7.5% SDS-PAGE gels followed by autoradiography.

Indirect immunofluorescence

Cells cultured on coverslips were washed in serum-free medium and fixed for 8 min. in acetone at -20°C. Non-specific binding sites were blocked by incubating with 10% goat serum diluted in phosphate buffered saline. In the double immunofluorescence staining protocol, primary antibodies (anti- α 11 cyt (rabbit antibody) and anti-vinculin (mouse antibody)) were simultaneously incubated with fixed cells for 1.5 hours at +37°C. Specifically bound antibodies were detected using anti-rabbit Cy3 IgG and anti-mouse FITC IgG. Stained cells were mounted in Vectashield™ mounting medium (Vector Laboratories, Inc., Burlingame, CA, USA) and visualized and photographed under a Zeiss light microscope equipped with optics for observing fluorescence.

RESULTS AND DISCUSSION

20 cDNA cloning of a novel integrin α -chain

In order to determine the nature of the integrin chain that we had previously characterized on human fetal muscle cells and named α mt (38), a number of approaches were used. Applying PCR with mRNA from fetal muscle cells as template together with degenerate primers to conserved regions of integrin α subunits (43) we amplified cDNA for α 1, α 4, α 5, α 6 and α v integrin chains (data not shown), but failed to amplify the novel integrin. However, while searching through the literature we came across two integrin sequences obtained in a subtractive hybridization protocol comparing human primary myoblasts with the rhabdomyosarcoma cell line RD (44). After having confirmed that these sequences could be amplified by PCR from human fetal G6 myoblast cDNA, PCR was performed assuming that these sequences were derived from the same transcript. In this manner a 1.4 kb cDNA fragment with integrin-like sequence was obtained. Screening of a human fetal myo-

20

blast cDNA library and 5' RACE yielded additional 5' sequence. We determined the mRNA expression pattern in a number of human tissues (see below) and observed a high mRNA expression in the uterus. Screening of a uterus cDNA library resulted in the identification of the complete open reading frame. A schematic illustration of the cloning strategy is shown in figure 1.

cDNA sequence and predicted amino acid sequence of $\alpha 1$ integrin chain

10 By sequence analysis of cDNA clones and 5' RACE products we obtained a continuous sequence of 3983 nucleotides (nt) composed of 90 nt 5' non-coding sequence, 3564 nt open reading frame, and 326 nt 3' non-coding sequence. Translation of the sequence predicts an
15 integrin α -chain like precursor of 1188 amino acids including a 22 amino acid long signal peptide (fig. 2, GenBank accession No. AF137378). The mature 1166 amino acid long peptide is larger than any other currently identified integrin α -chain (the closest being αE , composed of 1160 amino acids (45). The 1115 amino acid long
20 predicted extracellular domain contains 7 FG-GAP repeats in the amino-terminal end with an inserted I-domain between repeats 2 and 3. The I-domain consists of 195 amino acids and includes a conserved MIDAS motif. In
25 addition to the metal chelating site in the I-domain, three additional potential divalent cation binding motifs with the consensus sequence DXD/NXDXXXD are present in repeats 5-7. A total of 20 cysteines are located in the extracellular domain. Of these, 16 are conserved in the
30 most closely related integrin $\alpha 10$ and $\alpha 1$ chains and they may contribute to intramolecular disulphide bonds. The two non-conserved cysteines found at positions Cys 606 and Cys 988 most likely represent free unpaired cysteines while the two non-conserved cysteines Cys 806 and Cys 817
35 may pair to form a disulphide bond. Mapping of the cysteines in the suggested β -propeller structure shows that the first three disulphide bonds are likely to stabilize

blades one and two of the β -propeller whereas the remaining bonds are found outside the propeller region, in the stalk region towards the transmembrane domain. 16 potential N-glycosylation sites are present in $\alpha 11$. A search for sequence motifs reveals the presence of a 22 amino acid leucine zipper motif starting at position 951, and a 17 amino acid sequence starting at position 1082, which is similar to sequences found in G-protein coupled receptors. These sequences might represent functional domains of importance for protein-protein interactions.

The transmembrane region (amino acids 1142-1164) is 23 amino acids long and is followed by a cytoplasmic tail of 24 amino acids. The cytoplasmic tail contains the sequence GFFRS instead of the conserved GFFKR sequence, found in all other α -chains except $\alpha 8$ - $\alpha 10$. It will be interesting to determine the importance of this sequence in defining the cytoplasmic domain as well as its possible ability to bind calreticulin and other intracellular components.

Comparison of integrin $\alpha 11$ chain with other integrin α chains

Alignment of the predicted $\alpha 11$ integrin amino acid sequence with other integrin sequences shows the highest overall identity with $\alpha 10$ (42% identity), $\alpha 1$ (37% identity), and $\alpha 2$ (35% identity), followed by the remaining I-domain containing integrin subunits. Of the non I-domain containing integrins, $\alpha 4$ and $\alpha 9$ are the most similar to $\alpha 11$. A distance tree shows that $\alpha 10$ and $\alpha 11$ form a separate branch from the most closely related $\alpha 1$ and $\alpha 2$ integrin chains (fig. 3). The similarity with other integrins is particularly high in the N-terminal β -propeller part but lower in the stalk region. Comparison of $\alpha 1$ integrin with $\alpha 2$ integrin has pointed to the presence of a 38-residue insert in the β -propeller region of $\alpha 1$ integrin chain (15). Like $\alpha 1$ chain, $\alpha 11$ also contains inserted amino acids not present in the other I-domain containing integrin chains. however, in the $\alpha 11$

chain these are found within the stalk region at amino acids 804-826. The exact border of the predicted insertion varies depending on the alignment method and the parameters chosen, but is predicted to span at least 22 amino acids. The insert shows no significant similarity to other integrin sequences and contains two cysteines likely to form a disulphide bond (see fig. 2). We do not believe that the predicted inserted sequence represents a cloning artifact since it is present in three independently analysed clones. Other examples of non I-domain inserted sequences are found in the *Drosophila* α PS2 chain, where developmentally regulated splicing in the ligand binding region modulates ligand affinity (46). In α 7 integrin chain, splicing in the extracellular domain between predicted blades 2 and 3 in the β -propeller generates X1 and X2 variants, affecting the binding to laminin-1 in a cell-specific manner (47). In the more closely related α 1 integrin chain the 38 extra amino acids are present in a position that is predicted to be in the beginning of the sixth blade of the 7-bladed propeller. So far there is no evidence that the extra amino acids in either α 1 or α 11 arise by alternative splicing. In α 11 the predicted inserted region is outside the β -propeller and most likely does not directly affect ligand binding, but might instead be involved in modifying protein-protein interactions or be important for outside-in or inside-out signalling. In this regard it is interesting to note that tetraspan proteins by binding to the stalk region of certain integrin α -chains can recruit PI-4 kinase and protein kinase C to integrin complexes (48). Likewise the extracellular membrane-proximal parts of certain integrin α -chains have been shown to be involved in Shc-mediated integrin signalling (49).

Analysis of sequences identified during screening for genes upregulated during tadpole regression revealed a partial sequence, which at the time was reported to show the highest similarity to integrin α 1 (41% identity)

(50). This sequence, when translated (amino acids 1-116), shows 71% identity to human $\alpha 11$ and thus most likely represents the *Xenopus* orthologue of $\alpha 11$ rather than that of the $\alpha 1$. These data suggest that $\alpha 11$ is well conserved during evolution.

Chromosomal localization of the integrin $\alpha 11$ gene

A fluorescent cDNA probe was used for in situ hybridization on metaphase chromosome spreads. The analysis shows that the integrin $\alpha 11$ gene (ITGA11) is located on chromosome 15q23 (fig. 4). The genes for I-domain containing integrins $\alpha 1$ and $\alpha 2$ are both present on chromosome 5 (51,52), just as the genes for the closely related $\beta 2$ integrin associated α -chains all map to chromosome 16 (53). Interestingly, the $\alpha 11$ gene and the closely related $\alpha 1$ and $\alpha 2$ genes, map to different chromosomes. It will be of evolutionary interest to determine the chromosomal localization of the integrin $\alpha 10$ gene. Curiously, a form of Bardet-Biedl syndrome characterized by retinitis pigmentosa, polydactyly, obesity, hypogenitalism, mental retardation, and renal anomalies maps to 15q22-23 (54). Future studies will clarify a possible linkage of ITGA11 to Bardet-Biedl syndrome.

Expression pattern of $\alpha 11$ mRNA in adult tissues

Northern blot analysis of mRNA from various adult human tissues shows the highest level of expression of $\alpha 11$ in adult human uterus. A strong signal is also noted in heart, while intermediate levels of $\alpha 11$ mRNA are present in skeletal muscle and intermediate to low levels in other adult tissues tested (fig. 5 and data not shown). For a comparison, the same blot was probed for the closely related $\alpha 1$ integrin mRNA (fig. 5). A striking difference in the expression levels of $\alpha 1$ and $\alpha 11$ was observed in the smooth muscle rich uterus, which appears to lack $\alpha 1$. Immunohistochemical analysis and in situ hybridizations will elucidate the detailed distribution of $\alpha 11$ protein and mRNA in muscle and other tissues. Neither $\alpha 1$ (33) nor $\alpha 2$ (55) are present in muscle fibers, and the

distribution of $\alpha 10$ in skeletal muscle tissues is not known (5). Hence, no I-domain containing integrin has so far been reported to be expressed in the skeletal muscle sarcolemma. Recently the gene for $\alpha 1$ integrin was inactivated in mice, resulting in mice with an apparently normal phenotype (56). More careful analysis revealed a phenotype characterized by a hypocellular skin (57) and aberrant regulation of collagen synthesis (58). It will be interesting to compare sites of overlapping expression between $\alpha 1$, $\alpha 2$ and $\alpha 10$ integrins, and use reagents to $\alpha 10$ and $\alpha 11$ to examine possible functional compensatory mechanisms in $\alpha 1$ integrin-deficient mice.

Biochemical characterization of $\alpha 11$ protein

Following the cloning of the full-length $\alpha 11$ integrin cDNA it was essential to determine if the predicted amino acid sequence was identical to the novel uncleaved $\beta 1$ integrin-associated α -chain that we had previously noted to be upregulated during in vitro differentiation of human myoblasts (38). To answer this question we raised antibodies to the cytoplasmic tail of the integrin $\alpha 11$ chain. Immunoprecipitation from the human satellite cells showed that the antibodies precipitated a 145 kDa $\alpha 11$ band associated with a 115 kDa $\beta 1$ band (fig. 6, panel A) in SDS-PAGE under non-reducing conditions. Under reducing conditions the $\alpha 11$ band migrated as 155 kDa (see fig. 6, panel B). From the translated amino acid sequence an Mr of 133 400 is predicted for the $\alpha 11$ chain. Taking the 16 potential glycosylation sites into account this fits well with the observed 155 kDa band in SDS-PAGE. Under non-reducing conditions the 145 kDa band is distinctly larger than $\alpha 2$ (fig. 6, panel A) and $\alpha 10$ integrin chains which co-migrate as 140 kDa bands and $\alpha 11$ migrates well below the 180 kDa integrin $\alpha 1$ band. The $\alpha 2$ (59) and $\alpha 10$ (5) chains both contain 10 potential glycosylation sites whereas $\alpha 1$ contains 26 glycosylation sites (60). The intermediate size of $\alpha 11$ in SDS-PAGE compared with $\alpha 1$

and $\alpha 2/\alpha 10$ is thus most likely a result of differential glycosylation.

To show that $\alpha 11$ is associated with the $\beta 1$ subunit a two-step immunoprecipitation procedure was performed.

- 5 Integrins were first precipitated with a monoclonal anti- $\beta 1$ integrin antibody and GammaBind G captured integrins were then dissociated by boiling in 1% SDS. In the second step, SDS was diluted tenfold and antibodies to $\alpha 11$ were added. As shown in fig. 6 panel A antibodies to $\alpha 11$ immunoprecipitate only the 145 kDa band from the dissociated precipitate initially captured with $\beta 1$ antibodies.

10 Induction of $\alpha 11$ mRNA and protein during myogenic differentiation in vitro

- It has previously been determined that αmt is the major integrin α -chain that is up-regulated during myogenic differentiation on human fetal myoblasts in vitro (38). To compare $\alpha 11$ levels in myoblasts and myotubes, immuno-precipitates were analyzed from myoblast cultures in pro-liferation medium, and from parallel cultures allowed to differentiate and form myotubes in differentiation medium for 7 days. Immunoprecipitation with both $\beta 1$ and $\alpha 11$ antibodies showed that $\alpha 11$, like αmt , is strongly up-regulated at the protein level in differentiation cultures of human fetal muscle cells and satellite cells (fig. 6, panel B). To determine if the up-regulation occurs at the mRNA or protein level we analyzed $\alpha 11$ mRNA from different differentiation stages (day 1, day 3 and day 7) (fig. 6, panel C). Already at day 3 in differentiation medium a strong up-regulation of $\alpha 11$ mRNA was noted, establishing that the up-regulation of $\alpha 11$ integrin protein occurs as a result of increased transcription or mRNA stability. Based on similar SDS-PAGE migration patterns, similar behavior under reducing conditions, association with $\beta 1$ integrin chain, and up-regulation during in vitro differentiation of human fetal myoblasts, the present data show that $\alpha 11$ integrin is identical with αmt .

Analysis of mRNA from the two rhabdomyosarcoma cell lines RD and A204 (fig. 6, panel C) did not provide evidence for the presence of $\alpha 11$ in either cell line. Based on the observed up-regulation of $\alpha 11\beta 1$ in human fetal muscle cells and the presence of $\alpha 11$ message in adult muscle we suggest that $\alpha 11$ integrin might be involved in early steps of muscle formation and that it in adult muscle tissues may fulfill a stabilizing role. The $\alpha 7$ integrin subunit is a major $\beta 1$ -associated integrin chain in muscle, but genetic deletion of $\alpha 7$ leads to a fairly mild muscular dystrophy (30).

Ligand binding specificity of $\alpha 11\beta 1$ integrin

So far identified I-domain containing integrins of the $\beta 1$ integrin subfamily all bind collagens (5,15,59). For $\alpha 1$ and $\alpha 2$ this binding capacity has been shown to reside within the I-domain (17,18). To determine if $\alpha 11\beta 1$ also binds collagen we performed collagen type I Sepharose chromatography of membrane proteins from surface-iodinated XXVI satellite cells. Direct analysis of the EDTA eluate revealed weak bands corresponding to the positions of $\alpha 1$, $\alpha 2$, $\alpha 11$ and $\beta 1$ in parallel immunoprecipitations (fig. 7, panel 1). The EDTA eluate was concentrated by immunoprecipitation with $\beta 1$ and $\alpha 11$ antibodies. As shown in figure 7, a prominent $\alpha 11$ band is present in the collagen I Sepharose eluate. The relatively weak $\beta 1$ band in the proteins captured with $\alpha 11$ antibodies indicates that the $\alpha 11\beta 1$ heterodimer partly dissociates in the presence of EDTA. To visualize the interaction of $\alpha 11\beta 1$ integrin with collagen I in intact cells, myogenic cells expressing $\alpha 11\beta 1$ were trypsinized and plated on collagen and fibronectin for 1 hour. The ability to form focal contacts was investigated by double immunofluorescence staining for $\alpha 11$ -chain and vinculin. As seen in panel 2 of figure 7 $\alpha 11$ localizes to vinculin positive focal contacts on collagen but not on fibronectin. Binding studies with $\alpha 11$ I-domain expressed as a bacterial GST-fusion protein also confirmed a specific

affinity for collagen I (unpublished M. Höök, R. Rich, R. Owens). Stable transfections of $\alpha 11$ cDNA into cells with various integrin backgrounds will allow a more detailed study of $\alpha 11\beta 1$ interactions with different collagen, and possibly also laminin, isoforms. Combined with in vivo distribution studies of $\alpha 11\beta 1$ this is likely to yield valuable information regarding the in vivo ligands for $\alpha 11\beta 1$ in different tissues.

$\alpha 11$ integrin protein distribution in human embryo

Morphologically normal human embryos (aged from 4 to 8 post-ovulatory weeks) were obtained from legal abortions induced by Mifepristone (RU486) at Hopital Broussais in Paris. All procedures were approved by the Ethical Committee of Saint-Vincent de Paul Hospital in Paris.

Each sample was first examined macroscopically during dissection under a stereo-microscope. The development stage of the embryos was determined using established criteria. Tissues were collected shortly after delivery and frozen within the first 24 h post mortem on dry ice and stored at -80°C until used. Seven micron-thick cryostat sections were mounted on slides previously coated with a 2% 3-aminopropyl-triethoxysilane solution in acetone. The cryosection was left unfixed prior to blocking of non-specific binding sites with 10% goat serum diluted in phosphate buffered saline. For immunofluorescence, the section was incubated with $\alpha 11$ antibodies 1.5 h at $+37^{\circ}\text{C}$. Specifically bound antibodies were detected using goat anti-rabbit Cy3 IgG (Jackson Immunoresearch). The stained tissue section was mounted in Vectashield™ mounting medium (Vector Laboratories Inc.) and visualized and photographed under a Zeiss light microscope equipped with optics for observing fluorescence.

The results obtained are shown in figure 8. High levels of $\alpha 11$ protein were noted around vertebrae (arrows), in intervertebrae disc (asterisks), around ribs

(thin arrows) and around forming cartilage in the
forelimb (arrowhead).

References

1. Hynes, R.O. (1992) *Cell* 69, 11-25
2. Adams, J., and Watt, F.M. (1993) *Development* 117,
5 1183-1198
3. Sastry, S.K., and Horwitz, A.F. (1996) *Dev. Biol.*
180, 455-467
4. Gullberg, D., Tiger, C.-F., Lohikangas, L., and
Velling, T. (1998) *Front. Biosci.* 3, 1028-1039
- 10 5. Camper, L., Hellman, U., and Lundgren-Akerlund, E.
(1998) *J. Biol. Chem.* 273, 20383-20389
6. Aumailley, M., Gerl, M., Sonnenberg, A., Deutzmann,
R., and Timpl, R. (1990) *FEBS Lett* 262, 82-86
7. Pfaff, M., Aumailly, M., Specks, U., Knolle, J.,
15 Zerwes, H.G., and Timpl, R. (1993) *Exp. Cell Res.*
206, 167-176
8. Davis, G.E. (1992) *Biochem. Biophys. Res. Commun.*
182, 1025-1031
9. Goodman, S.L., Aumailly, M., and von der Mark, H.
20 (1991) *J. Cell Biol.* 113, 931-941
10. Dickeson, S.K., and Santoro, S.A. (1998) *Cell. Mol.*
Life Sci. 54, 556-566
11. Takada, Y., Kamata, T., Irie, A., Puzon-McLaughlin,
W., and Zhang, X.P. (1997) *Matrix Biol.* 16, 143-151
- 25 12. Colombatti, A., Bonaldo, P., and Doliana, R. (1993)
Matrix 13, 297-306
13. Lee, J.-O., Rieu, P., Arnaout, M.A., and Liddington,
R.C. (1995) *Cell* 80, 631-635
14. Gullberg, D., Turner, D.C., Borg, T.K., Terracio,
30 L., and Rubin, K. (1990) *Exp. Cell Res.* 190, 254-264
15. Ignatius, M.J., Large, T.H., Houde, M., Tawil, J.W.,
Barton, A., Esch, F., Carbonetto, S., and Reichardt,
L.F. (1990) *J. Cell Biol.* 111, 709-720
16. Dickeson, S.K., Walsh, J.J., and Santoro, S.A.
35 (1997) *J. Biol. Chem.* 272, 7661-7668

17. Calderwood, D.A., Tuckwell, D.S., Eble, J., Kühn, K., and Humphries, M.J. (1997) *J. Biol. Chem.* 272, 12311-12317
18. Tuckwell, D., Calderwood, D.A., Green, L.J., and
5 Humphries, M.J. (1995) *J. Cell Sci.* 108, 1629-1637
19. Tuckwell, D.S., Reid, K.B.M., Barnes, M.J., and Humphries, M.J. (1996) *Eur. J. Biochem.* 241, 732-739
20. King, S.L., Kamata, T., Cunningham, J.A., Emsley, J., Liddington, R.C., Takada, Y., and Bergelson, J.M. (1997) *J. Biol. Chem.* 272, 28518-28522
- 10 21. Springer, T.A. (1997) *Proc. Natl. Acad. Sci. USA* 94, 65-72
22. Huang, C., and Springer, T.A. (1997) *Proc. Natl. Acad. Sci. USA* 94, 3162-3167
- 15 23. Oxvig, C., and Springer, T.A. (1998) *Proc. Natl. Acad. Sci. USA* 95, 4870-4875
24. Huang, C., Lu, C., and Springer, T.A. (1997) *Proc. Natl. Acad. Sci. USA* 94, 3156-3161
25. Rojiani, M.V., Finlay, B.B., Gray, V., and Dedhar, S. (1991) *Biochemistry* 30, 9859-9866
- 20 26. Wright, T.F. (1960) *J. Exp. Zool.* 143, 77-99
27. MacKrell, A.J., Blumberg, B., Haynes, S.R., and Fessler, J.H. (1988) *Proc. Natl. Acad. Sci. USA* 85, 2633-2637
- 25 28. Bloor, J.A., and Brown, N.H. (1998) *Genetics* 148, 1127-1142
29. Gullberg, D., Fessler, L.I., and Fessler, J.H. (1994) *Dev. Dynamics* 199, 116-128
30. Mayer, U., Saher, G., Fassler, R., Bornemann, A., Echtermeyer, F., von der Mark, H., Miosge, N., Poschl, E., and von der Mark, K. (1997) *Nat. Genet.* 17, 318-323
- 30 31. Hayashi, Y.K., Chou, F.-L., Engvall, E., Ogawa, M., Matsuda, C., Hirabayashi, S., Yokochi, K., Ziober, B.L., Kramer, R.H., Kaufman, S.J., Ozawa, E., Goto, Y., Nonaka, I., Tsukahara, T., Wang, J.-Z., Hoffman, E.P., and Arahata, K. (1998) *Nat. Genet.* 19, 94-97
- 35

32. Song, W.K., Wang, W., Foster, R.F., Biesler, D.A.,
and Kaufman, S.J. (1992) *J. Cell Biol.* 117, 643-657
33. Martin, P.T., Kaufman, S.J., Kramer, R.H., and
Sanes, J.R. (1996) *Dev. Biol.* 174, 125-139
- 5 34. Belkin, A., Zhidkova, N.I., Balzac, F., Altruda, F.,
Tomatis, D., Maier, A., Tarone, G., Koteliensky,
V.E., and Burrridge, K. (1996) *J. Cell Biol.* 132,
211-226
35. Miyagoe, Y., Hanaoka, K., Nonaka, I., Hayasaka, M.,
10 Nabeshima, Y., Arahata, K., Nabeshima, Y., and
Takeda, S. (1997) *FEBS Lett.* 415, 33-39
36. Xu, H., Christmas, P., Wu, X.R., Wewer, U.M., and
Engvall, E. (1994) *Prox. Natl. Acad. Sci. USA* 91,
5572-5576
- 15 37. Sunada, Y., Bernier, S.M., Kozak, C.A., Yamada, Y.,
and Campbell, K.P. (1994) *J. Biol. Chem.* 269, 13729-
13732
38. Gullberg, D., Velling, T., Sjöberg, G., and
Sejersén, T. (1995) *Dev. Dyn.* 204, 57-65
- 20 39. Jin, P., Farmer, K., Ringertz, N.R., and Sejersén,
T. (1993) *Differentiation* 54, 47-54
40. Sambrook, J., Fritsch, E., and Maniatis, T. (1989)
Molecular cloning: A Laboratory Manual, Second
Edition Ed., Cold Spring Laboratory Press, NY
- 25 41. Galtier, N., Gouy, M., and Gautier, C., (1996)
Comput. Appl. Biosci. 12, 543-548
42. Heng, H., Squire, J., and Tsui, L.-C (1992) *Proc.*
Natl. Acad. Sci. USA 89, 9509-9513
43. Pytela, R., Suzuki, S., Breuss, J., Erle, D.J., and
30 Sheppard, D. (1994) *Methods Enzymol.* 245, 420-451
44. Genini, M., Schwalbe, P., Scholl, F.A., and Schafer,
B.W. (1996) *Int. J. Cancer* 66, 571-577
45. Shaw, S.K., Cepek, K.L., Murphy, E.A., Russel, G.J.,
Brenner, M.B., and Parker, C.M. (1994) *J. Biol.*
35 *Chem.* 269, 6016-6025
46. Fogerty, F.J., Fessler, L.I., Bunch, T.A., Yaron,
Y., Parker, C.G., Nelson, R.E., Brower, D.L.,

- Gullberg, D., and Fessler, J.H. (1994) *Development* 120, 1747-1758
47. Ziober, B.L., Chen, Y., and Kramer, R.H. (1997) *Mol. Biol. of the Cell* 8, 1723-1734
- 5 48. Hemler, M.E. (1998) *Curr. Opin. Cell Biol.* 10, 578-585
49. Wary, K.K., Mainiero, F., Isakoff, S.J., Marcantonio, E.E., and Giancotto, F.G. (1996) *Cell* 87, 733-743
- 10 50. Brown, D.D., Wang, Z., Furlow, J.D., Kanamori, A., Schwartzman, R.A., Remo, B.F., and Pinder, A. (1996) *Proc. Natl. Acad. Sci. USA* 93, 1924-1929
51. Jaspers, M., Marynen, P., Aly, M.S., Cuppens, H., Hilliker, C., and Cassiman, J.J. (1991) *Somat. Cell Mol. Genet.* 17, 505-511
- 15 52. Seymour, A.B., Yanak, B.L., O'Brien, E.P., Rusiniak, M.E., Novak, E.K., Pinto, L.H., Swank, R.T., and Gorin, M.B. (1996) *Genome Res.* 6, 538-544
53. Wong, D.A., Davis, E.M., Le Beau, M., and Springer, T.A. (1996) *Gene* 171, 291-294
- 20 54. Bruford, E.A., Riise, R., Teague, P.W., Porter, K., Thomson, K.L., Moore, A.T., Jay, M., Warburg, M., Schinzel, A., Tommerup, N., Tornqvist, K., Rosenberg, T., Patton, M., Mansfield, D.C., and Wright, A.F. (1997) *Genomics*, 41, 93-99
- 25 55. Wu, J.E., and Santoro, S.A. (1994) *Dev. Dyn.* 199 (292-314)
56. Gardner, H., Kreidberg, J., Koteliensky, V., and Jaenisch, R. (1996) *Dev. Biol.* 175, 301-313
- 30 57. Pozzi, A., Wary, K.K., Giancotti, F.G., and Gardner, H.A. (1998) *J. Cell Biol.* 142, 587-594
58. Gardner, H., Broberg, A., Pozzi, A., Laato, M., and Heino, J. (1999) *J. Cell Sci.* 112, 263-272
59. Takada, Y., and Hemler, M.E. (1989) *J. Cell Biol.* 109, 397-407
- 35 60. Briesewitz, R., Epstein, M.R., and Marcantonio, E.E. (1993) *J. Biol. Chem.* 268, 2989-2996

FIGURE LEGENDS

Figure 1. Schematic organization of PCR fragments and cDNA clones representing different parts of the full length sequence of integrin $\alpha 11$ subunit

5 A. Clones 1.1-1.3 and 2.1-2.3 are from the first and the second round of screening, respectively. Fragment 0.0 represents a 5' RACE product as well as a clone obtained from screening of the G6 library. PCR fragments 1-3 and a
10 SacI fragment of a clone 1.3, λ 290, are marked with thick lines. Names and positions of all the clones on a scheme are shown in tabulated form in B.

B. Names of the PCR-amplified fragments and cDNA clones shown in A are in the left column, and their positions in
15 the full length cDNA of integrin $\alpha 11$ in the right column.

Figure 2. Nucleotide and deduced amino acid sequence of the human integrin $\alpha 11$ chain

The putative signal peptide is underlined in bold, I-domain is boxed, potential N-linked glycosylation sites
20 are marked with asterisks, cysteines are underlined, potential divalent cation binding motifs are double underlined and the transmembrane domain is underlined with dashes. A 22 amino acid insert is boxed in bold.

Figure 3. A distance tree of the I-domain containing α -integrin subfamily members

A tree was assembled based using ClustalW multiple alignment - based SEAVIEW and PHYLOWIN softwares. A scale at the bottom shows percent identity.

Figure 4. Chromosome mapping of ITGA11 gene by fluorescent in situ hybridization (FISH)

30 A. Left panel shows the FISH signals on human chromosome 15; right panel shows the same mitotic figure stained with 4',6-diamino-2-phenylindole to identify human chromosome 15.

35 B. Diagram of FISH mapping result for the probe PCR3 based on a detailed analyses of 10 different images. Each

dot represents the double FISH signals detected on human chromosome 15.

Figure 5. Expression of integrin $\alpha 11$ and $\alpha 1$ subunit mRNAs in adult human tissues

- 5 Integrin $\alpha 11$ mRNA and integrin $\alpha 1$ mRNA were analyzed on a membrane with RNA from various adult human tissues where mRNA loading was normalized with respect to β -actin. Probes used for hybridizations are marked on the left. Size of molecular weight standard is marked to the right.
- 10 Note that the β -actin probe reacts with 2 kb β/γ actin transcripts and the muscle specific 1.8 kb α -actin message.

Figure 6. Biochemical characterization of integrin $\alpha 11$ chain and upregulation of corresponding protein and mRNA in myogenic cells

- 15 A. $\alpha 11$ associates with $\beta 1$ integrin chain. Human XXVI and G6 muscle cells were metabolically labeled with [35 S] cysteine/methionine and integrins were immunoprecipitated with the indicated antibodies ($\beta 1$, $\alpha 2$ and $\alpha 11$). Evidence
- 20 for the association of integrin $\alpha 11$ with the $\beta 1$ subunit obtained by treating proteins precipitated with anti- $\beta 1$ antibodies with SDS followed by a second precipitation with $\alpha 11$ antibodies (ant- $\alpha 11$ +SDS). Precipitated proteins were resolved on 7.5% SDS-PAGE gels in the absence of
- 25 reducing agents, followed by fluorography.
- B. Induction of integrin $\alpha 11$ upon myogenic differentiation in vitro.

- G6 muscle cells were metabolically labeled with [35 S] cysteine/methionine when growing in proliferation medium
- 30 (mb-proliferating myoblasts) and after 7 days in differentiation medium) (mt-myotubes). Integrins were precipitated with antibodies to $\beta 1$ and $\alpha 11$ and the precipitates were resolved on 7.5% SDS-PAGE gels both under non-reducing (UNREDUCED) and reducing (REDUCED) conditions.
- 35 Lanes 1, 3, 5 and 7 are immunoprecipitations with the antibody to integrin $\beta 1$, and lanes 2, 4, 6 and 8 with the antibody to integrin $\alpha 11$.

C. Upregulation of integrin $\alpha 11$ mRNA in differentiated myogenic cells.

mRNA was extracted from G6 and XXVI cells growing under proliferating (p) or differentiating (d) conditions for 3 days (d3) or 7 days (d7). Total RNA was isolated from RD and A204 cells. Following separation of RNA on agarose gel and transfer to the membrane, the filter was hybridized with probes to $\alpha 11$ integrin ($\alpha 11$) and glyceraldehyde-3-phosphate dehydrogenase (GAPDH). Size of bands in RNA standard (in kb) are marked to the right.

Figure 7. Ligand binding properties of $\alpha 11\beta 1$ integrin panel 1: Collagen binding integrins on XXVI cells.

XXVI cells were surface iodinated and integrins were analyzed by immunoprecipitation and collagen I Sepharose affinity chromatography. Immunoprecipitation reveals the presence of $\beta 1$ integrins (lane 1), $\alpha 1\beta 1$ (lane 2), $\alpha 11\beta 1$ (lane 3) and $\alpha 2\beta 1$ (lane 4) at the surface of XXVI cells. EDTA eluted proteins bound to collagen I Sepharose contain weak band in the position of $\alpha 1$, $\alpha 11$, $\alpha 2$ and $\beta 1$ integrin chains (lane 5). Immunoprecipitations with $\beta 1$ integrin antibodies (lane 6) and $\alpha 11$ integrin antibodies (lane 7) confirm the presence of $\alpha 11$ and $\beta 1$ in the EDTA eluate.

panel 2: $\alpha 11\beta 1$ localizes to focal contacts on collagen.

Indirect immunofluorescent visualization of vinculin (A, B) and $\alpha 11$ integrin chain (C, D) in human XXVI satellite cells seeded on collagen type I (A and C) and fibronectin (B and D). Note the localization of integrin $\alpha 11$ chain to focal contacts of cells allowed to attach to collagen and its complete absence on cells seeded on fibronectin. Vinculin is found in focal contacts on both substrates. A and C show the same cell double stained for both antigens. Scale bar is 20 μ m.

Figure 8. α 11 integrin protein distribution at 8 weeks of gestation.

- Composite of immunohistochemical staining of sagittal section of human embryo at 8 weeks of gestation. Note
- 5 high levels of α 11 protein around vetrebrae (arrows), in intervertebral disc (asterisks), around ribs (thin arrows) and around forming cartilage in the forelimb (arrowhead).

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

It is to be understood that a reference herein to a prior art document does not constitute an admission that the document forms part of the common general knowledge in the art in Australia or in any other country.

B
B
B
B
B

B
B
B
B
B

EDITORIAL NOTE

APPLICATION NUMBER - 54355/00

The following Sequence Listing pages 1 to 10 are part of the description. The claims pages follow on pages 37 to 51.

<110> ACTIVE BIOTECH AB

<120> AN INTEGRIN HETERODIMER AND AN ALPHA SUBUNIT THEREOF

<130> 2001358

<140>

<141>

<150> SE 9902056-2

<151> 1999-06-03

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 3983

<212> DNA

<213> human

<220>

<221> CDS

<222> (91) .. (3654)

<400> 1

ggcacgaggc cgcgccgagg aggctgccgc tctggcttgc cagtcctcccg ccgccgctgc 60

acaccggacc cagccgccgt gccgcgggcc atg gac ctg ccc agg ggc ctg gtg 114
Met Asp Leu Pro Arg Gly Leu Val
1 5

gtg gcc tgg gcg ctc agc ctg tgg cca ggg ttc acg gac acc ttc aac 162
Val Ala Trp Ala Leu Ser Leu Trp Pro Gly Phe Thr Asp Thr Phe Asn
10 15 20

atg	gac	acc	agg	aag	ccc	cgg	gtc	atc	cct	ggc	tcc	agg	acc	gcc	ttc	210
Met	Asp	Thr	Arg	Lys	Pro	Arg	Val	Ile	Pro	Gly	Ser	Arg	Thr	Ala	Phe	
25					30					35					40	

ttt ggc tac aca gtg cag cag cac gac atc agt ggc aat aag tgg ctg 258
Phe Gly Tyr Thr Val Gln Gln His Asp Ile Ser Gly Asn Lys Trp Leu
45 50 55

gtc gtg ggc gcc cca ctg gaa acc aat ggc tac cag aag acg gga gac 306
Val Val Gly Ala Pro Leu Glu Thr Asn Gly Tyr Gln Lys Thr Gly Asp
60 65 70

gtg tac aag tgt cca gtg atc cac ggg aac tgc acc aaa ctc aac ctg 354
Val Tyr Lys Cys Pro Val Ile His Gly Asn Cys Thr Lys Leu Asn Leu
75 80 85

gga agg gtc acc ctg tcc aac gtg tcc gag cgg aaa gac aac atg cgc 402
Gly Arg Val Thr Leu Ser Asn Val Ser Glu Arg Lys Asp Asn Met Arg
90 95 100

ctc ggc ctt agt ctc gcc acc aac ccc aag gac aac agc ttc ctg gcc 450
Leu Gly Leu Ser Leu Ala Thr Asn Pro Lys Asp Asn Ser Phe Leu Ala
105 110 115 120

tgc agc ccc ctc tgg tct cat gag tgt ggg agc tcc tac tac acc aca	498
Cys Ser Pro Leu Trp Ser His Glu Cys Gly Ser Ser Tyr Tyr Thr Thr	
125 130 135	
ggg atg tgt tca aga gtc aac tcc aac ttc agg ttc tcc aag acc gtg	546
Gly Met Cys Ser Arg Val Asn Ser Asn Phe Arg Phe Ser Lys Thr Val	
140 145 150	
gcc cca gct ctc caa agg tgc cag acc tac atg gac atc gtc att gtc	594
Ala Pro Ala Leu Gln Arg Cys Gln Thr Tyr Met Asp Ile Val Ile Val	
155 160 165	
ctg gat ggc tcc aac agc atc tac ccc tgg gtg gag gtt cag cac ttc	642
Leu Asp Gly Ser Asn Ser Ile Tyr Pro Trp Val Glu Val Gln His Phe	
170 175 180	
ctc atc aac atc ctg aaa aag ttt tac att ggc cca ggg cag atc cag	690
Leu Ile Asn Ile Leu Lys Lys Phe Tyr Ile Gly Pro Gly Gln Ile Gln	
185 190 195 200	
gtt gga gtt gtg cag tat ggc gaa gat gtg gtg cat gag ttt cac ctc	738
Val Gly Val Val Gln Tyr Gly Glu Asp Val Val His Glu Phe His Leu	
205 210 215	
aac gac tac agg tct gta aaa gat gtg gtg gaa gct gcc agc cac att	786
Asn Asp Tyr Arg Ser Val Lys Asp Val Val Glu Ala Ala Ser His Ile	
220 225 230	
gag cag aga gga gga aca gag acc cgg acg gca ttt ggc att gaa ttt	834
Glu Gln Arg Gly Gly Thr Glu Thr Arg Thr Ala Phe Gly Ile Glu Phe	
235 240 245	
gca cgc tca gag gct ttc cag aag ggt gga agg aaa gga gcc aag aag	882
Ala Arg Ser Glu Ala Phe Gln Lys Gly Gly Arg Lys Gly Ala Lys Lys	
250 255 260	
gtg atg att gtc atc aca gat ggg gag tcc cac gac agc cca gac ctg	930
Val Met Ile Val Ile Thr Asp Gly Glu Ser His Asp Ser Pro Asp Leu	
265 270 275 280	
gag aag gtg atc cag caa agc gaa aga gac aac gta aca aga tat gcg	978
Glu Lys Val Ile Gln Gln Ser Glu Arg Asp Asn Val Thr Arg Tyr Ala	
285 290 295	
gtg gcc gtc ctg ggc tac tac aac cgc agg ggg atc aat cca gaa act	1026
Val Ala Val Leu Gly Tyr Tyr Asn Arg Arg Gly Ile Asn Pro Glu Thr	
300 305 310	
ttt cta aat gaa atc aaa tac atc gcc agt gac cct gat gac aag cac	1074
Phe Leu Asn Glu Ile Lys Tyr Ile Ala Ser Asp Pro Asp Asp Lys His	
315 320 325	
ttc ttc aat gtc act gat gag gct gcc ttg aag gac att gtc gat gcc	1122
Phe Phe Asn Val Thr Asp Glu Ala Ala Leu Lys Asp Ile Val Asp Ala	
330 335 340	
ctg ggg gac aga atc ttc agc ctg gaa ggc acc aac aag aac gag acc	1170
Leu Gly Asp Arg Ile Phe Ser Leu Glu Gly Thr Asn Lys Asn Glu Thr	
345 350 355 360	

tcc ttt ggg ctg gag atg tca cag acg ggc ttt tcc tcg cac gtg gtg	1218
Ser Phe Gly Leu Glu Met Ser Gln Thr Gly Phe Ser Ser His Val Val	
365 370 375	
gag gat ggg gtt ctg ctg gga gcc gtc ggt gcc tat gac tgg aat gga	1266
Glu Asp Gly Val Leu Leu Gly Ala Val Gly Ala Tyr Asp Trp Asn Gly	
380 385 390	
gct gtg cta aag gag acg agt gcc ggg aag gtc att cct ctc cgc gag	1314
Ala Val Leu Lys Glu Thr Ser Ala Gly Lys Val Ile Pro Leu Arg Glu	
395 400 405	
tcc tac ctg aaa gag ttc ccc gag gag ctc aag aac cat ggt gca tac	1362
Ser Tyr Leu Lys Glu Phe Pro Glu Glu Leu Lys Asn His Gly Ala Tyr	
410 415 420	
ctg ggg tac aca gtc aca tcg gtc gtg tcc tcc agg cag ggg cga gtg	1410
Leu Gly Tyr Thr Val Thr Ser Val Val Ser Ser Arg Gln Gly Arg Val	
425 430 435 440	
tac gtg gcc gga gcc ccc cgg ttc aac cac acg ggc aag gtc atc ctg	1458
Tyr Val Ala Gly Ala Pro Arg Phe Asn His Thr Gly Lys Val Ile Leu	
445 450 455	
ttc acc atg cac aac aac cgg agc ctc acc atc cac cag gct atg cgg	1506
Phe Thr Met His Asn Asn Arg Ser Leu Thr Ile His Gln Ala Met Arg	
460 465 470	
ggc cag cag ata ggc tct tac ttt ggg agt gaa atc acc tcg gtg gac	1554
Gly Gln Gln Ile Gly Ser Tyr Phe Gly Ser Glu Ile Thr Ser Val Asp	
475 480 485	
atc gac ggc gac ggc gtg act gat gtc ctg ctg gtg ggc gca ccc atg	1602
Ile Asp Gly Asp Gly Val Thr Asp Val Leu Leu Val Gly Ala Pro Met	
490 495 500	
tac ttc aac gag ggc cgt gag cga ggc aag gtg tac gtc tat gag ctg	1650
Tyr Phe Asn Glu Gly Arg Glu Arg Gly Lys Val Tyr Val Tyr Glu Leu	
505 510 515 520	
aga cag aac cgg ttt gtt tat aac gga acg cta aag gat tca cac agt	1698
Arg Gln Asn Arg Phe Val Tyr Asn Gly Thr Leu Lys Asp Ser His Ser	
525 530 535	
tac cag aat gcc cga ttt ggg tcc tcc att gcc tca gtt cga gac ctc	1746
Tyr Gln Asn Ala Arg Phe Gly Ser Ser Ile Ala Ser Val Arg Asp Leu	
540 545 550	
aac cag gat tcc tac aat gac gtg gtg gtg gga gcc ccc ctg gag gac	1794
Asn Gln Asp Ser Tyr Asn Asp Val Val Val Gly Ala Pro Leu Glu Asp	
555 560 565	
aac cac gca gga gcc atc tac atc ttc cac ggc ttc cga ggc agc atc	1842
Asn His Ala Gly Ala Ile Tyr Ile Phe His Gly Phe Arg Gly Ser Ile	
570 575 580	
ctg aag aca cct aag cag aga atc aca gcc tca gag ctg gct acc ggc	1890
Leu Lys Thr Pro Lys Gln Arg Ile Thr Ala Ser Glu Leu Ala Thr Gly	
585 590 595 600	

ctc cag tat ttt ggc tgc agc atc cac ggg caa ttg gac ctc aat gag Leu Gln Tyr Phe Gly Cys Ser Ile His Gly Gln Leu Asp Leu Asn Glu 605 610 615	1938
gat ggg ctc atc gac ctg gca gtg gga gcc ctt ggc aac gct gtg att Asp Gly Leu Ile Asp Leu Ala Val Gly Ala Leu Gly Asn Ala Val Ile 620 625 630	1986
ctg tgg tcc cgc cca gtg gtt cag atc aat gcc agc ctc cac ttt gag Leu Trp Ser Arg Pro Val Val Gln Ile Asn Ala Ser Leu His Phe Glu 635 640 645	2034
cca tcc aag atc aac atc ttc cac aga gac tgc aag cgc agt ggc agg Pro Ser Lys Ile Asn Ile Phe His Arg Asp Cys Lys Arg Ser Gly Arg 650 655 660	2082
gat gcc acc tgc ctg gcc gcc ttc ctc tgc ttc acg ccc atc ttc ctg Asp Ala Thr Cys Leu Ala Ala Phe Leu Cys Phe Thr Pro Ile Phe Leu 665 670 675 680	2130
gca ccc cat ttc caa aca aca act gtt ggc atc aga tac aac gcc acc Ala Pro His Phe Gln Thr Thr Thr Val Gly Ile Arg Tyr Asn Ala Thr 685 690 695	2178
atg gat gag agg cgg tat aca ccg agg gcc cac ctg gac gag ggc ggg Met Asp Glu Arg Arg Tyr Thr Pro Arg Ala His Leu Asp Glu Gly Gly 700 705 710	2226
gac cga ttc acc aac aga gcc gta ctg ctc tcc tcc ggc cag gag ctc Asp Arg Phe Thr Asn Arg Ala Val Leu Leu Ser Ser Gly Gln Glu Leu 715 720 725	2274
tgt gag cgg atc aac ttc cat gtc ctg gac act gct gac tac gtg aag Cys Glu Arg Ile Asn Phe His Val Leu Asp Thr Ala Asp Tyr Val Lys 730 735 740	2322
cca gtg acc ttc tca gtc gag tat tcc ctg gag gac cct gac cat ggc Pro Val Thr Phe Ser Val Glu Tyr Ser Leu Glu Asp Pro Asp His Gly 745 750 755 760	2370
ccc atg ctg gac gac ggc tgg ccc acc act ctc aga gtc tgc gtg ccc Pro Met Leu Asp Asp Gly Trp Pro Thr Thr Leu Arg Val Ser Val Pro 765 770 775	2418
ttc tgg aac ggc tgc aat gag gat gag cac tgt gtc cct gac ctt gtg Phe Trp Asn Gly Cys Asn Glu Asp Glu His Cys Val Pro Asp Leu Val 780 785 790	2466
ttg gat gcc cgg agt gac ctg ccc acg gcc atg gag tac tgc cag agg Leu Asp Ala Arg Ser Asp Leu Pro Thr Ala Met Glu Tyr Cys Gln Arg 795 800 805	2514
gtg ctg agg aag cct gcg cag gac tgc tcc gca tac acg ctg tcc ttc Val Leu Arg Lys Pro Ala Gln Asp Cys Ser Ala Tyr Thr Leu Ser Phe 810 815 820	2562
gac acc aca gtc ttc atc ata gag agc aca cgc cag cga gtg gcg gtg Asp Thr Thr Val Phe Ile Ile Glu Ser Thr Arg Gln Arg Val Ala Val 825 830 835 840	2610

gag gcc aca ctg gag aac agg ggc gag aac gcc tac agt acg gtc cta	2658
Glu Ala Thr Leu Glu Asn Arg Gly Glu Asn Ala Tyr Ser Thr Val Leu	
845 850 855	
aat atc tcg cag tca gca aac ctg cag ttt gcc agc ttg atc cag aag	2706
Asn Ile Ser Gln Ser Ala Asn Leu Gln Phe Ala Ser Leu Ile Gln Lys	
860 865 870	
gag gac tca gac ggt agc att gag tgt gtg aac gag gag agg agg ctc	2754
Glu Asp Ser Asp Gly Ser Ile Glu Cys Val Asn Glu Glu Arg Arg Leu	
875 880 885	
cag aag caa gtc tgc aac gtc agc tat ccc ttc ttc cgg gcc aag gcc	2802
Gln Lys Gln Val Cys Asn Val Ser Tyr Pro Phe Phe Arg Ala Lys Ala	
890 895 900	
aag gtg gct ttc cgt ctt gat tcc gag ttc agc aaa tcc atc ttc cta	2850
Lys Val Ala Phe Arg Leu Asp Ser Glu Phe Ser Lys Ser Ile Phe Leu	
905 910 915 920	
cac cac ctg gag atc gag ctc gct gca ggc agt gac agt aat gag cgg	2898
His His Leu Glu Ile Glu Leu Ala Ala Gly Ser Asp Ser Asn Glu Arg	
925 930 935	
gac agc acc aag gaa gac aac gtg gcc ccc tta cgc ttc cac ctc aaa	2946
Asp Ser Thr Lys Glu Asp Asn Val Ala Pro Leu Arg Phe His Leu Lys	
940 945 950	
tac gag gct gac gtc ctc ttc acc agg agc agc agc ctg agc cac tac	2994
Tyr Glu Ala Asp Val Leu Phe Thr Arg Ser Ser Ser Leu Ser His Tyr	
955 960 965	
gag gtc aag ctc aac agc tcg ctg gag aga tac gat ggt atc ggg cct	3042
Glu Val Lys Leu Asn Ser Ser Leu Glu Arg Tyr Asp Gly Ile Gly Pro	
970 975 980	
ccc ttc agc tgc atc ttc agg atc cag aac ttg ggc ttg ttc ccc atc	3090
Pro Phe Ser Cys Ile Phe Arg Ile Gln Asn Leu Gly Leu Phe Pro Ile	
985 990 995 1000	
cac ggg atg atg atg aag atc acc att ccc atc gcc acc agg agc ggc	3138
His Gly Met Met Met Lys Ile Thr Ile Pro Ile Ala Thr Arg Ser Gly	
1005 1010 1015	
aac cgc cta ctg aag ctg agg gac ttc ctc acg gac gag gcg aac acg	3186
Asn Arg Leu Leu Lys Leu Arg Asp Phe Leu Thr Asp Glu Ala Asn Thr	
1020 1025 1030	
tcc tgt aac atc tgg ggc aat agc act gag tac cgg ccc acc cca gtg	3234
Ser Cys Asn Ile Trp Gly Asn Ser Thr Glu Tyr Arg Pro Thr Pro Val	
1035 1040 1045	
gag gaa gac ttg cgt cgt gct cca cag ctg aat cac agc aac tct gat	3282
Glu Glu Asp Leu Arg Arg Ala Pro Gln Leu Asn His Ser Asn Ser Asp	
1050 1055 1060	
gtc gtc tcc atc aac tgc aat ata cgg ctg gtc ccc aac cag gaa atc	3330
Val Val Ser Ile Asn Cys Asn Ile Arg Leu Val Pro Asn Gln Glu Ile	
1065 1070 1075 1080	

```

aat ttc cat cta ctg ggg aac ctg tgg ttg agg tcc cta aaa gca ctc 3378
Asn Phe His Leu Leu Gly Asn Leu Trp Leu Arg Ser Leu Lys Ala Leu
      1085                      1090                      1095

aag tac aaa tcc atg aaa atc atg gtc aac gca gcc ttg cag agg cag 3426
Lys Tyr Lys Ser Met Lys Ile Met Val Asn Ala Ala Leu Gln Arg Gln
      1100                      1105                      1110

ttc cac agc ccc ttc atc ttc cgt gag gag gat ccc agc cgc cag atc 3474
Phe His Ser Pro Phe Ile Phe Arg Glu Asp Pro Ser Arg Gln Ile
      1115                      1120                      1125

gag ttt gag atc tcc aag caa gag gac tgg cag gtc ccc atc tgg atc 3522
Glu Phe Glu Ile Ser Lys Gln Glu Asp Trp Gln Val Pro Ile Trp Ile
      1130                      1135                      1140

att gta ggc agc acc ctg ggg ggc ctc cta ctg ctg gcc ctg ctg gtc 3570
Ile Val Gly Ser Thr Leu Gly Gly Leu Leu Leu Leu Ala Leu Leu Val
      1145                      1150                      1155                      1160

ctg gca ctg cgg aag ctc ggc ttc ttt aga agt gcc agg cgc agg agg 3618
Leu Ala Leu Arg Lys Leu Gly Phe Phe Arg Ser Ala Arg Arg Arg Arg
      1165                      1170                      1175

gag cct ggt ctg gac ccc acc ccc aaa gtg ctg gag tgaggctcca 3664
Glu Pro Gly Leu Asp Pro Thr Pro Lys Val Leu Glu
      1180                      1185

gaggagactt tgagttgatg ggggccagga caccagtcca ggtagtgttg agaccaggc 3724

ctgtggcccc accgagctgg agcggagagg aagccagctg gctttgcact tgacctcacc 3784

tcccgagcaa tggcgccctgc tcctccaga atggaactca agctggtttt aagtggaaact 3844

gcctactggg agactgggac acctttacac agacccttag ggatttaaag ggacaccctt 3904

acacacaccc aggccacgc caaggcctcc ctcaggctct gtggagggca tttgctgccc 3964

cagctactaa ggtgctagg 3983

```

<210> 2
 <211> 1188
 <212> PRT
 <213> human

```

<400> 2
Met Asp Leu Pro Arg Gly Leu Val Val Ala Trp Ala Leu Ser Leu Trp
  1              5              10              15

Pro Gly Phe Thr Asp Thr Phe Asn Met Asp Thr Arg Lys Pro Arg Val
  20              25              30

Ile Pro Gly Ser Arg Thr Ala Phe Phe Gly Tyr Thr Val Gln Gln His
  35              40              45

Asp Ile Ser Gly Asn Lys Trp Leu Val Val Gly Ala Pro Leu Glu Thr
  50              55              60

Asn Gly Tyr Gln Lys Thr Gly Asp Val Tyr Lys Cys Pro Val Ile His
  65              70              75              80

```

Gly Asn Cys Thr Lys Leu Asn Leu Gly Arg Val Thr Leu Ser Asn Val
 85 90 95
 Ser Glu Arg Lys Asp Asn Met Arg Leu Gly Leu Ser Leu Ala Thr Asn
 100 105 110
 Pro Lys Asp Asn Ser Phe Leu Ala Cys Ser Pro Leu Trp Ser His Glu
 115 120 125
 Cys Gly Ser Ser Tyr Tyr Thr Thr Gly Met Cys Ser Arg Val Asn Ser
 130 135 140
 Asn Phe Arg Phe Ser Lys Thr Val Ala Pro Ala Leu Gln Arg Cys Gln
 145 150 155 160
 Thr Tyr Met Asp Ile Val Ile Val Leu Asp Gly Ser Asn Ser Ile Tyr
 165 170 175
 Pro Trp Val Glu Val Gln His Phe Leu Ile Asn Ile Leu Lys Lys Phe
 180 185 190
 Tyr Ile Gly Pro Gly Gln Ile Gln Val Gly Val Val Gln Tyr Gly Glu
 195 200 205
 Asp Val Val His Glu Phe His Leu Asn Asp Tyr Arg Ser Val Lys Asp
 210 215 220
 Val Val Glu Ala Ala Ser His Ile Glu Gln Arg Gly Gly Thr Glu Thr
 225 230 235 240
 Arg Thr Ala Phe Gly Ile Glu Phe Ala Arg Ser Glu Ala Phe Gln Lys
 245 250 255
 Gly Gly Arg Lys Gly Ala Lys Lys Val Met Ile Val Ile Thr Asp Gly
 260 265 270
 Glu Ser His Asp Ser Pro Asp Leu Glu Lys Val Ile Gln Gln Ser Glu
 275 280 285
 Arg Asp Asn Val Thr Arg Tyr Ala Val Ala Val Leu Gly Tyr Tyr Asn
 290 295 300
 Arg Arg Gly Ile Asn Pro Glu Thr Phe Leu Asn Glu Ile Lys Tyr Ile
 305 310 315 320
 Ala Ser Asp Pro Asp Asp Lys His Phe Phe Asn Val Thr Asp Glu Ala
 325 330 335
 Ala Leu Lys Asp Ile Val Asp Ala Leu Gly Asp Arg Ile Phe Ser Leu
 340 345 350
 Glu Gly Thr Asn Lys Asn Glu Thr Ser Phe Gly Leu Glu Met Ser Gln
 355 360 365
 Thr Gly Phe Ser Ser His Val Val Glu Asp Gly Val Leu Leu Gly Ala
 370 375 380
 Val Gly Ala Tyr Asp Trp Asn Gly Ala Val Leu Lys Glu Thr Ser Ala
 385 390 395 400

Gly Lys Val Ile Pro Leu Arg Glu Ser Tyr Leu Lys Glu Phe Pro Glu
 405 410 415
 Glu Leu Lys Asn His Gly Ala Tyr Leu Gly Tyr Thr Val Thr Ser Val
 420 425 430
 Val Ser Ser Arg Gln Gly Arg Val Tyr Val Ala Gly Ala Pro Arg Phe
 435 440 445
 Asn His Thr Gly Lys Val Ile Leu Phe Thr Met His Asn Asn Arg Ser
 450 455 460
 Leu Thr Ile His Gln Ala Met Arg Gly Gln Gln Ile Gly Ser Tyr Phe
 465 470 475 480
 Gly Ser Glu Ile Thr Ser Val Asp Ile Asp Gly Asp Gly Val Thr Asp
 485 490 495
 Val Leu Leu Val Gly Ala Pro Met Tyr Phe Asn Glu Gly Arg Glu Arg
 500 505 510
 Gly Lys Val Tyr Val Tyr Glu Leu Arg Gln Asn Arg Phe Val Tyr Asn
 515 520 525
 Gly Thr Leu Lys Asp Ser His Ser Tyr Gln Asn Ala Arg Phe Gly Ser
 530 535 540
 Ser Ile Ala Ser Val Arg Asp Leu Asn Gln Asp Ser Tyr Asn Asp Val
 545 550 555 560
 Val Val Gly Ala Pro Leu Glu Asp Asn His Ala Gly Ala Ile Tyr Ile
 565 570 575
 Phe His Gly Phe Arg Gly Ser Ile Leu Lys Thr Pro Lys Gln Arg Ile
 580 585 590
 Thr Ala Ser Glu Leu Ala Thr Gly Leu Gln Tyr Phe Gly Cys Ser Ile
 595 600 605
 His Gly Gln Leu Asp Leu Asn Glu Asp Gly Leu Ile Asp Leu Ala Val
 610 615 620
 Gly Ala Leu Gly Asn Ala Val Ile Leu Trp Ser Arg Pro Val Val Gln
 625 630 635 640
 Ile Asn Ala Ser Leu His Phe Glu Pro Ser Lys Ile Asn Ile Phe His
 645 650 655
 Arg Asp Cys Lys Arg Ser Gly Arg Asp Ala Thr Cys Leu Ala Ala Phe
 660 665 670
 Leu Cys Phe Thr Pro Ile Phe Leu Ala Pro His Phe Gln Thr Thr Thr
 675 680 685
 Val Gly Ile Arg Tyr Asn Ala Thr Met Asp Glu Arg Arg Tyr Thr Pro
 690 695 700
 Arg Ala His Leu Asp Glu Gly Gly Asp Arg Phe Thr Asn Arg Ala Val
 705 710 715 720

Leu Leu Ser Ser Gly Gln Glu Leu Cys Glu Arg Ile Asn Phe His Val
 725 730 735
 Leu Asp Thr Ala Asp Tyr Val Lys Pro Val Thr Phe Ser Val Glu Tyr
 740 745 750
 Ser Leu Glu Asp Pro Asp His Gly Pro Met Leu Asp Asp Gly Trp Pro
 755 760 765
 Thr Thr Leu Arg Val Ser Val Pro Phe Trp Asn Gly Cys Asn Glu Asp
 770 775 780
 Glu His Cys Val Pro Asp Leu Val Leu Asp Ala Arg Ser Asp Leu Pro
 785 790 795 800
 Thr Ala Met Glu Tyr Cys Gln Arg Val Leu Arg Lys Pro Ala Gln Asp
 805 810 815
 Cys Ser Ala Tyr Thr Leu Ser Phe Asp Thr Thr Val Phe Ile Ile Glu
 820 825 830
 Ser Thr Arg Gln Arg Val Ala Val Glu Ala Thr Leu Glu Asn Arg Gly
 835 840 845
 Glu Asn Ala Tyr Ser Thr Val Leu Asn Ile Ser Gln Ser Ala Asn Leu
 850 855 860
 Gln Phe Ala Ser Leu Ile Gln Lys Glu Asp Ser Asp Gly Ser Ile Glu
 865 870 875 880
 Cys Val Asn Glu Glu Arg Arg Leu Gln Lys Gln Val Cys Asn Val Ser
 885 890 895
 Tyr Pro Phe Phe Arg Ala Lys Ala Lys Val Ala Phe Arg Leu Asp Ser
 900 905 910
 Glu Phe Ser Lys Ser Ile Phe Leu His His Leu Glu Ile Glu Leu Ala
 915 920 925
 Ala Gly Ser Asp Ser Asn Glu Arg Asp Ser Thr Lys Glu Asp Asn Val
 930 935 940
 Ala Pro Leu Arg Phe His Leu Lys Tyr Glu Ala Asp Val Leu Phe Thr
 945 950 955 960
 Arg Ser Ser Ser Leu Ser His Tyr Glu Val Lys Leu Asn Ser Ser Leu
 965 970 975
 Glu Arg Tyr Asp Gly Ile Gly Pro Pro Phe Ser Cys Ile Phe Arg Ile
 980 985 990
 Gln Asn Leu Gly Leu Phe Pro Ile His Gly Met Met Met Lys Ile Thr
 995 1000 1005
 Ile Pro Ile Ala Thr Arg Ser Gly Asn Arg Leu Leu Lys Leu Arg Asp
 1010 1015 1020
 Phe Leu Thr Asp Glu Ala Asn Thr Ser Cys Asn Ile Trp Gly Asn Ser
 1025 1030 1035 1040

Thr Glu Tyr Arg Pro Thr Pro Val Glu Glu Asp Leu Arg Arg Ala Pro
 1045 1050 1055
 Gln Leu Asn His Ser Asn Ser Asp Val Val Ser Ile Asn Cys Asn Ile
 1060 1065 1070
 Arg Leu Val Pro Asn Gln Glu Ile Asn Phe His Leu Leu Gly Asn Leu
 1075 1080 1085
 Trp Leu Arg Ser Leu Lys Ala Leu Lys Tyr Lys Ser Met Lys Ile Met
 1090 1095 1100
 Val Asn Ala Ala Leu Gln Arg Gln Phe His Ser Pro Phe Ile Phe Arg
 1105 1110 1115 1120
 Glu Glu Asp Pro Ser Arg Gln Ile Glu Phe Glu Ile Ser Lys Gln Glu
 1125 1130 1135
 Asp Trp Gln Val Pro Ile Trp Ile Ile Val Gly Ser Thr Leu Gly Gly
 1140 1145 1150
 Leu Leu Leu Leu Ala Leu Leu Val Leu Ala Leu Arg Lys Leu Gly Phe
 1155 1160 1165
 Phe Arg Ser Ala Arg Arg Arg Glu Pro Gly Leu Asp Pro Thr Pro
 1170 1175 1180
 Lys Val Leu Glu
 1185

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A recombinant or isolated integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID NO. 1, and homologues and fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein.

2. A process of producing a recombinant integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID NO. 1, and homologues and fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which process comprises the steps of

a) isolating a polynucleotide comprising a nucleotide sequence coding for an integrin subunit $\alpha 11$, or for homologues and fragments thereof, wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein,

b) constructing an expression vector comprising the isolated polynucleotide,

c) transforming a host cell with said expression vector,

d) culturing said transformed host cell in a culture medium under conditions suitable for expression of said integrin subunit $\alpha 11$, of said homologues and fragments wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, in said transformed host cell, and, optionally,

e) isolating the integrin subunit $\alpha 11$, or homologues and fragments thereof, from said transformed host cell or said culture medium.

3. A process according to claim 2, step c, said transforming being an *in vitro* or *in situ* process.

4. A process according to claim 2, step c, said transforming being an *in vivo* process.

5. A process of providing an integrin subunit $\alpha 11$, or homologues or fragments thereof wherein the fragments include at least one binding site of the

integrin subunit $\alpha 11$ protein, as defined in claim 1, whereby said subunit is isolated from a cell in which it is naturally present.

6. An isolated polynucleotide or oligonucleotide comprising a nucleotide coding for an integrin subunit $\alpha 11$, or for homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which polynucleotide or oligonucleotide comprises essentially the nucleotide sequence shown in SEQ ID NO. 1 or suitable parts thereof.

7. An isolated polynucleotide or oligonucleotide which hybridises to a polynucleotide or oligonucleotide as defined in claim 6, whereby said isolated polynucleotide or oligonucleotide fails to hybridise to a polynucleotide or oligonucleotide encoding an integrin subunit $\alpha 10$.

8. A vector comprising a polynucleotide or oligonucleotide as defined in claim 6 or 7.

9. A cell containing the vector as defined in claim 8.

10. A cell, as generated by the process in steps a) to c) of claim 2, in which a polynucleotide or oligonucleotide coding for an integrin subunit $\alpha 11$, or for homologues and fragments thereof, said polynucleotide or oligonucleotide comprising essentially the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof, has been stably integrated in the cell genome.

11. An isolated protein comprising a binding site of the amino acid sequence of the integrin subunit $\alpha 11$, or of homologues and fragments thereof, as defined in claim 1, said binding sites having the capability of binding specifically to entities chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

12. Binding entities having the capability of binding specifically to integrin subunit $\alpha 11$, or to homologues or fragments thereof, as defined in claim 1,

which entities are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

13. A recombinant or isolated integrated heterodimer comprising a subunit $\alpha 11$ and a subunit β , in which the subunit $\alpha 11$ comprises essentially the amino acid sequence shown in SEQ ID NO. 1 or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein.

14. A recombinant or isolated integrin heterodimer according to claim 13, wherein the subunit β is $\beta 1$.

15. A process of producing a recombinant integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , in which the subunit $\alpha 11$ comprises essentially the amino acid sequence shown in SEQ ID NO. 1, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which process comprises the steps of

a) isolating one polynucleotide or oligonucleotide comprising a nucleotide sequence coding for said subunit $\alpha 11$ of said integrin heterodimer, or for said homologues or fragments thereof, and, optionally, another polynucleotide comprising a nucleotide sequence coding for said subunit β of an integrin heterodimer, or for homologues or fragments thereof,

b) constructing an expression vector comprising said isolated polynucleotides or oligonucleotides

c) transforming a host cell with said expression vector or vectors,

d) culturing said transformed host cell in a culture medium under conditions suitable for expression of said integrin heterodimer, or said homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, in said transformed host cell, and, optionally,

e) isolating said integrin heterodimer, or said

homologues or fragments thereof, from said transformed host cell or said culture medium.

16. A process according to claim 15, step c, said transforming being an *in vitro* process.

17. A process according to claim 15, step c, said transforming being an *in vivo* process.

18. A process of providing an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , as defined in claim 13 or 14, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, whereby said integrin heterodimer is isolated from a cell in which it is naturally present.

19. A cell containing

i) a first vector, said first vector comprising a polynucleotide or oligonucleotide coding for a subunit $\alpha 11$ of an integrin heterodimer, or for homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which polynucleotide or oligonucleotide comprises essentially the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof, and

ii) a second vector, said second vector comprising a polynucleotide or oligonucleotide coding for a subunit β of said integrin heterodimer.

20. An isolated protein comprising binding sites of an integrin heterodimer as defined in claim 13 or 14, or of homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, said binding sites having the capability of binding specifically to entities chosen among the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

21. Binding entities having the capability of binding specifically to an integrin heterodimer as defined in claim 13 or 14, or to homologues or fragments

thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, said binding entities being chosen among the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, polyclonal and monoclonal antibodies, and fragments thereof.

22. A fragment of an integrin subunit $\alpha 11$, which integrin subunit $\alpha 11$ comprises essentially the amino acid sequence shown in SEQ ID NO. 1, said fragment being a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the extracellular extension region wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein.

23. A fragment according to claim 22, said fragment being a peptide from the cytoplasmic domain comprising essentially the amino acid sequence KLGFFRSARRRREPGLDPTPKVLE.

24. A fragment according to claim 22, which is a peptide comprising essentially the amino acid sequence of the extracellular domain, from about amino acid No. 804 to about amino acid No. 826 of SEQ ID NO. 1.

25. A fragment according to claim 22, which is a peptide comprising essentially the amino acid sequence of the I-domain, from about amino acid No. 159 to about amino acid No. 355 of SEQ ID NO. 1.

26. A method of producing a fragment of the integrin subunit $\alpha 11$ as defined in any one of claims 22-25, which method comprises a sequential addition of amino acids.

27. A polynucleotide or oligonucleotide coding for a fragment of the integrin subunit $\alpha 11$ as defined in any one of claims 22-25.

28. An isolated protein comprising binding sites of an integrin subunit $\alpha 11$ fragment as defined in any one of claims 22-25, said binding sites having the capability of binding specifically to entities chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, monoclonal and

polyclonal antibodies, and fragments thereof.

29. Binding entities having the capability of binding specifically to an integrin subunit $\alpha 11$ fragment as defined in any one of claims 22-25, which binding entities are chosen from the group comprising proteins, peptides, carbohydrates, lipids, natural integrin binding ligands, monoclonal and polyclonal antibodies and fragments thereof.

30. A process of using an integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID NO. 1 or an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β , or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as a marker or target molecule of cells or tissues expressing said integrin subunit $\alpha 11$, which cells or tissues are of animal including human origin.

31. A process according to claim 30, which is a process for determining the differentiation-state of cells during differentiation, development, in pathological conditions, in tissue regeneration, in transplantation, or in therapeutic and physiological reparation of tissues.



32. A process according to claim 31, which process is used during pathological conditions involving said subunit a11.

5 33. A process according to claim 31, which pathological conditions are comprised within the group comprising damage of muscles, muscle dystrophy, fibrosis and wound healing.

10 34. A process according to claim 31, which pathological conditions are comprised within the group comprising damage of cartilage and/or bone, and cartilage and/or bone diseases.

15 35. A process according to claim 31, which pathological conditions are comprised within the group comprising trauma, rheumatoid arthritis, osteoarthritis and osteoporosis.

36. A process according to claim 30, which is a process for detecting the formation of cartilage during embryonic development.

20 37. A process according to claim 30, which is a process for detecting physiological or therapeutic repair of cartilage and/or muscle.

25 38. A process according to claim 30, which is a process for selection and analysis, or for sorting, isolating or purification of chondrocytes and/or muscle cells.

39. A process according to claim 30, which is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes, respectively, or of muscle or muscle cells
30 during transplantation of muscle or muscle cells, respectively.

40. A process according to claim 30, which is a process for studies of differentiation of chondrocytes or muscle cells.

35 41. A process according to any one of claims 30-40, which is an *in vitro* process.

AMENDED SHEET

42. A process according to any one of claims 30-40, which is an *in situ* process.

43. A process according to any one of claims 30-40, which is an *in vivo* process.

5 44. A process according to any one of claims 30-43, whereby a fragment of said integrin subunit $\alpha 11$ is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the extracellular extension region.

10 45. A process according to claim 44, whereby said fragment is a peptide comprising essentially the amino acid sequence KLGFFRSARRRRREGLDPTPKVLE from the cytoplasmic domain.

15 46. A process according to claim 44, whereby said fragment is a peptide comprising essentially the amino acid sequence of the extracellular domain, from about amino acid No. 804 to about amino acid no. 826 of SEQ ID No. 1.

20 47. A process according to claim 44, whereby said fragment is a peptide comprising essentially the amino acid sequence of the I-domain, from about amino acid No. 159 to about amino acid no. 355 of SEQ ID No. 1.

48. A process according to any one of claims 30-47, whereby a subunit β of the integrin heterodimer is $\beta 1$.

25 49. A process according to claim 30, whereby said cells are chosen from the group comprising fibroblasts, muscle cells, chondrocytes, osteoblasts, mesenchymally derived cells and stem cells.

30 50. A process of using binding entities having the capability of binding specifically to binding sites of an integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID NO. 1, or an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β ,
35 or to homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as markers or target molecules of cells or tissues expressing said integrin subunit $\alpha 11$, which cells or tissues are of animal including human origin.

51. A process according to claim 50, which is a process for detecting the presence of an integrin subunit $\alpha 11$ comprising essentially the amino acid sequence shown in SEQ ID No. 1, or of an integrin heterodimer comprising
5 said subunit $\alpha 11$ and a subunit β , or of homologues or fragments thereof.

52. A process according to claim 50, which is a process for determining the differentiation-state of cells during differentiation, development, in
10 pathological conditions, in tissue regeneration, in transplantation, or in therapeutic and physiological repair of tissues.

53. A process according to claim 52, which process is used during pathological conditions involving said
15 subunit $\alpha 11$.

54. A process according to claim 52, which pathological conditions are comprised within the group comprising damage of muscles, muscle dystrophy, fibrosis and wound healing.

20 55. A process according to claim 52, which pathological conditions are comprised within the group comprising damage of cartilage and/or bone, and cartilage and/or bone diseases.

56. A process according to claim 52, which pathological conditions are comprised within the group comprising trauma, rheumatoid arthritis, osteoarthritis and osteoporosis.

57. A process according to claim 52 which is a process for detecting the formation of cartilage during
30 embryonic development.

58. A process according to claim 52, which is a process for detecting physiological or therapeutic reparation of cartilage and/or muscle.

59. A process according to claim 52, which is a
35 process for selection and analysis, or for sorting, isolating or purification of chondrocytes and/or muscle cells.

AMENDED SHEET

60. A process according to claim 52, which is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes, respectively, or of muscle or muscle cells during transplantation of muscle or muscle cells, respectively.

61. A process according to claim 52, which is a process for studies of differentiation of chondrocytes or muscle cells.

62. A process according to any one of claims 50-61, which is an *in vitro* process.

63. A process according to any one of claims 50-61, which is an *in situ* process.

64. A process according to any one of claims 50-61, which is an *in vivo* process.

65. A process according to any one of claims 50-61, whereby a fragment of said integrin subunit $\alpha 11$ is a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the extracellular extension region.

66. A process according to claim 65, whereby said fragment is a peptide comprising essentially the amino acid sequence KLGFFRSARRRREGLDPTPKVLE from the cytoplasmic domain.

67. A process according to claim 65, whereby said fragment is a peptide comprising essentially the amino acid sequence of the extracellular domain, from about amino acid No. 804 to about amino acid no. 826 of SEQ ID No. 1.

68. A process according to claim 65, whereby said fragment is a peptide comprising essentially the amino acid sequence of the I-domain, from about amino acid No. 159 to about amino acid no. 355 of SEQ ID No. 1.

69. A process according to any one of claims 50-68, whereby a subunit β of the integrin heterodimer is $\beta 1$.

70. A process according to claim 50, whereby said cells are chosen from the group comprising fibroblasts,

AMENDED SHEET

muscle cells, chondrocytes, osteoblasts, mesenchymally derived cells and stem cells.

71. A process for detecting the presence of an integrin subunit $\alpha 11$, or of homologues or fragments of said integrin subunit, on cells, whereby a polynucleotide or oligonucleotide chosen from the group comprising a polynucleotide or oligonucleotide having essentially the nucleotide sequence as shown in SEQ ID NO. 1, or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, is used as a marker under hybridisation conditions, wherein said polynucleotide or oligonucleotide fails to hybridise to a polynucleotide or oligonucleotide encoding an integrin subunit $\alpha 10$.

72. A process according to claim 71, which is a process for determining the differentiation-state of cells during differentiation, development, in pathological conditions, in tissue regeneration, in transplantation, or in therapeutic and physiological reparation of tissues.

73. A process according to claim 72, which process is used during pathological conditions involving said subunit $\alpha 11$.

74. A process according to claim 72, which pathological conditions are comprised within the group comprising damage of muscles, muscle dystrophy, fibrosis and wound healing.

75. A process according to claim 72, which pathological conditions are comprised within the group comprising damage of cartilage and/or bone, and cartilage and/or bone diseases.

76. A process according to claim 72, which pathological conditions are comprised within the group comprising trauma, rheumatoid arthritis, osteoarthritis and osteoporosis.

77. A process according to claim 72, which is a process for detecting the formation of cartilage during embryonic development.



78. A process according to claim 72, which is a process for detecting physiological or therapeutic reparation of cartilage and/or muscle.

5 79. A process according to claim 72, which is a process for selection and analysis, or for sorting, isolating or purification of chondrocytes and/or muscle cells.

10 80. A process according to claim 72, which is a process for detecting regeneration of cartilage or chondrocytes during transplantation of cartilage or chondrocytes, respectively, or of muscle or muscle cells during transplantation of muscle or muscle cells, respectively.

15 81. A process according to claim 72, which is a process for studies of differentiation of chondrocytes or muscle cells.

82. A process according to any one of claims 71-81, which is an *in vitro* process.

20 83. A process according to any one of claims 71-81, which is an *in situ* process.

84. A process according to any one of claims 71-81, which is an *in vivo* process.

25 85. A process according to any one of claims 71-84, whereby said polynucleotide or oligonucleotide is a polynucleotide or oligonucleotide coding for a peptide chosen from the group comprising peptides of the cytoplasmic domain, the I-domain and the extracellular extension region.

30 86. A process according to claim 85, whereby said peptide is a peptide comprising essentially the amino acid sequence KLGFFRSARRRREPGLDPTPKVLE from the cytoplasmic domain.

35 87. A process according to claim 85, whereby said peptide is a peptide comprising essentially the amino acid sequence of the extracellular domain, from about amino acid No. 804 to about amino acid no. 826 of SEQ ID No. 1.

AMENDED SHEET

88. A process according to claim 85, whereby said peptide is a peptide comprising essentially the amino acid sequence of the I-domain, from about amino acid No. 159 to about amino acid No. 355 of SEQ ID NO. 1.

89. A process according to any one of claims 71-88, whereby a subunit β of the integrin heterodimer is $\beta 1$.

90. A process according to claim 71, whereby said cells are chosen from the group comprising fibroblasts, muscle cells, chondrocytes, osteoblasts, mesenchymally derived cells and stem cells.

91. A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit $\alpha 11$, as a target molecule.

92. A pharmaceutical composition comprising as an active ingredient a pharmaceutical agent or an antibody which is capable of stimulating cell surface expression or activation of an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit $\alpha 11$.

93. A pharmaceutical composition according to claim 92, for use in stimulating, inhibiting or blocking the formation of muscles, cartilage, bone or blood vessels.

94. A vaccine comprising as an active ingredient at least one member of the group comprising an integrin heterodimer, which heterodimer comprises a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, and homologues or fragments of said integrin or subunit $\alpha 11$, wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, and a polynucleotide and an oligonucleotide coding for said integrin subunit $\alpha 11$.

95. A method of gene therapy, whereby a vector



comprising a polynucleotide or oligonucleotide coding for a subunit $\alpha 11$ of an integrin heterodimer, or for homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, which polynucleotide or oligonucleotide comprises essentially the nucleotide sequence shown in SEQ ID NO. 1 or parts thereof, and optionally a second vector comprising a polynucleotide or oligonucleotide coding for a subunit β of said integrin heterodimer, is administered to a subject suffering from pathological conditions involving said subunit $\alpha 11$.

96. A method of using binding entities having the capability of binding specifically to binding sites of a integrin subunit $\alpha 11$ comprising substantially the amino acid sequence shown in SEQ ID NO. 1, or of an integrin heterodimer comprising said subunit $\alpha 11$ and a subunit β , or to homologues or fragments thereof, wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein for promoting adhesion of cells.

97. A method of using an integrin heterodimer comprising an integrin subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit $\alpha 11$, as a target for anti-adhesive drugs or molecules in tissues where adhesion impairs the function of the tissue.

98. A method of in vitro detecting the presence of integrin binding entities, comprising interaction of an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit, with a sample, thereby causing said integrin, subunit $\alpha 11$, or homologue or fragment thereof, to modulate the binding to its natural ligand or other integrin binding proteins present in said sample.

99. A method of in vitro studying consequences of the interaction of a human heterodimer integrin comprising a subunit $\alpha 11$ and a subunit β , or the subunit



α 11 thereof, or homologues or fragments of said integrin or subunit, with an integrin binding entity and thereby initiate integrin signalling.

100. A method according to claim 99, whereby the consequences of said interactions are measured as alterations in cellular functions.

101. A method of using a polynucleotide or oligonucleotide encoding an integrin subunit α 11 or homologues or fragments thereof wherein the fragments include at least one binding site of the integrin subunit α 11 protein, as a target molecule.

102. A method according to claim 101, comprising hybridising a polynucleotide or oligonucleotide to the DNA or RNA encoding the integrin subunit α 11 or homologue or fragment thereof wherein the fragments include at least one binding site of the integrin subunit α 11 protein, which polynucleotide or oligonucleotide fails to hybridise to a polynucleotide or oligonucleotide encoding an integrin subunit α 10.

103. A method of using binding entities having the capability of binding specifically to an integrin subunit α 11 comprising the amino acid sequence shown in SEQ ID NO. 1 or SEQ ID NO. 2, or an integrin heterodimer comprising said subunit α 11 and a subunit β , or to homologues or fragments thereof having similar biological activity, for promoting adhesion of chondrocytes and/or osteoblasts to surfaces of implants to stimulate osseointegration.

104. A method of using an integrin heterodimer comprising an integrin subunit α 11 and a subunit β , or the subunit α 11 thereof, or homologues or fragments of said integrin or subunit α 11 wherein the fragments include at least one binding site of the integrin subunit α 11 protein, as a target for anti-adhesive drugs or molecules in tendon, ligament, skeletal muscle or other tissues where adhesion impairs the function of the tissue.

105. A method of stimulating, inhibiting or

blocking the formation of cartilage or bone, comprising administration to a subject a suitable amount of a pharmaceutical agent or an antibody which is capable of using an integrin heterodimer comprising a subunit $\alpha 11$ and a subunit β , or the subunit $\alpha 11$ thereof, or homologues or fragments of said integrin or subunit $\alpha 11$ wherein the fragments include at least one binding site of the integrin subunit $\alpha 11$ protein, as a target molecule.

Dated this 9th day of December 2003

CARTELA AB

By their Patent Attorneys

GRIFFITH HACK



1/11




Fig. 1

SUBSTITUTE SHEET (RULE 26)


[illegible]

TTITCTCGACGTGGTGGAGATGGGGTTCTGCTGGGAGCGCTGGTGCCATATGCTGATATGGAGCTGTCTAAAGGAGACAGATGCCGGGAGGTCTTCTCTCCGCGAGTCTTAC 11320
 S S H V V E D G V L L G A V A Y D W N G A V L K E T S A G K V I P L R E S Y 410
 CTGAAGAAGTCTCCCGAGGAGCTCAAGAACCATGGTGACATCTGGGGTACACAGTCTCATCTGGCTTCCAGGCAAGCGCCAGCTGAGTGGCGGAGGCCCCCGGCTTATCAACCA 1440
 L K E F E L K N H G A Y L G G T V C A T S T V S V S R Q G R V V A G A P C T R F N H 450
 AGGGGCAAGGTGCATCTGTTCACCATGCAACAACACCGACCGCTCCACCATCCACAGGCTATGCGGGCGACGATAGGCTTCTACTTTGGGAGTGAATCACTCGGTGGAGCATCGAC 1560
 TTGTGKVYILFTIMHN¹RS²LT³IHQAMRGQOIGSYFGSEITSV⁴DL⁵ 490
 GCGGCACGGGTGACTGTCTCTGGTGGGCGACCCATGTACTTCAAGAGGCGGTGAGGAGGCAAGGTGATCTATGAGCTGAGACAGACGGTGTGTTATATACGGCAAGC 1680
 DCGV⁶TDV⁷TL⁸VT⁹TL¹⁰GTCTGGTGGGCGACCTGTACTTCAAGAGGCGGTGAGGAGGCAAGGTGATCTATGAGCTGAGACAGACGGTGTGTTATATACGGCAAGC 530
 LK¹¹DS¹²HS¹³YQN¹⁴AR¹⁵FG¹⁶SS¹⁷IAS¹⁸VR¹⁹DL²⁰N²¹OD²²SY²³N²⁴D²⁵V²⁶V²⁷VG²⁸AG²⁹AP³⁰LED³¹N³²H³³ 570
 AGAGACGACCATCTACATCTTACAGGGCTTCCGAGGAGCAGCTGTGAGACAGCTCAACAGGATCTCATAGCTGGTGGTGGGAGCGCCCTGGAGAGCAGCAC 1800
 GAGAIAIT³⁴CH³⁵GF³⁶RG³⁷GS³⁸IL³⁹KT⁴⁰PK⁴¹QRI⁴²ATAS⁴³EL⁴⁴AT⁴⁵GL⁴⁶Q⁴⁷Y⁴⁸FG⁴⁹CS⁵⁰I⁵¹H⁵²G⁵³ 610
 CAATTTGGACCTCAATGAGATGGGCTCATCGACCTGGCAGTGGAGGCCCTTGGCAACGGCTGTGATCTGTGTCGGCGGCGATGTGTGCAGATCATGACAGCTCTCCACTTTGAGCGCATCC 2040
 L⁵⁴DL⁵⁵N⁵⁶E⁵⁷D⁵⁸GL⁵⁹LD⁶⁰LA⁶¹V⁶²GA⁶³L⁶⁴NA⁶⁵V⁶⁶IL⁶⁷WS⁶⁸RV⁶⁹VI⁷⁰NA⁷¹AS⁷²AL⁷³CAATGACAGCTCTCCACTTTGAGCGCATCC 650
 AGAGATCAACATCTTCCACAGAGCTGCAAGCGGAGTGGCAGGATGCCACCTGCTGGCGGCTTCTCTGCTTCAGCGGCTTCTTCTGGCAGCCCATCTTCTGGCAGCCCATTTCCAAACAACAAGTGTGGC 2160
 X⁷⁴IN⁷⁵IF⁷⁶HD⁷⁷CK⁷⁸SG⁷⁹AD⁸⁰AT⁸¹CL⁸²AA⁸³FL⁸⁴CT⁸⁵PI⁸⁶FL⁸⁷AP⁸⁸H⁸⁹F⁹⁰QT⁹¹TT⁹²V⁹³G⁹⁴ 690
 ATGAGATACACGGCACATCGATGAGAGGGGTATACACGAGGCGCCACCTGACGAGGCGGGGACCAATCCACACAGACCGCTACTGCTCTCTCGGCGAGGAGCTCTGTGAG 7200
 L⁹⁵RY⁹⁶NA⁹⁷IT⁹⁸HD⁹⁹ER¹⁰⁰Y¹⁰¹TP¹⁰²RA¹⁰³HL¹⁰⁴DE¹⁰⁵GG¹⁰⁶DR¹⁰⁷FN¹⁰⁸RA¹⁰⁹VL¹¹⁰SL¹¹¹SS¹¹²Q¹¹³EL¹¹⁴CE¹¹⁵E¹¹⁶ 7380
 GGGATCAGTTCATCTCTGGACACTGCTGACTAGTGAGCGATGACCTTCTCATGTGAGTATCTCCCTGGAGGAGCCCTGACCATGGCCCATGCTGGACAGCGCTGGCGCCACCATC 2400
 L¹¹⁷RI¹¹⁸N¹¹⁹F¹²⁰HL¹²¹DT¹²²AD¹²³Y¹²⁴V¹²⁵K¹²⁶PT¹²⁷FS¹²⁸E¹²⁹YS¹³⁰LE¹³¹DP¹³²D¹³³HP¹³⁴HL¹³⁵LD¹³⁶D¹³⁷GH¹³⁸PT¹³⁹TT¹⁴⁰ 770
 TTCAGATCTCGGTGCCCTTCTGGAAAGGCTGCAATGAGATGAGCACTGTGTCTCTGACCTTTGTGTGGATGCCCGGAGTGACCTGCCACGGCCATGGATGACTGGCAGAGGTGGTGTG 2520
 L¹⁴¹RV¹⁴²SV¹⁴³VP¹⁴⁴FW¹⁴⁵NG¹⁴⁶CE¹⁴⁷DE¹⁴⁸HE¹⁴⁹CV¹⁵⁰LD¹⁵¹LV¹⁵²LD¹⁵³AR¹⁵⁴SS¹⁵⁵DL¹⁵⁶PT¹⁵⁷AM¹⁵⁸E¹⁵⁹YC¹⁶⁰OR¹⁶¹V¹⁶²L¹⁶³ 810
 AGGAAGCCTGGCGAGAGCTGCTCCGCAATACACGCTGTCTCTTCAGACACCGACCTTCTATCATACAGAGCAGCGCCGAGGAGTGGGAGGCCACATCTGGAAACAGAGGGCGCGAGAAC 2860
 RK¹⁶⁴PA¹⁶⁵Q¹⁶⁶DC¹⁶⁷SA¹⁶⁸YT¹⁶⁹IL¹⁷⁰SF¹⁷¹DI¹⁷²TV¹⁷³FI¹⁷⁴IES¹⁷⁵TQR¹⁷⁶V¹⁷⁷AVE¹⁷⁸AT¹⁷⁹LEN¹⁸⁰GR¹⁸¹GA¹⁸² 8450

Fig 2c

SUBSTITUTE SHEET (RULE 26)

5/11

**Fig. 3**

SUBSTITUTE SHEET (RULE 26)



-73-

7/11




Fig. 5

SUBSTITUTE SHEET (RULE 26)

8/11




Fig. 6

SUBSTITUTE SHEET (RULE 26)

9/11




Fig. 7 panel 1

SUBSTITUTE SHEET (RULE 26)

10/11




Fig. 7 panel 2

SUBSTITUTE SHEET (RULE 26)



FIG. 8

SUBSTITUTE SHEET (RULE 26)