



(51) International Patent Classification:

C07K 14/725 (2006.01) C07K 16/32 (2006.01)
C07K 14/73 (2006.01) C12N 5/0783 (2006.01)
C07K 16/28 (2006.01)

(21) International Application Number:

PCT/US2020/030818

(22) International Filing Date:

30 April 2020 (30.04.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/841,753 01 May 2019 (01.05.2019) US
62/841,748 01 May 2019 (01.05.2019) US

(71) Applicant: PACT PHARMA, INC. [US/US]; 2 Corporate Drive, South San Francisco, CA 94080 (US).

(72) Inventors: SENNINO, Barbara; c/o PACT PHARMA, INC., 2 Corporate Drive, South San Francisco, CA 94080

(US). JACOBY, Kyle; c/o PACT PHARMA, INC., 2 Corporate Drive, South San Francisco, CA 94080 (US). MANDL-CASHMAN, Stefanie; c/o PACT PHARMA, INC., 2 Corporate Drive, South San Francisco, CA 94080 (US). DUBREUIL, Michael, M.; c/o PACT PHARMA, INC., 2 Corporate Drive, South San Francisco, CA 94080 (US). GAGNON, John; c/o PACT PHARMA, INC., 2 Corporate Drive, South San Francisco, CA 94080 (US). FRANZUSOFF, Alex; c/o PACT PHARMA, INC., 2 Corporate Drive, South San Francisco, CA 94080 (US).

(74) Agent: LENDARIS, Steven, P. et al.; Baker Botts LLP, 30 Rockefeller Plaza, New York, NY 10112-4498 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,

(54) Title: COMPOSITIONS AND METHODS FOR THE TREATMENT OF CANCER USING A CDB ENGINEERED T CELL THERAPY




FIGURE 17B

(57) Abstract: Compositions comprising and methods for the treatment of cancer using a neoTCR based cell therapy with a CD8 expression construct. In certain embodiments, the presently disclosed subject matter provides a cell, comprising an exogenous T cell receptor (TCR), and an exogenous CD8. In certain embodiments, the exogenous CD8 comprises at least one monomer. In certain embodiments, the at least one monomer of the exogenous CD8 comprises an extracellular domain, a transmembrane domain, an intracellular domain, fragments thereof, or combinations thereof.



OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

**(84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

**COMPOSITIONS AND METHODS FOR THE TREATMENT OF CANCER  
USING A CD8 ENGINEERED T CELL THERAPY**

**CROSS-REFERENCE TO RELATED APPLICATIONS**

5           This application claims priority to U.S. Provisional Application No. 62/841,748, filed on May 1, 2019, and U.S. Provisional Application No. 62/841,753, filed on May 1, 2019, the content of which are incorporated in their entirety, and to which priority is claimed.

**SEQUENCE LISTINGS**

10           The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on April 30, 2020, is named 087520\_0145\_SL.txt and is 120,294 bytes in size.

**BACKGROUND OF THE INVENTION**

15           Activation of T cells requires signaling through the T cell receptor (TCR) and its coreceptor molecules. CD4 and CD8 are membrane proteins that are expressed on T helper cells and cytotoxic T lymphocytes that serve as coreceptors that augment TCR signaling by stabilizing the interactions between the peptide-major histocompatibility (pMHC) ligands and the TCR (Li QJ, et al. (2004) CD4 enhances T cell sensitivity to antigen by  
20           coordinating Lck accumulation at the immunological synapse. Nat Immunol 5:791–799; Holler PD, Kranz DM (2003) Quantitative analysis of the contribution of TCR/pepMHC affinity and CD8 to T cell activation. Immunity 18:255–264). Specifically, the CD4 and CD8 coreceptors are essential for the initiation of signaling because they facilitate the recruitment of a kinase to the TCR-pMHC complex. Furthermore, research has shown  
25           that while the CD4 and CD8 coreceptors both augment T cell sensitivity to its ligands, only CD8 plays a role in the stabilization of TCR-pMHC interactions.

          Furthermore, while naturally occurring MHC-I TCRs are presumed to require concurrent CD8 co-receptor help to stabilize peptide-MHC binding, higher affinity TCRs drive CD8-independent target binding and T cell activation. CD4 T cells, when engineered  
30           with high affinity NeoTCRs, are thus able to recognize peptide-MHC-I targets and trigger

effector T cell functions. However, lower affinity TCRs are dependent on CD8 co-receptors to trigger T cell activation.

Accordingly, a NeoTCR cell therapy that is engineered to have CD8 co-receptor expression could stabilize the TCR-pMHC interactions and increase the efficacy of NeoTCR cell therapies that comprise low affinity TCRs that are dependent on such CD8 co-receptors.

## SUMMARY OF THE INVENTION

In certain embodiments, the presently disclosed subject matter provides a cell, comprising an exogenous T cell receptor (TCR), and an exogenous CD8. In certain 10 embodiments, the exogenous CD8 comprises at least one monomer. In certain embodiments, the at least one monomer of the exogenous CD8 comprises an extracellular domain, a transmembrane domain, an intracellular domain, fragments thereof, or combinations thereof. In certain embodiments, the extracellular domain comprises a CD8 $\alpha$  extracellular domain or a CD8 $\beta$  extracellular domain. In certain embodiments, the 15 transmembrane a CD8 $\alpha$  transmembrane domain or a CD8 $\beta$  transmembrane domain. In certain embodiments, the intracellular domain comprises a CD8 $\alpha$  intracellular domain or a CD8 $\beta$  intracellular domain. In certain embodiments, the intracellular domain comprises a CD4 intracellular domain.

In certain embodiments, the at least one monomer comprises a CD8 $\alpha$  extracellular 20 domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain. In certain embodiments, the at least one monomer comprises a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain. In certain embodiments, the at least one monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain. In certain embodiments, the at least one 25 monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain. In certain embodiments, the at least one monomer comprises a signal peptide. In certain embodiments, the signal peptide is a CD8 signal peptide.

In certain embodiments, the extracellular domain comprises the amino acid sequence set forth in SEQ ID NO: 140, or SEQ ID NO: 145. In certain embodiments, the 30 transmembrane domain comprises the amino acid sequence set forth in SEQ ID NO: 141, or SEQ ID NO: 146. In certain embodiments, the intracellular domain comprises the amino acid sequence set forth in SEQ ID NO: 142, SEQ ID NO: 147, or SEQ ID NO: 148.

In certain embodiments, the signal peptide comprises the amino acid sequence set forth in SEQ ID NO: 139, or SEQ ID NO: 144.

In certain embodiments, the exogenous CD8 comprises a 2A sequence. In certain embodiments, the exogenous CD8 comprises a linker. In certain embodiments, the linker  
5 comprises the amino acid sequence set forth in SEQ ID NO: 137. In certain embodiments, the exogenous CD8 comprises a protease cleavage site. In certain embodiments, the protease cleavage site is a Furin cleavage site.

In certain embodiments, the exogenous TCR is a patient derived TCR. In certain embodiments, the exogenous TCR comprises a signal sequence, a first and second 2A  
10 sequence, and a TCR polypeptide sequence. In certain embodiments, the exogenous TCR recognizes a cancer antigen. In certain embodiments, the cancer antigen is a neoantigen. In certain embodiments, the cancer antigen is a patient specific antigen.

In certain embodiments, the cell is a primary cell. In certain embodiments, the cell is a patient-derived cell. In certain embodiments, the cell is a lymphocyte. In certain  
15 embodiments, the cell is a T cell. In certain embodiments, the cell is a young T cell. In certain embodiments, the cell is CD45RA+, CD62L+, CD28+, CD95-, CCR7+, and CD27+. In certain embodiments, the cell is CD45RA+, CD62L+, CD28+, CD95+, CD27+, CCR7+. In certain embodiments, the cell is CD45RO+, CD62L+, CD28+, CD95+, CCR7+, CD27+, CD127+.

In certain embodiments, the cell further comprises a gene modification to enhance  
20 cell persistence and/or enhances memory cell differentiation. In certain embodiments, killing activity of the cell is increased between about 10% to about 500% as compared to killing activity of a cell that does not have the exogenous CD8. In certain embodiments, proliferation of the cell upon binding of the TCR to the antigen is increased between about  
25 10% to about 500% as compared to proliferation of a cell that does not have the exogenous CD8. In certain embodiments, secretion of pro-inflammatory cytokine upon binding of the TCR to the antigen by the cell is increased between about 10% to about 500% as compared to secretion by a cell that does not have the exogenous CD8. In certain  
30 embodiments, LCK affinity of the cell is increased between about 10% to about 500% as compared to LCK affinity of a cell that does not have the exogenous CD8. In certain embodiments, persistence of the cell is increased between about 10% to about 500% as compared to persistence of a cell that does not have the exogenous CD8. In certain  
embodiments, tumor infiltration ability of the cell is increased between about 10% to about

500% as compared to tumor infiltration ability of a cell that does not have the exogenous CD8.

In certain embodiments, wherein the exogenous TCR is a CD8-dependent TCR. In certain embodiments, the exogenous TCR is a CD8-independent TCR. In certain  
5 embodiments, the exogenous CD8 is encoded by a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4. In certain embodiments, the exogenous CD8 comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a  
10 CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain. In certain embodiments, the exogenous CD8 comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID  
15 NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular  
20 domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO:  
25 148).

In certain embodiments, the presently disclosed subject matter provides a method of modifying a cell, the method comprising introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises first and second homology arms homologous to first and second target nucleic acid  
30 sequences, a TCR gene sequence positioned between the first and second homology arms, a CD8 gene sequence positioned between the first and the second homology arms, and recombining the HR template nucleic acid into an endogenous locus of the cell. In certain embodiments, the HR template comprises a first 2A-coding sequence positioned upstream of the CD8 gene sequence, a second 2A-coding sequence positioned downstream of the

CD8 gene sequence and upstream of the TCR gene sequence, and a third 2A-coding sequence positioned downstream of the TCR gene sequence; wherein the first, second, and third 2A-coding sequences code for the same amino acid sequence and are codon-diverged relative to each other. In certain embodiments, the HR template comprises a sequence coding for the amino acid sequence Gly Ser Gly positioned immediately upstream of the first, second, and/or third 2A-coding sequences. In certain embodiments, the HR template further comprises a sequence coding for a Furin cleavage site positioned upstream of the first, second, and/or third 2A-coding sequences. In certain embodiments, the HR template further comprises a sequence encoding a signal sequence positioned immediately upstream of the TCR gene sequence and/or the CD8 gene sequence.

In certain embodiments, the HR template comprises a second TCR sequence positioned between the third 2A-coding sequence and the second homology arm. In certain embodiments, the HR template comprises a sequence encoding a first signal sequence positioned immediately upstream the first TCR gene sequence, and a sequence encoding a second signal sequence positioned immediately upstream of the second TCR gene sequence. In certain embodiments, the HR template comprises a second CD8 gene sequence positioned between the first CD8 gene sequence and the second 2A-coding sequence. In certain embodiments, a 2A coding sequence is positioned between the first and second CD8 gene sequence. In certain embodiments, a sequence coding for the amino acid sequence Gly Ser Gly is positioned between the first and second CD8 gene sequences. In certain embodiments, a sequence coding for a Furin cleavage site is positioned between the first and second CD8 gene sequences.

In certain embodiments, the CD8 gene sequence comprises a sequence encoding an extracellular domain, a sequence encoding an intracellular domain, a sequence encoding an intracellular domain, fragments thereof, or combinations thereof. In certain embodiments, the sequence encoding an extracellular domain comprises a sequence encoding a CD8 $\alpha$  extracellular domain or a CD8 $\beta$  extracellular domain. In certain embodiments, the sequence encoding a transmembrane domain comprises a sequence encoding a CD8 $\alpha$  transmembrane domain or a CD8 $\beta$  transmembrane domain. In certain embodiments, the sequence encoding an intracellular domain comprises a sequence encoding a CD8 $\alpha$  intracellular domain or a CD8 $\beta$  intracellular domain. In certain embodiments, the sequence encoding an intracellular domain comprises a sequence encoding a CD4 intracellular domain.

In certain embodiments, the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain. In certain embodiments, the CD8 gene sequence comprises a sequence encoding a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain. In certain embodiments, the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain. In certain embodiments, the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

In certain embodiments, the HR template comprises a sequence encoding a first signal sequence positioned immediately upstream the first CD8 gene sequence, and a sequence encoding a second signal sequence positioned immediately upstream of the second CD8 gene sequence. In certain embodiments, the signal sequence is a CD8 signal sequence, a human growth hormone signal sequence, fragments thereof, or combinations thereof. In certain embodiments, the first and second homology arms of the HR template are each from about 300 bases to about 2,000 bases in length. In certain embodiments, the first and second homology arms of the HR template are each from about 600 bases to about 2,000 bases in length.

In certain embodiments, the exogenous TCR is a patient derived TCR. In certain embodiments, the exogenous TCR comprises a signal sequence, a first and second 2A sequence, and a TCR polypeptide sequence. In certain embodiments, the exogenous TCR recognizes a cancer antigen. In certain embodiments, the cancer antigen is a neoantigen. In certain embodiments, the cancer antigen is a patient specific antigen. In certain embodiments, the HR template is non-viral. In certain embodiments, the HR template is a circular DNA. In certain embodiments, the HR template is a linear DNA. In certain embodiments, the introducing occurs via electroporation.

In certain embodiments, the recombining comprises cleavage of the endogenous locus by a nuclease, and recombination of the HR template nucleic acid sequence into the endogenous locus by homology directed repair. In certain embodiments, the nuclease is a Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) family nuclease, or derivative thereof. In certain embodiments, the nuclease further comprises a gRNA.

In certain embodiments, the method further comprises culturing the cell. In certain embodiments, the culturing is conducted in the presence of at least one cytokine. In certain embodiments, the culturing is conducted in the presence of IL2, IL7, IL15, or any

combination thereof. In certain embodiments, the culturing is conducted in the presence of IL7 and IL15. In certain embodiments, the method comprises a gene modification to enhance cell persistence and/or enhances memory cell differentiation.

In certain embodiments, the cell is a primary cell. In certain embodiments, the cell is a patient-derived cell. In certain embodiments, the cell is a lymphocyte. In certain 5  
embodiments, the cell is a T cell. In certain embodiments, the cell is a young T cell. In certain embodiments, the cell is CD45RA+, CD62L+, CD28+, CD95-, CCR7+, and CD27+. In certain embodiments, the cell is CD45RA+, CD62L+, CD28+, CD95+, CD27+, CCR7+. In certain embodiments, the cell is CD45RO+, CD62L+, CD28+,  
10 CD95+, CCR7+, CD27+, CD127+.

In certain embodiments, killing activity of the cell is increased between about 10% to about 500% as compared to killing activity of a cell that does not have the CD8 gene sequence. In certain embodiments, proliferation of the cell upon binding of the TCR to the antigen is increased between about 10% to about 500% as compared to proliferation  
15 of a cell that does not have the CD8 gene sequence. In certain embodiments, secretion of pro-inflammatory cytokine upon binding of the TCR to the antigen by the cell is increased between about 10% to about 500% as compared to secretion by a cell that does not have the CD8 gene sequence. In certain embodiments, LCK affinity of the cell is increased between about 10% to about 500% as compared to LCK affinity of a cell that does not  
20 have the CD8 gene sequence. The method of any one of claims 43-95, wherein persistence of the cell is increased between about 10% to about 500% as compared to persistence of a cell that does not have the CD8 gene sequence. In certain embodiments, tumor infiltration ability of the cell is increased between about 10% to about 500% as compared to tumor infiltration ability of a cell that does not have the CD8 gene sequence.

In certain embodiments, the TCR gene encodes a CD8-dependent TCR. In certain 25  
embodiments, the TCR gene encodes a CD8-independent TCR. In certain embodiments, the CD8 gene sequence is encoded by a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4. In certain embodiments, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a  
30 CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, CD4 intracellular domain. In certain embodiments, the

CD8 gene sequence comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

In certain embodiments, the presently disclosed subject matter provides a cell modified by any of the methods disclosed herein.

In certain embodiments, the presently disclosed subject matter provides a composition comprising an effective amount of a cell disclosed herein. In certain embodiments, the composition is a pharmaceutical composition that further comprises a pharmaceutically acceptable excipient. In certain embodiments, the composition is administered to a patient in need thereof for the treatment of cancer. In certain embodiments, the composition comprises a cryopreservation agent. In certain embodiments, the composition comprises serum albumin. In certain embodiments, the composition comprises Plasma-Lyte A, HSA, and CryoStor CS10.

In certain embodiments, the presently disclosed subject matter provides a method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell or a composition disclosed herein.

In certain embodiments, prior to administering the therapeutically effective amount of cells disclosed herein, a non-myeloablative lymphodepletion regimen is administered to the subject. In certain embodiments, the cancer is a solid tumor. In certain embodiments, the cancer is a liquid tumor. In certain embodiments, the solid tumor is selected from the group consisting of melanoma, thoracic cancer, lung cancer, ovarian cancer, breast cancer, pancreatic cancer, head and neck cancer, prostate cancer, gynecological cancer, central nervous system cancer, cutaneous cancer, HPV+ cancer, esophageal cancer, thyroid cancer, gastric cancer, hepatocellular cancer,

cholangiocarcinomas, renal cell cancers, testicular cancer, sarcomas, and colorectal cancer. In certain embodiments, the liquid tumor is selected from the group consisting of follicular lymphoma, leukemia, and multiple myeloma.

5 In certain embodiments, the presently disclosed subject matter provides a kit comprising a cell disclosed herein, reagents for performing a method disclosed herein, or a composition disclosed herein. In certain embodiments, the kit further comprises written instructions for treating a cancer.

10 In certain embodiments, the presently disclosed subject matter provides a cell, comprising: an exogenous T cell receptor (TCR); and an exogenous CD8, comprising: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

15 In certain embodiments, the presently disclosed subject matter provides a cell, comprising: an exogenous T cell receptor (TCR); and an exogenous CD8, comprising: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

20 In certain embodiments, the presently disclosed subject matter provides a method of modifying a cell, the method comprising: introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises: first and second homology arms homologous to first and second target nucleic acid sequences; a TCR gene sequence positioned between the first and second homology arms;

a CD8 gene sequence positioned between the first and the second homology arms; and recombining the HR template nucleic acid into an endogenous locus of the cell, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

In certain embodiments, the presently disclosed subject matter provides a method of modifying a cell, the method comprising: introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises: first and second homology arms homologous to first and second target nucleic acid sequences; a TCR gene sequence positioned between the first and second homology arms; a CD8 gene sequence positioned between the first and the second homology arms; and recombining the HR template nucleic acid into an endogenous locus of the cell, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

In certain embodiments, the presently disclosed subject matter provides a composition comprising a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$

intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

In certain embodiments, the presently disclosed subject matter provides a composition comprising a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO:142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO:142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO:147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO:147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO:148).

In certain embodiments, the presently disclosed subject matter provides a method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

In certain embodiments, the presently disclosed subject matter provides a method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO:

142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

### BRIEF DESCRIPTION OF THE DRAWINGS

**Figures 1A-1C.** **Figures 1A-1C** show an example of a NeoE TCR cassette and gene editing methods that can be used to make NeoTCR Products. **Figure 1A** shows a schematic representing the general targeting strategy used for integrating neoantigen-specific TCR constructs (neoTCRs) into the TCR $\alpha$  locus. **Figures 1B** and **1C** show a neoantigen-specific TCR construct design used for integrating a NeoTCR into the TCR $\alpha$  locus wherein the cassette is shown with signal sequences (“SS”), protease cleavage sites (“P”), and 2A peptides (“2A”). **Figure 1B** shows a target TCR $\alpha$  locus (endogenous TRAC, top panel) and its CRISPR Cas9 target site (horizontal stripes, cleavage site designated by the arrow), and the circular plasmid HR template (bottom panel) with the polynucleotide encoding the neoTCR, which is located between left and right homology arms (“LHA” and “RHA” respectively) prior to integration. **Figure 1C** shows the integrated neoTCR in the TCR $\alpha$  locus (top panel), the transcribed and spliced neoTCR mRNA (middle panel), and translation and processing of the expressed neoTCR (bottom panel).

**Figures 2A-2D.** **Figures 2A-2D** show the circular plasmids used to encode CD8 constructs 1, 2, 3, and 4. **Figure 2A** shows CD8 Construct 1 used to produce the CD8 Product 1. **Figure 2B** shows CD8 Construct 2 used to produce the CD8 Product 2. **Figure 2C** shows CD8 Construct 3 used to produce the CD8 Product 3. **Figure 2D** shows CD8 Construct 4 used to produce the CD8 Product 4. As shown in **Figures 2A-2D**, SS stands for a signal sequence. As described in **Figures 6, 7, 8, and 9**, the SS may be HGH; however other signal sequences may be used as needed for appropriate trafficking. As shown in

**Figures 2A-2D**, P stands for a protease cleavage site. As described in **Figures 6, 7, 8, and 9**, the P may be Furin; however other protease cleavage sites may be used as appropriate to provide the cleavage action described herein. As shown in **Figures 2A-2D**, 2A stands for the 2A peptide. As described in **Figures 6, 7, 8, and 9**, the 2A may be the P2A peptide; however, other 2A peptides may be used.

**Figure 3.** **Figures 3A-3D** show the transcription/splicing and translation processing of each of CD8 Constructs 1, 2, 3, and 4 to yield CD8 Products 1, 2, 3, and 4. As shown in **Figures 3A-3D**, SS stands for a signal sequence. As described in **Figures 6, 7, 8, and 9**, the SS may be HGH; however other signal sequences may be used as needed for appropriate trafficking. As shown in **Figures 3A-3D**, P stands for a protease cleavage site. As described in **Figures 6, 7, 8, and 9**, the P may be Furin; however other protease cleavage sites may be used as appropriate to provide the cleavage action described herein. As shown in **Figures 3A-3D**, 2A stands for the 2A peptide. As described in **Figures 6, 7, 8, and 9**, the 2A may be the P2A peptide; however, other 2A peptides may be used.

**Figures 4A and 4B.** **Figure 4A** shows translated products of CD8 Product 1 and CD8 Product 2. **Figure 4B** shows the translated products of CD8 Product 3 and CD8 Product 4.

**Figure 5.** **Figure 5** provides an exemplary DNA sequence of the NeoTCR construct described in **Figures 1A-1C**. Conservative substitutions of nucleic acids can be used throughout to result in the same translated product. Furthermore, where amino acids can be substituted without a change in function, substitutions of the nucleic acids provided in **Figure 5** can also be used to achieve the substituted amino acids conferring a substantially similar or identical function of the translated proteins.

**Figure 6.** **Figure 6** provides an exemplary DNA sequence of the CD8 Construct 1 (and translated CD8 Product 1) described in **Figures 2A, 3A, and 4A**. Conservative substitutions of nucleic acids can be used throughout to result in the same translated product. Furthermore, where amino acids can be substituted without a change in function, substitutions of the nucleic acids provided in **Figure 6** can also be used to achieve the substituted amino acids conferring a substantially similar or identical function of the translated proteins.

**Figure 7.** **Figure 7** provides an exemplary DNA sequence of the CD8 Construct 2 (and translated CD8 Product 2) described in **Figures 2B, 3B, and 4A**. Conservative substitutions of nucleic acids can be used throughout to result in the same translated product. Furthermore, where amino acids can be substituted without a change in function,

substitutions of the nucleic acids provided in **Figure 7** can also be used to achieve the substituted amino acids conferring a substantially similar or identical function of the translated proteins.

**Figure 8.** **Figure 8** provides an exemplary DNA sequence of the CD8 Construct 3 (and translated CD8 Product 3) described in **Figures 2C, 3C, and 4B**. Conservative substitutions of nucleic acids can be used throughout to result in the same translated product. Furthermore, where amino acids can be substituted without a change in function, substitutions of the nucleic acids provided in **Figure 8** can also be used to achieve the substituted amino acids conferring a substantially similar or identical function of the translated proteins.

**Figure 9.** **Figure 9** provides an exemplary DNA sequence of the CD8 Construct 4 (and translated CD8 Product 4) described in **Figures 2D, 3D, and 4B**. Conservative substitutions of nucleic acids can be used throughout to result in the same translated product. Furthermore, where amino acids can be substituted without a change in function, substitutions of the nucleic acids provided in **Figure 9** can also be used to achieve the substituted amino acids conferring a substantially similar or identical function of the translated proteins.

**Figure 10.** **Figure 10** presents a visual depiction of CD8 Products 1, 2, 3, and 4 along with the predicted LCK activity of each of CD8 Products 1, 2, 3, and 4.

**Figures 11A-11D.** **Figure 11A** provides an exemplary expression construct of CD8 Product 1. **Figure 11B** provides an exemplary expression construct of CD8 Product 2. **Figure 11C** provides an exemplary expression construct of CD8 Product 3. **Figure 11D** provides an exemplary expression construct of CD8 Product 4.

**Figure 12.** **Figure 12** diagrams the design of an Incucyte experiment to show the killing ability of the CD8 Products.

**Figures 13A and 13B.** **Figures 13A and 13B** show the increased tumor killing ability of CD8 Product 4 compared to a NeoTCR Product with the same NeoTCR as the CD8 Product 4 in a population of CD4 T cells. Specifically, both the CD8 Product 4 and the NeoTCR Product express TCR097. However, the CD8 Product 4 also expresses CD8 Construct 4 which comprises the extracellular domain of CD8 $\alpha$  and the intracellular domain of CD4. The effector:target cell ratio (E:T Ratio) was 1:1 (**Figure 13A**) or 2:1 (**Figure 13B**) and each were normalized for gene editing per cell line. The SW620 COX6C R20Q heterozygous tumor cells used in this experiment was a cell line that expresses the cognate antigen for TCR097 (a CD8 dependent TCR). This shows that the

low affinity binding of TCR097 can be saved by co-expressing the extracellular domain of CD8 $\alpha$  and the intracellular domain of CD4.

**Figure 14.** **Figure 14** shows the increased tumor killing ability of CD8 Product 4 compared to a NeoTCR Product with the same NeoTCR as the CD8 Product 4 in a population of CD4 T cells. Specifically, both the CD8 Product 4 and the NeoTCR Product express TCR097. However, the CD8 Product 4 also expresses CD8 Construct 4 which comprises the extracellular domain of CD8 $\alpha$  and the intracellular domain of CD4. The effector:target cell ratio (E:T Ratio) was 1:1, 1:2, or 1:4 and each were normalized for gene editing per cell line. The SW620 COX6C R20Q homozygous tumor cells used in this experiment was a cell line that expresses the cognate antigen for TCR097. An increased killing can be seen in this experiment compared to that shown in **Figures 13A** and **13B** because the cell line is homozygous for the cognate antigen expression; thus, even though TCR97 is a low affinity TCR, the increased amount of cognate antigen overcame the low affinity limitations.

**Figures 15A and 15B.** **Figures 15A and 15B** provide exemplary control experiments showing that there is an increased tumor killing ability of CD8 Product 4 compared to a NeoTCR Product with the same NeoTCR as the CD8 Product 4 in a population of CD4 T cells (top graphs in **Figures 15A and 15B**) or CD8 T cells (bottom graphs in **Figures 15A and 15B**) and that CD8 Product 4 is more effective at tumor killing than simply expressing the NeoTCR construct and CD8 Product 4 in CD8 T cells. **Figure 15A** shows an E:T ratio of 2:1 and **Figure 15B** shows an E:T ratio of 1:1.

**Figure 16.** **Figure 16** shows the surface expression of CD8 Construct 4. Peak #1 is a NeoTCR Product in CD4<sup>+</sup>/CD8<sup>-</sup> T cells (expressing TCR089). Peak #2 is a CD8 Product 4 in CD8<sup>+</sup>/CD4<sup>+</sup> T cells. Peak #3 is a NeoTCR Product in CD8<sup>+</sup>/CD4<sup>-</sup> T cells. As shown CD8 Construct 4 exhibited proper and comparable surface expression as the NeoTCR Products. Similar results were achieved with the other CD8 Constructs (data not shown).

**Figures 17A and 17B.** **Figures 17A and 17B** shows that CD8 $\alpha$  expression boosts CD4 T cell sensitivity while maintaining specificity. **Figure 17A** shows that the expression of CD8 Constructs 1-4 increases the sensitivity of the T cells and does not change the specificity of the CD8 Product to the NeoTCR. As shown, co-expression of the CD8 Constructs decreases the EC50 which shows the increased sensitivity to the NeoTCR. **Figure 17B** shows that the CD8 Products 1-4 are specific for the cognate antigen to NeoTCR 097 because there is only INF $\gamma$  production by the CD8 Cells of the

CD8 Products 1-4 in the presence of cognate antigen (i.e., no INF $\gamma$  production by the CD8 Cells of the CD8 Products 1-4 in the presence of mismatched antigen). The same experiment as shown in **Figures 17A** and **17B** were performed with CD107a instead of INF $\gamma$  and the same increased sensitivity with maintained specificity to the NeoTCR was shown (data not shown) to confirm the results with INF $\gamma$ .

**Figure 18.** **Figure 18** shows that CD8 $\alpha$  expression boosts CD4 T cell sensitivity among CD8-independent NeoTCRs. The figure shows data from the CD8-independent NeoTCR089. This shows that even when a NeoTCR is CD8-independent, there is an increase in sensitivity of CD8 Product 4. This was a surprising result to find that sensitivity can be increased even for CD8-independent TCRs. Similar results were achieved with CD8 Products 1-3 (data not shown). Thus, CD8-dependent (e.g., NeoTCR097) and CD8-independent (e.g., NeoTCR089) exhibit increased sensitivity compared to NeoTCR Products expressing the same NeoTCRs.

## DETAILED DESCRIPTION

### DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art. The following references provide one of skill with a general definition of many of the terms used in the presently disclosed subject matter: Singleton et al., Dictionary of Microbiology and Molecular Biology (2nd ed. 1994); The Cambridge Dictionary of Science and Technology (Walker ed., 1988); The Glossary of Genetics, 5th Ed., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, The Harper Collins Dictionary of Biology (1991). As used herein, the following terms have the meanings ascribed to them below, unless specified otherwise.

It is understood that aspects and embodiments of the invention described herein include "comprising," "consisting," and "consisting essentially of" aspects and embodiments. The terms "comprises" and "comprising" are intended to have the broad meaning ascribed to them in U.S. Patent Law and can mean "includes", "including" and the like.

As used herein, the term "about" or "approximately" means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the

measurement system. For example, “about” can mean within 3 or more than 3 standard deviations, per the practice in the art. Alternatively, “about” can mean a range of up to 20%, e.g., up to 10%, up to 5%, or up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, e.g., within 5-fold or within 2-fold, of a value.

The term "antibody" as used herein is used in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific and tri-specific antibodies), and antibody fragments (e.g., bis-Fabs) so long as they exhibit the desired antigen-binding activity. "Antibody Fragment" as used herein refers to a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to bis-Fabs; Fv; Fab; Fab, Fab'-SH; F(ab')<sub>2</sub>; diabodies; linear antibodies; single-chain antibody molecules (e.g., scFv); and multispecific antibodies formed from antibody fragments.

The terms “Cancer” and “Tumor” are used interchangeably herein. As used herein, the terms “Cancer” or “Tumor” refer to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. The terms are further used to refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth/proliferation. Cancer can affect a variety of cell types, tissues, or organs, including but not limited to an organ selected from the group consisting of bladder, bone, brain, breast, cartilage, glia, esophagus, fallopian tube, gallbladder, heart, intestines, kidney, liver, lung, lymph node, nervous tissue, ovaries, pancreas, prostate, skeletal muscle, skin, spinal cord, spleen, stomach, testes, thymus, thyroid, trachea, urogenital tract, ureter, urethra, uterus, and vagina, or a tissue or cell type thereof. Cancer includes cancers, such as sarcomas, carcinomas, or plasmacytomas (malignant tumor of the plasma cells). Examples of cancer include, but are not limited to, those described herein. The terms “Cancer” or “Tumor” and “Proliferative Disorder” are not mutually exclusive as used herein.

“CD8” is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules.

“CD8 Cells” as used herein means one or more cells precision engineered to express one or more NeoTCRs and a CD8 Construct.

“CD8 Construct” as used herein means any one of a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4.

“CD8 Product” as used herein means a product comprising CD8 Cells.

“CD8 Construct 1” and “CD8 Product 1” refer to a construct that comprises a NeoTCR and CD8 $\alpha$  (CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\alpha$  intracellular domain) and the resulting product that comprises an expressed NeoTCR and CD8 $\alpha$ . Non-limiting examples of a CD8 Product 1 is provided in **Figures 2A, 3A, and 6**. A non-limiting example of a CD8 Product 1 is provided in **Figure 4A**.

“CD8 Construct 2” and “CD8 Product 2” refer to a construct that comprises a NeoTCR, CD8 $\alpha$  (CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\alpha$  intracellular domain), and CD8 $\beta$  (CD8 $\beta$  extracellular domain, CD8 $\beta$  transmembrane domain, and CD8 $\beta$  intracellular domain) and the resulting product that comprises an expressed NeoTCR, CD8 $\alpha$ , and CD8 $\beta$ . Non-limiting examples of a CD8 Product 2 is provided in **Figures 2B, 3B, and 7**. A non-limiting example of a CD8 Product 1 is provided in **Figure 4A**.

“CD8 Construct 3” and “CD8 Product 3” refer to a construct that comprises a NeoTCR, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\beta$  intracellular domain and the resulting product that comprises an expressed NeoTCR, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\beta$  intracellular domain. Non-limiting examples of a CD8 Product 3 is provided in **Figures 2C, 3C, and 8**. A non-limiting example of a CD8 Product 1 is provided in **Figure 4B**.

“CD8 Construct 4” and “CD8 Product 4” refer to a construct that comprises a NeoTCR, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD4 intracellular domain) and the resulting product that comprises an expressed NeoTCR, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD4 intracellular domain. Non-limiting examples of a CD8 Product 4 is provided in **Figures 2D, 3D, and 9**. A non-limiting example of a CD8 Product 1 is provided in **Figure 4B**.

A “conservative substitution” or a “conservative amino acid,” refers to the substitution of an amino acid with a chemically or functionally similar amino acid. Conservative substitution tables providing similar amino acids are well known in the art.

In certain embodiments, acidic amino acids D and E are conservative substitutions for one another; basic amino acids K, R, and H are conservative substitutions for one another; hydrophilic uncharged amino acids S, T, N, and Q are conservative substitutions for one another; aliphatic uncharged amino acids G, A, V, L, and I are conservative

substitutions for one another; non-polar uncharged amino acids C, M, and P are conservative substitutions for one another; aromatic amino acids F, Y, and W are conservative substitutions for one another; A, S, and T are conservative substitutions for one another; D and E are conservative substitutions for one another; N and Q are conservative substitutions for one another; R and K are conservative substitutions for one another; I, L, and M are conservative substitutions for one another; F, Y, and W are conservative substitutions for one another; A and G are conservative substitutions for one another; D and E are conservative substitutions for one another; N and Q are conservative substitutions for one another; R, K and H are conservative substitutions for one another; I, L, M, and V are conservative substitutions for one another; F, Y and W are conservative substitutions for one another; S and T are conservative substitutions for one another; and C and M are conservative substitutions for one another.

Additional conservative substitutions may be found, for example, in Creighton, *Proteins: Structures and Molecular Properties* 2nd ed. (1993) W. H. Freeman & Co., New York, NY.

“Treat,” “Treatment,” and “treating” are used interchangeably and as used herein mean obtaining beneficial or desired results including clinical results. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, the NeoTCR Product of the invention is used to delay development of a proliferative disorder (e.g., cancer) or to slow the progression of such disease.

“Dextramer” as used herein means a multimerized neoepitope-HLA complex that specifically binds to its cognate NeoTCR.

As used herein, the terms “neoantigen”, “neoepitope” or “neoE” refer to a newly formed antigenic determinant that arises, e.g., from a somatic mutation(s) and is recognized as “non-self.” A mutation giving rise to a “neoantigen”, “neoepitope” or “neoE” can include a frameshift or non-frameshift indel, missense or nonsense substitution, splice site alteration (e.g., alternatively spliced transcripts), genomic rearrangement or gene fusion, any genomic or expression alterations, or any post-translational modifications.

“NeoTCR”, “NeoE TCR” and “exogenous TCR” as used herein mean a neoepitope-specific T cell receptor that is introduced into a T cell, e.g., by gene editing

methods. As used herein, the term “TCR gene sequence” refers to a NeoTCR gene sequence.

“NeoTCR cells” as used herein means one or more cells precision engineered to express one or more NeoTCRs. In certain embodiments, the cells are T cells. In certain  
5 embodiments, the T cells are CD8<sup>+</sup> and/or CD4<sup>+</sup> T cells. In certain embodiments, the CD8<sup>+</sup> and/or CD4<sup>+</sup> T cells are autologous cells from the patient for whom a NeoTCR Product will be administered. The terms “NeoTCR cells” and “NeoTCR-P1 T cells” and “NeoTCR-P1 cells” are used interchangeably herein.

“NeoTCR Product” as used herein means a pharmaceutical formulation comprising  
10 one or more NeoTCR cells. NeoTCR Product consists of autologous precision genome-engineered CD8<sup>+</sup> and CD4<sup>+</sup> T cells. Using a targeted DNA-mediated *non-viral* precision genome engineering approach, expression of the endogenous TCR is eliminated and replaced by a patient-specific NeoTCR isolated from peripheral CD8<sup>+</sup> T cells targeting the tumor-exclusive neoepitope. In certain embodiments, the resulting engineered CD8<sup>+</sup>  
15 or CD4<sup>+</sup> T cells express NeoTCRs on their surface of native sequence, native expression levels, and native TCR function. The sequences of the NeoTCR external binding domain and cytoplasmic signaling domains are unmodified from the TCR isolated from native CD8<sup>+</sup> T cells. Regulation of the NeoTCR gene expression is driven by the native endogenous TCR promoter positioned upstream of where the NeoTCR gene cassette is  
20 integrated into the genome. Through this approach, native levels of NeoTCR expression are observed in unstimulated and antigen-activated T cell states.

The NeoTCR Product manufactured for each patient represents a defined dose of autologous CD8<sup>+</sup> and/or CD4<sup>+</sup> T cells that are precision genome engineered to express a single neoE-specific TCR cloned from neoE-specific CD8<sup>+</sup> T cells individually isolated  
25 from the peripheral blood of that same patient.

“NeoTCR Viral Product” as used herein has the same definition of NeoTCR Product except that the genome engineering is performed using viral mediated methods.

"Pharmaceutical Formulation" refers to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be effective, and  
30 which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered. For clarity, DMSO at quantities used in a NeoTCR Product is not considered unacceptably toxic.

A "subject," "patient," or an "individual" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo,

sports, or pet animals, such as dogs, horses, cats, cows, etc. Preferably, the mammal is human.

“TCR” as used herein means T cell receptor.

The term “tumor antigen” as used herein refers to an antigen (e.g., a polypeptide) that is uniquely or differentially expressed on a tumor cell compared to a normal or non-neoplastic cell. In certain embodiments, a tumor antigen includes any polypeptide expressed by a tumor that is capable of activating or inducing an immune response via an antigen-recognizing receptor or capable of suppressing an immune response via receptor-ligand binding.

“2A” and “2A peptide” are used interchangeably herein and mean a class of 18-22 amino acid long, viral, self-cleaving peptides that are able to mediate cleavage of peptides during translation in eukaryotic cells.

Four well-known members of the 2A peptide class are T2A, P2A, E2A, and F2A. The T2A peptide was first identified in the *Thosea asigna* virus 2A. The P2A peptide was first identified in the porcine teschovirus-1 2A. The E2A peptide was first identified in the equine rhinitis A virus. The F2A peptide was first identified in the foot-and-mouth disease virus.

The self-cleaving mechanism of the 2A peptides is a result of ribosome skipping the formation of a glycyl-prolyl peptide bond at the C-terminus of the 2A. Specifically, the 2A peptides have a C-terminal conserved sequence that is necessary for the creation of steric hindrance and ribosome skipping. The ribosome skipping can result in one of three options: 1) successful skipping and recommencement of translation resulting in two cleaved proteins (the upstream of the 2A protein which is attached to the complete 2A peptide except for the C-terminal proline and the downstream of the 2A protein which is attached to one proline at the N-terminal; 2) successful skipping but ribosome fall-off that results in discontinued translation and only the protein upstream of the 2A; or 3) unsuccessful skipping and continued translation (i.e., a fusion protein).

The term “endogenous” as used herein refers to a nucleic acid molecule or polypeptide that is normally expressed in a cell or tissue.

The term “exogenous” as used herein refers to a nucleic acid molecule or polypeptide that is not endogenously present in a cell. The term “exogenous” would therefore encompass any recombinant nucleic acid molecule or polypeptide expressed in a cell, such as foreign, heterologous, and over-expressed nucleic acid molecules and polypeptides. By “exogenous” nucleic acid is meant a nucleic acid not present in a native

wild-type cell; for example, an exogenous nucleic acid may vary from an endogenous counterpart by sequence, by position/location, or both. For clarity, an exogenous nucleic acid may have the same or different sequence relative to its native endogenous counterpart; it may be introduced by genetic engineering into the cell itself or a progenitor thereof, and  
5 may optionally be linked to alternative control sequences, such as a non-native promoter or secretory sequence.

“Young” or “Younger” or “Young T cell” as it relates to T cells means memory stem cells ( $T_{MSC}$ ) and central memory cells ( $T_{CM}$ ). These cells have T cell proliferation upon specific activation and are competent for multiple cell divisions. They also have the  
10 ability to engraft after re-infusion, to rapidly differentiate into effector T cells upon exposure to their cognate antigen and target and kill tumor cells, as well as to persist for ongoing cancer surveillance and control.

### NeoTCR PRODUCTS

In some embodiments, using the gene editing technology and neoTCR isolation  
15 technology described in PCT/US2020/17887 and PCT/US2019/025415, which are incorporated herein in their entireties, NeoTCRs are cloned in autologous CD8+ and CD4+ T cells from the same patient with cancer by precision genome engineered (using a DNA-mediated (non-viral) method as described in **Figures 1A-1C**) to express the neoTCR. In other words, the NeoTCRs that are tumor specific are identified in cancer patients, such  
20 NeoTCRs are then cloned, and then the cloned NeoTCRs are inserted into the cancer patient’s T cells. NeoTCR expressing T cells are then expanded in a manner that preserves a “young” T cell phenotypes, resulting in a NeoTCR-P1 product (i.e., a NeoTCR Product) in which the majority of the T cells exhibit T memory stem cell and T central memory phenotypes.

25 These ‘young’ or ‘younger’ or less-differentiated T cell phenotypes are described to confer improved engraftment potential and prolonged persistence post-infusion. Thus, the administration of NeoTCR Product, consisting significantly of ‘young’ T cell phenotypes, has the potential to benefit patients with cancer, through improved engraftment potential, prolonged persistence post-infusion, and rapid differentiation into  
30 effector T cells to eradicate tumor cells throughout the body.

*Ex vivo* mechanism-of-action studies were also performed with NeoTCR Product manufactured with T cells from patients with cancer. Comparable gene editing efficiencies

and functional activities, as measured by antigen-specificity of T cell killing activity, proliferation, and cytokine production, were observed demonstrating that the manufacturing process described herein is successful in generating products with T cells from patients with cancer as starting material.

5 In certain embodiments, the NeoTCR Product manufacturing process involves electroporation of dual ribonucleoprotein species of CRISPR-Cas9 nucleases bound to guide RNA sequences, with each species targeting the genomic TCR $\alpha$  and the genomic TCR $\beta$  loci. The specificity of targeting Cas9 nucleases to each genomic locus has been previously described in the literature as being highly specific. Comprehensive testing of  
10 the NeoTCR Product was performed *in vitro* and *in silico* analyses to survey possible off-target genomic cleavage sites, using COSMID and GUIDE-seq, respectively. Multiple NeoTCR Product or comparable cell products from healthy donors were assessed for cleavage of the candidate off-target sites by deep sequencing, supporting the published evidence that the selected nucleases are highly specific.

15 Further aspects of the precision genome engineering process have been assessed for safety. No evidence of genomic instability following precision genome engineering was found in assessing multiple NeoTCR Products by targeted locus amplification (TLA) or standard FISH cytogenetics. No off-target integration anywhere into the genome of the NeoTCR sequence was detected. No evidence of residual Cas9 was found in the cell  
20 product.

The comprehensive assessment of the NeoTCR Product and precision genome engineering process indicates that the NeoTCR Product will be well tolerated following infusion back to the patient.

The genome engineering approach described herein enables highly efficient  
25 generation of bespoke NeoTCR T cells (i.e., NeoTCR Products) for personalized adoptive cell therapy for patients with solid and liquid tumors. Furthermore, the engineering method is not restricted to the use in T cells and has also been applied successfully to other primary cell types, including natural killer and hematopoietic stem cells.

## **CD8 PRODUCTS**

30 Coexpression of MHC class I-restricted neoTCRs and ectopic CD8 receptors in precision genome engineered CD4 T cells significantly potentiates antigen-specific effector functions.

Neopeptides from tumor-exclusive mutations represent compelling targets for personalized neoE-specific autologous TCR-T cell therapies for patients with solid tumors. The imPACT Isolation Technology as described in PCT/US2020/17887, which is incorporated by reference in its entirety, is an ultra-sensitive and high-throughput process for capturing neoE-specific CD8 T cells from the blood of patients with solid cancers. Leveraging this technology, neopeptide-specific MHC class I-restricted TCRs (“MHC-I neoTCRs”) were cloned from individually captured CD8 T cells. Using DNA-mediated (non-viral) gene editing as described in **Example 1**, fresh CD8 and CD4 T cells from the same patient with cancer were engineered to express the MHC-I neoTCR (concomitant with elimination of the endogenous TCR).

While naturally occurring MHC-I TCRs were presumed to require concurrent CD8 co-receptor help to stabilize peptide-MHC binding, higher affinity TCRs were able to drive CD8-independent target binding and T cell activation. CD4 T cells, when engineered with high affinity neoTCRs, were thus able to recognize peptide-MHC-I targets and trigger effector T cell functions. However, lower affinity TCRs were dependent on CD8 co-receptors to trigger T cell activation. By precision genome engineering CD8 co-receptor genes together with the neoTCR into CD4 T cells, MHC-I neoTCRs were made competent to trigger antigen-specific effector T cell function.

CD8 stabilizes TCR-pMHC interactions and synergizes with TCRs for avidity. CD8 also serves to enhance avidity from low affinity TCRs, while the intracellular domain of CD8 $\alpha$  is critical for enhanced T cell activation. Expression of CD8 may enhance CD4 T cell responses, which may not respond to physiological concentrations of pMHC. Disruption of CD8 binding to MHC can convert catch-bond TCR-pMHC into slip-bonds, highlighting the importance of CD8-MHC interactions even for TCRs that bind pMHC independent of CD8.

CD8 $\alpha$  has a lower affinity to LCK than CD8 $\beta$ , suggesting that either co-expression of both CD8 $\alpha$  and CD8 $\beta$  or generation of a chimeric CD8 $\alpha$ -CD8 $\beta$  molecule which contains the extracellular CD8 $\alpha$  and intracellular domain of CD8 $\beta$  can improve effectiveness. The CD8 Constructs described herein are:

1. CD8 $\alpha$  homodimer (CD8 Construct 1)
2. CD8 $\alpha$ -P2A-CD8 $\beta$  (CD8 Construct 2)
3. CD8 $\alpha$  with CD8 $\beta$  intracellular domain (CD8 Construct 3)
4. CD8 $\alpha$  homodimer with CD4 intracellular domain (CD8 Construct 4)

In some embodiments, the NeoTCR Products described above include an additional modification to include the expression of CD8 Construct 1, CD8 Construct 2, CD8 Construct 3, or CD8 Construct 4 (each a CD8 Product). Specifically, using the gene editing technology and neoTCR isolation technology described in PCT/US2020/17887 and PCT/US2019/025415, which are incorporated herein in their entireties, NeoTCRs are cloned in autologous CD8<sup>+</sup> and CD4<sup>+</sup> T cells from the same patient with cancer by precision genome engineered (using a DNA-mediated (non-viral) method as described in **Figures 1A-1C**) to express the neoTCR.

Each of the CD8 Constructs, when expressed, result in CD8 Products. **Table 1** provides a description of each construct and product.

**Table 1. CD8 Products and CD8 Constructs**

CD8 Construct	CD8 Product	Expression components	Exemplary constructs	Exemplary products
CD8 Construct 1	CD8 Product 1	NeoTCR, CD8 $\alpha$ extracellular domain, CD8 $\alpha$ transmembrane domain, CD8 $\alpha$ intracellular domain	<b>Figure 2A, Figure 6</b>	<b>Figure 3A, Figure 4A</b>
CD8 Construct 2	CD8 Product 1	NeoTCR, CD8 $\alpha$ extracellular domain, CD8 $\alpha$ transmembrane domain, CD8 $\alpha$ intracellular domain, CD8 $\beta$ extracellular domain, CD8 $\beta$ transmembrane domain, CD8 $\beta$ intracellular domain	<b>Figure 2B, Figure 7</b>	<b>Figure 3B, Figure 4A</b>
CD8 Construct 3	CD8 Product 1	NeoTCR, CD8 $\alpha$ extracellular domain, CD8 $\alpha$ transmembrane domain, CD8 $\beta$ intracellular domain	<b>Figure 2C, Figure 8</b>	<b>Figure 3C, Figure 4B</b>
CD8 Construct 4	CD8 Product 1	NeoTCR, CD8 $\alpha$ extracellular domain, CD8 $\alpha$ transmembrane domain, CD4 intracellular domain	<b>Figure 2D, Figure 9</b>	<b>Figure 3D, Figure 4B</b>

In certain embodiments, the CD8 Product 1 comprises a NeoTCR and a CD8 homodimer. In certain embodiments, the CD8 Product 1 comprises the expression of a NeoTCR, a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain. In certain embodiments, the CD8 Product 1 further includes the expression of a CD8 $\alpha$  signal peptide. In certain embodiments, the CD8 Product 1 comprises the translated elements presented in **Figure 3A**. In a non-limiting exemplary

embodiment, the CD8 Product 1 comprises a NeoTCR, a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142). In certain embodiments, sequence modifications of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\alpha$  intracellular domain can be made that conserve or substantially conserve function of each element. In certain embodiments, such sequence modifications are conservative substitutions of amino acids.

In a non-limiting embodiment, the CD8 Product 1 is manufactured from a CD8 Construct 1 provided in **Figure 6**. In certain embodiments, the sequence of CD8 Construct 1 provided in **Figure 6** can be modified in any number of ways so long as the translation of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\alpha$  intracellular domain leave each element with conserved function. In certain embodiments, the order of each element of the CD8 Construct 1 in **Figure 6** remains the same but the sequences of each individual element can be changed so long as the amino acids that the nucleic acid encodes remain the same or only comprise conservative substitutions. In certain embodiments, the order of each element of the CD8 Construct 1 in **Figure 6** remains the same but the sequences of each individual element can be changed so long as the function of the encoded proteins remains substantially unchanged.

In certain embodiments, the CD8 Product 2 comprises a NeoTCR, a CD8 $\alpha$ , and aCD8 $\beta$ . In certain embodiments, the CD8 $\alpha$  and CD8 $\beta$  are separated by a protease cleavage site and a 2A peptide in the CD8 Product 2 construct for expression. In certain embodiments, the CD8 Product 2 comprises the expression of a NeoTCR, a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain. In certain embodiments, the CD8 Product 2 further comprises the expression of a CD8 $\alpha$  signal peptide. In certain embodiments, the CD8 Product further comprises the expression of a CD8 $\beta$  signal peptide. In certain embodiments, the CD8 Product further comprises the expression of a CD8 $\alpha$  signal peptide and a CD8 $\beta$  signal peptide. In certain embodiments, the CD8 Product 2 comprises the translated elements presented in **Figure 3B**. In a non-limiting exemplary embodiment, the CD8 Product 2 comprises a NeoTCR, a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular

domain (SEQ ID NO: 147). In certain embodiments, sequence modifications of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, CD8 $\alpha$  intracellular domain, CD8 $\beta$  extracellular domain, CD8 $\beta$  transmembrane domain, and CD8 $\beta$  intracellular domain can be made that conserve or substantially conserve function of each element. In certain embodiments, such sequence modifications are conservative substitutions of amino acids.

In a non-limiting embodiment, the CD8 Product 2 is manufactured from a CD8 Construct 2 provided in **Figure 7**. In certain embodiments, the sequence of CD8 Construct 2 provided in **Figure 7** can be modified in any number of ways so long as the translation of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, CD8 $\alpha$  intracellular domain, CD8 $\beta$  extracellular domain, CD8 $\beta$  transmembrane domain, and CD8 $\beta$  intracellular domain leave each element with conserved function. In certain embodiments, the order of each element of the CD8 Construct 2 in **Figure 7** remains the same but the sequences of each individual element can be changed so long as the amino acids that the nucleic acid encodes remain the same or only comprise conservative substitutions. In certain embodiments, the order of each element of the CD8 Construct 2 in **Figure 7** remains the same but the sequences of each individual element can be changed so long as the function of the encoded proteins remains substantially unchanged.

In certain embodiments, the CD8 Product 3 comprises a NeoTCR and a CD8 $\alpha$  with CD8 $\beta$  intracellular domain. In certain embodiments, the CD8 Product 3 comprises the expression of a NeoTCR, a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain. In certain embodiments, the CD8 Product 3 further includes the expression of a CD8 $\alpha$  signal peptide. In certain embodiments, the CD8 Product 3 comprises the translated elements presented in **Figure 3C**. In a non-limiting exemplary embodiment, the CD8 Product 3 comprises a NeoTCR, a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147). In certain embodiments, sequence modifications of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\beta$  intracellular domain can be made that conserve or substantially conserve function of each element. In certain embodiments, such sequence modifications are conservative substitutions of amino acids.

In a non-limiting embodiment, the CD8 Product 3 is manufactured from a CD8 Construct 3 provided in **Figure 8**. In certain embodiments, the sequence of CD8 Construct 3 provided in **Figure 8** can be modified in any number of ways so long as the translation

of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD8 $\beta$  intracellular domain leave each element with conserved function. In certain embodiments, the order of each element of the CD8 Construct 3 in **Figure 8** remains the same but the sequences of each individual element can be changed so long as the amino acids that the nucleic acid encodes remain the same or only comprise conservative substitutions. In certain embodiments, the order of each element of the CD8 Construct 3 in **Figure 8** remains the same but the sequences of each individual element can be changed so long as the function of the encoded proteins remains substantially unchanged.

In certain embodiments, the CD8 Product 4 comprises a NeoTCR and a CD8 $\alpha$  homodimer with a CD4 intracellular domain. In certain embodiments, the CD8 Product 4 comprises the expression of a NeoTCR, a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain. In certain embodiments, the CD8 Product 4 further includes the expression of a CD8 $\alpha$  signal peptide. In certain embodiments, the CD8 Product 4 comprises the translated elements presented in **Figure 3D**. In a non-limiting exemplary embodiment, the CD8 Product 4 comprises a NeoTCR, a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148). In certain embodiments, sequence modifications of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD4 intracellular domain can be made that conserve or substantially conserve function of each element. In certain embodiments, such sequence modifications are conservative substitutions of amino acids.

In a non-limiting embodiment, the CD8 Product 4 is manufactured from a CD8 Construct 4 provided in **Figure 9**. In certain embodiments, the sequence of CD8 Construct 4 provided in **Figure 9** can be modified in any number of ways so long as the translation of the CD8 $\alpha$  signal peptide, CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, and CD4 intracellular domain leave each element with conserved function. In certain embodiments, the order of each element of the CD8 Construct 4 in **Figure 9** remains the same but the sequences of each individual element can be changed so long as the amino acids that the nucleic acid encodes remain the same or only comprise conservative substitutions. In certain embodiments, the order of each element of the CD8 Construct 4 in **Figure 9** remains the same but the sequences of each individual element can be changed so long as the function of the encoded proteins remains substantially unchanged.

In certain embodiments, the CD8 Products comprise a TET2 knockout or TET2 knockdown. In certain embodiments, the cells of the CD8 Product are further engineered

to knockout the TET2 gene using non-viral methods. In certain embodiments, the cells of the CD8 Product are further engineered to knockout the TET2 gene using viral methods. In certain embodiments, the cells of the CD8 Product are further engineered to increase T cell persistence by knocking out, knocked down, or modifying the function of a gene associated with T cell persistence using non-viral methods. In certain embodiments, the cells of the CD8 Product are further engineered to increase T cell persistence by knocking out, knocking down, or modifying the function of a gene associated with T cell persistence using viral methods.

In some embodiments, the CD8 Products comprise cells that were engineered to express a NeoTCR and a CD8 Construct using viral methods (i.e., CD8 Viral Product). In certain embodiments, the cells of the CD8 Viral Product are further engineered to knockout the TET2 gene using non-viral methods. In certain embodiments, the cells of the CD8 Viral Product are further engineered to knockout the TET2 gene using viral methods. In certain embodiments, the cells of the CD8 Viral Product are further engineered to increase T cell persistence by knocking out, knocking down, or modifying the function of a gene associated with T cell persistence using non-viral methods. In certain embodiments, the cells of the CD8 Viral Product are further engineered to increase T cell persistence by knocking out, knocking down, or modifying the function of a gene associated with T cell persistence using viral methods.

In certain embodiments, the T cell persistence gene that is knocked out, knocked down, or with a modified function is a gene that confers downregulation of T cell activity. In certain embodiments, the gene that is knocked out, knocked down, or modified is a gene that downregulates T cell memory function. In certain embodiments, the gene that is knocked out, knocked down, or modified is a gene that decreases T-cell function, proliferation, and/or survival.

In certain embodiments, additional modifications to the CD8 Products and CD8 Cells thereof include modifications to increase tumor microenvironment resilience, increase T cell activity, increase tumor microenvironment homing/retention, increase T cell persistence, and increase ectopic effector functions of T cells. In certain embodiments, the tumor microenvironment resilience includes but is not limited to converting/counteracting negative environmental signals. In certain embodiments, TCR-mediated signal enhancements include but are not limited to increasing T cell activity. In certain embodiments, the tumor microenvironment homing/retention includes but is not limited to enhancing tumor infiltration. In certain embodiments, the functional T cell

persistence includes but is not limited to metabolic and transcriptional regulation. In certain embodiments, the ectopic effector functions include but is not limited to TCR-induced antibody, cytokine, or peptide secretion.

In certain embodiments, functional T cell persistence can be accomplished by genetic engineering as described herein, by co-administration of a pharmaceutical agent that improves the functional T cell persistence of a modified T cells (modified to incorporate at least a neoTCR as described herein), and by manufacturing and culture conditions of the modified T cells (modified to incorporate at least a neoTCR as described herein).

In certain embodiments, the TCR-induced antibodies used to improve ectopic effector functions are any one of the antibodies or functional fragments thereof described herein. In certain embodiments, the TCR-induced cytokines used to improve ectopic effector functions are naturally occurring cytokines, modified cytokines, fusion proteins of the cytokines, or any combination thereof. In certain embodiments, the TCR-induced cytokines are not cytokines but rather other co-factors or expression elements that induce endogenous cytokine production.

In certain embodiments, additional modifications to the CD8 Products and CD8 Cells thereof include modifications to knock in one or more additional genes and/or functional proteins. In certain embodiments, the genes and/or functional proteins knocked in include but are not limited c-Myb, dominant negative FAS, FAS truncations, FBXW7, CTP1A, OPA1, GLUT1, CA-STAT5A, dominant negative TGF $\beta$ R, DNMT3a, dominant negative PD-1R, dominant negative PD-1 or PD-L1, or PD-L2, dominant negative SHIP-1 protein, integrins, chemokine receptors, cytokines, and interleukins. In certain embodiments, the dominant negative form of a gene is an antibody or functional fragment thereof that is an antagonist of the gene. For example, a dominant negative PD-1 can be an anti-PD-1 antibody or functional fragment thereof. In certain embodiments, any of the knock in genes and/or functional proteins is a functional fragment (including but not limited to truncations) of the gene and/or protein.

In certain embodiments, additional modifications to the CD8 Products and CD8 Cells thereof include modifications to knock out or knock down of one or more additional genes and/or functional proteins. In certain embodiments, the genes and/or functional proteins knocked out or knocked down include but are not limited to TET2, IFNGR1, RICTOR, NR4A1, DNMT3A, SUV39H1, PPP2RD, adenosine 2A receptor, PP2A3, and PP2A4.

In certain embodiments, instead of knocking in a gene for the expression of a functional protein, a different gene can be modulated through genetic engineering described herein that upregulates the expression of the gene that would otherwise be knocked in.

5 In certain embodiments, instead of knocking out or knocking out a gene, a different gene can be modulated through genetic engineering described herein that downregulates the expression of the gene that would otherwise be knocked out.

The CD8 Product manufacturing process involves electroporation of dual ribonucleoprotein species of CRISPR-Cas9 nucleases bound to guide RNA sequences, with each species targeting the genomic TCR $\alpha$  and the genomic TCR $\beta$  loci. The specificity of targeting Cas9 nucleases to each genomic locus has been previously described in the literature as being highly specific. Comprehensive testing of the CD8 Product was performed *in vitro* and *in silico* analyses to survey possible off-target genomic cleavage sites, using COSMID and GUIDE-seq, respectively. Multiple CD8 Product or comparable cell products from healthy donors were assessed for cleavage of the candidate off-target sites by deep sequencing, supporting the published evidence that the selected nucleases are highly specific.

In certain embodiments, the CD8 Products described herein can be T cells, NK cells, NKT cells, macrophages, hematopoietic stem cells (HSCs), cells derived from HSCs, or dendritic/antigen-presenting cells.

In certain embodiments, CD8 Cells are expanded in a manner that preserves a “young” T cell phenotypes, resulting in a CD8 Product in which the majority of the T cells exhibit T memory stem cell and T central memory phenotypes

These ‘young’ or ‘younger’ or less-differentiated T cell phenotypes are described to confer improved engraftment potential and prolonged persistence post-infusion. Thus, the administration of CD8 Product, consisting significantly of ‘young’ T cell phenotypes, has the potential to benefit patients with cancer, through improved engraftment potential, prolonged persistence post-infusion, and rapid differentiation into effector T cells to eradicate tumor cells throughout the body.

In certain embodiments, the CD8 Cells of the CD8 Products predominantly comprise memory stem cells (T<sub>msc</sub>) and/or central memory cells (T<sub>cm</sub>). In certain embodiments, at least 25% of the CD8 Cells of the CD8 Products comprise memory stem cells (T<sub>msc</sub>) and/or central memory cells (T<sub>cm</sub>). In certain embodiments, at least 30% of the CD8 Cells of the CD8 Products comprise memory stem cells (T<sub>msc</sub>) and/or central

memory cells (Tcm). In certain embodiments, at least 35% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 40% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 45% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 50% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 55% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 60% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 65% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 70% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, at least 75% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). In certain embodiments, greater than 75% of the CD8 Cells of the CD8 Products comprise memory stem cells (Tmsc) and/or central memory cells (Tcm). Tmsc are characterized as cells that are CD45RA+CD62L+, CD28+CD95+, and CCR7+CD27+. Tcm are characterized as cells that are CD45RO+CD62L+, CD28+CD95+, and CCR7+CD27+CD127+. Both Tmsc and Tcm are characterized as having weak effector T cell function, robust proliferation, robust engraftment, and long telomeres.

In certain embodiments, the CD8 cells disclosed herein show improved properties (e.g., killing activity, cell proliferation, secretion of cytokines, LCK affinity, persistence, tumor infiltration ability) about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 100%, about 150%, about 200%, about 250%, about 300%, about 350%, about 400%, about 450%, or about 500% as compared to the cells that do not have a CD8 construct.

## **METHODS OF PRODUCING CD8 PRODUCTS WITH A YOUNG PHENOTYPE**

In certain embodiments, the present disclosure relates, in part, on the production of engineered “young” T cells. In certain embodiments, the present disclosure comprises methods for producing antigen-specific cells, e.g., T cells, *ex vivo*, comprising activating,

engineering, and expanding antigen-specific cells originally obtained from a subject or isolated from such sample.

In certain embodiments, the methods for activating cells comprise the steps of activating the TCR/CD3 complex. For example, without limitation, the T cells can be  
5 incubated and/or cultured with CD3 agonists, CD28 agonists, or a combination thereof.

In certain embodiments, engineered activated antigen-specific cells, e.g., engineered activated T cells, can be expanded by culturing the engineered activated antigen-specific cells, e.g., T cells, with cytokines, chemokine, soluble peptides, or combination thereof. In certain embodiments, the engineered activated antigen-specific  
10 cells, e.g., engineered activated T cells, can be cultured with one or more cytokines. In certain embodiments, the cytokines can be IL2, IL7, IL15, or combinations thereof. For example, engineered activated antigen-specific cells, e.g., engineered activated T cells, can be cultured with IL7 and IL15. In certain embodiments, the cytokine used in connection with the engineered activated antigen-specific cell, e.g., engineered activated T cell,  
15 culture can be present at a concentration from about 1 pg/ml to about 1 g/ml, from about 1 ng/ml to about 1 g/ml, from about 1 µg/ml to about 1 g/ml, or from about 1 mg/ml to about 1g/ml, and any values in between.

## **PHARMACEUTICAL FORMULATIONS.**

Pharmaceutical formulations of the CD8 Product are prepared by combining the  
20 CD8 Cells in a solution that can preserve the 'young' phenotype of the cells in a cryopreserved state. Table 1 provides an example of one such pharmaceutical formulation. Alternatively, pharmaceutical formulations of the CD8 Product can be prepared by combining the CD8 Cells in a solution that can preserve the 'young' phenotype of the cells without the need to freeze or cryopreserve the product (*i.e.*, the CD8 Product is maintained  
25 in an aqueous solution or as a non-frozen/cryopreserved cell pellet).

Additional pharmaceutically acceptable carriers, buffers, stabilizers, and/or preservatives can also be added to the cryopreservation solution or the aqueous storage solution (if the CD8 Product is not cryopreserved). Any cryopreservation agent and/or media can be used to cryopreserve the CD8 Product, including but not limited to CryoStor,  
30 CryoStor CS5, CELLBANKER, and custom cryopreservation media that optionally include DMSO.

## GENE-EDITING METHODS

In certain embodiments, the present disclosure involves, in part, methods of engineering human cells, e.g., engineered T cells or engineered human stem cells. In certain embodiments, the present disclosure involves, in part, methods of engineering  
5 human cells, e.g., NK cells, NKT cells, macrophages, hematopoietic stem cells (HSCs), cells derived from HSCs, or dendritic/antigen-presenting cells. In certain embodiments, such engineering involves genome editing. For example, but not by way of limitation, such genome editing can be accomplished with nucleases targeting one or more endogenous loci, e.g., TCR alpha (TCR $\alpha$ ) locus and TCR beta (TCR $\beta$ ) locus. In certain  
10 embodiments, the nucleases can generate single-stranded DNA nicks or double-stranded DNA breaks in an endogenous target sequence. In certain embodiments, the nuclease can target coding or non-coding portions of the genome, e.g., exons, introns. In certain embodiments, the nucleases contemplated herein comprise homing endonuclease, meganuclease, megaTAL nuclease, transcription activator-like effector nuclease  
15 (TALEN), zinc-finger nuclease (ZFN), and clustered regularly interspaced short palindromic repeats (CRISPR)/Cas nuclease. In certain embodiments, the nucleases can themselves be engineered, e.g., via the introduction of amino acid substitutions and/or deletions, to increase the efficiency of the cutting activity.

In certain embodiments, a CRISPR/Cas nuclease system is used to engineer human  
20 cells. In certain embodiments, the CRISPR/Cas nuclease system comprises a Cas nuclease and one or more RNAs that recruit the Cas nuclease to the endogenous target sequence, e.g., single guide RNA. In certain embodiments, the Cas nuclease and the RNA are introduced in the cell separately, e.g. using different vectors or compositions, or together, e.g., in a polycistronic construct or a single protein-RNA complex. In certain  
25 embodiments, the Cas nuclease is Cas9 or Cas12a. In certain embodiments, the Cas9 polypeptide is obtained from a bacterial species including, without limitation, *Streptococcus pyogenes* or *Neisseria meningitidis*. Additional examples of CRISPR/Cas systems are known in the art. See Adli, Mazhar. "The CRISPR tool kit for genome editing and beyond." Nature communications vol. 9,1 1911 (2018), herein incorporated by  
30 reference for all that it teaches.

In certain embodiments, genome editing occurs at one or more genome loci that regulate immunological responses. In certain embodiments, the loci include, without limitation, TCR alpha (TCR $\alpha$ ) locus, TCR beta (TCR $\beta$ ) locus, TCR gamma (TCR $\gamma$ ), and

TCR delta (TCR $\delta$ ). In certain embodiments, the loci for inserting a CD8 Construct is anywhere in the genome. In certain embodiments, the loci for inserting a CD8 Construct is the TRAC locus. In certain embodiments, the loci for inserting a CD8 Construct is one of the two TRBC loci. In certain embodiments, the locus for inserting a CD8 Construct is a locus other than the TRAC locus or TRAB loci. In certain embodiments, the loci for inserting a CD8 Construct is inserted into a gene locus wherein such gene is knocked out. By way of a non-limiting example, if the desired phenotype of a CD8 Product is the expression of a NeoTCR, the expression of a CD8 Construct, and the knockout of the TET2 gene or the AAVS1 gene, the CD8 Construct can be inserted at the TET2 locus or AAVS1 locus. In certain embodiments, the insertion of the CD8 Construct is in tandem with the NeoTCR insertion. In certain embodiments, the insertion of the CD8 Construct is a separate locus than the NeoTCR insertion.

In certain embodiments, genome editing is performed by using non-viral delivery systems. For example, a nucleic acid molecule can be introduced into a cell by administering the nucleic acid in the presence of lipofection (Feigner et al., Proc. Natl. Acad. Sci. U.S.A. 84:7413, 1987; Ono et al., Neuroscience Letters 17:259, 1990; Brigham et al., Am. J. Med. Sci. 298:278, 1989; Staubinger et al., Methods in Enzymology 101:512, 1983), asialoorosomucoid-polylysine conjugation (Wu et al., Journal of Biological Chemistry 263:14621, 1988; Wu et al., Journal of Biological Chemistry 264:16985, 1989), or by micro-injection under surgical conditions (Wolff et al., Science 247:1465, 1990). Other non-viral means for gene transfer include transfection in vitro using calcium phosphate, DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell. Transplantation of normal genes into the affected tissues of a subject can also be accomplished by transferring a normal nucleic acid into a cultivatable cell type ex vivo (e.g., an autologous or heterologous primary cell or progeny thereof), after which the cell (or its descendants) are injected into a targeted tissue or are injected systemically.

In certain embodiments, genome editing is performed by using viral delivery systems. In certain embodiments, the viral methods include targeted integration (including but not limited to AAV) and random integration (including but not limited to lentiviral approaches). In certain embodiments, the viral delivery would be accomplished without integration of the nuclease. In such embodiments, the viral delivery system can be Lentiflash or another similar delivery system.

## HOMOLOGY RECOMBINATION TEMPLATES

In certain embodiments, the present disclosure provides genome editing of a cell by introducing and recombining a homologous recombination (HR) template nucleic acid sequence into an endogenous locus of a cell. In certain embodiments, the HR template nucleic acid sequence is linear. In certain embodiments, the HR template nucleic acid sequence is circular. In certain embodiments, the circular HR template can be a plasmid, minicircle, or nanoplasmid. In certain embodiments, the HR template nucleic acid sequence comprises a first and a second homology arms. In certain embodiments, the homology arms can be of about 300 bases to about 2,000 bases. For example, each homology arm can be 1,000 bases. In certain embodiments, the homology arms can be homologous to a first and second endogenous sequences of the cell. In certain embodiments, the endogenous locus is a TCR locus. For example, the first and second endogenous sequences are within a TCR alpha locus or a TCR beta locus. In certain embodiments, the HR template comprises a TCR gene sequences. In non-limiting embodiments, the TCR gene sequence is a patient specific TCR gene sequence. In non-limiting embodiments, the TCR gene sequence is tumor-specific. In non-limiting embodiments, the TCR gene sequence can be identified and obtained using the methods described in PCT/US2020/017887, the content of which is herein incorporated by reference. In certain embodiments, the HR template comprises a TCR alpha gene sequence and a TCR beta gene sequence.

In certain embodiments, the HR template is a polycistronic polynucleotide. In certain embodiments, the HR template comprises sequences encoding for flexible polypeptide sequences (e.g., Gly-Ser-Gly sequence). In certain embodiments, the HR template comprises sequences encoding an internal ribosome entry site (IRES). In certain embodiments, the HR template comprises a 2A peptide (e.g., P2A, T2A, E2A, and F2A). Additional information on the HR template nucleic acids and methods of modifying a cell thereof can be found in International Patent Application no. PCT/US2018/058230, the content of which is herein incorporated by reference.

## METHODS OF TREATMENT

The presently disclosed subject matter provides methods for inducing and/or increasing an immune response in a subject in need thereof. The CD8 Products can be

used for treating and/or preventing a cancer in a subject. The CD8 Products can be used for prolonging the survival of a subject suffering from a cancer. The CD8 Products can also be used for treating and/or preventing a cancer in a subject. The CD8 Products can also be used for reducing tumor burden in a subject. Such methods comprise administering the CD8 Products in an amount effective or a composition (*e.g.*, a pharmaceutical composition) comprising thereof to achieve the desired effect, be it palliation of an existing condition or prevention of recurrence. For treatment, the amount administered is an amount effective in producing the desired effect. An effective amount can be provided in one or a series of administrations. An effective amount can be provided in a bolus or by continuous perfusion.

In certain embodiments, the CD8 Products can be used for treating viral or bacterial diseases. In certain embodiments, the CD8 Products can be used for treating autoimmune diseases.

In certain embodiments, an effective amount of the CD8 Products are delivered through IV administration. In certain embodiments, the CD8 Products are delivered through IV administration in a single administration. In certain embodiments, the CD8 Products are delivered through IV administration in multiple administrations. In certain embodiments, the CD8 Products are delivered through IV administration in two or more administrations. In certain embodiments, the CD8 Products are delivered through IV administration in two administrations. In certain embodiments, the CD8 Products are delivered through IV administration in three administrations.

The presently disclosed subject matter provides methods for treating and/or preventing cancer in a subject. In certain embodiments, the method comprises administering an effective amount of CD8 Products to a subject having cancer.

Non-limiting examples of cancer include blood cancers (*e.g.* leukemias, lymphomas, and myelomas), ovarian cancer, breast cancer, bladder cancer, brain cancer, colon cancer, intestinal cancer, liver cancer, lung cancer, pancreatic cancer, prostate cancer, skin cancer, stomach cancer, glioblastoma, throat cancer, melanoma, neuroblastoma, adenocarcinoma, glioma, soft tissue sarcoma, and various carcinomas (including prostate and small cell lung cancer). Suitable carcinomas further include any known in the field of oncology, including, but not limited to, astrocytoma, fibrosarcoma, myxosarcoma, liposarcoma, oligodendroglioma, ependymoma, medulloblastoma, primitive neural ectodermal tumor (PNET), chondrosarcoma, osteogenic sarcoma, pancreatic ductal adenocarcinoma, small and large cell lung adenocarcinomas, chordoma,

angiosarcoma, endotheliosarcoma, squamous cell carcinoma, bronchoalveolarcarcinoma, epithelial adenocarcinoma, and liver metastases thereof, lymphangiosarcoma, lymphangi endotheliosarcoma, hepatoma, cholangiocarcinoma, synovioma, mesothelioma, Ewing's tumor, rhabdomyosarcoma, colon carcinoma, basal cell carcinoma, sweat gland carcinoma, papillary carcinoma, sebaceous gland carcinoma, papillary adenocarcinoma, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, testicular tumor, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, neuroblastoma, retinoblastoma, leukemia, multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease, breast tumors such as ductal and lobular adenocarcinoma, squamous and adenocarcinomas of the uterine cervix, uterine and ovarian epithelial carcinomas, prostatic adenocarcinomas, transitional squamous cell carcinoma of the bladder, B and T cell lymphomas (nodular and diffuse) plasmacytoma, acute and chronic leukemias, malignant melanoma, soft tissue sarcomas and leiomyosarcomas. In certain embodiments, the neoplasia is selected from the group consisting of blood cancers (e.g. leukemias, lymphomas, and myelomas), ovarian cancer, prostate cancer, breast cancer, bladder cancer, brain cancer, colon cancer, intestinal cancer, liver cancer, lung cancer, pancreatic cancer, prostate cancer, skin cancer, stomach cancer, glioblastoma, and throat cancer. In certain embodiments, the presently disclosed young T cells and compositions comprising thereof can be used for treating and/or preventing blood cancers (e.g., leukemias, lymphomas, and myelomas) or ovarian cancer, which are not amenable to conventional therapeutic interventions.

In certain embodiments, the neoplasia is a solid cancer or a solid tumor. In certain embodiments, the solid tumor or solid cancer is selected from the group consisting of glioblastoma, prostate adenocarcinoma, kidney papillary cell carcinoma, sarcoma, ovarian cancer, pancreatic adenocarcinoma, rectum adenocarcinoma, colon adenocarcinoma, esophageal carcinoma, uterine corpus endometrioid carcinoma, breast cancer, skin cutaneous melanoma, lung adenocarcinoma, stomach adenocarcinoma, cervical and endocervical cancer, kidney clear cell carcinoma, testicular germ cell tumors, and aggressive B-cell lymphomas.

The subjects can have an advanced form of disease, in which case the treatment objective can include mitigation or reversal of disease progression, and/or amelioration of side effects. The subjects can have a history of the condition, for which they have already

been treated, in which case the therapeutic objective will typically include a decrease or delay in the risk of recurrence.

Suitable human subjects for therapy typically comprise two treatment groups that can be distinguished by clinical criteria. Subjects with “advanced disease” or “high tumor burden” are those who bear a clinically measurable tumor. A clinically measurable tumor is one that can be detected on the basis of tumor mass (e.g., by palpation, CAT scan, sonogram, mammogram or X-ray; positive biochemical or histopathologic markers on their own are insufficient to identify this population). A pharmaceutical composition is administered to these subjects to elicit an anti-tumor response, with the objective of palliating their condition. Ideally, reduction in tumor mass occurs as a result, but any clinical improvement constitutes a benefit. Clinical improvement includes decreased risk or rate of progression or reduction in pathological consequences of the tumor.

#### **ARTICLES OF MANUFACTURE**

The CD8 Products can be used in combination with articles of manufacture. Such articles of manufacture can be useful for the prevention or treatment of proliferative disorders (e.g., cancer). Examples of articles of manufacture include but are not limited to containers (e.g., infusion bags, bottles, storage containers, flasks, vials, syringes, tubes, and IV solution bags) and a label or package insert on or associated with the container. The containers may be made of any material that is acceptable for the storage and preservation of the CD8 Cells within the CD8 Products. In certain embodiments, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle. For example, the container may be a CryoMACS freezing bag. The label or package insert indicates that the CD8 Products are used for treating the condition of choice and the patient of origin. The patient is identified on the container of the CD8 Product because the CD8 Products is made from autologous cells and engineered as a patient-specific and individualized treatment.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; and 2) a second container with the same CD8 Product as the first container contained therein. Optionally, additional containers with the same CD8 Product as the first and second containers may be prepared and made. Optionally, additional

containers containing a composition comprising a different cytotoxic or otherwise therapeutic agent may also be combined with the containers described above.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; and 2) a second container with a composition contained therein, wherein  
5 the composition comprises a further cytotoxic or otherwise therapeutic agent.

The article of manufacture may comprise: 1) a first container with two CD8 Products contained therein; and 2) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent.

10 The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; and 3) optionally a third container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. In certain embodiments, the first and second CD8 Products are different CD8 Products. In certain  
15 embodiments, the first and second CD8 Products are the same CD8 Products.

The article of manufacture may comprise: 1) a first container with three CD8 Products contained therein; and 2) optionally a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent.

20 The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; 3) a third container with a third CD8 Product contained therein; and 4) optionally a fourth container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. In certain embodiments, the first, second,  
25 and third CD8 Products are different CD8 Products. In certain embodiments, the first, second, and third CD8 Products are the same CD8 Products. In certain embodiments, two of the first, second, and third CD8 Products are the same CD8 Products.

The article of manufacture may comprise: 1) a first container with four CD8 Products contained therein; and 2) optionally a second container with a composition  
30 contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; 3) a third container with a third CD8 Product contained therein; 4) a fourth container with a

fourth CD8 Product contained therein; and 5) optionally a fifth container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. In certain embodiments, the first, second, third, and fourth CD8 Products are different CD8 Products. In certain embodiments, the first, second, third, and fourth CD8 Products are the same NeoTCR Products. In certain embodiments, two of the first, second, third, and fourth CD8 Products are the same NeoTCR Products. In certain embodiments, three of the first, second, third, and fourth CD8 Products are the same CD8 Products.

The article of manufacture may comprise: 1) a first container with five or more CD8 Products contained therein; and 2) optionally a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; 3) a third container with a third CD8 Product contained therein; 4) a fourth container with a fourth CD8 Product contained therein; 5) a fifth container with a fifth CD8 Product contained therein; 6) optionally a sixth or more additional containers with a sixth or more CD8 Product contained therein; and 7) optionally an additional container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. In certain embodiments, all of the containers of CD8 Products are different CD8 Products. In certain embodiments, all of the containers of CD8 Products are the same CD8 Products. In certain embodiments, there can be any combination of same or different CD8 Products in the five or more containers based on the availability of detectable CD8s in a patient's tumor sample(s), the need and/or desire to have multiple CD8 Products for the patient, and the availability of any one CD8 Product that may require or benefit from one or more container.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; and 3) a third container with a third CD8 Product contained therein.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; 3) a third container with a third CD8 Product contained therein; and 4) optionally a fourth container with a fourth CD8 Product contained therein.

The article of manufacture may comprise: 1) a first container with a CD8 Product contained therein; 2) a second container with a second CD8 Product contained therein; 3) a third container with a third CD8 Product contained therein; 4) a fourth container with a fourth CD8 Product contained therein; and 5) optionally a fifth container with a fourth CD8 Product contained therein.

The article of manufacture may comprise a container with one CD8 Product contained therein. The article of manufacture may comprise a container with two CD8 Products contained therein. The article of manufacture may comprise a container with three CD8 Products contained therein. The article of manufacture may comprise a container with four CD8 Products contained therein. The article of manufacture may comprise a container with five CD8 Products contained therein.

The article of manufacture may comprise 1) a first container with one CD8 Product contained therein, and 2) a second container with two CD8 Products contained therein. The article of manufacture may comprise 1) a first container with two CD8 Products contained therein, and 2) a second container with one CD8 Product contained therein. In the examples above, a third and/or fourth container comprising one or more additional CD8 Products may be included in the article of manufacture. Additionally, a fifth container comprising one or more additional CD8 Products may be included in the article of manufacture.

Furthermore, any container of CD8 Product described herein can be split into two, three, or four separate containers for multiple time points of administration and/or based on the appropriate dose for the patient.

In certain embodiments, the CD8 Products are provided in a kit. The kit can, by means of non-limiting examples, contain package insert(s), labels, instructions for using the CD8 Product(s), syringes, disposal instructions, administration instructions, tubing, needles, and anything else a clinician would need in order to properly administer the CD8 Product(s).

## **THERAPEUTIC COMPOSITIONS AND METHODS OF MANUFACTURING**

As described herein, plasmid DNA-mediated precision genome engineering process for Good Manufacturing Practice (GMP) manufacturing of CD8 Products was developed. Targeted integration of the patient-specific neoTCR was accomplished by electroporating CRISPR endonuclease ribonucleoproteins (RNPs) together with the

personalized neoTCR gene cassette, encoded by the plasmid DNA. In addition to the neoTCR, the CD8 Constructs were inserted by incorporating them into the neoTCR vector and then electroporating with CRISPR endonuclease ribonucleoproteins (RNPs) as described above.

5           The CD8 Products can be formulated into a drug product using the clinical manufacturing process. Under this process, the CD8 Products are cryopreserved in CryoMACS Freezing Bags. One or more bags may be shipped to the site for each patient depending on patient needs. The product is composed of apheresis-derived, patient-autologous, CD8 and CD4 T cells that have been precision genome engineered to express  
10 one or more autologous neoTCRs targeting a neoepitope complexed to one of the endogenous HLA receptors presented exclusively on the surface of that patient’s tumor cells.

The final product will contain 5% dimethyl sulfoxide (DMSO), human serum albumin, and Plasma-Lyte. The final cell product will contain the list of components  
15 provided in **Table 2. Composition of the CD8 Product**

Component	Specification/Grade
Total nucleated NeoTCR cells	cGMP manufactured
Plasma-Lyte A	USP
Human Serum Albumin in 0.02 - 0.08 M sodium caprylate and sodium tryptophanate	USP
CryoStor CS10	cGMP manufactured with USP grade materials

**COMPOSITIONS AND VECTORS**

The presently disclosed subject matter provides compositions comprising cells (e.g., immunoresponsive cells) disclosed herein.

20           In certain embodiments, the presently disclosed subject matter provides nucleic acid compositions comprising a polynucleotide encoding the NeoTCR disclosed herein. In certain embodiments, the nucleic acid compositions disclosed herein comprise a polynucleotide encoding a CD8 Construct disclosed herein. Also provided are cells comprising such nucleic acid compositions.

25           In certain embodiments, the nucleic acid composition further comprises a promoter that is operably linked to the NeoTCR disclosed herein. In certain embodiments, the

nucleic acid composition further comprises a promoter that is operably linked to the CD8 Construct disclosed herein.

In certain embodiments, the promoter is endogenous or exogenous. In certain embodiments, the exogenous promoter is selected from the group consisting of an elongation factor (EF)-1 promoter, a CMV promoter, a SV40 promoter, a PGK promoter, a long terminal repeat (LTR) promoter and a metallothionein promoter. In certain  
5  
embodiments, the promoter is an inducible promoter. In certain embodiments, the inducible promoter is selected from the group consisting of a NFAT transcriptional response element (TRE) promoter, a CD69 promoter, a CD25 promoter, an IL-2 promoter,  
10  
an IL-12 promoter, a p40 promoter, and a Bcl-xL promoter.

The compositions and nucleic acid compositions can be administered to subjects or and/delivered into cells by art-known methods or as described herein. Genetic modification of a cell (e.g., a T cell) can be accomplished by transducing a substantially homogeneous cell composition with a recombinant DNA construct. In certain  
15  
embodiments, a retroviral vector (either a gamma-retroviral vector or a lentiviral vector) is employed for the introduction of the DNA construct into the cell. Non-viral vectors may be used as well.

Possible methods of transduction also include direct co-culture of the cells with producer cells, e.g., by the method of Bregni, *et al.* (1992) *Blood* 80:1418-1422, or  
20  
culturing with viral supernatant alone or concentrated vector stocks with or without appropriate growth factors and polycations, e.g., by the method of Xu, *et al.* (1994) *Exp. Hemat.* 22:223-230; and Hughes, *et al.* (1992) *J. Clin. Invest.* 89:1817.

Other transducing viral vectors can be used to modify a cell. In certain  
25  
embodiments, the chosen vector exhibits high efficiency of infection and stable integration and expression (see, e.g., Cayouette *et al.*, *Human Gene Therapy* 8:423-430, 1997; Kido *et al.*, *Current Eye Research* 15:833-844, 1996; Bloomer *et al.*, *Journal of Virology* 71:6641-6649, 1997; Naldini *et al.*, *Science* 272:263-267, 1996; and Miyoshi *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 94:10319, 1997). Other viral vectors that can be used include, for example, adenoviral, lentiviral, and adena-associated viral vectors, vaccinia virus, a bovine papilloma virus, or a herpes virus, such as Epstein-Barr Virus (also see, for example, the  
30  
vectors of Miller, *Human Gene Therapy* 15-14, 1990; Friedman, *Science* 244:1275-1281, 1989; Eglitis *et al.*, *BioTechniques* 6:608-614, 1988; Tolstoshev *et al.*, *Current Opinion in Biotechnology* 1:55-61, 1990; Sharp, *The Lancet* 337:1277-1278, 1991; Cornetta *et al.*, *Nucleic Acid Research and Molecular Biology* 36:311-322, 1987; Anderson, *Science*

226:401-409, 1984; Moen, Blood Cells 17:407-416, 1991; Miller et al., Biotechnology 7:980-990, 1989; LeGal La Salle et al., Science 259:988-990, 1993; and Johnson, Chest 107:77S- 83S, 1995). Retroviral vectors are particularly well developed and have been used in clinical settings (Rosenberg et al., N. Engl. J. Med 323:370, 1990; Anderson et al.,  
5 U.S. Pat. No. 5,399,346).

Non-viral approaches can also be employed for genetic modification of a cell. For example, a nucleic acid molecule can be introduced into a cell by administering the nucleic acid in the presence of lipofection (Feigner et al., Proc. Natl. Acad. Sci. U.S.A. 84:7413, 1987; Ono et al., Neuroscience Letters 17:259, 1990; Brigham et al., Am. J. Med. Sci.  
10 298:278, 1989; Staubinger et al., Methods in Enzymology 101:512, 1983), asialoorosomuroid-polylysine conjugation (Wu et al., Journal of Biological Chemistry 263:14621, 1988; Wu et al., Journal of Biological Chemistry 264:16985, 1989), or by micro-injection under surgical conditions (Wolff et al., Science 247:1465, 1990). Other non-viral means for gene transfer include transfection *in vitro* using calcium phosphate,  
15 DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell. Transplantation of normal genes into the affected tissues of a subject can also be accomplished by transferring a normal nucleic acid into a cultivatable cell type *ex vivo* (e.g., an autologous or heterologous primary cell or progeny thereof), after which the cell (or its descendants) are injected into a targeted tissue  
20 or are injected systemically.

Polynucleotide therapy methods can be directed from any suitable promoter (e.g., the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element or intron (e.g. the elongation factor 1a enhancer/promoter/intron structure). For example, if desired,  
25 enhancers known to preferentially direct gene expression in specific cell types can be used to direct the expression of a nucleic acid. The enhancers used can include, without limitation, those that are characterized as tissue- or cell-specific enhancers. Alternatively, if a genomic clone is used as a therapeutic construct, regulation can be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a  
30 heterologous source, including any of the promoters or regulatory elements described above.

The resulting cells can be grown under conditions similar to those for unmodified cells, whereby the modified cells can be expanded and used for a variety of purposes.

## KITS

The presently disclosed subject matter provides kits for inducing and/or enhancing an immune response and/or treating and/or preventing a cancer or a pathogen infection in a subject. In certain embodiments, the kit comprises an effective amount of presently disclosed cells or a pharmaceutical composition comprising thereof. In certain  
5       embodiments, the kit comprises a sterile container; such containers can be boxes, ampules, bottles, vials, tubes, bags, pouches, blister-packs, or other suitable container forms known in the art. Such containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding medicaments. In certain non-limiting embodiments,  
10       the kit includes an isolated nucleic acid molecule encoding a presently disclosed HR template.

If desired, the cells and/or nucleic acid molecules are provided together with instructions for administering the cells or nucleic acid molecules to a subject having or at risk of developing a cancer or pathogen or immune disorder. The instructions generally  
15       include information about the use of the composition for the treatment and/or prevention of a cancer or a pathogen infection. In certain embodiments, the instructions include at least one of the following: description of the therapeutic agent; dosage schedule and administration for treatment or prevention of a neoplasia, pathogen infection, or immune disorder or symptoms thereof; precautions; warnings; indications; counter-indications;  
20       over-dosage information; adverse reactions; animal pharmacology; clinical studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder supplied in or with the container. The resulting cells can be grown under conditions similar to those  
25       of purposes.

## EXEMPLARY EMBODIMENTS

A. In certain non-limiting embodiments, the presently disclosed subject matter provides for a cell, comprising an exogenous T cell receptor (TCR), and an exogenous  
30       CD8.

- A1. The foregoing of A, wherein the exogenous CD8 comprises at least one monomer.
- A2. The foregoing cell of A1, wherein the at least one monomer of the exogenous CD8 comprises an extracellular domain, a transmembrane domain, an intracellular domain, fragments thereof, or combinations thereof.
- 5 A3. The foregoing cell of A2, wherein the extracellular domain comprises a CD8 $\alpha$  extracellular domain or a CD8 $\beta$  extracellular domain.
- A4. The foregoing cell of A2 or A3, wherein the transmembrane a CD8 $\alpha$  transmembrane domain or a CD8 $\beta$  transmembrane domain.
- 10 A5. The foregoing cell of A2-A4, wherein the intracellular domain comprises a CD8 $\alpha$  intracellular domain or a CD8 $\beta$  intracellular domain.
- A6. The foregoing cell of A2-A4, wherein the intracellular domain comprises a CD4 intracellular domain.
- A7. The foregoing cell of A1-A5, wherein the at least one monomer comprises a
- 15 CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain.
- A8. The foregoing cell of A1-A5, wherein the at least one monomer comprises a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain.
- 20 A9. The foregoing cell of A1-A5, wherein the at least one monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain.
- A10. The foregoing cell of A1-A6, wherein the at least one monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular
- 25 domain.
- A11. The foregoing cell of A1-A10, wherein the at least one monomer comprises a signal peptide.
- A12. The foregoing cell of A11, wherein the signal peptide is a CD8 signal peptide.
- 30 A13. The foregoing cell of A2-A12, wherein the extracellular domain comprises the amino acid sequence set forth in SEQ ID NO: 140, or SEQ ID NO: 145.
- A14. The foregoing cell of A2-A13, wherein the transmembrane domain comprises the amino acid sequence set forth in SEQ ID NO: 141, or SEQ ID NO: 146.

A15. The foregoing cell of A2-A14, wherein the intracellular domain comprises the amino acid sequence set forth in SEQ ID NO: 142, SEQ ID NO: 147, or SEQ ID NO: 148.

5 A16. The foregoing cell of A11-A15, wherein the signal peptide comprises the amino acid sequence set forth in SEQ ID NO: 139, or SEQ ID NO: 144.

A17. The foregoing cell of A-A16, wherein the exogenous CD8 comprises a 2A sequence.

A18. The foregoing cell of A-A17, wherein the exogenous CD8 comprises a linker.

10 A19. The foregoing cell of A18, wherein the linker comprises the amino acid sequence set forth in SEQ ID NO: 137.

A20. The foregoing cell of A-A19, wherein the exogenous CD8 comprises a protease cleavage site.

15 A21. The foregoing cell of A-A20, wherein the protease cleavage site is a Furin cleavage site.

A22. The foregoing cell of A-A21, wherein the exogenous TCR is a patient derived TCR.

A23. The foregoing cell of A-A22, wherein the exogenous TCR comprises a signal sequence, a first and second 2A sequence, and a TCR polypeptide sequence.

20 A24. The foregoing cell of A-A23, wherein the exogenous TCR recognizes a cancer antigen.

A25. The foregoing cell of A24, wherein the cancer antigen is a neoantigen.

A26. The foregoing cell of A24, wherein the cancer antigen is a patient specific antigen.

25 A27. The foregoing cell of A-A26, wherein the cell is a primary cell.

A28. The foregoing cell of A-A26, wherein the cell is a patient-derived cell.

A29. The foregoing cell of A-A26, wherein the cell is a lymphocyte.

A30. The foregoing cell of A-A26, wherein the cell is a T cell.

A31. The foregoing cell of A-A26, wherein the cell if a young T cell.

30 A32. The foregoing cell of A31, wherein the cell is CD45RA+, CD62L+, CD28+, CD95-, CCR7+, and CD27+.

A33. The foregoing cell of A31, wherein the cell is CD45RA+, CD62L+, CD28+, CD95+, CD27+, CCR7+.

A34. The foregoing cell of A31, wherein the cell is CD45RO+, CD62L+, CD28+, CD95+, CCR7+, CD27+, CD127+.

A35. The foregoing cell of A-A34, further comprising a gene modification to enhance cell persistence and/or enhances memory cell differentiation

5 A36. The foregoing cell of A-A35, wherein killing activity of the cell is increased between about 10% to about 500% as compared to killing activity of a cell that does not have the exogenous CD8.

10 A37. The foregoing cell of A-A36, wherein proliferation of the cell upon binding of the TCR to the antigen is increased between about 10% to about 500% as compared to proliferation of a cell that does not have the exogenous CD8.

A38. The foregoing cell of A-A37, wherein secretion of pro-inflammatory cytokine upon binding of the TCR to the antigen by the cell is increased between about 10% to about 500% as compared to secretion by a cell that does not have the exogenous CD8.

15 A39. The foregoing cell of A-A38, wherein LCK affinity of the cell is increased between about 10% to about 500% as compared to LCK affinity of a cell that does not have the exogenous CD8.

20 A40. The foregoing cell of A-A39, wherein persistence of the cell is increased between about 10% to about 500% as compared to persistence of a cell that does not have the exogenous CD8.

A41. The foregoing cell of A-A40, wherein tumor infiltration ability of the cell is increased between about 10% to about 500% as compared to tumor infiltration ability of a cell that does not have the exogenous CD8.

25 A42. The foregoing cell of A-A41, wherein the exogenous TCR is a CD8-dependent TCR.

A43. The foregoing cell of A-A41, wherein the exogenous TCR is a CD8-independent TCR.

A44. The foregoing cell of A-A43, wherein the exogenous CD8 is encoded by a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4.

30 A45. The foregoing cell of A-A43, wherein the exogenous CD8 comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane

domain, and aCD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

A46. The foregoing cell of A-A43, wherein the exogenous CD8 comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140),  
5 a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular  
15 domain (SEQ ID NO: 148).

B. In certain non-limiting embodiments, the presently disclosed subject matter provides for a method of modifying a cell, the method comprising introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises first and second homology arms homologous to first and second target  
20 nucleic acid sequences, a TCR gene sequence positioned between the first and second homology arms, and a CD8 gene sequence positioned between the first and the second homology arms, and recombining the HR template nucleic acid into an endogenous locus of the cell.

B1. The foregoing method of B, wherein the HR template comprises a first 2A-coding sequence positioned upstream of the CD8 gene sequence, a second 2A-coding sequence positioned downstream of the CD8 gene sequence and upstream of the TCR gene sequence, and a third 2A-coding sequence positioned downstream of the TCR gene sequence; wherein the first, second, and third 2A-coding sequences code for the same amino acid sequence and are codon-diverged relative to each other.  
25

B2. The foregoing method of B or B1, wherein the HR template comprises a sequence coding for the amino acid sequence Gly Ser Gly positioned immediately upstream of the first, second, and/or third 2A-coding sequences.  
30

B3. The foregoing method of B1 or B2, wherein the HR template further comprises a sequence coding for a Furin cleavage site positioned upstream of the first, second, and/or third 2A-coding sequences.

5 B4. The foregoing method of B-B3, wherein the HR template further comprises a sequence encoding a signal sequence positioned immediately upstream of the TCR gene sequence and/or the CD8 gene sequence.

B5. The foregoing method of B-B4, wherein the HR template comprises a second TCR sequence positioned between the third 2A-coding sequence and the second homology arm.

10 B6. The foregoing method of B5, wherein the HR template comprises a sequence encoding a first signal sequence positioned immediately upstream the first TCR gene sequence; and a sequence encoding a second signal sequence positioned immediately upstream of the second TCR gene sequence.

15 B7. The foregoing method of B-B6, wherein the HR template comprises a second CD8 gene sequence positioned between the first CD8 gene sequence and the second 2A-coding sequence.

B8. The foregoing method of B7, wherein a 2A coding sequence is positioned between the first and second CD8 gene sequence.

20 B9. The foregoing method of B7 or B8, wherein a sequence coding for the amino acid sequence Gly Ser Gly is positioned between the first and second CD8 gene sequences.

B10. The foregoing method of B7-B9, wherein a sequence coding for a Furin cleavage site is positioned between the first and second CD8 gene sequences.

25 B11. The foregoing method of B-B10, wherein the CD8 gene sequence comprises a sequence encoding an extracellular domain, a sequence encoding an intracellular domain, a sequence encoding an intracellular domain, fragments thereof, or combinations thereof.

B12. The foregoing method of B11, wherein the sequence encoding an extracellular domain comprises a sequence encoding a CD8 $\alpha$  extracellular domain or a CD8 $\beta$  extracellular domain.

30 B13. The foregoing method of B11 or B12, wherein the sequence encoding a transmembrane domain comprises a sequence encoding a CD8 $\alpha$  transmembrane domain or a CD8 $\beta$  transmembrane domain.

B14. The foregoing method of B11-B13, wherein the sequence encoding an intracellular domain comprises a sequence encoding a CD8 $\alpha$  intracellular domain or a CD8 $\beta$  intracellular domain.

5 B15. The foregoing method of B11-B14, wherein the sequence encoding an intracellular domain comprises a sequence encoding a CD4 intracellular domain.

B16. The foregoing method of B11-B15, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain.

10 B17. The foregoing method of B11-B15, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain.

B18. The foregoing method of B11-B15, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain.

15 B19. The foregoing method of B11-B15, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

20 B20. The foregoing method of B7-B19, wherein the HR template comprises a sequence encoding a first signal sequence positioned immediately upstream the first CD8 gene sequence; and a sequence encoding a second signal sequence positioned immediately upstream of the second CD8 gene sequence.

B21. The foregoing method of B4-B20, wherein the signal sequence is a CD8 signal sequence, a human growth hormone signal sequence, fragments thereof, or combinations thereof.

25 B22. The foregoing method of B-B21, wherein the first and second homology arms of the HR template are each from about 300 bases to about 2,000 bases in length.

B23. The foregoing method of B-B22, wherein the first and second homology arms of the HR template are each from about 600 bases to about 2,000 bases in length.

30 B24. The foregoing method of B-B23, wherein the exogenous TCR is a patient derived TCR.

B25. The foregoing method of B-B24, wherein the exogenous TCR comprises a signal sequence, a first and second 2A sequence, and a TCR polypeptide sequence.

B26. The foregoing method of B-B25, wherein the exogenous TCR recognizes a cancer antigen.

B27. The foregoing method of B26, wherein the cancer antigen is a neoantigen.

B28. The foregoing method of B26, wherein the cancer antigen is a patient specific antigen.

B29. The foregoing method of B-B28, wherein the HR template is non-viral.

5 B30. The foregoing method of B-B29, wherein the HR template is a circular DNA.

B31. The foregoing method of B-B29, wherein the HR template is a linear DNA.

B32. The foregoing method of B-B31, wherein the introducing occurs via electroporation.

10 B33. The foregoing method of B-B32, wherein the recombining comprises cleavage of the endogenous locus by a nuclease; and recombination of the HR template nucleic acid sequence into the endogenous locus by homology directed repair.

B34. The foregoing method of B33, wherein the nuclease is a Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) family nuclease, or derivative thereof.

B35. The foregoing method of B34, further comprising a gRNA.

15 B36. The foregoing method of B-B35, wherein the process further comprises culturing the cell.

B37. The foregoing method of B36, wherein the culturing is conducted in the presence of at least one cytokine.

20 B38. The foregoing method of B36 or B37, wherein the culturing is conducted in the presence of IL2, IL7, IL15, or any combination thereof.

B39. The foregoing method of B36 or B37, wherein the culturing is conducted in the presence of IL7 and IL15.

B40. The foregoing method of B-B39, further comprising a gene modification to enhance cell persistence and/or enhances memory cell differentiation.

25 B41. The foregoing method of B-B40, wherein the cell is a primary cell.

B42. The foregoing method of B-B40, wherein the cell is a patient-derived cell.

B43. The foregoing method of B-B40, wherein the cell is a lymphocyte.

B44. The foregoing method of B-B40, wherein the cell is a T cell.

B45. The foregoing method of B-B40, wherein the cell is a young T cell.

30 B46. The foregoing method of B45, wherein the cell is CD45RA+, CD62L+, CD28+, CD95-, CCR7+, and CD27+.

B47. The foregoing method of B45, wherein the cell is CD45RA+, CD62L+, CD28+, CD95+, CD27+, CCR7+.

B48. The foregoing method of B45, wherein the cell is CD45RO+, CD62L+, CD28+, CD95+, CCR7+, CD27+, CD127+.

5 B49. The foregoing method of B-B48, wherein killing activity of the cell is increased between about 10% to about 500% as compared to killing activity of a cell that does not have the CD8 gene sequence.

B50. The foregoing method of B-B49, wherein proliferation of the cell upon binding of the TCR to the antigen is increased between about 10% to about 500% as compared to proliferation of a cell that does not have the CD8 gene sequence.

10 B51. The foregoing method of B-B50, wherein secretion of pro-inflammatory cytokine upon binding of the TCR to the antigen by the cell is increased between about 10% to about 500% as compared to secretion by a cell that does not have the CD8 gene sequence.

15 B52. The foregoing method of B-B51, wherein LCK affinity of the cell is increased between about 10% to about 500% as compared to LCK affinity of a cell that does not have the CD8 gene sequence.

B53. The foregoing method of B-B52, wherein persistence of the cell is increased between about 10% to about 500% as compared to persistence of a cell that does not have the CD8 gene sequence.

20 B54. The foregoing method of B-B53, wherein tumor infiltration ability of the cell is increased between about 10% to about 500% as compared to tumor infiltration ability of a cell that does not have the CD8 gene sequence.

B55. The foregoing method of B-B54, wherein the TCR gene encodes a CD8-dependent TCR.

25 B56. The foregoing method of B-B54, wherein the TCR gene encodes a CD8-independent TCR.

B57. The foregoing method of B-B56, wherein the CD8 gene sequence is encoded by a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4.

30 B58. The foregoing method of B-B56, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, CD4 intracellular domain.

B59. The foregoing method of B-B56, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

C. In certain non-limiting embodiments, the presently disclosed subject matter provides for a cell modified by the method of B-B57.

D. In certain non-limiting embodiments, the presently disclosed subject matter provides for a composition comprising an effective amount of a cell of A-A46 or a cell of C.

D1. The foregoing composition of D, wherein the composition is a pharmaceutical composition that further comprises a pharmaceutically acceptable excipient.

D2. The foregoing composition of D or D1, wherein the composition is administered to a patient in need thereof for the treatment of cancer.

D3. The foregoing composition of D-D2, wherein the composition comprises a cryopreservation agent.

D4. The foregoing composition of D-D3, wherein the composition comprises serum albumin.

D5. The foregoing composition of D-D4, wherein the composition comprises Plasma-Lyte A, HSA, and CryoStor CS10.

E. In certain non-limiting embodiments, the presently disclosed subject matter provides for a method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell of A-A46, a cell of claim C, or a composition of D-D5.

E1. The foregoing method of E, wherein prior to administering the therapeutically effective amount of cells, a non-myeloablative lymphodepletion regimen is administered to the subject.

E2. The foregoing method of E or E1, wherein the cancer is a solid tumor.

5 E3. The foregoing method of E or E1, wherein the cancer is liquid tumor.

E4. The foregoing method of E2, wherein the solid tumor is selected from the group consisting of melanoma, thoracic cancer, lung cancer, ovarian cancer, breast cancer, pancreatic cancer, head and neck cancer, prostate cancer, gynecological cancer, central nervous system cancer, cutaneous cancer, HPV+ cancer, esophageal cancer, thyroid  
10 cancer, gastric cancer, hepatocellular cancer, cholangiocarcinomas, renal cell cancers, testicular cancer, sarcomas, and colorectal cancer.

E5. The foregoing method of E3, wherein the liquid tumor is selected from the group consisting of follicular lymphoma, leukemia, and multiple myeloma.

F. In certain non-limiting embodiments, the presently disclosed subject matter  
15 provides for a kit comprising a cell of A-A46, reagents for performing the method of B-B57, a cell of C, or a composition of D-D5.

F1. The foregoing kit of F, wherein the kit further comprises written instructions for treating a cancer.

G. In certain non-limiting embodiments, the presently disclosed subject matter  
20 provides for a cell, comprising: an exogenous T cell receptor (TCR); and an exogenous CD8, comprising: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane  
25 domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

H. In certain non-limiting embodiments, the presently disclosed subject matter provides for a cell, comprising: an exogenous T cell receptor (TCR); and an exogenous CD8, comprising: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain  
30 (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane

domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

I. In certain non-limiting embodiments, the presently disclosed subject matter provides for a method of modifying a cell, the method comprising: introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises: first and second homology arms homologous to first and second target nucleic acid sequences; a TCR gene sequence positioned between the first and second homology arms; a CD8 gene sequence positioned between the first and the second homology arms; and recombining the HR template nucleic acid into an endogenous locus of the cell, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

J. In certain non-limiting embodiments, the presently disclosed subject matter provides for a method of modifying a cell, the method comprising: introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises: first and second homology arms homologous to first and second target nucleic acid sequences; a TCR gene sequence positioned between the first and second homology arms; a CD8 gene sequence positioned between the first and the second homology arms; and recombining the HR template nucleic acid into an endogenous locus of the cell, wherein the CD8 gene sequence comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular

domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

K. In certain non-limiting embodiments, the presently disclosed subject matter provides for a composition comprising a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

L. In certain non-limiting embodiments, the presently disclosed subject matter provides for a composition comprising a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

M. In certain non-limiting embodiments, the presently disclosed subject matter provides for a method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the

exogenous CD8 comprises: a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain; a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

N. A method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises: a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

## EXAMPLES

The following are examples of methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

### Example 1. Generation of NeoTCR Products

Neoepitope-specific TCRs identified by the imPACT Isolation Technology described in PCT/US2020/17887 (which is herein incorporated by reference in its entirety) were used to generate homologous recombination (HR) DNA templates. These HR templates were transfected into primary human T cells in tandem with site-specific nucleases (*see Figures 1A-1C*). The single-step non-viral precision genome engineering

resulted in the seamless replacement of the endogenous TCR with the patient's neoepitope-specific TCR, expressed by the endogenous promoter. The TCR expressed on the surface is entirely native in sequence.

The precision of neoTCR-T cell genome engineering was evaluated by Targeted  
5 Locus Amplification (TLA) for off-target integration hot spots or translocations, and by next generation sequencing based off-target cleavage assays and found to lack evidence of unintended outcomes.

As shown in **Figures 1A-1C**, constructs containing genes of interest were inserted  
10 into endogenous loci. This was accomplished with the use of homologous repair templates containing the coding sequence of the gene of interest flanked by left and right HR arms. In addition to the HR arms, the gene of interest was sandwiched between 2A peptides, a protease cleavage site that is upstream of the 2A peptide to remove the 2A peptide from the upstream translated gene of interest, and signal sequences (**Figure 1B**). Once  
15 integrated into the genome, the gene of interested expression gene cassette was transcribed as single messenger RNA. During the translation of this gene of interest in messenger RNA, the flanking regions were unlinked from the gene of interest by the self-cleaving 2A peptide and the protease cleavage site was cleaved for the removal of the 2A peptide upstream from the translated gene of interest (**Figure 1C**). In addition to the 2A peptide and protease cleavage site, a gly-ser-gly (GSG) linker was inserted before each 2A peptide  
20 to further enhance the separation of the gene of interest from the other elements in the expression cassette.

It was determined that P2A peptides were superior to other 2A peptides for Cell  
Products because of its efficient cleavage. Accordingly, two (2) P2A peptides and codon  
25 divergence were used to express the gene of interest without introducing any exogenous epitopes from remaining amino acids on either end of the gene of interest from the P2A peptide. The benefit of the gene edited cell having no exogenous epitopes (i.e., no flanking P2A peptide amino acids on either side of the gene of interest) is that immunogenicity is drastically decreased and there is less likelihood of a patient infused with a Cell Product containing the gene edited cell to have an immune reaction against the gene edited cell.

30 As described in PCT/US/2018/058230, NeoTCRs were integrated into the TCR $\alpha$  locus of T cells. Specifically, a homologous repair template containing a NeoTCR coding sequence flanked by left and right HR Arms was used. In addition, the endogenous TCR $\beta$  locus was disrupted leading to the expression of only TCR sequences encoded by the

NeoTCR construct. The general strategy was applied using circular HR templates as well as with linear templates.

The target TCR $\alpha$  locus (C $\alpha$ ) is shown along with the plasmid HR template, and the resulting edited sequence and downstream mRNA/protein products in **Figures 1B and 1C**.

5 The target TCR $\alpha$  locus (endogenous TRAC) and its CRISPR Cas9 target site (horizontal stripe, cleavage site designated by arrow) are shown (**Figures 1A-1C**). The circular plasmid HR template with the polynucleotide encoding the NeoTCR is located between left and right homology arms (“LHA” and “RHA” respectively). The region of the TRAC introduced by the HR template that was codon optimized is shown (vertical stripe). The

10 TCR $\beta$  constant domain was derived from TRBC2, which is indicated as being functionally equivalent to TRBC1. Other elements in the NeoTCR cassette include: 2A = 2A ribosome skipping element (by way of non-limiting example, the 2A peptides used in the cassette are both P2A sequences that are used in combination with codon divergence to eliminate any otherwise occurring non-endogenous epitopes in the translated product); P = protease

15 cleavage site upstream of 2A that removes the 2A tag from the upstream TCR $\beta$  protein (by way of non-limiting example the protease cleavage site can be a furin protease cleavage site); SS = signal sequences (by way of non-limited example the protease cleavage site can be a human growth hormone signal sequence). The HR template of the NeoTCR expression gene cassette includes two flanking homology arms to direct insertion into the

20 TCR $\alpha$  genomic locus targeted by the CRISPR Cas9 nuclease RNP with the TCR $\alpha$  guide RNA. These homology arms (LHA and RHA) flank the neoE-specific TCR sequences of the NeoTCR expression gene cassette. While the protease cleavage site used in this example was a furin protease cleavage site, any appropriate protease cleavage site known to one of skill in the art could be used. Similarly, while HGH was the signal sequence

25 chosen for this example, any signal sequence known to one of skill in the art could be selected based on the desired trafficking and used.

Once integrated into the genome (**Figure 1C**), the NeoTCR expression gene cassette is transcribed as a single messenger RNA from the endogenous TCR $\alpha$  promoter, which still includes a portion of the endogenous TCR $\alpha$  polypeptide from that individual T

30 cell (**Figure 1C**). During ribosomal polypeptide translation of this single NeoTCR messenger RNA, the NeoTCR sequences are unlinked from the endogenous, CRISPR-disrupted TCR $\alpha$  polypeptide by self-cleavage at a P2A peptide (**Figure 1C**). The encoded NeoTCR $\alpha$  and NeoTCR $\beta$  polypeptides are also unlinked from each other through cleavage by the endogenous cellular human furin protease and a second self-cleaving P2A sequence

motifs included in the NeoTCR expression gene cassette (**Figure 1C**). The NeoTCR $\alpha$  and NeoTCR $\beta$  polypeptides are separately targeted by signal leader sequences (derived from the human growth hormone, HGH) to the endoplasmic reticulum for multimer assembly and trafficking of the NeoTCR protein complexes to the T cell surface. The inclusion of the furin protease cleavage site facilitates the removal of the 2A sequence from the upstream TCR $\beta$  chain to reduce potential interference with TCR $\beta$  function. Inclusion of a gly-ser-gly linker before each 2A (not shown) further enhances the separation of the three polypeptides.

Additionally, three repeated protein sequences are codon diverged within the HR template to promote genomic stability. The two P2A are codon diverged relative to each other, as well as the two HGH signal sequences relative to each other, within the TCR gene cassette to promote stability of the introduced NeoTCR cassette sequences within the genome of the *ex vivo* engineered T cells. Similarly, the re-introduced 5' end of TRAC exon 1 (vertical stripe) reduces the likelihood of the entire cassette being lost over time through the removal of intervening sequence of two direct repeats.

In addition to NeoTCR Products, this method can be used for any CD8 Product.

In-Out PCR was used to confirm the precise target integration of the NeoE TCR cassette. Agarose gels show the results of a PCR using primers specific to the integration cassette and site generate products of the expected size only for cells treated with both nuclease and DNA template (KOKI and KOKIKO), demonstrating site-specific and precise integration.

Furthermore, Targeted Locus Amplification (TLA) was used to confirm the specificity of targeted integration. Crosslinking, ligation, and use of primers specific to the NeoTCR insert were used to obtain sequences around the site(s) of integration. The reads mapped to the genome are binned in 10 kb intervals. Significant read depths were obtained only around the intended site the integration site on chromosome 14, showing no evidence of common off-target insertion sites.

Antibody staining for endogenous TCR and peptide-HLA staining for neoTCR revealed that the engineering results in high frequency knock-in of the NeoTCR, with some TCR- cells and few WT T cells remaining. Knock-in is evidenced by neoTCR expression in the absence of an exogenous promoter. Engineering was carried out multiple times using the same neoTCR with similar results. Therefore, efficient and consistent expression of the NeoTCR and knockout of the endogenous TCR in engineered T cells was achieved.

## Example 2. Generation of CD8 Product 1

*T cell Isolation and Editing.* CD4 and CD8 T cells were isolated from healthy donor PBMCs using the Miltenyi Prodigy or Miltenyi MACS separation columns according to the manufacturers' instructions. Positively-selected CD4 and CD8 T cells (using Miltenyi antibodies and isolation column) were used fresh or cryopreserved in 1% human serum albumin (Gemini), 49% plasmalyte (Baxter), and 50% CS10 (Sigma). Cryopreserved cells were thawed, washed in TexMACS (Miltenyi) + 10% human AB serum (Valley Biomedical), and seeded at a density of  $2 \times 10^6$  cells per mL in TexMACS + 3% human AB serum (culture medium). One day after thaw, or immediately if used fresh, the cells were washed and re-seeded at a density of  $1.46 \times 10^6$  cells per mL in culture medium + 12.5 ng/mL IL7 + 12.5 ng/mL IL15 + 1:17.5 ratio of TransACT T cell activation reagent (all reagents from Miltenyi) by volume. Two days after activation, T cells were electroporated with i) a plasmid for the production of a NeoTCR Product (*see, e.g., Figure 1B*) or ii) a CD8 Construct 1 (*e.g., consisting of the coding sequence of CD8 $\alpha$  flanked by P2A sites upstream of the neoTCR beta and alpha sequences and gRNA-Cas9 RNPs targeting the TCR alpha and beta loci; see, e.g., Figure 2A*). An exemplary expression construct of CD8 Construct 1 is shown in **Figure 11A**. T cells were electroporated using the Lonza X-unit in 100  $\mu$ L cuvettes and program EO-115. T cells are expanded in culture medium supplemented with 12.5 ng/mL IL7 + 12.5 ng/mL IL15. Supplemented medium was exchanged every 2-3 days until the end of study, 13 days after activation.

*comPACT and comPACT-Dextramer preparation.* Neoantigen-specific peptide-HLA complex polypeptides (each a "comPACT") were prepared according to the method as described in PCT/US2019/025415, hereby incorporated by reference in its entirety. A comPACT-dextramer complex was made for the labeling of neoTCR expressing T cells. Biotinylated comPACT protein was incubated with a streptavidin-conjugated fluorophore for 10 min at room temperature (RT). Biotin-40-dextran (NANOCS) was added to the mixture and incubated at RT for an additional 10 minutes. The comPACT-Dextramer was stored at 4°C.

*Confirmation of comPACT binding to neoTCR edited T cells.* T cells were stained for flow cytometry. Cells were first stained with viability dye for 20 minutes at 4°C, then washed and stained with the comPACT-dextramer for 10 minutes at 4°C. Surface antibodies (anti-CD8 $\alpha$ , anti-CD8 $\beta$ , anti-CD4) were added to the suspension of cells and comPACT-dextramer, and the cells are incubated for an additional 20 minutes at 4°C. Cells

were then washed and fixed in intracellular fixation buffer (BD Biosciences). All cells were acquired on an Attune NxT Flow Cytometer (ThermoFisher Scientific) and data analyzed with either FCS Express or FlowJo.

*Cytometric Bead Array (CBA).* Streptavidin coated plates (Eagle Biosciences) were washed 3 times with wash buffer (PBS supplemented with 1% BSA and 0.05% tween20) and then coated with comPACTs at different concentrations ranging from 100-0.01 ng/well. Wells with no comPACT and wells coated with mismatched comPACT were used as controls. The plates were incubated for 2 hr at room temperature, washed three times with wash buffer, and then washed three times with TexMACS supplemented with 3% human AB serum to remove the tween20. T cells were given two washes with TexMACS supplemented with 3% human AB serum and resuspended at 1 million cells/mL in TexMACS supplemented with 3% human AB serum and 1X penicillin-streptomycin solution. T cells were plated onto the comPACT coated plate at 100  $\mu$ L/well and incubated at 37°C, 5% CO<sub>2</sub>. After 24h the supernatant was collected, and the cytokine concentrations were analyzed using the BD Cytometric Bead Array (CBA) Human Th1/Th2 Cytokine Kit II (Catalog No. 551809) following the manufacturer's protocol. Capture beads were mixed with culture supernatant, incubated with the detection reagent for 3 hr at RT protected from light, washed, and resuspended in wash buffer. Samples were assayed on an Attune NxT Flow Cytometer and data analyzed with FlowJo. The EC50 represents the concentration of cognate comPACT that elicits 50% of the maximum response and is calculated utilizing a least-squares fit of IFN $\gamma$  secretion over a range of comPACT concentrations.

*Intracellular Staining.* T cells were stained for flow cytometry on the indicated days. T cells are first stained with viability dye for 20 minutes at 4°C, then washed and incubated with surface antibodies (anti-CD8 $\alpha$ , anti-CD8 $\beta$ , anti-CD4) for an additional 20 minutes at 4°C. T cells are then washed and permeabilized for intracellular staining. T cells are stained with anti-2A peptide or with anti-IFN $\gamma$ , anti-TNF, or anti-IL2 in permeabilization buffer for 20 minutes at 4°C. T cells are fixed in intracellular fixation buffer (BD Biosciences). Samples are assayed on an Attune NxT Flow Cytometer (ThermoFisher Scientific) and data analyzed with either FCS Express or FlowJo.

*T cell Proliferation Assay.* Edited CD4 and CD8 T cells are labeled with the e450 proliferation dye (eBioscience) according to the manufacturer's instructions. Labeled cells were stimulated on comPACT coated plates with a range of concentrations as described

above. T cells were harvested over 48-96 hours and analyzed for proliferation as measured by dilution of the e450 dye.

*T cell Killing Assay.* HLA-matched cell lines were pulsed with the cognate neoantigen peptide or mismatched peptide for 1h at 37°C, 5% CO<sub>2</sub>. The cells were washed 3 times with media to remove any unbound peptide and then co-cultured with edited CD4 and CD8 T cells that are labeled with the e450 proliferation dye described above. Co-cultures were incubated for 48h at 37°C with 5% CO<sub>2</sub> before harvest. Cells were washed and stained with a fixable viability dye to determine killing efficiency. The e450 proliferation dye was used to distinguish edited T cells from target cells.

*Generation and Validation of CD8 Product 1.* An expression construct consisting of the coding sequence (CDS) of CD8 $\alpha$  flanked by P2A sites upstream of the neoTCR beta and alpha sequences is synthesized. Briefly, the CDS of human CD8 $\alpha$  is synthesized with a GSG-linker and P2A site upstream and flanked with restriction sites. The neoTCR expression vector of interest and the synthesized CD8 $\alpha$  construct were incubated with restriction enzymes and ligated together to create the final HDR construct. The CD8 Construct 1 was electroporated along with gRNA-Cas9 RNPs targeting the TCR $\alpha$  and  $\beta$  loci. Model neoTCRs known to bind dextramer among CD8 T cells but not CD4 T cells (*e.g.* TCR097) were used to demonstrate that expression of the CD8 $\alpha$  transgene enables these TCRs to bind dextramer among CD4 T cells.

To test the efficiency of the gene transfer and expression of CD8 $\alpha$  on the surface of CD4 T cells, engineered CD4 T cells were stained with anti-CD8 $\alpha$  antibodies and surface expression of the transgene is confirmed by flow cytometry as described above. NeoTCRs known to bind dextramer among CD8 T cells only (*e.g.*, TCR 097) were used to demonstrate that expression of the CD8 $\alpha$  transgene enables these TCRs to bind dextramer among CD4 T cells. *See, Figures 13A, 13B, 14, 15A, and 15B.* Furthermore, CD8 $\alpha$  expression on wild type and genetically engineered CD8 $\alpha$  T cells was assessed to determine whether the addition of the CD8 $\alpha$  transgene increased surface levels of CD8 $\alpha$  on CD8 T cells. CD4 T cells engineered to express the CD8 $\alpha$  transgene were double-positive for CD4 and CD8 $\alpha$ .

CD8 T cells were also engineered to express the CD8 $\alpha$  transgene and characterized as described above. **Figures 15A and 15B.** Relative CD8 $\alpha$  gene expression was also quantified via RT-qPCR and compared to control non-engineered CD8 T cells. CD8 T cells expressing the CD8 $\alpha$  transgene had higher than endogenous levels of CD8 $\alpha$  expression.

*Effect of CD8 expression on T cell proliferation upon encounter of cognate antigen.* Edited T cells were stained with a proliferation dye as described above. After staining, T cells were stimulated with a range of concentrations of cognate comPACT proteins. 48-72 hours later, T cells were harvested and stained with anti-CD4, anti-CD8, and anti-2A peptide as described above. Edited T cells were identified by 2A expression. Proliferation of the CD4 and CD8 T cells was determined by quantifying the dilution of the proliferative dye. NeoTCR-expressing T cells lacking the CD8 $\alpha$  transgene were used as a negative control. NeoTCR CD4 T cells expressing the CD8 $\alpha$  transgene proliferated in response to lower concentrations of cognate comPACT than CD4 T cells lacking CD8 $\alpha$  expression.

*Effect of CD8 $\alpha$  expression on cytokine production upon encounter of cognate antigen.* In addition to proliferation, effector cytokine production via intracellular cytokine staining was measured. In this assay, NeoTCR Products and CD8 Product 1 were be stimulated with various concentrations of cognate comPACT for 5 hours in the presence of brefeldin A. After stimulation, T cells were stained with anti-CD4 and anti-CD8 $\alpha$ . Cells were be permeabilized and stained with anti-P2A peptide, anti-IFN $\gamma$ , anti-TNF, and anti-IL2. NeoTCR CD4 T cells expressing the CD8 $\alpha$  transgene produced effector cytokines in response to lower concentrations of cognate comPACT than neoTCR CD4 T cells lacking CD8 $\alpha$  expression.

*Effect of CD8 $\alpha$  expression on killing activity upon encounter of cognate antigen.* To assess effector function, edited CD4 and CD8 T cells were cultured with HLA-matched target cells pulsed with cognate peptide as described above. CD4 and CD8 T cells were edited separately to evaluate the ability of CD4 T cells expressing CD8 $\alpha$  to kill target cells. NeoTCR CD4 T cells expressing the CD8 $\alpha$  transgene killed a greater fraction of target cells presenting cognate peptide than neoTCR CD4 T cells lacking CD8 $\alpha$  expression.

### **Example 3. Generation of CD8 Product 2**

CD4 T cells engineered to express CD8 $\alpha$  lack CD8 $\beta$  expression. However, CD8 $\beta$  has a higher affinity for LCK (Irie et al., 1998, J. Immunol, 161(1), 183–191). Therefore, T cells were edited to co-express CD8 $\beta$  with CD8 $\alpha$  (i.e., a CD8 Product 2). An additional construct containing CD8 $\beta$  flanked by P2A sites, CD8 $\alpha$  flanked by P2A sites, followed by the TRB and TRA alleles as previously described is generated. CD4 and CD8 T cells expressing CD8 $\alpha$  and CD8 $\beta$  are evaluated using the same assays described above. An

exemplary expression construct with the CD8 $\alpha$  and CD8  $\beta$  sequences is shown in **Figure 11B**.

CD8 $\beta$  expression was also assessed in the edited CD4 T cells. NeoTCR CD4 T cells expressing the CD8 $\alpha$  and CD8 $\beta$  transgenes proliferated in response to lower concentrations of cognate comPACT than CD4 T cells expressing CD8 $\alpha$  transgene alone. NeoTCR CD4 T cells expressing the CD8 $\alpha$  and CD8 $\beta$  transgenes also produced effector cytokines in response to lower concentrations of cognate comPACT than neoTCR CD4 T cells expressing the CD8 $\alpha$  transgene alone. And neoTCR CD4 T cells expressing the CD8 $\alpha$  and CD8 $\beta$  transgenes killed a greater fraction of target cells presenting cognate peptide than neoTCR CD4 T cells expressing CD8 $\alpha$  transgene alone.

#### **Example 4. Generation of chimeric CD8 $\alpha$ and CD8 $\beta$ Constructs and CD8 Products 3 and 4**

To ensure efficient editing of the T cells and expression of the neoTCR, chimeric proteins made up of the coding sequences for the extracellular and transmembrane domains of CD8 $\alpha$  linked to the intracellular domain of CD8 $\beta$  were generated. An exemplary expression construct with the extracellular and transmembrane domains of CD8 $\alpha$  linked to the intracellular domain of CD8 $\beta$  sequences is (*i.e.*, a CD8 Product 3) shown in **Figure 11C**.

Furthermore, the intracellular domain of CD4 has an even higher affinity for LCK than CD8 (Irie et al., 1998). Therefore, a second chimeric protein was generated containing the coding sequences for the extracellular and transmembrane domains of CD8 $\alpha$  linked to the intracellular domain of CD4. An exemplary expression construct with the extracellular and transmembrane domains of CD8 $\alpha$  linked to the intracellular domain of CD4 sequences (*i.e.*, a CD8 Product 4) is shown in **Figure 11D**.

CD4 and CD8 T cells expressing the CD8 Products 3 and 4 were evaluated using the same assays described above.

NeoTCR CD4 T cells expressing the CD8 $\alpha$ -CD8 $\beta$ -ID transgene proliferated in response to lower concentrations of cognate comPACT than CD4 T cells expressing CD8 $\alpha$  transgene alone. NeoTCR CD4 T cells expressing the CD8 $\alpha$ -CD8 $\beta$ -ID transgene produced effector cytokines in response to lower concentrations of cognate comPACT than neoTCR CD4 T cells expressing the CD8 $\alpha$  transgene alone. NeoTCR CD4 T cells expressing the

CD8 $\alpha$ -CD8 $\beta$ -ID transgene killed a greater fraction of target cells presenting cognate peptide than neoTCR CD4 T cells expressing CD8 $\alpha$ -CD8 $\beta$ -ID transgene.

NeoTCR CD4 T cells expressing the CD8 $\alpha$ -CD4-ID transgene proliferated in response to lower concentrations of cognate comPACT than neoTCR CD4 T cells expressing CD8 $\alpha$ -CD8 $\beta$ -ID transgene. NeoTCR CD4 T cells expressing the CD8 $\alpha$ -CD4 transgene produced effector cytokines in response to lower concentrations of cognate comPACT than neoTCR CD4 T cells expressing the CD8 $\alpha$ -CD8 $\beta$ -ID transgene.

NeoTCR CD4 T cells expressing the CD8 $\alpha$ -CD4-ID transgene killed a greater fraction of target cells presenting cognate peptide than neoTCR CD4 T cells expressing CD8 $\alpha$ -CD8 $\beta$ -ID transgene. NeoTCR CD8 T cells expressing the CD8 $\alpha$ -CD4-ID transgene proliferated in response to lower concentrations of cognate comPACT than neoTCR CD8 T cells lacking the transgene. NeoTCR CD8 T cells expressing the CD8 $\alpha$ -CD4 transgene produced effector cytokines in response to lower concentrations of cognate comPACT than neoTCR CD8 T cells lacking the transgene. NeoTCR CD8 T cells expressing the CD8 $\alpha$ -CD4-ID transgene killed a greater fraction of target cells presenting cognate peptide than neoTCR CD8 T cells lacking the transgene.

**Example 5. CD8 Products have increased sensitivity to neoE-HLA target recognition and trigger pro-inflammatory and cytotoxic function**

MHC-I neoTCRs were cloned from neoE-specific T cells captured from the blood of a patient with colorectal cancer. Healthy donor CD8 and CD4 T cells were precision genome engineered to express the cloned MHC-I neoTCRs alone or to include engineering of ectopic CD8 co-receptors in the gene-edited T cells. Flow cytometric analysis was used to evaluate surface expression of neoTCRs and ectopic CD8 co-receptors (i.e., the CD8 and CD4 components of the CD8 Constructs 1-4), respectively. Rescue of neoTCR binding among CD4 T cells for lower affinity, CD8-dependent neoTCRs was observed. Importantly, in response to stimulation with cognate antigen, CD107a and intracellular IFN $\gamma$  staining revealed 10-100-fold increases in the sensitivity of MHC-I neoTCR-induced effector functions by CD4 T cells, with no effect on specificity. No change in functionality or sensitivity was seen on CD8 T cells by the expression of additional CD8 co-receptor.

These results demonstrate that simultaneous precision genome engineering of the CD8 co-receptor together with CD8-dependent MHC-I neoTCRs into CD4 T cells (i.e., CD8 Products 1-4) significantly increases their sensitivity to neoE-HLA target recognition

as well as triggering pro-inflammatory and cytotoxic function, yet without compromising antigen-specificity.

### **Example 6. Generation and design of CD8 Products that have varying LCK affinities**

As described herein, four classes of CD8 Products were generated:

- 5           1. CD8 $\alpha$  homodimer (CD8 Construct 1)
2. CD8 $\alpha$ -P2A-CD8 $\beta$  (CD8 Construct 2)
3. CD8 $\alpha$  with CD8 $\beta$  intracellular domain (CD8 Construct 3)
4. CD8 $\alpha$  homodimer with CD4 intracellular domain (CD8 Construct 4)

As shown in **Figure 10**, these CD8 Constructs and resulting CD8 Products were  
10   designed to allow for varying degrees of LCK affinity. As predicted, CD8 Product 1 was  
shown to have the lowest LCK affinity, followed by CD8 Product 2, CD8 Product 3, and  
CD8 Product 4 (in that order with CD8 Product 4 having the highest LCK affinity).

Based on the high affinity of CD8 Product 4, this product was used in cell killing  
assays to exemplify the increased cell killing ability of CD8 Products 1-4 compared to  
15   NeoTCR Products.

CD4<sup>+</sup> T cells were engineered as described herein to express the CD8 Product 4  
(with TCR097 as the NeoTCR in the product) described in **Figured 2D** and **3D**. SW620  
cell lines that were engineered to heterologously express the R20Q mutation (the cognate  
antigen to TCR097). The CD8 Product 4 expressing NeoTCR097 was combined with the  
20   SW620 heterologous cells. As shown in **Figures 13A** and **13B**, the CD8 Product 4  
provided substantially better killing of the cognate antigen expressing SW620 cells than  
the NeoTCR Product also expressing the TCR097. The experiment shown in **Figure 13A**  
was done with an E:T ratio of 1:1 and as shown in the graph, the NeoTCR Product  
expressing TCR097 did not show any efficacy at killing of the cognate antigen expressing  
25   SW620 cells. The experiment shown in **Figure 13B** was done with an E:T ratio of 2:1 and  
while the NeoTCR Product expressing TCR097 showed some ability to kill the cognate  
antigen expressing SW620 cells, it was clear from the experiment that the CD8 Product 4  
expressing the TCR097 had superior efficacy.

The same engineered CD4<sup>+</sup> cells that were used in the experiment above with data  
30   provided in **Figures 13A** and **13B**, were also tested on SW620 cell lines that were  
engineered to homozygously express the R20Q mutation. In this experiment, the high

expression of the cognate antigen was able to compensate for the low affinity NeoTCR097 and both the NeoTCR Product expressing TCR097 and the CD8 Product 4 expressing the TCR097 showed efficacy at killing the SW620 cells. However, the high expression of the cognate antigen in the homozygous SW620 cells is not physiologically relevant and this experiment serves to highlight the ability to rescue NeoTCR Products with low affinity TCRs that cannot effectively engage with and kill tumor cells by further engineering them to also include a CD8 $\alpha$  homodimer with CD4 intracellular domain (*i.e.*, a CD8 Product 4).

As a final control and proof of the efficacy of the CD8 Products, CD8 T cells were also transfected to express the NeoTCR097 and a CD8 $\alpha$  homodimer with CD4 intracellular domain (*i.e.*, a CD8 Product 4 using CD8 T cells instead of CD4 T cells). As shown in the top graphs in **Figures 15A** and **15B**, the CD8 Product 4 provided substantially better killing of the SW620 cells than the NeoTCR Product when the products were made from CD4 T cells. However, when the CD8 Product 4 and NeoTCR Product were made from CD8 T cells (the bottom graphs in **Figures 15A** and **15B**) the ability of the NeoTCR Product was rescued because of the endogenous CD8 expression in the CD8 T cells. While no overlay of the graphs is shown, it is also of note that the CD8 Product 4 in CD4 T cells shown the top graphs in **Figures 15A** and **15B**, appear to have better efficacy than the CD8 Product 4 in CD8 T cells in the bottom two graphs; suggesting a superior ability of the CD8 Products described herein to engage with the cognate antigens of tumor cells leading to tumor death and effective treatment of patients with cancer in need of treatment.

#### **Example 6. CD8 Products have increased CD4 T cell sensitivity while maintaining NeoTCR sensitivity**

In order to confirm that the CD8 Constructs expressed properly, CD8 Products 1, 2, 3, and 4 were tested to determine surface expression of CD8 $\alpha$ . It was shown that each of CD8 Products 1, 2, 3, and 4 exhibited normal CD8 $\alpha$  surface expression. Representative data from CD8 Product 4 is shown in **Figure 16**.

It was also important to determine how the CD8 Constructs affected CD4 T cell sensitivity and specificity for the cognate antigen of the expressed NeoTCR in the CD8 Products. Sensitivity experiments were performed and it was shown that CD8 Products 1-4 exhibited an increased CD4 T cell sensitivity (**Figure 17A**). Specificity experiments were also performed on CD8 Products 1-4. CD8 Products 1, 2, 3, and 4 were made with NeoTCR097. Experiments were performed to test the CD8 Products 1, 2, 3, and 4

(expressing NeoTCR097) to assess the specificity of these products to the cognate antigen to NeoTCR097. As shown in **Figure 17B**, CD8 Products 1, 2, 3, and 4 (expressing NeoTCR097) were specific for the cognate antigen to NeoTCR097 and showed no activity when exposed to a mismatched antigen. The specificity was determined by INF $\gamma$  and CD107 production which are evidence of T cell activation. Thus, the CD8 Products described herein have an increased sensitivity to CD4 T cells and maintain their specificity to cognate antigen to the expressed NeoTCR compared to NeoTCR Products expressing the same NeoTCR.

Lastly, experiments were performed to investigate the impact the CD8 Constructs have on CD8-dependent and CD8-independent NeoTCRs. It was expected that CD8-dependent NeoTCRs would show an increased sensitivity to cognate antigen because the CD8 Cells would be engineered to express CD8 $\alpha$  and that CD8-independent NeoTCRs would not show an increased sensitivity because of the independent nature of the NeoTCR. However, it was shown that NeoTCR Products 1, 2, 3, and 4 surprisingly increased the sensitivity of CD8-dependent (e.g., NeoTCR097) and CD8-independent (e.g., NeoTCR089) NeoTCRs. Accordingly, it was shown that the CD8 Constructs 1, 2, 3, and 4 can improve NeoTCR engagement and T cell killing of tumor cells with cognate NeoTCR antigens for all NeoTCRs regardless of whether they are CD8-dependent or CD8-independent.

20

While the present invention has been described at some length and with some particularity with respect to the several described embodiments, it is not intended that it should be limited to any such particulars or embodiments or any particular embodiment, but it is to be construed with references to the appended claims so as to provide the broadest possible interpretation of such claims in view of the prior art and, therefore, to effectively encompass the intended scope of the invention.

All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, section headings, the materials, methods, and examples are illustrative only and not intended to be limiting.

30

**WHAT IS CLAIMED IS:**

1. A cell, comprising:
  - a. an exogenous T cell receptor (TCR); and
  - b. an exogenous CD8.
- 5 2. The cell of claim 1, wherein the exogenous CD8 comprises at least one monomer.
3. The cell of claim 2, wherein the at least one monomer of the exogenous CD8 comprises an extracellular domain, a transmembrane domain, an intracellular domain, fragments thereof, or combinations thereof.
4. The cell of claim 3, wherein the extracellular domain comprises a CD8 $\alpha$   
10 extracellular domain or a CD8 $\beta$  extracellular domain.
5. The cell of claim 3 or 4, wherein the transmembrane a CD8 $\alpha$  transmembrane domain or a CD8 $\beta$  transmembrane domain.
6. The cell of any one of claims 3-5, wherein the intracellular domain comprises a CD8 $\alpha$  intracellular domain or a CD8 $\beta$  intracellular domain.
- 15 7. The cell of any one of claims 3-5, wherein the intracellular domain comprises a CD4 intracellular domain.
8. The cell of any one of claims 2-6, wherein the at least one monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain.
- 20 9. The cell of any one of claims 2-6, wherein the at least one monomer comprises a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain.
10. The cell of any one of claims 2-6, wherein the at least one monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$   
25 intracellular domain.
11. The cell of any one of claims 2-7, wherein the at least one monomer comprises a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.
12. The cell of any one of claims 2-11, wherein the at least one monomer comprises a  
30 signal peptide.
13. The cell of claim 12, wherein the signal peptide is a CD8 signal peptide.
14. The cell of any one of claims 3-13, wherein the extracellular domain comprises the amino acid sequence set forth in SEQ ID NO: 140, or SEQ ID NO: 145.

15. The cell of any one of claims 3-14, wherein the transmembrane domain comprises the amino acid sequence set forth in SEQ ID NO: 141, or SEQ ID NO: 146.
16. The cell of any one of claims 3-15, wherein the intracellular domain comprises the amino acid sequence set forth in SEQ ID NO: 142, SEQ ID NO: 147, or SEQ ID NO: 148.
17. The cell of any one of claims 12-16, wherein the signal peptide comprises the amino acid sequence set forth in SEQ ID NO: 139, or SEQ ID NO: 144.
18. The cell of any one of claims 1-17, wherein the exogenous CD8 comprises a 2A sequence.
19. The cell of any one of claims 1-18, wherein the exogenous CD8 comprises a linker.
20. The cell of claim 19, wherein the linker comprises the amino acid sequence set forth in SEQ ID NO: 137.
21. The cell of any one of claims 1-20, wherein the exogenous CD8 comprises a protease cleavage site.
22. The cell of claim 21, wherein the protease cleavage site is a Furin cleavage site.
23. The cell of any one of claims 1-22, wherein the exogenous TCR is a patient derived TCR.
24. The cell of any one of claims 1-23, wherein the exogenous TCR comprises a signal sequence, a first and second 2A sequence, and a TCR polypeptide sequence.
25. The cell of any one of claims 1-24, wherein the exogenous TCR recognizes a cancer antigen.
26. The cell of claim 25, wherein the cancer antigen is a neoantigen.
27. The cell of claim 25, wherein the cancer antigen is a patient specific antigen.
28. The cell of any one of claims 1-27, wherein the cell is a primary cell.
29. The cell of any one of claims 1-27, wherein the cell is a patient-derived cell.
30. The cell of any one of claims 1-27, wherein the cell is a lymphocyte.
31. The cell of any one of claims 1-27, wherein the cell is a T cell.
32. The cell of any one of claims 1-27, wherein the cell if a young T cell.
33. The cell of claim 32, wherein the cell is CD45RA+, CD62L+, CD28+, CD95-, CCR7+, and CD27+.
34. The cell of claim 32, wherein the cell is CD45RA+, CD62L+, CD28+, CD95+, CD27+, CCR7+.
35. The cell of claim 32, wherein the cell is CD45RO+, CD62L+, CD28+, CD95+, CCR7+, CD27+, CD127+.

36. The cell of any one of claims 1-35, further comprising a gene modification to enhance cell persistence and/or enhances memory cell differentiation
37. The cell of any one of claims 1-36, wherein killing activity of the cell is increased between about 10% to about 500% as compared to killing activity of a cell that does not have the exogenous CD8.
38. The cell of any one of claims 1-37, wherein proliferation of the cell upon binding of the TCR to the antigen is increased between about 10% to about 500% as compared to proliferation of a cell that does not have the exogenous CD8.
39. The cell of any one of claims 1-38, wherein secretion of pro-inflammatory cytokine upon binding of the TCR to the antigen by the cell is increased between about 10% to about 500% as compared to secretion by a cell that does not have the exogenous CD8.
40. The cell of any one of claims 1-39, wherein LCK affinity of the cell is increased between about 10% to about 500% as compared to LCK affinity of a cell that does not have the exogenous CD8.
41. The cell of any one of claims 1-40, wherein persistence of the cell is increased between about 10% to about 500% as compared to persistence of a cell that does not have the exogenous CD8.
42. The cell of any one of claims 1-41, wherein tumor infiltration ability of the cell is increased between about 10% to about 500% as compared to tumor infiltration ability of a cell that does not have the exogenous CD8.
43. The cell of any one of claims 1-42, wherein the exogenous TCR is a CD8-dependent TCR.
44. The cell of any one of claims 1-42, wherein the exogenous TCR is a CD8-independent TCR.
45. The cell of any one of claims 1-44, wherein the exogenous CD8 is encoded by a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4.
46. The cell of any one of claims 1-44, wherein the exogenous CD8 comprises:
- a. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain;
  - b. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain;

- c. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or
- d. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.

5 47. The cell of any one of claims 1-44, wherein the exogenous CD8 comprises:

- a. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142);
- b. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);
- 10 c. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or
- d. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

- 15 48. A method of modifying a cell, the method comprising:
- a. introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises:
    - i. first and second homology arms homologous to first and second target nucleic acid sequences;
    - 25 ii. a TCR gene sequence positioned between the first and second homology arms;
    - iii. a CD8 gene sequence positioned between the first and the second homology arms; and
  - 30 b. recombining the HR template nucleic acid into an endogenous locus of the cell.

49. The method of claim 48, wherein the HR template comprises a first 2A-coding sequence positioned upstream of the CD8 gene sequence, a second 2A-coding sequence positioned downstream of the CD8 gene sequence and upstream of the

- TCR gene sequence, and a third 2A-coding sequence positioned downstream of the TCR gene sequence; wherein the first, second, and third 2A-coding sequences code for the same amino acid sequence and are codon-diverged relative to each other.
50. The method of claim 48 or 49, wherein the HR template comprises a sequence coding for the amino acid sequence Gly Ser Gly positioned immediately upstream of the first, second, and/or third 2A-coding sequences.
51. The method of claim 49 or 50, wherein the HR template further comprises a sequence coding for a Furin cleavage site positioned upstream of the first, second, and/or third 2A-coding sequences.
52. The method of any one of claims 48-51, wherein the HR template further comprises a sequence encoding a signal sequence positioned immediately upstream of the TCR gene sequence and/or the CD8 gene sequence.
53. The method of any one of claims 48-52, wherein the HR template comprises a second TCR sequence positioned between the third 2A-coding sequence and the second homology arm.
54. The method of claim 53, wherein the HR template comprises:
- a. a sequence encoding a first signal sequence positioned immediately upstream the first TCR gene sequence; and
  - b. a sequence encoding a second signal sequence positioned immediately upstream of the second TCR gene sequence.
55. The method of any one of claims 48-54, wherein the HR template comprises a second CD8 gene sequence positioned between the first CD8 gene sequence and the second 2A-coding sequence.
56. The method of claims 55, wherein a 2A coding sequence is positioned between the first and second CD8 gene sequence.
57. The method of claim 55 or 56, wherein a sequence coding for the amino acid sequence Gly Ser Gly is positioned between the first and second CD8 gene sequences.
58. The method of any one of claims 55-57, wherein a sequence coding for a Furin cleavage site is positioned between the first and second CD8 gene sequences.
59. The method of any one of claims 48-58, wherein the CD8 gene sequence comprises a sequence encoding an extracellular domain, a sequence encoding an intracellular domain, a sequence encoding an intracellular domain, fragments thereof, or combinations thereof.

60. The method of claim 59, wherein the sequence encoding an extracellular domain comprises a sequence encoding a CD8 $\alpha$  extracellular domain or a CD8 $\beta$  extracellular domain.
61. The method of claim 59 or 60, wherein the sequence encoding a transmembrane domain comprises a sequence encoding a CD8 $\alpha$  transmembrane domain or a CD8 $\beta$  transmembrane domain.
62. The method of any one of claims 59-61, wherein the sequence encoding an intracellular domain comprises a sequence encoding a CD8 $\alpha$  intracellular domain or a CD8 $\beta$  intracellular domain.
63. The method of any one of claims 59-61, wherein the sequence encoding an intracellular domain comprises a sequence encoding a CD4 intracellular domain.
64. The method of any one of claims 59-62, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain.
65. The method of any one of claims 59-62, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain.
66. The method of any one of claims 59-62, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain.
67. The method of any one of claims 59-63, wherein the CD8 gene sequence comprises a sequence encoding a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.
68. The method of any one of claims 55-67, wherein the HR template comprises:
- a. a sequence encoding a first signal sequence positioned immediately upstream the first CD8 gene sequence; and
  - b. a sequence encoding a second signal sequence positioned immediately upstream of the second CD8 gene sequence.
69. The method of any one of claims 52-68, wherein the signal sequence is a CD8 signal sequence, a human growth hormone signal sequence, fragments thereof, or combinations thereof.
70. The method of any one of claims 48-69, wherein the first and second homology arms of the HR template are each from about 300 bases to about 2,000 bases in length.

71. The method of any one of claims 48-70, wherein the first and second homology arms of the HR template are each from about 600 bases to about 2,000 bases in length.
72. The method of any one of claims 48-71, wherein the exogenous TCR is a patient derived TCR.
73. The method of any one of claims 48-72, wherein the exogenous TCR comprises a signal sequence, a first and second 2A sequence, and a TCR polypeptide sequence.
74. The method of any one of claims 48-73, wherein the exogenous TCR recognizes a cancer antigen.
75. The method of claim 74, wherein the cancer antigen is a neoantigen.
76. The method of claim 74, wherein the cancer antigen is a patient specific antigen.
77. The method of any one of claims 48-76, wherein the HR template is non-viral.
78. The method of any one of claims 48-77, wherein the HR template is a circular DNA.
79. The method of any one of claims 48-77, wherein the HR template is a linear DNA.
80. The method of any one of claims 48-79, wherein the introducing occurs via electroporation.
81. The method of any one of claims 48-80, wherein the recombining comprises:
- a. cleavage of the endogenous locus by a nuclease; and
  - b. recombination of the HR template nucleic acid sequence into the endogenous locus by homology directed repair.
82. The method of claim 81, wherein the nuclease is a Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) family nuclease, or derivative thereof.
83. The method of claim 82, further comprising a gRNA.
84. The method of any one of claims 48-83, wherein the process further comprises culturing the cell.
85. The method of claim 84, wherein the culturing is conducted in the presence of at least one cytokine.
86. The method of claim 84 or 85, wherein the culturing is conducted in the presence of IL2, IL7, IL15, or any combination thereof.
87. The method of claim 84 or 85, wherein the culturing is conducted in the presence of IL7 and IL15.
88. The method of any one of claims 48-87, further comprising a gene modification to enhance cell persistence and/or enhances memory cell differentiation.

89. The method of any one of claims 48-88, wherein the cell is a primary cell.
90. The method of any one of claims 48-88, wherein the cell is a patient-derived cell.
91. The method of any one of claims 48-88, wherein the cell is a lymphocyte.
92. The method of any one of claims 48-88, wherein the cell is a T cell.
- 5 93. The method of any one of claims 48-88, wherein the cell is a young T cell.
94. The method of claim 93, wherein the cell is CD45RA+, CD62L+, CD28+, CD95-, CCR7+, and CD27+.
95. The method of claim 93, wherein the cell is CD45RA+, CD62L+, CD28+, CD95+, CD27+, CCR7+.
- 10 96. The method of claim 93, wherein the cell is CD45RO+, CD62L+, CD28+, CD95+, CCR7+, CD27+, CD127+.
97. The method of any one of claims 48-96, wherein killing activity of the cell is increased between about 10% to about 500% as compared to killing activity of a cell that does not have the CD8 gene sequence.
- 15 98. The method of any one of claims 48-97, wherein proliferation of the cell upon binding of the TCR to the antigen is increased between about 10% to about 500% as compared to proliferation of a cell that does not have the CD8 gene sequence.
99. The method of any one of claims 48-98, wherein secretion of pro-inflammatory cytokine upon binding of the TCR to the antigen by the cell is increased between
- 20 about 10% to about 500% as compared to secretion by a cell that does not have the CD8 gene sequence.
100. The method of any one of claims 48-99, wherein LCK affinity of the cell is increased between about 10% to about 500% as compared to LCK affinity of a cell that does not have the CD8 gene sequence.
- 25 101. The method of any one of claims 48-100, wherein persistence of the cell is increased between about 10% to about 500% as compared to persistence of a cell that does not have the CD8 gene sequence.
102. The method of any one of claims 48-101, wherein tumor infiltration ability of the cell is increased between about 10% to about 500% as compared to tumor infiltration ability of a cell that does not have the CD8 gene sequence.
- 30 103. The method of any one of claims 48-102, wherein the TCR gene encodes a CD8-dependent TCR.
104. The method of any one of claims 48-102, wherein the TCR gene encodes a CD8-independent TCR.

105. The method of any one of claims 48-104, wherein the CD8 gene sequence is encoded by a CD8 Construct 1, a CD8 Construct 2, a CD8 Construct 3, or a CD8 Construct 4.
106. The method of any one of claims 48-104, wherein the CD8 gene sequence  
5 comprises:
- a. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain;
  - b. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain;  
10
  - c. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or
  - d. a CD8 $\alpha$  extracellular domain, CD8 $\alpha$  transmembrane domain, CD4 intracellular domain.
- 15 107. The method of any one of claims 48-104, wherein the CD8 gene sequence comprises:
- a. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142);
  - b. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);  
20
  - c. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or
  - d. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).  
25
- 30 108. A cell modified by the method of any one of claims 48-107.
109. A composition comprising an effective amount of a cell of any one of claims 1-47 or a cell of claim 108.

110. The composition of claim 109, wherein the composition is a pharmaceutical composition that further comprises a pharmaceutically acceptable excipient.
111. The composition of claim 109 or 110, wherein the composition is administered to a patient in need thereof for the treatment of cancer.
- 5 112. The composition of any one of claims 109-111, wherein the composition comprises a cryopreservation agent.
113. The composition of any one of claims 109-112, wherein the composition comprises serum albumin.
114. The composition of any one of claims 109-113, wherein the composition  
10 comprises Plasma-Lyte A, HSA, and CryoStor CS10.
115. A method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell of any one of claims 1-47, a cell of claim 108, or a composition of any one of claims 109-114.
116. The method of claim 115, wherein prior to administering the  
15 therapeutically effective amount of cells, a non-myeloablative lymphodepletion regimen is administered to the subject.
117. The method of claim 115 or 116, wherein the cancer is a solid tumor.
118. The method of claim 115 or 116, wherein the cancer is liquid tumor.
119. The method of claim 117, wherein the solid tumor is selected from the  
20 group consisting of melanoma, thoracic cancer, lung cancer, ovarian cancer, breast cancer, pancreatic cancer, head and neck cancer, prostate cancer, gynecological cancer, central nervous system cancer, cutaneous cancer, HPV+ cancer, esophageal cancer, thyroid cancer, gastric cancer, hepatocellular cancer, cholangiocarcinomas, renal cell cancers, testicular cancer, sarcomas, and colorectal cancer.
- 25 120. The method of claim 118, wherein the liquid tumor is selected from the group consisting of follicular lymphoma, leukemia, and multiple myeloma.
121. A kit comprising a cell of any one of claims 1-47, reagents for performing the method of any one of claims 48-107, a cell of claim 108, or a composition of any one of claims 109-114.
- 30 122. The kit of claim 121, wherein the kit further comprises written instructions for treating a cancer.
123. A cell, comprising:
- a. an exogenous T cell receptor (TCR); and
  - b. an exogenous CD8, comprising:

- 5
- i. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain;
  - ii. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain;
  - iii. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or
  - iv. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.
- 10 124. A cell, comprising:
- a. an exogenous T cell receptor (TCR); and
  - b. an exogenous CD8, comprising:
    - 15 i. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142);
    - ii. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);
    - 20 iii. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);
    - 25 or
    - iv. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).
125. A method of modifying a cell, the method comprising:
- 30 a. introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises:
    - i. first and second homology arms homologous to first and second target nucleic acid sequences;

- ii. a TCR gene sequence positioned between the first and second homology arms;
    - iii. a CD8 gene sequence positioned between the first and the second homology arms; and
  - 5 b. recombining the HR template nucleic acid into an endogenous locus of the cell,
    - wherein the CD8 gene sequence comprises:
      - i. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain;
      - 10 ii. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain;
      - iii. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\beta$  intracellular domain; or
      - 15 iv. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.
- 126. A method of modifying a cell, the method comprising:
  - a. introducing into the cell a homologous recombination (HR) template nucleic acid sequence, wherein the HR template comprises:
    - 20 i. first and second homology arms homologous to first and second target nucleic acid sequences;
    - ii. a TCR gene sequence positioned between the first and second homology arms;
    - iii. a CD8 gene sequence positioned between the first and the second homology arms; and
    - 25 b. recombining the HR template nucleic acid into an endogenous locus of the cell,
      - wherein the CD8 gene sequence comprises:
        - i. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142);
        - 30 ii. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a

- CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);
- 5           iii. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);  
or
- 10           iv. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).
127. A composition comprising a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises:
- 15           a. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain;
- b. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain,
- 20           c. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or
- d. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.
128. A composition comprising a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8
- 25           comprises:
- a. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142);
- 30           b. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);

- c. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or
- d. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).
- 5
129. A method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises:
- 10
- a. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\alpha$  intracellular domain;
- b. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, a CD8 $\alpha$  intracellular domain, a CD8 $\beta$  extracellular domain, a CD8 $\beta$  transmembrane domain, and a CD8 $\beta$  intracellular domain,
- 15
- c. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD8 $\beta$  intracellular domain; or
- d. a CD8 $\alpha$  extracellular domain, a CD8 $\alpha$  transmembrane domain, and a CD4 intracellular domain.
- 20
130. A method of treating cancer in a subject in need thereof, the method comprising administering a therapeutically effective amount of a cell, wherein the cell comprises an exogenous T cell receptor (TCR) and an exogenous CD8, wherein the exogenous CD8 comprises:
- 25
- a. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142);
- b. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), a CD8 $\alpha$  intracellular domain (SEQ ID NO: 142), a CD8 $\beta$  signal peptide (SEQ ID NO:144), a CD8 $\beta$  extracellular domain (SEQ ID NO:145), a CD8 $\beta$  transmembrane domain (SEQ ID NO:146), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147);
- 30

- 5
- c. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD8 $\beta$  intracellular domain (SEQ ID NO: 147); or
  - d. a CD8 $\alpha$  signal peptide (SEQ ID NO:139), a CD8 $\alpha$  extracellular domain (SEQ ID NO:140), a CD8 $\alpha$  transmembrane domain (SEQ ID NO:141), and a CD4 intracellular domain (SEQ ID NO: 148).

FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 2A






FIGURE 2C





FIGURE 2D



CD8 Product 4

**FIGURE 3A**  
(CD8A\_CD8 Product 1)





FIGURE 3B

(CD8alpha-P2A-CD8beta\_CD8 Product 2)



**FIGURE 3C**

(CD8A[EC]-CD8B[ID]\_CD8 Product 3)




FIGURE 3D

(CD8a[EC]-CD4[ID]\_CD8 Product 4)







FIGURE 4B



**FIGURE 5**

**NeoTCR Product Nucleic Acid Sequence**

ACATTAACAACAAAATCCTACGGAAATACTGAAGATGAGTCTCAGCACTAAGGAAAAGC  
 CTCCAGCAGCTCTGCTTCTGAGGGTGAAGGATAGACGCTGTGGCTCTGCATGACTCACTA  
 GCACTCTATCACGGCCATATCTGGCAGGGTCAGTGGCTCCAACCTAACATTTGTTGGTACTT  
 TACAGTTTATAATAGATGTTTATATGGAGAAGCTCTCATTTCTTCTCAGAAGAGCCTGGCT  
 AGGAAAGGTGGATGAGGCACCATAATTCATTTGCAGGTGAAATTCCTGAGATGTAAGGAGCT  
 GCTGTGACTTGCTCAAGGCCTTATATCGAGTAAACGGTAGTGTGGGGCTTAGACGCAGGTG  
 TTCTGATTTATAGTTCAAAACCTCTATCAATGAGAGCAATCTCCTGGTAATGTGATAGATTT  
 CCCAACTTAATGCCAACATACCATAAACCTCCCATTCGCTAATGCCCAGCCTAAGTTGGGGA  
 GACCACTCCAGATTCCAAAGATGTACAGTTTGTCTGGGCTTTTCCCATGCCTGCCTTTA  
 CTCTGCCAGAGTTATATTGCTGGGTTTTGAAGAAGATCCTATTAAATAAAAGAATAAGCAGT  
 ATTATTAGTAGCCCTGCATTTCAAGTTTCCCTTGAAGTGGCAGGCCAGGCCCTGGCCGTGAACG  
 TTCACTGAAATCATGGCCTCTTGGCCAAAGATTGATAGCTTGTGCCTGTCCCTGAGTCCCAGTC  
 CATCACGAGCAGCTGTTTCTAAGATGCTATTTCCCGTATAAAGCATGAGACCGTGACTTGCC  
 AGCCCCACAGAGCCCCCTTGTCCATCACTGGCATCTGGACTCCAGCCTGGGTTGGGGCA  
 AAGAGGGAAATGAGATCATGTCTTAACCCCTGATCCTCTTGTCCCACAG

**Left HR Arm (SEQ ID NO:1)**

**TRAC CDS (SEQ ID NO:2 )**

**GSG Linker 1 (SEQ ID NO:3)**

**P2A (SEQ ID NO:4)**

ATATCCAGAACCCCTGACCCCTGCCGTGTACCAGCTGAGAGACTCTAAATCCAGTG  
 ACAAGTCTGTCTGCCCTATTC

GAATTCGGCTCCGGA

GCCACTAACTTCTCCCTGTTGAAACAGGGCTGGCGATGTTGAAGAAAACCCCGG  
 TCCT

**FIGURE 5 Cont.**

**HGH/SS/2 (SEQ ID NO:5)**

ATGGCCACGGCTCTAGAACAAGCCTGCTCGCTTTTGGCCTGCTCTGCCTCCCATGGCTC  
CAAGAAGGATCTGCT

**Insertion site for neoTCR  
TRB\_VDJ sequence**

[[TRB\_VDJ Sequence, specific for the NeoTCR]]

**TCR-beta/constant (SEQ  
ID NO:7)**

CTGAAAACGTGTTCCCTCCAAAAGTGGCCGTGTTTCGAGCCTTCTGAGGCCGAGATCAGCC  
ACACACAGAAAGCCACACTCGTGTCTGGCTACCGGCTTCTACCCCGATCACGTGGAACCTG  
TCTTGGTGGTCAACGGCAAAAGAGGTGCACAGCGGCTCAGCACAGATCCCCAGCCTCTGA  
AAGAACAGCCCGCTCTGAACGACAGCCGCTACTGCCCTGTCTAGCAGACTGAGAGTGTCCGC  
CACCTTCTGGCAGAACCCCAAGAACCTTCAGATGCCAGGTCCAGTTCTACGGCCTGAGCG  
AGAACGATGAGTGGACCCAGGACAGAGCCAAAGCCTGTGACACAGATCGTGTCTGCCGAAG  
CCTGGGCAGAGCCGATTGTGGCTTTACCAGCGAGTCATACCCAGCGGCGTGTCTGTCTGC  
CACCATCTGTATGAGATCCTGCTCGGCAAGGCCACACTGTACGCTGTGCTGGTGTCTGTCTCT  
GGTGTGATGGCTATGCTCTCCCGGAGCGCATCCCCGAGGGC

**Furin Cleavage Site (SEQ ID NO: 8)**

CGGGCCAAAGCGG

**GSG Linker (SEQ ID NO:9)**

GGCAGCGGC

**P2A (SEQ ID NO:10)**

GCCACCAACTCAGCCTGCTGAAGCAGGCCGCGACGTGGAGGAGAAACCCCGGCCCT

**HGH SS (SEQ ID NO:11)**

ATGGCCACAGGCAGACAACATCTCTGTCTGGCCTTCGGACTGCTGTGTCTGCCTTGGCT  
GCAAGAGGGTTCCGCC



**FIGURE 5 Cont.**

CGCGTTGCTGGCGTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAATAATCGACGCT  
 CAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATAACCAGGCGTTTCCCCCTGGAAG  
 CTCCTCGTGGCTCTCCTGTTCCGACCCCTGCCGCTTACCGGATACCTGTCCGCCCTTCTCCCT  
 TCGGGAAGCGTGGCGCTTCTCATAGCTACGCTGAGGTATCTCAGTTCGGGTAGGTCGTT  
 CGTCCAAGCTGGCTGTGTGCACGAAACCCCGTTACGCCGACCCGCTGCCCTTATCCGG  
 TAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATGCCACTGGCAGCAGCCACTG  
 GTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGTACAGAGTCTTGAAGTGGTGCC  
 TAACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTT  
 CGGAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACCAACCCGCTGGTAGCGGTGGTTTT  
 TTTGTTGCAAGCAGCAGATTACGGCGAGAAAAGGATCTCAAGAAAGATCCTTTTGATCT

**pBR322\_origin (SEQ ID NO:17)**

TTAGAAAACCTCATCGAGCATCAAAATGAAACTGCAATTTATTATATCAGGATTATCAATACCA  
 TATTTTTGAAAAGCCGTTTCTGTAATGAAGGAGAAAACCTCACCGAGGCGAGTTCATAGGAT  
 GGCAAGATCCTGGTATCGGCTGCGATTCCGACTCGTCCAACATCAATACAACCTTAAATTC  
 CCTCGTCAAAAATAAGGTTATCAAGTGAGAAATCACCATGAGTGACGACTGAATCCGGTGA  
 GAATGGCAAAAAGTTTATGCAITTTCTCCAGACTTGTCAACAGGCCAGCCATTACGCTCGTC  
 ATCAAAATCACTCGCATCAACCAACCGTTATTCATTCGTGATTGCGCCTGAGCCAGACGAAA  
 TACGCGATCGCTGTTAAAAGGACAATTACAACAGGAATCGAATGCAACCCGGCGCAGGAAC  
 ACTGCCAGCGCATCAACAATATTTTCACTGAATCAGGATATTTCTAATACCTGGAATGCTG  
 TTTTTCCGGGATCGCAGTGGTGAATAACCATGCATCATCAGGAGTACGGATAAATGCTTG  
 ATGGTCGGAAGAGGCATAAATCCGTCAGCCAGTTTAGTCTGACCATCTCATCTGTAACATCA  
 TTGGCAACGCTACCTTTGCCATGTTTCAGAAAACAACCTGGCGCATCGGCTTCCCATACAAG  
 CGATAGATTGTCACCTGATTGCCCGACATTATCGCGAGCCCATTTATACCCCATATAAATCAG  
 CATCCATGTTGGAATTAATCGCGGCTCGACGTTTCCCGTTGAAATATGGCTCAT

**Kanamycin resistant gene (KanR2) (SEQ ID NO:18)**

AACACCCCTTGTTACTGTTTATGTAAGCAGACAGTTTTATTGTTTCATGATGATATATTTTATC  
 TTGTGCAATGTAACATCAGAGATTTTGAGACAC

**Kanamycin Promoter (SEQ ID NO:19)**

**FIGURE 6**

**CD8 Product 1 Product Nucleic Acid Sequence**

ACATTAACAACAAAATCCTACGGAAATACTGAAGATGAGTCTCAGCAGCTAAGGAAAAGC  
 CTCCAGCAGCTCCTGCTTCTGAGGGTGAAAGGATAGACGCTGTGGCTCTGCATGACTCACTA  
 GCACTCTATCAGGCCATATCTGGCAGGGTCAAGTGGCTCCAACATAACATTTGTTGGTACTT  
 TACAGTTTATTAATAGATGTTTATATGGAGAAGCTCTCATTTCTTCTCAGAAAGAGCCTGGCT  
 AGGAAGGTGGATGAGGCACCATAATTCATTTGCAGGTGAAATTCCTGAGATGTAAGGAGCT  
 GCTGTACTTGTCAAGCCCTTATATCGAGTAAACGGTAGTGTGGGCTTAGACGCAGGTG  
 TTCTGATTTATAGTTCAAAACCTCTATCAATGAGAGAGCAATCTCCTGGTAATGTGATAGATTT  
 CCCAACTTAATGCCAACATACCATAAACCTCCCATTCTGCTAATGCCAGCCTAAGTTGGGGA  
 GACCACTCCAGATCCAAAGATGTACAGTTTGTCTGGGCTTTTCCCATGCCTGCCTTTA  
 CTCTGCCAGAGTTATATTGCTGGGGTTTTGAAGAAGATCCTATTAAATAAAAAGAAATAGCAGT  
 ATTATTAAGTAGCCCTGCATTTCAAGTTTCCCTTGTAGTGGCAGGCCAGCCCTGGCCGTGAACG  
 TTCACTGAAATCATGGCCCTTTGGCCAAGATTGATAGCTTGTCCCTGTCCCTGAGTCCCAGTC  
 CATCAGGAGCAGCTGGTTTCTAAGATGCTATTTCCCGTATAAAGCATGAGACCCGTGACTTGCC  
 AGCCCCACAGAGCCCCCCTTGTCCATCACTGGCATCTGGACTCCAGCCTGGGTTGGGGCA  
 AAGAGGAAATGAGATCATGTCCCTAACCCCTGATCCTCTTGTCCCCACAG

**Left HR Arm (SEQ ID NO:22)**

**TRAC CDS (SEQ ID NO:23)**

**GSG Linker 1 (SEQ ID NO:24)**

**P2A (SEQ ID NO:25)**

ATATCCAGAACCCCTGACCCTGCCGTGTACCAGCTGAGAGACTCTAAATCCAGTGACAAAGTCTG  
 TCTGCCTATTC

GAATTCGGCTCCGGA

GCCACTAACTTCAGCCTGTTGAAGCAGGCCGGGACGTTGAGGAAAACCCCGGTCTCT

**FIGURE 6 Cont.**

<p>ATGGCCTTACCAGTGACCGCCTTGCTCCTGCCGCTGGCCTTGCTGTCTCCACGCGCCAGGGCC G</p>	<p><b>CD8A Signal peptide (SEQ ID NO:26)</b></p>
<p>AGCCAGTTCGGGTGTCGCCGCTGGATCGGACCTGGAACCTGGCGGAGACAGTGAGCTG AAGTGCCAGGTGCTGTCCAAACCGACGTCGGGCTGCTCGTGCTCTCCAGCCGCGCG GGCCCGCCAGTCCCACCTTCTCCTATACCTTCCCAAACAAGCCCAAGGCGGCCGAG GGGCTGGACACCCAGCGGTTCTCGGGCAAGAGGTTGGGGACACCTTCGTCTCACCCCTG AGCGACTTCCGCCGAGAGAACGAGGGCTACTATTCTGCTCGGCCCTGAGCAACTCCATCAT GTACTTCAGCCACTTCGTGCCGTCTTCTGCCAGCGAAGCCACCACGACGCCAGCGCCCGC GACCACCAACCGGCCACCATCGCTCGCAGCCCTGTCCCTGCGCCAGAGGCGGTG CCGGCCAGCGGGGGGCGCAGTGACACGAGGGGCTGGACTTCGCCCTGTGAT</p>	<p><b>CD8A Extracellular domain (SEQ ID NO:27)</b></p>
<p>ATCTACATCTGGGGCCCTTGGCCGGACTTGTGGGTCTTCTCCTGTCACTGGTTATCACC</p>	<p><b>CD8A transmembrane domain (SEQ ID NO:28)</b></p>
<p>CTTTACTGCAACCACAGGAACCGAAGACGTGTTGCAAAATGTCCCCGGCCTGTGGTCAAATC GGGAGACAAGCCAGCCTTTCGGCGGATACGTC</p>	<p><b>CD8A Intracellular domain (SEQ ID NO:29)</b></p>
<p>AGGGCTAAACGG</p>	<p><b>Furin cleavage site (SEQ ID NO:30)</b></p>
<p>GAATTCGGCTCCGGA</p>	<p><b>GSG Linker (SEQ ID NO:31)</b></p>
<p>GCCACTAACTTCCCTGTTGAAACAGGCTGGCGATGTTGAGAAACCCCGTCTCT</p>	<p><b>P2A (SEQ ID NO:32)</b></p>
<p>ATGGCCACCGGCTTAGAACAAGCCTGCTCGCTTTTGGCCCTGCTCTGCCCTCCCATGGCTC CAAGAAGGATCTGCT</p>	<p><b>HGH/SS/2 (SEQ ID NO:33)</b></p>

**FIGURE 6 Cont.**

<p>[[TRB_VDJ Sequence, specific for the NeoTCR]]</p> <p>CTGAAAACGTGTTCCCTCCAAAAGTGCCGGTGTTCGAGCCTTCTGAGCCGAGATCAGCC          ACACACAGAAAGCCACACTCGTGTGTGGCTACCGGCTTACCCCCGATCACGTGGAAC TG          TCTTGGTGGTCAACGGCAAAGAGGTGCACAGCGGCGTCAGCACAGATCCCCAGCCTCTGA          AAGAACAGCCCGCTCTGAACGACAGCCGCTACTGCCTGTCTAGCAGACTGAGAGTGTCCGC          CACCTTCTGGCAGAACCCAGAACCCACTTCAGATGCCAGGTCCAGTTCTACGGCCTGAGCG          AGAACGATGAGTGGACCCAGGACAGAGCCAGCCGTGTGACACAGATCGTGTGCCGAAG          CCTGGGCAGAGCCGATTGTGGCTTTACCAGCGAGTCATACCAGGCGGTGCTGTCTGC          CACCATCTGTATGAGATCCTGCTCGGCAAGCCACACTGTACGCTGTGCTGGTGTCTGCTCT          GGTGCTGATGGCTATGGTCTCCCGGAGCGCATCCCCGAGGCC</p>	<p><b>Insertion site for neoTCR          TRB_VDJ sequence</b></p>	<p><b>TCR-beta/constant (SEQ ID          NO:35)</b></p>
<p>CGGGCCAAAGCGG</p> <p>GGCAGCGGC</p> <p>GCCACCAACTCAGCCTGCTGAAGCAGGCGCGGACGTGGAGGAGAACCCCGGCCCT</p> <p>ATGGCCACAGGCAGCAACATCTCTGCTGTGGCCTTCGGACTGCTGTCTGCCTTGGCT          GCAAGAGGGTTCCGCC</p>	<p><b>Furin cleavage site (SEQ ID NO:36)</b></p> <p><b>GSG linker (SEQ ID NO:37)</b></p> <p><b>P2A (SEQ ID NO:38)</b></p> <p><b>HGH/SS (SEQ ID NO:39)</b></p>	<p><b>Insertion site for neoTCR TRA_VDJ sequence</b></p> <p><b>TCR-alpha/constant (SEQ ID          NO:41)</b></p>
<p>[[TRA_VDJ Sequence, specific for the NeoTCR]]</p> <p>ATATTAGAACCCCGATCCTGCTGTGTATCAGCTGCGCGACAGCAAGAGCAGCGACAAAGAGC          GTGTGTTGTTC</p>		

**FIGURE 6 Cont.**

**TRAC CDS/right HR arm  
(SEQ ID NO:42)**

ACCGATTTTGATTCTCAACAATAATGTGCACAAAGTAAGGATTCTGATGTATATACAGAC  
AAAACGTGCTAGACATGAGGCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCA  
ACAAATCTGACTTTGTCATGTGCAAAACGCCCTTCAACAACAGCATTATCCAGAAGACACCTTCT  
TCCCCAGCCCCAGG

**Right HR arm (SEQ ID  
NO:43)**

TAAGGGCAGCTTTGGTGCCTTCGCAGGCTGTTTCCTTGCCTTCAGGAATGGCCAGGTTCTGCC  
CAGAGCTCTGGTCAATGATGTCTAAACTCCTCTGATTGGTGTCTCGGCCTTATCCATTGCC  
ACCAAAACCCCTCTTTTACTAAGAAACAGTGAGCCTTGTCTGGCAGTCCAGAGAATGACAC  
GGGAAAAAAGCAGATGAGAGAAAGGTGGCAGGAGAGGGCACGTGGCCAGCCTCAGTCT  
CTCCAACTGAGTTCCTGCCTGCCTTTGCTCAGACTGTTTGCCCTTACTGCTCTTAGG  
CCTCATTCTAAGCCCCCTCTCCAAGTTGCCCTCCTTATTTCTCCCTGTCTGCCAAAATCTTT  
CCCAGTCACTAAGTCAGTCTCACGCAGTCACTATAACCCCACTACTGATTGTGCCGG  
CACATGAATGCACCCAGGTGTTGAAGTGGAGGAATTAATAAGTCAAGATGAGGGGTGTCGCCA  
GAGGAAGCACCATCTAGTTGGGGAGCCCCATCTGTCACTGGGAAAGTCCAAATAACTT  
CAGATTGGAATGTGTTTTAACTCAGGGTTGAGAAAACAGCTACCTTCAGGACAAAAGTCAG  
GGAAGGGCTCTCTGAAGAATGCTACTTGAAGATACCAGCCCTACCAAGGGCAGGGAGAG  
GACCCTATAGAGCCCTGGGACAGGAGCTCAATGAGAAAGGAGAGAGCAGCAGGCGCATGA  
GTTGAATGAAGGAGGCCAGGGCCGGTCCAGGGCCCTTCTAGGCCCATGAGAGGGTAGACAG

**(SEQ ID NO:44)**

GCTAGC

FIGURE 6 Cont.

pBR322\_origin (SEQ ID NO:45)

CGCGTTGCTGGCGTTTTCCATAGGCTCCGCCCTGACGAGCATCACAAAATCGACGCT
CAAGTCAGAGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTTCCCCCTGGAAG
CTCCCTCGTGGCTCTCCCTGTTCCGACCCTGCCGTTACCGGATACCTGTCCGCCCTTCTCCCT
TCGGGAAGCGTGGCGTTTTTCATAGCTACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTT
CGTCCAAAGCTGGGTGTGCACGAAACCCCGTTACGCCGACCCGCTGCGCCTTATCCGG
TAACTATCGTCTTGAGTCCAACCCGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTG
GTACAGGATTAGCAGAGCGAGGTATGTAGCGGTGTACAGAGTTCTTGAAGTGGTGGCC
TAACTACGGCTACACTAGAAAGAACAGTATTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTT
CGGAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACCAACCCGCTGGTAGCGGTGGTTTT
TTTGTGCAAGCAGCAGATTACGCGCAGAAAAGGATCTCAAGAAAGATCCTTTTGATCT

Kanamycin resistance (KanR2) (SEQ ID NO:46)

TTAGAAAACCTCATCGAGCATCAAATGAACTGCAATTTATTCATATCAGGATATCAATACCA
TATTTTTGAAAAGCCGTTTCTGTAATGAAGGAGAAAACCTCACCGAGGCAGTTCATAGGAT
GGCAAGATCCTGGTATCGGTCTGCCGATCCGACTCGTCCAAACATCAATACAACCTTAAATTC
CCCTCGTCAAAAATAAGTTATCAAGTGAGAAATCACCATGAGTGACGACTGAATCCGGTGA
GAATGGCAAAAGTTATGCATTTCTTCCAGACTTGTTC AACAGGCCAGCCATTACGCTCGTC
ATCAAATCACTCGCATCAACCAACCCGTTATTCATTCTGATTGCGCCTGAGCCAGACGAAA
TAGCGATCGCTGTAAAAGGACAATTACAACAGGAATCGAATGCAACCCGGCAGGAAAC
ACTGCCAGCGCATCAACAATATTTTACCCTGAATCAGGATATCTTCTAATACCTGGAAATGCTG
TTTTCCGGGATCGCAGTGGTGAGTAAACCATGCATCATCAGGAGTACGGATAAAATGCTTG
ATGTCGGAAGAGGCATAAATCCGTCAGCCAGTTAGTCTGACCATCTCATCTGTAACATCA
TTGGCAAACGCTACCTTTGCCAIGTTTCAGAAACAACCTGGCGCATCGGGCTTCCCATACAAG
CGATAGATTGTCGCACCTGATTGCCCGACATTATCGGAGCCCAITATACCATATAAATCAG
CATCCATGTTGGAATTTAATCGCGGCCCTCGACGTTTCCCGTTGAATATGGCTCAT

Kanamycin promoter (SEQ ID NO:47)

AACACCCCTTGTTACTGTTTATGTAAGCAGACAGTTTTATTTGTTTCATGATGATATATTTTATC
TTGTGCAATGTAAACATCAGAGATTTTGAGACAC

**FIGURE 7**

**CD8 Product 2 Product Nucleic Acid Sequence**

ACATTAAAACACAAAATCCTACGGAAATACTGAAGAATGAGTCTCAGCAGCTAAGGAAAAGC  
 CTCAGCAGCTCCTGCTTCTGAGGGTGAAGGATAGACGCTGTGGCTCTGCATGACTCACTA  
 GCACTCTATCAGGCCATATCTGGCAGGGTCAAGTGGCTCCAACATAACATTTGTTGGTACTT  
 TACAGTTTATTAATAGATGTTTATATGGAAAGCTCTCATTCTTCTCAGAAGAGCCTGGCT  
 AGGAAGGTGGATGAGGCACCATAATTCATTTGCAGGTGAAATTCCTGAGATGTAAGGAGCT  
 GCTGTACTTGCTCAAAGGCTTATATCGAGTAAACGGTAGTGCTGGGCTTAGACGCAGGTG  
 TTCTGATTTATAGTTCAAAACCTCTATCAATGAGAGAGCAATCTCTGGTAATGTGATAGATTT  
 CCCAACTTAATGCCAACATACCATAAACCTCCCATCTGTAATGCCAGCCTAAGTTGGGGA  
 GACCACTCAGATTCCAAAGATGTACAGTTTGTCTGGGCTTTTCCCATGCCTGCCTTTTA  
 CTCTGCCAGAGTTATATTGCTGGGTTTTGAAGAAGATCCTATTAATAAAGAATAAGCAGT  
 ATTATTAAGTAGCCCTGCATTTCCAGGTTTCTTGAGTGGCAGGCCAGCCGTGGCCGTGAACG  
 TTCACTGAAATCATGGCCTCTTGCCCAAGATTGATAGCTTGTGCCTGTCCCTGAGTCCCCAGTC  
 CACTCAGCAGCTGGTTCTAAGATGCTATTTCCCGTATAAAGCATGAGACCCGTGACTTGCC  
 AGCCCCACAGAGCCCCCTTGTCCATCACTGGCATCTGGACTCCAGCCTGGGTTGGGGCA  
 AAGAGGAAATGAGATCATGTCTAACCCTGATCCTCTTGTCACACAG

**Left HR Arm (SEQ ID NO:49)**

**TRAC CDS (SEQ ID NO:50)**

**GSG Linker 1 (SEQ ID NO:51)**

**P2A (SEQ ID NO:52)**

ATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGACTCTAAATCCAGTGACAAGTCTG  
 TCTGCCCTATTC

GAATTCGGCTCCGGA

GCCACTAACTTCAGCCTGTTGAAGCAGGCCGGCAGCTTGAGGAAAACCCCGGTCCT

**FIGURE 7 Cont.**

<p>ATGGCCTTACCAGTGACCGCCTTGCTCCTGCCGCTGGCCTTGCTGTCTCCACGCCCGCCAGGCC G</p>	<p><b>CD8A Signal peptide (SEQ ID NO:53)</b></p>
<p>AGCCAGTTCGGGTGTCGCCGCTGGATCGGACCTGGAACCTGGGCGGAGACAGTGGAGCTG AAGTGCCAGGTGCTGTCCAACCCGACGTCGGGTGCTCGTGGCTCTTCCAGCCGCGG GCGCCCGCCAGTCCCACCTTCTCTATACCTCTCCAAAACAAGCCCAAGGCGGCCGAG GGGCTGGACACCCAGCGGTTCTCGGGCAAAGAGGTTGGGGACACCTTCGTCTCACCCCTG AGCGACTTCGCGGAGAGAACGAGGGCTACTATTTCTGCTCGGCCCTGAGCAACTCCATCAT GTACTTCAGCCACTTCGTGCCGTCTTCTGCCAGCGAAGCCCAACACGACCCAGCGGCCGC GACCACCAACCGGCCACCATCGCTCGCAGCCCTGTCCCTGCGCCAGAGGCGGTG CCGGCCAGCGGGGGCGCAGTGACACAGGGGGCTGGACTTCGCCTGTGAT</p>	<p><b>CD8A Extracellular domain (SEQ ID NO:54)</b></p>
<p>ATCTACATCTGGGCGCCCTTGGCCGGACTTGTGGGTCCTTCTCTGTCACCTGGTTATCACC</p>	<p><b>CD8A transmembrane domain (SEQ ID NO:55)</b></p>
<p>CTTTACTGCAACCACAGGAACCGAAGACGTGTTTGCAAATGTCCCCGGCCTGTGGTCAAATC GGGAGACAAAGCCAGCCCTTCGGCGGAGATACGTC</p>	<p><b>CD8A Intracellular domain (SEQ ID NO:56)</b></p>
<p>AGAGCAAAGCGG</p>	<p><b>Furin cleavage site (SEQ ID NO:57)</b></p>
<p>GGCTCCGGA</p>	<p><b>GSG Linker (SEQ ID NO:58)</b></p>
<p>GCTACCAATTTAGCCTCTGAAGCAGGCTGGCGATGTTGAGGAAAACCCTGGTCCC</p>	<p><b>P2A (SEQ ID NO:59)</b></p>
<p>ATGCGCGCGGCTGTGGCTCCTCTTGGCCGCGCAGCTGACAGTTCTCCATGGCAACTCAGT C</p>	<p><b>CD8B Signal Peptide (SEQ ID NO:60)</b></p>

**FIGURE 7 Cont.**

<p>CTCAGCAGACCCCTGCATACATAAAGGTGCAAAACCAACAAGATGGTGATGCTGTCTGCGGA GGTAAATCTCCCTCAGTAACATGCGCATCTACTGGCTGAGACAGGCCAGGCCACCGAGCA GTGACAGTCAACCAGATTCCCTGCCCTCTGGGATCCGCAAAAGGGACTATCCACGGTGA AGAGTGGAACACAGGAGAGATAGCTGTGTTTCGGGATGCAAGCCGGTTCATTCTCAATCTC ACAAGCGTGAAGCCGGAAGACAGTGGCATCTACTTCTGCATGATCGTGGGAGCCCGGAGC TGACCTCGGGAAGGAACTCAGCTGAGTGTGGTTGATTTCCCTCCACCACTGCCCGAGCCCC ACCAAGAAGTCCACCCTCAAGAAAGAGAGTGTGCCGTTACCCAGGCCAGAGACCCAGAAAG GGCCCACTTTGTAGCCCC</p>	<p><b>CD8B Extracellular domain (SEQ ID NO:61)</b></p>
<p>ATCACCCTTGGCCTGCTGGTGGCTGGCGTCCCTGGTTCTGCTGGTTCCCTGGGAGTGCCCAT C</p>	<p><b>CD8B Transmembrane domain (SEQ ID NO:62)</b></p>
<p>CACCTGTGCTGCCGGGAGGAGAGCCCGGCTTCGTTTCATGAACAATTTACAAA AGGGCTAAACGG GAATTCGGCTCCGGA GCCACTAATTCTCCCTGTTGAAACAGGCTGGCGATGTTGAAGAAAACCCCGTCTCT ATGGCCACCGGCTCTAGAACAAGCCTGCTGCTCGCTTTTGGCCTGCTCTGCCCTCCCATGGCTC CAAGAAGGATCTGCT</p>	<p><b>CD8B Intracellular domain (SEQ ID NO:63)</b></p> <p><b>Furin cleavage site (SEQ ID NO:64)</b></p> <p><b>GSG Linker (SEQ ID NO:65)</b></p> <p><b>P2A (SEQ ID NO:66)</b></p> <p><b>HGH/SS/2 (SEQ ID NO:67)</b></p>

**FIGURE 7 Cont.**

[[TRB\_VDJ Sequence, specific for the NeoTCR]]

CTGAAAAACGTGTTCCCTCCAAAAGTGGCCGTGTTTCGAGCCTTCTGAGGCCGAGATCAGCC  
ACACACAGAAAGCCACACTCGTGTGTCTGGCTACCCGGCTTACCCCGATCACGTGGAACGTG  
TCTTGGTGGTCAACGGCAAAGAGGTGCACAGCGGGCTCAGCACAGATCCCCAGCCTCTGA  
AAGAACAGCCCGCTCTGAACGACAGCCGCTACTGCCGTCTAGCAGACTGAGAGTGTCCGC  
CACCTTCTGGCAGAACCCAGAAACCACTTCAGATGCCAGGTCACAGTTCTACGGCCTGAGCG  
AGAACGATGAGTGGACCCAGGACAGAGCCAAAGCCCTGTGACACAGATCGTGTCTGCCGAAG  
CCTGGGCAGAGCCGATTGTGGCTTACCAGCGAGTCATACCAGCGGGCGTGTCTGTCTGC  
CACCATCTGTATGATCCTGCTCGGCAAGGCCACACTGTACGCTGTGCTGGTGTCTGCTCT  
GGTGCTGATGGCTATGGTCTCCCGGGAGCGCATCCCCGAGGCC

**Insertion site for neoTCR  
TRB\_VDJ sequence**

**TCR-beta/constant (SEQ ID  
NO:69)**

CGGGCCAAGCGG

GGCAGCGGC

GCCACCAACTCAGCCTGCTGAAGCAGGCCCGGCGACGTGGAGGAGAAACCCCGGCCCT P2A (SEQ ID NO:72)

ATGGCCACAGGCAGAGAACATCTCTGCTGTGGCCTTCGGACTGCTGTCTGCCCTTGCT  
GCAAGAGGGTCCCGCC HGH/SS/2 (SEQ ID NO:73)

[[TRA\_VDJ Sequence, specific for the NeoTCR]]

ATATTCAGAAACCCCGATCCTGCTGTGTATCAGCTGCCGACAGCAAGAGCAGCGACAAGAGC  
GTGTGTTTGTTC

**Insertion site for neoTCR  
TRA\_VDJ sequence**

**TCR-alpha/constant (SEQ ID  
NO:75)**

**Furin cleavage site (SEQ ID NO:70)**

**GSG Linker (SEQ ID NO:71)**

**FIGURE 7 Cont.**

**TRAC CDS/right HR arm  
(SEQ ID NO:76)**

ACCGATTTTGATTCTCAAACAATGTGTACAAAGTAAAGGATTCTGATGTGTATATCACAGAC  
AAAACGTGTCTAGACATGAGGCTATGGACTTCAAGAGCAACAGTGTCTGGCCTGGAGCA  
ACAAATCTGACTTTGCATGTGCAAACGCCCTTCAACAACAGCATTATCCAGAAAGACACCTTCT  
TCCCCAGCCCCAGG

**Right HR arm (SEQ ID  
NO:77)**

TAAGGCAGCTTTGGTGCCTTCGCAGGCTGTTTCCTTGCTTCAGGAATGCCAGGTTCTGCC  
CAGAGCTCTGGTCAATGATGCTAAACTCCTCTGATTGGTGTCTCGGCCTTATCCATTGCC  
ACCAAACCCCTCTTTTACTAAGA AACAGT GAGCCTTGTTCTGGCAGTCCAGAGAATGACAC  
GGGAAAAAAGCAGATGAAAGAGAGGTTGGCAGGAGGGCACGTGGCCAGCCTCAGTCT  
CTCCAACTGAGTTCCTGCCTGCCCTTTGCTCAGACTGTTTGCCCTTACTGCTCTTCTAGG  
CCTCATTCTAAGCCCCCTTCTCCAAGTTGCCCTCCTTATTTCTCCCTGCTGCCAAAAATCTTT  
CCCAGCTCACTAAGTCAGTCTACGCAGTCACTCATTAACCCACCAATCACTGATTGTGCCGG  
CACATGAATGCCACCAGGTTTGAAGTGAGGAATTA AAAAGTCAGATGAGGGGTGTGCCCA  
GAGGAAGCACCATCTAGTTGGGGGAGCCCATCTGTCAGCTGGGAAAAGTCCAAATAACTT  
CAGATTGGAATGTTTTAACTCAGGTTGAGAAAACAGTACCTTCAGGACAAAAGTCAG  
GGAAGGCTCTGAAAGAAATGCTACTTGAAGATACCAGCCCTACCAAGGCAGGGAGAG  
GACCCATAGAGGCCCTGGGACAGGAGCTCAATGAGAAAAGGAGAAAGAGCAGCAGGCATGA  
GTTGAATGAAGGAGGCGCCGGTCCAGGGCCTTCTAGGCCATGAGAGGGGTAGACAG

**(SEQ ID NO:78)**

GCTAGC

**FIGURE 7 Cont.**

**pBR322\_origin (SEQ ID NO:79)**

CGGGTTGCTGGCGTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAATAATCGACGCT  
CAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATAACACAGGCGTTTCCCCCTGGAAG  
CTCCCTCGTGGCTCTCCTGTTCCGACCCCTGCCCTACCGGATAACCTGTCCGCCCTTCTCCCT  
TCGGGAAGCGTGGCGCTTCTCATAGCTCACGCTGAGGTATCTCAGTTCGGTG TAGGTCGTT  
CGTCCAAGCTGGGCTGTGTCACGAACCCCGTTAGCCCGACCGCTGCGCCTTATCCGG  
TAACTATCGTCTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTG  
GTAAACAGGATTAGCAGAGCGAGGTATGAGCGGTGTACAGAGTCTTGAAGTGGTGGCC  
TAACTACGGCTACACTAGAAAGAACAGTATTGGTATCTGCGCTGTGTAAGCCAGTTACCTT  
CGGAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACACCCGCTGTGAGCGGTGGTTTT  
TTTGTTCGCAAGCAGCAGATTACGCGCAGAAAAGGATCTCAAGAAAGATCCTTTGATCT

**Kanamycin resistance  
(KanR2) (SEQ ID NO:80)**

TTAGAAAACATCGAGCATCAAATGAAACTGCAATTTATTCAATACGAGATTATCAATACCA  
TATTTTTGAAAAAGCCGTTCTGTAATGAAGGAGAAAACCTACCGAGGCAGTTCATAGGAT  
GGCAAGATCCTGGTATCGGCTGCGATCCGACTCGTCCAAACATCAATACAACCTATTAATTC  
CCCTCGTCAAAAATAAGTTATCAAGTGAGAAATCACCATGATGACGACTGAATCCGGTGA  
GATGGCAAAAAGTTATGCATTTCTTCCAGACTTGTCAACAGGCCAGCCATTACGCTCGTC  
ATCAAAATCACTCGCATCAACCAACCGTTATTCATTCGTGATTGCGCCTGAGCCAGACGAAA  
TACGCGATCGCTGTAAAAGGACAATTAACAACAGGAATCGAATGCAACCCGGCGCAGGAAC  
ACTGCCAGCGCATCAACAATATTTACCTGAATCAGGATATCTTCTAATACCTGGAATGCTG  
TTTTTCCGGGATCGCAGTGGTGAATACCATGCATCATCAGGAGTACGGATAAAATGCTTG  
ATGTCGGAAGAGGCATAAATCCGTCAGCCAGTTAGTCTGACCATCTCATCTGTAACATCA  
TTGGCAACGCTACCTTTGCCATGTTTCAGAAACAACCTGGCGCATCGGGCTTCCCATACAAG  
CGATAGATTGTCGCACCTGATTGCCCGACATATCGCGAGCCCATTTATACCCATATAAATCAG  
CATCCATGTTGGAATTAATCGCGGCCTCGACGTTTCCCGTTGAATATGGCTCAT

**Kanamycin promoter (SEQ ID NO:81)**

AACACCCCTTGATTACTGTTTATGTAAGCAGACAGTTTTATTGTTTCATGATGATATATTTTATC  
TTGTGCAATGTAACATCAGAGATTTTGAGACAC

**FIGURE 8**

**CD8 Product 3 Product Nucleic Acid Sequence**

ACATTAAACACAAAATCCTACGGAAATACTGAAGAATGAGTCTCAGCCTAAGGAAAAGC  
 CTCAGCAGCTCCTGCTTTCTGAGGGTGAAGGATAGACCGCTGTGGCTCTGCATGACTCACTA  
 GCACTCTACAGGCCATATTCTGCAGGGTCAAGTGGCTCCAACATAACATTTGTTTGGTACTT  
 TACAGTTTATAAATAGATGTTTATATGGAAGCTCTCATTCTTCTCAGAAAGAGCCTGGCT  
 AGAAAGGTGATGAGGCCACCATATTCATTTGCAAGGTGAAATTCCTGAGATGTAAGGAGCT  
 GCTGTGACTTGTCAAGGCCTTATCGAGTAAACGGTAGTGTGGCTTAGACGCAGGTG  
 TTCTGATTTATAGTTCAAAACCTCTATCAATGAGAGAGCAATCCTGGTAATGTGATAGATTT  
 CCCAACTTAATGCCAACATACCATAAACCTCCCATCTGCTAATGCCAGCCTAAGTTGGGGA  
 GACCACTCCAGATCCAAGATGTACAGTTTGTCTTGTGGCCTTTTCCCATGCCTGCCCTTAA  
 CTCTGCCAGAGTTATATTGCTGGGTTTTGAAGAAGATCCTATTAATAAAGAATAAGCAGT  
 ATTATTAAGTAGCCCTGCATTTCCAGTTTCCCTGAGTGGCAGGCCAGCCGTGGCCGTGAACG  
 TTCACTGAAATCATGGCCTCTTGCCCAAGATTGATAGCTTGTGCCTGTCCCTGAGTCCCAGTC  
 CATCAGAGCAGCTGGTTTCTAAGATGCTATTTCCCGTATAAAGCATGAGACCCGTGACTTGCC  
 AGCCCCACAGAGCCCCCCTTGTCCATCACTGGCATCTGGACTCCAGCCCTGGGTTGGGGCA  
 AAGAGGAAATGAGATCATGTCCTAACCCCTGATCCTCTTGTCCCCACAG

**Left HR Arm (SEQ ID NO:83)**

**TRAC CDS (SEQ ID NO:84)**

**GSG Linker 1 (SEQ ID NO:85)**

**P2A (SEQ ID NO:86)**

ATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGACTCTAAATCCAGTGACAAGTCTG  
 TCTGCCATTTC  
 GAATTCGGCTCCGGA  
 GCCACTAACTTCAGCCTGTTGAAGCAGGCCCGGACGTTGAGGAAAACCCCGGTCTC

**FIGURE 8 Cont.**

ATGGCCTTACCAGTGACCGCCTTGCTCCTGCCGCTGGCCTTGCTGTCTCCACGCCCGCAGGCC G	<b>CD8A Signal peptide (SEQ ID NO:87)</b>
AGCCAGTCCGGGTGTCGCCGCTGGATCGGACCTGGAACCTGGCGGAGACAGTGGAGCTG AAGTGCCAGGTGCTGTCCAACCCGACGTGGGCTGCTCGTGGCTCTTCCAGCCGCGCG GCCCGCCCGCAGTCCACCTTCTCTATACCTCTCCAAACAAGCCCAAGGCCGCGGAG GGCTGGACACCCAGCGGTTCTCGGGCAAGAGGTTGGGGACACCTTCGTCTCACCCCTG AGCGACTTCGCCGAGAGAACGAGGGCTACTATTCTGCTCGGCCCTGAGCAACTCCATCAT GTACTTCAGCCACTTCGTCCGGTCTTCTGCCAGCGAAGCCACCACGACGCCAGCCCGC GACCACCAACCGGCCACCATCGCGTCGAGCCCTGTCCCTGGCCCAAGAGCGGTG CCGGCCAGCGGGGGCGCAGTGCACACGAGGGGCTGGACTTCGCCTGTGAT	<b>CD8A Extracellular domain (SEQ ID NO:88)</b>
ATCTACATCTGGCGCCCTTGCCGGACTTGTGGGTCTCTCTGCTCACTGGTTATCACC	<b>CD8A transmembrane domain (SEQ ID NO:89)</b>
CACCTGTGCTGCCGGGAGGAGAGCCCGCTTCGTTTCATGAACAATTTTACAAA	<b>CD8B Intracellular domain (SEQ ID NO:90)</b>
AGGGCTAAACGG	<b>Furin cleavage site (SEQ ID NO:91)</b>
GAATTCGGCTCCGGA	<b>GSG Linker (SEQ ID NO:92)</b>
GCCACTA ACTTCTCCTGTTGAACAGGCTGGGATGTTGAAGAAACCCCGGTCTCT	<b>P2A (SEQ ID NO:93)</b>
ATGGCCACCGGCTAGAACAAAGCCTGCTCGCTTTTGGCCCTGCTGCTCCCATGGCTC CAAGAAGGATCTGCT	<b>HGH/SS/2 (SEQ ID NO:94)</b>

**FIGURE 8 Cont.**

<p>[[TRB_VDJ Sequence, specific for the NeoTCR]]</p> <p>CTGAAAACGTGTTCCCTCCAAAAGTGGCCGTGTTGAGCCCTTCTGAGGCCGAGATCAGCC          ACACACAGAAAGCCACACTCGTGTGTGCTACCGGCTTACCCCGATCACGTGGAACCTG          TCTTGGTGGTCAACGGCAAAGAGGTGCACAGCGGCGTACAGACAGATCCCCAGCCTCTGA          AAGAACAGCCCGCTCTGAACGACAGCCGCTACTGCCCTGTCTAGCAGACTGAGAGTGTCCGC          CACCTTCTGGCAGAACCCAGAAACCACTTCAGATGCCAGGTCACAGTTCTACGGCCTGAGCG          AGAACGATGAGTGGACCCAGGACAGAGCCAAAGCCTGTGACACAGATCGTGTCTGCCGAAG          CCTGGGCAGAGCCGATTGTGGCTTTACCAGCGAGTCATACCAAGCAGGGCGTGTCTGTGC          CACCATCCTGTATGATCCTGTCTCGCAAGGCCACACTGTACGCTGTGTGTGTCTGTCTCT          GGTGCTGATGGCTATGGTCTCCCGGGAGGCGCATCCCCGAGGCC</p>	<p><b>Insertion site for neoTCR TRB_VDJ sequence</b></p>
<p>CGGGCCAAGCGG</p> <p>GGCAGCGGC</p> <p>GCCACCAACTCAGCCTGCTGAAGCAGGCCGCGACGTGGAGGAGAACCCCGGCCCT P2A (SEQ ID NO:99)</p> <p>ATGGCCACAGGCAGCAGAACATCTCTGTCTGTGGCCTTCGGACTGCTGTGTCTGCCTTGGCT          GCAAGAGGGTCCGCC</p>	<p><b>TCR-beta/constant (SEQ ID NO:96)</b></p> <p><b>Furin cleavage site (SEQ ID NO:97)</b></p> <p><b>GSG Linker (SEQ ID NO:98)</b></p> <p><b>HGH/SS/2 (SEQ ID NO:100)</b></p>
<p>[[TRA_VDJ Sequence, specific for the NeoTCR]]</p> <p>ATATTCAGAACCCCGATCCTGCTGTGTATCAGCTGCGGACAGCAAGAGCAGCGACAAGAGC          GTGTGTTTGTTC</p>	<p><b>Insertion site for neoTCR TRA_VDJ sequence</b></p> <p><b>TCR-alpha/constant (SEQ ID NO:102)</b></p>

**FIGURE 8 Cont.**

**TRAC CDS/right HR arm  
(SEQ ID NO:103)**

ACCGATTTTGATTCTCAACAATGTGTCAAAAGTAAGGATTCTGATGTGTATATACAGAC  
AAAAGTGTAGACATGAGGTCTATGGACTTCAAGAGCAACAGTGTGGCCTGGAGCA  
ACAAATCTGACTTTGCATGTGCAAAACGCCCTTCAACAACAGCATTATCCAGAAGACACCTTCT  
TCCCCAGCCCCAGG

**Right HR arm (SEQ ID  
NO:104)**

TAAGGGCAGCTTTGGTGCCTTCGCAGGGTGTTTCCTTGCTTCAGGAATGGCCAGGTTCTGCC  
CAGAGCTCTGGTCAATGATGTCTAAACTCCTCTGATTGGTGTCTCGGCCTTATCCATTGCC  
ACCAAAACCCCTCTTTTACTAAGAAACAGTGAGCCCTGTTCTGGCAGTCCAGAGAATGACAC  
GGGAAAAAGCAGATGAAGAGAAAGGTGGCAGGAGGGCACGTGGCCAGCCTCAGTCT  
CTCCAACTGAGTTCCTGCCTGCCTTTGCTCAGACTGTTTGCCCTTACTGCTCTTCTAGG  
CCTCATTCTAAGCCCCCTTCTCCAAGTTGCCCTCTCCTTATTTCTCCCTGTCTGCCAAAAATCTTT  
CCCAGTCACTAAGTCAGTCTCACGCAGTCACTATAACCCACCAATCACTGATTGTGCCGG  
CACATGAATGCACCCAGGTGTTGAAGTGGAGGAATTAATAAGTCAAGATGAGGGGTGTGCCCA  
GAGGAAGCACCATCTAGTTGGGGAGCCCCATCTGTCAAGTGGGAAAAGTCCAAATAACTT  
CAGATTGGAATGTGTTTTAACTCAGGGTTGAGAAAACAGTACCTTCAGGACAAAAGTCAG  
GGAAGGGCTCTCTGAAGAAATGCTACTTGAAGATACCAGCCCTACCAAGGGCAGGGAGAG  
GACCCTATAGAGGCCCTGGGACAGGAGCTCAATGAGAAAGGAGAGAGCAGCAGGCATGA  
GTTGAATGAAGGAGGCCAGGGCCGCTCACAGGGCTTCTAGGCCCATGAGAGGGGTAGACAG

**(SEQ ID NO:105)**

GCTAGC

**FIGURE 8 Cont.**

**pBR322\_origin (SEQ ID NO:106)**

CGCGTTGCTGGCGTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAATAATCGACGCT  
CAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTTCCCCCTGGAAG  
CTCCCTCGTGGCTCTCCTGTTCCGACCCCTGCCGCTTACCGGATACCTGTCCGCCCTTCTCCCT  
TCGGGAAGCGTGGCGCTTCTCATAGCTCACGCTGAGGTATCTCAGTTCGGTG TAGGTCGTT  
CGTCCAAGCTGGGCTGTGACGAAACCCCGTTAGCCCGACCGCTGGCCCTATCCGG  
TAACTATCGTCTTGAGTCCAACCCGGTAAAGACACGACTTATCGCCACTGGCAGCAGCCACTG  
GTAAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTCTACAGAGTTCTTGAAGTGGTGGCC  
TAACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTT  
CGGAAAAGAGTTGGTAGCTCTTGATCCGGCAACAACCCAGCCGCTGGTAGCGGTGTTTTT  
TTTGTGTTGCAAGCAGCAGATTACGGCGCAGAAAAGGATCTCAAGAAAGATCCTTTGATCT

**Kanamycin resistance (KanR2) (SEQ ID NO:107)**

TTAGAAAACCTCATCGATCAAAATGAAACTGCAATTTATTCATATCAGGATTATCAATACCA  
TATTTTTGAAAAAGCCGTTTTCTGTAATGAAGGAGAAAACCTCACCGAGGCGAGTTCCTATAGGAT  
GGCAAGATCCTGGTATCGGCTCGGATCCGACTCGTCCAAACATCAATACAACCTATTAATTC  
CCCTCGTCAAAAATAAGTTATCAAGTGAGAAATCACCATGAGTGACGACTGAATCCCGGTGA  
GAATGGCAAAAAGTTTTATGCATTTCTTCCAGACTTGTCAACAGGCCAGCCATTACGCTCGTC  
ATCAAAATCACTCGCATCAACCAACCCGTTATTCATTCGTGATTGCGCCTGAGCCAGACGAAA  
TACGGATCGCTGTAAAGGACAATTAACAACAGGAATCGAATGCAACCCGGCCAGGAAC  
ACTGCCAGCGCATCAACAATATTTCCACCTGAATCAGGATATCTTCTAATACCTGGAATGCTG  
TTTTTCCGGGATCGCAGTGGTGAGTAAACCATGCATCATCAGGAGTACGGATAAAATGCTTG  
ATGGTCGGAAGAGGCATAAATCCGTCAGCCAGTTTAGTCTGACCATCTCATCTGTAACATCA  
TTGGCAACGCTACCTTTGCCATGTTTCAGAAAACAACCTGGCGCATCGGGCTTCCCATACAAG  
CGATAGATTGTCACCTGATTGCCCGACATTATCGCGAGCCCAATTTATACCCCATATAAATCAG  
CATCCATGTTGGAATTAATCGCGGCCCTCGACGTTTCCCGTTGAAATATGGCTCAT

**Kanamycin promoter (SEQ ID NO:108)**

AACACCCCTTGTAATCTGTTTATGTAAGCAGACAGTTTTTATTGTTCAIGATGATAIATTTTTATC  
TTGTGCAATGTAACATCAGAGATTTTGAGACAC

**FIGURE 9**

**CD8 Product 4 Product Nucleic Acid Sequence**

ACATTAAAACAAAAATCCTACGGAAATACTGAAGAATGAGTCTCAGCACTAAGGAAAAGC  
 CTCAGCAGCTCCTGCTTCTGAGGGTGAAGGATAGACGCTGTGGCTCTGCATGACTCACTA  
 GCACTCTATCAGGCCATATCTGGCAGGTCAGTGGCTCCAATAACATTTGTTGGTACTT  
 TACAGTTTATTAATAGATGTTTATATGGAGAAGCTCTCATTTCTTCTCAGAAAGAGCCTGGCT  
 AGGAAGGTGGATGAGGCACCATAATTCATTTTGCAGGTGAAATTCCTGAGATGTAAGGAGCT  
 GCTGTACTTGCTCAAGGCTTATATCGAGTAAACGGTAGTGTGGGGCTTAGACGCAGGTG  
 TTCTGATTTATAGTTCAAAACCTCTATCAATGAGAGCAATCTCCTGGTAAATGTGATAGATTT  
 CCCAACTTAATGCCAACATACCATAAACCTCCCATTTCTGTAATGCCAGCCTAAGTTGGGGA  
 GACCACTCCAGATTCCAAGATGTACAGTTTGTCTGGGCTTTTCCCATGCCCTGCCCTTTA  
 CTCGTCCAGAGTTATATTGCTGGGTTTTGAAGAAGATCCTATTAAATAAAAGAATAAGCAGT  
 ATTATTAAGTAGCCCTGCATTTCAAGTTTCTTGTAGTGGCAGGCCAGGCTGGCCGTGAACG  
 TTCACTGAAATCATGGCCTCTTGGCCAAGATTGATAGCTTGTGCCCTGTCCCTGAGTCCCGATC  
 CATCAGAGCAGCTGTTTCTAAGATGCTATTTCCCGTATAAAGCATGAGACCGTGACTTGCC  
 AGCCCCACAGAGCCCCCTTGTCCATCACTGGCATCTGGACTCCAGCCTGGGTTGGGGCA  
 AAGAGGGAATGAGATCATGTCTTAACCCCTGATCCTCTTGTCCCCACAG

**Left HR Arm (SEQ ID NO:110)**

**TRAC CDS (SEQ ID NO:111)**

**GSG Linker 1 (SEQ ID NO:112)**

**P2A (SEQ ID NO:113)**

ATATCCAGAACCCCTGACCCCTGCCGTGTACCAGCTGAGAGACTCTAAATCCAGTGACAAAGTCTG  
 TCTGCCTATTC

GAATTCGGCTCCGGA

GCCACTAACTTCAGCCTGTTGAAGCAGGCGCCGGCAGGTTGAGGAAAACCCCGGTCTCT

**FIGURE 9 Cont.**

<p>ATGGCCTTACCAGTGACCGCCTTGCTCCTGCCGCTGGCCTTGCTGTCTCCACGCCCGCAGGCC G</p>	<p><b>CD8A Signal peptide (SEQ ID NO:114)</b></p>
<p>AGCCAGTTCGGGTGTCGCCGCTGGATCGGACCTGGAACCTGGCGGAGACAGTGGAGCTG AAGTGCCAGGTGCTGTCCAAACCCGACGTGGGCTGCTGGTCTTCCAGCCGCGCG GCGCCGCCAGTCCCACCTTCTCTATACCTTCCCAAACAAGCCCAAGGCGGCCGAG GGGTGGACACCCAGCGGTTCTCGGGCAAGAGGTTGGGGACACCTTCGTCTCACCCCTG AGCGACTTCGCCGAGAGAACGAGGGCTACTATTCTGCTCGGCCCTGAGCAACTCCATCAT GTACTTCAGCCACTTCGTGCCGGTCTTCTGCCCAGCGAAGCCCAACCCAGCCAGCCGCCG GACCAACAACCGCGCCACCATCGCGTCGAGCCCTGTCCCTGGCCCCAGAGGCGTG CCGGCCAGCGGGGGCGCAGTGCACACGAGGGGGCTGGACTTCGCCTGTGAT</p>	<p><b>CD8A Extracellular domain (SEQ ID NO:115)</b></p>
<p>ATCTACATCTGGGCGCCCTTGGCCGGACTTGTGGGTCTCTCTGTCACCTGGTTATCACC</p>	<p><b>CD8A transmembrane domain (SEQ ID NO:116)</b></p>
<p>TGTGTCAAGTGCCGGCACCGAAGGCGCCCAAGCAGAGCGGATGTCTCAGATCAAGAGACTCC TCAGTGAGAAGAAGACCTGCCAGTGTCTCACCAGTTTCAGAAAGACATGTAGCCCCATT</p>	<p><b>CD4 Intracellular domain (SEQ ID NO:117)</b></p>
<p>AGGGCTAAACGG</p>	<p><b>Furin cleavage site (SEQ ID NO:118)</b></p>
<p>GAATTCGGCTCCGGA</p>	<p><b>GSG Linker (SEQ ID NO:119)</b></p>
<p>GCCACTA ACTTCTCCCTGTTGAAACAGGCTGGCGATGTTGAAGAAACCCCGTCTCT</p>	<p><b>P2A (SEQ ID NO:120)</b></p>
<p>ATGGCCACCGGCTCTAGAACAAGCCTGTGCTCGCTTTTGGCCCTGCTCTGCCCTCCCATGGCTC CAAGAAGGATCTGCT</p>	<p><b>HGH/SS/2 (SEQ ID NO:121)</b></p>

**FIGURE 9 Cont.**

<p>[[TRB_VDJ Sequence, specific for the NeoTCR]]</p> <p>CTGAAAACGTGTTCCCTCCAAAAGTGGCCGTGTTGAGCCCTTCTGAGGCCGAGATCAGCC          ACACACAGAAAGCCACACTCGTGTGCTGGCTACCGGCTTACCCCGATCACGTGGAAC TG          TCTTGGTGGTCAACGGCAAAGAGGTGCACAGCGGGCTCAGCACAGATCCCCAGCCTCTGA          AAGAACAGCCCGCTCTGAACGACAGCCGCTACTGCCCTGTCTAGCAGACTGAGAGTGTCCGC          CACCTTCTGGCAGAACCCAGAAACCACTTCAGATGCCAGGTCAGTTCTACGGCCTGAGCG          AGAACGATGAGTGGACCCAGGACAGAGCCAAAGCCCTGTGACACAGATCGTGTGCCGAAG          CCTGGGCAGAGCCGATTGTGGCTTACCAGCGAGTCATACCAGCGGGCGTGTCTGTGC          CACCATCCTGATGATCCTGCTCGGCAAGGCCACACTGTACGCTGTGCTGTGCTGTGCTCT          GGTGCTGATGGCTATGGTCTCCCGGAGCGCATCCCCGAGGCC</p>	<p><b>Insertion site for neoTCR          TRB_VDJ sequence</b></p>
<p>CGGGCCAAAGCGG</p> <p>GGCAGCGGC</p> <p>GCCACCAACTTCAGCCTGCTGAAGCAGGCCCGGCGACGTGGAGGAGAAACCCCGGCCCT P2A (SEQ ID NO:126)</p> <p>ATGGCCACAGGCAGAGAACATCTCTGCTGCTGGCCTTCGGACTGCTGTGCTGCCTTGCT          GCAAGAGGGTCCCGC</p>	<p><b>Furin cleavage site (SEQ ID NO:124)</b></p> <p><b>GSG Linker (SEQ ID NO:125)</b></p> <p><b>HGH/SS/2 (SEQ ID NO:127)</b></p>
<p>[[TRA_VDJ Sequence, specific for the NeoTCR]]</p> <p>ATATTAGAAACCCCGATCCTGCTGTGTATCAGCTGGCGGACAGCAAGAGCAGCGACAAGAGC          GTGTGTTGTTC</p>	<p><b>Insertion site for neoTCR          TRA_VDJ sequence</b></p> <p><b>TCR-alpha/constant (SEQ ID          NO:129)</b></p>

**FIGURE 9 Cont.**

**TRAC CDS/right HR arm  
(SEQ ID NO:130)**

ACCGATTTGATTCTCAACAATAATGTGTCACAAAGTAAGGATTCTGATGTGTATATCACAGAC  
AAACTGTCTAGACATGAGGTCTATGGACTTCAAGAGCAACAGTGTGTGGCCTGGAGCA  
ACAAATCTGACTTTGCATGTGCAAAACGCCCTTCAACAACAGCATTATCCAGAAGACACCTTCT  
TCCCCAGCCCAGG

**Right HR arm (SEQ ID  
NO:131)**

TAAGGGCAGCTTTGGTGCCCTTCGCAGGGCTGTTCCCTTGCTTCAGGAATGGCCAGGTTCTGCC  
CAGAGCTCTGTCATGATGTCTAAACTCCTCTGATTGGTGGTCTCGGCCTTATCCATTGCC  
ACCAAAACCCCTTTTTACTAAGAAACAGTGAGCCTTGTTCTGGCAGTCCAGAGAATGACAC  
GGGAAAAAAGCAGATGAAGAGAAAGGTGGCAGGAGGGCACCGTGGCCAGCCTCAGTCT  
CTCCAAGTCTGCTGCCCTGCTCAGACTGTTTGCCCTTACTGCTCTTCTAGG  
CCTCATTCTAAGCCCCCTTCTCCAAAGTTGCCCTCTCCTTATTTCTCCCTGTGCCAAAATCTTT  
CCCAGCTCACTAAGTCAGTCTCACGCAGTCACTCATTAAACCCACCACCAATCACTGATTGTGCCGG  
CACATGAATGCACCAGGTGTTGAAGTGGAGGAATTAATAAGTCAGATGAGGGGTGTGCCCA  
GAGGAAGCACCATTCTAGTTGGGGGAGCCCCATCTGTGAGTGGGAAAAGTCCAAATAACTT  
CAGATTGGAATGTTTTAACTCAGGGTTGAGAAAACAGTACCTTCAGGACAAAAGTCAG  
GGAAGGGCTCTGAAGAAATGCTACTTGAAGATACCAGCCCTACCAAGGGCAGGGAGAG  
GACCCATAGAGGCCCTGGGACAGGAGCTCAATGAGAAAAGGAGAAGAGCAGCGGCATGA  
GTTGAATGAAGGAGGCGGGGTCACAGGGCCTTCTAGGCCATGAGAGGGGTAGACAG

**(SEQ ID NO:132)**

GCTAGC

**FIGURE 9 Cont.**

**pBR322\_origin (SEQ ID NO:133)**

CGCGTTGCGGTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAATAATCGACGCT  
CAAGTCAGAGGTGGCAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAG  
CTCCCTGCGCTCTCTGTTCCGACCTGCCGTTACCGGATACCTGTCCGCCCTTCTCCCT  
TCGGAAAGCGTGGCCTTCTCATAGCTCACGCTGAGGTATCTCAGTTCGGGTAGGTCGTT  
CGTCCAAGCTGGCTGTGACGAACCCCTTCCAGCCGACCCGCTGCGCCTTATCCGG  
TAACTATCGTCTTAGTCCAACCCGGTAGACACGACTTATCGCCACTGGCAGCAGCCACTG  
GTAACAGGATTAGCAGAGCGAGGTATGTAGCGGTGTACAGAGTCTTGAAGTGGTGCC  
TAACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTT  
CGGAAAAGAGTTGGTAGCTTTGATCCGGCAAAACCAACCAGCTGGTAGCGGTTT  
TTTGTGGCAAGCAGCAGATTACGGCGAGAAAAGGATCTCAAGAAAGATCCTTTGATCT

**Kanamycin resistance  
(KanR2) (SEQ ID NO:134)**

TTAGAAAACATCGAGCATCAAATGAAACTGCAATTTATTCATATCAGGATTATCAATACCA  
TATTTTTGAAAAGCCGTTTCTGTAATGAAGGAGAAAACCTCACCAGGCGAGTCCATAGGAT  
GGCAAGATCCTGGTATCGGTCTGGATTCCGACTGTCACATCAATCAACACCTATTAATTC  
CCCTCGTCAAAAATAAGTTATCAAGTGAGAAATCACCATGAGTGACGACTGAATCCGGTGA  
GAATGGCAAAAAGTTATGCATTTCTTCCAGACTTGTCAACAGGCCAGCCATTACGCTCGTC  
ATCAAAATCACTCGCATCAACCAACCGTTATTTCATTCGTGATTGCGCCTGAGCCAGACGAAA  
TACGCGATCGCTGTTAAAAGGACAATTACAACAGGAATCGAATGCAACCCGGCGCAGGAAC  
ACTGCCAGCGCATCAACAATATTTTCCCTGAATCAGGATATCTTCTAATACCTGGAAATGCTG  
TTTTTCCGGGATCGCAGTGGTAGTAACCATGCATCATCAGGAGTACGGATAAATGCTTG  
ATGGTCGGAAGAGGCATAAATCCGTCAGCCAGTTTAGTCTGACCATCTCATCTGTAACATCA  
TTGGCAACGCTACCTTTGCCATGTTTCAGAAACAACCTCTGGCGCATCGGCTTCCCATACAAG  
CGATAGATTGTCGCACCTGATTGCCCGACATTATCGCGAGCCCAATTTATACCCATATAAATCAG  
CATCCATGTTGGAATTTAATCGCGCCCTCGACGTTTCCCGTTGAATATGGCTCAT

**Kanamycin promoter (SEQ ID NO:135)**

AACACCCCTGTACTGTTTATGTAAGCAGACAGTTTTATTGTTTCATGATGATATATTTTTATC  
TTGTGCAATGTAACATCAGAGATTTTGAGACAC






FIGURE 10



**FIGURE 11A**



**FIGURE 11B**





FIGURE 11C



**FIGURE 11D**




FIGURE 12




FIGURE 13A




FIGURE 13B






FIGURE 14





FIGURE 15B





FIGURE 15A




FIGURE 16

50/51




FIGURE 17B

FIGURE 17A




FIGURE 18

## SEQUENCE LISTING

<110> PACT PHARMA, INC.

<120> COMPOSITIONS AND METHODS FOR THE TREATMENT OF CANCER USING A CD8  
ENGINEERED T CELL THERAPY

<130> 087520.0145

<140>

<141>

<150> 62/841,748

<151> 2019-05-01

<150> 62/841,753

<151> 2019-05-01

<160> 148

<170> PatentIn version 3.5

<210> 1

<211> 926

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Left Homology Arm\_NeoTCR Product

<400> 1

```

acattaaaa cacaaatcc tacggaaata ctgaagaatg agtctcagca ctaaggaaaa      60
gcctccagca gctcctgctt tctgagggtg aaggatagac gctgtggctc tgcatgactc     120
actagcactc tatcacggcc atattctggc agggctcagtg gctccaacta acatttgttt     180
ggtactttac agtttattaa atagatgttt atatggagaa gctctcattt ctttctcaga     240
agagcctggc taggaagggt gatgaggcac catattcatt ttgcaggtga aattcctgag     300
atgtaaggag ctgctgtgac ttgctcaagg ctttatatcg agtaaacggt agtgctgggg     360
cttagacgca ggtgttctga tttatagttc aaaacctcta tcaatgagag agcaatctcc     420
tggtaatgtg atagatttcc caacttaatg ccaacatacc ataaacctcc cattctgcta     480
atgcccagcc taagttgggg agaccactcc agattccaag atgtacagtt tgctttgctg     540
ggcctttttc ccatgcctgc ctttactctg ccagagttat attgctgggg ttttgaagaa     600
gatcctatta aataaaagaa taagcagtat tattaagtag ccctgcattt caggtttcct     660
tgagtggcag gccaggcctg gccgtgaacg ttcactgaaa tcatggcctc ttggccaaga     720
ttgatagctt gtgcctgtcc ctgagtccca gtccatcacg agcagctggt ttctaagatg     780
ctatttcccg tataaagcat gagaccgtga cttgccagcc ccacagagcc ccgcccttgt     840
ccatcactgg catctggact ccagcctggg ttggggcaaa gagggaaatg agatcatgtc     900
ctaaccctga tcctcttgtc ccacag                                           926

```

<210> 2

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
TRAC CDS I\_NeoTCR Product

<400> 2

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt 60

ctgtctgcct attc 74

<210> 3

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_NeoTCR Product

<400> 3

gaattcggct ccgga 15

<210> 4

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
P2A 1\_NeoTCR Product

<400> 4

gccactaact tctccctggt gaaacaggct ggcgatgttg aagaaaaccc cggctcct 57

<210> 5

<211> 78

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
HGH Signal Sequence and Furin sequence\_NeoTCR Product

<400> 5

atggccaccg gctctagaac aagcctgctg ctcgcttttg gcctgctctg cctcccatgg 60

ctccaagaag gatctgct 78

<210> 6

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Exemplary TRB\_VDJ (TCR097)\_NeoTCR Product

<400> 6  
aatgctgggtg tcaactcagac cccaaaattc cgcatcctga agataggaca gagcatgaca 60  
ctgcagtgtgta cccaggatat gaaccataac tacatgtact ggtatcgaca agaccaggc 120  
atggggctga agctgattta ttattcagtt ggtgctggta tcaactgataa aggagaagtc 180  
ccgaatggct acaacgtctc cagatcaacc acagaggatt tcccgtcag gctggagttg 240  
gctgctccct cccagacatc tgtgtacttc tgtgccagct ccctacaggt tccctacaat 300  
gagcagttct tcgggccagg gacacggctc accgtgctag aggac 345

<210> 7  
<211> 534  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TRBC Constant Region\_NeoTCR Product

<400> 7  
ctgaaaaacg tgttcctcc aaaagtggcc gtgttcgagc cttctgaggc cgagatcagc 60  
cacacacaga aagccacact cgtgtgtctg gctaccggct tctaccccga tcacgtggaa 120  
ctgtcttggg gggtaacgg caaagagggtg cacagcggcg tcagcacaga tccccagcct 180  
ctgaaagaac agcccgtctt gaacgacagc cgctactgcc tgtctagcag actgagagtg 240  
tccgccacct tctggcagaa cccagaaac cacttcagat gccaggcca gttctacggc 300  
ctgagcgaga acgatgagtg gaccaggac agagccaagc ctgtgacaca gatcgtgtct 360  
gccgaagcct ggggcagagc cgattgtggc tttaccagcg agtcatacca gcagggcgtg 420  
ctgtctgcca ccatcctgta tgagatcctg ctcggcaagg ccacactgta cgctgtgctg 480  
gtgtctgctc tgggtgctgat ggctatggtc tcccgggagc gcatccccga ggcc 534

<210> 8  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Furin cleavage site\_NeoTCR Product

<400> 8  
cgggccaagc gg 12

<210> 9  
<211> 9  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_NeoTCR Product

<400> 9  
ggcagcggc

9

<210> 10  
<211> 57  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
P2A\_NeoTCR Product

<400> 10  
gccaccaact tcagcctgct gaagcaggcc ggcgacgtgg aggagaacct cggccct 57

<210> 11  
<211> 78  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
HGH\_SS\_NeoTCR Product

<400> 11  
atggccacag gcagcagaac atctctgctg ctggccttcg gactgctgtg tctgccttgg 60

ctgcaagagg gttccgcc 78

<210> 12  
<211> 337  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Exemplary TRA-VDJ (TCR097)\_NeoTCR Product

<400> 12  
gctcagacag tcaactcagtc tcaaccagag atgtctgtgc aggaggcaga gaccgtgacc 60

ctgagctgca catatgacac cagtgagagt gattattatt tattctggta caagcagcct 120

cccagcaggc agatgattct cgttattcgc caagaagcct ataagcaaca gaatgcaaca 180

gagaatcggt tctctgtgaa cttccagaaa gcagccaaat cttcagtct caagatctca 240

gactcacagc tgggggatgc cgcatgtat ttctgtgctt ttgggaactt caacaaattt 300

tactttggat ctgggaccaa actcaatgta aaaccaa 337

<210> 13  
<211> 74  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TCR-alpha/constant\_NeoTCR Product

&lt;400&gt; 13

atattcagaa ccccgatcct gctgtgtatc agctg'gcgca cagcaagagc agcgacaaga 60

gcgtgtgttt gttc 74

&lt;210&gt; 14

&lt;211&gt; 200

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
TRAC CDS/Right HR arm\_NeoTCR Product

&lt;400&gt; 14

accgattttg attctcaaac aaatgtgtca caaagtaagg attctgatgt gtatatcaca 60

gacaaaactg tgctagacat gaggtctatg gacttcaaga gcaacagtgc tgtggcctgg 120

agcaacaaat ctgactttgc atgtgcaaac gccttcaaca acagcattat tccagaagac 180

accttcttcc ccagcccagg 200

&lt;210&gt; 15

&lt;211&gt; 800

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Right HR arm\_NeoTCR Product

&lt;400&gt; 15

taagggcagc tttggtgcct tcgcaggctg tttccttgct tcaggaatgg ccaggttctg 60

cccagagctc tggatcaatga tgtctaaaac tcctctgatt ggtggtctcg gccttatcca 120

ttgccaccaa aaccctcttt ttactaagaa acagttagcc ttgttctggc agtccagaga 180

atgacacggg aaaaaagcag atgaagagaa ggtggcagga gagggcacgt ggcccagcct 240

cagtctctcc aactgagttc ctgcctgcct gcctttgctc agactgtttg ccccttactg 300

ctcttctagg cctcattcta agccccttct ccaagttgcc tctccttatt tctccctgtc 360

tgccaaaaaa tctttcccag ctactaagt cagtctcag cagtactca ttaaccacc 420

aatcactgat tgtgccggca catgaatgca ccaggtgttg aagtggagga attaaaaagt 480

cagatgaggg gtgtgcccag aggaagcacc attctagttg ggggagccca tctgtcagct 540

gggaaaagt caaataactt cagattggaa tgtgttttaa ctgagggttg agaaaacagc 600

taccttcagg acaaaagtca gggaagggtc ctctgaagaa atgctacttg aagataccag 660

ccctaccaag ggcagggaga ggaccctata gaggcctggg acaggagctc aatgagaaag 720

gagaagagca gcaggcatga gttgaatgaa ggaggcaggg ccgggtcaca gggccttcta 780

ggccatgaga gggtagacag 800

&lt;210&gt; 16

<211> 6  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Artificial Sequence\_NeoTCR Product

<400> 16  
 gctagc 6

<210> 17  
 <211> 620  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 pBR322\_origin\_NeoTCR Product

<400> 17  
 cgcggttgctg gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg 60  
 ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt tccccctgg 120  
 aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc tgtccgcctt 180  
 tctcccttcg ggaagcgtgg cgcttttctca tagctcacgc tntaggtatc tcagttcggg 240  
 gtaggtcggt cgctccaagc tgggctgtgt gcacgaacc cccgttcagc ccgaccgctg 300  
 cgcttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact 360  
 ggtagcagcc actggtaaca ggattagcag agcgaggat gtaggcgggtg ctacagagtt 420  
 cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttgta tctgcgctct 480  
 gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca acaaaaccac 540  
 cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgagaa aaaaggatc 600  
 tcaagaagat cctttgatct 620

<210> 18  
 <211> 810  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Kanamycin resistance gene (KanR2) \_NeoTCR Product

<400> 18  
 ttgaaaaaac tcatcgagca tcaaatgaaa ctgcaattta tcatatcag gattatcaat 60  
 accatatttt tgaaaaagcc gtttctgtaa tgaaggagaa aactcaccga ggtagttcca 120  
 taggatggca agatcctggt atcggctctgc gattccgact cgtccaacat caatacaacc 180  
 tattaatttc ccctcgtcaa aaataagggt atcaagttag aatcaccat gtagtgacgac 240  
 tgaatccggt gagaatggca aaagtttatg catttcttc cagacttggt caacaggcca 300

gccattacgc tcgtcatcaa aatcactcgc atcaacaaa ccgttattca ttcgtgattg 360  
cgctgagcc agacgaaata cgcgatcgcgt gttaaaagga caattacaaa caggaatcga 420  
atgcaaccgg cgcaggaaca ctgccagcgc atcaacaata ttttcacctg aatcagata 480  
ttcttctaata acctggaatg ctgtttttcc ggggatcgcga gtggtgagta accatgcatc 540  
atcaggagta cggataaaat gcttgatggt cggaagaggc ataaattccg tcagccagtt 600  
tagtctgacc atctcatctg taacatcatt ggcaacgcta cctttgccat gtttcagaaa 660  
caactctggc gcatcgggct tccatacaa gcgatagatt gtcgcacctg attgcccgcac 720  
attatcgcga gcccatttat acccatataa atcagcatcc atggttgaat ttaatcgcgg 780  
cctcgacgtt tcccgttgaa tatggctcat 810

<210> 19

<211> 99

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Kanamycin promoter \_NeoTCR Product

<400> 19

aacaccccctt gtattactgt ttatgtaagc agacagtttt attgttcatg atgatatatt 60

tttatcttgt gcaatgtaac atcagagatt ttgagacac 99

<210> 20

<211> 5137

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Complete NeoTCR Product construct with TCR97 insertion\_NeoTCR  
Product

<400> 20

ggtaccacat taaaaacaca aaatcctacg gaaatactga agaatgagtc tcagcactaa 60

ggaaaagcct ccagcagctc ctgctttctg aggggtgaagg atagacgctg tggctctgca 120

tgactcacta gactctatc acggccatat tctggcaggg tcagtggctc caactaacat 180

ttgtttggta ctttacagtt tattaatag atgtttatat ggagaagctc tcatttcttt 240

ctcagaagag cctggctagg aagggtgatg aggcaccata ttcattttgc aggtgaaatt 300

cctgagatgt aaggagctgc tgtgacttgc tcaaggcctt atatcgagta aacggtagtg 360

ctggggctta gacgcaggtg ttctgattta tagttcaaaa cctctatcaa tgagagagca 420

atctcctggg aatgtgatag atttcccaac ttaatgcca cataccataa acctccatt 480

ctgctaatagc ccagcctaag ttggggagac cactccagat tccaagatgt acagtttgct 540

ttgctgggcc tttttccat gcctgccttt actctgccag agttatattg ctggggtttt 600

gaagaagatc ctattaaata aaagaataag cagtattatt aagtagccct gcatttcagg 660  
tttccttgag tggcaggcca ggctggccg tgaacgtca ctgaaatcat ggctcttg 720  
ccaagattga tagcttgtgc ctgtccctga gtcccagtc atcacgagca gctggtttct 780  
aagatgctat ttcccgtata aagcatgaga ccgtgacttg ccagccccac agagccccgc 840  
cctgtccat cactggcatc tggactccag cctgggttg ggcaaagagg gaaatgagat 900  
catgtcctaa ccctgatcct cttgtcccac agatatccag aaccctgacc ctgccgtgta 960  
ccagctgaga gactctaaat ccagtgacaa gtctgtctgc ctattcgaat tcggctccgg 1020  
agccactaac ttctccctgt tgaacagggc tggcgatgtt gaagaaaacc ccggtcctat 1080  
ggccaccggc tctagaacaa gcctgctgct cgcttttggc ctgctctgcc tccatggct 1140  
ccaagaagga tctgctaag ctggtgtcac tcagaccca aaattccgca tcctgaagat 1200  
aggacagagc atgacactgc agtgtacca ggatatgaac cataactaca tgtactggta 1260  
tcgacaagac ccaggcatgg ggctgaagct gatttattat tcagttggtg ctggtatcac 1320  
tgataaagga gaagtccga atggctacaa cgtctccaga tcaaccacag aggatttccc 1380  
gctcaggctg gagttggctg ctccctcca gacatctgtg tacttctgtg ccagctcct 1440  
acaggttccc tacaatgagc agttcttcgg gccagggaca cggctcaccg tgctagagga 1500  
cctgaaaaac gtgttcctc caaaagtggc cgtgttcgag ctttctgagg ccgagatcag 1560  
ccacacacag aaagccacac tcgtgtgtct ggctaccggc ttctaccccg atcacgtgga 1620  
actgtcttgg tgggtcaacg gcaaagaggt gcacagcggc gtcagcacag atccccagcc 1680  
tctgaaagaa cagcccgctc tgaacgacag ccgctactgc ctgtctagca gactgagagt 1740  
gtccgccacc ttctggcaga accccagaaa ccacttcaga tgccaggctc agttctacgg 1800  
cctgagcgag aacgatgagt ggaccagga cagagccaag cctgtgacac agatcgtgtc 1860  
tgccgaagcc tggggcagag ccgattgtgg ctttaccagc gagtcatacc agcagggcgt 1920  
gctgtctgcc accatcctgt atgagatcct gctcggcaag gccacactgt acgctgtgct 1980  
ggtgtctgct ctggtgctga tggctatggt ctcccgggag cgcatccccg agggccgggc 2040  
caagcggggc agcggcgcca ccaacttcag cctgctgaag caggccggcg acgtggagga 2100  
gaaccccgcc cctatggcca caggcagcag aacatctctg ctgctggcct tcggactgct 2160  
gtgtctgcct tggctgcaag agggttccgc cgctcagaca gtactcagt ctcaaccaga 2220  
gatgtctgtg caggaggcag agaccgtgac cctgagctgc acatatgaca ccagtgagag 2280  
tgattattat ttattctggt acaagcagcc tcccagcagg cagatgattc tcgttattcg 2340  
ccaagaagct tataagcaac agaatgcaac agagaatcgt ttctctgtga acttccagaa 2400  
agcagccaaa tccttcagtc tcaagatctc agactcacag ctgggggatg ccgcgatgta 2460  
tttctgtgct tttgggaact tcaacaaatt ttactttgga tctgggacca aactcaatgt 2520  
aaaaccaa attcagaacc ccgatcctgc tgtgtatcag ctgcgcgaca gcaagagcag 2580

cgacaagagc gtgtgtttgt tcaccgattt tgattctcaa acaaatgtgt cacaaagtaa 2640  
ggattctgat gtgtatatca cagacaaaac tgtgctagac atgaggctta tggacttcaa 2700  
gagcaacagt gctgtggcct ggagcaacaa atctgacttt gcatgtgcaa acgccttcaa 2760  
caacagcatt attccagaag acaccttctt cccagccca ggtaaggga gctttgggtgc 2820  
cttcgaggc tgtttccttg cttcaggaat ggccaggctt tgcccagagc tctggtcaat 2880  
gatgtctaaa actcctctga ttggtggctt cggccttatt cattgccacc aaaaccctct 2940  
ttttactaag aaacagttag ccttgctctg gcagtccaga gaatgacacg ggaaaaaagc 3000  
agatgaagag aagggtggcag gagagggcac gtggcccagc ctgagtctct ccaactgagt 3060  
tcctgcctgc ctgcctttgc tcagactggt tgccccttac tgctcttcta ggcctcattc 3120  
taagcccctt ctccaagttg cctctcctta tttctccctg tctgcaaaa aatctttccc 3180  
agctcactaa gtcagtctca cgcagtcaact cattaacca ccaatcactg attgtgccgg 3240  
cacatgaatg caccaggtgt tgaagtggag gaattaaana gtcagatgag ggggtgtgcc 3300  
agaggaagca ccattctagt tgggggagcc catctgtcag ctgggaaaag tccaataac 3360  
ttcagattgg aatgtgtttt aactcagggt tgagaaaaca gctaccttca ggacaaaagt 3420  
caggggaagg ctctctgaag aatgctact tgaagatacc agccctacca agggcaggga 3480  
gaggacccta tagaggcctg ggacaggagc tcaatgagaa aggagaagag cagcaggcat 3540  
gagttgaatg aaggaggcag ggccgggtca cagggccttc taggcatga gaggtagac 3600  
aggctagccg cgttgctggc gtttttccat aggctccgcc cccctgacga gcatcaciaa 3660  
aatcgacgct caagtcaag gtggcgaaac ccgacaggac tataaagata ccaggcgttt 3720  
ccccctggaa gctccctcgt gcgctctcct gttccgacc tgccgcttac cggatacctg 3780  
tccgcctttc tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc 3840  
agttcggtgt aggtcgttcg ctccaagctg ggctgtgtgc acgaacccc cgttcagccc 3900  
gaccgctgcg ctttatccgg taactatcgt cttgagtcca acccgtaag acacgactta 3960  
tcgccactgg cagcagccac tggtaacagg attagcagag cgaggtatgt aggcggtgct 4020  
acagagttct tgaagtgggt gcctaactac ggctacacta gaagaacagt atttggatc 4080  
tgcgctctgc tgaagccagt taccttcgga aaaagagttg gtagctcttg atccggcaaa 4140  
caaaccaccg ctggtagcgg tggttttttt gtttgcaagc agcagattac gcgcagaaaa 4200  
aaaggatctc aagaagatcc tttgatcttt agaaaaactc atcgagcatc aatgaaact 4260  
gcaatttatt catatcagga ttatcaatac catatttttg aaaaagccgt ttctgtaatg 4320  
aaggagaaaa ctcaccgagg cagttccata ggatggcaag atcctggtat cggctctcga 4380  
ttccgactcg tccaacatca atacaaccta ttaatttccc ctctcaaaa ataaggttat 4440  
caagtgagaa atcaccatga gtgacgactg aatccggtga gaatggcaaa agtttatgca 4500

tttctttcca gacttgttca acaggccagc cattacgctc gtcatcaaaa tctctgcat 4560  
caaccaaac gttattcatt cgtgattgcg cctgagccag acgaaatacg cgatcgctgt 4620  
taaaaggaca attacaaaca ggaatcgaat gcaaccggcg caggaacact gccagcgcac 4680  
caacaatatt ttcacctgaa tcaggatatt cttctaatac ctggaatgct gtttttccgg 4740  
ggatcgcagt ggtgagtaac catgcatcat caggagtacg gataaaatgc ttgatggtcg 4800  
gaagaggcat aaattccgtc agccagtta gtctgaccat ctcatctgta acatcattgg 4860  
caacgctacc tttgcatgt ttcagaaaca actctggcgc atcgggcttc ccatacaagc 4920  
gatagattgt cgcacctgat tgcccacat tatcgcgagc ccatttatac ccatataaat 4980  
cagcatccat gttggaattt aatcgcggcc tcgacgtttc ccgttgaata tggctcataa 5040  
cacccttgt attactgttt atgtaagcag acagttttat tgttcatgat gatataat 5100  
tatcttgtgc aatgtaacat cagagatttt gagacac 5137

<210> 21  
<211> 5140  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Complete NeoTCR Product construct with TCR89 insertion\_NeoTCR  
Product

<400> 21  
ggtaccacat taaaaacaca aaatcctacg gaaatactga agaatgagtc tcagcactaa 60  
ggaaaagcct ccagcagctc ctgctttctg aggggtgaagg atagacgctg tggctctgca 120  
tgactcacta gcactctatc acggccatat tctggcaggg tcagtggctc caactaacat 180  
ttgtttgta ctttacagtt tattaatag atgtttatat ggagaagctc tcatttcttt 240  
ctcagaagag cctggctagg aaggtggatg aggcaccata ttcattttgc aggtgaaatt 300  
cctgagatgt aaggagctgc tgtgacttgc tcaaggcctt atatcgagta aacggtagtg 360  
ctggggctta gacgcaggtg ttctgattta tagttcaaaa cctctatcaa tgagagagca 420  
atctcctggt aatgtgatag atttccaac ttaatgcaa cataccataa acctcccatt 480  
ctgctaattgc ccagcctaag ttggggagac cactccagat tccaagatgt acagtttgct 540  
ttgctgggcc tttttcccat gcctgccttt actctgccag agttatattg ctggggtttt 600  
gaagaagatc ctattaaata aaagaataag cagtattatt aagtagcct gcatttcagg 660  
ttccttgag tggcaggcca ggcctggccg tgaacgttca ctgaaatcat ggcctcttgg 720  
ccaagattga tagcttgtgc ctgtccctga gtcccagtcc atcacgagca gctggtttct 780  
aagatgctat ttcccgtata aagcatgaga ccgtgacttg ccagccccac agagccccgc 840  
cctgtccat cactggcatc tggactccag cctgggttgg ggcaaagagg gaaatgagat 900  
catgtcctaa cctgatcct cttgtccac agatatccag aaccctgacc ctgccgtgta 960

ccagctgaga gactctaaat ccagtgacaa gtctgtctgc ctattcgaat tcggctccgg 1020  
agccactaac ttctccctgt tgaacaggc tggcgatgtt gaagaaaacc ccggtcctat 1080  
ggccaccggc tctagaacaa gcctgctgct cgcttttggc ctgctctgcc tcccatggct 1140  
ccaagaagga tctgctgatg ctggaatcac ccagagccca agatacaaga tcacagagac 1200  
aggaaggcag gtgaccttga tgtgtcacca gacttggagc cacagctata tgttctggta 1260  
tcgacaagac ctgggacatg ggctgaggct gatctattac tcagcagctg ctgatattac 1320  
agataaagga gaagtctccg atggctatgt tgtctccaga tccaagacag agaatttccc 1380  
cctcactctg gagtcagcta cccgctccca gacatctgtg tatttctgcg ccagcagtga 1440  
ggacagttac gagcagtact tcgggccggg caccaggctc acggtcacag aggacctgaa 1500  
aaacgtgttc cctccaaaag tggccgtgtt cgagccttct gaggccgaga tcagccacac 1560  
acagaaagcc acactcgtgt gtctggctac cggcttctac cccgatcacg tggaaactgtc 1620  
ttggtggggtc aacggcaaag aggtgcacag cggcgtcagc acagatcccc agcctctgaa 1680  
agaacagccc gctctgaacg acagccgcta ctgcctgtct agcagactga gagtgtccgc 1740  
caccttctgg cagaacccca gaaaccactt cagatgccag gtccagttct acggcctgag 1800  
cgagaacgat gaggggacc aggacagagc caagcctgtg acacagatcg tgtctgccga 1860  
agcctggggc agagccgatt gtggcctttac cagcagatca taccagcagg gcgtgctgtc 1920  
tgccaccatc ctgtatgaga tcctgctcgg caaggccaca ctgtacgctg tgctggtgtc 1980  
tgctctgggtg ctgatggcta tggctctccg ggagcgcac cccgaggccc gggccaagcg 2040  
gggcagcggc gccaccaact tcagcctgct gaagcaggcc ggcgacgtgg aggagaacct 2100  
cggccctatg gccacaggca gcagaacatc tctgctgctg gccttcggac tgctgtgtct 2160  
gccttggctg caagagggtt ccgccccca gtcagtacc cagcctgaca tccacatcac 2220  
tgtctctgaa ggagcctcac tggagttgag atgtaactat tcctatgggg caacacctta 2280  
tctcttctgg tatgtccagt cccccggcca aggcctccag ctgctcctga agtacttttc 2340  
aggagacact ctggttcaag gcattaaagg ctttgaggct gaatttaaga ggagtcaatc 2400  
ttccttcaat ctgaggaaac cctctgtgca ttggagtgat gctgctgagt acttctgtgc 2460  
tgtgggtgaa ttggacacag gctttcagaa acttgtatct ggaactggca cccgacttct 2520  
ggtcagtcca aatattcaga accccgatcc tgctgtgtat cagctgcgcg acagcaagag 2580  
cagcgacaag agcgtgtgtt ttttcaccga ttttgattct caaacaatg tgcacaaag 2640  
taaggattct gatgtgtata tcacagacaa aactgtgcta gacatgaggt ctatggactt 2700  
caagagcaac agtgctgtgg cctggagcaa caaatctgac tttgcatgtg caaacgcctt 2760  
caacaacagc attattccag aagacacctt cttccccagc ccaggttaagg gcagctttgg 2820  
tgccctcgca ggctgtttcc ttgcttcagg aatggccagg ttctgcccag agctctggtc 2880

aatgatgtct aaaactcctc tgattggtgg tctcggcctt atccattgcc accaaaaccc 2940  
tctttttact aagaacagtg gagccttgtt ctggcagtcc agagaatgac acgggaaaaa 3000  
agcagatgaa gagaaggtgg caggagaggg cacgtggccc agcctcagtc tctccaactg 3060  
agttcctgcc tgctgcctt tgctcagact gtttgcccct tactgctctt ctaggcctca 3120  
ttctaagccc cttctcaag ttgcctctcc ttatttctcc ctgtctgcca aaaaatcttt 3180  
cccagctcac taagttagtc tcacgcagtc actcattaac ccaccaatca ctgattgtgc 3240  
cggcacatga atgcaccagg tgttgaagtg gaggaattaa aaagtcagat gaggggtgtg 3300  
cccagaggaa gcaccattct agttggggga gcccatctgt cagctgggaa aagtccaaat 3360  
aacttcagat tggaatgtgt ttttaactcag ggttgagaaa acagctacct tcaggacaaa 3420  
agtcagggaa gggctctctg aagaaatgct acttgaagat accagcccta ccaaggcag 3480  
ggagaggacc ctatagaggc ctgggacagg agctcaatga gaaaggagaa gagcagcagg 3540  
catgagttga atgaaggagg cagggccggg tcacagggcc ttctaggcca tgagagggta 3600  
gacaggctag ccgcgttgct ggcgttttcc cataggctcc gccccctga cgagcatcac 3660  
aaaaatcgac gctcaagtca gaggtggcga aaccgcagc gactataaag ataccaggcg 3720  
tttcccctg gaagctccct cgtgcgctct cctgttccga ccctgccgct taccggatac 3780  
ctgtccgctt ttctccctc gggaaagctg gcgctttctc atagctcacg ctgtaggat 3840  
ctcagttcgg ttaggtcgt tcgctccaag ctgggctgtg tgacgaacc cccggttcag 3900  
cccgaccgct gcgccttatc cggttaactat cgtcttgagt ccaaccggt aagacacgac 3960  
ttatcgccac tggcagcagc cactggtaac aggattagca gagcgaggta ttaggcggt 4020  
gctacagagt tcttgaagtg gtggcctaac tacggctaca ctagaagaac agtatttgg 4080  
atctgcgctc tgctgaagcc agttacctc ggaaaaagag ttgtagctc ttgatccggc 4140  
aaacaaacca ccgctggtag cggtggtttt tttgtttgca agcagcagat tacgcgcaga 4200  
aaaaaaggat ctcaagaaga tcctttgatc tttagaaaaa ctcatcgagc atcaaatgaa 4260  
actgcaatth atcatatca ggattatcaa taccatattt ttgaaaaagc cgtttctgta 4320  
atgaaggaga aaactcaccg aggcagttcc ataggatggc aagatcctgg taccggtctg 4380  
cgattccgac tcgtccaaca tcaatacaac ctattaattt cccctcgtca aaaataaggt 4440  
tatcaagtga gaaatcacca tgagtacga ctgaatccgg tgagaatggc aaaagtttat 4500  
gcatttcttt ccagacttgt tcaacaggcc agccattacg ctctcatca aatcactcg 4560  
catcaaccaa accgttattc attcgtgatt gcgcctgagc cagacgaaat acgcgatcgc 4620  
tgtaaaaagg acaattacaa acaggaatcg aatgcaaccg gcgcaggaac actgccagcg 4680  
catcaacaat attttcacct gaatcaggat attcttctaa tacctggaat gctgtttttc 4740  
cggggatcgc agtggtgagt aacctgcat catcaggagt acggataaaa tgcttgatgg 4800  
tcggaagagg cataaattcc gtcagccagt ttagtctgac catctcatct gtaacatcat 4860

tggcaacgct acctttgcca tgtttcagaa acaactctgg cgcatcgggc ttccataca 4920  
agcgatagat tgtcgcacct gattgcccgga cattatcgcg agcccattta tacccatata 4980  
aatcagcatc catgttggaa tttaatcgcg gcctcgacgt ttcccgttga atatggctca 5040  
taacaccct tgtattactg tttatgtaag cagacagttt tattgttcat gatgatatat 5100  
ttttatcttg tgcaatgtaa catcagagat tttgagacac 5140

<210> 22  
<211> 926  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Left HR arm\_CD8 Product 1

<400> 22  
acattaaaa cacaaaatcc tacggaaata ctgaagaatg agtctcagca ctaaggaaaa 60  
gcctccagca gtcctgctt tctgagggtg aaggatagac gctgtggctc tgcattgactc 120  
actagcactc tatcacggcc atattctggc agggctcagtg gctccaacta acatttgttt 180  
ggtactttac agtttattaa atagatgttt atatggagaa gctctcattt ctttctcaga 240  
agagcctggc taggaagggtg gatgaggcac catattcatt ttgcagggtga aattcctgag 300  
atgtaaggag ctgctgtgac ttgctcaagg ccttatatcg agtaaacggt agtctggtggg 360  
cttagacgca ggtgttctga tttatagttc aaaacctcta tcaatgagag agcaatctcc 420  
tggtaatgtg atagatttcc caacttaatg ccaacatacc ataaacctcc cattctgcta 480  
atgccagcc taagttgggg agaccactcc agattccaag atgtacagtt tgctttgctg 540  
ggcctttttc ccatgcctgc ctttactctg ccagagttat attgctgggg tttgaagaa 600  
gatcctatta aataaaagaa taagcagtat tattaagtag ccctgcattt caggtttcct 660  
tgagtggcag gccaggcctg gccgtgaacg ttactgaaa tcatggcctc ttggccaaga 720  
ttgatagctt gtgcctgtcc ctgagtcca gtccatcacg agcagctggt ttctaagatg 780  
ctatttcccg tataaagcat gagaccgtga cttgccagcc ccacagagcc ccgcccttgt 840  
ccatcactgg catctggact ccagcctggg ttggggcaaa gagggaaatg agatcatgtc 900  
ctaaccctga tcctcttgtc ccacag 926

<210> 23  
<211> 74  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TRAC CDS\_CD8 Product 1

<400> 23

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt 60

ctgtctgcct attc 74

<210> 24

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_CD8 Product 1

<400> 24

gaattcggct ccgga 15

<210> 25

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 1

<400> 25

gccactaact tcagcctggt gaagcaggcc ggcgacgttg aggaaaaccc cggctcct 57

<210> 26

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8A Signal peptide\_CD8 Product 1

<400> 26

atggccttac cagtgaccgc cttgctcctg ccgctggcct tgctgctcca cgccgccagg 60

ccg 63

<210> 27

<211> 483

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8A Extracellular domain\_CD8 Product 1

<400> 27

agccagttcc ggggtgtgcc gctggatcgg acctggaacc tgggcgagac agtggagctg 60

aagtgccagg tgctgctgtc caaccgacg tcgggctgct cgtggctctt ccagccgcgc 120

ggcgcgcccg ccagtccac cttcctccta tacctctccc aaaacaagcc caaggcggcc 180

gaggggctgg acaccagcg gttctcgggc aagaggttg gggacacctt cgtcctcacc 240

ctgagcgact tccgccgaga gaacgagggc tactatttct gctcggcct gagcaactcc 300

atcatgtact tcagccactt cgtgccggtc ttctgccag cgaagccac cacgaccca 360  
 gcgccgcgac caccaacacc ggcgccacc atcgcgtcgc agccctgtc cctgcgcca 420  
 gaggcgtgcc ggccagcggc gggggcgca gtgcacacga gggggctgga cttgcctgt 480  
 gat 483

<210> 28  
 <211> 63  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8A transmembrane domain\_CD8 Product 1

<400> 28  
 atctacatct gggcgccctt ggccgggact tgtgggtcc ttctctgtc actggttatc 60  
 acc 63

<210> 29  
 <211> 96  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8A Intracellular domain\_CD8 Product 1

<400> 29  
 ctttactgca accacaggaa ccgaagacgt gtttgcaaat gtccccggcc tgtggtcaaa 60  
 tcgggagaca agcccagcct ttcggcgaga tacgtc 96

<210> 30  
 <211> 12  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Furin cleavage site\_CD8 Product 1

<400> 30  
 agggctaaac gg 12

<210> 31  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 GSG Linker\_CD8 Product 1

<400> 31  
 gaattcggct ccgga 15

<210> 32  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 P2A\_CD8 Product 1

<400> 32  
 gccactaact tctccctggt gaaacaggct ggcgatgttg aagaaaaccc cggtcct 57

<210> 33  
 <211> 78  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 HGH/SS/2\_CD8 Product 1

<400> 33  
 atggccaccg gctctagaac aagcctgctg ctcgcttttg gcctgctctg cctcccatgg 60  
 ctccaagaag gatctgct 78

<210> 34  
 <211> 345  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Exemplary TRB\_VDJ (TCR097) \_CD8 Product 1

<400> 34  
 aatgctggtg tcaactcagac cccaaaattc cgcacacctga agataggaca gagcatgaca 60  
 ctgcagtgtgta cccaggatat gaaccataac tacatgtact ggtatcgaca agaccaggc 120  
 atggggctga agctgattta ttattcagtt ggtgctggta tcaactgataa aggagaagtc 180  
 ccgaatggct acaacgtctc cagatcaacc acagaggatt tcccgtcag gctggagttg 240  
 gctgctccct cccagacatc tgtgtacttc tgtgccagct ccctacaggt tcctacaat 300  
 gagcagttct tcgggccagg gacacggctc accgtgctag aggac 345

<210> 35  
 <211> 534  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 TCR-beta/constant\_CD8 Product 1

<400> 35  
 ctgaaaaacg tgttcctcc aaaagtggcc gtgttcgagc cttctgaggc cgagatcagc 60  
 cacacacaga aagccacact cgtgtgtctg gctaccggct tctaccccga tcactggaa 120

ctgtcttggg gggcaacgg caaagagggt cacagcggcg tcagcacaga tccccagcct 180  
ctgaaagaac agcccgctct gaacgacagc cgctactgcc tgtctagcag actgagagtg 240  
tccgccacct tctggcagaa cccagaaac cacttcagat gccaggcca gttctacggc 300  
ctgagcgaga acgatgagtg gacccaggac agagccaagc ctgtgacaca gatcgtgtct 360  
gccgaagcct ggggcagagc cgattgtggc tttaccagcg agtcatacca gcagggcgtg 420  
ctgtctgcca ccatcctgta tgagatcctg ctcggcaagg ccacactgta cgctgtgctg 480  
gtgtctgctc tgggtctgat ggctatggtc tcccgggagc gcatccccga ggcc 534

<210> 36

<211> 12

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Furin cleavage site\_CD8 Product 1

<400> 36

cgggccaagc gg

12

<210> 37

<211> 9

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
GSG linker\_CD8 Product 1

<400> 37

ggcagcggc

9

<210> 38

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 1

<400> 38

gccaccaact tcagcctgct gaagcaggcc ggcgacgtgg aggagaacct cggccct

57

<210> 39

<211> 78

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
HGH/SS\_CD8 Product 1

<400> 39

atggccacag gcagcagaac atctctgctg ctggccttcg gactgctgtg tctgccttgg 60

ctgcaagagg gttccgcc 78

<210> 40

<211> 337

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Exemplary TRA-VDJ (TCR097) \_CD8 Product 1

<400> 40

gctcagacag tcaactcagtc tcaaccagag atgtctgtgc aggaggcaga gaccgtgacc 60

ctgagctgca catatgacac cagtgagagt gattattatt tattctggta caagcagcct 120

cccagcaggc agatgattct cgttattcgc caagaagctt ataagcaaca gaatgcaaca 180

gagaatcggt tctctgtgaa cttccagaaa gcagccaaat cttcagctt caagatctca 240

gactcacagc tgggggatgc cgcatgtat ttctgtgctt ttgggaactt caacaaattt 300

tactttggat ctgggaccaa actcaatgta aaaccaa 337

<210> 41

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
TCR-alpha/constant\_CD8 Product 1

<400> 41

atattcagaa ccccgatcct gctgtgtatc agctgcgca cagcaagagc agcgacaaga 60

gcgtgtgttt gttc 74

<210> 42

<211> 200

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
TRAC CDS/right HR arm\_CD8 Product 1

<400> 42

accgattttg attctcaaac aaatgtgtca caaagtaagg attctgatgt gtatatcaca 60

gacaaaactg tgctagacat gaggtctatg gacttcaaga gcaacagtgc tgtggcctgg 120

agcaacaaat ctgactttgc atgtgcaaac gccttcaaca acagcattat tccagaagac 180

accttcttcc ccagcccagg 200

<210> 43

<211> 800

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Right HR arm\_CD8 Product 1

<400> 43

taagggcagc tttggtgcct tgcaggctg tttccttgct tcaggaatgg ccaggttctg	60
cccagagctc tggccaatga tgtctaaac tcctctgatt ggtggtctcg gccttatcca	120
ttgccaccaa aaccctcttt ttactaagaa acagtgagcc ttgttctggc agtccagaga	180
atgacacggg aaaaaagcag atgaagagaa ggtggcagga gagggcacgt ggcccagcct	240
cagtctctcc aactgagttc ctgcctgcct gcctttgctc agactgtttg ccccttactg	300
ctcttctagg cctcattcta agccccttct ccaagttgcc tctccttatt tctccctgtc	360
tgccaaaaaa tctttccag ctactaagt cagtctcacg cagtactca ttaaccacc	420
aatcactgat tgtgccggca catgaatgca ccagggtgtg aagtggagga attaaaaagt	480
cagatgaggg gtgtgcccag aggaagcacc attctagttg ggggagccca tctgtcagct	540
gggaaaagtc caaataactt cagattggaa tgtgttttaa ctcagggttg agaaaacagc	600
taccttcagg acaaaagtca gggaagggtc ctctgaagaa atgctacttg aagataccag	660
ccctaccaag ggcagggaga ggaccctata gaggcctggg acaggagctc aatgagaaaag	720
gagaagagca gcaggcatga gttgaatgaa ggaggcaggg ccgggtcaca gggccttcta	780
ggccatgaga gggtagacag	800

<210> 44

<211> 6

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Artificial Sequence\_CD8 Product 1

<400> 44

gctagc 6

<210> 45

<211> 620

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
pBR322\_origin\_CD8 Product 1

<400> 45

gcggttgctg gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg	60
ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt ttccccctgg	120
aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc tgtccgcctt	180

tctcccttcg ggaagcgtgg cgctttctca tagctcacgc tgtaggtatc tcagttcggg 240  
 gtaggtcggt cgctccaagc tgggctgtgt gcacgaacct cccgttcagc cggaccgctg 300  
 cgccttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact 360  
 ggcagcagcc actggtaaca ggattagcag agcgaggat gtaggcggtg ctacagagtt 420  
 cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttgta tctgcgctct 480  
 gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca acaaaaccac 540  
 cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgagaa aaaaaggatc 600  
 tcaagaagat cctttgatct 620

<210> 46  
 <211> 810  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Kanamycin resistance (KanR2) \_CD8 Product 1

<400> 46  
 ttgaaaaaac tcatcgagca tcaaatgaaa ctgcaattta ttcatatcag gattatcaat 60  
 accatatttt tgaaaaagcc gtttctgtaa tgaaggagaa aactcaccga ggcagttcca 120  
 taggatggca agatcctggt atcggctctgc gattccgact cgtccaacat caatacaacc 180  
 tattaatttc cctcgtcaa aaataagggt atcaagttag aaatcacatc gatgacgac 240  
 tgaatccggt gagaatggca aaagtttatg catttctttc cagacttggt caacaggcca 300  
 gccattacgc tcgtcatcaa aatcactcgc atcaaccaa ccgttattca ttcgtgattg 360  
 cgcctgagcc agacgaaata cgcgatcgc gttaaaagga caattacaaa caggaatcga 420  
 atgcaaccgg cgcaggaaca ctgccagcgc atcaacaata tttcacctg aatcaggata 480  
 ttcttcta atacctggaat ctgtttttcc ggggatcgc gtggtgagta accatgcatc 540  
 atcaggagta cggataaaat gcttgatggt cggaagaggc ataaattccg tcagccagtt 600  
 tagtctgacc atctcatctg taacatcatt ggcaacgcta cctttgcat gtttcagaaa 660  
 caactctggc gcatcgggct tccatacaa gcgatagatt gtcgcacctg attgcccagc 720  
 attatcgca gccatttat accatataa atcagcatcc atgttggaa ttaatcggg 780  
 cctcgacgtt tccggttga tatggctcat 810

<210> 47  
 <211> 99  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Kanamycin promoter\_CD8 Product 1

&lt;400&gt; 47

aacacccctt gtattactgt ttatgtaagc agacagtttt attgttcatg atgatatatt 60

tttatcttgt gcaatgtaac atcagagatt ttgagacac 99

&lt;210&gt; 48

&lt;211&gt; 5926

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
 Complete CD8 Product 1 construct with TCR97 insertion\_CD8  
 Product 1

&lt;400&gt; 48

ggtaccacat taaaaacaca aaatcctacg gaaatactga agaatgagtc tcagcactaa 60

ggaaaagcct ccagcagctc ctgctttctg aggggtgaagg atagacgctg tggctctgca 120

tgactcacta gcaactctatc acggccatat tctggcaggg tcagtggctc caactaacat 180

ttgtttggta ctttacagtt tattaatatag atgtttatat ggagaagctc tcatttcttt 240

ctcagaagag cctggctagg aaggtgatg aggcaccata ttcatthtgc aggtgaaatt 300

cctgagatgt aaggagctgc tgtgacttgc tcaaggcctt atatcgagta aacggtagtg 360

ctggggctta gacgcaggtg ttctgattta tagttcaaaa cctctatcaa tgagagagca 420

atctcctggt aatgtgatag atttccaac ttaatgcaa cataccataa acctcccatt 480

ctgctaattgc ccagcctaag ttggggagac cactccagat tccaagatgt acagtttgct 540

ttgctgggcc tttttcccat gcctgccttt actctgccag agttatattg ctggggtttt 600

gaagaagatc ctattaaata aaagaataag cagtattatt aagtagccct gcatttcagg 660

tttccttgag tggcaggcca ggcctggccg tgaacgttca ctgaaatcat ggctcttg 720

ccaagattga tagcttgtgc ctgtccctga gtcccagtc atcacgagca gctggtttct 780

aagatgctat ttcccgtata aagcatgaga ccgtgacttg ccagccccac agagccccgc 840

cctgtccat cactggcatc tggactccag cctgggttgg ggcaaagagg gaaatgagat 900

catgtcctaa ccctgatcct cttgtcccac agatatccag aaccctgacc ctgccgtgta 960

ccagctgaga gactctaaat ccagtgacaa gtctgtctgc ctattcgaat tcggctccgg 1020

agccactaac ttcagcctgt tgaagcaggc cggcgacgtt gaggaaaacc ccggtcctat 1080

ggccttacca gtgaccgctt tgctcctgcc gctggccttg ctgctccag ccgccaggcc 1140

gagccagttc cgggtgtcgc cgctggatcg gacctggaac ctgggcgaga cagtggagct 1200

gaagtgccag gtgctgctgt ccaaccgac gtcgggctgc tcgtggctct tccagccg 1260

cggcgccgcc gccagtccca cttcctcctt atacctctcc caaaacaagc ccaaggcggc 1320

cgaggggctg gacaccagc ggttctcggg caagaggttg ggggacacct tcgtcctcac 1380

cctgagcgac ttccgccgag agaacgaggg ctactatttc tgctcgccc tgagcaactc 1440

catcatgtac ttcagccact tcgtgccggt cttcctgcc gcaagccca ccacgacgcc 1500  
agcgcgcga ccaccaacac cggcgccac catcgcgtc cagcccctgt ccctgcgcc 1560  
agaggcgtgc cggccagcgg cggggggcgc agtgcacac agggggctgg acttcgctg 1620  
tgatatctac atctggggcg ccttgccgg gacttgtgg gtccttctcc tgtcactggt 1680  
tataccctt tactgcaacc acaggaacc aagacgtgt tgcaaatgt cccggcctgt 1740  
ggtcaaatcg ggagacaag ccagccttc ggcgagata gtcagggcta aacgggaatt 1800  
cggctccgga gccactaact tctccctgt gaaacaggct ggcatgttg aaaaaacc 1860  
cggctcctat gccaccggct ctagaacaag cctgctgct gcttttgcc tgctctgcct 1920  
cccatggctc caagaaggat ctgctaagc tgggtgact cagacccaa aattccgat 1980  
cctgaagata ggacagagca tgactgca gtgtaccag gatatgaacc ataactacat 2040  
gtactggtat cgacaagacc caggcatggg gctgaagct atttattatt cagttggtgc 2100  
tggatcact gataaaggag aagtcccga tggctacaac gtctccagat caaccacaga 2160  
ggatttccc ctcaggctgg agttggctgc tccctccag acatctgtgt acttctgtgc 2220  
cagctcccta caggttccct acaatgagca gttcttcggg ccaggacac ggctcaccgt 2280  
gctagaggac ctgaaaaac tgttccctc aaaagtggc gtgttcgagc cttctgaggc 2340  
cgagatcagc cacacacaga aagccacact cgtgtgtct gctaccggct tctaccccga 2400  
tcacgtggaa ctgtcttggg gggtaacgg caaagaggt cacagcggcg tcagcacaga 2460  
tccccagcct ctgaaagaac agcccgtct gaacgacag cgctactgcc tgtctagcag 2520  
actgagagt tccgccacct tctggcagaa cccagaaac cacttcagat gccaggcca 2580  
gttctacggc ctgagcgaga acgatgagt gaccaggac agagccaagc ctgtgacaca 2640  
gatcgtgtct gccgaagcct ggggcagagc cgattgtggc tttaccagc agtcatacca 2700  
gcagggcgtg ctgtctgcca ccacctgta tgagatcctg ctggcaagg ccactctgta 2760  
cgctgtgctg gtgtctgctc tgggtctgat ggctatggc tccgggagc gcatccccga 2820  
ggccggggc aagcggggca gcggcgccac caacttcagc ctgtgaagc agccggcga 2880  
cgtggaggag aacccggcc ctatggccac aggcagcaga acatctctgc tgctggcctt 2940  
cggactgctg tgtctgcctt ggctgcaaga gggttccgcc gctcagacag tctcagtc 3000  
tcaaccagag atgtctgtgc aggaggcaga gaccgtgacc ctgagctgca catatgacac 3060  
cagtgagagt gattattatt tattctgcta caagcagcct cccagcaggc agatgattct 3120  
cgttattcgc caagaagcct ataagcaaca gaatgcaaca gagaatcgtt tctctgtgaa 3180  
ctccagaaa gcagccaaat ccttcagtct caagatctca gactcacagc tgggggatgc 3240  
cgcgatgtat ttctgtgctt ttgggaact caacaaatt tactttggat ctgggaccaa 3300  
actcaatgta aaaccaaata ttcagaacc cgatcctgct gtgtatcagc tgcgcgacag 3360  
caagagcagc gacaagagcg tgtgtttgt caccgattt gattctcaa caatgtgtc 3420

acaaagtaag gattctgatg tgtatatcac agacaaaact gtgctagaca tgaggtctat 3480  
ggacttcaag agcaacagtg ctgtggcctg gagcaacaaa tctgactttg catgtgcaaa 3540  
cgccttcaac aacagcatta ttccagaaga caccttcttc cccagcccag gtaagggcag 3600  
ctttggtgcc ttcgcaggct gtttccttgc ttcaggaatg gccaggttct gccagagct 3660  
ctggtcaatg atgtctaaaa ctctctgat tgggtgtctc ggccttatcc attgccacca 3720  
aaaccctctt ttactaaga aacagtgagc cttgttctgg cagtccagag aatgacacgg 3780  
gaaaaaagca gatgaagaga aggtggcagg agagggcacg tggcccagcc tcagtctctc 3840  
caactgagtt cctgcctgcc tgcctttgct cagactgttt gcccttact gctcttctag 3900  
gcctcattct aagccccttc tccaagttgc ctctccttat ttctccctgt ctgcaaaaa 3960  
atctttccca gctcactaag tcagtctcac gcagtcactc attaaccac caatcactga 4020  
ttgtgccggc acatgaatgc accaggtgtt gaagtggagg aattaanaag tcagatgagg 4080  
ggtgtgcccc gaggaagcac cattctagtt gggggagccc atctgtcagc tgggaaaagt 4140  
ccaataact tcagattgga atgtgtttta actcagggtt gagaaaacag ctaccttcag 4200  
gacaaaagtc agggaagggc tctctgaaga aatgctactt gaagatacca gccctaccaa 4260  
gggcagggag aggaccctat agaggcctgg gacaggagct caatgagaaa ggagaagagc 4320  
agcaggcatg agttgaatga aggaggcagg gccgggtcac agggccttct aggccatgag 4380  
agggtagaca ggctagccgc gttgctggcg tttttccata ggctccgcc ccctgacgag 4440  
catcacaaaa atcgacgctc aagtcagagg tggcgaacc cgacaggact ataaagatac 4500  
caggcgtttc cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc 4560  
ggatacctgt ccgcttttct cccttcggga agcgtggcgc tttctcatag ctcacgctgt 4620  
aggatatctca gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc 4680  
gttcagcccc accgctgctc cttatccggt aactatcgtc ttgagtcaa cccgtaaga 4740  
cacgacttat cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta 4800  
ggcggtgcta cagagttctt gaagtgtggt cctaactacg gctacactag aagaacagta 4860  
tttggtatct gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga 4920  
tccggcaaac aaaccaccgc tggtagcggg ggtttttttg tttgcaagca gcagattacg 4980  
cgcagaaaaa aaggatctca agaagatcct ttgatcttta gaaaaactca tcgagatca 5040  
aatgaaactg caatttattc atatcaggat tatcaatacc atatttttga aaaagccgtt 5100  
tctgtaatga aggagaaaac tcaccgaggc agttccatag gatggcaaga tcctggtatc 5160  
ggtctgcatg tccgactcgt ccaacatcaa tacaacctat taatttccc tcgtcaaaaa 5220  
taaggttatc aagtgagaaa tcaccatgag tgacgactga atccggtgag aatggcaaaa 5280  
gtttatgcat ttctttccag acttgttcaa caggccagcc attacgctcg tcatcaaat 5340

cactcgcac aaccaaaccg ttattcattc gtgattgcgc ctgagccaga cgaataacgc 5400  
 gatcgctggt aaaaggacaa ttacaacag gaatcgaatg caaccggcgc aggaacactg 5460  
 ccagcgcac aacaatattt tcacctgaat caggatattc ttctaatacc tggaatgctg 5520  
 tttttccggg gatcgcagt gtgagtaacc atgcatcacc aggagtacgg ataaaatgct 5580  
 tgatggctgg aagaggcata aattccgtca gccagtttag tctgaccatc tcatctgtaa 5640  
 catcattggc aacgctacct ttgccatgtt tcagaaacaa ctctggcgca tcgggcttcc 5700  
 catacaagcg atagattgtc gcacctgatt gcccgacatt atcgcgagcc catttatacc 5760  
 catataaatc agcatccatg ttggaattta atcgcggcct cgacgtttcc cgttgaatat 5820  
 ggctcataac accccttgta ttactgttta tgtaagcaga cagttttatt gttcatgatg 5880  
 atatatTTTT atcttgtgca atgtaacatc agagattttg agacac 5926

<210> 49

<211> 926

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

Left HR Arm\_CD8 Product 2

<400> 49

acattaaaa cacaaaatcc tacggaaata ctgaagaatg agtctcagca ctaaggaaaa 60  
 gcctccagca gtcctgctt tctgagggtg aaggatagac gctgtggctc tgcatgactc 120  
 actagcactc tatcacggcc atattctggc agggctcagt gctccaacta acatttgttt 180  
 ggtactttac agtttattaa atagatgttt atatggagaa gctctcattt ctttctcaga 240  
 agagcctggc taggaagggt gatgaggcac catattcatt ttgcaggatga aattcctgag 300  
 atgtaaggag ctgctgtgac ttgctcaagg ctttatatcg agtaaaccgt agtgctgggg 360  
 cttagacgca ggtgttctga tttatagttc aaaacctcta tcaatgagag agcaatctcc 420  
 tggtaatgtg atagatttcc caacttaatg ccaacatacc ataaacctcc cattctgcta 480  
 atgcccagcc taagttgggg agaccactcc agattccaag atgtacagtt tgctttgctg 540  
 ggctttttc ccatgcctgc ctttactctg ccagagtat attgctgggg tttgaagaa 600  
 gatcctatta aataaaagaa taagcagtat tattaagtag ccctgcattt caggtttcct 660  
 tgagtggcag gccaggcctg gccgtgaacg ttactgaaa tcatggcctc ttggccaaga 720  
 ttgatagctt gtgcctgtcc ctgagtccca gtccatcacg agcagctggt ttctaagatg 780  
 ctatttcccg tataaagcat gagaccgtga cttgccagcc ccacagagcc ccgcccttgt 840  
 ccatcactgg catctggact ccagcctggg ttggggcaaa gagggaaatg agatcatgtc 900  
 ctaaccctga tcctcttgtc ccacag 926

<210> 50

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
TRAC CDS\_CD8 Product 2

<400> 50

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt 60

ctgtctgcct attc 74

<210> 51

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_CD8 Product 2

<400> 51

gaattcggct ccgga 15

<210> 52

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 2

<400> 52

gccactaact tcagcctggt gaagcaggcc ggcgacgttg aggaaaaccc cggtcct 57

<210> 53

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8A Signal Peptide\_CD8 Product 2

<400> 53

atggccttac cagtgaccgc cttgctcctg ccgctggcct tgctgctcca cgccgccagg 60

ccg 63

<210> 54

<211> 483

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8A Extracellular domain\_CD8 Product 2

<400> 54

agccagttcc ggggtgctgcc gctggatcgg acctggaacc tgggcgagac agtggagctg 60  
aagtgccagg tgctgctgtc caacccgacg tcgggctgct cgtggctctt ccagccgcgc 120  
ggcgccgccg ccagtccac cttcctccta tacctctccc aaaacaagcc caaggcggcc 180  
gaggggctgg acaccagcg gttctcgggc aagaggttg gggacacctt cgtcctcacc 240  
ctgagcgact tccgccgaga gaacgagggc tactatttct gctcggcctt gagcaactcc 300  
atcatgtact tcagccactt cgtgccggtc ttctgcccag cgaagcccac cacgaccca 360  
gcgccgcgac caccaacacc ggcgcccacc atcgcgtcgc agcccctgtc cctgcgcca 420  
gaggcgtgcc ggccagcggc gggggcgca gtgcacacga gggggctgga cttgcctgt 480  
gat 483

<210> 55  
<211> 63  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD8A Transmembrane domain\_CD8 Product 2

<400> 55  
atctacatct gggcgccctt ggccgggact tgtgggttcc ttctcctgtc actggttatc 60  
acc 63

<210> 56  
<211> 96  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD8A intracellular domain\_CD8 Product 2

<400> 56  
ctttactgca accacaggaa ccgaagacgt gtttgcaaat gtccccggcc tgtggtcaaa 60  
tcgggagaca agcccagcct ttcggcgaga tacgtc 96

<210> 57  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Furin cleavage site\_CD8 Product 2

<400> 57  
agagcaaagc gg 12

<210> 58  
<211> 9  
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_CD8 Product 2

<400> 58

ggctccgga

9

<210> 59

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 2

<400> 59

gctaccaatt ttagcctcct gaagcaggct ggcgatgttg aggaaaaccc tggcccc

57

<210> 60

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8B Signal Peptide\_CD8 Product 2

<400> 60

atgcgccgc ggctgtggct cctcttggcc gcgcagctga cagttctcca tggcaactca

60

gtc

63

<210> 61

<211> 447

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8B Extracellular domain\_CD8 Product 2

<400> 61

ctccagcaga ccctgcata cataaaggct caaaccaaca agatggatgat gctgtcctgc

60

gaggctaaaa tctccctcag taacatgcgc atctactggc tgagacagcg ccaggcaccg

120

agcagtgaca gtcaccacga gttcctggcc ctctgggatt ccgcaaaagg gactatccac

180

ggtgaagagg tggaacagga gaagatagct gtgtttcggg atgcaagccg gttcattctc

240

aatctcacia gcgtgaagcc ggaagacagt ggcattact tctgcatgat cgtcgggagc

300

cccagactga ccttcgggaa gggaactcag ctgagtgtgg ttgatttctt tcccaccact

360

gcccagccca ccaagaagtc caccctcaag aagagagtgt gccggttacc caggccagag

420

accagaagg gccactttg tagcccc

447

<210> 62  
 <211> 63  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8B Transmembrane domain\_CD8 Product 2

<400> 62  
 atcacccttg gcctgctggt ggctggcgtc ctggttctgc tggtttcct gggagtggcc 60  
 atc 63

<210> 63  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8B Intracellular domain\_CD8 Product 2

<400> 63  
 cacctgtgct gccggcggag gagagccgg cttcgtttca tgaacaatt ttacaaa 57

<210> 64  
 <211> 12  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Furin cleavage site\_CD8 Product 2

<400> 64  
 agggctaaac gg 12

<210> 65  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 GSG Linker\_CD8 Product 2

<400> 65  
 gaattcggct ccgga 15

<210> 66  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 P2A\_CD8 Product 2

<400> 66  
 gccactaact tctccctggt gaaacaggct ggcgatgttg aagaaaaccc cggtcct 57

<210> 67  
 <211> 78  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 HGH/SS/2\_CD8 Product 2

<400> 67  
 atggccaccg gctctagaac aagcctgctg ctcgcttttg gcctgctctg cctcccatgg 60  
 ctccaagaag gatctgct 78

<210> 68  
 <211> 345  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Exemplary TRB\_VDJ (TCR097) \_CD8 Product 2

<400> 68  
 aatgctgggtg tcaactcagac cccaaaattc cgcatacctga agataggaca gagcatgaca 60  
 ctgcagtgta cccaggatat gaaccataac tacatgtact ggtatcgaca agaccaggc 120  
 atggggctga agctgattta ttattcagtt ggtgctggta tcaactgataa aggagaagtc 180  
 ccgaatggct acaacgtctc cagatcaacc acagaggatt tcccgctcag gctggagttg 240  
 gctgctccct cccagacatc tgtgtacttc tgtgccagct ccctacaggt tcctacaat 300  
 gagcagttct tcgggccagg gacacggctc accgtgctag aggac 345

<210> 69  
 <211> 534  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 TCR-beta/constant\_CD8 Product 2

<400> 69  
 ctgaaaaacg tgttcctcc aaaagtggcc gtgttcgagc cttctgaggc cgagatcagc 60  
 cacacacaga aagccacact cgtgtgtctg gctaccggct tctaccccga tcacgtggaa 120  
 ctgtcttggt gggcaacgg caaagagtg cacagcggcg tcagcacaga tccccagcct 180  
 ctgaaagaac agcccgtct gaacgacagc cgctactgcc tgtctagcag actgagagtg 240  
 tccgccacct tctggcagaa cccagaaac cacttcagat gccaggtcca gttctacggc 300  
 ctgagcgaga acgatgagtg gaccaggac agagccaagc ctgtgacaca gatcgtgtct 360  
 gccgaagcct ggggcagagc cgattgtggc tttaccagcg agtcatacca gcagggcgtg 420  
 ctgtctgcca ccatcctgta tgagatcctg ctcggcaagg ccacactgta cgctgtgctg 480

gtgtctgctc tgggtgctgat ggctatggtc tcccgggagc gcatccccga gcc 534

<210> 70  
 <211> 12  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Furin cleavage site\_CD8 Product 2

<400> 70  
 cgggccaagc gg 12

<210> 71  
 <211> 9  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 GSG linker\_CD8 Product 2

<400> 71  
 ggcagcggc 9

<210> 72  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 P2A\_CD8 Product 2

<400> 72  
 gccaccaact tcagcctgct gaagcaggcc ggcgacgtgg aggagaaccc cggccct 57

<210> 73  
 <211> 78  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 HGH/SS\_CD8 Product 2

<400> 73  
 atggccacag gcagcagaac atctctgctg ctggccttcg gactgctgtg tctgccttgg 60

ctgcaagagg gttccgcc 78

<210> 74  
 <211> 337  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic

## Exemplary TRA-VDJ (TCR097) \_CD8 Product 2

<400> 74  
gctcagacag tcaactcagtc tcaaccagag atgtctgtgc aggaggcaga gaccgtgacc 60  
ctgagctgca catatgacac cagtgagagt gattattatt tattctggta caagcagcct 120  
cccagcaggc agatgattct cgttattcgc caagaagctt ataagcaaca gaatgcaaca 180  
gagaatcggt tctctgtgaa cttccagaaa gcagccaaat ccttcagtct caagatctca 240  
gactcacagc tgggggatgc cgcgatgat ttctgtgctt ttgggaactt caacaaattt 300  
tactttggat ctgggaccaa actcaatgta aaaccaa 337

<210> 75  
<211> 74  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TCR-alpha/constant\_CD8 Product 2

<400> 75  
atattcagaa ccccgatcct gctgtgtatc agctgcgcga cagcaagagc agcgacaaga 60  
gcgtgtgttt gttc 74

<210> 76  
<211> 200  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TRAC CDS/right HR arm\_CD8 Product 2

<400> 76  
accgattttg attctcaaac aatgtgtca caaagtaagg attctgatgt gtatatcaca 60  
gacaaaactg tgctagacat gaggtctatg gacttcaaga gcaacagtgc tgtggcctgg 120  
agcaacaaat ctgactttgc atgtgcaaac gccttcaaca acagcattat tccagaagac 180  
accttcttcc ccagcccagg 200

<210> 77  
<211> 800  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Right HR arm\_CD8 Product 2

<400> 77  
taagggcagc tttggcct tcgcaggctg tttccttgct tcaggaatgg ccaggttctg 60  
cccagagctc tggcfaatga tgtctaaaac tcctctgatt ggtggtctcg gccttatcca 120  
ttgccaccaa aaccctcttt ttactaagaa acagttagcc ttgttctggc agtccagaga 180

atgacacggg aaaaaagcag atgaagagaa ggtggcagga gagggcacgt ggcccagcct 240  
 cagtctctcc aactgagttc ctgcctgcct gcctttgctc agactgtttg ccccttactg 300  
 ctcttctagg cctcattcta agccccctct ccaagttgcc tctccttatt tctccctgtc 360  
 tgccaaaaaa tctttccag ctactaagt cagtctcacg cagtactca ttaaccacc 420  
 aatcactgat tgtgccgga catgaatgca ccagggttg aagtggagga attaaaaagt 480  
 cagatgaggg gtgtgccag aggaagcacc attctagttg ggggagccca tctgtcagct 540  
 gggaaaagtc caaataactt cagattggaa tgtgttttaa ctcagggttg agaaaacagc 600  
 taccttcagg acaaaagtca gggaagggt ctctgaagaa atgctacttg aagataccag 660  
 ccctaccaag ggcagggaga ggaccctata gaggcctggg acaggagctc aatgagaaag 720  
 gagaagagca gcaggcatga gttgaatgaa ggaggcaggg ccgggtcaca gggccttcta 780  
 ggccatgaga gggtagacag 800

<210> 78

<211> 6

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Artificial Sequence\_CD8 Product 2

<400> 78

gctagc

6

<210> 79

<211> 620

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
pBR322\_origin\_CD8 Product 2

<400> 79

cgcgttgctg gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg 60  
 ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt ttccccctgg 120  
 aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc tgtccgcctt 180  
 tctccccttg ggaagcgtgg cgcttttctca tagctcacgc tgtaggtatc tcagttcggt 240  
 gtaggtcgtt cgctccaagc tgggctgtgt gcacgaacc cccgttcagc ccgaccgctg 300  
 gccttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact 360  
 ggacagcagcc actggtaaca ggattagcag agcgaggat gtaggcggtg ctacagagtt 420  
 cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttggtg tctgcgctct 480  
 gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca acaaaccac 540

cgctggtagc ggtgggtttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaggatc 600

tcaagaagat cctttgatct 620

<210> 80

<211> 810

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Kanamycin resistance (KanR2) \_CD8 Product 2

<400> 80

ttagaaaaac tcatcgagca tcaaatgaaa ctgcaattta ttcatatcag gattatcaat 60

accatatttt tgaaaaagcc gtttctgtaa tgaaggagaa aactcaccga ggcagttcca 120

taggatggca agatcctggt atcggctctgc gattccgact cgtccaacat caatacaacc 180

tattaatttc ccctcgtcaa aaataaggtt atcaagttag aatcaccat gaggtagcgc 240

tgaatccggt gagaatggca aaagtttatg catttctttc cagacttggt caacaggcca 300

gccattacgc tcgtcatcaa aatcactcgc atcaaccaa cgttattca ttcgtgattg 360

cgctgagcc agacgaaata cgcgatcgct gttaaaagga caattacaaa caggaatcga 420

atgcaaccgg cgcaggaaca ctgccagcgc atcaacaata ttttcacctg aatcaggata 480

ttcttcta atctggaatg ctgtttttcc ggggatcgca gtggtgagta accatgcatc 540

atcaggagta cggataaaat gcttgatggt cggaagaggc ataaattccg tcagccagtt 600

tagtctgacc atctcatctg taacatcatt ggcaacgcta cctttgcat gtttcagaaa 660

caactctggc gcacgggct tccatacaa gcgatagatt gtcgcacctg attgcccgc 720

attatcgca gccatttat accatataa atcagcatcc atgttgaat ttaatcgcg 780

cctcgacgtt tccggtgaa tatggctcat 810

<210> 81

<211> 99

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Kanamycin promoter\_CD8 Product 2

<400> 81

aacaccctt gtattactgt ttatgtaagc agacagttt attgttcatg atgatatt 60

tttatcttgt gcaatgtaac atcagagatt ttgagacac 99

<210> 82

<211> 6634

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
 Complete CD8 Product 2 construct with TCR97 insertion\_CD8  
 Product 2

<400> 82

```

ggtagccat taaaaacaca aaatcctacg gaaatactga agaatgagtc tcagcactaa      60
ggaaaagcct ccagcagctc ctgctttctg aggggtgaagg atagacgctg tggctctgca     120
tgactcacta gcactctatc acggccatat tctggcaggg tcagtggctc caactaacat      180
ttgtttggta ctttacagtt tattaatatag atgtttatat ggagaagctc tcatttcttt     240
ctcagaagag cctggctagg aagggtgatg aggcaccata ttcatTTTgc aggtgaaatt      300
cctgagatgt aaggagctgc tgtgacttgc tcaaggcctt atatcgagta aacggtagtg     360
ctggggctta gacgcaggtg ttctgattta tagttcaaaa cctctatcaa tgagagagca     420
atctcctggg aatgtgatag atttcccaac ttaatgcaa cataccataa acctcccatt      480
ctgctaattg ccagcctaag ttggggagac cactccagat tccaagatgt acagtttgct     540
ttgctgggcc tttttcccat gcctgccttt actctgccag agttatattg ctggggTTTT     600
gaagaagatc ctattaaata aaagaataag cagtattatt aagtagccct gcatttcagg     660
tttcttgag tggcaggcca ggcctggccg tgaacgttca ctgaaatcat ggcctcttgg      720
ccaagattga tagcttgtgc ctgtccctga gtcccagtc atcacgagca gctggtttct     780
aagatgctat ttcccgtata aagcatgaga ccgtgacttg ccagccccac agagccccgc     840
ccttgtccat cactggcatc tggactccag cctgggttgg ggcaaagagg gaaatgagat     900
catgtcctaa ccctgatcct cttgtccac agatatccag aaccctgacc ctgccgtgta     960
ccagctgaga gactctaaat ccagtgacaa gtctgtctgc ctattcgaat tcggctccgg    1020
agccactaac ttcagcctgt tgaagcaggc cggcgacgtt gaggaaaacc ccggctcctat    1080
ggccttacca gtgaccgctt tgctcctgcc gctggccttg ctgctccacg ccgccaggcc    1140
gagccagttc cgggtgtcgc cgctggatcg gacctggaac ctgggcgaga cagtggagct    1200
gaagtgccag gtgctgctgt ccaaccgac gtcgggctgc tcgtggctct tccagccgcg    1260
cggcgccgcc gccagtccca ctttctcct atacctctcc caaaacaagc ccaaggcggc    1320
cgaggggctg gacaccagc ggttctcggg caagaggttg ggggacacct tcgtcctcac    1380
cctgagcgac ttccgccgag agaacgaggg ctactatttc tgctcggccc tgagcaactc    1440
catcatgtac ttcagccact tcgtgccggt cttcctgccg gcgaagccca ccacgacgcc    1500
agcggcgga ccaccaacac cggcgccac catcgctcg cagcccctgt ccctgcgcc    1560
agaggcgtgc cggccagcgg cggggggcgc agtgcacacg agggggctgg acttcgctg    1620
tgatatctac atctgggagc ctttggccgg gacttgtggg gtccttctcc tgtcactggt    1680
tatcacctt tactgcaacc acaggaaccg aagacgtgtt tgcaaatgtc cccggcctgt    1740
ggtcaaatcg ggagacaagc ccagcctttc ggcgagatac gtcagagcaa agcggggctc    1800

```

cggagctacc aatTTtagcc tcctgaagca ggctggcgat gttgaggaaa accctggtcc 1860  
catgCGGCCG cggctgtggc tcctcttggc cgCGcagctg acagtTctcc atggcaactc 1920  
agtCctccag cagaccctg catacataaa ggtgcaaacc aacaagatgg tgatgctgtc 1980  
ctgCGaggct aaaatctccc tcagtaacat gcgcatctac tggctgagac agCGccaggc 2040  
accgagcagt gacagtcacc acgagttcct ggccctctgg gattccgcaa aagggactat 2100  
ccacggtgaa gaggtggaac aggagaagat agctgtgttt cgggatgcaa gccggttcat 2160  
tctcaatctc acaagcgtga agccggaaga cagtggcatc tacttctgca tgatcgctcg 2220  
gagccccgag ctgaccttcg ggaagggaac tcagctgagt gtggttgatt tccttccac 2280  
cactgcccag cccaccaaga agtccaccct caagaagaga gtgtGCCggt taccaggcc 2340  
agagaccag aaggGCCac tttgtagccc catcaccctt ggctgctgg tggctggcgt 2400  
cctggttctg ctggtttccc tgggagtggc catccacctg tgctGCCggc ggaggagagc 2460  
cggcttctg ttcatgaaac aatTTTaca aaggGctaaa cgggaattcg gctccggagc 2520  
cactaacttc tcctgttga aacaggctgg cgatgttga gaaaacccg gtcctatggc 2580  
caccggctct agaacaagcc tgctgctgc ttttggcctg ctctgcctcc catggctcca 2640  
agaaggatct gctaatgctg gtgtcactca gacccaaaa ttccgatcc tgaagatagg 2700  
acagagcatg aactgCagt gtaccagga tatgaaccat aactacatgt actggtatcg 2760  
acaagacca ggcaggggc tgaagctgat ttattatca gttggtgctg gtatcactga 2820  
taaaggagaa gtcccgaatg gctacaacgt ctccagatca accacagagg atttcccgct 2880  
caggctggag ttggctgctc cctccagac atctgtgtac ttctgtgcca gctccctaca 2940  
ggttccctac aatgagcagt tcttcgggcc agggacacgg ctaccgtgc tagaggacct 3000  
gaaaaacgtg ttccctcaa aagtggccgt gttcGagcct tctgaggccg agatcagcca 3060  
cacacagaaa gccacactcg tgtgtctggc taccggctt caccgatc acgtggaact 3120  
gtcttggtgg gtcaacggca aagaggTgca cagCGcgctc agcacagatc cccagcctct 3180  
gaaagaacag cccgctctga acgacagccg ctactgcctg tctagcagac tgagagtgtc 3240  
cgccaccttc tggcagaacc ccagaaacca cttcagatgc caggTccagt tctacggcct 3300  
gagCGagaac gatgagtgga cccaggacag agccaagcct gtgacacaga tcgtgtctgc 3360  
cgaagcctgg ggcagagccg attgtggctt taccagCGag tcataccagc agggCGtgct 3420  
gtctGCCacc atcctgtatg agatcctgct cggcaaggcc aactgtacg ctgtgctggt 3480  
gtctgctctg gtgctgatgg ctatggtctc ccgggagcgc atccccgagg cccggGCCaa 3540  
gcggggcagc ggcGCCacca acttcagcct gctgaagcag gccggCGacg tggaggagaa 3600  
ccccggccct atggccacag gcagcagaac atctctgctg ctggccttcg gactgctgtg 3660  
tctgccttgg ctgcaagagg gttccGCCgc tcagacagtc actcagtctc aaccagagat 3720  
gtctgtgcag gaggcagaga ccgtgaccct gagctgcaca tatgacacca gtgagagtga 3780

ttattattta	ttctggtaca	agcagcctcc	cagcaggcag	atgattctcg	ttattcgcca	3840
agaagcttat	aagcaacaga	atgcaacaga	gaatcgtttc	tctgtgaact	tccagaaagc	3900
agccaaatcc	ttcagtctca	agatctcaga	ctcacagctg	ggggatgccg	cgatgtattt	3960
ctgtgctttt	gggaacttca	acaaatttta	ctttggatct	gggaccaaac	tcaatgtaaa	4020
accaaataat	cagaaccccc	atcctgctgt	gtatcagctg	cgcgacagca	agagcagcga	4080
caagagcgtg	tgtttgttca	ccgattttga	ttctcaaaca	aatgtgtcac	aaagtaagga	4140
ttctgatgtg	tatatcacag	acaaaactgt	gctagacatg	aggtctatgg	acttcaagag	4200
caacagtgct	gtggcctgga	gcaacaaatc	tgactttgca	tgtgcaaacg	ccttcaacaa	4260
cagcattatt	ccagaagaca	ccttcttccc	cagcccaggt	aagggcagct	ttggtgcctt	4320
cgcaggctgt	ttccttgctt	caggaatggc	caggttctgc	ccagagctct	ggtcaatgat	4380
gtctaaaact	cctctgattg	gtggtctcgg	ccttatccat	tgccaccaa	accctctttt	4440
tactaagaaa	cagtgagcct	tgttctggca	gtccagagaa	tgacacggga	aaaaagcaga	4500
tgaagagaag	gtggcaggag	agggcacgtg	gcccagcctc	agtctctcca	actgagttcc	4560
tgctgcctg	cctttgctca	gactgtttgc	cccttactgc	tcttctaggc	ctcattctaa	4620
gccccttctc	caagttgcct	ctccttattt	ctccctgtct	gccaaaaaat	ctttcccagc	4680
tcactaagtc	agtctcacgc	agtcactcat	taaccaccca	atcactgatt	gtgccggcac	4740
atgaatgcac	caggtgttga	agtggaggaa	ttaaaaagtc	agatgagggg	tgtgcccaga	4800
ggaagcacca	ttctagttagg	gggagcccat	ctgtcagctg	ggaaaagtcc	aaataacttc	4860
agattggaat	gtgttttaac	tcagggttga	gaaaacagct	accttcagga	caaaagtcag	4920
ggaagggctc	tctgaagaaa	tgctacttga	agataccagc	cctaccaagg	gcagggagag	4980
gacctatag	aggcctggga	caggagctca	atgagaaagg	agaagagcag	caggcatgag	5040
ttgaatgaag	gaggcagggc	cgggtcacag	ggccttctag	gccatgagag	ggtagacagg	5100
ctagccgctg	tgctggcggt	tttccatagg	ctccgcccc	ctgacgagca	tcacaaaaat	5160
cgacgtcaa	gtcagagggtg	gcgaaacccg	acaggactat	aaagatacca	ggcgtttccc	5220
cctggaagct	ccctcgtgcg	ctctcctggt	ccgaccctgc	cgcttaccgg	atacctgtcc	5280
gcctttctcc	cttcgggaag	cgtggcgctt	tctcatagct	cacgctgtag	gtatctcagt	5340
tcgggtgtagg	tcgttcgctc	caagctgggc	tgtgtgcacg	aacccccgt	tcagcccagc	5400
cgctgcgctt	tatccggtaa	ctatcgtctt	gagtccaacc	cgtaagaca	cgacttatcg	5460
ccactggcag	cagccactgg	taacaggatt	agcagagcga	ggtatgtagg	cggtgctaca	5520
gagttcttga	agtgggtggc	taactacggc	tacactagaa	gaacagtatt	tggtatctgc	5580
gctctgctga	agccagttac	cttcggaaaa	agagttggta	gctcttgatc	cggcaaacaa	5640
accaccgctg	gtagcgggtg	tttttttgtt	tgcaagcagc	agattacgcg	cagaaaaaaa	5700

ggatctcaag aagatccttt gatctttaga aaaactcatc gagcatcaaa tgaactgca 5760  
 atttattcat atcaggatta tcaataccat atttttgaaa aagccgtttc tgtaatgaag 5820  
 gagaaaactc accgaggcag ttccatagga tggcaagatc ctggtatcgg tctgcgattc 5880  
 cgactcgtcc aacatcaata caacctatta atttcccctc gtcaaaaata aggttatcaa 5940  
 gtgagaaatc accatgagtg acgactgaat ccggtgagaa tggcaaaagt ttatgcattt 6000  
 cttccagac ttgttcaaca ggccagccat tacgctcgtc atcaaaatca ctcgcatcaa 6060  
 ccaaaccgtt attcattcgt gattgcgctt gagccagacg aaatacgcga tcgctgttaa 6120  
 aaggacaatt acaaacagga atcgaatgca accggcgagc gaactgccc agcgcatcaa 6180  
 caatattttc acctgaatca ggatattctt ctaatacctg gaatgctgtt tttccgggga 6240  
 tcgcagtggg gagtaacat gcatcatcag gagtacggat aaaatgcttg atggtcggaa 6300  
 gaggcataaa ttccgtcagc cagtttagtc tgaccatctc atctgtaaca tcattggcaa 6360  
 cgctaccttt gccatgtttc agaaacaact ctggcgcagc gggcttcca tacaagcgat 6420  
 agattgtcgc acctgattgc ccgacattat cgcgagccca tttatacca tataaatcag 6480  
 catccatggt ggaatttaac cgcggcctcg acgtttcccg ttgaatatgg ctcataacac 6540  
 cccttgatt actgtttatg taagcagaca gttttattgt tcatgatgat atatttttat 6600  
 cttgtgcaat gtaacatcag agattttgag acac 6634

<210> 83

<211> 926

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Left HR Arm\_CD8 Product 3

<400> 83

acattaaaaa cacaaaatcc tacggaaata ctgaagaatg agtctcagca ctaaggaaaa 60  
 gcctccagca gctcctgctt tctgagggtg aaggatagac gctgtggctc tgcatgactc 120  
 actagcactc tatcacggcc atattctggc agggtcagtg gctccaacta acatttgttt 180  
 ggtactttac agtttattaa atagatgttt atatggagaa gctctcattt ctttctcaga 240  
 agagcctggc taggaagggt gatgaggcac catattcatt ttgcaggtga aattcctgag 300  
 atgtaaggag ctgctgtgac ttgctcaagg ctttatatcg agtaaaccgt agtgctgggg 360  
 cttagacgca ggtgttctga tttatagttc aaaacctcta tcaatgagag agcaatctcc 420  
 tgtaaatgtg atagatttcc caacttaatg ccaacatacc ataaacctcc cattctgcta 480  
 atgcccagcc taagtggggg agaccactcc agattccaag atgtacagtt tgctttgctg 540  
 ggcttttttc ccatgcctgc ctttactctg ccagagttat attgctgggg ttttgaagaa 600  
 gatcctatta aataaaagaa taagcagtat tattaagtag ccctgcattt caggtttcct 660

tgagtggcag gccaggcctg gccgtgaacg ttcactgaaa tcatggcctc ttggccaaga 720  
 ttgatagctt gtgcctgtcc ctgagtccca gtccatcacg agcagctggt ttctaagatg 780  
 ctattttccg tataaagcat gagaccgtga cttgccagcc ccacagagcc ccgcccttgt 840  
 ccatcactgg catctggact ccagcctggg ttggggcaaa gagggaaatg agatcatgtc 900  
 ctaaccctga tcctcttgtc ccacag 926

<210> 84  
 <211> 74  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 TRAC (part of homology arm) \_CD8 Product 3

<400> 84  
 atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt 60  
 ctgtctgcct attc 74

<210> 85  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 GSG Linker\_CD8 Product 3

<400> 85  
 gaattcggct ccgga 15

<210> 86  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 P2A\_CD8 Product 3

<400> 86  
 gccactaact tcagcctggt gaagcaggcc ggcgacgttg aggaaaaccc cggctcct 57

<210> 87  
 <211> 63  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8A Signal Peptide\_CD8 Product 3

<400> 87  
 atggccttac cagtgaccgc cttgctcctg ccgctggcct tgctgctcca cgccgccagg 60  
 ccg 63

<210> 88  
 <211> 483  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8A Extracellular domain\_CD8 Product 3

<400> 88  
 agccagttcc ggggtgctgcc gctggatcgg acctggaacc tgggagagac agtggagctg 60  
 aagtgccagg tgctgctgtc caaccgacg tcgggctgct cgtggctctt ccagccgcgc 120  
 ggcgcccgcg ccagtcccac cttcctccta tacctctccc aaaacaagcc caaggcggcc 180  
 gaggggctgg acaccagcg gttctcgggc aagaggttg gggacacctt cgtcctcacc 240  
 ctgagcgact tccgccgaga gaacgagggc tactatttct gctcggccct gagcaactcc 300  
 atcatgtact tcagccactt cgtgccggtc ttctgcccag cgaagcccac cacgacgcca 360  
 gcgccgcgac caccaacacc ggcgcccacc atcgcgtcgc agcccctgtc cctgcgcca 420  
 gaggcgtgcc ggccagcggc ggggggcgca gtgcacacga gggggctgga cttgcctgt 480  
 gat 483

<210> 89  
 <211> 63  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8A Transmembrane domain\_CD8 Product 3

<400> 89  
 atctacatct gggcgccctt ggccgggact tgtggggtcc ttctcctgtc actggttatc 60  
 acc 63

<210> 90  
 <211> 57  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8B intracellular domain\_CD8 Product 3

<400> 90  
 cacctgtgct gccggcggag gagagcccgg cttcgtttca tgaacaatt ttacaaa 57

<210> 91  
 <211> 12  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Furin cleavage site\_CD8 Product 3

<400> 91  
agggctaaac gg

12

<210> 92  
<211> 15  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_CD8 Product 3

<400> 92  
gaattcggct ccgga

15

<210> 93  
<211> 57  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 3

<400> 93  
gccactaact tctccctggt gaaacaggct ggcatgttg aagaaaaccc cggctct

57

<210> 94  
<211> 78  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
HGH/SS/2\_CD8 Product 3

<400> 94  
atggccaccg gctctagaac aagcctgctg ctcgcttttg gcctgctctg cctcccatgg

60

ctccaagaag gatctgct

78

<210> 95  
<211> 345  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TRB\_VDJ (TCR097)\_CD8 Product 3

<400> 95  
aatgctgggtg tcaactcagac cccaaaattc cgcatcctga agataggaca gagcatgaca

60

ctgcagtgtgta cccaggatat gaaccataac tacatgtact ggtatcgaca agaccaggc

120

atggggctga agctgattta ttattcagtt ggtgctggta tcaactgataa aggagaagtc

180

ccgaatggct acaacgtctc cagatcaacc acagaggatt tcccgctcag gctggagttg

240

gctgtccct cccagacatc tgtgtacttc tgtgccagct ccctacaggt tcctacaat 300

gagcagttct tcgggccagg gacacggctc accgtgctag aggac 345

<210> 96

<211> 534

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
TCR-beta/constant\_CD8 Product 3

<400> 96

ctgaaaaacg tgttcctcc aaaagtggcc gtgttcgagc cttctgaggc cgagatcagc 60

cacacacaga aagccacact cgtgtgtctg gctaccggct tctaccccga tcacgtggaa 120

ctgtcttggg gggtaacgg caaagagggt cacagcggcg tcagcacaga tccccagcct 180

ctgaaagaac agcccgtct gaacgacagc cgctactgcc tgtctagcag actgagagtg 240

tccgccacct tctggcagaa cccagaaac cacttcagat gccaggcca gttctacggc 300

ctgagcgaga acgatgagt gacccaggac agagccaagc ctgtgacaca gatcgtgtct 360

gccgaagcct ggggcagagc cgattgtggc tttaccagcg agtcatacca gcagggcgtg 420

ctgtctgcca ccatcctgta tgagatcctg ctcggcaagg ccacactgta cgctgtgctg 480

gtgtctgctc tgggtgctgat ggctatggtc tcccgggagc gcatccccga ggcc 534

<210> 97

<211> 12

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Furin cleavage site\_CD8 Product 3

<400> 97

cgggcccaagc gg 12

<210> 98

<211> 9

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
GSG linker\_CD8 Product 3

<400> 98

ggcagcggc 9

<210> 99

<211> 57

<212> DNA

<213> Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 3

&lt;400&gt; 99

gccaccaact tcagcctgct gaagcaggcc ggcgacgtgg aggagaaccc cggccct 57

&lt;210&gt; 100

&lt;211&gt; 78

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
HGH/SS\_CD8 Product 3

&lt;400&gt; 100

atggccacag gcagcagaac atctctgctg ctggccttcg gactgctgtg tctgccttgg 60

ctgcaagagg gttccgcc 78

&lt;210&gt; 101

&lt;211&gt; 337

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Exemplary TRA-VDJ (TCR097) \_CD8 Product 3

&lt;400&gt; 101

gctcagacag tcaactcagtc tcaaccagag atgtctgtgc aggaggcaga gaccgtgacc 60

ctgagctgca catatgacac cagtgagagt gattattatt tattctggta caagcagcct 120

cccagcaggc agatgattct cgttattcgc caagaagcct ataagcaaca gaatgaaca 180

gagaatcggt tctctgtgaa cttccagaaa gcagccaaat ccttcagtct caagatctca 240

gactcacagc tgggggatgc cgcatgtat ttctgtgctt ttgggaactt caacaaattt 300

tactttggat ctgggaccaa actcaatgta aaaccaa 337

&lt;210&gt; 102

&lt;211&gt; 74

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
TCR-alpha/constant\_CD8 Product 3

&lt;400&gt; 102

atattcagaa cccgatcct gctgtgtatc agctgcgcga cagcaagagc agcgacaaga 60

gcgtgtgttt gttc 74

&lt;210&gt; 103

&lt;211&gt; 200

&lt;212&gt; DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
TRAC CDS/right HR arm\_CD8 Product 3

<400> 103

accgattttg attctcaaac aaatgtgtca caaagtaagg attctgatgt gtatatcaca	60
gacaaaactg tgctagacat gaggtctatg gacttcaaga gcaacagtgc tgtggcctgg	120
agcaacaaat ctgactttgc atgtgcaaac gccttcaaca acagcattat tccagaagac	180
accttcttcc ccagcccagg	200

<210> 104

<211> 800

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Right HR arm\_CD8 Product 3

<400> 104

taagggcagc tttggtgcct tcgcaggctg tttccttgct tcaggaatgg ccaggttctg	60
cccagagctc tggatcaatga tgtctaaaac tcctctgatt ggtggtctcg gccttatcca	120
ttgccaccaa aaccctcttt ttactaagaa acagttagcc ttgttctggc agtccagaga	180
atgacacggg aaaaaagcag atgaagagaa ggtggcagga gagggcacgt ggcccagcct	240
cagtctctcc aactgagttc ctgcctgcct gcctttgctc agactgtttg ccccttactg	300
ctcttctagg cctcattcta agccccctct ccaagttgcc tctccttatt tctccctgtc	360
tgccaaaaaa tctttccag ctactaagt cagtctcag cagtactca ttaaccacc	420
aatcactgat tgtgccggca catgaatgca ccagggttg aagtggagga attaaaaagt	480
cagatgaggg gtgtgcccag aggaagcacc attctagttg ggggagccca tctgtcagct	540
gggaaaagtc caaataactt cagattggaa tgtgttttaa ctcagggttg agaaaacagc	600
taccttcagg acaaaagtca gggaaggct ctctgaagaa atgctacttg aagataccag	660
ccctaccaag ggcagggaga ggaccctata gaggcctggg acaggagctc aatgagaaag	720
gagaagagca gcaggcatga gttgaatgaa ggaggcaggg ccgggtcaca gggccttcta	780
ggccatgaga gggtagacag	800

<210> 105

<211> 6

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Artificial Sequence\_CD8 Product 3

<400> 105

gctagc

6

&lt;210&gt; 106

&lt;211&gt; 620

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
pBR322\_origin\_CD8 Product 3

&lt;400&gt; 106

cgcggttgctg gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg 60  
ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt ttccccctgg 120  
aagctccctc gtgcgctctc ctgttccgac cctgcccgtt accggatacc tgtccgcctt 180  
tctcccttcg ggaagcgtgg cgcttttctca tagctcacgc tgtaggtatc tcagttcggg 240  
gtaggtcggt cgctccaagc tgggctgtgt gcacgaacc cccgttcagc ccgaccgctg 300  
cgcttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact 360  
ggcagcagcc actggtaaca ggattagcag agcgaggat gtaggcggtg ctacagagtt 420  
cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttggtg tctgcgctct 480  
gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca acaaaaccac 540  
cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgagaa aaaaggatc 600  
tcaagaagat cctttgatct 620

&lt;210&gt; 107

&lt;211&gt; 810

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Kanamycin resistance (KanR2) \_CD8 Product 3

&lt;400&gt; 107

ttagaaaaac tcatcgagca tcaaatgaaa ctgcaattta ttcatatcag gattatcaat 60  
accatatttt tgaaaaagcc gtttctgtaa tgaaggagaa aactcaccga ggcagttcca 120  
taggatggca agatcctggt atcggctctgc gattccgact cgtccaacat caatacaacc 180  
tattaatttc cctcgtcaaa aaataagggt atcaagttag aatcaccat gagtgcgac 240  
tgaatccggt gagaatggca aaagtttatg catttctttc cagacttggt caacaggcca 300  
gccattacgc tcgtcatcaa aatcactcgc atcaacaaa ccgttattca ttcgtgattg 360  
cgctgagcc agacgaaata cgcgatcgtt gttaaaagga caattacaaa caggaatcga 420  
atgcaaccgg cgcaggaaca ctgccagcgc atcaacaata tttcacctg aatcaggata 480  
ttcttctaata acctggaatg ctgtttttcc ggggatcgcga gtggtgagta accatgcatc 540  
atcaggagta cggataaaat gcttgatggt cggaagaggc ataaattccg tcagccagtt 600

tagtctgacc atctcatctg taacatcatt ggcaacgcta cctttgccat gtttcagaaa 660  
caactctggc gcatcgggct tcccatacaa gcgatagatt gtcgcacctg attgcccgac 720  
attatcgga gccatttat acccatataa atcagcatcc atgttgaat ttaatcgcg 780  
cctcgacgtt tccggttga tatggctcat 810

<210> 108  
<211> 99  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Kanamycin promoter\_CD8 Product 3

<400> 108  
aacacccctt gtattactgt ttatgtaagc agacagtttt attgttcatg atgatatt 60  
tttatcttgt gcaatgtaac atcagagatt ttgagacac 99

<210> 109  
<211> 5887  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Complete CD8 Product 3 construct with TCR97 insertion\_CD8  
Product 3

<400> 109  
ggtaccacat taaaaacaca aaatcctacg gaaatactga agaatgagtc tcagcactaa 60  
ggaaaagcct ccagcagctc ctgctttctg aggggtgaagg atagacgctg tggctctgca 120  
tgactcacta gactctatc acggccatat tctggcaggg tcagtggctc caactaacat 180  
ttgtttggta ctttacagtt tattaatag atgtttatat ggagaagctc tcatttcttt 240  
ctcagaagag cctggctagg aagggtgatg aggcaccata ttcatthtgc aggtgaaatt 300  
cctgagatgt aaggagctgc tgtgacttgc tcaaggcctt atatcgagta aacggtagtg 360  
ctggggctta gacgcaggtg ttctgattta tagttcaaaa cctctatcaa tgagagagca 420  
atctcctggg aatgtgatag atttcccaac ttaatgcaa cataccataa acctccatt 480  
ctgctaagtc ccagcctaag ttggggagac cactccagat tccaagatgt acagtttgct 540  
ttgctgggcc tttttccat gcctgccttt actctgccag agttatattg ctggggtttt 600  
gaagaagatc ctattaaata aaagaataag cagtattatt aagtagccct gcatttcagg 660  
tttcttgag tggcaggcca ggcctggccg tgaacgttca ctgaaatcat ggcctcttgg 720  
ccaagattga tagcttgtgc ctgtccctga gtcccagtc atcacgagca gctggtttct 780  
aagatgctat ttcccgtata aagcatgaga ccgtgacttg ccagccccac agagccccgc 840  
cctgtccat cactggcatc tggactccag cctgggttgg ggcaaagagg gaaatgagat 900

catgtcctaa	ccctgatcct	cttgtccac	agatatccag	aaccctgacc	ctgccgtgta	960
ccagctgaga	gactctaaat	ccagtgacaa	gtctgtctgc	ctattcgaat	tcggctccgg	1020
agccactaac	ttcagcctgt	tgaagcaggc	cggcgacgtt	gaggaaaacc	ccggtcctat	1080
ggccttacca	gtgaccgct	tgctcctgcc	gctggccttg	ctgctccacg	ccgccaggcc	1140
gagccagttc	cgggtgtcgc	cgctggatcg	gacctggaac	ctgggcgaga	cagtggagct	1200
gaagtgccag	gtgctgtgt	ccaacccgac	gtcgggctgc	tcgtggctct	tccagccgcg	1260
cggcgccgcc	gccagtccca	ccttcctcct	atacctctcc	caaaacaagc	ccaaggcggc	1320
cgaggggctg	gacaccagc	ggttctcggg	caagaggttg	ggggacacct	tcgtcctcac	1380
cctgagcgac	ttccgccgag	agaacgaggg	ctactatttc	tgctcggccc	tgagcaactc	1440
catcatgtac	ttcagccact	tcgtgccggt	cctcctgcc	gcgaagccca	ccacgacgcc	1500
agcgcgcga	ccaccaacac	cggcgccac	catcgcgtcg	cagcccctgt	ccctgcgcc	1560
agaggcgtgc	cggccagcgg	cggggggcgc	agtgcacacg	agggggctgg	acttcgctg	1620
tgatatctac	atctgggcgc	ccttggccgg	gacttgtggg	gtccttctcc	tgtcactggt	1680
tatcaccac	ctgtgctgcc	ggcggaggag	agcccggctt	cgtttcatga	aacaatttta	1740
caaaagggct	aaacgggaat	tcggctccgg	agccactaac	ttctccctgt	tgaaacaggc	1800
tggcgatggt	gaagaaaacc	ccggctctat	ggccaccggc	tctagaacaa	gcctgctgct	1860
cgcttttggc	ctgctctgcc	tcccatggct	ccaagaagga	tctgctaata	ctggtgtcac	1920
tcagacccca	aaattccgca	tcctgaagat	aggacagagc	atgacactgc	agtgtacca	1980
ggatatgaac	cataactaca	tgtactggta	tcgacaagac	ccaggcatgg	ggctgaagct	2040
gatttattat	tcagttgggt	ctggtatcac	tgataaagga	gaagtcccga	atggctacaa	2100
cgctctcaga	tcaaccacag	aggatttccc	gctcaggctg	gagttggctg	ctcctccca	2160
gacatctgtg	tacttctgtg	ccagctccct	acaggttccc	tacaatgagc	agttcttcgg	2220
gccagggaca	cggctcaccg	tgctagagga	cctgaaaaac	gtgttccctc	caaaagtggc	2280
cgtgttcgag	ccttctgagg	ccgagatcag	ccacacacag	aaagccacac	tcgtgtgtct	2340
ggctaccggc	ttctaccccg	atcacgtgga	actgtcttgg	tgggtcaacg	gcaaagaggt	2400
gcacagcggc	gtcagcacag	atccccagcc	tctgaaagaa	cagcccgtc	tgaacgacag	2460
ccgctactgc	ctgtctagca	gactgagagt	gtccgccacc	ttctggcaga	acccagaaa	2520
ccacttcaga	tgccaggtcc	agttctacgg	cctgagcag	aacgatgagt	ggaccagga	2580
cagagccaag	cctgtgacac	agatcgtgtc	tgccgaagcc	tggggcagag	ccgattgtgg	2640
ctttaccagc	gagtcatacc	agcagggcgt	gctgtctgcc	accatcctgt	atgagatcct	2700
gctcggcaag	gccacactgt	acgctgtgct	ggtgtctgct	ctggtgctga	tggctatggt	2760
ctcccgggag	cgcattcccc	aggcccgggc	caagcggggc	agcggcgcca	ccaacttcag	2820

cctgctgaag caggccggcg acgtggagga gaacccggc cctatggcca caggcagcag 2880  
aacatctctg ctgctggcct tcggactgct gtgtctgcct tggctgcaag agggttccgc 2940  
cgctcagaca gtcactcagt ctcaaccaga gatgtctgtg caggaggcag agaccgtgac 3000  
cctgagctgc acatatgaca ccagtgagag tgattattat ttattctggt acaagcagcc 3060  
tcccagcagg cagatgattc tcgttattcg ccaagaagct tataagcaac agaatgcaac 3120  
agagaatcgt ttctctgtga acttccagaa agcagccaaa tccttcagtc tcaagatctc 3180  
agactcacag ctgggggatg ccgcgatgta tttctgtgct tttgggaact tcaacaaatt 3240  
ttactttgga tctgggacca aactcaatgt aaaaccaaatt attcagaacc cccgatcctgc 3300  
tgtgtatcag ctgctgcgaca gcaagagcag cgacaagagc gtgtgtttgt tcaccgattt 3360  
tgattctcaa acaaatgtgt cacaaagtaa ggattctgat gtgtatatca cagacaaaac 3420  
tgtgctagac atgaggctca tggacttcaa gagcaacagt gctgtggcct ggagcaacaa 3480  
atctgacttt gcatgtgcaa acgccttcaa caacagcatt attccagaag acaccttctt 3540  
ccccagccca ggtaagggca gctttggtgc cttcgcaggc tgtttccttg cttcaggaat 3600  
ggccaggttc tgcccagagc tctggtcaat gatgtctaaa actcctctga ttggtggtct 3660  
cggccttatt cattgccacc aaaaccctct tttactaag aaacagtgag ccttgttctg 3720  
gcagtccaga gaatgacacg ggaaaaaagc agatgaagag aaggaggcag gagagggcac 3780  
gtggcccagc ctgactctct ccaactgagt tcctgcctgc ctgcctttgc tcagactgtt 3840  
tgccccttac tgctcttcta ggctcattc taagcccctt ctccaagttg cctctcctta 3900  
tttctccctg tctgccaaaa aatctttccc agctactaa gtcagtctca cgcagtcact 3960  
cattaacca ccaatcactg attgtgccgg cacatgaatg caccaggtgt tgaagtggag 4020  
gaattaaaaa gtcagatgag ggggtgtgcc agaggaagca ccattctagt tgggggagcc 4080  
catctgtcag ctgggaaaag tccaaataac ttcagattgg aatgtgtttt aactcagggt 4140  
tgagaaaaca gctaccttca ggacaaaagt cagggaaggg ctctctgaag aatgctact 4200  
tgaagatacc agccctacca agggcagggg gaggacccta tagaggcctg ggacaggagc 4260  
tcaatgagaa aggagaagag cagcaggcat gagttgaatg aaggaggcag ggccgggtca 4320  
cagggccttc taggcatga gagggtagac aggctagccg cgttgctggc gtttttccat 4380  
aggctccgcc cccctgacga gcatcacaaa aatcgacgct caagtcagag gtggcgaaac 4440  
ccgacaggac tataaagata ccaggcgttt cccctggaa gctccctcgt gcgctctcct 4500  
gttccgacct tgccgcttac cggatacctg tccgcctttc tcccttcggg aagcgtggcg 4560  
ctttctcata gctcacgctg taggtatctc agttcgggtg aggtcgttcg ctccaagctg 4620  
ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg cttatccgg taactatcgt 4680  
cttgagtcca acccggtaag acacgactta tcgccactgg cagcagccac tggtaacagg 4740  
attagcagag cgaggatgt aggcgggtgct acagagtctt tgaagtggtg gcctaactac 4800

ggctacacta gaagaacagt atttggtatc tgcgctctgc tgaagccagt taccttcgga 4860  
 aaaagagttg gtagctcttg atccggcaaa caaaccaccg ctggtagcgg tggttttttt 4920  
 gtttgcaagc agcagattac gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt 4980  
 agaaaaactc atcgagcatc aaatgaaact gcaatttatt catatcagga ttatcaatac 5040  
 catatTTTTG aaaaagccgt ttctgtaatg aaggagaaaa ctcaccgagg cagttccata 5100  
 ggatggcaag atcctggtat cggctctgca ttccgactcg tccaacatca atacaacct 5160  
 ttaatttccc ctcgtcaaaa ataaggttat caagtgagaa atcaccatga gtgacgactg 5220  
 aatccggtga gaatggcaaa agtttatgca tttctttcca gacttgttca acaggccagc 5280  
 cattacgctc gtcatcaaaa tcaactcgcat caaccaaac gttattcatt cgtgattgcg 5340  
 cctgagccag acgaaatacg cgatcgctgt taaaaggaca attacaaca ggaatcgaat 5400  
 gcaaccggcg caggaacact gccagcgcat caacaatatt ttcacctgaa tcaggatatt 5460  
 cttctaatac ctggaatgct gtttttccgg ggatcgcagt ggtgagtaac catgcatcat 5520  
 caggagtacg gataaaatgc ttgatggtcg gaagaggcat aaattccgtc agccagttta 5580  
 gtctgaccat ctcatctgta acatcattgg caacgctacc tttgccatgt ttcagaaaca 5640  
 actctggcgc atcgggcttc ccatacaagc gatagattgt cgcacctgat tgcccacat 5700  
 tatcgcgagc ccatttatac ccatataaat cagcatccat gttggaattt aatcgcggcc 5760  
 tcgacgtttc ccgttgaata tggctcataa cacccttgt attactgttt atgtaagcag 5820  
 acagttttat tgttcatgat gatatatttt tatcttgtgc aatgtaacat cagagatttt 5880  
 gagacac 5887

<210> 110

<211> 926

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
Left HR Arm\_CD8 Product 4

<400> 110

acattaaaa cacaaatcc tacggaaata ctgaagaatg agtctcagca ctaaggaaaa 60  
 gcctccagca gctcctgctt tctgaggggtg aaggatagac gctgtggctc tgcatgactc 120  
 actagcactc tatcacggcc atattctggc agggtcagtg gctccaacta acatttgttt 180  
 ggtactttac agtttattaa atagatgttt atatggagaa gctctcattt ctttctcaga 240  
 agagcctggc taggaagggtg gatgaggcac catattcatt ttgcaggatga aattcctgag 300  
 atgtaaggag ctgctgtgac ttgctcaagg ctttatatcg agtaaacggt agtgctgggg 360  
 cttagacgca ggtgttctga tttatagttc aaaacctcta tcaatgagag agcaatctcc 420  
 tgtaaatgtg atagatttcc caacttaatg ccaacatacc ataacctcc cattctgcta 480

atgccagcc taagttgggg agaccactcc agattccaag atgtacagtt tgctttgctg 540  
 gccctttttc ccatgcctgc ctttactctg ccagagttat attgctgggg ttttgaagaa 600  
 gatcctatta aataaaagaa taagcagtat tattaagtag ccctgcattt caggtttctt 660  
 tgagtggcag gccaggcctg gccgtgaacg ttcactgaaa tcatggcctc ttggccaaga 720  
 ttgatagctt gtgcctgtcc ctgagtccca gtccatcacg agcagctggt ttctaagatg 780  
 ctatttcccg tataaagcat gagaccgtga cttgccagcc ccacagagcc ccgcccttgt 840  
 ccatcactgg catctggact ccagcctggg ttggggcaaa gagggaaatg agatcatgtc 900  
 ctaaccctga tcctcttgtc ccacag 926

<210> 111

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
 TRAC CDS\_CD8 Product 4

<400> 111

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt 60

ctgtctgcct attc 74

<210> 112

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
 GSG Linker\_CD8 Product 4

<400> 112

gaattcggct ccgga 15

<210> 113

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
 P2A\_CD8 Product 4

<400> 113

gccactaact tcagcctggt gaagcaggcc ggcgacgttg aggaaaaccc cggtcct 57

<210> 114

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
CD8A Signal peptide\_CD8 Product 4

<400> 114  
atggccttac cagtgaccgc cttgctcctg ccgctggcct tgctgctcca cgccgccagg 60  
ccg 63

<210> 115  
<211> 483  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD8A Extracellular Domain\_CD8 Product 4

<400> 115  
agccagttcc ggggtgctgcc gctggatcgg acctggaacc tgggagagac agtggagctg 60  
aagtgccagg tgctgctgtc caaccgacg tcgggctgct cgtggctctt ccagccgcgc 120  
ggcgcgccc ccagtccac cttcctccta tacctctccc aaaacaagcc caaggcggcc 180  
gaggggctgg acaccagcg gttctcgggc aagaggttgg gggacacctt cgtcctcacc 240  
ctgagcgact tccgccgaga gaacgagggc tactatttct gctcggcctt gagcaactcc 300  
atcatgtact tcagccactt cgtgccggtc ttctgcccag cgaagcccac cacgacgcca 360  
gcgccgcgac caccaacacc ggcgcccacc atcgcgtcgc agcccctgtc cctgcgccc 420  
gaggcgtgcc ggccagcggc ggggggcgca gtgcacacga gggggctgga cttgcctgt 480  
gat 483

<210> 116  
<211> 63  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD8A transmembrane domain\_CD8 Product 4

<400> 116  
atctacatct gggcgccctt ggccgggact tgtgggtcc ttctctgtc actggttatc 60  
acc 63

<210> 117  
<211> 120  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD4 intracellular domain\_CD8 Product 4

<400> 117  
tgtgtcaggt gccggcaccg aaggcgccaa gcagagcgga tgtctcagat caagagactc 60

ctcagtgaga agaagacctg ccagtgctct caccggtttc agaagacatg tagcccccatt 120

<210> 118  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Furin Cleavage site\_CD8 Product 4

<400> 118  
agggctaaac gg 12

<210> 119  
<211> 15  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_CD8 Product 4

<400> 119  
gaattcggct ccgga 15

<210> 120  
<211> 57  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 4

<400> 120  
gccactaact tctccctggt gaaacaggct ggcgatgttg aagaaaaccc cggctct 57

<210> 121  
<211> 78  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
HGH/SS/2\_CD8 Product 4

<400> 121  
atggccaccg gctctagaac aagcctgctg ctcgcttttg gcctgctctg cctcccatgg 60

ctccaagaag gatctgct 78

<210> 122  
<211> 345  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TRB\_VDJ (TCR097) \_CD8 Product 4

<400> 122  
aatgctgggtg tcaactcagac cccaaaattc cgcatcctga agataggaca gagcatgaca 60  
ctgcagtgta cccaggatat gaaccataac tacatgtact ggtatcgaca agaccaggc 120  
atggggctga agctgattta ttattcagtt ggtgctggta tcaactgataa aggagaagtc 180  
ccgaatggct acaacgtctc cagatcaacc acagaggatt tcccgtcag gctggagttg 240  
gctgctccct cccagacatc tgtgtacttc tgtgccagct ccctacaggt tccctacaat 300  
gagcagttct tcgggccagg gacacggctc accgtgctag aggac 345

<210> 123  
<211> 534  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TCR-beta/constant\_CD8 Product 4

<400> 123  
ctgaaaaacg tgttcctcc aaaagtggcc gtgttcgagc cttctgaggc cgagatcagc 60  
cacacacaga aagccacact cgtgtgtctg gctaccggct tctaccccga tcacgtggaa 120  
ctgtcttggg gggtaacgg caaagaggtg cacagcggcg tcagcacaga tccccagcct 180  
ctgaaagaac agcccgtctt gaacgacagc cgctactgcc tgtctagcag actgagagtg 240  
tccgccacct tctggcagaa cccagaaac cacttcagat gccaggcca gttctacggc 300  
ctgagcgaga acgatgagtg gaccaggac agagccaagc ctgtgacaca gatcgtgtct 360  
gccgaagcct ggggcagagc cgattgtggc tttaccagcg agtcatacca gcagggcgtg 420  
ctgtctgcca ccatcctgta tgagatcctg ctcggcaagg ccacactgta cgctgtgctg 480  
gtgtctgctc tgggtgctgat ggctatggtc tcccgggagc gcatccccga ggcc 534

<210> 124  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Furin cleavage site\_CD8 Product 4

<400> 124  
cgggccaagc gg 12

<210> 125  
<211> 9  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
GSG linker\_CD8 Product 4

<400> 125  
ggcagcggc

9

<210> 126  
<211> 57  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Product 4

<400> 126  
gccaccaact tcagcctgct gaagcaggcc ggcgacgtgg aggagaacct cggccct

57

<210> 127  
<211> 78  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
HGH/SS\_CD8 Product 4

<400> 127  
atggccacag gcagcagaac atctctgctg ctggccttcg gactgctgtg tctgccttgg

60

ctgcaagagg gttccgcc

78

<210> 128  
<211> 337  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
Exemplary TRA-VDJ (TCR097) \_CD8 Product 4

<400> 128  
gctcagacag tcaactcagtc tcaaccagag atgtctgtgc aggaggcaga gaccgtgacc

60

ctgagctgca catatgacac cagtgagagt gattattatt tattctggta caagcagcct

120

cccagcaggc agatgattct cgttattcgc caagaagctt ataagcaaca gaatgcaaca

180

gagaatcggt tctctgtgaa cttccagaaa gcagccaaat cttcagtct caagatctca

240

gactcacagc tgggggatgc cgcatgtat ttctgtgctt ttgggaactt caacaaattt

300

tactttggat ctgggaccaa actcaatgta aaaccaa

337

<210> 129  
<211> 74  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
TCR-alpha/constant\_CD8 Product 4

<400> 129  
 atattcagaa ccccgatcct gctgtgtatc agctg'gcgca cagcaagagc agcgacaaga 60  
 gcgtgtgttt gttc 74

<210> 130  
 <211> 200  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 TRAC CDS/right HR arm\_CD8 Product 4

<400> 130  
 accgattttg attctcaaac aaatgtgtca caaagtaagg attctgatgt gtatatcaca 60  
 gacaaaactg tgctagacat gaggtctatg gacttcaaga gcaacagtgc tgtggcctgg 120  
 agcaacaaat ctgactttgc atgtgcaaac gccttcaaca acagcattat tccagaagac 180  
 accttcttcc ccagcccagg 200

<210> 131  
 <211> 800  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Right HR arm\_CD8 Product 4

<400> 131  
 taagggcagc tttggtgcct tcgcaggctg tttccttgct tcaggaatgg ccaggttctg 60  
 cccagagctc tggatcaatga tgtctaaaac tcctctgatt ggtggtctcg gccttatcca 120  
 ttgccaccaa aaccctcttt ttactaagaa acagttagcc ttgttctggc agtccagaga 180  
 atgacacggg aaaaaagcag atgaagagaa ggtggcagga gagggcacgt ggcccagcct 240  
 cagtctctcc aactgagttc ctgcctgcct gcctttgctc agactgtttg ccccttactg 300  
 ctcttctagg cctcattcta agccccttct ccaagttgcc tctccttatt tctccctgtc 360  
 tgccaaaaaa tctttcccag ctactaagt cagtctcag cagtactca ttaaccacc 420  
 aatcactgat tgtgccggca catgaatgca ccagggtgtg aagtggagga attaaaaagt 480  
 cagatgaggg gtgtgccag aggaagcacc attctagttg ggggagccca tctgtcagct 540  
 gggaaaagt caaataactt cagattggaa tgtgttttaa ctgagggtg agaaaacagc 600  
 taccttcagg acaaaagtca gggaagggtc ctctgaagaa atgctacttg aagataccag 660  
 ccctaccaag ggcagggaga ggaccctata gaggcctggg acaggagctc aatgagaaag 720  
 gagaagagca gcaggcatga gttgaatgaa ggaggcaggg ccgggtcaca gggccttcta 780  
 ggccatgaga gggtagacag 800

<210> 132

<211> 6  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Artificial Sequence\_CD8 Product 4

<400> 132  
 gctagc 6

<210> 133  
 <211> 620  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 pBR322\_origin\_CD8 Product 4

<400> 133  
 cgcggttgctg gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg 60  
 ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt tccccctgg 120  
 aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc tgtccgcctt 180  
 tctcccttcg ggaagcgtgg cgcttttctca tagctcacgc tgtaggtatc tcagttcggg 240  
 gtaggtcggt cgctccaagc tgggctgtgt gcacgaacc cccgttcagc ccgaccgctg 300  
 cgcttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact 360  
 ggtagcagcc actggtaaca ggattagcag agcgaggat gtaggcgggtg ctacagagtt 420  
 cttgaagtgg tggcctaact acggctacac tagaagaaca gtatttgta tctgcgctct 480  
 gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaccac 540  
 cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaggatc 600  
 tcaagaagat cctttgatct 620

<210> 134  
 <211> 810  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Kanamycin resistance (KanR2) \_CD8 Product 4

<400> 134  
 ttgaaaaaac tcatcgagca tcaaatgaaa ctgcaattta ttcatatcag gattatcaat 60  
 accatatttt tgaaaaagcc gtttctgtaa tgaaggagaa aactcaccga ggtagttcca 120  
 taggatggca agatcctggg atcggctctgc gattccgact cgtccaacat caatacaacc 180  
 tattaatttc ccctcgtcaa aaataagggt atcaagtggag aatcaccat gaggtagcagc 240  
 tgaatccggt gagaatggca aaagtttatg catttcttc cagacttggt caacaggcca 300

gccattacgc tcgtcatcaa aatcactcgc atcaacaaa ccggtattca ttcgtgattg 360  
 cgctgagcc agacgaaata cgcgatcgcgt gttaaaagga caattacaaa caggaatcga 420  
 atgcaaccgg cgcaggaaca ctgccagcgc atcaacaata ttttcacctg aatcagata 480  
 ttcttctaata acctggaatg ctgtttttcc ggggatcgcga gtggtgagta accatgcatc 540  
 atcaggagta cggataaaat gcttgatggt cggaagaggc ataaattccg tcagccagtt 600  
 tagtctgacc atctcatctg taacatcatt ggcaacgcta cctttgccat gtttcagaaa 660  
 caactctggc gcatcgggct tcccatacaa gcgatagatt gtcgcacctg attgcccgcac 720  
 attatcgcga gcccatttat acccatataa atcagcatcc atggttgaat ttaatcgcgg 780  
 cctcgacgtt tcccgttgaa tatggctcat 810

<210> 135  
 <211> 99  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8 Product 4 Kanamycin promoter

<400> 135  
 aacacccctt gtattactgt ttatgtaagc agacagtttt attggtcatg atgatatatt 60  
 tttatcttgt gcaatgtaac atcagagatt ttgagacac 99

<210> 136  
 <211> 5950  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 Complete CD8 Product 4 construct with TCR97 insertion\_CD8  
 Product 4

<400> 136  
 ggtaccacat taaaaacaca aaatcctacg gaaatactga agaatgagtc tcagcactaa 60  
 ggaaaagcct ccagcagctc ctgctttctg aggggtgaagg atagacgctg tggctctgca 120  
 tgactcacta gactctatc acggccatat tctggcaggg tcagtggctc caactaacat 180  
 ttgtttggta ctttacagtt tattaatag atgtttatat ggagaagctc tcatttcttt 240  
 ctcagaagag cctggctagg aagggtgatg aggcaccata ttcattttgc aggtgaaatt 300  
 cctgagatgt aaggagctgc tgtgacttgc tcaaggcctt atatcgagta aacggtagtg 360  
 ctggggctta gacgcaggtg ttctgattta tagttcaaaa cctctatcaa tgagagagca 420  
 atctcctggg aatgtgatag atttcccaac ttaatgcca cataccataa acctccatt 480  
 ctgctaatac ccagcctaag ttggggagac cactccagat tccaagatgt acagtttgct 540  
 ttgctgggcc tttttccat gcctgccttt actctgccag agttatattg ctggggtttt 600

gaagaagatc ctattaaata aaagaataag cagtattatt aagtagccct gcatttcagg 660  
tttccttgag tggcaggcca ggctggccg tgaacgtca ctgaaatcat ggctcttg 720  
ccaagattga tagcttgtgc ctgtccctga gtcccagtc atcacgagca gctggtttct 780  
aagatgctat ttcccgtata aagcatgaga ccgtgacttg ccagccccac agagccccgc 840  
cctgtccat cactggcatc tggactccag cctgggttg ggcaaagagg gaaatgagat 900  
catgtcctaa ccctgatcct cttgtcccac agatatccag aaccctgacc ctgccgtgta 960  
ccagctgaga gactctaaat ccagtgacaa gtctgtctgc ctattcgaat tcggctccgg 1020  
agccactaac ttcagcctgt tgaagcaggc cggcgacgtt gaggaaaacc ccggtcctat 1080  
ggccttacca gtgaccgct tgcctctgcc gctggccttg ctgctccacg ccgccaggcc 1140  
gagccagttc cgggtgtcgc cgctggatcg gacctggaac ctgggcgaga cagtggagct 1200  
gaagtgccag gtgctgctgt ccaacccgac gtcgggctgc tcgtggctct tccagccgcg 1260  
cggcgccgcc gccagtccca ctttctctc atacctctcc caaaacaagc ccaaggcggc 1320  
cgaggggctg gacaccagc ggttctcggg caagaggttg ggggacacct tcgtcctcac 1380  
cctgagcgac ttccgccgag agaacgaggg ctactatttc tgctcggccc tgagcaactc 1440  
catcatgtac ttcagccact tcgtgccggt cttctgccca gcgaagcca ccacgacgcc 1500  
agcgcgcga ccaccaacac cggcgcccac catcgcgtcg cagcccctgt ccctgcgcc 1560  
agaggcgtgc cggccagcgg cggggggcgc agtgcacacg agggggctgg acttcgcctg 1620  
tgatatctac atctgggcgc cttggccgg gacttgtggg gtccttctcc tgtcactggt 1680  
tatcacctgt gtcagggtcc ggcaccgaag gcgccaagca gagcggatgt ctcatgcaa 1740  
gagactcctc agtgagaaga agacctgcca gtgtcctcac cggtttcaga agacatgtag 1800  
cccattagg gctaaacggg aattcggctc cggagccact aacttctccc tgttgaaca 1860  
ggctggcgat gttgaagaaa accccggtcc tatggccacc ggctctagaa caagcctgct 1920  
gctcgctttt ggctgctct gcctcccatg gctccaagaa ggatctgcta atgctggtgt 1980  
cactcagacc caaaattcc gcatcctgaa gataggacag agcatgacac tgcagtgtac 2040  
ccaggatatg aaccataact acatgtactg gtatcgacaa gaccaggca tggggctgaa 2100  
gctgatttat tattcagttg gtgctggtat cactgataaa ggagaagtcc cgaatggcta 2160  
caacgtctcc agatcaacca cagaggattt cccgctcagg ctggagttgg ctgctccctc 2220  
ccagacatct gtgtacttct gtgccagctc cctacaggtt ccctacaatg agcagttctt 2280  
cgggccaggg acacggctca ccgtgctaga ggacctgaaa aacgtgttcc ctccaaaagt 2340  
ggccgtgttc gagccttctg aggccgagat cagccacaca cagaaagcca cactcgtgtg 2400  
tctggctacc ggcttctacc ccgatcacgt ggaactgtct tggtaggtca acggcaaaga 2460  
ggtgcacagc ggcgtcagca cagatcccca gcctctgaaa gaacagcccg ctctgaacga 2520  
cagccgctac tgctgtctta gcagactgag agtgtccgcc accttctggc agaaccag 2580

aaaccacttc agatgccagg tccagttcta cggcctgagc gagaacgatg agtggacca 2640  
ggacagagcc aagcctgtga cacagatcgt gtctgccgaa gcctggggca gagccgattg 2700  
tggctttacc agcgagtcat accagcaggg cgtgctgtct gccaccatcc tgtatgagat 2760  
cctgctcggc aaggccacac tgtacgctgt gctggtgtct gctctggtgc tgatggctat 2820  
ggtctcccgg gagcgcaccc ccgaggcccg ggccaagcgg ggacagcggc ccaccaactt 2880  
cagcctgctg aagcaggccg gcgacgtgga ggagaacccc ggccctatgg ccacaggcag 2940  
cagaacatct ctgctgctgg ccttcggact gctgtgtctg ccttggtgc aagagggttc 3000  
cgccgctcag acagtcactc agtctcaacc agagatgtct gtgcaggagg cagagaccgt 3060  
gacctgagc tgcacatatg acaccagtga gagtgattat tatttattct ggtacaagca 3120  
gcctcccagc aggcagatga ttctcgttat tcgccaagaa gcttataagc aacagaatgc 3180  
aacagagaat cgtttctctg tgaacttcca gaaagcagcc aaatccttca gtctcaagat 3240  
ctcagactca cagctggggg atgccgcatg gtatttctgt gcttttggga acttcaacia 3300  
atttacttt ggatctggga ccaaactcaa tgtaaaacca aatattcaga accccgatcc 3360  
tgctgtgtat cagctgcgag acagcaagag cagcgacaag agcgtgtgtt tgttcaccga 3420  
ttttgattct caaaciaaat gtgcacaaag taaggattct gatgtgtata tcacagacia 3480  
aactgtgcta gacatgaggt ctatggactt caagagcaac agtgcgtgtg cctggagcaa 3540  
caaatctgac tttgcatgtg caaacgcctt caacaacagc attattccag aagacacctt 3600  
cttcccagc ccaggtaagg gcagctttgg tgccttcgca ggctgtttcc ttgcttcagg 3660  
aatggccagg ttctgcccag agctctggtc aatgatgtct aaaactcctc tgattggtgg 3720  
tctcggcctt atccattgcc accaaaacc tcttttact aagaaacagt gagccttgtt 3780  
ctggcagtcc agagaatgac acgggaaaaa agcagatgaa gagaagggtg caggagaggg 3840  
cacgtggccc agcctcagtc tctcactg agttcctgcc tgcctgcctt tgctcagact 3900  
gtttcccct tactgctctt ctaggcctca ttctaagccc cttctccaag ttgcctctcc 3960  
ttatttctcc ctgtctgcca aaaaatctt cccagctcac taagtcagtc tcacgcagtc 4020  
actcattaac ccaccaatca ctgattgtgc cggcacatga atgcaccagg tgttgaagtg 4080  
gaggaattaa aaagtcagat gaggggtgtg cccagaggaa gcaccattct agttggggga 4140  
gccatctgt cagctgggaa aagtccaaat aacttcagat tggaatgtgt ttttaactcag 4200  
ggttgagaaa acagctacct tcaggacaaa agtcagggaa gggctctctg aagaaatgct 4260  
acttgaagat accagcccta ccaaggcag ggagaggacc ctatagaggc ctgggacagg 4320  
agctcaatga gaaaggagaa gagcagcagg catgagttga atgaaggagg cagggccggg 4380  
tcacagggcc ttctaggcca tgagagggta gacaggctag ccgcttgct ggcgtttttc 4440  
cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 4500

aacccgacag gactataaag ataccaggcg tttccccctg gaagctcct cgtgcgctct 4560  
cctgttccga ccctgccgct taccggatac ctgtccgctt ttctcccttc gggaaagcgtg 4620  
gcgctttctc atagctcacg ctgtaggtat ctgagttcgg tgtaggtcgt tcgctccaag 4680  
ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct gcgccttata cggttaactat 4740  
cgtcttgagt ccaacccggt aagacacgac ttatcgccac tggcagcagc cactggtaac 4800  
aggattagca gagcgaggta tgtaggcggt gctacagagt tcttgaagtg gtggcctaac 4860  
tacggctaca ctagaagaac agtatttggg atctgcgctc tgctgaagcc agttaccttc 4920  
ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggtttt 4980  
ttgttttgca agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc 5040  
tttagaaaaa ctcatcgagc atcaaatgaa actgcaatth attcatatca ggattatcaa 5100  
taccatattt ttgaaaaagc cgtttctgta atgaaggaga aaactcaccg aggcagttcc 5160  
ataggatggc aagatcctgg tatcggtctg cgattccgac tcgtccaaca tcaatacaac 5220  
ctattaatth ccctcgtca aaaataaggt tatcaagtga gaaatcacca tgagtgcga 5280  
ctgaatccgg tgagaatggc aaaagttat gcatttcttt ccagacttgt tcaacaggcc 5340  
agccattacg ctcgatca aaatcactcg catcaacca accgttattc attcgtgatt 5400  
gcgcctgagc cagacgaaat acgcgatcgc tgttaaaagg acaattaca acaggaatcg 5460  
aatgcaaccg gcgcaggaac actgccagcg catcaacaat atttcacct gaatcaggat 5520  
attcttctaa tacctggaat gctgtttttc cggggatcgc agtggtgagt aacctgcat 5580  
catcaggagt acggataaaa tgcttgatgg tcggaagagg cataaattcc gtcagccagt 5640  
ttagtctgac catctcatct gtaacatcat tggcaacgct acctttgcca tgtttcagaa 5700  
acaactctgg cgcatcgggc ttccataca agcgatagat tgtcgcacct gattgcccga 5760  
cattatcgcg agcccattta taccatata aatcagcatc catgttgga tttaatcgcg 5820  
gcctcgacgt ttcccgttga atatggctca taacaccct tgtattactg tttatgtaag 5880  
cagacagttt tattgttcat gatgatatat ttttatcttg tgcaatgtaa catcagagat 5940  
tttgagacac 5950

<210> 137  
<211> 3  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
GSG Linker\_CD8 Products 1, 2, 3, 4

<400> 137  
Gly Ser Gly  
1

<210> 138  
<211> 19  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
P2A\_CD8 Products 1, 2, 3, 4

<400> 138  
Ala Thr Asn Phe Ser Leu Leu Lys Gln Ala Gly Asp Val Glu Glu Asn  
1 5 10 15

Pro Gly Pro

<210> 139  
<211> 21  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD8alpha Signal Peptide\_CD8 Products 1, 2, 3, 4

<400> 139  
Met Ala Leu Pro Val Thr Ala Leu Leu Leu Pro Leu Ala Leu Leu Leu  
1 5 10 15

His Ala Ala Arg Pro  
20

<210> 140  
<211> 161  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
CD8alpha Extracellular Domain\_CD8 Products 1, 2, 3, 4

<400> 140  
Ser Gln Phe Arg Val Ser Pro Leu Asp Arg Thr Trp Asn Leu Gly Glu  
1 5 10 15

Thr Val Glu Leu Lys Cys Gln Val Leu Leu Ser Asn Pro Thr Ser Gly  
20 25 30

Cys Ser Trp Leu Phe Gln Pro Arg Gly Ala Ala Ala Ser Pro Thr Phe  
35 40 45

Leu Leu Tyr Leu Ser Gln Asn Lys Pro Lys Ala Ala Glu Gly Leu Asp  
50 55 60

Thr Gln Arg Phe Ser Gly Lys Arg Leu Gly Asp Thr Phe Val Leu Thr  
65 70 75 80



&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
Furin Cleavage Site\_CD8 Products 1, 2, 3, 4

&lt;400&gt; 143

Arg Ala Lys Arg  
1

&lt;210&gt; 144

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
CD8beta Signal Peptide\_CD8 Product 2

&lt;400&gt; 144

Met Arg Pro Arg Leu Trp Leu Leu Leu Ala Ala Gln Leu Thr Val Leu  
1 5 10 15His Gly Asn Ser Val  
20

&lt;210&gt; 145

&lt;211&gt; 149

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Synthetic  
CD8beta Extracellular Domain\_CD8 Product 2

&lt;400&gt; 145

Leu Gln Gln Thr Pro Ala Tyr Ile Lys Val Gln Thr Asn Lys Met Val  
1 5 10 15Met Leu Ser Cys Glu Ala Lys Ile Ser Leu Ser Asn Met Arg Ile Tyr  
20 25 30Trp Leu Arg Gln Arg Gln Ala Pro Ser Ser Asp Ser His His Glu Phe  
35 40 45Leu Ala Leu Trp Asp Ser Ala Lys Gly Thr Ile His Gly Glu Glu Val  
50 55 60Glu Gln Glu Lys Ile Ala Val Phe Arg Asp Ala Ser Arg Phe Ile Leu  
65 70 75 80Asn Leu Thr Ser Val Lys Pro Glu Asp Ser Gly Ile Tyr Phe Cys Met  
85 90 95Ile Val Gly Ser Pro Glu Leu Thr Phe Gly Lys Gly Thr Gln Leu Ser  
100 105 110

Val Val Asp Phe Leu Pro Thr Thr Ala Gln Pro Thr Lys Lys Ser Thr  
 115 120 125

Leu Lys Lys Arg Val Cys Arg Leu Pro Arg Pro Glu Thr Gln Lys Gly  
 130 135 140

Pro Leu Cys Ser Pro  
 145

<210> 146  
 <211> 21  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8beta Transmembrane Domain\_CD8 Product 2

<400> 146  
 Ile Thr Leu Gly Leu Leu Val Ala Gly Val Leu Val Leu Leu Val Ser  
 1 5 10 15

Leu Gly Val Ala Ile  
 20

<210> 147  
 <211> 19  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD8beta Intracellular Domain\_CD8 Products 2 and 3

<400> 147  
 His Leu Cys Cys Arg Arg Arg Arg Ala Arg Leu Arg Phe Met Lys Gln  
 1 5 10 15

Phe Tyr Lys

<210> 148  
 <211> 40  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 CD4 Intracellular Domain\_CD8 Product 4

<400> 148  
 Cys Val Arg Cys Arg His Arg Arg Arg Gln Ala Glu Arg Met Ser Gln  
 1 5 10 15

Ile Lys Arg Leu Leu Ser Glu Lys Lys Thr Cys Gln Cys Pro His Arg  
 20 25 30

Phe Gln Lys Thr Cys Ser Pro Ile  
35 40