



US00PP14453P39

(12) United States Plant Patent Peterson

(10) Patent No.: **US PP14,453 P3**
(45) Date of Patent: **Jan. 13, 2004**

(54) PAWPAW TREE NAMED 'AIDFIEVATE'

(50) Latin Name: *Asimina triloba (L.) Dunal*
Varietal Denomination: Aidfievate

(76) Inventor: **Robert Neal Peterson**, P.O. Box 1277,
Franklin, WV (US) 26807

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: 09/954,140

(22) Filed: **Sep. 17, 2001**

(65) Prior Publication Data

US 2003/0056253 P1 Mar. 20, 2003

(51) Int. Cl.⁷ **A01H 5/00**

(52) U.S. Cl. **Plt./156**

(58) Field of Search **Plt./156**

Primary Examiner—Bruce R. Campell
Assistant Examiner—Susan B. McCormick

(57) ABSTRACT

Disclosed herein is a new and distinct variety of pawpaw tree, which has been given the name 'Aidfievate.' This variety is distinguished by good yields (atypical of pawpaw), by excellent harvesting characteristics and by good fruit quality. The leaf aspect is unique among pawpaws, being horizontal rather than drooping, which allows fruit to be more easily seen and picked. Fruit display a dependable, readily visible color break at ripeness which speeds picking. The number of fruit per cluster is low, often in singles, which simplifies harvest. Fruit quality is high as fruit are medium to large in size with a low seed-to-fruit ratio, are very consistent in size and shape, and possess a very good flavor. Skin is medium-thick and fruit texture is firm which helps in shipping and handling. This variety is one of three varieties newly identified as having potential to establish a commercial pawpaw industry.

8 Drawing Sheets

1

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct and superior variety of the pawpaw which is of interest for its fruit.

The species—*Asimina triloba (L.) Dunal*—is the largest native edible fruit of North America, a member of the Annonaceae family, and thus a relative of the cherimoya, sweetsop, guanabana and custard apple (*Annona cherimola*, *A. squamosa*, *A. muricata* and *A. reticulata*), all of which are popular fruits that are widely cultivated in tropical and subtropical regions of the world, including southern Florida and southern California. The pawpaw is the only truly temperate member of the Annonaceae, being indigenous to a region stretching from the Great Lakes to the Gulf Coastal Plain and from the Chesapeake Bay to the Great Plains.

Currently, the pawpaw is considered to be semi-domesticated. Native Americans casually cultivated the tree, as did the white settlers who displaced them. The selection, propagation and naming of pawpaw varieties from the wild has been practiced for more than a century, and the backyard cultivation of pawpaws for personal use is not uncommon in Appalachia and parts of the Midwest. Numerous unregistered, unpatented varieties are available in the mail-order nursery trade.

Commercial cultivation of pawpaw has not developed, however. The fruit is fragile and highly perishable which makes transport difficult. And scientific attention towards improving the fruit, its culture and its postharvest handling has received little attention until recently. The lack of high quality cultivars that meet the requirements of producers and consumers is the foremost reason that commercial cultivation has not been undertaken.

The present invention, named 'Aidfievate' by the developer, is the second of three new and distinct varieties of pawpaw, each of which represents a great improvement over existing pawpaw varieties because of higher yields, superior flavor, fewer seeds and firmer flesh. The other two varieties

2

are 'Wansevwan', which is the subject of co-pending application Ser. No. 09/954,536, and 'Levfiv', which is the subject of co-pending application Ser. No. 09/954,186. These varieties possess the minimum overall qualities required for the development of a commercial pawpaw industry. The primary use of this pawpaw variety will be for fresh eating as a dessert fruit. Secondary use will be in processed products such as ice cream, yogurt, juice and cosmetics. Of the three, the variety 'Aidfievate' is outstanding for ease of harvest, and for consistency of size and shape.

ORIGIN

The variety 'Aidfievate' was developed by R. Neal Peterson as the result of a breeding project to improve the pawpaw, which he began in 1980. The project was conducted during a period when Peterson was employed as an economist with the Economic Research Service of the United States Department of Agriculture in Washington, D.C. However, because Peterson was not employed in any capacity as a horticulturist or other researcher in the biological sciences, and no Department of Agriculture plant stock, facilities or information was used, the United States government has no interest in the rights to the claimed variety.

The germplasm for Peterson's breeding project came principally from the surviving remnants of five historic collections of pawpaw dating to the early twentieth century that were the work of the most prominent pawpaw collectors and breeders of the time, and whose named material was no longer propagated or otherwise available. In 1982 germ plasm (open-pollinated seed) was collected from these remnants and included with open-pollinated seed from named cultivars, which became the germ plasm for his own breeding and selection work.

In 1983 the seed was germinated and in the following spring 808 accessions were planted at the University of Maryland experiment station, the Wye Research & Educa-

tion Center, Queenstown, Md. Since then, the orchard has been supervised by the inventor, with basic staff support from the university. Basic tasks such as fertilizing, spraying and mowing were conducted by the station staff. The inventor pruned, weeded, and collected data on growth rates, flowering, fruit set, yields, cluster size, fruit size and fruit quality, including data from taste panels which he organized.

In 1991, the inventor analyzed four years of data, and concluded in identifying eleven trees as superior for further study. Nine of these are involved in regional variety trials around the country and have been termed advanced numbered selections. After nine additional years of observation, the original nine were narrowed to three that were consistently of the highest quality. One of these bears the accession number PPF 8-58 and is the variety that is the subject of this application. The original tree PPF 8-58 is 18 years old. This new variety, named 'Aidflevate,' originated as an open-pollinated seedling from a pawpaw tree on the grounds of the Blandy Experimental Farm, Boyce, Va., which tree was given the label BEF-30 by the inventor. Pawpaw BEF-30 is believed to have been a catalogued accession of the Blandy Experimental Farm's collection of *Asimina triloba* (circa 1922-1955) although records to support this contention are now missing. To our knowledge BEF-30 has never been asexually propagated.

ASEXUAL REPRODUCTION OF THE VARIETY

In 1994, Peterson began topworking the Wye orchard to his eleven advanced numbered selections, plus named varieties. Topworking was accomplished using bark-inlay grafting and chip-budding. Grafting of 'Aidflevate' was easy by either method, and gave a medium-high percentage of success. Replicates of 'Aidflevate' now number more than 15 at the Wye, having been propagated through successive cycles of grafting over seven years. The rootstock for these grafts were mature seedling pawpaw trees growing in the orchard, and were a portion of the original accessions from 1983, described previously. Six grafted trees of 'Aidflevate' have now been in bearing since 1997 and demonstrate that asexual reproduction of this new and distinct variety preserves the desirable characteristics of the variety and establishes and stably transmits those characteristics through successive propagation at the Queenstown location.

SUMMARY OF THE VARIETY

'Aidflevate' is outstanding in its ease of harvesting and excels in vigor, yields, flavor, and small cluster size. The fruit is medium in size, on average 200 gm for well-pollinated fruit, and may exceed 280 gm. It is usually borne in clusters of one, two or three fruits. Fruit size and shape is relatively uniform and consistent as compared to most pawpaws. Fruit shape is typically ovate to nearly round. Skin thickness is medium-thin, slightly thicker than is typical for the species. Skin color is pale green and glaucous, and at the stage for picking exhibits a noticeable color-break as the skin becomes paler and more yellowish. This color break is reliable and can be easily discerned with practice. The composition of the fruit is moderately fleshy, as measured by the seed-to-fruit ratio. The size of seeds is medium. The aroma of the fruit before and after cutting is pleasant and quite mild. The flavor is very good, though not as good as 'Wansevwan'—mild, sweet, free of bitterness or astringency, with a pleasant aftertaste. The pungent asiminoous component that is uniquely pawpaw and that many people find objectionable is weak to absent. The flesh is

firmer than average and very smooth with no detectable fiber or grit, although a certain rubberiness in texture sometimes detracts from an otherwise pleasant mouth-feel. Fruit cracking is a problem on vigorous rootstocks. The ripening period at Queenstown has been mid-season, September 6 to 14, more or less, depending on the degree of heat in the preceding months.

Several non-fruit characters also serve to distinguish 'Aidflevate' from other pawpaws. The aspect of the leaf in full sun is horizontal, a character that is unique to this variety and that contrasts with the typical drooping aspect of pawpaw leaves. This horizontal leaf aspect allows for easy inspection of the tree's interior where the fruit develops, and this simplifies picking. The branching habit is more vertical than is typical of the species. Flower measurements when taken as a whole may also identify 'Aidflevate.' Its flower is borne on a longer peduncle than average, is larger than average, and is more loosely closed in the female stage than is typical for pawpaw. For exact data, see the detailed description and accompanying photographs.

This variety is susceptible to *Talponia plummeriana*, the pawpaw peduncle moth, is susceptible to *Eurytides marcellus*, the zebra swallowtail butterfly, and is believed to be susceptible to pawpaw decline disease, the same in all cases as the species. This variety responds very vigorously to pruning and is difficult to control in size with pruning alone. It should be grafted onto less vigorous rootstocks both to control ultimate tree size and to correct the tendency towards fruit cracking.

BRIEF DESCRIPTION OF THE DRAWINGS

In the accompanying drawings, leaves, flowers, fruits and seeds are shown in color that is as nearly true as is reasonably possible to make in color photographs of this nature. The deep maroon hues of the flowers are the most difficult for photoemulsions to render accurately.

FIG. 1 Shows the fruit of the new variety, hanging on the tree.

FIG. 2 Shows the fruit of the new variety in dissection, revealing the color, fleshiness and seeds.

FIG. 3 Shows the flower of the new variety in the female stage, in profile and dissection.

FIG. 4 Shows the flowers of the new variety in the male stage, in profile and dissection.

FIG. 5 Shows the habit of a pruned tree of the new variety.

FIG. 6 Shows a close-up of the leaves of the new variety.

BOTANICAL DESCRIPTION

Because no variety of *Asimina triloba* has had application made to the Patent Office, we include a botanical description of the species drawn from two taxonomic authorities, C. S. Sargent and R. Kral.

Shrub or small tree 1.5-11 (~14) m tall from a stout, sometimes branched taproot, with a straight trunk seldom exceeding 30 cm in diameter; the bark of older trees gray-brown, shallowly furrowed, and marked with large ash-colored blotches; that of new shoots moderately to copiously dark brown-hairy toward the summit, aging smooth, gray-brown; winter vegetative buds naked, without stipules, acuminate and dark brown to rusty brown-hairy, 2-5 mm long, and tightly appressed against the stem; winter flower buds globose, dark brown-hairy, 2.5-5 mm in diameter; leaves membranaceous, oblong-ovate to lanceolate, 15-30 cm long; apex acute to acuminate; base more or less

gradually attenuate to the short (0.5–1 cm) petiole; margin flat or scarcely revolute; young surfaces sparsely appressed reddish-pubescent above; densely so beneath, becoming glabrous above and sparsely hairy on the veins beneath; flowers green initially, then turning brown to maroon to deep vinous red, 2–5 cm broad with a faintly fetid aroma, on densely dark brown-hairy, nodding peduncles 1–2.5 cm long which develop from the axils of the prominent leaf scars; calyx 8–12 mm long, of three triangular-deltoid sepals which are striate with brown hairs on the outside, glabrous within, outer petals 1.5–2.5 cm long, oblong-elliptic, with ascending bases and slightly to conspicuously recurved tips, copiously appressed-hairy along the veins outside, glabrous and impressed-veiny within; inner petals $\frac{1}{3}$ – $\frac{1}{2}$ the length of the outer, elliptic, saccate-based, recurved tipped, glabrate without, glabrous and impressed-veiny within, with a corrugated nectary zone usually of a lighter color; androecium globular, 0.5–1 cm broad, pale green at anthesis; gynoecium of 3–10 fusiform appressed-red-hairy carpels; fruits oblong-cylindric, 5–15 cm long, yellow-green to brownish when ripe, attached obliquely to the enlarged torus of the peduncle in clusters of variable number; seeds 1.5–2.5 cm long, brown to castaneous, shiny when mature, bean-shaped, somewhat laterally compressed, contained within an aril that is confluent with the pericarp from which the seed readily separates.

DETAILED DESCRIPTION OF THE INVENTION

Following is a detailed description of the new variety of pawpaw tree with color terminology in accordance with The Royal Horticulture Society (R.H.S.) Colour Chart (Ed. No. 2, 1986) except where general color terms of ordinary dictionary significance are used.

Tree:

Size.—Medium. 4.2 m on its own roots but taller on more vigorous rootstock.

Vigor.—High. Approximately 62 cm growth on strong primary laterals under standard fertilization of 50 lbs of N per acre.

Habit.—Rounded with age whether pruned or unpruned.

Branching pattern.—Vertical and spreading with time. Crotch angles vary from narrow to wide. Branch spacing varies from widely to closely spaced.

Apical dominance.—High.

Trunk.—Diameter 13.4 cm measured at 30 cm above ground level.

Bark.—Smooth with small raised horizontal lenticels, typical of the species. Color between R.H.S. Greyed-Green 197B and Grey 201B on both trunk and branches.

Leaf buds:

Length.—Axillary buds growing on branches of normal vigor are small, 2.1 mm average with a range of 1.5 to 2.6 mm. Buds growing on branches of high vigor larger, 2.7 mm average with a range of 2.5 to 2.9 mm.

Leaves:

Shape.—Oblanceolate with acuminate tip and attenuate to the base, as is typical for the species. Average ratio of width to length is 0.444.

Size.—Measurements are from mature leaves attached at midpoint of actively growing shoots of current season's growth. Size is 10.4 cm wide, 23.8 cm long.

Color.—Upper surface R.H.S. Green 147A. Lower surface R.H.S. Yellow-Green 146A. Both colors typical for the species.

Aspect.—Horizontal or nearly so. Leaves on the vertical shoots of the upper portion of the tree curled in a strongly concave fashion. Both features very atypical of the species.

Margin.—Entire, and slightly undulating which is atypical of the species.

Texture.—Smooth to only slightly impressed veiny, typical for the species.

Arrangement.—Alternate opposite.

Petiole.—Color yellow-green similar to lower leaf surface. Average diameter 3.3 mm at the petiole midpoint. Length ranges from 16.7 to 18.9 mm with an average of 17.9 mm. NOTE: The structure of the pawpaw leaf lamina is such, in the way that it tapers gradually to the petiole, that visual demarcation of the petiole terminus is imprecise and the resultant measurement of petiole length has a high margin of error. The measurements reported here are based on a precise tactile method. By means of running the back of one's thumbnail down the midvein (on the underside of the leaf) towards the petiole one encounters a sudden change in curvature, diameter, and hardness. This point of change is the petiole terminus and is easy to detect and replicate.

Flower buds:

Size.—Length 4.9 mm average. Width 3.6 mm average.

Appearance.—Very dark brown, between R.H.S. Brown 200A and Black 202A. Surface densely pubescent, velvety.

Flower:

Size.—Large. As the flower matures from female to male stage, the petals reflex, and therefore measured flower size is specific to flower stage, which data are presented below.

Outer petals.—Average of 28 mm wide, 35 mm long. Average ratio of width to length 0.80.

Inner petals.—Average of 11 mm wide, 19 mm long. Average ratio of width to length 0.58.

Color.—Maroon.

Form of female stage flower.—More loosely closed than typical.

Size of the female stage flower.—Diameter 40 mm average.

Nectary band of female stage flower.—Reddish-maroon color, not as dark as surrounding inner petal.

Form of male stage flower.—Outer and inner petals flaring and recurved, in typical fashion for the species.

Size of the male stage flower.—Diameter 53 mm average.

Nectary band of male stage flower.—A deep maroon color, basically indistinguishable from the inner petal color.

Peduncle.—Average length 20 mm. Lightly pubescent with brown hairs.

Fruit:

Dates of picking.—Mid-season, generally from September 5 to 14 at Queenstown, but dates can vary by more than a week depending on degree of summer heat.

Cluster size.—Few fruited. One, two and three-fruited clusters are most prevalent, and this small cluster size is atypical of pawpaw. Larger clusters also

occur. Fruit tends towards a uniform size, atypical of the species.

Fruit shape.—Ovate to nearly round. Shape varies considerably; but large well-formed fruit in single-fruited cluster tend towards length-to-width-to-depth proportions of 100:70:70.

Fruit size.—Medium, 200 gm average for well-pollinated fruit. Larger than typical pawpaw. Size varies from small (~23 gm) to large (~280 gm), dependent on the number of fertilized seed present in the fruit. Lengths normally range 4.4–10.8 cm, widths 3.1–7.6 cm, and depths 3.1–7.6 cm.

Suture.—Typically absent to indistinct.

Abscission type.—Primary mode is for peduncle to abscise from the branch remaining attached to the fruit (for single-fruited clusters). Abscission may also occur at the point of peduncle attachment to the fruit.

Abscission scar.—Medium, 9.2 mm in diameter. A character of low variance.

Peduncle.—Length 24.3 mm on average, ranging from 21.0 to 29.0 mm. Diameter extremely variable, varying in proportion to the cluster fruit mass, as is typical of the species. In comparison to the range of peduncle sizes of pawpaw cultivars, and relative to the total cluster fruit weight, the diameter is medium. Peduncle pubescence rufus, medium density. The point of peduncle attachment to single-fruited clusters is aligned with the central axis of the fruit, or only slightly offset to the dorsal side.

Skin color.—Under-ripe fruit pale green and glaucous, R.H.S. Yellow-Green 146D. Ripe fruit pale yellowish green and glaucous, R.H.S. Yellow-Green 151A. After the fruit passes its climacteric peak and falls from the tree, brown blotches appear which spread to cover the entire skin surface, as is typical for the species.

Color break.—Subtle but reliable color change at the time when the fruit may be picked. Exact color change noted in preceding paragraph.

Skin thickness.—Medium-thick.

Aril.—Thin, tender, and edible.

Fleshiness.—Moderately high. Percent seed is 5.7 on average. The average quantity of pulp per seed is 19.3 gm.

Flesh color.—Creamy yellow and uniform throughout. Color ranges from R.H.S. Yellow 13C to Yellow-Orange 15C.

Aroma of uncut fruit.—Pleasant. Low power.

Aroma of cut fruit.—Pleasant. Low power.

Flavor.—Very sweet, Brix 19%, mild, nonbitter, and nonastringent. Pungent asimino component low to absent.

Aftertaste.—Pleasant, long lingering, no negative components.

Consistency.—Pleasant mouth-feel. Flesh is firm (atypical of pawpaw) and smooth but sometimes with a slightly rubbery texture. No detectable fiber or grit.

Use.—Principally for fresh eating as a dessert fruit. Secondary in processed products.

Seed:

Size.—Medium. 1.33 gm average weight. Dimensions 25.2 mm long, 14.0 mm wide, 7.1 thick on average.

Color.—Dark brown. R.H.S. Brown 200A.

Number per fruit.—9 per average fruit of 210 gm. As fruit size varies greatly in pawpaw, the seed number per fruit is not a stable character, unlike the seed-to-fruit ratio (percent seed) which is stable.

Physiological and ecological characters:

Graftability.—Very easy to graft by virtually all methods. Percent take is medium to high for pawpaw, in vicinity of 85–95%.

Habit of tree after grafting.—Vigorous. Time to flowering on established rootstock is roughly three years.

It appears that common seedling rootstocks have more vigor than the tree's own roots.

Pruning.—The tree responds vigorously to pruning, sometimes with too much vegetative vigor, depending on rootstock. Flowering and general vigor increased by pruning.

Flower count.—Medium to high. Average of 5.3 blossoms per branch on vigorous branches, but flower number varies considerably depending on the vigor of the branch.

Self-fruitfulness.—Requires cross pollination.

Bearing.—Annual and consistent.

Fruit set.—Low, less than 20 percent in most years.

Yields.—High. From mature trees approximately 30 lbs. of fruit per tree. At orchard densities of 330 trees per acre, this is 10,000 lbs per acre.

Keeping quality of fruit (normal storage, 24° C.).—Short. Three days when ripe. Typical for the species.

Keeping quality of fruit (cold storage, 2° C.).—Moderate. Three weeks when picked at the proper stage of color break.

Shipping quality of fruit.—Medium if shipped refrigerated with adequate cushioning. Poor otherwise because of the rapid ripening which is typical for pawpaw.

Drought and heat tolerance.—Good, similar to the species. Drought tolerance superior to peach.

Hardiness.—Fully hardy at Queenstown location, USDA Zone 7. Believed hardy to Zone 5 (average annual minimum temperature –30° C.), the same as for the species.

*Resistance to *Talponia plummeriana**.—Susceptible.

*Resistance to *Eurytides marcellus**.—Susceptible.

Resistance to pawpaw decline disease.—Believed to be susceptible.

Variance in botanical details: The pawpaw tree and its fruit described herein will vary due to climate, soils, growing conditions and culture.

I claim:

1. A new and distinct variety of pawpaw tree obtained as an open-pollinated seedling of BEF-30 (unpatented), substantially as shown and described herein.

* * * * *



FIG 1




FIG 2



FIG 3A




FIG 3B




FIG 4A




FIG 4B



FIG 5



FIG 6