



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/04/10
(87) Date publication PCT/PCT Publication Date: 2019/10/17
(85) Entrée phase nationale/National Entry: 2020/10/09
(86) N° demande PCT/PCT Application No.: NL 2019/050212
(87) N° publication PCT/PCT Publication No.: 2019/199165
(30) Priorité/Priority: 2018/04/10 (EP18166498.8)

(51) Cl.Int./Int.Cl. *C07K 14/705* (2006.01),
A61K 35/17 (2015.01), *A61K 48/00* (2006.01),
C07K 19/00 (2006.01), *C12N 15/62* (2006.01),
C12N 5/0783 (2010.01)
(71) Demandeur/Applicant:
STICHTING SANQUIN BLOEDVOORZIENING, NL
(72) Inventeur/Inventor:
AMSEN, DERK, NL
(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : RECEPTEURS NOTCH CHIMERIQUES
(54) Title: CHIMERIC NOTCH RECEPTORS

(57) **Abrégé/Abstract:**

The invention relates to chimeric receptors comprising an intracellular domain, and transmembrane domain of a Notch receptor and a heterologous extracellular ligand-binding domain and to uses thereof, specifically in improving T cell function and/or T cell survival, more particularly in cancer therapy.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)**(19) World Intellectual Property****Organization**

International Bureau

(43) International Publication Date

17 October 2019 (17.10.2019)

**(10) International Publication Number****WO 2019/199165 A1****(51) International Patent Classification:**

C07K 14/705 (2006.01) C12N 5/0783 (2010.01)

C07K 19/00 (2006.01) A61K 35/17 (2015.01)

C12N 15/62 (2006.01) A61K 48/00 (2006.01)

(21) International Application Number:

PCT/NL2019/050212

(22) International Filing Date:

10 April 2019 (10.04.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

18166498.8 10 April 2018 (10.04.2018) EP

(71) Applicant: STICHTING SANQUIN BLOED-VOORZIENING [NL/NL]; Plesmanlaan 125, 1066 CX Amsterdam (NL).

(72) Inventor: AMSEN, Derk; c/o Plesmanlaan 125, 1066 CX Amsterdam (NL).

(74) Agent: JANSEN, C.M.; P.O. Box 87930, 2508 DH Den Haag (NL).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: CHIMERIC NOTCH RECEPTORS

(57) Abstract: The invention relates to chimeric receptors comprising an intracellular domain, and transmembrane domain of a Notch receptor and a heterologous extracellular ligand-binding domain and to uses thereof, specifically in improving T cell function and/or T cell survival, more particularly in cancer therapy.



WO 2019/199165 A1

Title: Chimeric Notch receptors

Field of the invention

5 The invention relates to the field of therapy, specifically cancer therapy, more specifically adoptive T cell immunotherapy.

Background

Remarkable successes have been obtained in tumor therapy by adoptive
10 transfer of *in vitro* expanded Tumor Infiltrating Lymphocytes (TIL) or T cells expressing chimeric antigen receptors (CAR). CARs contain an ectodomain (a portion of an antibody) specific for antigens found on tumors, coupled to the signaling domains of CD3 ζ and a costimulatory receptor, such as CD28 or 4-1BB (Figure 1). Expression of CARs in T cells leads to their activation by tumor
15 antigens. Up to 90% complete remissions have been obtained with CAR T cells in certain hematological malignancies. Much less success has been obtained in the treatment of solid tumors. Hence, still many patients are not cured by such treatments. Major hurdles are the suboptimal persistence of transferred T cells and blockade of T cell function by multiple inhibitory receptors (a phenomenon known
20 as exhaustion), which must all be targeted for maximal therapeutic effect. Ideally, anti-tumor T cells would be broadly impervious to suppressive mechanisms and live long enough to achieve complete tumor eradication.

Notch is a cell surface receptor that responds to membrane bound
ligands. It signals through a strikingly direct pathway, in which the intracellular
25 domain is cleaved off from the plasma membrane by a γ -secretase and migrates to the nucleus to act as a transcription factor (Figure 2). Notch is a major regulator of both CD4 and CD8 T cell effector differentiation. It also promotes long term survival of CD4 memory T cells as well of Tissue Resident Memory CD8 T cells, which are emerging as the most effective T cell type against solid tumors.
30 Furthermore, Notch is a major regulator of the CD8 effector T cell gene expression program. Among its direct target genes are those encoding IFN γ , Granzyme B and Perforin, as well as the transcription factors T-bet and Eomesodermin. Mice with T cell specific deficiencies in the Notch pathway are unable to reject model tumors.

Vice versa, deliberate activation of Notch promoted tumor rejection in mice. Tumor associated myeloid-derived suppressor cells (MDSC) downregulate Notch expression in T cells, presumably helping tumors escape effective T cell-mediated rejection. Expression of an active Notch allele rendered CD8 T cells insensitive to MDSC mediated suppression.

Recent studies (Morsut et al. 2016 and Roybal et al. 2016) created chimeric receptors containing the transmembrane region and a small part of the extracellular region of Notch. These were coupled to ligand-binding domains from unrelated surface receptors, while the intracellular part of Notch was replaced by an unrelated transactivator (Gal4). Ligand binding by these receptors resulted in γ -secretase mediated release of Gal4, which then activated transcription of artificial response genes. Hence, in these receptors both the intracellular effector domain of Notch and the extracellular ligand-binding domain of Notch, and consequently Notch signaling, are no longer present.

There remains a need in the art for new compositions and methods for immunotherapy of tumors, either or not to be used in combination with existing immunotherapy.

Summary of the invention

It is an object of the invention to provide methods for improving T cell function in general, and specifically in tumor immunotherapy.

The invention therefore provides a chimeric receptor comprising an intracellular domain and transmembrane domain of a Notch receptor and a heterologous extracellular ligand-binding domain. The chimeric receptor further preferably comprises a heterodimerization domain and a Lin-12-Notch (LNR) repeats domain of the Notch receptor.

The chimeric receptor according to the invention is capable of Notch signaling, preferably Notch1, Notch2, Notch3 and/or Notch4 signaling, more preferably Notch1 and/or Notch2 signaling, when the heterologous extracellular ligand-binding domain is bound a ligand.

In a further aspect, the invention provides a nucleic acid molecule comprising a sequence encoding a chimeric receptor according to the invention.

In a further aspect, the invention provides a vector comprising a nucleic acid molecule according to the invention.

In a further aspect, the invention provides an isolated cell comprising the nucleic acid molecule according to the invention. In a further aspect, the invention provide a population of such cells.

In a further aspect, the invention provides an isolated cell expressing a chimeric receptor according to the invention. In a further aspect, the invention provide a population of such cells.

In a further aspect, the invention provides a genetically modified T lymphocyte, which is transduced by the nucleic acid molecule or vector of the invention.

In a further aspect, the invention provides a pharmaceutical composition comprising a nucleic acid molecule, vector or cell according the invention and a pharmaceutically acceptable carrier, diluent or excipient.

In a further aspect, the invention provides a method for improving T cell function and/or T cell survival in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a chimeric receptor, a nucleic acid molecule, a vector or a cell according to the invention.

In a further aspect, the invention provides a chimeric receptor, a nucleic acid molecule, a vector or a cell according to the invention for use in a method for improving T cell function and/or T cell survival in a subject.

In a further aspect, the invention provides a method of immunotherapy in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a chimeric receptor, a nucleic acid molecule, a vector or a cell according to the invention.

In a further aspect, the invention provides a chimeric receptor, a nucleic acid molecule, a vector or a cell according to the invention for use in therapy, preferably immunotherapy.

In a further aspect, the invention provides a method for enhancing efficacy of an antibody-based immunotherapy in a subject suffering from cancer and being treated with said antibody, the method comprising administering to the subject a therapeutically effective amount of T cells expressing the chimeric receptor according to the invention.

In a further aspect, the invention provides T cells expressing a chimeric receptor according to the invention for use in a method for enhancing efficacy of an antibody-based immunotherapy in a subject suffering from cancer and being treated with said antibody.

5 In a further aspect, the invention provides a method of treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of T cells comprising a nucleic acid sequence encoding the chimeric receptor according to the invention.

10 In a further aspect, the invention provides T cells comprising a nucleic acid sequence encoding the chimeric receptor according to the invention for use in a method of treating cancer in a subject.

In a further aspect, the invention provides a method of producing a population of cells according to the invention, comprising

- providing cells, preferably human T-cells,
- 15 - providing said cells with a nucleic acid molecule or vector according to the invention, and
- allowing expression of the chimeric antigen receptor according to the invention.

Detailed description

20 The present invention is concerned with a chimeric receptor with functioning Notch signaling following ligand binding which receptor is created from a combination of the intracellular effector and transmembrane domains of Notch and a heterologous extracellular ligand binding domain. The present inventors found that Notch signaling suppresses expression of T cell specific inhibitory

25 receptors such as PD1 (programmed death protein 1) and LAG3 (lymphocyte activation gene 3) on T cells. Tumors often escape immune destruction by reducing the anti-tumor T cell response through upregulation of such inhibitory molecules. Therefore, therapeutic activation of Notch is an attractive target to enhance T cell responses against tumors in human patients. So far, therapeutic use of Notch has

30 been precluded by two problems. First, Notch functions in many cell types and its systemic activation is likely to elicit many side effects. Second, excessive Notch signaling can be oncogenic. Now that the present inventors found that Notch signaling is maintained when combining the intracellular effector domain of Notch

with a heterologous extracellular binding domain, these drawbacks are avoided because activation of Notch signaling can be regulated, both in time and location in the body. This is because the chimeric receptor of the invention responds to a heterologous ligand of choice. In the examples the preparation of a chimeric Notch
5 receptor consisting of an ScFv antibody domain directed against human CD19 fused to the 5'end of the human NOTCH1 protein is described.

Hence, the invention provides a chimeric receptor comprising an intracellular domain, and transmembrane domain of a Notch receptor and a
10 heterologous extracellular ligand-binding domain. The chimeric receptor further preferably comprises a heterodimerization domain and a Lin-12-Notch (LNR) repeats domain of the Notch receptor.

The Notch receptors Notch1, Notch2, Notch3 and Notch4 and their
15 sequences are well known in the art, as well as the different domains in these receptors and their sequence, including the Notch intracellular domain, transmembrane domain, heterodimerization domain, Lin-12-Notch (LNR) repeats domain and negative regulatory region (NRR). Hence, a skilled person is well capable of selecting the appropriate domain when making or using a chimeric
20 receptor according to the invention.

An "intracellular domain of a Notch receptor" as used herein refers to an intracellular domain that is capable of initiating Notch1, Notch2, Notch3 or Notch4 signaling, preferably Notch1 or Notch2 signaling. The chimeric receptor
25 according to the present invention is thus capable of Notch signaling, preferably Notch1, Notch2, Notch3 and/or Notch4 signaling, more preferably Notch1 and/or Notch2 signaling. Notch signaling, preferably Notch1, Notch2, Notch3 and/or Notch4 signaling, more preferably Notch1 and/or Notch2 signaling, is induced when the heterologous extracellular ligand-binding domain is bound a ligand.
30 Hence, "capable of Notch signaling" means that Notch signaling is induced when the heterologous extracellular ligand-binding domain of the chimeric receptor is bound a ligand. The Notch intracellular domain is well known to a person skilled in the art. Preferably it comprises the Notch intracellular domain (NICD), this is the

domain that is cleaved of by γ -secretase after ligand binding to the Notch extracellular domain of an intact Notch receptor, preferably the NICD of Notch1 or Notch2, more preferably of human Notch1, or a Notch signaling pathway initiating part of the NICD. Said part is capable of initiating Notch signaling. The chimeric
5 receptor furthermore in a preferred embodiment comprises the entire intracellular domain of Notch1, including the C-terminal transactivation domain, the RAM domain and the ankyrin repeats.

The NICD can be used including or lacking the C-terminal PEST region. Truncation of this region results in a more stable NICD protein, which elicits
10 stronger and more sustained signals. Hence, in a particularly preferred embodiment, the intracellular domain of a Notch receptor comprises a sequence of amino acids 1744 to 2424 of the sequence shown in figure 8, or the corresponding sequence of a Notch receptor other than Notch 1, or a sequence that is at least 90% identical to said sequence. Said sequence is preferably capable of initiating Notch
15 signaling. Said sequence is preferably at least 95% identical to amino acids 1744 to 2424 of said sequence shown in figure 8, more preferably at least 97%, more preferably at least 98%, more preferably at least 99%. In a particularly preferred embodiment, the intracellular domain of a Notch receptor comprises amino acids 1744 to 2424, of the sequence shown in figure 8, more preferably it consists of
20 amino acids 1744 to 2424 of the sequence shown in figure 8. It is preferred that the intracellular domain comprises the indicated sequence of Notch1, and thus amino acids 1744 to 2424, of the sequence shown in figure 8.

In another preferred embodiment, the entire NICD is used, and the intracellular domain of a Notch receptor comprises a sequence of amino acids 1744
25 to 2555 of the sequence shown in figure 8, or the corresponding sequence of a Notch receptor other than Notch 1, or a sequence that is at least 90% identical to said sequence. Said sequence is preferably capable of initiating Notch signaling. Said sequence is preferably at least 95% identical to amino acids 1744 to 2555 of said sequence shown in figure 8, more preferably at least 97%, more preferably at least
30 98%, more preferably at least 99%. In a particularly preferred embodiment, the intracellular domain of a Notch receptor comprises amino acids 1744 to 2555, of the sequence shown in figure 8, more preferably it consists of amino acids 1744 to 2555 of the sequence shown in figure 8. It is preferred that the intracellular

domain comprises the indicated sequence of Notch1, and thus amino acids 1744 to 2555 of the sequence shown in figure 8.

A “transmembrane domain” (TMD) of a Notch receptor” as used herein
5 refers to a transmembrane domain of Notch1, Notch2, Notch3 or Notch4,
preferably of Notch1 or Notch2. The Notch transmembrane domain is well known
to a person skilled in the art. In a particularly preferred embodiment, the
transmembrane domain of a Notch receptor comprises a sequence of amino acids
1736 to 1743 of the sequence shown in figure 8, or the corresponding sequence of a
10 Notch receptor other than Notch 1, or a sequence that is at least 90% identical to
said sequence. Said sequence is preferably capable of initiating cleavage of the
NICD by a γ -secretase. Said sequence is further preferably at least 95% identical to
amino acids 1736 to 1743 of said sequence shown in figure 8, more preferably at
least 97%, more preferably at least 98%, more preferably at least 99%. In a
15 particularly preferred embodiment, the transmembrane domain of a Notch receptor
comprises amino acids 1736 to 1743 of the sequence shown in figure 8, more
preferably it consists of amino acids 1736 to 1743 of the sequence shown in figure 8.
It is preferred that the TMD comprises the indicated sequence of Notch1, and thus
amino acids 1736 to 1743 of the sequence shown in figure 8.

20

The heterodimerization domain and Lin-12-Notch (LNR) repeats
domain of a Notch receptor together form the negative regulatory region (NRR) of
the receptor. The Notch LNR domain, heterodimerization domain and NRR are
well known to a person skilled in the art. The heterodimerization domain and the
25 LNR repeats are located between the heterologous extracellular ligand-binding
domain and the transmembrane domain in a chimeric receptor of the invention.
The order of domains is preferably the following: heterologous extracellular ligand-
binding domain - LNR domain - heterodimerization domain - transmembrane
domain. Canonical Notch signaling is initiated when a ligand binds to the Notch
30 receptor. This leads to ADAM metalloprotease mediated cleavage of the
extracellular fragment of the heterodimer. The membrane tethered fragment is
then cleaved by a γ -secretase to release the intracellular fragment of Notch (NICD).
The heterodimerization domain and the LNR domain are located in the NRR of the

Notch receptor, which is located between the ligand binding domain and the transmembrane domain. The LNRs participate in maintaining the receptor in resting conformation, i.e. prevent or inhibit cleavage by ADAM metalloprotease, in the absence of ligand binding. In a preferred embodiment, the chimeric receptor
5 comprises the entire negative regulatory region (NRR) of the Notch receptor. Preferably this NRR comprises amino acids 1447 to 1735 of the sequence shown in figure 8, or the corresponding sequence of a Notch receptor other than Notch 1, or a sequence that is at least 90% identical to said sequence. Said sequence is further preferably at least 95% identical to amino acids 1447 to 1735 of said sequence
10 shown in figure 8, more preferably at least 97%, more preferably at least 98%, more preferably at least 99%. In a further preferred embodiment this NRR comprises amino acids 1396 to 1735 of the sequence shown in figure 8 or the corresponding sequence of a Notch receptor other than Notch 1, or a sequence that is at least 90% identical to said sequence. Said sequence is further preferably at least 95%
15 identical to amino acids 1447 to 1735 of said sequence shown in figure 8, more preferably at least 97%, more preferably at least 98%, more preferably at least 99%. In this sequence, the extracellular portion of the Notch sequence is extended up till proline 1396 (see Figure 8), as this yields a receptor that is more reliably silent in the absence of ligand binding than shorter constructs. The chimeric
20 receptor of the invention further optionally comprises a signal peptide that directs the receptor to the cell membrane. It is preferred that the NRR comprises the indicated sequence of Notch1, and thus amino acids 1447 to 1735 or 1396 to 1735 of the sequence shown in figure 8.

25 In a particularly preferred embodiment, a chimeric receptor of the invention comprises an intracellular domain, a transmembrane domain, a heterodimerization domain and a Lin-12-Notch (LNR) repeats domain of a Notch receptor and a heterologous extracellular ligand-binding domain, preferably in the indicated order. Hence, a preferred chimeric receptor of the invention comprises
30 amino acids 1447 to 2424 of the sequence shown in figure 8, or the corresponding sequence of Notch receptor other than Notch 1. In a further particularly preferred embodiment, a chimeric receptor of the invention comprises amino acids 1447 to 2555 of the sequence shown in figure 8, or the corresponding sequence of Notch

receptor other than Notch 1. In a further particularly preferred embodiment, a chimeric receptor of the invention comprises amino acids 1396 to 2424 of the sequence shown in figure 8, or the corresponding sequence of Notch receptor other than Notch 1. In a further particularly preferred embodiment, a chimeric receptor
5 of the invention comprises amino acids 1396 to 2555 of the sequence shown in figure 8, or the corresponding sequence of Notch receptor other than Notch 1. It is preferred that the chimeric receptor comprises said sequences of Notch1, and thus of the sequence shown in figure 8.

10 The term “heterologous ligand-binding domain” as used herein refers to a domain other than the ligand-binding domain of a Notch receptor, i.e. a domain other than the extracellular-ligand binding domain of Notch1, Notch2, Notch3 or Notch4. The heterologous ligand-binding domain can be any domain that can be bound by a ligand of choice. In particular, the ligand-binding domain can be the
15 binding partner of any cell surface antigen or any soluble ligand. The versatility in the heterologous ligand-binding domain allows to select an appropriate ligand for any specific application. This way, activation of Notch signaling by the chimeric receptor of the invention can be induced at a selected time, a selected location / cell type, or both. Preferred examples of suitable extracellular ligand-binding domains
20 are a ligand binding domain specific for a soluble ligand, a ligand binding domain specific for a cell surface antigen and a combination thereof. More preferred examples are:

- an antibody or antigen binding part of an antibody, such as a single chain variable fragment (scFv), specific for a cell surface antigen;
- 25 • an antibody or antigen binding part of an antibody, such as a single chain variable fragment (scFv), specific for an epitope in an antibody, a Fab fragment, a F(ab)2 fragment directed against a cell surface antigen;
- an extracellular Fc-binding domain of an Fc receptor or a ligand-binding fragment thereof,
- 30 • an extracellular domain that comprises an epitope for an antibody that can crosslink the chimeric receptor without involvement of a surface molecule;
- an extracellular domain that comprises a moiety, such as biotin, that can be crosslinked by an agent with multiple binding sites for that moiety, such as

streptavidin (resulting in clustering of multiple chimeric receptors upon addition of said agent).

In principle the following types of surface antigens can be used in accordance with the invention:

1. tumor specific antigens;
2. antigens that have a higher level of expression on tumor cells as compared to the expression level on non-tumor cells;
3. antigens that are expressed on both tumor cells and non-tumor cells, but where activation of T cells expressing the chimeric receptor of the invention induced by non-tumor cells results in side-effects that are acceptable, such as CD19 and CD20;
4. antigens that are expressed on both tumor cells and non-tumor cells, but that are specific for tumor cells in combination with one or more other antigens, such as a T cell epitope; and
5. antigens expressed on cells surrounding a tumor, such as PDL1 and PDL2.

In a preferred embodiment, a cell surface antigen is a tumor antigen and the heterologous extracellular ligand-binding domain is an antibody or antigen binding part of an antibody specific for said tumor antigen. Preferred examples of tumor antigens are TAG-72, calcium-activated chloride channel 2, 9D7, Ep-CAM, EphA3, Her2/neu, mesothelin, SAP-1, BAGE family, MC1R, prostate-specific antigen, CML66, TGF- β RII, MUC1, CD5, CD19, CD20, CD30, CD33, CD47, CD52, CD152 (CTLA-4), CD274 (PD-L1), CD273 (PD-L2) CD340 (ErbB-2), GD2, TPBG, CA-125, MUCL, immature laminin receptor and ErbB-1.

A skilled person is well capable of identifying soluble ligand and their binding partners that can be used in a chimeric antigen receptor according to the invention. Examples of suitable soluble ligands are antibodies directed against an epitope in the extracellular domain of the chimeric Notch receptor or molecules such as streptavidin in combination with biotinylated extracellular domains of the chimeric Notch receptor. A combination of a ligand binding domain specific for a soluble ligand and a ligand binding domain specific for a cell surface antigen is also possible. In that case Notch signaling will only be induced if both the soluble ligand

and the cell surface antigen are present. For instance, an ectodomain can consist of an antibody to a peptide neo-epitope or to a Biotin or FITC moiety that is itself incorporated in another antibody (a "switch" antibody) directed to a surface antigen on a tumor. As a consequence, activation of the Chimeric Notch receptor will only occur if, in addition to the cell surface antigen targeted by the switch antibody, the switch antibody itself is also present. This set up is described in Ma et al 2016, which is incorporated herein by reference, and permits temporary control of the receptor (turning it on and off only when desired) as well as quantitative control (by in- or decreasing the concentration of the switch antibody).

10

The chimeric receptor of the invention further optionally comprises a linking sequence located between the transmembrane domain and the heterologous extracellular ligand-binding domain. Such linking sequence preferably comprises up to 30 amino acids, such as 2, 3, 4, 5, 6, 7, 8, 9 or 10 amino acids.

15

The percentage of identity of an amino acid sequence or nucleic acid sequence, or the term "% sequence identity", is defined herein as the percentage of residues of the full length of an amino acid sequence or nucleic acid sequence that is identical with the residues in a reference amino acid sequence or nucleic acid sequence after aligning the two sequences and introducing gaps, if necessary, to achieve the maximum percent identity. Methods and computer programs for the alignment are well known in the art, for example "Align 2".

20

In amino acid sequences as depicted herein amino acids are denoted by single-letter symbols. These single-letter symbols and three-letter symbols are well known to the person skilled in the art and have the following meaning: A (Ala) is alanine, C (Cys) is cysteine, D (Asp) is aspartic acid, E (Glu) is glutamic acid, F (Phe) is phenylalanine, G (Gly) is glycine, H (His) is histidine, I (Ile) is isoleucine, K (Lys) is lysine, L (Leu) is leucine, M (Met) is methionine, N (Asn) is asparagine, P (Pro) is proline, Q (Gln) is glutamine, R (Arg) is arginine, S (Ser) is serine, T (Thr) is threonine, V (Val) is valine, W (Trp) is tryptophan, Y (Tyr) is tyrosine.

25

30

As used herein the terms "specific for" and "specifically binds" or "capable of specifically binding" refer to the non-covalent interaction between a ligand and a ligand-binding domain, such as an antibody or an antigen binding

part thereof and its antigen or a soluble ligand and its binding partner. It indicates that the ligand preferentially binds to said ligand-binding domain over other domains.

An “antigen binding part of an antibody” is defined herein as a part of
5 an antibody that is capable of specifically binding the same antigen as the antibody, although not necessarily to the same extent. The part does not necessarily need to be present as such in the antibody and includes different fragments of the antibody that together are capable of binding the antigen, such as a single-chain variable fragment (ScFv), a fusion protein of the variable regions of
10 the heavy and light chains of an antibody.

A “cell surface antigen” as used herein refers to an antigen or molecule that is expressed at the extracellular surface of a cell.

As used herein “tumor antigen” refers to an antigen expressed on cells of a tumor. A tumor antigen is also referred to as a tumor-associated antigen
15 (TAA).

A “soluble ligand” as used herein refers to a water-soluble ligand for which a binding partner can be used as extracellular domain of the chimeric receptor of the invention. It is preferred that the soluble ligand can be administered to a subject, e.g. by injection, such as intravenous injection, or orally.
20

Also provided is a nucleic acid molecule comprising a sequence encoding a chimeric receptor according to the invention. Also provided is a vector comprising the nucleic acid molecule according to the invention. In a preferred embodiment, the vector is a viral vector, e.g., a lentiviral vector or a retroviral vector. In another
25 preferred embodiment, the vector comprises or is a transposon. Said nucleic acid molecule or vector may additionally comprise other components, such as means for high expression levels such as strong promoters, for example of viral origin, that direct expression in the specific cell in which the vector is introduced, and signal sequences. In a preferred embodiment, the nucleic acid molecule or vector
30 comprises one or more of the following components: a promoter that drives expression in T cells, such as the EF1a promoter or the 5' LTR of MSCV, a C-terminal signal peptide such as from the GM-CSF protein or the CD8 protein for targeting to the plasma membrane and a polyadenylation signal.

Also provided is an isolated cell, comprising the nucleic acid molecule or vector according to the invention. The isolated cell is preferably an immune cell, such as natural killer cell, macrophage, neutrophil, eosinophil, or T cell. The nucleic acid molecule or vector may be introduced into the cell, preferably immune cells, by any method known in the art, such as by lentiviral transduction, retroviral transduction, DNA electroporation, or RNA electroporation. The nucleic acid molecule or vector is either transiently, or, preferably, stably provided to the cell. Methods for transduction or electroporation of cells with a nucleic acid are known to the skilled person.

10

In general, the chimeric receptors of the invention are advantageously used to improve T cell function and/or T cell survival, preferably of T cells reactive against tumors. Provided is therefore a method for improving T cell function and/or T cell survival in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a chimeric receptor, a nucleic acid molecule, a vector or a cell, preferably a T cell, according to the invention. Improving T cell function and/or T cell survival preferably comprises preventing or inhibiting T cell exhaustion. In a preferred aspect the subject is suffering from cancer. Said cell is preferably a T cell, preferably an autologous T cell of a subject suffering from cancer, such as a tumor derived T cell or a tumor infiltrating lymphocyte (TIL) or a T cell isolated from blood of the subject.

15

Also provided is a chimeric receptor, nucleic acid molecule or vector according to the invention, or a cell comprising the nucleic acid molecule or vector according to the invention for use in therapy. Preferably, said therapy is immunotherapy, more preferably tumor immunotherapy. In a preferred embodiment said tumor immunotherapy comprises adoptive cell transfer, more preferably adoptive T cell transfer.

20

Also provided is therefore a method for immunotherapy in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a chimeric receptor, a nucleic acid molecule, a vector or a cell according to the invention. In a preferred embodiment, such method comprises administration of a cell or population of cells according to the invention.

25

30

“Adoptive cell transfer” refers to the transfer of cells into a patient. In particular, “adoptive T cell transfer” refers to the transfer of T cells into a patient. The cells may have originated from the patient itself or may have come from another individual. Adoptive T cell transfer preferably comprises transfer of tumor
5 infiltrating lymphocytes (TILs) or T cells isolated from blood, preferably derived from the subject or patient to be treated. If T cells isolated from blood are used, the T cells further preferably express a chimeric antigen receptor (CAR) or tumor specific T cell receptor.

“TILs” refers to autologous T cells found in or around the tumor of the
10 patient to be treated. The T cells are expanded *in vitro*, e.g. cultured with cytokines such as interleukin-2 (IL-2) and anti-CD3 antibodies, and transferred back into the patient. Upon administration *in vivo*, TILs reinfiltate the tumor and target tumor cells. Prior to TIL treatment, patients can be given nonmyeloablative
15 lymphodepletion is completed, patients are then infused with TILs, optionally in combination with IL-2. Procedures for immunotherapy with adoptive T cell transfer including TILs, are well known in the art. In a preferred embodiment, TILs used in accordance with the invention are provided with a nucleic acid molecule or vector according to the invention after isolation from the patient. It is
20 further preferred that the TILs express a chimeric receptor according to the invention.

“Immunotherapy” as used herein refers to treatment of an individual suffering from a disease or disorder by inducing or enhancing an immune response in said individual. Tumor immunotherapy relates to inducing or enhancing an
25 individual’s immune response against a tumor and/or cells of said tumor. Immunotherapy according to the invention can be either for treatment or prevention. “Treatment” means that the immune response induced or enhanced by the immunotherapy component ameliorates or inhibits an existing tumor. “Prevention” means that the immunotherapy component induces a protective
30 immune response that protects an individual against developing cancer.

Also provided is a method of treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of T cells comprising a nucleic acid sequence encoding the chimeric receptor according to the

invention. Said T cells are preferably autologous T cells, such as TILs or T cell isolated from blood of the subject.

Tumors that can be treated or prevented using therapy based on a chimeric receptor according to the invention and/or a cell, preferably T cell, more preferably autologous T cells, such as TILs or T cells isolated from blood, provided with a nucleic acid molecule encoding a chimeric antigen receptor according to the invention or expressing a chimeric antigen receptor according to the invention can be any type of tumor, including primary tumors, secondary tumors, advanced tumors and metastases. Non-limiting examples tumors that can be treated or prevented in accordance with the invention are acute myeloid leukemia (AML), chronic myeloid leukemia (CML), chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), chronic myelomonocytic leukemia (CMML), lymphoma, multiple myeloma, eosinophilic leukemia, hairy cell leukemia, Hodgkin lymphoma, non-Hodgkin lymphoma, large cell immunoblastic lymphoma, plasmacytoma, lung tumors, small cell lung carcinoma, non-small cell lung carcinoma, pancreatic tumors, breast tumors, liver tumors, brain tumors, skin tumors, bone tumors, colon tumors, rectal tumors, anal tumors, tumors of the small intestine, stomach tumors, gliomas, endocrine system tumors, thyroid tumors, esophageal tumors, gastric tumors, uterine tumors, urinary tract tumors and urinary bladder tumors, kidney tumors, renal cell carcinoma, prostate tumors, gall bladder tumors, tumors of the head or neck, ovarian tumors, cervical tumors, glioblastoma, melanoma, chondrosarcoma, fibrosarcoma, endometrial, esophageal, eye or gastrointestinal stromal tumors, liposarcoma, nasopharyngeal, thyroid, vaginal and vulvar tumors.

A “subject” as used herein is preferably a mammal, more preferably a human.

“T cells” or “TILs” referred to herein can be either CD4⁺ or CD8⁺ T cells or TILs or a combination of CD4⁺ or CD8⁺ T cells or TILs. In a preferred embodiment T cell or TILs are CD8⁺ T cells or TILs.

The invention also provides a genetically modified T cell, which is transduced by the nucleic acid molecule or vector of the invention. Said modified T cell is preferably a tumor derived T cell or a tumor infiltrating lymphocyte (TIL).

Further, an isolated cell according to the invention is preferably a T cell, more preferably a tumor derived T cell or a TIL. In a particularly preferred embodiment, the T cell is an autologous T cell isolated from a patient suffering from cancer, i.e. an autologous TIL or an autologous T cell isolated from blood. It is further
5 preferred that the T cell expresses a chimeric antigen receptor according to the invention.

In one aspect, treatment based on a chimeric receptor according to the invention is combined with at least one further immunotherapy component. Such
10 further immunotherapy component can be any immunotherapy component known in the art. Preferably, said further immunotherapy component is selected from the group consisting of cellular immunotherapy, antibody therapy, cytokine therapy, vaccination and/or small molecule immunotherapy, or combinations thereof.

In a preferred embodiment, treatment with a chimeric receptor is
15 combined with antibody-based immunotherapy, preferably comprising treatment using antibodies directed against a co-inhibitory T cell molecule. Co-inhibitory T cell molecules are also referred to as immune checkpoints. Preferred examples of co-inhibitory T cell molecules are cytotoxic T-lymphocyte antigen-4 (CTLA-4), programmed death-1 (PD-1), PD-ligand 1 (PD-L1), PD-L2, Signal-regulatory
20 protein alpha (SIRP α), T-cell immunoglobulin- and mucin domain-3-containing molecule 3 (TIM3), lymphocyte-activation gene 3 (LAG3), killer cell immunoglobulin-like receptor (KIR), CD276, CD272, A2AR, VISTA and indoleamine 2,3 dioxygenase (IDO). An antibody against a co-inhibitory T cell
25 molecule that is combined with a chimeric receptor or cell comprising a chimeric receptor according to the invention is therefore preferably selected from the group consisting of an anti-CTLA4 antibody, an anti-PD-1 antibody, an anti-PD-L1 antibody, an anti-PD-L2 antibody, an anti-SIRP α antibody, an anti-TIM3 antibody, an anti-LAG3 antibody, an anti-CD276 antibody, an anti-CD272 antibody, an anti-KIR antibody, an anti-A2AR antibody, an anti-VISTA antibody, anti TIGIT
30 antibody and an anti-IDO antibody. Suitable antibodies used as a further immunotherapy component are nivolumab, pembrolizumab, lambrolizumab, ipilimumab and lirilumab.

As demonstrated in the Examples, Notch signaling decreases expression of co-inhibitory T cell molecules. Also provided is therefore a method for enhancing efficacy of an antibody-based immunotherapy as defined herein in a subject suffering from cancer and being treated with said antibody, the method comprising
5 administering to the subject a therapeutically effective amount of T cells expressing the chimeric receptor according to the invention. Said T cells are preferably autologous T cells, such as autologous TILs or T cells isolated from blood of the subject.

In a further preferred embodiment, treatment with a chimeric receptor
10 is combined with treatment involving a chimeric antigen receptor (CAR) or tumor specific T cell receptor. Preferably cells comprising and/or expressing a chimeric receptor according to the invention that further comprise a chimeric antigen receptor (CAR) are used. This is in particular preferred if T cells other than TILs, such as autologous T cells isolated from blood, are used. CARs are antigen-targeted
15 receptors composed of intracellular T-cell signaling domains fused to extracellular tumor-binding moieties, mostly single-chain variable fragments (scFvs) from monoclonal antibodies. CARs specifically recognize (tumor) cell surface antigens, independent of MHC-mediated antigen presentation. CARs preferably contain an ectodomain (such as an antigen binding portion of an antibody) specific for a tumor
20 associated antigen, coupled to a signaling domain, preferably of CD3 ζ , and a costimulatory receptor, such as CD28 or 4-1BB. Said cells are preferably T cells, more preferably autologous T cells derived from the subject to be treated, such as from blood or the tumor.

25 Features may be described herein as part of the same or separate aspects or embodiments of the present invention for the purpose of clarity and a concise description. It will be appreciated by the skilled person that the scope of the invention may include embodiments having combinations of all or some of the features described herein as part of the same or separate embodiments.

30

The invention will be explained in more detail in the following, non-limiting examples.

Brief description of the drawings

Figure 1: Schematic of a Chimeric Antigen Receptor (CAR).

Shown is an scFv (single chain) ligand binding portion of an antibody, which is
 5 linked to the intracellular signaling domains of either the 4-1BB or the CD28
 costimulatory receptor and to the CD3 zeta chain.

Figure 2: Notch signaling pathway.

Shown light blue and in red are Jagged and Delta, two membrane bound ligands of
 10 Notch. The Notch receptor itself is depicted in orange. After ligand binding the
 intracellular domain of Notch (NICD) is cleaved off the membrane and translocates
 to the nucleus, where it forms a transcriptional activator in complex with CSL and
 MAML proteins.

15 **Figure 3: Notch deficiency leads to reduced effector functions in antiviral
 CD8 T cells. (A)** Flow chart of experiment. Wild type ($\text{Notch1}^{\text{flox/flox}}\text{Notch2}^{\text{flox/flox}}$) or
 T cell specific Notch1/2 knock out mice ($\text{Notch1}^{\text{flox/flox}}\text{Notch2}^{\text{flox/flox}}\text{CD4-Cre}$) were
 infected intranasally with HkX31 influenza virus and after 10 days T cells (results
 shown from spleen) were isolated and stained for CD8 and binding to the $\text{D}^{\text{b}}\text{NP}_{366-374}$
 20 MHC tetramer **(B)**. **(C)** Number of $\text{D}^{\text{b}}\text{NP}_{366-374}$ -specific CD8^+ T cells in wild type
 (black bars) or Notch1/2KO mice (open bars). Percentage $\text{IFN}\gamma$ **(D)** or Granzyme B
 producing cells **(E)**-blue histogram-wild type; red histogram N1/2ko) among
 $\text{D}^{\text{b}}\text{NP}_{366-374}$ -specific CD8^+ T cells. **(F)** Relative mRNA levels for Granzyme B and
 Perforin in FACSsorted $\text{D}^{\text{b}}\text{NP}_{366-374}$ -specific CD8^+ T cells. **(G)** HkX31 viral loads **(H)**
 25 mouse weight curves and **(I)** influenza-neutralizing antibody titers in blood of
 infected mice. All results from Backer et al. 2014.

**Figure 4: CD8 T cell-intrinsic requirement for Notch in generation of
 effective memory.** Wild type or Notch1/2 knock out mice were first infected
 30 intranasally with HkX31 influenza virus and then reinfected after 43 days with
 PR8 influenza. **(A)** Percentages of $\text{D}^{\text{b}}\text{NP}_{366-374}$ MHC tetramer binding CD8^+ T cells
 in blood 8 days after reinfection. **(B)** Numbers of $\text{D}^{\text{b}}\text{NP}_{366-374}$ MHC tetramer binding
 CD8^+ T cells in spleens and lungs. **(C)** Rag1 deficient mice were reconstituted with

CD45.1⁺ WT bone marrow (BM) mixed with CD45.2⁺ WT BM (black bars) or mixed with CD45.2⁺ Notch1/2KO BM (white bars). Mice were then infected and reinfected as in A. Shown on the left are responses of CD45.1⁺ CD8⁺ T cells and on the right responses of CD45.2⁺ CD8⁺ T cells. Also shown are responses of mice reconstituted with CD45.2⁺ KO BM only (grey bars). Results were normalized against the corresponding WT controls. **(D)** Percentage IFN γ , TNF α and Granzyme B producing CD8 T cells isolated from lungs and restimulated *in vitro* with NP₃₆₆₋₃₇₄ peptide and wild type splenic antigen presenting cells (note that the number of influenza specific T cells was similar in lungs-see Figure 4B).

10

Figure 5: Notch deficiency leads to reduced effector functions in antiviral CD8 T cells. **(A)** Gene Set Enrichment Analysis of differentially expressed genes (obtained by RNAseq) between influenza specific effector CD8 T cells from wild type or T cell specific Notch1/2 knock out mice. **(B)** mRNA levels for PD1 and Lag3 in wild type or Notch1/2ko effector T cells. **(C)** 10⁴ CD45.2 wild type or Notch1/2ko OT1 T cells were transferred into CD45.1 wild type congenic mice, which were subsequently infected with Ovalbumin NP₃₆₆₋₃₇₄ peptide expressing influenza. Representative FACS histogram (left) and MFI (right) for PD1 on influenza-specific memory CD8 T cells in lungs 30 days after infection. **(D)** Flow chart for experiment: CD45.2 OT1 T cells were transduced with empty vector or NICD (Notch intracellular domain) encoding retroviral vector and transferred into CD45.1 wild type mice infected as in (C). After 7 days, T cells were isolated and analyzed by FACS for PD1 levels **(E)**.

25

Figure 6: Physiological Notch responses are very sensitive to NICD. **(A)** Activation of the Notch responsive HES1-luciferase reporter induced by different levels of nuclear release of mER-NICD1 or constitutive NICD1 expression. U2OS cells were transfected with reporter plasmids expressing Firefly luciferase, a plasmid constitutively expressing Renilla luciferase and an empty vector control, mER-NICD or NICD1, respectively. Tamoxifen (4-HT) was added at the indicated concentrations. Firefly luciferase activities were normalized to Renilla luciferase activities from the same samples and are displayed as fold of empty vector control samples (mean + SD). Note that MER-NICD elicits 15.2-fold leaky induction in the

30

absence of 4-HT. **(B, C)** Flow cytometric analysis of thymocytes after 2 weeks of co-culture on control OP9 cells. CD34⁺CD1a⁻ progenitors were transduced with NICD1, mERNICD1 or an empty vector control prior to co-culture. Tamoxifen was added to mER- NICD1 and empty vector transduced cultures at the concentrations indicated. **(B)** Transduced cells were analyzed for surface expression of CD4 and CD8 to assess T cell differentiation. **(C)** ILC2 differentiation as determined by expression of CRTH2 on transduced lineage- cells.

Figure 7: The anti-TA-chNotch receptor. The LNR, heterodimerization, transmembrane and intracellular domains of Notch are fused to an antibody neo-ectodomain directed against a surface molecule on an adjacent cell, such as a tumor antigen (TA). Binding of the antibody neo-ectodomain to a ligand on an opposing cell, such as a tumor cell, will induce cleavage by TACE and γ -secretase, resulting in translocation of NICD to the nucleus and transactivation of endogenous Notch target genes. The anti-TA-chNotch receptor is inactive in the absence of the activating surface antigen.

Figure 8: Amino acid sequence of Notch1 receptor. Sequence of UniProtKB/Swiss-Prot: P46531.4.

20

Figure 9. Notch can protect CD8 T cells from developing hallmarks of exhaustion.

(A) OT-1 CD8⁺ T cells were activated and transduced with viruses expressing EV or NICD coupled to IRES-Thy1.1 and rested for 5 days. Subsequently, cells were co-cultured overnight with B16-F10 melanoma cells (not expressing Ovalbumin) and then stained for Thy1.1 (to identify transduced cells) and Granzyme B and analyzed by flow cytometry. Note that Thy1.1⁻ cells were gated out of the analysis. Note furthermore that the expression level of Thy1.1 differs between EV and the NICD construct due to the size of the NICD insert. **(B)** OT-1 T cells were activated and transduced as in (A). Five days after transduction, cells were cultured for an additional 6 days and fresh B16-F10 melanoma cells expressing Ovalbumin (B16-Ova) were added daily for repeated TCR stimulation leading to exhaustion. Cells were then stained for Thy1.1 and PD1 and analyzed by flow cytometry. **(C)** OT-1

30

CD8⁺ T cells were treated as in (B) and the percentages of Thy1.1⁺ cells were analyzed by flow cytometry after different times of coculture with B16-Ova, as indicated in the figure. (D) OT-1 CD8⁺ T cells were activated and transduced with viruses expressing EV or mER-NICD (a tamoxifen inducible version of NICD) and cultured with B16-Ova as in (C) without or with 0.05mM (+) or 0.5mM (++) tamoxifen. Thy1.1⁺ cells were then analyzed by flow cytometry for IFN γ , IL10, Granzyme B and PD1 expression.

Figure 10: Generation and expression of a chimeric Notch receptor (CNR) directed against CD19. (A) schematic of experiment. The CNR contains an extracellular ScFv domain specific for human CD19. A human CD19 protein, fused to a human IgG1 Fc portion, was used to detect surface expression of the CNR. A fluorescently labeled anti-human antibody was then used to detect the hCD19-Ig fusion protein. PEST = Notch PEST domain; AF647 = Alexa Fluor 647. (B) HEK293T cells were transfected with a CNR expression construct or control and subsequently stained without or with different concentrations of hCD19-Ig, followed by fluorescently labeled anti-human antibody.

Examples

20

Example 1

Results

To examine the role of Notch in CD8 T cell responses, in Backer et al. 2014 mice carrying T cell-specific deletions in the *Notch1* and *Notch2* genes (Notch1/2ko) were infected with influenza virus. At the peak of the response, influenza-specific CD8 T cells were detected using D^b tetramers loaded with an immuno-dominant peptide of influenza (**Figure 3a,b**). Although the magnitude of the influenza-specific CD8 T cell response was similar in wild type (WT) and Notch1/2ko mice (**Figure 3C** and not shown), Notch1/2 deficient T cells produced less IFN γ and Granzyme B than WT CD8 T cells (**Figure 3d,e,f**). Notch1/2ko mice were also less able to clear the influenza virus and exhibited delayed recovery (**Figure 3g,h**). Titers of neutralizing antibodies were, if anything, elevated in Notch1/2ko mice (**Figure 3i**),

suggesting that their inability to clear the virus was caused by their ineffective CD8 T cell response.

Memory responses to influenza were affected even more severely by Notch1/2
5 deficiency in all anatomical locations examined (**Figure 4a,b**). Defective memory activity was a consequence of a CD8 T cell-intrinsic function of Notch, as shown by the inability of Notch1/2ko CD8 T cells to expand even in mixed bone marrow chimeras (**Figure 4c**). Surprisingly, normal numbers of Notch1/2ko memory CD8 T cells were found in lungs (**Figure 4b**), but these hardly produced effector molecules
10 (**Figure 4d**).

The profound unresponsiveness of Notch1/2ko CD8 T cells is reminiscent of “exhaustion”: inability to fully respond due to expression of inhibitory receptors, such as PD1 and Lag3 (Wherry and Kurachi, 2015). This notion was reinforced by
15 whole transcriptome analysis of Notch1/2ko CD8 effector T cells. Among differentially expressed genes between Notch1/2ko and WT effector T cells, the most significantly enriched gene set was derived from a comparison between acute and chronic infection with LCMV (**Figure 5a**), the prototypical model used to study T cell exhaustion (Wherry and Kurachi, 2015). Indeed, mRNA levels for both
20 PD1 and Lag3 were elevated in Notch1/2ko CD8 effector T cells (**Figure 5b**).

Importantly, expression of PD1 was elevated on the surface of Notch1/2 deficient OT1 T cells transferred into WT congenic recipient mice that were infected with Influenza-Ova (to which the OT1 T cell receptor responds) (**Figure 5c**). The
25 endogenous repertoire of T and B cells effectively clears influenza virus in these mice, excluding viral persistence as an explanation for the elevated PD1 expression selectively on Notch1/2ko T cells. Furthermore, expression of an activated Notch1 allele (NICD) specifically in Notch1/2ko OT1 T cells strongly suppressed PD1 expression (**Figure 5e**). This demonstrates that Notch suppresses expression of
30 PD1 in a CD8 T cell-intrinsic manner.

Expression of the intracellular domain of Notch (NICD) mimics activation of Notch, both in CD4 T cells and CD8 T cells (Helbig et al. 2012; Backer et al. 2014; Amsen

et al. 2007). Notch signaling is exquisitely sensitive and the number of nuclear NICD molecules obtained by overexpression of an NICD construct likely vastly exceeds the number of molecules obtained after ligand-mediated activation. This is illustrated by experiments using tamoxifen inducible MER-NICD alleles in thymic progenitor cells. Culturing CD34⁺CD1a⁻ human thymic progenitor cells on OP9 stromal cells only resulted in differentiation if NICD was expressed (**Figure 6b**). Strikingly, maximal differentiation of CD4⁺CD8⁺ double positive cells was already obtained by the leaky activity of MER-NICD in the absence of tamoxifen (**Figure 6b**), conditions that result in very weak transactivation of a luciferase reporter construct (**Figure 6a**). Furthermore, increasing activity of MER-NICD by addition of tamoxifen resulted in a gradual conversion of differentiation from double positive thymocytes into CRTH2⁺ ILC2 cells (**Figure 6c**). These results emphasize the exquisite sensitivity of endogenous response programs to NICD. Furthermore, they show that the strength of Notch signaling sometimes qualitatively affects the biological response to this receptor. (*These results have been published in Gentek et al. 2013*)

Materials and methods

Mice. All mice were on a C57BL/6 background. *Notch1^{fllox/fllox}Notch2^{fllox/fllox}Cd4-Cre* mice were used (Amsen et al. 2014; Amsen et al. 2004). Cre-negative littermates were used in all experiments. Transgenic mice expressing the OT-I TCR (003831) are available from Jackson Laboratories. Mice were bred and housed in specific pathogen-free conditions at the Animal Centers of the Academic Medical Center (AMC, Amsterdam, The Netherlands). Mice (both male and female) were between 8–16 weeks of age at the start of the experiment. During infection experiments, wild-type and Notch1-2-KO mice were housed together to avoid cage bias. No intentional method for randomization was used. No formal method for blinding was used, except for determination of viral loads and hemagglutination assay, where the operator did not know mouse genotypes. Mixed-bone marrow (BM) chimeras containing wild-type and Notch1-2-KO BM at a 1:1 ratio were generated via intravenous injection of 5–10 × 10⁶ donor BM cells into lethally irradiated RAG1-deficient mice. Wild-type and Notch1-2-KO cells of donor origin were identified with the congenic CD45.1/2 markers. BM chimeras were used at 12 weeks after

engraftment. All mice were used in accordance of institutional and national animal experimentation guidelines. All procedures were approved by the local Animal Ethics Committees.

5 **Media, reagents and mAbs for mouse studies.** Culture medium was Iscove's modified Dulbecco's medium (IMDM; Lonza) supplemented with 10% heat-inactivated FCS (Lonza), 200 U/ml penicillin, 200 µg/ml streptomycin (Gibco), GlutaMAX (Gibco) and 50 µM β-mercaptoethanol (Invitrogen) (IMDMc). All directly conjugated monoclonal antibodies used for flow cytometry were purchased
10 from eBioscience, San Diego, CA, unless stated otherwise: anti-CD3ε (clone 145-2C11), anti-CD4 (clone GK1.5), anti-CD8α (Ly-2, clone 53-6.7), anti-CD28 (clone 37.51), anti-CD44 (clone IM7), anti-CD45.1 (clone A20, BD Biosciences), anti-CD45.2 (clone 104), anti-CD127 (anti-IL7Rα, clone A7R34), anti-Granzyme B (clone GB-11, Sanquin PeliCluster), anti-IL-2 (clone JES6-5H4), anti-IFN-γ (clone
15 XMG1.2), anti-KLRG-1 (clone 2F1), and anti-TNFα (clone MP6-XT22), isotype control (cat. #3900S) (Cell Signaling Technology).

Influenza infection. Mice were intranasally infected with 100–200 × 50% tissue culture effective dose (TCID₅₀) of the H3N2 influenza A virus HKx31 (Belz et al.
20 2000), influenza A/WSN/33, A/WSN/33–OVA(I) (Topham et al. 2001), A/PR/8/34 (H1N1) or the recombinant A/PR/8/34 expressing the LCMV gp₃₃₋₄₁ epitope (Mueller et al. 2010). Stocks and viral titers were obtained by infecting MDCK or LLC-MK2 cells as described previously (Van der Sluijs et al. 2004). At indicated time
25 intervals, blood samples were drawn from the tail vein or mice were sacrificed and organs were collected to determine numbers of influenza-specific CD8⁺ T cells. Influenza-specific CD8⁺ T cells were enumerated using anti-CD8 (53-6.7) and PE- or APC-conjugated tetramers of H-2D^b containing the influenza-A-derived nucleocapsid protein (NP) peptide NP₃₆₆₋₃₇₄ ASNENMETM (produced at the Sanquin Laboratory for Blood Research). A/PR/8/34 viral loads in lungs of infected
30 mice were determined by isolating lung mRNA and detection of viral mRNA by quantitative PCR using the following primers and probe specific for the A/PR/8/34 M gene. Sense primer: 5'-CAAAGCGTCTACGCTGCAGTCC-3'; antisense primer:

5'-TTTGTGTTTCACGCTCACCGTGCC-3'; Probe: 5'-
AAGACCAATCCTGTACCTCTGA-3'.

Sera were tested for the presence of neutralizing antibodies to this virus by hemagglutination inhibition (HI) assay as described previously using four
5 hemagglutinating units of virus and turkey erythrocytes (Palmer et al. 1975). Values represent the maximum serum dilution at which agglutination was completely inhibited.

Flow cytometry and cell sorting. For intracellular cytokine and granzyme B
10 staining, splenocytes and total lung samples were stimulated with 1 µg/ml of the MHC class I restricted influenza-derived peptide NP₃₆₆₋₃₇₄ ASNENMETM for 4 h in the presence of 10 µg/ml brefeldin A (Sigma) to prevent cytokine release. Cells were stained with the relevant fluorochrome-conjugated mAbs for 30 min at 4 °C in PBS containing 0.5% BSA and 0.02% NaN₃. For intracellular staining, cells were fixed
15 and permeabilized using the Cytotfix/Cytoperm (BD Biosciences). Data acquisition and analysis was done on a FACSCanto (Becton Dickinson) and FlowJo software. To isolate H-2 Db–NP tetramer-positive CD8⁺ T cells from influenza infected mice, single cell suspensions of spleens were stained with influenza-specific tetramers and various markers. Cells were sorted using FACS Aria cell sorters (BD
20 Biosciences).

For analysis of human thymocytes, distinction of live and dead cells was based on staining with 7-Aminoactinomycin D (7-AAD, eBiosciences) or fixable live/dead dyes (Invitrogen). Data were acquired on a LSR Fortessa flow cytometer (BD
25 Bioscience) and analyzed using FlowJo software (TreeStar). Single cell suspensions were stained with antibodies directly labeled with Fluorescein Isothiocyanate (FITC), Phycoerythrin (PE), Phycoerythrin- Cyanine 5 (PE-Cy5), PE-Cy5.5, PE-Cy7, PerCP-Cy5.5, Allophycocyanin (APC)/Alexa Fluor 647, APC-Cy7, AF700 (all BD Bioscience, Biolegend or MACS Miltenyi), Horizon V500 (HV500, BD
Bioscience), Brilliant Violet 421 (BV421), BV711 and BV785 (all Biolegend).
30 Antibodies specific for the following human antigens were used: CD1a, CD3, CD4, CD7, CD8, CD11c, CD14, CD19, CD25, CD34, CD45, CD56, CD94, CD117 (cKit), CD123, CD127 (IL-7R α), CD161, CD294 (CRTH2), CD303 (BDCA2), CD336 (Nkp44), CD278 (ICOS), TCR $\alpha\beta$, TCR $\gamma\delta$ and FcER1. Anti-mouse CD90.1 (Thy1.1) -

FITC, -PE or -APC-eFluor 780 (eBioscience) were used to detect cells transduced with MSCV - IRES-Thy1.1 retroviruses.

Retroviral transductions and adoptive transfers of mouse CD8⁺ T cells.

5 Virus was produced in PlatE cells as described (Amsen et al. 2004). Total splenocytes from CD45.2⁺ OT-I wild-type or OT-I Notch1-2-KO mice were incubated with 1 nM OVA₂₅₇₋₂₆₄ peptide, and next day cells were spin-infected (700 × g for 90 min at 37°C) with viral supernatant (with 8 µg/ml polybrene), followed by 5 h at 37°C. Medium was replaced and next day, live T cells were isolated by
10 density centrifugation (Lymphoprep, Axis-shield PoC) and between 7.5 × 10² and 5 × 10⁴ cells were transferred into timed influenza-OVA infected CD45.1⁺ mice. Donor OT-1 T cells were detected 5–10 days after transfer as CD45.2⁺CD8⁺ and Thy1.1 or GFP triple positive cells.

15 **Virus production and transduction of human thymocytes.** For virus production, Phoenix GALV packaging cells were transiently transfected using FuGene HD (Promega). Virus containing supernatant was harvested 48h after transfection, snap frozen on dry ice and stored at -80°C until use. For transduction, cells were incubated with virus supernatant in plates coated with Retronectin
20 (Takara Biomedicals) for 6-8h at 37°C the following day.

Retroviral constructs used for human thymocyte experiments. The human NICD1-IRES-Thy1.1-MSCV construct has been described before (Amsen et al. 2004). To generate the mER-NICD fusion, an N-terminal mER domain was PCR
25 amplified using the following primers:

GATCAGGAATTCCACACCATGGGAGATCCACGAAATGAA and
GATCAGGATATCCACCTTCCTCTTCTTCTTGG and cloned into the EcOR1 and EcORV sites of pBluescript (pBS) to create mER-pBS. Human NICD1 lacking a translation initiation signal was PCR amplified using these primers:
30 ATCGGAGGTTCTCGCAAGCGCCGGCGGCAGCAT and
GATCAGAAGCTTGAATTCTTACTTGAAGGCCTCCGGAATG and subsequently cloned into the EcORV and HindIII sites of mER-pBS. The mER-NICD1 fusion insert was then cloned into IRES-Thy1.1-MSCV using BamH1 and Cla1.

Gene expression profiling mouse studies. H-2 D^b-NP₃₆₆₋₃₇₄⁺CD8⁺ T cells were isolated from spleens of influenza infected mice by flow cytometry. Total RNA was extracted with TRIzol reagent (Invitrogen) according to the manufacturer's protocol. For Deep sequencing analysis, total RNA was further purified by

5 nucleospin RNAII columns (Macherey-Nagel) and RNA was amplified using the Superscript RNA amplification system (Invitrogen) and labeled with the ULS system (Kreatech), using either Cy3 or Cy5 dyes (Amersham). Sequences were obtained by pooling 10 samples in one lane on a HiSeq2000 machine. Between 17 and 27 million reads were obtained per sample.

10 Read mapping (TopHat) and determining differentially expressed genes (DESeq) was done as described in (Anders et al. 2013). Reads were mapped against the mouse reference genome (build mm9) using TopHat (version 1.4.0), which allows to span exon-exon junctions. TopHat was supplied with a known set of gene models (NCBI build 37, version 64). In order to obtain per sample gene counts HTSeq-count

15 was used. This tool generates gene counts for each gene that is present in the provided Gene Transfer Format (GTF) file. Genes that have zero counts across all samples were removed from the dataset. Statistical analysis was performed using the R package DESeq. Differentially expressed genes were determined between the SLEC and MPEC samples, and between the wild type and knock-out samples.

20 DESeq assumes that gene counts can be modelled by a negative binomial distribution. For sample normalisation the 'size factors' were determined from the count data. The empirical dispersion was determined with the 'pooled' method, which used the samples from all conditions with replicates to estimate a single pooled dispersion value. Subsequently, a parametric fit determines the dispersion-mean relationship for the expression values resulting in two dispersion estimates

25 for each gene (the empirical estimated, and the fitted value). Using the 'maximum sharingMode' we selected the maximum of these two values to be more conservative. Finally, p-values and FDR corrected p-values were calculated. To highlight biological processes that are over-represented in the set of

30 differentially expressed genes we used Bioconductor package Goseq (Young et al. 2010), which was developed for the analysis of RNA-seq data. First we selected all genes with an FDR<0.5 from the SLEC-MPEC and WT-KO comparisons. Subsequently, the GO 'Biological Processes' gene sets were used to determine over-

represented processes. In addition we used the 'C7' gene set from the Molecular Signatures Database (MSigDB; <http://www.broadinstitute.org/gsea>), which is a collection of annotated gene sets. Gene set C7 comprises immunologic signatures composed of gene sets that represent cell types, states, and perturbations within the immune system. The signatures were generated by manual curation of published microarray studies in human and mouse immunology. This gene set was generated as part of the Human Immunology Project Consortium (HIPC; <http://www.immuneprofilling.org>). An in-house R script was developed to convert the C7 gene set into a format that could be used by GSeq.

10

Statistical analysis. Figures represent means and error bars denote standard error of the mean (s.e.m.). Standard Student's *t*-tests (unpaired, two-tailed) was applied with GraphPadPrism software. If 3 or more groups were compared One-way ANOVA with Bonferroni correction was used. $P < 0.05$ was considered statistically significant.

15

Isolation of human thymic hematopoietic progenitors. Postnatal thymic (PNT) tissue specimens were obtained from children undergoing open heart surgery (LUMC, Leiden, the Netherlands); their use was approved by the AMC ethical committee in accordance with the declaration of Helsinki. Cell suspensions were prepared by mechanical disruption using the Stomacher 80 Biomaster (Seward). After overnight incubation at 4°C, thymocytes were isolated from a Ficoll-Hypaque (Lymphoprep; Nycomed Pharma) density gradient. Single cell suspensions were enriched for CD34⁺ cells by MACS (Miltenyi Biotec), stained with fluorescently labeled antibodies and subsequently FACS sorted on a FACS Aria (BD Bioscience) as CD34⁺CD1a⁻CD3⁻CD56⁻BDCA2⁻ or CD34⁺CD1a⁺CD3⁻CD56⁻BDCA2⁻, respectively (referred to in this study as CD34⁺CD1a⁻ and CD34⁺CD1a⁺). Purity of the sorted populations was > 99%.

20

25

***In vitro* differentiation of thymic progenitors.** Sorted thymic progenitors were cultured overnight in Yssel's medium containing 5% normal human serum, SCF (20ng/ml) and IL-7 (10ng/ml, both PeproTech). OP9 cells were mitotically inactivated by irradiation with 30Grey and seeded at a density of $5 \times 10^3/\text{cm}^2$ one

30

day prior to initiation of co-cultures. After transduction, thymic progenitors were added to pre-seeded OP9 cells. Co-cultures were performed in MEM α (Invitrogen) with FCS (20% Fetal Clone I, Hyclone) and IL-7 (5ng/ml). In some cases, Flt3l (5ng/ml, PeproTech) was added to the medium. Cultures were refreshed every 3-4
5 days. Differentiation assays for innate lymphoid cells were typically analyzed after 1week, unless stated otherwise. Cells were harvested by forceful pipetting and passed through 70 mm nylon mesh filters (Spectrum Labs).

Reporter gene assays. U2OS cells were transiently transfected using the FuGene HD transfection reagent (Promega). Cells were co-transfected with a NOTCH-responsive promoter and either NICD1 – MSCV Th1.1, mER- NICD1 – MSCV Th1.1 or an empty vector control. To correct for differences in transfection efficiency, the pRL-CMV control vector was co- transfected, from which Renilla luciferase is expressed constitutively. Transfections were performed in triplicate.
10
15 Where applicable, 4-Hydroxy- Tamoxifen (Sigma) was added after overnight incubation to induce nuclear translocation of mER-NICD1. Cells were lysed 48h post transfection and luciferase activity was measured using the Dual Luciferase Reporter Assay System (Promega) on a Synergy HT microplate reader (Syntek). Two different Notch responsive reporter constructs were used, which have been
20 described previously (Nam et al. 2007).

The Chimeric Notch receptor (ChNR) system. To generate a Chimeric Notch receptor the extracellular domain of Notch except the heterodimerization domain is replaced by a heterologous ligand binding domain consisting of an scFv antibody
25 domain fused to the heterodimerization domain of Notch. This receptor will be activated by binding to the cognate ligand of the scFv antibody on the surface of an adjacent cell, but will remain silent when this surface antigen is not present (**Figure 7**). ChNR can be expressed in CD4 T cells via retroviral transduction or other methods. If such modified T cells are adoptively transferred into patients,
30 Notch can specifically be turned on only in these T cells.

The ChNR will typically not by itself be sufficient to fully activate T cells. For that, additional T cell receptor signals (or mimics thereof) are required. For instance, T

cells can be derived from primary tumors (Tumor infiltrating lymphocytes-TIL) after selection for tumor reactivity. Also, ChNR can be used in conjunction with recombinant T cell receptors against tumor antigens or in T cells engineered to express traditional chimeric antigen receptors (CAR).

5

Many variations of this basic concept are possible. As ectodomain any antibody that recognizes a surface antigen can in principle be used and any surface antigen expressed on the surface of tumor cells can in principle be targeted. Finally, even ectodomains activated by soluble ligands are an option. For instance, an
10 ectodomain can consist of an antibody to a peptide neo-epitope (as described in Rodgers et al. 2016) or to a Biotin or FITC moiety (as described in Ma et al. 2016) that is itself incorporated in another antibody (a switch antibody) directed to a surface antigen on a tumor. As a consequence, activation of the Chimeric Notch receptor will only occur if, in addition to the cell surface antigen targeted by the
15 switch antibody, the switch antibody itself is also present. This set up would permit temporary control of the receptor (turning it on and off only when desired) as well as quantitative control (by in- or decreasing the concentration of the switch antibody. In all these situations, however, liberation of the intracellular domain of Notch from the Chimeric Notch receptors remains the central goal.

20

The preparation of an exemplary Chimeric notch receptor is described in example 2.

Example 2

25

Results

T cell exhaustion occurs when T cells are chronically stimulated via their T cell receptor. The results in example 1 show that CD8 T cells responding to an infection with influenza virus are protected from activation of this exhaustion program by
30 Notch. Influenza infection does not, however, normally cause chronic stimulation of T cells. We therefore asked whether deliberate activation of Notch can also prevent exhaustion under conditions that normally do lead to exhaustion. To this end, we resorted to an in vitro system in which an activated Notch allele (NICD) can be

introduced in T cells that are then subjected to repeated TCR stimulation. NICD was expressed in OT-1 CD8 T cells (which recognize the SIINFEKL peptide from the Ovalbumin protein in H2-K^b) using a retroviral expression system. An IRES-Thy1.1 sequence in this retroviral construct allows discrimination between the transduced T cells (Thy1.1⁺) and the untransduced T cells (Thy1.1⁻). Expression of
5 NICD in CD8⁺ OT-1 T cells strongly enhanced effector functions, as evidenced for instance by the spontaneous production of the cytolytic effector protein Granzyme B (**Figure 9A**). Transduced OT-1 cells were then repeatedly stimulated by daily addition of B16F10 melanoma cells expressing Ovalbumin (B16-Ova). These
10 conditions result in prominent expression of the check-point molecule (and hallmark of exhaustion) PD1 on the surface of OT-1 T cells transduced with a control virus (Empty Vector-EV) (**Figure 9B**, left). Expression of NICD, however, nearly completely prevented expression of PD1 (**Figure 9B**, right). Expression of NICD also afforded a competitive advantage to the OT-1 T cells: the proportion of
15 Th1.1⁺ cells in the population transduced with NICD gradually increased over time, whereas the Th1.1⁺ population remained stable when cells had been transduced with Empty Vector (**Figure 9C**).

The concentration of active Notch molecules that is obtained after expression of the
20 NICD allele is probably unphysiologically high. Moreover, it may not be possible to achieve similarly high levels of such active Notch molecules using the ChNR. To test whether the protective effects on CD8 T cells can also be obtained with weaker Notch stimulation, we made use of a Tamoxifen inducible version of NICD (also used in example 1, Figure 6). This construct consists of NICD coupled at the N-
25 terminus to the ligand binding domain of the Estrogen Receptor (ER), which has been mutated such that it responds only to Tamoxifen and no longer to Estrogen. This mutated ER domain (mER) sequesters NICD molecules in the cytoplasm by binding to heat shock proteins and thereby keeps it inactive. Upon addition of
tamoxifen, the mER-NICD fusion protein however dissociates from these heat
30 shock proteins, allowing NICD to become active. As shown by luciferase reporter assays (**Figure 6A**), this fusion protein reaches much lower maximal levels of Notch activity than NICD itself and its activity can be controlled quantitatively by titration of Tamoxifen. Finally, this mER-NICD possesses some "leaky" Notch

activity even in the absence of Tamoxifen, which is almost undetectable in luciferase reporter assays, yet can elicit physiological functions of Notch such as induction of differentiation of CD4⁺CD8⁺ thymocytes from thymic precursor cells (**Figure 6B**). We therefore used this mER-NICD construct to examine the signal strength requirements for protection against exhaustion in CD8 T cells, again using the repetitive stimulation model with B16-Ova melanoma cells (as in A-C). Stimulation of mER-NICD with 0.5 or even 0.05mM of tamoxifen indeed resulted in reduced expression of PD1 and production of the tolerogenic cytokine IL10 (**Figure 9D**). It also mobilized production of effector molecules such as IFN γ and Granzyme B. Remarkably, some of these effects were obtained even by the very low leaky NICD activity that is elicited by mER-NICD in the absence of tamoxifen. We thus conclude that Notch can protect CD8 T cells from developing hallmarks of exhaustion (expression of PD1, loss of production of effector molecules) even at relatively modest levels of Notch activity.

15

Generation of chimeric Notch receptor

A chimeric Notch receptor consisting of an ScFv antibody domain directed against human CD19 was generated (ScFv as described in Molecular Immunology 1997;34:1157–1165 and used in a CAR construct in J Immunother. 2009 Sep; 32(7): 689–702). This ScFv was fused in frame to the 5'end of the human NOTCH1 protein truncated upstream of the extracellular heterodimerization domain (**Figure 10A**).

Specifically, The GMCSF leader sequence (MLLLVTSLLL CELPHPAFLL) was fused in frame to the Igk light chain Variable domain followed by the Ig heavy chain Variable domain of FMC63-28Z anti CD19 ScFv (IPDIQMTQTTSSLSASLGDRVTISCRASQDISKYLNWYQQKPDGTVKLLIYHTSR LHSQVPSRFSGSGSGTDYSLTISNLEQEDIATYFCQQGNTLPYTFGGGTKLEITG STSGSGKPGSGEGSTKGEVKLQESGPGLVAPSQSLSVTCTVSGVSLPDYGVSWI RQPPRKGLEWLGVIWGSETTYYN SALKSRLTIHKDNSKSQVFLKMNSLQTDDTA IYYCAKHYYYGGSYAMDYWGQGTSVTVSSAAA), which was fused in frame with the C-terminus from the human full length NOTCH1 protein starting at Isoleucine 1427 till Lysine 2555 (of the sequence as depicted in figure 8).

In an alternative construct, the C terminus of human NOTCH1 sequence used starts at Proline 1390. Both variants (beginning with Ile 1427 or Proline 1390, see sequence of figure 8) are made also with a deletion of the C-terminal PEST domain of human NOTCH1 (ending at Alanine 2424 of the human NOTCH1 protein, see
5 sequence of figure 8).

The fusion protein was then expressed from the pHEFTIG lentiviral expression vector (described in *J Immunol* 2009; 183:7645-7655 as “modified pCDH1”, and as “pHEF” in *PNAS* August 9, 2011 108 (32) 13224-13229) after transfection into
10 HEK293T cells and its presence at the cell surface was documented by staining with recombinant human CD19-Ig protein (**Figure 10B**).

Materials and methods

Mice. Female or male OT-1 TCR transgenic mice (C57BL/6 strain) with transgenic
15 inserts for TCR α -V2 and TCR β -V5 genes that are specifically designed to target the ovalbumin residues 257-264 presented by H2-Kb, were bred and maintained in the animal facility of the Netherlands Cancer Institute (NKI, Amsterdam, The Netherlands). All animal experiments were performed according to protocols in compliance with institutional guidelines and approved by the Animal Ethics
20 Committee of the NKI.

Cell lines and reagents. B16-F10 and B16-OVA tumor cell lines were cultured in Iscove's Modified Dulbecco's Medium (IMDM) with HEPES supplemented with 10% heat-inactivated Fetal Calf Serum (Bodingo BV), 5% L-glutamine (Lonza,
25 Belgium) and 5% Penicillin/Streptomycin (Sigma, 10.000 U Penicillin and 10 mg Streptomycin). Platinum-Eco cells and HEK293T cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) with HEPES supplemented with 10% heat-inactivated Fetal Calf Serum (Bodingo BV) and 5% L-glutamine (Lonza, Belgium). All cells were incubated at 37 °C, 5% CO₂.

30

Cell purification. A single cell suspension was obtained from the spleen and lymph nodes from OT-1 mice. CD8⁺ T cells were enriched and purified by Magnetic-Activated Cell Sorting (MACS). CD8 α^+ T cell Isolation Kit, mouse (Miltenyi

Biotech) was used for the negative selection of CD8 α ⁺ T cells. The cells were then cultured up to two weeks with IMDM supplemented with 10% heat-inactivated Fetal Calf Serum (Bodingo BV), 5% L-glutamine (Lonza, Belgium), 5% Penicillin/Streptomycin (Sigma, 10.000 U Penicillin and 10 mg Streptomycin) and
5 50 μ M β -mercapto-ethanol (Sigma Aldrich).

Retroviral transductions of murine CD8⁺ T cells. Retroviral stocks were generated by transfection of Platinum-Eco cells with the construct using FuGENE[®] HD reagent (Promega) according to the manufacturer's instructions. 3 x
10 10⁶ cells were plated in a 100 mm dish one day prior to transfection. 56 μ l of FuGENE[®] HD reagent was added to 879 μ l of plasmid solution (0.020 μ g/ μ l in OptiMEM (Gibco by Life Technologies)) and subsequently incubated for 10 minutes at RT. The complex solution was then added to the cells and incubated o/n at 37 °C. Viral supernatant was collected and filtered with a 0.45 μ m syringe filter to remove
15 cell debris. Virus supernatants were made from pMSCV-EV and pMSCV-NICD. Retroviral vectors contained an IRES sequence enabling cap-independent translation and a Thy1.1 (CD90.1) selection marker, which was used for positive transduction selection. Activated CD8⁺ T cells purified from OT-1 mice were infected with virus with an addition of 10 μ g/ml Polybrene (Merck) in a 24-well
20 plate (1x10⁶ cells/well). The cells were spun at 2000 RPM for 90 min. at RT followed by incubation for 4 h at 37 °C and 5% CO₂.

Transfection HEK293T cells. Cells were transfected with CNR-pHEFTIG or pHEFTIG empty vector in 6 well plate using Fugene HD reagent following
25 manufacturer's instructions. After 48 hours, expression was analyzed by Flow Cytometry.

CD8⁺ T cell activation and re-stimulation. For efficient in vitro activation of the T cells, an engineered APC cell line MEC.B7.SigOVA (SAMBcd8+OK) that
30 encodes the OVA257-264 (SIINFEKL) peptide was used. Following CD8⁺ T cell purification, 10⁶ CD8⁺ T cells were co-cultured with 105 SAMBOK cells in a 24-well plate for 24 hours. Cells were then collected and transduced. Cells were maintained at a cell density of \pm 1.5 x 10⁶ cells/ml until re-stimulation. Five days

after transduction, 300.000 CD8⁺ T cells were co- cultured with 50.000 B16-F10/B16-OVA in a 96-flat bottom well plate (**Fig. 5**). T cells were removed from the adherent B16 cells and were seeded to new B16 cells every 24 hours. Four hours before each desired re-stimulation time point, Brefeldin A (1000x, Invitrogen, USA) was added. Cytokine production and expression of inhibitory receptors were assessed via flow cytometry.

Flow cytometry and antibodies. All samples were measured on the BD FACSymphony A5 (BD Biosciences). Prior to flow cytometry measurement, cells were stained extracellularly (in PBS containing 1.5% FCS at 4 °C) and were fixated and permeabilized using Cytofix/Cytoperm (BD Pharmingen). Cells were then stained intracellularly (in 1X PermWash at 4 °C). Human CD19 protein, fused to a human IgG1 Fc portion (R&D Systems), was used to detect surface expression of the CNR. A fluorescently labeled anti-human antibody (Invitrogen) was then used to detect the hCD19-Ig fusion protein.

References

- Amsen, D., Blander, J. M., Lee, G. R., Tanigaki, K., Honjo, T., and Flavell, R. A. (2004) Instruction of distinct CD4 T helper cell fates by different notch ligands on antigen-presenting cells. *Cell* 117, 515-526
- Amsen, D., Antov, A., Jankovic, D., Sher, A., Radtke, F., Souabni, A., Busslinger, M., McCright, B., Gridley, T., and Flavell, R. A. (2007) Direct regulation of Gata3 expression determines the T helper differentiation potential of Notch. *Immunity* 27, 89-99
- Anders, S., McCarthy, D. J., Chen, Y., Okoniewski, M., Smyth, G. K., Huber, W., and Robinson, M. D. (2013) Count-based differential expression analysis of RNA sequencing data using R and Bioconductor. *Nat Protoc* 8, 1765-1786
- Backer, R. A., Helbig, C., Gentek, R., Kent, A., Laidlaw, B. J., Dominguez, C. X., de Souza, Y. S., van Trierum, S. E., van Beek, R., Rimmelzwaan, G. F., ten Brinke, A., Willemsen, A. M., van Kampen, A. H., Kaech, S. M., Blander, J. M., van Gisbergen, K., and Amsen, D. (2014) A central role for Notch in effector CD8(+) T cell differentiation. *Nat Immunol* 15, 1143-1151

- Belz, G. T., Xie, W., Altman, J. D., and Doherty, P. C. (2000) A previously unrecognized H-2D(b)-restricted peptide prominent in the primary influenza A virus-specific CD8(+) T-cell response is much less apparent following secondary challenge. *J Virol* 74, 3486-3493
- 5 • Gentek R, Munneke JM, Helbig C, Blom B, Hazenberg MD, Spits H, Amsen D. Modulation of Signal Strength Switches Notch from an Inducer of T Cells to an Inducer of ILC2. *Front Immunol.* 2013 Oct 22;4:334.
- Helbig, C., Gentek, R., Backer, R. A., de Souza, Y., Derks, I. A., Eldering, E., Wagner, K., Jankovic, D., Gridley, T., Moerland, P. D., Flavell, R. A., and
- 10 Amsen, D. (2012) Notch controls the magnitude of T helper cell responses by promoting cellular longevity. *Proceedings of the National Academy of Sciences of the United States of America* 109, 9041-9046
- Kochenderfer JN. et al. Construction and preclinical evaluation of an anti-CD19 chimeric antigen receptor. *J Immunother.* 2009 Sep; 32(7): 689–702.
- 15 • Legrand N. et al. Functional CD47/signal regulatory protein alpha (SIRPα) interaction is required for optimal human T- and natural killer- (NK) cell homeostasis in vivo. *J Immunol* 2009; 183:7645-7655
- Ma, J. S., Kim, J. Y., Kazane, S. A., Choi, S. H., Yun, H. Y., Kim, M. S., Rodgers, D. T., Pugh, H. M., Singer, O., Sun, S. B., Fonslow, B. R.,
- 20 Kochenderfer, J. N., Wright, T. M., Schultz, P. G., Young, T. S., Kim, C. H., and Cao, Y. (2016) Versatile strategy for controlling the specificity and activity of engineered T cells. *Proc Natl Acad Sci U S A* 113, E450-458
- Mueller, S. N., Langley, W. A., Carnero, E., Garcia-Sastre, A., and Ahmed, R. (2010) Immunization with live attenuated influenza viruses that express
- 25 altered NS1 proteins results in potent and protective memory CD8+ T-cell responses. *J Virol* 84, 1847-1855
- Morsut, L., Roybal, K. T., Xiong, X., Gordley, R. M., Coyle, S. M., Thomson, M., and Lim, W. A. (2016) Engineering Customized Cell Sensing and Response Behaviors Using Synthetic Notch Receptors. *Cell* 164, 780-791
- 30 • Nam, Y., Sliz, P., Pear, W. S., Aster, J. C., and Blacklow, S. C. (2007) Cooperative assembly of higher-order Notch complexes functions as a switch to induce transcription. *Proceedings of the National Academy of Sciences of the United States of America* 104, 2103-2108.

- Nicholson, I.C. et al. Construction and characterisation of a functional CD19 specific single chain Fv fragment for immunotherapy of B lineage leukaemia and lymphoma. *Molecular Immunology* 1997;34:1157–1165
- Palmer, D. F., Dowdle, W. R., Coleman, M. T., and Schild, G. C. (1975)
5 Advanced laboratory techniques for influenza diagnosis: procedural guide. Haemagglutination inhibition test. in Atlanta: US Department of Health, Education and Welfare, Public Health Service, US Department of Health, Education and Welfare, Public Health Service. pp 25-62
- Rodgers, D. T., Mazagova, M., Hampton, E. N., Cao, Y., Ramadoss, N. S.,
10 Hardy, I. R., Schulman, A., Du, J., Wang, F., Singer, O., Ma, J., Nunez, V., Shen, J., Woods, A. K., Wright, T. M., Schultz, P. G., Kim, C. H., and Young, T. S. (2016) Switch-mediated activation and retargeting of CAR-T cells for B-cell malignancies. *Proc Natl Acad Sci U S A* 113, E459-468
- Roybal, K. T., Rupp, L. J., Morsut, L., Walker, W. J., McNally, K. A., Park,
15 J. S., and Lim, W. A. (2016) Precision Tumor Recognition by T Cells With Combinatorial Antigen-Sensing Circuits. *Cell* 164, 770-779
- Topham, D. J., Castrucci, M. R., Wingo, F. S., Belz, G. T., and Doherty, P. C. (2001) The role of antigen in the localization of naive, acutely activated, and memory CD8(+) T cells to the lung during influenza pneumonia. *J Immunol*
20 167, 6983-6990
- Van der Sluijs, K. F., van Elden, L. J., Nijhuis, M., Schuurman, R., Pater, J. M., Florquin, S., Goldman, M., Jansen, H. M., Lutter, R., and van der Poll, T. (2004) IL-10 is an important mediator of the enhanced susceptibility to pneumococcal pneumonia after influenza infection. *J Immunol* 172, 7603-
25 7609
- van Lent AU, et al. IL-7 enhances thymic human T cell development in “human immune system” Rag2^{-/-}-IL-2R^{gammac}^{-/-} mice without affecting peripheral T cell homeostasis. *J Immunol.* 2009;183:7645–7655
- Wherry, E. J., and Kurachi, M. (2015) Molecular and cellular insights into T
30 cell exhaustion. *Nat Rev Immunol* 15, 486-499
- Young, M. D., Wakefield, M. J., Smyth, G. K., and Oshlack, A. (2010) Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome biology* 11, R14.

Claims

1. A chimeric receptor comprising an intracellular domain, a transmembrane domain, a heterodimerization domain and a Lin-12-Notch (LNR) repeats domain of a Notch receptor, and a heterologous extracellular ligand-binding domain.
5
2. The chimeric receptor according to claim 1 wherein the receptor is capable of Notch signaling.
3. The chimeric receptor according to claim 1 or 2 wherein said heterologous
10 extracellular ligand-binding domain is selected from the group consisting of:
 - a ligand binding domain specific for a soluble ligand;
 - a ligand binding domain specific for a cell surface antigen, such as a ScFv antibody domain, preferably a ScFv antibody domain that is specific for a tumor cell surface antigen;
 - 15 • an extracellular ligand-binding domain of an Fc receptor or a ligand-binding fragment thereof;
 - an extracellular domain that comprises an epitope for an antibody that can crosslink the chimeric receptor without involvement of a surface molecule.
 - an extracellular domain that comprises a moiety, such as biotin, that can be
20 crosslinked by an agent with multiple binding sites for that moiety, such as streptavidin.
4. The chimeric receptor according to any one of claims 1 to 3 further comprising a linking sequence located between the LNR domain and the heterologous
25 extracellular ligand-binding domain.
5. A nucleic acid molecule comprising a sequence encoding a chimeric receptor according to any one of claims 1 to 4.
- 30 6. A vector comprising a nucleic acid molecule according to claim 5.

7. An isolated cell comprising the nucleic acid molecule according to claim 5 or the vector of claim 6.
8. The cell according to claim 7, wherein said cell is a T cell, such as a tumor
5 derived T cell or a tumor infiltrating lymphocyte (TIL).
9. The cell according to claim 7 or 8 wherein said T cell is an autologous T cell isolated from a patient suffering from cancer.
- 10 10. The cell according to claim 8 or 9 wherein said T cell expresses a chimeric antigen receptor.
11. A genetically modified T cell, which is transduced by the nucleic acid molecule or vector according to claim 5 or 6.
15
12. A pharmaceutical composition comprising the nucleic acid molecule according to claim 5, the vector according to claim 6 or the cell according to any one of claims 7 to 11 and a pharmaceutically acceptable carrier, diluent or excipient.
- 20 13. A method for improving T cell function and/or T cell survival in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a chimeric receptor, a nucleic acid molecule, a vector or a cell according to any one of claims 1 to 11.
- 25 14. A chimeric receptor, a nucleic acid molecule, a vector or a cell according to any one of claim 1 to 11 for use in a method for improving T cell function and/or T cell survival in a subject.
- 30 15. The method or chimeric receptor, nucleic acid molecule, vector or cell for use according to claim 13 or 14, wherein said method comprises preventing or inhibiting T cell exhaustion.

16. A method of immunotherapy in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a chimeric receptor, a nucleic acid molecule, a vector or a cell according to any one of claims 1 to 11.

5

17. A chimeric receptor, a nucleic acid molecule, a vector or a cell according to any one of claim 1 to 11 for use in therapy, preferably immunotherapy,

18. The method or chimeric receptor, nucleic acid molecule, vector or cell for use according to claim 16 or 17 wherein said therapy or immunotherapy further comprises antibody-based immunotherapy.

10

19. The method or the chimeric receptor, nucleic acid molecule, vector or cell for use according to any one of claims 13 to 18 wherein said subject is suffering from cancer.

15

20. A method for enhancing efficacy of an antibody-based immunotherapy in a subject suffering from cancer and being treated with said antibody, the method comprising administering to the subject a therapeutically effective amount of T cells expressing the chimeric receptor according to any one of claims 1 to 4.

20

21. T cells expressing the chimeric receptor according to any one of claims 1 to 4 for use in a method for enhancing efficacy of an antibody-based immunotherapy in a subject suffering from cancer and being treated with said antibody.

25

22. A method of treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of T cells comprising a nucleic acid sequence encoding the chimeric receptor according to any one of claims 1 to 4.

30

23. T cells comprising a nucleic acid sequence encoding the chimeric receptor according to any one of claims 1 to 4 for use in a method of treating cancer in a subject.

24. The method or T cells for use according to claim 22 or 23 wherein said method comprises:

- isolating T cells from the subject;
- modifying said T cells by providing them with a nucleic acid sequence encoding
5 the chimeric receptor according to any one of claims 1 to 4;
- returning the modified T cells to the subject.

25. A method of producing a population of cells according to any one of claims 7 to 11, comprising

- 10 - providing cells, preferably human T-cells,
- providing said cells with a nucleic acid molecule or vector according to claim 5 or 6, and
- allowing expression of the chimeric antigen receptor according to any one of
15 claims 1 to 4.




Figure 1




Figure 2




Figure 3




Figure 3 continued




Figure 4




Figure 4 continued




Figure 5 continued




Figure 6




Figure 7

```

1   mppllapllc lallpalaar gprcsqpget clnggkceaa ngteacvcgg afvgprcqdp
61  npc1stpckn agtchvvdrr gvadyacsca lgfsgplclt pldnacltnp crnggtcdll
121 tlteykrccp pgwsgkscqq adpcasnpc nngqclpfea syichcpps f hgptcrqdv n
181 ecgqkpglcr hgg tchnevg syrcvrath tgpncerpyv pcspspcng g tcrptgdvt
241 hecac1pgft gqnceenidd cpgnncngg acvdgvntyn croppewtgg yctedvdecq
301 lmpnacqngg tchnthggyn cvcvngwtge dcseniddca saacfhgac hdrvasfyce
361 cphgrtgllc hlndacisnp cnegsnctdn pvngkaictc psgy tgpacs qdvdecslga
421 npcehagkci ntlgsfecqc lqgytgprce idvnecvsnp cqndatcldq igefqciemp
481 gyegvhcevn tdecasspcl hngrcl dkin efqcecptgf tghlcqydvd ecastpckng
541 akcldgpnty tcvctegy t thcevdidec dpdpchygsc kdgvatftcl crpgy tghhc
601 etninecssq pcrhgg tccq rdnaylcfcl kgttgpncei nlddcasspc dsgtcl dki d
661 gycacepgy tgs mcninid ecagnpchn g gtcedgingf tcrcepyhd ptclsevne c
721 nsnpcvhgac rds lngykcd cdp gswgtnc dinnnecesn pcvnggtckd mtsgyvctcr
781 egfsgpncqt ninecasnpc lnggtciddv agykcnc1lp ytgatcevvl apcasp crn
841 ggecrqsedy esfscvcptg wgggtcevd i necvlspcrh gascqnthgg yrchc qagys
901 grncetdidd crpnpc hngg sctd gintaf cdclpgfrgt fceedin eca sdpcrnganc
961 tdcvdsytct cpagfsgihc enntpdctes scfnggtcvd ginsftclcp pgftg sycqh
1021 dvnecdspc lhgg tccqdc gsyrc tccpg ytgpncqnlv hwc dsspckn g gkewqhtq
1081 yrcecpsgwt glycdvpsvs cevaaqrqgv dvarlcqhgg lcvdagnthh crcqagytgs
1141 ycedlvdecs pspc qngatc tdy lggysck cvagyhg vnc seeideclsh pcqnggtcl d
1201 lpntykscsp rgtqgvhcei nvddcnppvd pvsrspkcf n ngtcvdqvgg ysctcp pggfv
1261 gercegdvne clsnpcdarg tqncvqrvnd fhcecraght grrcesving ck gkpc kngg
1321 tcavasntar gfickcpagf egatcendar tcgslrclng gtcisgprsp tclclgpftg
1381 pecqfpassp clgg npcynq gtceptsesp fyrclcpakf ngllchildy sfgggagr di
1441 ppplieeace lpecqedagn kvcs lqcnnh acgwdggdcs lnfn dpwkcnc tqslqcwkyf
1501 sdghcdsqcn sagclfdgfd cqraegqcnp lyd qyckdhf sdghcdqgc n saecewdgld
1561 caehvperla agtlvvvvlm ppeqlrnssf hflrelsrvl ht n vfk rda hqqmifpyy
1621 greelrkhp ikraaegwaa pdallgqvka sllpggseg g rrrreldpmd vrgsivylei
1681 dnrqcvqass qcfqsatdva aflgalaslg slnipykiea vqsetveppp paqlhfmyva
1741 aaafvllffv gcgvllsrkr rrqh gglwfp egfkvseask kkrreplged svglkplkna
1801 sdgalmdnq newgdedlet kkfrfeepvv lpdlddqt dh rwtqqhlda adlrmsamap
1861 tppqgevdad cmdvnvrgpd gftplmiasc sgggletgns eeedapavi sdfiyqgasl
1921 hnqtdrtget alhlaarysr sdaakrllea sadaniqdnm grtplhaavs adaqgvf qil
1981 irnratdlda rmhdgttpli laarlavegm ledlinshad vnavddlgks alhwaavnn
2041 vdaavvllkn gankdmqnr eetplflaar eg syetakvl ldhfanrdit dhmdrlprdi
2101 aqermhdiv rlldeynlvr spqlhgaplg gtptlspplc spngylgslk pgvqgkvrk
2161 psskg lacgs keakdlkarr kksqd gkgcl ldssgmlspv dslesphgyl sdvasppllp
2221 spfqqspsvp lnhlpgmpdt hlgighlnva akpemaalgg ggrlafetgp prlshlpvas
2281 gtstvl gsss ggalnftv gg stslngqcew lsrlqsgmvp nqynplrgsv appplstqap
2341 slqhgvmgpl hsslaasals qmmsyqglps trlatqphlv qtqqvqpqn l qmqqqnlqpa
2401 niqqqqslqp pppppqphlg vssaasghlg rsflsgepsq advqplgpss lavhtilp qe
2461 spalptslps slvppvtaa q fltpps qhsy sspvdntpsh qlqvpehpfl tpspespdqw
2521 sssps hsnvs dwsegvsspp tsmqsqiari peafk

```

Figure 8




Figure 9




Figure 9 continued




Figure 10

B

CD19-Ig




Figure 10 continued