



US00PP29719P2

(12) **United States Plant Patent**
Volmary

(10) **Patent No.:** **US PP29,719 P2**

(45) **Date of Patent:** **Oct. 2, 2018**

(54) **THUNBERGIA PLANT NAMED ‘VOLTHU 7828’**

(50) Latin Name: *Thunbergia alata*
Varietal Denomination: **VOLTHU 7828**

(71) Applicant: **Hubertus Volmary**, Muenster (DE)

(72) Inventor: **Hubertus Volmary**, Muenster (DE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/731,225**

(22) Filed: **May 5, 2017**

(51) **Int. Cl.**
A01H 5/02 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./226**

(58) **Field of Classification Search**
USPC **Plt./226**
See application file for complete search history.

Primary Examiner — Susan Beth McCormick Ewoldt

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Thunbergia* plant named ‘VOLTHU 7828’, characterized by its upright and vining plant habit; freely branching habit; dense and bushy plant form; freely flowering habit; large red orange and red bi-colored flowers; and good garden performance.

2 Drawing Sheets

1

Botanical designation: *Thunbergia alata*.
Cultivar denomination: ‘VOLTHU 7828’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Thunbergia* plant, botanically known as *Thunbergia alata*, and hereinafter referred to by the name ‘VOLTHU 7828’.

The new *Thunbergia* plant is a product of a planned breeding program conducted by the Inventor in Muenster, Germany. The objective of the breeding program is to create new freely-flowering *Thunbergia* plants with unique flower colors and long flowering period.

The new *Thunbergia* plant originated from an open-pollination during the summer of 2013 in Muenster, Germany of *Thunbergia alata* ‘Sunny Suzy Red-Orange’, not patented, as the female, or seed, parent with an unknown selection of *Thunbergia alata* as the male, or pollen, parent. The new *Thunbergia* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated open-pollination in a controlled environment in Muenster, Germany during the summer of 2014.

Asexual reproduction of the new *Thunbergia* plant by terminal cuttings in a controlled greenhouse environment in Muenster, Germany since the late summer of 2014 has shown that the unique features of this new *Thunbergia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Thunbergia* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘VOLTHU

2

7828’. These characteristics in combination distinguish ‘VOLTHU 7828’ as a new and distinct *Thunbergia* plant:

1. Upright and vining plant habit.
2. Freely branching habit; dense and bushy plant form.
3. Freely flowering habit.
4. Large orange and red bi-colored flowers.
5. Good garden performance.

Plants of the new *Thunbergia* can be compared to plants of the female parent, ‘Sunny Suzy Red-Orange’. Plants of the new *Thunbergia* differ from plants of ‘Sunny Suzy Red-Orange’ primarily in the following characteristics:

1. Plants of the new *Thunbergia* are more compact than plants of ‘Sunny Suzy Red-Orange’.
2. Plants of the new *Thunbergia* have shorter leaf petioles than plants of ‘Sunny Suzy Red-Orange’.
3. Plants of the new *Thunbergia* are more freely flowering than plants of ‘Sunny Suzy Red-Orange’.
4. Plants of the new *Thunbergia* and ‘Sunny Suzy Red-Orange’ differ in flower color as plants of ‘Sunny Suzy Red-Orange’ have orange-colored flowers.

Plants of the new *Thunbergia* can be compared to plants of *Thunbergia alata* ‘Orange Wonder’ U.S. Plant Pat. No. 14,767. In side-by-side comparisons, plants of the new *Thunbergia* differ from plants of ‘Orange Wonder’ in the following characteristics:

1. Plants of the new *Thunbergia* are more compact than plants of ‘Orange Wonder’.
2. Leaves of plants of the new *Thunbergia* are darker green in color than leaves of plants of ‘Orange Wonder’.
3. Plants of the new *Thunbergia* are more freely flowering than plants of ‘Orange Wonder’.
4. Plants of the new *Thunbergia* have larger flowers than plants of ‘Orange Wonder’.
5. Plants of the new *Thunbergia* and ‘Orange Wonder’ differ in flower color as plants of ‘Orange Wonder’ have bright orange-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Thunbergia* plant showing the

colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Thunbergia* plant.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'VOLTHU 7828' grown in an outdoor nursery.

The photograph on the second sheet is a close-up view of a typical flowering plant of 'VOLTHU 7828'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the spring in three-liter containers in a glass-covered greenhouse and in an outdoor nursery in Muenster, Germany and under cultural practices typical of commercial *Thunbergia* production. During the production of the plants, day temperatures ranged from 18° C. to 20° C., night temperatures ranged from 15° C. to 18° C. and light levels ranged from 5,000 to 40,000 lux. Plants were pinched one time about three weeks after planting. Plants were 16 weeks old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2015 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Thunbergia alata* 'VOLTHU 7828'.
Parentage:

Female, or seed, parent.—*Thunbergia alata* 'Sunny Suzy Red-Orange', not patented.

Male or pollen parent.—Unknown selection of *Thunbergia alata*, not patented.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About one to two weeks at temperatures about 20° C. to 25° C.

Time to initiate roots, winter.—About two to three weeks at temperatures about 15° C. to 23° C.

Time to develop roots, summer.—About three weeks at temperatures about 20° C. to 25° C.

Time to develop roots, winter.—About 30 days at temperatures about 15° C. to 23° C.

Root description.—Fine, fibrous; typically yellowish white in color, actual color of the roots is dependent on substrate composition, water quality, fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; medium density.

Plant description:

Plant and growth habit.—Upright and vining plant habit; plants require support; moderately vigorous to vigorous growth habit; moderately rapid growth rate.

Branching habit.—Freely branching habit with lateral branches potentially developing at every node; pinching enhances lateral branch development; bushy and dense habit.

Plant height.—About 80 cm to 100 cm.

Plant diameter (area of spread).—About 35 cm.

Lateral branch description:

Length.—About 50 cm.

Diameter.—About 1.5 mm.

Internode length.—About 14.5 cm.

Texture.—Slightly pubescent.

Strength.—Strong, flexible.

Color.—Close to 143C tinged with close to N77A.

Leaf description:

Arrangement.—Opposite, simple.

Length.—About 10 cm.

Width.—About 7.5 cm.

Shape.—Roughly deltoid.

Apex.—Acute to cuspidate.

Base.—Sagittate.

Margin.—Entire.

Aspect.—Horizontal to somewhat downward.

Texture and luster, upper and lower surfaces.—Pubescent; dull.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Close to 146A to 146B. Developing leaves, lower surface: Close to 147A. Fully developed leaves, upper surface: Close to 146A to 146B; venation, close to 145C. Fully developed leaves, lower surface: Close to 137B to 137C; venation, close to 145C.

Petioles.—Length: About 3 cm. Diameter: About 3 mm. Texture, upper and lower surfaces: Pubescent. Color, upper and lower surfaces: Close to 146A.

Flower description:

Flower type and habit.—Single axillary salverform flowers; freely flowering habit with numerous flowers developing per plant; flowers face outward to slightly drooping.

Natural flowering season.—Plants flower continuously from April until frost in Germany; plants begin flowering about ten to twelve weeks after planting.

Flower longevity.—Flowers last about five days on the plant; flowers persistent.

Fragrance.—None detected.

Flower diameter.—About 4.5 cm to 5 cm.

Flower length (height).—About 4.5 cm to 5 cm.

Flower buds.—Length: About 3 cm. Diameter: About 1.8 cm. Shape: Lanceolate to deltoid. Color: Close to 146C.

Petals.—Quantity and arrangement: Corolla consists of five petals in a single whorl and fused towards the base. Length: About 1.8 cm. Width: About 1.6 cm. Shape: Broadly elliptic to rounded. Apex: Broadly emarginate. Base: Fused. Margin: Entire, occasionally incised. Texture and luster, upper and lower surfaces: Smooth, glabrous; matte. Color: When opening, upper surface: Close to N25C and 44B; throat, close to N77A. When opening, lower surface: Close to 26C; tube, close to N77A. Fully opened, upper surface: Close to N25C and 44B; throat, close to N77A; color becoming closer to 14A with development. Fully opened, lower surface: Close to 24C; tube, close to N77A.

Flower bracts.—Quantity and arrangement: Two, opposite. Length: About 3 cm. Width: About 1.8 cm. Shape: Spear-shaped. Apex: Obtuse. Base: Cordate. Margin: Entire. Texture and luster, upper and lower surfaces: Slightly pubescent; matte. Color, upper and lower surfaces: Close to 143B.

Sepals.—Sepal development has not been observed on plants of the new *Thunbergia*.

Peduncles.—Length: About 2.3 cm. Diameter: About 1 mm. Angle: About 15° from stem axis. Strength: Strong, flexible. Texture: Densely pubescent. Color: Close to 143C.

Reproductive organs.—Androecium: Stamen number: About four. Filament length: About 5 mm. Filament color: Close to 11D. Anther length: About 1 cm. Anther shape: Ovate. Anther color: Close to 155D. Amount of pollen: None observed. Gynoecium: Pistil number: One. Style length: About 1.7 cm. Style color: Close to NN155D. Stigma appearance: Biparted. Stigma color: Close to NN155D. Ovary color: Close to NN155D.

Seeds and fruits.—Seed and fruit development has not been observed on plants of the new *Thunbergia* to date.

Garden performance: Plants of the new *Thunbergia* have been observed to have good garden performance and to tolerate rain, wind and temperatures ranging from about 5° to about 38° C.

5 Pathogen & pest resistance: Plants of the new *Thunbergia* have not been shown to be resistant to pathogens and pests common to *Thunbergia* plants.

It is claimed:

10 1. A new and distinct *Thunbergia* plant named 'VOLTHU 7828' as illustrated and described.

* * * * *



