



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2223176 C 2007/08/14

(11)(21)

2 223 176

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1996/06/03
(87) Date publication PCT/PCT Publication Date: 1996/12/19
(45) Date de délivrance/Issue Date: 2007/08/14
(85) Entrée phase nationale/National Entry: 1997/12/03
(86) N° demande PCT/PCT Application No.: US 1996/008871
(87) N° publication PCT/PCT Publication No.: 1996/040206
(30) Priorité/Priority: 1995/06/07 (US08/472,219)

(51) Cl.Int./Int.Cl. *C07K 7/06* (2006.01),
A61K 38/08 (2006.01), *A61K 38/33* (2006.01),
C07K 14/665 (2006.01), *C07K 5/103* (2006.01),
C07K 5/107 (2006.01), *A61K 38/00* (2006.01)
(72) Inventeurs/Inventors:
DOOLEY, COLETTE T., US;
HOUGHTEN, RICHARD A., US
(73) Propriétaire/Owner:
TORREY PINES INSTITUTE FOR MOLECULAR
STUDIES, US
(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : NOUVEAUX PEPTIDES OPIOIDES SELECTIFS VIS-A-VIS DU RECEPTEUR KAPPA
(54) Title: NOVEL KAPPA RECEPTOR SELECTIVE OPIOID PEPTIDES

(57) **Abrégé/Abstract:**

The present invention provides novel opioid peptides which are selective for the kappa opiate receptor. In one embodiment, the kappa-selective opioid peptides have the general structure Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂, (SEQ ID NO. 28), wherein A1 is Tyr or Arg, B2 is Arg or Phe, and C3 is Thr, Phe, or Met. In yet other embodiments, the kappa-selective opioid peptides have the general structure (D)Phe-D4-E5-F6 (SEQ ID NO. 29) or (D)Nle-D4-E5-F6 (SEQ ID NO. 30), where in both of these formulae D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/07, 38/08, C07K 5/103, 5/107, 7/06	A1	(11) International Publication Number: WO 96/40206 (43) International Publication Date: 19 December 1996 (19.12.96)
--	-----------	---

(21) International Application Number: PCT/US96/08871

(22) International Filing Date: 3 June 1996 (03.06.96)

(30) Priority Data:
08/472,219 7 June 1995 (07.06.95) US

(71) Applicant: TORREY PINES INSTITUTE FOR MOLECULAR STUDIES [US/US]; 3550 General Atomics Court, San Diego, CA 92121 (US).

(72) Inventors: DOOLEY, Colette, T.; 844 Sapphire Street, San Diego, CA 92109 (US). HOUGHTEN, Richard, A.; 4939 Rancho Viejo, Del Mar, CA 92014 (US).

(74) Agents: PERKINS, Susan, M. et al.; Campbell & Flores L.L.P., Suite 700, 4370 La Jolla Village Drive, San Diego, CA 92122 (US).

(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published
With international search report.

(54) Title: NOVEL KAPPA RECEPTOR SELECTIVE OPIOID PEPTIDES

(57) Abstract

The present invention provides novel opioid peptides which are selective for the kappa opiate receptor. In one embodiment, the kappa-selective opioid peptides have the general structure Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂, (SEQ ID NO. 28), wherein A1 is Tyr or Arg, B2 is Arg or Phe, and C3 is Thr, Phe, or Met. In yet other embodiments, the kappa-selective opioid peptides have the general structure (D)Phe-D4-E5-F6 (SEQ ID NO. 29) or (D)Nle-D4-E5-F6 (SEQ ID NO. 30), where in both of these formulae D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla.

NOVEL KAPPA RECEPTOR SELECTIVE OPIOID PEPTIDES

BACKGROUND OF THE INVENTIONFIELD OF THE INVENTION

The present invention relates generally to the
5 field of peptide chemistry and, more specifically, to
novel opioid peptides that can inhibit ligand binding to
an opioid receptor.

BACKGROUND INFORMATION

There are at least three known subtypes of
10 opioid receptors, mu (μ), delta (δ), and kappa (κ), to
which morphine, the enkephalins, and the dynorphins,
respectively, bind. The three receptor subtypes possess
analgesic properties. However, the type of pain
inhibited and the secondary functions vary with each
15 receptor type. The κ receptor is potent in affecting
analgesia in response to pain, including chemical
stimuli. The κ receptor also induces diuresis and
sedation and is associated with dysphoric and
psychometric effects as well as drug or other physical
20 dependence. The μ receptor is generally regarded as the
one associated with pain relief, respiratory depression,
intestinal motility, antidiuresis, an immune response,
and drug or other physical dependence. The δ receptor,
on the other hand, is associated with thermal analgesia
25 and, to a lesser extent, respiration and addiction.
These differences in the opioid receptor functions

encourage the search for drugs which produce analgesia without deleterious side effects.

The use of synthetic peptides has been instrumental in the delineation of these subtypes and in providing analogues that can be used for studying the interactions of ligands specific to these receptor systems in both *in vitro* and *in vivo* systems. Certain opioid compounds are agonists (bind to the receptor and produce an effect) while others are antagonists (bind to the receptor but do not produce an effect). Most previously known agonists and antagonists of the opioid receptors are analogues of the enkephalins and related peptides, including the dynorphins, the dermenkephalins and the casomorphins. The compounds of the present invention have little to no sequence homology with any of these known opioid peptides.

Recent advances in methods for the preparation and screening of large numbers of individual peptides have led to the identification of numerous peptides useful in all areas of biomedical research, including research regarding the interaction of a ligand to the opiate receptor. Both receptor-specific agonists and antagonists are needed as pharmacological tools and as therapeutic agents. Even with these advances, however, basic research and drug discovery has been limited by the availability of the requisite large number of diverse opiate agonists and antagonists required to ascertain the relationship between a ligand for a particular opiate receptor subtype. Thus, a need exists for large numbers

of individual peptides for use in biomedical research, including those for the study of opiate ligand-receptor interactions. As well there is a need for opioid peptides which have therapeutic value. This invention
5 satisfies these needs and provides related advantages as well.

SUMMARY OF THE INVENTION

The present invention provides novel opioid peptides which are selective for the κ opiate receptor.
10 In one embodiment, the opioid peptides have the general structure Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂, (SEQ ID NO. 28), wherein A1 is Tyr or Arg, B2 is Arg or Phe, and C3 is Thr, Phe, or Met. In yet other embodiments, the opioid peptides have the general
15 structure (D)Phe-D4-E5-F6 (SEQ ID NO. 29) or (D)Nle-D4-E5-F6 (SEQ ID NO. 30), where in both of these formulae D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla.

DETAILED DESCRIPTION OF THE INVENTION

20 The present invention provides novel opioid peptides which are capable of inhibiting the binding of the κ -selective opioid ligand [³H]-U69,593. In one embodiment, the peptides have the general structure Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂, (SEQ ID NO.
25 28), wherein A1 is Tyr or Arg, B2 is Arg or Phe, and C3 is Thr, Phe, or Met. Exemplary peptides encompassed by this formula, include the following: Ac-Tyr-Arg-Thr-Arg-

Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂ (SEQ ID NO. 1); Ac-Tyr-Arg-Phe-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂ (SEQ ID NO. 2); Ac-Arg-Phe-Phe-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂ (SEQ ID NO. 3); Ac-Tyr-Arg-Met-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂ (SEQ ID NO. 4); and Ac-Arg-Arg-Phe-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂ (SEQ ID NO. 5).

In another embodiment, the novel peptides are those encompassed by the formula (D)Phe-D4-E5-F6 (SEQ ID NO. 29), wherein D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla. Exemplary peptides within this embodiment have any one of the following sequences: (D)Phe-(D)NapAla-(D)Nle-(D)Arg (SEQ ID NO. 6); (D)Phe-(D)NapAla-(D)Nle-(D)ChAla (SEQ ID NO. 7); (D)Phe-(D)NapAla-Trp-(D)Arg (SEQ ID NO. 8); (D)Phe-(D)NapAla-Trp-(D)ChAla (SEQ ID NO. 9); (D)Phe-(D)NapAla-(D)Ile-(D)Arg (SEQ ID NO. 10); (D)Phe-(D)NapAla-(D)Ile-(D)ChAla (SEQ ID NO. 11); (D)Phe-(D)Phe-(D)Nle-(D)Arg (SEQ ID NO. 12); (D)Phe-(D)Phe-(D)Nle-(D)ChAla (SEQ ID NO. 13); (D)Phe-(D)Phe-Trp-(D)Arg (SEQ ID NO. 14); (D)Phe-(D)Phe-Trp-(D)ChAla (SEQ ID NO. 15); (D)Phe-(D)Phe-(D)Ile-(D)Arg (SEQ ID NO. 16); and (D)Phe-(D)Phe-(D)Ile-(D)ChAla (SEQ ID NO. 17).

In yet another embodiment of the invention, the peptides have the structure (D)Nle-D4-E5-F6 (SEQ ID NO. 30), wherein D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla. Exemplary peptides which fall within the scope of this formula include the following sequences: (D)Nle-(D)NapAla-(D)Nle-(D)Arg (SEQ ID NO. 18); (D)Nle-(D)NapAla-(D)Nle-(D)ChAla

(SEQ ID NO. 19); (D)Nle-(D)NapAla-Trp-(D)Arg (SEQ ID NO. 20); (D)Nle-(D)NapAla-(D)Ile-(D)Arg (SEQ ID NO. 21); (D)Nle-(D)NapAla-(D)Ile-(D)ChAla (SEQ ID NO. 22); (D)Nle-(D)Phe-(D)Nle-(D)Arg (SEQ ID NO. 23); (D)Nle-(D)Phe-Trp-(D)Arg (SEQ ID NO. 24); (D)Nle-(D)Phe-Trp-(D)ChAla (SEQ ID NO. 25); (D)Nle-(D)Phe-(D)Ile-(D)Arg (SEQ ID NO. 26); and (D)Nle-(D)Phe-(D)Ile-(D)ChAla (SEQ ID NO. 27).

The following standard abbreviations are used herein to identify amino acid residues.

10

	Amino Acid	Three-letter Abbreviation	One-letter Symbol
	Alanine	Ala	A
15	Arginine	Arg	R
	CyclohexylAlanine	ChAla	-
	Isoleucine	Ile	I
	Leucine	Leu	L
	Methionine	Met	M
20	NaphthylAlanine	NapAla	-
	NorLeucine	Nle	-
	Phenylalanine	Phe	F
	Threonine	Thr	T
	Tryptophan	Trp	W
25	Tyrosine	Tyr	Y

The amino acids are indicated by these commonly known codes as provided above and (D) designates an amino acid having the "D" configuration, as opposed to the

naturally occurring L-amino acids. Where no specific configuration is indicated, one skilled in the art would understand the amino acid to be an (L)-amino acid.

One skilled in the art would know that one or
5 more amino acids within the exemplified peptides could be modified or substituted, as for example, by a conservative amino acid substitution of one or more of the specific amino acids shown in the exemplified peptides. A conservative amino acid substitution change
10 can include, for example, the substitution of one basic amino acid for another basic amino acid, of one hydrophobic amino acid for another hydrophobic amino acid or other conservative substitutions well known in the art, including the use of non-naturally occurring amino
15 acids, such as ornithine (Orn) or homoArginine (homoArg) for Arg. Such substitutions include the corresponding (D)-amino acid where one in the (L) configuration is described, and visa-versa.

In addition to the above types of modifications
20 or substitutions, a mimic of one or more amino acids, otherwise known as a peptide mimetic or peptidomimetic, can also be used. As used herein, the term "mimic" means an amino acid or an amino acid analog that has the same or similar functional characteristic of an amino acid.
25 Thus, for example, a (D)arginine analog can be a mimic of (D)arginine if the analog contains a side chain having a positive charge at physiological pH, as is characteristic of the guanidinium side chain reactive group of arginine. A peptide mimetic or peptidomimetic is an organic

molecule that retains similar peptide chain pharmacophore groups as are present in the corresponding peptide.

The substitution of amino acids by non-naturally occurring amino acids and peptidomimetics as described above can enhance the overall activity or other properties of an individual peptide based on the modifications to the side chain functionalities. For example, these types of alterations to the exemplified peptides can enhance the peptide's stability to enzymatic breakdown or increase biological activity or decrease immunogenicity.

One skilled in the art, using the above formulae, can easily synthesize the peptides of this invention. Standard procedures for preparing synthetic peptides are well known in the art. The novel opioid peptides can be synthesized using the solid phase peptide synthesis (SPPS) method of Merrifield (J. Am. Chem. Soc., 85:2149 (1964)),

or modifications of SPPS, or the peptides can be synthesized using standard solution methods well known in the art (see, for example, 'Bodanzsky, M., Principles of Peptide Synthesis 2nd revised ed. (Springer-Verlag, 1988 and 1993)).

Alternatively, simultaneous multiple peptide synthesis (SMPS) techniques well known in the art can be used. Peptides prepared by the method of Merrifield can be synthesized using an automated peptide synthesizer such as the Applied Biosystems 431A-01 Peptide Synthesizer (Mountain View, CA) or using the manual

peptide synthesis technique described by Houghten, Proc. Natl. Acad. Sci. USA 82:5131 (1985).

Peptides can be synthesized using amino acids
5 or amino acid analogs, the active groups of which are
protected as necessary using, for example, a
t-butyldicarbonate (t-BOC) group or a fluorenylmethoxy
carbonyl (Fmoc) group. Amino acids and amino acid
analogues can be purchased commercially (Sigma Chemical
10 Co.; Advanced Chemtec) or synthesized using methods known
in the art. Peptides synthesized using the solid phase
method can be attached to resins including 4-methyl-
benzhydrylamine (MBHA), 4-(oxymethyl)-phenylacetamido
methyl and 4-(hydroxymethyl)phenoxymethyl-copoly(styrene-
15 1% divinylbenzene) (Wang resin), all of which are
commercially available, or to p-nitrobenzophenone oxime
polymer (oxime resin), which can be synthesized as
described by De Grado and Kaiser, J. Org. Chem. 47:3258
(1982).

20 In the exemplified peptides, "Ac" indicates an
acetyl group at the amino terminus and "NH₂" means an
amide group is at the carboxy terminus. Peptides can be
manipulated, for example, while still attached to a resin
to obtain N-terminal modified compounds such as an
25 acetylated peptide or can be removed from the resin using
hydrogen fluoride or an equivalent cleaving reagent and
then modified. Compounds synthesized containing the C-
terminal carboxy group (Wang resin) can be modified after
cleavage from the resin or, in some cases, prior to

solution phase synthesis. Methods for modifying the N-terminus or C-terminus such as methods for acetylation of the N-terminus or methods for amidation of the C-terminus are well known in the art.

5

A newly synthesized peptide can be purified using a method such as reverse phase high performance liquid chromatography (RP-HPLC), chromatofocusing, or other methods of separation based on the size or charge of the peptide, or by immuno-purification techniques. Furthermore, the purified peptide can be characterized using these and other well known methods such as amino acid analysis and mass spectrometry.

After manufacture, the peptides can be assayed for κ receptor binding activity using the κ radioreceptor assay as provided in Example I. For additional data on the specificity of the peptides, results from a κ radioreceptor assay can be compared with those of μ and δ radioreceptor assays as provided in Example II. To test the *in vitro* agonist or antagonist activity of the peptides, well known *in vivo* assays, such as the guinea-pig ileum assay as described in Example III can be routinely performed. In addition, several other reliable *in vivo* tests are known in the art which are useful for testing the opioid peptides and compositions of the present invention. Exemplary tests include the mouse tail-flick, tail-pinch and HCl writhing tests, as well as the rat warm-plate and hot plate assays, and rat tail-immersion, tail-flick and air writhing assays, all of which are described in Vonvoigtlander et al., Journal of

30

Pharmacology and Experimental Therapeutics, 224(1):7-12 (1983). One

of the more commonly used assays of those described above, the mouse tail-flick *in vivo* animal model, is also
5 described by Dixon et al., Drug Res., 27:1968 (1977),
Jiang et al., J. Pharmacol. Exp. Ther., 262:526 (1992),
and Dooley et al., Science, 266:2019 (1994).

Because the peptides of the present invention
10 bind to the κ receptor, they can be used in *in vitro*
assays to study the opiate receptor subtypes. For
example, in a sample receptor of unknown type or origin,
the peptides, after being labeled with a detectable
marker such as a radioisotope, can be contacted with the
15 receptor sample under conditions which specifically favor
binding to a particular receptor subtype, such as the κ
subtype. Unbound receptor and peptide can be removed,
for example, by washing with a saline solution, and bound
receptor can then be detected using methods well known to
20 those skilled in the art. Therefore, the peptides of the
present invention are useful *in vitro* for the diagnosis
of relevant opioid receptor subtypes, and in particular
the κ type, in brain and other tissue samples.

In addition to their utility in *in vitro*
25 screening methods, the peptides are also useful *in vivo*.
For example, the opioid peptides can be used *in vivo*
diagnostically to localize opioid receptor subtypes. The
peptides are also useful as drugs, namely as analgesics,
or to treat pathologies associated with other compounds

which interact with the opioid receptor system.
Therefore, the peptides of the present invention can be used in medicaments for treating pathologies associated with the κ receptor and as analgesics. For example, it
5 can be envisioned that these peptides can be used for therapeutic purposes to block the peripheral effects of a centrally acting pain killer. For instance, morphine is a centrally acting pain killer. Morphine, however, has a number of deleterious effects in the periphery which are
10 not required for the desired analgesic effects, such as constipation and pruritus (itching). While it is known that the many peptides do not readily cross the blood-brain barrier and, therefore, elicit no central effect, the subject peptides can have value in blocking the
15 periphery effects of morphine, such as constipation and pruritus.

The novel peptides claimed can be incorporated into pharmaceutical compositions. Pharmaceutically acceptable carriers are well known in the art and include
20 aqueous solutions such as physiologically buffered saline or other buffers or solvents or vehicles such as glycols, glycerol, oils such as olive oil or injectable organic esters. A pharmaceutically acceptable carrier can contain physiologically acceptable compounds that act,
25 for example, to stabilize the opioid peptide or increase the absorption of the peptide. Such physiologically acceptable compounds include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low
30 molecular weight proteins or other stabilizers or

excipients. One skilled in the art would know that the choice of a pharmaceutically acceptable carrier, including a physiologically acceptable compound, depends, for example, on the route of administration and on the particular physio-chemical characteristics of the specific opioid peptide.

Methods of administering a pharmaceutical are well known in the art and include but are not limited to administration orally, intravenously, intramuscularly or intraperitoneally. Administration can be effected continuously or intermittently and will vary with the subject and is dependent on the type of treatment and potency of the peptide used.

The following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Identification Of Kappa Selective Opioid Peptides By Radioreceptor Assay

This example describes the identification of individual peptides as inhibitors of the κ -selective ligand, [^3H]-U69,593. Individual peptides were identified as capable of inhibiting [^3H]-U69,593 by a radioreceptor assay.

Synthetic combinatorial libraries (SCLs) made up of mixtures of tens of millions of different peptides can be used to rapidly identify individual, active

compounds. Since the libraries are in solution (i.e., not attached to a bead, pin, phage, glass, etc.) they can be screened in virtually any assay system. All opioid peptides were initially prepared and contained within a positional scanning synthetic combinatorial library (PS-SCL). The libraries were used in conjunction with an iterative selection process to identify individual peptides capable of inhibiting [³H]-U69,593 in the radioreceptor assay.

10. As detailed below, the peptide libraries were screened in a radioreceptor assay using rat brain homogenates and [³H]-U69,593 as radioligand and IC_{50} values determined for peptides which significantly inhibited the binding of [³H]-U69,593.

15 Radioreceptor Assays Selective For The Kappa Receptor

Guinea pig brains, frozen in liquid nitrogen, were obtained from Harlan Bioproducts for Science (Indianapolis, IN). Frozen brains were thawed, the cerebella and cortex were removed and weighed. The combined cortex and cerebella were homogenized in 40 ml Tris-HCl buffer (50 mM, pH 7.4, 4°C) containing 100 μ M phenylmethylsulfonyl fluoride (PMSF), 5 mM MgCl₂, and 1 mg/ml bovine serum albumin (BSA) and centrifuged (39000 x g) (Model J2-HC; Beckman Instruments, Fullerton, CA) for 10 min at 4°C. The pellets were resuspended in fresh Tris-HCl buffer and incubated at 37°C for 40 min. Following incubation, the suspensions were centrifuged as above, the resulting pellets resuspended in 100 volumes

of Tris buffer and the suspensions combined. Membrane suspensions were prepared and used in the same day. Protein content of the crude homogenates ranged from 0.1-0.3 mg/ml as determined using the method described by
5 Bradford in Anal. Biochem. 72:248-254 (1976).

Binding assays were carried out in polypropylene tubes. Each tube contained 3 nM of the κ -selective ligand [^3H]-U69,593 (specific activity 62
10 Ci/mmol) (Amersham; Arlington Heights, IL), peptide mixture (a range of 3×10^4 to 3×10^5 M concentration for individual peptides in the mixture), and tissue homogenates prepared from guinea pig brains (cortex and cerebellum) using Tris buffer containing 100 μM PMSF, 5
15 mM MgCl_2 , and 1 mg/ml BSA, pH 7.4. Assay tubes were incubated for 2.5 hr at 25°C. The reaction was terminated by filtration through GF-B filters (Wallac, Gaithersburg, MD). The filters were subsequently washed with 6 ml Tris-HCl buffer at 4°C. Bound radioactivity
20 was counted on a Beta-plate Liquid Scintillation Counter (Life Technologies, Gaithersburg, MD) and expressed in counts per minute (cpm). Standard curves were prepared using 0.05-6300 nM naloxone (Sigma, St. Louis, MO). Competitive inhibition assays were performed as above
25 using serial dilutions of the peptide mixtures. IC_{50} values were then calculated using the software GRAPHPAD (ISI, San Diego, CA). The average standard deviation for IC_{50} values was $\pm 20\%$.

15

IC₅₀ values of less than 1000 nM are indicative of highly active opioid peptides which bind to the κ receptor, with particularly active compounds having IC₅₀ values of 100 nM or less, and the most active compounds 5 with values of less than 10 nM.

Opioid peptides identified from the assay screening of one SCL have the general structure Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂, (SEQ ID NO. 28), wherein A1 is Tyr or Arg, B2 is Arg or Phe, and C3 is 10 Thr, Phe, or Met. IC₅₀ values of peptides falling within the scope of this formula are provided in Table 1.

Table 1		
Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH ₂ (SEQ ID NO. 28)		
SEQ ID NO.	PEPTIDE	IC ₅₀ (nM)
1	Ac-Y-R-T-R-Y-R-Y-R-R-R-NH ₂	10
2	Ac-Y-R-F-R-Y-R-Y-R-R-R-NH ₂	18
3	Ac-R-F-F-R-Y-R-Y-R-R-R-NH ₂	22
4	Ac-Y-R-M-R-Y-R-Y-R-R-R-NH ₂	36
5	Ac-R-R-F-R-Y-R-Y-R-R-R-NH ₂	142

Also identified as κ -selective peptides are those of the general formula (D)Phe-D4-E5-F6 (SEQ ID NO. 29), wherein D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, 25 Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla. Table 2

provides the respective IC₅₀ values for each of the peptides within this formula.

Table 2		
(D) Phe-D4-E5-F6 (SEQ ID NO. 29)		
SEQ ID NO.	PEPTIDE	IC ₅₀ (nM)
6	(D) Phe- (D) NapAla- (D) Nle- (D) Arg	0.2
7	(D) Phe- (D) NapAla- (D) Nle- (D) ChAla	8
8	(D) Phe- (D) NapAla-Trp- (D) Arg	7584
9	(D) Phe- (D) NapAla-Trp- (D) ChAla	24984
10	(D) Phe- (D) NapAla- (D) Ile- (D) Arg	0.7
11	(D) Phe- (D) NapAla- (D) Ile- (D) ChAla	86
12	(D) Phe- (D) Phe- (D) Nle- (D) Arg	0.8
13	(D) Phe- (D) Phe- (D) Nle- (D) ChAla	3
14	(D) Phe- (D) Phe-Trp- (D) Arg	4953
15	(D) Phe- (D) Phe-Trp- (D) ChAla	52430
16	(D) Phe- (D) Phe- (D) Ile- (D) Arg	2
17	(D) Phe- (D) Phe- (D) Ile- (D) ChAla	169

Peptides of the general formula (D)Nle-D4-E5-F6 (SEQ ID NO. 30), wherein D4 is (D)NapAla or (D)Phe, E5 is (D)Nle, Trp, or (D)Ile, and F6 is (D)Arg or (D)ChAla were also identified as inhibitors of [³H]-U69,593. The IC₅₀ values of peptides within this formula are provided in Table 3 below.

Table 3		
(D)Nle-D4-E5-F6 (SEQ ID NO. 30)		
SEQ ID NO.	PEPTIDE	IC ₅₀ (nM)
18	(D)Nle- (D)NapAla- (D)Nle- (D)Arg	3
19	(D)Nle- (D)NapAla- (D)Nle- (D)ChAla	6
20	(D)Nle- (D)NapAla-Trp- (D)Arg	743
21	(D)Nle- (D)NapAla- (D)Ile- (D)Arg	2
22	(D)Nle- (D)NapAla- (D)Ile- (D)ChAla	42
23	(D)Nle- (D)Phe- (D)Nle- (D)Arg	10
24	(D)Nle- (D)Phe-Trp- (D)Arg	10985
25	(D)Nle- (D)Phe-Trp- (D)ChAla	42846
26	(D)Nle- (D)Phe- (D)Ile- (D)Arg	16
27	(D)Nle- (D)Phe- (D)Ile- (D)ChAla	1447

The results of Tables 1 through 3 above provide three structurally unique series of opioid peptides which are all selective for the κ opioid receptor.

EXAMPLE II

Kappa Receptor Specificity Of Opioid Peptides

This example demonstrates how to obtain additional data on the specificity of the novel opioid peptides for the κ receptors as compared to the μ and δ opiate receptors.

Individual peptides can be synthesized by procedures well known in the art as discussed above. The

activity of the individual, synthesized peptides in radioreceptor assays selective for the μ and δ receptors, as detailed below, can be compared to the results of the radioreceptor assay selective for the κ receptors from
5 Example I.

Radioreceptor assays selective for μ receptors can be performed using rat brain homogenates as above. Sample polypropylene tubes are prepared so as to contain approximately 0.5 ml of membrane suspension, 3 nM of the
10 μ -selective opioid peptide [^3H]-DAMGO (specific activity 36 Ci/mmol), 0.08 mg/ml peptide mixture and Tris-HCl buffer in a total volume of 0.65 ml. Assay tubes are generally incubated for 60 min at 25°C and the reaction terminated by filtration through, for example, GF-B
15 filters. The filters are subsequently washed with 6 ml Tris-HCl buffer at 4°C. Bound radioactivity is then counted on, for example, a Beta-plate Liquid Scintillation Counter (Life Technologies, Gaithersburg, MD) and expressed in counts per minute (cpm). Standard
20 curves can be determined by incubation of [^3H]-DAMGO in the presence of 0.13-3900 nM of unlabeled DAMGO.

Radioreceptor assays selective for δ receptors can be performed using rat brain homogenates as above and [^3H]-naltrindole (0.5 nM, specific activity 34.7 Ci/mmol)
25 as radioligand in Tris buffer containing 100 μM phenylmethanesulfonyl fluoride (PMSF), 5 mM MgCl_2 and 1 mg/ml bovine serum albumin (BSA), pH 7.4. Samples are generally incubated for 2.5 hr. Standard curves are

prepared using 0.10-3200 nM [³H]-D-Pen₂, Pen₅]-enkephalin ([³H]-DPDPE). All other steps are as described above.

Tritiated ligands, [³H]-DAMGO and [³H]-DPDPE can be obtained from the National Institute on Drug Abuse (NIDA) repository, as prepared by Multiple Peptide Systems (San Diego, CA), and [³H]-naltrindole from DuPont NEN Research Products (Los Angeles, CA). The average standard deviation for IC₅₀ values was ±20%.

The comparative data in the respective κ, μ, and δ radioreceptor assays can further demonstrate the selectivity of the novel opioid peptides for the κ receptors over the μ and δ opiate receptors.

EXAMPLE III

Kappa Receptor Agonist Or Antagonist Activity

15 By Guinea-Pig Ileum Assay

This example demonstrates that certain novel opioid peptides can be identified as agonists or antagonists of the κ receptor by the guinea-pig ileum assay.

20 Guinea-Pig Ileum Assay

A guinea-pig ileum (GPI) bioassay can be carried out to determine whether a peptide is an opioid agonist. Again, such information cannot be determined from the above-described radioreceptor binding assays.

The GPI assay (Kosterlitz et al., Br. J. Pharmacol. 39:398-418 (1970)),

is one of the most widely used assays for the determination of opioid activities in vitro. The assay is based on the ability of opioid agonists to inhibit electrically stimulated contraction in tissue. GPI is not a "clean" assay insofar as the ileum preparation contains both μ and κ receptors. However, κ receptor mediated effects in the GPI can be distinguished from μ receptor mediated effects by determining K_e values for the antagonist (κ effects: $K_e \sim 20-30$ nM; μ effects: $K_e \sim 1-3$ nM) or by blocking μ receptors with a specific μ antagonist, such as CTAP ((D)Tca-Cys-Tyr-(D)Trp-Arg-Thr-Pen-Thr-NH₂, wherein Tca is a cyclic (D)Trp analogue).

The guinea pigs are killed by decapitation, the ileum removed and longitudinal muscle-myenteric plexus preparations placed in Krebs solution (NaCl 118 mM, KCl 4.75 mM, NaH₂PO₄ 1 mM, NaHCO₃ 25 mM, MgSO₄ 1.2 mM, glucose 11 mM and CaCl₂ 2.5 mM). The solution is gassed with 95% O₂ and 5% CO₂ and maintained at 37°C. The tissue is suspended under a final tension of 1 g in a 10 ml organ bath and stabilized for 1 hr. Electrical stimulation via a straight platinum electrode is applied, 0.4 ms pulses of supra maximal voltage, delivered at a rate of 0.1 Hz. Isometric contractions are then measured via strain gauge force transducers and recorded on stripchart recorders.

Although the invention has been described with reference to the examples provided above, it should be understood that various modifications can be made by those skilled in the art without departing from the invention. Accordingly, the invention is set out in the following claims.

WE CLAIM:

1. A peptide consisting of the structure:

Ac-A1-B2-C3-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂,
(SEQ ID NO. 28)

wherein A1 is Tyr or Arg;

B2 is Arg or Phe; and

C3 is Thr, Phe, or Met.

2. The peptide of claim 1, consisting of a sequence
selected from the group consisting of:

Ac-Tyr-Arg-Thr-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂
(SEQ ID NO. 1);

Ac-Tyr-Arg-Phe-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂
(SEQ ID NO. 2);

Ac-Arg-Phe-Phe-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂
(SEQ ID NO. 3);

Ac-Tyr-Arg-Met-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂
(SEQ ID NO. 4);

and

Ac-Arg-Arg-Phe-Arg-Tyr-Arg-Tyr-Arg-Arg-Arg-NH₂
(SEQ ID NO. 5).