



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C12N 15/12, 15/64, C12P 21/08</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/21988</p> <p>(43) International Publication Date: 6 May 1999 (06.05.99)</p>
<p>(21) International Application Number: PCT/CN97/00115</p> <p>(22) International Filing Date: 29 October 1997 (29.10.97)</p> <p>(71) Applicant (for all designated States except US): SHANGHAI SECOND MEDICAL UNIVERSITY [CN/CN]; 197 Rui-Jin Road II, Shanghai 200025 (CN).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): WU, Ji-Sheng [CN/CN]; Room 302, No. 33, Lane 230, Guan Sheng Yuan Road, Shanghai 200233 (CN). WANG, Ya-Xin [CN/CN]; Room 1209, 381 He Fei Road, Shanghai 200025 (CN).</p> <p>(74) Agent: CHINA PATENT AGENCY (H.K.) LTD.; 22/F, Great Angle Centre, 23 Harbour Road, Wanchai, Hong-kong (CN).</p>		<p>(81) Designated States: CA, CN, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: THE HUMAN VESICLE TRAFFICKING PROTEIN SEC22b GENE OF CBFBBA01</p>		
<p>(57) Abstract</p> <p>CBFBBA01 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing CBFBA01 polypeptides and polynucleotides in the design of protocols for the treatment of cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, among others, and diagnostic assays for such conditions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

THE HUMAN VESICLE TRAFFICKING PROTEIN SEC22b GENE OF CFBFBA01**FIELD OF INVENTION**

5 This invention relates to newly identified polynucleotides, polypeptides encoded by them and to the use of such polynucleotides and polypeptides, and to their production. More particularly, the polynucleotides and polypeptides of the present invention relate to the SEC22b family, hereinafter referred to as CFBFBA01. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

10

BACKGROUND OF THE INVENTION

 The murine sec22b protein is a cytoplasmically-oriented integral membrane protein located in the ER/Golgi membrane and represents a vesicle-trafficking intermediate in ER-to-Golgi transfer reactions. This indicates that the SEC22b family has an established, proven history as therapeutic
15 targets. Clearly there is a need for identification and characterization of further members of the family which can play a role in preventing, ameliorating or correcting dysfunctions or diseases, including, but not limited to, cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis.

SUMMARY OF THE INVENTION

20 In one aspect, the invention relates to CFBFBA01 polypeptides and recombinant materials and methods for their production. Another aspect of the invention relates to methods for using such CFBFBA01 polypeptides and polynucleotides. Such uses include the treatment of cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the
25 invention, and treating conditions associated with CFBFBA01 imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate CFBFBA01 activity or levels.

DESCRIPTION OF THE INVENTION**30 Definitions**

 The following definitions are provided to facilitate understanding of certain terms used frequently herein.

 "CBFBA01" refers, among others, generally to a polypeptide having the amino acid sequence set forth in SEQ ID NO:2 or an allelic variant thereof.

"CBFBBA01 activity or CBFBBA01 polypeptide activity" or "biological activity of the CBFBBA01 or CBFBBA01 polypeptide" refers to the metabolic or physiologic function of said CBFBBA01 including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said

5 CBFBBA01.

"CBFBBA01 gene" refers to a polynucleotide having the nucleotide sequence set forth in SEQ ID NO:1 or allelic variants thereof and/or their complements.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab
10 or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting
15 materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and
20 double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified"
25 bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

30 "Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by

natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Posttranslational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62.

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or

more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and
5 polypeptides may be made by mutagenesis techniques or by direct synthesis.

“Identity” is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. “Identity” *per se* has an art-recognized meaning and can be calculated using published techniques. See, e.g.:
(COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New
10 York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W.,
ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA,
PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE
ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and
SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press,
15 New York, 1991). While there exist a number of methods to measure identity between two
polynucleotide or polypeptide sequences, the term “identity” is well known to skilled artisans
(Carrillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly
employed to determine identity or similarity between two sequences include, but are not limited to,
those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego,
20 1994, and Carrillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073. Methods to
determine identity and similarity are codified in computer programs. Preferred computer program
methods to determine identity and similarity between two sequences include, but are not limited to,
GCS program package (Devereux, J., *et al.*, *Nucleic Acids Research* (1984) 12(1):387), BLASTP,
BLASTN, FASTA (Atschul, S.F. *et al.*, *J Molec Biol* (1990) 215:403).

25 As an illustration, by a polynucleotide having a nucleotide sequence having at least, for
example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 is intended that the
nucleotide sequence of the polynucleotide is identical to the reference sequence except that the
polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the
reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having
30 a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the
nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a
number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted
into the reference sequence. These mutations of the reference sequence may occur at the 5 or 3
terminal positions of the reference nucleotide sequence or anywhere between those terminal

positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

15

Polypeptides of the Invention

In one aspect, the present invention relates to CBFBBA01 polypeptides (or CBFBBA01 proteins). The CBFBBA01 polypeptides include the polypeptide of SEQ ID NO:2; as well as polypeptides comprising the amino acid sequence of SEQ ID NO: 2; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Also included within CBFBBA01 polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and still more preferably at least 95% identity to SEQ ID NO:2. Furthermore, those with at least 97-99% are highly preferred. Preferably CBFBBA01 polypeptide exhibit at least one biological activity of CBFBBA01.

The CBFBBA01 polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

Fragments of the CBFBBA01 polypeptides are also included in the invention. A fragment is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino

acid sequence of the aforementioned CBFBBA01 polypeptides. As with CBFBBA01 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-5 80, 81-100, and 101 to the end of CBFBBA01 polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of CBFBBA01 polypeptides, except for deletion of a continuous series of residues that 10 includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, 15 hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate CBFBBA01 activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

20 Preferably, all of these polypeptide fragments retain the biological activity of the CBFBBA01, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and 25 Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

The CBFBBA01 polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, 30 synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the Invention

Another aspect of the invention relates to CBFBBA01 polynucleotides. CBFBBA01 polynucleotides include isolated polynucleotides which encode the CBFBBA01 polypeptides and fragments, and polynucleotides closely related thereto. More specifically, CBFBBA01 polynucleotide of the invention include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 encoding a CBFBBA01 polypeptide of SEQ ID NO: 2, and polynucleotide having the particular sequence of SEQ ID NO:1. CBFBBA01 polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the CBFBBA01 polypeptide of SEQ ID NO:2, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to of SEQ ID NO:1 over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under CBFBBA01 polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO:1 to hybridize under conditions useable for amplification or for use as a probe or marker. The invention also provides polynucleotides which are complementary to such CBFBBA01 polynucleotides.

CBFBBA01 of the invention is structurally related to other proteins of the family, as shown by the results of sequencing the cDNA of Table 1 (SEQ ID NO:1) encoding human CBFBBA01. The cDNA sequence of SEQ ID NO:1 contains an open reading frame (nucleotide number 65 to 709) encoding a polypeptide of 215 amino acids of SEQ ID NO:2. The amino acid sequence of Table 2 (SEQ ID NO:2) has identity with mouse vesicle trafficking protein sec22b (J.C.Hay et al., Cell, 89:149-158,1997). The nucleotide sequence of Table 1 (SEQ ID NO:1) has about 54.7% identity (using FASTA) in 1462 nucleotide residues with mouse vesicle trafficking protein sec22b (J.C.Hay et al., Cell, 89:149-158,1997). Thus, CBFBBA01 polypeptides and polynucleotides of the present invention are expected to have, inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1^a

GGAGGAAGTGAGGACGGCGCCAAGGGCCTTCCGGGCCAGTGTGGATCCCTGTAGTTTGT
GAAGATGGTGTGTTGCTAACAATGATCGCCCGAGTGGCGGACGGGCTCCCGCTGGCCGCTC
GATGCAGGAGGACGAACAGTCTGGCCGGGACCTTCAACAGTATCAGAGTCAGGCTAAGCA
ACTCTTTTCGAAAGTTGAATGAACAGTCCCCCTACCAGATGTACCTTGAAGCAGGAGCCAT
GACTTTTCACTACATTATTGAGCAGGGGTGTGTTATTTGGTTTTATGTGAAGCTGCCTT
CCCTAAGAAGTTGGCTTTTGCCTACCTAGAAGATTTGCACTCAGAAATTTGATGAACAGCA
TGGAAAGAAGGTGCCCACTGTGTCCCGACCCTATTCCTTTATTGAATTTGATACTTTTCAT

```

TCAGAAAACCAAGAAGCTCTACATTGACAGTCGTGCTCGAAGAAATCTAGGCTCCATCAA
CACTGAATTGCAAGATGTGCAGAGGATCATGGTGGCCAATATTGAAGAAGTGTACCAACG
AGGAGAAGCACTCTCAGCATTGGATTCAAAGGCTAACAAATTTGTCCAGTCTGTCCAAGAA
ATACCGCCAGGATGCGAAGTACTTGAACATGCGTTCCACTTATGCCAAACTGCAGCAGT
AGCTGTATTTTTTCATCATGTTAATAGTGTATGTCCGATTCTGGTGGCTGTGAAATAATGA
ATACAGTCACTGGTAAGGGAGAACCTAGAACCCAGTAGGTGTATATTTTCAGGAAACTGA
GCTCACAGAGATGTGTATTAGAATCCAAGTGGAACTTCTGCCCTCTAAAGACCTTGCAAG
AAAAGAGATGCCCTGAAAATGAAAGGTTGCACCTCATTTAATGAAGCTTAACCCTATGTA
GAAAGTCTCTTTTCGGGGGCAGAGGCTTTCTCTGGGTGCCAAGCCATATATATTAGGGAAT
AGTAGATTGTTAATTTTCGTTTTTTCCCTCCCAGTGCATTTTAAAAACAGCACTGGCTGGG
GCATTCCTCATTCTCTGATGGAGCCATCAATGAGATTTAACTTAGTCAACCTGTGCCAGCA
ACATTCTGAAATTCCTTCAAAGAAGGCAGTCCCTTGGGAAGGTGTTTTTTTTTTTTTTTT
TTTTTTGACTCTAATCAACATTCTTTTTGTTGGTGACATTTGTGATTTTCAGTAATCTGA
GTTTTTGATGGCCTTTTAAACAAGACTCCAGTATGTGAAGGTTAATTGCTGTGCTCCACA
GATCTCGTCTATTGGCCCCCTGTAGAAAGTTAACCTTTGTTGTTTTCTTTTATAAATTTGC
TTATTGCACAATTGCTTTAGGGTAAGTGAATTATATTAAGATGCCTTGAAATTATAGCAC
TCCTTGATTAAGAAGCTAAAATGTTTCTCTCATTACTCCTTAAACAAAAGACAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAA
    
```

^a A nucleotide sequence of a human CBFBA01 (SEQ ID NO: 1).

Table 2^b

```

MVLITMIARVADGLPLAASMQEDEQSGRDLQYQSQAKQLFRKLNEQSPTRCTLEAGAMT
PHYIIEQGVCYLVLCFAAFPKLAFAYLEDLHSEFDEQHGKVPVSRPYSFIEFDTFIQ
KTKKLYIDSRARRNLGSINTELQDVQRIMVANIEEVLQRGEALSALDSKANNLSSLSKKY
RQDAKYLNMRSITYAKLAAVAVFFIMLIVYVRFWWL
    
```

5 ^b An amino acid sequence of a human CBFBA01 (SEQ ID NO: 2).

10 One polynucleotide of the present invention encoding CBFBA01 may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells of human cord blood using the expressed sequence tag (EST) analysis (Adams, M.D., *et al. Science* (1991) 252:1651-1656; Adams, M.D. *et al., Nature*, (1992) 355:632-634; Adams, M.D., *et al., Nature* (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding CBFBBBA01 polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in Table 1 (nucleotide number 65 to 709 of SEQ ID NO:1), or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

5 When the polynucleotides of the invention are used for the recombinant production of CBFBBBA01 polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker
10 sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gents *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and
15 sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding CBFBBBA01 variants comprise the amino acid sequence CBFBBBA01 polypeptide of Table 2 (SEQ ID NO:2) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

20 The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

25 Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1 or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding CBFBBBA01 polypeptide and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to the CBFBBBA01
30 gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding CBFBA01 polypeptide, including homologs and orthologs from species other than human, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO: 1 or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Thus in another aspect, CBFBA01 polynucleotides of the present invention further include a nucleotide sequence comprising a nucleotide sequence that hybridize under stringent condition to a nucleotide sequence having SEQ ID NO: 1 or a fragment thereof. Also included with CBFBA01 polypeptides are polypeptide comprising amino acid sequence encoded by nucleotide sequence obtained by the above hybridization condition. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY* (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and

Aspergillus cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL (supra)*.

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the CBFBBA01 polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If CBFBBA01 polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

CBFBBA01 polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

This invention also relates to the use of CBFBBA01 polynucleotides for use as diagnostic reagents. Detection of a mutated form of CBFBBA01 gene associated with a dysfunction will provide a

diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of CBFBBA01. Individuals carrying mutations in the CBFBBA01 gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled CBFBBA01 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers *et al.*, *Science* (1985) 230:1242. Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. See Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85: 4397-4401. In another embodiment, an array of oligonucleotides probes comprising CBFBBA01 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

The diagnostic assays offer a process for diagnosing or determining a susceptibility to cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis through detection of mutation in the CBFBBA01 gene by the methods described.

In addition, cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of CBFBBA01 polypeptide or CBFBBA01 mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as an CBFBBA01 polypeptide, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease, particularly cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, which comprises:

- 5 (a) a CFBBA01 polynucleotide, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof ;
 - (b) a nucleotide sequence complementary to that of (a);
 - (c) a CFBBA01 polypeptide, preferably the polypeptide of SEQ ID NO: 2, or a fragment thereof, or
 - (d) an antibody to a CFBBA01 polypeptide, preferably to the polypeptide of SEQ ID NO: 2.
- 10 It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome
15 identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for
20 example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

The differences in the cDNA or genomic sequence between affected and unaffected
25 individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

30 Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the CFBBA01 polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the CFBBA01 polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against CFBBA01 polypeptides may also be employed to treat cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, among others.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with CFBBA01 polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, among others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering CFBBA01 polypeptide via a vector directing expression of CFBBA01 polynucleotide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a CFBBA01 polypeptide wherein the composition comprises a CFBBA01 polypeptide or CFBBA01 gene. The vaccine formulation may further comprise a suitable carrier. Since CFBBA01 polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile

suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

The CBFBBA01 polypeptide of the present invention may be employed in a screening process for compounds which activate (agonists) or inhibit activation of (antagonists, or otherwise called inhibitors) the CBFBBA01 polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess identify agonist or antagonists from, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These agonists or antagonists may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide of the present invention; or may be structural or functional mimetics of the polypeptide of the present invention. See Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991). CBFBBA01 polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate CBFBBA01 polypeptide on the one hand and which can inhibit the function of CBFBBA01 polypeptide on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions as cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions as cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis.

In general, such screening procedures may involve using appropriate cells which express the CBFBBA01 polypeptide or respond to CBFBBA01 polypeptide of the present invention. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. Cells which express the CBFBBA01 polypeptide (or cell membrane containing the expressed polypeptide) or respond to CBFBBA01 polypeptide are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response. The ability of the cells which were contacted with the candidate compounds is compared with the same cells which were not contacted for CBFBBA01 activity.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the CBFBBA01 polypeptide is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled

competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the CBFBBBA01 polypeptide, using detection systems appropriate to the cells bearing the CBFBBBA01 polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a CBFBBBA01 polypeptide to form a mixture, measuring CBFBBBA01 activity in the mixture, and comparing the CBFBBBA01 activity of the mixture to a standard.

The CBFBBBA01 cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of CBFBBBA01 mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of CBFBBBA01 protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of CBFBBBA01 (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The CBFBBBA01 protein may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the CBFBBBA01 is labeled with a radioactive isotope (eg 125I), chemically modified (eg biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. In addition to being used for purification and cloning of the receptor, these binding assays can be used to identify agonists and antagonists of CBFBBBA01 which compete with the binding of CBFBBBA01 to its receptors, if any. Standard methods for conducting screening assays are well understood in the art.

Examples of potential CBFBBBA01 polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the CBFBBBA01 polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for CBFBBBA01 polypeptides; or compounds which decrease or enhance the production of CBFBBBA01 polypeptides, which comprises:

- (a) a CFBFBA01 polypeptide, preferably that of SEQ ID NO:2;
- (b) a recombinant cell expressing a CFBFBA01 polypeptide, preferably that of SEQ ID NO:2;
- (c) a cell membrane expressing a CFBFBA01 polypeptide; preferably that of SEQ ID NO: 2; or
- (d) antibody to a CFBFBA01 polypeptide, preferably that of SEQ ID NO: 2.

5 It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Prophylactic and Therapeutic Methods

10 This invention provides methods of treating abnormal conditions such as, cancer, autoimmune disease, diabetes mellitus, and multiple sclerosis, related to both an excess of and insufficient amounts of CFBFBA01 polypeptide activity.

If the activity of CFBFBA01 polypeptide is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as hereinabove described along with a pharmaceutically acceptable carrier in an amount effective to inhibit the function
15 of the CFBFBA01 polypeptide, such as, for example, by blocking the binding of ligands, substrates, receptors, enzymes, etc., or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of CFBFBA01 polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc. in competition with endogenous CFBFBA01 polypeptide may be administered. Typical embodiments of such competitors comprise fragments of the CFBFBA01
20 polypeptide.

In still another approach, expression of the gene encoding endogenous CFBFBA01 polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. See, for example, O'Connor, *J Neurochem* (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors
25 of Gene Expression, CRC Press, Boca Raton, FL (1988). Alternatively, oligonucleotides which form triple helices with the gene can be supplied. See, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360. These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

For treating abnormal conditions related to an under-expression of CFBFBA01 and its activity,
30 several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates CFBFBA01 polypeptide, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of CFBFBA01 by the relevant cells in the subject. For example, a polynucleotide of the invention may

be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer
5 cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in *Human Molecular Genetics*, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of CBFBBA01 polypeptides in combination with a suitable pharmaceutical carrier.

10

Formulation and Administration

Peptides, such as the soluble form of CBFBBA01 polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a
15 pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

20

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

25

Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and
transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

30

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending
practitioner. Suitable dosages, however, are in the range of 0.1-100 $\mu\text{g}/\text{kg}$ of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these

dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

5 Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

10 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

SEQUENCE LISTING

- 5 (1) GENERAL INFORMATION
- (i) APPLICANT: WU, JI-SHENG
WANG, YA-XIN
- 10 (ii) TITLE OF THE INVENTION: THE HUMAN VESICLE TRAFFICKING
PROTEIN SEC22b GENE OF CBFBA01
- (iii) NUMBER OF SEQUENCES: 2
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: RATNER & PRESTIA
(B) STREET: P.O. BOX 980
(C) CITY: VALLEY FORGE
(D) STATE: PA
20 (E) COUNTRY: USA
(F) ZIP: 19482
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Diskette
25 (B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
30 (A) APPLICATION NUMBER: TO BE ASSIGNED
(B) FILING DATE:
(C) CLASSIFICATION: UNKNOWN
- (vii) PRIOR APPLICATION DATA:
35 (A) APPLICATION NUMBER:
(B) FILING DATE:
- 40 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: PRESTIA, PAUL F
(B) REGISTRATION NUMBER: 23,031
(C) REFERENCE/DOCKET NUMBER: GP-70313

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 610-407-0700
- (B) TELEFAX: 610-407-0701
- (C) TELEX: 846169

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1462 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

GGAGGAAGTG AGGACGGCGC CAAGGGCCTT CCGGGCCAGT GTTGGATCCC TGTAGTTTGT      60
20 GAAGATGGTG TTGCTAACAA TGATCGCCCG AGTGGCGGAC GGGCTCCCGC TGGCCGCCTC      120
GATGCAGGAG GACGAACAGT CTGGCCGGGA CCTTCAACAG TATCAGAGTC AGGCTAAGCA      180
ACTCTTTCGA AAGTTGAATG AACAGTCCCC TACCAGATGT ACCTTGAAG CAGGAGCCAT      240
GACTTTTCAC TACATTATTG AGCAGGGGGT GTGTTATTTG GTTTTATGTG AAGCTGCCTT      300
CCCTAAGAAG TTGGCTTTTG CCTACCTAGA AGATTTGCAC TCAGAATTTG ATGAACAGCA      360
25 TGGAAAGAAG GTGCCCACTG TGTCCCGACC CTATTCCTTT ATTGAATTTG ATACTTTTCAT      420
TCAGAAAACC AAGAAGCTCT ACATTGACAG TCGTGCTCGA AGAAATCTAG GCTCCATCAA      480
CACTGAATTG CAAGATGTGC AGAGGATCAT GGTGGCCAAT ATTGAAGAAG TGTTACAACG      540
AGGAGAAGCA CTCTCAGCAT TGGATTCAA GCGTAACAAT TTGTCCAGTC TGTCCAAGAA      600
ATACCGCCAG GATGCGAAGT ACTTGAACAT GCGTTCCACT TATGCCAAAC TTGCAGCAGT      660
30 AGCTGTATTT TTCATCATGT TAATAGTGTA TGTCCGATTC TGGTGGCTGT GAAATAATGA      720
ATACAGTCAC TGGTAAGGGA GAACCTAGAA CCCAGTAGGT GTATATTTTC AGGAAACTGA      780
GCTCACAGAG ATGTGTATTA GAATCCAAGT GGAACCTCTG CCCTCTAAAG ACCTTGCAAG      840
AAAAGAGATG CCCTGAAAAT GAAAGGTTGC ACCTCATTTA ATGAAGCTTA ACCCTATGTA      900
GAAAGTCTCT TTCGGGGGCA GAGGCTTTCT CTGGGTGCCA AGCCATATAT ATTAGGGAAT      960
35 AGTAGATTGT TAATTTGTTT TTTTCCCTCC CAGTGCATTT TAAAAACAGC ACTGGCTGGG      1020
GCATTCTCAT TCTCTGATGG AGCCATCAAT GAGATTTAAC TTAGTCAACC TGTGCCAGCA      1080
ACATTCTGAA ATTCCTTCAA AGAAGGCAGT CCTTTGGGAA GGTGTTTTTT TTTTTTTTTT      1140
TTTTTTGACT CTAATCAACA TTCCTTTTGT TGGTGACATT TGTGATTTTC AGTAATCTGA      1200
GTTTTTGATG GCCTTTTAAA CAAGACTCCA GTATGTGAAG GTTAATTGCT GTGCTCCACA      1260
40 GATCTCGTCT ATTGGCCCCT GTAGAAAGTT AACCTTTGTT GTTTTCCCTT TATAATTTGC      1320
TTATTGCACA ATTGCTTTAG GGTAAGTGAA TTATATTAAG ATGCCTTGAA ATTATAGCAC      1380
TCCTTGATTA AGAAGCTAAA ATGTTTCTCT CATTTACTCC TTAAACAAAA GACAAAAAAA      1440
AAAAAAAAAA AAAAAAAAAA AA
                                                                                                                                 1462
    
```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 215 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Val Leu Leu Thr Met Ile Ala Arg Val Ala Asp Gly Leu Pro Leu
 1           5           10           15
15 Ala Ala Ser Met Gln Glu Asp Glu Gln Ser Gly Arg Asp Leu Gln Gln
    20           25           30
Tyr Gln Ser Gln Ala Lys Gln Leu Phe Arg Lys Leu Asn Glu Gln Ser
    35           40           45
Pro Thr Arg Cys Thr Leu Glu Ala Gly Ala Met Thr Phe His Tyr Ile
20    50           55           60
Ile Glu Gln Gly Val Cys Tyr Leu Val Leu Cys Glu Ala Ala Phe Pro
65           70           75           80
Lys Lys Leu Ala Phe Ala Tyr Leu Glu Asp Leu His Ser Glu Phe Asp
    85           90           95
25 Glu Gln His Gly Lys Lys Val Pro Thr Val Ser Arg Pro Tyr Ser Phe
    100          105          110
Ile Glu Phe Asp Thr Phe Ile Gln Lys Thr Lys Lys Leu Tyr Ile Asp
    115          120          125
Ser Arg Ala Arg Arg Asn Leu Gly Ser Ile Asn Thr Glu Leu Gln Asp
30    130          135          140
Val Gln Arg Ile Met Val Ala Asn Ile Glu Glu Val Leu Gln Arg Gly
145          150          155          160
Glu Ala Leu Ser Ala Leu Asp Ser Lys Ala Asn Asn Leu Ser Ser Leu
    165          170          175
35 Ser Lys Lys Tyr Arg Gln Asp Ala Lys Tyr Leu Asn Met Arg Ser Thr
    180          185          190
Tyr Ala Lys Leu Ala Ala Val Ala Val Phe Phe Ile Met Leu Ile Val
    195          200          205
Tyr Val Arg Phe Trp Trp Leu
40    210          215
    
```


What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80%
5 identity over its entire length to a nucleotide sequence encoding the CBFBBAA01 polypeptide of SEQ ID NO:2; or a nucleotide sequence complementary to said isolated polynucleotide.
2. The polynucleotide of claim 1 wherein said polynucleotide comprises the
nucleotide sequence contained in SEQ ID NO:1 encoding the CBFBBAA01 polypeptide of SEQ ID
10 NO2.
3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide
sequence that is at least 80% identical to that of SEQ ID NO: 1 over its entire length.
- 15 4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: 1.
5. The polynucleotide of claim 1 which is DNA or RNA.
6. A DNA or RNA molecule comprising an expression system, wherein said
20 expression system is capable of producing a CBFBBAA01 polypeptide comprising an amino acid
sequence of SEQ ID NO:2 when said expression system is present in a compatible host cell.
7. A host cell comprising the expression system of claim 6.
- 25 8. A process for producing a CBFBBAA01 polypeptide comprising culturing a host of
claim 7 under conditions sufficient for the production of said polypeptide and recovering the
polypeptide from the culture.
9. A process for producing a cell which produces a CBFBBAA01 polypeptide thereof
30 comprising transforming or transfecting a host cell with the expression system of claim 6 such that
the host cell, under appropriate culture conditions, produces a CBFBBAA01 polypeptide.
10. A CBFBBAA01 polypeptide comprising an amino acid sequence of SEQ ID NO:2.

11. An antibody immunospecific for the CFBBA01 polypeptide of claim 10.
12. A method for the treatment of a subject in need of enhanced activity or expression
5 of CFBBA01 polypeptide of claim 10 comprising:
(a) administering to the subject a therapeutically effective amount of an agonist to said polypeptide; and/or
(b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence
10 that has at least 80% identity to a nucleotide sequence encoding the CFBBA01 polypeptide of SEQ ID NO:2 over its entire length; or a nucleotide sequence complementary to said nucleotide sequence in a form so as to effect production of said polypeptide activity *in vivo*.
13. A method for the treatment of a subject having need to inhibit activity or
expression of CFBBA01 polypeptide of claim 10 comprising:
15 (a) administering to the subject a therapeutically effective amount of an antagonist to said polypeptide; and/or
(b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said polypeptide; and/or
(c) administering to the subject a therapeutically effective amount of a polypeptide
20 that competes with said polypeptide for its ligand, substrate, or receptor.
14. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of CFBBA01 polypeptide of claim 10 in a subject comprising:
(a) determining the presence or absence of a mutation in the nucleotide sequence
25 encoding said CFBBA01 polypeptide in the genome of said subject; and/or
(b) analyzing for the presence or amount of the CFBBA01 polypeptide expression in a sample derived from said subject.
15. A method for identifying compounds which inhibit (antagonize) or agonize the
30 CFBBA01 polypeptide of claim 10 which comprises:
(a) contacting a candidate compound with cells which express the CFBBA01 polypeptide (or cell membrane expressing CFBBA01 polypeptide) or respond to CFBBA01 polypeptide; and


(b) observing the binding, or stimulation or inhibition of a functional response; or comparing the ability of the cells (or cell membrane) which were contacted with the candidate compounds with the same cells which were not contacted for CBFBBBA01 polypeptide activity.

- 5 16. An agonist identified by the method of claim 15.
17. An antagonist identified by the method of claim 15.
18. A recombinant host cell produced by a method of Claim 9 or a membrane thereof
- 10 expressing a CBFBBBA01 polypeptide.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN 97/00115

<p>A. CLASSIFICATION OF SUBJECT MATTER</p> <p>Int.Cl⁶ C12N15/12, C12N15/64, C12P21/08</p> <p>According to International Patent Classification (IPC) or to both national classification and IPC</p>														
<p>B. FIELDS SEARCHED</p> <p>Minimum documentation searched (classification system followed by classification symbols)</p> <p>Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched</p> <p>Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)</p> <p>Genebank , Dgene , Caplus , Biosis , Biotechds</p>														
<p>C. DOCUMENTS CONSIDERED TO BE RELEVANT</p> <table border="1"> <thead> <tr> <th>Category *</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>Y</td> <td>Annual Meeting of the 6th International congress on Cell Biology and the 36th. American Society for Cell Biology, San Francisco, California, USA December 7—11, 1996, Molecular Biology of the Cell 7 (SUPPL.) 1996, 74A Hay J C; et al "Mammalian vesicle trafficking proteins of the endoplasmic reticulum and Golgi" See the abstract.</td> <td>1—18</td> </tr> <tr> <td>Y</td> <td>Journal of Biological Chemistry 271(10). 1996, 56771—5679 Hay J C et al "Mammalian vesicle trafficking proteins of the endoplasmic reticulum and Golgi apparatus". See the abstract.</td> <td>1—18</td> </tr> <tr> <td>Y</td> <td>Cell 89(1) 1997, 149—158 Hay J C et al, "Protein interactions regulating vesicle transport between the endoplasmic reticulum and Golgi apparatus in mammalian cells" See the abstract.</td> <td>1—18</td> </tr> </tbody> </table>			Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	Y	Annual Meeting of the 6th International congress on Cell Biology and the 36th. American Society for Cell Biology, San Francisco, California, USA December 7—11, 1996, Molecular Biology of the Cell 7 (SUPPL.) 1996, 74A Hay J C; et al "Mammalian vesicle trafficking proteins of the endoplasmic reticulum and Golgi" See the abstract.	1—18	Y	Journal of Biological Chemistry 271(10). 1996, 56771—5679 Hay J C et al "Mammalian vesicle trafficking proteins of the endoplasmic reticulum and Golgi apparatus". See the abstract.	1—18	Y	Cell 89(1) 1997, 149—158 Hay J C et al, "Protein interactions regulating vesicle transport between the endoplasmic reticulum and Golgi apparatus in mammalian cells" See the abstract.	1—18
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
Y	Annual Meeting of the 6th International congress on Cell Biology and the 36th. American Society for Cell Biology, San Francisco, California, USA December 7—11, 1996, Molecular Biology of the Cell 7 (SUPPL.) 1996, 74A Hay J C; et al "Mammalian vesicle trafficking proteins of the endoplasmic reticulum and Golgi" See the abstract.	1—18												
Y	Journal of Biological Chemistry 271(10). 1996, 56771—5679 Hay J C et al "Mammalian vesicle trafficking proteins of the endoplasmic reticulum and Golgi apparatus". See the abstract.	1—18												
Y	Cell 89(1) 1997, 149—158 Hay J C et al, "Protein interactions regulating vesicle transport between the endoplasmic reticulum and Golgi apparatus in mammalian cells" See the abstract.	1—18												
<p><input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.</p>														
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claims (s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>														
<p>Date of the actual completion of the international search</p> <p>17. Mar. 1998(17. 03. 98)</p>		<p>Date of mailing of the international search report</p> <p>02 APR 1998 (02. 04. 98)</p>												
<p>Name and mailing address of the ISA/ Chinese Patent Office, 6 Xitucheng Rd. Jimen Bridge, Haidian District, 100088 Beijing, China Facsimile No. (86—10)62019451</p>		<p>Authorized officer</p> <p>Telephone No. 62093733</p> 												

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN 97/00115

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>Annual Meeting of the 6th International congress on cell Biology and the 36th American Society for cell Biology, San Francisco, California, USA, December 7—11, 1996, Molecular Biology of the cell 7 (SUPPL), 1996. 82A</p> <p>Yoon Y; McNiven M. A. "Identification of a novel dynamin-like protein expressed in rat tissues".</p> <p>See the abstract.</p>	1—18