

(19)



Europäisches
Patentamt
European
Patent Office
Office européen
des brevets



(11)

EP 2 862 871 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
22.04.2015 Bulletin 2015/17

(51) Int Cl.:

C07K 7/06 (2006.01)

A61K 49/14 (2006.01)

A61K 38/08 (2006.01)

G01N 33/574 (2006.01)

G01N 33/68 (2006.01)

C07K 16/18 (2006.01)

(21) Application number: 14190585.1

(22) Date of filing: 14.04.2009

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK TR

- Weissleder, Ralph
Peabody, MA 01960 (US)
- Bardeesy, Nabeel
Framingham, MA 01701 (US)

(30) Priority: 14.04.2008 US 44818

(74) Representative: Donald, Jenny Susan
Forresters
Skygarden
Erika-Mann-Strasse 11
80636 München (DE)

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
09733422.1 / 2 265 630

(71) Applicant: The General Hospital Corporation
Boston, MA 02114 (US)

Remarks:

- This application was filed on 28-10-2014 as a
divisional application to the application mentioned
under INID code 62.
- Claims filed after the date of filing of the application
(Rule 68(4) EPC).

(72) Inventors:
• Kelly, Kimberley
Crozet, VA 22932 (US)

(54) Plectin-1 targeted agents for detection and treatment of pancreatic ductal adenocarcinoma

(57) Described herein are compositions and methods for cancer cell biomarkers, such as pancreatic ductal adenocarcinoma (PDAC) cell biomarkers, and binding molecules for diagnosis and treatment of cancer, e.g., PDAC. Methods of identifying "accessible" proteomes

are disclosed for identifying cancer biomarkers, such as plectin-1, a PDAC biomarker. Additionally, imaging compositions are provided comprising magnetofluorescent nanoparticles conjugated to peptide ligands for identifying PDACs.

EP 2 862 871 A1

Description**CLAIM OF PRIORITY**

5 [0001] This application claims the benefit of U.S. Provisional Patent Application Serial No. 61/044,818, filed on April 14, 2008, the entire contents of which are hereby incorporated by reference.

GOVERNMENT SUPPORT

10 [0002] This invention was made with government support from the United States National Institutes of Health (NIH) under grant numbers P50-CA86355, PO1-CA117969-01, K01 CA104647-03, EB004626, ROI-HL078641, ROI-HL36436, HL080731, and P01-AI 054904. The United States Government may have certain rights in the invention.

FIELD OF THE INVENTION

15 [0003] The present invention relates to compositions and methods for providing cancer cell biomarkers, such as pancreatic ductal adenocarcinoma (PDAC) cell biomarkers, and binding molecules for diagnosis and treatment. Specifically, methods of identifying "accessible" proteomes are disclosed for providing biomarkers, such as plectin-1, for identifying PDACs. Additionally, imaging compositions are provided comprising magnetofluorescent nanoparticles conjugated to peptide ligands for identifying PDACs. Finally, methods of treating PDAC are also discussed.

BACKGROUND OF THE INVENTION

20 [0004] Pancreatic ductal adenocarcinoma (PDAC) is the 4th leading cause of cancer death in the United States showing a rapid clinical course leading to death. Once diagnosed, PDAC has a median survival of 6 months and a 5-year survival rate of only 3 percent (Li et al., Lancet 363:1049-1057 (2004)).

25 [0005] As chemotherapy and radiotherapy have only modest benefits, and surgery is only possible in 20% of patients, early detection that allows surgical resection offers the best hope for longer survival (Yeo et al., Ann Surg 222:580-588 (1995); discussion 588-592). Indeed, the detection of PDAC or high-grade precursors in high-risk patient groups (e.g., 30 hereditary cancer syndromes, chronic pancreatitis, and new-onset diabetes) represents a critical unmet need in the cancer diagnostic portfolio (Brentnall et al., Ann. Intern. Med. 131:247-255 (1999); Canto et al., Clin. Gastroenterol. Hepatol. 2:606-621 (2004)).

35 [0006] Currently, serum CA-19-9 is the clinically used biomarker; however, it lacks the sensitivity needed to detect early-stage PDAC (Goggins, J. Clin. Oncol. 23:4524-4531 (2005)). In addition, cross-sectional abdominal imaging has proven to be unreliable to detect early-stage PDAC in high-risk patients (Pelaez-Luna et al., Am J Gastroenterol 102:2157-2163 (2007)).

40 [0007] Thus a high priority in this field of medicine is the identification of biomarkers for the development of binding ligands as diagnostics, such as imaging probes for detecting pre-neoplastic/early invasive lesions and for use in treatments.

SUMMARY OF THE INVENTION

45 [0008] The present invention relates, at least in part, to compositions and methods for providing cancer cell biomarkers, such as pancreatic ductal adenocarcinoma (PDAC) cell biomarkers, and binding molecules for diagnosis and treatment. Specifically, methods of identifying "accessible" proteomes are disclosed for providing biomarkers, such as plectin-1, for identifying PDACs. Additionally, imaging compositions are provided comprising magnetofluorescent nanoparticles conjugated to peptide ligands for identifying PDACs.

50 [0009] Described herein are biomarkers for identifying cancer cells, as opposed to noncancer cells, for use in diagnostics and treatments, such as for providing binding partners (i.e., ligands) such as peptides, small molecules, peptide mimetics, nonpeptide mimetics, antibodies and the like.

55 [0010] In one aspect, the invention provides biomarkers of cancer cells, wherein the biomarker comprises a plectin-1 fragment and is located on the external membrane of the cancer cell. In some embodiments, the biomarker fragment includes any one of the sequences set forth in SEQ ID NOs: 9-23. Further, the present inventions are not limited by the type of cancer cell. Indeed a variety of cancer cell types are contemplated, including but not limited to a gastrointestinal cancer cell, a hepatobiliary cancer cell, a gall bladder cancer cell, a pancreatic cancer cell, a lung cancer cell, a mesothelioma cancer cell, a bladder cancer cell, a prostate cancer cell, a breast cancer cell, a head cancer cell, a neck cancer cell, a thyroid cancer cell, a uterine cancer cell, a cervix cancer cell, a uterine-cervix cancer cell, a blood cancer cell, a white blood cancer cell, a bone marrow cancer cell, a pleura cancer cell, a pleural fluid cancer cell, a prostate

cancer cell, and the like. In some embodiments, the cancer cell is selected from the group consisting of a pancreatic cancer cell and a pancreatic ductal adenocarcinoma cell (PDAC). In some embodiments, the biomarker comprises SEQ ID NO:24 or an isoform thereof. In some embodiments, the biomarker comprises a peptide or fragment thereof encoded by human mRNA sequences, such as those described in the GenBank Database at Acc. Nos. NM_000445.2; NM_201378.1; NM_201379.1; NM_201380.2; M_201381.1; NM_201382.1; NM_201383.1; and/or NM_201384.1. In some embodiments, the biomarker binds to a phage-displayed peptide.

[0011] In an additional aspect, the present invention features plectin-1 ligands including a first portion including a plectin-1 binding moiety, coupled to a second portion that includes a detectable moiety or a therapeutic agent. In some embodiments, the plectin-1 binding moiety is an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, or 4 - 8, or a peptidomimetic thereof. In some embodiments, the plectin-1 binding moiety is an anti-plectin-1 antibody or antigen-binding fragment thereof, a small molecule, or an aptamer. In some embodiments, the plectin-1 binding moiety is coupled to a nanoparticle, a microparticle, or a solid phase reagent. The detectable moiety can be selected from the group consisting of a radioactive isotope, a magnetic compound, an x-ray absorber, a chemical compound, a biological tag, and a fluorescent molecule. The therapeutic agent can be, e.g., a cytotoxic moiety or an immunomodulatory moiety (e.g., a compound that enhances the immune response to the tumor, e.g., an inflammatory cytokine such as interleukin-1 (1L-1), and tumour necrosis factor-alpha (TNF- α)).

[0012] In some embodiments, there is a linker between the first portion and the second portion, e.g., a flexible amino acid sequence, e.g., a photolinker.

[0013] In some embodiments, the second portion includes a physiologically inert nanoparticle, e.g., in addition to or as the detectable moiety or therapeutic agent.

[0014] The peptide ligand of claim 8, wherein the nanoparticle is magnetic, fluorescent, or radioactive. Exemplary nanoparticles include crosslinked iron oxide nanoparticles (CLIOs), superparamagnetic iron oxide nanoparticles (SP-TONs), and cross-linked superparamagnetic iron oxide nanoparticles.

[0015] In some embodiments, the second portion comprises a fluorochrome, e.g., with an excitation maxima range of 500 nm - 1000 nm. In some embodiments, the fluorochrome is a near infrared fluorochrome, e.g., with an excitation maxima range of 650-680 nm, e.g., a cyanine derivative such as cyanine 5.5.

[0016] In some, the second portion comprises a crosslinked iron oxide nanoparticle conjugated to a NIRF, e.g., cyanine 5.5 (CLIO-Cy5.5).

[0017] In some embodiments, the invention features peptide ligands including or consisting essentially of SEQ ID NO:1 coupled to a nanoparticle, optionally with a linker, e.g., a fluorescently labeled linker, and a nanoparticle, e.g., a magnetofluorescent nanoparticle. In some embodiments, one or both of the linker and the particle is fluorescent. In some embodiments, the magnetofluorescent nanoparticle comprises a near infrared (NIR) fluorochrome (NIRF).

[0018] In another aspect, plectin-1 binding moiety is the invention provides ligands for a biomarker, wherein the ligand is selected from the group consisting of a peptide ligand, a mimetic, a small molecule, and an antibody. In some embodiments, the ligand is a peptide. Thus, in some embodiments, the invention provides a peptide ligand including any one of amino acid SEQ ID NOs:1-8, for binding to a pancreatic ductal adenocarcinoma cell. In some embodiments, the peptide ligand binds to a biomarker for a pancreatic ductal adenocarcinoma cell molecule, such as plectin-1. Thus, in some embodiments, the invention provides a peptide ligand including amino acid SEQ ID NO:1, for binding to a pancreatic ductal adenocarcinoma cell. In some embodiments, the ligand binds to a receptor on a pancreatic cancer cell. In some embodiments, the ligand binds to a biomarker of a pancreatic cancer cell. In some embodiments, the biomarker is expressed differently in a cancer cell than in a noncancer cell. In some embodiments, the ligand has a different binding pattern to receptors of a cancer cell compared to a noncancer cell. In some embodiments, the ligand binds to biomarkers in a different location in a cancer cell than in a noncancer cell. In some embodiments, the ligand binds to a greater number of receptors of a cancer cell than of a noncancer cell. In some embodiments, the ligand binds to a greater number of cancer cells than to noncancer cells. In some embodiments, the ligand identifies a biomarker of a cancer cell. In some embodiments, the ligand binds to a biomarker of a cancer cell. In some embodiments, the biomarker is plectin-1. In some embodiments, the ligand derives from a random phage-displayed peptide library. In some embodiments, the peptide ligand derives from a random phage-displayed peptide library. In some embodiments, the ligand derives from a phage-displayed peptide. In some embodiments, the ligand is synthetic.

[0019] In another aspect, the invention provides bacteriophage displaying a peptide ligand bound to a pancreatic ductal adenocarcinoma cell. In some embodiments, the bacteriophage includes a fluorescent molecule, e.g., a fluorescein isothiocyanate.

[0020] In yet another aspect, the invention features isolated bacteriophage displaying a peptide ligand eluted from a pancreatic ductal adenocarcinoma cell, e.g., an isolated bacteriophage displaying a peptide ligand comprising SEQ ID NO:1.

[0021] In a further aspect, the invention provides diagnostic peptides, e.g., a peptide including SEQ ID NO: 1. In some embodiments, the diagnostic peptides are coupled to a label, e.g., a label selected from the group consisting of a radioactive isotope, a chemical compound, a biological tag, and a fluorescent molecule. In some embodiments, the label

is selected from the group consisting of I¹²⁵, biotin, histidine tag, a fluorochrome derived from a fluorochrome-hydro-succinimide ester, and a fluorescein isothiocyanate. In some embodiments, the diagnostic peptide further comprises a linker. In some embodiments, the linker is a photolinker. In some embodiments, the photolinker is sulfosuccinimidyl 2-[7-amino-4-methylcoumarin-3-acetamido]ethyl-1,3 dithiopropionate. In some embodiments, the linker is labeled, e.g., is fluorescently labeled. In some embodiments, the fluorescently labeled linker is a GGSK(Fluorescein isothiocyanate (FITC))C linker.

[0022] In some embodiments, the peptide is conjugated to a physiologically inert nanoparticle. In some embodiments, the nanoparticle is selected from the group consisting of a crosslinked iron oxide nanoparticle (CLIO), a superparamagnetic iron oxide nanoparticle (SPIONs), a cross-linked superparamagnetic iron oxide nanoparticle, et cetera. In some embodiments, the nanoparticle further comprises a fluorophore, e.g., a near infrared (NIR) fluorochrome, e.g., with an excitation/emission maxima at 785 nm/810 nm or an excitation/emission maxima at 675 nm/694 nm. In some embodiments, the near infrared (NIR) fluorochrome is cyanine 5.5 or a derivative thereof.

[0023] In some embodiments, the diagnostic peptide is linked to a magnetofluorescent nanoparticle, e.g., a crosslinked iron oxide nanoparticle conjugated to cyanine 5.5 (CLIO-Cy5.5).

[0024] In another aspect, the invention provides diagnostic compositions including peptides including SEQ ID NO: 1, and optionally one or both of a fluorescently labeled linker and a magnetofluorescent nanoparticle. In some embodiments, the diagnostic compositions include a peptide ligand including SEQ ID NO:1 coupled to a fluorescently labeled linker molecule conjugated to a crosslinked iron oxide nanoparticle, wherein the nanoparticle is conjugated to a near infrared (NIR) fluorochrome (NIRF).

[0025] In a further aspect, the invention provides diagnostic compositions including a peptide, e.g., SEQ ID NO:2 (i.e., KTLPTPGGSK), e.g., a FITC-labeled peptide including SEQ ID NO:2 (i.e., KTLPTPGGSK(Fluorescein isothiocyanate (FITC))C), conjugated to a crosslinked iron oxide nanoparticle, wherein the nanoparticle is further conjugated to a fluorochrome, e.g., a NIRF, e.g., a cyanine derivative such as cyanine 5.5 (CLIO-Cy5.5).

[0026] Also provided herein is the use of the plectin-1 binding compounds described herein for the diagnosis or treatment of pancreatic ductal adenocarcinoma, and in the manufacture of a medicament for the diagnosis or treatment of pancreatic ductal adenocarcinoma.

[0027] In another aspect, the invention provides methods for detecting a cancer cell in a subject. The methods include providing cells or tissue from the subject (e.g., from a biopsy, or from plasma or blood, i.e., circulating tumor cells from the subject); and detecting the presence or subcellular localization of plectin-1 protein in the cells or tissue, wherein the presence of cell membrane expression of plectin-1 indicates the presence of a cancer cell. In some embodiments, the absence of plectin-1, or the presence of only cytoplasmic and/or nuclear expression of plectin-1, indicates that there are no cancer cells in the cells or tissue.

[0028] In some embodiments, detecting subcellular localization of plectin-1 expression in the sample includes contacting the sample with an agent that binds to plectin-1 protein, optionally comprising a detectable moiety; and detecting subcellular localization of the agent; wherein the subcellular localization of the agent indicates the subcellular localization of plectin-1 expression. In some embodiments, the agent that binds to plectin-1 protein is a peptide ligand as described herein, or an antibody or antigen-binding portion thereof that is specific for plectin-1. In some embodiments, subcellular localization is detected using laser scanning microscopy, immunohistochemistry, fluorescent microscopy, and/or radiography. Other methods, including Raman spectroscopy, optical coherence tomography (OCT), detection of radiation (e.g., x-ray) scattering or absorption, ultrasound, and isotope detection, can also be used.

[0029] In yet another aspect, the invention provides methods for detecting pancreatic ductal adenocarcinoma (PDAC) or precursor pancreatic intraepithelial neoplasia (PanINs) in a subject *in vivo*. The methods include identifying a subject who is at risk for, or suspected of having, PDAC; administering to the subject a diagnostic composition as described herein; and detecting the presence of the peptide ligand in the pancreas of the subject using an *in vivo* imaging device. The presence of the peptide ligand in the pancreas indicates that the subject has PDAC. In some embodiments, the peptide ligand is administered via intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, oral, and/or gastric routes. In some embodiments, the *in vivo* imaging device is selected from the group consisting of magnetic resonance imagers (MRI), intravital laser scanning microscopes, endoscopes, and radiographic imagers.

[0030] Also provided herein are methods for treating pancreatic ductal adenocarcinoma (PDAC) in a subject. The methods include identifying a subject who is at risk for, or suspected of having, PDAC; administering to the subject a diagnostic composition comprising a diagnostic composition as described herein; detecting localization of the peptide ligand in the pancreas of the subject using an *in vivo* imaging device, wherein the localization of the peptide ligand indicates the localization of PDAC cells; and surgically removing the PDAC cells.

[0031] In a further aspect, the invention provides methods for treating pancreatic ductal adenocarcinoma (PDAC) in a subject. The methods include identifying a subject who is at risk for, or suspected of having, PDAC; and administering to the subject a therapeutically effective amount of a therapeutic composition as described herein, e.g., comprising a plectin-1 binding moiety coupled to a therapeutic agent. In some embodiments, the composition is administered by intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, oral, and/or gastric routes.

[0032] In some embodiments, the therapeutic agent is a cytotoxic or cytostatic moiety selected from the group consisting of a therapeutic drug, a radioisotope, molecules of plant, fungal, or bacterial origin, and biological proteins. In some embodiments, the therapeutic agent is a phototoxic or immunomodulatory compound.

5 [0033] Also provided herein are peptides consisting essentially of an amino acid sequence selected from the group consisting of any of SEQ ID NOs: 9-23.

[0034] In yet another aspect, the invention provides methods for identifying a pancreatic cancer cell peptide ligand. The methods include a) providing i) a phage-displayed random combinatorial peptide library, wherein the peptide ranges from 7 - 12 mer, ii) a receptor for the peptide, wherein the receptor derives from a pancreatic cancer cell's "accessible" proteome, and iii) a phage-displayed peptide-receptor binding assay; b) performing a phage-displayed peptide-receptor binding assay, and c) identifying a pancreatic cancer cell biomarker. In some embodiments, the cancer cell is a pancreatic ductal adenocarcinoma cell. In some embodiments, the phage-displayed random combinatorial peptide library comprises amino acid SEQ ID NO: 1. In some embodiments, the phage further comprises a fluorochrome-label. In some embodiments, the label is selected from the group consisting of a fluorochrome-hydrosuccinimide ester, cyanine 5.5, and fluorescein isothiocyanate. In some embodiments, the pancreatic cancer cell's "accessible" proteome includes components selected from the group consisting of proteins, subcellular fractions, cell lysate, and whole cells. In some embodiments, the receptor is plectin-1. In some embodiments, the peptide-receptor binding assay is selected from the group consisting of Enzyme-Linked ImmunoSorbent Assay (ELISA), a High Performance Liquid Chromatography (HPLC) assay, a competitive binding assay, an immunofluorescence assay, a radioactive assay, an intact cell binding assay, an SDS/PAGE gel assay, a fluorescent Microscopy assay and flow Cytometry assay. In some embodiments, the identifying step includes but is not limited to detecting differential gene expression, protein processing, carbohydrate processing, trafficking, intracellular location, cell surface expression, binding pattern, and binding amount.

10 [0035] In an additional aspect, the invention provides methods for identifying a peptide ligand for a pancreatic cancer cell. The methods include a) providing i) a pancreatic cancer cell, wherein the cancer cell expresses a receptor, ii) a non-cancer cell, iii) a phage-displayed peptide ligand, and, and iv) a peptide ligand - receptor binding assay; b) adding the peptide ligand to a cancer cell and to a non-cancer cell; and c) performing a peptide binding assay to distinguish the cancer cell from the non-cancer cell. In some embodiments, the method further comprises eluting the peptide ligand from the receptor. In some embodiments, the method further comprises sequencing the phage displayed peptide. In some embodiments, the receptor is an immobilized receptor-binding partner. In some embodiments, the immobilized receptor is immobilized on a support material selected from the group consisting of a biopsy specimen, a bead, a membrane, a gel, a membrane, and a plastic.

15 [0036] In another aspect, the invention provides methods for diagnosing a cancer in a subject. The methods include providing a sample, e.g., a biopsy sample, from a patient, and a diagnostic composition including a peptide ligand, wherein the peptide ligand includes SEQ ID NO: 1, optionally conjugated to a detectable moiety, adding the peptide to the sample, and detecting the diagnostic composition, e.g., by detecting the detectable moiety in the sample. In some embodiments, the imaging includes but is not limited to laser scanning microscopy, immunohistochemistry, fluorescent microscopy, radiographic imaging and the like.

20 [0037] In a further aspect, the invention provides *in vivo* methods for diagnosing a cancer. The methods include identifying a subject at risk for or suspected of having pancreatic cell cancer; administering a diagnostic composition comprising a peptide ligand of SEQ ID NO:1 conjugated to an imaging molecule to the subject, and imaging the imaging molecule within the subject using *in vivo* imaging. In some embodiments, the pancreatic cell cancer is a pancreatic ductal adenocarcinoma. In some embodiments, the imaging molecule is a magnetofluorescent particle. In some embodiments, the magnetofluorescent particle comprises a near infrared (NIR) fluorochrome (NIRF). In some embodiments, the composition is administered via route selected from the group consisting of intradermal, subcutaneous, intraperitoneal, intravenous, intraarterial, oral, and gastric routes. In some embodiments, the *in vivo* imaging includes but is not limited to magnetic resonance imaging (MRI), intravital laser scanning microscopy, endoscopy, and radiographic imaging.

25 [0038] In another embodiment, the present invention provides methods for surgically removing pancreatic cancer cells. The methods include a) providing, i) a composition comprising a peptide for distinguishing a pancreatic cancer cell from a pancreatic non-cancer cell, wherein the peptide is SEQ ID NO:1, ii) a subject known to have pancreatic cancer, iii) an *in vivo* imaging device, and b) administering the composition to a subject, c) imaging pancreatic cancer cells *in vivo* with the imaging device, and d) removing pancreatic cancer cells from the subject. In some embodiments, the administering is an intravenous injection of 30 mg Fe/kg (milligram Iron/kilogram). In some embodiments, the administering is an intravenous injection of 2.6 mg/kg.

30 [0039] The present invention provides a method of treating a patient with cancer, comprising, a) providing, i) a cancer patient in need of treatment, ii) a pharmaceutical composition comprising a ligand, wherein the ligand binds to a biomarker of the present invention, and b) administering the treatment composition to the patient. In some embodiments, the pharmaceutical composition further comprises a therapeutic agent. In some embodiments, the therapeutic agent is selected from the group consisting of a fusion protein, a toxin, a drug. The present inventions are not limited by the type of cancer. Indeed, various types of cancer are contemplated for use with the detection methods of the present inventions

including but not limited to lung cancer, bladder cancer, head and/or neck cancer, breast cancer, esophageal cancer, mouth cancer, tongue cancer, gum cancer, skin cancer (e.g., melanoma, basal cell carcinoma, Kaposi's sarcoma, etc.), muscle cancer, heart cancer, liver cancer, bronchial cancer, cartilage cancer, bone cancer, stomach cancer, prostate cancer, testicular cancer, ovarian cancer; cervical cancer, endometrial cancer, uterine cancer, pancreatic cancer, colon cancer, colorectal, gastric cancer, kidney cancer, bladder cancer, lymphoma cancer, spleen cancer, thymus cancer, thyroid cancer, brain cancer, neuronal cancer, mesothelioma, gall bladder cancer, ocular cancer (e.g., cancer of the cornea, cancer of uvea, cancer of the choroids, cancer of the macula, vitreous humor cancer, etc.), joint cancer (such as synovium cancer), glioblastoma, white blood cell cancer (e.g., lymphoma, leukemia, etc.), hereditary non-polyposis cancer (HNPC), colitis-associated cancer, etc. Cancers are further exemplified by sarcomas (such as osteosarcoma and Kaposi's sarcoma).

[0040] The present invention provides a method of treating a patient with pancreatic cancer, comprising, a) providing, i) a cancer patient in need of treatment, ii) a pharmaceutical composition comprising a ligand, wherein the ligand binds to plectin-1 or a fragment thereof, and b) administering the treatment composition to the patient. In some embodiments, the pharmaceutical composition further comprises a therapeutic agent. In some embodiments, the therapeutic agent is selected from the group consisting of a fusion protein, a toxin, and a drug.

[0041] In an additional aspect, the present invention provides methods of treating a subject who has pancreatic cancer. The methods include identifying a subject in need of treatment, e.g., on the basis that they have pancreatic cancer, and administering a therapeutically effective amount of a pharmaceutical composition including a peptide including SEQ ID NO:1 linked to a cytotoxic agent, e.g., a toxin or a drug.

[0042] In another aspect, the present invention provides methods for identifying a cancer cell-binding partner (receptor) having selective affinity for a peptide ligand. The methods include selectively immobilizing a diverse population of binding molecules to a solid support, contacting (e.g., simultaneously contacting) the diverse population immobilized on the solid support with one or more peptide ligands and determining at least one binding molecule which selectively binds to one or more of the peptide ligands, including those expressed by a bacteriophage. Also provided are methods for identifying peptide ligands having selective affinity for a tumor antigen (binding molecule). The methods include selectively immobilizing a tumor antigen to a solid support, contacting (e.g., simultaneously contacting) the immobilized tumor antigen on the solid support with one or more peptide ligands and identifying at least one peptide ligand which selectively binds to one or more of the tumor antigens. Also provided are isolated binding peptides ("peptide ligands") that are selective for a tumor antigen, in particular peptide ligands for plectin-1.

[0043] Also described herein are rapid and efficient methods for the identification of binding molecules that exhibit selective affinity for one or more peptide ligands of interest. The methods are advantageous in that they allow the simultaneous screening of multiple binding molecules against multiple peptide ligands of interest. Moreover, very little information is required regarding the identity or function of either the binding molecule or the ligand for use in the present inventions. For example, diverse populations of binding molecules can be simultaneously screened against diverse populations of peptide ligands to rapidly identify numerous molecules exhibiting a desired binding specificity. The methods described herein can therefore be advantageously applied for the discovery of specific reagents, such as peptide ligands and biomarkers, for diagnosis and treatment of human diseases.

Definitions

[0044] To facilitate an understanding of the present invention, a number of terms and phrases are defined below:

[0045] The use of the article "a" or "an" is intended to include one or more. As used in this application, the singular form "a," "an," and "the" include plural references unless the context clearly dictates otherwise. For example, the term "an agent" includes a plurality of agents, including mixtures thereof.

[0046] As used herein, the term "patient" or "subject" is an individual having symptoms of, or at risk for, pancreatic cancer or other malignancy. Patients may be human or non-human and may include, for example, animal strains or species used as "model systems" for research purposes, such a mouse model as described herein. A patient may include either adults or juveniles (e.g., children). The term "patient", further refers to any living organism, preferably a mammal (e.g., human or non-human) that may benefit from the administration of compositions contemplated herein.

[0047] As used herein, the term "animal" refers to any animal, preferably a mammal, and more preferably, the mammal includes, without limitation, human and non-human animals such simians, rodents, ovines, bovines, ruminants, lagomorphs, porcines, caprines, equines, canines, felines, aves, etc. Preferred non-human animals are members of the Order Rodentia (e.g., mouse and rat), sheep, pig, rabbit or cattle.

[0048] As used herein, the term "benefit" refers to but is not limited to a diagnostic assay, a diagnosis, a surgical tool, and the like.

[0049] As used herein, the term "subject suspected of having cancer" refers to a subject that presents one or more symptoms indicative of a cancer (e.g., a noticeable lump or mass) or is being screened for a cancer (e.g., during a routine physical). A subject suspected of having cancer may also have one or more risk factors. A subject suspected of having

cancer has generally not been tested for cancer. However, a "subject suspected of having cancer" encompasses an individual who has received an initial diagnosis but for whom the stage of cancer is not known. The term further includes people who once had cancer (e.g., an individual in remission).

[0050] As used herein, the term "subject at risk for cancer" refers to a subject with one or more risk factors for developing a specific cancer. Risk factors include, but are not limited to, gender, age, genetic predisposition, environmental exposé, and previous incidents of cancer, preexisting non-cancer diseases, and lifestyle.

[0051] As used herein, the term "suffering from disease" refers to a subject (e.g., a human) that is experiencing a particular disease. It is not intended that the present invention be limited to any particular signs or symptoms, nor disease. Thus, it is intended that the present invention encompass subjects that are experiencing any range of disease, from sub-clinical to full-blown disease, wherein the subject exhibits at least some of the indicia (e.g., signs and symptoms) associated with the particular disease.

[0052] As used herein, the terms "sample" and "specimen" are used in their broadest sense and encompass samples or specimens obtained from any source, including buffer solutions, saline solutions, cell culture media, etc.

[0053] As used herein, the term "biological samples" refers to samples or specimens obtained from animals (including humans, domestic animals, as well as feral or wild animals, such as ungulates, bear, fish, lagomorphs, rodents, etc.), and encompasses cells, fluids, solids, tissues, and gases. In preferred embodiments of this invention, biological samples include tissues (e.g., biopsy material), cell lines, cells isolated from tissue (whether or not the isolated cells are cultured after isolation from tissue), fixed cells (e.g., fixed for histological and/or immunohistochemical analysis), cerebrospinal fluid (CSF), serous fluid, blood, and blood products such as plasma, serum and the like. However, these examples are not to be construed as limiting the types of samples find use with the present invention.

[0054] As used herein, the term "biopsy tissue" refers to a sample of tissue that is removed from a subject for the purpose of determining if the sample contains cancerous tissue. In some embodiment, biopsy tissue is obtained because a subject is suspected of having cancer. The biopsy tissue is then examined for the presence or absence of cancer.

[0055] As used herein, the terms "peptide," "polypeptide" and "protein" all refer to a primary sequence of amino acids that are joined by covalent "peptide linkages." In general, a peptide consists of fewer amino acids than a full-length protein, typically from 2-50 amino acids.

[0056] As used herein, the term "polypeptide" may encompass both peptides and proteins. A peptide, polypeptide or protein may be synthetic, recombinant or naturally occurring. A synthetic peptide is a peptide that is produced by artificial means *in vitro* (e.g., was not produced *in vivo*). A "protein", "peptide", or "polypeptide" amino acid sequence may also comprise chemical compounds.

[0057] As used herein, the term "phage" or "φ" is synonymous to a "bacteriophage" refers to a virus, for example, a M13, T4, and the like, that infects a bacterium, such as an *E. coli*. Phage may also refer to an individual virion.

[0058] As used herein, the term "phage displayed peptide" or "bacteriophage displayed peptide" or "phage-displayed peptide" refer to a bacteriophage viron expressing one peptide nucleic acid sequence where the peptide is located on the outside surface of the viron. One virion may display one peptide or multiple copies of the peptide, such as five copies, seven copies, et cetera.

[0059] As used herein, the term "phage display" in reference to a library refers to a selection technique in which a library of peptides or protein or variants thereof expressed on the outside of a phage viron, while the genetic material encoding each variant resides on the inside variants, for example, a commercial Ph.D.™-7 Phage Display Peptide Library (New England Biolabs) provides 7 mer peptides in excess of two billion independent clones, a Ph.D.-12 Phage Display Peptide Library (New England Biolabs) provides 12 mer peptides, a noncommercial Phage Display Peptide Library provides a variety of peptide sizes from a several sources, randomly generated or chosen, such as for a certain cell type or ligand, and the like.

[0060] The terms "isolated" and "purified" in the context of "peptide sequences" refer to the separation of the desired peptide sequence(s) from non-desired peptide sequences and other contaminants (e.g. phage virions not expressing the desired peptide sequence, truncated or misformed synthetic peptides, lipids, carbohydrates, nuclei acids, etc.). The terms "isolated" and "purified" do not necessarily mean isolated and purified to 100% homogeneity, although this is also contemplated. Rather, the terms mean isolated and purified to at least 50% homogeneity. In a preferred embodiment, the peptide sequences are isolated and purified to at least 75% homogeneity. In a more preferred embodiment, the peptide sequences are isolated and purified to at least 90% homogeneity. After isolation and/or purification, the peptide sequences can then be conjugated to, mixed with or added to other compounds or molecules.

[0061] As used herein, the term "isolated" in reference to a "phage" or "phage virion" refers to a phage displaying a peptide obtained after one or more screening assays, such as a peptide - receptor screening assay of the present inventions, wherein a population of phage displayed peptides of a phage display library are screened for binding to a cancer cell receptor, such that a bound phage is "isolated" from a phage library upon binding to a cancer cell molecule. An "isolated phage" also refers to a phage virion eluted from a bound phage.

[0062] As used herein, the term "isolated" in reference to a peptide ligand refers to a peptide ligand encoded by an isolated phage expressing that peptide, such that the term includes but is not limited to the peptide ligand as expressed

by and including the phage displayed peptide, the nucleic acid sequence of the isolated peptide, the amino acid sequence of the isolated peptide, natural and synthetic forms of the isolated peptide ligand and the like. Thus an exemplary "isolated peptide ligand" of the present inventions includes but is not limited to a peptide expressed by any one of Clones 4, 15 and 27.

5 [0063] As used herein, the term "derives from" or "derived from" in reference to a peptide ligand, in particular an isolated peptide ligand, refers to a peptide ligand sequence obtained from an isolated phage virion displaying that peptide. However, a peptide, peptide derivative, or peptide mimetic, analogues and mimetic compounds are also intended to be included within the definition of this term.

10 [0064] As used herein, the term "mimetic" refers to any molecule that mimics the binding of a ligand to a biomarker (receptor) including a peptide, a non-peptide mimetic, a small molecule mimetic and the like.

15 [0065] As used herein, the term "small organic molecule" as used herein, refers to any molecule of a size comparable to those organic molecules generally used in pharmaceuticals. The term excludes biological macromolecules (e.g., proteins, nucleic acids, etc.). Preferred small organic molecules range in size from approximately 10 Da up to about 5000 Da, more preferably up to 2000 Da, and most preferably up to about 1000 Da.

20 [0066] As used herein, the term "desired" in reference to a peptide ligand refers to a peptide ligand capable of identifying a cancer cell. As used herein, the term "desired" in reference to a binding partner for a peptide ligand refers to a biomarker, including but not limited to a cancer cell biomarker.

25 [0067] As used herein, the term "fluorochrome derived from" in reference to a fluorochrome refers to any fluorochrome with a similarity of structure, for example, a fluorochrome with an ester group and the fluorochrome after loosing the charged ester group in a covalent linkage to a peptide, any of a family of fluorochrome compounds, such as a fluorescein (IUPAC: 3',6'-dihydroxyspiro[2-benzofuran-3,9'-xanthene]-1-one) family of compounds, including related and similar compounds, but not limited to fluorescein isothiocyanate, fluorescein isothiocyanate isomer I, and the like.

30 [0068] As used herein, the term "biomarker" refers to generally a molecule or substance that is an indicator of a biologic state, such as protein or chemical for distinguishing a specific stage of development, for example, distinguishing a pluripotent cell, a stem cell, a differentiated cell, such as a nerve cell, from other cells. A biomarker is an indicator of a normal biologic process, such as maturation, a pathogenic process, such as cancer cell development, or a pharmacological response to a therapeutic intervention, such as indication of the resolution of a disease or loss of cancer cells where a biomarker may be lost or gained. In particular, a biomarker of the present inventions refers to a molecule with altered expression, such as increased expression, altered location, in a cancer cell compared to a noncancer cell of a similar lineage. When referring to "distinguishing a cancer cell," in particular a cancer cell from a noncancer cell, distinguishing refers to reagents and assays used to detect the expression of one or more proteins, peptides, or genes in each cell type (e.g., including but not limited to, the cancer markers of the present invention). Examples of suitable reagents include but are not limited to, peptide ligands, nucleic acid probes capable of specifically hybridizing to the gene of interest, aptamers, PCR primers capable of specifically amplifying the gene of interest, and antibodies capable of specifically binding to proteins expressed by a gene of interest including but not limited to a biomarker. Other non-limiting examples can be found in the description and examples below. The term "biomarker" in reference to a cellular molecule refers to a molecule of the present inventions for identifying a subset of cells, for example, a biomarker for identifying a pancreatic cancer from a noncancer pancreatic cell, such as plectin-1.

35 [0069] As used herein, the term "plectin-1" refers to any molecule comprising a fragment or portion of plectin-1, such as a fragment of the plectin-1 amino acid sequence, for example, any one of SEQ ID NOs:8-23 or 24, and fragments thereof, as well as isolated nucleic acids encoding those amino acid sequences.

40 [0070] As used herein, the term "peptide ligand" (or the word "ligand" in reference to a peptide) refers to a protein fragment that specifically binds to a molecule, such as a protein, carbohydrate, and the like. A receptor can be essentially any type of molecule such as polypeptide, nucleic acid, carbohydrate, lipid, or any organic derived compound. Specific examples of ligands are peptide ligands of the present inventions.

45 [0071] As used herein, the term "selective" or "selectively" when referring to the binding of a receptor molecule to a ligand refers to an interaction that is discriminated from unwanted or non-specific interactions. Discrimination includes but is not limited to affinity of ligand for a receptor molecule, for example, determined by ligand-receptor binding assays, such as phage displayed peptide ligand - cell binding assays, phage displayed peptide ligand biopsy screening, affinity purification, such as High-performance liquid chromatography (HPLC) and the like, competitive binding assays, panning assays, affinity assays, avidity assays, ELISA assays, etc., either qualitatively or quantitatively. For example, affinity may be measured qualitatively by a panning assay, ELISA, and the like, as shown by heat maps where affinity was depicted as mean absorbance values of indicated clones in ELISA assay) and specificity was determined by a ratio of clones' affinity to tumor cells versus normal ductal cells, for example, a specific peptide ligand binding interaction showed a 2 fold higher ratio of absorbance for PDAC cells versus normal ductal cells.

50 [0072] Further, affinity may be measured quantitatively by calculating an affinity constant (association constant) (K_a) by measuring the strength of binding of the components in a complex, such that components A and B where a binding equilibrium is represented by A (ligand) + B (receptor molecule) = AB , the association constant is given by $[AB]/[A][B]$,

and becomes larger with tighter binding between A and B and smaller with looser binding between A and B. As opposed to a dissociation constant (Kd) referring to a measure of the tendency of a complex to dissociate such that a dissociation constant is represented by $[A][B]/[AB]$, where a tighter binding results in a smaller Kd while a looser binding results in a larger Kd. For example, an association constant Kd for a selective binding molecule interaction with a ligand may range from 10^{-3} M to a picoM value, for example, an association constant Kd for a selective binding interaction is generally greater than 10^{-3} M, is preferably greater than 10^{-4} M, is more preferably greater than 10^{-5} M, and further more preferably greater than 10^{-6} M. High affinity interactions are generally greater than 10^{-8} M to 10^{-9} M, and more preferably greater than 10^{-9} M.

[0073] As used herein, the term "avidity" refers to a total binding strength of a ligand with a receptor molecule, such that the strength of an interaction comprises multiple independent binding interactions between partners, which can be derived from multiple low affinity interactions or a small number of high affinity interactions.

[0074] As used herein, the term "attach", or "attachment", or "attached", or "attaching", used herein interchangeably with "bind", or "binding" or "binds" or "bound" refers to any physical relationship between molecules that results in forming a stable complex, such as a physical relationship between a ligand, such as a peptide or small molecule, with a "binding partner" or "receptor molecule." The relationship may be mediated by physicochemical interactions including, but not limited to, a selective noncovalent association, ionic attraction, hydrogen bonding, covalent bonding, Van der Waals forces or hydrophobic attraction.

[0075] The terms "specific binding" or "specifically binding" when used in reference to the interaction of a peptide (ligand) and a receptor (molecule) also refers to an interaction that is dependent upon the presence of a particular structure (i.e., an amino sequence of a ligand or a ligand binding domain within a protein); in other words the peptide comprises a structure allowing recognition and binding to a specific protein structure within a binding partner rather than to molecules in general. For example, if a ligand is specific for binding pocket "A," in a reaction containing labeled peptide ligand "A" (such as an isolated phage displayed peptide or isolated synthetic peptide) and unlabeled "A" in the presence of a protein comprising a binding pocket A the unlabeled peptide ligand will reduce the amount of labeled peptide ligand bound to the binding partner, in other words a competitive binding assay.

[0076] "Specifically binds" means that one molecule, such as a binding moiety, e.g., an oligonucleotide or antibody, binds preferentially to another molecule, such as a target molecule, e.g., a nucleic acid or a protein, in the presence of other molecules in a sample.

[0077] As used herein, the term "amino acid sequence," as used herein, refers to the primary (i.e., linear) structure of a protein, peptide, or polypeptide wherein the individual amino acids are linked by peptide bonds.

[0078] As used herein, the term "receptor," as used herein, refers to any "binding molecule" or "binding partner" (e.g., a cancer cell protein "recognized" or "bound" or "eluted from" a peptide ligand) including but not limited to a peptide, protein or glycoprotein to which an amino acid peptide sequence of the present invention appears to interact with or to specifically bind. For example, a binding molecule may reside on a cell surface or within a cell. An exemplary binding molecule is a plectin-1 molecule that interacts with SEQ ID NO:1 of the present invention. As used herein, the term "binding molecule" also refers to a molecule of sufficient size and complexity expressed by a cancer cell or tumor cell so as to be capable of selectively binding a peptide ligand. Such molecules are generally macromolecules, such as polypeptides, however include nucleic acids, carbohydrates and lipids. The size of a binding molecule is not important so long as the molecule exhibits or can be made to exhibit binding activity with a peptide ligand.

[0079] As used herein, the term "target cell" in reference to a "cancer cell" or "cell" or "host cell" refers to any cell or molecule used as a target for a peptide ligand in any of the assays of the present invention. "Target cell" also refers to any cell that either naturally expresses particular biomarkers of interest or is genetically altered so as to produce normal or mutated biomarkers.

[0080] As used herein, the term "target binding molecule" or "target receptor" refers to a binding partner molecule, both known and unknown molecules, of a peptide ligand of interest, such as an isolated peptide ligand.

[0081] The terms "variant" and "mutant" when used in reference to a polypeptide refer to an amino acid sequence that differs by one or more amino acids from another, usually a related polypeptide. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties. One type of conservative amino acid substitution refers to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. More rarely, a variant may have "non-conservative" changes (e.g., replacement of a glycine with a tryptophan). Similar minor variations may also include amino acid deletions or insertions (i.e., additions), or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing biological activity may

be found using computer programs well known in the art, for example, DNASTar software. Variants can be tested in functional assays. Preferred variants have less than 10%, preferably less than 5% and still more preferably less than 2% changes (whether substitutions, deletions, and so on).

[0082] As used herein, the term "nucleic acid molecule" refers to any nucleic acid containing molecule, including but not limited to, DNA or RNA. The term encompasses sequences that include any of the known base analogs of DNA and RNA including, but not limited to, 4-acetylcytosine, 8-hydroxy-N-6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudoouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N-6-isopentenyladenine, uracil-5-oxyacetic acid metlylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiacytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiacytosine, and 2,6-diaminopurine.

[0083] As used herein, the term "gene" refers to a nucleic acid (e.g., DNA) sequence that comprises coding sequences necessary for the production of a polypeptide, precursor, or RNA (e.g., rRNA, tRNA and the like). The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired activity or functional properties (e.g., enzymatic activity, ligand binding, signal transduction, immunogenicity, etc.) of the full-length or fragment are retained. The term also encompasses the coding region of a structural gene and the sequences located adjacent to the coding region on both the 5' and 3' ends for a distance of about 1 kb or more on either end such that the gene corresponds to the length of the full-length mRNA. Sequences located 5' of the coding region and present on the mRNA are referred to as 5' non-translated sequences. Sequences located 3' or downstream of the coding region and present on the mRNA are referred to as 3' non-translated sequences.

[0084] As used herein, the term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences." Introns are segments of a gene that are transcribed into nuclear RNA (hnRNA); introns may contain regulatory elements such as enhancers. Introns are removed or "spliced out" from the nuclear or primary transcript; introns therefore are absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

[0085] As used herein, the term "gene expression" refers to the process of converting genetic information encoded in a gene into RNA (e.g., mRNA, rRNA, tRNA, or snRNA) through "transcription" of the gene (e.g., via the enzymatic action of an RNA polymerase), and for protein encoding genes, into protein through "translation" of mRNA. Gene expression can be regulated at many stages in the process. "Up-regulation" or "activation" refers to regulation that increases the production of gene expression products (e.g., RNA or protein), while "down-regulation" or "repression" refers to regulation that decrease production. Molecules (e.g., transcription factors) that are involved in up-regulation or down-regulation are often called "activators" and "repressors," respectively.

[0086] As used herein, the term "label" as used herein refers to any atom or molecule that can be used to provide a detectable (preferably quantifiable) signal, and that can be attached to a nucleic acid or protein. Labels may provide signals detectable by fluorescence, radioactivity, colorimetry, gravimetry, X-ray diffraction or absorption, magnetism, enzymatic activity, and the like. A label may be a charged moiety (positive or negative charge) or alternatively, may be charge neutral. Labels can include or consist of nucleic acid or protein sequence, so long as the sequence comprising the label is detectable.

[0087] As used herein, the term "linker" refers to a molecule or sequence, such as an amino acid sequence, that attaches, as in a bridge, one molecule or sequence to another molecule or sequence. "Linked," "conjugated," or "coupled" means attached or bound by covalent bonds, or non-covalent bonds, or other bonds, such as van der Waals forces.

[0088] As used herein, the term "cell culture" refers to any *in vitro* culture of cells. Included within this term are continuous cell lines (e.g., with an immortal phenotype), primary cell cultures, finite cell lines (e.g., non-transformed cells), and any other cell population maintained *in vitro*.

[0089] As used herein, the term "epithelial cell" refers to a cuboidal-shaped, nucleated cell generally located on the surface of a tissue, such as a pancreatic ductal cell. A layer of epithelial cells generally functions to provide a protective lining and/or surface that may also be involved in transport processes. An epithelial cell is readily distinguished from a non-epithelial cell (e.g., muscle cell, nerve cell, secretory cell, etc.) using histological methods well known in the art.

[0090] As used herein, the term "endothelial cells", as used herein, refers to any cell that may provide a lining for a bodily organ comprising a lumen (e.g., blood vessels, intestines, lymphatic vessels or ducts etc.). Usually, endothelial cells provide physical and chemical protection as well a selective absorption of nutrients or other metabolically active compounds.

[0091] As used herein, "ductal cell", in reference to a pancreas, refers to any cell that forms or has the capability to form or originated from the ductal lining of ducts within and exiting from the pancreas.

[0092] As used herein, "pancreas" in reference to an organ refers to a collection of a plurality of cell types held together by connective tissue, such that the plurality of cells include but are not limited to acini cells, ductal cells and islet cells. The "acini" produce many of the enzymes, such as lipase, which are needed to digest food in the duodenum. The enzymes produced by the acini are carried to the duodenum by small channels called ducts. Typically, ductal cells are held in place by connective tissue in close proximity to vascular cells and nerve cells. Islets of Langerhans are typically embedded between exocrine acini units of the pancreas. Examples of islet endocrine cells are Alpha cells that secrete glucagon which counters the action of insulin while Beta cells secrete insulin, which helps control carbohydrate metabolism.

[0093] As used herein, "pancreatic cancer" refers to cancers that originate in the tissue that comprises a pancreas, such as a pancreatic ductal adenocarcinoma cell.

[0094] As used herein, "adenocarcinoma" refers to a cancerous tumor as opposed to an "adenoma" which refers to a benign (non-cancerous) tumor made up of cells that form glands (collections of cells surrounding an empty space).

[0095] As used herein, "pancreatic ductal adenocarcinoma cell" refers to a cancerous cell that had the capability to form or originated from the ductal lining of the pancreas. A pancreatic ductal adenocarcinoma cell may be found within the pancreas forming a gland, or found within any organ as a metastasized cell or found within the blood stream of lymphatic system.

[0096] As used herein, the term "characterizing cancer in subject" refers to the identification of one or more properties of a cancer sample in a subject, including but not limited to, the presence of benign, pre-cancerous or cancerous tissue, the stage of the cancer, and the subject's prognosis. Cancers may be characterized by the identification of the expression of one or more cancer marker genes, including but not limited to, the cancer markers disclosed herein.

[0097] As used herein, the term "stem cell cancer markers" refers to a gene or peptide expressed by the gene whose expression level, alone or in combination with other genes, is correlated with the presence of tumorigenic cancer cells. The correlation may relate to either an increased or decreased expression of the gene (e.g. increased or decreased levels of mRNA or the peptide encoded by the gene).

[0098] As used herein, the term "instructions for using the kit for detecting cancer in the subject" includes instructions for using the reagents contained in the kit for the detection and characterization of cancer in a sample from a subject.

[0099] As used herein, the term "providing a prognosis" refers to providing information regarding the impact of the presence of cancer (e.g., as determined by the diagnostic methods of the present invention) on a subject's future health (e.g., expected morbidity or mortality, the likelihood of getting cancer, and the risk of metastasis).

[0100] As used herein, the term "post surgical tumor tissue" refers to cancerous tissue (e.g., biopsy tissue) that has been removed from a subject (e.g., during surgery).

[0101] As used herein, the term "subject diagnosed with a cancer" refers to a subject who has been tested and found to have cancerous cells. The cancer may be diagnosed using any suitable method, including but not limited to, biopsy, x-ray, blood test, and the diagnostic methods of the present invention.

[0102] As used herein, the term "non-cancerous" in reference to a pancreatic cell refers to a cell demonstrating regulatable cell growth and functional physiology relative to its developmental stage and activity.

[0103] As used herein, the term "tumor" refers to an abnormal mass of tissue that results from excessive cell division that is uncontrolled and progressive. It is also called a neoplasm. Tumors may be either benign (not cancerous) or malignant.

[0104] As used herein, the term "tumor cell", as used herein, refers to any mass of cells that exhibits any uncontrolled growth patterns or altered physiology. Tumor cells may be derived from any tissue within an organism (e.g., a pancreatic ductal tumor cell).

[0105] As used herein, the term "cancer" is a general term for more than 100 diseases that are characterized by an uncontrolled, abnormal growth of cells. Cancer cells can spread locally or can intravasate and spread via the bloodstream and lymphatic system to other parts of the body and form metastases. Cancer cells that spread are called "malignant."

[0106] As used herein, the terms "cancer" and "cancerous" in reference to a physiological condition in mammals is typically characterized by unregulated cell growth. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer.

[0107] As used herein, the term "malignant" refers to having the properties of anaplasia, penetrance, such as into nearby areas or the vasculature, and metastasis.

[0108] As used herein, the term "invasive," or "metastasis" as used herein, refers to any migration of cells, especially to invasive cancer cells or tumor cells. The term applies to normally invasive cells such as wound-healing fibroblasts and also to cells that migrate abnormally. Although the term is not to be limited by any mechanistic rationale, such cells

are thought to migrate by defeating the body's means for keeping them sufficiently "in place" to function normally. Such cells are "invasive" if they migrate abnormally within a tissue or tumor, or escape the tissue, or invade other tissues.

[0109] As used herein, the term "cell migration" refers to the movement of a population of cells from one place to another. Such movement of cells may be normal as in the movement of neural crest cells during morphogenesis or it may be not normal such as with the movement of malignant cancer cells away from primary sites into nearby areas, the vasculature and thence, into new or secondary sites in the originating or other organs.

[0110] As used herein, the term "in vitro" refers to an artificial environment and to processes or reactions that occur within an artificial environment. In vitro environments consist of, but are not limited to, controlled laboratory conditions.

[0111] As used herein, the term "in vivo" refers to the natural environment (e.g., within an organism or a cell) and to processes or reactions that occur within that natural environment. Alternatively the term "in vitro" refers to performing a given experiment in a controlled environment outside of a living organism, such as an experiment within a laboratory Petri dish.

[0112] As used herein, the term "inhibits" in reference to a peptide ligand, for example, "a peptide ligand which inhibits malignant cell migration" or "a peptide ligand which inhibits cancer cell growth" refers to the partial or total inhibition of migration or growth respectively.

[0113] As used herein, the term "reducing cancer in a patient" in reference to a treatment refers to any treatment for decreasing the number of cancer cells in a patient, slowing the growth of cancer cells in a patient, reducing the metastasis of cancer cells in a patient and includes any type of response for either relieving cancer symptoms or increasing the life-span of a patient.

[0114] As used herein, the term "ELISA" refers to enzyme-linked immunosorbent assay. Numerous methods and applications for carrying out an ELISA are well known in the art, and provided in many sources (See, e.g., Crowther, "Enzyme-Linked Immunosorbent Assay (ELISA)," in Molecular Biomethods Handbook, Rapley et al. [eds.], pp. 595-617, Humana Press, Inc., Totowa, N.J. [1998]; Harlow and Lane (eds.), Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press [1988]; and Ausubel et al. (eds.), Current Protocols in Molecular Biology, Ch. 11, John Wiley & Sons, Inc., New York [1994]; and Newton, et al., (2006) *Neoplasia*. 8:772-780). In some embodiments of the present invention, a "direct ELISA" protocol is provided, where a target-binding molecule, such as a cell, cell lysate, or isolated protein, is first bound and immobilized to a microtiter plate well. In an alternative embodiment, a "sandwich ELISA" is provided, where a target-binding molecule is attached to the substrate by capturing it with an antibody that has been previously bound to the microtiter plate well. The ELISA method detects an immobilized ligand - receptor complex (binding) by use of fluorescent detection of fluorescently labeled ligands or an antibody-enzyme conjugate, where the antibody is specific for the antigen of interest, such as a phage virion, while the enzyme portion allows visualization and quantitation by the generation of a colored or fluorescent reaction product. The conjugated enzymes commonly used in the ELISA include horseradish peroxidase, urease, alkaline phosphatase, glucoamylase or O-galactosidase. The intensity of color development is proportional to the amount of antigen present in the reaction well.

[0115] As used herein, a "pharmaceutical composition" is a composition comprising a sequence or sequences of the present invention. The pharmaceutical composition may further comprise a carrier, a pharmaceutically acceptable excipient, and the like. The terms "pharmaceutical composition" and "therapeutic composition" are used herein interchangeably. It is not intended that the pharmaceutical compositions be limited to any particular carrier or excipient or other ingredient.

[0116] As used herein, the term "therapeutic agent" refers to chemicals or drugs or proteins that are able to inhibit cell function, inhibit cell replication or kill mammalian cells, preferably human cells.

[0117] As used herein, the term "kit" is used in reference to a combination of reagents, in particular a peptide ligand of the present inventions, and other materials. It is contemplated that the kit may include reagents such as phage displayed peptides, isolated peptide ligands, peptide ligands conjugated to any one of a fluorescent marker, a nanoparticle, a conjugate for MRI, a conjugate for therapeutics, antibodies, control proteins, as well as testing containers (e.g., microtiter plates, etc.). It is not intended that the term "kit" be limited to a particular combination of reagents and/or other materials.

[0118] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

[0119] Other features and advantages of the invention will be apparent from the following detailed description and figures, and from the claims.

DESCRIPTION OF THE DRAWINGS

[0120]

5 Fig. 1A is a heat map showing affinity (mean absorbance values of indicated clones in ELISA assay) and specificity (ratio of clones' affinity to tumor cells versus normal ductal cells) of selected clones. Data are displayed in terms of higher rankings (green) to lower rankings (red).

10 FIG. 1B is a histogram showing specificity of Clone 27 for pancreatic ductal adenocarcinoma cells validated via flow cytometry (FACS) showing FITC-labeled clone 27 bound specifically to mouse PDAC cells (FITC-27 was incubated with PDAC or normal mouse ductal cells then analyzed via FACS).

15 FIG. 1C is a bar graph showing FACS analysis of binding of FITC-27 or FITC unrelated phage clone (negative control) to the indicated cells. Data plotted are the mean fluorescence units obtained from FACS analysis.

20 FIG. 2 is a set of 15 panels showing exemplary Clone 27 *in vivo* detection of human and mouse pancreatic ductal adenocarcinoma cells (PDAC). Fluorescein isothiocyanate (FITC) labeled-clone 27 or wild type phage (no peptide insert) were incubated with frozen sections of indicated tissue, mouse and human, pancreas sections stained with Hematoxylin and Eosin (H & E) Clone 27 (PDAC-27).

25 FIGs. 3A-C are each four panels showing exemplary *in vivo* validation of Clone-27. 3A, wild type; 3B, 29-week old Kras/p16 ^{+/-}; and 3C, 12-week old Kras/p53 ^{L/+} mice were injected with Cyanine 5.5 labeled phage Clone 27 and SYTOX green (nonspecific cell-labeling agent) then imaged via intravital confocal microscopy. Correlative Histology: Pancreata from intravital imaging experiments were embedded in Optimal Cutting Temperature (OCT) compound, frozen, and stained with anti-M13 antibody (third column) Hematoxylin and Eosin (H & E) (fourth column). Black boxes correspond to regions of PanINs (3B) or PDAC (3C) and are magnified in the anti-M13 photomicrograph. Upper boxes in 3B and 3C, HE and inset in anti-M13 photomicrograph (third and fourth columns) correspond to uninvolved adjacent regions.

30 Fig. 4A is two panels. The left panel shows a silver stained gel of mouse PDAC cells incubated with either sulfo-SAED conjugated phage Clone 27 and biotin modified phage Clone 27 or control phage, exposed to light, lysates were incubated with streptavidin coated beads. Precipitated protein was eluted with (Dithiothreitol) DTT then run on an SDS PAGE gel. The right panel is a Far Western; PDAC lysates were loaded onto an SDS PAGE gel then transferred and analyzed with clone-27 or control (no insert) biotinylated phage.

35 FIG. 4B is a list of tryptic digest products. A band corresponding to clone-27 affinity purified protein from 4A was cut from the gel, digested with trypsin and analyzed via mass spectrometry (SEQ ID NOs: 9-23).

40 FIG. 4C is a blot of affinity purified protein from 4A that was run on a sodium dodecyl sulfate polyacrylamide (SDS) PAGE gel, transferred to PVDF membrane and analyzed for the presence of plectin-1.

45 FIG. 4D is a set of six blots of 293T cells, human umbilical vein endothelial cells (HUVECs), mouse PDAC cells, mouse normal duct cells, Paca-2 cells (human PDAC), and normal human duct cells that were subcellularly fractionated, and the components probed for the presence of plectin-1.

50 FIG. 4E is a set of three photomicrographs of pancreas tissue from wild type (left), 29-week old Kras/p16 ^{+/-} (center), and 12-week old Kras/p53 (right) mice. The tissues were embedded in OCT, frozen and stained with anti-plectin-1 antibody.

55 FIG. 4F is a histogram showing the results of a competition experiment in which mouse PDAC cells were incubated with FITC labeled clone 27 and either plectin-1 or vehicle then analyzed via FACS.

Fig. 5A is a schematic of conjugation of PTP to NP. Control-NP is synthesized the same way with substitution of control peptide for PTP

FIG. 5B is a set of four exemplary intravital confocal microscope images of early pancreatic lesions imaged using PTP-NP (top) or control-NP (bottom) and AF750-labeled bloodpool agent.

FIG. 5C is a pair of low-magnification views of pancreatic fluorescence shows distribution of PTP-NP in distinct areas of the pancreas. White light overlay provides anatomic correlation (left). Dotted line outlines the pancreas.

FIG. 5D is a histogram showing biodistribution of PTP-NP and control-NP.

Fig. 6A is a white light image of the abdominal cavity of injected 9-week old Kras/p53 ^{L/+} animals showing anatomic detail.

FIG. 6B is an intravital confocal microscope image of early pancreatic lesions imaged using PTP-NP (dark grey) and AF750-labeled bloodpool agent (light grey) demonstrates focal uptake distinct from the vasculature.

FIG. 6C is a low magnification view of pancreatic fluorescence showing distribution of PTP-NP in distinct areas of the pancreas.

FIG. 6D is a white light overlay image providing anatomic correlation. Dotted line outlines the pancreas.

Figs. 7A-C show an exemplary Magnetic resonance imaging (MRI) and correlative histology. 7A, Three adjacent slices from an *ex vivo* MRI of the pancreas from a 9-week old Kras/p53^{L/+} mouse demonstrates focal nanoparticle uptake (yellow arrow), which corresponds to tumor seen on correlated Hematoxylin and Eosin (H & E) sections (7B)

but not to regions of ductal metaplasia or normal pancreas (labeled). (7C) Fluorescence microscopy of adjacent sections demonstrate uptake of Cy5.5-labeled plectin-1 targeted peptide conjugated nanoparticles (PTP-NP) in regions of tumor (left) but not in adjacent tissue (right).

5 FIG. 8A is a line graph showing CLIO-Cy5.5 absorbance spectroscopy used for quantitation of number of Cy5.5/nanoparticle.

FIG. 8B is a dot graph of size distribution of CLIO via light scattering.

FIG. 8C is a line graph showing the results of absorbance spectroscopy of PTP-NP-Cy5.5 used to quantitate the number of peptides/nanoparticle. Notice background absorbance from nanoparticle below 500 nm, whose contribution is subtracted via reference to unreacted CLIO.

10 Fig. 9 is a bar graph showing an exemplary phage clone validation Via ELISA. After selection and subtraction, 30 individual phage clones were picked, amplified, and analyzed for affinity and specificity via ELISA.

Fig. 10A is a set of three photomicrographs of Cy5.5-labeled phage clone 27 and RITC-labeled phage clone 15 that were coinjected into Kras/p53L/L, and tumor binding analyzed via intravital confocal microscopy.

15 FIG. 10B is a photomicrograph of an unrelated phage clone injected into Kras/p53L/b and analyzed via intravital confocal microscopy.

FIG. 10C is a photomicrograph of Cy5.5-labeled phage clone 27 injected into a wild-type mouse and analyzed via intravital confocal microscopy.

FIGs. 11A-B show an exemplary amino acid sequence for human plectin-1, SEQ ID NO:24.

20 DETAILED DESCRIPTION OF THE INVENTION

[0121] The present invention relates, at least in part, to compositions and methods for providing cancer cell biomarkers, such as biomarkers of pancreatic ductal adenocarcinoma (PDAC) cell biomarkers, and binding molecules for diagnosis and treatment of cancer. "Accessible" proteomes are disclosed, and methods of use thereof for selecting biomarkers, 25 such as plectin-1, for identifying PDACs. Imaging compositions that include magnetofluorescent nanoparticles conjugated to peptide ligands are described, and methods of use thereof for identifying PDACs. Finally, therapeutic compositions comprising antibodies and peptide ligands that bind to plectin-1 coupled to cytotoxic agents, and methods of use thereof for the treatment of cancer, e.g., PDAC, are also described herein.

[0122] Specifically, as described herein a phage display screen and early passage PDAC cell lines isolated from 30 mouse models were used to identify peptides that distinguish both human and murine PDAC cells from normal pancreatic ductal cells.

[0123] Described herein are peptide ligands identified using phage displayed peptides, and then developed targeted imaging agents through conjugation of the isolated peptides to a magnetofluorescent nanoparticle; the results set forth herein demonstrate that these agents can effectively detect emerging tumors in vivo and in vitro. In this approach, the 35 binding partners of the phage displayed peptides represent a snapshot of the "accessible" proteome rather than an extensive list of overexpressed cellular proteins. Using these methods, the present inventors identified plectin-1 as a novel PDAC biomarker; expression levels of plectin-1 are modestly upregulated in cancer cells as compared to normal cells, and in addition the protein is aberrantly distributed to the cell membrane of cancerous cells and is thus accessible for probing with imaging agents as described herein.

[0124] As described herein, a multimodal nanoparticle-based targeted imaging agent (referred to herein as "PTP-NP") 40 was developed that was capable of identifying PDAC cells in a background of normal, mucinous, and ductal metaplasia of the pancreas. In some embodiments, these imaging agents are contemplated for use in both MRI and endoscopy in high-risk patients.

45 I. Methods for Identifying Pancreatic Cancer Cells

[0125] Pancreatic cancer is a leading cause of cancer-related death in the United States. When pancreatic cancer is found early, surgical removal of the tumor can sometimes provide a cure. Unfortunately, this cancer rarely causes any 50 symptoms in its early stages and the symptoms it does eventually cause include jaundice, abdominal pain, back pain, and weight loss, which are also seen in other illnesses, making early diagnosis difficult.

[0126] Magnetic resonance imaging (MRI) and other noninvasive imaging techniques are used to look at the pancreas, however by the time pancreatic tumors are large enough to show up on MRI scans, they have often already spread. Consequently, in most patients, pancreatic cancer is advanced by the time a diagnosis is made, hence surgery is no longer useful. These patients are given radiotherapy and chemotherapy but these treatments are rarely curative and 55 most patients die within a year of diagnosis.

[0127] Thus, pancreatic ductal adenocarcinoma (PDAC) is often considered an intractable clinical problem, typically presenting with metastasis at the time of diagnosis and exhibiting profound resistance to existing therapies. Because current detection methods are unreliable, considerable ongoing efforts aimed at identifying new PDAC detection biomar-

kers are currently being pursued using a variety of approaches including serum proteomics, expression profiling of tumor tissue, genetic analysis of pancreatic fluid, and methods using combinatorial chemistry (see, e.g., Goggins, *J Clin Oncol* 23:4524-4531 (2005); Misek, et al., *Methods Mol Med* 103:175-187 (2005); Bloomston, et al., *Cancer Res* 66:2592-2599 (2006); Yates, et al., *Anal Chem* 67:1426-1436 (1995); and Joyce, et al., *Cancer Cell* 4:393-403 (2003)).

[0128] Early work to develop better diagnostic and therapeutic molecules focused on the use of antibodies for tumor recognition and drug delivery (see, e.g., Folli, et al., *Cancer Res* 54:2643-2649 (1994); Neri, et al., *Nat Biotechnol* 15:1271-1275 (1997)). However, antibody targeting in the case of molecular imaging often does not have ideal pharmacokinetics, has a limited target-to-background ratio, and furthermore has limited capacity for carrying magnetic resonance (MR)-detectable imaging agents unless extensively modified. Peptides are useful as targeting moieties with various high-throughput screening methods being utilized to select for ideal specificity, affinity, and pharmacokinetics. To their detriment as imaging agents, peptides generally have very short vascular half-lives (approximately 5 minutes) and a lower affinity than their multivalent counterparts. Therefore, a combination of multimodal nanoparticles with targeting peptides may circumvent some of these issues since they can be designed as platforms with optimized pharmacokinetics, pharmacokinetics, allow multivalent peptide attachment, are small enough for targeting, and can be internalized into the cell resulting in signal amplification through intracellular trapping (see, e.g., Kelly, et al., *Circ Res* 96: 327-336 (2005)).

[0129] Moreover, there is a substantial challenge in studying the early molecular changes in PDAC because of the typical presentation of PDAC at advanced stage and the corresponding lack of suitable tissue specimens. Therefore, the present discovery made use of a series of related genetically engineered mouse models of PDAC that harbor the signature gene mutations of the human disease, including Kras activation and deletion of the p53 or Ink4a/Arf tumor suppressors (see Bardeesy et al. *Proc Natl Acad Sci U.S.A.* 103:5947-5952 (2006); and Aguirre et al., *Genes Dev* 17: 3112-3126 (2003)). The tumors in these models exhibit the characteristic multistage histopathological progression (from precursor pancreatic intraepithelial neoplasia (PanINs) (Hansel et al., *Annu Rev Genomics Hum Genet* 4:237-256 (2003)) to metastatic cancer) that defines PDAC in humans, providing tractable model systems for both biological and preclinical studies (Bardeesy et al., *Proc. Natl. Acad. Sci. U.S.A.* 103:5947-5952 (2006)). From these mouse models, primary cell lines were derived from emerging PDAC. These early-stage cancer cell lines, in conjunction with normal pancreatic ductal cells from wild-type mice (described in Schreiber et al., *Gastroenterology* 127:250-260 (2004)), facilitated screening for biomarkers and imaging agents using combinatorial chemistry-based approaches (see the Examples herein).

[0130] In order to overcome these limitations, the present methods include the use of novel molecular markers and imaging probes for incipient PDAC that enable earlier detection and guide the development of interventive therapies. Described herein are the use of peptide phage display and early passage PDAC cell lines isolated from mouse models to identify peptides that distinguish both human and murine PDAC cells from normal pancreatic ductal cells *in vitro*. In addition to the generation of imaging agents, the binding partners of the surface proteins identified in this approach represent a snapshot of the proteome in aberrant cells and may be useful for the delineation of the underlying signal transduction pathways important to disease progression. Further, these imaging probes are contemplated to provide effective treatments for pancreatic and other cancers. The methods described herein can be used for detecting tumors and pancreatic cancer cells at any stage, including early stages, prior to spreading.

[0131] Although biomarkers were reported for PDAC cells, these markers were problematic in that they appeared to extensively overlap with other cell types, were not conjugated to nanoparticles, or were not plectin-1. Numerous attempts for identifying PDACs were described in publications, for example, in methods for detecting and diagnosing pancreatic cancer, including but not limited to determining the expression level of pancreatic associated genes that discriminate between cancerous and normal cells both *in vivo* and *in vitro*, methods of screening therapeutic agents for treating pancreatic cancer, and methods of treating pancreatic cancer, (Nakamura, et al. "Method For Diagnosing Pancreatic Cancer," United States Patent Application Number 20050260639), a method of qualifying pancreatic cancer status in a subject by measuring at least one biomarker and correlating that measurement with the cancer's status (Chan et al. "Identification Of Biomarkers For Detecting Pancreatic Cancer," United States Patent Application Number 20050095611). None of these references describe plectin-1. Further, in methods and systems for identification of abnormal cell growth, particularly the presence of pancreatic cancer or susceptibility to pancreatic cancer where an identifying candidate agent for treatment of pancreatic cancer is obtained from Affymetrix GeneChip analysis (Hruban et al. "Pancreatic Cancer Diagnosis And Therapies," United States Patent Application Number 20030180747). In this last reference, plectin-1 was identified as one of 97 genes differentially overexpressed in pancreatic cancer cells compared to normal pancreatic cells however this reference does not discuss aberrant cell membrane distribution of plectin-1, or the detection of plectin-1 with a peptide. Moreover, previous attempts at using a peptide marker for identifying Pancreatic adenocarcinoma cells were published, where synthetic peptides of Peptide tyrosine tyrosine (termed "YY" or "PYY" or Pancreatic Peptide "PP"), a 36 amino acid residue peptide amide and fragments thereof, including PYY3-36 and YPIKPEAPGEDASPEEL-NRYYASLRHYLNVLTRQRY (SEQ ID NO:25), peptide YY14-36, in particular, (U.S. Patent No. 5,574,010) showed a high specific binding to pancreatic cancer cells and delivered fluorescent dyes to cancer cells. It was contemplated but not shown, that a strategy of using biotinylated peptides to deliver avidin-dye complexes to cancer cells will allow imaging of pancreatic tumors and delivery of therapeutic agents. However, further publications describe the use of these and

similar peptides for identifying neuronal cells and used for both diagnosing and treating obesity (WO/2004/056314) and treating cancer, for example, colon adenocarcinoma, pancreatic adenocarcinoma, or breast cancer. Additionally, a receptor for PYY was reported as a Y2 receptor, (for example, Patent No. 5,574,010; U.S. Patent No. 5,604,203; U.S. Patent No. 5,696,093; U.S. Patent No. 6,046,167; Gehlert et. al., Proc Soc Exp Biol Med 218:7-22 (1998); Sheikh et al.

5 Am J Physiol, 261:701-15 (1991); Fournier et al., Mol. Pharmacol. 45:93-101 (1994); Kirby et al., J Med Chem 38:4579-4586 (1995); Rist et al., Eur. J. Biochem 247:1019-1028 (1997); Kirby et al., J Med ClzeTn 36:3802-3808 (1993); Grundemar et al., Regulatory Peptides 62:131-136 (1996); U.S. Patent No. 5,696,093 (examples of PYY agonists), and U.S. Patent No. 6,046,167; all of which are herein incorporated by reference in their entirety). However, none of these references disclose the use of a peptide ligand that recognized plectin-1 for identifying pancreatic cancer cells.

10 [0132] As described herein, phage display was used to identify peptides that distinguish mouse and human PDAC cells from normal pancreatic duct cells in vitro. The inventors subsequently conjugated 2 peptides with the highest affinities and specificities for PDACs to magnetofluorescent nanoparticles (CLIO-VT680) and demonstrate that these agents can effectively detect emerging tumors and pre-neoplastic lesions in a relevant transgenic mouse model via intravital confocal microscopy (Olympus IV100) and optical/MR imaging (OV-100, Bruker Pharmascan). Correlative 15 histology confirmed the specific temporal localization of the PDAC targeted agents. Additionally, the peptide-binding partners identified from this approach represent a snapshot of the proteome in aberrant cells and also potential PDAC biomarkers. Using affinity chromatography, binding partners for several phage displayed peptides were identified, and their validity as biomarkers was demonstrated, in particular for Clones 27 and 15. These specific and sensitive probes are contemplated to have clinical utility in the diagnosis and management of PDAC in humans.

20 II. Peptide Ligands for Identifying "Accessible" Receptors in Proteomes of Pancreatic Cancer Cells

[0133] Differential protein processing and/or trafficking, which can be identified using proteomic approaches, represents 25 a potential class of biomarkers that would be missed if looking at cDNA expression data only or using whole-cell proteomics methods. For example, the binding partners of clone 27 and clone 15 identified by the methods described herein represent cancer biomarkers and may shed light on aberrant molecular pathways contributing to PDAC pathogenesis.

[0134] Clone 27 permitted the identification of membrane-localized plectin-1 as a potential new biomarker for PDAC. 30 As described herein, plectin-1 levels are low in normal pancreatic ductal cells, but expression of plectin-1 is upregulated in PanINs and remains elevated in PDAC. Plectin-1 exhibited distinct cytoplasmic and nuclear localization in normal fibroblasts, whereas aberrant expression on the cell membrane is observed in PDAC. In some embodiments, mechanisms 35 of protein upregulation, differential trafficking, and whether a biomarker contributes to disease progression are contemplated for additional use in biomarker diagnostics and treatments of cancer cells. Notable in this regard, recent publications illustrate that plectin-1 can be recruited to the membrane during epithelial cell transformation (Raymond et al., Mol Biol Cell 18: 4210-4221 (2007)). Altered subcellular localization of plectin-1 is also observed in the autoimmune condition, 40 paraneoplastic pemphigus, and in the associated lymphoproliferative neoplasm, Castleman disease (Aho et al., J Invest Dermatol 113: 422-423 (1999)). Plectin-1 has a number of important roles in signal transduction, influencing Rho activity (Andra et al., Genes Dev 12: 3442-3451 (1998)), and serving as a scaffold for proteins involved in protein kinase C (PKC) (Osmanagic-Myers et al., J Biol Chem 279: 18701-18710 (2004)) and AMP-activated protein kinase signaling pathways (Gregor et al., J Cell Sci 119: 1864-1875 (2006)). Thus, plectin-1 in PDAC may have an impact on signaling pathways that regulate cell migration, polarity, and energy metabolism.

Plectin-1 Expression in the Pancreas

[0135] Plectin-1 is a high molecular weight protein (500 kDa) that links intermediate filaments to microtubules and 45 microfilaments, in addition to anchorin the cytoskeleton the plasma and nuclear membranes (reviewed in Sonnenberg, et al., Exp Cell Res 313:2189-2203 (2007)).

[0136] As described herein, plectin-1 levels are low in normal pancreatic ductal cells but its expression is upregulated 50 in PanINs and remains elevated in PDACs. Plectin-1 exhibited distinct cytoplasm and nuclear localization in normal fibroblasts, whereas an aberrant expression on the cell membrane is observed in PDACs. Altered subcellular localization of plectin-1 was also observed in an autoimmune condition, paraneoplastic pemphigus and in the associated lymphoproliferative neoplasm, Castleman's disease (Aho et al., J Invest Dermatol 113:422-423 (1999)).

[0137] Studying the mechanisms of plectin-1 protein upregulation, differential trafficking, and contributions to disease 55 progression are important contemplated experiments. As noted above, plectin-1 has important roles in signal transduction. Thus, plectin-1 in PDACs may have an impact on signaling pathways that regulate cell migration, polarity and energy metabolism related to carcinogenesis.

[0138] Further, publications describe significant overexpression of plectin-1 in a variety of pancreatic cancer tissues and cells compared to their non-cancerous counterparts (such as Hruban, et al. "Pancreatic Cancer Diagnosis And Therapies," United States Patent Application Number 20030180747), where methods and systems were described for

identification of abnormal cell growth, particularly the presence of pancreatic cancer or susceptibility to pancreatic cancer. This reference further discloses methods for identifying candidate agents for treatment of pancreatic cancers. Affymetrix GeneChip analysis identified plectin-1 as one of 97 genes differentially overexpressed in pancreatic cancer cells compared to normal pancreatic cells. Iacobuzio-Donahue et al., Am J Pathol. 160(4):1239-1249 (2002) discloses Affymetrix GeneChip arrays to identify genes differentially expressed in resected pancreas cancer tissues and pancreas cancer cell lines as compared to normal pancreas and gastrointestinal mucosa cells. Table 1 identifies plectin-1 as one of the 97 known genes expressed at least five-fold (6.69-fold) in pancreatic cancers. Sato et al., Am J Pathol. 164(3):903-914 (2004) discloses Affymetrix GeneChip analysis of intraductal papillary mucinous neoplasms (IPMNs) of the pancreas that identifies plectin-1 as one of 673 transcripts significantly overexpressed relative to nonneoplastic pancreatic ductal epithelium. Finally, Johnson et al., Molecular Carcinogenesis 45:814-827 (2006) employs DNA array technology to identify genes differentially expressed in pancreatic tumors (11 PDACs) as compared to non-malignant pancreatic tissues (14 non-malignant bulk pancreatic duct specimens). The results listed in Table 2, of Sato et al., *supra*, confirmed previous findings regarding plectin-1 overexpression (4.5-fold) in PDAC tissues compared to non-cancerous tissues. Further, a potential natural ligand for plectin was identified using anti-plectin antibodies for immunoprecipitation of plectin-periplakin complexes in Boczonadi et al., Experimental Cell Research 313(16):3579-3591 (2007)). However these references do not discuss aberrant cell membrane distribution of plectin-1, or the detection of plectin-1 with a peptide or using a fluorescently labeled peptide probe (e.g., conjugated to nanoparticles) specific for plectin-1 to detect emerging PDACs.

III. The "Accessible" Proteome as a Source of Cancer Cell Biomarkers

[0139] As described herein, a phage display approach was used to screen for peptides that specifically bind to cell surface antigens on PDAC cells. These screens yielded a motif that distinguishes PDAC cells from normal pancreatic duct cells in vitro which, upon proteomic analysis, identified plectin-1 as a novel biomarker of PDAC. To assess their utility for in vivo imaging, the plectin-1 targeted peptides (PTP) were conjugated to magnetofluorescent nanoparticles. In conjunction with intravital confocal microscopy and MRI, these nanoparticles enabled detection of small PDAC and precursor lesions in engineered mouse models. The inventors developed a specific imaging probe based upon a peptide ligand, Clone 27, and discovered plectin-1 as a biomarker for pancreatic cancer cells contemplated for clinical utility in the diagnosis, management and treatment of PDAC in humans.

[0140] Phage display was used to identify peptides that distinguish mouse and human PDAC cells from normal pancreatic duct cells in vitro. The two peptides with the highest affinities and specificities were conjugated to magnetofluorescent nanoparticles (CLIO-VT680), and were demonstrated to effectively detect emerging tumors and pre-neoplastic lesions in a relevant transgenic mouse model via intravital confocal microscopy (Olympus IV100) and optical/MR imaging (OV-100, Bruker Pharmascan). Correlative histology confirmed the specific temporal localization of the PDAC targeted agents. Additionally, the peptide-binding partners identified from this approach represent a snapshot of the proteome in aberrant cells and also potential PDAC biomarkers. Using affinity chromatography, the binding partners for several peptides were identified, and their validity as biomarkers was demonstrated. These specific and sensitive probes are useful in the diagnosis and management of PDAC, e.g., in humans.

[0141] Methods for determining the sequence of identified binding molecules are also described herein. For example, when the binding molecules are produced in an expression library, encoding nucleic acids can be isolated from selected clones expressing binding molecules identified in a screen, such a ligand - target receptor binding assay. The encoding nucleic acids of the virions that bound to a target can then be sequenced using methods known to those skilled in the art.

[0142] Also described herein are methods for characterizing biomarkers that are selectively bound by a binding molecule, such as a peptide ligand. Once a binding partner (e.g., a peptide ligand) has been identified that is selective for a biomarker, the biomarker can then be isolated by, for example, affinity methods known in the art, and characterized. This characterization can be beneficial if the biomarkers used in the screen are not well characterized. Characterization of the biomarker includes such techniques as determining its apparent molecular weight by gel electrophoresis. Other methods applicable for characterizing biomarkers include, for example, high performance liquid chromatography (HPLC), mass spectrometry, or other methods that provide information about the physical, biochemical or functional properties of the biomarker, e.g., sequence and identity. Numerous methods are available for such characterization of biomarkers.

[0143] In preferred embodiments, biomarkers are derived from the cell surface of tumor cells. Cell surface molecules can be labeled, for example, with a detectable moiety such as a radioisotope or biotin or a fluorescence label. This labeling provides a source of biomarker where the only characteristic that needs to be known is that it is on the surface of a cell. As described in the Examples, cancer cell peptide ligands were prepared by labeling bacteriophage expressing cell surface peptides with FITC and used to identify binding molecules selective for the cancer cell surface polypeptides. Biomarker populations derived from other cell or subcellular compartments can similarly be used in the methods of the invention to obtain a binding partner peptide ligand that exhibits selective affinity for at least one biomarker within the initial population. Thus, the methods of the invention are applicable to a large variety of biomarker populations in which selective binding affinity is required for the therapeutic treatment or diagnosis of a disease.

[0144] The methods of the invention comprise selectively immobilizing a diverse population of binding molecules to a solid support. For selective immobilization, either an inherent characteristic of the binding molecules that comprise the population is exploited to provide selective immobilization or, alternatively, the molecules are engineered to contain a specific characteristic to be used for selective immobilization. For example, a binding partner itself may contain a hydrophobic chemical group or domain or may be fused to a hydrophobic chemical group or domain that causes the binding partner to be immobilized to a hydrophobic solid support such as plastic. In another example, the solid support can be coated with a chemical moiety or a biomolecule such that it is able to bind to and selectively immobilize only the binding molecules that make up the binding partner population. For example, the solid support can be coated with a biomolecule that selectively binds to a domain or sequence that is common to the binding molecules. The use of such biomolecules as linkers or tethers should be selected so that they do not interfere with the biomarker binding to the peptide ligands.

IV. Engineered Mouse Models of PDAC

[0145] There is a significant challenge in studying the early molecular changes in PDAC due to the typical presentation of PDAC at advanced stage and the corresponding lack of suitable tissue specimens. Therefore, the inventors elected to exploit a series of related genetically engineered mouse models of PDAC that harbor the signature gene mutations of the human disease, including Kras activation and deletion of the p53 or In k4a/Arf tumor suppressors (Bardeesy et al., Proc. Natl. Acad. Sci. U.S.A. 103:5947-5952 (2006); Aguirre et al., Genes Dev 17:3112-3126 (2003)). The tumors in these models exhibit the characteristic multistage histopathological progression (from precursor pancreatic intraepithelial neoplasia (PanINs) (Hansel et al., Annu Rev Genomics Hum Genet 4:237-256 (2003)) to metastatic cancer) that defines PDAC in humans, providing tractable model systems for both biological and preclinical studies (Bardeesy et al., (2006) *supra*). From these mouse models, the inventors have been able to generate primary cell lines derived from emerging PDAC. These early stage cancer cell lines, in conjunction with normal pancreatic ductal cells from wild-type mice (e.g., as described in Schreiber et al., Gastroenterology 127:250-260 (2004)), facilitate screening for biomarkers and imaging agents using combinatorial chemistry based approaches.

[0146] Genetically engineered mouse models of human cancers effectively recapitulate many of the molecular, biological, and clinical features of the human disease (Bardeesy et al., (2006) *supra*; Aguirre et al., (2003) *supra*).

[0147] Recent genomic studies of mouse and human cancers have established that cross-species *in vivo* analysis can serve as an effective filter in identifying recurrent chromosomal alterations associated with metastatic potential of melanoma and lymphoma-prone mice (Kim et al., Cell 125:1269-1281 (2006); Maser et al., Nature 447:966-971 (2007)). Using such an approach a 15-mer binding peptide ligand for identifying human prostate cancer cells was obtained by *in vivo* screening (Newton et al., Neoplasia 8:772-780 (2006)). Specifically, Newton describes identifying a phage clone displaying a 15 amino acid peptide, where the phage clone was labeled with the nearinfrared fluorophore (NIRF) AlexaFluor 680 (AF680), injected into a mouse at 10⁹ TU/ml (transducing/transfection units) of phage, and used for optical imaging of human prostate carcinoma in a mouse.

[0148] Studies described herein show the utility of mouse - human cancer models developed further for providing cancer cells at specific stages of development. Moreover phage clone screening methods were modified and extended to the development of assays for phage displaying smaller 7-mer peptides and use of such phage for sequencing and isolation of peptide ligands binding to molecules expressed by cancer cells at specific stages. These phage displaying 7-mer peptides and phage derived isolated 7-mer peptide ligands were used as molecularly targeted imaging agents while corresponding binding partners were identified and evaluated as biomarkers for pancreatic ductal adenocarcinoma.

[0149] Using these extended mouse models and phage displayed peptide binding screening methods, the inventors identified conserved biomarkers of early disease in screens that took advantage of mouse cell lines derived from early stages of cancer development and primary pancreatic ductal cells. Furthermore, the known kinetics of tumor progression of these mouse models facilitated testing of the imaging probes at defined stages of tumorigenesis. The approaches described herein are contemplated to be broadly applicable to the discovery of cancer biomarkers predictive of disease stage, prognosis, and the presence of specific genetic alterations associated with cancer cell development and prognosis.

V. Nanoparticle Based Imaging Labels and Diagnostic Methods for PDAC

[0150] Noninvasive imaging has particular applications in high risk groups, for example, hereditary PDAC kindreds and new-onset diabetes patients, who are candidates for screening for pancreatic cancer. Despite the increased cancer risk in these individuals, the incidence of actual pancreatic cancer is estimated to be only about 0.4%-0.6% (Chari, Semin Oncol 34: 284-294 (2007)), hence prophylactic surgery, which is associated with substantial morbidity and mortality, is not typically carried out. Traditional imaging methods such as CT scan or MRI often do not detect PDAC lesions until they have reached a size at which many tumors have already metastasized, thus rendering surgery ineffective. There is consequently a considerable need for a new imaging modality that would accurately identify the presence of PDAC at an earlier point in its development when surgery is effective.

[0151] Other settings for noninvasive imaging of incipient cancers include patients with cystic neoplasms, intraductal papillary mucinous neoplasms (IPMN) and mucinous cystic neoplasms (MCN). Thus in some embodiments, the inventors contemplate identifying peptide probes for biomarkers of incipient cancer cells. These tumors are often benign, however a subset of them progress to PDAC. Thus in some embodiments, the inventors contemplate methods for identifying and using peptide probes for biomarkers of neoplastic cells progressing to PDACs. New approaches using more accurate imaging molecules in postsurgical screening for recurrence and screening prior to surgery to determine exact tumor extension more accurately are contemplated. The new imaging molecules provided herein would be clinically valuable in differential diagnosis, i.e., patients presenting with pancreatitis, jaundice, or upper abdominal pain. Further, in screening of high-risk groups, the inventor contemplate an imaging molecule for distinguishing low-grade PanINs, which are present in many healthy individuals, from high-grade PanINs and carcinoma *in situ*. Probes are contemplated that would recognize lesions of PanIN-3 and higher since these are believed to have very high potential for progressing to invasive PDAC. Embodiments comprising translational studies are contemplated for conducting in patients undergoing resection because the rapid homing of the molecules described herein to tumors and subsequent clearance from the body makes this a technically feasible method.

[0152] In some embodiments, the peptide ligands described herein are linked to nanoparticle based imaging labels. These nanoparticles were developed and used for imaging cancer cells, such as the numerous types of magnetic nanoparticles and their magnetofluorescent analogues (see, e.g., Weissleder et al., *Nat. Biotechnol.*, 19:316-317 (2001); McCarthy et al., *Nanomedicine*, 2:153-167 (2007); Hogemann et al., *Bioconjug. Chem.*, 11:941-946 (2000), and Josephson et al., *Bioconjug. Chem.*, 10:186-191 (1999)) which are contemplated for use with isolated peptide ligands and phage displayed peptides. Multimodal nanoparticles are known that incorporate both magnetic and fluorescent molecules within the same molecule (e.g., as described herein, i.e., PTP-NP) and used for fluorescent microscopy (which detects the fluorescent part of this very small particle) and MRI (which detects its magnetic portion). In some embodiments, the imaging probes/ligands described herein include optical imaging probes such as NanoSPARKS™(VisEn Medical) and the like.

[0153] In some embodiments, the particles are conjugated to a fluorescent moiety, e.g., as described in U.S. Patent No. 5,492,814; Hogemann et al., *Bioconjug. Chem.*, 11:941-946 (2000).

[0154] The particles can be provided in any suitable form, e.g., lyophilized or in a liquid, e.g., a sterile carrier that is suitable for administration *in vivo*, e.g., sterile saline. Lyophilized particles can be reconstituted, e.g., in normal sterile saline, or in liquid carrier. In some embodiments, the methods use Combidex® (ferumoxtran-10), a molecular imaging agent consisting of iron oxide nanoparticles, available from Advanced Magnetics, Inc., Cambridge, MA.

[0155] The multimodal nanoparticle imaging probes described herein home to neoplasms while showing no appreciable colocalization with adjacent areas or acinar-ductal metaplasia. This high level of specificity is expected to reduce false positives in diagnostic tests. Further, these new imaging probes bound to PanINs as well as to advanced cancer cells. The capacity to detect such premalignant lesions enable the development of new approaches in the management of this disease. Although liver and kidney uptake is high, the tomographic imaging techniques that would be used with this probe (e.g., MRI, single photon emission computed tomography (SPECT)/CT, optical, etc.) would allow the resolution of the pancreas in the context of both organs, allowing differentiation therebetween.

[0156] The contrast agents are administered to the subject, e.g., by intravenous, intraarterial, subcutaneous, intramuscular, intraparenchymal, intracavity, topical, ocular, oral or rectal administration, with intravenous injection being preferred.

[0157] In addition to the development of novel molecularly targeted imaging agents, phage displayed peptide screening for bidding to molecules expressed by cancer cell and modified immunoprecipitation permitted the identification of membrane-localized plectin-1 as a new specific biomarker for PDAC. Significantly, differential protein processing and/or trafficking of plectin-1, identified using proteomic approaches, represents a potential class of biomarkers which is missed when merely evaluating cDNA expression data only or using whole-cell proteomics methods. In particular, although overexpression of plectin-1 was observed by gene chip analysis, it was one of 97 overexpressed genes and thus its relationship or use as a cancer cell biomarker was not known (see Iacobuzio-Donahue et al., *Am J Pathol.* 160:1239-1249 (2002); and United States Patent Application Publication Number 20030180747). The binding partners of additional clones identified herein from screening methods represent additional biomarkers contemplated for use in diagnostic methods and treatments in addition to shedding light on aberrant molecular pathways contributing to PDAC pathogenesis.

[0158] Thus the peptide ligands described herein can include one or more detectable moieties linked to a plectin-1 binding moiety that binds specifically to plectin-1, e.g., a peptide consisting essentially of SEQ ID NO: 1,2, or 4-8. The detectable moiety can be or include a fluorophore, e.g., a near infrared fluorophore (NIRF). A number of NIRFs useful in the methods and compositions described herein are known in the art, e.g., including Cy5.5, Cy5 and Cy7 (Amersham, Arlington Hts., IL; IRD41 and IRD700 (LI-COR, Lincoln, Nebr.); NIR-1, (Dejindo, Kumamoto, Japan); LaJolla Blue (Diatron, Miami, Fla.); indocyanine green (ICG) and its analogs (Licha et al., 1996, SPIE 2927:192-198; Ito et al., U.S. Pat. No. 5,968,479); indotricarbocyanine (ITC; WO 98/47538); and chelated lanthanide compounds. Fluorescent lanthanide metals include europium and terbium. Fluorescence properties of lanthanides are described in Lackowicz, 1999, *Principles*

of Fluorescence Spectroscopy, 2.sup.nd Ed., Kluwar Academic, New York. The fluorophores can be covalently linked to the plectin-1 binding moiety, or to a nanoparticle, e.g., via a fluorochrome attachment moiety, backbone, or spacer using any suitable reactive group on the fluorochrome and a compatible functional group on the fluorochrome attachment moiety, backbone, or spacer. For example, a carboxyl group (or activated ester) on a fluorochrome can be used to form an amide linkage with a primary amine such as the epsilon-amino group of the lysyl side chain on polylysine. Alternatively or in addition, the fluorophores can be linked directly to the backbone or linked to the backbone through nonbiodegradable spacers. See, e.g., US P.G.Pub. 20060275775.

[0159] The peptide ligands can be linked to the detectable moieties directly, e.g., as a fusion protein with protein or peptide detectable moieties (with or without an optional linking sequence, e.g., a flexible linker sequence) or via a chemical coupling moiety. A number of such coupling moieties are known in the art, e.g., a peptide linker or a chemical linker, e.g., as described in International Patent Application Publication No. WO 2009/036092.

VI. Ligands as a Targeting Moiety to Deliver Therapeutic Payloads

[0160] Ligand compositions, such as peptide ligands, used as therapeutics have advantages over other types of therapeutics, such as having a readily diffusible capability, low immunogenicity, and high specificity for target cells, in addition to flexibility in engineering novel additional elements, such as adding specific types of payloads, adding membrane permeabilizing factors, and the like. Thus, phage displaying peptide ligands and isolated peptide ligands as described herein are contemplated for use as a targeting moiety for selective delivery of therapeutic payloads, such as a radionuclide, cytokine, chemical drug, chemotherapy drug, and a therapeutic gene, to cancer cells. For the purposes of the present inventions, a "therapeutic payload" or "therapeutic cargo" includes a "therapeutic agent" and is intended to include any compound intended for extracellular or intracellular delivery to reduce the number of cancer cells or slow the growth of cancer cells or reduce the metastasis of cancer cells in a patient. Examples of types of payloads are drugs, small molecules, proteins, peptides, oligonucleotides, RNA and DNA, in other words any payload for reducing cancer in a patient.

[0161] A peptide ligand as described herein can be attached to a toxin, such as a diphtheria toxin (DTA) (for example, see, United States Patent No. 5827934 for DT fragments, and an exemplary fusion protein DAB_{3S9}EGF in Mishra, et al., 2003, Expert Opinion on Biological Therapy 3:1173-1180).

[0162] A drug attached to a peptide of the present invention can also include agents that are derived from, or that beneficially modulate host biological processes, such as interferons, tumor growth factors, tumor necrosis factors, growth factors such as GM-CSF and G-CSF and interleukins, for example, interleukin-2, interleukin-6, interleukin-7 and interleukin-12, and the like. A drug attached to a peptide of the present invention may comprise an agent which damages DNA and/or prevent cells from multiplying, such as genotoxins. A genotoxin includes but is not limited to alkylating agents, antimetabolites, DNA cutters, DNA binders, topoisomerase poisons and spindle poisons. Examples of alkylating agents are lomustine, carmustine, streptozocin, mechlorethamine, melphalan, uracil nitrogen mustard, chlorambucil, cyclophamide, iphosphamide, cisplatin, carboplatin, mitomycin, thiotepa, dacarbazine, procarbazine, hexamethyl melamine, triethylene melamine, busulfan, pipobroman, mitotane and other platine derivatives.

[0163] The peptide ligand as described herein can be used to deliver a variety of therapeutic agents, e.g., a cytotoxic moiety, e.g., a therapeutic drug, a radioisotope, molecules of plant, fungal, or bacterial origin, or biological proteins (e.g., protein toxins) or particles (e.g., a recombinant viral particles, e.g., via a viral coat protein), or mixtures thereof. The therapeutic agent can be an intracellularly active drug or other agent, such as short-range radiation emitters, including, for example, short-range, high-energy α -emitters, as described herein. In some embodiments, the peptide ligand can be coupled to a molecule of plant or bacterial origin (or derivative thereof), e.g., a maytansinoid (e.g., maytansinol or the DM1 maytansinoid). DM1 is a sulphydryl-containing derivative of maytansine that can be linked to the peptide, e.g., via a disulfide linker that releases DM1 when inside target cells. The disulfide linkers display greater stability in storage and in serum than other linkers. Maytansine is a cytotoxic agent that effects cell killing by preventing the formation of microtubules and depolymerization of extant microtubules. It is 100- to 1000-fold more cytotoxic than anticancer agents such as doxorubicin, methotrexate, and vinca alkyloid, which are currently in clinical use. Alternatively, the peptide ligand as described herein can be coupled to a taxane, a calicheamicin, a proteosome inhibitor, or a topoisomerase inhibitor. [(1R)-3-methyl-1-[(2S)-1-oxo-3-phenyl-2-[(3-mercaptopropyl) amino]propyl]amino]butyl] Boronic acid is a suitable proteosome inhibitor. N,N'-bis[2-(9-methylphenazine-1-carboxamido)ethyl]-1,2-ethanediamine is a suitable topoisomerase inhibitor.

[0164] Enzymatically active toxins and fragments thereof are exemplified by diphtheria toxin A fragment, nonbinding active fragments of diphtheria toxin, exotoxin A (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, α -sacrin, certain *Aleurites fordii* proteins, certain Dianthin proteins, *Phytolacca americana* proteins (PAP, PAPI and PAP-S), *Morodica charantia* inhibitor, curcin, crotin, *Saponaria officinalis* inhibitor, gelonin, mitogillin, restrictocin, phenomycin, and enomycin. In some embodiments, the peptide ligand is conjugated to maytansinoids, e.g., maytansinol (see US Patent No. 5,208,020), CC-1065 (see US Patent Nos. 5,475,092, 5,585,499, 5,846,545). Procedures for pre-

paring enzymatically active polypeptides of the immunotoxins are described in WO84/03508 and WO85/03508, which are hereby incorporated by reference. Examples of cytotoxic moieties that can be conjugated to the antibodies include adriamycin, chlorambucil, daunomycin, methotrexate, neocarzinostatin, and platinum.

[0165] To kill or ablate cancerous cells, a peptide ligand can be conjugated with a prodrug that is activated only when in close proximity with a prodrug activator. The prodrug activator is conjugated with a second peptide ligand, e.g., a second peptide ligand according to the present invention, preferably one that binds to a non-competing site on the same receptor (e.g., plectin-1) or cell. Whether two peptide ligand bind to competing or non-competing binding sites can be determined by conventional competitive binding assays. Drug-prodrug pairs suitable for use are known in the art, see, e.g., in Blakely et al., *Cancer Research* 56:3287-3292 (1996).

[0166] Alternatively, the peptide ligand can be coupled to high energy radiation emitters, for example, a radioisotope, such as ^{131}I , a γ -emitter, which, when localized at the tumor site, results in a killing of several cell diameters. See, e.g., Order, "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy", in *Monoclonal Antibodies for Cancer Detection and Therapy*, R.W. Baldwin et al. (eds.), pp 303-316 (Academic Press 1985). Other suitable radioisotopes include α -emitters, such as ^{212}Bi , ^{213}Bi , and ^{211}At , and β -emitters, such as ^{186}Re and ^{90}Y . Lu^{117} may also be used as both an imaging and cytotoxic agent.

[0167] Radioimmunotherapy (RIT) using peptide ligand labeled with ^{131}I , ^{90}Y , and ^{177}Lu is under intense clinical investigation. There are significant differences in the physical characteristics of these three nuclides and as a result, the choice of radionuclide can be important in order to deliver maximum radiation dose to the tumor. The higher beta energy particles of ^{90}Y may be good for bulky tumors, but it may not be necessary for small tumors and especially bone metastases, (e.g. those common to prostate cancer). The relatively low energy beta particles of ^{131}I are ideal, but *in vivo* dehalogenation of radioiodinated molecules is a major disadvantage for internalizing peptide ligands. In contrast, ^{177}Lu has low energy beta particle with only 0.2-0.3 mm range and delivers much lower radiation dose to bone marrow compared to ^{90}Y . In addition, due to longer physical half-life (compared to ^{90}Y), the tumor residence times are higher. As a result, higher activities (more mCi amounts) of ^{177}Lu labeled agents can be administered with comparatively less radiation dose to marrow. There have been several clinical studies investigating the use of ^{177}Lu labeled antibodies in the treatment of various cancers (see, e.g., Mulligan et al., *Clin Cancer Res.* 1: 1447-1454 (1995); Meredith et al., *J Nucl Med* 37:1491-1496 (1996); Alvarez et al., *Gynecologic Oncology* 65: 94-101 (1997)).

[0168] The peptide ligands of the invention can also be conjugated or fused to viral surface proteins present on viral particles. For example, a peptide ligand could be fused (e.g., to form a fusion protein) to a viral surface protein. Alternatively, a peptide ligand could be chemically conjugated (e.g., via a chemical linker) to a viral surface protein. Preferably, the virus is one that fuses with endocytic membranes, e.g., an influenza virus, such that the virus is internalized along with the peptide ligand and thereby cancer cells. The virus can be genetically engineered as a cellular toxin. For example, the virus could express or induce the expression of genes that are toxic to cells, e.g., cell death promoting genes. Preferably, such viruses would be incapable of viral replication.

[0169] The peptide ligands can be linked to the therapeutic agent directly, e.g., as a fusion protein with protein or peptide toxins (with or without an optional linking sequence, e.g., a flexible linker sequence) or via a chemical coupling moiety. A number of such coupling moieties are known in the art, e.g., a peptide linker or a chemical linker, e.g., as described in International Patent Application Publication No. WO 2009/036092.

[0170] The peptide ligands (e.g., linked to an imaging moiety or a therapeutic agent) described herein can be incorporated into pharmaceutical compositions. Such compositions typically include the compound (i.e., as an active agent) and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carriers" includes saline, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration.

[0171] Pharmaceutical compositions are typically formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide.

[0172] Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, e.g., gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a

disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[0173] Systemic administration of a therapeutic compound as described herein can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[0174] For administration by inhalation, the compounds are typically delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer. Such methods include those described in U.S. Patent No. 6,468,798.

[0175] The therapeutic compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[0176] In one embodiment, the therapeutic compounds are prepared with carriers that will protect the therapeutic compounds against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Such formulations can be prepared using standard techniques, or obtained commercially, e.g., from Alza Corporation and Nova Pharmaceuticals, Inc.

[0177] Liposomal suspensions (including liposomes targeted to selected cells with monoclonal antibodies to cellular antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[0178] Dosage, toxicity and therapeutic efficacy of the therapeutic compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit high therapeutic indices are preferred.

[0179] The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0180] Concentration ranges of a drug in vitro in which peptide ligand targeting may enhance the ability of a drug to selectively kill cancer cells depend, usually, on the drug used. For example, genotoxin is used usually at a concentration in vitro between 0.1 to 100, μ M, preferably between 0. 15 to 30 μ M/kg.

[0181] Therapeutic compositions of peptide ligands that deliver drugs to cancer cells may increase the sensitivity of cancer cells to additional anti-cancer treatments, for example, peptide ligands conjugated to paclitaxel, in the range of 150 mg/m² of body surface, increases cancer cell sensitivity to certain drugs, while cisplatin in the range of 20 mg/m²/day increases cancer cell sensitivity to radiation.

[0182] Peptide ligands that bind to the external surface of cells may comprise a "membrane fusion component" intended to include a domain or molecule that facilitates transport of a payload into a cell. The membrane fusion component may contain a membrane permeant motif. The membrane fusion component can be isolated from a naturally occurring protein, or may be a synthetic molecule based in whole or in part on a naturally occurring domain, for example, human immunodeficiency virus type 1 (HIV-1) glycoprotein-120 (GP120), human immunodeficiency virus type 1 (HIV-1) glycoprotein-42 (GP41), human immunodeficiency virus (HIV-1) (transactivator of transcription (Tat) protein), human parainfluenza virus, hemagglutinin (HA) of influenza virus (termed HA2), Ebola virus transmembrane fusion sequence, helical coil-coils, alpha-hemolysin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), Melittin (active component of bee ven-

om), a hydrophobic segment, a synthetic membrane transporter, and the like. Suitable membrane fusion components are known in the art, e.g., HIV-derived TAT peptide, penetratins, transportans, or hCT derived cell-penetrating peptides, see, e.g., Caron et al., (2001) Mol Ther. 3(3):310-8; Langel, Cell-Penetrating Peptides: Processes and Applications (CRC Press, Boca Raton FL 2002); El-Andaloussi et al., (2005) Curr Pharm Des. 11(28):3597-611; and Deshayes et al., (2005) Cell Mol Life Sci. 62(16):1839-49.

5 [0183] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

VIII. Antibodies

10 [0184] The present invention provides isolated antibodies (e.g., polyclonal or monoclonal) having affinity for a cancer biomarker. In some embodiments, the cancer comprises pancreatic cancer. In other embodiments, the cancer includes, but is not limited to, lung cancer, breast cancer, prostate cancer, skin cancer, brain cancer, liver cancer, bone cancer, or kidney cancer. In some embodiments, the present invention provides monoclonal antibodies that specifically bind to 15 an isolated polypeptide comprised of at least five amino acid residues of a plectin biomarker (e.g., SEQ ID NO:1). These antibodies find use in the imaging methods described herein.

20 [0185] An antibody against a protein of the present invention may be any monoclonal or polyclonal antibody, as long as it can recognize the protein. Antibodies can be produced by using a protein of the present invention as the antigen according to a conventional antibody or antiserum preparation process.

25 [0186] The present invention contemplates the use of both monoclonal and polyclonal antibodies. Any suitable method may be used to generate the antibodies used in the methods and compositions of the present invention, including but not limited to, those disclosed herein. For example, for preparation of a monoclonal antibody, protein, as such, or together with a suitable carrier or diluent is administered to an animal (e.g., a mammal) under conditions that permit the production of antibodies. For enhancing the antibody production capability, complete or incomplete Freund's adjuvant may be 30 administered. Normally, the protein is administered once every 2 weeks to 6 weeks, in total, about 2 times to about 10 times. Animals suitable for use in such methods include, but are not limited to, primates, rabbits, dogs, guinea pigs, mice, rats, sheep, goats, etc.

35 [0187] For preparing monoclonal antibody-producing cells, an individual animal whose antibody titer has been confirmed (e.g., a mouse) is selected, and 2 days to 5 days after the final immunization, its spleen or lymph node is harvested and antibody-producing cells contained therein are fused with myeloma cells to prepare the desired monoclonal antibody producer hybridoma. Measurement of the antibody titer in antiserum can be carried out, for example, by reacting the labeled protein, as described hereinafter and antiserum and then measuring the activity of the labeling agent bound to 40 the antibody. The cell fusion can be carried out according to known methods, for example, the method described by Kohler and Milstein (Nature 256:495 (1975)). As a fusion promoter, for example, polyethylene glycol (PEG) or Sendai virus (HVJ), preferably PEG is used.

45 [0188] Examples of myeloma cells include NS-1, P3U1, SP2/0, AP-1 and the like. The proportion of the number of antibody producer cells (spleen cells) and the number of myeloma cells to be used is preferably about 1:1 to about 20:1. PEG (preferably PEG 1000-PEG 6000) is preferably added in concentration of about 10% to about 80%. Cell fusion can be carried out efficiently by incubating a mixture of both cells at about 20°C to about 40°C, preferably about 30°C to 50 about 37°C for about 1 minute to 10 minutes.

55 [0189] Various methods may be used for screening for a hybridoma producing the antibody (e.g., against a tumor antigen or autoantibody of the present invention). For example, where a supernatant of the hybridoma is added to a solid phase (e.g., microplate) to which antibody is adsorbed directly or together with a carrier and then an anti-immunoglobulin antibody (if mouse cells are used in cell fusion, anti-mouse immunoglobulin antibody is used) or Protein A labeled with a radioactive substance or an enzyme is added to detect the monoclonal antibody against the protein bound to the solid phase. Alternately, a supernatant of the hybridoma is added to a solid phase to which an anti-immunoglobulin antibody or Protein A is adsorbed and then the protein labeled with a radioactive substance or an enzyme is added to detect the monoclonal antibody against the protein bound to the solid phase.

50 [0190] Selection of the monoclonal antibody can be carried out according to any known method or its modification. Normally, a medium for animal cells to which HAT (hypoxanthine, aminopterin, thymidine) are added is employed. Any selection and growth medium can be employed as long as the hybridoma can grow. For example, RPMI 1640 medium containing 1% to 20%, preferably 10% to 20% fetal bovine serum, GIT medium containing 1% to 10% fetal bovine serum, a serum free medium for cultivation of a hybridoma (SFM-101, Nissui Seiyaku) and the like can be used. Normally, the cultivation is carried out at 20°C to 40°C., preferably 37°C for about 5 days to 3 weeks, preferably 1 week to 2 weeks under about 5% CO₂ gas. The antibody titer of the supernatant of a hybridoma culture can be measured according to the same manner as described above with respect to the antibody titer of the anti-protein in the antiserum.

VIII. Designing Mimetics

[0191] Compounds mimicking the necessary conformation for recognition and binding to a biomarker, such as plectin-1, are contemplated as within the scope of this invention. For example, In some embodiments, mimetics of SEQ ID NOS: 1-8 and all of the peptides of the present invention are contemplated. A variety of designs for such mimetics are possible. United States Patent No. 5,192,746 to Lobl, et al., United States Patent No. 5,169,862 to Burke, Jr., et al., United States Patent No. 5,539,085 to Bischoff, et al., United States Patent No. 5,576,423 to Aversa, et al., United States Patent No. 5,051,448 to Shashoua, and United States Patent No. 5,559,103 to Gaeta, et al., all hereby incorporated by reference, describe multiple methods for creating such compounds.

[0192] Synthesis of nonpeptide compounds that mimic peptide sequences is also known in the art. Eldred, et al, (J. Med. Chem. 37:3882 (1994)) describe nonpeptide antagonists that mimic the Arg-Gly-Asp sequence. Likewise, Ku, et al, (J. Med. Chem. 38:9 (1995)) give further elucidation of the synthesis of a series of such compounds. Such nonpeptide compounds that mimic, for example, KTLLPTP [SEQ ID NO: 1] peptides (or, of any one or more of the polypeptides of the present invention) are specifically contemplated by the present invention.

[0193] The present invention also contemplates synthetic mimicking compounds that are multimeric compounds that repeat relevant peptide sequences. As is known in the art, peptides can be synthesized by linking an amino group to a carboxyl group that has been activated by reaction with a coupling agent, such as dicyclohexylcarbodiimide (DCC). The attack of a free amino group on the activated carboxyl leads to the formation of a peptide bond and the release of dicyclohexylurea. It can be necessary to protect potentially reactive groups other than the amino and carboxyl groups intended to react. For example, the α -amino group of the component containing the activated carboxyl group can be blocked with a tertbutyloxycarbonyl group. This protecting group can be subsequently removed by exposing the peptide to dilute acid, which leaves peptide bonds intact.

[0194] With this method, peptides can be readily synthesized by a solid phase method by adding amino acids stepwise to a growing peptide chain that is linked to an insoluble matrix, such as polystyrene beads. The carboxyl-terminal amino acid (with an amino protecting group) of the desired peptide sequence is first anchored to the polystyrene beads. The protecting group of the amino acid is then removed. The next amino acid (with the protecting group) is added with the coupling agent. This is followed by a washing cycle. The cycle is repeated as necessary.

[0195] In some embodiments, the mimetics of the present invention are peptides having sequence homology to the above-described plectin protein ligands. One common methodology for evaluating sequence homology, and more importantly statistically significant similarities, is to use a Monte Carlo analysis using an algorithm written by Lipman and Pearson to obtain a Z value. According to this analysis, a Z value greater than 6 indicates probable significance, and a Z value greater than 10 is considered to be statistically significant. See, e.g., Pearson and Lipman, Proc. Natl. Acad. Sci. (USA) 85:2444-2448 (1988); Lipman and Pearson, Science 227:1435-1441 (1985). In the present invention, synthetic polypeptides useful in tumor therapy and in blocking invasion are those peptides with statistically significant sequence homology and similarity (Z value of Lipman and Pearson algorithm in Monte Carlo analysis exceeding 6).

[0196] The present invention also contemplates peptide sequence derivatives of SEQ ID NOS: 1-8 (or any one or more of the polypeptides sequences of the present invention) identified by means of an amino acid pairing technique. See, e.g., Root-Bernstein, J. Theor. Biol. 94:885-859 (1982); and Stefanowicz et al., Letters in Peptide Science 5:329-331 (1998). To identify peptide sequence derivates useful in the present invention, the methodology is adapted to identify sequences that are in some wise complementary to seprase, but that specifically inhibit seprase activity without, at the same time, themselves potentiating the migration of cells.

[0197] The peptide ligands described herein can be protease resistant and can include one or more types of protecting groups such as an acyl group, an amide group, a benzyl or benzoyl group, or a polyethylene glycol. More specifically, a peptide, including the modified peptides described above, can be N-terminally acetylated and/or C-terminally amidated.

[0198] Where non-naturally occurring or modified amino acid residues are included they can be selected from the following or many others available in the art: 4-hydroxyproline, gamma-carboxyglutamic acid, o-phosphoserine, o-phosphotyrosine, or delta-hydroxylysine. Other examples include naphthylalanine, which can be substituted for tryptophan to facilitate synthesis, L-hydroxypropyl, L-3,4-dihydroxyphenylalanine, alpha-amino acids such as L-alpha-hydroxylysyl and D-alpha-methylalanine, L-alpha-methylalanine, beta-amino acids, and isoquinolyl. Peptides having non-naturally occurring amino acid residues may be referred to as synthetic peptides and constitute one type of variant as described herein. Other variants include peptides in which a naturally occurring side chain of an amino acid residue (in either the L- or D-form) is replaced with a non-naturally occurring side chain.

[0199] In some embodiments, the peptides can have three extra amino acids (Met-Gly-Ser) at either terminus (or both) (e.g., at the N-terminus) and seven to eight extra amino acids (e.g., Thr-Ser-His-His-His-His-His-Cys (SEQ ID NO:26)) at either terminus (or both) (e.g., at the C-terminus).

[0200] In some embodiments, the peptides can be PEGylated by methods known in the art.

[0201] For guidance on peptide modification by reduction/alkylation and/or acylation, one can consult Tarr, Methods of Protein Microcharacterization, Silver ed., Humana Press, Clifton N.J. (1986) 155-194; for guidance on chemical

coupling to an appropriate carrier, one can consult Mishell and Shiigi, eds, Selected Methods in Cellular Immunology, WH Freeman, San Francisco, Calif. (1980) and U.S. Pat. No. 4,939,239; and for guidance on mild formalin treatment, one can consult Marsh, Int. Arch. Allergy Appl. Immunol., (1971) 41:199-215.

[0202] Peptidomimetics of the peptide ligands can also be used. Peptide ligands disclosed herein can be modified according to methods known in the art for producing peptidomimetics. See, e.g., Kazmierski, W.M., ed., Peptidomimetics Protocols, Human Press (Totowa NJ 1998); Goodman et al., eds., Houben-Weyl Methods of Organic Chemistry: Synthesis of Peptides and Peptidomimetics, Thiele Verlag (New York 2003); and Mayo et al., J. Biol. Chem., 278:45746, (2003). In some cases, these modified peptidomimetic versions of the peptides and fragments disclosed herein exhibit enhanced stability *in vivo*, relative to the non-peptidomimetic peptides. Methods for creating a peptidomimetic include substituting one or more, e.g., all, of the amino acids in a peptide sequence with D-amino acid enantiomers. Such sequences are referred to herein as "retro" sequences. In another method, the N-terminal to C-terminal order of the amino acid residues is reversed, such that the order of amino acid residues from the N-terminus to the C-terminus of the original peptide becomes the order of amino acid residues from the C-terminus to the N-terminus in the modified peptidomimetic. Such sequences can be referred to as "inverso" sequences. Peptidomimetics can be both the retro and inverso versions, i.e., the "retro-inverso" version of a peptide disclosed herein. The new peptidomimetics can be composed of D-amino acids arranged so that the order of amino acid residues from the N-terminus to the C-terminus in the peptidomimetic corresponds to the order of amino acid residues from the C-terminus to the N-terminus in the original peptide.

[0203] Other methods for making a peptidomimetics include replacing one or more amino acid residues in a peptide with a chemically distinct but recognized functional analog of the amino acid, i.e., an artificial amino acid analog. Artificial amino acid analogs include β -amino acids, β -substituted β -amino acids (" β^3 -amino acids"), phosphorous analogs of amino acids, such as α -amino phosphonic acids and α -amino phosphinic acids, and amino acids having non-peptide linkages. Artificial amino acids can be used to create peptidomimetics, such as peptoid oligomers (e.g., peptoid amide or ester analogues), β -peptides, cyclic peptides, oligourea or oligocarbamate peptides; or heterocyclic ring molecules.

These sequences can be modified, e.g., by biotinylation of the amino terminus and amidation of the carboxy terminus.

[0204] Any of the peptides described herein, including the variant forms described herein, can further include a heterologous polypeptide. The heterologous polypeptide can be a polypeptide that increases the circulating half-life of the peptide to which it is attached (e.g., fused, as in a fusion protein). The heterologous polypeptide can be an albumin (e.g., a human serum albumin or a portion thereof) or a portion of an immunoglobulin (e.g., the Fc region of an IgG). The heterologous polypeptide can be a mitochondrial-penetrating moiety.

[0205] Compounds mimicking the necessary conformation of the peptides described herein are contemplated as within the scope of this invention. A variety of designs for such mimetics are possible. U.S. Patent No. 5,192,746; U.S. Patent No. 5,169,862; U.S. Patent No. 5,539,085; U.S. Patent No. 5,576,423; U.S. Patent No. 5,051,448; and U.S. Patent No. 5,559,103, all hereby incorporated by reference, describe multiple methods for creating such compounds. Non-peptidic compounds that mimic peptide sequences are known in the art (see, e.g., Meli et al. J. Med. Chem., 49:7721-7730 (2006), describing methods of identifying nonpeptide small molecule mimics of shepherin). Synthesis of non-peptide compounds that mimic peptide sequences is also known in the art (see, e.g., Eldred et al. J. Med. Chem., 37:3882, (1994); Ku et al. J. Med. Chem., 38:9, (1995); Meli et al. J. Med. Chem., 49:7721-7730 (2006)). Such nonpeptide compounds that mimic the sequences described herein that bind plectin-1 are specifically contemplated by the present invention.

[0206] The present invention also contemplates synthetic mimicking compounds. As is known in the art, peptides can be synthesized by linking an amino group to a carboxyl group that has been activated by reaction with a coupling agent, such as dicyclohexylcarbodiimide (DCC). The attack of a free amino group on the activated carboxyl leads to the formation of a peptide bond and the release of dicyclohexylurea. It can be necessary to protect potentially reactive groups other than the amino and carboxyl groups intended to react. For example, the (α -amino group of the component containing the activated carboxyl group can be blocked with a tertbutyloxycarbonyl group. This protecting group can be subsequently removed by exposing the peptide to dilute acid, which leaves peptide bonds intact. With this method, peptides can be readily synthesized by a solid phase method by adding amino acids stepwise to a growing peptide chain that is linked to an insoluble matrix, such as polystyrene beads. The carboxyl-terminal amino acid (with an amino protecting group) of the desired peptide sequence is first anchored to the polystyrene beads. The protecting group of the amino acid is then removed. The next amino acid (with the protecting group) is added with the coupling agent. This is followed by a washing cycle. The cycle is repeated as necessary.

[0207] In some embodiments, the mimetics of the present invention are peptides having sequence homology to the herein-described chaperone inhibitor peptides. These mimetics include, but are not limited to, peptides in which L-amino acids are replaced by their D-isomers. One common methodology for evaluating sequence homology, and more importantly statistically significant similarities, is to use a Monte Carlo analysis using an algorithm written by Lipman and Pearson to obtain a Z value. According to this analysis, a Z value greater than 6 indicates probable significance, and a Z value greater than 10 is considered to be statistically significant (Pearson and Lipman, Proc. Natl. Acad. Sci. (USA),

85:2444-2448, (1988); Lipman and Pearson, *Science*, 227:1435-1441, (1985). More generally, the peptide ligands described herein and the mimetics described above can be synthesized using any known methods, including tea-bag methodology or solid phase peptide synthesis procedures described by Merrifield et al., *Biochemistry*, 21:5020-5031, (1982); Houghten Wellings, *Proc. Natl. Acad. Sci. (USA)*, 82:5131-5135, (1985); Atherton, *Methods in Enzymology*, 289:44-66, (1997), or Guy and Fields, *Methods in Enzymology*, 289:67-83, (1997), or using a commercially available automated synthesizer.

IX. Small Molecule Drugs

[0208] In some embodiments, the present invention provides drugs (e.g., small molecule drugs) that reduce or eliminate cancer by binding to a cancer biomarker (e.g., plectin). In some embodiments, small molecule drugs are identified using the drug screening methods described herein. In preferred embodiments, the small molecule drugs of the present invention result in the death of cancer, but not normal cells. In some embodiments, small molecule drugs are identified using the drug screens described herein (e.g., in Section III above).

[0209] In some embodiments, the present invention provides drug screening assays (e.g., to screen for anticancer drugs). The present invention is not limited to a particular mechanism. Indeed, and understanding of the mechanism is not necessary to practice the present invention. The present invention provides drug screening methods for identifying compounds that bind to a cancer biomarker expressed on a cell surface membrane (e.g., a tumor tissue). The present invention further provides methods of identifying chemotherapeutic agents that are active in plectin-1 expressing cancers.

In some embodiments, candidate compounds such as small molecules are directed against plectin-1.

[0210] In some embodiments, the invention contemplates a method for identifying an effective nonpeptide small-molecule inhibitor that blocks/inhibits/prevents/disrupts a cancer biomarker (e.g., plectin). These molecules may be discovered using any one of several high-throughput screening methods. See, e.g., Stockwell, *Nature* 432:846-854 (2004); Kay et al., *Mol. Diversity* 1:139-140 (1996); Pfleger et al., *Cell Signaling* 18:1664-1670 (2006); Jung et al., *Proteomics* 5: 4427-4431 (2005); Nieuwenhuijsen et al., *J. Biomol. Screen* 8:676-684 (2003); and Berg, *Angew. Chem. Int. Ed. Engl.* 42:2462-2481 (2003).

EXAMPLES

[0211] The following examples serve to illustrate certain embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

[0212] In the experimental disclosures which follow, the following abbreviations apply: N (normal); M (molar); mM (millimolar); μ M (micromolar); mol (moles); mmol (millimoles); μ mol (micromoles); nmol (nanomoles); pmol (picomoles); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); pg (picograms); L and I (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); U (units); min (minute); s and sec (second); k (kilometer); deg (degree); °C (degrees Centigrade/Celsius), colony-forming units (cfu), plaque forming units (PFU), optical density (OD; o.d.), internal diameter (i.d.), and polymerase chain reaction (PCR).

EXAMPLE I

[0213] This example describes exemplary materials and methods for assays used during the development of the present inventions.

Cell Culture

[0214] Primary mouse pancreatic ductal cells from wildtype mice were isolated and cultivated using published methods (Schreiber, et al., (2004) *Gastroenterology* 127:250-260). Early passage PDAC cell lines were isolated from tumors arising in Pdx1-Cre LSL-KrasG12D p53^{L/L} mice (designated Kras/p53^{L/L}) (Bardeesy, et al. (2006) *Proc. Natl. Acad. Sci. U.S.A.* 103:5947-5952). For the phage display experiments, PDAC cells were first grown in the primary duct cell media (F12 medium supplemented with 5 mg/mL D-glucose (Sigma), 0.1 mg/mL soybean trypsin inhibitor type I (Sigma), 5 mL/L insulin-transferrin-selenium (ITS+; BD Biosciences, Palo Alto, CA), 25 μ g/mL bovine pituitary extract (BD Biosciences), 20 ng/mL epidermal growth factor (BD Biosciences), 5 nmol/L 3,3',5-triiodo-L-thyronine (Sigma), 1 μ mol/L dexamethasone (Sigma), 100 ng/mL cholera toxin (Sigma), 10 mmol/L nicotinamide (Sigma), 5% Nu-serum IV culture supplement (Collaborative Biomedical Products), and antibiotics (penicillin G 100 U/mL, streptomycin 100 μ g/mL, amphotericin B 0.25 μ g/mL; Gibco-BRL, Grand Island, N.Y.). Human PDAC cell lines (MNA, 8988, SW1990, MIA-PaCa-2, ASPC) were purchased from ATCC and cultured according to established protocols. NIH-3T3 cells (mouse fibroblasts) are purchased from ATCC. Murine heart endothelial cells (MHEC) were isolated from mice according to previously published protocols (Allport et al., *J Leukoc Biol* 71:821-828 (2002)) and used after the second subculture. Human

umbilical vein endothelial cells (HUVECs) were purchased from Clonetics and cultured according to the manufacturer's protocol.

Mouse cohorts

[0215] Imaging studies were performed in Pdxl-Cre LSL-KrasG12Dp53^{L/+} (Kras/p53^{L/+}), Pdxl-Cre LSL-KrasG12D p16^{+/−} (Kras/p16^{+/−}), Pdxl-Cre LSL-KrasG12D (Kras), and wild type mice (Bardeesy et al., Proc. Natl. Acad. Sci. U.S.A. 103:5947-5952 (2006)). Breeding, genotyping and analysis were performed as previously published (Bardeesy, et al., Proc. Natl. Acad. Sci. U.S.A. 103:5947-5952 (2006); and Aguirre et al., Genes Dev 17:3112-3126 (2003)). Mice were housed in a pathogen-free environment at the Massachusetts General Hospital (MGH). The mice were handled in strict accord with good animal practice as defined by the Office of Laboratory Animal Welfare, and all animal work was done with Institutional Animal Care and Use Committee approval.

Phage Selection

[0216] Phage positive selection and negative selection were achieved by incubating mouse PDAC cells isolated from the Kras/p53 mouse with phage (1x10¹¹ PFU), which displayed a randomized linear 7 amino acid peptide library (phD7, New England Biolabs, Beverly, MA) for 1 hour at 37° C to allow time for phage to be internalized into the PDAC cells. Screening for cell-internalizing phage affords a type of signal amplification by concentrating the imaging agent inside the cell with the additional benefit that the agent is not subject to k_{off} (off rate), further increasing the effective affinity (Kelly et al., Circ Res 96: 327-336 (2005)). To remove unbound phage and non-specific binding phage, the cells were first washed with DPBS supplemented with 1% BSA and 0.05% Tween-20. Cell surface bound phage were removed by washing with 0.1 M glycine (pH 2) for 8 minutes. Following a second glycine wash, the internalized phage were recovered by lysing the cells with 0.1% triethanolamine (Sigma, St. Louis, MO) in PBS (pH 7.6) for 5 min at room temperature. The internalized phage pool was neutralized with 100 μ L of 0.5 M Tris-HCl (pH 7). The counter-selection was done by incubating the internalized phage pool with normal pancreatic cells for three 30 minute cycles to effectively subtract all clones that bind to both normal pancreatic ductal cells and PDAC (Kelly et al., Neoplasia 5: 437-444 (2003)). The internalized phage were amplified in Escherichia coli, titered, and subjected to three additional rounds for a total of 4 rounds of positive selection on the PDAC cells. From this selection, 30 clones were selected for sequencing and analyzed by ELISA (see below).

Enzyme-Linked ImmunoSorbent Assay (ELISA) and Multidimensional Analysis

[0217] ELISA and multidimensional analysis were used to facilitate choosing appropriate clones for further study (Kelly et al., Mol Imaging 5:24-30 (2006)). Specifically, Pancreatic ductal adenocarcinoma (PDAC) and normal cells (noncancerous) were grown to 100% confluence in a 96-well plate and incubated sequentially at 37° C with 30 phage clones (10⁷ and 10¹⁰ PFU, 1 h) in triplicate, washed with PBS containing 0.1% Tween-20, incubated with biotinylated anti-M13 antibody (1:40, 1 h), detected with streptavidin- Horseradish peroxidase (HRP) (1:500), developed with tetra-methylbenzidine and absorbance at 650 was determined (Emax, Molecular Devices).

[0218] Raw plate-reader outputs corresponding to PDAC or normal ductal cells were unpivoted to afford a denormalized table of values, and each well position was then associated with similar arrays of metadata labels. Values from each well were background subtracted using the median value of mock-treatment wells (wild-type phage) from each assay plate. Background-subtracted (Bsub) values for mock-treatment wells were accumulated across multiple assay plates to afford two mocktreatment distributions reflecting assay noise, one corresponding to PDAC cells and one corresponding to normal ductal cells, and trimmed according to Chauvenet's criterion as previously described (Kelly et al., Circ Res 96: 327-336 (2005)). These mock-treatment distributions were used to normalize independently each value corresponding to a phage-treated well, affording Z-normalized (Znorm) values for each well. All data formatting, manipulation, and normalization were implemented using Pipeline Pilot (Scitegic) and data visualizations (heat map) were prepared using DecisionSite (Spotfire).

50

Phage Labeling

[0219] For in vitro and in vivo validation experiments, phage were fluorochrome-labeled as previously described (Kelly et al., Neoplasia 8:1011-1018 (2006)). Briefly, approximately 1 x 10¹² PFU of phage was suspended in 100 μ L of 0.3 M NaHCO₃ (pH 8.6) then depending upon the experiment the NaHCO₃ solution contained one of the following dyes: 1 mg/mL of fluorochrome-hydrosuccinimide ester (conjugated to either Cy5.5 or AF750), 0.25 mg/mL of FITC, 0.25 mg/mL of RITC (rhodamine isothiocyanate). The labeling reaction was allowed to continue in the dark at room temperature (RT) with gentle agitation. After 1 hour, the reaction mixture was diluted to 1 mL, in DPBS and the labeled-phage was purified

by PEG precipitation (3x). The fluorochrome-labeled phage was resuspended in 200 μ L DPBS. Plaque-forming units were determined by titer analysis and the concentration of the fluorochrome was determined spectrophotometrically (Varian Cary 11, Varian, Palo Alto, C.A.).

5 **Phage Detection by Fluorescent Microscopy and Flow Cytometry**

[0220] Mouse Pancreatic ductal adenocarcinoma (PDAC) cells, human PDAC cells (i.e. MNA, 8988, SW1990, PaCa-2, ASPC), normal human ductal cells and normal pancreatic cells were incubated with 1 mM (FITC) FITC-labeled phage clone 27 or unrelated phage clone (amino acid sequence SNLHPSD, negative control (SEQ ID NO:XX)) for 1 hour at 10 37° C, washed 3X with DPBS and visualized by fluorescent microscopy (Nikon Eclipse TE2000-S, Insight QE, 40X objective). The cells were then harvested by incubation with trypsin, centrifuged, and analyzed (10,000 cells/sample) by flow cytometry on a Beckton Dickinson FACSCalibur (San Jose, CA). Samples of PDAC cells showed a single narrow peak of fluorescent intensity that was higher than from normal cells (e.g., Figure 1B). Mean fluorescence was plotted 15 against cell number to show relative uptake between cell types.

15 **Ex vivo Biopsy Specimens**

[0221] Pancreatic ductal adenocarcinoma (PDAC) -specific peptides identified by phage studies were tested for binding 20 on histology sections. In particular, ex vivo mouse and human tissue sections from biopsy specimens were snap frozen, embedded in OCT, cut into 5 μ m sections, and then attached to slides. Slides with tissues were incubated with 1 μ M of FITC labeled Phage Clone 27 or FITC labeled control phage (no insert) for 1 hour at 37° C, washed 3x with PBS, fixed with 2% paraformaldehyde, and then visualized by fluorescence microscopy (Nikon Eclipse TE2000-S, Insight QE, 40X objective).

25 **Identification of a binding partner for a phage expressed peptide**

[0222] Phage were labeled with a photo linker (Sulfo-SAED (Sulfosuccinimidyl 2-[7-amino-4-methylcoumarin-3-acetamido]ethyl-1,3 dithiopropionate; Pierce, Waltham, MA) and biotin tag using the same NHS chemistry used to conjugate 30 fluorochromes to phage (Kelly et al., Neoplasia 8: 1011-1018 (2006)). Two Petri dishes (10 cm, Fisher scientific, Waltham, MA) were plated with the target cell line and grown to until confluence where the cells covered the plate. One confluent plate was incubated with 1 mL of the modified phage (roughly 10¹⁰ PFU/uL). For a negative control, the second plate 35 was incubated with control (no insert) phage. Both plates were incubated in the dark at 4°C for one hour. The cells were then again washed several times with DPBS, placed on ice, and photolyzed 30 min using a 15 watt 365 nm UV lamp (Spectroline, Westbury, NY), and lysed using 1% triton x-100 in PBS with mammalian protease inhibitor cocktail added (Sigma, St. Louis, MO). The cell lysates were incubated 1 hr with 100 μ L of Dynal Streptavidin beads (Invitrogen, Carlsbad, CA) which were pre-blocked with 5% BSA in PBS. The beads were washed twice with 1% triton x-100 in 10x PBS, then 40 incubated overnight at 4°C with a buffer containing DTT to reverse the chemical crosslink and release the precipitated protein. Half of the eluate was transferred to PVDF membrane and probed with plectin-1 antibody (Santa Cruz Biotechnology, Santa Cruz, CA). The other half of the eluate was loaded onto a SDS/PAGE gel (Biorad Criterion system, Hercules, CA) and stained using a mass spectroscopic compatible silver stain (Invitrogen, Carlsbad, CA).

[0223] The silver stained band was then excised and sent for tryptic digest/mass spec analysis (Tufts Peptide Core Facility). Nanobore electrospray columns were constructed from 360 mm o.d., 75 mm i.d. fused silica capillary with the 45 column tip tapered to a 15 mm opening. The columns were packed with 200 Å 5 um C18 beads (Michrom Bioresearches, Auburn, CA.), a reverse-phase packing material, to a length of 10 cm. The flow through the column was split precolumn to achieve a flow rate of 350 nL/min. The mobile phase used for gradient elution consisted of (A) 0.3% acetic acid 99.7 % water and (B) 0.3% acetic acid 99.7 % Acetonitrile. Tandem mass spectra were acquired on a Thermo LTQ ion trap 50 mass spectrometer (Thermo Corp., San Jose, CA). Needle voltage was set at 3 kV. Ion signals above a predetermined threshold automatically triggered the instrument to switch from mass spectrometry (MS) to tandem mass spectrometry (MS/MS) mode for generating fragmentation spectra. The MS/MS spectra were searched against the NCBI nonredundant protein sequence database using the SEQUEST computer algorithm (Yates et al., Anal Chem 67: 1426-1436 (1995)).

Verification of Clone 27 (Panc 27) binding to plectin-1

[0224] Subcellular fractionation: PDAC, PaCa-2, NIH-293T, and the normal ductal cells were cultured overnight in two 55 wells of a six well plate. The cells were harvested via scraping with 500 μ L of cell lysis buffer (CLB; 10 mM HEPES/10 mM NaCl/1mM KH₂PO₄/5mM NaHCO₃/1 mM CaCl₂/0.5 mM MgCl₂)/5m EDTA/10 μ g/ml aprotinin+10 μ g/ml leupeptin + 1 ug/ml pepstatin. The harvested cells were allowed to swell for five minutes, homogenized 50 times then centrifuged at 7500 rpm for five minutes. The pellet was suspended in 1 ml TSE/0.1% NP40/PI and homogenized for 30 minutes

followed with centrifugation at 5000 rpm for 5 minutes. The pellet was washed twice and suspended in 50 μ l of TSE/0.1%NP40/PI, leaving pure nuclei. The supernatant containing the cytosol with plasma membrane was centrifuged in a SWT0 rotor at 70,000 rpm for one hour. The pellet was resuspended and washed 2x with CLB to remove contaminating cytoplasmic proteins. Protein concentration of each fraction was determined via BCA assay (PIERCE Biotechnology), and equal amounts of protein from each fraction were size-fractionated by SDS-PAGE. Fractions were analyzed for plectin-1 expression via western blotting.

[0225] Competition experiments: mouse PDAC cells were incubated with FITC-labeled phage Clone 27 and either anti-plectin-1 antibody or vehicle for 1 hour at 37° C, washed, detached, then analyzed via flow cytometry (Becton Dickinson FACsCalibur (San Jose, CA)).

10 Peptide synthesis

[0226] Plectin-1 Targeted Peptide (PTP) (amino acid sequence: KTLLPTP (SEQ ID NO: 1)) and control peptide, see above, were synthesized with a GGSK(FITC)C linker conjugation of the peptide to a model fluorescent nanoparticle (crosslinked iron oxides [CLIO]-Cy5.5).

[0227] CLIO-Cy5.5 was synthesized in bulk using established procedures (see Montet et al., Bioconjug Chem 17:905-911 (2006); Reynolds et al., Bioconjug Chem 16:1240-1245 (2005); Wunderbaldinger et al., Acad Radiol 9 Suppl 2:S304-S306 (2002); and Schellenberger et al., Bioconjug Chem 15:1062-1067 (2004)), and aliquots used for the synthesis of the various nanoparticle conjugates. Briefly, T-10 dextran was dissolved in water mixed with ferric chloride and degassed by nitrogen purging. Ferric chloride solution was added to the mixture and the pH brought to 10 with ammonium hydroxide. The resulting particles were crosslinked with epichlorohydrin and ammonia to provide stability and amine groups for conjugation of fluorochromes and peptides. NHS-Cy5.5 was reacted with amino-CLIO in PBS overnight at 4° C and purified by size exclusion chromatography. Determination of Cy5.5 loading onto CLIO was done by absorbance spectroscopy at 680 nm using unreacted CLIO as a reference (Figure 8A). CLIO-Cy5.5 had the following physical properties: (a) size 38.7 nm (Figure 8B), (b) relaxation time constants R1-21.1 and R2-62.6 mM/ s, and (c) an average of 2.3 Cy5.5 per CLIO nanoparticle.

[0228] To produce plectin-1 targeted or control nanoparticles, succinimidyl iodoacetic acid was reacted with CLIO-Cy5.5 for 15 min, purified by size exclusion chromatography, then reacted with peptidyl-cysteine for 1 hour. Peptide-conjugated nanoparticles (PTP-NP) or controls (control-NP) were purified again using size exclusion chromatography and the ratio of peptides to nanoparticles was quantified at 497 nm by absorbance spectroscopy using unreacted CLIO as a reference (Figure 8C).

Intravital Laser Scanning Microscopy

[0229] Laser scanning microscopes with far red and near infrared imaging capabilities (IV 100, Olympus, Tokyo) have been described in details elsewhere (see, e.g., Alencar et al., Int. J. Cancer 117:335-339 (2005)). During all imaging sessions, mice were anesthetized (2% isoflurane in 2 1/minute O₂) and a small midline incision performed to expose the pancreas. As an in vivo screening approach, the inventors have used phage as targeted nanoparticles for imaging by labeling the phage coat proteins with a near infrared fluorochrome (Kelly et al., Neoplasia 8:1011-1018 (2006); and Newton et al., Neoplasia 8:772-780 (2006)). The Cy5.5 labeled phage were injected LV. 4 hours prior to imaging for both the distribution and tumor imaging studies. SYTOX green was injected 10 minutes prior to imaging (Fig. 3). Subsequent to imaging, tumors were removed for histological analysis. Serial frozen sections were hematoxylin and eosin (H&E) stained and also stained for the presence of M13 phage. For PTP-NP imaging (Figs. 5 and 6), the agent was injected 24 hours prior to IV 100 imaging. Angiosense (Visen Medical, Woburn, MA) was injected 10 minutes prior to imaging to visualize microvasculature. Images were acquired using appropriate dual excitation (561 nm for RITC, 633 nm for Cy5.5 and 748 nm for Angiosense-750). After fluorescence imaging, the pancreas was removed and embedded in degassed 1% low melting point agar in PBS to prevent susceptibility interfaces subsequent MRI imaging.

Biodistribution

[0230] Mice were maintained on a nonfluorescent diet (Harlen-Teklad) for 3 days prior to imaging and received an intravenous injection of PTP-NP or control probe (15 mg Fe/kg body weight), coupled to Cy5.5 for fluorescent imaging, 24 hours before biodistribution studies were carried out. Excised tissues were rinsed in PBS and imaged on the Siemens Bonsai system and Olympus OV100 system using Cy5.5 filters. Probe accumulation in tissues was compared to free probe, and biodistribution data were expressed as a percentage of injected dose. Fluorescence differences between the tissues were corrected by imaging tissues/organs from animals with no probe injected then subtracting this background from the total signal.

Magnetic resonance imaging (MRI)

[0231] Pancreata imaged optically *in vivo* were then embedded, and *ex vivo* MRI studies performed to directly correlate intrapancreatic signal intensity changes with histology. Imaging of resected and agar-embedded specimen was performed using a Bruker 4.7T Pharmascan magnet, with a 38 mm diameter transmit-receive radiofrequency coil. Scout and localizer images were obtained, followed by high resolution fast spin echo (FSE) and gradient echo sequences (GE). Specifically, for the T2 weighted FSE sequence, the following parameters were used: FOV 4.94 x 5.46 cm, matrix size of 512 x 512, slice thickness of 0.5 mm, RARE factor of 8, TE (effective) of 40 ms, TR 2811 ms, NEX of 50 for a total acquisition time of 2 hr 29 minutes. For the T2* weighted GE sequence, the parameters were: FOV 3x3 cm, matrix size 512 x 512, slice thickness of 0.5 mm, TE of 6.8 ms, TR of 398 ms, flip angle of 30 degrees, NEX of 50, for a total acquisition time of 2 hours 49 minutes. Fiducial markers were included to subsequently co-register high resolution MRI data sets with histologic sections.

Histology and Immunohistochemistry

[0232] Pancreas and PDAC specimens were isolated and either fixed in 10% paraformaldehyde or frozen in optimum cutting temperature (O.C.T.) compound as previously described (Bardeesy et al., Proc Natl Acad Sci USA 103:5947-5952 (2006)). The histology and immunohistochemical analyses were done as previously-described (Bardeesy et al., (2006) *supra*).

[0233] Serial frozen sections were stained with H&E or for the presence of M 13 (Amersham Biosciences, Piscataway, NJ) (Fig. 3) and plectin-1 (Fig. 4). Digital images were taken using a Nikon Eclipse E400 upright microscope (x20 and x43 objective) equipped with an Insight color camera. Serial frozen sections were stained with HE or imaged via fluorescence microscopy for the presence of PTPNP- Cy5.5 using a Nikon Eclipse 80i inverted microscope (203 objective) equipped with a 512 Photometrics Cascade CCD camera (Nikon).

EXAMPLE II**In vitro selection and validation of PDAC-specific peptides**

[0234] A genetically engineered mouse model of PDAC that recapitulates many of the histopathological, genomic and molecular features of the human disease was used (Carriere et al., Proc Natl Acad Sci USA. 104(11):4437-42 (2007); and Bardeesy et al. Proc Natl Acad Sci U S A 103:5947-5952 (2006)). The Kras/p53 L/L model (Bardeesy et al., Proc Natl Acad Sci U S A 103:5947-5952 (2006)) and wildtype controls served as a source of well-defined early passage PDAC cell lines and normal pancreatic ductal cells, respectively, for use in phage display selection and subtraction procedures to identify a pool of phage peptides specific for PDAC cells. Subsequent to selection procedures, the inventors isolated thirty individual phage plaques and performed an ELISA to identify the most selective phage for PDAC cells. The results of two experiments performed in triplicate are presented in the heat map shown in Figure 1A and in the bar graph shown in Figure 9. The heat map depicts affinity (mean absorbance values of indicated clones from the ELISA assay) and specificity (ratio of absorbance for PDAC cells versus normal ductal cells). Of the thirty phage clones analyzed, 16 phage clones (53%) had specificity for PDAC cells greater than 2 fold. Seven clones (Nos. 1, 5, 9, 15, 17, 22, and 27) were chosen for sequencing on the basis of ELISA and multidimensional analysis. For exemplary ELISA and sequencing methods see, e.g., Kelly et al., Circ. Res. 96:327-336 (2005); Kelly et al., Neoplasia 8:1011-1018 (2006)). Sequence results showed that Clones 27 and 5 shared identical peptide sequences (KTLLPTP, SEQ ID NO:1) and demonstrated ideal high affinity and specificity for the target PDAC cells.

[0235] Validation of Clone 27 as a PDAC marker (Figure 1B) was done in experiments described herein, in addition to testing clones 1 (SGVEFLH, SEQ ID NO:4), 9 (SKKDTHH, SEQ ID NO:5), 15 (TMAPSIK, SEQ ID NO:6), 17 (TQHQVTA, SEQ ID NO:7), and 22 (VNDRNVK, SEQ ID NO:8). The phage coat proteins were fluorescein labeled then the extent of phage clone binding and specificity quantified for mouse PDAC and normal ductal cells via flow cytometry (Fig. 1B). The results showed that Clone 27 was highly specific for mouse PDAC cells having a 112-fold specificity over normal ductal cells (Fig. 1C). Thus Clone 27 (SEQ ID NO:1 - KTLLPTP) demonstrated ideal affinity and specificity for the target PDAC cells (Kelly, et al., (2006) Neoplasia 8:1011-1018; Kelly et al. (2005) Circ Res 96: 327-336, all of which are herein incorporated by reference). Phage clone 15 was second in affinity with the remainder having nearly identical specificity.

[0236] Together, these data demonstrate the effectiveness of a cancer model-based phage screen as described herein for the identification and validation of phage clones with high affinity and specificity for cancer cells, such as mouse PDACs.

EXAMPLE III**Determining specificity of peptides for human PDACs**

5 [0237] The specificity of Clone 27 for human PDAC cells was evaluated as described in this example. Clone 27 and an unrelated phage clone were labeled with FITC, producing FITC-27 and a FITC-unrelated phage clone (negative control). Mouse PDAC cells as a positive control along with five human PDAC cell lines and with normal human ductal cells were incubated with each type of clone and then uptake of the clone was analyzed via fluorescence-activated cell sorter (FACS). FITC-27 (Clone 27) had an average specificity for PDAC cell lines of 141 (ratio of clone 27/unrelated 10 clone mean fluorescence) when compared with unrelated phage. In addition, the two phage clones (FITC-27 and FITC-unrelated) showed nearly identical, weak binding to normal human ductal cells (specificity = 0.85) (Fig. 1C).

EXAMPLE IV**Utilizing peptides for identifying human PDACs**

15 [0238] The use of the identified phage for the detection of mouse and human PDACs was demonstrated as follows. Phage 27 labeled with a fluorochrome was used as a probe to test binding to frozen sections of normal pancreas, 20 pancreases containing focal PanINs, and pancreata with PDAC. While no binding was observed in wild-type mouse pancreata or in normal regions adjacent to lesions, prominent binding was observed in PanINs and PDAC lesions. Control phage failed to detect any lesions (Figure 2, bottom row). Significantly, phage Clone 27 was able to specifically 25 detect human PDAC, whereas control phage failed to stain human PDAC specimens (Figure 2, (far right)). These results demonstrate that the phage probes bind to evolving mouse and human PDAC, supporting the utility of our models-based screening approach for the generation of candidate PDAC-specific diagnostic agents.

EXAMPLE V**Tumor localization of PDAC targeted phage**

30 [0239] Since phage clones 27 and 15 had the most favorable binding characteristics in vitro, these clones were further tested in vivo binding in wild-type animals, animals harboring PanINs, and animals harboring palpable pancreatic tumors. These animals were injected via tail vein with 1 nanomole of fluorescently labeled phage clone 27 and phage clone 15-alone 35 or in combination-and then imaged via intravital confocal microscopy 4 h postinjection (Figure 10A). Clone 27 illuminated PanINs and PDAC with a strong fluorescent signal, suggesting phage binding to tumors cells, whereas only a weak scattered signal was observed in the pancreas of wild-type mice (Figures 3A, 3B, and Figure 10C). The fluorescent signal was virtually absent when control phage was injected into animals harboring emerging or advanced PDAC (Figure 10B). Clones 15 and 27 showed distinctly different distributions within the pancreas and also different peptide sequences (Figure 10A), suggesting they target unique proteins. However while clone 15 did localize to the pancreas, total signal was less than that of clone 27, thus further experiments were focused on Clone 27.

40 [0240] Further documentation of the specificity of phage Clone 27 binding in vivo, pancreata from clone 27-injected animals were fixed and analyzed by immunohistochemistry using antibodies specific to phage coat proteins (Figure 3). In areas with PanINs or PDAC (black boxes in Figure 3B and 3C), there was strong uptake of phage whereas in regions of ductal metaplasia or normal pancreas (red boxes in Figure 3B and 3C) phage were undetectable. These studies supported the specificity of Clone 27 for cancer cells showing that phage clone 27 was localized to PanINs and PDAC 45 while absent in normal pancreatic tissues or regions of ductal metaplasia, which are low-grade neoplasms or reactive lesions associated with pancreatic damage (Figure 3) (Murtaugh et al., Cancer Cell 11: 211-213 (2007)).

EXAMPLE VI**Identification of Plectin-1 as the binding partner for peptide 27**

50 [0241] Since clone 27 showed specificity for human and mouse PDAC in vitro and in vivo, the next step was to determine its cellular binding partner. Using the phage as an affinity ligand, a unique 500 kDa band was identified in the mouse PDAC cell lysates via pulldown assay (Fig. 4A, left panel). In addition, far western analysis of PDAC lysates with biotinylated phage as the probe identified a band of similar molecular weight that was not recognized by control phage (Fig. 4A, right panel). Mass spectroscopic analysis of the isolated band revealed plectin-1, an intermediate, filament and important cross-linking element of the cytoskeleton (Fig.4B) (Sonnenberg, et al. (2007) Exp Cell Res 313: 2189-2203). Western blot using lysates from the phage pull down confirmed the presence of a band that crossreacts with the plectin-

1 antibody (Fig. 4C). Plectin-1 was found to be present in the cellular membrane as well as the nucleus and cytoplasm of both murine and human PDAC cells (Fig. 4D). Normal mouse pancreatic ductal cells showed low levels of plectin-1 expression, whereas normal human pancreatic ductal cells showed plectin-1 expression in the cytoplasm and nucleus but not on the membrane (Figure 4D). In contrast, HUVECs showed very low levels of plectin-1 expression in the nucleus.

5 NIH-3T3 cells were used as a control for cellular locations of plectin-1 expression because they were known to have plectin-1 in the cytoplasm and nucleus but not on the cell surface (Sonnenberg et al., Exp Cell Res 313: 2189-2203 (2007)). As was expected in the control cells, plectin-1 was absent from the membrane but present in the cytoplasmic and nuclear fractions of fibroblasts (Figure 4D).

10 [0242] Immunohistochemical analysis of sections from normal, PanIN, and PDAC-harboring mice corroborated the Western analysis findings. Normal animals had scattered plectin-1 staining, whereas in PanINs and PDAC, plectin-1 was expressed in the lesions but not in the surrounding tissue (Figure 4E). The plectin-1 staining patterns were nearly identical to that observed by PDAC-targeted phage shown in Figure 3 and Figure 4E. Finally, in a competition experiment, coincubation of anti-plectin-1 antibody and FITC-labeled phage clone 27 with PDAC cells resulted in 96.9% abrogation 15 of binding (Figure 4F).

EXAMPLE VII

Development of plectin-1 targeted PDAC imaging agents

20 [0243] In order to develop a nonbiologic, synthetic imaging agent with translational potential, the inventors chemically synthesized and attached PTP to a magnetofluorescent nanoparticle (PTP-NP) (schematic, Figure 5A).

25 [0244] The resultant MRI/optically detectable agent was tested in 9-wk-old Kras/p53L/p mice (Fig. 6A). At that age, these mice do not exhibit outward signs of illness but typically harbor small, focal PDAC, as well as regions of normal pancreas, ductal metaplasia, and fibrosis. Twenty-four hours after IV administration of the targeted nanoparticle, intravital confocal microscopy detected discrete areas of fluorescence in the abdominal region of these mice, suggestive of agent uptake (Figure 5B upper left and 6B, left). The agent was specifically present in the tumor tissue as a vasculature agent administered 10 min before injection failed to colocalize (Figure 5B upper right and 6B, right). The in vivo fluorescence correlated with surface reflectance imaging of the excised pancreas where discrete foci of signal were found (Figure 30 5C, 6C and 6D). In contrast, control-NP failed to highlight any regions of the pancreas (Figure 5B, lower left), although these tumors were similarly vascularized (Figure 5B, lower right).

[0245] Biodistribution studies revealed specific uptake in tumors with minimal uptake in muscle or skin, two tissues with reported plectin-1 expression (Figure 5D). In addition, tumor uptake relative to normal pancreas was 10.1-fold higher.

35 [0246] Similarly, MRI showed a reduction in magnetic resonance (MR) signal indicative of agent presence in focal regions of the pancreas (Figure 7A). From the biodistribution data, 3.13%-injected dose of material was present in the tumors. Using previously established thresholds for direct MRI sensitivity of 10 ng of Fe/g of tissue (Weissleder et al. (1997) J Magn Reson Imaging 7: 258-263 (1997)), the resulting signal was calculated at 20-fold over the threshold of detection. In addition, it is likely that current sensitivity is higher than what was previously published given available motion correction and multi-echo chemical sequences.

40 [0247] Histological analysis confirmed that the loss of signal associated with PTP-NP uptake was primarily in regions of PDAC but not in normal regions or regions of ductal metaplasia (Figure 7B). Fluorescence microscopy of the sections demonstrated PTP-NP accumulation in areas of PDAC (Figure 7C, left) but not in areas of normal pancreas (Figure 7C, right).

OTHER EMBODIMENTS

45 [0248] All publications and patents mentioned in the above specification are herein incorporated by reference in their entirety. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention that are obvious to those skilled in medicine, therapeutics, pharmaceuticals, MRI, in vivo imaging, molecular biology, biochemistry, chemistry, and cell biology or related fields are intended to be within the scope of the following claims. All references cited herein are incorporated in their entirety.

Clauses

55 [0249] The invention will now be defined by the following clauses:

1. A peptide ligand comprising:

5 a first portion comprising a plectin-1 binding moiety, coupled to a second portion comprising a detectable moiety or a therapeutic agent.

2. The peptide ligand of clause 1, wherein the plectin-1 binding moiety is an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, or 4 - 8, or a peptidomimetic thereof.

10 3. The peptide ligand of clause 1, wherein the detectable moiety is selected from the group consisting of a radioactive isotope, a magnetic compound, an x-ray absorber, a chemical compound, a biological tag, and a fluorescent molecule.

4. The peptide ligand of clause 1, wherein the therapeutic agent is a cytotoxic moiety or an immunomodulatory moiety.

15 5. The peptide ligand of clause 1, further comprising a linker between the first portion and the second portion.

6. The peptide ligand of clause 5, wherein the linker is a flexible amino acid sequence.

7. The peptide ligand of clause 4, wherein the linker is a photolinker.

20 8. The peptide ligand of clause 1, wherein the second portion further comprises a physiologically inert nanoparticle.

9. The peptide ligand of clause 8, wherein the nanoparticle is magnetic, fluorescent, or radioactive.

25 10. The peptide ligand of clause 1, wherein the second portion comprises a fluorochrome.

11. The peptide ligand of clause 10, wherein the fluorochrome is a near infrared fluorochrome.

30 12. The peptide ligand of clause 1, wherein the second portion comprises a crosslinked iron oxide nanoparticle conjugated to a NIRF.

13. A peptide ligand according to clause 1 comprising SEQ ID NO: 1 coupled to a nanoparticle.

35 14. The peptide ligand of clause 13, wherein the nanoparticle is a magnetofluorescent nanoparticle, preferably wherein the magnetofluorescent nanoparticle comprises a near infrared (NIR) fluorochrome (NIRF).

15. The peptide ligand of any of clauses 1-14 for the diagnosis or treatment of pancreatic ductal adenocarcinoma.

40 16. Use of the peptide ligand of any of clauses 1-14 for the diagnosis or treatment of pancreatic ductal adenocarcinoma, or in the manufacture of a medicament for the diagnosis or treatment of pancreatic ductal adenocarcinoma.

45

50

55

SEQUENCE LISTING

<110> The General Hospital Corporation
5
<120> PLECTIN-1 TARGETED AGENTS FOR DETECTION
AND TREATMENT OF PANCREATIC DUCTAL ADENOCARCINOMA

10
<130> 00786-0676WO1
<140> PCT/US2009/040480
<141> 2009-04-14

15
<150> US 61/044,818
<151> 2008-04-14
<160> 26

20
<170> FastSEQ for Windows Version 4.0
<210> 1
<211> 7
<212> PRT
<213> Artificial Sequence

25
<220>
<223> Synthetic peptide

30
<400> 1
Lys Thr Leu Leu Pro Thr Pro
1 5

35
<210> 2
<211> 11
<212> PRT
<213> Artificial Sequence
<220>
<223> Synthetic peptide

40
<400> 2
Lys Thr Leu Leu Pro Thr Pro Gly Gly Ser Lys
1 5 10

45
<210> 3
<211> 12
<212> PRT
<213> Artificial Sequence
<220>
<223> Synthetic peptide

50
<220>
<221> MOD_RES
<222> 11
<223> Conjugated Fluorescein isothiocyanate (FITC)

55
<400> 3
Lys Thr Leu Leu Pro Thr Pro Gly Gly Ser Lys Cys
1 5 10

5 <210> 4
<211> 7
<212> PRT
<213> Artificial Sequence

10 <220>
<223> Synthetic peptide

15 <400> 4
Ser Gly Val Glu Phe Leu His
1 5

20 <210> 5
<211> 7
<212> PRT
<213> Artificial Sequence

25 <220>
<223> Synthetic peptide

30 <400> 5
Ser Lys Lys Asp Thr His His
1 5

35 <210> 6
<211> 7
<212> PRT
<213> Artificial Sequence

40 <220>
<223> Synthetic peptide

45 <400> 6
Thr Met Ala Pro Ser Ile Lys
1 5

50 <210> 7
<211> 7
<212> PRT
<213> Artificial Sequence

55 <220>
<223> Synthetic peptide

60 <400> 7
Thr Gln His Gln Val Thr Ala
1 5

65 <210> 8
<211> 7
<212> PRT
<213> Artificial Sequence

70 <220>
<223> Synthetic peptide

75 <400> 8

Val Asn Asp Arg Asn Val Lys
1 5

5 <210> 9
<211> 14
<212> PRT
<213> Artificial Sequence

10 <220>
<223> Synthetic peptide

<400> 9
Asp Ser Gln Asp Ala Gly Gly Phe Pro Gly Pro Glu Asp Arg
1 5 10

15 <210> 10
<211> 16
<212> PRT
<213> Artificial Sequence

20 <220>
<223> Synthetic peptide

<400> 10
Glu Ala Gln Ala Val Pro Ala Thr Leu Gln Glu Leu Glu Ala Thr Lys
1 5 10 15

25 <210> 11
<211> 14
<212> PRT
<213> Artificial Sequence

30 <220>
<223> Synthetic peptide

<400> 11
Ala Gln Ala Glu Ala Gln Gln Pro Val Phe Asn Thr Leu Arg
1 5 10

35 <210> 12
<211> 11
<212> PRT
<213> Artificial Sequence

40 <220>
<223> Synthetic peptide

<400> 12
Gly Gly Ala Glu Gly Glu Leu Gln Ala Leu Arg
1 5 10

45 <210> 13
<211> 9
<212> PRT
<213> Artificial Sequence

50 <220>
<223> Synthetic peptide

<400> 13
Leu Ala Ala Glu Gln Glu Leu Ile Arg
1 5

<210> 14
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 14
Leu Arg Ala Glu Thr Glu Gln Gly Glu Gln Gln Arg
1 5 10

<210> 15
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 15
Ser Lys Glu Gln Ala Glu Leu Glu Ala Ala Arg
1 5 10

<210> 16
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 16
Ala Gln Val Glu Gln Glu Leu Thr Thr Leu Arg
1 5 10

<210> 17
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 17
Ser Gln Val Glu Glu Glu Leu Phe Ser Val Arg
1 5 10

<210> 18
<211> 10
<212> PRT
<213> Artificial Sequence

5 <220>
<223> Synthetic peptide

10 <400> 18
Leu Ser Val Ala Ala Gln Glu Ala Ala Arg
1 5 10

15 <210> 19
<211> 11
<212> PRT
<213> Artificial Sequence

20 <220>
<223> Synthetic peptide

25 <400> 19
Gln Leu Ala Glu Glu Asp Leu Ala Gln Gln Arg
1 5 10

30 <210> 20
<211> 9
<212> PRT
<213> Artificial Sequence

35 <220>
<223> Synthetic peptide

40 <400> 20
Leu Ser Val Tyr Thr Ala Leu Gln Arg
1 5

45 <210> 21
<211> 16
<212> PRT
<213> Artificial Sequence

50 <220>
<223> Synthetic peptide

55 <400> 21
Leu Leu Asp Ala Gln Leu Ser Thr Gly Gly Ile Val Asp Pro Ser Lys
1 5 10 15

60 <210> 22
<211> 10
<212> PRT
<213> Artificial Sequence

65 <220>
<223> Synthetic peptide

70 <400> 22
Leu Thr Ala Glu Asp Leu Tyr Glu Ala Arg
1 5 10

75 <210> 23
<211> 9
<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide

5

<400> 23

Val Ser Leu Asp Glu Ala Leu Gln Arg
1 5

10

<210> 24

<211> 4684

<212> PRT

<213> Homo sapiens

15 <400> 24

Met Val Ala Gly Met Leu Met Pro Arg Asp Gln Leu Arg Ala Ile Tyr

1 5 10 15

Glu Val Leu Phe Arg Glu Gly Val Met Val Ala Lys Lys Asp Arg Arg
20 25 30Pro Arg Ser Leu His Pro His Val Pro Gly Val Thr Asn Leu Gln Val
35 40 45Met Arg Ala Met Ala Ser Leu Arg Ala Arg Gly Leu Val Arg Glu Thr
50 55 60Phe Ala Trp Cys His Phe Tyr Trp Tyr Leu Thr Asn Glu Gly Ile Ala
65 70 75 8020 His Leu Arg Gln Tyr Leu His Leu Pro Pro Glu Ile Val Pro Ala Ser
85 90 95Leu Gln Arg Val Arg Arg Pro Val Ala Met Val Met Pro Ala Arg Arg
100 105 11030 Thr Pro His Val Gln Ala Val Gln Gly Pro Leu Gly Ser Pro Pro Lys
115 120 125Arg Gly Pro Leu Pro Thr Glu Glu Gln Arg Val Tyr Arg Arg Lys Glu
130 135 140Leu Glu Glu Val Ser Pro Glu Thr Pro Val Val Pro Ala Thr Thr Gln
145 150 155 160Arg Thr Leu Ala Arg Pro Gly Pro Glu Pro Ala Pro Ala Thr Asp Glu
165 170 17535 Arg Asp Arg Val Gln Lys Lys Thr Phe Thr Lys Trp Val Asn Lys His
180 185 190Leu Ile Lys Ala Gln Arg His Ile Ser Asp Leu Tyr Glu Asp Leu Arg
195 200 205Asp Gly His Asn Leu Ile Ser Leu Leu Glu Val Leu Ser Gly Asp Ser
210 215 22040 Leu Pro Arg Glu Lys Gly Arg Met Arg Phe His Lys Leu Gln Asn Val
225 230 235 240Gln Ile Ala Leu Asp Tyr Leu Arg His Arg Gln Val Lys Leu Val Asn
245 250 25545 Ile Arg Asn Asp Ile Ala Asp Gly Asn Pro Lys Leu Thr Leu Gly
260 265 270Leu Ile Trp Thr Ile Ile Leu His Phe Gln Ile Ser Asp Ile Gln Val
275 280 285Ser Gly Gln Ser Glu Asp Met Thr Ala Lys Glu Lys Leu Leu Leu Trp
290 295 30050 Ser Gln Arg Met Val Glu Gly Tyr Gln Gly Leu Arg Cys Asp Asn Phe
305 310 315 320Thr Ser Ser Trp Arg Asp Gly Arg Leu Phe Asn Ala Ile Ile His Arg
325 330 335His Lys Pro Leu Leu Ile Asp Met Asn Lys Val Tyr Arg Gln Thr Asn
340 345 35055 Leu Glu Asn Leu Asp Gln Ala Phe Ser Val Ala Glu Arg Asp Leu Gly
355 360 365

Val Thr Arg Leu Leu Asp Pro Glu Asp Val Asp Val Pro Gln Pro Asp

	370	375	380
	Glu Lys Ser Ile Ile Thr Tyr Val Ser Ser Leu Tyr Asp Ala Met Pro		
5	385 390 395 400		
	Arg Val Pro Asp Val Gln Asp Gly Val Arg Ala Asn Glu Leu Gln Leu		
	405 410 415 420 425 430		
	Arg Trp Gln Glu Tyr Arg Glu Leu Val Leu Leu Leu Gln Trp Met		
	435 440 445 450 455 460		
10	Glu Glu Ile Glu Ile Leu Trp Ser Gln Phe Leu Lys Phe Lys Glu Met		
	465 470 475 480 485 490 495		
	Gln Ser Leu Glu Gly Ala Val Gln Ala Gly Gln Leu Lys Val Pro Pro		
15	500 505 510 515 520 525 530 535 540 545 550 555 560		
	Gly Tyr His Pro Leu Asp Val Glu Lys Glu Trp Gly Lys Leu His Val		
	555 560 565 570 575 580 585 590 595 600 605 610 615 620		
20	Ala Ile Leu Glu Arg Glu Lys Gln Leu Arg Ser Glu Phe Glu Arg Leu		
	625 630 635 640 645 650 655 660 665 670 675 680 685 690 695 700		
	Glu Cys Leu Gln Arg Ile Val Thr Lys Leu Gln Met Glu Ala Gly Leu		
25	705 710 715 720 725 730 735 740 745 750 755 760 765 770 775 780 785 790 795 800		
	Cys Glu Glu Gln Leu Asn Gln Ala Asp Ala Leu Leu Gln Ser Asp Val		
	805 810 815 820 825 830 835 840 845 850 855 860 865 870 875 880		
30	Arg Leu Leu Ala Ala Gly Lys Val Pro Gln Arg Ala Gly Glu Val Glu		
	885 890 895 900 905 910 915 920 925 930 935 940 945 950 955 960 965 970 975 980 985 990		
35	Arg Asp Leu Asp Lys Ala Asp Ser Met Ile Arg Leu Leu Phe Asn Asp		
	995 1000 1005 1010 1015 1020 1025 1030 1035 1040 1045 1050 1055 1060 1065 1070 1075 1080 1085 1090 1095 1100 1105 1110 1115 1120 1125 1130 1135 1140 1145 1150 1155 1160 1165 1170 1175 1180 1185 1190 1195 1200 1205 1210 1215 1220 1225 1230 1235 1240 1245 1250 1255 1260 1265 1270 1275 1280 1285 1290 1295 1300 1305 1310 1315 1320 1325 1330 1335 1340 1345 1350 1355 1360 1365 1370 1375 1380 1385 1390 1395 1400 1405 1410 1415 1420 1425 1430 1435 1440 1445 1450 1455 1460 1465 1470 1475 1480 1485 1490 1495 1500 1505 1510 1515 1520 1525 1530 1535 1540 1545 1550 1555 1560 1565 1570 1575 1580 1585 1590 1595 1600 1605 1610 1615 1620 1625 1630 1635 1640 1645 1650 1655 1660 1665 1670 1675 1680 1685 1690 1695 1700 1705 1710 1715 1720 1725 1730 1735 1740 1745 1750 1755 1760 1765 1770 1775 1780 1785 1790 1795 1800 1805 1810 1815 1820 1825 1830 1835 1840 1845 1850 1855 1860 1865 1870 1875 1880 1885 1890 1895 1900 1905 1910 1915 1920 1925 1930 1935 1940 1945 1950 1955 1960 1965 1970 1975 1980 1985 1990 1995 2000 2005 2010 2015 2020 2025 2030 2035 2040 2045 2050 2055 2060 2065 2070 2075 2080 2085 2090 2095 2100 2105 2110 2115 2120 2125 2130 2135 2140 2145 2150 2155 2160 2165 2170 2175 2180 2185 2190 2195 2200 2205 2210 2215 2220 2225 2230 2235 2240 2245 2250 2255 2260 2265 2270 2275 2280 2285 2290 2295 2300 2305 2310 2315 2320 2325 2330 2335 2340 2345 2350 2355 2360 2365 2370 2375 2380 2385 2390 2395 2400 2405 2410 2415 2420 2425 2430 2435 2440 2445 2450 2455 2460 2465 2470 2475 2480 2485 2490 2495 2500 2505 2510 2515 2520 2525 2530 2535 2540 2545 2550 2555 2560 2565 2570 2575 2580 2585 2590 2595 2600 2605 2610 2615 2620 2625 2630 2635 2640 2645 2650 2655 2660 2665 2670 2675 2680 2685 2690 2695 2700 2705 2710 2715 2720 2725 2730 2735 2740 2745 2750 2755 2760 2765 2770 2775 2780 2785 2790 2795 2800 2805 2810 2815 2820 2825 2830 2835 2840 2845 2850 2855 2860 2865 2870 2875 2880 2885 2890 2895 2900 2905 2910 2915 2920 2925 2930 2935 2940 2945 2950 2955 2960 2965 2970 2975 2980 2985 2990 2995 3000 3005 3010 3015 3020 3025 3030 3035 3040 3045 3050 3055 3060 3065 3070 3075 3080 3085 3090 3095 3100 3105 3110 3115 3120 3125 3130 3135 3140 3145 3150 3155 3160 3165 3170 3175 3180 3185 3190 3195 3200 3205 3210 3215 3220 3225 3230 3235 3240 3245 3250 3255 3260 3265 3270 3275 3280 3285 3290 3295 3300 3305 3310 3315 3320 3325 3330 3335 3340 3345 3350 3355 3360 3365 3370 3375 3380 3385 3390 3395 3400 3405 3410 3415 3420 3425 3430 3435 3440 3445 3450 3455 3460 3465 3470 3475 3480 3485 3490 3495 3500 3505 3510 3515 3520 3525 3530 3535 3540 3545 3550 3555 3560 3565 3570 3575 3580 3585 3590 3595 3600 3605 3610 3615 3620 3625 3630 3635 3640 3645 3650 3655 3660 3665 3670 3675 3680 3685 3690 3695 3700 3705 3710 3715 3720 3725 3730 3735 3740 3745 3750 3755 3760 3765 3770 3775 3780 3785 3790 3795 3800 3805 3810 3815 3820 3825 3830 3835 3840 3845 3850 3855 3860 3865 3870 3875 3880 3885 3890 3895 3900 3905 3910 3915 3920 3925 3930 3935 3940 3945 3950 3955 3960 3965 3970 3975 3980 3985 3990 3995 4000 4005 4010 4015 4020 4025 4030 4035 4040 4045 4050 4055 4060 4065 4070 4075 4080 4085 4090 4095 4100 4105 4110 4115 4120 4125 4130 4135 4140 4145 4150 4155 4160 4165 4170 4175 4180 4185 4190 4195 4200 4205 4210 4215 4220 4225 4230 4235 4240 4245 4250 4255 4260 4265 4270 4275 4280 4285 4290 4295 4300 4305 4310 4315 4320 4325 4330 4335 4340 4345 4350 4355 4360 4365 4370 4375 4380 4385 4390 4395 4400 4405 4410 4415 4420 4425 4430 4435 4440 4445 4450 4455 4460 4465 4470 4475 4480 4485 4490 4495 4500 4505 4510 4515 4520 4525 4530 4535 4540 4545 4550 4555 4560 4565 4570 4575 4580 4585 4590 4595 4600 4605 4610 4615 4620 4625 4630 4635 4640 4645 4650 4655 4660 4665 4670 4675 4680 4685 4690 4695 4700 4705 4710 4715 4720 4725 4730 4735 4740 4745 4750 4755 4760 4765 4770 4775 4780 4785 4790 4795 4800 4805 4810 4815 4820 4825 4830 4835 4840 4845 4850 4855 4860 4865 4870 4875 4880 4885 4890 4895 4900 4905 4910 4915 4920 4925 4930 4935 4940 4945 4950 4955 4960 4965 4970 4975 4980 4985 4990 4995 5000 5005 5010 5015 5020 5025 5030 5035 5040 5045 5050 5055 5060 5065 5070 5075 5080 5085 5090 5095 5100 5105 5110 5115 5120 5125 5130 5135 5140 5145 5150 5155 5160 5165 5170 5175 5180 5185 5190 5195 5200 5205 5210 5215 5220 5225 5230 5235 5240 5245 5250 5255 5260 5265 5270 5275 5280 5285 5290 5295 5300 5305 5310 5315 5320 5325 5330 5335 5340 5345 5350 5355 5360 5365 5370 5375 5380 5385 5390 5395 5400 5405 5410 5415 5420 5425 5430 5435 5440 5445 5450 5455 5460 5465 5470 5475 5480 5485 5490 5495 5500 5505 5510 5515 5520 5525 5530 5535 5540 5545 5550 5555 5560 5565 5570 5575 5580 5585 5590 5595 5600 5605 5610 5615 5620 5625 5630 5635 5640 5645 5650 5655 5660 5665 5670 5675 5680 5685 5690 5695 5700 5705 5710 5715 5720 5725 5730 5735 5740 5745 5750 5755 5760 5765 5770 5775 5780 5785 5790 5795 5800 5805 5810 5815 5820 5825 5830 5835 5840 5845 5850 5855 5860 5865 5870 5875 5880 5885 5890 5895 5900 5905 5910 5915 5920 5925 5930 5935 5940 5945 5950 5955 5960 5965 5970 5975 5980 5985 5990 5995 6000 6005 6010 6015 6020 6025 6030 6035 6040 6045 6050 6055 6060 6065 6070 6075 6080 6085 6090 6095 6100 6105 6110 6115 6120 6125 6130 6135 6140 6145 6150 6155 6160 6165 6170 6175 6180 6185 6190 6195 6200 6205 6210 6215 6220 6225 6230 6235 6240 6245 6250 6255 6260 6265 6270 6275 6280 6285 6290 6295 6300 6305 6310 6315 6320 6325 6330 6335 6340 6345 6350 6355 6360 6365 6370 6375 6380 6385 6390 6395 6400 6405 6410 6415 6420 6425 6430 6435 6440 6445 6450 6455 6460 6465 6470 6475 6480 6485 6490 6495 6500 6505 6510 6515 6520 6525 6530 6535 6540 6545 6550 6555 6560 6565 6570 6575 6580 6585 6590 6595 6600 6605 6610 6615 6620 6625 6630 6635 6640 6645 6650 6655 6660 6665 6670 6675 6680 6685 6690 6695 6700 6705 6710 6715 6720 6725 6730 6735 6740 6745 6750 6755 6760 6765 6770 6775 6780 6785 6790 6795 6800 6805 6810 6815 6820 6825 6830 6835 6840 6845 6850 6855 6860 6865 6870 6875 6880 6885 6890 6895 6900 6905 6910 6915 6920 6925 6930 6935 6940 6945 6950 6955 6960 6965 6970 6975 6980 6985 6990 6995 7000 7005 7010 7015 7020 7025 7030 7035 7040 7045 7050 7055 7060 7065 7070 7075 7080 7085 7090 7095 7100 7105 7110 7115 7120 7125 7130 7135 7140 7145 7150 7155 7160 7165 7170 7175 7180 7185 7190 7195 7200 7205 7210 7215 7220 7225 7230 7235 7240 7245 7250 7255 7260 7265 7270 7275 7280 7285 7290 7295 7300 7305 7310 7315 7320 7325 7330 7335 7340 7345 7350 7355 7360 7365 7370 7375 7380 7385 7390 7395 7400 7405 7410 7415 7420 7425 7430 7435 7440 7445 7450 7455 7460 7465 7470 7475 7480 7485 7490 7495 7500 7505 7510 7515 7520 7525 7530 7535 7540 7545 7550 7555 7560 7565 7570 7575 7580 7585 7590 7595 7600 7605 7610 7615 7620 7625 7630 7635 7640 7645 7650 7655 7660 7665 7670 7675 7680 7685 7690 7695 7700 7705 7710 7715 7720 7725 7730 7735 7740 7745 7750 7755 7760 7765 7770 7775 7780 7785 7790 7795 7800 7805 7810 7815 7820 7825 7830 7835 7840 7845 7850 7855 7860 7865 7870 7875 7880 7885 7890 7895 7900 7905 7910 7915 7920 7925 7930 7935 7940 7945 7950 7955 7960 7965 7970 7975 7980 7985 7990 7995 8000 8005 8010 8015 8020 8025 8030 8035 8040 8045 8050 8055 8060 8065 8070 8075 8080 8085 8090 8095 8100 8105 8110 8115 8120 8125 8130 8135 8140 8145 8150 8155 8160 8165 8170 8175 8180 8185 8190 8195 8200 8205 8210 8215 8220 8225 8230 8235 8240 8245 8250 8255 8260 8265 8270 8275 8280 8285 8290 8295 8300 8305 8310 8315 8320 8325 8330 8335 8340 8345 8350 8355 8360 8365 8370 8375 8380 8385 8390 8395 8400 8405 8410 8415 8420 8425 8430 8435 8440 8445 8450 8455 8460 8465 8470 8475 8480 8485 8490 8495 8500 8505 8510 8515 8520 8525 8530 8535 8540 8545 8550 8555 8560 8565 8570 8575 8580 8585 8590 8595 8600 8605 8610 8615 8620 8625 8630 8635 8640 8645 8650 8655 8660 8665 8670 8675 8680 8685 8690 8695 8700 8705 8710 8715 8720 8725 8730 8735 8740 8745 8750 8755 8760 8765 8770 8775 8780 8785 8790 8795 8800 8805 8810 8815 8820 8825 8830 8835 8840 8845 8850 8855 8860 8865 8870 8875 8880 8885 8890 8895 8900 8905 8910 8915 8920 8925 8930 8935 8940 8945 8950 8955 8960 8965 8970 8975 8980 8985 8990 8995 9000 9005 9010 9015 9020 9025 9030 9035 9040 9045 9050 9055 9060 9065 9070 9075 9080 9085 9090 9095 9100 9105 9110 9115 9120 9125 9130 9135 9140 9145 9150 9155 9160 9165 9170 9175 9180 9185 9190 9195 9200 9205 9210 9215 9220 9225 9230 9235 9240 9245 9250 9255 9260 9265 9270 9275 9280 9285 9290 9295 9300 9305 9310 9315 9320 9325 9330 9335 9340 9345 9350 9355 9360 9365 9370 9375 9380 9385 9390 9395 9400 9405 9410 9415 9420 9425 9430 9435 9440 9445 9450 9455 9460 9465 9470 9475 9480 9485 9490 9495 9500 9505 9510 9515 9520 9525 9530 9535 9540 9545 9550 9555 9560 9565 9570 9575 9580 9585 9590 9595 9600 9605 9610 9615 9620 9625 9630 9635 9640 9645 9650 9655 9660 9665 9670 9675 9680 9685 9690 9695 9700 9705 9710 9715 9720 9725 9730 9735 9740 9745 9750 9755 9760 9765 9770 9775 9780 9785 9790 9795 9800 9805 9810 9815 9820 9825 9830 9835 9840 9845 9850 9855 9860 9865 9870 9875 9880 9885 9890 9895 9900 9905 9910 9915 9920 9925 9930 9935 9940 9945 9950 9955 9960 9965 9970 9975 9980 9985 9990 9995 10000 10005 10010 10015 10020 10025 10030 10035 10040 10045 10050 10055 10060 10065 10070 10075 10080 10085 10090 10095 10100 10105 10110 10115 10120 10125 10130 10135 10140 10145 10150 10155 10160 10165 10170 10175 10180 10185 10190 10195 10200 10205 10210 10215 10220 10225 10230 10235 10240 10245 10250 10255 10260 10265 10270 10275 10280 10285 10290 10295 10300 10305 10310 10315 10320 10325 10330 10335 10340 10345 10350 10355 10360 10365 10370 10375 10380 10385 10390 10395 10400 10405 10410 10415 10420 10425 10430 10435 10440 10445 10450 10455 10460 10465 10470 10475 10480 10485 10490 10495 10500 10505 10510 10515 10520 10525 10530 10535 10540 10545 10550 10555 10560 10565 10570 10575 10580 10585 10590 10595 10600 10605 10610 10615 10620 10625 10630 10635 10640 10645 10650 10655 10660 1		

	Ala Leu Arg Arg Lys Tyr Ser Cys Asp Arg Ser Ala Thr Val Thr Arg		
	885	890	895
	Leu Glu Asp Leu Leu Gln Asp Ala Gln Asp Glu Lys Glu Gln Leu Asn		
5	900	905	910
	Glu Tyr Lys Gly His Leu Ser Gly Leu Ala Lys Arg Ala Lys Ala Val		
	915	920	925
	Val Gln Leu Lys Pro Arg His Pro Ala His Pro Met Arg Gly Arg Leu		
10	930	935	940
	Pro Leu Leu Ala Val Cys Asp Tyr Lys Gln Val Glu Val Thr Val His		
	945	950	955
15	960		
	Lys Gly Asp Glu Cys Gln Leu Val Gly Pro Ala Gln Pro Ser His Trp		
	965	970	975
	Lys Val Leu Ser Ser Ser Gly Ser Glu Ala Ala Val Pro Ser Val Cys		
20	980	985	990
	Phe Leu Val Pro Pro Pro Asn Gln Glu Ala Gln Glu Ala Val Thr Arg		
	995	1000	1005
25	Leu Glu Ala Gln His Gln Ala Leu Val Thr Leu Trp His Gln Leu His		
	1010	1015	1020
	Val Asp Met Lys Ser Leu Leu Ala Trp Gln Ser Leu Arg Arg Asp Val		
	1025	1030	1035
30	1040		
	Gln Leu Ile Arg Ser Trp Ser Leu Ala Thr Phe Arg Thr Leu Lys Pro		
	1045	1050	1055
	Glu Glu Gln Arg Gln Ala Leu His Ser Leu Glu Leu His Tyr Gln Ala		
	1060	1065	1070
35	Phe Leu Arg Asp Ser Gln Asp Ala Gly Gly Phe Gly Pro Glu Asp Arg		
	1075	1080	1085
	Leu Met Ala Glu Arg Glu Tyr Gly Ser Cys Ser His His Tyr Gln Gln		
40	1090	1095	1100
	Leu Leu Gln Ser Leu Glu Gln Gly Ala Gln Glu Glu Ser Arg Cys Gln		
	1105	1110	1115
45	1120		
	Arg Cys Ile Ser Glu Leu Lys Asp Ile Arg Leu Gln Leu Glu Ala Cys		
	1125	1130	1135
50	Glu Thr Arg Thr Val His Arg Leu Arg Leu Pro Leu Asp Lys Glu Pro		
	1140	1145	1150
	Ala Arg Glu Cys Ala Gln Arg Ile Ala Glu Gln Gln Lys Ala Gln Ala		
	1155	1160	1165
55	Glu Val Glu Gly Leu Gly Lys Gly Val Ala Arg Leu Ser Ala Glu Ala		
	1170	1175	1180
	Glu Lys Val Leu Ala Leu Pro Glu Pro Ser Pro Ala Ala Pro Thr Leu		
	1185	1190	1195
	Arg Ser Glu Leu Glu Leu Thr Leu Gly Lys Leu Glu Gln Val Arg Ser		
	1205	1210	1215
	Leu Ser Ala Ile Tyr Leu Glu Lys Leu Lys Thr Ile Ser Leu Val Ile		
	1220	1225	1230
60	Arg Gly Thr Gln Gly Ala Glu Glu Val Leu Arg Ala His Glu Glu Gln		
	1235	1240	1245
	Leu Lys Glu Ala Gln Ala Val Pro Ala Thr Leu Pro Glu Leu Glu Ala		
	1250	1255	1260
	Thr Lys Ala Ser Leu Lys Leu Arg Ala Gln Ala Glu Ala Gln Gln		
	1265	1270	1275
65	1280		
	Pro Thr Phe Asp Ala Leu Arg Asp Glu Leu Arg Gly Ala Gln Glu Val		
	1285	1290	1295
	Gly Glu Arg Leu Gln Gln Arg His Gly Glu Arg Asp Val Glu Val Glu		
	1300	1305	1310
70	Arg Trp Arg Glu Arg Val Ala Gln Leu Leu Glu Arg Trp Gln Ala Val		
	1315	1320	1325
	Leu Ala Gln Thr Asp Val Arg Gln Arg Glu Leu Glu Gln Leu Gly Arg		
	1330	1335	1340
75	Gln Leu Arg Tyr Tyr Arg Ser Ala Asp Pro Leu Gly Ala Trp Leu		
	1345	1350	1355
	1360		
	Gln Asp Ala Arg Arg Gln Glu Gln Ile Gln Ala Met Pro Leu Ala		
	1365	1370	1375
80	Asp Ser Gln Ala Val Arg Glu Gln Leu Arg Gln Glu Gln Ala Leu Leu		

	1380	1385	1390
	Glu Glu Ile Glu Arg His Gly Glu Lys Val Glu Glu Cys Gln Arg Phe		
	1395	1400	1405
5	Ala Lys Gln Tyr Ile Asn Ala Ile Lys Asp Tyr Glu Leu Gln Leu Val		
	1410	1415	1420
	Thr Tyr Lys Ala Gln Leu Glu Pro Val Ala Ser Pro Ala Lys Lys Pro		
	1425	1430	1435
	1440		
	Lys Val Gln Ser Gly Ser Glu Ser Val Ile Gln Glu Tyr Val Asp Leu		
	1445	1450	1455
10	Arg Thr His Tyr Ser Glu Leu Thr Thr Ser Gln Tyr Ile Lys		
	1460	1465	1470
	Phe Ile Ser Glu Thr Leu Arg Arg Met Glu Glu Glu Arg Leu Ala		
	1475	1480	1485
	Glu Gln Gln Arg Ala Glu Glu Arg Glu Arg Leu Ala Glu Val Glu Ala		
	1490	1495	1500
15	Ala Leu Glu Lys Gln Arg Gln Leu Ala Glu Ala His Ala Gln Ala Lys		
	1505	1510	1515
	1520		
	Ala Gln Ala Glu Arg Glu Ala Lys Glu Leu Gln Gln Arg Met Gln Glu		
	1525	1530	1535
	Glu Val Val Arg Arg Glu Glu Ala Ala Val Asp Ala Gln Gln Lys		
	1540	1545	1550
20	Arg Ser Ile Gln Glu Glu Leu Gln Gln Leu Arg Gln Ser Ser Glu Ala		
	1555	1560	1565
	Glu Ile Gln Ala Lys Ala Arg Gln Ala Glu Ala Ala Glu Arg Ser Arg		
	1570	1575	1580
	Leu Arg Ile Glu Glu Glu Ile Arg Val Val Arg Leu Gln Leu Glu Ala		
	1585	1590	1595
	1600		
25	Thr Glu Arg Gln Arg Gly Gly Ala Glu Gly Glu Leu Gln Ala Leu Arg		
	1605	1610	1615
	Ala Arg Ala Glu Glu Ala Glu Ala Gln Lys Arg Gln Ala Gln Glu Glu		
	1620	1625	1630
30	Ala Glu Arg Leu Arg Arg Gln Val Gln Asp Glu Ser Gln Arg Lys Arg		
	1635	1640	1645
	Gln Ala Glu Val Glu Leu Ala Ser Arg Val Lys Ala Glu Ala Glu Ala		
	1650	1655	1660
	Ala Arg Glu Lys Gln Arg Ala Leu Gln Ala Leu Glu Glu Leu Arg Leu		
	1665	1670	1675
	1680		
35	Gln Ala Glu Glu Ala Glu Arg Arg Leu Arg Gln Ala Glu Val Glu Arg		
	1685	1690	1695
	Ala Arg Gln Val Gln Val Ala Leu Glu Thr Ala Gln Arg Ser Ala Glu		
	1700	1705	1710
	Ala Glu Leu Gln Ser Lys Arg Ala Ser Phe Ala Glu Lys Thr Ala Gln		
	1715	1720	1725
40	Leu Glu Arg Ser Leu Gln Glu Glu His Val Ala Val Ala Gln Leu Arg		
	1730	1735	1740
	Glu Glu Ala Glu Arg Arg Ala Gln Gln Ala Glu Ala Glu Arg Ala		
	1745	1750	1755
	1760		
	Arg Glu Glu Ala Glu Arg Glu Leu Glu Arg Trp Gln Leu Lys Ala Asn		
	1765	1770	1775
45	Glu Ala Leu Arg Leu Arg Leu Gln Ala Glu Glu Val Ala Gln Gln Lys		
	1780	1785	1790
	Ser Leu Ala Gln Ala Glu Ala Glu Lys Gln Lys Glu Glu Ala Glu Arg		
	1795	1800	1805
	Glu Ala Arg Arg Arg Gly Lys Ala Glu Glu Gln Ala Val Arg Gln Arg		
	1810	1815	1820
50	Glu Leu Ala Glu Gln Glu Leu Glu Lys Gln Arg Gln Leu Ala Glu Gly		
	1825	1830	1835
	1840		
	Thr Ala Gln Gln Arg Leu Ala Ala Glu Gln Glu Leu Ile Arg Leu Arg		
	1845	1850	1855
	Ala Glu Thr Glu Gln Gly Glu Gln Gln Arg Gln Leu Leu Glu Glu Glu		
	1860	1865	1870
55	Ile Ala Arg Leu Gln Arg Glu Ala Ala Ala Ala Thr Gln Lys Arg Gln		
	1875	1880	1885

	Glu	Leu	Glu	Ala	Glu	Leu	Ala	Lys	Val	Arg	Ala	Glu	Met	Glu	Val	Leu	
	1890							1895					1900				
5	Leu	Ala	Ser	Lys	Ala	Arg	Ala	Glu	Glu	Ser	Arg	Ser	Thr	Ser	Glu		
	1905							1910					1915			1920	
	Lys	Ser	Lys	Gln	Arg	Leu	Glu	Ala	Glu	Gly	Arg	Phe	Arg	Glu	Leu		
						1925			1930				1935				
10	Ala	Glu	Glu	Ala	Ala	Arg	Leu	Arg	Ala	Leu	Ala	Glu	Glu	Ala	Lys	Arg	
						1940			1945				1950				
	Gln	Arg	Gln	Leu	Ala	Glu	Glu	Asp	Ala	Ala	Arg	Gln	Arg	Ala	Glu	Ala	
						1955			1960				1965				
15	Glu	Arg	Val	Leu	Ala	Glu	Lys	Leu	Ala	Ile	Gly	Glu	Ala	Thr	Arg		
						1970			1975				1980				
	Leu	Lys	Thr	Glu	Ala	Glu	Ile	Ala	Leu	Lys	Glu	Lys	Glu	Ala	Asn		
	1985					1990				1995				2000			
20	Glu	Arg	Leu	Arg	Arg	Leu	Ala	Glu	Asp	Glu	Ala	Phe	Gln	Arg	Arg	Arg	
						2005				2010				2015			
	Leu	Glu	Glu	Gln	Ala	Ala	Gln	His	Lys	Ala	Asp	Ile	Glu	Arg	Leu		
						2020				2025				2030			
25	Ala	Gln	Leu	Arg	Lys	Ala	Ser	Asp	Ser	Glu	Leu	Glu	Arg	Gln	Lys	Gly	
						2035			2040				2045				
	Leu	Val	Glu	Asp	Thr	Leu	Arg	Gln	Arg	Arg	Gln	Val	Glu	Glu	Ile		
						2050			2055				2060				
30	Leu	Ala	Leu	Lys	Ala	Ser	Phe	Glu	Lys	Ala	Ala	Gly	Lys	Ala	Glu		
	2065					2070				2075				2080			
	Leu	Glu	Leu	Glu	Leu	Gly	Arg	Ile	Arg	Ser	Asn	Ala	Glu	Asp	Thr	Leu	
						2085			2090				2095				
35	Arg	Ser	Lys	Glu	Gln	Ala	Glu	Leu	Glu	Ala	Ala	Arg	Gln	Arg	Gln	Leu	
						2100			2105				2110				
	Ala	Ala	Glu	Glu	Glu	Arg	Arg	Arg	Arg	Glu	Ala	Glu	Glu	Arg	Val	Gln	
						2115			2120				2125				
40	Lys	Ser	Leu	Ala	Ala	Glu	Glu	Glu	Ala	Ala	Arg	Gln	Arg	Lys	Ala	Ala	
						2130			2135				2140				
	Leu	Glu	Glu	Val	Glu	Arg	Leu	Lys	Ala	Lys	Val	Glu	Glu	Ala	Arg	Arg	
						2145			2150				2155			2160	
45	Leu	Arg	Glu	Arg	Ala	Glu	Gln	Glu	Ser	Ala	Arg	Gln	Leu	Gln	Leu	Ala	
						2165			2170				2175				
	Gln	Glu	Ala	Ala	Gln	Lys	Arg	Leu	Gln	Ala	Glu	Glu	Lys	Ala	His	Ala	
						2180			2185				2190				
50	Phe	Ala	Val	Gln	Gln	Lys	Glu	Gln	Glu	Leu	Gln	Gln	Thr	Leu	Gln	Gln	
						2195			2200				2205				
	Glu	Gln	Ser	Val	Leu	Asp	Gln	Leu	Arg	Gly	Glu	Ala	Glu	Ala	Ala	Arg	
						2210			2215				2220				
55	Arg	Ala	Ala	Glu	Glu	Ala	Glu	Glu	Ala	Arg	Val	Gln	Ala	Glu	Arg	Glu	
						2225			2230				2235			2240	
	Ala	Ala	Gln	Ser	Arg	Arg	Gln	Val	Glu	Ala	Glu	Arg	Leu	Lys	Gln		
						2245			2250				2255				
	Ser	Ala	Glu	Glu	Gln	Ala	Gln	Ala	Arg	Ala	Gln	Ala	Gln	Ala	Ala	Ala	
						2260			2265				2270				
	Glu	Lys	Leu	Arg	Lys	Glu	Ala	Glu	Gln	Glu	Ala	Ala	Arg	Arg	Ala	Gln	
						2275			2280				2285				
55	Ala	Glu	Gln	Ala	Ala	Leu	Arg	Gln	Lys	Gln	Ala	Ala	Asp	Ala	Glu	Met	
						2290			2295				2300				
	Glu	Lys	His	Lys	Lys	Phe	Ala	Glu	Gln	Thr	Leu	Arg	Gln	Lys	Ala	Gln	
						2305			2310				2315			2320	
	Val	Glu	Gln	Glu	Leu	Thr	Thr	Leu	Arg	Leu	Gln	Leu	Glu	Thr	Asp		
						2325			2330				2335				
	His	Gln	Lys	Asn	Leu	Leu	Asp	Glu	Glu	Leu	Gln	Arg	Leu	Lys	Ala	Glu	
						2340			2345				2350				
	Ala	Thr	Glu	Ala	Ala	Arg	Gln	Arg	Ser	Gln	Val	Glu	Glu	Leu	Phe		
						2355			2360				2365				
55	Ser	Val	Arg	Val	Gln	Met	Glu	Glu	Leu	Ser	Lys	Leu	Lys	Ala	Arg	Ile	
						2370			2375				2380				
	Glu	Ala	Glu	Asn	Arg	Ala	Ile	Leu	Arg	Asp	Lys	Asp	Asn	Thr	Gln		

2385	2390	2395	2400
Arg Phe Leu Gln Glu Glu Ala Glu Lys Met Lys Gln Val Ala Glu Glu			
2405	2410	2415	
Ala Ala Arg Leu Ser Val Ala Ala Gln Glu Ala Ala Arg Leu Arg Gln			
2420	2425	2430	
Ieu Ala Glu Glu Asp Leu Ala Gln Gln Arg Ala Leu Ala Glu Lys Met			
2435	2440	2445	
Leu Lys Glu Lys Met Gln Ala Val Gln Glu Ala Thr Arg Leu Lys Ala			
2450	2455	2460	
Glu Ala Glu Leu Leu Gln Gln Lys Glu Ieu Ala Gln Glu Gln Ala			
2465	2470	2475	2480
Arg Arg Leu Gln Glu Asp Lys Glu Gln Met Ala Gln Gln Leu Ala Glu			
2485	2490	2495	
Glu Thr Gln Gly Phe Gln Arg Thr Leu Glu Ala Glu Arg Gln Arg Gln			
2500	2505	2510	
Leu Glu Met Ser Ala Glu Ala Glu Arg Leu Lys Leu Arg Val Ala Glu			
2515	2520	2525	
Met Ser Arg Ala Gln Ala Arg Ala Glu Glu Asp Ala Gln Arg Phe Arg			
2530	2535	2540	
Lys Gln Ala Glu Glu Ile Gly Glu Lys Leu His Arg Thr Glu Leu Ala			
2545	2550	2555	2560
Thr Gln Glu Lys Val Thr Leu Val Gln Thr Leu Glu Ile Gln Arg Gln			
2565	2570	2575	
Gln Ser Asp His Asp Ala Glu Arg Leu Arg Glu Ala Ile Ala Glu Leu			
2580	2585	2590	
Glu Arg Glu Lys Glu Lys Leu Gln Gln Glu Ala Lys Leu Leu Gln Leu			
2595	2600	2605	
Lys Ser Glu Glu Met Gln Thr Val Gln Gln Glu Gln Leu Leu Gln Glu			
2610	2615	2620	
Thr Gln Ala Leu Gln Gln Ser Phe Leu Ser Glu Lys Asp Ser Leu Leu			
2625	2630	2635	2640
Gln Arg Glu Arg Phe Ile Glu Gln Glu Lys Ala Lys Leu Glu Gln Leu			
2645	2650	2655	
Phe Gln Asp Glu Val Ala Lys Ala Gln Gln Leu Arg Glu Glu Gln Gln			
2660	2665	2670	
Arg Gln Gln Gln Met Glu Gln Glu Arg Gln Arg Leu Val Ala Ser			
2675	2680	2685	
Met Glu Glu Ala Arg Arg Gln His Glu Ala Glu Glu Gly Val Arg			
2690	2695	2700	
Arg Lys Gln Glu Glu Leu Gln Gln Leu Glu Gln Gln Arg Arg Gln Gln			
2705	2710	2715	2720
Glu Glu Leu Leu Ala Glu Glu Asn Gln Arg Leu Arg Glu Gln Leu Gln			
2725	2730	2735	
Leu Leu Glu Gln His Arg Ala Ala Leu Ala His Ser Glu Glu Val			
2740	2745	2750	
Thr Ala Ser Gln Val Ala Ala Thr Lys Thr Leu Pro Asn Gly Arg Asp			
2755	2760	2765	
Ala Leu Asp Gly Pro Ala Ala Glu Ala Glu Pro Glu His Ser Phe Asp			
2770	2775	2780	
Gly Leu Arg Arg Lys Val Ser Ala Gln Arg Leu Gln Glu Ala Gly Ile			
2785	2790	2795	2800
Leu Ser Ala Glu Glu Leu Gln Arg Leu Ala Gln Gly His Thr Thr Val			
2805	2810	2815	
Asp Glu Leu Ala Arg Arg Glu Asp Val Arg His Tyr Leu Gln Gly Arg			
2820	2825	2830	
Ser Ser Ile Ala Gly Leu Leu Lys Ala Thr Asn Glu Lys Leu Ser			
2835	2840	2845	
Val Tyr Ala Ala Leu Gln Arg Gln Leu Leu Ser Pro Gly Thr Ala Leu			
2850	2855	2860	
Ile Leu Leu Glu Ala Gln Ala Ala Ser Gly Phe Leu Leu Asp Pro Val			
2865	2870	2875	2880
Arg Asn Arg Arg Leu Thr Val Asn Glu Ala Val Lys Glu Gly Val Val			
2885	2890	2895	

	Gly Pro Glu Leu His His Lys Leu Leu Ser Ala Glu Arg Ala Val Thr			
	2900	2905	2910	
	Gly Tyr Lys Asp Pro Tyr Thr Gly Gln Gln Ile Ser Leu Phe Gln Ala			
	2915	2920	2925	
5	Met Gln Lys Gly Leu Ile Val Arg Glu His Gly Ile Arg Leu Leu Glu			
	2930	2935	2940	
	Ala Gln Ile Ala Thr Gly Gly Val Ile Asp Pro Val His Ser His Arg			
	2945	2950	2955	2960
	Val Pro Val Asp Val Ala Tyr Arg Arg Gly Tyr Phe Asp Glu Glu Met			
	2965	2970	2975	
10	Asn Arg Val Leu Ala Asp Pro Ser Asp Asp Thr Lys Gly Phe Phe Asp			
	2980	2985	2990	
	Pro Asn Thr His Glu Asn Leu Thr Tyr Leu Gln Leu Leu Glu Arg Cys			
	2995	3000	3005	
	Val Glu Asp Pro Glu Thr Gly Leu Cys Leu Leu Pro Leu Thr Asp Lys			
	3010	3015	3020	
15	Ala Ala Lys Gly Gly Glu Leu Val Tyr Thr Asp Ser Glu Ala Arg Asp			
	3025	3030	3035	3040
	Val Phe Glu Lys Ala Thr Val Ser Ala Pro Phe Gly Lys Phe Gln Gly			
	3045	3050	3055	
	Lys Thr Val Thr Ile Trp Glu Ile Ile Asn Ser Glu Tyr Phe Thr Ala			
	3060	3065	3070	
20	Glu Gln Arg Arg Asp Leu Leu Arg Gln Phe Arg Thr Gly Arg Ile Thr			
	3075	3080	3085	
	Val Glu Lys Ile Ile Lys Ile Ile Thr Val Val Glu Glu Gln Glu			
	3090	3095	3100	
25	Gln Lys Gly Arg Leu Cys Phe Glu Gly Leu Arg Ser Leu Val Pro Ala			
	3105	3110	3115	3120
	Ala Glu Leu Leu Glu Ser Arg Val Ile Asp Arg Glu Leu Tyr Gln Gln			
	3125	3130	3135	
	Leu Gln Arg Gly Glu Arg Ser Val Arg Asp Val Ala Glu Val Asp Thr			
	3140	3145	3150	
30	Val Arg Arg Ala Leu Arg Gly Ala Asn Val Ile Ala Gly Val Trp Leu			
	3155	3160	3165	
	Glu Glu Ala Gly Gln Lys Leu Ser Ile Tyr Asn Ala Leu Lys Lys Asp			
	3170	3175	3180	
	Leu Leu Pro Ser Asp Met Ala Val Ala Leu Leu Glu Ala Gln Ala Gly			
	3185	3190	3195	3200
35	Thr Gly His Ile Ile Asp Pro Ala Thr Ser Ala Arg Leu Thr Val Asp			
	3205	3210	3215	
	Glu Ala Val Arg Ala Gly Leu Val Gly Pro Glu Phe His Glu Lys Leu			
	3220	3225	3230	
	Leu Ser Ala Glu Lys Ala Val Thr Gly Tyr Arg Asp Pro Tyr Thr Gly			
	3235	3240	3245	
40	Gln Ser Val Ser Leu Phe Gln Ala Leu Lys Lys Gly Leu Ile Pro Arg			
	3250	3255	3260	
	Glu Gln Gly Leu Arg Leu Leu Asp Ala Gln Leu Ser Thr Gly Gly Ile			
	3265	3270	3275	3280
	Val Asp Pro Ser Lys Ser His Arg Val Pro Leu Asp Val Ala Cys Ala			
	3285	3290	3295	
45	Arg Gly Cys Leu Asp Glu Glu Thr Ser Arg Ala Leu Ser Ala Pro Arg			
	3300	3305	3310	
	Ala Asp Ala Lys Ala Tyr Ser Asp Pro Ser Thr Gly Glu Pro Ala Thr			
	3315	3320	3325	
50	Tyr Gly Glu Leu Gln Gln Arg Cys Arg Pro Asp Gln Leu Thr Gly Leu			
	3330	3335	3340	
	Ser Leu Leu Pro Leu Ser Glu Lys Ala Ala Arg Ala Arg Gln Glu Glu			
	3345	3350	3355	3360
	Leu Tyr Ser Glu Leu Gln Ala Arg Glu Thr Phe Glu Lys Thr Pro Val			
	3365	3370	3375	
55	Glu Val Pro Val Gly Gly Phe Lys Gly Arg Thr Val Thr Val Trp Glu			
	3380	3385	3390	
	Leu Ile Ser Ser Glu Tyr Phe Thr Ala Glu Gln Arg Gln Glu Leu Leu			

	3395	3400	3405
	Arg Gln Phe Arg Thr Gly Lys Val Thr Val Glu Lys Val Ile Lys Ile		
	3410	3415	3420
5	Leu Ile Thr Ile Val Glu Val Glu Thr Leu Arg Gln Glu Arg Leu		
	3425	3430	3435
	Ser Phe Ser Gly Leu Arg Ala Pro Val Pro Ala Ser Glu Leu Leu Ala		
	3445	3450	3455
	Ser Gly Val Leu Ser Arg Ala Gln Phe Glu Gln Leu Lys Asp Gly Lys		
	3460	3465	3470
10	Thr Thr Val Lys Asp Leu Ser Glu Leu Gly Ser Val Arg Thr Leu Leu		
	3475	3480	3485
	Gln Gly Ser Gly Cys Leu Ala Gly Ile Tyr Leu Glu Asp Thr Lys Glu		
	3490	3495	3500
	Lys Val Ser Ile Tyr Glu Ala Met Arg Arg Gly Leu Leu Arg Ala Thr		
	3505	3510	3515
15	Thr Ala Ala Leu Leu Glu Ala Gln Ala Ala Thr Gly Phe Leu Val		
	3525	3530	3535
	Asp Pro Val Arg Asn Gln Arg Leu Tyr Val His Glu Ala Val Lys Ala		
	3540	3545	3550
	Gly Val Val Gly Pro Glu Leu His Glu Gln Leu Leu Ser Ala Glu Lys		
	3555	3560	3565
20	Ala Val Thr Gly Tyr Arg Asp Pro Tyr Ser Gly Ser Thr Ile Ser Leu		
	3570	3575	3580
	Phe Gln Ala Met Gln Lys Gly Leu Val Leu Arg Gln His Gly Ile Arg		
	3585	3590	3595
	Leu Leu Glu Ala Gln Ile Ala Thr Gly Gly Ile Ile Asp Pro Val His		
	3605	3610	3615
25	Ser His Arg Val Pro Val Asp Val Ala Tyr Gln Arg Gly Tyr Phe Ser		
	3620	3625	3630
	Glu Glu Met Asn Arg Val Leu Ala Asp Pro Ser Asp Asp Thr Lys Gly		
	3635	3640	3645
	Phe Phe Asp Pro Asn Thr His Glu Asn Leu Thr Tyr Arg Gln Leu Leu		
30	3650	3655	3660
	Glu Arg Cys Val Glu Asp Pro Glu Thr Gly Leu Arg Leu Leu Pro Leu		
	3665	3670	3675
	Lys Gly Ala Glu Lys Ala Glu Val Val Glu Thr Thr Gln Val Tyr Thr		
	3685	3690	3695
35	Glu Glu Glu Thr Arg Arg Ala Phe Glu Glu Thr Gln Ile Asp Ile Pro		
	3700	3705	3710
	Gly Gly Gly Ser His Gly Gly Ser Thr Met Ser Leu Trp Glu Val Met		
	3715	3720	3725
	Gln Ser Asp Leu Ile Pro Glu Glu Gln Arg Ala Gln Leu Met Ala Asp		
	3730	3735	3740
40	Phe Gln Ala Gly Arg Val Thr Lys Glu Arg Met Ile Ile Ile Ile Ile		
	3745	3750	3755
	Glu Ile Ile Glu Lys Thr Glu Ile Ile Arg Gln Gln Gly Leu Ala Ser		
	3765	3770	3775
	Tyr Asp Tyr Val Arg Arg Arg Leu Thr Ala Glu Asp Leu Phe Glu Ala		
	3780	3785	3790
45	Arg Ile Ile Ser Leu Glu Thr Tyr Asn Leu Leu Arg Glu Gly Thr Arg		
	3795	3800	3805
	Ser Leu Arg Glu Ala Leu Glu Ala Glu Ser Ala Trp Cys Tyr Leu Tyr		
	3810	3815	3820
	Gly Thr Gly Ser Val Ala Gly Val Tyr Leu Pro Gly Ser Arg Gln Thr		
50	3825	3830	3835
	Leu Ser Ile Tyr Gln Ala Leu Lys Lys Gly Leu Leu Ser Ala Glu Val		
	3845	3850	3855
	Ala Arg Leu Leu Glu Ala Gln Ala Ala Thr Gly Phe Leu Leu Asp		
	3860	3865	3870
55	Pro Val Lys Gly Glu Arg Leu Thr Val Asp Glu Ala Val Arg Lys Gly		
	3875	3880	3885
	Ile Val Gly Pro Glu Leu His Asp Arg Leu Leu Ser Ala Glu Arg Ala		
	3890	3895	3900

	Val Thr Gly Tyr Arg Asp Pro Tyr Thr Glu Gln Thr Ile Ser Leu Phe			
	3905	3910	3915	3920
	Gln Ala Met Lys Lys Glu Leu Ile Pro Thr Glu Glu Ala Leu Arg Leu			
	3925	3930	3935	
5	Leu Asp Ala Gln Leu Ala Thr Gly Gly Ile Val Asp Pro Arg Leu Gly			
	3940	3945	3950	
	Phe His Leu Pro Leu Glu Val Ala Tyr Gln Arg Gly Tyr Leu Asn Lys			
	3955	3960	3965	
10	Asp Thr His Asp Gln Leu Ser Glu Pro Ser Glu Val Arg Ser Tyr Val			
	3970	3975	3980	
	Asp Pro Ser Thr Asp Glu Arg Leu Ser Tyr Thr Gln Leu Leu Arg Arg			
	3985	3990	3995	4000
	Cys Arg Arg Asp Asp Gly Thr Gly Gln Leu Leu Leu Pro Leu Ser Asp			
	4005	4010	4015	
15	Ala Arg Lys Leu Thr Phe Arg Gly Leu Arg Lys Gln Ile Thr Met Glu			
	4020	4025	4030	
	Glu Leu Val Arg Ser Gln Val Met Asp Glu Ala Thr Ala Leu Gln Leu			
	4035	4040	4045	
	Arg Glu Gly Leu Thr Ser Ile Glu Glu Val Thr Lys Asn Leu Gln Lys			
	4050	4055	4060	
20	Phe Leu Glu Gly Thr Ser Cys Ile Ala Gly Val Phe Val Asp Ala Thr			
	4065	4070	4075	4080
	Lys Glu Arg Leu Ser Val Tyr Gln Ala Met Lys Lys Gly Ile Ile Arg			
	4085	4090	4095	
	Pro Gly Thr Ala Phe Glu Leu Leu Glu Ala Gln Ala Ala Thr Gly Tyr			
	4100	4105	4110	
25	Val Ile Asp Pro Ile Lys Gly Leu Lys Leu Thr Val Glu Glu Ala Val			
	4115	4120	4125	
	Arg Met Gly Ile Val Gly Pro Glu Phe Lys Asp Lys Leu Leu Ser Ala			
	4130	4135	4140	
	Glu Arg Ala Val Thr Gly Tyr Lys Asp Pro Tyr Ser Gly Lys Leu Ile			
	4145	4150	4155	4160
30	Ser Leu Phe Gln Ala Met Lys Lys Gly Leu Ile Leu Lys Asp His Gly			
	4165	4170	4175	
	Ile Arg Leu Leu Glu Ala Gln Ile Ala Thr Gly Gly Ile Ile Asp Pro			
	4180	4185	4190	
	Glu Glu Ser His Arg Leu Pro Val Glu Val Ala Tyr Lys Arg Gly Leu			
	4195	4200	4205	
35	Phe Asp Glu Glu Met Asn Glu Ile Leu Thr Asp Pro Ser Asp Asp Thr			
	4210	4215	4220	
	Lys Gly Phe Phe Asp Pro Asn Thr Glu Glu Asn Leu Thr Tyr Leu Gln			
	4225	4230	4235	4240
	Leu Met Glu Arg Cys Ile Thr Asp Pro Gln Thr Gly Leu Cys Leu Leu			
	4245	4250	4255	
40	Pro Leu Lys Glu Lys Lys Arg Glu Arg Lys Thr Ser Ser Lys Ser Ser			
	4260	4265	4270	
	Val Arg Lys Arg Arg Val Val Ile Val Asp Pro Glu Thr Gly Lys Glu			
	4275	4280	4285	
	Met Ser Val Tyr Glu Ala Tyr Arg Lys Gly Leu Ile Asp His Gln Thr			
45	4290	4295	4300	
	Tyr Leu Glu Leu Ser Glu Gln Glu Cys Glu Trp Glu Glu Ile Thr Ile			
	4305	4310	4315	4320
	Ser Ser Ser Asp Gly Val Val Lys Ser Met Ile Ile Asp Arg Arg Ser			
	4325	4330	4335	
50	Gly Arg Gln Tyr Asp Ile Asp Asp Ala Ile Ala Lys Asn Leu Ile Asp			
	4340	4345	4350	
	Arg Ser Ala Leu Asp Gln Tyr Arg Ala Gly Thr Leu Ser Ile Thr Glu			
	4355	4360	4365	
	Phe Ala Asp Met Leu Ser Gly Asn Ala Gly Gly Phe Arg Ser Arg Ser			
	4370	4375	4380	
55	Ser Ser Val Gly Ser Ser Ser Ser Tyr Pro Ile Ser Pro Ala Val Ser			
	4385	4390	4395	4400
	Arg Thr Gln Leu Ala Ser Trp Ser Asp Pro Thr Glu Glu Thr Gly Pro			

	4405	4410	4415
	Val Ala Gly Ile Leu Asp Thr Glu Thr Leu Glu Lys Val Ser Ile Thr		
	4420	4425	4430
5	Glu Ala Met His Arg Asn Leu Val Asp Asn Ile Thr Gly Gln Arg Leu		
	4435	4440	4445
	Ile Glu Ala Gln Ala Cys Thr Gly Gly Ile Ile Asp Pro Ser Thr Gly		
	4450	4455	4460
	Glu Arg Phe Pro Val Thr Asp Ala Val Asn Lys Gly Leu Val Asp Lys		
	4465	4470	4475
10	Ile Met Val Asp Arg Ile Asn Leu Ala Gln Lys Ala Phe Cys Gly Phe		
	4485	4490	4495
	Glu Asp Pro Arg Thr Lys Thr Lys Met Ser Ala Ala Gln Ala Leu Lys		
	4500	4505	4510
	Lys Gly Trp Leu Tyr Tyr Glu Ala Gly Gln Arg Phe Leu Glu Val Gln		
	4515	4520	4525
15	Tyr Leu Thr Gly Gly Leu Ile Glu Pro Asp Thr Pro Gly Arg Val Pro		
	4530	4535	4540
	Ile Asp Glu Ala Leu Gln Arg Gly Thr Val Asp Ala Arg Thr Ala Gln		
	4545	4550	4555
	Lys Leu Arg Asp Val Gly Ala Tyr Ser Lys Tyr Leu Thr Cys Pro Lys		
	4565	4570	4575
20	Thr Lys Leu Lys Ile Ser Tyr Lys Asp Ala Leu Asp Arg Ser Met Val		
	4580	4585	4590
	Glu Glu Gly Thr Gly Leu Arg Leu Leu Glu Ala Ala Gln Ser Thr		
	4595	4600	4605
	Lys Gly Tyr Tyr Ser Pro Tyr Ser Val Ser Gly Ser Gly Ser Thr Ala		
25	4610	4615	4620
	Gly Ser Arg Thr Gly Ser Arg Thr Gly Ser Arg Ala Gly Ser Arg Arg		
	4625	4630	4635
	Gly Ser Phe Asp Ala Thr Gly Ser Gly Phe Ser Met Thr Phe Ser Ser		
	4645	4650	4655
30	Ser Ser Tyr Ser Ser Ser Gly Tyr Gly Arg Arg Tyr Ala Ser Gly Ser		
	4660	4665	4670
	Ser Ala Ser Leu Gly Gly Pro Glu Ser Ala Val Ala		
	4675	4680	

35 <210> 25
 <211> 36
 <212> PRT
 <213> Artificial Sequence

40 <220>
 <223> Synthetic peptide
 <400> 25
 Tyr Pro Ile Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu
 1 5 10 15
 Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr
 20 25 30
 Arg Gln Arg Tyr
 35

50 <210> 26
 <211> 9
 <212> PRT
 <213> Artificial Sequence

55 <220>
 <223> Synthetic peptide
 <400> 26

Thr Ser His His His His His His Cys
1 5

5


Claims

1. A pharmaceutical composition comprising a ligand comprising a first portion comprising a plectin-1 binding moiety coupled to a second portion comprising a detectable moiety or a therapeutic agent for the diagnosis or treatment of cancer.
2. Use of a ligand comprising a first portion comprising a plectin-1 binding moiety coupled to a second portion comprising a detectable moiety or a therapeutic agent in the manufacture of a medicament for the diagnosis or treatment of cancer.
3. The composition or use of claim 1 or claim 2, wherein the detectable moiety is selected from the group consisting of a radioactive isotope, a magnetic compound, an x-ray absorber, a chemical compound, a biological tag, and a fluorescent molecule.
4. The composition or use of claim 1 or claim 2, wherein the therapeutic agent is a cytotoxic moiety or an immunomodulatory moiety.
5. The composition or use of claim 1 or claim 2, further comprising a linker between the first portion and the second portion.
6. The composition or use of claim 5, wherein the linker is a flexible amino acid sequence.
7. The composition or use of claim 5, wherein the linker is a photolinker.
8. The composition or use of claim 1 or 2 wherein the second portion further comprises a physiologically inert nanoparticle.
9. The composition or use of claim 8, wherein the nanoparticle is magnetic, fluorescent, or radioactive.
10. The composition or use of claim 1 or 2, wherein the second portion comprises a fluorochrome.
11. The composition or use of claim 1 or 2, wherein the ligand is a plectin-1 specific antibody.
12. The composition or use of claim 11, wherein the antibody is a monoclonal antibody.
13. The composition or use of claim 1 or 2, wherein the cancer is a pancreatic ductal adenocarcinoma.

45

50

55



FIGURES 1A-C







FIGURE 2



FIGURES 3A-C





FIGURES 4A-F




FIGURES 5A-D

FIGURES 6A-D





FIGURES 7A-C



FIGURES 8A-C





FIGURE 9

A Kras p53L/L Mouse



B Kras p53L/L Mouse



C Wild Type Mouse



FIGURES 10A-C

Figure 11A

SEQ ID NO:24

MVAGMLMPRD	QLRAIYEVLF	REGVMVAKKD	RRPRSLHPHV	PGVTNLQVMR	50
AMASLRARGL	VRETFAWCHF	FWYLTNEGIA	HLRQYLHLPP	EIVAASLQRV	100
RRPVAMVMPA	RRTPHVQAVQ	GPLGSPPKRG	PLPTEEQRLY	RRKELEEVSP	150
ETPVVPATTQ	RTLARPGPEP	APATDERDRV	QKKTFTKWN	KHLIKAQRHI	200
SDLYEDLRDG	HNLISLLEV	SGDSLPREKG	RMRFHKLQNV	QIALDYLRR	250
QVKLVNIRND	DIADGNPKLT	LGLIWTIILH	FQISDIQVSG	QSEDMTAKEK	300
LLLWSQRMVE	GYQGLRCDFN	TSSWRDGRFLP	NAIIHRHKPL	LIDMNKVYRQ	350
TNLENLDQAF	SVAERDLGVT	RLLDPEDVDV	PQPDEKSIIT	YVSSLYDAMP	400
RVPDVQDGVR	ANELQLRWQE	YRELVLLLQ	WMRRHHTAAFE	ERRFPSSFEE	450
IEILWSQPLK	FKEMELPAKE	ADKNRSKG	YQQSLEGAVQAG	QLKVPPGYHP	500
LDVEKEWGKL	HVAILEREKQ	LRSEFERLEC	LQRIVTKLQM	EAGLCEEQLN	550
QADALLQSDV	RLLAAGKVPQ	RAGEVERDLD	KADSMIRLLF	NDVQTLKDGR	600
HPQGEQMYRR	VYRLHERLVA	IRTEYNLRLK	AGVAAPATQV	AQVTLQSVQR	650
RPELEDSTLR	YLQDLLAWVE	ENQHRVGDAE	WGVDLPSVEA	QLGSHRGLHQ	700
SIEEFRAKIE	RARSDEGQLS	PATRGAYRDC	LGRLDLQYAK	LLNNSSKARLR	750
SLESLHSFVA	AATKELMWLN	EKEEEEVGFD	WSDRNTNMTA	KKESYSALMR	800
ELELKEKKIK	ELQNAGDRLL	REDHPARPTV	ESFQAAALQTO	WSWMLQLCCC	850
IEAHLKENAA	YFQFFSDVRE	AEGQLQKLQE	ALRRKYSYCDR	SATVTRLEDL	900
LQDAQDEKEQ	LNEYKGHLG	LAKRAKAVVQ	LKPRRHPAHPM	RGRPLPLAVC	950
DYKQEVETVH	KGDECQLVGP	AQPSHWKVLS	SSGSEAAVPS	VCFLVPPPNQ	1000
EAQEAVTRLE	AQHOALVTLW	HQLHVDMKSL	LAQOSLRRDV	QLIRWSLAT	1050
FRTLKPEEQR	QALHSLELHY	QAFLRDSQDA	GGFGPEDRJM	AEREYGSCH	1100
HYQQQLLQSLE	QGAQEEESRCQ	RCISELKDIR	LQLEACETRT	VHRLRLPLDK	1150
EPARECAQRI	AEQQKAQAEV	EGLGKGVARL	SAEAEKVLA	PEPSPAAPTL	1200
RSELELTGK	LEQVRSLSAI	YLEKLKTISL	VIRGTQGAAE	VLRAHEEQLK	1250
EAQAVPATLP	ELEATKASLK	KLRAQAEAQ	PTFDALRDEL	RGAQEVGRL	1300
QQRHGERDVE	VERWRERVAQ	LLERWQAVLA	QTDVRQRELE	QLGRQLRYYR	1350
ESADPLGAWL	QDARRRQEIQ	QAMPLADSQA	VREQLRQEQA	LLEBIEHRGE	1400
KVEECQRFK	QYINAIDYE	LQLVTYKAQL	EPVASPAKKP	KVQSGSESVI	1450
QEYVDLRTHY	SELTTLTSQY	IKPISETLRR	MEEBEERLAEQ	QRAEERERLA	1500
EVEAALEKQR	QLAEAAHQAK	AQAEREAKEL	QORMQEEVVR	REEAADVDAQQ	1550
QKRSIQEELQ	QLRQSSEAEI	QAKARQAEAA	ERSRLRIEEE	IRVURLQLEA	1600
TERQRGGAEG	ELQALRARA	EAAEAKRQAO	EEAERLRRQV	QDESQRKRQA	1650
EVELASRVKA	EAAEAREKQR	ALQALEELRL	QAEAEERRRL	QAEVERARQV	1700
QVALETAQRS	AEAEQLSKRA	SFAEKTAQLE	RSLQEEHVAV	AQLREEAERR	1750
AQQQAEAAERA	REEAERELE	WQLKANEALR	LRLQAEVEAQ	QKSLAQAEAE	1800
KQKEEAAREA	RRRGKAEEQA	VRQRELAEQE	LEKQRQLAEG	TAQQLAAEQ	1850
BLIRLRAETE	QGEQQRQLLE	BELARLQREA	AAATQKRQEL	EAELAKVRAE	1900
MEVLLASKAR	AEESRSTSE	KSQQLRLEAA	GRFRELABEA	ARLRLALABEA	1950
KRQRQLAEE	AARQRAEAER	VLAEKLAAG	EATRLKTEAE	IALKEKEAEN	2000
ERLRLRAEDE	AFQRRRLEEQ	AAQHKADIEE	RLAQLRKASD	SELERQKGLV	2050
EDTLRQRQV	EEEILALKAS	FEKAAAGKAE	LELELGRIRS	NAEDTLRSKE	2100
QAELEAARQR	QLAEEBERR	REAEERVQKS	LAAEEEARQ	RKAALEEVER	2150
LKANVEEARR	LRERAEQESA	RQLQLAQEAA	QKRLQAEKA	HAFAVQQKEQ	2200
ELQOTLQQEQ	SVLDQLRGEA	EAARRAAEEA	EEARVQAERE	AAQARROQVEE	2250
AERLKQSAEB	QAQARAQQAQ	AAEKLKEA	QEAARRAQAE	QAALRQKQAA	2300
DAEMEKHKKF	AEQTLRQKAQ	VEQELTTLRL	QLEETDHQKN	LLDEELQRLK	2350
AEATEAARQR	SQVEEELFSV	RVQMEELSKL	KARIEAENRA	LILRDKDNTQ	2400
RFLQEEAEM	KQVAEEAARL	SVAAQEAA	RQLAEEDLAQ	QRALAEKMLK	2450
EKMQAVQEAT	RLKAEAEELLQ	QOKELAQEQ	RRQLEQDEQM	AQQLAEETQG	2500
FQRTLEAERQ	RQLEMSAEAE	RLKLRVAEMS	RAQARAEEDA	QRFRKOABEI	2550
GEKLHRTELA	TQEKTBLVQT	LEIQRQQSDH	DAERLREAIA	ELEREKEKLQ	2600
QEAKLQLKS	EEMOTVQOEQ	LLQETQALQQ	SFLSEKDSL	QRERFIEQEK	2650
AKLEQLFQDE	VAKAQQLREE	OQRQQQQMEQ	ERQLVASME	EARRRQHEAE	2700

FIGURE 11B

EGVRRKQEEL QOLEQQRQQ EELLAENQR LREQQLLEE QHRAALAHSE 2750
 EVTASQVAAT KTLPGNRRDAL DGPAEEAEPE HSFDGLRRKV SAQRLQEAGI 2800
 LSAEEELQRLA QGHTTVDELA RREDVRHYLQ GRSSIAGLLL KATNEKLSVY 2850
 AALQRQLLSP GTALILLEAQ AASGPPLDPV RNRRRTVNEA VKEGVVGPEL 2900
 HHKLLSAERA VTGYKDPYTG QQISLFQAMQ KGLIVREHGI RLLEAQIATG 2950
 CVIDPVHSHR VPVDVAYRRG YFDEEMNRVL ADPSDDTKGF FDPNTHENLT 3000
 YLQLLERCVE DPETGLCLLP LTDKAAGGGE LVYTDSEARD VFEKATVSAP 3050
 FGKFQGKTVT IWEIINSEYF TAEQRDLLR QFRTGRITVE KIIKIIITVV 3100
 EEQEQQKGRRLC FEGLRSLVPA AELLESRVID RELYQQQLQRG ERSVRDVAEV 3150
 DTVRALRGA NVIAGVWLEE AGQKLSIYNA LKKDLLPSDM AVALLEAQAG 3200
 TGHIIDPATS ARLTVDDEAVR AGLVGPEFHE KLLSAEKAVT GYRDPYTQGS 3250
 VSLFQALKKG LIPREQGLRL LDAQLSTGGI VDPSKSHRVP LDVACARGCL 3300
 DEETSRLSA PRADAKAYSD PSTGEPATYG ELQQRCPDQ LTGLSLLPLS 3350
 EKAARARQEE LYSELQARET FEKTPVEVPV GGFKGRTVTW WELISSEYFT 3400
 AEQRQELLRQ FRTGKVTVEK VIKILITIVE EVETLRQERL SFSGLRAPVP 3450
 ASELLASGVL SRAQFEQLKD GKTTVKDLSE LGSVRTLLQG SGCLAGIYLE 3500
 DTKEKVSIYE AMRRGLLRAT TAALLLEAQ AATGFLVDPVR NQRLYVHEAV 3550
 KAGVVGPELH EQLLSAEKAV TGYRDPYSGS TISLFQAMQK GLVLRQHGIR 3600
 LLEAQIATGG IIDPVHSHRV PVDVAYQRGY FSEEMNRVL A DPSDDTKGFF 3650
 DPNTHENLTY RQLLERCVED PETGLRLLPL KGAEKAEVVE TTQVYTEEET 3700
 RRAFEETQID IPGGGSHGGS TMSLWEVMQS DLIPEEQRAQ LMADFQAGRV 3750
 TKERMIIII EIEKTEIIR QQGLASYDYV RRRRLTAEDLF EARIISLETY 3800
 NLLREGTRSL REALEAESAW CYLYGTGSVA GVYLPGRQQT LSIYQALKKG 3850
 LLSAEVARLL LEAQAAATGFL LDPVKGERLT VDEAVRKGLV GPELHDRLLS 3900
 AERAVTGYRD PYTEQTISLP QAMKKELIPT EEAIRLLDAQ LATGGIVDPR 3950
 LGPHLPLEVA YQRGYLNKDT HDQLSEPSEV RSYVDPSTDE RLSYTQLRR 4000
 CRRDDGTGQL LLPLSDARKL TFRGLRKQIT MEELVRSQVM DEATALQLRE 4050
 GLTSIEEVTK NLQKFLEGTS CIAGVFVDAK KERLSVYQAM KKGIIIRPGTA 4100
 FELLEAQAAAT GYVIDPIKGL KLTVEAVRM GIVGPEFKDK LLSAERAVTG 4150
 YKDPYSGKLI SLFQAMKKGL ILKDHGIRLL EAQIATGGII DPEESHRLPV 4200
 EVAYKRGFLD EEMNEILTD P SDDTKGFFDP NTEENLTYLQ LMERCITDPQ 4250
 TGLCLLPLKE KKRERKRTSSK SSVRKRRVVI VPETGKEMS VYEAYRKGLI 4300
 DHQTYLELSE QCEWEEBITI SSSDGVVKSM IIDRRSGRQY DIDDIAIKNL 4350
 IDRSLADQYR AGTLSITEFA DMLSGNAGGP RSRSSSVGSS SSYPISPAS 4400
 RTQLASWSDP TEETGPVAGI LDTETLEKVS ITEAMHRNLV DNITGQLLE 4450
 AQACTGGIID PSTGERFPVT DAVNKGVLVK IMVDRINLAQ KAFCGFEDPR 4500
 TKTMSAAQA LKKGWLYYE A GQRFLEVQYL TCGGLIEPDTP GRVPLDEALQ 4550
 RGTVDARTAQ KLRDVGAYSK YLTCPKTKLK ISYKDALDRS MVEEGTGLRL 4600
 LEAAAQSTKG YYSPYSVSGS GSTAGSRTGS RTGSRAGSRR GSFDATGSGF 4650
 SMTFSSSSYS SSGYGRYYAS GSSASLGGPE SAVA 4684



EUROPEAN SEARCH REPORT

Application Number

EP 14 19 0585

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10 X	WO 02/062838 A1 (PETER MACCALLUM CANCER INST [AU]; ST VINCENTS INST MED RES [AU]; BOWTE) 15 August 2002 (2002-08-15) * p. 10, lines 23-32, Fig. 34, p. 28, lines 11-23 * -----	1-3,5,6	INV. C07K7/06 A61K49/14 A61K38/08 G01N33/574 G01N33/68 C07K16/18
15 X	FONTAO LIONEL ET AL: "The interaction of plectin with actin: Evidence for cross-linking of actin filaments by dimerization of the actin-binding domain of plectin", JOURNAL OF CELL SCIENCE, vol. 114, no. 11, June 2001 (2001-06), pages 2065-2076, XP002683983, ISSN: 0021-9533 * the whole document *	1-3,5,6,10	
20 Y,D	WO 03/030725 A2 (UNIV JOHNS HOPKINS [US]; HRUBAN RALPH H [US]; ARGANI PEDRAM [US]; IACO) 17 April 2003 (2003-04-17) * p. 14, line 6 - p. 44, line 20, claims 1-68 *	1-13	
25	-----		TECHNICAL FIELDS SEARCHED (IPC)
30 Y	LEE, K. Y., LIU, Y.-H., HO, C.-C., PEI, R.-J., YEH, K.-T., CHENG, C.-C., LAI, Y.-S.: "An Early Evaluation Of Malignant Tendency With Plectin Expression In Human Colorectal Adenoma And Adenocarcinoma", JOURNAL OF MEDICINE, vol. 35, no. 1-6, 2004, pages 141-149, XP009163041, * the whole document *	1-13	A61K C07K G01N
35	-----		
40	-----	-/-	
45			
2	The present search report has been drawn up for all claims		
50	Place of search Munich	Date of completion of the search 16 March 2015	Examiner R. von Eggelkraut-G.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



EUROPEAN SEARCH REPORT

Application Number

EP 14 19 0585

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	Y CHIUNG-CHI CHENG ET AL: "The influence of plectin deficiency on stability of cytokeratin18 in hepatocellular carcinoma", JOURNAL OF MOLECULAR HISTOLOGY, KLUWER ACADEMIC PUBLISHERS, D0, vol. 39, no. 2, 24 November 2007 (2007-11-24), pages 209-216, XP019573299, ISSN: 1567-2387 * the whole document * -----	1-13	
15	Y,D US 5 208 020 A (CHARI RAVI J [US] ET AL) 4 May 1993 (1993-05-04) * col. 3, line 60 - col. 6, line 29, col. 10, line 64 - col. 14, line 25 *	1-13	
20	Y,D US 5 968 479 A (ITO SUSUMU [JP] ET AL) 19 October 1999 (1999-10-19) * col. 2, line 35 - col. 12, line 36 *	1-13	
25	Y WO 2007/109809 A1 (INVITROGEN CORP [US]; MAURO JOHN MATTHEW [US]; NYHUS JULIE KAY [US]; S) 27 September 2007 (2007-09-27) * p. 4, line 34 - p. 21, line 3 *	1-13	
30	X,P KAM LEUNG: "Lys-Thr-Leu-Leu-Pro-Thr-Pro-cross-linked iron oxide-Cy5.5 [PTP-CLIO-Cy5.5]", INTERNET CITATION, 3 September 2008 (2008-09-03), XP008143627, Retrieved from the Internet: URL: http://www.ncbi.nlm.nih.gov/bookshelf/br.fcgi?book=micad&part=KTLLPTP-CLIO-Cy55 [retrieved on 2011-09-29] * the whole document *	1-13	
35		1-13	
40		1-13	
45		1-13	
50	2 The present search report has been drawn up for all claims	1-13	
55	Place of search Munich	Date of completion of the search 16 March 2015	Examiner R. von Eggelkraut-G.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			



EUROPEAN SEARCH REPORT

Application Number

EP 14 19 0585

5

10

15

20

25

30

35

40

45

EPO FORM 1503 03/82 (P04C01)

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X, P	KIMBERLY A KELLY ET AL: "Targeted nanoparticles for imaging incipient pancreatic ductal adenocarcinoma", PLOS MEDICINE, PUBLIC LIBRARY OF SCIENCE, US, vol. 5, no. 4, 15 April 2008 (2008-04-15), pages 657-668, XP008143633, ISSN: 1549-1277 * the whole document * -----	1-13	
X, P	D. BAUSCH ET AL: "Plectin-1 as a Novel Biomarker for Pancreatic Cancer", CLINICAL CANCER RESEARCH, vol. 17, no. 2, 15 January 2011 (2011-01-15), pages 302-309, XP055038259, ISSN: 1078-0432, DOI: 10.1158/1078-0432.CCR-10-0999 * the whole document * -----	1-13	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (IPC)
2	Place of search Munich	Date of completion of the search 16 March 2015	Examiner R. von Eggelkraut-G.
50	CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
55			

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 14 19 0585

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

16-03-2015

10

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 02062838	A1	15-08-2002	US	2005079503 A1		14-04-2005
			WO	02062838 A1		15-08-2002
WO 03030725	A2	17-04-2003	AU	2002342053 A1		22-04-2003
			US	2003180747 A1		25-09-2003
			WO	03030725 A2		17-04-2003
US 5208020	A	04-05-1993	US	5208020 A		04-05-1993
			US	5416064 A		16-05-1995
US 5968479	A	19-10-1999	AT	264694 T		15-05-2004
			AU	4497096 A		21-08-1996
			CA	2211470 A1		08-08-1996
			CN	1180316 A		29-04-1998
			DE	69632244 D1		27-05-2004
			DE	69632244 T2		14-04-2005
			EP	0800831 A1		15-10-1997
			KR	100397020 B1		29-04-2004
			NO	973484 A		30-09-1997
			US	5968479 A		19-10-1999
			WO	9623525 A1		08-08-1996
WO 2007109809	A1	27-09-2007	EP	2005187 A1		24-12-2008
			JP	2009531332 A		03-09-2009
			US	2009311193 A1		17-12-2009
			US	2012253160 A1		04-10-2012
			WO	2007109809 A1		27-09-2007

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61044818 A [0001]
- US 20050260639 A [0131]
- US 20050095611 A [0131]
- US 20030180747 A [0131] [0138] [0157]
- US 5574010 A [0131]
- WO 2004056314 A [0131]
- US 5604203 A [0131]
- US 5696093 A [0131]
- US 6046167 A [0131]
- US 5492814 A [0153]
- US 5968479 A, Ito [0158]
- WO 9847538 A [0158]
- US 20060275775 A [0158]
- WO 2009036092 A [0159] [0169]
- US 5827934 A [0161]
- US 5208020 A [0164]
- US 5475092 A [0164]
- US 5585499 A [0164]
- US 5846545 A [0164]
- WO 8403508 A [0164]
- WO 8503508 A [0164]
- US 6468798 B [0174]
- US 4522811 A [0176]
- US 5192746 A, Lobl [0191] [0205]
- US 5169862 A, Burke, Jr. [0191] [0205]
- US 5539085 A, Bischoff [0191] [0205]
- US 5576423 A, Aversa [0191] [0205]
- US 5051448 A, Shashoua [0191] [0205]
- US 5559103 A, Gaeta [0191] [0205]
- US 4939239 A [0201]

Non-patent literature cited in the description

- LI et al. *Lancet*, 2004, vol. 363, 1049-1057 [0004]
- YEO et al. *Ann Surg*, 1995, vol. 222, 588-592 [0005]
- BRENTNALL et al. *Ann. Intern. Med.*, 1999, vol. 131, 247-255 [0005]
- CANTO et al. *Clin. Gastroenterol. Hepatol.*, 2004, vol. 2, 606-621 [0005]
- GOGGINS. *J. Clin. Oncol.*, 2005, vol. 23, 4524-4531 [0006]
- PELAEZ-LUNA et al. *Am J Gastroenterol*, 2007, vol. 102, 2157-2163 [0006]
- Enzyme-Linked Immunosorbent Assay (ELISA). CROWTHER et al. Molecular Biomethods Handbook. Humana Press, Inc, 1998, 595-617 [0114]
- Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1988 [0114]
- Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1994 [0114]
- NEWTON et al. *Neoplasia*, 2006, vol. 8, 772-780 [0114] [0147] [0229]
- GOGGINS. *J Clin Oncol*, 2005, vol. 23, 4524-4531 [0127]
- MISEK et al. *Methods Mol Med*, 2005, vol. 103, 175-187 [0127]
- BLOOMSTON et al. *Cancer Res*, 2006, vol. 66, 2592-2599 [0127]
- YATES et al. *Anal Chem*, 1995, vol. 67, 1426-1436 [0127] [0223]
- JOYCE et al. *Cancer Cell*, 2003, vol. 4, 393-403 [0127]
- FOLLI et al. *Cancer Res*, 1994, vol. 54, 2643-2649 [0128]
- NERI et al. *Nat Biotechnol*, 1997, vol. 15, 1271-1275 [0128]
- KELLY et al. *Circ Res*, 2005, vol. 96, 327-336 [0128] [0216] [0218] [0235]
- BARDEESY et al. *Proc Natl Acad Sci U.S.A.*, 2006, vol. 103, 5947-5952 [0129]
- AGUIRRE et al. *Genes Dev*, 2003, vol. 17, 3112-3126 [0129] [0145] [0215]
- HANSEL et al. *Annu Rev Genomics Hum Genet*, 2003, vol. 4, 237-256 [0129] [0145]
- BARDEESY et al. *Proc. Natl. Acad. Sci. U.S.A.*, 2006, vol. 103, 5947-5952 [0129] [0145] [0214] [0215]
- SCHREIBER et al. *Gastroenterology*, 2004, vol. 127, 250-260 [0129] [0145] [0214]
- GEHLERT. *Proc Soc Exp Biol Med*, 1998, vol. 218, 7-22 [0131]
- SHEIKH et al. *Am J Physiol*, 1991, vol. 261, 701-15 [0131]
- FOURNIER et al. *Mol. Pharmacol.*, 1994, vol. 45, 93-101 [0131]
- KIRBY et al. *J Med Chem*, 1995, vol. 38, 4579-4586 [0131]
- RIST et al. *Eur. J. Biochem*, 1997, vol. 247, 1019-1028 [0131]
- KIRBY et al. *J Med Chz Tn*, 1993, vol. 36, 3802-3808 [0131]

- **GRUNDEMAR** et al. *Regulatory Peptides*, 1996, vol. 62, 131-136 [0131]
- **RAYMOND** et al. *Mol Biol Cell*, 2007, vol. 18, 4210-4221 [0134]
- **AHO** et al. *J Invest Dermatol*, 1999, vol. 113, 422-423 [0134] [0136]
- **ANDRA** et al. *Genes Dev*, 1998, vol. 12, 3442-3451 [0134]
- **OSMANAGIC-MYERS** et al. *J Biol Chem*, 2004, vol. 279, 18701-18710 [0134]
- **GREGOR** et al. *J Cell Sci*, 2006, vol. 119, 1864-1875 [0134]
- **SONNENBERG** et al. *Exp Cell Res*, 2007, vol. 313, 2189-2203 [0135] [0241]
- **IACOBUIZO-DONAHUE** et al. *Am J Pathol*, 2002, vol. 160 (4), 1239-1249 [0138]
- **SATO** et al. *Am J Pathol*, 2004, vol. 164 (3), 903-914 [0138]
- **JOHNSON** et al. *Molecular Carcinogenesis*, 2006, vol. 45, 814-827 [0138]
- **BOCZONADI** et al. *Experimental Cell Research*, 2007, vol. 313 (16), 3579-3591 [0138]
- **KIM** et al. *Cell*, 2006, vol. 125, 1269-1281 [0147]
- **MASER** et al. *Nature*, 2007, vol. 447, 966-971 [0147]
- **CHARI**. *Semin Oncol*, 2007, vol. 34, 284-294 [0150]
- **WEISSLEDER** et al. *Nat. Biotechnol*, 2001, vol. 19, 316-317 [0152]
- **MCCARTHY** et al. *Nanomedicine*, 2007, vol. 2, 153-167 [0152]
- **HOGEMANN** et al. *Bioconjug. Chem.*, 2000, vol. 11, 941-946 [0152] [0153]
- **JOSEPHSON** et al. *Bioconjug. Chem.*, 1999, vol. 10, 186-191 [0152]
- **IACOBUIZO-DONAHUE** et al. *Am J Pathol*, 2002, vol. 160, 1239-1249 [0157]
- **LICHA** et al. *SPIE*, 1996, vol. 2927, 192-198 [0158]
- **LACKOWICZ**. *Principles of Fluorescence Spectroscopy*. Kluwar Academic, 1999 [0158]
- **MISHRA** et al. *Expert Opinion on Biological Therapy*, 2003, vol. 3, 1173-1180 [0161]
- **BLAKELY** et al. *Cancer Research*, 1996, vol. 56, 3287-3292 [0165]
- Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy. **ORDER** et al. *Monoclonal Antibodies for Cancer Detection and Therapy*. Academic Press, 1985, 303-316 [0166]
- **MULLIGAN** et al. *Clin Cancer Res.*, 1995, vol. 1, 1447-1454 [0167]
- **MEREDITH** et al. *J Nucl Med*, 1996, vol. 37, 1491-1496 [0167]
- **ALVAREZ** et al. *Gynecologic Oncology*, 1997, vol. 65, 94-101 [0167]
- **CARON** et al. *Mol Ther*, 2001, vol. 3 (3), 310-8 [0182]
- **LANGEL**. *Cell-Penetrating Peptides: Processes and Applications*. CRC Press, 2002 [0182]
- **EL-ANDALOUSSI** et al. *Curr Pharm Des*, 2005, vol. 11 (28), 3597-611 [0182]
- **DESHAYES** et al. *Cell Mol Life Sci.*, 2005, vol. 62 (16), 1839-49 [0182]
- **KOEHLER** ; **MILSTEIN**. *Nature*, 1975, vol. 256, 495 [0187]
- **ELDRED** et al. *J. Med. Chem.*, 1994, vol. 37, 3882 [0192] [0205]
- **KU** et al. *J. Med. Chem.*, 1995, vol. 38, 9 [0192] [0205]
- **PEARSON** ; **LIPMAN**. *Proc. Natl. Acad. Sci. (USA)*, 1988, vol. 85, 2444-2448 [0195] [0207]
- **LIPMAN** ; **PEARSON**. *Science*, 1985, vol. 227, 1435-1441 [0195] [0207]
- **ROOT-BERNSTEIN**. *J. Theor. Biol.*, 1982, vol. 94, 885-859 [0196]
- **STEFANOWICZ** et al. *Letters in Peptide Science*, 1998, vol. 5, 329-331 [0196]
- **TARR**. *Methods of Protein Microcharacterization*. Humana Press, 1986, 155-194 [0201]
- Selected Methods in Cellular Immunology. WH Freeman, 1980 [0201]
- **MARSH**. *Int. Arch. Allergy Appl. Immunol.*, 1971, vol. 41, 199-215 [0201]
- Peptidomimetics Protocols. Human Press, 1998 [0202]
- Houben-Weyl Methods of Organic Chemistry: Synthesis of Peptides and Peptidomimetics. Thiele Verlag, 2003 [0202]
- **MAYO** et al. *J. Biol. Chem.*, 2003, vol. 278, 45746 [0202]
- **MELI** et al. *J. Med. Chem.*, 2006, vol. 49, 7721-7730 [0205]
- **MERRIFIELD** et al. *Biochemistry*, 1982, vol. 21, 5020-5031 [0207]
- **HOUGHTEN WELLINGS**. *Proc. Natl. Acad. Sci. (USA)*, 1985, vol. 82, 5131-5135 [0207]
- **ATHERTON**. *Methods in Enzymology*, 1997, vol. 289, 44-66 [0207]
- **GUY** ; **FIELDS**. *Methods in Enzymology*, 1997, vol. 289, 67-83 [0207]
- **STOCKWELL**. *Nature*, 2004, vol. 432, 846-854 [0210]
- **KAY** et al. *Mol. Diversity*, 1996, vol. 1, 139-140 [0210]
- **PFLEGER** et al. *Cell Signaling*, 2006, vol. 18, 1664-1670 [0210]
- **JUNG** et al. *Proteomics*, 2005, vol. 5, 4427-4431 [0210]
- **NIEUWENHUIJSEN** et al. *J. Biomol. Screen*, 2003, vol. 8, 676-684 [0210]
- **BERG**. *Angew. Chem. Int. Ed. Engl.*, 2003, vol. 42, 2462-2481 [0210]
- **ALLPORT** et al. *J Leukoc Biol*, 2002, vol. 71, 821-828 [0214]
- **KELLY** et al. *Neoplasia*, 2003, vol. 5, 437-444 [0216]
- **KELLY** et al. *Mol Imaging*, 2006, vol. 5, 24-30 [0217]
- **KELLY** et al. *Neoplasia*, 2006, vol. 8, 1011-1018 [0219] [0222] [0229] [0234] [0235]
- **MONTEL** et al. *Bioconjug. Chem.*, 2006, vol. 17, 905-911 [0227]

- **REYNOLDS et al.** *Bioconjug Chem*, 2005, vol. 16, 1240-1245 [0227]
- **WUNDERBALDINGER et al.** *Acad Radiol*, 2002, vol. 9 (2), S304-S306 [0227]
- **SCHELLENBERGER et al.** *Bioconjug Chem*, 2004, vol. 15, 1062-1067 [0227]
- **ALENCAR et al.** *Int. J. Cancer*, 2005, vol. 117, 335-339 [0229]
- **BARDEESY et al.** *Proc Natl Acad Sci USA*, 2006, vol. 103, 5947-5952 [0232]
- **CARRIERE et al.** *Proc Natl Acad Sci USA*, 2007, vol. 104 (11), 4437-42 [0234]
- **BARDEESY et al.** *Proc Natl Acad Sci U S A*, 2006, vol. 103, 5947-5952 [0234]
- **KELLY et al.** *Circ. Res.*, 2005, vol. 96, 327-336 [0234]
- **MURTAUGH et al.** *Cancer Cell*, 2007, vol. 11, 211-213 [0240]
- **WEISSLEDER et al.** *J Magn Reson Imaging*, 1997, vol. 7, 258-263 [0246]

摘要

在此描述的是癌細胞生物標記，如胰腺導管腺癌（PDAC）細胞生物標記，的組成和方法及用以診斷和治療癌症，如PDAC，的結合分子。辨認“可到達”蛋白質組的方法被披露以辨認癌症生物標記，如網蛋白-1，一PDAC生物標記。此外，包含結合至肽配的磁熒光納米粒子的影像組合物被披露以辨認PDAC。