



HU000033085T2

(19) **HU**(11) Lajstromszám: **E 033 085**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 13 305517**(51) Int. Cl.: **C07J 43/00** (2006.01)(22) A bejelentés napja: **2013. 04. 19.**

(96) Az európai bejelentés bejelentési száma:

EP 20130305517

(97) Az európai bejelentés közzétételi adatai:

EP 2792682 A1 **2014. 10. 22.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2792682 B1 **2016. 11. 09.**

(72) Feltaláló(k):
Poirier, Patricia, 49370 SAINT CLEMENT DE LA PLACE (FR)
Derrien, Yvon, 49770 LA MEIGNANNE (FR)
Forcato, Massimiliano, 49240 AVRILLE (FR)
COTARCA, Livius, 33052 Cervignano del Friuli (Udine) (IT)
Morice, Pierrick, 49070 BEAUCOUZE (FR)

(73) Jogosult(ak):
Zach System, 49240 Avrille (FR)

(74) Képviselő:
Danubia Szabadalmi és Jogi Iroda Kft., Budapest

(54) **Eljárás szilárd halmazállapotú abirateron-acetát előállítására**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 792 682 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
09.11.2016 Bulletin 2016/45

(51) Int Cl.:
C07J 43/00 (2006.01)

(21) Application number: **13305517.8**

(22) Date of filing: **19.04.2013**

(54) Process for producing a solid form of abiraterone acetate

Verfahren zur Herstellung einer festen Form von Abirateron-Azetat

Procédé pour la préparation d'un forme solide de l'actetate de l'abiraterone

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
 Designated Extension States:
BA ME

(43) Date of publication of application:
22.10.2014 Bulletin 2014/43

(73) Proprietor: **Zach System**
49240 Avrille (FR)

(72) Inventors:

- **Poirier, Patricia**
49370 SAINT CLEMENT DE LA PLACE (FR)
- **Derrien, Yvon**
49770 LA MEIGNANNE (FR)
- **Forcato, Massimiliano**
49240 AVRILLE (FR)
- **Cotarca, Livius**
33052 CERVIGNANO DEL FRIULI (UDINE) (IT)

- **Morice, Pierrick**
49070 BEAUCOUZE (FR)

(74) Representative: **Nevant, Marc et al**
Cabinet Beau de Loménie
158, rue de l'Université
75340 Paris Cedex 07 (FR)

(56) References cited:
WO-A1-2006/021776 CN-A- 101 768 199
CN-A- 102 336 801 CN-A- 102 558 275

- **POTTER G A ET AL: "A CONVENIENT, LARGE-SCALE SYNTHESIS OF ABIRATERONE ACETATE**
Ä3.BETA.-ACETOXY-17-(3-PYRIDYL)ANDROST A-5, 16-DIENEÜ, A POTENTIAL NEW DRUG FOR THE TREATMENT OF PROSTATE CANCER", ORGANIC PREPARATIONS AND PROCEDURES INTERNATIONAL, ORGANIC PREPARATION AND PROCEDURES CO., NEWTON HIGHLANDS, MA, US, vol. 290, no. 1, 1 January 1997 (1997-01-01), pages 123-128, XP008056111, ISSN: 0030-4948

EP 2 792 682 B1


Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to a process for the preparation of 17-substituted steroids and, more particularly, to an improved method of preparing micro size abiraterone or derivatives thereof in high yield and purity by means of a spherical agglomeration process.

BACKGROUND OF THE INVENTION

[0002] Abiraterone acetate, chemically designated as (3 β)-17-(3-pyridinyl)-androsta-5,16-dien-3-yl acetate of formula



is a prodrug which is converted *in vivo* to abiraterone, 17-(3-pyridyl)-androsta-5,16-dien-3 β -ol.

[0003] Abiraterone is a potent inhibitor of human cytochrome P450_{17 α} , a potential target enzyme in the treatment of hormone-dependent prostatic carcinoma.

[0004] Abiraterone acetate is the active ingredient of the approved drug (Zytiga[®]) which is administered in a solid oral dosage form (250 mg tablet).

[0005] Zytiga[®] in combination with prednisone is indicated for the treatment of patients with metastatic castration resistant prostate cancer (CRPC) who have received prior chemotherapy containing docetaxel.

[0006] Literature reports several processes for the preparation of abiraterone or derivatives thereof.

[0007] Abiraterone was first described in patent application EP 0633893 (BTG International Ltd.) covering 16,17-ene-17-(3-pyridyl) steroids as a class of compounds useful in the treatment of androgen- and oestrogen-dependent disorders.

[0008] Particularly, in the specific experimental work, abiraterone acetate is prepared from dehydroepiandrosterone-3-acetate and, then, converted into abiraterone. Chromatography is required to give the desired acetyl ester by elution with petroleum-diethyl ether and crystallization from hexane. Abiraterone is obtained by elution with toluene-diethyl ether and crystallization from toluene.

[0009] However, it has been observed that the overall yield of the process is low (around 48%) and chromatography appears to be the only tool able to provide a substantially pure product which is, then, crystallised from apolar solvents and, optionally, used in the downstream of the process.

[0010] EP 0721461 (BTG International Ltd.) describes an improved method for the preparation of (3 β)-acyloxy-16,17-ene-17-(3-pyridyl) steroids; especially, the preferred compound (3 β)-acetoxy-17-(3-pyridyl)-androsta-5,16-diene via acylation of (3 β)-hydroxy derivative and crystallization from ethanol/water and, then, hexane is prepared.

[0011] However, the estimated overall yield starting from dehydroepiandrosterone is low (around 41%) and, mainly, a final purification by reverse phase chromatography is required.

[0012] Org. Prepn. Proc. Int. Vol 29(1) pp 123-128 (1997) discloses the purification of abiraterone acetate by dissolution in acetonitrile, concentration and addition of water.

[0013] WO 2006/021776 (BTG International Ltd.) describes novel salt forms of C₂-C₄ acyl esters of abiraterone or a derivative thereof and to a process for the preparation of abiraterone or a salt or derivative thereof. The preferred salt is abiraterone acetate methanesulfonate which is, preferably, recovered from methyl tert-butyl ether.

[0014] WO 2006/021777 (BTG International Ltd.) describes a process for the preparation of abiraterone or C₂-C₄ acyl esters of abiraterone or a derivative thereof.

[0015] Coupling as well as isolation steps to abiraterone acetate are carried out by following the teachings of the International application WO '776 above.

[0016] However, the estimated overall yield of the process starting from dehydroepiandrosterone 3-acetate is very low (around 32%) with a purity around 97%; isolation as, inter alia, the mesylate salt entails an additional neutralization and optionally crystallization step(s) with further loss in yield.

[0017] Furthermore, Chinese application CN 101768199 discloses abiraterone acetate polymorphs A, B, C and D;

methods of preparing said polymorphs comprise recrystallizing abiraterone acetate from different solvents such as ethanol-hexane or ethanol-water mixtures, ethyl acetate or acetone, isopropanol as well as acetonitrile. Chinese application CN 102731605 provides for an abiraterone acetate purification method, said method comprises salification of crude abiraterone acetate with phosphoric acid to give a pale yellow crystal, i.e. abiraterone acetate phosphate, neutralization and further recrystallization from methanol yielding the acetate product.

[0018] Chinese application CN 102030798 provides for a purification method of abiraterone acetate. The method comprises reacting trifluoromethanesulfonic acid as a salt forming reagent with crude abiraterone acetate to obtain dry abiraterone acetate trifluoromethanesulfonic salt with a purity of over 97%; and performing a neutralization reaction with alkali in dichloromethane to give abiraterone acetate. Chinese application CN 102558275 discloses α -Type abiraterone acetate polymorph; the preparation method comprises dissolving abiraterone acetate in a solvent under heating condition to form a solution, filtering, cooling, crystallizing and drying; solvents used in the method are water, methanol, ethanol, isopropanol, isopropyl ether, acetonitrile, THF, ethyl acetate, chloroform, dichloromethane, toluene, hexane, acetone and mixtures thereof.

[0019] Chinese application CN 102321142 discloses abiraterone acetate crystalline form E; the preparation method comprises dissolving abiraterone acetate in a solvent, crystallizing under cooling or water addition and solid separation; solvents used in the method are methanol, ethanol, isopropanol, acetone, acetonitrile and/or THF.

[0020] Co-pending international patent publication No. WO-A-2013/053691 in the name of the same Applicant relates to a process for the preparation of 17-substituted steroids and, more particularly, to an improved method of synthesizing abiraterone or derivatives thereof in high yield and purity by means of a key 3 β -formyloxy intermediate. Abiraterone is, eventually, crystallized from alcoholic solvents and, optionally, converted into its 3 β -acetoxy ester in accordance with known techniques which is, in turn, crystallized preferably from a heptane/ethanol solution.

PURPOSE OF THE INVENTION

[0021] Abiraterone acetate is a white to off-white powder practically insoluble in aqueous media (pH range 2.0 to 12.9) very slightly soluble in 0.1N HCl solution and soluble to freely soluble in organic solvents. It is classified as BCS class IV and prepared exclusively as a single polymorphic form, namely, Form A.

[0022] EMA Assessment Report for Zytiga[®] states that the effect on particle size on manufacturability and tablet harness has been evaluated.

[0023] Comparative dissolution of tablets manufactured with varying API particle sizes demonstrate that tablet hardness was found to decrease with increasing drug substance particle size and for API D50 controlled between 3-10 μ m little effect on dissolution performance could be observed.

[0024] Solid state properties of active pharmaceutical ingredients have a decisive impact on dosage form development as well as *in vivo* performance of the drug. Micrometric properties of drug particles such as shape and size are essential for the formulation of solid dose unit. Particularly, the particle size of poorly soluble ingredients is commonly recognized as an issue due to its impact on dissolution properties.

[0025] It results from the art that the 16,17-ene-17-(3-pyridyl) steroid derivative, abiraterone acetate, endowed with a specific particle size distribution (PSD) allows to control tablet dissolution performance.

[0026] To comply with EMA reported PSD range, micronization is required.

[0027] It is known in the art that conventional dry size reduction of pharmaceutical ingredients is accomplished by impact size reduction, especially, via fluid energy impact equipments i.e. by micronizers.

[0028] However, said conventional micronization method appears not to be suitable for a reliable and economic industrial application since it suffers from significant cost and procedural drawbacks.

[0029] Micronizers are characterized by low efficiency and mechanical stress; high power input during micronization step gives rise to increased free surface energy, electrostatic tendencies and, thus, poor flowability and compressibility of powders which make them difficult to use in downstream processing in the pharmaceutical industry.

[0030] Mainly, it is worth noting that abiraterone acetate is a steroid and, thus, confinement issues must be taken into consideration when developing such a fluid energy impact size reduction.

[0031] Moreover, micronized particles tend to agglomerate and the increase in surface area is not always reflected in improved dissolution, thus, desired.

[0032] Hence, it would be desirable to study improved, efficient methods for preparing pure micro size abiraterone acetate with high yields and under conditions more favourable from the industrial application point of view.

SUMMARY OF THE INVENTION

[0033] We have now, surprisingly, found an easy and efficient process to prepare micro size abiraterone acetate via a spherical agglomeration process which allows to overcome the drawbacks of the processes described in the prior art.

[0034] The invention is illustrated by reference to the accompanying drawing described below:

Fig. 1 shows a powder X-ray diffractogram of abiraterone acetate crystalline Form A according to example 1.

DETAILED DESCRIPTION OF THE INVENTION

5 **[0035]** The present invention provides for an efficient process to prepare micro size abiraterone acetate via a spherical agglomeration process.

[0036] Common techniques for the preparation of micro-size drugs are the mechanical comminution of previously formed larger particles by micronizers or fluid-energy impact equipments.

10 **[0037]** We have surprisingly found that micro-size abiraterone acetate can be prepared by first precipitating the active ingredient thin crystals out of a solution; agglomerating said crystals in order to isolate them from the reaction mixture; then, disagglomerating by mild milling to get particles endowed with the desired PSD. The key step of the invention consists essentially in a spherical agglomeration of abiraterone acetate crystals by using a three solvent system. It involves dissolution of abiraterone acetate in a good solvent, crystallization and agglomeration by means of an antisolvent in the presence of a bridging liquid.

15 **[0038]** Therefore, an object of the present invention is a process for preparing micro size abiraterone acetate which comprises:

- a) dissolving abiraterone acetate in a polar aprotic solvent;
- b) precipitating abiraterone acetate crystals out of the solution by addition of an antisolvent;
- 20 c) agglomerating said crystals in the presence of a bridging liquid, wherein said bridging liquid is an ether;
- d) isolating the resulting spherical agglomerates of abiraterone acetate crystals; and
- e) milling the agglomerates.

25 **[0039]** Abiraterone acetate is a well known commercially available active pharmaceutical ingredient whose preparation is extensively described in the art.

[0040] According to a preferred embodiment of the invention abiraterone acetate is prepared by following the teachings of Co-pending international patent publication No. WO-A-2013/053691 in the name of the same Applicant.

[0041] The present invention provides for an efficient process to prepare micro size abiraterone acetate via a spherical agglomeration process.

30 **[0042]** Abiraterone acetate is first dissolved in a good solvent (step a).

[0043] Operatively, abiraterone acetate is contacted with a polar aprotic solvent to give a solution which is optionally maintained at a temperature suitable to avoid crystallization.

35 **[0044]** Preferred polar aprotic solvents are dimethylacetamide (DMA), dimethylsulfoxide (DMSO) and dimethylformamide (DMF) the latter being the preferred one. Preferably, the amount of good solvent is comprised between 10-15 l/Kg, more preferably is 12 l/Kg.

[0045] Preferably, the concentration of the solution is comprised between 70-90 mg/ml, more preferably is 84 mg/ml.

[0046] In a preferred embodiment of the invention the solution obtained in step a) is maintained at a temperature between 25-40°C, preferably around 30°C.

[0047] Abiraterone acetate crystals are precipitated out of the solution by addition of an antisolvent (step b).

40 **[0048]** The general procedure comprises precipitating said crystals by addition of a suitable amount of antisolvent.

[0049] In one embodiment of the invention said addition occurs over a period of around 1 hour; the temperature of the reaction mixture is maintained between 15-30°C to give a thick slurry; and, when the addition is completed, said mixture is kept under stirring for a few minutes at a temperature comprised between 15-30°C.

45 **[0050]** Preferred antisolvents are water or water in admixture with water miscible organic solvent(s) such as, for instance, methanol, ethanol, isopropanol and acetone; water being preferred.

[0051] Preferably, the reaction mixture is kept at a temperature comprised between 20-25°C.

[0052] Preferably the amount of antisolvent is comprised between 10-20 l/Kg, more preferably between 12-15 l/Kg.

[0053] Preferably, the ratio of the antisolvent with respect to the polar aprotic solvent is comprised between 1 and 1.25 v/v.

50 **[0054]** Preferably, the solution is added to the antisolvent over a period of around 30 min. to 2 hours, more preferably over a period of 1 hour.

[0055] In a preferred embodiment of the invention, a solution of pure abiraterone acetate in dimethylformamide at 30°C is added to purified water over a period of 1 hour, by maintaining the temperature of the mixture between 20 and 25°C. Abiraterone acetate is precipitated by giving a thick slurry. When the addition is completed, the mixture is stirred for 15 minutes at around 20-25°C.

55 **[0056]** The crystallization conditions of the invention allows obtaining very thin abiraterone acetate crystals.

[0057] Agglomeration occurs in the presence of a bridging liquid (step c).

[0058] Operatively, a suitable amount of bridging liquid is added to the organic layer coming from step b) to give

EP 2 792 682 B1

spherical agglomerates of abiraterone acetate crystals. The bridging liquid is an ether such as, for instance, diethyl ether, di-isopropyl ether, cyclopentyl-methyl ether (CPME) and tert-butyl methyl ether (MTBE); the latter being preferred.

[0059] Preferably, the bridging liquid is added to the slurry as in step b) over a period comprised between 15 min. to 1 hour, 30 min. being more preferred.

[0060] Preferably, the bridging liquid is added to the slurry as in step b) by keeping the temperature of the reaction mixture comprised between 15-30°C, 20-25°C being more preferred.

[0061] Preferably the amount of bridging liquid is comprised between 2-3 l/Kg, more preferably between 2.5-2.7 l/Kg.

[0062] Preferably, the ratio of the bridging liquid with respect to the polar aprotic solvent is comprised between 0.21-0.23 v/v.

[0063] Preferably, when the addition is completed, the reaction mixture is stirred over a period of 15-30 min.

[0064] Preferably, when the addition is completed, the reaction mixture is kept at a temperature comprised between 15-30°C, more preferably between 20-25°C.

[0065] In a preferred embodiment of the invention tert-butyl methyl ether is added to the slurry obtained in step b) over a period of 30 minutes by maintaining the temperature at around 20-25°C. When the addition is completed agglomerates are observed. The mixture is then stirred for 15 to 30 minutes at 20-25°C temperature.

[0066] Spherical agglomerates of abiraterone acetate crystals are then isolated (step d).

[0067] In one embodiment of the process of the invention, the step d) isolation is carried out by filtration.

[0068] Operatively, the slurry coming from step c) is filtered and the resulting solid product is washed with a suitable solvent; the wet solid is, then, dried to give the desired product.

[0069] Preferably, said solid product is washed with water; more preferably, several washes are carried out in order to eliminate any residual solvent.

[0070] Preferably, said wet solid is dried under vacuum.

[0071] Micro size abiraterone acetate is, eventually, obtained by disagglomerating the isolated dry solid (step e).

[0072] Operatively, the obtained wet solid is dried under vacuum and, then, milled to give quantitatively abiraterone acetate as very thin particles.

[0073] No micronization step by micronizers or fluid-energy impact equipments is required. In a preferred embodiment of the invention, abiraterone acetate is dissolved in dimethylformamide, then, precipitated by water addition to give a thick slurry which is agglomerated by adding tert-butyl methyl ether.

[0074] The filtration of the slurry is very easy. After water washes, the wet solid is dried under vacuum, then, milled to give quantitatively micro size abiraterone acetate. The micro size abiraterone acetate according to the invention complies with the desired particle size distribution of $D(0.5) = 3-10 \mu\text{m}$.

[0075] As above reported, it is known in the art that said specific PSD allows to control tablet dissolution performance.

[0076] The agglomerating step c) allows isolating abiraterone acetate crystals from the reaction mixture (thick slurry) prepared according to the precipitation conditions. Spherical agglomerates of abiraterone acetate crystals obtained by the process according to the invention are, thus, useful intermediates in the preparation of micro size abiraterone acetate.

[0077] Since the invention is aimed to prepare few micron-size particles, said agglomerates need to be milled.

[0078] The inventors have observed that disagglomeration takes place even during a drying procedure, however, mild milling allows getting a homogeneous suitable particle size distribution.

[0079] It is worth noting that the agglomeration process of the invention does not have an impact on the polymorphic form of the crystals of the active ingredient used as substrate.

[0080] Therefore, a further object of the invention is a process for preparing micro size abiraterone acetate as described above wherein spherical agglomerates of abiraterone acetate crystalline Form A are isolated.

[0081] It is thus evident how the method object of the invention is suitable for industrial production, and constitutes an efficient and economic synthetic improvement for the preparation of abiraterone and derivatives thereof.

[0082] To the best of the inventors' knowledge, the spherical agglomeration process carried out on abiraterone acetate is neither known in the art nor suggested by any prior art reference.

[0083] The inventors have developed an original method which avoids micronization via conventional physical means i.e. by micronizers or fluid-energy impact equipments and allows obtaining the API, abiraterone acetate, complying with the suitable particle size distribution range (D_{50} between 3-10 μm).

[0084] The product is obtained in quantitative yields and with very high purity; no change of the solid polymorph form as well as the quality profile during and after the agglomeration process of the invention is observed.

[0085] Combined precipitation and spherical agglomeration according to the invention allows isolating abiraterone acetate by means of common techniques such as filtration; then, disagglomeration by mild milling allowed the inventors to prepare the active ingredient in batch as a very thin crystalline solid endowed with the desired PSD and without any high energy physical stress.

[0086] In substance, the process of the invention provides for:

- a) quantitative conversion into highly pure spherical agglomerates;

EP 2 792 682 B1

- b) easy and efficient isolation of the API;
- c) no micronization by micronizers or fluid-energy impact equipments needed;
- d) higher efficiency in terms of feed/mill rate (kg/h) for specific API;
- e) controlled PSD distribution (3-10 μm);
- f) reduced physical stress of particles;
- g) reduced confinement concerns; and
- h) no polymorphic conversion concerns.

[0087] A practical embodiment of the process object of the present invention comprises solubilising abiraterone acetate in a polar aprotic solvent; precipitating it by antisolvent addition to give a thick slurry; agglomerating said slurry by addition of an ether (bridging liquid); filtrating and washing; so obtained wet solid is dried under vacuum, then, milled to give quantitatively micro size abiraterone acetate as very thin particles [D(0.5)=3-10 μm].

[0088] A preferred practical embodiment of the process object of the present invention comprises solubilising abiraterone acetate in DMF; said solution is maintained at around 30°C; said solution is, then, added to purified water in around 1 hour by maintaining the temperature of the mixture at around 20-25°C to give a thick slurry; MTBE is then added in around 30 min. by maintaining the temperature of the slurry at around 20-25°C; slurry is, then, filtered and the solid washed with water; so obtained wet solid is dried under vacuum and, then, milled to give quantitatively micro size abiraterone acetate as very thin particles [D(0.5)=3-10 μm].

[0089] For better illustrating the invention the following example is now given.

Example 1

[0090] A solution of 16 g of pure abiraterone acetate in 192 ml of dimethylformamide, maintained at 30°C, was added to 192 ml of purified water over a period of 1 hour, maintaining the temperature of the mixture between 20 and 25°C. Abiraterone acetate was precipitated by giving a thick slurry. When the addition was completed, the mixture was stirred for 15 minutes at 20-25°C. 40 ml of tert-butyl methyl ether were then added to the slurry over a period of 30 minutes, maintaining the temperature at 20-25°C. When the addition was completed, agglomerates were observed. The mixture was then stirred for 15 to 30 minutes at 20-25°C. The slurry was filtered and the solid washed three times with 32 ml of purified water. The wet solid was then dried at 50-55°C under vacuum to give 15,7g of dried product which was then milled.

Measured Particle Size Distribution : D(0.5) = 4.9 μm

Expected: D(0.5) = 3-10 μm

[0091] PSD was determined by laser diffraction-dry powder method with a Malvern Mastersizer 2000 device.

[0092] The X-ray diffraction pattern of the milled dried product was measured on a D8 ADVANCE® diffractometer (Bruker) equipped with a Cu K alpha-1 radiation source and a VANTEC-1 detector (Bruker). The diffractogram is shown in Figure 1 and comprises the peaks listed in table 1 below. It can be seen from the diffractogram that micro size abiraterone acetate of example 1 has all the characteristic peaks of abiraterone acetate crystalline form A.

[0093] It should be borne in mind that the relative intensity of the X-ray powder diffraction peaks can vary depending upon sample preparation technique, sample mounting procedure and the particular instrument employed.

Table 1

Caption	Angle	d value	Intensity	Intensity %
	2-Theta °	Angstrom	Cps	%
2th=5.806 °	5.806	15.21083	26.1	10.7
2th=9.485 °	9.485	9.31672	13.2	5.4
2th=10.755 °	10.755	8.21974	3.62	1.5
2th=12.049 °	12.049	7.33922	92.7	37.8
2th=12.580 °	12.58	7.03074	16.4	6.7
2th=13.064 °	13.064	6.77145	1.44	0.6
2th=14.813 °	14.813	5.97542	153	62.5
2th=15.094 °	15.094	5.86486	138	56.3
2th=15.919 °	15.919	5.56284	106	43.2

EP 2 792 682 B1

(continued)

5
10
15
20
25
30
35
40
45
50

Caption	Angle	d value	Intensity	Intensity %	
	2-Theta °	Angstrom	Cps	%	
	2th=16.566 °	16.566	5.34683	2.92	1.2
	2th=17.209 °	17.209	5.14872	33.5	13.6
	2th=18.387 °	18.387	4.8214	245	100
	2th=18.935 °	18.935	4.68289	164	66.9
	2th=19.759 °	19.759	4.48945	33.1	13.5
	2th=20.162 °	20.162	4.4008	1.59	0.6
	2th=21.679 °	21.679	4.09604	143	58.5
	2th=22.448 °	22.448	3.95747	54.3	22.1
	2th=23.044 °	23.044	3.85651	83.4	34
	2th=23.411 °	23.411	3.79686	24.2	9.8
	2th=23.672 °	23.672	3.75553	14.1	5.7
	2th=24.276 °	24.276	3.66342	21.5	8.8
	2th=25.380 °	25.38	3.50654	35.7	14.6
	2th=25.968 °	25.968	3.42851	21.8	8.9
	2th=26.802 °	26.802	3.32361	22.4	9.1
	2th=27.517 °	27.517	3.23891	58	23.6
	2th=27.921 °	27.921	3.19294	14.6	6
	2th=29.031 °	29.031	3.07332	3.65	1.5
	2th=29.875 °	29.875	2.98835	29.6	12.1
	2th=30.475 °	30.475	2.93087	8.68	3.5
	2th=30.964 °	30.964	2.8857	6.72	2.7
	2th=31.168 °	31.168	2.86732	5.21	2.1
	2th=32.211 °	32.211	2.77682	16.3	6.7
	2th=33.438 °	33.438	2.67764	4.57	1.9
	2th=34.480 °	34.48	2.59905	15.6	6.4
	2th=34.788 °	34.788	2.57674	10.6	4.3
	2th=35.828 °	35.828	2.50432	4.73	1.9
	2th=36.796 °	36.796	2.44065	26.5	10.8
	2th=37.640 °	37.64	2.38779	10.5	4.3
	2th=38.440 °	38.44	2.33995	12.3	5
	2th=38.759 °	38.759	2.3214	19.2	7.8
	2th=39.297 °	39.297	2.29085	4.47	1.8

Claims

1. A process for preparing micro size abiraterone acetate which comprises:

a) dissolving abiraterone acetate in a polar aprotic solvent;

EP 2 792 682 B1

- b) precipitating abiraterone acetate crystals out of the solution by addition of an antisolvent;
- c) agglomerating said crystals in the presence of a bridging liquid, wherein said bridging liquid is an ether;
- d) isolating so obtained spherical agglomerates of abiraterone acetate crystals; and
- e) milling the agglomerates.

- 5
2. The process according to claim 1, wherein the polar aprotic solvent of step a) is selected from DMA, DMSO and DMF.
3. The process according to claim 2, wherein the polar aprotic solvent is DMF.
- 10
4. The process according to any of claims 1 to 3, wherein the solution obtained in step a) is maintained at a temperature comprised between 25-40°C.
5. The process according to any of claims 1 to 4, wherein the antisolvent of step b) is selected from water and water in admixture with methanol, ethanol, isopropanol or acetone.
- 15
6. The process according to claim 5, wherein the antisolvent is water.
7. The process according to any of claims 1 to 6, wherein step b) is carried out at a temperature comprised between 20-25°C.
- 20
8. The process according to any one of claims 1 to 7, wherein the ether is selected from diethyl ether, di-isopropyl ether, cyclopentyl-methyl ether (CPME) and tert-butyl methyl ether (MTBE).
9. The process according to claim 8, wherein the ether is MTBE.
- 25
10. The process according to any of claims 1 to 9, wherein step c) is carried out at a temperature comprised between 15-30°C.
11. The process according to any of claims 1 to 10, wherein the ratio of the antisolvent with respect to the polar aprotic solvent is comprised between 1-1.25 v/v.
- 30
12. A process according to any of claims 1 to 11, wherein spherical agglomerates of abiraterone acetate crystalline Form A are isolated.

35

Patentansprüche

1. Verfahren zum Herstellen von Abirateronacetat in Mikrogröße, das umfasst:
- 40
- a) Lösen von Abirateronacetat in einem polaren aprotischen Lösungsmittel,
 - b) Ausfällen von Abirateronacetatkristallen aus der Lösung durch Hinzufügen eines Antilösungsmittels,
 - c) Agglomerieren der Kristalle in Gegenwart einer brückenbildenden Flüssigkeit, wobei die brückenbildende Flüssigkeit Ether ist,
 - d) Isolieren der so erhaltenen sphärischen Agglomerate aus Abirateronacetatkristallen und
 - e) Mahlen der Agglomerate.
- 45
2. Verfahren nach Anspruch 1, wobei das polare aprotische Lösungsmittel in Schritt a) aus DMA, DMSO und DMF ausgewählt ist.
- 50
3. Verfahren nach Anspruch 2, wobei das polare aprotische Lösungsmittel DMF ist.
4. Verfahren nach einem der Ansprüche 1 bis 3, wobei die in Schritt a) erhaltene Lösung bei einer zwischen 25 und 40 °C liegenden Temperatur gehalten wird.
- 55
5. Verfahren nach einem der Ansprüche 1 bis 4, wobei das Antilösungsmittel in Schritt b) aus Wasser und Wasser mit Beimischungen von Methanol, Ethanol, Isopropanol oder Aceton ausgewählt ist.
6. Verfahren nach Anspruch 5, wobei das Antilösungsmittel Wasser ist.

EP 2 792 682 B1

7. Verfahren nach einem der Ansprüche 1 bis 6, wobei Schritt b) bei einer zwischen 20 und 25 °C liegenden Temperatur durchgeführt wird.
- 5 8. Verfahren nach einem der Ansprüche 1 bis 7, wobei der Ether aus Diethylether, Diisopropylether, Cyclopentylmethylether (CPME) und Tertbutylmethylether (MTBE) ausgewählt ist.
9. Verfahren nach Anspruch 8, wobei der Ether MTBE ist.
- 10 10. Verfahren nach einem der Ansprüche 1 bis 9, wobei Schritt c) bei einer zwischen 15 und 30 °C liegenden Temperatur durchgeführt wird.
11. Verfahren nach einem der Ansprüche 1 bis 10, wobei das Verhältnis des Antilösungsmittels in Bezug auf das polare aprotische Lösungsmittel zwischen 1 und 1,25 Vol.-% liegt.
- 15 12. Verfahren nach einem der Ansprüche 1 bis 11, wobei sphärische Agglomerate der Abirateronacetatkristallform A isoliert werden.

Revendications

- 20 1. Procédé de préparation d'acétate d'abiratérone de taille microscopique qui comprend :
- 25 a) dissoudre de l'acétate d'abiratérone dans un solvant aprotique polaire ;
b) précipiter des cristaux d'acétate d'abiratérone à partir de la solution par addition d'un antisolvant ;
c) agglomérer lesdits cristaux en présence d'un liquid de pontage, ledit liquid de pontage étant un éther ;
d) isoler les agglomérats sphériques de cristaux d'acétate d'abiratérone ainsi obtenus ; et
e) broyer les agglomérats.
- 30 2. Procédé selon la revendication 1, dans lequel le solvant aprotique polaire de l'étape a) est choisi parmi le DMA, le DMSO et le DMF.
3. Procédé selon la revendication 2, dans lequel le solvant aprotique polaire est le DMF.
- 35 4. Procédé selon l'une des revendications 1 à 3, dans lequel la solution obtenue à l'étape a) est maintenue à une température comprise entre 25 et 40°C.
5. Procédé selon l'une des revendications 1 à 4, dans lequel l'antisolvant de l'étape b) est choisi parmi l'eau et un mélange d'eau et de méthanol, éthanol, isopropanol ou acétone.
- 40 6. Procédé selon la revendication 5, dans lequel l'antisolvant est l'eau.
7. Procédé selon l'une des revendications 1 à 6, dans lequel l'étape b) est mise en oeuvre à une température comprise entre 20 et 25°C.
- 45 8. Procédé selon l'une des revendications 1 à 7, dans lequel l'éther est choisi parmi l'éther éthylique, l'éther isopropylique, l'éther de cyclopentyle et de méthyle et l'éther de tert-butyle et de méthyle (MTBE).
9. Procédé selon la revendication 8, dans lequel l'éther est le MTBE.
- 50 10. Procédé selon l'une des revendications 1 à 9, dans lequel l'étape c) est mise en oeuvre à une température comprise entre 15 et 30°C.
11. Procédé selon l'une des revendications 1 à 10, dans lequel le rapport de l'antisolvant au solvant aprotique polaire est compris entre 1 et 1.25 v/v.
- 55 12. Procédé selon l'une des revendications 1 à 11, dans lequel on isole des agglomérats sphériques de la forme cristalline A de l'acétate d'abiratérone.




FIG. 1

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0633893 A [0007]
- EP 0721461 A [0010]
- WO 2006021776 A [0013]
- WO 2006021777 A [0014]
- CN 101768199 [0017]
- CN 102731605 [0017]
- CN 102030798 [0018]
- CN 102558275 [0018]
- CN 102321142 [0019]
- WO 2013053691 A [0020] [0040]

Non-patent literature cited in the description

- *Org. Prepn. Proc. Int.*, 1997, vol. 29 (1), 123-128 [0012]

ELJÁRÁS SZILÁRD HALMAZÁLLAPOTÚ ABIRATERON-ACETÁT ELŐÁLLÍTÁSÁRA

Szabadalmi igénypontok

1. Eljárás mikroméretű abirateron-acetát előállítására, amely tartalmazza a következőket:

- a) abirateron-acetát feloldását poláris aprotikus oldószerben;
- b) abirateron-acetát kristályok kicsapását az oldatból ellenoldat hozzáadásával;
- c) a kristályokból agglomerátum képzését áthidaló folyadék jelenlétében, ahol az áthidaló folyadék éter;
- d) az abirateron-acetát kristályai így kapott szférikus agglomerátumainak izolálását; és
- e) az agglomerátumok őrlését.

2. Az 1. igénypont szerinti eljárás, ahol az a) lépés szerinti, poláris aprotikus oldószer DMA, DMSO és DMF közül választott.

3. A 2. igénypont szerinti eljárás, ahol a poláris aprotikus oldószer DMF.

4. Az 1-3. igénypontok bármelyike szerinti eljárás, ahol az a) lépésben kapott oldatot 25-40°C hőmérsékleten tartjuk fenn.

5. Az 1-4. igénypontok bármelyike szerinti eljárás, ahol a b) lépés szerinti ellenoldat a következők közül választott: víz, valamint metanollal, etanollal, izopropanollal vagy acetonnal kevert víz.

6. Az 5. igénypont szerinti eljárás, ahol az ellenoldat víz.

7. Az 1-6. igénypontok bármelyike szerinti eljárás, ahol a b) lépést 20-25°C hőmérsékleten végezzük.

8. Az 1-7. igénypontok bármelyike szerinti eljárás, ahol az éter a következők közül választott: dietil-éter, diizopropil-éter, ciklopentil-metil-éter (CPME) és terc-butil-metil-éter (MTBE).

9. A 8. igénypont szerinti eljárás, ahol az éter MTBE.

10. Az 1-9. igénypontok bármelyike szerinti eljárás, ahol a c) lépést 15-30°C közötti hőmérsékleten végezzük.

11. Az 1-10. igénypontok bármelyike szerinti eljárás, ahol az ellenoldat aránya a poláris aprotikus oldószerhez képest 1-1,25 térfogat%.

12. Az 1-11. igénypontok bármelyike szerinti eljárás, ahol A kristályformájú abirateron-acetát szférikus agglomerátumait izoláljuk.

