

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 July 2009 (09.07.2009)

PCT

(10) International Publication Number
WO 2009/086191 A1

(51) International Patent Classification:
CI2N 5/02 (2006.01) *CI2N 5/08* (2006.01)

(74) Agent: **PILLAI, Suresh, B.**; McDonnell Boehnen Hulbert & Berghoff LLP, 300 S. Wacker Drive, Suite 3100, Chicago, IL 60606 (US).

(21) International Application Number:
PCT/US2008/087818

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
19 December 2008 (19.12.2008)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/015,764 21 December 2007 (21.12.2007) US
20076604 21 December 2007 (21.12.2007) NO

Published:
— with international search report

(71) Applicant (for all designated States except US): **CELL-CURA, INC.** [US/US]; 505 South Rosa Road, Suite 115, Madison, WI 53719 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ALI BIN M. ABDULLAH, Jaffar** [MY/MY]; A4-3 Kondo Danau Murni, Taman Danau Desa, 58100 Kuala Lumpur (MY).

WO 2009/086191 A1

(54) Title: PROTEIN-FREE GAMETE AND EMBRYO HANDLING AND CULTURE MEDIA PRODUCTS

(57) Abstract: This invention discloses a protein-free cell culture medium for assisted reproductive technologies containing 14 kDa methylcellulose.

PROTEIN-FREE GAMETE AND EMBRYO HANDLING AND CULTURE MEDIA PRODUCTS

FIELD OF THE INVENTION

5

The present invention relates to protein-free media (PFM) specialized and optimized for human reproduction and fertility programs. The substantially protein-free media products described herein are useful in preventing transmission of protein-bound pathogens including *inter alia* and in particular prions to patients undergoing infertility treatment, to newborns conceived from Assisted Reproduction Technologies (ART) and to workers in these areas of endeavor, specifically relating to application in human ART.

BACKGROUND OF THE INVENTION

15

Conventional commercially-available embryo culture media for human ART procedures contain human serum albumin (HSA) obtained from human blood and tissue sources. In some laboratories bovine serum albumin (BSA) is also used as a source of protein in human embryo culture procedures (Loutradis *et al.*, 1992; Quinn, 20 1994), similarly obtained from blood and tissue sources from cows. The efficiency of media containing HSA and BSA was reported to be similar (Staessen *et al.*, 1998). The use in culture medium of protein obtained from donors (human or bovine) has the potential to transmit diseases to patients undergoing assisted conception treatment. The generation of viable human embryos in a chemically-defined culture system 25 (devoid of added protein or albumin, or animal or human-derived protein extract) beginning from oocyte collection, followed by sperm washing, insemination, fertilization up to the cleaved embryo stage and finally embryo transfer has not been described, although chemically-defined media for mouse, rabbit and primates have been reported in recent years (Spindle, 1995; Li 30 *et al.*, 1996; Schramm and Bavister, 1996). Previous claims of a chemically defined protein-free medium (PFM) for human application, in fact, were not truly protein-free, because the sperm meant for fertilization was still prepared in medium that contained serum proteins (Mohr & Trounson, 1986; Serta and Kiessling 1997 (Abst); Parinaud *et al.* 1999). A totally protein-free media for ART procedures thus has not been 35 described hitherto.

The availability of a chemically defined medium used for generating viable early cleavage stage human embryos is needed in the art to ensure the safety of patients undergoing assisted reproduction treatment by avoiding the use of potentially hazardous donor protein. Concern over possible transmission of pathogenic diseases, 5 in particular viral diseases, such as human acquired immunodeficiency disease syndrome (AIDS) and hepatitis, or Creutzfeldt-Jakob disease (CJD) transmitted by prions or others in blood-derived products has led a number of providers of healthcare services in the area of ART worldwide to seek alternative(s) to donor protein for their embryo culture and handling procedures.

10

The transmission of a deadly viral disease (AIDS) to hemophiliacs through blood-derived products is well documented (See, for example, Craven *et al.*, 1997, *Med. Sci. Law* 37: 215-227; Keshavjee *et al.*, 2001, *Soc. Sci. Med.* 53: 1081-1094; Weinberg *et al.*, 2002, *Ann. Intern. Med.* 136: 312-319; Evatt, 2006, *Semin. Hematol.* 43: S4-9).

15

Human growth hormone extracted from the pituitary was found to be capable of transmitting CJD to humans (Esmonde *et al.*, 1994) and human gonadotropin injections could also transmit CJD from person to person (CDC, 1985). CJD can be transmitted through blood (although the titer of CJD prions is low in blood; Heye *et al.*, 1994). In the past an epidemic of hepatitis B occurred in about 200 IVF patients

20

that received embryos cultured in medium containing pooled sera contaminated with hepatitis B virus (van Os *et al.*, 1991). Recently the scientific community was confronted with the dilemma of having to inform their patients that a commercial preparation of a culture medium used for embryo culture and handling may be contaminated with albumin donated by a person who later died of CJD (Kemmann, 25 1998).

30

There are a number of reports of development of mouse blastocysts from zygote and cleaved stages in a protein-free medium. The earliest reports include those of Brinster (1965) and, Cholewa and Whitten (1970) who used polyvinylpyrrolidone (PVP), which is a high molecular weight colloid, as a lubricant and for increasing the viscosity of the medium. Subsequent to this work a number of other workers have successfully cultured mouse as well as other mammalian embryos in protein-free media (Dandekar and Glass, 1990; Spindle, 1995; Li *et al.*, 1996; Schramm and Bavister, 1996). In recent years, in addition to PVP, polyvinyl alcohol (PVA)

(Bavister, 1995) has also been used to replace serum protein in culture medium. For example, Biggers et al. (1997) investigated the effect of replacing bovine serum albumin (BSA) with polyvinyl alcohol (PVA) and/or amino acids on mouse zygote development. They observed that PVA could not be substituted completely for BSA

5 in the mouse embryo culture medium. The effect of PVA on rate of blastocyst development was only slightly less than with BSA but the rate of partial hatching was significantly less. Substitution of BSA with PVA lowered the overall response but did not lead to major perturbations.

10 In addition to its many biological roles, serum proteins confer useful physical attributes such as lubrication and viscosity in the culture medium. Increased viscosity and lubrication in the culture medium is required for ease of handling and manipulation of the embryo and to prevent it from adhering to the walls of the culture dish and embryo transfer catheters. The incorporation of PVP and PVA merely serve

15 to duplicate the physical attributes of serum proteins. However, PVP and PVA are not sources of fixed nitrogen and they do not perform the various biological roles of protein. In addition, the teratological properties of PVP and PVA have not been fully examined, which make their use in human therapeutic assisted reproduction questionable (Gardner and Lane, 1998a).

20 Some investigators have attempted to generate viable human embryos in a protein-free culture system, most recently Serta et al. (1997) and Parinaud et al. (1998a). However Serta and co-workers (1997) prepared their spermatozoa for insemination by swim-down through a column of BSA (although subsequent culture was performed in

25 a protein-free medium). These workers achieved a pregnancy rate of 31% (n=45) some of which have proceeded to term with the birth of normal offspring. Parinaud et al. (1998a) obtained fertilizations with spermatozoa prepared in a protein-free medium when insemination was performed in the same medium. However the resultant zygotes were cultured in BM1 medium, and although this reference did not

30 specify whether their BM1 medium contained protein, a previous publication from the same group suggested BM1 medium contained 1% HSA (Parinaud et al., 1998b).

Some workers have shown that replacement of serum protein with a single antioxidant and chelator such as EDTA (Mehta and Kiessling, 1990; Serta et al., 1997) does not

impair fertilization and cleavage of viable embryos in the mouse and human. However Serta et al. (1997) suggested that the embryo transfer catheter be rinsed extensively with the protein-free medium, which implies the tendency of embryos to stick to the inner wall of embryo transfer catheter. Serum protein also has a role in 5 maintaining pH in culture medium (Moessner et al., 1993). Beside its role as a nutrient in biological systems, protein has a number of other roles such as a chain-breaking antioxidant and a chelator of metal ions (Barber, 1961; Vidlakova et al., 1972; Wayner et al., 1987).

10 The physiological functions of albumin and plasma proteins in general are well documented. The role of albumin in preventing membrane peroxidation indicates a direct role in membrane stability. It is involved in capillary membrane permeability and in osmoregulation. Albumin provides 80% of the total colloid osmotic pressure in plasma. Albumin is involved in the transport of carbon dioxide and acts as a pH 15 buffer; albumin accounts for the greatest (95%) portion of the non-bicarbonate buffer value of plasma. Proteins also serve as a source of energy. Deaminated alanine is pyruvate, which can be either converted to acetyl-CoA or glucose and glycogen. Albumin may help solubilize lipids and transports hormones, vitamins and metals. It serves as reservoirs for the release and use of these components.

20 Any attempt at substituting serum albumin in culture medium should therefore take into consideration these *in vivo* roles and physical attributes which are useful for embryo handling and manipulation *in vitro*. A single component may not fulfill all the functions of serum protein.

25 Although protein-free media that supports development of a number of animal species has been described previously, no such protein-free media has been successfully used in humans, nor could such media be presumed to support or be optimal for human embryo development. Thus there exists a distinct need for a defined, protein-free 30 growth medium especially adapted for human ART and IVF.

BACKGROUND OF THE RELATED ART

There are previously known putatively “protein-free” media for treating and cultivating mammalian cells, particularly cells from rodents (mice, rats, guinea pigs, etc.) but also for fertilization, embryo development and pregnancy in humans for *in vitro* fertilization.

5 Caro *et al.* have described one such “protein-free” medium for human application; however they were not able to overcome the need for proteins in the sperm preparation medium because proteins are required to induce capacitation of the sperm without which the sperm will not acquire the capacity to penetrate and/or fertilize the egg. Thus, Caro and co-worker’s media system is not completely protein-free.

10 Likewise, Serta *et al.* (1997; abstract) and Parinaud *et al.* (1998a, abstract; 1999) have also described “protein-free” media for human ART but they, like Caro *et al.* (1986) before them used proteins in at least one stage of their culture system.

Washing sperm previously prepared with medium containing added proteins as described in these references does not remove contaminating infectious agents with complete certainty, especially viruses and prions. In addition, this is also disadvantageous because it subjects the spermatozoa to unnecessary stressful washing conditions, and can impose excessive demands on manpower and resources.

20 “Protein-free” media for growth of mammalian or particularly human cells outside the IVF context has been disclosed, *inter alia*, in Kovár *et al.*, 1987, *Biotechnology Letters*, vol. 9 no. 4, p. 259-264 “Iron Compounds at high Concentrations Enable Hybridoma Growth in a Protein-free Medium”; Keen, 1995, *Cytotechnology*, vol. 17: 193-202 “The culture of rat myeloma and rat hybridoma cells in a protein-free medium”; Stoll *et al.*, 1996, *J. Biotechnology*, vol. 45, p. 111-123 “Systematic improvement of a chemically defined protein-free medium for hybridoma growth and monoclonal antibody production.” This work is related specifically to monoclonal antibody production and does not disclose adapting such media for IVF or ART uses. Other publications disclosing protein-free growth media, but not adapted or suitable for human IVF procedures are: Zang *et al.*, 1995, *Biotechnology*, vol. 13, p. 389-392, “Production of Recombinant Proteins in Chinese Hamster Ovary Cells Using A Protein-Free Cell Culture Medium”; and International patent application Publication no. WO 2005/120576 A2.

SUMMARY OF THE INVENTION

The present invention relates to novel protein-free embryo culture media formulations for retrieval of the human egg (i. Flushing medium), handling and storage of

5 spermatozoa (ii. Gamete Handling Medium), fertilization of the human egg (iii.

Embryo Culture Medium), gamete and embryo handling (iv. Gamete and Embryo

Handling Medium), and as well as development of the resulting zygote in culture (v.

Embryo Culture Medium) through 1 to 2 cleavage cycles for human application in the treatment and alleviation of sub-fertility and infertility using ART procedures. The

10 present invention includes the formulation of a single medium solution that can

replace and/or be used as a replacement/medium for all of the above novel media

solutions. The substantially protein-free media of the invention are distinct from the

protein-free media (termed ART-7 and ART-7b series therein) previously reported by

the same inventor (Ali, 1997; Ali et al., 2000). In previous reports, the inventor has

15 described the systematic investigation that led to the formulation of three precursor

media, namely, ART-1, ART-2 and ART-3. In the same report the inventor has

described how the substantially protein-free media series ART-7 and ART-7b evolved

from the ART-1 medium. The embryos generated by intra-cytoplasmic sperm

injection (ICSI) in the substantially protein-free ART-7 and ART-7b media series was

20 successfully used to treat 11 of 21 (52.4%) women of 39 years of age or lower to

achieve clinical pregnancies. The formulation of ART-1, ART-7 and ART-7b media

series has not been disclosed. The present invention provides substantially protein-

free media that has been formulated from, but is different from, the precursor ART-3

medium. The development of the precursor ART-3 medium, but not its composition,

25 has been described by the inventor (Ali, 1997; Ali et al., 2000).

In exemplary embodiments of this invention, a series of substantially protein-free media are specifically disclosed, termed the PFM-11 series herein. These media have the specific advantage of being of uniform composition devoid of potentially

30 hazardous non-uniform biological components that may harm gametes and embryos,

and could potentially transmit deadly presently known and hitherto unknown diseases

to the patients undergoing ART treatment, the babies conceived through such

treatment and to healthcare workers involved in ART treatment procedures, as well as

to comply with rigorous new norms and regulations governing in vitro fertilization techniques and methods.

The completely defined nature of the substantially protein-free media formulations

5 according to the present invention also is useful in facilitating research into pre-implantation embryo metabolism that was previously hampered due to the undefined nature and non-uniform composition of currently available embryo culture media formulations. The quality of day 2 human embryos generated in these substantially protein-free media (PFM) series using the continuous ultramicrodroplet (cUMD)

10 culture technique was statistically comparable to or better than embryos that developed in control commercial media formulations containing serum proteins. The PFM media were not toxic to human spermatozoa. Fertilization and subsequent development of human embryos by conventional IVF and intra-cytoplasmic sperm injection (ICSI) was comparable to controls.

15 The substantially protein-free media of this invention also are suitable for use in substantially protein-free human embryo culture and handling media and will ensure that protein-bound diseases are not transmitted to patients and newborns. The substantially protein-free media of the invention can be stored frozen at -20°C for up to 2 years without loss of efficacy.

The present invention has successfully overcome the need for added donor proteins in the culture system and provides a substantially protein-free media system for human ART application that utilizes substantially protein-free media products beginning

25 from egg retrieval, sperm preparation, insemination, fertilization, culture of resultant embryo and embryo transfer. This feature advantageously distinguishes the media of this invention from previously described “protein-free” culture systems, which utilized protein-containing media products at some point during the procedure and were thus not completely protein-free. As a consequence, these prior culture media can be contaminated with infectious agents and may face possible regulatory restrictions. The compositions, ranges, preferred ranges and particular specifications 30 of the various components of the present invention are set forth herein.

The present invention is the product of studies on the effect, tolerance and determination of optimal levels of individual components such as amino acids, antioxidants and chelators, osmolytes, vitamins, nutrients and alternate energy sources that could substitute in part the various roles of protein in vivo and in vitro and which exemplifies the functions of proteins in various protein-free handling media (e.g. sperm and oocyte handling media, and ICSI medium) and embryo culture media of this invention that is free of donor serum proteins. The optimal concentrations of the mentioned components were utilized to formulate a number of embryo culture media. Subsequently the best of these media showing optimal embryo development and significantly higher blastocyst hatching rate were identified and further refined to support human embryo development in the absence of added serum proteins. The substantially protein-free media of the invention are capable of supporting fertilization of human oocytes with spermatozoa prepared in the same medium. The resultant zygotes subsequently developed in the substantially protein-free medium to form viable early cleavage-stage human embryos. The transfer of these embryos led to normal pregnancies and live births.

The present invention thus successfully provides media useful throughout IVF/ART procedures (egg retrieval, sperm preparation, insemination, fertilization, culture of resultant embryo and embryo transfer) that overcome the need for added proteins in the culture system.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE

INVENTION

The present invention provides a series of nutrient solutions generically referred to as embryo handling and culture media that are devoid of any protein or protein-like components. These media are useful in the fertilization of human eggs and the subsequent development of fertilized eggs for up to 2 to 6 days outside the body in vitro. These nutrient solutions are called "embryo culture media". In addition to IVF and ART, other methodologies and techniques where it would be advantageous to use substantially protein-free nutrient growth media include stem cell technology and

therapy, cell/tissue regeneration and transplantation treatment procedures. The skilled worker will appreciate how to adapt the media set forth herein for such uses.

As disclosed in further detail below, use of the substantially protein-free media of this

5 invention for human gametes and embryos increased the efficacy of IVF and ART.

For example, the clinical pregnancy rate obtained using conventional IVF with substantially protein-free (PFM-11) media is increased to 50% (14 of 28) in all age groups and up to 53.8% (14 of 26) in women below 40 years of age; when compared to currently available embryo media containing proteins (33% pregnancies for all age

10 groups combined, as cited in Ali, 2004). The ICSI clinical pregnancy rate with substantially protein-free media of this invention is likewise increased to 46.2% (12 of 26) in all age groups and up to 54.5 % (12 of 22) in women below 40 years of age.

This difference is statistically significant in favor of the present substantially protein-free media, indicating that the substantially protein-free media of this invention is at

15 least equivalent to (and can be superior to) currently available commercial protein-containing medium currently available in the market. The number of babies born from embryos generated in the PFM-11 media by conventional IVF was 12 and by ICSI, 12. The babies born from embryos generated in the PFM-11 of the invention are apparently normal, intelligent, talkative and active (according to parental reports).

20 These statistical data were produced from a sample of at least 24 children from such births.

As used herein, the term “protein-free,” “essentially protein-free” and “substantially

25 protein-free” is intended to mean that the media was prepared using non-protein containing components (as described in further detail herein) and that no protein or protein-containing components were added to the media.

Exemplary embodiments of the present invention include a series of substantially protein-free media termed the PFM-11 series. These media series have a range of

30 specific media products, namely:

1. Substantially Protein-Free Flushing Medium
2. Substantially Protein-Free Gamete (Sperm) Handling Medium
3. Substantially Protein-Free Gamete (Oocyte/Embryo) Handling medium

4. Substantially Protein-Free Embryo Culture Medium

These media are used according to the methods set forth below, which are intended to illustrate the use of these media but are not limiting to any additional uses of the

5 media, *inter alia*, in IVF or ART methods known to those with skill in the art. In developing the media of the invention in experiments as set forth herein, each procedure was performed in parallel using conventional protein-containing media and exemplary PFM of the invention.

10 *Human Spermatozoa Preparation*

Male patients produced human semen by masturbation. Spermatozoa were prepared by the standard swim-up technique and occasionally with density gradient centrifugation. In the density gradient procedure, the recovered pellet was washed

15 twice and resuspended in culture medium. Spermatozoa preparations were made using test or control media as appropriate. Specifically, spermatozoa for ICSI or IVF were prepared in the substantially protein-free (PFM-11) medium for the experiments on the effect of protein deficiency on embryo development. The harvested spermatozoa were incubated under gaseous phase of 5% CO₂ in air at 37°C until use.

20

Ovarian Stimulation and Oocyte Retrieval

Ovarian stimulation was induced by subcutaneous injections of gonadotrophin-releasing hormone agonist (Buserelin; Suprefact; Hoescht, Frankfurt, Germany)

25 starting in the mid-luteal phase until menstruation or down-regulation was achieved. Down-regulation was considered to have been achieved when endometrial thickness was 4 mm or less, and when blood estradiol, progesterone and LH levels reached baseline values. Injections of follicle stimulating hormone (FSH; Metrodin; Serono, Rome, Italy) were administered for three days to initiate recruitment of follicles.

30 Thereafter, human menopausal gonadotrophin (Pergonal 500; Serono, Rome, Italy) was administered according to the response of the patient to stimulate follicular development. Ovulation was induced by an injection of a 10,000 IU of human chorionic gonadotrophin (hCG; Pregnyl; Organon, Oss, Holland) after achieving

follicular size of 16mm or more. Oocyte retrieval (OR) was performed 36h later by ultrasound guided vaginal aspiration.

Insemination and Culture Techniques

5

In Vitro fertilization (IVF)

Oocytes were individually inseminated with motile spermatozoa in micro-droplets of equilibrated culture medium at a concentration of 100,000/mL under equilibrated and embryo tested mineral oil (M8410, Sigma Chemicals, and USA). The inseminated

10 oocytes were cultured in an atmosphere of 5% CO₂ in air at 37°C. The following morning the oocytes were denuded with denuding pipettes (Stripper®, MidAtlantic Diagnostics, USA).

15 Intracytoplasmic sperm injection (ICSI)

In an alternative procedure (used, for example, for overcoming human male factor subfertility), ICSI was performed. Oocytes were prepared for ICSI by exposure to hyaluronidase solution for 30 seconds (80 IU/mL; Cat No. 10110010, Medi-Cult A/S,

20 Lersø Parkalle 42, 2100 Copenhagen, Denmark) and then transferred into equilibrated Medi-Cult™ or substantially protein-free media of the invention. Human oocytes were incubated for 5-7 minutes in the culture medium under gaseous phase of 5% CO₂ in air at 37°C and then denuded with a denuding pipette (ART No.1670, International Medical Products BV, Zutphen, Holland). The denuded human oocytes

25 were washed and finally incubated in culture medium for a further 30-60 minutes.

Commercially available ICSI pipettes were used for micromanipulation (Injection Needles, Cat No.130340B; Holding pipette, Cat No. 13030013; Laboratoire CCD, 60 Rue Pierre Charron, 75008 Paris, France or Injection needle, Cat. No.10-MIC; holding pipette, Cat. No. 10-MPH-120; Humagen, Charlottesville, Virginia 22911, USA). A 5µL ultra micro-droplet of PVP (Cat No. 1089001, Medi-Cult A/S, Denmark) was thinly spread at the centre of a petri dish (Cat No. 1006, Falcon Plastics, Becton Dickinson, Rutherford, New Jersey, 07070, USA). The PVP spread was surrounded by up to a maximum of five 10 µL micro-droplets of HEPES-buffered

IVF medium (Gamete-1009 Scandinavian IVF Sciences AB, Gothenberg, Sweden). The micro-droplets were overlaid with equilibrated and embryo tested mineral oil (M8410, Sigma Chemicals, USA). Sperm preparation (1 to 2 μ L) was introduced at the centre of the PVP spread. ICSI of mature oocytes were performed conventionally (for example, as set forth in Palermo *et al.*, 1992).

Ultra micro-droplet culture

Ultra micro-droplets (UMD; 1.5 to 2 μ L for culture of 3 to 7 human embryos) of culture medium were made in 4-well dishes previously filled with equilibrated mineral oil. The dishes were incubated under a gaseous phase of 5% CO₂ in air at 37°C. Medium was changed daily (UMD technique) with a finely drawn out Pasteur pipette using equilibrated culture medium in preliminary experiments but not changed (cUMD; or continuous UMD) until day 3 of culture in latter experiments.

15

Determination of Fertilization

Fertilization was determined 18h to 20h after ICSI or IVF. The oocytes were considered fertilized when two distinct pronuclei were visible. Fertilized oocytes were cultured in micro-droplets or ultra micro-droplets of equilibrated medium in an atmosphere of 5% CO₂ in air at 37°C. Cleavage and embryo quality were assessed after 24h.

Zygote Arrest rate

25

This parameter was utilized to determine whether the culture medium according to the present inventions causes developmental blocks at the 1-cell stage. The efficacy of the medium can be determined using this assay. Higher proportion of developmental blocks at the 1-cell stage will suggest the medium to be deficient or less efficacious.

30

Determination of Day 2 Cleaved Embryo Quality

Two parameters were employed to determine the embryo quality of day 2 embryos, viz.: the average blastomere score and the average embryo grade. These were determined as follows:

5 (i) Average blastomere score =
$$\frac{\text{Total number of blastomeres}}{\text{Total number of embryos}}$$

(ii) Average embryo grade =
$$\frac{\text{Total of embryo grades}}{\text{Total number of embryos}}$$

10

Since most healthy day 2 human embryos generally attain the 4-cell stage, an average blastomere score 4 and above (in a range of 2 to 6) was considered excellent. The embryos were graded according to a scale of 1 to 4, where the numerical 1 denoted poor quality, and 4, excellent quality. Embryos were graded collectively by a minimum of three (but preferably four) laboratory personnel. In addition, two other parameters included to establish differences in the rate of development of cleavage stage embryos. These were: (i) the proportion of embryos at or above 4-cell stage and (ii) the proportion of embryos that were grade 3 and above, on day 2 of culture in human studies.

15

Media Formulation

The formulation of an exemplary substantially protein-free media of the invention described herein was as follows. Albumin is known to be a source of fixed nitrogen and nutrients, an antioxidant, and to also have a number of other roles including membrane stabilization. Thus, the substantially protein-free media of the invention substituted albumin with other media components that perform, individually or collectively, the functions of albumin in culture. Individual embryo culture medium components used for these purposes include amino acids (including but not limited to alanine, asparagine, aspartate, cystine, glutamate, glutamine, glycine, histidine, isoluecine, leucine, lysine, methionine, phenylalanine, taurine, thereonine, tryptophan, tyrosine, valine, serine), antioxidants and chelators (*for example*, EDTA, reduced glutathione, tocopherol), alternate energy sources (such as fructose, glutamine, sodium pyruvate), osmolytes (including mannitol ands myoinositol), vitamins (ascorbic acid, cyanocobalamin, folic acid, tocopherol, etc) and elemental iron.


The concentration of individual components that supported the highest blastocyst development was considered optimal. Three media were formulated (ART-1, ART-2 and ART-3) in the development of the present invention. The inventor has previously described the various experiments undertaken to formulate the substantially protein-free media the ART-7 and ART-7b series from the culture medium ART-1 (Ali, 1997; Ali et al., 2000). The present invention, the PFM-11 protein-free media series were formulated from the culture medium ART-3 (Ali, 1997; Ali et al., 2000).

10 The final formulations for the various media products of the PFM-11 series are given in tables as set forth below. The concentrations of HEPES and bicarbonate in these media preparations vary, as set forth herein.

15 The substantially protein-free media of the invention comprise mineral salts, amino acids, antioxidants, antibiotics, energy components and buffer components (HEPES and bicarbonate for incubation under CO₂-supplemented conditions) that are similar to but distinct in their particulars from commercially-available, protein-containing media. The substantially protein-free media of the invention uniquely comprise a macromolecular species (methylcellulose and related polymers) and an optionally an un-metabolized sugar alcohol (D-mannitol).

20

The macromolecule comprising the substantially protein-free media of the invention is methyl cellulose of Formula I:



25 wherein

each R is independently CH₃ or H and n is between about 34 and about 43.

In the formula of methylcellulose above, R can be either H or CH₃. The extent of methoxy substitution ranges between 27.5-31.5% by weight. Degree of substitution (D.S., average number of substituent groups attached to the ring hydroxyls) is 1.5-1.9. This range gives maximum water solubility. Lower methoxy substitution results in 5 higher solubility in water and is preferred. The code “n” designates the level of polymerization, and optimally corresponds to a molecular weight of 14,000 Daltons relating to an approximate viscosity index of 15 cPS for a 2% solution in water at 20 °C (methylcellulose having these specifications is commercially available from Sigma Chemical Co., St. Louis, MO USA; www.sigmaaldrich.com); this corresponds to a 10 value of “n” that is between about 34 and about 43, preferably 36 to 39. The range of this component in the PFM is 0.01 to 0.15 g/L and the most preferred range is 0.09 to 0.1g/L. The optimal concentration is 0.1 g/L.

Methylcellulose can act as an antioxidant and osmolyte, it is non-toxic, enzyme 15 resistant and not cell permeable (Stewart et al., 1995). It helps to protect gametes and embryos against environmental insults, in particular attack by free radicals and osmotic pressure changes. It protects the cellular membrane from damage and helps maintain homeostasis. It is also a surfactant and lubricant. It contributes to increased viscosity of the medium, so that the gametes and embryos do not stick to sides of 20 dishes and inside pipettes and catheters. These physical attributes are supplied in the absence of serum proteins that normally perform these functions. This substance is inert and safe for human application.


Methylcellulose has been used elsewhere in human pharmaceutical and food industry 25 for well over 25 years without any side effects in human or animal studies. FAO/WHO and EU directives allow consumption of methylcellulose, and it has been used as a negative control in cancer research as it is known to be non-carcinogenic.

Methylcellulose is also used as a thickener and emulsifier in various food and 30 cosmetic products, and has medicinal uses, such as for treatment of constipation. It is not digestible, non-toxic, and not allergenic. Its pharmacological/clinical uses are as excipients and a carrier material. It is used in eye drops, as a bulking agent and laxative, used for diarrhea in functional bowel disease, to control ileostomy output

and as absorbent of toxic substances that causes infective diarrhea. It is also an antioxidant.

The substantially protein-free media of the invention also comprises D-mannitol. This 5 compound is poorly absorbed and is excreted almost unchanged in urine. As an antioxidant and an osmolyte this substance can protect the embryo from harmful environmental effects. It thus can further substitute for serum proteins, in part, to protect the embryos against adverse conditions. It is of interest to note that D-mannitol exerts a positive effect on mouse blastocyst development *in vitro* even in the presence 10 of protein in the media.

In the exemplary substantially protein-free media of this invention (PFM-11), D-mannitol is present at a preferred concentration of 2.8 micromolar (the most preferred concentration) within the range of 0.056 to 6.9 micromolar. The more preferred range 15 is 1.4 to 5.5 micromolar; even more preferred within the range of 2.5 to 3.0 micromolar. The most preferred concentration is about 2.7 micromolar. The empirical formula of D-mannitol is C₆H₁₄O₆ and has the structural formula:



20 D-mannitol is a food additive, used in cakes, confectionaries and sweets; being sweeter than sucrose, it is considered an alternative sweetener for diabetics. It is an osmolyte and antioxidant. It has been used in high concentrations to treat acute stroke

for well over 30 years. Hypermolar concentrations of this compound are used to treat severe brain damage and elevated intracranial pressure. It is a diuretic and is used for diuresis in instances of poisoning, or to measure extracellular fluid compartment. It also has laxative effects in mammals including man. No adverse effects in man have 5 been reported as a result of clinical application of D-mannitol as a therapeutic agent. It is also non-cytotoxic and non-mutagenic in several species. D-mannitol is poorly absorbed and is excreted (Milde, 1965; Widdowson and Dickerson, 1965). The inclusion of this substance in culture medium increased blastocyst development in the mouse even in the presence of serum proteins, suggesting a positive role in embryo 10 culture as an antioxidant and osmolyte (Ali, unpublished observations). Its use is permitted by the FDA and the EU, and FAO/WHO has concluded it to be safe for human consumption.

In the solutions and formulations of the invention, D-mannitol of Formula II is present 15 at a concentration of from about 0.05 micromolar to about 6.9 micromolar. The more preferred range is 1.4 to 5.5 micromolar. The most preferred concentration is about 2.8 micromolar.

The skilled worker will appreciate that the role of serum proteins are numerous, and 20 in its absence, one or two components by themselves may not provide a complete substitute for proteins in the medium. Thus, the skilled worker will recognize that the invention does not merely provide media supplemented with methylcellulose and D-mannitol. Rather, the invention provides media comprising in addition a complex mixture of additional components, preferably in optimal concentrations, and the 25 selective exclusion of commonly used media components that have proven detrimental to embryos in culture. Some of these include:

- (i) Including unique concentrations for two amino acids (L-taurine and glutamine) that are the principal providers of nutrition and osmotic balance. 30 These amino acids are provided at concentrations within the range of 1mM to 30mM L-taurine and 1mM to 50mM L-glutamine, more preferably 10mM to 30mM L-taurine and 10mM to 30mM L-glutamine and most preferably 20mM each of L-taurine and L-glutamine.

(ii) Including as energy sources any one or plurality of compounds (including D-glucose, fructose, or pyruvate) that are less likely to cause developmental blocks in human embryos. Optimal concentrations of fructose are from about 0.5 mM to 6.0mM fructose, with a more preferred concentration being from about 1mM to 5.6mM, and a most preferred concentration of about 5.1mM. The optimal concentrations of the remaining two energy sources are given elsewhere in this document.

10 (iii) Including the following concentrations of certain amino acids.

Name	Optimum range Conc. in PFM (mM)	Preferred range (mM)	Most preferred Conc. (mM)
15 L-alanine	0.1-10	0.45-0.55	0.5
L-arginine	0.018-0.18	0.072-0.125	0.072
L-cystine.2HCl	0.0025-0.025	0.01-0.02	0.01
L-glutamate	0.01-1.0	0.45-0.55	0.5
20 L-glycine	0.1-1.0	0.2-0.3	0.25
L-histidineHCl.H ₂ O	0.005-0.05	0.02-0.04	0.02
L-isoleucine	0.01-0.1	0.04-0.08	0.04
L-leucine	0.01-0.1	0.04-0.08	0.04
L-lysine HCl	0.0125-0.125	0.05-0.1	0.05
25 L-methionine	0.0025-0.025	0.01-0.02	0.01
L-phenylalanine	0.005-0.05	0.02-0.04	0.02
L-threonine	0.01-0.1	0.04-0.08	0.04
L-tryptophan	0.00125-0.0125	0.005-0.01	0.005
L-tyrosine.2Na2H ₂ O	0.005-0.05	0.02-0.04	0.02
30 L-valine	0.01-0.1	0.04-0.08	0.04

(iv) Including the following concentrations of water soluble vitamins.

Name	Optimum range Conc.	Preferred range (mM)	Most preferred (mM)
35 Choline chloride	0.004 - 0.007	0.004 - 0.005	0.004

D-Biotin	0.0024 - 0.004	0.0024 - 0.003	0.0024
Folic acid	0.0014 - 0.0023	0.0014 - 0.0016	0.0014
Myoinositol	0.0067- 0.011	0.0067- 0.0078	0.0067
Niacinamide	0.005 - 0.008	0.005 - 0.0057	0.005
5 D-pantothenic Acid.%Ca	0.0025 - 0.004	0.0025 - 0.003	0.0025
Pyridoxine HCl	0.003 - 0.005	0.003 - 0.0034	0.003
Riboflavin	0.00016 - 0.00027	0.00016 - 0.00019	0.00016
Thiamine HCl	0.0018 - 0.003	0.0018 - 0.002	0.0018

10

—
The range for vitamin B12 is as follows:

15

Name	Optimum range Cone. in PFM (pM)	Preferred range (pM)	Most preferred Conc. (pM)
Vitamin B12	443 – 885	590-738	616

20

—
(v) Including vitamin E (Vitamin E Type 6, Sigma Chemical Co.) at a concentration of from 5 micromolar to 20 micromolar, more preferably 8 micromolar to 12 micromolar, even more preferably 10 micromolar.

25

(vi) Excluding L-asparagine, L-aspartate and L-serine that can be detrimental to the development of zygotes and early cleavage stage embryos.

30

(vii) Excluding elemental iron, that can be detrimental to embryo development.

35

(viii) Including reduced glutathione (GSH) at a concentration of 60 micromolar to 500 micromolar, and more preferably 250 micromolar to 350 micromolar, and even more preferably 300 micromolar.

Other compounds

Concentration in final solution

	Description	Range	Preferred Range	Most Preferred Conc.
5	D-glucose	0.75-1.0	0.75-0.90	0.78 g/L
	Sodium chloride	6.12-6.95	6.12-6.19	6.171 g/L
10	Potassium chloride	0.35-0.4	0.35-0.36	0.355 g/L
	Calcium chloride –	0.23-0.27	0.23-0.24	0.235 g/L
	Magnesium sulfate	0.086-0.098	0.086-0.087	0.087 g/L
	Sodium dihydrogen phosphate	0.107-0.122	0.107-0.109	0.108 g/L
	EDTA, Sodium	0.0416-0.043	0.0416-0.042	0.0418 g/L
15	Sodium bicarbonate*	2.2	2.2	2.2 g/L
	Sodium lactate, 60% syrup	1.9	1.9	1.9 ml/L
	Phenol red	0.011	0.011	0.011 g/L

***Sodium bicarbonate**

20	In Fertilization and Culture media	2.2 g/L (26.2 mM) mM)	2.2 g/L (26.2 mM)	2.2 g/L (26.2
	In Gamete Handling Media	2.2 g/L (26.2 mM)	2.2 g/L (26.2 mM)	2.2 g/L (26.2
	In Flushing medium	2.2 g/L (26.2 mM)	2.2 g/L (26.2 mM)	2.2 g/L (26.2

25 mM)

**** HEPES**

30	In Fertilization and Embryo Culture media	0 mM
	In Gamete Handling Media	15 mM
	In Flushing medium	25 mM

35 The PFM-11 media series set forth herein differs from conventional embryo culture media in at least the following aspects:

1. Absence of adding donor serum proteins to media – this eliminates the potential for disease transmission, making the PFM-11 media series according to the present invention non-hazardous.

2. The presence of methylcellulose and D-mannitol.
3. Alterations in the composition of the media regarding species and concentrations of amino acids, antioxidants, vitamins, energy sources and mineral salts, all optimized for embryo culture.

5 Among the advantages of the media of the invention are reduced risks to patients/healthcare workers. The substantially protein-free media of the invention is also advantageous because it reduces risks to patients, their babies and healthcare 10 workers exposed to the media. While certain risks remain these are minor in comparison with the infectious risks avoided using the substantially protein-free media of the invention. These risks include allergies to the components used. The chance of this occurring is very unlikely because the ingredients except for antibiotics are mostly inert and non-reactive. All ingredients are in minute concentrations not 15 likely to elicit an allergic response.

Clinical Trial

20 A clinical trial was approved for the transfer of embryos generated in the final formulations of the PFM-11 media series. The clinical trial also investigated the difference in efficacy of the PFM-11 media series when fertilization was performed either by conventional IVF and ICSI.

Statistical Analysis

25 Statistical comparisons were performed with the 2-sample *t*-test or two by two tables. A value of *p*<0.05 or less was considered statistically significant.

RESULTS

Culture Characteristics of Sibling Human Oocytes When Inseminated By Conventional IVF In PFM-11 Protein-Free Medium And MediCult™ Embryo Culture Media

The fertilization rate of sibling oocytes after conventional IVF was similar in the control MediCult™ and the PFM-11 media (Table A). A higher proportion of zygotes arrested in the control MediCult™ medium (8.1%) compared to (2.2%) the PFM but this was not statistically significant. The quality of cleaved day 2 human cIVF sibling embryos generated in the PFM-11 in ultramicroplet culture was significantly superior in terms of blastomere number and grade compared to those generated in the control medium. In the PFM-11 medium the average blastomere number was 3.4 and the average embryo grade was 3.1 (n=116). In the control medium (MediCult™) the average blastomere number was similar (p=0.865) at 3.4 and the average embryo grade was significantly lower (p=0.001) at 2.7 (n=118). The proportion of embryos that had attained the ≥ 4-stage was similar in both groups but the proportion of embryos that were above grade ≥3 was significantly higher the PFM-11 protein-free group.

15 **Table A: Quality of day 2 human sibling embryos generated by conv. IVF in
protein-free PFM-11 medium**

20	Medium	Fertil %	Arrested @ 1-cell stage %	Blast number Mean (1SD)	Grade Mean (1SD)	% ≥ 4 cells	% ≥3 Grade
25	PFM-11 (- protein)	85.3 (116/136)	2.2	3.4 (1.0)	3.1 (0.9)	58.4	74.3
	MediCult™ (+ protein)	79.2 (118/149)	8.1	3.4 (1.0)	2.7 (0.8)	56.2	58.1
30	Significance	p=0.235	p=0.052	p=0.865		p=0.001	p=0.847 p=0.017

—
[Embryo grade: 4=excellent; 3=good; 2= fair; 1=poor]; Control embryos generated by ICSI or IVF; Blast = blastomere; **Findings:** Test medium superior to control

35 **Culture Characteristics of Sibling Human Oocytes When Inseminated By
Conventional IVF In PFM-11 Protein-Free Medium And MediCult™ Embryo
Culture Media**

The fertilization rate of sibling oocytes after ICSI was statistically higher in the protein-free PFM-11 medium compared to the control MediCult™ (77.8% vs 69.4%, p=0.043;

Table B). The developmental arrest at the zygote stage was lower in the PFM-11

5 medium compared to the MediCult™ medium but this difference was not statistically significant (2.8 vs 6.3% respectively, p=0.088). The quality of cleaved day 2 human ICSI sibling embryos generated in the PFM-11 medium cultured in ultra micro-droplets was significantly superior in terms of blastomere score compared to those generated in the control medium (3.8 vs 3.3 respectively, p=0.001). The average 10 embryo grade of embryos generated in the PFM-11 medium was similar to that of embryos generated in the control medium (2.9 vs 2.8, p=0.080). The proportion of embryos that had attained the ≥ 4-stage was similar in both groups but the proportion of embryos that were above grade ≥3 was significantly higher the PFM-11 protein-free group.

15

Table B: Quality of day 2 human sibling embryos generated by ICSI in protein free PFM-11 medium

Medium	Fertil %	Arrested @ 1-cell stage %	Blast number Mean	Grade Mean (1SD)	% ≥ 4 cells	% ≥3
PFM-11 (- protein)	77.8 (196/252)	2.8	3.8 (1.2)	2.9 (0.7)	71.4	63.5
MediCult™ (+ protein)	69.4 (175/252)	6.3	3.3 (1.1)	2.8 (0.8)	54.8	58.6
Significance	<u>p=0.043</u> p=0.413		<u>p=0.088</u>	<u>p=0.001</u>	p=0.080	<u>p=0.002</u>

25
30
35
(Embryo grade: 4=excellent; 3=good; 2= fair; 1=poor); Blast = blastomere; Control embryos generated by ICSI/IVF; **Findings:** test medium superior to control

Clinical Trial

40 The conventional IVF clinical pregnancy rate with substantially protein-free PFM-11 media is increased to 50% (14 of 28) in all age groups and up to 53.8% (14 of 26) in women below 40 years of age; when compared to currently available embryo media

containing proteins (33% pregnancies for all age groups combined; cited from Ali, 2004). The ICSI clinical pregnancy rate with substantially protein-free PFM-11 media is likewise increased to 46.2% (12 of 26) in all age groups and up to 54.5 % (12 of 22) in women below 40 years of age. This difference is statistically significant in favor of 5 the present substantially protein-free PFM-11 media, indicating that the substantially protein-free media of this invention is at least equivalent to (and can be superior to) currently-available commercial protein-containing medium currently available in the market. The number of babies born from embryos generated in the PFM-11 media by conventional IVF was 12 babies (30.8%, that is 8 of 26 women treated delivered; or 10 61.5% 8 of 13 of the pregnant women delivered; one pregnancy lost to follow-up) and by ICSI, also 12 babies (36%, that is 8 of 22 women treated delivered or 66.7% (8 of 12) or of the pregnant women delivered. The babies born from embryos generated in the PFM-11 of the invention are apparently normal, intelligent, talkative and active (according to parental reports). These statistical data were produced from a sample of 15 at least 24 children from such births.

SIGNIFICANCE OF RESULTS

The availability of a chemically defined medium for generating viable early cleavage 20 stage human embryos improves the safety of patients undergoing assisted reproduction treatment (ART) because, *inter alia*, potentially hazardous donor protein can be eliminated from the media. This is a particular concern with regard to transmission of pathogenic diseases, in particular viral diseases, such as human acquired immunodeficiency disease syndrome (AIDS) and hepatitis, or Creutzfeldt- 25 Jakob disease (CJD) transmitted by prions, or others in blood-derived products has led a number of embryologists worldwide to seek (unsuccessfully) alternative(s) to donor protein for their embryo culture and handling procedures. The danger is certainly real: in the past an epidemic of hepatitis B occurred in about 200 IVF patients that received embryos cultured in medium containing pooled sera contaminated with hepatitis B 30 virus (van Os et al., 1991). More recently the scientific community was confronted with the dilemma of having to inform their patients that a commercial preparation of a culture medium used for embryo culture and handling may be contaminated with albumin donated by a person who later died of CJD (Kemmann, 1998).

In addition to pathogens and prions, serum or proteins derived from serum such as albumin may contain embryotoxic factors, as in patients with endometriosis, recurrent abortion and those suffering from unexplained infertility (Miller et al., 1995). These embryotoxic factors, the nature of which remains to be elucidated, have been shown

5 to be detrimental to embryos in vitro (Miller et al., 1995; Fein et al., 1995). Moreover, the use of serum or albumin raises the problem of reproducibility, because batch-to-batch variation in sera or albumin has been well recognized. An inability to control quality of different batches of sera can affect quality of embryos generated. Problems associated with embryotoxicity and/or non-reproducibility of the quality of serum
10 proteins are avoided, of course, when chemically defined medium is used instead of culture media containing donor proteins.

The quality of embryos generated in the PFM-11 culture medium under ultra microdroplet culture conditions was similar to or somewhat better than those

15 generated in the control medium containing proteins when performed under similar culture conditions. More importantly the fertilization rate, the zygote arrest rate and the quality of embryos generated in the novel substantially protein-free medium PFM-11 medium is not inferior to that observed in the control medium containing proteins for any of the parameters tested. On the other hand the control medium has proven
20 inferior in some of the parameter tested. The clinical trial showed that the PFM-11 medium is as efficacious or better than the control embryo culture medium containing proteins for day 2 embryos. Moreover it is completely safe as it is devoid of hazardous donor proteins of biological origin.

25 Improved quality of embryos generated in micro-droplets have been earlier shown to be due to autocrine and paracrine effects of certain growth factors released by the embryo into the medium (Paria and Dey, 1990; Paria et al., 1991; Canesco et al, 1992). Indeed, the addition of hIGF-1 to culture medium was found to enhance human pre-implantation development, in particular, blastocyst formation was significantly
30 improved (Lighten et al., 1998). Recently communal culture of human embryos was reported to result in higher implantation and pregnancy rate in human assisted reproduction (Almagor et al., 1996). The effect of growth factors in pre-implantation development has been extensively reviewed by Kane et al. (1997). Evidence from previous studies on the effect of growth factors indicates a lack of clear growth

pattern on embryo development in vitro. Autocrine growth factors may be essential for embryo development under stressful culture conditions. Growth factors that may show positive effects on embryo pre-implantation development include CSF-1; IGF-I; IGF-II; TGF- α , and possibly LIF.

5

Fertilization in human oocytes was found to occur in PFM-11 with spermatozoa prepared in a PFM-11. Capacitation was not affected. PFM-11 has been shown herein to be efficacious in promoting fertilization in the absence of serum proteins in the medium. The high fertilization rate during conventional IVF using PFM-11 indicated that the absence of protein in the medium did not impair sperm capacitation, fertilization, and embryo development *in vitro* and subsequent viability *in utero*. In spite of a deficiency of proteins in the medium, sperm capacitation, penetration of sperm into the oocyte and fertilization occurred.

15 The clinical pregnancy, delivery and abortion rates were significantly better in the test group (using substantially protein-free media of the invention) compared to the control (using conventional protein-containing media). Also, the proportion of pregnant women in the test group that delivered was similar to the world average reported for cleavage stage embryo transfers (DeMouzon and Lancaster, 1997).

20

The results of the experiments disclosed herein revealed that: (i) penetration of spermatozoa into oocytes during conventional IVF, gamete interaction and subsequent fertilization in the human following both conventional IVF or ICSI were not impaired using substantially protein-free media PFM-11 of the invention, (ii) the resultant early 25 cleavage stage embryos were viable and capable of eliciting clinical pregnancies, (iii) the quality of early cleavage stage embryos generated in the PFM-11 was in general similar or somewhat superior to that generated in control medium containing serum proteins.

30

Stringent purification and sterilization measures employed for preparations of animal origin cannot exclude with absolute certainty the possibility of transmission of unknown pathogens (Truyen et al., 1995). With the formulation of substantially protein-free media and the substitution of hyaluronidase with a recombinant hyaluronidase such as Cumulase®, it is now possible to perform the entire laboratory

assisted reproduction procedures with no apparent risk of transmission of disease to IVF patients.

In conclusion, the substantially protein-free PFM-11 medium of the present invention was demonstrated to be efficacious, with sperm penetration into oocyte during conventional IVF, and fertilization following ICSI or conventional IVF, being uncompromised. Pregnancies were also not compromised when embryos generated in the PFM were transferred, with a clinical pregnancy rate of above 50% in women 39 years and below. Overall the PFM-11 medium was equally efficacious if not superior to medium containing serum proteins in the generation of viable human embryos. The routine use of PFM in human assisted conception procedures can eliminate the potential risk of transmission of diseases through protein-bound pathogens or dangerous prions. This study showed that it is possible to generate viable human embryos in a medium devoid of added protein in a cell-free culture system.

15

PREPARATION OF PFM-11 OF THE INVENTION

Basal Salt Solution (BSS) Stock Solution 1:

20 Preparation of stock solution of basal salts (Basal Salt Solution – BSS) to be used is a final formulation of PFM Culture medium, Gamete handling medium and Flushing medium.

	Raw material component	Quantity (g/L)
1.	Calcium chloride · 2 H ₂ O	2.65 (23.8 mM)
2.	Magnesium sulphate (anhyd.)	0.9767 (8.1 mM)
3.	Potassium chloride	4.0 (53.7 mM)
4.	Sodium chloride	69.53 (1.2 M)
5.	Sodium phosphate monobasic (anhyd.)	1.22 (10 mM)
6.	D-glucose	10.517 (58.4 mM)
7.	Phenol red	0.11 (0.31 mM)
8.	WFI (Water for injection)	950 mL

Preparation procedure:

- 5 1. Rinse mixing container with WFI (Water for injection, 18.2 MegaOhms resistivity) before preparation of stock solution.
- 10 2. Add component 1 to 1000 mL of WFI water as it is extremely hydroscopic.
- 15 3. Add components 2-7 in order, mixing continually and make up final volume to 1000 mL using WFI.
4. Sterile filtration to be carried out immediately after all solutes are fully dissolved. Do not filter if solutes remain. 0.1 micron filters can be used but avoid excessive pressure when filtering. The Stock 1 solution can be pre-filtered with 0.2 micron filter followed by 0.1 micron filter.
- 20 5. There should be no precipitate or cloudiness post-filtration.
- 25 6. Fill into containers of suitable volume, e.g. bottles.
7. Cap bottles (preferably tamper-evident seal bottles).
8. Store in the dark between 2 and 6 degrees Celsius.

Storage and shelf life.

The Basal Salt Solution (BSS) stock was stable for two months when stored between

2 and 6 degrees Celsius. Always check stored BSS stock carefully for precipitates or

30 30 cloudiness before use. Discard if precipitates occur or the solution has turned cloudy during storage.

Inclusion volumes of BSS stock 1 in final product:

35 35 Where sodium bicarbonate is the predominantly active buffer component in the formulation (Culture media products), 70 ml** of BSS must be present in every 1000 ml of final formulation prepared.

Where HEPES is the predominantly active buffer component in the formulation (sperm/flushing media products), 50 mL of BSS is added for every 1000 mL of final formulation prepared.

5

Adjusting the osmolality of final medium with BSS or WFI water to increase or decrease (respectively) the osmolality of the medium:

Should the osmolality of the final medium be 285 mOsmols when final volume is less

10 than 1000 ml (i.e. after adding BSS, BAAS and BVS); then separately dilute a small amount of BSS with WFI. Use 1 volume of BSS with 9 volumes of WFI water to give a working BSS (WBSS). Adjust osmolality of WBSS to 285 mOsmols. Make up final volume of final medium to 1000 ml with adjusted WBSS.

15 For composition of final formulation from all stock solutions – see Stock 4 instructions.

Basal Amino Acid Solution (BAAS) Stock Solution2:

20 This protocol was used to prepare a 1 litre stock solution of basal amino acids in solution (Basal 5 Amino Acid Solution – BAAS) for inclusion in final formulations of PFM Culture medium, Gamete Handling medium and Flushing medium.

Reagents

25

	Raw material component	Quantity (g/L)
1.	L-arginine HCl	1.26 (6.0 mM)
2.	L-cystine (L cystine.2HCl)*	0.313 (1.0 mM)
3.	L-histidine HCl.H2O	0.42 (2.0 mM)
4.	L-isoleucine	0.52 (4.0 mM)
5.	L-leucine	0.52 (4.0 mM)
6.	L-lysine.HCl	0.752 (4.1 mM)
7.	L-methionine	0.15 (1.0 mM)

8.	L-phenylalanine	0.32 (1.9 mM)
9.	L-threonine	0.48 (4.0 mM)
10.	L-tryptophan	0.1 (0.49 mM)
11.	L-tyrosine (L-tyrosine.2Na 2H ₂ O)*	0.519 (2.0 mM)
12.	L-valine	0.46 (3.9 mM)
13.	WFI (Water for injection)	1000 mL

Instructions for preparation of Stock 2 BAAS

5 1. Rinse mixing containers with WFI (Water for injection, 18.2 MegaOhms resistivity)

2. Dissolve components 1, 2 and 4-11 in 400 mLs of WFI.

Compliance with Supplier's storage recommendations is required as difficulties with

10 solubility may be experienced with less than optimal conditions. If a precipitate occurs discard and start again using 25 mL of 0.11 N NaOH and 275 mL of WFI. Sodium hydroxide is only used as last option.

15 3. L-histidine HCl-H₂O and L-valine can be difficult to dissolve and in such

circumstances solubility can be achieved by using 1N HCl. Components 3 and 12 should be dissolved in their own separate 300 mL volumes of WFI. Use only the smallest volume possible of 1N HCl to achieve solubilization. The maximum amount of 1N HCl (embryo tested) that should be used is 25 mL in 300 mL of WFI. Avoid heating water, alkalis and acids above 37 degrees Celsius to dissolve all components.

20

4. Once components 3 and 12 are fully dissolved, make up final volume of 1000 mL by gradually adding in a stepwise manner, a 200 mL portion of the water soluble amino acids made in step 2 to:

25 (a) 300 mL solution of L-histidine HCl.H₂O giving 500 mL in total volume.

(b) 300 mL solution of L-valine giving 500 mL in total volume.

5. Combine both 500 mL volumes, gradually in a stepwise manner with constant mixing to avoid precipitation.

6. Sterile filter solution with 0.2 micron filter immediately.

5

7. Fill into containers of suitable volume (e.g. bottles).

8. Cap bottles (preferably tamper-evident seal bottles).

10 9. Store in the dark between 2-6 degrees Celsius.

Storage and shelf life.

15 The Basal Amino Acid Solution (BAAS) stock will keep for two months when stored between 2 and 6 degrees Celsius. Always check stored BAAS stock carefully for precipitates or cloudiness before use. Discard if precipitates or cloudiness occurs during storage.

20

Inclusion volumes of BAA stock in final product:

10 ml of BAA stock solution should be used per 1000 ml in the preparation of all final formulations.

25

L-Tyrosine.2Na 2H₂O and L-Cystine.2HCl were used because free form L-Tyrosine and

L-Cystine were not commercially available.

30 Basal Vitamin Solution (BVS) Stock Solution 3.

The protocol according to this example prepares a 1 litre stock solution of basal vitamins in solution (Basal Vitamin Solution – BVS) for inclusion in final formulations of PFM Culture medium, Gamete handling medium and Flushing medium.

35

Reagents

	Raw material component	Quantity (g/L)
1.	Choline chloride	0.1 (0.7 mM)
2.	D-biotin	0.1 (0.4 mM)
3.	Myoinositol	0.2 (1.1 mM)
4.	Niacinamide	0.1 (0.82 mM)
5.	D-pantothenic acid	0.1 (0.46 mM)
6.	Pyridoxine HCl	0.1 (0.49 mM)
7.	Riboflavin	0.01 (26.5 µM)
8.	Thiamine HCl	0.1 (0.33 mM)
9.	Folic acid	0.1 (0.23 mM)
10.	WFI (Water for injection)	1000 mL

5 Instructions for preparation of Stock 3 (BVS)

1. Dissolve components 1-7 in 800 mL of WFI water (Water for injection, 18.2 MegaOhms resistivity). Should any component not dissolve, small amounts of 0.1 N or
 - 10 1 N NaOH can be used to achieve solvation as a last option.
2. Dissolve component 8 in 2.5 mL of 0.1 N NaOH and add carefully in a stepwise manner, with constant mixing to 800 mL containing components 1-7.
- 15 3. Make up volume to 1000 mL using WFI water with constant mixing and sterile filter immediately with 0.2 micron filter into suitable containers (e.g. 60 mL Nalgene bottles, tamper-evident bottles).
4. Cap bottles.

5. Store in the dark at -20 degrees Celsius.

Storage and shelf-life.

5 The Basal Vitamin Solution (BVS) stock can be stored for two months when stored at -20 degrees Celsius in 60 mL aliquots. Always check thawed BVS stock 3 carefully for precipitates or cloudiness before use. Discard if precipitates appear or the thaw solution appears cloudy.

10 Inclusion volumes of BVS stock into final product:

6.0 mL of BVS stock solution should be used per 1000 mL in the preparation of all final formulations prepared – see Stock 4.

15

Stock 4 solution and final formulation combinations.

20 The preparation of Stock 4 permits the final addition of the unique chemicals in this Formulation and the addition of specific volumes taken from Stock 1 Basal Salts Solution (BSS), Stock 2 Basal Amino Acid Solution (BAAS) and Stock 3 Basal Vitamin Solution (BVS). The protocol describes the addition of buffers and antibiotics that are specific to the media formulation being prepared, i.e. Culture Gamete handling and Flush media.

25

The following protocol prepares 1 litre of final formulation of Culture medium, Gamete handling or Flush medium. Larger batches may be prepared by appropriate scaling of quantities.

30 Reagents

	Raw material component	Quantity (g/l)
1.	Sodium pyruvate	0.02975 (0.27 mM)

2.	Fructose	0.92125 (5.1 mM)
3.	Glycine	0.01875 (0.25 mM)
4.	Glutathione (Reduced)	0.09225 (0.30 mM)
5.	D-mannitol	0.0911 (0.5 mM)
6.	EDTA (sodium tetra)	0.04175 (0.10 mM)
7.	L-alanine	0.04455 (0.5 mM)

	Raw material component			
		Quantity (g/l)		
		Quantity g/L	Quantity g/L	Quantity g/L
		Culture medium	Gamete handling Medium	Flush medium
8.	L-taurine	1.251 (1.0 mM) 20mM		
9.	L-glutamic acid mono sodium	0.0735 (0.39 mM)		
10.	Lactic acid Sodium salt	1.9 mL per liter		
11.	Vitamin B-12	100 µL/L		
12.	Vitamin E Type 6	333 µL/L		
13.	Stock 1 BSS	70 mL		
14.	Stock 2 BAAS	10 mL		
15.	Stock 3 BVS	6 mL		
16.	Gentamycin sulphate	1.5	4 (sperm) 1.5 (oocyte)	1.5
17.	Penicillin G	75	0 (sperm) 75 (oocyte)	75
18.	L-glutamine	1.461 (20mM)	1.461 (20mM)	1.461 (20mM)
19.	Sodium bicarbonate	2.2 (26.2 mM)	2.2 (26.2 mM)	2.2 (26.2 mM)

20.	HEPES	0.0	3.5745 (15 mM)	5.9575 (25 mM)
21.	Methyl cellulose	0.1	0.1	0.1
22.	WFI (Water for injection)	1000 mL	1000 mL	1000 mL

Instructions for preparation of Stock 4 and final formulation combination.

- 5 1. Rinse mixing vessel with WFI (Water for injection, 18.2 MegaOhms resistivity).
2. Dissolve components 1-11 in 700 mL of WFI mixing continuously whilst the additions are made.
3. Add components 12-15 and make up to 900 mL with WFI and continue mixing.
4. Add components 16-21 depending on preparation of the basic formulation type i.e.
- 10 PFM Culture media, PFM Gamete media handling medium and PFM Flush medium.
5. Adjust osmolality of all three media and separately filter sterilize using 0.2 microns pore-size filters, 0.1 micron filter can also be used but avoid using pressure. Do not use
- 0.1 microns filter if high pressure required for filtering the solution.
- 15 6. Sterile filter immediately into final packaging (Nalgene bottles).:
 - PFM Culture media is filled into 60 mL bottles
 - PFM Gamete handling media is filled into 60 mL bottles.
 - PFM Flush medium is filled into 125 mL bottles.
 (Bottles used are preferably tamper-evident bottles.)
- 20 7. Cap bottles.
8. Label bottles.
9. Store in the dark at between 2 and 6 degrees Celsius.

Footnote for Stock 4 and final formulation combination.

25 Check osmolality. If the osmolality is high, then adjust to 285 mOsmols by adding WFI pure water. The amount to be added is calculated as follows:

30
$$[\text{Osmolality of medium} - \text{Desired osmolality (i.e., 285)} / \text{Osmolality of medium}] \times$$

 volume of medium

Example: If osmolality of medium is 300 and volume of medium is 900 ml as above, then calculate as shown below:

5 $[300 - 285/300] \times 900 = 15/300 \times 900 = 45$

Therefore if you add 45 ml of water to the medium and then measure osmolality again, you should theoretically have an osmolality of about 285 plus/minus 2 or 3 units. Be very careful not to add too much water because the important ingredients in
10 the medium will become diluted and will affect the efficacy of the medium, even if you bring back the osmolality of the medium by adding more Stock 1 BSS solution.

However, if osmolality is lower than 285 mOsmols, add Stock 1 BSS only to the medium, approximately 2 to 3 mOsmols increases per ml of Stock 1 BSS but this may
15 not always be predictable. So be very careful not to add too much. Add a little at a time and measure osmolality.

REFERENCES

20 Ali J. (1997) Formulation of a protein-free culture system for the culture of human embryos: preliminary findings and pregnancies. In: Programme and abstract book of the
16th Annual Scientific meeting of the Fertility Society of Australia, 2-4 December
25 1997, Adelaide, Australia, p34 (Abstract)

Ali J. (2000) Investigation into the nutrient requirement of the human embryos: Successful formulation and clinical trial of a novel protein-free embryo culture medium.
30 Emirates Med J. 18:195-202

Ali, J (2004) Generation of viable human embryos in a protein-free embryo culture (ART-7b) medium enhances clinical pregnancy rate and prevents disease transmission in assisted reproduction. Middle East Fertil Soc. J. 9:118-127

Ali, J., Abdulkader, A., Shahata, M.A.M. et al. (1998) Term deliveries from human embryos conceived in a novel culture medium devoid of added protein. Middle East Fertil. Soc. J. 3, 40 (Abstract)

5

Ali, J., Whitten, W.K. and Shelton, J.N. (1993). Effect of culture systems on mouse early embryo development. Hum. Reprod. 8, 1110-1114

Almagor, M., Bejar, C., Kafka, I. et al. (1996) Pregnancy rates after communal

10 growth of pre-implantation human embryos in vitro. Fertil. Steril. 66, 394-397

Balakier, H. (1993) Tripromuclear human zygotes: the first cell cycle and subsequent development. Hum. Reprod. 8, 1892-1897

15 Barber, A. A. (1961) Inhibition of lipid peroxide formation by vertebrate blood serum. Arch. Biochem. Biophys. 92, 38-43

Barnes, RI-, Crombie, A., Gardner, D.K. et al. (1995) Blastocyst development and pregnancy after in vitro maturation of human primary oocytes, intracytoplasmic sperm injection and assisted hatching. Hum. Reprod. 10, 3243-3247

20 Bavister, B.D. (1995) Culture of pre-implantation embryos: facts and artifacts. Hum. Reprod. Update 1, 91-148

25 Biggers, J.D., Summers, M.C. and McGinnis, K.L. (1997) Polyvinyl alcohol and amino acids as substitutes for bovine serum albumin in culture media for mouse pre-implantation embryos. Hum. Reprod. Update 3, 125-135

30 Boyers, S.P., Diamond, M.P., Lavy, G. et al. (1987) The effect of polyploidy on embryo cleavage after in vitro fertilization in humans. Fertil. Steril. 48, 624-627

Canesco, R.S., Sparks, A., Pearson, R.E. et al. (1992) Embryo density and medium volume effects on early marine embryo development. J. Assist. Reprod. Genet. 9, 454-

457

Caro, C.M. and Trounson, A. (1986) Successful Fertilization, Embryo Development, and Pregnancy in Human in Vitro Fertilization (IVF) Using a Chemically Defined

5 Culture Medium Containing No Protein. *Journal of in Vitro Fertilization and Embryo Transfer*, vol. 3, no. 4. p. 215-217)

Carrillo, A.J., Lane, 13., Pridham, D.D. et al. (1998) Improved clinical outcomes for in vitro fertilization with delay of embryo transfer from 48 to 72 hours after oocyte

10 retrieval: use of glucose- and phosphate-free media. *Fertil. Steril.* 69,329-334

Centers for Disease Control. (1985) Fatal degenerative neurologic disease in patients who received pituitary derived human growth hormone. *MMWR Morb Mortal Wkly Rep* 34:359-60, 365-6.

15

Cholewa, J.A. and Whitten, W.K. (1965) Development of 2-cell mouse embryos in the absence of a fixed nitrogen source. *J. Reprod. Fertil.* 22, 553-555

Cockle SM, Aitken A, Beg F, Morrell JM, Smyth DG. (1989) The TRH-related

20 peptide by pyroglutamylglutamylprolinamide is present in human semen. *FEBS Lett* 1989;

252:13-7

Dandekar, P.V. and Glass, R.H. (1990) Development of two-cell mouse embryos in

25 protein-free and protein-supplemented media. *J. In Vitro Fertil. Embryo Transfer* 7, 107113

De Mouzon, J. and Lancaster, P. (1997). World collaborative report on in vitro fertilization. Preliminary data for 1995. *J. Assist. Reprod. Genet. (suppl.)* 14, 250S-

30 265S

Drakakis, P., Loutradis, D., Milingos, S. et al. (1996) The in vitro development of mouse embryos beyond the blastocyst stage into the hatching and outgrowth stage using different energy sources. *J. Assist. Reprod. Genet.* 13, 786-792

Du, Z.F. and Wales, R.G. (1993) Effect of culture from the zygote stage on the metabolism of glucose and glutamine by 2-cell embryos and blastocysts recovered from outbred or H female mice. *Reprod. Fertil. Dev.* 5, 555-565

5 Esmonde T, Lueck CJD, Symon L, Duchen LW, Will RG. (1994) Creutzfeldt-Jakob disease and lyophilised dura mater grafts: report of two cases. *J. Neurol Neurosurg Psychiat.* 56:999-1000.

Gardner, D.K. (1994) Mammalian embryo culture in the absence of serum or somatic 10 cell support. *Cell Biol. Int.* 18, 1163-1179

Gardner, D.K. and Lane, M. (1998a) Culture of viable human blastocysts in defined sequential serum-free media. *Hum. Reprod. (suppl.)* 13, 148-160

15 Gardner, D.K., Schooler, W.B., Wagley, L. et al. (1998b) A prospective randomized trial of blastocyst culture and transfer in in-vitro fertilization. *Hum. Reprod.* 13, 3434-3440

20 Green CM, Cockle SM, Watson PP et al. (1996) A possible mechanism of action for fertilization promoting peptide, a TRH-related tripeptide that promotes capacitation and fertilizing ability in mammalian spermatozoa. *Mol. Reprod. Dev.* 45:244-52.

Heyc, N., Henson, S., and Muller, N. (1994) Creutzfeldt-Jakob disease and blood 25 transfusion. *Lancet* 343:298-299.

Kane, KT., Morgan, P.M. and Coonan, C. (1997) Peptide growth factors and preimplantation development. *Hum. Reprod. Update* 3:137-157

30 Kemmann, E. (1998) Creutzfeldt-Jakob disease (CJD) and assisted reproductive technology (ART). *Hum. Reprod.* 13, 1777

Lane, M. and Gardner, D.K. (1997) Non-essential amino acids and glutamine decrease the time of the first three cleavage divisions and increase compaction of mouse zygotes in vitro. *J. Assist. Reprod. Genet.* 14,398-403

5 Li, J., Foote, R.H., Liu, Z. et al. (1996) Development of rabbit zygotes into blastocysts in defined protein-free medium and offspring born following culture and embryo transfer. *Theriogenology* 47,1103-1113

Lighten, A.D., Moore, GI., Winston, et al. (1998). Routine addition of human insulinlike growth factor-1 ligand could benefit clinical in-vitro fertilization culture. *Hum. Reprod.* 13, 3144-3150

10 Loutradis, D., Kallianidis, K., Drakkis, P. et al. (1992) Successful pregnancy in human IVF using BSA as a protein source in the transfer medium. *Assist. Reprod. Technol.* *Androl.* 3, 233-238

15 Mehta, T.S. and Kicssling, A.A. (1990) Development potential of mouse embryos conceived in vitro and cultured in ethylenediaminetetraacetic acid with and without amino acids or serum. *Biol. Reprod.* 43, 600-606

20 Milde, M.D. (1965). *Ann. Rev. Pharmac.* 5: 125

25 Moessner, J. and Dodson, W.C. (1993) The role of growth factors and proteins in media for in vitro fertilization. *Assist. Reprod. Rev.* 3, 63-67

30 Palermo, G., Joris, H., Devroey, P. and Van Steirteghem, A (1992) Pregnancies after intracytoplasmic injection of a single spermatozoon into oocyte. *Lancet.* 340:17-18

Paria, B.C. and Dey, S.K. (1990) Pre-implantation embryo development in vitro: cooperative interactions among embryos and the role of growth factors. *Proc. Natl. Acad. Sci. USA*, 87, 4756-4760

Paria, B.C, Tsukamura, H. and Iley, S.K. (1991) Epidermal growth factor-specific protein tyrosine phosphorylation in pre-implantation embryo development. *Biol. Reprod.* 45, 711-718

5 Parinaud, J., Milhet, P., Vieitez, G. et al. (1998a) Human sperm capacitation and in vitro fertilization in a chemically defined and protein-free medium (SMART1). *Proceedings of the 16th World Congress on Fertility and Sterility and the 54th Annual Meeting of the American Society for Reproductive Medicine*. October 4-9, 1998, San Francisco, California, United States of America, *Fertil. Stern.* (suppl.) p.S195-S196
10 (Abstract)

Parinaud, J., Vieitez, Milhet, P., et al. (1998b) Use of plant enzyme preparation (coronase) instead of hyaluronidase for cumulus cell removal before intracytoplasmic sperm injection. *Hum. Reprod.* 13, 1933-1935

15 Poiley, S.M. (1960) A systematic method of breeder rotation for non-inbred laboratory animal colonies. *Proc. Anim. Care Panel*, 10, 159-166

20 Quinn, P. (1994) Use of co-culture with cumulus cells in insemination medium in human in vitro fertilization (IVF). *J. Assist. Reprod. Genet.* 11, 270-277

Saito, H., Hirayama, T., Koike, K. et al. (1994) Cumulus mass maintains embryo quality. *Steril.* 62, 555-558.

25 Schramm, R.D. and Bavister, B.D. (1996) Development of in-vitro-fertilized primate embryos into blastocysts in a chemically defined, protein-free culture medium. *Hum. Reprod.* 11, 1690-1697.

30 Serta, R.S., Sakellariou, M., Kiessling, A.A. et al. (1997) Outcome of human embryos conceived and cleaved in protein-free culture conditions. *Proceedings of the 53rd Annual Meeting of the American Society for Reproductive Medicine*. October 1.

Stewart GJ, Wang Y, Niewiarowski S. (1995) Methylcellulose protects the ability of anchorage-dependant cells to adhere following isolation and holding in suspension.

Biotechniques. 19(4):598-604

Meeting of the American Society for Reproductive Medicine. October 188-22, 1997, Cincinnati, Ohio, United States of America, (suppl.) 5214-5215 (Abstract)

5

Spindle, A. (1995) Beneficial effect of taurine on mouse zygotes developing in protein-free culture medium. Theriogenology 44,761-772

10 Staessen, C., Janssenswillen, C., De Clerck, E. et al. (1998) Controlled comparison of commercial media for human in-vitro fertilization: Menezo B2 medium versus Medi-Cult universal and BM1 medium. Hum. Reprod. 13, 2548-2554

15 Tesarik, J., Pilka, L., Drahorad, J. et al. (1988) The role of cumulus cell-secreted proteins in the development of human sperm fertilizing ability: implications in IVF. Hum. Reprod. 3,129-132

Parinaud, J., Milhet, P., Vieitez, G. and Richoilley, G. (1999). Use of a medium devoid of 5 any human or animal compound (SMART2I) for embryo culture in intracytoplasmic sperm injection. J. Assist. Reprod. Genet. 16: 13-16.

20

Truyen, U., Parrish, C.R., Harder, T.C. et al. (1995) There is nothing permanent except change. The emergence of new viral diseases. Vet. Microbiol. 43, 103-122. Cited in: Parinaud et al., 1998b

25 Van Os, H.C., Drogendijk, Aat, C. Fetter, W.P.F. et al. (1991) The influence of contamination of culture medium with hepatitis B virus on the outcome of in vitro fertilization pregnancies. Am. J. Obstet. Gynecol. 165, 152-159


30 Vidlakova, Erazimova, J., Horky, J. et al. (1972) Relationship of serum antioxidative activity to tocopherol and serum inhibitor of lipid peroxidation, Chim. Acta 36, 61-66

Wayner, Dam, Burton, G.W., Ingold, K.U. et al. (1987) The relative contribution of vitamin F, mate, ascorbate and proteins to total peroxy radical•trapping antioxidant activity of human blood plasma. *Biochim. Biophys. Acta* 924, 408-419

5 Widdowson, E.M. and Dickerson, J.W.T. (1964). In: Comar, L.C. and Brommer, F. (eds.), *Mineral Metabolism*, New York – London, Vol. IIA, p. 13.

We claim:

1. A substantially protein-free cell culture medium for human reproductive cells,
5 suitable for use throughout *in vitro* treatment of human reproductive cells to be used in *in vitro* fertilization and other assisted reproduction technologies, comprising mineral salts, amino acids, antioxidants, vitamins, nutrients, antibiotics, D-mannitol, and methylcellulose that has a molecular weight of 14,000 Daltons.
- 10 2. A substantially protein-free medium according to claim 1, wherein said methylcellulose is characterized so that a 2% solution has a viscosity of 15 centipoise at 25°C.
- 15 3. A substantially protein-free medium according to claim 1, wherein said methylcellulose is of formula I:



wherein each R is independently H or CH₃ and n is an integer having a value from about 34 to about 43 and wherein methoxy substitution is from 27.5% to 31.5% by weight.

20 4. A substantially protein-free medium according to claim 4, wherein average number of CH₃ substituents attached to each sugar moiety of the compound of formula I is 1.5 to 1.9.

5. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein said methylcellulose is present in the solution at a concentration of from 0.01 g/L (0.71 micromolar) to 0.5 g/L (0.036 mM).

5

6. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein said methylcellulose is present in the solution at a concentration of from 0.01 g/L (0.71 micromolar) to 0.15 g/L (0.00011 mM).

10 7. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein said methylcellulose is present in the solution at a concentration of about 0.1 g/L (0.0071 mM)).

15 8. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein the amino acids are L-arginine, L-cystine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-threonine, L-tryptophan. L-tyrosine, L-valine, L-alanine, L-taurine, L-glutamic acid, L-glutamine or glycine, or any combination thereof.

20 9. The substantially protein-free medium according to claim 8, wherein the amino acids are present at concentrations between: about 0.018 mM and about 0.18 mM for L-arginine HCl; about 0.0025 mM and about 0.025 mM for L-cystine.2HCl; about 0.005 mM and about 0.05 mM for L-histidine HCl.H₂O; about 0.01 mM and about 0.1 mM for L-isoleucine; about 0.01 mM and about 0.1 mM for L-leucine; about 0.0125 mM and about 0.125 mM for L-lysine.HCl; about 0.0025 mM and about 0.025 mM for L-methionine; about 0.005 mM and about 0.05 mM for L-phenylalanine; about 0.01 mM and about 0.1 mM for L-threonine; about 0.00125 and about 0.0125 mM L-tryptophan; about 0.005 and about 0.05 mM L-tyrosine.2Na2H₂O; about 0.01 mM and about 0.1 mM for L-valine; about 1.0 mM and about 10 mM for L-alanine; about 1.0 mM and about 30 mM for L-taurine; about 0.01 mM and about 1.0 mM for glutamic acid; about 1.0 mM and about 50 mM for L-glutamine or about 0.1 mM and about 1.0 mM for L-glycine.

30 10. The substantially protein-free medium according to claim 9, wherein the amino

acids are present at a concentration of: 0.5 mM L-arginine; 0.01 mM L-cystine.2HCl; 0.02 mM L-histidine, 0.04 mM L-isoleucine; 0.04 mM L-leucine; 0.05 mM L-lysine.HCl; 0.01 mM L-methionine; 0.02 mM L-phenylalanine; 0.04 mM L-threonine; 0.005 mM L-tryptophan; 0.02 mM L-tyrosine.2Na2H₂O; 0.04 mM L-valine; 0.5 mM L-alanine; 20 mM for L-taurine; 0.5 mM glutamic acid; and 20 mM L-glutamine or 0.25 mM L-Glycine.

11. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein the mineral salts comprising the medium are calcium chloride, magnesium

10 sulfate, potassium chloride, sodium chloride and sodium phosphate.

12. The substantially protein-free medium according to claim 11, wherein the mineral salts comprising the medium are present at concentrations between: about 3.0 mM (0.23 g/L) and 3.6 mM (0.27 g/L) for calcium chloride; about 0.7 mM (0.086 g/L) and 15 about 0.81 mM (0.098 g/L) for magnesium sulfate; about 4.7 mM (0.35 g/L) and about 5.4 mM (0.4 g/L) for potassium chloride; about 0.1 M (6.12 g/L) and about 0.12 M (6.95 g/L) for sodium chloride; and about 0.89 mM (0.107 g/L) to about 1.0 mM (0.122 g/L) for sodium dihydrogen phosphate.

20 13. The substantially protein-free medium according to claim 12, wherein the mineral salts comprising the medium are present at a concentration of: 3.1 mM (0.235 g/L) for calcium chloride; 0.72 mM (0.087 g/L) for magnesium sulfate; 4.8 mM (0.355 g/L) for potassium chloride; 0.11 M (6.171 g/L) for sodium chloride; and 0.88 mM (0.108 g/L) for sodium dihydrogen phosphate.

25

14. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein the vitamins comprising the medium are choline chloride, myoinositol, niacinamide, D-pantothenic acid, pyridoxine HCl, riboflavin, thiamine HCl, folic acid, vitamin B12, vitamin E, or any combination thereof.

30

15. The substantially protein-free medium according to claim 14, wherein the vitamins

comprising the medium are present at concentrations between: about 0.004 mM (0.0005 g/L) and 0.005 mM (0.0007 g/L) for choline chloride; about 0.0024 mM

(0.0006 g/L) and 0.0028 (0.0007 g/L) D-biotin ;about 0.0067 mM (0.0012 g/L) and 0.0078 mM (0.0014 g/L) for myoinositol; about 0.005 mM (0.0006 g/L) and 0.0057 mM for niacinamide (0.0007 g/L); about 0.0025 mM (0.0005 g/L) and 0.003 mM (0.0007 g/L) for D-pantothenic acid; about 0.003 (0.0006 g/L) mM and 0.0034 mM (0.0007 g/L) for pyridoxine HCl; about 0.00016 mM (0.00006 g/L) and 0.00019 mM (0.00007 g/L) for riboflavin; about 0.0018 mM (0.0005 g/L) and 0.0021 mM (0.0006 g/L) for thiamine HCl; about 0.0014 mM (0.006 g/L) and 0.0016 mM (0.0007 g/L) for folic acid; about 443 pM (600 ng/L) and 885 pM (1.2 mg/L) for vitamin B12; and 0.008 mM (0.003 g/L) and 0.012 mM (0.005 g/L) for vitamin E.

10

16. The substantially protein-free medium according to claim 15, wherein the vitamins

comprising the medium are present at a concentration of 0.004 mM (0.0005 g/L) for choline chloride; 0.0024 mM (0.0006 g/L) D-biotin ; 0.0067 mM (0.0012 g/L) for myoinositol; 0.005 mM (0.0006 g/L) for niacinamide; 0.0025 mM (0.0005 g/L) for D-pantothenic acid; 0.003 (0.0006 g/L) for pyridoxine HCl; 0.00016 mM (0.00006 g/L) for riboflavin; 0.0018 mM (0.0005 g/L) for thiamine HCl; 0.0014 mM (0.006 g/L) for folic acid; 616 pM (800 ng/L) for vitamin B12; and 0.010 mM (0.004 g/L) vitamin E.

15

17. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein the nutrients comprising the medium are D-glucose, pyruvate, fructose, lactic acid, or any combination thereof.

25

18. The substantially protein-free medium according to claim 17, wherein the nutrients

comprising the solution are present at a concentration of 4.3 mM (0.78 g/L) for D-glucose; 0.3 mM (0.02975 g/L) for sodium pyruvate; 5.1 mM (0.92 g/L) for fructose; and 10mM (1.13 g/L) for sodium lactate.

30

19. The substantially protein-free medium according to claim 18, wherein D-glucose is present at a concentration between 3.0 mM (0.75 g/L) and 5.6 mM (1.0 g/L).

20. The substantially protein-free medium according to claim 19, wherein fructose is present at a concentration between 1 mM (0.18 g/L) and 5.6 mM (1.01 g/L).

5 21. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein the antioxidant in the solution is glutathione.

22. The substantially protein-free medium according to claim 21, wherein the concentration of glutathione is between about 0.25 mM (0.077 g/L) and 0.35 mM (0.11 g/L).

10

23. The substantially protein-free medium according to claim 22, wherein the concentration of glutathione is 0.3 mM (0.092 g/L).

15

24. A substantially protein-free medium according to claims 1, 2, 3, or 4, further comprising EDTA.

25. The substantially protein-free medium according to claim 24, wherein the concentration of EDTA is between about 0.1 mM (0.0416 g/L) and 0.103 mM (0.043 g/L).

20

26. The substantially protein-free medium according to claim 25, wherein the concentration of EDTA is 0.1 mM (0.0418 g/L).

25

27. A substantially protein-free medium according to claim 1,2 ,3, or 4, further comprising HEPES.

30

28. The substantially protein-free medium according to claim 27, wherein the concentration of HEPES is 15 mM (3.6 g/L) or 25mM (5.96 g/L), and the concentration of D-Glucose is between about 3.3 mM (0.59 g/L) to about 3.6 mM (0.65 g/L).

29. A substantially protein-free medium according to claim 1,2 ,3, or 4, further comprising phenol red or other pH indicator.

30. The substantially protein-free medium according to claim 29, wherein the concentration of phenol red is 0.031 mM (0.011 g/L).

31. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein 5 gentamycin sulfate is present in the solution at a concentration from about 1.5 mg/L to about 4 mg/L.

32. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein Penicillin G is present at a concentration of 75 mg/L.

10

33. A substantially protein-free medium according to claims 1, 2, 3, or 4, wherein D-mannitol is present at a concentration from about 0.056 micromolar to about 6.9 micromolar.

15

34. The substantially protein-free medium of claim 33, wherein D-mannitol is present at a concentration of 2.8 micromolar.

35. A substantially protein-free medium of claims 1, 2, 3, or 4, further comprising sodium bicarbonate.

20

36. The substantially protein-free medium of claim 35, wherein sodium bicarbonate is present at a concentration of 26.2 mM (2.2 g/L).

25

37. The substantially protein-free medium of claim 36, wherein sodium bicarbonate is present at a concentration of between 4.0 mM (0.336 g/L) and 26.2 mM (2.2 g/L).

38. Use of a medium for assisted reproductive technology comprising the step of handling human reproductive cells with the substantially protein-free culture media of claim 36.

30

39. Use of a medium for assisted reproductive technology according to claim 38, wherein the substantially protein-free culture media is the media of claim 36.

40. Use of a medium for assisted reproductive technology according to claim

38, wherein the human reproductive cell is an unfertilized ovum.

41. Use of a medium for assisted reproductive technology according to claim 38, wherein the human reproductive cell is a plurality of spermatozoa.

5

42. Use of a medium for assisted reproductive technology according to claim 38, wherein the human reproductive cell is a fertilized ovum.

43. Use of a medium for assisted reproductive technology according to claim 38,

10 wherein the medium is specific for inter-uterine insemination.

44. A substantially protein-free cell culture medium for human reproductive cells according to claim 1, prepared from at least one or a plurality of stock solutions of mineral salts, amino acids, antioxidants, vitamins, nutrients, antibiotics, D-mannitol, and methylcellulose having a molecular weight of 14,000 Daltons, wherein said stock solutions are diluted with water to form the substantially protein-free cell culture medium.

45. A substantially protein-free cell culture medium for human reproductive cells,

20 suitable for use throughout *in vitro* treatment of human reproductive cells to be used in *in vitro* fertilization and other assisted reproduction technologies, comprising 0.5 mM L-arginine; 0.01 mM L-cystine.2HCl; 0.02 mM L-histidine, 0.04 mM L-isoleucine; 0.04 mM L-leucine; 0.05 mM L-lysine.HCl; 0.01 mM L-methionine; 0.02 mM L-phenylalanine; 0.04 mM L-threonine; 0.005 mM L-tryptophan; 0.02 mM L-tyrosine.2Na2H₂O; 0.04 mM L-valine; 0.5 mM L-alanine; 20 mM L-taurine; 0.5 mM glutamic acid (0.39 mM); 20 mM L-glutamine or 0.25 mM L-Glycine; 3.1 mM (0.235 g/L) calcium chloride; 0.72 mM (0.087 g/L) magnesium sulfate; 4.8 mM (0.355 g/L) potassium chloride; 0.11 M (6.171 g/L) sodium chloride; 0.88 mM (0.108 g/L) for sodium dihydrogen phosphate; 0.004 mM (0.0005 g/L) choline chloride;

25 0.0024 mM (0.0006 g/L) D-biotin; 0.0067 mM (0.0012 g/L) myoinositol; 0.005 mM (0.0006 g/L) niacinamide; 0.0025 mM (0.0005 g/L) D-pantothenic acid; 0.003 (0.0006 g/L) pyridoxine HCl; 0.00016 mM (0.00006 g/L) riboflavin; 0.0018 mM (0.0005 g/L) thiamine HCl; 0.0014 mM (0.006 g/L) folic acid; 616 pM (800 ng/L) vitamin B12; 0.010 mM (0.004 g/L) vitamin E; 4.3 mM (0.78 g/L) D-glucose; 0.3

mM (0.02975 g/L) sodium pyruvate; 5.1 mM (0.92 g/L) fructose; 10mM (1.13 g/L) sodium lactate; 0.3 mM (0.092 g/L) glutathione; 0.1 mM (0.0418 g/L) EDTA; 0.031 mM (0.011 g/L) phenol red; 2.8 micromolar D-mannitol; 26.2 mM (2.2 g/L) sodium bicarbonate; 75mg/L Penicillin G; and 1.5 mg/L gentamycin sulfate.

5

46. The substantially protein-free cell culture medium of claim 45, wherein the concentration of gentamycin sulfate is 4 mg/L.
47. The substantially protein-free cell culture medium of claim 45, wherein the concentration of D-glucose is between 3.3 mM (0.59 g/L) to about 3.6 mM (0.65 g/L) and further comprising 15 mM (3.6 g/L) HEPES.
48. The substantially protein-free cell culture medium of claim 45, wherein the concentration of D-glucose is between 3.3 mM (0.59 g/L) to about 3.6 mM (0.65 g/L) and further comprising 25 mM (5.96 g/L) HEPES.
49. The substantially protein-free cell culture medium of claim 46, wherein the concentration of D-glucose is between 3.3 mM (0.59 g/L) to about 3.6 mM (0.65 g/L) and further comprising 15 mM (3.6 g/L) HEPES.
50. The substantially protein-free cell culture medium of claim 46, wherein the concentration of D-glucose is between 3.3 mM (0.59 g/L) to about 3.6 mM (0.65 g/L) and further comprising 25mM (5.96 g/L) HEPES.

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2008/087818A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N5/02 C12N5/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BIGGERS J D ET AL: "Polyvinyl alcohol and amino acids as substitutes for bovine serum albumin in culture media for mouse preimplantation embryos."</p> <p>HUMAN REPRODUCTION UPDATE, vol. 3, no. 2, March 1997 (1997-03), pages 125-135, XP002520128</p> <p>ISSN: 1355-4786</p> <p>cited in the application</p> <p>-----</p> <p style="text-align: center;">-/-</p>	

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

19 March 2009

30/03/2009

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Teyssier, Bertrand

INTERNATIONAL SEARCH REPORT

International application No PCT/US2008/087818

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SCHRAMM R D ET AL: "Development of in-vitro-fertilized primate embryos into blastocysts in a chemically defined, protein-free culture medium." <i>HUMAN REPRODUCTION</i>, vol. 11, no. 8, August 1996 (1996-08), pages 1690-1697, XP002520129 ISSN: 0268-1161 cited in the application</p> <p>-----</p>	
A	<p>GARDNER D K & LANE M: "Culture of viable human blastocysts in defined sequential serum-free media." <i>HUMAN REPRODUCTION</i>, vol. 13 Suppl 3, June 1998 (1998-06), pages 148-159, XP001026692 ISSN: 0268-1161 cited in the application</p> <p>-----</p>	
A	<p>STEWART G J ET AL: "Methylcellulose protects the ability of anchorage-dependent cells to adhere following isolation and holding in suspension." <i>BIOTECHNIQUES</i>, vol. 19, no. 4, October 1995 (1995-10), pages 598-604, XP002520130 ISSN: 0736-6205 cited in the application</p> <p>-----</p>	
A	<p>WO 2005/108555 A (E.N.V. NANTES [FR]; TAINTURIER [FR]; MARTAL J [FR]; NEIRA J A [CO]) 17 November 2005 (2005-11-17)</p> <p>-----</p>	
A	<p>STAESSEN C ET AL: "Controlled comparison of commercial media for human in-vitro fertilization: Ménézo B2 medium versus Medi-Cult universal and BM1 medium." <i>HUMAN REPRODUCTION</i>, vol. 13, no. 9, September 1998 (1998-09), pages 2548-2554, XP002520131 ISSN: 0268-1161 cited in the application</p> <p>-----</p>	
A	<p>ALI J: "Investigation into the nutrient requirement of the human embryo: Successful formulation and clinical trial of a novel protein-free embryo culture medium" <i>EMIRATES MEDICAL JOURNAL</i>, vol. 18, no. 3, 2000, pages 195-202, XP008103636 cited in the application</p> <p>-----</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/US2008/087818

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 2005108555	A 17-11-2005	EP FR	1756266 A2 2869043 A1	28-02-2007 21-10-2005