Office de la Propriété Intellectuelle du Canada
Un organisme d'Industrie Canada

Canadian Intellectual Property Office
An agency of Industry Canada

CA 2454613 A1 2003/01/16

(21) 2 454 613

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2002/07/03
(87) Date publication PCT/PCT Publication Date: 2003/01/16
(85) Entrée phase nationale/National Entry: 2003/12/22
(86) N° demande PCT/PCT Application No.: US 2002/021063
(87) N° publication PCT/PCT Publication No.: 2003/004027
(30) Priorités/Priorities: 2001/07/05 (09/899,794) US;
2002/01/09 (10/042,582) US

(51) Cl.Int.^{7}/Int.Cl.^{7} A61K 31/445, C07D 211/94,
C07D 211/30, C07D 403/12, C07D 401/12,
C07D 403/06, C07D 401/06

(71) Demandeur/Applicant:
SYNAPTIC PHARMACEUTICAL CORPORATION, US

(72) Inventeurs/Inventors:
MARZABADI, MOHAMMAD R., US;
WETZEL, JOHN, US;
DELEON, JOHN E., US;
JIANG, YU, US;
CHEN, CHIEN-AN, US;
LU, KAI, US;
LAGU, BHARAT, US

(74) Agent: GOUDREAU GAGE DUBUC

(54) Titre : PIPERIDINES ANILINIQUES SUBSTITUEES, UTILISEES COMME ANTAGONISTES SELECTIFS DE LA MCH

(54) Title: SUBSTITUTED ANILINIC PIPERIDINES AS MCH SELECTIVE ANTAGONISTS

(57) Abrégé/Abstract:
This invention is directed to compounds which are selective antagonists for melanin concentrating hormone-1 (MCH1) receptors. The invention provides a pharmaceutical composition comprising a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier. This invention provides a pharmaceutical composition made by combining a therapeutically effective amount of the compound of this invention and a pharmaceutically acceptable carrier. This invention further provides a process for making a pharmaceutical composition comprising combining a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004027 A1

(51) International Patent Classification[7]: A61K 31/445, C07D 211/30, 211/94, 401/06, 401/12, 403/06, 403/12

(21) International Application Number: PCT/US02/21063

(22) International Filing Date: 3 July 2002 (03.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

09/899,794	5 July 2001 (05.07.2001)	US
10/042,582	9 January 2002 (09.01.2002)	US

(71) Applicant: **SYNAPTIC PHARMACEUTICAL CORPORATION** [US/US]; 215 College Road, Paramus, NJ 07652 (US).

(72) Inventors: **MARZABADI, Mohammad R.**; 153 Woodland Avenue, Ridgewood, NJ 07450 (US). **WETZEL, John**; 23-17 Morlot Avenue, Fairlawn, NJ 07410 (US). **DELEON, John, E.**; 7427 Boulevard East, Apt. 5, North Bergen, NJ 07047 (US). **JIANG, Yu**; 179 Manhattan Avenue, Apt. 7B, Jersey, City, NJ 07307 (US).

(74) Agent: **WHITE, John, P.**; Cooper & Dunahm LLP, 1185 Avenue of the Americas, New York, NY 10036 (US).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/004027 A1

(54) Title: SUBSTITUTED ANILINIC PIPERIDINES AS MCH SELECTIVE ANTAGONISTS

(57) Abstract: This invention is directed to compounds which are selective antagonists for melanin concentrating hormone-1 (MCH1) receptors. The invention provides a pharmaceutical composition comprising a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier. This invention provides a pharmaceutical composition made by combining a therapeutically effective amount of the compound of this invention and a pharmaceutically acceptable carrier. This invention further provides a process for making a pharmaceutical composition comprising combining a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 474

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 474

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

SUBSTITUTED ANILINIC PIPERIDINES AS MCH SELECTIVE ANTAGONISTS

BACKGROUND OF THE INVENTION

This application is a continuation-in-part of U.S. Serial No. 10/042,582, filed January 9, 2002, and of U.S. Serial No. 09/899,794, filed July 5, 2001, the contents of both of which are hereby incorporated by reference into the subject application.

Throughout this application, various publications are referenced in parentheses by author and year. Full citations for these references may be found at the end of the specification immediately preceding the sequence listings and the claims. The disclosure of these publications in their entireties are hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains. Melanin-concentrating hormone (MCH) is a cyclic peptide originally isolated from salmonid (teleost fish) pituitaries (Kawauchi et al., 1983). In fish the 17 amino acid peptide causes aggregation of melanin within the melanophores and inhibits the release of ACTH, acting as a functional antagonist of α-MSH. Mammalian MCH (19 amino acids) is highly conserved between rat, mouse, and human, exhibiting 100% amino acid identity, but its physiological roles are less clear. MCH has been reported to participate in a variety of processes including feeding, water balance, energy metabolism, general arousal/attention state,

memory and cognitive functions, and psychiatric disorders (for reviews, see Baker, 1991; Baker, 1994; Nahon, 1994; Knigge et al., 1996). Its role in feeding or body weight regulation is supported by a recent *Nature* publication (Qu et al., 1996) demonstrating that MCH is overexpressed in the hypothalamus of *ob/ob* mice compared with *ob/+* mice, and that fasting further increased MCH mRNA in both obese and normal mice during fasting. MCH also stimulated feeding in normal rats when injected into the lateral ventricles (Rossi et al., 1997). MCH also has been reported to functionally antagonize the behavioral effects of α-MSH (Miller et al., 1993; Gonzalez et al, 1996; Sanchez et al., 1997); in addition, stress has been shown to increase POMC mRNA levels while decreasing the MCH precursor preproMCH (ppMCH) mRNA levels (Presse et al., 1992). Thus MCH may serve as an integrative neuropeptide involved in the reaction to stress, as well as in the regulation of feeding and sexual activity (Baker, 1991; Knigge et al., 1996).

Although the biological effects of MCH are believed to be mediated by specific receptors, binding sites for MCH have not been well described. A tritiated ligand ($[^3H]$-MCH) was reported to exhibit specific binding to brain membranes but was unusable for saturation analyses, so neither affinity nor B_{max} were determined (Drozdz and Eberle, 1995). Radioiodination of the tyrosine at position thirteen resulted in a ligand with dramatically reduced biological activity (see Drozdz and Eberle, 1995). In contrast, the radioiodination of the MCH analogue $[Phe^{13},Tyr^{19}]$-MCH was successful (Drozdz et al., 1995); the ligand retained biological activity and

exhibited specific binding to a variety of cell lines including mouse melanoma (B16-F1, G4F, and G4F-7), PC12, and COS cells. In G4F-7 cells, the K_D = 0.118nM and the B_{max} ~1100 sites/cell. Importantly, the binding was not inhibited by α-MSH but was weakly inhibited by rat ANF (Ki = 116 nM vs. 12 nM for native MCH) (Drozdz et al., 1995). More recently specific MCH binding was reported in transformed keratinocytes (Burgaud et al., 1997) and melanoma cells (Drozdz et al., 1998), where photo-crosslinking studies suggest that the receptor is a membrane protein with an apparent molecular weight of 45-50 kDaltons, compatible with the molecular weight range of the GPCR superfamily of receptors. No radioautoradiographic studies of MCH receptor localization using this ligand have been reported as yet.

The localization and biological activities of MCH peptide suggest that the modulation of MCH receptor activity may be useful in a number of therapeutic applications. The role of MCH in feeding is the best characterized of its potential clinical uses. MCH is expressed in the lateral hypothalamus, a brain area implicated in the regulation of thirst and hunger (Grillon et al., 1997); recently orexins A and B, which are potent orexigenic agents, have been shown to have very similar localization to MCH in the lateral hypothalamus (Sakurai et al., 1998). MCH mRNA levels in this brain region are increased in rats after 24 hours of food-deprivation (Hervé and Fellman, 1997); after insulin injection, a significant increase in the abundance and staining intensity of MCH immunoreactive perikarya and fibres was observed concurrent with a

significant increase in the level of MCH mRNA (Bahjaoui-Bouhaddi et al., 1994). Consistent with the ability of MCH to stimulate feeding in rats (Rossi et al., 1997) is the observation that MCH mRNA levels are upregulated in the hypothalami of obese *ob/ob* mice (Qu et al., 1996), and decreased in the hypothalami of rats treated with leptin, whose food intake and body weight gains are also decreased (Sahu, 1998). MCH appears to act as a functional antagonist of the melanocortin system in its effects on food intake and on hormone secretion within the HPA (hypothalamopituitary/adrenal axis) (Ludwig et al., 1998). Together these data suggest a role for endogenous MCH in the regulation of energy balance and response to stress, and provide a rationale for the development of specific compounds acting at MCH receptors for use in the treatment of obesity and stress-related disorders.

In all species studied to date, a major portion of the neurons of the MCH cell group occupies a rather constant location in those areas of the lateral hypothalamus and subthalamus where they lie and may be a part of some of the so-called "extrapyramidal" motor circuits. These involve substantial striato- and pallidofugal pathways involving the thalamus and cerebral cortex, hypothalamic areas, and reciprocal connections to subthalamic nucleus, substantia nigra, and mid-brain centers (Bittencourt et al., 1992). In their location, the MCH cell group may offer a bridge or mechanism for expressing hypothalamic visceral activity with appropriate and coordinated motor activity. Clinically it may be of some value to consider the involvement of this MCH system in movement disorders, such as

Parkinson's disease and Huntingdon's Chorea in which extrapyramidal circuits are known to be involved.

Human genetic linkage studies have located authentic hMCH loci on chromosome 12 (12q23-24) and the variant hMCH loci on chromosome 5 (5q12-13) (Pedeutour et al., 1994). Locus 12q23-24 coincides with a locus to which autosomal dominant cerebellar ataxia type II (SCA2) has been mapped (Auburger et al., 1992; Twells et al., 1992). This disease comprises neurodegenerative disorders, including an olivopontocerebellar atrophy. Furthermore, the gene for Darier's disease, has been mapped to locus 12q23-24 (Craddock et al., 1993). Dariers' disease is characterized by abnormalities I keratinocyte adhesion and mental illnesses in some families. In view of the functional and neuroanatomical patterns of the MCH neural system in the rat and human brains, the MCH gene may represent a good candidate for SCA2 or Darier's disease. Interestingly, diseases with high social impact have been mapped to this locus. Indeed, the gene responsible for chronic or acute forms of spinal muscular atrophies has been assigned to chromosome 5q12-13 using genetic linkage analysis (Melki et al., 1990; Westbrook et al., 1992). Furthermore, independent lines of evidence support the assignment of a major schizophrenia locus to chromosome 5q11.2-13.3 (Sherrington et al., 1988; Bassett et al., 1988; Gilliam et al., 1989). The above studies suggest that MCH may play a role in neurodegenerative diseases and disorders of emotion.

Additional therapeutic applications for MCH-related compounds are suggested by the observed effects of MCH

in other biological systems. For example, MCH may regulate reproductive functions in male and female rats. MCH transcripts and MCH peptide were found within germ cells in testes of adult rats, suggesting that MCH may participate in stem cell renewal and/or differentiation of early spermatocytes (Hervieu et al., 1996). MCH injected directly into the medial preoptic area (MPOA) or ventromedial nucleus (VMN) stimulated sexual activity in female rats (Gonzalez et al., 1996). In ovariectomized rats primed with estradiol, MCH stimulated luteinizing hormone (LH) release while anti-MCH antiserum inhibited LH release (Gonzalez et al., 1997). The zona incerta, which contains a large population of MCH cell bodies, has previously been identified as a regulatory site for the pre-ovulatory LH surge (MacKenzie et al., 1984). MCH has been reported to influence release of pituitary hormones including ACTH and oxytocin. MCH analogues may also be useful in treating epilepsy. In the PTZ seizure model, injection of MCH prior to seizure induction prevented seizure activity in both rats and guinea pigs, suggesting that MCH-containing neurons may participate in the neural circuitry underlying PTZ-induced seizure (Knigge and Wagner, 1997). MCH has also been observed to affect behavioral correlates of cognitive functions. MCH treatment hastened extinction of the passive avoidance response in rats (McBride et al., 1994), raising the possibility that MCH receptor antagonists may be beneficial for memory storage and/or retention. A possible role for MCH in the modulation or perception of pain is supported by the dense innervation of the periaqueductal grey (PAG) by MCH-positive fibers. Finally, MCH may participate in the regulation of fluid

intake. ICV infusion of MCH in conscious sheep produced diuretic, natriuretic, and kaliuretic changes in response to increased plasma volume (Parkes, 1996). Together with anatomical data reporting the presence of MCH in fluid regulatory areas of the brain, the results indicate that MCH may be an important peptide involved in the central control of fluid homeostasis in mammals.

The identification of a G-protein coupled receptor for MCH has recently been published (Chambers et al., 1999; Saito et al., 1999). These groups identified MCH as the endogenous ligand for the human orphan G-protein coupled receptor SLC-1 (Lakaye et al., 1998). The rat homologue of this receptor (now called MCH-1) was reported to be localized in regions of the rat brain associated with feeding behavior (e.g. dorsomedial and ventromedial hypothalamus). The link between MCH-1 and the effects of MCH on feeding has been strengthened by recent reports on the phenotype of MCH-1 knockout mice. Two groups have shown independently (Marsh et al, 2002; Chen et al, 2002) that the targeted disruption of the MCH-1 receptor gene (MCH-1 knockout) in mice results in animals that are hyperphagic but are lean and have decreased body mass relative to wild-type littermates. The decrease in body mass is attributed to an increase in metabolism. Each group demonstrated that the MCH-1 knockout mice are resistant to diet-induced obesity, and generally exhibit weights similar to littermates maintained on regular chow.

Finally, synthetic antagonist molecules for the MCH-1 receptor have now been described in the literature. Bednarek et al. (2002) have reported on the synthesis of

high affinity peptide antagonists of MCH-1. In addition, a small molecule antagonist of MCH-1 has been described by Takekawa et al. (Takekawa et al., 2002). This compound, T-226296, exhibits high affinity for the MCH-1 receptor (~ 5-9 nM for rat and human MCH-1), and was shown to inhibit food intake induced by the intracerebroventricular application of MCH. These data validate the strategy of using an MCH-1 receptor antagonist to treat obesity.

Furthermore, in our own studies, we have tested MCH1 antagonists in several animal models that are well known as predictive for the efficacy of compounds in humans (Borowsky, et al., in press; unpublished data). These experiments indicate that MCH1 antagonists are useful to treat obesity, depression, anxiety, as well as urinary disorders.

As used in this invention, the term "antagonist" refers to a compound which binds to, and decreases the activity of, a receptor in the presence of an agonist. In the case of a G-protein coupled receptor, activation may be measured using any appropriate second messenger system which is coupled to the receptor in a cell or tissue in which the receptor is expressed. Some specific, but by no means limiting, examples of well-known second messenger systems are adenylate cyclase, intracellular calcium mobilization, ion channel activation, guanylate cyclase and inositol phospholipid hydrolysis. Conversely, the term "agonist" refers to a compound which binds to, and increases activity of, a receptor as compared with the activity of the receptor in the absence of any agonist.

In one embodiment of this invention, the synthesis of novel compounds which bind selectively to the cloned human melanin-concentrating hormone-1 (MCH1) receptor, compared to other cloned G-protein coupled receptors, and inhibit the activation of the cloned receptors as measured in *in vitro* assays is disclosed. The *in vitro* receptor binding assays described hereinafter were performed using various cultured cell lines, each transfected with and expressing only a single cloned receptor.

Furthermore, the compounds of the present invention may also be used to treat abnormal conditions such as feeding disorders (obesity, bulimia and bulimia nervosa), sexual/reproductive disorders, depression, anxiety, depression and anxiety, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleep disturbances, or any condition in which antagonism of an MCH1 receptor may be beneficial. In addition, the compounds of the present invention may be used to reduce the body mass of a subject. Furthermore, the compounds of the present invention may be used to treat urinary disorders.

Summary of the Invention

This invention provides a compound having the structure:



;

wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CH_3$, $-CF_3$, $-COR_2$, $-CO_2R_2$, phenyl, phenoxy or straight chained or branched C_1-C_7 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl;

wherein A is –H, –F, -Cl, -Br, -CN, $-NO_2$, $-COR_3$, $-CO_2R_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is O or NH; and

wherein n is an integer from 0 to 5 inclusive.

In one embodiment, R_1 is aryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-COR_2$, $-CO_2R_2$, straight chained or branched C_1-C_7 alkyl;

wherein R_3 is phenyl;

wherein A is H; and

wherein X is O.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



In one embodiment, compound has the structure:



In one embodiment, R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl; and wherein R_3 is aryl.

In one embodiment, R_2 is isopropyl; and A is hydrogen.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



The present invention also provides a compound having the structure:


;

wherein R_1 is aryl or heteroaryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-OCH_3$, phenoxy, fused cyclopentanyl, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein R_2 is straight-chained or branched C_1-C_4 alkyl or cyclopropyl;

wherein A is -H, -F, -Cl, - Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; and

wherein n is an integer from 1 to 5 inclusive.

In one embodiment, R_1 is aryl optionally substituted with one or more -F, -Cl, -Br, -I or straight chained or branched C_1-C_4 alkyl; and

wherein A is H.

In one embodiment, R_2 is isopropyl; and

wherein n is 2.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, R_1 is thienyl; and wherein A is H.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



The invention provides a compound having the structure:



;

wherein W is



or



;

wherein each R_1 is independently hydrogen, methyl or ethyl;

wherein R_2 is straight- chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is hydrogen, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -H, -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl.

wherein each A is independently -H, -F, -Cl, -Br, -CN, -NO_2, -COR_3, -CO_2R_3, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is O, NR_3, CO or may be absent; and

wherein Y is hydrogen, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl.

In one embodiment, W is



;

and wherein X is O or may be absent.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, W is



In one embodiment, A is -H, -F, -Cl, -Br.

In one embodiment, R_2 is isopropyl; and A is hydrogen.

In one embodiment, the compound has the structure:



This invention provides a compound having the structure:



;

wherein W is



or



wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-OCH_3$, $-CO_2CH_3$, $-CF_3$, phenyl, straight chained or branched C_1-C_7 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl or phenyl.

wherein each B is independently -H, -F, -Cl, -Br, -I,

-CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, - OCH_3, $-OCF_3$, $-CF_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl or aryl, phenoxy or benzyloxy, wherein the aryl, phenoxy or benzyloxy is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, $-OCH_3$, $-OCF_3$, $-CF_3$ or straight chained or branched C1 -C3 alkyl.

In one embodiment, W is



.

In one embodiment, R_1 is hydrogen or phenyl optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



.

In one embodiment, the compound has the structure:



This invention provides a compound having the structure:



;

wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-CF_3$, $-OCH_3$, straight chained or branched C_1-C_3 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is -H, -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CF_3$, $-OCH_3$, or straight chained or branched C_1-C_3 alkyl, monofluoroalkyl or polyfluoroalkyl, or a phenyl ring fused to C_6 and C_7 of the indole moiety;

wherein R_4 is hydrogen or aryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CF_3$, straight chained or branched C_1-C_3 alkyl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; and

wherein n is an integer from 2 to 4 inclusive.

In one embodiment, R_3 is -H, -F, -Cl, -Br, -I, -CN, $-NO_2$, $-OCF_3$ or $-OCH_3$; and

wherein R_4 is hydrogen or phenyl optionally substituted with one or more -F, -Cl or $-CF_3$.

In one embodiment, R_1 is hydrogen or phenyl optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-CF_3$, $-OCH_3$ or straight chained or branched C_1-C_3 alkyl;

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



25

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



This invention provides a compound having the structure:



;

wherein each R_1 is independently hydrogen or CH_3;

wherein R_2 is straight-chained or branched C_1-C_4 alkyl or cyclopropyl;

wherein R_3 is benzyl or phenyl, wherein the benzyl or phenyl is optionally substituted with a methylenenedioxy group or one or more -F or -Cl;

wherein A is –H, –F, -Cl, - Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is $(CH_2)_2$, $COCH_2$ or CONH;

In one embodiment, R_3 is phenyl optionally substituted with one or more –F; and

wherein A is hydrogen.

In one embodiment, X is CONH.

In one embodiment, R_2 is methyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



;

wherein each Y is independently hydrogen or –F.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, R_3 is benzyl optionally substituted with a methylenedioxy group or one or more -F or -Cl.

In one embodiment, the compound has the structure:



;

wherein each Y is independently hydrogen or -F.

In one embodiment, the compound has the structure:



In one embodiment, the compound is enantiomerically pure.

In one embodiment, the compound is diastereomerically pure.

In one embodiment, the compound is enantiomerically and diastereomerically pure.

This invention also provides a pharmaceutical composition comprising a therapeutically amount of a compound of the invention and a pharmaceutically acceptable carrier.

In one embodiment, the amount of the compound is from about 0.01mg to about 500mg.

In one embodiment, the amount of the compound is from about 0.1mg to about 60mg.

In one embodiment, the amount of the compound is from about 1mg to about 20mg.

In one embodiment, the pharmaceutical composition consists of a carrier which is a liquid and the composition is a solution.

In one embodiment, the pharmaceutical composition consists of a carrier which is a solid and the composition is a tablet.

In one embodiment, the pharmaceutical composition consists of a carrier which is a gel and the composition is a suppository.

The invention also provides a process for making a pharmaceutical composition comprising admixing a therapeutically effective amount of the compound of any of the invention and a pharmaceutically acceptable carrier.

This invention also provides the method of treating a subject suffering from a disorder selected from the group consisting of depression, anxiety, urge incontinence, or obesity comprising administering to the subject a therapeutically effective amount of the compound of the invention.

In one embodiment, the therapeutically effective amount is between about 0.03 and about 1000 mg per day.

In one embodiment, the therapeutically effective amount is between about 0.30 and about 300 mg per day.

In one embodiment, the therapeutically effective amount is between about 1.0 and about 100 mg per day.

In one embodiment, the disorder is depression.

In one embodiment, the disorder is anxiety.

In one embodiment, the disorder is obesity.

In one embodiment, the disorder is urge incontinence.

The invention provides the method of reducing the body mass of a subject, which comprises administering to the subject an amount of a compound of the invention effective to reduce the body mass of the subject.

The invention provides the method of treating a subject suffering from depression, which comprises administering to the subject an amount of a compound of any of claims of the invention effective to treat the subject's depression.

The invention provides the method of treating a subject suffering from anxiety, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's anxiety.

The invention provides the method of alleviating urge urinary incontinence in a subject suffering from an overactive bladder, which comprises administering to the subject an amount of the compound of the invention effective to alleviate the subject's urge urinary incontinence.

The invention provides the method of managing obesity in a subject in need of weight loss, which comprises administering to the subject an amount of a compound of the invention effective to induce weight loss in the subject.

The invention provides the method of managing obesity in a subject who has experienced weight loss, which comprises administering to the subject an amount of a compound of the invention effective to maintain such weight loss in the subject.

The invention provides the method of treating overactive bladder in a subject, which comprises administering to the subject an amount of a compound of any of the invention effective to treat the subject's overactive bladder.

The invention provides the method of treating a disorder in a subject, wherein the symptoms of the subject can be alleviated by treatment with an MCH1 antagonist, wherein the MCH1 antagonist is the compound of the invention.

The invention provides the method of alleviating the symptoms of a disor4der in a subject, which comprises administering to the subject an amount of an MCH1 antagonist effective to alleviate the symptoms, wherein the MCH1 antagonist is the compound of the invention.

Detailed Description of the Invention

This invention provides a compound having the structure:



,



,



; or



;

wherein each V is independently phthalimide, aryl, phenoxy or heteroaryl, wherein the aryl, phenoxy or heteroaryl is optionally substituted with one or more F; Cl; Br; I; COR_5; CO_2R_5; $-OCOR_5$; $-CON(R_5)_2$; $-N(R_5)COR_5$; CN; $-NO_2$; $-N(R_5)_2$; $-OR_5$; $-SR_5$; $(CH_2)_qOR_5$; $(CH_2)_qR_5$; $(CH_2)_qSR_5$; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl, polyfluoroalkyl, aminoalkyl, or carboxamidoalkyl; straight chained or branched C_2-C_7 alkenyl, C_2-C_7 alkynyl; aryl; phenoxy; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein each W is independently aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more F; Cl; Br; I; COR_3; $-OCOR_3$; CO_2R_3; $-CON(R_3)_2$; $-N(R_3)COR_3$; CN; $-NO_2$; $-N(R_3)_2$; $-OR_3$; $-SR_3$; $(CH_2)_qOR_3$; $(CH_2)_qSR_3$; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl, polyfluoroalkyl, aminoalkyl, or carboxamidoalkyl; straight chained or branched C_2-C_7 alkenyl, C_2-C_7 alkynyl; aryl; phenoxy; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein X is hydrogen or - OR_3, provided that when X is $-OR_3$ the V geminal to X cannot be phthalimide;

wherein Y is hydrogen, =O (carbonyl oxygen), OR_3, OV, COV, =NNHV, $=NNR_5$, NZR_5, NZV, NCONV (ureas), $NCONR_5$, NR_3, carbazole, indole or phthalimide;

wherein each R is independently -H; -F; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl or alkynyl; $-N(R_3)_2$; $-NO_2$; -CN; $-CO_2R_3$; $-OCOR_3$; $-OR_3$; or $-N(R_3)COR_3$; $-CON(R_3)_2$;

wherein each R_3 is independently -H; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl or alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein each R_5 is -H; $-NO_2$; $-N_3$; -CN; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl or alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl; $-N(R_3)_2$; $-OR_3$; $-(CH_2)_pOR_3$; $-COR_3$; $-CO_2R_3$; $-OCOR_3$; $-CON(R_3)_2$; $-N(R_3)COR_3$; aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more F; Cl; Br; I; COR_6; CO_2R_3; $-OCOR_3$; $-CON(R_3)_2$; $-N(R_3)COR_3$; CN; $-NO_2$; $-N(R_3)_2$; $-OR_6$; $-SR_6$; $(CH_2)_qOR6$; $(CH_2)_qSR_6$; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl, polyfluoroalkyl, aminoalkyl, or carboxamidoalkyl; straight chained or branched C_2-C_7 alkenyl, C_2-C_7 alkynyl; C_3-C_7 cycloalkyl,

monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein R_6 is. -H; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl or alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl; $-N(R_3)_2$; $-OR_3$; $-(CH_2)_pOR_3$; $-COR_3$; $-CO_2R_3$; $-OCOR_3$; $-CON(R_3)_2$; $-N(R_3)COR_3$; aryl, benzyl or heteroaryl, optionally substituted with one or more F; Cl; Br; I; COR_3; CO_2R_3; $-OCOR_3$; $-CON(R_3)_2$; $-N(R_3)COR_3$, CN; $-NO_2$; $-N(R_3)_2$; $-OR_3$; $-SR_3$; $(CH_2)_qOR_3$; $(CH_2)_qSR_3$; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl, polyfluoroalkyl, aminoalkyl, or carboxamidoalkyl; aryl; benzyl; straight chained or branched C_2-C_7 alkenyl, C_2-C_7 alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein Z is CO, SO_2 or SO_2NR_6;

wherein each m is independently an integer from 0 to 3 inclusive;

wherein each n is independently an integer from 0 to 5 inclusive;

wherein each p is independently an integer from 1 to 7 inclusive; and

wherein q is an integer from 1 to 3 inclusive;

or a pharmaceutically acceptable salt thereof.

As used in the present invention, the term "cycloalkyl" includes C_3-C_7 cycloalky moities which may be substituted with one or more of the following: F; CN; $-NO_2$; straight chained or branched C_1-C_7 alkyl, straight chained or branched C_1-C_7 monofluoroalkyl,straight chained or branched C_1-C_7 polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl, straight chained or branched C_2-C_7 alkynyl; C_3-C_7 cycloalkyl, C_3-C_7 monofluorocycloalkyl, C_3-C_7 polyfluorocycloalkyl, C_5-C_7 cycloalkenyl, $-N(R_3)_2$; $-OR_3$; $-NCOR_3$; $-COR_3$; $-CO_2R_3$; $-CON(R_3)_2$ or $(CH_2)_p$-O-$(CH_3)_m$-CH_3.

In the present invention, the term "cycloalkenyl" includes C_5-C_7 cycloalkenyl moities which may be substituted with one or more of the following: -F; -Cl; -Br, -I; CN; $-NO_2$; straight chained or branched C_1-C_7 alkyl, straight chained or branched C_1-C_7 monofluoroalkyl, straight chained or branched C_1-C_7 polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl, straight chained or branched C_2-C_7 alkynyl; C_3-C_7 cycloalkyl, C_3-C_7 monofluorocycloalkyl, C_3-C_7 polyfluorocycloalkyl, C_5-C_7 cycloalkenyl,$-N(R_3)_2$; $-OR_3$; $-NCOR_3$; $-COR_3$; $-CO_2R_3$; $-CON(R_3)_2$ or $(CH_2)_p$-O-$(CH_3)_m$-CH_3.

As used in the present invention, the term "heteroaryl" is used to include five and six membered unsaturated rings that may contain one or more oxygen, sulfur, or nitrogen atoms. Examples of heteroaryl groups include, but are not limited to, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazzolyl, thiadiazolyl,

pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl.

In addition, the term "heteroaryl" is used to include fused bicyclic ring systems that may contain one or more heteroatoms such as oxygen, sulfur and nitrogen. Examples of such heteroaryl groups include, but are not limited to, indolizinyl, indolyl, isoindolyl, benzo[b]furanyl, benzo[b]thiophenyl, indazolyl, benzimidazolyl, purinyl, benzoxazolyl, benzisoxazolyl, benzo[b]thiazolyl, imidazo[2,1-b]thiazolyl, cinnolinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, quinolinyl, isoquinolinyl, phthalimidyl and 2,1,3-benzothiazolyl.

The term "heteroaryl" also includes those chemical moieties recited above which may be substituted with one or more of the following: -F; -Cl; -Br, -I; CN; $-NO_2$; straight chained or branched C_1-C_7 alkyl, straight chained or branched C_1-C_7 monofluoroalkyl, straight chained or branched C_1-C_7 polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl, straight chained or branched C_2-C_7 alkynyl; C_3-C_7 cycloalkyl, C_3-C_7 monofluorocycloalkyl, C_3-C_7 polyfluorocycloalkyl, C_5-C_7 cycloalkenyl, $-N(R_3)_2$; $-OR_3$; $-NCOR_3$; $-COR_3$; $-CO_2R_3$; $-CON(R_3)_2$ or $(CH_2)_p$-O-$(CH_3)_m$-CH_3.

In still another embodiment of the above described invention, the compound has the structure:



;



;



; or



.

In a further embodiment of the instant invention, R_6 is straight chained or branched C_1-C_7 alkyl; C_3-C_7 cycloalkyl; $-N(R_3)_2$; $-OR_3$; $-(CH_2)_pOR_3$; aryl, benzyl or heteroaryl, optionally substituted with one or more F; Cl; Br; I; $-OR_3$; $-(CH_2)_qOR_3$; or straight chained or branched C_1-C_7 alkyl.

In an embodiment of the present invention, the compound has the structure:



; or

.

In a further embodiment of the present invention, at least one V is phenyl optionally substituted with one or more F; Cl; Br; $-OR_3$; $(CH_2)_qOR_3$; straight chained or branched C_1-C_7 alkyl; C_1-C_7 polyfluoroalkyl; or phenoxy.

In one embodiment of the present invention, the compound is:



.

In one embodiment, the compound is:



.

In one embodiment, the compound is:



In another embodiment of the present invention, the compound has the structure:



; or



In a further embodiment of the present invention, at least one V is phenyl optionally substituted with one or more F; Cl; Br; $-OR_3$; $(CH_2)_qOR_3$; straight chained or branched C_1-C_7 alkyl; C_1-C_7 polyfluoroalkyl; or phenoxy.

In another embodiment of the present invention, the compound is



In one embodiment, the compound is



.

In a further embodiment of the present invention, the compound has the structure:



; or



.

OV

In another embodiment of the present invention, at least one V is phenyl optionally substituted with one or more F; Cl; Br; $-OR_3$; $-COR_3$; $(CH_2)_qOR_3$; straight chained or branched C_1-C_7 alkyl; C_1-C_7 polyfluoroalkyl; aryl or phenoxy.

In yet another embodiment of the present invention, the compound is



In one embodiment, the compound is



In one embodiment, the compound is



In one embodiment, the compound is



In one embodiment, the compound is



In one embodiment, the compound is



In an embodiment of the present invention, the compound has the structure:



; or



In a further embodiment of the present invention, at least one V is phenyl optionally substituted with one or more F; Cl; Br; $-OR_3$; $(CH_2)_qOR_3$; straight chained or branched C_1-C_7 alkyl; C_1-C_7 polyfluoroalkyl; or phenoxy.

In yet another embodiment of the present invention, the compound is



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In an additional embodiment of the present invention, Y is hydrogen and V is phthalimide.

In an additional embodiment of the present invention, R_6 is straight chained or branched C_1-C_7 alkyl; C_3-C_7 cycloalkyl; $-N(R_3)_2$; $-OR_3$; $-(CH_2)_pOR_3$; aryl, benzyl or heteroaryl, optionally substituted with one or more F; Cl; Br; I; $-OR_3$; $-(CH_2)_qOR_3$; or straight chained or branched C_1-C_7 alkyl.

In a further embodiment of the present invention, the compound is



In one embodiment, the compound has the structure:



In one embodiment of the compound, at least one V is phenyl or heteroaryl optionally substituted with one or more F; Cl; Br; I; R_5; $-OR_5$; $-(CH_2)_qOR_5$; $-(CH_2)_qR_5$; straight chained or branched C_1-C_7 alkyl; C_1-C_7 monoflouroalkyl or polyflouroalkyl; or phenoxy.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment of the compound, V is phenyl which is optionally substituted with one or more F; Cl; Br; $-OR_5$; $-(CH_2)_qOR_5$; $-(CH_2)_qR_5$; straight chained or branched C_1-C_7 alkyl; C_1-C_7 monoflouroalkyl or polyflouroalkyl; or phenoxy.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment of the compound, R_5 is straight chained or branched C_1-C_7 alkyl; C_3-C_7 cycloalkyl; $-N(R_6)_2$; $-OR_6$; $-(CH_2)_qOR_6$; $-CH(R_6)_2$; $-(CH_2)_qR_6$; or aryl, benzyl or heteroaryl, wherein the aryl, benzyl or heteroaryl is optionally substituted with one or more F; Cl; I; R_6; $-N(R_6)_2$; $-OR_6$; $-(CH_2)_qOR_6$; $-(CH_2)_qR_6$; or straight chained or branched C_1-C_7 alkyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment of the compound, X is hydrogen and Y is carbazole optionally substituted with one or more F; Cl; Br; R_5; $-OR_5$; $-(CH_2)_qOR_5$; $-(CH_2)_qR_5$; straight chained or branched C_1 - C_7 alkyl; or C_1-C_7 monoflouroalkyl or polyflouroalkyl; or phenoxy.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment of the compound, Y is hydrogen and V is an indole, which can be optionally substituted with one or more F; Cl; Br; R_5; $-CO_2R_5$; $-OR_5$; $-(CH_2)_qOR_5$; $-(CH_2)_qR_5$; straight chained or branched C_1 - C_7 alkyl; C_1-C_7 monoflouroalkyl or polyflouroalkyl; or phenoxy on the 1, 2, 3, 4 , 5, 6 or 7 positions.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



The present invention provides a compond having the srucuture:



;



;

or



wherein each X is independently O or S;

wherein q is 1 or 2;

wherein each R_2 is independently H; $-(CH_2)_tXR_3$; $-(CH_2)_tC(X)N(R_3)_2$; $-(CH_2)_tCO_2R_3$; $-CO_2R_3$; straight chained or branched C_1-C_7 alkyl optionally substituted with $-N(R_3)_2$; $-CON(R_3)_2$ or $-N(R_3)C(O)R_3$; straight chained or branched C_2-C_7 alkenyl, or alkynyl; or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

wherein each t is independently an integer from 1 to 4 inclusive;

wherein each R_3 is independently H; straight chained or branched C_1-C_7 alkyl, straight chained or branched C_2-C_7 alkenyl, or alkynyl; or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

wherein R_4 is aryl, heteroaryl, C_1-C_7 alkyl substituted with one or two aryl, or C_1-C_7 alkyl substituted with one or two heteroaryl; wherein the aryl or heteroaryl may be substituted with one or more of F, Cl, Br, I, -CN, $-NO_2$, $-N(R_3)_2$, $-COR_3$, $-(CH_2)_nXR_3$, $-(CH_2)_nC(X)NR_3$, $-(CH_2)_nN(R_3)C(X)R_3$, $-(CH_2)_nCO_2R_3$, $-(CH_2)_nOCOR_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl OR polyfluoroalkyl or straight chained or branched C_2-C_7 aminoalkyl, alkenyl or alkynyl, or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

wherein each n independently is an integer from 0 to 7 inclusive;

wherein R_5 is H; aryl, C_1-C_7 alkyl substituted with aryl, heteroaryl, or C_1-C_7 alkyl substituted with heteroaryl; wherein the aryl or heteroaryl may be substituted with one or more of F, Cl, Br, I, -CN, $-NO_2$, $-N(R_3)_2$, $-COR_3$, $-(CH_2)_nXR_3$, $-(CH_2)_nC(X)NR_3$, $-(CH_2)_nCO_2R_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl, polyfluoroalkyl or carboxamidoalkyl, or straight chained or branched C_2-C_7 aminoalkyl, alkenyl or alkynyl, or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

where R_5 and one R_2 on adjacent carbon atoms together may form aryl, heteroaryl, indane or tetrahydronaphthyl, C_3-C_7 cycloalkyl, or heterocycloalkyl wherein one or two heteroatoms may be O, N or S;

wherein R_1 is



, or



;

;

wherein each V is independently aryl, phenoxy or heteroaryl, wherein the aryl, phenoxy or heteroaryl is optionally substituted with one or more F; Cl; Br; I; COR_5; CO_2R_5; $-OCOR_5$; $-CON(R_5)_2$; $-N(R_5)COR_5$; CN; $-NO_2$; $-N(R_5)_2$; $-OR_5$; $-SR_5$; $(CH_2)_qOR_5$; $(CH_2)_qSR_5$; straight chained or branched C_1-C_7 alkyl optionally substituted with $-CON(R_5)_2$, $-N(R_5)C(O)R_3$ or $N(R_3)_2$, straight chained or branched monofluoroalkyl or polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl, C_2-C_7 alkynyl; phenoxy; or C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein each R_6 is independently H; $(CH_2)_tXR_3$; $(CH_2)_tC(X)NR_3$; $(CH_2)_tN(R_3)C(X)R_3$; $(CH_2)_tCO_2R_3$; $(CH_2)_tOCOR_3$; straight chained or branched C_1-C_7 alkyl optionally substituted with $-CON(R_3)_2$ or $-NC(O)R_3$; straight chained or branched C_2-C_7 alkyl substituted with $-N(R_3)_2$; straight

chained or branched C_2-C_7 alkenyl or alkynyl; or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

where each R_7 is independently H; F; Cl; Br; I; $-COR_3$; $-CO_2R_3$; $-(CH_2)_nXR_3$; $-(CH_2)_nN(R_3)C(O)R_3$; $(CH_2)_nC(X)N(R_3)_2$; $-(CH_2)_nCO_2R_3$; -CN; $-NO_2$; $-N(R_3)_2$; straight chained or branched C_1-C_7 alkyl, hydroxyalkyl, aminoalkyl, carboxamidoalkyl, alkoxyalkyl, monofluoroalkyl or polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl or alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or C_5-C_7 cycloalkenyl, wherein the alkyl, aminoalkyl, carboxamidoalkyl, hydroxyalkyl, alkoxyalkyl, alkenyl, alkynyl, cycloalkyl or cycloalkenyl may be substituted with one or more aryl or heteroaryl, wherein the aryl or heteroaryl may be substituted with one or more of F, Cl, Br, I, $-(CH_2)_nXR_3$, $-COR_3$, $-(CH_2)_nC(X)N(R_3)_2$,$-(CH_2)_nCO_2R_3$, -CN, $-NO_2$, $-(CH_2)_nN(R_3)C(O)R_3$; $-N(R_3)_2$, $-SO_2R_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl or alkynyl, or C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or C_5-C_7 cycloalkenyl; aryl or heteroaryl, wherein the aryl or heteroaryl may be substituted with one or more of F, Cl, Br, I, $-(CH_2)_nXR_3$, $-COR_3$, $-(CH_2)_nC(X)N(R_3)_2$

$-(CH_2)_nCO_2R_3$, $-(CH_2)_nN(R_3)C(O)R_3$; -CN, $-NO_2$, $-N(R_3)_2$, $-SO_2R_3$, straight chained or branched C_1-C_7 alkyl, straight chained or branched C_1-C_7 monofluoroalkyl or polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl or alkynyl, or C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or C_5-C_7 cycloalkenyl;

wherein B is CO, SO_2 or SO_2NR_6;

wherein R_8 is -H; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl or alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl; $-N(R_3)_2$; $-NR_3C(O)R_3$; $-OR_3$; $-(CH_2)_pOR_3$; $-COR_3$; $-CO_2R_3$; $-OCOR_3$;$-CON(R_3)_2$; aryl or heteroaryl, optionally substituted with one or more F; Cl; Br; I; COR_3; CO_2R_3; $-OCOR_3$; $-NR_3C(O)R_3$; $-CON(R_3)_2$; CN; $-NO_2$; $-N(R_3)_2$; $-OR_3$; $-SR_3$; $(CH_2)_qOR_3$; $(CH_2)_qSR_3$; straight chained or branched C_1-C_7 alkyl optionally substituted with $-CON(R_3)_2$, $-NR_3C(O)R_3$ or $-N(R_3)_2$; straight chained or branched monofluoroalkyl, polyfluoroalkyl; straight chained or branched C_2-C_7 alkenyl, C_2-C_7 alkynyl; C_3-C_7 cycloalkyl, monofluorocycloalkyl, polyfluorocycloalkyl or cycloalkenyl;

wherein each m independently is an integer from 0 to 3 inclusive;

wherein Z is



,



,



,



,



,



,



, or

;

or C_2-C_7 alkenyl, wherein the C_2-C_7 alkenyl may be unsubstituted or substituted with one or more R_9 groups;

wherein each R_9 is independently H; F; Cl; Br; I; $-(CH_2)_mXR_3$; $(CH_2)_mC(X)NR_3$; $(CH_2)_mCO_2R_3$; straight chained or branched C_1-C_7 alkyl, monofluoroalkyl, polyfluoroalkyl, aminoalkyl, or carboxamidoalkyl; straight chained or branched C_2-C_7 alkenyl, or alkynyl; or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

wherein R_{10} is H; $(CH_2)_tXR_3$; $(CH_2)_tC(X)NR_3$; $(CH_2)_tCO_2R_3$; straight chained or branched C_1-C_7 alkyl, carboxamidoalkyl; straight chained or branched C_2-C_7 aminoalkyl, alkenyl, or alkynyl; or C_3-C_7 cycloalkyl or C_5-C_7 cycloalkenyl;

wherein Y is S, O, or NR_{10};

wherein each p is independently an integer from 1 to 7 inclusive;

or a pharmaceutically acceptable salt thereof.

In a further embodiment of the present invention, the compound has the following structure:



or



In an additional embodiment of the present invention, the compound has the structure:



; or



.

In an additional embodiment of the present invention, the compound has the structure:



.

In one embodiment of the present invention, Z is:



.

In one embodiment of the present invention, Z is:



In an additional embodiment of the present invention, the compound has the structure:



In one embodiment of the present invention, the compound has the strucuture:



This invention provides a compound having the structure:



;

wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CH_3$, $-CF_3$, $-COCH_3$, $-CO_2R_2$, phenyl, phenoxy or straight chained or branched C_1-C_7 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, $-COR_3$, $-CO_2R_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is O or NH;

wherein n is an integer from 0 to 5 inclusive;

In one embodiment, R_1 is aryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-COCH_3$,

$-CO_2R_2$, straight chained or branched C_1-C_7 alkyl;

wherein R_3 is phenyl;

wherein A is H; and

wherein X is O.

In one embodiment, R_2 is isopropyl.

In a preferred embodiment, X is NH, R_1 is alkyl and n is 1 or 2.

In the most preferred embodiment, X is O, R_1 is 3-acetyl phenyl, R_2 is isopropyl, R_3 is phenyl and n is 1.

In one embodiment, the compound has the structure:



In one embodiment, compound has the structure:



In one embodiment, R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl; and wherein R_3 is aryl.

In one embodiment, R_2 is isopropyl; and A is hydrogen.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



The present invention also provides a compound having the structure:


;

wherein R_1 is aryl or heteroaryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-OCH_3$, phenoxy, fused cyclopentanyl, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein R_2 is straight-chained or branched C_1-C_4 alkyl or cyclopropyl;

wherein A is -H, -F, -Cl, -Br, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; and

wherein n is an integer from 1 to 5 inclusive.

In one embodiment, R_1 is aryl optionally substituted with one or more -F, -Cl, -Br, -I or straight chained or branched C_1-C_4 alkyl; and

wherein A is H.

In one embodiment, R_2 is isopropyl; and

wherein n is 2.

In a preferred embodiment, n is 2 and R_2 is isopropyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, R_1 is thienyl; and wherein A is H.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



The invention provides a compound having the structure:



wherein W is



or



;

wherein each R_1 is independently hydrogen, methyl or ethyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or

wherein R_2 is straight- chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is hydrogen, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -H, -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl.

wherein each A is independently -H, -F, -Cl, -Br, -CN, -NO_2, -COR_3, -CO_2R_3, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is O, NR_3, CO or may be absent; and

wherein Y is hydrogen, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl.

In one embodiment, W is



;

and wherein X is O or may be absent.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, W is



In one embodiment, A is -H, -F, -Cl, -Br.

In one embodiment, R_2 is isopropyl; and A is hydrogen.

In one embodiment, the compound has the structure:



This invention provides a compound having the structure:



wherein W is



or



wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-OCH_3$, $-CO_2CH_3$, $-CF_3$, phenyl, straight chained or branched C_1-C_7 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein A is –H, –F, -Cl, -Br, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl or phenyl.

wherein each B is independently –H, –F, -Cl, -Br, -I, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, $-OCH_3$, $-OCF_3$, $-CF_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl or aryl, phenoxy or benzyloxy, wherein the aryl, phenoxy or benzyloxy is optionally substituted with one or more –F, -Cl, -Br, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$,

$-OCH_3$, $-OCF_3$, $-CF_3$ or straight chained or branched C1 -C3 alkyl.

In one embodiment, W is



R_1 .

In one embodiment, R_1 is hydrogen or phenyl optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl.

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



This invention provides a compound having the structure:



;

wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-CF_3$, $-OCH_3$, straight chained or branched C_1-C_3 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is -H, -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CF_3$, $-OCH_3$, or straight chained or branched C_1-C_3 alkyl, monofluoroalkyl or polyfluoroalkyl, or a phenyl ring fused to C_6 and C_7 of the indole moiety;

wherein R_4 is hydrogen or aryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CF_3$, straight chained or branched C_1-C_3 alkyl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; and

wherein n is an integer from 2 to 4 inclusive.

In one embodiment, R_3 is - H, -F, -Cl, -Br, -I, -CN, -NO_2, -OCF_3 or -OCH_3; and

wherein R_4 is hydrogen or phenyl optionally substituted with one or more -F, -Cl or -CF_3.

In one embodiment, R_1 is hydrogen or phenyl optionally substituted with one or more -F, -Cl, -Br, -CN, -NO_2, -CF_3, -OCH_3 or straight chained or branched C_1-C_3 alkyl;

In one embodiment, R_2 is isopropyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



This invention provides a compound having the structure:



;

wherein each R_1 is independently hydrogen or CH_3;

wherein R_2 is straight-chained or branched C_1-C_4 alkyl or cyclopropyl;

wherein R_3 is benzyl or phenyl, wherein the benzyl or phenyl is optionally substituted with a methylenenedioxy group or one or more -F or -Cl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is $(CH_2)_2$, $COCH_2$ or CONH;

In one embodiment, R_3 is phenyl optionally substituted with one or more -F; and

wherein A is hydrogen.

In one embodiment, X is CONH.

In one embodiment, R_2 is methyl.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



;

wherein each Y is independently hydrogen or -F.

In one embodiment, the compound has the structure:



In one embodiment, the compound has the structure:



In one embodiment, R_3 is benzyl optionally substituted with a methylenedioxy group or one or more -F or -Cl.

In one embodiment, the compound has the structure:



;

wherein each Y is independently hydrogen or -F.

In one embodiment, the compound has the structure:



In one embodiment, the compound is enantiomerically pure.

In one embodiment, the compound is diastereomerically pure.

In one embodiment, the compound is enantiomerically and diastereomerically pure.

This invention also provides a pharmaceutical composition comprising a therapeutically amount of a

compound of the invention and a pharmaceutically acceptable carrier.

In one embodiment, the amount of the compound is from about 0.01mg to about 500mg.

In one embodiment, the amount of the compound is from about 0.1mg to about 60mg.

In one embodiment, the amount of the compound is from about 1mg to about 20mg.

In one embodiment, the pharmaceutical composition consists of a carrier which is a liquid and the composition is a solution.

In one embodiment, the pharmaceutical composition consists of a carrier which is a solid and the composition is a tablet.

In one embodiment, the pharmaceutical composition consists of a carrier which is a gel and the composition is a suppository.

The invention also provides a process for making a pharmaceutical composition comprising admixing a therapeutically effective amount of the compound of any of the invention and a pharmaceutically acceptable carrier.

This invention also provides the method of treating a subject suffering from a disorder selected from the group consisting of depression, anxiety, urge

incontinence, or obesity comprising administering to the subject a therapeutically effective amount of the compound of the invention.

In one embodiment, the therapeutically effective amount is between about 0.03 and about 1000 mg per day.

In one embodiment, the therapeutically effective amount is between about 0.30 and about 300 mg per day.

In one embodiment, the therapeutically effective amount is between about 1.0 and about 100 mg per day.

In one embodiment, the disorder is depression.

In one embodiment, the disorder is anxiety.

In one embodiment, the disorder is obesity.

In one embodiment, the disorder is urge incontinence.

The invention provides the method of reducing the body mass of a subject, which comprises administering to the subject an amount of a compound of the invention effective to reduce the body mass of the subject.

The invention provides the method of treating a subject suffering from depression, which comprises administering to the subject an amount of a compound of any of claims of the invention effective to treat the subject's depression.

The invention provides the method of treating a subject suffering from anxiety, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's anxiety.

The invention provides the method of alleviating urge urinary incontinence in a subject suffering from an overactive bladder, which comprises administering to the subject an amount of the compound of the invention effective to alleviate the subject's urge urinary incontinence.

The invention provides the method of managing obesity in a subject in need of weight loss, which comprises administering to the subject an amount of a compound of the invention effective to induce weight loss in the subject.

The invention provides the method of managing obesity in a subject who has experienced weight loss, which comprises administering to the subject an amount of a compound of the invention effective to maintain such weight loss in the subject.

The invention provides the method of treating overactive bladder in a subject, which comprises administering to the subject an amount of a compound of any of the invention effective to treat the subject's overactive bladder.

The invention provides the method of treating a disorder in a subject, wherein the symptoms of the subject can be

alleviated by treatment with an MCH1 antagonist, wherein the MCH1 antagonist is the compound of the invention.

The invention provides the method of alleviating the symptoms of a disor4der in a subject, which comprises administering to the subject an amount of an MCH1 antagonist effective to alleviate the symptoms, wherein the MCH1 antagonist is the compound of the invention

As used in the present invention, the term "heteroaryl" is used to include five and six membered unsaturated rings that may contain one or more oxygen, sulfur, or nitrogen atoms. Examples of heteroaryl groups include, but are not limited to, carbazole, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl.

In addition, the term "heteroaryl" is used to include fused bicyclic ring systems that may contain one or more heteroatoms such as oxygen, sulfur and nitrogen. Examples of such heteroaryl groups include, but are not limited to, indolizinyl, indolyl, isoindolyl, benzo[b]furanyl, benzo[b]thiophenyl, indazolyl, benzimidazolyl, purinyl, benzoxazolyl, benzisoxazolyl, benzo[b]thiazolyl, imidazo[2,1-b]thiazolyl, cinnolinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, quinolinyl, isoquinolinyl, phthalimidyl and 2,1,3-benzothiazolyl.

The term "heteroaryl" also includes those chemical moieties recited above which may be substituted with one or more of the following: -F, -Cl, -Br, -I, CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, straight chained or branched C_1-C_7 monofluoroalkyl, straight chained or branched C_1-C_7 polyfluoroalkyl, straight chained or branched C_2-C_7 alkenyl, straight chained or branched C_2-C_7 alkynyl; C_3-C_7 cycloalkyl, C_3-C_7 monofluorocycloalkyl, C_3-C_7 polyfluorocycloalkyl, C_5-C_7 cycloalkenyl,

The term "heteroaryl" further includes the N-oxides of those chemical moieties recited above which include at least one nitrogen atom.

In the present invention, the term "aryl" is phenyl or naphthyl.

The invention provides for each pure stereoisomer of any of the compounds described herein. Such stereoisomers may include enantiomers, diastereomers, or E or Z alkene or imine isomers. The invention also provides for stereoisomeric mixtures, including racemic mixtures, diastereomeric mixtures, or E/Z isomeric mixtures. Stereoisomers can be synthesized in pure form (Nógrádi, M.; Stereoselective Synthesis, (1987) VCH Editor Ebel, H. and Asymmetric Synthesis, Volumes 3 B 5, (1983) Academic Press, Editor Morrison, J.) or they can be resolved by a variety of methods such as crystallization and chromatographic techniques (Jaques, J.; Collet, A.; Wilen, S.; Enantiomer, Racemates, and Resolutions, 1981,

John Wiley and Sons and Asymmetric Synthesis, Vol. 2, 1983, Academic Press, Editor Morrison, J).

In addition the compounds of the present invention may be present as enantiomers, diasteriomers, isomers or two or more of the compounds may be present to form a racemic or diastereomeric mixture.

The compounds of the present invention are preferably 80% pure, more preferably 90% pure, and most preferably 95% pure. Included in this invention are pharmaceutically acceptable salts and complexes of all of the compounds described herein. The acids and bases from which these salts are prepared include but are not limited to the acids and bases listed herein. The acids include, but are not limited to, the following inorganic acids: hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and boric acid. The acids include, but are not limited to, the following organic acids: acetic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, maleic acid, citric acid, methanesulfonic acid, benzoic acid, glycolic acid, lactic acid and mandelic acid. The bases include, but are not limited to ammonia, methylamine, ethylamine, propylamine, dimethylamine, diethylamine, trimethylamine, triethylamine, ethylenediamine, hydroxyethylamine, morpholine, piperazine and guanidine. This invention further provides for the hydrates and polymorphs of all of the compounds described herein.

The present invention includes within its scope prodrugs of the compounds of the invention. In general, such prodrugs will be functional derivatives of the compounds

of the invention which are readily convertible in vivo into the required compound. Thus, in the present invention, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound in vivo after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in Design of Prodrugs, ed. H. Bundgaard, Elsevier, 1985.

The present invention further includes metabolites of the compounds of the present invention. Metabolites include active species produced upon introduction of compounds of this invention into the biological milieu.

This invention further provides a pharmaceutical composition comprising a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier. In one embodiment, the amount of the compound is from about 0.01 mg to about 800 mg. In another embodiment, the amount of the compound is from about 0.01 mg to about 500 mg. In yet another embodiment, the amount of the compound is from about 0.1 mg to about 250 mg. In another embodiment, the amount of the compound is from about 0.1 mg to about 60 mg. In yet another embodiment, the amount of the compound is from about 1 mg to about 20 mg. In a further embodiment, the carrier is a liquid and the composition is a solution. In another embodiment, the carrier is a solid and the composition is a tablet. In another embodiment, the carrier is a gel and the

composition is a capsule, suppository or a cream. In a further embodiment the compound may be formulated as a part of a pharmaceutically acceptable transdermal patch. In yet a further embodiment, the compound may be delivered to the subject by means of a spray or inhalant.

This invention also provides a pharmaceutical composition made by combining a therapeutically effective amount of the compound of this invention and a pharmaceutically acceptable carrier.

This invention provides a process for making a pharmaceutical composition comprising combining a therapeutically effective amount of the compound of this invention and a pharmaceutically acceptable carrier.

A solid carrier can include one or more substances which may also act as endogenous carriers (e.g. nutrient or micronutrient carriers), flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch,

gelatin, cellulose, polyvinylpyrrolidine, low melting waxes and ion exchange resins.

Liquid carriers are used in preparing solutions, suspensions, emulsions, syrups, elixirs and pressurized compositions. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, coloring agents, viscosity regulators, stabilizers or osmoregulators. Suitable examples of liquid carriers for oral and parenteral administration include water (partially containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate or isopropyl myristate. Sterile liquid carriers are useful in sterile liquid form compositions for parenteral administration. The liquid carrier for pressurized compositions can be a halogenated hydrocarbon or other pharmaceutically acceptable propellent.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by for example, intramuscular, intrathecal, epidural, intraperitoneal or subcutaneous injection. Sterile solutions can also be

administered intravenously. The compounds may be prepared as a sterile solid composition which may be dissolved or suspended at the time of administration using sterile water, saline, or other appropriate sterile injectable medium. Carriers are intended to include necessary and inert binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. The compound can be administered orally in the form of a sterile solution or suspension containing other solutes or suspending agents (for example, enough saline or glucose to make the solution isotonic), bile salts, acacia, gelatin, sorbitan monoleate, polysorbate 80 (oleate esters of sorbitol and its anhydrides copolymerized with ethylene oxide) and the like.

The compound can also be administered orally either in liquid or solid composition form. Compositions suitable for oral administration include solid forms, such as pills, capsules, granules, tablets, and powders, and liquid forms, such as solutions, syrups, elixirs, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions, and suspenions.

Optimal dosages to be administered may be determined by those skilled in the art, and will vary with the particular compound in use, the strength of the preparation, the mode of administration, and the advancement of the disease condition. Additional factors depending on the particular subject being treated will result in a need to adjust dosages, including subject age, weight, gender, diet, and time of administration.

In the subject application a "therapeutically effective amount" is any amount of a compound which, when administered to a subject suffering from a disease against which the compounds are effective, causes reduction, remission, or regression of the disease. In a subject application, a "subject" is a vertebrate, a mammal or a human.

This invention provides a method of treating a subject suffering from an abnormality wherein the abnormality is alleviated by decreasing the activity of an MCH1 receptor which comprises administering to the subject an amount of a compound of the invention which is an MCH1 receptor antagonist effective to treat the subject=s abnormality.

In separate embodiments, the abnormality is a regulation of a steroid or pituitary hormone disorder, an epinephrine release disorder, a gastrointestinal disorder, a cardiovascular disorder, an electrolyte balance disorder, hypertension, diabetes, a respiratory disorder, asthma, a reproductive function disorder, an immune disorder, an endocrine disorder, a musculoskeletal disorder, a neuroendocrine disorder, a cognitive disorder, a memory disorder such as Alzheimer=s disease, a sensory modulation and transmission disorder, a motor coordination disorder, a sensory integration disorder, a motor integration disorder, a dopaminergic function disorder such as Parkinson=s disease, a sensory transmission disorder, an olfaction disorder, a sympathetic innervation disorder, an affective disorder such as depression and anxiety, a

stress-related disorder, a fluid-balance disorder, a seizure disorder, pain, psychotic behavior such as schizophrenia, morphine tolerance, opiate addiction, migraine or a urinary disorder such as urinary incontinence.

The following description of depressive and anxiety disorders is for the purpose of illustrating the utility of the compounds of this invention. The definitions of depressive and anxiety disorders given below are those listed in Diagnostic and Statistical Manual of Mental Disorders. 4th ed. (DSM-IV; American Psychiatric Association, 1994a) or Diagnostic and Statistical Manual of Mental Disorders. 3rd ed. Revised (DSM-III-R; American Psychiatric Association, 1987). Additional information regarding these disorders can be found in this reference, as well as the others cited below, all of which are incorporated herein by reference.

Depressive disorders include major depressive disorder and dysthymic disorder (American Psychiatric Association, 1994a; American Psychiatric Association, 1994b). Major depressive disorder is characterized by the occurrence of one or more major depressive episodes without manic or hypomanic episodes. A major depressive episode is defined as a prominent and relatively persistent depressed or dysphoric mood that usually interferes with daily functioning (nearly every day for at least 2 weeks); it should include at least 4 of the following 8 symptoms: change in appetite, change in sleep, psychomotor agitation or retardation, loss of interest in usual activities or decrease in sexual drive, increased fatigue, feelings of guilt or

worthlessness, slowed thinking or impaired concentration, and a suicide attempt or suicidal ideation (Medical Economics Company, 2002). Dysthymic disorder involves a type of depression that is not severe enough to be called a major depressive episode, but that lasts much longer than major depressive disorder, without high phases.

It is contemplated that the compounds of this invention will be effective in treating depression in patients who have been diagnosed with depression by administration of any of the following tests: Hamilton Depression Rating Scale (HDRS), Hamilton depressed mood item, Clinical Global Impressions (CGI)-Severity of Illness. It is further contemplated that the compounds of the invention will be effective in inducing improvements in certain of the factors measured in these tests, such as the HDRS subfactor scores, including the depressed mood item, sleep disturbance factor and anxiety factor, and the CGI-Severity of Illness rating. It is also contemplated that the compounds of this invention will be effective in preventing relapse of major depressive episodes.

Anxiety disorders include panic disorder, agoraphobia with or without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, post-traumatic stress disorder, acute stress disorder and generalized anxiety disorder. It is contemplated that the compounds of this invention will be effective in treating any of all of these disorders in patients who have been diagnosed with these disorders.

Obsessive compulsive disorder is characterized by recurrent and persistent ideas, thoughts, impulses or images (obsessions) that are ego-dystonic and/or repetitive, purposeful and intentional behaviors (compulsions) that are recognized by the person as excessive or unreasonable (American Psychiatric Association, 1994a). The obsessions or compulsions cause marked distress, are time-consuming, or significantly interfere with social or occupational functioning.

It is contemplated that the compounds of this invention will be effective in treating obsessions and compulsions in patients who have been diagnosed with obsessive compulsive disorder by administration of appropriate tests, which may include, but are not limited to any of the following: Yale Brown Obsessive Compulsive Scale (YBOCS) (Goodman, 1989) (for adults), National Institute of Mental Health Global OCD Scale (NIMH GOCS), CGI-Severity of Illness scale. It is further contemplated that the compounds of the invention will be effective in inducing improvements in certain of the factors measured in these tests, such as a reduction of several points in the YBOCS total score. It is also contemplated that the compounds of this invention will be effective in preventing relapse of obsessive compulsive disorder.

Panic disorder is characterized by recurrent unexpected panic attacks and associated concern about having additional attacks, worry about the implications or consequences of the attacks, and/or a significant change in behavior related to the attacks (American Psychiatric Association, 1994a). A panic attack is defined as a

discrete period of intense fear or discomfort in which four (or more) of the following symptoms develop abruptly and reach a peak within 10 minutes: (1) palpitations, pounding heart, or accelerated heart rate; (2) sweating; (3) trembling or shaking; (4) sensations of shortness of breath or smothering; (5) feeling of choking; (6) chest pain or discomfort; (7) nausea or abdominal distress; (8) feeling dizzy, unsteady, lightheaded, or faint; (9) derealization (feelings of unreality) or depersonalization (being detached from oneself); fear of losing control; (11) fear of dying; (12) paresthesias (numbness or tingling sensations); (13) chills or hot flushes. Panic disorder may or may not be associated with agoraphobia, or an irrational and often disabling fear of being out in public.

It is contemplated that the compounds of this invention will be effective in treating panic disorder in patients who have been diagnosed with panic disorder on the basis of frequency of occurrence of panic attacks, or by means of the CGI-Severity of Illness scale. It is further contemplated that the compounds of the invention will be effective in inducing improvements in certain of the factors measured in these evaluations, such as a reduction in frequency or elimination of panic attacks, an improvement in the CGI-Severity of Illness scale or a CGI-Global Improvement score of 1 (very much improved), 2 (much improved) or 3 (minimally improved).. It is also contemplated that the compounds of this invention will be effective in preventing relapse of panic disorder.

Social anxiety disorder, also known as social phobia, is characterized by a marked and persistent fear of one or

more social or performance situations in which the person is exposed to unfamiliar people or to possible scrutiny by others (American Psychiatric Association, 1994a). Exposure to the feared situation almost invariably provokes anxiety, which may approach the intensity of a panic attack. The feared situations are avoided or endured with intense anxiety or distress. The avoidance, anxious anticipation, or distress in the feared situation(s) interferes significantly with the person's normal routine, occupational or academic functioning, or social activities or relationships, or there is marked distress about having the phobias. Lesser degrees of performance anxiety or shyness generally do not require psychopharmacological treatment.

It is contemplated that the compounds of this invention will be effective in treating social anxiety disorder in patients who have been diagnosed with social anxiety disorder by administration of any of the following tests: the Liebowitz Social Anxiety Scale (LSAS), the CGI-Severity of Illness scale, the Hamilton Rating Scale for Anxiety (HAM-A), the Hamilton Rating Scale for Depression (HAM-D), the axis V Social and Occupational Functioning Assessment Scale of DSM-IV, the axis II (ICD-10) World Health Organization Disability Assessment, Schedule 2 (DAS-2), the Sheehan Disability Scales, the Schneier Disability Profile., the World Health Organization Quality of Life-100 (WHOQOL-100), or other tests as described in Bobes, 1998, which is incorporated herein by reference. It is further contemplated that the compounds of the invention will be effective in inducing improvements as measured by these

tests, such as the a change from baseline in the Liebowitz Social Anxiety Scale (LSAS), or a CGI- Global Improvement score of 1 (very much improved), 2 (much improved) or 3 (minimally improved). It is also contemplated that the compounds of this invention will be effective in preventing relapse of social anxiety disorder.

Generalized anxiety disorder is characterized by excessive anxiety and worry (apprehensive expectation) that is persistent for at least 6 months and which the person finds difficult to control (American Psychiatric Association, 1994a). It must be associated with at least 3 of the following 6 symptoms: restlessness or feeling keyed up or on edge, being easily fatigued, difficulty concentrating or mind going blank, irritability, muscle tension, sleep disturbance. The diagnostic criteria for this disorder are described in further detail in DSM-IV, which is incorporated herein by reference (American Psychiatric Association, 1994a).

It is contemplated that the compounds of this invention will be effective in treating generalized anxiety disorder in patients who have been diagnosed with this disorder according to the diagnostic criteria described in DSM-IV. It is further contemplated that the compounds of the invention will be effective in reducing symptoms of this disorder, such as the following: excessive worry and anxiety, difficulty controlling worry, restlessness or feeling keyed up or on edge, being easily fatigued, difficulty concentrating or mind going blank, irritability, muscle tension, or sleep disturbance. It is also contemplated that the compounds

of this invention will be effective in preventing relapse of general anxiety disorder.

Post-traumatic stress disorder (PTSD), as defined by DSM-III-R/IV (American Psychiatric Association, 1987, American Psychiatric Association, 1994a), requires exposure to a traumatic event that involved actual or threatened death or serious injury, or threat to the physical integrity of self or others, and a response which involves intense fear, helplessness, or horror. Symptoms that occur as a result of exposure to the traumatic event include re-experiencing of the event in the form of intrusive thoughts, flashbacks or dreams, and intense psychological distress and physiological reactivity on exposure to cues to the event; avoidance of situations reminiscent of the traumatic event, inability to recall details of the event, and/or numbing of general responsiveness manifested as diminished interest in significant activities, estrangement from others, restricted range of affect, or sense of foreshortened future; and symptoms of autonomic arousal including hypervigilance, exaggerated startle response, sleep disturbance, impaired concentration, and irritability or outbursts of anger. A PTSD diagnosis requires that the symptoms are present for at least a month and that they cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.

It is contemplated that the compounds of this invention will be effective in treating PTSD in patients who have been diagnosed with PTSD by administration of any of the following tests: Clinician-Administered PTSD Scale Part

2 (CAPS), the patient-rated Impact of Event Scale (IES) (Medical Economics Company, 2002, p. 2752). It is further contemplated that the compounds of the invention will be effective in inducing improvements in the scores of the CAPS, IES, CGI-Severity of Illness or CGI-Global Improvement tests. It is also contemplated that the compounds of this invention will be effective in preventing relapse of PTSD.

In a preferred embodiment, the subject invention provides a method of treatment or management of the following indications: depressive disorders, anxiety disorders, eating/body weight disorders, and urinary disorders. Examples of depressive disorders are major depressive disorder or dysthymic disorder. Examples of anxiety disorders are panic disorder, agoraphobia without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, post-traumatic stress disorder, acute stress disorder or generalized anxiety disorder. Examples of eating/body weight disorders are obesity, weight gain, bulimia, bulimia nervosa or anorexia nervosa. Examples of urinary disorders include, but are not limited to urinary incontinence overactive bladder, urge incontinence, urinary frequency, urinary urgency, nocturia or enuresis. Overactive bladder and urinary urgency may or may not be associated with benign prostatic hyperplasia.

This invention provides a method of modifying the feeding behavior of a subject, which comprises administering to the subject an amount of a compound of the invention effective to decrease the consumption of

food by the subject. This invention also provides a method of treating an eating disorder in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the eating disorder. In an embodiment of the present invention, the eating disorder is obesity, bulimia, bulimia nervosa or anorexia nervosa.

The present invention further provides a method of reducing the body mass of a subject, which comprises administering to the subject an amount of a compound of the invention effective to reduce the body mass of the subject. This invention also provides a method of managing obesity in a subject in need of weight loss, which comprises administering to the subject an amount of a compound of the invention effective to induce weight loss in the subject. This invention also provides a method of managing obesity in a subject who has experienced weight loss, which comprises administering to the subject an amount of a compound of the invention effective to maintain such weight loss in the subject.

The present invention also provides a method of treating depression in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's depression. This invention also provides a method of treating anxiety in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's anxiety. This invention also provides a method of treating depression and anxiety in a subject, which comprises administering to the subject

an amount of a compound of the invention effective to treat the subject's depression and anxiety. This invention also provides a method of treating major depressive disorder in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's major depressive disorder. This invention also provides a method of treating dysthymic disorder in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's dysthymic disorder. This invention also provides a method of treating obsessions and compulsions in a subject with obsessive compulsive disorder, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's obsessions and compulsions. This invention also provides a method of treating panic disorder, with or without agoraphobia, in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's panic disorder. This invention also provides a method of treating social anxiety disorder in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's social anxiety disorder. This invention also provides a method of treating generalized anxiety disorder in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's generalized anxiety disorder. This invention also provides a method of treating post-traumatic stress disorder in a subject, which comprises administering to the subject an amount

of a compound of the invention effective to treat the subject's post-traumatic stress disorder.

It is contemplated that the compounds of this invention will be effective in treating obesity, including weight loss and maintenance of weight loss in patients, who have been diagnosed with obesity by the one or more of the following measurements: an increased body mass index, increased waist circumference (an indicator of intra-adominal fat), Dual Energy X-Ray Absorptiometry (DEXA) and trucal (android) fat mass. It is further contemplated that the compounds of the invention will be effective in inducing improvements in certain factors measured in these tests.

It is contemplated that the compounds of this invention will be effective in treating urinary disorders in patients who have urge or mixed (with a predominance of urge) incontinence as evidenced by the number of unnecessary episodes per week, the number of unnecessary micturitions per day and a low volume voided per micturition. It is further contemplated that the compounds of the invention will be effective in inducing improvements in certain factors measured in these tests.

The present invention also provides a method of treating a subject suffering from bipolar I or II disorder, schizoaffective disorder, a cognitive disorder with depressed mood, a personality disorder, insomnia, hypersomnia, narcolepsy, circadian rhythm sleep disorder, nightmare disorder, sleep terror disorder or sleepwalking disorder.

The present invention provides a method of treating overactive bladder with symptoms of urge urinary incontinence, urgency and/or frequency in a subject, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's overactive bladder. This invention also provides a method of alleviating urge urinary incontinence in a subject suffering from overactive bladder, which comprises administering to the subject an amount of a compound of the invention effective to alleviate the subject's urge urinary incontinence. This invention further provides a method of alleviating urinary urgency in a subject suffering from overactive bladder, which comprises administering to the subject an amount of a compound of the invention effective to alleviate the subject's urinary urgency. Additionally, this invention provides a method of alleviating urinary frequency in a subject suffering from overactive bladder, which comprises administering to the subject an amount of a compound of the invention effective to alleviate the subject's urinary frequency.

The present invention also provides a method of treating a subject suffering from a urinary disorder, which comprises administering to the subject an amount of a compound of the invention effective to treat the subject's urinary disorder. In some embodiments the urinary disorder is urinary incontinence, overactive bladder, urge incontinence, urinary frequency, urinary urgency, nocturia or enuresis.

The present invention provides a method of alleviating the symptoms of a disorder in a subject, which comprises administering to the subject an amount of an MCH1 antagonist effective to alleviate the symptoms, wherein the MCH1 antagonist is any of the compounds of the invention.

In an embodiment of the invention, the subject is a vertebrate, a mammal, a human or a canine. In another embodiment, the compound is administered orally. In yet another embodiment, the compound is administered in combination with food.

This invention will be better understood from the Experimental Details In a preferred embodiment, the subject invention provides a method of treatment for the following indications: depression, anxiety, eating/body weight disorders, and urinary disorders. Examples of eating/body weight disorders are obesity, bulimia, or bulimia nervosa. Examples of urinary disorders include, but are not limited to, urinary incontinence, overactive bladder, urge incontinence, urinary frequency, urinary urgency, nocturia, or enuresis. Overactive bladder and urinary urgency may or may not be associated with benign prostatic hyperplasia.

This invention provides a method of modifying the feeding behavior of a subject which comprises administering to the subject an amount of a compound of the invention effective to decrease the consumption of food by the subject.

This invention also provides a method of treating an eating disorder in a subject which comprises administering to the subject an amount of a compound of this invention effective to decrease the consumption of food by the subject. In an embodiment of the present invention, the eating disorder is bulimia, obesity or bulimia nervosa. In an embodiment of the present invention, the subject is a vertebrate, a mammal, a human or a canine. In a further embodiment, the compound is administered in combination with food.

The present invention further provides a method of reducing the body mass of a subject which comprises administering to the subject an amount of a compound of the invention effective to reduce the body mass of the subject.

The present invention also provides a method of treating a subject suffering from depression which comprises administering to the subject an amount of a compound of this invention effective to treat the subject's depression. The present invention further provides a method of treating a subject suffering from anxiety which comprises administering to the subject an amount of a compound of this invention effective to treat the subject's anxiety. The present invention also provides a method of treating a subject suffering from depression and anxiety which comprises administering to the subject an amount of a compound of this invention effective to treat the subject's depression and anxiety.

The present invention also provides a method of treating a subject suffering from major depressive disorder,

dysthymic disorder, bipolar I and II disorders, schizoaffective disorder, cognitive disorders with depressed mood, personality disorders, insomnia, hypersomnia, narcolepsy, circadian rhythm sleep disorder, nightmare disorder, sleep terror disorder, sleepwalking disorder, obsessive-compulsive disorder, panic disorder, with or without agoraphobia, posttraumatic stress disorder, social anxiety disorder, social phobia and generalized anxiety disorder.

The present invention also provides a method of treating a subject suffering from a urinary disorder which comprises administering to the subject an amount of a compound of this invention effective to treat the subject's a urinary disorder. In some embodiments, the urinary disorder is urinary incontinence, overactive bladder, urge incontinence, urinary frequency, urinary urgency, nocturia, or enuresis.

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

Experimental Section

I. Synthetic Methods for Examples

General Methods: All reactions (except for those done by parallel synthesis reaction arrays) were performed under an Argon atmosphere and the reagents, neat or in appropriate solvents, were transferred to the reaction vessel via syringe and cannula techniques. The parallel synthesis reaction arrays were performed in vials (without an inert atmosphere) using J-KEM heating shakers (Saint Louis, MO). Anhydrous solvents were purchased from Aldrich Chemical Company and used as received. The examples described in the patent were named using ACD/Name program (version 2.51, Advanced Chemistry Development Inc., Toronto, Ontario, M5H2L3, Canada). Unless otherwise noted, the ^{1}H spectra were recorded at 300 and 400 MHz (QE Plus and Brüker respectively) with tetramethylsilane as internal standard. s = singlet; d = doublet; t = triplet; q = quartet; p = pentet; sext; sept; br = broad; m = multiplet. Elemental analyses were performed by Robertson Microlit Laboratories, Inc. Unless otherwise noted, mass spectra were obtained using low-resolution electrospray (ESMS) and MH^{+} is reported. Thin-layer chromatography (TLC) was carried out on glass plates precoated with silica gel 60 F_{254} (0.25 mm, EM Separations Tech.). Preparative thin-layer chromatography was carried out on glass sheets precoated with silica gel GF (2 mm, Analtech). Flash column chromatography was performed on Merck silica gel 60 (230 - 400 mesh). Melting points (mp) were determined in

open capillary tubes on a Mel-Temp apparatus and are uncorrected.

Piperidine Side Chain Intermediates

***TERT*-BUTYL 4-{[(TRIFLUOROMETHYL)SULFONYL]OXY}-1,2,3,6-TETRAHYDRO-1-PYRIDINECARBOXYLATE:** *n*-Butyl lithium (17.6 mL, 44.2 mmol, 2.5 M in hexanes) was added to a solution of diisopropyl amine (96.2 mL, 44.2 mmol) in 40 mL of dry THF at 0 °C and stirred for 20 minutes. The reaction mixture was cooled to -78 °C and *tert*-butyl 4-oxo-1-piperidinecarboxylate (Aldrich Chemical Company, 40.0 mmol) in THF (40 mL) was added dropwise to the reaction mixture and stirred for 30 minutes. Tf_2NPh (42.0 mmol, 15.0 g) in THF (40 mL) was added dropwise to the reaction mixture and stirred at °C overnight. The reaction mixture was concentrated *in vacuo*, re-dissolved in hexanes:EtOAc (9:1), passed through a plug of alumina and the alumina plug was washed with hexanes:EtOAc (9:1). The combined extracts were concentrated to yield 16.5 g of the desired product that was contaminated with some starting Tf_2NPh.

^1H NMR (400 MHz, 400 MHz, $CDCl_3$) δ 5.77 (s, 1 H), 4.05 (dm, 2 H, J=3.0 Hz), 3.63 (t, 2 H, J=5.7 Hz), 2.45 (m, 2 H), 1.47 (s, 9 H).

***TERT*-BUTYL 4-[3-(AMINO)PHENYL]-1,2,3,6-TETRAHYDRO-1-PYRIDINECARBOXYLATE:** A mixture of 2 M aqueous Na_2CO_3 solution (4.2 mL), *tert*-butyl 4-{[(trifluoromethyl)sulfonyl]oxy}-1,2,3,6-tetrahydro-1-pyridine-carboxylate (0.500 g, 1.51 mmol), 3-aminophenylboronic acid hemisulfate (0.393 g, 2.11 mmol), lithium chloride (0.191 g, 4.50 mmol) and

tetrakis-triphenylphosphine palladium (0) (0.080 g, 0.075 mmol) in dimethoxyethane (5 mL) was heated at reflux temperature for 3 hours, under an inert atmosphere (an initial degassing of the mixture is recommended to prevent the formation of triphenylphosphine oxide). The organic layer of the cooled reaction mixture was separated and the aqueous layer was washed with ethyl acetate (3X). The combined organic extracts were dried and concentrated *in vacuo*. The crude product was chromatographed (silica, hexanes:EtOAc:dichloromethane (6:1:1) with 1% added isopropylamine to protect the BOC group from hydrolysis) to give 0.330 g of the desired product in 81% yield. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.12 (t, 1H, J= 7.60 Hz), 6.78 (d, 1H, J= 8.4 Hz), 6.69 (t, 1H, J= 2.0 Hz), 6.59 (dd, 1H, J= 2.2, 8.0 Hz), 6.01 (m, 1H), 4.10 - 4.01 (d, 2H, J= 2.4 Hz), 3.61 (t, 2H, J= 5.6 Hz), 2.52 - 2.46 (m, 2H), 1.49 (s, 9H); ESMS m/e : 275.2 $(M + H)^+$. Anal. Calc. for $C_{16}H_{24}N_2O_2$: C, 70.04; H, 8.08; N, 10.21. Found: C, 69.78; H, 7.80; N, 9.92.

***TERT*-BUTYL 4-[3-(AMINO)PHENYL]-1-PIPERIDINECARBOXYLATE:**

A mixture of 3.10 g of tert-butyl 4-(3-aminophenyl)-1,2,3,6-tetrahydropyridine-1-carboxylate (11.3 mmol) and 1.0 g of 10% Pd/C in 200 mL of ethanol was hydrogenated at room temperature using the balloon method for 2 days. The reaction mixture was filtered and washed with ethanol. The combined ethanol extracts were concentrated *in vacuo* and the residue was chromatographed on silica (dichloromethane: methanol 95:5 with 1% isopropylamine added to protect the BOC group from hydrolysis) to give 2.63 g of the desired product (84%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.10 (t, 1H, J=

7.60 Hz), 6.62 (d, 1H, J= 8.4 Hz), 6.60 - 6.59 (m, 2H), 4.27 - 4.18 (m, 2H), 3.62 - 3.58 (m, 2H), 2.80 - 2.72 (m, 2H), 2.62 - 2.59 (m, 1H), 1.89 - 1.52 (m, 4H), 1.49 (s, 9H); ESMS m/e : 277.2 $(M + H)^+$.

***TERT*-BUTYL 4-[3-(ACETYLAMINO)PHENYL]-1,2,3,6-TETRAHYDRO-1-PYRIDINECARBOXYLATE:** A mixture of saturated aqueous Na_2CO_3 solution (25 mL), *tert*-butyl 4-{[(trifluoromethyl)sulfonyl]oxy}-1,2,3,6-tetrahydro-1-pyridine-carboxylate (20 mmol), 3-acetamidophenylboronic acid (30 mmol) and tetrakis-triphenylphosphine palladium (0) (1.15 g) and dimethoxyethane (40 mL) was heated at reflux temperature overnight. The organic layer of the cooled reaction mixture was separated and the aqueous layer was washed with ethyl acetate (3X). The combined organic extracts were dried and concentrated *in vacuo*. The crude product was chromatograghed, giving the desired product: ^1H NMR ($CDCl_3$) δ 8.11 (br s, 1 H), 7.57 (br s, 1 H), 7.41 (br d , 1 H, J=7.8 Hz), 7.25 (apparent t, 1 H, J=7.8 Hz), 7.08 (br d, 1 H, J=7.8 Hz), 5.99 (br s, 1 H), 4.03 (br m, 2 H, J=2.7 Hz), 3.59 (t, 2 H, J=5.7 Hz), 2.46 (m, 2 H,), 2.16 (s, 3 H), 1.49 (s, 9 H).

***N*1-[3-(1,2,3,6-TETRAHYDRO-4-PYRIDINYL)PHENYL]ACETAMIDE:** A solution of 4 M HCl in dioxane (10 mL) was added to *tert*-butyl 4-[3-(acetylamino)phenyl]-1,2,3,6-tetrahydro-1-pyridinecarboxylate (8.25 mmol) in dichloromethane (30 mL). The reaction mixture was stirred at room temperature overnight, concentrated *in vacuo*, giving the desired product as the hydrochloride salt (2.1 g): ^1H NMR ($CDCl_3$) δ 7.41-7.00 (m, 4 H), 6.10 (br, 1 H), 3.55 (m, 2

H), 3.16 (t, 2 H, J = 5.7 Hz), 2.44 (m, 2 H), 2.19 (s, 3 H).

***TERT*-BUTYL *N*-(3-BROMOPROPYL)CARBAMATE:** Prepared from 3-bromopropylamine hydrobromide and BOC_2O in the presence of base in dichloromethane, 9.89 mmol: ^1H NMR ($CDCl_3$) δ 5.07 (br, 1 H), 3.31 (t, 2 H, J=6.6 Hz), 3.12 (apparent br q, 2 H, J=6.0 Hz), 1.92 (p, 2 H, J=6.6 Hz), 1.30 (s, 9H).

***TERT*-BUTYL *N*-(3-{4-[3-(ACETYLAMINO)PHENYL]-1,2,3, 6-TETRAHYDRO-1-PYRIDINYL}PROPYL)CARBAMATE:** A solution of *N*1-[3-(1,2,3,6-tetrahydro-4-pyridinyl)phenyl]acetamide.HCl (8.24 mmol), *tert*-butyl *N*-(3-bromopropyl)carbamate and potassium carbonate (33 mmol) in dry dioxane (30 mL) was heated at reflux temperature overnight. The solids were removed by filtration, the solution was concentrated *in vacuo* and the product was chromatograghed, giving the desired product (110 mg). ^1H NMR ($CDCl_3$) δ 7.65 (s, 1 H), 6.98 (s, 1 H), 7.45 (d, 1 H, J=7.8 Hz), 7.16 (apparent t, 1 H, J=7.8 Hz), 7.10 (d, 1 H, J=7.8 Hz), 6.02 (s, 1 H), 5.23 (b, 1 H), 3.40 (b, 2 H), 3.30-1.80 (m, 10 H), 2.18 (s, 3 H), 1.45 (s, 9 H).

***N*1-{3-[1-(3-AMINOPROPYL)-1,2,3,6-TETRAHYDRO-4-PYRIDINYL]PHENYL}ACETAMIDE:** A 1:1 solution of TFA:CH_2Cl_2 (5 mL) was added to *tert*-butyl *N*-(3-{4-[3-(acetylamino)phenyl]-1,2,3,6-tetrahydro-1-pyridinyl}propyl)carbamate in dichloromethane (5 mL). The resulting solution was stirred at room temperature for 1-3 days, saturated $NaHCO_3$ was added until pH > 6, the organic layer was separated, and dried *in vacuo*,

giving the desired product (45 mg): ^{1}H NMR ($CDCl_3$) δ 7.68 (br, 1 H), 7.35 (dm, 1 H, J=7.8 Hz), 7.25 (apparent t, 1 H, J=7.8 Hz), 7.15 (dm, 1 H, J=7.8 Hz), 6.12 (m, 1 H), 3.22 (m, 2 H), 3.03 (t, 2 H, J=7.3 Hz), 2.78 (t, 2 H, J=5.5 Hz), 2.70-2.50 (m, 4 H), 2.10 (s, 3 H), 1.87 (p, 2 H, J=7.3 Hz).

***TERT*-BUTYL 4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINECARBOXYLATE:** A mixture *tert*-butyl 4-[3-(acetylamino)phenyl]-1,2,3,6-tetrahydro-1-pyridinecarboxylate (710 mg) and 5% Pd/C (100 mg) in EtOH (10 mL) was hydrogenated (balloon technique) at room temperature overnight. The reaction mixture was passed through a pad of Celite 545 and the pad of Celite was washed with ethanol. The combined ethanol extracts were concentrated and chromatograghed, giving the desired product (660 mg): ^{1}H NMR ($CDCl_3$) δ 7.80 (s, 1 H), 7.41-7.20 (m, 3 H), 6.94 (d, 1 H, J=7.5 Hz), 4.21 (m, 2 H), 2.75 (m, 2 H), 2.62 (m, 1 H), 2.16 (s, 3 H), 1.78 (m, 2 H), 1.56 (m, 2 H), 1.48 (s, 9 H).

***N*1-[3-(4-PIPERIDYL)PHENYL]ACETAMIDE:** A solution of HCl in dioxane (4N, 5 mL) was added to *tert*-butyl 4-[3-(acetylamino)phenyl]-1-piperidinecarboxylate (660 mg) in dry dichloromethane (15 mL). The reaction mixture was stirred at room temperature overnight and concentrated *in vacuo*, giving the desired product (550 mg): mp 102-104 °C; ^{1}H NMR ($CDCl_3$) δ 2.02 (d, J=13.2 Hz, 2H), 2.11-2.45 (m, 5H), 2.67-2.77 (m, 1H), 3.00-3.10 (m, 2H), 3.51 (d, J=10.5 Hz, 2H), 6.94 (d, J=7.5 Hz, 1H), 7.20-7.46 (m, 3H), 7.60 (s, 1H); Anal. Calcd. For $C_{13}H_{19}N_2OCl$+0.86 CH_2Cl_2: C, 50.78; H, 6.37; N, 8.55. Found: C, 50.80; H, 7.55; N, 7.01.

***TERT*-BUTYL *N*-(3-{4-[3-(ACETYLAMINO)PHENYL]PIPERIDINO}PROPYL)CARBAMATE:** A solution of *N*1-[3-(4-piperidyl)phenyl]acetamide (550 mg, 0.210 mmol), *tert*-butyl *N*-(3-bromopropyl)carbamate (550 mg, 0.230 mmol), K_2CO_3 (1.10 g, 0.890 mmol), diisopropylethyl amine (1.50 mL) and a few crystals of KI in dioxane (20 mL) was heated at reflux temperature for 2 days. The precipitated salts were removed by filtration, concentrated *in vacuo* and the crude product was chromatographed, giving the desired product (340 mg): ^1H NMR ($CDCl_3$) δ 8.15 (s, 1 H), 7.47-7.44 (m, 2 H), 7.22 (t, 1 H, J=7.8 Hz), 6.94 (d, 1 H, J=7.8 Hz), 5.53 (b, 1 H), 3.23 (b, 6 H), 2.80-1.60 (m, 9 H), 2.20 (s, 3 H), 1.45 (s, 9 H).

***N*1-{3-[1-(3-AMINOPROPYL)-4-PIPERIDYL]PHENYL}ACETAMIDE:** TFA (1.0 mL) was added to a solution of *tert*-butyl *N*-(3-{4-[3-(acetylamino)phenyl]piperidino}propyl)carbamate (340 mg) in dry dichloromethane (10 mL) and stirred at room temperature for 5 h. A 10% aqueous solution of KOH was added to the reaction mixture until pH > 6 and then the dichloromethane was removed *in vacuo*. The aqueous layer was frozen and lyophilized to give a solid, which was extracted with methanol. Removal of the solvent gave the desired product (120 mg) as an oil: ^1H NMR ($CDCl_3$) δ 7.23-7.16 (apparent t, 1H, J=7.5 Hz), 6.95-6.92 (m, 1H), 3.03-2.99 (m, 2H), 2.77-2.73 (t, 2H, J = 6.6 Hz), 2.50-1.60 (m, 10 H), 2.13 (s, 3 H).

TERT-BUTYL 4-(3-NITROPHENYL)-3,6-DIHYDRO-1(2H)-PYRIDINECARBOXYLATE: According to the procedure used for the synthesis of *tert*-butyl 4-[3-(amino)phenyl]-1,2,3,6-tetrahydro-1-pyridinecarboxylate,a mixture of 2 M aqueous Na_2CO_3 solution (2.2 mL), *tert*-butyl 4-{[(trifluoromethyl)sulfonyl]oxy}-1,2,3,6-tetrahydro-1-pyridine-carboxylate (0.500 g, 1.51 mmol), 3-nitrophenylboronic acid (0.353 g, 2.11 mmol), lithium chloride (0.191 g, 4.50 mmol) and tetrakis-triphenylphosphine palladium (0) (0.080 g, 0.075 mmol) in dimethoxyethane (5 mL) afforded 0.380g of the desired product.

^1H NMR (400 MHz, $CDCl_3$) δ 8.23 (s, 1H), 8.11 (d, 1H, J=8.0 Hz), 7.69 (d, 1H, J=8.0 Hz), 7.51 (t, 1H, J=8.0 Hz), 6.20 (m, 1H), 4.17-4.08 (m, 2H), 3.67 (t, 2H, J=5.6 Hz), 2.61-2.52 (m, 2H), 1.50 (s, 9H); ESMS m/e : 249.1 $(M + H - C_4H_8)^+$.

1,2,3,6-TETRAHYDRO-4-(3-NITROPHENYL)PYRIDINE: Into a stirred solution of 5.00 g (16.0 mmol) of *tert*-butyl 1,2,3,6-tetrahydro-4-(3-nitrophenyl)pyridine-1-carboxylate in 100 ml of 1,4-dioxane at 0°C was bubbled HCl gas for 10 minutes. The reaction mixture was allowed to warm to room temperature and the bubbling of the HCl gas was continued for an additional 1 hour. The solvent was removed *in vacuo*, the residue was dissolved in 50 mL of water and was neutralized by the addition of KOH pellets. The aqueous solution was extracted with 3 X 80 mL of dichloromethane and the combined organic extracts were dried ($MgSO_4$), filtered and concentrated *in vacuo*. The residue was purified by column chromatography (silica, 9 : 1 ,dichloromethane :

methanol + 1% isopropyl amine) to afford 2.85 g (87.5% yield) of the desired product: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.24 (s, 1H), 8.09 (d, 1H, J=8.4 Hz), 7.71 (d, 1H, J=8.0 Hz), 7.49 (t, 1H, J=8.0 Hz), 6.35-6.25 (m, 1H), 3.58 (apparent q, 2H, J=3.0 Hz), 3.14 (t, 2H, J=5.6 Hz), 2.54-2.46 (m, 2H).

TERT-BUTYL 3-(4-(3-NITROPHENYL)-3,6-DIHYDRO-1(2H)-PYRIDINYL)PROPYLCARBAMATE: A mixture of 2.80 g (14.0 mmol) of 1,2,3,6-tetrahydro-4-(3-nitrophenyl)pyridine, 3.60 g (15.0 mmol) of *tert*-butyl N-(3-bromopropyl)carbamate, 11.6 g (84.0 mmol) of K_2CO_3, 14.6 mL (84.0 mmol) of diisopropylethylamine and 0.78 g (2.00 mmol) of tetrabutylammonium iodide in 250 mL of 1,4-dioxane was heated at reflux temperature for 14 hours. The reaction mixture was filtered and the filtrate was dried ($MgSO_4$), concentrated *in vacuo* and the residue was purified by column chromatography (silica, 9:1, dichloromethane: methanol + 1% isopropyl amine) to afford 4.35 g (85.7% yield) of the desired product: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.24 (t, 1H, J=1.9 Hz), 8.09 (dd, 1H, J=1.9, 8.0 Hz), 7.70 (apparent d, 1H, J=8.0 Hz), 7.49 (t, 1H, J=8.0 Hz), 6.23 (m, 1H), 3.29-3.18 (m, 4H), 2.75 (t, 2H, J=5.6 Hz), 2.64-2.54 (m, 4H), 1.82-1.70 (m, 2H), 1.44 (s, 9H); ESMS m/e : 362.2 $(M + H)^+$.

3-(4-(3-NITROPHENYL)-3,6-DIHYDRO-1(2H)-PYRIDINYL)-1-PROPANAMINE: Into a stirred solution of 4.35 (12.0 mmol) of tert-butyl 3-(4-(3-nitrophenyl)-3,6-dihydro-1(2H)-pyridinyl)propylcarbamate in 100 ml of 1,4-dioxane at 0°C was bubbled HCl gas for 10 minutes. The reaction mixture was allowed to warm to room temperature and the

bubbling was continued for an additional 1 hour. The solvent was removed *in vacuo*, the residue was dissolved in 50 mL of water and was neutralized by the addition of KOH pellets. .The aqueous solution was extracted with 3 X 80 mL of dichloromethane, the combined organic extracts were dried ($MgSO_4$), filtered and concentrated *in vacuo*. The residue was purified by column chromatography (silica, 9 : 1 ,dichloromethane : methanol + 1% isopropyl amine) to afford 3.05 g (97.0% yield) of the desired product: ^1H NMR (400 MHz, $CDCl_3$) δ 8.24 (t, 1H, J=1.8 Hz), 8.09 (dd, 1H, J=1.8, 8.2 Hz), 7.69 (dd, 1H, J=1.8, 8.2 Hz), 7.48 (t, 1H, J=8.2 Hz), 6.24 (m, 1H), 3.21 (d, 2H, J=3.6 Hz), 2.84 (t, 2H, J=6.6 Hz), 2.75 (t, 2H, J=5.8 Hz), 2.64-2.54 (m, 4H), 1.76 (m, 2H); ESMS m/e : 262.2 $(M + H)^+$; *Anal. Calc.* for $C_{14}H_{19}N_3O_2$ (0.06 $CHCl_3$): C, 62.90; H, 7.16; N, 15.65. *Found:* C, 63.20; H, 7.16; N, 15.65.

METHYL (4S)-3-[({3-[4-(3-AMINOPHENYL)-1-PIPERIDINYL]PROPYL}AMINO)CARBONYL]-4-(3,4-DIFLUOROPHENYL)-6-(METHOXYMETHYL)-2-OXO-1,2,3,4-TETRAHYDRO-5-PYRIMIDINECARBOXYLATE: A mixture of 3.02 g (6.33 mmol) 5-methyl 1-(4-nitrophenyl) (6*S*)-6-(3,4-difluorophenyl)-4-(methoxymethyl)-2-oxo-3,6-dihydro-1,5(2*H*)-pyrimidinedicarboxylate, 1.50 g (5.80 mmol) of 3-(4-(3-nitrophenyl)-3,6-dihydro-1(2H)-pyridinyl)-1-propanamine, 7.94 g (75.5 mmol) of K_2CO_3 and 1.00 mL of methanol in 200 mL dichloromethane (under argon) was stirred at room temperature for 1 hour. The reaction mixture was filtered and concentrated *in vacuo*. The residue was dissolved in 100 mL of ethyl acetate and washed 3 X 50 mL of 5% aqueous NaOH solution, the organic layer was dried ($MgSO_4$) and concentrated *in*

vacuo. The residue was dissolved in 100 mL of anhydrous ethanol containing 0.50 g 10% Pd/C and the reaction mixture was stirred under a hydrogen balloon for 24 hours. The reaction mixture was passed through a column of Celite 545 filtering agent, washed with ethanol, the filtrate was dried ($MgSO_4$) and concentrated *in vacuo*. The residue was purified by column chromatography (silica, 9.5 : 0.5 ,dichloromethane : methanol + 1% isopropyl amine) to afford 1.65 g (52.0% yield) of the desired product: ^1H NMR (400 MHz, $CDCl_3$) δ 7.80 (s, 1H), 7.22-7.02 (m, 2H), 6.95 (t, J = 8.70 Hz, 1H), 6.63-6.44 (m, 4H), 4.56 (Abq, 2H), 3.62 (s, 3H), 3.33 (s, 3H), 3.32-3.20 (m, 4H), 2.96 (br s, 2H), 2.33 (t, J = 7.50 Hz, 2H), 2.11-1.94 (m, 3H), 1.81-1.64 (m, 4H); ESMS m/e : 572.3 $(M + H)^+$;

TERT-BUTYL 4-[3-(ISOBUTYRYLAMINO)PHENYL]-3,6-DIHYDRO-1(2H)-PYRIDINECARBOXYLATE: Into a solution of 4.00 g (16.0 mmol) of tert-butyl 4-(3-aminophenyl)-3,6-dihydro-1(2H)-pyridinecarboxylate and 5.60 mL (32.0 mmol) of diisopropylethylamine in 100 mL dichloromethane was slowly added 1.90 mL (19.0 mmol) of isobutyryl chloride. The reaction mixture was stirred at room temperature for 2 hours, washed with water, dried ($MgSO_4$), and concentrated *in vacuo*. The residue was purified by column chromatography (silica, 50 : 46 : 3 : 1, hexanes : dichloromethane : methanol : isopropyl amine) to afford 2.90 g (52.0% yield) of the desired product: ^1H NMR (400 MHz, $CDCl_3$) δ 7.69 (s, 1 H), 7.34 (d, 1 H, J=7.8 Hz), 7.27 (t, 1H, J=7.8 Hz), 7.11 (d, 1H, J=7.8 Hz), 6.04 (s, 1H), 4.05 (s, 2H), 3.62 (apparent t, 2 H, J=4.9 Hz), 2.51 (m, 3H), 1.49 (s, 9H), 1.25 (d, 6H, J=7.4 Hz); ESMS *m/e*: 345.5 $(M + H)^+$. Anal. Calc. for

$C_{20}H_{28}N_2O_3+0.175$ $CHCl_3$: C, 66.33; H, 7.77; N, 7.67. Found: C, 66.20; H, 7.41; N, 7.88

***TERT*-BUTYL 4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINECARBOXYLATE**: A mixture of 2.90 g (8.40 mmol) of tert-butyl 4-[3-(isobutyrylamino)phenyl]-3,6-dihydro-1(2H)-pyridinecarboxylate and 0.80 g of 10% yield Pd/C in 100 mL of ethanol was stirred under a hydrogen balloon for 24 hours. The reaction mixture was passed through a column of Celite 545 filtering agent, the filtrate was dried ($MgSO_4$) and concentrated *in vacuo*. The residue was purified by column chromatography (silica, 9.5 : 0.5 ,dichloromethane : methanol + 1% isopropyl amine) to afford 2.40 g (84.0% yield) of the desired product: ^1H NMR (400 MHz, $CDCl_3$) δ 7.49-7.44 (m, 2H), 7.24 (t, 1H, J=7.6 Hz), 6.93 (d, 1H, J=7.6 Hz), 4.20-4.10 (m, 2H), 2.86-2.45 (m, 4H), 1.86-1.75 (m, 4H), 1.48 (s, 9H), 1.24 (d, 6H, J=6.8 Hz); ESMS m/e : 345.2 $(M + H)^+$; Anal. Calc. for $C_{20}H_{30}N_2O_3+0.3H_2O$: C, 68.27; H, 8.77; N, 7.96. Found: C, 68.25; H, 8.54; N, 7.84.

2-METHYL-*N*-[3-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: Into a stirred solution of 2.20 (6.50 mmol) of *tert*-butyl 4-[3-(isobutyrylamino)phenyl]-1-piperidinecarboxylate in 100 ml of 1,4-dioxane at 0 °C was bubbled HCl gas for 10 minutes. The reaction mixture was allowed to warm to room temperature and the bubbling of the HCl gas was continued for 1 hour. The solvent was removed *in vacuo*, the residue was dissolved in 50 mL of water and was neutralized by the addition of KOH pellets. The aqueous solution was extracted with 3 X 80 mL of dichloromethane, the combined organic extracts were dried ($MgSO_4$), filtered and concentrated *in vacuo*. The

residue was purified by column chromatography (silica, 9 : 1 ,dichloromethane : methanol + 1% isopropyl amine) to afford 0.700 g (46.0% yield) of the desired product: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.47 (s, 1H), 7.40 (d, 1H, J=7.8 Hz), 7.24 (t, 1H, J=7.8 Hz), 7.00 (d, 1H, J=7.8 Hz), 3.23-3.14 (m, 5H), 2.82-2.57 (m, 4H), 1.20 (d, 6H, J=6.8 Hz); ESMS m/e : 247.2 $(M + H)^+$; The hydrochloride salt was used for the combustion analysis: Anal. Calc. for $C_{15}H_{22}N_2O$+HCl+0.15 $CHCl_3$: C, 60.51; H, 7.76; N, 9.32. Found: C, 60.57; H, 7.83; N, 8.88.

3-(4-PIPERIDINYL)ANILINE: A solution of 4 M HCl in dioxane (25 mL) was added to *tert*-butyl 4-[3-(amino)phenyl]-1-piperidinecarboxylate (2.60 g, 9.00 mmol) in dichloromethane (250 mL). The reaction mixture was stirred at room temperature overnight, concentrated *in vacuo*, and the residue was dissolved in water (50 mL). The mixture was nuetralized using KOH pellets and extracted with methylene chloride (3 X 50 mL). The combined organic extracts were dried ($MgSO_4$), concentrated and chromatographed to give the desired product (1.03 g). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.01 (t, 1H, J=7.6 Hz), 6.62-6.54 (m, 3H), 3.16 (br d, 2H, J=10.3 Hz), 2.75 (dt, 2H, J=2.7, 12.3 Hz), 2.56 (tt, 1H, J=3.6, 12.3 Hz), 1.81 (br d, 2H, J=12.3 Hz), 1.65 (dq, 2H, J=4.0, 12.3 Hz); ESMS m/e : 177.2 $(M + H)^+$.

***TERT*-BUTYL 4-(4-NITROPHENYL)-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** To a 25-mL RB flask, equipped with a condensor, was added *tert*-butyl 4-{[(trifluoromethyl)sulfonyl]oxy}-3,6-dihydro-1(2*H*)-pyridinecarboxylate (1.0 g), 4-nitrophenylboronic acid (0.71 g), sodium carbonate (0.430 mL of 2M solution),

lithium chloride (0.382 g), tetrakis(triphenylphosphine)- palladium (0) (0.173 g) and ethylene glycol dimethyl ether (10 mL). The reaction mixture was flushed with Argon three times, then the reaction mixture was heated to 100 °C for 3 hrs. After cooling to room temperature, the reaction mixture was diluted with methylene chloride (30 mL) and water (30 mL) and the organic layer was separated. The aqueous layer was extracted with methylene chloride (3x20 mL) and the combined organic extracts were washed with sat NH_4Cl (20 mL) and brine (20 mL), dried over $MgSO_4$ and concentrated under reduced pressure. The residue was purified by chromatography (6:1=hexane:ethyl acetate with 1% NH_3) to afford the product (0.55 g, 59.9%) as a yellow oil. The compound is not stable at room temperature and should be used as promptly as practical: ^1H NMR (400 MHz, $CDCl_3$) δ 8.20 (d, 2H, J=8.6 Hz), 7.51 (d, 2H, J=8.6 Hz), 6.24 (m, 1H), 4.13 (m, 2H), 3.67 (apparent t, 2H, J=5.5 Hz), 2.55 (m, 2H), 1.49 (s, 9H).

4-(4-NITROPHENYL)-1,2,3,6-TETRAHYDROPYRIDINE: 4-(4-Nitrophenyl)-1,2,3,6-tetrahydropyridine was prepared by a similar procedure to that used for the preparation of 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide using HCl gas and *tert*-Butyl 4-(4-Nitrophenyl)-3,6-dihydro-1(2*H*)-pyridinecarboxylate (130 mg) in dioxane (5.0 mL) at room temperature. The reaction mixture was concentrated *in vacuo* to give the crude product (69.8 mg) which used in the next reaction without further purification.

Oxazolidinone Intermediates:

AMINO-(3,4-DIFLUOROPHENYL)-ACETONITRILE: Through a solution of 3,4-difluorobenzaldehyde (25.0 g, 0.176 mol) in MeOH (500 mL) in a round bottom flask, was bubbled ammonia gas for two hours at room temperature. The flask was then cooled to 0 ^{0}C and trimethylsilyl cyanide was then added slowly. The reaction mixture was stirred for 2 h, at which time TLC analysis indicated that the reaction was complete (R_f = 0.35, 3:2 hexane/EtOAc). The solvent was removed *in vacuo* and the residue was subjected to flash column chromatography on silica gel to obtain the desired product, which was used in the next step without purification.

AMINO-(3,4-DIFLUOROPHENYL)-ACETIC ACID METHYL ESTER: Into a well-stirred solution of amino-(3,4-difluorophenyl)-acetonitrile (22.0 g, 0.130 mol), a solution of HCl in MeOH (200 mL) was added at room temperature. The resulting yellow solution was stirred at room temperature for 10 h and was heated at reflux temperature for 1.5 h. After cooling, the solvent was removed *in vacuo* and the resulting yellow solid was dissolved in water (200 mL). The aqueous solution was then carefully basified with 20% NaOH solution to pH 9. The aqueous layer was extracted with CH_2Cl_2 (3 x 100 mL). The organic layer was separated and dried over Na_2SO_4, filtered and the solvent was removed *in vacuo* to obtain the desired product which was used in the next step without purification.

2-AMINO-2-(3,4-DIFLUOROPHENYL)-ETHANOL: Into a well-stirred suspension of $LiAlH_4$ (4.7 g, 0.125 mol) in THF (120 mL) in a 3-necked round bottom flask fitted with a

condenser and a dropping funnel, was added a solution of amino-(3,4-difluorophenyl)-acetic acid methyl ester (10.0 g, 0.05 mol) in THF (100 mL) dropwise at 0 ^{0}C. The resulting greenish brown suspension was heated at reflux temperature for 2 h. The reaction mixture was cooled to 0 ^{0}C and then carefully quenched sequentially with 5 mL of water, 5 mL of 3N NaOH followed by 15 mL of water. The resulting suspension was filtered through a fritted glass funnel. To the filter cake was added 100 mL Et_2O and the suspension was heated at reflux temperature for 20 min. The suspension was filtered and the combined filtrates were dried over $MgSO_4$, filtered and the solvent was removed *in vacuo*. 2-Amino-2-(3,4-difluorophenyl)-ethanol was obtained as a yellow glassy syrup which was used in the next step without further purification.

[1-(3,4-DIFLUOROPHENYL)-2-HYDROXY-ETHYL]-CARBAMIC ACID-*TERT*-BUTYL ESTER: Into a solution of 2-amino-2-(3,4-difluorophenyl)-ethanol (8.6 g, 49.7 mmol) in $CHCl_3$ (150 mL) at 0 ^{0}C was added a solution of *di-tert*-butyl dicarbonate (11.4 g, 52.0 mmol) in $CHCl_3$ (50 mL) in one portion and the resulting solution was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue was subjected to column chromatography on silica gel (2:1 hexane-EtOAc followed by EtOAc) to obtain [1-(3,4-difluorophenyl)-2-hydroxy-ethyl]-carbamic acid-*tert*-butyl ester as a white solid (10.0 g, 74% yield).

(+)-4-(3,4-DIFLUOROPHENYL)-OXAZOLIDIN-2-ONE: Into a well-stirred suspension of NaH (1.1 g, 45.8 mmol) in THF (40 mL) at R.T. was added a solution of [1-(3,5-

difluorophenyl)-2-hydroxy- ethyl]-carbamic acid-*tert*-butyl ester (5.0 g, 18.3 mmol) in THF (20 mL) via a dropping funnel at room temperature. The resulting suspension was stirred for 3 h and then quenched carefully with 10 mL of water. The biphasic mixture was extracted with 100 mL of Et_2O, washed with brine, filtered and the solvent was removed *in vacuo*. The gummy residue thus obtained was purified by column chromatography over silica gel (R_f = 0.15, 3:2 hexane-EtOAc) to obtain 4-(3,5-difluorophenyl)-oxazolidin-2-one as a white flaky solid (2.8 g, 77% yield). M.P. 81-83 ^{0}C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.23-7.03 (m, 3H), 6.08 (br s, 1H), 4.94 (dd, J=6.6 Hz, J=8.7 Hz, 1 H), 4.73 (t, J=8.7 Hz, 1 H), 4.13 (dd, J=6.6 Hz, J=8.7 Hz, 1 H). The enantiomers were separated by HPLC on a Chiralcel OD (20 x 250 mm) column using 80% hexane/20% isopropyl alcohol as the eluting system at 12.0 mL/min (U.V. 254 nm). The retention times for the two isomers were 16.19 min and 20.08 min respectively.

4-NITROPHENYL (4S)-4-(3,4-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXYLATE: Into a suspension of NaH (0.14 g, 5.30 mmol) in 20 mL of anhydrous THF under argon, a solution of (+)-4-(3,4-difluorophenyl)-oxazolidin-2-one (0.88 g, 4.42 mmol) in THF was added dropwise (dropping funnel). The resulting suspension was stirred at room temperature for 30 min. This suspension was then added dropwise *via* cannula into another round bottom flask containing a solution of 4-nitrophenylchloroformate (1.11 g, 5.30 mmol) in 25 mL of THF and cooled at -78 ^{0}C over a period of 15 min. The stirring was continued for 2 h after which the solvent was removed and the residue was purified by column

chromatography on silica gel with 1:1 hexane/CH_2Cl_2 followed by CH_2Cl_2 (R_f= 0.4, CH_2Cl_2) to obtain the desired product as a white solid (1.55 g, 86% yield).

Similarly, following the above procedure, 4-(3,5-difluorophenyl)-2-oxo-oxazolidine-3-carboxylic acid-4-nitro-phenyl ester and 4-(3,4,5-trifluorophenyl)-2-oxo-oxazolidine-3-carboxylic acid-4-nitro-phenyl ester were obtained by substituting 3,4-diflourobenzaldehyde in the first step with 3,5-diflourobenzaldehyde or 3,4,5-triflourobenzaldehyde, respectively. The oxazolidinone enantiomers were resolved by HPLC on a Chiralcel OD column (as in the previous example) and the 4-nitro-phenyl carbamates were prepared using 4-nitrophenyl chloroformate.

4-NITROPHENYL (4S)-4-(3,5-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXYLATE: Following the procedure for the synthesis of 4-(3,4-difluorophenyl)-2-oxo-oxazolidine-3-carboxylic acid-4-nitro-phenyl ester, 3,5-diflourobenzaldehyde yielded the desired product.

^1H NMR (400 MHz, $CDCl_3$) δ 8.26 (d, 2H, J= 9.3 Hz), 7.33 - 6.81 (m, 5H), 5.41 (dd, 1H, J=4.1, 8.7 Hz), 4.81 (t, 1H, J=9.3 Hz), 4.33 (dd, 1H, J=4.1, 9.3 Hz); Anal. Calc. for $C_{16}H_{10}F_2N_2O_6$+0.2EtOAc: C, 52.84; H, 3.06; N, 7.34. Found: C, 53.26; H, 2.83; N, 7.73

4-NITROPHENYL (4S)-2-OXO-4-(3,4,5-TRIFLUOROPHENYL)-1,3-OXAZOLIDINE-3-CARBOXYLATE: Following the procedure for the synthesis of 4-(3,4-difluorophenyl)-2-oxo-oxazolidine-3-carboxylic acid-4-nitro-phenyl ester, 3,4,5-triflourobenzaldehyde yielded the desired product.

^{1}H NMR (400 MHz, $CDCl_3$) δ 8.27 (d, 2H, J=9.0 Hz), 7.31 (d, 2H, J=9.0 Hz), 7.11-7.02 (m, 2H), 5.37 (dd, 1H, J=4.1, 9.0 Hz), 4.81 (apparent t, 1H, J=9.0 Hz), 4.33 (dd, 1H, J=4.1, 9.0 Hz); Anal. Calc. for $C_{16}H_9F_3N_2O_6$: C, 50.27; H, 2.37; N, 7.33. Found: C, 50.56; H, 2.50; N, 7.49.

1-(3,4-DIFLUOROPHENYL)-2-METHYL-2-HYDROXYPROPYLAMINE:

Into a well-stirred solution of methyl 2-amino-2-(3,4-difluorophenyl)acetate (10.5 g, 52.19 mmol) in anhydrous ether (200 mL) at 0 °C a solution of methylmagnesium bromide (3 M, 87 mL, 261 mmol) in ether was added over 10 minutes. The reaction mixture was stirred at 0 °C for 2.5 h and allowed to warm to room temperature. After 12 h, the reaction mixture was carefully poured onto a mixture of ice (300 g) and saturated aqueous ammonium chloride (50 g). The ether layer was separated and the aqueous layer was extracted with more ether (4 X 200 mL). The combined extracts were dried with magnesium sulfate and the solvent evaporated. The crude product was purified by column chromatography on silica gel using chloroform/methanol/2M ammonia in methanol (1000:20:10, 1000:40:20, 1000:80:40) as the eluent to give the product as an oil (6.5 g, 62% yield) which was used in the next step without further purification.

4-(3,4-DIFLUOROPHENYL)-5,5-DIMETHYL-2-OXO-OXAZOLIDINE: A mixture of 1-(3,4-difluorophenyl)-2-methyl-2-hydroxypropylamine (3.00 g, 14.9 mmol) and carbonyldiimidazole (2.418 g, 14.9 mmol) in dichloromethane (150 mL) was heated at reflux temperature for 36 h and the solvent evaporated. The residue was purified by column chromatography on silica

gel using chloroform/ethyl acetate (9:1) to give the product as a viscous oil which solidified on standing (1.80 g, 50% yield). The product was used in the next step without further characterization.

4-NITROPHENYL 4-(3,4-DIFLUOROPHENYL)-5,5-DIMETHYL-2-OXO-1,3-OXAZOLIDINE-3-CARBOXYLATE: Into a stirred suspension of sodium hydride (60% suspension in paraffin 203 mg, 1.4 eq.) in THF (20 mL) at 0 °C, a solution of 4-(3,4-difluorophenyl)-5,5-dimethyl-2-oxo-oxazolidine (870 mg, 3.622 mmol) in THF (5 mL) was added followed by stirring for 30 minutes. This suspension was added to a solution of 4-nitrophenyl chloroformate (950 mg, 4.71 mmol) in THF (20 mL) at -78 °C under argon and the stirring was continued for 2 h. It was slowly warmed to room temperature and after 4 h the solvent was evaporated. The residue was mixed with dichloromethane (150 mL), washed with 0.05 N sodium hydroxide (3 X 10 mL), and dried (sodium sulfate). The solvent was evaporated and the residue was purified by column chromatography on silica gel using chloroform/ethyl acetate (9:1) as the eluent to give the product as a white powder (860 mg, 59% yield).

^{1}H NMR (400 MHz, $CDCl_3$) δ 8.24 (d, 2H, J=9 Hz), 7.29 - 6.97 (m, 5H), 5.04 (s, 1H), 1.09 (s, 6H); Anal. Calc. for $C_{18}H_{14}F_2N_2O_6$+0.2% H_2O: C, 54.61; H, 3.67; N, 7.08. Found: C, 54.89; H, 3.59; N, 7.41.

(3,4-DIFLOUROPHENYL)-N(DIPHENYLMETHYLENE)METHANAMINE: Into a solution of 3,4-difluorobenzylamine (9.8 g, 69 mmol) and benzophenone (13.0 g, 71.0 mmol) in toluene (200 mL) was added a catalytic amount of $BF_3.OEt_2$ and the resulting solution was heated at reflux temperature for

12 h. The reaction mixture was concentrated *in vacuo*, yielding an oil (21 g, >95%), which was characterized by NMR analysis and subjected to the following reaction without any further purification. ^1H NMR ($CDCl_3$) δ 4.57 (s, 2H), 7.80-6.80 (m, 13H).

1-(3,4-DIFLOUROPHENYL)-1-[(DIPHENYLMETHYLENE)AMINO]PROPAN-2-OL: Into a solution of the benzhydrylindene-(3,4-difluoro-benzyl)-amine (21 g, 69 mmol) in 250 ml of dry THF was added *tert*-butyllithium (1.7 M, 60 ml) dropwise and the resulting solution was stirred at -78 °C for 0.5 h. To the solution was added acetaldehyde (10 ml, 180 mmol) in 100 ml of THF and the solution was stirred at -78 °C for 2 h and 25 °C for 1 h. The reaction mixture was quenched by addition of brine. The reaction mixture was diluted with 500 ml of Et_2O and washed with brine. The organic layer was dried over Na_2SO_4 and concentrated *in vacuo* to give an oil, which was taken to the next step without any further purification. ^1H NMR ($CDCl_3$) δ 1.04 (d, 3H), 2.77 (broad s. 1H), 4.08(m, 1H), 4.15 (d, 1H), 7.80-6.80 (m, 13H).

1-AMINO-1-(3,4-DIFLUORO-PHENYL)-PROPAN-2-OL: A solution of crude product from the previous procedure and $MeONH_2.HCl$ (10 g, 120 mmol) was diluted in 200 ml of MeOH and stirred for 12 h. The reaction mixture was concentrated *in vacuo*, yielding an oily residue, which was re-dissolved in 200 ml of EtOAc and washed with brine. The organic layer was concentrated *in vacuo* to produce an oily mixture, which was subjected to column chromatography [5% NH_3 (2.0 M in MeOH) in $CHCl_3$] to yield the desired product (8.8 g, 68% yield from 3,4-

difluorobenzylamine) as a mixture of diastereomers. ^{1}H NMR ($CDCl_3$) (~ 4:1 mixture of the diastereomers) δ 1.02 (d, J=6.0 Hz, 3 H), 1.04 (d, J=6.3 Hz, 3 H), 2.10 (br, 6 H), 3.56-3.69 (m, 2 H), 3.88-3.92 (m, 2 H), 7.02-7.17 (m, 6 H).

[1-(3,4-DIFLUOROPHENYL)-2-HYDROXY-PROPYL]-CARBAMIC ACID-*TERT*-BUTYL ESTER: Into a solution of 1-amino-1-(3,4-difluorophenyl)-propan-2-ol (13.1 g, 70.1 mmol) in $CHCl_3$ (150 mL) at 0 ^{0}C was added a solution of *di-tert*-butyl dicarbonate (19.3 g, 87.6 mmol) in $CHCl_3$ (50 mL) in one portion and the resulting solution was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue was subjected to column chromatography on silica gel (2:1 hexane-EtOAc followed by EtOAc) to obtain [1-(3,4-difluorophenyl)-2-hydroxy-propyl]-carbamic acid-*tert*-butyl ester as a viscous oil (18.4 g, 91% yield). ^{1}H NMR ($CDCl_3$) (~ 4:1 mixture of the diastereomers) δ 1.05 (d, J=6.6 Hz, 3 H), 1.25 (d, J=6.0 Hz, 3 H), 1.41 (br, 20 H), 3.92-4.19 (br, 2 H), 4.45-4.60 (m, 2 H), 5.41-5.49 (br, 2 H), 7.02-7.17 (m, 6 H).

4-(3,4-DIFLUOROPHENYL)-5-METHYL-OXAZOLIDIN-2-ONE: Into a well-stirred solution of [1-(3,4-difluorophenyl)-2-hydroxy-propyl]-carbamic acid-*tert*-butyl ester (0.43 g, 1.5 mmol) in THF (20 mL) was added 95% NaH (0.09 g, 3.8 mmol) at room temperature. When the reaction was carried out on a larger (> 5 g) scale, 1.0 equivalent of KH and 1.5 eq. of NaH was used as the base. The resulting suspension was stirred for 3 h at about 35 ^{0}C (warm water bath) and then quenched carefully with ice. The biphasic mixture was extracted with 100 mL of EtOAc, washed with brine, dried over Na_2SO_4, filtered and the

solvent was removed *in vacuo*. The two diastereomers were separated by column chromatography over silica gel (First isomer: 0.16 g, R_f = 0.6, 3:1 hexane-EtOAc; second isomer: 0.18 g, R_f = 0.5, 3:1 hexane-EtOAc). NOE experiments suggested that the first diastereomer had the methyl and the aryl group in trans configuration while the second diastereomer had cis relationship between the two groups. The ^{1}H NMR spectrum for the *trans* diastereomer is as follows. ^{1}H NMR ($CDCl_3$) δ 1.49 (d, J = 6.0 Hz, 3H), 4.37 (dq, J = 6.0 Hz, J = 7.2 Hz, 1H), 4.45 (d, J = 7.2 Hz, 1H), 6.63 (br s, 1H), 7.08-7.28 (m, 3H).

The ^{1}H NMR spectrum for the *cis* diastereomer is as follows. ^{1}H NMR ($CDCl_3$) δ 0.96 (d, J = 6.6 Hz, 3H), 4.91 (d, J = 8.1 Hz, 1H), 4.99 (dq, J = 6.6 Hz, J = 8.1 Hz, 1H), 6.63 (br s, 1H), 7.08-7.28 (m, 3H).

4-(3,4-DIFLUOROPHENYL)-5-METHYL-2-OXO-OXAZOLIDINE-3-CARBOXYLIC ACID-4-NITRO-PHENYL ESTER : Into a solution of one of the two diastereomers of 4-(3,4-difluorophenyl)-5-methyl-oxazolidin-2-one (0.97 g, 4.55 mmol) in 60 mL THF was added a solution of n-butyllithium in hexane (3.06 mmol, 4.9 mmol) dropwise via a syringe under argon atmosphere at -78 ^{0}C. The resulting yellow solution was stirred at -78 ^{0}C for 40 min. This solution was then added dropwise via a cannula into another round bottom flask containing a solution of 4-nitrophenylchloroformate (1.03 g, 5.1 mmol) in 60 mL of THF, cooled at -78 ^{0}C, over a period of 15 min. After five minutes, the flask was removed from the cooling bath and stirring was continued for 1 h. The reaction mixture was quenched by adding ice and it was

extracted with EtOAc. The organic extracts were washed with brine and the organic layer was dried over Na_2SO_4. The solvent was removed after filtration and the residue was purified by column chromatography on silica gel with 1:1 hexane/CH_2Cl_2 followed by CH_2Cl_2 to give the desired product.

The relative configurations of the cis and trans isomers were assigned on the basis of ^1H NMR analysis of the respective p-nitrophenyloxycarbonyl derivatives. For the trans isomer, an NOE was observed between the protons of the C-5 methyl group and the proton at C-4. No NOE was observed between the protons at the C-4 and C-5 positions of this isomer, which was thus assigned *trans* stereochemistry. For the cis isomer, no NOE was observed between the protons of the C-5 methyl group and the proton at C-4. However, a NOE was observed between the protons at the C-4 and C-5 positions, leading us to assign this isomer *cis* stereochemistry. The vicinal coupling constants of the C-4 protons of cis (J = 7.8 Hz) and trans (J = 5.1 Hz) are also consistent with the values reported for similar oxazolidinones, and were thus helpful in making the stereochemical assignments (Dondoni, A.; Perrone, D.; Semola, T. *Synthesis 1995*, 181).

Enantiomers of the diastereomers were separated by HPLC by using a Chiralcel OD column (20 x 250 mm) with 80% hexane/20% isopropyl alcohol/ 0.1 % diethylamine as the eluting system (12 mL/min) under isocratic conditions (U.V. 254 nm).

In order to assign the absolute configurations at the

stereogenic centers of the oxazolidinone rings, a new synthetic route was designed which employed an enantiomerically pure substrate derived from the chiral pool. Commercially available (*S*)-(+)-methyl lactate was converted into its pyrrolidine amide according to the method of *Martin et al* (Martin, R.; Pascual, O.; Romea, P.; Rovira, R.; Urpi, F.; Vilarrasa, J. *Tetrahedron Lett. 1997, 38,* 1633). Following the protection of the hydroxy group of (2*S*)-1-oxo-1-(1-pyrrolidinyl)-2-propanol to a TBDMS group, treatment of *tert*-butyl(dimethyl)silyl (1*S*)-1-methyl-2-oxo-2-(1-pyrrolidinyl)ethyl ether with 3,4-difluorophenyllithium yielded (2*S*)-2-{[*tert*-butyl(dimethyl)silyl]oxy}-1-(3,4-difluorophenyl)-1-propanone as the sole product, which was then converted to (2*S*)-2-{[*tert*-butyl(dimethyl)silyl]oxy}-1-(3,4-difluorophenyl)-1-propanone oxime. Reduction of the (2*S*)-2-{[*tert*-butyl(dimethyl)silyl]oxy}-1-(3,4-difluorophenyl)-1-propanone oxime with $LiAlH_4$, N-acylation, and base induced cyclization provided oxazolidinone diastereomers, which were separated by flash column chromatography. The enantiomeric purity of these isomers was confirmed by chiral HPLC analysis and their relative configurations were assigned by comparison of their ^1H NMR spectra with those of the racemic isomers. As the absolute configuration at C-5 of the lactic acid derived oxazolidinone described above is (*S*), the C-4 center in *trans* compounds also has the (*S*) configuration. Accordingly, the absolute configurations for the stereogenic centers in the *cis* compounds are assigned accordingly (*4R,5S*).

4-NITROPHENYL (4S,5R)-4-(3,4-DIFLUOROPHENYL)-5-METHYL-2-OXO-1,3-OXAZOLIDINE-3-CARBOXYLATE: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.25 (d, 2H, J=8.8 Hz), 7.30 - 6.99 (m, 5H), 5.35 (d, 1H, J=7.7 Hz), 5.07 (apparent quintet, 1H), 1.17 (d, 3H, J=6.5 Hz); Anal. Calc. for $C_{17}H_{12}F_2N_2O_6$+0.5H_2O: C, 52.72; H, 3.38; N, 7.23. Found: C, 53.09; H, 3.19; N, 7.50.

(+)-2-AMINO-3-(3,4-DIFLUORO)-PHENYL-PROPAN-1-OL: (+)-3,4-difluorophenyl alanine (1.0 g, 5.0 mmol) was added in small portions to a stirring suspension of $LiAlH_4$ (0.480 g, 12.5 mmol) in THF (30 mL) at 0 ^{0}C. The resulting gray suspension was then heated at reflux for 2 h. The reaction mixture was cooled to 0 ^{0}C and then carefully quenched sequentially with water (0.5 mL), 3 N NaOH (0.5 mL), and water (1.50 mL). The resulting suspension was filtered through a fritted glass funnel. Ether (50 mL) was added to the filter cake and the suspension was heated at reflux temperature for 20 min. The suspension was filtered and was combined with the previous filtrate. The combined organics were dried over $MgSO_4$, filtered and the solvent was removed *in vacuo*. 2-Amino-3-(3,4-difluoro)-phenyl-propan-1-ol was obtained as a white solid (0.500 g, 100%) which was used in the next step without further purification.

(+)-[1-(3,4-DIFLUOROBENZYL)-2-HYDROXY-ETHYL]-CARBAMIC ACID-*TERT*-BUTYL ESTER: A solution of di-*tert*-butyl dicarbonate (0.640 g, 2.90 mmol) in $CHCl_3$ (10 mL) was added in one portion to a solution of (+)-2-amino-3-(3,4-difluoro)-phenyl-propan-1-ol (0.500 g, 2.62 mmol) in $CHCl_3$ (20 mL) at 0 ^{0}C and the resulting solution was stirred overnight at room temperature. The solvent was

removed *in vacuo* and the residue was chromatographed (2:1 hexane-EtOAc, followed by EtOAc), giving (+)-[1-(3,4-difluorobenzyl)-2-hydroxy-ethyl]-carbamic acid-*tert*-butyl ester as a white solid (0.640 g, 99%).

(+)-4-(3,4-DIFLUORO-BENZYL)-OXAZOLIDIN-2-ONE: A solution of (+)-[1-(3,4-difluorobenzyl)-2-hydroxy-ethyl]-carbamic acid-*tert*-butyl ester (1.00 g, 4.00 mmol) in THF (10 mL) was added *via* a dropping funnel to a stirring suspension of 95% NaH (0.12 g, 5.0 mmol) in THF (20 mL) at room temperature. The resulting suspension was stirred for 3 h and then quenched carefully with water (10 mL). The biphasic mixture was extracted with Et_2O (50 mL), washed with brine, filtered and the solvent was removed *in vacuo*. The resulting gummy residue was purified by column chromatography (R_f = 0.25, 3:2 hexane-EtOAc), to give the desired product as a white solid (0.320 g, 76%).

(+)-4-(3,4-DIFLUORO-BENZYL)-OXAZOLIDIN-2-ONE-3-CARBOXYLIC ACID-4-NITRO-PHENYL ESTER: A solution of (+)-4-(3,4-difluoro-benzyl)-oxazolidin-2-one (0.210 g, 1.0 mmol) in THF (10 mL) was added dropwise *via* a dropping funnel to a stirring suspension of NaH (30.0 mg, 1.30 mmol) in anhydrous THF (10 mL) under argon. The resulting suspension was stirred at room temperature for 30 min. This suspension was then added dropwise *via* cannula to a solution of 4-nitrophenylchloroformate (0.300 g, 1.50 mmol) in THF (20 mL) at -78 ^{0}C over 15 min. Stirring was continued for 2 h after which the solvent was removed and the residue was purified by column chromatography (1:1 hexane/CH_2Cl_2, followed by

CH_2Cl_2; R_f= 0.4, CH_2Cl_2), to give the desired product as a yellow solid (0.350 g, 82%).

Similarly, following the above procedure, 4-nitrophenyl 4-(4-fluorobenzyl)-2-oxo-1,3-oxazolidine-3-carboxylate was obtained by substituting (+)-3,4-diflourophenyl alanine with *p*-fluorophenyl alanine:

4-NITROPHENYL 4-(4-FLUOROBENZYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXYLATE: ^1H NMR (400 MHz, $CDCl_3$) δ 8.32 (d, 2H, J=9.3 Hz), 7.42 (d, 2H, J=8.9 Hz), 7.24-6.99 (m, 4H), 4.69 - 4.59 (m, 1H), 4.35 (t, 1H, J=8.6 Hz), 4.23 (dd, 1H, J=2.7, 9.3 Hz), 3.37 (dd, 1H, J=3.8, 13.6 Hz), 2.94 (dd, 1H, J=9.3, 13.6 Hz); Anal. Calc. for $C_{17}H_{13}FN_2O_6$: C, 56.67; H, 3.64; N, 7.77. Found: C, 56.94; H, 3.76; N, 7.71.

2-[6-(4-PHENYL-1-PIPERIDINYL)HEXYL]-1H-ISOINDOLE-1,3(2H)-DIONE: To the 500 ml RB-flask was added 4-phenylpiperidine hydrochloride (5 g, 25 mmol), N-(6-bromohexyl)phthalimide (15.5 g, 50 mmol), N,N-diisopropylethylamine (21.8 ml, 125 mmol), tetrabutylammonium iodide (0.2 g), and dioxane (250 ml) at room temperature. The reaction mixture was stirred at 100 °C for 72 h. The solvent was removed *in vacuo* and the crude product was purified by flash chromatography (98:2 = Chloroform : 2N ammonia in methanol) to afford 7.67 g of the desired product (77% yield): ^1H NMR (400 MHz, $CDCl_3$) δ 7.78-7.79 (m, 2H), 7.74-7.65 (m, 2H), 7.32-7.14 (m, 5H), 3.69 (t, 2H, J=7.35 Hz), 3.06 (d, 2H, J=11.0 Hz), 2.49 (quintet, 1H, J=7.6 Hz), 2.36 (t, 2H, J=7.6 Hz), 2.02 (t, 2H, J=12.5 Hz), 1.82 (br s, 4H), 1.69 (t, 2H, J=6.3 Hz), 1.54 (br s, 2H), 1.37 (br s,

4H); ESMS m/e: 391.3 (M + H)$^+$; Anal. Calc. for $C_{25}H_{30}N_2O_2$+0.2H_2O: C, 76.19; H, 7.77; N, 7.11. Found: C, 76.14; H, 7.38; N, 7.13.

METHOD I. General procedure for the Preparation of the substituted 4-[4-(3-aminophenyl)-1-piperidinyl]-1-(phenyl)-1-butanones: A mixture of 4-(3-aminophenyl)piperidine (2.0 mmol), 2.4 mmol of the appropriate substituted phenyl butyryl chloride (e.g. 4-chloro-4'-phenoxybutyrophenone, 4-chloro-3',4'-dimethylbutyrophenone, 4-chloro-4'-chlorobutyrophenone, γ-chlorobutyrophenone, 4-chloro-3',4'-dimethoxybutyrophenone), 3.0 mmol of K_2CO_3, and 10 mg of 18-crown-6 in 5 mL of toluene were heated at 110 °C for 2.5 days. The reaction mixture was concentrated and chromatographed on silica (5% methanol in dichloromethane) to give the desired compound:

4-[4-(3-AMINOPHENYL)-1-PIPERIDINYL]-1-(4-PHENOXYPHENYL)-1-BUTANONE: Using Method I, the desired product was obtained. 305 mg; ESMS m/e : 415.4 (M + H)$^+$.

4-[4-(3-AMINOPHENYL)-1-PIPERIDINYL]-1-(3,4-DIMETHYLPHENYL)-1-BUTANONE: Using Method I, the desired product was obtained. 320 mg; ESMS m/e : 351.3 (M + H)$^+$.

4-[4-(3-AMINOPHENYL)-1-PIPERIDINYL]-1-(4-CHLOROPHENYL)-1-BUTANONE: Using Method I, the desired product was obtained. 500 mg; Anal. Calc for $C_{21}H_{25}ClN_2O$+0.3H_2O: C, 69.62; H, 7.12; N, 7.73. Found: C, 69.63; H, 7.34; N, 7.60; ESMS m/e : 357.3 (M + H)$^+$.

4-[4-(3-AMINOPHENYL)-1-PIPERIDINYL]-1-PHENYL-1-BUTANONE: Using Method I, the desired product was obtained. 250 mg; Anal. Calc for $C_{21}H_{26}N_2O+0.2H_2O$: C, 77.36; H, 8.16; N, 8.59. Found: C, 77.55; H, 8.12; N, 8.75; ESMS m/e : 323.3 $(M + H)^+$.

4-[4-(3-AMINOPHENYL)-1-PIPERIDINYL]-1-(2,4-DIMETHOXYPHENYL)-1-BUTANONE: Using Method I, the desired product was obtained. 330 mg; Anal. Calc for $C_{23}H_{30}N_2O_3+0.5H_2O$: C, 70.56; H, 7.98; N, 7.16. Found: C, 70.69; H, 7.87; N, 6.99; ESMS m/e : 383.3 $(M + H)^+$.

METHOD II. General Procedure for the Acylation or Sulfonylation of the Substituted 4-[4-(3-Aminophenyl)-1-piperidinyl]-1-(4-phenyl)-1-butanones: A mixture of 1 equivalent of a substituted 4-[4-(3-aminophenyl)-1-piperidinyl]-1-(4-phenyl)-1-butanone, 1.5 equivalent of an acid chloride or a sulfonyl chloride, and 5 equivalents of diisopropylethylamine, in dichloromethane was stirred at room temperature for two days. The reaction mixture was applied to a preparative TLC plate and eluted with dichloromethane: methanol (15:1, containing 1% isopropyl amine) to give the desired product.

METHOD III. General procedure for the Preparation of the substituted 4-N-(3-{1-[4-(phenyl)-4-oxobutyl]-4-piperidinyl}phenyl)acetamides: A mixture of N-[3-(4-piperidinyl)phenyl]acetamide (1.0 eq) and an aryl substituted chlorobutyrophenone (2.0 eq), K_2CO_3 (5.0 eq), diisopropylethylamine (3.0 eq) and tetrabutylammonium iodide (cat. 5-10%) in dioxane (0.5 to 1.0 M) were heated at reflux temperature for 16 h. The reaction

mixture was filtered and concentrated *in vacuo*. The crude product was chromatographed using silica preparative TLC (chloroform : methanol containing 0.5% isopropyl amine) to give the desired product.

Example 1

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^{1}H NMR ($CDCl_3$) δ 7.75 (s, 1H), 7.71 (d, 1H, J=7.6 Hz), 7.45 (d, 2H, J=7.2 Hz), 7.35 (s, 1H), 7.26-7.22 (m, 2H), 6.93 (d, 1H, J=7.6 Hz), 3.24-3.21 (m, 2H), 3.04 (t, 2H, J=7.0 Hz), 2.67-2.63 (m, 2H), 2.59-2.48 (m, 1H), 2.32 (s, 6H), 2.30-2.27 (m, 2H), 2.18 (s, 3H), 2.14-2.06 (m, 2H), 2.00-1.80 (m, 4H); ESMS m/e : 393.3 $(M + H)^+$.

Example 2

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of 0.0500 g (0.200 mmol) of 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide, 0.100 g (0.480 mmol) of 4-chloro-3',4'-dimethylbutyrophenone, 0.080 g (0.600 mmol) of K_2CO_3 and 0.090 g (0.600 mmol) of NaI in 5 mL of DMF was heated at reflux temperature for 18 hours. The reaction mixture was filtered, the filtrate was poured into 5 mL of water and washed with 3 X 5 mL of ethyl acetate. The combined organic extracts were dried ($MgSO_4$), concentrated *in vacuo* and purified by preparative TLC (silica; 9.5 : 0.5, dichloromethane : methanol + 1% isopropyl amine) to afford 0.067 g (80.0% yield) of the desired product: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.72 (d, 1H, J=8.0 Hz), 7.44 (s, 1H), 7.38 (d, 1H, J=8.0 Hz), 7.23-7.20 (m, 2H), 7.16 (s, 1H), 6.95 (d, 1H, J=6.8

Hz), 3.13-3.11 (m, 2H), 3.02 (t, 2H, J=7.0 Hz), 2.56-2.40 (m, 4H), 2.32 (s, 6H), 2.17-2.15 (m, 2H), 2.04-1.78 (m, 6H), 1.25 (d, 6H, J=6.8 Hz); ESMS m/e : 421.3 $(M + H)^+$.

Example 3

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOHEXANECARBOXAMIDE: Using Method II, the desired compound was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.80-6.81 (m, 7H), 3.41-3.00 (m, 4H), 2.95-2.41 (m, 4H), 2.32 (s, 6H), 2.22-1.05 (m, 18H); ESMS m/e : 461.4 $(M + H)^+$.

Example 4

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-PHENYLACETAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.85-7.65 (m, 2H), 7.45-6.92 (m, 10H), 3.76 (s, 2H), 3.10-2.90 (m, 4H), 2.50-2.35 (m, 3H), 2.32 (s, 6H), 2.10-1.85 (m, 4H), 1.80-1.60 (m, 4H); ESMS m/e : 469.4 $(M + H)^+$.

Example 5

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-(3-METHOXYPHENYL)ACETAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.76-7.65 (m, 2H), 7.38-7.12 (m, 6H), 6.95-6.80 (m, 3H), 3.82 (s, 3H), 3.70 (s, 2H), 3.10-2.90 (m, 4H), 2.50-2.38 (m, 3H), 2.32 (s, 6H), 2.10-1.85 (m, 4H), 1.80 -1.60 (m, 4H); ESMS m/e : 499.4 $(M + H)^+$.

Example 6

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-METHOXYACETAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.80-7.75 (m, 2H), 7.50-7.38 (m, 2H), 7.34-6.90 (m, 3H), 4.00 (s, 2H), 3.51 (s, 3H), 3.30-2.95 (m, 4H), 2.70-2.50 (m, 3H), 2.32 (s, 6H), 2.15 -1.80 (m, 8H); ESMS m/e : 423.3 $(M + H)^+$.

Example 7

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)METHANESULFONAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.82-7.10 (m, 7H), 3.41 (s, 3H), 3.40-2.85 (m, 4H), 2.82-2.35 (m, 5H), 2.32 (s, 6H), 2.22-1.80 (m, 6H); ESMS m/e : 429.3 $(M + H)^+$.

Example 8

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ETHANESULFONAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.75 (s, 1H), 7.71 (d, 1H, J=7.6 Hz), 7.30-7.09 (m, 4H), 7.02 (d, 1H, J=7.2 Hz), 3.36-3.05 (m, 6H), 2.77-2.52 (m, 3H), 2.32 (s, 6H), 2.15-1.82 (m, 8H), 1.37 (t, 3H, J=7.4 Hz); ESMS m/e : 443.3 $(M + H)^+$

Example 9

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (d, 2H, J=8.8 Hz), 7.55-7.40 (m, 3H), 7.35 (s, 1H), 7.22 (t, 1H, J=8.0 Hz), 6.92 (d, 1H, J=8.0 Hz), 3.30-3.27 (m, 2H), 3.09 (t, 2H, J=7.0 Hz), 2.76-2.39 (m, 5H),

2.20 (s, 3H), 2.17-1.85 (m, 6H); ESMS m/e : 399.3 $(M + H)^+$.

Example 10

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.93 (d, 2H, J=8.6 Hz), 7.45 (d, 2H, J=8.6 Hz), 7.39 (d, 1H, J=7.2 Hz), 7.32 (s, 1H), 7.24 (t, 1H, J=7.8 Hz), 6.94 (d, 1H, J=8.4 Hz), 3.21-3.18 (m, 2H), 3.05 (t, 2H, J=7.0 Hz), 2.64-2.51 (m, 4H), 2.28-1.86 (m, 8H), 1.26 (d, 6H, J=6.8 Hz); ESMS m/e : 427.3 $(M + H)^+$.

Example 11

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOHEXANECARBOXAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.93 (d, 2H, J=8.4 Hz), 7.55-7.19 (m, 5H), 6.93 (d, 1H, J=7.6 Hz), 3.25-3.00 (m, 4H), 2.65-2.45 (m, 4H), 2.30-1.50 (m, 18H); ESMS m/e : 467.3 $(M + H)^+$.

Example 12

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-PHENYLACETAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.92 (d, 2H, J=8.4 Hz), 7.46-7.26 (m, 9H), 7.20 (t, 1H, J=7.6 Hz), 6.92 (d, 1H, J=7.6 Hz), 3.75 (s, 2H), 3.15-3.13 (m, 2H), 3.03 (t, 2H, J=7.0 Hz), 2.64-2.46 (m, 3H), 2.22-1.60 (m, 8H); ESMS m/e : 475.3 $(M + H)^+$.

Example 13

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-(3-METHOXYPHENYL)ACETAMIDE: Using

Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (d, 2H, J=8.4 Hz), 7.44 (d, 2H, J=8.4 Hz) 7.38 (s, 1H), 7.35-7.25 (m, 3H), 7.19 (t, 1H, J=7.8 Hz), 6.94-6.86 (m, 3H), 3.81 (s, 3H), 3.72 (s, 2H), 3.12-3.09 (m, 2H), 3.02 (t, 2H, J=6.8 Hz), 2.57-2.44 (m, 3H), 2.20-1.60 (m, 8H); ESMS m/e : 505.3 $(M + H)^+$.

Example 14

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-METHOXYACETAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.93 (d, 2H, J=8.4 Hz), 7.50-7.25 (m, 5H), 6.98 (d, 1H, J=7.8 Hz), 4.01 (s, 2H), 3.57 (s, 3H), 3.30-3.15 (m, 2H), 3.06 (t, 2H, J=6.8 Hz), 2.70-2.50 (m, 3H), 2.35-1.80 (m, 8H); ESMS m/e : 429.3 $(M + H)^+$.

Example 15

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)METHANESULFONAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.95-6.96 (m, 8H), 3.48 (s, 3H), 3.28-2.90 (m, 6H), 2.80-2.57 (m, 3H), 2.38-1.86 (m, 6H); ESMS m/e : 435.2 $(M + H)^+$.

Example 16

N-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ETHANESULFONAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.93 (d, 2H, J=8.2 Hz), 7.45 (d, 2H, J=8.2 Hz), 7.30-7.08 (m, 3H), 6.99 (d, 1H, J=7.6 Hz), 3.26-3.02 (m, 6H), 2.69-2.45 (m, 3H), 2.32-1.75 (m, 8H), 1.36 (t, 3H, J=7.4 Hz); ESMS m/e : 449.3 $(M + H)^+$.

Example 17

N-{3-[1-(4-OXO-4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}ACETAMIDE: Using Method III, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.10-6.80 (m, 9H), 3.40-2.95 (m, 4H), 2.85-2.20 (m, 3H), 2.19 (s, 3H), 2.15-1.70 (m, 8H); ESMS m/e : 365.3 (M + H)$^+$.

Example 18

2-METHYL-N-{3-[1-(4-OXO-4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.99 (d, 2H, J=7.4 Hz), 7.57 (t, 1H, J=7.4 Hz), 7.48 (t, 2H, J=7.4 Hz), 7.45-7.20 (m, 2H), 7.24 (t, 1H, J=8.0 Hz), 6.94 (d, 1H, 8.0 Hz), 3.24-3.21 (m, 2H), 3.09 (t, 2H, J=7.0 Hz), 2.57-2.25 (m, 4H), 2.31-1.84 (m, 8H), 1.26 (d, 6H, J=7.2 Hz); ESMS m/e : 393.3 (M + H)$^+$.

Example 19

N-{3-[1-(4-OXO-4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}-2-PHENYLACETAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.98 (d, 2H, J=7.6 Hz), 7.65-7.15 (m, 11H), 6.92 (d, 2H, J=7.2 Hz), 3.74 (s, 2H), 3.20-2.95 (m, 4H), 2.65-2.40 (m, 3H), 2.25-1.70 (m, 8H); ESMS m/e : 441.3 (M + H)$^+$.

Example 20

2-(3-METHOXYPHENYL)-N-{3-[1-(4-OXO-4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}ACETAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.98 (d, 2H, J=7.6 Hz), 7.56 (t, 1H, J=7.62 Hz), 7.46 (t, 2H, J=7.6 Hz), 7.40 (s, 1H), 7.37-7.26 (m, 2H), 7.19

(t, 1H, J=7.8 Hz), 6.94- 6.86 (m, 3H), 3.81 (s, 3H), 3.71 (s, 3H), 3.12-3.03 (m, 4H), 2.57-2.44 (m, 3H), 2.16-1.77 (m, 8H); ESMS m/e : 471.3 $(M + H)^+$.

Example 21

N-(3-{1-[4-(2,4-DIMETHOXYPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.82 (d, 1H, J=8.8 Hz), 7.54 (d, 1H, J=7.6 Hz), 7.33 (s, 1H), 7.22 (t, 1H, J=7.6 Hz), 6.93 (d, 1H, J=7.6 Hz), 6.53 (d, 1H, J=8.8 Hz), 6.46 (s, 1H), 3.90 (s, 3H), 3.86 (s, 3H), 3.48-3.27 (m, 2H), 3.05 (t, 2H, J=6.8 Hz), 2.90-2.68 (m, 2H), 2.65-2.38 (m, 3H), 2.25 (s, 3H), 2.18-1.80 (m, 6H); ESMS m/e : 425.3 $(M + H)^+$.

Example 22

N-(3-{1-[4-(2,4-DIMETHOXYPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.98 (d, 1H, J=8.6 Hz), 7.41-7.37 (m, 2H), 7.24 (t, 1H, J=7.8 Hz), 6.96 (d, 1H, J=7.8 Hz), 6.54 (d, 1H, J=8.6 Hz), 6.46 (s, 1H), 3.89 (s, 3H), 3.86 (s, 3H), 3.11-3.08 (m, 2H), 2.98 (t, 2H, J=7.2 Hz), 2.53-2.46 (m, 4H), 2.13-1.79 (m, 8H), 1.25 (d, 6H, J=6.8 Hz); ESMS m/e : 453.3 $(M + H)^+$.

Example 23

N-(3-{1-[4-(2,4-DIMETHOXYPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-PHENYLACETAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.85 (m, 12H), 3.89 (s, 3H), 3.86 (s, 3H), 3.74 (s, 2H), 3.22-2.90 (m, 4H), 2.64-2.40 (m, 3H), 2.25-1.70 (m, 8H); ESMS m/e : 501.3 $(M + H)^+$.

Example 24

N-(3-{1-[4-(2,4-DIMETHOXYPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-(3-METHOXYPHENYL)ACETAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.82 (d, 1H, J=8.8 Hz), 7.48-7.15 (m, 5H), 6.95-6.80 (m, 3H), 6.58-6.45 (m, 2H), 3.89 (s, 3H), 3.86 (s, 3H), 3.81 (s, 3H), 3.72 (s, 2H), 3.25-2.95 (m, 4H), 2.65-2.40 (m, 3H), 2.30-1.95 (m, 4H), 1.93-1.72 (m, 4H); ESMS m/e : 531.3 $(M + H)^+$.

Example 25

N-(3-{1-[4-OXO-4-(4-PHENOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained.

^{1}H NMR (400 MHz, $CDCl_3$) δ 8.15-6.75 (m, 13H), 3.30-2.80 (m, 4H), 2.75-2.10 (m, 5H), 2.03 (s, 3H), 2.00-1.60 (m, 6H); ESMS m/e : 457.3 $(M + H)^+$.

Example 26

2-METHYL-N-(3-{1-[4-OXO-4-(4-PHENOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.96 (d, 2H, J=8.8 Hz), 7.43-7.15 (m, 6H), 7.10-6.93 (m, 5H), 3.42-2.95 (m, 4H), 2.80-2.45 (m, 4H), 2.20-1.80 (m, 8H), 1.14 (d, 6H, J=6.8 Hz); ESMS m/e : 485.4 $(M + H)^+$.

Example 27

2-(3-METHOXYPHENYL)-N-(3-{1-[4-OXO-4-(4-PHENOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method II, the desired product was obtained. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.97 (d, 2H, J=8.8 Hz), 7.41-7.18 (m, 7H), 7.08-6.99 (m, 5H), 6.94-6.87 (m, 3H), 3.82 (s, 3H), 3.70 (s, 2H), 3.10-2.95 (m, 4H), 2.55-2.40 (m, 3H),

2.15-1.95 (m, 4H), 1.81- 1.70 (m, 4H); ESMS m/e : 563.4 $(M + H)^+$.

Example 28

N'-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-N,N-DIMETHYLSULFAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.93 (d, 2H, J=8.8 Hz), 7.44 (d, 2H, J=8.8 Hz), 7.27 (s, 1H), 7.25-7.10 (m, 2H), 6.94 (d, 1H, J=7.6 Hz), 3.30-3.10 (m, 2H), 3.04 (t, 2H, J=6.8 Hz), 2.83 (s, 6H), 2.68-2.45 (m, 3H), 2.30-1.75 (m, 8H); ESMS m/e : 464.3 $(M + H)^+$.

Example 29

N-(3-{1-[4-OXO-4-(2-THIENYL)BUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.90-6.78 (m, 7H), 3.22-2.88 (m, 4H), 2.69-2.25 (m, 5H), 2.02 (s, 3H), 2.00-1.64 (m, 6H); ESMS m/e : 371.2 $(M + H)^+$.

Example 30

N-(3-{1-[4-(4-ISOPROPYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 8.00-6.78 (m, 8H), 3.15-2.98 (m, 4H), 2.77-2.15 (m, 4H), 2.03 (s, 3H), 2.00-1.62 (m, 8H), 0.927 (d, 6H, J=6.0 Hz); ESMS m/e : 407.3 $(M + H)^+$.

Example 31

N-(3-{1-[4-(4-METHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ

7.90-6.80 (m, 8H), 3.10- 2.45 (m, 7H), 2.32 (S, 3H), 2.02 (s, 3H), 2.01-1.68 (m, 8H); ESMS m/e : 379.3 $(M + H)^+$.

Example 32

N-(3-{1-[4-(4-BROMOPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)ACETAMIDE: Using Method III, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.90-6.80 (m, 8H), 3.30-3.05 (m, 4H), 2.70-2.45 (m, 3H), 2.05 (s, 3H), 1.98-1.65 (m, 8H); ESMS m/e : 444.0 $(M + H)^+$.

EXAMPLE 33

N-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-2-PROPANESULFONAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.75 (s, 1H), 7.71 (d, 1H, J=7.6 Hz), 7.27-7.00 (m, 5H), 3.32-3.24 (m, 3H), 3.10-3.02 (m, 2H), 2.78-2.50 (m, 3H), 2.32 (s, 6H), 2.19-1.84 (m, 8H), 1.39 (d, 6H, J=6.8 Hz); ESMS m/e : 457.4 $(M + H)^+$.

Example 34

N-(3-{1-[4-OXO-4-(4-PHENOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-PROPANESULFONAMIDE: Using Method II, the desired product was obtained. ^1H NMR (400 MHz, $CDCl_3$) δ 7.97 (d, 2H, J=7.6 Hz), 7.44 (t, 2H, J=7.6 Hz), 7.27-7.00 (m, 9H), 3.35-2.96 (m, 5H), 2.69-2.45 (m, 3H), 2.14-1.79 (m, 8H), 1.39 (d, 6H, J=6.8 Hz); ESMS m/e : 521.4 $(M + H)^+$.

Example 35

N-(3-{1-[3-(4-CHLOROPHENYL)-3-METHOXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of 3-

methoxy-3-(p-chlorophenyl)-1-chloropropane (27.4 mg, 0.125 mmol), 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.125 mmol), diisopropylethylamine (0.50 mL) and catalytic amount of tetrabutylammonium iodide in dioxane (2.0 mL) was stirred at 90 °C for 72 hrs. The reaction mixture was concentrated to a small volume and chromatographed using preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave N-(3-{1-[3-(4-chlorophenyl)-3-methoxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (39.5 mg, 73.8% yield) as a thick oil: ^1H NMR δ 7.48 (S, 1 H), 7.34-7.3 (m, 2H), 7.25 (m, 4H), 6.96 (d, 1H, J=7.4 Hz), 4.20 (apparent dd, 1H, J=5.9, 7.6 Hz), 3.2 (s, 3H), 3.04 (d, 1H, J=10.1 Hz), 2.99 (d, 1H, J=10.1 Hz), 2.49 (h, 4H, J=6.6 Hz), 2.20-2.10 (m, 4H), 1.82 (m, 4H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 429.4 $(M + H)^+$.

Example 36

N-(3-{1-[6-(1,3-DIOXO-1,3-DIHYDRO-2H-ISOINDOL-2-YL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: The synthetic method is the same as described for 2-[6-(4-phenyl-1-piperidinyl)hexyl]-1H-isoindole-1,3(2H)-dione. N-(3-{1-[6-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)hexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: 506 mg (56% yield); ^1H NMR (400 MHz, $CDCl_3$) δ 7.86-7.80 (m, 2H), 7.73-7.68 (m, 2H), 7.44 (s, 1H), 7.37 (d, 1H, J=8.3 Hz), 7.22 (t, 1H, J=7.7 Hz), 6.96 (d, 1H, J=7.7 Hz), 3.69 (t, 2H, J=7.2 Hz), 3.01 (apparent d, 2H, J=11.3 Hz), 2.58-2.40 (m, 2H), 2.33 (m, 2H) 1.98 (dt, 2H, J=3.2, 11.3 Hz), 1.84-1.64 (m, 4H), 1.51 (q, 2H, J=7.1 Hz), 1.43-1.30 (m, 6H), 1.24 (d, 6H, J=6.8 Hz); ESMS m/e: 476.4 $(M + H)^+$.

Example 37

N-{3-[1-(3-METHOXY-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: A mixture of 3-methoxy-3-phenyl-1-chloropropane (23.1 mg, 0.126 mmol), 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.126 mmol), diisopropylethylamine (0.50 mL) and catalytic amount of tetrabutylammonium iodide in dioxane (2.0 mL) was stirred at 90 °C for 72 hrs. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave N-{3-[1-(3-methoxy-3-phenylpropyl)-4-piperidinyl]phenyl}-2-methylpropanamide (45.4 mg, 91.2% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.45 (S, 1 H), 7.34-7.25 (m, 5H), 7.25 (m, 2H), 6.96 (d, 1H, J=7.4 Hz), 4.20 (apparent dd, 1H, J=5.9, 7.6 Hz), 3.2 (s, 3H), 3.04 (d, 1H, J=10.1 Hz), 2.99 (d, 1H, J=10.1Hz), 2.49 (apparent sept, partially hidden, 4H, J=6.6 Hz), 2.3-2.1(m, 4H), 1.82 (m, 4H), 1.25 (d, 6H, J=7.1 Hz); ESMS m/e: 395.4 $(M + H)^+$.

Example 38

N-(3-{1-[4-(1,3-DIOXO-1,3-DIHYDRO-2H-ISOINDOL-2-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: The synthetic method is the same as described for 2-[6-(4-phenyl-1-piperidinyl)hexyl]-1H-isoindole-1,3(2H)-dione. N-(3-{1-[4-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide: 664 mg (74% yield); ^1H NMR (400 MHz, $CDCl_3$) δ 7.87-7.78 (m, 2H), 7.76-7.64 (m, 2H), 7.47 (s, 1H), 7.39 (d, 1H, J=7.6 Hz), 7.21 (t, 1H, J=8.1 Hz), 6.94 (d, 1H, J=7.6 Hz), 3.72 (t, 2H, J=6.8 Hz), 3.37-3.22 (m, 2H), 3.0 (apparent d, 2H, J=10.7 Hz), 2.75 (q, 2H, J=7.0 Hz), 2.64-2.33 (m, 4H), 1.99 (dt, 2H, J=2.6, 11.7 Hz), 1.86-1.65 (m, 2H), 1.63-1.50 (m, 2H), 1.23 and 1,21 (two d, 6H, J=5.5 Hz);

ESMS m/e: 448.4 $(M + H)^+$; Anal. Calc. for $C_{27}H_{34}N_3ClO_3+0.4H_2O$: C, 66.02; H, 7.14; N, 8.55. Found: C, 66.07; H, 6.78; N, 8.65.

Example 39

***N*-(3-{1-[4-(1,3-DIOXO-1,3-DIHYDRO-2*H*-ISOINDOL-2-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** The synthetic method is the same as described for 2-[6-(4-phenyl-1-piperidinyl)hexyl]-1H-isoindole-1,3(2H)-dione. N-(3-{1-[5-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)pentyl]-4-piperidinyl}phenyl)-2-methylpropanamide: 614 mg (64% yield); ^1H NMR (400 MHz, $CDCl_3$) δ 7.87-7.8 (m, 2H), 7.76-7.68 (m, 2H), 7.48 (s, 1H), 7.41 (d, 1H, J=7.6 Hz), 7.21 (t, 1H, J=7.6 Hz), 6.95 (d, 1H, J=7.6 Hz), 3.69 (t, 2H, J=7.2 Hz), 3.39-3.28 (m, 2H), 3.02 (apparent d, 2H, J=11.6 Hz), 2.78 (q, 2H, J=7.2 Hz), 2.64-2.52 (m, 1H), 2.52-2.40 (m, 1H), 2.40-2.31 (m, 2H), 2.01 (dt, 2H, J=3.7, 11.1 Hz), 1.85-1.64 (m, 2H), 1.58 (q, 2H, J=7.6 Hz), 1.45-1.32 (m, 2H), 1.23 (d, 6H, J=6.9 Hz); ESMS m/e: 462.4 $(M + H)^+$; Anal. Calc. for $C_{28}H_{36}N_3ClO_3$: C, 67.52; H, 7.29; N, 8.44. Found: C, 67.04; H, 7.06; N, 8.38.

Example 40

2-METHYL-N-{3-[1-(4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: A mixture of 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.100 mmol), 4-phenyl-1-chlorobutane (21.1 mg, 0.125 mmol), diisopropylethylamine (0.50 mL), catalytic amount of tetrabutylammonium iodide and dioxane (2.0 mL) was heated at reflux temperature for 3 days. The reaction mixture was concentrated and chromatographed using preparative TLC plates [2.5% of NH_3 (2.0 M in methanol)

in $CHCl_3$] afforded the product, 2-methyl-N-{3-[1-(4-phenylbutyl)-4-piperidinyl]phenyl}propanamide (9.50 mg, 25.1% yield) as a thick oil: ^1H NMR δ 7.37 (s, 1H), 7.29 (apparent d, 1H, J=7.9 Hz), 7.18 (m, 3H), 7.11 (m, 3H), 6.90 (apparent d, 1H, J=7.9 Hz), 3.02 (d, 2H, J=6.8 Hz), 2.41 (m, 4H, partially hidden), 2.01 (m, 2H), 1.78 (m, 4H), 1.57 (m, 4H), 1.18 (d, 6H, J=7.7 Hz); ESMS *m/e*: 379.4 $(M + H)^+$.

Example 41

N-(3-{1-[3-(1,3-DIOXO-1,3-DIHYDRO-2H-ISOINDOL-2-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:

The synthetic method is the same as described for 2-[6-(4-phenyl-1-piperidinyl)hexyl]-1H-isoindole-1,3(2H)-dione. N-(3-{1-[3-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide: 810 mg (93% yield); ^1H NMR (400 MHz, $CDCl_3$) δ 7.87-7.82 (m, 2H), 7.73-7.68 (m, 2H), 7.57 (s, 1H), 7.36 (d, 1H, J=8.5 Hz), 7.18 (t, 1H, J=7.7 Hz), 6.79 (d, 1H, J=7.1 Hz), 3.78 (t, 2H, J=6.8 Hz), 3.06 (quintet, 2H, J=6 Hz), 2.95 (apparent d, 2H, J=12.2 Hz), 2.58-2.31 (m, 4H), 1.96-1.83 (m, 2H), 1.70 (apparent d, 2H, J=12.1 Hz), 1.52 (dt, 2H, J=3.5, 12.5 Hz), 1.03 (d, 6H, J=6.5 Hz); ESMS m/e: 434.4 $(M + H)^+$.

Example 42

N-(3-{1-[(3S)-3-HYDROXY-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of (S)-(-)-3-chloro-1-phenyl-1-propanol (0.426 g, 2.50 mmol, 99%ee), 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (0.565 g, 2.00 mmol), diisopropylethylamine (1.29 g, 10.0 mmol), dioxane (5.0 mL) and catalytic amount of tetrabutylammonium iodide

was stirred at 90 °C for 72 hrs. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (306 mg, 39.3 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.46 (S, 1 H), 7.42 (d, 4H, J=8.1 Hz), 7.35 (m, 1 H), 7.30 (d, 1 H, J=8.0 Hz), 7.23 (t, 1H, J=8.1 Hz), 7.12 (s, 1H), 6.96 (apparent dd, 1H, J=8.0 Hz), 5.0 (apparent dd, 1H, J=4.4, 8.3 Hz), 3.18 (apparent dd, 2H, J=2.5, 12.5 Hz), 2.74 (m, 2 H), 2.50 (m, 2H), 2.3-2.1 (m, 6H), 1.8 (m, 2H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 389.2 $(M + H)^+$.

Example 43

N-(3-{1-[3-METHOXY-3-(4-METHYLPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of 3-methoxy-3-(p-tolyl)-1-chloropropane (24.9 mg, 0.126 mmol), 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.126 mmol), diisopropylethylamine (0.50 mL) and catalytic amount of tetrabutylammonium iodide in dioxane (2.0 mL) was stirred at 90 °C for 72 hrs. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (10.9 mg, 21.2 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.44 (s, 1 H), 7.38 (m, 1H), 7.3-7.1 (m, 5 H), 6.96 (d, 1H, J=7.4 Hz), 4.18 (apparent dd, 1H, J=5.6, 7.9 Hz), 3.24 (d, 1H, J=8.2 Hz), 3.2 (s, 3H), 3.11 (m, 2H, J=10.1Hz), 2.49 (m, 4H), 2.35 (s, 3H), 2.3-2.1(m, 3H), 1.92 (d, 4H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 409.4 $(M + H)^+$.

Example 44

N-{3-[1-(3-ISOPROPOXY-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: A mixture of 3-

isopropyl-3'-phenyl-1-chloropropane (26.6 mg, 0.126 mmol), 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.126 mmol), diisopropylethylamine (0.50 mL) and catalytic amount of tetrabutylammonium iodide in dioxane (2.0 mL) was stirred at 90 °C for 72 hrs. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (14.1 mg, 26.5% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1H), 7.43-7.37 (m, 2H), 7.33 (m, 3H), 7.23 (m, 2H), 6.95 (d, 1H, J=8.4 Hz), 4.46 (apparent dd, 1H, J=5.0, 8.3 Hz), 3.49 (apparent sept, 1H, J=7.1 Hz), 3.10 (s, 2H), 2.70 (m, 2H), 2.52 (apparent sept, partially hidden, 4H, J=6.6 Hz), 2.30-2.10 (m, 2H), 1.90-1.80 (d, 4H), 1.25 (d, 6H, J=7.1 Hz), 1.15 (d, 3H, J=6.4 Hz), 1.08 (d, 3H, J=6.4 Hz); ESMS *m/e*: 423.4 $(M + H)^+$.

Example 45

N-(3-{1-[4,4-BIS(4-FLUOROPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of 4,4-bis(4-fluoro-phenyl)-1-chloro-butane (39.0 mg, 0.126 mmol), 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.126 mmol), diisopropylethylamine (0.50 mL) and catalytic amount of tetrabutylammonium iodide in dioxane (2.0 mL) was stirred at 90 °C for 72 hrs. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (15.9 mg, 25.2 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.02 (s, 1H), 7.41 (s, 1H), 7.3-7.15 (m, 4H), 7.10 (m, 3H), 6.89 (apparent t, 5H), 3.81 (t, 1H, J=7.8 Hz), 3.30 (s, 1H), 2.91 (d, 1H, J=12,5 Hz), 2.80 (m, 1H), 2.40 (m, 2H), 2.31 (t, 1H, J=8.0 Hz), 1.93 (apparent q, 3H, J=8.0 Hz), 1.72 (m, 3H), 1.40 (m, 2H),

1.20 (m, 2H), 1.15 (d, 6H, J=8.1 Hz); ESMS *m/e*: 491.4 $(M + H)^+$

EXAMPLE 46

N-{3-[1-(3-METHOXYBENZYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: A mixture of 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.100 mmol), 3-methoxybenzyl chloride (19.6 mg, 0.125 mmol), diisopropylethylamine (0.50 mL), catalytic amount of tetrabutylammonium iodide and dioxane (2.0 mL). Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] afforded the desired product (10.2 mg, 27.9% yield) as a yellow solid: ^1H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1H), 7.35 (apparent d, 1H, J=8.3 Hz), 7.27-7.21 (m, 2H), 6.95 (apparent t, 3H, J=6.9 Hz), 6.82 (apparent dd, 1H, J=2.4, 8.3 Hz), 3.84 (m, 3H), 3.56 (s, 2H), 3.05 (d, 2H, J=10.5 Hz), 2.51 (apparent sept, partially hidden, 4H, J=7.2 Hz), 2.13 (apparent t, 2H, J=9.7 Hz), 1.88 (m, 2H), 1.25 (d, 6H, J=6.7 Hz); ESMS *m/e*: 367.3 $(M + H)^+$.

Example 47

N-(3-{1-[3,5-BIS(TRIFLUOROMETHYL)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (28.3 mg, 0.100 mmol), 3,5-bis(trifluoromethyl)benzyl bromide (38.4 mg, 0.125 mmol), diisopropylethylamine (0.50 mL), catalytic amount of tetrabutylammonium iodide and dioxane (2.0 mL). Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (12.2 mg, 25.8% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.83 (s, 2H), 7.77 (s, 1H), 7.53 (s, 1H), 7.30-7.21 (m, 2H), 7.16 (s,

1H), 6.98 (apparent d, 1H, J=7.6 Hz), 3.62 (s, 2H), 2.94 (d, 2H, J=9.4 Hz), 2.51 (apparent sept, partially hidden, 2H, J=6.6 Hz), 2.14 (m, 2H), 1.82 (m, 4H), 1.25 (d, 6H, J=6.6 Hz); ESMS *m/e*: 473.2 $(M + H)^+$.

Example 48

N-(3-{1-[(3R)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE

Method A

4-{[(1*R*)-3-chloro-1-phenylpropyl]oxy}-1,2-dimethoxybenzene: A mixture of 3,4-dimethoxyphenol (4.07 g, 26.4 mmol), (S)-(-)-3-chloro-phenyl-1-propanol (4.50 g, 26.4 mmol, 99%ee, Aldrich Chemical Co.), triphenylphosphine (6.92 g, 26.4 mmol) and diethyl azodicarboxylate (4.59 g, 26.4 mmol) in THF (110 mL) was stirred at room temperature for 24 h. The reaction mixture was concentrated *in vacuo*. At this point, the residue can either be washed with pentane (x3) and the combined pentane extracts were concentrated and chromatographed (silica with hexanes-EtOAc 8:1 as the eluent) to give the desired product (as described as a general procedure by: Srebnik, M.; Ramachandran, P.V.; Brown, H.C. *J. Org. Chem. 1988, 53*, 2916-2920). This procedure was performed on a smaller scale reaction and only a 40% yield of the product was realized.

Alternatively, on a larger scale (26.4 mmol), the crude product was triturated with a small amount of dichloromethane and the precipitated triphenylphosphine oxide was filtered. The filtrate was concentrated and the crude product was chromatographed to give the desired product as a thick yellow oil (7.30 g, 88.9%

yield): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39-7.32 (m, 4H), 7.20 (m, 1H), 6.64 (d, 1H, J=8.7 Hz), 6.51 (d, 1H, J=2.7 Hz), 6.30 (dd, 1H, J=2.7, 8.7 Hz), 5.27 (apparent dd, 1H, J=4.5, 8.7 Hz), 3.79 (s, 3H), 3.77 (s, 3H), 3.61 (m, 1H), 2.45 (m, 1 H), 2.20 (m, 1H), 1.80 (s, 1H); ESMS *m/e*: 307.11 $(M+H)^+$.

N-(3-{1-[(3R)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of potassium carbonate (321 mg, 2.32 mmol), sodium iodide (522 mg, 3.48 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (570 mg, 2.32 mmol) and 4-{[(1*R*)-3-chloro-1-phenylpropyl]oxy}-1,2-dimethoxybenzene (712 mg, 2.32 mmol) in DMF (5.0 mL) was stirred at 100 °C for 3 hrs, at which time TLC indicated that the reaction was complete. The reaction mixture was poured into water (50 mL) and the aqueous layer was extracted with methylene chloride (3x30 mL). The combined organic extracts were washed with brine (30 mL), dried over $MgSO_4$ and concentrated under reduced pressure. The crude product was purified by Prep-TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to afford the product (970 mg, 90.1%) as a thick oil.

Method B

Into a 25-mL RB-flask was added triphenylphosphine (9.80 mg, 0.0375 mmol), diethyl azodicarboxylate (5.22 mg, 0.0300 mmol), N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 3,4-dimethoxyphenol (7.70 mg, 0.050 mmol) and THF (1.0 mL) at room temperature. The reaction mixture was stirred at room temperature overnight (16 hrs). The solvent was removed under reduced pressure and the

residue was purified by preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to afford the desired product (4.4 mg, 34.1 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1 H), 7.40-7.30 (m, 4H), 7.25 (m, 3H), 6.97 (d, 1H, J=7.8 Hz), 6.64 (d, 1H, J=9.1 Hz), 6.51 (d, 1H, J=2.6 Hz), 6.29 (d, 1H, J=2.6, 9.1 Hz), 5.20 (apparent dd, 1H, J=4.4, 8.5 Hz), 3.80 (s, 3H), 3.77 (s, 3H), 3.23 (m, 2H), 2.77 (m, 2 H), 2.5 (m, 2H), 2.3-2.1(m, 6H), 1.80 (m, 2H), 1.25 (d, 6H, J=7.9 Hz); ESMS *m/e*: 517.4 $(M + H)^+$.

Example 49

2-METHYL-N-(3-{1-[(3S)-3-PHENOXY-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), phenol (4.70 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (2.7 mg, 23.6 % yield) as a thick oil: ^1H NMR δ 7.46 (s, 2H), 7.40-7.30 (m, 4H), 7.25 (m, 3 H), 7.20 (m, 2H), 6.97 (apparent d, 1H, J=7.4 Hz), 6.89 (apparent tt, 1H, J=0.8, 7.6 Hz), 6.84 (apparent dt, 1H, J=0.8, 8.0 Hz), 5.20 (apparent dd, 1H, J=4.4, 8.5 Hz), 3.35 (m, 2H), 2.91 (m, 2H), 2.60 (m, 2H), 2.30-2.10 (m, 6H), 1.90 (m, 2H), 1.25 (d, 6H, J=7.9 Hz); ESMS *m/e*: 457.4 $(M + H)^+$;

Example 50

N-(3-{1-[(3S)-3-(4-METHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-

(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 4-methoxyphenol (6.20 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.2 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.6 mg, 37.9 % yield) as a thick oil. ^1H NMR (400 MHz, $CDCl_3$) δ 7.38-7.14 (m, 8H), 6.90 (apparent d, 1H, J=7.7 Hz), 6.72-6.46 (m, 4H), 5.09 (apparent dd, 1H, J=4.8, 8.1 Hz), 3.64 (s, 3H), 3.18 (m, 2H), 2.73 (m, 2H), 2.50 (m, 2H), 2.37-1.72 (m, 8H), 1.25 (d, 6H, J=7.4 Hz); ESMS *m/e*: 487.4 $(M + H)^+$.

Example 51

N-(3-{1-[(3S)-3-(3-CHLOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 3-chlorophenol (6.40 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.9 mg, 40.0 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.39 (s, 1H), 7.35-7.10 (m, 7H), 7.02 (t, 1H, J=8.0 Hz), 6.90 (d, 1H, J=7.6 Hz), 6.84-6.75 (m, 2H), 6.65 (m, 1H), 5.09 (apparent dd, 1H, J=4.99, 8.1 Hz), 3.10 (m, 2H), 2.60 (m, 2H), 2.50 (m, 2H), 2.30-1.70 (m, 8H), 1.18 (d, 6H, J=6.8 Hz); ESMS *m/e*: 491.4 $(M + H)^+$.

Example 52

N-(3-{1-[(3S)-3-(4-CHLOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 4-chlorophenol (6.40 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (3.3 mg, 26.9 % yield) as a thick oil: ^1H NMR δ 7.36 (s, 1H), 7.35-7.22 (m, 7H), 7.12 (m, 2H), 6.97 (apparent d, 1H, J=7.2 Hz), 6.77 (m, 2H), 5.23 (m, 1H), 3.18 (m, 2H), 2.70 (m, 2H), 2.50 (m, 2H), 2.40-1.80 (m, 8H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 491.4 $(M + H)^+$.

Example 53

2-METHYL-N-[3-(1-{(3S)-3-PHENYL-3-[4-(TRIFLUOROMETHYL) PHENOXY] PROPYL}-4-PIPERIDINYL) PHENYL] PROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 4-trifluoromethylphenol (8.100 mg, 0.050 mmol), triphenylphosphine (9.8 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (5.10 mg, 38.9 % yield) as a thick oil: ^1H NMR δ 8.06 (s, 1H), 7.49 (s, 1H), 7.44 (apparent d, 2H, J=.6 Hz), 7.38-7.30 (m, 4H), 7.30-7.20 (m, 3H), 6.96

(apparent d, 1H, J=7.6 Hz), 6.91 (apparent d, 2H, J=8.6 Hz), 5.34 (m, 1H), 3.19 (m, 2H), 2.72 (m, 2H), 2.53 (m, 2H), 2.40-1.80 (m, 8H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 525.4 $(M + H)^+$.

Example 54

N-(3-{1-[(3R)-3-(2,5-DIFLUOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 2,5-difluorophenol (6.50 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (3.60 mg, 29.3 % yield) as a thick oil: ^1H NMR δ 7.46 (s, 1H), 7.40-7.32 (m, 4H), 7.31-7.20 (m, 2H), 7.17 (s, 1H), 7.01-6.92 (m, 2H), 6.65-6.42 (m, 2H), 5.27 (m, 1H), 3.13 (m, 2H), 2.64 (m, 2H), 2.51 (m, 2H), 2.28-1.80 (m, 8 H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 493.4 $(M + H)^+$.

Example 55

N-(3-{1-[(3R)-3-(3,4-DICHLOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 3,4-dichlorophenol (8.20 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5%

of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (5.20 mg, 39.7 % yield) as a thick oil: ^1H NMR ($CDCl_3$) δ 7.70-7.63 (m, 2H), 7.55 (m, 1H), 7.47-7.43 (m, 3H), 7.40-7.19 (m, 3H), 7.00-6.50 (m, 2H), 6.69 (dd, 1H, J=2.2, 8.8 Hz), 5.25 (m, 1H), 3.20 (m, 2H), 2.70 (m, 2H), 2.53 (m, 2H), 2.40-2.20 (m, 4H), 2.10-1.80 (m, 4H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 525.4 $(M + H)^+$.

Example 56

2-METHYL-N-(3-{1-[(3R)-3-PHENOXY-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), phenol (4.70 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.0 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.1 mg, 36.0 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.45 (s, 1H), 7.40-7.15 (m, 10H), 6.97 (d, 1H, J=7.6 Hz), 6.88-6.82 (m, 2H), 5.26 (m, 1H), 3.18 (m, 2H), 2.75 (m, 2H), 2.53 (m, 2H), 2.40-2.10 (m, 4H), 2.10-1.80 (m, 4H), 1.25 (d, 6H, J=6.9 Hz); ESMS *m/e*: 457.4 $(M + H)^+$.

Example 57

N-(3-{1-[(3R)-3-HYDROXY-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE

Method A

Into a 25-mL RB-flask was added (R)-(+)-3-chloro-1-phenyl-1-propanol (0.545 g, 3.19 mmol, 99%ee, Aldrich Chemical Co.), 2-methyl-*N*-[3-(4-

piperidinyl)phenyl]propanamide (0.748 g, 3.04 mmol), potassium carbonate (0.420 g, 3.04 mmol) and sodium iodide (0.684 g, 4.56 mmol) and DMF (6.0 mL) at room temperature. After stirring at 100 °C for 3 hrs, the TLC showed the reaction was complete. The reaction mixture was poured into water (50 mL) and the aqueous layer was extracted with methylene chloride (3x20 mL). The combined organic extracts were washed with brine (20 mL), dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by flash chromatography (1:1= hexane: ethyl acetate with 1% isopropylamine) to afford the desired product (1.09 g, 94.3 % yield) as light-yellow solid: ^1H NMR (400 MHz, $CDCl_3$) δ 8.10 (s, 1H), 7.46-7.35 (m, 6H), 7.27 (m, 2H), 6.98 (apparent d, 1H, J=7.6 Hz), 5.02 (apparent dd, 1H, J=4.4, 8.1 Hz), 3.18 (apparent dd, 2H, J=2.5, 12.5 Hz), 2.74 (m, 2 H), 2.50 (m, 2H), 2.30-2.10 (m, 6H), 1.80 (m, 2H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 381.2 $(M + H)^+$.

The hydrochloric salt was prepared by addition of a slight excess of 1 N HCl in ether (1.2 eq.) to a solution of the free base in dichloromethane. The solvent was removed under reduced pressure, the residue was washed with ether and dried under reduced pressure: Anal. Calc. for $C_{24}H_{32}N_2O_2$+HCl+0.8H_2O: C, 66.82; H, 8.08; N, 6.49; Cl, 8.22. Found: C, 66.90; H, 7.78; N, 6.63; Cl, 8.52.

Method B

Into a 25-mL RB-flask was added (R)-(+)-3-chloro-1-phenyl-1-propanol (0.426 g, 2.50 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (0.565 g, 2.00 mmol),

diisopropylethylamine (1.29 g, 10.0 mmol), dioxane (5.0 mL) and catalytic amount of tetrabutylammonium iodide at room temperature. After stirring at 90 °C for 72 hrs, the reaction mixture was poured into water (50 mL) and the aqueous layer was extracted with methylene chloride (3x20 mL). The combined organic extracts were washed with brine (20 mL), dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by preparative TLC plates (1:5:100=isopropylamine:methanol:ethyl acetate) to afford the desired product (0.260 g, 34.2 % yield) as light-yellow solid.

Example 58

N-(3-{1-[(3S)-3-(4-CYANO-PHEONXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: N-(3-{1-[(3S)-3-(4-cyanophenoxy)-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide

A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 4-cyanophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.70 mg, 71.3 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.54 (m, 2H), 7.48 (d, 2H, J=8.4 Hz), 7.30-7.20 (m, 3H), 7.20 (m, 3H), 6.97 (apparent d, 1H, J=8.4 Hz), 6.92 (apparent d, 2H, J=8.4 Hz), 5.36 (apparent dd, 1H, J=3.9, 7.6 Hz), 3.12 (m, 2H), 2.61 (m, 2H), 2.53 (apparent sept, partially hidden, 2H, J=7.6 Hz), 2.30-2.10 (m, 6H), 1.82

(m, 2H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 482.2 $(M + H)^+$.

Example 59

N-(3-{1-[(3S)-3-(4-FLUOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 4-fluorophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.20 mg, 64.7% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.40 (m, 2H), 7.30-7.20 (m, 5H), 7.20 (m, 3H), 6.97 (apparent d, 1H, J=7.7 Hz), 6.87 (m, 1H), 6.76 (m, 1H), 5.26 (apparent dd, 1H, J=4.0, 8.1 Hz), 3.09 (m, 2H), 2.66 (m, 2H), 2.51 (m, 2H), 2.3-2.1 (m, 6H), 1.82 (m, 2H), 1.25 (d, 6H, overlapped); ESMS *m/e*: 475.2 $(M + H)^+$.

Example 60

N-(3-{1-[(3S)-3-(4-BROMOPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 4-bromophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] the desired product (0.70 mg, 9.6% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1H), 7.48

(m, 2H), 7.30-7.20 (m, 5H), 7.20 (m, 3H), 6.97 (apparent d, 1H, J=8.5 Hz), 6.73 (apparent d, 2H, J=8.5 Hz), 5.22 (apparent dd, 1H, J=4.9, 7.8 Hz), 3.15 (m, 2H), 2.65 (m, 2H), 2.51 (apparent sept, partially hidden, 2H, J=7.6 Hz), 2.30-2.10 (m, 6H), 1.82 (m, 2H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 535.1 $(M + H)^+$.

Example 61

N-(3-{1-[(3S)-3-(3-METHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 3-methoxyphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (3.1 mg, 46.6 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.47 (d, 1H, J=6.7 Hz), 7.42 (s, 1H), 7.3-7.20 (m, 3H), 7.20 (m, 3H), 7.07 (t, 1H, J=8.4 Hz), 6.97 (apparent d, 1H, J=6.7 Hz), 6.40 (m, 3H), 5.27 (apparent dd, 1H, J=5.3, 8.0 Hz), 3.74 (s, 3H), 3.38 (m, 2H), 2.93 (m, 2H), 2.61 (s, 1H), 2.53 (apparent sept, partially hidden, 1H, J=6.5 Hz), 2.30-2.10 (m, 6H), 1.82 (m, 2H), 1.25 (d, 6H, J=6.9 Hz); ESMS *m/e*: 487.3 $(M + H)^+$.

Example 62

N-(3-{1-[(3S)-3-(4-CYANO-2-METHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-methoxy-4-cyanophenol (100 mg),

triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (5.50 mg, 76.5 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.51 (s, 1H), 7.38 (s, 1H), 7.37 (d, 2H, J=2.4 Hz), 7.20 (m, 4H), 7.10 (d, 1H, J=2.4 Hz), 7.08 (s, 1H), 6.99 (apparent d, 1H, J=8.3 Hz), 6.76 (apparent d, 1H, J=8.3 Hz), 5.43 (apparent dd, 1H, J=5.1, 8.0 Hz), 3.91 (s, 3H), 3.34 (m, 2H), 2.63 (m, 2H), 2.63 (s, 1H), 2.53 (apparent sept, partially hidden, 1H, J=7.7 Hz), 2.30-2.10 (m, 6H), 1.82 (m, 2H), 1.28 (d, 6H, J=6.8 Hz); ESMS *m/e*: 512.2 $(M + H)^+$.

Example 63

N-(3-{1-[(3S)-3-(5-ACETYL-2-METHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:

A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-methoxy-5-acetylphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (1.60 mg, 22.2 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.52 (d, 2H, J=2.4 Hz), 7.3-7.2 (m, 5H), 7.20 (m, 3H), 6.97 (apparent d, 1H, J=6.7 Hz), 6.69 (apparent d, 1H, J=8.0 Hz), 5.47 (apparent dd, 1H, J=4.3, 7.8 Hz), 3.95 (s, 3H), 3.38 (m, 2H), 2.93 (m, 2H), 2.61 (s, 1H), 2.53 (apparent sept, partially hidden, 1H, J=7.6 Hz), 2.50 (s, 3H), 2.30-2.10 (m, 6H),

1.82 (m, 2H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 529.6 $(M + H)^+$.

Example 64

N-(3-{1-[(3R)-3-(2-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.2 mg, 0.0137 mmol), 2-acetylphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (1.70 mg, 24.9 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.65 (m, 1H), 7.55 (s, 1H), 7.30-7.20 (m, 5H), 7.20 (m, 3H), 6.97 (m, 2H), 6.76 (apparent d, 1H), 5.49 (apparent dd, 1H, J=4.3, 8.0 Hz), 3.38 (m, 2H), 2.93 (m, 2H), 2.71 (s, 3H), 2.60 (s, 1H), 2.53 (apparent sept, partially hidden, 1H, J=7.6 Hz), 2.30-2.10 (m, 6H), 1.82 (m, 2H), 1.25 (d, 6H, J=6.9 Hz); ESMS *m/e*: 498.8 (M^+).

Example 65

N-[3-(1-{(3R)-3-[2-FLUORO-5-(TRIFLUOROMETHYL)PHENOXY]-3-PHENYLPROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-fluoro-5-trifluoromethylphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired

product (2.50 mg, 33.7 % yield) as a thick oil: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.07 (s, 1H), 7.67 (m, 1H), 7.54 (m, 1H), 7.45 (m, 2H), 7.30-7.10 (m, 6H), 7.14 (d, 1H, J=7.4 Hz), 6.97 (apparent d, 1H, J=7.7 Hz), 5.37 (apparent dd, 1H, J=5.0, 8.5 Hz), 3.4 (m, 2H), 2.8 (m, 2H), 2.6 (s, 1H), 2.53 (apparent sept, partially hidden, 1H, J=7.4 Hz), 2.30-2.10 (m, 6H), 1.80 (m, 2H), 1.25 (d, 6H, J=7.1 Hz, overlapped); ESMS *m/e*: 542.6 (M^+), 543.54 $(M + H)^+$.

Example 66

N-[3-(1-{(3S)-3-[2-FLUORO-5-(TRIFLUOROMETHYL)PHENOXY]-3-PHENYLPROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-fluoro-5-trifluoromethylphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (3.00 mg, 40.4% yield) as a thick oil: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1H), 7.67 (m, 2H), 7.55 (m, 2H), 7.50-7.40 (m, 3H), 7.30-7.10 (m, 3H), 7.17 (d, 1H, J=8.9 Hz), 7.07 (apparent d, 1H, J=6.7 Hz), 6.97 (apparent d, 1H, J=7.8 Hz), 5.37 (apparent dd, 1H, J=4.2, 8.1 Hz), 3.37 (m, 2H), 2.93 (m, 2H), 2.63 (s, 1H), 2.50 (apparent sept, partially hidden, 1H, J=7.9 Hz), 2.30-2.10 (m, 6H), 1.85 (m, 2H), 1.25 (d, 6H, J=6.9 Hz); ESMS *m/e*: 542.7 $(M + H)^+$.

Example 67

N-(3-{1-[(3S)-3-(2,5-DIFLUOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2,5-difluorophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (2.70 mg, 40.1 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1H), 7.40-7.30 (m, 4H), 7.20 (m, 2H), 7.17 (s, 1H), 6.97 (m, 2H), 6.58 (m, 1H), 6.51 (m, 1H), 5.27 (apparent dd, 1H, J=5.1, 8.2 Hz), 3.13 (apparent d, J=9.7 Hz, 2H), 2.64 (m, 2H), 2.51 (m, 2H), 2.34 (apparent sept, partially hidden, J=7.1 Hz, 1H), 2.17 (m, 3H), 1.90-1.80 (m, 4H), 1.25 (d, 6H, J=7.1 Hz); ESMS *m/e*: 493.1 $(M + H)^+$.

Example 68

N-(3-{1-[(3R)-3-(3-CHLOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 3-chlorophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (2.4 mg, 35.8% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.30 (m, 2H), 7.30-7.20 (m, 3H), 7.20 (m, 3H), 6.90 (apparent d, 1H, J=7.7 Hz), 6.71 (apparent d, 1H, J=2.9 Hz), 6.69

(apparent t, 1H, J=2.9 Hz), 6.67 (apparent t, 1H, J=2.9 Hz), 6.65 (apparent d, 1H, J=2.9 Hz), 5.09 (apparent dd, 1H, J=4.8, 8.1 Hz), 3.18 (m, 2H), 2.73 (m, 2H), 2.50 (apparent sept, partially hidden, 2H, J=7.1 Hz), 2.30-2.10 (m, 6H), 1.89 (m, 2H), 1.25 (d, 6H, overlapped); ESMS *m/e*: 491.1 $(M + H)^+$.

Example 69

(1S)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL 1-NAPHTHOATE: Into a 25-mL RB-flask was added N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 1-naphthalenecarbonyl chloride (100 mg), diisopropylethylamine (0.30 mL) in THF (0.50 mL) at room temperature. After stirring for 16 hrs at room temperature, the reaction mixture was concentrated under reduced pressure. The residue was purified using preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.70 mg, 71.3 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.90 (d, 1H, J=8.9 Hz), 8.28 (apparent dd, 1H, J=1.5, 7.2 Hz), 8.03 (d, 1H, J=8.7 Hz), 7.88 (dm, 2H, J=8.7 Hz), 7.60-7.48 (m, 7H), 7.40-7.32 (m, 3H), 7.25 (m, 1H), 6.90 (apparent d, 1H, J=7.4 Hz), 6.18 (apparent dd, 1H, J=5.7, 7.8 Hz), 3.42 (m, 2H), 2.84 (m, 2H), 2.53 (m, 2H), 2.44 (apparent sept, partially hidden, 4H, J=7.5 Hz), 2.30-2.10 (m, 2H), 1.82 (m, 2H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 535.6 $(M + H)^+$.

Example 70

N-(3-{1-[(3S)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-

piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-acetylphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (1.50 mg, 22.0% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.65 (m, 1H), 7.55 (s, 1H), 7.30-7.20 (m, 5H), 7.20 (m, 3H), 6.97 (m, 2H), 6.76 (apparent d, 1H), 5.49 (apparent dd, 1H, J=4.3, 8.0 Hz), 3.38 (m, 2H), 2.93 (m, 2H), 2.75 (s, 3H), 2.53 (apparent sept, partially hidden, 2H, J=7.6 Hz), 2.30-2.10 (m, 6H), 1.92 (m, 2H), 1.25 (d, 6H, J=6.9 Hz); ESMS *m/e*: 498.81 (M^+), 499.6 $(M + H)^+$.

Example 71

N-(3-{1-[(3S)-3-(2-FLUOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-fluorophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (3.5 mg, 53.9% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.07 (s, 1H), 7.65 (m, 1H), 7.41 (s, 1H), 7.40-7.10 (m, 5H), 7.05 (m, 2H), 6.97 (apparent d, 1H, J=8.7 Hz), 6.86 (m, 2H), 6.79 (apparent dt, 1H, J=2.4, 7.9 Hz), 5.31 (apparent dd, 1H, J=4.5, 8.0 Hz), 3.39 (m, 2H), 2.97 (m, 2H), 2.53 (apparent sept, partially hidden, 2H, J=7.5 Hz), 2.3-2.1

(m, 6H), 1.92 (m, 2H), 1.25 (d, 6H, J=6.7 Hz); ESMS *m/e*: 475.7 $(M + H)^+$.

Example 72

(4S)-N-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-4-(3,5-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXAMIDE

Method: Into a 20 ml vial was added *N*1-{3-[1-(3-aminopropyl)-4-piperidyl]phenyl}acetamide (15 mg, 0.054 mmol), (4*S*)-4-(3,5-difluorophenyl)-2-oxo-oxazolidine-3-carboxylic acid-4-nitro-phenyl ester (39.3 mg, 1.08 mmol, 2 eq) and dichloromethane with 0.6% of methanol (3 ml) at room temperature. After stirring at room temperature for 3 hrs, the reaction mixture was filtered, and purified by preparative silica TLC (19:1 = chloroform : methanol) to afford the desired product (18.3 mg, 68% yield); ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.09 (br s, 1H), 7.40 (d, 1H, J=8.0 Hz), 7.36-7.28 (m, 2H), 7.24 (t, 1H, J=8.0 Hz), 6.99 (d, 1H, J=8.0 Hz), 6.86-6.82 (m, 2H), 5.41 (dd, 1H, J=4.1, 9.0 Hz), 4.72 (t, 1H, J=9.0 Hz), 4.22 (dd, 1H, J=3.9, 9.1 Hz), 3.42-3.29 (m, 2H); 3.02 (d, 2H J=11.1 Hz), 2.52-2.38 (m, 3H), 2.16 (s, 3H), 2.08-1.98 (m, 2H), 1.86-1.70 (m, 6H); ESMS m/e: 501.2 $(M + H)^+$; Anal. Calc. for $C_{26}H_{30}F_2N_4O_4+0.5H_2O$: C, 60.64; H, 6.18; N, 10.88. Found: C, 60.67; H, 5.79; N, 10.86.

Example 73

The synthetic method is the same as described for the synthesis of (4S)-*N*-(3-{4-[3-(acetylamino)phenyl]-1-piperidinyl}propyl)-4-(3,5-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxamide.

(4S)-N-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-OXO-4-(3,4,5-TRIFLUOROPHENYL)-1,3-OXAZOLIDINE-3-CARBOXAMIDE: 18.8 mg (67% yield); ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.09 (br s, 1H), 7.41-7.20 (m, 3H), 7.02-6.91 (m, 3H), 5.37 (dd, 1H, J=3.8, 8.9 Hz), 4.71 (t, 1H, J=9 Hz), 4.21 (dd, 1H, J=4, 9.3 Hz), 3.43-3.27 (m, 2H), 3.02 (d, 2H, J=11.0 Hz), 2.53-2.37 (m, 3H), 2.16 (s, 3H), 2.08-1.97 (m, 2H), 1.85-1.69 (m, 6H); ESMS m/e: 519.2 $(M + H)^+$; Anal. Calc. for $C_{26}H_{29}F_3N_4O_4+0.5H_2O$: C, 59.20; H, 5.73; N, 10.62. Found: C, 59.40; H, 5.35; N, 10.65.

Example 74

The synthetic method is the same as described for the synthesis of (4S)-*N*-(3-{4-[3-(acetylamino)phenyl]-1-piperidinyl}propyl)-4-(3,5-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxamide.

N-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-4-(3,4-DIFLUOROPHENYL)-5,5-DIMETHYL-2-OXO-1,3-OXAZOLIDINE-3-CARBOXAMIDE: 19.6 mg (68% yield); ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.18 (t, 1H, J=5.9 Hz), 7.41 (d, 1H, J=8.8 Hz), 7.33 (s, 1H), 7.27-7.14 (m, 2H), 7.02-6.88 (m, 3H), 5.04 (s, 1H), 3.34 (qm, 2H, J=6.3 Hz), 3.02 (dm, 2H, J=10.9 Hz), 2.53-2.38 (m, 3H), 2.16 (s, 3H), 2.07-1.96 (m, 2H), 1.87-1.69 (m, 6H), 1.62 (s, 3H), 1.02 (s, 3H); ESMS m/e: 529.3 $(M + H)^+$; Anal. Calc. for $C_{28}H_{34}F_2N_4O_4$: C, 63.62; H, 6.48; N, 10.60. Found: C, 63.15; H, 6.27; N, 10.48.

Example 75

The synthetic method is the same as described for the synthesis of (4S)-*N*-(3-{4-[3-(acetylamino)phenyl]-1-

piperidinyl}propyl)-4- (3,5-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxamide.

(4S,5R)-N-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-4-(3,4-DIFLUOROPHENYL)-5-METHYL-2-OXO-1,3-OXAZOLIDINE-3-CARBOXAMIDE: 20.5 mg (74% yield); ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.14 (t, 1H, J=5.5 Hz), 7.40 (d, 1H, J=7.8 Hz), 7.37-6.89 (m, 6H), 5.35 (d, 1H, J=7.5 Hz), 5.02-4.93 (m, 1H), 3.41-3.25 (m, 2H), 3.02 (d, 2H, J=10.8 Hz), 2.53-2.37 (m, 3H), 2.16 (s, 3H), 2.07 (m, 2H), 1.89-1.68 (m, 6H), 1.04 (d, 3H, J=6.4 Hz); ESMS m/e: 515.3 $(M + H)^+$; Anal. Calc. for $C_{27}H_{32}F_2N_4O_4$+0.5H_2O: C, 61.94; H, 6.35; N, 10.70. Found: C, 61.90; H, 6.13; N, 10.64.

Example 76

The synthetic method is the same as described for the synthesis of (4S)-*N*-(3-{4-[3-(acetylamino)phenyl]-1-piperidinyl}propyl)-4-(3,5-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxamide.

N-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-4-(4-FLUOROBENZYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXAMIDE: 17.4 mg (65% yield); ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.08 (t, 1H, J=5.6 Hz), 7.4 (d, 1H, J=7.2 Hz), 7.34 (s, 1H), 7.28-7.14 (m, 3H), 7.05-6.95 (m, 3H), 4.69-4.60 (m, 1H), 4.26 (t, 1H, J=8.8 Hz), 4.15 (dd, 1H, J=3.2, 9 Hz), 3.43 (q, 2H, J=6.2 Hz), 3.3 (dm 1H, J=13.6 Hz), 3.04 (dm, 2H, J=11 Hz), 2.87 (dd, 1H, J=9.3, 14.4 Hz), 2.53-2.42 (m, 3H), 2.16 (s, 3H), 2.09-1.99 (m, 2H), 1.87-1.65 (m, 6H); ESMS m/e: 497.3 $(M + H)^+$; Anal. Calc. for $C_{27}H_{33}FN_4O_4$+0.5H_2O: C, 64.14; H, 6.78; N, 11.08. Found: C, 64.26; H, 6.39; N, 11.12.

Example 77

2-METHYL-N-(3-{1-[(3R)-3-(2-NITROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-nitrophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (2.37 mg, 34.5% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.84 (d, 1H), 7.90 (m, 1H), 7.45 (m 1H), 7.30-7.20 (m, 5H), 7.20 (m, 2H), 6.98 (m, 2H), 6.89 (apparent d, 1H, J=7.7 Hz), 5.62 (apparent dd, 1H, J=4.1, 8.9 Hz), 3.10 (m, 2H), 2.60 (m, 2H), 2.53 (m, 2H), 2.30-2.10 (m, 6H), 1.90 (m, 2H), 1.25 (d, 6H, overlapped); ESMS *m/e*: 502.3 $(M + H)^+$.

Example 78

N-(3-{1-[(3S)-3-([1,1'-BIPHENYL]-4-YLOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 4-phenylphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (3.00 mg, 41.2% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1H), 7.48 (m, 2H), 7.40-7.30 (m, 8H), 7.30-7.25 (m, 4H), 6.97 (apparent d, 1H, J=7.6 Hz), 6.91 (apparent d, 2H, J=8.7

Hz), 5.34 (apparent dd, 1H, J=4.4, 8.0 Hz), 3.40 (m, 2H), 2.98 (m, 2H), 2.53 (apparent sept, partially hidden, 1H, J=8.1 Hz), 2.44 (m, 1H), 2.30-2.10 (m, 6H), 1.93 (d, 2H), 1.26 (d, 6H, J=6.9 Hz); ESMS *m/e*: 533.4 (M + H)$^+$.

Example 79

2-METHYL-N-(3-{1-[(3R)-3-(3-NITROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of N-(3-{1-[(3S)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 3-nitrophenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (2.80 mg, 40.8 % yield) as a thick oil: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (dm, 1H), 7.71 (t, 1H, J=1.8 Hz), 7.50-7.40 (m, 2H), 7.40-7.25 (m, 7H), 7.17 (apparent dd, 1H, J=2.4, 8.2), 6.97 (apparent d, 1H, J=7.7 Hz), 5.45 (apparent dd, 1H, J=5.0, 8.1 Hz), 3.45 (m, 2H), 2.89 (m, 2H), 2.53 (apparent sept, partially hidden, 2H, J=8.3 Hz), 2.30-2.10 (m, 6H), 1.92 (m, 2H), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 502.3 (M + H)$^+$.

Example 80

N-(3-{1-[(3S)-3-(2-ETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 2-ethoxyphenol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg,

0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (1.16 mg, 15.5% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1H), 7.52 (s, 1H), 7.40-7.33 (m, 4H), 7.30-7.20 (m, 3H), 6.97 (apparent d, 1H, J=7.7 Hz), 6.88 (m, 2H), 6.68 (m, 2H), 5.21 (m, 1H), 4.11 (q, 2H, J=7.3 Hz), 3.37 (m, 2H), 2.71 (m, 2H), 2.53 (apparent sept, partially hidden, 2H, J=7.6 Hz), 2.30-2.10 (m, 6H), 1.89 (m, 2H), 1.49 (t, 3H, J=7.3 Hz), 1.25 (d, 6H, J=6.8 Hz); ESMS *m/e*: 501.4 $(M + H)^+$.

Example 81

2-METHYL-N-(3-{1-[(3S)-3-(1-NAPHTHYLOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.20 mg, 0.0137 mmol), 1-naphthol (100 mg), triphenylphosphine (30.0 mg, 0.115 mmol) and diethyl azodicarboxylate (7.42 mg, 0.0426 mmol) in THF (0.50 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (4.30 mg, 66.2% yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1H), 7.72 (d, 1H, J=8.5 Hz), 7.59 (d, 1H, J=8.5 Hz), 7.5 (m, 2H), 7.45-7.30 (m, 6H), 7.25 (m, 3H), 7.17 (apparent dd, 1H, J=2.6, 9.0 Hz), 7.01 (apparent d, 1H, J=2.6 Hz), 6.97 (apparent d, 1H, J=7.9 Hz), 5.46 (apparent dd, 1H, J=4.5, 8.1 Hz), 3.12 (m, 2H), 2.61 (m, 2H), 2.53 (apparent sept, partially hidden, 2H, J=7.9 Hz), 2.30-2.10 (m, 6H), 1.90 (m, 2H), 1.25 (d, 6H, J=7.3 Hz, overlapped); ESMS *m/e*: 507.2 $(M + H)^+$.

Example 82

N-(3-{1-[(3S)-3-(1,3-DIOXO-1,3-DIHYDRO-2H-ISOINDOL-2-YL)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE

Step 1:

2-[(1*S*)-3-CHLORO-1-PHENYLPROPYL]-1*H*-ISOINDOLE-1,3(2*H*)-DIONE: According to the general procedure descibed in Srebnik, M.; Ramachandran, P.V.; Brown, H.C. *J. Org. Chem.* **1988**, *53*, 2916-2920, a mixture of phthalimide (0.147 g, 1.0 mmol), (R)-(+)-3-chloro-phenyl-1-propanol (0.171 g, 1.0 mmol), triphenylphosphine (0.262 g, 1.0 mmol) and diethyl azodicarboxylate (0.174 g, 1.0 mmol) in 5.0 mL of THF was stirred at room temperature for 24 h. The reaction mixture was concentrated *in vacuo*. The residue was washed with pentane (x3) and the combined pentane extracts were concentrated and chromatographed (silica with hexanes-EtOAc 8:1 as the eluent) to give the desired product (0.121 g, 50.2 %) as a yellow solid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.82 (apparent dd, 2H, J=2.9 Hz), 7.70 (apparent dd, 2H, J=2.9 Hz), 7.56 (m, 2H), 7.39-7.27 (m, 3H), 5.64 (apparent dd, 1H, J=7.0, 9.2 Hz), 3.57 (m, 2H), 3.05 (m, 1H), 2.82 (apparent sept, 1H, J=7.0 Hz); ESMS *m/e*: 300.13 $(M+H)^+$.

Step 2:

N-(3-{1-[(3S)-3-(1,3-DIOXO-1,3-DIHYDRO-2H-ISOINDOL-2-YL)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of potassium carbonate

(29.2 mg, 0.211 mmol), sodium iodide (47.5 mg, 0.317 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (51.8 mg, 0.211 mmol) 2-[(1*S*)-3-chloro-1-phenylpropyl]-1*H*-isoindole-1,3(2*H*)-dione (63.1 mg, 0.211 mmol) in DMF (5.0 mL) was stirred at 100 °C for 3 hrs, at which time TLC indicated that the reaction was complete. The reaction mixture was poured into water (50 mL) and the aqueous layer was extracted with methylene chloride (3x30 mL). The combined organic extracts were washed with brine (30 mL), dried over $MgSO_4$ and concentrated under reduced pressure. The crude product was purified by Prep-TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to give the desired product (74.1 mg, 77.1 %) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.83 (apparent dd, 2H, J=2.9 Hz), 7.69 (apparent dd, 2H, J=2.9 Hz), 7.56 (apparent dt, 3H, J=2.9, 7.3 Hz), 7.33 (m, 4H), 7.21 (t, 1H, J=7.8 Hz), 7.09 (s, 1H), 6.81 (apparent d, 1H, J=7.8 Hz), 5.49 (apparent dd, 1H, J=5.5, 9.5 Hz), 2.98 (d, 1H, J=9.5 Hz), 2.87 (m, 2H), 2.50 (apparent sept, 1H, J=6.7 Hz), 2.40-2.35 (m, 4H), 1.94 (m, 2H), 1.70-1.50 (m, 4H), 1.25 (d, 6H, J=7.9 Hz); ESMS *m/e*: 510.37 $(M+H)^+$.

Example 83

2-METHYL-N-(3-{1-[(3S)-3-(4-PHENOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE

STEP 1:

4-{[(1*S*)-3-CHLORO-1-PHENYLPROPYL]OXY}-(4-PHENOXY)BENZENE: A mixture of 4-phenoxyphenol (1.86 g, 10.0 mmol), (R)-(-)-3-chloro-phenyl-1-propanol (1.70 g, 10.0 mmol), triphenylphosphine (2.62 g, 10.0 mmol),

diethyl azodicarboxylate (1.57 mL, 10.0 mmol) in 5.0 mL of THF was stirred at room temperature for 24 h. The reaction mixture was concentrated *in vacuo*. The residue was washed with pentane (x3) and the combined pentane extracts were concentrated and chromatographed (silica with hexanes-EtOAc 97:3 as the eluent) to give the desired product as a thick oil which solidified on standing (2.51 g, 75.7 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.4-7.23 (m, 7H), 7.03 (apparent t, 1H, J=7.3 Hz), 6.91 (apparent dm, 2H, J=7.8 Hz), 6.93 (apparent q, 4H, J=7.8 Hz), 5.31 (apparent dd, 1H, J=4.5, 8.6 Hz), 3.82 (m, 1H), 3.62 (apparent quintet, 1H, J=5.6 Hz), 2.47 (m, 1H), 2.20 (m, 1H).

Step 2:

2-METHYL-N-(3-{1-[(3S)-3-(4-PHENOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (65.5 mg, 0.266 mmol), 4-{[(1*S*)-3-chloro-1-phenylpropyl]oxy}-(4-phenoxy)benzene (0.100 mg, 0.296 mmol), potassium carbonate (40.9 mg, 0.296 mmol) and sodium iodide (67.0 mg, 0.444 mmol) in DMF (1.0 mL) at 100 °C for 3 hours. The reaction mixture was poured into water (50 mL) and the aqueous layer was extracted with methylene chloride (3x30 mL). The combined organic extracts were washed with brine (30 mL), dried over $MgSO_4$ and concentrated under reduced pressure. The crude product was purified by Prep-TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to give the desired product (0.109 g, 74.6 %) as a thick oil: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48 (s, 1H), 7.40-7.30 (m, 4H), 7.20-7.10 (m, 6 H), 7.09 (s, 1H), 6.99 (apparent d, 1H,

J=7.8 Hz), 6.98 (apparent t, 1H, J=7.8 Hz), 6.93 (apparent d, 2H, J=8.4 Hz), 6.84 (m, 2H), 5.20 (apparent dd, 1H, J=4.4, 8.5 Hz), 3.03 (m, 2H), 2.51 (m, 4H), 2.24 (apparent sept, 1H, J=7.8 Hz), 2.20-2.10 (m, 3H), 1.90 (m, 4H), 1.25 (d, 6H, J=7.9 Hz); ESMS *m/e*: 549.41 $(M+H)^+$; Anal. Calc. for $C_{36}H_{40}N_2O_3$: C, 78.80; H, 7.35; N, 5.11. Found: C, 78.58; H, 7.48; N, 5.09.

Example 84

***N*-(4-{1-[(3*R*)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE**

Step 1:

1-[(3*R*)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-(4-NITROPHENYL)-1,2,3,6-TETRAHYDROPYRIDINE: A mixture of potassium carbonate (24.0 mg, 0.174 mmol), sodium iodide (39.0 mg, 0.260 mmol), 4-(4-nitrophenyl)-1,2,3,6-tetrahydropyridine (35.4 mg, 0.174 mmol) and 4-{[(1*R*)-3-chloro-1-phenylpropyl]oxy}-1,2-dimethoxybenzene (53.4 mg, 0.174 mmol) in DMF (0.5 mL) was stirred at 100 °C for 3 hrs, at which time TLC indicated that the reaction was complete. The reaction mixture was poured into water (5.0 mL) and the aqueous layer was extracted with methylene chloride (3x30 mL). The combined organic extracts were washed with brine (30 mL), dried over $MgSO_4$ and concentrated under reduced pressure. The crude product was purified by Prep-TLC plates [1:1=hexane:ethyl acetate with 1% NH_3] afforded the product (63.1 mg, 76.6 %) as a yellow oil. The product was used in next reaction without further purification.

Step 2:

4-{1-[(3*R*)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}ANILINE: A 25-mL RB flask, equipped with a hydrogen-filled balloon, was charged with 1-[(3*R*)-3-(3,4-dimethoxyphenoxy)-3-phenylpropyl]-4-(4-nitrophenyl)-1,2,3,6-tetrahydropyridine (63.0 mg, 0.133 mmol), palladium on carbon (5.0 mol-eq%, 0.00665 mmol, 7.04 mg) and ethanol (2.0 mL) at room temperature. After 1 hr the reaction mixture was filtered through a plug of Celite 545 and concentrated under reduced pressure. The crude product (54.1 mg, 89.4%) was used in next reaction without further purification.

STEP 3:

***N*-(4-{1-[(3*R*)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of 4-{1-[(3*R*)-3-(3,4-dimethoxyphenoxy)-3-phenylpropyl]-4-piperidinyl}aniline (5.31 mg, 0.0119 mmol), isobutyryl chloride (2.08 mg, 0.019 mmol), N,N-diisopropylethylamine (8.40 mg, 0.0650 mmol) in methylene chloride (1.0 mL) was stirred at room temperature for 24 hours. The reaction mixture was concentrated and chromatographed using a preparative TLC plate [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to give the product (3.5 mg, 56.5 %) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.38 (d, 1H, J=8.6 Hz), 7.30-7.20 (m, 4H), 7.20(m, 1H), 7.11 (d, 2H, J=8.6 Hz), 7.04 (s, 1H), 6.57 (d, 1H, J=8.3 Hz), 6.44 (d, 1H, J=2.6 Hz), 6.22 (dd, 1H, J=2.6, 8.3 Hz), 5.09 (apparent dd, 1H, J=4.4, 8.1 Hz), 3.72 (s, 3H), 3.70 (s, 3H), 3.08 (m, 2H), 2.57 (m, 2 H), 2.43 (apparent sept, partially hidden, 2H, J=6.8 Hz),

2.30-2.10 (m, 6H), 1.80 (m, 2H), 1.25 (d, 6H, J=7.9 Hz); ESMS *m/e*: 517.3 $(M+H)^+$.

Example 85

N-(3-{1-[(3S)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Into a 25-mL RB-flask was added triphenylphosphine (9.80 mg, 0.0375 mmol), diethyl azodicarboxylate (5.22 mg, 0.0300 mmol), N-(3-{1-[(3R)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 3-hydroxyacetophenone (100 mg) and THF (1.0 mL) at room temperature. The reaction mixture was stirred at room temperature overnight (16 hrs). The solvent was removed under reduced pressure and the residue was purified by preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to afford the desired product (2.73 mg, 39.9%) as a thick oil: ^1H NMR ($CDCl_3$) δ 7.70-7.64 (m, 2H), 7.54 (m, 2H), 7.49-7.44 (m, 6H), 7.25 (m, 1H), 7.05 (d, 1H, J=8.3 Hz), 6.96 (apparent d, 1H, J=7.7 Hz), 5.34 (apparent dd, 1H, J=4.8, 8.2 Hz), 3.15 (m, 2H), 2.67 (m, 2H), 2.52 (s, 3H), 2.53 (apparent sept, partially hidden, 2H, J=7.6 Hz), 2.30-2.10 (m, 6H), 1.89 (m, 2H), 1.25 (d, 6H, J=6.9 Hz); ESMS *m/e*: 499.4 $(M + H)^+$.

Scheme A. Synthesis of tert-Butyl 4-(3-aminophenyl)-1-piperidinecarboxylate

a. n-BuLi, diisopropylamine, THF, $PhN(Tf)_2$, -78 °C to room temperature, 81%

b. 3-aminophenylboronic acid hemisulfate, LiCl, tetrakis-triphenylphosphine -palladium (0), Na_2CO_3, DME-H_2O, reflux, 81%

c. 10% Pd/C, ethanol, H_2, room temperature, balloon method, 84%

Scheme B1. A General Synthesis of the MCH Antagonists



Scheme B2. A General Synthesis of the MCH Antagonists



X = C, S(=O) halide = Cl, Br

Scheme C1. Specific Examples of the Syntheses of the MCH Antagonists





Scheme C2. Specific Examples of the Syntheses of the MCH Antagonists





Scheme D1. Specific Examples of the Syntheses of the MCH Antagonists



Scheme D2. Specific Examples of the Syntheses of the MCH Antagonists



Scheme E: General Synthesis of the MCH Antagonists



a. dioxane, diisopropylethylamine, Bu_4NI, reflux
or DMF, Ki, Na_2CO_3, 90-100 °C
or toluene, 110 °C, 18-crown-6

b. diisopyropylethylamine, dichloromethane

X = S(=O), C

R_1 = Aromatic, substituted aromatic or heterocyclic

R_2 = aliphatic oraromatic

Scheme F. General Synthesis of the MCH Antagonists



If R = $(CH_2)_nCHOH$-Ar, then,





5

Scheme G. General Synthesis of the MCH Antagonists



5

Scheme H: Synthesis of Oxazolidinones



a. NH_3, then TMS-CN; b. HCl in MeOH (room temperature to reflux);
c. LAH, THF, reflux; d. $(BOC)_2O$, chloroform; e. NaH, THF; f. Chiralcel OD column
g. NaH, p-nitrophenyl chloroformate, THF;
h. an amine such as N-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide

Ar = 3,4-difluorophenyl, 3,5-difluorophenyl or 3,4,5-trifluorophenyl

Scheme I: Synthesis of gem-Dialkyl Substituted Oxazolidinones



a. methyl magnesium bromide, THF; b. N,N-carbonyldiimidazole, DCM; c. NaH, THF, p-nitrophenylchloroformate; d. an amine such as N-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide

Scheme J: Synthesis and Chiral Resolution of Oxazolidinones



[a] (a) *t*-BuLi, THF, RCHO (b) $CH_3ONH_2.HCl$, MeOH, 50-68% over 2 steps (c) Boc_2O, $CHCl_3$, >90% (d) NaH,THF, 76-92% (e) separate diastereomers by column chromatography and separate enantiomers by chiral phase HPLC, 10-16% (f) n-BuLi, THF, 4-nitrophenylchloroformate, ~75%
(g) THF, >80%, an amine such as N-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide

Scheme K: Synthesis Oxazolidinones from Amino Acids



a. LAH, THF; b. $(BOC)_2O$, $CHCl_3$; c. NaH, THF; d. p-nitrophenylchloroformate, NaH, THF; h. an amine such as N-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide

Ar = aromatic such as 4-fluorophenyl or 3,4-difluorophenyl

Scheme L: Determination of the Absolute Stereochemistry of the Di-Substituted Oxazolidinones Using Lactic Acid Derivatives



a. pyrrolidine, methanol, heat; b. t-butyldimethylsilyl chloride; c. LAH, ether, reflux
d. $(BOC)_2O$, chloroform; e. NaH, THF; h. silica gel chromatography

For more details, See: Lagu, B.; Wetzel, J. M.; Forray, C.; Patane, M. A.; Bock, M. G. "Determination of the Relative and Absolute Stereochemistry of a Potent α1A Selective Adrenoceptor Antagonist" Bioorg. Med. Chem. Lett. 2000, 10, 2705.

Scheme M



Example



n=2, R1=H, R2=Ph, R3=H
n=5, R1=H, R2=H, R3=5-OMe
n=1, R1=H, R2=Ph, R3=H
n=4, R1=H, R2=H, R3=5-OMe

Scheme N

Example



R1=6-Cl, R2=H
R1=H, R2=4'-tolyl

Scheme P



Example



Scheme O



R1=H, R2=4'-Me

Scheme Q



Example



7a: R1 = p-OMe
7b: R1 = p-F
7c: R1 = p-CO_2Me
7d: R1= p-Me

Scheme R



Example



EXPERIMENTAL SECTION

The following additional abbreviations are used: HOAc, acetic acid; DMF, *N*,*N*-dimethylformamide; EtOAc, ethyl acetate; MeOH, methanol; NMP, 1-methyl-2-pyrrolidinone; TEA, triethylamine; THF, tetrahydrofuran; All solvent ratios are volume/volume unless stated otherwise.

1-(4-METHYLPHENYL)1*H*-INDOLE: A mixture of 1-*H*-indole (58.5 mg, 0.500 mmol), 1-(iodo)-4-methylbenzene (0.218 g, 1.00 mmol), copper powder (32.0 mg, 0.500 mmol), and K_2CO_3 (0.138 g, 1.00 mmol) in 1-methyl-2-pyrrolidinone (1 mL) was heated at 150 °C for 12 h under argon. The resulting mixture was diluted with H_2O (6 mL). The aqueous layer was extracted with CH_2Cl_2 (3 X 10 mL). The combined organic extracts were washed with brine (10 mL), dried over $MgSO_4$, and concentrated *in vacuo*. The residue was purified by preparative TLC using EtOAc/hexane (1:4) to give the desired product (82 mg, 79%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.67 (d, 1H, *J* = 7.7 Hz), 7.52 (d, 1H, *J* = 7.4 Hz), 7.38 (d, 2H, *J* = 8.4 Hz), 7.34-7.29 (m, 3H), 7.21 (t, 1H, *J* = 7.0 Hz), 7.15 (t, 1H, *J* = 7.0 Hz), 6.66 (d, 1H, 3.3 Hz), 2.43 (s, 3H); ESMS *m/e*: 208.0 $(M + H)^+$.

Example 86

***N*-(3-{1-[(6-CHLORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A solution of 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (0.369 g, 1.50 mmol) and 37 wt % aqueous formaldehyde (30.0 mg, 1.50 mmol) in 1 mL of HOAc:dioxane (1:4) was added to 6-chloro-1-*H*-indole (0.152 g, 1.00 mmol) and the reaction mixture was stirred for 12 h at room

temperature. The resulting mixture was diluted with H_2O (10 mL). The aqueous layer was extracted with CH_2Cl_2 (3 X 100 mL). The combined organic extracts were washed with brine (10 mL), dried over $MgSO_4$, and concentrated *in vacuo*. The residue was purified by preparative TLC on silica using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (79 mg, 42%). ^1H NMR (400 MHz, $CDCl_3$) δ 9.14 (s, 1H), 8.04 (s, 1H), 7.52 (t, 2H, *J* = 8.1 Hz), 7.35 (d, 2H, *J* = 13.3 Hz), 7.18 (t, 1H, *J* = 7.9 Hz), 7.09 (dd, 1H, *J* = 1.9, 8.5 Hz), 6.85 (d, 1H, *J* = 7.4 Hz), 5.18 (s, 1H), 4.01 (s, 2H), 2.55 (septet, 1H, *J* = 6.8 Hz), 2.48-2.34 (m, 3H), 2.08-1.95 (m, 4H), 1.78 (d, 2H, *J* = 12.8 Hz), 1.22 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 410.1 $(M + H)^+$.

Example 87

2-METHYL-*N*-[3-(1-{[1-(4-METHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: According to the procedure used for the synthesis of *N*-(3-{1-[(6-chloro-1*H*-indol-3-yl)methyl]-4-piperidinyl}phenyl)-2-methylpropanamide, 1-(4-methylphenyl)-1*H*-indole (0.207 g, 1.00 mmol) provided 2-methyl-*N*-[3-(1-{[1-(4-methylphenyl)-1*H*-indol-3-yl]methyl}-4-piperidinyl)phenyl]propanamide (0.441 g, 78%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.90 (s, 1H), 7.73 (d, 1H, *J* = 7.2 Hz), 7.58-7.51 (m, 2H), 7.43-7.36 (m, 3H), 7.35-7.29 (m, 3H), 7.26-7.15 (m, 3H), 6.89 (d, 1H, *J* = 7.7 Hz), 4.07 (s, 2H), 3.36 (d, 2H, *J* = 11.6 Hz), 2.59-2.39 (m, 6H), 2.55 (sept, 1H, *J* = 6.7 Hz), 2.10-1.98 (m, 2H), 1.83 (d, 2H, *J* = 12.9 Hz), 1.23 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 466.2 $(M + H)^+$.

2-[(1*S*)-3-CHLORO-1-PHENYLPROPYL]-1*H*-ISOINDOLE-1,3(2*H*)-DIONE: Triphenylphosphine (5.25 g, 20.0 mmol) and diethyl azodicarboxylate (3.58 g, 20.0 mmol) were added to a solution of (1*R*)-3-chloro-1-phenyl-1-propanol (3.42 g, 20.0 mmol) and phthalimide (2.94 g, 20.0 mmol) in THF (100 mL). The reaction mixture was stirred for 4 h at room temperature. The solvent was removed under reduced pressure and the residue was triturated with pentane (3 X 50 mL). The combined pentane fractions were concentrated *in vacuo* and the crude product was purified by chromatography on silica using EtOAc/hexane (3:97) to give the desired product (4.40 g, 74%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.82 (d, 1H, *J* = 5.7 Hz), 7.81 (d, 1H, *J* = 5.5 Hz), 7.70 (d, 1H, *J* = 5.4 Hz), 7.69 (d, 1H, *J* = 5.8 Hz), 7.55 (d, 2H, *J* = 7.2 Hz), 7.38-7.28 (m, 3H), 5.64 (dd, 1H, *J* = 6.8, 9.2 Hz), 3.56 (t, 2H, *J* = 6.4 Hz), 3.11-3.02 (m, 1H), 2.85-2.75 (m, 1H); ESMS *m/e*: 300.1 $(M + H)^+$.

***N*-(3-{1-[(3*S*)-3-(1,3-DIOXO-1,3-DIHYDRO-2*H*-ISOINDOL-2-YL)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of 2-[(1*S*)-3-chloro-1-phenylpropyl]-1*H*-isoindole-1,3(2*H*)-dione (4.50 g, 15.0 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (4.26 g, 15.0 mmol), K_2CO_3 (4.16 g, 30.0 mmol), and NaI (3.40 g, 20.0 mmol) in DMF (40 mL) was stirred at 90 °C for 12 hrs. The reaction mixture was diluted with water (50 mL), extracted with CH_2Cl_2 (3 X 50 mL), and the combined organic extracts were washed with brine (50 mL), dried over $MgSO_4$, and concentrated under reduced pressure. The residue was purified by chromatography on silica using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (5.10 g, 74%). ^{1}H NMR (400 MHz,

$CDCl_3$) δ 7.83 (d, 1H, *J* = 5.5 Hz), 7.82 (d, 1H, *J* = 5.5 Hz), 7.71 (d, 1H, *J* = 5.5 Hz), 7.70 (d, 1H, *J* = 5.4 Hz), 7.56 (d, 2H, *J* = 7.1 Hz), 7.35-7.27 (m, 5H), 7.22 (t, 1H, *J* = 7.5 Hz), 7.09 (s, 1H), 6.81 (d, 1H, *J* = 7.8 Hz), 5.49 (dd, 1H, *J* = 5.5, 9.6 Hz), 2.97 (d, 1H, *J* = 10.1 Hz), 2.92-2.82 (m, 2H), 2.44 (sept, 1H, *J* = 6.7 Hz), 2.40-2.29 (m, 3H), 2.00-1.83 (m, 2H), 1.79-1.39 (m, 5H), 1.26 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 510.4 $(M + H)^+$.

***N*-(3-{1-[(3*S*)-3-AMINO-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of *N*-(3-{1-[(3*S*)-3-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (4.60 g, 9.06 mmol) and hydrazine (3.62 g, 72.4 mmol) in ethanol (150 mL) was refluxed for 12 h. The resulting white precipitate was filtered out and the filtrate was concentrated under vacuum. The residue was washed with CH_2Cl_2/EtOAc (1:1, 3 X 50 mL) and the combined organic fractions were concentrated *in vacuo* to give the desired product (2.90 g, 95%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.45 (s, 1H), 7.39-7.30 (m, 6H), 7.29-7.19 (m, 2H), 6.95 (d, 1H, *J* = 7.2), 4.01 (t, 1H, *J* = 6.8 Hz), 3.04 (t, 2H, *J* = 10.6 Hz), 2.62-2.30 (m, 6H), 2.05-1.70 (m, 8H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 380.4 $(M + H)^+$.

Example 88

2-METHYL-*N*-(3-{1-[(3*S*)-3-PHENYL-3-(PROPIONYLAMINO)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: According to the procedure used for the synthesis of *N*-(3-{1-[(3*S*)-3-(acetylamino)-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide, *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (11.0 mg, 0.0280 mmol) and propionyl chloride (3.80 mg, 0.0420 mmol) provided 2-methyl-*N*-(3-{1-[(3*S*)-3-phenyl-3-(propionylamino)propyl]-4-piperidinyl}phenyl)propanamide (12 mg, 97% yield). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.05 (s, 1H), 7.59 (s, 1H), 7.40-7.20 (m, 7H), 6.96 (s, 1H), 5.19-5.12 (m, 1H), 3.18 (d, 1 H, *J* = 12.0 Hz), 2.99 (d, 1H, *J* = 10.4 Hz), 2.93-2.86 (m, 1H), 2.61-2.40 (m, 3H), 2.38-2.23 (m, 3H), 2.19-1.75 (m, 8H), 1.25 (d, 6H, *J* = 6.9 Hz), 1.22-1.08 (m, 3H); ESMS *m/e*: 436.4 $(M + H)^+$.

Example 89

N-{3-[1-((3*S*)-3-{[(4-FLUOROPHENYL)ACETYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: A mixture of N-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (11.0 mg, 0.0280 mmol) and (4-fluorophenyl)acetyl chloride (7.20 mg, 0.0420 mmol) in THF (5 mL) was stirred at room temperature for 4 h. The solvent was removed under reduced pressure and the residue was purified by preparative TLC using Hexane:EtOAc (2:1) to give the desired product (13 mg, 90% yield). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.89 (d, 1H, *J* = 8.4 Hz), 7.59 (s, 1H), 7.31-6.93 (m, 13H), 5.13 (q, 1H, *J* = 6.0 Hz), 3.56 (s, 2H), 3.07 (d, 1H, *J* = 11.7 Hz), 2.91 (d, 1H, *J* = 11.0 Hz), 2.62-2.42 (m, 2H), 2.40-2.30 (m, 1H), 2.12-1.54 (m, 9H), 1.24 (d, 6H, *J* = 6.7 Hz); ESMS *m/e*: 515.3 $(M + H)^+$.

Example 90

N-(3-{1-[3-(1,2-DIPHENYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: A mixture of 1,1-diphenylhydrazine hydrochloride (10.3 mg, 0.0470 mmol), 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide (14.7 mg, 0.0362 mmol), $ZnCl_2$ (14.85 mg, 0.109 mmol), and HOAc (0.5 mL) was heated for 4 h at 80 °C. The resulting crude mixture was diluted with water (10 mL), the aqueous layer was neutralized with saturated K_2CO_3 and extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified by preparative TLC using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product *N*-(3-{1-[3-(1,2-diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide (4.1 mg, 37%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.71-7.65 (m, 1H), 7.42 (d, 1H, *J* = 7.4 Hz), 7.39 (s, 1H), 7.36-7.15 (m, 15H), 6.94 (d, 1H, *J* = 7.8 Hz), 3.12 (d, 2H, *J* = 11.2 Hz), 2.90 (t, 2H, *J* = 7.8 Hz), 2.59-2.45 (m, 3H), 2.19-1.91 (m, 7H), 1.82 (d, 2H, *J* = 13.5 Hz), 1.24 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 555.3 $(M + H)^+$.

Example 91

***N*-(3-{1-[3-(5-METHOXY-2-PHENYL-1*H*-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** According to the procedure used for the synthesis of *N*-(3-{1-[3-(1,2-diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide, 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide (15.6 mg, 38.2 mmol), and 1-(4-methoxyphenyl)hydrazine hydrochloride (8.00 mg,

0.0458 mmol) provided *N*-(3-{1-[3-(5-methoxy-2-phenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide (3.9 mg, 20%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1H), 7.55 (d, 2H, *J* = 7.4 Hz), 7.43-7.39 (m, 3H), 7.38-7.35 (m, 2H), 7.27-7.19 (m, 3H), 7.08 (d, 1H, *J* = 7.4 Hz), 6.94 (d, 1H, *J* = 7.6 Hz), 6.87 (dd, 1H, *J* = 4.0, 6.6 Hz), 3.88 (s, 3H), 3.80-3.69 (m, 1H), 2.99 (d, 2H, *J* = 11.7 Hz), 2.89 (t, 2H, *J* = 7.3), 2.55-2.39 (m, 4H), 2.02-1.88 (m, 3H), 1.82-1.68 (m, 4H), 1.24 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 510.3 $(M + H)^+$.

Example 92

***N*-(3-{1-[4-(5-METHOXY-2-PHENYL-1*H*-INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** According to the procedure used for the synthesis of *N*-(3-{1-[3-(1,2-diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide, 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide (14.3 mg, 0.0339 mmol) and 1-(4-methoxyphenyl)hydrazine hydrochloride (7.10 mg, 0.0407 mmol) provided *N*-(3-{1-[4-(5-methoxy-2-phenyl-1*H*-indol-3-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.8 mg, 33%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.95 (d, 2H, *J* = 7.8 Hz), 7.61-7.15 (m, 11H), 6.97 (d, 1H, *J* = 7.0 Hz), 3.88 (s, 3H), 3.09 (d, 2H, *J* = 11.3 Hz), 2.99 (t, 2H, *J* = 7.0 Hz), 2.55-2.35 (m, 4H), 2.12-1.70 (m, 6H), 1.68-1.52 (m, 2H), 1.48-1.34 (m, 2H), 1.25 (d, 6H, *J* = 6.7 Hz); ESMS *m/e*: 524.3 $(M + H)^+$.

Example 93

2-METHYL-*N*-(3-{1-[(1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: According to the procedure used for the synthesis of *N*-(3-{1-[3-(1,2-

diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide, *N*-{3-[1-(3,3-dimethoxypropyl)-4-piperidinyl]phenyl}-2-methylpropanamide (15.2 mg, 0.0436 mmol) and 1,1-diphenylhydrazine hydrochloride (11.6 mg, 0.0524 mmol) provided 2-methyl-*N*-(3-{1-[(1-phenyl-1*H*-indol-3-yl)methyl]-4-piperidinyl}phenyl)propanamide (11 mg, 56%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, 1H, *J* = 7.8 Hz), 7.57 (d, 1H, *J* = 7.7 Hz), 7.54-7.47 (m, 4H), 7.43-7.32 (m, 4H), 7.25-7.16 (m, 4H), 6.95 (d, 1H, *J* = 7.8 Hz), 3.87 (s, 2H), 2.53-2.47 (m, 2H), 2.21 (dt, 2H, *J* = 3.0, 10.5 Hz), 2.12-1.77 (m, 6H), 1.24 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 451.3 $(M + H)^+$.

Example 94

2-METHYL-*N*-(3-{1-[(4*E*)-4-PHENYL-4-(2-PYRIDINYLHYDRAZONO)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: According to the procedure used for the synthesis of *N*-(3-{1-[3-(1,2-diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide, 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide (8.70 mg, 0.0223 mmol) and 2-hydrazinopyridine (2.92 mg, 0.0268 mmol) provided 2-methyl-*N*-(3-{1-[(4*E*)-4-phenyl-4-(2-pyridinylhydrazono)butyl]-4-piperidinyl}phenyl)propanamide (2.5 mg, 24%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.97 (d, 1H, *J* = 8.6 Hz), 7.85 (d, 1H, *J* = 7.3 Hz), 7.64-7.27 (m, 9H), 7.09 (d, 1H, *J* = 8.0 Hz), 6.97 (d, 1H, *J* = 8.4 Hz), 6.73 (q, 1H, *J* = 6.6 Hz), 3.52-3.48 (m, 2H), 3.20-3.10 (m, 2H), 2.85-1.75 (m, 13H), 1.26 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 484.4 $(M + H)^+$.

Example 95

***N*-(3-{1-[3-(5-METHOXY-1*H*-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** According to the procedure used for the synthesis of *N*-(3-{1-[3-(1,2-diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide, *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide (23.5 mg, 0.0628 mmol) and 1-(4-methoxyphenyl)hydrazine hydrochloride (13.2 mg, 0.0774 mmol) provided *N*-(3-{1-[3-(5-methoxy-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide (11 mg, 42%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.86 (s, 1H), 7.45 (s, 1H), 7.32 (d, 1H, *J* = 8.4 Hz), 7.28-7.21 (m, 2H), 7.10 (s, 1H), 7.05 (d, 1H, *J* = 2.3 Hz), 7.00-6.91 (m, 2H), 6.85 (dd, 1H, *J* = 2.7, 9.0 Hz), 3.87 (s, 3H), 3.06 (d, 2H, *J* = 11.6 Hz), 2.75 (t, 2H, *J* = 7.2 Hz), 2.55-2.42 (m, 4H), 2.08-1.90 (m, 4H), 1.88-1.74 (m, 4H), 1.25 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 434.2 $(M + H)^+$.

***TERT*-BUTYL 4-[3-(PROPIONYLAMINO)PHENYL]-1-PIPERIDINECARBOXYLATE:** Propionyl chloride (5.53 g, 0.0597 mol) was added dropwise to a solution of *tert*-butyl 4-(3-aminophenyl)-1-piperidinecarboxylate (15.0 g, 0.0543 mol) and TEA (16.5 g, 0.163 mol) in THF (200 mL) and the mixture was stirred at room temperature for 3 h. Water (50 mL) was added to the reaction mixture, the aqueous layer was extracted with CH_2Cl_2 (3 X 100 mL), and the combined organic extracts were washed with brine (50 mL), dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by chromatography on silica using hexane/EtOAc (10:1) to afford the product (18.8 g, 99%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48 (s, 1H),

7.34-7.21 (m, 3H), 6.93 (d, 1H, *J* = 7.4 Hz), 2.77 (t, 2H, *J* = 11.5 Hz), 2.68-2.58 (m, 1H), 2.38 (q, 2H, *J* = 7.6 Hz), 1.87-1.67 (m, 4H), 1.67-1.54 (m, 2H), 1.48 (s, 9H), 1.25 (t, 3H, *J* = 7.5 Hz); ESMS *m/e*: 333.4 (M + H)$^+$.

***N*-[3-(4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Into a stirred solution of *tert*-butyl 4-[3-(propionylamino)phenyl]-1-piperidinecarboxylate (18.8 g, 0.0543 mmol) in dioxane (100 mL) at 5 °C was bubbled HCl gas for 2 h. The solvent was removed *in vacuo*, the residue was dissolved in water (100 mL) and neutralized by adding 10% KOH aqueous solution. The aqueous layer was extracted (3 X 200 mL) with a mixture of $CHCl_3$/isopropyl alcohol (3:1), and the combined organic layers were washed with brine (100 mL), dried over Na_2SO_4, filtered and concentrated *in vacuo*. The residue was purified by column chromatography on silica using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to afford the desired product (12.6 g, 99%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.44 (s, 1H), 7.32 (d, 1H, *J* = 7.2 Hz), 7.28-7.21 (m, 1H), 7.09 (s, 1H), 6.97 (d, 1H, *J* = 7.6 Hz), 3.18 (d, 2H, *J* = 12.6 Hz), 2.73 (dt, 2H, *J* = 2.2, 11.2 Hz), 2.65-2.57 (m, 1H), 2.38 (q, 2H, *J* = 7.4 Hz), 1.83 (d, 2H, *J* = 12.1 Hz), 1.70-1.61 (m, 3H), 1.25 (t, 3H, *J* = 7.5 Hz); ESMS *m/e*: 233.1 (M + H)$^+$.

***TERT*-BUTYL 4-{3-[(CYCLOPROPYLCARBONYL)AMINO]PHENYL}-1-PIPERIDINECARBOXYLATE:** According to the procedure used for the synthesis of *tert*-butyl 4-[3-(propionylamino)phenyl]-1-piperidinecarboxylate, *tert*-butyl 4-(3-aminophenyl)-1-piperidinecarboxylate (16.47 g 0.0596 mol) and cyclopropanecarbonyl chloride (6.27 g,

0.0597 mol) provided the *tert*-butyl 4-{3-[(cyclopropylcarbonyl)amino]phenyl}-1-piperidinecarboxylate (18.1 g, 100%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.55-7.46 (m, 2H), 7.29-7.21 (m, 2H), 6.96-6.89 (m, 1H), 2.79 (t, 2H, *J* = 12.1 Hz), 2.68-2.58(m, 1H), 1.84 (d, 2H, *J* = 12.6 Hz), 1.83-1.76 (m, 4H), 1.48 (s, 9H), 1.19-1.12 (m, 1H), 1.09-1.05 (m, 2H), 0.89-0.75 (m, 2H); ESMS *m/e*: 345.5 $(M + H)^+$.

N-[3-(4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE: According to the procedure used for the synthesis of *N*-[3-(4-piperidinyl)phenyl]propanamide, *tert*-butyl 4-{3-[(cyclopropylcarbonyl)amino]phenyl}-1-piperidinecarboxylate (18.9 g, 0.0543 mol) provided *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide (13.2 g, 100 %). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1H), 7.36-7.22 (m, 3H), 7.23 (d, 1H, *J* = 6.9 Hz), 3.17 (d, 2H, *J* = 11.9 Hz), 2.72 (dt, 2H, *J* = 2.6, 12.2 Hz), 2.65-2.55 (m, 1H), 1.82 (d, 2H, *J* = 13.9 Hz), 1.63 (dt, 3H, *J* = 4.1, 12.5 Hz), 1.53-1.45 (m, 1H), 1.11-1.06 (m, 2H), 0.87-0.81 (m, 2H); ESMS *m/e*: 245.03 $(M + H)^+$.

1-(6-CHLOROHEXYL)-1*H*-INDOLE: To a mixture of NaH (0.249 g, 10.0 mmol) in DMF (5 mL) at 0 °C was added a solution of 1-*H*-indole (0.585 g, 5.00 mmol) in DMF (2 mL). The reaction mixture was stirred for 30 minutes and warmed up to room temperature. Then 1-bromo-6-chlorohexane (0.998 g, 5.00 mmol) was added dropwise by syringe and the reaction mixture was stirred overnight. The reaction mixture was diluted with EtOAc (30 mL), washed with water (3 X 10 mL), dried over $MgSO_4$, concentrated *in vacuo* and purified by chromatography using hexane/EtOAc (97.5:2.5) to give the desired product (0.900 g, 76 %).

^{1}H NMR ($CDCl_3$) δ 7.76-7.54 (m, 1H), 7.47-6.96 (m, 4H), 6.60-6.34 (m, 1H), 4.13 (t, 2H, *J* = 6.8 Hz), 3.50 (t, 2H, *J* = 5.6 Hz), 1.98-1.79 (m, 2H), 1.79-1.64 (m, 2H), 1.54-1.17 (m, 4H).

1-(5-CHLOROPENTYL)-1*H*-INDOLE: According to the procedure used for the synthesis of 1-(6-chlorohexyl)-1*H*-indole, 1-*H*-indole (0.585 g, 5.00 mmol) and 1-bromo-5-chloropentane (0.928 g, 5.00 mmol) gave the desired product (0.890 g, 80%). ^{1}H NMR ($CDCl_3$) δ 7.76-7.51 (m, 1H), 7.44-6.96 (m, 4H), 6.60-6.38 (m, 1H), 4.11 (t, 2H, *J* = 6.8 Hz), 3.47 (t, 2H, *J* = 6.4 Hz), 1.97-1.79 (m, 2H), 1.79-1.61 (m, 2H), 1.58-1.32 (m, 2H).

Example 96

***N*-(3-{1-[6-(1*H*-INDOL-1-YL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** 1-(6-Chlorohexyl)-1*H*-indole (23.6 mg, 0.100 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (24.6 mg, 0.100 mmol), K_2CO_3 (27.6 mg, 0.200 mmol), NaI (22.5 mg, 0.150 mmol) and DMF (1.00 mL) were combined and stirred overnight at 100 °C. The reaction mixture was cooled to room temperature and the crude material was purified by preparative TLC using 5 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product as a yellow solid (40 mg, 90%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.08-6.52 (m, 11H), 4.17 (t, 2H, *J* = 7.2 Hz), 3.26 (d, 2H, *J* = 11.6 Hz), 2.74-2.52 (m, 4H), 2.44-2.28 (m, 2H), 2.20-2.02 (m, 2H), 1.98-1.82 (m,4H), 1.78-1.62 (m, 2H), 1.43-1.28 (m, 4H), 1.28 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 446.5 $(M + H)^+$.

Example 97

***N*-(3-{1-[5-(1*H*-INDOL-1-YL)PENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared as above, using 1-(5-chloropentyl)-1*H*-indole (22.2 mg, 0.100 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (24.6 mg, 0.100 mmol), K_2CO_3 (27.6 mg, 0.200 mmol), NaI (23.0 mg, 0.150 mmol) and DMF (1.00 mL), giving the desired product as a yellow oil (36 mg, 81%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.08-6.52 (m, 11H), 4.19 (t, 2H, *J* = 7.2 Hz), 3.26-3.10 (m, 2H), 2.71-2.55 (m, 2H), 2.55-2.42 (m, 2H), 2.35-2.12 (m, 2H), 2.12-1.80 (m, 6H), 1.80-1.57 (m, 2H), 1.51-1.34 (m, 2H), 1.31 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 432.2 $(M + H)^+$.

Example 98

***N*-(4-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** According to the procedure used for the synthesis of *N*-(3-{1-[4-(4-CHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE (Example 108) *N*-(3-{1-[3-(1,2-diphenyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide, 9-ethyl-9*H*-carbazole-3-carbaldehyde (22.3 mg, 0.100 mmol) and 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide (24.6 mg, 0.100 mmol) provided *N*-(4-{1-[(9-ethyl-9*H*-carbazol-3-yl)methyl]-4-piperidinyl}phenyl)-2-methylpropanamide. The product was obtained as a white crystalline solid (20 mg, 44%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.21-7.09 (m, 12H), 4.38 (q, 2H, *J* = 7.2 Hz), 3.81 (s, 2H), 3.25-3.03 (m, 2H), 2.60-2.38 (m, 2H), 2.31-2.09 (m, 2H), 1.98-1.69 (m, 4H), 1.44 (t, 3H, *J* = 7.2 Hz), 1.23 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 454.3 $(M + H)^+$.

Example 99

***N*-(3-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** According to the procedure used for the synthesis of *N*-(3-{1-[4-(4-CHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE (Example 108) *N*-(4-{1-[(9-ethyl-9*H*-carbazol-3-yl)methyl]-4-piperidinyl}phenyl)-2-methylpropanamide, 9-ethyl-9*H*-carbazole-3-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide afforded *N*-(3-{1-[(9-ethyl-9*H*-carbazol-3-yl)methyl]-4-piperidinyl}phenyl)-2-methylpropanamide (37 mg, 95%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.24-6.29 (m, 12H), 4.37 (q, 2H, *J* = 7.2 Hz), 3.82 (s, 2H), 3.23-3.06 (m, 2H), 2.65-2.38 (m, 2H), 2.31-2.11 (m, 2H), 2.01-1.73 (m, 4H), 1.43 (t, 3H, *J* = 7.2 Hz), 1.25 (d, 6H, *J* = 4.0 Hz); ESMS *m/e*: 454.3 $(M + H)^+$.

Example 100

***N*-[3-(1-{[1-(4-METHOXYPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** According to the procedure used for the synthesis of 1-(4-methylphenyl)1*H* -indole, *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide (37.5 mg, 0.100 mmol) and 1-iodo-4-methoxybenzene (46.8 mg, 0.200 mmol) gave the desired product (27 mg, 56%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.70-6.58 (m, 14H), 3.88 (s, 3H), 3.67 (s, 2H), 3.14-3.01 (m, 2H), 2.57-2.41 (m, 2H), 2.25-2.01 (m, 2H), 1.93-1.69 (m, 4H), 1.24 (d, 6H, *J* = 7.2 Hz); ESMS m/e: 482.2 $(M + H)^+$.

Example 101

***N*-[3-(1-{[1-(4-FLUOROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** According to

the procedure used for the synthesis of 1-(4-methylphenyl)1*H*-indole, *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide (37.5 mg, 0.100 mmol) and 1-fluoro-4-iodobenzene (44.4 mg, 0.200 mmol) gave the desired product (21 mg, 45%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.71-6.60 (m, 14H), 3.69 (s, 2H), 3.19-2.99 (m, 2H), 2.62-2.41 (m, 2H), 2.22-2.07 (m, 2H), 1.94-1.70 (m, 4H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 470.2 $(M + H)^+$.

Example 102

METHYL-4-[5-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-IPERIDINYL}METHYL)-1*H*-INDOL-1-YL]BENZOATE: According to the procedure used for the synthesis of 1-(4-methylphenyl)1*H*-indole, *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide (37.5 mg, 0.100 mmol) and methyl 4-iodobenzoate (52.4 mg, 0.200 mmol) gave the desired product (11 mg, 22%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.31-6.64 (m, 14H), 3.96 (s, 3H), 3.67 (s, 2H), 3.16-2.96 (m, 2H), 2.57-2.41 (m, 2H), 2.18-2.02 (m, 2H), 1.91—1.73 (m, 4H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 510.2 $(M + H)^+$.

Example 103

2-METHYL-*N*-[3-(1-{[1-(3-METHYLPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: According to the procedure used for the synthesis of 1-(4-methylphenyl)1*H*-indole, *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide (37.5 mg, 0.100 mmol) and 1-iodo-3-methylbenzene (43.6 mg, 0.200 mmol) gave the desired product (28 mg, 60%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.68-6.60 (m, 14H), 3.66 (s, 2H), 3.16-2.96 (m, 2H), 2.59-2.44 (m, 2H), 2.44 (s, 3H), 2.18-2.01 (m, 2H),

1.91-1.68 (m, 4H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 466.2 $(M + H)^+$.

Example 104

***N*-{3-[1-(3-{[(4-CHLORO-3-NITROPHENYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** A mixture of *N*-{3[1-(2-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide (10.0 mg, 0.0350 mmol), 4-chloro-3-nitrobenzenesulfonyl chloride (9.90 mg, 0.0380 mmol), and TEA (7.00 mg, 0.0700 mmol) in THF (2 mL) was stirred for 12 h at room temperature. The crude product was purified by preparative TLC (CH_2Cl_2/MeOH/isopropyl amine = 19:1:0.2) to give the desired product (16 mg, 86%). ^1H NMR (400 MHz, $CDCl_3$) δ 8.45-8.38 (m, 1H), 8.02 (d, 1H, *J* = 8.4 Hz), 7.72 (d, 1H, *J* = 8.8 Hz), 7.48-7.40 (m, 3H), 7.29-7.24 (m, 2H), 6.96 (d, 1H, *J* = 7.5 Hz), 3.17-3.09 (m, 4H), 2.63-2.48 (m, 4H), 2.15 (t, 2H, *J* = 11.8 Hz), 1.96-1.72 (m, 6H), 1.25 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 523.2 $(M + H)^+$.

Example 105

***N*-[3-(1-{5-[4-(3,4-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDIN-3-YL]PENTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** A mixture of 3-(5-bromopentyl)-4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one (38.0 mg, 0.110 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (26.0 mg, 0.100 mmol), NaI (23.0 mg, 0.150 mmol), and K_2CO_3 (14.0 mg, 0.100 mmol) in DMF (2 mL) was heated for 1 h at 50°C. The crude product was purified by preparative TLC using CH_2Cl_2/MeOH/isopropyl amine (19:1:0.2) to give the desired product (21 mg, 41%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.49 (s, 1H), 7.39-7.32 (m, 2H), 7.26-7.20 (m, 2H),

7.18-7.11 (m, 1H), 7.10- 7.03 (m, 1H), 6.96 (d, 1H, *J* = 7.6 Hz), 4.80-4.73 (m, 1H), 4.62 (t, 1H, *J* = 7.9 Hz), 4.09-4.04 (m, 1H), 3.51-3.42 (m, 1H), 3.03 (d, 2H, *J* = 11.7 Hz), 2.82-2.72 (m, 1H), 2.51-2.42 (m, 2H), 2.32 (t, 2H, *J* = 7.9 Hz), 2.11 (s, 1H), 2.03-1.97 (m, 2H), 1.85-1.70 (m, 4H), 1.49 (m, 4H), 1.31-1.27 (m, 1H), 1.24 (d, 6H, *J* = 6.9 Hz); ESMS *m/e*: 514.4 $(M + H)^+$.

Example 106

3-(2,6-DICHLOROPHENYL)-*N*-(5-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PENTYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: A mixture of 3-(2,6-dichlorophenyl)-4-formyl-5-isoxazolecarbonyl chloride (69.0 mg, 0.250 mmol), *N*-{3-[1-(5-aminopentyl)-4-piperidinyl]phenyl}-2-ethylpropanamide (44.0 mg, 0.150 mmol), TEA (30.0 mg, 0.300 mmol) in THF (2 mL) was stirred for 12 h at room temperature. The crude product was purified by preparative TLC using CH_2Cl_2/MeOH/ isopropyl amine (19:1:0.2) to give the desired product (52 mg, 67%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.52-7.49 (m, 2H), 7.49-7.41 (m, 2H), 7.39-7.31 (m, 2H), 7.29-7.21 (m, 2H), 6.92 (d, 1H, *J* = 7.6 Hz), 3.25-3.11 (m, 5H), 2.81-2.74 (m, 4H), 2.58-2.44 (m, 4H), 2.30-2.19 (m, 2H), 1.93- 1.78 (m, 4H), 1.56-1.44 (m, 2H), 1.31-1.28 (m, 2H), 1.24 (d, 6H, *J* = 6.6 Hz); ESMS *m/e*: 585.2 $(M + H)^+$.

Example 107

***N*-[3-(1-{2-[(DIPHENYLACETYL)AMINO]ETHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** A mixture of *N*-{3[1-(2-aminoethyl)-4-piperidinyl]phenyl}-2-methylpropanamide (20.0 mg, 0.0700 mmol), diphenylacetyl chloride (23.0 mg, 0.110 mmol), and TEA (20.0 mg, 0.140 mmol) in THF (2 mL) was stirred overnight at 23 °C. The

crude product was purified by preparative TLC using CH_2Cl_2/MeOH/isopropyl amine (19:1:0.2) to give the desired product (8.0 mg, 47%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.53 (s, 1H), 7.37-7.20 (m, 13H), 6.97-6.92 (m, 1H), 6.67 (s, 1H), 4.98 (s, 1H), 3.43 (q, 2H, *J* = 5.9 Hz), 2.90 (d, 2H, *J* = 11.6 Hz), 2.57-2.42 (m, 4H), 2.11 (t, 2H, *J* = 10.4 Hz), 1.75 (d, 2H, *J* = 12.4 Hz), 1.70-1.58 (m, 2H), 1.25 (d, 6H, *J* = 6.7 Hz); ESMS *m/e*: 484.2 $(M + H)^+$.

Example 108

***N*-(3-{1-[4-(4-CHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** 4-(4-chlorophenoxy)benzaldehyde (0.119 g, 0.510 mmol) and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (0.126 g, 0.510 mmol) were mixed in 1,2-dichloroethane (5 mL) and then treated with sodium triacetoxyborohydride (0.424 g, 2.00 mmol) and HOAc (0.03 mL, 0.5 mmol). The mixture was stirred overnight at room temperature. The reaction mixture was neutralized with saturated $NaHCO_3$ aqueous solution and the aqueous layer was extracted with CH_2Cl_2 (3 X 10 mL). The combined organic layers were washed with brine, dried over $MgSO_4$, concentrated *in vacuo*, and purified by preparative TLC using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (53 mg, 23%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.50 (s, 1H), 7.34-7.19 (m, 7H), 6.98-6.87 (m, 5H), 3.50 (s, 2H), 2.98 (d, 2H, *J* = 11.8 Hz), 2.58-2.44 (m, 2H), 2.10-1.98 (m, 2H), 1.83-1.76 (m, 4H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 463.2 $(M + H)^+$.

Example 109

***N*-{3-[1-({2,5-DIMETHYL-1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-PYRROL-3-YL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by the procedure described in example 108, substituting 2,5-dimethyl-1-[3-(trifluoromethyl)phenyl]-1*H*-pyrrole-3-carbaldehyde (0.136 g, 0.510 mmol) for 4-(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.69-7.56 (m, 2H), 7.53-7.32 (m, 4H), 7.28-7.18 (m, 2H), 6.99 (s, 1H), 5.98 (s, 1H), 3.43 (s, 2H), 3.16-3.06 (m, 2H), 2.57-2.42 (m, 2H), 2.07-1.95 (m, 8H), 1.89-1.76 (m, 4H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 498.2 $(M + H)^+$.

Example 110

***N*-(3-{1-[4-(3,4-DIFLUOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by the procedure described in example 108, substituting 4-(3,4-difluorophenoxy)benzaldehyde (0.119 g, 0.510 mmol) for 4-(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.52 (s, 1H), 7.32 (d, 2H, *J* = 8.4 Hz), 7.28-7.21 (m, 2H), 7.14-7.06 (m, 2H), 6.98-6.94 (m, 3H), 6.86-6.79 (m, 1H), 6.76-6.69 (m, 1H), 3.51 (s, 2H), 2.99 (d, 2H, *J* = 11.7 Hz), 2.55-2.44 (m, 2H), 2.12-2.02 (m, 2H), 1.86-1.74 (m, 4H), 1.25 (d, 6H, *J* = 7.0 Hz); ESMS *m/e*: 465.2 $(M + H)^+$.

Example 111

***N*-(3-{1-[(5-CHLORO-3-METHYL-1-PHENYL-1*H*-PYRAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by the procedure described in example 108, substituting 5-chloro-3-methyl-1-phenyl-1*H*-pyrazole-4-carbaldehyde (0.113 g, 0.510 mmol) for 4-(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62-7.19 (m, 9H), 6.97 (s, 1H), 3.43 (s, 2H), 3.08-2.98

211

(m, 2H), 2.58-2.43 (m, 2H), 2.39-2.32 (m, 3H), 2.18-1.71 (m, 6H), 1.24 (d, 6H, J = 6.9 Hz); ESMS m/e: 451.2 $(M + H)^+$.

Example 112

***N*-(3-{1-[4-(3,4-DICHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by the procedure described in example 108, substituting 4-(3,4-dichlorophenoxy)benzaldehyde (0.136 g, 0.510 mmol) for 4-(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.53 (s, 1H), 7.36-7.18 (m, 6H), 7.08 (d, 1H, J = 1.8 Hz), 6.96 (d, 3H, J = 6.8 Hz), 6.84 (dd, 1H, J = 2.8, 8.9 Hz), 3.51 (s, 2H), 2.99 (d, 2H, J = 11.5 Hz), 2.55-2.42 (m, 2H), 2.12-2.02 (m, 2H), 1.84-1.73 (m, 4H), 1.24 (d, 6H, J = 7.0 Hz); ESMS m/e: 497.1 $(M + H)^+$.

Example 113

2-METHYL-*N*-(3-{1-[(2-PHENYL-1*H*-IMIDAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by the procedure described in example 108, substituting 2-phenyl-1*H*-imidazole-4-carbaldehyde (88.0 mg, 0.510 mmol) for 4-(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (d, 2H, J = 7.4 Hz), 7.65-7.31 (m, 6H), 7.28-7.18 (m, 2H), 7.12-7.05 (m, 1H), 6.95-6.88 (m, 1H), 3.69 (s, 2H), 3.17-3.05 (m, 2H), 2.62-2.45 (m, 2H), 2.28-2.18 (m, 2H), 1.88-1.70 (m, 4H), 1.25 (d, 6H, J = 6.8 Hz); ESMS m/e: 403.2 $(M + H)^+$.

Example 114

***N*-(3-{1-[4-(DIPHENYLAMINO)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by the procedure described in example 108, substituting 4-(diphenylamino)benzaldehyde (0.139 g, 0.510 mmol) for 4-

(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.49 (s, 1H), 7.39-6.92 (m, 18H), 3.49 (s, 2H), 3.02-2.99 (m, 2H), 2.59-2.43 (m, 2H), 2.15-2.03 (m, 2H), 1.92-1.76 (m, 4H), 1.23 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 504.2 $(M + H)^+$.

Example 115

N-[3-(1-{[4-BROMO-1-(4-CHLOROBENZYL)-1*H*-PYRAZOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by the procedure described in example 108, substituting 4-bromo-1-(4-chlorobenzyl)-1*H*-pyrazole-5-carbaldehyde (0.153 g, 0.510 mmol) for 4-(4-chlorophenoxy)benzaldehyde. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (s, 1H), 7.36 (d, 1H, *J* = 8.8 Hz), 7.34-7.30 (m, 3H), 7.29-7.26 (m, 1H), 7.22 (t, 1H, *J* = 7.8 Hz), 7.16 (d, 2H, *J* = 8.6 Hz), 6.95 (d, 1H, *J* = 7.5 Hz), 5.24 (s, 2H), 3.61 (s, 2H), 3.09 (d, 2H, *J* = 11.9 Hz), 2.55-2.42 (m, 2H), 2.19 (dt, 2H, *J* = 4.4, 11.4 Hz), 1.89-1.76 (m, 4H), 1.24 (d, 6H, *J* = 6.7 Hz); ESMS *m/e*: 529.1 $(M + H)^+$.

1-(3-[{(1*R*)-3-CHLORO-PHENYLPROPYL]OXY}PHENYL)ETHANONE:

Azodicarboxylate (5.37 g, 0.0310 mol) was added to a solution of triphenylphosphine (8.09 g, 0.0308 mol), 1*S*-3-chloro-1-phenyl-1-propanol (4.20 g, 0.031 mol) and, 1-(3-hydroxyphenyl)ethanone in THF (150 mL). The reaction mixture was stirred for 4 days at 23 °C. The solvent was removed under reduced pressure and the residue was triturated with ether/hexane (1:2, (3 X 100 mL). The combined organic fractions were concentrated *in vacuo* and the crude product was purified by chromatography using EtOAc/hexane (1:14) to give the desired product (6.55 g, 74%). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48-7.31 (m,

6H), 7.26 (t, 2H, *J* = 8.2 Hz), 7.04 (d, 1H, *J* = 8.1 Hz), 5.44 (dd, 1H, *J* = 4.4, 8.1 Hz), 3.83-3.74 (m, 1H), 3.63-3.56 (m, 1H), 2.51 (s, 3H), 2.51-2.45 (m, 1H), 2.29-2.17 (m, 1H); ESMS *m/e*: 289.0 $(M + H)^+$.

Example 116

***N*-(3-{1-[(3*R*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of 1-(3-{[(1*R*)-3-chloro-1-phenylpropyl]oxy}phenyl)ethanone (58.5 mg, 0.200 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (56.8 mg, 0.200 mmol), NaI (34.0 mg, 0.200 mmol) and K_2CO_3 (55.5 mg, 0.400 mmol) in DMF (1 mL) was stirred at 100 °C for 3 h. The solvent was removed under reduced pressure and the residue was purified by chromatography on silica using 5 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (98 mg, 98%). ^1H NMR (400 MHz, $CDCl_3$) δ 8.01 (s, 1H), 7.49-7.21 (m, 11H), 7.09-7.03 (m, 1H), 6.96 (d, 1H, *J* = 7.9 Hz), 5.32 (dd, 1H, *J* = 5.0, 7.9 Hz), 3.08-2.98 (m, 2H), 2.57-2.43 (m, 6H), 2.11-1.72 (m, 9H), 1.25 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 499.4 $(M + H)^+$.

Procedures:

Procedure A (see also example 48)

***N*-(3-{1-[(3*R*)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:**



Method A

4-{[(1*R*)-3-CHLORO-1-PHENYLPROPYL]OXY}-1,2-DIMETHOXYBENZENE: A mixture of 3,4-dimethoxyphenol (4.07 g, 26.4 mmol), (*S*)-(-)-3-chloro-phenyl-1-propanol (4.50 g, 26.4 mmol, 99% ee, Aldrich Chemical Co.), triphenylphosphine (6.92 g, 26.4 mmol) and diethyl azodicarboxylate (4.59 g, 26.4 mmol) in THF (110 mL) was stirred at room temperature for 24 h. The reaction mixture was concentrated *in vacuo*. At this point, the residue can either be washed with pentane and the combined pentane extracts were concentrated and chromatographed with hexane:EtOAc (8:1) as the eluent to give the desired product (as described as a general procedure by: Srebnik, M.; Ramachandran, P.V.; Brown, H.C. *J. Org. Chem.* 1988, *53*, 2916-2920). This procedure was performed on a smaller scale reaction and only a 40% yield of the product was realized.

Alternatively, on a larger scale (26.4 mmol), the crude product was triturated with a small amount of dichloromethane and the precipitated triphenylphosphine oxide was filtered. The filtrate was concentrated and the crude product was chromatographed to give the desired product as a thick yellow oil (7.30 g, 88.9% yield): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39-7.32 (m, 4H), 7.20 (m, 1H), 6.64 (d, 1H, J = 8.7 Hz), 6.51 (d, 1H, J = 2.7 Hz), 6.30 (dd, 1H, J = 2.7, 8.7 Hz), 5.27 (apparent dd, 1H, J = 4.5, 8.7 Hz), 3.79 (s, 3H), 3.77 (s, 3H), 3.61 (m, 1H), 2.45 (m, 1 H), 2.20 (m, 1H), 1.80 (s, 1H); ESMS m/e: 307.1 $(M + H)^+$.

N-(3-{1-[(3R)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:

A mixture of potassium carbonate (321 mg, 2.32 mmol), sodium iodide (522 mg, 3.48 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (570 mg, 2.32 mmol) and 4-{[(1*R*)-3-chloro-1-phenylpropyl]oxy}-1,2-dimethoxybenzene (712 mg, 2.32 mmol) in DMF (5.00 mL) was stirred at 100 °C for 3 h, at which time TLC indicated that the reaction was complete. The reaction mixture was poured into water (50 mL) and the aqueous layer was extracted with methylene chloride (3 x 30 mL). The combined organic extracts were washed with brine (30 mL), dried over $MgSO_4$ and concentrated under reduced pressure. The crude product was purified by Preparatory TLC [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to afford the product (970 mg, 90.1%) as a thick oil.

Method B

Into a 25-mL RB-flask was added triphenylphosphine (9.80 mg, 0.0375 mmol), diethyl azodicarboxylate (5.22 mg, 0.0300 mmol), *N*-(3-{1-[(3*S*)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), 3,4-dimethoxyphenol (7.70 mg, 0.0500 mmol) and THF (1.00 mL) at room temperature. The reaction mixture was stirred at room temperature overnight (16 h). The solvent was removed under reduced pressure and the residue was purified by preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] to afford the desired product (4.40 mg, 34.1 % yield) as a thick oil: ^1H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1 H), 7.40-7.30 (m, 4H), 7.25 (m, 3H), 6.97 (d, 1H, J = 7.8 Hz), 6.64 (d, 1H, J = 9.1 Hz), 6.51 (d, 1H, J = 2.6 Hz), 6.29 (d, 1H, J = 2.6, 9.1 Hz), 5.20 (apparent dd, 1H, J = 4.4, 8.5 Hz), 3.80

(s, 3H), 3.77 (s, 3H), 3.23 (m, 2H), 2.77 (m, 2H), 2.5 (m, 2H), 2.3-2.1 (m, 6H), 1.80 (m, 2H), 1.25 (d, 6H, J = 7.9 Hz); ESMS *m/e*: 517.4 $(M + H)^+$.

Procedure B (see also example 49)

2-METHYL-*N*-(3-{1-[(3*S*)-3-PHENOXY-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of *N*-(3-{1-[(3*R*)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide (9.53 mg, 0.0250 mmol), phenol (4.70 mg, 0.050 mmol), triphenylphosphine (9.80 mg, 0.0375 mmol) and diethyl azodicarboxylate (5.22 mg, 0.0300 mmol) in THF (1.00 mL) was stirred at room temperature for 3 days. Chromatography using silica preparative TLC plates [2.5% of NH_3 (2.0 M in methanol) in $CHCl_3$] gave the desired product (2.70 mg, 23.6 % yield) as a thick oil: ^1H NMR δ 7.46 (s, 2H), 7.40-7.30 (m, 4H), 7.25 (m, 3H), 7.20 (m, 2H), 6.97 (apparent d, 1H, J = 7.4 Hz), 6.89 (apparent tt, 1H, J = 0.8, 7.6 Hz), 6.84 (apparent dt, 1H, J = 0.8, 8.0 Hz), 5.20 (apparent dd, 1H, J = 4.4, 8.5 Hz), 3.35 (m, 2H), 2.91 (m, 2H), 2.60 (m, 2H), 2.30-2.10 (m, 6H), 1.90 (m, 2H), 1.25 (d, 6H, J = 7.9 Hz); ESMS *m/e*: 457.4 $(M + H)^+$;

Procedure C

Scheme O



R_1=H, R_2=4'-Me

1-(4-METHYLPHENYL)1H-INDOLE: A mixture of 1-H-indole (58.5 mg, 0.500 mmol), 1-iodo-4-methylbenzene (0.218 g, 1.00 mmol), copper powder (32.0 mg, 0.500 mmol), and K_2CO_3 (0.138 g, 1.00 mmol) in 1-methyl-2-pyrrolidinone (1.00 mL) was heated at 150 °C for 12 h under argon. The resulting mixture was diluted with H_2O (6 mL). The aqueous layer was extracted with CH_2Cl_2 (3 X 10 mL). The combined organic extracts were washed with brine (10 mL), dried over $MgSO_4$, and concentrated *in vacuo*. The residue was purified by preparative TLC using EtOAc:hexane (1:4) to give the desired product (82.0 mg, 79.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.67 (d, 1H, J = 7.7 Hz), 7.52 (d, 1H, J = 7.4 Hz), 7.38 (d, 2H, J = 8.4 Hz), 7.34-7.29 (m, 3H), 7.21 (t, 1H, J = 7.0 Hz), 7.15 (t, 1H, J = 7.0 Hz), 6.66 (d, 1H, J = 3.3 Hz), 2.43 (s, 3H); ESMS *m/e*: 208.0 $(M + H)^+$.

Procedure D **(see also example 86)**

Scheme N



Example



R_1=6-Cl, R_2=H
R_1=H, R_2=4'-tolyl

***N*-(3-{1-[(6-CHLORO-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A solution of 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (0.369 g, 1.50 mmol) and 37 wt % aqueous formaldehyde (30.0 mg, 1.50 mmol) in 1.00 mL of HOAc:dioxane (1:4) was added to 6-chloro-1-H-indole (0.152 g, 1.00 mmol) and the reaction mixture was stirred for 12 h at room temperature. The resulting mixture was diluted with H_2O (10 mL). The aqueous layer was extracted with CH_2Cl_2 (3 X 100 mL). The combined organic extracts were washed with brine (10 mL), dried over $MgSO_4$, and concentrated *in vacuo*. The residue was purified by preparative TLC plates using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (79.0 mg, 42.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 9.14 (s, 1H), 8.04 (s, 1H), 7.52 (t, 2H, J = 8.1 Hz), 7.35 (d, 2H, J = 13.3 Hz), 7.18 (t, 1H, J = 7.9 Hz), 7.09 (dd, 1H, J = 1.9, 8.5 Hz), 6.85 (d, 1H, J

= 7.4 Hz), 5.18 (s, 1H), 4.01 (s, 2H), 2.55 (septet, 1H, J = 6.8 Hz), 2.48-2.34 (m, 3H), 2.08-1.95 (m, 4H), 1.78 (d, 2H, J = 12.8 Hz), 1.22 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 410.1 $(M + H)^+$.

Procedure E **(see also example 90)**

Scheme M



Example



n=2, R_1=H, R_2=Ph, R_3=H
n=5, R_1=H, R_2=H, R_3=5-OM
n=1, R_1=H, R_2=Ph, R_3=H
n=4, R_1=H, R_2=H, R_3=5-OM

***N*-(3-{1-[3-(1,2-DIPHENYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of 1,1-diphenylhydrazine hydrochloride (10.3 mg, 0.0470 mmol), 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide (14.7 mg, 0.0362 mmol), $ZnCl_2$ (14.8 mg, 0.109 mmol), and HOAc (0.500 mL) was heated for 4 h at 80 °C. The resulting crude mixture was diluted with water (10 mL), the aqueous layer was neutralized with saturated K_2CO_3 (10 mL) and extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified

by preparative TLC plates using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product *N*-(3-{1-[3-(1,2-diphenyl-1H-indol-3-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide (4.10 mg, 37.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.71-7.65 (m, 1H), 7.42 (d, 1H, J = 7.4 Hz), 7.39 (s, 1H), 7.36-7.15 (m, 15H), 6.94 (d, 1H, J = 7.8 Hz), 3.12 (d, 2H, J = 11.2 Hz), 2.90 (t, 2H, J = 7.8 Hz), 2.59-2.45 (m, 3H), 2.19-1.91 (m, 7H), 1.82 (d, 2H, J = 13.5 Hz), 1.24 (d, 6H, J = 6.9 Hz); ESMS *m/e*: 555.3 $(M + H)^+$.

Procedure F **(see also example 108)**

Scheme R



Example



***N*-(3-{1-[4-(4-CHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A solution of 4-(4-chlorophenoxy)benzaldehyde (0.119 g, 0.510 mmol) and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (0.126 g, 0.510 mmol) in 1,2-dichloroethane (5.00 mL)

was treated with sodium triacetoxyborohydride (0.424 g, 2.00 mmol) and HOAc (0.0300 mL, 0.500 mmol) at room temperature. The mixture was stirred overnight at room temperature. The reaction mixture was neutralized with saturated $NaHCO_3$ aqueous solution (10 mL) and the aqueous layer was extracted with CH_2Cl_2 (3 x 10 mL). The combined organic layers were washed with brine, dried over $MgSO_4$, concentrated *in vacuo* and purified by preparative TLC plates using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (53.0 mg, 23.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.50 (s, 1H), 7.34-7.19 (m, 7H), 6.98-6.87 (m, 5H), 3.50 (s, 2H), 2.98 (d, 2H, J = 11.8 Hz), 2.58-2.44 (m, 2H), 2.10-1.98 (m, 2H), 1.83-1.76 (m, 4H), 1.24 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 463.2 $(M + H)^+$.

Procedure G **(see also example 116)**

Scheme F



***N*-(3-{1-[(3R)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of 1-(3-{[(1*R*)-3-chloro-1-phenylpropyl]oxy}phenyl)ethanone (58.5 mg, 0.200 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (56.8 mg, 0.200 mmol), NaI (34.0 mg, 0.200 mmol) and K_2CO_3 (55.5 mg, 0.400 mmol) in DMF (1.00 mL) was stirred at 100 °C for 3 h. The

solvent was removed under reduced pressure and the residue was purified by chromatography on silica using 5 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (98.0 mg, 98.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 8.01 (s, 1H), 7.49-7.21 (m, 11H), 7.09-7.03 (m, 1H), 6.96 (d, 1H, J = 7.9 Hz), 5.32 (dd, 1H, J = 5.0, 7.9 Hz), 3.08-2.98 (m, 2H), 2.57-2.43 (m, 6H), 2.11-1.72 (m, 9H), 1.25 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 499.4 $(M + H)^+$.

Scheme S



Procedure H



2-METHYL-*N*-(3-{1-[3-(1-METHYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: A mixture of *N*-(3-{1-[4-

(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide (100 mg, 0.270 mmol), 1-methyl-1-phenylhydrazine (106 mg, 0.870 mmol), $ZnCl_2$ (119 mg, 0.870 mmol), and HOAc (1.00 mL) was heated for 12 h at 80 °C. The resulting crude mixture was diluted with water (20 mL), the aqueous layer was neutralized with saturated K_2CO_3 solotion (10 mL) and extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified by preparative TLC using 3 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product 2-methyl-*N*-(3-{1-[3-(1-methyl-1H-indol-3-yl)propyl]-4-piperidinyl}phenyl)propanamide (20.7 mg, 18.7 %):

^1H NMR (400 MHz, $CDCl_3$) δ 7.60 (d, 1H, J = 8.1 Hz), 7.45 (s, 1H), 7.35 (d, 1H, J = 7.4 Hz), 7.26-7.24 (m, 4H), 7.09 (t, 1H, J = 7.3 Hz), 6.97 (d, 1H, J = 7.3 Hz), 6.86 (s, 1H), 3.75 (s, 3H), 3.11 (d, 2H, J = 11.6 Hz), 2.79 (t, 2H, J = 7.3 Hz), 2.51-2.50 (m, 4H), 2.12-1.81 (m, 8H), 1.25 (d, 6H, J = 7.1 Hz); Anal. Calcd for $C_{27}H_{35}N_3O+0.225CHCl_3$: C, 73.57; H, 7.99; N, 9.45. Found: C, 73.93; H, 7.90; N, 9.23; ESMS *m/e*: 418.2 $(M + H)^+$.

Procedure I

Scheme T





7-(2-FLUOROPHENYL)-1H-INDOLE: A mixture of 2-fluorophenylboronic acid (83.4 mg, 0.600 mmol), 7-bromo-1*H*-indole (98.0 mg, 0.500 mmol), LiCl (42.0 mg, 1.00 mmol), Na_2CO_3 (2.0 M, 0.100 mL), $Pd(PPh_3)_4$ (115 mg, 0.100 mmol) and DME (2.00 mL) was heated at 75 °C for 12 h under Argon. The resulting crude mixture was diluted with water (40 mL), the aqueous layer was extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were washed with brine (30 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The residue was purified by preparative TLC using hexane:EtOAc (8:1) to give the desired product 7-(2-fluorophenyl)-1*H*-indole (108 mg, 100 %): ^1H NMR (400 MHz, $CDCl_3$) 8.21 (br s, 1H), 7.71 (dm, 1H, J = 7.3), 7.55 (dt, 1H, J = 7.3, 1.6 Hz), 7.39 (m, 1 H), 7.30-7.19 (m, 5H), 6.62 (dd, 1H, J = 2.1-3.3 Hz); ESMS *m/e*: 211.9 $(M + H)^+$.

Procedure J

Scheme U





5-(4-METHYLPHENOXY)-1H-INDOLE: A mixture of 5-bromo-1*H*-indole (98.0 mg, 0.500 mmol), *p*-cresol (108 mg, 1.00 mmol), Cu (32.0 mg, 0.500 mmol), K_2CO_3 (138 mg, 1.00 mL) and DMF (1.00 mL) was heated at 160 °C for 12 h. The resulting crude mixture was diluted with water (40 mL), the aqueous layer was extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were washed with brine (30 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The residue was purified by preparative TLC using hexane:EtOAc (4:1) to give the desired product 5-(4-methylphenoxy)-1*H*-indole (57.5 mg, 51.5 %): ESMS *m/e*: 224.0 $(M + H)^+$.

Procedure K



Scheme AN



***N*-(3-{1-[7-(2-FLUOROPHENYL)-7-OXOHEPTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A 50-mL round-bottom flask was charged with a solution of 7-chloro-1-oxo-1(2-fluorophenyl)heptane (2.42 g, 10.0 mmol), 2-methyl-*N*-[3-(4-piperidyl)phenyl] propanamide (2.46 g, 10.0 mmol), K_2CO_3 (2.76 g, 20.0 mmol) and NaI (2.25 g, 15.0 mmol) in DMF (25.0 mL). The mixture was stirred for 10 min at 25 °C and then heated at 100 °C for 12 h, cooled to 25 °C and diluted with EtOAc (100 mL). The reaction mixture was washed with water (4 X 50 mL) and the aqueous layer was extracted with EtOAc (100 mL). The organic layers were washed with brine (50 mL), dried over $MgSO_4$, concentrated *in vacuo* and the crude product was purified by chromatography (EtOAc:MeOH 97:3) to give the desired product (3.70 g, 82.0 %).

Procedure L



Scheme AN



***N*-(3-{1-[7-(2-FLUOROPHENYL)-7-HYDROXYHEPTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** To a 50-mL round-bottomed flask charged with *N*-(3-{1-[7-(2-fluorophenyl)-7-oxoheptyl]-4-piperidinyl}phenyl)-2-methylpropanamide (5.0 mmol) and methanol (20 mL) was added $NaBH_4$ (7.5 mmol) at 0 °C in an ice-bath. The reaction mixture was warmed to 25 °C and stirred for 2 h. The reaction was monitored by TLC (EtOAc:MeOH 95:5). If necessary, another 5.0 mmol of $NaBH_4$ was added to the reaction mixture and the reaction mixture was refluxed for 1 h. The reaction was quenched with water (5.0 mL) and diluted with EtOAc (10 mL). The organic layer was separated, washed with saturated $NaHCO_3$ solution (10 mL), dried over $MgSO_4$ and concentrated *in vacuo*. The crude

product was purified by chromatography (EtOAc:MeOH 97:3) to give the desired product (90%).

Procedure M

Scheme A



Step 1: If reacted individually, a solution of the amine or aniline (1.00 eq), diisopropylethylamine or TEA (2.00 eq) and an electrophile (1.50 eq) in CH_2Cl_2 was stirred for 24 h at 23 °C. The solvent was removed *in vacuo* and the crude product was chromatographed (silica) to give the final product.



***TERT*-BUTYL 4-{3-[(4-CHLOROBUTANOYL)AMINO]PHENYL}-1-PIPERIDINECARBOXYLATE** (3.32 g, 87.4 %) was synthesized according to Scheme A and Procedure M: ^1H NMR (400 MHz, $CDCl_3$) δ 7.55 (s, 1H), 7.47 (s, 1H), 7.37 (m, 1H), 7.28 (m, 1H), 6.97 (d, 1H, J = 7.6 Hz), 3.89 (t, 1H, J = 6.4

Hz), 3.74 (m, 2H), 2.79- 2.75 (m, 4H), 2.64 (m, 2H), 1.88-1.77 (m, 4H), 1.60-1.59 (m, 4H), 1.48 (s, 9H).



Step B:

***TERT*-BUTYL 4-[3-(2-OXO-1-PYRROLIDINYL)PHENYL]-1-PIPERIDINECARBOXYLATE:** To a solution of *tert*-butyl 4-[3-(2-oxo-1-pyrrolidinyl)phenyl]-1-piperidinecarboxylate (0.429 g, 16.9 mmol) in dioxane (100 mL) was bubbled HCl gas for 1 h at 25 °C. The resulting crude mixture was basified with 10% KOH solution (100 mL), the aqueous layer was extracted with 3:1 $CHCl_3$:iso-propyl alcohol (3 X 150 mL). The combined organic layers were washed with brine (100 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The residue was purified by preparative TLC using 20% NH_3 (2.0 M in MeOH) in CH_2Cl_2 solution to give the desired product *tert*-butyl 4-[3-(2-oxo-1-pyrrolidinyl)phenyl]-1-piperidinecarboxylate (245 mg, 78.7 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.52 (t, 1H, J = 1.8 Hz), 7.41 (ddd, 1H, J = 8.1, 2.3, 0.9 Hz), 7.30 (t, 1H, J = 7.9 Hz), 7.02 (d, 1H, J = 7.9 Hz), 3.86 (t, 2H, J = 7.3 Hz), 3.21 (dt, 2H, J = 11.9, 2.9 Hz), 2.76 (dt, 2H, J = 12.1, 2.4 Hz), 2.65 (tt, 1H, J = 11.9, 3.5 Hz), 2.61 (t, 2H, J = 8.3 Hz), 2.22 (br s, 1H), 2.16 (qt, 2H, J = 7.5 Hz), 1.85 (d, 2H, J = 12.4 Hz), 1.67 (dq, 2H, J = 12.5, 4.0 Hz).



***TERT*-BUTYL 4-(4-AMINOPHENYL)-1-PIPERIDINECARBOXYLATE:** Available from Arch Chemical Company, NJ.

2-METHYL-*N*-[4-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: To a solution of *tert*-butyl 4-(4-aminophenyl)-1-piperidinecarboxylate (8.20 g, 29.7 mmol) and triethylamine (8.4 mL, 60 mmol) in dry THF (100 mL) at 0 °C was slowly added a solution of 2-methylpropanoyl chloride (3.84 g, 36.0 mmol) in THF (50 mL). The reaction mixture was then warmed up to room temperature and stirred for 2 h. After removing the solvent *in vacuo*, the crude product was purified by recrystallization (hexane/THF), affording the desired amide, *tert*-butyl 4-[4-(isobutyrylamino)phenyl]-1-piperidinecarboxylate, as a white solid (8.60 g, 84%). The *tert*-butyl 4-[4-(isobutyrylamino)phenyl]-1-piperidinecarboxylate was dissolved in CH_2Cl_2 (50 mL) at room temperature, TFA (13.68 g, 120 mmol, 5 equiv.) was added by syringe. The reaction mixture was stirred for 3 or 4 h and another 5 equivalents of TFA was added and the mixture was stirred for 2 or 3 more hours. The reaction solution was then basified to pH > 14 by KOH

(aq, 2 M). The solution was extracted with CH_2Cl_2 (8 x 200 mL). The combined organic layer was dried over K_2CO_3. Removal of solvent under reduced pressure gave the free amine, 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide, as a brownish solid (5.99 g, 98%). ^1H NMR (400 MHz, $CDCl_3$) δ 7.55-7.35 (m, 2H), 7.35-6.9 (m, 3H), 3.26-2.98 (m, 2H), 2.84-2.64 (m, 2H), 2.64-2.53 (m, 1H), 2.53-2.32 (m, 1H), 1.90-1.68 (m, 2H), 1.68-1.36 (m, 3H), 1.22 (d, 6H, J = 6.0 Hz); ESMS *m/e*: 247.1 $(M + H)^+$.

_N_-[4-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by the procedure for 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide using *tert*-butyl 4-(4-aminophenyl)-1-piperidinecarboxylate and propanoyl chloride: ESMS *m/e*: 233.1 $(M + H)^+$.

_N_-[4-(4-PIPERIDINYL)PHENYL]BUTANAMIDE: Prepared by the procedure for 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide using *tert*-butyl 4-(4-aminophenyl)-1-piperidinecarboxylate and butanoyl chloride: ESMS *m/e*: 247.2 $(M + H)^+$.

_N_-[3-(4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE: Prepared by the procedure for 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide using *tert*-butyl 4-(3-aminophenyl)-1-piperidinecarboxylate and cyclopropanecarbonyl chloride: Anal. Calcd for $C_{15}H_{20}N_2O+0.15CH_2Cl_2$: C, 70.8; H, 7.87; N, 10.9. found: C, 70.9; H, 7.68; N, 11.1; ESMS *m/e*: 245.0 $(M + H)^+$.

***N*-[3-(4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by the procedure for 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide using *tert*-butyl 4-(3-aminophenyl)-1-piperidinecarboxylate and propanoyl chloride: Anal. Calcd for $C_{14}H_{20}N_2O$: C, 72.2; H, 8.63; N, 12.1. found: C, 72.4; H, 8.68; N, 12.1; ESMS *m/e*: 233.1.

Procedure N

Scheme AV



The library was constructed in polypropylene Robbins 46 well plates Reactor Blocks. In the initial incubation period, each well was charged with PS-TBD resin (from Argonaut Technologies, 0.280 mmol, 2.50 eq, 200 mg) and piperidine (0.120 mmol, 1.10 eq) in acetonitrile (0.500 mL) and agitated for 1 h. A solution of benzyl iodide or bromide (0.110 mmol, 1.00 eq) in acetonitrile (0.500 mL) was added to each well followed by additional acetonitrile (1.00 mL) to make a total volume of 2.00 mL and the mixture was rotated in a Robbins rotating oven at room temperature for 16 h. Then AP-Isocyanate resin (Argonaut Technologies, 250 mg, 0.430 mmol, 4.00 eq) was added to each well and reacted further at room temperature for another 12 h. The mixture was filtered

and the filtrate was concentrated *in vacuo* to obtain the desired product that was characterized via LC-MS.

Procedure O

Alkylation of Piperidines Using Alcohols and PS-TSCl Resin in Robbins 48 well "Reactor Blocks"

Scheme W



The library was constructed in polypropylene Robbins "Reactor Blocks", 46 well plates. PS-TSCl resin (100 mg, 1.00 eq, purchased from Argonaut Technologies) was placed in each well of the "Reactor Blocks" 46 well plates. To each well was added an alcohol (1.50 mmol) in 3.00 mL of CH_2Cl_2 and pyridine (1:1). The mixture was stirred for 5 h and the resin was washed with CH_2Cl_2 (3 x 4mL), DMF (5 x 4.0 mL), DMF/H_2O (3:1, 5 x 4.0 mL), THF (3 x 4.0 mL), CH_2Cl_2 (3 x 4.0 mL), acetonitrile (2 x 4.0 mL) and dried under reduced pressure. A solution of an amine (0.0750 mmol, 0.500 eq) and N,N-diisopropylethyl amine (19.0 mg, 0.150 mmol, 1.00 eq) in acetonitrile (3.00 mL) was added to the well containing the derivatized resin and the mixture was reacted at 70 °C for 16 h. Finally, AP-Isocyanate resin (120 mg, 0.150 mmol, 1.00 eq) and THF (2.00 mL) was added to the reaction vessel and reacted at room temperature for another 3 h. The solution was filtered into the Robbins receiving plates and concentrated *in vacuo* to give the desired tertiary amine, which was analyzed via LC-MS.

Procedure P

Scheme AB



***N*-{3-[1-(3-{[(4-FLUOROANILINO)CARBONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** A solution of *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide (26.4 mg, 0.0870 mol), 1-fluoro-4-isocyanatobenzene (11.9 mg, 0.0870 mmol), in THF (1.00 mL) was stirred for 12 h at 25 °C. The resulting crude mixture was diluted with water (10 mL), the aqueous layer was extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified by preparative TLC using 2.5 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product *N*-{3-[1-(3-{[(4-fluoroanilino)carbonyl]amino}propyl)-4-piperidinyl]phenyl}-2-methylpropanamide (4.18 mg, 10.9 %): ^1H NMR (400 MHz, $CDCl_3$) 7.45 (q, 2H, J = 4.7 Hz), 7.23-7.21 (m, 4H), 7.05 (t, 4H, J = 7.8 Hz), 6.75 (m, 1H), 4.05 (m, 1H), 3.19 (s, 1H), 2.71 (m, 1H),

2.53 (m, 1H), 2.26-2.21 (m, 3H), 1.80-1.60 (m, 9H), 1.25 (d, 6H, J = 6.4 Hz); ESMS *m/e*: 439.4 $(M + H)^+$.

Procedure Q_1

Scheme AT



If reacted individually, a solution of the amine (1.0 eq), an electrophile (1.5 eq), diisopropylethylamine (2.0 eq) in CH_2Cl_2 was stirred for 1 day. The solvent was removed *in vacuo* and the crude product was chromatographed to give the final product.



2-METHYL-*N*-{3-[1-(3-{[(4-METHYLPHENYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: A solution of 4-methylbenzenesulfonyl chloride (16.6 mg, 0.0870 mmol), *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide (26.4 mg, 0.0870 mmol), TEA (10.0 mg, 0.174 mmol) in THF (1.00 mL) was stirred for 12 h at 25 °C. The resulting crude mixture was diluted with water (20 mL), the aqueous layer was extracted with CH_2Cl_2 (2 X 20 mL). The combined organic layers were concentrated

in vacuo and the residue was purified by preparative TLC using 2.5 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product 2-methyl-*N*-{3-[1-(3-{[(4-methylphenyl)sulfonyl]amino} propyl)-4-piperidinyl]phenyl}propanamide (17.3 mg, 43.6 %): ^1H NMR (400 MHz, $CDCl_3$) δ 8.19 (s, 1H), 7.53 (s, 1H), 7.41 (s, 1H), 7.32-7.21 (m, 4H), 7.16 (s, 1H), 6.97 (d, 1H, J = 7.9 Hz), 3.44 (t, 2H, J = 6.3 Hz), 3.15 (d, 2H, J = 9.8 Hz), 2.62-2.45 (m, 4H), 2.15 (m, 3H), 2.05 (s, 3H), 1.95-1.71 (m, 5H), 1.26 (d, 6H, J = 6.6 Hz); ESMS *m/e*: 458.2 $(M + H)^+$.

Procedure Q_2

The Capture and Release Method for the Synthesis and Purification of the Piperidine Library

The commercially obtained Amberlyst 15 exchange resin (Aldrich) was activated using the following procedure:

1. The resin was shaken in methanol for 24 hr.
2. The resin was filtered and washed with methanol on a fritted funnel.
3. The resin was neutralized with 2N NH_3 in MeOH (pH checked) - shaken for 1 hr.
4. The neutralized resin was acidified with 3M HCl in MeOH (pH checked) - shaken for 1 hr.
5. The resin was captured on a fritted funnel and washed with MeOH.
6. The resin was dried *in vacuo* and stored.

Synthesis (Acylation of the Amines):

The library was constructed in polypropylene Robbins "Reactor Blocks", 46 well plates. In each plate an array of 5 amines (0.10 mmol) and 8 electrophiles (acid

chlorides, sulfonyl chlorides, 1.5 eq.) in the presence of triethylamine (2.0 eq) in THF/DCM 3:1 (2.0 mL) were reacted overnight to give 40 compounds/plate. The reactions were rigorously monitored via TLC to the depletion of the starting amine due to the ensuing purification methodology via the acidic Amberlyst 15 resin. Following the disappearance of the starting amine, the desired products were captured and then released using the process outlined below.

Purification of the Piperidine Products: Activated Amberlyst 15 ion-exchange resin (0.90 g, Aldrich) was added to each well, and the plates were rotated for 2 hours in a Robbins rotating oven to capture the desired final product from the reaction mixture. The solvent was filtered and the resin was washed with CH_3OH and CH_2Cl_2 (x 3) alternately with each of the solvents (for 10 minutes each time). After the last filtration, 2 N ammonia in methanol was added to the resin (2 mL to each well) and the reaction blocks were rotated for 2 hours to release the desired compounds from the resin. The final compounds were filtered into Robbins' "Receiving Blocks", the solvent was removed and the compounds were analyzed via LC-MS.

Procedure R

Scheme Z



[(3-CHLOROPROPYL)SULFANYL]BENZENE: A mixture of benzenethiol (0.550 g, 5.00 mmol), 1-bromo-3-chloropropane (106 mg, 5.50 mmol), TEA (1.01 g, 10.0 mmol) and THF (10.0 mL) was stirred for 12 h at 25 °C. The resulting crude mixture was diluted with water (40 mL), the aqueous layer was extracted with CH_2Cl_2 (3 X 30 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified by preparative TLC using hexane:EtOAc (10:1) to give the desired product [(3-chloropropyl)sulfanyl]benzene (1.05 g, 100 %).

Scheme AA



Procedure S

3-CHLOROPROPYL 4-FLUOROPHENYL SULFOXIDE: A solution of 3-chloropropyl 4-fluorophenyl sulfide (77.5 mg, 0.380 mmol) in CH_2Cl_2 (2.00 mL) was cooled to 0 °C. To this solution *m*-CPBA (78.7 mg, 0.460 mmol) was added. The reaction mixture was stirred at 0 °C for 30 min, then at

23 °C for 4 h. The resulting crude mixture was diluted with 10% aqueous Na_2SO_3 (10 mL), the aqueous layer was extracted with CH_2Cl_2 (2 X 15 mL). The combined organic layers were washed with brine (10 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The residue was purified by preparative TLC using 2.5 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product 3-chloropropyl 4-fluorophenyl sulfoxide (47.8 mg, 57.0 %).

Procedure T

Scheme AD





***N*-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-*N*,2-DIMETHYLPROPANAMIDE:** A mixture of *N*-(3-{1-[4-(3,4-dimethylphenyl)-4-oxobutyl]-4-piperidinyl}phenyl)-2-methylpropanamide (15.0 mg, 0.0357 mmol), MeI (5.07 mg, 0.0357 mmol), NaO*t*Bu (6.86 mg, 0.0714 mmol) and THF (1.00 mL) was stirred for 5 h at 25 °C. The resulting crude mixture was diluted with water (10 mL), the aqueous layer was extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified by preparative TLC using 4.0 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to afford the desired product *N*-(3-{1-[4-(3,4-

dimethylphenyl)-4-oxobutyl]-4-piperidinyl}phenyl)-*N*,2-dimethylpropanamide (13.8 mg, 89.1 %): ^{1}H NMR (400 MHz, $CDCl_3$) 7.76 (s, 1H), 7.72 (dd, 1H, J = 1.8, 7.7 Hz), 7.33(t, 1H, J = 8.8 Hz), 7.22 (d, 1H, J = 7.8 Hz), 7.18(d, 1H, J = 8.8 Hz), 7.01 (m, 2H), 3.24 (s, 3H), 3.10 (d, 1H, J = 10.6 Hz), 3.00 (t, 1H, J = 7.6 Hz), 2.49-2.44 (m, 4H), 2.33 (s, 6H), 2.112.10 (m, 2H), 1.99 (m, 1H), 1.79-1.77 (m, 4H), 1.26 (t, 2H, J = 7.6 Hz), 1.02 (d, 6H, J = 7.6 Hz); ESMS *m/e*: 435.2 $(M + H)^+$.

Procedure U

Scheme AK





1-[3-(3-CHLOROPROPOXY)PHENYL]ETHANONE: To a suspension of NaH (50.5 mg, 2.00 mmol) in DMF (1.00 mL) was added 1-(3-hydroxyphenyl)ethanone (136 mg, 1.00 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 1 h. To this mixture was added a solution of 1-bromo-3-chloropropane (188 mg, 1.20 mmol) in DMF (0.500 mL). The reaction mixture was stirred at room temperature for 5 h. The resulting crude mixture was diluted with water (20 mL), the aqueous layer was extracted with CH_2Cl_2 (3 x

20 mL). The combined organic layers were washed with brine (20 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The residue was purified by preparative TLC using hexane:EtOAc (4:1) to afford the desired product 1-[3-(3-chloropropoxy)phenyl]ethanone (235 mg, 55.2 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.7 (d, 1H, J = 6.6 Hz), 7.52(s, 1H), 7.25 (t, 1H, J = 6.6 Hz), 7.01 (m, 1H), 4.11 (t, 2H, J = 7.9 Hz), 3.69 (t, 2H, J = 7.9 Hz), 2.61 (s, 3H), 1.95-1.92 (m, 2H).

Procedure V

Scheme AE



1-[(2,2-DIMETHYLPROPANOYL)OXY]-4-(4,4,5,5-TETRAMETHYL-1,3,2-DIOXABOROLAN-2-YL)-1,2,3,6-TETRAHYDROPYRIDINE:

To a 50-mL RB-flask, charged with bis(pinacolato)diboron (422 mg, 1.66 mmol), KOAc (444 mg, 4.53 mmol) and $PdCl_2$dppf (37.0 mg, 3.00 mol%), dppf (25.0 mg, 3.00 mol%), was added a solution of 1-[(2,2-dimethylpropanoyl)oxy]-1,2,3,6-tetrahydro-4-pyridinyl trifluoromethanesulfonate (500 mg, 1.51 mmol) in 1,4-dioxane (10.0 mL) at room temperature under argon. The mixture was heated at 80 °C overnight. After cooled to room temperature, the mixture was filtered through celite and the celite was washed with EtOAc (3 x 20 mL). The filtrates were concentrated *in vacuo*. The resulting residue was dissolved in EtOAc and washed with H_2O and

brine, dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was purified by flash chromatography (1:9 EtOAc:hexane) to give 1-[(2,2-dimethylpropanoyl)oxy]-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,6-tetrahydropyridine (355 mg, 76.0 %).

Procedure W

Scheme AF



***TERT*-BUTYL 4-[5-(ISOBUTYRYLAMINO)-2-METHYLPHENYL]-3,6-DIHYDRO-1(2H)-PYRIDINECARBOXYLATE:** To a 50-mL RB flask containing 1-[(2,2-dimethylpropanoyl)oxy]-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,6-tetrahydropyridine (500 mg, 1.62 mmol), K_2CO_3 (670 mg, 4.86 mmol) and $PdCl_2dppf$ (155 mg) was added a solution of N-(3-bromo-4-methylphenyl)-2-methylpropanamide (415mg, 1.62 mmol) in DMF (10.0 mL) at room temperature under argon. The mixture was heated to 80 °C under argon overnight. After cooled to room temperature, the mixture was filtered through celite and the celite was

washed with EtOAc (3 x 20 mL). The filtrates were washed with H_2O (20 mL), brine (20 mL), dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was purified flash chromatography (20% EtOAc/ hexane) to give tert-butyl 4-[5-(isobutyrylamino)-2-methylphenyl]-3,6-dihydro-1(2H)-pyridinecarboxylate (360 mg, 62.0 %).

Scheme AG



Procedure X

***TERT*-BUTYL 4-[5-(ISOBUTYRYLAMINO)-2-METHYLPHENYL]-1-PIPERIDINECARBOXYLATE:** A solution of tert-butyl 4-[5-(isobutyrylamino)-2-methylphenyl]-3,6-dihydro-1(2H)-pyridinecarboxylate (335 mg, 0.93 mmol) and 10% Pd/C (35.0 mg) in EtOH (20.0 mL) was hydrogenated at room temperature overnight using the hydrogen balloon method. The reaction mixture was filtered through celite and washed with ethanol (3 x 10 mL). The combined extracts were concentrated *in vacuo* to afford tert-butyl 4-[5-(isobutyrylamino)-2-methylphenyl]-1-piperidinecarboxylate (335 mg, 100 %).

Procedure Y

Scheme AH



2-METHYL-*N*-[4-METHYL-3-(4-PIPERIDINYL)PHENYL] PROPANAMIDE: Into a solution of tert-butyl 4-[5-(isobutyrylamino)-2-methylphenyl]-1-piperidinecarboxylate (335 mg, 0.930 mmol) in CH_2Cl_2 (10.0 mL) was added TFA (10.0 mL) at room temperature. The reaction mixture was stirred for 2 h and concentrated *in vacuo*. The residue was dissolved in 20 mL of $CHCl_3$/i-PrOH (3:1) and was basified with 5% KOH solution (10 mL). The aqueous layer was extracted with $CHCl_3$/i-PrOH (3:1, 3 x 10 mL). The combined organic extracts were washed with brine, dried over $MgSO_4$, filtered and concentrated *in vacuo* to give 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide (190 mg, 78.0 %).

Procedure Z

Scheme AI





***N*-(3-{1-[4,4-BIS(4-FLUOROPHENYL)BUTYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** A solution of 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide (49.0 mg, 0.190 mmol), 1-[4-chloro-1-(4-fluorophenyl)butyl]-4-fluorobenzene (58.0 mg, 0.210 mmol), NaI (42.0 mg, 0.280 mmol) and K_2CO_3 (52.0 mg, 0.380 mmol) in DMF (10.0 mL) was heated at 95 °C overnight. The mixture was diluted with water (20 mL) and the aqueous layer was extracted with EtOAc (3 x 20 mL). The combined organic layers were washed with brine, dried over $MgSO_4$ and concentrated *in vacuo*. The crude product was purified by flash chromatography [5% NH_3 (2.0 M in MeOH) in CH_2Cl_2] to afford *N*-(3-{1-[4,4-bis(4-fluorophenyl)butyl]-4-piperidinyl}-4-methylphenyl)-2-methylpropanamide (37.0 mg, 38.0 %).

Procedure AA

Scheme AJ





***N*-(3-{1-[4-(3,4-DIFLUOROPHENOXY)BENZYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** To a solution of 4-(3,4-Difluorophenoxy)benzaldehyde (41.0 mg, 0.170 mmol) and 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide (45.0 mg, 0.170 mmol) in 1,2-dichloroethane (5.00 mL) was added sodium triacetoxyborohydride (110 mg, 0.520 mmol) and AcOH (10.0 µL, 0.170 mmol) at room temperature. The mixture was stirred overnight. The reaction mixture was quenched by saturated $NaHCO_3$ solution (10 mL) and extracted with CH_2Cl_2 (3 x 10 mL). The combined organic layers were washed with brine, dried over $MgSO_4$, concentrated *in vacuo*. The crude product was purified by preparative TLC using 5% NH_3 {2.0 M in MeOH) in CH_2Cl_2 to give the desired product *N*-(3-{1-[4-(3,4-difluorophenoxy)benzyl]-4-piperidinyl}-4-methylphenyl)-2-methylpropanamide (44.0 mg, 54.0 %).

Procedure AC

Scheme AT: Synthesis of Amides using PS-Carbodiimide Resin





A mixture of a carboxylic acid (0.0800 mmol) and PS-Carbodiimide Resin (2.00 eq, 80.0 mg, 1.34 mmol/g) in DCM:DMF (10:1, 3.00 mL) was shaken for 30 min. To the reaction mixture was added amine (0.0540 mmol) and the resulting mixture was shaken for 12 h at room temperature. The reaction mixture was filtered and the resin was washed with CH_2Cl_2. The combined organic extracts were concentrated to a small volume, applied to a preparative TLC plate and eluted with 6 % NH_3 (2.0 M in MeOH) in CH_2Cl_2 to give the desired product.

Procedure AD

Scheme X







***TERT*-BUTYL *N*-(3-BROMOPROPYL)CARBAMATE:** Prepared from 3-bromopropylamine hydrobromide and BOC_2O in the presence of base in CH_2Cl_2: ^1H NMR (300 MHz) δ 5.07 (br, 1 H), 3.31 (t, 2 H, J = 6.6 Hz), 3.12 (apparent br q, 2 H, J = 6.0 Hz), 1.92 (p, 2 H, J = 6.6 Hz), 1.30 (s, 9H).

Step 1. To a solution of piperidine (19.3 mmol) in dioxane (20.0 mL) was *N*-(*tert*-butoxycarbonyl)-3-bromopropylamine (21.2 mmol) and potassium carbonate (38.7 mmol) at room temperature and the mixture was heated at reflux temperature for 24 h. The reaction mixture was cooled to room temperature, concentrated *in vacuo* and partitioned between $CHCl_3$ (40 mL) and water (5 mL). The organic layer was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by column chromatography (ethyl acetate: methanol 9:1) to yield the required product *tert*-butyl 3-{4-[3-(acetylamino)phenyl]-1-

piperidinyl}propylcarbamate as a colorless oil: ESMS *m/e*: 376.2 $[M+H]^+$.

Step 2. HCl gas was bubbled into a solution of the boc-protected amine (12.1 mmol) in dioxane (5.00 mL) for 10-20 minutes at 0-5 °C. The resulting solution was stirred at 0-5 °C for 1 h, concentrated, neutralized with 10 % KOH solution (10 mL) and extracted into CH_2Cl_2 (25 mL). The organic extract was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The crude product was chromatographed to give the desired product *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide: ESMS *m/e*: 276.1 $[M+H]^+$.

Procedure AE

Scheme Y



Step 1: A mixture of piperidine (1.00 eq, 0.0226 mmol), *N*-(bromoalkyl)phthalimide (1.50 eq, 0.0338 mmol), Bu_4NI (200 mg) and diisopropylethylamine (5.00 eq, 0.113 mmol)

in dioxane (200 mL) was heated at 99 °C for 24 h. The reaction was followed by TLC analysis (95:5 CH_2Cl_2:methanol). If necessary additional 0.0113 mmol of the appropriate bromoalkylphthalimides was added to each reaction mixture and the heating was continued for additional 48 h. The reaction mixture was cooled to room temperature, the ammonium salts were filtered out and the solvent was removed under reduced pressure. The crude product was chromatographed to give the desired product.



ESMS m/e: 420.2
$[M+H]^+$



ESMS m/e: 434.4
$[M+H]^+$



ESMS m/e: 448.4
$[M+H]^+$



ESMS m/e: 462.4
$[M+H]^+$



ESMS m/e: 476.4
$[M+H]^+$

Step 2: Deprotection of the resulting phthalimides was conducted by heating a solution of phthaliamide-protected amines with excess hydrazine hydrate (10 eq) in ethanol (0.5-1.0 M) at 90 °C for 4 h. The reaction mixture was monitored by TLC to completion. Upon the reaction was completed, the mixture was cooled to room temperature, the insoluble by-products were filtered out through celite and the solvent was removed *in vacuo*. The crude product was chromatographed (dichloromethane-methanol-isoprpylamine) to give the desired products.



ESMS m/e: 290.2 $[M+H]^+$



ESMS m/e: 304.1 $[M+H]^+$



ESMS m/e: 318.2 $[M+H]^+$



ESMS m/e: 332.2 $[M+H]^+$



ESMS m/e: 346.3 $[M+H]^+$

Procedure AF





Scheme H



n = 0-4

(4*R*)-4-(3,4-DIFLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXAMIDE was synthesized according to Scheme H and Procedure AF: To a solution of (4*R*)-4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one (this compound and analogs were prepared according to *J. Med. Chem* **2000,** *43,* 2775) (0.300 mol, 60.0 mg) in THF (5.00 mL) was added LDA (2.0 M in THF, 0.390 mmol, 0.200 mL) at -78 °C under argon. After 30 min at -78 °C, to the mixture was added a solution of 4-nitrophenyl chloroformate (0.330 mmol, 51.2 mg) in THF (0.500 mL) at -78 °C. After stirring for 30 min at -78 °C the reaction

mixture was diluted with a saturated Na_2CO_3 solution (5.0 mL) and the aqueous layer was extracted with CH_2Cl_2 (3 x 10 mL). The combined organic layers were washed with brine (10 mL), dried over Na_2SO_4 and concentrated *in vacuo*. The residue was purified by preparative TLC plates (10:1 hexane:ethyl acetate) to afford 4-nitrophenyl (4*R*)-4-(3,4-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxylate (51.5 mg, 54.0 %).

4-Nitrophenyl (4*R*)-4-(3,4-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxylate (169 mg, 0.465 mmol), *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide (141 mg, 0.465 mmol), K_2CO_3 (0.193 g, 1.39 mmol), CH_2Cl_2 (10 mL), and methanol (0.1 mL) were combined in a flask. The mixture was stirred overnight at room temperature, the solvent was removed *in vacuo*, and the residue was purified by chromatography [2.5% of NH_3 (2.0 M in methanol) in CH_2Cl_2] to afford the desired product (26.1 mg, 10.6 %): ^1H NMR (400 MHz, $CDCl_3$) δ 8.08 (t, 1H, J = 5.5 Hz), 7.45 (S, 2H), 7.38 (d, 1H, J = 8.6 Hz), 7.24-7.12 (m, 3H), 7.06 (m, 1H), 6.97 (d, 1H, J = 8.6 Hz), 5.40 (dd, 1H, J = 3.9-8.8 Hz), 4.71 (t, 1H, J = 8.8 Hz), 4.23 (dd, 1H, J = 4.4, 9.1 Hz), 3.32 (qt, 2H, J = 6.1 Hz), 2.99 (d, 2H, J = 11.0 Hz), 2.49 (qt, 2H, J = 7.0 Hz), 2.41(t, 2H, J = 7.0 Hz), 1.99-1.97 (m, 2H), 1.82-1.68 (m, 6H), 1.23 (d, 6H, J = 7.3 Hz); Anal. Calcd. for $C_{28}H_{34}F_2N_4O_4$+HCl+0.185$CHCl_3$: C, 57.6; H, 6.04; N, 9.54. Found: C, 58.5; H, 6.08; N, 9.47; ESMS *m/e*: 529.1 $(M + H)^+$.

Procedure AG

Scheme AR:



Step 1: A solution of ketoester (10 mmol), Meldrum's acid (10 mmol), aldehyde (10 mmol) and an ammonium acetate (11 mmol) in HOAc (10 mL) was heated at reflux temperature for 18 h.[1] The cooled reaction mixture was poured over ice (100 g). The precipitated oils were collected and dried under reduced pressure. The benzyl ester protected analogs solidified upon trituration with a mixture of ether/hexane.



1.05 g, 29.0 %



523 mg, 15.0%

[1] MORALES, A.; OCHOA, E.; SUAREZ, M.; VERDECIA, Y.; GONZALEZ, L.; MARTIN, N.; QUINTEIRO, M.; SEOANE, C.; SOTO, J. L.; *J. Heterocycl. Chem.* [JHTCAD] **1996**, *33 (1)*, 103-107.

Step 2: A mixture of a benzyl ester and 10% Pd/C in methanol was hydrogenated using the balloon method at room temperature. The reaction mixture was monitored (TLC) to completion, filtered through Celite 545 and the Celite filter cake was washed with methanol (3 x 10 mL). The combined methanol extracts were concentrated *in vacuo* to give the desired carboxylic acid that was used in the next step without any further purification.



4-(2,4-DIFLUOROPHENYL)-2-METHYL-6-OXO-1,4,5,6-TETRAHYDRO-3-PYRIDINECARBOXYLIC ACID was synthesized according to Procedure AG and Scheme AR: ^{1}H NMR ($CDCl_3$, 400 MHz) δ 7.82 (s, 1H), 7.00-6.72 (m, 3H), 4.51 (d, 1H, J = 8.4 Hz), 2.90 (dd, 1H, J = 8.4, 16.3 Hz), 2.68 (d, 1H, J = 16.3 Hz), 2.46 (s, 3H).



4-(3,4-DIFLUOROPHENYL)-2-METHYL-6-OXO-1,4,5,6-TETRAHYDRO-3-PYRIDINECARBOXYLIC ACID was synthesized according to Procedure AG and Scheme AR: ^{1}H NMR ($CDCl_3$, 300 MHz) δ 7.40-6.80 (m, 4 H), 4.23 (d, 1 H, J = 7.5 avg. Hz), 2.93 (dd, 1 H, J = 16.8, 7.5 avg. Hz), 2.68 (d, 1 H, J = 16.5 avg. Hz), 2.45 (s, 3 H).

Procedure AH



1-(6-CHLOROHEXYL)-1*H*-INDOLE: To a mixture of NaH (0.249 g, 10.0 mmol) in DMF (5.00 mL) was added a solution of 1-*H*-indole (0.585 g, 5.00 mmol) in DMF (2.00 mL) at 0 °C. The reaction mixture was stirred for 30 minutes at 0 °C and warmed up to room temperature. To the reaction mixture 1-bromo-6-chlorohexane (0.998 g, 5.00 mmol) was added dropwise via syringe and the reaction mixture was stirred overnight. The reaction mixture was diluted with EtOAc (30 mL), washed with water (3 X 10 mL), brine (10 mL), dried over $MgSO_4$, concentrated *in vacuo* and purified by chromatography using hexane:EtOAc (97.5:2.5) to give the desired product (0.900 g, 76.0 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76-7.54 (m, 1H), 7.47-6.96 (m, 4H), 6.60-6.34 (m, 1H), 4.13 (t, 2H, *J* = 6.8 Hz), 3.50 (t, 2H, *J* = 5.6 Hz), 1.98-1.79 (m, 2H), 1.79-1.64 (m, 2H), 1.54-1.17 (m, 4H).

***N*-(3-{1-[6-(1*H*-INDOL-1-YL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** A mixture of 1-(6-Chlorohexyl)-1*H*-indole (23.6 mg, 0.100 mmol), 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (24.6 mg, 0.100 mmol), K_2CO_3 (27.6 mg, 0.200 mmol), NaI (22.5 mg, 0.150 mmol) and DMF (1.00 mL) was heated at 100 °C for 12 h. The reaction mixture was cooled to room temperature and the crude material was purified by preparative TLC using 5 %

of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product as a yellow solid (40 mg, 90 %): ^1H NMR (400 MHz, $CDCl_3$) δ 8.08-6.52 (m, 11H), 4.17 (t, 2H, *J* = 7.2 Hz), 3.26 (d, 2H, *J* = 11.6 Hz), 2.74-2.52 (m, 4H), 2.44-2.28 (m, 2H), 2.20-2.02 (m, 2H), 1.98-1.82 (m, 4H), 1.78-1.62 (m, 2H), 1.43-1.28 (m, 4H), 1.28 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 446.5 $(M + H)^+$.

Procedure AI:

Scheme AU: Preparation of tert-Piperdines Usingd PS-SO2Cl Resin



The library was constructed in polypropylene Robbins "Reactor Blocks", 48 well plates. PS-TSCl resin (100 mg, 1.00 eq, purchased from Argonaut Technologies) was placed in each well of the "Reactor Blocks" 48 well plates. To each well was added 2-10 eq of an alcohol in dichloromethane:pyridine (1:1, 3.00 mL). The mixture was stirred at room temperature for 5 h and the resin was washed with dichloromethane (3 x 4.00 mL), DMF (5 x 4.00 mL), DMF/H_2O (3:1, 5 x 4.00 mL), THF (3 x 4.00 mL), dichloromethane (3 x 4.00 mL), acetonitrile (2 x 4.00 mL) and dried under reduced pressure. A solution of an amine (0.0750 mmol, 0.500 eq) and N,N-diisopropylethyl amine (19.0 mg, 0.150 mmol, 1.00 eq) in acetonitrile

(3.00 mL) was added to the well containing the derivatized resin and the mixture was reacted at 70 °C for 16 h in the Robbins rotating oven. After cooling, AP-isocyanate resin (120 mg, 0.150 mmol, 1.00 eq) and THF (2.00 mL) was added to the each reaction vessel and reacted at room temperature for additional 3 h. The solution was filtered into the Robbins® receiving plates and concentrated *in vacuo* to give the desired tertiary amines which were analyzed via LC-MS.

Procedure AJ:

Scheme AV: Preparation of tert-Piperidines Using Piperdines,





The library was constructed in polypropylene Robbins® 48 well plates Reactor Blocks. In the initial incubation period, each well was charged with PS-TBD resin (from Argonaut Technologies, 200 mg, 0.280 mmol, 2.50 eq) and piperidine (0.120 mmol, 1.10 eq) in acetonitrile (0.500 mL) and agitated for 1 h. A solution of benzyl iodide or bromide (0.110 mmol, 1.00 eq) in acetonitrile (0.500 mL) was added to each well followed by additional acetonitrile (1.00 mL) to make a total volume of 2 mL and the mixture was rotated in a Robbins rotating oven at room temperature for 16 h. Then AP-Isocyanate resin (Argonaut Technologies, 250 mg (0.430 mmol, 4.00 eq) was

added to each well and reacted further at room temperature for another 12 h. The mixture was filtered and the filtrate was concentrated *in vacuo* to obtain the desired product that was characterized via LC-MS.

Scheme AX



Example 117

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-OXOPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Procedure K (KI) and Scheme E (K_2CO_3) using 1-(4-bromophenyl)-3-chloro-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 457.1 (M + H)$^+$.

Example 118

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-OXOPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Procedure K (KI) and Scheme E (K_2CO_3) using 3-chloro-1-(4-chlorophenyl)-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.1 (M + H)$^+$.

Example 119

***N*-(3-{1-[3-(4-METHOXYPHENYL)-3-OXOPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Procedure K (KI) and Scheme E (K_2CO_3) using 3-chloro-1-(4-

260

methoxyphenyl)-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 409.2 (M + H)$^+$.

Example 120

***N*-(3-{1-[3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)-3-OXOPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Procedure K (KI) and Scheme E (K_2CO_3) using 3-chloro-1-(2,3-dihydro-1*H*-inden-5-yl)-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 419.2 (M + H)$^+$.

Example 121

2-METHYL-*N*-{3-[1-(3-OXO-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Procedure K (KI) and Scheme E (K_2CO_3) using 3-chloro-1-phenyl-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 379.2 (M + H)$^+$.

Example 122

2-METHYL-*N*-(3-{1-[3-(4-METHYLPHENYL)-3-OXOPROPYL]-4-PIPERIDINYL}PHENYL) PROPANAMIDE: Procedure K (KI) and Scheme E (K_2CO_3) using 3-chloro-1-(4-methylphenyl)-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 393.2 (M + H)$^+$.

Example 123

***N*-(3-{1-[3-(4-FLUOROPHENYL)-3-OXOPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Procedure K (KI) and Scheme E (K_2CO_3) using 3-chloro-1-(4-fluorophenyl)-1-propanone and 2-methyl-*N*-[3-(4-

piperidinyl)phenyl]propanamide: ESMS *m/e*: 397.2 (M + H)$^+$.

Example 124

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-HYDROXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-oxopropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 415.1 (M + H)$^+$.

Example 125

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-(3,4-DIFLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3,4-difluorophenol: ESMS *m/e*: 526.8 (M + H)$^+$.

Example 126

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-(2-METHYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and *o*-cresol: ESMS *m/e*: 505.4 (M + H)$^+$.

Example 127

***N*-(3-{1-[3-(4-FLUOROPHENYL)-3-HYDROXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-oxopropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 399.2 (M + H)$^+$.

Example 128

***N*-(3-{1-[3-HYDROXY-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[3-(4-methoxyphenyl)-3-oxopropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 411.2 $(M + H)^+$.

Example 129

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-HYDROXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-oxopropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 459.1 $(M + H)^+$.

Example 130

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-(4-METHOXYPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenol: ESMS *m/e*: 520.8 $(M + H)^+$.

Example 131

***N*-(3-{1-[3-(4-CHLOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-chlorophenol: ESMS *m/e*: 509.1 $(M + H)^+$.

Example 132

***N*-(3-{1-[3-(4-FLUOROPHENYL)-3-(2,3,4,5,6-PENTAFLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN

using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2,3,4,5,6-pentafluorophenol: ESMS *m/e*: 564.7 $(M + H)^+$.

Example 133

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(2-METHYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-methylphenol: ESMS *m/e*: 548.8 $(M + H)^+$.

Example 134

***N*-(3-{1-[3-(3,4-DIFLUOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3,4-difluorophenol: ESMS *m/e*: 511.1 $(M + H)^+$.

Example 135

***N*-(3-{1-[3-(4-BROMOPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-bromophenol: ESMS *m/e*: 553.0 $(M + H)^+$.

Example 136

***N*-(3-{1-[3-(3,4-DICHLOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN

using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3,4-dichlorophenol: ESMS *m/e*: 542.7 $(M + H)^+$.

Example 137

***N*-[3-(1-{3-(4-FLUOROPHENYL)-3-[4-(TRIFLUOROMETHYL)PHENOXY]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethyl)phenol: ESMS *m/e*: 543.1 $(M + H)^+$.

Example 138

***N*-(3-{1-[3-(3-BROMOPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-bromophenol: ESMS *m/e*: 552.7 $(M + H)^+$.

Example 139

***N*-(3-{1-[3-(4-FLUOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-fluorophenol: ESMS *m/e*: 493.2 $(M + H)^+$.

Example 140

***N*-(3-{1-[3-(3-FLUOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 3-fluorophenol: ESMS *m/e*: 492.9 $(M + H)^+$.

Example 141

***N*-(3-{1-[3-(2,6-DICHLOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2,6-dichlorophenol: ESMS *m/e*: 543.0 $(M + H)^+$.

Example 142

***N*-(3-{1-[3-(2,5-DIFLUOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2,5-difluorophenol: ESMS *m/e*: 511.5 $(M + H)^+$.

Example 143

***N*-(3-{1-[3-(3-CHLOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-chlorophenol: ESMS *m/e*: 509.1 $(M + H)^+$.

Example 144

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(3-METHYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 3- methylphenol: ESMS *m/e*: 549.1 $(M + H)^+$.

Example 145

***N*-(3-{1-[3-([1,1'-BIPHENYL]-4-YLOXY)-3-(4-BROMOPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-phenylphenol: ESMS *m/e*: 611.2 $(M + H)^+$.

Example 146

***N*-(3-{1-[3-(2,4-DIFLUOROPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2,4-difluorophenol: ESMS *m/e*: 511.1 $(M + H)^+$.

Example 147

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(3-METHOXYPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-methoxyphenol: ESMS *m/e*: 564.6 $(M + H)^+$.

Example 148

METHYL 4-(1-(4-BROMOPHENYL)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPOXY)BENZOATE: Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-

2-methylpropanamide and methyl 4-hydroxybenzoate: ESMS *m/e*: 593.0 $(M + H)^+$.

Example 149

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(4-PHENOXYPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-phenoxyphenol: ESMS *m/e*: 626.6 $(M + H)^+$.

Example 150

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(2-CHLORO-4-METHYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-chloro-4-methylphenol: ESMS *m/e*: 583.0 $(M + H)^+$.

Example 151

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-PHENOXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenol: ESMS *m/e*: 535.0 $(M + H)^+$.

Example 152

***N*-[3-(1-{3-(4-BROMOPHENYL)-3-[4-(TRIFLUOROMETHYL)PHENOXY]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethyl)phenol: ESMS *m/e*: 603.1 $(M + H)^+$.

Example 153

***N*-(3-{1-[3-(2-ACETYLPHENOXY)-3-(4-BROMOPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-acetylphenol: ESMS *m/e*: 576.6 $(M + H)^+$.

Example 154

***N*-(3-{1-[3-(3-ACETYLPHENOXY)-3-(4-BROMOPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-acetylphenol: ESMS *m/e*: 576.9 $(M + H)^+$.

Example 155

***N*-(3-{1-[3-(3-ACETYLPHENOXY)-3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-acetylphenol: ESMS *m/e*: 539.2 $(M + H)^+$.

Example 156

***N*-(3-{1-[3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)-3-PHENOXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenol: ESMS *m/e*: 497.2 $(M + H)^+$.

Example 157

***N*-(3-{1-[3-(2-ACETYLPHENOXY)-3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide 2-acetylphenol: ESMS *m/e*: 539.1 $(M + H)^+$.

Example 158

***N*-(3-{1-[3-(4-BROMOPHENOXY)-3-(4-BROMOPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-bromophenol: ESMS *m/e*: 612.7 $(M + H)^+$.

Example 159

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(4-CHLOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-chlorophenol: ESMS *m/e*: 568.7 $(M + H)^+$.

Example 160

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(4-FLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-fluorophenol: ESMS *m/e*: 552.8 $(M + H)^+$.

Example 161

***N*-(3-{1-[3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)-3-(4-METHOXYPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenol: ESMS *m/e*: 527.3 $(M + H)^+$.

Example 162

***N*-(3-{1-[3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)-3-(4-FLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-fluorophenol: ESMS *m/e*: 515.2 $(M + H)^+$.

Example 163

***N*-(3-{1-[3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)-3-HYDROXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE**

Prepared by Procedure L and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-oxopropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 421.2 $(M + H)^+$.

Example 164

***N*-[3-(1-{3-(2,3-DIHYDRO-1*H*-INDEN-5-YL)-3-[4-(TRIFLUOROMETHYL)PHENOXY]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-trifluoromethylphenol: ESMS *m/e*: 565.0 $(M + H)^+$.

Example 165

N-(3-{1-[3-(4-BROMOPHENOXY)-3-(2,3-DIHYDRO-1H-INDEN-5-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-bromophenol: ESMS *m/e*: 577.4 $(M + H)^+$.

Example 166

N-(3-{1-[3-(3-ACETYLPHENOXY)-3-(4-CHLOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-acetylphenol: SMS *m/e*: 533.1 $(M + H)^+$.

Example 167

N-(3-{1-[3-(4-METHOXYPHENOXY)-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenol: ESMS *m/e*: 517.4 $(M + H)^+$.

Example 168

N-(3-{1-[3-(4-CHLOROPHENOXY)-3-(2,3-DIHYDRO-1H-INDEN-5-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(2,3-dihydro-1*H*-inden-5-yl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-chlorophenol: ESMS *m/e*: 531.1 $(M + H)^+$.

Example 169

***N*-(3-{1-[3-(2-ACETYLPHENOXY)-3-(4-CHLOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-acetylphenol: ESMS *m/e*: 533.4 $(M + H)^+$.

Example 170

***N*-(3-{1-[3-(4-BROMOPHENYL)-3-(4-METHOXYPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-bromophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenol: ESMS *m/e*: 565.0 $(M + H)^+$.

Example 171

***N*-(3-{1-[3-(4-BROMOPHENOXY)-3-(4-CHLOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-bromophenol: ESMS *m/e*: 568.8 $(M + H)^+$.

Example 172

***N*-(3-{1-[3-(4-CHLOROPHENOXY)-3-(4-CHLOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-chlorophenol: ESMS *m/e*: 525.0 $(M + H)^+$.

Example 173

***N*-(3-{1-[3-(4-METHOXYPHENYL)-3-PHENOXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenol: ESMS *m/e*: 487.4 $(M + H)^+$.

Example 174

***N*-(3-{1-[3-(4-FLUOROPHENYL)-3-PHENOXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenol: ESMS *m/e*: 475.6 $(M + H)^+$.

Example 175

***N*-(3-{1-[3-(2-ACETYLPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-acetylphenol: ESMS *m/e*: 517.1 $(M + H)^+$.

Example 176

***N*-(3-{1-[3-(3-ACETYLPHENOXY)-3-(4-FLUOROPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-acetylphenol: ESMS *m/e*: 516.9 $(M + H)^+$.

Example 177

***N*-(3-{1-[3-(4-FLUOROPHENYL)-3-(4-METHOXYPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by

Procedure A and Scheme AN using *N*-(3-{1-[3-(4-fluorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenol: ESMS *m/e*: 505.2 $(M + H)^+$.

Example 178

***N*-(3-{1-[3-(4-CHLOROPHENOXY)-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-cholorophenol: ESMS *m/e*: 521.5 $(M + H)^+$.

Example 179

***N*-(3-{1-[3-(3-ACETYLPHENOXY)-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-acetylphenol: ESMS *m/e*: 529.0 $(M + H)^+$.

Example 180

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-PHENOXYPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenol. ESMS *m/e*: 490.9 $(M + H)^+$.

Example 181

***N*-(3-{1-[3-(4-BROMOPHENOXY)-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 4- bromophenol: ESMS *m/e*: 564.9 $(M + H)^+$.

Example 182

***N*-[3-(1-{3-(4-METHOXYPHENYL)-3-[4-(TRIFLUOROMETHYL)PHENOXY]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-trifluoromethyphenol: ESMS *m/e*: 555.1 $(M + H)^+$.

Example 183

***N*-(3-{1-[3-(4-CHLOROPHENYL)-3-(4-FLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-fluorophenol: ESMS *m/e*: 509.1 $(M + H)^+$.

Example 184

***N*-(3-{1-[3-(4-FLUOROPHENOXY)-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-fluorophenol: ESMS *m/e*: 505.5 $(M + H)^+$.

Example 185

***N*-(3-{1-[3-(2-ACETYLPHENOXY)-3-(4-METHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-hydroxy-3-(4-methoxyphenyl)propyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 2- acetylphenol: ESMS *m/e*: 529.2 $(M + H)^+$.

Example 186

***N*-[3-(1-{3-(4-CHLOROPHENYL)-3-[4-(TRIFLUOROMETHYL)PHENOXY]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using *N*-(3-{1-[3-(4-chlorophenyl)-3-hydroxypropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-trifluoromethylphenol: SMS *m/e*: 559.1 $(M + H)^+$.

Example 187

***N*-(3-{1-[(3*S*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-(3-{[(1*S*)-3-chloro-1-phenylpropyl]oxy}phenyl)ethanone and 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 513.0 $(M + H)^+$.

2-(ISOPENTYLOXY)-1-NAPHTHALDEHYDE: 2-Hydroxy-1-naphthaldehyde (1.72 g, 10.0 mmol) and THF (50 ml) were combined in a flask. NaH (312 mg, 13 mmol) was added, followed by 1-bromo-3-methylbutane (1.20 mL, 10.0 mmol). The solution was stirred at room temperature overnight, the solvent was removed *in vacuo*, and the residue was purified by chromatography (5-10 % ethyl acetate / hexane): ^1H NMR (400 MHz, $CDCl_3$) δ 10.9 (s, 1H), 9.28 (dd, 1H, *J* = 0.7 Hz, 8.6 Hz), 8.02 (d, 1H, *J* = 9.1 Hz), 7.75 (d, 1H, *J* = 8.1 Hz), 7.63-7.59 (m, 1H), 7.43-7.39 (m, 1H), 7.27 (d, 1H, *J* = 9.2 Hz), 4.25 (t, 2H, *J* = 6.5 Hz), 1.98-1.84 (m, 1H), 1.80-1.75 (m, 2H), 0.99 (d, 6H, *J* = 6.6 Hz); ESMS *m/e*: 242.8 $(M + H)^+$.

Example 188

***N*-[3-(1-{[2-(ISOPENTYLOXY)-1-NAPHTHYL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2-(isopentyloxy)-1-naphthaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.3 $(M + H)^+$.

2-PROPOXY-1-NAPHTHALDEHYDE: Prepared according to the Procedure for 2-(isopentyloxy)-1-naphthaldehyde using 2-hydroxy-1-naphthaldehyde and 1-bromopropane.

Example 189

2-METHYL-*N*-(3-{1-[(2-PROPOXY-1-NAPHTHYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 2-propoxy-1-naphthaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 445.2 $(M + H)^+$.

4-{[(1-FORMYL-2-NAPHTHYL)OXY]METHYL}BENZONITRILE: Prepared according to the Procedure for 2-(isopentyloxy)-1-naphthaldehyde using 2-hydroxy-1-naphthaldehyde and 4-(bromomethyl)benzonitrile.

Example 190

***N*-{3-[1-({2-[(4-CYANOBENZYL)OXY]-1-NAPHTHYL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-{[(1-formyl-2-naphthyl)oxy]methyl}benzonitrile and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 518.2 $(M + H)^+$.

[(1-FORMYL-2-NAPHTHYL)OXY]ACETONITRILE: Prepared according to the Procedure for 2-(isopentyloxy)-1-

naphthaldehyde using 2-hydroxy-1-naphthaldehyde and bromoacetonitrile.

Example 191

***N*-[3-(1-{[2-(CYANOMETHOXY)-1-NAPHTHYL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using [(1-formyl-2-naphthyl)oxy]acetonitrile and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 442.2 $(M + H)^+$.

2-[(3-CHLOROBENZYL)OXY]-1-NAPHTHALDEHYDE: Prepared according to the Procedure for 2-(isopentyloxy)-1-naphthaldehyde using 2-hydroxy-1-naphthaldehyde and 1-(bromomethyl)-3-chlorobenzene.

Example 192

***N*-{3-[1-({2-[(3-CHLOROBENZYL)OXY]-1-NAPHTHYL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2-[(3-chlorobenzyl)oxy]-1-naphthaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 527.2 $(M + H)^+$.

Example 193

***N*-(3-{1-[4-(4-CHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(4-chlorophenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.50 (s, 1H), 7.34-7.19 (m, 7H), 6.98-6.87 (m, 5H), 3.50 (s, 2H), 2.98 (d, 2H, *J* = 11.8 Hz), 2.58-2.44 (m, 2H), 2.10-1.98 (m, 2H), 1.83-1.76 (m, 4H), 1.24 (d, 6H, *J* = 6.8 Hz); ESMS *m/e*: 463.2 $(M + H)^+$.

Example 194

***N*-(3-{1-[4-(3,4-DIFLUOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(3,4-difluorophenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 465.2 $(M + H)^+$.

4-(ISOPENTYLOXY)-1-NAPHTHALDEHYDE: Prepared according to the Procedure for 2-(isopentyloxy)-1-naphthaldehyde using 4-hydroxy-1-naphthaldehyde and 1-bromo-3-methylbutane.

Example 195

***N*-[3-(1-{[4-(ISOPENTYLOXY)-1-NAPHTHYL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(isopentyloxy)-1-naphthaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.3 $(M + H)^+$.

Example 196

***N*-(3-{1-[4-(4-METHOXYPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(4-methoxyphenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 459.2 $(M + H)^+$.

4-PROPOXY-1-NAPHTHALDEHYDE: Prepared according to the Procedure for 2-(isopentyloxy)-1-naphthaldehyde using 4-hydroxy-1-naphthaldehyde and 1-bromopropane.

Example 197

2-METHYL-*N*-(3-{1-[(4-PROPOXY-1-NAPHTHYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F

and Scheme R using 4- propoxy-1-naphthaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 445.2 $(M + H)^+$.

Example 198

***N*-(3-{1-[4-(3,4-DICHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(3,4-dichlorophenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*:497.1 $(M + H)^+$.

Example 199

***N*-(3-{1-[4-(DIPHENYLAMINO)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(diphenylamino)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 504.2 $(M + H)^+$.

Example 200

***N*-{3-[1-({2,5-DIMETHYL-1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-PYRROL-3-YL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2,5-dimethyl-1-[3-(trifluoromethyl)phenyl]-1*H*-pyrrole-3-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 498.2 $(M + H)^+$.

Example 201

2-METHYL-*N*-(3-{1-[1-(2-PHENYL-1,3-THIAZOL-4-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 1-(2-phenyl-1,3-thiazol-4-yl)ethanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 434.2 $(M + H)^+$.

Example 202

***N*-(3-{1-[(5-CHLORO-3-METHYL-1-PHENYL-1*H*-PYRAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 5-chloro-3-methyl-1-phenyl-1*H*-pyrazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 451.2 $(M + H)^+$.

Example 203

2-METHYL-*N*-(3-{1-[(2-PHENYL-1*H*-IMIDAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 2-phenyl-1*H*-imidazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 403.2 $(M + H)^+$.

Example 204

***N*-[3-(1-{[4-BROMO-1-(4-CHLOROBENZYL)-1*H*-PYRAZOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-bromo-1-(4-chlorobenzyl)-1*H*-pyrazole-5-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 529.1 $(M + H)^+$.

Example 205

2-METHYL-*N*-{3-[1-(3-PHENOXYBENZYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-phenoxybenzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.2 $(M + H)^+$.

Example 206

***N*-(3-{1-[3-(3,4-DICHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by

Procedure F and Scheme R using 3-(3,4-dichlorophenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 497.15 (M + H)$^+$.

Example 207

***N*-(3-{1-[3-(3,5-dichlorophenoxy)benzyl]-4-piperidinyl}phenyl)-2-methylpropanamide:** Prepared by Procedure F and Scheme R using 3-(3,5-dichlorophenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 497.2 (M + H)$^+$.

Example 208

2-METHYL-*N*-(3-{1-[3-(4-METHYLPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-(4-methylphenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 443.2 (M + H)$^+$.

Example 209

2-METHYL-*N*-[3-(1-{3-[3-(TRIFLUOROMETHYL)PHENOXY]BENZYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-[3-(trifluoromethyl)phenoxy]benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 497.2 (M + H)$^+$.

Example 210

***N*-(3-{1-[3-(4-CHLOROPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3-(4-chlorophenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 463.2 (M + H)$^+$.

Example 211

***N*-(3-{1-[3-(DIMETHYLAMINO)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3-(dimethylamino)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 380.2 $(M + H)^+$.

Example 212

***N*-(3-{1-[3-(4-METHOXYPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3-(4-methoxyphenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 459.2 $(M + H)^+$.

Example 213

***N*-(3-{1-[3-(4-*TERT*-BUTYLPHENOXY)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3-(4-*tert*-butylphenoxy)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 485.3 $(M + H)^+$.

Example 214

2-METHYL-*N*-(3-{1-[3-NITRO-4-(1-PIPERIDINYL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-nitro-4-(1-piperidinyl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 465.2 $(M + H)^+$.

Example 215

***N*-(3-{1-[(3,4-DIMETHYLTHIENO[2,3-*B*]THIEN-2-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3,4-dimethylthieno[2,3-*b*]thiophene-2-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 427.1 $(M + H)^+$.

Example 216

2-METHYL-*N*-{3-[1-({3-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-PYRAZOL-4-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-[4-(trifluoromethyl)phenyl]-1*H*-pyrazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 471.1 $(M + H)^+$.

Example 217

2-METHYL-*N*-(3-{1-[4-(1*H*-1,2,4-TRIAZOL-1-YL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 404.1 $(M + H)^+$.

Example 218

2-METHYL-*N*-(3-{1-[(5-METHYL-1-PHENYL-1*H*-PYRAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-methyl-1-phenyl-1*H*-pyrazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 417.1 $(M + H)^+$.

Example 219

2-METHYL-*N*-(3-{1-[4-(4-MORPHOLINYL)-3-NITROBENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-(4-morpholinyl)-3-nitrobenzaldehyde

and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 467.1 $(M + H)^+$.

Example 220

N-{3-[1-({5-[2-CHLORO-4-(TRIFLUOROMETHYL)PHENYL]-2-FURYL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure F and Scheme R using 5-[2-chloro-4-(trifluoromethyl)phenyl]-2-furaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 505.0 $(M + H)^+$.

Example 221

ETHYL 4-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-2,5-DIMETHYL-1-PHENYL-1*H*-PYRROLE-3-CARBOXYLATE: Prepared by Procedure F and Scheme R using ethyl 4-formyl-2,5-dimethyl-1-phenyl-1*H*-pyrrole-3-carboxylate and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 502.2 $(M + H)^+$.

Example 222

ETHYL 5-(4-CHLOROPHENYL)-2-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-3-FUROATE: Prepared by Procedure F and Scheme R using ethyl 5-(4-chlorophenyl)-2-formyl-3-furoate and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 509.0 $(M + H)^+$.

Example 223

***N*-{3-[1-(2,3-DIHYDRO-1,4-BENZODIOXIN-6-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2,3-dihydro-1,4-benzodioxine-6-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 395.1 $(M + H)^+$.

Example 224

2-METHYL-*N*-(3-{1-[(6-PHENOXY-3-PYRIDINYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 6-phenoxynicotinaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 430.1 $(M + H)^+$.

Example 225

2-METHYL-*N*-[3-(1-{[5-(2-PYRIDINYL)-2-THIENYL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-(2-pyridinyl)-2-thiophenecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 420.1 $(M + H)^+$.

Example 226

2-METHYL-*N*-{3-[1-({5-[1-METHYL-3-(TRIFLUOROMETHYL)-1*H*-PYRAZOL-5-YL]-2-THIENYL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-[1-methyl-3-(trifluoromethyl)-1*H*-pyrazol-5-yl]-2-thiophenecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 491.0 $(M + H)^+$.

Example 227

2-METHYL-*N*-[3-(1-{[1-(PHENYLSULFONYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure F and Scheme R using 1-(phenylsulfonyl)-1*H*-indole-3-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 516.1 $(M + H)^+$.

Example 228

***N*-(3-{1-[(1,5-DIMETHYL-3-OXO-2-PHENYL-2,3-DIHYDRO-1*H*-PYRAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 1,5-dimethyl-3-oxo-2-phenyl-2,3-dihydro-1*H*-pyrazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 447.2 $(M + H)^+$.

Example 229

***N*-(3-{1-[4-(4-*TERT*-BUTYL-1,3-THIAZOL-2-YL)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(4-*tert*-butyl-1,3-thiazol-2-yl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide.

Example 230

***N*-{3-[1-(2,3-DIHYDRO-1-BENZOFURAN-5-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2,3-dihydro-1-benzofuran-5-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 379.1 $(M + H)^+$.

Example 231

2-METHYL-*N*-(3-{1-[(4-METHYL-2-PHENYL-5-PYRIMIDINYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-methyl-2-phenyl-5-pyrimidinecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.2 $(M + H)^+$.

Example 232

***N*-{3-[1-(2,1,3-BENZOTHIADIAZOL-5-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2,1,3-benzothiadiazole-5-

carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 395.1 $(M + H)^+$.

Example 233

2-METHYL-*N*-(3-{1-[(5-PHENYL-2-THIENYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-phenyl-2-thiophenecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 419.1 $(M + H)^+$.

Example 234

***N*-{3-[1-(3,4-DIHYDRO-2*H*-1,5-BENZODIOXEPIN-7-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3,4-dihydro-2*H*-1,5-benzodioxepine-7-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 409.2 $(M + H)^+$.

Example 235

2-METHYL-*N*-[3-(1-{[3-(2-THIENYL)-1*H*-PYRAZOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-(2-thienyl)-1*H*-pyrazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 409.1 $(M + H)^+$.

Example 236

***N*-{3-[1-([1,1'-BITHIENYL]-4-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 2,2'-Bithiophene-5-carboxaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.0 $(M + H)^+$.

Example 237

N-(3-{1-[(2,2-DIMETHYL-3,4-DIHYDRO-2H-CHROMEN-6-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure F and Scheme R using 2,2-dimethyl-6-chromanecarbaldehyde and 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 421.2 $(M + H)^+$.

Example 238

2-METHYL-N-{3-[1-({5-[1-METHYL-5-(TRIFLUOROMETHYL)-1H-PYRAZOL-3-YL]-2-THIENYL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-[1-methyl-5-(trifluoromethyl)-1H-pyrazol-3-yl]-2-thiophenecarbaldehyde and 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 491.1 $(M + H)^+$.

Example 239

2-METHYL-N-(3-{1-[(2-PHENYL-1,3-THIAZOL-4-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 2-phenyl-1,3-thiazole-4-carbaldehyde and 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 420.0 $(M + H)^+$.

Example 240

2-METHYL-N-(3-{1-[(3-PHENOXY-2-THIENYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 3-phenoxy-2-thiophenecarbaldehyde and 2-methyl-N-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 435.0 $(M + H)^+$.

Example 241

N-{3-[1-({2-[(4-CHLOROPHENYL)SULFANYL]-3-THIENYL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure F and Scheme R

290

using 2-[(4- chlorophenyl)sulfanyl]-3-thiophenecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 485.0 $(M + H)^+$.

Example 242

N-[3-(1-{[1-(4-CHLOROPHENYL)-1*H*-PYRROL-2-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure F and Scheme R using 1-(4-chlorophenyl)-1*H*-pyrrole-2-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 436.0 $(M + H)^+$.

Example 243

2-METHYL-*N*-{3-[1-({5-[2-(TRIFLUOROMETHOXY)PHENYL]-2-FURYL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-[2-(trifluoromethoxy)phenyl]-2-furaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 487.1 $(M + H)^+$.

Example 244

2-METHYL-*N*-(3-{1-[2-(4-MORPHOLINYL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 2-(4-morpholinyl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 422.2 $(M + H)^+$.

Example 245

***N*-[3-(1-{[3-(4-METHOXYPHENYL)-1*H*-PYRAZOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3-(4-methoxyphenyl)-1*H*-pyrazole-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 433.1 $(M + H)^+$.

Example 246

2-METHYL-*N*-(3-{1-[4-(1*H*-PYRAZOL-1-YL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-(1*H*-pyrazol-1-yl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 402.8 $(M + H)^+$.

Example 247

2-METHYL-*N*-{3-[1-(4-QUINOLINYLMETHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-quinolinecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 388.1 $(M + H)^+$.

Example 248

2-METHYL-*N*-(3-{1-[4-(4-MORPHOLINYL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-(4-morpholinyl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 422.5 $(M + H)^+$.

Example 249

2-METHYL-*N*-(3-{1-[4-(2-THIENYL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-(2-thienyl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 419.1 $(M + H)^+$.

Example 250

2-METHYL-*N*-(3-{1-[(2-METHYL-5-PHENYL-3-FURYL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 2-methyl-5-phenyl-3-furaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 417.2 $(M + H)^+$.

Example 251

***N*-(3-{1-[3-(CYCLOPENTYLOXY)-4-METHOXYBENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 3-(cyclopentyloxy)-4-methoxybenzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 451.1 $(M + H)^+$.

Example 252

2-METHYL-*N*-{3-[1-({5-[4-(TRIFLUOROMETHOXY)PHENYL]-2-FURYL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-[4-(trifluoromethoxy)phenyl]-2-furaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 487.1 $(M + H)^+$.

Example 253

***N*-{3-[1-(1-BENZOTHIEN-2-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 1-benzothiophene-2-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 393.2 $(M + H)^+$.

Example 254

2-METHYL-*N*-{3-[1-({5-[3-(TRIFLUOROMETHOXY)PHENYL]-2-FURYL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 5-[3-(trifluoromethoxy)phenyl]-2-furaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 487.2 $(M + H)^+$.

Example 255

2-METHYL-*N*-{3-[1-(2-QUINOLINYLMETHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 2-quinolinecarbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 388.1 $(M + H)^+$.

Example 256

***N*-(3-{1-[4-(1*H*-IMIDAZOL-1-YL)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(1*H*-imidazol-1-yl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 403.2 $(M + H)^+$.

Example 257

***N*-{3-[1-(9*H*-FLUOREN-2-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 9*H*-fluorene-2-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 425.1 $(M + H)^+$.

Example 258

METHYL 3-[5-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-2-FURYL]-2-THIOPHENECARBOXYLATE: Prepared by Procedure F and Scheme R using methyl 3-(5-formyl-2-furyl)-2-thiophenecarboxylate and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 467.1 $(M + H)^+$.

Example 259

2-METHYL-*N*-{3-[1-(4-PHENOXYBENZYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-phenoxybenzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 429.2 $(M + H)^+$.

Example 260

***N*-{3-[1-([1,1'-BIPHENYL]-4-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using [1,1'-biphenyl]-4-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 413.2 $(M + H)^+$.

Example 261

***N*-(3-{1-[4-(DIBUTYLAMINO)BENZYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 4-(dibutylamino)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 464.6 $(M + H)^+$.

Example 262

2-METHYL-*N*-[3-(1-{4-[(4-METHYLPHENYL)SULFANYL]-3-NITROBENZYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-[(4-methylphenyl)sulfanyl]-3-nitrobenzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/:e* 504.2 $(M + H)^+$.

Example 263

2-METHYL-*N*-(3-{1-[4-(1,2,3-THIADIAZOL-4-YL)BENZYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R using 4-(1,2,3-thiadiazol-4-yl)benzaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e:* 421.1 $(M + H)^+$.

1-(3-{[(1*S*)-3-CHLORO-1-PHENYLPROPYL]OXY}PHENYL)ETHANONE: (1*R*)-3-Chloro-1-phenyl-1-propanol (1.000 g, 5.86 mmol), 1-(3-hydroxyphenyl)ethanone (0.797 g, 5.86 mmol), triphenylphosphine (1.54 g, 5.86 mmol) and

diethylazodicarboxylate (1.53 g, 8.79 mmol) were combined in a flask, which was immediately flushed with argon. THF (20 mL) was added and the mixture was stirred overnight under argon. THF was removed *in vacuo*, the crude product was dissolved in 50 mL of CH_2Cl_2/H_2O (1:1) and the organic layer was separated and dried over $MgSO_4$. After removing the solvent *in vacuo*, the residue was purified by flash chromatography using 10 % ethyl acetate/hexane to yield the desired product (900 mg, 76.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.49-7.46 (m, 2H), 7.40-7.26 (m, 6H), 7.07-7.04 (m, 1H), 5.46-5.43 (dd, 1H, *J* = 4.4 Hz, 8.8 Hz), 3.84-3.78 (m, 1H), 3.64-3.59 (m, 1H), 2.52 (s, 3H), 2.51-2.46 (m, 1H), 2.29-2.22 (m, 1H).

4-(3,4-DIFLUOROPHENOXY)BENZALDEHYDE: 4-Fluorobenzaldehyde (5.32 mL, 49.6 mmol), 3,4-difluorophenol (7.10 g, 54.6 mmol) and K_2CO_3 (8.31 g, 60.1 mmol) were combined in a flask, which was immediately flushed with argon. DMF (50.0 mL) was added and the mixture was heated at reflux under argon for 6 h. Upon cooling to room temperature, EtOAc (100 mL) and H_2O (100 mL) were added; the ethyl acetate layer was separated and washed with H_2O (2 X 100 mL). The combined organic layers were washed with brine, dried over $MgSO_4$, and the solvent was removed *in vacuo*. The desired product was obtained (11.4 g, 98.0 %): ^1H NMR (400 MHz, $CDCl_3$) δ 9.95 (s, 1H), 7.88 (dd, 2H, *J* = 0.8 Hz, 8.8 Hz), 7.24-7.17 (m, 1H), 7.07 (d, 2H, *J* = 8.8 Hz), 6.97-6.92 (m, 1H), 6.86-6.82 (m, 1H); ESMS *m/e*: 235.0 $(M + H)^+$.

***TERT*-BUTYL 4-(4,4,5,5-TETRAMETHYL-1,3,2-DIOXABOROLAN-2-YL)-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** To a flask

were added bis(pinacolato)diboron (422 mg, 1.66 mmol), KOAc (444 mg, 4.53 mmol), $PdCl_2dppf$ (37.0 mg, 3.00 mol%), dppf (25.0 mg, 3.00 mol%) and the flask was flushed with argon. A solution of *tert*-butyl 4-{[(trifluoromethyl)sulfonyl]oxy}-1,2,3,6-tetrahydro-1-pyridinecarboxylate (500 mg, 1.51 mmol) in 1,4-dioxane (10.0 ml) was added and the mixture was stirred at 80 °C overnight. The mixture was filtered through Celite and the filtrate was evaporated *in vacuo*. The resulting residue was dissolved in EtOAc and washed with H_2O, followed by brine. The organic layer was dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was purified by flash chromatography (10% EtOAC/hexane) to give *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-1(2*H*)-pyridinecarboxylate (355 mg, 76.0%): ^1H NMR (400 MHz, $CDCl_3$) δ 6.44 (br s, 1H), 3.93 (br s, 2H), 3.42 (br s, 2H), 2.21 (br s, 2H), 1.45 (s, 9H), 1.25 (s, 12H); ESMS *m/e*: 310.4 $(M + H)^+$.

N-(6-BROMO-2-PYRIDINYL)-2-METHYLPROPANAMIDE: Prepared by Procedure Q1 using 2-methylpropanoyl chloride and 6-bromo-2-pyridinamine: ESMS *m/e*: 242.8 $(M + H)^+$.

***TERT*-BUTYL 4-[6-(ISOBUTYRYLAMINO)-2-PYRIDINYL]-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** Prepared by Procedure W and Scheme AF using *N*-(6-bromo-2-pyridinyl)-2-methylpropanamide and *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 245.8 $(M - 100)^+$.

2-METHYL-*N*-[6-(4-PIPERIDINYL)-2-PYRIDINYL]PROPANAMIDE: Prepared by Procedures X and Y, Schemes AG and AH,

respectively using *tert*- butyl 4-[6-(isobutyrylamino)-2-pyridinyl]-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 248.1 $(M + H)^+$.

Example 264

***N*-(6-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}-2-PYRIDINYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 4-chloro-1-(3,4-dimethylphenyl)-1-butanone and 2-methyl-*N*-[6-(4-piperidinyl)-2-pyridinyl]propanamide: ESMS *m/e*: 422.1 $(M + H)^+$.

Example 265

***N*-(6-{1-[4,4-BIS(4-FLUOROPHENYL)BUTYL]-4-PIPERIDINYL}-2-PYRIDINYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-[4-chloro-1-(4-fluorophenyl)butyl]-4-fluorobenzene and 2-methyl-*N*-[6-(4-piperidinyl)-2-pyridinyl]propanamide: ESMS *m/e*: 492.2 $(M + H)^+$.

Example 266

***N*-(6-{1-[4-(3,4-DIFLUOROPHENOXY)BENZYL]-4-PIPERIDINYL}-2-PYRIDINYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AA and Scheme AJ using 4-(3,4-difluorophenoxy)benzaldehyde and 2-methyl-*N*-[6-(4-piperidinyl)-2-pyridinyl]propanamide: ESMS *m/e*: 466.0 $(M + H)^+$.

***N*-(3-BROMO-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 using 2-methylpropanoyl chloride and 3-bromo-4-methylaniline: ESMS *m/e*: 255.9 $(M + H)^+$.

***TERT*-BUTYL 4-[5-(ISOBUTYRYLAMINO)-2-METHYLPHENYL]-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** Prepared by Procedure W and Scheme AF using *N*-(3-bromo-4-methylphenyl)-2-methylpropanamide and *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 259.1 (M - 100)$^+$.

2-METHYL-*N*-[4-METHYL-3-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedures X and Y, Schemes AG and AH, respectively using *tert*-butyl 4-[5-(isobutyrylamino)-2-methylphenyl]-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 261.0 (M + H)$^+$.

Example 267

***N*-(3-{1-[4-(3,4-DIFLUOROPHENOXY)BENZYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AA and Scheme AJ using 4-(3,4-difluorophenoxy)benzaldehyde and using 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 479.1 (M + H)$^+$.

***N*-(5-BROMO-2-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 using 2-methylpropanoyl chloride and 5-bromo-2-methylaniline: ESMS *m/e*: 255.9 (M + H)$^+$.

***TERT*-BUTYL 4-[3-(ISOBUTYRYLAMINO)-4-METHYLPHENYL]-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** Prepared by Procedure W and Scheme AF using *N*-(5-bromo-2-methylphenyl)-2-methylpropanamide and *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-

dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 259.1 $(M - 100)^+$.

2-METHYL-*N*-[2-METHYL-5-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedures X and Y, Schemes AG and AH, respectively using *tert*-butyl 4-[3-(isobutyrylamino)-4-methylphenyl]-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 261.0 $(M + H)^+$.

Example 268

***N*-(5-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}-2-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AA and Scheme AJ using 9-ethyl-9*H*-carbazole-3-carbaldehyde and 2-methyl-*N*-[2-methyl-5-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 468.1 $(M + H)^+$.

Example 269

***N*-(5-{1-[4-(3,4-DIFLUOROPHENOXY)BENZYL]-4-PIPERIDINYL}-2-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AA and Scheme AJ using 4-(3,4-difluorophenoxy)benzaldehyde and 2-methyl-*N*-[2-methyl-5-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 479.2 $(M + H)^+$.

Example 270

***N*-(3-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AA and Scheme AJ using 9-ethyl-9*H*-carbazole-3-carbaldehyde and 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 468.1 $(M + H)^+$.

Example 271

2-METHYL-*N*-[2-METHYL-5-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure G and Scheme AI using 1-[4-chloro-1-(4-fluorophenyl)butyl]-4-fluorobenzene and 2-methyl-*N*-[2-methyl-5-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 505.1 $(M + H)^+$.

Example 272

***N*-(3-{1-[(3*S*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-(3-{[(1*S*)-3-chloro-1-phenylpropyl]oxy}phenyl)ethanone and 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 513.0 $(M + H)^+$.

Example 273

***N*-(5-{1-[(3*S*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}-2-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-(3-{[(1*S*)-3-chloro-1-phenylpropyl]oxy}phenyl)ethanone and 2-methyl-*N*-[2-methyl-5-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 512.9 $(M + H)^+$.

***N*-(2-IODOPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 using 2-methylpropanoyl chloride and 2-iodoaniline: ESMS *m/e*: 289.9 $(M + H)^+$.

***TERT*-BUTYL 4-[2-(ISOBUTYRYLAMINO)PHENYL]-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** Prepared by Procedure W and Scheme AF using *N*-(2-iodophenyl)-2-methylpropanamide and *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-

301

yl)-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 245.1 $(M - 100)^+$.

2-METHYL-*N*-[2-(4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedures X and Y, Schemes AG and AH, respectively using *tert*-butyl 4-[2-(isobutyrylamino)phenyl]-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 247.1 $(M + H)^+$.

Example 274

***N*-(2-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AA and Scheme AJ using 9-ethyl-9*H*-carbazole-3-carbaldehyde and 2-methyl-*N*-[2-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 454.1 $(M + H)^+$.

Example 275

***N*-(3-{1-[4,4-BIS(4-FLUOROPHENYL)BUTYL]-4-PIPERIDINYL}-4-METHYLPHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-[4-chloro-1-(4-fluorophenyl)butyl]-4-fluorobenzene and 2-methyl-*N*-[4-methyl-3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 505.0 $(M + H)^+$.

Example 276

***N*-(2-{1-[4,4-BIS(4-FLUOROPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-[4-chloro-1-(4-fluorophenyl)butyl]-4-fluorobenzene and 2-methyl-*N*-[2-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 490.9 $(M + H)^+$.

***N*-[2-BROMO-4-(TRIFLUOROMETHOXY)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 using 2-methylpropanoyl chloride and 2-bromo-4-(trifluoromethoxy)aniline: ESMS *m/e*: 325.9 $(M + H)^+$.

***TERT*-BUTYL 4-[2-(ISOBUTYRYLAMINO)-5-(TRIFLUOROMETHOXY)PHENYL]-3,6-DIHYDRO-1(2*H*)-PYRIDINECARBOXYLATE:** Prepared by Procedure W and Scheme AF using *N*-[2-bromo-4-(trifluoromethoxy)phenyl]-2-methylpropanamide and *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 329.0 $(M - 100)^+$.

2-METHYL-*N*-[2-(4-PIPERIDINYL)-4-(TRIFLUOROMETHOXY)PHENYL]PROPANAMIDE: Prepared by Procedures X and Y, Schemes AG and AH, respectively using *tert*-butyl 4-[2-(isobutyrylamino)-5-(trifluoromethoxy)phenyl]-3,6-dihydro-1(2*H*)-pyridinecarboxylate: ESMS *m/e*: 330.9 $(M + H)^+$.

Example 277

***N*-[2-{1-[4,4-BIS(4-FLUOROPHENYL)BUTYL]-4-PIPERIDINYL}-4-(TRIFLUOROMETHOXY)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme AI using 1-[4-chloro-1-(4-fluorophenyl)butyl]-4-fluorobenzene and 2-methyl-*N*-[2-(4-piperidinyl)-4-(trifluoromethoxy)phenyl]propanamide: ESMS *m/e*: 574.8 $(M + H)^+$.

***N*-{3-[1-(4-HYDROXYBUTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 4-chloro-1-butanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 319.3 $(M + H)^+$.

***N*-{3-[1-(5-HYDROXYPENTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 5-chloro-1-pentanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 333.3 $(M + H)^+$.

***N*-{3-[1-(6-HYDROXYHEXYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 6-chloro-1-hexanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 347.3 $(M + H)^+$.

***N*-{3-[1-(3-HYDROXYPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 3-chloro-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 305.3 $(M + H)^+$.

***N*-(3-{1-[(2*S*)-2-HYDROXY-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using (1*S*)-2-chloro-1-phenylethanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 367.2 $(M + H)^+$.

***N*-(3-{1-[(2*R*)-2-HYDROXY-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using (1*R*)-2-chloro-1-phenylethanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 367.2 $(M + H)^+$.

***N*-(3-{1-[(2*S*)-3-HYDROXY-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by

Procedure G and Scheme B1 using (2*R*)-3-chloro-2-methyl-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 319.2 $(M + H)^+$.

N-(3-{1-[(2R)-3-HYDROXY-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure G and Scheme B1 using (2*S*)-3-chloro-2-methyl-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 319.2 $(M + H)^+$.

Example 278

N-(3-{1-[(3R)-3-HYDROXY-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE: Prepared by Procedure G and Scheme B1 using (1*R*)-3-chloro-1-phenyl-1-propanol and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 379.2 $(M + H)^+$.

Example 279

N-{3-[1-(4-HYDROXY-4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure L and Scheme AN, Step 1 using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide: Anal. Calcd for $C25H34N2O2+0.08CHCl_3$: C, 74.5; H, 8.50; N, 6.93. Found: C, 74.5; H, 8.63; N, 6.81; ESMS *m/e*: 395.2 $(M + H)^+$.

Example 280

N-{3-[1-(5-HYDROXY-5-PHENYLPENTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure L and Scheme AN, Step 1 using 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide: Anal. Calcd for

$C_{26}H_{36}N_2O_2+0.25CHCl_3$: C, 71.9; H, 8.33; N, 6.39. Found: C, 71.3; H, 8.96; N, 6.86; ESMS *m/e*: 409.2 $(M + H)^+$.

Example 281

***N*-{3-[1-(6-HYDROXY-6-PHENYLHEXYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN, Step 1 using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide: Anal. Calcd for $C_{27}H_{38}N_2O_2+0.1CHCl_3$: C, 75.5; H, 8.93; N, 6.50. Found: C, 75.3; H, 8.52; N, 6.00; ESMS *m/e*: 423.2 $(M + H)^+$.

Example 282

***N*-{3-[1-(7-HYDROXY-7-PHENYLHEPTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN, Step 1 using 2-methyl-*N*-{3-[1-(7-oxo-7-phenylheptyl)-4-piperidinyl]phenyl}propanamide: Anal. Calcd for $C_{28}H_{40}N_2O_2+0.1CHCl_3$: C, 75.8; H, 9.10; N, 6.29. Found: C, 75.1; H, 9.24; N, 6.51; ESMS *m/e*: 437.1 $(M + H)^+$.

Example 283

***N*-(3-{1-[4-(4-FLUOROPHENYL)-4-HYDROXYBUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN, Step 1 using *N*-(3-{1-[4-(4-fluorophenyl)-4-oxobutyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 413.1 $(M + H)^+$.

Example 284

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLBUTYL 3-(2,6-DICHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxy-4-phenylbutyl)-

4-piperidinyl]phenyl}-2- methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.56 (m, 1H), 7.47 (m, 2H), 7.44-7.39 (m, 3H), 7.25 (m, 2H), 7.09 (s, 1H), 7.03 (m, 2H), 6.95 (m, 1H), 6.83 (m, 1H), 5.75 (t, 1H, J = 7.1 Hz), 3.03 (t, 2H, J = 7.2 Hz), 2.93 (m, 2H), 2.78 (s, 3H), 2.48 (m, 3H), 2.25 (m, 2H), 1.48 (m, 3H), 1.77 (m, 2H), 1.54 (m, 2H), 1.25 (d, 6H, J = 7.3 Hz); ESMS *m/e*: 647.7 $(M + H)^+$.

Example 285

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLBUTYL (4-FLUOROPHENYL)ACETATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxy-4-phenylbutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and (4-fluorophenyl)acetyl chloride: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.45 (s, 1H), 7.34-7.19 (m, 8H), 7.11 (m, 1H), 6.98 (m, 3H), 5.75 (t, 1H, J = 6.8 Hz), 3.61 (s, 2H), 2.92 (d, 2H, J = 8.1 Hz), 2.48 (m, 2H), 2.31 (m, 2H), 1.99-1.84 (m, 4H), 1.84-1.67 (m, 5H), 1.55-1.35 (m, 2H), 1.25 (d, 6H, J = 6.9 Hz); ESMS *m/e*: 531.1 $(M + H)^+$.

Example 286

3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL (4-FLUOROPHENYL)ACETATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(3-hydroxypropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and (4-fluorophenyl)acetyl chloride: ESMS *m/e*: 441.3 $(M + H)^+$.

Example 287

3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL 3-(2-CHLORO-6-FLUOROPHENYL)-5-METHYL-4-

ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(3-hydroxypropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chloro-6-fluorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 542.2 $(M + H)^+$.

Example 288

3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL 3-(2,6-DICHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(3-hydroxypropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 558.2 $(M + H)^+$.

Example 289

3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL 3-(2-CHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(3-hydroxypropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 524.2 $(M + H)^+$.

Example 290

(1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL 3-(2,6-DICHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-(3-{1-[(3*S*)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 633.6 $(M + H)^+$.

Example 291

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL 3-(2-CHLORO-6-FLUOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxybutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chloro-6-fluorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: Anal. Calcd for $C_{30}H_{35}ClFN_3O_4+CH_2Cl_2$: C, 63.3; H, 6.23; N, 7.33. Found: C, 63.0; H, 6.39; N, 7.03; ESMS *m/e*: 556.2 $(M + H)^+$.

Example 292

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL 3-(2-CHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxybutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 538.2 $(M + H)^+$.

Example 293

3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL 5-METHYL-3-PHENYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(3-hydroxypropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-methyl-3-phenyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 490.3 $(M + H)^+$.

Example 294

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL 3-(2,6-DICHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxybutyl)-4-piperidinyl]phenyl}-2-

309

methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 572.2 $(M + H)^+$.

Example 295

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLBUTYL 3-(2-CHLORO-6-FLUOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxy-4-phenylbutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chloro-6-fluorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: Anal. Calcd for $C_{36}H_{39}ClFN_3O_4+0.54CHCl_3$: C, 63.0; H, 5.72; N, 6.03. Found: C, 63.0; H, 5.54; N, 6.05; ESMS *m/e*: 632.2 $(M + H)^+$.

Example 296

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL 5-METHYL-3-PHENYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxybutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-methyl-3-phenyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 504.3 $(M + H)^+$.

Example 297

6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL 3-(2,6-DICHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(6-hydroxyhexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 600.0 $(M + H)^+$.

Example 298

6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL 5-METHYL-3-PHENYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(6-hydroxyhexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-methyl-3-phenyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 532.1 $(M + H)^+$.

Example 299

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL (4-FLUOROPHENYL)ACETATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxybutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and (4-fluorophenyl)acetyl chloride: ESMS *m/e*: 455.3 $(M + H)^+$.

Example 300

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLBUTYL 3-(2-CHLOROPHENYL)-5-METHYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxy-4-phenylbutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 614.2 $(M + H)^+$.

Example 301

4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLBUTYL 5-METHYL-3-PHENYL-4-ISOXAZOLECARBOXYLATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-{3-[1-(4-hydroxy-4-phenylbutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-methyl-3-phenyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 580.0 $(M + H)^+$.

Example 302

(1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL (4-FLUOROPHENYL)ACETATE: Prepared by Procedure Q1 and Scheme C2 (TEA) using *N*-(3-{1-[(3*S*)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and (4-fluorophenyl)acetyl chloride: Anal. Calcd for $C_{32}H_{37}FN_2O_3+0.07CHCl_3$: C, 73.4; H, 7.12; N, 5.34. Found: C, 73.4; H, 6.96; N, 5.14; ESMS *m/e*: 517.1 $(M + H)^+$.

Example 303

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)BENZAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and benzoyl chloride: Anal. Calcd for $C_{31}H_{37}N_3O_2+0.55CHCl_3$: C, 69.0; H, 6.89; N, 7.65. Found: C, 69.7; H, 6.73; N, 6.03; ESMS *m/e*: 484.4 $(M + H)^+$.

Example 304

***N*-[3-(1-{(3*S*)-3-[(DIPHENYLACETYL)AMINO]-3-PHENYLPROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and diphenylacetyl chloride: ESMS *m/e*: 574.3 $(M + H)^+$.

Example 305

3-CHLORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)BENZAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 3-chlorobenzoyl chloride: ESMS *m/e*: 518.3 $(M + H)^+$.

Example 306

3,5-DICHLORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)BENZAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3,5-dichlorobenzoyl chloride: ESMS *m/e*: 552.3 $(M + H)^+$.

Example 307

2-(ETHYLSULFANYL)-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)NICOTINAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-(ethylsulfanyl)nicotinoyl chloride: ESMS *m/e*: 545.3 $(M + H)^+$.

Example 308

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)[1,1'-BIPHENYL]-4-CARBOXAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and [1,1'-biphenyl]-4-carbonyl chloride: ESMS *m/e*: 560.3 $(M + H)^+$.

Example 309

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-PYRIDINECARBOXAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 2- pyridinecarbonyl chloride: ESMS *m/e*: 484.6 $(M + H)^+$.

Example 310

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-METHOXYBENZAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-methoxybenzoyl chloride: ESMS *m/e*: 514.1 $(M + H)^+$.

Example 311

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-1-NAPHTHAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthoyl chloride: ESMS *m/e*: 533.7 $(M + H)^+$.

Example 312

2,4-DIFLUORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)BENZAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2,4-difluorobenzoyl chloride: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 313

3-(2-CHLORO-6-FLUOROPHENYL)-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide

314

and 3-(2-chloro-6- fluorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 617.2 $(M + H)^+$.

Example 314

3-CHLORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-THIOPHENECARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3-chloro-2-thiophenecarbonyl chloride: ESMS *m/e*: 524.2 $(M + H)^+$.

Example 315

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-PHENOXYNICOTINAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-phenoxynicotinoyl chloride: ESMS *m/e*: 577.3 $(M + H)^+$.

Example 316

1-(4-CHLOROPHENYL)-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-3-PROPYL-1*H*-PYRAZOLE-4-CARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(4-chlorophenyl)-3-propyl-1*H*-pyrazole-4-carbonyl chloride: ESMS *m/e*: 626.3 $(M + H)^+$.

Example 317

4-CHLORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-1,3-DIMETHYL-1*H*-PYRAZOLO[3,4-*B*]PYRIDINE-5-CARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-

315

3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-chloro-1,3-dimethyl-1*H*-pyrazolo[3,4-*b*]pyridine-5-carbonyl chloride: ESMS *m/e*: 587.3 $(M + H)^+$.

Example 318

5-(3,5-DICHLOROPHENOXY)-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-1*H*-PYRROLE-2-CARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 5-(3,5-dichlorophenoxy)-1*H*-pyrrole-2-carbonyl chloride: ESMS *m/e*: 634.2 $(M + H)^+$.

Example 319

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)NICOTINAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and nicotinoyl chloride: ESMS *m/e*: 485.3 $(M + H)^+$.

Example 320

3,4-DIFLUORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)BENZAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3,4-difluorobenzoyl chloride: ESMS *m/e*: 520.3 $(M + H)^+$.

Example 321

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-1-PHENYL-3-PROPYL-1*H*-PYRAZOLE-4-CARBOXAMIDE:** Prepared by Procedure Q1 and Scheme AC

316

using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-phenyl-3-propyl-1*H*-pyrazole-4-carbonyl chloride: ESMS *m/e*: 592.2 $(M + H)^+$.

Example 322

4-(DIMETHYLAMINO)-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)BENZAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(dimethylamino)benzoyl chloride: ESMS *m/e*: 527.3 $(M + H)^+$.

Example 323

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-THIOPHENECARBOXAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 2-thiophenecarbonyl chloride: ESMS *m/e*: 490.2 $(M + H)^+$.

Example 324

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-5-NITRO-2-FURAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 5-nitro-2-furoyl chloride: ESMS *m/e*: 519.2 $(M + H)^+$.

Example 325

***N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-5-METHYL-3-PHENYL-4-**

ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-methyl-3-phenyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 489.1 $(M + H)^+$.

Example 326

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-FURAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-furoyl chloride: ESMS *m/e*: 474.2 $(M + H)^+$.

Example 327

***N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-1-(4-NITROPHENYL)-5-(TRIFLUOROMETHYL)-1*H*-PYRAZOLE-4-CARBOXAMIDE:** Prepared by Procedure Q1 and Scheme AC using *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(4-nitrophenyl)-5-(trifluoromethyl)-1*H*-pyrazole-4-carbonyl chloride: ESMS *m/e*: 663.2 $(M + H)^+$.

Example 328

3-(2-CHLORO-6-FLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chloro-6-fluorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 541.2 $(M + H)^+$.

Example 329

***N*-[3-(1-{3-[(DIPHENYLACETYL)AMINO]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by

Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and diphenylacetyl chloride: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.51 (s, 1H), 7.33-7.21 (m, 13H), 6.94 (m, 2H), 4.88 (s, 1H), 3.39 (t, 2H, J = 5.6 Hz), 2.93 (d, 2H, J = 11.3 Hz), 2.52-2.36 (m, 4H), 1.97 (t, 2H, J = 11.3 Hz), 1.83-1.58 (m, 6H), 1.24 (d, 6H, J = 7.6 Hz); Anal. Calcd for $C_{32}H_{39}N_3O_2$+HCl+0.19$CHCl_3$: C, 69.44; H, 7.27; N, 7.55. Found: C, 69.44; H, 7.43; N, 7.43; ESMS *m/e*: 498.4 (M + H)$^+$.

Example 330

***N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-1-BENZOTHIOPHENE-3-CARBOXAMIDE:**

Prepared by Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 1-benzothiophene-3-carbonyl chloride: ESMS *m/e*: 464.2 (M + H)$^+$.

Example 331

3-(2-CHLOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE:

Prepared by Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 523.1 (M + H)$^+$.

Example 332

3-(2,6-DICHLOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2,6-

dichlorophenyl)-5-methyl- 4-isoxazolecarbonyl chloride: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.50 (d, 1H, J = 2.3 Hz), 7.48 (s, 1H), 7.4 (m, 1H), 7.39 (s, 1H), 7.37 (m, 2H), 7.24 (t, 1H, J = 7.2 Hz), 6.92 (d, 1H, J = 7.9 Hz), 6.06 (s, 1H), 3.31 (q, 2H, J = 6.4 Hz), 2.94 (d, 2H, J = 10.8 Hz), 2.79 (s, 3H), 2.53 (q, 1H, J = 6.1), 2.47 (tt, 1H, J = 4.2, 11.4 Hz), 2.29 (t, 2H, J = 7.2 Hz), 1.99 (t, 2H, J = 11.4 Hz), 1.81 (m, 2H), 1.69 (dt, 2H, J = 2.4, 11.6), 1.59 (q, 2H, J = 6.6 Hz), 1.24 (d, 6H, J = 6.5 Hz); ESMS *m/e*: 557.0 $(M + H)^+$.

1-[3-(3-CHLOROPROPOXY)PHENYL]ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-3-chloropropane.

1-(3-CHLOROPROPOXY)-2-FLUOROBENZENE: Prepared by Procedure U and Scheme AK using 2-fluorophenol and 1-bromo-3-chloropropane.

1-CHLORO-3-(3-CHLOROPROPOXY)BENZENE: Prepared by Procedure U and Scheme AK using 3-chlorophenol and 1-bromo-3-chloropropane.

1-CHLORO-4-(3-CHLOROPROPOXY)BENZENE: Prepared by Procedure U and Scheme AK using 4-chlorophenol and 1-bromo-3-chloropropane.

1-(3-CHLOROPROPOXY)-3-FLUOROBENZENE: Prepared by Procedure U and Scheme AK using 3-fluorophenol and 1-bromo-3-chloropropane.

1-(3-CHLOROPROPOXY)-4-FLUOROBENZENE: Prepared by Procedure U and Scheme AK using 4-fluorophenol and 1-bromo-3-chloropropane.

1-CHLORO-2-(3-CHLOROPROPOXY)BENZENE: Prepared by Procedure U and Scheme AK using 2-chlorophenol and 1-bromo-3-chloropropane.

4-(3-CHLOROPROPOXY)-1,2-DIMETHYLBENZENE: Prepared by Procedure U and Scheme AK using 3,4-dimethylphenol and 1-bromo-3-chloropropane.

1-BROMO-2-(3-CHLOROPROPOXY)BENZENE: Prepared by Procedure U and Scheme AK using 2-bromophenol and 1-bromo-3-chloropropane.

1-BROMO-3-(3-CHLOROPROPOXY)BENZENE: Prepared by Procedure U and Scheme AK using 3-bromophenol and 1-bromo-3-chloropropane.

1-BROMO-4-(3-CHLOROPROPOXY)BENZENE: Prepared by Procedure U and Scheme AK using 4-bromophenol and 1-bromo-3-chloropropane.

1-(3-CHLOROPROPOXY)-4-METHYLBENZENE: Prepared by Procedure U and Scheme AK using *p*-cresol and 1-bromo-3-chloropropane.

4-BROMOPHENYL (2*R*)-3-CHLORO-2-METHYLPROPYL ETHER: Prepared by Procedure U and Scheme AK using 4-bromophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}-2,4,5-TRIFLUOROBENZENE: Prepared by Procedure U and Scheme AK

using 2,4,5- trifluorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-CHLORO-3-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}BENZENE:

Prepared by Procedure U and Scheme AK using 3-chlorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}-4-FLUOROBENZENE:

Prepared by Procedure U and Scheme AK using 4-fluorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}-3-FLUOROBENZENE:

Prepared by Procedure U and Scheme AK using 3-fluorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-CHLORO-2-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}BENZENE:

Prepared by Procedure U and Scheme AK using 2-chlorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}-2-FLUOROBENZENE:

Prepared by Procedure U and Scheme AK using 2-fluorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-CHLORO-4-{[(2*R*)-3-CHLORO-2-METHYLPROPYL]OXY}BENZENE:

Prepared by Procedure U and Scheme AK using 4-chlorophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

3-BROMOPHENYL (2*R*)-3-CHLORO-2-METHYLPROPYL ETHER:

Prepared by Procedure U and Scheme AK using 3-bromophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

2-BROMOPHENYL (2*R*)-3-CHLORO-2-METHYLPROPYL ETHER:

Prepared by Procedure U and Scheme AK using 2-bromophenol and (2*S*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*S*)-3-CHLORO-2-METHYLPROPYL]OXY}-3-FLUOROBENZENE:
Prepared by Procedure U and Scheme AK using 3-fluorophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*S*)-3-CHLORO-2-METHYLPROPYL]OXY}-4-FLUOROBENZENE:
Prepared by Procedure U and Scheme AK using 4-fluorophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

1-{[(2*S*)-3-CHLORO-2-METHYLPROPYL]OXY}-2-FLUOROBENZENE:
Prepared by Procedure U and Scheme AK using 2-fluorophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

1-CHLORO-2-{[(2*S*)-3-CHLORO-2-METHYLPROPYL]OXY}BENZENE:
Prepared by Procedure U and Scheme AK using 2-chlorophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

1-CHLORO-4-{[(2*S*)-3-CHLORO-2-METHYLPROPYL]OXY}BENZENE:
Prepared by Procedure U and Scheme AK using 4-chlorophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

4-BROMOPHENYL (2*S*)-3-CHLORO-2-METHYLPROPYL ETHER:
Prepared by Procedure U and Scheme AK using 4-bromophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

3-BROMOPHENYL (2*S*)-3-CHLORO-2-METHYLPROPYL ETHER:
Prepared by Procedure U and Scheme AK using 3-bromophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

2-BROMOPHENYL (2*S*)-3-CHLORO-2-METHYLPROPYL ETHER:
Prepared by Procedure U and Scheme AK using 2-bromophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

1-CHLORO-3-{[(2*S*)-3-METHYLPROPYL]OXY}BENZENE: Prepared by Procedure U and Scheme AK using 3-chlorophenol and (2*R*)-1-bromo-3-chloro-2-methylpropane.

1-[3-(4-CHLOROBUTOXY)PHENYL]ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-4-chlorobutane.

1-[3-(4-CHLOROBUTOXY)PHENYL]ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-4-chlorobutane.

1-(4-CHLOROBUTOXY)-3-METHOXYBENZENE: Prepared by Procedure U and Scheme AK using 3-methoxyphenol and 1-bromo-4-chlorobutane.

1-(4-CHLOROBUTOXY)-4-METHOXYBENZENE: Prepared by Procedure U and Scheme AK using 4-methoxyphenol and 1-bromo-4-chlorobutane.

1-(4-CHLOROBUTOXY)-2-METHOXYBENZENE: Prepared by Procedure U and Scheme AK using 2-methoxyphenol and 1-bromo-4-chlorobutane.

4-(4-CHLOROBUTOXY)-1,2-DIMETHYLBENZENE: Prepared by Procedure U and Scheme AK using 3,4-dimethylphenol and 1-bromo-4-chlorobutane.

1-{3-[(5-CHLOROPENTYL)OXY]PHENYL}ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-5-chloropentane.

1-{3-[(5-CHLOROPENTYL)OXY]PHENYL}ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-5-chloropentane.

1-{3-[(6-CHLOROHEXYL)OXY]PHENYL}ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-6-chlorohexane.

1-{3-[(6-CHLOROHEXYL)OXY]PHENYL}ETHANONE: Prepared by Procedure U and Scheme AK using 1-(3-hydroxyphenyl)ethanone and 1-bromo-6-chlorohexane.

Example 333

***N*-(3-{1-[(2*S*)-2-(3-ACETYLPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 1-(3-hydroxyphenyl)ethanone and *N*-(3-{1-[(2*R*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 485.0 $(M + H)^+$.

Example 334

***N*-(3-{1-[(2*S*)-2-(2-ACETYLPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 1-(2-hydroxyphenyl)ethanone and *N*-(3-{1-[(2*R*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 485.2 $(M + H)^+$.

Example 335

***N*-(3-{1-[(2*S*)-2-(3-CHLOROPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 3-chlorophenol and *N*-(3-

{1-[(2*R*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 477.1 $(M + H)^+$.

Example 336

***N*-(3-{1-[(2*S*)-2-(3,4-DIMETHOXYPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 3,4-dimethoxyphenol and *N*-(3-{1-[(2*R*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 503.2 $(M + H)^+$.

Example 337

***N*-(3-{1-[(2*R*)-2-(4-FLUOROPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 4-fluorophenol and *N*-(3-{1-[(2*S*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 461.2 $(M + H)^+$.

Example 338

***N*-(3-{1-[(2*R*)-2-(3-METHOXYPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 3-methoxyphenol and *N*-(3-{1-[(2*S*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 472.9 $(M + H)^+$.

Example 339

***N*-(3-{1-[(2*R*)-2-(3-CHLOROPHENOXY)-2-PHENYLETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme B1 using 3-chlorophenol and *N*-(3-{1-[(2*S*)-2-hydroxy-2-phenylethyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 478.5 $(M + H)^+$.

***N*-{3-[1-(3,3-DIMETHOXYPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 3-bromo-1,1-dimethoxypropane and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 349.2 $(M + H)^+$.

Example 340

***N*-(3-{1-[(3*S*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure B and Scheme B1 using 1-(3-hydroxyphenyl)ethanone and *N*-(3-{1-[(3*R*)-3-hydroxy-3-phenylpropyl]-4-piperidinyl}phenyl)cyclopropanecarboxamide: ESMS *m/e*: 497.1 $(M + H)^+$.

Example 341

***N*-(3-{1-[3-(3-ACETYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-[3-(3-chloropropoxy)phenyl]ethanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 423.2 $(M + H)^+$.

Example 342

***N*-(3-{1-[3-(3-ACETYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-[3-(3-chloropropoxy)phenyl]ethanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 421.2 $(M + H)^+$.

Example 343

***N*-(3-{1-[3-(2-FLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by

Procedure G and Scheme B1 using 1-(3-chloropropoxy)-2-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 399.2 $(M + H)^+$.

Example 344

***N*-(3-{1-[3-(3-CHLOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-3-(3-chloropropoxy)benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 415.2 $(M + H)^+$.

Example 345

***N*-(3-{1-[3-(4-CHLOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-4-(3-chloropropoxy)benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.71 (dd, 1H, J = 3.2, 5.7 Hz), 7.53 (dd, 1H, J = 3.2, 5.7 Hz), 7.50 (m, 1H), 7.31 (m, 1H), 7.24-7.20 (m, 2H), 6.94 (d, 1H, J = 7.9 Hz), 6.85-6.82 (m, 2H), 4.00 (t, 2H, J = 6.1 Hz), 3.07 (d, 2H, J = 10.9 Hz), 2.55 (m, 3H), 2.50 (sept, 1H, J = 6.2 Hz), 2.08 (dt, 2H, J = 3.1, 10.9 Hz), 2.00 (m, 2H), 1.83 (m, 3H), 1.69 (qt, 1H, J = 6.2 Hz), 1.24 (d, 6H, J = 6.8 Hz); Anal. Calcd for $C_{24}H_{31}ClN_2O_2$+HCl: C, 63.8; H, 7.09; N, 6.21. Found: C, 63.3; H, 7.04; N, 6.27; ESMS *m/e*: 415.2 $(M + H)^+$.

Example 346

***N*-(3-{1-[3-(3-FLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-(3-chloropropoxy)-3-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 399.2 $(M + H)^+$.

Example 347

***N*-(3-{1-[3-(4-FLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-(3-chloropropoxy)-4-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 399.2 $(M + H)^+$.

Example 348

***N*-(3-{1-[3-(2-CHLOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-2-(3-chloropropoxy)benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 415.2 $(M + H)^+$.

Example 349

***N*-(3-{1-[3-(3,4-DIMETHYLPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 4-(3-chloropropoxy)-1,2-dimethylbenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 409.2 $(M + H)^+$.

Example 350

***N*-(3-{1-[3-(2-BROMOPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-bromo-2-(3-chloropropoxy)benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.53 (dd, 1H, J = 1.6, 7.9 Hz), 7.48 (s, 1H), 7.32 (m, 1H), 7.28-7.22 (m, 3H), 7.17 (s, 1H), 6.98 (d, 1H, J = 7.7 Hz), 6.93 (dd, 1H, J = 1.4, 8.4 Hz), 6.82 (dt, 1H, J = 7.6, 1.4 Hz), 4.11 (t, 2H, J = 6.3 Hz), 3.07 (d, 2H, J = 11.3 Hz), 2.61 (t, 2H, J = 6.9 Hz), 2.50 (m, 3H), 2.07

(m, 1H), 1.8-1.75 (m, 5H), 1.25 (d, 6H, J = 6.7 Hz); Anal. Calcd for $C_{24}H_{31}BrN_2O_2.HCl+0.2\ CHCl_3$: C, 55.9; H, 6.24; N, 5.39. Found: C, 55.8; H, 6.23; N, 5.47; ESMS *m/e*: 459.1 $(M + H)^+$.

Example 351

***N*-(3-{1-[3-(3-BROMOPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-bromo-3-(3-chloropropoxy)benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 459.1 $(M + H)^+$.

Example 352

***N*-(3-{1-[3-(4-BROMOPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-bromo-4-(3-chloropropoxy)benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.51 (s, 1H), 7.37 (d, 2H, J = 7.6 Hz), 7.26 (m, 3H), 6.97 (d, 1H, J = 7.7 Hz), 6.79 (d, 2H, J = 7.7 Hz), 4.01 (t, 2H, J = 5.6 Hz), 3.08 (d, 2H, J = 9.4 Hz), 2.53 (m, 4H), 2.05 (m, 4H), 1.84 (m, 4H), 1.24 (d, 6H, J = 5.9 Hz); Anal. Calcd for $C_{24}H_{31}BrN_2O_2.HCl+0.34CHCl_3$: C, 54.5; H, 6.08; N, 5.22. Found: C, 54.5; H, 6.22; N, 5.22; ESMS *m/e*: 459.1 $(M + H)^+$.

Example 353

***N*-(3-{1-[(3*R*)-3-(3,4-DIMETHOXYPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-N,2-DIMETHYLPROPANAMIDE:** Prepared by Procedure T and Scheme AD using *N*-(3-{1-[(3*R*)-3-(3,4-dimethoxyphenoxy)-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and methyl iodide: ESMS *m/e*: 531.2 $(M + H)^+$.

Example 354

***N*-(3-{1-[(3*R*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-N,2-DIMETHYLPROPANAMIDE:** Prepared by Procedure T and Scheme AD using *N*-(3-{1-[(3*R*)-3-(3-acetylphenoxy)-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and methyl iodide: ESMS *m/e*: 513.2 (M + H)$^+$.

Example 355

***N*-(3-{1-[(3*S*)-3-(3-ACETYLPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-N,2-DIMETHYLPROPANAMIDE:** Prepared by Procedure T and Scheme AD using *N*-(3-{1-[(3*S*)-3-(3-acetylphenoxy)-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide and methyl iodide: ESMS *m/e*: 513.2 (M + H)$^+$.

Example 356

***N*-(3-{1-[(2*S*)-3-(4-BROMOPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 4-bromophenyl (2*R*)-3-chloro-2-methylpropyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.0 (M + H)$^+$.

Example 357

2-METHYL-*N*-(3-{1-[(2*S*)-2-METHYL-3-(2,4,5-TRIFLUOROPHENOXY)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure G and Scheme B1 using 1-{[(2*R*)-3-chloro-2-methylpropyl]oxy}-2,4,5-trifluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 449.2 (M + H)$^+$.

Example 358

***N*-(3-{1-[(2*S*)-3-(3-CHLOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-3-{[(2*R*)-3-chloro-2-methylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.2 $(M + H)^+$.

Example 359

***N*-(3-{1-[(2*S*)-3-(4-FLUOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{[(2*R*)-3-chloro-2-methylpropyl]oxy}-4-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.2 $(M + H)^+$.

Example 360

***N*-(3-{1-[(2*S*)-3-(3-FLUOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{[(2*R*)-3-chloro-2-methylpropyl]oxy}-3-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.2 $(M + H)^+$.

Example 361

***N*-(3-{1-[(2*S*)-3-(2-CHLOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-2-{[(2*R*)-3-chloro-2-methylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.1 $(M + H)^+$.

Example 362

***N*-(3-{1-[(2*S*)-3-(2-FLUOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{[(2*R*)-3-chloro-2-methylpropyl]oxy}-2-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.2 $(M + H)^+$.

Example 363

***N*-(3-{1-[(2*S*)-3-(4-CHLOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-4-{[(2*R*)-3-chloro-2-methylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.2 $(M + H)^+$.

Example 364

***N*-(3-{1-[(2*S*)-3-(3-BROMOPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 3-bromophenyl (2*R*)-3-chloro-2-methylpropyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 474.0 $(M + H)^+$.

Example 365

***N*-(3-{1-[(2*S*)-3-(2-BROMOPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 2-bromophenyl (2*R*)-3-chloro-2-methylpropyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.0 $(M + H)^+$.

Example 366

***N*-(3-{1-[(2*R*)-3-(3-FLUOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{[(2*S*)-3-chloro-2-methylpropyl]oxy}-3-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.2 $(M + H)^+$.

Example 367

***N*-(3-{1-[(2*R*)-3-(4-FLUOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{[(2*S*)-3-chloro-2-

methylpropyl]oxy}-4-fluorobenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.8 $(M + H)^+$.

Example 368

***N*-(3-{1-[(2*R*)-3-(2-CHLOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-2-{[(2*S*)-3-chloro-2-methylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.1 $(M + H)^+$.

Example 369

***N*-(3-{1-[(2*R*)-3-(4-CHLOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-4-{[(2*S*)-3-chloro-2-methylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.1 $(M + H)^+$.

Example 370

***N*-(3-{1-[(2*R*)-3-(4-BROMOPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 4-bromophenyl (2*S*)-3-chloro-2-methylpropyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.0 $(M + H)^+$.

Example 371

***N*-(3-{1-[(2*R*)-3-(3-BROMOPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 3-bromophenyl (2*S*)-3-chloro-2-methylpropyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.0 $(M + H)^+$.

Example 372

***N*-(3-{1-[(2*R*)-3-(2-BROMOPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 2-bromophenyl (2*S*)-3-chloro-2-methylpropyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 473.0 $(M + H)^+$.

Example 373

***N*-(3-{1-[(2*R*)-3-(3-CHLOROPHENOXY)-2-METHYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-chloro-3-{[(2*S*)-3-chloro-2-methylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 429.1 $(M + H)^+$.

Example 374

***N*-(3-{1-[3-(5,5-DIMETHYL-1,3-DIOXAN-2-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 2-(3-bromopropyl)-5,5-dimethyl-1,3-dioxane and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 403.2 $(M + H)^+$

Example 375

***N*-(3-{1-[4-(3-ACETYLPHENOXY)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-[3-(4-chlorobutoxy)phenyl]ethanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 437.2 $(M + H)^+$.

Example 376

***N*-(3-{1-[4-(3-METHOXYPHENOXY)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-(4-chlorobutoxy)-3-methoxybenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 377

***N*-(3-{1-[4-(4-METHOXYPHENOXY)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-(4-chlorobutoxy)-4-methoxybenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 378

***N*-(3-{1-[4-(2-METHOXYPHENOXY)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-(4-chlorobutoxy)-2-methoxybenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 379

***N*-(3-{1-[4-(3,4-DIMETHYLPHENOXY)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 4-(4-chlorobutoxy)-1,2-dimethylbenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 423.2 $(M + H)^+$.

Example 380

***N*-(3-{1-[4-(1,3-DIOXOLAN-2-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 2-(4-chlorobutyl)-1,3-dioxolane and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 375.2 $(M + H)^+$

Example 381

***N*-(3-{1-[5-(3-ACETYLPHENOXY)PENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{3-[(5-

chloropentyl)oxy]phenyl}ethanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 451.3 $(M + H)^+$.

Example 382

***N*-(3-{1-[5-(3-ACETYLPHENOXY)PENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{3-[(5-chloropentyl)oxy]phenyl}ethanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 449.2 $(M + H)^+$.

Example 383

***N*-(3-{1-[6-(3-ACETYLPHENOXY)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{3-[(6-chlorohexyl)oxy]phenyl}ethanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 465.3 $(M + H)^+$.

Example 384

***N*-(3-{1-[6-(3-ACETYLPHENOXY)HEXYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure G and Scheme B1 using 1-{3-[(6-chlorohexyl)oxy]phenyl}ethanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 463.3 $(M + H)^+$.

Example 385

***N*-(3-{1-[4-(4-CHLOROPHENOXY)-4-(4-CHLOROPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure B and Scheme AN using 4-chlorophenol and *N*-(3-{1-[4-(4-chlorophenyl)-4-hydroxybutyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 562.9 $(M + 23)^+$.

Example 386

2-METHYL-N-(3-{1-[2-(1-METHYL-2-PHENYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 480.3 $(M + H)^+$.

Example 387

2-METHYL-*N*-(3-{1-[2-(2-PHENYL-1*H*-BENZO[*G*]INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide and 1-(1-naphthyl)hydrazine hydrochloride: ESMS *m/e*: 516.4 $(M + H)^+$.

Example 388

2-METHYL-*N*-(3-{1-[3-(2-PHENYL-1*H*-BENZO[*G*]INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide and 1-(1-naphthyl)hydrazine hydrochloride: ESMS *m/e*: 530.2 $(M + H)^+$.

Example 389

2-METHYL-*N*-[3-(1-{3-[2-PHENYL-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide and 1-[4-(trifluoromethoxy)phenyl]hydrazine hydrochloride: ESMS *m/e* : 564.2 $(M + H)^+$.

Example 390

2-METHYL-N-[3-(1-{4-[2-PHENYL-5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]BUTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1-[4-(trifluoromethoxy)phenyl]hydrazine hydrochloride: ESMS *m/e*: 578.2 $(M + H)^+$.

Example 391

2-METHYL-N-(3-{1-[3-(1-METHYL-2-PHENYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 495.3 $(M + H)^+$.

Example 392

N-(3-{1-[4-(1,2-DIPHENYL-1H-INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: $(M + H)^+$. 570.3**

Example 393

2-METHYL-N-[3-(1-{5-[2-PHENYL-5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(7-oxo-7-phenylheptyl)-4-piperidinyl]phenyl}propanamide and 1-[4-(trifluoromethoxy)phenyl]hydrazine hydrochloride: ESMS *m/e*: 592.3 $(M + H)^+$.

Example 394

N-(3-{1-[5-(1,2-DIPHENYL-1H-INDOL-3-YL)PENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(7-oxo-7-phenylheptyl)-4-piperidinyl]phenyl}propanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 584.3 (M + H)$^+$.**

Example 395

2-METHYL-N-(3-{1-[5-(1-METHYL-2-PHENYL-1H-INDOL-3-YL)PENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(7-oxo-7-phenylheptyl)-4-piperidinyl]phenyl}propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 522.3 (M + H)$^+$.**

Example 396

2-METHYL-N-(3-{1-[4-(2-PHENYL-1H-BENZO[G]INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1-(1-naphthyl)hydrazine hydrochloride: ESMS *m/e*: 544.3 (M + H)$^+$.**

Example 397

2-METHYL-N-(3-{1-[4-(1-METHYL-2-PHENYL-1H-INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 508.3 (M + H)$^+$.

Example 398

2-METHYL-N-(3-{1-[5-(2-PHENYL-1H-BENZO[G]INDOL-3-YL)PENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(7-**

oxo-7-phenylheptyl)-4-piperidinyl]phenyl}propanamide and 1-(1-naphthyl)hydrazine hydrochloride: ESMS *m/e*: 558.2 (M + H)$^+$.

Example 399

2-METHYL-N-(3-{1-[2-(5-METHYL-2-PHENYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide and 1-(4-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 480.2 (M + H)$^+$.**

Example 400

N-(3-{1-[2-(7-METHOXY-2-PHENYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide and 1-(2-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 496.2 (M + H)$^+$.**

Example 401

2-METHYL-N-(3-{1-[2-(7-METHYL-2-PHENYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 480.2 (M + H)$^+$.**

Example 402

N-(3-{1-[3-(7-METHOXY-2-PHENYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: **Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide and 1-**

5-phenylpentyl)-4-piperidinyl]phenyl}propanamide and 1-(2-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 510.2 $(M + H)^+$.

Example 403

2-METHYL-N-(3-{1-[4-(7-METHYL-2-PHENYL-1H-INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 508.3 $(M + H)^+$.

Example 404

N-(3-{1-[2-(5-METHOXY-2-PHENYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(4-oxo-4-phenylbutyl)-4-piperidinyl]phenyl}propanamide and 1-(4-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 496.2 $(M + H)^+$.

Example 405

2-METHYL-N-(3-{1-[3-(5-METHYL-2-PHENYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(5-oxo-5-phenylpentyl)-4-piperidinyl]phenyl}propanamide and 1-(4-methylphenyl)hydrazine hydrochloride: ESMS ***m/e*: 494.3 $(M + H)^+$.**

Example 406

N-(3-{1-[4-(7-METHOXY-2-PHENYL-1H-INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE

Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1-(2-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 524.3 $(M + H)^+$.

Example 407

2-METHYL-N-(3-{1-[3-(1-PHENYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 480.2 $(M + H)^+$.

Example 408

2-METHYL-N-(3-{1-[2-(1-PHENYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 409

2-METHYL-N-(3-{1-[2-(7-METHYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 404.2 $(M + H)^+$.

Example 410

2-METHYL-N-(3-{1-[2-(1-METHYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure

343

H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 404.2 $(M + H)^+$.

Example 411

2-METHYL-N-(3-{1-[2-(5-METHYL-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(4-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 404.2 $(M + H)^+$.

Example 412

2-METHYL-N-[3-(1-{2-[5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]ETHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-[4-(trifluoromethoxy)phenyl]hydrazine hydrochloride: ESMS *m/e*: 474.2 $(M + H)^+$.

Example 413

N-(3-{1-[3-(1H-BENZO[G]INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(1-naphthyl)hydrazine hydrochloride: ESMS 454.2 *m/e*: $(M + H)^+$.

Example 414

2-METHYL-*N*-(3-{1-[3-(1-METHYL-1*H*-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S. A mixture of *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide (100

mg, 0.270 mmol), 1-methyl- 1-phenylhydrazine (106 mg, 0.870 mmol), $ZnCl_2$ (119 mg, 0.870 mmol) and HOAc (1.00 mL) was heated for 12 h at 80 °C. The resulting crude mixture was diluted with water (20 mL), the aqueous layer was neutralized with a saturated K_2CO_3 solution (10 mL) and extracted with CH_2Cl_2 (3 X 20 mL). The combined organic layers were concentrated *in vacuo* and the residue was purified by preparative TLC using 3 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product 2-methyl-*N*-(3-{1-[3-(1-methyl-1*H*-indol-3-yl)propyl]-4-piperidinyl}phenyl)propanamide (20.7 mg, 18.7 %): ^1H NMR (400 MHz, $CDCl_3$) δ 7.60 (d, 1H, J = 8.1 Hz), 7.45 (s, 1H), 7.35 (d, 1H, J = 7.4 Hz), 7.25 (m, 4H), 7.09 (t, 1H, J = 7.3 Hz), 6.97 (d, 1H, J = 7.3 Hz), 6.86 (s, 1H), 3.75 (s, 3H), 3.11 (d, 2H, J = 11.6 Hz), 2.79 (t, 2H, J = 7.3 Hz), 2.51 (m, 4H), 2.12-1.81 (m, 8H), 1.25 (d, 6H, J = 7.1 Hz); Anal. Calcd for $C_{27}H_{35}N_3O$+0.225 $CHCl_3$: C, 73.57; H, 7.99; N, 9.45. Found: C, 73.93; H, 7.90; N, 9.23; ESMS *m/e*: 418.2 $(M + H)^+$.

Example 415

2-METHYL-N-(3-{1-[3-(5-METHYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(4-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 418.2 $(M + H)^+$.

Example 416

2-METHYL-N-[3-(1-{3-[5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide

345

and 1-[4-(trifluoromethoxy)phenyl]hydrazine hydrochloride: ESMS *m/e*: 488.2 $(M + H)^+$.

Example 417

2-METHYL-N-(3-{1-[3-(7-METHYL-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 418.2 $(M + H)^+$.

Example 418

N-(3-{1-[3-(7-METHOXY-1H-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[4-(1,3-dioxolan-2-yl)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 434.0 $(M + H)^+$.

Example 419

N-(3-{1-[2-(7-METHOXY-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 420.2 $(M + H)^+$.

Example 420

N-(3-{1-[2-(5-METHOXY-1H-INDOL-3-YL)ETHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure H and Scheme S using *N*-(3-{1-[3-(1,3-dioxolan-2-yl)propyl]-4-piperidinyl}phenyl)-2-methylpropanamide

and 1-(4-methoxyphenyl)hydrazine hydrochloride: ESMS *m/e*: 420.2 $(M + H)^+$.

Example 421

2-METHYL-N-(3-{1-[4-(5-METHYL-2-PHENYL-1H-INDOL-3-YL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-{3-[1-(6-oxo-6-phenylhexyl)-4-piperidinyl]phenyl}propanamide and 1-(4-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 508.3 $(M + H)^+$.

Example 422

2-METHYL-N-[4-(1-{[1-(4-METHYLPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide and 1-(4-methylphenyl)-1*H*-indole: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 423

N-[4-(1-{[1-(4-METHYLPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]BUTANAMIDE: Prepared by Procedure D and Scheme N using *N*-[4-(4-piperidinyl)phenyl]butanamide and 1-(4-methylphenyl)-1*H*-indole: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 424

N-[3-(1-{[2-(2-AMINOPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 2-(1*H*-indol-2-yl)aniline: ESMS *m/e*: 467.2 $(M + H)^+$.

Example 425

ETHYL 3-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-1H-INDOLE-2-CARBOXYLATE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and ethyl 1*H*-indole-2-carboxylate: ESMS *m/e*: 448.2 $(M + H)^+$.

Example 426

2-METHYL-N-(3-{1-[(1-METHYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1-methyl-1*H*-indole: ESMS *m/e*: 390.2 $(M + H)^+$.

Example 427

N-(3-{1-[(5-METHOXY-2-METHYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 5-methoxy-2-methyl-1*H*-indole: ESMS *m/e*: 420.2 $(M + H)^+$.

Example 428

2-METHYL-N-(3-{1-[(1-METHYL-2-PHENYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1-methyl-2-phenyl-1*H*-indole: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 429

2-METHYL-N-(3-{1-[(5-NITRO-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 5-nitro-1*H*-indole: ESMS *m/e*: 421.1 $(M + H)^+$.

Example 430

2-METHYL-N-(3-{1-[(2-METHYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 2-methyl-1*H*-indole: ESMS *m/e*: 390.2 $(M + H)^+$.

Example 431

N-(3-{1-[(4-BROMO-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 4-bromo-1*H*-indole: ESMS *m/e*: 455.0 $(M + H)^+$.

Example 432

N-[3-(1-{[2-(4-FLUOROPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: **Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 2-(4-fluorophenyl)-1*H*-indole: ESMS *m/e*: 470.0 $(M + H)^+$.**

Example 433

N-(3-{1-[(1,2-DIPHENYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: **Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1,2-diphenyl-1*H*-indole: ESMS *m/e*: 528.2 $(M + H)^+$.**

Example 434

N-[3-(1-{[2-(4-CHLOROPHENYL)-1-ETHYL-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-

[3-(4-piperidinyl)phenyl]propanamide and 2-(4-chlorophenyl)-1-ethyl-1*H*-indole: ESMS *m/e*: 514.1 $(M + H)^+$.

Example 435

N-(3-{1-[(5-CHLORO-2-METHYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 5-chloro-2-methyl-1*H*-indole: ESMS *m/e*: 424.1$(M + H)^+$.

Example 436

N-(3-{1-[(5-CYANO-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1*H*-indole-5-carbonitrile: ESMS *m/e*: 401.1 $(M + H)^+$.

Example 437

2-METHYL-N-(3-{1-[(5-METHYL-2-PHENYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure D and Scheme N using** 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 5-methyl-2-phenyl-1*H*-indole: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 438

2-METHYL-N-[3-(1-{[1-(4-NITROPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1-(4-nitrophenyl)-1*H*-indole: ESMS *m/e*: 497.2 $(M + H)^+$.

Example 439

N-[3-(1-{[1-(2-FLUOROPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1-(2-fluorophenyl)-1*H*-indole: ESMS *m/e*: 470.1 $(M + H)^+$.

Example 440

N-(3-{1-[(5,6-DIMETHOXY-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 5,6-dimethoxy-1*H*-indole: ESMS *m/e*: 436.2 $(M + H)^+$.

Example 441

2-METHYL-N-[3-(1-{[1-(3-METHYLPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1-(3-methylphenyl)-1*H*-indole: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 442

2-METHYL-N-{3-[1-({1-[3-(TRIFLUOROMETHYL)PHENYL]-1H-INDOL-3-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 1-[3-(trifluoromethyl)phenyl]-1*H*-indole: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 443

N-[3-(1-{[1-(4-METHOXYPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-

piperidinyl)phenyl]propanamide and 5-methoxy-2-phenyl-1*H*-indole: ESMS *m/e*: 482.2 $(M + H)^+$.

Example 445

2-METHYL-N-(3-{1-[(5-METHYL-1H-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: **Prepared by Procedure D and Scheme N using 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide and 5-methyl-1*H*-indole: ESMS *m/e*: 390.2 $(M + H)^+$.**

Example 446

***N*-[3-(1-{[1-(2-NITROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(2-nitrophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 497.2 $(M + H)^+$.

Example 447

***N*-[3-(1-{[1-(2-METHOXYPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(2-methoxyphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 482.2 $(M + H)^+$.

Example 448

2-METHYL-*N*-{3-[1-({1-[2-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure D and Scheme N using 1-[2-(trifluoromethyl)phenyl]-1*H*-indole, and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 449

***N*-(3-{1-[(5-METHOXY-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1*H*-indol-5-yl methyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 406.2 $(M + H)^+$.

Example 450

***N*-[3-(1-{[1-(4-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(4-fluorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 470.2 $(M + H)^+$.

Example 451

***N*-[3-(1-{[1-(3-METHOXYPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(3-methoxyphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 482.2 $(M + H)^+$.

Example 452

2-METHYL-*N*-[3-(1-{[1-(2-METHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 1-(2-methylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 453

ETHYL 3-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-5-METHOXY-1*H*-INDOLE-2-CARBOXYLATE: Prepared by Procedure D and Scheme N using ethyl 5-methoxy-1*H*-indole-2-carboxylate and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 478.2 $(M + H)^+$.

Example 454

***N*-(3-{1-[(5-FLUORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 5-fluoro-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 394.2 $(M + H)^+$.

1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and iodobenzene: ESMS *m/e*: 193.9 $(M + H)^+$.

1-(4-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-chloro-4-iodobenzene: ESMS *m/e*: 227.9 $(M + H)^+$.

1-(3-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-chloro-3-iodobenzene: ESMS *m/e*: 227.9 $(M + H)^+$.

1-(2-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-chloro-2-iodobenzene: ESMS *m/e*: 227.9 $(M + H)^+$.

1-[2-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-2-(trifluoromethyl)benzene: ESMS *m/e*: 262.0 $(M + H)^+$.

4-(1*H*-INDOL-1-YL)BENZONITRILE: Prepared by Procedure C and Scheme O using 1*H*-indole and 4-iodobenzonitrile: ESMS *m/e*: 219.0 $(M + H)^+$.

1-(4-NITROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-4-nitrobenzene: ESMS *m/e*: 238.2 $(M + H)^+$.

1-(2-NITROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-2-nitrobenzene: ESMS *m/e*: 238.2 $(M + H)^+$.

Example 455

***N*-[3-(1-{[1-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(4-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 472.1 $(M + H)^+$.

Example 456

***N*-[3-(1-{[1-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(3-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 472.1 $(M + H)^+$.

Example 457

***N*-[3-(1-{[1-(2-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 1-(2-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 484.1 $(M + H)^+$.

Example 458

***N*-[3-(1-{[1-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(3-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 486.1 $(M + H)^+$.

Example 459

***N*-[3-(1-{[1-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(4-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 460

***N*-[3-(1-{[1-(2-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(2-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 461

***N*-[3-(1-{[1-(2-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-(2-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 472.1 $(M + H)^+$.

Example 462

***N*-[3-(1-{[1-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 1-(4-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 484.1 $(M + H)^+$.

Example 463

***N*-[3-(1-{[1-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by

Procedure D and Scheme N using 1-(3-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 484.1 $(M + H)^+$..

Example 464

***N*-(3-{1-[(1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-phenyl-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 438.2 $(M + H)^+$.

Example 465

***N*-(3-{1-[(1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 1-phenyl-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 450.2 $(M + H)^+$.

6-CHLORO-1-(4-NITROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-4-nitrobenzene: ESMS *m/e*: 272.6 $(M + H)^+$.

6-CHLORO-1-(2,3-DICHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1,2-dichloro-3-iodobenzene: ESMS *m/e*: 296.5 $(M + H)^+$.

6-CHLORO-1-(3-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-3-methylbenzene: ESMS *m/e*: 241.9 $(M + H)^+$.

6-CHLORO-1-(2-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-2-methylbenzene: ESMS *m/e*: 241.9 $(M + H)^+$.

2-(6-CHLORO-1*H*-INDOL-1-YL)PHENYL METHYL ETHER: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-2-methoxybenzene: ESMS *m/e*: 257.9 $(M + H)^+$.

6-CHLORO-1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-3-(trifluoromethyl)benzene: ESMS *m/e*: 295.6 $(M + H)^+$.

6-CHLORO-1-(2-FLUOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-fluoro-2-iodobenzene: ESMS *m/e*: 245.9 $(M + H)^+$.

6-CHLORO-1-(3-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-chloro-3-iodobenzene: ESMS *m/e*: 261.9 $(M + H)^+$.

6-CHLORO-1-(4-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-chloro-4-iodobenzene: ESMS *m/e*: 262.9 $(M + H)^+$.

6-CHLORO-1-(2-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-chloro-2-iodobenzene: ESMS *m/e*: 262.9 $(M + H)^+$.

3-(6-CHLORO-1*H*-INDOL-1-YL)PHENYL METHYL ETHER: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-3-methoxybenzene: ESMS *m/e*: 257.9 $(M + H)^+$.

6-CHLORO-1-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-

indole and 1-iodo-4-(trifluoromethyl)benzene ESMS *m/e*: 295.6 $(M + H)^+$.

6-CHLORO-1-(4-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-iodo-4-methylbenzene: ESMS *m/e*: 241.9 $(M + H)^+$.

6-CHLORO-1-(4-FLUOROPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and 1-fluoro-4-iodobenzene: ESMS *m/e*: 245.9 $(M + H)^+$.

Example 466

***N*-[3-(1-{[6-CHLORO-1-(4-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-fluorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 502.1 $(M + H)^+$.

Example 467

***N*-[3-(1-{[6-CHLORO-1-(4-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-fluorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 490.1 $(M + H)^+$.

Example 468

***N*-(3-{1-[(6-FLUORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-fluoro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 380.1 $(M + H)^+$.

Example 469

***N*-(3-{1-[(6-FLUORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 6-fluoro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 392.1 $(M + H)^+$.

Example 470

***N*-(3-{1-[(6-FLUORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-fluoro-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 394.1 $(M + H)^+$.

Example 471

***N*-[3-(1-{[6-CHLORO-1-(4-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-fluorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 504.1 $(M + H)^+$.

Example 472

***N*-[3-(1-{[6-CHLORO-1-(2-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(2-fluorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 490.1 $(M + H)^+$.

Example 473

***N*-[3-(1-{[6-CHLORO-1-(2-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(2-fluorophenyl)-1*H*-indole and *N*-[3-(4-

piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 502.1 $(M + H)^+$.

Example 474

***N*-[3-(1-{[6-CHLORO-1-(2-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(2-fluorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide ESMS *m/e*: 504.1 $(M + H)^+$.

Example 475

***N*-[3-(1-{[6-CHLORO-1-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide ESMS *m/e*: 506.1 $(M + H)^+$.

Example 476

***N*-[3-(1-{[6-CHLORO-1-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide ESMS *m/e*: 518.1 $(M + H)^+$.

Example 477

***N*-[3-(1-{[6-CHLORO-1-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide ESMS *m/e*: 520.1 $(M + H)^+$.

Example 478

***N*-[3-(1-{[6-CHLORO-1-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(3-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 506.1 $(M + H)^+$.

Example 479

***N*-[3-(1-{[6-CHLORO-1-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(3-chlorophenyl)-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.72 (d, 1H, J = 8.4 Hz), 7.68 (s, 1H), 7.49 (m, 2H), 7.44 (d, 2H, J = 7.9 Hz), 7.49-7.25 (m, 4H), 7.21 (d, 1H, J = 7.9 Hz), 7.17 (d, 1H, J = 7.9 Hz), 6.93 (d, 1H, J = 7.9 Hz), 3.79 (s, 2H), 3.13 (d, 2H, J = 9.4 Hz), 2.48 (sept, 1H, J = 7.5 Hz), 2.16 (m, 2H), 1.80 (m, 4H), 1.51 (s, 1H), 1.06 (m, 2H), 0.806 (m, 2H); Anal. Calcd for $C_{30}H_{29}Cl_2N_3O+HCl+1.4H_2O$: C, 62.11; H, 5.70; N, 7.24. Found: C, 62.19; H, 6.21; N, 7.06; ESMS *m/e*: 519.2 $(M + H)^+$.

Example 480

***N*-[3-(1-{[6-CHLORO-1-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(3-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 520.1 $(M + H)^+$.

Example 481

***N*-(3-{1-[(5-FLUORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by

Procedure D and Scheme N using 5-fluoro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 392.1 $(M + H)^+$.

Example 482

_N_-[3-(1-{[6-CHLORO-1-(2-CHLOROPHENYL)-1_H_-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure D and Scheme N using 6-chloro-1-(2-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 483

_N_-[3-(1-{[6-CHLORO-1-(3-METHOXYPHENYL)-1_H_-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure D and Scheme N using 3-(6-chloro-1*H*-indol-1-yl)phenyl methyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 516.2 $(M + H)^+$.

Example 484

_N_-[3-(1-{[6-CHLORO-1-(2-METHOXYPHENYL)-1_H_-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure D and Scheme N using 2-(6-chloro-1*H*-indol-1-yl)phenyl methyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 516.2 $(M + H)^+$.

Example 485

_N_-[3-(1-{[6-CHLORO-1-(2,3-DICHLOROPHENYL)-1_H_-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure D and Scheme N using 6-chloro-1-(2,3-dichlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 555.1 $(M + H)^+$.

Example 486

***N*-[3-(1-{[6-CHLORO-1-(4-METHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(4-methylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 500.2 $(M + H)^+$.

Example 487

***N*-{3-[1-({6-CHLORO-1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-[3-(trifluoromethyl)phenyl]-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 554.2 $(M + H)^+$.

Example 488

***N*-{3-[1-({6-CHLORO-1-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}METHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-[4-(trifluoromethyl)phenyl]-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 554.2 $(M + H)^+$.

Example 489

***N*-[3-(1-{[6-CHLORO-1-(2-METHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-(2-methylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 500.2 $(M + H)^+$.

Example 490

***N*-[3-(1-{[6-CHLORO-1-(3-METHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-

(3-methylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 500.2 $(M + H)^+$.

Example 491

***N*-(3-{1-[(7-CHLORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 7-chloro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 408.1 $(M + H)^+$.

Example 492

***N*-(3-{1-[(7-CHLORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 7-chloro-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 410.1 $(M + H)^+$.

Example 493

***N*-(3-{1-[(4-FLUORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure D and Scheme N using 4-fluoro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 380.2 $(M + H)^+$.

Example 494

***N*-(3-{1-[(7-CHLORO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure D and Scheme N using 7-chloro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 396.1 $(M + H)^+$.

Example 495

2-METHYL-*N*-(3-{1-[(6-METHYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 6-methyl-1*H*-indole and 2-methyl-*N*-[3-

(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 390.2 $(M + H)^+$.

Example 496

***N*-[3-(1-{[6-(BENZYLOXY)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-(benzyloxy)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 482.2 $(M + H)^+$.

Example 497

***N*-(3-{1-[(6-METHOXY-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1*H*-indol-6-yl methyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 406.2 $(M + H)^+$.

Example 498

METHYL 3-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-1*H*-INDOLE-6-CARBOXYLATE: Prepared by Procedure D and Scheme N using methyl 1*H*-indole-6-carboxylate and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 434.2 $(M + H)^+$.

Example 499

2-METHYL-*N*-[3-(1-{[6-(TRIFLUOROMETHYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 6-(trifluoromethyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1H), 7.66 (s, 1H), 7.63 (s, 2H), 7.44 (d, 1H, J = 8.4 Hz), 7.39 (s, 2H), 7.32 (d, 1H, J = 8.4 Hz), 7.16 (t, 1H, J = 8.4 Hz), 6.84 (d, 1H, J = 8.4 Hz), 4.06 (s,

2H), 3.27 (d, 2H, J = 11.6 Hz), 2.56 (sept, 1H, J = 6.8 Hz), 2.37 (m, 3H), 1.93 (m, 2H), 1.75 (m, 2H), 1.22 (d, 6H, J = 6.8 Hz); Anal. Calcd for $C_{25}H_{28}F_3N_3O$+2HCl+0.5EtOAc: C, 57.8; H, 6.11; N, 7.50. Found: C, 56.5; H, 6.46; N, 7.77; ESMS *m/e*: 444.2 $(M + H)^+$.

1-(2-PYRIDINYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 2-iodopyridine and 1*H*-indole: ESMS *m/e*: 195.0 $(M + H)^+$.

1-(3-PYRIDINYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 3-iodopyridine and 1*H*-indole: ESMS *m/e*: 195.0 $(M + H)^+$.

Example 500

2-METHYL-*N*-[3-(1-{[1-(3-PYRIDINYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 1-(3-pyridinyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 501

2-METHYL-*N*-[3-(1-{[1-(2-PYRIDINYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 1-(2-pyridinyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 502

***N*-(3-{1-[(6-FLUORO-1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE**: Prepared by Procedure D and Scheme N using 6-fluoro-1-phenyl-1*H*-

indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 470.2 $(M + H)^+$.

Example 503

***N*-(3-{1-[(6-CHLORO-1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-chloro-1-phenyl-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

7-METHYL-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 7-methyl-1*H*-indole and iodobenzene: ESMS *m/e*: 208.1 $(M + H)^+$.

METHYL 1-PHENYL-1*H*-INDOLE-6-CARBOXYLATE: Prepared by Procedure C and Scheme O using methyl 1*H*-indole-6-carboxylate and iodobenzene: ESMS *m/e*: 252.0 $(M + H)^+$.

6-METHYL-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-methyl-1*H*-indole and iodobenzene: ESMS *m/e*: 208.0 $(M + H)^+$.

7-CHLORO-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 7-chloro-1*H*-indole and iodobenzene: ESMS *m/e*: 228.0 $(M + H)^+$.

6-NITRO-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-nitro-1*H*-indole and iodobenzene: ESMS *m/e*: 238.2 $(M + H)^+$.

6-METHOXY-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indol-6-yl methyl ether and iodobenzene: ESMS *m/e*: 224.0 $(M + H)^+$.

BENZYL 1-PHENYL-1*H*-INDOL-6-YL ETHER: Prepared by Procedure C and Scheme O using 6-(benzyloxy)-1*H*-indole and iodobenzene: ESMS *m/e*: 300.0 $(M + H)^+$.

1-PHENYL-1*H*-INDOL-6-YL TRIFLUOROMETHYL ETHER: Prepared by Procedure C and Scheme O using 6-(trifluoromethoxy)-1*H*-indole and iodobenzene: ESMS *m/e*: 278.0 $(M + H)^+$.

7-METHOXY-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indol-7-yl methyl ether and iodobenzene: ESMS *m/e*: 224.0 $(M + H)^+$.

1-PHENYL-6-(TRIFLUOROMETHYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-(trifluoromethyl)-1*H*-indole and iodobenzene: ESMS *m/e*: 262.0 $(M + H)^+$.

1-(4-PYRIDINYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 4-iodopyridine: ESMS *m/e*: 195 $(M + H)^+$.

Example 504

***N*-[3-(1-{[6-(BENZYLOXY)-1-PHENYL-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using benzyl 1-phenyl-1*H*-indol-6-yl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 558.0 $(M + H)^+$.

Example 505

2-METHYL-*N*-(3-{1-[(6-METHYL-1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 6-methyl-1-phenyl-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$) δ

7.66 (s, 1H), 7.64 (d, 1H, J = 7.8 Hz), 7.51 (d, 1H, J = 3.9 Hz), 7.50 (m, 3H), 7.4 (m, 2H), 7.36-7.32 (m, 2H), 7.31 (s, 1H), 7.19 (t, 1H, J= 7.8 Hz), 7.04 (d, 1H, J = 7.8 Hz), 6.91 (d, 1H, J = 7.8 Hz), 3.94 (s, 2H), 3.25 (d, 2H, J = 9.2 Hz), 2.52 (sept, 1H, J = 6.4 Hz), 2.46 (s, 3H), 2.28 (dt, 2H, J = 11.8, 2.6 Hz), 1.89 (dq, 2H, J = 2.9 Hz), 1.80 (m, 3H), 1.22 (d, 6H, J = 6.9 Hz); Anal. Calcd for $C_{31}H_{35}N_3O$+HCl+0.6EtOAc: C, 72.2; H, 7.41; N, 7.57. Found: C, 71.0; H, 7.40; N, 7.66; ESMS *m/e*: 466 $(M + H)^+$.

Example 506

METHYL 3-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-1-PHENYL-1*H*-INDOLE-6-CARBOXYLATE: Prepared by Procedure D and Scheme N using methyl 1-phenyl-1*H*-indole-6-carboxylate and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 510.0 $(M + H)^+$.

Example 507

2-METHYL-*N*-(3-{1-[(6-NITRO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 6-nitro-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 421.0 $(M + H)^+$.

Example 508

2-METHYL-*N*-[3-(1-{[1-PHENYL-6-(TRIFLUOROMETHYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 1-phenyl-6-(trifluoromethyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 520.0 $(M + H)^+$.

Example 509

2-METHYL-*N*-(*3*-{1-[(7-METHYL-1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 7-methyl-1-phenyl-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 466.0 $(M + H)^+$.

Example 510

***N*-(*3*-{1-[(7-METHOXY-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1*H*-indol-7-yl methyl ether and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 406.0 $(M + H)^+$.

Example 511

***N*-(*3*-{1-[(7-METHOXY-1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 7-methoxy-1-phenyl-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 482.0 $(M + H)^+$.

Example 512

***N*-(*3*-{1-[(7-CHLORO-1-PHENYL-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 7-chloro-1-phenyl-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 488.6 $(M + H)^+$.

Example 513

2-METHYL-*N*-(*3*-{1-[(7-NITRO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure D and Scheme N using 7-nitro-1*H*-indole and 2-methyl-*N*-[3-

(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 421.1 $(M + H)^+$.

Example 514

***N*-(3-{1-[(7-NITRO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure D and Scheme N using 7-nitro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 419.5 $(M + H)^+$.

Example 515

***N*-(3-{1-[(7-NITRO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure D and Scheme N using 7-nitro-1*H*-indole and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 407.3 $(M + H)^+$.

7-(2-FLUOROPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 7-bromo-1*H*-indole and 2-fluorophenylboronic acid: ESMS *m/e*: 211.9 $(M + H)^+$.

Example 516

***N*-[3-(1-{[7-(2-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N. A solution of 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide (23.3 mg, 0.0948 mmol) and 37 wt % aqueous formaldehyde (11.4 mg, 0.142 mmol) in 1.00 mL of HOAc:dioxane (1:4) was added to 7-(2-fluorophenyl)-1*H*-indole (20.0 mg, 0.0948 mmol) and the reaction mixture was stirred for 12 h at room temperature. The resulting mixture was diluted with H_2O (10 mL). The aqueous layer was extracted with CH_2Cl_2 (3 X 10 mL). The combined organic extracts were washed with brine (10 mL), dried over $MgSO_4$, and concentrated *in*

vacuo. The residue was purified by preparative TLC on silica using 4 % of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product (56.1 mg, 100%): ^1H NMR (400 MHz, $CDCl_3$) δ 8.58 (s, 1H), 7.73 (dd, 1H, J = 2.8, 6.3 Hz), 7.69 (s, 1H), 7.53 (dt, 1H, J = 1.8, 7.6 Hz), 7.44 (d, 1H, J = 8.1 Hz), 7.38 (m, 2H), 7.32 (s, 1H), 7.27-7.21 (m, 4H), 7.17 (t, 1H, J = 7.6 Hz), 6.88 (d, 1H, J = 7.6 Hz), 3.92 (s, 2H); 3.20 (d, 1H, J = 11.6 Hz), 2.51 (qt, 1H, J = 6.7 Hz), 2.42 (m, 1H), 2.25 (dt, 2H, J = 2.2, 11.6 Hz), 1.89-1.72 (m, 5H), 1.22 (d, 6H, J = 7.3 Hz); ESMS *m/e*: 470.1 $(M + H)^+$.

7-(4-ETHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 7-bromo-1*H*-indole and 4-ethylphenylboronic acid: ESMS *m/e*: 222.0 $(M + H)^+$.

7-(2-NAPHTHYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 7-bromo-1*H*-indole and 2-naphthylboronic acid: ESMS *m/e*: 244.0 $(M + H)^+$.

7-(3-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 7-bromo-1*H*-indole and 3-chlorophenylboronic acid: ESMS *m/e*: 227.9 $(M + H)^+$.

6-(2-FLUOROPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 6-bromo-1*H*-indole and 2-fluorophenylboronic acid: ESMS *m/e*: 211.9 $(M + H)^+$.

7-(3-NITROPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 7-bromo-1*H*-indole and 3-nitrophenylboronic acid: ESMS *m/e*: 238.9 $(M + H)^+$.

1-[4-(1*H*-INDOL-7-YL) PHENYL] ETHANONE: Prepared by Procedure I and Scheme T using 7-bromo-1*H*-indole and 4-acetylphenylboronic acid: ESMS *m/e*: 235.2 $(M + H)^+$.

6-(2-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 6-bromo-1*H*-indole and 2-methylphenylboronic acid: ESMS *m/e*: 207.9 $(M + H)^+$.

6-(3-CHLOROPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 6-bromo-1*H*-indole and 3-chlorophenylboronic acid: ESMS *m/e*: 227.9 $(M + H)^+$.

1-[4-(1*H*-INDOL-6-YL) PHENYL] ETHANONE: Prepared by Procedure I and Scheme T using 6-bromo-1*H*-indole and 4-acetylphenylboronic acid: ESMS *m/e*: 235.8 $(M + H)^+$.

7-(2-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 7-bromo-1H-indole and 2-methylphenylboronic acid: ESMS *m/e*: 208 $(M + H)^+$.

6-(4-ETHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 6-bromo-1*H*-indole and 4-ethylphenylboronic acid: ESMS *m/e*: 221.9 $(M + H)^+$.

Example 517

2-METHYL-*N*-[3-(1-{[7-(2-NAPHTHYL)-1*H*-INDOL-3-YL] METHYL}-4-PIPERIDINYL) PHENYL] PROPANAMIDE: Prepared by Procedure D and Scheme N using 7-(2-naphthyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 502.2 $(M + H)^+$.

Example 518

N-[3-(1-{[7-(4-ETHYLPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 7-(4-ethylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 480.2 $(M + H)^+$.

Example 519

2-METHYL-N-[3-(1-{[6-(2-METHYLPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 6-(2-methylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 8.2 (s, 1H), 7.53 (m, 4H), 7.41 (d, 1H, J = 8.4 Hz), 7.34 (m, 2H), 7.27-7.12 (m, 5H), 6.81 (d, 1H, J = 8.4 Hz), 4.09 (s, 2H), 3.32 (d, 2H, J = 11.4 Hz), 2.57 (q, 2H, J = 7.6 Hz), 2.43 (m, 3H), 2.08 (s, 3H), 1.98 (m, 1H), 1.75 (m, 2H), 1.22 (d, 6H, J = 6.3 Hz); Anal. Calcd for $C_{31}H_{35}N_3O+CHCl_3+DMF$: C, 57.0; H, 6.09; N, 8.06. Found: C, 56.5; H, 5.94; N, 7.76; ESMS *m/e*: 466.2 $(M + H)^+$.

Example 520

N-[3-(1-{[7-(3-CHLOROPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure D and Scheme N using 7-(3-chlorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 486.1 $(M + H)^+$.

Example 521

2-METHYL-N-[3-(1-{[7-(3-NITROPHENYL)-1H-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 7-(3-nitrophenyl)-1*H*-

indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 497.0 $(M + H)^+$.

Example 522

***N*-[3-(1-{[7-(4-ACETYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 1-[4-(1*H*-indol-7-yl)phenyl]ethanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 493.6 $(M + H)^+$.

Example 523

***N*-[3-(1-{[6-(4-ETHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-(4-ethylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 480.1 $(M + H)^+$.

Example 524

2-METHYL-*N*-[3-(1-{[7-(2-METHYLPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 7-(2-methylphenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 466.1 $(M + H)^+$.

Example 525

***N*-[3-(1-{[6-(2-FLUOROPHENYL)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 6-(2-fluorophenyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 470.2 $(M + H)^+$.

5-(4-METHYLPHENOXY)-1*H*-INDOLE: Prepared by Procedure J and Scheme U using 5-bromo-1*H*-indole and *p*-cresol: ESMS *m/e*: 224.0 $(M + H)^+$.

Example 526

***N*-(3-{1-[(5-BROMO-1*H*-INDOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure D and Scheme N using 5-bromo-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 454.0 $(M + H)^+$.

1-(4-PYRIDINYL)-6-(TRIFLUOROMETHYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-(trifluoromethyl)-1*H*-indole and 4-iodopyridine: ESMS *m/e*: 262.9 $(M + H)^+$.

Example 527

2-METHYL-*N*-[3-(1-{[5-(4-METHYLPHENOXY)-1*H*-INDOL-3-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure D and Scheme N using 5-(4-methylphenoxy)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 481.9 $(M + H)^+$.

1-(4-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-4-methylbenzene: ESMS *m/e*: 208.0 $(M + H)^+$.

1-(3-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-3-methylbenzene: ESMS *m/e*: 208.0 $(M + H)^+$.

1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-3-(trifluoromethyl)benzene: ESMS *m/e*: 262.0 $(M + H)^+$.

1-(4-METHOXYPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-4-methoxybenzene: ESMS *m/e*: 224.0 $(M + H)^+$.

1-(2-METHOXYPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-2-methoxybenzene: ESMS *m/e*: 224.0 $(M + H)^+$.

1-(3-METHOXYPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-3-methoxybenzene: ESMS *m/e*: 224.0 $(M + H)^+$.

1-(2-METHYLPHENYL)-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 1*H*-indole and 1-iodo-2-methylbenzene: ESMS *m/e*: 208.0 $(M + H)^+$.

6-FLUORO-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-fluoro-1*H*-indole and iodobenzene: ESMS *m/e*: 212.0 $(M + H)^+$.

6-CHLORO-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 6-chloro-1*H*-indole and iodobenzene: ESMS *m/e*: 228.0 $(M + H)^+$.

7-CHLORO-1-PHENYL-1*H*-INDOLE: Prepared by Procedure C and Scheme O using 7-chloro-1*H*-indole and iodobenzene: ESMS *m/e*: 228.0 $(M + H)^+$.

6-(2-FLUOROPHENYL)-1*H*-INDOLE: Prepared by Procedure I and Scheme T using 6-bromo-1*H*-indole and 2-fluorophenylboronic acid: ESMS *m/e*: 211.9 $(M + H)^+$.

Example 528

2-METHYL-*N*-{3-[1-(7-OXO-7-PHENYLHEPTYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure K and Scheme B1 using 7-chloro-1-phenyl-1-heptanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 435.1 $(M + H)^+$.

Example 529

2-METHYL-*N*-{3-[1-(6-OXO-6-PHENYLHEXYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure K and Scheme B1 using 6-chloro-1-phenyl-1-hexanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: Anal. Calcd for $C_{27}H_{36}N_2O_2+0.1CHCl_3$: C, 75.3; H, 8.39; N, 6.46. Found: C, 75.4; H, 7.89; N, 6.18; ESMS *m/e*: 421.1 $(M + H)^+$.

Example 530

2-METHYL-*N*-{3-[1-(5-OXO-5-PHENYLPENTYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure K and Scheme B1 using 5-chloro-1-phenyl-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 407.1 $(M + H)^+$.

Example 531

***N*-(3-{1-[4-(4-METHOXYPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-methoxyphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 409.2 $(M + H)^+$.

Example 532

***N*-(3-{1-[4-(4-CHLOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-chlorophenyl)-1-

butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 413.1 $(M + H)^+$.

Example 533

***N*-(3-{1-[4-(4-BROMOPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 1-(4-bromophenyl)-4-chloro-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 457.1 $(M + H)^+$.

Example 534

***N*-(3-{1-[4-(4-*TERT*-BUTYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 1-(4-*tert*-butylphenyl)-4-chloro-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 435.2 $(M + H)^+$.

Example 535

***N*-(3-{1-[4-(4-FLUOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-fluorophenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 397.2 $(M + H)^+$.

Example 536

***N*-(3-{1-[4-OXO-4-(4-PHENOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-phenoxyphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 471.2 $(M + H)^+$.

Example 537

***N*-(3-{1-[4-(4-ISOPROPYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-isopropylphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 433.2 $(M + H)^+$.

Example 538

***N*-(3-{1-[4-(4-METHOXYPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-methoxyphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 421.2 $(M + H)^+$.

Example 539

***N*-(3-{1-[4-OXO-4-(4-PHENOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-phenoxyphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 483.2 $(M + H)^+$.

Example 540

***N*-(3-{1-[4-(4-ISOPROPYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-isopropylphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 421.3 $(M + H)^+$.

Example 541

***N*-(3-{1-[4-(4-*TERT*-BUTYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 1-(4-*tert*-butylphenyl)-4-chloro-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 447.2 $(M + H)^+$.

Example 542

***N*-(3-{1-[4-(4-METHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-methylphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 393.2 $(M + H)^+$.

Example 543

***N*-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(3,4-dimethylphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 407.2 $(M + H)^+$.

Example 544

***N*-(3-{1-[4-(4-BROMOPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 1-(4-bromophenyl)-4-chloro-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 469.1 $(M + H)^+$.

Example 545

***N*-(3-{1-[5-(4-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(4-fluorophenyl)-1-

pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 411.2 $(M + H)^+$.

Example 546

***N*-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(3,4-dimethylphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 419.2 $(M + H)^+$.

Example 547

***N*-(3-{1-[4-(4-METHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-methylphenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 405.2 $(M + H)^+$.

Example 548

***N*-(3-{1-[4-(4-FLUOROPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 4-chloro-1-(4-fluorophenyl)-1-butanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 409.2 $(M + H)^+$.

Example 549

***N*-(3-{1-[5-(3-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(3-fluorophenyl)-1-pentanone and *N*-[3-(4-

piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 423.2 $(M + H)^+$.

Example 550

***N*-[3-(1-{5-OXO-5-[4-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-[4-(trifluoromethyl)phenyl]-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 461.2 $(M + H)^+$.

Example 551

***N*-(3-{1-[5-(4-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(4-fluorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 423.2 $(M + H)^+$.

Example 552

***N*-(3-{1-[5-(3-NITROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(3-nitrophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 438.2 $(M + H)^+$.

Example 553

***N*-(3-{1-[5-(3-NITROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(3-nitrophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 450.2 $(M + H)^+$.

Example 554

***N*-(3-{1-[5-(2-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(2-fluorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 411.2 $(M + H)^+$.

Example 555

***N*-(3-{1-[5-(3-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(3-fluorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 411.2 $(M + H)^+$.

Example 556

***N*-(3-{1-[5-(4-NITROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(4-nitrophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 438.1 $(M + H)^+$.

Example 557

***N*-(3-{1-[5-(4-NITROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(4-nitrophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 450.1 $(M + H)^+$.

Example 558

***N*-(3-{1-[5-(4-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(4-

385

chlorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 439.1 $(M + H)^+$.

Example 559

***N*-[3-(1-{5-OXO-5-[2-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-[2-(trifluoromethyl)phenyl]-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 461.2 $(M + H)^+$.

Example 560

***N*-[3-(1-{5-OXO-5-[2-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-[2-(trifluoromethyl)phenyl]-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 473.2 $(M + H)^+$.

Example 561

***N*-(3-{1-[5-(4-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(4-chlorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 427.1 $(M + H)^+$.

Example 562

***N*-(3-{1-[5-(3-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(3-chlorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 427.1 $(M + H)^+$.

Example 563

***N*-(3-{1-[5-(2-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(2-fluorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 423.1 $(M + H)^+$.

Example 564

***N*-(3-{1-[5-(3-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(3-chlorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 439.1 $(M + H)^+$.

Example 565

***N*-[3-(1-{5-OXO-5-[4-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-[4-(trifluoromethyl)phenyl]-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 473.2 $(M + H)^+$.

Example 566

***N*-(3-{1-[5-(2-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(2-chlorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 427.1 $(M + H)^+$.

Example 567

***N*-(3-{1-[5-(2-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-(2-chlorophenyl)-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 439.1 $(M + H)^+$.

Example 568

***N*-[3-(1-{5-OXO-5-[3-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-[3-(trifluoromethyl)phenyl]-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 473.2 $(M + H)^+$.

Example 569

***N*-(3-{1-[4-(3,4-DIMETHYLPHENYL)-4-OXOBUTYL]-4-PIPERIDINYL}PHENYL)-*N*,2-DIMETHYLPROPANAMIDE:** Prepared by Procedure T and Scheme AD using *N*-(3-{1-[4-(3,4-dimethylphenyl)-4-oxobutyl]-4-piperidinyl}phenyl)-2-methylpropanamide and methyl iodide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.76 (s, 1H), 7.72 (dd, 1H, J = 1.8, 7.7 Hz), 7.33 (t, 1H, J = 8.8 Hz), 7.22 (d, 1H, J = 7.8 Hz), 7.18 (d, 1H, J = 8.8 Hz), 7.01 (m, 2H), 3.24 (s, 3H), 3.10 (d, 1H, J = 10.6 Hz), 3.00 (t, 1H, J = 7.6 Hz), 2.49 (m, 4H), 2.33 (s, 6H), 2.11 (m, 3H), 1.99 (m, 1H), 1.79 (m, 4H), 1.26 (t, 2H, J = 7.6 Hz), 1.02 (d, 6H, J = 7.6 Hz); ESMS *m/e*: 435.2 $(M + H)^+$.

Example 570

2-METHYL-*N*-{3-[1-(1-METHYL-4-OXO-4-PHENYLBUTYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure K

388

and Scheme B1 using 4-chloro-1-phenyl-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 407.2 $(M + H)^+$.

Example 571

***N*-[3-(1-{5-OXO-5-[3-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:** Prepared by Procedure K and Scheme B1 using 5-chloro-1-[3-(trifluoromethyl)phenyl]-1-pentanone and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 461.2 $(M + H)^+$.

3-(5-CHLOROPENTANOYL)-4-(3,4-DIFLUOROPHENYL)-1,3-OXAZOLIDIN-2-ONE: Prepared by Procedure AF and Scheme H using 4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one and 5-chloropentanoyl chloride.

3-(5-CHLOROPENTYL)-4-(3,4-DIFLUOROPHENYL)-1,3-OXAZOLIDIN-2-ONE: Prepared by Procedure G and Scheme C1 using 4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one and 1-bromo-5-chloropentane.

Example 572

***N*-[3-(1-{5-[(4*R*)-4-(3,4-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDIN-3-YL]-5-OXOPENTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using (4*R*)-3-(5-chloropentanoyl)-4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 573

(4*R*)-4-(3,4-DIFLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-OXO-

1,3-OXAZOLIDINE-3-CARBOXAMIDE: Prepared by Procedure AF and Scheme H using 4-nitrophenyl (4*R*)-4-(3,4-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxylate and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.08 (t, 1H, J = 5.5 Hz), 7.45 (S, 2H), 7.38 (d, 1H, J = 8.6 Hz), 7.24-7.12 (m, 3H), 7.06 (m, 1H), 6.97 (d, 1H, J = 8.6 Hz), 5.40 (dd, 1H, J = 3.9,8.8 Hz), 4.71 (t, 1H, J = 8.8 Hz), 4.23 (dd, 1H, J = 4.4, 9.1 Hz), 3.32 (qt, 2H, J = 6.1 Hz), 2.99 (d, 2H, J = 11.0 Hz), 2.49 (qt, 2H, J = 7.0 Hz), 2.41 (t, 2H, J = 7.0 Hz), 1.99 (m, 2H), 1.82-1.68 (m, 6H), 1.23 (d, 6H, J = 7.3 Hz); ESMS *m/e*: 529.1 $(M + H)^+$.

(4*S*)-3-(5-CHLOROPENTYL)-4-(3,4-DIFLUOROPHENYL)-1,3-OXAZOLIDIN-2-ONE: Prepared by Procedure G and Scheme C1 using (4*S*)-4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one and 1-bromo-5-chloropentane.

Example 574

***N*-[3-(1-{5-[(4*S*)-4-(3,4-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDIN-3-YL]PENTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using (4*S*)-3-(5-chloropentyl)-4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48 (s, 1H), 7.32 (d, 1H, J = 8.6 Hz), 7.26-7.21 (m, 2H), 7.20-7.12 (m, 2H), 7.06 (m, 1H), 6.97 (d, 1H, J = 6.96 Hz), 4.76 (dd, 1H, J = 6.3, 8.3 Hz), 4.62 (t, 1H, J = 9.0 Hz), 4.06 (dd, 1H, J = 6.4, 8.7 Hz), 3.46 (m, 1H), 3.0 (d, 2H, J = 9.0 Hz), 2.77 (q, 1H, J = 6.8 Hz), 2.50 (q, 2H, J = 6.8 Hz), 2.31 (t, 2H, J = 6.8 Hz), 2.01 (m, 4H), 1.81 (m, 4H), 1.48 (m, 4H), 1.26 (d, 6H, J = 7.3

Hz); Anal. Calcd for $C_{28}H_{37}F_2N_3O_3$+HCl+0.25$CHCl_3$: C, 60.6; H, 6.65; N, 7.25. Found: C, 60.7; H, 6.91; N, 7.05; ESMS *m/e*: 514.2 $(M + H)^+$.

Example 575

***N*-[3-(1-{5-[(4*S*)-4-(3,4-DIFLUOROPHENYL)-2-OXO-1,3-OXAZOLIDIN-3-YL]-5-OXOPENTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using (4*S*)-3-(5-chloropentanoyl)-4-(3,4-difluorophenyl)-1,3-oxazolidin-2-one and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 528.1 $(M + H)^+$.

Example 576

(4*S*)-4-(3,4-DIFLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-OXO-1,3-OXAZOLIDINE-3-CARBOXAMIDE: Prepared by Procedure AF and Scheme H using 4-nitrophenyl (4*S*)-4-(3,4-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxylate and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 529.1 $(M + H)^+$.

Example 577

(4*S*)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-OXO-4-(3,4,5-TRIFLUOROPHENYL)-1,3-OXAZOLIDINE-3-CARBOXAMIDE: Prepared by Procedure AF and Scheme H using 4-nitrophenyl (4*S*)-4-(3,4-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxylate and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 547.1 $(M + H)^+$.

Example 578

(4*S*)-4-(3,5-DIFLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-OXO-

391

1,3-OXAZOLIDINE-3-CARBOXAMIDE: Prepared by Procedure AF and Scheme H using 4-nitrophenyl (4*S*)-4-(3,4-difluorophenyl)-2-oxo-1,3-oxazolidine-3-carboxylate and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 529.2 $(M + H)^+$.

Example 579

***N*-(3-{1-[3-(PHENYLSULFANYL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure G and Scheme B1 using [(3-chloropropyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 382.9 $(M + H)^+$.

Example 580

***N*-(3-{1-[3-(PHENYLSULFANYL)PROPYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure G and Scheme B1 using [(3-chloropropyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 395.1 $(M + H)^+$.

Example 581

2-METHYL-*N*-(3-{1-[3-(PHENYLSULFANYL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure G and Scheme B1 using [(3-chloropropyl)sulfanyl]benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.63 (s, 1H), 7.48 (s, 1H), 7.33 (m, 3H), 7.27 (t, 2H, J = 7.5 Hz), 7.20 (t, 1H, J = 7.9 Hz), 7.15 (tt, 1H, J = 7.2, 1.4 Hz), 6.95 (d, 1H, J = 7.6 Hz), 2.97 (t, 4H, J = 7.3 Hz), 2.46 (m, 4H), 1.99 (dt, 2H, J = 11.4, 3.0 Hz), 1.84 (qt, 2H, J = 7.3 Hz), 1.77 (m, 4H), 1.21 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 396.8 $(M + H)^+$.

Example 582

***N*-(3-{1-[6-(PHENYLSULFANYL)HEXYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure G and Scheme B1 using [(6-chlorohexyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 437.4 $(M + H)^+$.

Example 583

***N*-(3-{1-[4-(PHENYLSULFANYL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure G and Scheme B1 using [(4-chlorobutyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 396.8 $(M + H)^+$.

Example 584

***N*-(3-{1-[4-(PHENYLSULFANYL)BUTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure G and Scheme B1 using [(4-chlorobutyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 409.5 $(M + H)^+$.

Example 585

2-METHYL-*N*-(3-{1-[4-(PHENYLSULFANYL)BUTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure G and Scheme B1 using [(4-chlorobutyl)sulfanyl]benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 410.6 $(M + H)^+$.

Example 586

2-METHYL-*N*-(3-{1-[5-(PHENYLSULFANYL)PENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure G

and Scheme B1 using [(5-chloropentyl)sulfanyl]benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.1 $(M + H)^+$.

Example 587

***N*-(3-{1-[5-(PHENYLSULFANYL)PENTYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure G and Scheme B1 using [(5-chloropentyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 423.1 $(M + H)^+$.

[(6-CHLOROHEXYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using benzenethiol and 1-bromo-6-chlorohexane.

[(4-CHLOROBUTYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using benzenethiol and 1-bromo-4-chlorobutane.

Example 588

***N*-(3-{1-[6-(PHENYLSULFANYL)HEXYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure G and Scheme B1 using [(6-chlorohexyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.4 $(M + H)^+$.

[(5-CHLOROPENTYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using benzenethiol and 1-bromo-5-chloropentane.

[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using benzenethiol and 1-bromo-

3-chloropropane: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.37-7.34 (m, 2H), 7.32-7.26 (m, 2H), 7.19 (tt, 1H, J = 1.4, 7.3 Hz), 3.67 (t, 2H, J = 6.6 Hz), 3.08 (t, 2H, J = 6.6 Hz), 2.06 (qt, 2H, J = 6.6 Hz).

Example 589

***N*-(3-{1-[5-(PHENYLSULFANYL)PENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure G and Scheme B1 using [(5-chloropentyl)sulfanyl]benzene and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 411.1 $(M + H)^+$.

3-CHLOROPROPYL 4-FLUOROPHENYL SULFIDE: Prepared by Procedure R and Scheme Z using 4-fluorobenzenethiol and 1-bromo-3-chloropropane.

1-BROMO-2-[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using 2-bromobenzenethiol and 1-bromo-3-chloropropane.

3-CHLOROPROPYL 4-FLUOROPHENYL SULFOXIDE: Prepared by Procedure S and Scheme AA using 3-chloropropyl 4-fluorophenyl sulfide and 1 eq *m*-CPBA: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.65-7.62 (m, 2H), 7.28-7.21 (m, 2H), 3.65 (m, 2H), 2.94 (m, 2H), 2.28 (m, 1H), 2.06 (m, 1H); ESMS *m/e*: 220.9 $(M + H)^+$.

3-CHLOROPROPYL 3-FLUOROPHENYL SULFIDE: Prepared by Procedure R and Scheme Z using 3-fluorobenzenethiol and 1-bromo-3-chloropropane.

3-CHLOROPROPYL 2-FLUOROPHENYL SULFIDE: Prepared by Procedure R and Scheme Z using 2-fluorobenzenethiol and 1-bromo-3-chloropropane.

1-BROMO-2-[(3-CHLOROPROPYL)SULFINYL]BENZENE: Prepared by Procedure S and Scheme AA using 1-bromo-2-[(3-chloropropyl)sulfanyl]benzene and 1 eq *m*-CPBA: ESMS *m/e*: 282.8 $(M + H)^+$.

1-CHLORO-2-[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using 2-chlorobenzenethiol and 1-bromo-3-chloropropane.

1-CHLORO-3-[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using 3-chlorobenzenethiol and 1-bromo-3-chloropropane.

1-CHLORO-4-[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using 4-chlorobenzenethiol and 1-bromo-3-chloropropane.

1-BROMO-3-[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using 3-bromobenzenethiol and 1-bromo-3-chloropropane.

1-BROMO-4-[(3-CHLOROPROPYL)SULFANYL]BENZENE: Prepared by Procedure R and Scheme Z using 4-bromobenzenethiol and 1-bromo-3-chloropropane.

3-CHLOROPROPYL 3,4-DIMETHYLPHENYL SULFIDE: Prepared by Procedure R and Scheme Z using 3,4-dimethylbenzenethiol and 1-bromo-3-chloropropane.

Example 590

***N*-[3-(1-{3-[(4-FLUOROPHENYL)SULFINYL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 3-chloropropyl 4-

fluorophenyl sulfoxide and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.64 (m, 2H), 7.53 (s, 1H), 7.24 (m, 5H), 6.94 (d, 1H, J = 7.7 Hz), 2.89 (m, 4H), 2.45 (m, 4H), 1.99 (m, 3H), 1.77 (m, 5H), 1.24 (d, 6H, J = 6.8 Hz); Anal. Calcd for $C_{24}H_{31}FN_2O_2S$+0.6EtOAc: C, 65.5; H, 7.45; N, 5.79. Found: C, 65.4; H, 7.30; N, 5.73; ESMS *m/e*: 431.1 (M + H)$^+$.

Example 591

***N*-[3-(1-{3-[(2-BROMOPHENYL)SULFINYL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure G and Scheme B1 using 1-bromo-2-[(3-chloropropyl)sulfinyl]benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: Anal. Calcd for $C_{24}H_{31}BrN_2O_2S$+0.3$CHCl_3$: ESMS *m/e*: 491.0 (M + H)$^+$.

Example 592

***N*-{3-[1-((3*S*)-3-{[(3,4-DIFLUOROPHENYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using 3,4-difluorobenzenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 556.2 (M + H)$^+$.

Example 593

3-CHLORO-*N*-((1*S*)-3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}-1-PHENYLPROPYL)-2-THIOPHENECARBOXAMIDE: Prepared by Procedure Q1 and Scheme AC using 3-chloro-2-thiophenecarbonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 524.2 (M + H)$^+$.

Example 594

***N*-(3-{1-[(3*S*)-3-({[5-(DIMETHYLAMINO)-1-NAPHTHYL]SULFONYL}AMINO)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using 5-(dimethylamino)-1-naphthalenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 613.3 $(M + H)^+$.

Example 595

2-METHYL-*N*-{3-[1-((3*S*)-3-{[(4-METHYLPHENYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure Q1 and Scheme AC using 4-methylbenzenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 534.2 $(M + H)^+$.

Example 596

***N*-{3-[1-((3*S*)-3-{[(3,5-DICHLORO-2-HYDROXYPHENYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared Procedure Q1 and Scheme AC using 3,5-dichloro-2-hydroxybenzenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 605.4 $(M + H)^+$.

Example 597

2-METHYL-*N*-[3-(1-{(3*S*)-3-[(METHYLSULFONYL)AMINO]-3-PHENYLPROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure Q1 and Scheme AC using methanesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 458.6 $(M + H)^+$.

Example 598

***N*-{3-[1-((3*S*)-3-{[(4-FLUOROPHENYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using 4-fluorobenzenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 538.1 $(M + H)^+$.

Example 599

***N*-{3-[1-((3*S*)-3-{[(4-*TERT*-BUTYLPHENYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using 4-*tert*-butylbenzenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 576.2 $(M + H)^+$.

Example 600

***N*-{3-[1-((3*S*)-3-{[(2,5-DICHLOROPHENYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using 2,5-dichlorobenzenesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 588.0 $(M + H)^+$.

Example 601

2-METHYL-*N*-[3-(1-{(3*S*)-3-PHENYL-3-[(PROPYLSULFONYL)AMINO]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure Q1 and Scheme AC using 1-propanesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 602

***N*-{3-[1-((3*S*)-3-{[(3,5-DIMETHYL-4-ISOXAZOLYL)SULFONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 and Scheme AC using 3,5-dimethyl-4-isoxazolesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.53 (s, 2H), 7.3-7.1 (m, 5H), 7.05 (t, 2H, J = 6.5 Hz), 6.81 (d, 1H, J = 7.1 Hz), 4.65 (dd, 1H, J = 6.3, 2.2 Hz), 3.11 (t, 2H, J = 7.2 Hz), 2.4 (m, 4H), 2.2 (s, 3H), 2.05 (m, 2H), 2.01 (s, 3H), 2.0-1.8 (m, 7H), 1.21 (d, 6H, J = 7.1 Hz); ESMS *m/e*: 539.5 $(M + H)^+$.

Example 603

METHYL 3-{[(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)AMINO]SULFONYL}-2-THIOPHENECARBOXYLATE: Prepared Procedure Q1 and Scheme AC using methyl 3-(chlorosulfonyl)-2-thiophenecarboxylate and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: Anal. Calcd for $C_{24}H_{33}N_3O_5S.HCl$: C, 6.00; H, 5.30; N, 7.72. Found: C, 52.9; H, 6.04; N, 7.59; ESMS *m/e*: 508.2 $(M + H)^+$.

Example 604

2-METHYL-*N*-{3-[1-((3*S*)-3-{[(4-PHENOXYANILINO)CARBONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isocyanato-4-phenoxybenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 591.3 $(M + H)^+$.

PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure Q1 and Scheme AC using 3,5-dimethyl-4-isoxazolesulfonyl chloride and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.53 (s, 2H), 7.3-7.1 (m, 5H), 7.05 (t, 2H, J = 6.5 Hz), 6.81 (d, 1H, J = 7.1 Hz), 4.65 (dd, 1H, J = 6.3, 2.2 Hz), 3.11 (t, 2H, J = 7.2 Hz), 2.4 (m, 4H), 2.2 (s, 3H), 2.05 (m, 2H), 2.01 (s, 3H), 2.0-1.8 (m, 7H), 1.21 (d, 6H, J = 7.1 Hz); ESMS *m/e*: 539.5 $(M + H)^+$.

Example 603

METHYL 3-{[(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)AMINO]SULFONYL}-2-THIOPHENECARBOXYLATE: Prepared Procedure Q1 and Scheme AC using methyl 3-(chlorosulfonyl)-2-thiophenecarboxylate and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: Anal. Calcd for $C_{24}H_{33}N_3O_5S.HCl$: C, 6.00; H, 5.30; N, 7.72. Found: C, 52.9; H, 6.04; N, 7.59; ESMS *m/e*: 508.2 $(M + H)^+$.

Example 604

2-METHYL-*N*-{3-[1-((3*S*)-3-{[(4-PHENOXYANILINO)CARBONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isocyanato-4-phenoxybenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 591.3 $(M + H)^+$.

Example 605

***N*-[3-(1-{(3*S*)-3-[(ANILINOCARBONYL)AMINO]-3-PHENYLPROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure P and Scheme AB using isocyanatobenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 499.2 $(M + H)^+$.

Example 606

N-{3-[1-((3*S*)-3-{[(*TERT*-BUTYLAMINO)CARBOTHIOYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure P and Scheme AB using 2-isothiocyanato-2-methylpropane and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 495.1 $(M + H)^+$.

Example 607

N-{3-[1-((3*S*)-3-{[(2-FLUOROANILINO)CARBONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-fluoro-2-isocyanatobenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 517.0 $(M + H)^+$.

Example 608

2-METHYL-N-[3-(1-{(3*S*)-3-PHENYL-3-[(2-TOLUIDINOCARBOTHIOYL)AMINO]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isothiocyanato-2-methylbenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 529.1 $(M + H)^+$.

Example 609

N-{3-[1-((3*S*)-3-{[(BENZYLAMINO)CARBONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:

phenylpropyl]-4- piperidinyl}phenyl)-2-methylpropanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.44 (s, 1H), 7.67 (d, 1H, J = 7.9 Hz), 7.31-7.13 (m, 13H), 6.38 (s, 1H), 6.80 (d, 1H, J = 7.9 Hz), 5.54 (m, 1H), 4.81 (m, 1H), 4.41 (dd, 1H, J = 14.8, 6.2 Hz), 4.29 (dd, 1H, J = 14.9, 5.4 Hz), 2.99 (d, 1H, J = 11.2 Hz), 2.87 (d, 1H, J = 11.2 Hz), 2.67 (q, 1H, J = 6.2 Hz), 2.3 (m, 3H), 2.0-1.5 (m, 7H), 1.23 (d, 6H, J = 6.7 Hz); ESMS *m/e*: 513.2 $(M + H)^+$.

Example 610

2-METHYL-*N*-{3-[1-((3*S*)-3-{[(2-NITROANILINO)CARBONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isocyanato-2-nitrobenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 543.6 $(M + H)^+$.

Example 611

***N*-{3-[1-((3*S*)-3-{[(3,4-DICHLOROANILINO)CARBONYL]AMINO}-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using 1,2-dichloro-4-isocyanatobenzene and *N*-(3-{1-[(3*S*)-3-amino-3-phenylpropyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 567.1 $(M + H)^+$.

Example 612

2-METHYL-*N*-(3-{1-[(3*S*)-3-({[2-(METHYLSULFANYL)ANILINO]CARBONYL}AMINO)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isocyanato-2-(methylsulfanyl)benzene and *N*-(3-{1-[(3*S*)-3-amino-3-

phenylpropyl]-4- piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 545.0 $(M + H)^+$.

Example 613

***N*-{3-[1-(3-{[(4-FLUOROANILINO)CARBONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using 1-fluoro-4-isocyanatobenzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.45 (q, 2H, J = 4.7 Hz), 7.23 (m, 4H), 7.05 (t, 4H, J = 7.8 Hz), 6.75 (m, 1H), 4.05 (m, 1H), 3.19 (s, 1H), 2.71 (m, 1H), 2.53 (m, 1H), 2.25 (m, 3H), 1.8 (m, 9H), 1.25 (d, 6H, J = 6.4 Hz); ESMS *m/e*: 441.1 $(M + H)^+$.

Example 614

***N*-{3-[1-(3-{[(3,4-DICHLOROANILINO)CARBONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using 1,2-dichloro-4-isocyanatobenzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 493.2 $(M + H)^+$.

Example 615

2-METHYL-*N*-[3-(1-{3-[(2-TOLUIDINOCARBOTHIOYL)AMINO]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isothiocyanato-2-methylbenzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 616

***N*-{3-[1-(3-{[(BENZYLAMINO)CARBONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using (isocyanatomethyl)benzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 437.2 $(M + H)^+$.

Example 617

***N*-{3-[1-(3-{[(4-ETHOXYANILINO)CARBONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using 1-ethoxy-4-isocyanatobenzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 467.2 $(M + H)^+$.

Example 618

***N*-[3-(1-{3-[(ANILINOCARBONYL)AMINO]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using isocyanatobenzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 422.9 $(M + H)^+$.

Example 619

2-METHYL-*N*-(3-{1-[3-({[2-(METHYLSULFANYL)ANILINO]CARBONYL}AMINO)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isocyanato-2-(methylsulfanyl)benzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 469.1 $(M + H)^+$.

Example 620

***N*-{3-[1-(3-{[(*TERT*-BUTYLAMINO)CARBOTHIOYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure P and Scheme AB using 2-isothiocyanato-2-methylpropane and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 419.0 $(M + H)^+$.

Example 621

2-METHYL-*N*-{3-[1-(3-{[(4-PHENOXYANILINO)CARBONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure P and Scheme AB using 1-isocyanato-4-phenoxybenzene and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 515.5 $(M + H)^+$.

Example 622

***N*-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-4-(2,4-DIFLUOROPHENYL)-2-METHYL-6-OXO-1,4,5,6-TETRAHYDRO-3-PYRIDINECARBOXAMIDE:** Prepared by Procedure AC and Scheme AM using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide and 4-(2,4-difluorophenyl)-2-methyl-6-oxo-1,4,5,6-tetrahydro-3-pyridinecarboxylic acid: ESMS *m/e*: 525.2 $(M + H)^+$.

Example 623

***N*-(3-{4-[3-(ACETYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-4-(3,4-DIFLUOROPHENYL)-2-METHYL-6-OXO-1,4,5,6-TETRAHYDRO-3-PYRIDINECARBOXAMIDE:** Prepared by Procedure AC and Scheme AM using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}acetamide and 4-(3,4-difluorophenyl)-2-methyl-6-oxo-1,4,5,6-tetrahydro-3-pyridinecarboxylic acid: ESMS *m/e*: 525.2 $(M + H)^+$.

Example 624

***N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-1-(4-NITROPHENYL)-5-(TRIFLUOROMETHYL)-1*H*-PYRAZOLE-4-CARBOXAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 1-(4-nitrophenyl)-5-(trifluoromethyl)-1*H*-pyrazole-4-carbonyl chloride: ESMS *m/e*: 629.2 $(M + H)^+$.

Example 625

***N*-[3-(1-{6-[(DIPHENYLACETYL)AMINO]HEXYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and diphenylacetyl chloride: ESMS *m/e*: 540.3 $(M + H)^+$.

Example 626

5-(3,5-DICHLOROPHENOXY)-*N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-2-FURAMIDE:

Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-(3,5-dichlorophenoxy)-2-furoyl chloride: ESMS *m/e*: 600.2 $(M + H)^+$.

Example 627

***N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-2-PHENOXYNICOTINAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-phenoxynicotinoyl chloride: ESMS *m/e*: 543.3 $(M + H)^+$.

Example 628

***N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-2-NAPHTHAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-naphthoyl chloride: ESMS *m/e*: 500.3 $(M + H)^+$.

Example 629

1-BENZYL-3-*TERT*-BUTYL-*N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-1*H*-PYRAZOLE-5-CARBOXAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 1-benzyl-3-*tert*-butyl-1*H*-pyrazole-5-carbonyl chloride: ESMS *m/e*: 586.3 $(M + H)^+$.

Example 630

3-CHLORO-*N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-4-(ISOPROPYLSULFONYL)-2-THIOPHENECARBOXAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-chloro-4-(isopropylsulfonyl)-2-thiophenecarbonyl chloride: ESMS *m/e*: 596.2 $(M + H)^+$.

Example 631

***N*-[3-(1-{6-[(ANILINOCARBONYL)AMINO]HEXYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and phenyl isocyanate : ESMS *m/e*: 465.2 $(M + H)^+$.

Example 632

***N*-{3-[1-(6-{[(2,4-DICHLOROANILINO)CARBONYL]AMINO}HEXYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2,4-dichlorophenyl isocyanate: ESMS *m/e*: 533.2 $(M + H)^+$.

Example 633

***N*-(6-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}HEXYL)-1-PHENYL-5-PROPYL-1*H*-PYRAZOLE-4-CARBOXAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 1-phenyl-5-propyl-1*H*-pyrazole-4-carbonyl chloride: ESMS *m/e*: 558.3 $(M + H)^+$.

Example 634

2-METHYL-*N*-{3-[1-(6-{[(1-NAPHTHYLAMINO)CARBONYL]AMINO}HEXYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 1-naphthyl isocyanate: ESMS *m/e*: 515.3 $(M + H)^+$.

Example 635

***N*-{3-[1-(6-{[([1,1'-BIPHENYL]-4-YLAMINO)CARBONYL]AMINO}HEXYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 4-biphenyl isocyanate: ESMS *m/e*: 541.3 $(M + H)^+$.

Example 636

2-METHYL-*N*-{3-[1-(6-{[(2-NAPHTHYLAMINO)CARBONYL]AMINO}HEXYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(6-aminohexyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-naphthyl isocyanate: ESMS *m/e*: 515.3 $(M + H)^+$.

Example 637

***N*-{3-[1-(3-{[(3,4-DIMETHOXYPHENYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3,4-dimethoxybenzenesulfonyl chloride: ESMS *m/e*: 504.2 $(M + H)^+$.

Example 638

***N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-5-METHYL-3-PHENYL-4-ISOXAZOLECARBOXAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-methyl-3-phenyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 489.3 $(M + H)^+$.

Example 639

***N*-{3-[1-(3-{[(4-FLUOROPHENYL)ACETYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and (4-fluorophenyl)acetyl chloride: ESMS *m/e*: 440.3 $(M + H)^+$.

Example 640

***N*-{3-[1-(3-{[(4-CHLORO-3-NITROPHENYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 4-chloro-3-nitrobenzenesulfonyl chloride: ESMS *m/e*: 523.1 $(M + H)^+$.

Example 641

2-(4-CHLOROPHENOXY)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)NICOTINAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-(4-chlorophenoxy)nicotinoyl chloride: ESMS *m/e*: 535.2 $(M + H)^+$.

Example 642

5-(3,5-DICHLOROPHENOXY)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-FURAMIDE:

Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-(3,5-dichlorophenoxy)-2-furoyl chloride: ESMS *m/e*: 558.2 $(M + H)^+$.

Example 643

***N*-{3-[1-(3-{[(2-FLUOROPHENYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-fluorobenzenesulfonyl chloride: ESMS *m/e*: 462.2 $(M + H)^+$.

Example 644

***N*-{3-[1-(3-{[(3,5-DIMETHYL-4-ISOXAZOLYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3,5-dimethyl-4-isoxazolesulfonyl chloride: ESMS *m/e*: 463.2 $(M + H)^+$.

Example 644

***N*-{3-[1-(3-{[(4-*TERT*-BUTYLPHENYL)SULFONYL]AMINO}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 4-*tert*-butylbenzenesulfonyl chloride: ESMS *m/e*: 500.3 $(M + H)^+$.

Example 646

***N*-{3-[1-(6-AMINOHEXYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure AE and Scheme Y using *N*-(3-{1-[6-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)hexyl]-4-piperidinyl}phenyl)-2-methylpropanamide and hydrazine hydrate: ESMS *m/e*: 346.2 $(M + H)^+$.

Example 647

***N*-{3-[1-(2-{[([1,1'-BIPHENYL]-4-YLAMINO)CARBONYL]AMINO}ETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(2-aminoethyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 4-biphenyl isocyanate: ESMS *m/e*: 485.2 $(M + H)^+$.

Example 648

5-(3,5-DICHLOROPHENOXY)-*N*-(2-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}ETHYL)-3-FURAMIDE:

Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(2-aminoethyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-(3,5-dichlorophenoxy)-3-furoyl chloride: ESMS *m/e*: 544.1 $(M + H)^+$.

Example 649

N-[3-(1-{2-[(DIPHENYLACETYL)AMINO]ETHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(2-aminoethyl)-4-piperidinyl]phenyl}-2-methylpropanamide and diphenylacetyl chloride: ESMS *m/e*: 484.2 $(M + H)^+$.

Example 650

***N*-(2-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}ETHYL)-2-NAPHTHAMIDE:** Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(2-aminoethyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-naphthoyl chloride: ESMS *m/e*: 444. 2 $(M + H)^+$.

Example 651

3-(2,6-DICHLOROPHENYL)-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE:

Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 571.2 $(M + H)^+$.

Example 652

3-(2,6-DICHLOROPHENYL)-N-(5-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PENTYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q1 (THF) and Scheme AT using *N*-{3-[1-(5-aminopentyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2,6-dichlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride. ESMS *m/e*: 585.2 $(M + H)^+$.

Example 653

***N*-[3-(1-{4-[(DIPHENYLACETYL)AMINO]BUTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure Q2(THF/DCM, 1:3)) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and diphenylacetyl chloride: ESMS *m/e*: 512.0 $(M + H)^+$.

Example 654

***N*-[3-(1-{5-[(DIPHENYLACETYL)AMINO]PENTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure Q2(THF/DCM, 1:3)) and Scheme AT using *N*-{3-[1-(5-aminopentyl)-4-piperidinyl]phenyl}-2-methylpropanamide and diphenylacetyl chloride: ESMS *m/e*: 526.0 $(M + H)^+$.

Example 655

3,5-DICHLORO-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)BENZAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3,5-dichlorobenzoyl chloride: ESMS *m/e*: 490.0 $(M + H)^+$.

Example 656

5-(3,5-DICHLOROPHENOXY)-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)-2-FURAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 5-(3,5-dichlorophenoxy)-2-furoyl chloride: ESMS *m/e*: 572.0 $(M + H)^+$.

Example 657

3-CHLORO-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)BENZAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-chlorobenzoyl chloride: ESMS *m/e*: 456.0 $(M + H)^+$.

Example 658

3,4-DIFLUORO-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)BENZAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3,4-difluorobenzoyl chloride: ESMS *m/e*: 458.0 $(M + H)^+$.

Example 659

***N*-{3-[1-(4-{[(3,5-DICHLOROANILINO)CARBONYL]AMINO}BUTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-(3-{1-[4-(formylamino)butyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 3,5-dichlorophenyl isocyanate: ESMS *m/e*: 505.0 $(M + H)^+$.

Example 660

***N*-{3-[1-(4-{[([1,1'-BIPHENYL]-4-YLAMINO)CARBONYL]AMINO}BUTYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 4-biphenyl isocyanate: ESMS *m/e*: 513.0 $(M + H)^+$.

Example 661

2-METHYL-*N*-(3-{1-[5-(4-NITROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure K and Scheme B1 (K_2CO_3) using 5-chloro-1-(4-nitrophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 452.2 $(M + H)^+$.

Example 662

***N*-(3-{1-[5-(4-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 5-chloro-1-(4-fluorophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 663

2-METHYL-*N*-[3-(1-{5-OXO-5-[2-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure K and Scheme B1 (K_2CO_3) using 5-chloro-1-[2-(trifluoromethyl)phenyl]-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 475.2 $(M + H)^+$.

Example 664

***N*-(3-{1-[5-(3-BROMOPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by

Procedure K and Scheme B1 (K_2CO_3) using 1-(3-bromophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 485.1 $(M + H)^+$.

Example 665

2-METHYL-*N*-(3-{1-[5-(3-NITROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure K and Scheme B1 (K_2CO_3) using 5-chloro-1-(3-nitrophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 452.2 $(M + H)^+$.

Example 666

***N*-(3-{1-[5-(3-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(3-chlorophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 441.1 $(M + H)^+$.

Example 667

***N*-(3-{1-[5-(4-BROMOPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(4-bromophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 485.1 $(M + H)^+$.

Example 668

***N*-(3-{1-[5-(2-IODOPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(2-iodophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 533.0 $(M + H)^+$.

Example 669

***N*-(3-{1-[5-(3-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(3-fluorophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 670

2-METHYL-*N*-[3-(1-{5-OXO-5-[3-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-[3-(trifluoromethyl)phenyl]-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 475.2 $(M + H)^+$.

Example 671

***N*-(3-{1-[5-(2-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(2-fluorophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 672

***N*-(3-{1-[5-(3-IODOPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(3-iodophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 533.0 $(M + H)^+$.

Example 673

***N*-(3-{1-[5-(2-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by

Procedure K and Scheme B1 (K_2CO_3) using 1-(2-chlorophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 441.1 $(M + H)^+$.

Example 674

2-METHYL-*N*-[3-(1-{5-OXO-5-[4-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-[4-(trifluoromethyl)phenyl]-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 475.2 $(M + H)^+$.

Example 675

***N*-(3-{1-[5-(4-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(4-chlorophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 441.1 $(M + H)^+$.

Example 676

***N*-(3-{1-[5-(4-IODOPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(4-iodophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 533 $(M + H)^+$.

Example 677

***N*-(3-{1-[5-(2-BROMOPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme B1 (K_2CO_3) using 1-(2-bromophenyl)-5-chloro-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 485.1 $(M + H)^+$.

Example 678

2-(4-CHLOROPHENOXY)-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)NICOTINAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 2-(4-chlorophenoxy)nicotinoyl chloride: ESMS *m/e*: 549.0 $(M + H)^+$.

Example 679

***N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)-3,4-DIMETHOXYBENZAMIDE:** Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3,4-dimethoxybenzoyl chloride: ESMS *m/e*: 482.0 $(M + H)^+$.

Example 680

3-(2-CHLOROPHENYL)-*N*-(4-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}BUTYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(4-aminobutyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 537.0 $(M + H)^+$.

Example 681

3-(2-CHLOROPHENYL)-*N*-(5-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PENTYL)-5-METHYL-4-ISOXAZOLECARBOXAMIDE: Prepared by Procedure Q2 (THF/DCM, 1:3) and Scheme AT using *N*-{3-[1-(5-aminopentyl)-4-piperidinyl]phenyl}-2-methylpropanamide and 3-(2-chlorophenyl)-5-methyl-4-isoxazolecarbonyl chloride: ESMS *m/e*: 551.0 $(M + H)^+$.

Example 682

2-METHYL-*N*-{3-[1-(3-{1-METHYL-2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 562.2 $(M + H)^+$.

Example 683

2-METHYL-*N*-{3-[1-(3-{1-METHYL-2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 632.2 $(M + H)^+$.

Example 684

2-METHYL-*N*-{3-[1-(3-{2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and phenylhydrazine: ESMS *m/e*: 548.2 $(M + H)^+$.

Example 685

2-METHYL-*N*-{3-[1-(3-{1-PHENYL-2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-

piperidinyl)phenyl]propanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 624.2 $(M + H)^+$.

Example 686

2-METHYL-*N*-{3-[1-(3-{2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-BENZO[*G*]INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 598.2 $(M + H)^+$.

Example 687

2-METHYL-*N*-{3-[1-(3-{7-METHYL-2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 562.2(M + H)$^+$.

Example 688

2-METHYL-*N*-{3-[1-(3-{5-METHYL-2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 562.2(M + H)$^+$.

Example 689

***N*-{3-[1-(3-{5-METHOXY-2-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-3-YL}PROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 578.2 $(M + H)^+$.

Example 690

***N*-[3-(1-{3-[2-(3-FLUOROPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 691

***N*-[3-(1-{3-[2-(4-CHLOROPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 692

***N*-[3-(1-{3-[2-(4-FLUOROPHENYL)-5-METHOXY-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 528.2$(M + H)^+$.

Example 693

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 498.2 $(M + H)^+$.

Example 694

***N*-[3-(1-{3-[2-(3-FLUOROPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 582.2 $(M + H)^+$.

Example 695

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 582.2 $(M + H)^+$.

Example 696

***N*-[3-(1-{3-[2-(4-FLUOROPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 548.2 $(M + H)^+$.

Example 697

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 547.7 $(M + H)^+$.

Example 698

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 512.2($M + H)^+$.

Example 699

***N*-[3-(1-{3-[2-(3-FLUOROPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 548.2 $(M + H)^+$.

Example 700

***N*-[3-(1-{3-[2-(4-FLUOROPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 701

N-[3-(1-{3-[2-(3-FLUOROPHENYL)-5-METHOXY-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 702

N-[3-(1-{3-[2-(3-FLUOROPHENYL)-1-PHENYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 574.2 $(M + H)^+$.

Example 703

N-[3-(1-{3-[2-(4-CHLOROPHENYL)-5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 598.2 $(M + H)^+$.

Example 704

N-[3-(1-{3-[2-(3-FLUOROPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 498.2 $(M + H)^+$.

Example 705

***N*-[3-(1-{3-[2-(3-FLUOROPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 706

***N*-[3-(1-{3-[2-(3-FLUOROPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 707

***N*-[3-(1-{3-[2-(4-CHLOROPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 564.2 $(M + H)^+$.

Example 708

***N*-[3-(1-{3-[2-(4-CHLOROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-phenylhydrazine hydrochloride: ESMS *m/e*: 514.2 $(M + H)^+$.

Example 709

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 710

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 711

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 574.2 $(M + H)^+$.

Example 712

***N*-[3-(1-{3-[2-(2-FLUOROPHENYL)-5-METHOXY-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 713

N-[3-(1-{3-[2-(4-CHLOROPHENYL)-5-METHOXY-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 544.2 $(M + H)^+$.

Example 714

N-[3-(1-{3-[2-(4-FLUOROPHENYL)-1H-BENZO[G]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 548.2 $(M + H)^+$.

Example 715

N-[3-(1-{3-[2-(4-FLUOROPHENYL)-5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 582.9 $(M + H)^+$.

Example 716

N-[3-(1-{3-[2-(4-FLUOROPHENYL)-7-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 717

***N*-[3-(1-{3-[2-(4-FLUOROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 498.2 $(M + H)^+$.

Example 718

***N*-[3-(1-{3-[2-(4-FLUOROPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 512.2 $(M + H)^+$.

Example 719

***N*-[3-(1-{3-[2-(4-CHLOROPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 720

***N*-[3-(1-{3-[2-(4-CHLOROPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 721

***N*-[3-(1-{3-[2-(4-CHLOROPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 590.2 $(M + H)^+$.

Example 722

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 528.1 $(M + H)^+$.

Example 723

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 598.2 $(M + H)^+$.

Example 724

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 725

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 590.3 $(M + H)^+$.

Example 726

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-5-METHOXY-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 544.3 $(M + H)^+$.

Example 727

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 728

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 564.2 $(M + H)^+$.

Example 729

***N*-[3-(1-{3-[2-(3-CHLOROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 514.2 $(M + H)^+$.

Example 730

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 514.2 $(M + H)^+$.

Example 731

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 598.2 $(M + H)^+$.

Example 732

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 564.2 $(M + H)^+$.

Example 733

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 734

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 590.2 $(M + H)^+$.

Example 735

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 736

***N*-[3-(1-{3-[2-(2-CHLOROPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 528.2 $(M + H)^+$.

Example 737

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 606.2 $(M + H)^+$.

Example 738

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 620.2 $(M + H)^+$.

Example 739

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 682.2 $(M + H)^+$.

Example 740

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 656.2 $(M + H)^+$.

Example 741

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 690.2 $(M + H)^+$.

Example 742

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 620.2 $(M + H)^+$.

Example 743

***N*-[3-(1-{3-[2-(3-IODOPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 620.2 $(M + H)^+$.

Example 744

***N*-[3-(1-{3-[2-(4-IODOPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-

(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 690.1 $(M + H)^+$.

Example 745

N-[3-(1-{3-[2-(4-IODOPHENYL)-5-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 620.1 $(M + H)^+$.

Example 746

N-[3-(1-{3-[2-(4-IODOPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 620.1 $(M + H)^+$.

Example 747

N-[3-(1-{3-[2-(4-IODOPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 682.1 $(M + H)^+$.

Example 748

N-[3-(1-{3-[2-(4-IODOPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 620.1 $(M + H)^+$.

Example 749

***N*-[3-(1-{3-[2-(4-IODOPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 656.1 $(M + H)^+$.

Example 750

***N*-[3-(1-{3-[2-(4-IODOPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 606.1 $(M + H)^+$.

Example 751

***N*-[3-(1-{3-[2-(3-BROMOPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 642.0 $(M + H)^+$.

Example 752

***N*-[3-(1-{3-[2-(4-BROMOPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 608.0 $(M + H)^+$.

Example 753

***N*-[3-(1-{3-[2-(4-BROMOPHENYL)-7-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 572 $(M + H)^+$.

Example 754

***N*-[3-(1-{3-[2-(4-BROMOPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 642 $(M + H)^+$.

Example 755

***N*-[3-(1-{3-[2-(3-BROMOPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 608.0 $(M + H)^+$.

Example 756

***N*-[3-(1-{3-[2-(4-BROMOPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-

439

methylpropanamide and phenylhydrazine: ESMS *m/e*: 558.1 $(M + H)^+$.

Example 757

***N*-[3-(1-{3-[2-(3-BROMOPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 634.0 $(M + H)^+$.

Example 758

***N*-[3-(1-{3-[2-(3-BROMOPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 572.0 $(M + H)^+$.

Example 759

***N*-[3-(1-{3-[2-(4-BROMOPHENYL)-1-METHYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 572.0 $(M + H)^+$.

Example 760

N-[3-(1-{3-[2-(4-BROMOPHENYL)-1-PHENYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 634.0 $(M + H)^+$.

Example 761

N-[3-(1-{3-[2-(4-BROMOPHENYL)-5-METHOXY-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 588.1 $(M + H)^+$.

Example 762

N-[3-(1-{3-[2-(3-BROMOPHENYL)-7-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 572 $(M + H)^+$.

Example 763

N-[3-(1-{3-[2-(3-BROMOPHENYL)-5-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 572 $(M + H)^+$.

Example 764

N-[3-(1-{3-[2-(4-BROMOPHENYL)-5-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(4-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-

methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 572.0 $(M + H)^+$.

Example 765

N-[3-(1-{3-[2-(3-BROMOPHENYL)-5-METHOXY-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(3-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 588.0 $(M + H)^+$.

Example 766

2-METHYL-*N*-[3-(1-{3-[2-(3-NITROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and phenylhydrazine: ESMS *m/e*: 525.2 $(M + H)^+$.

Example 767

2-METHYL-*N*-[3-(1-{3-[2-(3-NITROPHENYL)-1*H*-BENZO[*G*]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 575.1 $(M + H)^+$.

Example 768

2-METHYL-*N*-[3-(1-{3-[2-(3-NITROPHENYL)-5-(TRIFLUOROMETHOXY)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 4-

(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 609.1 $(M + H)^+$.

Example 769

2-METHYL-*N*-[3-(1-{3-[5-METHYL-2-(3-NITROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 539.2 $(M + H)^+$.

Example 770

***N*-[3-(1-{3-[5-METHOXY-2-(3-NITROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:**

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 555.2 $(M + H)^+$.

Example 771

2-METHYL-*N*-[3-(1-{3-[2-(3-NITROPHENYL)-1-PHENYL-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 601.1 $(M + H)^+$.

Example 772

2-METHYL-*N*-[3-(1-{3-[1-METHYL-2-(3-NITROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-

443

(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 539.2 $(M + H)^+$.

Example 773

2-METHYL-*N*-[3-(1-{3-[7-METHYL-2-(3-NITROPHENYL)-1*H*-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(3-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 539.2 $(M + H)^+$.

Example 774

N-[3-(1-{3-[5-METHOXY-2-(4-NITROPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 4-methoxyphenylhydrazine hydrochloride: ESMS *m/e*: 555.6 $(M + H)^+$.

Example 775

N-[3-(1-{3-[2-(2-BROMOPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 557.9 $(M + H)^+$.

Example 776

2-METHYL-N-[3-(1-{3-[5-METHYL-2-(4-NITROPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

444

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 539.1 $(M + H)^+$.

Example 777

2-METHYL-N-[3-(1-{3-[2-(4-NITROPHENYL)-1H-BENZO[G]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 574.7 $(M + H)^+$.

Example 778

2-METHYL-N-(3-{1-[(5E)-5-(4-NITROPHENYL)-5-(PHENYLHYDRAZONO)PENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:
Prepared by Procedure E and Scheme AX using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and phenylhydrazine: ESMS *m/e*: 542.4 $(M + H)^+$.

Example 779

2-METHYL-N-[3-(1-{3-[7-METHYL-2-(4-NITROPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:
Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 538.8 $(M + H)^+$.

Example 780

2-METHYL-N-{3-[1-((5E)-5-(4-NITROPHENYL)-5-{[4-(TRIFLUOROMETHOXY)PHENYL]HYDRAZONO}PENTYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 626.2 $(M + H)^+$.

Example 781

N-[3-(1-{3-[2-(2-BROMOPHENYL)-1H-BENZO[G]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 608.0 $(M + H)^+$.

Example 782

N-[3-(1-{3-[2-(2-BROMOPHENYL)-5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 641.9 $(M + H)^+$.

Example 783

N-[3-(1-{3-[2-(2-BROMOPHENYL)-7-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE: Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 572.0 $(M + H)^+$.

Example 784

N-[3-(1-{3-[2-(2-BROMOPHENYL)-1-PHENYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 634 $(M + H)^+$.

Example 785

N-[3-(1-{3-[2-(2-BROMOPHENYL)-5-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 572.0 $(M + H)^+$.

Example 786

N-[3-(1-{3-[2-(2-IODOPHENYL)-5-(TRIFLUOROMETHOXY)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-(trifluoromethoxy)phenylhydrazine hydrochloride: ESMS *m/e*: 690.0 $(M + H)^+$.

Example 787

N-[3-(1-{3-[2-(2-IODOPHENYL)-5-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 4-methylphenylhydrazine hydrochloride: ESMS *m/e*: 620.2 $(M + H)^+$.

Example 788

2-METHYL-N-[3-(1-{3-[1-METHYL-2-(4-NITROPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 539.6 $(M + H)^+$.

Example 789

2-METHYL-N-[3-(1-{3-[2-(4-NITROPHENYL)-1-PHENYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE:

Prepared by Procedure E and Scheme M using 2-methyl-*N*-(3-{1-[5-(4-nitrophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)propanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 601.6 $(M + H)^+$.

Example 790

N-[3-(1-{3-[2-(2-IODOPHENYL)-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and phenylhydrazine: ESMS *m/e*: 606.1 $(M + H)^+$.

Example 791

N-[3-(1-{3-[2-(2-IODOPHENYL)-1H-BENZO[G]INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-naphthylhydrazine hydrochloride: ESMS *m/e*: 656.1 $(M + H)^+$.

Example 792

N-[3-(1-{3-[2-(2-IODOPHENYL)-1-PHENYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1,1-diphenylhydrazine hydrochloride: ESMS *m/e*: 682.1 $(M + H)^+$.

Example 793

N-[3-(1-{3-[2-(2-IODOPHENYL)-7-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-iodophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-(2-methylphenyl)hydrazine hydrochloride: ESMS *m/e*: 619.6 $(M + H)^+$.

Example 794

N-[3-(1-{3-[2-(2-BROMOPHENYL)-1-METHYL-1H-INDOL-3-YL]PROPYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:

Prepared by Procedure E and Scheme M using *N*-(3-{1-[5-(2-bromophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide and 1-methyl-1-phenylhydrazine: ESMS *m/e*: 572 $(M + H)^+$.

Example 795

4-(3,4-DIFLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-METHYL-6-OXO-1,4,5,6-TETRAHYDRO-3-PYRIDINECARBOXAMIDE:

Prepared by Procedure AC and Scheme AM using 4-(3,4-difluorophenyl)-2-methyl-6-oxo-1,4,5,6-tetrahydro-3-pyridinecarboxylic acid and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 553.0 $(M + H)^+$.

Example 796

4-(2,4-DIFLUOROPHENYL)-*N*-(3-{4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}PROPYL)-2-METHYL-6-OXO-1,4,5,6-TETRAHYDRO-3-PYRIDINECARBOXAMIDE: Prepared by Procedure AC and Scheme AM using 4-(2,4-difluorophenyl)-2-methyl-6-oxo-1,4,5,6-tetrahydro-3-pyridinecarboxylic acid and *N*-{3-[1-(3-aminopropyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 553.0 $(M + H)^+$.

Example 797

***N*-(3-{1-[4-(4-METHOXYPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure O and Scheme W using 4-(4-methoxyphenyl)-1-butanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 409 $(M + H)^+$.

Example 798

***N*-(4-{1-[3-(1,2-DIPHENYL-1*H*-INDOL-3-YL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure O and Scheme W using 3-(1,2-diphenyl-*1H*-indol-3-yl)-1-propanol and *N*-[4-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 542.0 $(M + H)^+$.

Example 799

***N*-{4-[1-(3,3-DIPHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE:** Prepared by Procedure O and Scheme W using 3,3-diphenyl-1-propanol and *N*-[4-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 427.0 $(M + H)^+$.

Example 800

2-METHYL-*N*-(3-{1-[4-(4-NITROPHENYL)BUTYL]-4-PIPERIDINYL}PHENYL): Prepared by Procedure O and Scheme W using 4-(4-nitrophenyl)-1-butanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 424.2 $(M + H)^+$.

Example 801

2-METHYL-*N*-(3-{1-[2-(1-NAPHTHYL)ETHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE Prepared by Procedure O and Scheme W using 2-(1-naphthyl)ethanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 401.2 $(M + H)^+$.

Example 802

***N*-{3-[1-(3,3-DIPHENYLPROPYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure O and Scheme W using 3,3-diphenyl-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 441.2 $(M + H)^+$.

Example 803

***N*-(3-{1-[3-(3,4-DIMETHOXYPHENYL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure O and Scheme W using 3-(3,4-dimethoxyphenyl)-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 425.2 $(M + H)^+$.

Example 804

2-METHYL-*N*-{3-[1-(3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure O and Scheme W using 3-phenyl-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 365.2 $(M + H)^+$.

Example 805

2-METHYL-*N*-(3-{1-[3-(4-PYRIDINYL)PROPYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure O and Scheme W using 3-(4-pyridinyl)-1-propanol and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 366.2 $(M + H)^+$.

Example 806

***N*-{3-[1-(4-*TERT*-BUTYLBENZYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure AJ and Scheme AV using 1-bromomethyl)-4-tert-butylbenzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 393.0 $(M + H)^+$.

Example 807

***N*-{3-[1-(4-BENZOYLBENZYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure AJ and Scheme AV using [4-(bromomethyl)phenyl](phenyl)methanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS m/e: 441.0 $(M + H)^+$.

1,2-DICHLORO-4-{[(1*S*)-3-CHLORO-1-PHENYLPROPYL]OXY}BENZENE: Prepared by Procedure A using 3,4-dichlorophenol and (1*R*)-3-chloro-1-phenyl-1-propanol.

Example 808

***N*-(3-{1-[(3*S*)-3-(3,4-DICHLOROPHENOXY)-3-PHENYLPROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A using 1,2-dichloro-4-{[(1*S*)-3-chloro-1-phenylpropyl]oxy}benzene and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 525.3 $(M + H)^+$.

Example 809

***N*-(3-{1-[6-(2-FLUOROPHENYL)-6-HYDROXYHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[6-(2-fluorophenyl)-6-oxohexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 441.3 $(M + H)^+$.

Example 810

***N*-[3-(1-{5-HYDROXY-5-[4-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using 2-methyl-*N*-[3-(1-{5-oxo-5-[4-(trifluoromethyl)phenyl]pentyl}-4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 477.2 $(M + H)^+$.

Example 811

***N*-(3-{1-[5-(4-FLUOROPHENYL)-5-HYDROXYPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[5-(4-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 427.2 $(M + H)^+$.

Example 812

***N*-(3-{1-[7-(2-FLUOROPHENYL)-7-HYDROXYHEPTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[7-(2-fluorophenyl)-7-oxoheptyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 455.2 $(M + H)^+$.

Example 813

***N*-(3-{1-[6-(3-FLUOROPHENYL)-6-HYDROXYHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[6-(3-

453

fluorophenyl)-6-oxohexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 441.2 $(M + H)^+$.

Example 814

***N*-(3-{1-[5-(2-FLUOROPHENYL)-5-HYDROXYPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[5-(2-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 427.2 $(M + H)^+$.

Example 815

***N*-(3-{1-[5-(3-FLUOROPHENYL)-5-HYDROXYPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[5-(3-fluorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 427.2 $(M + H)^+$.

Example 816

***N*-(3-{1-[5-(3-CHLOROPHENYL)-5-HYDROXYPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[5-(3-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 443.1 $(M + H)^+$.

Example 817

***N*-(3-{1-[6-(4-FLUOROPHENYL)-6-HYDROXYHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[6-(4-fluorophenyl)-6-oxohexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 441.2 $(M + H)^+$.

Example 818

***N*-(3-{1-[6-(4-CHLOROPHENYL)-6-HYDROXYHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[6-(4-chlorophenyl)-6-oxohexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 456.9 $(M + H)^+$.

Example 819

***N*-(3-{1-[5-(4-CHLOROPHENYL)-5-HYDROXYPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure L and Scheme AN using *N*-(3-{1-[5-(4-chlorophenyl)-5-oxopentyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 443.0 $(M + H)^+$.

Example 820

***N*-(4-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)BUTANAMIDE:** Prepared by Procedure F and Scheme R, without HOAc, using 9-ethyl-9*H*-carbazole-3-carbaldehyde and *N*-[4-(4-piperidinyl)phenyl]butanamide: ESMS *m/e*: 454.2 $(M + H)^+$.

Example 821

***N*-(3-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure F and Scheme R, without HOAc, using 9-ethyl-9*H*-carbazole-3-carbaldehyde and *N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 440.5 $(M + H)^+$.

Example 822

***N*-(3-{1-[(1,9-DIMETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R, without HOAc, using 1,9-dimethyl-9*H*-carbazole-3-carbaldehyde and 2-methyl-*N*-[3-

(4-piperidinyl)phenyl]propanamide: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.05-6.77 (m, 10H), 5.20-5.12 (m, 1H), 4.04 (s, 3H), 3.93 (s, 2H), 3.34-3.24 (m, 2H), 2.79 (s, 3H), 2.56-2.38 (m, 2H), 2.38-2.26 (m, 2H), 2.08-1.88 (m, 2H), 1.82-1.70 (m, 2H), 1.16 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 454.2 $(M + H)^+$.

Example 823

***N*-(3-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)CYCLOPROPANECARBOXAMIDE:** Prepared by Procedure F and Scheme R, without HOAc, using 9-ethyl-9*H*-carbazole-3-carbaldehyde and *N*-[3-(4-piperidinyl)phenyl]cyclopropanecarboxamide: ESMS *m/e*: 452.6 $(M + H)^+$.

Example 824

1-(3-{1-[(9-ETHYL-9*H*-CARBAZOL-3-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-PYRROLIDINONE: Prepared by Scheme R and Procedure F. A solution of 1-(9-ethyl-9*H*-carbazol-3-yl)ethanone (22.3 mg, 0.100 mmol) and 1-[3-(4-piperidinyl)phenyl]-2-pyrrolidinone (27.2 mg, 0.100 mmol) in 1,2-dichloroethane (1.00 mL) was treated with sodium triacetoxyborohydride (63.6 mg, 0.300 mmol) and HOAc (5.70 uL, 0.100 mmol). The mixture was stirred overnight at room temperature. The reaction mixture was treated with a saturated aqueous $NaHCO_3$ solution (10 mL). The aqueous layer was extracted with CH_2Cl_2 (3 X 10 mL) and the combined organic layers were washed with brine (10 mL), dried over $MgSO_4$ and concentrated *in vacuo*. The residue was purified by preparative TLC using 5% of NH_3 (2.0 M in methanol) in CH_2Cl_2 to give the desired product 1-(3-{1-[(9-ethyl-9*H*-carbazol-3-yl)methyl]-4-

piperidinyl}phenyl)-2- pyrrolidinone (4.60 mg, 9.43 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.04 (d, 1H, J = 7.4 Hz), 7.99 (s, 1H), 7.43-7.28 (m, 5H), 6.96 (d, 1H, J = 7.4 Hz), 4.31 (q, 2H, *J* = 6.8 Hz), 3.77 (t, 2H, J = 7.3 Hz), 3.70 (s, 2H), 3.06 (d, 2H, *J* = 10.6 Hz), 2.56-2.42 (m, 3H), 2.07 (m, 4H), 1.77 (m, 4H), 1.36 (m, 3H); ESMS *m/e*: 452.5 $(M + H)^+$.

N-{3-[1-(1*H*-INDOL-5-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE: Prepared by Procedure F and Scheme R, without HOAc, using 1*H*-indole-5-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 376.2 $(M + H)^+$.

1-(4-CHLOROBUTYL)-1*H*-INDOLE: Prepared by Procedure AH, and Scheme P using 1*H*-indole and 1-bromo-4-chlorobutane: ^{1}NMR (400 MHz, $CDCl_3$) δ 7.72-7.02 (m, 5H), 6.49 (d, 1H, *J* = 2.8 Hz), 4.13 (t, 2H, *J* = 6.8 Hz), 3.48 (t, 2H, *J* = 6.8 Hz), 2.06-1.92 (m, 2H), 1.80-1.70 (m, 2H).

1-(3-CHLOROPROPYL)-1*H*-INDOLE: Prepared by Procedure AH, and Scheme P using 1*H*-indole and 1-bromo-3-chloropropane: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.70-7.04 (m, 5H), 6.50 (d, 1H, *J* = 2.8 Hz), 4.31 (t, 2H, *J* = 6.8 Hz), 3.42 (t, 2H, *J* = 6.4 Hz), 2.28-2.20 (m, 2H).

Example 825

***N*-(4-{1-[5-(1*H*-INDOL-1-YL)PENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(5-chloropentyl)-1*H*-indole and 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 432.3 $(M + H)^+$.

Example 826

***N*-(4-{1-[5-(1*H*-INDOL-1-YL)PENTYL]-4-PIPERIDINYL}PHENYL)BUTANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(5-chloropentyl)-1*H*-indole and *N*-[4-(4-piperidinyl)phenyl]butanamide: ESMS *m/e*: 432.3 (M + H)$^+$.

Example 827

***N*-(4-{1-[5-(1*H*-INDOL-1-YL)PENTYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(5-chloropentyl)-1*H*-indole and *N*-[4-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 418.2 (M + H)$^+$.

Example 828

***N*-(4-{1-[6-(1*H*-INDOL-1-YL)HEXYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(6-chlorohexyl)-1*H*-indole and *N*-[4-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 432.3 (M + H)$^+$.

Example 829

2-METHYL-*N*-(3-{1-[(1-METHYL-1*H*-INDOL-2-YL)METHYL]-4-PIPERIDINYL}PHENYL)PROPANAMIDE: Prepared by Procedure F and Scheme R, without HOAc, using 1-methyl-1*H*-indole-2-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 390.3 (M + H)$^+$.

Example 830

***N*-{3-[1-(1*H*-INDOL-4-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R, without HOAc, using 1*H*-indole-4-carbaldehyde and 2-

methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 376.2 $(M + H)^+$.

Example 831

***N*-(4-{1-[6-(1*H*-INDOL-1-YL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(6-chlorohexyl)-1*H*-indole and 2-methyl-*N*-[4-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 446.3 $(M + H)^+$.

Example 832

***N*-{3-[1-(1*H*-INDOL-7-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R, without HOAc, using 1*H*-indole-7-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 376.2 $(M + H)^+$.

Example 833

***N*-[3-(1-{[1-(4-METHOXYPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-4-methoxybenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 482.0 $(M + H)^+$.

Example 834

METHYL 4-[4-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-1*H*-INDOL-1-YL]BENZOATE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using methyl 4-iodobenzoate and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 510.3 $(M + H)^+$.

Example 835

2-METHYL-*N*-[3-(1-{[1-(3-METHYLPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-methylbenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 466.3 $(M + H)^+$.

Example 836

***N*-[3-(1-{[1-(4-FLUOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-fluoro-4-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.66-6.92 (m, 12H), 6.65 (d, 1H, J = 3.2 Hz), 3.69 (s, 2H), 3.15-3.02 (m, 2H), 2.58-2.40 (m, 2H), 2.20-2.04 (m, 2H), 1.94-1.76 (m, 4H), 1.25 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 470.6 $(M + H)^+$.

Example 837

***N*-(3-{1-[4-(1*H*-INDOL-1-YL)BUTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(4-chlorobutyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 418.3 $(M + H)^+$.

Example 838

***N*-[3-(1-{[1-(4-CHLOROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-chloro-4-iodobenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 839

***N*-[3-(1-{[1-(3-METHOXYPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-methoxybenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 482.2 $(M + H)^+$.

Example 840

***N*-(4-{1-[4-(1*H*-INDOL-1-YL)BUTYL]-4-PIPERIDINYL}PHENYL)BUTANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(4-chlorobutyl)-1*H*-indole and *N*-[4-(4-piperidinyl)phenyl]butanamide: ESMS *m/e*: 418.2 $(M + H)^+$.

Example 841

***N*-[3-(1-{[1-(2-METHOXYPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-2-methoxybenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 482.2 $(M + H)^+$.

Example 842

***N*-[3-(1-{[1-(3-CHLOROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-chloro-3-iodobenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 843

METHYL 2-[5-({4-[3-(ISOBUTYRYLAMINO)PHENYL]-1-PIPERIDINYL}METHYL)-1*H*-INDOL-1-YL]BENZOATE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using methyl 2-iodobenzoate and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 510.2 $(M + H)^+$.

Example 844

***N*-(3-{1-[3-(1*H*-INDOL-1-YL)PROPYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure AH and Scheme P using 1-(3-chloropropyl)-1*H*-indole and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 404.2 $(M + H)^+$.

Example 845

2-METHYL-*N*-{3-[1-({1-[4-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-5-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-4-(trifluoromethyl)benzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 846

***N*-(3-{1-[(1-[1,1'-BIPHENYL]-2-YL-1*H*-INDOL-5-YL)METHYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 2-iodo-1,1'-biphenyl and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 528.3 $(M + H)^+$.

Example 847

2-METHYL-*N*-[3-(1-{[1-(2-METHYLPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-2-methylbenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 466.2 $(M + H)^+$.

Example 848

2-METHYL-*N*-[3-(1-{[1-(4-METHYLPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-4-methylbenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 466.3 $(M + H)^+$.

Example 849

***N*-[3-(1-{[1-(2-CHLOROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-chloro-2-iodobenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 850

2-METHYL-*N*-{3-[1-({1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-5-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-(trifluoromethyl)benzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ^1H NMR (400 MHz, $CDCl_3$) δ 7.80-6.94 (m, 12H), 6.69 (d, 1H, J = 3.6 Hz), 3.36 (s, 2H), 3.10-3.00 (m, 2H), 2.58-2.42 (m, 2H), 2.16-2.02 (m, 2H),

1.85-1.75 (m, 4H), 1.25 (d, 6H, J = 7.2 Hz); ESMS *m/e*: 520.2 $(M + H)^+$.

Example 851

2-METHYL-*N*-[3-(1-{[1-(2-NITROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-2-nitrobenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 497.2 $(M + H)^+$.

Example 852

***N*-[3-(1-{[1-(2-FLUOROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-fluoro-2-iodobenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 470.2 $(M + H)^+$.

Example 853

2-METHYL-*N*-[3-(1-{[1-(1-NAPHTHYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodonaphthalene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 502.2 $(M + H)^+$.

Example 854

***N*-[3-(1-{[1-(2,3-DICHLOROPHENYL)-1*H*-INDOL-5-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1,2-dichloro-3-iodobenzene and *N*-{3-[1-(1*H*-indol-5-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ^1H

NMR (400 MHz, $CDCl_3$) δ 7.68-6.94 (m, 12H), 6.68 (d, 1H, J = 2.8 Hz), 3.69 (s, 2H), 3.15-3.02 (m, 2H), 2.54-2.42 (m, 2H), 2.18-2.02 (m, 2H), 1.88-1.76 (m, 4H), 1.25 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 520.1 $(M + H)^+$.

Example 855

***N*-[3-(1-{[1-(2,3-DICHLOROPHENYL)-1*H*-INDOL-7-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1,2-dichloro-3-iodobenzene and *N*-{3-[1-(1*H*-indol-7-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 856

***N*-[3-(1-{[1-(3-METHOXYPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-methoxybenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 482.3 $(M + H)^+$.

Example 857

***N*-[3-(1-{[1-(2,3-DICHLOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1,2-dichloro-3-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 858

***N*-[3-(1-{[1-(3-CHLOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu,

465

using 1-chloro-3- iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 486.2 $(M + H)^+$.

Example 859

2-METHYL-*N*-[3-(1-{[1-(3-METHYLPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-methylbenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 466.3 $(M + H)^+$.

Example 860

***N*-[3-(1-{[1-(3-METHOXYPHENYL)-1*H*-INDOL-7-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-methoxybenzene and *N*-{3-[1-(1*H*-indol-7-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 482.3 $(M + H)^+$.

Example 861

2-METHYL-*N*-{3-[1-({1-[3-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-4-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-3-(trifluoromethyl)benzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.2 $(M + H)^+$.

Example 862

***N*-[3-(1-{[1-(3,4-DIMETHYLPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-

piperidinyl]phenyl}-2-methylpropanamide and 4-iodo-1,2-dimethylbenzene: ESMS *m/e*: 480.0 $(M + H)^+$.

Example 863

***N*-[3-(1-{[1-(3,4-DIFLUOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1,3-dichloro-5-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.0 $(M + H)^+$.

Example 864

***N*-[3-(1-{[1-(3,4-DICHLOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1,2-dichloro-4-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.0 $(M + H)^+$.

Example 865

***N*-[3-(1-{[1-(2-CHLORO-4-FLUOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 2-chloro-4-fluoro-1-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 504.0 $(M + H)^+$.

Example 866

***N*-[3-(1-{[1-(2,4-DIFLUOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 2,4-difluoro-1-iodobenzene and *N*-{3-[1-(1*H*-indol-

4-ylmethyl)-4- piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 488.0 $(M + H)^+$.

Example 867

2-METHYL-*N*-[3-(1-{[1-(3-PYRIDINYL)-1*H*-INDOL-7-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 3-iodopyridine and *N*-{3-[1-(1*H*-indol-7-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 453.1 $(M + H)^+$.

Example 868

***N*-{3-[1-(1*H*-INDOL-6-YLMETHYL)-4-PIPERIDINYL]PHENYL}-2-METHYLPROPANAMIDE:** Prepared by Procedure F and Scheme R using 1*H*-indole-6-carbaldehyde and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 376.2 $(M + H)^+$.

Example 869

2-METHYL-*N*-[3-(1-{[1-(4-PYRIDINYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 4-iodopyridine and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 870

2-METHYL-*N*-[3-(1-{[1-(2-PYRIDINYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 2-iodopyridine and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 871

***N*-[3-(1-{[1-(2-FLUOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-fluoro-2-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 470.1 $(M + H)^+$.

Example 872

***N*-[3-(1-{[1-(4-CHLOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-chloro-4-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 486.1 $(M + H)^+$.

Example 873

2-METHYL-*N*-[3-(1-{[1-(3-PYRIDINYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 3-iodopyridine and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 874

***N*-[3-(1-{[1-(2,3-DIMETHYLPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-2,3-dimethylbenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 480.1 $(M + H)^+$.

Example 875

***N*-[3-(1-{[1-(3-FLUOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-fluoro-3-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 470.1 $(M + H)^+$.

Example 876

2-METHYL-*N*-{3-[1-({1-[2-(TRIFLUOROMETHYL)PHENYL]-1*H*-INDOL-4-YL}METHYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-2-(trifluoromethyl)benzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 520.1 $(M + H)^+$.

Example 877

***N*-[3-(1-{[1-(2-CHLOROPHENYL)-1*H*-INDOL-4-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-chloro-2-iodobenzene and *N*-{3-[1-(1*H*-indol-4-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 486.1 $(M + H)^+$.

Example 878

***N*-[3-(1-{[1-(2,3-DIMETHYLPHENYL)-1*H*-INDOL-7-YL]METHYL}-4-PIPERIDINYL)PHENYL]-2-METHYLPROPANAMIDE:** Prepared by Procedure C and Scheme Q1, with CuBr in place of Cu, using 1-iodo-2,3-dimethylbenzene and *N*-{3-[1-(1*H*-indol-7-ylmethyl)-4-piperidinyl]phenyl}-2-methylpropanamide: ESMS *m/e*: 480.0 $(M + H)^+$.

470

2-METHYL-*N*-[3-(1-{5-OXO-5-[4-(TRIFLUOROMETHYL)PHENYL]PENTYL}-4-PIPERIDINYL)PHENYL]PROPANAMIDE: Prepared by Procedure K and Scheme E using 5-chloro-1-[4-(trifluoromethyl)phenyl]-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 475.1 (M + H)$^+$.

***N*-(3-{1-[5-(4-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using 5-chloro-1-(4-fluorophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 (M + H)$^+$.

***N*-(3-{1-[5-(3-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using 5-chloro-1-(3-fluorophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 (M + H)$^+$.

***N*-(3-{1-[5-(3-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using 5-chloro-1-(3-chlorophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 441.1 (M + H)$^+$.

***N*-(3-{1-[5-(4-CHLOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using 5-chloro-1-(4-chlorophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 441.1 (M + H)$^+$.

Example 879

2-METHYL-*N*-{3-[1-(3-OXO-3-PHENYLPROPYL)-4-PIPERIDINYL]PHENYL}PROPANAMIDE: Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 3-chloro-1-phenyl-1-propanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 379.3 $(M + H)^+$.

Example 880

***N*-(3-{1-[7-(2-FLUOROPHENYL)-7-OXOHEPTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 7-chloro-1-(2-fluorophenyl)-1-heptanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ^1H NMR (400 MHz, $CDCl_3$), δ 8.17 (s, br, 1H), 8.06-6.88 (m, 8H), 3.08-2.94 (m, 4H), 2.62-2.48 (m, 1H), 2.48-2.38 (m, 1H), 2.38-2.15 (m, 2H), 2.02-1.92 (m, 2H), 1.84-1.77 (m, 4H), 1.77-1.66 (m, 2H), 1.62-1.46 (m, 2H), 1.46-1.29 (M, 4H), 1.21 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 453.2 $(M + H)^+$.

Example 881

***N*-(3-{1-[5-(2-FLUOROPHENYL)-5-OXOPENTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 5-chloro-1-(2-fluorophenyl)-1-pentanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 425.2 $(M + H)^+$.

Example 882

***N*-(3-{1-[6-(3-FLUOROPHENYL)-6-OXOHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and

NaI instead of KI and 6-chloro-1-(3-fluorophenyl)-1-hexanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 439.2 $(M + H)^+$.

Example 883

***N*-(3-{1-[6-(2-FLUOROPHENYL)-6-OXOHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 6-chloro-1-(2-fluorophenyl)-1-hexanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 439.2 $(M + H)^+$.

Example 884

***N*-(3-{1-[7-(4-FLUOROPHENYL)-7-OXOHEPTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 7-chloro-1-(4-fluorophenyl)-1-heptanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 453.2 $(M + H)^+$.

Example 885

***N*-(3-{1-[6-(4-CHLOROPHENYL)-6-OXOHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 6-chloro-1-(4-chlorophenyl)-1-hexanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 455.1 $(M + H)^+$.

Example 886

***N*-(3-{1-[7-(4-CHLOROPHENYL)-7-OXOHEPTYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 7-chloro-1-(4-chlorophenyl)-1-

heptanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 469.1 $(M + H)^+$.

Example 887

***N*-(3-{1-[6-(4-FLUOROPHENYL)-6-OXOHEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure K and Scheme E using K_2CO_3 instead of Na_2CO_3 and NaI instead of KI and 6-chloro-1-(4-fluorophenyl)-1-hexanone and 2-methyl-*N*-[3-(4-piperidinyl)phenyl]propanamide: ESMS *m/e*: 439.1 $(M + H)^+$.

Example 888

***N*-(3-{1-[6-(3-ACETYLPHENOXY)-6-(2-FLUOROPHENYL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using 1-(3-hydroxyphenyl)ethanone and *N*-(3-{1-[6-(2-fluorophenyl)-6-hydroxyhexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 559.5 $(M + H)^+$.

Example 889

***N*-(3-{1-[6-(2-FLUOROPHENOXY)-6-(2-FLUOROPHENYL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using 2-fluorophenol and *N*-(3-{1-[6-(2-fluorophenyl)-6-hydroxyhexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 535.1 $(M + H)^+$.

Example 890

***N*-(3-{1-[6-(4-FLUOROPHENOXY)-6-(2-FLUOROPHENYL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using 4-fluorophenol and *N*-(3-{1-[6-(2-fluorophenyl)-6-hydroxyhexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ^1H NMR (400 MHz,

$CDCl_3$), HCl salt δ 7.72- 6.72 (m, 12H), 5.42-5.34 (m, 1H), 3.68-3.58 (m, br, 2H), 3.02-2.92 (m, 2H), 2.80-2.46 (m, 6H), 2.05-1.78 (m, 6H), 1.68-1.56 (m, 1H), 1.56-1.38 (m, 3H), 1.25 (d, 6H, J = 6.8 Hz); ESMS *m/e*: 535.1 $(M + H)^+$.

Example 891

***N*-(3-{1-[6-(2-FLUOROPHENYL)-6-(2-METHOXYPHENOXY)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using 2-methoxyphenol and *N*-(3-{1-[6-(2-fluorophenyl)-6-hydroxyhexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 547.0 $(M + H)^+$.

Example 892

***N*-(3-{1-[6-(2-FLUOROPHENYL)-6-(4-METHOXYPHENOXY)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using 4-methoxyphenol and *N*-(3-{1-[6-(2-fluorophenyl)-6-hydroxyhexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 547.1 $(M + H)^+$.

Example 893

***N*-(3-{1-[6-(4-ACETYLPHENOXY)-6-(2-FLUOROPHENYL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-METHYLPROPANAMIDE:** Prepared by Procedure A and Scheme AN using 1-(4-hydroxyphenyl)ethanone and *N*-(3-{1-[6-(2-fluorophenyl)-6-hydroxyhexyl]-4-piperidinyl}phenyl)-2-methylpropanamide: ESMS *m/e*: 559.2 $(M + H)^+$.

Example 894

***N*-(3-{1-[6-(3,4-DIMETHOXYPHENOXY)-6-(2-FLUOROPHENYL)HEXYL]-4-PIPERIDINYL}PHENYL)-2-**

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 474

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 474

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

What is claimed is:

1. A compound having the structure:



;

wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, -NO_2, -CH_3, -CF_3, -$COCH_3$, -CO_2R_2, phenyl, phenoxy or straight chained or branched C_1-C_7 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl;

wherein A is -H, -F, -Cl, -Br, -CN, -NO_2, -COR_3, -CO_2R_3, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is O or NH; and

wherein n is an integer from 0 to 5 inclusive.

2. The compound of claim 1, wherein R_1 is aryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-COCH_3$, $-CO_2R_2$, straight chained or branched C_1-C_7 alkyl;

wherein R_3 is phenyl;

wherein A is H; and

wherein X is O.

3. The compound of claim 2, wherein R_2 is isopropyl.

4. The compound of claim 3, wherein the compound has the structure:



5. The compound of claim 3, wherein the compound has the structure:



6. The compound of claim 1, wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl; and wherein R_3

is aryl.

7. The compound of claim 6, wherein R_2 is isopropyl; and A is hydrogen.

8. The compound of claim 7, wherein the compound has the structure:



9. The compound of claim 7, wherein the compound has the structure:



10. A compound having the structure:



wherein R_1 is aryl or heteroaryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-OCH_3$, phenoxy, fused cyclopentanyl, straight chained or

branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein R_2 is straight-chained or branched C_1-C_4 alkyl or cyclopropyl;

wherein A is -H, -F, -Cl, -Br, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; and

wherein n is an integer from 1 to 5 inclusive.

11. The compound of claim 10, wherein R_1 is aryl optionally substituted with one or more -F, -Cl, -Br, -I or straight chained or branched C_1-C_4 alkyl; and

wherein A is H.

12. The compound of claim 11, wherein R_2 is isopropyl; and

wherein n is 2.

13. The compound of claim 12, wherein the compound has the structure:



14. The compound of claim 12, wherein the compound has the structure:



15. The compound of claim 12, wherein the compound has the structure:



16. The compound of claim 10, wherein R_1 is thienyl; and wherein A is H.

17. The compound of claim 16, wherein R_2 is isopropyl.

18. The compound of claim 17, wherein the compound has the structure:



19. A compound having the structure:



wherein W is



or



;

wherein each R_1 is independently hydrogen, methyl or ethyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is hydrogen, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -H, -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl.

wherein each A is independently -H, -F, -Cl, -Br, -CN, -NO_2, -COR_3, -CO_2R_3, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is O, NR_3, CO or may be absent; and

wherein Y is hydrogen, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -I, -CN, -NO_2, straight chained or branched C_1-C_7 alkyl.

20. The compound of claim 19, wherein W is

756



;

and wherein X is O or may be absent.

21. The compound of claim 20, wherein R_2 is isopropyl.

22. The compound of claim 21, wherein the compound has the structure:



23. The compound of claim 21, wherein the compound has the structure:



24. The compound of claim 19, wherein W is



25. The compound of claim 24, wherein A is -H, -F, -Cl, -Br.

26. The compound of claim 25, wherein R_2 is isopropyl; and A is hydrogen.

27. The compound of claim 26, wherein the compound has the structure:



28. A compound having the structure:



;

wherein W is



or



wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-OCH_3$, $-CO_2CH_3$, $-CF_3$, phenyl, straight chained or branched C_1-C_7 alkyl;

wherein R_2 is straight- chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl or phenyl.

wherein each B is independently -H, -F, -Cl, -Br, -I, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, $-OCH_3$, $-OCF_3$, $-CF_3$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl or aryl, phenoxy or benzyloxy, wherein the aryl, phenoxy or benzyloxy is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-COR_1$, $-CO_2R_1$, $-OCH_3$, $-OCF_3$, $-CF_3$ or straight chained or branched C_1-C_7 alkyl.

29. The compound of claim 28, wherein W is



30. The compound of claim 29, wherein R_1 is hydrogen or phenyl optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl.

31. The compound of claim 30, wherein R_2 is isopropyl.

32. The compound of claim 31, wherein the compound has the structure:



33. The compound of claim 31, wherein the compound has the structure:



34. A compound having the structure:

wherein R_1 is hydrogen, straight chained or branched C_1-C_7 alkyl, aryl or heteroaryl, wherein the aryl or heteroaryl is optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-CF_3$, $-OCH_3$, straight chained or branched C_1-C_3 alkyl;

wherein R_2 is straight-chained or branched C_3-C_4 alkyl or cyclopropyl;

wherein R_3 is -H, -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CF_3$, -

OCH_3, or straight chained or branched C_1-C_3 alkyl, monofluoroalkyl or polyfluoroalkyl, or a phenyl ring fused to C_6 and C_7 of the indole moiety;

wherein R_4 is hydrogen or aryl optionally substituted with one or more -F, -Cl, -Br, -I, -CN, $-NO_2$, $-CF_3$, straight chained or branched C_1-C_3 alkyl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl; and

wherein n is an integer from 2 to 4 inclusive.

35. The compound of claim 34, wherein R_3 is -H, -F, -Cl, -Br, -I, -CN, $-NO_2$, $-OCF_3$ or $-OCH_3$; and

wherein R_4 is hydrogen or phenyl optionally substituted with one or more -F, -Cl or $-CF_3$.

36. The compound of claim 35, wherein R_1 is hydrogen or phenyl optionally substituted with one or more -F, -Cl, -Br, -CN, $-NO_2$, $-CF_3$, $-OCH_3$ or straight chained or branched C_1-C_3 alkyl;

37. The compound of claim 36, wherein R_2 is isopropyl.

38. The compound of claim 37, wherein the compound has the structure:

761



39. The compound of claim 37, wherein the compound has the structure:



40. The compound of claim 37, wherein the compound has the structure:



41. A compound having the structure:



wherein each R_1 is independently hydrogen or CH_3;

wherein R_2 is straight-chained or branched C_1-C_4 alkyl or cyclopropyl;

wherein R_3 is benzyl or phenyl, wherein the benzyl or phenyl is optionally substituted with a methylenenedioxy group or one or more -F or -Cl;

wherein A is -H, -F, -Cl, -Br, -CN, $-NO_2$, straight chained or branched C_1-C_7 alkyl, monofluoroalkyl or polyfluoroalkyl;

wherein X is $(CH_2)_2$, $COCH_2$ or CONH;

42. The compound of claim 41, wherein R_3 is phenyl optionally substituted with one or more -F; and

wherein A is hydrogen.

43. The compound of claim 42, wherein X is CONH.

44. The compound of claim 43, wherein R_2 is methyl.

45. The compound of claim 44, wherein the compound has the structure:



46. The compound of claim 44, wherein the compound has the structure:



wherein each Y is independently hydrogen or -F.

47. The compound of claim 46, wherein the compound has the structure:



48. The compound of claim 46, wherein the compound has the structure:



49. The compound of claim 41, wherein R_3 is benzyl optionally substituted with a methylenedioxy group or one or more -F or -Cl.

50. The compound of claim 49, wherein the compound has the structure:



wherein each Y is independently hydrogen or -F.

51. The compound of claim 50, wherein the compound has the structure:

765



52. A compound of claims 1 to 51, wherein the compound is enantiomerically pure.

53. A compound of claims 1 to 51, wherein the compound is diastereomerically pure.

54. The compound of claims 52 and 53, wherein the compound is enantiomerically and diastereomerically pure.

55. A pharmaceutical composition comprising a therapeutically amount of a compound of any of claims 1 to 51 and a pharmaceutically acceptable carrier.

56. The pharmaceutical composition of claim 55, wherein the amount of the compound is from about 0.01mg to about 500mg.

57. The pharmaceutical composition of claim 56, wherein the amount of the compound is from about 0.1mg to about 60mg.

58. The pharmaceutical composition of claim 57, wherein the amount of the compound is from about 1mg to about 20mg.

59.The pharmaceutical composition of claim 55, wherein the carrier is a liquid and the composition is a solution.

60. The pharmaceutical composition of claim 55, wherein the carrier is a solid and the composition is a tablet.

61. The pharmaceutical composition of claim 55, wherein the carrier is a gel and the composition is a suppository.

62. A process for making a pharmaceutical composition comprising admixing a therapeutically effective amount of the compound of any of claims 1 to 51 and a pharmaceutically acceptable carrier.

63. A method of treating a subject suffering from a disorder selected from the group consisting of depression, anxiety, urge incontinence, or obesity comprising administering to the subject a therapeutically effective amount of the compound of any of claims 1 to 51.

64. The method of claim 63, wherein the therapeutically effective amount is between about 0.03 and about 1000 mg per day.

65. The method of claim 64, wherein the therapeutically effective amount is between about 0.30 and about 300 mg per day.

66. The method of claim 65, wherein the therapeutically effective amount is between about 1.0 and about 100 mg per day.

67. The method of claim 63, wherein the disorder is depression.

68. The method of claim 63, wherein the disorder is anxiety.

69. The method of claim 63, wherein the disorder is obesity.

70. The method of claim 63, wherein the disorder is urge incontinence.

71.A method of reducing the body mass of a subject, which comprises administering to the subject an amount of a compound of any of claims 1 to 51 effective to reduce the body mass of the subject.

72. A method of treating a subject suffering from depression, which comprises administering to the subject an amount of a compound of any of claims 1 to 51 effective to treat the subject's depression.

73. A method of treating a subject suffering from anxiety, which comprises administering to the subject an amount of a compound of any of claims 1 to 51 effective to treat the subject's anxiety.

74.A method of alleviating urge urinary incontinence in a subject suffering from an overactive bladder, which

comprises administering to the subject an amount of the compound of any of claims 1 to 51 effective to alleviate the subject's urge urinary incontinence.

74. A method of managing obesity in a subject in need of weight loss, which comprises administering to the subject an amount of a compound of any of claims 1 to 51 effective to induce weight loss in the subject.

75. A method of managing obesity in a subject who has experienced weight loss, which comprises administering to the subject an amount of a compound of any of claims 1 to 51 effective to maintain such weight loss in the subject.

76. A method of treating overactive bladder in a subject, which comprises administering to the subject an amount of a compound of any of claims 1 to 51 effective to treat the subject's overactive bladder.

78. A method of treating a disorder in a subject, wherein the symptoms of the subject can be alleviated by treatment with an MCH1 antagonist, wherein the MCH1 antagonist is the compound of any of claims 1 to 51.

79. A method of alleviating the symptoms of a disorder in a subject, which comprises administering to the subject an amount of an MCH1 antagonist effective to alleviate the symptoms, wherein the MCH1 antagonist is the compound of any of claims 1 to 51.