



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/82, C12Q 1/68, G01N 33/53, C12N 5/40, A61K 48/00, 39/395, 38/17	A2	(11) International Publication Number: WO 96/39516 (43) International Publication Date: 12 December 1996 (12.12.96)
(21) International Application Number: PCT/US96/09286 (22) International Filing Date: 5 June 1996 (05.06.96) (30) Priority Data: 08/463,660 5 June 1995 (05.06.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/463,660 (CIP) Filed on 5 June 1995 (05.06.95) (71) Applicant (for all designated States except US): CALIFORNIA PACIFIC MEDICAL RESEARCH INSTITUTE [US/US]; 2485 Clay Street, San Francisco, CA 94115 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SMITH, Helene [US/US]; 99 Anderson, San Francisco, CA 94110 (US). CHEN, Ling- Chun [US/US]; 510 Lowell Place, Fremont, CA 94536 (US). (74) Agents: SCHIFF, J., Michael et al.; Morrison & Foerster L.L.P., 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TARGETS FOR BREAST CANCER DIAGNOSIS AND TREATMENT		
(57) Abstract cDNA sequences derived from four novel genes associated with breast cancer are provided. In over about 60 % of the cancer cell lines tested, RNA hybridizing with the cDNAs were substantially more abundant than in normal cells. Most of the cell lines also showed a duplication of the corresponding gene, which probably contributed to the increased level of RNA in the cell. However, for each of the four genes, there were some cell lines which had RNA overabundance without gene duplication. This suggests that the gene product is sufficiently important to the cancer process that cells will use several alternative mechanisms to achieve increased expression. The polynucleotides, polypeptides, and antibodies provided by this invention are useful in diagnosis or treatment of breast cancer.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

TARGETS FOR BREAST CANCER DIAGNOSIS AND TREATMENT

STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

This invention was made in part during work supported by a grant from the National Cancer Institute (P01-CA44768). The Government has certain rights in the invention.

TECHNICAL FIELD

The present invention relates generally to the field of human genetics. More specifically, it relates to the identification of novel genes associated with overabundance of RNA in human cancer, particularly breast cancer. It pertains especially to those genes and the products thereof which may be important in diagnosis and treatment.

BACKGROUND ART

One of the most pressing health issues today is breast cancer. In the industrial world, about one woman in every nine can expect to develop breast cancer in her lifetime. In the United States, it is the most common cancer amongst women, with an annual incidence of about 175,000 new cases and nearly 50,000 deaths.

Despite an ongoing improvement in our understanding of the disease, breast cancer has remained resistant to medical intervention. Most clinical initiatives are focused on early diagnosis, followed by conventional forms of intervention, particularly surgery and chemotherapy. Such interventions are of limited success, particularly in patients where the tumor has undergone metastasis. There is a pressing

need to improve the arsenal of therapies available to provide more precise and more effective treatment in a less invasive way.

A promising area for the development of new modalities has emerged from recent understanding of the genetics of cancer. Alteration of gene expression is intimately related to the uncontrolled growth and de-differentiation that are hallmarks of the disease. There are two classes of alterations that take place (Bishop). The first type is the decreased expression of recessive genes, known as tumor suppresser genes, that apparently act to prevent malignant growth. The second type is the increased expression of dominant genes, such as oncogenes, that act to promote malignant growth, or to provide some other phenotype critical for malignancy. Thus, alteration in the expression of either type of gene is a potential diagnostic indicator. Furthermore, a treatment strategy might seek to reinstate the expression of suppresser genes, or reduce the expression of dominant genes. The latter strategy is a focus of the present invention.

The most frequently studied mechanism for gene overexpression in cancer cells is generally referred to in the literature as amplification. This is a process whereby the gene is duplicated within the chromosomes of the ancestral cell into multiple copies. The process involves unscheduled replications of the region of the chromosome comprising the gene, followed by recombination of the replicated segments back into the chromosome (Alitalo et al.). As a result, 50 or more copies of the gene may be produced. The duplicated region is sometimes referred to as an "amplicon". The level of expression of the gene (that is, the amount of messenger RNA produced) escalates in the transformed cell in the same proportion as the number of copies of the gene that are made (Alitalo et al.).

Several human oncogenes have been described, some of which are duplicated in a significant proportion of breast tumors. A prototype is the *erbB2* gene (also known as *HER-2/neu*), which encodes a 185 kDa membrane growth factor receptor homologous to the epidermal growth factor receptor. *erbB2* is duplicated in 61 of 283 tumors (22%) tested in a recent survey (Adnane et al.). Other oncogenes duplicated in breast cancer are the *bcl* gene, duplicated in 34 out of 286 (12%); the *flg* gene, duplicated in 37 out of 297 (12%), the *myc* gene, duplicated in 43 out of 275 (16%) (Adnane et al.).

Work with other oncogenes, particularly those described for neuroblastoma, suggested that gene duplication of the proto-oncogene was an event involved in the more malignant forms of cancer, and could act as a predictor of clinical outcome (reviewed by Schwab et al. and Alitalo et al.). In breast cancer, duplication of the *erbB2* gene has been reported as correlating both with reoccurrence of the disease and decreased survival times (Slamon et al.). There is some evidence that *erbB2* helps identify tumors that are responsive to adjuvant chemotherapy with cyclophosphamide, doxorubicin, and fluorouracil (Muss et al.).

It is clear that only a proportion of the genes that can undergo gene duplication in breast cancer have been identified. First, chromosome abnormalities, such as double minute (DM) chromosomes and homogeneously stained regions (HSRs), are abundant in cancer cells. HSRs are chromosomal regions that appear in karyotype analysis with intermediate density Giemsa staining throughout their length, rather than with the normal pattern of alternating dark and light bands. They correspond to multiple gene repeats. HSRs are particularly abundant in breast cancers, showing up in 60-65% of tumors surveyed (Dutrillaux et al., Zafrani et al.). When such regions are checked by in situ hybridization with probes for any of 16 known human oncogenes, including *erbB2* and *myc*, only a proportion of tumors show any hybridization to HSR regions. Furthermore, only a proportion of the HSRs within each karyotype are implicated.

Second, comparative genomic hybridization (CGH) has revealed the presence of copy number increases in tumors, even in chromosomal regions outside of HSRs. In CGH, whole chromosome spreads are stained simultaneously with DNA fragments from normal cells and from cancer cells, using two different fluorochromes. The images are computer-processed for the fluorescence ratio, revealing chromosomal regions that have undergone amplification or deletion in the cancer cells (Kallioniemi et al. 1992). DNA sequence copy number increases have been detected in breast cancer cell lines (Kallioniemi et al. 1994). Although the approximate areas of affected chromosomes were shown, possible duplicated genes near the location were not identified.

Cloning genes that undergo duplication in cancer is a formidable challenge. The known human oncogenes have generally been identified by hybridizing with

probes for other known growth-promoting genes, particularly known oncogenes in other species. For example, the *erbB2* gene was identified using a probe from a chemically induced rat neuroglioblastoma (Slamon et al.). Genes with novel sequences and functions will evade this type of search.

5 Furthermore, a gene can be overexpressed so as to support tumor malignancy without being duplicated within the chromosome. Increased expression can come about through a higher level of transcription of the gene; for example, by up-regulation of the promoter or substitution with an alternative promoter. It can also occur if the transcription product is able to persist longer in the cell; for example, by
10 increasing the resistance to cytoplasmic RNase or by reducing the level of such cytoplasmic enzymes. Two examples are the epidermal growth factor receptor, overexpressed in 45% of breast cancer tumors (Klijn et al.), and the IGF-1 receptor, overexpressed in 50-93% of breast cancer tumors (Berns et al.). In almost all cases, the overexpression of each of these receptors is by a mechanism other than gene
15 duplication.

 The traditional way of examining overexpression at the messenger RNA level is by subtractive hybridization. It involves producing positive and negative cDNA strands from two RNA preparations, and looking for cDNA which is not completely hybridized by the opposing preparation. Alternatively, expression can be examined
20 by differential display (Liang et al. 1992). In this technique, cDNA is prepared from only a subpopulation of each RNA preparation, and expanded via the polymerase chain reaction using primers of particular specificity. RNA preparations are compared by gel autoradiography to detect expression differences in their respective sources. In order to survey the RNA preparations entirely, the assay must be
25 repeated with a comprehensive set of PCR primers. The method has recently been applied to breast cancer cell lines, highlighting a number of expression differences (Liang et al. 1994). By excising the corresponding region of the separating gel, it is possible to recover and sequence the cDNA.

 Problems remain in terms of applying such techniques in the search for new
30 cancer genes. Because these are tests for RNA levels, any phenotypic difference between cell lines will be detected, not just those attributable to the cancer. This leads to a number of "false positive" identifications. In addition, a number of

adjustments are made to gene expression levels when a cell becomes cancerous that are not central to the malignant transformation. Thus, even when a novel sequence is obtained from the differential display, it is far from certain that the corresponding gene is at the root of the disease process.

5

DISCLOSURE OF THE INVENTION

It is an objective of this invention to provide isolated polynucleotides, polypeptides, and antibodies derived from four novel genes which are associated with
10 breast cancer. The genes are designated CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1. These designations refer to both strands of the cDNA and fragments thereof; and to the respective corresponding messenger RNA, including splice variants, allelic variants, and fragments of any of these forms. These genes show RNA overabundance in a majority of cancer cell lines tested. A majority of the
15 cells showing RNA overabundance also have duplication of the corresponding gene. Another object of this invention is to provide materials and methods based on these polynucleotides, polypeptides, and antibodies for use in the diagnosis and treatment of cancer, particularly breast cancer.

Accordingly, one embodiment of this invention is an isolated polynucleotide
20 comprising a linear sequence contained in a polynucleotide selected from the group consisting of CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1. The linear sequence is contained in a duplicated gene or overabundant RNA in cancerous cells. The RNA may be overabundant due to gene duplication, increased RNA transcription or processing, increased RNA persistence, any combination thereof, or by any other
25 mechanism, in a proportion of breast cancer cells. Preferably, the RNA is overabundant in at least about 20% of a representative panel of breast cancer cell lines, such as the panels listed herein; more preferably, it is overabundant in at least about 40% of the panel; even more preferably, it is overabundant in at least 60% or more of the panel. Preferably, the RNA is overabundant in at least about 5% of
30 spontaneously occurring breast cancer tumors; more preferably, it is overabundant in at least about 10% of such tumors; more preferably, it is overabundant in at least about 20% of such tumors; more preferably, it is overabundant in at least about 30%

of such tumors; even more preferably, it is overabundant in at least about 50% of such tumors.

Preferably, a sequence of at least 10 nucleotides is essentially identical between the isolated polynucleotide of the invention and a cDNA from CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1; more preferably, a sequence of at least about 15 nucleotides is essentially identical; more preferably, a sequence of at least about 20 nucleotides is essentially identical; more preferably, a sequence of at least about 40 nucleotides is essentially identical; even more preferably, a sequence of at least about 70 nucleotides is essentially identical; still more preferably, a sequence of about 100 nucleotides or more is essentially identical. A further embodiment of this invention is an isolated polynucleotide comprising a linear sequence essentially identical to a sequence selected from the group consisting of SEQ. ID NO:15, SEQ. ID NO:18, SEQ. ID NO:21, SEQ. ID NO:23, SEQ. ID NO:26, SEQ. ID NO:29, and SEQ. ID NO:31. These embodiments include an isolated polynucleotide which is a DNA polynucleotide, an RNA polynucleotide, a polynucleotide probe, or a polynucleotide primer.

This invention also provides an isolated polypeptide comprising a sequence of amino acids essentially identical to the polypeptide encoded by or translated from a polynucleotide selected from the group consisting of CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1. Preferably, a sequence of at least about 5 amino acids is essentially identical between the polypeptide of this invention and that encoded by the polynucleotide; more preferably, a sequence of at least about 10 amino acids is essentially identical; more preferably, a sequence of at least 15 amino acids is essentially identical; even more preferably, a sequence of at least 20 amino acids is essentially identical; still more preferably, a sequence of about 30 amino acids or more is essentially identical. Preferably, the polypeptide comprises a linear sequence of at least 15 amino acids essentially identical to a sequence encoded by said polynucleotide. Another embodiment of this invention is a polypeptide comprising a linear sequence essentially identical to a sequence selected from the group consisting of SEQ. ID NO:17, SEQ. ID NO:20, SEQ. ID NO:25, SEQ. ID NO:28, SEQ. ID NO:30, and SEQ. ID NO:32.

A further embodiment of this invention is an antibody specific for a polypeptide embodied in this invention. This encompasses both monoclonal and isolated polyclonal antibodies.

5 A further embodiment of this invention is a method of using the polynucleotides of this invention for detecting or measuring gene duplication in cancerous cells, especially but not limited to breast cancer cells, comprising the steps of reacting DNA contained in a clinical sample with a reagent comprising the polynucleotide, said clinical sample having been obtained from an individual suspected of having cancerous cells; and comparing the amount of complexes formed
10 between the reagent and the DNA in the clinical sample with the amount of complexes formed between the reagent and DNA in a control sample.

A further embodiment is a method of using the polynucleotides of this invention for detecting or measuring overabundance of RNA in cancerous cells, especially but not limited to breast cancer cells, comprising the steps of reacting RNA
15 contained in a clinical sample with a reagent comprising the polynucleotide, said clinical sample having been obtained from an individual suspected of having cancerous cells; and comparing the amount of complexes formed between the reagent and the RNA in the clinical sample with the amount of complexes formed between the reagent and RNA in a control sample.

20 Another embodiment of this invention is a diagnostic kit for detecting or measuring gene duplication or RNA overabundance in cells contained in an individual as manifest in a clinical sample, comprising a reagent and a buffer in suitable packaging, wherein the reagent comprises a polynucleotide of this invention.

Another embodiment of this invention is a method of using a polypeptide of
25 this invention for detecting or measuring specific antibodies in a clinical sample, comprising the steps of reacting antibodies contained in the clinical sample with a reagent comprising the polypeptide, said clinical sample having been obtained from an individual suspected of having cancerous cells, especially but not limited to breast cancer cells; and comparing the amount of complexes formed between the reagent and
30 the antibodies in the clinical sample with the amount of complexes formed between the reagent and antibodies in a control sample.

Another embodiment of this invention is a method of using an antibody of this invention for detecting or measuring altered protein expression in a clinical sample, comprising the steps of reacting a polypeptide contained in the clinical sample with a reagent comprising the antibody, said clinical sample having been obtained from an individual suspected of having cancerous cells, especially but not limited to breast cancer cells; and comparing the amount of complexes formed between the reagent and the polypeptide in the clinical sample with the amount of complexes formed between the reagent and a polypeptide in a control sample. Further embodiments of this invention are diagnostic kits for detecting or measuring a polypeptide or antibody present in a clinical sample, comprising a reagent and a buffer in suitable packaging, wherein the reagent respectively comprises either an antibody or a polypeptide of this invention.

Yet another embodiment of this invention is a host cell transfected by a polynucleotide of this invention. A further embodiment of this invention is a method for using a polynucleotide for screening a pharmaceutical candidate, comprising the steps of separating progeny of the transfected host cell into a first group and a second group; treating the first group of cells with the pharmaceutical candidate; not treating the second group of cells with the pharmaceutical candidate; and comparing the phenotype of the treated cells with that of the untreated cells.

This invention also embodies a pharmaceutical preparation for use in cancer therapy, comprising a polynucleotide or polypeptide embodied by this invention, said preparation being capable of reducing the pathology of cancerous cells, especially for but not limited to breast cancer cells. Further embodiments of this invention are methods for treating an individual bearing cancerous cells, such as breast cancer cells, comprising administering any of the aforementioned pharmaceutical preparations.

Still another embodiment of this invention is a pharmaceutical preparation or active vaccine comprising a polypeptide embodied by this invention in an immunogenic form and a pharmaceutically compatible excipient. A further embodiment is a method for treatment of cancer, especially but not limited to breast cancer, either prophylactically or after cancerous cells are present in an individual being treated, comprising administration of the aforementioned pharmaceutical preparation.

BRIEF DESCRIPTION OF THE DRAWINGS

- 5 **Figure 1** is a half-tone reproduction of an autoradiogram of a differential display experiment, in which radiolabeled cDNA corresponding to a subset of total messenger RNA in different cells are compared. This is used to select cDNA corresponding to particular RNA that are overabundant in breast cancer.
- 10 **Figure 2** is a half-tone reproduction of an autoradiogram of electrophoresed DNA digests from a panel of breast cancer cell lines probed with a CH8-2a13-1 insert (Panel A) or a loading control (Panel B).
- 15 **Figure 3** is a half-tone reproduction of an autoradiogram of electrophoresed total RNA from a panel of breast cancer cell lines probed with a CH8-2a13-1 insert (Panel A) or a loading control (Panel B).
- 20 **Figure 4** is a half-tone reproduction of an autoradiogram of electrophoresed DNA digests from a panel of breast cancer cell lines probed with a CH13-2a12-1 insert.
- 25 **Figure 5** is a half-tone reproduction of an autoradiogram of electrophoresed total RNA from a panel of breast cancer cell lines probed with a CH13-2a12-1 insert.
- 30 **Figure 6** is a map of cDNA fragments obtained for the breast cancer associated genes CH1-9a11-2, CH8-2a13-1, CH13-2a12-1 and CH14-2a16-1. Regions of the fragments used to deduce sequence data listed in the application are indicated by shading. Nucleotide positions are numbered from the left-most residue for which double-strand sequence data has been obtained, which is not necessarily the 5' terminus of the corresponding message.
- Figure 7** is a listing of primers used for obtaining the cDNA sequence data for CH1-9a11-2.

Figure 8 is a listing of cDNA sequence obtained for CH1-9a11-2.

Figure 9 is a listing of the amino acid sequence corresponding to the longest open reading frame of the DNA sequence of CH1-9a11-2 shown in Figure 8. The single-letter amino acid code is used. Stop codons are indicated by a dot (●). The upper panel shows the complete amino acid translation; the lower panel shows the predicted gene product protein sequence. A possible transmembrane region is indicated by underlining.

Figure 10 is a listing of primers used for obtaining the cDNA sequence data for CH8-2a13-1.

Figure 11 is a listing of cDNA sequence obtained for CH8-2a13-1.

Figure 12 is a listing of the amino acid sequence corresponding to the longest open reading frame of the DNA sequence of CH8-2a13-1 shown in Figure 11. The upper panel shows the complete amino acid translation; the lower panel shows the predicted gene product protein sequence.

Figure 13 is a listing of the nucleotide sequence predicted for a full-length CH8-2a13-1 cDNA.

Figure 14 is a listing of the amino acid sequence corresponding to the longest open reading frame of the DNA sequence of CH8-2a13-1 shown in Figure 13.

Figure 15 is a listing of primers used for obtaining the cDNA sequence data for CH13-2a12-1.

Figure 16 is a listing of cDNA sequence obtained for CH13-2a12-1.

Figure 17 is a listing of the amino acid sequence corresponding to the longest open reading frame of the DNA sequence of CH13-2a12-1 shown in Figure 16. The upper panel shows the complete amino acid translation; the lower panel shows the predicted gene product protein sequence.

5

Figure 18 is a listing of primers used for obtaining cDNA sequence data for CH13-2a12-1..

Figure 19 is a listing of the cDNA sequence data obtained by two-directional sequencing for CH14-2a16-1.

Figure 20 is a listing of the amino acid sequence corresponding to the longest open reading frame of the DNA sequence of CH14-2a16-1 shown in Figure 19. The upper panel shows the complete amino acid translation; the lower panel shows the predicted gene product protein sequence. Residues corresponding to three zinc finger motifs are underlined, indicating that the protein may have DNA or RNA binding activity.

Figure 21 is a listing of additional DNA sequence data towards the 5' end of CH14-2a16-1 obtained by one-directional sequencing of the fragment pCH14-1.3. First two panels show nucleotide and amino acid sequence from the 5' end of the fragment; the second two panels show nucleotide and amino acid sequence from the 3' end of the fragment. Regions of overlap with pCH14-800 are underlined.

Figure 22 is a listing of the nucleotide sequences of initial fragments obtained corresponding to the four breast cancer associated genes, along with their amino acid translations.

BEST MODE FOR CARRYING OUT THE INVENTION

We have discovered and characterized four novel genes associated with breast cancer. The cDNA of these genes, and their sequences as disclosed below, provide the basis of a series of reagents that can be used in diagnosis and therapy.

Using a panel of about 15 cancer cell lines, each of the four genes was found to be duplicated in 40-60% of the cells tested. We were surprised to find that each of the four genes was duplicated in at least one cell line where studies using comparative genomic hybridization had not revealed any amplification of the corresponding chromosomal region.

Levels of expression at the mRNA level were tested in a similar panel for two of these four genes. In addition to those cell lines showing gene duplication, 17 to 37% of the lines showed RNA overabundance without gene duplication, indicating that the malignant cells had used some mechanism *other than gene duplication* to promote the abundance of RNA corresponding to these genes. All four of the breast cancer genes have open reading frames, and likely are transcribed at various levels in different cell types. Overabundance of the corresponding RNA in a cancerous cell is likely associated with overexpression of the protein gene product. Such overexpression may be manifest as increased secretion of the protein from the cell into blood or the surrounding environment, an increased density of the protein at the cell surface, or an increased accumulation the protein within the cell, in comparison to the typical level in noncancerous cells of the same tissue type.

Different tumors bear different genotypes and phenotypes, even when derived from the same tissue. Gene therapy in cancer is more likely to be effective if it is aimed at genes that are involved in supporting the malignancy of the cancer. This invention discloses genes that achieve RNA overabundance by several mechanisms, because they are more likely to be directly involved in the pathogenic process, and therefore suitable targets for pharmacological manipulation.

Features of the four novel genes, the respective mRNA, and the cDNA used to find them are provided in Table 1.

TABLE 1: Characteristics of 4 Novel Breast Cancer Genes			
Chromosome	Designation	mRNA Observed	Exemplary cDNA Fragments Cloned
1	CH1-9a11-2	5.5kb, 4.5kb	1.1 kb, 2.5 kb
8	CH8-2a13-1	4.2kb	0.6 kb (two), 3.0 kb, 4.0 kb
13	CH13-2a12-1	3.5kb, 3.2kb	1.6 kb, 3.5 kb
14	CH14-2a16-1	3.8kb, 3kb	0.8 kb, 1.3 kb, 1.6 kb, 2.5 kb

All four genes sequences are unrelated to other genes known to be overexpressed in breast cancer, including the *erbB2* gene (Adnane et al.), tissue factor (Chen et al.), mammaglobulin (Watson et al.), and *DD96* (Kocher et al.).

5 The four mRNA sequences each comprise an open reading frame. The CH1-9a11-2 gene is expressed at the mRNA level at relatively elevated levels in pancreas and testis. The CH8-2a13-1 gene is expressed at relatively elevated levels in adult heart, spleen, thymus, small intestine, colon, and tissues of the reproductive system; and at higher levels in certain tissues of the fetus. The CH13-2a12-1 gene is
 10 expressed at relatively elevated levels in heart, skeletal muscle, and testis. The CH14-2a16-1 gene is expressed at relatively elevated levels in testis. The level of expression of all four genes is especially high in a substantial proportion of breast cancer cell lines.

The CH1-9a11-2 gene encodes a protein with a putative transmembrane region, and may be expressed as a surface protein on cancer cells. The CH13-2a12-1 gene is distantly related to a *C. elegans* gene implicated in cell cycle regulation, and may play a role in the regulation of cell proliferation. The protein encoded by CH13-2a12-1 is distantly related to a vasopressin-activated calcium binding receptor, and may have Ca^{++} binding activity. The CH14-2a16-1 comprises at least five domains of a zinc finger binding motif and is distantly related to a yeast RNA binding protein.

The CH14-2a16-1 gene product is suspected of having DNA or RNA binding activity, which may relate to a role in cancer pathogenesis.

Definitions

Terms used in this application include the following:

The term "polynucleotide" refers to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof.

Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown. The following are non-limiting examples of polynucleotides: a gene or gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component.

The term polynucleotide, as used herein, refers interchangeably to double- and single-stranded molecules. Unless otherwise specified or required, any embodiment of the invention described herein that is a polynucleotide encompasses both the double-stranded form, and each of two complementary single-stranded forms known or predicted to make up the double-stranded form.

In the context of polynucleotides, a "linear sequence" or a "sequence" is an order of nucleotides in a polynucleotide in a 5' to 3' direction in which residues that neighbor each other in the sequence are contiguous in the primary structure of the polynucleotide. A "partial sequence" is a linear sequence of part of a polynucleotide which is known to comprise additional residues in one or both directions.

"Hybridization" refers to a reaction in which one or more polynucleotides react to form a complex that is stabilized via hydrogen bonding between the bases of the nucleotide residues. The hydrogen bonding may occur by Watson-Crick base pairing, Hoogsteen binding, or in any other sequence-specific manner. The complex may comprise two strands forming a duplex structure, three or more strands forming a multi-stranded complex, a single self-hybridizing strand, or any combination of these. A hybridization reaction may constitute a step in a more extensive process, such as the initiation of a PCR, or the enzymatic cleavage of a polynucleotide by a ribozyme.

Hybridization reactions can be performed under conditions of different "stringency". Relevant conditions include temperature, ionic strength, time of incubation, the presence of additional solutes in the reaction mixture such as formamide, and the washing procedure. Higher stringency conditions are those conditions, such as higher temperature and lower sodium ion concentration, which require higher minimum complementarity between hybridizing elements for a stable hybridization complex to form. Conditions that increase the stringency of a hybridization reaction are widely known and published in the art: see, for example, "Molecular Cloning: A Laboratory Manual", Second Edition (Sambrook, Fritsch & Maniatis, 1989).

When hybridization occurs in an antiparallel configuration between two single-stranded polynucleotides, those polynucleotides are described as "complementary". A double-stranded polynucleotide can be "complementary" to another polynucleotide, if hybridization can occur between one of the strands of the first polynucleotide and the second. Complementarity (the degree that one polynucleotide is complementary with another) is quantifiable in terms of the proportion of bases in opposing strands that are expected to form hydrogen bonding with each other, according to generally accepted base-pairing rules.

A linear sequence of nucleotides is "identical" to another linear sequence, if the order of nucleotides in each sequence is the same, and occurs without substitution, deletion, or material substitution. It is understood that purine and pyrimidine nitrogenous bases with similar structures can be functionally equivalent in terms of Watson-Crick base-pairing; and the inter-substitution of like nitrogenous bases, particularly uracil and thymine, or the modification of nitrogenous bases, such as by methylation, does not constitute a material substitution. An RNA and a DNA polynucleotide have identical sequences when the sequence for the RNA reflects the order of nitrogenous bases in the polyribonucleotides, the sequence for the DNA reflects the order of nitrogenous bases in the polydeoxyribonucleotides, and the two sequences satisfy the other requirements of this definition. Where one or both of the polynucleotides being compared is double-stranded, the sequences are identical if one strand of the first polynucleotide is identical with one strand of the second polynucleotide.

A linear sequence of nucleotides is "essentially identical" to another linear sequence, if both sequences are capable of hybridizing to form a duplex with the same complementary polynucleotide. Sequences that hybridize under conditions of greater stringency are more preferred. It is understood that hybridization reactions can accommodate insertions, deletions, and substitutions in the nucleotide sequence. Thus, linear sequences of nucleotides can be essentially identical even if some of the nucleotide residues do not precisely correspond or align. In general, essentially identical sequences of about 40 nucleotides in length will hybridize at about 30°C in 10 x SSC (0.15 M NaCl, 15 mM citrate buffer); preferably, they will hybridize at about 40°C in 6 x SSC; more preferably, they will hybridize at about 50°C in 6 x SSC; even more preferably, they will hybridize at about 60°C in 6 x SSC, or at about 40°C in 0.5 x SSC, or at about 30°C in 6 x SSC containing 50% formamide; still more preferably, they will hybridize at 40°C or higher in 2 x SSC or lower in the presence of 50% or more formamide. It is understood that the rigor of the test is partly a function of the length of the polynucleotide; hence shorter polynucleotides with the same homology should be tested under lower stringency and longer polynucleotides should be tested under higher stringency, adjusting the conditions accordingly. The relationship between hybridization stringency, degree of sequence

identity, and polynucleotide length is known in the art and can be calculated by standard formulae; see, e.g., Meinkoth et al. Sequences that correspond or align more closely to the invention disclosed herein are comparably more preferred. Generally, essentially identical sequences are at least about 50% identical with each other, after alignment of the homologous regions. Preferably, the sequences are at least about 60% identical; more preferably, they are at least about 70% identical; more preferably, they are at least about 80% identical; more preferably, the sequences are at least about 90% identical; even more preferably, they are at least 95% identical; still more preferably, the sequences are 100% identical. Percent identity is calculated as the percent of residues in the sequence being compared that are identical to those in the reference sequence, which is usually one of those listed or described in this application, unless stated otherwise. No penalty is imposed for introduction of gaps in the reference or comparison sequence for purposes of alignment, but the resulting fragments must be rationally derived — small gaps may not be introduced to trivially improve the identity score.

In determining whether polynucleotide sequences are essentially identical, a sequence that preserves the functionality of the polynucleotide with which it is being compared is particularly preferred. Functionality may be established by different criteria, such as ability to hybridize with a target polynucleotide, and whether the polynucleotide encodes an identical or essentially identical polypeptides. Thus, nucleotide substitutions which cause a non-conservative substitution in the encoded polypeptide are preferred over nucleotide substitutions that create a stop codon; nucleotide substitutions that cause a conservative substitution in the encoded polypeptide are more preferred, and identical nucleotide sequences are even more preferred. Insertions or deletions in the polynucleotide that result in insertions or deletions in the polypeptide are preferred over those that result in the down-stream coding region being rendered out of phase. The relative importance of hybridization properties and the polypeptide encoded by a polynucleotide depends on the application of the invention.

A “reagent” polynucleotide, polypeptide, or antibody, is a substance provided for a reaction, the substance having some known and desirable parameters for the reaction. A reaction mixture may also contain a “target”, such as a polynucleotide,

antibody, or polypeptide that the reagent is capable of reacting with. For example, in some types of diagnostic tests, the amount of the target in a sample is determined by adding a reagent, allowing the reagent and target to react, and measuring the amount of reaction product. In the context of clinical management, a "target" may also be a
5 cell, collection of cells, tissue, or organ that is the object of an administered substance, such as a pharmaceutical compound.

"cDNA" or "complementary DNA" is a single- or double-stranded DNA polynucleotide in which one strand is complementary to a messenger RNA. "Full-length cDNA" is cDNA comprised of a strand which is complementary to an
10 entire messenger RNA molecule. A "cDNA fragment" as used herein generally represents a sub-region of the full-length form, but the entire full-length cDNA may also be included. Unless explicitly specified, the term cDNA encompasses both the full-length form and the fragment form.

Different polynucleotides are said to "correspond" to each other if one is
15 ultimately derived from another. For example, messenger RNA corresponds to the gene from which it is transcribed. cDNA corresponds to the RNA from which it has been produced, such as by a reverse transcription reaction, or by chemical synthesis of a DNA based upon knowledge of the RNA sequence. cDNA also corresponds to the gene that encodes the RNA. Polynucleotides may be said to correspond even
20 when one of the pair is derived from only a portion of the other.

A "probe" when used in the context of polynucleotide manipulation refers to a polynucleotide which is provided as a reagent to detect a target potentially present in a sample of interest by hybridizing with the target. Usually, a probe will comprise a label or a means by which a label can be attached, either before or subsequent to the
25 hybridization reaction. Suitable labels include, but are not limited to radioisotopes, fluorochromes, chemiluminescent compounds, dyes, and enzymes.

A "primer" is a short polynucleotide, generally with a free 3' -OH group, that binds to a target potentially present in a sample of interest by hybridizing with the target, and thereafter promoting polymerization of a polynucleotide complementary to
30 the target. A "polymerase chain reaction" ("PCR") is a reaction in which replicate copies are made of a target polynucleotide using one or more primers, and a catalyst of polymerization, such as a reverse transcriptase or a DNA polymerase, and

particularly a thermally stable polymerase enzyme. Methods for PCR are taught in U.S. Patent Nos. 4,683,195 (Mullis) and 4,683,202 (Mullis et al.). All processes of producing replicate copies of the same polynucleotide, such as PCR or gene cloning, are collectively referred to herein as "replication."

5 An "operon" is a genetic region comprising a gene encoding a protein and functionally related 5' and 3' flanking regions. Elements within an operon include but are not limited to promoter regions, enhancer regions, repressor binding regions, transcription initiation sites, ribosome binding sites, translation initiation sites, protein encoding regions, introns and exons, and termination sites for transcription and
10 translation. A "promoter" is a DNA region capable under certain conditions of binding RNA polymerase and initiating transcription of a coding region located downstream (in the 3' direction) from the promoter. "Operably linked" refers to a juxtaposition of genetic elements, wherein the elements are in a relationship permitting them to operate in the expected manner. For instance, a promoter is
15 operably linked to a coding region if the promoter helps initiate transcription of the coding sequence. There may be intervening residues between the promoter and coding region so long as this functional relationship is maintained.

 "Gene duplication" is a term used herein to describe the process whereby an increased number of copies of a particular gene or a fragment thereof is present in a
20 particular cell or cell line. "Gene amplification" generally is synonymous with gene duplication.

 "Expression" is defined alternately in the scientific literature either as the transcription of a gene into an RNA polynucleotide, or as the transcription and subsequent translation into a polypeptide. As used herein, "expression" or "gene
25 expression" generally refers to the production of the RNA unless specified or required otherwise. Thus, "RNA overexpression" reflects the presence of more RNA (as a proportion of total RNA) from a particular gene in a cell being described, such as a cancerous cell, in relation to that of the cell it is being compared with, such as a non-cancerous cell. The protein product of the gene may or may not be produced in
30 normal or abnormal amounts. "Protein overexpression" similarly reflects the presence of relatively more protein present in or produced by, for example, a cancerous cell.

“Abundance” of RNA refers to the amount of a particular RNA present in a particular cell type. Thus, “RNA overabundance” or “overabundance of RNA” describes RNA that is present in greater proportion of total RNA in the cell type being described, compared with the same RNA as a proportion of the total RNA in a control
5 cell. A number of mechanisms may contribute to RNA overabundance in a particular cell type: for example, gene duplication, increased level of transcription of the gene, increased persistence of the RNA within the cell after it is produced, or any combination of these.

The terms “polypeptide”, “peptide” and “protein” are used interchangeably
10 herein to refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a
15 labeling component.

In the context of polypeptides, a “linear sequence” or a “sequence” is an order of amino acids in a polypeptide in an N-terminal to C-terminal direction in which residues that neighbor each other in the sequence are contiguous in the primary structure of the polypeptide. A “partial sequence” is a linear sequence of part of a
20 polypeptide which is known to comprise additional residues in one or both directions.

A linear sequence of amino acids is “essentially identical” to another sequence if the two sequences have a substantial degree of sequence identity. It is understood that the folding and the biological function of proteins can accommodate insertions, deletions, and substitutions in the amino acid sequence. Thus, linear sequences of
25 amino acids can be essentially identical even if some of the residues do not precisely correspond or align. Sequences that correspond or align more closely to the invention disclosed herein are more preferred. It is also understood that some amino acid substitutions are more easily tolerated. For example, substitution of an amino acid with hydrophobic side chains, aromatic side chains, polar side chains, side chains with
30 a positive or negative charge, or side chains comprising two or fewer carbon atoms, by another amino acid with a side chain of like properties can occur without disturbing the essential identity of the two sequences. Methods for determining

homologous regions and scoring the degree of homology are well known in the art; see for example Altschul et al. and Henikoff et al. Well-tolerated sequence differences are referred to as "conservative substitutions". Thus, sequences with conservative substitutions are preferred over those with other substitutions in the same positions; sequences with identical residues at the same positions are still more preferred. In general, amino acid sequences that are essentially identical are at least about 15% identical, and comprise at least about another 15% which are either identical or are conservative substitutions, after alignment of homologous regions. More preferably, essentially identical sequences comprise at least about 50% identical residues or conservative substitutions; more preferably, they comprise at least about 70% identical residues or conservative substitutions; more preferably, they comprise at least about 80% identical residues or conservative substitutions; more preferably, they comprise at least about 90% identical residues or conservative substitutions; more preferably, they comprise at least about 95% identical residues or conservative substitutions; even more preferably, they contain 100% identical residues.

In determining whether polypeptide sequences are essentially identical, a sequence that preserves the functionality of the polypeptide with which it is being compared is particularly preferred. Functionality may be established by different parameters, such as enzymatic activity, the binding rate or affinity in a receptor-ligand interaction, the binding affinity with an antibody, and X-ray crystallographic structure.

An "antibody" (interchangeably used in plural form) is an immunoglobulin molecule capable of specific binding to a target, such as a polypeptide, through at least one antigen recognition site, located in the variable region of the immunoglobulin molecule. As used herein, the term encompasses not only intact antibodies, but also fragments thereof, mutants thereof, fusion proteins, humanized antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen recognition site of the required specificity.

The term "antigen" refers to the target molecule that is specifically bound by an antibody through its antigen recognition site. The antigen may, but need not be chemically related to the immunogen that stimulated production of the antibody. The antigen may be polyvalent, or it may be a monovalent hapten. Examples of kinds of

antigens that can be recognized by antibodies include polypeptides, polynucleotides, other antibody molecules, oligosaccharides, complex lipids, drugs, and chemicals.

An "immunogen" is an antigen capable of stimulating production of an antibody when injected into a suitable host, usually a mammal. Compounds may be rendered
5 immunogenic by many techniques known in the art, including crosslinking or conjugating with a carrier to increase valency, mixing with a mitogen to increase the immune response, and combining with an adjuvant to enhance presentation.

An "active vaccine" is a pharmaceutical preparation for human or animal use, which is used with the intention of eliciting a specific immune response. The immune
10 response may be either humoral or cellular, systemic or secretory. The immune response may be desired for experimental purposes, for the treatment of a particular condition, for the elimination of a particular substance, or for prophylaxis against a particular condition or substance.

An "isolated" polynucleotide, polypeptide, protein, antibody, or other
15 substance refers to a preparation of the substance devoid of at least some of the other components that may also be present where the substance or a similar substance naturally occurs or is initially obtained from. Thus, for example, an isolated substance may be prepared by using a purification technique to enrich it from a source mixture. Enrichment can be measured on an absolute basis, such as weight per
20 volume of solution, or it can be measured in relation to a second, potentially interfering substance present in the source mixture. Increasing enrichments of the embodiments of this invention are increasingly more preferred. Thus, for example, a 2-fold enrichment is preferred, 10-fold enrichment is more preferred, 100-fold enrichment is more preferred, 1000-fold enrichment is even more preferred. A
25 substance can also be provided in an isolated state by a process of artificial assembly, such as by chemical synthesis or recombinant expression.

A polynucleotide used in a reaction, such as a probe used in a hybridization reaction, a primer used in a PCR, or a polynucleotide present in a pharmaceutical preparation, is referred to as "specific" or "selective" if it hybridizes or reacts with
30 the intended target more frequently, more rapidly, or with greater duration than it does with alternative substances. Similarly, an antibody is referred to as "specific" or "selective" if it binds via at least one antigen recognition site to the intended target

more frequently, more rapidly, or with greater duration than it does to alternative substances. A polynucleotide or antibody is said to "selectively inhibit" or "selectively interfere with" a reaction if it inhibits or interferes with the reaction between particular substrates to a greater degree or for a greater duration than it does with the reaction between alternative substrates. An antibody is capable of "specifically delivering" a substance if it conveys or retains that substance near a particular cell type more frequently or for a greater duration compared with other cell types.

The "effector component" of a pharmaceutical preparation is a component which modifies target cells by altering their function in a desirable way when administered to a subject bearing the cells. Some advanced pharmaceutical preparations also have a "targeting component", such as an antibody, which helps deliver the effector component more efficaciously to the target site. Depending on the desired action, the effector component may have any one of a number of modes of action. For example, it may restore or enhance a normal function of a cell, it may eliminate or suppress an abnormal function of a cell, or it may alter a cell's phenotype. Alternatively, it may kill or render dormant a cell with pathological features, such as a cancer cell. Examples of effector components are provided in a later section.

A "pharmaceutical candidate" or "drug candidate" is a compound believed to have therapeutic potential, that is to be tested for efficacy. The "screening" of a pharmaceutical candidate refers to conducting an assay that is capable of evaluating the efficacy and/or specificity of the candidate. In this context, "efficacy" refers to the ability of the candidate to effect the cell or organism it is administered to in a beneficial way: for example, the limitation of the pathology of cancerous cells.

A "cell line" or "cell culture" denotes higher eukaryotic cells grown or maintained in vitro. It is understood that the descendants of a cell may not be completely identical (either morphologically, genotypically, or phenotypically) to the parent cell. Cells described as "uncultured" are obtained directly from a living organism, and have been maintained for a limited amount of time away from the organism: not long enough or under conditions for the cells to undergo substantial replication.

“Genetic alteration” refers to a process wherein a genetic element is introduced into a cell other than by mitosis or meiosis. The element may be heterologous to the cell, or it may be an additional copy or improved version of an element already present in the cell. Genetic alteration may be effected, for example, by transfecting a cell with a recombinant plasmid or other polynucleotide through any process known in the art, such as electroporation, calcium phosphate precipitation, or contacting with a polynucleotide-liposome complex. Genetic alteration may also be effected, for example, by transduction or infection with a DNA or RNA virus or viral vector.

10 A “host cell” is a cell which has been genetically altered, or is capable of being genetically altered, by administration of an exogenous polynucleotide.

The terms “cancerous cell” or “cancer cell”, used either in the singular or plural form, refer to cells that have undergone a malignant transformation that makes them pathological to the host organism. Malignant transformation is a single- or multi-step process, which involves in part an alteration in the genetic makeup of the cell and/or the expression profile. Malignant transformation may occur either spontaneously, or via an event or combination of events such as drug or chemical treatment, radiation, fusion with other cells, viral infection, or activation or inactivation of particular genes. Malignant transformation may occur in vivo or in vitro, and can if necessary be experimentally induced.

20 A frequent feature of cancer cells is the tendency to grow in a manner that is uncontrollable by the host, but the pathology associated with a particular cancer cell may take another form, as outlined infra. Primary cancer cells (that is, cells obtained from near the site of malignant transformation) can be readily distinguished from non-cancerous cells by well-established techniques, particularly histological examination. The definition of a cancer cell, as used herein, includes not only a primary cancer cell, but any cell derived from a cancer cell ancestor. This includes metastasized cancer cells, and in vitro cultures and cell lines derived from cancer cells.

30 The “pathology” caused by a cancer cell within a host is anything that compromises the well-being or normal physiology of the host. This may involve (but is not limited to) abnormal or uncontrollable growth of the cell, metastasis, release of

cytokines or other secretory products at an inappropriate level, manifestation of a function inappropriate for its physiological milieu, interference with the normal function of neighboring cells, aggravation or suppression of an inflammatory or immunological response, or the harboring of undesirable chemical agents or invasive
5 organisms.

“Treatment” of an individual or a cell is any type of intervention in an attempt to alter the natural course of the individual or cell. For example, treatment of an individual may be undertaken to decrease or limit the pathology caused by a cancer cell harbored in the individual. Treatment includes (but is not limited to)
10 administration of a composition, such as a pharmaceutical composition, and may be performed either prophylactically, or subsequent to the initiation of a pathologic event or contact with an etiologic agent.

A “control cell” is an alternative source of cells or an alternative cell line used in an experiment for comparison purposes. Where the purpose of the experiment is to
15 establish a base line for gene copy number or expression level, it is generally preferable to use a control cell that is not a cancer cell.

The term “cancer gene” as used herein refers to any gene which is yielding transcription or translation products at a substantially altered level or in a substantially altered form in cancerous cells compared with non-cancerous cells, and which may
20 play a role in supporting the malignancy of the cell. It may be a normally quiescent gene that becomes activated (such as a dominant proto-oncogene), it may be a gene that becomes expressed at an abnormally high level (such as a growth factor receptor), it may be a gene that becomes mutated to produce a variant phenotype, or it may be a gene that becomes expressed at an abnormally low level (such as a tumor suppresser
25 gene). The present invention is particularly directed towards the discovery of genes in the first two categories.

It is understood that a “clinical sample” encompasses a variety of sample types obtained from a subject and useful in an in vitro procedure, such as a diagnostic test. The definition encompasses solid tissue samples obtained as a surgical removal, a
30 pathology specimen, or a biopsy specimen, tissue cultures or cells derived therefrom and the progeny thereof, and sections or smears prepared from any of these sources. Non-limiting examples are samples obtained from breast tissue, lymph nodes, and

tumors. The definition also encompasses blood, spinal fluid, and other liquid sample of biologic origin, and may refer to either the cells or cell fragments suspended therein, or to the liquid medium and its solutes.

5 The term "relative amount" is used where a comparison is made between a test measurement and a control measurement. Thus, the relative amount of a reagent forming a complex in a reaction is the amount reacting with a test specimen, compared with the amount reacting with a control specimen. The control specimen may be run separately in the same assay, or it may be part of the same sample (for example, normal tissue surrounding a malignant area in a tissue section).

10 A "differential" result is generally obtained from an assay in which a comparison is made between the findings of two different assay samples, such as a cancerous cell line and a control cell line. Thus, for example, "differential expression" is observed when the level of expression of a particular gene is higher in one cell than another. "Differential display" refers to a display of a component,
15 particularly RNA, from different cells to determine if there is a difference in the level of the component amongst different cells. Differential display of RNA is conducted, for example, by selective production and display of cDNA corresponding thereto. A method for performing differential display is provided in a later section.

A polynucleotide derived from or corresponding to CH1-9a11-2, CH8-2a13-1,
20 CH13-2a12-1, or CH14-2a16-1 is any of the following: the respective cDNA fragments, the corresponding messenger RNA, including splice variants and fragments thereof, both strands of the corresponding full-length cDNA and fragments thereof, and the corresponding gene. Isolated allelic variants of any of these forms are included. This invention embodies any polynucleotide corresponding to CH1-
25 9a11-2, CH8-2a13-1, CH13-2a12-1, or CH14-2a16-1 in an isolated form. It also embodies any such polynucleotide that has been cloned or transfected into a cell line.

General methods

30 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained

fully in the literature. See, for example, "Molecular Cloning: A Laboratory Manual", Second Edition (Sambrook, Fritsch & Maniatis, 1989), "Oligonucleotide Synthesis" (M.J. Gait, ed., 1984), "Animal Cell Culture" (R.I. Freshney, ed., 1987); the series "Methods in Enzymology" (Academic Press, Inc.); "Handbook of
5 Experimental Immunology" (D.M. Weir & C.C. Blackwell, Eds.), "Gene Transfer Vectors for Mammalian Cells" (J.M. Miller & M.P. Calos, eds., 1987), "Current Protocols in Molecular Biology" (F.M. Ausubel et al., eds., 1987); and "Current Protocols in Immunology" (J.E. Coligan et al., eds., 1991). All patents, patent applications, articles and publications mentioned herein, both supra and infra, are
10 hereby incorporated herein by reference.

How breast cancer associated genes were identified

To identify particular RNA that is overabundant in cancer cells, RNA was
15 prepared from both breast cancer and control cells. The amount of total cellular RNA obtained from cancer cells was compared with that from control cells using a modified differential display method. Primers were used for the 3' region of the mRNAs which have an oligo-dT sequence, followed by two other nucleotides (TiNM, where $i \approx 11$, $N \in \{A, C, G\}$, and $M \in \{A, C, G, T\}$). Random or arbitrary primers of about
20 10 nucleotides were used for replication towards what corresponds in the sequence to the 5' region of the mRNA. The labeled amplification product was then separated by molecular weight on a polyacrylamide sequencing gel.

Particular mRNAs were chosen that were present in a higher proportion of the RNA in cancerous cells, compared with control cells, according to the proportional
25 intensity amongst neighboring cDNA bands. The cDNAs was recovered directly from the gel and amplified to provide a probe for screening. Candidate polynucleotides were screened by a number of criteria, including both Northern and Southern analysis to determine if the corresponding genes were duplicated or responsible for to RNA overabundance in breast cancer cells. Sequence data of the
30 polynucleotides was obtained and compared with sequences in GenBank. Novel polynucleotides with the desired expression patterns were used to probe for longer cDNA inserts in a λ gt10 library constructed from the breast cancer cell line BT474,

which were then sequenced. Further illustration of methods for obtaining polynucleotides and sequence data for CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1 are provided in the Example section.

5 ***Preparation of polynucleotides, polypeptides and antibodies***

Polynucleotides based on the cDNA of CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, CH14-2a16-1, can be rescued from cloned plasmids and phage provided as part of this invention. They may also be obtained from breast cancer cell libraries or
10 mRNA preparations, or from normal human tissues such as placenta, by judicious use of primers or probes based on the sequence data provided herein. Alternatively, the sequence data provided herein can be used in artificial synthesis to produce a polynucleotide with an identical sequence, or that incorporates occasional variations.

Polypeptides encoded by the corresponding mRNA can be prepared by several
15 different methods, all of which will be known to a practitioner of ordinary skill. For example, the appropriate strand of the full-length cDNA can be operably linked to a suitable promoter, and transfected into a suitable host cell. The host cell is then cultured under conditions that allow transcription and translation to occur, and the polypeptide is subsequently recovered. Another convenient method is to determine
20 the polynucleotide sequence of the cDNA, and predict the polypeptide sequence according to the genetic code. A polypeptide can then be prepared directly, for example, by chemical synthesis, either identical to the predicted sequence, or incorporating occasional variations.

Antibodies against polypeptides of this invention may be prepared by any
25 method known in the art. For stimulating antibody production in an animal, it is often preferable to enhance the immunogenicity of a polypeptide by such techniques as polymerization with glutaraldehyde, or combining with an adjuvant, such as Freund's adjuvant. The immunogen is injected into a suitable experimental animal: preferably a rodent for the preparation of monoclonal antibodies; preferably a larger
30 animal such as a rabbit or sheep for preparation of polyclonal antibodies. It is preferable to provide a second or booster injection after about 4 weeks, and begin harvesting the antibody source no less than about 1 week later.

Sera harvested from the immunized animals provide a source of polyclonal antibodies. Detailed procedures for purifying specific antibody activity from a source material are known within the art. Unwanted activity cross-reacting with other antigens, if present, can be removed, for example, by running the preparation over
5 adsorbants made of those antigens attached to a solid phase, and collecting the unbound fraction. If desired, the specific antibody activity can be further purified by such techniques as protein A chromatography, ammonium sulfate precipitation, ion exchange chromatography, high-performance liquid chromatography and immunoaffinity chromatography on a column of the immunizing polypeptide coupled
10 to a solid support.

Alternatively, immune cells such as splenocytes can be recovered from the immunized animals and used to prepare a monoclonal antibody-producing cell line. See, for example, Harrow & Lane (1988), U.S. Patent Nos. 4,491,632 (J.R. Wands et al.), U.S. 4,472,500 (C. Milstein et al.), and U.S. 4,444,887 (M.K. Hoffman et
15 al.)

Briefly, an antibody-producing line can be produced inter alia by cell fusion, or by transfecting antibody-producing cells with Epstein Barr Virus, or transforming with oncogenic DNA. The treated cells are cloned and cultured, and clones are selected that produce antibody of the desired specificity. Specificity testing can be
20 performed on culture supernatants by a number of techniques, such as using the immunizing polypeptide as the detecting reagent in a standard immunoassay, or using cells expressing the polypeptide in immunohistochemistry. A supply of monoclonal antibody from the selected clones can be purified from a large volume of tissue culture supernatant, or from the ascites fluid of suitably prepared host animals injected
25 with the clone.

Effective variations of this method include those in which the immunization with the polypeptide is performed on isolated cells. Antibody fragments and other derivatives can be prepared by methods of standard protein chemistry, such as subjecting the antibody to cleavage with a proteolytic enzyme. Genetically engineered
30 variants of the antibody can be produced by obtaining a polynucleotide encoding the antibody, and applying the general methods of molecular biology to introduce mutations and translate the variant.

Use in diagnosis

Novel cDNA sequences corresponding to genes associated with cancer are
5 potentially useful as diagnostic aids. Similarly, polypeptides encoded by such genes,
and antibodies specific for these polypeptides, are also potentially useful as diagnostic
aids.

More specifically, gene duplication or overabundance of RNA in particular
cells can help identify those cells as being cancerous, and thereby play a part in the
10 initial diagnosis. Increased levels of RNA corresponding to CH1-9a11-2,
CH8-2a13-12, CH13-2a12-1, and CH14-2a16-1 are present in a substantial proportion
of breast cancer cell lines and primary breast tumors. In addition, preliminary
Northern analysis using probes for CH8-2a13-12, CH13-2a12-1, and CH14-2a16-1
indicates that these genes may be duplicated or be associated with RNA
15 overabundance in certain cell lines derived from cancers other than breast cancer,
including colon cancer, lung cancer, prostate cancer, glioma, and ovarian cancer.

For patients already diagnosed with cancer, gene duplication or overabundance
of RNA can assist with clinical management and prognosis. For example,
overabundance of RNA may be a useful predictor of disease survival, metastasis,
20 susceptibility to various regimens of standard chemotherapy, the stage of the cancer,
or its aggressiveness. See generally the article by Blast, U.S. Patent No. 4,968,603
(Slamon et al.) and PCT Application WO 94/00601 (Levine et al.). All of these
determinations are important in helping the clinician choose between the available
treatment options.

25 A particularly important diagnostic application contemplated in this invention
is the identification of patients suitable for gene-specific therapy, as outlined in the
following section. For example, treatment directed against a particular gene or gene
product is appropriate in cancers where the gene is duplicated or there is RNA
overabundance. Given a particular pharmaceutical that is directed at a particular
30 gene, a diagnostic test specific for the same gene is important in selecting patients
likely to benefit from the pharmaceutical. Given a selection of such pharmaceuticals

specific for different genes, diagnostic tests for each gene are important in selecting which pharmaceutical is likely to benefit a particular patient.

The polynucleotide, polypeptide, and antibodies embodied in this invention provide specific reagents that can be used in standard diagnostic procedures. The actual procedures for conducting diagnostic tests are extensively known in the art, and are routine for a practitioner of ordinary skill. See, for example, U.S. Patent No. 4,968,603 (Slamon et al.), and PCT Applications WO 94/00601 (Levine et al.) and WO 94/17414 (K. Keyomarsi et al.). What follows is a brief non-limiting survey of some of the known procedures that can be applied.

Generally, to perform a diagnostic method of this invention, one of the compositions of this invention is provided as a reagent to detect a target in a clinical sample with which it reacts. Thus, the polynucleotide of this invention can be used as a reagent to detect a DNA or RNA target, such as might be present in a cell with duplication or RNA overabundance of the corresponding gene. The polypeptide can be used as a reagent to detect a target for which it has a specific binding site, such as an antibody molecule or (if the polypeptide is a receptor) the corresponding ligand. The antibody can be used as a reagent to detect a target it specifically recognizes, such as the polypeptide used as an immunogen to raise it.

The target is supplied by obtaining a suitable tissue sample from an individual for whom the diagnostic parameter is to be measured. Relevant test samples are those obtained from individuals suspected of containing cancerous cells, particularly breast cancer cells. Many types of samples are suitable for this purpose, including those that are obtained near the suspected tumor site by biopsy or surgical dissection, in vitro cultures of cells derived therefrom, blood, and blood components. If desired, the target may be partially purified from the sample or amplified before the assay is conducted. The reaction is performed by contacting the reagent with the sample under conditions that will allow a complex to form between the reagent and the target.

The reaction may be performed in solution, or on a solid tissue sample, for example, using histology sections. The formation of the complex is detected by a number of techniques known in the art. For example, the reagent may be supplied with a label and unreacted reagent may be removed from the complex; the amount of remaining

label thereby indicating the amount of complex formed. Further details and alternatives for complex detection are provided in the descriptions that follow.

To determine whether the amount of complex formed is representative of cancerous or non-cancerous cells, the assay result is compared with a similar assay
5 conducted on a control sample. It is generally preferable to use a control sample which is from a non-cancerous source, and otherwise similar in composition to the clinical sample being tested. However, any control sample may be suitable provided the relative amount of target in the control is known or can be used for comparative purposes. Where the assay is being conducted on tissue sections, suitable control
10 cells with normal histopathology may surround the cancerous cells being tested. It is often preferable to conduct the assay on the test sample and the control sample simultaneously. However, if the amount of complex formed is quantifiable and sufficiently consistent, it is acceptable to assay the test sample and control sample on different days or in different laboratories.

15 A polynucleotide embodied in this invention can be used as a reagent for determining gene duplication or RNA overabundance that may be present in a clinical sample. The binding of the reagent polynucleotide to a target in a clinical sample generally relies in part on a hybridization reaction between a region of the polynucleotide reagent, and the DNA or RNA in a sample being tested.

20 If desired, the nucleic acid may be extracted from the sample, and may also be partially purified. To measure gene duplication, the preparation is preferably enriched for chromosomal DNA; to measure RNA overabundance, the preparation is preferably enriched for RNA. The target polynucleotide can be optionally subjected to any combination of additional treatments, including digestion with restriction
25 endonucleases, size separation, for example by electrophoresis in agarose or polyacrylamide, and affixed to a reaction matrix, such as a blotting material.

Hybridization is allowed to occur by mixing the reagent polynucleotide with a sample suspected of containing a target polynucleotide under appropriate reaction conditions. This may be followed by washing or separation to remove unreacted
30 reagent. Generally, both the target polynucleotide and the reagent must be at least partly equilibrated into the single-stranded form in order for complementary

sequences to hybridize efficiently. Thus, it may be useful (particularly in tests for DNA) to prepare the sample by standard denaturation techniques known in the art.

The minimum complementarity between the reagent sequence and the target sequence for a complex to form depends on the conditions under which the complex-forming reaction is allowed to occur. Such conditions include temperature, ionic strength, time of incubation, the presence of additional solutes in the reaction mixture such as formamide, and washing procedure. Higher stringency conditions are those under which higher minimum complementarity is required for stable hybridization to occur. It is generally preferable in diagnostic applications to increase the specificity of the reaction, minimizing cross-reactivity of the reagent polynucleotide alternative undesired hybridization sites in the sample. Thus, it is preferable to conduct the reaction under conditions of high stringency: for example, in the presence of high temperature, low salt, formamide, a combination of these, or followed by a low-salt wash.

In order to detect the complexes formed between the reagent and the target, the reagent is generally provided with a label. Some of the labels often used in this type of assay include radioisotopes such as ^{32}P and ^{33}P , chemiluminescent or fluorescent reagents such as fluorescein, and enzymes such as alkaline phosphatase that are capable of producing a colored solute or precipitant. The label may be intrinsic to the reagent, it may be attached by direct chemical linkage, or it may be connected through a series of intermediate reactive molecules, such as a biotin-avidin complex, or a series of inter-reactive polynucleotides. The label may be added to the reagent before hybridization with the target polynucleotide, or afterwards.

To improve the sensitivity of the assay, it is often desirable to increase the signal ensuing from hybridization. This can be accomplished by replicating either the target polynucleotide or the reagent polynucleotide, such as by a polymerase chain reaction. Alternatively, a combination of serially hybridizing polynucleotides or branched polynucleotides can be used in such a way that multiple label components become incorporated into each complex. See U.S. Patent No. 5,124,246 (Urdea et al.).

An antibody embodied in this invention can also be used as a reagent in cancer diagnosis, or for determining gene duplication or RNA overabundance that may be

present in a clinical sample. This relies on the fact that overabundance of RNA in affected cells is often associated with increased production of the corresponding polypeptide. Several of the genes up-regulated in cancer cells encode for cell surface receptors – for example, *erbB-2*, *c-myc* and epidermal growth factor. Alternatively,
5 the RNA may encode a protein kept inside the cell, or it may encode a protein secreted by the cell into the surrounding milieu.

Any such protein product can be detected in solid tissue samples and cultured cells by immunohistological techniques that will be obvious to a practitioner of ordinary skill. Generally, the tissue is preserved by a combination of techniques
10 which may include cooling, exchanging into different solvents, fixing with agents such as paraformaldehyde, or embedding in a commercially available medium such as paraffin or OCT. A section of the sample is suitably prepared and overlaid with a primary antibody specific for the protein.

The primary antibody may be provided directly with a suitable label. More
15 frequently, the primary antibody is detected using one of a number of developing reagents which are easily produced or available commercially. Typically, these developing reagents are anti-immunoglobulin or protein A, and they typically bear labels which include, but are not limited to: fluorescent markers such as fluorescein, enzymes such as peroxidase that are capable of precipitating a suitable chemical
20 compound, electron dense markers such as colloidal gold, or radioisotopes such as ¹²⁵I. The section is then visualized using an appropriate microscopic technique, and the level of labeling is compared between the suspected cancer cell and a control cell, such as cells surrounding the tumor area or those taken from an alternative site.

The amount of protein corresponding to the cancer-associated gene may be
25 detected in a standard quantitative immunoassay. If the protein is secreted or shed from the cell in any appreciable amount, it may be detectable in plasma or serum samples. Alternatively, the target protein may be solubilized or extracted from a solid tissue sample. Before quantitating, the protein may optionally be affixed to a solid phase, such as by a blot technique or using a capture antibody.

30 A number of immunoassay methods are established in the art for performing the quantitation. For example, the protein may be mixed with a pre-determined non-limiting amount of the reagent antibody specific for the protein. The reagent

antibody may contain a directly attached label, such as an enzyme or a radioisotope, or a second labeled reagent may be added, such as anti-immunoglobulin or protein A.

For a solid-phase assay, unreacted reagents are removed by washing. For a liquid-phase assay, unreacted reagents are removed by some other separation
5 technique, such as filtration or chromatography. The amount of label captured in the complex is positively related to the amount of target protein present in the test sample.

A variation of this technique is a competitive assay, in which the target protein competes with a labeled analog for binding sites on the specific antibody. In this case, the amount of label captured is negatively related to the amount of target protein
10 present in a test sample. Results obtained using any such assay on a sample from a suspected cancer-bearing source are compared with those from a non-cancerous source.

A polypeptide embodied in this invention can also be used as a reagent in cancer diagnosis, or for determining gene duplication or RNA overabundance that
15 may be present in a clinical sample. Overabundance of RNA in affected cells may result in the corresponding polypeptide being produced by the cells in an abnormal amount. On occasion, overabundance of RNA may occur concurrently with expression of the polypeptide in an unusual form. This in turn may result in stimulation of the immune response of the host to produce its own antibody molecules
20 that are specific for the polypeptide. Thus, a number of human hybridomas have been raised from cancer patients that produce antibodies against their own tumor antigens.

To use the polypeptide in the detection of such antibodies in a subject suspected of having cancer, an immunoassay is conducted. Suitable methods are generally the same as the immunoassays outlined in the preceding paragraphs, except
25 that the polypeptide is provided as a reagent, and the antibody is the target in the clinical sample which is to be quantified. For example, human IgG antibody molecules present in a serum sample may be captured with solid-phase protein A, and then overlaid with the labeled polypeptide reagent. The amount of antibody would then be proportional to the label attached to the solid phase. Alternatively, cells or
30 tissue sections expressing the polypeptide may be overlaid first with the test sample containing the antibody, and then with a detecting reagent such as labeled anti-immunoglobulin. The amount of antibody would then be proportional to the label

attached to the cells. The amount of antibody detected in the sample from a suspected cancerous source would be compared with the amount detected in a control sample.

These diagnostic procedures may be performed by diagnostic laboratories, experimental laboratories, practitioners, or private individuals. This invention provides diagnostic kits which can be used in these settings. The presence of cancer cells in the individual may be manifest in a clinical sample obtained from that individual as an alteration in the DNA, RNA, protein, or antibodies contained in the sample. An alteration in one of these components resulting from the presence of cancer may take the form of an increase or decrease of the level of the component, or an alteration in the form of the component, compared with that in a sample from a healthy individual. The clinical sample is optionally pre-treated for enrichment of the target being tested for. The user then applies a reagent contained in the kit in order to detect the changed level or alteration in the diagnostic component.

Each kit necessarily comprises the reagent which renders the procedure specific: a reagent polynucleotide, used for detecting target DNA or RNA; a reagent antibody, used for detecting target protein; or a reagent polypeptide, used for detecting target antibody that may be present in a sample to be analyzed. The reagent is supplied in a solid form or liquid buffer that is suitable for inventory storage, and later for exchange or addition into the reaction medium when the test is performed. Suitable packaging is provided. The kit may optionally provide additional components that are useful in the procedure. These optional components include buffers, capture reagents, developing reagents, labels, reacting surfaces, means for detection, control samples, instructions, and interpretive information.

25 *Use in pharmaceutical development*

Embodied in this invention are modes of treating subjects bearing cancer cells that have overabundance of the particular RNA described. The strategy used to obtain the cDNAs provided in this invention was deliberately focused on genes that achieve RNA overabundance by gene duplication in some cells, and by alternative mechanisms in other cells. These alternative mechanisms may include, for example, translocation or enhancement of transcription enhancing elements near the coding

region of the gene, deletion of repressor binding sites, or altered production of gene regulators. Such mechanisms would result in more RNA being transcribed from the same gene. Alternatively, the same amount of RNA may be transcribed, but may persist longer in the cell, resulting in greater abundance. This could occur, for example, by reduction in the level of ribozymes or protein enzymes that degrade RNA, or in the modification of the RNA to render it more resistant to such enzymes or spontaneous degradation.

Thus, different cells make use of at least two different mechanisms to achieve a single result – the overabundance of a particular RNA. This suggests that RNA overabundance of these genes is central to the cancer process in the affected cells. Interfering with the specific gene or gene product would consequently modify the cancer process. It is an objective of this invention to provide pharmaceutical compositions that enable therapy of this kind.

One way this invention achieves this objective is through screening candidate drugs. The general screening strategy is to apply the candidate to a manifestation of a gene associated with cancer, and then determine whether the effect is beneficial and specific. For example, a composition that interferes with a polynucleotide or polypeptide corresponding any of the novel cancer-associated genes described herein has the potential to block the associated pathology when administered to a tumor of the appropriate phenotype. It is not necessary that the mechanism of interference be known; only that the interference be preferential for cancerous cells (or cells near the cancer site) but not other cells.

A preferred method of screening is to provide cells in which a polynucleotide related to a cancer gene has been transfected. See, for example, PCT application WO 93/08701. A practitioner of ordinary skill will be well acquainted with techniques for transfecting eukaryotic cells, including the preparation of a suitable vector, such as a viral vector; conveying the vector into the cell, such as by electroporation; and selecting cells that have been transformed, such as by using a reporter or drug sensitivity element.

A cell line is chosen which has a phenotype desirable in testing, and which can be maintained well in culture. The cell line is transfected with a polynucleotide corresponding to one of the cancer-associated genes identified herein. Transfection is

performed such that the polynucleotide is operably linked to a genetic controlling element that permits the correct strand of the polynucleotide to be transcribed within the cell. Successful transfection can be determined by the increased abundance of the RNA compared with an untransfected cell. It is not necessary that the cell previously
5 be devoid of the RNA, only that the transfection result in a substantial increase in the level observed. RNA abundance in the cell is measured using the same polynucleotide, according to the hybridization assays outlined earlier.

Drug screening is performed by adding each candidate to a sample of transfected cells, and monitoring the effect. The experiment includes a parallel
10 sample which does not receive the candidate drug. The treated and untreated cells are then compared by any suitable phenotypic criteria, including but not limited to microscopic analysis, viability testing, ability to replicate, histological examination, the level of a particular RNA or polypeptide associated with the cells, the level of enzymatic activity expressed by the cells or cell lysates, and the ability of the cells to
15 interact with other cells or compounds. Differences between treated and untreated cells indicates effects attributable to the candidate. In a preferred method, the effect of the drug on the cell transfected with the polynucleotide is also compared with the effect on a control cell. Suitable control cells include untransfected cells of similar ancestry, cells transfected with an alternative polynucleotide, or cells transfected with
20 the same polynucleotide in an inoperative fashion. Optimally, the drug has a greater effect on operably transfected cells than on control cells.

Desirable effects of a candidate drug include an effect on any phenotype that was conferred by transfection of the cell line with the polynucleotide from the cancer-associated gene, or an effect that could limit a pathological feature of the gene in a
25 cancerous cell. Examples of the first type would be a drug that limits the overabundance of RNA in the transfected cell, limits production of the encoded protein, or limits the functional effect of the protein. The effect of the drug would be apparent when comparing results between treated and untreated cells. An example of the second type would be a drug that makes use of the transfected gene or a gene
30 product to specifically poison the cell. The effect of the drug would be apparent when comparing results between operably transfected cells and control cells.

Use in treatment

This invention also provides gene-specific pharmaceuticals in which each of the polynucleotides, polypeptides, and antibodies embodied herein as a specific active
5 ingredient in pharmaceutical compositions. Such compositions may decrease the pathology of cancer cells on their own, or render the cancer cells more susceptible to treatment by the non-specific agents, such as classical chemotherapy or radiation.

An example of how polynucleotides embodied in this invention can be effectively used in treatment is gene therapy. See, for example, Morgan et al.,
10 Culver et al., and U.S. Patent No. 5,399,346 (French et al.). The general principle is to introduce the polynucleotide into a cancer cell in a patient, and allow it to interfere with the expression of the corresponding gene, such as by complexing with the gene itself or with the RNA transcribed from the gene. Entry into the cell is facilitated by
15 suitable techniques known in the art as providing the polynucleotide in the form of a suitable vector, or encapsulation of the polynucleotide in a liposome. The polynucleotide may be provided to the cancer site by an antigen-specific homing mechanism, or by direct injection.

A preferred mode of gene therapy is to provide the polynucleotide in such a way that it will replicate inside the cell, enhancing and prolonging the interference
20 effect. Thus, the polynucleotide is operably linked to a suitable promoter, such as the natural promoter of the corresponding gene, a heterologous promoter that is intrinsically active in cancer cells, or a heterologous promoter that can be induced by a suitable agent. Preferably, the construct is designed so that the polynucleotide sequence operably linked to the promoter is complementary to the sequence of the
25 corresponding gene. Thus, once integrated into the cellular genome, the transcript of the administered polynucleotide will be complementary to the transcript of the gene, and capable of hybridizing with it. This approach is known as anti-sense therapy. See, for example, Culver et al. and Roth.

The use of antibodies embodied in this invention in the treatment of cancer
30 partly relies on the fact that genes that show RNA overabundance in cancer frequently encode cell-surface proteins. Location of these proteins at the cell surface may correspond to an important biological function of the cancer cell, such as their

interaction with other cells, the modulation of other cell-surface proteins, or triggering by an incoming cytokine.

These mechanisms suggest a variety of ways in which a specific antibody may be effective in decreasing the pathology of a cancer cell. For example, if the gene encodes for a growth receptor, then an antibody that blocks the ligand binding site or causes endocytosis of the receptor would decrease the ability of the receptor to provide its signal to the cell. It is unnecessary to have knowledge of the mechanism beforehand; the effectiveness of a particular antibody can be predicted empirically by testing with cultured cancer cells expressing the corresponding protein. Monoclonal antibodies may be more effective in this form of cancer therapy if several different clones directed at different determinants of the same cancer-associate gene product are used in combination: see PCT application WO 94/00136 (Kasprzyk et al.). Such antibody treatment may directly decrease the pathology of the cancer cells, or render them more susceptible to non-specific cytotoxic agents such as platinum (Lippman).

Another example of how antibodies can be used in cancer therapy is in the specific targeting of effector components. The protein product of the cancer-associated gene is expected to appear in high frequency on cancer cells compared to unaffected cells, due to the overabundance of the corresponding RNA. The protein therefore provides a marker for cancer cells that a specific antibody can bind to. An effector component attached to the antibody therefore becomes concentrated near the cancer cells, improving the effect on those cells and decreasing the effect on non-cancer cells. This concentration would generally occur not only near the primary tumor, but also near cancer cells that have metastasized to other tissue sites. Furthermore, if the antibody is able to induce endocytosis, this will enhance entry of the effector into the cell interior.

For the purpose of targeting, an antibody specific for the protein of the cancer-associated gene is conjugated with a suitable effector component, preferably by a covalent or high-affinity bond. Suitable effector components in such compositions include radionuclides such as ^{131}I , toxic chemicals such as vincristine, and toxic peptides such as diphtheria toxin. Other suitable effector components include peptides or polynucleotides capable of altering the phenotype of the cell in a desirable fashion:

for example, installing a tumor suppresser gene, or rendering them susceptible to immune attack.

In most applications of antibody molecules in human therapy, it is preferable to use human monoclonals, or antibodies that have been humanized by techniques known in the art. This helps prevent the antibody molecules themselves from becoming a target of the host's immune system.

An example of how polypeptides embodied in this invention can be effectively used in treatment is through vaccination. The growth of cancer cells is naturally limited in part due to immune surveillance. This refers to the recognition of cancer cells by immune recognition units, particularly antibodies and T cells, and the consequent triggering of immune effector functions that limit tumor progression. Stimulation of the immune system using a particular tumor-specific antigen enhances the effect towards the tumor expressing the antigen. Thus, an active vaccine comprising a polypeptide encoded by the cDNA of this invention would be appropriately administered to subjects having overabundance of the corresponding RNA. There may also be a prophylactic role for the vaccine in a population predisposed for developing cancer cells with overabundance of the same RNA.

Ways of increasing the effectiveness of cancer vaccines are known in the art (Beardsley, MacLean et al.). For example, synthetic antigens are conjugated to a carrier like keyhole limpet hemocyanin (KLH), and then combined with an adjuvant such as DETOX™, a mixture of mycobacterial cell walls and lipid A. Any polypeptide encoded by the four novel genes described in this invention can be used in analogous compositions.

Methods for preparing and administering polypeptide vaccines are known in the art. Peptides may be capable of eliciting an immune response on their own, or they may be rendered more immunogenic by chemical manipulation, such as cross-linking or attaching to a protein carrier like KLH. Preferably, the vaccine also comprises an adjuvant, such as alum, muramyl dipeptides, liposomes, or DETOX™. The vaccine may optionally comprise auxiliary substances such as wetting agents, emulsifying agents, and organic or inorganic salts or acids,. It also comprises a pharmaceutically acceptable excipient which is compatible with the active ingredient and appropriate for the route of administration. The desired dose for peptide vaccines

is generally from 10 µg to 1 mg, with a broad effective latitude. The vaccine is preferably administered first as a priming dose, and then again as a boosting dose, usually at least four weeks later. Further boosting doses may be given to enhance the effect. The dose and its timing are usually determined by the person responsible for the treatment.

Sequence data and deposits

The foregoing detailed description provides, inter alia, a detailed explanation of how genes associated with cancer can be identified and their cDNA obtained. Polynucleotide sequences for CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1 are provided.

The sequence data listed in this application was obtained by two-directional sequencing, except where indicated otherwise. The data are believed to be accurate — nevertheless, it is readily appreciated that the techniques of the art as used herein have the potential of introducing occasional and infrequent sequence errors. Clones and inserts obtained via PCR may also comprise occasional errors introduced during amplification. Nucleotide sequences predicted from database compilations, and sequence data obtained by one-directional sequencing may also contain occasional errors in accordance with the limitations of the underlying techniques. In addition, allelic variations to both nucleotide and amino acid sequences may occur naturally or be deliberately induced. Differences of any of these types between the sequences provided herein and the invention as practiced may be present without departing from the spirit of the invention.

Sequence data for CH8-2a13-1 and CH13-2a12-1 cDNA are believed to comprise the entire translated coding sequence, and 5' and 3' untranslated regions corresponding to those found in typical mRNA transcripts. Multiple mRNA transcripts may be found depending on the patterns of transcript processing in various cell types of interest. Sequence data for CH1-9a11-2 and CH14-2a16-1 cDNA comprise a portion of the coding sequence and 3' untranslated regions. Additional sequence is typically present in the corresponding mRNA transcripts, comprising an

additional coding region in the N-terminal direction of the protein, and possibly a 5' untranslated region.

Certain embodiments of this invention may be practiced by polynucleotide synthesis according to the data provided herein, by rescuing an appropriate insert
5 corresponding to the gene of interest from one of the deposits listed below, or by isolating a corresponding polynucleotide from a suitable tissue source. Various useful probes and primers for use in polynucleotide isolation are provided herein, or may be designed from the sequence data.

Three deposits have been made on May 31, 1996 with the American Type
10 Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852 under terms of the Budapest treaty. The deposits are described in Table 2:

TABLE 2: ATCC Deposits			
BCGF1 Accession No. [pending]	Mixture of <i>E. coli</i> with recombinant plasmids of cDNA fragments of genes associated with breast cancer. The 8 recombinant plasmids may be separated by plating on Ampicillin plates and selecting single colonies for analysis by PCR using SP6 and T7 primers.		
	Gene	Subclone	Expected size of PCR product
	CH1-9a11-2	pch1-1.1	1.1 kb
		pch1-2.5	2.5 kb
	CH8-2a13-1	pch8-600	600 bp
		pch8-3k	3.0 kb
		pch8-4k	4.0 kb
	CH14-2a16-1	pch14-800	800 kb
		pch14-1.6	1.6 kb
		pch14-1.3	1.3 kb
BCGF 2 Accession No. [pending]	Mixture of λ gt10 recombinant phages with cDNA inserts of genes associated with breast cancer. The 2 phages may be separated by growing in the <i>E. coli</i> host (strain NM514) and plating out for single plaques. These plaques can be distinguished by PCR using λ gt10 reverse and forward primers.		
	Gene	Phage	Expected size of PCR product
	CH13-2a12-1	λ ch13-3.5	3.5 kb
	CH14-2a16-1	λ ch14-2.5	2.5 kb
λBCBT474 Accession No. [pending]	cDNA library derived from breast cancer cell line BT474 in λ gt10 vector, supplemented with a cDNA library from breast cancer cell line 600PE in λ gt10 vector. The cDNA insert sizes range from about 0.5 to 5 kb. λ BCBT474 is a source of additional cDNA inserts corresponding to CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, or CH14-2a16-1 not present in BCGF-1 or BCGF-2.		

The examples presented below are provided as a further guide to a practitioner
 5 of ordinary skill in the art, and are not meant to be limiting in any way.

EXAMPLES

Example 1: Selecting cDNA for messenger RNA that is overabundant in breast cancer cells

5

Total RNA was isolated from each breast cancer cell line or control cell by centrifugation through a gradient of guanidine isothiocyanate/CsCl. The RNA was treated with RNase-free DNase (Promega, Madison, WI). After extraction with phenol-chloroform, the RNA preparations were stored at -70°C. Oligo-dT
10 polynucleotides for priming at the 3' end of messenger RNA with the sequence T₁₁NM (where N ∈ {A,C,G} and M ∈ {A,C,G,T}) were synthesized according to standard protocols. Arbitrary decamer polynucleotides (OPA01 to OPA20) for priming towards the 5' end were purchased from Operon Biotechnology, Inc., Alameda, CA.

15 The RNA was reverse-transcribed using AMV reverse transcriptase (obtained from BRL) and an anchored oligo-dT primer in a volume of 20 µL, according to the manufacturer's directions. The reaction was incubated at 37°C for 60 min and stopped by incubating at 95°C for 5 min. The cDNA obtained was used immediately or stored frozen at -70°C.

20 Differential display was conducted according to the following procedure: 1 µL cDNA was replicated in a total volume of 10 µL PCR mixture containing the appropriate T₁₁NM sequence, 0.5 µM of a decamer primer, 200 µM dNTP, 5 µCi [³⁵S]-dATP (Amersham), Taq polymerase buffer with 2.5 mM MgCl₂ and 0.3 unit Taq polymerase (Promega). Forty cycles were conducted in the following sequence:
25 94°C for 30 sec, 40°C for 2 min, 72°C for 30 sec; and then the sample was incubated at 72°C for 5 min. The replicated cDNA was separated on a 6% polyacrylamide sequencing gel. After electrophoresis, the gel was dried and exposed to X-ray film.

The autoradiogram was analyzed for labeled cDNA that was present in larger relative amount in all of the lanes corresponding to breast cancer cells, compared with
30 all of the lanes corresponding to control cells. Figure 1 provides an example of an autoradiogram from such an experiment. Lane 1 is from non-proliferating normal breast cells; lane 2 is from proliferating normal breast cells; lanes 3 to 5 are from

breast cancer cell lines BT474, SKBR3, and MCF7. The left and right side shows the pattern obtained from experiments using the same T₁₁NM sequence (T₁₁AC), but two different decamer primers. The arrows indicate the cDNA fragments that were more abundant in all three tumor lines compared with controls.

5 The assay illustrated in Figure 1 was conducted using different combinations of oligo-dT primers and decamer primers. A number of differentially expressed bands were detected when different primer combinations were used. 33 cDNA fragments were isolated from 15 displays.

10 ***Example 2: Sub-selecting cDNA that corresponds to genes that are duplicated in breast cancer cells***

cDNA fragments that were differentially expressed in the fashion described in Example 1 were excised from the dried gel and extracted by boiling at 95°C for 10
15 min. Eluted cDNA was recovered by ethanol precipitation, and replicated by PCR. The product was cloned into the pCRII vector using the TA cloning system (Invitrogen).

*Eco*RI digested placenta DNA, and *Eco*RI digested DNA from the breast cancer cell lines BT474, SKBR3 and ZR-75-30 were used to prepare Southern blots to
20 screen the cloned cDNA fragments. The cloned cDNA fragments were labeled with [32P]-dCTP, and used individually to probe the blots. A larger relative amount of binding of the probe to the lanes corresponding to the cancer cell DNA indicated that the corresponding gene had been duplicated in the cancer cells. The labeled cDNA probes were also used in Northern blots to verify that the corresponding RNA was
25 overabundant in the appropriate cell lines. Partial nucleotide sequences were obtained using M13 primers, and compared with known sequences in GenBank. Partial sequences for CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1 showed no significant identity with any known sequences.

To obtain longer cDNA corresponding to the cDNA fragments with
30 novel sequences, the fragments were used as probes to screen a cDNA library from breast cancer cell line BT474, constructed in λ -GT10 (alternatively referred to herein

as the λ -BT474 library). The longer cDNA obtained from λ -GT10 were sequenced using λ -GT10 forward and reverse primers:

λ -GT10 forward: (SEQ. ID NO:35)

5

5'-AGCAAGTTCAGCCTGGTTAAG-3'

λ -GT10 reverse: (SEQ. ID NO:36)

5'-CTTATGAGTATTTCTTCCAGGGTA-3'

The chromosomal locations of the cDNAs were determined using panels of somatic cell hybrids.

10

The probes used to identify the 4 new breast cancer genes are shown in Table

3.

TABLE 3 Primers used for Differential Display		
cDNA	Oligo-dT primer	Arbitrary primer
CH1-9a11-2	T ₁₁ CC (SEQ ID NO: 9)	SEQ ID NO:11
CH8-2a13-1	T ₁₁ AC (SEQ ID NO:10)	SEQ ID NO:12
CH13-2a12-1	T ₁₁ AC (SEQ ID NO:10)	SEQ ID NO:13
CH14-2a16-1n	T ₁₁ AC (SEQ ID NO:10)	SEQ ID NO:14

15

Example 3: Using the cDNA to test panels of breast cancer cells

To determine the proportion of breast cancers in which the putative breast cancer genes were duplicated, or showed RNA overabundance without gene duplication, the four cDNA obtained according to the selection procedures described

20

were used to probe a panel of breast cancer cell lines and primary tumors.

Gene duplication was detected either by Southern analysis or slot-blot analysis.

For Southern analysis, 10 µg of *Eco*RI digested genomic DNA from different cell lines was electrophoresed on 0.8% agarose and transferred to a HYBOND™ N+ membrane (Amersham). The filters were hybridized with 32P-labeled cDNA for the putative breast cancer gene. After an autoradiogram was obtained, the probe was stripped and the blot was re-probed using a reference probe to adjust for differences in sample loading. Either chromosome 2 probe D2S5 or chromosome 21 probe D21S6 was used as a reference. Densities of the signals on the autoradiograms were obtained using a densitometer (Molecular Dynamics). The density ratio between the breast cancer gene and the reference gene was calculated for each sample. Two samples of placental DNA digests were run in each Southern analysis as a control.

For slot-blot analysis, 1 µg of genomic DNA was denatured and slotted on the HYBOND™ membrane. D21S5 or human repetitive sequences were used as reference probes for slot blots. The density ratio between the breast cancer gene and the reference gene was calculated for each sample. 10-15 samples of placental DNA digests were used as control. Amongst the control samples, the highest density ratio was set at 1.0. The density ratio of the tumor cell lines were standardized accordingly. An arbitrary cut-off for the standardized ratio (typically 1.3) was defined to identify samples in which the putative gene had been duplicated. Each of the cell lines in the breast cancer panel was scored positively or negatively for duplication of the gene being tested.

Some of the cell lines in the panel were known to have duplicated chromosomal regions from comparative genomic hybridization analysis. In instances where the cDNA being used as probe mapped to the known amplified region, the cDNA indicated that the corresponding gene had also been duplicated. However, duplicated genes were also detected using each of the four cDNAs in instances where comparative genomic hybridization had not revealed any amplification.

Because of the nature of the technique, the standardized ratio calculated as described underestimates the gene copy number, although it is expected to rank in the same order. For example, the standardized ratio obtained for the *c-myc* gene in the SKBR3 breast cancer cell was 5.0. However, it is known that SKBR3 has approximately 50 copies of the *c-myc* gene.

To test for overabundance of RNA, 10 µg of total RNA from breast cancer cell lines or primary breast cancer tumors were electrophoresed on 0.8% agarose in the presence of the denaturant formamide, and then transferred to a nylon membrane.

5 The membrane was probed first with 32P-labeled cDNA corresponding to the putative breast cancer gene, then stripped and reprobed with 32P-labeled cDNA for the beta-actin gene to adjust for differences in sample loading. Ratios of densities between the candidate gene and the beta-actin gene were calculated. RNA from three different cultured normal epithelial cells were included in the analysis as a control for the normal level of gene expression. The highest ratio obtained from the normal cell
10 samples was set at 1.0, and the ratios in the various tumor cells were standardized accordingly.

Example 4: Chromosome 1 gene CH1-9a11-2

15 One of the cDNA obtained through the selection procedures of Examples 1 and 2 corresponded to a gene that mapped to Chromosome 1.

Table 4 summarizes the results of the analysis for gene duplication and RNA overabundance. Both quantitative and qualitative assessment is shown. The numbers shown were obtained by comparing the autoradiograph intensity of the hybridizing band in each sample with that of the controls. Several control samples were used for the gene duplication experiments, consisting of different preparations of placental DNA. The control sample with the highest level of intensity was used for standardizing the other values. Other sources used for this analysis were breast cancer cell lines with the designations shown. For reasons stated in Example 3, the quantitative number is not a direct indication of the gene copy number, although it is expected to rank in the same order. Similarly, up to 6 control samples were used for the RNA overabundance experiments, consisting of different preparations of breast cell organoids which had been maintained briefly in tissue culture until the experiment was performed. The control sample with the highest level of intensity was used for standardizing the other values. Each cell line was scored + or - according to an arbitrary cut-off value.

TABLE 4: Chromosome 1 Gene in Breast Cancer Cell Lines						
Source	CH1-9a11-2 Gene Duplication		CH1-9a11-2 RNA Overabundance			
			5.2kb		4.4kb	
Normal	-	1.00*	-	1.00**	-	1.0**
BT474	+	2.70	+	1.57	+	3.7
ZR-75-30	+	2.65		nd		nd
MDA453	+	2.86	+	5.79	+	6.2
MDA435	+	3.72	-	0.89	+	2.4
SKBR3	+	1.86	-	0.94	+	2.9
600PE	+	1.72	+	4.47	+	6.8
MDA157	+	1.49	-	1.08	+	1.4
MCF7	+	1.95		nd		nd
DU4475	+	2.02	-	1.13	+	1.5
MDA231	-	1.23	+	1.47	-	
BT20	-	1.09	-	0.83	+	1.9
T47D	-	1.05		nd		nd
UACC812	-	0.67	+	1.57	+	1.8
MDA134	-	1.19	+	5.04	+	7.1
CAMA-1	-	1.02	+	2.51	+	7.2
Incidence (%)	9/15 (60%)		7/12 (58%)		11/12 (92%)	

Gene duplication or RNA overabundance; - no duplication or overabundance; nd = not done

* Degree of gene duplication is reported relative to placental DNA preparations.

** Degree of RNA overabundance is reported relative to the highest level observed for several cultures of normal epithelial cells. Two hybridizing species of RNA are calculated and reported separately.

The gene corresponding to the CH1-9a11-2 cDNA was duplicated in 9 out of 15 (60%) of the breast cancer cell lines tested, compared with placental DNA digests (P3 and P12). The sequence of the 115 bases from the 5' end of the cDNA fragment (SEQ. ID NO:1) is shown in Figure 22. There was no substantial homology to any known gene in GenBank. One of the three possible reading frames was found to be open, with the predicted amino acid shown in Figure 22 (SEQ. ID NO:2).

The CH1-9a11-2 gene was further characterized by obtaining additional sequence information. A λ -GT10 cDNA library from the breast cancer cell line BT474 (Example 2) was screened using the initial cDNA insert, and a clone with a 2.5 kilobase insert was identified. The identified clone was subcloned into plasmid vector pCRII. T7 and Sp6 primers for regions flanking the cDNA inserts were used as initial sequencing primers:

T7 primer: (SEQ. ID NO:37)

5'-TAATACGACTCACTATAGGGAGA-3'

Sp6 primer: (SEQ. ID NO:38)

5'-CATACGATTTAGGTGACACTATAG-3'

Sequencing continued by walking along the region of interest by standard techniques, using sequencing primers based on data already obtained. Primers used in sequencing are designated 1-16 in Figure 7.

A second clone (designated pCH1-1.1) overlapping on the 5' end was obtained using CLONTECH Maration™ cDNA Amplification Kit. A map showing the overlapping regions is provided in Figure 6. Briefly, two DNA primers designated CH1a and CH1b (Figure 7) were synthesized. Polyadenylated RNA from breast cancer cell line 600PE was reverse transcribed using CH1b primer. After second strand synthesis, adaptor DNA provided in the kit was ligated to the double-stranded cDNA. The 5' end cDNA of CH1-9a11-2 was then amplified by PCR using primers CH1a and AP1 (provided in the kit). To increase the specificity of the PCR products, the first PCR products were PCR reamplified using nested primers CH1a and AP2 (provided in the kit). The PCR products were cloned into pCRII vector (Invitrogen) and screened with CH1-9a11-2 probe.

The sequence of 3452 base pairs between the 5' end of pCH1-1.1 and the poly-A tail of CH1-9a11-2 was determined by standard sequencing techniques. The DNA sequence is shown in Figure 8 (SEQ. ID NO:15). The longest open reading frame is in frame 1 (bases 1-1875), and codes for 624 amino acids before the stop codon. The corresponding amino acid sequence of this frame is shown in the upper panel of Figure 9 (SEQ. ID NO:16). The partial sequence predicted for the translated protein is listed the low panel of Figure 9 (SEQ. ID NO:17). Bases 1876 to the end of the sequence are believed to be a 3' untranslated region. A hydrophobicity analysis identified a putative membrane insertion or membrane spanning region at about amino acids 382-400, indicated in Figure 9 by underlining.

A GENINFO® BLAST search of nucleotide and peptide sequence databases was performed through the National Center for Biotechnology Information on February 23, 1996. Short segments of homology with other reported human sequences were found at the nucleotide level (<500 base pairs), but none with any ascribed function in the respective identifier. At the amino acid level, no identity higher than 30% was found with any reported eukaryotic sequences.

A CH1-9a11-2 cloned insert has been used to probe the level of relative expression in polyadenylated RNA from a panel of tissue sources. The RNA was obtained already prepared for Northern blot analysis (CLONTECH Catalog # 7759-1, 7760-1 and 7756-1.) The manufacturer produced the blots from approximately 2 µg of poly-A RNA per lane, run on a denaturing formaldehyde 1-2% agarose gel, transferred to a nylon membrane, and fixed by UV irradiation. The relative CH1-9a11-2 expression observed at the RNA level is shown in Table 5:

TABLE 5: Northern blot analysis	
Tissue	CH1-9a11-2 mRNA
heart	++
brain	+
placenta	++
lung	+/-
liver	+/-
skeletal muscle	+
kidney	+/-
pancreas	+++
spleen	+
thymus	+
prostate	++
testis	+++
ovary	++
small intestine	+
colon	+/-
peripheral blood	+/-
++++ Very high +++ High ++ Medium + Low +/- Very low	

Relatively elevated levels of expression were observed in heart, placenta, pancreas, prostate, testis and ovary. We believe that the level of expression in breast cancer cell lines is also relatively high (about ++++ on the scale), since the Northern analysis performed on these lines (described above) was conducted on *total* cellular RNA, of which polyadenylated RNA constitutes only about 5%. It is likely that the CH1-9a11-2 gene is involved in a biological process that is typical to the tissue types showing medium to high levels of expression, which may relate to increased tissue growth or metabolism.

Since the obtained sequence is shorter than the apparent size of mRNA observed in Northern analysis (Table 1), an additional polynucleotide segment is

believed to be present at the 5' end of the sequence shown in SEQ. ID NO:15. Further sequence data at the 5' end is deduced by obtaining additional cloned cDNA using standard techniques. Briefly, in one approach, mRNA from breast cancer cell lines MDA-453 and/or 600PE are cloned and screened using primers based on
5 sequence data from SEQ. ID NO:15. Two nested primers of about 20 nucleotides are prepared, the innermost about 150 base pairs from the 5' end, and the outermost about 170 base pairs from the 5' end. The outermost primer is used to synthesize a first cDNA strand complementary to the mRNA in the upstream direction. Second strand synthesis is performed using reagents in a CLONTECH Marathon™ cDNA
10 amplification kit according to manufacturer's directions. The double-stranded DNA is then ligated at the 5' end of the coding sequence with the double-stranded adaptor fragment provided in the kit. A first PCR amplification (about 30 cycles) is performed using the first adapter primer from the kit and the outermost RNA-specific primer, and a second amplification (about 30 cycles) is performed using the second
15 adapter primer and the innermost RNA-specific primer. In an alternative approach, a CLONTECH RACE-READY single-stranded cDNA from human placenta is PCR amplified using nested 5' anchor primers in combination with the outermost and innermost RNA-specific primers. Amplified DNA obtained using either approach is analyzed by gel electrophoresis, and cloned into plasmid vector pCRII. Clones are
20 screened, as necessary, using the 2.5 kilobase CH1-9a11-2 insert. Clones corresponding to full-length mRNA (4.5 kb or 5.5 kb; Table 1), or cDNA fragments overlapping at the 5' end are selected for sequencing. Compared with the 4.5 kb form, additional polynucleotide segments may be present in the 5.5 kb form within the encoding region, or in the 5' or 3' untranslated region.

25

Example 5: Chromosome 8 gene CH8-2a13-1

One of the cDNA obtained corresponded to a gene that mapped to Chromosome 8. Figure 2 shows the Southern blot analysis for the corresponding
30 gene in various DNA digests. Lane 1 (P12) is the control preparation of placental DNA; the rest show DNA obtained from human breast cancer cell lines. Panel A shows the pattern obtained using the 32P-labeled CH8-2a13-1 cDNA probe. Panel B

shows the pattern obtained with the same blot using the ³²P-labeled D2S6 probe as a loading control. The sizes of the restriction fragments are indicated on the right.

Figure 3 shows the Northern blot analysis for RNA overabundance. Lanes 1-3 show the level of expression in cultured normal epithelial cells. Lanes 4-19 show the level of expression in human breast cancer cell lines. Panel A shows the pattern obtained using the CH8-2a13-1 probe; panel B shows the pattern obtained with beta-actin cDNA, a loading control.

The results are summarized in Table 6. The scoring method is the same as for Example 4.

10

TABLE 6: Chromosome 8 Genes in Breast Cancer Cell Lines						
Source	CH8-2a13-1 Gene Duplication		CH8-2a13-1 RNA Overabundance		c-myc Gene Duplication	
Normal	-	1.00*	-	1.00**	-	1.00*
SKBR3	+	4.25	+	4.30	+	4.73
ZR-75-30	+	3.82	nd		+	2.24
BT474	+	1.53	+	1.72	+	1.76
MDA157	+	2.02	+	3.39	+	1.39
MCF7	+	1.84	+	4.92	+	3.10
CAMA-1	+	3.62	+	2.14	+	1.61
MDA361	+	2.00	+	1.74	nd	
MDA468	nd		+	4.50	nd	
T47D	+	1.41	+	1.58	-	1.02
MDA453	+	1.83	+	3.10	-	0.90
MDA134	+	1.30	+	3.70	-	0.88
MDA435	+	2.15	+	4.94	-	1.00
600PE	-	0.95	+	2.04	-	0.54
UACC812	+	1.25	+	2.40	-	0.74
MDA231	-	0.80	+	1.28	+	1.27
DU4475	-	0.85	-	0.88	-	0.50
BT468	-	0.37	-	0.70	-	0.23
BT20	-	0.95	-	0.82	-	
Incidence (%)	12/17 (71%)		14/17 (82%)		7/16 (44%)	

- + Gene duplication or RNA overabundance; - no duplication or overabundance; nd = not done.
 * Degree of gene duplication is reported relative to placental DNA preparations.
 ** Degree of RNA overabundance is reported relative to the highest level observed for several cultures of normal epithelial cells.

5

The gene corresponding to CH8-2a13-1 showed clear evidence of duplication in 12 out of 17 (71%) of the cells tested. RNA overabundance was observed in 14 out of 17 (82%). Thus, 11% of the cells had achieved RNA overabundance by a mechanism other than gene duplication.

10

Since the known oncogene c-myc is located on Chromosome 8, the Southern analysis was also conducted using a probe for c-myc. At least 2 of the breast cancer cells showing duplication of the gene corresponding to CH8-2a13-1 gene did not show duplication of c-myc. This indicates that the gene corresponding to CH8-2a13-1 is not part of the myc amplicon.

The sequence of 150 bases from the 5' end of the cDNA fragment is shown in Figure 22 (SEQ ID NO:3). There was no substantial homology to any known gene in GenBank. One of the three possible reading frames was found to be open, with the amino acid sequence shown in Figure 22 (SEQ ID NO:4).

The CH8-2a13-1 gene was further characterized by obtaining additional sequence information. A λ -GT10 cDNA library from the breast cancer cell line BT474 (Example 2) was screened using the initial cDNA insert, and clones with a 3.0 kb and a 4.0 kb insert were identified. The two identified clones were subcloned into plasmid vector pCRII. T7 and Sp6 primers for regions flanking the cDNA inserts were used as initial sequencing primers. Sequencing continued by walking along the region of interest by standard techniques, using sequencing primers based on data already obtained. The two inserts were found to overlap (Figure 6). Primers used are those designated 1-25 in Figure 10.

A third clone of about 600 bp (designated pCH8-600) overlapping on the 5' end (Figure 6) was obtained using CLONTECH Maration™ cDNA Amplification Kit.

Briefly, two DNA primers CH8a and CH8b (Figure 10) were synthesized. Polyadenylated RNA from breast cancer cell line BT474 was reverse transcribed using CH8b primer. After second strand synthesis, adaptor DNA provided in the kit was ligated to the double-stranded cDNA. The 5' end cDNA of CH8-2a13-1 was then amplified by PCR using primers CH8a and AP1 (provided in the kit). To increase the specificity of the PCR products, the first PCR products were PCR reamplified using nested primers CH8a and AP2 (provided in the kit). The PCR products were cloned into pCRII vector (Invitrogen) and screened with CH8-2a13-1 probe.

By sequencing relevant portions of the three clones, a nucleic acid sequence of 3982 base pairs between the 5' end and the poly-A tail of CH8-2a13-1 was determined. The DNA sequence is shown in Figure 11 (SEQ. ID NO:18). Bases 1-

152 are believed to be a 5' untranslated region. The longest open reading frame is in frame 3 from base 153 to 3911, and codes for 1252 amino acids before the stop codon. The corresponding amino acid sequence of this frame is shown in the upper panel of Figure 12 (SEQ. ID NO:19). The sequence predicted for the translated
5 protein is shown in the lower panel of Figure 12(SEQ. ID NO:20).

A GENINFO® BLAST search of nucleotide and peptide sequence databases was performed through the National Center for Biotechnology Information on March 26, 1996. The sequences were found to be about 99% identical at the nucleotide and amino acid level with bases 343-4103 of KIAA0196 protein (N. Nomura et al., in
10 press; sequence submitted to the DDBJ/EMBL/GenBank databases on March 4, 1996). The KIAA0196 was one of 200 different cDNA cloned at random from an immature male human myeloblast cell line. KIAA0196 has no known biological function, and is described by Nomura et al. as being ubiquitously expressed.

A fourth clone of about 600 bp overlapping pCH8-600 at the 5' end has also
15 been obtained. Briefly, a DNA primer was synthesized corresponding to about the first 20 nucleotides at the 5' of the predicted cDNA sequence, and used along with a primer based on the pCH8-600 sequence to reverse-transcribe RNA from breast cancer cell line BT474. The product was cloned into pCRII vector (Invitrogen) and screened with a CH8-2a13-1 probe. The new clone is sequenced along both strands
20 to obtain additional 5' untranslated sequence data for the cDNA. The predicted compiled cDNA nucleotide sequence of CH8-2a13-1 cDNA is shown in Figure 13 (SEQ. ID NO:21). The corresponding amino acid sequence of this frame is shown in Figure 14 (SEQ. ID NO:22). A polynucleotide comprising the compiled sequence is assembled by joining the insert of this fourth clone to pCH8-4k within the shared
25 region. Briefly, CH8-4k is cut with *Xba*I and *Not*I. The fourth clone is cut with *Bam*HI and *Xba*I. The ligated polynucleotide is then inserted into pCRII cut with *Bam*HI and *Not*I.

A CH8-2a13-1 cloned insert has been used to probe the level of relative expression in polyadenylated RNA from a panel of tissue sources obtained from
30 CLONTECH, as in Example 4. The relative CH8-2a13-12 expression observed at the mRNA level is shown in Table 7:

TABLE 7: Northern blot analysis	
Tissue	CH1-9a11-2 mRNA
heart	++
brain	+
placenta	+
lung	+
liver	+/-
skeletal muscle	+/-
kidney	+/-
pancreas	+/-
spleen	+
thymus	+
prostate	+
testis	++
ovary	+
small intestine	+
colon	+
peripheral blood	+/-
++++ Very high +++ High ++ Medium + Low +/- Very low	

Relative levels of expression observed were as follows: Low levels of expression were observed in adult peripheral blood leukocytes (PBL), brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Medium levels of expression were observed in adult heart, spleen, thymus, prostate, testis, ovary, small intestine, and colon. High levels of expression were observed in four fetal tissues tested: brain, lung, liver and kidney. We believe that the level of expression in breast cancer cell lines is relatively high (about ++++ on the scale), since the Northern analysis performed on these lines was conducted on *total* cellular RNA. It is likely that the CH8-2a13-1 gene is involved in a biological process that is typical to the tissue types

showing medium to high levels of expression, which may relate to increased tissue growth or metabolism.

Example 6: Chromosome 13 gene CH13-2a12-1

5

One of the cDNA obtained corresponded to a gene that mapped to Chromosome 13. Figure 4 shows the Southern blot analysis for the corresponding gene in various DNA digests. Lanes 1 and 2 are control preparations of placental DNA; the rest show DNA obtained from human breast cancer cell lines. Panel A shows the pattern obtained using the CH13-2a12-1 cDNA probe; panel B shows the pattern using D2S6 probe as a loading control. The sizes of the restriction fragments are indicated on the right.

Figure 5 shows the Northern blot analysis for RNA overabundance of the CH13-2a12-1 gene. Lanes 1-3 show the level of expression in cultured normal epithelial cells. Lanes 4-19 show the level of expression in human breast cancer cell lines. Panel A shows the pattern obtained using the CH13-2a12-1 probe; panel B shows the pattern obtained with beta-actin cDNA, a loading control. The apparent size of the mRNA varied depending upon conditions of electrophoresis. Full-length mRNA is believed to occur at sizes of about 3.2 and 3.5 kb.

The results of the RNA abundance comparison are summarized in Table 8. The scoring method is the same as for Example 4.

TABLE 8: Chromosome 13 Gene in Breast Cancer Cell Lines		
Source	CH13-2a12-1 Gene duplication	CH13-2a12-1 RNA Overabundance
Normal	- 1.00*	- 1.00**
600PE	+ 2.18	+ 5.57
BT474	+ 1.60	+ 3.20
SKBR3	+ 1.58	+ 4.25
MDA157	+ 2.21	+ 3.76
CAMA-1	+ 1.41	+ 1.99
MDA231	+ 1.65	+ 2.09
T47D	+ 1.23	+ 1.20
MDA468	nd	+ 6.90
MDA361	nd	+ 2.59
MDA435	- 0.59	+ 3.41
MDA134	- 0.53	+ 2.59
DU4475	- 0.75	+ 1.79
MDA453	- 0.89	+ 1.97
BT20	- 0.37	- 1.04
MCF7	- 0.29	- 1.03
UACC812	- 0.30	- 0.39
BT468	- 0.47	nd
ZR-75-30	- 0.70	nd
Incidence (%)	7/16 (44%)	13/16 (81%)

+ Gene duplication or RNA overabundance; - no duplication or overabundance; nd = not done

* Degree of gene duplication is reported relative to placental DNA preparations.

** Degree of RNA overabundance is reported relative to the highest level observed for several cultures of normal epithelial cells.

5

The gene corresponding to CH13-2a12-1 was duplicated in 7 out of 16 (44%) of the cells tested. Three of the positive cell lines (600PE, BT474, and MDA435) had been studied previously by comparative genomic hybridization, but had not

shown amplified chromatin in the region where CH13-2A12-1 has been mapped in these studies.

RNA overabundance was observed in 13 out of 16 (81%) of the cell lines tested. Thus, 37% of the cells had achieved RNA overabundance by a mechanism other than gene duplication. We have also obtained cells from primary breast tumors, and analyzed them for duplication of the chromosome 13 gene. Ten of the 82 tumors analyzed (12%) were positive, confirming that duplication of this gene is not an artifact of in vitro culture.

The sequence of 107 bases from the 5' end of the 1.5 kb cDNA fragment is shown in Figure 22 (SEQ ID NO:5). There was no substantial homology to any known gene in GenBank. One of the three possible reading frames was found to be open, with the predicted amino acid sequence shown in Figure 22 (SEQ ID NO:6).

The CH13-2a12-1 gene was further characterized by obtaining additional sequence information. A λ -GT10 cDNA library from the breast cancer cell line BT474 (Example 2) was screened using the initial cDNA insert, and clones with a 3.5 kilobase and a 1.6 kilobase insert were identified. The two identified clones were subcloned into plasmid vector pCRII. T7 and Sp6 primers for regions flanking the cDNA inserts were used as initial sequencing primers. Sequencing continued by walking along the region of interest by standard techniques, using sequencing primers based on data already obtained. The two inserts were found to overlap (Figure 6). Primers used during sequencing are shown in Figure 15.

By sequencing relevant portions of the 3.5 and 1.6 kb clones, a nucleic acid sequence of 3339 base pairs between the 5' end and the poly-A tail of CH13-2a12-1 was determined. The DNA sequence is shown in Figure 16 (SEQ. ID NO:23). Bases 1-520 are believed to be a 5' untranslated region. The longest open reading frame is in frame 2 from base 521 to 1838, and codes for 611 amino acids before the stop codon. The corresponding amino acid sequence of this frame is shown in the upper panel of Figure 17 (SEQ. ID NO:24). The sequence predicted for the translated protein is shown in the lower panel of Figure 17 (SEQ. ID NO:25). Bases 1838 to 3339 of the nucleotide sequence are believed to be a 3' untranslated region, which is present in the 3.5 kb insert. The 3.5 kb insert appears to be a splice variant (Figure 6), in which the 3' untranslated region consists of bases 1838-2797 in the sequence.

A GENINFO® BLAST search of nucleotide and peptide sequence databases was performed through the National Center for Biotechnology Information on March 26, 1996. Short segments of homology with other reported human sequences were found at the nucleotide level (<500 base pairs), but none with any ascribed function in the respective identifier. At the amino acid level, the sequence was found to share 33% identities and 54% positives with 228 residues of the *lin 19* protein of *Caenorhabditis elegans*. This protein has been implicated in regulating the cell cycle of *C. elegans* (ET Kipreos, W He & EM Hedgecock). The CH13-2a12-1 gene is suspected of a role in controlling cell proliferation. "Controlling cell proliferation" in this context means that an abnormally high or low level of gene expression at the RNA or protein level results in a higher or lower rate of cell proliferation, or vice versa, compared with cells with an otherwise similar phenotype. There is also a low-level homology between CH13-2a12-1 and VACM-1, a vasopressin-activated, calcium-mobilizing receptor from rabbit kidney medulla (Burnatowska-Hledin et al). VACM-1 has a transmembrane sequence, whereas none has been detected in CH13-2a12-1. Nevertheless, it is possible that the CH13-2a12-1 protein product has a Ca^{++} binding or Ca^{++} mobilizing function.

A CH13-2a12-1 cloned insert has been used to probe the level of relative expression in polyadenylated RNA from a panel of tissue sources obtained from CLONTECH, as in Example 4. The relative CH13-2a12-1 expression observed at the mRNA level is shown in Table 9:

TABLE 9: Northern blot analysis	
Tissue	CH13-2a12-1 mRNA
heart	++++
brain	+
placenta	++
lung	+
liver	++
skeletal muscle	++++
kidney	+
pancreas	++
spleen	++
thymus	++
prostate	++
testis	+++
ovary	++
small intestine	++
colon	+
peripheral blood	+
++++ Very high +++ High ++ Medium + Low +/- Very low	

Relatively elevated levels of expression were observed in heart, skeletal muscle and testis. We believe that the level of expression in breast cancer cell lines is relatively high (about ++++ on the scale), since the Northern analysis performed on these lines was conducted on *total* cellular RNA. It is likely that the CH13-2a12-1 gene is involved in a biological process that is typical to the tissue types showing medium to high levels of expression, which may relate to increased tissue growth or metabolism.

Example 7: Chromosome 14 gene CH14-2a16-1

One of the cDNA obtained corresponded to a gene that mapped to Chromosome 14. Results of the analysis are summarized in Table 10. The scoring
5 method is the same as for Example 4.

TABLE 10: Chromosome 14 Gene in Breast Cancer Cell Lines		
Source	CH14-2a16-1 Gene duplication	CH14-2a16-1 RNA Overabundance
Normal	- 1.00*	- 1.00**
BT474	+ 2.89	+ 2.57
MCF7	+ 1.35	+ 1.88
SKBR3	+ 2.58	+ 2.19
T47D	+ 2.28	nd
MDA157	+ 1.52	+ 2.52
UACC812	+ 2.23	nd
MDA361	- 0.97	+ 1.43
MDA453	+ 1.58	+ 5.92
BT20	-	- 1.07
600PE	- 0.94	+ 2.00
MDA231	+ 1.66	+ 2.19
CAMA-1	- 0.92	- 0.71
DU4475	- 0.87	+ 1.33
BT468	- 0.46	nd
MDA134	- 0.77	+ 7.17
Incidence (%)	8/15 (53%)	10/12 (83%)

+ Gene duplication or overabundance; - no duplication or overabundance; nd = not done

* Degree of gene duplication is reported relative to placental DNA preparations.

** Degree of RNA overabundance is reported relative to the highest level observed for several cultures of normal epithelial cells.

5

The gene corresponding to CH14-2a16-1 was duplicated in 8 out of 15 (53%) of the cells tested. The sequence of 114 bases from the 5' end of the cDNA fragment is shown in Figure 22 (SEQ ID NO:7). There was no substantial homology to any known gene in GenBank. One of the three possible reading frames was found to be open, with the predicted amino acid sequence shown in Figure 22 (SEQ ID NO:8).

10

The CH14-2a16-1 gene was further characterized by obtaining additional sequence information. A λ -GT10 cDNA library from the breast cancer cell line BT474 (Example 2) was screened using the initial cDNA insert, and two clones were identified: one with a 1.6 kb insert, and the other with a 2.5 kb insert. The identified clones were subcloned into plasmid vector pCRII. The 1.6 kb insert was sequenced by using T7 and Sp6 primers for regions flanking the cDNA inserts as initial sequencing primers. Sequencing continued by walking along the region of interest by standard techniques, using sequencing primers based on data already obtained. Primers used are those designated 1-11 in Figure 18.

A third clone (designated pCH14-800) overlapping on the 5' end (Figure 6) was obtained using CLONTECH Maration™ cDNA Amplification Kit. Briefly, DNA primers CH14a, CH14b, CH14c and CH14d (Figure 18) were prepared. Polyadenylated RNA from breast cancer cell line MDA453 was reverse transcribed using 14b primer. After second strand synthesis, adaptor DNA provided in the kit was ligated to the double-stranded cDNA. The 5' end cDNA of CH14-2a16-1 was then amplified by PCR using primers CH14b (or CH14c) and AP1 (provided in the kit). To increase the specificity of the PCR products, the first PCR products were PCR reamplified using nested primers CH14a (or CH14d) and AP2 (provided in the kit). The PCR products were cloned into pCRII vector (Invitrogen) and screened with CH14-2a16-1 probe.

By sequencing pCH14-1.6 and pCH14-800, a nucleic acid sequence of 2021 base pairs between the 5' end and the poly-A tail of CH14-2a16-1 has been determined. The DNA sequence is shown in Figure 19 (SEQ. ID NO:26). The longest open reading frame is in frame 1 from base 1 to 792, and codes for 263 amino acids before the stop codon. The corresponding amino acid sequence of this frame is shown in the upper panel of Figure 20 (SEQ. ID NO:27). The partial sequence predicted for the translated protein is shown in the lower panel of Figure 20 (SEQ. ID NO:28). The 2.1 kb clone has not been sequenced, but is believed to consist about the same region of the CH14-2a16-1 cDNA as pCH14-1.6 and pCH14-800 combined.

A GENINFO® BLAST search of nucleotide and peptide sequence databases was performed through the National Center for Biotechnology Information on March 26, 1996. Short segments of homology with other reported human sequences were

found at the nucleotide level (<500 base pairs), but none with any ascribed function in the respective identifier. At the amino acid level, the sequence was found to share homologies within the first 106 residues with an RNA binding protein from *Saccharomyces cerevisiae* with the designation NAB2. NAB2 is one of the major proteins associated with nuclear polyadenylated RNA in yeast cells, as detected by UV light-induced cross-linking and immunofluorescence. NAB2 is strongly and specifically associated with nuclear poly(A)+ RNA in vivo. Gene knock-out experiments have shown that this protein is essential to yeast cell survival (Anderson et al.). Accordingly, the protein encoded by CH14-2a16-1 is suspected of having DNA or RNA binding activity.

A fourth clone (pCH14-1.3) has been obtained that overlaps the pCH14-800 clone at the 5' end (Figure 6). The method of isolation was similar to that for pCH14-800, using primers based on the pCH14-800 sequence. Partial sequence data for pCH14-1.3 has been obtained by one-directional sequencing from the 5' and 3' ends of the pCH14-1.3 clone. Figure 21 shows the nucleotide sequence of the sequence of the 5' end (SEQ. ID NO:29) and the amino acid translation of the likely open reading frame (SEQ. ID NO:30); the nucleotide sequence of the 3' end (SEQ. ID NO:31) and the likely open reading frame (SEQ. ID NO:32). This data is confirmed and additional sequence between SEQ. ID NOS.29 and 31 is obtained by fully sequencing both strands of pCH14-1.3. Once compiled, the sequence data from pCH14-1.3, pCH14-800 and pCH14-1.6 may be shorter than the apparent size of mRNA observed in Northern analysis (Table 1). If necessary, further sequence data at the 5' end is deduced by obtaining additional cloned cDNA according to approaches described in this Example or Example 4.

A CH14-2a16-1 cloned insert has been used to probe the level of relative expression in polyadenylated RNA from a panel of tissue sources obtained from CLONTECH, as in Example 4. The relative CH14-2a16-1 expression observed at the mRNA level is shown in Table 11:

TABLE 11: Northern blot analysis	
Tissue	CH14-2a16-1 mRNA
heart	+
brain	+
placenta	+
lung	+
liver	+
skeletal muscle	+
kidney	+/-
pancreas	+
spleen	+
thymus	+
prostate	+
testis	++++
ovary	+
small intestine	+
colon	+
peripheral blood	+/
++++ Very high +++ High ++ Medium + Low +/- Very low	

CH14-2a16-1 mRNA was particularly high in testis. We believe that the level of expression in breast cancer cell lines is also quite high, since the Northern analysis performed on these lines was conducted on *total* cellular RNA. It is likely that the CH14-2a16-1 gene is involved in a biological process that is typical to the tissue types showing medium to high levels of expression, which may relate to increased tissue growth or metabolism.

Five motifs corresponding to a zinc finger protein have been found in the CH14-2a16-1 nucleotide sequence. Further zinc finger motifs may be present in CH14-2a16-1 in the upstream direction. Zinc finger motifs are present, for example,

in RNA polymerases I, II, and III from *S. cerevisiae*, and are related to the zinc knuckle family of RNA/ssDNA-binding proteins found in the HIV nucleocapsid protein. The actual sequence observed in each of the five zinc finger motifs of CH14-2a16-1 is:

5

Cys-(Xaa)₅-Cys-(Xaa)₄-Cys-(Xaa)₃-His or (SEQ. ID NO:33)

Cys-(Xaa)₅-Cys-(Xaa)₅-Cys-(Xaa)₃-His (SEQ. ID NO:34)

- 10 which is indicated in Figure 20 by underlining. This is identical to the 7 zinc finger motifs of NAB2, which make up an RNA/ssDNA binding region (Anderson et al.). Accordingly, the CH14-2a16-1 gene product is suspected of having DNA or RNA binding activity, and may be specific for polyadenylated RNA. It may very well play a role in the regulation of gene replication, transcription, the processing of hnRNA
- 15 into mature mRNA, the export of mRNA from the nucleus to the cytoplasm, or translation into protein. This role in turn may be closely implicated in cell growth or proliferation, particularly as manifest in tumor cells.

REFERENCES

Articles

- 5 Adnane J. et al. (1991), "BEK and FLG, two receptors to members of the FGF family, are amplified in subsets of human breast cancers", *Oncogene* 6:659-663.
- Alitalo K. et al. (1986), "Oncogene amplification in tumor cells", *Adv. Cancer Res.* 47:235-281.
- 10 Altschul et al. (1986), *Bull. Math. Bio.* 48:603-616.
- Anderson J.T., Wilson S.M., Datar K.V. and Swanson M.S. (1993). "NAB2: a yeast nuclear polyadenylated RNA-binding protein essential for cell viability", *Mol.*
- 15 *Cell. Biol.* 13:2730-2741.
- Bauer D. et al. (1993), "Identification of differentially expressed mRNA species by an improved display technique (DDRT-PCR)", *Nucl. Acids Res.* 21:4272-4280.
- 20 Beardsley T. (1994), "Crabshoot: manufacturers gamble on cancer vaccines — again", *Scientific American*, Sept:102.
- Berns E.M. et al. (1992), "Sporadic amplification of the insulin-like growth factor 1 receptor gene in human breast tumors", *Cancer Res.* 52:1036-1039.
- 25 Bishop J.M. (1991), "Molecular themes in oncogenesis", *Cell* 64:235-248.
- Blast R.C. Jr. (1993), "Perspectives on the future of cancer markers", *Clin Chem.* 31:2444-2451.
- 30 Brison O. (1993), "Gene amplification and tumor progression", *Biochim. Biophys. Acta* 1155:25-41.

Burnatowska-Hledin, M.A. et al (1995). "Expression cloning of an AVP-activated, calcium mobilizing receptor from rabbit kidney medulla", Am. J. Physiol. 268:F1198-F1210.

5

Chen Z. et al. (1995). "Differential expression of human tissue factor in normal mammary epithelial cells and in carcinomas". Molec. Med. 1:153-160.

Culver K.W. et al. (1994), "Gene therapy for cancer," Trends Genet. 10:174-178.

10

Dutrillaux B. et al. (1990), "Characterization of chromosomal anomalies in human breast cancer", Cancer Genet Cytogenet 49:203-217.

Henikoff et al. (1992), Proc. Natl. Acad. Sci. USA 89:10915-10919.

15

Kallioniemi A. et al. (1992), "Comparative genomic hybridization for molecular cytogenetic analysis of solid tumors", Science 258:818-821.

20

Kallioniemi A. et al. (1994), "Detection and mapping of amplified DNA sequences in breast cancer by comparative genomic hybridization", Proc. Natl. Acad. Sci. USA 91:2156-2160.

25

Kocher O. et al. (1995), "Identification of a novel gene, selectively up-regulated in human carcinomas, using the differential display technique", Clin. Cancer Res. 1:1209-1215.

Lippman M.E. (1993), "The development of biological therapies for breast cancer", Science 259:631-632.

30

Liang P. et al. (1992), "Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction", Science 257:967-971.

- Liang P. et al. (1993), "Distribution and cloning of eukaryotic mRNAs by means of differential display: refinements and optimization", Nucl. Acids Res. 21:3269-3275.
- MacLean G.D. et al. (1992), "The immune system, cancer antigens and immunotherapy", Contemp. Oncol. Aug/Sept. pp 1-7.
- Muss H.B. et al. (1994), "c-*erbB*-2 expression and response to adjuvant therapy in women with node-positive early breast cancer", New Engl. J. Med. 330:1260-1266.
- Morgan R.A. et al. (1993), "Human gene therapy," Annu. Rev. Biochem. 62:191-217.
- Roth J.A. (1994), "Modulation of oncogene and tumor-suppressor gene expression: a novel strategy for cancer prevention and treatment", Ann. Surg. Oncol. 1:79-86.
- Saint-Ruf C. et al. (1990), "Proto-oncogene amplification and homogeneously staining regions in human breast carcinomas", Genes Chromosomes Cancer 2:18-26.
- Slamon D.J. et al. (1987), "Human breast cancer: correlation of relapse and survival with amplification of the HER-2/*neu* oncogene", Science 235:178-182.
- Schwab M. et al. (1990), "Amplification of cellular oncogenes: a predictor of clinical outcome of human cancer", Genes Chromosomes Cancer 1:181-193.
- Trentmann S.M. et al. (1995), "Alternatives to ³⁵S as a label for the differential display of eukaryotic messenger RNA", Science 267:1186-1187.
- Unsigned (1994), "Synthetic vaccine stabilizes advanced cancer, prolongs survival", Oncol. News Int. 3:1.

Watson M.A. et al. (1995). "Mammaglobulin, a mammary-specific member of the uteroglobuline gene family, is overexpressed in human breast cancer", *Cancer Res.* 56:860-885.

- 5 Zafrani B. et al. (1992), "Cytogenetic study of breast cancer: Clinicopathologic significance of homogenously staining regions in 84 patients", *Hum. Pathol* 23:542-547.

U.S. Patents

10

US 4,444,887	4/1984	Hoffman M.K.	(mAB method)
US 4,472,500	9/1984	Milstein C. et al.	(mAB cell)
US 4,491,632	1/1985	Wands J.R. et al.	(HBV mAb)
US 4,683,195	7/1987	Mullis K.B.	(PCR)
15 US 4,683,202	7/1987	Mullis K.B. et al.	(PCR)
US 4,968,603	11/1990	Slamon D.J. et al.	(<i>erbB2</i> in
diagnosis)			
US 5,124,246	6/1992	Urdea M.S. et al.	(branched DNA)
US 5,399,346	21/1995	Anderson W.F. et al.	(gene therapy)

20

Other Patent Publications

WO 93/08701	5/1993	Goldstein, J.A. et al.	(<i>c-myc</i>)
WO 94/00136	1/1994	Kasprzyk P.G. et al.	(anti- <i>erb</i> in
25 therapy)			
WO 94/00601	1/1994	Levine A.J. et al.	(mAB in
diagnosis)			
WO 94/17414	8/1994	Keyomarsi K. et al.	(detection)
WO 94/28127	12/1994	Sikora K. et al.	(<i>erb</i> promoter)

30

SEQUENCE LISTINGS:

SEQ. ID NO	Designation	Description	Type	
1	CH1-9a11-2	152 bp sequence (fragment)	dsDNA	Figure 22
2		translation	amino acid	Figure 22
3	CH8-2a13-1	163 bp sequence (fragment)	dsDNA	Figure 22
4		translation	amino acid	Figure 22
5	CH13-2a12-1	107 bp sequence (fragment)	dsDNA	Figure 22
6		translation	amino acid	Figure 22
7	CH14-2a16-1	114 bp sequence (fragment)	dsDNA	Figure 22
8		translation	amino acid	Figure 22
9 to 14	Primers		ssDNA	Table 2
15	CH1-9a11-2	3.5 kb nucleotide sequence	dsDNA	Figure 8
16		translation	amino acid	Figure 9
17		protein	amino acid	Figure 9
18	CH8-2a13-1	4.0 kb nucleotide sequence	dsDNA	Figure 11
19		translation	amino acid	Figure 12
20		protein	amino acid	Figure 12
21		4.1 kb sequence (predicted)	dsDNA	Figure 13
22		translation	amino acid	Figure 14
23	CH13-2a12-1	3.3 kb nucleotide sequence	dsDNA	Figure 16
24		translation	amino acid	Figure 17
25		protein	amino acid	Figure 17
26	CH14-2a16-1	2.0 kb nucleotide sequence	dsDNA	Figure 19
27		translation	amino acid	Figure 20
28		protein	amino acid	Figure 20
29		0.6 kb nucleotide sequence	ssDNA	Figure 21
30		translation	amino acid	Figure 21
31		0.3 kb nucleotide sequence	ssDNA	Figure 21
32		translation	amino acid	Figure 21
33 & 34	Motif	Zinc-finger binding domain	dsDNA	text
35-38	Primers		ssDNA	text
39 & up	Primers		ssDNA	Figures 7, 10, 15, 18

CLAIMS

We claim:

- 5 1. An isolated polynucleotide comprising a linear sequence of at least 10 nucleotides identical to a linear sequence contained in a polynucleotide selected from the group consisting of CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1.
- 10 2. An isolated polynucleotide comprising a linear sequence of at least 15 nucleotides identical to a linear sequence contained in a polynucleotide selected from the group consisting of CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1.
- 15 3. The isolated polynucleotide of claim 2, wherein said linear sequence is contained in a duplicated gene or overabundant RNA in cancerous cells.
- 20 4. The isolated polynucleotide of claim 2, comprising a linear sequence of at least 40 nucleotides essentially identical to a sequence contained in the polynucleotide selected from said group.
- 25 5. The isolated polynucleotide of claim 2, wherein the polynucleotide selected from said group is a CH13-2a12-1 polynucleotide, and is contained in an encoding region for a protein or RNA molecule that controls cell proliferation.
- 30 6. The isolated polynucleotide of claim 2, wherein the polynucleotide selected from said group is a CH14-2a16-1 polynucleotide, and is contained in an encoding region for a protein with DNA or RNA binding activity.
7. The isolated polynucleotide of claim 2, present in a recombinant plasmid deposited under Accession No. _____.

8. The isolated polynucleotide of claim 2, present in a recombinant phage deposited under Accession No. ____.
- 5 9. The isolated polynucleotide of claim 2, present in the λ BCBT474 cDNA library deposited under Accession No. ____.
- 10 10. An isolated polynucleotide comprising a linear sequence of polynucleotides essentially identical to a sequence selected from the group consisting of SEQ. ID NO:15, SEQ. ID NO: 18, SEQ. ID NO:21, SEQ. ID NO:23, SEQ. ID NO:26, SEQ. ID NO:29, and SEQ. ID NO:31.
- 15 11. An isolated polypeptide comprising a linear sequence of at least 5 amino acid residues identical to a sequence encoded by a polynucleotide selected from the group consisting of CH1-9a11-2, CH8-2a13-1, CH13-2a12-1, and CH14-2a16-1.
- 20 12. The isolated polypeptide of claim 11, wherein said linear sequence is encoded in a duplicated gene or overabundant RNA in cancerous cells.
- 20 13. The isolated polypeptide of claim 11, which is overexpressed in cancerous cells.
- 25 14. The isolated polypeptide of claim 11, comprising a linear sequence of at least 15 amino acids essentially identical to a sequence encoded by said polynucleotide.
- 30 15. The isolated polypeptide of claim 11, wherein the polynucleotide selected from said group is a CH1-9a11-2 polynucleotide, and the polypeptide is a transmembrane polypeptide.

16. The isolated polypeptide of claim 11, wherein the polynucleotide selected from said group is a CH13-2a12-1 polynucleotide, and the polypeptide is contained in a protein that controls cell proliferation.
- 5 17. The isolated polypeptide of claim 11, wherein the polynucleotide selected from said group is a CH14-2a16-1 polynucleotide, and the polypeptide is contained in a protein with DNA or RNA binding activity.
- 10 18. An isolated polypeptide comprising a linear sequence of amino acids essentially identical to a sequence selected from the group consisting of SEQ. ID NO:17, SEQ. ID NO:20, SEQ. ID NO:25 and SEQ. ID NO: 28.
- 15 19. A monoclonal or isolated polyclonal antibody specific for the polypeptide of claim 18.
- 20 20. A method of using a polynucleotide of claim 4 for detecting gene duplication in cancerous cells, comprising the steps of:
- 20 a) reacting DNA contained in a clinical sample with a reagent comprising the polynucleotide of claim 4, said clinical sample having been obtained from an individual suspected of having cancerous cells; and
- 25 21. A method of using a polynucleotide of claim 4 for detecting overabundance of RNA in cancerous cells, comprising the steps of:
- 30 a) reacting RNA contained in a clinical sample with a reagent comprising the polynucleotide of claim 4, said clinical sample having been obtained from an individual suspected of having cancerous cells; and
- b) comparing the amount of any complexes formed between the reagent and the RNA in the clinical sample with the amount of any complexes formed between the reagent and RNA in a control sample.

22. A diagnostic kit for detecting gene duplication or RNA overabundance in cells contained in an individual as manifest in a clinical sample, comprising a reagent and a buffer in suitable packaging, wherein the reagent comprises the polynucleotide of claim 4.
23. A method for using the antibody of claim 19 for detecting altered protein expression in cancerous cells, comprising the steps of:
- a) reacting a polypeptide contained in a clinical sample with a reagent comprising the antibody of claim 19, said clinical sample having been obtained from an individual suspected of having cancerous cells; and
 - b) comparing the amount of any complexes formed between the reagent and the polypeptide in the clinical sample with the amount of any complexes formed between the reagent and a polypeptide in a control sample.
24. A diagnostic kit for detecting a polypeptide present in a clinical sample, comprising a reagent and a buffer in suitable packaging, wherein the reagent comprises the antibody of claim 19.
25. A host cell genetically altered by the polynucleotide of claim 4.
26. A method of using the cell of claim 19 for screening a pharmaceutical candidate, comprising the steps of:
- a) separating progeny of the cell of claim 19 into a first group and a second group;
 - b) treating the first group of cells with the pharmaceutical candidate;
 - c) not treating the second group of cells with the pharmaceutical candidate; and
 - d) comparing the phenotype of the treated cells with that of the untreated cells.

27. A pharmaceutical preparation for use in cancer therapy, comprising the polynucleotide of claim 4, said preparation being capable of reducing the pathology of cancerous cells.
- 5 28. A method for treating an individual bearing cancerous cells, comprising administering the pharmaceutical preparation of claim 27.
29. A pharmaceutical preparation for use in cancer therapy, comprising the antibody of claim 19, said preparation being capable of reducing the pathology
10 of cancerous cells.
30. A method for treating an individual bearing cancerous cells, comprising administering the pharmaceutical preparation of claim 29.
- 15 31. A pharmaceutical preparation comprising the polypeptide of claim 11 in an immunogenic form, and a pharmaceutically compatible excipient.
32. A method for treatment of cancer, comprising administration of the pharmaceutical preparation of claim 31.


Figure 1


Figure 2


Figure 3

Figure 4





Figure 5





Figure 6**CH1-9a11-2****CH8-2a13-1****CH13-2a12-1****CH14-2a16-1**

Figure 7

+ strand (sense)		sequence (5'-->3')
	1st base	
1. pch1-t7-1f	1123	CGG GAG GTT TCA GAT CGA C
2. pch1-t7-2f	1437	GCG CTG CAA GTA CAA AAT TG
3. pch1-t7-3f	1729	TCT AAA GTC CAA GAC CAA GG
4. pch1-t7-4f	1987	CAG AAA TTA TGG TTT CTA CC
5. pch1-t7-5f	2266	CaG GAA GAG GAG GGA TAA C (T)
6. pch1-sp6-3fb	2684	AAA CAT ACA CAA TAA ACA C
7. pch1-sp6-2rb	2966	TTG GCA GCG ACT GTA TTT G
8. pch1-sp6-1rb	3283	CCT GAT TTT ATA GAA GCC CC
- strand (antisense)		
9. pch1-sp6-1f	3302	GGG GCT TCT ATA AAA TCA GG
10. pch1-sp6-2f	2987	ATT CAA ATA CAG TTG CTG C
11. pch1-sp6-3f	2705	TTA GTG TTT ATT GTG TAT G
12. pch1-sp6-4f	2458	AGT GTT CAT TTC CAG TGA G
13. pch1-sp6-5f	2066	CTT TGT TCT TGG ACT TTA G
14. pch1-t7-3fb	1748	CCT TGG TCT TGG ACT TTA G
15. pch1-t7-2rb	1445	AAT TTT GTA CTT GCA GCG C
16. pch1-t7-1rb	1141	GTC GAT CTG AAA CCT CCC G
17. CH1a	1063	GTG CCT GTA GCA ACT GGA TGG C
18. CH1b	1079	GTC ATG TTG GTC AGC TGT GCC

Figure 8(A)

1	GAATACATAT	ATAAATGGTG	TTCAGTTAGA	GTTGCTCTTT	ATCGGCAGCG
51	CAGCCGAAC	GCTTTGAGTA	AAGGAAAAGA	TTATCTTG TG	TTAGCTCAAC
101	CACCCCTACT	ACTTCCTGCG	GAATCAGTAG	ATGTTTCAGT	ATTGCAACCT
151	CTGAGTGGAG	AATTGGAAAA	TACGAATATA	GAAAGGGAAG	CTGAAACTGT
201	TGTTCTGGGT	GATTTAAGTA	GTAGTATGCA	CCAGGATGAC	TTGGTGAATC
251	ACACTGTAGA	TGCAGTTGAA	CTTGAACCAA	GCCATTCTCA	AACTCTTTCT
301	CAGTCTCTTC	TTTTAGATAT	TACCCAGAA	ATCAATCCCT	TGCCTAAAAT
351	AGAAGTATCT	GAGTCTGTTG	AATATGAGGC	AGGACATATA	CCATCACCAG
401	TGATTCCTCCA	AGAGAGTTCT	GTTGAGATCG	ATAATGAAAC	AGAACAAAAG
451	TCTGAGAGCT	TTAGTTCTAT	AGAGAAACCA	TCTATTACCT	ATGAAACAAA
501	TAAAGTTAAT	GAGTTAATGG	ATAATATTAT	AAAAGAAGAT	ATGAACTCCA
551	TGCAAAATTTT	CACAAAGCTG	TCTGAAACAA	TAGTGCCACC	AATAAATACA
601	GCCACTGTAC	CCGACAATGA	AGATGGGGAA	GCCAAAATGA	ATATAGCTGA
651	CACAGCAAAG	CAAACTTTGA	TTTCTGTTGT	GGATTCTTCT	TCATTACCTG
701	AAGTAAAAGA	AGAAGAACAG	TCTCCAGAAG	ATGCCCTTTT	GAGAGGGTTA
751	CAGAGGACAG	CTACAGATTT	TTATGCTGAA	TTGCAAAATT	CTACAGATCT
801	AGGATATGCT	AATGGAAATC	TTGTACATGG	ATCAAACCAA	AAGGAGTCAG
851	TATTTATGAG	ACTTAATAAT	CGTATTAAAG	CCTTAGAAGT	TAACATGTCT
901	CTCAGTGGTC	GCTATCTGGA	GGAGCTTAGC	CAAAGGTACC	GAAAACAAAT
951	GGAAGAAATG	CAAAGGCTT	TCAACAAAAC	AATCGTGAAA	CTTCAGAATA
1001	CTTCAAGAAT	AGCAGAGGAG	CAGGATCAGC	GGCAAACTGA	AGCCATCCAG
1051	TTGCTACAGG	CACAGCTGAC	CAACATGACA	CAGCTTGTTT	CAAATTTATC
1101	AGCAACAGTA	GCAGAAATGA	AACGGGAGGT	TTCAGATCGA	CAAAGCTATC
1151	TTGTCAATATC	TTTGGTTCTT	TGTGTTGTCT	TGGGACTGAT	GCTTTGTATG
1201	CAGCGTTGTC	GAAATACTTC	TCAATTTGAT	GGAGATTATA	TTTCAAACT
1251	TCCTAAAAGT	AATCAGTATC	CAAGCCCTAA	AAGGTGTTTC	TCCTCCTATG
1301	ATGATATGAA	TTTGAAAAGA	AGAACTTCAT	TCCCACTCAT	GAGATCCAAG
1351	TCTCTACAGT	TAACTGGCAA	AGAAGTAGAC	CCAAATGATT	TGTACATTGT
1401	AGAACCCCTC	AAGTTTCTC	CAGAAAAGAA	GAAGAAGCGC	TGCAAGTACA
1451	AAATTGAAAA	AATTGAGACC	ATAAAGCCTG	AAGAACCATT	GCACCCATA
1501	GCCAATGGCG	ACATAAAAGG	AAGAAAGCCC	TTTACGAACC	AGAGAGATTT
1551	TTCTAATATG	GGAGAAGTTT	ATCACTCTTC	TTATAAAGGT	CCTCCATCTG
1601	AAGGAAGCTC	AGAACTTCA	TCACAGTCAG	AAGAGTCCTA	TTTTTGTGGC
1651	ATTTTCAGCTT	GCACAAGTCT	GTGCAATGGA	CAGTCTCAAA	AGACAAAAAC
1701	TGAGAAGAGG	GCTTTAAAAAC	GAAGACGATC	TAAAGTCCAA	GACCAAGGAA
1751	AATTGATAAA	AACTCTAATA	CAGACTAAGT	CGGGATCATT	GCCGAGCCTG
1801	CATGACATAA	TCAAAGGAAA	CAAAGAGATC	ACCGTGGGAA	CATTTGGTGT
1851	TACAGCAGTC	TCGGGACATA	TCTAAAATTA	ATTGAACTTT	TCATACAGAA
1901	GACTTTTTTG	TTGTGTGTTCT	TTGAAGAACA	GTCTGTAGTA	TTTGAAGGGT
1951	TTGGGGGAGG	GAGAAAATAT	TAATGGGAAA	GGCATTTCAGA	AATTATGGTT
2001	TCTACCTTTT	TAAAAAGTAG	ATGGGATTGT	GCTCAATCTT	GGTTAATGAG
2051	CTACAGTTTT	ACAAAGCTGA	TCATTTCCTA	TAAGGACAAT	GGTAGACATT
2101	TTATAAAGAT	GTTTTTTCAC	AAGATTAAAT	ACTGGGACAA	AAGTAATTTG
2151	GAAGCCAGT	TCCTTAGGTG	GGATAGGAAT	GAAAGCCTAA	ACCTCTTCCT
2201	TTAGCTTTGT	TCCTATTTCT	TGCACCTTCC	CATATTTATG	TGCCTTTTGT
2251	CTATTTTATA	TGCCACTGGA	AGAGGAGGGA	TAACTTTTTT	TGTTATTTGA
2301	TTTCTTTTAT	AACTTTGTTA	GGTTTTTGAA	GCTGCAACAA	CTACAATGCT
2351	TTGAGGGGGT	CTGTGCCTGA	AGCTCAGGAG	TGTGGATCAG	ACAGTCTAAA
2401	GATCCTAAAA	ACTTGCCAAC	TGGATCTTTG	TTTAGCAAAC	TCACTGGAAA
2451	TGAACACTTA	ATGGAATTTT	TAAGTCTGTT	CTGTTAGGTA	GATGGTGATG
2501	CTCTTGTTAT	TTTCACTTAT	TCAGGCTGGA	TTACTTCTTA	CTTAGTTACT
2551	AACTCAATGA	GGAAAAAATC	CCTACAGGAT	CTTTTTTTGC	AAACAACTGA

Figure 8(B)

```

2601 TATATGCAGA CAAATTTTGG ACAAATTCAC CTTTAAACA CGACGTTAAC
2651 CGATTTGTGA AGGTTTTCTT TAGCTTACAT TTTAAACATA CACAATAAAC
2701 ACTAATCCTC CAAACTTTCA CTGTTTTTAT TAGTATGAAT ATAAAATTTG
2751 AAGGTTTGGC CAATTAGTAC AAGTCTCATG ATATAATCAC AGCCTGCATA
2801 CATATGCACA GATCCAGTTA GTGAGTTTGT CAAGCTTAAT CTAATTGGTT
2851 AAGTCTAAAG AGATTATTAT TCCTTGATGT TTGCTTTGTA TTGGCTACAA
2901 ATGTGCAGAG GTAATACATA TGTGATGTCG ATGTCTCTGT CTTTTTTTTT
2951 GTCTTTAAAA AATAATTGGC AGCAACTGTA TTTGAATAAA ATGATTTCTT
3001 AGTATGATTG TACAGTAATG AATGAAAGTG GAACATGTTT CTTTTTGAAA
3051 GGGAGAGAAT TGACCATTTA TTGTTGTGAT GTTTAAGTTA TAACTTATTG
3101 AGCACTTTTA GTAGTGATAA CTGTTTTTAA ACTTGCCTAA TACCTTTCTT
3151 GGGTATTGTT TGTAATGTGA CTATTTTAAC GCCTTCCTTG TTTGTTTAAG
3201 TTGCTGCTTT AGGTTAACAG CGTGTTTTAG AAGATTTAAA TTTCTTTCTT
3251 GTCTGCACAA TTAGCTATTC AGAGCAAGAG GGCCTGATTT TATAGAAGCC
3301 CCTTGAAAAG AGGTCCAGAT GAGAGCAGAG ATACAGTGAG AAATTATGTG
3351 ATCTGTGTGT TGTGGGAAGA GAATTTTCAA TATGTAACTA CGGAGCTGTA
3401 GTGCCATTAG AAACGTGAA TTTCCAAATA AATCTGAACA CTTGTCTTTA
3451 TT

```

Figure 9

```

1      EYIYKWCSVR VALYRQRSRT ALSKGKDYLVAQPPLLLPA ESVDVSVLQP
51     LSGELENTINI EREAETVVLG DLSSSMHQDD LVNHTVDAVE LEPSHSQTL
101    QSLLLDITPE INPLPKIEVS ESVEYEAGHI PSPVIPQESS VEIDNETEQK
151    SESFSSIEKP SITYETNKVN ELMDNIIKED MNSMQIFTKL SETIVPPINT
201    ATVPDNEDGE AKMNIADTAK QTLISVVDSS SLPEVKEEEQ SPEDALLRGL
251    QRTATDFYAE LQNSTDLGYA NGNLVHGSNQ KESVFMRLNN RIKALEVNMS
301    LSGRYLEELS QRYRKQMEEM QKAFNKTIVK LQNTSRIAE QDQRQTEAIQ
351    LLQAQLTNMT QLVSNLSATV AELKREVSDR QSYLVISLVL CVVLGLMLCM
401    QRCRNTSQFD GDYISKLPKS NQYPSPKRCF SSYDDMNLKR RTSFPLMR
451    SLQLTGKEVD PNDLYIVEPL KFSPEKKKKR CKYKIEKIET IKPEEPLHPI
501    ANGDIGRKP FTNQDFSNM GEVYHSSYKG PPSEGSSETS SQSEESYFCG
551    ISACTSLCNG QSQKTKTEKR ALKRRRSKVQ DQGLIKTLI QTKSGSLPSL
601    HDIIKGNKEI TVGTFGVTAV SGHI•N•LNF SYRRLFCCCS LKNSL•YK
651    LGEGENINGK GIQKLWFLPF •KVDGIVLNL G••ATVLQS• SLPIRTMVDI
701    L•RCFFTRLI TGTKVIWKPS SLGGIGMKA• TSSFSFVPIS CTFPYLCAFC
751    LFIMPLEEEG •LFLLFDFFY NFVRFLKLQT LQCFEGVCA• SSGVWIRQSK
801    DPKNLPTGSL FSKLTGNEHL MEFLSLFC•V DGDALVIFTY SGWITSYLV
851    NSMRKSLQD LFLQTTDICR QIFDKFTF•T RR•PICEGFL •LTF•TYTIN
901    TNPPNFHCFY •YEYKI•RFG QLVQVS•YNH SLHTYAQIQL VSLSSLI•LV
951    KSKEIIP•C LLCIGYKCAE VIH•CRCLC LFFCL•KIIG SNCI•IK•FL
1001   SMIVQ••MKV EHVSF•KGEN •PFIVVMFKL •LIEHF•••• LFLNLPNTFL
1051   GYCL•CDLFN AFFVCLSCCF RLTACFRRFK FLSCLN•LF RARGPDFTEA
1101   P•KEVQMAE IQ•EIM•SVC CGKRIFNM•L RSCSAIRNCE FPNKSEHLSL

```

```

1      EYIYKWCSVR VALYRQRSRT ALSKGKDYLVAQPPLLLPA ESVDVSVLQP
51     LSGELENTINI EREAETVVLG DLSSSMHQDD LVNHTVDAVE LEPSHSQTL
101    QSLLLDITPE INPLPKIEVS ESVEYEAGHI PSPVIPQESS VEIDNETEQK
151    SESFSSIEKP SITYETNKVN ELMDNIIKED MNSMQIFTKL SETIVPPINT
201    ATVPDNEDGE AKMNIADTAK QTLISVVDSS SLPEVKEEEQ SPEDALLRGL
251    QRTATDFYAE LQNSTDLGYA NGNLVHGSNQ KESVFMRLNN RIKALEVNMS
301    LSGRYLEELS QRYRKQMEEM QKAFNKTIVK LQNTSRIAE QDQRQTEAIQ
351    LLQAQLTNMT QLVSNLSATV AELKREVSDR QSYLVISLVL CVVLGLMLCM
401    QRCRNTSQFD GDYISKLPKS NQYPSPKRCF SSYDDMNLKR RTSFPLMR
451    SLQLTGKEVD PNDLYIVEPL KFSPEKKKKR CKYKIEKIET IKPEEPLHPI
501    ANGDIGRKP FTNQDFSNM GEVYHSSYKG PPSEGSSETS SQSEESYFCG
551    ISACTSLCNG QSQKTKTEKR ALKRRRSKVQ DQGLIKTLI QTKSGSLPSL
601    HDIIKGNKEI TVGTFGVTAV SGHI

```

Figure 10

+ strand (sense)		sequence (5'-->3')
1. pch8-sp6-1f	369	GCT AAG CCA GAG CTA CAG G
2. pch8-sp6-2f	677	tCT GAT CTT CTG CTG ATT C
3. pch8-1fa	1238	(CTC) TCT GAA CTG CCT GAG AGA C
4. pch8-2f	1462	CCA AAT GGG AGC ATT ACA AG
5. pch8-3f	1745	TCA TCA AAT GAT CAG AAC C
6. pch8-4f	1995	ATT CTG GAG AGT TGG TAT CC
7. pch8-5f	2277	GGA ATA AGG AAA GAG CTT G
8. pch8-6f	2559	TCC ACT CAT ATT CCA ATA CC
9. pch8-5rb	2849	CCT GAG AGA CAG AAC TGT TC
10. pch8-4rb	3090	GGA CCC TTC ACT TCC TTA C
11. pch8-3rb	3370	GGC CAC CAC TTG TCC TGG G
12. pch8-2rb	3517	CAG AAC AGT GCT CTA ACT G
13. pch8-1rb	3970	GTA CTG CCT CTC TTA AAT G
- strand (antisense)		sequence (5'-->3')
14. pch8-2r	3617	CAG TTA CAG CAC TGT TCT G
15. pch8-3r	3360	CCC AGG ACA AGT GGT GGC C
16. pch8-4r	3140	GTA AGG AAG TGA AGG GTC C
17. pch8-5r	3849	GAA CAG TTC TGT CTC TCA GG
18. pch8-6r	3563	CTT GGG TAT TGG AAT ATG AG
19. pch8-5fb	2277	CAA GCT CTT TCC TTA TTC C
20. pch8-4fb	1999	ATA GGA TAC CAA CTC TCC AG
21. pch8-3fb	1746	TGG TTC TGA TCA TTT GAT G
22. pch8-2fb	1462	CTT GTA ATG CTC CCA TTT GG
23. pch8-1fb	1238	GTC TCT CAG GCA GTT CAG A
24. pch8-fb-1f	941	GTA GAG AAT CAC GTA CAG C
25. pch8-fb-2f	612	CAA TGA CCA GTA GCA TAA C
26. CH8-3670	3891	CAG CAT TTA AGA GAG GCA G
27. CH8a	387	CCT GTA GCT CTG GCT TAG CAT CC
28. CH8b	510	CCC CTT CAT TGA GAT CAT CTA G

Figure 11(A)

1	GTGCGCCGTG	GCGCGGCCCG	GCTGACAGGT	TCTTTAATGG	AGGAGCCAAT
51	CTCTCTGCAC	ACCTGGTTTC	ATCTAATAAT	ATACAGACAC	CAGCTCTGAG
101	GCCAGTTAAT	CATCCCCAGT	GTCCAGGCAC	AGAGTAGTCG	GTCCGCCTCA
151	CAATGTTGGA	CTTCTAGCC	GAGAACAACC	TCTGTGGCCA	AGCAATCCTA
201	AGGATTGTTT	CCTGTGGTAA	TGCCATCATT	GCTGAACTTT	TGAGACTCTC
251	TGAGTTTATT	CCTGCTGTGT	TCAGGTAAA	AGACAGAGCT	GATCAACAGA
301	AATATGGAGA	TATCATATTT	GATTTTCAGCT	ATTTTAAGGG	TCCAGAATTA
351	TGGGAAAGCA	AACTGGATGC	TAAGCCAGAG	CTACAGGATT	TAGATGAAGA
401	ATTTTCGTGAA	AACAACATAG	AAATTGTGAC	CAGATTTTAT	TTAGCATTTTC
451	AAAGTGATCA	TAAATATATT	GTAGACTTAA	ACAGATATCT	AGATGATCTC
501	AATGAAGGGG	TTTATATTCA	GCAAACCTTA	GAAACTGTGC	TTCTCAATGA
551	AGATGGAAAA	CAACTTCTAT	GTGAAGCACT	GTACTTATAT	GGAGTTATGC
601	TACTGGTCAT	TGACCAAAAG	ATTGAAGGAG	AAGTCAGAGA	GAGGATGCTG
651	GTTTCTTACT	ACCGATACAG	TGCTGCTCGA	TCTTCTGCTG	ATTCAAATAT
701	GGACGATATT	TGTAAGCTGC	TTCGAAGTAC	AGGTTATTCT	AGCCAACCAG
751	GTGCCAAAAG	ACCATCCAAC	TATCCCGAGA	GCTATTTCCA	GAGAGTGCCT
801	ATCAACGAAT	CCTTCATCAG	TATGGTCATT	GGTCGACTGA	GATCTGATGA
851	TATTTACAAC	CAGGTCTCAG	CGTATCCTTT	GCCGGAGCAT	CGCAGCACAG
901	CCCTGGCAAA	CCAAGCTGCC	ATGCTGTACG	TGATTCTCTA	CTTTGAGCCT
951	TCCATCCTTC	ACACCCATCA	AGCAAAAATG	AGAGAGATAG	TGGATAAATA
1001	CTTTCCAGAT	AATTGGGTAA	TTAGTATTTA	CATGGGGATC	ACAGTTAATC
1051	TAGTAGATGC	TTGGGAACCT	TACAAAGCTG	CAAAAACCTG	TTTAAATAAT
1101	ACCCTGGACC	TTTCAAATGT	CAGAGAACAG	GCAAGCAGAT	ATGCTACTGT
1151	CAGTGAAAGA	GTGCATGCTC	AAGTCAGCA	ATTTCTAAAA	GAAGGTTATT
1201	TAAGGGAGGA	GATGGTTCTG	GACAATATCC	CAAAGCTTCT	GAAGTGCCTG
1251	AGAGACTGCA	ATGTTGCCAT	CCGATGGCTG	ATGCTTCATA	CAGCAGACTC
1301	AGCCTGTGAC	CCAAACAACA	AACGCCTTCG	TCAAATCAAG	GACCAGATTG
1351	TAAAGACTC	TCGGTACAAT	CCCAGGATCC	TCTTCCAGCT	GCTGTTAGAT
1401	ACTGCACAAT	TTGAGTTTAT	ACTCAAAGAG	ATGTTCAAGC	AAATGCTTTC
1451	AGAAAAGCAA	ACCAAATGGG	AGCATTACAA	GAAAGAGGGT	TCCGAGCCGA
1501	TGACTGAGCT	TGCTGATGTC	TTTTTCAGGAG	TGAAACCCCT	AACCAGAGTG
1551	GAGAAAAATG	AAAACCTTCA	AGCTTGGTTC	AGAGAGATCT	CAAAACAAAT
1601	ATTGTCTTTA	AATTATGATG	ATTCTACTGC	TGCGGGCAGA	AAAACCTGTAC
1651	AACTGATACA	AGCTTTGGAA	GAGGTTCAAG	AATTCCACCA	GTTGGAATCC
1701	AATCTGCAAG	TATGTCAGTT	TCTTGCCGAT	ACTCGAAAGT	TTCTTCATCA
1751	AATGATCAGA	ACCATTAACA	TTAAAGAGGA	GGTTCTGATC	ACAATGCAGA
1801	TCGTTGGGGA	CCTTCTTTTC	GCTTGGCAGT	TGATTGACAG	TTTCACATCC
1851	ATCATGCAAG	AAAGCATAAG	GGTAAATCCA	TCCATGGTTA	CTAAACTCAG
1901	AGCTACCTTC	CTAAAGCTTG	CCTCTGCCCT	CGATCTGCCC	CTTCTTCGTA
1951	TTAATCAGGC	AAATCGCCCC	GACCTGCTCA	GCGTGTCA	GTACTATTCT
2001	GGAGAGTTGG	TATCCTATGT	GAGAAAAGTT	TTGCAGATCA	TCCAGAAAG
2051	CATGTTTACA	TCTCTTCTAA	AGATCATAAA	GCTTCAGACC	CACGACATTA
2101	TTGAAGTGCC	TACCCGCCTG	GACAAAGACA	AGCTGAGGGA	CTATGCTCAG
2151	CTAGGCCCCAC	GATACGAGGT	TGCCAAGCTT	ACTCATGCTA	TTTCCATTTT
2201	TACTGAAGGC	ATCTTAATGA	TGAAAACGAC	TTTGGTTGGC	ATCATCAAGG
2251	TGGATCCAAA	GCAGTTGCTG	GAAGATGGAA	TAAGGAAAGA	GCTTGTGAAG
2301	CGCGTTGCCT	TTGCCCTGCA	TAGGGGACTG	ATATTCAACC	CTCGAGCCAA
2351	GCCAAGTGAA	TTGATGCCCA	AGCTGAAAAG	GTTGGGAGCG	ACCATGGATG
2401	GATTCCATCG	TTCTTTTGAA	TACATACAGG	ACTATGTCAA	CATTTATGGT
2451	CTGAAGATTT	GGCAGGAAGA	AGTATCTCGT	ATCATAAATT	ACAACGTGGA
2501	GCAAGAGTGT	AATAACTTTC	TAAGAACGAA	GATTCAAGAT	TGGCAAAGCA
2551	TGTACCAGTC	CACTCATATT	CCAATACCCA	AGTTTACCCC	TGTGGATGAG

Figure 11(B)

2601	TCTGTAACGT	TTATTGGTCG	ACTCTGCAGA	GAAATCCTGC	GGATCACAGA
2651	CCCCAAAATG	ACATGTCACA	TAGACCAGCT	GAACACTTGG	TATGATATGA
2701	AAACTCATCA	GGAAGTGACC	AGCAGCCGCC	TCTTCTCAGA	AATCCAGACC
2751	ACCTTGGGAA	CCTTTGGTCT	AAATGGCTTA	GACAGGCTTC	TGTGCTTTAT
2801	GATTGTAAAA	GAGTTACAGA	ATTTCTCAG	TATGTTTCAG	AAAATTATCC
2851	TGAGAGACAG	AACTGTTTCA	GACACTTTAA	AAACCCTCAT	GAATGCTGTC
2901	AGTCCCCTAA	AAAGTATTGT	CGCAAATTCA	AATAAAATTT	ATTTTTCCGC
2951	CATTGCCAAA	ACACAGAAGA	TTTGGACTGC	GTATCTCGAG	GCTATAATGA
3001	AGGTTGGGCA	GATGCAGATT	CTGAGGCAAC	AGATTGCCAA	TGAATTAAAT
3051	TATTCTTGTC	GGTTTGATTC	TAAACATCTG	GCAGCTGCTC	TGGAGAATCT
3101	CAATAAGGCT	CTCCTAGCAG	ACATTGAAGC	CCACTATCAG	GACCCTTCAC
3151	TTCTTACCC	CAAAGAAGAT	AACACACTTT	TATATGAAAT	CACAGCCTAT
3201	CTGGAGGCAG	CTGGCATTCA	CAACCCACTG	AATAAGATAT	ACATAACAAC
3251	AAAGCGCTTA	CCCTATTTTC	CAATTGTAAA	CTTTCTATTT	TTGATCGCTC
3301	AGTTGCCAAA	ACTTCAATAC	AACAAAATC	TGGGAATGGT	CTGCCGAAAA
3351	CCGACCGACC	CGGTTGATTG	GCCACCACTT	GTCTGGGAC	TGCTCACTCT
3401	GCTGAAGCAG	TTCCATTCCC	GGTACACCGA	GCAGCTCCTG	GCGCTGATTG
3451	GCCAGTTTAT	CTGCTCCACG	GTGGAGCAGT	GTACAAGCCA	GAAGATACCT
3501	GAAATTCTCTG	CAGATGTTGT	GGGTGCCCTT	CTGTTCTGG	AGGATTATGT
3551	TCGGTACACA	AAGCTACCCA	GGAGGGTTGC	TGAAGCACAT	GTGCCTAATT
3601	TCATTTTGA	TGAGTTCAGA	ACAGTGCTGT	AACTGTTTTT	CCTACTTCTT
3651	CAATGGAAGG	ATTGTCCTTA	GATCTTCCCA	CCATCACAAA	TGAATTTGAA
3701	GATGAAAAAG	AACTCAGTTG	CTCATAACAAC	TGCATTTTTT	CTGTCTATTA
3751	TGGGAAACAT	CAGACGTTAT	GAGTAAGATA	TATCTCATGG	CATTAGTTAA
3801	TATAACTGAT	ATTGTTTAAA	TCATGGTATT	ACATGCAATT	TATATCAGAT
3851	AAAAGCAGAA	CACATTTTGG	TACTGCCTCT	CTTAAATGCT	GAATGTAAC
3901	GTTATGTATA	AATCCATTTA	GTMTTATGTT	CTAAAGAACT	ATTTGTGCAA
3951	CTCCAGATTT	TCAGTAAAT	AGTATTACTA	GT	

Figure 12(A)

APWRGPADRF FNGGANLSAH LVSSNNIOTP ALRPVNHPOC PGTE•SVRLT
 MLDFLAENNL CGQAILRIVS CGNAIIAELL RLSEFIPAVF RLKDRADQOK
 YGDIIFDFS Y FKGPPELWESK LDKPELQDL DEEFRENNIE IVTRFYLAFO
 SVHKYIVDLN RYLDDLNEG V YIQQTLETVL LNEDGKQLLC EALYLYGVML
 LVIDQKIEGE VRERMLVSY RYSAARSSAD SNMDDICKLL RSTGYSSQPG
 AKRPSNYPE S YFQRPINES FISMVIGRLR SDDIYNQVSA YPLPEHRSTA
 LANQAAMLV Y ILYFEPSILH THQAKMREIV DKYFPDNWVI STYMGITVNL
 VDAWEPYKAA KTALNNTLDL SNVREQASRY ATVSEVHAQ VQOFLKEGYL
 REEMVLDNIP KLLNCLRDCN VAIRWMLMHT ADSACDPNNK RLRQIKDQIL
 TDSRYNPRIL FQLLLDTAQF EFILKEMFKQ MLSEKQTKWE HYKKEGSERM
 TELADVFSGV KPLTRVEKNE NLQAWFREIS KQILSLNYDD STAAGRKTVO
 LIQALEEVQE FHQLESNLQV CQFLADTRKF LHQMIRTINI KEEVLITMQI
 VGDLSFAWQL IDSFTSIMQE SIRVNPSMVT KLRATFLKLA SALDPLLR I
 NQANRPDLLS VSQYYSSELV SYVRKVLQII PESMFTSLLK IIKLQTHDII
 EVPTRLDKDK LRDYAQLGPR YEVAKLTHAI SIFTEGILMM KTTLVGIIKV
 DPKQLLEDGI RKELVKRVAF ALHRGLIFNP RAKPSELMPK LKELGATMDG
 FHRSFEYIQD YVNIYGLKIW QEEVSRIINY NVEQECNNFL RTKIQDWQSM
 YQSTHIPIPK FTPVDES VTF IGRLCREILR ITDPKMTCHI DQLNTWYDMK
 THQEVTSRL FSEIQTTLGT FGLNGLDRLL CFMIVKELQN FL SMFQKIIL
 RDRTVQDTLK TLMNAVSPK SIVANSNKIY FSAIAKTQKI WTAYLEAIMK
 VGOMQILRQQ IANELNYSR FDSKHLAAAL ENLNKALLAD IEAHYQDPSL
 PYPKEDNTLL YEITAYLEAA GIHNPLNKIY ITTKRLPYFP IVNFLFLIAQ
 LPKLQYNKNL GMVCRKPTDP VDWPPVLVGL LTLLKQFHRS YTEQLLALIG
 QFICSTVEQC TSQKIPEIPA DVVGALLFLE DYVRYTKLPR RVAEAHVPNF
 IFDEFRTVL• LFFLLLQWKD CP•IFPPSQM NLKMKRNSVA HTTAFFLSIM
 GNIRRYE•DI SHGIS•YN•Y CLNHGITCNL YQIKAEHIFV LPLLNAECNC
 YV•IHLVLC S KELFVQLQIF SKIVLL

Figure 12(B)

MLDFLAENNL CGQAILRIVS CGNAIIAELL RLSEFIPAVF RLKDRADQOK
 YGDIIFDFS Y FKGPPELWESK LDKAPELQDL DEEFRENNIE IVTRFYLAFO
 SVHKYIVDLN RYLDLNEG V YIQQTLETVL LNEDGKQLLC EALYLYGVML
 LVIDQKIEGE VRERMLVSYY RYSAARSSAD SNMDDICKLL RSTGYSSQPG
 AKRPSNYPES YFQRPINES FISMVIGRLR SDDIYNQVSA YPLPEHRSTA
 LANQAAMLYV ILYFEP SILH THQAKMREIV DKYFPDNWVI SIYMGITVNL
 VDAWEPYKAA KTALNNTLDL SNVREQASRY ATVSEVHAQ VQOFLKEGYL
 REEMVLNIP KLLNCLRDCN VAIRWMLHT ADSACDPNNK RLRQIKDQIL
 TDSRYNPRIL FQLLLDTAQF EFILKEMFKQ MLSEKQTKWE HYKKEGSERM
 TELADVFSGV KPLTRVEKNE NLQAWFREIS KQILSLNYDD STAAGRKTVO
 LIQALEEVQE FHQLESNLQV CQFLADTRKF LHQMIRTINI KEEVLITMQI
 VGDLSFAWQL IDSFTSIMQE SIRVNPSMVT KLRATFLKLA SALDPLLRRI
 NQANRPDLLS VSQYYSSELV SYVRKVLQII PESMFTSLK IIKLQTHDII
 EVPTRLDKDK LRDYAQLGPR YEVAKLTHAI SIFTEGILMM KTTLVGIIKV
 DPKQLLEDGI RKELVKRVAF ALHRGLIFNP RAKPSELMPK LKELGATMDG
 FHRSFEYIQD YVNIYGLKIW QEEVSRIINY NVEQECNNFL RTKIQDWQSM
 YQSTHIPIPK FTPVDES VTF IGRLCREILR ITDPKMTCHI DQNTWYDMK
 THQEVTSRL FSEIQTTLGT FGLNGLDRLL CFMTVKELQN FLMSFQKIIL
 RDRTVQDTLK TLMNAVSPK SIVANSNKIY FSAIAKTQKI WTAYLEAIMK
 VGOMQILRQQ IANELNYSCR FDSKHLAAAL ENLNKALLAD IEAHYQDPSL
 PYPKEDNTLL YEITAYLEAA GIHNPLNKIY ITTKRLPYFP IVNFLFLIAQ
 LPKLQYNKNL GMVCRKPTDP VDWPPVLGL LTLLKQFHSR YTEQLLALIG
 QFICSTVEQC TSQKIPEIPA DVVGALLFLE DYVRYTKLPR RVAEAHVNF
 IFDEFRTVL

Figure 13(A)

AGG GGC GGA AGT CGG GGT CTG ACC CGC TCC AGG TCC GGG ACT GCG GAT
 AGA AGA GGA CCG CCG CCT TGA GGG AGG GGT GGA AAC TGG GTG CCG GCT
 CCG CGC GCG ACC TCC GGC CCT GCG CGT GCG CCG TGG CGC GGC CCG GCT
 GAC AGG TTC TTT AAT GGA GGA GCC AAT CTC TCT GCA CAC CTG GTT TCA
 TCT AAT AAT ATA CAG ACA CCA GCT CTG AGG CCA GTT AAT CAT CCC CAG
 TGT CCA GGC ACA GAG TAG TCG GTC CGC CTC ACA ATG TTG GAC TTT CTA
 GCC GAG AAC AAC CTC TGT GGC CAA GCA ATC CTA AGG ATT GTT TCC TGT
 GGT AAT GCC ATC ATT GCT GAA CTT TTG AGA CTC TCT GAG TTT ATT CCT
 GCT GTG TTC AGG TTA AAA GAC AGA GCT GAT CAA CAG AAA TAT GGA GAT
 ATC ATA TTT GAT TTC AGC TAT TTT AAG GGT CCA GAA TTA TGG GAA AGC
 AAA CTG GAT GCT AAG CCA GAG CTA CAG GAT TTA GAT GAA GAA TTT CGT
 GAA AAC AAC ATA GAA ATT GTG ACC AGA TTT TAT TTA GCA TTT CAA AGT
 GTA CAT AAA TAT ATT GTA GAC TTA AAC AGA TAT CTA GAT GAT CTC AAT
 GAA GGG GTT TAT ATT CAG CAA ACC TTA GAA ACT GTG CTT CTC AAT GAA
 GAT GGA AAA CAA CTT CTA TGT GAA GCA CTG TAC TTA TAT GGA GTT ATG
 CTA CTG GTC ATT GAC CAA AAG ATT GAA GGA GAA GTC AGA GAG AGG ATG
 CTG GTT TCT TAC TAC CGA TAC AGT GCT GCT CGA TCT TCT GCT GAT TCA
 AAT ATG GAC GAT ATT TGT AAG CTG CTT CGA AGT ACA GGT TAT TCT AGC
 CAA CCA GGT GCC AAA AGA CCA TCC AAC TAT CCC GAG AGC TAT TTC CAG
 AGA GTG CCT ATC AAC GAA TCC TTC ATC AGT ATG GTC ATT GGT CGA CTG
 AGA TCT GAT GAT ATT TAC AAC CAG GTC TCA GCG TAT CCT TTG CCG GAG
 CAT CGC AGC ACA GCC CTG GCA AAC CAA GCT GCC ATG CTG TAC GTG ATT
 CTC TAC TTT GAG CCT TCC ATC CTT CAC ACC CAT CAA GCA AAA ATG AGA
 GAG ATA GTG GAT AAA TAC TTT CCA GAT AAT TGG GTA ATT AGT ATT TAC
 ATG GGG ATC ACA GTT AAT CTA GTA GAT GCT TGG GAA CCT TAC AAA GCT
 GCA AAA ACT GCT TTA AAT AAT ACC CTG GAC CTT TCA AAT GTC AGA GAA
 CAG GCA AGC AGA TAT GCT ACT GTC AGT GAA AGA GTG CAT GCT CAA GTG
 CAG CAA TTT CTA AAA GAA GGT TAT TTA AGG GAG GAG ATG GTT CTG GAC
 AAT ATC CCA AAG CTT CTG AAC TGC CTG AGA GAC TGC AAT GTT GCC ATC
 CGA TGG CTG ATG CTT CAT ACA GCA GAC TCA GCC TGT GAC CCA AAC AAC
 AAA CGC CTT CGT CAA ATC AAG GAC CAG ATT CTA ACA GAC TCT CGG TAC
 AAT CCC AGG ATC CTC TTC CAG CTG CTG TTA GAT ACT GCA CAA TTT GAG
 TTT ATA CTC AAA GAG ATG TTC AAG CAA ATG CTT TCA GAA AAG CAA ACC
 AAA TGG GAG CAT TAC AAG AAA GAG GGT TCG GAG CGG ATG ACT GAG CTT
 GCT GAT GTC TTT TCA GGA GTG AAA CCC CTA ACC AGA GTG GAG AAA AAT
 GAA AAC CTT CAA GCT TGG TTC AGA GAG ATC TCA AAA CAA ATA TTG TCT
 TTA AAT TAT GAT GAT TCT ACT GCT GCG GGC AGA AAA ACT GTA CAA CTG
 ATA CAA GCT TTG GAA GAG GTT CAA GAA TTC CAC CAG TTG GAA TCC AAT
 CTG CAA GTA TGT CAG TTT CTT GCC GAT ACT CGA AAG TTT CTT CAT CAA
 ATG ATC AGA ACC ATT AAC ATT AAA GAG GAG GTT CTG ATC ACA ATG CAG
 ATC GTT GGG GAC CTT TCT TTC GCT TGG CAG TTG ATT GAC AGT TTC ACA
 TCC ATC ATG CAA GAA AGC ATA AGG GTA AAT CCA TCC ATG GTT ACT AAA
 CTC AGA GCT ACC TTC CTA AAG CTT GCC TCT GCC CTC GAT CTG CCC CTT
 CTT CGT ATT AAT CAG GCA AAT CGC CCC GAC CTG CTC AGC GTG TCA CAG
 TAC TAT TCT GGA GAG TTG GTA TCC TAT GTG AGA AAA GTT TTG CAG ATC
 ATC CCA GAA AGC ATG TTT ACA TCT CTT CTA AAG ATC ATA AAG CTT CAG
 ACC CAC GAC ATT ATT GAA GTG CCT ACC CGC CTG GAC AAA GAC AAG CTG
 AGG GAC TAT GCT CAG CTA GGC CCA CGA TAC GAG GTT GCC AAG CTT ACT
 CAT GCT ATT TCC ATT TTT ACT GAA GGC ATC TTA ATG ATG AAA ACG ACT

Figure 13(B)

TTG GTT GGC ATC ATC AAG GTG GAT CCA AAG CAG TTG CTG GAA GAT GGA
ATA AGG AAA GAG CTT GTG AAG CGC GTT GCC TTT GCC CTG CAT AGG GGA
CTG ATA TTC AAC CCT CGA GCC AAG CCA AGT GAA TTG ATG CCC AAG CTG
AAA GAG TTG GGA GCG ACC ATG GAT GGA TTC CAT CGT TCT TTT GAA TAC
ATA CAG GAC TAT GTC AAC ATT TAT GGT CTG AAG ATT TGG CAG GAA GAA
GTA TCT CGT ATC ATA AAT TAC AAC GTG GAG CAA GAG TGT AAT AAC TTT
CTA AGA ACG AAG ATT CAA GAT TGG CAA AGC ATG TAC CAG TCC ACT CAT
ATT CCA ATA CCC AAG TTT ACC CCT GTG GAT GAG TCT GTA ACG TTT ATT
GGT CGA CTC TGC AGA GAA ATC CTG CGG ATC ACA GAC CCA AAA ATG ACA
TGT CAC ATA GAC CAG CTG AAC ACT TGG TAT GAT ATG AAA ACT CAT CAG
GAA GTG ACC AGC AGC CGC CTC TTC TCA GAA ATC CAG ACC ACC TTG GGA
ACC TTT GGT CTA AAT GGC TTA GAC AGG CTT CTG TGC TTT ATG ATT GTA
AAA GAG TTA CAG AAT TTC CTC AGT ATG TTT CAG AAA ATT ATC CTG AGA
GAC AGA ACT GTT CAG GAC ACT TTA AAA ACC CTC ATG AAT GCT GTC AGT
CCC CTA AAA AGT ATT GTC GCA AAT TCA AAT AAA ATT TAT TTT TCC GCC
ATT GCC AAA ACA CAG AAG ATT TGG ACT GCG TAT CTC GAG GCT ATA ATG
AAG GTT GGG CAG ATG CAG ATT CTG AGG CAA CAG ATT GCC AAT GAA TTA
AAT TAT TCT TGT CGG TTT GAT TCT AAA CAT CTG GCA GCT GCT CTG GAG
AAT CTC AAT AAG GCT CTC CTA GCA GAC ATT GAA GCC CAC TAT CAG GAC
CCT TCA CTT CCT TAC CCC AAA GAA GAT AAC ACA CTT TTA TAT GAA ATC
ACA GCC TAT CTG GAG GCA GCT GGC ATT CAC AAC CCA CTG AAT AAG ATA
TAC ATA ACA ACA AAG CGC TTA CCC TAT TTT CCA ATT GTA AAC TTT CTA
TTTT TTG ATC GCT CAG TTG CCA AAA CTT CAA TAC AAC AAA AAT CTG GGA
ATG GTC TGC CGA AAA CCG ACC GAC CCG GTT GAT TGG CCA CCA CTT GTC
CTG GGA CTG CTC ACT CTG CTG AAG CAG TTC CAT TCC CGG TAC ACC GAG
CAG CTC CTG GCG CTG ATT GGC CAG TTT ATC TGC TCC ACG GTG GAG CAG
TGT ACA AGC CAG AAG ATA CCT GAA ATT CCT GCA GAT GTT GTG GGT GCC
CTT CTG TTC CTG GAG GAT TAT GTT CGG TAC ACA AAG CTA CCC AGG AGG
GTT GCT GAA GCA CAT GTG CCT AAT TTC ATT TTT GAT GAG TTC AGA ACA
GTG CTG TAA CTG TTT TTC CTA CTT CTT CAA TGG AAG GAT TGT CCT TAG
ATC TTC CCA CCA TCA CAA ATG AAT TTG AAG ATG AAA AGA AAC TCA GTT
GCT CAT ACA ACT GCA TTT TTT CTG TCT ATT ATG GGA AAC ATC AGA CGT
TAT GAG TAA GAT ATA TCT CAT GGC ATT AGT TAA TAT AAC TGA TAT TGT
TTA AAT CAT GGT ATT ACA TGC AAT TTA TAT CAG ATA AAA GCA GAA CAC
ATT TTT GTA CTG CCT CTC TTA AAT GCT GAA TGT AAC TGT TAT GTA TAA
ATC CAT TTA GTT TTA TGT TCT AAA GAA CTA TTT GTG CAA CTC CAG ATT
TTC AGT AAA ATA GTA TTA CTA GT

Figure 14(A)

Arg Gly Gly Ser Arg Gly Leu Thr Arg Ser Arg Ser Gly Thr Ala Asp
 Arg Arg Gly Pro Pro Pro * Gly Arg Gly Gly Asn Trp Val Pro Ala
 Pro Arg Ala Thr Ser Gly Pro Ala Arg Ala Pro Trp Arg Gly Pro Ala
 Asp Arg Phe Phe Asn Gly Gly Ala Asn Leu Ser Ala His Leu Val Ser
 Ser Asn Asn Ile Gln Thr Pro Ala Leu Arg Pro Val Asn His Pro Gln
 Cys Pro Gly Thr Glu * Ser Val Arg Leu Thr Met Leu Asp Phe Leu
 Ala Glu Asn Asn Leu Cys Gly Gln Ala Ile Leu Arg Ile Val Ser Cys
 Gly Asn Ala Ile Ile Ala Glu Leu Leu Arg Leu Ser Glu Phe Ile Pro
 Ala Val Phe Arg Leu Lys Asp Arg Ala Asp Gln Gln Lys Tyr Gly Asp
 Ile Ile Phe Asp Phe Ser Tyr Phe Lys Gly Pro Glu Leu Trp Glu Ser
 Lys Leu Asp Ala Lys Pro Glu Leu Gln Asp Leu Asp Glu Glu Phe Arg
 Glu Asn Asn Ile Glu Ile Val Thr Arg Phe Tyr Leu Ala Phe Gln Ser
 Val His Lys Tyr Ile Val Asp Leu Asn Arg Tyr Leu Asp Asp Leu Asn
 Glu Gly Val Tyr Ile Gln Gln Thr Leu Glu Thr Val Leu Leu Asn Glu
 Asp Gly Lys Gln Leu Leu Cys Glu Ala Leu Tyr Leu Tyr Gly Val Met
 Leu Leu Val Ile Asp Gln Lys Ile Glu Gly Glu Val Arg Glu Arg Met
 Leu Val Ser Tyr Tyr Arg Tyr Ser Ala Ala Arg Ser Ser Ala Asp Ser
 Asn Met Asp Asp Ile Cys Lys Leu Leu Arg Ser Thr Gly Tyr Ser Ser
 Gln Pro Gly Ala Lys Arg Pro Ser Asn Tyr Pro Glu Ser Tyr Phe Gln
 Arg Val Pro Ile Asn Glu Ser Phe Ile Ser Met Val Ile Gly Arg Leu
 Arg Ser Asp Asp Ile Tyr Asn Gln Val Ser Ala Tyr Pro Leu Pro Glu
 His Arg Ser Thr Ala Leu Ala Asn Gln Ala Ala Met Leu Tyr Val Ile
 Leu Tyr Phe Glu Pro Ser Ile Leu His Thr His Gln Ala Lys Met Arg
 Glu Ile Val Asp Lys Tyr Phe Pro Asp Asn Trp Val Ile Ser Ile Tyr
 Met Gly Ile Thr Val Asn Leu Val Asp Ala Trp Glu Pro Tyr Lys Ala
 Ala Lys Thr Ala Leu Asn Asn Thr Leu Asp Leu Ser Asn Val Arg Glu
 Gln Ala Ser Arg Tyr Ala Thr Val Ser Glu Arg Val His Ala Gln Val
 Gln Gln Phe Leu Lys Glu Gly Tyr Leu Arg Glu Glu Met Val Leu Asp
 Asn Ile Pro Lys Leu Leu Asn Cys Leu Arg Asp Cys Asn Val Ala Ile
 Arg Trp Leu Met Leu His Thr Ala Asp Ser Ala Cys Asp Pro Asn Asn
 Lys Arg Leu Arg Gln Ile Lys Asp Gln Ile Leu Thr Asp Ser Arg Tyr
 Asn Pro Arg Ile Leu Phe Gln Leu Leu Leu Asp Thr Ala Gln Phe Glu
 Phe Ile Leu Lys Glu Met Phe Lys Gln Met Leu Ser Glu Lys Gln Thr
 Lys Trp Glu His Tyr Lys Lys Glu Gly Ser Glu Arg Met Thr Glu Leu
 Ala Asp Val Phe Ser Gly Val Lys Pro Leu Thr Arg Val Glu Lys Asn
 Glu Asn Leu Gln Ala Trp Phe Arg Glu Ile Ser Lys Gln Ile Leu Ser
 Leu Asn Tyr Asp Asp Ser Thr Ala Ala Gly Arg Lys Thr Val Gln Leu
 Ile Gln Ala Leu Glu Glu Val Gln Glu Phe His Gln Leu Glu Ser Asn
 Leu Gln Val Cys Gln Phe Leu Ala Asp Thr Arg Lys Phe Leu His Gln
 Met Ile Arg Thr Ile Asn Ile Lys Glu Glu Val Leu Ile Thr Met Gln
 Ile Val Gly Asp Leu Ser Phe Ala Trp Gln Leu Ile Asp Ser Phe Thr
 Ser Ile Met Gln Glu Ser Ile Arg Val Asn Pro Ser Met Val Thr Lys
 Leu Arg Ala Thr Phe Leu Lys Leu Ala Ser Ala Leu Asp Leu Pro Leu
 Leu Arg Ile Asn Gln Ala Asn Arg Pro Asp Leu Leu Ser Val Ser Gln
 Tyr Tyr Ser Gly Glu Leu Val Ser Tyr Val Arg Lys Val Leu Gln Ile
 Ile Pro Glu Ser Met Phe Thr Ser Leu Leu Lys Ile Ile Lys Leu Gln
 Thr His Asp Ile Ile Glu Val Pro Thr Arg Leu Asp Lys Asp Lys Leu
 Arg Asp Tyr Ala Gln Leu Gly Pro Arg Tyr Glu Val Ala Lys Leu Thr
 His Ala Ile Ser Ile Phe Thr Glu Gly Ile Leu Met Met Lys Thr Thr
 Leu Val Gly Ile Ile Lys Val Asp Pro Lys Gln Leu Leu Glu Asp Gly
 Ile Arg Lys Glu Leu Val Lys Arg Val Ala Phe Ala Leu His Arg Gly

Figure 14(B)

Leu Ile Phe Asn Pro Arg Ala Lys Pro Ser Glu Leu Met Pro Lys Leu
 Lys Glu Leu Gly Ala Thr Met Asp Gly Phe His Arg Ser Phe Glu Tyr
 Ile Gln Asp Tyr Val Asn Ile Tyr Gly Leu Lys Ile Trp Gln Glu Glu
 Val Ser Arg Ile Ile Asn Tyr Asn Val Glu Gln Glu Cys Asn Asn Phe
 Leu Arg Thr Lys Ile Gln Asp Trp Gln Ser Met Tyr Gln Ser Thr His
 Ile Pro Ile Pro Lys Phe Thr Pro Val Asp Glu Ser Val Thr Phe Ile
 Gly Arg Leu Cys Arg Glu Ile Leu Arg Ile Thr Asp Pro Lys Met Thr
 Cys His Ile Asp Gln Leu Asn Thr Trp Tyr Asp Met Lys Thr His Gln
 Glu Val Thr Ser Ser Arg Leu Phe Ser Glu Ile Gln Thr Thr Leu Gly
 Thr Phe Gly Leu Asn Gly Leu Asp Arg Leu Leu Cys Phe Met Ile Val
 Lys Glu Leu Gln Asn Phe Leu Ser Met Phe Gln Lys Ile Ile Leu Arg
 Asp Arg Thr Val Gln Asp Thr Leu Lys Thr Leu Met Asn Ala Val Ser
 Pro Leu Lys Ser Ile Val Ala Asn Ser Asn Lys Ile Tyr Phe Ser Ala
 Ile Ala Lys Thr Gln Lys Ile Trp Thr Ala Tyr Leu Glu Ala Ile Met
 Lys Val Gly Gln Met Gln Ile Leu Arg Gln Gln Ile Ala Asn Glu Leu
 Asn Tyr Ser Cys Arg Phe Asp Ser Lys His Leu Ala Ala Ala Leu Glu
 Asn Leu Asn Lys Ala Leu Leu Ala Asp Ile Glu Ala His Tyr Gln Asp
 Pro Ser Leu Pro Tyr Pro Lys Glu Asp Asn Thr Leu Leu Tyr Glu Ile
 Thr Ala Tyr Leu Glu Ala Ala Gly Ile His Asn Pro Leu Asn Lys Ile
 Tyr Ile Thr Thr Lys Arg Leu Pro Tyr Phe Pro Ile Val Asn Phe Leu
 Phe Leu Ile Ala Gln Leu Pro Lys Leu Gln Tyr Asn Lys Asn Leu Gly
 Met Val Cys Arg Lys Pro Thr Asp Pro Val Asp Trp Pro Pro Leu Val
 Leu Gly Leu Leu Thr Leu Leu Lys Gln Phe His Ser Arg Tyr Thr Glu
 Gln Leu Leu Ala Leu Ile Gly Gln Phe Ile Cys Ser Thr Val Glu Gln
 Cys Thr Ser Gln Lys Ile Pro Glu Ile Pro Ala Asp Val Val Gly Ala
 Leu Leu Phe Leu Glu Asp Tyr Val Arg Tyr Thr Lys Leu Pro Arg Arg
 Val Ala Glu Ala His Val Pro Asn Phe Ile Phe Asp Glu Phe Arg Thr
 Val Leu * Leu Phe Phe Leu Leu Leu Gln Trp Lys Asp Cys Pro *
 Ile Phe Pro Pro Ser Gln Met Asn Leu Lys Met Lys Arg Asn Ser Val
 Ala His Thr Thr Ala Phe Phe Leu Ser Ile Met Gly Asn Ile Arg Arg
 Tyr Glu * Asp Ile Ser His Gly Ile Ser * Tyr Asn * Tyr Cys
 Leu Asn His Gly Ile Thr Cys Asn Leu Tyr Gln Ile Lys Ala Glu His
 Ile Phe Val Leu Pro Leu Leu Asn Ala Glu Cys Asn Cys Tyr Val *
 Ile His Leu Val Leu Cys Ser Lys Glu Leu Phe Val Gln Leu Gln Ile
 Phe Ser Lys Ile Val Leu Leu

Figure 15

+ strand (sense)		sequence (5'-->3')
	1st base	
1. pch13-sp6-1f	370	TTT ACT TCT AAC GCT TAT TC
2. pch13-sp6-2f	726	TGA AGG AGT CCT TTG AGA CG
3. T7.1	1140	TCA CAA TGG GCT ACT GG
4. T7.2	1361	TTC AAC GAG GGA GAT GG
5. T7.3	1602	TTA GCA CCA CTG AGA GA
6. T7.4	2041	GTT CTT TTA GGC ATT TA
7. ch13-2480	2486	GCT GCG TCT GTT CGT CAG C
- strand (antisense)		
8. SP6.1	2746	CCT CTG CTT CAC AAC AT
9. SP6.2	2490	GCA GtA GGG CGG ACA CC
10. SP6.3	2213	(C) AGG GTC TTC TTC ATT GT
11. SP6.4	1812	GGA TTG TCT TTG TCT CT
12.pch13-t7-1f	1165	AGT GCA CTT CCA TGG GCG TG
13.pch13-t7-1fa	712	CCT TCA TCA GGT TGA CGA AC
14.pch13-t7-2fa	286	GCG GCA ATC AGA AAC GGA AG
15.CH13-AS-1	536	TGA ACA CGT GGT ACA T

Figure 16(A)

1	CTTCCCTGAG	CCCTTCTGTC	CTGTGTAGGA	AGCAGAAGGC	GGAATGTCCG
51	CTCTGCCCTT	CTCCGTAAGA	TGGTGCATTA	AAACGTTCCCT	TATAAACTGG
101	AAATGAAGGC	TTGGGAAGAT	GGCTAAAATC	AGCAATCCTT	GGAATAACGC
151	AGAAGCATCC	CTGCTTCCCT	GGGCCCCGCC	GTGGGCCTGC	TTGTGCTGTT
201	CAGTAGGTGG	TTTTTAGAAA	GGGCTTCCTT	CAGCGTCATT	AGCAACAGGA
251	GTGTCGTCC	GTTTGCATGA	GGAAATGTTT	TTAACCTTCC	GTTTCTGATT
301	GCCTCTAGAC	TGCATCTGTC	ATAGACAAAT	GCCCCATCT	TTTACAGAGA
351	ACCAGTCTCT	TCTTTAAACT	TTACTTCTAA	CGCTTATTCT	TTTTACCTTA
401	TATAGGAAAC	CACTGATTGC	TTGTGTGGAG	AAACAGCTAT	TAGGAGAACA
451	TTTAACAGCA	ATTCTGCAGA	AAGGGCTCGA	CCACTTACTG	GATGAGAACA
501	GAGTGCCGGA	CCTCGCACAG	ATGTACCAGC	TGTTTCAGCCG	GGTGAGGGGC
551	GGGCAGCAGG	CGCTGCTGCA	GCACTGGAGC	GAGTACATCA	AGACTTTTGG
601	AACAGCGATC	GTAATCAATC	CTGAGAAAGA	CAAAGACATG	GTCCAAGACC
651	TGTTGGACTT	CAAGGACAAG	GTGGACCACG	TGATCGAGGT	CTGCTTCCAG
701	AAGAATGAGC	GGTTCGTCAA	CCTGATGAAG	GAGTCCTTTG	AGACGTTTAT
751	CAACAAGAGA	CCCAACAAGC	CTGCAGAACT	GATCGCAAAG	CATGTGGATT
801	CAAAGTTAAG	AGCAGGCAAC	AAAGAAGCCA	CAGACGAGGA	GCTGGAGCGG
851	ACGTTGGACA	AGATCATGAT	CCTGTTTCAGG	TTTATCCACG	GTAAAGATGT
901	CTTTGAAGCA	TTTTTATAAA	AAGATTTGGC	AAAAAGACTC	CTTGTGTTGG
951	AAAGTGCCTC	AGTCGATGCT	GAAAAGTCTA	TGTTGTCAAA	GCTCAAGCAT
1001	GAGTGCCGTG	CAGCCTTCAC	CAGCAAGCTG	GAAGGCATGT	TCAAGGACAT
1051	GGAGCTTTTC	AAGGACATCA	TGGTTTCATT	CAAGCAGCAT	ATGCAGAATC
1101	AGAGTGACTC	AGGCCCTATA	GACCTCACAG	TGAACATACT	CACAATGGGC
1151	TACTGGCCAA	CATACACGCC	CATGGAAGTG	CACTTAACCC	CAGAAATGAT
1201	TAAACTTCAG	GAGTATTTTA	AGGCATTTTA	TCTTGGAAG	CACAGTGGTC
1251	GAAAACCTTCA	GTGGCAAACT	ACTTTGGGAC	ATGCTGTTTT	AAAAGCGGAG
1301	TTTAAAGAAG	GGAAGAAGGA	ATTCCAGGTG	TCCCTCTTCC	AGACACTGGT
1351	GCTCCTCATG	TTCAACGAGG	GAGATGGCTT	CAGCTTTGAG	GAGATAAAAA
1401	TGGCCACGGG	GATAGAGGAT	AGTGAATTGC	GCAGAACGCT	GCAGTCCCTG
1451	GCCTGTGGCA	AAGCACGTGT	GCTGATTAAA	AGTCCCAAAG	GAAAGGAAGT
1501	GGAAGATGGA	GACAAGTTCA	TTTTTAAATG	AGAGTTCAAG	CACAAGTTGT
1551	TTAGAATAAA	GATCAATCAA	ATTCCAGATGA	AGGAAACTGT	TGAGGAACAG
1601	GTTAGCACCA	CTGAGAGAGT	GTTTCAGGAT	AGACAATATC	AGATTGATGC
1651	TGCTATCGTC	AGAATAATGA	AGATGAGAAA	GACTCTTGGT	CATAATCTTC
1701	TAGTTTCTGA	ATTATATAAT	CAGCTGAAAT	TTCCAGTAAA	GCCTGGAGAT
1751	TTGAAAAAGA	GAATTGAATC	TCTGATAGAC	AGAGACTATA	TGGAGAGAGA
1801	CAAAGACAAT	CCGAATCAGT	ACCACTACGT	GGCCTGACGC	ATCTGCAGAC
1851	GGTTCCCTTT	CATGAAACAC	TAGAATGTAC	CCTCAGAGCA	GGAAGCACAC
1901	CTGTGCCATT	TCTGGGACTC	TGATTGATCC	AGCTGTGGAC	ATTGGAAGGC
1951	GAAGGAAGGG	AGGTGGCTCC	TGGGTCTACT	TTTCAAGGC	TCAAGACTTC
2001	AACCTGCAGA	TGTATCTTTT	TCCCTCCAGT	TTTTCTCTTA	GTTCTTTTAG
2051	GCATTTAAAT	TGTTTCTGTT	ACTCTGTGCA	AAATAACTTT	GAGATTGGAC
2101	AAGAAGATGT	TACTAAAGAG	AAGTTCCCTT	AAAAGGTCTT	GTTCTTGTGT
2151	CAAAAAGCTG	CAAGTTTGGT	TTGTTCTCGT	GTGTGATCAT	GAGTGACAAA
2201	TGAAGAAGAC	CCTAGATGCT	GCATTTTTTA	GCTCTGAAGA	TTCCCTTAGGT
2251	ATCCCTGAAG	ACAGCTCGCT	CAGATGATCA	GCATTTAGAG	TGAAAACAAG
2301	GGCCCTTCAT	GGGTGAACAT	TAGAAAGAGC	CAGGGTTCAA	AGCTGGCGAA
2351	TGGATGACGC	ACCCTAGCCA	CTGGCCCCCT	CCTGTTTCAT	GTATTTCCAA
2401	AAGTTGTAAA	CTTTGCTGGC	TGATTTTTTC	TAAGTCAGGT	TTCTAAGTGA
2451	GCTCCCTGAG	GTGCCAAGGC	CATGGTGTCC	GCCCTGCTGC	GTCTGTTCGT
2501	CAGCTGAGTT	CCTTGTGAAT	CTCTGTTTTA	GGGGTTGGGG	CTAGTGTGTT

Figure 16(B)

2551	TGTGTTTCCA	TTCTAAGATT	GAGTCTGGCA	GTCCCTGTTT	TTTTGCATTG
2601	GGGTAACTGC	TCTTTGATTT	TTTTTAATTG	CAGTATTTGT	GTGATTGCAA
2651	TAATAAAGTT	TGGTTTGGTT	TTTACAGTCA	TGCGCAGGGA	CGATCCTTGT
2701	TCTCTGCTGT	AAACTGTAAA	AAGTTTATGG	AGACCTAAAG	TCTTGATGTT
2751	GTGAAGCAGA	GGTTATTTTG	TGGAAAGATT	AAAAGGATTT	TGTTGGTACC
2801	TGGTTTTGTG	TGTGTATAT	ATACATGAGG	TTGAACAGTG	AAAGGAAAGT
2851	TCAGTAGTGA	TGTTAGAAGG	GTAACATGA	CAAAGATACT	TTTGAGATAA
2901	CATTTAAAAG	TACTTTATAT	TTTACATAAT	AGCATGTTTC	ATTTTGATTA
2951	AAAGCTACCA	AAGGAATTTT	GATCATGGCA	TAAGTGTTTA	AAGCAATATT
3001	TTCTGGAATA	TACCAAGTTT	ATATAATTTG	ATTTTGTGCT	AAATTATTAA
3051	GAGTCTCTTT	TTGAAACATG	CGGGTTTGAA	ATATGACACC	TTGTGGGTTT
3101	CCATATTAAA	ATCCTCACTC	TTTAATTGTC	ATTTTATCT	TTGAAAATTT
3151	TCATTTATGA	GTTCCATGAT	ATGTGGTCTA	AGAAAGACCA	AACAGATTTT
3201	TATTTTTTTT	TCTTATAAGT	TCGTTGTGTC	TAGAGATTGT	TAATATTGTA
3251	ATTTAATGTA	GACTTACTTT	GAATAAAATT	AGTTTAATTG	GCCTTAAAAT
3301	TACATTAATA	AAACTTTGTG	ATATGCAAAT	GACACATTC	

Figure 17

```

1      FPEPFLPV•E AEGGMSALPF SVRWCIKTFL INWK•RLGKM AKISNPWNNA
51     EASLLPWARP WACLCCSVGG F•KGLPSASL ATGVVVRLE EMFLTFRF•L
101    PLDCICHRQM PPSFTENQSL L•TLLLTLIL FTLYRKPLIA CVEKQLLGEH
151    LTAILQKGLD HLLDENRVPD LAQMYQLFSR VRGGQQALLQ HWSEYIKTFG
201    TAIVINPEKD KDMVQDLLDF KDKVDHVIEV CFQKNERFVN LMKESFETFI
251    NKRPNKPAEL IAKHVDSKLR AGNKEATDEE LERTLDKIMI LFRFIHGKDV
301    FEA FYKKDLA KRLLVGKSAS VDAEKSMLSK LKHECGAAFT SKLEGFMFKDM
351    ELSKDIMVHF KQHMQNQSDS GPIDLTVNIL TMGYWPTYTP MEVHLTPEMI
401    KLQEVFKAFY LGKHSGRKLQ WQTTLGHAVL KAEFKEGKKE FQVSLFQTLV
451    LLMFNEGDGF SFEEIKMATG IEDSELRRTL QSLACGKARV LIKSPKGKEV
501    EDGDKFIFNG EFKHKLFRK INQIQMKETV EEQVSTTERV FQDRQYQIDA
551    AIVRIMMRK TLGHNLLVSE LYNQLKFPVK PGDLKKRIES LIDRDYMERD
601    KDNPNQYHYV A•RICRRFPF MKH•NVPSEQ EAHLCFWDSD D•SSCGHWKA
651    KEGRWLLGHL SQGSRLQPAD VSFSLQFFL• FF•AFKLFL LCAK•L•DWT
701    RRCY•REVPL KGLVLVSKSC KFGLFSCVIM SAQ•RRP•ML HFLALKIP•V
751    SLKTARSDDQ HLE•KQGPFM GEH•KEPGFK AGEWMTHPSH WPLPVSCISK
801    SCKLWWLIFR KSGF•VSSLR CQGHGVRPAA SVRQLSSL•I SVLGVGASVF
851    VFPF•D•VWQ SLFFCIGVTA L•FFLIADFV •LQ••SLVWF LQSCAGTILV
901    LCCKL•KVYG DLKS•CCEAE VILWKD•KDF VGTWFCVVI YMLNLSERKV
951    Q••C•KGNVD KDTFEITFKS TLYFT••HVS F•LKATKGIL IMA•VFKAIF
1001   SGIYQVYII• FCAKLLRVSF •NMRV•NMTP CGFPY•NPHS LIVIFIFENF
1051   HL•VP•YVV• ERPNRFLFFF LISSLCLEIV NIVI•CRLTL NKISLIGLKI
1101   TLIKLCMQM TH

```

```

201    TAIVINPEKD KDMVQDLLDF KDKVDHVIEV CFQKNERFVN LMKESFETFI
251    NKRPNKPAEL IAKHVDSKLR AGNKEATDEE LERTLDKIMI LFRFIHGKDV
301    FEA FYKKDLA KRLLVGKSAS VDAEKSMLSK LKHECGAAFT SKLEGFMFKDM
351    ELSKDIMVHF KQHMQNQSDS GPIDLTVNIL TMGYWPTYTP MEVHLTPEMI
401    KLQEVFKAFY LGKHSGRKLQ WQTTLGHAVL KAEFKEGKKE FQVSLFQTLV
451    LLMFNEGDGF SFEEIKMATG IEDSELRRTL QSLACGKARV LIKSPKGKEV
501    EDGDKFIFNG EFKHKLFRK INQIQMKETV EEQVSTTERV FQDRQYQIDA
551    AIVRIMMRK TLGHNLLVSE LYNQLKFPVK PGDLKKRIES LIDRDYMERD
601    KDNPNQYHYV A

```

Figure 18

+ strand (sense)		sequence (5'-->3')
	1st base	
1. pch14-sp6-1f	686	GGC TTA ACA CTC AAT GTA C
2. pch14-sp6-2f	1005	CTA TGA AAA GAC AGC TTA AG
3. pch14-SP6-3f	1315	ATT TAG TTT GAA AAG CAT G
4. pch14-sp6-4f	1589	CAG ACT TTA AAG TCA CAA G
5. pch14-sp6-5f	1808	CAA AGA CTT GGT GTA TAG TG
- strand(antisense)		sequence (5'-->3')
6. pch14-sp6-6fb	2020	GCA GTT TAA TTT GGT CCT G
7. pch14-sp6-5fb	1757	CTG TAA TTA TAG TTC TGT C
8. pch14-sp6-4fb	1607	CTT GTG ACT TTA AAG TCT G
9. pch14-sp6-3fb	1339	ATA ATC ATG CTT TTC AAA C
10. pch14-sp6-2rb	1023	TTA AGC TGT CTT TTC ATA G
11. pch14-sp6-1rb	704	GTA CAT TGA GTG TTA AAC C
12. CH14a	629	CGG CAG AGC TGA CTA CTG GAA GG
13. CH14b	644	CAA GCA GGG AAG TAA CGG CAG
14. CH14c	109	CTT GTT AGC TTG TTT AGA AGG TGG AAG AG
15. CH14d	90	GGT GGA AGA GAA GGT CTC CTT TCA GGC

Figure 19

1	GAAGATGATG	ATTACGGGTC	TCGAACAGGA	AGCATCTCCA	GCAGTGTGTC
51	TGTGCCTGCA	AAGCCTGAAA	GGAGACCTTC	TCTTCCACCT	TCTAAACAAG
101	CTAACAAGAA	TCTGATTTTG	AAGGCTATAT	CTGAAGCTCA	AGAATCCGTA
151	ACAAAAACAA	CTAACTACTC	TACAGTTCCA	CAGAAACAGA	CACTTCCAGT
201	TGCTCCCAGA	ACTCGAACTT	CTCAAGAAGA	ATTGCTAGCA	GAAGTGGTCC
251	AGGGACAAAG	TAGGACCCCC	AGAATAAGTC	CCCCCATTA	AGAAGAGGAA
301	ACAAAAGGAG	ATTCTGTAGA	AAAAAATCAA	GCTGAGATGA	GTGAACTGAG
351	TGTGGCACAG	AAACCAGAAA	AACTTTTGGA	GCGCTGCAAG	TACTGGCCTG
401	CTTGTAAGAA	TGGGGATGAG	TGTGCCTACC	ATCACCCCAT	CTCACCCCTGC
451	AAAGCCTTCC	CCAATTGTAA	ATTTGCTGAA	AAATGTTTGT	TTGTTTCAACC
501	AAATTGTAAA	TATGATGCAA	AGTGACTATA	ACCAGATTGT	CCCTTCACTC
551	ATGTGAGTAG	AAGAATTCCA	GTACTGTCTC	CAAAACCACT	TGCACCACCA
601	GCACCACCTT	CCAGTAGTCA	GCTCTGCCGT	TACTTCCCTG	CTTGTAAGAA
651	GATGGAATGT	CCCTTCTATC	ATCCAAAACA	TTGTAGGTTT	AACACTCAAT
701	GTACAAGTCC	GGACTGCACA	TTCTACCATC	CCACCATTAA	TGTCCCACCA
751	CGACATGCCT	TGAAATGGAT	TCGACCTCAA	ACCAGCGAAT	AGCACCCAGT
801	CCTGCCTGGC	AGAAGATCAT	GCAGTTTGGA	AGTTTTTCATG	TACTGATGAA
851	AGATACTCTA	CAGAACTTGT	CAAATCTTTG	AAACTTGGAA	TATATTGCTT
901	TCATAATATG	AAGTTTTATT	GCCTATCTAT	CTGAAGTGTC	TAATTTTTTCA
951	AGTTTGTAA	TTTATTATGT	GGTTTAAACA	TTGGGTGTTT	TTGTTTTTGT
1001	TTTACTATGA	AAAGACAGCT	TAAGGAAGAG	CTAAATTCTG	TTAAAATATT
1051	TGGGGCATGT	TTGTGCACTG	CTGTTGTGAG	GATCAGCATA	TGAAATTGAC
1101	ATCATGGTTA	GTCATGGTAC	TGCAGCTTAG	GGGGCTACAC	GGTTGCTGTG
1151	TGAGTGGAGA	GATGCAGTGA	GGCAGTTGTC	ATTATTCTAA	AAATTGTACT
1201	ACTTTCACCT	TTCCCAAAGA	TTATATAATG	TTCATATCC	ACCATGAAAA
1251	CAGCATTGGC	CAAAGGTACT	GAGGCTGCTT	AAAATATTCA	ATTCTGCTTT
1301	TTAGTTTTTA	AGTGAATTTA	GTTTGAAAAG	CATGATTATA	CAGGCCTCTC
1351	GAGGCTGAGT	GCTACTTTTCG	GTAAAGTTCC	AGTTTTCCAG	CCTTCTGTGA
1401	CAGGATGAAT	GAGGTGGGTA	TGGACAGTGG	AGGCAGCTGG	AATGGCAAGT
1451	GCAGAAAATA	GGAACAGTTC	TATACAGTGC	TCTCATTTAC	TAATAACATA
1501	ATGCCCTTCTA	AATAATTTTT	TTGGGAAACT	ACATTATCAC	AAAATTATAC
1551	AAATTTTTTT	ACAAGTATTT	ACATACTGTA	TCTGAAAACA	GACTTTAAAG
1601	TCACAAGATT	ATAAATGTAC	ATATGTATTTC	TCACATTCTG	AAAAATAACA
1651	TTCTCAGAAT	CCACAGAAAA	TATACTTAGT	TACTACTGAA	GATAATTTTT
1701	GAAATGTAAA	AATTAGATTT	AAATAGTATA	TTTTAAATGA	CAGAACTATA
1751	ATTACAGAGA	TCAGATCAGA	TAGGTAAACT	GCAAGATAGA	TAGGATGAAA
1801	CTTTTGGCCT	ACTGTATTAC	TTACAGAGTT	TTTTTGTGTG	TGGTTTTTTAA
1851	AACTGTTAAG	GCAAGAAGTG	TCAAATGCTT	TAGAGTTAAA	TAACAGATCA
1901	CTGATTTCAA	AGACTTGGTG	TATAGTGTTA	AAAATTAAAG	CTTAAAAGGT
1951	GGTTAGAAAA	GTGGATTAAT	GCAAAAGGGG	TAATAAAGAC	TGCAACATTC
2001	TCAGGACCAA	ATTAACTGCT	T		

Figure 20

```

1      EDDDYGSRTG SISSSVSVPA KPERRPSLPP SKQANKNLIL KAISEAQESV
51     TKTINYSTVP QKQTLFVAPR TRTSQEELLA EVVQGQSRTP RISPPIKEEE
101    TKGDSVEKNQ AEMSELSVAQ KPEKLLERCK YWPACKNGDE CAYHHPISPC
151    KAFPNCFAE KCLFVHPNCK YDAKCTKPCD PFTHVSRRIIP VLSPKPVAPP
201    APPSSSQLCR YFPACKKMEC PFYHPKHCRF NTQCTSPDCT FYHPTINVPP
251    RHALKWIRPQ TSE•HPVLPQ RRSCSLEVFM Y••KILYRTC QIFETWNILL
301    S•YEVLLPIY LKCLIFQVCK FIMWF•HWVF LFCFYYEKTA •GRAKFC•NI
351    WGMFVHCCCE DQHMKLTSWL VMVLQLRGLH GCCVSGEMQ• GSCHYSKNCT
401    TFTFPKDYIM FIIHHENSIG QRY•GCLKYS ILLFSF•VNL V•KA•LYRPL
451    EAECYFR•SS SFPAFCDRMN EVGMDSGGSW NGKCRK•EQF YTVLSFTNNI
501    MPSK•FFWET TLSQNYTNFF TSIYILYLKT DFKVTRL•MY ICILTF•KIT
551    FSESTENILS YY•R•FLKCK N•I•IVYFK• QNYNYRDQIR •VNCKIDRMK
601    LLAYCITYRV FLCVVFCTVK ARSVKCFRVK •QITDFKDLV YSVKN•SLKG
651    G•KSGLMQKG ••RLQHSQDQ IKL

```

```

EDDDYGSRTG SISSSVSVPA KPERRPSLPP SKQANKNLIL KAISEAQESV
TKTINYSTVP QKQTLFVAPR TRTSQEELLA EVVQGQSRTP RISPPIKEEE
TKGDSVEKNQ AEMSELSVAQ KPEKLLERCK YWPACKNGDE CAYHHPISPC
KAFPNCFAE KCLFVHPNCK YDAKCTKPCD PFTHVSRRIIP VLSPKPVAPP
APPSSSQLCR YFPACKKMEC PFYHPKHCRF NTQCTSPDCT FYHPTINVPP
RHALKWIRPQ TSE

```

Figure 21

```

1      AAAACTTTTCG GAAGAGAAAG TTGCCTGTGG TAAGTTCAGT TGTAAAGTA
51     AAAAAATTCA ATCATGATGG AGAAGAGGAG GAAGAAGATG ATGATTACGG
101    GTCTCGAACA GGAAGCATCT CCAGCAGTGT GTCTGTGCCT GCAAAGCCTG
151    AAAGGAGACC TTCTCTTCCA CCTTCTAAAC AAGCTAACAA GAATCTGATT
201    TTGAAGGCTA TATCTGAAGC TCAAGAATCC GTAACAAAAA CAACTAACTA
251    CTCTACAGTT CCACAGAAAC AGACACTTCC AGTTGCTCCC AGAACTCGAA
301    CTTCTCAAGA AGAATTGCTA GCAGAAGTGG TCCAGGGGAC AAAGTAGGAC
351    CCCCAGAATA AGTCCCCCCA TTAAAGAAGA GGAAACAAAA GGAGATTCTG
401    TAGAAAAAAA TCAAGATTAC TATGACATGG AATCCATGGT CCATGCAGAC
451    ACAAGATCAT TTATTCTGAA GAAGCCAAAG CTGTCTGAGG AAGTANTAGT
501    GGCACCAAAC CAAGANTCGG GGATGAAGAC TGCAGATTCC CTTCCGGGTTT
551    TTTTCAGGGAC CCTTATGCAG ACACNAGATC TTGTTCAACC AGATAAACCT
601    GCAAGTCCCA AG

```

```

1      KTFGRESCLW •VQLLK•KNS IMMEKRRKKM MITGLEQEAS PAVCLCLQSL
51     KGDLLFHLN KLTRI•F•RL YLKLKNP•QK QLITLQFHRN RHFQLLPELE
101    LLKKNC•QKW SRGQSRTPRI SPPIKEEETK GDSVEKNQDY YDMESMVHAD
151    TRSFILKKPK LSEEVXVAPN QXSGMKTADS LRVLSGTLMQ TXDLVQPDKP
201    ASPK

```

```

1      NAGCTGCTCT GACGGGNAGN GGAATGNATG GNGGCTTGTT CNGAAACNNG
51     CCAGATGGCG NGAGGGGGAC AAGTAGCGGC GTGATTNAGA AGAGGGAGGT
101    GAGGGTNCTC ACATCACCNC ATCTNACCAT GNCNGGCCNT CCCCANTANT
151    AANANTGATG ATAGNNGGAA GTGGGCCCAC CCAGAAGCNT GATTGAGCGG
201    CCGCCAGTAN GAAACNNGTT TGTCCANTTA GNCATACNNA TNGTAGGGTT
251    CNAGCNGCGT CCCCGGCACC NGCANANNNN CNNCNGGGAC NACNGCCCN
301    NNNNTNNGTTA NNCNGNGNAG NNAAAAAATT CAATCATGAT GGAGAAGAGG
351    AGGAAGAAGA TGATGATTAC GGGTCTCGAA CAGGAAGCAT CTCCAGCAGT
401    GTGTCTGTGC CTGCAAA

```

Untitled translated in RF 2

```

1      SCSDGXXNXW XLVXXARWX EGDK•RRDXE EGGE GXHITX SXHXXXSPXX
51     XXMIXGSGPT QKXD•AAASX KXVCPXXHXX XRVXXASPAX AXXXXGXXPX
101    XXLXXXXKKF NHDGEEEEED DDYGSRTGSI SSSVSVPA

```

Figure 22**CH1-9a11-2**

GA AAA CAA ATG GAA GAA ATG CAA AAG GCT TTC AAT AAA ACA ATC GTG
 AAA CTT CAG AAT ACT TCA AGA ATA GCA GAG GAG CAG GAT CAG CGG CAA
 ACT GAA GCC ATC CAG TTG CTA CAG GCA CAG CTG ACC AAC ATG ACA CAG
 CTT GTT CAA

Lys Gln Met Glu Glu Met Gln Lys Ala Phe Asn Lys Thr Ile Val Lys
 Leu Gln Asn Thr Ser Arg Ile Ala Glu Glu Gln Asp Gln Arg Gln Thr
 Glu Ala Ile Gln Leu Leu Gln Ala Gln Leu Thr Asn Met Thr Gln Leu
 Val Gln

CH8-2a13-1

GAA CAG GCA AGC AGA TAT GCT ACT GTC AGT GAA AGA GTG CAT GCT CAA
 GTG CAG CAA TTT CTA AAA GAA GGT TAT TTA AGG GAG GAG ATG GTT CTG
 GAC AAT ATC CCA AAG CTT CTG AAC TGC CTG AGA GAC TGC AAT GTT GCC
 ATC CGA TGG CTG ATG CTT C

Glu Gln Ala Ser Arg Tyr Ala Thr Val Ser Glu Arg Val His Ala Gln
 Val Gln Gln Phe Leu Lys Glu Gly Tyr Leu Arg Glu Glu Met Val Leu
 Asp Asn Ile Pro Lys Leu Leu Asn Cys Leu Arg Asp Cys Asn Val Ala
 Ile Arg Trp Leu Met Leu

CH13-2a12-1

CTC ACA ATG GGC TAC TGG CCA ACA TAC ACG CCC ATG GAA GTG CAC TTA
 ACC CCA GAA ATG ATT AAA CTT CAG GAA GTA TTT AAG GCA TTT TAT CTT
 GGA AAG CAC AG

Leu Thr Met Gly Tyr Trp Pro Thr Tyr Thr Pro Met Glu Val His Leu
 Thr Pro Glu Met Ile Lys Leu Gln Glu Val Phe Lys Ala Phe Tyr Leu
 Gly Lys His

CH14-2a16-1

TG TTT GTT CAC CCA AAT TGT AAA TAT GAT GCA AAG TGT ACT AAA CCA
 GAT TGT CCC TTC ACT CAT GTG AGT AGA AGA ATT CCA GTA CTG TCT CCA
 AAA CCA GTT GCA CCA CCA G

Phe Val His Pro Asn Cys Lys Tyr Asp Ala Lys Cys Thr Lys Pro Asp
 Cys Pro Phe Thr His Val Ser Arg Arg Ile Pro Val Leu Ser Pro Lys
 Pro Val Ala Pro Pro