

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2012293463 B2

(54) Title
Injectable solution of at least one type of basal insulin

(51) International Patent Classification(s)
A61K 38/28 (2006.01) **A61K 47/36** (2006.01)
A61K 9/00 (2006.01)

(21) Application No: **2012293463** (22) Date of Filing: **2012.08.09**

(87) WIPO No: **WO13/021143**

(30) Priority Data

(31) Number	(32) Date	(33) Country
11/57291	2011.08.10	FR
11/62445	2011.12.23	FR
61/579,966	2011.12.23	US
61/522,031	2011.08.10	US

(43) Publication Date: **2013.02.14**
(44) Accepted Journal Date: **2016.08.04**

(71) Applicant(s)
Adocia

(72) Inventor(s)
Soula, Olivier;Soula, Gerard;Tonnar, Jeff;Geissler, Alexandre

(74) Agent / Attorney
MINTER ELLISON, Attention: Grant Williams Rialto Towers 525 Collins Street, Melbourne, VIC, 3000

(56) Related Art
WO2007/121256

(12) DEMANDE INTERNATIONALE PUBLIÉE EN VERTU DU TRAITÉ DE COOPÉRATION EN MATIÈRE DE BREVETS (PCT)

(19) Organisation Mondiale de la Propriété Intellectuelle
Bureau international



(43) Date de la publication internationale
14 février 2013 (14.02.2013)

WIPO | PCT

(10) Numéro de publication internationale

WO 2013/021143 A1

(51) Classification internationale des brevets :
A61K 38/28 (2006.01) A61K 9/00 (2006.01)
A61K 47/36 (2006.01)

Avenue Félix Faure, F-69003 Lyon (FR). GEISSLER ALEXANDRE [FR/FR]; 28 Rue Camille Roy, 69007 LYON (FR).

(21) Numéro de la demande internationale :
PCT/FR2012/051880

(74) Mandataire : CABINET INES TRIPPOZ; 52 Rue du Colombier, F-69007 Lyon (FR).

(22) Date de dépôt international :
9 août 2012 (09.08.2012)

(81) États désignés (sauf indication contraire, pour tout titre de protection nationale disponible) : AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Langue de dépôt : français
(26) Langue de publication : français
(30) Données relatives à la priorité :
11/57291 10 août 2011 (10.08.2011) FR
61/522,031 10 août 2011 (10.08.2011) US
11/62445 23 décembre 2011 (23.12.2011) FR
61/579,966 23 décembre 2011 (23.12.2011) US

(84) États désignés (sauf indication contraire, pour tout titre de protection régionale disponible) : ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), eurasien (AM, AZ, BY, KG, KZ, RU, TJ, TM), européen (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, ZW).

(71) Déposant (pour tous les États désignés sauf US) : ADO-

CIA [FR/FR]; 115 Avenue Lacassagne, F-69003 Lyon

(FR).



(72) Inventeurs; et

(75) Inventeurs/Déposants (pour US seulement) : SOULA, Olivier [FR/FR]; 115 Avenue du Carreau, F-69330 Meyzieu (FR). SOULA, Gérard [FR/FR]; 33 Rue Nungesser, F-69330 Meyzieu (FR). TONNAR, Jeff [FR/FR]; 203

[Suite sur la page suivante]

(54) Title : INJECTABLE SOLUTION OF AT LEAST ONE TYPE OF BASAL INSULIN

(54) Titre : SOLUTION INJECTABLE D'AU MOINS UNE INSULINE BASALE



(57) Abstract : The invention relates to an injectable solution having a pH of 7, and including at least one type of basal insulin, the pI of which is between 5.8 and 8.5. The invention relates to a composition in the form of an injectable aqueous solution, the pH of which is between 6.0 and 8.0, including at least: a) one type of basal insulin, the isoelectric point (pI) of which is between 5.8 and 8.5; and b) a dextran substituted with radicals having carboxylate fillers and hydrophobic radicals of formula I or II, where R is -OH or is selected from the group consisting of the radicals $\text{o}-(\text{f}-[\text{A}]-\text{COOH})$, $\text{o}-(\text{g}-[\text{B}]-\text{k}-[\text{D}])_m$, D comprising at least one alkyl chain comprising at least 8 carbon atoms, where R is -OH or a radical $-(\text{f}-[\text{A}]-\text{COOH})_n$, and R' is selected from the group consisting of the radicals $\text{o}-\text{CH}_2\text{NH}-[\text{E}](-\text{o}-[\text{F}])_l$ and $\text{o}-\text{C}(\text{O})\text{NH}-[\text{E}](-\text{o}-[\text{F}])_l$.

(57) Abrégé : Solution injectable à PH 7 Comportant au moins une insuline basale dont le PI est compris entre 5,8 et 8,5. L'invention concerne une composition sous forme d'une

[Suite sur la page suivante]



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, —

ML, MR, NE, SN, TD, TG). — avant l'expiration du délai prévu pour la modification des revendications, sera republiée si des modifications sont reçues (règle 48.2.h))

Publiée :

— avec rapport de recherche internationale (Art. 21(3))

solution aqueuse injectable, dont le pH est compris entre 6,0 et 8,0, comprenant au moins : • a) une insuline basale dont le point isoélectrique pl est compris entre 5,8 et 8,5 et; • b) un dextrane substitué par des radicaux porteurs de charges carboxylates et des radicaux hydrophobes de formule I ou II : dans laquelle, • R est -OH ou choisi dans le groupe constitué par les radicaux : • o - (f-[A]-COOH), • o -(g-[B]-k-[D])_m, D comportant au moins une chaîne alkyle comportant au moins 8 atomes de carbone, dans laquelle, • R est -OH ou un radical -(f-[A]-COOH)_n • R' est choisi dans le groupe constitué par les radicaux : • o -CH₂NH-[E](-o-[F])_t • o -C(O)NH-[E](-o-[F])_t

INJECTABLE SOLUTION OF AT LEAST ONE TYPE OF BASAL INSULIN

5 [0001] The invention relates to therapies by injection of insulin(s) for treating diabetes.

[0002] Insulin therapy, or diabetes therapy by injecting insulin, has made remarkable progress in recent years by virtue especially of the development of novel insulins that offer correction of patients' glycaemia, which enables better simulation of 10 the physiological activity of the pancreas.

[0003] To cover his daily insulin needs, a diabetic patient currently has available, schematically, two types of insulin with complementary actions: prandial insulins (or rapid-acting insulins) and basal insulins (or slow-acting insulins).

[0004] Prandial insulins allow rapid management (metabolization and/or storage) of 15 the glucose provided by meals and snacks. The patient must inject a prandial insulin before each food intake, i.e. about 2 to 3 injections per day. The prandial insulins most commonly used are: the recombinant human insulin, NovoLog® (insulin aspart from Novo Nordisk), Humalog® (insulin lispro from Eli Lilly) and Apidra® (insulin glulisine from Sanofi-Aventis).

20 [0005] Basal insulins maintain the patient's glycaemic homeostasis, outside periods of food intake. They act essentially by blocking the endogenous production of glucose (hepatic glucose). The daily dose of basal insulin generally corresponds to 40-50% of the total daily insulin needs. Depending on the basal insulin used, this dose is dispensed in 1 or 2 injections, regularly distributed throughout the day. The basal 25 insulins most commonly used are Levemir® (insulin detemir from Novo Nordisk) and Lantus® (insulin glargine from Sanofi-Aventis).

[0006] It will be noted, for the sake of completeness, that NPH (NPH insulin, for Neutral Protamine Hagedorn; Humuline NPH®, Insulatard®) is the oldest basal insulin. This formulation is the result of a precipitation of human insulin (anionic at neutral pH) 30 with a cationic protein, protamine. These microcrystals are dispersed in an aqueous suspension and dissolve slowly after subcutaneous injection. This slow dissolution ensures sustained release of the insulin. However, this release does not ensure a constant concentration of insulin over time. The release profile is bell-shaped and lasts only between 12 and 16 hours. It is thus injected twice a day. This

NPH basal insulin is much less efficient than the modern basal insulins Levemir® and Lantus®. NPH is an intermediate-acting basal insulin.

[0007] The principle of NPH changed with the appearance of rapid insulin analogues to give "Premix" products that afford both rapid action and intermediate action. NovoLog Mix® (Novo Nordisk) and Humalog Mix® (Eli Lilly) are formulations comprising a rapid insulin analogue, Novolog® and Humalog®, partially complexed with protamine. These formulations thus contain microcrystals of insulin whose action is said to be intermediate, and a proportion of insulin which has remained soluble, whose action is rapid. These formulations do indeed offer the advantage of a rapid insulin, but they also have the defect of NPH, i.e. a duration of action limited to between 12 and 16 hours and a bell-shaped release of insulin. However, these products enable the patient to carry out a single injection of an intermediate-acting basal insulin with a rapid-acting prandial insulin. In point of fact, many patients are desirous to reduce their number of injections.

15

[0008] The basal insulins currently marketed and currently under clinical development may be categorized as a function of the technical solution for obtaining sustained action, and two approaches are used at the present time.

20 [0009] The first, that of insulin detemir, is the binding to albumin *in vivo*. It is an analogue, soluble at pH 7, which comprises a fatty acid (tetradecanoyl) side chain attached to position B29 which, *in vivo*, enables this insulin to combine with albumin. Its sustained action is mainly due to this affinity for albumin after subcutaneous injection.

25 [00010] However, its pharmacokinetic profile does not make it possible to cover a whole day, which means that it is usually used in two injections per day.

[00011] Other basal insulins which are soluble at pH 7, such as Degludec®, are currently under development. Degludec® also comprises a fatty acid side chain attached to insulin (hexadecanoyl- γ -L-Glu).

30 [00012] The second, that of insulin glargine, is the precipitation at physiological pH. This is an analogue of human insulin obtained by elongation of the C-terminal part of the B chain of human insulin with two arginine residues, and by substitution of the asparagine residue A21 with a glycine residue (US 5 656 722). The addition of the two arginine residues was conceived to adjust the pI (isoelectric point) of insulin glargine at physiological pH, and thus make this insulin analogue insoluble in physiological medium.

35 [00013] The substitution of A21 was conceived in order to make insulin glargine stable at acidic pH and thus to be able to formulate it in the form of an injectable solution at acidic pH.

During subcutaneous injection, the passage of insulin glargine from an acidic pH (pH 4-4.5) to a physiological pH (neutral pH) brings about its precipitation under the skin. The slow redissolution of the insulin glargine microparticles ensures a slow and sustained action.

5 [00014] The hypoglycaemic effect of insulin glargine is virtually constant over a period of 24 hours, which allows the majority of patients to limit themselves to a single injection per day.

[00015] Insulin glargine is nowadays considered as the best basal insulin marketed.

10 [00016] However, the acidic pH of the formulations of basal insulins, the isoelectric point of which is between 5.8 and 8.5, of insulin glargine type, prevents any pharmaceutical combination with other proteins and in particular prandial insulins, since the latter are unstable at acidic pH.

15 [00017] However, no one has hitherto sought to dissolve these basal insulins, the isoelectric point of which is between 5.8 and 8.5, of insulin glargine type, at neutral pH, while maintaining a difference in solubility between the *in vitro* medium (containing it) and the *in vivo* medium (under the skin), independently of the pH.

20 [00018] Specifically, the operating principle of basal insulins, of insulin glargine type, outlined above, namely that they are soluble at acidic pH and precipitate at physiological pH, dissuades a person skilled in the art from any solution in which insulin of insulin glargine type would be dissolved at pH 6-8 while maintaining its essential property, which is that of precipitating in subcutaneous medium.

25 [00019] Furthermore, the impossibility of formulating a prandial insulin, at acidic pH, arises from the fact that a prandial insulin undergoes, under these conditions, a deamidation side reaction in position A21; this does not make it possible to satisfy the requirement of the US Pharmacopeia, namely less than 5% of by-product after 4 weeks at 30°C.

30 [00020] Furthermore, this acidic pH of formulations of basal insulins, the isoelectric point of which is between 5.8 and 8.5, of insulin glargine type, even prevents any extemporaneous combination with prandial insulins at neutral pH.

35 [00021] A recent clinical study, presented at the 69th Scientific Sessions of the American Diabetes Association, New Orleans, Louisiana, 5-9 June 2009, verified this limitation of use of insulin glargine. A dose of insulin glargine and a dose of prandial insulin (in the case in point, insulin lispro) were mixed together just before injection (E. Cengiz et al., 2010, Diabetes Care, 33(5), 1009-12). This experiment made it possible to demonstrate a significant delay in the pharmacokinetic and pharmacodynamic profiles of prandial insulin, which may give rise

to postprandial hyperglycaemia and to nocturnal hypoglycaemia. This study indeed confirms the incompatibility of insulin glargine with the currently marketed rapid-acting insulins.

5 [00022] Moreover, the instructions for use for Lantus®, the commercial product based on insulin glargine from the company Sanofi-Aventis, explicitly indicates to users that they should not mix it with a solution of prandial insulin, whichever it may be, on account of the serious risk of modifying the pharmacokinetics and pharmacodynamics of the insulin glargine and/or of the mixed prandial insulin.

10 [00023] However, from a therapeutic point of view, clinical studies made public during the 70th annual scientific sessions of the *American Diabetes Association* (ADA) of 2010, abstract 2163-PO and abstract number 0001-LB, in particular those conducted by the company Sanofi-Aventis, showed that treatments which combine Lantus®, insulin glargine, and a prandial insulin are much more efficient than 15 treatments based on products of the "Premix", Novolog Mix® or Humalog Mix® type.

[00024] As regards combinations of insulin glargine and rapid insulin, the company Biodel has notably described, in patent application US 7 718 609, compositions comprising a basal insulin and a prandial insulin at a pH of between 3.0 and 4.2 in the presence of a chelating agent and polyacids. This patent teaches how to make a 20 prandial insulin compatible at acidic pH in the presence of an insulin of insulin glargine type. It does not teach how to prepare a combination of insulin of insulin glargine type and of a prandial insulin at neutral pH.

[00025] From the analysis of the compositions described in the literature and the 25 patents, it appears that the insolubility at pH 7 of basal insulins of the insulin glargine type is a prerequisite for having its slow action. As a result, all the solutions proposed for combining them with other products, such as prandial insulins, are based on tests of dissolution or stabilization of prandial insulins at acidic pH; see, for example, WO 2007/121256 and WO 2009/021955.

30 Surprisingly, the compositions according to the invention make it possible to dissolve, at pH 7, a basal insulin whose isoelectric point is between 5.8 and 8.5.

Surprisingly, the compositions according to the invention allow maintenance of the 35 duration of the hypoglycaemic activity of the basal insulin whose isoelectric point is between 5.8 and 8.5, despite its dissolution at pH 7 before injection. This noteworthy property arises from the fact that the insulin of insulin glargine type

dissolved at pH 7 in the composition of the invention precipitates in subcutaneous medium by means of a change of composition of the medium. The factor triggering the precipitation of the insulin of insulin glargine type is no longer the pH modification, but a modification of composition of the environment during the passage of the

5 pharmaceutical composition into the physiological medium.

[00026] By solving this problem of solubility at pH 7, the present invention makes it possible:

- to propose an injectable composition, intended for treating diabetes, comprising a basal insulin whose isoelectric point is between 5.8 and 8.5, in the form of a

10 homogeneous solution at pH 7, while retaining its biological activity and its action profile;

- to propose a composition in the form of a formulation comprising a combination of a basal insulin whose isoelectric point is between 5.8 and 8.5 and a prandial insulin without modification of the activity profile of the prandial insulin which is soluble at

15 pH 6-8 and unstable at acidic pH, while maintaining the basal action profile intrinsic to the basal insulin;

- to propose an injectable composition, intended for treating diabetes, additionally comprising a combination of a basal insulin whose isoelectric point is between 5.8 and 8.5 and a derivative or an analogue of a gastrointestinal hormone such as


20 GLP-1 or "glucagon-like peptide-1";

- for patients to reduce their number of injections;
- for the said compositions to satisfy the requirements of the American Pharmacopeia and European Pharmacopoeia.

25 [00027] Surprisingly, in insulin combinations of insulin glargine type with a prandial insulin, which are subject-matters of the invention, the rapid action of the prandial insulin is preserved despite the precipitation of the insulin of insulin glargine type in subcutaneous medium.

[00028] The invention relates to a composition in the form of an injectable aqueous solution, the pH of which is between 6.0 and 8.0, comprising at least:

5 a) a basal insulin, the isoelectric point π I of which is between 5.8 and 8.5;
 b) a dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals of formula I or of formula II:



in which:

10 • R is $-\text{OH}$ or chosen from the group consisting of the radicals:
 ○ $-(f\text{-}[A]\text{-COOH})_n$;
 ○ $-(g\text{-}[B]\text{-}k\text{-}[D])_m$, D comprising at least one alkyl chain comprising at least 8 carbon atoms;

15 • n represents the degree of substitution of the glucoside units by $-f\text{-}[A]\text{-COOH}$ and $0.1 \leq n \leq 2$;

20 • m represents the degree of substitution of the glucoside units by $-g\text{-}[B]\text{-}k\text{-}[D]$ and $0 < m \leq 0.5$;

25 • q represents the degree of polymerization as glucoside units, that is to say the mean number of glucoside units per polysaccharide chain, and $3 \leq q \leq 50$;

30 • $-(f\text{-}[A]\text{-COOH})_n$:
 ○ $-A$ - is a linear or branched radical comprising from 1 to 4 carbon atoms; the said $-A$ - radical:
 ○ being bonded to a glucoside unit *via* a functional group f chosen from the group consisting of ether, ester and carbamate functional groups;

 • $-(g\text{-}[B]\text{-}k\text{-}[D])_m$:
 ○ $-B$ - is a linear or branched, at least divalent, radical comprising from 1 to 4 carbon atoms; the said $-B$ - radical:
 ○ being bonded to a glucoside unit *via* a functional group g chosen from the group consisting of ether, ester and carbamate functional groups;

- being bonded to a -D radical *via* a functional group *k*; *k* chosen from the group consisting of ester, amide and carbamate functional groups; the said -D radical:

5


- being an $-X(-I-Y)_p$ radical, *X* being an at least divalent radical comprising from 1 to 12 atoms chosen from the group consisting of C, N and O atoms, optionally carrying carboxyl or amine functional groups and/or resulting from an amino acid, a dialcohol, a diamine or a mono- or polyethylene glycol mono- or diamine; *Y* being a linear or cyclic alkyl group, an alkylaryl or an arylalkyl, of 8 to 30 carbon atoms, optionally substituted by one or more C_1 to C_3 alkyl groups; $p \geq 1$ and *I* a functional group chosen from the group consisting of ester, amide and carbamate functional groups;

10

20

15

- *f*, *g* and *k* being identical or different;
- the free acid functional groups being in the form of salts of alkali metal cations chosen from the group consisting of Na^+ and K^+ ;
- and, when $p = 1$, if *Y* is a C_8 to C_{14} alkyl, then $q^*m \geq 2$, if *Y* is a C_{15} alkyl, then $q^*m \geq 2$; and if *Y* is a C_{16} to C_{20} alkyl, then $q^*m \geq 1$;
- and, when $p \geq 2$, if *Y* is a C_8 to C_9 alkyl, then $q^*m \geq 2$ and, if *Y* is a C_{10} to C_{16} alkyl, then $q^*m \geq 0.2$;



25 in which:

- *R* is $-OH$ or a $-(f-[A]-COOH)_n$ radical:
 - $-A-$ is a linear or branched radical comprising from 1 to 4 carbon atoms; the said radical $-A-$:
 - being bonded to a glucoside unit *via* a functional group *f* chosen from the group consisting of ether, ester or carbamate functional groups;

30

- n represents the degree of substitution of the glucoside units by $-f-[A]-COOH$ and $0.1 \leq n \leq 2$;
- R' is chosen from the group consisting of the radicals:

5 ○ $-C(O)NH-[E]-(o-[F])_t$;

○ $-CH_2N(L)_z-[E]-(o-[F])_t$;

in which:

- z is a positive integer equal to 1 or 2,
- L is chosen from the group consisting of:

10 ▪ $-H$ and z is equal to 1, and/or

▪ $-[A]-COOH$ and z is equal to 1 or 2, if f is an ether functional group,

▪ $-CO-[A]-COOH$ and z is equal to 1 if f is an ester functional group, and

15 ▪ $-CO-NH-[A]-COOH$ and z is equal to 1 if f is a carbamate functional group;

○ $-[E]-(o-[F])_t$:

- $-E-$ is a linear or branched, at least divalent, radical comprising from 1 to 8 carbon atoms and optionally comprising heteroatoms, such as O, N or S;
- $-F-$ is a linear or cyclic alkyl group, an alkylaryl or an arylalkyl, of 12 to 30 carbon atoms, optionally substituted by one or more C_1 to C_3 alkyl groups;
- o is a functional group chosen from the group consisting of ether, ester, amide or carbamate functional groups;
- t is a positive integer equal to 1 or 2;

20 • q represents the degree of polymerization as glucoside units, that is to say the mean number of glucoside units per polysaccharide chain, and $3 \leq q \leq 50$;

25 • the free acid functional groups being in the form of salts of alkali metal cations chosen from the group consisting of Na^+ and K^+ ;

30 • when $z = 2$, the nitrogen atom is in the form of a quaternary ammonium.

[00029] In one embodiment, when $p = 1$, if Y is a C_{21} to C_{30} group, then $q*m \geq 1$.

[00030] In one embodiment, when $p = 1$, if Y is a C_{21} to C_{30} group, then $q*m \geq 0.1$.

[00031] In one embodiment, the $-(f-[A]-COOH)_n$ radical is such that:

- $-A-$ is a radical comprising one carbon atom; the said $-A-$ radical being bonded to a glucoside unit *via* a ether functional group f .


5 [00032] In one embodiment, the $-(g-[B]-k-[D])_m$ radical is such that:

- $-B-$ is a radical comprising one carbon atom; the said $-B-$ radical being bonded to a glucoside unit *via* an ether functional group g , and
- X is a radical resulting from an amino acid.

[00033] In one embodiment, the $-(f-[A]-COOH)_n$ radical is such that:

- $-A-$ is a radical comprising one carbon atom; the said $-A-$ radical being bonded to a glucoside unit *via* an ether functional group f , and
- the $-(g-[B]-k-[D])_m$ radical is such that:
- $-B-$ is a radical comprising one carbon atom; the said $-B-$ radical being bonded to a glucoside unit *via* an ether functional group g , and
- X is a radical resulting from an amino acid,
- k is an amide functional group.

[00034] In one embodiment, the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is of formula III:




20 Formula III

in which:

- R is $-OH$ or chosen from the group consisting of the radicals:
 - $-(f-[A]-COOH)_n$;
 - $-(g-[B]-k-[D])_m$, D comprising at least one alkyl chain comprising at least 8 carbon atoms;
- n represents the degree of substitution of the glucoside units by $-f-[A]-COOH$ and $0.1 \leq n \leq 2$;
- m represents the degree of substitution of the glucoside units by $-g-[B]-k-[D]$ and $0 < m \leq 0.5$;

- q represents the degree of polymerization as glucoside units, that is to say the mean number of glucoside units per polysaccharide chain, and $3 \leq q \leq 50$;
- 5 • $-(f-[A]-COOH)_n$:
 - $-A-$ is a linear or branched radical comprising from 1 to 4 carbon atoms; the said $-A-$ radical:
 - being bonded to a glucoside unit *via* a functional group *f* chosen from the group consisting of ether, ester and carbamate functional groups;
- 10 • $-(g-[B]-k-[D])_m$:
 - $-B-$ is a linear or branched, at least divalent, radical comprising from 1 to 4 carbon atoms; the said $-B-$ radical:
 - being bonded to a glucoside unit *via* a functional group *g* chosen from the group consisting of ether, ester and carbamate functional groups;
 - 15 ◦ being bonded to a $-D$ radical *via* a functional group *k*; *k* chosen from the group consisting of ester, amide and carbamate functional groups; the said $-D$ radical:
 - being an $-X(-I-Y)_p$ radical, *X* being an at least divalent radical comprising from 1 to 12 atoms chosen from the group consisting of C, N and O atoms, optionally carrying carboxyl or amine functional groups and/or resulting from an amino acid, a dialcohol, a diamine or a mono- or polyethylene glycol mono- or diamine; *Y* being a linear or cyclic alkyl group, an alkylaryl or an arylalkyl, of 8 to 20 carbon atoms, optionally substituted by one or more C_1 to C_3 alkyl groups; *p* ≥ 1 and *I* a functional group chosen from the group consisting of ester, amide and carbamate functional groups;
- 20 • *f, g and k* being identical or different;
- 30 • the free acid functional groups being in the form of salts of alkali metal cations chosen from the group consisting of Na^+ and K^+ ;
- and, when *p* = 1, if *Y* is a C_8 to C_{14} alkyl, then $q*m \geq 2$, if *Y* is a C_{15} alkyl, then $q*m \geq 2$; and if *Y* is a C_{16} to C_{20} alkyl, then $q*m \geq 1$;
- and, when *p* ≥ 2 , if *Y* is a C_8 to C_{11} alkyl, then $q*m \geq 2$ and, if *Y* is a C_{12} to C_{16} alkyl, then $q*m \geq 0.3$.

[00035] In one embodiment, the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is of formula IV:



in which:

- 5 • R is $-\text{OH}$ or chosen from the group consisting of the radicals:
 - $-(f\text{-}[A]\text{-COOH})_n$;
 - $-(g\text{-}[B]\text{-}k\text{-}[D])_m$, D comprising at least one alkyl chain comprising at least 8 carbon atoms;
- 10 • n represents the degree of substitution of the hydroxyl $-\text{OH}$ functional groups by $-f\text{-}[A]\text{-COOH}$ per glucoside unit; and $0.1 \leq n \leq 2$;
- 15 • m represents the degree of substitution of the hydroxyl $-\text{OH}$ functional groups by $-g\text{-}[B]\text{-}k\text{-}[D]$ per glucoside unit; and $0 < m \leq 0.5$;
- 20 • q represents the degree of polymerization as glucoside units, that is to say the mean number of glucoside units per polysaccharide chain, and $3 \leq q \leq 50$;
- 25 • $-(f\text{-}[A]\text{-COOH})_n$:
 - $-A-$ is a linear or branched radical comprising from 1 to 4 carbon atoms; the said $-A-$ radical:
 - being bonded to a glucoside unit *via* a functional group *f* chosen from the group consisting of ether, ester and carbamate functional groups;
- 30 • $-(g\text{-}[B]\text{-}k\text{-}[D])_m$:
 - $-B-$ is a linear or branched, at least divalent, radical comprising from 1 to 4 carbon atoms; the said $-B-$ radical:
 - being bonded to a glucoside unit *via* a functional group *g* chosen from the group consisting of ether, ester and carbamate functional groups;
 - being bonded to a $-D$ radical *via* a functional group *k*; *k* chosen from the group consisting of ester, amide and carbamate functional groups; the said $-D$ radical:
 - being an $-X(-I\text{-}Y)_p$ radical, X being an at least divalent radical comprising from 1 to 12 atoms chosen from the group consisting of C, N and O atoms, optionally carrying carboxyl or amine functional groups and/or resulting from an amino acid, a dialcohol, a diamine or a mono- or polyethylene glycol mono- or

5 diamine; Y being a linear or cyclic alkyl group, an alkylaryl or an arylalkyl, of 8 to 30 carbon atoms, optionally substituted by one or more C₁ to C₃ alkyl groups; p ≥ 1 and / a functional group chosen from the group consisting of ester, amide and carbamate functional groups;

10

- f, g and k being identical or different;
- the free acid functional groups being in the form of salts of alkali metal cations chosen from the group consisting of Na⁺ and K⁺;
- and, when p = 1, if Y is a C₈ to C₁₄ alkyl, then q*m ≥ 2, if Y is a C₁₅ alkyl, then q*m ≥ 2; and if Y is a C₁₆ to C₃₀ alkyl, then q*m ≥ 1;
- and, when p ≥ 2, if Y is a C₈ to C₉ alkyl, then q*m ≥ 2 and, if Y is a C₁₀ to C₁₆ alkyl, then q*m ≥ 0.2.

15 [00036] The structure drawn corresponds to the representation commonly used to represent dextran, which is a polysaccharide predominantly composed of (1,6) linkages between glucoside units, which is the representation adopted. Dextran also comprises (1,3) linkages at approximately 5% in general, which are deliberately not represented but which are, of course, included within the scope of the invention.

20 [00037] The term "basal insulin" whose isoelectric point is between 5.8 and 8.5 is understood to mean an insulin which is insoluble at pH 7 and which has a duration of action of between 8 and 24 hours or more in the standard models of diabetes.

25 [00038] These basal insulins whose isoelectric point is between 5.8 and 8.5 are recombinant insulins, the primary structure of which has been modified mainly by introduction of basic amino acids, such as arginine or lysine. They are described, for example, in the following patents, patent applications or publications: WO 2003/053339, WO 2004/096854, US 5 656 722 and US 6 100 376.

[00039] In one embodiment, the basal insulin whose isoelectric point is between 5.8 and 8.5 is insulin glargine.

30 [00040] In one embodiment, the compositions according to the invention comprise 100 IU/ml (i.e., approximately 3.6 mg/ml) of basal insulin whose isoelectric point is between 5.8 and 8.5.

[00041] In one embodiment, the compositions according to the invention comprise 40 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5.

[00042] In one embodiment, the compositions according to the invention comprise 200 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5.

5 [00043] In one embodiment, the compositions according to the invention comprise 300 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5.

[00044] In one embodiment, the compositions according to the invention comprise 400 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5.

[00045] In one embodiment, the compositions according to the invention comprise

10 500 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5.

[00046] In one embodiment, the ratio by weight of the basal insulin whose isoelectric point is between 5.8 and 8.5 to the substituted dextran, i.e. substituted dextran/basal insulin, is between 0.2 and 5.

15 [00047] In one embodiment, the ratio by weight of the basal insulin whose isoelectric point is between 5.8 and 8.5 to the substituted dextran, i.e. substituted dextran/basal insulin, is between 0.2 and 4.

[00048] In one embodiment, the ratio by weight of the basal insulin whose isoelectric point is between 5.8 and 8.5 to the substituted dextran, i.e. substituted dextran/basal

20 insulin, is between 0.2 and 3.

[00049] In one embodiment, the ratio by weight of the basal insulin whose isoelectric point is between 5.8 and 8.5 to the substituted dextran, i.e. substituted dextran/basal insulin, is between 0.5 and 3.

25 [00050] In one embodiment, the ratio by weight of the basal insulin whose isoelectric point is between 5.8 and 8.5 to the substituted dextran, i.e. substituted dextran/basal insulin, is between 0.8 and 3.

[00051] In one embodiment, the ratio by weight of the basal insulin whose isoelectric point is between 5.8 and 8.5 to the substituted dextran, i.e. substituted dextran/basal insulin, is between 1 and 3.

30 [00052] In one embodiment, the concentration of substituted dextran is between 1 and 100 mg/ml.

[00053] In one embodiment, the concentration of substituted dextran is between 1 and 80 mg/ml.

[00054] In one embodiment, the concentration of substituted dextran is between 1 and 60 mg/ml.

[00055] In one embodiment, the concentration of substituted dextran is between 1 and 50 mg/ml.

5 [00056] In one embodiment, the concentration of substituted dextran is between 1 and 30 mg/ml.

[00057] In one embodiment, the concentration of substituted dextran is between 1 and 20 mg/ml.

10 [00058] In one embodiment, the concentration of substituted dextran is between 1 and 10 mg/ml.

[00059] In one embodiment, the concentration of polysaccharide is between 5 and 20 mg/ml.

[00060] In one embodiment, the concentration of polysaccharide is between 5 and 10 mg/ml.

15 [00061]

[00062] In one embodiment, the compositions according to the invention additionally comprise a prandial insulin. Prandial insulins are soluble at pH 7.

20 [00063] The term "prandial insulin" is understood to mean a "rapid" or "regular" insulin.

[00064] "Rapid" prandial insulins are insulins which must meet the needs brought about by the ingestion of proteins and carbohydrates during a meal; they have to act in less than 30 minutes.

25 [00065] In one embodiment, "regular" prandial insulins are chosen from the group consisting of Humulin® (human insulin) and Novolin® (human insulin).

[00066] "Fast-acting" prandial insulins are insulins which are obtained by recombination and which are modified in order to reduce their action time.

30 [00067] In one embodiment, "fast-acting" prandial insulins are chosen from the group consisting of insulin lispro (Humalog®), insulin glulisine (Apidra®) and insulin aspart (NovoLog®).

[00068] In one embodiment, the compositions according to the invention comprise, in total, 100 IU/ml (i.e., approximately 3.6 mg/ml) of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

[00069] In one embodiment, the compositions according to the invention comprise, in total, 40 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

5 [00070] In one embodiment, the compositions according to the invention comprise, in total, 200 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

[00071] In one embodiment, the compositions according to the invention comprise, in total, 300 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

10 [00072] In one embodiment, the compositions according to the invention comprise, in total, 400 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

15 [00073] In one embodiment, the compositions according to the invention comprise, in total, 500 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

[00074] In one embodiment, the compositions according to the invention comprise, in total, 600 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

20 [00075] In one embodiment, the compositions according to the invention comprise, in total, 700 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

[00076] In one embodiment, the compositions according to the invention comprise, in total, 800 IU/ml of insulin with a combination of prandial insulin and basal insulin whose isoelectric point is between 5.8 and 8.5.

25

[00077] The proportions between the basal insulin whose isoelectric point is between 5.8 and 8.5 and the prandial insulin, expressed as percentage with respect to the total amount of insulin, are, for example, 25/75, 30/70, 40/60, 50/50, 60/40, 70/30, 80/20 and 90/10 for the formulations as described above from 40 to 800 IU/ml. However, 30 any other proportion can be produced.

[00078] For a formulation comprising 100 IU/ml as total insulin, the proportions between the basal insulin whose isoelectric point is between 5.8 and 8.5 and the prandial insulin are, for example, in IU/ml, 25/75, 30/70, 40/60, 50/50, 60/40, 70/30, 80/20 or 90/10. However, any other proportion can be produced.

35

[00079] In one embodiment, the composition according to the invention additionally comprises a GLP-1, a GLP-1 analogue or a GLP-1 derivative.

[00080] In one embodiment, the GLP-1 analogues or derivatives are chosen from the group consisting of exenatide or Byetta®, developed by Eli Lilly & Co and Amylin Pharmaceuticals, liraglutide or Victoza® developed by Novo Nordisk, or lixisenatide or Lyxumia® developed by Sanofi, their analogues or derivatives and their pharmaceutically acceptable salts.

[00081] In one embodiment, the GLP-1 analogue or derivative is exenatide or Byetta®, its analogues or derivatives and their pharmaceutically acceptable salts.

10 [00082] In one embodiment, the GLP-1 analogue or derivative is liraglutide or Victoza®, its analogues or derivatives and their pharmaceutically acceptable salts.

[00083] In one embodiment, the GLP-1 analogue or derivative is lixisenatide or Lyxumia®, its analogues or derivatives and their pharmaceutically acceptable salts.

[00084] The term "analogue" is understood to mean, when it is used with reference to a peptide or a protein, a peptide or a protein in which one or more constituent amino acid residues have been replaced by other amino acid residues and/or from which one or more constituent amino acid residues have been deleted and/or to which one or more constituent amino acid residues have been added. The percentage of homology accepted for the present definition of an analogue is 50%.

15 [00085] The term "derivative" is understood to mean, when it is used with reference to a peptide or protein, a peptide or a protein or an analogue chemically modified by a substituent which is not present in the reference peptide or protein or analogue, that is to say a peptide or a protein which has been modified by creation of covalent bonds, in order to introduce substituents.

20 [00086] In one embodiment, the concentration of GLP-1, of GLP-1 analogue or of GLP-1 derivative is within a range from 0.01 to 10 mg/ml.

[00087] In one embodiment, the concentration of exenatide, its analogues or derivatives and their pharmaceutically acceptable salts is within a range from 0.05 to 0.5 mg/ml.

25 [00088] In one embodiment, the concentration of liraglutide, its analogues or derivatives and their pharmaceutically acceptable salts is within a range from 1 to 10 mg/ml.

[00089] In one embodiment, the concentration of lixisenatide, its analogues or derivatives and their pharmaceutically acceptable salts is within a range from 0.01 to 1 mg/ml.

[00090] In one embodiment, the compositions according to the invention are produced by mixing commercial solutions of basal insulin whose isoelectric point is between 5.8 and 8.5 and commercial solutions of GLP-1, of GLP-1 analogue or of GLP-1 derivative in ratios by volume within a range from 10/90 to 90/10.

5

[00091] In one embodiment, the composition according to the invention comprises a daily dose of basal insulin and a daily dose of GLP-1, GLP-1 analogue or GLP-1 derivative.

[00092] In one embodiment, the compositions according to the invention comprise
10 500 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 to 0.5 mg/ml of exenatide.

[00093] In one embodiment, the compositions according to the invention comprise 500 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 1 to 10 mg/ml of liraglutide.

15 [00094] In one embodiment, the compositions according to the invention comprise 500 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 to 0.5 mg/ml of lixisenatide.

[00095] In one embodiment, the compositions according to the invention comprise
20 100 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 to 0.5 mg/ml of exenatide.

[00096] In one embodiment, the compositions according to the invention comprise 100 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 1 to 10 mg/ml of liraglutide.

[00097] In one embodiment, the compositions according to the invention comprise
25 100 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 to 0.5 mg/ml of lixisenatide.

[00098] In one embodiment, the compositions according to the invention comprise 40 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 à 0.5 mg/ml of exenatide.

30 [00099] In one embodiment, the compositions according to the invention comprise 40 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 1 to 10 mg/ml of liraglutide.

[000100] In one embodiment, the compositions according to the invention comprise 40 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 à 0.5 mg/ml of lixisenatide.

35

[000101] In one embodiment, the compositions according to the invention comprise 200 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 to 0.5 mg/ml of exenatide.

5 [000102] In one embodiment, the compositions according to the invention comprise 200 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 1 to 10 mg/ml of liraglutide.

[000103] In one embodiment, the compositions according to the invention comprise 200 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5 and from 0.05 to 0.5 mg/ml of lixisenatide.

10

[000104] In one embodiment, the compositions according to the invention additionally comprise zinc salts at a concentration of between 0 and 5000 μ M.

[000105] In one embodiment, the compositions according to the invention additionally comprise zinc salts at a concentration of between 50 and 4000 μ M.

15 [000106] In one embodiment, the compositions according to the invention additionally comprise zinc salts at a concentration of between 200 and 3000 μ M.

[000107] In one embodiment, the compositions according to the invention additionally comprise zinc salts at a concentration of between 0 and 1000 μ M.

20 [000108] In one embodiment, the compositions according to the invention additionally comprise zinc salts at a concentration of between 20 and 600 μ M.

[000109] In one embodiment, the compositions according to the invention additionally comprise zinc salts at a concentration of between 50 and 500 μ M.

25 [000110] In one embodiment, the compositions according to the invention comprise buffers chosen from the group consisting of Tris, citrates and phosphates at concentrations of between 0 and 100 mM, preferably between 0 and 50 mM or between 15 and 50 mM.

[000111] In one embodiment, the compositions according to the invention additionally comprise preservatives.

30 [000112] In one embodiment, the preservatives are chosen from the group consisting of m-cresol and phenol, alone or as a mixture.

[000113] In one embodiment, the concentration of the preservatives is between 10 and 50 mM.

[000114] In one embodiment, the concentration of the preservatives is between 10 and 40 mM.

[000115] The compositions according to the invention can additionally comprise additives, such as tonicity agents, such as glycerol, NaCl, mannitol and glycine.

5

[000116] The compositions according to the invention can additionally comprise additives in accordance with the pharmacopoeias, such as surfactants, for example polysorbate.

10 [000117] The compositions according to the invention can additionally comprise all the excipients in accordance with the pharmacopoeias which are compatible with the insulins used at the concentrations of use.

[000118] In one embodiment, $0.3 \leq n \leq 1.7$.

[000119] In one embodiment, $0.7 \leq n \leq 1.5$.

15 [000120] In one embodiment, $0.9 \leq n \leq 1.2$.

[000121] In one embodiment, $0.01 \leq m \leq 0.5$.

[000122] In one embodiment, $0.02 \leq m \leq 0.4$.

[000123] In one embodiment, $0.03 \leq m \leq 0.3$.

20 [000124] In one embodiment, $0.05 \leq m \leq 0.2$.

[000125] In one embodiment, $3 \leq q \leq 50$.



[000126] In one embodiment, $3 \leq q \leq 40$.

[000127] In one embodiment, $3 \leq q \leq 30$.

25 [000128] In one embodiment, $3 \leq q \leq 20$.

[000129] In one embodiment, $3 \leq q \leq 10$.

[000130] In one embodiment, the $-(f-[A]-COOH)_n$ radical is chosen from the group consisting of the following sequences, f having the meaning given above:



5 [000131] In one embodiment, the $-(g-[B]-k-[D])_m$ radical is chosen from the group consisting of the following sequences, g , k and D having the meanings given above:



[000132] In one embodiment, D is such that the X radical is an at least divalent radical resulting from an amino acid.

10 [000133] In one embodiment, D is such that the X radical is an at least divalent radical resulting from an amino acid chosen from the group consisting of glycine, leucine, phenylalanine, lysine, isoleucine, alanine, valine, aspartic acid and glutamic acid.

15 [000134] The radicals resulting from amino acids can be either laevorotatory or dextrorotatory.

[000135] In one embodiment, D is such that the X radical is an at least divalent radical resulting from a mono- or polyethylene glycol.

[000136] In one embodiment, D is such that the X radical is an at least divalent radical resulting from ethylene glycol.

20 [000137] In one embodiment, D is such that the X radical is an at least divalent radical resulting from a polyethylene glycol chosen from the group consisting of diethylene glycol and triethylene glycol.

[000138] In one embodiment, D is such that the X radical is an at least divalent radical resulting from a mono- or polyethylene glycol amine.

25 [000139] In one embodiment, D is such that the X radical is an at least divalent radical resulting from a mono- or polyethylene glycol amine chosen from the group consisting of ethanolamine, diethylene glycol amine and triethylene glycol amine.

[000140] In one embodiment, D is such that the X radical is an at least divalent radical resulting from a mono- or polyethylene glycol diamine.

30 [000141] In one embodiment, D is such that the X radical is an at least divalent radical resulting from ethylenediamine.

[000142] In one embodiment, D is such that the X radical is an at least divalent radical resulting from a mono- or polyethylene glycol diamine chosen from the group consisting of diethylene glycol diamine and triethylene glycol diamine.

5 [000143] In one embodiment, D is such that the Y group is an alkyl group resulting from a hydrophobic alcohol.

[000144] In one embodiment, D is such that the Y group is an alkyl group resulting from a hydrophobic alcohol chosen from the group consisting of octanol (capryl alcohol), 3,7-dimethyloctan-1-ol, decanol (decyl alcohol), dodecanol (lauryl alcohol), tetradecanol (myristyl alcohol) and hexadecanol (cetyl alcohol).

10 [000145] In one embodiment, D is such that the Y group is an alkyl group resulting from a hydrophobic acid.

[000146] In one embodiment, D is such that the Y group is an alkyl group resulting from a hydrophobic acid chosen from the group consisting of decanoic acid, dodecanoic acid, tetradecanoic acid and hexadecanoic acid.

15 [000147] In one embodiment, D is such that the Y group is a group resulting from a sterol.

[000148] In one embodiment, D is such that the Y group is a group resulting from a sterol chosen from the group consisting of cholesterol and its derivatives.

20 [000149] In one embodiment, D is such that the Y group is a group resulting from a tocopherol.

[000150] In one embodiment, D is such that the Y group is a group resulting from a tocopherol derivative chosen from the racemate, the L isomer or the D isomer of α -tocopherol.

25 [000151] In one embodiment, D is such that the X radical results from glycine, p = 1, the Y group results from octanol and the functional group / is an ester functional group.

[000152] In one embodiment, D is such that the X radical results from glycine, p = 1, the Y group results from dodecanol and the functional group / is an ester functional group.

30 [000153] In one embodiment, D is such that the X radical results from glycine, p = 1, the Y group results from hexadecanol and the functional group / is an ester functional group.

[000154] In one embodiment, D is such that the X radical results from phenylalanine, p = 1, the Y group results from octanol and the functional group / is an ester functional group.

5 [000155] In one embodiment, D is such that the X radical results from phenylalanine, p = 1, the Y group results from 3,7-dimethyloctan-1-ol and the functional group / is an ester functional group.

[000156] In one embodiment, D is such that the X radical results from aspartic acid, p = 2, the Y groups result from octanol and the functional groups / are ester functional groups.

10 [000157] In one embodiment, D is such that the X radical results from aspartic acid, p = 2, the Y groups result from decanol and the functional groups / are ester functional groups.

15 [000158] In one embodiment, D is such that the X radical results from aspartic acid, p = 2, the Y groups result from dodecanol and the functional groups / are ester functional groups.

[000159] In one embodiment, D is such that the X radical results from ethylenediamine, the Y group results from dodecanoic acid and the functional group / is an amide functional group.

20 [000160] In one embodiment, D is such that the X radical results from diethylene glycol amine, p = 1, the Y group results from dodecanoic acid and the functional group / is an ester functional group.

[000161] In one embodiment, D is such that the X radical results from triethylene glycol diamine, p = 1, the Y group results from dodecanoic acid and the functional group / is an amide functional group.

25 [000162] In one embodiment, D is such that the X radical results from triethylene glycol diamine, p = 1, the Y group results from hexadecanoic acid and the functional group / is an amide functional group.

30 [000163] In one embodiment, D is such that the X radical results from leucine, p = 1, the Y group results from cholesterol and the functional group / is an ester functional group.

[000164] In one embodiment, D is such that X results from ethylenediamine, p = 1, the Y group results from cholesterol and the functional group / is a carbamate functional group.

[000165] In one embodiment, the E radical is an at least divalent radical resulting from an amino acid chosen from the group consisting of glycine, leucine, phenylalanine, lysine, isoleucine, alanine, valine, serine, threonine, aspartic acid and glutamic acid.

5 [000166] The radicals resulting from amino acids can be either laevorotatory or dextrorotatory.

[000167] In one embodiment, the E radical is an at least divalent radical resulting from a mono- or polyethylene glycol amine.

10 [000168] In one embodiment, the E radical is an at least divalent radical resulting from a mono- or polyethylene glycol amine chosen from the group consisting of ethanolamine, diethylene glycol amine and triethylene glycol amine.

[000169] In one embodiment, the E radical is an at least divalent radical resulting from a mono- or polyethylene glycol diamine.

15 [000170] In one embodiment, the E radical is an at least divalent radical resulting from ethylenediamine.

[000171] In one embodiment, the E radical is an at least divalent radical resulting from a mono- or polyethylene glycol diamine chosen from the group consisting of diethylene glycol diamine and triethylene glycol diamine.

20 [000172] In one embodiment, the F group is an alkyl group resulting from a hydrophobic alcohol.

[000173] In one embodiment, the F group is a group resulting from a hydrophobic alcohol chosen from the group consisting of dodecanol (lauryl alcohol), tetradecanol (myristyl alcohol) and hexadecanol (cetyl alcohol).

25 [000174] In one embodiment, the F group is a group resulting from a hydrophobic acid.

[000175] In one embodiment, the F group is a group resulting from a hydrophobic acid chosen from the group consisting of dodecanoic acid, tetradecanoic acid and hexadecanoic acid.

[000176] In one embodiment, the F group is a group resulting from a sterol.

30 [000177] In one embodiment, the F group is a group resulting from a sterol chosen from the group consisting of cholesterol and its derivatives.

[000178] In one embodiment, the F group is a group resulting from a tocopherol.

[000179] In one embodiment, the F group is a group resulting from a tocopherol derivative chosen from the racemate, the L isomer or the D isomer of α -tocopherol.

[000180] In one embodiment, the E radical results from ethylenediamine, $t = 1$, o is a carbamate functional group and the F group results from cholesterol.

[000181] In one embodiment:

5 o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;

o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from glycine, l is an ester functional group and Y results from octanol;

10 o $q = 38$, $n = 0.9$ and $m = 0.2$.

[000182] In one embodiment:

o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;

o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from glycine, $p = 1$, l is an ester functional group and Y results from hexadecanol;

15 o $q = 19$, $n = 1.0$ and $m = 0.1$.

[000183] In one embodiment:

20 o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;

o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from phenylalanine, $p = 1$, l is an ester functional group and Y results from octanol;

25 o $q = 38$, $n = 1.0$ and $m = 0.1$.

[000184] In one embodiment:

o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;

30 o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from phenylalanine, $p = 1$, l is an ester functional group and Y results from octanol;

o $q = 19$, $n = 1.0$ and $m = 0.2$.

[000185] In one embodiment:

- o $-(f-[A]-COOH)n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from phenylalanine, p = 1, I is an ester functional group and Y results from 3,7-dimethyloctan-1-ol;
- o q = 38, n = 1.0 and m = 0.1.

[000186] In one embodiment:

- o $-(f-[A]-COOH)n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, p = 2, I are ester functional groups and Y result from octanol;
- o q = 38, n = 1.05 and m = 0.05.

[000187] In one embodiment:

- o $-(f-[A]-COOH)n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, p = 2, I are ester functional groups and Y result from decanol;
- o q = 38, n = 1.05 and m = 0.05.

[000188] In one embodiment:

- o $-(f-[A]-COOH)n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, p = 2, I are ester functional groups and Y result from dodecanol;
- o q = 19, n = 1.05 and m = 0.05.

[000189] In one embodiment:

- o $-(f-[A]-COOH)n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from

ethylenediamine, $p = 1$, l is an amide functional group and Y results from dodecanoic acid;

- $q = 38$, $n = 1.0$ and $m = 0.1$.

[000190] In one embodiment:

- 5 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-CH_2-$ radical and f is an ester functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ester functional group, B is the $-CH_2-CH_2-$ radical, k is an amide functional group and D is such that X results from glycine, $p = 1$, l is an ester functional group and Y results from dodecanol;
- 10 ○ $q = 38$, $n = 1.3$ and $m = 0.1$.

[000191] In one embodiment:

- 15 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is a carbamate functional group;
- $-(g-[B]-k-[D])_m$ is such that g is a carbamate functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, $p = 2$, l are ester functional groups and Y result from octanol;
- $q = 38$, $n = 1.3$ and $m = 0.1$.

20 [000192] In one embodiment:

- $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, $p = 2$, l are ester functional groups and Y result from dodecanol;
- 25 ○ $q = 4$, $n = 0.96$ and $m = 0.07$.

[000193] In one embodiment:

- 30 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from diethylene glycol amine, $p = 1$, l is an ester functional group and Y results from dodecanoic acid;
- 35 ○ $q = 38$, $n = 1.0$ and $m = 0.1$.

[000194] In one embodiment:

- $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from triethylene glycol diamine, p = 1, l is an amide functional group and Y results from dodecanoic acid;
- q = 38, n = 1.0 and m = 0.1.

[000195] In one embodiment:

- 10 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from triethylene glycol diamine, p = 1, l is an amide functional group and Y results from hexadecanoic acid;
- 15 ○ q = 38, n = 1.05 and m = 0.05.

[000196] In one embodiment:

- 20 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from glycine, p = 1, l is an ester functional group and Y results from hexadecanol;
- q = 19, n = 1.05 and m = 0.05.

25 [000197] In one embodiment:

- $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from glycine, p = 1, l is an ester functional group and Y results from hexadecanol;
- 30 ○ q = 38, n = 0.37 and m = 0.05.

[000198] In one embodiment:

- 35 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;

- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-\text{CH}_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- $q = 19$, $n = 1.61$ and $m = 0.04$.

5 [000199] In one embodiment:

- $-(f-[A]-\text{COOH})_n$ is such that A is the $-\text{CH}_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-\text{CH}_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- $q = 19$, $n = 1.06$ and $m = 0.04$.

[000200] In one embodiment:

- $-(f-[A]-\text{COOH})_n$ is such that A is the $-\text{CH}_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-\text{CH}_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- $q = 19$, $n = 0.66$ and $m = 0.04$.

[000201] In one embodiment:

- $-(f-[A]-\text{COOH})_n$ is such that A is the $-\text{CH}_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-\text{CH}_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- $q = 19$, $n = 0.46$ and $m = 0.04$.

[000202] In one embodiment:

- $-(f-[A]-\text{COOH})_n$ is such that A is the $-\text{CH}_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-\text{CH}_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- $q = 4$, $n = 1.61$ and $m = 0.05$.

[000203] In one embodiment:

- $-(f-[A]-\text{COOH})_n$ is such that A is the $-\text{CH}_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-\text{CH}_2-$ radical, k is an amide functional group and D is such that X results from

ethylenediamine, $p = 1$, l is a carbamate functional group and Y results from cholesterol;

- o $q = 19$, $n = 1.61$ and $m = 0.04$.

[000204] In one embodiment:

- 5 o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is a carbamate functional group;
- o $-(g-[B]-k-[D])_m$ is such that g is a carbamate functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- 10 o $q = 19$, $n = 1.96$ and $m = 0.04$.

[000205] In one embodiment:

- o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- 15 o $-[E]-(o-[F])_t$ is such that E results from ethylenediamine, o is a carbamate functional group and F results from cholesterol;
- o $q = 19$ and $n = 1.65$.

[000206] In one embodiment:

- 20 o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from leucine, $p = 1$, l is an ester functional group and Y results from cholesterol;
- o $q = 38$, $n = 0.99$ and $m = 0.05$.

25 [000207] In one embodiment:

- o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, $p = 2$, l are ester functional groups and Y result from dodecanol;
- 30 o $q = 4$, $n = 1.41$ and $m = 0.16$.

[000208] In one embodiment:

- 35 o $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- o $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from

aspartic acid, $p = 2$, l are ester functional groups and Y result from dodecanol;

- $q = 4$, $n = 1.50$ and $m = 0.07$.

[000209] In one embodiment:

- 5 ○ $-(f-[A]-COOH)_n$ is such that A is the $-CH_2-$ radical and f is an ether functional group;
- $-(g-[B]-k-[D])_m$ is such that g is an ether functional group, B is the $-CH_2-$ radical, k is an amide functional group and D is such that X results from aspartic acid, $p = 2$, l are ester functional groups and Y result from decanol;
- 10 ○ $q = 4$, $n = 1.05$ and $m = 0.05$.

[000210] In one embodiment, the compositions according to the invention comprise a dextran chosen from the group consisting of the following dextrans of formula I, III or IV:

- 15 - Sodium dextranmethylcarboxylate modified by octyl glycinate,
- Sodium dextranmethylcarboxylate modified by cetyl glycinate,
- Sodium dextranmethylcarboxylate modified by octyl phenylalaninate,
- Sodium dextranmethylcarboxylate modified by 3,7-dimethyl-1-octyl phenylalaninate,
- 20 - Sodium dextranmethylcarboxylate modified by dioctyl aspartate,
- Sodium dextranmethylcarboxylate modified by didecyl aspartate,
- Sodium dextranmethylcarboxylate modified by dilauryl aspartate,
- Sodium dextranmethylcarboxylate modified by N-(2-aminoethyl)dodecanamide,
- 25 - Sodium dextransuccinate modified by lauryl glycinate,
- N-(sodium methylcarboxylate) dextran carbamate modified by dioctyl aspartate,
- Sodium dextranmethylcarboxylate modified by dilauryl aspartate,
- Sodium dextranmethylcarboxylate modified by 2-(2-aminoethoxy)ethyl dodecanoate,
- 30 - Sodium dextranmethylcarboxylate modified by 2-(2-{2-[dodecanoylamino]ethoxy}ethoxy)ethylamine,

- Sodium dextranmethylcarboxylate modified by 2-(2-[2-[hexadecanoylamino]ethoxy]ethoxy)ethylamine,
- Sodium dextranmethylcarboxylate modified by cholesteryl leucinate,
- Sodium dextranmethylcarboxylate modified by cholesteryl 5 1-ethylenediaminecarboxylate,
- N-(sodium methylcarboxylate) dextran carbamate modified by cholesteryl leucinate.

[000211] In one embodiment, the compositions according to the invention comprise a dextran chosen from the group consisting of the dextran of the following formula II:

10 - Sodium dextranmethylcarboxylate modified by cholesteryl 1-ethylenediaminecarboxylate grafted by reductive amination to the reducing chain end.

[000212] The invention also relates to single-dose formulations at a pH of between 6.6 and 7.8 comprising a basal insulin whose isoelectric point is between 5.8 and 8.5 and 15 a prandial insulin.

[000213] The invention also relates to single-dose formulations at a pH of between 7 and 7.8 comprising a basal insulin whose isoelectric point is between 5.8 and 8.5 and a prandial insulin.

20 [000214] In one embodiment, the basal insulin whose isoelectric point is between 5.8 and 8.5 is insulin glargine.

[000215] In one embodiment, the prandial insulin is chosen from the group consisting of Humulin® (human insulin) and Novolin® (human insulin).

[000216] In one embodiment, the prandial insulin is chosen from the group consisting of insulin lispro (Humalog®), insulin glulisine (Apidra®) and insulin aspart (NovoLog®).

25 [000217] The dissolution at a pH of between 6.6 and 7.8 of the basal insulins whose isoelectric point is between 5.8 and 8.5 by the polysaccharides of formula I, II, III or IV can be observed and controlled in a simple way, with the naked eye, by virtue of a change in appearance of the solution

30 [000218] The dissolution at a pH of between 7 and 7.8 of the basal insulins whose isoelectric point is between 5.8 and 8.5 by the polysaccharides of formula I, II, III or IV can be observed and controlled in a simple way, with the naked eye, by virtue of a change in appearance of the solution.

35 [000219] Furthermore and just as importantly, the Applicant Company has been able to confirm that a basal insulin whose isoelectric point is between 5.8 and 8.5,

dissolved in the presence of a polysaccharide of formula I, II, III or IV, had lost none of its slow insulin action.

[000220] The preparation of a composition according to the invention exhibits the advantage of being able to be carried out by simple mixing of an aqueous solution of basal insulin whose isoelectric point is between 5.8 and 8.5, of a solution of prandial insulin and of a polysaccharide of formula I, II, III or IV in aqueous solution or in the lyophilized form. If necessary, the pH of the preparation is adjusted to pH 7.

[000221] The preparation of a composition according to the invention exhibits the advantage of being able to be carried out by simple mixing of an aqueous solution of basal insulin whose isoelectric point is between 5.8 and 8.5, of a polysaccharide of formula I, II, III or IV in aqueous solution or in the lyophilized form, and of a prandial insulin in aqueous solution or in the lyophilized form.

[000222] In one embodiment, the mixture of basal insulin and polysaccharide is concentrated by ultrafiltration before mixing with the prandial insulin in aqueous solution or in the lyophilized form.

[000223] If necessary, the composition of the mixture is adjusted in excipients, such as glycerol, m-cresol, zinc chloride and tween, by addition of concentrated solutions of these excipients to the mixture. If necessary, the pH of the preparation is adjusted to 7.

20

[000224] Description of the figures:

[000225] Figures 1 to 6 present the results obtained in the form of pharmacodynamic curves for glucose. The axis of the ordinates represents the Dglucose (expressed in mM) as a function of the post-injection time (expressed in minutes).

25

[000226] Figure 1: Mean + standard deviation of the mean curves for the sequential administrations of Apidra® and Lantus® (□) in comparison with a composition according to the invention Polysaccharide 4/Lantus®/Apidra® (75/25) (•).

[000227] Figure 2: Apidra® Lantus® individual curves (tested on six pigs).

30

[000228] Figure 3: Polysaccharide 4/Apidra®/Lantus® individual curves (tested on six pigs).

[000229] Figure 4: Mean + standard deviation of the mean curves for the sequential administration of Humalog® and Lantus® (□) in comparison with the administration of a composition according to the invention Polysaccharide 4/Humalog®/Lantus® (•).

[000230] Figure 5: Humalog® Lantus® individual curves (tested on six pigs).

35

[000231] Figure 6: Polysaccharide 4/Humalog®/Lantus® individual curves (tested on five pigs).

[000232] Figures 7 to 12 present the results obtained in the form of pharmacodynamic curves for glucose. The axis of the ordinates represents the Dglucose (expressed in mM) as a function of the post-injection time (expressed in hours).

5 [000233] Figure 7: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) (···), in comparison with a composition according to the invention described in Example B28 (0.53 IU/kg) (—).

10 [000234] Figure 8: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) (···), in comparison with a composition according to the invention described in Example B27 (0.47 IU/kg) (—).

15 [000235] Figure 9: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) (···), in comparison with a composition according to the invention described in Example B29 (0.53 IU/kg) (—).

[000236] Figure 10: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) (···), in comparison with a composition according to the invention described in Example B31 (0.48 IU/kg) (—).

20 [000237] Figure 11: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.24 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) (···), in comparison with a composition according to the invention described in Example B30 (0.64 IU/kg) (—).





25 [000238] Figure 12: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) (···), in comparison with a composition according to the invention described in Example B32 (0.53 IU/kg) (—).







Examples





Part A: Polysaccharides

[000239] Table 1 below presents, in a nonlimiting way, examples of polysaccharides which can be used in the compositions according to the invention.

POLYSACCHARIDES	SUBSTITUENTS -f-A-COONa -g-B-k-D	COMMON NAME
Polysaccharide 1 q: 38 n: 0.9 m: 0.2	 	Sodium dextranmethylcarboxylate modified by octyl glycinate
Polysaccharide 2 q: 19 n: 1.0 m: 0.1 Polysaccharide 16 q: 19 n: 1.05 m: 0.05 Polysaccharide 17 q: 38 n: 0.37 m: 0.05	 	Sodium dextranmethylcarboxylate modified by cetyl glycinate
Polysaccharide 3 q: 38 n: 1.0 m: 0.1 Polysaccharide 4 q: 19 n: 1.0 m: 0.2	 	Sodium dextranmethylcarboxylate modified by octyl phenylalaninate

Polysaccharide 5 q: 38 n: 1.0 m: 0.1		Sodium dextranmethylcarboxylate modified by 3,7-dimethyl-1-octyl phenylalaninate
Polysaccharide 6 q: 38 n: 1.05 m: 0.05		Sodium dextranmethylcarboxylate modified by dioctyl aspartate
Polysaccharide 7 q: 38 n: 1.05 m: 0.05		Sodium dextranmethylcarboxylate modified by didecyl aspartate
Polysaccharide 29		
Polysaccharide 8 q: 19 n: 1.05 m: 0.05		
Polysaccharide 27 q: 4 n: 1.41 m: 0.16		Sodium dextranmethylcarboxylate modified by dilauryl aspartate
Polysaccharide 28 q: 4 n: 1.50 m: 0.07		

Polysaccharide 9 q: 38 n: 1.0 m: 0.1		Sodium dextranmethylcarboxylate modified by N-(2-aminoethyl)dodecanamide
Polysaccharide 10 q: 38 n: 1.3 m: 0.1		Sodium dextranmethylcarboxylate modified by lauryl glycinate
Polysaccharide 11 q: 38 n: 1.3 m: 0.1		N-(sodium methylcarboxylate) dextran carbamate modified by diethyl aspartate
Polysaccharide 12 q: 4 n: 0.96 m: 0.07		Sodium dextranmethylcarboxylate modified by dilauryl aspartate
Polysaccharide 13 q: 38 n: 1.0 m: 0.1		Sodium dextranmethylcarboxylate modified by 2-(2-aminoethoxy)ethyl dodecanoate
Polysaccharide 14 q: 38 n: 1.0 m: 0.1		Sodium dextranmethylcarboxylate modified by 2-(2-{dodecanoylamino}ethoxy)ethylamine

Polysaccharide 15 q: 38 n: 1.05 m: 0.05		Sodium dextranmethylcarboxylate modified by 2-(2-{2-[hexadecanoylamino]ethoxy}ethoxy)ethylamine
Polysaccharide 18 q: 19 n: 1.61 m: 0.04		
Polysaccharide 19 q: 19 n: 1.06 m: 0.04		
Polysaccharide 20 q: 19 n: 0.66 m: 0.04		Sodium dextranmethylcarboxylate modified by cholestryl leucinate
Polysaccharide 21 q: 19 n: 0.46 m: 0.04		
Polysaccharide 22 q: 4 n: 1.61 m: 0.04		
Polysaccharide 26 q: 38 n: 0.99 m: 0.05		
Polysaccharide 23 q: 19 n: 1.61 m: 0.04		Sodium dextranmethylcarboxylate modified by cholestryl 1-ethylenediaminecarboxylate
Polysaccharide 24 q: 19 n: 1.96 m: 0.04		N-(sodium methylcarboxylate) dextran carbamate modified by cholestryl leucinate


POLYSACCHARIDES	SUBSTITUENTS -f-A-COONa -[E]-o-[F]	COMMON NAME
Polysaccharide 25 q: 19 n: 1.65		Sodium dextranmethylcarboxylate modified by cholestryl 1-ethylenediaminecarboxylate grafted by reductive amination to the reducing chain end

Table 1

Example A1: Preparation of Polysaccharide 1

[000240] 16 g (i.e., 296 mmol of hydroxyls) of dextran with a weight-average molar mass of approximately 10 kg/mol (q = 38, Pharmacosmos) are dissolved in water at 420 g/l. 30 ml of 10N NaOH (296 mmol) are added to the solution. The mixture is brought to 35°C and then 46 g (396 mmol) of sodium chloroacetate are added. The temperature of the reaction mixture is brought to 60°C at 0.5°C/min and then maintained at 60°C for 100 minutes. The reaction medium is diluted with 200 ml of

water, neutralized with acetic acid and purified by ultrafiltration through a 5 kDa PES membrane against 6 volumes of water. The final solution is assayed by dry extract, to determine the concentration of polysaccharide, and then assayed by acid/base titration in 50/50 (V/V) water/acetone, to determine the mean number of methylcarboxylate units per glucoside unit.

[000241] According to the dry extract: [polysaccharide] = 31.5 mg/g.

[000242] According to the acid/base titration: the mean number of methylcarboxylate units per glucoside unit is 1.1.

[000243] The sodium dextranmethylcarboxylate solution is passed through a Purolite resin (anionic) in order to obtain dextranmethylcarboxylic acid, which is subsequently lyophilized for 18 hours.

[000244] Octyl glycinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000245] 10 g of dextranmethylcarboxylic acid (44.86 mmol of methylcarboxylic acid) are dissolved in DMF at 60 g/l and then cooled to 0°C. 3.23 g of octyl glycinate, para-toluenesulphonic acid salt, (8.97 mmol) are suspended in DMF at 100 g/l. 0.91 g of triethylamine (8.97 mmol) is subsequently added to the suspension. Once the polysaccharide solution is at 0°C, a solution of NMM (5.24 g, 51.8 mmol) in DMF

(530 g/l) and 5.62 g (51.8 mmol) of EtOCOCl are subsequently added. After reacting for 10 minutes, the octyl glycinate suspension is added. The medium is subsequently maintained at 10°C for 45 minutes. The medium is subsequently heated to 30°C. A solution of imidazole (10.38 g in 17 ml of water) and 52 ml of water are added to the reaction medium. The polysaccharide solution is ultrafiltered through a 10 kDa PES membrane against 15 volumes of 0.9% NaCl solution and 5 volumes of water. The concentration of the polysaccharide solution is determined by dry extract. A fraction of solution is lyophilized and analyzed by ^1H NMR in D_2O in order to determine the degree of substitution of the methylcarboxylates to give octyl glycinate per glucoside unit.

5 [000246] According to the dry extract: [Polysaccharide 1] = 36.4 mg/g
10 [000247] According to the acid/base titration: n = 0.9
[000248] According to the ^1H NMR: m = 0.2.

Example A2: Preparation of Polysaccharide 2

15 [000249] Cetyl glycinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.
[000250] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 5 kg/mol (q = 19, Pharmacosmos), modified by cetyl glycinate, is 20 obtained by a process similar to that described in Example A1.
[000251] According to the dry extract: [Polysaccharide 2] = 15.1 mg/g
[000252] According to the acid/base titration: n = 1.05
[000253] According to the ^1H NMR: m = 0.05.

25 Example A3: Preparation of Polysaccharide 3

[000254] Octyl phenylalaninate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

30 [000255] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol (q = 38, Pharmacosmos), modified by octyl phenylalaninate, is obtained by a process similar to that described in Example A1.

[000256] According to the dry extract: [Polysaccharide 3] = 27.4 mg/g
[000257] According to the acid/base titration: n = 1.0
[000258] According to the ^1H NMR: m = 0.1.

35

Example A4: Preparation of Polysaccharide 4

[000259] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of

approximately 5 kg/mol ($q = 19$, Pharmacosmos), modified by octyl phenylalaninate, is obtained by a process similar to that described in Example A3.

[000260] According to the dry extract: [Polysaccharide 4] = 21.8 mg/g

[000261] According to the acid/base titration: $n = 1.0$

5 [000262] According to the ^1H NMR: $m = 0.2$.

Example A5: Preparation of Polysaccharide 5

[000263] 3,7-dimethyl-1-octyl phenylalaninate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

10 [000264] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by 3,7-dimethyl-1-octyl phenylalaninate, is obtained by a process similar to that described in Example A1.

[000265] According to the dry extract: [Polysaccharide 5] = 24.3 mg/g

15 [000266] According to the acid/base titration: $n = 1.0$

[000267] According to the ^1H NMR: $m = 0.1$.

Example A6: Preparation of Polysaccharide 6

[000268] Dioctyl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

20 [000269] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by dioctyl aspartate, is obtained by a process similar to that described in Example A1.

25 [000270] According to the dry extract: [Polysaccharide 6] = 22.2 mg/g

[000271] According to the acid/base titration: $n = 1.05$

[000272] According to the ^1H NMR: $m = 0.05$.

Example A7: Preparation of Polysaccharide 7

30 [000273] Didecyl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000274] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by didecyl aspartate, is obtained by a process similar to that described in Example A1.

[000275] According to the dry extract: [Polysaccharide 7] = 19.8 mg/g

[000276] According to the acid/base titration: $n = 1.05$

[000277] According to the ^1H NMR: $m = 0.05$.

Example A8: Preparation of Polysaccharide 8

[000278] Dilauryl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

5 [000279] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 5 kg/mol ($q = 19$, Pharmacosmos), modified by dilauryl aspartate is obtained by a process similar to that described in Example A1.

[000280] According to the dry extract: [Polysaccharide 8] = 22.8 mg/g

[000281] According to the acid/base titration: $n = 1.05$

10 [000282] According to the ^1H NMR: $m = 0.05$.

Example A9: Preparation of Polysaccharide 9

[000283] N-(2-aminoethyl)dodecanamide is obtained according to the process described in Patent US 2 387 201, from the methyl ester of dodecanoic acid (Sigma) and ethylenediamine (Roth).

15 [000284] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by N-(2-aminoethyl)dodecanamide, is obtained by a process similar to that described in Example A1.

20 [000285] According to the dry extract: [Polysaccharide 9] = 23.8 mg/g

[000286] According to the acid/base titration: $n = 1.0$

[000287] According to the ^1H NMR: $m = 0.1$.

25 Example A10: Preparation of Polysaccharide 10

[000288] Sodium dextransuccinate is obtained from a dextran with a weight-average molar mass of approximately 10 kg/mol ($q = 38$, Pharmacosmos) according to the method described in the paper by Sanchez-Chaves et al., 1998 (Manuel et al., Polymer, 1998, 39 (13), 2751-2757). According to the ^1H NMR in $\text{D}_2\text{O}/\text{NaOD}$, the mean number of succinate groups per glucoside unit is 1.4.

30 [000289] Lauryl glycinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000290] A sodium dextransuccinate modified by lauryl glycinate is obtained by a process similar to that described in Example A1.

35 [000291] According to the dry extract: [Polysaccharide 10] = 16.1 mg/g

[000292] According to the acid/base titration: $n = 1.3$

[000293] According to the ^1H NMR: $m = 0.1$.

Example A11: Preparation of Polysaccharide 11

[000294] Dioctyl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000295] 12 g (i.e., 0.22 mol of hydroxyls) of dextran with a weight-average molar mass of approximately 10 kg/mol ($q = 38$, Pharmacosmos) are dissolved in a DMF/DMSO mixture. The mixture is brought to 80°C with stirring. 3.32 g (0.03 mol) of 1,4-diazabicyclo[2.2.2]octane and then 14.35 g (0.11 mol) of ethyl isocyanatoacetate are gradually introduced. After reacting for 5 h, the medium is diluted in water and purified by diafiltration through a 5 kDa PES membrane against 0.1N NaOH, 0.9% NaCl and water. The final solution is assayed by dry extract, to determine the concentration of polysaccharide; and then assayed by acid/base titration in 50/50 (V/V) water/acetone, to determine the mean number of N-methylcarboxylate carbamate units per glucoside unit.

[000296] According to the dry extract: [polysaccharide] = 30.5 mg/g

[000297] According to the acid/base titration: the mean number of N-methylcarboxylate carbamate units per glucoside unit is 1.4.

[000298] An N-(sodium methylcarboxylate) dextran carbamate modified by dioctyl aspartate is obtained by a process similar to that described in Example A1.

[000299] According to the dry extract: [Polysaccharide 11] = 17.8 mg/g

[000300] According to the acid/base titration: $n = 1.3$

[000301] According to the ^1H NMR: $m = 0.1$.

Example A12: Preparation of Polysaccharide 12

[000302] Dilauryl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000303] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 1 kg/mol ($q = 4$, Pharmacosmos), modified by dilauryl aspartate, is obtained by a process similar to that described in Example A1.

[000304] According to the dry extract: [Polysaccharide 12] = 12.3 mg/g

[000305] According to the acid/base titration: $n = 0.96$

[000306] According to the ^1H NMR: $m = 0.07$.

Example A13: Preparation of Polysaccharide 13

[000307] 2-(2-aminoethoxy)ethyl dodecanoate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000308] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of

approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by 2-(2-aminoethoxy)ethyl dodecanoate is obtained, by a process similar to that described in Example A1.

[000309] According to the dry extract: [Polysaccharide 13] = 25.6 mg/g

5 [000310] According to the acid/base titration: $n = 1.0$

[000311] According to the ^1H NMR: $m = 0.1$.

Example A14: Preparation of Polysaccharide 14

[000312] 2-(2-{2-[dodecanoylamino]ethoxy}ethoxy)ethylamine is obtained according to the process described in Patent US 2 387 201, from the methyl ester of dodecanoic acid (Sigma) and triethylene glycol diamine (Huntsman).

[000313] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by 2-(2-{2-[dodecanoylamino]ethoxy}ethoxy)ethylamine, is obtained by a process similar to that described in Example A1.

[000314] According to the dry extract: [Polysaccharide 14] = 24.9 mg/g

[000315] According to the acid/base titration: $n = 1.0$

[000316] According to the ^1H NMR: $m = 0.1$.

20

Example A15: Preparation of Polysaccharide 15

[000317] 2-(2-{2-[hexadecanoylamino]ethoxy}ethoxy)ethylamine is obtained, according to the process described in Patent US 2 387 201, from the methyl ester of palmitic acid (Sigma) and triethylene glycol diamine (Huntsman).

25 [000318] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol ($q = 38$, Pharmacosmos), modified by 2-(2-{2-[hexadecanoylamino]ethoxy}ethoxy)ethylamine is obtained by a process similar to that described in Example A1.

30 [000319] According to the dry extract: [Polysaccharide 15] = 22.2 mg/g

[000320] According to the acid/base titration: $n = 1.05$

[000321] According to the ^1H NMR: $m = 0.05$.

Example A16: Preparation of Polysaccharide 16

35 [000322] Cetyl glycinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000323] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of

approximately 5 kg/mol ($q = 19$, Pharmacosmos), modified by cetyl glycinate is obtained by a process similar to that described in Example A1.

[000324] According to the dry extract: [Polysaccharide 16] = 23 mg/g

[000325] According to the acid/base titration: $n = 1.05$

5 [000326] According to the ^1H NMR: $m = 0.05$.

Example A17: Preparation of Polysaccharide 17

[000327] 10 g (i.e., 185 mmol of hydroxyls) of dextran with a weight-average molar mass of approximately 10 kg/mol ($q = 38$, Pharmacosmos) are dissolved in water at

10 420 g/l. 19 ml of 10N NaOH (185 mmol) are added to the solution. The mixture is brought to 35°C and then 8.6 g (74 mmol) of sodium chloroacetate are added. The temperature of the reaction mixture is brought to 60°C at 0.5°C/min and is then maintained at 60°C for 100 minutes. The reaction medium is diluted with 200 ml of water, neutralized with acetic acid and purified by ultrafiltration through a 5 kDa PES 15 membrane against 6 volumes of water. The final solution is assayed by dry extract, to determine the polysaccharide concentration, and then assayed by acid/base titration in 50/50 (V/V) water/acetone, to determine the mean number of methylcarboxylate units per glucoside unit.

[000328] According to the dry extract: [polysaccharide] = 35.1 mg/g

20 [000329] According to the acid/base titration: the mean number of methylcarboxylate units per glucoside unit is 0.42.

[000330] The sodium dextranmethylcarboxylate solution is passed through a Purolite resin (anionic) in order to obtain dextranmethylcarboxylic acid, which is subsequently lyophilized for 18 hours.

25 [000331] Cetyl glycinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000332] A sodium dextranmethylcarboxylate modified by cetyl glycinate is obtained by a process similar to that described in Example A1.

[000333] According to the dry extract: [Polysaccharide 17] = 18 mg/g

30 [000334] According to the acid/base titration: $n = 0.37$

[000335] According to the ^1H NMR: $m = 0.05$.

Example A18: Preparation of Polysaccharide 18

10 g of sodium dextranmethylcarboxylate, characterized by a degree of substitution as

35 methylcarboxylate of 1.10 per glucoside unit, are synthesized from a dextran with a weight-average molar mass of 5 kg/mol ($q = 19$, Pharmacosmos), according to a process similar to that described for Polysaccharide 1, and then lyophilized.

8 g (i.e., 64 mmol of hydroxyls) of sodium dextranmethylcarboxylate, characterized by a degree of substitution as methylcarboxylate of 1.05 per glucoside unit, are dissolved in water at 1000 g/l. 6 ml of 10N NaOH (64 mmol) are added. The mixture is heated to 35°C and 7.6 g of sodium chloroacetate (65 mmol) are added. The 5 mixture is gradually brought to a temperature of 60°C, and is maintained at this temperature for an additional 100 minutes. The mixture is diluted with water, neutralized with acetic acid and then purified by ultrafiltration through a 5 kDa PES membrane against water. The final solution is assayed by dry extract, to determine the polysaccharide concentration, and then assayed by acid/base titration in 50/50 10 (V/V) water/acetone, to determine the mean number of methylcarboxylate units per glucoside unit.

According to the dry extract: [polysaccharide] = 45.8 mg/g

According to the acid/base titration: the mean number of methylcarboxylate units per glucoside unit is 1.65.

15 The sodium dextranmethylcarboxylate solution is passed through a Purolite resin (anionic) in order to obtain dextranmethylcarboxylic acid, which is subsequently lyophilized for 18 hours.

Cholesteryl leucinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

20 A sodium dextranmethylcarboxylate modified by cholesteryl leucinate, is obtained by a process similar to that described in Example A1.

According to the dry extract: [Polysaccharide 18] = 21 mg/g

According to the acid/base titration: n = 1.61

According to the ¹H NMR: m = 0.04.

25

Example A19: Preparation of Polysaccharide 19

A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 5 kg/mol (q = 19, Pharmacosmos), modified by cholesteryl leucinate, is obtained by a 30 process similar to that described in Example A1.

According to the dry extract: [Polysaccharide 19] = 19.4 mg/g

According to the acid/base titration: n = 1.06

According to the ¹H NMR: m = 0.04.

35 Example A20: Preparation of Polysaccharide 20

16 g (i.e., 296 mmol of hydroxyls) of dextran with a weight-average molar mass of approximately 5 kg/mol (q = 19, Pharmacosmos) are dissolved in water at 420 g/l. 30 ml of 10N NaOH (296 mmol) are added to this solution. The mixture is brought to

35°C and then 26 g (222 mmol) of sodium chloroacetate are added. The temperature of the reaction medium is gradually brought to 60°C and then maintained at 60°C for 100 minutes. The reaction medium is diluted with water, neutralized with acetic acid and purified by ultrafiltration through a 5 kDa PES membrane against water. The final 5 solution is assayed by dry extract, to determine the polysaccharide concentration, and then assayed by acid/base titration in 50/50 (V/V) water/acetone, to determine the mean number of methylcarboxylate units per glucoside unit.

According to the dry extract: [polysaccharide] = 33.1 mg/g
According to the acid/base titration: the mean number of methylcarboxylate units per 10 glucoside unit is 0.70.

The sodium dextranmethylcarboxylate solution is passed through a Purolite resin (anionic) in order to obtain dextranmethylcarboxylic acid, which is subsequently lyophilized for 18 hours.

A sodium dextranmethylcarboxylate modified by cholesteryl leucinate is obtained by a 15 process similar to that described in Example A1.

According to the dry extract: [Polysaccharide 20] = 18.9 mg/g
According to the acid/base titration: n = 0.66
According to the ¹H NMR: m = 0.04.

20 Example A21: Preparation of Polysaccharide 21

16 g (i.e., 296 mmol of hydroxyls) of dextran with a weight-average molar mass of approximately 5 kg/mol (q = 19, Pharmacosmos) are dissolved in water at 420 g/l. 30 ml of 10N NaOH (296 mmol) are added to this solution. The mixture is brought to 35°C and then 18 g (158 mmol) of sodium chloroacetate are added. The temperature 25 of the reaction medium is gradually brought to 60°C and then maintained at 60°C for 100 minutes. The reaction medium is diluted with water, neutralized with acetic acid and purified by ultrafiltration through a 1 kDa PES membrane against water. The final solution is assayed by dry extract, to determine the polysaccharide concentration, and then assayed by acid/base titration in 50/50 (V/V) water/acetone, to determine the 30 mean number of methylcarboxylate units per glucoside unit.

According to the dry extract: [polysaccharide] = 52.6 mg/g
According to the acid/base titration: the mean number of methylcarboxylate units per glucoside unit is 0.50.

The sodium dextranmethylcarboxylate solution is passed through a Purolite resin 35 (anionic) in order to obtain dextranmethylcarboxylic acid, which is subsequently lyophilized for 18 hours.

A sodium dextranmethylcarboxylate modified by cholesteryl leucinate is obtained by a process similar to that described in Example A1.

According to the dry extract: [Polysaccharide 21] = 18.9 mg/g

According to the acid/base titration: n = 0.46

According to the ^1H NMR: m = 0.04.

5 Example A22: Preparation of Polysaccharide 22

A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A18 using a dextran with a weight-average molecular weight of approximately 1 kg/mol (q = 4, Pharmacosmos), modified by cholesteryl leucinate, is obtained by a process similar to that described in Example A18.

10 According to the dry extract: [Polysaccharide 22] = 20.2 mg/g

According to the acid/base titration: n = 1.61

According to the ^1H NMR: m = 0.04.

Example A23: Preparation of Polysaccharide 23

15 Cholesteryl 1-ethylenediaminecarboxylate hydrochloride is obtained according to the process described in the patent (Akiyoshi, K et al., WO 2010/053140).

A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A18 using a dextran with a weight-average molecular weight of approximately 5 kg/mol (q = 19, Pharmacosmos), modified by cholesteryl

20 1-ethylenediaminecarboxylate, is obtained by a process similar to that described in Example A18.

According to the dry extract: [Polysaccharide 23] = 20.1 mg/g

According to the acid/base titration: n = 1.61

According to the ^1H NMR: m = 0.04.

25

Example A24: Preparation of Polysaccharide 24

12 g (i.e., 0.22 mol of hydroxyls) of dextran with a weight-average molar mass of approximately 5 kg/mol (q = 19, Pharmacosmos) are dissolved in a DMF/DMSO mixture. The mixture is brought to 80°C with stirring. 3.32 g (0.03 mol) of 1,4-

30 diazabicyclo[2.2.2]octane and then 26.8 g (0.21 mol) of ethyl isocyanatoacetate are gradually introduced. After reacting for 5 h, the medium is diluted in water and purified by diafiltration through a 5 kDa PES membrane against 0.1N NaOH, 0.9%

NaCl and water. The final solution is assayed by dry extract, to determine the concentration of polysaccharide; and then assayed by acid/base titration in 50/50 (V/V) water/acetone, to determine the mean number of N-methylcarboxylate carbamate units per glucoside unit.

According to the dry extract: [polysaccharide] = 30.1 mg/g

According to the acid/base titration: the mean number of N-methylcarboxylate carbamate units per glucoside unit is 2.0.

An N-(sodium methylcarboxylate) dextran carbamate modified by cholestryl leucinate is obtained by a process similar to that described in Example A1.

5 According to the dry extract: [Polysaccharide 24] = 17.9 mg/g
 [000336] According to the acid/base titration: n = 1.96
 [000337] According to the ^1H NMR: m = 0.04.

Example A25: Preparation of Polysaccharide 25

10 Cholestryl 1-ethylenediaminecarboxylate hydrochlorate is obtained according to the process described in the patent (Akiyoshi, K et al., WO 2010/053140).
 10 g of dextran with a weight-average molar mass of approximately 5 kg/mol (q = 19, Pharmacosmos, 3.2 mmol of chain ends) are dissolved in DMSO at 80°C. 4.8 g of cholestryl 1-ethylenediaminecarboxylate hydrochloride (9.5 mmol), 0.96 g of triethylamine (9.5 mmol) and 2.0 g of sodium cyanoborohydride (32 mmol) are added to the reaction medium, which is stirred at 80°C for 24 hours. After cooling, the mixture is precipitated from dichloromethane and then from acetone, and dried under vacuum. According to the ^1H NMR, a dextran modified at the chain end by cholestryl 1-ethylenediaminecarboxylate is obtained. A sodium dextranmethylcarboxylate
 15 characterized by a degree of substitution as methylcarboxylate of 1.65 per glucoside unit and modified at the chain end by cholestryl 1-ethylenediaminecarboxylate was synthesized by a process similar to that described in Example A18 using the dextran modified at the chain end by cholestryl 1-ethylenediaminecarboxylate.

According to the dry extract: [Polysaccharide 25] = 13.7 mg/g

25 According to the acid/base titration: n = 1.65
 According to the ^1H NMR: each polymer chain carries a cholestryl 1-ethylenediaminecarboxylate group grafted to the reducing chain end.

Example A26: Preparation of Polysaccharide 26

30 [000338] Cholesterol leucinate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.
 [000339] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 10 kg/mol (q = 38, Pharmacosmos), modified by cholesterol leucinate,
 35 is obtained by a process similar to that described in Example A1.
 [000340] According to the dry extract: [Polysaccharide 26] = 26.6 mg/g
 [000341] According to the acid/base titration: n = 0.99
 [000342] According to the ^1H NMR: m = 0.05.

Example A27: Preparation of Polysaccharide 27

Dilauryl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

5 A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A18 using a dextran with a weight-average molecular weight of approximately 1 kg/mol (q = 4, Pharmacosmos), modified by dilauryl aspartate, is obtained by a process similar to that described in Example A1.

[000343] According to the dry extract: [Polysaccharide 27] = 16.7 mg/g

10 [000344] According to the acid/base titration: n = 1.41

[000345] According to the ^1H NMR: m = 0.16.

Example A28: Preparation of Polysaccharide 28

[000346] Dilauryl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000347] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A18 using a dextran with a weight-average molecular weight of approximately 1 kg/mol (q = 4, Pharmacosmos), modified by dilauryl aspartate, is obtained by a process similar to that described in Example A1.

20 [000348] According to the dry extract: [Polysaccharide 28] = 25 mg/g

[000349] According to the acid/base titration: n = 1.50

[000350] According to the ^1H NMR: m = 0.07.

Example A29: Preparation of Polysaccharide 29

25 [000351] Didecyl aspartate, para-toluenesulphonic acid salt, is obtained according to the process described in Patent US 4 826 818.

[000352] A sodium dextranmethylcarboxylate, synthesized according to the process described in Example A1 using a dextran with a weight-average molecular weight of approximately 1 kg/mol (q = 4, Pharmacosmos), modified by didecyl aspartate, is obtained by a process similar to that described in Example A1.

30 [000353] According to the dry extract: [Polysaccharide 29] = 15 mg/g

[000354] According to the acid/base titration: n = 1.05

[000355] According to the ^1H NMR: m = 0.05.

[000356] Examples

Part B: Demonstration of the properties of the compositions according to the invention

5

Example B1: Solution of rapid-acting insulin analogue (NovoLog®) at 100 IU/ml

[000357] This solution is a commercial solution of insulin aspart, sold by the company Novo Nordisk under the name of NovoLog® in the USA and Novorapid® in Europe. This product is a rapid-acting insulin analogue.

10

Example B2: Solution of rapid-acting insulin analogue (Humalog®) at 100 IU/ml

[000358] This solution is a commercial solution of insulin lispro, sold by the company Eli Lilly under the name of Humalog®. This product is a rapid-acting insulin analogue.

15

Example B3: Solution of rapid-acting insulin analogue (Apidra®) at 100 IU/ml

[000359] This solution is a commercial solution of insulin glulisine, sold by the company Sanofi-Aventis under the name of Apidra®. This product is a rapid-acting insulin analogue.

20

Example B4: Solution of slow-acting insulin analogue (Lantus®) at 100 IU/ml

[000360] This solution is a commercial solution of insulin glargine, sold by the company Sanofi-Aventis under the name of Lantus®. This product is a slow-acting insulin analogue.

25

Example B5: Solution of human insulin (ActRapid®) at 100 IU/ml

[000361] This solution is a commercial solution from Novo Nordisk, sold under the name of Actrapid®. This product is a human insulin.

30

Example B6: Dissolution of Lantus® at 100 IU/ml and at pH 7 using a substituted dextran

[000362] 20 mg of Polysaccharide 4 described in Example A4 are weighed out accurately. This lyophilisate is taken up in 2 ml of Lantus® in its commercial formulation. A transient precipitate appears but the solution becomes clear after approximately 30 minutes. The pH of this solution is 6.3. The pH is adjusted to 7 with a 0.1N sodium hydroxide solution. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C.

Example B7: Preparation of a substituted dextran/Lantus®/Apidra® 75/25 composition at pH 7

[000363] 0.25 ml of Apidra® (in its commercial formulation) is added to 0.75 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6, in order to form 1 ml of a composition at pH 7. The composition is clear, testifying to the good solubility of Lantus® and Apidra® under these formulation conditions. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C.

Example B8: Preparation of a substituted dextran/Lantus®/Humalog® 75/25 composition at pH 7

[000364] 0.25 ml of Humalog® (in its commercial formulation) is added to 0.75 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6, in order to form 1 ml of a composition at pH 7. The composition is clear, testifying to the good solubility of Lantus® and Humalog® under these formulation conditions. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C.

Example B9: Preparation of a substituted dextran/Lantus®/NovoLog® 75/25 composition at pH 7

[000365] 0.25 ml of NovoLog® (in its commercial formulation) is added to 0.75 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6, in order to form 1 ml of a composition at pH 7. The composition is clear, testifying to the good solubility of Lantus® and NovoLog® under these formulation conditions. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C.

Example B10: Preparation of a substituted dextran/Lantus®/ActRapid® 75/25 composition at pH 7

[000366] 0.25 ml of ActRapid® (in its commercial formulation) is added to 0.75 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6, in order to form 1 ml of a composition at pH 7. The composition is clear, testifying to the good solubility of Lantus® and ActRapid® under these formulation conditions. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C.

Example B11: Preparation of a substituted dextran/Lantus®/Apidra® 60/40 composition at pH 7

[000367] 0.4 ml of Apidra® (in its commercial formulation) is added to 0.6 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6, in order to form 1 ml of a composition at pH 7. The composition is clear, testifying to the good solubility of

Lantus® and Apidra® under these formulation conditions. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C.

Example B12: Preparation of a substituted dextran/Lantus®/Apidra® 40/60
5 composition at pH 7

[000368] 0.6 ml of Apidra® (in its commercial formulation) is added to 0.4 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6, in order to form 1 ml of a composition at pH 7. The composition is clear, testifying to the good solubility of Lantus® and Apidra® under these formulation conditions. This clear solution is filtered
10 through a 0.22 µm filter and is then placed at +4°C.

Example B13: Precipitation of Lantus®

[000369] 1 ml of Lantus® is added to 2 ml of a PBS solution containing 20 mg/ml of BSA (bovine serum albumin). The PBS/BSA mixture simulates the composition of the
15 subcutaneous medium. A precipitate appears, which is in good agreement with the mechanism of operation of Lantus® (precipitation on injection due to the increase in the pH).

[000370] Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the
20 supernatent. The result of this is that 86% of Lantus® is found in a precipitated form.

Example B14: Precipitation of a substituted dextran/Lantus® composition

[000371] 1 ml of Polysaccharide 4/Lantus® solution prepared in Example B6 is added to 2 ml of a PBS solution containing 20 mg/ml of BSA (bovine serum albumin). The
25 PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

[000372] Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. The result of this is that 85% of Lantus® is found in a precipitated form.
30 This percentage of precipitation of Lantus® is identical to that obtained for the control described in Example B13.

[000373] Dissolution and precipitation tests identical to those described in Examples B6 and B14 were carried out with other substituted dextrans at the same
35 concentration of 10 mg/ml of polysaccharide per 100 IU/ml of Lantus®. 20 mg of polysaccharide in the lyophilisate form are weighed out accurately. This lyophilisate is taken up in 2 ml of Lantus® in its commercial formulation. A transient precipitate appears but the solution becomes clear after approximately 30 minutes to a few hours

(depending on the nature of the polysaccharide). The pH of this solution is 6.3. The pH is adjusted to 7 with a 0.1N sodium hydroxide solution. This clear solution is filtered through a 0.22 µm filter and is then placed at +4°C. The results are collated in Table 2.

5

Polysaccharide No.	Dissolution of Lantus®	Precipitation of Lantus®	% of precipitation
2	Yes	Yes	85
1	Yes	Yes	Not measured
4	Yes	Yes	87
3	Yes	Yes	Not measured
5	Yes	Yes	94
6	Yes	Yes	Not measured
7	Yes	Yes	Not measured
8	Yes	Yes	Not measured
9	Yes	Yes	94
10	Yes	Yes	Not measured
15	Yes	Yes	Not measured
14	Yes	Yes	Not measured
13	Yes	Yes	Not measured
12	Yes	Yes	Not measured
11	Yes	Yes	Not measured
16	Yes	Yes	Not measured
17	Yes	Yes	Not measured
18	Yes	Yes	Not measured
19	Yes	Yes	Not measured
20	Yes	Yes	Not measured
21	Yes	Yes	Not measured
22	Yes	Yes	Not measured
23	Yes	Yes	Not measured
24	Yes	Yes	Not measured
25	Yes	Yes	Not measured
26	Yes	Yes	Not measured

Table 2

Example B15: Precipitation of a substituted dextran/Lantus®/Apidra® 75/25 composition at pH 7

1 ml of the substituted dextran/Lantus®/Apidra® 75/25 composition (containing 7.5 mg/ml of polysaccharide, 75 IU/ml of Lantus® and 25 IU/ml of Apidra®) prepared in Example B7 is added to 2 ml of a PBS solution containing 20 mg/ml of BSA (bovine serum albumin). The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

[000374] Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. The percentages of precipitation of Lantus® are similar to the control described in Example B13.

Example B16: Precipitation of various compositions, the nature of the substituted dextran being varied

[000375] Other tests under the same conditions as those of Example B15 were carried out in the presence of other substituted dextrans.

[000376] The results are combined in the following Table 3 and it is observed that the dissolution and the precipitation of Lantus® are retained.

Polysaccharide No.	Dissolution Lantus®/Apidra® 75/25	Percentage of precipitation of Lantus®
2	Yes	85
1	Yes	Not measured
4	Yes	87
3	Yes	Not measured
5	Yes	86
6	Yes	Not measured
7	Yes	Not measured
8	Yes	Not measured
9	Yes	86
10	Yes	85
15	Yes	87
14	Yes	86
13	Yes	88
12	Yes	91
18	Yes	Not measured
19	Yes	Not measured
20	Yes	Not measured
21	Yes	Not measured
22	Yes	Not measured
23	Yes	Not measured
24	Yes	Not measured
25	Yes	Not measured
26	Yes	Not measured

Table 3

5 Example B17: Precipitation of various compositions, the nature of the prandial insulin being varied

[000377] Compositions are prepared by mixing 0.75 ml of the Polysaccharide 4/Lantus® solution prepared in Example B6 with 0.25 ml of a prandial insulin in order to form 1 ml of substituted dextran/Lantus®/prandial insulin composition (containing

10 7.5 mg/ml of polysaccharide, 75 IU/ml of Lantus® and 25 IU/ml of prandial insulin).

[000378] This composition is added to 2 ml of PBS solution containing 20 mg/ml of BSA (bovine serum albumin). The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

[000379] Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. In the presence of the four prandial insulins tested, Lantus® precipitates to approximately 90%. This percentage of precipitation of Lantus® is similar to the control described in Example B13; the results are collated in Table 4.

Nature of the prandial insulin	Dissolution Lantus®/prandial insulin 75/25	Percentage of precipitation of Lantus®
Apidra®	Yes	88
NovoLog®	Yes	92
Humalog®	Yes	89
ActRapid®	Yes	90

Table 4

Example B18: Preparation of a concentrated solution of slow-acting insulin analogue (glargine).

A commercial solution of insulin glargine, sold by the company Sanofi-Aventis under the name of Lantus®, is concentrated by ultrafiltration through a 3 kDa membrane made of regenerated cellulose (Amicon® Ultra-15, sold by the company Millipore). On conclusion of this ultrafiltration stage, the concentration of insulin glargine is assayed in the retentate by reverse phase liquid chromatography (RP-HPLC). The final concentration of insulin glargine is subsequently adjusted by the addition of commercial 100 IU/ml glargine solution in order to obtain the desired final concentration. This process makes it possible to obtain concentrated solutions of glargine, denoted C_{glargine} , at various concentrations of greater than 100 IU/ml, such as $C_{\text{glargine}} = 200, 250, 300$ and 333 IU/ml. The concentrated solutions are filtered through a $0.22\text{ }\mu\text{m}$ filter and then stored at $+4^\circ\text{C}$.

Example B19: Dialysis of a commercial solution of rapid-acting insulin analogue (lispro)

A commercial solution of insulin lispro, sold by the company Lilly under the name of Humalog® is dialyzed by ultrafiltration through a 3 kDa membrane made of regenerated cellulose (Amicon® Ultra-15, sold by the company Millipore). The dialysis is carried out in a 1 mM phosphate buffer at pH 7. On conclusion of this dialysis stage, the concentration $C_{\text{dialyzed Humalog}}$ of lispro in the retentate is determined by reverse phase liquid chromatography (RP-HPLC). The dialyzed solution is stored in a freezer at -80°C .

Example B20: Lyophilization of a solution of rapid-acting insulin analogue (lispro) in its commercial form

5 A volume V_{Humalog} of a solution of rapid-acting insulin lispro at a concentration of 100 IU/ml in its commercial form is placed in a Lyoguard® sterilized beforehand in an autoclave. The Lyoguard® is placed in a freezer at -80°C for approximately 1 h before being subjected to lyophilization overnight at a temperature of 20°C and a pressure of 0.31 mbar.

The sterile lyophilisate thus obtained is stored at ambient temperature.

10 Example B21: Lyophilization of a dialyzed commercial solution of rapid-acting insulin analogue (lispro)

A volume $V_{\text{dialyzed Humalog}}$ of a solution of rapid-acting insulin lispro obtained according to Example B19 at a concentration of $C_{\text{dialyzed Humalog}}$ is placed in a Lyoguard® sterilized beforehand in an autoclave. The Lyoguard® is placed in a freezer at -80°C for 15 approximately 1 h before being subjected to lyophilization overnight at a temperature of 20°C and a pressure of 0.31 mbar.

The sterile lyophilisate thus obtained is stored at ambient temperature.

20 Example B22: Preparation of a substituted dextran/glargine composition at pH 7 using a substituted dextran, according to a process using glargine in the liquid form (in solution) and a polysaccharide in the solid form (lyophilized)

A weight $w_{\text{polys.}}$ of Polysaccharide 18 is weighed out accurately. This lyophilisate is taken up in a volume V_{glargine} of a concentrated solution of glargine prepared according to Example B18, so as to obtain a composition exhibiting a concentration of 25 polysaccharide $C_{\text{polys.}}(\text{mg/ml}) = w_{\text{polys.}}/V_{\text{glargine}}$ and a concentration of glargine C_{glargine} (IU/ml). The solution is opalescent. The pH of this solution is approximately 6.3. The pH is adjusted to 7 by addition of concentrated NaOH and then the solution is placed under static conditions in an oven at 37°C for approximately 1 hour. A volume $V_{\text{polys./glargine}}$ of this visually clear solution is placed at +4°C.

30

Example B23: Preparation of a substituted dextran/glargine composition at pH 7 using a substituted dextran, according to a process using glargine in the liquid form (in solution) and a polysaccharide in the liquid form (in solution)

Concentrated solutions of m-cresol, glycerol and tween 20 are added to a mother 35 solution of polysaccharide 20 at pH 7 exhibiting a concentration $C_{\text{polys. mother}}$, so as to obtain a polysaccharide solution having the concentration $C_{\text{polys. mother/excipients}}$ (mg/ml) in the presence of these excipients at contents equivalent to those described in the Lantus® commercial solution in a 10 ml bottle.

A volume V_{Lantus} of a commercial solution of slow-acting insulin glargine, sold under the name of Lantus® at a concentration of 100 IU/ml, is added to a volume $V_{\text{polys. mother/excipients}}$ of a polysaccharide solution at the concentration $C_{\text{polys. mother/excipients}}$ (mg/ml) in a sterile flask. A cloudiness appears. The pH is adjusted to pH 7 by 5 addition of 1M NaOH and the solution is placed under static conditions in an oven at 37°C for approximately 1 h. This visually clear solution is placed at +4°C.

Example B24: Preparation of a concentrated polysaccharide/glargine composition at pH=7 using a substituted dextran, according to a process for the concentration of a 10 dilute composition

A dilute Polysaccharide 20/glargine composition at pH 7 described in Example B23 is concentrated by ultrafiltration through a 3 kDa membrane made of regenerated cellulose (Amicon® Ultra-15, sold by the company Millipore). On conclusion of this ultrafiltration stage, the retentate is clear and the concentration of insulin glargine in 15 the composition is assayed by reverse phase chromatography (RP-HPLC). If necessary, the concentration of insulin glargine in the composition is subsequently adjusted to the desired value by diluting in a solution of m-cresol/glycerol/tween 20 excipients exhibiting, for each entity, a concentration equivalent to that described in the Lantus® commercial solution (in a 10 ml bottle). This solution at pH 7, which is 20 visually clear, exhibiting a glargine concentration C_{glargine} (IU/ml) and a polysaccharide concentration $C_{\text{polys.}}$ (mg/ml), is placed at +4°C.

Example B25: Preparation of a substituted dextran/glargine/lispro composition at pH 7, starting from a rapid-acting insulin lispro in its commercial form

25 A volume $V_{\text{polysach./glargine}}$ of polysaccharide/glargine solution, pH 7, exhibiting a concentration of glargine C_{glargine} (IU/ml) and a concentration of Polysaccharide 18 $C_{\text{polys.}}$ (mg/ml), prepared according to Example B22, is added to an insulin lispro lyophilisate obtained by lyophilization of a volume V_{lispro} , the preparation of which is described in Example B19, so that the ratio $V_{\text{polysach./glargine}} / V_{\text{lispro}} = 100 / C_{\text{lispro}}$, where 30 C_{lispro} is the concentration of lispro (IU/ml) targeted in the composition. The solution is clear. The zinc content of the formulation is adjusted to the desired concentration C_{zinc} (μM) by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargine and lispro under 35 these formulation conditions. This solution is filtered through a 0.22 μm filter and is placed at +4°C.

Example B26: Preparation of a substituted dextran/glargine/lispro composition at pH 7, starting from a rapid-acting insulin lispro obtained by dialysis of a commercial solution

A volume $V_{\text{polysach./glargine}}$ of polysaccharide/glargine solution, pH 7, exhibiting a concentration of glargine C_{glargine} (IU/ml) and a concentration of Polysaccharide 20 $C_{\text{polys.}}$ (mg/ml), prepared according to Example B24, is added to an insulin lispro lyophilisate obtained by lyophilization of a volume $V_{\text{dialyzed Humalog}}$, the preparation of which is described in Example B21, so that the ratio $V_{\text{polysach./glargine}} / V_{\text{dialyzed Humalog}} = C_{\text{dialyzed Humalog}} / C_{\text{lispro}}$, where $C_{\text{dialyzed Humalog}}$ is the concentration of lispro (IU/ml) obtained on conclusion of the dialysis of the commercial solution, which stage is described in Example B19, and C_{lispro} is the concentration of lispro (IU/ml) targeted in the composition. The solution is clear. The zinc content of the formulation is adjusted to the desired concentration C_{zinc} (μM) by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargine and lispro under these formulation conditions. This solution is filtered through a 0.22 μm filter and is placed at +4°C.

Example B27: Preparation of a substituted dextran/glargine/lispro composition at pH 7

exhibiting a concentration of glargine of 200 IU/ml and a concentration of lispro of 33 IU/ml (proportion as percentage of insulin: 85/15 as glargine/lispro)

A concentrated 200 IU/ml glargine solution is prepared according to Example B18. A Polysaccharide 18 (13 mg/ml)/glargine 300 IU/ml composition, pH 7, is prepared from Polysaccharide 18 according to the method of preparation described in Example B22.

This Polysaccharide 18/glargine 200 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue in its commercial form, according to the method of preparation described in Example B25. The solution is clear. The zinc content of the formulation is adjusted to the concentration C_{zinc} (μM) = 750 μM by addition of a concentrated zinc chloride solution.

The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

This composition is described in Table 5.

Example B28: Preparation of a substituted dextran/glargine/lispro composition at pH 7

exhibiting a concentration of glargine of 200 IU/ml and a concentration of lispro of 66

IU/ml (proportion as percentage of insulin: 75/25 as glargine/lispro)

A concentrated 200 IU/ml glargine solution is prepared according to Example B18. A Polysaccharide 18 (13 mg/ml)/glargine 300 IU/ml composition, pH 7, is prepared from Polysaccharide 18 according to the method of preparation described in Example B22.

This Polysaccharide 18/glargine 200 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue in its commercial form, according to the method of preparation described in Example B25. The solution is clear. The zinc content of the formulation is adjusted to the 5 concentration C_{zinc} (μM) = 1500 μM by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargin and lispro under these formulation conditions. This solution is filtered through a 0.22 μm filter and is placed at +4°C.

10 This composition is described in Table 5.

Example B29: Preparation of a substituted dextran/glargine/lispro composition at pH 7 exhibiting a concentration of glargin of 300 IU/ml and a concentration of lispro of 100 IU/ml (proportion as percentage of insulin: 75/25 as glargin/lispro)

15 A concentrated 300 IU/ml glargin solution is prepared according to Example B18. A Polysaccharide 18 (23 mg/ml)/glargin 300 IU/ml composition, pH 7, is prepared from Polysaccharide 18 according to the method of preparation described in Example B22. This Polysaccharide 18/glargine 300 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue in its 20 commercial form, according to the method of preparation described in Example B25. The solution is clear. The zinc content of the formulation is adjusted to the concentration C_{zinc} (μM) = 2000 μM by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargin and lispro under 25 these formulation conditions. This solution is filtered through a 0.22 μm filter and is placed at +4°C.

This composition is described in Table 5.

Example B30: Preparation of a substituted dextran/glargine/lispro composition at pH 7 30 exhibiting a concentration of glargin of 250 IU/ml and a concentration of lispro of 150 IU/ml (proportion as percentage of insulin: 63/37 as glargin/lispro)

A concentrated 300 IU/ml glargin solution is prepared according to Example B18. A Polysaccharide 18 (19 mg/ml)/glargin 300 IU/ml composition, pH 7, is prepared from Polysaccharide 18 according to the method of preparation described in Example B22. 35 This Polysaccharide 18/glargine 250 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue in its commercial form, according to the method of preparation described in Example B25. The solution is clear. The zinc content of the formulation is adjusted to the

concentration C_{zinc} (μ M) = 1500 μ M by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargine and lispro under these formulation conditions. This solution is filtered through a 0.22 μ m filter and is placed at +4°C.

This composition is described in Table 5.

Example B31: Preparation of a substituted dextran/glargine/lispro composition at pH 7 exhibiting a concentration of glargine of 333 IU/ml and a concentration of lispro of 67 IU/ml (proportion as percentage of insulin: 83/17 as glargine/lispro)

A concentrated 333 IU/ml glargine solution is prepared according to Example B18. A Polysaccharide 18 (20 mg/ml)/glargine 300 IU/ml composition, pH 7, is prepared from Polysaccharide 18 according to the method of preparation described in Example B22. This Polysaccharide 18/glargine 333 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue in its commercial form, according to the method of preparation described in Example B25. The solution is clear. The zinc content of the formulation is adjusted to the concentration C_{zinc} (μ M) = 2000 μ M by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargine and lispro under these formulation conditions. This solution is filtered through a 0.22 μ m filter and is placed at +4°C.

This composition is described in Table 5.

Example B32: Preparation of a substituted dextran/glargine/lispro composition at pH 7 exhibiting a concentration of glargine of 300 IU/ml and a concentration of lispro of 100 IU/ml (proportion as percentage of insulin: 75/25 as glargine/lispro)

A concentrated 300 IU/ml glargine solution is prepared according to Example B18. A Polysaccharide 19 (23 mg/ml)/glargine 300 IU/ml composition, pH 7, is prepared from Polysaccharide 19 according to the method of preparation described in Example B22. This Polysaccharide 19/glargine 300 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue under its dialyzed form, according to the method of preparation described in Example B26. The solution is clear. The zinc content of the formulation is adjusted to the concentration C_{zinc} (μ M) = 3000 μ M by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

The formulation is clear, testifying to the good solubility of glargine and lispro under these formulation conditions. This solution is filtered through a 0.22 µm filter and is placed at +4°C.

This composition is described in Table 5.

5

Example B33: Preparation of a substituted dextran/glargine/lispro composition at pH 7 exhibiting a concentration of glargine of 300 IU/ml and a concentration of lispro of 100 IU/ml (proportion as percentage of insulin: 75/25 as glargine/lispro)

10 A Polysaccharide 20 (23 mg/ml)/glargine 300 IU/ml composition, pH 7, is prepared from Polysaccharide 20 according to the method of preparation described in Example B23. This Polysaccharide 20/glargine 300 IU/ml composition is added to an insulin lispro lyophilisate obtained by lyophilization of the solution of rapid-acting analogue resulting from the dialysis of a commercial solution, according to the method of preparation described in Example B26. The solution is clear. The zinc content of the 15 formulation is adjusted to the concentration C_{zinc} (µM) = 1500 µM by addition of a concentrated zinc chloride solution. The final pH is adjusted to 7 by addition of concentrated NaOH or HCl.

20 The formulation is clear, testifying to the good solubility of glargine and lispro under these formulation conditions. This solution is filtered through a 0.22 µm filter and is placed at +4°C.

This composition is described in Table 5.

Example No.	Polysaccharide No.	$C_{\text{polysach.}}$ (mg/ml)	C_{glargine} (IU/ml)	C_{lispro} (IU/ml)	$C_{\text{glargine}}/C_{\text{lispro}}$ (%/%)	pH
B27	18	13	200	33	85/15	7
B28	18	13	200	66	75/25	7
B29	18	23	300	100	75/25	7
B30	18	19	250	150	63/37	7
B31	18	20	333	67	83/17	7
B32	19	23	300	100	75/25	7
B33	20	23	300	100	75/25	7

Table 5: Substituted dextran/glargine/lispro compositions at pH 7

25 Example B34: Precipitation of various substituted dextran/glargine/lispro compositions at pH 7 exhibiting different concentrations of insulin glargine and insulin lispro and different relative proportions of the two insulins

1 ml of substituted dextran/Lantus®/Humalog® composition prepared in Examples B27 to B33 is added to 2 ml of a PBS solution containing 20 mg/ml of BSA (bovine serum albumin). The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

5 Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. The percentages of precipitation of Lantus® are similar to the control described in Example B13. The results are summarized in Table 6.

Example No.	Polysaccharide No.	C _{polysach.} (mg/ml)	C _{glargine} (IU/ml)	C _{lispro} (IU/ml)	C _{glargine/C lispro} (%/%)	Dis-solution of glargine and of lispro at pH 7	Precipitation of glargine	% Precipitation
B27	18	13	200	33	85/15	YES	YES	96
B28	18	13	200	66	75/25	YES	YES	86
B29	18	23	300	100	75/25	YES	YES	91
B30	18	19	250	150	63/37	YES	YES	90
B31	18	20	333	67	83/17	YES	YES	93
B32	19	23	300	100	75/25	YES	YES	98
B33	20	23	300	100	75/25	YES	YES	Not measured

10

Table 6

Example B35: Chemical stability of the compositions

[000380] The substituted dextran/Lantus®/prandial insulin compositions described in Examples B7, B27, B28 and B29 and the corresponding controls are placed at 30°C for 15 4 weeks. Regulations require 95% of native (nondegraded) insulin after 4 weeks at 30°C.

[000381] After 4 weeks, the formulations studied meet the specifications defined by the regulations. The results are combined in Table 7.

20

Compositions	Percentage of native glargine after 4 weeks at 30°C	Percentage of native prandial insulin after 4 weeks at 30°C
Lantus® (commercial formulation)	97	na
Apidra® (commercial formulation)	na	95
Humalog® (commercial formulation)	na	98
B7	96	98
B27	97	99
B28	95	97
B29	98	100

Table 7

5 [000382] Thus, whatever the formulation studied, a percentage of native insulin of greater than 95% is obtained, which is in accordance with regulatory requirements.

Example B36: Injectability of the solutions

10 [000383] All the compositions prepared can be injected with the normal systems for the injection of insulin. The solutions described in Examples B7 to B12 and the compositions described in Examples B27 à B33 are injected without any difficulty, both with insulin syringes equipped with 31-gauge needles and with insulin pens from Novo Nordisk, sold under the name of Novopen®, equipped with 31-gauge needles.

15 Example B37: Protocol for measuring the pharmacodynamics of the insulin solutions. Preclinical studies were carried out on pigs for the purpose of evaluating two compositions according to the invention:
 [000384] Lantus® / Apidra® (75/25), formulated with Polysaccharide 4 (6 mg/ml), described in Example B7, and
 20 [000385] Lantus® / Humalog® (75/25), formulated with Polysaccharide 4 (6 mg/ml), described in Example B8.

25 [000386] The hypoglycaemic effects of these compositions were compared with respect to injections carried out with simultaneous but separate injections of Lantus® (pH 4) and then of a prandial insulin Apidra® or Humalog®.

[000387] Six domesticated pigs weighing approximately 50 kg, catheterized beforehand in the jugular vein, are deprived of food for 2 to 3 hours before the beginning of the experiment. In the hour preceding the injection of insulin, three blood samples are taken in order to determine the basal glucose level.

5 [000388] Injection of insulin at a dose of 0.4 IU/kg is carried out by subcutaneous injection into the neck, under the ear of the animal, using the Novopen® insulin pen equipped with a 31 G needle.

[000389] Blood samples are subsequently taken after 4, 8, 12, 16, 20, 30, 40, 50, 60, 90, 120, 240, 360, 480, 600, 660 and 720 minutes. After each sample has been

10 taken, the catheter is rinsed with a dilute heparin solution.

[000390] A drop of blood is taken in order to determine the glycaemia using a glucose meter. The pharmacodynamic curves of glucose are subsequently plotted.

[000391] The results obtained are presented in the form of pharmacodynamic curves for glucose represented in Figures 1 to 6.

15 [000392] Lantus® / Apidra® (75/25), formulated with Polysaccharide 4 (6 mg/ml).

[000393] Figure 1: Mean + standard deviation of the mean curves for the sequential administrations of Apidra® and Lantus®, in comparison with a composition according to the invention Polysaccharide 4/ Lantus® /Apidra® (75/25).

[000394] Figure 2: Apidra® Lantus® individual curves.

20 [000395] Figure 3: Polysaccharide 4/Apidra®/Lantus® individual curves.

[000396] Figure 1 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the pigs tested for each formulation. The drop in glycaemia in the first 30 minutes is similar for the two formulations, indicating that the presence of a polysaccharide does not interfere with the rapid-acting nature of Apidra®.

25 [000397] On the other hand, between 90 min and 10 h (600 minutes), the sequential administration of Apidra® and Lantus® brings about a heterogeneous drop in glucose with a homogeneous plateau response with regard to three pigs and a heterogeneous response with regard to the other three pigs (Figure 2). In contrast, the six pigs tested with the Polysaccharide 4/Apidra®/Lantus® formulation have a homogeneous response

30 (Figure 3). This is reflected by the analysis of the coefficients of variation (CV) between 60 min and 10 h, which are on average 54% (between 21% and 113%) for the Apidra® Lantus® control and 12% (between 5% and 25%) for Polysaccharide 4/Apidra®/Lantus®.

35 [000398] Lantus®/Humalog® (75/25), formulated with Polysaccharide 4 (6 mg/ml).

[000399] Figure 4: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® and Lantus® in comparison with the administration of a composition according to the invention Polysaccharide 4/Humalog®/Lantus®.

[000400] Figure 5: Humalog® Lantus® individual curves.

[000401] Figure 6: Polysaccharide 4/Humalog®/Lantus® individual curves.

[000402] Figure 4 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the pigs tested with regard to each formulation. The drop in

5 glycaemia in the first 30 minutes is similar for the two formulations, indicating that the presence of Polysaccharide 4 does not interfere with the rapid-acting nature of Humalog®. On the other hand, between 60 min and 8 h (480 minutes), the sequential administration of Humalog® and Lantus® brings about a heterogeneous drop in glucose with a homogeneous plateau response with regard to four pigs and a

10 heterogeneous response with regard to the other two pigs (Figure 5). In contrast, the 5 pigs tested with the Polysaccharide 4/Humalog®/Lantus® formulation have a homogeneous response (Figure 6). This is reflected by the analysis of the coefficients of variation (CV) with regard to the data for drop in glycaemia between 60 min and 8 h, which are on average 54% (between 31% and 72%) for the Humalog® Lantus®

15 control and 15% (between 6% and 28%) for Polysaccharide 4/Humalog®/Lantus®. The presence of Polysaccharide 4 thus greatly reduced the variability of Lantus® with regard to the drop in glycaemia.

Example B38: Protocol for measuring the pharmacodynamics of the insulin solutions

20 Preclinical studies were carried out on dogs for the purpose of evaluating six compositions according to the invention:

[000403] The hypoglycaemic effects of these compositions were compared with respect to injections carried out with simultaneous but separate injections of 100 IU/ml Lantus® (pH 4) and then of a prandial insulin 100 IU/ml Humalog®.

25

[000404] Ten domesticated dogs (beagles) weighing approximately 12 kg are deprived of food for 18 hours before the beginning of the experiment. In the hour preceding the injection of insulin, three blood samples are taken in order to determine the basal glucose level.

30 [000405] The injection of insulin at a dose of 0.53 IU/kg (unless otherwise mentioned in the examples below) is carried out by subcutaneous injection into the neck of the animal using the Novopen® insulin pen equipped with a 31 G needle.

[000406] Blood samples are subsequently taken after 10, 20, 30, 40, 50, 60, 90, 120,

35 180, 240, 300, 360, 420, 480, 540, 600, 660, 720, 780, 840, 900 and 960 minutes. The first samples are taken with a catheter (up to 180 min) and then directly from the jugular vein. After each sample has been taken, the catheter is rinsed with a dilute heparin solution.

[000407] A drop of blood is taken in order to determine the glycaemia by means of a glucose meter. The pharmacodynamic curves for glucose are subsequently plotted.

[000408] The results obtained are presented in the form of pharmacodynamic curves for glucose represented in Figures 7 to 12.

5

The solution of Example B28

[000409] Figure 7: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg), in comparison with a composition according to the invention described in Example B28 (0.53 IU/kg).

[000410] Figure 7 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the dogs tested for each formulation. The two curves are similar up to 12 hours with a rapid drop in glycaemia, indicating that the polysaccharide does not influence the rapid-acting effect of Humalog®, a marked return between the peak due to Humalog® and the plateau due to glargine, and then a plateau of the glargine up to 12 h, indicating that the basal effect of glargine is well retained.

The solution of Example B27

[000411] Figure 8: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) in comparison with a composition according to the invention described in Example B27 (0.47 IU/kg).

[000412] Figure 8 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the dogs tested for each formulation. In this comparison, the dose of basal insulin (Lantus®) is identical whereas the dose of Humalog® is half for the composition, with respect to the control. The drop in glucose is greater in the case of the formulation B27 in comparison with the control, whereas this control contains twice as much Humalog®. On the other hand, the duration of the plateau is shorter in the case of the combination, with respect to the control. This indicates that, in this composition, a portion of Lantus® is not precipitated on injection and acts with Humalog®.

The solution of Example B29

[000413] Figure 9: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) in comparison with a composition according to the invention described in Example B29 (0.53 IU/kg).

[000414] Figure 9 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the dogs tested for each formulation. The two curves are similar, with a rapid drop in glycaemia, indicating that the polysaccharide does not influence the rapid-acting effect of Humalog®, a marked return between the peak due to Humalog® and the plateau due to Lantus®, and then a plateau of the Lantus® up to 13 h, indicating that the basal effect of glargin is well retained.

The solution of Example B31

[000415] Figure 10: Mean + standard deviation of the mean curves for the sequential 10 administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg), in comparison with a composition according to the invention described in Example B31 (0.48 IU/kg).

[000416] Figure 10 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the dogs tested for each formulation. In this comparison, 15 the dose of basal insulin (Lantus®) is identical, whereas the dose of Humalog® is half, for the composition, with respect to the control. The drop in glucose is greater in the case of the control, with respect to the combination corresponding to Example B31. This response was expected, in view of the concentration of Humalog® in the combination, which was half that of the control. Furthermore, the duration of the 20 Lantus® plateau is identical in the case of the combination, with respect to the control. This indicates that, in this composition and by comparison with the composition described in Example B29 (Figure 9), it is possible to adjust the amount of Humalog® in the combination without modifying the Lantus® basal effect.

25 The solution of Example B30

[000417] Figure 11: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.24 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) in comparison with a composition according to the invention described in Example B30 (0.64 IU/kg).

30 [000418] Figure 11 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the dogs tested for each formulation. The two curves are similar, with a rapid drop in glycaemia, indicating that the polysaccharide does not influence the rapid-acting effect of Humalog®, a marked return between the peak due to Humalog® and the plateau due to Lantus®, and then a plateau of the Lantus® up to 35 10 h, indicating that the glargin basal effect is well retained.

The solution of Example B32

[000419] Figure 12: Mean + standard deviation of the mean curves for the sequential administrations of Humalog® (100 IU/ml, 0.13 IU/kg) and Lantus® (100 IU/ml, 0.4 IU/kg) in comparison with a composition according to the invention described in Example B32 (0.53 IU/kg).

[000420] Figure 12 presents the mean curves for drop in glycaemia and the standard deviations of the mean for the dogs tested for each formulation. The two curves are similar up to 10 hours, with a rapid drop in glycaemia, indicating that the polysaccharide does not influence the rapid-acting effect of Humalog®, a marked return between the peak due to Humalog® and the plateau due to Lantus®, and then a glargin plateau, indicating that the glargin basal effect is retained up to 10 h.

In conclusion, Figures 7 to 12 show that, by adjusting the composition of the polysaccharide and the concentrations of lispro and glargin, it is possible to obtain profiles identical to a double injection with different proportions of rapid-acting insulin and basal insulin. It is also possible to adjust the duration of the basal insulin without impacting the rapid-acting insulin or to adjust the amount of rapid-acting insulin without impacting the effect of the basal insulin.

20 [000421] Examples

Part C: Demonstration of the properties of the compositions comprising a GLP-1 analogue or derivative according to the invention

Example C1: 0.25 mg/ml solution of GLP-1 analogue Exenatide (Byetta®)

[000422] This solution is an exenatide solution marketed by the company Eli Lilly and Company under the name of Byetta®.

Example C2: 6 mg/ml solution of GLP-1 derivative Liraglutide (Victoza®)

[000423] This solution is a liraglutide solution marketed by the company Novo Nordisk under the name of Victoza®.

30

Example C3: Dissolution of Lantus® at 100 IU/ml and at pH 7 using a substituted dextran at a concentration of 10 mg/ml

[000424] 20 mg of a substituted dextran chosen from those described in Table 1 are weighed out accurately. This lyophilisate is taken up in 2 ml of the insulin glargin solution of Example B4 in order to obtain a solution whose polysaccharide concentration is equal to 10 mg/ml. After mechanically stirring on rolls at ambient temperature, the solution becomes clear. The pH of this solution is 6.3. The pH is

adjusted to 7 with a 0.1N sodium hydroxide solution. This clear solution is filtered through a membrane (0.22 µm) filter and is then placed at +4°C.

[000425] Generalization: Clear solutions of Lantus at 100 IU/ml and at pH 7 were also obtained with concentrations of substituted dextrans at 20 and 40 mg/ml by following the same protocol as that described in Example C3. Thus, a weight of lyophilized polysaccharide among those described in Table 1 is weighed out accurately. This lyophilisate is taken up in the insulin glargine solution of Example B4, so as to obtain a solution whose concentration of substituted dextran is 20 or 40 mg/ml, as described in Table 8. After mechanical stirring on rolls at ambient temperature, the solution becomes clear. The pH of this solution is less than 7. The pH is subsequently adjusted to 7 with a 0.1N sodium hydroxide solution. This clear final solution is filtered through a membrane (0.22 µm) and is then placed at +4°C.

Final concentration of substituted dextran (mg/ml)	Weight of substituted dextran weighed out (mg)	Volume of the insulin glargine solution of Example B4 added (ml)
10	20	2
20	40	2
40	80	2

Table 8: Preparation of a solution of Lantus® at 100 IU/ml and at pH 7 using a substituted dextran at a concentration of 10, 20 or 40 mg/ml

Example C4: Preparation of a Lantus®/Byetta® 70/30 composition at pH 7.5

[000426] 0.09 ml of the exenatide solution of Example C1 is added to 0.21 ml of the insulin glargine solution of Example B4, in order to obtain 0.3 ml of composition whose pH is 4.5 on mixing. The composition, which contains 70 IU/ml of Lantus® and 0.075 mg/ml of Byetta®, is clear, testifying to the good solubility of Lantus® and Byetta® under these formulation conditions (pH 4.5). The pH is subsequently adjusted to 7.5 with a 0.1N sodium hydroxide solution. The composition then becomes cloudy, testifying to the poor solubility of the Lantus®/Byetta® composition at pH 7.5.

[000427] 70/30 Lantus®/Byetta® compositions were also prepared at pH 4.5, 5.5, 6.5, 8.5 and 9.5 by following a protocol similar to that described in Example C4. For each of these compositions, 0.09 ml of the exenatide solution of Example C1 is added to 0.21 ml of the insulin glargine solution of Example B4, in order to obtain 0.3 ml of a composition whose pH is 4.5 on mixing. The composition is clear, testifying to the good solubility of Lantus® and Byetta® under these formulation conditions (pH 4.5). The pH is adjusted to 5.5 or 6.5 or 8.5 or 9.5 with a 0.1N sodium hydroxide solution.

After adjusting the pH, the composition at 5.5 is slightly cloudy, the compositions at 6.5 – 7.5 and 8.5 are very cloudy and the composition at pH 9.5 is clear. These compositions are placed at +4°C for 48 h. After 48 h at +4°C, only the composition at pH 4.5 remains clear. The visual appearance after 48 h of the 70/30 Lantus®/Byetta® 5 compositions at different pH values is summarized in Table 9.

70/30 Lantus®/Byetta® compositions at different pH values	
pH	Visual appearance at t = 48h
4.5	Clear
5.5	Presence of a precipitate
6.5	Presence of a precipitate
7.5	Presence of a precipitate
8.5	Presence of a precipitate
9.5	Presence of a precipitate

Table 9: Visual appearance after 48 h of the 70/30 Lantus®/Byetta® compositions at different pH values

10 Example C5: Preparation of the 70/30 Lantus®/Victoza® composition at pH 7.5 [000428] 0.09 ml of the liraglutide solution of Example C2 is added to 0.21 ml of the insulin glargine solution of Example B4, in order to obtain 0.3 ml of a composition whose pH is 7 on mixing. The composition, which contains 70 IU/ml of glargine and 15 1.8 mg/ml of exenatide, is cloudy, testifying to the poor solubility of the Lantus®/Victoza® composition under these formulation conditions. The pH is adjusted to 7.5 with a 0.1N sodium hydroxide solution. After adjusting the pH, the composition remains cloudy. This composition is placed at +4°C for 48 h.

[000429] 70/30 Lantus®/Victoza® compositions were also prepared at pH 4.5 – 5.5 – 20 6.5 – 8.5 and 9.5 by following a protocol similar to that described in Example C5. For each of these compositions, 0.09 ml of the liraglutide solution of Example C1 is added to 0.21 ml of the insulin glargine solution of Example B4, in order to obtain 0.3 ml of a composition whose pH is 7. The composition is cloudy, testifying to the poor solubility of the Lantus®/Victoza® composition under these formulation conditions (pH 7). The 25 pH is adjusted to 4.5 or 5.5 or 6.5 with a 0.1N hydrochloric acid solution or to pH 9,5 with a 0.1N sodium hydroxide solution. After adjusting the pH, the compositions at pH 4.5 – 5.5 and 6.5 are cloudy, testifying to the poor solubility of the Lantus®/Victoza® composition under these formulation conditions. These compositions are placed at +4°C for 48 h. After 48 h at 4°C, only the composition at pH 9.5 is clear. The visual

appearance after 48 h of the 70/30 Lantus®/Victoza® compositions of different pH values is summarized in Table 10.

70/30 Lantus®/Victoza® compositions at different pH values	
pH	Visual appearance at t = 48h
4.5	Presence of a precipitate
5.5	Presence of a precipitate
6.5	Presence of a precipitate
7.5	Presence of a precipitate
8.5	Presence of a precipitate
9.5	Clear

Table 10: Visual appearance after 48 h of the 70/30 Lantus®/Victoza® compositions at different pH values

5

Example C6: Preparation of a substituted dextran-70/30 Lantus®/Byetta® composition at pH 7

[000430] 0.09 ml of the exenatide solution of Example C1 is added to 0.21 ml of the substituted dextran/Lantus® solution prepared in Example C3, in order to obtain 0.3 ml of a composition at pH 5.3. The pH is adjusted to 7 with a 0.1N sodium hydroxide solution. The composition, which contains 7 mg/ml of polysaccharide, 70 IU/ml of Lantus® and 0.075 mg/ml of Byetta® is clear, testifying to the good solubility of Lantus® and Byetta® in the presence of the substituted dextran at pH 7.

15 This clear solution is placed at +4°C.

[000431] Generalization: Substituted dextran-Lantus®/Byetta® compositions at pH 7 were also prepared at ratios by volume $V_{\text{Lantus}}/V_{\text{Byetta}}$ of 90/10, 50/50, 30/70 and 10/90 by following the same protocol as that described in Example C6. Thus, a volume V_{Byetta} 20 of the exenatide solution of Example C1 is added to a volume V_{Lantus} of the substituted dextran/Lantus® solution prepared in Example C3, in order to obtain a composition whose pH is adjusted at 7 with a 0.1N sodium hydroxide solution. The compositions obtained (see Table 11) are clear, testifying to the good solubility of Lantus® and Byetta® in the presence of a substituted dextran at pH 7. These clear solutions are 25 placed at +4°C.

Example C7: Preparation of a substituted dextran-100/50 Lantus®/Byetta® composition at pH 7

[000432] 0.150 ml of the exenatide solution of Example C1 is lyophilized and then 30 0.3 ml of a substituted dextran/Lantus® solution prepared in Example C3 are added to

the lyophilisate in order to obtain a composition whose pH is adjusted to 7 with a 0.1N sodium hydroxide solution. The composition, which contains 10 mg/ml of polysaccharide, 100 IU/ml of Lantus® and 0.125 mg/ml of Byetta®, is clear, testifying to the good solubility of Lantus® and Byetta® in the presence of the substituted dextran at pH 7. This clear solution is placed at +4°C.

	Lantus®		[Polysaccharide No.] (mg/ml)	Byetta® (mg/ml)
	IU/ml	mg/ml		
100/50	100	3.5	10	0.125
90/10	90	3.15	9	0.025
70/30	70	2.45	7	0.075
50/50	50	1.75	5	0.125
30/70	30	1.05	3	0.175

Table 11: Final concentrations of Lantus®, substituted dextran and Byetta® of the compositions obtained in Examples C6 and C7

10 Example C8: Preparation of a substituted dextran-70/30 Lantus®/Victoza® composition at pH 7
 [000433] 0.09 ml of the liraglutide solution of Example C2 is added to 0.21 ml of the substituted dextran/Lantus® solution prepared in Example C3, in order to obtain 0.3 ml of a composition at pH 7.6. The pH is adjusted to 7 with a 0.1N hydrochloric acid solution. The composition, which contains 7 mg/ml of polysaccharide, 70 IU/ml of Lantus® and 1.8 mg/ml of Victoza®, is clear, testifying to the good solubility of Lantus® and Victoza® in the presence of the substituted dextran at pH 7. This clear solution is placed at +4°C.

15
 20 [000434] Generalization: Substituted dextran-Lantus®/Victoza® compositions at pH 7 have also been prepared at $V_{\text{Lantus}}/V_{\text{Victoza}}$ ratios by volume of 90/10, 50/50,

30/70, and 90/10 by following the same protocol as that described in Example C6. Thus, a volume V_{Victoza} of the liraglutide solution of Example C2 is added to a volume V_{Lantus} of the substituted dextran/Lantus® solution prepared in Example B3, in order to obtain a composition whose pH is adjusted to 7 with a 0.1N hydrochloric acid solution.

5 [000435] The compositions obtained (see Table 12) are clear, testifying to the good solubility of Lantus® and Victoza® in the presence of a substituted dextran at pH 7. These clear solutions are placed at +4°C.

10 Example C9: Preparation of a substituted dextran-100/50 Lantus®/Victoza® composition at pH 7

15 [000436] 0.150 ml of the liraglutide solution of Example C2 is lyophilized and then 0.3 ml of a substituted dextran/Lantus® solution prepared in Example C3 is added to the lyophilisate, in order to obtain a composition whose pH is adjusted to 7 with a 0.1N sodium hydroxide solution. The composition, which contains 10 mg/ml of polysaccharide, 100 IU/ml of Lantus® and 3 mg/ml of Victoza®, is clear, testifying to the good solubility of Lantus® and Victoza® in the presence of the substituted dextran at pH 7. This clear solution is placed at +4°C.

	Lantus®		[Polysaccharide No.] (mg/ml)	Victoza® (mg/ml)
	IU/ml	mg/ml		
100/50	100	3.5	10	3
90/10	90	3.15	9	0.6
70/30	70	2.45	7	1.8
50/50	50	1.75	5	3
30/70	30	1.05	3	4.2

20 Table 12: Final concentrations of Lantus®, substituted dextran and Victoza® in the composition obtained in Examples C8 and C9

Example C10: Preparation of a substituted dextran-60/20/20 Lantus®/Apidra®/Byetta® composition at pH 7

25 [000437] 20 mg of lyophilized Polysaccharide 4 described in Example A3 are weighed out accurately. This lyophilisate is taken up in 2 ml of the insulin glargine solution of Example B4. After mechanical stirring on rolls at ambient temperature, the solution becomes clear. The pH of this solution is 6.3. The pH is adjusted to 7 with a 0.1N sodium hydroxide solution. 0.2 ml of the exenatide solution of Example C1 and 0.2 ml of the insulin glulisine solution of Example B3 are added to 0.6 ml of the substituted dextran/Lantus® solution prepared above, in order to obtain 1 ml

[000438] of a composition at pH 7. The composition, which contains 6 mg/ml of polysaccharide, 60 IU/ml of Lantus®, 20 IU/ml Apidra® and 0.05 mg/ml of Byetta®, is clear, testifying to the good solubility of Lantus®, Apidra® and Byetta® in the presence of substituted dextran at pH 7. This clear solution is filtered through a membrane 5 (0.22 µm) and is then placed at +4°C.

Example C11: Precipitation of Lantus®

[000439] 0.250 ml of Lantus® is added to 0.5 ml of a PBS (Phosphate Buffer Solution) 10 solution containing 20 mg/ml of BSA (Bovine Serum Albumin). The PBS/BSA mixture simulates the composition of the subcutaneous medium.

[000440] A precipitate appears, which is in good agreement with the mechanism of operation of Lantus® (precipitation on injection due to the increase in the pH).

[000441] Centrifuging at 4000 rev/min is carried out in order to separate the 15 precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. The result of this is that 90% of Lantus® is found in a precipitated form.

Example C12: Precipitation of a substituted dextran/Lantus® composition

[000442] 0.250 ml of substituted dextran/Lantus® solution prepared in Example C3 is 20 added to 0.5 ml of a PBS solution containing 20 mg/ml of BSA. The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

[000443] Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. The result of this is that 90% of Lantus® is found in a precipitated form. 25 This percentage of precipitation of Lantus® is identical to that obtained for the control described in Example C11.

Example C13: Precipitation of a substituted dextran-Lantus®/Byetta® composition

[000444] 0.250 ml of the substituted dextran-Lantus®/Byetta® composition prepared 30 in Example C6 is added to 0.5 ml of a PBS solution containing 20 mg/ml of BSA. The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

[000445] Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® and Byetta® are assayed in the supernatent. The percentage of precipitation of Lantus® is similar to the control 35 described in Example C11.

Example C14: Precipitation of a substituted dextran-70/30 Lantus®/Victoza® composition

[000446] 0.250 ml of the substituted dextran-Lantus®/Victoza® composition prepared in Example C8 is added to 0.5 ml of a PBS solution containing 20 mg/ml of BSA (bovine serum albumin). The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® and Victoza® are assayed in the supernatent.

10 The percentage of precipitation of Lantus® is similar to the control described in Example C11.

Example C15: Precipitation of different compositions, the nature of the substituted dextran being varied

15 [000447] Other tests under the same conditions as those of Examples C13 and C14 were carried out in the presence of other dextrans.

[000448] Results with at most 20 mg/ml of substituted dextran and a 70/30 Lantus®/Byetta® composition are combined in the following Table 13. It is observed that the dissolution and the precipitation of Lantus® are retained.

20

Polysaccharide No.	Dissolution 70/30 Lantus®/Byetta®	Percentage of precipitation of Lantus®
1	Yes	94
2	Yes	96
5	Yes	88
7	Yes	95
10	Yes	Not measured
11	Yes	81
14	Yes	Not measured
16	Yes	96
26	Yes	81
27	Yes	96
28	Yes	96
29	Yes	95

Table 13: Results of the dissolution and precipitation tests obtained with at most 20 mg/ml of substituted dextran and a 70/30 Lantus®/Byetta® composition

[000449] Results with at most 20 mg/ml of substituted dextran and various Lantus®/Byetta® compositions are combined in the following Table 14. It is observed that the dissolution and the precipitation of Lantus® are retained.

Polysaccharide No.	Ratio Lantus®/Byetta®	Dissolution Lantus®/Byetta®	Percentage of precipitation of Lantus®
4	100/50	Yes	95
4	90/10	Yes	94
4	70/30	Yes	95
4	50/50	Yes	90
4	30/70	Yes	82
8	100/50	Yes	96
8	90/10	Yes	94
8	70/30	Yes	96
8	50/50	Yes	90
8	30/70	Yes	81

5 Table 14: Results of the dissolution and precipitation tests obtained with at most
20 mg/ml of substituted dextran and various Lantus®/Byetta® compositions

[000450] Results with at most 40 mg/ml of substituted dextran and a 70/30 Lantus®/Victoza® composition are combined in the following Table 15. It is observed
10 that the dissolution and the precipitation of Lantus® are retained.

[000451]

Polysaccharide No.	Dissolution 70/30 Lantus®/Victoza®	Percentage of precipitation of Lantus®
1	Yes	95
2	Yes	97
5	Yes	Not measured
7	Yes	97
10	Yes	Not measured
11	Yes	Not measured
14	Yes	90
16	Yes	97
26	Yes	74
27	Yes	96
28	Yes	95
29	Yes	94

Table 15: Results of the dissolution and precipitation tests obtained with at most 40 mg/ml of substituted dextran and a 70/30 Lantus®/Victoza® composition

5

[000452] Results with at most 20 mg/ml of substituted dextran and various Lantus®/Victoza® compositions are combined in the following Table 16. It is observed that the dissolution and the precipitation of Lantus® are retained.

Polysaccharide No.	Ratio Lantus®/Victoza®	Dissolution Lantus®/Victoza®	Percentage of precipitation of Lantus®
4	90/10	Yes	94
4	70/30	Yes	Not measured
4	50/50	Yes	90
4	30/70	Yes	86
8	100/50	Yes	93
8	90/10	Yes	95
8	70/30	Yes	98
8	50/50	Yes	89
8	30/70	Yes	85

10 Table 16: Results of the dissolution and precipitation tests obtained with at most 20 mg/ml of substituted dextran and various Lantus®/Victoza® compositions

Example C16: Precipitation of a substituted dextran-60 /20/20 Lantus®/Apidra®/Byetta® composition at pH 7.

[000453] 0.250 ml of the substituted dextran-Lantus®/Apidra®/Byetta® composition prepared in Example C10 is added to 0.5 ml of a PBS solution containing 20 mg/ml of BSA. The PBS/BSA mixture simulates the composition of the subcutaneous medium. A precipitate appears.

Centrifuging at 4000 rev/min is carried out in order to separate the precipitate from the supernatent. Subsequently, Lantus® is assayed in the supernatent. The percentage of precipitation of Lantus® is similar to the control described in Example C11.


[000454] In this specification where a document, act or item of knowledge is referred to or discussed, this reference or discussion is not an admission that the document, act or item of knowledge or any combination thereof was at the priority date publicly available, known to the public, part of the common general knowledge or known to be relevant to an attempt to solve any problem with which this specification is concerned.

[000455] The word 'comprising' and forms of the word 'comprising' as used in this description and in the claims does not limit the invention claimed to exclude any variants or additions.

CLAIMS

1. Composition in the form of an injectable aqueous solution, the pH of which is between 6.0 and 8.0, comprising at least:


- a) a basal insulin, the isoelectric point pI of which is between 5.8 and 8.5;
- b) a dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals of formula I or of formula II:



in which:

- R is $-\text{OH}$ or chosen from the group consisting of the radicals:
 - $-(f\text{-}[A]\text{-COOH})_n$;
 - $-(g\text{-}[B]\text{-}k\text{-}[D])_m$, D comprising at least one alkyl chain comprising at least 8 carbon atoms;
- n represents the degree of substitution of the glucoside units by $-\text{f-}[A]\text{-COOH}$ and $0.1 \leq n \leq 2$;
- m represents the degree of substitution of the glucoside units by $-\text{g-}[B]\text{-}k\text{-}[D]$ and $0 < m \leq 0.5$;
- q represents the degree of polymerization as glucoside units, that is to say the mean number of glucoside units per polysaccharide chain, and $3 \leq q \leq 50$;
- $-(f\text{-}[A]\text{-COOH})_n$:
 - -A- is a linear or branched radical comprising from 1 to 4 carbon atoms; the said -A- radical:
 - being bonded to a glucoside unit via a functional group f chosen from the group consisting of ether, ester and carbamate functional groups;
- $-(g\text{-}[B]\text{-}k\text{-}[D])_m$:
 - -B- is a linear or branched, at least divalent, radical comprising from 1 to 4 carbon atoms; the said -B- radical:
 - being bonded to a glucoside unit via a functional group g chosen from the group consisting of ether, ester and carbamate functional groups;

- being bonded to a -D radical via a functional group *k*; *k* chosen from the group consisting of ester, amide and carbamate functional groups; the said -D radical:
 - being an $-X(-I-Y)_p$ radical, *X* being an at least divalent radical comprising from 1 to 12 atoms chosen from the group consisting of C, N and O atoms, optionally carrying carboxyl or amine functional groups and/or resulting from an amino acid, a dialcohol, a diamine or a mono- or polyethylene glycol mono- or diamine; *Y* being a linear or cyclic alkyl group, an alkylaryl or an arylalkyl, of 8 to 30 carbon atoms, optionally substituted by one or more C_1 to C_3 alkyl groups; $p \geq 1$ and *I* a functional group chosen from the group consisting of ester, amide and carbamate functional groups;
 - *f*, *g* and *k* being identical or different;
 - the free acid functional groups being in the form of salts of alkali metal cations chosen from the group consisting of Na^+ and K^+ ;
 - and, when $p = 1$, if *Y* is a C_8 to C_{14} alkyl, then $q^*m \geq 2$, if *Y* is a C_{15} alkyl, then $q^*m \geq 2$; and if *Y* is a C_{16} to C_{20} alkyl, then $q^*m \geq 1$;
 - and, when $p \geq 2$, if *Y* is a C_8 to C_9 alkyl, then $q^*m \geq 2$ and, if *Y* is a C_{10} to C_{16} alkyl, then $q^*m \geq 0.2$;



in which:

- *R* is $-OH$ or a $-(f-[A]-COOH)_n$ radical:
 - $-A-$ is a linear or branched radical comprising from 1 to 4 carbon atoms; the said radical $-A-$:
 - being bonded to a glucoside unit *via* a functional group *f* chosen from the group consisting of ether, ester or carbamate functional groups;
 - *n* represents the degree of substitution of the glucoside units by $-f-[A]-COOH$ and $0.1 \leq n \leq 2$;
- *R'* is chosen from the group consisting of the radicals:
 - $-C(O)NH-[E]-(o-[F])_t$;
 - $-CH_2N(L)_z-[E]-(o-[F])_t$;

in which:

- z is a positive integer equal to 1 or 2,
- L is chosen from the group consisting of:
 - -H and z is equal to 1, and/or
 - -[A]-COOH and z is equal to 1 or 2, if f is an ether functional group,
 - -CO-[A]-COOH and z is equal to 1 if f is an ester functional group, and
 - -CO-NH-[A]-COOH and z is equal to 1 if f is a carbamate functional group;
- -[E]-(o-[F])_t:
 - -E- is a linear or branched, at least divalent, radical comprising from 1 to 8 carbon atoms and optionally comprising heteroatoms, such as O, N or S;
 - -F is a linear or cyclic alkyl group, an alkylaryl or an arylalkyl, of 12 to 30 carbon atoms, optionally substituted by one or more C₁ to C₃ alkyl groups;
 - o is a functional group chosen from the group consisting of ether, ester, amide or carbamate functional groups;
 - t is a positive integer equal to 1 or 2;
- q represents the degree of polymerization as glucoside units, that is to say the mean number of glucoside units per polysaccharide chain, and 3 ≤ q ≤ 50;
- the free acid functional groups being in the form of salts of alkali metal cations chosen from the group consisting of Na⁺ and K⁺;
- when z = 2, the nitrogen atom is in the form of a quaternary ammonium.

2. Composition according to Claim 1, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula I.


3. Composition according to Claim 1, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula II.

4. Composition according to either one of Claims 1 and 2, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula I in which the -(f-[A]-COOH)_n radical is

chosen from the group consisting of the following sequences, *f* having the meaning given above:



5. Composition according to any one of Claims 1, 2 and 4, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula I in which the $-(g-[B]-k-[D])_m$ radical is chosen from the group consisting of the following sequences, *g*, *k* and *D* having the meanings given above:



6. Composition according to any one of Claims 1, 2 and 4, 5, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula I in which the $-(g-[B]-k-[D])_m$ radical is such that:

- $-B-$ is a radical comprising one carbon atom; the said $-B-$ radical being bonded to a glucoside unit *via* an ether functional group *g*, and *X* is a radical resulting from an amino acid.

7. Composition according to any one of Claims 1, 2 and 4 to 6, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula I in which the *X* radical is an at least divalent radical resulting from an amino acid chosen from the group consisting of glycine, leucine, phenylalanine, lysine, isoleucine, alanine, valine, aspartic acid and glutamic acid.

8. Composition according to any one of Claims 1, 2 and 4 to 7, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula I in which the *Y* group is chosen from the group consisting of a hydrophobic alcohol, a hydrophobic acid, a sterol or a tocopherol.

9. Composition according to any one of Claims 1, 2 and 4 to 8, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic

radicals is chosen from the dextrans of formula I in which the Y group is a sterol chosen from cholesterol derivatives.

10. Composition according to either one of Claims 1 and 3, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula II in which the R' group is such that the -E- radical results from a diamine.

11. Composition according to any one of Claims 1, 3 and 10, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the dextrans of formula II in which the R' group is such that the -F group results from a cholesterol derivative.

12. Composition according to any one of Claims 1, 2 and 4 to 9, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the group consisting of the following dextrans of formula I:

- Sodium dextranmethylcarboxylate modified by octyl glycinate,
- Sodium dextranmethylcarboxylate modified by cetyl glycinate,
- Sodium dextranmethylcarboxylate modified by octyl phenylalaninate,
- Sodium dextranmethylcarboxylate modified by 3,7-dimethyl-1-octyl phenylalaninate,
- Sodium dextranmethylcarboxylate modified by dioctyl aspartate,
- Sodium dextranmethylcarboxylate modified by didecyl aspartate,
- Sodium dextranmethylcarboxylate modified by dilauryl aspartate,
- Sodium dextranmethylcarboxylate modified by N-(2-aminoethyl)dodecanamide,
- Sodium dextransuccinate modified by lauryl glycinate,
- N-(sodium methylcarboxylate) dextran carbamate modified by dioctyl aspartate,
- Sodium dextranmethylcarboxylate modified by dilauryl aspartate,
- Sodium dextranmethylcarboxylate modified by 2-(2-aminoethoxy)ethyl dodecanoate,
- Sodium dextranmethylcarboxylate modified by 2-(2-{2-[dodecanoylamino]ethoxy}ethoxy)ethylamine,

- Sodium dextranmethylcarboxylate modified by 2-(2-{2-[hexadecanoylamino]ethoxy}ethoxy)ethylamine,
- Sodium dextranmethylcarboxylate modified by cholesteryl leucinate,
- Sodium dextranmethylcarboxylate modified by cholesteryl 1-ethylenediaminecarboxylate,
- N-(sodium methylcarboxylate) dextran carbamate modified by cholesteryl leucinate.

13. Composition according to any one of Claims 1, 3, 10 and 11, characterized in that the dextran substituted by radicals carrying carboxylate charges and hydrophobic radicals is chosen from the group consisting of the dextrans of formula II and is:

- Sodium dextranmethylcarboxylate modified by cholesteryl 1-ethylenediaminecarboxylate grafted by reductive amination to the reducing chain end.

14. Composition according to any one of the preceding claims, characterized in that the basal insulin whose isoelectric point is between 5.8 and 8.5 is insulin glargine.

15. Composition according to any one of the preceding claims, characterized in that it comprises between 40 IU/ml and 500 IU/ml of basal insulin whose isoelectric point is between 5.8 and 8.5.

16. Composition according to any one of the preceding claims, characterized in that it additionally comprises a prandial insulin.

17. Composition according to any one of the preceding claims, characterized in that it comprises between 40 and 800 IU/ml of total insulin.

18. Composition according to either one of Claims 16 and 17, characterized in that it comprises proportions, expressed as percentage, between the basal insulin whose isoelectric point is between 5.8 and 8.5 and a prandial insulin of 25/75, 30/70, 40/60, 50/50, 60/40, 70/30, 80/20 and 90/10.

19. Composition according to any one of Claims 1 to 15, characterized in that it additionally comprises a GLP-1, a GLP-1 analogue or a GLP-1 derivative.

20. Composition according to any one of the preceding claims, characterized in that it additionally comprises zinc salts at a concentration of between 0 and 5000 μ M.

21. Composition according to any one of the preceding claims, characterized in that it comprises buffers chosen from the group consisting of Tris, citrates and phosphates at concentrations of between 0 and 100 mM, preferably between 0 and 50 mM.

22. Composition according to any one of Claims 16 to 18, characterized in that the said prandial insulin is chosen from the group formed by human insulin, insulin glulisine, insulin lispro and insulin aspart.

23. Single-dose formulation comprising a composition according to any one of Claims 1 to 18 and 20 to 22, at a pH of between 6.6 and 7.8, and a prandial insulin.

24. Single-dose formulation comprising a composition according to any one of Claims 1 to 15 and 19 to 21, at a pH of between 6.6 and 7.8, and a GLP-1, a GLP-1 derivative or a GLP-1 analogue.

25. Single-dose formulation according to Claim 23, characterized in that the prandial insulin is chosen from the group comprising of human insulin.

26. Single-dose formulation according to Claim 23, characterized in that the prandial insulin is chosen from the group comprising of insulin lispro, insulin glulisine and insulin aspart.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

4/6




Figure 7




Figure 8

5/6




Figure 9




Figure 10

6/6




Figure 11




Figure 12