


(51) International Patent Classification:  
*B01D 15/38* (2006.01) *C07K 1/16* (2006.01)(21) International Application Number:  
PCT/US2013/066319(22) International Filing Date:  
23 October 2013 (23.10.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/717,880 24 October 2012 (24.10.2012) US(71) Applicant: GENZYME CORPORATION [US/US]; 500  
Kendall Street, Cambridge, Massachusetts 02142 (US).(72) Inventors: GODAWAT, Rahul; c/o SANOFI U.S., 55  
Corporate Drive, Mail Stop: 55A-505A, Bridgewater, New  
Jersey 08807 (US). CUMMINGS, Daniel; c/o SANOFI  
U.S., 55 Corporate Drive, Mail Stop: 55A-505A, Bridge-  
water, New Jersey 08807 (US). WARIKOO, Veena; c/o  
SANOFI U.S., 55 Corporate Drive, Mail Stop: 55A-505A,  
Bridgewater, New Jersey 08807 (US).(74) Agents: SZCZEPANIK, Ann Marie et al.; Sanofi U.S.,  
55 Corporate Drive, Mail Code 55A-505A, Bridgewater,  
New Jersey 08807 (US).(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,  
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,  
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,  
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,  
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,  
ZW.(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,  
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,  
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: ELUTION OF BIOMOLECULES FROM MULTI-MODAL RESINS USING MES AND MOPS AS MOBILE PHASE  
MODIFIERS(57) Abstract: The present invention relates to methods for  
purifying or enriching a biomolecule using multimodal res-  
ins and an elution buffer containing a Good's buffer.

**Declarations under Rule 4.17:****Published:**

- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
  - *of inventorship (Rule 4.17(iv))*
- *with international search report (Art. 21(3))*

**ELUTION OF BIOMOLECULES FROM MULTI-MODAL RESINS USING MES AND MOPS AS MOBILE  
PHASE MODIFIERS**

**CROSS-REFERENCE TO RELATED APPLICATIONS**

5           This application claims priority to U.S. Application Serial No. 61/717,880, filed  
October 24, 2012, which is incorporated herein by reference in its entirety.

**FIELD OF THE INVENTION**

10           The present invention relates to methods for purifying or enriching a biomolecule  
using multimodal resins and an elution buffer containing a Good's buffer.

**BACKGROUND OF THE INVENTION**

15           Multimodal resins (also termed mixed mode resins) are chromatography media  
that contain at least two different groups that interact with the target molecule by at least  
two different mechanisms. For example, multimodal ion exchangers carry a charged  
group (e.g., a positively charged amino group, a positively charged quarternary  
ammonium group; a negatively charged carboxylic acid group, a negatively charged  
sulfonic acid group) which may interact by charge-charge interactions with the target  
molecule. These multimodal ion exchangers additionally carry another group that may  
20           interact with the target molecule by different modes of action, such as hydrophobic  
interactions, van der Waals interactions, dipole interactions, cation-pi interactions, or  
hydrogen bonding.

25           EP 2 167 526 B1 describes the purification of coagulation Factor VIII using a  
multimodal resin containing ligands which comprise a hydrophobic part and a negatively  
charged part. The elution is carried out with an elution buffer containing a high  
concentration of salt ( $> 1.5$  M), at least 40% (w/v) ethylene glycol, propylene glycol, or a  
mixture thereof, and calcium ions.

30           US 2011/0160435 A1 describes another process for the purification of  
coagulation factor VIII using a multimodal resin. The elution buffer comprises at least one  
amino acid that is positively charged at pH 6 to 8. Preferably, the amino acid is selected  
from lysine, arginine and histidine and is used in concentrations above 0.4 M, especially  
above 0.5 M.

35           US 2011/0166332 A1 describes a method for separating at least one non-  
aggregated protein from a liquid protein preparation by contacting said preparation with a  
multimodal anion exchanger. The experimental section describes the separation of non-

aggregated IgG using an elution buffer containing 1 M glycine in 20 mM Tris, 20 mM Hepes, 20 mM MES, 50 mM NaCl with a linear gradient from pH 9.0 to pH 4.5.

### SUMMARY OF THE INVENTION

- 5 In a first aspect the present invention relates to a method for purifying or enriching a biomolecule, said method comprising the steps:
- (a) providing an aqueous solution containing said biomolecule;
  - (b) contacting said aqueous solution with a multimodal resin;
  - (c) optionally washing the multimodal resin with an aqueous washing buffer;
  - 10 (d) eluting the biomolecule from the multimodal resin with an aqueous buffer solution, wherein said aqueous buffer solution comprises, essentially consists of or consists of: at least 15 mM at least one Good's buffer; between 0 and 400 mM in total concentration of one or more of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, 15 phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 0 and 400 mM in total concentration of one or more of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, 20 potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 0% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol; and
  - (e) collecting biomolecule containing fractions, thereby obtaining the biomolecule in 25 purified or enriched form.

In one embodiment, the aqueous buffer solution comprises 15 mM to 2.5 mM Good's buffer, such as at least 20 mM, 50 mM, 100 mM, 150 mM, 200 mM or more of a Good's buffer. In another embodiment, the aqueous buffer comprises between 10 mM and 400 mM of one or more of alanine, asparagine, aspartic acid, cysteine, glutamine, 30 glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 10 mM and 400 mM of one or more of guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium 35 phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and

between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

In one embodiment, the Good's buffer in the aqueous buffer is MES, MOPS or Tris.

5 In a second aspect, the invention relates to a purified biomolecule, such as a protein, obtainable by the method according to the first aspect.

In a third aspect the present invention relates to a preparation comprising:

- (a) a biomolecule; and
- (b) an aqueous buffer solution comprising, essentially consisting of or consisting of at  
10 least 15 mM of at least one Good's buffer; between 0 and 400 mM in total concentration of one or more of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 0 and 400 mM in total concentration of one or more of urea, guanidinium chloride, guanidinium  
15 thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 0% and  
20 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

In one embodiment, the aqueous buffer solution comprises 15 mM to 2.5 mM of a Good's buffer, such as at least 20 mM, 50 mM, 100 mM, 150 mM, 200 mM or more of a Good's buffer. In another embodiment, the aqueous buffer comprises between 10 mM  
25 and 400 mM of one or more of alanine, asparagine, aspartic acid, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 10 mM and 400 mM of one or more of guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium  
30 phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

35 In one embodiment, the Good's buffer in the aqueous buffer is MES, MOPS or Tris.

In a fourth aspect, the invention relates to a use of an aqueous buffer solution for desorbing a biomolecule from a multimodal resin, wherein said aqueous buffer solution comprises, essentially consists of or consists of at least 15 mM of at least one Good's buffer; between 0 and 400 mM in total concentration of one or more of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 0 and 400 mM in total concentration of one or more of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 0% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

For example, in certain embodiments, the aqueous buffer solution comprises 15 mM to 2.5 mM of a Good's buffer, such as at least 20 mM, 50 mM, 100 mM, 150 mM, 200 mM or more of a Good's buffer. In another embodiment, the aqueous buffer comprises between 10 mM and 400 mM of one or more of alanine, asparagine, aspartic acid, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 10 mM and 400 mM of one or more of guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

In one embodiment, the Good's buffer in the aqueous buffer is MES, MOPS or Tris.

In a fifth aspect, the present invention relates to use of an aqueous buffer solution for increasing the recovery rate of a biomolecule adsorbed on a multimodal resin by contacting (a) the multimodal resin onto which said biomolecule has been adsorbed with (b) the aqueous buffer solution, wherein the aqueous buffer solution comprises, essentially consists of or consists of at least 15 mM of at least one Good's buffer; between 0 and 400 mM in total concentration of one or more of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine,

tryptophan, tyrosine, and valine; between 0 and 400 mM in total concentration of one or more of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 0% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. For example, the aqueous buffer solution for increasing the recovery rate of a biomolecule adsorbed on a multimodal resin can include at least 15 mM, at least 100 mM, or at least 200 mM of a Good's buffer; between 10 mM and 400 mM of one or more of alanine, asparagine, aspartic acid, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 10 mM and 400 mM of one or more of guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

In one embodiment, the solution comprising the biomolecule recovered from the multimodal resin is applied to one or more of ultrafiltration/diafiltration (UF/DF), tangential flow filtration (TFF), microfiltration (MF) and hydrophobic interaction chromatography (HIC). In one embodiment, the solution comprising the biomolecule recovered from the multimodal resin is applied to HIC.

In a sixth aspect, the present invention relates to an article of manufacture comprising:

- (a) a packaging material or container;
- (b) an aqueous buffer solution comprising, essentially consisting of or consisting of:
  - at least 15 mM of at least one Good's buffer;
  - between 0 and 400 mM in total concentration of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine;
  - between 0 and 400 mM in total concentration of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate,

sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and

- 5     – between 0% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol;

and

(c) a data carrier containing instructions for carrying out a method according to the first aspect.

10 In one embodiment, the aqueous buffer solution in the article of manufacture includes at least 15 mM, at least 100 mM, or at least 200 mM of a Good's buffer; between 10 mM and 400 mM of one or more of alanine, asparagine, aspartic acid, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; between 10  
15 mM and 400 mM of one or more of guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium  
20 phosphate, and taurine; and between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

Other features and advantages will be apparent from the accompanying drawings and description, and from the claims.

25 **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 is a contour plot for yield for an experiment in which agalsidase beta is desorbed from the multimodal chromatographic resin Capto™ Adhere (GE Healthcare) by an elution buffer containing MES and arginine. The contour plot is calculated for a pH value of 6.25.

Fig. 2 is a contour plot for yield for an experiment in which agalsidase beta is desorbed from the multimodal chromatographic resin Capto™ Adhere (GE Healthcare) by an elution buffer containing MOPS and arginine. The contour plot is calculated for a pH value of 7.0.

Fig. 3 is a contour plot for yield for an experiment in which agalsidase beta is  
35 desorbed from the multimodal chromatographic resin Capto™ Adhere (GE Healthcare)  
by an elution buffer containing MES and glycol.

Fig. 4 is the multimodal ligand of Capto™ Adhere. The leftmost part of the molecule forms the linkage to the support matrix.

Fig. 5 is the multimodal ligand of Capto™ MMC.

Fig. 6 is the multimodal ligands of Pall PPA HyperCel™, Pall HEA HyperCel™, and Pall MEP HyperCel™. Fig. 6A shows the phenylpropyl substituent of Pall PPA HyperCel™, which is bound to an amine group (not shown). Fig. 6B shows the n-hexyl substituent of Pall HEA HyperCel™, which is bound to an amine group (not shown). Fig. 6C shows the 4-mercapto-ethyl-pyridine ligand of Pall MEP HyperCel™.

Fig. 7 is the multimodal ligand of Eshmuno® HCX resin obtainable from Merck Millipore, Germany.

#### DETAILED DESCRIPTION OF THE INVENTION

The invention is based on the discovery that both MES and MOPS, which are typically used as buffering agents, are surprisingly effective in disrupting multimodal interactions. Thus, by using MES, MOPS or other Good's buffers, it is possible to lower the concentration of arginine or related mobile phase modifiers with high conductivity or to use elution buffers free, or substantially free, from arginine and related mobile phase modifiers when desorbing biomolecules from multimodal chromatographic resins. Furthermore, by using an increased concentration of MES, MOPS or other Good's buffers in accordance with the present invention, it is possible to significantly reduce or even completely avoid the presence of chaotropic agents in the elution buffer.

Desorption of biomolecules bound to a multimodal chromatographic resin can be challenging due to the complex nature of interactions that combine ionic, hydrogen bonding and hydrophobic interactions. Traditional mobile phase additives, such as simple salts (e.g., NaCl) and organic solvents, may not work well for desorption of biomolecules due to opposing effects on individual modes of interaction. For example, addition of NaCl weakens ionic interactions but strengthens hydrophobic interactions and, vice versa for organic solvents. Therefore, special mobile phase modifiers are needed to disrupt multimodal interactions collectively. Use of arginine, guanidine, urea and sodium caprylate have been reported in the literature (Arakawa *et al*, Protein Expression and Purification 63 2009 pp. 158-163, Kaleas *et al*, J. Chromatogr. A, 1217, 2010 pp. 235-242 and Holstein *et al*, Biotechnol. Bioeng., 109 Jan 2012).

Solutions of the above-mentioned mobile phase modifiers (e.g., arginine or other amino acids) typically exhibit a high conductivity, which is disadvantageous for subsequent purification steps (e.g., ion exchange steps) that are often carried out after the multimodal resin chromatography. More specifically, a high conductivity may require

large dilutions or extended times of dialysis. Furthermore, several of the commonly used mobile phase modifiers can denature the quaternary structure or even the tertiary structure of biomolecules (e.g., proteins), if such mobile phase modifiers are present in high concentration. The risk of denaturation is particularly high if the mobile phase  
5 modifier is a chaotropic agent, such as urea or a guanidinium salt.

Therefore, it would be desirable to reduce the concentration of arginine or other additives with high conductivity and to identify mobile phase modifiers that exhibit low conductivity. It is further desirable to reduce the concentration of chaotropic agents (such as urea, guanidinium chloride, guanidinium thiocyanate) in the elution buffer and to  
10 identify mobile phase modifiers that exert a weak effect, or have no denaturing effect, on biomolecules.

Definitions. The present invention is not limited to the particular methodology, protocols and reagents described herein as these may vary. Further, the terminology used herein is for the purpose of describing particular embodiments only, and is not  
15 intended to limit the scope of the present invention which will be limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs.

The terms used herein are defined as described in "A multilingual glossary of  
20 biotechnological terms: (IUPAC Recommendations)", Leuenberger, H.G.W, Nagel, B. and Kölbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland).

Throughout this specification and the claims that follow, unless the context requires otherwise, the word "comprise," and variations such as "comprises" and "comprising," will be understood to imply the inclusion of a stated integer or step or group  
25 of integers or steps but not the exclusion of any other integer or step or group of integer or step.

Several documents (for example: patents, patent applications, scientific publications, manufacturer's specifications, instructions, GenBank Accession Number sequence submissions etc.) are cited throughout the text of this specification. Nothing  
30 herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention. Some of the documents cited herein are characterized as being "*incorporated by reference*". In the event of a conflict between the definitions or teachings of such incorporated references and definitions or teachings recited in the present specification, the text of the present specification takes  
35 precedence.

The term "Good's buffer" originally referred to one of twelve buffering agents that were selected and described by Norman Good and colleagues in 1966 as particularly suitable for biological research. In the context of the present specification, the "Good's buffer" does not only encompass the originally described twelve buffering agents but further encompasses structurally related buffering agents. Accordingly, as used herein, the term "Good's buffer" comprises the following chemical compounds: ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, MES, MOPS, PIPES, TAPS, TES, and Tris.

ADA is an abbreviation for N-(2-Acetamido)iminodiacetic acid.

ACES is an abbreviation for N-(2-Acetamido)-2-aminoethanesulfonic acid.

BES is an abbreviation for N,N-Bis(2-hydroxyethyl)-2-aminoethanesulfonic.

Bicine is an abbreviation for N,N-Bis(2-hydroxyethyl)glycine

CAPS is an abbreviation for 3-(Cyclohexylamino)-1-propanesulfonic acid.

CAPSO is an abbreviation for 3-(Cyclohexylamino)-2-hydroxy-1-propanesulfonic acid.

CHES is an abbreviation for 2-(Cyclohexylamino)ethanesulfonic acid.

HEPES is an abbreviation for 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid.

MES is an abbreviation for 2-(N-Morpholino)ethanesulfonic acid.

MOPS is an abbreviation for 3-(N-Morpholino)propanesulfonic acid.

PIPES is an abbreviation for 1,4-Piperazinediethanesulfonic acid.

TAPS is an abbreviation for N-[Tris(hydroxymethyl)methyl]-3-aminopropanesulfonic acid.

TES is an abbreviation for 2-[(2-Hydroxy-1,1-bis(hydroxymethyl)ethyl)amino]ethane-sulfonic acid.

Tris is an abbreviation for Tris(hydroxymethyl)aminomethane.

Pall MEP HyperCel™ is a chromatography sorbent composed of a rigid cellulose matrix to which 4-Mercapto-Ethyl-Pyridine (4-MEP) is linked (see Fig. 6C). The cellulose bead confers high porosity, chemical stability and low non-specific interaction. The average bead diameter is 80 µm to 100 µm. Pall MEP HyperCel™ operates by a mixed-mode or multi-mode mechanism also described as Hydrophobic Charge Induction Chromatography (HCIC). HCIC is based on the pH-dependent behavior of ionizable, dual-mode ligands.

Pall PPA HyperCel™ is a chromatography mixed-mode sorbent. The sorbent material is based on high porosity cross-linked cellulose and contains phenylpropylamine groups as ligands (see Fig. 6A).

Pall HEA HyperCel™ is a chromatography mixed-mode sorbent. The sorbent material is based on high porosity crossed-linked cellulose and contains hexylamine groups as ligands (see Fig. 6B).

Capto™ Adhere is a chromatography sorbent employing the ligand  
5 N-benzyl-N-methyl-ethanolamine immobilized on porous agarose particles. It is obtainable from GE Healthcare Life Sciences. The detailed chemical structure of this ligand is shown in Fig. 4.

Capto™ MMC is a chromatography sorbent employing a ligand derived from N-benzoyl-methionine immobilized on porous agarose particles. It is obtainable from GE  
10 Healthcare Life Sciences. The detailed chemical structure of this ligand is shown in Fig. 5.

Eshmuno® HCX media is a mixed-mode resin that couples a tentacle structure (of the ligands) with a hydrophilic polyvinyl ether base matrix. More specifically, the base matrix is a surface grafted rigid polyvinyl ether hydrophilic polymer. The mean particle  
15 size ( $d_{50}$ ) is 75  $\mu\text{m}$  to 95  $\mu\text{m}$ . The detailed chemical structure of the ligand employed in Eshmuno® HCX media is shown in Fig. 7. The Eshmuno® HCX media is obtainable from Merck Millipore, Germany, or EMD Millipore, USA.

#### Alpha-galactosidase

20 Alpha-galactosidase is a glycoside hydrolase enzyme that hydrolyses the terminal alpha-galactosyl moieties from glycolipids and glycoproteins. It is encoded by the GLA gene. This enzyme is a homodimeric glycoprotein that hydrolyses the terminal alpha-galactosyl moieties from glycolipids and glycoproteins. It predominantly hydrolyzes ceramide trihexoside, and it can catalyze the hydrolysis of melibiose into galactose and  
25 glucose.

A variety of mutations in this gene affect the synthesis, processing, and stability of this enzyme, which causes Fabry's disease, a rare lysosomal storage disorder and sphingolipidosis that results from a failure to catabolize alpha-D-galactosyl glycolipid moieties. Two enzyme replacement therapies are available to functionally compensate  
30 for alpha-galactosidase deficiency. Agalsidase alpha and beta are both recombinant forms of the human  $\alpha$ -galactosidase A enzyme and both have the same amino acid sequence as the native enzyme. Agalsidase alpha and beta differ in the structures of their oligosaccharide side chains. The pharmaceutical company Shire manufactures agalsidase alpha under the brand name Replagal® as a treatment for Fabry's disease.  
35 The pharmaceutical company Genzyme produces synthetic agalsidase beta under the brand name Fabrazyme® for treatment of Fabry's disease.

Each of the below aspects may be combined with any other aspect or aspects unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous, unless clearly indicated to the contrary. The present invention is especially directed to those embodiments, in which particularly preferred features are combined with one or more other particularly preferred features.

In a first aspect the present invention is directed to a method for purifying or enriching a biomolecule, said method comprising the steps:

- (a) providing an aqueous solution containing the biomolecule;
- 10 (b) contacting the aqueous solution with a multimodal resin;
- (c) optionally washing the multimodal resin with an aqueous washing buffer;
- (d) eluting the biomolecule from the multimodal resin by an aqueous buffer solution, wherein the aqueous buffer solution comprises, essentially consists of or consists of:
  - at least 15 mM of at least one Good's buffer, such as between 15 mM and 2.5 M of at least one Good's buffer, between 50 mM and 2.0 M, between 75 mM and 1.5 M, between 100 mM and 1.25 M, between 200 mM and 1.0 M, between 300 mM and 900 mM, or between 400 mM and 750 mM of at least one Good's buffer;
  - between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM or between 125 mM and 150 mM), in total concentration of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine;
  - between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM or between 125 mM and 150 mM ) in total concentration of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and
  - between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol;

and

(e) collecting biomolecule containing fractions, thereby obtaining the biomolecule in purified or enriched form.

The method according to the first aspect provides several advantages as compared to previous methods for purifying or enriching biomolecules. Previous methods of purifying or enriching biomolecules using multimodal chromatography typically employ elution buffers containing either high concentrations of arginine, glycine, or other amino acids (e.g., 1 M glycine, cf. example 1 of US 2011/0166332 A1; or 0.8 mol/kg L-arginine, cf. example 10 of US 2011/0160435 A1) or high concentrations of salt (e.g., up to 2.5 M of  $\text{NH}_4\text{Cl}$  or  $\text{NaCl}$ , cf. example 6 of EP 2 167 526 B1). However, arginine and other additives exhibit a high conductivity which is disadvantageous in subsequent purification steps (e.g., ion exchange steps) that are often carried out after the multimodal resin chromatography. In particular, a high conductivity may require large dilutions or extended times of dialysis. In addition, high concentrations of salts can denature the quaternary structure or even the tertiary structure of biomolecules (e.g., proteins). The risk of denaturation is particularly high if the salt is a chaotropic agent (e.g., guanidinium salts or lithium perchlorate).

The method according to the first aspect provides the advantage that the desorption of the biomolecule is effected by an elution buffer containing only low concentrations of amino acids (such as between 0 and 400 mM, such as between 10 mM and 300 mM) and only low concentrations of salts or chaotropic agents (such as between 0 and 400 mM, such as between 10 mM and 300 mM), thereby reducing the risk of denaturing the biomolecule and facilitating subsequent purification steps. This advantage is achieved by increasing the concentration of the Good's buffer in the elution buffer to values of at least 15 mM, at least 50 mM, at least 100 mM or at least 200 mM. Good's buffers are well known for their use in biological research and typically have mild to no effects on biomolecules. Accordingly, the purification or enrichment method of the present invention is gentler than methods traditionally employed.

In summary, desorption of a biomolecule, such as a protein, such as an enzyme, is effected in accordance with the invention either by a Good's buffer in a concentration of 15 mM or more (such as 50 mM, 100 mM or 200 mM or more) in combination with an amino acid in a concentration below 400 mM or by a Good's buffer in a concentration of 15 mM or more, such as 50 mM, 100 mM or 200 mM, in combination with ethylene glycol and/or propylene glycol in a total concentration of not more than 60% (v/v). Of course, the amino acid and the glycol may also be present at the same time.

In some embodiments, the aqueous buffer solution contains at least one amino acid (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid,

glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine), or ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. In other words, in these embodiments the concentrations of ethylene glycol and propylene glycol are each 0%, if the aqueous buffer solution contains one or more amino acids. Conversely, in these embodiments the total concentration of said one or more amino acids is 0 mM, if the aqueous buffer solution contains ethylene glycol or propylene glycol or both.

In some embodiments, the aqueous buffer solution comprises between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol.

In those embodiments in which the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, asparagine, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, and valine. In other words, in these embodiments aspartic acid, glutamic acid, tryptophan and tyrosine are absent. In those embodiments of the first aspect in which the aqueous buffer solution contains at least one amino acid, the at least one amino acid is selected from the group consisting of alanine, cysteine, glycine, histidine, lysine, proline, serine, threonine, and valine. In other words, in these embodiments, asparagine, aspartic acid, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, tryptophan and tyrosine are typically absent. In those embodiments of the first aspect in which the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, glycine, histidine, and lysine. In other words, in these embodiments, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine are typically absent.

In one embodiment, the at least one Good's buffer is selected from the group consisting of MES, MOPS, Tris, ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, PIPES, TAPS, and TES. In one embodiment, the at least one Good's buffer is MES, and in another embodiment, the at least one Good's buffer is MOPS.

In one embodiment, the biomolecule is a protein (such as an enzyme, a structural protein, a transport protein, a signaling protein, a transmembrane protein, a peptide hormone or an antibody) or a fragment thereof, or a nucleic acid (e.g., a single-stranded oligonucleotide, a double-stranded oligonucleotide, a single-stranded polynucleotide, or a double-stranded polynucleotide). The nucleic acid can be, for example, DNA, RNA or a hybrid double-strand formed from DNA and RNA. In one embodiment, the biomolecule is

an enzyme (e.g., an oxidoreductase, a transferase, a hydrolase, a lyase, an isomerase, or a ligase). For example, in one embodiment, the enzyme is a hydrolase, such as a glycosidase (often also termed glycoside hydrolase). In one embodiment, the glycosidase is an alpha-galactosidase, such as an alpha-galactosidase selected from the group consisting of agalsidase alpha (e.g., Replagal® manufactured by Shire, United Kingdom) and agalsidase beta (e.g., Fabrazyme® manufactured by Genzyme, Cambridge, MA). In one embodiment, the alpha-galactosidase is agalsidase beta. In another embodiment, the biomolecule is a peptide hormone, such as thyroid-stimulating hormone (also known as TSH or thyrotropin). TSH is a glycoprotein consisting of two subunits: the alpha subunit and the beta subunit. A recombinant form of human TSH-alpha is obtainable from Genzyme Corp. under the tradename "Thyrogen". In yet another embodiment, the biomolecule is an antibody, such as a monoclonal antibody. In certain embodiments, the biomolecule is a glycosylated protein.

In one embodiment of the first aspect, the multimodal resin is selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP HyperCel™, and Eshmuno® HCX media. The ligands used in these multimodal resins on the filing day of the present application are shown in Figures 4, 5, 6A, 6B, 6C, and 7. In one embodiment, the multimodal resin is Capto™ Adhere.

In a second aspect, the present invention is directed to a purified biomolecule obtainable or obtained by the method according to the first aspect. A biomolecule can be purified or enriched by a method that is gentler than the purification or enrichment methods typically known in the art. Accordingly, it is expected that the purified biomolecule is in a state that is more native than that of a purified biomolecule obtainable by methods of the prior art.

In some embodiments, the aqueous buffer solution contains either at least one amino acid (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine), or ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. In other words, in these embodiments the concentrations of ethylene glycol and propylene glycol are each 0%, if the aqueous buffer solution contains one or more amino acids. Conversely, in these embodiments the total concentration of said one or more amino acids is 0 mM, if the aqueous buffer solution contains ethylene glycol or propylene glycol or both.

In some embodiments, the aqueous buffer solution comprises between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol.

In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, asparagine, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, and valine. In other words, in these  
5       embodiments aspartic acid, glutamic acid, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, cysteine, glycine, histidine, lysine, proline, serine, threonine, and valine. In other words, in these embodiments, asparagine, aspartic acid, glutamic acid, glutamine,  
10       isoleucine, leucine, methionine, phenylalanine, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, glycine, histidine, and lysine. In other words, in these embodiments, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, isoleucine, leucine, methionine,  
15       phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine are typically absent.

In one embodiment, the at least one Good's buffer is selected from the group consisting of MES, MOPS, Tris, ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, PIPES, TAPS, and TES. In one embodiment, the at least one Good's buffer is  
20       MES, and in another embodiment, the at least one Good's buffer is MOPS.

In one embodiment, the biomolecule is a protein (e.g., an enzyme, a structural protein, a transport protein, a signaling protein, a transmembrane protein, a peptide hormone or an antibody) or a fragment thereof, or a nucleic acid (e.g., a single-stranded oligonucleotide, a double-stranded oligonucleotide, a single-stranded polynucleotide, or  
25       a double-stranded polynucleotide). The nucleic acid can be, for example, DNA, RNA or a hybrid double-strand formed from DNA and RNA. In one embodiment, the biomolecule is an enzyme (e.g., an oxidoreductase, a transferase, a hydrolase, a lyase, an isomerase, or a ligase). For example, the enzyme can be a hydrolase, such as a glycosidase (often also termed glycoside hydrolase). In one embodiment, the glycosidase is an alpha-  
30       galactosidase, such as an alpha-galactosidase selected from the group consisting of agalsidase alpha (e.g., Replagal® manufactured by Shire, United Kingdom) and agalsidase beta (e.g., Fabrazyme® manufactured by Genzyme, Cambridge, MA). In one embodiment, the alpha-galactosidase is agalsidase beta. In another embodiment, the biomolecule is a peptide hormone, such as thyroid-stimulating hormone (also known as  
35       TSH or thyrotropin). TSH is a glycoprotein consisting of two subunits: the alpha subunit and the beta subunit. A recombinant form of human TSH-alpha is obtainable from Genzyme Corp. under the tradename "Thyrogen." In yet another embodiment, the

biomolecule is an antibody, such as a monoclonal antibody. In certain embodiments, the biomolecule is a glycosylated protein.

In one embodiment, the multimodal resin is selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP  
 5 HyperCel™, and Eshmuno® HCX media. The ligands used in these multimodal resins on the filing day of the present application are shown in Figures 4, 5, 6A, 6B, 6C, and 7. In one embodiment, the multimodal resin is Capto™ Adhere.

In a third aspect, the present invention is directed to a preparation comprising:

- (a) a biomolecule; and
- 10 (b) an aqueous buffer solution comprising, essentially consisting of or consisting of
  - at least 15 mM of at least one Good's buffer, such as between 15 mM and 2.5 M of at least one Good's buffer, such as between 50 mM and 2.0 M, between 75 mM and 1.5 M, between 100 mM and 1.25 M, between 200 mM and 1.0 M, between 300 mM and 900 mM, or between 400 mM and 750 mM of at  
 15 least one Good's buffer;
  - between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 mM and 150 mM), in total concentration of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid,  
 20 glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine;
  - between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 mM and 150 mM) in total concentration of  
 25 urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate,  
 30 magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and
  - between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol or propylene glycol or a  
 35 mixture of ethylene glycol and propylene glycol.

As explained above in the section discussing the first aspect, the aqueous buffer solution described herein involves a reduced risk of denaturing a biomolecule as compared to elution buffer solutions described in the art. Therefore, it is expected that the preparation according to the third aspect contains the biomolecule in a state that is more native than that of biomolecules in preparations described in the art.

In some embodiments, the aqueous buffer solution contains either at least one amino acid (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine), or ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. In other words, in these embodiments the concentrations of ethylene glycol and propylene glycol are each 0%, if the aqueous buffer solution contains one or more amino acids. Conversely, in these embodiments the total concentration of said one or more amino acids is 0 mM, if the aqueous buffer solution contains ethylene glycol or propylene glycol or both.

In some embodiments, the aqueous buffer solution comprises between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol.

In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, asparagine, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, and valine. In other words, in these embodiments aspartic acid, glutamic acid, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, cysteine, glycine, histidine, lysine, proline, serine, threonine, and valine. In other words, in these embodiments, asparagine, aspartic acid, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is selected from the group consisting of alanine, glycine, histidine, and lysine. In other words, in these embodiments, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine are typically absent.

In one embodiment, the at least one Good's buffer is selected from the group consisting of MES, MOPS, Tris, ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, PIPES, TAPS, and TES. In one embodiment, the at least one Good's buffer is MES, and in another embodiment, the at least one Good's buffer is MOPS.

In one embodiment, the biomolecule is a protein (e.g., an enzyme, a structural protein, a transport protein, a signaling protein, a transmembrane protein, a peptide hormone or an antibody) or a fragment thereof, or a nucleic acid (e.g., a single-stranded oligonucleotide, a double-stranded oligonucleotide, a single-stranded polynucleotide, or a double-stranded polynucleotide). The nucleic acid can be, for example, DNA, RNA or a hybrid double-strand formed from DNA and RNA. In one embodiment, the biomolecule is an enzyme (e.g., an oxidoreductase, a transferase, a hydrolase, a lyase, an isomerase, or a ligase). In one embodiment, the enzyme is a hydrolase, such as a glycosidase (often also termed glycoside hydrolase). In one embodiment, the glycosidase is an alpha-galactosidase, such as an alpha-galactosidase selected from the group consisting of agalsidase alpha (e.g., Replagal® manufactured by Shire, United Kingdom) and agalsidase beta (e.g., Fabrazyme® manufactured by Genzyme, Cambridge, MA). In one embodiment, the alpha-galactosidase is agalsidase beta, and in another embodiment, the biomolecule is a peptide hormone, such as thyroid-stimulating hormone (also known as TSH or thyrotropin). TSH is a glycoprotein consisting of two subunits: the alpha subunit and the beta subunit. A recombinant form of human TSH-alpha is obtainable from Genzyme Corp. under the tradename "Thyrogen". In yet another embodiment, the biomolecule is an antibody, such as a monoclonal antibody. In certain embodiments, the biomolecule is a glycosylated protein.

In a fourth aspect, the present invention is directed to a use of an aqueous buffer solution for desorbing a biomolecule from a multimodal resin, wherein said aqueous buffer solution comprises, essentially consists of or consists of:

- at least 15 mM of at least one Good's buffer, such as between 15 mM and 2.5 M of at least one Good's buffer, between 50 mM and 2.0 M, between 75 mM and 1.5 M, between 100 mM and 1.25 M, between 200 mM and 1.0 M, between 300 mM and 900 mM, or between 400 mM and 750 mM of at least one Good's buffer;
- between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 mM and 150 mM), in total concentration of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine;
- between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 mM and 150 mM) in total concentration of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate,

sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and

- 5                   – between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

As will be illustrated in the experimental section below, the aqueous buffer  
10 solution described herein is particularly well suited for desorbing biomolecules from a multimodal resin. It was especially unexpected that Good's buffers could be effective in desorbing biomolecules from a multimodal resin to an extent that is comparable to that of arginine.

In some embodiments, the aqueous buffer solution contains either at least one  
15 amino acid (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine), or ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. In other words, in these embodiments the concentrations of ethylene glycol and propylene glycol are each  
20 0%, if the aqueous buffer solution contains one or more amino acids. Conversely, in these embodiments the total concentration of said one or more amino acids is 0 mM, if the aqueous buffer solution contains ethylene glycol or propylene glycol or both.

In some embodiments of the fourth aspect, the aqueous buffer solution comprises  
between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40%  
25 (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol.

In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, asparagine, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine,  
30 methionine, phenylalanine, proline, serine, threonine, and valine. In other words, in these embodiments aspartic acid, glutamic acid, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, cysteine, glycine, histidine, lysine, proline, serine, threonine, and valine. In other  
35 words, in these embodiments, asparagine, aspartic acid, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one

amino acid, the at least one amino acid is typically selected from the group consisting of alanine, glycine, histidine, and lysine. In other words, in these embodiments, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine are typically  
5 absent.

In one embodiment, the at least one Good's buffer is selected from the group consisting of MES, MOPS, Tris, ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, PIPES, TAPS, and TES. In one embodiment, the at least one Good's buffer is MES, and in another embodiment, the at least one Good's buffer is MOPS.

10 In one embodiment, the biomolecule is a protein (e.g., an enzyme, a structural protein, a transport protein, a signaling protein, a transmembrane protein, a peptide hormone or an antibody) or a fragment thereof, or a nucleic acid (e.g., a single-stranded oligonucleotide, a double-stranded oligonucleotide, a single-stranded polynucleotide, or a double-stranded polynucleotide). The nucleic acid can be, for example, DNA, RNA or a  
15 hybrid double-strand formed from DNA and RNA. In one embodiment, the biomolecule is an enzyme (e.g., an oxidoreductase, a transferase, a hydrolase, a lyase, an isomerase, or a ligase). In one embodiment, the enzyme is a hydrolase, such as a glycosidase (often also termed glycoside hydrolase). In another embodiment, the glycosidase is an alpha-galactosidase, such as agalsidase alpha (e.g., Replagal® manufactured by Shire,  
20 United Kingdom) or agalsidase beta (e.g., Fabrazyme® manufactured by Genzyme, Cambridge, MA). In one, the alpha-galactosidase is agalsidase beta. In another embodiment, the biomolecule is a peptide hormone, such as thyroid-stimulating hormone (also known as TSH or thyrotropin). TSH is a glycoprotein consisting of two subunits: the alpha subunit and the beta subunit. A recombinant form of human TSH-alpha is  
25 obtainable from Genzyme Corp. under the tradename "Thyrogen". In yet another embodiment, the biomolecule is an antibody, such as a monoclonal antibody. In certain embodiments, the biomolecule is a glycosylated protein.

In one embodiment, the multimodal resin is selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP  
30 HyperCel™, and Eshmuno® HCX media. The ligands used in these multimodal resins on the filing day of the present application are shown in Figs. 4, 5, 6A, 6B, 6C, and 7. In one embodiment, the multimodal resin is Capto™ Adhere.

In a fifth aspect, the present invention is directed to a use of an aqueous buffer solution for increasing the recovery rate of a biomolecule adsorbed on a multimodal resin  
35 by contacting (a) said multimodal resin onto which said biomolecule has been adsorbed with (b) said aqueous buffer solution, wherein said aqueous buffer solution comprises, essentially consists of or consists of:

- at least 15 mM of at least one Good's buffer, such as between 15 mM and 2.5 M of at least one Good's buffer, between 50 mM and 2.0 M, between 75 mM and 1.5 M, between 100 mM and 1.25 M, between 200 mM and 1.0 M, between 300 mM and 900 mM, or between 400 mM and 750 mM of at least one Good's buffer;
- 5 – between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 mM and 150 mM), in total concentration of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, 10 tryptophan, tyrosine, and valine;
- between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 mM and 150 mM) in total concentration of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium 15 phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and
- 20 – between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

As will be illustrated in the experimental section below, the aqueous buffer 25 solution described herein is particularly well-suited for desorbing biomolecules from a multimodal chromatography resin at a high recovery rate. The inventors achieved recovery rates of up to 99% using aqueous buffer solutions featured in the invention (see Table 1).

In some embodiments, the aqueous buffer solution contains either at least one 30 amino acid (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine), or ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. In other words, in these embodiments the concentrations of ethylene glycol and propylene glycol are each 35 0%, if the aqueous buffer solution contains one or more amino acids. Conversely, in

these embodiments the total concentration of said one or more amino acids is 0 mM, if the aqueous buffer solution contains ethylene glycol or propylene glycol or both.

In some embodiments of the fifth aspect, the aqueous buffer solution comprises between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), or between 15% and 20% (v/v)) of ethylene glycol.

In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, asparagine, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, and valine. In other words, in these 10 embodiments aspartic acid, glutamic acid, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, cysteine, glycine, histidine, lysine, proline, serine, threonine, and valine. In other 15 words, in these embodiments, asparagine, aspartic acid, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, tryptophan and tyrosine are typically absent. In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, glycine, histidine, and lysine. In other words, in these most embodiments, 20 asparagine, aspartic acid, cysteine, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine are typically absent.

In one embodiment, the at least one Good's buffer is selected from the group consisting of MES, MOPS, Tris, ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, 25 HEPES, PIPES, TAPS, and TES. In one embodiment, the at least one Good's buffer is MES, and in another embodiment, the at least one Good's buffer is MOPS.

In one embodiment, the biomolecule is a protein (e.g., an enzyme, a structural protein, a transport protein, a signaling protein, a transmembrane protein, a peptide hormone or an antibody) or a fragment thereof, or a nucleic acid (e.g., a single-stranded 30 oligonucleotide, a double-stranded oligonucleotide, a single-stranded polynucleotide, or a double-stranded polynucleotide). The nucleic acid can be, for example, DNA, RNA or a hybrid double-strand formed from DNA and RNA. In one embodiment, the biomolecule is an enzyme (e.g., an oxidoreductase, a transferase, a hydrolase, a lyase, an isomerase, or a ligase). In another embodiment, the enzyme is a hydrolase, such as a glycosidase 35 (often also termed glycoside hydrolase). In one embodiment, the glycosidase is an alpha-galactosidase, such as an alpha-galactosidase selected from the group consisting of agalsidase alpha (e.g., Replagal® manufactured by Shire, United Kingdom) and

agalsidase beta (e.g., Fabrazyme® manufactured by Genzyme, Cambridge, MA). In one embodiment, the alpha-galactosidase is agalsidase beta. In another embodiment, the biomolecule is a peptide hormone, such as thyroid-stimulating hormone (also known as TSH or thyrotropin). TSH is a glycoprotein consisting of two subunits: the alpha subunit and the beta subunit. A recombinant form of human TSH-alpha is obtainable from Genzyme Corp. under the tradename "Thyrogen". In yet another embodiment, the biomolecule is an antibody, such as a monoclonal antibody. In other embodiments, the biomolecule is a glycosylated protein.

In one embodiment, the multimodal resin is selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP HyperCel™, and Eshmuno® HCX media. The ligands used in these multimodal resins on the filing day of the present application are shown in Figures 4, 5, 6A, 6B, 6C, and 7. In one embodiment, the multimodal resin is Capto™ Adhere.

In a sixth aspect, the present invention is directed to an article of manufacture comprising:

- (a) a packaging material or container;
- (b) an aqueous buffer solution comprising, essentially consisting of or consisting of:
  - at least 15 mM of at least one Good's buffer, such as between 15 mM and 2.5 M of at least one Good's buffer, between 50 mM and 2.0 M, between 75 mM and 1.5 M, between 100 mM and 1.25 M, between 200 mM and 1.0 M, between 300 mM and 900 mM, and between 400 mM and 750 mM of at least one Good's buffer;
  - between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 and 150 mM), in total concentration of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine;
  - between 0 and 400 mM (such as 0 mM, between 10 mM and 350 mM, between 25 mM and 300 mM, between 50 mM and 250 mM, between 100 mM and 200 mM, or between 125 and 150 mM) in total concentration of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, and taurine; and

- between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), between 15% and 20% (v/v)) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol;

5 and

- (c) a data carrier containing instructions for carrying out a method described herein.

The aqueous buffer solution featured in the invention involves a reduced risk of denaturing a biomolecule as compared to elution buffer solutions described in the prior art. The article of manufacture according to the sixth aspect provides this advantageous aqueous buffer solution in combination with a container or packaging material and with a data carrier containing instructions for carrying out a method that makes use of this advantageous aqueous buffer solution.

Therefore, it is expected that the preparation according to the present aspect contains the biomolecule in a state that is more native than that of biomolecules in preparations described in the prior art.

In one embodiment, the data carrier is selected from the group consisting of a label present on the packaging material or container (e.g., in the form of an adhesive label, a tag, a chip, or an RFID tag) and a packaging insert (e.g., a leaflet; a booklet; a chip; or a computer-readable storage medium, such as a floppy disk, a CD-ROM, a CD-R, a DVD, or a blue-ray disk). In certain embodiments, the data carrier comprises written information, visual information (e.g., pictures), or computer-readable information (e.g., a program or a bar code).

In one embodiment, the data carrier further comprises one or more of the following information:

- (i) instructions for use of aqueous buffer solution;
- (ii) quality information such as information about the lot/batch number, the manufacturing or assembly site or the expiry or sell-by date of the aqueous buffer solution; or information concerning the correct storage or handling of the article;
- (iii) information concerning the composition of the aqueous buffer solution;
- (iv) information concerning the interpretation of information obtained when performing methods according to the first aspect;
- (v) a warning concerning possible misinterpretations or wrong results when applying unsuitable methods; and/or
- (vi) a warning concerning possible misinterpretations or wrong results when using unsuitable reagent(s) and/or buffer(s).

In one embodiment of the sixth aspect, the article of manufacture further comprises one or more of the following components:

- (d) a chromatography medium;
- (e) a chromatography column;
- (f) a reference biomolecule usable for calibrating a method for purifying or enriching a biomolecule, such as for calibrating a method according to the first aspect.

5 In some embodiments, the chromatography column is prepacked with the chromatography medium.

In one embodiment, the chromatography medium is a multimodal chromatography resin. The multimodal chromatography resin is typically selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA  
 10 HyperCel™, Pall MEP HyperCel™, and Eshmuno® HCX media. The ligands used in these multimodal resins are shown in Figures 4, 5, 6A, 6B, 6C, and 7. In one embodiment, the multimodal resin is Capto™ Adhere.

In one embodiment, the reference biomolecule is selected from the group consisting of a protein (e.g., an enzyme, a structural protein, a transport protein, a  
 15 signaling protein, a transmembrane protein, a peptide hormone or an antibody) and a nucleic acid (e.g., a single-stranded oligonucleotide, a double-stranded oligonucleotide, a single-stranded polynucleotide, or a double-stranded polynucleotide). The nucleic acid can be DNA, RNA or a hybrid double-strand formed from DNA and RNA. In one embodiment, the reference biomolecule is an enzyme (e.g., an oxidoreductase, a  
 20 transferase, a hydrolase, a lyase, an isomerase, or a ligase). The reference enzyme can be, for example, a hydrolase, such as a glycosidase (often also termed glycoside hydrolase). In one embodiment, the reference glycosidase is an alpha-galactosidase, such as an agalsidase alpha (e.g., Replagal® manufactured by Shire, United Kingdom) or an agalsidase beta (e.g., Fabrazyme® manufactured by Genzyme, Cambridge, MA).  
 25 In one embodiment, the reference alpha-galactosidase is agalsidase beta. In another embodiment, the reference biomolecule is a peptide hormone, such as thyroid-stimulating hormone (also known as TSH or thyrotropin). TSH is a glycoprotein consisting of two subunits: the alpha subunit and the beta subunit. A recombinant form of human TSH-alpha is obtainable from Genzyme Corp. under the tradename "Thyrogen".  
 30 In yet another embodiment, the reference biomolecule is an antibody, such as a monoclonal antibody. In certain embodiments, the reference biomolecule is a glycosylated protein.

In some embodiments, the aqueous buffer solution contains either at least one amino acid (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid,  
 35 glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine), or ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol. In other words, in

these embodiments the concentrations of ethylene glycol and propylene glycol are each 0%, if the aqueous buffer solution contains one or more amino acids. Conversely, in these embodiments the total concentration of said one or more amino acids is 0 mM, if the aqueous buffer solution contains ethylene glycol or propylene glycol or both.

5 In some embodiments of the sixth aspect, the aqueous buffer solution comprises between 0% and 60% (v/v) (such as between 1% and 50% (v/v), between 5% and 40% (v/v), between 8% and 30% (v/v), between 10% and 25% (v/v), between 15% and 20% (v/v)) of ethylene glycol.

10 In those embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, asparagine, cysteine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, and valine. In other words, in these embodiments aspartic acid, glutamic acid, tryptophan and tyrosine are absent. In those  
15 embodiments where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is selected from the group consisting of alanine, cysteine, glycine, histidine, lysine, proline, serine, threonine, and valine. In other words, in these embodiments, asparagine, aspartic acid, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, tryptophan and tyrosine are absent. In those embodiments  
20 where the aqueous buffer solution contains at least one amino acid, the at least one amino acid is typically selected from the group consisting of alanine, glycine, histidine, and lysine. In other words, in these embodiments, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine are typically absent.

In one embodiment of the sixth aspect, the at least one Good's buffer is selected  
25 from the group consisting of MES, MOPS, Tris, ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, PIPES, TAPS, and TES. In one embodiment, the at least one Good's buffer is MES. In another embodiment, the at least one Good's buffer is MOPS.

### EXAMPLES

30 The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions featured in the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with  
35 respect to numbers used but some experimental errors and deviations should be accounted for.

**Example 1:** Desorption of a biomolecule using a combination of a Good's buffer and an amino acid

In this experiment, a biomolecule, agalsidase beta (Fabrazyme® manufactured by Genzyme, Cambridge, MA), was bound on multimodal chromatographic resin Capto™ Adhere (GE Healthcare) and desorbed using MES and MOPS (see **Table 1** and **Figures 1** and **2**). The table shows the detailed experimental design, where a range of MES concentrations from 50 mM to 200 mM and MOPS concentrations from 20 mM to 200 mM was tested in the presence of 150 mM to 400 mM arginine within a pH range of 5.5 to 7.5. Yield was monitored in these experiments as an indicator of desorption from the chromatographic resin. Under a given buffer condition, 0% yield means no desorption and 100% means complete desorption of the biomolecule. The experimental design and analysis was done using Design-Expert®.

**Table 1:** Experimental design and results for elution with MES and MOPS

| MES (mM) | Arginine (mM) | pH  | Yield (%) | MOPS (mM) | Arginine (mM) | pH  | Yield (%) |
|----------|---------------|-----|-----------|-----------|---------------|-----|-----------|
| 200      | 400           | 7   | 99        | 20        | 400           | 7.5 | 61        |
| 50       | 400           | 5.5 | 82        | 20        | 150           | 6.5 | 0         |
| 200      | 150           | 7   | 68        | 200       | 400           | 7.5 | 78        |
| 200      | 150           | 5.5 | 55        | 200       | 150           | 6.5 | 23        |
| 50       | 400           | 7   | 80        | 20        | 400           | 6.5 | 74        |
| 50       | 150           | 5.5 | 16        | 20        | 150           | 7.5 | 0         |
| 200      | 400           | 5.5 | 85        | 200       | 400           | 6.5 | 84        |
| 50       | 150           | 7   | 0         | 200       | 150           | 7.5 | 24        |

**Fig. 1** and **Fig. 2** show contour plots for yield with MES at pH 6.25 and MOPS at pH 7.0, respectively. As expected from literature, for a given pH and concentration of MES, yield increases monotonically with increasing Arginine concentration. Surprisingly, for a given pH and Arginine concentration, yield increases monotonically with MES and MOPS concentration as well. Quantitatively, MES was found to be at least as effective as arginine in desorption, whereas, MOPS was slightly less effective than arginine. In summary, both MES and MOPS act similar to additives like arginine in desorbing the biomolecule from the multimodal resin.

MES and MOPS are considered to be particularly suitable additives for elution of biomolecules in downstream processing. This is because (i) MES and MOPS solutions are of lower conductivity than arginine and guanidine, allowing easier processing on a following ion exchange step; and (ii) solutions containing MES and MOPS are expected

to be less disruptive for biomolecules than other additives, because biomolecular solutions of MES and MOPS are regularly used in biochemistry as part of Good's buffer systems.

Just like other additives (such as NaCl, urea, guanidinium salts etc.) can be applied to many different multimodal chromatographic resins, MES and MOPS can be used in desorption of biomolecules from other multimodal chromatographic resins available, such as Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP HyperCel™, Capto™ MMC, and potentially various others multimodal resins which combine hydrophobic, ionic and/or hydrogen bonding interactions.

Furthermore, it is expected that other Good's buffers (e.g., ADA, ACES, BES, Bicine, CAPS, CAPSO, CHES, HEPES, MES, MOPS, PIPES, TAPS, TES, and Tris) exhibit similar advantageous properties as MES and MOPS in the desorption of biomolecules from multimodal chromatographic resins.

**Example 2:** Desorption of a biomolecule using a combination of a Good's buffer and glycol

The experiment from Example 1 was modified in that arginine was replaced by ethylene glycol. The contour plot of this experiment is shown in **Fig. 3**.

**Table 2:** Experimental design and results for elution with MES and glycol

| MES (mM) | Glycol (% v/v) | Yield (%) |
|----------|----------------|-----------|
| 225      | 58             | 95        |
| 402      | 30             | 108*      |
| 100      | 10             | 0         |
| 225      | 30             | 29        |
| 225      | 30             | 14        |
| 225      | 2              | 2         |
| 350      | 50             | 95        |
| 225      | 30             | 13        |
| 350      | 10             | 96        |
| 225      | 30             | 23        |
| 48       | 30             | 10        |
| 100      | 50             | 14        |
| 225      | 30             | 33        |

\* more than 100% yield is not possible and is only a result of measurement errors. It is, however, expected to be close to 100%.

Other embodiments are in the claims.

## CLAIMS

1. A method for purifying or enriching a biomolecule, said method comprising the steps:
  - 5 (a) providing an aqueous solution containing said biomolecule;
  - (b) contacting said aqueous solution with a multimodal resin;
  - (c) optionally washing the multimodal resin with an aqueous washing buffer;
  - (d) eluting the biomolecule from the multimodal resin by an aqueous buffer solution,
    - 10 wherein said aqueous buffer solution comprises one or more of:
      - (i) at least 15 mM of a Good's buffer;
      - (ii) between 0 and 400 mM alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine;
      - 15 (iii) between 0 and 400 mM guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, or taurine; and
      - 20 (iv) between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol;
      - 25 and
  - (e) collecting biomolecule containing fractions, thereby obtaining the biomolecule in purified or enriched form.
2. The method according to claim 1, wherein the aqueous buffer solution comprises
  - 30 15 mM to 2.5 M Good's buffer.
3. The method according to claim 1 or claim 2, wherein the Good's buffer is MES, MOPS, or Tris.
- 35 4. The method according to claim 1 or claim 2, wherein the Good's buffer is MES.
5. The method according to claim 1 or claim 2, wherein the Good's buffer is MOPS.

6. The method according to any one of claims 1 to 5, wherein the biomolecule is a protein.
- 5 7. The method according to claim 6, wherein the protein is an enzyme.
8. The method according to claim 7, wherein the enzyme is a glycosidase.
9. The method according to claim 8, wherein the glycosidase is an alpha-galactosidase.
- 10 10. The method according to claim 9, wherein the alpha-galactosidase is selected from the group consisting of agalsidase alpha and agalsidase beta.
- 15 11. The method according to any one of claims 1 to 10, wherein the multimodal resin is selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP HyperCel™, and Eshmuno® HCX.
- 20 12. A purified biomolecule obtained by the method according to any one of claims 1 to 11.
13. A preparation comprising:
  - (a) a biomolecule; and
  - 25 (b) an aqueous buffer solution comprising one or more of:
    - (i) at least 15 mM Good's buffer;
    - (ii) between 0 and 400 mM alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan,
    - 30 tyrosine, or valine;
    - (iii) between 0 and 400 mM guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride,
    - 35 potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, or taurine; and

(iv) between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.

14. The preparation according to claim 13, wherein the aqueous buffer solution comprises 15 mM and 2.5 M Good's buffer.
15. The preparation according to claim 13 or 14, wherein the at least one Good's buffer is selected from the group consisting of MES, MOPS, or Tris.
16. The preparation according to claim 13 or 14, wherein the Good's buffer is MES.
17. The preparation according to claim 13 or 14, wherein the Good's buffer is MOPS.
18. The preparation according to any one of claims 13 to 17, wherein the biomolecule is a protein.
19. The preparation according to claim 18, wherein the protein is an enzyme.
20. The preparation according to claim 19, wherein the enzyme is a glycosidase.
21. The preparation according to claim 20, wherein the glycosidase is an alpha-galactosidase.
22. The preparation according to claim 21, wherein the alpha-galactosidase is selected from the group consisting of agalsidase alpha and agalsidase beta.
23. A method for desorbing a biomolecule from a multimodal resin, comprising contacting a biomolecule bound to a multimodal resin with an aqueous buffer, wherein said aqueous buffer solution comprises one or more of:
  - (i) at least 15 mM Good's buffer;
  - (ii) between 0 and 400 mM alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine;
  - (iii) between 0 and 400 mM guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium

- caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, or taurine; and
- 5 (iv) between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol.
24. A method for increasing the recovery rate of a biomolecule adsorbed on a multimodal resin by contacting said multimodal resin onto which said biomolecule
- 10 has been adsorbed with an aqueous buffer solution, wherein said aqueous buffer solution comprises one or more of:
- (i) at least 15 mM Good's buffer;
- (ii) between 0 and 400 mM alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine,
- 15 lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine;
- (iii) between 0 and 400 mM guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium
- 20 caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate, calcium phosphate, or taurine; and
- (iv) between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a
- 25 mixture of ethylene glycol and propylene glycol.
25. The method of claim 23 or claim 24, wherein the aqueous buffer solution comprises between 15 mM and 2.5 M Good's buffer.
- 30 26. The method of any one of claims 23 to 25, wherein the Good's buffer is selected from the group consisting of MES, MOPS, and Tris.
27. The method of any one of claims 23 to 25, wherein the Good's buffer is MES.
- 35 28. The method of any one of claims 23 to 25, wherein the Good's buffer is MOPS.
29. The method of any one of claims 23 to 28, wherein the biomolecule is a protein.

30. The method of claim 29, wherein the protein is an enzyme.
31. The method of claim 30, wherein the enzyme is a glycosidase.
- 5 32. The method of claim 31, wherein the glycosidase is an alpha-galactosidase.
33. The method of claim 32, wherein the alpha-galactosidase is selected from the group consisting of agalsidase alpha and agalsidase beta.
- 10 34. The method of any one of claims 23 to 33, wherein the multimodal resin is selected from the group consisting of Capto™ Adhere, Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP HyperCel™, and Eshmuno® HCX.
- 15 35. An article of manufacture comprising:
- (a) a packaging material or container;
- (b) an aqueous buffer solution comprising one or more of:
- (i) at least 15 mM Good's buffer;
- (ii) between 0 and 400 mM alanine, arginine, asparagine, aspartic  
20 acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine;
- (iii) between 0 and 400 mM in total concentration of urea, guanidinium chloride, guanidinium thiocyanate, ammonium chloride, ammonium  
25 sulfate, ammonium phosphate, ammonium acetate, sodium chloride, sodium citrate, sodium caprylate, sodium phosphate, sodium iodide, potassium chloride, potassium citrate, potassium sulfate, potassium phosphate, potassium iodide, magnesium chloride, magnesium sulfate, magnesium phosphate, calcium chloride, calcium sulfate,  
30 calcium phosphate, or taurine; and
- (iv) between 1% and 60% (v/v) of ethylene glycol or propylene glycol or a mixture of ethylene glycol and propylene glycol;
- and
- (c) a data carrier containing instructions for carrying out a method according  
35 to any one of claims 1 to 11.

36. The article of manufacture according to claim 35, wherein the data carrier is a label present on the packaging material or container or a packaging insert.
37. The article of manufacture according to claim 35 or claim 36, wherein the data carrier comprises written information, visual information, or computer-readable information.
38. The article of manufacture according to any one of claims 35 to 37, wherein the data carrier further comprises one or more of the following information:
- (i) instructions for use of aqueous buffer solution;
  - (ii) quality information such as information about the lot/batch number, the manufacturing or assembly site or the expiry or sell-by date of the aqueous buffer solution; or information concerning the correct storage or handling of the article;
  - (iii) information concerning the composition of the aqueous buffer solution;
  - (iv) information concerning the interpretation of information obtained when performing methods according to any one of claims 1 to 11;
  - (v) a warning concerning possible misinterpretations or wrong results when applying unsuitable methods; and
  - (vi) a warning concerning possible misinterpretations or wrong results when using unsuitable reagent(s) and/or buffer(s).
39. The article of manufacture according to any one of claims 35 to 38, further comprising one or more of the following components:
- (d) a chromatography medium;
  - (e) a chromatography column; and
  - (f) a reference biomolecule usable for calibrating a method for purifying or enriching a biomolecule.
40. The article of manufacture according to claim 39, wherein said chromatography column is prepacked with said chromatography medium.
41. The article of manufacture according to any one of claims 39 to 40, wherein said chromatography medium is a multimodal chromatography resin.
42. The article of manufacture according to claim 41, wherein the multimodal chromatography resin is selected from the group consisting of Capto™ Adhere,

Capto™ MMC, Pall PPA HyperCel™, Pall HEA HyperCel™, Pall MEP HyperCel™, and Eshmuno® HCX.

- 5      43.      The article of manufacture according to any one of claims 39 to 42, wherein the reference biomolecule is a protein or a nucleic acid.
44.      The article of manufacture according to any one of claims 35 to 43, wherein the aqueous buffer solution comprises between 15 mM and 2.5 M Good's buffer.
- 10    45.      The article of manufacture according to any one of claims 35 to 44, wherein the at least one Good's buffer is selected from the group consisting of MES, MOPS, and Tris.
- 15    46.      The article of manufacture according to any one of claims 35 to 45, wherein the Good's buffer is MES.
47.      The article of manufacture according to any one of claims 35 to 45, wherein the Good's buffer is MOPS.

20

FIG. 1






FIG. 2



Design-Expert® Software

Factor Coding: Actual

Yield (%)



X1 = A: MOPS (mM)

X2 = B: Arg (mM)

Actual Factor

C: pH = 7.00

FIG. 3




FIG. 4




FIG. 5




FIG. 6A




FIG. 6B



FIG. 6C





FIG. 7



# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2013/066319

A. CLASSIFICATION OF SUBJECT MATTER  
INV. B01D15/38 C07K1/16  
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
B01D C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EP0-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages                                                                                               | Relevant to claim No. |
|-----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X         | WO 2012/059495 A1 (HOFFMANN LA ROCHE [CH];<br>FALKENSTEIN ROBERTO [DE]; LEMM THORSTEN<br>[DE];) 10 May 2012 (2012-05-10)<br>examples 2-4                                         | 1-47                  |
| Y         | -----<br>EP 2 287 174 A1 (BIO RAD LABORATORIES<br>[US]) 23 February 2011 (2011-02-23)<br>examples                                                                                | 1-47                  |
| Y         | -----<br>US 2006/246544 A1 (KANG YUN [US] ET AL)<br>2 November 2006 (2006-11-02)<br>paragraph [0158]<br>paragraph [0168] - paragraph [0171]<br>paragraph [0303]<br>-----<br>-/-- | 1-47                  |



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 January 2014

Date of mailing of the international search report

23/01/2014

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040,  
Fax: (+31-70) 340-3016

Authorized officer

Fourgeaud, Damien

# INTERNATIONAL SEARCH REPORT

International application No

PCT/US2013/066319

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages                                                                                             | Relevant to claim No. |
|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| A         | US 2011/178276 A1 (CUMMINGS LARRY J [US]<br>ET AL) 21 July 2011 (2011-07-21)<br>paragraph [0022] - paragraph [0024]<br>paragraph [0057]; example 8<br>claims<br>-----          | 1-47                  |
| A         | WO 2011/121031 A1 (OCTAPHARMA AG [CH];<br>GILLJAM GUSTAV [SE]; WINGE STEFAN [SE];<br>TIEMEYER M) 6 October 2011 (2011-10-06)<br>page 8, line 1 - line 17<br>example 3<br>----- | 1-47                  |
| X,P       | WO 2013/067301 A1 (GENENTECH INC [US];<br>HOFFMANN LA ROCHE [CH])<br>10 May 2013 (2013-05-10)<br>paragraph [0091]<br>paragraph [0137] - paragraph [0145]<br>-----              | 1-47                  |

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2013/066319

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s)                                                                                                                                                              | Publication<br>date                                                                                                                      |
|-------------------------------------------|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------|
| WO 2012059495 A1                          | 10-05-2012          | CA 2815282 A1<br>CN 103201288 A<br>EP 2635600 A1<br>KR 20130067306 A<br>US 2013217866 A1<br>WO 2012059495 A1                                                                            | 10-05-2012<br>10-07-2013<br>11-09-2013<br>21-06-2013<br>22-08-2013<br>10-05-2012                                                         |
| EP 2287174 A1                             | 23-02-2011          | EP 2238154 A1<br>EP 2242762 A1<br>EP 2275431 A2<br>EP 2287174 A1<br>JP 2011510303 A<br>US 2009186396 A1<br>US 2009187005 A1<br>US 2012077961 A1<br>WO 2009092010 A1<br>WO 2009092014 A1 | 13-10-2010<br>27-10-2010<br>19-01-2011<br>23-02-2011<br>31-03-2011<br>23-07-2009<br>23-07-2009<br>29-03-2012<br>23-07-2009<br>23-07-2009 |
| US 2006246544 A1                          | 02-11-2006          | EP 1866427 A2<br>JP 2008538181 A<br>US 2006246544 A1<br>WO 2006105426 A2                                                                                                                | 19-12-2007<br>16-10-2008<br>02-11-2006<br>05-10-2006                                                                                     |
| US 2011178276 A1                          | 21-07-2011          | CA 2787009 A1<br>EP 2524215 A1<br>JP 2013517284 A<br>US 2011178276 A1<br>WO 2011088225 A1                                                                                               | 21-07-2011<br>21-11-2012<br>16-05-2013<br>21-07-2011<br>21-07-2011                                                                       |
| WO 2011121031 A1                          | 06-10-2011          | AU 2011234532 A1<br>CA 2793143 A1<br>CN 102858797 A<br>EP 2552948 A1<br>JP 2013523690 A<br>KR 20130010897 A<br>US 2013096279 A1<br>WO 2011121031 A1                                     | 20-09-2012<br>06-10-2011<br>02-01-2013<br>06-02-2013<br>17-06-2013<br>29-01-2013<br>18-04-2013<br>06-10-2011                             |
| WO 2013067301 A1                          | 10-05-2013          | AU 2012323995 A1<br>WO 2013067301 A1                                                                                                                                                    | 16-05-2013<br>10-05-2013                                                                                                                 |