

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2007 (15.11.2007)

PCT

(10) International Publication Number
WO 2007/129895 A9

(51) International Patent Classification:
C07K 16/28 (2006.01)

(21) International Application Number:
PCT/NL2007/050202

(22) International Filing Date: 9 May 2007 (09.05.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
06076028.7 9 May 2006 (09.05.2006) EP
60/798,819 9 May 2006 (09.05.2006) US

(71) Applicant (for all designated States except US): **PAN-GENETICS B.V.** [NL/NL]; Yalelaan 46, NL-3584 CM Utrecht (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DEN HARTOG, Marcel, Theodorus** [NL/NL]; Ossenvenne 36, NL-1141 RS Monnickendam (NL). **VAN NEERVEN, Ruprecht, Jules, Joost** [NL/NL]; Prozastraat 90, NL-1321 HC Almere (NL). **JOHNSON, Kevin, Stuart** [GB/GB]; Bower Cottage, 24 East St., Lilley Hertfordshire LU2 8LW (GB). **CASSON, Robert, Duncan** [GB/GB]; The Paddock, Pedlers Lane, Therfield, Royston Hertfordshire SG8 9PX (GB).

(74) Agent: **VAN LOON, C., J., J.**; Johan de Wittlaan 7, Verenigde, NL-2517 JR Den Haag (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
31 December 2008

(48) Date of publication of this corrected version:
19 February 2009

(15) Information about Correction:
see Notice of 19 February 2009

(54) Title: DEIMMUNIZED ANTAGONISTIC ANTI-HUMAN CD40 MONOCLONAL ANTIBODY FROM THE CH5D12 ANTIBODY

(57) Abstract: The present invention provides novel antagonistic anti-human CD40 monoclonal antibodies, methods for generating them and uses thereof.



WO 2007/129895 A9

DEIMMUNIZED ANTAGONISTIC ANTI-HUMAN CD40 MONOCLONAL ANTIBODY
FROM THE CH5D12 ANTIBODY

The invention relates to human, humanized and monoclonal antibodies with decreased immunogenicity, uses and methods of production thereof. The invention in particular relates to antagonistic anti-human CD40 monoclonal antibodies.

5

The CD40 molecule is a 50 kDa type I membrane glycoprotein and is expressed on B cells, monocytes/macrophages, dendritic cells (DCs) and activated endothelial cells.¹⁻⁶ Under certain conditions, CD40 can also be found on fibroblasts, epithelial cells and keratinocytes.⁷ CD40 ligand (CD40L, CD154), a 32 kDa type II integral membrane glycoprotein, is transiently expressed on activated CD4⁺ T cells and a small population of activated CD8⁺ T cells.^{8, 9} In addition, CD40L has been found on a number of other cell types after activation, including mast cells, basophils, B cells, eosinophils, DCs and platelets.^{10, 11}

15

Studies in murine models have clearly demonstrated the involvement of the CD40L-CD40 interaction in the pathophysiology of various autoimmune diseases (for review, see reference¹²). Evidence from CD40L transgenic mice, which acquire lethal inflammatory bowel disease, provided the first evidence that CD40-CD40L interactions might also play a role in the pathogenesis of inflammatory bowel diseases.¹³ An anti-mouse CD40L monoclonal antibody (Mab) effectively prevents mucosal inflammation and interferon- γ production by lamina propria CD4⁺ T cells in TNBS-induced colitis.¹⁴ In a Severe Combined Immunodeficiency (SCID) mouse inflammatory bowel disease model it was shown that treatment with anti-CD40L from the day of T-cell reconstitution completely prevented clinical and histological appearance of experimental colitis.¹⁵ Furthermore, anti-CD40L administration from week 5 after T-cell reconstitution could still prevent progression of the disease and treated animals showed improvement in disease symptoms and histology

25

compared with control animals.¹⁵ In addition, reconstitution of SCID mice with T cells from CD40L knock-out mice, further demonstrated the essential role of CD40L-expressing T cells in disease development and interleukin-12 production.¹⁶

5

The CD40-CD40L interaction can be antagonized with monoclonal antibodies (Mabs) against either CD40L or CD40. The expression of CD40L on activated platelets has resulted in thrombo-embolic events during treatment of humans with IgG₁ anti-human CD40L Mabs at higher dose levels and
10 termination of the development of these Mabs.^{17, 19} Antagonizing CD40 therefore seems a more attractive approach. The non-stimulatory antagonistic activity of Mab 5D12 (anti-human CD40) was demonstrated in various in vitro studies using different CD40-bearing cell types^{20, 22} and chimeric 5D12 (ch5D12) antagonist activity was validated in vivo using various non-human
15 primate disease models.^{23, 27} ch5D12 is a molecularly engineered human IgG₄ antibody containing the murine variable domains of the heavy and light chains of 5D12 and was constructed to reduce the potential for immunogenicity and to enhance the in vivo half-life of the murine 5D12 Mab when used in humans.

20 Patients with Crohn's disease suffer from a debilitating inflammatory disorder of the gastrointestinal tract of which the precise aetiology and pathogenesis remain elusive.^{28, 29} The disease is characterized by an influx into diseased mucosa of activated T cells, B cells and macrophages,^{30, 31} local production of soluble mediators of inflammation, and damage of involved
25 tissues.^{28, 29} Mucosal CD4⁺ T cells and macrophages and cytokines such as tumour necrosis factor (TNF)- α and IL-12 have been shown to play a central role in initiating an inflammatory loop in Crohn's disease.^{32, 38} T cells from inflamed mucosa exhibit a higher proliferative capacity,^{28, 29} and secrete increased amounts of IFN- γ and IL-2. Increased levels of T-cell associated
30 cytokine mRNA transcripts have been found in mucosal biopsies from Crohn's

disease patients.³³ A dominant role of CD40L on the activated CD4⁺ T cells has been suggested in our studies on CD40/CD40L expression in Crohn's disease lesions.³⁹ CD40L can mediate a strong activation of CD40-bearing cells, predominantly B cells and macrophages, thus resulting in increased

5 production of TNF- α and IL-12 in lesions. Using immunohistochemistry, increased staining with 5D12 was found in all samples of diseased areas from Crohn's disease patients compared to non-diseased areas. Double staining for CD40 and CD20 (B cells) or CD68 (macrophages) indicated that in the sections

10 from patients with Crohn's disease, CD40⁺ cells were mainly B cells in the lymphoid follicles and macrophages in the lamina propria. Lamina propria T cells from inflamed mucosa of Crohn's disease patients induced monocytes to produce significant amounts of IL-12 and TNF- α after 48 h of co-culture. Addition of 5D12 resulted in reduced IL-12 and TNF- α production; levels of production were reduced to the levels observed using control lamina propria T

15 cells both in the absence and presence of IFN- γ .³⁹

It is an object of the invention to provide alternative molecules that share at least the kind of safety and/or efficacy of 5D12 *in vivo*, not necessarily the amount of said safety and/or efficacy. The antibody 5D12, or at least the

20 variable domains thereof, have a murine background. The present invention provides variants of the heavy and light chain variable domains of 5D12. To this end the invention provides a polypeptide comprising an amino acid sequence of formula (I)

25	1	11	21	31
	GFSX ₁ S RYSVY WX ₂ RQP PGKGX ₃ EWX ₄ GM MWGGG STDYS			
	41	51	61	
30				

TSLKS RLTIS KDTSK SQVX₅L KMNSL RTDDT AMYYC

71

|

5 VRTDG DY

wherein:

X₁ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;

X₂ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;

X₃ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;

10 X₄ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H; and

X₅ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H.

Said polypeptide comprises extensive sequence identity with the heavy chain variable domain of the 5D12 antibody, however, the polypeptide is less immunogenic in a human individual administered as such or in the context of an antibody comprising said polypeptide. At the indicated positions X₁-X₅ several amino acids may be present as indicated. A binding molecule comprising a polypeptide of the invention has good CD40 binding properties. It has been noted that production of the antibody in a cell varies somewhat with the kind of amino acid at the indicated positions. This will be detailed elsewhere herein below.

The invention further provides a polypeptide according to the invention comprising an amino acid sequence

25

1	11	21	31

QVKLQ ESGPG LVKPS ETL SI TCTVS GFSX₁S RYSVY WX₂RQP

30

41	51	61	71
----	----	----	----

PGKGX₃ EWX₄GM MWGGG STDYS TSLKS RLTIS KDTSK

	81	91	101	111
5				
	SQVX ₅ L KMNSL RTDDT AMYYC VRTDG DYWGQ GTTVT VSS.			

The abovementioned polypeptide essentially spans the variable domain of the heavy chain of the murine 5D12 antibody. The amino acid sequence is altered at several positions with respect to murine sequence. The altered polypeptide couples good binding properties, while at the same time being well tolerated by humans provided with the polypeptide. At the indicated positions various amino acids can be inserted without dramatically reducing and/or altering at least immunological properties of the polypeptide when compared to the original murine polypeptide. It is preferred that a polypeptide of the invention comprises a G, A, V, L, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H at position X₁. In this way at least the production properties of the polypeptide in a mammalian cell are not dramatically reduced and/or altered when compared to the original murine or chimeric polypeptide. In a particularly preferred embodiment

X_1 is G, A, V, L, P, F or M;
 X_2 is G, A, V, L, I, P, F or M;
 X_3 is G, A, V, L, I, P, F, M;
 X_4 is G, A, V, L, I, P, F, M; and
25 X_5 is G, A, V, L, I, P, F, M, W, C, N, Q, S, T or Y.

These polypeptides are better suited for high level production in a mammalian cell. Particularly in the context of an antibody.

The invention further provides a polypeptide according to the
30 invention, wherein:

X₁ is G, A, V, L or M;

X₂ is G, A, V, L, I or M;

X₃ is G, A, V, L, I, P, F, M;

X₄ is G, A, V, L, I or M; and

5 X₅ is P, F, W, N, Q, S, T or Y.

These polypeptides are better suited for high level antibody production while at the same time exhibiting good tolerance properties in a human. The invention further provides a polypeptide of formula (I) wherein X₁ is L; X₂ is I; X₃ is P; X₄ is M; and/or X₅ is S. This polypeptide is particularly preferred
10 because of its excellent production properties in the context of an antibody that mimics the binding and pharmacological properties of antibody ch5D12 whereas immunological properties are improved in humans when compared to the murine or ch5D12 antibody and wherein at least production of the polypeptide is not dramatically reduced when compared to the murine or
15 chimeric counterpart.

The invention further provides a polypeptide of formula (I) wherein X₁ is I; X₂ is V; X₃ is P; X₄ is M; and/or X₅ is S. This polypeptide is particularly preferred because of its excellent production properties in the context of an antibody that mimics the binding and pharmacological properties of antibody ch5D12
20 whereas immunological properties are improved in humans when compared to the murine or ch5D12 antibody and wherein at least production of the polypeptide is not dramatically reduced when compared to the murine or chimeric counterpart.

25

An antibody that deviates at least at one position X₁-X₅ when compared to chimeric 5D12 at the corresponding positions, has better immunological properties than the chimeric 5D12 antibody in humans. In a preferred embodiment of such antibodies the invention preferably provides a
30 polypeptide according to the invention, wherein: X₁ is I and X₂ is V; X₁ is I and

X₂ is I; X₁ is L and X₂ is I; or X₁ is L and X₂ is V. The polypeptides are particularly preferred in combination with an X₃ is P; X₄ is M; and X₅ is either F or S; In one embodiment the invention provides a polypeptide according to the invention, wherein: X₁ is L; X₂ is V; X₃ is L; X₄ is L and X₅ is F. Production of antibodies comprising said polypeptide is very good, while simultaneously providing improved immunological properties in humans when compared to ch5D12. The invention further provides a polypeptide comprising amino acid sequence GFSX₁S RYSVY WX₂R, wherein: X₁ is L and X₂ is I; or X₁ is I and X₂ is V. This polypeptide comprises a modified CDR1 of a 5D12 heavy chain variable fragment. This CDR1 comprises at least one different amino acid when compared to the CDR1 of the heavy chain variable fragment of antibody 5D12. This amino acid change both leads to improved immunological properties of a modified 5D12 or ch5D12 antibody wherein the modification comprises at least a replacement of the corresponding sequence of said polypeptide in 5D12 or ch5D12 with a polypeptide of the invention, whereas at the same time enabling good production of the antibody in a mammalian cell.

The invention further provides a heavy chain variable domain comprising a polypeptide of formula (I). Said variable domain comprises preferably 90-130, more preferably 100-120, more preferably 105-115, most preferably 113 amino acids. The polypeptide can be generated synthetically or by a cell. Preferably said heavy chain variable domain is produced by a cell. In nature at least five types of heavy chain exist: γ , δ , α , μ and ϵ , wherein each type defines a class of immunoglobulins. A polypeptide of the invention may be used as a binding body directly or may be incorporated into an antibody. When incorporated into an antibody the polypeptide is preferably combined with a constant part of an antibody heavy chain. To this end the invention further provides an antibody heavy chain comprising a polypeptide of formula (I). The art knows many derivatives and analogues of variable domain antibodies. However, currently many different parts, derivatives and/or analogues of antibodies are in use.

Non-limiting examples of such parts, derivatives and/or analogues are, single chain Fv-fragments, monobodies, VHH, Fab-fragments, artificial binding proteins such as for example avimers, and the like. A common denominator of such specific binding bodies is the presence of a heavy chain variable domain.

- 5 Thus the invention further provides a binding body comprising a polypeptide of formula (I).

A preferred binding body of the invention is an antibody, as an antibody comprises a naturally occurring structure. Therefore, the invention in a preferred embodiment provides an antibody comprising a polypeptide

- 10 according to the invention.

- A binding body according to the invention is preferably a binding body that is well tolerated in an animal. Tolerance of an animal for a polypeptide is governed by many different aspects. Immunity, be it T-cell mediated, B-cell
15 mediated or other is one of the variables that are encompassed in tolerance of the animal for a polypeptide. As mentioned above, the 5D12 antibody has a murine background. The polypeptide of formula (I) has a reduced immunogenicity in human. It is therefore sometimes referred to as a deimmunized variant of the heavy chain variable domain of 5D12. Thus in an
20 aspect the invention provides an antibody comprising an epitope specificity of a 5D12 antibody, wherein the heavy chain of said antibody is a polypeptide of formula (I). Deimmunized as used herein is defined as less immunogenic in an animal than the original antibody. A polypeptide of formula (I) is deimmunized when compared to the heavy chain in 5D12 through the removal of known
25 human T cell epitopes. T cell epitopes are amino acid sequences within proteins with the capacity to bind to MHC class II molecules. By removal of the T cell epitopes the antibody is less immunogenic. Preferably a variable domain of the invention is further humanized, such as for instance veneered. By using veneering techniques, exterior residues which are readily
30 encountered by the immune system are selectively replaced with human

residues to provide a hybrid molecule that comprises either a weakly immunogenic or substantially non-immunogenic veneered surface. An animal as used in the invention is preferably a mammal, more preferably a primate, most preferably a human.

5

An antibody according to the invention preferably comprises a constant region of a human antibody. According to differences in their heavy chain constant domains, antibodies are grouped into five classes, or isotypes: IgG, IgA, IgM, IgD, and IgE. These classes or isotypes comprise at least one of
10 said heavy chains that is named with a corresponding Greek letter. In a preferred embodiment the invention provides an antibody according to the invention wherein said constant region is selected from the group of IgG, IgA, IgM, IgD, and IgE constant regions, more preferably said constant region comprises an IgG constant region, more preferably an IgG₁ constant region,
15 preferably a mutated IgG₁ constant region, most preferably said constant region is an IgG₄ constant region. Furthermore, said IgG₄ constant region is preferably a human IgG₄ constant region. Preferably, the IgG₄ antibody of the invention comprises the constant regions of the heavy and light chain amino acid sequence as depicted in figure 18. Preferably, the IgG₄ antibody of the
20 invention comprises the heavy and light chain amino acid sequence as depicted in figure 18. Some variation in the constant region of IgG₄ occurs in nature and/or is allowed without changing the immunological properties of the resulting antibody. Typically between about 1-5 amino acid substitutions are allowed in the constant region. An antibody with an IgG₄ constant region or a
25 mutated IgG₁ constant region has at least most of the pharmacological properties of an antibody but does not bind complement, and will thus not induce depletion of the cell its binds to *in vivo*. Preferably said constant region is a constant region of a human antibody.

In one embodiment the invention provides a nucleic acid encoding a
30 polypeptide according to the invention, and/or a binding body according to the

invention, and/or an antibody according to the invention. A nucleic acid as used in the invention is typically but not exclusively a ribonucleic acid (RNA) or a deoxyribonucleic acid (DNA). Alternative nucleic acids are available for a person skilled in the art, such as for instance peptide nucleic acids (PNA). A
5 nucleic acid according to the invention is for instance comprised in a cell. When said nucleic acid is expressed in said cell, said cell produces a polypeptide and/or a binding body and/or an antibody according to the invention. Therefore, the invention in one embodiment provides a cell comprising a polypeptide according to the invention, a binding body according
10 to the invention, an antibody according to the invention and/or a nucleic acid according to the invention. Said cell is preferably an animal cell, more preferably a mammal cell, more preferably a primate cell, most preferably a human cell. For the purposes of the invention a suitable cell is any cell capable of comprising and preferably of producing a polypeptide according to the
15 invention, a binding body according to the invention, an antibody according to the invention and/or a nucleic acid according to the invention.

The invention further provides a cell comprising an antibody according to the invention. Preferably said cell produces said antibody. In a preferred embodiment said cell is a hybridoma cell, a Chinese hamster ovary
20 (CHO) cell, an NS0 cell or a PER-C6TM cell. In a particularly preferred embodiment said cell is a CHO cell. Further provided is a cell culture comprising a cell according to the invention. Various institutions and companies have developed cell lines for the large scale production of antibodies, for instance for clinical use. Non-limiting examples of such cell
25 lines are CHO cells, NS0 cells or PER.C6TM cells. These cells are also used for other purposes such as the production of proteins. Cell lines developed for industrial scale production of proteins and antibodies are herein further referred to as industrial cell lines. Thus in a preferred embodiment the invention provides the use of a cell line developed for the large scale

production of protein and/or antibody for the production of an antibody of the invention.

The invention further provides a method for producing an antibody comprising culturing a cell of the invention and harvesting said antibody from said

5 culture. Preferably said cell is cultured in a serum free medium. Preferably said cell is adapted for suspension growth. Further provided is an antibody obtainable by a method for producing an antibody according to the invention. The antibody is preferably purified from the medium of the culture. Preferably said antibody is affinity purified.

10 A cell of the invention is for instance a hybridoma cell line, a CHO cell, an NS0 cell or another cell type known for its suitability for antibody production for clinical purposes. In a particularly preferred embodiment said cell is a human cell. Preferably a cell that is transformed by an adenovirus E1 region or a functional equivalent thereof. A preferred example of such a cell line is the
15 PER.C6TM cell line or equivalent thereof. In a particularly preferred embodiment said cell is a CHO cell or a variant thereof. Preferably a variant that makes use of a Glutamine synthetase (GS) vector system for expression of an antibody.

20 It has been noted that some amino acids at the mentioned positions X₁-X₅ are less suited for high level production of an antibody comprising a polypeptide of formula (I) in an antibody producing cell. In a preferred embodiment a polypeptide of formula (I) in said antibody comprises X₁-X₅ wherein:

25 X₁ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H. In a particularly preferred embodiment X₁ is G, A, V, L, I, P, F or M; X₂ is G, A, V, L, I, P, F or M; X₃ is G, A, V, L, I, P, F, M; X₄ is G, A, V, L, I, P, F, M; and X₅ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T or Y. More preferably, said antibody comprises a polypeptide of formula I wherein X₁ is G, A, V, L, I or M;
30 X₂ is G, A, V, L, I or M; X₃ is G, A, V, L, I, P, F, M; X₄ is G, A, V, L, I or M;

and X₅ is P, F, W, N, Q, S, T or Y. More preferably X₁ is L; X₂ is I; X₃ is P; X₄ is M; and/or X₅ is S. Particularly preferred is when X₁ is I and X₂ is V; X₁ is I and X₂ is I; X₁ is L and X₂ is I; X₁ is L and X₂ is L; X₁ is V and X₂ is I; X₁ is V and X₂ is V; X₁ is L and X₂ is L; X₁ is V and X₂ is L; or X₁ is L and X₂ is V. These latter
5 polypeptides are particularly preferred in combination with an X₃ is P; X₄ is M; and X₅ is either F or S, preferably S; In one embodiment the invention provides a polypeptide according to the invention, wherein: X₁ is L; X₂ is V; X₃ is L; X₄ is L and X₅ is F. Production of antibodies comprising said polypeptide is good, while simultaneously providing improved immunological properties in humans
10 when compared to ch5D12.

In another preferred embodiment the invention provides a polypeptide according to the invention wherein at least one of X₁, X₂, X₃, X₄ or X₅ is the same as an amino acid at the corresponding position in a sequence
15 that is shown by the invention to yield specifically good expression levels and wherein further at least one of X₁, X₂, X₃, X₄ or X₅ is the same as an amino acid at the corresponding position in a 5D12 amino acid sequence. An advantage of a polypeptide according to the invention wherein at least one of X₁, X₂, X₃, X₄ or X₅ is the same as an amino acid at the corresponding position in a 5D12
20 amino acid sequence, is that a binding body according to the invention comprising such a polypeptide shows a better expression level than a polypeptide according to the invention wherein none of X₁, X₂, X₃, X₄ or X₅ are the same as an amino acid at the corresponding position in a 5D12 amino acid sequence. Without being bound by theory, said better expression level is
25 believed to be due to the fact that an amino acid at the position of X₁, X₂, X₃, X₄ and/or X₅ contributes to a proper assembly of a binding body according to the invention when said amino acid is the same as an amino acid at the corresponding position in a 5D12 amino acid sequence.

An antibody comprising a polypeptide of formula (I) shows non-stimulatory antagonistic activity. As the CD40L-CD40 interaction is involved in the pathophysiology of various inflammatory disorders such as autoimmune diseases and graft rejection, a polypeptide according to the invention is

5 therefore particularly suited for ameliorating a symptom of an inflammatory disorder. In one embodiment an antibody comprises a binding body of the invention. In a preferred embodiment said antibody is a monoclonal antibody. Monoclonal antibody technology allows for production of large amounts of essentially pure antibodies, thus obtaining predictable products. Therefore, the

10 invention in one embodiment provides an antagonistic anti-human CD40 monoclonal antibody comprising a polypeptide according to the invention. A binding body according to the invention is all the more suitable for that purpose since it is in one embodiment deimmunized compared to a mouse 5D12 and/or a chimeric 5D12. Therefore, a binding body of the invention has a

15 reduced immunogenicity and an enhanced half-life in humans compared to a mouse 5D12 and/or a chimeric 5D12. As a result thereof, a binding body of the invention has a sustainable pharmaceutical potential against various inflammatory disorders. Thus, in a preferred embodiment the invention provides a deimmunized antagonistic anti-human CD40 monoclonal antibody

20 according to the invention.

As referred to previously, the present invention provides 5D12 like molecules that comprise amino acid alterations with respect to the 5D12 amino acid sequence, wherein the alterations are at least in the heavy chain variable

25 domain and preferably also in the light chain variable domain. In this context the invention further provides a binding body according to the invention comprising an amino acid sequence of formula (II)

	1	11	21	31
30	X ₆ LGX ₇ X ₈ ASISC RSSQS LX ₉ NSN GNTYL HWYLQ RPGQS			

14

41 51 61 71
 | | | |
 PRLLI YKVSN RFSGV PDRFS GSGSG TDFTL KISRV EAEDX₁₀

5 81 91
 | |
 GVYX₁₁C SQSTH VPWT

wherein:

- X₆ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
 10 X₇ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
 X₈ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
 X₉ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
 X₁₀ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H; and
 X₁₁ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H.
- 15 Said binding body is preferably an antagonistic anti-human CD40 monoclonal antibody according to the invention. In a preferred embodiment an X selected from X₆, X₇, X₈, X₉, X₁₀ or X₁₁ is selected from a group resembling an amino acid at a corresponding position in a 5D12 amino acid sequence and/or resembling an amino acid at a corresponding position in a sequence that is
- 20 shown by the invention to yield good expression levels, as shown in example 2. Therefore, a binding body according to the invention preferably comprises an amino acid sequence of formula (II) wherein:
- X₆ is N, Q, S, T, Y, W or C;
 X₇ is D, E, N, Q, S, T, Y, W or C;
 25 X₈ is N, Q, S, T, Y, G, A, V, L, I, P, F, M, W or C;
 X₉ is G, A, V, L, I, P, F, M;
 X₁₀ is G, A, V, L, I, P, F, M; and
 X₁₁ is N, Q, S, T, Y, G, A, V, L, I, P, F, M, W or C.

In another embodiment the invention provides an antagonistic anti-human CD40 monoclonal antibody according to the invention comprising a polypeptide of formula (II) of the invention comprising an amino acid sequence as follows.

	1	11	21	31
5				
	ELQLT QSPLS LPVX ₆ L GX ₇ X ₈ AS ISCRS SQSLX ₉ NSNGN TYLHW			
	41	51	61	71
10	YLQRP GQSPR LLIYK VSNRF SGVPD RFSGS GSGTD FTLKI			
	81	91	101	111
	SRVEA EDX ₁₀ GV YX ₁₁ CSQ STHVP WTFGG GTKLE IKR.			

- 15 In a preferred embodiment the invention provides a polypeptide of formula (II) of the invention wherein X₆ is T or S, X₇ is D or Q, X₈ is Q or P, X₉ is V or A, X₁₀ is V or L and X₁₁ is F or Y. More preferably wherein: X₆ is T, X₇ is Q, X₈ is P, X₉ is A, X₁₀ is V and X₁₁ is Y. 5D12-like antibodies comprising a polypeptide of formula (I) and the preferred polypeptide of formula (II) combine good
- 20 expression levels in a producing cell with good tolerance properties and pharmacodynamic properties in a human.

An anti human-CD40 antagonist antibody of the invention preferably comprises a heavy chain variable domain amino acid sequence of formula (I)

25 and a light chain variable domain amino acid sequence of formula (II). Such an antibody has good characteristics. It is of course possible to generate variants of such an original antibody by modifying one or more amino acids therein. Many of such variants will behave more or less similar when compared to said original. Such variants are also included in the scope of the invention. There

30 are many ways to modify an antibody of the invention. A non-limiting example

of such a modification is an antibody comprising a pyro-glutamate instead of a glutamate. Other non-limiting examples of such modifications are an insertion, deletion, inversion and/or substitution of one or more amino acids when compared to said original antibody. The present invention provides means and methods for generating such a variant. It also provides tests for determining the characteristics of said a variant. In a preferred embodiment the invention provides a variant of an original antibody of the invention said variant comprising an insertion, deletion, inversion and/or substitution of between about 1 to 10 amino acids when compared to the amino acid sequence of said original antibody. Preferably said insertion, deletion, inversion and/or substitution does not comprise the amino acids at position X₁ and position X₂ of the heavy chain variable domain of formula (I) of the original antibody.

In a preferred embodiment the invention provides a method for selecting an anti-human CD40 antagonist antibody comprising generating a first cell line that produces an original anti-human CD40 antagonist antibody and determining the amount of original antibody that is produced by said first cell line, said original antibody comprising the heavy chain variable domain amino acid sequence

20	1	11	21	31
	QVKLQ ESGPG LVKPS ETL SI TCTVS GFSX ₁ S RYSVY WX ₂ RQP			
	41	51	61	71
25				
	PGKGP EWMGM MWGGG STDYS TSLKS RLTIS KDTSK			
	81	91	101	111
30	SQVSL KMNSL RTDDT AMYYC VRTDG DYWGQ GTT VT VSS			

wherein X_1 and X_2 are pair wise selected from the group consisting of $X_1 = I$ and $X_2 = V$; $X_1 = L$ and $X_2 = I$; $X_1 = V$ and $X_2 = V$; $X_1 = L$ and $X_2 = L$; or $X_1 = L$ and $X_2 = V$,

said method further comprising

- 5 generating at least one further cell line that produces a variant of said original antibody wherein said variant antibody is a modified original antibody comprising an insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids when compared to said original antibody, wherein said modification does not consist of a modification of the amino acids at the
 - 10 position(s) identified by X_1 and X_2 , and determining the amount of variant antibody that is produced by said at least one further cell line,
- said method further comprising selecting a variant antibody that was produced at an amount that was at least 50% of the amount of original antibody.

Preferably said original antibody comprises the light chain amino acid

15 sequence

1	11	21	31
ELQLT QSPLS LPV T L G Q P AS ISCRS SQSL A NSNGN TYLHW			

20	41	51	61	71
	YLQRP GQSPR LLIYK VSNRF SGVPD RFSGS GSGTD FTLKI			

25	81	91	101	111
	SRVEA ED V GV Y Y CSQ STHVP WTFGG GTKLE IKR.			

- Said insertion, deletion, inversion, and/or substitution of between about 1 to 5
- 30 amino acids can be in any part of the antibody not being or involving the

amino acids at positions X₁ and X₂. Preferably, said insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids is in said heavy chain amino acid sequence or said light chain amino acid sequence when compared to the corresponding chain amino acid sequence in said original antibody. Preferably said insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids is in said heavy chain amino acid sequence when compared to said heavy chain sequence of said original antibody. Preferably said method further comprises generating an antibody producer cell line that produces said selected antibody. This producer cell line can be said further cell line, or yet another cell line that produces said selected antibody. Preferably said method further comprises collecting said selected antibody. The invention further provides an isolated and/or recombinant anti-human CD40 antagonist antibody obtainable by a method according to the invention. In a preferred embodiment said anti-human CD40 antagonist antibody comprises a modification of the heavy chain amino acid sequence

	1	11	21	31
	QVKLQ ESGPG LVKPS ETL SI TCTVS GFSX ₁ S RYSVY WX ₂ RQP			
20	41	51	61	71
	PGKGP EWMGM MWGGG STDYS TSLKS RLTIS KDT SK			
	81	91	101	111
25				
	SQVSL KMNSL RTDDT AMYYC VRTDG DYWGQ GTTVT VSS			

wherein X₁ and X₂ are pair wise selected from the group consisting of X₁ = I and X₂ = V; X₁ = L and X₂ = I; X₁ = V and X₂ = V; X₁ = L and X₂ = L; or X₁ = L and X₂ = V,

said modification comprising an insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids when compared to said heavy chain amino acid sequence and wherein said modification does not consist of a modification of the amino acids at the position(s) identified by X₁ and X₂.

5

In one embodiment the invention provides a pharmaceutical composition comprising a polypeptide according to the invention, a binding body according to the invention, an antibody according to the invention, a nucleic acid according to the invention and/or a cell according to the invention.

10 Also, the invention provides a polypeptide according to the invention, a binding body according to the invention, an antibody according to the invention, a nucleic acid according to the invention and/or a cell according to the invention for use as a medicament. Preferably a medicament for ameliorating a symptom of an autoimmune disorder and/or an inflammatory disorder and/or for
15 reducing graft rejection and/or for the treatment of CD40 positive cancers. In a preferred embodiment said autoimmune and/or an inflammatory disorder is selected from the group of inflammatory bowel disease, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, psoriasis, bullous pemphigoides and atopic dermatitis.

20

As a polypeptide according to the invention is particularly suited for ameliorating a symptom of an inflammatory disorder because of its non-stimulatory CD40 antagonizing properties, a polypeptide according to the invention is suitable for ameliorating a symptom of several disorders. An
25 inflammatory disorder as used in the invention is defined as any disease that involves an inflammatory component. For the interest of the invention, an inflammatory disorder specifically includes an autoimmune disorder and/or graft rejection. The central role of CD40-CD40L interaction in the initiation, amplification and prolongation of immune responses makes a polypeptide of

the invention specifically suitable for immune modulation in an autoimmune disorder.

Herein below information is provided on CD40 and CD40L in order to illustrate the role of CD40 and its ligand in inflammatory disorders. The CD40 molecule is a 50 kDa type I membrane glycoprotein and is expressed on B cells, monocytes/macrophages, and dendritic cells (DCs)⁴⁵⁻⁵⁰. Moreover, under pathological conditions CD40 can be found on endothelial cells (ECs), fibroblasts, epithelial cells and keratinocytes⁵¹. CD40 ligand (gp39, TBAM, TRAP, CD40L, CD154), a 32 kDa type II integral membrane glycoprotein, is transiently expressed on activated CD4⁺ T cells and a small population of activated CD8⁺ T cells^{52,53}. In addition, CD40L has been found on a number of other cell types after activation, including mast cells, basophils, B cells, eosinophils, DCs and platelets^{54,55}.

Engagement of CD40 by CD40L triggers a number of biological events in B cells including proliferation, expression of activation markers, immunoglobulin (Ig) production, isotype switching, homotypic adhesion and rescue from apoptosis^{56,57}. However, as described above, the distribution of the CD40 molecule is not restricted to B cells as was originally postulated. Freshly isolated human monocytes express low levels of the CD40 molecule, which can be up-regulated by culture in the presence of IFN- γ ^{47-49,58}. CD40 ligation of monocyte/macrophages induces the secretion of large amounts of pro-inflammatory mediators such as IL-1, TNF- α and IL-12, which induce inflammatory responses and tumoricidal activity^{47-49,58}, and rescue them from apoptosis⁴⁸. CD40 ligation also causes DCs to enhance their differentiation and activation, to enhance expression of costimulatory molecules such as CD86, CD80 and CD58, to increase cytokine production, and to inhibit apoptosis^{50,59}. Furthermore, when expressed under inflammatory conditions, CD40 signaling can induce expression of intercellular adhesion molecule 1 (ICAM-1), vascular

cell adhesion molecule 1 (VCAM-1) and E-selectin on ECs⁵⁵. These results suggest that signaling through CD40 during T cell-EC interactions may be an important step in the regulation of EC activation and leukocyte recruitment into non-lymphoid tissues. In vivo studies have indicated the importance of the
5 CD40-CD40L interactions in the generation of humoral immune responses^{60,61}, in the priming and activation of antigen-specific T cells⁶², in the temporal activation of macrophages⁶³, as well as in the protective cell-mediated immune responses through T-cell mediated macrophage activation against intracellular parasite infections such as *Pneumocystis*, *Cryptosporidium*, and *Leishmania*⁶⁴
10 ⁶⁶.

A specific role for CD40 and its ligand is demonstrated in murine models of autoimmune disease. Animal model studies have clearly indicated the involvement of the CD40L-CD40 interaction in the pathophysiology of
15 various autoimmune diseases. In these studies, using mice suffering from spontaneous or experimental autoimmune diseases, interfering with CD40L-CD40 interactions had clear beneficial effects. A Mab to murine CD40L was shown to prevent or reduce disease symptoms in collagen-induced arthritis, experimental allergic encephalomyelitis (EAE; an animal model for MS), in
20 (SWR x NZB)F1 lupus mice and in non-obese diabetic (NOD) mice that spontaneously develop T-cell dependent autoimmune diabetes. Evidence indicates that CD40-CD40L interactions also play a role in the pathogenesis of inflammatory bowel diseases, which includes Crohn's disease and ulcerative colitis. CD40L transgenic mice with high transgene copy numbers were shown
25 to acquire lethal inflammatory bowel disease marked by infiltration of CD40⁺ cells and CD40L⁺ T cells into diseased tissues⁶⁷. Anti-CD40L Mab effectively prevent mucosal inflammation and IFN- γ production by lamina propria CD4⁺ T cells in animals with colitis that had been induced by 2,4,6-trinitrobenzene sulfonic acid⁴⁴. Very recently a direct comparison was made between anti-TNF- α treatment and interference of the CD40-CD40L pathway using an anti-
30

CD40L Mab in a SCID mouse experimental inflammatory bowel disease model. In this model, syngeneic CD45RB^{high} CD4⁺ cells are injected into SCID mice that subsequently develop diarrhea or soft stools and show progressive weight loss starting 3-5 weeks after T-cell reconstitution, as symptoms of experimental inflammatory bowel disease. Treatment with either anti-TNF- α or anti-CD40L from the day of T-cell reconstitution completely prevented clinical and histological appearance of experimental inflammatory bowel disease. Furthermore, anti-CD40L administration from week 5 after T-cell reconstitution could still prevent progression of the disease and treated animals showed improvement in disease symptoms and histology compared to control animals (unpublished observations).

Recent work has also demonstrated that interference with the CD40-CD40L pathway is strongly immunosuppressive in transplantation models. Combined treatment with allogeneic small lymphocytes or T cell-depleted small lymphocytes plus an antibody to mouse CD40L permitted indefinite pancreatic islet allograft survival in 37 out of 40 recipients that differed in major and minor histocompatibility loci⁶⁸. From these experiments it was concluded that the effective interference of the CD40L-CD40 interaction most likely had resulted in preventing the induction of costimulatory molecules on the small resting lymphocytes by the alloreactive host T cells. In another recent study, it was demonstrated that administration of a Mab to mouse CD40L at the time of transplantation markedly prolonged survival of fully disparate murine cardiac allografts in both naive and sensitized hosts. However, when anti-CD40L therapy was delayed until postoperative day 5, anti-CD40L failed to prolong graft survival. From this study, it was concluded that anti-CD40L therapy inhibited allograft rejection primarily by interfering with T cell help for effector functions. It has further been shown that interfering with CD80/CD86-CD28 and CD40-CD40L pathways simultaneously effectively aborts T cell clonal expansion in vitro and in vivo,

promotes long-term survival of fully allogeneic skin grafts, and inhibits the development of chronic vascular rejection of primarily vascularized cardiac allografts. Furthermore, interfering with a CD40-CD40L pathway, optionally in combination with interference with a CD80/CD86-CD28 pathway, prevents kidney allograft rejection in a rhesus monkey kidney allograft model^{69,70}. For further information on the effects of 5D12 in inflammatory disorders, see for instance references ⁷¹⁻⁸¹.

Multiple sclerosis is an autoimmune disease of the central (cerebrospinal) nervous system. In this disorder, the white matter surrounding nerve fibers becomes hardened. The term multiple sclerosis (MS) literally means "many scars." The hardened areas of nerve tissue are called plaques. The symptoms, severity, and course of multiple sclerosis are highly variable, depending partly on the sites of the plaques and the extent of the deterioration of the white matter. The deterioration of white matter in the nervous system slows nerve impulses leading to nervous system incoordination.

Experimental autoimmune encephalomyelitis (EAE) in the common marmoset (*Callithrix jacchus*) is a useful preclinical model of multiple sclerosis (reviewed in ^{86, 87, 102, 104}). The central nervous system (CNS) white matter lesions that develop in the various versions of this EAE model share pathomorphological, radiological and immunological features with MS ^{95, 98, 101}. Hence, the marmoset EAE model can bridge the wide immunological gap between humans and rodents that hampers the selection of promising treatments in the drug development pipeline at a preclinical stage ^{96, 99, 103}.

Marmosets immunized with rhMOG, a recombinant protein representing the extracellular fragment of human MOG (amino acids 1–125) develop EAE in 100% of the cases, which is due to the presence of the monomorphic MHC class II susceptibility element C_αA-DRB*W1201 in the repertoire of each monkey ^{82,84,85,107}. A particularly useful aspect of this model for therapy development is

that lesions developing in the brain white matter can be visualized and tentatively characterized with clinically relevant magnetic resonance imaging techniques^{92,105}. Longitudinal analysis of the brain white matter lesions with magnetic resonance imaging (MRI) showed a progressive increment of the volume and persistent inflammatory activity in the majority of the lesions. Furthermore, the characterization of the CNS pathology with MRI and previously described histological criteria¹⁰¹ revealed that the majority of the lesions are in an early active stage¹⁰⁶.

10 The rhMOG-induced EAE model has been used to test whether antibodies targeting co-stimulatory molecules of antigen presenting cells (APC) and T-cells are a potential treatment for MS. The interaction of CD40 with its ligand CD154 plays an important role in various immunopathogenic processes that operate in EAE, including B-cell activation, antigen-presenting cell (APC)
15 activation, initiation of antigen-specific T-cell responses and induction of macrophage effector functions^{90,93,97,7}. A study performed in 1996 confirmed that mice treated with an antibody against CD154 are protected against EAE⁸⁸. However, a clinical trial in MS patients with antibody against CD154 was stopped due to unexpected side-effects that were not observed in the
20 animal experiments¹⁷.

The mouse monoclonal antibody (Mab) 5D12 (mu5D12) has been raised against human CD40. The 5D12 antibody appeared a potent inhibitor of CD40-CD40L mediated activation on several cell types and, unlike most other
25 anti-CD40 Mabs, does not exert CD40 stimulatory activity^{21,22,39}. Both the mouse anti-human CD40 antibody mu5D12 and the chimeric version of this antibody, ch5D12, displayed strong suppressive effects on the development of CNS white matter lesions and neurological deficit in the marmoset EAE model and showed no marked side-effects^{23,24}. The same studies showed that
30 intravenously injected anti-CD40 Mab into EAE affected common marmosets

can gain access to the brain white matter lesions where CD40 molecules are prominently expressed on infiltrated macrophages and activated microglia⁹⁵, as was found earlier in MS⁸⁸. This has raised the question whether ch5D12 also has a therapeutic effect on already existing lesions.

5

't Hart et al monitored brain lesion development in 7 rhMOG-immunized monkeys by serial magnetic resonance imaging (MRI) at 2 weeks interval ⁷⁶. The results of this study demonstrated suppression of lesion inflammation in all 3 ch5D12-treated monkeys while lesion enlargement was diminished in 2 of
10 the 3 ch5D12-treated monkeys.

Prevention of CD40 engagement with its ligand CD154 on activated T cells early in the disease process has a significant impact on the clinical and neuropathological expression of EAE in rodents ^{88,89,91,94,100} and non-human
15 primate models ^{23,24}. CD40 is prominently expressed within the CNS white matter lesions of MS patients as well as EAE-affected rodents ⁸⁸ and non-human primates ⁹⁵. That CD40 bearing APC within the CNS, such as infiltrated macrophages as well as perivascular and parenchymal glia cells, contribute significantly to the pathogenesis of EAE has been elegantly shown
20 in bone marrow chimeric mice ⁸³.

The results of 't Hart et al, Laman et al, and Boon et al. ^{76, 79, 81} indicate that antibody blockade of CD40 is a potentially effective treatment of MS. Importantly, the ch5D12 Mab has no apparent side effects in the marmoset
25 EAE model nor in other primate species ⁸⁰. The beneficial clinical effect of anti-CD40 antibody was demonstrated in placebo-controlled experiments in two EAE models in marmosets, namely induced with human myelin ⁷⁹ or rhMOG ⁸¹. In addition, 't Hart et al have shown an inhibitory effect of anti-CD40 antibody treatment on already existing lesions ⁷⁶.

30

Psoriasis is an inflammatory skin disease afflicting 1-2% of the population. In this disease, T cells and keratinocytes in the lesions are activated and express activation markers and co-stimulatory molecules. It is thought that some co-stimulatory molecules expressed on keratinocytes and T cells interact with each other and that these interactions contribute to disease activity¹⁰⁸⁻¹¹⁰. One such set of molecules may be CD40, which is expressed on activated keratinocytes, and CD154 (CD40 ligand), which is transiently expressed on activated CD4⁺ T cells. CD40-CD154 ligation between T cells and keratinocytes may release from these cells inflammatory mediators that are seen in abundance in psoriatic lesions. CD40, CD154, and CD40-CD154 interactions have recently been reviewed¹¹¹.

Cultured keratinocytes also express CD40; expression is enhanced by IFN- γ treatment. Ligation of highly expressed CD40 on IFN- γ treated keratinocytes (referred to as CD40⁺⁺ keratinocytes throughout this paper) with CD154 results in up-regulation of ICAM and increased production of cytokines¹¹²⁻¹¹⁴.

So far, only one report has appeared that implicates this ligation in the pathogenesis of psoriasis¹¹³. The cell types that express CD40 in psoriatic lesions were not identified in the latter study and their incidence and lesional status were not described. The presence of CD154⁺ T cells in psoriatic lesions was also not studied. Thus, it remains unknown whether CD154 acts as one of the signals for production by keratinocytes of chemokines and complement, which are found in abundance in psoriatic lesions¹¹⁵.

Pasch et al recently demonstrated the presence and localization of CD40⁺ and CD154⁺ cells by immunohistochemistry in lesional and non-lesional skin from ten psoriasis patients¹¹⁶. Increased positivity for CD40 was observed on clusters of keratinocytes, and high expression of CD40 was shown on almost all CD1a⁺ Langerhans cells in normal, non-lesional, and lesional epidermis. In addition, high expression of CD40 was present on almost all CD83⁺ cells in

psoriatic lesions; they were rarely seen in non-lesional and normal skin¹¹⁶. Also, a small proportion of T cells showed CD154 expression, in most patients in juxtaposition to CD40+ cells. These results raised the possibility that CD154+ T cells may ligate with CD40+ keratinocytes, Langerhans cells, and CD83+ dendritic cells, and release mediators from them in the lesion. In addition, they demonstrated that CD40 ligation induces the release of chemokines (IL-8, RANTES, and MCP-1)¹¹⁶. In the same publication, Pasch et al. showed that the CD40-related release of the chemokines IL-8, MCP-1 and to a lesser extent of RANTES was inhibited by the antagonistic anti-CD40 antibody 5D12¹¹⁶. These data suggest that antagonist anti-CD40 mAb 5D12 may at least in part have an effect on the inflammation seen in psoriatic lesions.

US2003/0165499 discloses a measurable anti-psoriatic effect of the 5D12 and other antagonistic anti-CD40 antibodies in a SCID mouse xenogeneic transplant model system that is used as a model for psoriasis treatment, showing that antagonistic anti-CD40 antibodies may be used for the treatment of psoriasis. A therapeutic effect of 5D12 in this in vivo system was demonstrated

In a particularly preferred embodiment said autoimmune and/or an inflammatory disorder comprises inflammatory bowel disease. In another preferred embodiment a use of the invention is provided wherein said inflammatory bowel disease comprises ulcerative colitis (UC) or Crohn's disease (CD).

25

As illustrated above, a polypeptide according to the invention is suited for treatment of diverse inflammatory disorders, including autoimmune disorders and graft rejection. Therefore, the invention in one embodiment provides a method for ameliorating a symptom of an autoimmune disorder and/or an inflammatory disorder and/or for reducing graft rejection. In a

30

further embodiment the invention provides use of a polypeptide according to the invention, a binding body according to the invention, an antibody according to the invention, a nucleic acid according to the invention and/or a cell according to the invention for the manufacture of a medicament for

5 ameliorating a symptom of an autoimmune disorder and/or an inflammatory disorder and/or for reducing graft rejection. Ameliorating a symptom as used in the invention is defined as ameliorating at least one symptom of a disorder at least partially. The invention is of specific interest for autoimmune and/or inflammatory disorders for which at present no efficient treatment exists.

10 Examples of these disorders are rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, psoriasis, inflammatory bowel disease, bullous pemphigoides and atopic dermatitis. A polypeptide according to the invention, optionally comprised in a binding body or a cell of the invention and/or encoded by a nucleic acid of the invention, is specifically suited for

15 ameliorating a symptom of an autoimmune and/or inflammatory disorder as defined properties of a polypeptide according to the invention allow for interfering with a CD40-CD40L pathway in a specific manner. Furthermore, since a binding body according to the invention is preferably deimmunized, a binding body according to the invention is present for a sustained period of

20 time and thus shows its antagonizing activity in a patient for a considerable period of time. Therefore, the invention in one embodiment provides a use of a polypeptide according to the invention, a binding body according to the invention, an antibody according to the invention, a nucleic acid according to the invention and/or a cell according to the invention for the manufacture of a

25 medicament for ameliorating a symptom of an autoimmune disorder and/or an inflammatory disorder and/or for reducing graft rejection, wherein said autoimmune and/or said inflammatory disorder is selected from the group of rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, psoriasis, inflammatory bowel disease, bullous pemphigoides and atopic

30 dermatitis.

Brief description of the drawings

Figure 1. ch5D12 serum concentration after single administration at day 0 at the four dose levels as indicated by the symbols. Values are given as μg ch5D12/mL serum and were determined by enzyme-linked immunosorbent assay as detailed in Ref²⁷.

Figure 2. Changes in Crohn's Disease Activity Index (CDAI) scores during the 28-day period after the ch5D12 infusion. The shaded area indicates the period that ch5D12 levels exceed the $10 \mu\text{g/mL}$ concentration (see also Figure 1) which was found to be a functional antagonizing serum level in primates.²⁴⁻²⁷ In the 0.3 mg/kg cohort, serum levels never exceeded the $10 \mu\text{g/mL}$ level.

Figure 3. Histological disease activity scores at day 0 and day 28. Maximum activity score is 16 and activity score was performed according to Ref.⁴⁰ Samples from the ileum (diamonds) from nine subjects and samples from the colon (squares) from 11 subjects were obtained at day 0 and day 28 and results are presented per dose level of ch5D12 [(a) 0.3 mg/kg; (b), 1.0 mg/kg; (c), 3.0 mg/kg and (d), 10.0 mg/kg].

Figure 4. Colon biopsies from patient 012 (3.0 mg/kg) and ileum biopsies from patient 011 (3.0 mg/kg) are shown as examples for the reduction of the inflammatory response. Samples were stained with antibodies recognizing all T lymphocytes (CD3), B cells (CD19), macrophages (CD68) and CD40(+) cells before (a, c) and on day 28 after ch5D12 administration (b, d). HE, haematoxylin-eosin.

Figure 5. Consensus DNA and deduced amino acid sequence of the murine 5D12 VH and VL region.

Figure 6. Amino acid comparison of murine V regions with veneered and deimmunised V regions of 5D12.

Figure 7. Alignment of the 12 tested amino acid variants (Q5E, K13A, E16Q, T17S, I29L, I37V, P45L, M48L, STS60NSA, T68S, S79F, and T108S) of deimmunised 5D12 VH compared with the parent murine sequence (ch5D12) and the fully deimmunised sequence (DI5D12).

Figure 8. FACS analysis using JY cells. PER.C6 cell supernatant of the transient expressed variants (Q5E, K13A, E16Q, T17S, I29L, I37V, P45L, M48L, STS60NSA, T68S, S79F, and T108S) of 5D12 was harvested after 48h. As control also supernatant was harvested of cells transfected with either ch5D12 or DI5D12 together with supernatant from mock (no plasmid) transfected cells. Binding of expressed antibodies was tested by FACS using JY cells together with anti-human FITC labelled secondary antibody (1/100 diluted). As FACS control JY cells were incubated with only the secondary FITC labelled antibody.

Figure 9. Alignment of the additional V-L-I variants on positions 29 and 37 (29I-37V, 29V-37I, 29I-37L, 29V-37V, 29L-37L, 29V-37L, 29L-37V) of deimmunised 5D12 VH compared with the parent chimeric sequence (ch5D12), the fully deimmunised sequence (DI5D12; 29I-37I) and PG102 (29L-37I)

Figure 10. Distribution of antibody productivity for GS-CHO cell lines producing the PG102 antibody and 'overgrown' 24-well plate cultures. Number of cell lines is plotted on the X-axis, antibody concentration ranges are plotted on the Y-axis (antibody concentration ranges are distributed in ascending groups starting at 0-25 ug/ml).

Figure 11. Growth and antibody accumulation profiles for cell lines A) L107 (DC1; top panel), B) L25 (DC2; middle panel) and C) M95 (DC3; lower panel) in CDACF fed-batch shake flask culture.

- 5 Figure 12. Comparison of the binding of PG102(red) and ch5D12 (PG100)(blue) at 10µg/mL to CD40 immobilised at 3 different levels.

Figure 13. Inhibition of CD40 mAb binding to JY cells expressing human CD40 by the anti-idiotypic mAb 173-36-1. Anti-CD40 mAbs were assayed at a
10 concentration of 1 µg/ml. Data indicate mean ± s.e. mean of 4 separate determinations for each antibody.

Figure 14. CD40-Fc ELISA. Concentration-related binding of ch5D12 (PG100) and PG102 to human CD40-Fc. Plates were coated overnight with 250 ng/ml
15 human CD40-Fc. The isotype control mAb, chFUN-1, is directed against human CD86. Data indicate mean ± s.e. mean (all n=3).

Figure 15. FACS quantification of anti-CD40 mAb binding to JY cells. ch5D12 (PG100) and PG102 displayed comparable half-maximal binding
20 concentrations. The anti-CD86 mAb, chFUN-1, also showed binding to JY cells in these experiments due to the surface expression of human CD86 by this cell line. Data indicate the mean of duplicate determinations for each antibody.

Figure 16. Inhibition of ch5D12 (PG100)-PE and PG102-PE binding to JY cells
25 by unlabelled ch5D12 (PG100) and PG102 antibodies. Labelled ch5D12 (PG100) (A) and PG102 (B) were incubated at 1 µg/ml in the presence of increasing concentration of competing unlabelled antibody. Labelled antibody binding was determined as mean fluorescence intensity (MFI) by flow cytometry. Maximal MFI for ch5D12 (PG100)PE and PG102PE binding was

369 fluorescence units (A) and 305 fluorescence units (B), respectively. Data indicate mean $\pm$ s.e. mean of 4 separate experiments.

Figure 17. Inhibition of IL-8 release by THP-1 cells after co-culture with
5 Jurkat cells. THP-1 cells, pulsed with IFN γ , are co-cultivated with
Jurkat cells in the presence of increasing concentrations ch5D12 (PG100) or
PG102. The IL-8 release from THP-1 cells, induced by binding of Jurkat cells
to THP-1 cells via CD40-CD40L, was measured by ELISA. Data indicate the
result from a single experiment

10

Figure 18. Nucleic acid and amino acid sequence of an IgG₄ PG102 heavy and
light chain.

Examples

Example 1

5 *Materials and methods*

Eighteen adult subjects (18-60 years of age) with a clinical diagnosis of Crohn's disease confirmed by radiological, endoscopic or histological evidence, and with a Crohn's Disease Activity Index (CDAI) score of at least 220 but not more
10 than 450 (scored over 7 days prior to study drug administration) were selected for study inclusion. Subjects were allowed to have the following treatments before and during the study: mesalamine treatment for 8 or more weeks with the dose remaining stable for the 4 weeks before screening; a maximum of 30 mg of corticosteroids per day (or 9 mg budesonide per day) for 8 or more weeks,
15 with the dose remaining stable 2 weeks before screening; mercaptopurine or azathioprine for 4 or more months with the dose remaining stable for 8 weeks before screening. Subjects could not have received treatment with cyclosporin A or methotrexate within 3 months before screening nor were they allowed to have prior exposure to treatment with Mabs. The average age of all subjects
20 enrolled was 35.8 years of whom seven were males and 11 were females. All subjects were of Caucasian ethnicity. No apparent differences between patients in the different cohorts at inclusion were observed (Table 1), with the exception that in the lower three dose groups the majority of the subjects were females, while the highest dose group (10.0 mg/kg) consisted of only male
25 subjects. There were no notable differences between the four cohorts with regard to baseline electrocardiogram (ECG), vital signs, physical examinations, history of Crohn's disease signs and symptoms, and baseline laboratory values. There were no significant differences regarding baseline characteristics amongst the four dose cohorts. The highest dose cohort however also showed
30 the highest CDAI score at baseline. This study was approved by the Medical

Ethical Boards of The University Hospitals Leuven, Belgium, Leiden University Medical Center, The Netherlands, Medizinische Klinik, Kiel, Germany, and Hadassah Medical Center, Jerusalem, Israel. All patients gave informed consent to the study.

5

Study design and treatment protocol

ch5D12 is a molecularly engineered human IgG4 antibody containing the variable domains of the heavy and light chains of Mab 5D12 parent version.

- 10 This Mab has been shown to bind to CD40-bearing cells and to antagonize CD40 mediated activation of various cells²⁰⁻²⁷. ch5D12 was administered in an open label, single dose, multi-centre trial, studying four dose levels. There were five subjects in each treatment group except for the final dose cohort in which only three subjects were enrolled. Single administration at dose levels of
- 15 0.3, 1.0, 3.0, and 10.0 mg/kg of ch5D12 was done intravenously. After completing recruitment into one dose group, recruitment into the next group was only started when safety had been established at the current dose level. Clinical disease activity was assessed at each visit weekly for the first 28-day period and a subsequent last visit at day 56. Two subjects (3.0 mg/kg dose
- 20 cohort) withdrew from the study after the day 28 assessments but their data are included in the evaluation. Subjects were to remain stable on the same dose of their current medication during the course of the study. No un-allowed concomitant medication was used during the study. Response to ch5D12 treatment was defined as a decrease in CDAI of ≥ 100 points and clinical
- 25 disease remission was defined as a CDAI of ≤ 150 (total score) and a decrease of CDAI of at least 100. All subjects who underwent an endoscopy at screening and at day 28 ($n = 11$) were analysed for a decrease in their Crohn's Disease Endoscopic Index of Severity (CDEIS) scoring index. Biopsies were taken from these 11 patients prior to and at day 28 after the ch5D12 administration for
- 30 histopathology and immunohistochemistry. Safety evaluations included

physical examinations, vital signs, ECG, and laboratory data (chemistry, haematology and urinalysis) including, anti-dsDNA, pANCA, and human anti-chimeric antibody (HACA) evaluations. HACA were determined by enzyme-linked immunosorbent assay (ELISA) as explained in Ref. (27).

5

Pharmacokinetics

Serum concentrations of ch5D12 were determined by ELISA as previously described²⁷. To determine coating of CD40 by the injected ch5D12, blood was
10 collected in heparin tubes and diluted twofold in PBS. Peripheral blood mononuclear cells (PBMC) were isolated by gradient centrifugation (Lymphoprep, Nycomed, Roskilde, Denmark) and 500 000 cells were stained with FITC-labelled ch5D12 and PerCP-labelled anti-CD20 (Becton-Dickinson, Mountain View, CA, USA) and incubated on ice for 30 min. As a background
15 control for 5D12-FITC binding, a separate tube of PBMC was stained only with PerCP-labelled anti-CD20 antibody. Unbound antibodies were washed away, and cells were analysed using a flow cytometer (FACSort; Becton-Dickinson). Acquisition of events was conducted by gating and acquiring CD20-expressing cells for a total of 5000 events. If the number of CD20-expressing cells was
20 lower for a sample set, care was taken to acquire the same number of events in both the unstained and stained preparation.

Histology and immunohistochemistry

25 Mucosal biopsies from ileum and colon obtained during ileocolonoscopy prior to treatment on day 0 and on day 28 with a standard forceps were fixed in formalin 6% for routine analysis. Additional samples were immediately snap-frozen in Tissue-Tek optimum cutting temperature compound (Miles Laboratories Inc, Naperville, IL, USA) in liquid nitrogen cooled isopentane.
30 Samples were stored at -80 °C until further use. Formalin fixed samples were

routinely paraffin processed. Five-micrometer thick sections were prepared and stained with haematoxylin and eosin. The sections were analysed using a Leitz Wetzlar microscope (Wetzler, Germany). A total of four semi-serial sections for each sample were analysed. The frozen samples were used for immunohistochemical analysis, which was performed using a panel of Mabs to assess the presence of different subsets of lymphocytes and monocytes/macrophages. The panel was completed with Mabs directed against CD40 and CD40L. Immunohistochemical staining was performed on cryostat sections, dried overnight at room temperature, and fixed in absolute acetone for 10 min. Re-hydrated slides were incubated for 30 min with the following Mabs; CD3 (clone: UCHT1, 1/10 dilution) (Dako, Glostrup, Denmark), CD4 (clone: MT310, 1/10 dilution) (Dako), CD8 (clone: 144B, 1/20 dilution) (Dako), CD19 (clone: HD37, 1/30 dilution) (Dako), CD40 (clone: 5D12, 1/100 dilution) (PanGenetics, Amsterdam, The Netherlands), CD68 (clone: Kp1, 1/50 dilution) (Dako), CD154 (clone: M90, 1/10 dilution) (Serotec, Oxford, UK). The secondary Mab was a biotin-labelled anti-mouse immunoglobulin (1/400 dilution; Dako) applied for 30 min. To efficiently block endogenous peroxidase, sections were also incubated in methanol solution containing 0.3% (v/v) H₂O₂ for 20 min. After three washes with PBS, the avidin/biotin peroxidase-labelled complex (Dako) was added. In between the incubations, the sections were washed in phosphate buffered saline at pH 7 for 15 min. The reaction product was visualized by incubating the section for 10 min in 0.05 M acetate buffer at pH 4.9, containing 0.05% 3-amino-9-ethyl-carbazole (Janssen, Beerse, Belgium) and 0.01% H₂O₂, resulting in a bright-red immunoreactive site. Afterwards, the slides were faintly counterstained with Harris haematoxylin, rinsed with distilled water and a cover-slip was applied with glycerol. Controls, which were negative, consisted of an omission of a primary or secondary antibody, use of chromogen alone, and use of irrelevant isotype-matched mouse IgG (vimentin; Dako).

Paraffin sections stained with haematoxylin and eosin, were all analysed by the same pathologist (K. G.) blinded for the origin of the samples. Disease activity was assessed using a histological numerical score⁴⁰ of which the maximum activity corresponded to 16. The immunohistochemically stained sections were also analysed in a blinded fashion. For 15 sets of biopsies, positively staining cells were counted in high power fields (magnification $\times 40$). The fields were chosen according to the highest density of stromal cells. The number was expressed as a percentage of the total number of mononuclear cells in the stroma. These counts were used for the construction of a scoring system whereby the intensity of the positively staining cells was subdivided into four categories: -, +/-, + and ++. The normal value for CD3+, CD4+, CD8+, CD19+ and CD68+ for both the ileum and colon is +. The normal value for CD40+ and CD154+ cells is negative (-).

15 *Immunologic safety parameters*

To assess any non-specific immunosuppression by ch5D12, proliferation of peripheral blood T cells in response to the phytohemagglutinin mitogen (PHA) was evaluated. PHA stimulation (1 $\mu\text{g/mL}$) was performed on isolated PBMC as previously reported.⁴¹ In addition, percentages of circulating CD3+, CD4+, CD8+ and CD20+ cells were measured to exclude depletion of circulating cells. For flow cytometry, blood was collected at indicated times in heparin tubes, of which 100 μL whole blood in each test-tube was stained as follows (all Mabs were purchased from Becton-Dickinson): anti-CD3-FITC, anti-CD19-PE, anti-CD45-PerCP, anti-CD45-FITC, anti-CD4-PE, anti-CD8-PE and anti-CD14-PE. The stained cells were then analysed in a FACScan flow cytometer (Becton-Dickinson). This protocol was conducted to calculate the ratio of CD4+ to CD8+ cells and to determine the proportion of B cells within the lymphocytic population. The percentages reported for CD3 and CD19 were percentages of T cells and B cells within the lymphocytic population, whereas, CD4 and CD8

cells were reported as percentages of CD3 cells. Only CD14 (monocytic population) is shown as a percentage of the total CD45 population.

Results

5

Pharmacokinetics

The mean peak levels of ch5D12 after a single i.v. injection were dose-dependent and dose-proportional (Figure 1). After 24 h, approximately 50% of the administrated ch5D12 could be detected in serum in the highest three dose cohorts. For the 0.3 mg/kg cohort, only 15% was present in serum after 24 h. From non-human primate studies, a $t_{1/2\beta}$ of 8-10 days has been calculated²⁷ which was confirmed by the current human data. Complete coating of CD40 on peripheral blood B cells was not achieved by the infusion of 0.3 mg/kg as determined by an ex vivo competition assay using FITC-labelled ch5D12 (not shown). For the 1.0 mg/kg group, approximately 1 week of coating of CD40 on circulating B cells could be observed, while in the two highest dose groups this period increased to 2-3 weeks. This indicates that in the lowest dose group, antagonism of CD40 was incomplete, while complete antagonism was achieved in the 1.0 mg/kg cohort for 1 week and even longer in the higher dose groups.

15
20

Clinical responses

The analysis of CDAI evolution indicates that 13 of 18 subjects (72%) experienced a favourable response after the ch5D12 infusion. Similarly, four of 18 subjects (22%) experienced a remission during that period. When evaluated at day 21, a favourable response was recorded in 10 of 18 subjects. Cohorts 2 and 4 showed the largest mean decrease in CDAI, and no clear dose-effect relations were observed (Figure 2; Table 2). A post hoc repeated measure anova showed a statistically significant decrease in CDAI over the 56-day

25
30

observation period ($P < 0.001$). The difference between the cohorts was not statistically significant. Only in cohorts 3 and 4, CDEIS could be evaluated (as endoscopy was performed in only three subjects in cohort 1 and 2). In cohort 3, two of five subjects had a decrease in CDEIS, whereas in cohort 4, two of three
5 subjects had a decrease. Remaining subjects in these two cohorts showed no change (not shown).

Histological changes

10 In the patients from whom biopsies were taken ($n = 11$), a single investigator performed evaluation of changes in histopathology and immunohistochemistry. The assessments performed on biopsy material show a clear influence of treatment upon microscopic activity of the disease as well as upon the intensity of the lamina propria mononuclear cell infiltrate.

15 The change from the baseline to day 28 histopathology activity score is shown in Figure 3. Samples from the ileum of nine patients and samples from the colon of 11 patients at day 0 were available. In four cases, the ileum was not involved at screening while five patients had active ileal disease. The mean
20 histology score at day 0 for those with active disease was 4.6 (range 3-7). The mean histology score for the five patients decreased to 1.0 (range 0-3) at day 28. In four of 11 patients, the colon was not involved at day 0 and the colonic samples from these patients remained normal at day 28. In seven patients the colon was clearly involved with a mean histology score at day 0 of 4.5 (range 2-
25 12). This decreased at day 28 to a mean score of 1.7 (range 0-7). In five of seven patients, the score at day 28 was either 0 indicating a normal biopsy or 1 indicating only architectural abnormalities. In the lowest dose cohort (panel a), the colonic score of one subject decreased, but the other colonic scores in this group remained high and the ileum score even increased. In the higher dose
30 cohorts (panels b-d), either the ileum or colon or both decreased in the activity

score at day 28 after treatment with ch5D12. Conclusively, a positive response [as defined in Ref. (41)] was observed in 81% (9/11) of subjects. In addition, seven subjects with active disease at day 0 showed reduced neutrophil activity to the point that neutrophils were not present in their biopsies at day 28.

5

To assess which cells were targeted by treatment with ch5D12, an immuno-histochemical evaluation was performed per protocol on the available biopsies using T-cell (CD3, CD4 and CD8) and B-cell (CD19) markers and a macrophage marker (CD68). Results are listed in Table 3. No reduction was observed at day 28 in the lowest dose cohort while in all other cohorts decreased infiltration was observed. In the samples from the ileum at baseline, there was an increased CD3+ T cell number in six of nine cases. In the colon, an increased CD3+ T cell number was present in four of 11 patients, all of whom also had active disease on routine histology. At day 28, the CD3+ T-cell numbers had normalized in all but one patient. In this patient (patient 001; lowest dose group; 0.3 mg/kg) there was a persistent CD3+ cell increase in the ileum and colon, in parallel with persistent inflammatory activity and granulomatous inflammation on routine histology. The CD4+ and CD8+ cells followed a similar pattern as CD3+.

20

The pattern for the CD19+ B cells was comparable with that observed for the T cells. At day 0, there was an elevation of B cells in six of nine patients of whom ileal biopsies were available and in four of 11 samples from the colon. At day 28, the amount of the B cells had normalized in all cases except for the one patient from the low-dose group.

25

A similar decrease on day 28 was observed for CD68+ monocyte/macrophage cells. At day 0, there was an increased amount of CD68+ monocyte/macrophage cells in five of nine patients of whom ileal biopsies were available and in four of 11 samples from the colon. Again all increases had

30

normalized on day 28, except for one low-dose cohort patient that remained high for CD68+ monocyte/macrophage cells in the colonic sample and even an increase in the ileac biopsy, showing this patient had still active disease at day 28.

5

Figure 4 shows representative examples of the decrease of all three major cell types (T cells, B cells and monocyte/macrophage cells) in ileum biopsies of patient 011 (activity score: 7) and colonic biopsies of patient 012 (activity score: 12) before ch5D12 administration and at day 28 after ch5D12 administration.

10

Safety

No subjects withdrew from the study because of adverse events (AEs) with the majority of these AEs being mild to moderate in severity. An evaluation of AEs occurring at $\geq 5\%$ of the total number of AEs revealed the most common AEs being pyrexia, arthralgia, myalgia and headache. Arthralgia, occurring frequently both at baseline and during the follow-up period, decreased in frequency over the course of the follow-up period. Flu-like symptoms and pruritic rashes occurred in four subjects, of whom only one subject experienced a moderate rash starting shortly after study drug administration (3.0 mg/kg). Overall, there is no clear dose response for any single AE, with the possible exception of headache/migraine.

There were six serious adverse events (SAEs) noted in six subjects; no events occurred in the 0.3 mg/kg cohort, three events in the 1.0 mg/kg cohort, two events in the 3.0 mg/kg cohort and one event in the 10.0 mg/kg cohort. Five of the six SAEs were considered possibly study drug related, one event was considered not related. All but one event were gastrointestinal events (two sub-ileus both in 1.0 mg/kg; two Crohn's disease aggravated, 1.0 and 3.0 mg/kg subjects; and one abdominal pain aggravated in a 3.0 mg/kg subject). Two of

these events however, occurred at such a time-point that a causal relationship to the study drug is very unlikely (one subject on the day of infusion and another subject after undergoing endoscopy 29 days after study drug administration). None of the observed abnormalities appear study drug or dose related.

Immunological safety parameters

As ch5D12 inhibits B-cell activation in vitro, total immunoglobulin levels were determined and found to be stable over the course of the follow-up period. To exclude the possibility that ch5D12 is able to deplete B cells from the circulation we performed subset analysis of PBMC over the course of the study. Percentages and absolute numbers of CD3, CD4, CD8 and CD14 positive cells were not affected by ch5D12 administration. Absolute counts of circulating CD19+ B cells transiently decreased shortly after ch5D12 administration and recovered to their original values after ch5D12 had been cleared from the circulation (not shown).

To study whether treatment with ch5D12 would cause a general immune suppressive effect, PBMC isolated at various time-points prior and after treatment were stimulated by the polyclonal T-cell activator PHA. Unfortunately, cells from the 10.0 mg/kg dose cohort were not frozen appropriately and were therefore unable to respond to PHA, including the cells obtained before ch5D12 administration. The stimulation index induced by incubation with PHA was highly variable but no trend towards a decrease after ch5D12 treatment was detected (data not shown).

Anti-ch5D12 antibodies were not detectable in the two lower dose cohorts throughout the entire study. In the 3.0 mg/kg cohort, one subject was found to have mounted an anti-ch5D12 response, although the signal was very low and

could only be detected in the undiluted day 28 sample of this patient. In the highest dose group, another subject had anti-ch5D12 antibodies on day 28 and day 56 of the study. These could be detected with a 1:64 dilution. Therefore, only two subjects had a possible immune response to ch5D12 administration.

5

Discussion

In this open label, multi-centre trial, four dose levels of ch5D12 were administrated to subjects with moderate to severe Crohn's disease. Eighteen
10 subjects were enrolled, with five subjects in each treatment group except for the final dose cohort in which only three subjects were enrolled. ch5D12 serum levels increased dose-dependently and the long serum half-life that was observed in cynomolgus monkeys²⁷ was confirmed in humans. This study shows the feasibility of antagonizing CD40-mediated cell activation using
15 ch5D12, in the absence of major side effects. The CDAI analysis indicates that 72% of subjects experienced a clinical response and 22% experienced a remission during the study. These results are promising; however the open-label design and the absence of a control group make it impossible to definitely attribute the observed changes to administration of the study drug. However,
20 we do have several arguments in favour of anti-inflammatory effect of ch5D12. First, the observed decrease in CDEIS amongst subjects in the two highest dose cohorts supports treatment efficacy. Secondly, the results of the histological and immunohistochemistry analyses done on biopsies, show that ch5D12 reduces inflammatory activity in colon or ileum. In the low-dose cohort
25 no clear effect was observed. Subjects in the 10.0 mg/kg cohort had the lowest histological activity scores at day 0 and the reduction in activity score is therefore limited in this cohort. The highest scores at baseline were found in the 1.0 mg/kg and the 3.0 mg/kg groups and in these patients there was a clear reduction in histological scores at day 28.

30

Although two patients were identified having anti-ch5D12 antibodies in the current study, the levels of HACA antibodies remained very low and were only detectable at very low serum dilutions. It remains to be established whether anti-ch5D12 responses will be detected in subsequent studies.

5

This trial represents the first administration of anti-CD40 Mab in general and of ch5D12 in particular to humans. However, evidence of biological activity of Mab 5D12 has previously been obtained in vivo in the experimental autoimmune encephalomyelitis (EAE) model in marmoset monkeys.

- 10 Administration of 5D12 starting 2-4 weeks after induction of disease had clear beneficial effects²³. This in vivo proof of concept initiated recombinant humanization and further development of 5D12. The biological activity of ch5D12 was also evaluated in the marmoset EAE model. At day 50 after MOG-immunization, disease incidence in the placebo group was 100% (50% of the
- 15 animals were killed because of their severity of EAE), while no disease signs were observed in the ch5D12-treated group²⁴. Furthermore ch5D12 has shown functional activity in a rhesus monkey renal transplantation model²⁵ and when used as immune suppressive strategy to prevent immune responses against adenoviral particles and their products²⁶. A tissue cross-reactivity study on
- 20 human and cynomolgus tissues showed that ch5D12 bound to the cell surface of B cells and DCs in lymphoid organs. No unexpected cross-reactivity was observed on either human or cynomolgus tissues. A safety and tolerability evaluation for ch5D12 was performed in cynomolgus monkeys, in which weekly administration of ch5D12 for 4 weeks was shown to be safe and without any
- 25 side-effects in all monkeys. In this study, functional evidence was obtained that ch5D12 can prevent B-cell activation and proliferation²⁷. Collectively these studies showed that antagonist anti-human CD40 Mab ch5D12 has no unexpected cross-reactivity, is safe, limits immune reactivity in cynomolgus monkeys, and inhibits a prototype chronic inflammatory disease of the central

nervous system in marmoset monkeys. Hence, the preclinical assessment supported further development for use in patients.

The CD40-CD154 (CD40L) co-stimulatory pathway has been validated in animal models as a promising clinical target for treatment of autoimmune disease and transplantation rejection.¹⁵ Potentially the CD40-CD154 interaction can be inhibited by targeting either CD40 or CD154. All previous studies have focused on antagonizing CD154, which is selectively expressed on activated T cells and platelets.⁸⁻¹¹ Human clinical studies using an anti-human CD154 Mab have been halted because of thromboembolic events¹²⁻¹⁴. As this anti-CD154 Mab is constructed as a humanized IgG1, which binds extremely well to Fc-receptors, the anti-CD154 Mab may cross-link CD154 to Fc-receptors resulting in the formation of blood clots. Recently, CD40 was reported to be constitutively expressed on platelets and found to be functionally important using soluble CD154 as stimulus for platelet activation. These stimulatory effects that resulted in activation of platelets were abrogated by addition of anti-CD154 or anti-CD40 Mabs⁴². We confirmed the non-stimulatory characteristics of ch5D12 also in experiments with human platelets, as addition of ch5D12 to prestimulated platelets using sub-optimal concentration of ADP or collagen was without any effect on the activation status of the platelets (M.F. Hoylaerts, unpublished observations, Leuven, Belgium). Furthermore, the IgG₄ Fc part of the ch5D12 Mab backbone has reduced capability to bind to Fc-receptors expressed on platelets as compared with an IgG₁ backbone. Further discrimination between the two approaches comes from the fact that ch5D12 targets B cells, monocytes, macrophages and DC, while an anti-CD154 Mab targets mainly activated T cells, resulting in markedly different clinical strategies. No indications of thromboembolism in any of the patients treated in this study were observed.

The effect of ch5D12 is largely based on the interference with CD40 mediated cell activation. CD40 triggering induces cytokine and chemokine secretion by macrophages and DC, and enhances the antigen presenting capacity of the latter. As ch5D12 antagonizes the CD40 receptor through which TNF- α production is induced³⁹, it might also antagonize TNF- α secretion which undoubtedly is an important cytokine in Crohn's disease⁴³. However, also a number of other inflammatory cytokines, such as IL-8 and IL-12³⁹, are induced by CD40 triggering and therefore ch5D12 might have anti-inflammatory properties in addition to the prevention of TNF- α production. Whether the effects observed depend mainly on a reduced T-cell activation, or a reduced release of cytokines and chemokines, awaits further investigation.

This first-into-man study with ch5D12 shows promising clinical benefit in the absence of serious side-effects. Further clinical studies will be needed to establish the role and optimal dosing scheme for ch5D12 in maintenance of remissions and in the safety monitoring of multiple infusions. Obviously, therapeutic benefit by antagonizing CD40 will not be limited to Crohn's disease and can potentially be extended to many other inflammatory disorders.

20 Example 2

Material and Methods

Generation of chimeric version of 5D12

Starting from the 5D12 hybridoma cell line the murine variable regions of 5D12 were cloned. Briefly, from lysated cells using Qiagen total RNA was isolated. By RT-PCR using the oligo's 5' CAG GTS MAR CTS SAG SAG TC W GG 3' and 5' GCA TGT ACT AGT AAT TTT TVT TGT CCA CYT TGG TGC T 3' to amplify the VH region and using the oligo's 5'CGA TAC GAS MTY CAG CTG ACC CAG TCT CCA 3' and 5'GAC TCA TCT AGA TAC ACT CAT TCC

TGT TGA AGC TCT TG 3' to amplify the VL region of mu5D12. After sequence analysis the consensus sequences for the V regions were determined.

- Subsequently the V-regions were recloned in pcDNA3002 (Marissen et al, J. of Virol, 2005: 79:4672-4678), together with genomic sequences encoding human
- 5 IgG₄ and human kappa to construct a chimeric IgG₄ immunoglobulin expression plasmid.

Generation of deimmunized version of 5D12

In collaboration with Biovation using their proprietary DeImmunisation technology within the V regions potential T-cell epitopes were identified.

- 10 Helper T cell epitopes comprise short amino acid sequences within proteins with the capacity to bind to MHC class II molecules. By removal of the T cell epitopes the antibody can no longer trigger T cell help and the subsequent immunogenicity against it. Identification was done using Peptide Threading. The Peptide Threading approach is a computer-based method for predicting
- 15 the binding of peptides (small amino acid sequences) to 18 different human MHC class II molecules based on known structures solved by X-ray crystallography. Furthermore an analysis was done to search the V regions for the presence of known human T cell binding peptides from a database (www.wehil.wehi.edu.au).
- 20 Based on the primary murine V region amino acid sequences, surface humanized (veneered) V regions were designed. Within these veneered VH sequence 8 potential T-cell epitopes and within the veneered VL region 4 T-cell epitopes were identified. Identified T-cell epitopes were removed by amino acid substitution. Based on this analysis the DNA sequences encoding the
- 25 deimmunized VH and VL region were synthesized and recloned in the pcDNA3002-hIgG₄-kappa expression plasmid.

Site-Directed Mutagenesis

To restore the murine sequences within the VH region of deimmunized pcDNA3002 expression plasmid the QuickChange Site-Directed Mutagenesis kit of Stratagene was used according to the manufacturer's protocol. In table 4 the used oligonucleotides are shown. Briefly, the plasmid was denatured followed by annealing of the sense and anti-sense primers containing the desired mutation. By cycling (30 sec 95°C; 30 sec 55°C; 600 sec 68°C) for 16 times using a Biometra T gradient PCR machine the mutagenic sequences were incorporated. The non-mutated parental plasmid was digested using DpnI, which only cuts methylated DNA. Subsequently the PCR mixture containing the mutated plasmid was transformed in XL1-Blue competent cells. Colonies were picked and analyzed by DNA sequences for the correct plasmid. A second set of experiments focused on generating additional variants of positions 29 and 37 specifically by substitution of I for structurally related V or L. In a similar way as described above, plasmids were generated using the oligonucleotides as shown in table 6. Two mutagenesis rounds were required to create double mutants (Table 7). Plasmids were used to transform XL1-blue competent cells.

Expression in PER.C6TM

20

Expression of the various versions of 5D12 was done transiently in PER.C6TM (Crucell). Briefly, the day before the transfection PER.C6TM were trypsinized and counted. Cells were plated in DMEM substituted with 10% FCS at a density of 4×10^5 cells per well in a T24 well plate. Next day medium was replaced with 0.5 ml fresh medium. For each well 1 µg of plasmid DNA was diluted into 50 µl of OPTI-MEM I, which was combined with equal volume OPTI-MEM I containing 3 µl LipofectAMINE 2000 (LF2000) reagent (Invitrogen). Mixture was incubated for 20 min at room-temperature to allow DNA-LF2000 reagent complexes to form. Subsequently the DNA-LF2000 reagent complexes (100 µl) were directly added to each well. The transfected

cells were incubated at 37°C in a CO₂ incubator. After 48h the supernatants were analyzed for antibody expression.

ELISA

5

The amount of expressed antibody was measured by a sandwich ELISA.

Briefly, 96 wells EIA/RIA plates (Corning 3590) were coated o/n at 4°C with polyclonal antibody anti-human IgG (Jackson 109-005-088) diluted 1/1000 in PBS (100 µl/well). After blocking with PBS containing 0.05% Tween and 1%

- 10 BSA, plates were 3x washes with PBS-0.05% Tween. Subsequently as standard purified chimeric 5D12 was applied in titration starting at 400 ng/ml to zero in triplo. Samples (transient supernatants) also applied in titration. All dilutions were done in PBS-0.05% Tween. Plates incubated for 1 hour at 37°C. After 3 times washing 1/5000 diluted detection antibody (anti-human kappa
- 15 alkaline phosphatase labelled (Southern Biotech Associates 2060-04)) was applied to each well. Plates incubated for 1 hour at 37°C. After removal of incubation plates were 5 times washed. Staining was done with PNP substrate (Sigma N-2765) (1 mg PNP/ml substrate buffer - 100 mM Tris/HCl, 100 mM NaCl, 5 mM MgCl₂·6H₂O pH 9.5 -). OD405 nm was measured on a BioRad 550
- 20 Titertek using 655 nm as reference.

FACS

- By FACS analysis the antigen specificity of the expressed antibodies was
- 25 analyzed. JY cells are EBV transformed human B-cells, which express CD40. Briefly, 100,000 JY cells were incubated with 100 µl supernatant from the transient expression experiment at 4°C for 20 minutes. To remove unbound antibody cells were washed with PBS containing 0.05% NaAzide and 1% BSA. As secondary antibody 1/100 diluted FITC labeled anti-human IgG (Jackson

109-095-127) was used. After 20 minutes incubation at 4°C cells were washed. Finally cell samples were measured on a FACScan (BD).

Results

5

After analysis of several clones the consensus sequence was determined for the murine V regions of 5D12 (figure 5).

10

By human surfacing the V regions were redesigned, retaining critical murine amino acids. By the Peptide Threading method and the database search potential T-cell epitopes within these redesigned sequences were determined. Within the VH region 8 (first base of potential epitope: position 27, 30, 43, 46, 57, 61, 77 and 83) and within the VL region 4 (position 7, 13, 28 and 86) potential epitopes were identified. Based on the T-cell epitope mapping and the veneering process the optimal deimmunized VH and VL amino acid sequences were defined (figure 6). To remove the T-cell epitope at position 61 in the VH region special attention was taken not to introduce a potential N-glycosylation site (N-X-T/S): NSA was substituted with STS. In this final design 12 positions in the VH region were changed (counting position 60-62 as 1 change). For the VL region 8 positions were changed.

15

From preliminary experiments comparing the expression levels of fully deimmunized 5D12 (DI5D12) compared to ch5D12 (renamed PG100) it was found that due to the introduced mutations DI5D12 was expressed at lower levels as compared to the parent sequence (data not shown).

25

In principal each substitution can influence the characteristics of the expressed antibody. Therefore to investigate the influence of each changed position in the VH region on the characteristics of the antibody 12 additional

variants were designed (figure 7). In each variant 1 position was changed back to the original murine amino acid sequence.

To test the expression levels and the functionality of the various versions 14
5 pcDNA3002 expression plasmids were constructed. Each plasmid contains the human IgG₄ backbone to which the VH regions were linked and the human kappa region to which the VL regions were linked. One plasmid encoded for ch5D12, in which the murine V regions were used. The other 13 plasmids are deimmunized versions, in which the deimmunized VL region is used in
10 combination with the deimmunized VH region or one of the 12 VH variants shown in figure 7. Subsequently PER.C6TM cells were transfected with the 14 plasmids using LipoFectamine2000. Also included were negative mock transfections (no plasmid). After 48 hour the transient supernatants from each transfection were harvested and analysed for antibody expression. In figure 8
15 the FACS data are shown. Clearly all tested constructs bind to JY cells, indicating that the CD40 binding specificity of 5D12 is preserved.

Furthermore the FACS data indicate that the produced antibody levels are different. Clearly there is a difference in expression level between DI5D12 and
20 ch5D12 (PG100). Variants Q5E, K13A, E16Q, T17S, STS60NSA and T68S show the same expression characteristics as DI5D12. The other variants seem to increase the expression levels. To exactly determine the various expression levels, all supernatants were analysed by quantitative ELISA measuring antibody levels. As shown in table 5, the tested variants can be divided in 3
25 groups: no or low improvement on the expression level (Q5E, K13A, E16Q, T17S and STS60NSA), variants P45L, M48L, S79F and T108S restore the expression levels as compared with ch5D12 (PG100) up to 40%, while the I29L and I37V variants restore the level above 50%. Especially the substitution at position 29 (I to L) seems to have large influence on the expression levels. The
30 I29L variant was subsequently renamed PG102 antibody.

The observation that substitution at position 29 (I to L) and 37 (I to V) both have large influence on the expression levels was among others unexpected because amino acids I, L and V are structurally closely related and the 3 together form the group of branched-chain amino acids. In a subsequent study
5 it was further established what the effect of combined substitutions of the amino acids I at positions 29 and 37 with structurally related amino acids V or L was.

Figure 9 shows the V-L-I variants that were designed in alignment with ch5D12 (PG100), DI5D12 and PG102 (I29L variant).

10

To test the expression levels and the functionality of these V-L-I variants, additional pcDNA3002 expression plasmids were constructed that contain the human IgG₄ backbone to which the VH region of one of the variants was linked and the human kappa region to which the VL regions were linked, as described
15 above. Subsequently PER.C6™ cells were transfected with the plasmids using LipoFectamine2000. After 48 hour the transient supernatants from each transfection were harvested and analysed for antibody expression. To exactly determine the various expression levels, all supernatants were analysed by quantitative ELISA measuring antibody levels. As shown in table 8, the
20 various combinations of a V, L or I on positions 29 and 37 are associated with variable production levels. This further demonstrates that not just all V, L and I substitutions ultimately lead to higher production levels.

25 Discussion

By reducing the immunogenicity of a therapeutic antibody the characteristics can change. In this study it is shown that the deimmunization design result in an antibody which still has binding capacity to its antigen, however the
30 antibody could only be produced at low levels. Most probably upon expression

there is a problem in the assembly of the deimmunized VH and VL region. By changing each substituted position back to the original murine sequence within the VH region, it was revealed that some positions were essential for a proper expression of the antibody. Especially the sites which were based on the peptide threading approach had large influence on the expression level. The N-terminal changes, all introduced based on veneering had little influence. Remarkably the substitution of I by L in position 29 had the largest influence, which was unexpected as both amino acids are structurally closely related. The I29L variant was subsequently renamed PG102 antibody.

Subsequent studies demonstrated that various combinations of a V, L or I at positions 29 and 37 led to variations in production levels, demonstrating that although the 3 branched-chain amino acids V, L and I are structurally closely related, substitution of each of these on positions 29 and 37 has an impact on production levels of the complete antibodies.

Example 3

Materials and Methods

Construction, selection and development of a GS-CHO cell line for the production of PG102 (I29L variant).

Lonza Biologics (Lonza) was requested by PanGenetics BV to undertake the construction, selection and evaluation of a GS-CHO cell line expressing the human recombinant IgG₄/kappa antibody PG102 (I29L variant).

Cell lines were constructed by transfecting CHOK1SV host cells with vectors generated using Lonza's Glutamine Synthetase (GS) Gene Expression System (WO2003054172). The gene sequences were supplied by PanGenetics. The cell line CHOK1SV is a suspension variant of the Chinese hamster ovary (CHO) cell line CHOK1, which has been adapted to chemically defined, animal

component-free (CDACF) medium (Bebbington et al (1192) Biotechnology 10:169-75; de la Cruz Edmonds MC et al (2006) Mol. Biotechnol. 34:179-90).

5 All reagents, media components and materials were obtained from accredited suppliers. The formulation of the feeds SF40 and SF41, medium CM42, and the medium supplement SPE were obtained from Lonza. The medium CD-CHO (Invitrogen) was also used with the addition of selective agent L-methionine sulfoximine (MSX) for routine subculture of CDACF adapted cell lines and with Phenol Red for transfection. The selective agent MSX was
10 supplied by Sigma. These media and feeds do not contain antibiotics. The construction of the GS expression vector encoding the genes for the PG102 was also performed at Lonza. The CHOK1SV host cells were obtained from a vial of Lonza's cryopreserved working cell bank 269-W.

15 *Cell culture, concentration and viability*

Cells were revived from vials of cryopreserved stocks by rapidly warming to 37 °C and diluting into approximately 50 mL of growth medium. The cryoprotectants were removed by centrifuging the cells, discarding the supernatant and resuspending the cells in fresh growth medium.
20 For static culture, cells were grown in T25 flasks with a culture volume of 5 to 15 mL. These static cultures were placed in an incubator at 35.5 to 37.0 °C containing an atmosphere of 10% v/v CO₂ in air. For suspension culture, cultures were expanded in a suitable volume of growth medium and subcultured every 4 days for growth in CDACF medium. The
25 containers used were: 125 mL shake flasks (for a culture volume of 10 to 30 mL), 250 mL shake flasks (for a culture volume of 30 to 50 mL), 500 mL shake flasks (for a culture volume of 50 to 100 mL), 1 L roller bottles (for a culture volume of 100 to 200 mL), or 2 L roller bottles (for a culture volume of 200 to 400 mL). The headspace of each culture was gassed with 5% v/v CO₂ in

air and the flasks sealed. The cultures were then incubated at 35.5 to 37.0 °C on a shaking platform rotating at 140 ± 5 rpm.

Total and viable cell concentrations were obtained by aseptically removing a sample of culture, staining the non-viable cells with Trypan Blue and

5 microscopic examination using a modified Fuchs-Rosenthal haemocytometer. When appropriate, samples were diluted prior to counting.

Alternatively, the total and viable cell concentrations were determined using a CASY-1 particle counter or a Vi-CELL XR automated cell counter. For the CASY-1 counter, a sample of culture was aseptically removed from each flask,

10 diluted with Casyton buffer and the number of viable and total cells determined. Viable and non-viable cells were distinguished by size.

Percentage viability was calculated as the ratio of viable cells to total cells multiplied by 100. For the Vi-CELL XR counter, 0.7 mL of cell culture was aseptically removed and an automated Trypan Blue count was performed. A

15 series of camera pictures and a computer analysis algorithm are used to identify and count the viable and non-viable cells. For all counting methods, percentage viability was calculated as the ratio of viable to total cells multiplied by 100.

20 *Transfection of CHOK1SV host cells*

On the day of transfection, cells growing in non-selective medium were harvested by centrifugation and resuspended at a concentration of 1.43×10^7 viable cells/mL. For each transfection, approximately 0.7 mL of the cell suspension and 40 µg of plasmid DNA were added to an electroporation

25 cuvette. The electroporation cuvette was then placed in the electroporation apparatus and a single pulse of 300 V, 900 µF was delivered. Following electroporation, the cells from the cuvette were distributed into 96-well plates at approximately 5000 host cells/well, using the medium CD-CHO/phenol red. The plates were incubated at 35.5 to 37.0 °C in an atmosphere of 10% v/v CO₂

30 in air. The day after transfection, 150 µL of the selective medium CD-

CHO/phenol red/66.6 μ M MSX, was added to each well. The final concentration of MSX in each well was 50 μ M. The plates were monitored to determine when the non-transfected cells died to leave foci of transfected cells. Foci of transfected cells became apparent approximately 3 to 4 weeks after
5 transfection. All the transfectants examined and progressed further came from wells containing only a single colony, as determined by visual assessment.

Assessment of productivity of cell lines in static culture

10 The 96-well plates were incubated for approximately 3 to 4 weeks, following transfection to allow colony formation. The colonies were examined microscopically to verify that they were of a suitable size (covering greater than 60% of the bottom of the well), and that only one colony was present in each well. The culture supernatant was then assayed for antibody production
15 using Lonza's ELISA assay for assembled antibody, described under "assembly ELISA". The percentage confluence of the well was assessed at the time of sampling. The value obtained by dividing the assay result by percentage confluence was used to rank the cell lines.

Cultures of high ranking cell lines were expanded into 24-well plates in the
20 medium CD-CHO/phenol red/25 μ M MSX and were left to reach confluence. On reaching confluence, this culture was used to inoculate a T25 flask in the medium CD-CHO/phenol red/25 μ M MSX, whilst the remaining culture was re-fed with fresh CD-CHO/phenol red/25 μ M MSX, and incubated for a further 14 days. At this point, culture supernatant was collected from the 24-well
25 plates and the antibody concentration was measured using Lonza's ELISA assay for assembled antibody. A selection of the most productive cell lines was made. On reaching confluence in the T25 flasks, the selected cell lines were allowed to multilayer and then re-fed with CD-CHO/25 μ M MSX. After feeding, the cultures were returned to the incubator for a further four to seven

days until the medium turned orange/yellow again and the cells lifted from the flasks.

Expansion of cell lines to CDACF suspension culture

Suspension cultures were initiated from confluent T25 flask cultures. The suspension cultures were inoculated at a concentration of 0.05 to 0.2×10^6 viable cells/mL in 125 mL shake flasks containing CD-CHO/25 μ M MSX medium. If the viable cell concentration did not reach 0.05×10^6 viable cells/mL after a maximum of 14 days in T25 flasks, 15 mL of each culture was automatically progressed into 15 mL of CD-CHO/25 μ M MSX in 125 mL shake flasks. The cell lines were then serially subcultured into fresh CD-CHO/25 μ M MSX medium, at an inoculation concentration of 0.2×10^6 viable cells/mL, on a 4 day subculture regime, until acceptable and reproducible growth characteristics were obtained.

*Assessment of growth and productivity of cell lines in suspension culture***Fed-batch shake flask culture**

Singlet cultures of each selected cell line were prepared in 500 mL shake flasks with 100 mL of cell suspension using CM42/4 x SPE medium. The cultures were inoculated at 0.2×10^6 viable cells/mL and the headspace of each culture was equilibrated with 5% v/v CO₂ in air. The cultures were incubated at 35.5 to 37.0 °C on a shaking platform rotating at 140 ± 5 rpm until the viable cell concentration, post peak, was less than or equal to 0.6×10^6 viable cells/mL or day 15 was reached ('overgrown'). At this point the cultures were harvested. The cell concentration was determined daily using a Vi-CELL automated cell counter. At a cell concentration of 1.4 to 2.2×10^6 viable cells/mL, a bolus addition of the nutrient feed SF40 was made. Further bolus additions were made at approximately 24, 48 and 72 hours after the initial addition. A second feed (SF41) was applied once cell concentrations greater than 4.0×10^6 viable cells/mL were obtained. Samples of cultures were taken at appropriate intervals, centrifuged and the supernatants stored at -20 ± 5 °C until assayed for antibody concentration by Protein A HPLC. Harvest supernatants from

each overgrown culture were stored at -20 ± 5 °C. The ten overgrown cultures with the highest antibody concentrations were further assessed for quality of the antibody produced. Samples were assessed for aggregate levels using the gel permeation high performance liquid chromatography (GP-HPLC) method before Protein A purification and antibody characterisation by reduced and non-reduced sodium dodecylsulphate poly acrylamide gel electrophoresis (SDS-PAGE) and isoelectric focusing (IEF) analysis and matrix assisted laser desorption ionisation time of flight mass spectrometry (MALDI TOF-MS) oligosaccharide analysis.

10

Cryopreservation

Cells were cryopreserved at key points throughout the work programme. The cells in growing cultures were recovered by centrifugation and resuspended in a cryopreservation mixture. This consisted of the appropriate growth medium (92.5% v/v) and dimethylsulphoxide (7.5% v/v).

15

Aliquots of cells were then distributed into labelled vials, each aliquot containing approximately 1.5 mL of cryopreservation medium and approximately 0.5 to 1.5×10^7 viable cells. The vials were then frozen in an automatic programmable cell freezing machine and subsequently transferred to liquid nitrogen refrigerators.

20

Generation number

For each of the selected cell lines, the generation number of the cells was defined as zero at the point the culture was passaged into suspension. The generation number was subsequently calculated at each subculture using a procedure derived from the following equation (rounded down to the nearest 0.5 generation).

25

$$N_f = N_i + \left\{ \frac{1}{\ln 2} \cdot \ln \frac{X_f}{X_i} \right\}$$

Where: N_f = Generation number at time f

N_i = Generation number at time i

5 X_f = Viable cell concentration (cells/mL) at time f

X_i = Viable cell concentration (cells/mL) at time i

and $f > i$

Data analysis

10 The time integral of the area under the growth curve (the time integral of the viable cell concentration (IVC); 10^9 cells.h/L) was calculated by summing the area (approximated to a right angle trapezium) between adjacent cell concentration values. The areas were calculated by multiplying the mean of the two viable cell concentrations (cells/mL) by the elapsed time between the two determinations (h). This method is based on that described by Renard *et al.* (1988) in *Biotechnology Letters* 10 (2) pages 91-96.

Where appropriate the overall specific production rate (q_p overall:

mg/ 10^9 cells/h) was calculated by linear regression analysis of the antibody concentration (mg/L) against the time integral of viable cell concentration.

20 The slope of this line is equal to q_p overall. The harvest specific production rate (q_p harvest) was calculated by dividing the harvest antibody concentration by the value of the IVC at harvest.

Assembly ELISA

25 The antibody concentration of supernatant samples was determined using an ELISA that measures assembled human IgG. This involved capture of assembled antibody in samples and standard onto a 96-well plate coated with goat anti-human IgG Fc. Bound antibody was revealed with goat anti-human

light chain linked to horseradish peroxidase and the chromogenic substrate tetramethylbenzidine. Colour development was proportional to the mass of antibody present in the sample. The concentration was determined using a standard curve generated using an IgG standard (Lot number L07387/5/2)

5

Protein A HPLC

Product concentration was quantified by Protein A High Performance Liquid Chromatography (HPLC) on an Agilent 1100 HPLC. Antibody was selectively bound to a POROS Protein A immunodetection column. Unbound material was eluted from the column and the remaining bound antibody was released by decreasing the pH of the solvent. Elution was monitored by absorbance at 280 nm using a multiple-wavelength detector. Antibody was quantified (using Chemstation software) against an antibody standard (Lot L07385/4/10) and corrected for an extinction coefficient of $E_{280nm}^{0.1\%} = 1.52$.

15

Protein A affinity purification

Gravity fed rmp Protein A Sepharose columns (5 mL) were prepared. The columns were cleaned before use with 6 M guanidine HCl and equilibrated with 50 mM glycine/glycinate-250 mM NaCl buffer, pH 8.0. The pH of the column load material was adjusted to pH 8.0 ± 0.1 before application to the column. The rmp Protein A affinity column was washed with equilibration buffer and eluted with 0.1 M glycine buffer, pH 3.5. The column eluates were neutralised to approximately pH 7.0 with 2 M Tris Base and dialysed against Dulbecco's phosphate buffered saline in preparation for IEF and SDS PAGE analyses.

25

SDS PAGE analysis of Protein A affinity purified antibody

The cell culture samples were prepared for analysis by protein A purification. Electrophoresis was carried out using 4 to 20% precast Novex polyacrylamide gels. For reducing SDS PAGE, samples were reduced using 2-mercaptoethanol

30

and denatured by SDS at pH 8.0. Samples were heated for 2.0 ± 0.5 minutes before they were loaded onto the gel (10 μg per lane). Non-reduced SDS PAGE was performed using the same gels and Novex non-reducing sample buffer (2-mercaptoethanol is not added). Samples were heated for a standard time of

5 1.0 ± 0.5 minutes before they were loaded onto the gel (4 μg per lane). Following electrophoresis, proteins were visualised by staining for 75 minutes with Coomassie brilliant blue R250 and destained with methanol/acetic acid.

IEF analysis of Protein A affinity purified antibody

The cell culture samples were prepared for analysis by protein A purification.

IEF analysis was carried out using pre-cast agarose IEF gels, pH 3 to pH 10.

- 5 Approximately 10 µg protein sample per lane was loaded onto the gel and then pre-focused for 70 Vh. After removal of the sample application strip, electrophoresis was continued for 1500 Vh (1500 V, 20 mA, 25 W per gel). The IEF gels were fixed in a trichloroacetic acid: sulphosalicylic acid: methanol solution and then stained with Coomassie Brilliant Blue R250 (Lot numbers of
- 10 Coomassie Blue 003/L11905/118 and 003/L11905/131). The gels were then destained and dried. (Lot numbers of destain 003/L11905/117 and 003/L11905/121).

GP-HPLC

- 15 The GP-HPLC method separates protein components according to their size. Large components, such as protein aggregates, are too large to penetrate the matrix particles to any great extent and therefore eluted from the column ahead of smaller components, such as protein monomer. Smaller components, such as fragments, penetrate the matrix more easily and eluted after the
- 20 monomer. This technique was therefore used to separate antibody monomer from both aggregates and fragments. The monomeric component was identified by its characteristic retention time and position, relative to calibration markers.

- Undiluted samples were injected to give 50 to 250 µg loads where the product
- 25 concentration was 1 to 5 mg/mL (validated load range of Lonza's assay). The molecular mass range for the GF-250 column was 4 to 400 kDa.

- Sample components were detected at 280 nm and peak chromatograms were analysed using Agilent Chemstation software. Integration was performed using the perpendicular drop method for aggregates and tangent skim for
- 30 fragments.

The proportion of sample components was determined by calculation of the peak areas of each component relative to the total integrated peak area.

MALDI TOF-MS analysis

- 5 The oligosaccharide analysis of the PG102 antibody produced by each cell line was performed by MALDI TOF-MS using a Micromass™ MALDI-LR coupled with delayed extraction. The MALDI TOF-MS was operated in reflectron configuration with positive ion mode. The instrument was calibrated using a mixed N-glycan standard.
- 10 Antibody was purified from harvest supernatants using a POROS Protein A column. The purified fractions were collected and the disulphide bonds were reduced with dithiothreitol followed by alkylation of the reduced thiols using iodoacetate. Oligosaccharides were released using the enzyme N-glycanase (PNGaseF) and were sandwiched between two layers of super-dihydroxy
- 15 benzoic acid (super-DHB) matrix on the target plate for MALDI TOF-MS analysis.

Results

Transfection of CHOK1SV host cells

- 20 The PG102 gene sequence was used by Lonza to construct the double gene vector pPG102/DGV (not shown) that was used to transfect the CHOK1SV host cell line by electroporation. Six sets of transfections were performed using the vector pPG102/DGV and supernatants of 255 transfectants were screened for antibody production in 96-well plates by assembly ELISA. Detectable
- 25 antibody levels were produced by 99.6% of the screened transfectants. Forty three high ranking transfectants were selected for further evaluation in 24-well plates.

Expansion of static cultures and suspension adaptation

Cultures of the selected GS-CHO cell lines were first expanded into 24-well plates and then expanded into T25 static flasks. The remaining cells in the 24-well plates were fed with fresh medium and 'overgrown' before assessing the antibody productivity. The selected cell lines displayed a range of antibody concentrations from 8.3 to 120 mg/L (Figure 10).

Cultures of the 23 cell lines with the highest product concentrations, as determined by assembly ELISA, were selected for further evaluation. The selected cultures were progressed into CDACF medium in suspension culture. All cell lines were successfully adapted to suspension culture and were selected for further evaluation in fed-batch cultures.

Evaluation of the growth and productivity characteristics of selected cell lines in fed-batch shake flask culture

The growth and productivity characteristics of the 23 selected cell lines were evaluated in fed-batch shake flask culture. Singlet cultures were prepared for each cell line. Two feeds, SF40 and SF41, were added to the cultures when the viable cell concentration met the feeding criteria. The protocol used for the feed regime mimicks as closely as technically possible, Lonza's generic GS-CHO fermentation process. Growth and productivity data for the 23 selected cell lines are presented in Table 9. Protein A HPLC analysis of the harvested supernatants of the fed-batch shake flask cultures showed that four cell lines produced antibody concentrations of greater than 1000 mg/L. Cell line L107 produced the highest antibody concentration, 1674 mg/L.

Characterisation of the antibody purified from fed-batch shake flask culture of selected CHO cell lines

Harvest samples of the ten cell lines with the highest antibody concentration were purified using rmp Protein A Sepharose chromatography prior to analysis. The Protein A purified antibodies were then analysed using SDS PAGE and IEF. Visual analysis of the Protein A purified samples analysed by SDS PAGE under non-reducing conditions showed all samples were comparable to each other, exhibiting an intact antibody band at approximately 200 kDa (data not shown). Additional minor bands were visible, although they may not all be visible on the gel image. Three bands with molecular mass between 116 and 200 kDa, one band with a molecular weight between 66 and 97 kDa, one band with a molecular weight between 37 and 55 kDa and two bands with molecular weights between 22 and 31 kDa were also observed. The extra band observed between 66 and 97 kDa is the half antibody, typical of IgG₄ antibodies under non-reducing conditions. The same half antibody is also observed in the IgG₄ inter assay control (IAC).

Protein A purified samples analysed by SDS PAGE under reducing conditions were also comparable to each other, exhibiting the heavy chain band at approximately 50 kDa and the light chain band at approximately 25 kDa (Data not shown). IEF analysis showed that the integrity of the Protein A purified antibody samples from the selected cell lines demonstrated comparable profiles, with six primary bands (3 major and 3 minor bands) in the pI range 8.15 to 9.30 for all cell lines except L97, M92 and M59. These three cell lines exhibited three major and two minor bands. Additional bands were observed in samples J3 and J4, although these may not all be visible on the gel image.

Oligosaccharide analysis of antibody

The oligosaccharide analysis of the antibody produced by each of the ten cell lines selected for further evaluation in fed-batch shake flask cultures was

determined by MALDI TOF-MS. The predominant oligosaccharide structures in the antibody obtained from the ten cell lines were G0F and G1F, which are typical N-linked oligosaccharide structures observed on antibodies. Relatively low levels of oligomannose structures were detected with the highest level of 6.7% of total glycans measured for antibody derived from cell line L52. Oligomannose-5 (man-5) structures were observed in all samples at a range of 1.1 to 4.6% of total glycans. All of the samples analysed contained comparable levels of oligosaccharides and relatively low concentrations of man-5 (data not shown).

10

GP-HPLC

When antibody produced by each of the ten cell lines selected for further evaluation was analysed by GP-HPLC, the lower molecular weight components (LMWC) were the least critical but a level of under 25% was preferable. All samples displayed levels of under 25%, with the exception of L52, which displayed a level of 27.1% (Table 10).

Two different calculations of proportions of aggregate would not usually be used, but was deemed necessary as LMWC levels varied so much between samples.

Further analysis of the data, excluding the LMWC and looking only at monomer and larger peaks, demonstrated aggregate peak areas of less than 2.2% for all samples (Table 11). Levels of under 5% are considered acceptable.

Selection of cell lines

Three cell lines were selected for preparation of pre-seed stocks (PSS). The selection criteria used were a combination of high harvest antibody concentration, acceptable growth characteristics and acceptable product quality characteristics.

Cell line L107 was selected as the lead cell line as it exhibited the highest harvest antibody concentration of the 23 cell lines evaluated, along with acceptable growth and product quality characteristics (Figure 11A).

5 Cell line L25 exhibited the second highest harvest antibody concentration of the twenty three cell lines evaluated, along with acceptable growth and product quality characteristics (Figure 11B). Therefore cell line L25 was selected as the first back-up cell line.

10 Cell line M95 exhibited the third highest harvest antibody concentration of the twenty three cell lines evaluated, and had acceptable growth and product quality characteristics (Figure 11C). On this basis cell line M95 was selected as the second back-up cell line.

Cryopreservation of pre-seed stocks

15 Pre-seed stocks (20 vials per stock) were cryopreserved for each of the three selected cell lines. These stocks were stored in vapour phase liquid nitrogen refrigerators. Cell lines L107, L25 and M95 were renamed DC1, DC2 and DC3 respectively.

Discussion

20 Lonza Biologics was requested to undertake the construction, selection and evaluation of a GS-CHO cell line expressing the human recombinant IgG₄/kappa antibody PG102.

25 The gene sequence was supplied by PanGenetics and vectors were generated using Lonza's Glutamine Synthetase (GS) Gene Expression System. The cell line CHOK1SV, a suspension variant of the Chinese hamster ovary (CHO) cell line CHO-K1 adapted to chemically defined, animal component-free medium, was transfected by electroporation.

30 From the CHOK1SV transfections, 254 cell lines were obtained that secreted the antibody. Forty three cell lines were evaluated for growth and productivity in 24-well plates. Antibody concentrations of up to 120 µg/mL were obtained.

Twenty three cell lines were evaluated in singlet fed-batch shake flask cultures. Harvest antibody concentrations in the range of 121 to 1674 µg/mL were obtained. Harvest antibody concentrations in excess of 1000 mg/L were achieved for four cell lines. The ten cell lines with highest antibody concentrations were selected for product quality analysis. The product quality of purified antibody from each of the selected cell lines was comparable to each other by reduced and non-reduced sodium dodecylsulphate polyacrylamide gel electrophoresis and isoelectric focusing analysis. Oligosaccharide analysis of purified antibody from each of the ten selected cell lines showed relatively low (<4.6%) levels of oligomannose-5.

GP-HPLC analysis showed that aggregate levels were below 1% for each of the top three cell lines.

Three cell lines (L107, L25 and M95) were selected for further evaluation based on growth, productivity and product quality data. Pre-seed stocks of 20 vials of each cell line were cryopreserved. The three cell lines were renamed DC1, DC2 and DC3 respectively.

Example 4

Determination of the kinetics of ch5D12 (PG100) and PG102 (I29L variant) binding to human CD40-Fc by surface plasmon resonance (Biacore)

Materials and Methods

The kinetics of ch5D12 (PG100) and PG102 (I29L variant) binding to human CD40 were compared by surface plasmon resonance at a constant temperature of 25.0 °C, using a Biacore 3000 instrument. Recombinant human CD40 extracellular domain fused to human IgG1-Fc domain (huCD40-Fc; R&D Systems catalogue number 149-CD-50) was used as the target antigen in these experiments. HuCD40-Fc (1µg/mL in 10mM acetate buffer, pH 5.0) was

coupled via the amine groups of Lys residues to carbodiimide-activated Biacore CM5 chips, resulting in an antigen immobilization level of 127.4 RU. For kinetic analyses, dilution series of ch5D12 (PG100) and PG102 (range 10.0 $\mu\text{g/mL}$ to 0.313 $\mu\text{g/mL}$) were prepared in Biacore Running Buffer (HEPES-buffered saline containing EDTA and surfactant; HBS-EP). Kinetic analyses were performed at a flow rate of 30 $\mu\text{L/min}$. Samples were allowed to bind for 4 min and to dissociate for 5 min, at the same flow rate. The chip surface was then regenerated for 30 seconds using 30mM NaOH and a baseline re-established in HBS-EP for 2.5 min before injection of the next sample. Alternate samples of PG102 and ch5D12 (PG100) were injected, increasing in concentration for the first set of replicates at each dilution and then decreasing for the second set of replicates.

BIAcore control software (Version 3.2) was used for data collection and BIAevaluation software (Version 4.1) was used for analysis of kinetic data.

Binding parameters were calculated using two different models of molecular interaction: a bivalent analyte model and Langmuir 1:1 binding model.

Results

Maximum levels of observed binding for ch5D12 (PG100) and PG102 antibodies were 152.5 RU and 140.2 RU, respectively.

The calculated kinetic rate constants according to the Langmuir and bivalent binding models (k_a and k_d , or k_{a1} and k_{d1} , respectively) differed slightly for the two antibodies. Thus, PG102 displayed faster association and dissociation rates than ch5D12 (PG100) (Figure 12), although the overall affinity (K_D) of the two antibodies did not differ appreciably within models. Kinetic parameters from these experiments are summarised below:

Langmuir (1:1) model:

	ch5D12 (PG100)	PG102
ka	$1.88 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$	$2.51 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$
5 kd	$5.07 \times 10^{-4} \text{ s}^{-1}$	$5.96 \times 10^{-4} \text{ s}^{-1}$
KD	$2.71 \times 10^{-10} \text{ M}$	$2.37 \times 10^{-10} \text{ M}$

Bivalent binding (1:2) model:

	ch5D12 (PG100)	PG102
10 ka1	$6.41 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$	$7.82 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$
kd1	$1.60 \times 10^{-3} \text{ s}^{-1}$	$2.47 \times 10^{-3} \text{ s}^{-1}$
ka2	$1.21 \times 10^{-3} \text{ RU s}^{-1}$	$1.31 \times 10^{-3} \text{ RU s}^{-1}$
kd2	$2.55 \times 10^{-3} \text{ s}^{-1}$	$2.24 \times 10^{-3} \text{ s}^{-1}$
15 KD	$5.29 \times 10^{-9} \text{ M}$	$5.41 \times 10^{-9} \text{ M}$

Analysis of the Goodness of Fit for the two methods indicated that the data is better described by the bivalent binding model.

The calculated affinities of the two antibody variants differed between models, however, the estimates of the dissociation constants (KD) obtained for each antibody with the same model were comparable; $2.71 \times 10^{-10} \text{ M}$ and $2.37 \times 10^{-10} \text{ M}$ for ch5D12 (PG100) and PG102 respectively using the Langmuir model and $5.25 \times 10^{-9} \text{ M}$ and $5.41 \times 10^{-9} \text{ M}$ using the bivalent analyte model.

25 Example 5

Inhibition of ch5D12 (PG100) and PG102 binding to JY cells using an anti-idiotypic antibody

To demonstrate that PG102 (I29L variant) binds to the same epitope of CD40 as its parent molecule ch5D12 (PG100), an anti-idiotypic antibody was used to

inhibit binding of PG102 or ch5D12 to CD40-expressing JY cells, measured by FACS.

Materials and Methods

5 JY cells expressing human CD40 were blocked with 5% human serum for 30 min at room temperature. The anti-idiotypic antibody to murine mAb 5D12 (clone 173-36-1) was pre-incubated at varying concentration (0 – 10 $\mu\text{g/mL}$) with either ch5D12 (PG100) (1 $\mu\text{g/mL}$), PG102 (1 $\mu\text{g/mL}$), or negative control chimaeric anti-huCD86 antibody (chFUN-1; 1 $\mu\text{g/mL}$) in a total volume of 50
10 μL /tube and incubated for 15 min. JY cells were washed with FACS buffer (1x PBS, 1% BSA and 0.05 % sodium azide) and supernatant was discarded. JY cells were resuspended in FACS buffer to a concentration of 2×10^6 cells/mL. 50 μL of the JY cell suspension was added to the pre-incubated antibody mixes to give a final volume of 100 μL /tube. Cells were incubated for 30 min at 4 °C
15 before washing with 4 ml FACS buffer/tube and the supernatant discarded. The cell pellet was resuspended in 100 μL of diluted goat-anti-human IgG-FITC (Jackson ImmunoResearch Labs. Cat. No. 109-095-127) in FACS buffer (1:100) and incubated for 30 min at 4 °C. Cells were washed with 4 ml FACS buffer and supernatant was discarded. The pellet was resuspended in 200 μL FACS
20 buffer and bound antibody was measured using a FACScan flow cytometer (Becton Dickinson).

Results

The anti-idiotypic mAb 173-36-1 inhibited PG102 and ch5D12 (PG100) binding
25 to huCD40 expressed on JY cells in a concentration-dependent manner (Figure 13). The binding of chFUN-1 to CD86 that is also expressed on JY cells was not influenced by mAb 173-36-1. There was no significant difference in the blocking effect of anti-idiotypic mAb 173-36-1 on the binding of CD40-specific antibodies to CD40 on JY cells. For example, the calculated log IC₅₀ values for
30 mAb 173-3-1-mediated inhibition of ch5D12 (PG100) (Batch 1) and PG102

(Batch 2) were -5.74 ± 0.14 and -5.76 ± 0.11 , respectively ($p > 0.05$; both $n=4$). These values correspond to anti-idiotypic concentrations of approximately 1.8 $\mu\text{g/mL}$ (~ 12 nM). Table 12 shows a summary of calculated $-\log\text{IC}_{50}$ values for anti-idiotypic-mediated inhibition of ch5D12 (PG100) and PG102 binding to JY

5 cells.

Example 6

*Binding of ch5D12 (PG100) and PG102 to CD-40Fc determined by ELISA***Material and Methods**

- 5 ELISA plates (Costar EIA/RIA plate, Corning catalogue number 3590) were coated with 100 μ L/well huCD40-muIg (Ancell, catalogue number 504-020), diluted to 250 ng/mL in PBS and incubated overnight at 4 °C in a humid environment. Plates were washed 3 times with 200 μ L/well wash buffer (0.05% Tween-20 in PBS). Subsequently, plates were blocked with 200 μ L/well
- 10 blocking buffer (5% BSA Fraction V [Roche, catalogue number 735094] in wash buffer) and incubated for 1 h at 37 °C (humid environment). Plates were washed 3 times with wash buffer. Test antibodies (ch5D12 (PG100), PG102, and negative control chimaeric anti-huCD86 antibody chFUN-1) were diluted in the range 0-1200 ng/mL in blocking buffer and transferred to assay plates in
- 15 a final volume of 100 μ L/well, followed by incubation for 1 h at 37 °C (humid environment). Plates were washed 3 times with wash buffer followed by addition of 100 μ L/well goat anti-human kappa-AP detection antibody (Southern Biotech Associates catalogue number 2060-0) diluted 1:1000 in blocking buffer. Plates were incubated for 1 h at 37 °C in a humid environment
- 20 before washing 3 times with wash buffer and a further single wash with PBS. PNP substrate was prepared by adding 1 tablet p-Nitrophenyl Phosphate tablets (Sigma, catalogue number N-2765) in 15 mL PNP substrate buffer (12.1 g Tris, 5.84 g NaCl, 1.02 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ in 1L H_2O (adjusted to pH 9.6 with HCl). PNP substrate was added to assay plates at 100 μ L/well. Plates
- 25 were incubated for several minutes (max 30 min) at 37 °C to allow colour development before the reaction was stopped by addition of 3M NaOH. Colour intensity was determined at 405 nm using a micro plate reader (Biorad, model 550).

Results

In a number of independent experiments serially-diluted ch5D12 (PG100) and PG102 (I29L variant) were found to bind comparably to immobilized CD40-Fc (Figure 14), whilst the anti-CD86 control antibody did not display appreciable binding to CD40-Fc. Calculated half-maximal binding concentrations for all experiments are summarized in Table 13. No significant difference was observed in the half-maximal binding concentration for any of the anti-CD40 antibody batches evaluated.

Example 7

Binding of ch5D12 (PG100) and PG102 to huCD40 expressed on JY cells determined by FACS

5 Materials and Methods

CD40-expressing JY cells were washed with FACS buffer (1x PBS, 1 % BSA and 0.05 % sodium azide). Supernatant was discarded and JY cells were resuspended in FACS buffer to a concentration of 2×10^6 cells/mL. Thereafter, 50 μ L of the JY cell suspension was added to 50 μ L of each of the test
10 antibodies (ch5D12 (PG100), PG102, and negative control chimaeric anti-huCD86 antibody chFUN-1) prepared in the concentration range 0-900 ng/mL and incubated for 30 min at 4 °C. Cells were washed with 4 ml FACS
buffer/tube and supernatant was discarded. Cell pellets were resuspended in 100 μ L of goat anti-human IgG-FITC (Jackson Immunoresearch Labs.
15 Catalogue number 109-095-127) diluted 1:100 in FACS buffer and incubated for 30 min at 4 °C. Cells were washed with 4 ml FACS buffer and supernatant was discarded. The pellet was resuspended in 200 μ L FACS buffer and bound antibody was measured using a FACScan flow cytometer (Becton Dickinson).

20 Results

All tested batches of ch5D12 (PG100) and PG102 showed similar characteristics of binding to CD40 expressed on JY cells (Figure 15) with comparable half-maximal antibody binding concentrations (Table 14). chFUN-1 also bound to JY cells, albeit with lower mean fluorescence intensity (MFI)
25 than ch5D12 (PG100) and PG102. This is because JY cells also express human CD86.

Discussion

The CD40-binding characteristics of various PG102 (I29L variant) and ch5D12 (PG100) batches have been determined using in vitro cellular and ELISA-based assays. Binding of PG102 and ch5D12 (PG100) to JY cells expressing CD40 was inhibited with equal potency by the anti-idiotypic mAb 173-36-1. Thus, the range of calculated mean $-\log IC_{50}$ values for PG102 and CH5D12 (PG100) batches was 5.51 to 6.11 (n=8), and 5.43 to 6.04 (n=10), respectively (p>0.05). Antibody 173-36-1 is directed against the variable regions of the ch5D12 (PG100) precursor, murine mAb 5D12, and did not inhibit binding of an anti-CD86 isotype control antibody. In further experiments the half-maximal binding concentration of PG102 and ch5D12 (PG100) was determined by ELISA analysis of antibody binding to human CD40-Fc, and by FACS analysis of binding to JY cells. In ELISA assays there was no significant difference between the half-maximal binding concentration of PG102 and ch5D12 (PG100) for any batch tested. For example, calculated mean $-\log$ half-maximal binding concentrations for the PG102 and ch5D12 (PG100) batches were 7.47 ± 0.04 and 7.54 ± 0.03 , respectively (P>0.05, both n=4), corresponding to approximately 30 ng/ml. Similar half-maximal binding concentrations were observed in FACS experiments of PG102 and ch5D12 (PG100) binding to CD40 expressed on JY cells. These results indicate that PG102 and ch5D12 (PG100) share the same CD40-binding paratope, as demonstrated by the comparable potency characteristics of the anti-idiotypic antibody for inhibition of PG102 and ch5D12 (PG100) binding to CD40. Furthermore, both antibodies display similar in vitro potencies for binding to human CD40 in ELISA and cell based assays. Together, the data presented suggest that the binding properties of PG102 and ch5D12 (PG100) to human CD40 are the same.

Example 8

Competitive inhibition of ch5D12 (PG100)-PE and PG102-PE binding to JY cells expressing human CD40 using unlabelled anti-CD40 antibody

5

Materials and Methods

JY cells (Epstein–Barr virus-transformed human B lymphoblastoid cell line) were grown in Iscove's Modified Dulbecco's Medium (IMDM) containing 10% heat-inactivated foetal calf serum (FCS), 2 mM L-glutamine and 50 µg/mL gentamycin in an air-humidified atmosphere containing 5% CO₂, at 37°C. Cells were harvested on the day of flow cytometric measurements. Unlabelled ch5D12 (PG100) (Batch 4) and PG102 (Batch 2) antibodies were used in these experiments. Both antibodies were also used in PE-labelled form for direct determination of binding to JY cells by flow cytometry. Antibodies were custom-labelled with PE by AbD Serotec (Oxford, UK). For each FACS analysis the following protocol was used. JY cells were harvested from cell culture and counted. Subsequently 1×10^5 CD40-expressing JY cells /200 µL incubation buffer (PBS, 1% BSA, 0.05% sodium azide) were incubated with 1µg/mL labelled ch5D12 (PG100) or PG102 antibodies, in the presence of varying concentration (0-10 µg/mL) of unlabelled competing ch5D12 (PG100) or PG102, for 30 min at 4-8°C. Cells were washed followed by flow cytometric analysis in which the MFI at every concentration was determined using a FACScan flow cyometer (Becton & Dickinson). Five thousand events/sample were measured and analyzed using CellQuest® software (Becton & Dickinson).

25

Results

Unlabelled ch5D12 (PG100) and PG102 antibodies displayed similar inhibitory characteristics in these assays (Figure 16). Mean $-\log IC_{50}$ values for inhibition of 1 µg/mL ch5D12 (PG100) binding by unlabelled ch5D12 (PG100) and PG102 were 7.50 ± 0.03 and 7.63 ± 0.03 , respectively (both n=4), whilst mean –

30

log IC₅₀ values for inhibition of 1 ug/ml PG102 binding by unlabelled ch5D12 (PG100) and PG102 were 7.54 ± 0.01 and 7.65 ± 0.02 , respectively (both n=4). These -log IC₅₀ values correspond to PG102 and ch5D12 (PG100) concentrations of approximately 20-30 ng/mL (i.e. ~130-200pM).

5

Example 9

Inhibition of IL-8 release from THP-1 cells

Materials and Methods

10 The functionality of PG102 was measured using a cell-based functional bioassay. Briefly, on day 1, THP-1 and Jurkat 39.8/50 human cells were cultured in Iscove's Modified Dulbecco's Medium (IMDM, BioWhittaker, catalogue number BE12-722F supplemented with 10 % foetal bovine serum (Gibco, ref 10270-106) and 50 µg/mL gentamycin (Invitrogen, catalogue
15 number 15750-045). THP-1 cells were pulsed with rhuIFN-γ at 1000 units/mL of cell culture.

On day 3 of the bioassay THP-1 and Jurkat 39.8/50 cells were required at a concentration of 4×10^5 cells/mL. The THP-1 and J39.8/50 cells were counted and their viability determined. Cells were then diluted to a concentration of
20 approximately 1×10^6 cells/mL, Cell suspensions were mixed 1:1 and incubated at 37 °C in a humidified 5 % CO₂ atmosphere for 48 h.

On day 3 the THP-1 and J39.8/50 cells were washed as follows (In the case of THP-1 cells the wash step was to remove the IFN-γ): 5 mL of IFN-γ pulsed
25 THP-1 and J39.8/50 cell suspension were transferred to a 50 mL falcon tube. 40 mL of Hank's Balanced Salt Solution (HBSS) was added, and the cells were centrifuged at 1500 rpm (using IEC centrifuge) for 6 min. The supernatant was discarded. The cell pellet was resuspended in pre-warmed IMDM

supplemented with 10 % FCS. The THP-1 cells and J39.8/50 cells were adjusted to a concentration of 4×10^5 cells/mL.

For inhibition assays the test sample ch5D12 (PG100) or PG102 (I29L variant) was serially diluted in warmed IMDM medium supplemented with 10 % foetal bovine serum to obtain final assay concentrations in the range 0 – 160 ng/mL. The THP-1, J39.8/50 and test sample(s) were added in triplicate to round bottomed cell culture plates (NuncTM) in the following order: 50 μ L of THP-1 cells (equivalent to 2×10^4 cells per well), 50 μ L of the test sample and 50 μ L J39.8/50 cells. The total volume was 150 μ L per well. The cell culture plates were wrapped in porous cling film and incubated at 37 °C in a humidified 5 % CO₂ atmosphere for 48 h.

On day 5, after a culture period of 48 h, 70 μ L of cell culture supernatants were aspirated and transferred to a round bottomed microtitre plate. The harvested cell culture supernatants were assayed for IL-8 content using a commercial ELISA (Biosource, Human IL-8 cytoset, catalogue number CHC1304) in accordance with the manufacturer's instructions.

20 Results

PG102 (I29L variant) and ch5D12 (PG100) showed similar inhibitory effects on evoked IL-8 release by THP-1 cells (Figure 17). Calculated $-\log IC_{50}$ values for the two antibodies were 8.42 and 8.28, respectively. These values correspond to IC_{50} concentrations of approximately 30pM.

25

Example 10

A previous tissue cross-reactivity study on human and cynomolgus tissues showed that ch5D12 (PG100) bound to the cell surface of B cells and DCs in lymphoid organs. No unexpected cross-reactivity was observed on either

30

human or cynomolgus tissues. This study was repeated for PG102 and similar results were obtained (data not shown) indicating that ch5D12 (PG100) and PG102 bind to a variety of tissue sections in a similar fashion.

- 5 Also a previous safety and tolerability evaluation for ch5D12 was performed in cynomolgus monkeys, in which weekly administration of ch5D12 for 4 weeks was shown to be safe and without any side-effects in all monkeys. In this study, functional evidence was obtained that ch5D12 can prevent B-cell activation and proliferation²⁷. Safety studies were repeated for PG102 using a
- 10 more prolonged protocol. Briefly, a 13 week intravenous toxicity study in cynomolgus monkey with a 14 week recovery period was designed, using 3 dose levels (0, 25 and 100 mg/kg i.v.) and 26 monkeys (6 monkeys (3M, 3F) per treatment group, plus 4 monkeys (2M, 2F) as recovery groups in each active treatment arm). The following measurements were done: TK, anti-PG102
- 15 response, flow cytometry (including CD40 coating of PBMC), lymph node biopsy and standard toxicology assessments such as haematology and immunohistochemistry.

The results of this study demonstrated that there is no toxicity observed for any of the dose levels tested. PG102 was safe and well-tolerated.

20

Collectively these studies demonstrate that antagonist anti-human CD40 Mab PG102, as its parent antibody ch5D12 (PG100), has no unexpected cross-reactivity, and is safe and well tolerated in vivo.

Table 1. Baseline demographics by cohort

Cohort	Dose of ch5D12 (mg/kg)	Gender M/F	Mean age	Mean weight (kg)	Mean length (cm)	Mean BMI	Mean CDAI	Mean CDEIS
1	0.3	1/4	30	57	164	21	301	16 (n = 2)
2	1.0	2/3	43	72	169	25	276	14 (n = 1)
3	3.0	1/4	34	65	168	23	292	15 (n = 5)
4	10.0	3/0	34	70	170	24	331	15 (n = 3)
Total		7/11	35	65	167	23	300	15 (n = 11)

BMI, body mass index; CDAI, Crohn's Disease Activity Index; CDEIS, Crohn's Disease Endoscopic Index of Severity.

Table 1

Table 2

Table 2. Decreases in Crohn's Disease Activity Index scores after ch5D12 treatment

Subject	CDAI at screening	Decrease in CDAI at day 28	Max CDAI decrease ^a	Day of max CDAI decrease	Remission (yes/no) ^b	Day of remission
Cohort 1: 0.3 mg/kg ch5D12						
001	253	-30	41	7	No	-
002	274	57	107	21	No	-
003	320	86	129	14	No	-
004	392	58	66	7	No	-
005	265	117	149	14	Yes	14
Mean	301	58	98	12.6		
Cohort 2: 1 mg/kg ch5D12						
006	232	-43	69	7	No	-
007	228	133	175	21	Yes	21
008	394	144	172	21	No	-
009	288	56	123	21	No	-
010 ^c	241	-42	No decrease	-	No	-
Mean	277	50	108	14		
Cohort 3: 3 mg/kg ch5D12						
011	299	27	27	28	No	-
012	308	194	194	28	Yes	28
013	280	106	106	28	No	-
014	327	-84	147	14	No	-
015	249	103	103	28	No	-
Mean	293	69	115	25.2		
Cohort 4: 10 mg/kg ch5D12						
016	357	-65	109	21	No	-
017	358	302	301	28	Yes	14
018	280	20	102	21	No	-
Mean	332	86	171	23.3		

^a Maximal decrease in CDAI over the 28-day observation period after ch5D12 infusion.^b Remission was defined as a total CDAI score of ≤ 150 and a decrease of >100 .^c All subjects showed a decrease in their CDAI score except for subject 010 (1.0 mg/kg cohort). This subject had a screening score of 241.4 and the CDAI score never dropped below the value obtained at screening.

CDAI, Crohn's Disease Activity Index.

Table 3

Table 3. Immunohistochemistry scoring^a

Subject ID	Cohort (mg/kg)	Sample origin	CD3		CD4		CD8		CD19		CD68	
			0	28	0	28	0	28	0	28	0	28
001	0.3	Ileum	++	++	++	++	++	++	++	++	+	++
		Colon	++	++	++	++	++	++	++	++	++	++
005	0.3	Ileum	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
		Colon	++	+	++	+	++	+	++	+	++	+
009	1.0	Ileum	++	+	++	+	++	+	++	+	++	+
		Colon	+	+	+	+	+	+	+	+	+	+
011	3.0	Ileum	++	+	++	+	++	+	++	+	++	+
		Colon	+	+	+	+	+	+	+	+	+	+
012	3.0	Ileum	++	+	++	+	++	+	++	+	++	+
		Colon	++	+	++	+	++	+	++	+	++	+
013	3.0	Ileum	++	+	++	+	++	+	++	+	++	+
		Colon	+	+	+	+	+	+	+	+	+	+
014	3.0	Ileum	++	+	++	+	++	+	++	+	++	+
		Colon	+	+	+	+	+	+	+	+	+	+
015	3.0	Ileum	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
		Colon	+	+	+	+	+	+	+	+	+	+
016	10	Ileum	+	+	+	+	+	+	+	+	+	+
		Colon	+	+	+	+	+	+	+	+	+	+
017	10	Ileum	+	+	+	+	+	+	+	+	+	+
		Colon	+	+	+	+	+	+	+	+	+	+
018	10	Ileum	+	+	+	+	+	+	+	+	+	+
		Colon	++	+	++	+	++	+	++	+	++	+

Lymphocyte results: 0 = screening, 28 = day 28.

NS, no sample.

^a All available biopsies were stained with antibodies recognizing all T lymphocytes (CD3), T-helper cells (CD4), cytotoxic T cells (CD8), B cells (CD19) and macrophages (CD68). The value for CD3⁺, CD4⁺, CD8⁺, CD19⁺ and CD68⁺ for both the ileum and colon from non-inflamed tissue is single positive (+).

Table 4

primer	sequence 5' - 3'
Q5E	ctcccagggttaagcttgaggagtctggacctgg
Q5Erev	ccagggtccagactcctcaagcttaacctgggag
K13A	gacctggcctggtggcaccctcagagaccc
K13Arev	gggtctctgaggggtgccaccaggccaggtc
E16Q	cctggtgaaaccctcacagaccctgtccatcac
E16Qrev	gtgatggacagggctctgtgagggtttcaccagg
T17S	gtgaaaccctcagagagcctgtccatcacatgc
T17Srev	gcatgtgatggacaggctctctgagggtttcac
I29L	gcactgtctctgggttctcactctccagatatagtgtatac
I29Lrev	gtatacactatatctggagagtgagaacccagagacagtgc
I37V	gatatagtgtatactgggttcgccagcctccagg
I37Vrev	cctggaggctggcgaaccagtatatacactatatac
P45L	cctccaggaaagggctctggagtggatgggaatg
P45Lrev	cattcccattccactccagaccctttcctggagg
M48L	gggtccggagtggctgggaatgatgtg
M48Lrev	cacatcattcccagccactccggaccc
STS60NSA	gtggtggttccacagactataattcagctctcaaataccagactgacc
STS60NSArev	ggtcagtctggatttgagagctgaattatagtctgtggaaccaccac
T68S	ctcaaataccagactgagcatcagcaaggacacc
T68Srev	ggtgtccttgctgatgctcagtctggatttgag
S79F	cacctcgaagagccaggctcttcttaaaaatgaacagtctgc
S79Frev	gcagactgttcatttttaagaagacctggctcttcgaggtg
T108S	gggtcaaggaacctcggtcaccgtctc
T108Srev	gagacggtgaccgaggttccttgaccc

Mutagenic oligonucleotides. Oligonucleotides used to introduce the mutations at amino acid position 5, 13, 16, 17, 29, 37, 45, 48, 60, 68, 79 and 108. For each position is a sense and a antisense ("rev") oligonucleotide necessary.

Table 5

	FACS	ELISA	ratio
variant	MFI	ng/ml	
Q5E	39	444	20
K13A	31	267	12
E16Q	39	420	19
T17S	33	482	22
I29L	108	1331	61
I37V	96	1175	54
P45L	82	566	26
M48L	80	525	24
STS60NSA	62	308	14
T68S	35	295	13
S79F	96	869	40
T108S	99	648	30
CH5D12	103	2187	100
DI5D12	50	392	18
mock	11	n.d.	
blank	10	n.d.	

Overview expression data as measured by FACS and quantitative ELISA. For each 5D12 variant (Q5E, K13A, E16Q, T17S, I29L, I37V, P45L, M48L, STS60NSA, T68S, S79F, and T108S) together with ch5D12 and DI5D12 the MFI value (as determined by FACS) and the antibody concentration in the harvested supernatant (as determined by ELISA) is shown. Also MFI values shown for mock transfection and medium control. Ratio is calculated using the ELISA data compared with ch5D12 value (stated as 100%).

Table 6: mutagenic oligonucleotides. Oligonucleotides used to introduce the mutations at amino acid positions 29 and 37. Sense and antisense (rev) oligonucleotides are listed.

Primer	sequence 5' - 3'
I29V	gcactgtctctgggttctcagtctctagatatagtgtatac
I29Vrev	gtatacactatatctagagactgagaacccagagacagtgc
I29L	gcactgtctctgggttctcactctccagatatagtgtatac
I29Lrev	gtatacactatatctggagagtgagaacccagagacagtgc
I37L	gatatagtgtatactggctgcgccagcctccagg
I37Lrev	cctggaggctggcgccagccagtatacactatatac
I37V	gatatagtgtatactgggttcgccagcctccagg
I37Vrev	cctggaggctggcgaacccagtatacactatatac

Table 7: The creation of double mutants required 2 mutagenesis rounds.

Below it is listed which primer sets were used in step 1 and step 2 respectively to come to the additional DI5D12 variants.

Mutant	Step 1	Step 2
PG102 [29L-37I (LI)]	I29L+I29Lrev	n.a.
29I-37V (IV)	I37V+I37Vrev	n.a.
29V-37I (VI)	I29V+I29Vrev	n.a.
29I-37L (IL)	I37L+I37Lrev	n.a.
29V-37V (VV)	I29V+I29Vrev	I37V+I37Vrev
29L-37L (LL)	I29L+I29Lrev	I37L+I37Lrev
29V-37L (VL)	I29V+I29Vrev	I37L+I37Lrev
29L-37V (LV)	I29L+I29Lrev	I37V+I37Vrev

Table 8. Expression data as measured by quantitative ELISA. For each additional 5D12 variant (29I-37V, 29V-37I, 29I-37L, 29V-37V, 29L-37L, 29V-37L and 29L-37V), together with ch5D12 (PG100), DI5D12 and PG102 (I29L variant), the antibody concentration in the harvested supernatant is shown.

variant	name	ELISA (ng/ml)
29L-37V	ch5D12 (PG100)	2846
29I-37I	DI5D12	587
29L-37I	PG102	2226
29I-37V		1728
29V-37I		928
29I-37L		729
29V-37V		2027
29L-37L		1115
29V-37L		891
29L-37V		4263

Table 9: Summary of growth & productivity data of CHO cell lines in CDACF fed-batch shake flask cultures

Culture ID	Maximum Viable Cell Concentration 10 ⁶ cells/mL	Time Integral of Viable Cell Concentration 10 ⁹ cells.h/L	Product Concentration at Harvest mg/L	Specific Production Rate (q _p harvest) mg/10 ⁹ cells/h	Specific Production Rate (q _p overall) mg/10 ⁹ cells/h
L107 (DC1)	9.66	2201	1674	0.76	0.78
L25 (DC2)	11.06	2610	1258	0.48	0.48
M95 (DC3)	9.58	2146	1173	0.55	0.58
L52	13.45	2989	1001	0.33	0.34
L73	5.95	1325	936	0.71	0.77
L45	10.48	2096	915	0.44	0.45
J3	9.20	2012	888	0.44	0.48
J4	11.02	2037	871	0.43	0.45
L97	7.55	1821	819	0.45	0.53
M92	7.82	1665	807	0.48	0.53
M59	14.65	3270	792	0.24	0.24
L102	6.20	1282	733	0.57	0.60
M58	8.78	1970	662	0.34	0.39
G20	5.73	1391	658	0.47	0.52
L65	3.74	878	651	0.74	0.90
L56	8.52	1728	546	0.32	0.30
M63	10.79	2465	444	0.18	0.18
G7	9.87	2058	376	0.18	0.20
G11	5.04	1145	310	0.27	0.30
L1	5.78	1163	263	0.23	0.21
M15	11.23	2654	121	0.05	0.04
M54	9.95	2278	276	0.12	0.15
M56	11.56	2347	27	0.01	0.01

Table 10: GP-HPLC Analysis (including LMWC)

Sample Number	Culture ID	Relative Peak Area (%)			
		Total Aggregate	Monomer	Fragment	LMWC
1	L107	0.62	84.8	ND	14.58
2	L25	0.65	76.5	ND	22.88
3	M95	0.74	84.7	ND	14.60
4	L52	0.94	72.0	ND	27.07
5	L73	1.48	87.9	ND	10.63
6	J3	1.01	92.5	ND	6.48
7	J4	1.68	73.4	0.04	24.91
8	L97	1.66	84.4	ND	13.92
9	M92	1.18	89.4	ND	9.42
10	M59	1.55	84.8	ND	13.63

ND=Not Detected

Table 11: GP-HPLC Analysis (IgG products only)

Sample Number	Culture ID	Relative Peak Area (%) without LMWC		
		Total Aggregate	Monomer	Fragment
1	L107	0.72	99.3	ND
2	L25	0.84	99.2	ND
3	M95	0.87	99.1	ND
4	L52	1.28	98.7	ND
5	L73	1.66	98.3	ND
6	J3	1.08	98.9	ND
7	J4	2.24	97.7	0.05
8	L97	1.93	98.1	ND
9	M92	1.30	98.7	ND
10	M59	1.79	98.2	ND

ND=Not Detected

Anti-CD40 mAb (batch)	-log IC₅₀	n
ch5D12 (PG100) (Batch 1)	5.74 ± 0.14	4
ch5D12 (PG100) (Batch 2)	5.64 ± 0.01	2
ch5D12 (PG100) (Batch 3)	5.55 ± 0.06	2
ch5D12 (PG100) (Batch 4)	5.63 ± 0.02	2
PG102 (Batch 1)	5.85 ± 0.12	4
PG102 (Batch 2)	5.76 ± 0.11	4

Table 12: Summary of calculated $-\log IC_{50}$ values for anti-idiotypic-mediated inhibition of ch5D12 (PG100) and PG102 binding to JY cells. Data indicate the mean $\pm$ s.e. mean of n determinations. There was no significant difference in the inhibitory potency of the anti-idiotypic with any of the antibody batches tested ($p > 0.05$, ANOVA followed by Tukey's test for multiple comparisons).

Anti-CD40 mAb (batch)	-log half-maximal binding concentration	n
ch5D12 (PG100) (Batch 1)	7.54 ± 0.03	4
ch5D12 (PG100) (Batch 2)	7.53 ± 0.02	2
ch5D12 (PG100) (Batch 3)	7.48 ± 0.10	2
ch5D12 (PG100) (Batch 4)	7.37 ± 0.10	2
PG102 (Batch 1)	7.45 ± 0.04	4
PG102 (Batch 2)	7.47 ± 0.04	4

Table 13. Summary of calculated -log half-maximal binding concentration values for ch5D12 (PG100) and PG102 binding to human CD40-Fc, as

- 5 determined by ELISA. Data indicate the mean ± s.e. mean of n determinations. There was no significant difference in the binding of any antibody batch tested ($p > 0.05$, ANOVA followed by Tukey's test for multiple comparisons). For reference, a -log half-maximal binding concentration of 7.45 corresponds to a concentration of approximately 35 ng/ml (~ 200 pmol).

Anti-CD40 mAb (batch)	-log half-maximal binding concentration	n
ch5D12 (PG100) (Batch 1)	7.26	2
ch5D12 (PG100) (Batch 2)	7.34	2
ch5D12 (PG100) (Batch 3)	7.49	2
ch5D12 (PG100) (Batch 4)	7.59	2
PG102 (Batch 1)	7.44	2
PG102 (Batch 2)	7.48	2

Table 14: Summary of calculated -log half-maximal binding concentration values for ch5D12 (PG100) and PG102 binding to JY cells expressing human CD40, as determined by ELISA. Data indicate the mean of duplicate determinations. For reference, a -log half-maximal binding concentration of 7.45 corresponds to a concentration of 35 ng/ml (~ 200 pmol).

References:

- 1 Ranheim EA, Kipps TJ. Activated T cells induce expression of B7/BB1 on
5 normal or leukemic B cells through a CD40-dependent signal. *J Exp Med*
(1993); 177: 925-35.
- 2 Hasbold J, Johnson-Leger C, Atkins CJ, Clark EA, Klaus GGB. Properties
of mouse CD40: cellular distribution of CD40 and B cell activation by
10 monoclonal anti-mouse CD40 antibodies. *Eur J Immunol* (1994); 24: 1835-42.
- 3 Alderson MR, Armitage RJ, Tough TW, Strockbine L, Fanslow WC, Spriggs
MK. CD40 expression by monocytes: regulation by cytokines and activation of
monocytes by the ligand for CD40. *J Exp Med* (1993); 178: 669-74.
15
- 4 Kiener PA, Moran-Davis P, Rankin BM, Wahl AF, Aruffo A, Hollenbaugh
D. Stimulation of CD40 with purified soluble gp39 induces proinflammatory
responses in human monocytes. *J Immunol* (1995); 155: 4917-25.
- 20 5 Shu U, Kiniwa M, Wu CY, et al. Activated T cells induce interleukin-12
production by monocytes via CD40-CD40 ligand interaction. *Eur J Immunol*
(1995); 25: 1125-8.
- 6 Cella M, Scheidegger D, Palmer-Lehmann K, Lane P, Lanzavecchia A,
25 Alber G. Ligation of CD40 on dendritic cells triggers production of high levels
of interleukin-12 and enhances T cell stimulatory capacity: T-T help via APC
activation. *J Exp Med* (1996); 184: 747-52.
- 7 van Kooten C, Banchereau J. Functions of CD40 on B cells, dendritic cells
30 and other cells. *Curr Opin Immunol* (1997); 9: 330-7.

- 8 Cayabyab M, Phillips JH, Lanier LL. CD40 preferentially costimulates activation of CD4+ T lymphocytes. *J Immunol* (1994); 152: 1523-31.
- 5 9 Hermann P, Van-Kooten C, Gaillard C, Banchereau J, Blanchard D. CD40 ligand-positive CD8+ T cell clones allow B cell growth and differentiation. *Eur J Immunol* (1995); 25: 2972-7.
- 10 10 Grewal IS, Flavell RA. CD40 and CD154 in cell-mediated immunity. *Annu Rev Immunol* (1998); 16: 111-35.
- 11 Henn V, Slupsky JR, Grafe M, et al. CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature* (1998); 391: 591-4.
- 15 12 van Kooten C, Banchereau J. CD40-CD40 ligand. *J Leukoc Biol* (2000); 67: 2-17.
- 13 Clegg CH, Rulffes JT, Haugen HS, et al. Thymus dysfunction and chronic inflammatory disease in gp39 transgenic mice. *Int Immunol* (1997); 9: 1111-22.
- 20 14 Stüber E, Strober W, Neurath M. Blocking the CD40L-CD40 interaction in vivo specifically prevents the priming of T helper 1 cells through the inhibition of interleukin 12 secretion. *J Exp Med* (1996); 183: 693-8.
- 25 15 Liu Z, Geboes K, Colpaert S, et al. Prevention of experimental colitis in SCID mice reconstituted with CD45RB^{high} CD4+ T cells by blocking the CD40-CD154 interactions. *J Immunol* (2000); 164: 6005-14.

- 16 De Jong YP, Comiskey M, Kalled SL, et al. Chronic murine colitis is dependent on the CD154/CD40 pathway and can be attenuated by anti-CD154 administration. *Gastroenterology* (2000); 119: 715-23.
- 5 17 Kawai T, Andrews D, Colvin RB, Sachs DH, Cosimi AB. Thromboembolic complications after treatment with monoclonal antibody against CD40 ligand. *Nat Med* (2000); 6: 114.
- 18 Buhler L, Alwayn IP, Appel JZ III, Robson SC, Cooper DK. Anti-CD154
10 monoclonal antibody and thromboembolism. *Transplantation* (2001); 71: 491.
- 19 Knosalla C, Gollackner B, Cooper DK. Anti-CD154 monoclonal antibody and thromboembolism revisited. *Transplantation* (2002); 74: 416-17.
- 15 20 de Boer M, Conroy J, Min HY, Kwekkeboom J. Generation of monoclonal antibodies to human lymphocyte cell surface antigens using insect cells expressing recombinant proteins. *J Immunol Meth* (1992); 152: 15-23.
- 21 Kwekkeboom J, de Boer M, Tager JM, de Groot C. CD40 plays an essential
20 role in the activation of human B cells by murine EL4B5 cells. *Immunology* (1993); 79: 439-44.
- 22 Kwekkeboom J, de Rijk D, Kasran A, Barcy S, de Groot C, de Boer M. Helper effector function of human T cells stimulated by anti-CD3 Mab can be
25 enhanced by co-stimulatory signals and is partially dependent on CD40-CD40 ligand interaction. *Eur J Immunol* (1994); 24: 508-17.
- 23 Laman JD, 't Hart BA, Brok, HPM, et al. Protection of marmoset monkeys against EAE by treatment with a murine antibody blocking CD40 (mu5D12).
30 *Eur J Immunol* (2002); 32: 2218-28.

- 24 Boon L, Brok HPM, Bauer J, et al. Prevention of experimental autoimmune encephalomyelitis in the common marmoset (*Callithrix jacchus*) using a chimeric antagonist Mab against human CD40 is associated with altered B-cell responses. *J Immunol* (2001); 167: 2942-9.
- 25 Haanstra KG, Ringers J, Sick EA, et al. Prevention of kidney allograft rejection using anti-CD40 and anti-CD86 in primates. *Transplantation* (2003); 75: 637-43.
- 26 Haegel-Kronenberger H, Haanstra K, Ziller-Remy C, et al. Inhibition of costimulation allows for repeated systemic administration of adenoviral vector in rhesus monkeys. *Gene Ther* (2004); 11: 241-52.
- 27 Boon L, Laman JD, Ortiz-Buijsse A, et al. Preclinical assessment of anti-CD40 Mab 5D12 in cynomolgus monkeys. *Toxicology* (2002); 174: 53-65.
- 28 Sartor RB. Pathogenesis and immune mechanisms of chronic inflammatory bowel disease. *Am J Gastroenterol* (1997); 92: 5S-11S.
- 29 Fiocchi C. Inflammatory bowel disease: etiology and pathogenesis. *Gastroenterology* (1998); 115: 182-205.
- 30 Burgio VL, Fais S, Boirivant M, Perrone A, Pallone F. Peripheral monocyte and naive T-cell recruitment and activation in Crohn's disease. *Gastroenterology* (1995); 109: 1029-38.
- 31 Rugtveit J, Brandtzaeg P, Halstensen TS, Fausa O, Scott H. Increased macrophage subset in inflammatory bowel disease: apparent recruitment from peripheral blood monocytes. *Gut* (1994); 35: 669-74.

- 32 Fuss IJ, Neurath M, Boirivant M, et al. Disparate CD4+ lamina propria (LP) lymphokine secretion profiles in inflammatory bowel disease. *J Immunol* (1996); 157: 1261-70.
- 5
- 33 Mullin GE, Maycon ZR, Braun-Elwert L, et al. Inflammatory bowel disease mucosal biopsies have specialized lymphokine mRNA profiles. *Inflam Bowel Dis* (1996); 2: 16-26.
- 10
- 34 Pospai, D, Rene E, Fiasse R, et al. Crohn's disease stable remission after human immunodeficient virus infection. *Dig Dis Sci* (1998); 43: 412-9.
- 35 Stronkhorst A, Radema S, Yong SL, et al. CD4 antibody treatment in patients with active Crohn's disease: a phase 1 dose finding study. *Gut* (1997); 40: 320-7.
- 15
- 36 Sadlack B, Merz H, Schorle H, Schimpl A, Feller AC, Horak I. Ulcerative colitis-like disease in mice with a disrupted interleukin-2 gene. *Cell* (1993); 75: 253-61.
- 20
- 37 Kuhn R, Lohler J, Rennick D, Rajewsky K, Muller K. Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* (1993); 75: 263-74.
- 38 Powrie F, Leach MM, Mauze S, Menon S, Caddle LB, Coffman RA.
- 25 Inhibition of Th1 responses prevents inflammatory bowel disease in SCID mice reconstituted with CD45RBhi CD4+ T cells. *Immunity* (1994); 1: 553-62.
- 39 Liu Z, Colpaert S, D'Haens GR, et al. Hyperexpression of CD40 Ligand (CD154) in inflammatory bowel disease and its role in pathogenic cytokine
- 30 production. *J Immunol* (1999); 163: 4049-57.

- 40 D'Haens G, Geboes K, Peeters M, Baert F, Penninckx F, Rutgeerts P. Early lesions caused by infusion of intestinal contents in excluded ileum in Crohn's disease. *Gastroenterology* (1998); 114: 262-7.
- 5
- 41 Cornillie F, Shealy D, D'Haens G, et al. Infliximab induces potent anti-inflammatory and local immunomodulatory activity but no systemic immune suppression in patients with Crohn's disease. *Aliment Pharmacol Ther* (2001); 15: 2041-2.
- 10
- 42 Inwald DP, McDowall A, Peters MJ, Callard RE, Klein NJ. CD40 is constitutively expressed on platelets and provides a novel mechanism for platelet activation. *Circ Res* (2003); 92: 1041-8.
- 15
- 43 Van Assche G, Rutgeerts P. Anti-TNF agents in Crohn's disease. *Expert Opin Investig Drugs* (2000); 9: 103-11.
- 20
- 44 Stüber E, Strober W, Neurath M. Blocking the CD40L-CD40 interaction in vivo specifically prevents the priming of T helper 1 cells through the inhibition of interleukin 12 secretion. *J Exp Med* (1996);183:693-698.
- 25
- 45 Ranheim EA, Kipps TJ. Activated T cells induce expression of B7/BB1 on normal or leukemic B cells through a CD40-dependent signal. *J Exp Med* (1993);177:925-935.
- 46 Hasbold J, Johnson-Leger C, Atkins CJ, Clark EA, Klaus GGB. Properties of mouse CD40: cellular distribution of CD40 and B cell activation by monoclonal anti-mouse CD40 antibodies. *Eur J Immunol* (1994);24:1835-1842.

- 47 Alderson MR, Armitage RJ, Tough TW, Strockbine L, Fanslow WC, Spriggs MK. CD40 expression by monocytes: regulation by cytokines and activation of monocytes by the ligand for CD40. *J Exp Med* (1993);178:669-674.
- 5 48 Kiener PA, Moran-Davis P, Rankin BM, Wahl AF, Aruffo A, Hollenbaugh D. Stimulation of CD40 with purified soluble gp39 induces proinflammatory responses in human monocytes. *J Immunol* (1995);155:4917-4925.
- 49 Shu U, Kiniwa M, Wu CY, Maliszewski C, Vezzio N, Hakimi J, Gately M, Delespesse G. Activated T cells induce interleukin-12 production by monocytes via CD40-CD40 ligand interaction. *Eur J Immunol* (1995);25:1125-1128.
- 10 50 Cella M, Scheidegger D, Palmer-Lehmann K, Lane P, Lanzavecchia A, Alber G. Ligation of CD40 on dendritic cells triggers production of high levels of interleukin-12 and enhances T cell stimulatory capacity: T-T help via APC activation. *J Exp Med* (1996);184:747-752.
- 15 51 van Kooten C, Banchereau J. Functions of CD40 on B cells, dendritic cells and other cells. *Curr Opin Immunol* (1997);9:330-337.
- 20 52 Cayabyab M, Phillips JH, Lanier LL. CD40 preferentially costimulates activation of CD4+ T lymphocytes. *J Immunol* (1994);152:1523-1531.
- 53 Hermann P, Van-Kooten C, Gaillard C, Banchereau J, Blanchard D. CD40 ligand-positive CD8+ T cell clones allow B cell growth and differentiation. *Eur J Immunol* (1995);25:2972-2977.
- 25 54 Grewal IS, Flavell RA. CD40 and CD154 in cell-mediated immunity. *Annu Rev Immunol* (1998);16:111-135.

- 55 Henn V, Slupsky JR, Grafe M, Anagnostopoulos I, Forster R, Muller-Berghaus G, KroczeK RA. CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature* (1998);391:591-594.
- 5 56 Banchereau J, Bazan F, Blanchard D, Briere F, Galizzi JP, van Kooten C, Liu YJ, Saeland S. The CD40 antigen and its ligand. *Annu Rev Immunol* (1994);12:881-922.
- 10 57 Foy TM, Aruffo A, Bajorath J, Buhlmann JE, Noelle RJ. Immune regulation by CD40 and its ligand gp39. *Annu Rev Immunol* (1996);14:591-617.
- 15 58 Kato T, Hakamada R, Yamane H, Nariuchi H. Induction of IL-12 p40 messenger RNA expression and IL-12 production of macrophages via CD40-CD40 ligand interaction. *J Immunol* (1996);156:3932-3938.
- 20 59 Ludewig B, Graf D, Gelderblom HR, Becker Y, KroczeK RA, Pauli G. Spontaneous apoptosis of dendritic cells is efficiently inhibited by TRAP (CD40-ligand) and TNF- α , but strongly enhanced by interleukin-10. *Eur J Immunol* (1995);25:1943-1950
- 25 60 Allen RC, Armitage RJ, Conley ME, Rosenblatt H, Jenkins NA, Copeland NG, Bedell MA, Edelhoff S, Disteché CM, Simoneaux DK, Fanshow WC, Belmont J, Spriggs MK. CD40 ligand gene defects responsible for X-linked hyper-IgM syndrome. *Science* (1993);259:990-993.
- 61 Xu J, Foy TM, Laman JD, Elliott EA, Dunn JJ, Waldschmidt TJ, Elsemore J, Noelle RJ, Flavell RA. Mice deficient for the CD40 ligand. *Immunity* (1994);1:423-431.

- 62 Grewal IS, Xu J, Flavell RA. Impairment of antigen-specific T-cell priming in mice lacking CD40 ligand. *Nature* (1995);378:617-620.
- 63 Stout RD, Suttles J, Xu J, Grewal IS, Flavell RA. Impaired T cell-mediated
5 macrophage activation in CD40 ligand-deficient mice. *J Immunol* (1996);156:8-11.
- 64 Levy J, Espanol-Boren T, Thomas C, Fischer A, Tovo P, Bordigoni P, Resnick I, Fasth A, Baer M, Gomez L, Sanders EA, Tabone MD, Plantaz D,
10 Etzioni A, Monafo V, Abinun M, Hammarstrom L, Abrabamsen T, Jones A, Finn A, Klemola T, DeVries E, Sanal O, Peitsch MC, Notarangelo LD. Clinical spectrum of X-linked hyper-IgM syndrome. *J Pediatr* (1997);131:47-54.
- 65 Soong L, Xu JC, Grewal IS, Kima P, Sun J, Longley BJ Jr, Ruddle NH,
15 McMahon-Pratt D, Flavell RA. Disruption of CD40-CD40 ligand interactions results in an enhanced susceptibility to *Leishmania amazonensis* infection. *Immunity* (1996);4:263-273.
- 66 Campbell KA, Ovendale PJ, Kennedy MK, Fanslow WC, Reed SG,
20 Maliszewski CR. CD40 ligand is required for protective cell-mediated immunity to *Leishmania major*. *Immunity* (1996);4:283-289.
- 67 Clegg CH, Rulffes JT, Haugen HS, Hoggatt IH, Aruffo A, Durham SK, Farr AG, Hollenbaugh D. Thymus dysfunction and chronic inflammatory
25 disease in gp39 transgenic mice. *Int Immunol* (1997);9:1111-1122.
- 68 Parker et al., *Proc. Nat. Acad. Sci. USA* (1995) 92:9560

- 69 Haanstra KG, Ringers J, Sick EA, Ramdien-Murli S, Kuhn EM, Boon L, Jonker M. Prevention of kidney allograft rejection using anti-CD40 and anti-CD86 in primates. *Transplantation*. (2003);75(5):637-43.
- 5 70 Haanstra KG, Sick EA, Ringers J, Wubben JAM, Kuhn EM, Boon L, Jonker M. Costimulation blockade followed by a 12-week period of cyclosporine A facilitates prolonged drug-free survival of Rhesus Monkey kidney allografts. *Transplantation*. (2005); 79: 1623-1626.
- 10 71 Boon L, Brok HP, Bauer J, Ortiz-Buijsse A, Schellekens MM, Ramdien-Murli S et al. Prevention of experimental autoimmune encephalomyelitis in the common marmoset (*Callithrix jacchus*) using a chimeric antagonist monoclonal antibody against human CD40 is associated with altered B cell responses. *J Immunol* (2001); 167:2942-2949.
- 15 72 Laman JD, 't Hart BA, Brok H, Meurs M, Schellekens MM, Kasran A et al. Protection of marmoset monkeys against EAE by treatment with a murine antibody blocking CD40 (mu5D12). *Eur J Immunol* (2002); 32:2218-2228.
- 20 73 Boon L, Laman JD, Ortiz-Buijsse A, den Hartog MT, Hoffenberg S, Liu P et al. Preclinical assessment of anti-CD40 Mab 5D12 in cynomolgus monkeys. *Toxicology* (2002); 174:53-65.
- 25 74 Kasran A, Boon L, Wortel CH, Hogezaand RA, Schreiber S, Goldin E, Boer M, Geboes K, Rutgeerts P, Ceuppens JL. Safety and tolerability of antagonist anti-human CD40 Mab ch5D12 in patients with moderate to severe Crohn's disease. *Aliment Pharmacol Ther*. (2005) ;22:111-22.

- 75 Haanstra KG, Sick EA, Ringers J, Wubben JA, Kuhn EM, Boon L, Jonker M. Costimulation blockade followed by a 12-week period of cyclosporine A facilitates prolonged drug-free survival of rhesus monkey kidney allografts. *Transplantation*. (2005);79:1623-6.
- 5
- 76 't Hart BA, Blezer EL, Brok HP, Boon L, de Boer M, Bauer J, Laman JD. Treatment with chimeric anti-human CD40 antibody suppresses MRI-detectable inflammation and enlargement of pre-existing brain lesions in common marmosets affected by MOG-induced EAE. *J Neuroimmunol*. (2005);163:31-9.
- 10
- 77 Koenen HJ, den Hartog MT, Heerkens S, Fasse E, Ortiz-Buijsse A, van Neerven RJ, Simons PJ, Joosten I, Boon L. A novel bispecific antihuman CD40/CD86 fusion protein with t-cell tolerizing potential. *Transplantation*. (2004);78:1429-38.
- 15
- 78 de Vos AF, Melief MJ, van Riel D, Boon L, van Eijk M, de Boer M, Laman JD. Antagonist anti-human CD40 antibody inhibits germinal center formation in cynomolgus monkeys. *Eur J Immunol*. (2004);34:3446-55.
- 20
- 79 Laman JD, 't Hart BA, Brok H, Meurs M, Schellekens MM, Kasran A, Boon L, Bauer J, Boer M, Ceuppens J. Protection of marmoset monkeys against EAE by treatment with a murine antibody blocking CD40 (mu5D12). *Eur J Immunol*. (2002);32:2218-28.
- 25
- 80 Boon L, Laman JD, Ortiz-Buijsse A, den Hartog MT, Hoffenberg S, Liu P, Shiao F, de Boer M. Preclinical assessment of anti-CD40 Mab 5D12 in cynomolgus monkeys. *Toxicology*. (2002);174:53-65.

- 81 Boon L, Brok HP, Bauer J, Ortiz-Buijsse A, Schellekens MM, Ramdien-Murli S, Blezer E, van Meurs M, Ceuppens J, de Boer M, 't Hart BA, Laman JD. Prevention of experimental autoimmune encephalomyelitis in the common marmoset (*Callithrix jacchus*) using a chimeric antagonist monoclonal antibody against human CD40 is associated with altered B cell responses. J Immunol. (2001);167:2942-9.
- 5
- 82 S.G. Antunes, N.G. de Groot, H. Brok, G. Doxiadis, A.A. Menezes, N. Otting and R.E. Bontrop, The common marmoset: a new world primate species with limited Mhc class II variability, Proc. Natl. Acad. Sci. U. S. A. (1998);95:, 11745–11750.
- 10
- 83 B. Becher, B.G. Durell, A.V. Miga, W.F. Hickey and R.J. Noelle, The clinical course of experimental autoimmune encephalomyelitis and inflammation is controlled by the expression of CD40 within the central nervous system, J. Exp. Med. (2001);193:. 967–974.
- 15
- 84 R.E. Bontrop, N. Otting, N.G. de Groot and G.G. Doxiadis, Major histocompatibility complex class II polymorphisms in primates, Immunol. Rev. (1999);167:. 339–350.
- 20
- 85 H.P. Brok, A. Uccelli, N. Kerlero De Rosbo, R.E. Bontrop, L. Roccatagliata, N.G. de Groot, E. Capello, J.D. Laman, K. Nicolay, G.L. Mancardi, A. Ben-Nun and B.A. 't Hart, Myelin/oligodendrocyte glycoprotein-induced autoimmune encephalomyelitis in common marmosets: the encephalitogenic T cell epitope pMOG24-36 is presented by a monomorphic MHC class II molecule, J. Immunol. (2000);165: 1093–1101.
- 25

- 86 H.P. Brok, J. Bauer, M. Jonker, E. Blezer, S. Amor, R.E. Bontrop, J.D. Laman and B.A. 't Hart, Non-human primate models of multiple sclerosis, *Immunol. Rev.* (2001);183: 173–185.
- 5 87 C.P. Genain and S.L. Hauser, Experimental allergic encephalomyelitis in the New World monkey *Callithrix jacchus*, *Immunol. Rev.* (2001);183: 159–172.
- 88 K. Gerritse, J.D. Laman, R.J. Noelle, A. Aruffo, J.A. Ledbetter, W.J. Boersma and E. Claassen, CD40–CD40 ligand interactions in experimental
10 allergic encephalomyelitis and multiple sclerosis, *Proc. Natl. Acad. Sci. U. S. A.* (1996);93: 2499–2504.
- 89 A.M. Girvin, M.C. Dal Canto and S.D. Miller, CD40/CD40L interaction is essential for the induction of EAE in the absence of CD28-mediated co-
15 stimulation, *J. Autoimmun.* (2002);18:83–94.
- 90 I.S. Grewal and R.A. Flavell, A central role of CD40 ligand in the regulation of CD4+ T-cell responses, *Immunol. Today* (1996);17:410–414.
- 20 91 L.M. Howard, A.J. Miga, C.L. Vanderlugt, M.C. Dal Canto, J.D. Laman, R.J. Noelle and S.D. Miller, Mechanisms of immunotherapeutic intervention by anti-CD40L (CD154) antibody in an animal model of multiple sclerosis, *J. Clin. Invest.* (1999);103:281–290.
- 25 92 E.K. Jordan, H.I. McFarland, B.K. Lewis, N. Tresser, M.A. Gates, M. Johnson, M. Lenardo, L.A. Matis, H.F. McFarland and J.A. Frank, Serial MR imaging of experimental autoimmune encephalomyelitis induced by human white matter or by chimeric myelin-basic and proteolipid protein in the common marmoset, *AJNR Am. J. Neuroradiol.* (1999);20:965–976.

- 93 J.D. Laman, E. Claassen and R.J. Noelle, Functions of CD40 and its ligand, gp39 (CD40L), *Crit. Rev. Immunol.* (1996);16:59–108.
- 5 94 J.D. Laman, C.B. Maassen, M.M. Schellekens, L. Visser, M. Kap, E. de Jong, M. van Puijenbroek, M.J. van Stipdonk, M. van Meurs, C. Schwarzler and U. Gunthert, Therapy with antibodies against CD40L (CD154) and CD44-variant isoforms reduces experimental autoimmune encephalomyelitis induced by a proteolipid protein peptide, *Mult. Scler.* (1998);4: 147–153.
- 10 95 J.D. Laman, M. van Meurs, M.M. Schellekens, M. de Boer, B. Melchers, L. Massacesi, H. Lassmann, E. Claassen and B.A. 't Hart, Expression of accessory molecules and cytokines in acute EAE in marmoset monkeys (*Callithrix jacchus*), *J. Neuroimmunol.* (1998);86:30-45.
- 15 96 J. Mestas and C.C. Hughes, Of mice and not men: differences between mouse and human immunology, *J. Immunol.* (2004);172: 2731–2738.
- 20 97 S.A. Quezada, L.Z. Jarvinen, E.F. Lind and R.J. Noelle, CD40/CD154 interactions at the interface of tolerance and immunity, *Annu. Rev. Immunol.* (2004);22:307–328.
- 25 98 C.S. Raine, B. Cannella, S.L. Hauser and C.P. Genain, Demyelination in primate autoimmune encephalomyelitis and acute multiple sclerosis lesions: a case for antigen-specific antibody mediation, *Ann. Neurol.* (1999);46:144–160.
- 99 D.H. Sachs, Tolerance: of mice and men, *J. Clin. Invest.* (2003);111:1819–1821.

- 100 E.B. Samoilova, J.L. Horton, H. Zhang and Y. Chen, CD40L blockade prevents autoimmune encephalomyelitis and hampers TH1 but not TH2 pathway of T cell differentiation, *J. Mol. Med.* (1997);75: 603–608.
- 5 101 B.A. 't Hart, J. Bauer, H.J. Muller, B. Melchers, K. Nicolay, H. Brok, R.E. Bontrop, H. Lassmann and L. Massacesi, Histopathological characterization of magnetic resonance imaging-detectable brain white matter lesions in a primate model of multiple sclerosis: a correlative study in the experimental autoimmune encephalomyelitis model in common marmosets (*Callithrix*
- 10 *jacchus*), *Am. J. Pathol.* (1998);153:649–663.
- 102 B.A. 't Hart, M. van Meurs, H.P. Brok, L. Massacesi, J. Bauer, L. Boon, R.E. Bontrop and J.D. Laman, A new primate model for multiple sclerosis in the common marmoset, *Immunol. Today*(2000);21:290–297.
- 15 103 B. 't Hart, S. Amor and M. Jonker, Evaluating the validity of animal models for research into therapies for immune-based disorders, *Drug Discov. Today* (2004);9: 517–524.
- 20 104 't Hart et al., 2004b B.A. 't Hart, J.D. Laman, J. Bauer, E.D. Blezer, Y. van Kooyk and R.Q. Hintzen, Modelling of multiple sclerosis: lessons learned in a non-human primate, *Lancet Neurol* (2004);3: 589–597.
- 25 105 B.A. 't Hart, J. Vogels, J. Bauer, H.P. Brok and E. Blezer, Non-invasive measurement of brain damage in a primate model of multiple sclerosis, *Trends Mol. Med.* (2004);10: 85–91.
- 30 106 B.A. 't Hart, J.T. Vogels, J. Bauer, H.P.M. Brok and E. Blezer, Non-invasive measurement of brain damage in a primate model of multiple sclerosis, *Trends Mol. Med.* (2004);10: 85–91.

- 107 P. Villoslada, K. Abel, N. Heald, R. Goertsches, S.L. Hauser and C.P. Genain, Frequency, heterogeneity and encephalitogenicity of T cells specific for myelin oligodendrocyte glycoprotein in naive outbred primates, *Eur. J. Immunol.* (2001);31: 2942-2950.
- 108 Bata-Csorgo Z, Hammerberg C, Voorhees JJ, Cooper KD. Intralesional T-lymphocyte activation as a mediator of psoriatic epidermal hyperplasia. *J Invest Dermatol* (1995); 105(1 Suppl): 89S-94S.
- 109 Bos JD, De Rie MA. The pathogenesis of psoriasis: immunological facts and speculations. *Immunol Today* (1999); 20: 40-46.
- 110 Lebwohl M. Psoriasis. *Lancet* (2003); 361: 1197-1204.
- 111 van Kooten C, Banchereau J. CD40-CD40 ligand. *J Leukoc Biol* (2000); 67: 2-17.
- 112 Peguet-Navarro J, Dalbiez-Gauthier C, Moulon C, et al. CD40 ligation of human keratinocytes inhibits their proliferation and induces their differentiation. *J Immunol* (1997); 158: 144-152.
- 113 Denfeld RW, Hollenbaugh D, Fehrenbach A, et al. CD40 is functionally expressed on human keratinocytes. *Eur J Immunol* (1996); 26: 2329-2334.
- 114 Gaspari AA, Sempowski GD, Chess P, Gish J, Phipps RP. Human epidermal keratinocytes are induced to secrete interleukin-6 and co-stimulate T lymphocyte proliferation by a CD40-dependent mechanism. *Eur J Immunol* (1996); 26: 1371-1377.

- 115 Pasch MC, Bos JD, Asghar SS. Activation of complement from psoriasis. Clin Exp Dermatol (1998); 23: 189-190.
- 116 Pasch MC, Timar, K, van Meus M, Heyendaal VMR, Bos JD, Laman JD,
5 Ashgar SS. In situ demonstration of CD40- and CD154-positive cells in psoriatic lesions and keratinocyte production of chemokines by CD40 ligation in vitro. J. Pathol (2004); 203:839-848.

111

Claims

1. A polypeptide comprising an amino acid sequence

5	1	11	21	31
	GFSX ₁ S RYSVY WX ₂ RQP PGKGX ₃ EWX ₄ GM MWGGG STDYS			
	41	51	61	
10				
	TSLKS RLTIS KDTSK SQVX ₅ L KMNSL RTDDT AMYYC			
	71			
15	VRTDG DY			

wherein:

	X ₁ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
	X ₂ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
	X ₃ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H;
20	X ₄ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H; and
	X ₅ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H.

2. A polypeptide according to claim 1 comprising an amino acid sequence

25

1	11	21	31

QVKLQ ESGPG LVKPS ETL SI TCTVS GFSX₁S RYSVY WX₂RQP

41 51 61 71
| | | |

5 PGKGX₃ EWX₄GM MWGGG STDYS TSLKS RLTIS KDTSK

81 91 101 111
| | | |
SQVX₅L KMNSL RTDDT AMYYC VRTDG DYWGQ GTTVT VSS.

10

3. A polypeptide according to claim 1 or claim 2, wherein:

X₁ is G, A, V, L, P, F, M, W, C, N, Q, S, T, Y, D, E, K, R or H.

4. A polypeptide according to any one of claims 1-3, wherein:

X₁ is G, A, V, L, P, F or M;

15 X₂ is G, A, V, L, I, P, F or M;

X₃ is G, A, V, L, I, P, F, M;

X₄ is G, A, V, L, I, P, F, M; and

X₅ is G, A, V, L, I, P, F, M, W, C, N, Q, S, T or Y.

5. A polypeptide according to any one of claims 1-4, wherein:

20 X₁ is G, A, V, L or M;

X₂ is G, A, V, L, I or M;

X₃ is G, A, V, L, I, P, F, M;

X₄ is G, A, V, L, I or M; and

X₅ is P, F, W, N, Q, S, T or Y.

25 6. A polypeptide according to claim 5, wherein:

X₁ is L;

X₂ is I;

X₃ is P;

X₄ is M; or X₅ is S.

30 7. A polypeptide according to claim 1 or claim 2, wherein:

X₁ is I and X₂ is V;

X₁ is L and X₂ is I;

X₁ is V and X₂ is V;

X₁ is L and X₂ is L; or

5 X₁ is L and X₂ is V.

8. A polypeptide according to claim 7, wherein

X₃ is P;

X₄ is M; and

X₅ is F or X₅ is S, preferably S;

10 9. A polypeptide according to any one of claims 1-7, wherein:

X₁ is L; X₂ is V; X₃ is L; X₄ is L and X₅ is F.

10. A polypeptide comprising amino acid sequence

15 1 11
 | |
 GFSX₁S RYSVY WX₂R

wherein:

X₁ is L and X₂ is I; or

20 X₁ is I and X₂ is V.

11. A binding body comprising a polypeptide according to any one of claims 1-10.

12. An antibody comprising a polypeptide according to any one of claims 1-10.

13. An antibody according to claim 12, comprising a constant region of a
25 human antibody, preferably an IgG constant region.

14. An antibody according to claim 12 or 13, wherein said constant region is a region that is deficient in complement activation, preferably human IgG₄ constant region or a mutated human IgG₁ constant region.

15. A nucleic acid encoding a polypeptide according to any one of claims 1-10, and/or a binding body according to claim 11, and/or an antibody according to any one of claims 12-14.

16. A cell comprising a polypeptide according to any one of claims 1-10, a
5 binding body according to claim 11, and/or an antibody according to any one of claims 12-14 and/or a nucleic acid according to claim 15.

17. An antagonistic anti-human CD40 monoclonal antibody comprising a polypeptide according to any one of claims 1-10.

18. A deimmunized antagonistic anti-human CD40 monoclonal antibody
10 according to claim 17.

19. An antagonistic anti-human CD40 monoclonal antibody according to claim 17 or claim 18 comprising an amino acid sequence

	1	11	21	31
15	X ₆ LGX ₇ X ₈ ASISC RSSQS LX ₉ NSN GNTYL HWYLQ RPGQS			
	41	51	61	71
	PRLLI YKVSN RFSGV PDRFS GSGSG TDFTL KISRV EAEDX ₁₀			
20				
	81	91		
	GVYX ₁₁ C SQSTH VPWT			

25 wherein:

X₆ is N, Q, S, T, Y, W or C;

X₇ is D, E, N, Q, S, T, Y, W or C;

X₈ is N, Q, S, T, Y, G, A, V, L, I, P, F, M, W or C;

X₉ is G, A, V, L, I, P, F, M;

30 X₁₀ is G, A, V, L, I, P, F, M; and

X₁₁ is N, Q, S, T, Y, G, A, V, L, I, P, F, M, W or C.

20. An antagonistic anti-human CD40 monoclonal antibody according to claim 19 comprising an amino acid sequence

5	1	11	21	31
	ELQLT QSPLS LPVX ₆ L GX ₇ X ₈ AS ISCRS SQSLX ₉ NSNGN TYLHW			
	41	51	61	71
10				
	YLQRP GQSPR LLIYK VSNRF SGVPD RFSGS GSGTD FTLKI			
	81	91	101	111
15				
	SRVEA EDX ₁₀ GV YX ₁₁ CSQ STHVP WTFGG GTKLE IKR			

21. An antibody according to claim 19 or claim 20 wherein

X₆ is T or S, X₇ is D or Q, X₈ is Q or P, X₉ is V or A, X₁₀ is V or L and X₁₁ is F or Y.

20 22. An antibody according to any one of claims 19-21, wherein:

X₆ is T, X₇ is Q, X₈ is P, X₉ is A, X₁₀ is V and X₁₁ is Y.

23. A cell comprising an antibody according to any one of claims 19-22.

24. A cell according to claim 23, that produces said antibody.

25 25. A cell according to claim 16, 23 or claim 24, that is a hybridoma cell, a Chinese hamster ovary cell, an NS0 cell or a PER-C6TM cell.

26. A cell according to claim 25 deposited under number ..., at the ... collection.

27. A cell culture comprising a cell according to any one of claims 23-26.

28. A method for producing an antibody comprising culturing a cell according to any one of claims 16, 23-26 and harvesting said antibody from said culture.

30 29. An antibody obtainable by a method according to claim 28.

30. An antibody according to claim 29 that is purified.

31. A pharmaceutical composition comprising a polypeptide according to any one of claims 1-10, a binding body according to claim 11, an antibody according to any one of claims 12-14, 17-22, 29 or 30, a nucleic acid according to claim 15
5 and/or a cell according to claim 16, 23-26.

32. A polypeptide according to any one of claims 1-10, a binding body according to claim 11, an antibody according to any one of claims 12-14, 17-22, 29 or 30, a nucleic acid according to claim 15 and/or a cell according to claim 16, 23-26 for use as a medicament.

10 33. Use of a polypeptide according to any one of claims 1-10, a binding body according to claim 11, an antibody according to any one of claims 12-14, 17-22, 29 or 30, a nucleic acid according to claim 15 and/or a cell according to claim 16, 23-26 for the manufacture of a medicament for ameliorating a symptom of an autoimmune disorder and/or an inflammatory disorder and/or for reducing
15 graft rejection and/or for the treatment of CD40 positive cancers.

34. Use according to claim 33, wherein said autoimmune and/or an inflammatory disorder is selected from the group of rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, psoriasis, bullous pemphigoides and atopic dermatitis.

20 35. Use according to claim 33, wherein said autoimmune and/or an inflammatory disorder comprises inflammatory bowel disease.

36. Use according to claim 35, wherein said inflammatory bowel disease comprises ulcerative colitis (UC) or Crohn disease (CD).

37. A method for providing an individual with an antibody comprising
25 providing said individual with an antibody according to any one of claims 12-14, 17-22, 29 or 30.

38. A method for selecting an anti-human CD40 antagonist antibody comprising generating a first cell line that produces an original anti-human CD40 antagonist antibody and determining the amount of original antibody

117

that is produced by said first cell line, said original antibody comprising the heavy chain variable domain amino acid sequence

	1	11	21	31
5	QVKLQ ESGPG LVKPS ETL SI TCTVS GFSX ₁ S RYSVY WX ₂ RQP			
	41	51	61	71
	PGKGP EWMGM MWGGG STDYS TSLKS RLTIS KDTSK			

10

	81	91	101	111
	SQVSL KMNSL RTDDT AMYYC VRTDG DYWGQ GTTVT VSS			

wherein X₁ and X₂ are pair wise selected from the group consisting of X₁ = I and X₂ = V; X₁ = L and X₂ = I; X₁ = V and X₂ = V; X₁ = L and X₂ = L; or X₁ = L and X₂ = V,

15

said method further comprising

generating at least one further cell line that produces a variant of said original antibody wherein said variant antibody is a modified original antibody

20 comprising an insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids when compared to said original antibody, wherein said modification does not consist of a modification of the amino acids at the position(s) identified by X₁ and X₂, and determining the amount of variant antibody that is produced by said at least one further cell line,

25 said method further comprising selecting a variant antibody that was produced at an amount that was at least 50% of the amount of original antibody.

39. A method according to claim 38, wherein said original antibody comprises the light chain amino acid sequence

	1	11	21	31
30				

ELQLT QSPLS LPV T L G Q P AS ISCRS SQSL A NSNGN TYLHW

41 51 61 71
| | | |

5 YLQRP GQSPR LLIYK VSNRF SGVPD RFSGS GSGTD FTLKI

81 91 101 111
| | | |

10 SRVEA ED V GV Y Y CSQ STHVP WTFGG GTKLE IKR.

40. A method according to claim 38 or 39, wherein said insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids is in said heavy chain amino acid sequence or said light chain amino acid sequence when compared to the corresponding chain amino acid sequence in said original antibody.

41. A method according to claim 40, wherein said insertion, deletion, inversion, and/or substitution of between about 1 to 5 amino acids is in said heavy chain amino acid sequence when compared to said heavy chain sequence of said original antibody.

20 42. A method according to any one of claims 38-41, further comprising generating an antibody producer cell line producing said selected antibody.

43. A method according to claim 42, further comprising collecting said selected antibody.

25 44. An anti-human CD40 antagonist antibody obtainable by a method according to claim 43.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

Mu5D12 VH region

Q V K L	E E S	G P G	L V A P	S Q S	L S I
10	20	30	40	50	60
CAGGTCAAGC	TCGAGGAGTC	TGGACCTGGC	CTGGTGGCAC	CCTCACAGAG	CCTGTCCATC
GTCCAGTTCTG	AGCTCCTCAG	ACCTGGACCG	GACCACCGTG	GGAGTGTCTC	GGACAGGTAG
T C T V	S G F	S L S	R Y S V	Y W V	R Q P
70	80	90	100	110	120
ACATGCACTG	TCTCTGGGTT	CTCATTATCC	AGATATAGTG	TATACTGGGT	TCGCCAGCCT
TGTACGTGAC	AGAGACCCAA	GAGTAATAGG	TCTATATCAC	ATATGACCCA	AGCGGTCGGA
P G K G	L E W	L G M	M W G G	G S T	D Y N
130	140	150	160	170	180
CCAGGAAAGG	GTCTGGAGTG	GCTGGGAATG	ATGTGGGGTG	GTGGATCCAC	AGACTATAAT
GGTCCTTTCC	CAGACCTCAC	CGACCCTTAC	TACACCCAC	CACCTAGGTG	TCTGATATTA
S A L K	S R L	S I S	K D T S	K S Q	V F L
190	200	210	220	230	240
TCAGCTCTCA	AATCCAGACT	GAGCATCAGC	AAGGACACCT	CGAAGAGCCA	GGTCTTCTTA
AGTCGAGAGT	TTAGGTCTGA	CTCGTAGTCG	TTCCTGTGGA	GCTTCTCGGT	CCAGAAGAAT
K M N S	L R T	D D T	A M Y Y	C V R	T D G
250	260	270	280	290	300
AAAATGAACA	GTCTGCGAAC	TGATGACACA	GCCATGTACT	ACTGTGTCAG	AACCGATGGG
TTTACTTGT	CAGACGCTTG	ACTACTGTGT	CGGTACATGA	TGACACAGTC	TTGGCTACCC
D Y W G	Q G T	S V T	V S S		
310	320	330	339		
GACTACTGGG	GTCAAGGAAC	CTCAGTCACC	GTCTCCTCA		
CTGATGACCC	CAGTTCCTTG	GAGTCAGTGG	CAGAGGAGT		

Mu5D12 VL region

E L Q L	T Q S	P L S	L P V S	L G D	Q A S
10	20	30	40	50	60
GAGCTCCAGC	TGACCCAGTC	TCCACTCTCC	CTGCCTGTCA	GTCTTGAGGA	TCAAGCCTCC
CTCGAGGTCTG	ACTGGGTCAG	AGGTGAGAGG	GACGGACAGT	CAGAACCTCT	AGTTCGGAGG
I S C R	S S Q	S L V	N S N G	N T Y	L H W
70	80	90	100	110	120
ATCTCTTGCA	GATCTAGTCA	GAGCCTTGTA	AACAGTAATG	GAAACACCTA	TTTACATTGG
TAGAGAACGT	CTAGATCAGT	CTCGGAACAT	TTGTCATTAC	CTTTGTGGAT	AAATGTAACC
Y L Q K	P G Q	S P K	L L I Y	K V S	N R F
130	140	150	160	170	180
TACCTGCAGA	AGCCAGGCCA	GTCTCCAAAG	CTCCTGATCT	ACAAAGTTTC	CAACCGATTT
ATGGACGTCT	TGGTCCGGT	CAGAGGTTTC	GAGGACTAGA	TGTTTCAAAG	GTTGGCTAAA
S G V P	D R F	S G S	G S G T	D F T	L K I
190	200	210	220	230	240
TCTGGGGTCC	CAGACAGGTT	CAGTGGCAGT	GGATCAGGGA	CAGATTTTAC	ACTCAAGATT
AGACCCCAGG	GTCTGTCCAA	GTCACCGTCA	CCTAGTCCCT	GTCTAAAGTG	TGAGTTCTAA
S R V E	A E D	L G V	Y F C S	Q S T	H V P
250	260	270	280	290	300
AGCAGAGTGG	AGGCTGAGGA	TCTGGGAGTT	TATTTCTGCT	CTCAAAGTAC	ACATGTTCCG
TCGTCTCACC	TCCGACTCCT	AGACCCCTCAA	ATAAAGACGA	GAGTTTCATG	TGTACAAGGC
W T F G	G G T	K L E	I K R		
310	320	330	339		
TGGACGTTCTG	GTGGAGGCAC	CAAGCTGGAA	ATCAAACGT		
ACCTGCAAGC	CACCTCCGTG	GTTCGACCTT	TAGTTTGCA		

Figure 6





Figure 7

12 variants DI5D12VH


1.	Q5E	1	QVKLEESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
2.	K13A	1	QVKLQESGPGGLVAPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
3.	E16Q	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
4.	T17S	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
5.	I29L	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
6.	I37V	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
7.	P45L	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
8.	M48L	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWLGMMWGGGSTDYS	60
9.	ST86NSA	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYN	60
10.	T68S	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
11.	S79F	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
12.	T108S	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
	CH5D12	1	QVKLEESGPGGLVAPSQSLSITCTVSGFSLSRYSVYWIROPFGKGPPEWLGMMWGGGSTDYN	60
	DI5D12	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYWIROPFGKGPPEWMGMMWGGGSTDYS	60
1.	Q5E	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
2.	K13A	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
3.	E16Q	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
4.	T17S	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
5.	I29L	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
6.	I37V	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
7.	P45L	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
8.	M48L	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
9.	ST86NSA	61	SALKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
10.	T68S	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
11.	S79F	61	TSLKSRLLTISKDTSKSQVFLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113
12.	T108S	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTSTVTVSS	113
	CH5D12	61	SALKSRLLTISKDTSKSQVFLKMNSLRTDDTAMYYCVRTDGDYWGQGTSTVTVSS	113
	DI5D12	61	TSLKSRLLTISKDTSKSQVSLKMNSLRTDDTAMYYCVRTDGDYWGQGTTVTVSS	113

8/21

Figure 8 FACS data
Q5E


% Gated	Mean	Median
100.00	39.05	37.52

K13A




% Gated	Mean	Median
100.00	31.22	29.96

E16Q




% Gated	Mean	Median
100.00	39.48	38.20

T17S




% Gated	Mean	Median
100.00	33.43	31.91

I29L



% Gated	Mean	Median
100.00	108.28	91.81


I37V



% Gated	Mean	Median
100.00	96.05	82.05


9/21

P45L




% Gated	Mean	Median
100.00	82.53	74.32

M48L




% Gated	Mean	Median
100.00	80.30	72.34

STS60NSA




% Gated	Mean	Median
100.00	62.37	58.82

T68S




% Gated	Mean	Median
100.00	35.21	33.98

S79F



% Gated	Mean	Median
100.00	96.40	82.05

T108S




% Gated	Mean	Median
100.00	99.04	85.82


10/21

Ch5D12

DI5D12




% Gated	Mean	Median
100.00	102.97	88.96




% Gated	Mean	Median
100.00	50.44	47.40

Mock Control

FACS control



% Gated	Mean	Median
100.00	11.25	8.74



% Gated	Mean	Median
100.00	10.18	8.74

Figure 9: V-L-I variants of DI5D12VH

1.	IV	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYVVRQPPGKGPEWMGMMWGGGSTDYSTS	62
2.	VI	1	QVKLQESGPGGLVKPSETLSITCTVSGFVSRYSVYVIRQPPGKGPEWMGMMWGGGSTDYSTS	62
3.	IL	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYVLRQPPGKGPEWMGMMWGGGSTDYSTS	62
4.	VV	1	QVKLQESGPGGLVKPSETLSITCTVSGFVSRYSVYVVRQPPGKGPEWMGMMWGGGSTDYSTS	62
5.	LL	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYVLRQPPGKGPEWMGMMWGGGSTDYSTS	62
6.	VL	1	QVKLQESGPGGLVKPSETLSITCTVSGFVSRYSVYVLRQPPGKGPEWMGMMWGGGSTDYSTS	62
7.	LV	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYVVRQPPGKGPEWMGMMWGGGSTDYSTS	62
CH5D12	(PG100)	1	QVKLEESGPGGLVAPSQSLSITCTVSGFSLSRYSVYVVRQPPGKGLEWLCMMWGGGSTDYNBSA	62
DI5D12	(II)	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYVIRQPPGKGPEWMGMMWGGGSTDYSTS	62
PG102	(LI)	1	QVKLQESGPGGLVKPSETLSITCTVSGFSLSRYSVYVIRQPPGKGPEWMGMMWGGGSTDYSTS	62
1.	IV	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
2.	VI	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
3.	IL	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
4.	VV	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
5.	LL	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
6.	VL	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
7.	LV	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
CH5D12	(PG102)	63	LKSRLSISKDTSKSQVFLKMNSLRDDETAMYYCVRTDGDYWGQGTISVTVSS	113
DI5D12	(II)	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113
PG102	(LI)	63	LKSRLTISKDTSKSQVSLKMNSLRDDETAMYYCVRTDGDYWGQGTITVTVSS	113

Figure 10: Distribution of antibody productivity for GS-CHO cell lines producing the PG102 antibody and 'overgrown' 24-well plate cultures




Figure 11: Growth and antibody accumulation profiles for cell lines L107 (DC1; top panel), L25 (DC2; middle panel) and M95 (DC3; lower panel) in CDACF fed-batch shake flask culture





Figure 12: Comparison of the binding of PG102(red) and PG100(blue) at 10µg/mL to CD40 immobilised at 3 different levels






Figure 13 Inhibition of CD40 mAb binding to JY cells expressing human CD40 by the anti-idiotypic mAb 173-36-1. Anti-CD40 mAbs were assayed at a concentration of 1 μ g/ml. Data indicate mean $\pm$ s.e. mean of 4 separate determinations for each antibody.




Figure 14: CD40-Fc ELISA. Concentration-related binding of PG100 and PG102 to human CD40-Fc. Plates were coated overnight with 250 ng/ml human CD40-Fc. The isotype control mAb, chFUN-1, is directed against human CD86. Data indicate mean $\pm$ s.e. mean (all n=3).




Figure 15: FACS quantification of anti-CD40 mAb binding to JY cells. PG100 and PG102 displayed comparable half-maximal binding concentrations. The anti-CD86 mAb, chFUN-1, also showed binding to JY cells in these experiments due to the surface expression of human CD86 by this cell line. Data indicate the mean of duplicate determinations for each antibody.




Figure 16. Inhibition of PG100PE and PG102PE binding to JY cells by unlabelled PG100 and PG102 antibodies. Labelled PG100 (A) and PG102 (B) were incubated at 1 μ g/ml in the presence of increasing concentration of competing unlabelled antibody. Labelled antibody binding was determined as mean fluorescence intensity (MFI) by flow cytometry. Maximal MFI for PG100PE and PG102PE binding was 369 fluorescence units (A) and 305 fluorescence units (B), respectively. Data indicate mean $\pm$ s.e. mean of 4 separate experiments.

Figure 17: Inhibition of IL-8 release by THP-1 cells after co-culture with Jurkat cells.




Figure 18

Protein sequence of PG102 heavy and light chain

Signal sequences are underlined, variable regions in bold.

Heavy chain:

MEWSWVFLFFLSVTTGVHS**QVKLQESGPGLVKPSETLSITCTVSGFSLSRYSVIWIRQPPGKGPE**
WMGMMWGGGSTDYSTSLKSRLTISKDTSKSQVSLKMNSLRDDTAMYYCVRTDGDYWGQGTTVTV
SSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY
SLSSVVTVPSSSLGTKYTCNVDHKPSNTKVDKRVESKYGPPCPSCPAPEFLGGPSVFLFPPKPK
DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDW
LNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIA
VEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSL
SLGK

Light chain:

MSVPTQVLGLLLLWLTDARCELQLTQSPLSLPVTLGQPASISCRSSQSLANSNGNTYLHWYLQRP
GQSPRLLIYKVSNRFSGVDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSTHVPWTFGGGTKLE
IKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKD
STYLSSTLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC

DNA sequences encoding PG102 heavy and light chains

Signal sequences is indicated as underlined, variable regions in bold, and stop codons in lower case.

Heavy chain:

ATGGAGTGGTCTTGGGTGTTCCCTGTTCTTCCTGTCTGTGACAACAGGAGTGCACCTCTCAGGTCAA
GCTGCAGGAGTCTGGACCAGGACTGGTGAAGCCATCTGAGACCCTGAGCATCACCTGTACAGTGA
GCGGCTTCAGCCTCTCTAGATACAGCGTGTACTGGATCAGACAGCCACCTGGAAAGGGACCAGAG
TGGATGGGAATGATGTGGGGAGGAGGATCTACAGACTACAGCACCAGCCTGAAGTCTAGACTGAC
CATCAGCAAGGACACCTCTAAGTCTCAGGTCTCCCTGAAGATGAACCTCTCTGAGAACAGACGACA
CCGCCATGTACTACTGTGTGAGAACCAGCGAGATTATTGGGGACAGGGCACAACAGTGACAGTG
TCCTCTGCCTCTACAAAGGGACCATCTGTGTTTCCACTGGCCCCATGTTCTAGATCTACCAGCGA
GTCTACAGCTGCTCTGGGATGTCTGGTGAAGGACTACTTTCAGAGCCTGTGACAGTGTCTTGGA
ATAGTGGAGCCCTGACATCTGGAGTGCACACATTTCCAGCTGTGCTGCAGTCTAGCGGACTGTAT
TCTCTGTCCAGCGTGGTGACAGTGCCATCTTCTTCTCTGGGCACCAAGACCTACACATGTAACGT
GGACCACAAGCCATCTAACACCAAGGTGGACAAGAGAGTGGAGTCTAAGTACGGACCACCATGCC
CATCTTGTCCAGCTCCAGAGTTTCTGGGAGGACCTAGCGTGTTTCTGTTCCCCCAAAGCCAAAG
GATACCCCTGATGATCTCTAGAACCCAGAGGTGACATGTGTGGTGGTGGATGTGTCTCAGGAGGA
TCCAGAGGTCCAGTTTAACTGGTACGTGGATGGAGTGGAGGTGCACAACGCTAAGACAAAGCCAA
GAGAGGAGCAGTTCAACAGCACATACAGAGTGGTGTCTGTGCTGACAGTGCATCAGGATTGG
CTGAACGGCAAGGAATACAAGTGTAAAGTCTCCAACAAGGGCCTGCCATCTTCTATCGAGAAAAC
CATCTCTAAGGCTAAGGGACAGCCAAGGGAGCCACAGGTGTACACACTGCCACCATCTCAAGAGG
AGATGACCAAGAACCAGGTGTCCCTGACATGCCTGGTGAAGGGATTCTACCCATCTGATATCGCT
GTGGAGTGGGAGTCTAATGGACAGCCCAGAACTACAAGACCACACCACAGTGTCTGGATTCT
TGACGGCTCTTTCTTCTGTACAGCAGACTGACAGTGGACAAGTCTAGATGGCAGGAGGGAAACG
TCTTTAGCTGTAGCGTGATGCATGAGGCTCTGCACAACCACTACACCCAGAAGTCTCTGTCTCTG
AGTCTGGGAAAAtgatatag

Light chain:

ATGTCTGTGCCAACACAGGTGCTGGGACTGCTGCTGCTGTGGCTGACAGATGCTAGATGTGAGCT
GCAGCTGACACAGTCTCCACTGTCTCTGCCAGTGACACTGGGACAGCCAGCTAGCATCAGCTGTA
GAAGCTCTCAGTCTCTGGCCAACTCTAACGGCAACACATACCTGCATTGGTATCTGCAGAGACCA
GGACAGTCTCCAAGACTGCTGATCTACAAGGTGTCCAACAGATTCTCTGGAGTGCCAGACAGATT
TTCTGGCTCTGGCTCTGGAACAGACTTCACCCTGAAGATCTCTAGAGTGGAGGCTGAGGATGTGG
GAGTGTACTACTGCTCTCAGTCTACACATGTGCCATGGACATTTCGGAGGAGGAACAAAGCTGGAG
ATCAAGAGAACAGTGGCTGCCCCATCTGTGTTTATCTTCCCCCATCTGATGAGCAGCTGAAGTC
TGGAACAGCTTCTGTGGTGTGTCTGCTGAACAACCTTCTACCCAAGGGAGGCTAAGGTGCAGTGGA
AGGTGGACAATGCTCTGCAGTCTGGAACTCTCAGGAGTCTGTCTACAGAGCAGGACAGCAAGGAC
TCTACCTACTCTCTGAGCAGCACACTGACACTGTCTAAGGCCGACTACGAGAAGCATAAGGTGTA
CGCCTGTGAGGTGACACATCAGGGACTGTCTAGCCCAGTGACCAAGTCTTTCAACAGAGGCGAGT
GCTgatatag

INTERNATIONAL SEARCH REPORT

International application No
PCT/NL2007/050202

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K16/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KASRAN A ET AL: "Safety and tolerability of antagonist anti-human CD40 Mab ch5D12 in patients with moderate to severe Crohn's disease." ALIMENTARY PHARMACOLOGY & THERAPEUTICS 15 JUL 2005, vol. 22, no. 2, 15 July 2005 (2005-07-15), pages 111-122, XP002416781 ISSN: 0269-2813 page 119, left-hand column	10-44
Y	WO 98/52976 A (BIOVATION LTD [GB]; CARR FRANCIS J [GB]) 26 November 1998 (1998-11-26) examples 1-6	10-44

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

12 October 2007

Date of mailing of the international search report

24/10/2007

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Le Flao, Katell

INTERNATIONAL SEARCH REPORT

International application No
PCT/NL2007/050202

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>BOON L ET AL: "Preclinical assessment of anti-CD40 Mab 5D12 in cynomolgus monkeys" TOXICOLOGY, LIMERICK, IR, vol. 174, 2002, pages 53-65, XP002954468 ISSN: 0300-483X</p> <p>page 60, left-hand column, paragraph 2 - page 61, right-hand column, line 2 page 64, left-hand column, paragraph 2 - right-hand column, paragraph 1</p>	10-44
A	<p>ROQUE-NAVARRO LOURDES ET AL: "Humanization of predicted T-cell epitopes reduces the immunogenicity of chimeric antibodies: New evidence supporting a simple method." HYBRIDOMA AND HYBRIDOMICS, vol. 22, no. 4, August 2003 (2003-08), pages 245-257, XP002397469 ISSN: 1536-8599 the whole document</p>	10-44

INTERNATIONAL SEARCH REPORT

International application No.
PCT/NL2007/050202

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-9
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.2

Claims Nos.: 1-9

The present claims 1-9 relate to an extremely large number of possible compounds. Support and disclosure in the sense of Article 6 and 5 PCT is to be found however for only a very small proportion of the compounds claimed, see example 2 from page 40 to page 46 (a deimmunized antibody 5D12 and 12 variants of it have been cloned). The non-compliance with the substantive provisions is to such an extent, that a meaningful search of the whole claimed subject-matter of the claim could not be carried out (PCT Guidelines 9.19 and 9.23). The extent of the search was consequently limited.

The search of claim 1-9 was restricted to those claimed compounds which appear to be supported and a generalisation of their structural formulae, thus to deimmunized anti human CD40 monoclonal antibodies.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/NL2007/050202

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9852976	A	26-11-1998	
		AT 319745 T	15-03-2006
		AU 736549 B2	02-08-2001
		AU 7539398 A	11-12-1998
		CA 2290485 A1	26-11-1998
		DE 69833755 T2	28-12-2006
		EP 0983303 A1	08-03-2000
		GB 2339430 A	26-01-2000
		JP 2002512624 T	23-04-2002
<hr/>			