



US00PP07869P

United States Patent [19]

Izsak et al.

[11] Patent Number: Plant 7,869

[45] Date of Patent: May 19, 1992

[54] STRAWBERRY PLANT DORIT

[75] Inventors: Eva Izsak; Shamai Izhar, both of Rehovot, Israel
[73] Assignee: State of Israel, Ministry of Agriculture, The Volcani Center, Bet Dagan, Israel

[21] Appl. No.: 735,968

[22] Filed: Jul. 25, 1991

Related U.S. Application Data

[63] Continuation of Ser. No. 489,440, Mar. 6, 1990, abandoned.

[51] Int. Cl.⁵ A01H 5/00

[52] U.S. Cl. Plt./48

[58] Field of Search Plt./48, 49

[56] References Cited

U.S. PATENT DOCUMENTS

P.P. 4,487 11/1979 Bringhurst et al. Plt. 49
P.P. 5,262 7/1984 Voth et al. Plt. 48
P.P. 5,263 7/1984 Voth et al. Plt. 48
P.P. 5,264 7/1984 Bringhurst et al. Plt. 48
P.P. 5,268 8/1984 Voth et al. Plt. 48
P.P. 6,578 1/1989 Voth et al. Plt. 48
P.P. 7,024 9/1989 Johnson et al. Plt. 49
P.P. 7,127 2/1990 Voth et al. Plt. 49

1

This is a continuation of application Ser. No. 07/489,440 filed Mar. 6, 1990, now abandoned.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety of strawberry (*Fragaria L.*) called "Dorit". The variety was developed from an organized scientifically designated breeding program carried out at the Agricultural Research Organization, the Volcani Center, Bet Dagan, Israel. The variety is the product of selection of seedlings resulting from crosses between the strawberry varieties "Dover A" and "Nurit". The variety was asexually vegetatively propagated through runners and the reproduction ran true.

SUMMARY OF THE INVENTION

The new variety "Dorit" is able to grow in September and produce fruit starting in November and lasting until summer. The production of fruit beginning in November is two months earlier than any known variety of *Fragaria L.* The fruit of the "Dorit" variety is characterized by good taste, good shape and size as well as a long shelf life.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Photograph of the "Dorit" variety illustrating the foliage and fruit.

FIG. 2. Photograph of the "Dorit" variety illustrating the fruit.

FIG. 3. Photograph of the "Dorit" variety illustrating the entire plant with both flowers and fruit.

FOREIGN PATENT DOCUMENTS

370/82 11/1984 Israel .

OTHER PUBLICATIONS

Howard, C. M. and Albrechts, E. E., 1980; HortScience 15(4):540.

Dubinsky, V., 1985; Mc.Sc. Thesis, The Hebrew University of Jerusalem; Rehovot, Israel (Hebrew original and English translation).

Bringhurst, R. S. and Voth, V., 1989; Fruit Var. J 43(1):12-19.

Izsak, E. (1978) Hassadeh 59:443-446.

Izsak, E. and Izhar, S. (1983) Hassadeh 63:2100-2103.

Izsak, E. and Izhar, S. (1984) Hassadeh 64:1774-1777.

Izsak, E. and Izhar, S. (1984) Hassadeh, 64:1778-1781.

Primary Examiner—James R. Feyrer

Attorney, Agent, or Firm—Pennie & Edmonds

[57]

ABSTRACT

A new and distinct variety of strawberry (*Fragaria L.*) called "Dorit" is disclosed. The variety is a cross between "Dover A" and "Nurit", which results in a variety that flowers several months earlier than other known strawberry varieties.

2 Drawing Sheets

2

DETAILED BOTANICAL DESCRIPTION OF THE INVENTION

The "Dorit" variety was grown in winter under polyethylene tunnels in Israel. "Dorit" is a short day variety that flowers earlier than other known and available short day length strawberry varieties. Flowering and fruit production is not affected by the use of polyethylene wind tunnels. This production procedure is utilized in normal agricultural practices by the skilled artisan and does not involve temperature or light control. Mother plants were stored at 0° C. from January through April. They were then planted in the nursery without further treatment. Runners with plantlets were produced during summer. These young plantlets were collected from the nursery in September and transferred to raised beds. Average temperatures at that time of the year are 30° C. during the day and 22° C. at night. Water and fertilizers were applied through drip irrigation. An example of an optimum planting date is between September 20th to the 25th with the approximate date of flowering on October 27, and the approximate date of first fruiting on November 27. "Dorit" flowering is not induced by chilling, but by natural exposure to short day length (long nights) characteristic of late fall and early winter. Color readings described herein were taken under natural light conditions and color identifications were made by reference to The Royal Horticultural Society Colour Chart (R.H.S.C.C.) except where common terms of color definition are employed.

The pertinent characteristics of the present invention are presented in Table 1 and Table 2. Additionally, the

TABLE 1-continued

MORPHOLOGICAL TRAIT	PLANT CHARACTERISTICS OF "DORIT"		COMPARABLE VARIETY ^b			
	DESCRIPTION ^a	DESCRIPTION ^a				
Pistil	Normal					
^a The description of "Dorit" is based on the test guidelines for <i>Fragaria</i> L. of the International Union for the Protection of New Plant Varieties. (UPOV).						
^b Only characteristics which are relevant for comparing varieties are listed. For example, there are no varietal differences acknowledged in the characteristic "color of lower side of leaf".						
10	The flower size of the "Dorit" variety is also larger than flowers of the variety "Gorella".					
15	"Dorit" flowers at the end of October. One of the earliest known varieties for comparison is "Karina", which flowers in January.					
20	"Dorit" still produces more runners than the comparable variety, "Gorella".					
TABLE 2						
FRUIT CHARACTERISTICS OF "DORIT" ^a		DESCRIPTION				
CHARACTERISTIC		DESCRIPTION				
Time of ripening		Early				
Ratio of length/maximum width		Longer than broad				
Primary Fruit ^b		Bi-conical				
Length		43-50 mm				
Width		30-36 mm				
Secondary Fruit		Conical				
Length		33-37 mm				
Width		27-32 mm				
Tertiary Fruit		Ovoid				
length		28-30 mm				
width		25-26 mm				
Size		Large				
Band without achenes		Narrow				
Unevenness of surface		Weak				
Color		Orange red				
Evenness of color		Even				
Glossiness		Medium				
Insertion of achenes		Level with surface				
Insertion of calyx		Level with surface				
Pose of calyx segments		Reflexed				
Size of calyx in relation to fruit diameter		Same size				
Adherence of calyx		Strong				
Firmness		Medium to firm				
Color of Flesh		Light red				
Evenness of flesh color		Even				
Sweetness ^c		Strong				
Color		42 BC circa (RHSCC)				
Taste ^c		Good				
^a The shape of "Dorit" fruit is similar to that of the variety "Pajaro".						
45	^b There is a marked difference between the shape of the primary, secondary and tertiary fruit.					
50	^c See Table 4.					
TABLE 3						
COMPARATIVE YIELD OF "DORIT" ^a						
	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.
Dorit	30	70	100	100	100	100
Sharon ^b	40	60	80	80	80	80
55	50	70	80	100	100	70
Shalom ^c	50	70	100	100	100	50
Smadar ^d	0	70	100	100	100	100
Saaid ^e	0	0	40	150	150	150
Douglas ^f	0	0	30	150	150	120
Chandler ^g						
^a Average yield in g/m ² in Ramat Hadar, Israel (1989-90). The time of ripening for "Dorit" fruit is early.						
60	^b U.S. Application Ser. No. 07/735,969. The time of ripening for "Sharon" fruit is very early.					
65	^c U.S. Application Ser. No. 07/735,695. The time of ripening for "Shalom" fruit is early.					
	^d U.S. Application Ser. No. 07/735,967. The time of ripening for "Smadar" fruit is very early to early.					
	^e U.S. Application Ser. No. 07/735,970. The time of ripening for "Saaid" fruit is early to medium.					
	^f U.S. Plant Pat. #4,487. The time of ripening for "Douglas" fruit is late.					
	^g U.S. Plant Pat. #5,262. The time of ripening for "Chandler" fruit is late.					

Plant 7,869

5

TABLE 4

COMPARATIVE FRUIT CHARACTERISTICS OF "DORIT"

	T.S.S. ^a in %	Acidity ^b in %	Aroma	Taste
Dorit	8.5-9.5	1.0	5	Good
Sharon ^c	6.5-7.0	1.0	3	Slightly Acidic
Shalom ^d	8.0-9.0	1.0	4	Normal
Smadar ^e	8.5-9.5	1.0	5	Good
Saaid ^f	8.0-9.0	1.0	3	Normal
Douglas ^g	6.5-7.0	0.8	3	Slightly Acidic
Chandler ^h	6.5-7.0	0.8	3	Slightly

6

TABLE 4-continued

5	COMPARATIVE FRUIT CHARACTERISTICS OF "DORIT"			
	T.S.S. ^a in %	Acidity ^b in %	Aroma	Taste

^aTotal Soluble Solids expresses fruit sweetness and was determined with a refractometer.

^bPercent of acidity was determined as follows: 2 cc of juice extract was mixed with 20 cc of water. Five drops of fenolthaleinen was added and the mixture was titrated with NaOH. The percent acidity is calculated as the quantity of NaOH (cc) \times 0.32.

^cU.S. Application Ser. No. 07/735,969.

^dU.S. Application Ser. No. 07/735,695.

^eU.S. Application Ser. No. 07/735,967.

^fU.S. Application Ser. No. 07/735,970.

^gU.S. Plant Pat. #4,487.

^hU.S. Plant Pat. #5,262.

10 15 What is claimed is:

1. A new distinct variety of strawberry substantially as illustrated and described and distinguished as being able to grow in September and produce fruit starting in November and lasting until summer, with fruit having a good taste and shape and a long shelf life.

* * * * *

25

30

35

40

45

50

55

60

65




FIG. 1




FIG. 2

U.S. Patent

May 19, 1992

Sheet 2 of 2

Plant 7,869



FIG. 3