



(86) Date de dépôt PCT/PCT Filing Date: 2001/04/06
(87) Date publication PCT/PCT Publication Date: 2001/10/18
(45) Date de délivrance/Issue Date: 2012/06/19
(85) Entrée phase nationale/National Entry: 2002/10/02
(86) N° demande PCT/PCT Application No.: US 2001/011294
(87) N° publication PCT/PCT Publication No.: 2001/077354
(30) Priorité/Priority: 2000/04/07 (US60/196,001)

(51) Cl.Int./Int.Cl. *C07K 14/415* (2006.01),
A01H 5/00 (2006.01), *C12N 15/29* (2006.01),
C12N 15/82 (2006.01), *C12N 5/10* (2006.01)

(72) Inventeurs/Inventors:
DA COSTA E SILVA, OSWALDO, US;
BOHNERT, HANS J., US;
VAN THIELEN, NOCHA, US;
CHEN, RUOYING, US;
SARRIA-MILLAN, RODRIGO, US

(73) Propriétaire/Owner:
BASF PLANT SCIENCE GMBH, DE

(74) Agent: ROBIC

(54) Titre : PROTEINES ASSOCIEES AU STRESS DU CYCLE CELLULAIRE ET METHODES D'UTILISATION DANS
LES PLANTES

(54) Title: CELL CYCLE STRESS-RELATED PROTEINS AND METHODS OF USE IN PLANTS

(57) Abrégé/Abstract:

A transgenic plant transformed by a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid sequence in the plant results in increased tolerance to environmental stress as compared to a wild type variety of the plant. Also provided are agricultural products, including seeds, produced by the transgenic plants. Also provided are isolated CCSRPs, and isolated nucleic acid coding CCSRPs, and vectors and host cells containing the latter.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 October 2001 (18.10.2001)

PCT

(10) International Publication Number
WO 01/077354 A3

- (51) **International Patent Classification⁷:** **C12N 15/82**, A01H 5/00, 5/10, C07K 14/415, C12N 5/10, 15/29
- (74) **Agents:** COBERN, Lisa, M. et al.; Sutherland Asbill & Brennan, LLP, 999 Peachtree Street, N.E., Atlanta, GA 30309-3996 (US).
- (21) **International Application Number:** PCT/US01/11294
- (22) **International Filing Date:** 6 April 2001 (06.04.2001)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
60/196,001 7 April 2000 (07.04.2000) US
- (81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) **Applicant (for all designated States except US):** **BASF PLANT SCIENCE GMBH** [DE/DE]; 67056 Ludwigshafen (DE).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **COSTA E SILVA, Oswaldo da** [BR/US]; 203 Littleford Lane, Apex, NC 27502 (US). **BOHNERT, Hans, J.** [US/US]; 2514 Pembroke Point, Champaign, IL 61820 (US). **VAN THIELEN, Nocha** [CA/US]; 206-T Hyde Park Court, Cary, SC 27513 (US). **CHEN, Ruoying** [CN/US]; 105 Rustic Pine Court, Apex, NC 27502 (US). **SARRIA-MILLAN, Rodrigo** [CO/US]; 2324 Winter Walk Circle, Morrisville, NC 27560 (US).
- Published:**
— with international search report
- (88) **Date of publication of the international search report:**
12 September 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) **Title:** CELL CYCLE STRESS-RELATED PROTEINS AND METHODS OF USE IN PLANTS

(57) **Abstract:** A transgenic plant transformed by a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid sequence in the plant results in increased tolerance to environmental stress as compared to a wild type variety of the plant. Also provided are agricultural products, including seeds, produced by the transgenic plants. Also provided are isolated CCSRPs, and isolated nucleic acid coding CCSRPs, and vectors and host cells containing the latter.



WO 01/077354 A3

CELL CYCLE STRESS-RELATED PROTEINS AND METHODS OF USE IN PLANTS

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention relates generally to nucleic acid sequences encoding proteins that are associated with abiotic stress responses and abiotic stress tolerance in plants. In particular, this invention relates to nucleic acid sequences encoding proteins that confer drought, cold, and/or salt tolerance to plants.

Background Art

[0002] Abiotic environmental stresses, such as drought stress, salinity stress, heat stress, and cold stress, are major limiting factors of plant growth and productivity. Crop losses and crop yield losses of major crops such as rice, maize (corn) and wheat caused by these stresses represent a significant economic and political factor and contribute to food shortages in many underdeveloped countries.

[0003] Plants are typically exposed during their life cycle to conditions of reduced environmental water content. Most plants have evolved strategies to protect themselves against these conditions of desiccation. However, if the severity and duration of the drought conditions are too great, the effects on plant development, growth and yield of most crop plants are profound. Furthermore, most of the crop plants are very susceptible to higher salt concentrations in the soil. Continuous exposure to drought and high salt causes major alterations in the plant metabolism. These great changes in metabolism ultimately lead to cell death and consequently yield losses.

[0004] Developing stress-tolerant plants is a strategy that has the potential to solve or mediate at least some of these problems. However, traditional plant breeding strategies to develop new lines of plants that exhibit resistance (tolerance) to these types of stresses are relatively slow and require specific resistant lines for crossing with the desired line. Limited germplasm resources for stress tolerance and incompatibility in crosses between distantly related plant species represent significant problems encountered in conventional breeding. Additionally, the cellular processes leading to drought, cold and salt tolerance in model, drought- and/or salt-tolerant plants are complex in nature and involve multiple mechanisms of cellular adaptation and numerous metabolic pathways. This multi-component nature of stress tolerance has not only made breeding for tolerance largely unsuccessful, but has also limited the ability to genetically engineer stress tolerance plants using biotechnological methods.

[0005] Therefore, what is needed is the identification of the genes and proteins involved in these multi-component processes leading to stress tolerance. Elucidating the function of genes expressed in stress tolerant plants will not only advance our understanding of plant adaptation and tolerance to environmental stresses, but also may provide important information for designing new strategies for crop improvement.

[0006] One group of proteins that seems to be associated with stress responses in plants and animals are proteins related to cell division and the cell cycle. Evidence of this relationship is found in the fact that different environmental stresses related to soil water content, incident light and varying leaf temperature all result in a change in the cell division rate. For example, in sunflower leaves, there is a lengthening of the cell cycle associated with stress and aging. Cells in the sunflower leaves tend to arrest at the G1 phase of cell cycle, with no change in the duration of the S-G2-M phases of the cell cycle. Similar observations have been made in other plant species placed under environmental stresses such as sucrose starvation (Van't Hof 1973 Bookhaven Symp. 25:152), oxidative stress (Reichheld et al. 1999 Plant J. 17:647), and water depletion (Schuppler et al. 1998 Plant Physiol. 117:667). These observations and results suggest that there is an important "checkpoint" in the regulation of the cell cycle at the G1-S transition, while other studies suggest another "checkpoint" at the G2-M transition with an arrest of the cells at the G2 phase. It has also been determined that temperature affects the duration of all cell cycle phases by a similar proportion without the preferential arrest in any particular phase of the cell cycle (Tardieu and Granier, 2000 Plant Mol. Biol. 43:555). Although these aforementioned responses to

environmental stresses differ, each of these results demonstrates the interconnectivity of the stress response system of these plants and the cell division proteins therein.

[0007] The prior art describes several proteins associated with cell division and the cell cycle. One such protein, or protein complex, is the cyclin-CDK complex, which controls progression of the cell cycle phases. Cyclin-dependent protein kinases (CDKs) require cyclin binding for activity and are now widely recognized players at the checkpoints of the eukaryotic cell cycle. The widespread importance of CDKs was realized nearly ten years ago from independent genetic approaches in yeast and biochemical studies of mitotic controls in fertilized eggs of marine invertebrates.

[0008] One known component of a CDK complex is a Cdc2 protein termed P34^{cdc2}, which is required at both control points (G1-S and G2-M). Several other Cdc2 homologs have been isolated from human and plant species including yeast. One such yeast homolog is Cdc48, which plays a role in the spindle pole body separation in *Saccharomyces cerevisiae*. Another Cdc2 homolog has been described in *Arabidopsis* (Feiler et al. 1995 EMBO J. 14:5626) that is highly expressed in the proliferating cells of the vegetative shoot, root, floral inflorescence and flowers and in rapidly growing cells. The *Arabidopsis* Cdc48 gene is up regulated in the developing microspores and ovules and down regulated in most differentiated cell types. In addition, this gene has been localized to the nucleus and during cytokinesis to the fragmoplast.

[0009] Another group of proteins involved in cell division and the cell cycle are pRB proteins. Growing evidence suggests that pRB-like proteins in plants might be among the nuclear targets of plant CDKs. In mammals, the pRB is central to the regulation of the G1-to-S transition. Phosphorylation of mammalian pRB by cyclin D- and cyclin E-dependent kinases renders the pRB inactive and thereby represses the S phase and promotes DNA replication. Significantly, the pRB-binding motif LXCXE (where X denotes any amino acid) is found in all known plant D cyclins. In plants, LXCXE-dependent interactions between D cyclins from *Arabidopsis* and maize pRB proteins have been demonstrated *in vitro* and in a yeast two-hybrid assay. The role of pRBs in plant cell division related signaling cascades is further supported by the fact that several pRBs, and in particular, the maize pRB, contain multiple putative CDK phosphorylation sites and are efficiently phosphorylated *in vitro* by mammalian G1- and S-specific CDKs. Maize pRB proteins are also known to undergo changes in phosphorylation during the transition to endoreduplication in the endosperm. However, phosphorylation by plant CDKs remains to be demonstrated.

[0010] Therefore, although several plant cell cycle proteins have been elucidated, the prior art has yet to describe the plant cell cycle signal transduction cascades in detail. The prior art also fails to describe the relation between plant cell cycle proteins and a plant's response to environmental stress such that a stress tolerant plant can be generated. Accordingly, there is a need to identify genes expressed in stress tolerant plants that have the capacity to confer stress resistance to its host plant and to other plant species. Newly generated stress tolerant plants will have many advantages, such as increasing the range that crop plants can be cultivated by, for example, decreasing the water requirements of a plant species.

SUMMARY OF THE INVENTION

[0011] This invention fulfills in part the need to identify new, unique cell cycle proteins capable of conferring stress tolerance to plants upon over-expression. The present invention provides a transgenic plant cell transformed by a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid sequence in the plant cell results in increased tolerance to environmental stress as compared to a wild type variety of the plant cell. Namely, described herein are the cell cycle proteins 1) Cell Cycle-1 (CC-1); 2) Cell Cycle-2 (CC-2) and 3) Cell Cycle-3 (CC-3), all from *Physcomitrella patens*.

[0012] The invention provides in some embodiments that the CCSRPs and coding nucleic acid are that found in members of the genus *Physcomitrella*. In another preferred embodiment, the nucleic acid and protein are from a *Physcomitrella patens*. The invention provides that the environmental stress can be salinity, drought, temperature, metal, chemical, pathogenic and oxidative stresses, or combinations thereof. In preferred embodiments, the environmental stress can be drought or cold temperature.

[0013] The invention further provides a seed produced by a transgenic plant transformed by a CCSRPs coding nucleic acid, wherein the plant is true breeding for increased tolerance to environmental stress as compared to a wild type variety of the plant. The invention further provides a seed produced by a transgenic plant expressing a CCSRPs, wherein the plant is true breeding for increased tolerance to environmental stress as compared to a wild type variety of the plant.

[0014] The invention further provides an agricultural product produced by any of the below-described transgenic plants, plant parts or seeds. The invention further provides an isolated CCSRPs as described below. The invention further provides an isolated CCSRPs

coding nucleic acid, wherein the CCSRP coding nucleic acid codes for a CCSRP as described below.

[0015] The invention further provides an isolated recombinant expression vector comprising a CCSRP coding nucleic acid as described below, wherein expression of the vector in a host cell results in increased tolerance to environmental stress as compared to a wild type variety of the host cell. The invention further provides a host cell containing the vector and a plant containing the host cell.

10 [0016] The invention further provides a method of producing a transgenic plant with a CCSRP coding nucleic acid, wherein expression of the nucleic acid in the plant results in increased tolerance to environmental stress as compared to a wild type variety of the plant comprising: (a) transforming a plant cell with an expression vector comprising a CCSRP coding nucleic acid, and (b) generating from the plant cell a transgenic plant with an increased tolerance to environmental stress as compared to a wild type variety of the plant. In preferred embodiments, the CCSRP and CCSRP coding nucleic acid are as described below.

20 [0017] The present invention further provides a method of identifying a novel CCSRP, comprising (a) raising a specific antibody response to a CCSRP, or fragment thereof, as described below; (b) screening putative CCSRP material with the antibody, wherein specific binding of the antibody to the material indicates the presence of a potentially novel CCSRP; and (c) identifying from the bound material a novel CCSRP in comparison to known CCSRP. Alternatively, hybridization with nucleic acid probes as described below can be used to identify novel CCSRP nucleic acids.

[0018] The present invention also provides methods of modifying stress tolerance of a plant comprising, modifying the expression of a CCSRP in the plant, wherein the CCSRP is as described below. The invention provides that this method can be performed such that the stress tolerance is either increased or decreased. Preferably, stress tolerance is increased in a plant via increasing expression of a CCSRP.

[0018A] An aspect of the invention provides for an isolated Cell Cycle Stress-Related Protein (CCSRP) wherein the CCSRP is

- a. a Cell Cycle-1 protein (CC-1) as defined in SEQ ID NO:7; or

- b. a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length; or
- c. a polypeptide encoded by a nucleic acid that has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4,

and wherein the CCSRP upon expression in a plant results in the plant's increased tolerance to drought or freezing stress as compared to a wild type variety of the plant.

[0018B] The invention further comprehends a use of the CCSRP, for increasing the tolerance of a transformed plant or plant cell or plant seed to drought or freezing stress, wherein the plant or plant cell or plant seed includes the CCSRP, the CCSRP is a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length.

[0018C] Yet further, the present invention comprehends an isolated Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein the CCSRP coding nucleic acid is a nucleic acid:

- a. having a sequence as defined in SEQ ID NO:4; or
- b. having a sequence of at least 90% identity to SEQ ID NO:4 over the entire coding region of SEQ ID NO:4; or
- c. encoding the CCSRP as defined in claim 8;

and wherein the expression in a plant increases the plants tolerance to drought or freezing stress as compared to a wild type variety of the plant.

[0018D] Further still, the invention comprehends a method of producing a transgenic plant containing a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid in the plant results in increased tolerance to drought or freezing stress as compared to a wild type variety of the plant, comprising transforming a plant cell with an expression vector comprising the nucleic acid, generating from the plant cell a transgenic plant with an increased

tolerance to the drought and freezing stress, as compared to a wild type variety of the plant, wherein

- a. the CCSRP is a Cell Cycle-1 protein (CC-1) as defined in SEQ ID NO:7; or
- b. the CCSRP is a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length; or
- c. the CCSRP coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

10 **[0018E]** Further still, the invention comprehends a transgenic plant cell transformed by a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid in the plant cell results in increased tolerance to a drought or freezing stress as compared to a wild type variety of the plant cell, wherein

- a. the CCSRP is a cell cycle protein-1 (CC-1) protein as defined in SEQ ID NO:7; or
- b. a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length, or
- c. the CCSRP coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

20 **[0018F]** The invention also comprehends a method of modifying drought or freezing stress tolerance of a transgenic plant comprising, modifying the expression of a Cell Cycle Stress-Related Protein (CCSRP) in the transgenic plant as compared to a wild type variety of the plant, wherein

- a. the CCSRP is a cell cycle protein-1 (CC-1) protein as defined in SEQ ID NO:7; or
- b. a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length, or
- c. the CCSRP coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

The invention further comprehends an isolated recombinant expression vector comprising the CCSRP protein as defined above as well as the use of such vector or CCSRP protein in a host cell for increasing tolerance to drought or freezing stress as compared to the wild type variety of the host cell.


The invention still comprehends a use of the isolated CCSRP coding nucleic acid as defined above for identification or cloning of CC-1 homologs in other cell types and organisms.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019] Figures 1(A-C) show the partial cDNA sequences of CC-1 (SEQ ID NO:1), CC-2 (SEQ ID NO:2) and CC-3 (SEQ ID NO:3) from *Physcomitrella patens*.

[0020] Figures 2(A-C) show the full-length cDNA sequences of CC-1 (SEQ ID NO:4), CC-2 (SEQ ID NO:5) and CC-3 (SEQ ID NO:6) from *Physcomitrella patens*.

[0021] Figures 3(A-H) show the deduced amino acid sequences of CC-1 (SEQ ID NO:7), CC-2 (SEQ ID NO:8) and CC-3 (SEQ ID NO:9) from *Physcomitrella patens*.



[0022] Figure 4 shows a diagram of the plant expression vector ppBPSsc022 containing the super promoter driving the expression of SEQ ID NOs: 4, 5 and 6 ("Desired Gene"). The components are: NPTII kanamycin resistance gene, AtAct2-i promoter, OCS3 terminator, NOSpA terminator (Jefferson et al., 1987 EMBO J 6:3901-7).

[0023] Figure 5 shows the results of a drought stress test with over-expressing PpCC-1 transgenic plants and wild-type *Arabidopsis* lines. The transgenic lines display a tolerant phenotype. Individual transformant lines are shown.

[0024] Figure 6 shows the results of a freezing stress test with over-expressing PpCC-1 transgenic plants and wild-type *Arabidopsis* lines. The transgenic lines display a tolerant phenotype. Individual transformant lines are shown.

[0025] Figure 7 shows the results of a drought stress test with over-expressing PpCC-2 transgenic plants and wild-type *Arabidopsis* lines. The transgenic lines display a tolerant phenotype. Individual transformant lines are shown.

[0026] Figure 8 shows the results of a drought stress test with over-expressing PpCC-3 transgenic plants and wild-type *Arabidopsis* lines. The transgenic lines display a tolerant phenotype. Individual transformant lines are shown.

DETAILED DESCRIPTION OF THE INVENTION

[0027] The present invention may be understood more readily by reference to the following detailed description of the preferred embodiments of the invention and the Examples included herein. However, before the present compounds, compositions, and methods are disclosed and described, it is to be understood that this invention is not limited to specific nucleic acids, specific polypeptides, specific cell types, specific host cells, specific conditions, or specific methods, etc., as such may, of course, vary, and the numerous modifications and variations therein will be apparent to those skilled in the art. It is also to be understood that the terminology used herein is for the purpose of describing specific embodiments only and is not intended to be limiting. In particular, the designation of the amino acid sequences as protein "Cell Cycle Stress-Related Proteins" (CCSRPs), in no way limits the functionality of those sequences.

[0028] The present invention provides a transgenic plant cell transformed by a CCSRPs coding nucleic acid, wherein expression of the nucleic acid sequence in the plant cell results in increased tolerance to environmental stress as compared to a wild type variety of the plant cell. The invention further provides transgenic plant parts and transgenic plants containing the plant cells described herein. Also provided is a plant seed produced by a

transgenic plant transformed by a CCSRP coding nucleic acid, wherein the seed contains the CCSRP coding nucleic acid, and wherein the plant is true breeding for increased tolerance to environmental stress as compared to a wild type variety of the plant. The invention further provides a seed produced by a transgenic plant expressing a CCSRP, wherein the seed contains the CCSRP, and wherein the plant is true breeding for increased tolerance to environmental stress as compared to a wild type variety of the plant. The invention also provides an agricultural product produced by any of the below-described transgenic plants, plant parts and plant seeds.

[0029] As used herein, the term “variety” refers to a group of plants within a species that share constant characters that separate them from the typical form and from other possible varieties within that species. While possessing at least one distinctive trait, a variety is also characterized by some variation between individuals within the variety, based primarily on the Mendelian segregation of traits among the progeny of succeeding generations. A variety is considered “true breeding” for a particular trait if it is genetically homozygous for that trait to the extent that, when the true-breeding variety is self-pollinated, a significant amount of independent segregation of the trait among the progeny is not observed. In the present invention, the trait arises from the transgenic expression of one or more DNA sequences introduced into a plant variety.

[0030] The present invention describes for the first time that the *Physcomitrella patens* CCSRPs, CC-1, CC-2 and CC-3, are useful for increasing a plant’s tolerance to environmental stress. Accordingly, the present invention provides isolated CCSRPs selected from the group consisting of CC-1, CC-2 and CC-3, and homologs thereof. In preferred embodiments, the CCSRP is selected from 1) a Cell Cycle-1 (CC-1) protein as defined in SEQ ID NO:7; 2) a Cell Cycle-2 (CC-2) protein as defined in SEQ ID NO:8; and 3) a Cell Cycle-3 (CC-3) protein as defined in SEQ ID NO:9 and homologs and orthologs thereof. Homologs and orthologs of the amino acid sequences are defined below.

[0031] The CCSRPs of the present invention are preferably produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the protein is cloned into an expression vector (as described below), the expression vector is introduced into a host cell (as described below) and the CCSRP is expressed in the host cell. The CCSRP can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Alternative to recombinant expression, a CCSRP polypeptide, or peptide can be synthesized chemically using standard peptide synthesis techniques. Moreover, native CCSRP can be isolated from cells (e.g., *Physcomitrella patens*), for

example using an anti-CCSRP antibody, which can be produced by standard techniques utilizing a CCSRП or fragment thereof.

[0032] The invention further provides an isolated CCSRП coding nucleic acid. The present invention includes CCSRП coding nucleic acids that encode CCSRПs as described herein. In preferred embodiments, the CCSRП coding nucleic acid is selected from 1) a Cell Cycle-1 (CC-1) nucleic acid as defined in SEQ ID NO:4; 2) a Cell Cycle-2 (CC-2) nucleic acid as defined in SEQ ID NO:5; and 3) a Cell Cycle-3 (CC-3) nucleic acid as defined in SEQ ID NO:6 and homologs and orthologs thereof. Homologs and orthologs of the nucleotide sequences are defined below. In one preferred embodiment, the nucleic acid and protein are isolated from the plant genus *Physcomitrella*. In another preferred embodiment, the nucleic acid and protein are from a *Physcomitrella patens* (*P. patens*) plant.

[0033] As used herein, the term “environmental stress” refers to any sub-optimal growing condition and includes, but is not limited to, sub-optimal conditions associated with salinity, drought, temperature, metal, chemical, pathogenic and oxidative stresses, or combinations thereof. In preferred embodiments, the environmental stress can be salinity, drought, or temperature, or combinations thereof, and in particular, can be high salinity, low water content or low temperature. It is also to be understood that as used in the specification and in the claims, “a” or “an” can mean one or more, depending upon the context in which it is used. Thus, for example, reference to “a cell” can mean that at least one cell can be utilized.

[0034] As also used herein, the terms “nucleic acid” and “nucleic acid molecule” are intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. This term also encompasses untranslated sequence located at both the 3’ and 5’ ends of the coding region of the gene: at least about 1000 nucleotides of sequence upstream from the 5’ end of the coding region and at least about 200 nucleotides of sequence downstream from the 3’ end of the coding region of the gene. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

[0035] An “isolated” nucleic acid molecule is one that is substantially separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an “isolated” nucleic acid is free of some of the sequences which naturally flank the nucleic acid (i.e., sequences located at the 5’ and 3’ ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated CCSRП nucleic acid molecule can contain less than about

5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived (e.g., a *Physcomitrella patens* cell). Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be free from some of the other cellular material with which it is naturally associated, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized.

[0036] A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule having a nucleotide sequence of SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, a *P. patens* CCSRP cDNA can be isolated from a *P. patens* library using all or portion of one of the sequences of SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:3. Moreover, a nucleic acid molecule encompassing all or a portion of one of the sequences of SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:3 can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon this sequence. For example, mRNA can be isolated from plant cells (e.g., by the guanidinium-thiocyanate extraction procedure of Chirgwin et al., 1979 Biochemistry 18:5294-5299) and cDNA can be prepared using reverse transcriptase (e.g., Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, MD; or AMV reverse transcriptase, available from Seikagaku America, Inc., St. Petersburg, FL). Synthetic oligonucleotide primers for polymerase chain reaction amplification can be designed based upon one of the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:3. A nucleic acid molecule of the invention can be amplified using cDNA or, alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid molecule so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to a CCSRP nucleotide sequence can be prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

[0037] In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6. These cDNAs comprise sequences encoding the CCSRPs (i.e., the "coding region", indicated in Table 1), as well as 5' untranslated sequences and 3' untranslated sequences. It is to be understood that SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6 comprise both coding regions and 5' and 3' untranslated regions. Alternatively, the nucleic acid molecules of the present invention can comprise only the coding region of any of the

sequences in SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6 or can contain whole genomic fragments isolated from genomic DNA. A coding region of these sequences is indicated as an "ORF position". The present invention also includes CCSRP coding nucleic acids that encode CCSRPs as described herein. Preferred is a CCSRP coding nucleic acid that encodes a CCSRP selected from the group consisting of, CC-1 (SEQ ID NO:7), CC-2 (SEQ ID NO:8) and CC-3 (SEQ ID NO:9).

[0038] Moreover, the nucleic acid molecule of the invention can comprise only a portion of the coding region of one of the sequences in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6, for example, a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a CCSRP. The nucleotide sequences determined from the cloning of the CCSRP genes from *P. patens* allow for the generation of probes and primers designed for use in identifying and/or cloning CCSRP homologs in other cell types and organisms, as well as CCSRP homologs from other mosses and related species.

[0039] Portions of proteins encoded by the CCSRP nucleic acid molecules of the invention are preferably biologically active portions of one of the CCSRPs described herein. As used herein, the term "biologically active portion of" a CCSRP is intended to include a portion, e.g., a domain/motif, of a CCSRP that participates in a stress tolerance response in a plant, has an activity as set forth in Table 1, or participates in the transcription of a protein involved in a stress tolerance response in a plant. To determine whether a CCSRP, or a biologically active portion thereof, can participate in transcription of a protein involved in a stress tolerance response in a plant, a stress analysis of a plant comprising the CCSRP may be performed. Such analysis methods are well known to those skilled in the art, as detailed in Example 7. More specifically, nucleic acid fragments encoding biologically active portions of a CCSRP can be prepared by isolating a portion of one of the sequences in SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9, expressing the encoded portion of the CCSRP or peptide (e.g., by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the CCSRP or peptide.

[0040] Biologically active portions of a CCSRP are encompassed by the present invention and include peptides comprising amino acid sequences derived from the amino acid sequence of a CCSRP, e.g., an amino acid sequence of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9, or the amino acid sequence of a protein homologous to a CCSRP, which include fewer amino acids than a full length CCSRP or the full length protein which is homologous to a CCSRP, and exhibit at least one activity of a CCSRP. Typically, biologically active portions (e.g., peptides which are, for example, 5, 10, 15, 20, 30, 35, 36,

37, 38, 39, 40, 50, 100 or more amino acids in length) comprise a domain or motif with at least one activity of a CCSRP. Moreover, other biologically active portions in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the activities described herein. Preferably, the biologically active portions of a CCSRP include one or more selected domains/motifs or portions thereof having biological activity.

[0041] The invention also provides CCSRP chimeric or fusion proteins. As used herein, a CCSRP "chimeric protein" or "fusion protein" comprises a CCSRP polypeptide operatively linked to a non-CCSRP polypeptide. A CCSRP polypeptide refers to a polypeptide having an amino acid sequence corresponding to a CCSRP, whereas a non-CCSRP polypeptide refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the CCSRP, e.g., a protein that is different from the CCSRP and is derived from the same or a different organism. Within the fusion protein, the term "operatively linked" is intended to indicate that the CCSRP polypeptide and the non-CCSRP polypeptide are fused to each other so that both sequences fulfill the proposed function attributed to the sequence used. The non-CCSRP polypeptide can be fused to the N-terminus or C-terminus of the CCSRP polypeptide. For example, in one embodiment, the fusion protein is a GST-CCSRP fusion protein in which the CCSRP sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant CCSRPs. In another embodiment, the fusion protein is a CCSRP containing a heterologous signal sequence at its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a CCSRP can be increased through use of a heterologous signal sequence.

[0042] Preferably, a CCSRP chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see, for example, Current Protocols in

Molecular Biology, Eds. Ausubel et al. John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A CCSRP encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the CCSRP .

[0043] In addition to fragments and fusion proteins of the CCSRPs described herein, the present invention includes homologs and analogs of naturally occurring CCSRPs and CCSRP encoding nucleic acids in a plant. "Homologs" are defined herein as two nucleic acids or proteins that have similar, or "homologous", nucleotide or amino acid sequences, respectively. Homologs include allelic variants, orthologs, paralogs, agonists and antagonists of CCSRPs as defined hereafter. The term "homolog" further encompasses nucleic acid molecules that differ from one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6 (and portions thereof) due to degeneracy of the genetic code and thus encode the same CCSRP as that encoded by the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. As used herein a "naturally occurring" CCSRP refers to a CCSRP amino acid sequence that occurs in nature. Preferably, a naturally occurring CCSRP comprises an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9.

[0044] An agonist of the CCSRP can retain substantially the same, or a subset, of the biological activities of the CCSRP. An antagonist of the CCSRP can inhibit one or more of the activities of the naturally occurring form of the CCSRP. For example, the CCSRP antagonist can competitively bind to a downstream or upstream member of the cell membrane component metabolic cascade that includes the CCSRP, or bind to a CCSRP that mediates transport of compounds across such membranes, thereby preventing translocation from taking place.

[0045] Nucleic acid molecules corresponding to natural allelic variants and analogs, orthologs and paralogs of a CCSRP cDNA can be isolated based on their identity to the *Physcomitrella patens* CCSRP nucleic acids described herein using CCSRP cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. In an alternative embodiment, homologs of the CCSRP can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of the CCSRP for CCSRP agonist or antagonist activity. In one embodiment, a variegated library of CCSRP variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of CCSRP variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into

gene sequences such that a degenerate set of potential CCSRP sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display) containing the set of CCSRP sequences therein. There are a variety of methods that can be used to produce libraries of potential CCSRP homologs from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene is then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential CCSRP sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, S.A., 1983 Tetrahedron 39:3; Itakura et al., 1984 Annu. Rev. Biochem. 53:323; Itakura et al., 1984 Science 198:1056; Ike et al., 1983 Nucleic Acid Res. 11:477).

[0046] In addition, libraries of fragments of the CCSRP coding regions can be used to generate a variegated population of CCSRP fragments for screening and subsequent selection of homologs of a CCSRP. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of a CCSRP coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA, which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the CCSRP.

[0047] Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of CCSRP homologs. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique that enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify CCSRP homologs (Arkin and Yourvan, 1992 PNAS 89:7811-

7815; Delgrave et al., 1993 Protein Engineering 6(3):327-331). In another embodiment, cell based assays can be exploited to analyze a variegated CCSRP library, using methods well known in the art. The present invention further provides a method of identifying a novel CCSRP, comprising (a) raising a specific antibody response to a CCSRP, or a fragment thereof, as described above; (b) screening putative CCSRP material with the antibody, wherein specific binding of the antibody to the material indicates the presence of a potentially novel CCSRP; and (c) analyzing the bound material in comparison to known CCSRP, to determine its novelty.

[0048] To determine the percent homology of two amino acid sequences (e.g., one of the sequences of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9 and a mutant form thereof), the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of one protein or nucleic acid for optimal alignment with the other protein or nucleic acid). The amino acid residues at corresponding amino acid positions or nucleotide positions are then compared. When a position in one sequence (e.g., one of the sequences of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9) is occupied by the same amino acid residue as the corresponding position in the other sequence (e.g., a mutant form of the sequence selected from the polypeptide of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9), then the molecules are homologous at that position (i.e., as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The same type of comparison can be made between two nucleic acid sequences.

[0049] The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = numbers of identical positions/total numbers of positions x 100). Preferably, the amino acid sequences included in the present invention are at least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence shown in SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:9. In yet another embodiment, at least about 50-60%, preferably at least about 60-70%, and more preferably at least about 70-80%, 80-90%, 90-95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence encoded by a nucleic acid sequence shown in SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. In other embodiments, the preferable length of sequence comparison for proteins is at least 15 amino acid residues, more preferably at least 25 amino acid residues, and most preferably at least 35 amino acid residues.

[0050] In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which is at least about 50-60%, preferably at least about 60-70%, more preferably at least about 70-80%, 80-90%, or 90-95%, and even more preferably at least about 95%, 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence shown in SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6, or a portion thereof. The preferable length of sequence comparison for nucleic acids is at least 75 nucleotides, more preferably at least 100 nucleotides and most preferably the entire coding region of the nucleic acid.

[0051] It is also preferable that the homologous nucleic acid molecule of the invention encodes a protein or portion thereof which includes an amino acid sequence which is sufficiently homologous to an amino acid sequence of SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:9 such that the protein or portion thereof maintains the same or a similar function as the amino acid sequence to which it is compared. Functions of the CCSRP amino acid sequences of the present invention include the ability to participate in a stress tolerance response in a plant, or more particularly, to participate in the transcription of a protein involved in a stress tolerance response in a *Physcomitrella patens* plant. Examples of such activities are described in Table 1.

[0052] In addition to the above-described methods, a determination of the percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990 Proc. Natl. Acad. Sci. USA 90:5873-5877). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990 J. Mol. Biol. 215:403-410).

[0053] BLAST nucleic acid searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleic acid sequences homologous to the CCSRP nucleic acid molecules of the invention. Additionally, BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to CCSRPs of the present invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997 Nucleic Acids Res. 25:3389-3402). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. Another preferred non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (CABIOS 1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) that is part of the GCG

sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4 can be used to obtain amino acid sequences homologous to the CCSRPs of the present invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997 Nucleic Acids Res. 25:3389-3402). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. Another preferred non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (CABIOS 1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) that is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4 can be used.

[0054] Finally, homology between nucleic acid sequences can also be determined using hybridization techniques known to those of skill in the art. Accordingly, an isolated nucleic acid molecule of the invention comprises a nucleotide sequence which hybridizes, e.g., hybridizes under stringent conditions, to one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6, or a portion thereof. More particularly, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. In other embodiments, the nucleic acid is at least 30, 50, 100, 250 or more nucleotides in length.

[0055] As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 65%, more preferably at least about 70%, and even more preferably at least about 75% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, 6.3.1-6.3.6, John Wiley & Sons, N.Y. (1989). A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to a sequence of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6 corresponds to a naturally occurring nucleic acid molecule. As used herein, a "naturally occurring" nucleic acid molecule refers to an RNA or

DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein). In one embodiment, the nucleic acid encodes a naturally occurring *Physcomitrella patens* CCSRP.

[0056] Using the above-described methods, and others known to those of skill in the art, one of ordinary skill in the art can isolate homologs of the CCSRPs comprising amino acid sequences shown in SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:9. One subset of these homologs is allelic variants. As used herein, the term “allelic variant” refers to a nucleotide sequence containing polymorphisms that lead to changes in the amino acid sequences of a CCSRP and that exist within a natural population (e.g., a plant species or variety). Such natural allelic variations can typically result in 1-5% variance in a CCSRP nucleic acid. Allelic variants can be identified by sequencing the nucleic acid sequence of interest in a number of different plants, which can be readily carried out by using hybridization probes to identify the same CCSRP genetic locus in those plants. Any and all such nucleic acid variations and resulting amino acid polymorphisms or variations in a CCSRP that are the result of natural allelic variation and that do not alter the functional activity of a CCSRP, are intended to be within the scope of the invention.

[0057] Moreover, nucleic acid molecules encoding CCSRPs from the same or other species such as CCSRP analogs, orthologs and paralogs, are intended to be within the scope of the present invention. As used herein, the term “analogs” refers to two nucleic acids that have the same or similar function, but that have evolved separately in unrelated organisms. As used herein, the term “orthologs” refers to two nucleic acids from different species, but that have evolved from a common ancestral gene by speciation. Normally, orthologs encode proteins having the same or similar functions. As also used herein, the term “paralogs” refers to two nucleic acids that are related by duplication within a genome. Paralogs usually have different functions, but these functions may be related (Tatusov, R.L. et al. 1997 Science 278(5338):631-637). Analogs, orthologs and paralogs of a naturally occurring CCSRP can differ from the naturally occurring CCSRP by post-translational modifications, by amino acid sequence differences, or by both. Post-translational modifications include *in vivo* and *in vitro* chemical derivatization of polypeptides, e.g., acetylation, carboxylation, phosphorylation, or glycosylation, and such modifications may occur during polypeptide synthesis or processing or following treatment with isolated modifying enzymes. In particular, orthologs of the invention will generally exhibit at least 80-85%, more preferably 90%, and most preferably 95%, 96%, 97%, 98% or even 99% identity or homology with all or part of a naturally occurring CCSRP amino acid sequence and will exhibit a function similar to a CCSRP.

Orthologs of the present invention are also preferably capable of participating in the stress response in plants. In one embodiment, the CCSRP orthologs maintain the ability to participate in the metabolism of compounds necessary for the construction of cellular membranes in *Physcomitrella patens*, or in the transport of molecules across these membranes.

[0058] In addition to naturally-occurring variants of a CCSRP sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into a nucleotide sequence of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6, thereby leading to changes in the amino acid sequence of the encoded CCSRP, without altering the functional ability of the CCSRP. For example, nucleotide substitutions leading to amino acid substitutions at “non-essential” amino acid residues can be made in a sequence of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. A “non-essential” amino acid residue is a residue that can be altered from the wild-type sequence of one of the CCSRPs without altering the activity of said CCSRP, whereas an “essential” amino acid residue is required for CCSRP activity. Other amino acid residues, however, (e.g., those that are not conserved or only semi-conserved in the domain having CCSRP activity) may not be essential for activity and thus are likely to be amenable to alteration without altering CCSRP activity.

[0059] Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding CCSRPs that contain changes in amino acid residues that are not essential for CCSRP activity. Such CCSRPs differ in amino acid sequence from a sequence contained in SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:9, yet retain at least one of the CCSRP activities described herein. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 50% homologous to an amino acid sequence of SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:9. Preferably, the protein encoded by the nucleic acid molecule is at least about 50-60% homologous to one of the sequences of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9, more preferably at least about 60-70% homologous to one of the sequences of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9, even more preferably at least about 70-80%, 80-90%, 90-95% homologous to one of the sequences of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9, and most preferably at least about 96%, 97%, 98%, or 99% homologous to one of the sequences of SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9. The preferred CCSRP homologs of the present invention are preferably capable of participating in the a stress tolerance response in a plant, or more particularly, participating in the

transcription of a protein involved in a stress tolerance response in a *Physcomitrella patens* plant, or have one or more activities set forth in Table 1.

[0060] An isolated nucleic acid molecule encoding a CCSRP homologous to a protein sequence of SEQ ID NO:7, SEQ ID NO:8 or SEQ ID NO:9 can be created by introducing one or more nucleotide substitutions, additions or deletions into a nucleotide sequence of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6 such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into one of the sequences of SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain.

[0061] Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a CCSRP is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a CCSRP coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for a CCSRP activity described herein to identify mutants that retain CCSRP activity. Following mutagenesis of one of the sequences of SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6, the encoded protein can be expressed recombinantly and the activity of the protein can be determined by analyzing the stress tolerance of a plant expressing the protein as described in Example 7.

[0062] In addition to the nucleic acid molecules encoding the CCSRPs described above, another aspect of the invention pertains to isolated nucleic acid molecules that are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be complementary to an entire CCSRP coding strand, or to

only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding a CCSRP. The term "coding region" refers to the region of the nucleotide sequence comprising codons that are translated into amino acid residues (e.g., the entire coding region of ,,,, comprises nucleotides 1 to ...). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding a CCSRP. The term "noncoding region" refers to 5' and 3' sequences that flank the coding region that are not translated into amino acids (i.e., also referred to as 5' and 3' untranslated regions).

[0063] In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6, or a portion thereof. A nucleic acid molecule that is complementary to one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6 is one which is sufficiently complementary to one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6 such that it can hybridize to one of the nucleotide sequences shown in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6, thereby forming a stable duplex.

[0064] Given the coding strand sequences encoding the CCSRPs disclosed herein (e.g., the sequences set forth in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of CCSRP mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of CCSRP mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of CCSRP mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length.

[0065] An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-

(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

[0066] The antisense nucleic acid molecules of the invention are typically administered to a cell or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a CCSRP to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. The antisense molecule can be modified such that it specifically binds to a receptor or an antigen expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecule to a peptide or an antibody which binds to a cell surface receptor or antigen. The antisense nucleic acid molecule can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong prokaryotic, viral, or eukaryotic (including plant) promoter are preferred.

[0067] In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier et al., 1987 Nucleic Acids. Res. 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-

methyribonucleotide (Inoue et al., 1987 Nucleic Acids Res. 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue et al., 1987 FEBS Lett. 215:327-330).

[0068] In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes described in Haselhoff and Gerlach, 1988 Nature 334:585-591) can be used to catalytically cleave CCSRP mRNA transcripts to thereby inhibit translation of CCSRP mRNA. A ribozyme having specificity for a CCSRP-encoding nucleic acid can be designed based upon the nucleotide sequence of a CCSRP cDNA, as disclosed herein (i.e., SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6) or on the basis of a heterologous sequence to be isolated according to methods taught in this invention. For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a CCSRP-encoding mRNA. See, e.g., Cech et al. U.S. Patent No. 4,987,071 and Cech et al. U.S. Patent No. 5,116,742. Alternatively, CCSRP mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel, D. and Szostak, J.W., 1993 Science 261:1411-1418.

[0069] Alternatively, CCSRP gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of a CCSRP nucleotide sequence (e.g., a CCSRP promoter and/or enhancer) to form triple helical structures that prevent transcription of a CCSRP gene in target cells. See generally, Helene, C., 1991 Anticancer Drug Des. 6(6):569-84; Helene, C. et al., 1992 Ann. N.Y. Acad. Sci. 660:27-36; and Maher, L.J., 1992 Bioassays 14(12):807-15.

[0070] In addition to the CCSRP nucleic acids and proteins described above, the present invention encompasses these nucleic acids and proteins attached to a moiety. These moieties include, but are not limited to, detection moieties, hybridization moieties, purification moieties, delivery moieties, reaction moieties, binding moieties, and the like. A typical group of nucleic acids attached to a moiety are probes and primers. Probes and primers typically comprise a substantially isolated oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense strand of one of the sequences set forth in SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6, an anti-sense sequence of one of the sequences set forth in SEQ ID

NO:4, SEQ ID NO:5 and SEQ ID NO:6, or naturally occurring mutants thereof. Primers based on a nucleotide sequence of SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6 can be used in PCR reactions to clone CCSRP homologs. Probes based on the CCSRP nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, e.g. the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a genomic marker test kit for identifying cells which express a CCSRP, such as by measuring a level of a CCSRP-encoding nucleic acid, in a sample of cells, e.g., detecting CCSRP mRNA levels or determining whether a genomic CCSRP gene has been mutated or deleted.

[0071] In particular, a useful method to ascertain the level of transcription of the gene (an indicator of the amount of mRNA available for translation to the gene product) is to perform a Northern blot (for reference see, for example, Ausubel et al., 1988 Current Protocols in Molecular Biology, Wiley: New York). This information at least partially demonstrates the degree of transcription of the transformed gene. Total cellular RNA can be prepared from cells, tissues or organs by several methods, all well-known in the art, such as that described in Bormann, E.R. et al., 1992 Mol. Microbiol. 6:317-326. To assess the presence or relative quantity of protein translated from this mRNA, standard techniques, such as a Western blot, may be employed. These techniques are well known to one of ordinary skill in the art. (See, for example, Ausubel et al., 1988 Current Protocols in Molecular Biology, Wiley: New York).

[0072] The invention further provides an isolated recombinant expression vector comprising a CCSRP nucleic acid as described above, wherein expression of the vector in a host cell results in increased tolerance to environmental stress as compared to a wild type variety of the host cell. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are

operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0073] The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (e.g., in an *in vitro* transcription/ translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990) or see: Gruber and Crosby, in: Methods in Plant Molecular Biology and Biotechnology, eds. Glick and Thompson, Chapter 7, 89-108, CRC Press: Boca Raton, Florida, including the references therein. Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cells and those that direct expression of the nucleotide sequence only in certain host cells or under certain conditions. It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., CCSRPs, mutant forms of CCSRPs, fusion proteins, etc.).

[0074] The recombinant expression vectors of the invention can be designed for expression of CCSRPs in prokaryotic or eukaryotic cells. For example, CCSRP genes can be expressed in bacterial cells such as *C. glutamicum*, insect cells (using baculovirus expression vectors), yeast and other fungal cells (see Romanos, M.A. et al., 1992 Foreign gene

expression in yeast: a review, *Yeast* 8:423-488; van den Hondel, C.A.M.J.J. et al., 1991 Heterologous gene expression in filamentous fungi, in: *More Gene Manipulations in Fungi*, J.W. Bennet & L.L. Lasure, eds., p. 396-428: Academic Press: San Diego; and van den Hondel, C.A.M.J.J. & Punt, P.J., 1991 Gene transfer systems and vector development for filamentous fungi, in: *Applied Molecular Genetics of Fungi*, Peberdy, J.F. et al., eds., p. 1-28, Cambridge University Press: Cambridge), algae (Falciatore et al., 1999 *Marine Biotechnology* 1(3):239-251), ciliates of the types: Holotrichia, Peritrichia, Spirotrichia, Suctoria, Tetrahymena, Paramecium, Colpidium, Glaucoma, Platyophrya, Potomacus, Pseudocohnilembus, Euplotes, Engelmaniella, and Stylonychia, especially of the genus *Stylonychia lemnae* with vectors following a transformation method as described in PCT Application No. WO 98/01572 and multicellular plant cells (see Schmidt, R. and Willmitzer, L., 1988 High efficiency *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* leaf and cotyledon explants, *Plant Cell Rep.* 583-586); *Plant Molecular Biology and Biotechnology*, C Press, Boca Raton, Florida, chapter 6/7, S.71-119 (1993); F.F. White, B. Jenes et al., *Techniques for Gene Transfer*, in: *Transgenic Plants*, Vol. 1, Engineering and Utilization, eds. Kung und R. Wu, 128-43, Academic Press: 1993; Potrykus, 1991 *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42:205-225 and references cited therein) or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press: San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

[0075] Expression of proteins in prokaryotes is most often carried out with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein but also to the C-terminus or fused within suitable regions in the proteins. Such fusion vectors typically serve three purposes: 1) to increase expression of a recombinant protein; 2) to increase the solubility of a recombinant protein; and 3) to aid in the purification of a recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.

[0076] Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S., 1988 *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. In one embodiment, the coding sequence of the CCSRP is cloned into a pGEX expression vector to create a vector encoding a fusion protein comprising, from the N-terminus to the C-terminus, GST-thrombin cleavage site-X protein. The fusion protein can be purified by affinity chromatography using glutathione-agarose resin. Recombinant CCSRP unfused to GST can be recovered by cleavage of the fusion protein with thrombin.

[0077] Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann et al., 1988 *Gene* 69:301-315) and pET 11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 *gn10*-lac fusion promoter mediated by a co-expressed viral RNA polymerase (T7 *gn1*). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 *gn1* gene under the transcriptional control of the lacUV 5 promoter.

[0078] One strategy to maximize recombinant protein expression is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in the bacterium chosen for expression, such as *C. glutamicum* (Wada et al., 1992 *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

[0079] In another embodiment, the CCSRP expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, et al., 1987 *EMBO J.* 6:229-234), pMFa (Kurjan and Herskowitz, 1982 *Cell* 30:933-943), pJRY88 (Schultz et al., 1987 *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA). Vectors and methods for the construction of vectors appropriate for use in other fungi, such as the filamentous fungi, include those detailed in: van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in:

Applied Molecular Genetics of Fungi, J.F. Peberdy, et al., eds., p. 1-28, Cambridge University Press: Cambridge.

[0080] Alternatively, the CCSRPs of the invention can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith et al., 1983 Mol. Cell Biol. 3:2156-2165) and the pVL series (Lucklow and Summers, 1989 Virology 170:31-39).

[0081] In yet another embodiment, a CCSRP nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B., 1987 Nature 329:840) and pMT2PC (Kaufman et al., 1987 EMBO J. 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

[0082] In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert et al., 1987 Genes Dev. 1:268-277), lymphoid-specific promoters (Calame and Eaton, 1988 Adv. Immunol. 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989 EMBO J. 8:729-733) and immunoglobulins (Banerji et al., 1983 Cell 33:729-740; Queen and Baltimore, 1983 Cell 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle, 1989 PNAS 86:5473-5477), pancreas-specific promoters (Edlund et al., 1985 Science 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally regulated promoters are also encompassed, for example, the murine hox promoters (Kessel and Gruss, 1990 Science 249:374-379) and the fetoprotein promoter (Campes and Tilghman, 1989 Genes Dev. 3:537-546).

[0083] In another embodiment, the CCSRPs of the invention may be expressed in unicellular plant cells (such as algae) (see Falciatore et al., 1999 Marine Biotechnology

1(3):239-251 and references therein) and plant cells from higher plants (e.g., the spermatophytes, such as crop plants). Examples of plant expression vectors include those detailed in: Becker, D., Kemper, E., Schell, J. and Masterson, R., 1992 New plant binary vectors with selectable markers located proximal to the left border, *Plant Mol. Biol.* 20: 1195-1197; and Bevan, M.W., 1984 Binary *Agrobacterium* vectors for plant transformation, *Nucl. Acid. Res.* 12:8711-8721; Vectors for Gene Transfer in Higher Plants; in: *Transgenic Plants, Vol. 1, Engineering and Utilization*, eds.: Kung and R. Wu, Academic Press, 1993, S. 15-38.

[0084] A plant expression cassette preferably contains regulatory sequences capable of driving gene expression in plant cells and operably linked so that each sequence can fulfill its function, for example, termination of transcription by polyadenylation signals. Preferred polyadenylation signals are those originating from *Agrobacterium tumefaciens* t-DNA such as the gene 3 known as octopine synthase of the Ti-plasmid pTiACH5 (Gielen et al., 1984 *EMBO J.* 3:835) or functional equivalents thereof but also all other terminators functionally active in plants are suitable.

[0085] As plant gene expression is very often not limited on transcriptional levels, a plant expression cassette preferably contains other operably linked sequences like translational enhancers such as the overdrive-sequence containing the 5'-untranslated leader sequence from tobacco mosaic virus enhancing the protein per RNA ratio (Gallie et al., 1987 *Nucl. Acids Research* 15:8693-8711).

[0086] Plant gene expression has to be operably linked to an appropriate promoter conferring gene expression in a timely, cell or tissue specific manner. Preferred are promoters driving constitutive expression (Benfey et al., 1989 *EMBO J.* 8:2195-2202) like those derived from plant viruses like the 35S CAMV (Franck et al., 1980 *Cell* 21:285-294), the 19S CaMV (see also U.S. Patent No. 5352605 and PCT Application No. WO 8402913) or plant promoters like those from Rubisco small subunit described in U.S. Patent No. 4,962,028.

[0087] Other preferred sequences for use in plant gene expression cassettes are targeting-sequences necessary to direct the gene product in its appropriate cell compartment (for review see Kermode, 1996 *Crit. Rev. Plant Sci.* 15(4):285-423 and references cited therein) such as the vacuole, the nucleus, all types of plastids like amyloplasts, chloroplasts, chromoplasts, the extracellular space, mitochondria, the endoplasmic reticulum, oil bodies, peroxisomes and other compartments of plant cells.

[0088] Plant gene expression can also be facilitated via an inducible promoter (for review see Gatz, 1997 Annu. Rev. Plant Physiol. Plant Mol. Biol. 48:89-108). Chemically inducible promoters are especially suitable if gene expression is wanted to occur in a time specific manner. Examples of such promoters are a salicylic acid inducible promoter (PCT Application No. WO 95/19443), a tetracycline inducible promoter (Gatz et al., 1992 Plant J. 2:397-404) and an ethanol inducible promoter (PCT Application No. WO 93/21334).

[0089] Also, suitable promoters responding to biotic or abiotic stress conditions are those such as the pathogen inducible PRP1-gene promoter (Ward et al., 1993 Plant. Mol. Biol. 22:361-366), the heat inducible hsp80-promoter from tomato (U.S. Patent No. 5187267), cold inducible alpha-amylase promoter from potato (PCT Application No. WO 96/12814) or the wound-inducible pinII-promoter (European Patent No. 375091). For other examples of drought, cold, and salt-inducible promoters, such as the RD29A promoter, see Yamaguchi-Shinozaki et al. (1993 Mol. Gen. Genet. 236:331-340).

[0090] Especially preferred are those promoters that confer gene expression in specific tissues and organs, such as guard cells and the root hair cells. Suitable promoters include the napin-gene promoter from rapeseed (U.S. Patent No. 5,608,152), the USP-promoter from *Vicia faba* (Baeumlein et al., 1991 Mol Gen Genet. 225(3):459-67), the oleosin-promoter from *Arabidopsis* (PCT Application No. WO 98/45461), the phaseolin-promoter from *Phaseolus vulgaris* (U.S. Patent No. 5,504,200), the Bce4-promoter from Brassica (PCT Application No. WO 91/13980) or the legumin B4 promoter (LeB4; Baeumlein et al., 1992 Plant Journal, 2(2):233-9) as well as promoters conferring seed specific expression in monocot plants like maize, barley, wheat, rye, rice, etc. Suitable promoters to note are the lpt2 or lpt1-gene promoter from barley (PCT Application No. WO 95/15389 and PCT Application No. WO 95/23230) or those described in PCT Application No. WO 99/16890 (promoters from the barley hordein-gene, rice glutelin gene, rice oryzin gene, rice prolamin gene, wheat gliadin gene, wheat glutelin gene, maize zein gene, oat glutelin gene, Sorghum kasirin-gene and rye secalin gene).

[0091] Also especially suited are promoters that confer plastid-specific gene expression since plastids are the compartment where lipid biosynthesis occurs. Suitable promoters are the viral RNA-polymerase promoter described in PCT Application No. WO 95/16783 and PCT Application No. WO 97/06250 and the clpP-promoter from *Arabidopsis* described in PCT Application No. WO 99/46394.

[0092] The invention further provides a recombinant expression vector comprising a CCSRP DNA molecule of the invention cloned into the expression vector in an antisense

orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner that allows for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to a CCSRP mRNA. Regulatory sequences operatively linked to a nucleic acid molecule cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types. For instance, viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus wherein antisense nucleic acids are produced under the control of a high efficiency regulatory region. The activity of the regulatory region can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. et al., Antisense RNA as a molecular tool for genetic analysis, Reviews - Trends in Genetics, Vol. 1(1) 1986 and Mol et al., 1990 FEBS Letters 268:427-430.

[0093] Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but they also apply to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

[0094] A host cell can be any prokaryotic or eukaryotic cell. For example, a CCSRP can be expressed in bacterial cells such as *C. glutamicum*, insect cells, fungal cells or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells), algae, ciliates, plant cells, fungi or other microorganisms like *C. glutamicum*. Other suitable host cells are known to those skilled in the art.

[0095] Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation", "transfection", "conjugation" and "transduction" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, natural competence, chemical-mediated transfer and electroporation. Suitable methods for transforming or transfecting host cells including plant cells can be found in Sambrook, et al. (Molecular Cloning: A Laboratory Manual. 2nd, ed.,

Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) and other laboratory manuals such as *Methods in Molecular Biology*, 1995, Vol. 44, *Agrobacterium* protocols, ed: Gartland and Davey, Humana Press, Totowa, New Jersey. As biotic and abiotic stress tolerance is a general trait wished to be inherited into a wide variety of plants like maize, wheat, rye, oat, triticale, rice, barley, soybean, peanut, cotton, rapeseed and canola, manihot, pepper, sunflower and tagetes, solanaceous plants like potato, tobacco, eggplant, and tomato, Vicia species, pea, alfalfa, bushy plants (coffee, cacao, tea), Salix species, trees (oil palm, coconut), perennial grasses and forage crops, these crop plants are also preferred target plants for a genetic engineering as one further embodiment of the present invention.

[0096] In particular, the invention provides a method of producing a transgenic plant with a CCSRP coding nucleic acid, wherein expression of the nucleic acid(s) in the plant results in increased tolerance to environmental stress as compared to a wild type variety of the plant comprising: (a) transforming a plant cell with an expression vector comprising a CCSRP nucleic acid, and (b) generating from the plant cell a transgenic plant with a increased tolerance to environmental stress as compared to a wild type variety of the plant. The invention also provides a method of increasing expression of a gene of interest within a host cell as compared to a wild type variety of the host cell, wherein the gene of interest is transcribed in response to a CCSRP, comprising: (a) transforming the host cell with an expression vector comprising a CCSRP coding nucleic acid, and (b) expressing the CCSRP within the host cell, thereby increasing the expression of the gene transcribed in response to the CCSRP, as compared to a wild type variety of the host cell.

[0097] For such plant transformation, binary vectors such as pBinAR can be used (Höfgen and Willmitzer, 1990 *Plant Science* 66:221-230). Construction of the binary vectors can be performed by ligation of the cDNA in sense or antisense orientation into the T-DNA. 5-prime to the cDNA a plant promoter activates transcription of the cDNA. A polyadenylation sequence is located 3-prime to the cDNA. Tissue-specific expression can be achieved by using a tissue specific promoter. For example, seed-specific expression can be achieved by cloning the napin or LeB4 or USP promoter 5-prime to the cDNA. Also, any other seed specific promoter element can be used. For constitutive expression within the whole plant, the CaMV 35S promoter can be used. The expressed protein can be targeted to a cellular compartment using a signal peptide, for example for plastids, mitochondria or endoplasmic reticulum (Kermode, 1996 *Crit. Rev. Plant Sci.* 4 (15):285-423). The signal peptide is cloned 5-prime in frame to the cDNA to archive subcellular localization of the

fusion protein. Additionally, promoters that are responsive to abiotic stresses can be used with, such as the *Arabidopsis* promoter RD29A, the nucleic acid sequences disclosed herein. One skilled in the art will recognize that the promoter used should be operatively linked to the nucleic acid such that the promoter causes transcription of the nucleic acid which results in the synthesis of a mRNA which encodes a polypeptide. Alternatively, the RNA can be an antisense RNA for use in affecting subsequent expression of the same or another gene or genes.

[0098] Alternate methods of transfection include the direct transfer of DNA into developing flowers via electroporation or *Agrobacterium* mediated gene transfer. *Agrobacterium* mediated plant transformation can be performed using for example the GV3101(pMP90) (Koncz and Schell, 1986 Mol. Gen. Genet. 204:383-396) or LBA4404 (Clontech) *Agrobacterium tumefaciens* strain. Transformation can be performed by standard transformation and regeneration techniques (Deblaere et al., 1994 Nucl. Acids. Res. 13:4777-4788; Gelvin, Stanton B. and Schilperoort, Robert A, Plant Molecular Biology Manual, 2nd Ed. - Dordrecht : Kluwer Academic Publ., 1995. - in Sect., Ringbuc Zentrale Signatur: BT11-P ISBN 0-7923-2731-4; Glick, Bernard R.; Thompson, John E., Methods in Plant Molecular Biology and Biotechnology, Boca Raton : CRC Press, 1993 360 S., ISBN 0-8493-5164-2). For example, rapeseed can be transformed via cotyledon or hypocotyl transformation (Moloney et al., 1989 Plant cell Report 8:238-242; De Block et al., 1989 Plant Physiol. 91:694-701). Use of antibiotics for *Agrobacterium* and plant selection depends on the binary vector and the *Agrobacterium* strain used for transformation. Rapeseed selection is normally performed using kanamycin as selectable plant marker. *Agrobacterium* mediated gene transfer to flax can be performed using, for example, a technique described by Mlynarova et al., 1994 Plant Cell Report 13:282-285. Additionally, transformation of soybean can be performed using for example a technique described in European Patent No. 0424 047, U.S. Patent No. 5,322,783, European Patent No. 0397 687, U.S. Patent No. 5,376,543 or U.S. Patent No. 5,169,770. Transformation of maize can be achieved by particle bombardment, polyethylene glycol mediated DNA uptake or via the silicon carbide fiber technique. (See, for example, Freeling and Walbot "The maize handbook" Springer Verlag: New York (1993) ISBN 3-540-97826-7). A specific example of maize transformation is found in U.S. Patent No. 5,990,387 and a specific example of wheat transformation can be found in PCT Application No. WO 93/07256.

[0099] For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may

integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate or in plants that confer resistance towards a herbicide such as glyphosate or glufosinate. Nucleic acid molecules encoding a selectable marker can be introduced into a host cell on the same vector as that encoding a CCSRP or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid molecule can be identified by, for example, drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

[0100] To create a homologous recombinant microorganism, a vector is prepared which contains at least a portion of a CCSRP gene into which a deletion, addition or substitution has been introduced to thereby alter, e.g., functionally disrupt, the CCSRP gene. Preferably, the CCSRP gene is a *Physcomitrella patens* CCSRP gene, but it can be a homolog from a related plant or even from a mammalian, yeast, or insect source. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous CCSRP gene is functionally disrupted (i.e., no longer encodes a functional protein; also referred to as a knock-out vector). Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous CCSRP gene is mutated or otherwise altered but still encodes a functional protein (e.g., the upstream regulatory region can be altered to thereby alter the expression of the endogenous CCSRP). To create a point mutation via homologous recombination, DNA-RNA hybrids can be used in a technique known as chimeraplasty (Cole-Strauss et al., 1999 Nucleic Acids Research 27(5):1323-1330 and Kmiec, 1999 Gene therapy American Scientist. 87(3):240-247). Homologous recombination procedures in *Physcomitrella patens* are also well known in the art and are contemplated for use herein.

[0101] Whereas in the homologous recombination vector, the altered portion of the CCSRP gene is flanked at its 5' and 3' ends by an additional nucleic acid molecule of the CCSRP gene to allow for homologous recombination to occur between the exogenous CCSRP gene carried by the vector and an endogenous CCSRP gene, in a microorganism or plant. The additional flanking CCSRP nucleic acid molecule is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several hundreds of base pairs up to kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see e.g., Thomas, K.R., and Capecchi, M.R., 1987 Cell 51:503 for a

description of homologous recombination vectors or Strepp et al., 1998 PNAS, 95 (8):4368-4373 for cDNA based recombination in *Physcomitrella patens*). The vector is introduced into a microorganism or plant cell (e.g., via polyethylene glycol mediated DNA), and cells in which the introduced CCSRP gene has homologously recombined with the endogenous CCSRP gene are selected using art-known techniques.

[0102] In another embodiment, recombinant microorganisms can be produced that contain selected systems which allow for regulated expression of the introduced gene. For example, inclusion of a CCSRP gene on a vector placing it under control of the lac operon permits expression of the CCSRP gene only in the presence of IPTG. Such regulatory systems are well known in the art.

[0103] A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (i.e., express) a CCSRP. Accordingly, the invention further provides methods for producing CCSRPs using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding a CCSRP has been introduced, or into which genome has been introduced a gene encoding a wild-type or altered CCSRP) in a suitable medium until CCSRP is produced. In another embodiment, the method further comprises isolating CCSRPs from the medium or the host cell.

[0104] Another aspect of the invention pertains to isolated CCSRPs, and biologically active portions thereof. An "isolated" or "purified" protein or biologically active portion thereof is free of some of the cellular material when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of CCSRP in which the protein is separated from some of the cellular components of the cells in which it is naturally or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of a CCSRP having less than about 30% (by dry weight) of non-CCSRP material (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-CCSRP material, still more preferably less than about 10% of non-CCSRP material, and most preferably less than about 5% non-CCSRP material.

[0105] When the CCSRP or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation. The language "substantially free of chemical precursors or other chemicals" includes preparations of CCSRP in which the

protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language “substantially free of chemical precursors or other chemicals” includes preparations of a CCSRP having less than about 30% (by dry weight) of chemical precursors or non-CCSRP chemicals, more preferably less than about 20% chemical precursors or non-CCSRP chemicals, still more preferably less than about 10% chemical precursors or non-CCSRP chemicals, and most preferably less than about 5% chemical precursors or non-CCSRP chemicals. In preferred embodiments, isolated proteins, or biologically active portions thereof, lack contaminating proteins from the same organism from which the CCSRP is derived. Typically, such proteins are produced by recombinant expression of, for example, a *Physcomitrella patens* CCSRP in plants other than *Physcomitrella patens* or microorganisms such as *C. glutamicum*, ciliates, algae or fungi.

[0106] The nucleic acid molecules, proteins, protein homologs, fusion proteins, primers, vectors, and host cells described herein can be used in one or more of the following methods: identification of *Physcomitrella patens* and related organisms; mapping of genomes of organisms related to *Physcomitrella patens*; identification and localization of *Physcomitrella patens* sequences of interest; evolutionary studies; determination of CCSRP regions required for function; modulation of a CCSRP activity; modulation of the metabolism of one or more cell functions; modulation of the transmembrane transport of one or more compounds; and modulation of stress resistance.

[0107] The moss *Physcomitrella patens* represents one member of the mosses. It is related to other mosses such as *Ceratodon purpureus* which is capable of growth in the absence of light. Mosses like *Ceratodon* and *Physcomitrella* share a high degree of homology on the DNA sequence and polypeptide level allowing the use of heterologous screening of DNA molecules with probes evolving from other mosses or organisms, thus enabling the derivation of a consensus sequence suitable for heterologous screening or functional annotation and prediction of gene functions in third species. The ability to identify such functions can therefore have significant relevance, e.g., prediction of substrate specificity of enzymes. Further, these nucleic acid molecules may serve as reference points for the mapping of moss genomes, or of genomes of related organisms.

[0108] The CCSRP nucleic acid molecules of the invention have a variety of uses. Most importantly, the nucleic acid and amino acid sequences of the present invention can be used to transform plants, thereby inducing tolerance to stresses such as drought, high salinity and cold. The present invention therefore provides a transgenic plant transformed by a CCSRP nucleic acid, wherein expression of the nucleic acid sequence in the plant results in

increased tolerance to environmental stress as compared to a wild type variety of the plant. The transgenic plant can be a monocot or a dicot. The invention further provides that the transgenic plant can be selected from maize, wheat, rye, oat, triticale, rice, barley, soybean, peanut, cotton, rapeseed, canola, manihot, pepper, sunflower, tagetes, solanaceous plants, potato, tobacco, eggplant, tomato, Vicia species, pea, alfalfa, coffee, cacao, tea, Salix species, oil palm, coconut, perennial grass and forage crops, for example.

[0109] In particular, the present invention describes using the expression of CC-1, CC-2 and CC-3 of *Physcomitrella patens* to engineer drought-tolerant, salt-tolerant and/or cold-tolerant plants. This strategy has herein been demonstrated for *Arabidopsis thaliana*, Rapeseed/Canola, soybeans, corn and wheat but its application is not restricted to these plants. Accordingly, the invention provides a transgenic plant containing a CCSRP selected from CC-1 (SEQ ID NO:7), CC-2 (SEQ ID NO:8) and CC-3 (SEQ ID NO:9), wherein the environmental stress is drought, increased salt or decreased or increased temperature. In preferred embodiments, the environmental stress is drought or decreased temperature.

[0110] The present invention also provides methods of modifying stress tolerance of a plant comprising, modifying the expression of a CCSRP in the plant. The invention provides that this method can be performed such that the stress tolerance is either increased or decreased. In particular, the present invention provides methods of producing a transgenic plant having an increased tolerance to environmental stress as compared to a wild type variety of the plant comprising increasing expression of a CCSRP in a plant.

[0111] The methods of increasing expression of CCSRPs can be used wherein the plant is either transgenic or not transgenic. In cases when the plant is transgenic, the plant can be transformed with a vector containing any of the above described CCSRP coding nucleic acids, or the plant can be transformed with a promoter that directs expression of native CCSRP in the plant, for example. The invention provides that such a promoter can be tissue specific. Furthermore, such a promoter can be developmentally regulated. Alternatively, non-transgenic plants can have native CCSRP expression modified by inducing a native promoter.

[0112] The expression of CC-1 (SEQ ID NO:7), CC-2 (SEQ ID NO:8) or CC-3 (SEQ ID NO:9) in target plants can be accomplished by, but is not limited to, one of the following examples: (a) constitutive promoter, (b) stress-inducible promoter, (c) chemical-induced promoter, and (d) engineered promoter over-expression with for example zinc-finger derived transcription factors (Greisman and Pabo, 1997 Science 275:657). The later case involves identification of the CC-1 (SEQ ID NO:7), CC-2 (SEQ ID NO:8) or CC-3 (SEQ ID NO:9)

homologs in the target plant as well as from its promoter. Zinc-finger-containing recombinant transcription factors are engineered to specifically interact with the CC-1 (SEQ ID NO:7), CC-2 (SEQ ID NO:8) or CC-3 (SEQ ID NO:9) homolog and transcription of the corresponding gene is activated.

[0113] In addition to introducing the CCSRP nucleic acid sequences into transgenic plants, these sequences can also be used to identify an organism as being *Physcomitrella patens* or a close relative thereof. Also, they may be used to identify the presence of *Physcomitrella patens* or a relative thereof in a mixed population of microorganisms. The invention provides the nucleic acid sequences of a number of *Physcomitrella patens* genes; by probing the extracted genomic DNA of a culture of a unique or mixed population of microorganisms under stringent conditions with a probe spanning a region of a *Physcomitrella patens* gene which is unique to this organism, one can ascertain whether this organism is present.

[0114] Further, the nucleic acid and protein molecules of the invention may serve as markers for specific regions of the genome. This has utility not only in the mapping of the genome, but also in functional studies of *Physcomitrella patens* proteins. For example, to identify the region of the genome to which a particular *Physcomitrella patens* DNA-binding protein binds, the *Physcomitrella patens* genome could be digested, and the fragments incubated with the DNA-binding protein. Those fragments that bind the protein may be additionally probed with the nucleic acid molecules of the invention, preferably with readily detectable labels. Binding of such a nucleic acid molecule to the genome fragment enables the localization of the fragment to the genome map of *Physcomitrella patens*, and, when performed multiple times with different enzymes, facilitates a rapid determination of the nucleic acid sequence to which the protein binds. Further, the nucleic acid molecules of the invention may be sufficiently homologous to the sequences of related species such that these nucleic acid molecules may serve as markers for the construction of a genomic map in related mosses.

[0115] The CCSRP nucleic acid molecules of the invention are also useful for evolutionary and protein structural studies. The metabolic and transport processes in which the molecules of the invention participate are utilized by a wide variety of prokaryotic and eukaryotic cells; by comparing the sequences of the nucleic acid molecules of the present invention to those encoding similar enzymes from other organisms, the evolutionary relatedness of the organisms can be assessed. Similarly, such a comparison permits an assessment of which regions of the sequence are conserved and which are not, which may aid

in determining those regions of the protein that are essential for the functioning of the enzyme. This type of determination is of value for protein engineering studies and may give an indication of what the protein can tolerate in terms of mutagenesis without losing function.

[0116] Manipulation of the CCSRP nucleic acid molecules of the invention may result in the production of CCSRPs having functional differences from the wild-type CCSRPs. These proteins may be improved in efficiency or activity, may be present in greater numbers in the cell than is usual, or may be decreased in efficiency or activity.

[0117] There are a number of mechanisms by which the alteration of a CCSRP of the invention may directly affect stress response and/or stress tolerance. In the case of plants expressing CCSRPs, increased transport can lead to improved salt and/or solute partitioning within the plant tissue and organs. By either increasing the number or the activity of transporter molecules which export ionic molecules from the cell, it may be possible to affect the salt tolerance of the cell.

[0118] The effect of the genetic modification in plants, *C. glutamicum*, fungi, algae, or ciliates on stress tolerance can be assessed by growing the modified microorganism or plant under less than suitable conditions and then analyzing the growth characteristics and/or metabolism of the plant. Such analysis techniques are well known to one skilled in the art, and include dry weight, wet weight, protein synthesis, carbohydrate synthesis, lipid synthesis, evapotranspiration rates, general plant and/or crop yield, flowering, reproduction, seed setting, root growth, respiration rates, photosynthesis rates, etc. (Applications of HPLC in Biochemistry in: Laboratory Techniques in Biochemistry and Molecular Biology, vol. 17; Rehm et al., 1993 Biotechnology, vol. 3, Chapter III: Product recovery and purification, page 469-714, VCH: Weinheim; Belter, P.A. et al., 1988 Bioseparations: downstream processing for biotechnology, John Wiley and Sons; Kennedy, J.F. and Cabral, J.M.S., 1992 Recovery processes for biological materials, John Wiley and Sons; Shaeiwitz, J.A. and Henry, J.D., 1988 Biochemical separations, in: Ulmann's Encyclopedia of Industrial Chemistry, vol. B3, Chapter 11, page 1-27, VCH: Weinheim; and Dechow, F.J. (1989) Separation and purification techniques in biotechnology, Noyes Publications).

[0119] For example, yeast expression vectors comprising the nucleic acids disclosed herein, or fragments thereof, can be constructed and transformed into *Saccharomyces cerevisiae* using standard protocols. The resulting transgenic cells can then be assayed for fail or alteration of their tolerance to drought, salt, and temperature stress. Similarly, plant expression vectors comprising the nucleic acids disclosed herein, or fragments thereof, can be constructed and transformed into an appropriate plant cell such as *Arabidopsis*, soy, rape,

maize, wheat, *Medicago truncatula*, etc., using standard protocols. The resulting transgenic cells and/or plants derived there from can then be assayed for fail or alteration of their tolerance to drought, salt, and temperature stress.

[0120] The engineering of one or more CCSRP genes of the invention may also result in CCSRPs having altered activities which indirectly impact the stress response and/or stress tolerance of algae, plants, ciliates or fungi or other microorganisms like *C. glutamicum*. For example, the normal biochemical processes of metabolism result in the production of a variety of products (e.g., hydrogen peroxide and other reactive oxygen species) which may actively interfere with these same metabolic processes (for example, peroxynitrite is known to nitrate tyrosine side chains, thereby inactivating some enzymes having tyrosine in the active site (Groves, J.T., 1999 Curr. Opin. Chem. Biol. 3(2):226-235). While these products are typically excreted, cells can be genetically altered to transport more products than is typical for a wild-type cell. By optimizing the activity of one or more CCSRPs of the invention which are involved in the export of specific molecules, such as salt molecules, it may be possible to improve the stress tolerance of the cell.

[0121] Additionally, the sequences disclosed herein, or fragments thereof, can be used to generate knockout mutations in the genomes of various organisms, such as bacteria, mammalian cells, yeast cells, and plant cells (Girke, T., 1998 The Plant Journal 15:39-48). The resultant knockout cells can then be evaluated for their ability or capacity to tolerate various stress conditions, their response to various stress conditions, and the effect on the phenotype and/or genotype of the mutation. For other methods of gene inactivation see U.S. Patent No. 6004804 "Non-Chimeric Mutational Vectors" and Puttaraju et al., 1999 Spliceosome-mediated RNA *trans*-splicing as a tool for gene therapy Nature Biotechnology 17:246-252.

[0122] The aforementioned mutagenesis strategies for CCSRPs resulting in increased stress resistance are not meant to be limiting; variations on these strategies will be readily apparent to one skilled in the art. Using such strategies, and incorporating the mechanisms disclosed herein, the nucleic acid and protein molecules of the invention may be utilized to generate algae, ciliates, plants, fungi or other microorganisms like *C. glutamicum* expressing mutated CCSRP nucleic acid and protein molecules such that the stress tolerance is improved.

[0123] The present invention also provides antibodies that specifically bind to a CCSRP, or a portion thereof, as encoded by a nucleic acid described herein. Antibodies can be made by many well-known methods (See, e.g. Harlow and Lane, "Antibodies; A

Laboratory Manual" Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1988)). Briefly, purified antigen can be injected into an animal in an amount and in intervals sufficient to elicit an immune response. Antibodies can either be purified directly, or spleen cells can be obtained from the animal. The cells can then fused with an immortal cell line and screened for antibody secretion. The antibodies can be used to screen nucleic acid clone libraries for cells secreting the antigen. Those positive clones can then be sequenced. (See, for example, Kelly et al., 1992 Bio/Technology 10:163-167; Bebbington et al., 1992 Bio/Technology 10:169-175).

[0124] The phrases "selectively binds" and "specifically binds" with the polypeptide refer to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bound to a particular protein do not bind in a significant amount to other proteins present in the sample. Selective binding of an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. A variety of immunoassay formats may be used to select antibodies that selectively bind with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies selectively immunoreactive with a protein. See *Harlow and Lane "Antibodies, A Laboratory Manual"* Cold Spring Harbor Publications, New York, (1988), for a description of immunoassay formats and conditions that could be used to determine selective binding.

[0125] In some instances, it is desirable to prepare monoclonal antibodies from various hosts. A description of techniques for preparing such monoclonal antibodies may be found in Stites et al., editors, "Basic and Clinical Immunology," (Lange Medical Publications, Los Altos, Calif., Fourth Edition) and references cited therein, and in Harlow and Lane ("Antibodies, A Laboratory Manual" Cold Spring Harbor Publications, New York, 1988).

[0126] Throughout this application, various publications are referenced. The disclosures of all of these publications and those references cited within those publications in their entireties may be referred to for further details in this application in order to more fully describe the state of the art to which this invention pertains.

[0127] It should also be understood that the foregoing relates to preferred embodiments of the present invention and that numerous changes may be made therein without departing from the scope of the invention. The invention is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon

the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof, which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention and/or the scope of the appended claims.

EXAMPLES

Example 1

Growth of Physcomitrella patens cultures

[0128] For this study, plants of the species *Physcomitrella patens* (Hedw.) B.S.G. from the collection of the genetic studies section of the University of Hamburg were used. They originate from the strain 16/14 collected by H.L.K. Whitehouse in Gransden Wood, Huntingdonshire (England), which was subcultured from a spore by Engel (1968, Am. J. Bot. 55, 438-446). Proliferation of the plants was carried out by means of spores and by means of regeneration of the gametophytes. The protonema developed from the haploid spore as a chloroplast-rich chloronema and chloroplast-low caulonema, on which buds formed after approximately 12 days. These grew to give gametophores bearing antheridia and archegonia. After fertilization, the diploid sporophyte with a short seta and the spore capsule resulted, in which the meiospores matured.

[0129] Culturing was carried out in a climatic chamber at an air temperature of 25°C and light intensity of 55 micromols^{-1m2} (white light; Philips TL 65W/25 fluorescent tube) and a light/dark change of 16/8 hours. The moss was either modified in liquid culture using Knop medium according to Reski and Abel (1985, Planta 165:354-358) or cultured on Knop solid medium using 1% oxoid agar (Unipath, Basingstoke, England). The protonemas used for RNA and DNA isolation were cultured in aerated liquid cultures. The protonemas were comminuted every 9 days and transferred to fresh culture medium.

Example 2

Total DNA isolation from plants

[0130] The details for the isolation of total DNA relate to the working up of one gram fresh weight of plant material. The materials used include the following buffers: CTAB buffer: 2% (w/v) N-cethyl-N,N,N-trimethylammonium bromide (CTAB); 100 mM Tris HCl pH 8.0; 1.4 M NaCl; 20 mM EDTA; N-Laurylsarcosine buffer: 10% (w/v) N-laurylsarcosine; 100 mM Tris HCl pH 8.0; 20 mM EDTA.

[0131] The plant material was triturated under liquid nitrogen in a mortar to give a fine powder and transferred to 2 ml Eppendorf vessels. The frozen plant material was then covered with a layer of 1 ml of decomposition buffer (1 ml CTAB buffer, 100 µl of N-laurylsarcosine buffer, 20 µl of β-mercaptoethanol and 10 µl of proteinase K solution, 10 mg/ml) and incubated at 60°C for one hour with continuous shaking. The homogenate obtained was distributed into two Eppendorf vessels (2 ml) and extracted twice by shaking with the same volume of chloroform/isoamyl alcohol (24:1). For phase separation, centrifugation was carried out at 8000 x g and room temperature for 15 minutes in each case. The DNA was then precipitated at -70°C for 30 minutes using ice-cold isopropanol. The precipitated DNA was sedimented at 4°C and 10,000 g for 30 minutes and resuspended in 180 µl of TE buffer (Sambrook et al., 1989, Cold Spring Harbor Laboratory Press: ISBN 0-87969-309-6). For further purification, the DNA was treated with NaCl (1.2 M final concentration) and precipitated again at -70°C for 30 minutes using twice the volume of absolute ethanol. After a washing step with 70% ethanol, the DNA was dried and subsequently taken up in 50 µl of H₂O + RNase (50 mg/ml final concentration). The DNA was dissolved overnight at 4°C and the RNase digestion was subsequently carried out at 37°C for 1 hour. Storage of the DNA took place at 4°C.

Example 3

Isolation of total RNA and poly-(A)⁺ RNA and cDNA library construction from Physcomitrella patens

[0132] For the investigation of transcripts, both total RNA and poly-(A)⁺ RNA were isolated. The total RNA was obtained from wild-type 9 day old protonemata following the GTC-method (Reski et al. 1994, Mol. Gen. Genet., 244:352-359). The Poly(A)⁺ RNA was isolated using Dyna Beads^R (Dyna, Oslo, Norway) following the instructions of the manufacturers protocol. After determination of the concentration of the RNA or of the poly(A)⁺ RNA, the RNA was precipitated by addition of 1/10 volumes of 3 M sodium acetate pH 4.6 and 2 volumes of ethanol and stored at -70°C.

[0133] For cDNA library construction, first strand synthesis was achieved using Murine Leukemia Virus reverse transcriptase (Roche, Mannheim, Germany) and oligo-d(T)-primers, second strand synthesis by incubation with DNA polymerase I, Klenow enzyme and RNaseH digestion at 12°C (2 hours), 16°C (1 hour) and 22°C (1 hour). The reaction was stopped by incubation at 65°C (10 minutes) and subsequently transferred to ice. Double

stranded DNA molecules were blunted by T4-DNA-polymerase (Roche, Mannheim) at 37°C (30 minutes). Nucleotides were removed by phenol/chloroform extraction and Sephadex G50 spin columns. EcoRI adapters (Pharmacia, Freiburg, Germany) were ligated to the cDNA ends by T4-DNA-ligase (Roche, 12°C, overnight) and phosphorylated by incubation with polynucleotide kinase (Roche, 37°C, 30 minutes). This mixture was subjected to separation on a low melting agarose gel. DNA molecules larger than 300 base pairs were eluted from the gel, phenol extracted, concentrated on Elutip-D-columns (Schleicher and Schuell, Dassel, Germany) and were ligated to vector arms and packed into lambda ZAPII phages or lambda ZAP-Express phages using the Gigapack Gold Kit (Stratagene, Amsterdam, Netherlands) using material and following the instructions of the manufacturer.

Example 4

Sequencing and function annotation of Physcomitrella patens ESTs

[0134] cDNA libraries as described in Example 3 were used for DNA sequencing according to standard methods, and in particular, by the chain termination method using the ABI PRISM Big Dye Terminator Cycle Sequencing Ready Reaction Kit (Perkin-Elmer, Weiterstadt, Germany). Random Sequencing was carried out subsequent to preparative plasmid recovery from cDNA libraries via *in vivo* mass excision, retransformation, and subsequent plating of DH10B on agar plates (material and protocol details from Stratagene, Amsterdam, Netherlands. Plasmid DNA was prepared from overnight grown *E. coli* cultures grown in Luria-Broth medium containing ampicillin (see Sambrook et al. 1989 Cold Spring Harbor Laboratory Press: ISBN 0-87969-309-6) on a Qiagen DNA preparation robot (Qiagen, Hilden) according to the manufacturer's protocols. Sequencing primers with the following nucleotide sequences were used:

5'-CAGGAAACAGCTATGACC-3'	SEQ ID NO:10
5'-CTAAAGGGAACAAAAGCTG-3'	SEQ ID NO:11
5'-TGTAACACGACGGCCAGT-3'	SEQ ID NO:12

[0135] Sequences were processed and annotated using the software package EST-MAX commercially provided by Bio-Max (Munich, Germany). The program incorporates practically all bioinformatics methods important for functional and structural characterization of protein sequences. For reference the website at pedant.mips.biochem.mpg.de. The most important algorithms incorporated in EST-MAX are: FASTA: Very sensitive sequence

database searches with estimates of statistical significance; Pearson W.R. (1990) Rapid and sensitive sequence comparison with FASTP and FASTA. *Methods Enzymol.* 183:63-98; BLAST: Very sensitive sequence database searches with estimates of statistical significance. Altschul S.F., Gish W., Miller W., Myers E.W., and Lipman D.J. Basic local alignment search tool. *Journal of Molecular Biology* 215:403-10; PREDATOR: High-accuracy secondary structure prediction from single and multiple sequences. Frishman, D. and Argos, P. (1997) 75% accuracy in protein secondary structure prediction. *Proteins*, 27:329-335; CLUSTALW: Multiple sequence alignment. Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22:4673-4680; TMAP: Transmembrane region prediction from multiply aligned sequences. Persson, B. and Argos, P. (1994) Prediction of transmembrane segments in proteins utilizing multiple sequence alignments. *J. Mol. Biol.* 237:182-192; ALOM2: Transmembrane region prediction from single sequences. Klein, P., Kanehisa, M., and DeLisi, C. Prediction of protein function from sequence properties: A discriminate analysis of a database. *Biochim. Biophys. Acta* 787:221-226 (1984). Version 2 by Dr. K. Nakai; PROSEARCH: Detection of PROSITE protein sequence patterns. Kolakowski L.F. Jr., Leunissen J.A.M., Smith J.E. (1992) ProSearch: fast searching of protein sequences with regular expression patterns related to protein structure and function. *Biotechniques* 13, 919-921; BLIMPS: Similarity searches against a database of ungapped blocks. J.C. Wallace and Henikoff S., (1992); PATMAT: A searching and extraction program for sequence, pattern and block queries and databases, *CABIOS* 8:249-254. Written by Bill Alford.

Example 5

Identification of Physcomitrella patens ORFs corresponding to CC-1, CC-2 and CC-3

[0136] The *Physcomitrella patens* partial cDNAs (ESTs) shown in Table 1 below were identified in the *Physcomitrella patens* EST sequencing program using the program EST-MAX through BLAST analysis. The Sequence Identification Numbers corresponding to these ESTs are as follows: CC-1 (SEQ ID NO:1), CC-2 (SEQ ID NO:2) and CC-3 (SEQ ID NO:3).

Table 1

Name	Functional categories	Function	Sequence code	ORF position
PpCC-1	Cell Cycle	suppressor protein	c_pp004062017r	116-886
PpCC-2	Cell Cycle	cell division control protein	c_pp004064149r	1-590
PpCC-3	Cell Cycle	cell division protein	s_pp002007053r	106-492

Table 2

[0137] Degree of Amino Acid Identity and Similarity of PpCC-1 and Other Homologous Proteins GCG Gap program was used: gap penalty: 10; gap extension penalty: 0.1; score matrix: blosum62)

Swiss-Prot #	Q9ZNT0	P46467	P52917	Q9UI03	Q09803
Protein name	PUTATIVE SKD1 PROTEIN	SKD1 PROTEIN	VACUOLAR PROTEIN SORTING-ASSOCIATED PROTEIN VPS4	SKD1 PROTEIN	SUPPRESSOR PROTEIN OF BEM1/BED5 DOUBLE MUTANTS
Species	Arabidopsis thaliana (Mouse-ear cress)	Mus musculus (Mouse)	Saccharomyces cerevisiae (Baker's yeast)	Homo sapiens (Human)	Schizosaccharomyces pombe (Fission yeast)
Identity %	82%	54%	55%	53%	52%
Similarity %	90%	66%	67%	65%	65%

Table 3

[0138] Degree of Amino Acid Identity and Similarity of PpCC-2 and Other Homologous Proteins GCG Gap program was used: gap penalty: 10; gap extension penalty: 0.1; score matrix: blosum62)

Swiss-Prot #	Q9SIV8	P29458	P30665	P49717	P33991
Protein name	PUTATIVE CDC21 PROTEIN	CDC21 PROTEIN	CELL DIVISION CONTROL PROTEIN 54	DNA REPLICATI ON LICENSING FACTOR MCM4	DNA REPLICATI ON LICENSING FACTOR MCM4
Species	Arabidopsis thaliana (Mouse-ear cress)	Schizosacc haromyces pombe (Fission yeast)	Saccharomy ces cerevisiae (Baker's yeast)	Mus musculus (Mouse)	Homo sapiens (Human)
Identity %	49%	41%	40%	43%	43%
Similarity %	61%	55%	54%	56%	57%

Table 4

[0139] Degree of Amino Acid Identity and Similarity of PpCC-3 and Other Homologous Proteins GCG Gap program was used: gap penalty: 10; gap extension penalty: 0.1; score matrix: blosum62)

Swiss-Prot #	O80860	Q9LXF7	P51327	O78516	Q55700
Protein name	PUTATIVE CELL DIVISION PROTEIN (ZINC DEPENDENT PROTEASE)	FTSH-LIKE PROTEIN PFTF	CELL DIVISION PROTEIN FTSH HOMOLOG	CELL DIVISION PROTEIN FTSH HOMOLOG	CELL DIVISION PROTEIN FTSH HOMOLOG 1
Species	Arabidopsis thaliana (Mouse-ear cress)	Arabidopsis thaliana (Mouse-ear cress)	Porphyra purpurea	Guillardia theta (Cryptomonas phi)	Synechocystis sp. (strain PCC 6803)
Identity %	77%	64%	60%	59%	56%
Similarity %	84%	75%	69%	69%	67%

Example 6

Cloning of the full-length Physcomitrella patens cDNA encoding for CC-1, CC-2 and CC-3

[0140] To isolate the clones encoding Pp CC-1 (SEQ ID NO:4), PpCC-2 (SEQ ID NO:5) and PpCC-3 (SEQ ID NO:6) from *Physcomitrella patens*, cDNA libraries were created with SMART RACE cDNA Amplification kit (Clontech Laboratories) following manufacturer's instructions. Total RNA isolated as described in Example 3 was used as the template. The cultures were treated prior to RNA isolation as follows: Salt Stress: 2, 6, 12, 24, 48 hours with 1-M NaCl-supplemented medium; Cold Stress: 4°C for the same time points as for salt; Drought Stress: cultures were incubated on dry filter paper for the same time points as for salt.

5' RACE Protocol

[0141] The EST sequences CC-2 (SEQ ID NO:2) and CC-3 (SEQ ID NO:3) identified from the database search as described in Example 4 were used to design oligos for

RACE (see Table 5). The extended sequences for these genes were obtained by performing Rapid Amplification of cDNA Ends polymerase chain reaction (RACE PCR) using the Advantage 2 PCR kit (Clontech Laboratories) and the SMART RACE cDNA amplification kit (Clontech Laboratories) using a Biometra T3 Thermocycler following the manufacturer's instructions. The sequences obtained from the RACE reactions corresponded to full-length coding regions of CC-2 and CC-3 and were used to design oligos for full-length cloning of the respective genes (see below full-length amplification).

Full-length Amplification

[0142] Full-length clones corresponding CC-1 (SEQ ID NO:4), CC-2 (SEQ ID NO:5), and CC-3 (SEQ ID NO:6) were obtained by performing polymerase chain reaction (PCR) with gene-specific primers (see Table 5) and the original EST as the template. The conditions for the reaction were standard conditions with PWO DNA polymerase (Roche). PCR was performed according to standard conditions and to manufacturer's protocols (Sambrook et al., 1989 Molecular Cloning, A Laboratory Manual. 2nd Edition. Cold Spring Harbor Laboratory Press. Cold Spring Harbor, N.Y., Biometra T3 Thermocycler). The parameters for the reaction were: five minutes at 94°C followed by five cycles of one minute at 94°C, one minute at 50°C and 1.5 minutes at 72°C. This was followed by twenty five cycles of one minute at 94°C, one minute at 65°C and 1.5 minutes at 72°C.

[0143] The amplified fragments were extracted from agarose gel with a QIAquick Gel Extraction Kit (Qiagen) and ligated into the TOPO pCR 2.1 vector (Invitrogen) following manufacturer's instructions. Recombinant vectors were transformed into Top10 cells (Invitrogen) using standard conditions (Sambrook et al. 1989. Molecular Cloning, A Laboratory Manual. 2nd Edition. Cold Spring Harbor Laboratory Press. Cold Spring Harbor, NY). Transformed cells were selected for on LB agar containing 100 µg/ml carbenicillin, 0.8 mg X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside) and 0.8 mg IPTG (isopropylthio-β-D-galactoside) grown overnight at 37°C. White colonies were selected and used to inoculate 3 ml of liquid LB containing 100 µg/ml ampicillin and grown overnight at 37°C. Plasmid DNA was extracted using the QIAprep Spin Miniprep Kit (Qiagen) following manufacturer's instructions. Analyses of subsequent clones and restriction mapping was performed according to standard molecular biology techniques (Sambrook et al., 1989 Molecular Cloning, A Laboratory Manual. 2nd Edition. Cold Spring Harbor Laboratory Press. Cold Spring Harbor, N.Y.).

Table 5

[0144] Scheme and primers used for cloning of full-length clones

Gene	Sites in final product	Isolation Method	Primers Race	Primer RT-PCR
PpCC-1	EcoRV/ SacI	5' RACE and RT-PCR for Full-length clone	N/A	RC409 (SEQ ID NO:13) GCGATATCGACC CAAGGTGTGTAG AGAAGGGGAT RC410 (SEQ ID NO:14) GCGAGCTCGCGA CTGATGCAACTCA CCCATGAA
PpCC-2	AscI/ SacI	5' RACE and RT-PCR for Full-length clone	RC793: (SEQ ID NO:15) CCCAAATGCTT GGAGCCAGCGA CCT RC794: (SEQ ID NO:16) GACGTGCACGC AATCGATGTAG GTC NVT: (SEQ ID NO:17) AGTCGACCCCT GTCAGACTTCTT GA	RC990: (SEQ ID NO:18) ATGGCGCGCCGC ACTCACGTGAGG AATTGCACCTCC RC991: (SEQ ID NO:19) GCGAGCTCTGGG AATTGCGCATGG GAAACTGG

PpCC-3	XmaI/ SacI	5' RACE and RT-PCR for Full-length clone	NVT: (SEQ ID NO:20) CTGGCTACCCA AACCGCGTCGG AGA	RC695: (SEQ ID NO:21) ATCCCGGGTGAT ACGTTGTGCATAT TTGGTGTTC RC696: (SEQ ID NO:22) GCGAGCTCTGAT ACAAGAGCGTCG ATGCTCCA
---------------	------------	---	---	--

Example 7

Engineering stress-tolerant Arabidopsis plants by over-expressing the genes CC-1, CC-2 and CC-3

Binary vector construction: kanamycin

[0145] The plasmid construct pACGH101 was digested with PstI (Roche) and FseI (NEB) according to manufacturers' instructions. The fragment was purified by agarose gel and extracted via the Qiaex II DNA Extraction kit (Qiagen). This resulted in a vector fragment with the Arabidopsis Actin2 promoter with internal intron and the OCS3 terminator. Primers for PCR amplification of the NPTII gene were designed as follows:

5'NPT-Pst:

GCG-CTG-CAG-ATT-TCA-TTT-GGA-GAG-GAC-ACG (SEQ ID NO:23)

3'NPT-Fse:

CGC-GGC-CGG-CCT-CAG-AAG-AAC-TCG-TCA-AGA-AGG-CG (SEQ ID NO:24)

[0146] The 0.9 kilobase NPTII gene was amplified via PCR from pCambia 2301 plasmid DNA [94oC 60sec, {94oC 60sec, 61oC (-0.1oC per cycle) 60sec, 72oC 2min} x 25 cycles, 72oC 10min on Biometra T-Gradient machine], and purified via the Qiaquick PCR Extraction kit (Qiagen) as per manufacturer's instructions. The PCR DNA was then subcloned into the pCR-BluntII TOPO vector (Invitrogen) pursuant to the manufacturer's instructions (NPT-Topo construct). These ligations were transformed into Top10 cells

(Invitrogen) and grown on LB plates with 50ug/ml kanamycin sulfate overnight at 37°C. Colonies were then used to inoculate 2ml LB media with 50ug/ml kanamycin sulfate and grown overnight at 37°C. Plasmid DNA was recovered using the Qiaprep Spin Miniprep kit (Qiagen) and sequenced in both the 5' and 3' directions using standard conditions. Subsequent analysis of the sequence data using VectorNTI software revealed no PCR errors present in the NPTII gene sequence.

[0147] The NPT-Topo construct was then digested with PstI (Roche) and FseI (NEB) according to manufacturers' instructions. The 0.9 kilobase fragment was purified on agarose gel and extracted by Qiaex II DNA Extraction kit (Qiagen). The Pst/Fse insert fragment from NPT-Topo and the Pst/Fse vector fragment from pACGH101 were then ligated together using T4 DNA Ligase (Roche) following manufacturer's instructions. The ligation was then transformed into Top10 cells (Invitrogen) under standard conditions, creating pBPSsc019 construct. Colonies were selected on LB plates with 50 µg/ml kanamycin sulfate and grown overnight at 37°C. These colonies were then used to inoculate 2ml LB media with 50 µg/ml kanamycin sulfate and grown overnight at 37°C. Plasmid DNA was recovered using the Qiaprep Spin Miniprep kit (Qiagen) following the manufacturer's instructions.

[0148] The pBPSSC019 construct was digested with KpnI and BsaI (Roche) according to manufacturer's instructions. The fragment was purified via agarose gel and then extracted via the Qiaex II DNA Extraction kit (Qiagen) as per its instructions, resulting in a 3 kilobase Act-NPT cassette, which included the Arabidopsis Actin2 promoter with internal intron, the NPTII gene and the OCS3 terminator.

[0149] The pBPSJH001 vector was digested with SpeI and ApaI (Roche) and blunt-end filled with Klenow enzyme and 0.1mM dNTPs (Roche) according to manufacture's instructions. This produced a 10.1 kilobase vector fragment minus the Gentamycin cassette, which was recircularized by self-ligating with T4 DNA Ligase (Roche), and transformed into Top10 cells (Invitrogen) via standard conditions. Transformed cells were selected for on LB agar containing 50µg/ml kanamycin sulfate and grown overnight at 37°C. Colonies were then used to inoculate 2ml of liquid LB containing 50µg/ml kanamycin sulfate and grown overnight at 37°C. Plasmid DNA was extracted using the QIAprep Spin Miniprep Kit (Qiagen) following manufacture's instructions. The recircularized plasmid was then digested with KpnI (Roche) and extracted from agarose gel via the Qiaex II DNA Extraction kit (Qiagen) as per manufacturers' instructions.

[0150] The Act-NPT Kpn-cut insert and the Kpn-cut pBPSJH001 recircularized vector were then ligated together using T4 DNA Ligase (Roche) and transformed into Top10 cells (Invitrogen) as per manufacturers' instructions. The resulting construct, pBPSsc022, now contained the Super Promoter, the GUS gene, the NOS terminator, and the Act-NPT cassette. Transformed cells were selected for on LB agar containing 50µg/ml kanamycin sulfate and grown overnight at 37°C. Colonies were then used to inoculate 2ml of liquid LB containing 50µg/ml kanamycin sulfate and grown overnight at 37°C. Plasmid DNA was extracted using the QIAprep Spin Miniprep Kit (Qiagen) following manufacturer's instructions. After confirmation of ligation success via restriction digests, pBPSsc022 plasmid DNA was further propagated and recovered using the Plasmid Midiprep Kit (Qiagen) following the manufacturer's instructions.

Subcloning of CC-1, CC-2 and CC-3 into the binary vector

[0151] The fragments containing the different *Physcomitrella patens* transcription factors were subcloned from the recombinant PCR2.1 TOPO vectors by double digestion with restriction enzymes (see Table 6) according to manufacturer's instructions. The subsequence fragment was excised from agarose gel with a QIAquick Gel Extraction Kit (QIAGEN) according to manufacture's instructions and ligated into the binary vectors pGMSG, cleaved with XmaI and Ecl136II and dephosphorylated prior to ligation. The resulting recombinant pGMSG contained the corresponding transcription factor in the sense orientation under the constitutive super promoter.

Table 6

[0152] Listed are the names of the various constructs of the *Physcomitrella patens* transcription factors used for plant transformation

Gene	Enzymes used to generate gene fragment	Enzymes used to restrict pBPSJH001	Binary Vector Construct
PpCC-1	EcoRV/SacI	SmaI/SacI	pBPSSY032
PpCC-2	AscI/SacI	XmaI/SacI	pBPSsc335
PpCC-3	XmaI/SacI	XmaI/SacI	pBP SLVM186

Agrobacterium Transformation

[0153] The recombinant vectors were transformed into *Agrobacterium tumefaciens* C58C1 and PMP90 according to standard conditions (Hoefgen and Willmitzer, 1990).

Plant Transformation

[0154] *Arabidopsis thaliana* ecotype C24 were grown and transformed according to standard conditions (Bechtold 1993, Acad. Sci. Paris. 316:1194-1199; Bent et al. 1994, Science 265:1856-1860).

Screening of Transformed Plants

[0155] T1 seeds were sterilized according to standard protocols (Xiong et al. 1999, Plant Molecular Biology Reporter 17: 159-170). Seeds were plated on ½ MS 0.6% agar supplemented with 1% sucrose, 50 µg/ml kanamycin (Sigma-Aldrich) and 2 µg/ml benomyl (Sigma-Aldrich). Seeds on plates were vernalized for four days at 4°C. The seeds were germinated in a climatic chamber at an air temperature of 22°C and light intensity of 40 micromols⁻¹m² (white light; Philips TL 65W/25 fluorescent tube) and 16 hours light and 8 hours dark day length cycle. Transformed seedlings were selected after 14 days and transferred to ½ MS 0.6% agar plates supplemented with 1% sucrose and allowed to recover for five-seven days.

Drought Tolerance Screening

[0156] T1 seedlings were transferred to dry, sterile filter paper in a petri dish and allowed to desiccate for two hours at 80% RH (relative humidity) in a Sanyo Growth Cabinet MLR-350H, micromols⁻¹m² (white light; Philips TL 65W/25 fluorescent tube). The RH was then decreased to 60% and the seedlings were desiccated further for eight hours. Seedlings were then removed and placed on ½ MS 0.6% agar plates supplemented with 2 µg/ml benomyl and scored after five days.

[0157] Under drought stress conditions, PpCC-1 over-expressing *Arabidopsis thaliana* plants showed a 67% (6 survivors from 9 stressed plants) survival rate to the stress screening; PpCC-2, 75% (6 survivors from 8 stressed plants) and PpCC-3, 92% (11 survivors from 12 stressed plants), whereas the control only showed a 11% survival rate (1 survivor from 9 stressed plants) (see Table 7). It is noteworthy that the analyses of these transgenic lines were performed with T1 plants, and therefore, the results will be better when a homozygous, strong expresser is found.

Table 7

[0158]

Summary of the drought stress tests

Gene Name	Drought Stress Test		
	Number of survivors	Total number of plants	Percentage of survivors
PpCC-1	6	9	67%
PpCC-2	6	8	75%
PpCC-3	11	12	92%

Freezing Tolerance Screening

[0159] Seedlings were moved to petri dishes containing ½ MS 0.6% agar supplemented with 2% sucrose and 2 µg/ml benomyl. After four days, the seedlings were incubated at 4°C for 1 hour and then covered with shaved ice. The seedlings were then placed in an Environmental Specialist ES2000 Environmental Chamber and incubated for 3.5 hours beginning at -1.0°C decreasing -1°C hour. The seedlings were then incubated at -5.0°C for 24 hours and then allowed to thaw at 5°C for 12 hours. The water was poured off and the seedlings were scored after 5 days.

[0160] Under freezing stress conditions, PpCC-1 over-expressing *Arabidopsis thaliana* plants showed a 89% (8 survivors from 9 stressed plants) survival rate to the stress screening, whereas the untransformed control only showed a 2% survival rate (1 survivor from 48 stressed plants) (see Table 8). It is noteworthy that the analyses of these transgenic lines were performed with T1 plants, and therefore, the results will be better when a homozygous, strong expresser is found.

Table 8

Summary of the freezing stress tests

Gene Name	Freezing Stress Test		
	Number of survivors	Total number of plants	Percentage of survivors
PpCC-1	8	9	89%
Control	1	48	2%

Salt Tolerance Screening

[0161] Seedlings were transferred to filter paper soaked in ½ MS and placed on ½ MS 0.6% agar supplemented with 2 µg/ml benomyl the night before the salt tolerance screening. For the salt tolerance screening, the filter paper with the seedlings was moved to stacks of sterile filter paper, soaked in 50 mM NaCl, in a petri dish. After two hours, the filter paper with the seedlings was moved to stacks of sterile filter paper, soaked with 200 mM NaCl, in a petri dish. After two hours, the filter paper with the seedlings was moved to stacks of sterile filter paper, soaked in 600mM NaCl, in a petri dish. After 10 hours, the seedlings were moved to petri dishes containing ½ MS 0.6% agar supplemented with 2 µg/ml benomyl. The seedlings were scored after 5 days.

[0162] The transgenic plants are screened for their improved salt tolerance demonstrating that transgene expression confers salt tolerance.

Example 8*Detection of the CC-1, CC-2 and CC-3 transgenes in the transgenic Arabidopsis lines*

[0163] One leaf from a wild type and a transgenic *Arabidopsis* plant was homogenized in 250 µl Hexadecyltrimethyl ammonium bromide (CTAB) buffer (2% CTAB, 1.4 M NaCl, 8mM EDTA and 20mM Tris pH 8.0) and 1 µl β-mercaptoethanol. The samples were incubated at 60-65°C for 30 minutes and 250 µl of Chloroform was then added to each sample. The samples were vortexed for 3 minutes and centrifuged for 5 minutes at 18,000 x g. The supernatant was taken from each sample and 150 µl isopropanol was added. The samples were incubated at room temperature for 15 minutes, and centrifuged for 10 minutes at 18,000 x g. Each pellet was washed with 70% ethanol, dried, and resuspended in 20 µl TE. 4 µl of above suspension was used in a 20 µl PCR reaction using *Taq* DNA polymerase (Roche Molecular Biochemicals) according to the manufacturer's instructions. The primers used in the reactions were:

PpCC-1

GCTGACACGCCAAGCCTCGCTAGTC

(SEQ ID NO:25)

GCGAGCTCGCGACTGATGCAACTCACCCATGAA

(SEQ ID NO:26)

[0164] The PCR program was as following: 30 cycles of 1 minute at 94°C, 1 minute at 62°C and 4 minutes at 72°C, followed by 10 minutes at 72°C. A 1.6 kb fragment was produced from the positive control and the transgenic plants.

PpCC-2

GCTGACACGCCAAGCCTCGCTAGTC (SEQ ID NO:25)

GCGAGCTCTGGGAATTGCGCATGGGAAACTGG (SEQ ID NO:27)

[0165] The primers were used in the first round of reactions with the following program: 30 cycles of 1 minute at 94°C, 1 minute at 62°C and 4 minutes at 72°C, followed by 10 minutes at 72°C. A 3 kb fragment was generated from the positive control and the T1 transgenic plants.

PpCC-3

GCTGACACGCCAAGCCTCGCTAGTC (SEQ ID NO:25)

GCGAGCTCTGATACAAGAGCGTCGATGCTCCA (SEQ ID NO:28)

[0166] The PCR program was as following: 30 cycles of 1 minute at 94°C, 1 minute at 62°C and 4 minutes at 72°C, followed by 10 minutes at 72°C. A 2.3 kb fragment was produced from the positive control and the transgenic plants..

[0167] The transgenes were successfully amplified from the T1 transgenic lines, but not from the wild type C24. This result indicates that the T1 transgenic plants contain at least one copy of the transgenes. There was no indication of existence of either identical or very similar genes in the untransformed *Arabidopsis thaliana* control that could be amplified by this method.

Example 9

Detection of the CC-1, CC-2 and CC-3 transgene mRNA in transgenic Arabidopsis lines

[0168] Transgene expression was detected using RT-PCR. Total RNA was isolated from stress-treated plants using a procedure adapted from (Verwoerd et al., 1989 NAR 17:2362). Leaf samples (50-100 mg) were collected and ground to a fine powder in liquid nitrogen. Ground tissue was resuspended in 500 µl of a 80°C, 1:1 mixture, of phenol to extraction buffer (100mM LiCl, 100 mM Tris pH8, 10 mM EDTA, 1% SDS), followed by

brief vortexing to mix. After the addition of 250 µl of chloroform, each sample was vortexed briefly. Samples were then centrifuged for 5 minutes at 12,000 x g. The upper aqueous phase was removed to a fresh eppendorf tube. RNA was precipitated by adding 1/10th volume 3M sodium acetate and 2 volumes 95% ethanol. Samples were mixed by inversion and placed on ice for 30 minutes. RNA was pelleted by centrifugation at 12,000 x g for 10 minutes. The supernatant was removed and pellets briefly air-dried. RNA sample pellets were resuspended in 10 µl DEPC treated water. To remove contaminating DNA from the samples, each was treated with RNase-free DNase (Roche) according to the manufacturer's recommendations. cDNA was synthesized from total RNA using the 1st Strand cDNA synthesis kit (Boehringer Mannheim) following manufacturer's recommendations.

[0169] PCR amplification of a gene-specific fragment from the synthesized cDNA was performed using *Taq* DNA polymerase (Roche) and gene-specific primers (see below for primers) in the following reaction: 1X PCR buffer, 1.5mM MgCl₂, 0.2 µM each primer, 0.2µM dNTPs, 1 unit polymerase, 5µl cDNA from synthesis reaction. Amplification was performed under the following conditions: Denaturation, 95°C, 1 minute; annealing, 62°C, 30 seconds; extension, 72°C, 1 minute, 35 cycles; extension, 72°C, 5 minutes; hold, 4°C, forever. PCR products were run on a 1% agarose gel, stained with ethidium bromide, and visualized under UV light using the Quantity-One gel documentation system (Bio-Rad).

[0170] Expression of the transgenes was detected in the T1 transgenic line. This result indicated that the transgenes are expressed in the transgenic lines and strongly suggested that their gene product improved plant stress tolerance in the transgenic line. On the other hand, no expression of identical or very similar endogenous genes could be detected by this method. These results are in agreement with the data from Example 7. This greatly supports our statement that the observed stress tolerance is due to the introduced transgene.

[0171] The following primer combination was used in the above-described reactions.

PpCC-1:

GATGGTGATGGTGACAGCGAGGATC	(SEQ ID NO:29)
CGACGTGAAGCCTCGCTCTCATTTCC	(SEQ ID NO:30)

PpCC-2:

CGATCTCCTTCAAGGGAACCTAGTGCTG	(SEQ ID NO:31)
GAGTTCACGCATGTGCACCGATGCT	(SEQ ID NO:32)

PpCC-3:

CGTAGCGTCTTCCAGGATGTCCTAC

(SEQ ID NO:33)

GCTTGGCCTCGTCTACTCCCGCAAC

(SEQ ID NO:34)

[0172] Amplification was performed under the following conditions: Denaturation, 95°C, 1 minute; annealing, 62°C, 30 seconds; extension, 72°C, 1 minute, 35 cycles; extension, 72°C, 5 minutes; hold, 4°C, forever. PCR products were run on a 1% agarose gel, stained with ethidium bromide, and visualized under UV light using the Quantity-One gel documentation system (Bio-Rad).

[0173] Expression of the transgenes was detected in the T1 transgenic line. These results indicated that the transgenes are expressed in the transgenic lines and strongly suggested that their gene product improved plant stress tolerance in the transgenic lines. In agreement with the previous statement, no expression of identical or very similar endogenous genes could be detected by this method. These results are in agreement with the data from Example 7.

Example 10

Engineering stress-tolerant soybean plants by over-expressing the CC-1, CC-2 and CC-3 gene

[0174] The constructs pBPSSY032, pBPSSc335 and pBPSSLVM186 were used to transform soybean as described below.

[0175] Seeds of soybean were surface sterilized with 70% ethanol for 4 minutes at room temperature with continuous shaking, followed by 20% (v/v) Clorox supplemented with 0.05% (v/v) Tween for 20 minutes with continuous shaking. Then, the seeds were rinsed 4 times with distilled water and placed on moistened sterile filter paper in a Petri dish at room temperature for 6 to 39 hours. The seed coats were peeled off, and cotyledons are detached from the embryo axis. The embryo axis was examined to make sure that the meristematic region is not damaged. The excised embryo axes were collected in a half-open sterile Petri dish and air-dried to a moisture content less than 20% (fresh weight) in a sealed Petri dish until further use.

[0176] *Agrobacterium tumefaciens* culture was prepared from a single colony in LB solid medium plus appropriate antibiotics (e.g. 100 mg/l streptomycin, 50 mg/l kanamycin) followed by growth of the single colony in liquid LB medium to an optical density at 600 nm of 0.8. Then, the bacteria culture was pelleted at 7000 rpm for 7 minutes at room temperature, and resuspended in MS (Murashige and Skoog, 1962) medium supplemented with 100 µM

acetosyringone. Bacteria cultures were incubated in this pre-induction medium for 2 hours at room temperature before use. The axis of soybean zygotic seed embryos at approximately 15% moisture content were imbibed for 2 hours at room temperature with the pre-induced *Agrobacterium* suspension culture. The embryos are removed from the imbibition culture and were transferred to Petri dishes containing solid MS medium supplemented with 2% sucrose and incubated for 2 days, in the dark at room temperature. Alternatively, the embryos were placed on top of moistened (liquid MS medium) sterile filter paper in a Petri dish and incubated under the same conditions described above. After this period, the embryos were transferred to either solid or liquid MS medium supplemented with 500 mg/L carbenicillin or 300mg/L cefotaxime to kill the agrobacteria. The liquid medium was used to moisten the sterile filter paper. The embryos were incubated during 4 weeks at 25°C, under $150 \mu\text{mol m}^{-2}\text{sec}^{-1}$ and 12 hours photoperiod. Once the seedlings produced roots, they were transferred to sterile metromix soil. The medium of the *in vitro* plants was washed off before transferring the plants to soil. The plants were kept under a plastic cover for 1 week to favor the acclimatization process. Then the plants were transferred to a growth room where they were incubated at 25°C, under $150 \mu\text{mol m}^{-2}\text{sec}^{-1}$ light intensity and 12 hours photoperiod for about 80 days.

[0177] The transgenic plants were then screened for their improved drought, salt and/or cold tolerance according to the screening method described in Example 7 demonstrating that transgene expression confers stress tolerance.

Example 11

Engineering stress-tolerant Rapeseed/Canola plants by over-expressing the CC-1, CC-2 and CC-3 genes

[0178] The constructs pBPSSY032, pBPSsc335 and pBPSLVM186 were used to transform rapeseed/canola as described below.

[0179] The method of plant transformation described herein is also applicable to *Brassica* and other crops. Seeds of canola are surface sterilized with 70% ethanol for 4 minutes at room temperature with continuous shaking, followed by 20% (v/v) Clorox supplemented with 0.05 % (v/v) Tween for 20 minutes, at room temperature with continuous shaking. Then, the seeds are rinsed 4 times with distilled water and placed on moistened sterile filter paper in a Petri dish at room temperature for 18 hours. Then the seed coats are removed and the seeds are air dried overnight in a half-open sterile Petri dish. During this period, the seeds lose approx. 85% of its water content. The seeds are then stored at room

temperature in a sealed Petri dish until further use. DNA constructs and embryo imbibition are as described in Example 10. Samples of the primary transgenic plants (T0) are analyzed by PCR to confirm the presence of T-DNA. These results are confirmed by Southern hybridization in which DNA is electrophoresed on a 1% agarose gel and transferred to a positively charged nylon membrane (Roche Diagnostics). The PCR DIG Probe Synthesis Kit (Roche Diagnostics) is used to prepare a digoxigenin-labelled probe by PCR, and used as recommended by the manufacturer.

[0180] The transgenic plants are then screened for their improved stress tolerance according to the screening method described in Example 7 demonstrating that transgene expression confers drought tolerance.

Example 12

Engineering stress-tolerant corn plants by over-expressing the CC-1, CC-2 and CC-3 genes

[0181] The constructs pBPSSY032, pBPSSc335 and pBPSSLVM186 were used to transform corn as described below.

[0182] Transformation of maize (*Zea Mays L.*) is performed with the method described by Ishida et al. 1996. Nature Biotech 14745-50. Immature embryos are co-cultivated with *Agrobacterium tumefaciens* that carry "super binary" vectors, and transgenic plants are recovered through organogenesis. This procedure provides a transformation efficiency of between 2.5% and 20%. The transgenic plants are then screened for their improved drought, salt and/or cold tolerance according to the screening method described in Example 7 demonstrating that transgene expression confers stress tolerance.

Example 13

Engineering stress-tolerant wheat plants by over-expressing the CC-1, CC-2 and CC-3 genes

[0183] The constructs pBPSSY032, pBPSSc335 and pBPSSLVM186 were used to transform wheat as described below.

[0184] Transformation of wheat is performed with the method described by Ishida et al. 1996 Nature Biotech. 14745-50. Immature embryos are co-cultivated with *Agrobacterium tumefaciens* that carry "super binary" vectors, and transgenic plants are recovered through organogenesis. This procedure provides a transformation efficiency between 2.5% and 20%. The transgenic plants are then screened for their improved stress tolerance according to the screening method described in Example 7 demonstrating that transgene expression confers drought tolerance.

Example 14

Identification of Homologous and Heterologous Genes

[0185] Gene sequences can be used to identify homologous or heterologous genes from cDNA or genomic libraries. Homologous genes (e. g. full-length cDNA clones) can be isolated via nucleic acid hybridization using for example cDNA libraries. Depending on the abundance of the gene of interest, 100,000 up to 1,000,000 recombinant bacteriophages are plated and transferred to nylon membranes. After denaturation with alkali, DNA is immobilized on the membrane by e. g. UV cross linking. Hybridization is carried out at high stringency conditions. In aqueous solution hybridization and washing is performed at an ionic strength of 1 M NaCl and a temperature of 68°C. Hybridization probes are generated by e. g. radioactive (^{32}P) nick transcription labeling (High Prime, Roche, Mannheim, Germany). Signals are detected by autoradiography.

[0186] Partially homologous or heterologous genes that are related but not identical can be identified in a manner analogous to the above-described procedure using low stringency hybridization and washing conditions. For aqueous hybridization, the ionic strength is normally kept at 1 M NaCl while the temperature is progressively lowered from 68 to 42°C.

[0187] Isolation of gene sequences with homologies (or sequence identity/similarity) only in a distinct domain of (for example 10-20 amino acids) can be carried out by using synthetic radio labeled oligonucleotide probes. Radio labeled oligonucleotides are prepared by phosphorylation of the 5-prime end of two complementary oligonucleotides with T4 polynucleotide kinase. The complementary oligonucleotides are annealed and ligated to form concatemers. The double stranded concatemers are then radiolabeled by, for example, nick transcription. Hybridization is normally performed at low stringency conditions using high oligonucleotide concentrations.

Oligonucleotide hybridization solution:

6 x SSC

0.01 M sodium phosphate

1 mM EDTA (pH 8)

0.5 % SDS

100 µg/ml denatured salmon sperm DNA

0.1 % nonfat dried milk

[0188] During hybridization, temperature is lowered stepwise to 5-10°C below the estimated oligonucleotide T_m or down to room temperature followed by washing steps and autoradiography. Washing is performed with low stringency such as 3 washing steps using 4 x SSC. Further details are described by Sambrook, J. *et al.* (1989), "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory Press or Ausubel, F.M. *et al.* (1994) "Current Protocols in Molecular Biology", John Wiley & Sons.

Example 15

Identification of Homologous Genes by Screening Expression Libraries with Antibodies

[0189] c-DNA clones can be used to produce recombinant protein for example in *E. coli* (e. g. Qiagen QLAexpress pQE system). Recombinant proteins are then normally affinity purified via Ni-NTA affinity chromatography (Qiagen). Recombinant proteins are then used to produce specific antibodies for example by using standard techniques for rabbit immunization. Antibodies are affinity purified using a Ni-NTA column saturated with the recombinant antigen as described by Gu *et al.*, 1994 *BioTechniques* 17:257-262. The antibody can then be used to screen expression cDNA libraries to identify homologous or heterologous genes via an immunological screening (Sambrook, J. *et al.* (1989), "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory Press or Ausubel, F.M. *et al.* (1994) "Current Protocols in Molecular Biology", John Wiley & Sons).

Example 16

In vivo Mutagenesis

[0190] *In vivo* mutagenesis of microorganisms can be performed by passage of plasmid (or other vector) DNA through *E. coli* or other microorganisms (e.g. *Bacillus* spp. or yeasts such as *Saccharomyces cerevisiae*) which are impaired in their capabilities to maintain the integrity of their genetic information. Typical mutator strains have mutations in the genes for the DNA repair system (e.g., *mutHLS*, *mutD*, *mutT*, etc.; for reference, see Rupp, W.D. (1996) DNA repair mechanisms, in: *Escherichia coli* and *Salmonella*, p. 2277-2294, ASM: Washington.) Such strains are well known to those skilled in the art. The use of such strains is illustrated, for example, in Greener, A. and Callahan, M. (1994) *Strategies* 7: 32-34. Transfer of mutated DNA molecules into plants is preferably done after selection and testing in microorganisms. Transgenic plants are generated according to various examples within the exemplification of this document.

Example 17

In vitro Analysis of the Function of Physcomitrella Genes in Transgenic Organisms

[0191] The determination of activities and kinetic parameters of enzymes is well established in the art. Experiments to determine the activity of any given altered enzyme must be tailored to the specific activity of the wild-type enzyme, which is well within the ability of one skilled in the art. Overviews about enzymes in general, as well as specific details concerning structure, kinetics, principles, methods, applications and examples for the determination of many enzyme activities may be found, for example, in the following references: Dixon, M., and Webb, E.C., (1979) *Enzymes*. Longmans: London; Fersht, (1985) *Enzyme Structure and Mechanism*. Freeman: New York; Walsh, (1979) *Enzymatic Reaction Mechanisms*. Freeman: San Francisco; Price, N.C., Stevens, L. (1982) *Fundamentals of Enzymology*. Oxford Univ. Press: Oxford; Boyer, P.D., ed. (1983) *The Enzymes*, 3rd ed. Academic Press: New York; Bisswanger, H., (1994) *Enzymkinetik*, 2nd ed. VCH: Weinheim (ISBN 3527300325); Bergmeyer, H.U., Bergmeyer, J., Graßl, M., eds. (1983-1986) *Methods of Enzymatic Analysis*, 3rd ed., vol. I-XII, Verlag Chemie: Weinheim; and Ullmann's *Encyclopedia of Industrial Chemistry* (1987) vol. A9, *Enzymes*. VCH: Weinheim, p. 352-363.

[0192] The activity of proteins which bind to DNA can be measured by several well-established methods, such as DNA band-shift assays (also called gel retardation assays). The effect of such proteins on the expression of other molecules can be measured using reporter gene assays (such as that described in Kolmar, H. et al. (1995) *EMBO J.* 14: 3895-3904 and references cited therein). Reporter gene test systems are well known and established for applications in both pro- and eukaryotic cells, using enzymes such as β -galactosidase, green fluorescent protein, and several others.

[0193] The determination of activity of membrane-transport proteins can be performed according to techniques such as those described in Gennis, R.B. Pores, Channels and Transporters, in *Biomembranes, Molecular Structure and Function*, pp. 85-137, 199-234 and 270-322, Springer: Heidelberg (1989).

Example 18

Purification of the Desired Product from Transformed Organisms

[0194] Recovery of the desired product from plant material (i.e., *Physcomitrella patens* or *Arabidopsis thaliana*), fungi, algae, ciliates, *C. glutamicum* cells, or other bacterial cells transformed with the nucleic acid sequences described herein, or the supernatant of the above-described cultures can be performed by various methods well known in the art. If the desired product is not secreted from the cells, can be harvested from the culture by low-speed centrifugation, the cells can be lysed by standard techniques, such as mechanical force or sonification. Organs of plants can be separated mechanically from other tissue or organs. Following homogenization cellular debris is removed by centrifugation, and the supernatant fraction containing the soluble proteins is retained for further purification of the desired compound. If the product is secreted from desired cells, then the cells are removed from the culture by low-speed centrifugation, and the supernate fraction is retained for further purification.

[0195] The supernatant fraction from either purification method is subjected to chromatography with a suitable resin, in which the desired molecule is either retained on a chromatography resin while many of the impurities in the sample are not, or where the impurities are retained by the resin while the sample is not. Such chromatography steps may be repeated as necessary, using the same or different chromatography resins. One skilled in the art would be well-versed in the selection of appropriate chromatography resins and in their most efficacious application for a particular molecule to be purified. The purified product may be concentrated by filtration or ultrafiltration, and stored at a temperature at which the stability of the product is maximized.

[0196] There is a wide array of purification methods known to the art and the preceding method of purification is not meant to be limiting. Such purification techniques are described, for example, in Bailey, J.E. & Ollis, D.F. *Biochemical Engineering Fundamentals*, McGraw-Hill: New York (1986). Additionally, the identity and purity of the isolated compounds may be assessed by techniques standard in the art. These include high-performance liquid chromatography (HPLC), spectroscopic methods, staining methods, thin layer chromatography, NIRS, enzymatic assay, or microbiologically. Such analysis methods are reviewed in: Patek et al., 1994 *Appl. Environ. Microbiol.* 60:133-140; Malakhova et al., 1996 *Biotechnologiya* 11:27-32; and Schmidt et al., 1998 *Bioprocess Engineer.* 19:67-70. *Ulmann's Encyclopedia of Industrial Chemistry*, (1996) vol. A27, VCH: Weinheim, p. 89-90, p. 521-540, p. 540-547, p. 559-566, 575-581 and p. 581-587; Michal, G. (1999) *Biochemical*

Pathways: An Atlas of Biochemistry and Molecular Biology, John Wiley and Sons; Fallon, A. et al. (1987) Applications of HPLC in Biochemistry in: Laboratory Techniques in Biochemistry and Molecular Biology, vol. 17.

SEQUENCE LISTING

<110> BASF PLANT SCIENCE GMBH

<120> CELL CYCLE STRESS-RELATED PROTEINS AND METHODS OF USE
IN PLANTS

<130> 16313-0035

<140> PCT/US01/11294

<141> 2001-04-06

<150> 60/196,001

<151> 2000-04-07

<160> 34

<170> PatentIn Ver. 2.1

<210> 1

<211> 887

<212> DNA

<213> *Physcomitrella patens*

<220>

<221> modified_base

<222> (71)

<223> a, t, c, g, other or unknown

<400> 1

cgggagttgg tgatcttcag catgctcgta gttgacaagg gaggtaggat tgtttgaccc 60
aaggtgtgta nagaagggga tagccatgta cagcaacttc aaagagcagg ccatagaata 120
ttgtgcgtca agccgtagcg gaagacaacg caggggaacta tgccaaagcg tttccgctgt 180
acatgaacgc gcttgagtac ttcaagacgc atctaaagta tgagaagaat cccaaaatca 240
aggaggccat cactcagaag ttcacggagt atttgaggag ggcggaggag attcgagccg 300
ttttggacga tggccccact ggaccctctg caaatggaga cgcggcagtt caagctaaac 360
cgaagtcgaa atcaggggaag aaggatggtg gcggggggtga tggatgatggt gacagcgagg 420
atccccacca gcagaagctg agatcagggc tgaactcggc aatcatacgg gaaaagccaa 480
atgttcggtg ggctgatggt gctggacttg aaagtgccaa gcaggcgttg caggaggcag 540
tgatcttgcc cgtgaagttt cccaatttt tcacaggga gcaagaaca tggcgagcat 600
ttttgtggtg tgggcccccc gggactggaa aatcgatatc tgcaaaagct gttgctacgg 660
aagctgattc tacattcttt agtatttcct cttcagactt ggtgtcaaag tggatgggag 720
agagtgagaa gcttggtgca aatctgtttc aaatggcccc tgaagctgct ccatccatca 780
tcttcataga cgagattgat tctttatgcg gtactcgagg tgaaggaaat gagagcgagg 840
cttcacgtcg tatcaagact gagttgctag ttcaaagca ggggtgct 887

<210> 2
 <211> 723
 <212> DNA
 <213> *Physcomitrella patens*

<400> 2
 cggcaccagg gagatcgtat tgaggtaaca ggagttttca aggccatggc agttcgagtt 60
 ggtccgaatc aacgaacatt acgagcattg tataagacct acatcgattg cgtgcacgtc 120
 aagaagtctg acaggggtcg actgcaaact gaagatccta tggagatgga taaggagaat 180
 gatatgtatg ctgggtatca tgaagtgat acttcagaag ctgctaata agcaaagatt 240
 caaaaactta aagagctgtc caagctcccg gacatttatg atagactttc aaggctcgctg 300
 gctccaagca tttgggagct tgaagatatt aaaaagggtc ttctttgcca gctctttggt 360
 gggaaggcta agaaaattcc atctggagca tctttccgag gtgacatcaa tgttttactt 420
 gttggggacc ctggtaccag taaatctcag ctgcttcagt atgtgcacaa gatagctcct 480
 cgtggaatct acactagtgg gcgaggaagt tcggcggttg ggctgacagc gtatgtaaac 540
 gaaggatcca gaaactcgag agacggtatt ggagagcgga gctttgggtc ttagtgatcg 600
 tgggatatgc tgtatcgatg agttcgacaa aatgtctgat aatgcccga gcatgcttca 660
 tgaggtaatg gagcaacaaa cggatatctg acccaagcgg ttcattgctcg tgaagccgag 720
 ttg 723

<210> 3
 <211> 566
 <212> DNA
 <213> *Physcomitrella patens*

<400> 3
 gcaccagccg ctttggaatc ccatccctcg gttgcataga cacaagggga ttcagtgtag 60
 tgatacgttg tgcataattg gtgttgcaag atttttgggt tcttgattgt tagctatggc 120
 gtctgcaaca gcggctacaa tggcgctcct cctcacgcct gggctctctcc gacgcggttt 180
 gggtagccag gaatcgtcga cccaatttgc tcccctagct ggtcctcgta agacatcagt 240
 ttcgcgtagg gtgactgcta gcgctagtgg gaagaacgac aatggagtcg tggaagatgt 300
 ggatatgggg aagcggggta tggtgaaagg cgtagcggga gctttggctg cagttctccc 360
 tgctgttatc gcgaagaaag cttcagcagc tgaggagcag ggcgtagcgt cttccaggat 420
 gtcctactcg aggtttttgg agtatattgga tatggaccgt gtgaagaagt tgacttgtat 480
 gaaaatggga ccatagcaat tgtggaggct gtatcccctg aattgggcaa cagagtgcaa 540
 cgcgtacgcg tgcagctccc cggaac 566

<210> 4
 <211> 1564
 <212> DNA
 <213> *Physcomitrella patens*

<400> 4
 gcgatatcga cccaaggtgt gtagagaagg ggatagccat gtacagcaac ttcaaagagc 60
 aggccataga atatgtgcgt caagccgtag cggaagacaa cgcagggaac tatgccaaag 120
 cgtttccgct gtacatgaac gcgcttgagt acttcaagac gcatctaaag tatgagaaga 180
 atccccaaat caaggaggcc atcactcaga agttcacgga gtatttgagg agggcggagg 240
 agattcgagc cgttttggac gatggcccca ctggaccctc tgcaaattga gacgcggcag 300
 ttcaagctaa accgaagtcg aaatcaggga agaaggatgg tggcgggggt gatggtgatg 360
 gtgacagcga ggatcccgag cagcagaagc tgagatcagg gctgaactcg gcaatcatac 420
 gggaaaagcc aaatgttcgg tgggctgatg ttgctggact tgaaagtgcc aagcaggcgt 480
 tgcaggaggc agtgatcttg cccgtgaagt ttccccaatt tttcacaggg aagcgaagac 540
 catggcgagc atttttgttg tatgggcccc ccgggactgg aaaatcgtat cttgcaaaag 600
 ctgttgctac ggaagctgat tctacattct ttagtatttc ctcttcagac ttggtgtcaa 660
 agtggatggg agagagtgag aagcttggtg caaatctgtt tcaaattggc cgtgaagctg 720
 ctccatccat catcttcata gacgagattg attctttatg cgg tactcga ggtgaaggaa 780
 atgagagcga ggcttcacgt cgtatcaaga ctgagttgct agttcaaatt caggggtgtcg 840
 gcaatcaaga cactaagggt cttgtgttag ctgctacaaa tacgccctac tctttggatc 900
 aggcggtgag gcgacgtttc gacaagcgta tctacatccc actaccggag tctaaggctc 960
 ggcagcacat gtttaagggt catttgggag atacgcaaaa caacctgact gaacgtgatt 1020
 atgaggatct ggctaggaag actgatgggt tttcaggctc ggatattgca gtttgtgtga 1080
 aagatgtact atttgagcct gttcgtaaga cccaagatgc tatgcatttc aaaagaatta 1140
 ataccaaaga aggagagatg tggatgcctt gtgggccccg agaaccagggt gctaggcaaa 1200
 ccactatgac ggagcttgcc gctgaagggc aggcacgaa gattttacca cctccaatca 1260
 caaaatcaga tttcgacaaa gtcctcgcaa agcagaggcc cactgtcagc aaaggcgacc 1320
 ttattattca agagaaattc accaaagaat ttggtgaaga aggttgaatg gtgcttcgag 1380
 ttaagaattt ggaaggttct gggttacgga agacagacga aatagaacgt cgtaggtacg 1440
 gtagcctaag agtaaattac ggaagttttc cgacttgcca gttgtgcact ctttcaacat 1500
 gaagggaagg aagctacttg tcggttgcc tttcatgggt gagttgcac agtcgcgagc 1560
 tcgc 1564

<210> 5
 <211> 4348
 <212> DNA
 <213> *Physcomitrella patens*

<400> 5
 atggcgcgcc gcactcacgt gaggaattgc acctccttgt tctgcgacgg ttccattctt 60
 tttggttttt agtttgcaaa tcttgatcgt ggagttgaga aaaagggcgg ttcgttgtct 120
 tgaggtgttc ttgttgattg ttcgtcatgg aaaataatga tgcacttgac attggagccg 180
 tgtcgtcccc atatccttcg caatctgaag gagtgtctac gccattgccg caagtaacat 240
 caccgagctt cgacaatgca gcctcacccg tggccggggc gagggccgta cggcagaccc 300
 ctacatctgc agttcgaagg agagggagag aaacggattc cgctcgtcgt aggaggagtc 360
 gatctcgcag tttaggcaat tctgtttata gttcccccta cgatgcgggg actcctggaa 420
 ctctggaac tccagtggct actccggttt acgctacccc agtcggtaca cctatgggta 480
 ccccatcgtt ccatcgtggc acgccacagt acaaacagcg cagtgcgctt ggttcccagg 540
 ggaagcctct acatcggaga cgtcgatctc aatccagaga acccgggcat cgatctcctt 600
 caaggggaacc tagtgctgat gggcgtccct ctgaatctgc tgagccagat gacactttgg 660
 gtggagaata tgcttatgtt tgggggacga atgttaacat tccagatgtg cttagggcga 720
 ttcgtcgatt tctccacaat tatcgttcga gtgctcatga tcttaattcc aagtacatcc 780
 agatcataga ggagactgtg gagcgtgagg aggatactct aaatatcgac atgtcagaca 840
 tttatgacca tgatcctgat ctatacgcaa aaattgttcg ataccactc gacatcatcc 900
 ccctgttgga cactgagtgt caggaagtgt ctacctctt actaccaacg tttgagaagc 960
 atattgaggg cagacctttc aatctcaaag catcggtgca catgcgtgaa ctcaaccctt 1020
 cagatataga caaattgggt tctgttaaag gaatgggttat ccggtgcagt tctatcatac 1080
 ctgaaattaa gggggccttc ttcaaagtgt tagtggtgtg tctactcgcct ccgctagtta 1140
 cagttgttaa agggcgggtt gaggagccaa caaggtgtga aaagccagaa tgtgcagcac 1200
 ggaatgctat gtctcttatt cacaatcgat gcacttttgc aaataagcag atagtgcgtc 1260
 ttcaagaaac tccagatgcc attcctgaag gagagactcc acacacagtc agcatgtgtt 1320
 tatacaacac tatggttgat gctgtgaagc ctggagatcg tattgaggta acaggagttt 1380
 tcaaggccat ggcagttcga gttgggtcatg gcgcgccgca ctcacgtgag gaattgcacc 1440
 tccttgttct gcgacgggtc cattcttttt ggtttttagt ttgcaaattc tgatcgtgga 1500
 gttgagaaaa agggcgggtc gttgtcttga ggtgttcttg ttgattgttc gtcattgaaa 1560
 ataatgatgc acttgacatt ggagccgtgt cgtccccata tccttcgcaa tctgaaggag 1620
 tgtctacgcc attgccgcaa gtaacatcac cgagcttcga caatgcagcc tcacccgtgg 1680

ccgggaggag ggccgtacgg cagaccccta catctgcagt tcgaaggaga gggagagaaa 1740
 cggattccgc tcgtcgtagg aggagtcgat ctgcagttt aggcaattct gtttatagtt 1800
 ccccttacga tgcggggact cctggaactc ctggaactcc agtggctact ccggtttacg 1860
 ctaccccagt cggtagacct atgggtaccc catcggtcca tcgtggcacg ccacagtaca 1920
 aacagcgcag tgagcttggt tcccagggga agcctctaca tcggagacgt cgatctcaat 1980
 ccagagaacc cgggcatcga tctccttcaa gggaaacctag tgctgatggg cgtccctctg 2040
 aatctgctga gccagatgac actttgggtg gagaatatgc ttatgtttgg gggacgaatg 2100
 ttaacattcc agatgtgctt agggcgattc gtcgatttct ccacaattat cgttcgagtg 2160
 ctcatgatct taattccaag tacatccaga tcatagagga gactgtggag cgtgaggagg 2220
 atactctaaa tatcgacatg tcagacattt atgaccatga tcctgatcta tacgcaaaaa 2280
 ttgttcgata cccactcgac atcatcccc tggtggacac tgagtgtcag gaagttgcta 2340
 cctctttact accaacgttt gagaagcata ttgaggccag acctttcaat ctcaaagcat 2400
 cgggtcacat gcgtgaactc aacccttcag atatagacaa attggtttct gttaaaggaa 2460
 tggttatccg gtgcagttct atcatacctg aaattaaggg ggccttcttc aaatgtttag 2520
 tgtgtggtca ctgcctccg ctagttacag ttgttaaagg gcgggttgag gagccaacaa 2580
 ggtgtgaaaa gccagaatgt gcagcacgga atgctatgtc tcttattcac aatcgatgca 2640
 cttttgcaaa taagcagata gtgcgtcttc aagaaactcc agatgccatt cctgaaggag 2700
 agactccaca cacagtcagc atgtgtttat acaacactat ggttgatgct gtgaagcctg 2760
 gagatcgtat tgaggtaaca ggagttttca aggccatggc agttcgagtt ggtccgaatc 2820
 aacgaacatt acgagcattg tataagacct acatcgattg cgtgcacgtc aagaagtctg 2880
 acaggggtcg actgcaaact gaagatccta tggagatgga taaggagaat gatatgtatg 2940
 ctgggtatca tgaaagtgat acttcagaag ctgctaataga agcaaagatt caaaaactta 3000
 aagagctgtc caagctcccg ggcatttatg atagactttc aaggctcgctg gctccaagca 3060
 tttgggagct tgaagatatt aaaaagggtc ttctttgcca gctctttggt gggaaggcta 3120
 agaaaattcc atctggagca tctttccgag gtgacatcaa tgttttactt gttggggacc 3180
 ctggtaccag taaatctcag ctgcttcagt atgtgcacaa gatagctcct cgtggaatct 3240
 acactagtgg gcgaggaagt tcggcggttg ggctgacagc gtatgtaacg aaggatccag 3300
 aaactcgaga gacggtattg gagagcggag ctttggttct tagtgatcgt gggatatgct 3360
 gtatcgatga gttcgacaaa atgtctgata atgcccgaag catgcttcat gaggtaatgg 3420
 agcaacaaac ggtatctgta gccaaagggg gtatcattgc ctgcgtgaac gctcggacgt 3480

ctgtccttgc atgtgcaa at cctagtgggt cccgatacaa tgcgcgcctt tctgtgattg 3540
 ataacatcca gcttcctcca actctacttt ctagatttga ttttaatttac ttaatgctcg 3600
 acaaaccaga cgagcaaaac gatcgtcgtc tcgccaggca tctcgtgggt ttacactatg 3660
 aaaactatga agtttcaaag caggacgcct tagatctaca aacacttacc gcgtatatca 3720
 cctatgctcg tcagcatgta catcctacat taagtgatga agctgctgaa gatttgatta 3780
 atggctatgt tgagatgcgc caaaagggca actttcctgg aagcagtaaa aaggtgataa 3840
 cagccacacc tcggcaactc gaaagtatga ttcgtatcag tgaagcccta gctcgaatga 3900
 gattttctga agtggttagag aaagttgatg cagcagaagc tgtgcgcctt ttagacgtcg 3960
 ctttgcagca atctgctact gatcatgcaa caggtacgat agacatggat cttatcacga 4020
 ctggagtgtc ggccagcgag cgtattcgtc gggccaactt gctagctgct ctgcgagagc 4080
 ttatagcaga taaaatttca cctggcagct cctctggctt gaagaccagt cagcttcttg 4140
 aggatatccg gagccaaagc agtgtggacg ttagtttgca ggatattaaa aatgctctgg 4200
 gtagcctcca aggagaaggc tttcttactg tccatggtga catagtcaag agagtttgag 4260
 acagtttcta actgttcgaa tccatgagct ataactctga acgaaaggga aaacctccag 4320
 tttcccatgc gcaattccca gagctcgc 4348

<210> 6
 <211> 2237
 <212> DNA
 <213> *Physcomitrella patens*

<400> 6
 atcccgggtg atacgttgtg catatttgggt gttgcaagtt ttttggtttc ttgattgtta 60
 gctatggcgt ctgcaacagc ggctacaatg gcgtccctcc tcacgcctgg gtctctccga 120
 cgcggtttgg gtagccagga atcgtcgacc caatttgctc ccctagctgg tcctcgtaag 180
 acatcagttt cgcgtagggg gactgctagc gctagtggga agaacgacaa tggagtcgtg 240
 gaagatgtgg atatggggaa gcgggggtatg ttgaaaggcg tagcgggagc tttggctgca 300
 gttctccctg ctgttatcgc gaagaaagct tcagcagctg aggagcaggg cgtagcgtct 360
 tccaggatgt cctactcgag gtttttggag tatttggata tggaccgtgt gaagaagggt 420
 gacttgtatg aaaatgggac catagcaatt gtggaggctg tatccctga attgggcaac 480
 agagtacaac gcgtacgcgt gcagctcccc ggaactagct ccgaattgtt gtcgaagttt 540
 agatcgaaga atgtagattt tgccgcacac agcccacaag aggactctgg ctctgtcatt 600
 ttgaacctca tcggaaattt ggctttcccc ttgttgctcg ttggaggtct gttcttcttg 660
 tctcgtagat cccaagggtg tatgggacct ggcggtcctg ggaacccgat ggccttcggg 720

aagtccaagg ccaagttcca gatggagccc aacacgggca ttacattcca agatgttgcg 780
ggagtagacg aggccaagca agacttcatg gaggttgtgg agttcttgaa gcggcctgag 840
agattcacag cagtgggcg ctaaaatccca aaggagtggt tgctgggttg accacccggt 900
accggaaga ctctattggc gaaggccatc gctggggaag ctggagtacc attcttctcc 960
atctctgggt ctgagttcgt ggaaatgttc gtgggagtgg gagcttcccg tgtgaggagac 1020
ttgttcaaga aggcgaaaga gaatgctccc tgcattgtgt ttgtggatga gattgatgcc 1080
gttggaagac agagaggaac tggaattgga ggaggcaatg atgagcgtga gcagacgttg 1140
aatcagttgt tgacggagat ggacggtttc gaaggaaaca ctggtgtgat tgtcattgct 1200
gccaccaaca gggctgatat tctcgacgct gccttgcttc gtctggaag attcgacaga 1260
caggtttccg tggatgttcc ggacgtgaag ggaaggactg acatcctcaa ggtgcatgct 1320
agtaacaaga agttcgccga tgatgtgtct ctggatatca ttgccatgag gacaccaggg 1380
ttcagtggag ctgacttggc caacttggtg aacgaagcag ctatcttgac cggaaggaga 1440
ggaaagacag ccatcagtgc taaggagatt gacgattcaa ttgacagaat cgttgccggg 1500
atggaaggta ccgtcatgac ggacggcaag agcaagagtc ttgttgcata ccacgaagtc 1560
ggccatgcta tctgcgggac ttgactccc ggacacgacg ccgtgcaaaa ggtcaccctc 1620
attcctagag gtcaggcccc tgggtctcacc tggtttattc ccggagaaga cccgactttg 1680
atttcaaagc agcaaattct tgcccgtatt gttggtgctc ttggtggtag ggctaccgag 1740
caggttgtct tcggtgatgc tgaagttacg actggagcgt ccagtgattt gcagcaagtt 1800
acttctatgg ccaagcagat ggtcacagta ttcggtatgt cagacatcgg cccatgggct 1860
ttgatggacc cttcatccca gggaggagat atgattatgc gtatgatggc acgtaactcc 1920
atgtccgaga agttggctga ggacatcgac aaggctgtga aggctatctc tgacgaggcc 1980
tacgaagtcg cactgggtca cattaggaac aaccgcacgg ccatggacaa gattgtagag 2040
gttctgcttg agaaggagac ttgtccggc gccgagttta gggctattct ttcggaatac 2100
acagagattc ctgctgaaaa ccgtgtatca gacaaccagg ccgcacctgt agctgtttga 2160
agtgatcaga agcaagggtg ttgtgaaca caatcgtgta ggttttggag catcgacgct 2220
cttgtatcag agctcgc

2237

<210> 7
 <211> 442
 <212> PRT
 <213> Physcomitrella patens

<400> 7

Met	Tyr	Ser	Asn	Phe	Lys	Glu	Gln	Ala	Ile	Glu	Tyr	Val	Arg	Gln	Ala
1				5					10					15	
Val	Ala	Glu	Asp	Asn	Ala	Gly	Asn	Tyr	Ala	Lys	Ala	Phe	Pro	Leu	Tyr
			20				25						30		
Met	Asn	Ala	Leu	Glu	Tyr	Phe	Lys	Thr	His	Leu	Lys	Tyr	Glu	Lys	Asn
		35					40					45			
Pro	Lys	Ile	Lys	Glu	Ala	Ile	Thr	Gln	Lys	Phe	Thr	Glu	Tyr	Leu	Arg
	50					55					60				
Arg	Ala	Glu	Glu	Ile	Arg	Ala	Val	Leu	Asp	Asp	Gly	Pro	Thr	Gly	Pro
65					70					75					80
Ser	Ala	Asn	Gly	Asp	Ala	Ala	Val	Gln	Ala	Lys	Pro	Lys	Ser	Lys	Ser
				85					90					95	
Gly	Lys	Lys	Asp	Gly	Gly	Gly	Gly	Asp	Gly	Asp	Gly	Asp	Ser	Glu	Asp
			100					105					110		
Pro	Glu	Gln	Gln	Lys	Leu	Arg	Ser	Gly	Leu	Asn	Ser	Ala	Ile	Ile	Arg
		115					120					125			
Glu	Lys	Pro	Asn	Val	Arg	Trp	Ala	Asp	Val	Ala	Gly	Leu	Glu	Ser	Ala
	130					135					140				
Lys	Gln	Ala	Leu	Gln	Glu	Ala	Val	Ile	Leu	Pro	Val	Lys	Phe	Pro	Gln
145					150					155					160
Phe	Phe	Thr	Gly	Lys	Arg	Arg	Pro	Trp	Arg	Ala	Phe	Leu	Leu	Tyr	Gly
				165					170					175	
Pro	Pro	Gly	Thr	Gly	Lys	Ser	Tyr	Leu	Ala	Lys	Ala	Val	Ala	Thr	Glu
			180					185					190		
Ala	Asp	Ser	Thr	Phe	Phe	Ser	Ile	Ser	Ser	Ser	Asp	Leu	Val	Ser	Lys
		195					200					205			
Trp	Met	Gly	Glu	Ser	Glu	Lys	Leu	Val	Ala	Asn	Leu	Phe	Gln	Met	Ala
	210					215					220				
Arg	Glu	Ala	Ala	Pro	Ser	Ile	Ile	Phe	Ile	Asp	Glu	Ile	Asp	Ser	Leu
225					230					235					240
Cys	Gly	Thr	Arg	Gly	Glu	Gly	Asn	Glu	Ser	Glu	Ala	Ser	Arg	Arg	Ile
				245					250					255	
Lys	Thr	Glu	Leu	Leu	Val	Gln	Met	Gln	Gly	Val	Gly	Asn	Gln	Asp	Thr
			260					265					270		
Lys	Val	Leu	Val	Leu	Ala	Ala	Thr	Asn	Thr	Pro	Tyr	Ser	Leu	Asp	Gln
		275					280					285			

Ala Val Arg Arg Arg Phe Asp Lys Arg Ile Tyr Ile Pro Leu Pro Glu
 290 295 300
 Ser Lys Ala Arg Gln His Met Phe Lys Val His Leu Gly Asp Thr Pro
 305 310 315 320
 Asn Asn Leu Thr Glu Arg Asp Tyr Glu Asp Leu Ala Arg Lys Thr Asp
 325 330 335
 Gly Phe Ser Gly Ser Asp Ile Ala Val Cys Val Lys Asp Val Leu Phe
 340 345 350
 Glu Pro Val Arg Lys Thr Gln Asp Ala Met His Phe Lys Arg Ile Asn
 355 360 365
 Thr Lys Glu Gly Glu Met Trp Met Pro Cys Gly Pro Arg Glu Pro Gly
 370 375 380
 Ala Arg Gln Thr Thr Met Thr Glu Leu Ala Ala Glu Gly Gln Ala Ser
 385 390 395 400
 Lys Ile Leu Pro Pro Pro Ile Thr Lys Ser Asp Phe Asp Lys Val Leu
 405 410 415
 Ala Lys Gln Arg Pro Thr Val Ser Lys Gly Asp Leu Ile Ile Gln Glu
 420 425 430
 Lys Phe Thr Lys Glu Phe Gly Glu Glu Gly
 435 440

<210> 8
 <211> 901
 <212> PRT
 <213> Physcomitrella patens

<400> 8
 Met Glu Asn Asn Asp Ala Leu Asp Ile Gly Ala Val Ser Ser Pro Tyr
 1 5 10 15
 Pro Ser Gln Ser Glu Gly Val Ser Thr Pro Leu Pro Gln Val Thr Ser
 20 25 30
 Pro Ser Phe Asp Asn Ala Ala Ser Pro Val Ala Gly Arg Arg Ala Val
 35 40 45
 Arg Gln Thr Pro Thr Ser Ala Val Arg Arg Arg Gly Arg Glu Thr Asp
 50 55 60
 Ser Ala Arg Arg Arg Arg Ser Arg Ser Arg Ser Leu Gly Asn Ser Val
 65 70 75 80
 Tyr Ser Ser Pro Tyr Asp Ala Gly Thr Pro Gly Thr Pro Gly Thr Pro
 85 90 95
 Val Ala Thr Pro Val Tyr Ala Thr Pro Val Gly Thr Pro Met Gly Thr
 100 105 110
 Pro Ser Phe His Arg Gly Thr Pro Gln Tyr Lys Gln Arg Ser Glu Leu
 115 120 125

Gly	Ser	Gln	Gly	Lys	Pro	Leu	His	Arg	Arg	Arg	Arg	Ser	Gln	Ser	Arg	130	135	140
Glu	Pro	Gly	His	Arg	Ser	Pro	Ser	Arg	Glu	Pro	Ser	Ala	Asp	Gly	Arg	145	150	155
Pro	Ser	Glu	Ser	Ala	Glu	Pro	Asp	Asp	Thr	Leu	Gly	Gly	Glu	Tyr	Ala	165	170	175
Tyr	Val	Trp	Gly	Thr	Asn	Val	Asn	Ile	Pro	Asp	Val	Leu	Arg	Ala	Ile	180	185	190
Arg	Arg	Phe	Leu	His	Asn	Tyr	Arg	Ser	Ser	Ala	His	Asp	Leu	Asn	Ser	195	200	205
Lys	Tyr	Ile	Gln	Ile	Ile	Glu	Glu	Thr	Val	Glu	Arg	Glu	Glu	Asp	Thr	210	215	220
Leu	Asn	Ile	Asp	Met	Ser	Asp	Ile	Tyr	Asp	His	Asp	Pro	Asp	Leu	Tyr	225	230	235
Ala	Lys	Ile	Val	Arg	Tyr	Pro	Leu	Asp	Ile	Ile	Pro	Leu	Leu	Asp	Thr	245	250	255
Glu	Cys	Gln	Glu	Val	Ala	Thr	Ser	Leu	Leu	Pro	Thr	Phe	Glu	Lys	His	260	265	270
Ile	Glu	Ala	Arg	Pro	Phe	Asn	Leu	Lys	Ala	Ser	Val	His	Met	Arg	Glu	275	280	285
Leu	Asn	Pro	Ser	Asp	Ile	Asp	Lys	Leu	Val	Ser	Val	Lys	Gly	Met	Val	290	295	300
Ile	Arg	Cys	Ser	Ser	Ile	Ile	Pro	Glu	Ile	Lys	Gly	Ala	Phe	Phe	Lys	305	310	315
Cys	Leu	Val	Cys	Gly	His	Ser	Pro	Pro	Leu	Val	Thr	Val	Val	Lys	Gly	325	330	335
Arg	Val	Glu	Glu	Pro	Thr	Arg	Cys	Glu	Lys	Pro	Glu	Cys	Ala	Ala	Arg	340	345	350
Asn	Ala	Met	Ser	Leu	Ile	His	Asn	Arg	Cys	Thr	Phe	Ala	Asn	Lys	Gln	355	360	365
Ile	Val	Arg	Leu	Gln	Glu	Thr	Pro	Asp	Ala	Ile	Pro	Glu	Gly	Glu	Thr	370	375	380
Pro	His	Thr	Val	Ser	Met	Cys	Leu	Tyr	Asn	Thr	Met	Val	Asp	Ala	Val	385	390	395
Lys	Pro	Gly	Asp	Arg	Ile	Glu	Val	Thr	Gly	Val	Phe	Lys	Ala	Met	Ala	405	410	415
Val	Arg	Val	Gly	Pro	Asn	Gln	Arg	Thr	Leu	Arg	Ala	Leu	Tyr	Lys	Thr	420	425	430
Tyr	Ile	Asp	Cys	Val	His	Val	Lys	Lys	Ser	Asp	Arg	Gly	Arg	Leu	Gln	435	440	445

Thr	Glu	Asp	Pro	Met	Glu	Met	Asp	Lys	Glu	Asn	Asp	Met	Tyr	Ala	Gly
450						455					460				
Tyr	His	Glu	Ser	Asp	Thr	Ser	Glu	Ala	Ala	Asn	Glu	Ala	Lys	Ile	Gln
465					470					475					480
Lys	Leu	Lys	Glu	Leu	Ser	Lys	Leu	Pro	Gly	Ile	Tyr	Asp	Arg	Leu	Ser
				485					490					495	
Arg	Ser	Leu	Ala	Pro	Ser	Ile	Trp	Glu	Leu	Glu	Asp	Ile	Lys	Lys	Gly
			500					505					510		
Leu	Leu	Cys	Gln	Leu	Phe	Gly	Gly	Lys	Ala	Lys	Lys	Ile	Pro	Ser	Gly
		515					520					525			
Ala	Ser	Phe	Arg	Gly	Asp	Ile	Asn	Val	Leu	Leu	Val	Gly	Asp	Pro	Gly
	530					535					540				
Thr	Ser	Lys	Ser	Gln	Leu	Leu	Gln	Tyr	Val	His	Lys	Ile	Ala	Pro	Arg
545					550					555					560
Gly	Ile	Tyr	Thr	Ser	Gly	Arg	Gly	Ser	Ser	Ala	Val	Gly	Leu	Thr	Ala
				565					570					575	
Tyr	Val	Thr	Lys	Asp	Pro	Glu	Thr	Arg	Glu	Thr	Val	Leu	Glu	Ser	Gly
			580					585					590		
Ala	Leu	Val	Leu	Ser	Asp	Arg	Gly	Ile	Cys	Cys	Ile	Asp	Glu	Phe	Asp
		595					600					605			
Lys	Met	Ser	Asp	Asn	Ala	Arg	Ser	Met	Leu	His	Glu	Val	Met	Glu	Gln
	610					615					620				
Gln	Thr	Val	Ser	Val	Ala	Lys	Gly	Gly	Ile	Ile	Ala	Ser	Leu	Asn	Ala
625					630					635					640
Arg	Thr	Ser	Val	Leu	Ala	Cys	Ala	Asn	Pro	Ser	Gly	Ser	Arg	Tyr	Asn
				645					650					655	
Ala	Arg	Leu	Ser	Val	Ile	Asp	Asn	Ile	Gln	Leu	Pro	Pro	Thr	Leu	Leu
			660					665					670		
Ser	Arg	Phe	Asp	Leu	Ile	Tyr	Leu	Met	Leu	Asp	Lys	Pro	Asp	Glu	Gln
		675					680					685			
Asn	Asp	Arg	Arg	Leu	Ala	Arg	His	Leu	Val	Ala	Leu	His	Tyr	Glu	Asn
	690					695					700				
Tyr	Glu	Val	Ser	Lys	Gln	Asp	Ala	Leu	Asp	Leu	Gln	Thr	Leu	Thr	Ala
705					710					715					720
Tyr	Ile	Thr	Tyr	Ala	Arg	Gln	His	Val	His	Pro	Thr	Leu	Ser	Asp	Glu
				725					730					735	
Ala	Ala	Glu	Asp	Leu	Ile	Asn	Gly	Tyr	Val	Glu	Met	Arg	Gln	Lys	Gly
			740					745					750		
Asn	Phe	Pro	Gly	Ser	Ser	Lys	Lys	Val	Ile	Thr	Ala	Thr	Pro	Arg	Gln
		755					760					765			

Leu Glu Ser Met Ile Arg Ile Ser Glu Ala Leu Ala Arg Met Arg Phe
 770 775 780
 Ser Glu Val Val Glu Lys Val Asp Ala Ala Glu Ala Val Arg Leu Leu
 785 790 795 800
 Asp Val Ala Leu Gln Gln Ser Ala Thr Asp His Ala Thr Gly Thr Ile
 805 810 815
 Asp Met Asp Leu Ile Thr Thr Gly Val Ser Ala Ser Glu Arg Ile Arg
 820 825 830
 Arg Ala Asn Leu Leu Ala Ala Leu Arg Glu Leu Ile Ala Asp Lys Ile
 835 840 845
 Ser Pro Gly Ser Ser Ser Gly Leu Lys Thr Ser Gln Leu Leu Glu Asp
 850 855 860
 Ile Arg Ser Gln Ser Ser Val Asp Val Ser Leu Gln Asp Ile Lys Asn
 865 870 875 880
 Ala Leu Gly Ser Leu Gln Gly Glu Gly Phe Leu Thr Val His Gly Asp
 885 890 895
 Ile Val Lys Arg Val
 900

<210> 9
 <211> 698
 <212> PRT
 <213> *Physcomitrella patens*

<400> 9
 Met Ala Ser Ala Thr Ala Ala Thr Met Ala Ser Leu Leu Thr Pro Gly
 1 5 10 15
 Ser Leu Arg Arg Gly Leu Gly Ser Gln Glu Ser Ser Thr Gln Phe Ala
 20 25 30
 Pro Leu Ala Gly Pro Arg Lys Thr Ser Val Ser Arg Arg Val Thr Ala
 35 40 45
 Ser Ala Ser Gly Lys Asn Asp Asn Gly Val Val Glu Asp Val Asp Met
 50 55 60
 Gly Lys Arg Gly Met Leu Lys Gly Val Ala Gly Ala Leu Ala Ala Val
 65 70 75 80
 Leu Pro Ala Val Ile Ala Lys Lys Ala Ser Ala Ala Glu Glu Gln Gly
 85 90 95
 Val Ala Ser Ser Arg Met Ser Tyr Ser Arg Phe Leu Glu Tyr Leu Asp
 100 105 110
 Met Asp Arg Val Lys Lys Val Asp Leu Tyr Glu Asn Gly Thr Ile Ala
 115 120 125
 Ile Val Glu Ala Val Ser Pro Glu Leu Gly Asn Arg Val Gln Arg Val
 130 135 140

Arg 145	Val	Gln	Leu	Pro	Gly 150	Thr	Ser	Ser	Glu	Leu 155	Leu	Ser	Lys	Phe	Arg 160
Ser	Lys	Asn	Val	Asp 165	Phe	Ala	Ala	His	Ser 170	Pro	Gln	Glu	Asp	Ser	Gly 175
Ser	Val	Ile	Leu 180	Asn	Leu	Ile	Gly	Asn 185	Leu	Ala	Phe	Pro	Leu 190	Leu	Leu
Val	Gly	Gly 195	Leu	Phe	Phe	Leu	Ser 200	Arg	Arg	Ser	Gln	Gly 205	Gly	Met	Gly
Pro	Gly 210	Gly	Pro	Gly	Asn	Pro 215	Met	Ala	Phe	Gly	Lys 220	Ser	Lys	Ala	Lys
Phe 225	Gln	Met	Glu	Pro	Asn 230	Thr	Gly	Ile	Thr	Phe 235	Gln	Asp	Val	Ala	Gly 240
Val	Asp	Glu	Ala	Lys 245	Gln	Asp	Phe	Met	Glu 250	Val	Val	Glu	Phe	Leu 255	Lys
Arg	Pro	Glu	Arg 260	Phe	Thr	Ala	Val	Gly 265	Ala	Lys	Ile	Pro	Lys 270	Gly	Val
Leu	Leu	Val 275	Gly	Pro	Pro	Gly	Thr 280	Gly	Lys	Thr	Leu	Leu 285	Ala	Lys	Ala
Ile 290	Ala	Gly	Glu	Ala	Gly	Val 295	Pro	Phe	Phe	Ser	Ile 300	Ser	Gly	Ser	Glu
Phe 305	Val	Glu	Met	Phe	Val 310	Gly	Val	Gly	Ala	Ser 315	Arg	Val	Arg	Asp	Leu 320
Phe	Lys	Lys	Ala	Lys 325	Glu	Asn	Ala	Pro	Cys 330	Ile	Val	Phe	Val	Asp 335	Glu
Ile	Asp	Ala	Val 340	Gly	Arg	Gln	Arg	Gly 345	Thr	Gly	Ile	Gly	Gly 350	Gly	Asn
Asp	Glu	Arg 355	Glu	Gln	Thr	Leu	Asn 360	Gln	Leu	Leu	Thr	Glu 365	Met	Asp	Gly
Phe 370	Glu	Gly	Asn	Thr	Gly	Val 375	Ile	Val	Ile	Ala	Ala 380	Thr	Asn	Arg	Ala
Asp 385	Ile	Leu	Asp	Ala	Ala 390	Leu	Leu	Arg	Pro	Gly 395	Arg	Phe	Asp	Arg	Gln 400
Val	Ser	Val	Asp 405	Val	Pro	Asp	Val	Lys	Gly 410	Arg	Thr	Asp	Ile	Leu 415	Lys
Val	His	Ala	Ser 420	Asn	Lys	Lys	Phe	Ala 425	Asp	Asp	Val	Ser	Leu 430	Asp	Ile
Ile	Ala	Met 435	Arg	Thr	Pro	Gly	Phe 440	Ser	Gly	Ala	Asp	Leu 445	Ala	Asn	Leu
Leu 450	Asn	Glu	Ala	Ala	Ile	Leu 455	Thr	Gly	Arg	Arg	Gly 460	Lys	Thr	Ala	Ile

Ser Ala Lys Glu Ile Asp Asp Ser Ile Asp Arg Ile Val Ala Gly Met
 465 470 475 480
 Glu Gly Thr Val Met Thr Asp Gly Lys Ser Lys Ser Leu Val Ala Tyr
 485 490 495
 His Glu Val Gly His Ala Ile Cys Gly Thr Leu Thr Pro Gly His Asp
 500 505 510
 Ala Val Gln Lys Val Thr Leu Ile Pro Arg Gly Gln Ala Arg Gly Leu
 515 520 525
 Thr Trp Phe Ile Pro Gly Glu Asp Pro Thr Leu Ile Ser Lys Gln Gln
 530 535 540
 Ile Phe Ala Arg Ile Val Gly Ala Leu Gly Gly Arg Ala Thr Glu Gln
 545 550 555 560
 Val Val Phe Gly Asp Ala Glu Val Thr Thr Gly Ala Ser Ser Asp Leu
 565 570 575
 Gln Gln Val Thr Ser Met Ala Lys Gln Met Val Thr Val Phe Gly Met
 580 585 590
 Ser Asp Ile Gly Pro Trp Ala Leu Met Asp Pro Ser Ser Gln Gly Gly
 595 600 605
 Asp Met Ile Met Arg Met Met Ala Arg Asn Ser Met Ser Glu Lys Leu
 610 615 620
 Ala Glu Asp Ile Asp Lys Ala Val Lys Ala Ile Ser Asp Glu Ala Tyr
 625 630 635 640
 Glu Val Ala Leu Gly His Ile Arg Asn Asn Arg Thr Ala Met Asp Lys
 645 650 655
 Ile Val Glu Val Leu Leu Glu Lys Glu Thr Leu Ser Gly Ala Glu Phe
 660 665 670
 Arg Ala Ile Leu Ser Glu Tyr Thr Glu Ile Pro Ala Glu Asn Arg Val
 675 680 685
 Ser Asp Asn Gln Ala Ala Pro Val Ala Val
 690 695

<210> 10

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 10

caggaaacag ctatgacc

18

<210> 11
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 11
 ctaaaggga caaaagctg 19

<210> 12
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 12
 tgtaaacga cggccagt 18

<210> 13
 <211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 13
 gcgatatcga cccaaggtgt gtagagaagg ggat 34

<210> 14
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 14
 gcgagctcgc gactgatgca actcaccat gaa 33

<210> 15
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 15
 cccaaatgct tggagccagc gacct 25

<210> 16
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 16
 gacgtgcacg caatcgatgt aggtc 25

<210> 17
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 17
 agtcgacccc tgtcagactt cttga 25

<210> 18
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 18
 atggcgcgcc gcactcacgt gaggaattgc acctcc 36

<210> 19
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 19
 gcgagctctg ggaattgcgc atgggaaact gg 32

<210> 20
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 20

ctggctaccc aaaccgcgtc ggaga

25

<210> 21

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 21

atccccgggtg atacgttgtg catatttggt gttgc

35

<210> 22

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 22

gcgagctctg atacaagagc gtcgatgctc ca

32

<210> 23

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 23

gcgctgcaga tttcatttgg agaggacacg

30

<210> 24

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 24

cgcggccggc ctcagaagaa ctcgtcaaga aggcg

35

<210> 25
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 25
 gctgacacgc caagcctcgc tagtc 25

<210> 26
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 26
 gcgagctcgc gactgatgca actcacccat gaa 33

<210> 27
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 27
 gcgagctctg ggaattgcgc atgggaaact gg 32

<210> 28
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 28
 gcgagctctg atacaagagc gtcgatgctc ca 32

<210> 29
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 29
 gatggtgatg gtgacagcga ggatc 25

<210> 30
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 30
cgacgtgaag cctcgctctc atttcc 26

<210> 31
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 31
cgatctcctt caagggaacc tagtgctg 28

<210> 32
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 32
gagttcacgc atgtgcaccg atgct 25

<210> 33
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 33
cgtagcgtct tccaggatgt cctac 25

<210> 34
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 34
gcttgacctc gtctactccc gcaac 25

WHAT IS CLAIMED IS:

1. A transgenic plant cell transformed by a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid in the plant cell results in increased tolerance to a drought or freezing stress as compared to a wild type variety of the plant cell, wherein
 - a. the CCSR is a cell cycle protein-1 (CC-1) protein as defined in SEQ ID NO:7; or
 - b. a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length, or
 - 10 c. the CCSR coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.
2. The transgenic plant cell of claim 1, wherein the CCSR is a CC-1 as defined in SEQ ID 7.
3. The transgenic plant cell of claim 1, wherein the CCSR coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.
- 20 4. The transgenic plant cell of claim 1, wherein the tolerance to drought stress is increased.
5. The transgenic plant cell of any one of claims 1 to 4, wherein the plant is a monocot.
6. The transgenic plant cell of any one of claims 1 to 4, wherein the plant is a dicot.

7. The transgenic plant cell of any one of claims 1 to 4, wherein the plant is maize, wheat, rye, oat, triticale, rice, barley, soybean, peanut, cotton, rapeseed, canola, manihot, pepper, sunflower, tagetes, solanaceous plants, potato, tobacco, eggplant, tomato, Vicia species, pea, alfalfa, coffee, cacao, tea, Salix species, oil palm, coconut, or perennial grass.

8. An isolated Cell Cycle Stress-Related Protein (CCSRP) wherein the CCSR is:

- a. a Cell Cycle-1 protein (CC-1) as defined in SEQ ID NO:7; or
- 10 b. a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length; or
- c. a polypeptide encoded by a nucleic acid that has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4,

and wherein the CCSR upon expression in a plant results in the plant's increased tolerance to drought or freezing stress as compared to a wild type variety of the plant.

9. Use of the CCSR as defined in claim 8, for increasing the tolerance of a
20 transformed plant or plant cell or plant seed to drought or freezing stress as compared to a wild type variety of the plant or plant cell or plant seed, wherein the plant or plant cell or plant seed comprises the CCSR.

10. The isolated Cell Cycle Stress-Related Protein of claim 8, wherein the CCSR is a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length.

11. An isolated Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein the CCSR coding nucleic acid, is a nucleic acid:

- a. having a sequence as defined in SEQ ID NO:4; or
- b. having a sequence of at least 90% identity to SEQ ID NO:4 over the entire coding region of SEQ ID NO:4; or
- c. encoding the CCSRP as defined in claim 8;

and wherein the expression in a plant increases the plant's tolerance to drought or freezing stress as compared to a wild type variety of the plant.

12. The isolated CCSRP coding nucleic acid of claim 11, wherein the CCSRP coding nucleic acid is as defined in SEQ ID NO:4 or a sequence of at least 90% identity to SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

13. The isolated CCSRP coding nucleic acid of claim 11, wherein the CCSRP coding nucleic acid encodes a polypeptide as defined in SEQ ID NO:7.

14. Use of an isolated CCSRP coding nucleic acid of any one of claims 11 to 13 for increasing the tolerance of a plant or plant cell or plant seed to drought or freezing stress, wherein the plant or plant cell or plant seed comprises the CCSRP.

15. Use of the isolated CCSRP coding nucleic acid of any one of claims 11 to 13 for identification or cloning of CC-1 homologs in other cell types and organisms.

16. An isolated recombinant expression vector comprising the nucleic acids of any one of claims 11 to 13, wherein the expression of the vector in a host cell results in increased tolerance to drought or freezing stress as compared to a wild type variety of the host cell.

17. A method of producing a transgenic plant containing a Cell Cycle Stress-Related Protein (CCSRP) coding nucleic acid, wherein expression of the nucleic acid in the plant results in increased tolerance to drought or freezing stress as

compared to a wild type variety of the plant, comprising transforming a plant cell with an expression vector comprising the nucleic acid, generating from the plant cell a transgenic plant with an increased tolerance to the drought and freezing stress, as compared to a wild type variety of the plant, wherein

- a. the CCSRP is a Cell Cycle-1 protein (CC-1) as defined in SEQ ID NO:7; or
- b. the CCSRP is a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length; or
- c. the CCSRP coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

10

18. The method of claim 17, wherein the CCSRP has at least 90% sequence identity with SEQ ID NO:7 over its entire length.

19. The method of claim 17, wherein the CCSRP coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

20. A method of modifying drought or freezing stress tolerance of a transgenic plant comprising, modifying the expression of a Cell Cycle Stress-Related Protein (CCSRP) in the transgenic plant as compared to a wild type variety of the plant, wherein

- a. the CCSRP is a cell cycle protein-1 (CC-1) protein as defined in SEQ ID NO:7; or
- b. a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length, or
- c. the CCSRP coding nucleic acid has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

21. The method of claim 20, wherein the CCSRP is a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length.
22. The method of claim 20, wherein the CCSRP is a protein as defined in SEQ ID NO:7.
23. The method of claim 20, wherein the CCSRP is encoded by a nucleic acid that has at least 90% sequence identity with SEQ ID NO:4 over the entire coding
10 region of SEQ ID NO:4.
24. The method of any one claims 20 to 23, wherein the drought or freezing stress tolerance is increased as compared to a wild type variety of the plant.
25. The method of claim 20, wherein the plant is transformed with a promoter that directs expression of the CCSRP.
26. The method of claim 25, wherein the promoter is tissue specific.
- 20 27. The method of claim 25, wherein the promoter is developmentally regulated.
28. The method of claim 20, wherein CCSRP expression is modified by administration of an antisense molecule that inhibits expression of CCSRP.
29. Use of claim 14, wherein the CCSRP is a polypeptide having at least 90% sequence identity with SEQ ID NO:7 over its entire length.
30. Use of claim 14, wherein the CCSRP is a protein as defined in SEQ ID NO:7.

31. Use of claim 14, wherein the CCSRP is encoded by a nucleic acid that has at least 90% sequence identity with SEQ ID NO:4 over the entire coding region of SEQ ID NO:4.

FIGURE 1A

Nucleotide sequence of the partial CC-1 from *Physcomitrella patens* (SEQ ID NO:1)

CGGGAGTTGGTGATCTTCAGCATGCTCGTAGTTGACAAGGGAGGTAGGATTGTTTGA
CCCAAGGTGTGTANAGAAGGGGATAGCCATGTACAGCAACTTCAAAGAGCAGGCCA
TAGAATATTGTGCGTCAAGCCGTAGCGGAAGACAACGCAGGGA ACTATGCCAAAGC
GTTTCCGCTGTACATGAACGCGCTTGAGTACTTCAAGACGCATCTAAAGTATGAGAA
GAATCCCAAATCAAGGAGGCCATCACTCAGAAGTTCACGGAGTATTTGAGGAGGG
CGGAGGAGATTCGAGCCGTTTTGGACGATGGCCCCACTGGACCCTCTGCAAATGGA
GACGCGGCAGTTCAAGCTAAACCGAAGTCGAAATCAGGGAAGAAGGATGGTGGCG
GGGGTGATGGTGATGGTGACAGCGAGGATCCCGACCAGCAGAAGCTGAGATCAGGG
CTGAACTCGGCAATCATACGGGAAAAGCCAAATGTTTCGGTGGGCTGATGTTGCTGG
ACTTGAAAGTGCCAAGCAGGCGTTGCAGGAGGCAGTGATCTTGCCCGTGAAGTTTCC
CCAATTTTTCACAGGGAAGCGAAGAACATGGCGAGCATTTTTGTGGTATGGGGCCCCC
CGGGACTGGAAAATCGTATCTTGCAAAAGCTGTTGCTACGGAAGCTGATTCTACATT
CTTTAGTATTTCTCTTCAGACTTGGTGTCAAAGTGGATGGGAGAGAGTGAGAAGCT
TGTTGCAAATCTGTTTCAAATGGCCCGTGAAGCTGCTCCATCCATCATCTTCATAGAC
GAGATTGATTCTTTATGCGGTACTCGAGGTGAAGGAAATGAGAGCGAGGCTTCACG
TCGTATCAAGACTGAGTTGCTAGTTCAAATGCAGGGTGTC

FIGURE 1B

Nucleotide sequence of the partial CC-2 from *Physcomitrella patens* (SEQ ID NO:2)

CGGCACCAGGGAGATCGTATTGAGGTAACAGGAGTTTTCAAGGCCATGGCAGTTTCG
AGTTGGTCCGAATCAACGAACATTACGAGCATTGTATAAGACCTACATCGATTGCGT
GCACGTCAAGAAGTCTGACAGGGGGTCGACTGCAAACCTGAAGATCCTATGGAGATGG
ATAAGGAGAATGATATGTATGCTGGGTATCATGAAAGTGATACTTCAGAAGCTGCT
AATGAAGCAAAGATTCAAAAACCTTAAAGAGCTGTCCAAGCTCCCGGACATTTATGA
TAGACTTTCAAGGTCGCTGGCTCCAAGCATTTGGGAGCTTGAAGATATTA AAAAAGGG
TCTTCTTTGCCAGCTCTTTGGTGGGAAGGCTAAGAAAATTCCATCTGGAGCATCTTTC
CGAGGTGACATCAATGTTTTACTTGTTGGGGACCCTGGTACCAGTAAATCTCAGCTG
CTTCAGTATGTGCACAAGATAGCTCCTCGTGGAATCTACACTAGTGGGCGAGGAAGT
TCGGCGGTTGGGCTGACAGCGTATGTAAACGAAGGATCCAGAAACTCGAGAGACGG
TATTGGAGAGCGGAGCTTTGGTTCTTAGTGATCGTGGGATATGCTGTATCGATGAGT
TCGACAAAATGTCTGATAATGCCCCGAAGCATGCTTCATGAGGTAATGGAGCAACAA
ACGGTATCTGGACCCAAGCGGTTCATGCTCGTGAAGCCGAGTTG

FIGURE 1C

Nucleotide sequence of the partial CC-3 from *Physcomitrella patens* (SEQ ID NO:3)

GCACCAGCCGCTTTGGAATCCCATCCCTCGGTTGCATAGACACAAGGGGATTCAGTG
TAGTGATACGTTGTGCATATTTGGTGTGCAAGATTTTTGGTTTCTTGATTGTTAGCT
ATGGCGTCTGCAACAGCGGCTACAATGGCGTCCCTCCTCACGCCTGGGTCTCTCCGA
CGCGGTTTGGGTAGCCAGGAATCGTCGACCCAATTTGCTCCCCTAGCTGGTCCTCGT
AAGACATCAGTTTCGCGTAGGGTGACTGCTAGCGCTAGTGGGAAGAACGACAATGG
AGTCGTGGAAGATGTGGATATGGGGAAGCGGGGTATGTTGAAAGGCGTAGCGGGAG
CTTTGGCTGCAGTTCTCCCTGCTGTTATCGCGAAGAAAGCTTCAGCAGCTGAGGAGC
AGGGCGTAGCGTCTTCCAGGATGTCCTACTCGAGGTTTTTGGAGTATTTGGATATGG
ACCGTGTGAAGAAGTTGACTTGTATGAAAATGGGACCATAGCAATTGTGGAGGCTG
TATCCCCTGAATTGGGCAACAGAGTGCAACGCGTACGCGTGCAGCTCCCCGGAAC

FIGURE 2A

Nucleotide sequence of the full-length CC-1 from *Physcomitrella patens* (SEQ ID NO:4)

GCGATATCGACCCAAGGTGTGTAGAGAAGGGGATAGCCATGTACAGCAACTTCAAA
GAGCAGGCCATAGAATATGTGCGTCAAGCCGTAGCGGAAGACAACGCAGGGAACTA
TGCCAAAGCGTTTCCGCTGTACATGAACGCGCTTGAGTACTTCAAGACGCATCTAAA
GTATGAGAAGAATCCCAAAATCAAGGAGGCCATCACTCAGAAGTTCACGGAGTATT
TGAGGAGGGCGGAGGAGATTCTGAGCCGTTTTGGACGATGGCCCCACTGGACCCTCT
GCAAATGGAGACGCGGCAGTTCAAGCTAAACCGAAGTCGAAATCAGGGAAGAAGG
ATGGTGGCGGGGGTGATGGTGATGGTGACAGCGAGGATCCCGAGCAGCAGAAGCTG
AGATCAGGGCTGAACTCGGCAATCATACGGGAAAAGCCAAATGTTTCGGTGGGCTGA
TGTTGCTGGACTTGAAAGTGCCAAGCAGGCGTTGCAGGAGGCAGTGATCTTGCCCGT
GAAGTTTCCCAATTTTTTCACAGGGAAGCGAAGACCATGGCGAGCATTTTTGTTGTA
TGGGCCCCCGGGACTGGAAAATCGTATCTTGCAAAAGCTGTTGCTACGGAAGCTG
ATTCTACATTCTTTAGTATTTCTCTTCAGACTTGGTGTCAAAGTGGATGGGAGAGA
GTGAGAAGCTTGTTGCAAATCTGTTTCAAATGGCCCGTGAAGCTGCTCCATCCATCA
TCTTCATAGACGAGATTGATTCTTTATGCGGTACTCGAGGTGAAGGAAATGAGAGCG
AGGCTTCACGTCGTATCAAGACTGAGTTGCTAGTTCAAATGCAGGGTGTCGGCAATC
AAGACACTAAGGTTCTTGTGTTAGCTGCTACAAATACGCCCTACTCTTTGGATCAGG
CGGTGAGGCGACGTTTCGACAAGCGTATCTACATCCCCTACCGGAGTCTAAGGCTC
GGCAGCACATGTTTAAGGTGCATTTGGGAGATACGCCAAACAACCTGACTGAACGT
GATTATGAGGATCTGGCTAGGAAGACTGATGGGTTTTTCAGGCTCGGATATTGCAGTT
TGTGTGAAAGATGTACTATTTGAGCCTGTTTCGTAAGACCCAAGATGCTATGCATTTC
AAAAGAATTAATACCAAAGAAGGAGAGATGTGGATGCCTTGTGGGCCCCGAGAACC

FIGURE 2A (Continued)

AGGTGCTAGGCAAACCACTATGACGGAGCTTGCCGCTGAAGGGCAGGCATCGAAGA
TTTTACCACCTCCAATCACAAAATCAGATTTCGACAAAGTCCTCGCAAAGCAGAGGC
CCTGTGTCAGCAAAGGCGACCTTATTATTCAAGAGAAATTCACCAAAGAATTTGGTG
AAGAAGGTTGAATGGTGCTTCGAGTTAAGAATTTGGAAGGTTCTGGGTACGGAAG
ACAGACGAAATAGAACGTCGTAGGTACGGTAGCCTAAGAGTAAATTACGGAAGTTT
TCCGACTTGCCAGTTGTGCACTCTTTCAACATGAAGGGAAGGAAGCTACTTGTCGGT
TGCCTTTTCATGGGTGAGTTGCATCAGTCGCGAGCTCGC

FIGURE 2B

Nucleotide sequence of the full-length CC-2 from *Physcomitrella patens* (SEQ ID NO:5)

ATGGCGCGCCGCACTCACGTGAGGAATTGCACCTCCTTGTTCTGCGACGGTTCCATT
CTTTTTGGTTTTTAGTTTGCAAATCTTGATCGTGGAGTTGAGAAAAAGGGCGGTTCTGT
TGTCTTGAGGTGTTCTTGTTGATTGTTTCGTCATGGAAAATAATGATGCACTTGACATT
GGAGCCGTGTCGTCCCCATATCCTTCGCAATCTGAAGGAGTGTCTACGCCATTGCCG
CAAGTAACATCACCGAGCTTCGACAATGCAGCCTCACCCGTGGCCGGGCGGAGGGC
CGTACGGCAGACCCCTACATCTGCAGTTCGAAGGAGAGGGAGAGAAACGGATTCCG
CTCGTCGTAGGAGGAGTCGATCTCGCAGTTTAGGCAATTCTGTTTATAGTTCCCCTTA
CGATGCGGGGACTCCTGGAACCTCCTGGAACCTCCAGTGGCTACTCCGGTTTACGCTAC
CCCAGTCGGTACACCTATGGGTACCCCATCGTTCCATCGTGGCACGCCACAGTACAA
ACAGCGCAGTGAGCTTGGTTCCCAGGGGAAGCCTCTACATCGGAGACGTCGATCTC
AATCCAGAGAACCCGGGCATCGATCTCCTTCAAGGGAACCTAGTGCTGATGGGCGT
CCCTCTGAATCTGCTGAGCCAGATGACACTTTGGGTGGAGAATATGCTTATGTTTGG
GGGACGAATGTTAACATTCCAGATGTGCTTAGGGCGATTTCGTCGATTTCTCCACAAT
TATCGTTCGAGTGCTCATGATCTTAATTCCAAGTACATCCAGATCATAGAGGAGACT
GTGGAGCGTGAGGAGGATACTCTAAATATCGACATGTCAGACATTTATGACCATGAT
CCTGATCTATACGCAAAAATTGTTTCGATACCCACTCGACATCATCCCCCTGTTGGAC
ACTGAGTGTCAGGAAGTTGCTACCTCTTTACTACCAACGTTTGAGAAGCATATTGAG
GCCAGACCTTTCAATCTCAAAGCATCGGTGCACATGCGTGAACCTCAACCCTTCAGAT
ATAGACAAATTGGTTTCTGTTAAAGGAATGGTTATCCGGTGCAGTTCTATCATACCT
GAAATTAAGGGGGCCTTCTTCAAATGTTTAGTGTGTGGTCACTCGCCTCCGCTAGTT
ACAGTTGTTAAAGGGCGGGTTGAGGAGCCAACAAGGTGTGAAAAGCCAGAATGTGC

FIGURE 2B (Continued)

AGCACGGAATGCTATGTCTCTTATTCACAATCGATGCACTTTTGCAAATAAGCAGAT
AGTGCGTCTTCAAGAACTCCAGATGCCATTCTGAAGGAGAGACTCCACACACAG
TCAGCATGTGTTTATACAACACTATGGTTGATGCTGTGAAGCCTGGAGATCGTATTG
AGGTAACAGGAGTTTTCAAGGCCATGGCAGTTCGAGTTGGTCATGGCGCGCCGCACT
CACGTGAGGAATTGCACCTCCTTGTTCTGCGACGGTTCATTCTTTTTGGTTTTTAGT
TTGCAAATCTTGATCGTGGAGTTGAGAAAAAGGGCGGTTTCGTTGTCTTGAGGTGTTC
TTGTTGATTGTTTCGTCATGGAAAATAATGATGCACTTGACATTGGAGCCGTGTCGTC
CCCATATCCTTCGCAATCTGAAGGAGTGTCTACGCCATTGCCGCAAGTAACATCACC
GAGCTTCGACAATGCAGCCTCACCCGTGGCCGGGCGGAGGGCCGTACGGCAGACCC
CTACATCTGCAGTTCGAAGGAGAGGGAGAGAAACGGATTCCGCTCGTCGTAGGAGG
AGTCGATCTCGCAGTTTAGGCAATTCTGTTTATAGTTCCCCTTACGATGCGGGGACTC
CTGGAACCTCCTGGAACCTCCAGTGGCTACTCCGGTTTACGCTACCCCAGTCGGTACAC
CTATGGGTACCCCATCGTTCCATCGTGGCACGCCACAGTACAAACAGCGCAGTGAG
CTTGGTTCCCAGGGGAAGCCTCTACATCGGAGACGTCGATCTCAATCCAGAGAACCC
GGGCATCGATCTCCTTCAAGGGAACCTAGTGCTGATGGGCGTCCCTCTGAATCTGCT
GAGCCAGATGACACTTTGGGTGGAGAATATGCTTATGTTTGGGGGACGAATGTTAAC
ATTCCAGATGTGCTTAGGGCGATTTCGTCGATTTCTCCACAATTATCGTTTCGAGTGCTC
ATGATCTTAATTCCAAGTACATCCAGATCATAGAGGAGACTGTGGAGCGTGAGGAG
GATACTCTAAATATCGACATGTCAGACATTTATGACCATGATCCTGATCTATACGCA
AAAATTGTTCGATACCCACTCGACATCATCCCCCTGTTGGACACTGAGTGTCAGGAA
GTTGCTACCTCTTTACTACCAACGTTTGAGAAGCATATTGAGGCCAGACCTTTCAAT
CTCAAAGCATCGGTGCACATGCGTGAACCTCAACCCTTCAGATATAGACAAATTGGTT

FIGURE 2B (Continued)

TCTGTTAAAGGAATGGTTATCCGGTGCAGTTCTATCATACCTGAAATTAAGGGGGGCC
TTCTTCAAATGTTTAGTGTGTGGTCACTCGCCTCCGCTAGTTACAGTTGTTAAAGGGC
GGGTTGAGGAGCCAACAAGGTGTGAAAAGCCAGAATGTGCAGCACGGAATGCTATG
TCTCTTATTCACAATCGATGCACTTTTGCAAATAAGCAGATAGTGCGTCTTCAAGAA
ACTCCAGATGCCATTCCTGAAGGAGAGACTCCACACACAGTCAGCATGTGTTTATAC
AACACTATGGTTGATGCTGTGAAGCCTGGAGATCGTATTGAGGTAACAGGAGTTTTTC
AAGGCCATGGCAGTTCGAGTTGGTCCGAATCAACGAACATTACGAGCATTGTATAA
GACCTACATCGATTGCGTGCACGTCAAGAAGTCTGACAGGGGGTCGACTGCAAACCTG
AAGATCCTATGGAGATGGATAAGGAGAATGATATGTATGCTGGGTATCATGAAAGT
GATACTTCAGAAGCTGCTAATGAAGCAAAGATTCAAAAACCTTAAAGAGCTGTCCAA
GCTCCCGGGCATTATGATAGACTTTCAAGGTCGCTGGCTCCAAGCATTGTTGGGAGCT
TGAAGATATTA AAAAAGGGTCTTCTTTGCCAGCTCTTTGGTGGGAAGGCTAAGAAAAT
TCCATCTGGAGCATCTTTCCGAGGTGACATCAATGTTTTACTTGTTGGGGACCCTGGT
ACCAGTAAATCTCAGCTGCTTCAGTATGTGCACAAGATAGCTCCTCGTGGAATCTAC
ACTAGTGGGCGAGGAAGTTCGGCGGTTGGGCTGACAGCGTATGTAACGAAGGATCC
AGAAACTCGAGAGACGGTATTGGAGAGCGGAGCTTTGGTTCTTAGTGATCGTGGGA
TATGCTGTATCGATGAGTTCGACAAAATGTCTGATAATGCCCCGAAGCATGCTTCATG
AGGTAATGGAGCAACAAACGGTATCTGTAGCCAAAGGGGGTATCATTGCCTCGCTG
AACGCTCGGACGTCTGTCCTTGCAATGTGCAAATCCTAGTGGGTCCCGATACAATGCG
CGCCTTTCTGTGATTGATAACATCCAGCTTCCTCCAACCTCTACTTTCTAGATTTGATT
TAATTTACTTAATGCTCGACAAACCAGACGAGCAAAACGATCGTCGTCTCGCCAGGC
ATCTCGTGGCTTTACACTATGAAAACCTATGAAGTTTCAAAGCAGGACGCCTTAGATC

FIGURE 2B (Continued)

TACAAACACTTACCGCGTATATCACCTATGCTCGTCAGCATGTACATCCTACATTAA
GTGATGAAGCTGCTGAAGATTTGATTAATGGCTATGTTGAGATGCGCCAAAAGGGC
AACTTTCCTGGAAGCAGTAAAAAGGTGATAACAGCCACACCTCGGCAACTCGAAAG
TATGATTCGTATCAGTGAAGCCCTAGCTCGAATGAGATTTTCTGAAGTGGTAGAGAA
AGTTGATGCAGCAGAAGCTGTGCGCCTTTTAGACGTCGCTTTGCAGCAATCTGCTAC
TGATCATGCAACAGGTACGATAGACATGGATCTTATCACGACTGGAGTGTCGGCCA
GCGAGCGTATTCGTCGGGCCAACTTGCTAGCTGCTCTGCGAGAGCTTATAGCAGATA
AAATTTACCTGGCAGCTCCTCTGGCTTGAAGACCAGTCAGCTTCTTGAGGATATCC
GGAGCCAAAGCAGTGTGGACGTTAGTTTGCAGGATATTAAAAATGCTCTGGGTAGC
CTCCAAGGAGAAGGCTTTCTTACTGTCCATGGTGACATAGTCAAGAGAGTTTGAGAC
AGTTTCTAACTGTTTCGAATCCATGAGCTATAACTCTGAACGAAAGGGAAAACCTCCA
GTTTCCCATGCGCAATTCCCAGAGCTCGC

FIGURE 2C

Nucleotide sequence of the full-length CC-3 from *Physcomitrella patens* (SEQ ID NO:6)

ATCCCGGGTGATACGTTGTGCATATTTGGTGTTGCAAGTTTTTTGGTTTCTTGATTGT
TAGCTATGGCGTCTGCAACAGCGGCTACAATGGCGTCCCTCCTCACGCCTGGGTCTC
TCCGACGCGGTTTGGGTAGCCAGGAATCGTCGACCCAATTTGCTCCCCTAGCTGGTC
CTCGTAAGACATCAGTTTCGCGTAGGGTGACTGCTAGCGCTAGTGGGAAGAACGAC
AATGGAGTCGTGGAAGATGTGGATATGGGGAAGCGGGGTATGTTGAAAGGCGTAGC
GGGAGCTTTGGCTGCAGTTCTCCCTGCTGTTATCGCGAAGAAAGCTTCAGCAGCTGA
GGAGCAGGGCGTAGCGTCTTCCAGGATGTCCTACTCGAGGTTTTTGGAGTATTTGGA
TATGGACCGTGTGAAGAAGGTTGACTTGTATGAAAATGGGACCATAGCAATTGTGG
AGGCTGTATCCCCTGAATTGGGCAACAGAGTACAACGCGTACGCGTGCAGCTCCCC
GGAAGTAGCTCCGAATTGTTGTCGAAGTTTAGATCGAAGAATGTAGATTTTGCCGCA
CACAGCCCACAAGAGGACTCTGGCTCTGTCATTTTGAACCTCATCGGAAATTTGGCT
TTCCCCTTGTTGCTCGTTGGAGGTCTGTTCTTCTTGTCTCGTAGATCCCAAGGTGGTA
TGGGACCTGGCGGTCCTGGGAACCCGATGGCCTTCGGGAAGTCCAAGGCCAAGTTC
CAGATGGAGCCCAACACGGGCATTACATTCCAAGATGTTGCGGGAGTAGACGAGGC
CAAGCAAGACTTCATGGAGGTTGTGGAGTTCTTGAAGCGGCCTGAGAGATTCACAG
CAGTGGGCGCTAAAATCCCAAAGGGAGTGTTGCTGGTTGGACCACCCGGTACCGGA
AAGACTCTATTGGCGAAGGCCATCGCTGGGGAAGCTGGAGTACCATTCTTCTCCATC
TCTGGGTCTGAGTTCGTGGAAATGTTTCGTGGGAGTGGGAGCTTCCCGTGTGAGGGAC
TTGTTCAAGAAGGCGAAAGAGAATGCTCCCTGCATTGTGTTTGTGGATGAGATTGAT
GCCGTTGGAAGACAGAGAGGAACTGGAATTGGAGGAGGCAATGATGAGCGTGAGC
AGACGTTGAATCAGTTGTTGACGGAGATGGACGGTTTCGAAGGAAACACTGGTGTG

FIGURE 2C (Continued)

ATTGTCATTGCTGCCACCAACAGGGCTGATATTCTCGACGCTGCCTTGCTTCGTCCTG
GAAGATTCGACAGACAGGTTTCCGTGGATGTTCCGGACGTGAAGGGAAGGACTGAC
ATCCTCAAGGTGCATGCTAGTAACAAGAAGTTCGCCGATGATGTGTCTCTGGATATC
ATTGCCATGAGGACACCAGGGTTCAGTGGAGCTGACTTGGCCAACCTTGTTGAACGA
AGCAGCTATCTTGACCGGAAGGAGAGGAAAGACAGCCATCAGTGCTAAGGAGATTG
ACGATTCAATTGACAGAATCGTTGCCGGGATGGAAGGTACCGTCATGACGGACGGC
AAGAGCAAGAGTCTTGTTGCATACCACGAAGTCGGCCATGCTATCTGCGGGACTTTG
ACTCCCGGACACGACGCCGTGCAAAAGGTCACCCTCATTCTAGAGGTCAGGCCCG
TGGTCTCACCTGGTTTATTCCCGGAGAAGACCCGACTTTGATTTCAAAGCAGCAAAT
CTTTGCCCGTATTGTTGGTGCTCTTGGTGGTAGGGCTACCGAGCAGGTTGTCTTCGGT
GATGCTGAAGTTACGACTGGAGCGTCCAGTGATTTGCAGCAAGTTACTTCTATGGCC
AAGCAGATGGTTCACAGTATTCGGTATGTCAGACATCGGCCCATGGGCTTTGATGGAC
CCTTCATCCCAGGGAGGAGATATGATTATGCGTATGATGGCACGTAACCTCCATGTCC
GAGAAGTTGGCTGAGGACATCGACAAGGCTGTGAAGGCTATCTCTGACGAGGCCTA
CGAAGTCGCACTGGGTCACATTAGGAACAACCGCACGGCCATGGACAAGATTGTAG
AGGTTCTGCTTGAGAAGGAGACTTTGTCCGGCGCCGAGTTTAGGGCTATTCTTTCGG
AATACACAGAGATTCCTGCTGAAAACCGTGTATCAGACAACCAGGCCGCACCTGTA
GCTGTTTGAAGTGATCAGAAGCAAGGGTGTTTGTGAACACAATCGTGTAGGTTTTGG
AGCATCGACGCTCTTGTATCAGAGCTCGC

FIGURE 3A

Deduced amino acid sequence of CC-1 from *Physcomitrella patens* (SEQ ID NO:7)

MYSNFKEQAIEYVRQAVAEDNAGNYAKAFPLYMNALEYFKTHLKYEKNPKEAITQK
FTEYLRRAEEIRAVLDDGPTGPSANGDAAVQAKPKSKSGKKDGGGGDGDGDSEDPEQQ
KLRSGLNSAIIREKPNVRWADVAGLESAKQALQEAVILPVKFPQFFTGKRRPWRAFLLY
GPPGTGKSYLAKAVATEADSTFFSISSDLVSKWMGESEKLVANLFQMAREAAPSIIFID
EIDSLCGTRGEGNESEASRRIKTELLVQMQGVGNQDTKVLVLAATNTPYSLDQAVRRRF
DKRIYIPLPESKARQHMFKVHLGDTPNNLTERDYEDLARKTDGFSGSDIAVCVKDVLFE
PVRKTQDAMHFKRINTKEGEMWMPCGPREGARQTTMTELA AEGQASKILPPPITKSDF
DKVLAKQRPTVSKGDLIIQEKFTKEFGEEG*

FIGURE 3B

Deduced amino acid sequence of CC-2 from *Physcomitrella patens* (SEQ ID NO:8)

MENNDALDIGAVSSPYPSQSEGVSTPLPQVTSPSFDNAASPVAGRRAVRQTPTSAVRRR
GRETDSARRRRRSRSLGNSVYSSPYDAGTPGTPGTPVATPVYATPVGTPMGTPSFHRG
TPQYKQRSELGSQGKPLHRRRRRSQSREPGHRSPSREPSADGRPSESAEPDDTLGGEYAYV
WGTVNVNIPDVLRAIRRFHNYRSSAHDLSNSKYIQIIETVEREEDTLNIDMSDIYDHDPDL
YAKIVRYPLDIPLLDTECQEVATSLLPTEKHIEARPFNLKASVHMRELNPSDIDKLVS
KGMVIRCSSIPEIKGAFFKCLVCGHSPPLVTVVKGRVEEPTRCCKPECAARNAMSLIHN
CTFANKQIVRLQETPD A IPEGETPHTVSMCLYNTMVDAVKPGDRIEVTGVFKAMAVRV
GPNQRTLRLALYKTYIDCVHVKKSDRGRLQTEDPMEMDKENDMYAGYHESDTSEAANE
AKIQKLKELSKLPGIYDRLSRSLAPSIWELEDIKKGLLCQLFGGKAKKIPSGASF
RGDINVLLVGDPGTSKSQLLQYVHKIAPRGIYTSGRGSSAVGLTAYVTKDPETRET
VLESGALVLSDRGICCIDFEFDKMSDNARSMLHEVMEQQTVSVAKGGIILASLNARTSV
LACANPSGSRYNARLSVIDNIQLPPTLLSRFDLIYLMMLDKPDEQNDRRLARHLVALHY
ENYEVSKQDALDLQTLTAYITYARQH V HPTLSDEAAEDLINGYVEMRQKGNFPGSSKK
VITATPRQLESMIRISEALARMRFSEVVEKVDAAEAVRLLDVALQQSATDHATGTIDMDL
ITTGV SASERIRRANLLAALRELIADKISPGSSSGLKTSQ LLEDIRSQSSVDVSLQDIKNA
LGSLQGEGFLT VHGDIVKRV*

FIGURE 3C

Deduced amino acid sequence of CC-3 from *Physcomitrella patens* (SEQ ID NO:9)

MASATAATMASLLTPGSLRRGLGSQESSTQFAPLAGPRKTSVSRRVTASASGKNDNGV
VEDVDMGKRGMLKGVAGALAAVLPAVIAKKASAAEEQGVASSRMSYSRFLEYLDMD
RVKKVDLYENGTIAIVEAVSPELGNRVQRVRVQLPGTSSELLSKFRSKNVDFAAHSPQE
DSGSVILNLIGNLAFPLLLVGGLFFLSRRSQGGMGPGGPGNPMAFGKSKAKFQMEPNTGI
TFQDVAGVDEAKQDFMEVVEFLKRPERFTA VGAKIPKGVLLVGPPGTGKTLLAKAIAAGE
AGVPFFSISGSEFVEMFVGVGASRVRDLFKKAKENAPCIVFVDEIDAVGRQRGTGIGGG
NDEREQTLNQLLTEMDFEGNTGVIVIAATNRADILDAALLRPGRFDRQVSVDVPDVKG
RTDILKVHASNKKFADDVSLDIAMRTPGFSGADLANLLNEAAILTGRRGKTAISAKEID
DSIDRIVAGMEGTVMTDGKSKSLVAYHEVGHAICGTLTPGHDAVQKVTLIPRGQARGLT
WFIPGEDPTLISKQQIFARIVGALGGRATEQVVFGDAEVTGASSDLQQVTSMAKQMVT
VFGMSDIGPWALMDPSSQGGDMIMRMMARNMSSEKLAEDIDKAVKAISDEAYEVALG
HIRNNRTAMDKIVEVLLEKETLSGAEFRAILSEYTEIPAENRVSDNQAAPVAV*





FIGURE 4

Figure 5

PpCC-1
Drought



Control
Drought





Figure 6










Figure 7





PpCC-2
Drought



Control
Drought

PpCC-3
Drought



Control
Drought






Figure 8