



US00PP13373P2

(12) **United States Plant Patent**
Dümmen

(10) **Patent No.:** **US PP13,373 P2**

(45) **Date of Patent:** **Dec. 17, 2002**

(54) **NEW GUINEA IMPATIENS PLANT NAMED**
‘DUESWEETWHITE’

(75) Inventor: **Marga Dümmen**, Rheinberg (DE)

(73) Assignee: **Dümmen Jungpflanzenkulturen**,
Rheinberg (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/104,378**

(22) Filed: **Mar. 22, 2002**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Search** **Plt./318**

(56) **References Cited**
PUBLICATIONS

UPOV-ROM hit on ‘Duesweetwhite’, GTI-Jouve Retrieval
software, Plant Variety Database 2001/05.*

* cited by examiner

Primary Examiner—Bruce R. Campell

Assistant Examiner—Anne Marie Grünberg

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea Impatiens plant
named ‘Duesweetwhite’, characterized by its compact,
upright, rounded and uniform plant habit; freely branching
and freely flowering habit; white-colored flowers that are
positioned above and beyond the leaves; and dark green-
colored leaves.

1 Drawing Sheet

1

BOTANICAL CLASSIFICATION/CULTIVAR
DENOMINATION

Impatiens hawkeri cultivar Duesweetwhite.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of New Guinea Impatiens plant, botanically known as
Impatiens hawkeri, and hereinafter referred to by the name
‘Duesweetwhite’.

The new Impatiens is a product of a planned breeding
program conducted by the Inventor in Rheinberg, Germany.
The objective of the breeding program is to develop freely
branching New Guinea Impatiens cultivars that flower early
and have large flowers.

The new Impatiens originated from a cross-pollination
made by the Inventor in 1996 of an unidentified proprietary
selection of *Impatiens hawkeri*, not patented, as the female,
or seed parent, with another unidentified proprietary selec-
tion of *Impatiens hawkeri*, not patented, as the male, or
pollen parent. The cultivar Duesweetwhite was discovered
and selected by the Inventor as a flowering plant within the
progeny of the stated cross-pollination in a controlled envi-
ronment in Rheinberg, Germany in 1997.

Asexual reproduction of the new cultivar by terminal
cuttings taken in Rheinberg, Germany since 1998, has
shown that the unique features of this new Impatiens are
stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of
‘Duesweetwhite’. These characteristics in combination dis-
tinguish ‘Duesweetwhite’ as a new and distinct Impatiens
cultivar:

1. Compact, upright, rounded and uniform plant habit.
2. Freely branching and freely flowering habit.
3. White-colored flowers that are positioned above and
beyond the leaves.

2

4. Dark green-colored leaves.

Plants of the new Impatiens differ from plants of the
parent selections primarily in flower color.

Plants of the new Impatiens can be compared to plants of
the cultivar Moorea, disclosed in U.S. Plant Pat. No. 9,147.
In side-by-side comparisons conducted in Rheinberg,
Germany, plants of the new Impatiens differed from plants
of the cultivar Moorea in the following characteristics:

1. Plants of the new Impatiens were shorter than plants of
the cultivar Moorea.
2. Plants of the new Impatiens had shorter internodes than
plants of the cultivar Moorea.
3. Plants of the new Impatiens had smaller leaves than
plants of the cultivar Moorea.
4. Plants of the new Impatiens were more freely flowering
than plants of the cultivar Moorea.
5. Plants of the new Impatiens had smaller flowers than
plants of the cultivar Moorea.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the
overall appearance of the new cultivar, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photograph may differ
from the color values cited in the detailed botanical descrip-
tion which accurately describe the colors of the new Impa-
tiens. The photograph comprises a side perspective view of
a typical flowering plant of ‘Duesweetwhite’ grown in a
container.

DETAILED BOTANICAL DESCRIPTION

The cultivar Duesweetwhite has not been observed under
all possible environmental conditions. The phenotype may
vary somewhat with variations in environment such as
temperature and light intensity without, however, any vari-
ance in genotype.

The aforementioned photograph and following observa-
tions and measurements describe plants grown in Rheinberg,
Germany during the spring, under commercial practice in a

glass-covered greenhouse. Plants were about 16 weeks from cuttings and were grown in 12-cm containers. During the production of the plants, day and night temperatures averaged 18° C. and light levels were about 4,500 lux.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* cultivar Duesweetwhite.

Parentage.

Female parent.—Unidentified proprietary selection of *Impatiens hawkeri*, not patented.

Male parent.—Unidentified proprietary selection of *Impatiens hawkeri*, not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—Summer: About 7 days at 22° C. Winter: About 10 days at 22° C.

Time to produce a rooted cutting.—Summer: About 18 days at 22° C. Winter: About 25 days at 22° C.

Root description.—Fine, fibrous and white in color.

Rooting habit.—Freely branching.

Plant description

General appearance.—Compact, upright, rounded and uniform plant growth habit; freely branching and flowering habit. Moderately vigorous.

Crop time.—From unrooted cuttings, about 16 weeks are required to produce finished flowering plants in 12-cm containers.

Plant height.—About 13 cm.

Plant diameter or spread.—About 25 cm.

Lateral branches.—Quantity per plant: About six. Length: About 8.5 cm. Diameter: About 5 mm. Internode length: About 2.3 cm. Color: 144A.

Foliage description.—Arrangement: Opposite or in whorls. Length: About 8.3 cm. Width: About 2.2 cm. Shape: Ovate. Apex: Apiculate. Base: Obtuse. Margin: Serrulate with ciliation. Texture, upper and lower surfaces: Smooth, glabrous; leathery. Venation pattern: Pinnate. Color: Young and fully expanded foliage, upper surface: 147A. Young and fully expanded foliage, lower surface: 147B. Venation, upper and lower surfaces: 144A.

Petiole.—Length: About 1.8 cm. Diameter: About 2.3 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 144B.

Flower description:

Flower type and flowering habit.—Single white-colored flowers. Freely and continuously flowering;

usually about seven flowers and flower buds per lateral branch. Flowers positioned above and beyond the foliage and typically face upward or outward. Petals self-cleaning; gynoecium persistent. Flowers not fragrant.

Flower longevity.—Flowers last about one week on the plant.

Flowering season.—Year-round under greenhouse conditions; in the garden, flowering from spring until fall. Plants begin flowering about eight weeks after planting.

Flower buds.—Length: About 1 cm. Diameter: About 1 cm. Shape: Ovoid. Color: 144B.

Flower diameter.—About 5 by 4.8 cm.

Flower depth.—About 1.5 cm.

Flower shape.—Rounded; slightly cupped.

Petals.—Quantity: Five per flower, imbricate. Length: About 2.5 cm. Width: About 2.5 cm. Shape: Obcordate. Apex: Emarginate, lobed. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth; satiny. Color: When opening and fully opened, upper surface: 155C. When opening and fully opened, lower surface: 155C.

Spur.—Quantity: One per flower. Length: About 4.5 cm. Diameter: At apex: Less than 1 mm. At flower: About 2.5 mm. Aspect: Curved downward. Color: 154D.

Peduncles.—Length: About 3.3 cm. Diameter: About 1 mm. Strength: Moderately strong; flexible. Color: 144C.

Reproductive organs.—Androecium: Stamen quantity/arrangement: Five fused at anthers, hooded; filaments free. Anther length: About 5 mm. Anther shape: Oval. Anther color: 155C. Pollen amount: Abundant. Pollen color: 11D. Gynoecium: Pistil quantity: One per flower. Pistil length: About 5 mm. Stigma color: 145D. Style length: Less than 1 mm. Style color: 145D. Ovary arrangement: Five-celled. Ovary color: 146A.

Seeds/fruits.—Seed and fruit development has not been observed.

Disease/pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

Temperature tolerance: Plants of the new *Impatiens* have been observed to be tolerant to temperatures ranging from 8 to 35° C.

It is claimed:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Duesweetwhite', as illustrated and described.

* * * * *

