

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199875832 B2**
(10) Patent No. **737862**

(54) Title
Methods and compositions to identify swine genetically resistant to F18 E. Coli associated diseases

(51)⁶ International Patent Classification(s)
C12Q 001/68 C07K 014/47

(21) Application No: **199875832** (22) Application Date: **1998.05.20**

(87) WIPO No: **WO98/53102**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/047181	1997.05.20	US

(43) Publication Date : **1998.12.11**
(43) Publication Journal Date : **1999.02.04**
(44) Accepted Journal Date : **2001.08.30**

(71) Applicant(s)
Biotechnology Research and Development Corporation; United States of America, as Represented by the Secretary of Agriculture, The; Swiss Federal Institute of Technology Zurich

(72) Inventor(s)
Brad T. Bosworth; Peter vogeli

(74) Agent/Attorney
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

(56) Related Art
IMMUNOGENETICS 44, PP76-79 (1996)

OPI DATE 11/12/98 APPLN. ID 75832/98
AQJP DATE 04/02/99 PCT NUMBER PCT/US98/10318



AU9875832

1)

(51) International Patent Classification ⁶ : C12Q 1/68, C07K 14/47		A1	(11) International Publication Number: WO 98/53102
			(43) International Publication Date: 26 November 1998 (26.11.98)
(21) International Application Number: PCT/US98/10318		(74) Agent: MARTIN, Alice, O.; Brinks Hofer Gilson & Lione, NBC Tower, Suite 3600, 455 North Cityfront Plaza Drive, Chicago, IL 60611-5599 (US).	
(22) International Filing Date: 20 May 1998 (20.05.98)			
(30) Priority Data: 60/047,181 20 May 1997 (20.05.97) US		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/047,181 (CIP) Filed on 20 May 1997 (20.05.97)		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicants (for all designated States except US): BIOTECHNOLOGY RESEARCH AND DEVELOPMENT CORPORATION [US/US]; 1815 N. University Street, Peoria, IL 61604 (US). U.S. DEPARTMENT OF AGRICULTURE [US/US]; Washington, DC 20250 (US). SWISS FEDERAL INSTITUTE OF TECHNOLOGY ZURICH [CH/CH]; Ramistrasse 101, ETH-Zentrum, CH-8092 Zurich (CH).		(71) The United States of America, as represented by the Secretary of Agriculture	
(72) Inventors; and (75) Inventors/Applicants (for US only): BOSWORTH, Brad, T. [US/US]; R.R. #1, Box 182, Nevada, IA 50201 (US). VO-GELI, Peter [CH/CH]; Adamenghessdhen #12, CH-8180 Buclad (CH).			
(54) Title: METHODS AND COMPOSITIONS TO IDENTIFY SWINE GENETICALLY RESISTANT TO F18 <i>E. COLI</i> ASSOCIATED DISEASES			
(57) Abstract The present invention provides non-invasive methods and compositions to differentiate, with a high level of sensitivity and specificity, swine that are genetically susceptible to diseases associated with F18 <i>E. coli</i> infection, from resistant swine. DNA polymorphisms in the swine alpha (1,2) fucosyltransferase 1 (<i>FUT1</i>) gene were used to differentiate resistant from susceptible swine. The invention includes a polypeptide with amino acid substitutions, encoded by the nucleotide polymorphisms, a molecular diagnostic assay, and a kit for the differentiation, of <i>E. coli</i> F18-adhesion resistant, heterozygous (carrier) and homozygous susceptible pigs. The molecular test identifies susceptibility to oedema disease and postweaning diarrhea with high sensitivity and specificity, therefore, is useful to swine breeders in their effort to enhance for resistance. Information on the polymorphisms of the present invention provides insight into causation and treatment of <i>E. coli</i> associated intestinal disorders.			



**METHODS AND COMPOSITIONS TO IDENTIFY
SWINE GENETICALLY RESISTANT TO
F18 *E. COLI* ASSOCIATED DISEASES**

5 Compositions and non-invasive methods are provided for the identification of swine genetically resistant to *E. coli* related diseases, in particular, intestinal diseases associated with a strain of *E. coli* bacteria supplied with fimbriae F18. DNA polymorphisms in swine alpha (1,2) fucosyltransferase (*FUT1*) gene were identified that differentiate resistant from susceptible swine and provide a diagnostic test useful for swine
10 breeders.

 A major problem in breeding swine is to keep them disease-free. Intestinal disorders post weaning are a particular problem. A limited number of serotypes of toxigenic *Escherichia (E.) coli* strains are the causative agents of oedema disease and postweaning diarrhea in swine which induce serious economic losses, especially among
15 piglets aged 4 to 12 weeks, in swine breeding farms all over the world. The typical clinical symptoms of oedema disease are neurological signs such as ataxia, convulsions and paralysis. At post mortem examination, oedema is typically present at characteristic sites such as eyelids and forehead, stomach wall and mesocolon. The diseases are caused by Shiga-like toxin-II variant and enterotoxins LT, STa, STb, respectively, produced by
20 *E. coli* that colonize the surface of the small intestine without affecting major morphological changes of the enterocytes (cells in the intestine). Certain types of bacterial *E. coli* strains, F18, F4 (K88) are major, lethal villains in this regard. "Oedema disease of pigs is an enterotoxaemia characterized by generalized vascular damage. The latter is caused by a toxin, Shiga-like toxin-II variant, produced by certain strains of *E. coli*."
25 (Bertschinger *et al.*, 1993). The *E. coli* are distinguished by their pili types. A group of adhesive fimbriae that are related are designated *e.g.*, K88 or F18 (Vögeli *et al.*, 1997).

 Not all swine succumb to *E. coli* infections. Colonization depends on adherence of the bacteria to the enterocytes, which is mediated by the bacterial fimbriae designated *e.g.*, K88 or F18. Susceptibility to adhesion, *i.e.* expression of receptors in swine for
30 binding the fimbriae, has been shown to be genetically controlled by the host and is inherited as a dominant trait with, in the case of F18, *B* being the susceptibility allele and *b* the resistance allele (Vögeli *et al.*, 1996; Meijerink *et al.*, 1997). The genetic locus for this *E. coli* F18-receptor (*ECF18R*) has been mapped to porcine chromosome 6 (SSC6),

based on its close genetic linkage to the S locus and other loci of the halothane (HAL) linkage group on chromosome 6. The receptor for K88 *E. coli* is on chromosome 13.

The mechanism for resistance appears to be that intestinal borders in resistant animals are not colonized by *E. coli*, i.e., the bacteria do not adhere to intestinal walls of resistant swine. Glycoprotein receptors in the brush border membrane of the intestine were shown to be responsible for the differences between adhesive and non-adhesive phenotypes related to some *E. coli*, therefore, the genotype of the host swine determines resistance. The fimbriated bacteria also have been studied (WO 9413811).

Current methods of identifying swine that are resistant to F18 *E. coli* associated diseases are either to 1) collect intestinal samples from swine at slaughter and perform the microscopic adhesion test, 2) challenge the animals with virulent *E. coli* ("colonization test"), or 3) perform blood typing of the A0 (S) blood group system. The first two methods are not practical for identifying resistant animals for use as breeding stock. Although the blood typing method does identify resistant animals, the test is unable to determine whether susceptible animals are homozygous or heterozygous for susceptibility. Knowledge of the genotype of animals with regard to these alleles (conditions of a gene) is essential to develop a successful breeding program. The purpose of the breeding program is to produce swine that resistant to F18 *E. coli* associated diseases that decimate stock post-weaning.

In one publication, the authors stated, in reference to oedema disease in swine, that, "Searches are underway for appropriate genetic markers.." (Bertschinger *et al.*, 1993, p. 87) and citing Walters and Sellwood, 1982:

Breeding resistant swine is an attractive method for prevention of diseases for which an effective prophylaxis is not available. The feasibility of this approach will depend on the prevalence of the gene(s) encoding resistance in the pig population, improved methods for the detection of resistant pigs, and absence of negative genetic traits co-selected with this resistance.

A genetic "marker" locus is a coding or non-coding locus that is close to a genetic locus of interest, but is not necessarily The locus itself. Detectable phenotypes include continuous or discontinuous traits, e.g. restriction length fragment polymorphisms, production traits, bacterial adhesion traits, colorimetric or enzymatic reactions, and antibiotic resistance. The S locus controls expression of The A and O blood group

antigens. Swine homozygous recessive at the *S* locus do not express either A or O blood antigens. A similar condition exists in humans and is due to mutations in the alpha (1,2) fucosyltransferase gene which encodes the human blood group H (Kelly *et al.*, 1994; see also WO 9628967). The porcine alpha (1,2) fucosyltransferase gene of swine has recently
5 been sequenced (Cohney *et al.*, 1996). This gene is very likely the gene present at the *S* locus in swine.

The blood group *H* and *Se* loci have been mapped genetically and physically to human chromosome 19q13.3. This region is evolutionarily conserved, containing genes homologous to the HAL linkage group of genes in pigs. The blood group H encoding
10 gene is the so called *FUT1* whereas the *Se* gene is equivalent to the *FUT2* gene. *FUT1* determines H antigen expression in the erythroid cell lineage, whereas *FUT2* regulates expression of the H antigen in the secretory epithelia and saliva. Conservation of the *FUT1* gene has been shown in lower mammals such as rat and rabbit, and The mRNA expression has been shown in rabbit brain tissue and rat colon. In all these species two
15 types of alpha (1,2) fucosyltransferase genes have been reported which are structurally very similar to the human *FUT1* and *FUT2* genes, but in particular the *FUT1* homologous genes show a species specific expression pattern. In humans the *FUT1* gene is responsible for synthesis of H antigens in the precursors of erythrocytes. However, in pigs erythrocytes passively adsorb H-like antigens from the serum, as is the case for the human
20 *Lewis* antigens. In pigs all H-like antigens are related to exocrine secretory tissues, and expression of The *FUT2* (*Secretor*) gene is seen in secretory tissue of other animal species. Therefore, expression of The porcine A-O blood group determinants which cross-react with anti-human blood group H and A antibodies might be influenced by the *FUT2* gene.

25 Further information about blood groups and *E. coli* swine diseases include that carbohydrate structures of blood group antigens mediate the adhesion of some pathogenic microorganisms to host tissues, *e.g.* *Helicobacter pylori* adhere to Lewis^b blood group antigens, and *E. coli* causing urinary tract infections adhere to blood group *P* substance. Genes encoding glycosyltransferases that are responsible for the formation of the blood
30 group specific carbohydrate structures, therefore, represent candidate genes for the control of bacterial colonization by the host. The localization of these genes is in the same chromosomal region as The locus responsible for adhesion/non-adhesion of F18 positive

E. coli in the swine small intestine. Swine do not express blood group antigens A and O until after weaning, this is the same time that they become susceptible to disease caused by F18 *E. coli*.

New methods of diagnosis and treatment are needed for *E. coli* related intestinal disease in swine. Detection of a genetic mutation was proposed as a diagnostic test for some swine disorders (malignant hyperthermia) (Fujii *et al.*, 1991; U.S. Patent No. 5,358,649), but polymorphic markers were not reported for diagnosis. Vaccines to develop resistance to *E. coli* colonization were described (U.S. Patent No. 5,552,144; WO 8604604), but are unlikely to be a preferred method to prevent the *E. coli* disease because of difficulties in administering live vaccine orally to newborn swine, and because of regulatory restrictions. Antibiotics are available for treatment, but there is no successful prophylaxis.

Summary of the Invention

According to a first embodiment of the invention, there is provided a method for identifying a swine that is resistant to intestinal colonization by *E. coli*, said method comprising:

- a. determining whether a genetic polymorphism at the alpha (1,2) fucosyltransferase locus that is associated with resistance to colonization, is in a biological sample from the swine; and
- b. inferring that the swine is resistant if the swine is homozygous for the polymorphism.

According to a second embodiment of the invention, there is provided a method for identifying a swine that is resistant to *E. coli* associated intestinal disorders, said method comprising:

- a. determining in a biological sample from the swine whether the only nitrogen base at position 307 in the alpha (1,2) fucosyltransferase gene 1 of the swine is adenine; and
- b. identifying the swine is resistant if the only nitrogen base at position 307 is adenine.

According to a third embodiment of the invention, there is provided a method for breeding swine that are resistant to *E. coli* associated diseases, said method comprising:

- a. selecting for breeding swine that have a genetic polymorphism in the alpha (1,2) fucosyltransferase 1 gene that identifies them as swine that are resistant to *E. coli* related intestinal diseases; and



b. breeding the selected swine.

According to a fourth embodiment of the invention, there is provided an isolated DNA molecule which is a polymorphic variant of the alpha (1,2) fucosyltransferase gene 1 in swine, wherein said polymorphic variant is associated with resistance to intestinal colonisation by *E. coli* and/or *E. coli* related intestinal diseases in swine.

According to a fifth embodiment of the invention, there is provided an isolated DNA molecule having a nucleotide sequence in accordance with FIG. 1 wherein said sequence is associated with resistance to intestinal colonisation by *E. coli* and/or *E. coli* related intestinal diseases in swine.

According to a sixth embodiment of the invention, there is provided a polypeptide encoded by the DNA molecule in accordance with the fourth or fifth embodiment of the present invention, said molecule having alpha (1,2) fucosyltransferase activity.

According to a seventh embodiment of the invention, there is provided a molecular assay for detecting *E. coli* receptors, said assay comprising isolating DNA from porcine nucleated cells, amplifying said DNA in a polymerase chain reaction using oligonucleotides as primers which are complementary to a DNA sequence of the porcine alpha (1,2) fucosyltransferase 1 gene, performing a restriction enzyme digest with at least one restriction enzyme, separating the resulting fragments by gel electrophoresis, determining the respective numbers and lengths of fragments, and determining from the numbers and lengths of fragments, which receptors are present.

According to an eighth embodiment of the invention, there is provided a kit for detecting *E. coli* F18 receptors, said kit comprising in separate containers oligonucleotides which are complementary to a DNA sequence of polymorphisms of the porcine alpha (1,2) fucosyltransferase gene 1.

The compositions and non-invasive methods of the present invention provide detection and elimination of swine that are susceptible to *E. coli* associated diseases. A non-invasive method for identifying a swine that is resistant to intestinal colonization by *E. coli* F18 includes the following steps: determining whether a genetic polymorphism associated with resistance to colonization is in a biological sample from the swine; and inferring that the swine is resistant if the swine is homozygous for the polymorphism¹.


More particularly, the method is determining in a biological sample from the swine whether the nitrogen base at position 307 in the alpha (1,2) fucosyltransferase gene of the swine is only adenine or has guanine; and identifying the swine as resistant if the only nitrogen base at position 307 is adenine.

¹ A polymorphism is a change in a nucleotide sequence that exists in a population due to mutation.



4b

To determine whether a polymorphism is present in a biological sample, restriction fragment length polymorphisms are analyzed on a gel that separates them by molecular weight. Restriction endonucleases are enzymes that reproducibly cut nucleic acid molecules at specific sites, resulting in nucleic acid fragments of different molecular weights, depending on the locations of the cuts.



The invention also relates to a method for breeding swine resistant to *E. coli* associated diseases by selecting for breeding swine that have a genetic polymorphism in the alpha (1,2) fucosyltransferase 1 gene that identifies them as swine that are resistant to *E. coli* related diseases; and breeding the selected swine.

5 An aspect of the invention is a DNA molecule which is polymorphic for the alpha (1,2) fucosyltransferase 1 gene in swine, in particular a sequence in accordance with FIG. 1. Other aspects of the invention are molecules with nucleotide sequences complementary to that of FIG. 1, and polymorphic at resistance producing nucleotide positions.

10 An aspect of the invention is an isolated DNA molecule with a substitution of adenine for guanine in position 307. This molecule may also have a substitution adenine for guanine in position 857. Other isolated DNA molecules of the present invention include those with a mutation at nucleotide position 229 of the sequence of FIG. 1, wherein the codon CTT is changed to TTT, encoding for the amino acid phenylalanine instead of leucine. A mutation at nucleotide position 714 is from GAT → GAC, but there
15 is no accompanying amino acid substitution in the encoded product.

Polypeptides encoded by the DNA molecules of the present invention and having alpha (1,2) fucosyltransferase activity are also aspects of the invention.

A molecular assay for detecting *E. coli* F18 receptors in swine is to (a) isolate
20 DNA from porcine nucleated cells; (b) amplify the DNA in a polymerase chain reaction (PCR) using oligonucleotides as primers which are complementary to a DNA sequence of the porcine alpha (1,2) fucosyltransferase gene 1; (c) perform a restriction enzyme digest with at least one restriction enzyme *e.g.*, CfoI; (d) separate the resulting fragments by gel electrophoresis; and (e) determine the respective numbers and lengths of fragments
25 on the gel; and (f) determine from the numbers and lengths of fragments of *FUT1*, which F18 receptor genotype is present in the porcine cells. Use of the larger amplified fragments disclosed herein for restriction length polymorphism analysis (RFLP), rather than smaller fragments, is less expensive because the DNA bands can be run on agarose gels of relatively low concentration. Also, to produce some of the fragments, only one
30 restriction enzyme is needed for a constant restriction site adjacent to the variable diagnostic site.

A kit for detecting polymorphisms associated with *E. coli* F18 receptors uses oligonucleotides in separate containers which are complementary to a DNA sequence of the porcine alpha (1,2) fucosyltransferase gene 1 that distinguishes resistant from sensitive swine. The test can be performed on swine of any age.

The polymorphisms are also useful to develop drugs to treat swine that have *E. coli*-associated disease. A mutated form of porcine alpha (1,2) fucosyltransferase could interfere with the normal enzyme, preventing it from producing the intestinal receptor for F18.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG.1 show the nucleotide sequence (*FUT1*) (below) and the predicted amino acid sequence (above) of the swine $\alpha 1 \rightarrow 2$ fucosyltransferase polymorphism of the present invention using the *one-letter* amino acid code. The *solid double line* below the amino acid sequences (=) is the putative transmembrane region; the *dotted line* below the amino acid sequence shows three potential N-linked glycosylation sites (...).

□ is where an adenine (A) is substituted for guanine (G) in resistant swine.

* Indicates the termination codon

Abbreviations for the amino acid residues are as follows: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp; and Y, Tyr.

DESCRIPTION OF THE PREFERRED EMBODIMENT

Molecular analysis of DNA polymorphisms associated with resistance of swine to *E. coli* associated diseases facilitated diagnostic assays to select resistant pigs for breeding. Resistant pigs differ from sensitive pigs at the *E. coli* F18 receptor locus as identified by the polymorphic markers of the present invention.

The present invention provides non-invasive methods and compositions to distinguish with a high level of sensitivity and specificity, swine that are genetically susceptible to diseases associated with F18 *E. coli* infection from resistant swine. A DNA polymorphism in the alpha (1,2) fucosyltransferase (*FUT1*) gene was identified that differentiates resistant from susceptible swine. The polymorphism arose by a mutation (change) in a nucleotide sequence leading to a new allele. An allele is a condition of a

gene. In a population there may be many alleles of a gene differing by nitrogen base substitutions, presumably caused by mutations in an ancestral DNA molecule. The coexistence in a population of more than one allele (sometimes referred to as a "variant") is called a genetic polymorphism. Loci at which more than one allele may exist as
5 apparently stable components of a population, is a polymorphic locus. Usually, one of the polymorphic loci is at a low frequency in the population.

As determined from a biological sample, preferably blood, the resistant swine have a polymorphism in their genomes in which the only base detected at position 307 (see FIG. 1) in the nucleotide sequence is adenine, whereas the base in the same position in
10 homozygous susceptible swine is guanine. Heterozygous swine will show both types of DNA and will be susceptible. The polymorphism is a variation of a porcine gene sequence reported by Cohney *et al.*, 1996.

Genetic linkage analysis was performed on families of swine and genetic associations between polymorphisms in FUT1 and disease resistance in outbreed swine were determined. According to the present invention, polymorphisms have been found
15 in the alpha (1,2) fucosyltransferase I gene (*FUT1*). A polymorphism that has a single nucleotide base substitution at position 307 was used to establish a close linkage between the fucosyltransferase gene and the S-system, the *ECF18R* locus and other loci of the *HAL* linkage group.

The detection of the close linkage of the mutation at *FUT1* and *ECF18R* allowed a molecular test to be developed for the identification of *E. coli* F18 adhesion resistant, heterozygous (carrier) and homozygous susceptible pigs. This diagnostic test identifies, with high sensitivity and specificity, pigs that are susceptible to oedema disease and postweaning diarrhea. The incidence of the polymorphisms of the present invention
25 differs among swine breeds. Vögel, et al. (1997) presented frequencies of the M307 allele in 5 pig breeds from herds that were not related to one another. The availability of the diagnostic test for the polymorphism of the present invention provides breeders with the opportunity to effectively eliminate the *ECF18R* susceptible allele from their swine herds, thereby eliminating a prerequisite for *E. coli* F18 bacterial causing oedema disease and
30 postweaning diarrhea.

The present invention further includes nucleotide sequences that are variants of a sequence of the alpha (1,2) fucosyltransferase gene 1 representing the various

polymorphisms at bp 307, and diagnostic molecular based kits to identify polymorphisms in the alpha (1,2) fucosyltransferase gene.

In order to obtain candidate genes for the *E. coli* F18 receptor locus (*ECF18R*) 5 cosmids and one genomic clone containing the gene were isolated containing the alpha (1,2) fucosyltransferase genes, *FUT1* and *FUT2* (Meijerink *et al.*, 1997), from a porcine genomic library. Mapping by fluorescence *in situ* hybridization placed all these clones in band q11 of porcine chromosome 6 (SSC6q11). Sequence analysis of the cosmids resulted in the characterization of (a) an open reading frame (ORF), 1098 base pairs in length, that is 82.3% identical to the human *FUT1* sequence, and (b) a second ORF, 1023 base pairs in length which is 85% identical to human *FUT2* sequence. The *FUT1* and *FUT2* loci therefore seem to be porcine equivalents of the blood group *H* and the *Secretor* locus. Direct sequencing of the two ORFs in swine either susceptible or resistant to adhesion and colonization by F18 fimbriated *E. coli* (*ECF18R*) revealed two polymorphisms at base pair 307 (M307) and at base pair 857 (M857) of *FUT1* ORF. The nucleotide positions are numbered from ATG (encoding methionine). The open reading frame (ORF) extends from there to the terminating codon, that is 1098 nucleotides in length. Nucleotides before and after this ORF may vary, *e.g.* as in FIG. 1. Analysis of these mutations in 34 matings of Landrace families with 221 progeny showed close linkage with locus controlling resistance and susceptibility to *E. coli* F18 adhesion and colonization in the small intestine (*ECF18R*) and with the locus of the blood group inhibitor *S*. Therefore, the M307 mutation is a good marker for marker-assisted selection of *E. coli* F18 adhesion resistant animals. Another mutation at nucleotide position 229 was found leading to a polymorphism in which the codon encoding leucine (CTT) was changed to TTT (encoding phenylalanine). A mutation at position 714 (GAT → GAC) (encoding aspartic acid) did not produce an amino acid substitution. No polymorphisms were identified in *FUT2* that differentiated susceptible and resistant pigs.

- 9 -

Examples

The following examples provide embodiments of the invention.

Example 1: An Assay for Resistant Swine

5 The polymorphisms of the present invention are easily identified using PCR-RFLP tests. One embodiment of the tests used a 160bp fragment of porcine alpha (1,2) fucosyltransferase 1 amplified using PCR with the following primers: 5'CCAACGCCTCCGATTCCTGT3' and 5'GTGCATGGCAGGCTGGATGA3' (Table 1). Preferred PCR conditions for this embodiment are 25 cycles at the following times and temperatures: 94°C, 30 sec; 60°C, 45 sec; 72°C, 90 sec. The amplified DNA from resistant swine was digested by the restriction enzyme HgaI, but was not digested by the restriction enzyme HinPI. The amplified DNA from homozygous susceptible swine was digested by the restriction enzyme HinPI. The amplified DNA from heterozygous susceptible swine was partially digested by both enzymes.

15 Alternatively, DNA was isolated from porcine nucleated cells according to standard procedures. Direct sequencing of porcine *FUT1* and *FUT2* sequences and their flanking regions in animals of different ECF18R genotype (Bb, bb) resulted in the identification of two G → A transitions at positions 307 and 857 (termed *M307* and *M857*, respectively) of *FUT1* ORF. The *M307* transition eliminates a restriction site for CfoI. Amplification of DNA isolated from porcine nucleated cells was performed according to standard procedures with primers P6 and P11 (3 min at 95°C, 30 cycles of 30 sec at 95°C, 30 sec at 56°C and 30 sec at 72°C, followed by a 7 min final extension at 72°C) followed by CfoI digestion and separation on a 3% agarose gel resulted in a restriction fragment length polymorphism (RFLP). Homozygous *M307^{AA}* animals showed 2 bands. 20 Homozygous *M307^{GG}* animals showed 93-, 241- and 87bp fragments. Heterozygous animals showed all four fragments.

Example 2: Sensitivity and Specificity of an Assay using Alpha (1,2) Fucosyltransferase in Detecting Swine Resistant to *F18 E. coli*

30 A study was conducted to determine the association between disease resistance and the polymorphism at position 307 of the *FUT1* gene. 183 weaned swine (ranging in ages 2-6 months) were obtained from six different breeding herds. Only one of these

herds was known to contain resistant animals before the start of the study, and this herd is known to have a high incidence of porcine stress syndrome. The other 5 herds had no evidence of porcine stress syndrome, and the incidence of disease resistance was unknown. Swine from each herd were randomly selected, humanely euthanized and spleens and samples of small intestine were removed. DNA was extracted from splenic tissue and used in a PCR-RFLP assay described in Example 1. Intestinal cells were purified by scraping the mucosal surface off the intestine, lysing the cells in a hypotonic EDTA solution and washing by centrifugation. The purified intestinal cell brush borders were incubated with F18 *E. coli*. This mixture was examined by phase contrast microscopy. This assay determined if swine were susceptible (intestinal samples had adhering bacteria) or resistant (intestinal samples had no adhering bacteria). The PCR-RFLP assay for the polymorphism correlated with the bacteria-intestinal cell binding assay in 53 of 53 resistant swine and 128 of 130 susceptible swine. Two swine that were determined susceptible using the bacteria-intestinal cell binding assay were incorrectly predicted to be resistant using the PCR-RFLP assay. Two of the six herds examined contained resistant pigs, while only one herd had porcine stress syndrome, demonstrating that the PCR-RFLP assay can identify disease resistant animals in animals that do not have porcine stress syndrome.

Example 3: Localization of *FUT1* on Chromosome 6 (SSC6)

Cosmids ETHs1, -s2, -s3, -s4 and -s6 were identified after screening of the cosmid library with a *FUT1* nucleotide probe obtained from porcine genomic DNA with primers P7 and P10 and were mapped by FISH and DISC-PCR to chromosome 6 in band q11.

Example 4: Identification of the Porcine *FUT1* ORF

Hybridizing KspI, EcoRI and KspI/EcoRI cosmid digests with radiolabelled porcine *FUT1* fragments P6 and P11 and with P7 and P10 for Southern blot analysis revealed identical autoradiography signals for ETHs2, -s4 and -s6, whereas different signals were obtained from cosmids ETHs1 and -s3. From cosmid ETHs2 KspI, subclones 940 bp and 6.2 kb in length were isolated, corresponding to the estimated length of hybridizing KspI fragments on the Southern blot. The sequence results of both subclones were combined to yield a 1501 bp sequence, which was in agreement with

results of direct sequencing of genomic PCR products. The 1501 bp sequence contains an open reading frame (ORF) of 1098 bp corresponding to the human *FUT1* ORF, with 82.3% nucleotide and 80.8% amino acid identity. The ORF encodes a polypeptide.

5 Example 5: Identification of a Porcine *FUT2* and a Pseudogene *FUTP*

ETHs1 has one DNA fragment (2.7 kb) that hybridizes to *FUT1* sequences, whereas ETHs3 has two (2.7 kb and 8.2 kb). (Meijerink *et al.*, 1997). Subcloning and partial sequencing of the 2.7 kb EcoRI fragment of ETHs1 and -s3 confirmed that these two fragments are identical. The sequence is highly similar to the human *FUT2* but
10 shows several changes in the NH₂ and -COOH terminal regions. These changes lead to frame shifts that are not compatible with a conserved ORF, therefore an assumption is that the sequence obtained from the 2.7 kb fragment represents a pseudogene (*FUT2P*). After subcloning of ETHs3 BamHI digests, the hybridizing sequences contained in the 8.2 kb EcoRI fragment were identified. The sequence of the subclones obtained represents a
15 1023 bp ORF and is 85% identical at the nucleotide- and 83% identical at the amino acid level to the human *FUT2* sequence. Many differences in the NH₂ and -COOH terminal regions were observed between the porcine *FUT2* sequence and the *FUT2P* sequence derived from the 2.7 kb fragment. The predicted amino acid sequence corresponds to the partially determined amino acid sequence of the porcine *Secretor* enzymes (Thurin and
20 Błaszczyk-Thurin, 1995). The porcine *FUT1*, *FUT2* and *FUTP* sequences obtained were submitted to GenBank and have accession numbers U70883, U70881 and U70882, respectively. The *FUT1* and *FUT2* genes have highly homologous sequences. This has to be considered in, for example, primer development. Furthermore, *FUT1* and *FUT2* enzyme activity need to be differentiated in further studies.

25

Example 6: Identification of M307 and M857 Mutations and Characterization of M307

DNA was isolated from porcine nucleated cells according to standard procedures. Direct sequencing of porcine *FUT1* and *FUT2* sequences and their flanking regions in animals of different *ECF18R* genotypes (*Bb*, *bb*) resulted in the identification of two G-A
30 transitions at positions 307 and 857 (termed *M307* and *M857*, respectively) of the *FUT1* ORF. The *M307* transition eliminates a restriction site for the enzyme CfoI. Amplification of DNA isolated from porcine nucleated cells was performed according to

- 12 -

standard procedures with primers P6 and P11 (3 min at 95°C, 30 cycles of 30 sec at 95°C, 30 sec at 56°C and 30 sec at 72°C, followed by a 7 min final extension at 72°C) followed by CfoI digestion and separation on a 3% agarose gel resulted in a restriction fragment length polymorphism (RFLP). Homozygous *M307^{AA}* animals showed 2 bands (93- and 328-bp fragments). Homozygous *M307^{AA}* animals showed 87-, 93-, and 241-bp fragments. Heterozygous animals showed all four fragments.

Example 7: Characterization of Mutation M857

The M857 mutation is a transition that eliminates an AclI site. Primer PBEST was designed to mismatch two additional AclI sites at positions 866 and 872. PCR with primers P7 and PBEST (3 min. at 95°C, 30 cycles of 30 sec at 95°C, 30 sec at 56°C and 30 sec at 72°C, followed by a 7 min final extension at 72°C) followed by AclI digestion enables PCR-RFLP analysis on a 3% agarose gel. Homozygous *M857^{AA}* animals show a 174 bp fragment while amplification products of *M857^{GG}* animals show 136- and 38- bp fragments.

Example 8: Genetic Mapping of the *FUT1* Gene

In Landrace swine families, recombination events between *M307* and the loci of the HAL linkage group (*S*, *ECF18R*, *RYR1*, *GP1*, *PGD*) revealed recombination fractions $\theta < 0.04$ (Table 2). The lodscores *Z* for the overall recombination fractions were between 24.5 and 50.6, showing strong evidence for linkage between these loci. These data allow genetic mapping of the *FUT1* gene to the HAL linkage group in close proximity of *S* and *ECF18R* which are both influenced by *FUT1*. In the experimental Landrace families, allelic association was found between *ECF18R* and *RYR1*. An excess of genotypes *RYR1^{TT}* at position 1843 in *RYR1* (halothane susceptible genotype) was observed among pigs resistant to oedema disease and postweaning diarrhea (genotype *ECF18R^{bb}* (Table 3). This allelic association is a result of linkage disequilibrium, that is, deviation of observed haplotype frequencies from expected haplotype frequencies under independent assortment of alleles. Therefore, linkage disequilibrium designates a non-random association of alleles belonging to linked loci. Owing to low recombination rates whoever, no locus order could be determined as being significantly better than others.

AMENDED SHEET

Example 9: Association of M307^A With *ECF18R^b* and M307^G With *ECF18R^b*

In Landrace (SL) and Large White (LW) parental pigs, *ECF18R^b* (the oedema and postweaning diarrhea resistance allele) is 100% associated with M307^A, and *ECF18R^b* (the oedema and postweaning diarrhea susceptibility allele) is 100% associated with M307^G (wherein A=adenine; G=guanine). In SL pigs 88% (30/34) of S⁺ accounted for all *ECF18R^b* and M307^A haplotypes, respectively. The corresponding values for both the S⁺-*ECF18R^b* and S⁺-M307^A haplotypes were 82% (9/11) in Large White pigs. In the experimental SL families, the occurrence of the M857^A allele at the *FUT1* locus was low, and even absent, in LW pigs. Therefore, a significant gametic association was not observed between the alleles of M857 and the alleles of the flanking genes. The G->A transitions at positions *FUT2* 307 and *FUT1* 857 were found with variable frequencies also in Duroc, Hampshire and Pietrain pigs, making it likely that those transitions also occur in other pig breeds.

Example 10: Distribution of *FUT1* Genotypes

Table 4 shows that the distribution of *FUT1* genotypes at nucleotide position 307 among *ECF18R* types was significantly different from the expected ratio under a hypothesis that the two are independent. Of the 119 oedema disease and postweaning diarrhea resistant *ECF18R^{bb}* animals, 118 were determined to have the genotype M307^{A/A} in the DNA-based test. One resistant animal had the genotype M307^{A/G}. Of the 131 susceptible pigs, 130 were M307^{A/G} or M307^{G/G}. One animal, susceptible to *E. coli* adhesion, was shown to be homozygous M307^{A/A} by the DNA-test. The data from this example and example 2, together with other studies suggested that the *FUT1* gene is the gene present at the S locus in swine and the *ECF18* locus. While 4 animals in this example and example 2 contradict this hypothesis, it is probable that these animals were incorrectly phenotyped in regards to disease resistance/susceptibility.

Example 11: Amino Acid Exchanges in Alpha (1,2) Fucosyltransferase

The G-A changes at bp+307 and bp+857 of the alpha (1,2) fucosyltransferase gene 1 results in a predicted amino acid substitution of threonine (neutral-polar) instead of alanine (neutral-nonpolar) and glutamine (neutral-polar) instead of arginine (basic),

- 14 -

respectively which may have functional consequences in the encoded product. A C-T change at bp 229 results in an amino acid substitution of leucine (neutral-nonpolar) instead of phenylalanine (neutral-nonpolar).

5

Table 1: Sequences of Forward-(F) and Reverse-(R) Primers And Their Relative Position To The Porcine *FUT1* And *FUT2* Start Codons²

	Primer Name	Primer Sequence	Position
5	FUT1 P6 (R)	5'-CTTCAGCCAGGGCTCCTTTAAG-3'	+489
	FUT1 P7 (F)	5'-TTACCTCCAGCAGGCTATGGAC-3'	+720
	FUT1 P10 (R)	5'-TCCAGAGTGGAGACAAGTCTGC-3'	+1082
	FUT1 P11 (F)	5'-CTGCCTGAACGTCTATCAAGATC-3'	+69
	FUT1 P16 (F)	5'-AGAGTTTCCTCATGCCACAGG-3'	-90
10	FUT1 P18 (R)	5'-CTGCTACAGGACCACCAGCATC-3'	+1203
	FUT1 PBEST (R)	5'-ACCAGCAGCGCAAAGTCCCTGAC GGGCACGGCCTC-3'	+893
	FUT2 P16 (R)	5'-CTCCCTGTGCCTTGAAGTGAT-3'	+1094
15	FUT2 P17 (F)	5'-AACTGCACTGCCAGCTTCATGC-3'	-83

Table 2: Overall Recombination Fractions (θ), Lodscores (Z) and Number of Informative Animals (N) For *M307* and Loci of the *HAL* Linkage Group In The Landrace Experimental Population

	Locus pair	N	θ	Z
20	<i>S-ECF18R</i>	183	0.01	50.6
	<i>M307-S</i>	183	0.01	50.6
	<i>M307-ECF18R</i>	216	0.01	57.1
	<i>M307-RYRI</i>	198	0.02	47.2
25	<i>M307-GPI</i>	147	0.03	34.2
	<i>M307-PGD</i>	147	0.04	24.5

²Primers *FUT1* P10 and *FUT1* P11 are derived from the human *FUT1* gene.

Table 3: Haplotype Frequencies At the Four Loci [*S-FUT1* (*M307*, *M857*)-*ECF18R-RYRI*] In The Landrace (SL) Experimental Population and Randomly Selected Large White (LW) Pigs.

5	Breed	Haplotype ³ at <i>S</i> , <i>FUT1</i> (<i>M307</i> , <i>M857</i>), <i>ECF18R</i> , <i>RYRI</i>	Frequency ⁴ (number)
	SL	<i>sAGbT</i>	70 (28)
		<i>sAGbC</i>	5 (2)
		<i>sGGBC</i>	15 (6)
		<i>sGABC</i>	10 (4)
10	LW	<i>sAGbC</i>	56 (9)
		<i>sGGBC</i>	31 (5)
		<i>sGGBC</i>	13 (2)

15

³*S*: Suppressor locus for A and O blood types (*S* and *s*).

FUT1 (*M307*): alteration of adenine (A) to guanine (G) at nucleotide 307 of the alpha (1,2) fucosyltransferase (*FUT1*) gene. *FUT1* (*M857*) alteration of adenine (A) to guanine (G) at nucleotide 857 of *FUT1* gene. *ECF18R*: E. coli F18 receptor. The dominant susceptible allele is indicated by *B* and the resistant allele by *b*. *RYRI*: skeletal muscle ryanodine receptor. C (cytosine) is the dominant resistant and T (thymine) the susceptible allele for malignant hyperthermia.

⁴Haplotype frequencies in % and absolute number of haplotypes between brackets.

CHDS01 AOM 46557

AMENDED SHEET

Table 4: Distribution Of The Genotypes, Tetrachoric Correlation (r) And Significance Of The Association (χ^2 And $w \times \chi^2$)⁵ Of The Associated Polymorphic *FUT1* (M307) And *ECF18R* Loci In Landrace (SL) Experimental Population And Randomly Selected Large White (LW) Swine.

Breed	Locus	<i>FUT1</i> M307			r	χ^2	$w \times \chi^2$
		Genotype	A/G	A/A			
SL	<i>ECF18R</i> ⁶	<i>b/b</i>	1	113	0.98	213.1	42.6***
		<i>B/b</i>	106	1			
LW	<i>ECF18R</i> ⁶	Genotype	A/G, G/G	A/A	1.00	29.0	11.6***
		<i>b/b</i>	0	5			
		<i>B/b, B/B</i>	24	0			

⁵A weight factor of $w = 0.2$ (SL) and 0.4 (LW) was applied to correct for the lack of precision resulting from inclusion of related animals in the data, according to Cotterman (1947). *** $p < 0.001$.

⁶Animals of genotype *b/b* at the *ECF18R* locus are resistant and those of genotype *B/b* and *B/B* are susceptible to adhesion of F18ab *E. coli* bacteria.

METHODS

1. Primers

Primers derived from the human *FUT1* gene were used for the amplification of its porcine counterpart from genomic DNA. From the resulting porcine sequences specific primers were designed which were used in further amplification and sequencing reactions (Table 1).

2. Screening of a Porcine Genomic Library

Porcine genomic libraries were screened with either a porcine *FUT1* probe obtained with primers P7 and P10 or a porcine *FUT1* cDNA. A porcine genomic library, constructed in SuperCos 1 (Stratagene, La Jolla, Ca, USA), was screened with an $\alpha^{32}\text{P}$ dATP labeled (Prime It II, Stratagene) *FUT1* probe obtained from porcine genomic DNA with primers P7 and P10. After hybridization of replica filters at 42°C for 15h (50% formamide, $6 \times \text{SSC}$, $5 \times \text{Denhardt's}$, 0.5% SDS, 0.1 mg/ml Salmon Sperm) and washing

twice at 65°C for 30 min. (1 x SSC, 0.1% SDS), positive colonies were identified after exposure (15h, -80°C) to X-ray film.

3. In situ Hybridization of Porcine Metaphase Chromosomes

5 Cosmid clones ETHs1, ETHs2, ETHs3, ETHs4 and ETHs6 were subjected to fluorescence *in situ* hybridization (FISH) (Solinas Toldo, *et al.* 1993) or direct *in situ* chromosomal PCT (DISC PCR) on porcine metaphases. Metaphase chromosomes were Q-banded and photographed before hybridization. The probes were labeled by random priming using biotin-16-dUTP. Signal detection and amplification was performed using
10 avidin-FITC and biotinylated anti-avidin. The chromosomes were counterstained with 4,6-diamidino-2-phenylindole, and the relative positions of the cosmids were determined as described by Solinas Toldo, 1993.

4. Subcloning

15 Enzymatic digests of probe positive genomic colonies were separated on agarose gel, transferred to a nylon membrane, and probe positive bands were subcloned into plasmids for *FUT1* sequencing. The sequence of *FUT1* derived from this method is shown in FIG. 1.

KspI-, EcoRI- and KspI/EcoRI digests of all cosmids were separated on a 0.8%
20 agarose gel and transferred to a Hybond N nylon membrane (Meijerink *et al.*, 1997). This blot was hybridized with $\alpha^{32}\text{P}$ dATP labeled porcine *FUT1* PCR products (primers P6-P11 and P7-P10). Based on the autoradiographic signals, ETHs1, -s2 and -s3 were subjected to further subcloning into pBluescript SK- (Stratagene), and *FUT* sequences were determined from subclones. The sequences of two *FUT*-like open reading frames (ORFs)
25 (*FUT1* and *FUT2*) obtained from cosmids ETHs2 and -s3 were compared in ECF18R positive (*BB/Bb*) and negative (*bb*) animals by direct sequencing of PCR products.

5. Polymerase Chain Reaction and Direct Sequencing

Using the Perkin Elmer Ready Reaction Dye Terminator kit (Perkin Elmer Cetus,
30 Norwalk, CT, USA) and 10 pmol of primer, cycle sequencing was performed with a thermal program consisting of an initial denaturation of 5 min at 95°C, followed by 25 cycles of 30 sec 95°C, 15 sec 50°C and 4 min 60°C. Primers used for amplification and

sequencing of the porcine alpha (1,2) fucosyltransferase genes are listed in Table 1. Additional primers were designed taking the possibility of cross-annealing of primers due to the high similarity of *FUT1*, *FUT2* and the *FUT2* pseudogene into account. Samples were analyzed on a 373A ABI sequencer (Applied Biosystems Inc.) and sequence analysis was performed with the GCG package (Devereux, 1984).

6. Production of Informative Offspring

Single nucleotide polymorphisms were analyzed in 221 Landrace swine produced from 4 boars and 16 sows, and in 29 Large White (LW) swine produced from 9 matings between unrelated swine. In order to produce a large number of informative offspring for the examination of linkage between porcine genes encoding ECF18 receptors and selected polymorphic loci, only informative Landrace matings of the type *B/b* x *b/b* were produced.

7. Colonization Test

In a study of Bertschinger *et al.*, 1993, the above mentioned Landrace swine were also tested for ECF18 susceptibility in a colonization test. For this, swine were inoculated shortly after weaning with bacteria of *E. coli* strain 124/76 of serotype O139:K12(B):H1:F18 (Rippinger *et al.*, 1995). Faecal shedding of the bacteria was monitored daily. The extent of colonization was calculated as the mean of the two highest faecal scores. Swine with a mean faecal score of 3.5, corresponding to 6.7 log colony forming units (CFU)/g or more, were considered susceptible to colonization. This limit was based on a lack of mortality below this value, and on scores obtained from completely resistant litters.

8. Linkage Analysis of Nucleotide Polymorphisms

The results of the single nucleotide polymorphisms were compared with typing data for ECF18R, which were identified in an *in vitro* adhesion assay described by Vögeli *et al.*, (1996), and with typing data for the *GPI*-, *PGD*-, α -1-B-glycoprotein-(A1BG), ryanodine receptor (*RYR1*), *EAH*- and *S*-loci as published by Vögeli *et al.* (1996). Pairwise linkage analysis and calculation of recombination fractions was performed using the CRI-MAP version 2.4 programme (Green *et al.*, 1990). Multipoint linkage analysis was performed by sequential insertion of the above loci into the map. Haplotype

frequencies were calculated from the parental animals, in the Landrace families and from the 8 parental Large White animals which were haplotyped for *ECF18R* from progeny information. Tetrachoric correlations of *ECF18R* and mutations in *FUT1* (*FUT1M307*) (polymorphisms) were calculated on all Landrace and Large White progeny.

5

9. Southern Blot Analysis

Southern blot analysis was performed of cosmids ETHs1, ETHs2 and ETHs3 after digestion with enzymes KspI, EcoRI and KspIEcoRI and separation of 0.8% agarose. Hybridization with an α^{32} PdATP labeled 5' *FUT1* fragment (primers P6-P11) results in the same hybridizing 940 bp band in both the KspI digest and the KspI/EcoRI digest. However, hybridization with a 3' *FUT1* fragment (primers P7-P10) (Meijerink *et al.*, 1997) shows a 6.2 kb KspI band and a 1.1 kb KspI/EcoRI band. Both the 5' and 3' *FUT1* fragments hybridize to the same 4.6 kb EcoRI fragment. This indicates the presence of a KspI site in the *FUT1* gene contained in cosmid ETHs2. Cross hybridization of the 3' *FUT1* fragment detects 2.7 kb and 8.2 kb bands, resulting in the identification of the *FUT2* pseudogene (incomplete ORF) and the *FUT2* gene sequences, respectively.

10

15

10. Restriction Fragment Polymorphisms

Detection of (A) the M307 G to A and (B) the M857 G to A mutation in the porcine *FUT1* gene was achieved by restriction length polymorphism analysis, using various restriction enzymes. Digestion of amplified *FUT1* fragments with CfoI (A) and AclI (B) results in a restriction fragment polymorphism (Meijerink *et al.*, 1997). (A) The *M307^{AA}* genotype generates 328 and 93 bp restriction fragments while the *M307^{GG}* genotype generates 93, 241 and 87 bp fragments and heterozygous *M307^{AG}* genotypes shows all four fragments.

25

(B) Digestion of the *M857^{AA}* genotype generates 174 bp fragments, while it generates 136 and 38 bp fragments in the *M857^{GG}* genotypes, and in *M857^{AG}* genotypes all three fragments are generated.

30

11. Source of Swine

Data of the Swiss Landrace experimental population came from two pedigrees, which were build up at the Institute of Veterinary Bacteriology, University of Zurich. All

- 21 -

other pigs of the Large White, Swiss Landrace, Duroc, Hampshire and Pietrain breeds came from different breeding herds of Switzerland. Other swine were randomly obtained from farms in the U.S. Midwest.

DOCUMENTS CITED:

- 5 Bertschinger *et al.* (1993) *Veterinary Microbiology* 35:79-89.
- Cottermann (1947) *Contrib. Lab. Verlebr. Biol. Univ. Mich.* 33:1-21.
- 5 Cohny *et al.* (1996) *Immunogenetics* 44:76-79.
- Devereux *et al.*, (1984) *Nucleic Acids Res.* 1:387-395.
- 10 Fujii *et al.* (1991) *Science* 253:448-451.
- Green, *et al.* (1990) Documentation for CRI-MAP, Version 2.4, St. Louis: Washington University School of Medicine.
- 15 Kelly *et al.* (1994), *Proc. Natl. Acad. Sci., USA.* 91:5843-5847.
- Meijerink, *et al.* (1997) *Mamm. Genome* 8:736-741.
- Rippinger (1995) *Vet. Microbial.* 45:281-295.
- 20 Solinas Toldo *et al.* (1993) *Mamm. Genome* 4:720-727.
- Thurin and Blaszczyk-Thurin, (1995) *J. Biol. Chem.* 270 (44):26577-26580.
- 25 Vögeli *et al.* (1996) *Animal Genetics* 27:321-328
- Vögeli *et al.* (1997) *Schweiz. Arch. Tierheilk.* 139:479-484.
- 30 U.S. Pat. No. 5,358,649, MacIennnon *et al.*
- U.S. Pat. No. 5,552,144, Valery *et al.*
- WO 8604604 987P, Inventor: Peterson.
- 35 WO 9628967, Inventor: Koike, C.
- WO 9413811, Inventors: Imbrechts and Lintermans.
- 40 TW 266264, Inventors: Jeng and Liou.

The claims defining the invention are as follows:

1. A method for identifying a swine that is resistant to intestinal colonisation by *E. coli*, said method comprising:
 - a. determining whether a polymorphism at the alpha (1,2) fucosyltransferase locus that is associated with resistance to colonisation, is in a biological sample from the swine; and
 - b. inferring that the swine is resistant if the swine is homozygous for the polymorphism.
2. A method for identifying a swine that is resistant to *E. coli* associated intestinal disorders, said method comprising:
 - a. determining in a biological sample from the swine whether the only nitrogen base at position 307 in the alpha (1,2) fucosyltransferase gene 1 of the swine is adenine; and
 - b. identifying the swine as resistant if the only nitrogen base at position 307 is adenine.
3. A method for breeding swine that are resistant to *E. coli* associated diseases, said method comprising:
 - a. selecting for breeding swine that have a genetic polymorphism in the alpha (1,2) fucosyltransferase 1 gene that identifies them as swine that are resistant to *E. coli* related intestinal diseases; and
 - b. breeding the swine.
4. The method of claim 1, 2, or 3, wherein *E. coli* is strain F18.
5. An isolated DNA molecule which is a polymorphic variant of the alpha (1,2) fucosyltransferase gene 1 in swine, wherein said polymorphic variant is associated with resistance to intestinal colonisation by *E. coli* and/or *E. coli* related intestinal diseases in swine.
6. An isolated DNA molecule having a nucleotide sequence in accordance with FIG. 1 wherein said sequence is associated with resistance to intestinal colonisation by *E. coli* and/or *E. coli* related intestinal diseases in swine.



7. The isolated DNA molecule of claim 6 wherein adenine is in nucleotide position 307 instead of guanine.

8. An isolated DNA molecule which is complementary to the nucleotide sequence of claim 6.

5 9. The isolated DNA molecule of claim 6 with a substitution of adenine for guanine in nucleotide position 857.

10. An isolated polypeptide encoded by the DNA molecule of claim 5 or 6, said molecule having alpha (1,2) fucosyltransferase activity.

11. The isolated polypeptide according to claim 10, wherein the DNA molecule
10 has a nucleotide sequence in accordance with Fig. 1, and wherein adenine is in position 307 instead of guanine.

12. The isolated polypeptide according to claim 10 or claim 11, wherein the DNA molecule has a nucleotide sequence in accordance with Fig. 1, and wherein adenine is in position 857 instead of guanine.

13. A portion of the isolated DNA molecule of claim 5 or 6, said portion
15 including the nucleotide sequence that distinguishes *E. coli* colonization resistant from sensitive swine.

14. The nucleotide sequence according to claim 13, wherein the DNA molecule
20 has a nucleotide sequence in accordance with Fig. 1, and wherein adenine is in position 307 instead of guanine.

15. The nucleotide sequence according to claim 13 or claim 14, wherein the DNA molecule has a nucleotide sequence in accordance with Fig. 1, and wherein adenine is in position 857 instead of guanine.

16. A molecular assay for detecting *E. coli* receptors, said assay comprising
25 isolating DNA from porcine nucleated cells, amplifying said DNA in a polymerase chain reaction using oligonucleotides as primers which are complementary to a DNA sequence of the porcine alpha (1,2) fucosyltransferase 1 gene, performing a restriction enzyme digest with at least one restriction enzyme, separating the resulting fragments by gel electrophoresis, determining the respective numbers and lengths of fragments, and
30 determining from the numbers and lengths of fragments, which receptors are present.

17. The molecular assay of claim 16 where the restriction enzyme is *CfoI*.

18. A kit for detecting *E. coli* F18 receptors, said kit comprising in separate containers oligonucleotides which are complementary to a DNA sequence of polymorphisms of the porcine alpha (1,2) fucosyltransferase gene 1.



19. A porcine polypeptide having alpha (1,2) fucosyltransferase activity and an amino acid substitution in position 103.

20. The polypeptide of claim 19, further characterized as having an amino acid substitution in position 286.

5 21. Use of nucleotide polymorphisms of porcine alpha (1,2) fucosyltransferase to develop drugs to treat swine that have *E. coli*-associated disease by mutating the nucleotide polymorphism to produce a molecule that interferes with the normal receptor of *E. coli*.

22. A method for identifying a swine that is resistant to intestinal colonisation by
10 *E. coli*, substantially as hereinbefore described with reference to any one of the examples.

23. A method for breeding swine that are resistant to *E. coli* associated diseases, substantially as hereinbefore described with reference to any one of the examples.

24. Swine identified in accordance with the method of any one of claims 1, 2, 4 or 22.

15 25. Swine bred in accordance with the method of any one of claims 3, 4 or 23.

Dated 2 July, 2001

**Biotechnology Research And Development Corporation
The United States Of America, As Represented By The Secretary
of Agriculture
20 Swiss Federal Institute of Technology Zurich**

**Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON**



FIGURE 1

	M	W	V	P	S	R	R	H	L	C	L	T	F	L	L	V	C			
CT	CGA	GCC	ATG	TGG	GTC	CCC	AGC	CGC	CGC	CAC	CTC	TGT	CTG	ACC	TTC	CTG	CTA	GTC	TGT	17
V	L	A	A	I	F	F	L	N	V	Y	Q	D	L	F	Y	S	G	L	D	51
GTT	TTA	GCA	GCA	ATT	TTC	TTC	CTG	AAC	GTC	TAT	CAA	GAC	CTC	TTT	TAC	AGT	GGC	TTA	GAC	37
L	L	A	L	C	P	D	H	N	V	V	S	S	P	V	A	I	F	C	L	111
CTG	CTG	GCC	CTG	TGT	CCA	GAC	CAT	AAC	GTG	GTA	TCA	TCT	CCC	GTG	GCC	ATA	TTC	TGC	CTG	57
A	G	T	P	V	H	P	N	A	S	D	S	C	P	K	H	P	A	S	F	171
GCG	GGC	ACG	CCG	GTA	CAC	CCC	AAC	GCC	TCC	GAT	TCC	TGT	CCC	AAG	CAT	CCT	GCC	TCC	TTT	77
S	G	T	W	T	I	Y	P	D	G	R	F	G	N	Q	M	G	Q	Y	A	231
TCC	GGG	ACC	TGG	ACT	ATT	TAC	CCG	GAT	GGC	CGG	TTT	GGG	AAC	CAG	ATG	GGA	CAG	TAT	GCC	97
T	L	L	A	L	A	Q	L	N	G	R	Q	A	F	I	Q	P	A	M	H	291
ACG	CTG	CTG	GCC	CTG	GCG	CAG	CTC	AAC	GGC	CGC	CAG	GCC	TTC	ATC	CAG	CCT	GCC	ATG	CAC	117
A	V	L	A	P	V	F	R	I	T	L	P	V	L	A	P	E	V	D	R	351
GCC	GTC	CTG	GCC	CCC	GTG	TTC	CGC	ATC	ACG	CTG	CCT	GTC	CTG	GCG	CCC	GAG	GTA	GAC	AGG	137
H	A	P	W	R	E	L	E	L	H	D	W	M	S	E	D	Y	A	H	L	411
CAC	GCT	CCT	TGG	CGG	GAG	CTG	GAG	CTT	CAC	GAC	TGG	ATG	TCC	GAG	GAT	TAT	GCC	CAC	TTA	157
K	E	P	W	L	K	L	T	G	F	P	C	S	W	T	F	F	H	H	L	471
AAG	GAG	CCC	TGG	CTG	AAG	CTC	ACC	GGC	TTC	CCC	TGC	TCC	TGG	ACC	TTC	TTC	CAC	CAC	CTC	177
R	E	Q	I	R	S	E	F	T	L	H	D	H	L	R	Q	E	A	Q	G	531
CGG	GAG	CAG	ATC	CGC	AGC	GAG	TTC	ACC	CTG	CAC	GAC	CAC	CTT	CGG	CAA	GAG	GCC	CAG	GGG	197
V	L	S	Q	F	R	L	P	R	T	G	D	R	P	S	T	F	V	G	V	591
GTA	CTG	AGT	CAG	TTC	CGT	CTA	CCC	CGC	ACA	GGG	GAC	CGC	CCC	AGC	ACC	TTC	GTG	GGG	GTG	217
H	V	R	R	G	D	Y	L	R	V	M	P	K	R	H	K	G	V	V	G	651
CAC	GTG	CGC	CGC	GGG	GAC	TAT	CTG	CGT	GTG	ATG	CCC	AAG	CGC	TGG	AAG	GGG	GTG	GTG	GGT	237
D	G	A	Y	L	Q	Q	A	M	D	W	F	R	A	R	Y	E	A	P	V	711
GAC	GGC	GCT	TAC	CTC	CAG	CAG	GCT	ATG	GAC	TGG	TTC	CGG	GCC	CGA	TAC	GAA	GCC	CCC	GTG	257
F	V	V	T	S	N	G	M	E	N	C	R	K	N	I	D	T	S	R	G	771
TTT	GTG	GTC	ACC	AGC	AAC	GGC	ATG	GAG	TGG	TGC	CGG	AAG	AAC	ATC	GAC	ACC	TCC	CGG	GGG	277
D	V	I	F	A	G	D	G	R	E	A	A	P	A	R	D	F	A	L	L	831
GAC	GTG	ATC	TTT	GCT	GCC	GAT	GGG	GAG	GCC	GCG	CCC	GCC	AGG	GAC	TTT	GCG	CTG	CTG		297
V	Q	C	N	H	T	I	M	T	I	G	T	F	G	F	W	A	A	Y	L	891
GTG	CAG	TGC	AAC	CAC	ACC	ATC	ATG	ACC	ATT	GGC	ACC	TTC	GGC	TTC	TGG	GCC	GCC	TAC	CTG	317
A	G	G	D	T	I	Y	L	A	N	F	T	L	P	T	S	S	F	L	K	951
GCT	GGT	GGA	GAT	ACC	ATC	TAC	TTG	GCT	AAC	TTC	ACC	CTG	CCC	ACT	TCC	AGC	TTC	CTG	AAG	337
I	F	K	P	E	A	A	F	L	P	E	W	V	G	I	N	A	D	L	S	1011
ATC	TTT	AAA	CCC	GAG	GCT	GCC	TTC	CTG	CCC	GAG	TGG	GTG	GGC	ATT	AAT	GCA	GAC	TTG	TCT	357
P	L	Q	M	L	A	G	P													1071
CCA	CTC	CAG	ATG	TTG	GCT	GGG	CCT	TGA	ACC	AGC	CAG	GAG	CCT	TTC	TGG	AAT	AGC	CTC	GGT	365
CAA	CCC	AGG	GCC	AGC	GTT	ATG	GGT	CTC	CGG	AAG	CCC	GAG	TAA	CTT	CCG	GAG	ATG	CTG	GTG	1131
GTC	CTG	TAG	CAG	GCT	GGA	CAC	TTA	TTT	CAA	GAG	TGA	TTC	TAA	TTG	GCT	GGA	CTC	AGA	GGA	1191
AAC	CCT	GCA	G																	1251
																				1261

AMENDED SHEET