



US 20100173864A1

(19) **United States**

(12) **Patent Application Publication**  
**Cheng et al.**

(10) **Pub. No.: US 2010/0173864 A1**

(43) **Pub. Date: Jul. 8, 2010**

(54) **COMPOSITIONS INCLUDING TRICIRIBINE  
AND ONE OR MORE PLATINUM  
COMPOUNDS AND METHODS OF USE  
THEREOF**

(76) Inventors: **Jin Q. Cheng**, Tampa, FL (US);  
**Said M. Sebti**, Tampa, FL (US)

Correspondence Address:  
**MORGAN LEWIS & BOCKIUS LLP**  
**1111 PENNSYLVANIA AVENUE NW**  
**WASHINGTON, DC 20004 (US)**

(21) Appl. No.: **12/118,868**

(22) Filed: **May 12, 2008**

**Related U.S. Application Data**

(63) Continuation-in-part of application No. 11/096,082,  
filed on Mar. 29, 2005.

(60) Provisional application No. 60/557,599, filed on Mar.  
29, 2004.



**Publication Classification**

(51) **Int. Cl.**  
**A61K 31/70** (2006.01)  
**A61P 35/04** (2006.01)


(52) **U.S. Cl.** ..... **514/49**

(57) **ABSTRACT**

This application encompasses combination therapies including triciribine and related compounds and one or more platinum compounds and compositions with reduced toxicity for the treatment and prevention of tumors, cancer, and other disorders associated with abnormal cell proliferation.




**FIG. 1A**




**FIG. 1B**

**FIG. 1D**


**FIG. 1C**




**FIG. 2A**




**FIG. 2B**



**FIG. 2C**



**FIG. 2D**









FIG. 3A

**FIG. 3B**

**FIG. 3B(cont)**



**FIG. 3C**









FIG. 3C(cont)




**FIG. 4A**




**FIG. 4C**



**FIG. 4D**



**FIG. 4E**






FIG. 5

atgaacgacgtagccattgtgaaggagggtggtgcacaaacgaggggaatatattaaa  
1 -----+-----+-----+-----+-----+ 60  
tacttgctgcatcggtaacacttcctccgcgcgacgtgttgctccocctataataatt  
a M N D V A I V K E G W L H K R G E Y I K -  
acctggcgccacgctacttcctcctcaagaacgatggcaacctttattggctacaaggaa  
61 -----+-----+-----+-----+-----+ 120  
tggaccgcgcggtgcgatgaaggaggagttcttgctacogtggaataaccgatgttcott  
a T W R P R Y F L L K N D G T F I G Y K E -  
cggcctcaggatgtggatcagcgagagtgcccaactcaacaacttctcagtggcacaatgc  
121 -----+-----+-----+-----+-----+ 180  
gocggagtgctacacctagtgcgtctcagggggtgagttggtgaagagtcacogtggtacg  
a R P Q D V D Q R E S P L N N F S V A Q C -  
PstI  
cagctgatgaagacagagcgcccaaggcccaacacctttatcatccogtgcctgcagtggt  
181 -----+-----+-----+-----+-----+ 240  
gtcgactacttctgtctcgccggttcgggttggtgaaatagtaggcgacggacgtcacc  
a Q L M K T E R P R P N T F I I R C L Q W -  
accacagtcattgagcgcaccttccatgtagaaacgocctgaggagcggaagaatgggccc  
241 -----+-----+-----+-----+-----+ 300  
tgggtgctcagtaactcgcgtggaaggtacatctttgcggactcctcgcccttcttaccgg  
a T T V I E R T F H V E T P E E R E E W A -  
accgccattcagactgtggcgatggactcaagaggcaggaagaagagacgatggacttc  
301 -----+-----+-----+-----+-----+ 360  
tggcggtaagtcgacacccgctacctgagttctccgtccttcttctctgctacctgaag  
a T A I Q T V A D G L K R Q E E E T M D F -  
cgatcaggctcaccacagtgaactcaggggctgaagagatggaggtgtccctggccaag  
361 -----+-----+-----+-----+-----+ 420  
gctagtcggagtgggtcactgttgagtcgccgacttctctacctccacagggacgggttc  
a R S G S P S D N S G A E E M E V S L A K -  
cccaagcacccgtgtgaccatgaacgagtttgagtacctgaaactactgggcaagggcacc  
421 -----+-----+-----+-----+-----+ 480  
gggttcgtggcacactggtaacttgctcaaaactcatggactttgatgacccggttcccggtg  
a P K H R V T M N E F E Y L K L L G K G T -

FIGURE 6a

tttgggaaagtgattcttggtaaaagagaaggccacaggccgctactatgccatgaagatc  
451 -----+-----+-----+-----+-----+-----+-----+ 540  
aaacccctttcactaagaccattttctcttccgggtgtccggcgatgatacggtaacttctag  
F G K V I L V K E K A T G R Y Y A M K I -  
ctcaagaaggaggtcatcgctcgccaaggatgaggttgcccacacgcttactgaaaaccgt  
541 -----+-----+-----+-----+-----+-----+ 600  
gagttcttctccagtagcagcggttctctactccaacgggtgtgcgaatgacttttgcca  
L K K E V I V A K D E V A H T L T E N R -  
PstI  
gtcctgcagaactctaggaatcccttcttaccggccctcaagtactcattccagacccac  
601 -----+-----+-----+-----+-----+-----+ 660  
caggacgtcttgagatccgtaggggaaggaaatgccgggagttcatgagtaaggctctgggtg  
V L Q N S R H P F L T A L K Y S F Q T H -  
XhoI  
gancgcctctgcttttgcacatggagtatgccaacggggcgagctcttcttccacctgtct  
661 -----+-----+-----+-----+-----+-----+ 720  
ctggcggagacgaaacagtaacctcatcaggttgcctcccgctcgagaagaaggtggacaga  
D R L C F V M E Y A N G G E L F F H L S -  
cgagagcgcgctgtttctccgaggacccggcccgcttctatggtgcggagattgtgtctgcc  
721 -----+-----+-----+-----+-----+-----+ 780  
gctctcgcgcacaaagaggctctctggcccgggcggaagataaccaagccctctaacacagacgg  
R E R V F S E D R A R F Y G A E I V S A -  
ctggactacttgcaactccgagaagaacgtggtgtaccgggacctgaagctagagaaacctc  
781 -----+-----+-----+-----+-----+-----+ 840  
yacctgatgaacgtgaggctcttcttgcaccacatggcccttgacttcgatctcttggag  
L D Y L H S E K N V V Y R D L K L E N L -  
atgctggacaaggacgggcacatcaagataacggacttcgggctgtgcaaggaggggatac  
841 -----+-----+-----+-----+-----+-----+ 900  
tacgacctgttctgcccgtgtagttctattgcctgaagcccgacacgttctccctag  
M L D K D G H I K I T D F G L C K E G I -  
aaggatggtgccactatgaagacattctgcggaaacgccggagtaacctggccctgaggtg  
901 -----+-----+-----+-----+-----+-----+ 960  
ttcctaccacggtgataacttctgtaagacgccttgccgctcatggaccggggaactccac  
K D G A T M K T F C G T P E Y L A P E V -  
ctggaagacaacgactacggccgtgcagtggaactgggtggggctgggctgtgtcatgtat  
961 -----+-----+-----+-----+-----+-----+ 1020  
gaccttctgttgcgtgatgccggcacgtcacctgaccaccccgaccgcaccagtagata  
L E D N D Y G R A V D W W G L G V V M Y -

FIGURE 6b

gagatgatgtgtggcgcctgcccttctacaaccaggaccaagagaagctgttcgagctg  
1021 -----+-----+-----+-----+-----+-----+-----+ 1080  
ctctactacacaccggcggaaggatgttggctcctggtgctcttcgacaagctcgac

a E M M C G R L P F Y N Q D H E K L F E L -

atcctcatggaggagatccgcttcccgccacactcggccctgaggccaagtccttgcctc  
1081 -----+-----+-----+-----+-----+-----+-----+ 1140  
taggagtagctcctctaggcgaaggcgcggtgtgagccgggactccggttcagggaagcag

a I L M E E I R F P R T L G P E A K S L L -

tcggggctgctcaagaaggaccctacacagaggctcgggtggggctctgaggatgccaag  
1141 -----+-----+-----+-----+-----+-----+-----+ 1200  
aggcccgacgagttcttctgggatgtgtctccgagccaccccgagactcctacggttc

a S G L L K K D P T Q R L G G G S E D A K -

gagatcatgcagcaccgggtcttttccaacatcgtgtggcaggatgtgtatgagaagaag  
1201 -----+-----+-----+-----+-----+-----+-----+ 1260  
ctctagtagctcgtggccaagaaacgggtgttagcacaccgtcctacacatactcttcttc

a E I M Q H R F F A N I V W Q D V Y E K K -

ctgagccacacctttcaagccccaggtcacctctgagactgacaccaggtatttcgatgag  
1261 -----+-----+-----+-----+-----+-----+-----+ 1320  
gactcgggtggaaagttcggggtccagtgagactctgactgtggtccataaagctactc

a L S P P F K P Q V T S E T D T R Y F D E -

gagttcacagctcagatgatcaccatcacgcgcctgatcaagatgacagcatggagtg  
1321 -----+-----+-----+-----+-----+-----+-----+ 1380  
ctcaagtgctgagctctactagtggttagtgcggcggactagttctactgtcgtacctcaca

a E F T A Q M I T I T P P D Q D D S M E C -

gtggacagtgagcgggagggcgaacttccccagttctcctactcagccagtgggcacagcc  
1381 -----+-----+-----+-----+-----+-----+-----+ 1440  
cacctgtcactcgcctccggcggtgaagggggtcaagaggatgagtcggtcacccgtgtcgg

a V D S E R R P H F P Q F S Y S A S G T A -

tga  
1441 --- 1443  
act

a \* -

FIGURE 6c

ATGAATGAGGTGTCTGTCATCAAAGAAGGCTGGCTCCACAAGCGTGGTGAATACATCAAG  
328 --+-----+-----+-----+-----+-----+-----+-----+----- 387  
TACTTACTCCACAGACAGTAGTTTCTCCGACCGAGGTGTTCCGCACCACTTATGTAGTTC

a M N E V S V I K E G W L H K R G E Y I K -

ACCTGGAGGCCACGGTACTTCCTGCTGAAGAGCGACGGCTCCTTCATTGGGTACAAGGAG  
388 --+-----+-----+-----+-----+-----+-----+-----+----- 447  
TGGACCTCCGGTGCCATGAAGGACGACTTCTCGCTGCCGAGGAAGTAACCCATGTTCCCTC

a T W R P R Y F L L K S D G S F I G Y K E -

AGGCCCGAGGCCCTGATCAGACTCTACCCCCCTTAAACAACCTTCTCCGTAGCAGAATGC  
448 --+-----+-----+-----+-----+-----+-----+-----+----- 507  
TCCGGGCTCCGGGGACTAGTCTGAGATGGGGGGAATTTGTTGAAGAGGCATCGTCTTACG

a R P E A P D Q T L P P L N N F S V A E C -

PstI  
|

CAGCTGATGAAGACCGAGAGGCCGCGACCCAAACACCTTTGTCATACGCTGCCTGCAGTGG  
508 --+-----+-----+-----+-----+-----+-----+-----+----- 567  
GTCGACTACTTCTGGCTCTCCGGCGCTGGGTTGTGGAAACAGTATGCGACGGACGTCACC

a Q L M K T E R P R P N T F V I R C L Q W -

ACCACAGTCATCGAGAGGACCTTCCACGTGGATTCTCCAGACGAGAGGGAGGAGTGGATG  
568 --+-----+-----+-----+-----+-----+-----+-----+----- 627  
TGGTGTCACTAGCTCTCCTGGAAGGTGCACCTAAGAGGTCTGCTCTCCCTCCTCACCTAC

a T T V I E R T F H V D S P D E R E E W M -

ApaI  
|

CGGGCCATCCAGATGGTCCGCCAACAGCCTCAAGCAGCGGGCCCCAGGCGAGGACCCCATG  
628 --+-----+-----+-----+-----+-----+-----+-----+----- 687  
GCCCCGTAGGTCTACCAGCGGTTGTCCGAGTTCTCGTCCCGGGGTCCGCTCCTGGGGTAC

a R A I Q M V A N S L K Q R A P G E D P M -

GACTACAAGTGTGGCTCCCCAGTGAAGTCTCCACGACTGAGGAGATGGAAGTGGCGGTC  
688 --+-----+-----+-----+-----+-----+-----+-----+----- 747  
CTGATGTTACACCGAGGGGTCCTGAGGAGGTGCTGACTCCTCTACCTTCACCGCCAG

a D Y K C G S P S D S S T T E E M E V A V -

AGCAAGGCACGGGCTAAAGTGACCATGAATGACTTCGACTATCTCAAACCTCCTTGSCAAG  
748 --+-----+-----+-----+-----+-----+-----+-----+----- 807  
TCGTTCCGTGCCCGATTTCACTGGTACTTACTGAAGCTGATAGAGTTTGAGGAACCGTTC

FIGURE 7a



FIGURE 7b

a V L E D N D Y G R A V D W W G L G V V M -  
TACGAGATGATGTGCGGCCGCTGCCCTTCTACAACCAGGACCACGAGCGCCTCTTCGAG  
1348 -----+-----+-----+-----+-----+----- 1407  
ATGCTCTACTACACGCCGCGGACGGGAAGATGTTGGTCCTGGTGCCTCGCGGAGAAAGCTC.

a Y E M M C G R L P F Y N Q D H E R L F E -  
SacI  
|  
CTCATCCTCATGGAAGAGATCCGCTTCCCGCGCAGGCTCAGCCCCGAGGCCAAGTCCCTG  
1408 -----+-----+-----+-----+-----+----- 1467  
GAGTAGGAGTACCTTCTCTAGGCGAAGGGCGCGTSCGAGTCCGGGGCTCCGCTTCAGGGAC  
|

a L I L M E E I R F P R T L S P E A K S L -  
ApaI  
|  
CTTGCTGGGCTGCTTAAGAAGGACCCCAAGCAGAGGCTTGGTGGGGGGCCAGCGATGCC  
1468 -----+-----+-----+-----+-----+----- 1527  
GAACGACCCGACGAATTCTTCT3GGGTTGCTCTCCGAACACCCCCGGGTCCGTACGG

a L A G L L K K D P K Q R L G G G P S D A -  
AAGGAGGTCATGGAGCACAGGTCTCTTCTCAGCATCACTGGCAGGACGTGGTCCAGAAG  
1528 -----+-----+-----+-----+-----+----- 1587  
TTCTCCAGTACCTCGTGTCCAAGAAGGAGTCTAGTTGACCGTCTGCACCAGGTCTTC

a K E V M E H R F F L S I N W Q D V V Q K -  
SalI  
|  
AAGCTCCTGCCACCCCTTCAAACCTCAGGTCACGTCGGAGGTCGACACAAGGTACTTCGAT  
1588 -----+-----+-----+-----+-----+----- 1647  
TTGAGGACGGTGGGAAGTTTGGAGTCCAGTGCAGGCTCCAGCTGTGTTCATGAAGCTA

a K L L P P F K P Q V T S E V D T R Y F D -  
GATGAATTTACCGCCAGTCCATCACAATCACACCCCTGACCGCTATGACAGCCTGGGC  
1648 -----+-----+-----+-----+-----+----- 1707  
CTACTTAATGGCGGGTCAGGTAGTGTAGTGTGGGGGACTGGCGATACTGTCCGACCCG

a D E F T A Q S I T I T P P D R Y D S L G -  
TTACTGGAGCTGGACAGCGGACCCACTTCCCCAGTTCTCTACTCGGCCAGCATCCGC  
1708 -----+-----+-----+-----+-----+----- 1767  
AATGACCTCGACCTGGTCGCCTGGGTGAAGGGGGTCAAGAGGATGAGCCGGTCTGAGGCG

a L L E L D Q R T H F P Q F S Y S A S I R -  
GAGTCAGCAGTCTCCCCACGCAGAGGACGCACGCTCGCTGCCATCACCGCTGGGTGGTTF  
1768 -----+-----+-----+-----+-----+----- 1827  
CTCACTCGTCAGACGGTGCGTCTCCTGCGTGCAGCGACGGTAGTGCGGACCCACCAA

a E \* A V C P R R G R T L A A I T A G W F -  
TTTACCCCTGCCCGG

FIGURE 7c

1828    --+-----+-    1842  
AAATGGGGACGGGCC  
a            F   T   P   A   R   -

FIGURE 7d

```

      gggctcagaggggagtcacatcatgagcgatgttaccattgtgaaagaaggttgggttcaga
1  -----+-----+-----+-----+-----+ 60
      ccgaggtctccccctcagtagtactcgctacaatggtaacactttcttccaccaccaagtct
c      A Q R G V I M S D V T I V K E G W V Q K -
      agaggggagaatatataaaaaactggaggccaagatacttccctttgaagacagatggct
61 -----+-----+-----+-----+-----+ 120
      tctccctcttatataatTTTTTgacctccggttcttatgaaggaaaacttctgtctaccga
c      R G E Y I K N W R P R Y F I L K T D G S -
      catlcataggatataaagagaaaacctcaagatgtggatttaccttatccctcaacaact
121 -----+-----+-----+-----+-----+ 180
      gtaagtatcctatatatttctcttttgaggttctacacctaaatgggaataggggagttgttga
c      F I G Y K E K P Q D V D L P Y P L N N F -
      tttcagtgggcaaaatgccagttaatgaaaacagaaacgaccaaaagccaaacacatttataa
181 -----+-----+-----+-----+-----+ 240
      aaagtcaccggtttacgggtcaattacttttgtcttctgtggtttcggtttgtgtaaatatt
c      S V A K C Q L M K T E R P K P N T F I I -
      tcagatgtctccagtggaactactgttatagagagaacatttcatgtagatactccagagg
241 -----+-----+-----+-----+-----+ 300
      aguctacagagggtcacctgatgacaatatctctctgttaaagtacatctatgaggtctcc
c      R C L Q W T T V I E R T F H V D T P E E -
                                     PstI
                                     |
      aaagggagaatggacagaagctatccaggctgtagcagacagactgcagagggaagaag
301 -----+-----+-----+-----+-----+ 360
      ttcccttcttacctgtcttcgatagggtccgacatcgtctgtctgacgtctccgttcttc
c      R E E W T E A I Q A V A D R L Q R Q E E -
      aggagagaatgaattgtagtccaacttcacaaatlgataatataggagaggaagagatgg
361 -----+-----+-----+-----+-----+ 420
      tcctctcttacttaacatcagggttgaagtgtttaactattatctctctctctctacc
c      E R M N C S P T S Q I D N I G E E E M D -
      atgcctctacaacccatcataaaagaagacaatgaatgattttgactatttgsaactac
421 -----+-----+-----+-----+-----+ 480
      tacggagatgttgggtagttatttctctctgttacttactaaaactgataaactttgatg
c      A S T T H H K R T M N D F D Y L K L L -
```

FIGURE 8a

taggtaaaggcacttttgggaaagtatttttggttcgagagaaggcaagtggaaaatact  
481 -----+-----+-----+-----+-----+ 540  
atccatttccgtgaaaaccctttcaataaaaccaagctctcttccgttcaccttttatga  
c G K G T F G K V I L V R E K A S G K Y Y -  
atgctatgaagattctgaagaagaagtcattattgcaaaggatgaagtggcacacactc  
541 -----+-----+-----+-----+-----+ 600  
taagatactttcaagacttctttcttcagtaataacgttttctacttcacogtgtgtgag  
c A M K I L K K E V I I A K D E V A H T L -  
taactgaaagcagagpattaaagaacactagacatccctttttaacatccttgaaatatt  
601 -----+-----+-----+-----+-----+ 660  
attgaactttcgtctcataattttctgtgatctgtagggaataattgttaggaactttataa  
c T E S R V L K N T R H P F L T S L K Y S -  
ccttcagacaaaagaccgttttctgttttctgtgatggaatatgttaattggggcgagctgt  
661 -----+-----+-----+-----+-----+ 720  
ggaaggctctgttttctggcaaacacaaaacactaccttatacaattaccccoctcgcaca  
c F Q T K D R L C F V M E Y V N G G E L F -  
ttttccattttgtcgagagagcggtgttctctgaggaccgcacacgtttctatggtgcag  
721 -----+-----+-----+-----+-----+ 780  
aaaaggtaaacagctctctcgcacacaagagactcctggcgtgtgcaaagataccacgtc  
c F H L S R E R V F S E D R T R F Y G A E -  
aaattgtctctgccttggactatctacattccggaaagattgtgtaccgtgatctcaagt  
781 -----+-----+-----+-----+-----+ 840  
tttaacagagacggaaacctgatagatglaaggcccttctaacacatggcactagagttca  
c I V S A L D Y L H S G K I V Y R D L K L -  
tggaagaatctaagtgtggacaagatggccacataaaaattacagattttggacttttga  
841 -----+-----+-----+-----+-----+ 900  
acctcttagattacgacctgtttctaccgggtgtatttttaattgtctaaaacctgaaacgt  
c E N L M L D K D G H I K I T D F G L C K -  
aagaagggatcacagatgcagccaccatgaagacattctgtggcactccagaatatctgg  
901 -----+-----+-----+-----+-----+ 960  
ttcttcctagtgtctacgtcgggtgtacttctgttaagacaccgtgaggtcttatagacc  
c E G I T D A A T M K T F C G T P E Y L A -  
caccagaggtgttagaagataatgactatggccgagcagtagactggtggggcctagggg  
961 -----+-----+-----+-----+-----+ 1020  
gtggtctccacaatcttctattactgataccggctcgtcatctgaccaccccgatcccc  
c P E V L E D N D Y G R A V D W W G L G V -  
ttgtcatgtatgaaatgatgtgtgggaggttacctttctacaaccaggaccatgagaaac  
1021 -----+-----+-----+-----+-----+ 1080

FIGURE 8b



FIGURE 8c




Figure 10






Figure 11




Figure 12





Figure 13



OV3




Figure 14




Figure 15







Figure 16













Figure 18







**COMPOSITIONS INCLUDING TRICIRIBINE  
AND ONE OR MORE PLATINUM  
COMPOUNDS AND METHODS OF USE  
THEREOF**

**[0001]** This application is a continuation-in-part of U.S. application Ser. No. 11/096,082, filed Mar. 25, 2005, which claims the benefit of U.S. provisional patent application No. 60/557,599 filed Mar. 29, 2004, which is incorporated herein by reference.

### 1. FIELD OF THE INVENTION

**[0002]** This application relates to combination therapies including triciribine compounds and one or more platinum compounds and compositions with reduced toxicity for the treatment and prevention of tumors, cancer, and other disorders associated with abnormal cell proliferation.

### 2. BACKGROUND OF THE INVENTION

**[0003]** Cancer is an abnormal growth of cells. Cancer cells rapidly reproduce despite restriction of space, nutrients shared by other cells, or signals sent from the body to stop reproduction. Cancer cells are often shaped differently from healthy cells, do not function properly, and can spread into many areas of the body. Abnormal growths of tissue, called tumors, are clusters of cells that are capable of growing and dividing uncontrollably. Tumors can be benign (noncancerous) or malignant (cancerous). Benign tumors tend to grow slowly and do not spread. Malignant tumors can grow rapidly, invade and destroy nearby normal tissues, and spread throughout the body.

**[0004]** Cancers are classified according to the kind of fluid or tissue from which they originate, or according to the location in the body where they first developed. In addition, some cancers are of mixed types. Cancers can be grouped into five broad categories, carcinomas, sarcomas, lymphomas, leukemias, and myelomas, which indicate the tissue and blood classifications of the cancer. Carcinomas are cancers found in body tissue known as epithelial tissue that covers or lines surfaces of organs, glands, or body structures. For example, a cancer of the lining of the stomach is called a carcinoma. Many carcinomas affect organs or glands that are involved with secretion, such as breasts that produce milk. Carcinomas account for approximately eighty to ninety percent of all cancer cases. Sarcomas are malignant tumors growing from connective tissues, such as cartilage, fat, muscle, tendons, and bones. The most common sarcoma, a tumor on the bone, usually occurs in young adults. Examples of sarcoma include osteosarcoma (bone) and chondrosarcoma (cartilage). Lymphoma refers to a cancer that originates in the nodes or glands of the lymphatic system, whose job it is to produce white blood cells and clean body fluids, or in organs such as the brain and breast. Lymphomas are classified into two categories: Hodgkin's lymphoma and non-Hodgkin's lymphoma. Leukemia, also known as blood cancer, is a cancer of the bone marrow that keeps the marrow from producing normal red and white blood cells and platelets. White blood cells are needed to resist infection. Red blood cells are needed to prevent anemia. Platelets keep the body from easily bruising and bleeding. Examples of leukemia include acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, and chronic lymphocytic leukemia. The

terms myelogenous and lymphocytic indicate the type of cells that are involved. Finally, myelomas grow in the plasma cells of bone marrow. In some cases, the myeloma cells collect in one bone and form a single tumor, called a plasmacytoma. However, in other cases, the myeloma cells collect in many bones, forming many bone tumors. This is called multiple myeloma.

**[0005]** Tumor induction and progression are often the result of accumulated changes in the tumor-cell genome. Such changes can include inactivation of cell growth inhibiting genes, or tumor suppressor genes, as well as activation of cell growth promoting genes, or oncogenes. Hundreds of activated cellular oncogenes have been identified to date in animal models, however, only a small minority of these genes have proven to be relevant to human cancers (Weinberg et al., 1989, *Oncogenes and the Molecular Origins of Cancer* Cold Spring Harbor, N.Y.; Stanbridge and Nowell, 1990, Cell 63: 867-874; Godwin et al., 1992, *Oncogenes and antioncogenes in gynecological malignancies*, in W J Hoskins, C A Perez and R C Young (eds), *Gynecological oncology: principles and practice*, pp 87-116, Lippincott, Philadelphia). The activation of oncogenes in human cancers can result from factors such as increased gene copy number or structural changes. These factors can cause numerous cellular effects, for example, they can result in overexpression of a gene product. Several oncogenes involved in human cancer can be activated through gene overexpression.

**[0006]** It has become apparent that the successive genetic aberrations acquired by cancer cells result in defects in regulatory signal transduction circuits that govern normal cell proliferation, differentiation and programmed cell death (Hanahan et al., 2000, Cell 100: 57-700). This in turn results in fundamental defects in cell physiology which dictate malignancy. These defects include: a) self sufficiency in growth signals (i.e. overexpression of growth factor receptor tyrosine kinases such as EGFR and aberrant activation of downstream signal transduction pathways such as Ras/Raf/Mek/Erk  $\frac{1}{2}$  and Ras/PI3K/Akt), b) resistance to anti-growth signals (i.e. lower expression of TGF $\beta$  and its receptor), c) evading apoptosis (i.e. loss of proapoptotic p53; overexpression of pro-survival Bcl-2; hyperactivation of survival pathways such as those mediated by PI3K/Akt), d) sustained angiogenesis (i.e. high levels of secretion of VEGF) and f) tissue invasion and metastasis (i.e. extracellular proteases and prometastatic integrins) (Hanahan et al., 2000, Cell 100: 57-700).

**[0007]** Receptor tyrosine kinases such as EGFR, ErbB2, VEGFR and insulin-like growth factor I receptor (IGF-1R) are intimately involved in the development of many human cancers including colorectal pancreatic, breast and ovarian cancers (Khaleghpour et al., 2004, Carcinogenesis 25: 241-8.; Sekharam et al., 2003, Cancer Res 63: 7708-16). Binding of ligands such as EGF, VEGF and IGF-1 to their receptors promotes stimulation of the intrinsic tyrosine kinase activity, autophosphorylation of specific tyrosines in the cytoplasmic domain of the receptors and recruitment of signaling proteins that trigger a variety of complex signal transduction pathways (Olayioye et al., 2000, Embo J 19: 3159-3167, Porter et al., 1998, Oncogene 17: 1343-52). This in turn leads to the activation of many tumor survival and oncogenic pathways such as the Ras/Raf/Mek/Erk  $\frac{1}{2}$ , JAK/STAT3 and PI3K/Akt pathways. Although all three pathways have been implicated in colon, pancreatic, breast and ovarian oncogenesis, those that are mediated by Akt have been shown to be critical in many steps of malignant transformation including cell prolifera-

tion, anti-apoptosis/survival, invasion and metastasis and angiogenesis (Datta et al., 1999, Genes Dev 13: 2905-2927).

**[0008]** Akt is a serine/threonine protein kinase (also known as PKB), which has 3 family members Akt1, Akt2 and Akt3. Stimulation of cells with growth or survival factors results in recruitment to the receptors of the lipid kinase phosphoinositide-3-OH-kinase (PI3K) which phosphorylates phosphoinositide-4,5-bisphosphate (PIP<sub>2</sub>) to PIP<sub>3</sub> which recruits Akt to the plasma membrane where it can be activated by phosphorylation on Thr308 and Ser473 (Akt1), Thr308 and Ser474 (Akt2) and Thr308 and Ser472 (Akt3) (Datta et al., 1999, Genes Dev 13: 2905-2927). Thus, PI3K activates Akt by phosphorylating PIP<sub>2</sub> and converting to PIP<sub>3</sub>. The phosphatase PTEN dephosphorylates PIP<sub>3</sub> to PIP<sub>2</sub> and hence prevents the activation of Akt.

**[0009]** The majority of human cancers contain hyperactivated Akt (Datta et al., 1999, Genes Dev 13: 2905-2927; Bellacosa et al., 1995, Int J Cancer 64: 280-285; Sun et al., 2001, Am J Pathol 159: 431-437). In particular, Akt is overexpressed and/or hyperactivated in 57%, 32%, 27% and 36% of human colorectal, pancreatic, breast and ovarian cancers, respectively (Roy et al., 2002, Carcinogenesis 23: 201-205; Altomare et al., 2003, J Cell Biochem 88: 470-476; Sun et al., 2001, Cancer Res 61: 5985-5991; Stal et al., 2003, Breast Cancer Res 5: R37-R44; Cheng et al., 1992, Proc Natl Acad Sci USA 89: 9267-9271; Yuan et al., 2000, Oncogene 19: 2324-2330). Hyperactivation of Akt is due to amplification and/or overexpression of Akt itself as well as genetic alterations upstream of Akt including overexpression of receptor tyrosine kinases and/or their ligands (Khaleghpour et al., 2004, Carcinogenesis 25: 241-248; Sekharam et al., 2003, Cancer Res 63: 7708-7716; Cohen et al., 1998, Biochem Soc Symp 63: 199-210; Muller et al., 1998, Biochem Soc Symp 63: 149-157; Miller et al., 1995, J. Virol 69: 4390-4398; Slamon et al., 1987, Science 235: 177-182; Andrulis et al., 1998, J Clin Oncol 16: 1340-1349) and deletion of the phosphatase PTEN. Proof-of-concept of the involvement of Akt in oncogenesis has been demonstrated preclinically by showing that ectopic expression of Akt induces malignant transformation and promotes cell survival (Sun et al., 2001, Am J Pathol 159: 431-437; Cheng et al., 1997, Oncogene 14: 2793-2801) and that disruption of Akt pathways inhibits cell growth and induces apoptosis (Jetzt et al., 2003, Cancer Res 63: 6697-6706).

**[0010]** Current treatments of cancer and related diseases have limited effectiveness and numerous serious unintended side effects. Despite demonstrated clinical efficacy of many anti-cancer drugs, severe systemic toxicity often halts the clinical development of promising chemotherapeutic agents. Further, overexpression of receptor tyrosine kinases such as EGFR and their ligands such as IGF-1, Akt overexpression and/or loss of PTEN (all of which result in hyperactivation of Akt) are associated with poor prognosis, resistance to chemotherapy and shortened survival time of cancer patients. Current research strategies emphasize the search for effective therapeutic modes with less risk.

**[0011]** Thus, a combination of a tricinibine compound or a derivative thereof and one or more platinum compounds holds promise as a potential combination therapy for treating tumors, cancer, and abnormal cell proliferation.


### 3. SUMMARY OF THE INVENTION

**[0012]** The present invention encompasses novel therapeutic regimens of tricinibine, tricinibine phosphate and related


compounds in combination with one or more platinum compounds to treat tumors or cancer in a subject while limiting systemic toxicity. The invention is based on the discovery that tumors or cancers, which overexpress Akt kinase are particularly sensitive to the cytotoxic effects of TCN and related compounds and a synergistic affect would arise with a combination of one or more platinum compounds. The inventors have determined, contrary to the prior art and experience, how to successfully use tricinibine and one or more platinum compounds to treat tumors and cancer by one or a combination of (i) administering tricinibine and one or more platinum compounds to patients who exhibit enhanced sensitivity to the tricinibine compound and/or the one or more platinum compounds; (ii) use of a described dosage level that minimizes the toxicity of the tricinibine compound and/or one or more platinum compounds but yet still exhibits efficacy; or (iii) use of a described dosage regimen that minimizes the toxicity of the tricinibine compound and/or one or more platinum compounds.

**[0013]** In one illustrative embodiment of the invention, the invention encompasses a composition including:


**[0014]** (i) a compound of the formula I-IV:




Formula I



Formula II



Formula III




Formula IV

[0015] wherein each  $R_2'$ ,  $R_3'$  and  $R_5'$  are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein  $R_2'$ ,  $R_3'$  or  $R_5'$  is independently H or mono-, di- or triphosphate;

[0016] wherein  $R^x$  and  $R^y$  are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid.

[0017]  $R_1$  and  $R_2$  each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl;

[0018] (ii) one or more one or more platinum compounds as in Formula V.



Platinum Compounds

Formula V

[0019] wherein

[0020]  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight


chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$ ;

[0021]  $R_5$  and  $R_6$  are each independently halogen, optionally substituted alkoxy, optionally substituted ester, optionally substituted alkyl amine, aromatic amine, heteroaromatic amine;


[0022] (iii) a pharmaceutically acceptable carrier.

[0023] In another illustrative embodiment, the invention encompasses a method of treating a tumor or cancer in a mammal including administering to the mammal an effective amount of a composition including:


[0024] (i) a compound of formula I-IV:




Formula I



Formula II



Formula III




Formula IV

wherein each  $R_2'$ ,  $R_3'$  and  $R_5'$  are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate pro-drug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein  $R_2'$ ,  $R_3'$  and  $R_5'$  is independently H or mono-, di- or tri-phosphate;

**[0025]** wherein  $R^x$  and  $R^y$  are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid; and

**[0026]**  $R_1$  and  $R_2$  each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl; and

**[0027]** (i) one or more one or more platinum compounds as in formula V.



Formula V

Platinum Compounds

**[0028]** wherein:

**[0029]**  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$ ;

**[0030]**  $R_5$  and  $R_6$  are each independently halogen, optionally substituted alkoxy, optionally substituted ester, optionally substituted alkyl amine, aromatic amine, heteroaromatic amine.

**[0031]** Methods are useful to treat tumors and cancers that are particularly susceptible to the toxic effects of TCN, TCN-P and/or related compounds. In another embodiment, methods are provided for treating a tumor in a mammal, particularly a human that includes (i) obtaining a biological sample from the tumor; (ii) determining whether the tumor overexpresses an Akt kinase, and (iii) treating the tumor that overexpresses Akt kinase with triciribine, triciribine phosphate or a related compound in combination with one or more platinum compounds, as described herein. In another embodiment, the level of Akt kinase expression can be determined by assaying the tumor or cancer for the presence of a phosphorylated Akt kinase, for example, by using an antibody that can detect the phosphorylated form. In another embodiment, the level of

Akt expression can be determined by assaying a tumor or cancer cells obtained from a subject and comparing the levels to a control tissue. In certain embodiments, the Akt can be overexpressed at least 2, 2.5, 3 or 5 fold in the cancer sample compared to the control. In certain embodiments, the overexpressed Akt kinase can be a hyperactivated and phosphorylated Akt kinase.

**[0032]** In another embodiment of the invention, dosing regimens are provided that limit the toxic side effects of TCN and related compounds. In another embodiment, such dosing regimens minimize or eliminate toxic side effects, including, but not limited to, hepatotoxicity, thrombocytopenia, hyperglycemia, vomiting, hypocalcemia, anemia, hypoalbuminemia, myelosuppression, hypertriglyceridemia, hyperamylasemia, diarrhea, stomatitis and/or fever. In another embodiment, the administration of TCN, TCN-P or related compounds provides at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15, 20, or 25% of the subjects.

**[0033]** In another embodiment, a method is provided to treat a subject who has been diagnosed with a tumor by administering to the subject an effective amount of TCN, TCN-P or a related compound and one or more platinum compounds according to a dosing schedule that includes administering the triciribine compound and one or more platinum compounds approximately one time per week for approximately three weeks followed by a one week period wherein the triciribine compound and one or more platinum compounds are not administered. In another embodiment, methods are provided to treat tumor or cancer in a subject by administering to the subject a dosing regimen of 10 mg/m<sup>2</sup> or less of TCN, TCN-P or a related compound and one or more platinum compounds each one time per week. In another embodiment, the triciribine compound and one or more platinum compounds can be administered as a single bolus dose over a short period of time, for example, about 5, 10 or 15 minutes. In further embodiments, dosing schedules are provided in which the triciribine compound and one or more platinum compounds are administered via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the continuous administration can be repeated at least once a week, once every two weeks and/or once a month. In other embodiments, the triciribine compound and one or more platinum compounds can be administered at least once every three weeks. In further embodiments, the compounds can be administered at least once a day for at least 2, 3, 4 or 5 days.

**[0034]** In further embodiments, the triciribine compound and one or more platinum compounds as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration the triciribine compound and one or more platinum compounds can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m<sup>2</sup> of the triciribine compound and one or more platinum compounds disclosed herein can be administered to a subject. The administration of the triciribine compound and one or more platinum compounds can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen can include administering less than 20 mg/m<sup>2</sup> of the triciribine compound and one or more platinum compounds. In one embodiment, less than 10 mg/m<sup>2</sup> of the triciribine compound and less than about 10

mg/kg of one or more platinum compounds can be administered once a week. In further embodiments, dosages of or less than 2 mg/m<sup>2</sup>, 5 mg/m<sup>2</sup>, 10 mg/m<sup>2</sup>, 15 mg/m<sup>2</sup>, 20 mg/m<sup>2</sup>, 25 mg/m<sup>2</sup>, 35 mg/m<sup>2</sup>, 45 mg/m<sup>2</sup>, 55 mg/m<sup>2</sup>, 65 mg/m<sup>2</sup>, 75 mg/m<sup>2</sup>, 85 mg/m<sup>2</sup>, and/or 95 mg/m<sup>2</sup> of the triciribine compound and dosages of or less than 2 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 35 mg/kg, 45 mg/kg, 55 mg/kg, 65 mg/kg, 75 mg/kg, 85 mg/kg, and/or 95 mg/kg one or more platinum compounds can be administered to a subject. In another embodiment, less than 100 mg/m<sup>2</sup> of the triciribine compound can be administered to a subject via continuous infusion once a day for at least five days. In particular embodiments, the triciribine compound and one or more platinum compounds as disclosed herein can be used for the treatment of pancreatic, prostate, colo-rectal and/or ovarian cancer.

**[0035]** In another embodiment, the triciribine compound and one or more platinum compounds and/or therapeutic regimens of the present invention can be used to prevent and/or treat a carcinoma, sarcoma, lymphoma, leukemia, and/or myeloma. In other embodiments of the invention, the triciribine compound and one or more platinum compounds can be used to treat solid tumors. In still further embodiments, the triciribine compound and one or more platinum compounds and compositions disclosed herein can be used for the treatment of a tumor or cancer, such as, but not limited to cancer of the following organs or tissues: breast, prostate, bone, lung, colon, including, but not limited to colorectal, urinary, bladder, non-Hodgkin lymphoma, melanoma, kidney, renal, pancreas, pharynx, thyroid, stomach, brain, and/or ovaries. In a particular embodiment, the triciribine compound and one or more platinum compounds can be used for the treatment of pancreatic, breast, colorectal and/or ovarian cancer. In further embodiments of the present invention, the triciribine compound and one or more platinum compounds disclosed herein can be used in the treatment of angiogenesis-related diseases. In certain embodiments, methods are provided to treat leukemia via continuous infusion of, the triciribine compound and one or more platinum compounds via continuous infusion for at least 24, 48, 72 or 96 hours. In other embodiments, the continuous infusion can be repeated, for example, at least once every two, three or four weeks.

**[0036]** In a particular embodiment, there is provided a method for the treatment of tumors, cancer, and others disorders associated with an abnormal cell proliferation in a host, the method includes administering to the host an effective amount of the triciribine compound and one or more platinum compounds optionally in combination with a pharmaceutically acceptable carrier.

**[0037]** In one embodiment, the triciribine compound and one or more platinum compounds and compositions can be administered in combination and can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

**[0038]** In other embodiments, the triciribine compound and one or more platinum compounds as disclosed herein can be used to treat tumors or cancers resistant to one or more known anti-cancer drugs, including the embodiments of tumors or cancers and compounds disclosed herein. In one embodiment, the triciribine compound and one or more platinum compounds as disclosed herein is administered in an effective amount for the treatment of a patient with a drug resistant tumor or cancer, for example, multidrug resistant tumors or

cancer including, but not limited to, those resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

**[0039]** In certain embodiments, a method is provided including administering to a host in need thereof an effective amount of a triciribine compound and one or more platinum compounds disclosed herein, or pharmaceutical composition including a triciribine compound and one or more platinum compounds, in an effective amount for the treatment of the treatment of tumors, cancer, and others disorders associated with an abnormal cell proliferation in a host.

**[0040]** In another embodiment, a method for the treatment of a tumor or cancer is provided including an effective amount of a compound disclosed herein, or a salt, isomer, prodrug or ester thereof, to an individual in need thereof, wherein the cancer is for example, carcinoma, sarcoma, lymphoma, leukemia, or myeloma. The compound, or salt, isomer, prodrug or ester thereof, is optionally provided in a pharmaceutically acceptable composition including the appropriate carriers, such as water, which is formulated for the desired route of administration to an individual in need thereof. Optionally the compound is administered in combination or alternation with at least one additional therapeutic agent for the treatment of tumors or cancer.

**[0041]** Also within the scope of the invention is the use of a compound disclosed herein or a salt, solvate, prodrug or ester thereof in the treatment of a tumor or cancer, optionally in a pharmaceutically acceptable carrier; and the use of a triciribine compound and one or more platinum compounds disclosed herein or a salt, prodrug or ester thereof in the manufacture of a medicament for the treatment of cancer or tumor, optionally in a pharmaceutically acceptable carrier.

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

**[0042]** FIG. 1 demonstrates the identification of API-2 (triciribine) as a candidate of Akt inhibitor from the NCI Diversity Set. A illustrates the chemical structure of API-2 (triciribine). B demonstrates that API-2 inhibits phosphorylation levels of AKT2 in AKT2-transformed NIH3T3 cells. Wild type AKT2-transformed NIH3T3 cells were treated with API-2 (1  $\mu$ M) for indicated times and subjected to immunoblotting analysis with anti-phospho-Akt-T308 and -S473 antibodies (top and middle panels). The bottom panel shows expression of total AKT2. In C, it is shown that API-2 inhibits three isoforms of Akt. HEK293 cells were transfected with HA-Akt 1, -AKT2 and -AKT3 and treated with API-2 (1  $\mu$ M) or wortmannin (15  $\mu$ M) prior to EGF stimulation, the cells were lysed and immunoprecipitated with anti-HA antibody. The immunoprecipitates were subjected to in vitro kinase assay (top) and immunoblotting analysis with anti-phospho-Akt-T308 (bottom) antibody. Middle panel shows expression of transfected Akt1, AKT2 and AKT3. D illustrates that API-2 did not inhibit Akt in vitro. In vitro kinase assay of constitutively active AKT2 recombinant protein in a kinase buffer containing 1  $\mu$ M API-2 (lane 3).

**[0043]** FIG. 2 demonstrates that API-2 does not inhibit PI3K, PDK1 and the closely related members of AGC kinase family. A demonstrates an in vitro PI3K kinase assay. HEK293 cells were serum-starved and treated with API-2 (1  $\mu$ M) or Wortmannin (15  $\mu$ M) for 30 minutes prior to EGF stimulation. Cells were lysed and immunoprecipitated with anti-p110 $\alpha$  antibody. The immunoprecipitates were subjected to in vitro kinase assay using PI-4-P as substrate. B illustrates the effect of API-2 on in vitro PDK1 activation (top



panel), closed circles show inhibition by API-2. Open circles show inhibition by the positive control staurosporine, which is a potent PDK1 inhibitor ( $IC_{50}=5$  nM). Bottom panels are immunoblotting analysis of HEK293 cells that were transfected with Myc-PDK1 and treated with wortmannin or API-2 prior to EGF stimulation. The immunoblots were detected with indicated antibodies. C illustrates an immunoblot analysis of phosphorylation levels of PKC $\alpha$  with anti-phospho-PKC $\alpha$ -T638 (top) and total PKC $\alpha$  (bottom) antibodies following treatment with API-2 or a nonselective PKC inhibitor Ro31-8220. D shows an in vitro SGK kinase assay. HEK293 cells were transfected with HA-SGK and treated with API-2 or wortmannin prior to EGF stimulation. In vitro kinase was performed with HA-SGK immunoprecipitates using MBP as substrate (top). Bottom panel shows the expression of transfected HA-SGK. E illustrates the results of a PKA kinase assay. Immuno-purified PKA was incubated in ADB buffer (Upstate Biotechnology Inc) containing indicated inhibitors (API-2 or PKAI) and substrate Kemptide. The kinase activity was quantified. In F, a western blot is shown. OVCAR3 cells were treated with API-2 for indicated times. Cell lysates were immunoblotted with indicated anti-phospho-antibodies (panels 1-4) and anti-actin antibody (bottom).

**[0044]** FIG. 3 demonstrates that API-2 inhibits Akt activity and cell growth and induces apoptosis in human cancer cells with elevated Akt. A is a western blot, following treatment with API-2, phosphorylation levels of Akt were detected with anti-phospho-Akt-T308 antibody in indicated human cancer cell lines. The blots were re-probed with anti-total Akt antibody (bottom panels). In B, a cell proliferation assay is shown. Cell lines as indicated in the figure were treated with different doses of API-2 for 24 h and 48 h and then analyzed with CellTiter 96 Cell Proliferation Assay kit (Promega). C provides an apoptosis analysis. Cells were treated with API-2 and stained with annexin V and PI and analyzed by FACSscan.

**[0045]** FIG. 4 shows that API-2 inhibits downstream targets of Akt and exhibits anti-tumor activity in cancer cell lines with elevated Akt in mouse xenograft. In A, it is demonstrated that API-2 inhibits Akt phosphorylation of tuberlin, Bad, AFX and GSK-3 $\beta$ . Following treatment with API-2, OVAR3 cells were lysed and immunoblotted with indicated antibodies. B shows that API-2 inhibits tumor growth. Tumor cells were subcutaneously injected into nude mice with low level of Akt cells on left side and elevated level of Akt cells on right side. When the tumors reached an average size of about 100-150 mm<sup>3</sup>, animals were treated with either vehicle or 1 mg/kg/day API-2. Each measurement represents an average of 10 tumors. C illustrates a representation of the mice with OVCAR3 (right) and OVCAR5 (left) xenograft treated with API-2 or vehicle (control). D shows examples of tumor size (bottom) and weight (top) at the end of experiment. In E, immunoblot analysis of tumor lysates was performed with anti-phospho-Akt-S473 (top) and anti-AKT2 (bottom) antibodies in OVCAR-3-derived tumors that were treated (T3 and T4) and untreated (T1 and T2) with API-2.

**[0046]** FIG. 5 shows that API-2 (triciribine) inhibits Akt kinase activity in vitro. In vitro kinase assay was performed with recombinant of PDK1 and Akt in a kinase buffer containing phosphatidylinositol-3,4,5-P3 (PIP3), API-2 and histone H2B as substrate. After incubation of 30 min, the reactions were separated by SDS-PAGE and exposed in a film.

**[0047]** FIG. 6 provides the mRNA and amino acid sequence of human Akt1, restriction enzyme sites are also noted.

**[0048]** FIG. 7 provides the mRNA and amino acid sequence of human Akt2 restriction enzyme sites are also noted.

**[0049]** FIG. 8 provides the mRNA and amino acid sequence of human Akt3 restriction enzyme sites are also noted.

**[0050]** FIG. 9 shows that cisplatin (CDDP) resistant cell lines endogenously express hyperactive Akt. FIG. 9A shows a graph of cell survival for two ovarian cancer cell lines (A2780S and OV2008) and their cisplatin resistant counterparts (A2780CP and C13, respectively) treated with increasing amounts of cisplatin. The cisplatin resistant variants, A2780CP and C13, demonstrated increased survival in response to cisplatin. FIG. 9B shows that a comparison of cisplatin sensitive ovarian cancer cells, A2780S and OV2008, to their cisplatin resistant counterparts. The upper panels illustrate hyperactive Akt in the cisplatin resistant cell lines C13 and A2780S. The lower panels illustrate comparable total amounts of Akt are present in each cell type.

**[0051]** FIG. 10 illustrates that triciribine (TCN) (API-2) overcomes cisplatin resistance. Cisplatin resistant cell lines, A2780S and C13, were treated with 0, 5, 10, and 20  $\mu$ M cisplatin. The white and striped bars represent treatment with cisplatin only. The black and grey bars represent treatment with cisplatin and 10  $\mu$ M TCN. The co-treatment with TCN decreased cell survival, indicating TCN overcomes cisplatin resistance.

**[0052]** FIG. 11 illustrates the synergistic effect seen with TCN and cisplatin on the cell survival of C13 ovarian cancer cells. C13 cells were treated with 0, 1, 5, 10 and 20  $\mu$ M TCN, with and without 10  $\mu$ M cisplatin. The combination of cisplatin and TCN synergistically reduced cell survival.

**[0053]** FIG. 12 illustrates that TCN enhances the ability of cisplatin to inhibit the growth of C13 ovarian cancels in nude mice xenografts. Nude mice received C13 xenografts, followed by treatment with vehicle (DMSO), cisplatin alone, TCN alone, or cisplatin and TCN, followed by analysis of tumor progression at 7, 14, 21, or 28 days post-inoculation. The left panel (A) shows a graph illustrating that the combination of cisplatin and TCN significantly reduces tumor growth. The right panel (B) depicts representative tumor mass in inoculated mice receiving the represented treatments.

**[0054]** FIG. 13 shows the rapid activation of Akt pathway by mTOR inhibitors in human ovarian cancer cells. The upper panels (A) show western blots of OV3 cell lysates following the treatment with 1 nM mTOR inhibitor rapamycin for the time indicated. Membranes were probed with antibodies specific to phosphorylated isoforms and to the total protein for comparison of Akt, p70S6K and FKHL1. The lower panels (B) illustrate western blots of MCF7 cell lysates following treatment with the mTOR inhibitor RAD001 for the time indicated. Membranes were probed with antibodies specific to phosphorylated isoforms and to the total protein for comparison of Akt and p70S6K.

**[0055]** FIG. 14 shows that activation of AKT pathway via mTOR inhibitors is attenuated by TCN. The left panel (A) illustrates the effect of TCN on Akt phosphorylation with rapamycin treatment in OV3 cells. The right panel (B) illustrates the same effect of TCN on Akt phosphorylation with RAD001 treatment in MCF7 cells.

**[0056]** FIG. 15 illustrates the enhanced effect on cell growth through the combination of TCN with RAD001 or rapamycin. The top left panel (A) is a graph illustrating cell number counts of DU-145 cells over six days treated with control, rapamycin alone, TCN alone, or TCN and rapamycin. The top right panel (B) is a graph illustrating cell number counts of MCF7 cells over six days treated with control, RAD001 alone, TCN alone, or TCN and RAD001. The bottom panel (C) is a graph illustrating cell number counts of OV3 cells over six days treated with control, rapamycin alone, TCN alone, or TCN and rapamycin.

**[0057]** FIG. 16 illustrates the induction of cell survival and chemoresistance through Aurora-A activation of Akt. The top panels (A) show successfully transfection and expression of Aurora-A in A2780S, A2780CP and OV2008 ovarian cancer cell lines. The second panel (B) is a graph illustrating that expression of Aurora-A in A2780S cells increases cell survival and resistance to paclitaxel. The third panel (C) is a graph illustrating that expression of Aurora-A in A2780S cells increases cell survival and resistance to cisplatin. The bottom panel (D) is a graph illustrating that expression of Aurora-A in OV2008 cells increases cell viability and resistance to cisplatin.

**[0058]** FIG. 17 illustrates the effect of Aurora-A on cytochrome c release and Akt activation. FIG. 17A shows immunofluorescence of cytochrome c in A2780S cells transfected with empty vector or Aurora-A treated with or without cisplatin. Cisplatin induced release of cytochrome c is inhibited by expression of Aurora-A. FIG. 17B is a western blot showing expression of Aurora-A increases phosphorylation of Akt in cisplatin sensitive cells. FIG. 17C shows that increased concentrations of Aurora-A increase the phosphorylation and activation of Akt.

**[0059]** FIG. 18 shows the effect of TCN (APT-2) on Aurora-A induced cisplatin resistance. The top two graphs (A and B) show A2780S cells transfected with empty vector or Aurora-A, followed by treatment with cisplatin alone, API-2 alone or API-2 and cisplatin. The bottom two graphs (C and D) show OV2008 cells transfected with empty vector or Aurora-A, followed by treatment with cisplatin alone, API-2 alone or API-2 and cisplatin. TCN diminishes the resistance to cisplatin afforded by Aurora-A expression in both A2780S and OV2008 cells.

**[0060]** FIG. 19 shows a schematic cartoon of the roles API-2, Aurora-A, p53 and Akt play in cell survival.

**[0061]** FIG. 20 shows the effect of the Src family tyrosine kinase inhibitor dasatinib (Sprycel®) on Src phosphorylation and Akt phosphorylation in A375 cells. Inhibition of Src tyrosine kinases increases phosphorylation of Akt.

## 5. DETAILED DESCRIPTION OF THE INVENTION

**[0062]** The inventors have determined, contrary to the prior art and experience, how to successfully use tricinibine compounds in combination with one or more platinum compounds to treat tumors and cancer by one or a combination of (i) administering tricinibine and one or more platinum compounds only to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the tricinibine compound and/or the one or more platinum compounds; (ii) using a described dosage level that minimizes the toxicity of the tricinibine compound and/or the one or more platinum compounds but yet still exhibits efficacy; or (iii) using a

described dosage regimen that minimizes the toxicity of the tricinibine compound and/or the one or more platinum compounds.

### 5.1. DEFINITIONS

**[0063]** As used herein, the term “compounds of the invention” refers to compounds encompassed by formulas I-IV, compounds of formula V, compounds of structures A-I and combinations thereof.

**[0064]** As used herein, the terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth, i.e., proliferative disorders. Examples of such proliferative disorders include cancers such as carcinoma, lymphoma, blastoma, sarcoma, and leukemia, as well as other cancers disclosed herein. More particular examples of such cancers include breast cancer, prostate cancer, colon cancer, squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, e.g., hepatic carcinoma, bladder cancer, colorectal cancer, endometrial carcinoma, kidney cancer, and thyroid cancer.

**[0065]** Other non-limiting examples of cancers are basal cell carcinoma, biliary tract cancer; bone cancer; brain and CNS cancer; choriocarcinoma; connective tissue cancer; esophageal cancer; eye cancer; cancer of the head and neck; gastric cancer; intra-epithelial neoplasm; larynx cancer; lymphoma including Hodgkin's and Non-Hodgkin's lymphoma; melanoma; myeloma; neuroblastoma; oral cavity cancer (e.g., lip, tongue, mouth, and pharynx); pancreatic cancer; retinoblastoma; rhabdomyosarcoma; rectal cancer; cancer of the respiratory system; sarcoma; skin cancer; stomach cancer; testicular cancer; uterine cancer; cancer of the urinary system, as well as other carcinomas and sarcomas.

**[0066]** As used herein, the term “tumor” refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. For example, a particular cancer may be characterized by a solid mass tumor. The solid tumor mass, if present, may be a primary tumor mass. A primary tumor mass refers to a growth of cancer cells in a tissue resulting from the transformation of a normal cell of that tissue. In most cases, the primary tumor mass is identified by the presence of a cyst, which can be found through visual or palpation methods, or by irregularity in shape, texture or weight of the tissue. However, some primary tumors are not palpable and can be detected only through medical imaging techniques such as X-rays (e.g., mammography), or by needle aspirations. The use of these latter techniques is more common in early detection. Molecular and phenotypic analysis of cancer cells within a tissue will usually confirm if the cancer is endogenous to the tissue or if the lesion is due to metastasis from another site.

**[0067]** The term alkyl, as used herein, unless otherwise specified, includes a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of for example C<sub>1</sub> to C<sub>24</sub>, and specifically includes methyl, trifluoromethyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, t-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. The alkyl is optionally substituted, e.g., with one or more substituents such as halo (F, Cl, Br or I), (e.g., CF<sub>3</sub>, 2-Br-ethyl, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>CF<sub>3</sub> or CF<sub>2</sub>CF<sub>3</sub>), hydroxyl (e.g. CH<sub>2</sub>OH), amino CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>NHCH<sub>3</sub> or CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>), alkylamino, arylamino,

alkoxy, aryloxy, nitro, azido  $\text{CH}_2\text{N}_3$ , cyano (e.g.,  $\text{CH}_2\text{CN}$ ), sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, hereby incorporated by reference.

**[0068]** The term lower alkyl, as used herein, and unless otherwise specified, refers to a  $\text{C}_1$  to  $\text{C}_4$  saturated straight, branched, or if appropriate, a cyclic (for example, cyclopropyl) alkyl group, including both substituted and unsubstituted forms.

**[0069]** The term alkylamino or arylamino includes an amino group that has one or two alkyl or aryl substituents, respectively.

**[0070]** The term amino acid includes naturally occurring and synthetic  $\alpha$ ,  $\beta$ ,  $\gamma$  or  $\delta$  amino acids, and includes but is not limited to, amino acids found in proteins, i.e. glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine. In a preferred embodiment, the amino acid is in the L-configuration. Alternatively, the amino acid can be a derivative of alanyl, valinyl, leucyl, isoleuccinyl, prolinyl, phenylalaninyl, tryptophanyl, methioninyl, glycyl, serinyl, threoninyl, cysteinyl, tyrosinyl, asparaginyl, glutaminyl, aspartoyl, glutaroyl, lysinyl, argininyl, histidinyl,  $\beta$ -alanyl,  $\beta$ -valinyl,  $\beta$ -leucinyl,  $\beta\beta$ -isoleuccinyl,  $\beta$ -prolinyl,  $\beta$ -phenylalaninyl,  $\beta$ -tryptophanyl,  $\beta$ -methioninyl,  $\beta$ -glycyl,  $\beta$ -serinyl,  $\beta$ -threoninyl,  $\beta$ -cysteinyl,  $\beta$ -tyrosinyl,  $\beta$ -asparaginyl,  $\beta$ -glutaminyl,  $\beta$ -aspartoyl,  $\beta$ -glutaroyl,  $\beta$ -lysinyl,  $\beta$ -argininyl or  $\beta$ -histidinyl. When the term amino acid is used, it is considered to be a specific and independent disclosure of each of the esters of a natural or synthetic amino acid, including but not limited to  $\alpha$ ,  $\beta$ ,  $\gamma$  or  $\delta$  glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine in the D and L-configurations.

**[0071]** The term "protected" as used herein and unless otherwise defined includes a group that is added to an oxygen, nitrogen, sulfur or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis (see Greene et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons).

**[0072]** The term aryl, as used herein, and unless otherwise specified, includes phenyl, biphenyl, or naphthyl, and preferably phenyl. The aryl group is optionally substituted with one or more moieties such as halo, hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons.

**[0073]** The term alkaryl or alkarylaryl includes an alkyl group with an aryl substituent. The term aralkyl or arylalkyl includes an aryl group with an alkyl substituent.

**[0074]** The term halo, as used herein, includes chloro, bromo, iodo, and fluoro.

**[0075]** The term acyl includes a carboxylic acid ester in which the non-carbonyl moiety of the ester group is selected from straight, branched, or cyclic alkyl or lower alkyl, alkoxyalkyl including methoxymethyl, aralkyl including

benzyl, aryloxyalkyl such as phenoxyethyl, aryl including phenyl optionally substituted with halogen,  $\text{C}_1$  to  $\text{C}_4$  alkyl or  $\text{C}_1$  to  $\text{C}_4$  alkoxy, sulfonate esters such as alkyl or aralkyl sulfonyl including methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxytrityl, substituted benzyl, trialkylsilyl (e.g. dimethyl-t-butylsilyl) or diphenylmethylsilyl. Aryl groups in the esters optimally comprise a phenyl group. The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

**[0076]** As used herein, the term "substantially free of" or "substantially in the absence of" with respect to enantiomeric purity, refers to a composition that includes at least 85% or 90% by weight, preferably 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the designated enantiomer. In a preferred embodiment, in the methods and compounds of this invention, the compounds are substantially free of other enantiomers.

**[0077]** Similarly, the term "isolated" refers to a compound composition that includes at least 85% or 90% by weight, preferably 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the compound, the remainder comprising other chemical species or enantiomers.

**[0078]** The term "independently" is used herein to indicate that the variable, which is independently applied, varies independently from application to application. Thus, in a compound such as  $\text{R}^n\text{XYR}^m$ , wherein  $\text{R}^n$  is "independently carbon or nitrogen," both  $\text{R}^n$  can be carbon, both  $\text{R}^n$  can be nitrogen, or one  $\text{R}^n$  can be carbon and the other  $\text{R}^n$  nitrogen.

**[0079]** The term "pharmaceutically acceptable salt or prodrug" is used throughout the specification to describe any pharmaceutically acceptable form (such as an ester, phosphate ester, salt of an ester or a related group) of a compound, which, upon administration to a patient, provides the compound. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. Pharmaceutically acceptable prodrugs refer to a compound that is metabolized, for example hydrolyzed or oxidized, in the host to form the compound of the present invention. Typical examples of prodrugs include compounds that have biologically labile protecting groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, dephosphorylated to produce the active compound.

**[0080]** The term "pharmaceutically acceptable esters" as used herein, unless otherwise specified, includes those esters of one or more compounds, which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of hosts without undue toxicity, irritation, allergic response and the like, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.


**[0081]** The term "subject" as used herein refers to an animal, preferably a mammal, most preferably a human. Mammals can include non-human mammals, including, but not limited to, pigs, sheep, goats, cows (bovine), deer, mules, horses, monkeys and other non-human primates, dogs, cats, rats, mice, rabbits or any other known or disclosed herein.

## 5.2. COMPOUNDS OF THE INVENTION

**[0082]** The present invention provides for the use of TCN, TCN-P and related compounds in combination with one or

more platinum compounds for use in particular therapeutic regimens for the treatment of proliferative disorders.

**[0083]** As used herein and unless otherwise indicated, the term “tricitabine compound(s)” and “tricitabine and related compounds” refers to compounds having the following structures:



**[0084]** wherein each  $R_2'$ ,  $R_3'$  and  $R_5'$  are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein  $R_2'$ ,  $R_3'$  or  $R_5'$  is independently H or mono-, di- or tri-phosphate;


**[0085]** wherein  $R^x$  and  $R^y$  are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally

substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid;

**[0086]**  $R_1$  and  $R_2$  each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl.

**[0087]** In one embodiment,  $R_2'$  and  $R_3'$  are hydrogen. In another embodiment,  $R_2'$  and  $R_5'$  are hydrogen. In yet another embodiment,  $R_2'$ ,  $R_3'$  and  $R_5'$  are hydrogen. In yet another embodiment,  $R_2'$ ,  $R_3'$ ,  $R_5'$ ,  $R_1$  and  $R_2$  are hydrogen.


**[0088]** In another embodiment, the tricitabine compound has the following structure:




**[0089]** wherein  $R_3$  is H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl,  $NH_2$ ,  $NHR^4$ ,  $N(R^4)_2$ , aryl, alkoxyalkyl, aryloxyalkyl, or substituted aryl; and


**[0090]** Each  $R^4$  independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl, or aryl. In a subembodiment,  $R_3$  is a straight chained C1-11 alkyl, iso-propyl, t-butyl, or phenyl.

**[0091]** In one embodiment, the tricitabine compounds provided herein have the following structure:






**[0092]** In another embodiment, the tricyribine compounds provided herein have the following structure:



**[0093]** In another embodiment, the tricyribine compounds provided herein have the following structure:




**[0094]** wherein  $R_6$  is H, alkyl, (including lower alkyl) alkenyl, alkynyl, alkoxyalkyl, hydroxyalkyl, arylalkyl, cycloalkyl,  $NH_2$ ,  $NHR^4$ ,  $NR^4R^4$ ,  $CF_3$ ,  $CH_2OH$ ,  $CH_2Cl$ ,  $CH_2CF_3$ ,  $C(Y^3)_3$ ,  $C(Y^3)_2C(Y^3)_3$ ,  $C(=O)OH$ ,  $C(=O)OR^4$ ,  $C(=O)$ -aryl,  $C(=O)$ -alkoxyalkyl,  $C(=O)NH_2$ ,  $C(=O)NHR^4$ ,  $C(=O)N(R^4)_2$ , where each  $Y^3$  is independently H or halo; and

**[0095]** each  $R^4$  independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl, or aryl.

**[0096]** In a subembodiment,  $R_6$  is ethyl,  $CH_2CH_2OH$ , or  $CH_2$ -phenyl.

**[0097]** In another embodiment, the tricyribine compounds provided herein have the following structure:





**[0098]** wherein  $R_7$  is H, halo, alkyl (including lower alkyl), alkenyl, alkynyl, alkoxy, alkoxyalkyl, hydroxyalkyl, cycloalkyl, nitro, cyano, OH,  $OR^4$ ,  $NH_2$ ,  $NHR^4$ ,  $NR^4R^4$ , SH,  $SR^4$ ,  $CF_3$ ,  $CH_2OH$ ,  $CH_2F$ ,  $CH_2Cl$ ,  $CH_2CF_3$ ,  $C(Y^3)_3$ ,  $C(Y^3)_2C(Y^3)_3$ ,  $C(=O)OR^4$ ,  $C(=O)$ -alkyl,  $C(=O)$ -aryl,  $C(=O)$ -alkoxyalkyl,  $C(=O)NHR^4$ ,  $C(=O)N(R^4)_2$ , or  $N_3$ , where each  $Y^3$  is independently H or halo; and


**[0099]** each  $R^4$  independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl.

**[0100]** In a subembodiment,  $R_7$  is methyl, ethyl, phenyl, chloro or  $NH_2$ .


**[0101]** In another embodiment, the tricyribine compounds provided herein have the following structure:



[0102] In another embodiment, the triciribine compounds provided herein have the following structure:



[0103] As used herein and unless otherwise indicated, the term “platinum compounds” refers to a compound of formula V.




Platinum Compounds

Formula V


[0104] wherein:

[0105] R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>.

[0106] In an illustrative embodiment, the platinum compounds include, but are limited to, platinum compounds of structures A-I:




A




B


-continued




C




D



E




F



G



H



I

[0107] It is to be understood that the compounds disclosed herein may contain chiral centers. Such chiral centers may be of either the (R) or (S) configuration, or may be a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, or be stereoisomeric or diastereomeric mixtures. It is understood that the disclosure of a compound herein encompasses any racemic, optically active, polymor-

phic, or stereoisomeric form, or mixtures thereof, which preferably possesses the useful properties described herein, it being well known in the art how to prepare optically active forms and how to determine activity using the standard tests described herein, or using other similar tests which are well known in the art. Examples of methods that can be used to obtain optical isomers of the compounds include the following:

[0108] i) physical separation of crystals—a technique whereby macroscopic crystals of the individual enantiomers are manually separated. This technique can be used if crystals of the separate enantiomers exist, i.e., the material is a conglomerate, and the crystals are visually distinct;

[0109] ii) simultaneous crystallization—a technique whereby the individual enantiomers are separately crystallized from a solution of the racemate, possible only if the latter is a conglomerate in the solid state;

[0110] iii) enzymatic resolutions—a technique whereby partial or complete separation of a racemate by virtue of differing rates of reaction for the enantiomers with an enzyme

[0111] iv) enzymatic asymmetric synthesis—a synthetic technique whereby at least one step of the synthesis uses an enzymatic reaction to obtain an enantiomerically pure or enriched synthetic precursor of the desired enantiomer;

[0112] v) chemical asymmetric synthesis—a synthetic technique whereby the desired enantiomer is synthesized from an achiral precursor under conditions that produce asymmetry (i.e., chirality) in the product, which may be achieved using chiral catalysts or chiral auxiliaries;

[0113] vi) diastereomer separations—a technique whereby a racemic compound is reacted with an enantiomerically pure reagent (the chiral auxiliary) that converts the individual enantiomers to diastereomers. The resulting diastereomers are then separated by chromatography or crystallization by virtue of their now more distinct structural differences and the chiral auxiliary later removed to obtain the desired enantiomer;

[0114] vii) first- and second-order asymmetric transformations—a technique whereby diastereomers from the racemate equilibrate to yield a preponderance in solution of the diastereomer from the desired enantiomer or where preferential crystallization of the diastereomer from the desired enantiomer perturbs the equilibrium such that eventually in principle all the material is converted to the crystalline diastereomer from the desired enantiomer. The desired enantiomer is then released from the diastereomer;

[0115] viii) kinetic resolutions—this technique refers to the achievement of partial or complete resolution of a racemate (or of a further resolution of a partially resolved compound) by virtue of unequal reaction rates of the enantiomers with a chiral, non-racemic reagent or catalyst under kinetic conditions;

[0116] ix) enantiospecific synthesis from non-racemic precursors—a synthetic technique whereby the desired enantiomer is obtained from non-chiral starting materials and where the stereochemical integrity is not or is only minimally compromised over the course of the synthesis;

[0117] x) chiral liquid chromatography—a technique whereby the enantiomers of a racemate are separated in a liquid mobile phase by virtue of their differing interactions with a stationary phase. The stationary phase can be made of chiral material or the mobile phase can contain an additional chiral material to provoke the differing interactions;

[0118] xi) chiral gas chromatography—a technique whereby the racemate is volatilized and enantiomers are separated by virtue of their differing interactions in the gaseous mobile phase with a column containing a fixed non-racemic chiral adsorbent phase;

[0119] xii) extraction with chiral solvents—a technique whereby the enantiomers are separated by virtue of preferential dissolution of one enantiomer into a particular chiral solvent;

[0120] xiii) transport across chiral membranes—a technique whereby a racemate is placed in contact with a thin membrane barrier. The barrier typically separates two miscible fluids, one containing the racemate, and a driving force such as concentration or pressure differential causes preferential transport across the membrane barrier. Separation occurs as a result of the non-racemic chiral nature of the membrane which allows only one enantiomer of the racemate to pass through.

[0121] In some embodiments, tricitabine, tricitabine phosphate (TCN-P), tricitabine 5'-phosphate (TCN-P), or the DMF adduct of tricitabine (TCN-DMF) are provided. TCN can be synthesized by any technique known to one skilled in the art, for example, as described in Tetrahedron Letters, 49: 4757-4760 (1971). TCN-P can be prepared by any technique known to one skilled in the art, for example, as described in U.S. Pat. No. 4,123,524. The synthesis of TCN-DMF is described, for example, in INSERM, 81: 37-82 (1978). Other compounds related to TCN as described herein can be synthesized, for example, according to the methods disclosed in Gudmundsson et al., 2001, Nucleosides Nucleotides Nucleic Acids 20: 1823-1830; Porcari et al., 2000, J Med Chem 43: 2457-2463; Porcari et al., 1999, Nucleosides Nucleotides, 18:2475-2497; Porcari et al., 2000, J Med Chem, 43: 2438-2448; Porcari et al., 2003, Nucleosides Nucleotides Nucleic Acids, 22: 2171-2193; Porcari et al., 2004, Nucleosides Nucleotides Nucleic Acids 23: 31-39; Schweinsberg et al., 1981, Biochem Pharmacol 30: 2521-2526; Smith et al., 2004, Bioorg Med Chem Lett 14: 3517-3520; Townsend et al., 1986, Nucleic Acids Symp Ser 1986: 41-44; and/or, Wotring et al., 1986, Cancer Treat Rep 70: 491-7.

### 5.3. PHARMACEUTICALLY ACCEPTABLE SALTS, HYDRATES AND PRODRUGS

[0122] In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compound as a pharmaceutically acceptable salt may be appropriate. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. In particular, examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids, which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartrate, succinate, benzoate, ascorbate,  $\alpha$ -ketoglutarate, and  $\alpha$ -glycero-

phosphate. Suitable inorganic salts may also be formed, including, sulfate, nitrate, bicarbonate, and carbonate salts.

**[0123]** In another embodiment, the tricitabine compound is a solvate, for example, tricitabine phosphate monohydrate (TCN-PM).

**[0124]** Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example, reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

**[0125]** Any of the nucleotides described herein can be administered as a nucleotide prodrug to increase the activity, bioavailability, stability or otherwise alter the properties of the nucleoside. A number of nucleotide prodrug ligands are known. In general, alkylation, acylation or other lipophilic modification of the mono, di or triphosphate of the nucleoside will increase the stability of the nucleotide. Examples of substituent groups that can replace one or more hydrogens on the phosphate moiety are alkyl, aryl, steroids, carbohydrates, including sugars, 1,2-diacylglycerol and alcohols. Many are described in R. Jones and N. Bischoffberger, 1995, *Antiviral Research* 27: 1-17. Any of these can be used in combination with the disclosed nucleosides to achieve a desired effect.

**[0126]** In one embodiment, the tricitabine or a related compound is provided as 5'-hydroxyl lipophilic prodrug. Nonlimiting examples of U.S. patents that disclose suitable lipophilic substituents that can be covalently incorporated into the nucleoside, preferably at the 5'-OH position of the nucleoside or lipophilic preparations, include U.S. Pat. Nos. 5,149,794 (Yatvin, et al.); 5,194,654 (Hostetler, et al.); 5,223,263 (Hostetler, et al.); 5,256,641 (Yatvin, et al.); 5,411,947 (Hostetler, et al.); 5,463,092 (Hostetler, et al.); 5,543,389 (Yatvin, et al.); 5,543,390 (Yatvin, et al.); 5,543,391 (Yatvin, et al.); and 5,554,728 (Basava, et al.), all of which are incorporated herein by reference.

**[0127]** Foreign patent applications that disclose lipophilic substituents that can be attached to the tricitabine or a related compound s of the present invention, or lipophilic preparations, include WO 89/02733, WO 90/00555, WO 91/16920, WO 91/18914, WO 93/00910, WO 94/26273, WO/15132, EP 0 350 287, EP 93917054.4, and WO 91/19721.

**[0128]** Additional nonlimiting examples of derivatives of tricitabine or a related compound s are those that contain substituents as described in the following publications. These derivatized tricitabine or a related compound s can be used for the indications described in the text or otherwise as antiviral agents, including as anti-HIV or anti-HBV agents. Ho, 1973, *Cancer Res* 33: 2816-2820; Holy, 1993, *Isopolar phosphorous-modified nucleotide analogues*, in: De Clercq (ed.), *Advances in Antiviral Drug Design*, Vol. I, JAI Press, pp. 179-231; Hong et al., 1976, *Biochem Biophys Res Commun* 88: 1223-1229; Hong et al., 1980, *J Med Chem* 28: 171-177; Hostetler et al., 1990, *J Biol Chem* 266: 11714-11717; Hostetler et al., 1994, *Antiviral Res* 24: 59-67; Hostetler et al., 1994, *Antimicrobial Agents Chemother* 38: 2792-2797; Hunston et al., 1984, *J Med Chem* 27: 440-444; Ji et al 1990, *J Med Chem* 33: 2264-2270; Jones et al., 1984, *J Chem Soc Perkin Trans I*: 1471-1474; Juedka et al., 1974, *Coll Czech Chem Comm* 39: 363-968; Kataoka et al., 1989, *Nucleic Acids Res Sym Ser* 21: 1-2; Kataoka et al., 1991, *Heterocycles* 32: 1351-1356; Kinchington et al., 1992, *Antiviral Chem Chemother* 3: 107-112; Kodama et al., 1989, *Jpn J*

*Cancer Res* 80: 679-685; Korty et al., 1979, *Naunyn-Schmiedeberg's Arch Pharmacol* 310: 103-111; Kumar et al., 1990, *J Med Chem* 33: 2368-2375; LeBec et al., 1991, *Tetrahedron Lett* 32: 6553-6556; Lichtenstein et al., 1960, *J Biol Chem* 235: 457-465; Luethy et al., 1981, *Mitt Geg Lebensmittelunters Hyg* 72: 131-133 (Chem. Abstr. 95, 127093); McGuigan et al., 1989, *Nucleic Acids Res* 17: 6065-6075; McGuigan et al., 1990, *Antiviral Chem Chemother* 1: 107-113; McGuigan et al., 1990, *Antiviral Chem Chemother* 1: 355-360; McGuigan et al., 1990, *Antiviral Chem Chemother* 1: 25-33; McGuigan et al., 1991, *Antiviral Res* 15: 255-263; McGuigan et al., 1992, *Antiviral Res* 17: 311-321; McGuigan et al., 1993, *Antiviral Chem Chemother* 4: 97-101; McGuigan et al., 1993, *J Med Chem* 36: 1048-1052.

**[0129]** Alkyl hydrogen phosphonate derivatives of the anti-HIV agent AZT may be less toxic than the parent nucleoside analogue. *Antiviral Chem Chemother* 5: 271-277; Meyer et al., 1973, *Tetrahedron Lett* 269-272; Nagyvary et al., 1973, *Biochem Biophys Res Commun* 55: 1072-1077; Namane et al., 1992, *J Med Chem* 35: 3939-3044; Nargeot et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80: 2395-2399; Nelson et al., 1987, *J Am Chem Soc* 109: 4058-4064; Nerbonne et al., 1984, *Nature* 301: 74-76; Neumann et al., 1989, *J Am Chem Soc* 111: 4270-4277; Ohno et al., 1991, *Oncology* 48: 451-455. Palomino et al., 1989, *J Med Chem* 32: 622-625; Perkins et al., 1993, *Antiviral Res* 20(Supp. I): 84; Piantadosi et al., 1991, *J Med Chem* 34: 1408-1414; Pompon et al., 1994, *Antiviral Chem Chemother* 5: 91-98; Postemark, 1974, *Arm Rev Pharmacol* 14: 23-33; Prisbe et al., 1986, *J Med Chem* 29: 671-675; Pucchi et al., 1993, *Antiviral Res* 22: 155-174; Pugaeva et al., 1969, *Gig Trf Prof Zabol* 13: 47-48 (Chem. Abstr. 72, 212); Robins, 1984, *Pharm Res* 11-18; Rosowsky et al., 1982, *J Med Chem* 25: 171-178; Ross, 1961, *Biochem Pharm* 8: 235-240; Rya et al., 1982, *J Med Chem* 25: 1322-1329; Saffhill et al., 1986, *Chem Biol Interact* 57: 347-355; Saneyoshi et al., 1980, *Chem Pharm Bull* 28: 2915-2923; Sastry et al., 1992, *Mol Pharmacol* 41: 441-445; Shaw et al., 1994, 9th Annual AAPS Meeting, San Diego, Calif. (Abstract); Shuto et al., 1987, *Tetrahedron Lett* 28: 199-202; Shuto et al., 1988, *Chem Pharm Bull* 36: 209-217. One preferred phosphate prodrug group is the S-acyl-2-thioethyl group, also referred to as "SATE."

**[0130]** Additional examples of prodrugs that can be used are those described in the following patents and patent applications: U.S. Pat. Nos. 5,614,548, 5,512,671, 5,770,584, 5,962,437, 5,223,263, 5,817,638, 6,252,060, 6,448,392, 5,411,947, 5,744,592, 5,484,809, 5,827,831, 5,696,277, 6,022,029, 5,780,617, 5,194,654, 5,463,092, 5,744,461, 4,444,766, 4,562,179, 4,599,205, 4,493,832, 4,221,732, 5,116,992, 6,429,227, 5,149,794, 5,703,063, 5,888,990, 4810,697, 5,512,671, 6,030,960, 2004/0259845, 6,670,341, 2004/0161398, 2002/082242, 5,512,671, 2002/0082242, and or PCT Publication Nos WO 90/11079, WO 96/39197, and/or WO 93/08807.

#### 5.4. IN VIVO EFFICACY/DOSING REGIMENS

**[0131]** In another aspect of the present invention, dosing regimens are provided that limit the toxic side effects of TCN and related compounds. In one embodiment, such dosing regimens minimize the following toxic side effects, including, but not limited to, hepatotoxicity, thrombocytopenia, hyperglycemia, vomiting, hypocalcemia, anemia, hypoalbuminemia, myelosuppression, hypertriglyceridemia, hyperamylasemia, diarrhea, stomatitis and/or fever.



[0132] In another embodiment, the administration of TCN, TCN-P or related compounds and one or more platinum compounds provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

[0133] In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM or a related compound and one or more platinum compounds according to a dosing schedule that includes administering the tricitriline compound and/or the one or more platinum compounds one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

[0134] In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the tricitriline compound and/or one or more platinum compounds are not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m<sup>2</sup> of TCN, TCN-P, TCN-PM or a related compound can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 mg/m<sup>2</sup> of TCN, TCN-P, TCN-PM or a related compound can be administered according to a 42 day cycle. In another particular embodiment, about 1 to about 50 mg of a platinum compound is administered. In a particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90 or 100 mg/kg of a platinum compound can be administered according to a 42 day cycle.

[0135] In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m<sup>2</sup> or less of TCN, TCN-P or a related compound and less than about 30 mg of one or more platinum compounds one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m<sup>2</sup> of TCN, TCN-P or a related compound as disclosed herein can be administered one time per week. In another particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, or 100 mg/kg of a platinum compound can be administered one time per week.

[0136] In embodiments of the present invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the tricitriline compound and/or one or more platinum compounds via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined

in any manner disclosed herein to create a dosing cycle. The tricitriline compound and/or one or more platinum compounds can be repeatedly administered via a certain dosing cycle, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The tricitriline compound and/or one or more platinum compounds can be administered according to any combination disclosed herein, for example, the tricitriline compound and/or one or more platinum compounds can be administered once a week every three weeks for 3 cycles.

[0137] In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the tricitriline compound and/or one or more platinum compounds are not administered.

[0138] The TCN, TCN-P, TCM-PM and related compounds and one or more platinum compounds as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P or related compounds and one or more platinum compounds can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m<sup>2</sup> of a tricitriline compound disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m<sup>2</sup> of TCN, TCN-P, TCN-PM or a related compound disclosed herein can be administered to a subject. In certain embodiments, 10, 20, 50, 100, 150, 200, 250, 300, 350, or 400 mg of a platinum compound can be administered to a subject.

[0139] The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m<sup>2</sup> of TCN and related compounds and less than about 100 mg of a platinum compound either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m<sup>2</sup> of TCN or related compounds can be administered once a week concurrently with less than about 100 mg of a platinum compound. In another embodiment, less than 20 mg/m<sup>2</sup> of TCN or related compounds can be administered once a week and less than about 100 mg of a platinum compound can be administered the following week.

[0140] In further embodiments, 2 mg/m<sup>2</sup>, 5 mg/m<sup>2</sup>, 10 mg/m<sup>2</sup>, and/or 15 mg/m<sup>2</sup> of TCN or a related compound and less than about 30, 25, 20, 50, or 100 mg of a platinum compound can be administered to a subject. In another embodiment, less than 10 mg/m<sup>2</sup> of a tricitriline compound and less than about 100 mg of a platinum compound can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

#### 5.5. SCREENING OF PATIENT POPULATIONS

[0141] In another aspect of the present invention, methods are provided to identify cancers or tumors that are susceptible

to the toxic effects of triciribine (TCN) and related compounds. In one embodiment, methods are provided to treat a cancer or tumor in a mammal by (i) obtaining a biological sample from the tumor; (ii) determining whether the cancer or tumor overexpresses Akt kinase or hyperactivated and phosphorylated Akt kinase, and (iii) treating the cancer or tumor with triciribine or a related compound as described herein. In one embodiment, the biological sample can be a biopsy. In other embodiments, the biological sample can be fluid, cells and/or aspirates obtained from the tumor or cancer.

**[0142]** The biological sample can be obtained according to any technique known to one skilled in the art. In one embodiment, a biopsy can be conducted to obtain the biological sample. A biopsy is a procedure performed to remove tissue or cells from the body for examination. Some biopsies can be performed in a physician's office, while others need to be done in a hospital setting. In addition, some biopsies require use of an anesthetic to numb the area, while others do not require any sedation. In certain embodiments, an endoscopic biopsy can be performed. This type of biopsy is performed through a fiberoptic endoscope (a long, thin tube that has a close-focusing telescope on the end for viewing) through a natural body orifice (i.e., rectum) or a small incision (i.e., arthroscopy). The endoscope is used to view the organ in question for abnormal or suspicious areas, in order to obtain a small amount of tissue for study. Endoscopic procedures are named for the organ or body area to be visualized and/or treated. The physician can insert the endoscope into the gastrointestinal tract (alimentary tract endoscopy), bladder (cystoscopy), abdominal cavity (laparoscopy), joint cavity (arthroscopy), mid-portion of the chest (mediastinoscopy), or trachea and bronchial system (laryngoscopy and bronchoscopy).

**[0143]** In another embodiment, a bone marrow biopsy can be performed. This type of biopsy can be performed either from the sternum (breastbone) or the iliac crest hipbone (the bone area on either side of the pelvis on the lower back area). The skin is cleansed and a local anesthetic is given to numb the area. A long, rigid needle is inserted into the marrow, and cells are aspirated for study; this step is occasionally uncomfortable. A core biopsy (removing a small bone "chip" from the marrow) may follow the aspiration.

**[0144]** In a further embodiment, an excisional or incisional biopsy can be performed on the mammal. This type of biopsy is often used when a wider or deeper portion of the skin is needed. Using a scalpel (surgical knife), a full thickness of skin is removed for further examination, and the wound is sutured (sewed shut with surgical thread). When the entire tumor is removed, it is referred to as an excisional biopsy technique. If only a portion of the tumor is removed, it is referred to as an incisional biopsy technique. Excisional biopsy is often the method usually preferred, for example, when melanoma (a type of skin cancer) is suspected.

**[0145]** In still further embodiments, a fine needle aspiration (FNA) biopsy can be used. This type of biopsy involves using a thin needle to remove very small pieces from a tumor. Local anesthetic is sometimes used to numb the area, but the test rarely causes much discomfort and leaves no scar. FNA is not, for example, used for diagnosis of a suspicious mole, but may be used, for example, to biopsy large lymph nodes near a melanoma to see if the melanoma has metastasized (spread). A computed tomography scan (CT or CAT scan) can be used to guide a needle into a tumor in an internal organ such as the lung or liver.

**[0146]** In other embodiments, punch shave and/or skin biopsies can be conducted. Punch biopsies involve taking a deeper sample of skin with a biopsy instrument that removes a short cylinder, or "apple core," of tissue. After a local anesthetic is administered, the instrument is rotated on the surface of the skin until it cuts through all the layers, including the dermis, epidermis, and the most superficial parts of the subcutis (fat). A shave biopsy involves removing the top layers of skin by shaving it off. Shave biopsies are also performed with a local anesthetic. Skin biopsies involve removing a sample of skin for examination under the microscope to determine if, for example, melanoma is present. The biopsy is performed under local anesthesia.

**[0147]** In particular embodiment, methods are provided to determine whether the tumor overexpresses an Akt kinase. Akt kinase overexpression can refer to the phosphorylation state of the kinase. Hyperphosphorylation of Akt can be detected according to the methods described herein. In one embodiment, a tumor biopsy can be compared to a control tissue. The control tissue can be a normal tissue from the mammal in which the biopsy was obtained or a normal tissue from a healthy mammal. Akt kinase overexpression or hyperphosphorylation can be determined if the tumor biopsy contains greater amounts of Akt kinase and/or Akt kinase phosphorylation than the control tissue, such as, for example, at least approximately 1.5, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 5.5, 6, 7, 8, 9, or 10-fold greater amounts of Akt kinase than contained in the control tissue.

**[0148]** In one embodiment, the present invention provides a method to detect aberrant Akt kinase expression in a subject or in a biological sample from the subject by contacting cells, cell extracts, serum or other sample from the subjects or said biological sample with an immunointeractive molecule specific for an Akt kinase or antigenic portion thereof and screening for the level of immunointeractive molecule-Akt kinase complex formation, wherein an elevated presence of the complex relative to a normal cell is indicative of an aberrant cell that expresses or overexpresses Akt. In one example, cells or cell extracts can be screened immunologically for the presence of elevated levels of Akt kinase.

**[0149]** In an alternative embodiment, the aberrant expression of Akt in a cell is detected at the genetic level by screening for the level of expression of a gene encoding an Akt kinase wherein an elevated level of a transcriptional expression product (i.e., mRNA) compared to a normal cell is indicative of an aberrant cell. In certain embodiments, real-time PCR as well as other PCR procedures can be used to determine transcriptional activity. In one embodiment, mRNA can be obtained from cells of a subject or from a biological sample from a subject and cDNA optionally generated. The mRNA or cDNA can then be contacted with a genetic probe capable of hybridizing to and/or amplifying all or part of a nucleotide sequence encoding Akt kinase or its complementary nucleotide sequence and then the level of the mRNA or cDNA can be detected wherein the presence of elevated levels of the mRNA or cDNA compared to normal controls can be assessed.

**[0150]** Yet another embodiment of the present invention contemplates the use of an antibody, monoclonal or polyclonal, to Akt kinase in a quantitative or semi-quantitative diagnostic kit to determine relative levels of Akt kinase in suspected cancer cells from a patient, which can include all the reagents necessary to perform the assay. In one embodiment, a kit utilizing reagents and materials necessary to per-

form an ELISA assay is provided. Reagents can include, for example, washing buffer, antibody dilution buffer, blocking buffer, cell staining solution, developing solution, stop solution, anti-phospho-protein specific antibodies, anti-Pan protein specific antibodies, secondary antibodies, and distilled water. The kit can also include instructions for use and can optionally be automated or semi-automated or in a form which is compatible with automated machine or software. In one embodiment, a phosphor-ser-473 Akt antibody that detects the activated form of AKT (Akt phosphorylated at serine 474) can be utilized as the antibody in a diagnostic kit. See, for example, Yuan et al., 2000, *Oncogene* 19: 2324-2330.

### 5.6. AKT KINASES

[0151] Akt, also named PKB<sup>3</sup>, represents a subfamily of the serine/threonine kinase. Three members, AKT1, AKT2, and AKT3, have been identified in this subfamily. Akt is activated by extracellular stimuli in a PI3K-dependent manner (Datta et al., 1999, *Genes Dev* 13: 2905-2927). Full activation of Akt requires phosphorylation of Thr<sup>308</sup> in the activation loop and Ser<sup>473</sup> in the C-terminal activation domain. Akt is negatively regulated by PTEN tumor suppressor. Mutations in PTEN have been identified in various tumors, which lead to activation of Akt pathway (Datta et al., 1999, *Genes Dev* 13: 2905-2927). In addition, amplification, overexpression and/or activation of Akt have been detected in a number of human malignancies (Datta et al., 1999, *Genes Dev* 13: 2905-2927; Cheng et al., 2001, *AKT signal transduction pathway in oncogenesis* in Schwab D (ed.) *Encyclopedic Reference of cancer* Springer pp 35-7). Ectopic expression of Akt, especially constitutively active Akt, induces cell survival and malignant transformation whereas inhibition of Akt activity stimulates apoptosis in a range of mammalian cells (Datta et al., 1999, *Genes Dev* 13: 2905-2927; Cheng et al., 2001, *AKT signal transduction pathway in oncogenesis* in Schwab D (ed.) *Encyclopedic Reference of cancer* Springer pp 35-7; Sun et al., 2001, *Am J Path*, 159: 431-437; Cheng et al., 1997, *Oncogene* 14: 2793-2801). Further, activation of Akt has been shown to associate with tumor invasiveness and chemoresistance (West et al., 2002, *Drug Resist Vpdat* 5: 234-248).

[0152] Activation of the Akt pathway plays a pivotal role in malignant transformation and chemoresistance by inducing cell survival, growth, migration, and angiogenesis. The present invention provides methods to determine levels of Akt kinase overexpression and/or hyperactivated and phosphorylated Akt kinase.

[0153] The Akt kinase can be any known Akt family kinase, or kinase related thereto, including, but not limited to Akt 1, Akt 2, Akt 3. The mRNA and amino acid sequences of human Akt1, Akt2, and Akt 3 are illustrated in FIGS. 6a-c, 7a-d, and 8a-c, respectively.

### 5.7. DIAGNOSTIC ASSAYS

[0154] Immunological Assays

[0155] In one embodiment, a method is provided for detecting the aberrant expression of an Akt kinase in a cell in a mammal or in a biological sample from the mammal, by contacting cells, cell extracts or serum or other sample from the mammal or biological sample with an immunointeractive molecule specific for an Akt kinase or antigenic portion thereof and screening for the level of immunointeractive mol-

ecule-Akt kinase complex formations and determining whether an elevated presence of the complex relative to a normal cell is present.

[0156] The immunointeractive molecule can be a molecule having specificity and binding affinity for an Akt kinase or its antigenic parts or its homologs or derivatives thereof. In one embodiment, the immunointeractive molecule can be an immunoglobulin molecule. In other embodiments, the immunointeractive molecules can be an antibody fragments, single chain antibodies, and/or deimmunized molecules including humanized antibodies and T-cell associated antigen-binding molecules (TABMs). In one particular embodiment, the antibody can be a monoclonal antibody. In another particular embodiment, the antibody can be a polyclonal antibody. The immunointeractive molecule can exhibit specificity for an Akt kinase or more particularly an antigenic determinant or epitope on an Akt kinase. An antigenic determinant or epitope on an Akt kinase includes that part of the molecule to which an immune response is directed. The antigenic determinant or epitope can be a B-cell epitope or where appropriate a T-cell epitope. In one embodiment, the antibody is a phosphor-ser 473 Akt antibody.

[0157] One embodiment of the present invention provides a method for diagnosing the presence of cancer or cancer-like growth in a mammal, in which aberrant Akt activity is present, by contacting cells or cell extracts from the mammal or a biological sample from the subject with an Akt kinase-binding effective amount of an antibody having specificity for the Akt kinase or an antigenic determinant or epitope thereon and then quantitatively or qualitatively determining the level of an Akt kinase-antibody complex wherein the presence of elevated levels of said complex compared to a normal cell is determined.

[0158] Antibodies can be prepared by any of a number of means known to one skilled in the art. For example, for the detection of human Akt kinase, antibodies can be generally but not necessarily derived from non-human animals such as primates, livestock animals (e.g. sheep, cows, pigs, goats, horses), laboratory test animals (e.g. mice, rats, guinea pigs, rabbits) and/or companion animals (e.g. dogs, cats). Antibodies may also be recombinantly produced in prokaryotic or eukaryotic host cells. Generally, antibody based assays can be conducted in vitro on cell or tissue biopsies. However, if an antibody is suitably deimmunized or, in the case of human use, humanized, then the antibody can be labeled with, for example, a nuclear tag, administered to a patient and the site of nuclear label accumulation determined by radiological techniques. The Akt kinase antibody can be a cancer targeting agent. Accordingly, another embodiment of the present invention provides deimmunized forms of the antibodies for use in cancer imaging in human and non-human patients.

[0159] In general, for the generation of antibodies to an Akt kinase, the enzyme is required to be extracted from a biological sample whether this be from animal including human tissue or from cell culture if produced by recombinant means. The Akt kinase can be separated from the biological sample by any suitable means. For example, the separation may take advantage of any one or more of the Akt kinase's surface charge properties, size, density, biological activity and its affinity for another entity (e.g. another protein or chemical compound to which it binds or otherwise associates). Thus, for example, separation of the Akt kinase from the biological fluid can be achieved by any one or more of ultra-centrifugation, ion-exchange chromatography (e.g. anion exchange

chromatography, cation exchange chromatography), electrophoresis (e.g. polyacrylamide gel electrophoresis, isoelectric focussing), size separation (e.g., gel filtration, ultra-filtration) and affinity-mediated separation (e.g. immunoaffinity separation including, but not limited to, magnetic bead separation such as Dynabead (trademark) separation, immunochromatography, immuno-precipitation). The separation of Akt kinase from the biological fluid can preserve conformational epitopes present on the kinase and, thus, suitably avoids techniques that cause denaturation of the enzyme. In a further embodiment, the kinase can be separated from the biological fluid using any one or more of affinity separation, gel filtration and/or ultra-filtration.

**[0160]** Immunization and subsequent production of monoclonal antibodies can be carried out using standard protocols known in the art, such as, for example, described by Kohler and Milstein (Kohler and Milstein, 1975, *Nature* 256: 495-499; Kohler and Milstein, 1976, *Eur J Immunol* 6: 511-519, Coligan et al. *Current Protocols in Immunology*, John Wiley & Sons, Inc) or Toyama et al., *Monoclonal Antibody, Experiment Manual*, Kodansha Scientific). Essentially, an animal is immunized with an Akt kinase-containing biological fluid or fraction thereof or a recombinant form of Akt kinase by standard methods to produce antibody-producing cells, particularly antibody-producing somatic cells (e.g. B lymphocytes). These cells can then be removed from the immunized animal for immortalization. In certain embodiment, a fragment of an Akt kinase can be used to generate antibodies. The fragment can be associated with a carrier. The carrier can be any substance of typically high molecular weight to which a non- or poorly immunogenic substance (e.g. a hapten) is naturally or artificially linked to enhance its immunogenicity.

**[0161]** Immortalization of antibody-producing cells can be carried out using methods which are well-known in the art. For example, the immortalization may be achieved by the transformation method using Epstein-Barr virus (EBV) (Kozbor et al, 1986, *Methods in Enzymology* 121: 140). In another embodiment, antibody-producing cells are immortalized using the cell fusion method (described in Coligan et al., supra), which is widely employed for the production of monoclonal antibodies. In this method, somatic antibody-producing cells with the potential to produce antibodies, particularly B cells, are fused with a myeloma cell line. These somatic cells may be derived from the lymph nodes, spleens and peripheral blood of primed animals, preferably rodent animals such as mice and rats. In a particular embodiment, mice spleen cells can be used. In other embodiments, rat, rabbit, sheep or goat cells can also be used. Specialized myeloma cell lines have been developed from lymphocytic tumours for use in hybridoma-producing fusion procedures (Kohler and Milstein, 1976, supra; Shulman et al., 1978, *Nature* 276: 269-270; Volk et al., 1982, *J Virol* 42: 220-227). Many myeloma cell lines can also be used for the production of fused cell hybrids, including, e.g. P3.times.63-Ag8, P3.times.63-AG8.653, P3/NS1-Ag-4-1 (NS-1), Sp2/0-Ag14 and S194/5.XXO.Bu.1. The P3.times.63-Ag8 and NS-1 cell lines have been described by Kohler and Milstein (1976, supra). Shulman et al. (1978, supra) developed the Sp2/0-Ag14 myeloma line. The S194/5.XXO.Bu.1 line was reported by Trowbridge (1978, *J Exp Med* 148: 313-323). Methods for generating hybrids of antibody-producing spleen or lymph node cells and myeloma cells usually involve mixing somatic cells with myeloma cells in a 10:1 proportion (although the proportion may vary from about 20:1 to about

1:1), respectively, in the presence of an agent or agents (chemical, viral or electrical) that promotes the fusion of cell membranes. Fusion methods have been described (Kohler and Milstein, 1975, supra; Kohler and Milstein, 1976, supra; Gefer et al., 1977, *Somatic Cell Genet* 3: 231-236; Volk et al., 1982, supra). The fusion-promoting agents used by those investigators were Sendai virus and polyethylene glycol (PEG). In certain embodiments, means to select the fused cell hybrids from the remaining unfused cells, particularly the unfused myeloma cells, are provided. Generally, the selection of fused cell hybrids can be accomplished by culturing the cells in media that support the growth of hybridomas but prevent the growth of the unfused myeloma cells, which normally would go on dividing indefinitely. The somatic cells used in the fusion do not maintain long-term viability in *in vitro* culture and hence do not pose a problem. Several weeks are required to selectively culture the fused cell hybrids. Early in this time period, it is necessary to identify those hybrids which produce the desired antibody, so that they may subsequently be cloned and propagated. Generally, around 10% of the hybrids obtained produce the desired antibody, although a range of from about 1 to about 30% is not uncommon. The detection of antibody-producing hybrids can be achieved by any one of several standard assay methods, including enzyme-linked immunoassay and radioimmunoassay techniques as, for example, described in Kennet et al., *Monoclonal Antibodies and Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, New York pp 376-384) and by FACS analysis (O'Reilly et al. 1998, *Biotechniques* 25: 824-830).

**[0162]** Once the desired fused cell hybrids have been selected and cloned into individual antibody-producing cell lines, each cell line may be propagated in either of two standard ways. A suspension of the hybridoma cells can be injected into a histocompatible animal. The injected animal will then develop tumours that secrete the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can be tapped to provide monoclonal antibodies in high concentration. Alternatively, the individual cell lines may be propagated *in vitro* in laboratory culture vessels. The culture medium containing high concentrations of a single specific monoclonal antibody can be harvested by decantation, filtration or centrifugation, and subsequently purified.

**[0163]** The cell lines can then be tested for their specificity to detect the Akt kinase of interest by any suitable immunodetection means. For example, cell lines can be aliquoted into a number of wells and incubated and the supernatant from each well is analyzed by enzyme-linked immunosorbent assay (ELISA), indirect fluorescent antibody technique, or the like. The cell line(s) producing a monoclonal antibody capable of recognizing the target LIM kinase but which does not recognize non-target epitopes are identified and then directly cultured *in vitro* or injected into a histocompatible animal to form tumours and to produce, collect and purify the required antibodies.

**[0164]** The present invention provides, therefore, a method of detecting in a sample an Akt kinase or fragment, variant or derivative thereof comprising contacting the sample with an antibody or fragment or derivative thereof and detecting the level of a complex containing the antibody and Akt kinase or fragment, variant or derivative thereof compared to normal controls wherein elevated levels of Akt kinase is determined. Any suitable technique for determining formation of the com-

plex may be used. For example, an antibody according to the invention, having a reporter molecule associated therewith, may be utilized in immunoassays. Such immunoassays include but are not limited to radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs) immunochromatographic techniques (ICTs), and Western blotting which are well known to those of skill in the art. Immunoassays can also include competitive assays. The present invention encompasses qualitative and quantitative immunoassays.

**[0165]** Suitable immunoassay techniques are described, for example, in U.S. Pat. Nos. 4,016,043; 4,424,279; and 4,018,653. These include both single-site and two-site assays of the non-competitive types, as well as the traditional competitive binding assays. These assays also include direct binding of a labeled antigen-binding molecule to a target antigen.

**[0166]** The invention further provides methods for quantifying Akt protein expression and activation levels in cells or tissue samples obtained from an animal, such as a human cancer patient or an individual suspected of having cancer. In one embodiment, the invention provides methods for quantifying Akt protein expression or activation levels using an imaging system quantitatively. The imaging system can be used to receive, enhance, and process images of cells or tissue samples, that have been stained with AKT protein-specific stains, in order to determine the amount or activation level of AKT protein expressed in the cells or tissue samples from such an animal. In embodiments of the methods of the invention, a calibration curve of AKT1 and AKT2 protein expression can be generated for at least two cell lines expressing differing amounts of AKT protein. The calibration curve can then be used to quantitatively determine the amount of AKT protein that is expressed in a cell or tissue sample. Analogous calibration curves can be made for activated AKT proteins using reagents specific for the activation features. It can also be used to determine changes in amounts and activation state of AKT before and after clinical cancer treatment.

**[0167]** In one particular embodiment of the methods of the invention, AKT protein expression in a cell or tissue sample can be quantified using an enzyme-linked immunoabsorbent assay (ELISA) to determine the amount of AKT protein in a sample. Such methods are described, for example, in U.S. Patent Publication No. 2002/0015974.

**[0168]** In other embodiments enzyme immunoassays can be used to detect the Akt kinase. In such assays, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. The substrates to be used with the specific enzymes are generally chosen for the production of upon hydrolysis by the corresponding enzyme, a detectable colour change. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates. The enzyme-labeled antibody can be added to the first antibody-antigen complex, allowed to bind, and then the excess reagent washed away. A solution containing the appropriate substrate can then be added to the complex of antibody-antigen-antibody. The substrate can react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of antigen which was present in the sample. Alternatively, fluorescent compounds, such as fluorescein, rhodamine and the lanthanide, europium (EU), can be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labeled antibody

adsorbs the light energy, inducing a state to excitability in the molecule, followed by emission of the light at a characteristic colour visually detectable with a light microscope. The fluorescent-labeled antibody is allowed to bind to the first antibody-antigen complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to light of an appropriate wavelength. The fluorescence observed indicates the presence of the antigen of interest. Immunofluorometric assays (IFMA) are well established in the art and are particularly useful for the present method. However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules can also be employed.

**[0169]** In a particular embodiment, antibodies to Akt kinase can also be used in ELISA-mediated detection of Akt kinase especially in serum or other circulatory fluid. This can be accomplished by immobilizing anti-Akt kinase antibodies to a solid support and contacting these with a biological extract such as serum, blood, lymph or other bodily fluid, cell extract or cell biopsy. Labeled anti-Akt kinase antibodies can then be used to detect immobilized Akt kinase. This assay can be varied in any number of ways and all variations are encompassed by the present invention and known to one skilled in the art. This approach can enable rapid detection and quantitation of Akt kinase levels using, for example, a serum-based assay.

**[0170]** In one embodiment, an Akt Elisa assay kit may be used in the present invention. For example, a Cellular Activation of Signaling ELISA kit for Akt S473 from SuperArray Bioscience can be utilized in the present invention. In one embodiment, the antibody can be an anti-pan antibody that recognizes Akt S473. Elisa assay kit containing an anti-Akt antibody and additional reagents, including, but not limited to, washing buffer, antibody dilution buffer, blocking buffer, cell staining solution, developing solution, stop solution, secondary antibodies, and distilled water.

**[0171]** Nucleotide Detection

**[0172]** In another embodiment, a method to detect Akt kinases is provided by detecting the level of expression in a cell of a polynucleotide encoding an Akt kinase. Expression of the polynucleotide can be determined using any suitable technique known to one skilled in the art. In one embodiment, a labeled polynucleotide encoding an Akt kinase can be utilized as a probe in a Northern blot of an RNA extract obtained from the cell. In other embodiments, a nucleic acid extract from an animal can be utilized in concert with oligonucleotide primers corresponding to sense and antisense sequences of a polynucleotide encoding the kinase, or flanking sequences thereof, in a nucleic acid amplification reaction such as RT PCR. A variety of automated solid-phase detection techniques are also available to one skilled in the art, for example, as described by Fodor et al. (1991, Science 251: 767-777) and Kazal et al. (1996, Nature Medicine 2: 753-759).

**[0173]** In other embodiments, methods are provided to detect akt kinase encoding RNA transcripts. The RNA can be isolated from a cellular sample suspected of containing Akt kinase RNA, e.g. total RNA isolated from human cancer tissue. RNA can be isolated by methods known in the art, e.g. using TRIZOL reagent (GIBCO-BRL/Life Technologies, Gaithersburg, Md.). Oligo-dT, or random-sequence oligonucleotides, as well as sequence-specific oligonucleotides can be employed as a primer in a reverse transcriptase reaction to prepare first-strand cDNAs from the isolated RNA.

Resultant first-strand cDNAs can then be amplified with sequence-specific oligonucleotides in PCR reactions to yield an amplified product.

**[0174]** Polymerase chain reaction or "PCR" refers to a procedure or technique in which amounts of a preselected fragment of nucleic acid, RNA and/or DNA, are amplified as described, for example, in U.S. Pat. No. 4,683,195. Generally, sequence information from the ends of the region of interest or beyond is employed to design oligonucleotide primers. These primers will be identical or similar in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific RNA sequences and cDNA transcribed from total cellular RNA. See generally Mullis et al. (1987, *Quant Biol* 51: 263; Erlich (ed.), *PCR Technology*, Stockton Press, NY). Thus, amplification of specific nucleic acid sequences by PCR relies upon oligonucleotides or "primers" having conserved nucleotide sequences wherein the conserved sequences are deduced from alignments of related gene or protein sequences, e.g. a sequence comparison of mammalian Akt kinase genes. For example, one primer is prepared which is predicted to anneal to the antisense strand and another primer prepared which is predicted to anneal to the sense strand of a cDNA molecule which encodes a Akt kinase. To detect the amplified product, the reaction mixture is typically subjected to agarose gel electrophoresis or other convenient separation technique and the relative presence of the Akt kinase specific amplified DNA detected. For example, Akt kinase amplified DNA may be detected using Southern hybridization with a specific oligonucleotide probe or comparing its electrophoretic mobility with DNA standards of known molecular weight. Isolation, purification and characterization of the amplified Akt kinase DNA can be accomplished by excising or eluting the fragment from the gel (for example, see references Lawn et al., 1981, *Nucleic Acids Res* 2: 6103; Goeddel et al., 1980, *Nucleic Acids Res* 8: 4057), cloning the amplified product into a cloning site of a suitable vector, such as the pCRII vector (Invitrogen), sequencing the cloned insert and comparing the DNA sequence to the known sequence of LIM kinase. The relative amounts of LIM kinase mRNA and cDNA can then be determined.

**[0175]** In one embodiment, real-time PCR can be used to determine transcriptional levels of Akt nucleotides. Determination of transcriptional activity also includes a measure of potential translational activity based on available mRNA transcripts. Real-time PCR as well as other PCR procedures use a number of chemistries for detection of PCR product including the binding of DNA binding fluorophores, the 5' endonuclease, adjacent liner and hairpin oligoprobes and the self-fluorescing amplicons. These chemistries and real-time PCR in general are discussed, for example, in Mackay et al., 2002, *Nucleic Acids Res* 30: 1292-1305; Walker, 2001, *J Biochem Mol Toxicology* 15: 121-127; Lewis et al., 2001, *J Pathol* 195: 66-71

**[0176]** In an alternate embodiment, the aberrant expression of Akt can be identified by contacting a nucleotide sequences isolated from a biological sample with an oligonucleotide probe having a sequence complementary to an Akt sequences selected from the nucleotide sequences of FIG. 6a-c, 7a-d, or 8a-c, or fragment thereof, and then detecting the sequence by hybridizing the probe to the sequence, and comparing the results to a normal sample. The hybridization of the probe to the biological sample can be detected by labeling the probe using any detectable agent. The probe can be labeled for example, with a radioisotope, or with biotin, fluorescent dye,

electron-dense reagent, enzyme, hapten or protein for which antibodies are available. The detectable label can be assayed by any desired means, including spectroscopic, photochemical, biochemical, immunochemical, radioisotopic, or chemical means. The probe can also be detected using techniques such as an oligomer restriction technique, a dot blot assay, a reverse dot blot assay, a line probe assay, and a 5' nuclease assay. Alternatively, the probe can be detected using any of the generally applicable DNA array technologies, including macroarray, microarray and DNA microchip technologies. The oligonucleotide probe typically includes approximately at least 14, 15, 16, 18, 20, 25 or 28 nucleotides that hybridize to the nucleotides selected from FIGS. 6a-c, 7a-d, and 8a-c, or a fragment thereof. It is generally not preferred to use a probe that is greater than approximately 25 or 28 nucleotides in length. The oligonucleotide probe is designed to identify an Akt nucleotide sequence.

#### **[0177] Kinase Assays**

**[0178]** The activity of the Akt kinases can be measured using any suitable kinase assay known in the art. For example, and not by way of limitation, the methods described in Hogg et al (1994, *Oncogene* 9: 98-96), Mills et al (1992, *J Biol Chem* 267: 16000-006) and Tomizawa et al. (2001, *FEBS Lett* 492: 221-7), Schmandt et al. (1994, *J Immunol* 152: 96-105) can be used. Further serine, threonine and tyrosine kinase assays are described in Ausubel et al. (*Short Protocols in Molecular Biology*, 1999, unit 17.6).

**[0179]** Akt kinase assays can generally use an Akt polypeptide, a labeled donor substrate, and a receptor substrate that is either specific or non-specific for Akt. In such assays Akt transfers a labeled moiety from the donor substrate to the receptor substrate, and kinase activity is measured by the amount of labeled moiety transferred from the donor substrate to the receptor substrate. Akt polypeptide can be produced using various expression systems, can be purified from cells, can be in the form of a cleaved or uncleaved recombinant fusion protein and/or can have non-Akt polypeptide sequences, for example a His tag or .beta.-galactosidase at its N- or C-terminus. Akt activity can be assayed in cancerous cells lines if the cancerous cell lines are used as a source of the Akt to be assayed. Suitable donor substrates for Akt assays include any molecule that is susceptible to dephosphorylation by Akt., such as, for example include .gamma.-labeled ATP and ATP analogs, wherein the label is <sup>33</sup>P, <sup>32</sup>P, <sup>35</sup>S or any other radioactive isotope or a suitable fluorescent marker. Suitable recipient substrates for Akt assays include any polypeptide or other molecule that is susceptible to phosphorylation by Akt. Recipient substrates can be derived from fragments of in vivo targets of Akt. Recipient substrates fragments can be 8 to 50 amino acids in length, usually 10 to 30 amino acids and particularly of about 10, 12, 15, 18, 20 and 25 amino acids in length. Further recipient substrates can be determined empirically using a set of different polypeptides or other molecules. Targets of Recipient substrates for TTK can be capable of being purified from other components of the reaction once the reaction has been performed. This purification is usually done through a molecular interaction, where the recipient substrates is biotinylated and purified through its interaction with streptavidin, or a specific antibody is available that can specifically recognize the recipient substrates. The reaction can be performed in a variety of conditions, such as on a solid support, in a gel, in solution or in living cells. The choice of detection methods depends on type of label used for the donor molecule and may include, for example, measurement of

incorporated radiation or fluorescence by autoradiography, scintillation, scanning or fluorography.

## 6. METHODS OF TREATMENT

**[0180]** The compounds and pharmaceutical compositions provided herein can be used in the treatment of a condition including tumors, cancer, and other disorders associated with abnormal cell proliferation. In one embodiment, the compounds of the present invention can be used to treat a carcinoma, sarcoma, lymphoma, leukemia, and/or myeloma. In other embodiments of the present invention, the compounds disclosed herein can be used to treat solid tumors.

**[0181]** The compounds of the present invention can be used for the treatment of cancer, such as, but not limited to cancer of the following organs or tissues: breast, prostate, lung, bronchus, colon, urinary, bladder, non-Hodgkin lymphoma, melanoma, kidney, renal, pancreas, pharynx, thyroid, stomach, brain, multiple myeloma, esophagus, liver, intrahepatic bile duct, cervix, larynx, acute myeloid leukemia, chronic lymphatic leukemia, soft tissue, such as heart, Hodgkin lymphoma, testis, small intestine, chronic myeloid leukemia, acute lymphatic leukemia, anus, anal canal, anorectal, thyroid, vulva, gallbladder, pleura, eye, nose nasal cavity, middle ear, nasopharynx, ureter, peritoneum, omentum, mesentery, and gastrointestinal, high grade glioma, glioblastoma,

colon, rectal, pancreatic, gastric cancers, hepatocellular carcinoma; head and neck cancers, carcinomas; renal cell carcinoma; adenocarcinoma; sarcomas; hemangioendothelioma; lymphomas; leukemias, mycosis fungoides. In additional embodiments, the compounds of the present invention can be used to treat skin diseases including, but not limited to, the malignant diseases angiosarcoma, hemangioendothelioma, basal cell carcinoma, squamous cell carcinoma, malignant melanoma and Kaposi's sarcoma, and the non-malignant diseases or conditions such as psoriasis, lymphangiogenesis, hemangioma of childhood, Sturge-Weber syndrome, verruca vulgaris, neurofibromatosis, tuberous sclerosis, pyogenic granulomas, recessive dystrophic epidermolysis bullosa, venous ulcers, acne, rosacea, eczema, molluscum contagiosum, seborrheic keratosis, and actinic keratosis.

**[0182]** Compositions including the compounds of the invention can be used to treat these cancers and other cancers at any stage from the discovery of the cancer to advanced stages. In addition, compositions including compounds of the invention can be used in the treatment of the primary cancer and metastases thereof.

**[0183]** In other embodiments of the invention, the compounds described herein can be used for the treatment of cancer, including, but not limited to the cancers listed in Table 1 below.

TABLE 1

Types of Cancer	
Acute Lymphoblastic Leukemia, Adult	Hairy Cell Leukemia
Acute Lymphoblastic Leukemia, Childhood	Head and Neck Cancer
Acute Myeloid Leukemia, Adult	Hepatocellular (Liver) Cancer, Adult (Primary)
Acute Myeloid Leukemia, Childhood	Hepatocellular (Liver) Cancer, Childhood (Primary)
Adrenocortical Carcinoma	Hodgkin's Lymphoma, Adult
Adrenocortical Carcinoma, Childhood	Hodgkin's Lymphoma, Childhood
AIDS-Related Cancers	Hodgkin's Lymphoma During Pregnancy
AIDS-Related Lymphoma	Hypopharyngeal Cancer
Anal Cancer	Hypothalamic and Visual Pathway Glioma, Childhood
Astrocytoma, Childhood Cerebellar	Intraocular Melanoma
Astrocytoma, Childhood Cerebral	Islet Cell Carcinoma (Endocrine Pancreas)
Basal Cell Carcinoma	Kaposi's Sarcoma
Bile Duct Cancer, Extrahepatic	Kidney (Renal Cell) Cancer
Bladder Cancer	Kidney Cancer, Childhood
Bladder Cancer, Childhood	Laryngeal Cancer
Bone Cancer, Osteosarcoma/Malignant	Laryngeal Cancer, Childhood
Fibrous Histiocytoma	Leukemia, Acute Lymphoblastic, Adult
Brain Stem Glioma, Childhood	Leukemia, Acute Lymphoblastic, Childhood
Brain Tumor, Adult	Leukemia, Acute Myeloid, Adult
Brain Tumor, Brain Stem Glioma, Childhood	Leukemia, Acute Myeloid, Childhood
Brain Tumor, Cerebellar Astrocytoma, Childhood	Leukemia, Chronic Lymphocytic
Brain Tumor, Cerebral	Leukemia, Chronic Myelogenous
Astrocytoma/Malignant Glioma, Childhood	Leukemia, B Cell
Brain Tumor, Ependymoma, Childhood	Lip and Oral Cavity Cancer
Brain Tumor, Medulloblastoma, Childhood	Liver Cancer, Adult (Primary)
Brain Tumor, Supratentorial Primitive Neuroectodermal Tumors, Childhood	Liver Cancer, Childhood (Primary)
Brain Tumor, Visual Pathway and Hypothalamic Glioma, Childhood	Lung Cancer, Non-Small Cell
Brain Tumor, Childhood	Lung Cancer, Small Cell
Breast Cancer	Lymphoma, AIDS-Related
Breast Cancer, Childhood	Lymphoma, Burkitt's
Breast Cancer, Male	Lymphoma, Cutaneous T-Cell, see Mycosis Fungoides and Sézary Syndrome
Bronchial Adenomas/Carcinoids, Childhood	Lymphoma, Hodgkin's, Adult
Burkitt's Lymphoma	Lymphoma, Hodgkin's, Childhood
	Lymphoma, Hodgkin's During Pregnancy
	Lymphoma, Non-Hodgkin's, Adult
	Lymphoma, Non-Hodgkin's, Childhood
	Lymphoma, Non-Hodgkin's During Pregnancy
	Lymphoma, Primary Central Nervous System

TABLE 1-continued

Types of Cancer	
Carcinoid Tumor, Childhood	Macroglobulinemia, Waldenström's
Carcinoid Tumor, Gastrointestinal	Malignant Fibrous Histiocytoma of
Carcinoma of Unknown Primary	Bone/Osteosarcoma
Central Nervous System Lymphoma, Primary	Medulloblastoma, Childhood
Cerebellar Astrocytoma, Childhood	Melanoma
Cerebral Astrocytoma/Malignant Glioma, Childhood	Melanoma, Intraocular (Eye)
Cervical Cancer	Merkel Cell Carcinoma
Childhood Cancers	Mesothelioma, Adult Malignant
Chronic Lymphocytic Leukemia	Mesothelioma, Childhood
Chronic Myelogenous Leukemia	Metastatic Squamous Neck Cancer with Occult Primary
Chronic Myeloproliferative Disorders	Multiple Endocrine Neoplasia Syndrome, Childhood
Colon Cancer	Multiple Myeloma/Plasma Cell Neoplasm
Colorectal Cancer, Childhood	Mycosis Fungoides
Cutaneous T-Cell Lymphoma, see	Myelodysplastic Syndromes
Mycosis Fungoides and Sézary Syndrome	Myelodysplastic/Myeloproliferative Diseases
Endometrial Cancer	Myelogenous Leukemia, Chronic
Ependymoma, Childhood	Myeloid Leukemia, Adult Acute
Esophageal Cancer	Myeloid Leukemia, Childhood Acute
Esophageal Cancer, Childhood	Myeloma, Multiple
Ewing's Family of Tumors	Myeloproliferative Disorders, Chronic
Extracranial Germ Cell Tumor, Childhood	Nasal Cavity and Paranasal Sinus Cancer
Extragenital Germ Cell Tumor	Nasopharyngeal Cancer
Extrahepatic Bile Duct Cancer	Nasopharyngeal Cancer, Childhood
Eye Cancer, Intraocular Melanoma	Neuroblastoma
Eye Cancer, Retinoblastoma	Non-Hodgkin's Lymphoma, Adult
Gallbladder Cancer	Non-Hodgkin's Lymphoma, Childhood
Gastric (Stomach) Cancer	Non-Hodgkin's Lymphoma During Pregnancy
Gastric (Stomach) Cancer, Childhood	Non-Small Cell Lung Cancer
Gastrointestinal Carcinoid Tumor	Oral Cancer, Childhood
Germ Cell Tumor, Extracranial, Childhood	Oral Cavity Cancer, Lip and Oropharyngeal Cancer
Germ Cell Tumor, Extragenital	Osteosarcoma/Malignant Fibrous Histiocytoma of Bone
Germ Cell Tumor, Ovarian	Ovarian Cancer, Childhood
Gestational Trophoblastic Tumor	Ovarian Epithelial Cancer
Glioma, Adult	Ovarian Germ Cell Tumor
Glioma, Childhood Brain Stem	Ovarian Low Malignant Potential Tumor
Glioma, Childhood Cerebral	Pancreatic Cancer
Astrocytoma	Pancreatic Cancer, Childhood
Glioma, Childhood Visual Pathway and Hypothalamic	Pancreatic Cancer, Islet Cell
Skin Cancer (Melanoma)	Paranasal Sinus and Nasal Cavity Cancer
Skin Carcinoma, Merkel Cell	Parathyroid Cancer
Small Cell Lung Cancer	Penile Cancer
Small Intestine Cancer	Pheochromocytoma
Soft Tissue Sarcoma, Adult	Pineoblastoma and Supratentorial Primitive Neuroectodermal Tumors, Childhood
Soft Tissue Sarcoma, Childhood	Pituitary Tumor
Squamous Cell Carcinoma, see Skin Cancer (non-Melanoma)	Plasma Cell Neoplasm/Multiple Myeloma
Squamous Neck Cancer with Occult Primary, Metastatic	Pleuropulmonary Blastoma
Stomach (Gastric) Cancer	Pregnancy and Breast Cancer
Stomach (Gastric) Cancer, Childhood	Pregnancy and Hodgkin's Lymphoma
Supratentorial Primitive Neuroectodermal Tumors, Childhood	Pregnancy and Non-Hodgkin's Lymphoma
T-Cell Lymphoma, Cutaneous, see Mycosis Fungoides and Sézary Syndrome	Primary Central Nervous System Lymphoma
Testicular Cancer	Prostate Cancer
Thymoma, Childhood	Rectal Cancer
Thymoma and Thymic Carcinoma	Renal Cell (Kidney) Cancer
Thyroid Cancer	Renal Cell (Kidney) Cancer, Childhood
Thyroid Cancer, Childhood	Renal Pelvis and Ureter, Transitional Cell Cancer
Transitional Cell Cancer of the Renal Pelvis and Ureter	Retinoblastoma
Trophoblastic Tumor, Gestational	Rhabdomyosarcoma, Childhood
Unknown Primary Site, Carcinoma of, Adult	Salivary Gland Cancer
Unknown Primary Site, Cancer of, Childhood	Salivary Gland Cancer, Childhood
Unusual Cancers of Childhood	Sarcoma, Ewing's Family of Tumors
	Sarcoma, Kaposi's
	Sarcoma, Soft Tissue, Adult
	Sarcoma, Soft Tissue, Childhood
	Sarcoma, Uterine
	Sézary Syndrome
	Skin Cancer (non-Melanoma)
	Skin Cancer, Childhood



TABLE 1-continued

Types of Cancer
Ureter and Renal Pelvis, Transitional Cell Cancer
Urethral Cancer
Uterine Cancer, Endometrial
Uterine Sarcoma
Vaginal Cancer
Visual Pathway and Hypothalamic
Glioma, Childhood
Vulvar Cancer
Waldenström's Macroglobulinemia
Wilms' Tumor

**[0184]** In further embodiments of the present invention, the compounds disclosed herein can be used in the treatment of angiogenesis-related diseases.

**[0185]** Antiangiogenic small molecules include thalidomide, which acts in part by inhibiting NFκB, 2-methoxyestradiol, which influences microtubule activation and hypoxia inducing factor (HIF1a) activation, cyclo-oxygenase 2 (COX2) inhibitors, and low doses of conventional chemotherapeutic agents, including cyclophosphamide, and vinca alkaloids (vincristine, vinblastine) (D'Amato et al., 1994, Proc. Natl. Acad. Sci. U.S.A. 91: 3964-3968; D'Amato et al., 1994, Proc. Natl. Acad. Sci. U.S.A. 91: 4082-4085). In addition, certain tyrosine kinase inhibitors indirectly decrease angiogenesis by decreasing production of VEGF and other proangiogenic factors by tumor and stromal cells. These drugs include Herceptin, imatinib (Gleevec), and Iressa (Bergers et al., 2003, J Clin Invest 111: 1287-1295; Ciardiello et al., 2001, Clin Cancer Res 7: 1459-1465; Plum et al., 2003, Clin Cancer Res 9: 4619-4626).

**[0186]** Recently, angiogenesis inhibitors have moved from animal models to human patients. Angiogenesis inhibitors represent a promising treatment for a variety of cancers. Recently, Avastin a high affinity antibody against vascular endothelial growth factor (VEGF), has been shown to prolong life as a single agent in advanced renal cell carcinoma and prolong life in combination with chemotherapy in advanced colon cancer (Yang et al., 2003, New Eng J Med 349: 427-434; Kabbavar et al., 2003, J Clin Oncol 21: 60-65).

**[0187]** Angiogenesis-related diseases include, but are not limited to, inflammatory, autoimmune, and infectious diseases; angiogenesis-dependent cancer, including, for example, solid tumors, blood born tumors such as leukemias, and tumor metastases; benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; rheumatoid arthritis; psoriasis; eczema; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; and wound granulation. In addition, compositions of this invention can be used to treat diseases such as, but not limited to, intestinal adhesions, atherosclerosis, scleroderma, warts, and hypertrophic scars (i.e., keloids). Compositions of this invention can also be used in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (Rochele minallia quintosa), ulcers (Helobacter pylori), tuberculosis, and leprosy.

### 6.1. Treatment of Drug Resistant Tumors or Cancers

**[0188]** The invention provides compounds that can be used to treat drug resistant cancer, including the embodiments of cancers and the tricitiribine compound and/or the one or more platinum compounds disclosed herein.

**[0189]** Multidrug resistance (MDR) occurs in human cancers and can be a significant obstacle to the success of chemotherapy. Multidrug resistance is a phenomenon whereby tumor cells in vitro that have been exposed to one cytotoxic agent develop cross-resistance to a range of structurally and functionally unrelated compounds. In addition, MDR can occur intrinsically in some cancers without previous exposure to chemotherapy agents. Thus, in one embodiment, the present invention provides methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer, by administration of TCN, TCN-P or a related compound and one or more one or more platinum compounds as disclosed herein. In certain embodiments, TCN, TCN-P and related compounds and one or more one or more platinum compounds can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

**[0190]** In one embodiment, TCN, TCN-P or a related compound and one or more one or more platinum compounds as disclosed herein can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

### 6.2. Combination Therapy

**[0191]** In one embodiment, the tricitiribine compounds and one or more platinum compounds of the invention can be administered together with other cytotoxic agents. In another embodiment, the tricitiribine compounds and one or more platinum compounds and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

**[0192]** In another embodiment of the present invention, the tricitiribine compounds and one or more platinum compounds and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

**[0193]** In one embodiment, the tricitiribine compounds and one or more platinum compounds disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and

administered together with other cytotoxic agents. In another embodiment, the tricycline compounds and one or more platinum compounds and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopryl, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angi-inhibins, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimetabolic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anticycline drugs adriamycin, doxorubicin, pharomurubicin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyprotrone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carboplatin, oxaliplatin, paraplalin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed below in Table 2.

TABLE 2

Chemotherapeutic Agents	
13-cis-Retinoic Acid	Neosar
2-Amino-6-Mercaptopurine	Neulasta
2-CdA	Neumega
2-Chlorodeoxyadenosine	Neupogen
5-fluorouracil	Nilandron
5-FU	Nilutamide
6-TG	Nitrogen Mustard
6-Thioguanine	Novaldex
6-Mercaptopurine	Novantrone
6-MP	Octreotide
Accutane	Octreotide acetate
Actinomycin-D	Oncospar
Adriamycin	Oncovin
Adrucil	Ontak

TABLE 2-continued

Chemotherapeutic Agents	
Agrylin	Onxal
Ala-Cort	Oprevelkin
Aldesleukin	Orapred
Alemtuzumab	Orasone
Alitretinoin	Oxaliplatin
Alkaban-AQ	Paclitaxel
Alkeran	Pamidronate
All-transretinoic acid	Panretin
Alpha interferon	Paraplatin
Altretamine	Pediapred
Amethopterin	PEG Interferon
Amifostine	Pegaspargase
Aminoglutethimide	Pegfilgrastim
Anagrelide	PEG-INTRON
Anandron	PEG-L-asparaginase
Anastrozole	Phenylalanine Mustard
Arabinosylcytosine	Platinol
Ara-C	Platinol-AQ
Aranesp	Prednisolone
Aredia	Prednisone
Arimidex	Prelone
Aromasin	Procarbazine
Arsenic trioxide	PROCRIT
Asparaginase	Proleukin
ATRA	Prolifeprospan 20 with Camustine implant
Avastin	Purinethol
BCG	Raloxifene
BCNU	Rheumatrex
Bevacizumab	Rituxan
Bexarotene	Rituximab
Bicalutamide	Roveron-A (interferon alfa-2a)
BiCNU	Rubex
Blenoxane	Rubidomycin hydrochloride
Bleomycin	Sandostatin
Bortezomib	Sandostatin LAR
Busulfan	Sargramostim
Busulfex	Solu-Cortef
C225	Solu-Medrol
Calcium Leucovorin	STI-571
Campath	Streptozocin
Camptosar	Tamoxifen
Camptothecin-11	Targretin
Capecitabine	Taxol
Carac	Taxotere
Carboplatin	Temodar
Carmustine	Temozolomide
Carmustine wafer	Teniposide
Casodex	TESPA
CCNU	Thalidomide
CDDP	Thalomid
CeeNU	TheraCys
Cerubidine	Thioguanine
cetuximab	Thioguanine Tabloid
Chlorambucil	Thiophosphamide
Cisplatin	Thiopex
Citrovorum Factor	Thiotepa
Cladribine	TICE
Cortisone	Toposar
Cosmegen	Topotecan
CPT-11	Toremifene
Cyclophosphamide	A platinum compound
Cytadren	Tretinoin
Cytarabine	Trexall
Cytarabine liposomal	Trisenox
Cytosar-U	TSPA
Cytoxan	VCR
Dacarbazine	Velban
Dactinomycin	Velcade
Darbepoetin alfa	VePesid
Daunomycin	Vesanoid
Daunorubicin	Viadur
Daunorubicin hydrochloride	Vinblastine
Daunorubicin liposomal	Vinblastine Sulfate
DaunoXome	Vincasar Pfs

TABLE 2-continued

Chemotherapeutic Agents	
Decadron	Vincristine
Delta-Cortef	Vinorelbine
Deltasone	Vinorelbine tartrate
Denileukin diftitox	VLB
DepoCyt	VP-16
Dexamethasone	Vumon
Dexamethasone acetate	Xeloda
dexamethasone sodium phosphate	Zanosar
Dexasone	Zevalin
Dexrazoxane	Zinecard
DHAD	Zoladex
DIC	Zoledronic acid
Diodex	Zometa
Docetaxel	Gliadel wafer
Doxil	Glivec
Doxorubicin	GM-CSF
Doxorubicin liposomal	Goserelin
	granulocyte - colony stimulating factor
Droxia	Granulocyte macrophage colony stimulating factor
DTIC	Halotestin
DTIC-Dome	Herceptin
Duralone	Hexadrol
Efudex	Hexalen
Eligard	Hexamethylmelamine
Ellence	HMM
Eloxatin	Hycamtin
Elspar	Hydrea
Emcyt	Hydrocort Acetate
Epirubicin	Hydrocortisone
Epoetin alfa	Hydrocortisone sodium phosphate
Erbix	Hydrocortisone sodium succinate
Erwinia L-asparaginase	Hydrocortone phosphate
Estramustine	Hydroxyurea
Ethylol	Ibritumomab
Etopophos	Ibritumomab Tiuxetan
Etoposide	Idamycin
Etoposide phosphate	Idarubicin
Eulexin	Ifex
Evista	IFN-alpha
Exemestane	Ifosfamide
Fareston	IL-2
Faslodex	IL-11
Femara	Imatinib mesylate
Filgrastim	Imidazole Carboxamide
Floxuridine	Interferon alfa
Fludara	Interferon Alfa-2b (PEG conjugate)
Fludarabine	Interleukin-2
Fluoroplex	Interleukin-11
Fluorouracil	Intron A (interferon alfa-2b)
Fluorouracil (cream)	Leucovorin
Fluoxymesterone	Leukeran
Flutamide	Leukine
Folinic Acid	Leuprolide
FUDR	Leurocristine
Fulvestrant	Leustatin
G-CSF	Liposomal Ara-C
Gefitinib	Liquid Pred
Gemcitabine	Lomustine
Gemtuzumab ozogamicin	L-PAM
Gemzar	L-Sarcosylsin
Gleevec	Meticorten
Lupron	Mitomycin
Lupron Depot	Mitomycin-C
Matulane	Mitoxantrone
Maxidex	M-Prednisol
Mechlorethamine	MTC
Mechlorethamine	MTX
Hydrochlorine	Mustargen
Medralone	Mustine
Medrol	Mutamycin
Megace	Myleran
Megestrol	Iressa

TABLE 2-continued

Chemotherapeutic Agents	
Megestrol Acetate	Irinotecan
Melphalan	Isotretinoin
Mercaptopurine	Kidrolase
Mesna	Lanacort
Mesnex	L-asparaginase
Methotrexate	LCR
Methotrexate Sodium	
Methylprednisolone	
Mylocel	
Letrozole	

**[0194]** In certain embodiments, interferons (IFNs) can be used in combinations with the compounds of the present invention. Suitable interferons include: interferon alpha-2a, interferon alpha-2b, pegylated interferon alpha, including interferon alpha-2a and interferon alpha 2b, interferon beta, interferon gamma, interferon tau, interferon omega, INFERGEN (interferon alphacon-1) by InterMune, OMNIFERON (natural interferon) by Viragen, ALBUFERON by Human Genome Sciences, REBIF (interferon beta-1a) by Ares-Serono, Omega Interferon by BioMedicine, Oral Interferon Alpha by Amarillo Biosciences, and interferon gamma, interferon tau, and/or interferon gamma-1 $\beta$  by InterMune.

**[0195]** In one embodiment TCN, TCN-P or a related compound and one or more platinum compounds as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 3, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextinidipine, valsopodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

## 7. PHARMACEUTICAL COMPOSITIONS

**[0196]** The compositions including tricycline compounds and one or more platinum compounds can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The tricycline compounds and in combination with one or more platinum compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with one or more platinum compounds.

**[0197]** Compositions including the tricycline compounds and one or more platinum compounds may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

[0198] The compositions may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association one or more compositions of the present invention and one or more pharmaceutical carriers or excipients.

[0199] The tricitiribine compounds and one or more platinum compounds and compositions thereof can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the tricitiribine compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel Introduction to Pharmaceutical Dosage Forms, Fourth Ed., 1985, p. 126).

[0200] In the compositions, effective concentrations of one or more compounds or pharmaceutically acceptable derivatives thereof may be mixed with one or more suitable pharmaceutical carriers. The compounds of the invention may be derivatized as the corresponding salts, esters, enol ethers or esters, acetals, ketals, orthoesters, hemiacetals, hemiketals, acids, bases, solvates, hydrates or prodrugs prior to formulation. The concentrations of the compounds in the compositions are effective for delivery of an amount, upon administration, that treats, prevents, or ameliorates one or more of the symptoms of the target disease or disorder. In one embodiment, the compositions are formulated for single dosage administration. To formulate a composition, the weight fraction of compound is dissolved, suspended, dispersed or otherwise mixed in a selected carrier at an effective concentration such that the treated condition is relieved, prevented, or one or more symptoms are ameliorated.

[0201] Compositions suitable for oral administration may be presented as discrete units such as, but not limited to, tablets, caplets, pills or dragees capsules, or cachets, each containing a predetermined amount of one or more of the compositions; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion or as a bolus, etc.

[0202] Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing a tricitiribine compound and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, solubilizing agents, pH buffering agents, preservatives, flavoring agents, and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa.

[0203] Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or

more tricitiribine compounds and one or more platinum compounds of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

[0204] The tablets, pills, capsules, troches and the like can contain one or more of the following ingredients, or compounds of a similar nature: a binder; a lubricant; a diluent; a glidant; a disintegrating agent; a coloring agent; a sweetening agent; a flavoring agent; a wetting agent; an emetic coating; and a film coating. Examples of binders include microcrystalline cellulose, gum tragacanth, glucose solution, acacia mucilage, gelatin solution, molasses, polyvinylpyrrolidone, povidone, crospovidones, sucrose and starch paste. Lubricants include talc, starch, magnesium or calcium stearate, *lycopodium* and stearic acid. Diluents include, for example, lactose, sucrose, starch, kaolin, salt, mannitol and dicalcium phosphate. Glidants include, but are not limited to, colloidal silicon dioxide. Disintegrating agents include crosscarmellose sodium, sodium starch glycolate, alginic acid, corn starch, potato starch, bentonite, methylcellulose, agar and carboxymethylcellulose. Coloring agents include, for example, any of the approved certified water soluble FD and C dyes, mixtures thereof; and water insoluble FD and C dyes suspended on alumina hydrate. Sweetening agents include sucrose, lactose, mannitol and artificial sweetening agents such as saccharin, and any number of spray dried flavors. Flavoring agents include natural flavors extracted from plants such as fruits and synthetic blends of compounds which produce a pleasant sensation, such as, but not limited to peppermint and methyl salicylate. Wetting agents include propylene glycol monostearate, sorbitan monoolate, diethylene glycol monolaurate and polyoxyethylene laural ether. Emetic-coatings include fatty acids, fats, waxes, shellac, ammoniated shellac and cellulose acetate phthalates. Film coatings include hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate.

[0205] Compositions suitable for topical administration to the skin may be presented as ointments, creams, gels, and pastes, having one or more of the compositions administered in a pharmaceutical acceptable carrier.

[0206] Compositions for rectal administration may be presented as a suppository with a suitable base including, for example, cocoa butter or a salicylate.

[0207] Compositions suitable for nasal administration, when the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to 500 microns which is administered in the manner in which snuff is taken, (i.e., by rapid inhalation through the nasal passage from a container of the powder held close up to the nose). When the carrier is a liquid (for example, a nasal spray or as nasal drops), one or more of the compositions can be admixed in an aqueous or oily solution, and inhaled or sprayed into the nasal passage.

[0208] Compositions suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing one or more of the compositions and appropriate carriers.

[0209] Compositions suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous ster-

ile suspensions which may include suspending agents and thickening agents. The compositions may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules, and tablets of the kind previously described above.

**[0210]** Pharmaceutical organic or inorganic solid or liquid carrier media suitable for enteral or parenteral administration can be used to fabricate the compositions. Gelatin, lactose, starch, magnesium stearate, talc, vegetable and animal fats and oils, gum, polyalkylene glycol, water, or other known carriers may all be suitable as carrier media.

**[0211]** Compositions including tricitiribine compounds and one or more platinum compounds may be used in combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

**[0212]** Additionally, the compositions including tricitiribine compounds and one or more platinum compounds may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

**[0213]** It will be understood, however, that the total daily usage of the compositions will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular host will depend upon a variety of factors, including for example, the disorder being treated and the severity of the disorder; activity of the specific composition employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration; route of administration; rate of excretion of the specific compound employed; the duration of the treatment; the tricitiribine compound and/or the one or more platinum compounds used in combination or coincidental with the specific composition employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the composition at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

**[0214]** Compositions including tricitiribine compounds and one or more platinum compounds are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

**[0215]** Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of the administered ingredient. For example, approximately 1-5 mg per day of a compound disclosed herein can reduce the volume of a solid tumor in mice.

**[0216]** The dosage will depend on host factors such as weight, age, surface area, metabolism, tissue distribution, absorption rate and excretion rate. In one embodiment, approximately 0.5 to 7 grams per day of a tricitiribine compound disclosed herein may be administered to humans. Optionally, approximately 1 to 4 grams per day of the compound can be administered to humans. In certain embodiments 0.001-5 mg/day is administered to a human. The therapeutically effective dose level will depend on many factors as noted above. In addition, it is well within the skill of the art to start doses of the composition at relatively low levels, and increase the dosage until the desired effect is achieved.

**[0217]** Compositions including tricitiribine compounds and one or more platinum compounds may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

**[0218]** The tricitiribine compounds and one or more platinum compounds may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

**[0219]** The tricitiribine compounds and one or more platinum compounds may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microfine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

**[0220]** Compositions including the tricitiribine compounds and one or more platinum compounds may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with one or more compositions of

the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

#### 7.1. Additional Embodiments

**[0221]** The pharmaceutical compositions including tricitiribine compounds and one or more platinum compounds can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, Remington's Pharmaceutical Sciences (Mack Publishing Company, Easton, Pa.) describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

**[0222]** The methods of the present invention, for example, for inhibiting the growth of a cancerous cell, can be advantageously combined with at least one additional therapeutic method, including but not limited to chemotherapy, radiation therapy, therapy that selectively inhibits Ras oncogenic signaling, or any other therapy known to those of skill in the art of the treatment and management of cancer, such as administration of an anti-cancer agent.

**[0223]** Administration of API-2 (tricitiribine) as a salt may be carried out. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, alpha-ketoglutarate, and alpha-glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

**[0224]** Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

**[0225]** The tricitiribine compounds and one or more platinum compounds can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

**[0226]** Thus, the tricitiribine compounds and one or more platinum compounds of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

**[0227]** The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the compounds of the invention, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the compounds of the invention may be incorporated into sustained-release preparations and devices.

**[0228]** The tricitiribine compounds and one or more platinum compounds may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

**[0229]** The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders including the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium including, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance

of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

**[0230]** Sterile injectable solutions are prepared by incorporating compounds of the invention in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

**[0231]** For topical administration, the tricyriline compounds and one or more platinum compounds may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

**[0232]** Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the compounds of the invention can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

**[0233]** Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user. Examples of useful dermatological compositions which can be used to deliver the compounds of the invention to the skin are disclosed in Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat. No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Woltzman (U.S. Pat. No. 4,820,508).

**[0234]** Useful dosages of the pharmaceutical compositions of the present invention can be determined by comparing their in vitro activity, and in vivo activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949.

**[0235]** In one non-limiting embodiment, the concentration of the active agent in a liquid composition, such as a lotion, can be from about 0.1-25 wt.-%, or from about 0.5-10 wt.-%. In one embodiment, the concentration in a semi-solid or solid composition such as a gel or a powder can be about 0.1-5 wt.-%, preferably about 0.5-2.5 wt.-%. In one embodiment, single dosages for injection, infusion or ingestion will generally vary between 5-1500 mg, and may be administered, i.e., 1-3 times daily, to yield levels of about 0.1-50 mg/kg, for adults. A non-limiting dosage of the present invention can be

between 7.5 to 45 mg per clay, administered orally, with appropriate adjustment for the body weight of an individual.

**[0236]** Accordingly, the present invention includes a pharmaceutical composition including tricyriline compounds and one or more platinum compounds or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of tricyriline compounds and one or more platinum compounds or a pharmaceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

**[0237]** A suitable dose is that which will result in a concentration of the tricyriline compounds and one or more platinum compounds in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

**[0238]** Mammalian species which benefit from the disclosed methods for the inhibition of cancer cell growth, include, but are not limited to, primates, such as apes, chimpanzees, orangutans, humans, monkeys; domesticated animals (e.g., pets) such as dogs, cats, guinea pigs, hamsters, Vietnamese pot-bellied pigs, rabbits, and ferrets; domesticated farm animals such as cows, buffalo, bison, horses, donkey, swine, sheep, and goats; exotic animals typically found in zoos, such as bear, lions, tigers, panthers, elephants, hippopotamus, rhinoceros, giraffes, antelopes, sloth, gazelles, zebras, wildebeests, prairie dogs, koala bears, kangaroo, opossums, raccoons, pandas, hyena, seals, sea lions, elephant seals, otters, porpoises, dolphins, and whales. The terms "patient" and "subject" are used herein interchangeably and are intended to include such human and non-human mammalian species. Likewise, in vitro methods of the present invention can be earned out on cells of such mammalian species.

**[0239]** Patients in need of treatment using the methods of the present invention can be identified using standard techniques known to those in the medical profession.

**[0240]** The following examples are offered by way of illustration and not by way of limitation.

## 8. EXAMPLES

### 8.1. Example 1

#### In Vitro Screening

**[0241]** Cell Lines and NCI Diversity Set. All cell lines can be purchased from ATCC or described previously (Cheng et al., 1997, *Oncogene* 14: 2793-2801; West et al., 2002, *Drug Resist Updat* 5: 234-248; Satyamoorthy et al., 2001, *Cancer Res* 61: 7318-7324). The NCI Structural Diversity Set is a library of 1,992 compounds selected from the approximately 140,000-compound NCI drug depository. In-depth data on the selection, structures, and activities of these diversity set compounds can be found on the NCI Developmental Therapeutics Program web site.

**[0242]** Screening for Inhibition of Akt-transformed Cell Growth. AKT2 transformed NIH3T3 cells or LXSN vector-transfected NIH3T3 control cells (Cheng et al., 1997, *Oncogene* 14: 2793-2801) are plated into 96-well tissue culture plate. Following treatment with 5  $\mu$ M of NCI Diversity Set compound, cell growth can be detected with CellTier 96 One Solution Cell Proliferation kit (Promega). Compounds that inhibit growth in AKT2-transformed but not LXSN-transfected NIH3T3 cells are considered as candidates of Akt inhibitor and subjected to further analysis.

**[0243]** In vitro Protein Kinase, Cell Survival and Apoptosis Assays. In vitro kinase is performed as previously described (see, for example, Jiang et al., 2000, *Mol Cell Biol* 20: 139-148). Cell survival is assayed with MTS (Promega). Apoptosis was detected with annexin V, which is performed as previously described (Jiang et al., 2000, *Mol Cell Biol* 20: 139-148). Recombinant Akt and PDK1 are purchased from Upstate Biotechnology Inc.

## Results

**[0244]** Identification of Small Molecule Inhibitor of Akt Signaling Pathway, API-2. Frequent alterations of Akt has been detected in human cancer and disruption of Akt pathway induces apoptosis and inhibits tumor growth (Jetzt et al., 2003, *Cancer Res* 63: 697-706). Thus, Akt is considered as an attractive molecular target for development of novel cancer therapeutics. To identify small molecule inhibitor(s) of Akt, a chemical library of 1,992-compounds from the NCI (the NCI Diversity Set) is evaluated for agents capable of inhibition of growth in AKT2-transformed but not empty vector LXSN-transfected NIH3T3 cells. Repeated experiments showed that 32 compounds inhibited growth only in AKT2-transformed cells. The most potent of these compounds, API-2 (NCI identifier: NSC 154020), can suppress cell growth at a concentration of 50 nM. FIG. 1A shows the chemical structure of API-2, which is also known as triciribine (Schweinsberg et al., 1981, *Biochem Pharmacol* 30: 2521-2526). The fact that API-2 inhibits selectively AKT2 transformed cells over untransformed parental cells prompted us to determine whether API-2 is an inhibitor of AKT2 kinase. To this end, AKT2 is immunoprecipitated with anti-AKT2 antibody from AKT2 transformed NIH3T3 cells following treatment with API-2. AKT2 immunoprecipitates were immunoblotted with anti-phospho-Akt antibodies. As shown in FIG. 1B, API-2 significantly inhibited AKT2 phosphorylation at both threonine-309 and serine-474, which are required for full activation of AKT2 (Datta et al., 1999, *Genes Dev* 13: 2905-2927). As three isoforms of Akt share high homology and similar structure, the effect of API-2 on their kinase activities is evaluated. HEK293 cells are transfected with HA-Akt1, -AKT2 and -AKT3, serum-starved overnight and treated with API-2 for 60 min prior to EGF (50 ng/ml) stimulation. Triple experiments showed that API-2 suppressed EGF-induced kinase activity and phosphorylation of Akt1, AKT2 and AKT3 (FIG. 1C). However, kinase activity of recombinant constitutively active AKT2 (Myr-AKT2) is not inhibited by API-2 in an in vitro kinase reaction (FIG. 1D), suggesting that API-2 does not directly inhibit Akt in vitro and that API-2 neither functions as ATP competitor nor as the substrate competitor that binds to active site of Akt.

**[0245]** API-2 Does Not Inhibit Known Upstream Activators of Akt. It has been well documented that Akt is activated by extracellular stimuli and intracellular signal molecules, such as active Ras and Src, through a PI3K-dependent man-

ner. Therefore, API-2 inhibition of Akt could result from targeting upstream molecule(s) of Akt. As PI3K and PDK1 are direct upstream regulators of Akt (Datta et al., 1999, *Genes Dev* 13: 2905-2927), whether API-2 inhibits PI3K and/or PDK1 is examined. HEK293 cells are serum-starved and then can be treated with API-2 or PI3K inhibitor, wortmannin, for 30 min prior to EGF stimulation. PI3K is immunoprecipitated with anti-p110 $\alpha$  antibody. The immunoprecipitates are subjected to in vitro PI3K kinase assay using PI-4-P as a substrate. As shown in FIG. 2A, the EGF-induced PI3K activity is inhibited by wortmannin but not by API-2. To evaluate the effect of API-2 on PDK1, an assay in which recombinant PDK1 promotes the threonine-309 phosphorylation of AKT2 peptides is used in the presence of lipid vesicles containing phosphatidylinositol. As shown in FIG. 2B, the assay is potently inhibited by the control PDK1 inhibitor staurosporine (1050=5 nM). In contrast, API-2 displayed only 21% inhibition of the assay at the highest concentration tested (5.1  $\mu$ M). To further evaluate the effect of API-2 on PDK1 activation, the autophosphorylation level of PDK1 at serine-241, a residue that is phosphorylated by itself and is critical for its activity is examined (Datta et al., 1999, *Genes Dev* 13: 2905-2927), following API-2 treatment of HEK293 cells. Triplicate experiments show that phosphorylation levels of PDK1 are not inhibited by API-2 (FIG. 2B). However, PI3K inhibitor wortmannin can inhibit EGF-stimulated PDK1 (FIG. 2B).

**[0246]** API-2 Is Highly Selective for the Akt over PKC, PKA, SGK, STAT, JNK, p38, and ERK Signaling Pathways. Akt belongs to AGC (PKA/PKG/PKC) kinase family, which also include PKA, PKC, serum- and glucocorticoid-inducible kinase (SGK), p90 ribosomal S6 kinase, p70<sup>S6K</sup>, mitogen- and stress-activated protein kinase and PKC-related kinase. Among AGC kinase family, protein structures of PKA, PKC and SGK are more close to Akt kinase than other members. Therefore, next examined are the effects of API-2 on the enzymatic activities of these 3 kinases. HEK293 cells are transfected with HA-tagged PKA, PKC $\alpha$  or SGK. In vitro kinase assay and immunoblotting analysis show that the kinase activities of PKA and PKC $\alpha$  are inhibited by PKAi and Ro 31-8220, a PKC inhibitor, respectively, whereas API-2 exhibits no effect on their activities (FIGS. 2C and 2E). Further, serum-induced SGK kinase activity is attenuated by wortmannin but not by API-2 (FIG. 2D). In addition, it is determined whether API-2 has effect on other oncogenic survival pathways. Western blotting analyses with commercially available anti-phospho-antibodies reveals that phosphorylation levels of Stat3, INK, p38 and Erk1/2 were not affected by API-2 treatment (FIG. 2F). These data indicate that API-2 specifically inhibits Akt signaling pathway.

**[0247]** API-2 Suppresses Cell Growth and Induces Apoptosis in Akt-overexpressing/activating Human Cancer Cell Lines. The ability of API-2 to selectively inhibit the Akt pathway suggests that it should inhibit proliferation and/or induces apoptosis preferentially in those tumor cells with aberrant expression/activation of Akt. As activation of Akt in human malignancies commonly results from overexpression of Akt or PTEN mutations, API-2 is used to treat the cells that express constitutively active Akt, caused by overexpression of AKT2 (OVCAR3, OVCAR8, PANC1 and AKT2-transformed NIH3T3) or mutations of the PTEN gene (PC-3, LNCaP, MDA-MB-468), and cells that do not (OVCAR5, DU-145, T47D, COLO357 and LXSN-NIH3T3) as well as melanoma cells that are activated by IGF-1 to activate Akt or



do not respond to growth stimulation by IGF-1 (Satyamoorthy et al., 2001, *Cancer Res* 61: 7318-7324). Immunoblotting analysis showed that phosphorylation levels of Akt are inhibited by API-2 only in the cells expressing elevated Akt or responding to IGF-1 stimulation (FIG. 3A). Accordingly, API-2 inhibited cell growth to a much higher degree in Akt-overexpressing/activating cells as compared to those with low levels of Akt. As shown in FIG. 3B, API-2 treatment inhibited cell proliferation by approximate 50-60% in Akt-overexpressing/activating cell lines, LNCaP, PC-3, OVCAR3, OVCAR8, PANC1, MDA-MB-468, and WM35, whereas only by about 10-20% in DU145, OVCAR5, COLO357, T47D and WM852 cells, which exhibit low levels of Akt or do not respond to growth stimulation by IGF-1. Moreover, API-2 induces apoptosis by 8-fold (OVCAR3), 6-fold (OVCAR8), 6-fold (PANC1), and 3-fold (AKT2-NIH3T3). No significant difference of apoptosis is observed between API-2 and vehicle (DMSO) treatment in OVCAR5, COLO357 and LXSNIH3T3 cells. Thus, API-2 inhibits cell growth and induces apoptosis preferentially in cells that express aberrant Akt.

**[0248]** API-2 Inhibits Downstream Targets of Akt. It has been shown that Akt exerts its cellular effects through phosphorylation of a number of proteins (Datta et al., 1999, *Genes Dev* 13: 2905-2927). More than 20 proteins have been identified as Akt substrates, including the members of Forkhead protein family (FKHR, AFX and FKHL1), tuberlin/TSC2, p70<sup>S6K</sup>, GSK-3 $\beta$ , p21<sup>WAF1/Cip1</sup>, p27<sup>Kip1</sup>, MDM2, Bad, ASK1 and IKK $\alpha$  etc. it is next examined whether API-2 inhibits downstream targets of Akt. As anti-phospho-tuberlin, -Bad, -AFX, and -GSK-3 $\beta$  antibodies are commercially available, therefore, the effect of API-2 on their phosphorylation induced by Akt was determined. Following API-2 (1  $\mu$ M) treatment, OVCAR3 cells were lysed and immunoblotted with the individual anti-phospho-antibody. FIG. 4A shows that API-2 considerably inhibited the phosphorylation levels of tuberlin leading to stabilization and upregulation of tuberlin (Dan et al., 2002, *J Biol Chem* 277: 35364-35370). The phosphorylation levels of Bad, GSK-3 $\beta$ , and AFX are partially attenuated by API-2. These data suggest that API-2 induces cell death and cell growth arrest by inhibiting phosphorylation of its downstream targets. API-2 inhibition of Akt downstream targets at different degrees could be due to the fact that phosphorylation sites of these targets are also regulated by other kinase(s), for instance, Bad serine-136 is phosphorylated by PAK1 in addition to Akt (Schurmann et al., 2000, *Mol Cell Biol* 20: 453-461).

## 8.2 Example 2

### Antitumor Activity in the Nude Mouse Tumor Xenograft Model

**[0249]** Tumor cells can be harvested, suspended in PBS, and can be injected s.c. into the right and left flanks ( $2 \times 10^6$  cells/flank) of 8-week-old female nude mice as reported previously (Sun et al., 1999, *Cancer Res* 59: 4919-4926). When tumors reach about 100-150 mm<sup>3</sup>, animals are randomized and dosed i.p. with 0.2 ml vehicle of the triciribine compound and/or one or more platinum compounds daily. Control animals receive DMSO (20%) vehicle, whereas treated animals can be injected with API-2 (1 mg/kg/day) in 20% DMSO.

**[0250]** API-2 Inhibits the Growth of Tumors in Nude Mice that Overexpress Akt. Frequent overexpression/activation and/or amplification of AKT1 and AKT2 in human ovarian

and pancreatic cancer was shown (Cheng et al., AKT signal transduction pathway in oncogenesis, in Schwab D(ed.) *Encyclopedic Reference of cancer*, Springer, pp 35-7). Inhibition of Akt pathway by inhibitors of PI3K, HSP70, Src and farnesyltransferase resulted in cell growth arrest and induction of apoptosis (Split et al., 2003, *Cancer Res* 63: 2139-2144; Xu et al., 2003, *Cancer Res* 63: 7777-7784). A recent study showed that the tumor growth of xenografts with elevated Akt was also significantly inhibited by intratumoral injection of adenovirus of dominant negative Akt (Jetzt et al., 2003, *Cancer Res* 63: 697-706). Because API-2 inhibits Akt signaling and induces apoptosis and cell growth arrest only in cancer cells with elevated levels of Akt (FIG. 3), the growth of tumors with elevated levels of Akt should be more sensitive to API-2 than that of tumors with low levels of Akt in nude mice. To this end, s.c. Akt-overexpressing cells (OVCAR3, OVCAR8 and PANC-1) are s.c. implanted into the right flank, and those cell lines that express low levels of Akt (OVCAR5 and COLO357) into the left flank of mice. When the tumors reach an average size of about 100-150 mm<sup>3</sup>, the animals are randomized and treated i.p. with either vehicle or API-2 (1 mg/kg/day). As illustrated in FIG. 4B, OVCAR-5 and COLO357 tumors treated with vehicle grew to about 800-1,000 mm<sup>3</sup> 49 days after tumor implantation. OVCAR3, OVCAR8 and PANC1 tumors treated with vehicle control grew to about 700-900 mm<sup>3</sup> 49 days after tumor implantation. API-2 inhibited OVCAR3, OVCAR8 and PANC1 tumor growth by 90%, 88% and 80%, respectively. In contrast, API-2 has little effect on the growth of OVCAR5 and COLO357 cells in nude mice (FIGS. 4B-4D and data not shown). At dose 1 mg/kg/day, API-2 had no effects on blood glucose level, body weight, activity and food intake of mice. In treated tumor samples, Akt activity was inhibited by API-2 without change of total Akt content (FIG. 4E). Taken together, these results indicate that API-2 selectively inhibits the growth of tumors with elevated levels of Akt.

## 8.3 Example 3

### TCN Directly Inhibits Wild Type Akt Kinase Activity

**[0251]** API-2 (TCN) can directly inhibit wild type Akt kinase activity induced by PDK1 in vitro (FIG. 1). This result supports that API-2 is a direct Akt inhibitor and that the underlying mechanism may be API-2 binding to PH domain and/or threonine-308 of Akt. An in vitro kinase assay is performed with recombinant of PDK1 and Akt in a kinase buffer containing phosphatidylinositol-3,4,5-P3 (PIP3), API-2 and histone H2B as substrate. After incubation of 30 min, the reactions were separated by SDS-PAGE and exposed in a film.

## 8.4 Example 4

### TCN is Effective in Cancer Resistant Cells

**[0252]** The effects of TCN (API-2) are tested in cisplatin, paclitaxel, and tamoxifen resistant A270CP, C-13, OVCAR433 and MCF7/TAM cells. API-2 overcame cisplatin, paclitaxel, and tamoxifen resistance in these cells

## 8.5 Example 5

### TCN Overcomes Cisplatin Resistance in Cisplatin Resistant Ovarian Cancer Cells

**[0253]** Ovarian cancer cell lines sensitive to cisplatin, A2780S and OV2008, and their cisplatin resistant derivatives,

A2780CP and C13, respectively, were treated with 0, 10, 20 and 30  $\mu$ M cisplatin, followed by an assessment of the percentage of cells surviving the cisplatin. The resistance to cisplatin toxicity was confirmed in the A2780S and C13 cell lines (FIG. 9A). Lysates from all four cell types were resolved by SDS-PAGE and probed with antibodies to phosphorylated Akt and total Akt. The cisplatin resistant cell lines endogenously express significantly higher levels phosphorylated Akt (FIG. 9B).

**[0254]** It was next assessed what effects TCN would have on cisplatin resistance. A2780CP and C13 ovarian cancer cells were treated with 0, 5, 10 and 20  $\mu\text{M}$ , with or without 10  $\mu\text{M}$  TCN, followed by measurement of cell survival. The addition of TCN significantly lowered the resistance to cisplatin in both cell lines (FIG. 10). C13 cells were then treated with 0, 1, 5, 10 and 20  $\mu\text{M}$  TCN with or without 10  $\mu\text{M}$  cisplatin, followed by measurement of cell survival. The combination of TCN and cisplatin synergistically reduced cell survival (FIG. 11).

**[0255]** C13 cells were injected subcutaneously in nude mice, followed by treatment with vehicle (DMSO), cisplatin alone (2 mg/kg/day), TCN alone (2 mg/kg/day) or TCN and cisplatin. Tumor mass was assessed at weekly intervals for four weeks. The combination of cisplatin and TCN enhanced the ability to suppress the growth and progression of the tumor (FIG. 12).

### 8.6 Example 6

### TCN Enhances the Effects of mTOR Inhibition

**[0256]** OV3 and MCF7 cells were treated with the mTOR inhibitors rapamycin and RAD001, respectively, over a 24 hour time course. Lysates were resolved by SDS-PAGE and transferred to a membrane for western blotting. Membranes were probed with antibodies to phospho-Akt, total Akt, phospho-p70S6K, and total p70S6K. Membranes with lysates derived from OV3 cells were additionally probed with antibodies to phospho-FKHRL 1 and total FKHRL 1. The use of mTOR inhibitors lead to rapid activation of Akt pathway in both cell lines (FIG. 13).

**[0257]** OV3 cells were then treated with either control, rapamycin, or rapamycin and TCN and MCF7 cells were treated with control, RAD001, or RAD001 and TCN. TCN markedly decreased the mTOR inhibitors' increase in Akt

phosphorylation without a subsequent increase in p70S6K activity (FIG. 14). DU-145, MCF7 and OV3 cell growth rates were then assessed over 6 days in the presence of control, mTOR inhibitor, TCN, or TCN and mTOR inhibitor. The inhibition on cell growth by mTOR inhibition was enhanced by TCN (FIG. 15).

### 8.7 Example 7

### TCN Overcomes Aurora-A Induced Cisplatin Resistance

**[0258]** Aurora-A is a serine/threonine kinase involved in progression of the cell cycle. The effect of Aurora-A on sensitivity to cisplatin was examined. A2780S and A2780CP cells, as well as OV2008 cells, were transfected with empty vector or with a construct encoding an HA-tagged Aurora-A protein. Expression of Aurora-A was confirmed by western blotting lysates resolved by SDS-PAGE and transferred to a membrane (FIG. 16, top panels). The viability of cells was assessed in the presence of paclitaxel and cisplatin. A2780S and OV2008 cells transfected with Aurora-A showed improved viability over A2780S and OV2008 cells transfected with empty vector only (FIG. 16).

**[0259]** It was next assessed what effect Aurora-A exhibited on the induction of apoptosis induced by cisplatin. A27805 cells were transfected with empty vector or with a construct encoding the Aurora-A protein. Cells were then treated with cisplatin. Immunofluorescent staining (FIG. 17A) shows that expression of Aurora-A inhibits cytochrome c release. Western blot analysis revealed that introduction of Aurora-A increases phosphorylation of Akt, as well as Akt kinase activity (FIGS. 17B and C).

**[0260]** Next, A27805 and OV2008 cells, transfected with either empty vector or Aurora-A, were treated with either cisplatin, TCN, or cisplatin and TCN. Both A2780S and OV2008 cells expressing Aurora-A showed markedly decreased resistance to cisplatin in the presence of TCN (FIG. 18).

[0261] This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention will be obvious to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of this invention.

## SEQUENCE LISTING

<160> NUMBER OF SEO ID NOS: 6

```
<210> SEQ ID NO 1
<211> LENGTH: 1443
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1443)
```

&lt;400&gt; SEQUENCE: 1

atg aac gac gta gcc att gtg aag gag ggc tgg ctg cac aaa cga ggg 48  
Met Asn Asp Val Ala Ile Val Lys Glu Gly Trp Leu His Lys Arg Gly  
1 5 10 15

qaa tat att aaa acc tqg cgg cca cgc tac ttc ctc ctc aaq aac gat 96

-continued

Glu	Tyr	Ile	Lys	Thr	Trp	Arg	Pro	Arg	Tyr	Phe	Leu	Leu	Lys	Asn	Asp	
			20					25					30			
ggc	acc	ttt	att	ggc	tac	aag	gaa	cgg	cct	cag	gat	gtg	gat	cag	cga	144
Gly	Thr	Phe	Ile	Gly	Tyr	Lys	Glu	Arg	Pro	Gln	Asp	Val	Asp	Gln	Arg	
		35					40					45				
gag	tcc	cca	ctc	aac	aac	ttc	tca	gtg	gca	caa	tgc	cag	ctg	atg	aag	192
Glu	Ser	Pro	Leu	Asn	Asn	Phe	Ser	Val	Ala	Gln	Cys	Gln	Leu	Met	Lys	
	50					55					60					
aca	gag	cgg	cca	agg	ccc	aac	acc	ttt	atc	atc	cgc	tgc	ctg	cag	tggt	240
Thr	Glu	Arg	Pro	Arg	Pro	Asn	Thr	Phe	Ile	Ile	Arg	Cys	Leu	Gln	Trp	
65					70				75						80	
acc	aca	gtc	att	gag	cgc	acc	ttc	cat	gta	gaa	acg	cct	gag	gag	cgg	288
Thr	Thr	Val	Ile	Glu	Arg	Thr	Phe	His	Val	Glu	Thr	Pro	Glu	Glu	Arg	
			85					90						95		
gaa	gaa	tggt	gcc	acc	gcc	att	cag	act	gtg	gcc	gat	gga	ctc	aag	agg	336
Glu	Glu	Trp	Ala	Thr	Ala	Ile	Gln	Thr	Val	Ala	Asp	Gly	Leu	Lys	Arg	
		100					105						110			
cag	gaa	gaa	gag	acg	atg	gac	ttc	cga	tca	ggc	tca	ccc	agt	gac	aac	384
Gln	Glu	Glu	Glu	Thr	Met	Asp	Phe	Arg	Ser	Gly	Ser	Pro	Ser	Asp	Asn	
		115				120						125				
tca	ggg	gct	gaa	gag	atg	gag	gtg	tcc	ctg	gcc	aag	ccc	aag	cac	cgt	432
Ser	Gly	Ala	Glu	Glu	Met	Glu	Val	Ser	Leu	Ala	Lys	Pro	Lys	His	Arg	
	130					135					140					
gtg	acc	atg	aac	gag	ttt	gag	tac	ctg	aaa	cta	ctg	ggc	aag	ggc	acc	480
Val	Thr	Met	Asn	Glu	Phe	Glu	Tyr	Leu	Lys	Leu	Leu	Gly	Lys	Gly	Thr	
145					150					155					160	
ttt	ggg	aaa	gtg	att	ctg	gta	aaa	gag	aag	gcc	aca	ggc	cgc	tac	tat	528
Phe	Gly	Lys	Val	Ile	Leu	Val	Lys	Glu	Lys	Ala	Thr	Gly	Arg	Tyr	Tyr	
			165					170						175		
gcc	atg	aag	atc	ctc	aag	aag	gag	gtc	atc	gtc	gcc	aag	gat	gag	gtt	576
Ala	Met	Lys	Ile	Leu	Lys	Lys	Glu	Val	Ile	Val	Ala	Lys	Asp	Glu	Val	
		180					185						190			
gcc	cac	acg	ctt	act	gaa	aac	cgt	gtc	ctg	cag	aac	tct	agg	cat	ccc	624
Ala	His	Thr	Leu	Thr	Glu	Asn	Arg	Val	Leu	Gln	Asn	Ser	Arg	His	Pro	
		195					200					205				
ttc	ctt	acg	gcc	ctc	aag	tac	tca	ttc	cag	acc	cac	gac	cgc	ctc	tggt	672
Phe	Leu	Thr	Ala	Leu	Lys	Tyr	Ser	Phe	Gln	Thr	His	Asp	Arg	Leu	Cys	
	210					215					220					
ttt	gtc	atg	gag	tat	gcc	aac	ggg	ggc	gag	ctc	ttc	ttc	cac	ctg	tct	720
Phe	Val	Met	Glu	Tyr	Ala	Asn	Gly	Gly	Glu	Leu	Phe	Phe	His	Leu	Ser	
225					230				235					240		
cga	gag	cgc	gtg	ttc	tcc	gag	gac	cgg	gcc	cgc	ttc	tat	ggt	gcg	gag	768
Arg	Glu	Arg	Val	Phe	Ser	Glu	Asp	Arg	Ala	Arg	Phe	Tyr	Gly	Ala	Glu	
			245						250					255		
att	gtg	tct	gcc	ctg	gac	tac	ttg	cac	tgc	gag	aag	aac	gtg	gtg	tac	816
Ile	Val	Ser	Ala	Leu	Asp	Tyr	Leu	His	Ser	Glu	Lys	Asn	Val	Val	Tyr	
			260					265					270			
cgg	gac	ctg	aag	cta	gag	aac	ctc	atg	ctg	gac	aag	gac	ggg	cac	atc	864
Arg	Asp	Leu	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	Gly	His	Ile	
		275					280					285				
aag	ata	acg	gac	ttc	ggg	ctg	tgc	aag	gag	ggg	atc	aag	gat	ggt	gcc	912
Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Lys	Asp	Gly	Ala	
		290					295				300					
act	atg	aag	aca	ttc	tgc	gga	acg	ccg	gag	tac	ctg	gcc	cct	gag	gtg	960
Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	Pro	Glu	Val	
305					310					315				320		
ctg	gaa	gac	aac	gac	tac	ggc	cgt	gca	gtg	gac	tgg	tgg	ggg	ctg	ggc	1008

-continued

Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	Gly	Leu	Gly		
			325						330					335			
gtg	gtc	atg	tat	gag	atg	atg	tgt	ggc	cgc	ctg	ccc	ttc	tac	aac	cag	1056	
Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	Tyr	Asn	Gln		
			340					345					350				
gac	cac	gag	aag	ctg	ttc	gag	ctg	atc	ctc	atg	gag	gag	atc	cgc	ttc	1104	
Asp	His	Glu	Lys	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Glu	Ile	Arg	Phe		
			355			360						365					
ccg	cgc	aca	ctc	ggc	cct	gag	gcc	aag	tcc	ctg	ctc	tcc	ggg	ctg	ctc	1152	
Pro	Arg	Thr	Leu	Gly	Pro	Glu	Ala	Lys	Ser	Leu	Leu	Ser	Gly	Leu	Leu		
			370			375						380					
aag	aag	gac	cct	aca	cag	agg	ctc	ggt	ggg	ggc	tct	gag	gat	gcc	aag	1200	
Lys	Lys	Asp	Pro	Thr	Gln	Arg	Leu	Gly	Gly	Gly	Ser	Glu	Asp	Ala	Lys		
			385		390				395					400			
gag	atc	atg	cag	cac	cgg	ttc	ttt	gcc	aac	atc	gtg	tgg	cag	gat	gtg	1248	
Glu	Ile	Met	Gln	His	Arg	Phe	Phe	Ala	Asn	Ile	Val	Trp	Gln	Asp	Val		
			405					410					415				
tat	gag	aag	aag	ctg	agc	cca	cct	ttc	aag	ccc	cag	gtc	acc	tct	gag	1296	
Tyr	Glu	Lys	Lys	Leu	Ser	Pro	Pro	Phe	Lys	Pro	Gln	Val	Thr	Ser	Glu		
			420					425					430				
act	gac	acc	agg	tat	ttc	gat	gag	gag	ttc	aca	gct	cag	atg	atc	acc	1344	
Thr	Asp	Thr	Arg	Tyr	Phe	Asp	Glu	Glu	Phe	Thr	Ala	Gln	Met	Ile	Thr		
			435			440						445					
atc	acg	ccg	cct	gat	caa	gat	gac	agc	atg	gag	tgt	gtg	gac	agt	gag	1392	
Ile	Thr	Pro	Pro	Asp	Gln	Asp	Asp	Ser	Met	Glu	Cys	Val	Asp	Ser	Glu		
			450			455					460						
cgg	agg	ccg	cac	ttc	ccc	cag	ttc	tcc	tac	tca	gcc	agt	ggc	aca	gcc	1440	
Arg	Arg	Pro	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser	Ala	Ser	Gly	Thr	Ala		
			465		470					475				480			
tga																1443	

<210> SEQ ID NO 2  
 <211> LENGTH: 480  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met	Asn	Asp	Val	Ala	Ile	Val	Lys	Glu	Gly	Trp	Leu	His	Lys	Arg	Gly
1			5					10					15		
Glu	Tyr	Ile	Lys	Thr	Trp	Arg	Pro	Arg	Tyr	Phe	Leu	Leu	Lys	Asn	Asp
			20				25					30			
Gly	Thr	Phe	Ile	Gly	Tyr	Lys	Glu	Arg	Pro	Gln	Asp	Val	Asp	Gln	Arg
			35			40					45				
Glu	Ser	Pro	Leu	Asn	Asn	Phe	Ser	Val	Ala	Gln	Cys	Gln	Leu	Met	Lys
			50		55					60					
Thr	Glu	Arg	Pro	Arg	Pro	Asn	Thr	Phe	Ile	Ile	Arg	Cys	Leu	Gln	Trp
			65		70				75				80		
Thr	Thr	Val	Ile	Glu	Arg	Thr	Phe	His	Val	Glu	Thr	Pro	Glu	Glu	Arg
			85				90						95		
Glu	Glu	Trp	Ala	Thr	Ala	Ile	Gln	Thr	Val	Ala	Asp	Gly	Leu	Lys	Arg
			100				105					110			
Gln	Glu	Glu	Glu	Thr	Met	Asp	Phe	Arg	Ser	Gly	Ser	Pro	Ser	Asp	Asn
			115			120					125				
Ser	Gly	Ala	Glu	Glu	Met	Glu	Val	Ser	Leu	Ala	Lys	Pro	Lys	His	Arg
			130			135					140				

-continued

---

Val	Thr	Met	Asn	Glu	Phe	Glu	Tyr	Leu	Lys	Leu	Leu	Gly	Lys	Gly	Thr	145	150	155	160
Phe	Gly	Lys	Val	Ile	Leu	Val	Lys	Glu	Lys	Ala	Thr	Gly	Arg	Tyr	Tyr	165	170	175	
Ala	Met	Lys	Ile	Leu	Lys	Lys	Glu	Val	Ile	Val	Ala	Lys	Asp	Glu	Val	180	185	190	
Ala	His	Thr	Leu	Thr	Glu	Asn	Arg	Val	Leu	Gln	Asn	Ser	Arg	His	Pro	195	200	205	
Phe	Leu	Thr	Ala	Leu	Lys	Tyr	Ser	Phe	Gln	Thr	His	Asp	Arg	Leu	Cys	210	215	220	
Phe	Val	Met	Glu	Tyr	Ala	Asn	Gly	Gly	Glu	Leu	Phe	Phe	His	Leu	Ser	225	230	235	240
Arg	Glu	Arg	Val	Phe	Ser	Glu	Asp	Arg	Ala	Arg	Phe	Tyr	Gly	Ala	Glu	245	250	255	
Ile	Val	Ser	Ala	Leu	Asp	Tyr	Leu	His	Ser	Glu	Lys	Asn	Val	Val	Tyr	260	265	270	
Arg	Asp	Leu	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	Gly	His	Ile	275	280	285	
Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Lys	Asp	Gly	Ala	290	295	300	
Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	Pro	Glu	Val	305	310	315	320
Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	Gly	Leu	Gly	325	330	335	
Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	Tyr	Asn	Gln	340	345	350	
Asp	His	Glu	Lys	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Glu	Ile	Arg	Phe	355	360	365	
Pro	Arg	Thr	Leu	Gly	Pro	Glu	Ala	Lys	Ser	Leu	Leu	Ser	Gly	Leu	Leu	370	375	380	
Lys	Lys	Asp	Pro	Thr	Gln	Arg	Leu	Gly	Gly	Gly	Ser	Glu	Asp	Ala	Lys	385	390	395	400
Glu	Ile	Met	Gln	His	Arg	Phe	Phe	Ala	Asn	Ile	Val	Trp	Gln	Asp	Val	405	410	415	
Tyr	Glu	Lys	Lys	Leu	Ser	Pro	Pro	Phe	Lys	Pro	Gln	Val	Thr	Ser	Glu	420	425	430	
Thr	Asp	Thr	Arg	Tyr	Phe	Asp	Glu	Glu	Phe	Thr	Ala	Gln	Met	Ile	Thr	435	440	445	
Ile	Thr	Pro	Pro	Asp	Gln	Asp	Asp	Ser	Met	Glu	Cys	Val	Asp	Ser	Glu	450	455	460	
Arg	Arg	Pro	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser	Ala	Ser	Gly	Thr	Ala	465	470	475	480

<210> SEQ ID NO 3  
 <211> LENGTH: 1515  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1446)

<400> SEQUENCE: 3

atg aat gag gtg tct gtc atc aaa gaa ggc tgg ctc cac aag cgt ggt  
 Met Asn Glu Val Ser Val Ile Lys Glu Gly Trp Leu His Lys Arg Gly

-continued

1	5	10	15	
gaa tac atc aag acc tgg agg cca cgg tac ttc ctg ctg aag agc gac				96
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Ser Asp				
	20	25	30	
ggc tcc ttc att ggg tac aag gag agg ccc gag gcc cct gat cag act				144
Gly Ser Phe Ile Gly Tyr Lys Glu Arg Pro Glu Ala Pro Asp Gln Thr				
	35	40	45	
cta ccc ccc tta aac aac ttc tcc gta gca gaa tgc cag ctg atg aag				192
Leu Pro Pro Leu Asn Asn Phe Ser Val Ala Glu Cys Gln Leu Met Lys				
	50	55	60	
acc gag agg ccg cga ccc aac acc ttt gtc ata cgc tgc ctg cag tgg				240
Thr Glu Arg Pro Arg Pro Asn Thr Phe Val Ile Arg Cys Leu Gln Trp				
	65	70	75	80
acc aca gtc atc gag agg acc ttc cac gtg gat tct cca gac gag agg				288
Thr Thr Val Ile Glu Arg Thr Phe His Val Asp Ser Pro Asp Glu Arg				
	85	90	95	
gag gag tgg atg cgg gcc atc cag atg gtc gcc aac agc ctc aag cag				336
Glu Glu Trp Met Arg Ala Ile Gln Met Val Ala Asn Ser Leu Lys Gln				
	100	105	110	
cgg gcc cca ggc gag gac ccc atg gac tac aag tgt ggc tcc ccc agt				384
Arg Ala Pro Gly Glu Asp Pro Met Asp Tyr Lys Cys Gly Ser Pro Ser				
	115	120	125	
gac tcc tcc acg act gag gag atg gaa gtg gcg gtc agc aag gca cgg				432
Asp Ser Ser Thr Thr Glu Glu Met Glu Val Ala Val Ser Lys Ala Arg				
	130	135	140	
gct aaa gtg acc atg aat gac ttc gac tat ctc aaa ctc ctt ggc aag				480
Ala Lys Val Thr Met Asn Asp Phe Asp Tyr Leu Lys Leu Leu Gly Lys				
	145	150	155	160
gga acc ttt ggc aaa gtc atc ctg gtg cgg gag aag gcc act ggc cgc				528
Gly Thr Phe Gly Lys Val Ile Leu Val Arg Glu Lys Ala Thr Gly Arg				
	165	170	175	
tac tac gcc atg aag atc ctg cga aag gaa gtc atc att gcc aag gat				576
Tyr Tyr Ala Met Lys Ile Leu Arg Lys Glu Val Ile Ile Ala Lys Asp				
	180	185	190	
gaa gtc gct cac aca gtc acc gag agc cgg gtc ctc cag aac acc agg				624
Glu Val Ala His Thr Val Thr Glu Ser Arg Val Leu Gln Asn Thr Arg				
	195	200	205	
cac ccg ttc ctc act gcg ctg aag tat gcc ttc cag acc cac gac cgc				672
His Pro Phe Leu Thr Ala Leu Lys Tyr Ala Phe Gln Thr His Asp Arg				
	210	215	220	
ctg tgc ttt gtg atg gag tat gcc aac ggg ggt gag ctg ttc ttc cac				720
Leu Cys Phe Val Met Glu Tyr Ala Asn Gly Gly Glu Leu Phe Phe His				
	225	230	235	240
ctg tcc cgg gag cgt gtc ttc aca gag gag cgg gcc cgg ttt tat ggt				768
Leu Ser Arg Glu Arg Val Phe Thr Glu Glu Arg Ala Arg Phe Tyr Gly				
	245	250	255	
gca gag att gtc tcg gct ctt gag tac ttg cac tcg cgg gac gtg gta				816
Ala Glu Ile Val Ser Ala Leu Glu Tyr Leu His Ser Arg Asp Val Val				
	260	265	270	
tac cgc gac atc aag ctg gaa aac ctc atg ctg gac aaa gat ggc cac				864
Tyr Arg Asp Ile Lys Leu Glu Asn Leu Met Leu Asp Lys Asp Gly His				
	275	280	285	
atc aag atc act gac ttt ggc ctc tgc aaa gag ggc atc agt gac ggg				912
Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile Ser Asp Gly				
	290	295	300	
gcc acc atg aaa acc ttc tgt ggg acc ccg gag tac ctg gcg cct gag				960
Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu Ala Pro Glu				

-continued

305	310	315	320	
gtg ctg gag gac aat gac tat ggc cgg gcc gtg gac tgg tgg ggg ctg				1008
Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp Trp Gly Leu	325	330	335	
ggt gtg gtc atg tac gag atg atg tgc ggc cgc ctg ccc ttc tac aac				1056
Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro Phe Tyr Asn	340	345	350	
cag gac cac gag cgc ctc ttc gag ctc atc ctc atg gaa gag atc cgc				1104
Gln Asp His Glu Arg Leu Phe Glu Leu Ile Leu Met Glu Ile Arg	355	360	365	
ttc cgg cgc acg ctc agc ccc gag gcc aag tcc ctg ctt gct ggg ctg				1152
Phe Pro Arg Thr Leu Ser Pro Glu Ala Lys Ser Leu Leu Ala Gly Leu	370	375	380	
ctt aag aag gac ccc aag cag agg ctt ggt ggg ggg ccc agc gat gcc				1200
Leu Lys Lys Asp Pro Lys Gln Arg Leu Gly Gly Gly Pro Ser Asp Ala	385	390	395	400
aag gag gtc atg gag cac agg ttc ttc ctc agc atc aac tgg cag gac				1248
Lys Glu Val Met Glu His Arg Phe Phe Leu Ser Ile Asn Trp Gln Asp	405	410	415	
gtg gtc cag aag aag ctc ctg cca ccc ttc aaa cct cag gtc acg tcc				1296
Val Val Gln Lys Lys Leu Leu Pro Pro Phe Lys Pro Gln Val Thr Ser	420	425	430	
gag gtc gac aca agg tac ttc gat gat gaa ttt acc gcc cag tcc atc				1344
Glu Val Asp Thr Arg Tyr Phe Asp Asp Glu Phe Thr Ala Gln Ser Ile	435	440	445	
aca atc aca ccc cct gac cgc tat gac agc ctg ggc tta ctg gag ctg				1392
Thr Ile Thr Pro Pro Asp Arg Tyr Asp Ser Leu Gly Leu Leu Glu Leu	450	455	460	
gac cag cgg acc cac ttc ccc cag ttc tcc tac tcg gcc agc atc cgc				1440
Asp Gln Arg Thr His Phe Pro Gln Phe Ser Tyr Ser Ala Ser Ile Arg	465	470	475	480
gag tga gcagtctgcc cacgcagagg acgcacgctc gctgccatca ccgctgggtg				
Glu				1496
gtttttttacc cctgcccg				1515
<210> SEQ ID NO 4				
<211> LENGTH: 481				
<212> TYPE: PRT				
<213> ORGANISM: Homo sapiens				
<400> SEQUENCE: 4				
Met Asn Glu Val Ser Val Ile Lys Glu Gly Trp Leu His Lys Arg Gly				
1	5	10	15	
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Ser Asp				
	20	25	30	
Gly Ser Phe Ile Gly Tyr Lys Glu Arg Pro Glu Ala Pro Asp Gln Thr				
	35	40	45	
Leu Pro Pro Leu Asn Asn Phe Ser Val Ala Glu Cys Gln Leu Met Lys				
	50	55	60	
Thr Glu Arg Pro Arg Pro Asn Thr Phe Val Ile Arg Cys Leu Gln Trp				
	65	70	75	80
Thr Thr Val Ile Glu Arg Thr Phe His Val Asp Ser Pro Asp Glu Arg				
	85	90	95	
Glu Glu Trp Met Arg Ala Ile Gln Met Val Ala Asn Ser Leu Lys Gln				
	100	105	110	

Arg	Ala	Pro	Gly	Glu	Asp	Pro	Met	Asp	Tyr	Lys	Cys	Gly	Ser	Pro	Ser	
		115					120					125				
Asp	Ser	Ser	Thr	Thr	Glu	Glu	Met	Glu	Val	Ala	Val	Ser	Lys	Ala	Arg	
		130					135					140				
Ala	Lys	Val	Thr	Met	Asn	Asp	Phe	Asp	Tyr	Leu	Lys	Leu	Leu	Gly	Lys	
		145					150					155				
Gly	Thr	Phe	Gly	Lys	Val	Ile	Leu	Val	Arg	Glu	Lys	Ala	Thr	Gly	Arg	
				165							170					
Tyr	Tyr	Ala	Met	Lys	Ile	Leu	Arg	Lys	Glu	Val	Ile	Ile	Ala	Lys	Asp	
				180											190	
Glu	Val	Ala	His	Thr	Val	Thr	Glu	Ser	Arg	Val	Leu	Gln	Asn	Thr	Arg	
				195											205	
His	Pro	Phe	Leu	Thr	Ala	Leu	Lys	Tyr	Ala	Phe	Gln	Thr	His	Asp	Arg	
				210											220	
Leu	Cys	Phe	Val	Met	Glu	Tyr	Ala	Asn	Gly	Gly	Glu	Leu	Phe	Phe	His	
				225											235	240
Leu	Ser	Arg	Glu	Arg	Val	Phe	Thr	Glu	Glu	Arg	Ala	Arg	Phe	Tyr	Gly	
				245											255	
Ala	Glu	Ile	Val	Ser	Ala	Leu	Glu	Tyr	Leu	His	Ser	Arg	Asp	Val	Val	
				260											270	
Tyr	Arg	Asp	Ile	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	Gly	His	
				275											285	
Ile	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Ser	Asp	Gly	
				290											300	
Ala	Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	Pro	Glu	
				305											310	320
Val	Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	Gly	Leu	
				325											335	
Gly	Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	Tyr	Asn	
				340											350	
Gln	Asp	His	Glu	Arg	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Glu	Ile	Arg	
				355											365	
Phe	Pro	Arg	Thr	Leu	Ser	Pro	Glu	Ala	Lys	Ser	Leu	Leu	Ala	Gly	Leu	
				370											380	
Leu	Lys	Lys	Asp	Pro	Lys	Gln	Arg	Leu	Gly	Gly	Gly	Pro	Ser	Asp	Ala	
				385											395	400
Lys	Glu	Val	Met	Glu	His	Arg	Phe	Phe	Leu	Ser	Ile	Asn	Trp	Gln	Asp	
				405											415	
Val	Val	Gln	Lys	Lys	Leu	Leu	Pro	Pro	Phe	Lys	Pro	Gln	Val	Thr	Ser	
				420											425	
Glu	Val	Asp	Thr	Arg	Tyr	Phe	Asp	Asp	Glu	Phe	Thr	Ala	Gln	Ser	Ile	
				435											445	
Thr	Ile	Thr	Pro	Pro	Asp	Arg	Tyr	Asp	Ser	Leu	Gly	Leu	Leu	Glu	Leu	
				450											460	
Asp	Gln	Arg	Thr	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser	Ala	Ser	Ile	Arg	
				465											475	480
Glu																

```
<210> SEQ ID NO 5
<211> LENGTH: 1510
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```



-continued

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: CDS

&lt;222&gt; LOCATION: (3) .. (1460)

&lt;400&gt; SEQUENCE: 5

```

gg gct cag agg gga gtc atc atg agc gat gtt acc att gtg aaa gaa      47
  Ala Gln Arg Gly Val Ile Met Ser Asp Val Thr Ile Val Lys Glu
    1             5             10             15

ggt tgg gtt cag aag agg gga gaa tat ata aaa aac tgg agg cca aga      95
Gly Trp Val Gln Lys Arg Gly Glu Tyr Ile Lys Asn Trp Arg Pro Arg
              20             25             30

tac ttc ctt ttg aag aca gat ggc tca ttc ata gga tat aaa gag aaa     143
Tyr Phe Leu Leu Lys Thr Asp Gly Ser Phe Ile Gly Tyr Lys Glu Lys
              35             40             45

cct caa gat gtg gat tta cct tat ccc ctc aac aac ttt tca gtg gca     191
Pro Gln Asp Val Asp Leu Pro Tyr Pro Leu Asn Asn Phe Ser Val Ala
              50             55             60

aaa tgc cag tta atg aaa aca gaa cga cca aag cca aac aca ttt ata     239
Lys Cys Gln Leu Met Lys Thr Glu Arg Pro Lys Pro Asn Thr Phe Ile
              65             70             75

atc aga tgt ctc cag tgg act act gtt ata gag aga aca ttt cat gta     287
Ile Arg Cys Leu Gln Trp Thr Thr Val Ile Glu Arg Thr Phe His Val
              80             85             90             95

gat act cca gag gaa agg gaa gaa tgg aca gaa gct atc cag gct gta     335
Asp Thr Pro Glu Glu Arg Glu Glu Trp Thr Glu Ala Ile Gln Ala Val
              100             105             110

gca gac aga ctg cag agg caa gaa gag gag aga atg aat tgt agt cca     383
Ala Asp Arg Leu Gln Arg Gln Glu Glu Glu Arg Met Asn Cys Ser Pro
              115             120             125

act tca caa att gat aat ata gga gag gaa gag atg gat gcc tct aca     431
Thr Ser Gln Ile Asp Asn Ile Gly Glu Glu Glu Met Asp Ala Ser Thr
              130             135             140

acc cat cat aaa aga aag aca atg aat gat ttt gac tat ttg aaa cta     479
Thr His His Lys Arg Lys Thr Met Asn Asp Phe Asp Tyr Leu Lys Leu
              145             150             155

cta ggt aaa ggc act ttt ggg aaa gtt att ttg gtt cga gag aag gca     527
Leu Gly Lys Gly Thr Phe Gly Lys Val Ile Leu Val Arg Glu Lys Ala
              160             165             170             175

agt gga aaa tac tat gct atg aag att ctg aag aaa gaa gtc att att     575
Ser Gly Lys Tyr Tyr Ala Met Lys Ile Leu Lys Lys Glu Val Ile Ile
              180             185             190

gca aag gat gaa gtg gca cac act cta act gaa agc aga gta tta aag     623
Ala Lys Asp Glu Val Ala His Thr Leu Thr Glu Ser Arg Val Leu Lys
              195             200             205

aac act aga cat ccc ttt tta aca tcc ttg aaa tat tcc ttc cag aca     671
Asn Thr Arg His Pro Phe Leu Thr Ser Leu Lys Tyr Ser Phe Gln Thr
              210             215             220

aaa gac cgt ttg tgt ttt gtg atg gaa tat gtt aat ggg ggc gag ctg     719
Lys Asp Arg Leu Cys Phe Val Met Glu Tyr Val Asn Gly Gly Glu Leu
              225             230             235

ttt ttc cat ttg tcg aga gag cgg gtg ttc tct gag gac cgc aca cgt     767
Phe Phe His Leu Ser Arg Glu Arg Val Phe Ser Glu Asp Arg Thr Arg
              240             245             250             255

ttc tat ggt gca gaa att gtc tct gcc ttg gac tat cta cat tcc gga     815
Phe Tyr Gly Ala Glu Ile Val Ser Ala Leu Asp Tyr Leu His Ser Gly
              260             265             270

aag att gtg tac cgt gat ctc aag ttg gag aat cta atg ctg gac aaa     863
Lys Ile Val Tyr Arg Asp Leu Lys Leu Glu Asn Leu Met Leu Asp Lys

```

-continued

275	280	285	
gat ggc cac ata aaa att aca gat ttt gga ctt tgc aaa gaa ggg atc Asp Gly His Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile 290 295 300			911
aca gat gca gcc acc atg aag aca ttc tgt ggc act cca gaa tat ctg Thr Asp Ala Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu 305 310 315			959
gca cca gag gtg tta gaa gat aat gac tat ggc cga gca gta gac tgg Ala Pro Glu Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp 320 325 330 335			1007
tgg ggc cta ggg gtt gtc atg tat gaa atg atg tgt ggg agg tta cct Trp Gly Leu Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro 340 345 350			1055
ttc tac aac cag gac cat gag aaa ctt ttt gaa tta ata tta atg gaa Phe Tyr Asn Gln Asp His Glu Lys Leu Phe Glu Leu Ile Leu Met Glu 355 360 365			1103
gac att aaa ttt cct cga aca ctc tct tca gat gca aaa tca ttg ctt Asp Ile Lys Phe Pro Arg Thr Leu Ser Ser Asp Ala Lys Ser Leu Leu 370 375 380			1151
tca ggg ctc ttg ata aag gat cca aat aaa cgc ctt ggt gga gga cca Ser Gly Leu Leu Ile Lys Asp Pro Asn Lys Arg Leu Gly Gly Gly Pro 385 390 395			1199
gat gat gca aaa gaa att atg aga cac agt ttc ttc tct gga gta aac Asp Asp Ala Lys Glu Ile Met Arg His Ser Phe Phe Ser Gly Val Asn 400 405 410 415			1247
tgg caa gat gta tat gat aaa aag ctt gta cct cct ttt aaa cct caa Trp Gln Asp Val Tyr Asp Lys Lys Leu Val Pro Phe Lys Pro Gln 420 425 430			1295
gta aca tct gag aca gat act aga tat ttt gat gaa gaa ttt aca gct Val Thr Ser Glu Thr Asp Thr Arg Tyr Phe Asp Glu Glu Phe Thr Ala 435 440 445			1343
cag act att aca ata aca cca cct gaa aaa tat gat gag gat ggt atg Gln Thr Ile Thr Ile Thr Pro Pro Glu Lys Tyr Asp Glu Asp Gly Met 450 455 460			1391
gac tgc atg gac aat gag agg cgg ccg cat ttc cct caa ttt tcc tac Asp Cys Met Asp Asn Glu Arg Arg Pro His Phe Pro Gln Phe Ser Tyr 465 470 475			1439
tct gca agt gga cga gaa taa gtctctttca ttctgtact tcaactgtcat Ser Ala Ser Gly Arg Glu 480 485			1490
cttcaattta ttactgaaaa			1510

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 485

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 6

Ala Gln Arg Gly Val Ile Met Ser Asp Val Thr Ile Val Lys Glu Gly  
1 5 10 15

Trp Val Gln Lys Arg Gly Glu Tyr Ile Lys Asn Trp Arg Pro Arg Tyr  
20 25 30

Phe Leu Leu Lys Thr Asp Gly Ser Phe Ile Gly Tyr Lys Glu Lys Pro  
35 40 45

Gln Asp Val Asp Leu Pro Tyr Pro Leu Asn Asn Phe Ser Val Ala Lys  
50 55 60

-continued

Cys	Gln	Leu	Met	Lys	Thr	Glu	Arg	Pro	Lys	Pro	Asn	Thr	Phe	Ile	Ile	65	70	75	80
Arg	Cys	Leu	Gln	Trp	Thr	Thr	Val	Ile	Glu	Arg	Thr	Phe	His	Val	Asp	85	90	95	
Thr	Pro	Glu	Glu	Arg	Glu	Glu	Trp	Thr	Glu	Ala	Ile	Gln	Ala	Val	Ala	100	105	110	
Asp	Arg	Leu	Gln	Arg	Gln	Glu	Glu	Glu	Arg	Met	Asn	Cys	Ser	Pro	Thr	115	120	125	
Ser	Gln	Ile	Asp	Asn	Ile	Gly	Glu	Glu	Glu	Met	Asp	Ala	Ser	Thr	Thr	130	135	140	
His	His	Lys	Arg	Lys	Thr	Met	Asn	Asp	Phe	Asp	Tyr	Leu	Lys	Leu	Leu	145	150	155	160
Gly	Lys	Gly	Thr	Phe	Gly	Lys	Val	Ile	Leu	Val	Arg	Glu	Lys	Ala	Ser	165	170	175	
Gly	Lys	Tyr	Tyr	Ala	Met	Lys	Ile	Leu	Lys	Lys	Glu	Val	Ile	Ile	Ala	180	185	190	
Lys	Asp	Glu	Val	Ala	His	Thr	Leu	Thr	Glu	Ser	Arg	Val	Leu	Lys	Asn	195	200	205	
Thr	Arg	His	Pro	Phe	Leu	Thr	Ser	Leu	Lys	Tyr	Ser	Phe	Gln	Thr	Lys	210	215	220	
Asp	Arg	Leu	Cys	Phe	Val	Met	Glu	Tyr	Val	Asn	Gly	Gly	Glu	Leu	Phe	225	230	235	240
Phe	His	Leu	Ser	Arg	Glu	Arg	Val	Phe	Ser	Glu	Asp	Arg	Thr	Arg	Phe	245	250	255	
Tyr	Gly	Ala	Glu	Ile	Val	Ser	Ala	Leu	Asp	Tyr	Leu	His	Ser	Gly	Lys	260	265	270	
Ile	Val	Tyr	Arg	Asp	Leu	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	275	280	285	
Gly	His	Ile	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Thr	290	295	300	
Asp	Ala	Ala	Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	305	310	315	320
Pro	Glu	Val	Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	325	330	335	
Gly	Leu	Gly	Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	340	345	350	
Tyr	Asn	Gln	Asp	His	Glu	Lys	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Asp	355	360	365	
Ile	Lys	Phe	Pro	Arg	Thr	Leu	Ser	Ser	Asp	Ala	Lys	Ser	Leu	Leu	Ser	370	375	380	
Gly	Leu	Leu	Ile	Lys	Asp	Pro	Asn	Lys	Arg	Leu	Gly	Gly	Gly	Pro	Asp	385	390	395	400
Asp	Ala	Lys	Glu	Ile	Met	Arg	His	Ser	Phe	Phe	Ser	Gly	Val	Asn	Trp	405	410	415	
Gln	Asp	Val	Tyr	Asp	Lys	Lys	Leu	Val	Pro	Pro	Phe	Lys	Pro	Gln	Val	420	425	430	
Thr	Ser	Glu	Thr	Asp	Thr	Arg	Tyr	Phe	Asp	Glu	Glu	Phe	Thr	Ala	Gln	435	440	445	
Thr	Ile	Thr	Ile	Thr	Pro	Pro	Glu	Lys	Tyr	Asp	Glu	Asp	Gly	Met	Asp	450	455	460	
Cys	Met	Asp	Asn	Glu	Arg	Arg	Pro	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser				

-continued


465                                      470                                      475                                      480

Ala Ser Gly Arg Glu  
485


What is claimed:

1. A composition comprising:


(i) a compound of the formula I-IV:




Formula I



Formula II



Formula III




Formula IV

wherein each  $R_2'$ ,  $R_3'$  and  $R_5'$  are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methane-sulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein  $R_2'$ ,  $R_3'$  or  $R_5'$  is independently H or mono-, di- or tri-phosphate;

wherein  $R^x$  and  $R^y$  are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid.

$R_1$  and  $R_2$  each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl;

(ii) one or more platinum compounds as in formula V,



Formula V

wherein:

$R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$ ;

$R_5$  and  $R_6$  are each independently halogen, optionally substituted alkoxy, optionally substituted ester, optionally substituted alkyl amine, aromatic amine, heteroaromatic amine;


(iii) a pharmaceutically acceptable carrier.

2. The composition of claim 1, wherein the compound of formula I is tricyribine.

3. The composition of claim 1, wherein the compound of formula I is tricyribine phosphate.

4. The composition of claim 1, wherein the compound of formula I is tricitriline phosphate monohydrate.

5. The composition of claim 1, wherein the platinum compounds of formula V is




wherein:


$R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$

$R_5$  and  $R_6$  are each independently halogen.


6. The composition of claim 1, wherein the one or more platinum compounds are selected from the group consisting of:




A




B



C




D




E

-continued




F




G



H



I



7. The composition of claim 1, wherein the compound of formula I is present in a dose amount of at least 10 mg/m<sup>2</sup>.

8. The composition of claim 1, wherein the compound of formula V is present in an amount of at least 1 mg/kg.

9. The composition of claim 1, suitable for parenteral administration.

10. The composition of claim 9, wherein the parenteral administration is intravenous administration.


11. The composition of claim 1, suitable for oral administration.


12. The composition of claim 1, suitable for topical administration.

13. A method of treating a tumor or cancer in a mammal comprising administering to the mammal an effective amount of a composition comprising:


a compound of formula I-IV:

Formula I






Formula II



Formula III



Formula IV


wherein each R<sub>2</sub>', R<sub>3</sub>' and R<sub>5</sub>' are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate prodrug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein R<sub>2</sub>', R<sub>3</sub>' or R<sub>5</sub>' is independently H or mono-, di- or tri-phosphate;

wherein R<sup>x</sup> and R<sup>y</sup> are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable

leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid.

R<sub>1</sub> and R<sub>2</sub> each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl; and

(ii) one or more platinum compounds of formula V,



Formula V

wherein

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>;

R<sub>5</sub> and R<sub>6</sub> are each independently halogen, optionally substituted alkoxy, optionally substituted ester, optionally substituted alkyl amine, aromatic amine, heteroaromatic amine.

**14.** The method of claim 13, wherein a compound of formula I-IV and a compound of formula V are administered simultaneously.

**15.** The method of claim 13, wherein a compound of formula I-IV is administered followed by the administration of a compound of formula V.

**16.** The method of claim 13, wherein a compound of formula V is administered followed by the administration of a compound of formula I-IV.

**17.** The method of claim 13, wherein the compound of formula I-IV is administered one time per week for three weeks followed by a one week period wherein the compound is not administered.

**18.** The method of claim 13, wherein the compound of formula V is administered one time per week for three weeks followed by a one week period wherein the compound of formula I-IV is not administered.

**19.** The method of claim 18, wherein the dosing schedule is repeated at least twice.

**20.** The method of claim 18, wherein the dosing schedule is repeated at least 4 times.

**21.** The method of claim 13, wherein the tumor treated is breast, pancreatic, ovarian and colorectal tumors.

**22.** The method of claim 13, wherein at least 10 g/m<sup>2</sup> of the compound of formula I-IV is administered.

**23.** The method of claim 13, wherein 100 mg/m<sup>2</sup> or less of the compound of formula I-IV is administered.

**24.** The method of claim 13, wherein at least 1 mg/kg of the compound of formula V is administered.

**25.** The method of claim 13, wherein 10 mg/kg or less of the compound of formula V is administered.

\* \* \* \* \*